

TABLE 3b. Estimated percent for time since first diagnosis among children¹ with active² asthma by state/territory - BRFSS Asthma Call-back Survey, United States, 2006-2010

State/Territory	Sample size [±]	5 or fewer years				More than 5 years			
		Sample size	%	SE [§]	95% CI [†]	Sample size	%	SE	95% CI
Arizona	150	69	59.0	6.1	(47.0 - 71.0)	81	41.0	6.1	(29.0 - 53.0)
California	215	104	48.9	4.6	(39.9 - 57.9)	111	51.1	4.6	(42.1 - 60.1)
Connecticut	426	223	52.2	3.2	(45.9 - 58.5)	203	47.8	3.2	(41.5 - 54.1)
District of Columbia	149	74	48.0	4.9	(38.4 - 57.6)	75	52.0	4.9	(42.4 - 61.6)
Georgia	416	196	52.8	3.5	(45.9 - 59.7)	220	47.2	3.5	(40.3 - 54.1)
Hawaii	453	212	51.6	3.0	(45.7 - 57.5)	241	48.4	3.0	(42.5 - 54.3)
Illinois	200	86	49.4	4.5	(40.6 - 58.2)	114	50.6	4.5	(41.8 - 59.4)
Indiana	397	200	57.0	3.2	(50.7 - 63.3)	197	43.0	3.2	(36.7 - 49.3)
Iowa	279	147	56.2	3.5	(49.3 - 63.1)	132	43.8	3.5	(36.9 - 50.7)
Kansas	671	331	54.5	2.3	(50.0 - 59.0)	340	45.5	2.3	(41.0 - 50.0)
Louisiana	122	50	47.3	5.5	(36.5 - 58.1)	72	52.7	5.5	(41.9 - 63.5)
Maine	343	161	49.6	3.1	(43.5 - 55.7)	182	50.4	3.1	(44.3 - 56.5)
Maryland	477	212	48.4	3.0	(42.5 - 54.3)	265	51.6	3.0	(45.7 - 57.5)
Massachusetts	323	141	47.9	4.2	(39.7 - 56.1)	182	52.1	4.2	(43.9 - 60.3)
Michigan	547	256	49.5	2.8	(44.0 - 55.0)	291	50.5	2.8	(45.0 - 56.0)
Mississippi	95	46	61.6	6.0	(49.8 - 73.4)	49	38.4	6.0	(26.6 - 50.2)
Missouri	259	134	53.8	4.6	(44.8 - 62.8)	125	46.2	4.6	(37.2 - 55.2)
Montana	257	130	56.0	3.6	(48.9 - 63.1)	127	44.0	3.6	(36.9 - 51.1)
Nebraska	551	279	54.7	3.6	(47.6 - 61.8)	272	45.3	3.6	(38.2 - 52.4)
New Hampshire	178	90	50.4	4.5	(41.6 - 59.2)	88	49.6	4.5	(40.8 - 58.4)
New Jersey	346	158	44.1	3.8	(36.7 - 51.5)	188	55.9	3.8	(48.5 - 63.3)
New Mexico	268	106	47.5	4.1	(39.5 - 55.5)	162	52.5	4.1	(44.5 - 60.5)
New York	347	160	52.8	3.7	(45.5 - 60.1)	187	47.2	3.7	(39.9 - 54.5)
North Dakota	124	64	57.8	5.2	(47.6 - 68.0)	60	42.2	5.2	(32.0 - 52.4)
Ohio	266	144	55.0	4.7	(45.8 - 64.2)	122	45.0	4.7	(35.8 - 54.2)
Oklahoma	261	133	54.7	3.8	(47.3 - 62.1)	128	45.3	3.8	(37.9 - 52.7)
Oregon	228	111	53.7	4.1	(45.7 - 61.7)	117	46.3	4.1	(38.3 - 54.3)
Pennsylvania	158	76	49.6	5.3	(39.2 - 60.0)	82	50.4	5.3	(40.0 - 60.8)
Rhode Island	189	85	51.0	4.3	(42.6 - 59.4)	104	49.0	4.3	(40.6 - 57.4)
Texas	576	268	56.1	3.5	(49.2 - 63.0)	308	43.9	3.5	(37.0 - 50.8)
Utah	426	221	54.4	3.3	(47.9 - 60.9)	205	45.6	3.3	(39.1 - 52.1)
Vermont	450	214	49.2	2.8	(43.7 - 54.7)	236	50.8	2.8	(45.3 - 56.3)
Virginia	101	49	51.2	8.0	(35.5 - 66.9)	52	48.8	8.0	(33.1 - 64.5)
Washington	397	190	52.7	3.2	(46.4 - 59.0)	207	47.3	3.2	(41.0 - 53.6)
West Virginia	223	98	48.4	3.9	(40.8 - 56.0)	125	51.6	3.9	(44.0 - 59.2)
Wisconsin	207	106	60.6	4.6	(51.6 - 69.6)	101	39.4	4.6	(30.4 - 48.4)
36 state total	11,075	5,324	52.1	1.1	(49.9 - 54.3)	5,751	47.9	1.1	(45.7 - 50.1)
Median			52.5				47.6		
Range			44.1-61.6				38.4-55.9		

¹ Aged < 18 years

² Reported any of the following during the past year: asthma medication use, asthma symptoms, or MD visits for asthma

[±] Sample size is for multiple years of participation. Responses of "Don't Know" and "Refused" are excluded.

[§] Standard error

[†] Confidence interval