

TABLE 8a. Estimated percent advised to make environmental changes for current¹ and former² asthma status among children³ ever diagnosed with asthma by state/territory - BRFSS Asthma Call-back Survey, United States, 2006-2010

State/Territory	Current				Former			
	Sample size [±]	%	SE [§]	95% CI [†]	Sample size	%	SE	95% CI
Arizona	125	34.8	6.5	(22.1 - 47.5)	66	N/A	N/A	(N/A - N/A)
California	178	45.6	4.9	(36.0 - 55.2)	97	33.7	7.5	(19.0 - 48.4)
Connecticut	356	47.8	3.6	(40.7 - 54.9)	173	40.8	4.7	(31.6 - 50.0)
District of Columbia	120	58.3	5.7	(47.1 - 69.5)	52	37.8	8.1	(21.9 - 53.7)
Georgia	355	42.3	3.5	(35.4 - 49.2)	174	30.3	5.1	(20.3 - 40.3)
Hawaii	426	34.7	2.9	(29.0 - 40.4)	254	28.5	3.6	(21.4 - 35.6)
Illinois	172	48.3	4.9	(38.7 - 57.9)	84	50.1	7.1	(36.2 - 64.0)
Indiana	351	44.0	3.5	(37.1 - 50.9)	134	36.9	5.5	(26.1 - 47.7)
Iowa	255	31.5	3.3	(25.0 - 38.0)	114	23.2	4.4	(14.6 - 31.8)
Kansas	608	40.8	2.4	(36.1 - 45.5)	203	33.5	4.2	(25.3 - 41.7)
Louisiana	106	33.9	5.3	(23.5 - 44.3)	51	47.0	7.9	(31.5 - 62.5)
Maine	292	41.5	3.4	(34.8 - 48.2)	135	35.7	5.2	(25.5 - 45.9)
Maryland	393	41.2	3.2	(34.9 - 47.5)	194	27.9	3.9	(20.3 - 35.5)
Massachusetts	272	45.0	4.4	(36.4 - 53.6)	137	35.1	5.6	(24.1 - 46.1)
Michigan	490	48.3	3.0	(42.4 - 54.2)	169	43.8	5.3	(33.4 - 54.2)
Mississippi	81	64.1	7.1	(50.2 - 78.0)	29	50.8	10.8	(29.6 - 72.0)
Missouri	234	47.6	4.8	(38.2 - 57.0)	88	50.1	7.3	(35.8 - 64.4)
Montana	212	39.4	4.3	(31.0 - 47.8)	114	25.6	4.7	(16.4 - 34.8)
Nebraska	488	39.5	3.8	(32.1 - 46.9)	207	32.7	5.8	(21.3 - 44.1)
New Hampshire	153	45.4	4.7	(36.2 - 54.6)	72	40.1	6.9	(26.6 - 53.6)
New Jersey	308	40.1	3.9	(32.5 - 47.7)	148	35.4	5.9	(23.8 - 47.0)
New Mexico	239	34.6	4.5	(25.8 - 43.4)	107	22.9	4.9	(13.3 - 32.5)
New York	284	51.8	3.9	(44.2 - 59.4)	147	35.2	5.1	(25.2 - 45.2)
North Dakota	102	38.8	5.6	(27.8 - 49.8)	51	36.1	7.7	(21.0 - 51.2)
Ohio	228	41.2	4.6	(32.2 - 50.2)	106	30.8	6.3	(18.5 - 43.1)
Oklahoma	254	34.1	3.6	(27.0 - 41.2)	74	30.4	6.0	(18.6 - 42.2)
Oregon	197	31.8	3.8	(24.4 - 39.2)	116	32.8	5.2	(22.6 - 43.0)
Pennsylvania	135	36.3	5.6	(25.3 - 47.3)	65	41.1	8.4	(24.6 - 57.6)
Rhode Island	160	41.3	4.9	(31.7 - 50.9)	64	34.9	7.0	(21.2 - 48.6)
Texas	492	34.3	3.2	(28.0 - 40.6)	261	40.4	5.1	(30.4 - 50.4)
Utah	377	26.3	2.9	(20.6 - 32.0)	170	28.7	5.2	(18.5 - 38.9)
Vermont	393	41.5	2.9	(35.8 - 47.2)	186	41.2	4.2	(33.0 - 49.4)
Virginia	87	41.7	8.6	(24.8 - 58.6)	53	28.8	7.6	(13.9 - 43.7)
Washington	350	39.5	3.4	(32.8 - 46.2)	219	37.3	4.5	(28.5 - 46.1)
West Virginia	203	40.2	4.0	(32.4 - 48.0)	67	24.0	5.6	(13.0 - 35.0)
Wisconsin	148	36.8	5.4	(26.2 - 47.4)	119	29.7	4.9	(20.1 - 39.3)
36 state total	9,624	42.4	1.1	(40.2 - 44.6)	4,500	36.1	1.8	(32.6 - 39.6)
Median		41.0				35.1		
Range		26.3-64.1				22.9-50.8		

¹ "Yes" response to "Do you still have asthma?"

² "No" response to "Do you still have asthma?"

³ Aged < 18 years

[±] Sample size is for multiple years of participation. Responses of "Don't Know" and "Refused" are excluded.

[§] Standard error

[†] Confidence interval

N/A: Estimate not available (N/A) if the unweighted sample size for the denominator was < 50 or if the relative standard error was > 0.30