

**TABLE 8b. Estimated percent advised to make environmental changes for active<sup>1</sup> and inactive<sup>2</sup> asthma status among children<sup>3</sup> ever diagnosed with asthma by state/territory - BRFSS Asthma Call-back Survey, United States, 2006-2010**

State/Territory	Active				Inactive			
	Sample size <sup>±</sup>	%	SE <sup>§</sup>	95% CI <sup>†</sup>	Sample size	%	SE	95% CI
Arizona	149	36.0	6.2	(23.8 - 48.2)	46	N/A	N/A	(N/A - N/A)
California	215	42.6	4.6	(33.6 - 51.6)	62	35.7	9.7	(16.7 - 54.7)
Connecticut	424	46.3	3.2	(40.0 - 52.6)	119	40.6	5.5	(29.8 - 51.4)
District of Columbia	147	53.8	5.1	(43.8 - 63.8)	32	53.5	12.7	(28.6 - 78.4)
Georgia	415	40.1	3.3	(33.6 - 46.6)	123	33.9	5.9	(22.3 - 45.5)
Hawaii	452	34.6	2.8	(29.1 - 40.1)	242	28.1	3.7	(20.8 - 35.4)
Illinois	200	49.4	4.5	(40.6 - 58.2)	57	47.8	9.2	(29.8 - 65.8)
Indiana	394	42.7	3.4	(36.0 - 49.4)	101	38.6	5.9	(27.0 - 50.2)
Iowa	282	29.4	3.1	(23.3 - 35.5)	91	27.0	5.4	(16.4 - 37.6)
Kansas	675	39.9	2.3	(35.4 - 44.4)	150	33.2	4.9	(23.6 - 42.8)
Louisiana	121	35.2	4.9	(25.6 - 44.8)	39	45.6	8.9	(28.2 - 63.0)
Maine	342	39.7	3.1	(33.6 - 45.8)	92	37.8	6.3	(25.5 - 50.1)
Maryland	473	39.1	2.9	(33.4 - 44.8)	139	26.8	4.6	(17.8 - 35.8)
Massachusetts	318	44.3	4.0	(36.5 - 52.1)	95	31.9	6.4	(19.4 - 44.4)
Michigan	546	50.2	2.8	(44.7 - 55.7)	130	34.6	6.2	(22.4 - 46.8)
Mississippi	95	61.8	6.5	(49.1 - 74.5)	16	58.5	13.1	(32.8 - 84.2)
Missouri	258	49.0	4.7	(39.8 - 58.2)	68	43.0	7.7	(27.9 - 58.1)
Montana	255	37.2	3.8	(29.8 - 44.6)	81	21.8	5.4	(11.2 - 32.4)
Nebraska	552	39.2	3.7	(31.9 - 46.5)	162	29.5	6.1	(17.5 - 41.5)
New Hampshire	176	44.2	4.4	(35.6 - 52.8)	53	39.1	7.8	(23.8 - 54.4)
New Jersey	350	39.5	3.8	(32.1 - 46.9)	108	35.1	6.5	(22.4 - 47.8)
New Mexico	267	34.2	4.2	(26.0 - 42.4)	85	22.9	5.6	(11.9 - 33.9)
New York	343	49.2	3.7	(41.9 - 56.5)	94	34.8	6.3	(22.5 - 47.1)
North Dakota	124	41.4	5.1	(31.4 - 51.4)	36	N/A	N/A	(N/A - N/A)
Ohio	265	42.7	4.4	(34.1 - 51.3)	79	N/A	N/A	(N/A - N/A)
Oklahoma	261	31.6	3.4	(24.9 - 38.3)	70	41.2	6.9	(27.7 - 54.7)
Oregon	226	30.2	3.4	(23.5 - 36.9)	100	38.4	5.8	(27.0 - 49.8)
Pennsylvania	158	40.0	5.3	(29.6 - 50.4)	50	26.9	7.8	(11.6 - 42.2)
Rhode Island	188	41.4	4.2	(33.2 - 49.6)	48	31.2	8.0	(15.5 - 46.9)
Texas	573	35.3	3.2	(29.0 - 41.6)	200	40.0	5.3	(29.6 - 50.4)
Utah	423	26.6	2.8	(21.1 - 32.1)	144	26.2	5.2	(16.0 - 36.4)
Vermont	448	42.3	2.8	(36.8 - 47.8)	147	38.0	4.5	(29.2 - 46.8)
Virginia	100	43.1	8.0	(27.4 - 58.8)	41	N/A	N/A	(N/A - N/A)
Washington	398	37.4	3.2	(31.1 - 43.7)	193	37.0	4.5	(28.2 - 45.8)
West Virginia	223	39.0	3.8	(31.6 - 46.4)	49	22.5	6.4	(10.0 - 35.0)
Wisconsin	207	34.4	4.3	(26.0 - 42.8)	64	33.4	7.3	(19.1 - 47.7)
36 state total	11,043	41.9	1.1	(39.7 - 44.1)	3,406	34.3	2.0	(30.4 - 38.2)
Median		40.0				35.0		
Range		26.6-61.8				21.8-58.5		

<sup>1</sup> Reported any of the following during the past year: asthma medication use, asthma symptoms, or MD visits for asthma

<sup>2</sup> Reported none of the following during the past year: asthma medication use, asthma symptoms, or MD visits for asthma

<sup>3</sup> Aged < 18 years

<sup>±</sup> Sample size is for multiple years of participation. Responses of "Don't Know" and "Refused" are excluded.

<sup>§</sup> Standard error

<sup>†</sup> Confidence interval

N/A: Estimate not available (N/A) if the unweighted sample size for the denominator was < 50 or if the relative standard error was > 0.30