

Table C2_1
Child Current Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2007

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *	Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	30,673	10.1	0.32	(9.5 - 10.8)	28,715	7.7	0.31	(7.1 - 8.4)
AK	474	9.3	1.90	(6.2 - 13.8)	463	4.5	1.17	(2.6 - 7.4)
AZ	669	9.9	1.95	(6.7 - 14.5)	567	4.9	1.08	(3.1 - 7.5)
CA	877	7.7	0.95	(6.0 - 9.7)	808	7.6	1.13	(5.7 - 10.1)
CT	981	11.3	1.21	(9.1 - 13.9)	920	7.7	1.05	(5.9 - 10.0)
DC	430	17.3	2.53	(12.9 - 22.9)	355	14.3	2.70	(9.7 - 20.4)
GA	1,068	11.5	1.32	(9.2 - 14.4)	995	8.0	1.16	(6.0 - 10.6)
HI	1,087	14.2	1.34	(11.7 - 17.0)	1,050	9.9	1.12	(7.9 - 12.3)
IL	737	8.8	1.40	(6.4 - 12.0)	747	6.7	1.13	(4.8 - 9.2)
IN	798	8.9	1.34	(6.6 - 11.9)	798	6.9	0.99	(5.2 - 9.1)
IA	789	5.2	0.90	(3.7 - 7.3)	713	5.1	0.82	(3.7 - 6.9)
KS	1,224	8.4	0.91	(6.8 - 10.3)	1,150	8.1	0.88	(6.5 - 10.0)
ME	817	10.8	1.25	(8.6 - 13.5)	768	6.1	0.93	(4.5 - 8.2)
MD	1,258	9.0	0.96	(7.3 - 11.0)	1,073	9.0	1.30	(6.8 - 11.9)
MA	608	12.5	1.71	(9.5 - 16.3)	557	8.5	1.47	(6.0 - 11.9)
MI	1,007	11.9	1.28	(9.7 - 14.7)	990	7.8	1.17	(5.8 - 10.4)
MS	1,082	12.4	1.29	(10.1 - 15.1)	1,077	8.4	1.05	(6.5 - 10.7)
MO	699	10.7	1.49	(8.1 - 14.0)	658	6.4	1.04	(4.6 - 8.8)
MT	764	7.4	1.14	(5.5 - 10.0)	728	6.5	1.11	(4.7 - 9.1)
NE	1,568	10.5	1.48	(7.9 - 13.8)	1,497	6.3	1.03	(4.6 - 8.7)
NV	571	6.8	1.15	(4.8 - 9.4)	591	5.9	1.32	(3.7 - 9.1)
NH	777	7.9	1.12	(6.0 - 10.4)	709	7.2	1.12	(5.3 - 9.7)
NJ	480	15.1	2.34	(11.0 - 20.3)	421	5.7	1.16	(3.9 - 8.5)
NM	964	10.2	1.21	(8.1 - 12.9)	857	6.9	1.11	(5.0 - 9.4)
NY	781	12.7	1.66	(9.8 - 16.3)	775	10.5	1.52	(7.9 - 13.9)
OH	1,483	10.3	0.96	(8.6 - 12.4)	1,364	7.1	0.87	(5.6 - 9.0)
OK	1,046	12.9	1.23	(10.7 - 15.5)	995	8.9	1.01	(7.1 - 11.1)

Notes:
* CI denotes confidence interval.
** Total includes data from 33 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table C2_1
Child Current Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2007

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
OR	625	7.7	1.22	(5.6 - 10.5)		610	9.0	1.35	(6.7 - 12.0)
PA	1,594	11.2	1.28	(9.0 - 14.0)		1,522	9.9	1.23	(7.8 - 12.6)
TX	1,343	10.2	1.13	(8.2 - 12.6)		1,224	8.0	1.13	(6.1 - 10.5)
UT	1,092	8.7	1.03	(6.9 - 10.9)		902	6.8	1.09	(5.0 - 9.3)
VT	886	7.7	1.08	(5.9 - 10.1)		862	7.8	1.01	(6.0 - 10.0)
VA	851	9.8	1.22	(7.6 - 12.5)		764	5.5	0.93	(4.0 - 7.6)
WV	544	10.6	1.51	(8.0 - 14.0)		516	9.8	1.48	(7.3 - 13.2)
WI	699	8.5	1.43	(6.1 - 11.7)		689	7.5	1.43	(5.1 - 10.8)
Territories									
PR	556	14.7	1.76	(11.6 - 18.5)		511	12.5	1.79	(9.4 - 16.5)

Notes:
* CI denotes confidence interval.
** Total includes data from 33 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.