

Table L2_1
Child Lifetime Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2007

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	30,794	15.7	0.40	(14.9 - 16.5)		28,803	11.3	0.37	(10.6 - 12.1)
AK	477	13.5	2.30	(9.6 - 18.7)		465	8.3	1.81	(5.4 - 12.6)
AZ	670	17.4	2.60	(12.8 - 23.1)		567	9.8	2.19	(6.3 - 15.0)
CA	879	14.9	1.39	(12.4 - 17.9)		808	11.4	1.34	(9.0 - 14.3)
CT	985	17.5	1.48	(14.7 - 20.6)		923	11.6	1.29	(9.3 - 14.4)
DC	432	22.7	2.82	(17.7 - 28.7)		355	16.9	2.82	(12.1 - 23.1)
GA	1,074	18.3	1.58	(15.4 - 21.6)		997	12.4	1.36	(10.0 - 15.3)
HI	1,091	21.1	1.54	(18.3 - 24.3)		1,054	15.1	1.32	(12.7 - 17.9)
IL	740	12.9	1.63	(10.0 - 16.4)		747	9.6	1.31	(7.3 - 12.5)
IN	800	13.2	1.52	(10.4 - 16.4)		799	9.3	1.15	(7.3 - 11.8)
IA	792	9.3	1.16	(7.3 - 11.8)		716	8.0	1.05	(6.2 - 10.3)
KS	1,227	12.8	1.10	(10.8 - 15.1)		1,152	9.6	0.96	(7.9 - 11.7)
ME	817	16.6	1.50	(13.8 - 19.7)		769	9.9	1.20	(7.8 - 12.5)
MD	1,264	15.0	1.28	(12.7 - 17.7)		1,078	12.4	1.42	(9.8 - 15.4)
MA	611	17.5	2.02	(13.9 - 21.8)		559	11.8	1.71	(8.8 - 15.6)
MI	1,011	17.1	1.45	(14.4 - 20.1)		994	11.2	1.38	(8.7 - 14.2)
MS	1,088	17.5	1.49	(14.7 - 20.6)		1,080	12.1	1.20	(9.9 - 14.6)
MO	701	15.2	1.68	(12.2 - 18.7)		659	10.8	1.46	(8.2 - 14.0)
MT	768	10.9	1.39	(8.5 - 14.0)		731	8.3	1.20	(6.2 - 11.0)
NE	1,575	14.2	1.62	(11.4 - 17.7)		1,500	7.9	1.14	(5.9 - 10.4)
NV	574	11.0	1.55	(8.4 - 14.5)		594	9.5	1.52	(6.9 - 12.9)
NH	781	13.1	1.41	(10.6 - 16.1)		712	9.5	1.27	(7.3 - 12.3)
NJ	482	20.3	2.55	(15.8 - 25.8)		422	8.7	1.53	(6.1 - 12.2)
NM	967	15.7	1.47	(13.0 - 18.8)		860	10.3	1.30	(8.0 - 13.1)
NY	785	18.0	1.81	(14.7 - 21.8)		778	14.2	1.69	(11.2 - 17.9)
OH	1,491	16.4	1.18	(14.2 - 18.8)		1,372	10.8	1.02	(9.0 - 13.0)
OK	1,049	18.0	1.41	(15.4 - 20.9)		997	12.9	1.21	(10.8 - 15.5)

Notes:
* CI denotes confidence interval.
** Total includes data from 33 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L2_1
Child Lifetime Asthma Prevalence Rate (Percent)
by Sex and State or Territory: BRFSS 2007

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI *		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI *
OR	628	12.1	1.50	(9.4 - 15.3)		612	11.7	1.50	(9.1 - 15.0)
PA	1,601	15.7	1.50	(13.0 - 18.9)		1,527	13.6	1.42	(11.0 - 16.6)
TX	1,348	15.1	1.34	(12.6 - 17.9)		1,228	12.1	1.33	(9.8 - 15.0)
UT	1,097	13.2	1.22	(11.0 - 15.8)		906	10.0	1.35	(7.6 - 13.0)
VT	889	12.7	1.32	(10.4 - 15.5)		867	12.0	1.20	(9.8 - 14.6)
VA	856	17.5	1.94	(14.0 - 21.6)		768	9.0	1.16	(6.9 - 11.5)
WV	544	14.8	1.70	(11.8 - 18.5)		517	12.5	1.66	(9.6 - 16.1)
WI	700	11.5	1.63	(8.6 - 15.1)		690	9.7	1.55	(7.0 - 13.2)
Territories									
PR	556	32.4	2.29	(28.1 - 37.0)		514	24.4	2.19	(20.4 - 29.0)

Notes:
* CI denotes confidence interval.
** Total includes data from 33 states plus the District of Columbia and excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.