

**Table L2\_1**  
**Child Lifetime Asthma Prevalence**  
**by Sex and State or Territory: BRFSS 2016**

State	Male Unweighted Number	Male Prevalence	Male Standard Error	Male 95% CI *		Female Unweighted Number	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	31,902	14.8	0.36	(14.1–15.5)		30,004	10.4	0.32	(9.8–11.0)
AZ	974	15.1	1.62	(12.2–18.5)		843	10.1	1.46	(7.5–13.3)
CA	1,342	15.5	1.19	(13.3–18.0)		1,241	10.8	1.07	(8.9–13.1)
CT	1,100	18.9	1.47	(16.2–22.0)		1,026	13.5	1.35	(11.0–16.3)
FL	3,420	14.1	1.07	(12.1–16.3)		3,172	11.9	1.05	(10.0–14.1)
GA	475	17.4	2.13	(13.6–22.0)		436	11.4	1.75	(8.4–15.2)
HI	837	19.5	1.75	(16.3–23.1)		788	14.8	1.64	(11.9–18.3)
IL	489	13.2	1.72	(10.2–17.0)		489	9.5	1.52	(6.9–12.9)
IN	1,080	15.0	1.36	(12.5–17.8)		1,021	11.3	1.21	(9.1–13.9)
KS	1,374	15.6	1.17	(13.4–18.0)		1,295	9.1	0.93	(7.4–11.1)
KY	1,060	15.7	1.53	(12.9–18.9)		1,091	9.0	1.16	(6.9–11.5)
ME	977	16.1	1.70	(13.1–19.8)		932	10.2	1.38	(7.8–13.2)
MA	366	17.6	2.62	(13.1–23.3)		326	10.4	2.07	(7.0–15.2)
MI	1,357	15.7	1.21	(13.5–18.2)		1,314	9.3	0.92	(7.7–11.3)
MN	1,950	12.1	0.96	(10.3–14.1)		1,783	9.0	0.81	(7.6–10.7)
MS	592	13.6	1.77	(10.4–17.4)		552	8.7	1.37	(6.4–11.8)
MO	654	16.0	2.00	(12.5–20.3)		654	9.6	1.52	(7.0–13.0)
MT	667	8.5	1.42	(6.1–11.7)		625	8.2	1.72	(5.4–12.2)
NE	1,844	11.4	1.17	(9.2–13.9)		1,787	6.6	0.82	(5.2–8.4)
NV	462	15.1	2.50	(10.9–20.7)		460	7.9	1.76	(5.1–12.1)
NJ	807	13.2	1.57	(10.4–16.6)		761	11.1	1.62	(8.3–14.7)
NM	706	15.2	1.90	(11.8–19.3)		616	12.4	1.79	(9.3–16.4)
NY	1,172	16.0	1.84	(12.7–20.0)		1,101	11.2	1.55	(8.5–14.6)
OH	1,205	12.0	1.23	(9.8–14.6)		1,180	9.5	1.24	(7.4–12.2)
OK	590	16.7	1.84	(13.4–20.6)		602	10.7	1.54	(8.0–14.1)
OR	559	13.3	1.62	(10.4–16.8)		508	9.0	1.45	(6.5–12.3)
PA	737	14.2	1.55	(11.4–17.5)		714	11.2	1.44	(8.6–14.3)
RI	560	16.3	2.12	(12.6–20.9)		551	12.4	1.92	(9.1–16.7)
UT	1,837	11.1	0.96	(9.3–13.1)		1,652	6.1	0.67	(4.9–7.6)
VT	676	14.5	1.75	(11.4–18.3)		575	11.4	2.40	(7.4–17.0)
WA	1,503	13.1	1.17	(11.0–15.6)		1,414	9.2	0.93	(7.5–11.2)

Notes:

\* CI denotes confidence interval.

\*\* Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table L2\_1**  
**Child Lifetime Asthma Prevalence**  
**by Sex and State or Territory: BRFSS 2016**

<b>State</b>	<b>Male Unweighted Number</b>	<b>Male Prevalence</b>	<b>Male Standard Error</b>	<b>Male 95% CI *</b>	<b>Female Unweighted Number</b>	<b>Female Prevalence</b>	<b>Female Standard Error</b>	<b>Female 95% CI *</b>
WI	530	14.0	2.27	(10.1–19.1)	495	9.3	1.83	(6.3–13.6)
Territory								
PR	616	26.8	2.50	(22.1–31.9)	550	21.5	2.53	(17.0–26.9)

Notes:

\* CI denotes confidence interval.

\*\* Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.