

**Table L3
Child Lifetime Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2016**

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
Total **	0-4	12,717	5.2	0.35	(4.5–5.9)		552,022	(476,920–627,123)
Total **	5-9	13,450	12.8	0.50	(11.8–13.8)		1,572,340	(1,444,086–1,700,593)
Total **	10-14	16,110	16.0	0.53	(15.0–17.1)		1,815,374	(1,689,595–1,941,153)
Total **	15-17	13,208	17.7	0.63	(16.5–19.0)		1,440,523	(1,330,395–1,550,651)
AZ	0-4	305	5.8	2.36	(2.6–12.6)		23,663	(3,934–43,393)
AZ	5-9	400	12.8	2.27	(9.0–17.9)		47,524	(30,015–65,034)
AZ	10-14	509	16.4	2.11	(12.7–21.0)		68,344	(49,985–86,702)
AZ	15-17	449	17.7	2.57	(13.2–23.3)		46,673	(32,146–61,201)
CA	0-4	594	4.6	1.07	(2.9–7.2)		71,907	(38,315–105,500)
CA	5-9	582	12.5	1.50	(9.9–15.8)		272,107	(205,111–339,104)
CA	10-14	650	17.6	1.87	(14.3–21.6)		304,239	(234,575–373,902)
CA	15-17	509	16.5	1.95	(13.0–20.7)		250,696	(187,551–313,841)
CT	0-4	351	8.2	1.90	(5.2–12.8)		14,585	(7,676–21,494)
CT	5-9	477	15.4	1.99	(11.9–19.8)		29,606	(21,600–37,612)
CT	10-14	603	21.0	2.10	(17.1–25.4)		34,745	(27,106–42,384)
CT	15-17	548	21.0	2.26	(17.0–25.8)		27,446	(20,992–33,900)
FL	0-4	1,334	6.4	1.15	(4.5–9.1)		46,891	(29,911–63,872)
FL	5-9	1,447	14.1	1.43	(11.5–17.1)		126,532	(100,419–152,645)
FL	10-14	1,602	14.6	1.68	(11.6–18.2)		132,678	(100,596–164,759)
FL	15-17	1,248	17.7	2.10	(14.0–22.2)		109,842	(81,705–137,978)
GA	0-4	145	3.9	1.71	(1.7–9.0)		22,604	(3,132–42,075)
GA	5-9	194	20.0	3.38	(14.2–27.5)		109,055	(69,441–148,669)
GA	10-14	244	16.5	2.87	(11.6–22.9)		77,476	(49,093–105,858)
GA	15-17	199	23.6	4.00	(16.7–32.3)		93,710	(57,428–129,992)
HI	0-4	369	6.0	1.38	(3.8–9.4)		3,421	(1,867–4,975)
HI	5-9	353	18.2	2.62	(13.6–23.9)		14,649	(10,091–19,208)
HI	10-14	413	21.0	2.52	(16.4–26.3)		14,664	(10,843–18,485)
HI	15-17	286	25.6	3.30	(19.7–32.6)		12,326	(8,767–15,886)
IL	0-4	178	4.1	1.66	(1.8–8.9)		27,598	(5,337–49,858)
IL	5-9	197	8.1	1.95	(5.0–12.8)		60,998	(31,749–90,248)
IL	10-14	275	18.9	2.74	(14.1–24.9)		134,654	(92,616–176,692)

Notes:

* CI denotes confidence interval.

** Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Child Lifetime Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2016

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
IL	15-17	252	16.4	2.84	(11.6–22.7)		76,146	(48,128–104,163)
IN	0-4	362	6.3	1.53	(3.9–10.0)		23,797	(12,145–35,448)
IN	5-9	445	14.5	2.04	(10.9–19.0)		61,867	(43,639–80,096)
IN	10-14	595	16.3	1.88	(13.0–20.4)		52,643	(39,836–65,449)
IN	15-17	488	18.5	2.16	(14.6–23.1)		48,704	(36,499–60,910)
KS	0-4	576	5.7	1.18	(3.7–8.5)		8,398	(4,892–11,904)
KS	5-9	565	12.3	1.57	(9.6–15.8)		24,032	(17,738–30,325)
KS	10-14	691	15.5	1.60	(12.6–18.9)		26,027	(20,383–31,671)
KS	15-17	563	16.9	1.81	(13.6–20.7)		18,358	(14,279–22,437)
KY	0-4	444	6.1	1.47	(3.8–9.7)		14,622	(7,626–21,618)
KY	5-9	434	11.1	1.88	(8.0–15.4)		25,606	(16,822–34,391)
KY	10-14	529	15.2	2.05	(11.6–19.7)		38,874	(28,251–49,498)
KY	15-17	568	20.8	2.77	(15.8–26.7)		29,356	(20,609–38,104)
ME	0-4	365	5.5	1.83	(2.8–10.4)		3,214	(1,045–5,383)
ME	5-9	429	12.2	1.98	(8.9–16.7)		7,889	(5,297–10,482)
ME	10-14	545	17.8	2.40	(13.6–23.0)		10,821	(7,681–13,961)
ME	15-17	438	22.5	3.15	(16.9–29.3)		9,786	(6,706–12,865)
MA	0-4	122	4.1	2.34	(1.3–12.1)		10,123	(0–21,689)
MA	5-9	154	8.8	2.53	(4.9–15.2)		26,625	(11,297–41,952)
MA	10-14	177	21.5	4.20	(14.4–30.8)		71,543	(40,492–102,593)
MA	15-17	151	22.9	4.62	(15.1–33.2)		51,711	(28,734–74,687)
MI	0-4	525	5.6	1.22	(3.7–8.6)		28,933	(16,382–41,484)
MI	5-9	592	11.6	1.52	(8.9–14.9)		63,668	(46,533–80,803)
MI	10-14	689	15.5	1.51	(12.8–18.7)		76,647	(61,321–91,974)
MI	15-17	653	19.0	1.95	(15.4–23.1)		65,518	(51,085–79,951)
MN	0-4	759	3.2	0.81	(1.9–5.2)		6,729	(3,285–10,173)
MN	5-9	819	8.7	1.17	(6.7–11.3)		30,850	(22,410–39,290)
MN	10-14	911	15.0	1.41	(12.4–18.0)		43,278	(34,765–51,792)
MN	15-17	746	15.7	1.77	(12.6–19.5)		33,344	(25,312–41,377)
MS	0-4	250	6.0	1.82	(3.3–10.7)		10,253	(4,027–16,479)
MS	5-9	289	10.8	1.97	(7.5–15.3)		17,815	(11,262–24,368)

Notes:

* CI denotes confidence interval.

** Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Child Lifetime Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2016

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
MS	10-14	276	15.5	2.95	(10.6–22.2)		25,765	(15,375–36,155)
MS	15-17	224	13.9	2.64	(9.5–19.9)		16,816	(10,253–23,380)
MO	0-4	252	6.1	1.59	(3.7–10.1)		18,618	(9,055–28,180)
MO	5-9	282	12.0	2.74	(7.6–18.5)		41,464	(21,669–61,259)
MO	10-14	367	15.6	2.68	(11.0–21.6)		51,749	(33,159–70,339)
MO	15-17	284	21.0	3.47	(15.0–28.6)		43,119	(27,041–59,198)
MT	0-4	296	3.2	1.38	(1.4–7.3)		1,572	(238–2,907)
MT	5-9	292	8.8	2.35	(5.1–14.6)		5,298	(2,433–8,164)
MT	10-14	333	11.1	2.52	(7.0–17.0)		5,899	(3,124–8,675)
MT	15-17	254	10.4	2.87	(5.9–17.5)		3,789	(1,638–5,939)
NE	0-4	862	3.6	1.00	(2.1–6.2)		3,799	(1,720–5,879)
NE	5-9	822	5.9	1.24	(3.9–8.9)		7,232	(4,171–10,293)
NE	10-14	921	12.8	1.54	(10.1–16.2)		15,795	(11,892–19,698)
NE	15-17	741	17.0	2.30	(12.9–22.0)		12,036	(8,532–15,540)
NV	0-4	183	3.5	1.75	(1.3–9.1)		5,473	(0–10,999)
NV	5-9	198	16.5	3.76	(10.4–25.2)		29,519	(15,298–43,739)
NV	10-14	223	14.8	3.57	(9.1–23.2)		17,898	(8,727–27,069)
NV	15-17	194	9.9	2.60	(5.8–16.3)		9,817	(4,657–14,976)
NJ	0-4	271	4.9	1.70	(2.5–9.5)		23,728	(7,308–40,147)
NJ	5-9	339	19.5	3.04	(14.2–26.2)		81,051	(53,748–108,354)
NJ	10-14	389	11.2	2.08	(7.7–16.0)		43,018	(26,701–59,334)
NJ	15-17	322	15.4	2.79	(10.7–21.7)		41,430	(25,909–56,951)
NM	0-4	247	9.4	2.72	(5.2–16.2)		12,495	(4,987–20,003)
NM	5-9	295	15.5	2.89	(10.6–22.0)		18,241	(10,994–25,488)
NM	10-14	366	19.3	2.77	(14.5–25.4)		19,410	(13,520–25,300)
NM	15-17	268	15.9	2.90	(11.0–22.4)		11,486	(7,129–15,843)
NY	0-4	504	4.8	1.75	(2.4–9.7)		43,786	(11,982–75,591)
NY	5-9	462	15.7	2.98	(10.7–22.5)		149,144	(89,877–208,411)
NY	10-14	587	14.0	2.22	(10.2–19.0)		134,399	(90,759–178,039)
NY	15-17	455	20.3	3.17	(14.7–27.2)		132,226	(88,448–176,004)
OH	0-4	446	5.3	1.35	(3.2–8.7)		30,848	(15,282–46,414)

Notes:

* CI denotes confidence interval.

** Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Child Lifetime Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2016

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
OH	5-9	504	9.1	1.72	(6.2–13.0)		52,306	(32,084–72,528)
OH	10-14	640	15.7	1.94	(12.2–19.9)		91,822	(67,939–115,705)
OH	15-17	503	13.1	1.93	(9.7–17.3)		55,588	(38,889–72,288)
OK	0-4	276	3.9	1.17	(2.1–6.9)		8,107	(3,266–12,949)
OK	5-9	292	17.6	2.77	(12.8–23.7)		45,828	(30,283–61,372)
OK	10-14	322	19.0	2.54	(14.5–24.5)		41,654	(29,939–53,370)
OK	15-17	226	18.6	3.16	(13.2–25.6)		27,509	(17,386–37,632)
OR	0-4	249	3.5	1.61	(1.4–8.5)		6,098	(550–11,646)
OR	5-9	253	9.2	1.94	(6.1–13.8)		18,336	(10,694–25,978)
OR	10-14	268	16.6	2.57	(12.1–22.2)		36,978	(25,105–48,850)
OR	15-17	200	16.2	2.85	(11.3–22.6)		25,134	(16,071–34,196)
PA	0-4	334	7.3	1.81	(4.4–11.7)		39,148	(19,540–58,755)
PA	5-9	320	12.8	2.26	(9.0–18.0)		86,502	(55,045–117,960)
PA	10-14	352	14.9	2.30	(11.0–20.0)		96,467	(65,739–127,194)
PA	15-17	306	19.7	2.76	(14.8–25.6)		85,252	(60,079–110,426)
RI	0-4	203	2.9	1.42	(1.1–7.4)		1,553	(54–3,051)
RI	5-9	226	18.4	3.40	(12.6–26.0)		10,054	(6,070–14,039)
RI	10-14	342	17.0	2.67	(12.4–22.9)		6,979	(4,650–9,309)
RI	15-17	244	21.1	3.65	(14.9–29.2)		8,056	(4,922–11,189)
UT	0-4	766	1.8	0.53	(1.0–3.2)		3,368	(1,468–5,268)
UT	5-9	698	7.1	1.18	(5.1–9.8)		18,444	(12,294–24,593)
UT	10-14	905	13.9	1.46	(11.3–17.0)		29,386	(22,963–35,808)
UT	15-17	731	13.1	1.56	(10.3–16.4)		17,902	(13,513–22,292)
VT	0-4	286	11.7	3.10	(6.8–19.2)		3,078	(1,345–4,811)
VT	5-9	256	10.1	2.20	(6.5–15.3)		3,148	(1,777–4,519)
VT	10-14	348	14.2	2.30	(10.2–19.3)		4,019	(2,691–5,347)
VT	15-17	296	19.9	4.90	(11.9–31.2)		3,994	(1,719–6,269)
WA	0-4	636	5.7	1.26	(3.7–8.8)		18,833	(10,483–27,183)
WA	5-9	609	11.7	1.65	(8.9–15.4)		53,338	(37,838–68,838)
WA	10-14	782	12.6	1.44	(10.0–15.7)		49,918	(38,029–61,808)
WA	15-17	627	16.5	2.03	(12.9–20.8)		40,586	(29,815–51,358)

Notes:

* CI denotes confidence interval.

** Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L3
Child Lifetime Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2016

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
WI	0-4	227	5.7	2.24	(2.6–12.0)		14,778	(2,918–26,638)
WI	5-9	225	10.4	3.12	(5.6–18.3)		33,611	(12,460–54,762)
WI	10-14	256	17.5	3.50	(11.7–25.4)		57,584	(32,750–82,418)
WI	15-17	235	15.4	2.92	(10.5–22.0)		32,165	(19,471–44,859)
Territory								
PR	0-4	251	22.1	4.08	(15.1–31.1)		44,469	(25,325–63,613)
PR	5-9	259	20.0	2.95	(14.8–26.4)		38,955	(27,607–50,302)
PR	10-14	300	30.8	3.65	(24.1–38.4)		56,572	(42,240–70,904)
PR	15-17	264	22.6	3.68	(16.2–30.6)		31,545	(20,264–42,826)

Notes:

* CI denotes confidence interval.

** Total includes data from 32 states based on cellphone and landline data combined excludes the territory, Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.