Table C1Child Current Asthma Prevalenceand Weighted Number by State or Territory: BRFSS 2018

State	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)		Weighted Number	95% CI * (Number)
Total **	53,869	7.2	0.22	(6.7–7.6)	П	3,243,220	(3,050,289–3,436,152)
CA	2,859	6.2	0.58	(5.1–7.4)	11	542,696	(441,417–643,976)
СТ	1,976	9.7	0.85	(8.2–11.5)	1	62,383	(51,432–73,334)
DC	712	10.9	1.40	(8.4–14.0)	1	12,688	(9,431–15,945)
FL	2,848	7.5	0.94	(5.8–9.5)	1	302,125	(227,331–376,919)
GA	2,052	7.6	0.72	(6.3–9.1)	1	177,862	(144,489–211,235)
HI	1,831	10.2	0.92	(8.5–12.2)	1	30,003	(24,541–35,464)
IL	1,252	6.0	0.82	(4.6–7.8)	1	163,940	(119,933–207,947)
IN	1,390	8.7	0.95	(7.0–10.7)	1	115,612	(90,392–140,831)
KS	2,340	7.6	0.67	(6.4–9.1)	1	51,731	(42,747–60,715)
KY	1,857	5.9	0.74	(4.6–7.5)	1	57,017	(43,281–70,753)
ME	1,685	7.4	0.96	(5.7–9.5)	1	17,157	(12,668–21,647)
MD	1,051	7.6	1.14	(5.6–10.1)	1	91,975	(64,418–119,532)
MA	590	6.4	1.10	(4.6–8.9)	1	79,567	(53,433–105,701)
MI	2,371	8.6	0.74	(7.2–10.2)	1	176,987	(146,738–207,235)
MN	4,051	5.3	0.47	(4.4–6.3)	1	66,701	(55,084–78,318)
MS	1,413	6.5	0.72	(5.2–8.1)	1	43,220	(33,869–52,571)
MO	990	8.4	1.22	(6.3–11.1)	1	108,246	(76,890–139,603)
MT	1,151	5.3	0.84	(3.9–7.2)		11,736	(8,131–15,342)
NE	3,437	5.0	0.57	(4.0–6.3)	1	23,551	(18,290–28,812)
NV	731	7.0	1.44	(4.6–10.4)	1	45,915	(27,107–64,723)
NH	977	8.5	1.05	(6.7–10.8)	1	19,479	(14,745–24,212)
NJ	383	7.2	1.67	(4.6–11.3)	11	133,141	(75,534–190,748)
NM	1,570	7.5	1.01	(5.8–9.8)	П	35,177	(25,742–44,612)
NY	2,416	6.2	0.76	(4.8–7.8)	11	207,867	(157,416–258,317)
ОН	2,512	7.8	0.78	(6.4–9.4)	1	189,425	(152,158–226,691)
OR	1,235	7.3	1.10	(5.4–9.7)	П	58,704	(40,825–76,583)
PA	1,274	9.7	1.36	(7.4–12.7)	1	242,183	(174,373–309,993)
RI	1,032	7.9	1.28	(5.7–10.8)	1	14,681	(9,832–19,531)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Asthma and Community Health Branch, National Center for Environmental Health Centers for Disease Control and Prevention

Table C1Child Current Asthma Prevalenceand Weighted Number by State or Territory: BRFSS 2018

State	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)		Weighted Number	95% CI * (Number)
UT	3,689	5.5	0.46	(4.7–6.5)	Ш	49,652	(41,527–57,778)
VT	1,217	8.8	1.15	(6.8–11.3)	П	8,771	(6,495–11,046)
WI	977	8.4	1.25	(6.3–11.2)		103,029	(72,919–133,139)
Territory							
PR	1,194	14.1	1.18	(12.0–16.6)	П	97,765	(81,279–114,251)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Asthma and Community Health Branch, National Center for Environmental Health Centers for Disease Control and Prevention