

**Table C2\_1**  
**Child Current Asthma Prevalence**  
**by Sex and State or Territory: BRFSS 2018**

State	Male Unweighted Number	Male Prevalence	Male Standard Error	Male 95% CI *		Female Unweighted Number	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	27,325	8.2	0.33	(7.6–8.9)		25,721	6.0	0.27	(5.5–6.6)
CA	1,445	6.7	0.86	(5.2–8.7)		1,387	5.1	0.71	(3.9–6.7)
CT	963	10.3	1.19	(8.2–12.9)		965	9.0	1.26	(6.9–11.8)
DC	349	10.3	1.95	(7.1–14.8)		353	11.6	2.03	(8.2–16.2)
FL	1,411	9.2	1.49	(6.7–12.6)		1,386	5.7	1.21	(3.7–8.6)
GA	1,070	9.8	1.19	(7.7–12.4)		946	5.4	0.80	(4.0–7.2)
HI	939	11.5	1.27	(9.2–14.2)		866	9.0	1.37	(6.7–12.1)
IL	641	6.0	1.16	(4.1–8.7)		592	6.0	1.17	(4.1–8.8)
IN	693	9.2	1.39	(6.8–12.3)		681	7.9	1.25	(5.8–10.7)
KS	1,171	8.7	1.04	(6.9–11.0)		1,131	6.6	0.88	(5.1–8.6)
KY	957	6.2	0.99	(4.5–8.4)		887	5.7	1.13	(3.8–8.3)
ME	845	10.5	1.66	(7.7–14.2)		810	4.3	0.92	(2.8–6.5)
MD	530	7.7	1.66	(5.0–11.7)		507	7.6	1.61	(5.0–11.4)
MA	291	6.9	1.44	(4.5–10.3)		279	6.3	1.78	(3.6–10.8)
MI	1,195	11.3	1.21	(9.2–13.9)		1,146	5.8	0.85	(4.4–7.7)
MN	2,127	6.0	0.72	(4.7–7.6)		1,847	4.6	0.60	(3.6–5.9)
MS	702	7.6	1.10	(5.7–10.0)		702	5.4	0.92	(3.9–7.5)
MO	512	11.0	2.06	(7.5–15.7)		465	5.8	1.30	(3.7–8.9)
MT	599	4.6	0.89	(3.1–6.6)		533	5.8	1.38	(3.6–9.1)
NE	1,785	5.0	0.78	(3.7–6.8)		1,613	5.2	0.84	(3.7–7.1)
NV	376	7.0	1.92	(4.1–11.9)		346	7.0	2.16	(3.8–12.6)
NH	486	9.6	1.64	(6.8–13.3)		474	7.2	1.31	(5.0–10.2)
NJ	202	8.8	2.61	(4.8–15.4)		176	5.9	1.86	(3.2–10.8)
NM	808	7.8	1.43	(5.5–11.1)		744	7.3	1.45	(4.9–10.7)
NY	1,241	6.5	1.07	(4.7–8.9)		1,137	5.7	1.07	(3.9–8.2)
OH	1,284	9.0	1.14	(7.0–11.5)		1,190	6.4	1.07	(4.6–8.9)
OR	645	6.1	0.97	(4.5–8.3)		565	8.2	1.99	(5.0–13.0)
PA	634	12.1	2.36	(8.2–17.5)		621	7.6	1.22	(5.5–10.3)
RI	528	8.8	1.63	(6.0–12.5)		496	6.7	1.95	(3.8–11.7)
UT	1,785	6.2	0.70	(5.0–7.7)		1,826	4.8	0.61	(3.7–6.1)
VT	603	8.7	1.60	(6.0–12.4)		592	9.1	1.67	(6.3–12.9)

Notes:

\* CI denotes confidence interval.

\*\* Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table C2\_1**  
**Child Current Asthma Prevalence**  
**by Sex and State or Territory: BRFSS 2018**

State	Male Unweighted Number	Male Prevalence	Male Standard Error	Male 95% CI *		Female Unweighted Number	Female Prevalence	Female Standard Error	Female 95% CI *
WI	508	11.1	2.09	(7.6–15.9)		458	5.9	1.36	(3.8–9.2)
Territory									
PR	606	14.6	1.60	(11.7–18.0)		581	13.8	1.73	(10.7–17.5)

Notes:

\* CI denotes confidence interval.

\*\* Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.