

Table C3
Child Current Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2018

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
Total **	0-4	12,398	3.4	0.31	(2.8–4.1)		415,025	(340,046–490,003)
Total **	5-9	12,218	7.6	0.45	(6.7–8.5)		953,213	(837,337–1,069,088)
Total **	10-14	17,592	9.0	0.43	(8.2–9.9)		1,131,950	(1,022,402–1,241,499)
Total **	15-17	11,661	9.3	0.54	(8.3–10.4)		743,033	(655,445–830,620)
CA	0-4	707	2.4	0.80	(1.2–4.6)		56,369	(18,379–94,358)
CA	5-9	712	6.5	1.15	(4.6–9.2)		159,113	(102,409–215,817)
CA	10-14	885	8.8	1.23	(6.7–11.5)		211,323	(151,096–271,549)
CA	15-17	555	7.5	1.50	(5.0–11.0)		115,892	(68,623–163,161)
CT	0-4	425	5.6	1.28	(3.6–8.7)		8,519	(4,652–12,387)
CT	5-9	391	9.0	1.65	(6.3–12.8)		16,499	(10,428–22,570)
CT	10-14	661	12.0	1.82	(8.9–16.1)		21,518	(14,714–28,322)
CT	15-17	499	12.3	1.93	(9.0–16.6)		15,847	(10,714–20,980)
DC	0-4	246	4.7	1.38	(2.6–8.3)		1,495	(622–2,367)
DC	5-9	168	9.5	2.47	(5.6–15.6)		3,748	(1,775–5,720)
DC	10-14	177	17.9	3.46	(12.1–25.7)		5,079	(2,981–7,177)
DC	15-17	121	14.1	4.02	(7.9–24.0)		2,367	(939–3,795)
FL	0-4	723	4.6	1.57	(2.3–8.8)		47,846	(14,879–80,814)
FL	5-9	611	6.7	1.69	(4.0–10.8)		76,662	(37,944–115,380)
FL	10-14	959	10.5	2.14	(7.0–15.5)		117,607	(67,671–167,543)
FL	15-17	555	8.2	2.13	(4.9–13.4)		60,010	(28,403–91,617)
GA	0-4	410	3.2	1.09	(1.6–6.1)		19,712	(6,220–33,203)
GA	5-9	453	9.2	1.48	(6.7–12.5)		60,509	(40,843–80,176)
GA	10-14	764	8.2	1.29	(6.0–11.1)		52,739	(35,845–69,634)
GA	15-17	425	10.7	1.96	(7.4–15.1)		44,901	(27,855–61,948)
HI	0-4	608	5.6	1.15	(3.7–8.3)		4,448	(2,603–6,293)
HI	5-9	456	12.8	1.93	(9.4–17.0)		10,869	(7,454–14,284)
HI	10-14	475	12.4	1.95	(9.1–16.7)		10,137	(6,839–13,436)
HI	15-17	292	9.7	2.34	(5.9–15.3)		4,548	(2,266–6,830)
IL	0-4	235	3.9	1.53	(1.8–8.3)		29,689	(6,531–52,847)
IL	5-9	326	4.8	1.26	(2.9–8.0)		35,683	(17,327–54,039)
IL	10-14	347	6.8	1.64	(4.2–10.8)		51,522	(26,478–76,565)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table C3
Child Current Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2018**

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
IL	15-17	344	10.0	2.28	(6.4–15.4)		47,046	(24,879–69,213)
IN	0-4	279	2.9	0.92	(1.5–5.4)		10,267	(3,873–16,661)
IN	5-9	320	13.4	2.33	(9.5–18.6)		50,925	(32,340–69,510)
IN	10-14	466	9.2	1.88	(6.2–13.6)		33,311	(19,344–47,277)
IN	15-17	325	9.0	1.81	(6.0–13.2)		21,109	(12,495–29,723)
KS	0-4	598	3.9	0.98	(2.4–6.4)		7,198	(3,644–10,752)
KS	5-9	527	7.7	1.44	(5.3–11.1)		14,870	(9,262–20,478)
KS	10-14	707	10.3	1.33	(8.0–13.2)		19,445	(14,384–24,506)
KS	15-17	508	8.9	1.66	(6.2–12.7)		10,218	(6,423–14,014)
KY	0-4	316	1.8	0.64	(0.9–3.6)		4,761	(1,427–8,095)
KY	5-9	375	3.0	0.94	(1.6–5.5)		8,089	(3,142–13,037)
KY	10-14	654	12.0	2.14	(8.4–16.8)		31,788	(20,156–43,419)
KY	15-17	512	7.6	1.53	(5.1–11.2)		12,379	(7,605–17,153)
ME	0-4	337	6.9	2.34	(3.5–13.1)		4,294	(1,305–7,282)
ME	5-9	390	4.1	1.32	(2.2–7.6)		2,574	(916–4,232)
ME	10-14	594	10.8	1.86	(7.7–15.0)		7,002	(4,519–9,484)
ME	15-17	364	7.7	2.00	(4.6–12.7)		3,288	(1,553–5,022)
MD	0-4	185	3.5	1.53	(1.4–8.1)		12,197	(1,503–22,891)
MD	5-9	249	9.4	2.25	(5.8–14.8)		34,199	(17,534–50,864)
MD	10-14	360	7.5	1.90	(4.5–12.2)		23,309	(11,284–35,334)
MD	15-17	257	11.9	3.93	(6.1–22.0)		22,270	(6,554–37,987)
MA	0-4	114	3.1				10,490	
MA	5-9	161	9.5	2.37	(5.8–15.3)		29,935	(15,172–44,699)
MA	10-14	163	5.6	1.76	(3.0–10.2)		19,792	(7,659–31,925)
MA	15-17	152	8.4	2.39	(4.8–14.4)		19,350	(8,407–30,292)
MI	0-4	487	4.2	1.16	(2.4–7.2)		23,294	(10,407–36,182)
MI	5-9	517	8.6	1.56	(6.0–12.2)		49,244	(31,159–67,328)
MI	10-14	818	11.5	1.47	(8.9–14.7)		63,502	(46,704–80,299)
MI	15-17	549	10.7	1.69	(7.8–14.4)		40,947	(27,656–54,238)
MN	0-4	756	2.8	0.72	(1.7–4.6)		10,025	(4,967–15,084)
MN	5-9	800	4.0	0.87	(2.6–6.1)		14,356	(8,140–20,573)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table C3
Child Current Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2018**

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
MN	10-14	1,654	6.8	0.93	(5.2–8.9)		22,717	(16,414–29,020)
MN	15-17	841	9.0	1.33	(6.7–12.0)		19,603	(13,682–25,523)
MS	0-4	320	2.3	0.80	(1.2–4.5)		4,062	(1,322–6,803)
MS	5-9	304	8.9	1.73	(6.0–12.9)		16,007	(9,770–22,245)
MS	10-14	479	6.5	1.05	(4.7–8.9)		12,341	(8,468–16,214)
MS	15-17	310	9.1	2.21	(5.6–14.5)		10,809	(5,400–16,219)
MO	0-4	182	7.8	2.50	(4.1–14.3)		28,878	(9,957–47,798)
MO	5-9	192	7.9	2.63	(4.1–14.8)		29,450	(9,512–49,389)
MO	10-14	344	8.9	1.90	(5.8–13.4)		29,292	(16,665–41,919)
MO	15-17	272	9.6	2.41	(5.8–15.5)		20,626	(10,019–31,232)
MT	0-4	296	1.8	0.74	(0.8–4.0)		1,085	(209–1,961)
MT	5-9	239	5.4	1.76	(2.8–10.1)		3,292	(1,148–5,436)
MT	10-14	363	6.1	1.58	(3.6–10.0)		3,660	(1,790–5,530)
MT	15-17	253	9.6	2.73	(5.4–16.4)		3,699	(1,575–5,823)
NE	0-4	955	2.5	0.84	(1.3–4.8)		3,273	(1,092–5,455)
NE	5-9	791	5.4	1.35	(3.3–8.8)		7,052	(3,499–10,605)
NE	10-14	965	5.1	0.89	(3.7–7.2)		6,670	(4,383–8,958)
NE	15-17	726	8.5	1.51	(5.9–11.9)		6,555	(4,171–8,939)
NV	0-4	149	0.6				1,091	
NV	5-9	141	11.3	4.06	(5.5–22.0)		21,775	(5,543–38,008)
NV	10-14	298	8.1	2.13	(4.8–13.4)		16,330	(7,757–24,903)
NV	15-17	143	8.9	3.16	(4.3–17.3)		6,720	(1,886–11,553)
NH	0-4	161	3.7	1.46	(1.7–7.9)		2,238	(479–3,996)
NH	5-9	247	8.2	2.34	(4.6–14.1)		5,138	(2,117–8,160)
NH	10-14	357	11.8	2.14	(8.2–16.7)		7,260	(4,561–9,960)
NH	15-17	212	11.2	2.31	(7.4–16.6)		4,843	(2,827–6,858)
NJ	0-4	81	3.1				15,124	
NJ	5-9	80	8.5	4.20	(3.2–21.1)		44,050	(505–87,594)
NJ	10-14	140	8.8	2.56	(4.9–15.3)		47,913	(21,867–73,959)
NJ	15-17	82	9.0	3.97	(3.7–20.4)		26,054	(2,718–49,391)
NM	0-4	331	2.4	1.07	(1.0–5.7)		3,056	(343–5,768)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table C3
Child Current Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2018**

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * Weighted Number
NM	5-9	366	10.7	2.38	(6.9–16.3)		13,861	(7,527–20,194)
NM	10-14	558	7.7	2.07	(4.5–12.9)		10,030	(4,500–15,559)
NM	15-17	315	10.2	2.31	(6.5–15.7)		8,231	(4,471–11,991)
NY	0-4	585	1.9	0.61	(1.0–3.5)		15,967	(5,986–25,948)
NY	5-9	585	6.6	1.66	(4.0–10.7)		61,401	(30,114–92,689)
NY	10-14	757	6.9	1.45	(4.5–10.3)		65,266	(37,679–92,853)
NY	15-17	489	10.1	2.15	(6.6–15.2)		65,232	(37,281–93,184)
OH	0-4	646	5.2	1.27	(3.2–8.3)		34,193	(17,510–50,875)
OH	5-9	528	8.2	1.75	(5.4–12.4)		54,818	(31,116–78,520)
OH	10-14	757	9.0	1.51	(6.5–12.5)		61,609	(40,862–82,356)
OH	15-17	581	8.9	1.59	(6.3–12.6)		38,804	(25,058–52,551)
OR	0-4	277	3.4	1.25	(1.6–6.9)		7,681	(2,097–13,265)
OR	5-9	304	4.8	1.29	(2.9–8.1)		10,851	(5,161–16,540)
OR	10-14	430	8.8	1.55	(6.2–12.3)		19,049	(12,407–25,692)
OR	15-17	224	14.9	4.74	(7.8–26.7)		21,123	(6,332–35,914)
PA	0-4	261	3.3	1.30	(1.5–7.1)		22,535	(5,006–40,063)
PA	5-9	315	11.7	3.36	(6.5–20.0)		76,948	(29,370–124,527)
PA	10-14	401	13.0	2.51	(8.8–18.8)		89,938	(53,597–126,278)
PA	15-17	297	11.6	2.94	(6.9–18.7)		52,762	(24,380–81,144)
RI	0-4	290	7.8	2.20	(4.4–13.3)		3,380	(1,439–5,321)
RI	5-9	216	8.1	3.21	(3.6–17.0)		4,451	(782–8,119)
RI	10-14	293	6.9	1.70	(4.2–11.1)		3,524	(1,801–5,247)
RI	15-17	233	8.9	2.61	(4.9–15.5)		3,326	(1,347–5,306)
UT	0-4	944	3.5	0.68	(2.4–5.1)		8,751	(5,346–12,155)
UT	5-9	890	5.9	0.91	(4.4–8.0)		14,882	(10,276–19,488)
UT	10-14	1,119	6.3	0.94	(4.7–8.4)		16,042	(11,216–20,867)
UT	15-17	736	6.8	1.16	(4.8–9.4)		9,978	(6,517–13,440)
VT	0-4	228	4.4	1.38	(2.4–8.0)		1,260	(493–2,027)
VT	5-9	329	12.6	2.74	(8.1–19.0)		3,459	(1,878–5,039)
VT	10-14	387	8.7	1.97	(5.5–13.4)		2,097	(1,141–3,052)
VT	15-17	273	10.0	2.84	(5.6–17.1)		1,955	(801–3,109)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table C3
Child Current Asthma Prevalence
and Weighted Number by Age and State or Territory: BRFSS 2018

State	Age Group	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)		Weighted Number	95% CI * Weighted Number
WI	0-4	266	3.8	2.55	(1.0–13.5)		11,848	(0–27,998)
WI	5-9	235	5.6	1.76	(3.0–10.2)		18,502	(6,903–30,101)
WI	10-14	260	14.1	2.64	(9.7–20.1)		50,140	(30,799–69,480)
WI	15-17	216	10.3	3.05	(5.7–18.0)		22,540	(9,079–36,001)
Territory								
PR	0-4	270	13.3	2.60	(9.0–19.3)		22,142	(12,946–31,338)
PR	5-9	282	17.9	2.44	(13.6–23.2)		34,615	(24,618–44,611)
PR	10-14	394	12.7	1.82	(9.6–16.7)		24,502	(17,352–31,651)
PR	15-17	248	11.8	2.56	(7.6–17.8)		16,506	(8,998–24,015)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.