

Table L1
Child Lifetime Asthma Prevalence
and Weighted Number by State or Territory: BRFSS 2018

State	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * (Number)
Total **	54,038	11.4	0.27	(10.9–11.9)		5,177,190	(4,936,486–5,417,894)
CA	2,869	10.7	0.72	(9.3–12.2)		940,371	(812,879–1,067,863)
CT	1,989	14.9	1.04	(13.0–17.1)		96,592	(82,959–110,225)
DC	714	16.4	1.69	(13.4–20.0)		19,187	(15,197–23,176)
FL	2,857	12.5	1.21	(10.3–15.1)		509,688	(410,653–608,723)
GA	2,060	12.2	0.88	(10.6–14.1)		288,252	(247,268–329,237)
HI	1,836	17.7	1.15	(15.5–20.1)		52,154	(45,224–59,084)
IL	1,252	9.3	0.96	(7.6–11.4)		254,492	(202,890–306,093)
IN	1,396	12.9	1.11	(10.9–15.2)		171,674	(141,875–201,473)
KS	2,350	11.1	0.78	(9.6–12.7)		75,061	(64,701–85,421)
KY	1,864	9.4	0.95	(7.7–11.5)		91,534	(73,819–109,249)
ME	1,694	11.8	1.14	(9.8–14.3)		27,656	(22,281–33,031)
MD	1,056	13.3	1.47	(10.6–16.4)		162,080	(125,789–198,370)
MA	593	13.1	1.64	(10.2–16.6)		163,365	(123,279–203,451)
MI	2,381	12.5	0.85	(10.9–14.2)		258,346	(223,682–293,009)
MN	4,059	7.4	0.52	(6.4–8.5)		93,544	(80,601–106,487)
MS	1,415	10.0	0.90	(8.4–11.9)		66,772	(55,015–78,528)
MO	993	13.5	1.54	(10.8–16.8)		174,603	(133,777–215,429)
MT	1,154	7.1	1.01	(5.4–9.4)		15,742	(11,361–20,123)
NE	3,442	7.8	0.68	(6.5–9.2)		36,324	(30,038–42,610)
NV	732	13.3	2.07	(9.7–17.9)		87,680	(60,009–115,352)
NH	980	11.2	1.18	(9.1–13.8)		25,808	(20,478–31,138)
NJ	386	11.3	2.10	(7.8–16.1)		210,811	(136,654–284,968)
NM	1,573	11.6	1.20	(9.5–14.2)		54,449	(43,139–65,758)
NY	2,425	9.9	0.94	(8.2–11.9)		333,053	(270,359–395,748)
OH	2,516	12.2	0.99	(10.4–14.2)		297,284	(249,020–345,548)
OR	1,236	11.6	1.43	(9.0–14.7)		94,170	(70,349–117,992)
PA	1,276	12.9	1.44	(10.4–16.0)		322,166	(249,899–394,433)
RI	1,037	14.3	1.80	(11.1–18.2)		26,772	(19,755–33,789)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L1
Child Lifetime Asthma Prevalence
and Weighted Number by State or Territory: BRFSS 2018

State	Unweighted Number	Prevalence (Percent)	Standard Error	95% CI * (Percent)	 	Weighted Number	95% CI * (Number)
UT	3,699	8.2	0.54	(7.2–9.3)		74,135	(64,432–83,838)
VT	1,225	13.0	1.34	(10.6–15.9)		13,022	(10,359–15,686)
WI	979	11.5	1.42	(9.0–14.6)		140,405	(106,073–174,736)
Territory							
PR	1,196	24.8	1.43	(22.1–27.7)		172,131	(151,742–192,521)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.