

**Table L2_1
Child Lifetime Asthma Prevalence
by Sex and State or Territory: BRFSS 2018**

State	Male Unweighted Number	Male Prevalence	Male Standard Error	Male 95% CI *	Female Unweighted Number	Female Prevalence	Female Standard Error	Female 95% CI *
Total **	27,429	13.2	0.41	(12.5–14.1)	25,782	9.5	0.33	(8.9–10.2)
CA	1,452	12.5	1.09	(10.5–14.8)	1,390	8.3	0.90	(6.7–10.3)
CT	971	15.7	1.43	(13.1–18.7)	969	13.7	1.55	(11.0–17.1)
DC	350	15.3	2.36	(11.2–20.5)	353	17.2	2.42	(13.0–22.5)
FL	1,417	16.5	2.01	(12.9–20.9)	1,388	8.7	1.38	(6.3–11.8)
GA	1,076	14.6	1.36	(12.1–17.4)	948	10.1	1.12	(8.1–12.5)
HI	942	19.6	1.64	(16.6–23.0)	868	16.1	1.65	(13.1–19.6)
IL	641	9.0	1.35	(6.7–12.1)	592	9.5	1.38	(7.1–12.6)
IN	695	14.0	1.64	(11.1–17.6)	685	11.5	1.47	(8.9–14.7)
KS	1,177	13.0	1.21	(10.8–15.6)	1,135	9.3	1.00	(7.6–11.5)
KY	961	10.8	1.43	(8.3–13.9)	890	8.2	1.27	(6.0–11.1)
ME	852	16.0	1.89	(12.6–20.0)	812	7.5	1.22	(5.4–10.3)
MD	532	13.7	2.21	(9.9–18.7)	509	12.8	1.97	(9.4–17.2)
MA	292	14.0	2.37	(10.0–19.3)	281	12.6	2.39	(8.6–18.1)
MI	1,202	15.7	1.34	(13.3–18.5)	1,149	9.2	1.03	(7.4–11.5)
MN	2,132	8.9	0.82	(7.5–10.7)	1,850	5.8	0.65	(4.7–7.2)
MS	704	12.4	1.43	(9.9–15.5)	702	7.5	1.06	(5.6–9.8)
MO	513	18.1	2.61	(13.6–23.8)	467	8.8	1.62	(6.1–12.6)
MT	601	6.1	0.99	(4.4–8.3)	534	7.6	1.69	(4.9–11.7)
NE	1,788	8.6	0.99	(6.8–10.7)	1,615	6.9	0.93	(5.2–8.9)
NV	377	13.5	2.64	(9.1–19.5)	346	13.3	3.21	(8.1–20.9)
NH	487	12.9	1.79	(9.7–16.8)	476	9.4	1.53	(6.8–12.9)
NJ	203	13.2	3.25	(8.0–21.0)	178	9.7	2.35	(6.0–15.4)
NM	810	13.5	1.83	(10.3–17.5)	745	9.9	1.58	(7.2–13.4)
NY	1,245	10.3	1.30	(8.1–13.2)	1,142	9.4	1.37	(7.0–12.4)
OH	1,287	14.6	1.52	(11.9–17.9)	1,191	9.6	1.28	(7.4–12.4)
OR	645	9.8	1.35	(7.4–12.8)	566	13.3	2.52	(9.1–19.0)
PA	636	15.2	2.39	(11.1–20.5)	621	11.0	1.49	(8.4–14.2)
RI	530	17.6	2.76	(12.9–23.7)	499	11.0	2.27	(7.3–16.3)
UT	1,792	9.8	0.85	(8.3–11.6)	1,829	6.6	0.69	(5.4–8.1)
VT	610	13.5	1.85	(10.3–17.6)	593	12.7	2.00	(9.2–17.1)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

Table L2_1
Child Lifetime Asthma Prevalence
by Sex and State or Territory: BRFSS 2018

State	Male Unweighted Number	Male Prevalence	Male Standard Error	Male 95% CI *		Female Unweighted Number	Female Prevalence	Female Standard Error	Female 95% CI *
WI	509	14.4	2.25	(10.5–19.4)		459	8.7	1.70	(5.9–12.7)
Territory									
PR	607	26.7	2.04	(22.9–30.9)		582	23.1	2.00	(19.4–27.2)

Notes:

* CI denotes confidence interval.

** Total includes data from 31 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.