

**Table C2\_1**  
**Child<sup>§</sup> Current Asthma Prevalence**  
**by Sex and State or Territory: BRFSS 2019**

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI <sup>¶</sup>		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI <sup>¶</sup>
Total *	24,325	8.1	0.38	(7.4–8.9)		22,678	6.6	0.37	(5.9–7.4)
CA	1,355	8.3	0.96	(6.6–10.4)		1,291	6.2	0.82	(4.8–8.0)
CT	761	8.8	1.27	(6.6–11.6)		705	8.9	1.45	(6.4–12.2)
DC	216	12.8	2.58	(8.5–18.7)		218	10.4	3.38	(5.4–19.1)
FL	1,354	8.4	1.45	(5.9–11.7)		1,172	5.1	1.06	(3.3–7.6)
GA	690	12.2	2.34	(8.3–17.6)		634	8.1	1.75	(5.3–12.3)
HI	877	8.1	1.10	(6.1–10.5)		787	6.9	1.08	(5.1–9.4)
IL	686	8.1	1.48	(5.7–11.6)		560	5.7	1.21	(3.8–8.6)
IN	817	7.9	1.10	(6.0–10.3)		799	5.4	0.81	(4.0–7.2)
KS	1,280	10.7	1.25	(8.5–13.4)		1,218	6.8	0.92	(5.2–8.8)
ME	741	11.1	1.84	(7.9–15.2)		710	8.0	1.43	(5.6–11.3)
MA	352	10.1	2.13	(6.6–15.1)		268	9.8	2.78	(5.6–16.8)
MI	1,083	8.7	1.15	(6.7–11.2)		1,020	7.0	1.07	(5.2–9.4)
MN	1,639	5.3	0.72	(4.1–7.0)		1,633	4.8	0.64	(3.7–6.2)
MS	539	9.6	1.84	(6.6–13.9)		525	5.9	1.21	(3.9–8.8)
MO	686	9.2	1.52	(6.6–12.7)		617	4.9	1.10	(3.1–7.6)
MT	717	6.7	1.20	(4.7–9.5)		681	6.9	1.15	(4.9–9.5)
NE	1,778	7.6	0.92	(6.0–9.6)		1,686	4.9	0.70	(3.7–6.5)
NV	351	10.3	2.92	(5.9–17.6)		305	5.6	1.77	(2.9–10.2)
NH	459	8.4	1.60	(5.7–12.1)		413	5.8	1.48	(3.5–9.5)
NM	639	4.6	1.02	(2.9–7.1)		568	6.5	1.77	(3.7–10.9)
NY	386	9.3	2.17	(5.8–14.5)		405	6.8	1.90	(3.9–11.6)
OH	1,130	8.7	1.33	(6.4–11.7)		989	6.5	1.16	(4.6–9.2)
OR	587	5.9	1.50	(3.5–9.6)		587	8.3	2.54	(4.5–14.9)
PA	713	9.2	1.23	(7.0–11.9)		647	6.8	1.16	(4.9–9.5)
RI	433	9.5	2.00	(6.2–14.2)		447	8.0	1.88	(5.0–12.6)
TX	1,262	5.7	1.14	(3.8–8.4)		1,080	7.9	1.60	(5.3–11.7)
UT	1,851	6.4	0.65	(5.2–7.7)		1,846	5.2	0.61	(4.1–6.5)
VT	543	7.7	1.49	(5.2–11.1)		503	8.9	1.73	(6.0–12.9)
WI	400	4.5	1.35	(2.5–8.0)		364	4.0	1.40	(2.0–7.9)

Notes:

<sup>§</sup>Aged 0-17

<sup>¶</sup>CI denotes confidence interval.

\* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Asthma and Community Health Branch, National Center for Environmental Health  
Centers for Disease Control and Prevention**

**Table C2\_1**  
**Child<sup>§</sup> Current Asthma Prevalence**  
**by Sex and State or Territory: BRFSS 2019**

State	Male Sample Size	Male Prevalence	Male Standard Error	Male 95% CI <sup>¶</sup>		Female Sample Size	Female Prevalence	Female Standard Error	Female 95% CI <sup>¶</sup>
Territory									
PR	716	14.8	1.48	(12.1–17.9)		689	15.4	1.58	(12.5–18.7)

Notes:

<sup>§</sup>Aged 0-17

<sup>¶</sup>CI denotes confidence interval.

\* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Asthma and Community Health Branch, National Center for Environmental Health  
Centers for Disease Control and Prevention**