

**Table C3**  
**Child<sup>§</sup> Current Asthma Prevalence**  
**and Weighted Number by Age and State or Territory: BRFSS 2019**

State	Age Group	Sample Size	Prevalence (Percent)	Standard Error	95% CI <sup>¶</sup> (Percent)	    	Weighted Number	95% CI <sup>¶</sup> Weighted Number
Total *	0-4	9,568	3.6	0.36	(2.9–4.4)		425,108	(339,030–511,186)
Total *	5-9	9,935	8.6	0.64	(7.4–10.0)		1,097,053	(928,388–1,265,719)
Total *	10-14	12,473	9.6	0.61	(8.5–10.9)		1,055,153	(916,116–1,194,191)
Total *	15-17	10,044	9.0	0.61	(7.9–10.3)		728,306	(628,479–828,133)
CA	0-4	607	3.9	1.00	(2.4–6.4)		77,015	(37,483–116,546)
CA	5-9	591	8.2	1.44	(5.8–11.5)		189,202	(121,049–257,356)
CA	10-14	675	9.7	1.40	(7.3–12.8)		197,251	(138,986–255,516)
CA	15-17	506	8.7	1.50	(6.2–12.1)		133,108	(86,718–179,499)
CT	0-4	242	3.8	1.31	(2.0–7.4)		6,814	(2,219–11,410)
CT	5-9	284	11.1	2.33	(7.3–16.6)		18,396	(10,433–26,359)
CT	10-14	411	9.5	1.77	(6.6–13.6)		14,732	(9,141–20,323)
CT	15-17	369	10.5	2.50	(6.5–16.5)		12,941	(6,493–19,388)
DC	0-4	84	8.9	4.79	(3.0–23.7)		3,742	(0–7,913)
DC	5-9	106	12.7	3.35	(7.4–20.8)		5,015	(2,324–7,705)
DC	10-14	125	13.6	3.35	(8.2–21.6)		3,154	(1,547–4,762)
DC	15-17	73	13.0	4.91	(6.0–25.9)		1,499	(301–2,697)
FL	0-4	492	2.3	0.75	(1.2–4.3)		26,706	(10,100–43,312)
FL	5-9	580	6.1	1.63	(3.6–10.2)		57,431	(27,326–87,536)
FL	10-14	686	10.0	2.41	(6.2–15.8)		73,543	(37,096–109,991)
FL	15-17	431	10.8	2.55	(6.7–16.9)		77,657	(40,697–114,617)
GA	0-4	268	9.7	3.38	(4.8–18.6)		60,582	(16,439–104,726)
GA	5-9	296	12.0	3.49	(6.7–20.7)		68,684	(26,107–111,260)
GA	10-14	317	9.5	2.08	(6.1–14.4)		51,004	(28,333–73,676)
GA	15-17	288	10.0	3.73	(4.7–20.0)		35,222	(7,731–62,713)
HI	0-4	371	3.9	1.31	(2.0–7.5)		2,321	(772–3,870)
HI	5-9	361	8.7	1.79	(5.8–12.9)		6,426	(3,740–9,112)
HI	10-14	397	7.4	1.49	(4.9–10.9)		5,094	(3,017–7,172)
HI	15-17	319	12.1	2.20	(8.4–17.1)		5,491	(3,431–7,551)
IL	0-4	248	5.0	1.75	(2.5–9.8)		38,079	(11,349–64,810)
IL	5-9	247	7.3	2.21	(4.0–13.0)		55,963	(21,321–90,605)
IL	10-14	350	8.2	1.66	(5.5–12.1)		47,941	(28,370–67,512)

Notes:

<sup>§</sup>Aged 0-17

<sup>¶</sup>CI denotes confidence interval.

\* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Table C3**  
**Child<sup>§</sup> Current Asthma Prevalence**  
**and Weighted Number by Age and State or Territory: BRFSS 2019**

State	Age Group	Sample Size	Prevalence (Percent)	Standard Error	95% CI <sup>¶</sup> (Percent)	    	Weighted Number	95% CI <sup>¶</sup> Weighted Number
IL	15-17	289	8.4	2.31	(4.8–14.2)		34,699	(15,068–54,329)
IN	0-4	299	2.7	1.00	(1.3–5.5)		10,126	(2,753–17,499)
IN	5-9	355	7.0	1.55	(4.5–10.7)		25,577	(14,054–37,099)
IN	10-14	429	8.9	1.55	(6.3–12.4)		28,227	(18,166–38,289)
IN	15-17	361	10.6	2.00	(7.3–15.2)		25,305	(15,399–35,211)
KS	0-4	528	4.1	0.98	(2.6–6.5)		7,452	(3,938–10,965)
KS	5-9	554	7.2	1.31	(5.0–10.2)		13,101	(8,282–17,921)
KS	10-14	667	13.0	1.77	(9.9–16.9)		19,615	(13,994–25,237)
KS	15-17	540	11.7	2.66	(7.4–18.0)		13,271	(6,811–19,732)
ME	0-4	296	2.9	1.46	(1.1–7.6)		1,453	(5–2,901)
ME	5-9	273	14.4	3.04	(9.4–21.4)		8,649	(4,780–12,519)
ME	10-14	435	13.0	2.29	(9.1–18.2)		6,862	(4,352–9,372)
ME	15-17	319	7.9	2.03	(4.7–12.9)		3,189	(1,538–4,839)
MA	0-4	118	5.2	3.52	(1.3–18.2)		16,183	(0–38,630)
MA	5-9	146	12.2	3.83	(6.5–21.9)		28,253	(9,884–46,621)
MA	10-14	148	11.5	3.24	(6.5–19.5)		32,830	(13,480–52,179)
MA	15-17	140	12.6	4.85	(5.7–25.5)		30,963	(5,127–56,800)
MI	0-4	429	3.3	0.90	(1.9–5.6)		16,345	(7,445–25,245)
MI	5-9	430	8.3	1.92	(5.2–12.9)		44,394	(23,556–65,232)
MI	10-14	552	12.1	1.98	(8.7–16.6)		61,746	(40,523–82,969)
MI	15-17	533	9.2	1.50	(6.7–12.6)		31,103	(20,940–41,267)
MN	0-4	674	1.1	0.38	(0.5–2.2)		3,853	(1,214–6,492)
MN	5-9	690	5.4	1.16	(3.5–8.2)		18,831	(10,602–27,060)
MN	10-14	856	7.3	1.01	(5.5–9.5)		16,574	(11,985–21,162)
MN	15-17	726	8.7	1.40	(6.3–11.9)		18,833	(12,658–25,007)
MS	0-4	240	1.9				2,473	
MS	5-9	229	11.8	2.97	(7.1–19.0)		20,889	(9,891–31,887)
MS	10-14	282	8.9	2.22	(5.4–14.3)		14,113	(6,928–21,298)
MS	15-17	181	7.8	2.48	(4.1–14.3)		8,507	(3,066–13,947)
MO	0-4	258	3.1	1.19	(1.4–6.5)		10,926	(2,651–19,202)
MO	5-9	271	7.5	2.03	(4.4–12.6)		24,157	(10,870–37,445)

Notes:

<sup>§</sup>Aged 0-17

<sup>¶</sup>CI denotes confidence interval.

\* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Asthma and Community Health Branch, National Center for Environmental Health  
Centers for Disease Control and Prevention**

**Table C3**  
**Child<sup>§</sup> Current Asthma Prevalence**  
**and Weighted Number by Age and State or Territory: BRFSS 2019**

State	Age Group	Sample Size	Prevalence (Percent)	Standard Error	95% CI <sup>¶</sup> (Percent)	    	Weighted Number	95% CI <sup>¶</sup> Weighted Number
MO	10-14	352	9.6	2.39	(5.8–15.4)		24,848	(11,982–37,714)
MO	15-17	272	8.2	2.14	(4.9–13.5)		18,437	(8,728–28,146)
MT	0-4	302	1.6				921	
MT	5-9	310	6.8	1.61	(4.3–10.7)		4,124	(2,164–6,083)
MT	10-14	367	11.0	2.21	(7.3–16.1)		5,370	(3,124–7,616)
MT	15-17	286	8.8	2.27	(5.3–14.4)		3,235	(1,515–4,955)
NE	0-4	741	3.4	0.92	(2.0–5.7)		3,782	(1,760–5,805)
NE	5-9	746	6.8	1.35	(4.5–9.9)		8,562	(5,077–12,048)
NE	10-14	941	7.9	1.23	(5.8–10.7)		9,122	(6,252–11,992)
NE	15-17	759	9.3	1.41	(6.8–12.4)		6,683	(4,609–8,757)
NV	0-4	125	3.5				6,676	
NV	5-9	133	14.2	4.56	(7.4–25.6)		26,491	(8,265–44,717)
NV	10-14	178	10.8	3.95	(5.1–21.3)		11,906	(2,700–21,112)
NV	15-17	135	2.3				1,950	
NH	0-4	156	1.8				1,139	
NH	5-9	191	7.5	2.77	(3.6–15.1)		4,637	(1,143–8,131)
NH	10-14	245	13.7	2.74	(9.1–20.0)		8,349	(4,879–11,818)
NH	15-17	217	7.2	1.87	(4.2–11.8)		3,107	(1,475–4,739)
NM	0-4	228	1.3				1,503	
NM	5-9	251	5.5	2.49	(2.3–13.0)		7,086	(615–13,558)
NM	10-14	327	6.4	2.02	(3.4–11.7)		5,718	(2,060–9,375)
NM	15-17	238	7.9	2.78	(3.9–15.3)		5,937	(1,638–10,236)
NY	0-4	144	5.1	2.31	(2.1–12.1)		38,911	(3,777–74,046)
NY	5-9	165	13.6	4.13	(7.3–23.8)		140,384	(48,604–232,163)
NY	10-14	206	7.1	2.26	(3.7–13.0)		64,283	(22,591–105,976)
NY	15-17	167	7.8	3.06	(3.5–16.3)		47,491	(9,408–85,573)
OH	0-4	418	4.0	1.70	(1.7–9.1)		20,887	(3,125–38,649)
OH	5-9	435	10.7	2.13	(7.1–15.6)		72,228	(42,518–101,937)
OH	10-14	580	9.3	1.82	(6.3–13.5)		54,176	(32,608–75,744)
OH	15-17	478	7.7	1.88	(4.7–12.3)		31,921	(15,971–47,872)
OR	0-4	250	3.7	2.07	(1.2–10.7)		6,049	(0–12,847)

Notes:

<sup>§</sup>Aged 0-17

<sup>¶</sup>CI denotes confidence interval.

\* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Asthma and Community Health Branch, National Center for Environmental Health  
Centers for Disease Control and Prevention**

**Table C3**  
**Child<sup>§</sup> Current Asthma Prevalence**  
**and Weighted Number by Age and State or Territory: BRFSS 2019**

State	Age Group	Sample Size	Prevalence (Percent)	Standard Error	95% CI <sup>¶</sup> (Percent)	    	Weighted Number	95% CI <sup>¶</sup> Weighted Number
OR	5-9	242	6.6	3.20	(2.5–16.3)		14,621	(0–29,346)
OR	10-14	301	11.2	4.12	(5.3–22.1)		19,532	(4,099–34,966)
OR	15-17	251	7.5	2.81	(3.5–15.2)		9,663	(2,202–17,124)
PA	0-4	285	3.3	1.03	(1.7–6.0)		19,566	(7,394–31,737)
PA	5-9	305	7.3	1.61	(4.7–11.1)		48,662	(27,554–69,770)
PA	10-14	340	9.6	1.81	(6.6–13.8)		57,303	(35,533–79,073)
PA	15-17	294	10.7	2.07	(7.3–15.5)		42,373	(25,848–58,897)
RI	0-4	164	7.4	3.04	(3.2–16.0)		3,985	(621–7,350)
RI	5-9	178	9.8	3.06	(5.2–17.6)		5,218	(1,862–8,574)
RI	10-14	235	10.8	2.84	(6.4–17.8)		3,944	(1,802–6,087)
RI	15-17	203	9.5	2.44	(5.7–15.5)		2,995	(1,451–4,538)
TX	0-4	481	1.3	0.49	(0.7–2.7)		21,885	(6,471–37,299)
TX	5-9	479	7.9	2.36	(4.3–13.9)		144,049	(55,532–232,565)
TX	10-14	588	11.4	2.75	(7.0–18.0)		190,621	(94,275–286,967)
TX	15-17	491	8.6	1.99	(5.4–13.4)		97,481	(52,194–142,768)
UT	0-4	780	2.7	0.71	(1.6–4.5)		6,395	(3,103–9,687)
UT	5-9	692	5.8	0.96	(4.2–8.0)		14,793	(9,914–19,673)
UT	10-14	971	7.5	0.95	(5.9–9.6)		12,834	(9,566–16,102)
UT	15-17	773	9.2	1.26	(7.0–12.0)		13,809	(9,937–17,680)
VT	0-4	203	5.2	2.07	(2.3–11.1)		1,548	(308–2,788)
VT	5-9	241	9.4	2.44	(5.6–15.4)		2,318	(1,101–3,535)
VT	10-14	309	11.1	2.23	(7.4–16.2)		2,963	(1,749–4,178)
VT	15-17	218	9.9	3.08	(5.3–17.8)		2,114	(751–3,478)
WI	0-4	137	2.4				7,788	
WI	5-9	154	5.9	2.48	(2.6–13.1)		18,912	(2,918–34,907)
WI	10-14	203	4.3	2.18	(1.6–11.2)		11,496	(0–23,081)
WI	15-17	187	5.5	2.09	(2.6–11.3)		9,322	(2,300–16,344)
Territory								
PR	0-4	295	12.6	2.15	(9.0–17.4)		21,558	(13,921–29,196)
PR	5-9	333	19.2	2.39	(15.0–24.3)		36,504	(26,820–46,188)

Notes:

<sup>§</sup>Aged 0-17

<sup>¶</sup>CI denotes confidence interval.

\* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Asthma and Community Health Branch, National Center for Environmental Health  
Centers for Disease Control and Prevention**

**Table C3**  
**Child<sup>§</sup> Current Asthma Prevalence**  
**and Weighted Number by Age and State or Territory: BRFSS 2019**

State	Age Group	Sample Size	Prevalence (Percent)	Standard Error	95% CI <sup>¶</sup> (Percent)	    	Weighted Number	95% CI <sup>¶</sup> Weighted Number
PR	10-14	367	14.8	2.01	(11.3–19.2)		20,801	(14,949–26,654)
PR	15-17	293	14.2	2.31	(10.2–19.3)		20,041	(13,170–26,913)

Notes:

<sup>§</sup>Aged 0-17

<sup>¶</sup>CI denotes confidence interval.

\* Total includes data from 29 states and DC, but excludes Puerto Rico.

If values for the standard error and the 95% confidence interval are not provided, the normal approximation to the binomial distribution does not apply due to small sample size.

When the sample size is less than 50, estimates are not precise and should be interpreted with caution.

**Asthma and Community Health Branch, National Center for Environmental Health  
Centers for Disease Control and Prevention**