

**Table L1
Child^a Lifetime Asthma^b Prevalence
and Weighted Number^c by State or Territory: BRFSS 2020**

State	Sample Size ^d	Prevalence (Percent)	Standard Error	95% CI ^e (Percent)	 	Weighted Number ^f	95% CI ^e (Number)
Total	47,356	11.3	0.38	(10.6–12.0)		5,420,952	(5,059,448–5,782,457)
CA	909	10.0	1.27	(7.7–12.7)		875,328	(653,509–1,097,147)
CT	1,855	18.0	1.69	(15.0–21.6)		124,416	(100,497–148,335)
DC	545	12.8	1.75	(9.7–16.6)		16,896	(12,232–21,561)
FL	1,726	10.1	1.34	(7.8–13.0)		408,997	(302,373–515,620)
GA	1,729	16.6	1.65	(13.6–20.1)		394,259	(313,986–474,532)
HI	1,572	13.6	1.19	(11.4–16.1)		39,424	(32,458–46,391)
IN	1,718	8.6	0.77	(7.2–10.2)		127,080	(104,439–149,721)
KS	2,342	10.7	0.79	(9.2–12.3)		71,330	(60,827–81,832)
KY	790	9.9	1.38	(7.5–13.0)		93,548	(68,092–119,005)
ME	1,520	12.2	1.24	(10.0–14.9)		27,383	(21,910–32,855)
MA	595	12.9	1.92	(9.6–17.2)		152,875	(106,259–199,492)
MI	1,457	12.4	1.08	(10.4–14.7)		251,379	(207,878–294,880)
MN	3,668	6.6	0.54	(5.7–7.8)		83,399	(70,124–96,674)
MS	1,357	13.9	1.40	(11.4–16.9)		91,764	(73,191–110,338)
MO	1,833	12.2	1.03	(10.4–14.4)		157,813	(130,939–184,686)
MT	1,388	9.4	0.99	(7.6–11.5)		20,695	(16,390–24,999)
NE	3,377	7.3	0.67	(6.1–8.8)		34,246	(28,112–40,380)
NV	448	8.6	1.78	(5.7–12.7)		56,073	(33,382–78,764)
NH	938	12.2	1.33	(9.8–15.0)		29,405	(23,047–35,763)
NJ	2,203	12.0	0.95	(10.3–14.0)		212,802	(179,112–246,492)
NM	1,475	10.7	1.32	(8.4–13.6)		48,389	(36,618–60,160)
NY	978	12.4	1.35	(10.0–15.3)		469,100	(368,807–569,393)
OH	2,864	10.2	0.89	(8.5–12.1)		243,263	(200,838–285,688)
PA	1,125	14.2	1.68	(11.2–17.8)		350,884	(265,704–436,063)
RI	1,004	15.0	1.64	(12.1–18.5)		29,233	(22,859–35,607)
TX	2,489	11.1	1.29	(8.8–13.9)		796,125	(612,020–980,230)
UT	3,292	8.7	0.62	(7.6–10.0)		80,562	(69,309–91,816)
VT	1,222	11.3	1.16	(9.3–13.8)		12,266	(9,821–14,710)
WI	937	10.0	1.55	(7.4–13.5)		122,018	(84,462–159,575)

Notes: For additional information on methods and variable description, refer to “Technical Information, available at: <https://www.cdc.gov/asthma/brfss/2020/child/brfsschildtechninfo.html>

^aAged 0-17

^bLifetime question: Has a doctor, nurse or other health professional ever said that the child has asthma?

^cWeighted number of respondents who answered the lifetime asthma question

^dUnweighted number of respondents who answered the lifetime asthma question

^eCI denotes confidence interval

^fTotal includes data from 30 states and DC, but excludes Puerto Rico

‘.-/blank’: Suppressed if the normal approximation to the binomial distribution does not apply due to small sample size or relative standard error (RSE)>30%

Table L1
Child^a Lifetime Asthma^b Prevalence
and Weighted Number^c by State or Territory: BRFSS 2020

State	Sample Size ^d	Prevalence (Percent)	Standard Error	95% CI ^e (Percent)	 	Weighted Number ^f	95% CI ^e (Number)
Territory							
PR	1,169	13.8	1.12	(11.7–16.1)		94,554	(79,214–109,894)

Notes: For additional information on methods and variable description, refer to “Technical Information, available at: <https://www.cdc.gov/asthma/brfss/2020/child/brfsschildtechinfo.html>

^aAged 0-17

^bLifetime question: Has a doctor, nurse or other health professional ever said that the child has asthma?

^cWeighted number of respondents who answered the lifetime asthma question

^dUnweighted number of respondents who answered the lifetime asthma question

^eCI denotes confidence interval

^fTotal includes data from 30 states and DC, but excludes Puerto Rico

‘.-/blank’: Suppressed if the normal approximation to the binomial distribution does not apply due to small sample size or relative standard error (RSE) > 30%