

# SLATE II (Simplified Algorithm for Treatment Eligibility)

Evidence-Based for Engagement to HIV Care

Evidence-Based for Viral Suppression

Evidence-Based Structural Intervention

## INTERVENTION DESCRIPTION

### Target Population

- People with HIV (PWH) who are presenting for antiretroviral treatment (ART) initiation including those who are re-initiating treatment, with mild TB symptoms

### Goal of Intervention

- Improve ART initiation and retention

### Brief Description

*SLATE II (Simplified Algorithm for Treatment Eligibility)* is a revised algorithm of [SLATE I](#), an evidence-based intervention, that focuses on a simple, accelerated algorithm for ART initiation that requires no point-of-care laboratory testing and may be administered entirely by primary care clinicians, including nurses with prescribing authority. The SLATE I algorithm consists of four screening tools: (1) symptom report, (2) medical history, (3) brief physical examination, and (4) patient readiness assessment, each as an evaluation area for eligibility requirements for same-day ART initiation. SLATE II uses the 4 screening tools and allows same-day ART initiation for patients with mild tuberculosis (TB) symptoms and other less serious reasons for delay. Participants determined to be eligible based on all four screening tools were offered ART initiation on the same day as study enrollment. Participants found to be ineligible on any of the screening tools were referred to routine care at the study site clinic for further management.

### Theoretical Basis

- None reported

### Intervention Duration

- Ongoing

### Intervention Settings

- High volume, public-sector primary health care clinics in South Africa

### Deliverer

- Primary care clinicians, primarily nurse practitioners able to dispense ART

### Delivery Methods

- Interviewing
- Mobile tablet

- Physical examination

### Structural Components

- Access— HIV health care
  - Increased access to HIV testing and referral to HIV medical care
- Capacity building— Provider/supervisor training
  - Trained staff on interviewing and data capture of eligible participants and on use of SLATE II algorithm

- Physical Structure – Integration of services
  - Integration of services for TB treatment
- Policy/Procedure—Institutional policy/procedure
  - Implemented same-day ART initiation for eligible patients who tested HIV positive

## INTERVENTION PACKAGE INFORMATION

**Contact study author for training and intervention materials.** Please contact **Mhairi Maskew**, Health Economics and Epidemiology Research Office, University of the Witwatersrand, School of Clinical Medicine, 1 Jan Smuts Avenue, Braamfontein, Johannesburg, 2000, South Africa.

**Email:** [mmaskew@heroza.org](mailto:mmaskew@heroza.org) for details on intervention materials.

## EVALUATION STUDY AND RESULTS

### Study Location Information

The original evaluation study was conducted in Johannesburg, South Africa between March 2018 through September 2018.

### Key Intervention Effects

- Increased ART initiation
- Increased engagement in HIV care
- Increased viral suppression

### Recruitment Settings

- High volume, public sector, primary care clinics (n=3) based in both urban formal and informal settings

### Study Sample

The baseline sample (N = 593) is characterized by the following:

#### Intervention arm (n = 296)

- 64% female persons, 36% male persons
- Median age of 35 years
- 90% living in peri-urban areas, 10% living in urban areas
- 58% single persons, 32% married persons or persons living with long-term partner, 10% divorced/ separated/widowed persons
- 47% persons presented with TB symptoms
- Median CD4 count of 298 cells/mm<sup>3</sup> 25<sup>th</sup>-75<sup>th</sup> percentile: 137-483 cells/mm<sup>3</sup>

#### Standard arm (n = 297)

- 62% female persons, 38% male persons
- Median age of 35 years
- 90% living in peri-urban areas, 10% living in urban areas
- 53% single persons, 36% married persons or persons living with long-term partner, 11% divorced/ separated/widowed persons
- 52% persons presented with TB symptoms
- Median CD4 count of 293 cells/mm<sup>3</sup>, 25<sup>th</sup> - 75<sup>th</sup> percentile: 127-514 cells/mm<sup>3</sup>

Note: Percentages may not add up to 100% due to rounding.

### Eligibility Criteria

Adults over the age of 18 years, who were not pregnant, confirmed HIV-positive test result, not currently on ART nor prescribed ART in the past 3 months, and presented at the study clinic for any reason that led to referral for HIV testing or care.

### Assignment Method

This study used an individually randomized, non-blinded, parallel-group design. Participants were randomized 1:1 to the intervention and standard (comparison) arm.

### Comparison

Participants in the standard group received usual care per national guidelines for ART initiation, which could include same day initiation.

### Relevant Outcomes Measured

- Engagement in HIV care was measured as having initiated ART within 28 days of study enrolment and an observed clinic visit in either the participant's paper record or the site's electronic patient register between 5 and 8 months after study enrollment. The PRS Project considers the outcome of ART initiation within 1 month and retention on ART 8 months after study enrollment as engagement in HIV care (i.e., 1 visit in a time period).
- Viral suppression was measured as having a viral load < 400 copies/mL.

### Participant Retention

Because participant retention is not a criterion for the Linkage to, Retention in and Re-engagement in HIV Care (LRC) chapter, the Prevention Research Synthesis project does not evaluate that information.

### Significant Findings on Relevant Outcomes

- A greater percentage of intervention arm participants-initiated ART  $\leq$  28 days and engaged in care at 8 months after study enrollment compared to standard arm participants. (74% v. 59%, respectively; Crude Relative Risk [CRR] = 1.26, 95% Confidence Interval [CI]: 1.12–1.42).
- A greater percentage of intervention arm participants-initiated ART and were known to be virally suppressed  $\leq$  8 months after study enrollment compared to standard arm participants (44% vs. 32%, respectively; CRR = 1.39, 95% CI: 1.12–1.71).
- A greater percentage of intervention arm participants-initiated ART after study enrollment compared to standard arm participants, respectively, within:
  - the same day (87% vs. 38%; CRR = 2.26, 95% CI: 1.95 – 2.63)
  - 7 days (91 % vs 68 %; CRR = 1.34, 95% CI: 1.23 – 1.46)
  - 14 days (93% vs. 77%; CRR = 1.21, 95% CI: 1.12 – 1.29)
  - 28 days (94% vs. 82%: CRR = 1.14, 95% CI: 1.08 – 1.22)

### Considerations

*Additional significant positive findings on non-relevant outcomes*

- None reported

*Non-significant findings on relevant outcomes*

- None reported

*Negative findings*

- None reported

#### *Other related findings*

- None reported

#### *Implementation research-related findings*

- None reported

#### *Process/study execution findings*

- None reported

#### *Adverse events*

- Six patients who initiated ART within  $\leq 28$  days of study enrollment were known to have died during study follow-up. One additional standard arm patient died without having initiated ART. Investigators concluded that none of these deaths were related to study procedures.
- One intervention arm patient who was eligible for same-day initiation and did start ART on the same day was found to have an elevated serum creatinine.
- Only about 1 out of 10 patients had serious TB symptoms or other reasons for not starting ART that day. None of the patients who had mild TB symptoms and started ART on the same day reported any serious TB-related illness after ART initiation.

#### **Funding**

The Bill & Melinda Gates Foundation under the terms of (#OPP1136158 to Boston University (SR)).

## REFERENCES AND CONTACT INFORMATION

Maskew, M., Brennan, A. T., Fox, M. P., Vezi, L., Venter, W. D., Ehrenkranz, P., & Rosen, S. (2020). [A clinical algorithm for same-day HIV treatment initiation in settings with high TB symptom prevalence in South Africa: The SLATE II individually randomized clinical trial](#). *PLoS Medicine*, 17(8), e1003226. doi: 10.1371/journal.pmed.1003226

Rosen, S., Maskew, M., Larson, B. A., Brennan, A. T., Tsikhutsu, I., Fox, M. P., Vezi, L., Bii, M., & Venter, W. D. (2019). [Simplified clinical algorithm for identifying patients eligible for same-day HIV treatment initiation \(SLATE\): Results from an individually randomized trial in South Africa and Kenya](#). *PLoS Medicine*, 16(9), e1002912. doi: 10.1371/journal.pmed.1002912

#### **Researcher: Mhairi Maskew, MBBCh, PhD**

School of Clinical Medicine  
University of the Witwatersrand  
1 Jan Smuts Avenue  
Braamfontein, Johannesburg, 2000, South Africa

**Email:** [mmaskew@heroza.org](mailto:mmaskew@heroza.org)

