

Data Brief 348. Secondhand Smoke Exposure Among Nonsmoking Youth: United States, 2013–2016

Data table for Figure 1. Percentage of youth with secondhand smoke exposure, by age and sex: United States, 2013–2016

Characteristic	Sample size	Percent	Standard error
3–17 years			
Both	4,515	35.4	2.0
Boys	2,304	36.6	2.3
Girls	2,211	34.2	2.1
3–11 years			
Both	2,833	38.1	2.1
Boys	1,452	38.0	2.2
Girls	1,381	38.2	2.5
12–17 years			
Both	1,682	32.1	2.2
Boys	852	34.8	2.8
Girls	830	29.2	2.5

NOTE: Secondhand smoke is defined as serum cotinine levels of 0.05–10.00 ng/mL (aged 3–17 years) and no reported tobacco use (aged 12–17 years).

SOURCE: NCHS, National Health and Nutrition Examination Survey, 2013–2016.

Data Brief 348. Secondhand Smoke Exposure Among Nonsmoking Youth: United States, 2013–2016

Data table for Figure 2. Percentage of youth with secondhand smoke exposure, by age and race and Hispanic origin: United States, 2013–2016

Characteristic	Sample size	Percent	Standard error
3–17 years			
Non-Hispanic white	1,154	34.3	3.3
Non-Hispanic black	1,051	61.8	3.0
Non-Hispanic Asian	377	18.3	2.9
Hispanic	1,641	24.9	1.9
3–11 years			
Non-Hispanic white	715	38.1	3.6
Non-Hispanic black	678	66.2	3.3
Non-Hispanic Asian	215	20.1	3.8
Hispanic	1,030	25.9	2.4
12–17 years			
Non-Hispanic white	439	30.1	3.6
Non-Hispanic black	373	51.6	4.0
Non-Hispanic Asian	162	16.0	3.3
Hispanic	611	23.4	2.1

NOTE: Secondhand smoke is defined as serum cotinine levels of 0.05–10.00 ng/mL (aged 3–17 years) and no reported tobacco use (aged 12–17 years).

SOURCE: NCHS, National Health and Nutrition Examination Survey, 2013–2016.

Data Brief 348. Secondhand Smoke Exposure Among Nonsmoking Youth: United States, 2013–2016

Data table for Figure 3. Percentage of youth with secondhand smoke exposure, by age and family income level: United States, 2013–2016

Family income level	Sample size	Percent	Standard error
3–17 years			
Less than 100% FPL	1,399	54.8	2.7
100% to less than 200% FPL	1,170	42.5	2.5
200% to less than 400% FPL	974	27.9	3.4
At or above 400% FPL	633	16.0	2.0
3–11 years			
Less than 100% FPL	939	55.1	2.9
100% to less than 200% FPL	729	45.2	3.8
200% to less than 400% FPL	570	30.9	3.6
At or above 400% FPL	392	15.1	2.3
12–17 years			
Less than 100% FPL	460	54.3	3.1
100% to less than 200% FPL	441	39.0	2.5
200% to less than 400% FPL	404	24.9	3.9
At or above 400% FPL	241	17.0	3.5

NOTES: FPL is federal poverty level. Secondhand smoke is defined as serum cotinine levels of 0.05–10.00 ng/mL (aged 3–17 years) and no reported tobacco use (aged 12–17 years).

SOURCE: NCHS, National Health and Nutrition Examination Survey, 2013–2016.

Data Brief 348. Secondhand Smoke Exposure Among Nonsmoking Youth: United States, 2013–2016

Data table for Figure 4. Percentage of youth with secondhand smoke exposure, by age and number of tobacco smokers living in their home: United States, 2013–2016

Number of tobacco smokers in the home	Sample size	Percent	Standard error
3–17 years			
No smokers	3,352	23.9	1.7
One smoker	697	67.2	2.3
Two smokers or more	397	80.9	4.0
3–11 years			
No smokers	2,087	26.1	1.7
One smoker	452	72.1	3.3
Two smokers or more	254	81.4	4.7
12–17 years			
No smokers	1,265	21.1	1.8
One smoker	245	60.2	3.6
Two smokers or more	143	80.3	5.0

NOTE: Secondhand smoke is defined as serum cotinine levels of 0.05–10.00 ng/mL (aged 3–17 years) and no reported tobacco use (aged 12–17 years).

SOURCE: NCHS, National Health and Nutrition Examination Survey, 2013–2016.