

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,403,351	100.0	873.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	710,760	29.6	258.2
2	Malignant neoplasms (C00-C97)	553,091	23.0	200.9
3	Cerebrovascular diseases (I60-I69)	167,661	7.0	60.9
4	Chronic lower respiratory diseases (J40-J47)	122,009	5.1	44.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	97,900	4.1	35.6
6	Diabetes mellitus (E10-E14)	69,301	2.9	25.2
7	Influenza and pneumonia (J10-J18)	65,313	2.7	23.7
8	Alzheimer's disease (G30)	49,558	2.1	18.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37,251	1.5	13.5
10	Septicemia (A40-A41)	31,224	1.3	11.3
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	29,350	1.2	10.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	26,552	1.1	9.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	18,073	0.8	6.6
14	Assault (homicide) (X85-Y09,Y87.1)	16,765	0.7	6.1
15	Pneumonitis due to solids and liquids (J69)	16,636	0.7	6.0
...	All other causes (Residual)	391,907	16.3	142.4
All races, both sexes, 1-4 years				
...	All causes	4,979	100.0	32.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,826	36.7	12.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	495	9.9	3.3
3	Malignant neoplasms (C00-C97)	420	8.4	2.8
4	Assault (homicide) (X85-Y09,Y87.1)	356	7.2	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	181	3.6	1.2
6	Influenza and pneumonia (J10-J18)	103	2.1	0.7
7	Septicemia (A40-A41)	99	2.0	0.7
8	Certain conditions originating in the perinatal period (P00-P96)	79	1.6	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.0	0.3
11	Cerebrovascular diseases (I60-I69)	47	0.9	0.3
12	Meningitis (G00,G03)	36	0.7	0.2
13	Anemias (D50-D64)	35	0.7	0.2
14	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.4	0.1
15	Meningococcal infection (A39)	19	0.4	*
...	All other causes (Residual)	1,157	23.2	7.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-9 years				
...	All causes	3,253	100.0	16.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,391	42.8	7.0
2	Malignant neoplasms (C00-C97)	489	15.0	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	6.1	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	140	4.3	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	3.3	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	1.9	0.3
7	Chronic lower respiratory diseases (J40-J47)	48	1.5	0.2
8	Influenza and pneumonia (J10-J18)	47	1.4	0.2
9	Septicemia (A40-A41)	38	1.2	0.2
10	Cerebrovascular diseases (I60-I69)	25	0.8	0.1
10	Certain conditions originating in the perinatal period (P00-P96)	25	0.8	0.1
12	Anemias (D50-D64)	24	0.7	0.1
13	Complications of medical and surgical care (Y40-Y84,Y88)	18	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.5	*
15	Meningococcal infection (A39)	12	0.4	*
15	Meningitis (G00,G03)	12	0.4	*
...	All other causes (Residual)	601	18.5	3.0
All races, both sexes, 10-14 years				
...	All causes	4,160	100.0	20.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,588	38.2	8.0
2	Malignant neoplasms (C00-C97)	525	12.6	2.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	300	7.2	1.5
4	Assault (homicide) (X85-Y09,Y87.1)	231	5.6	1.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	4.8	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	165	4.0	0.8
7	Chronic lower respiratory diseases (J40-J47)	91	2.2	0.5
8	Cerebrovascular diseases (I60-I69)	51	1.2	0.3
9	Influenza and pneumonia (J10-J18)	40	1.0	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.9	0.2
11	Human immunodeficiency virus (HIV) disease (B20-B24)	29	0.7	0.1
12	Anemias (D50-D64)	28	0.7	0.1
13	Septicemia (A40-A41)	25	0.6	0.1
14	Diabetes mellitus (E10-E14)	18	0.4	*
14	Meningitis (G00,G03)	18	0.4	*
...	All other causes (Residual)	813	19.5	4.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 15-19 years				
...	All causes	13,563	100.0	68.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,755	49.8	34.0
2	Assault (homicide) (X85-Y09,Y87.1)	1,914	14.1	9.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,621	12.0	8.2
4	Malignant neoplasms (C00-C97)	745	5.5	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	403	3.0	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	225	1.7	1.1
7	Chronic lower respiratory diseases (J40-J47)	86	0.6	0.4
8	Cerebrovascular diseases (I60-I69)	67	0.5	0.3
9	Influenza and pneumonia (J10-J18)	65	0.5	0.3
10	Diabetes mellitus (E10-E14)	48	0.4	0.2
11	Septicemia (A40-A41)	40	0.3	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.3	0.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	35	0.3	0.2
14	Meningococcal infection (A39)	34	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	0.2	0.2
...	All other causes (Residual)	1,453	10.7	7.3
All races, both sexes, 20-24 years				
...	All causes	17,744	100.0	96.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,358	41.5	39.8
2	Assault (homicide) (X85-Y09,Y87.1)	3,025	17.0	16.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,373	13.4	12.8
4	Malignant neoplasms (C00-C97)	968	5.5	5.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	628	3.5	3.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	216	1.2	1.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	144	0.8	0.8
8	Cerebrovascular diseases (I60-I69)	132	0.7	0.7
9	Influenza and pneumonia (J10-J18)	124	0.7	0.7
10	Diabetes mellitus (E10-E14)	114	0.6	0.6
11	Chronic lower respiratory diseases (J40-J47)	104	0.6	0.6
12	Pregnancy, childbirth and the puerperium (O00-099)	77	0.4	0.4
13	Septicemia (A40-A41)	60	0.3	0.3
14	Anemias (D50-D64)	58	0.3	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	0.3	0.3
...	All other causes (Residual)	2,307	13.0	12.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-29 years				
...	All causes	17,681	100.0	99.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,898	33.4	33.0
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,341	13.2	13.1
3	Assault (homicide) (X85-Y09,Y87.1)	2,312	13.1	13.0
4	Malignant neoplasms (C00-C97)	1,359	7.7	7.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,013	5.7	5.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	647	3.7	3.6
7	Cerebrovascular diseases (I60-I69)	209	1.2	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	1.2	1.2
9	Diabetes mellitus (E10-E14)	206	1.2	1.2
10	Influenza and pneumonia (J10-J18)	142	0.8	0.8
11	Chronic lower respiratory diseases (J40-J47)	125	0.7	0.7
12	Septicemia (A40-A41)	104	0.6	0.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	88	0.5	0.5
14	Pregnancy, childbirth and the puerperium (O00-099)	87	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	0.5	0.4
...	All other causes (Residual)	2,862	16.2	16.0
All races, both sexes, 30-34 years				
...	All causes	22,770	100.0	116.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,871	25.8	30.0
2	Malignant neoplasms (C00-C97)	2,557	11.2	13.1
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,451	10.8	12.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,945	8.5	9.9
5	Assault (homicide) (X85-Y09,Y87.1)	1,852	8.1	9.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,790	7.9	9.1
7	Diabetes mellitus (E10-E14)	417	1.8	2.1
8	Cerebrovascular diseases (I60-I69)	393	1.7	2.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	327	1.4	1.7
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	269	1.2	1.4
11	Influenza and pneumonia (J10-J18)	222	1.0	1.1
12	Septicemia (A40-A41)	176	0.8	0.9
13	Chronic lower respiratory diseases (J40-J47)	171	0.8	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	0.7	0.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	97	0.4	0.5
...	All other causes (Residual)	4,068	17.9	20.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 35-39 years				
...	All causes	36,140	100.0	162.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,471	20.7	33.5
2	Malignant neoplasms (C00-C97)	5,565	15.4	25.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,333	12.0	19.5
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,112	8.6	14.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,896	8.0	13.0
6	Assault (homicide) (X85-Y09,Y87.1)	1,732	4.8	7.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,083	3.0	4.9
8	Cerebrovascular diseases (I60-I69)	880	2.4	4.0
9	Diabetes mellitus (E10-E14)	684	1.9	3.1
10	Influenza and pneumonia (J10-J18)	405	1.1	1.8
11	Septicemia (A40-A41)	324	0.9	1.5
12	Chronic lower respiratory diseases (J40-J47)	310	0.9	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	278	0.8	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	277	0.8	1.2
15	Viral hepatitis (B15-B19)	219	0.6	1.0
...	All other causes (Residual)	6,571	18.2	29.5
All races, both sexes, 40-44 years				
...	All causes	53,658	100.0	237.3
1	Malignant neoplasms (C00-C97)	10,955	20.4	48.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,848	16.5	39.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,942	14.8	35.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,450	6.4	15.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	3,023	5.6	13.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,288	4.3	10.1
7	Cerebrovascular diseases (I60-I69)	1,719	3.2	7.6
8	Assault (homicide) (X85-Y09,Y87.1)	1,487	2.8	6.6
9	Diabetes mellitus (E10-E14)	1,242	2.3	5.5
10	Influenza and pneumonia (J10-J18)	663	1.2	2.9
11	Chronic lower respiratory diseases (J40-J47)	620	1.2	2.7
12	Viral hepatitis (B15-B19)	560	1.0	2.5
13	Septicemia (A40-A41)	553	1.0	2.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	462	0.9	2.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	292	0.5	1.3
...	All other causes (Residual)	9,554	17.8	42.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-49 years				
...	All causes	70,832	100.0	356.0
1	Malignant neoplasms (C00-C97)	18,803	26.5	94.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,313	20.2	71.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,870	9.7	34.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,374	4.8	17.0
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,987	4.2	15.0
6	Cerebrovascular diseases (I60-I69)	2,625	3.7	13.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,551	3.6	12.8
8	Diabetes mellitus (E10-E14)	1,917	2.7	9.6
9	Chronic lower respiratory diseases (J40-J47)	1,174	1.7	5.9
10	Assault (homicide) (X85-Y09,Y87.1)	1,059	1.5	5.3
11	Viral hepatitis (B15-B19)	1,044	1.5	5.2
12	Influenza and pneumonia (J10-J18)	783	1.1	3.9
13	Septicemia (A40-A41)	781	1.1	3.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	687	1.0	3.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	352	0.5	1.8
...	All other causes (Residual)	11,512	16.3	57.9
All races, both sexes, 50-54 years				
...	All causes	89,509	100.0	518.6
1	Malignant neoplasms (C00-C97)	29,231	32.7	169.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,167	23.6	122.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,408	6.0	31.3
4	Cerebrovascular diseases (I60-I69)	3,386	3.8	19.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,280	3.7	19.0
6	Diabetes mellitus (E10-E14)	3,037	3.4	17.6
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,450	2.7	14.2
8	Chronic lower respiratory diseases (J40-J47)	2,077	2.3	12.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,591	1.8	9.2
10	Septicemia (A40-A41)	1,064	1.2	6.2
11	Influenza and pneumonia (J10-J18)	991	1.1	5.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	964	1.1	5.6
13	Viral hepatitis (B15-B19)	850	0.9	4.9
14	Assault (homicide) (X85-Y09,Y87.1)	696	0.8	4.0
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	526	0.6	3.0
...	All other causes (Residual)	12,791	14.3	74.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 55-59 years				
...	All causes	106,751	100.0	801.8
1	Malignant neoplasms (C00-C97)	38,930	36.5	292.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,676	25.9	207.9
3	Cerebrovascular diseases (I60-I69)	4,145	3.9	31.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,021	3.8	30.2
5	Diabetes mellitus (E10-E14)	3,992	3.7	30.0
6	Chronic lower respiratory diseases (J40-J47)	3,985	3.7	29.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,947	2.8	22.1
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,708	1.6	12.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,269	1.2	9.5
10	Septicemia (A40-A41)	1,260	1.2	9.5
11	Influenza and pneumonia (J10-J18)	1,198	1.1	9.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	797	0.7	6.0
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	597	0.6	4.5
14	Aortic aneurysm and dissection (I71)	585	0.5	4.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	513	0.5	3.9
...	All other causes (Residual)	13,128	12.3	98.6
All races, both sexes, 60-64 years				
...	All causes	134,095	100.0	1,257.9
1	Malignant neoplasms (C00-C97)	50,075	37.3	469.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,723	26.6	335.1
3	Chronic lower respiratory diseases (J40-J47)	6,754	5.0	63.4
4	Cerebrovascular diseases (I60-I69)	5,811	4.3	54.5
5	Diabetes mellitus (E10-E14)	5,194	3.9	48.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,484	2.6	32.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,827	2.1	26.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,831	1.4	17.2
9	Influenza and pneumonia (J10-J18)	1,681	1.3	15.8
10	Septicemia (A40-A41)	1,639	1.2	15.4
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,237	0.9	11.6
12	Aortic aneurysm and dissection (I71)	908	0.7	8.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	843	0.6	7.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	711	0.5	6.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	442	0.3	4.1
...	All other causes (Residual)	14,935	11.1	140.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-69 years				
...	All causes	181,739	100.0	1,928.2
1	Malignant neoplasms (C00-C97)	65,843	36.2	698.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,178	27.1	521.8
3	Chronic lower respiratory diseases (J40-J47)	11,833	6.5	125.5
4	Cerebrovascular diseases (I60-I69)	8,671	4.8	92.0
5	Diabetes mellitus (E10-E14)	7,054	3.9	74.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,375	1.9	35.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,811	1.5	29.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2,751	1.5	29.2
9	Influenza and pneumonia (J10-J18)	2,533	1.4	26.9
10	Septicemia (A40-A41)	2,303	1.3	24.4
11	Aortic aneurysm and dissection (I71)	1,552	0.9	16.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,094	0.6	11.6
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,073	0.6	11.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,007	0.6	10.7
15	Alzheimer's disease (G30)	868	0.5	9.2
...	All other causes (Residual)	19,793	10.9	210.0
All races, both sexes, 70-74 years				
...	All causes	259,470	100.0	2,968.1
1	Malignant neoplasms (C00-C97)	84,288	32.5	964.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,227	28.2	837.6
3	Chronic lower respiratory diseases (J40-J47)	19,324	7.4	221.0
4	Cerebrovascular diseases (I60-I69)	14,978	5.8	171.3
5	Diabetes mellitus (E10-E14)	9,620	3.7	110.0
6	Influenza and pneumonia (J10-J18)	4,656	1.8	53.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,323	1.7	49.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,179	1.6	47.8
9	Septicemia (A40-A41)	3,401	1.3	38.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,731	1.1	31.2
11	Alzheimer's disease (G30)	2,563	1.0	29.3
12	Aortic aneurysm and dissection (I71)	2,460	0.9	28.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,690	0.7	19.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,539	0.6	17.6
15	Parkinson's disease (G20-G21)	1,482	0.6	17.0
...	All other causes (Residual)	29,009	11.2	331.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 75-79 years				
...	All causes	337,700	100.0	4,556.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	101,764	30.1	1,373.1
2	Malignant neoplasms (C00-C97)	90,655	26.8	1,223.2
3	Cerebrovascular diseases (I60-I69)	24,678	7.3	333.0
4	Chronic lower respiratory diseases (J40-J47)	24,309	7.2	328.0
5	Diabetes mellitus (E10-E14)	11,607	3.4	156.6
6	Influenza and pneumonia (J10-J18)	7,912	2.3	106.8
7	Alzheimer's disease (G30)	6,319	1.9	85.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,893	1.7	79.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,596	1.7	75.5
10	Septicemia (A40-A41)	4,745	1.4	64.0
11	Parkinson's disease (G20-G21)	3,322	1.0	44.8
12	Aortic aneurysm and dissection (I71)	3,043	0.9	41.1
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,507	0.7	33.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2,367	0.7	31.9
15	Pneumonitis due to solids and liquids (J69)	2,258	0.7	30.5
...	All other causes (Residual)	40,725	12.1	549.5
All races, both sexes, 80-84 years				
...	All causes	362,745	100.0	7,399.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	118,296	32.6	2,413.1
2	Malignant neoplasms (C00-C97)	74,444	20.5	1,518.6
3	Cerebrovascular diseases (I60-I69)	32,342	8.9	659.7
4	Chronic lower respiratory diseases (J40-J47)	23,413	6.5	477.6
5	Influenza and pneumonia (J10-J18)	11,909	3.3	242.9
6	Alzheimer's disease (G30)	10,934	3.0	223.0
7	Diabetes mellitus (E10-E14)	10,577	2.9	215.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,565	1.8	133.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,162	1.7	125.7
10	Septicemia (A40-A41)	5,193	1.4	105.9
11	Parkinson's disease (G20-G21)	4,330	1.2	88.3
12	Pneumonitis due to solids and liquids (J69)	3,247	0.9	66.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,116	0.9	63.6
14	Aortic aneurysm and dissection (I71)	2,792	0.8	57.0
15	Atherosclerosis (I70)	2,592	0.7	52.9
...	All other causes (Residual)	46,833	12.9	955.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85-89 years				
...	All causes	335,813	100.0	12,452.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	121,003	36.0	4,486.9
2	Malignant neoplasms (C00-C97)	49,155	14.6	1,822.7
3	Cerebrovascular diseases (I60-I69)	34,030	10.1	1,261.9
4	Chronic lower respiratory diseases (J40-J47)	17,085	5.1	633.5
5	Influenza and pneumonia (J10-J18)	13,752	4.1	509.9
6	Alzheimer's disease (G30)	13,337	4.0	494.6
7	Diabetes mellitus (E10-E14)	8,002	2.4	296.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,150	1.8	228.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,933	1.8	220.0
10	Septicemia (A40-A41)	4,900	1.5	181.7
11	Pneumonitis due to solids and liquids (J69)	3,825	1.1	141.8
12	Parkinson's disease (G20-G21)	3,567	1.1	132.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,273	1.0	121.4
14	Atherosclerosis (I70)	3,188	0.9	118.2
15	Aortic aneurysm and dissection (I71)	2,134	0.6	79.1
...	All other causes (Residual)	46,479	13.8	1,723.5
All races, both sexes, 90-94 years				
...	All causes	219,283	100.0	18,677.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	86,424	39.4	7,361.2
2	Cerebrovascular diseases (I60-I69)	23,205	10.6	1,976.5
3	Malignant neoplasms (C00-C97)	21,564	9.8	1,836.7
4	Influenza and pneumonia (J10-J18)	11,318	5.2	964.0
5	Alzheimer's disease (G30)	10,257	4.7	873.6
6	Chronic lower respiratory diseases (J40-J47)	7,823	3.6	666.3
7	Diabetes mellitus (E10-E14)	4,104	1.9	349.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,939	1.8	335.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,915	1.8	333.5
10	Septicemia (A40-A41)	2,949	1.3	251.2
11	Atherosclerosis (I70)	2,870	1.3	244.5
12	Pneumonitis due to solids and liquids (J69)	2,779	1.3	236.7
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,375	1.1	202.3
14	Parkinson's disease (G20-G21)	1,627	0.7	138.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,143	0.5	97.4
...	All other causes (Residual)	32,991	15.0	2,810.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 95-99 years				
...	All causes	84,641	100.0	23,443.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,605	42.1	9,861.6
2	Cerebrovascular diseases (I60-I69)	8,530	10.1	2,362.6
3	Malignant neoplasms (C00-C97)	5,653	6.7	1,565.7
4	Influenza and pneumonia (J10-J18)	5,116	6.0	1,417.0
5	Alzheimer's disease (G30)	4,014	4.7	1,111.8
6	Chronic lower respiratory diseases (J40-J47)	2,238	2.6	619.9
7	Atherosclerosis (I70)	1,518	1.8	420.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,443	1.7	399.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,419	1.7	393.0
10	Diabetes mellitus (E10-E14)	1,250	1.5	346.2
11	Pneumonitis due to solids and liquids (J69)	1,111	1.3	307.7
12	Septicemia (A40-A41)	1,079	1.3	298.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,017	1.2	281.7
14	Parkinson's disease (G20-G21)	369	0.4	102.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	314	0.4	87.0
...	All other causes (Residual)	13,965	16.5	3,867.9
All races, both sexes, 100 years and over				
...	All causes	18,434	100.0	28,877.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,210	44.5	12,861.1
2	Cerebrovascular diseases (I60-I69)	1,611	8.7	2,523.7
3	Influenza and pneumonia (J10-J18)	1,361	7.4	2,132.0
4	Malignant neoplasms (C00-C97)	764	4.1	1,196.8
5	Alzheimer's disease (G30)	701	3.8	1,098.1
6	Atherosclerosis (I70)	429	2.3	672.0
7	Chronic lower respiratory diseases (J40-J47)	350	1.9	548.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	1.6	476.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	269	1.5	421.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	242	1.3	379.1
11	Pneumonitis due to solids and liquids (J69)	239	1.3	374.4
12	Septicemia (A40-A41)	216	1.2	338.4
13	Diabetes mellitus (E10-E14)	200	1.1	313.3
14	Nutritional deficiencies (E40-E64)	90	0.5	141.0
15	Anemias (D50-D64)	78	0.4	122.2
...	All other causes (Residual)	3,370	18.3	5,279.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, all ages				
...	All causes	1,177,578	100.0	874.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	344,807	29.3	256.1
2	Malignant neoplasms (C00-C97)	286,082	24.3	212.5
3	Cerebrovascular diseases (I60-I69)	64,769	5.5	48.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63,817	5.4	47.4
5	Chronic lower respiratory diseases (J40-J47)	60,004	5.1	44.6
6	Diabetes mellitus (E10-E14)	31,602	2.7	23.5
7	Influenza and pneumonia (J10-J18)	28,658	2.4	21.3
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	23,618	2.0	17.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,811	1.5	13.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	17,214	1.5	12.8
11	Alzheimer's disease (G30)	14,438	1.2	10.7
12	Septicemia (A40-A41)	13,537	1.1	10.1
13	Assault (homicide) (X85-Y09,Y87.1)	12,820	1.1	9.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	10,868	0.9	8.1
15	Aortic aneurysm and dissection (I71)	9,661	0.8	7.2
...	All other causes (Residual)	177,872	15.1	132.1
All races, male, 1-4 years				
...	All causes	2,824	100.0	36.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,098	38.9	14.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	259	9.2	3.3
3	Malignant neoplasms (C00-C97)	233	8.3	3.0
4	Assault (homicide) (X85-Y09,Y87.1)	198	7.0	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	107	3.8	1.4
6	Influenza and pneumonia (J10-J18)	54	1.9	0.7
7	Septicemia (A40-A41)	53	1.9	0.7
8	Certain conditions originating in the perinatal period (P00-P96)	52	1.8	0.7
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	1.2	0.4
10	Chronic lower respiratory diseases (J40-J47)	28	1.0	0.4
11	Meningitis (G00,G03)	20	0.7	0.3
12	Cerebrovascular diseases (I60-I69)	19	0.7	*
13	Anemias (D50-D64)	16	0.6	*
14	Meningococcal infection (A39)	12	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
...	All other causes (Residual)	631	22.3	8.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 5-9 years				
...	All causes	1,850	100.0	18.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	824	44.5	8.1
2	Malignant neoplasms (C00-C97)	269	14.5	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	5.7	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	77	4.2	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	3.5	0.6
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.7	0.3
7	Influenza and pneumonia (J10-J18)	27	1.5	0.3
8	Chronic lower respiratory diseases (J40-J47)	26	1.4	0.3
9	Septicemia (A40-A41)	24	1.3	0.2
10	Anemias (D50-D64)	17	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	13	0.7	*
12	Cerebrovascular diseases (I60-I69)	12	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
...	All other causes (Residual)	333	18.0	3.3
All races, male, 10-14 years				
...	All causes	2,551	100.0	25.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,037	40.7	10.2
2	Malignant neoplasms (C00-C97)	300	11.8	2.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	238	9.3	2.3
4	Assault (homicide) (X85-Y09,Y87.1)	150	5.9	1.5
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	4.0	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	3.9	1.0
7	Chronic lower respiratory diseases (J40-J47)	54	2.1	0.5
8	Cerebrovascular diseases (I60-I69)	21	0.8	0.2
8	Influenza and pneumonia (J10-J18)	21	0.8	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.7	*
11	Septicemia (A40-A41)	14	0.5	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.5	*
13	Anemias (D50-D64)	12	0.5	*
14	Meningitis (G00,G03)	10	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	10	0.4	*
...	All other causes (Residual)	443	17.4	4.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-19 years				
...	All causes	9,697	100.0	94.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,757	49.1	46.6
2	Assault (homicide) (X85-Y09,Y87.1)	1,613	16.6	15.8
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,351	13.9	13.2
4	Malignant neoplasms (C00-C97)	446	4.6	4.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	248	2.6	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	1.4	1.3
7	Chronic lower respiratory diseases (J40-J47)	58	0.6	0.6
8	Cerebrovascular diseases (I60-I69)	38	0.4	0.4
9	Influenza and pneumonia (J10-J18)	35	0.4	0.3
10	Diabetes mellitus (E10-E14)	31	0.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.3	0.3
11	Legal intervention (Y35,Y89.0)	27	0.3	0.3
13	Meningococcal infection (A39)	24	0.2	0.2
14	Septicemia (A40-A41)	23	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.2	*
...	All other causes (Residual)	863	8.9	8.4
All races, male, 20-24 years				
...	All causes	13,374	100.0	142.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,703	42.6	60.6
2	Assault (homicide) (X85-Y09,Y87.1)	2,590	19.4	27.5
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,073	15.5	22.0
4	Malignant neoplasms (C00-C97)	576	4.3	6.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	390	2.9	4.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	1.0	1.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	80	0.6	0.8
8	Cerebrovascular diseases (I60-I69)	68	0.5	0.7
9	Influenza and pneumonia (J10-J18)	67	0.5	0.7
10	Diabetes mellitus (E10-E14)	63	0.5	0.7
11	Chronic lower respiratory diseases (J40-J47)	62	0.5	0.7
12	Legal intervention (Y35,Y89.0)	44	0.3	0.5
13	Septicemia (A40-A41)	37	0.3	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.3	0.4
15	Anemias (D50-D64)	30	0.2	0.3
...	All other causes (Residual)	1,425	10.7	15.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 25-29 years				
...	All causes	12,619	100.0	141.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,606	36.5	51.8
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,956	15.5	22.0
3	Assault (homicide) (X85-Y09,Y87.1)	1,918	15.2	21.6
4	Malignant neoplasms (C00-C97)	692	5.5	7.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	640	5.1	7.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	371	2.9	4.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	1.0	1.3
8	Diabetes mellitus (E10-E14)	118	0.9	1.3
9	Cerebrovascular diseases (I60-I69)	111	0.9	1.2
10	Influenza and pneumonia (J10-J18)	82	0.6	0.9
11	Chronic lower respiratory diseases (J40-J47)	72	0.6	0.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	63	0.5	0.7
13	Septicemia (A40-A41)	62	0.5	0.7
14	Legal intervention (Y35,Y89.0)	58	0.5	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.3	0.5
...	All other causes (Residual)	1,706	13.5	19.2
All races, male, 30-34 years				
...	All causes	15,271	100.0	157.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,453	29.2	45.9
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,982	13.0	20.4
3	Assault (homicide) (X85-Y09,Y87.1)	1,443	9.4	14.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,282	8.4	13.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,231	8.1	12.7
6	Malignant neoplasms (C00-C97)	1,162	7.6	12.0
7	Diabetes mellitus (E10-E14)	258	1.7	2.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	217	1.4	2.2
9	Cerebrovascular diseases (I60-I69)	197	1.3	2.0
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	159	1.0	1.6
11	Influenza and pneumonia (J10-J18)	122	0.8	1.3
12	Septicemia (A40-A41)	96	0.6	1.0
13	Chronic lower respiratory diseases (J40-J47)	93	0.6	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	85	0.6	0.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.3	0.5
...	All other causes (Residual)	2,439	16.0	25.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-39 years				
...	All causes	23,252	100.0	209.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,537	23.8	50.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,945	12.7	26.6
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,457	10.6	22.2
4	Malignant neoplasms (C00-C97)	2,402	10.3	21.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,172	9.3	19.6
6	Assault (homicide) (X85-Y09,Y87.1)	1,235	5.3	11.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	716	3.1	6.5
8	Cerebrovascular diseases (I60-I69)	430	1.8	3.9
9	Diabetes mellitus (E10-E14)	396	1.7	3.6
10	Influenza and pneumonia (J10-J18)	211	0.9	1.9
11	Septicemia (A40-A41)	162	0.7	1.5
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	154	0.7	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	0.7	1.4
14	Viral hepatitis (B15-B19)	151	0.6	1.4
15	Chronic lower respiratory diseases (J40-J47)	145	0.6	1.3
...	All other causes (Residual)	3,987	17.1	36.0
All races, male, 40-44 years				
...	All causes	34,045	100.0	303.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,341	18.6	56.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,907	17.4	52.6
3	Malignant neoplasms (C00-C97)	4,942	14.5	44.0
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,657	7.8	23.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	2,269	6.7	20.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,598	4.7	14.2
7	Assault (homicide) (X85-Y09,Y87.1)	1,084	3.2	9.7
8	Cerebrovascular diseases (I60-I69)	867	2.5	7.7
9	Diabetes mellitus (E10-E14)	755	2.2	6.7
10	Influenza and pneumonia (J10-J18)	419	1.2	3.7
11	Viral hepatitis (B15-B19)	389	1.1	3.5
12	Chronic lower respiratory diseases (J40-J47)	292	0.9	2.6
13	Septicemia (A40-A41)	289	0.8	2.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	277	0.8	2.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	162	0.5	1.4
...	All other causes (Residual)	5,797	17.0	51.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 45-49 years				
...	All causes	45,121	100.0	461.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,596	23.5	108.4
2	Malignant neoplasms (C00-C97)	9,243	20.5	94.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,103	11.3	52.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,515	5.6	25.7
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,307	5.1	23.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,008	4.5	20.5
7	Cerebrovascular diseases (I60-I69)	1,401	3.1	14.3
8	Diabetes mellitus (E10-E14)	1,124	2.5	11.5
9	Viral hepatitis (B15-B19)	787	1.7	8.1
10	Assault (homicide) (X85-Y09,Y87.1)	773	1.7	7.9
11	Chronic lower respiratory diseases (J40-J47)	617	1.4	6.3
12	Influenza and pneumonia (J10-J18)	464	1.0	4.7
13	Septicemia (A40-A41)	439	1.0	4.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	433	1.0	4.4
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	224	0.5	2.3
...	All other causes (Residual)	7,087	15.7	72.5
All races, male, 50-54 years				
...	All causes	55,277	100.0	658.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,331	27.7	182.6
2	Malignant neoplasms (C00-C97)	14,972	27.1	178.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,960	7.2	47.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,460	4.5	29.3
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,842	3.3	21.9
6	Cerebrovascular diseases (I60-I69)	1,834	3.3	21.8
7	Diabetes mellitus (E10-E14)	1,697	3.1	20.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,288	2.3	15.3
9	Chronic lower respiratory diseases (J40-J47)	1,044	1.9	12.4
10	Viral hepatitis (B15-B19)	638	1.2	7.6
11	Influenza and pneumonia (J10-J18)	602	1.1	7.2
12	Septicemia (A40-A41)	600	1.1	7.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	549	1.0	6.5
14	Assault (homicide) (X85-Y09,Y87.1)	511	0.9	6.1
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	331	0.6	3.9
...	All other causes (Residual)	7,618	13.8	90.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-59 years				
...	All causes	64,425	100.0	1,007.5
1	Malignant neoplasms (C00-C97)	20,912	32.5	327.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,456	30.2	304.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,804	4.4	43.9
4	Cerebrovascular diseases (I60-I69)	2,294	3.6	35.9
5	Diabetes mellitus (E10-E14)	2,173	3.4	34.0
6	Chronic lower respiratory diseases (J40-J47)	2,072	3.2	32.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,066	3.2	32.3
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,306	2.0	20.4
9	Influenza and pneumonia (J10-J18)	710	1.1	11.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	697	1.1	10.9
11	Septicemia (A40-A41)	657	1.0	10.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	655	1.0	10.2
13	Aortic aneurysm and dissection (I71)	438	0.7	6.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	352	0.5	5.5
15	Viral hepatitis (B15-B19)	338	0.5	5.3
15	Assault (homicide) (X85-Y09,Y87.1)	338	0.5	5.3
...	All other causes (Residual)	7,157	11.1	111.9
All races, male, 60-64 years				
...	All causes	78,896	100.0	1,565.5
1	Malignant neoplasms (C00-C97)	27,515	34.9	546.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,829	30.2	472.8
3	Chronic lower respiratory diseases (J40-J47)	3,492	4.4	69.3
4	Cerebrovascular diseases (I60-I69)	3,202	4.1	63.5
5	Diabetes mellitus (E10-E14)	2,770	3.5	55.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,329	3.0	46.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,939	2.5	38.5
8	Influenza and pneumonia (J10-J18)	970	1.2	19.2
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	959	1.2	19.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	954	1.2	18.9
11	Septicemia (A40-A41)	856	1.1	17.0
12	Aortic aneurysm and dissection (I71)	673	0.9	13.4
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	466	0.6	9.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	387	0.5	7.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	358	0.5	7.1
...	All other causes (Residual)	8,197	10.4	162.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 65-69 years				
...	All causes	103,935	100.0	2,399.3
1	Malignant neoplasms (C00-C97)	36,730	35.3	847.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,890	29.7	713.1
3	Chronic lower respiratory diseases (J40-J47)	6,149	5.9	141.9
4	Cerebrovascular diseases (I60-I69)	4,513	4.3	104.2
5	Diabetes mellitus (E10-E14)	3,556	3.4	82.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,117	2.0	48.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,729	1.7	39.9
8	Influenza and pneumonia (J10-J18)	1,447	1.4	33.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,444	1.4	33.3
10	Septicemia (A40-A41)	1,182	1.1	27.3
11	Aortic aneurysm and dissection (I71)	1,097	1.1	25.3
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	870	0.8	20.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	587	0.6	13.6
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	504	0.5	11.6
15	Alzheimer's disease (G30)	420	0.4	9.7
...	All other causes (Residual)	10,700	10.3	247.0
All races, male, 70-74 years				
...	All causes	143,473	100.0	3,705.4
1	Malignant neoplasms (C00-C97)	46,461	32.4	1,199.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,699	30.5	1,128.6
3	Chronic lower respiratory diseases (J40-J47)	10,061	7.0	259.8
4	Cerebrovascular diseases (I60-I69)	7,529	5.2	194.4
5	Diabetes mellitus (E10-E14)	4,670	3.3	120.6
6	Influenza and pneumonia (J10-J18)	2,566	1.8	66.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,522	1.8	65.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,120	1.5	54.8
9	Septicemia (A40-A41)	1,685	1.2	43.5
10	Aortic aneurysm and dissection (I71)	1,633	1.1	42.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,588	1.1	41.0
12	Alzheimer's disease (G30)	1,134	0.8	29.3
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,018	0.7	26.3
14	Parkinson's disease (G20-G21)	940	0.7	24.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	828	0.6	21.4
...	All other causes (Residual)	15,019	10.5	387.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-79 years				
...	All causes	173,327	100.0	5,591.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,421	31.4	1,755.5
2	Malignant neoplasms (C00-C97)	48,284	27.9	1,557.6
3	Chronic lower respiratory diseases (J40-J47)	12,305	7.1	396.9
4	Cerebrovascular diseases (I60-I69)	11,192	6.5	361.0
5	Diabetes mellitus (E10-E14)	5,288	3.1	170.6
6	Influenza and pneumonia (J10-J18)	4,084	2.4	131.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,039	1.8	98.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,039	1.8	98.0
9	Alzheimer's disease (G30)	2,433	1.4	78.5
10	Septicemia (A40-A41)	2,249	1.3	72.5
11	Parkinson's disease (G20-G21)	2,052	1.2	66.2
12	Aortic aneurysm and dissection (I71)	1,885	1.1	60.8
13	Pneumonitis due to solids and liquids (J69)	1,291	0.7	41.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,240	0.7	40.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,120	0.6	36.1
...	All other causes (Residual)	19,405	11.2	626.0
All races, male, 80-84 years				
...	All causes	166,892	100.0	8,956.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,272	33.1	2,966.4
2	Malignant neoplasms (C00-C97)	37,623	22.5	2,019.2
3	Cerebrovascular diseases (I60-I69)	12,754	7.6	684.5
4	Chronic lower respiratory diseases (J40-J47)	11,533	6.9	619.0
5	Influenza and pneumonia (J10-J18)	5,667	3.4	304.1
6	Diabetes mellitus (E10-E14)	4,450	2.7	238.8
7	Alzheimer's disease (G30)	3,795	2.3	203.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,174	1.9	170.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,024	1.8	162.3
10	Parkinson's disease (G20-G21)	2,478	1.5	133.0
11	Septicemia (A40-A41)	2,233	1.3	119.8
12	Pneumonitis due to solids and liquids (J69)	1,747	1.0	93.8
13	Aortic aneurysm and dissection (I71)	1,558	0.9	83.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,199	0.7	64.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,155	0.7	62.0
...	All other causes (Residual)	19,230	11.5	1,032.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 85-89 years				
...	All causes	128,877	100.0	14,604.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,756	35.5	5,185.1
2	Malignant neoplasms (C00-C97)	22,714	17.6	2,574.0
3	Cerebrovascular diseases (I60-I69)	10,864	8.4	1,231.1
4	Chronic lower respiratory diseases (J40-J47)	7,843	6.1	888.8
5	Influenza and pneumonia (J10-J18)	5,575	4.3	631.8
6	Alzheimer's disease (G30)	3,687	2.9	417.8
7	Diabetes mellitus (E10-E14)	2,789	2.2	316.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,672	2.1	302.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,563	2.0	290.4
10	Parkinson's disease (G20-G21)	1,879	1.5	212.9
11	Pneumonitis due to solids and liquids (J69)	1,839	1.4	208.4
12	Septicemia (A40-A41)	1,690	1.3	191.5
13	Atherosclerosis (I70)	1,090	0.8	123.5
14	Aortic aneurysm and dissection (I71)	1,063	0.8	120.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	991	0.8	112.3
...	All other causes (Residual)	15,862	12.3	1,797.5
All races, male, 90-94 years				
...	All causes	64,439	100.0	20,233.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,475	38.0	7,684.9
2	Malignant neoplasms (C00-C97)	8,506	13.2	2,670.8
3	Cerebrovascular diseases (I60-I69)	5,588	8.7	1,754.6
4	Influenza and pneumonia (J10-J18)	3,724	5.8	1,169.3
5	Chronic lower respiratory diseases (J40-J47)	3,165	4.9	993.8
6	Alzheimer's disease (G30)	2,101	3.3	659.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,473	2.3	462.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,391	2.2	436.8
9	Diabetes mellitus (E10-E14)	1,165	1.8	365.8
10	Pneumonitis due to solids and liquids (J69)	1,116	1.7	350.4
11	Septicemia (A40-A41)	812	1.3	255.0
12	Atherosclerosis (I70)	738	1.1	231.7
13	Parkinson's disease (G20-G21)	726	1.1	228.0
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	569	0.9	178.7
15	Aortic aneurysm and dissection (I71)	425	0.7	133.4
...	All other causes (Residual)	8,465	13.1	2,657.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 95-99 years				
...	All causes	18,552	100.0	22,992.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,430	40.0	9,208.3
2	Malignant neoplasms (C00-C97)	1,850	10.0	2,292.8
3	Cerebrovascular diseases (I60-I69)	1,564	8.4	1,938.3
4	Influenza and pneumonia (J10-J18)	1,373	7.4	1,701.6
5	Chronic lower respiratory diseases (J40-J47)	776	4.2	961.7
6	Alzheimer's disease (G30)	564	3.0	699.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	453	2.4	561.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	397	2.1	492.0
9	Pneumonitis due to solids and liquids (J69)	367	2.0	454.8
10	Atherosclerosis (I70)	283	1.5	350.7
11	Diabetes mellitus (E10-E14)	260	1.4	322.2
12	Septicemia (A40-A41)	199	1.1	246.6
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	155	0.8	192.1
14	Parkinson's disease (G20-G21)	115	0.6	142.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	86	0.5	106.6
...	All other causes (Residual)	2,680	14.4	3,321.4
All races, male, 100 years and over				
...	All causes	2,874	100.0	24,805.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,235	43.0	10,659.4
2	Influenza and pneumonia (J10-J18)	273	9.5	2,356.3
3	Cerebrovascular diseases (I60-I69)	196	6.8	1,691.7
4	Malignant neoplasms (C00-C97)	190	6.6	1,639.9
5	Chronic lower respiratory diseases (J40-J47)	92	3.2	794.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	73	2.5	630.1
7	Pneumonitis due to solids and liquids (J69)	65	2.3	561.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	2.2	535.1
9	Atherosclerosis (I70)	55	1.9	474.7
10	Alzheimer's disease (G30)	54	1.9	466.1
11	Septicemia (A40-A41)	35	1.2	302.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	32	1.1	276.2
13	Diabetes mellitus (E10-E14)	20	0.7	172.6
14	Anemias (D50-D64)	14	0.5	*
15	Nutritional deficiencies (E40-E64)	13	0.5	*
...	All other causes (Residual)	465	16.2	4,013.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,225,773	100.0	871.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	365,953	29.9	260.2
2	Malignant neoplasms (C00-C97)	267,009	21.8	189.9
3	Cerebrovascular diseases (I60-I69)	102,892	8.4	73.2
4	Chronic lower respiratory diseases (J40-J47)	62,005	5.1	44.1
5	Diabetes mellitus (E10-E14)	37,699	3.1	26.8
6	Influenza and pneumonia (J10-J18)	36,655	3.0	26.1
7	Alzheimer's disease (G30)	35,120	2.9	25.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,083	2.8	24.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19,440	1.6	13.8
10	Septicemia (A40-A41)	17,687	1.4	12.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11,178	0.9	7.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,338	0.8	6.6
13	Atherosclerosis (I70)	9,036	0.7	6.4
14	Pneumonitis due to solids and liquids (J69)	8,246	0.7	5.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,987	0.6	5.0
...	All other causes (Residual)	202,445	16.5	143.9
All races, female, 1-4 years				
...	All causes	2,155	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	728	33.8	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	236	11.0	3.2
3	Malignant neoplasms (C00-C97)	187	8.7	2.5
4	Assault (homicide) (X85-Y09,Y87.1)	158	7.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	3.4	1.0
6	Influenza and pneumonia (J10-J18)	49	2.3	0.7
7	Septicemia (A40-A41)	46	2.1	0.6
8	Cerebrovascular diseases (I60-I69)	28	1.3	0.4
9	Certain conditions originating in the perinatal period (P00-P96)	27	1.3	0.4
10	Chronic lower respiratory diseases (J40-J47)	23	1.1	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.9	0.3
12	Anemias (D50-D64)	19	0.9	*
13	Meningitis (G00,G03)	16	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.4	*
...	All other causes (Residual)	525	24.4	7.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-9 years				
...	All causes	1,403	100.0	14.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	567	40.4	5.9
2	Malignant neoplasms (C00-C97)	220	15.7	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	6.6	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	63	4.5	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.0	0.4
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	2.1	0.3
7	Chronic lower respiratory diseases (J40-J47)	22	1.6	0.2
8	Influenza and pneumonia (J10-J18)	20	1.4	0.2
9	Septicemia (A40-A41)	14	1.0	*
10	Cerebrovascular diseases (I60-I69)	13	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	12	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.6	*
13	Meningococcal infection (A39)	7	0.5	*
13	Anemias (D50-D64)	7	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.5	*
...	All other causes (Residual)	278	19.8	2.9
All races, female, 10-14 years				
...	All causes	1,609	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	551	34.2	5.7
2	Malignant neoplasms (C00-C97)	225	14.0	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	6.2	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	81	5.0	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	4.1	0.7
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	62	3.9	0.6
7	Chronic lower respiratory diseases (J40-J47)	37	2.3	0.4
8	Cerebrovascular diseases (I60-I69)	30	1.9	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.2	0.2
10	Influenza and pneumonia (J10-J18)	19	1.2	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	16	1.0	*
11	Anemias (D50-D64)	16	1.0	*
13	Septicemia (A40-A41)	11	0.7	*
14	Diabetes mellitus (E10-E14)	10	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.6	*
...	All other causes (Residual)	357	22.2	3.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 15-19 years				
...	All causes	3,866	100.0	40.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,998	51.7	20.7
2	Assault (homicide) (X85-Y09,Y87.1)	301	7.8	3.1
3	Malignant neoplasms (C00-C97)	299	7.7	3.1
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	270	7.0	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	155	4.0	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	2.3	0.9
7	Influenza and pneumonia (J10-J18)	30	0.8	0.3
8	Cerebrovascular diseases (I60-I69)	29	0.8	0.3
9	Chronic lower respiratory diseases (J40-J47)	28	0.7	0.3
10	Anemias (D50-D64)	18	0.5	*
10	Pregnancy, childbirth and the puerperium (O00-099)	18	0.5	*
12	Septicemia (A40-A41)	17	0.4	*
12	Diabetes mellitus (E10-E14)	17	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.4	*
...	All other causes (Residual)	568	14.7	5.9
All races, female, 20-24 years				
...	All causes	4,370	100.0	48.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,655	37.9	18.3
2	Assault (homicide) (X85-Y09,Y87.1)	435	10.0	4.8
3	Malignant neoplasms (C00-C97)	392	9.0	4.3
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	300	6.9	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	238	5.4	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	1.9	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	77	1.8	0.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	64	1.5	0.7
8	Cerebrovascular diseases (I60-I69)	64	1.5	0.7
10	Influenza and pneumonia (J10-J18)	57	1.3	0.6
11	Diabetes mellitus (E10-E14)	51	1.2	0.6
12	Chronic lower respiratory diseases (J40-J47)	42	1.0	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.7	0.3
14	Anemias (D50-D64)	28	0.6	0.3
15	Septicemia (A40-A41)	23	0.5	0.3
...	All other causes (Residual)	828	18.9	9.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-29 years				
...	All causes	5,062	100.0	56.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,292	25.5	14.4
2	Malignant neoplasms (C00-C97)	667	13.2	7.4
3	Assault (homicide) (X85-Y09,Y87.1)	394	7.8	4.4
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	385	7.6	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	373	7.4	4.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	276	5.5	3.1
7	Cerebrovascular diseases (I60-I69)	98	1.9	1.1
8	Diabetes mellitus (E10-E14)	88	1.7	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	1.7	1.0
10	Pregnancy, childbirth and the puerperium (O00-099)	87	1.7	1.0
11	Influenza and pneumonia (J10-J18)	60	1.2	0.7
12	Chronic lower respiratory diseases (J40-J47)	53	1.0	0.6
13	Septicemia (A40-A41)	42	0.8	0.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.8	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	0.7	0.4
...	All other causes (Residual)	1,085	21.4	12.1
All races, female, 30-34 years				
...	All causes	7,499	100.0	76.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,418	18.9	14.4
2	Malignant neoplasms (C00-C97)	1,395	18.6	14.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	663	8.8	6.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	559	7.5	5.7
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	469	6.3	4.8
6	Assault (homicide) (X85-Y09,Y87.1)	409	5.5	4.1
7	Cerebrovascular diseases (I60-I69)	196	2.6	2.0
8	Diabetes mellitus (E10-E14)	159	2.1	1.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	110	1.5	1.1
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	1.5	1.1
11	Influenza and pneumonia (J10-J18)	100	1.3	1.0
12	Pregnancy, childbirth and the puerperium (O00-099)	95	1.3	1.0
13	Septicemia (A40-A41)	80	1.1	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.1	0.8
15	Chronic lower respiratory diseases (J40-J47)	78	1.0	0.8
...	All other causes (Residual)	1,579	21.1	16.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 35-39 years				
...	All causes	12,888	100.0	115.1
1	Malignant neoplasms (C00-C97)	3,163	24.5	28.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,934	15.0	17.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,388	10.8	12.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	724	5.6	6.5
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	655	5.1	5.9
6	Assault (homicide) (X85-Y09,Y87.1)	497	3.9	4.4
7	Cerebrovascular diseases (I60-I69)	450	3.5	4.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	367	2.8	3.3
9	Diabetes mellitus (E10-E14)	288	2.2	2.6
10	Influenza and pneumonia (J10-J18)	194	1.5	1.7
11	Chronic lower respiratory diseases (J40-J47)	165	1.3	1.5
12	Septicemia (A40-A41)	162	1.3	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	1.0	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	1.0	1.1
15	Pregnancy, childbirth and the puerperium (O00-099)	86	0.7	0.8
...	All other causes (Residual)	2,566	19.9	22.9
All races, female, 40-44 years				
...	All causes	19,613	100.0	172.2
1	Malignant neoplasms (C00-C97)	6,013	30.7	52.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,507	12.8	22.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,035	10.4	17.9
4	Cerebrovascular diseases (I60-I69)	852	4.3	7.5
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	793	4.0	7.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	754	3.8	6.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	690	3.5	6.1
8	Diabetes mellitus (E10-E14)	487	2.5	4.3
9	Assault (homicide) (X85-Y09,Y87.1)	403	2.1	3.5
10	Chronic lower respiratory diseases (J40-J47)	328	1.7	2.9
11	Septicemia (A40-A41)	264	1.3	2.3
12	Influenza and pneumonia (J10-J18)	244	1.2	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	185	0.9	1.6
14	Viral hepatitis (B15-B19)	171	0.9	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	0.7	1.1
...	All other causes (Residual)	3,757	19.2	33.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-49 years				
...	All causes	25,711	100.0	254.0
1	Malignant neoplasms (C00-C97)	9,560	37.2	94.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,717	14.5	36.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,767	6.9	17.5
4	Cerebrovascular diseases (I60-I69)	1,224	4.8	12.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	859	3.3	8.5
6	Diabetes mellitus (E10-E14)	793	3.1	7.8
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	680	2.6	6.7
8	Chronic lower respiratory diseases (J40-J47)	557	2.2	5.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	543	2.1	5.4
10	Septicemia (A40-A41)	342	1.3	3.4
11	Influenza and pneumonia (J10-J18)	319	1.2	3.2
12	Assault (homicide) (X85-Y09,Y87.1)	286	1.1	2.8
13	Viral hepatitis (B15-B19)	257	1.0	2.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	254	1.0	2.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.5	1.3
...	All other causes (Residual)	4,420	17.2	43.7
All races, female, 50-54 years				
...	All causes	34,232	100.0	386.3
1	Malignant neoplasms (C00-C97)	14,259	41.7	160.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,836	17.0	65.9
3	Cerebrovascular diseases (I60-I69)	1,552	4.5	17.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,448	4.2	16.3
5	Diabetes mellitus (E10-E14)	1,340	3.9	15.1
6	Chronic lower respiratory diseases (J40-J47)	1,033	3.0	11.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	820	2.4	9.3
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	608	1.8	6.9
9	Septicemia (A40-A41)	464	1.4	5.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	415	1.2	4.7
11	Influenza and pneumonia (J10-J18)	389	1.1	4.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	303	0.9	3.4
13	Viral hepatitis (B15-B19)	212	0.6	2.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	195	0.6	2.2
15	Assault (homicide) (X85-Y09,Y87.1)	185	0.5	2.1
...	All other causes (Residual)	5,173	15.1	58.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 55-59 years				
...	All causes	42,326	100.0	611.8
1	Malignant neoplasms (C00-C97)	18,018	42.6	260.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,220	19.4	118.8
3	Chronic lower respiratory diseases (J40-J47)	1,913	4.5	27.6
4	Cerebrovascular diseases (I60-I69)	1,851	4.4	26.8
5	Diabetes mellitus (E10-E14)	1,819	4.3	26.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,217	2.9	17.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	881	2.1	12.7
8	Septicemia (A40-A41)	603	1.4	8.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	572	1.4	8.3
10	Influenza and pneumonia (J10-J18)	488	1.2	7.1
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	402	0.9	5.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	245	0.6	3.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	243	0.6	3.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	166	0.4	2.4
15	Viral hepatitis (B15-B19)	154	0.4	2.2
...	All other causes (Residual)	5,534	13.1	80.0
All races, female, 60-64 years				
...	All causes	55,199	100.0	982.0
1	Malignant neoplasms (C00-C97)	22,560	40.9	401.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,894	21.5	211.6
3	Chronic lower respiratory diseases (J40-J47)	3,262	5.9	58.0
4	Cerebrovascular diseases (I60-I69)	2,609	4.7	46.4
5	Diabetes mellitus (E10-E14)	2,424	4.4	43.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,155	2.1	20.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	888	1.6	15.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	877	1.6	15.6
9	Septicemia (A40-A41)	783	1.4	13.9
10	Influenza and pneumonia (J10-J18)	711	1.3	12.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	377	0.7	6.7
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	324	0.6	5.8
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	278	0.5	4.9
14	Aortic aneurysm and dissection (I71)	235	0.4	4.2
15	Alzheimer's disease (G30)	199	0.4	3.5
...	All other causes (Residual)	6,623	12.0	117.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-69 years				
...	All causes	77,804	100.0	1,527.5
1	Malignant neoplasms (C00-C97)	29,113	37.4	571.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,288	23.5	359.0
3	Chronic lower respiratory diseases (J40-J47)	5,684	7.3	111.6
4	Cerebrovascular diseases (I60-I69)	4,158	5.3	81.6
5	Diabetes mellitus (E10-E14)	3,498	4.5	68.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,367	1.8	26.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,258	1.6	24.7
8	Septicemia (A40-A41)	1,121	1.4	22.0
9	Influenza and pneumonia (J10-J18)	1,086	1.4	21.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,022	1.3	20.1
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	590	0.8	11.6
12	Aortic aneurysm and dissection (I71)	455	0.6	8.9
13	Alzheimer's disease (G30)	448	0.6	8.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	420	0.5	8.2
15	Pneumonitis due to solids and liquids (J69)	237	0.3	4.7
...	All other causes (Residual)	9,059	11.6	177.9
All races, female, 70-74 years				
...	All causes	115,997	100.0	2,381.8
1	Malignant neoplasms (C00-C97)	37,827	32.6	776.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,528	25.5	606.3
3	Chronic lower respiratory diseases (J40-J47)	9,263	8.0	190.2
4	Cerebrovascular diseases (I60-I69)	7,449	6.4	153.0
5	Diabetes mellitus (E10-E14)	4,950	4.3	101.6
6	Influenza and pneumonia (J10-J18)	2,090	1.8	42.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,059	1.8	42.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,801	1.6	37.0
9	Septicemia (A40-A41)	1,716	1.5	35.2
10	Alzheimer's disease (G30)	1,429	1.2	29.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,143	1.0	23.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	886	0.8	18.2
13	Aortic aneurysm and dissection (I71)	827	0.7	17.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	711	0.6	14.6
15	Parkinson's disease (G20-G21)	542	0.5	11.1
...	All other causes (Residual)	13,776	11.9	282.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 75-79 years				
...	All causes	164,373	100.0	3,812.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,343	28.8	1,098.1
2	Malignant neoplasms (C00-C97)	42,371	25.8	982.8
3	Cerebrovascular diseases (I60-I69)	13,486	8.2	312.8
4	Chronic lower respiratory diseases (J40-J47)	12,004	7.3	278.4
5	Diabetes mellitus (E10-E14)	6,319	3.8	146.6
6	Alzheimer's disease (G30)	3,886	2.4	90.1
7	Influenza and pneumonia (J10-J18)	3,828	2.3	88.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,854	1.7	66.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,557	1.6	59.3
10	Septicemia (A40-A41)	2,496	1.5	57.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,458	0.9	33.8
12	Parkinson's disease (G20-G21)	1,270	0.8	29.5
13	Aortic aneurysm and dissection (I71)	1,158	0.7	26.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,127	0.7	26.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,022	0.6	23.7
...	All other causes (Residual)	21,194	12.9	491.6
All races, female, 80-84 years				
...	All causes	195,853	100.0	6,444.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	63,024	32.2	2,073.9
2	Malignant neoplasms (C00-C97)	36,821	18.8	1,211.6
3	Cerebrovascular diseases (I60-I69)	19,588	10.0	644.6
4	Chronic lower respiratory diseases (J40-J47)	11,880	6.1	390.9
5	Alzheimer's disease (G30)	7,139	3.6	234.9
6	Influenza and pneumonia (J10-J18)	6,242	3.2	205.4
7	Diabetes mellitus (E10-E14)	6,127	3.1	201.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,391	1.7	111.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,138	1.6	103.3
10	Septicemia (A40-A41)	2,960	1.5	97.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,961	1.0	64.5
12	Parkinson's disease (G20-G21)	1,852	0.9	60.9
13	Pneumonitis due to solids and liquids (J69)	1,500	0.8	49.4
14	Atherosclerosis (I70)	1,495	0.8	49.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,304	0.7	42.9
...	All other causes (Residual)	27,431	14.0	902.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85-89 years				
...	All causes	206,936	100.0	11,405.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,247	36.4	4,147.4
2	Malignant neoplasms (C00-C97)	26,441	12.8	1,457.3
3	Cerebrovascular diseases (I60-I69)	23,166	11.2	1,276.8
4	Alzheimer's disease (G30)	9,650	4.7	531.9
5	Chronic lower respiratory diseases (J40-J47)	9,242	4.5	509.4
6	Influenza and pneumonia (J10-J18)	8,177	4.0	450.7
7	Diabetes mellitus (E10-E14)	5,213	2.5	287.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,478	1.7	191.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,370	1.6	185.7
10	Septicemia (A40-A41)	3,210	1.6	176.9
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,282	1.1	125.8
12	Atherosclerosis (I70)	2,098	1.0	115.6
13	Pneumonitis due to solids and liquids (J69)	1,986	1.0	109.5
14	Parkinson's disease (G20-G21)	1,688	0.8	93.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,222	0.6	67.4
...	All other causes (Residual)	30,466	14.7	1,679.2
All races, female, 90-94 years				
...	All causes	154,844	100.0	18,098.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	61,949	40.0	7,240.7
2	Cerebrovascular diseases (I60-I69)	17,617	11.4	2,059.1
3	Malignant neoplasms (C00-C97)	13,058	8.4	1,526.2
4	Alzheimer's disease (G30)	8,156	5.3	953.3
5	Influenza and pneumonia (J10-J18)	7,594	4.9	887.6
6	Chronic lower respiratory diseases (J40-J47)	4,658	3.0	544.4
7	Diabetes mellitus (E10-E14)	2,939	1.9	343.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,548	1.6	297.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,442	1.6	285.4
10	Septicemia (A40-A41)	2,137	1.4	249.8
11	Atherosclerosis (I70)	2,132	1.4	249.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,806	1.2	211.1
13	Pneumonitis due to solids and liquids (J69)	1,663	1.1	194.4
14	Parkinson's disease (G20-G21)	901	0.6	105.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	781	0.5	91.3
...	All other causes (Residual)	24,463	15.8	2,859.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 95-99 years				
...	All causes	66,089	100.0	23,572.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,175	42.6	10,049.6
2	Cerebrovascular diseases (I60-I69)	6,966	10.5	2,484.7
3	Malignant neoplasms (C00-C97)	3,803	5.8	1,356.5
4	Influenza and pneumonia (J10-J18)	3,743	5.7	1,335.1
5	Alzheimer's disease (G30)	3,450	5.2	1,230.6
6	Chronic lower respiratory diseases (J40-J47)	1,462	2.2	521.5
7	Atherosclerosis (I70)	1,235	1.9	440.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,022	1.5	364.5
9	Diabetes mellitus (E10-E14)	990	1.5	353.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	990	1.5	353.1
11	Septicemia (A40-A41)	880	1.3	313.9
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	862	1.3	307.5
13	Pneumonitis due to solids and liquids (J69)	744	1.1	265.4
14	Parkinson's disease (G20-G21)	254	0.4	90.6
15	Nutritional deficiencies (E40-E64)	250	0.4	89.2
...	All other causes (Residual)	11,263	17.0	4,017.3
All races, female, 100 years and over				
...	All causes	15,560	100.0	29,779.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,975	44.8	13,349.3
2	Cerebrovascular diseases (I60-I69)	1,415	9.1	2,708.1
3	Influenza and pneumonia (J10-J18)	1,088	7.0	2,082.3
4	Alzheimer's disease (G30)	647	4.2	1,238.3
5	Malignant neoplasms (C00-C97)	574	3.7	1,098.6
6	Atherosclerosis (I70)	374	2.4	715.8
7	Chronic lower respiratory diseases (J40-J47)	258	1.7	493.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	1.6	463.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	210	1.3	401.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	196	1.3	375.1
11	Septicemia (A40-A41)	181	1.2	346.4
12	Diabetes mellitus (E10-E14)	180	1.2	344.5
13	Pneumonitis due to solids and liquids (J69)	174	1.1	333.0
14	Nutritional deficiencies (E40-E64)	77	0.5	147.4
15	Anemias (D50-D64)	64	0.4	122.5
...	All other causes (Residual)	2,905	18.7	5,559.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, all ages				
...	All causes	2,071,287	100.0	915.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	621,719	30.0	274.8
2	Malignant neoplasms (C00-C97)	480,011	23.2	212.2
3	Cerebrovascular diseases (I60-I69)	144,580	7.0	63.9
4	Chronic lower respiratory diseases (J40-J47)	112,840	5.4	49.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82,592	4.0	36.5
6	Influenza and pneumonia (J10-J18)	57,914	2.8	25.6
7	Diabetes mellitus (E10-E14)	55,561	2.7	24.6
8	Alzheimer's disease (G30)	46,460	2.2	20.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29,598	1.4	13.1
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	26,475	1.3	11.7
11	Septicemia (A40-A41)	24,804	1.2	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	22,982	1.1	10.2
13	Parkinson's disease (G20-G21)	14,942	0.7	6.6
14	Pneumonitis due to solids and liquids (J69)	14,869	0.7	6.6
15	Aortic aneurysm and dissection (I71)	14,210	0.7	6.3
...	All other causes (Residual)	321,730	15.5	142.2
White, both sexes, 1-4 years				
...	All causes	3,494	100.0	29.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,335	38.2	11.1
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	347	9.9	2.9
3	Malignant neoplasms (C00-C97)	330	9.4	2.7
4	Assault (homicide) (X85-Y09,Y87.1)	199	5.7	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	3.6	1.0
6	Influenza and pneumonia (J10-J18)	69	2.0	0.6
7	Septicemia (A40-A41)	66	1.9	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	46	1.3	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.3	0.4
10	Cerebrovascular diseases (I60-I69)	30	0.9	0.2
11	Chronic lower respiratory diseases (J40-J47)	28	0.8	0.2
12	Meningitis (G00,G03)	23	0.7	0.2
13	Anemias (D50-D64)	15	0.4	*
14	Meningococcal infection (A39)	14	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.3	*
...	All other causes (Residual)	810	23.2	6.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 5-9 years				
...	All causes	2,359	100.0	15.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	996	42.2	6.4
2	Malignant neoplasms (C00-C97)	397	16.8	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	147	6.2	0.9
4	Assault (homicide) (X85-Y09,Y87.1)	98	4.2	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	71	3.0	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	1.9	0.3
7	Influenza and pneumonia (J10-J18)	36	1.5	0.2
8	Septicemia (A40-A41)	28	1.2	0.2
9	Chronic lower respiratory diseases (J40-J47)	23	1.0	0.1
10	Cerebrovascular diseases (I60-I69)	21	0.9	0.1
11	Certain conditions originating in the perinatal period (P00-P96)	18	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.5	*
13	Meningitis (G00,G03)	11	0.5	*
14	Anemias (D50-D64)	9	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.3	*
15	Pneumonitis due to solids and liquids (J69)	8	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (Residual)	422	17.9	2.7
White, both sexes, 10-14 years				
...	All causes	3,091	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,223	39.6	7.8
2	Malignant neoplasms (C00-C97)	421	13.6	2.7
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	248	8.0	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	4.7	0.9
5	Assault (homicide) (X85-Y09,Y87.1)	142	4.6	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	3.4	0.7
7	Chronic lower respiratory diseases (J40-J47)	40	1.3	0.3
8	Cerebrovascular diseases (I60-I69)	34	1.1	0.2
8	Influenza and pneumonia (J10-J18)	34	1.1	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	1.0	0.2
11	Septicemia (A40-A41)	21	0.7	0.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.4	*
13	Anemias (D50-D64)	12	0.4	*
13	Diabetes mellitus (E10-E14)	12	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	12	0.4	*
...	All other causes (Residual)	586	19.0	3.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-19 years				
...	All causes	10,273	100.0	65.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,731	55.8	36.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,354	13.2	8.6
3	Assault (homicide) (X85-Y09,Y87.1)	823	8.0	5.2
4	Malignant neoplasms (C00-C97)	576	5.6	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	264	2.6	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	159	1.5	1.0
7	Cerebrovascular diseases (I60-I69)	51	0.5	0.3
8	Influenza and pneumonia (J10-J18)	48	0.5	0.3
9	Chronic lower respiratory diseases (J40-J47)	40	0.4	0.3
10	Diabetes mellitus (E10-E14)	33	0.3	0.2
11	Meningococcal infection (A39)	29	0.3	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.3	0.2
13	Septicemia (A40-A41)	24	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.2	0.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.1	*
...	All other causes (Residual)	1,079	10.5	6.8
White, both sexes, 20-24 years				
...	All causes	12,745	100.0	86.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,092	47.8	41.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,961	15.4	13.3
3	Assault (homicide) (X85-Y09,Y87.1)	1,158	9.1	7.9
4	Malignant neoplasms (C00-C97)	757	5.9	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	392	3.1	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	173	1.4	1.2
7	Cerebrovascular diseases (I60-I69)	97	0.8	0.7
8	Influenza and pneumonia (J10-J18)	90	0.7	0.6
9	Diabetes mellitus (E10-E14)	72	0.6	0.5
10	Chronic lower respiratory diseases (J40-J47)	63	0.5	0.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.3	0.3
11	Pregnancy, childbirth and the puerperium (O00-099)	44	0.3	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.3	0.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	40	0.3	0.3
15	Septicemia (A40-A41)	36	0.3	0.2
...	All other causes (Residual)	1,683	13.2	11.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 25-29 years				
...	All causes	12,427	100.0	87.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,736	38.1	33.5
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,941	15.6	13.7
3	Malignant neoplasms (C00-C97)	1,081	8.7	7.6
4	Assault (homicide) (X85-Y09,Y87.1)	945	7.6	6.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	637	5.1	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	235	1.9	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	1.3	1.2
8	Cerebrovascular diseases (I60-I69)	160	1.3	1.1
9	Diabetes mellitus (E10-E14)	128	1.0	0.9
10	Influenza and pneumonia (J10-J18)	98	0.8	0.7
11	Chronic lower respiratory diseases (J40-J47)	67	0.5	0.5
12	Septicemia (A40-A41)	61	0.5	0.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	57	0.5	0.4
14	Pregnancy, childbirth and the puerperium (O00-099)	53	0.4	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	48	0.4	0.3
...	All other causes (Residual)	2,017	16.2	14.3
White, both sexes, 30-34 years				
...	All causes	16,292	100.0	103.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,783	29.4	30.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,143	13.2	13.6
3	Malignant neoplasms (C00-C97)	1,991	12.2	12.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,300	8.0	8.3
5	Assault (homicide) (X85-Y09,Y87.1)	869	5.3	5.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	748	4.6	4.8
7	Diabetes mellitus (E10-E14)	292	1.8	1.9
8	Cerebrovascular diseases (I60-I69)	260	1.6	1.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	255	1.6	1.6
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	228	1.4	1.4
11	Influenza and pneumonia (J10-J18)	163	1.0	1.0
12	Septicemia (A40-A41)	104	0.6	0.7
13	Chronic lower respiratory diseases (J40-J47)	99	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	0.5	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.5	0.5
...	All other causes (Residual)	2,887	17.7	18.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-39 years				
...	All causes	26,633	100.0	146.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,171	23.2	33.9
2	Malignant neoplasms (C00-C97)	4,316	16.2	23.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,083	11.6	16.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,796	10.5	15.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,402	5.3	7.7
6	Assault (homicide) (X85-Y09,Y87.1)	909	3.4	5.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	872	3.3	4.8
8	Cerebrovascular diseases (I60-I69)	558	2.1	3.1
9	Diabetes mellitus (E10-E14)	493	1.9	2.7
10	Influenza and pneumonia (J10-J18)	305	1.1	1.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	223	0.8	1.2
12	Septicemia (A40-A41)	204	0.8	1.1
13	Chronic lower respiratory diseases (J40-J47)	194	0.7	1.1
14	Viral hepatitis (B15-B19)	180	0.7	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	0.5	0.8
...	All other causes (Residual)	4,783	18.0	26.3
White, both sexes, 40-44 years				
...	All causes	39,863	100.0	213.3
1	Malignant neoplasms (C00-C97)	8,448	21.2	45.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,492	16.3	34.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,449	16.2	34.5
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	3,146	7.9	16.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,873	4.7	10.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,383	3.5	7.4
7	Cerebrovascular diseases (I60-I69)	1,087	2.7	5.8
8	Diabetes mellitus (E10-E14)	926	2.3	5.0
9	Assault (homicide) (X85-Y09,Y87.1)	816	2.0	4.4
10	Influenza and pneumonia (J10-J18)	462	1.2	2.5
11	Viral hepatitis (B15-B19)	445	1.1	2.4
12	Chronic lower respiratory diseases (J40-J47)	430	1.1	2.3
13	Septicemia (A40-A41)	374	0.9	2.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	268	0.7	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	236	0.6	1.3
...	All other causes (Residual)	7,028	17.6	37.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 45-49 years				
...	All causes	53,131	100.0	319.6
1	Malignant neoplasms (C00-C97)	14,743	27.7	88.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,645	20.0	64.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,554	10.5	33.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,817	5.3	16.9
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,762	5.2	16.6
6	Cerebrovascular diseases (I60-I69)	1,627	3.1	9.8
7	Diabetes mellitus (E10-E14)	1,344	2.5	8.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1,150	2.2	6.9
9	Chronic lower respiratory diseases (J40-J47)	882	1.7	5.3
10	Viral hepatitis (B15-B19)	827	1.6	5.0
11	Assault (homicide) (X85-Y09,Y87.1)	616	1.2	3.7
12	Influenza and pneumonia (J10-J18)	561	1.1	3.4
13	Septicemia (A40-A41)	495	0.9	3.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	376	0.7	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	245	0.5	1.5
...	All other causes (Residual)	8,487	16.0	51.1
White, both sexes, 50-54 years				
...	All causes	69,543	100.0	473.5
1	Malignant neoplasms (C00-C97)	23,735	34.1	161.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,346	23.5	111.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,347	6.3	29.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,698	3.9	18.4
5	Cerebrovascular diseases (I60-I69)	2,287	3.3	15.6
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,273	3.3	15.5
7	Diabetes mellitus (E10-E14)	2,170	3.1	14.8
8	Chronic lower respiratory diseases (J40-J47)	1,702	2.4	11.6
9	Influenza and pneumonia (J10-J18)	736	1.1	5.0
10	Septicemia (A40-A41)	724	1.0	4.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	706	1.0	4.8
12	Viral hepatitis (B15-B19)	635	0.9	4.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	560	0.8	3.8
14	Assault (homicide) (X85-Y09,Y87.1)	432	0.6	2.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	333	0.5	2.3
...	All other causes (Residual)	9,859	14.2	67.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-59 years				
...	All causes	85,840	100.0	749.8
1	Malignant neoplasms (C00-C97)	32,440	37.8	283.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,927	25.5	191.5
3	Chronic lower respiratory diseases (J40-J47)	3,469	4.0	30.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,334	3.9	29.1
5	Diabetes mellitus (E10-E14)	2,967	3.5	25.9
6	Cerebrovascular diseases (I60-I69)	2,930	3.4	25.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,497	2.9	21.8
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,604	1.9	14.0
9	Influenza and pneumonia (J10-J18)	932	1.1	8.1
10	Septicemia (A40-A41)	897	1.0	7.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	789	0.9	6.9
12	Aortic aneurysm and dissection (I71)	475	0.6	4.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	447	0.5	3.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	391	0.5	3.4
15	Viral hepatitis (B15-B19)	350	0.4	3.1
...	All other causes (Residual)	10,391	12.1	90.8
White, both sexes, 60-64 years				
...	All causes	109,701	100.0	1,197.7
1	Malignant neoplasms (C00-C97)	42,319	38.6	462.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,654	26.1	312.8
3	Chronic lower respiratory diseases (J40-J47)	6,063	5.5	66.2
4	Cerebrovascular diseases (I60-I69)	4,309	3.9	47.0
5	Diabetes mellitus (E10-E14)	3,912	3.6	42.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,941	2.7	32.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,451	2.2	26.8
8	Influenza and pneumonia (J10-J18)	1,326	1.2	14.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,220	1.1	13.3
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,171	1.1	12.8
11	Septicemia (A40-A41)	1,169	1.1	12.8
12	Aortic aneurysm and dissection (I71)	782	0.7	8.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	611	0.6	6.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	506	0.5	5.5
15	Alzheimer's disease (G30)	318	0.3	3.5
...	All other causes (Residual)	11,949	10.9	130.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 65-69 years				
...	All causes	152,597	100.0	1,871.7
1	Malignant neoplasms (C00-C97)	56,673	37.1	695.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,751	26.7	499.8
3	Chronic lower respiratory diseases (J40-J47)	10,793	7.1	132.4
4	Cerebrovascular diseases (I60-I69)	6,775	4.4	83.1
5	Diabetes mellitus (E10-E14)	5,393	3.5	66.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,855	1.9	35.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,434	1.6	29.9
8	Influenza and pneumonia (J10-J18)	2,030	1.3	24.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,966	1.3	24.1
10	Septicemia (A40-A41)	1,716	1.1	21.0
11	Aortic aneurysm and dissection (I71)	1,381	0.9	16.9
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,017	0.7	12.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	888	0.6	10.9
14	Alzheimer's disease (G30)	796	0.5	9.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	706	0.5	8.7
...	All other causes (Residual)	16,423	10.8	201.4
White, both sexes, 70-74 years				
...	All causes	224,389	100.0	2,906.9
1	Malignant neoplasms (C00-C97)	73,897	32.9	957.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62,698	27.9	812.2
3	Chronic lower respiratory diseases (J40-J47)	17,972	8.0	232.8
4	Cerebrovascular diseases (I60-I69)	12,449	5.5	161.3
5	Diabetes mellitus (E10-E14)	7,682	3.4	99.5
6	Influenza and pneumonia (J10-J18)	3,939	1.8	51.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,762	1.7	48.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,178	1.4	41.2
9	Septicemia (A40-A41)	2,702	1.2	35.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,491	1.1	32.3
11	Alzheimer's disease (G30)	2,343	1.0	30.4
12	Aortic aneurysm and dissection (I71)	2,270	1.0	29.4
13	Parkinson's disease (G20-G21)	1,397	0.6	18.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,351	0.6	17.5
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,200	0.5	15.5
...	All other causes (Residual)	25,058	11.2	324.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-79 years				
...	All causes	299,257	100.0	4,497.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	89,644	30.0	1,347.1
2	Malignant neoplasms (C00-C97)	80,805	27.0	1,214.3
3	Chronic lower respiratory diseases (J40-J47)	22,832	7.6	343.1
4	Cerebrovascular diseases (I60-I69)	21,385	7.1	321.4
5	Diabetes mellitus (E10-E14)	9,561	3.2	143.7
6	Influenza and pneumonia (J10-J18)	7,028	2.3	105.6
7	Alzheimer's disease (G30)	5,864	2.0	88.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,039	1.7	75.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,857	1.6	73.0
10	Septicemia (A40-A41)	3,859	1.3	58.0
11	Parkinson's disease (G20-G21)	3,163	1.1	47.5
12	Aortic aneurysm and dissection (I71)	2,795	0.9	42.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,223	0.7	33.4
14	Pneumonitis due to solids and liquids (J69)	2,006	0.7	30.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,962	0.7	29.5
...	All other causes (Residual)	36,234	12.1	544.5
White, both sexes, 80-84 years				
...	All causes	328,472	100.0	7,379.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	107,189	32.6	2,408.1
2	Malignant neoplasms (C00-C97)	67,321	20.5	1,512.4
3	Cerebrovascular diseases (I60-I69)	28,982	8.8	651.1
4	Chronic lower respiratory diseases (J40-J47)	22,193	6.8	498.6
5	Influenza and pneumonia (J10-J18)	10,783	3.3	242.2
6	Alzheimer's disease (G30)	10,260	3.1	230.5
7	Diabetes mellitus (E10-E14)	8,917	2.7	200.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,701	1.7	128.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,613	1.7	126.1
10	Septicemia (A40-A41)	4,364	1.3	98.0
11	Parkinson's disease (G20-G21)	4,143	1.3	93.1
12	Pneumonitis due to solids and liquids (J69)	2,942	0.9	66.1
13	Aortic aneurysm and dissection (I71)	2,603	0.8	58.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,573	0.8	57.8
15	Atherosclerosis (I70)	2,375	0.7	53.4
...	All other causes (Residual)	42,513	12.9	955.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 85-89 years				
...	All causes	307,495	100.0	12,522.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	111,070	36.1	4,523.3
2	Malignant neoplasms (C00-C97)	44,566	14.5	1,814.9
3	Cerebrovascular diseases (I60-I69)	30,998	10.1	1,262.4
4	Chronic lower respiratory diseases (J40-J47)	16,186	5.3	659.2
5	Influenza and pneumonia (J10-J18)	12,677	4.1	516.3
6	Alzheimer's disease (G30)	12,562	4.1	511.6
7	Diabetes mellitus (E10-E14)	6,905	2.2	281.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,570	1.8	226.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,390	1.8	219.5
10	Septicemia (A40-A41)	4,206	1.4	171.3
11	Pneumonitis due to solids and liquids (J69)	3,532	1.1	143.8
12	Parkinson's disease (G20-G21)	3,418	1.1	139.2
13	Atherosclerosis (I70)	2,984	1.0	121.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,772	0.9	112.9
15	Aortic aneurysm and dissection (I71)	1,988	0.6	81.0
...	All other causes (Residual)	42,671	13.9	1,737.8
White, both sexes, 90-94 years				
...	All causes	201,942	100.0	19,103.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,035	39.6	7,571.3
2	Cerebrovascular diseases (I60-I69)	21,356	10.6	2,020.3
3	Malignant neoplasms (C00-C97)	19,488	9.7	1,843.6
4	Influenza and pneumonia (J10-J18)	10,480	5.2	991.4
5	Alzheimer's disease (G30)	9,723	4.8	919.8
6	Chronic lower respiratory diseases (J40-J47)	7,360	3.6	696.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,693	1.8	349.4
8	Diabetes mellitus (E10-E14)	3,525	1.7	333.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,447	1.7	326.1
10	Atherosclerosis (I70)	2,693	1.3	254.8
11	Pneumonitis due to solids and liquids (J69)	2,562	1.3	242.4
12	Septicemia (A40-A41)	2,546	1.3	240.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,056	1.0	194.5
14	Parkinson's disease (G20-G21)	1,560	0.8	147.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,066	0.5	100.8
...	All other causes (Residual)	30,352	15.0	2,871.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 95-99 years				
...	All causes	77,167	100.0	24,507.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	32,666	42.3	10,374.6
2	Cerebrovascular diseases (I60-I69)	7,711	10.0	2,449.0
3	Malignant neoplasms (C00-C97)	4,995	6.5	1,586.4
4	Influenza and pneumonia (J10-J18)	4,728	6.1	1,501.6
5	Alzheimer's disease (G30)	3,765	4.9	1,195.7
6	Chronic lower respiratory diseases (J40-J47)	2,070	2.7	657.4
7	Atherosclerosis (I70)	1,431	1.9	454.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,317	1.7	418.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,258	1.6	399.5
10	Diabetes mellitus (E10-E14)	1,067	1.4	338.9
11	Pneumonitis due to solids and liquids (J69)	1,026	1.3	325.9
12	Septicemia (A40-A41)	875	1.1	277.9
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	856	1.1	271.9
14	Parkinson's disease (G20-G21)	350	0.5	111.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	291	0.4	92.4
...	All other causes (Residual)	12,761	16.5	4,052.8
White, both sexes, 100 years and over				
...	All causes	16,157	100.0	30,367.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,279	45.1	13,681.0
2	Cerebrovascular diseases (I60-I69)	1,397	8.6	2,625.7
3	Influenza and pneumonia (J10-J18)	1,216	7.5	2,285.5
4	Alzheimer's disease (G30)	633	3.9	1,189.7
5	Malignant neoplasms (C00-C97)	630	3.9	1,184.1
6	Atherosclerosis (I70)	383	2.4	719.9
7	Chronic lower respiratory diseases (J40-J47)	310	1.9	582.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	1.7	511.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	216	1.3	406.0
10	Pneumonitis due to solids and liquids (J69)	211	1.3	396.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	196	1.2	368.4
12	Septicemia (A40-A41)	163	1.0	306.4
13	Diabetes mellitus (E10-E14)	148	0.9	278.2
14	Nutritional deficiencies (E40-E64)	71	0.4	133.4
15	Anemias (D50-D64)	68	0.4	127.8
...	All other causes (Residual)	2,964	18.3	5,570.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,007,191	100.0	905.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	301,551	29.9	271.2
2	Malignant neoplasms (C00-C97)	247,403	24.6	222.5
3	Cerebrovascular diseases (I60-I69)	54,938	5.5	49.4
4	Chronic lower respiratory diseases (J40-J47)	54,816	5.4	49.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53,329	5.3	48.0
6	Diabetes mellitus (E10-E14)	26,009	2.6	23.4
7	Influenza and pneumonia (J10-J18)	25,002	2.5	22.5
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	21,293	2.1	19.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	15,002	1.5	13.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,385	1.4	12.9
11	Alzheimer's disease (G30)	13,524	1.3	12.2
12	Septicemia (A40-A41)	10,716	1.1	9.6
13	Aortic aneurysm and dissection (I71)	8,797	0.9	7.9
14	Parkinson's disease (G20-G21)	8,367	0.8	7.5
15	Pneumonitis due to solids and liquids (J69)	7,488	0.7	6.7
...	All other causes (Residual)	144,571	14.4	130.0
White, male, 1-4 years				
...	All causes	2,004	100.0	32.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	808	40.3	13.1
2	Malignant neoplasms (C00-C97)	190	9.5	3.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	9.0	2.9
4	Assault (homicide) (X85-Y09,Y87.1)	114	5.7	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	4.0	1.3
6	Septicemia (A40-A41)	35	1.7	0.6
7	Influenza and pneumonia (J10-J18)	30	1.5	0.5
7	Certain conditions originating in the perinatal period (P00-P96)	30	1.5	0.5
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.2	0.4
10	Meningitis (G00,G03)	14	0.7	*
10	Cerebrovascular diseases (I60-I69)	14	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	13	0.6	*
13	Meningococcal infection (A39)	8	0.4	*
13	Anemias (D50-D64)	8	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.4	*
...	All other causes (Residual)	447	22.3	7.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-9 years				
...	All causes	1,348	100.0	16.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	587	43.5	7.4
2	Malignant neoplasms (C00-C97)	215	15.9	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	5.3	0.9
4	Assault (homicide) (X85-Y09,Y87.1)	59	4.4	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.1	0.5
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.8	0.3
7	Influenza and pneumonia (J10-J18)	22	1.6	0.3
8	Septicemia (A40-A41)	19	1.4	*
9	Cerebrovascular diseases (I60-I69)	12	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	12	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.7	*
12	Anemias (D50-D64)	7	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.5	*
14	Meningitis (G00,G03)	6	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
...	All other causes (Residual)	248	18.4	3.1
White, male, 10-14 years				
...	All causes	1,907	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	791	41.5	9.9
2	Malignant neoplasms (C00-C97)	241	12.6	3.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	195	10.2	2.4
4	Assault (homicide) (X85-Y09,Y87.1)	89	4.7	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	3.9	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	3.6	0.9
7	Chronic lower respiratory diseases (J40-J47)	21	1.1	0.3
8	Influenza and pneumonia (J10-J18)	17	0.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.8	*
10	Cerebrovascular diseases (I60-I69)	14	0.7	*
11	Septicemia (A40-A41)	11	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	10	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
14	Diabetes mellitus (E10-E14)	6	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.3	*
15	Anemias (D50-D64)	5	0.3	*
...	All other causes (Residual)	335	17.6	4.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 15-19 years				
...	All causes	7,242	100.0	89.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,007	55.3	49.3
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,130	15.6	13.9
3	Assault (homicide) (X85-Y09,Y87.1)	659	9.1	8.1
4	Malignant neoplasms (C00-C97)	344	4.8	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	161	2.2	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	1.3	1.1
7	Cerebrovascular diseases (I60-I69)	33	0.5	0.4
8	Influenza and pneumonia (J10-J18)	27	0.4	0.3
9	Chronic lower respiratory diseases (J40-J47)	26	0.4	0.3
10	Diabetes mellitus (E10-E14)	22	0.3	0.3
11	Meningococcal infection (A39)	20	0.3	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.2	*
13	Septicemia (A40-A41)	15	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.2	*
14	Legal intervention (Y35,Y89.0)	11	0.2	*
...	All other causes (Residual)	665	9.2	8.2
White, male, 20-24 years				
...	All causes	9,626	100.0	127.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,763	49.5	63.1
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,723	17.9	22.8
3	Assault (homicide) (X85-Y09,Y87.1)	924	9.6	12.2
4	Malignant neoplasms (C00-C97)	454	4.7	6.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	252	2.6	3.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	1.1	1.4
7	Influenza and pneumonia (J10-J18)	52	0.5	0.7
8	Cerebrovascular diseases (I60-I69)	51	0.5	0.7
9	Diabetes mellitus (E10-E14)	42	0.4	0.6
10	Chronic lower respiratory diseases (J40-J47)	39	0.4	0.5
11	Legal intervention (Y35,Y89.0)	31	0.3	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	30	0.3	0.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.3	0.4
14	Septicemia (A40-A41)	24	0.2	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.2	0.3
...	All other causes (Residual)	1,081	11.2	14.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-29 years				
...	All causes	8,943	100.0	125.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,704	41.4	52.1
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,631	18.2	22.9
3	Assault (homicide) (X85-Y09,Y87.1)	721	8.1	10.1
4	Malignant neoplasms (C00-C97)	559	6.3	7.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	418	4.7	5.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	158	1.8	2.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	93	1.0	1.3
8	Cerebrovascular diseases (I60-I69)	89	1.0	1.3
9	Diabetes mellitus (E10-E14)	76	0.8	1.1
10	Influenza and pneumonia (J10-J18)	51	0.6	0.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	46	0.5	0.6
12	Chronic lower respiratory diseases (J40-J47)	36	0.4	0.5
13	Legal intervention (Y35,Y89.0)	35	0.4	0.5
14	Septicemia (A40-A41)	34	0.4	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.3	0.4
...	All other causes (Residual)	1,264	14.1	17.8
White, male, 30-34 years				
...	All causes	11,197	100.0	142.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,665	32.7	46.5
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,732	15.5	22.0
3	Malignant neoplasms (C00-C97)	941	8.4	11.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	898	8.0	11.4
5	Assault (homicide) (X85-Y09,Y87.1)	635	5.7	8.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	591	5.3	7.5
7	Diabetes mellitus (E10-E14)	191	1.7	2.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	178	1.6	2.3
9	Cerebrovascular diseases (I60-I69)	136	1.2	1.7
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	1.2	1.7
11	Influenza and pneumonia (J10-J18)	87	0.8	1.1
12	Septicemia (A40-A41)	61	0.5	0.8
13	Chronic lower respiratory diseases (J40-J47)	48	0.4	0.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.4	0.6
...	All other causes (Residual)	1,807	16.1	22.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 35-39 years				
...	All causes	17,529	100.0	191.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,608	26.3	50.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,197	12.5	24.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,185	12.5	23.9
4	Malignant neoplasms (C00-C97)	1,907	10.9	20.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,165	6.6	12.7
6	Assault (homicide) (X85-Y09,Y87.1)	614	3.5	6.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	588	3.4	6.4
8	Diabetes mellitus (E10-E14)	285	1.6	3.1
9	Cerebrovascular diseases (I60-I69)	282	1.6	3.1
10	Influenza and pneumonia (J10-J18)	161	0.9	1.8
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	0.7	1.4
12	Viral hepatitis (B15-B19)	123	0.7	1.3
13	Septicemia (A40-A41)	110	0.6	1.2
14	Chronic lower respiratory diseases (J40-J47)	85	0.5	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	0.5	0.9
...	All other causes (Residual)	3,012	17.2	32.9
White, male, 40-44 years				
...	All causes	25,849	100.0	275.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,854	18.8	51.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,835	18.7	51.6
3	Malignant neoplasms (C00-C97)	3,832	14.8	40.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,423	9.4	25.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,333	5.2	14.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,135	4.4	12.1
7	Assault (homicide) (X85-Y09,Y87.1)	588	2.3	6.3
8	Diabetes mellitus (E10-E14)	581	2.2	6.2
9	Cerebrovascular diseases (I60-I69)	560	2.2	6.0
10	Viral hepatitis (B15-B19)	316	1.2	3.4
11	Influenza and pneumonia (J10-J18)	294	1.1	3.1
12	Chronic lower respiratory diseases (J40-J47)	207	0.8	2.2
13	Septicemia (A40-A41)	201	0.8	2.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	0.6	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	0.5	1.4
...	All other causes (Residual)	4,395	17.0	46.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-49 years				
...	All causes	34,599	100.0	418.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,267	23.9	100.1
2	Malignant neoplasms (C00-C97)	7,292	21.1	88.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,143	12.0	50.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,160	6.2	26.2
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	2,128	6.2	25.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	977	2.8	11.8
7	Cerebrovascular diseases (I60-I69)	894	2.6	10.8
8	Diabetes mellitus (E10-E14)	808	2.3	9.8
9	Viral hepatitis (B15-B19)	636	1.8	7.7
10	Chronic lower respiratory diseases (J40-J47)	472	1.4	5.7
11	Assault (homicide) (X85-Y09,Y87.1)	423	1.2	5.1
12	Influenza and pneumonia (J10-J18)	325	0.9	3.9
13	Septicemia (A40-A41)	289	0.8	3.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	237	0.7	2.9
15	Aortic aneurysm and dissection (I71)	147	0.4	1.8
...	All other causes (Residual)	5,401	15.6	65.4
White, male, 50-54 years				
...	All causes	43,267	100.0	598.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,213	28.2	168.9
2	Malignant neoplasms (C00-C97)	12,061	27.9	166.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,179	7.3	44.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,045	4.7	28.3
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,706	3.9	23.6
6	Diabetes mellitus (E10-E14)	1,269	2.9	17.6
7	Cerebrovascular diseases (I60-I69)	1,244	2.9	17.2
8	Chronic lower respiratory diseases (J40-J47)	849	2.0	11.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	613	1.4	8.5
10	Viral hepatitis (B15-B19)	487	1.1	6.7
11	Influenza and pneumonia (J10-J18)	434	1.0	6.0
12	Septicemia (A40-A41)	404	0.9	5.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	321	0.7	4.4
14	Assault (homicide) (X85-Y09,Y87.1)	295	0.7	4.1
15	Aortic aneurysm and dissection (I71)	237	0.5	3.3
...	All other causes (Residual)	5,910	13.7	81.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 55-59 years				
...	All causes	52,048	100.0	936.0
1	Malignant neoplasms (C00-C97)	17,275	33.2	310.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,794	30.3	284.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,297	4.4	41.3
4	Chronic lower respiratory diseases (J40-J47)	1,791	3.4	32.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,775	3.4	31.9
6	Diabetes mellitus (E10-E14)	1,662	3.2	29.9
7	Cerebrovascular diseases (I60-I69)	1,607	3.1	28.9
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,225	2.4	22.0
9	Influenza and pneumonia (J10-J18)	536	1.0	9.6
10	Septicemia (A40-A41)	474	0.9	8.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	444	0.9	8.0
12	Aortic aneurysm and dissection (I71)	364	0.7	6.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	336	0.6	6.0
14	Viral hepatitis (B15-B19)	246	0.5	4.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	240	0.5	4.3
...	All other causes (Residual)	5,982	11.5	107.6
White, male, 60-64 years				
...	All causes	65,066	100.0	1,484.5
1	Malignant neoplasms (C00-C97)	23,182	35.6	528.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,628	30.2	447.8
3	Chronic lower respiratory diseases (J40-J47)	3,102	4.8	70.8
4	Cerebrovascular diseases (I60-I69)	2,391	3.7	54.5
5	Diabetes mellitus (E10-E14)	2,161	3.3	49.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,945	3.0	44.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,700	2.6	38.8
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	908	1.4	20.7
9	Influenza and pneumonia (J10-J18)	766	1.2	17.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	662	1.0	15.1
11	Septicemia (A40-A41)	623	1.0	14.2
12	Aortic aneurysm and dissection (I71)	592	0.9	13.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	324	0.5	7.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	291	0.4	6.6
15	Viral hepatitis (B15-B19)	186	0.3	4.2
...	All other causes (Residual)	6,605	10.2	150.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-69 years				
...	All causes	88,182	100.0	2,328.7
1	Malignant neoplasms (C00-C97)	31,634	35.9	835.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,246	29.8	693.1
3	Chronic lower respiratory diseases (J40-J47)	5,523	6.3	145.8
4	Cerebrovascular diseases (I60-I69)	3,540	4.0	93.5
5	Diabetes mellitus (E10-E14)	2,849	3.2	75.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,770	2.0	46.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,553	1.8	41.0
8	Influenza and pneumonia (J10-J18)	1,145	1.3	30.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,071	1.2	28.3
10	Aortic aneurysm and dissection (I71)	1,006	1.1	26.6
11	Septicemia (A40-A41)	892	1.0	23.6
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	823	0.9	21.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	517	0.6	13.7
14	Alzheimer's disease (G30)	389	0.4	10.3
15	Parkinson's disease (G20-G21)	372	0.4	9.8
...	All other causes (Residual)	8,852	10.0	233.8
White, male, 70-74 years				
...	All causes	125,197	100.0	3,632.1
1	Malignant neoplasms (C00-C97)	40,650	32.5	1,179.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,218	30.5	1,108.8
3	Chronic lower respiratory diseases (J40-J47)	9,263	7.4	268.7
4	Cerebrovascular diseases (I60-I69)	6,288	5.0	182.4
5	Diabetes mellitus (E10-E14)	3,890	3.1	112.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,207	1.8	64.0
7	Influenza and pneumonia (J10-J18)	2,169	1.7	62.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,698	1.4	49.3
9	Aortic aneurysm and dissection (I71)	1,527	1.2	44.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,453	1.2	42.2
11	Septicemia (A40-A41)	1,372	1.1	39.8
12	Alzheimer's disease (G30)	1,043	0.8	30.3
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	961	0.8	27.9
14	Parkinson's disease (G20-G21)	890	0.7	25.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	730	0.6	21.2
...	All other causes (Residual)	12,838	10.3	372.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 75-79 years				
...	All causes	154,451	100.0	5,521.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,669	31.5	1,739.7
2	Malignant neoplasms (C00-C97)	42,849	27.7	1,531.7
3	Chronic lower respiratory diseases (J40-J47)	11,418	7.4	408.1
4	Cerebrovascular diseases (I60-I69)	9,746	6.3	348.4
5	Diabetes mellitus (E10-E14)	4,508	2.9	161.1
6	Influenza and pneumonia (J10-J18)	3,623	2.3	129.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,712	1.8	96.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,567	1.7	91.8
9	Alzheimer's disease (G30)	2,271	1.5	81.2
10	Parkinson's disease (G20-G21)	1,943	1.3	69.5
11	Septicemia (A40-A41)	1,833	1.2	65.5
12	Aortic aneurysm and dissection (I71)	1,747	1.1	62.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,162	0.8	41.5
14	Pneumonitis due to solids and liquids (J69)	1,139	0.7	40.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,032	0.7	36.9
...	All other causes (Residual)	17,232	11.2	616.0
White, male, 80-84 years				
...	All causes	151,919	100.0	8,956.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,626	33.3	2,984.7
2	Malignant neoplasms (C00-C97)	33,865	22.3	1,996.5
3	Cerebrovascular diseases (I60-I69)	11,464	7.5	675.9
4	Chronic lower respiratory diseases (J40-J47)	10,836	7.1	638.8
5	Influenza and pneumonia (J10-J18)	5,119	3.4	301.8
6	Diabetes mellitus (E10-E14)	3,911	2.6	230.6
7	Alzheimer's disease (G30)	3,583	2.4	211.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,786	1.8	164.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,781	1.8	164.0
10	Parkinson's disease (G20-G21)	2,364	1.6	139.4
11	Septicemia (A40-A41)	1,904	1.3	112.3
12	Pneumonitis due to solids and liquids (J69)	1,607	1.1	94.7
13	Aortic aneurysm and dissection (I71)	1,480	1.0	87.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,136	0.7	67.0
15	Atherosclerosis (I70)	1,008	0.7	59.4
...	All other causes (Residual)	17,449	11.5	1,028.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85-89 years				
...	All causes	118,321	100.0	14,759.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,311	35.8	5,278.1
2	Malignant neoplasms (C00-C97)	20,503	17.3	2,557.6
3	Cerebrovascular diseases (I60-I69)	9,892	8.4	1,234.0
4	Chronic lower respiratory diseases (J40-J47)	7,340	6.2	915.6
5	Influenza and pneumonia (J10-J18)	5,128	4.3	639.7
6	Alzheimer's disease (G30)	3,466	2.9	432.4
7	Diabetes mellitus (E10-E14)	2,473	2.1	308.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,400	2.0	299.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,398	2.0	299.1
10	Parkinson's disease (G20-G21)	1,804	1.5	225.0
11	Pneumonitis due to solids and liquids (J69)	1,720	1.5	214.6
12	Septicemia (A40-A41)	1,448	1.2	180.6
13	Atherosclerosis (I70)	1,015	0.9	126.6
13	Aortic aneurysm and dissection (I71)	1,015	0.9	126.6
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	839	0.7	104.7
...	All other causes (Residual)	14,569	12.3	1,817.4
White, male, 90-94 years				
...	All causes	59,009	100.0	20,848.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,657	38.4	8,004.9
2	Malignant neoplasms (C00-C97)	7,611	12.9	2,689.0
3	Cerebrovascular diseases (I60-I69)	5,095	8.6	1,800.1
4	Influenza and pneumonia (J10-J18)	3,415	5.8	1,206.5
5	Chronic lower respiratory diseases (J40-J47)	2,938	5.0	1,038.0
6	Alzheimer's disease (G30)	1,974	3.3	697.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,320	2.2	466.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,290	2.2	455.8
9	Pneumonitis due to solids and liquids (J69)	1,028	1.7	363.2
10	Diabetes mellitus (E10-E14)	1,023	1.7	361.4
11	Septicemia (A40-A41)	692	1.2	244.5
12	Parkinson's disease (G20-G21)	689	1.2	243.4
13	Atherosclerosis (I70)	687	1.2	242.7
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	483	0.8	170.6
15	Aortic aneurysm and dissection (I71)	389	0.7	137.4
...	All other causes (Residual)	7,718	13.1	2,726.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 95-99 years				
...	All causes	16,624	100.0	24,296.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,708	40.4	9,804.0
2	Malignant neoplasms (C00-C97)	1,600	9.6	2,338.5
3	Cerebrovascular diseases (I60-I69)	1,367	8.2	1,997.9
4	Influenza and pneumonia (J10-J18)	1,261	7.6	1,843.0
5	Chronic lower respiratory diseases (J40-J47)	702	4.2	1,026.0
6	Alzheimer's disease (G30)	525	3.2	767.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	387	2.3	565.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	357	2.1	521.8
9	Pneumonitis due to solids and liquids (J69)	341	2.1	498.4
10	Atherosclerosis (I70)	261	1.6	381.5
11	Diabetes mellitus (E10-E14)	228	1.4	333.2
12	Septicemia (A40-A41)	161	1.0	235.3
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	123	0.7	179.8
14	Parkinson's disease (G20-G21)	109	0.7	159.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.5	119.8
...	All other causes (Residual)	2,412	14.5	3,525.2
White, male, 100 years and over				
...	All causes	2,455	100.0	26,563.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,059	43.1	11,458.6
2	Influenza and pneumonia (J10-J18)	241	9.8	2,607.7
3	Cerebrovascular diseases (I60-I69)	172	7.0	1,861.1
4	Malignant neoplasms (C00-C97)	149	6.1	1,612.2
5	Chronic lower respiratory diseases (J40-J47)	78	3.2	844.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	2.7	714.1
7	Pneumonitis due to solids and liquids (J69)	60	2.4	649.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	2.2	573.5
9	Atherosclerosis (I70)	48	2.0	519.4
10	Alzheimer's disease (G30)	45	1.8	486.9
11	Septicemia (A40-A41)	25	1.0	270.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	24	1.0	259.7
13	Diabetes mellitus (E10-E14)	16	0.7	*
14	Anemias (D50-D64)	13	0.5	*
15	Nutritional deficiencies (E40-E64)	11	0.4	*
...	All other causes (Residual)	395	16.1	4,274.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, all ages				
...	All causes	1,064,096	100.0	924.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	320,168	30.1	278.3
2	Malignant neoplasms (C00-C97)	232,608	21.9	202.2
3	Cerebrovascular diseases (I60-I69)	89,642	8.4	77.9
4	Chronic lower respiratory diseases (J40-J47)	58,024	5.5	50.4
5	Alzheimer's disease (G30)	32,936	3.1	28.6
6	Influenza and pneumonia (J10-J18)	32,912	3.1	28.6
7	Diabetes mellitus (E10-E14)	29,552	2.8	25.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,263	2.8	25.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,213	1.4	13.2
10	Septicemia (A40-A41)	14,088	1.3	12.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	8,655	0.8	7.5
12	Atherosclerosis (I70)	8,307	0.8	7.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,980	0.7	6.9
14	Pneumonitis due to solids and liquids (J69)	7,381	0.7	6.4
15	Parkinson's disease (G20-G21)	6,575	0.6	5.7
...	All other causes (Residual)	170,792	16.1	148.4
White, female, 1-4 years				
...	All causes	1,490	100.0	25.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	527	35.4	9.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	11.2	2.8
3	Malignant neoplasms (C00-C97)	140	9.4	2.4
4	Assault (homicide) (X85-Y09,Y87.1)	85	5.7	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	3.1	0.8
6	Influenza and pneumonia (J10-J18)	39	2.6	0.7
7	Septicemia (A40-A41)	31	2.1	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.3	*
9	Cerebrovascular diseases (I60-I69)	16	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	16	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	15	1.0	*
12	Meningitis (G00,G03)	9	0.6	*
13	Anemias (D50-D64)	7	0.5	*
14	Meningococcal infection (A39)	6	0.4	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (Residual)	362	24.3	6.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 5-9 years				
...	All causes	1,011	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	40.5	5.4
2	Malignant neoplasms (C00-C97)	182	18.0	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	7.4	1.0
4	Assault (homicide) (X85-Y09,Y87.1)	39	3.9	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	2.9	0.4
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	2.2	0.3
7	Influenza and pneumonia (J10-J18)	14	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
9	Septicemia (A40-A41)	9	0.9	*
9	Cerebrovascular diseases (I60-I69)	9	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.8	*
12	Meningitis (G00,G03)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
15	Meningococcal infection (A39)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (Residual)	184	18.2	2.4
White, female, 10-14 years				
...	All causes	1,184	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	432	36.5	5.7
2	Malignant neoplasms (C00-C97)	180	15.2	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	5.9	0.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	53	4.5	0.7
4	Assault (homicide) (X85-Y09,Y87.1)	53	4.5	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.1	0.5
7	Cerebrovascular diseases (I60-I69)	20	1.7	0.3
8	Chronic lower respiratory diseases (J40-J47)	19	1.6	*
9	Influenza and pneumonia (J10-J18)	17	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	1.3	*
11	Septicemia (A40-A41)	10	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.7	*
13	Anemias (D50-D64)	7	0.6	*
13	Diseases of appendix (K35-K38)	7	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
...	All other causes (Residual)	249	21.0	3.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-19 years				
...	All causes	3,031	100.0	39.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,724	56.9	22.6
2	Malignant neoplasms (C00-C97)	232	7.7	3.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	224	7.4	2.9
4	Assault (homicide) (X85-Y09,Y87.1)	164	5.4	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	3.4	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	2.2	0.9
7	Influenza and pneumonia (J10-J18)	21	0.7	0.3
8	Cerebrovascular diseases (I60-I69)	18	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	14	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
10	Diabetes mellitus (E10-E14)	11	0.4	*
10	Pregnancy, childbirth and the puerperium (O00-099)	11	0.4	*
13	Meningococcal infection (A39)	9	0.3	*
13	Septicemia (A40-A41)	9	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.3	*
...	All other causes (Residual)	405	13.4	5.3
White, female, 20-24 years				
...	All causes	3,119	100.0	43.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,329	42.6	18.6
2	Malignant neoplasms (C00-C97)	303	9.7	4.2
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	238	7.6	3.3
4	Assault (homicide) (X85-Y09,Y87.1)	234	7.5	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	140	4.5	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	2.1	0.9
7	Cerebrovascular diseases (I60-I69)	46	1.5	0.6
8	Pregnancy, childbirth and the puerperium (O00-099)	44	1.4	0.6
9	Influenza and pneumonia (J10-J18)	38	1.2	0.5
10	Diabetes mellitus (E10-E14)	30	1.0	0.4
11	Chronic lower respiratory diseases (J40-J47)	24	0.8	0.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.7	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.4	*
14	Septicemia (A40-A41)	12	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.3	*
...	All other causes (Residual)	571	18.3	8.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 25-29 years				
...	All causes	3,484	100.0	49.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,032	29.6	14.7
2	Malignant neoplasms (C00-C97)	522	15.0	7.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	310	8.9	4.4
4	Assault (homicide) (X85-Y09,Y87.1)	224	6.4	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	219	6.3	3.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	77	2.2	1.1
7	Cerebrovascular diseases (I60-I69)	71	2.0	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	2.0	1.0
9	Pregnancy, childbirth and the puerperium (O00-099)	53	1.5	0.8
10	Diabetes mellitus (E10-E14)	52	1.5	0.7
11	Influenza and pneumonia (J10-J18)	47	1.3	0.7
12	Chronic lower respiratory diseases (J40-J47)	31	0.9	0.4
13	Septicemia (A40-A41)	27	0.8	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.7	0.3
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.6	0.3
...	All other causes (Residual)	706	20.3	10.0
White, female, 30-34 years				
...	All causes	5,095	100.0	64.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,118	21.9	14.2
2	Malignant neoplasms (C00-C97)	1,050	20.6	13.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	411	8.1	5.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	402	7.9	5.1
5	Assault (homicide) (X85-Y09,Y87.1)	234	4.6	3.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	157	3.1	2.0
7	Cerebrovascular diseases (I60-I69)	124	2.4	1.6
8	Diabetes mellitus (E10-E14)	101	2.0	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	1.8	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	77	1.5	1.0
11	Influenza and pneumonia (J10-J18)	76	1.5	1.0
12	Pregnancy, childbirth and the puerperium (O00-099)	55	1.1	0.7
13	Chronic lower respiratory diseases (J40-J47)	51	1.0	0.6
14	Septicemia (A40-A41)	43	0.8	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.8	0.5
...	All other causes (Residual)	1,061	20.8	13.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-39 years				
...	All causes	9,104	100.0	100.5
1	Malignant neoplasms (C00-C97)	2,409	26.5	26.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,563	17.2	17.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	898	9.9	9.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	599	6.6	6.6
5	Assault (homicide) (X85-Y09,Y87.1)	295	3.2	3.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	284	3.1	3.1
7	Cerebrovascular diseases (I60-I69)	276	3.0	3.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	237	2.6	2.6
9	Diabetes mellitus (E10-E14)	208	2.3	2.3
10	Influenza and pneumonia (J10-J18)	144	1.6	1.6
11	Chronic lower respiratory diseases (J40-J47)	109	1.2	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	1.1	1.1
13	Septicemia (A40-A41)	94	1.0	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	0.7	0.7
15	Viral hepatitis (B15-B19)	57	0.6	0.6
...	All other causes (Residual)	1,771	19.5	19.6
White, female, 40-44 years				
...	All causes	14,014	100.0	150.4
1	Malignant neoplasms (C00-C97)	4,616	32.9	49.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,638	11.7	17.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,614	11.5	17.3
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	723	5.2	7.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	540	3.9	5.8
6	Cerebrovascular diseases (I60-I69)	527	3.8	5.7
7	Diabetes mellitus (E10-E14)	345	2.5	3.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	248	1.8	2.7
9	Assault (homicide) (X85-Y09,Y87.1)	228	1.6	2.4
10	Chronic lower respiratory diseases (J40-J47)	223	1.6	2.4
11	Septicemia (A40-A41)	173	1.2	1.9
12	Influenza and pneumonia (J10-J18)	168	1.2	1.8
13	Viral hepatitis (B15-B19)	129	0.9	1.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	0.7	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	0.7	1.1
...	All other causes (Residual)	2,633	18.8	28.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 45-49 years				
...	All causes	18,532	100.0	221.6
1	Malignant neoplasms (C00-C97)	7,451	40.2	89.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,378	12.8	28.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,411	7.6	16.9
4	Cerebrovascular diseases (I60-I69)	733	4.0	8.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	657	3.5	7.9
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	634	3.4	7.6
7	Diabetes mellitus (E10-E14)	536	2.9	6.4
8	Chronic lower respiratory diseases (J40-J47)	410	2.2	4.9
9	Influenza and pneumonia (J10-J18)	236	1.3	2.8
10	Septicemia (A40-A41)	206	1.1	2.5
11	Assault (homicide) (X85-Y09,Y87.1)	193	1.0	2.3
12	Viral hepatitis (B15-B19)	191	1.0	2.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	173	0.9	2.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	0.8	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	0.6	1.4
...	All other causes (Residual)	3,071	16.6	36.7
White, female, 50-54 years				
...	All causes	26,276	100.0	352.3
1	Malignant neoplasms (C00-C97)	11,674	44.4	156.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,133	15.7	55.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,168	4.4	15.7
4	Cerebrovascular diseases (I60-I69)	1,043	4.0	14.0
5	Diabetes mellitus (E10-E14)	901	3.4	12.1
6	Chronic lower respiratory diseases (J40-J47)	853	3.2	11.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	653	2.5	8.8
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	567	2.2	7.6
9	Septicemia (A40-A41)	320	1.2	4.3
10	Influenza and pneumonia (J10-J18)	302	1.1	4.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	239	0.9	3.2
12	Viral hepatitis (B15-B19)	148	0.6	2.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	141	0.5	1.9
14	Assault (homicide) (X85-Y09,Y87.1)	137	0.5	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	0.5	1.7
...	All other causes (Residual)	3,871	14.7	51.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-59 years				
...	All causes	33,792	100.0	574.0
1	Malignant neoplasms (C00-C97)	15,165	44.9	257.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,133	18.1	104.2
3	Chronic lower respiratory diseases (J40-J47)	1,678	5.0	28.5
4	Cerebrovascular diseases (I60-I69)	1,323	3.9	22.5
5	Diabetes mellitus (E10-E14)	1,305	3.9	22.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,037	3.1	17.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	722	2.1	12.3
8	Septicemia (A40-A41)	423	1.3	7.2
9	Influenza and pneumonia (J10-J18)	396	1.2	6.7
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	379	1.1	6.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	345	1.0	5.9
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	207	0.6	3.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	141	0.4	2.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	133	0.4	2.3
15	Aortic aneurysm and dissection (I71)	111	0.3	1.9
...	All other causes (Residual)	4,294	12.7	72.9
White, female, 60-64 years				
...	All causes	44,635	100.0	934.5
1	Malignant neoplasms (C00-C97)	19,137	42.9	400.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,026	20.2	189.0
3	Chronic lower respiratory diseases (J40-J47)	2,961	6.6	62.0
4	Cerebrovascular diseases (I60-I69)	1,918	4.3	40.2
5	Diabetes mellitus (E10-E14)	1,751	3.9	36.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	996	2.2	20.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	751	1.7	15.7
8	Influenza and pneumonia (J10-J18)	560	1.3	11.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	558	1.3	11.7
10	Septicemia (A40-A41)	546	1.2	11.4
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	287	0.6	6.0
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	263	0.6	5.5
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	215	0.5	4.5
14	Aortic aneurysm and dissection (I71)	190	0.4	4.0
15	Alzheimer's disease (G30)	180	0.4	3.8
...	All other causes (Residual)	5,296	11.9	110.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 65-69 years				
...	All causes	64,415	100.0	1,475.3
1	Malignant neoplasms (C00-C97)	25,039	38.9	573.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,505	22.5	332.2
3	Chronic lower respiratory diseases (J40-J47)	5,270	8.2	120.7
4	Cerebrovascular diseases (I60-I69)	3,235	5.0	74.1
5	Diabetes mellitus (E10-E14)	2,544	3.9	58.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,085	1.7	24.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	895	1.4	20.5
8	Influenza and pneumonia (J10-J18)	885	1.4	20.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	881	1.4	20.2
10	Septicemia (A40-A41)	824	1.3	18.9
11	Alzheimer's disease (G30)	407	0.6	9.3
12	Aortic aneurysm and dissection (I71)	375	0.6	8.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	371	0.6	8.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	366	0.6	8.4
15	Parkinson's disease (G20-G21)	213	0.3	4.9
...	All other causes (Residual)	7,520	11.7	172.2
White, female, 70-74 years				
...	All causes	99,192	100.0	2,321.8
1	Malignant neoplasms (C00-C97)	33,247	33.5	778.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,480	24.7	573.0
3	Chronic lower respiratory diseases (J40-J47)	8,709	8.8	203.9
4	Cerebrovascular diseases (I60-I69)	6,161	6.2	144.2
5	Diabetes mellitus (E10-E14)	3,792	3.8	88.8
6	Influenza and pneumonia (J10-J18)	1,770	1.8	41.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,555	1.6	36.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,480	1.5	34.6
9	Septicemia (A40-A41)	1,330	1.3	31.1
10	Alzheimer's disease (G30)	1,300	1.3	30.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,038	1.0	24.3
12	Aortic aneurysm and dissection (I71)	743	0.7	17.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	621	0.6	14.5
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	614	0.6	14.4
15	Parkinson's disease (G20-G21)	507	0.5	11.9
...	All other causes (Residual)	11,845	11.9	277.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-79 years				
...	All causes	144,806	100.0	3,754.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,975	28.3	1,062.4
2	Malignant neoplasms (C00-C97)	37,956	26.2	984.1
3	Cerebrovascular diseases (I60-I69)	11,639	8.0	301.8
4	Chronic lower respiratory diseases (J40-J47)	11,414	7.9	295.9
5	Diabetes mellitus (E10-E14)	5,053	3.5	131.0
6	Alzheimer's disease (G30)	3,593	2.5	93.2
7	Influenza and pneumonia (J10-J18)	3,405	2.4	88.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,327	1.6	60.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,290	1.6	59.4
10	Septicemia (A40-A41)	2,026	1.4	52.5
11	Parkinson's disease (G20-G21)	1,220	0.8	31.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,102	0.8	28.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,061	0.7	27.5
14	Aortic aneurysm and dissection (I71)	1,048	0.7	27.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	930	0.6	24.1
...	All other causes (Residual)	18,767	13.0	486.6
White, female, 80-84 years				
...	All causes	176,553	100.0	6,408.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	56,563	32.0	2,053.1
2	Malignant neoplasms (C00-C97)	33,456	18.9	1,214.4
3	Cerebrovascular diseases (I60-I69)	17,518	9.9	635.9
4	Chronic lower respiratory diseases (J40-J47)	11,357	6.4	412.2
5	Alzheimer's disease (G30)	6,677	3.8	242.4
6	Influenza and pneumonia (J10-J18)	5,664	3.2	205.6
7	Diabetes mellitus (E10-E14)	5,006	2.8	181.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,920	1.7	106.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,827	1.6	102.6
10	Septicemia (A40-A41)	2,460	1.4	89.3
11	Parkinson's disease (G20-G21)	1,779	1.0	64.6
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,613	0.9	58.5
13	Atherosclerosis (I70)	1,367	0.8	49.6
14	Pneumonitis due to solids and liquids (J69)	1,335	0.8	48.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,204	0.7	43.7
...	All other causes (Residual)	24,807	14.1	900.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 85-89 years				
...	All causes	189,174	100.0	11,438.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,759	36.3	4,157.5
2	Malignant neoplasms (C00-C97)	24,063	12.7	1,455.0
3	Cerebrovascular diseases (I60-I69)	21,106	11.2	1,276.2
4	Alzheimer's disease (G30)	9,096	4.8	550.0
5	Chronic lower respiratory diseases (J40-J47)	8,846	4.7	534.9
6	Influenza and pneumonia (J10-J18)	7,549	4.0	456.4
7	Diabetes mellitus (E10-E14)	4,432	2.3	268.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,170	1.7	191.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,992	1.6	180.9
10	Septicemia (A40-A41)	2,758	1.5	166.8
11	Atherosclerosis (I70)	1,969	1.0	119.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,933	1.0	116.9
13	Pneumonitis due to solids and liquids (J69)	1,812	1.0	109.6
14	Parkinson's disease (G20-G21)	1,614	0.9	97.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,142	0.6	69.1
...	All other causes (Residual)	27,933	14.8	1,689.0
White, female, 90-94 years				
...	All causes	142,933	100.0	18,465.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,378	40.1	7,412.8
2	Cerebrovascular diseases (I60-I69)	16,261	11.4	2,100.8
3	Malignant neoplasms (C00-C97)	11,877	8.3	1,534.4
4	Alzheimer's disease (G30)	7,749	5.4	1,001.1
5	Influenza and pneumonia (J10-J18)	7,065	4.9	912.7
6	Chronic lower respiratory diseases (J40-J47)	4,422	3.1	571.3
7	Diabetes mellitus (E10-E14)	2,502	1.8	323.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,403	1.7	310.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,127	1.5	274.8
10	Atherosclerosis (I70)	2,006	1.4	259.2
11	Septicemia (A40-A41)	1,854	1.3	239.5
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,573	1.1	203.2
13	Pneumonitis due to solids and liquids (J69)	1,534	1.1	198.2
14	Parkinson's disease (G20-G21)	871	0.6	112.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	725	0.5	93.7
...	All other causes (Residual)	22,586	15.8	2,917.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 95-99 years				
...	All causes	60,543	100.0	24,566.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,958	42.9	10,533.0
2	Cerebrovascular diseases (I60-I69)	6,344	10.5	2,574.2
3	Influenza and pneumonia (J10-J18)	3,467	5.7	1,406.8
4	Malignant neoplasms (C00-C97)	3,395	5.6	1,377.6
5	Alzheimer's disease (G30)	3,240	5.4	1,314.7
6	Chronic lower respiratory diseases (J40-J47)	1,368	2.3	555.1
7	Atherosclerosis (I70)	1,170	1.9	474.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	960	1.6	389.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	871	1.4	353.4
10	Diabetes mellitus (E10-E14)	839	1.4	340.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	733	1.2	297.4
12	Septicemia (A40-A41)	714	1.2	289.7
13	Pneumonitis due to solids and liquids (J69)	685	1.1	278.0
14	Parkinson's disease (G20-G21)	241	0.4	97.8
15	Nutritional deficiencies (E40-E64)	214	0.4	86.8
...	All other causes (Residual)	10,344	17.1	4,197.3
White, female, 100 years and over				
...	All causes	13,702	100.0	31,167.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,220	45.4	14,148.3
2	Cerebrovascular diseases (I60-I69)	1,225	8.9	2,786.4
3	Influenza and pneumonia (J10-J18)	975	7.1	2,217.8
4	Alzheimer's disease (G30)	588	4.3	1,337.5
5	Malignant neoplasms (C00-C97)	481	3.5	1,094.1
6	Atherosclerosis (I70)	335	2.4	762.0
7	Chronic lower respiratory diseases (J40-J47)	232	1.7	527.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	1.6	498.1
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	172	1.3	391.2
10	Pneumonitis due to solids and liquids (J69)	151	1.1	343.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	150	1.1	341.2
12	Septicemia (A40-A41)	138	1.0	313.9
13	Diabetes mellitus (E10-E14)	132	1.0	300.3
14	Nutritional deficiencies (E40-E64)	60	0.4	136.5
15	Anemias (D50-D64)	55	0.4	125.1
...	All other causes (Residual)	2,569	18.7	5,843.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	285,826	100.0	809.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,523	27.1	219.6
2	Malignant neoplasms (C00-C97)	61,945	21.7	175.5
3	Cerebrovascular diseases (I60-I69)	19,221	6.7	54.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,277	4.3	34.8
5	Diabetes mellitus (E10-E14)	12,021	4.2	34.1
6	Assault (homicide) (X85-Y09,Y87.1)	7,867	2.8	22.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7,848	2.7	22.2
8	Chronic lower respiratory diseases (J40-J47)	7,607	2.7	21.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,911	2.4	19.6
10	Influenza and pneumonia (J10-J18)	5,990	2.1	17.0
11	Septicemia (A40-A41)	5,928	2.1	16.8
12	Certain conditions originating in the perinatal period (P00-P96)	5,021	1.8	14.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3,887	1.4	11.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2,737	1.0	7.8
15	Alzheimer's disease (G30)	2,728	1.0	7.7
...	All other causes (Residual)	46,315	16.2	131.2
Black, both sexes, 1-4 years				
...	All causes	1,248	100.0	56.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	405	32.5	18.2
2	Assault (homicide) (X85-Y09,Y87.1)	145	11.6	6.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	10.1	5.7
4	Malignant neoplasms (C00-C97)	70	5.6	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	3.9	2.2
6	Influenza and pneumonia (J10-J18)	30	2.4	1.3
7	Certain conditions originating in the perinatal period (P00-P96)	27	2.2	1.2
8	Septicemia (A40-A41)	26	2.1	1.2
9	Chronic lower respiratory diseases (J40-J47)	21	1.7	0.9
10	Anemias (D50-D64)	19	1.5	*
11	Cerebrovascular diseases (I60-I69)	14	1.1	*
12	Meningitis (G00,G03)	13	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.6	*
...	All other causes (Residual)	274	22.0	12.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-9 years				
...	All causes	756	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	44.7	10.9
2	Malignant neoplasms (C00-C97)	72	9.5	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	5.8	1.4
4	Assault (homicide) (X85-Y09,Y87.1)	40	5.3	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	4.0	1.0
6	Chronic lower respiratory diseases (J40-J47)	24	3.2	0.8
7	Anemias (D50-D64)	15	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.2	*
10	Septicemia (A40-A41)	8	1.1	*
11	Meningococcal infection (A39)	7	0.9	*
11	Influenza and pneumonia (J10-J18)	7	0.9	*
13	Certain conditions originating in the perinatal period (P00-P96)	6	0.8	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
15	Diabetes mellitus (E10-E14)	3	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
...	All other causes (Residual)	133	17.6	4.3
Black, both sexes, 10-14 years				
...	All causes	887	100.0	28.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	293	33.0	9.2
2	Assault (homicide) (X85-Y09,Y87.1)	84	9.5	2.6
3	Malignant neoplasms (C00-C97)	78	8.8	2.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	5.3	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	5.2	1.5
5	Chronic lower respiratory diseases (J40-J47)	46	5.2	1.5
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	40	4.5	1.3
8	Anemias (D50-D64)	16	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	14	1.6	*
10	Cerebrovascular diseases (I60-I69)	13	1.5	*
11	Meningitis (G00,G03)	8	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.7	*
12	Diabetes mellitus (E10-E14)	6	0.7	*
14	Influenza and pneumonia (J10-J18)	4	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.5	*
...	All other causes (Residual)	182	20.5	5.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 15-19 years				
...	All causes	2,717	100.0	89.0
1	Assault (homicide) (X85-Y09,Y87.1)	1,028	37.8	33.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	737	27.1	24.1
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	172	6.3	5.6
4	Malignant neoplasms (C00-C97)	136	5.0	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	125	4.6	4.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	2.1	1.8
7	Chronic lower respiratory diseases (J40-J47)	44	1.6	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.8	0.7
9	Anemias (D50-D64)	16	0.6	*
10	Legal intervention (Y35,Y89.0)	15	0.6	*
11	Diabetes mellitus (E10-E14)	14	0.5	*
11	Influenza and pneumonia (J10-J18)	14	0.5	*
13	Cerebrovascular diseases (I60-I69)	13	0.5	*
14	Septicemia (A40-A41)	12	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
...	All other causes (Residual)	301	11.1	9.9
Black, both sexes, 20-24 years				
...	All causes	4,332	100.0	155.7
1	Assault (homicide) (X85-Y09,Y87.1)	1,791	41.3	64.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	958	22.1	34.4
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	308	7.1	11.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	208	4.8	7.5
5	Malignant neoplasms (C00-C97)	165	3.8	5.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	103	2.4	3.7
7	Anemias (D50-D64)	46	1.1	1.7
8	Diabetes mellitus (E10-E14)	41	0.9	1.5
9	Chronic lower respiratory diseases (J40-J47)	37	0.9	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	0.8	1.3
11	Pregnancy, childbirth and the puerperium (O00-099)	32	0.7	1.2
12	Cerebrovascular diseases (I60-I69)	30	0.7	1.1
12	Influenza and pneumonia (J10-J18)	30	0.7	1.1
14	Septicemia (A40-A41)	21	0.5	0.8
15	Legal intervention (Y35,Y89.0)	13	0.3	*
...	All other causes (Residual)	514	11.9	18.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-29 years				
...	All causes	4,541	100.0	175.6
1	Assault (homicide) (X85-Y09,Y87.1)	1,301	28.7	50.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	892	19.6	34.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	409	9.0	15.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	346	7.6	13.4
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	266	5.9	10.3
6	Malignant neoplasms (C00-C97)	215	4.7	8.3
7	Diabetes mellitus (E10-E14)	75	1.7	2.9
8	Chronic lower respiratory diseases (J40-J47)	51	1.1	2.0
9	Anemias (D50-D64)	47	1.0	1.8
10	Cerebrovascular diseases (I60-I69)	41	0.9	1.6
11	Septicemia (A40-A41)	40	0.9	1.5
12	Influenza and pneumonia (J10-J18)	38	0.8	1.5
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	0.8	1.5
14	Pregnancy, childbirth and the puerperium (O00-099)	31	0.7	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.6	1.1
...	All other causes (Residual)	722	15.9	27.9
Black, both sexes, 30-34 years				
...	All causes	5,698	100.0	214.9
1	Human immunodeficiency virus (HIV) disease (B20-B24)	1,023	18.0	38.6
2	Assault (homicide) (X85-Y09,Y87.1)	923	16.2	34.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	878	15.4	33.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	591	10.4	22.3
5	Malignant neoplasms (C00-C97)	434	7.6	16.4
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	210	3.7	7.9
7	Cerebrovascular diseases (I60-I69)	119	2.1	4.5
8	Diabetes mellitus (E10-E14)	114	2.0	4.3
9	Anemias (D50-D64)	77	1.4	2.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	1.2	2.7
11	Septicemia (A40-A41)	68	1.2	2.6
12	Chronic lower respiratory diseases (J40-J47)	65	1.1	2.5
13	Influenza and pneumonia (J10-J18)	55	1.0	2.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	0.7	1.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	36	0.6	1.4
...	All other causes (Residual)	995	17.5	37.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 35-39 years				
...	All causes	8,352	100.0	288.5
1	Human immunodeficiency virus (HIV) disease (B20-B24)	1,470	17.6	50.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,127	13.5	38.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,058	12.7	36.5
4	Malignant neoplasms (C00-C97)	1,017	12.2	35.1
5	Assault (homicide) (X85-Y09,Y87.1)	764	9.1	26.4
6	Cerebrovascular diseases (I60-I69)	283	3.4	9.8
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	214	2.6	7.4
8	Diabetes mellitus (E10-E14)	177	2.1	6.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	144	1.7	5.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	1.6	4.5
11	Septicemia (A40-A41)	109	1.3	3.8
12	Chronic lower respiratory diseases (J40-J47)	104	1.2	3.6
13	Influenza and pneumonia (J10-J18)	85	1.0	2.9
14	Anemias (D50-D64)	65	0.8	2.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	62	0.7	2.1
...	All other causes (Residual)	1,543	18.5	53.3
Black, both sexes, 40-44 years				
...	All causes	12,205	100.0	434.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,178	17.8	77.5
2	Malignant neoplasms (C00-C97)	2,091	17.1	74.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,610	13.2	57.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,229	10.1	43.7
5	Assault (homicide) (X85-Y09,Y87.1)	612	5.0	21.8
6	Cerebrovascular diseases (I60-I69)	553	4.5	19.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	307	2.5	10.9
8	Diabetes mellitus (E10-E14)	285	2.3	10.1
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	224	1.8	8.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	176	1.4	6.3
11	Influenza and pneumonia (J10-J18)	172	1.4	6.1
12	Chronic lower respiratory diseases (J40-J47)	171	1.4	6.1
13	Septicemia (A40-A41)	163	1.3	5.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	117	1.0	4.2
15	Viral hepatitis (B15-B19)	88	0.7	3.1
...	All other causes (Residual)	2,229	18.3	79.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-49 years				
...	All causes	15,735	100.0	677.5
1	Malignant neoplasms (C00-C97)	3,437	21.8	148.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,356	21.3	144.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,385	8.8	59.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,109	7.0	47.8
5	Cerebrovascular diseases (I60-I69)	878	5.6	37.8
6	Diabetes mellitus (E10-E14)	517	3.3	22.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	453	2.9	19.5
8	Assault (homicide) (X85-Y09,Y87.1)	397	2.5	17.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	300	1.9	12.9
10	Septicemia (A40-A41)	259	1.6	11.2
10	Chronic lower respiratory diseases (J40-J47)	259	1.6	11.2
12	Influenza and pneumonia (J10-J18)	197	1.3	8.5
13	Viral hepatitis (B15-B19)	172	1.1	7.4
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	171	1.1	7.4
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	156	1.0	6.7
...	All other causes (Residual)	2,689	17.1	115.8
Black, both sexes, 50-54 years				
...	All causes	17,554	100.0	971.3
1	Malignant neoplasms (C00-C97)	4,710	26.8	260.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,324	24.6	239.3
3	Cerebrovascular diseases (I60-I69)	958	5.5	53.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	879	5.0	48.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	872	5.0	48.2
6	Diabetes mellitus (E10-E14)	762	4.3	42.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	484	2.8	26.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	369	2.1	20.4
9	Chronic lower respiratory diseases (J40-J47)	340	1.9	18.8
10	Septicemia (A40-A41)	316	1.8	17.5
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	238	1.4	13.2
12	Assault (homicide) (X85-Y09,Y87.1)	231	1.3	12.8
13	Influenza and pneumonia (J10-J18)	228	1.3	12.6
14	Viral hepatitis (B15-B19)	164	0.9	9.1
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	119	0.7	6.6
...	All other causes (Residual)	2,560	14.6	141.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 55-59 years				
...	All causes	18,161	100.0	1,366.1
1	Malignant neoplasms (C00-C97)	5,499	30.3	413.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,133	28.3	386.1
3	Cerebrovascular diseases (I60-I69)	1,043	5.7	78.5
4	Diabetes mellitus (E10-E14)	894	4.9	67.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	566	3.1	42.6
6	Chronic lower respiratory diseases (J40-J47)	445	2.5	33.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	436	2.4	32.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	395	2.2	29.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	349	1.9	26.3
10	Septicemia (A40-A41)	328	1.8	24.7
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	242	1.3	18.2
12	Influenza and pneumonia (J10-J18)	240	1.3	18.1
13	Assault (homicide) (X85-Y09,Y87.1)	123	0.7	9.3
14	Aortic aneurysm and dissection (I71)	98	0.5	7.4
15	Viral hepatitis (B15-B19)	93	0.5	7.0
...	All other causes (Residual)	2,277	12.5	171.3
Black, both sexes, 60-64 years				
...	All causes	21,120	100.0	1,950.9
1	Malignant neoplasms (C00-C97)	6,620	31.3	611.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,252	29.6	577.5
3	Cerebrovascular diseases (I60-I69)	1,248	5.9	115.3
4	Diabetes mellitus (E10-E14)	1,121	5.3	103.6
5	Chronic lower respiratory diseases (J40-J47)	628	3.0	58.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	555	2.6	51.3
7	Septicemia (A40-A41)	440	2.1	40.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	431	2.0	39.8
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	317	1.5	29.3
10	Influenza and pneumonia (J10-J18)	298	1.4	27.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	284	1.3	26.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	241	1.1	22.3
13	Aortic aneurysm and dissection (I71)	102	0.5	9.4
14	Assault (homicide) (X85-Y09,Y87.1)	81	0.4	7.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.4	7.2
...	All other causes (Residual)	2,424	11.5	223.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-69 years				
...	All causes	25,064	100.0	2,662.8
1	Malignant neoplasms (C00-C97)	7,797	31.1	828.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,367	29.4	782.7
3	Cerebrovascular diseases (I60-I69)	1,582	6.3	168.1
4	Diabetes mellitus (E10-E14)	1,441	5.7	153.1
5	Chronic lower respiratory diseases (J40-J47)	881	3.5	93.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	756	3.0	80.3
7	Septicemia (A40-A41)	540	2.2	57.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	417	1.7	44.3
9	Influenza and pneumonia (J10-J18)	416	1.7	44.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	365	1.5	38.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	249	1.0	26.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	137	0.5	14.6
13	Aortic aneurysm and dissection (I71)	129	0.5	13.7
14	Pneumonitis due to solids and liquids (J69)	102	0.4	10.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	96	0.4	10.2
...	All other causes (Residual)	2,789	11.1	296.3
Black, both sexes, 70-74 years				
...	All causes	30,131	100.0	3,984.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,223	30.6	1,219.5
2	Malignant neoplasms (C00-C97)	8,839	29.3	1,168.8
3	Cerebrovascular diseases (I60-I69)	2,107	7.0	278.6
4	Diabetes mellitus (E10-E14)	1,677	5.6	221.7
5	Chronic lower respiratory diseases (J40-J47)	1,133	3.8	149.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	897	3.0	118.6
7	Septicemia (A40-A41)	649	2.2	85.8
8	Influenza and pneumonia (J10-J18)	581	1.9	76.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	451	1.5	59.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	441	1.5	58.3
11	Alzheimer's disease (G30)	194	0.6	25.7
12	Pneumonitis due to solids and liquids (J69)	188	0.6	24.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	185	0.6	24.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	152	0.5	20.1
14	Aortic aneurysm and dissection (I71)	152	0.5	20.1
...	All other causes (Residual)	3,262	10.8	431.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 75-79 years				
...	All causes	32,541	100.0	5,803.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,333	31.8	1,843.0
2	Malignant neoplasms (C00-C97)	8,338	25.6	1,487.1
3	Cerebrovascular diseases (I60-I69)	2,674	8.2	476.9
4	Diabetes mellitus (E10-E14)	1,781	5.5	317.7
5	Chronic lower respiratory diseases (J40-J47)	1,191	3.7	212.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	931	2.9	166.0
7	Septicemia (A40-A41)	824	2.5	147.0
8	Influenza and pneumonia (J10-J18)	700	2.2	124.8
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	520	1.6	92.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	424	1.3	75.6
11	Alzheimer's disease (G30)	400	1.2	71.3
12	Pneumonitis due to solids and liquids (J69)	219	0.7	39.1
13	Aortic aneurysm and dissection (I71)	190	0.6	33.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	150	0.5	26.8
15	Atherosclerosis (I70)	132	0.4	23.5
...	All other causes (Residual)	3,734	11.5	666.0
Black, both sexes, 80-84 years				
...	All causes	28,903	100.0	8,515.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,469	32.8	2,789.8
2	Malignant neoplasms (C00-C97)	6,023	20.8	1,774.5
3	Cerebrovascular diseases (I60-I69)	2,725	9.4	802.9
4	Diabetes mellitus (E10-E14)	1,434	5.0	422.5
5	Chronic lower respiratory diseases (J40-J47)	961	3.3	283.1
6	Influenza and pneumonia (J10-J18)	869	3.0	256.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	858	3.0	252.8
8	Septicemia (A40-A41)	766	2.7	225.7
9	Alzheimer's disease (G30)	608	2.1	179.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	462	1.6	136.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	354	1.2	104.3
12	Pneumonitis due to solids and liquids (J69)	253	0.9	74.5
13	Atherosclerosis (I70)	191	0.7	56.3
14	Aortic aneurysm and dissection (I71)	143	0.5	42.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	134	0.5	39.5
...	All other causes (Residual)	3,653	12.6	1,076.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85-89 years				
...	All causes	23,919	100.0	13,085.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,478	35.4	4,638.0
2	Malignant neoplasms (C00-C97)	3,897	16.3	2,131.9
3	Cerebrovascular diseases (I60-I69)	2,492	10.4	1,363.3
4	Diabetes mellitus (E10-E14)	962	4.0	526.3
5	Influenza and pneumonia (J10-J18)	827	3.5	452.4
6	Alzheimer's disease (G30)	682	2.9	373.1
7	Chronic lower respiratory diseases (J40-J47)	671	2.8	367.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	671	2.8	367.1
9	Septicemia (A40-A41)	635	2.7	347.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	440	1.8	240.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	280	1.2	153.2
12	Pneumonitis due to solids and liquids (J69)	236	1.0	129.1
13	Atherosclerosis (I70)	173	0.7	94.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	118	0.5	64.6
15	Aortic aneurysm and dissection (I71)	114	0.5	62.4
...	All other causes (Residual)	3,243	13.6	1,774.1
Black, both sexes, 90-94 years				
...	All causes	14,680	100.0	16,151.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,414	36.9	5,956.7
2	Malignant neoplasms (C00-C97)	1,784	12.2	1,962.8
3	Cerebrovascular diseases (I60-I69)	1,514	10.3	1,665.8
4	Influenza and pneumonia (J10-J18)	657	4.5	722.9
5	Diabetes mellitus (E10-E14)	507	3.5	557.8
6	Alzheimer's disease (G30)	464	3.2	510.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	413	2.8	454.4
8	Septicemia (A40-A41)	380	2.6	418.1
9	Chronic lower respiratory diseases (J40-J47)	360	2.5	396.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	283	1.9	311.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	1.3	210.1
12	Pneumonitis due to solids and liquids (J69)	180	1.2	198.0
13	Atherosclerosis (I70)	153	1.0	168.3
14	Aortic aneurysm and dissection (I71)	64	0.4	70.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.4	69.3
...	All other causes (Residual)	2,253	15.3	2,478.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 95-99 years				
...	All causes	6,407	100.0	17,451.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,503	39.1	6,817.6
2	Cerebrovascular diseases (I60-I69)	688	10.7	1,873.9
3	Malignant neoplasms (C00-C97)	591	9.2	1,609.7
4	Influenza and pneumonia (J10-J18)	315	4.9	858.0
5	Alzheimer's disease (G30)	219	3.4	596.5
6	Septicemia (A40-A41)	187	2.9	509.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	171	2.7	465.8
8	Diabetes mellitus (E10-E14)	160	2.5	435.8
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	142	2.2	386.8
10	Chronic lower respiratory diseases (J40-J47)	133	2.1	362.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	1.2	217.9
12	Atherosclerosis (I70)	76	1.2	207.0
13	Pneumonitis due to solids and liquids (J69)	74	1.2	201.6
14	Nutritional deficiencies (E40-E64)	41	0.6	111.7
15	Anemias (D50-D64)	28	0.4	76.3
...	All other causes (Residual)	999	15.6	2,721.0
Black, both sexes, 100 years and over				
...	All causes	2,032	100.0	24,024.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	834	41.0	9,860.5
2	Cerebrovascular diseases (I60-I69)	197	9.7	2,329.2
3	Influenza and pneumonia (J10-J18)	122	6.0	1,442.4
4	Malignant neoplasms (C00-C97)	118	5.8	1,395.1
5	Alzheimer's disease (G30)	58	2.9	685.7
6	Septicemia (A40-A41)	53	2.6	626.6
7	Diabetes mellitus (E10-E14)	49	2.4	579.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	2.2	520.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	41	2.0	484.7
10	Atherosclerosis (I70)	40	2.0	472.9
11	Chronic lower respiratory diseases (J40-J47)	31	1.5	366.5
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	1.4	331.0
13	Pneumonitis due to solids and liquids (J69)	21	1.0	248.3
14	Nutritional deficiencies (E40-E64)	16	0.8	*
15	Anemias (D50-D64)	10	0.5	*
...	All other causes (Residual)	370	18.2	4,374.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, all ages				
...	All causes	145,184	100.0	865.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,740	25.3	219.0
2	Malignant neoplasms (C00-C97)	32,817	22.6	195.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,531	5.9	50.9
4	Cerebrovascular diseases (I60-I69)	8,026	5.5	47.8
5	Assault (homicide) (X85-Y09,Y87.1)	6,482	4.5	38.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	5,400	3.7	32.2
7	Diabetes mellitus (E10-E14)	4,771	3.3	28.4
8	Chronic lower respiratory diseases (J40-J47)	4,238	2.9	25.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,074	2.1	18.3
10	Influenza and pneumonia (J10-J18)	2,915	2.0	17.4
11	Certain conditions originating in the perinatal period (P00-P96)	2,844	2.0	17.0
12	Septicemia (A40-A41)	2,587	1.8	15.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,730	1.2	10.3
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	1,636	1.1	9.8
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1,592	1.1	9.5
...	All other causes (Residual)	21,801	15.0	130.0
Black, male, 1-4 years				
...	All causes	692	100.0	61.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	34.5	21.1
2	Assault (homicide) (X85-Y09,Y87.1)	75	10.8	6.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	10.1	6.2
4	Malignant neoplasms (C00-C97)	34	4.9	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	3.5	2.1
6	Influenza and pneumonia (J10-J18)	21	3.0	1.9
7	Certain conditions originating in the perinatal period (P00-P96)	18	2.6	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.0	*
9	Septicemia (A40-A41)	13	1.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.2	*
10	Anemias (D50-D64)	8	1.2	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.9	*
12	Meningitis (G00,G03)	6	0.9	*
14	Meningococcal infection (A39)	4	0.6	*
15	Cerebrovascular diseases (I60-I69)	3	0.4	*
...	All other causes (Residual)	149	21.5	13.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 5-9 years				
...	All causes	431	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	209	48.5	13.3
2	Malignant neoplasms (C00-C97)	42	9.7	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	6.7	1.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	4.2	*
5	Assault (homicide) (X85-Y09,Y87.1)	16	3.7	*
6	Chronic lower respiratory diseases (J40-J47)	14	3.2	*
7	Anemias (D50-D64)	10	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.4	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.2	*
10	Septicemia (A40-A41)	4	0.9	*
10	Influenza and pneumonia (J10-J18)	4	0.9	*
12	Meningococcal infection (A39)	3	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
14	Diabetes mellitus (E10-E14)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	64	14.8	4.1
Black, male, 10-14 years				
...	All causes	543	100.0	33.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	37.8	12.7
2	Assault (homicide) (X85-Y09,Y87.1)	56	10.3	3.5
3	Malignant neoplasms (C00-C97)	45	8.3	2.8
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	35	6.4	2.2
5	Chronic lower respiratory diseases (J40-J47)	29	5.3	1.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.4	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	4.2	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.3	*
8	Anemias (D50-D64)	7	1.3	*
10	Meningitis (G00,G03)	6	1.1	*
11	Cerebrovascular diseases (I60-I69)	5	0.9	*
12	Influenza and pneumonia (J10-J18)	3	0.6	*
13	Septicemia (A40-A41)	2	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Legal intervention (Y35,Y89.0)	2	0.4	*
...	All other causes (Residual)	90	16.6	5.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-19 years				
...	All causes	2,045	100.0	131.6
1	Assault (homicide) (X85-Y09,Y87.1)	899	44.0	57.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	542	26.5	34.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	150	7.3	9.7
4	Malignant neoplasms (C00-C97)	82	4.0	5.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	75	3.7	4.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	1.9	2.5
7	Chronic lower respiratory diseases (J40-J47)	31	1.5	2.0
8	Legal intervention (Y35,Y89.0)	15	0.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.5	*
10	Diabetes mellitus (E10-E14)	8	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.3	*
11	Anemias (D50-D64)	7	0.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
14	Septicemia (A40-A41)	6	0.3	*
14	Influenza and pneumonia (J10-J18)	6	0.3	*
...	All other causes (Residual)	160	7.8	10.3
Black, male, 20-24 years				
...	All causes	3,273	100.0	237.6
1	Assault (homicide) (X85-Y09,Y87.1)	1,602	48.9	116.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	719	22.0	52.2
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	265	8.1	19.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	118	3.6	8.6
5	Malignant neoplasms (C00-C97)	97	3.0	7.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	49	1.5	3.6
7	Anemias (D50-D64)	23	0.7	1.7
8	Diabetes mellitus (E10-E14)	20	0.6	1.5
8	Chronic lower respiratory diseases (J40-J47)	20	0.6	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.6	*
11	Cerebrovascular diseases (I60-I69)	15	0.5	*
11	Influenza and pneumonia (J10-J18)	15	0.5	*
13	Legal intervention (Y35,Y89.0)	13	0.4	*
14	Septicemia (A40-A41)	12	0.4	*
15	Viral hepatitis (B15-B19)	5	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.2	*
...	All other causes (Residual)	276	8.4	20.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 25-29 years				
...	All causes	3,163	100.0	255.6
1	Assault (homicide) (X85-Y09,Y87.1)	1,146	36.2	92.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	693	21.9	56.0
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	224	7.1	18.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	210	6.6	17.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	201	6.4	16.2
6	Malignant neoplasms (C00-C97)	94	3.0	7.6
7	Diabetes mellitus (E10-E14)	40	1.3	3.2
8	Chronic lower respiratory diseases (J40-J47)	31	1.0	2.5
9	Septicemia (A40-A41)	27	0.9	2.2
9	Influenza and pneumonia (J10-J18)	27	0.9	2.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.8	1.9
12	Anemias (D50-D64)	20	0.6	1.6
13	Legal intervention (Y35,Y89.0)	19	0.6	*
14	Cerebrovascular diseases (I60-I69)	18	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
...	All other causes (Residual)	376	11.9	30.4
Black, male, 30-34 years				
...	All causes	3,587	100.0	287.9
1	Assault (homicide) (X85-Y09,Y87.1)	765	21.3	61.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	647	18.0	51.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	625	17.4	50.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	346	9.6	27.8
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	178	5.0	14.3
6	Malignant neoplasms (C00-C97)	168	4.7	13.5
7	Diabetes mellitus (E10-E14)	59	1.6	4.7
8	Cerebrovascular diseases (I60-I69)	53	1.5	4.3
9	Chronic lower respiratory diseases (J40-J47)	41	1.1	3.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.0	3.0
11	Anemias (D50-D64)	35	1.0	2.8
12	Septicemia (A40-A41)	32	0.9	2.6
12	Influenza and pneumonia (J10-J18)	32	0.9	2.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.6	1.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	15	0.4	*
...	All other causes (Residual)	532	14.8	42.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-39 years				
...	All causes	5,015	100.0	368.1
1	Human immunodeficiency virus (HIV) disease (B20-B24)	988	19.7	72.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	772	15.4	56.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	674	13.4	49.5
4	Assault (homicide) (X85-Y09,Y87.1)	590	11.8	43.3
5	Malignant neoplasms (C00-C97)	383	7.6	28.1
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	180	3.6	13.2
7	Cerebrovascular diseases (I60-I69)	127	2.5	9.3
8	Diabetes mellitus (E10-E14)	99	2.0	7.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	92	1.8	6.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	1.4	5.2
11	Chronic lower respiratory diseases (J40-J47)	52	1.0	3.8
12	Septicemia (A40-A41)	48	1.0	3.5
13	Influenza and pneumonia (J10-J18)	42	0.8	3.1
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	30	0.6	2.2
15	Anemias (D50-D64)	26	0.5	1.9
...	All other causes (Residual)	841	16.8	61.7
Black, male, 40-44 years				
...	All causes	7,236	100.0	548.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,335	18.4	101.1
2	Human immunodeficiency virus (HIV) disease (B20-B24)	1,108	15.3	83.9
3	Malignant neoplasms (C00-C97)	926	12.8	70.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	906	12.5	68.6
5	Assault (homicide) (X85-Y09,Y87.1)	450	6.2	34.1
6	Cerebrovascular diseases (I60-I69)	270	3.7	20.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	192	2.7	14.5
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	176	2.4	13.3
9	Diabetes mellitus (E10-E14)	155	2.1	11.7
10	Influenza and pneumonia (J10-J18)	110	1.5	8.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	1.4	7.9
12	Chronic lower respiratory diseases (J40-J47)	78	1.1	5.9
13	Septicemia (A40-A41)	77	1.1	5.8
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	70	1.0	5.3
15	Viral hepatitis (B15-B19)	56	0.8	4.2
...	All other causes (Residual)	1,223	16.9	92.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 45-49 years				
...	All causes	9,350	100.0	877.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,091	22.4	196.1
2	Malignant neoplasms (C00-C97)	1,664	17.8	156.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,016	10.9	95.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	817	8.7	76.6
5	Cerebrovascular diseases (I60-I69)	449	4.8	42.1
6	Assault (homicide) (X85-Y09,Y87.1)	315	3.4	29.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	289	3.1	27.1
8	Diabetes mellitus (E10-E14)	280	3.0	26.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	191	2.0	17.9
10	Septicemia (A40-A41)	136	1.5	12.8
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	129	1.4	12.1
12	Chronic lower respiratory diseases (J40-J47)	125	1.3	11.7
13	Influenza and pneumonia (J10-J18)	124	1.3	11.6
14	Viral hepatitis (B15-B19)	121	1.3	11.3
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	99	1.1	9.3
...	All other causes (Residual)	1,504	16.1	141.1
Black, male, 50-54 years				
...	All causes	10,563	100.0	1,300.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,746	26.0	338.2
2	Malignant neoplasms (C00-C97)	2,528	23.9	311.3
3	Human immunodeficiency virus (HIV) disease (B20-B24)	666	6.3	82.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	658	6.2	81.0
5	Cerebrovascular diseases (I60-I69)	520	4.9	64.0
6	Diabetes mellitus (E10-E14)	374	3.5	46.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	350	3.3	43.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	207	2.0	25.5
9	Assault (homicide) (X85-Y09,Y87.1)	193	1.8	23.8
10	Septicemia (A40-A41)	183	1.7	22.5
11	Chronic lower respiratory diseases (J40-J47)	169	1.6	20.8
12	Influenza and pneumonia (J10-J18)	148	1.4	18.2
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	134	1.3	16.5
14	Viral hepatitis (B15-B19)	120	1.1	14.8
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	90	0.9	11.1
...	All other causes (Residual)	1,477	14.0	181.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-59 years				
...	All causes	10,787	100.0	1,854.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,237	30.0	556.5
2	Malignant neoplasms (C00-C97)	3,114	28.9	535.4
3	Cerebrovascular diseases (I60-I69)	594	5.5	102.1
4	Diabetes mellitus (E10-E14)	438	4.1	75.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	434	4.0	74.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	310	2.9	53.3
7	Chronic lower respiratory diseases (J40-J47)	240	2.2	41.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	234	2.2	40.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	227	2.1	39.0
10	Septicemia (A40-A41)	169	1.6	29.1
11	Influenza and pneumonia (J10-J18)	157	1.5	27.0
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	137	1.3	23.6
13	Assault (homicide) (X85-Y09,Y87.1)	97	0.9	16.7
14	Viral hepatitis (B15-B19)	63	0.6	10.8
14	Aortic aneurysm and dissection (I71)	63	0.6	10.8
...	All other causes (Residual)	1,273	11.8	218.9
Black, male, 60-64 years				
...	All causes	11,891	100.0	2,573.7
1	Malignant neoplasms (C00-C97)	3,715	31.2	804.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,645	30.7	788.9
3	Cerebrovascular diseases (I60-I69)	664	5.6	143.7
4	Diabetes mellitus (E10-E14)	524	4.4	113.4
5	Chronic lower respiratory diseases (J40-J47)	355	3.0	76.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	316	2.7	68.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	262	2.2	56.7
8	Septicemia (A40-A41)	218	1.8	47.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	191	1.6	41.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	180	1.5	39.0
11	Influenza and pneumonia (J10-J18)	170	1.4	36.8
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	166	1.4	35.9
13	Assault (homicide) (X85-Y09,Y87.1)	63	0.5	13.6
14	Aortic aneurysm and dissection (I71)	62	0.5	13.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.4	10.6
...	All other causes (Residual)	1,311	11.0	283.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 65-69 years				
...	All causes	13,505	100.0	3,365.9
1	Malignant neoplasms (C00-C97)	4,371	32.4	1,089.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,973	29.4	990.2
3	Cerebrovascular diseases (I60-I69)	818	6.1	203.9
4	Diabetes mellitus (E10-E14)	593	4.4	147.8
5	Chronic lower respiratory diseases (J40-J47)	532	3.9	132.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	334	2.5	83.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	285	2.1	71.0
8	Septicemia (A40-A41)	265	2.0	66.0
9	Influenza and pneumonia (J10-J18)	249	1.8	62.1
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	154	1.1	38.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	146	1.1	36.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	101	0.7	25.2
13	Aortic aneurysm and dissection (I71)	68	0.5	16.9
14	Pneumonitis due to solids and liquids (J69)	67	0.5	16.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.4	15.0
...	All other causes (Residual)	1,489	11.0	371.1
Black, male, 70-74 years				
...	All causes	15,566	100.0	4,960.0
1	Malignant neoplasms (C00-C97)	4,950	31.8	1,577.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,699	30.2	1,497.3
3	Cerebrovascular diseases (I60-I69)	1,017	6.5	324.1
4	Chronic lower respiratory diseases (J40-J47)	677	4.3	215.7
5	Diabetes mellitus (E10-E14)	671	4.3	213.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	382	2.5	121.7
7	Influenza and pneumonia (J10-J18)	322	2.1	102.6
8	Septicemia (A40-A41)	290	1.9	92.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	258	1.7	82.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	191	1.2	60.9
11	Pneumonitis due to solids and liquids (J69)	123	0.8	39.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	112	0.7	35.7
13	Alzheimer's disease (G30)	83	0.5	26.4
13	Aortic aneurysm and dissection (I71)	83	0.5	26.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	71	0.5	22.6
...	All other causes (Residual)	1,637	10.5	521.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-79 years				
...	All causes	15,682	100.0	7,139.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,752	30.3	2,163.3
2	Malignant neoplasms (C00-C97)	4,584	29.2	2,086.9
3	Cerebrovascular diseases (I60-I69)	1,153	7.4	524.9
4	Chronic lower respiratory diseases (J40-J47)	691	4.4	314.6
5	Diabetes mellitus (E10-E14)	659	4.2	300.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	421	2.7	191.7
7	Septicemia (A40-A41)	387	2.5	176.2
8	Influenza and pneumonia (J10-J18)	365	2.3	166.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	253	1.6	115.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	197	1.3	89.7
11	Alzheimer's disease (G30)	137	0.9	62.4
12	Pneumonitis due to solids and liquids (J69)	132	0.8	60.1
13	Aortic aneurysm and dissection (I71)	101	0.6	46.0
14	Parkinson's disease (G20-G21)	80	0.5	36.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.5	33.2
...	All other causes (Residual)	1,697	10.8	772.6
Black, male, 80-84 years				
...	All causes	12,344	100.0	10,247.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,827	31.0	3,177.1
2	Malignant neoplasms (C00-C97)	3,183	25.8	2,642.5
3	Cerebrovascular diseases (I60-I69)	1,024	8.3	850.1
4	Chronic lower respiratory diseases (J40-J47)	532	4.3	441.7
5	Diabetes mellitus (E10-E14)	454	3.7	376.9
6	Influenza and pneumonia (J10-J18)	408	3.3	338.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	344	2.8	285.6
8	Septicemia (A40-A41)	297	2.4	246.6
9	Alzheimer's disease (G30)	185	1.5	153.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	1.5	153.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	163	1.3	135.3
12	Pneumonitis due to solids and liquids (J69)	114	0.9	94.6
13	Atherosclerosis (I70)	79	0.6	65.6
14	Parkinson's disease (G20-G21)	78	0.6	64.8
15	Aortic aneurysm and dissection (I71)	54	0.4	44.8
...	All other causes (Residual)	1,417	11.5	1,176.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 85-89 years				
...	All causes	8,428	100.0	14,613.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,776	32.9	4,813.3
2	Malignant neoplasms (C00-C97)	1,832	21.7	3,176.5
3	Cerebrovascular diseases (I60-I69)	730	8.7	1,265.7
4	Chronic lower respiratory diseases (J40-J47)	360	4.3	624.2
5	Influenza and pneumonia (J10-J18)	325	3.9	563.5
6	Diabetes mellitus (E10-E14)	260	3.1	450.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	234	2.8	405.7
8	Septicemia (A40-A41)	218	2.6	378.0
9	Alzheimer's disease (G30)	181	2.1	313.8
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	127	1.5	220.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	1.4	209.8
12	Pneumonitis due to solids and liquids (J69)	90	1.1	156.0
13	Atherosclerosis (I70)	63	0.7	109.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.5	78.0
15	Parkinson's disease (G20-G21)	44	0.5	76.3
...	All other causes (Residual)	1,022	12.1	1,772.0
Black, male, 90-94 years				
...	All causes	4,249	100.0	16,649.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,408	33.1	5,517.2
2	Malignant neoplasms (C00-C97)	742	17.5	2,907.5
3	Cerebrovascular diseases (I60-I69)	369	8.7	1,445.9
4	Influenza and pneumonia (J10-J18)	219	5.2	858.2
5	Chronic lower respiratory diseases (J40-J47)	173	4.1	677.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	3.0	493.7
7	Septicemia (A40-A41)	109	2.6	427.1
7	Diabetes mellitus (E10-E14)	109	2.6	427.1
9	Alzheimer's disease (G30)	99	2.3	387.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	1.9	309.6
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	75	1.8	293.9
12	Pneumonitis due to solids and liquids (J69)	67	1.6	262.5
13	Atherosclerosis (I70)	41	1.0	160.7
14	Parkinson's disease (G20-G21)	24	0.6	94.0
15	Anemias (D50-D64)	22	0.5	86.2
...	All other causes (Residual)	587	13.8	2,300.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 95-99 years				
...	All causes	1,532	100.0	16,990.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	548	35.8	6,077.4
2	Malignant neoplasms (C00-C97)	219	14.3	2,428.7
3	Cerebrovascular diseases (I60-I69)	145	9.5	1,608.1
4	Influenza and pneumonia (J10-J18)	86	5.6	953.8
5	Chronic lower respiratory diseases (J40-J47)	60	3.9	665.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	3.9	665.4
7	Alzheimer's disease (G30)	36	2.3	399.2
8	Septicemia (A40-A41)	34	2.2	377.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	2.1	354.9
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	27	1.8	299.4
11	Diabetes mellitus (E10-E14)	21	1.4	232.9
11	Pneumonitis due to solids and liquids (J69)	21	1.4	232.9
13	Atherosclerosis (I70)	19	1.2	*
14	Nutritional deficiencies (E40-E64)	11	0.7	*
15	Parkinson's disease (G20-G21)	6	0.4	*
...	All other causes (Residual)	207	13.5	2,295.7
Black, male, 100 years and over				
...	All causes	351	100.0	19,977.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	153	43.6	8,708.0
2	Malignant neoplasms (C00-C97)	38	10.8	2,162.8
3	Influenza and pneumonia (J10-J18)	25	7.1	1,422.9
4	Cerebrovascular diseases (I60-I69)	21	6.0	1,195.2
5	Septicemia (A40-A41)	10	2.8	*
6	Chronic lower respiratory diseases (J40-J47)	8	2.3	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	2.0	*
8	Alzheimer's disease (G30)	6	1.7	*
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	1.4	*
9	Atherosclerosis (I70)	5	1.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.4	*
12	Pneumonitis due to solids and liquids (J69)	4	1.1	*
13	Diabetes mellitus (E10-E14)	2	0.6	*
14	Anemias (D50-D64)	1	0.3	*
14	Nutritional deficiencies (E40-E64)	1	0.3	*
14	Parkinson's disease (G20-G21)	1	0.3	*
14	Peptic ulcer (K25-K28)	1	0.3	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	1	0.3	*
...	All other causes (Residual)	56	16.0	3,187.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	140,642	100.0	759.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,783	29.0	220.1
2	Malignant neoplasms (C00-C97)	29,128	20.7	157.2
3	Cerebrovascular diseases (I60-I69)	11,195	8.0	60.4
4	Diabetes mellitus (E10-E14)	7,250	5.2	39.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,837	2.7	20.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,746	2.7	20.2
7	Chronic lower respiratory diseases (J40-J47)	3,369	2.4	18.2
8	Septicemia (A40-A41)	3,341	2.4	18.0
9	Influenza and pneumonia (J10-J18)	3,075	2.2	16.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,448	1.7	13.2
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2,295	1.6	12.4
12	Certain conditions originating in the perinatal period (P00-P96)	2,177	1.5	11.8
13	Alzheimer's disease (G30)	1,952	1.4	10.5
14	Assault (homicide) (X85-Y09,Y87.1)	1,385	1.0	7.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,007	0.7	5.4
...	All other causes (Residual)	23,654	16.8	127.7
Black, female, 1-4 years				
...	All causes	556	100.0	50.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	29.9	15.2
2	Assault (homicide) (X85-Y09,Y87.1)	70	12.6	6.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	10.1	5.1
4	Malignant neoplasms (C00-C97)	36	6.5	3.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	4.5	2.3
6	Septicemia (A40-A41)	13	2.3	*
7	Anemias (D50-D64)	11	2.0	*
7	Cerebrovascular diseases (I60-I69)	11	2.0	*
9	Influenza and pneumonia (J10-J18)	9	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.6	*
11	Meningitis (G00,G03)	7	1.3	*
11	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.1	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.1	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	118	21.2	10.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-9 years				
...	All causes	325	100.0	21.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	39.7	8.5
2	Malignant neoplasms (C00-C97)	30	9.2	2.0
3	Assault (homicide) (X85-Y09,Y87.1)	24	7.4	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	4.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.7	*
6	Chronic lower respiratory diseases (J40-J47)	10	3.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	2.2	*
8	Anemias (D50-D64)	5	1.5	*
9	Meningococcal infection (A39)	4	1.2	*
9	Septicemia (A40-A41)	4	1.2	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.2	*
12	Influenza and pneumonia (J10-J18)	3	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.9	*
14	Cerebrovascular diseases (I60-I69)	2	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
...	All other causes (Residual)	69	21.2	4.5
Black, female, 10-14 years				
...	All causes	344	100.0	22.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	25.6	5.6
2	Malignant neoplasms (C00-C97)	33	9.6	2.1
3	Assault (homicide) (X85-Y09,Y87.1)	28	8.1	1.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	7.0	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	6.4	1.4
6	Chronic lower respiratory diseases (J40-J47)	17	4.9	*
7	Anemias (D50-D64)	9	2.6	*
8	Cerebrovascular diseases (I60-I69)	8	2.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	7	2.0	*
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	5	1.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.2	*
11	Diabetes mellitus (E10-E14)	4	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	1.2	*
14	Nutritional deficiencies (E40-E64)	2	0.6	*
14	Meningitis (G00,G03)	2	0.6	*
14	Aortic aneurysm and dissection (I71)	2	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.6	*
...	All other causes (Residual)	83	24.1	5.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 15-19 years				
...	All causes	672	100.0	44.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	29.0	13.0
2	Assault (homicide) (X85-Y09,Y87.1)	129	19.2	8.6
3	Malignant neoplasms (C00-C97)	54	8.0	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	7.4	3.3
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	22	3.3	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.5	*
7	Chronic lower respiratory diseases (J40-J47)	13	1.9	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.6	*
9	Anemias (D50-D64)	9	1.3	*
9	Cerebrovascular diseases (I60-I69)	9	1.3	*
11	Influenza and pneumonia (J10-J18)	8	1.2	*
12	Septicemia (A40-A41)	6	0.9	*
12	Diabetes mellitus (E10-E14)	6	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.6	*
15	Pregnancy, childbirth and the puerperium (O00-099)	4	0.6	*
...	All other causes (Residual)	130	19.3	8.7
Black, female, 20-24 years				
...	All causes	1,059	100.0	75.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	22.6	17.0
2	Assault (homicide) (X85-Y09,Y87.1)	189	17.8	13.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	90	8.5	6.4
4	Malignant neoplasms (C00-C97)	68	6.4	4.8
5	Human immunodeficiency virus (HIV) disease (B20-B24)	54	5.1	3.8
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	43	4.1	3.1
7	Pregnancy, childbirth and the puerperium (O00-099)	32	3.0	2.3
8	Anemias (D50-D64)	23	2.2	1.6
9	Diabetes mellitus (E10-E14)	21	2.0	1.5
10	Chronic lower respiratory diseases (J40-J47)	17	1.6	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	1.5	*
12	Cerebrovascular diseases (I60-I69)	15	1.4	*
12	Influenza and pneumonia (J10-J18)	15	1.4	*
14	Septicemia (A40-A41)	9	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	*
...	All other causes (Residual)	219	20.7	15.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-29 years				
...	All causes	1,378	100.0	102.2
1	Human immunodeficiency virus (HIV) disease (B20-B24)	199	14.4	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	14.4	14.8
3	Assault (homicide) (X85-Y09,Y87.1)	155	11.2	11.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	10.5	10.8
5	Malignant neoplasms (C00-C97)	121	8.8	9.0
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	42	3.0	3.1
7	Diabetes mellitus (E10-E14)	35	2.5	2.6
8	Pregnancy, childbirth and the puerperium (O00-099)	31	2.2	2.3
9	Anemias (D50-D64)	27	2.0	2.0
10	Cerebrovascular diseases (I60-I69)	23	1.7	1.7
11	Chronic lower respiratory diseases (J40-J47)	20	1.5	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.2	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.0	*
14	Septicemia (A40-A41)	13	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.9	*
...	All other causes (Residual)	325	23.6	24.1
Black, female, 30-34 years				
...	All causes	2,111	100.0	150.2
1	Human immunodeficiency virus (HIV) disease (B20-B24)	398	18.9	28.3
2	Malignant neoplasms (C00-C97)	266	12.6	18.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	245	11.6	17.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	231	10.9	16.4
5	Assault (homicide) (X85-Y09,Y87.1)	158	7.5	11.2
6	Cerebrovascular diseases (I60-I69)	66	3.1	4.7
7	Diabetes mellitus (E10-E14)	55	2.6	3.9
8	Anemias (D50-D64)	42	2.0	3.0
9	Septicemia (A40-A41)	36	1.7	2.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.6	2.4
10	Pregnancy, childbirth and the puerperium (O00-099)	34	1.6	2.4
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	32	1.5	2.3
13	Chronic lower respiratory diseases (J40-J47)	24	1.1	1.7
14	Influenza and pneumonia (J10-J18)	23	1.1	1.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	21	1.0	1.5
...	All other causes (Residual)	446	21.1	31.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 35-39 years				
...	All causes	3,337	100.0	217.8
1	Malignant neoplasms (C00-C97)	634	19.0	41.4
2	Human immunodeficiency virus (HIV) disease (B20-B24)	482	14.4	31.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	453	13.6	29.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	286	8.6	18.7
5	Assault (homicide) (X85-Y09,Y87.1)	174	5.2	11.4
6	Cerebrovascular diseases (I60-I69)	156	4.7	10.2
7	Diabetes mellitus (E10-E14)	78	2.3	5.1
8	Septicemia (A40-A41)	61	1.8	4.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	1.8	3.9
10	Chronic lower respiratory diseases (J40-J47)	52	1.6	3.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	52	1.6	3.4
12	Influenza and pneumonia (J10-J18)	43	1.3	2.8
13	Anemias (D50-D64)	39	1.2	2.5
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	34	1.0	2.2
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	32	1.0	2.1
...	All other causes (Residual)	702	21.0	45.8
Black, female, 40-44 years				
...	All causes	4,969	100.0	333.2
1	Malignant neoplasms (C00-C97)	1,165	23.4	78.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	843	17.0	56.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	502	10.1	33.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	323	6.5	21.7
5	Cerebrovascular diseases (I60-I69)	283	5.7	19.0
6	Assault (homicide) (X85-Y09,Y87.1)	162	3.3	10.9
7	Diabetes mellitus (E10-E14)	130	2.6	8.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	115	2.3	7.7
9	Chronic lower respiratory diseases (J40-J47)	93	1.9	6.2
10	Septicemia (A40-A41)	86	1.7	5.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	1.4	4.8
12	Influenza and pneumonia (J10-J18)	62	1.2	4.2
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	48	1.0	3.2
14	Anemias (D50-D64)	47	0.9	3.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	47	0.9	3.2
...	All other causes (Residual)	991	19.9	66.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-49 years				
...	All causes	6,385	100.0	508.2
1	Malignant neoplasms (C00-C97)	1,773	27.8	141.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,265	19.8	100.7
3	Cerebrovascular diseases (I60-I69)	429	6.7	34.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	369	5.8	29.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	292	4.6	23.2
6	Diabetes mellitus (E10-E14)	237	3.7	18.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	164	2.6	13.1
8	Chronic lower respiratory diseases (J40-J47)	134	2.1	10.7
9	Septicemia (A40-A41)	123	1.9	9.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	1.7	8.7
11	Assault (homicide) (X85-Y09,Y87.1)	82	1.3	6.5
12	Influenza and pneumonia (J10-J18)	73	1.1	5.8
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	72	1.1	5.7
14	Viral hepatitis (B15-B19)	51	0.8	4.1
15	Anemias (D50-D64)	34	0.5	2.7
...	All other causes (Residual)	1,178	18.4	93.8
Black, female, 50-54 years				
...	All causes	6,991	100.0	702.4
1	Malignant neoplasms (C00-C97)	2,182	31.2	219.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,578	22.6	158.5
3	Cerebrovascular diseases (I60-I69)	438	6.3	44.0
4	Diabetes mellitus (E10-E14)	388	5.5	39.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	3.2	22.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	206	2.9	20.7
7	Chronic lower respiratory diseases (J40-J47)	171	2.4	17.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	2.3	16.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	134	1.9	13.5
10	Septicemia (A40-A41)	133	1.9	13.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	104	1.5	10.4
12	Influenza and pneumonia (J10-J18)	80	1.1	8.0
13	Viral hepatitis (B15-B19)	44	0.6	4.4
14	Assault (homicide) (X85-Y09,Y87.1)	38	0.5	3.8
15	Anemias (D50-D64)	30	0.4	3.0
...	All other causes (Residual)	1,082	15.5	108.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 55-59 years				
...	All causes	7,374	100.0	986.1
1	Malignant neoplasms (C00-C97)	2,385	32.3	318.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,896	25.7	253.5
3	Diabetes mellitus (E10-E14)	456	6.2	61.0
4	Cerebrovascular diseases (I60-I69)	449	6.1	60.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	209	2.8	27.9
6	Chronic lower respiratory diseases (J40-J47)	205	2.8	27.4
7	Septicemia (A40-A41)	159	2.2	21.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	1.8	17.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	115	1.6	15.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	105	1.4	14.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	85	1.2	11.4
12	Influenza and pneumonia (J10-J18)	83	1.1	11.1
13	Aortic aneurysm and dissection (I71)	35	0.5	4.7
14	Viral hepatitis (B15-B19)	30	0.4	4.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	4.0
...	All other causes (Residual)	1,000	13.6	133.7
Black, female, 60-64 years				
...	All causes	9,229	100.0	1,487.3
1	Malignant neoplasms (C00-C97)	2,905	31.5	468.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,607	28.2	420.1
3	Diabetes mellitus (E10-E14)	597	6.5	96.2
4	Cerebrovascular diseases (I60-I69)	584	6.3	94.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	293	3.2	47.2
6	Chronic lower respiratory diseases (J40-J47)	273	3.0	44.0
7	Septicemia (A40-A41)	222	2.4	35.8
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	151	1.6	24.3
9	Influenza and pneumonia (J10-J18)	128	1.4	20.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	1.2	18.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	104	1.1	16.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	50	0.5	8.1
13	Aortic aneurysm and dissection (I71)	40	0.4	6.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.3	4.7
14	Complications of medical and surgical care (Y40-Y84,Y88)	29	0.3	4.7
...	All other causes (Residual)	1,102	11.9	177.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-69 years				
...	All causes	11,559	100.0	2,140.4
1	Malignant neoplasms (C00-C97)	3,426	29.6	634.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,394	29.4	628.5
3	Diabetes mellitus (E10-E14)	848	7.3	157.0
4	Cerebrovascular diseases (I60-I69)	764	6.6	141.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	422	3.7	78.1
6	Chronic lower respiratory diseases (J40-J47)	349	3.0	64.6
7	Septicemia (A40-A41)	275	2.4	50.9
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	211	1.8	39.1
9	Influenza and pneumonia (J10-J18)	167	1.4	30.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	1.1	24.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	103	0.9	19.1
12	Aortic aneurysm and dissection (I71)	61	0.5	11.3
13	Alzheimer's disease (G30)	37	0.3	6.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	36	0.3	6.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.3	6.7
...	All other causes (Residual)	1,298	11.2	240.4
Black, female, 70-74 years				
...	All causes	14,565	100.0	3,292.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,524	31.1	1,022.5
2	Malignant neoplasms (C00-C97)	3,889	26.7	879.0
3	Cerebrovascular diseases (I60-I69)	1,090	7.5	246.4
4	Diabetes mellitus (E10-E14)	1,006	6.9	227.4
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	515	3.5	116.4
6	Chronic lower respiratory diseases (J40-J47)	456	3.1	103.1
7	Septicemia (A40-A41)	359	2.5	81.1
8	Influenza and pneumonia (J10-J18)	259	1.8	58.5
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	250	1.7	56.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	193	1.3	43.6
11	Alzheimer's disease (G30)	111	0.8	25.1
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	81	0.6	18.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	73	0.5	16.5
14	Aortic aneurysm and dissection (I71)	69	0.5	15.6
15	Pneumonitis due to solids and liquids (J69)	65	0.4	14.7
...	All other causes (Residual)	1,625	11.2	367.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 75-79 years				
...	All causes	16,859	100.0	4,943.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,581	33.1	1,636.6
2	Malignant neoplasms (C00-C97)	3,754	22.3	1,100.8
3	Cerebrovascular diseases (I60-I69)	1,521	9.0	446.0
4	Diabetes mellitus (E10-E14)	1,122	6.7	329.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	510	3.0	149.6
6	Chronic lower respiratory diseases (J40-J47)	500	3.0	146.6
7	Septicemia (A40-A41)	437	2.6	128.1
8	Influenza and pneumonia (J10-J18)	335	2.0	98.2
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	323	1.9	94.7
10	Alzheimer's disease (G30)	263	1.6	77.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	1.0	50.1
12	Aortic aneurysm and dissection (I71)	89	0.5	26.1
13	Pneumonitis due to solids and liquids (J69)	87	0.5	25.5
14	Atherosclerosis (I70)	78	0.5	22.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.5	22.6
...	All other causes (Residual)	2,011	11.9	589.7
Black, female, 80-84 years				
...	All causes	16,559	100.0	7,562.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,642	34.1	2,576.7
2	Malignant neoplasms (C00-C97)	2,840	17.2	1,297.1
3	Cerebrovascular diseases (I60-I69)	1,701	10.3	776.9
4	Diabetes mellitus (E10-E14)	980	5.9	447.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	514	3.1	234.7
6	Septicemia (A40-A41)	469	2.8	214.2
7	Influenza and pneumonia (J10-J18)	461	2.8	210.5
8	Chronic lower respiratory diseases (J40-J47)	429	2.6	195.9
9	Alzheimer's disease (G30)	423	2.6	193.2
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	299	1.8	136.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	1.0	77.2
12	Pneumonitis due to solids and liquids (J69)	139	0.8	63.5
13	Atherosclerosis (I70)	112	0.7	51.2
14	Aortic aneurysm and dissection (I71)	89	0.5	40.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	88	0.5	40.2
...	All other causes (Residual)	2,204	13.3	1,006.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85-89 years				
...	All causes	15,491	100.0	12,380.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,702	36.8	4,557.2
2	Malignant neoplasms (C00-C97)	2,065	13.3	1,650.4
3	Cerebrovascular diseases (I60-I69)	1,762	11.4	1,408.2
4	Diabetes mellitus (E10-E14)	702	4.5	561.1
5	Influenza and pneumonia (J10-J18)	502	3.2	401.2
6	Alzheimer's disease (G30)	501	3.2	400.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	437	2.8	349.3
8	Septicemia (A40-A41)	417	2.7	333.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	313	2.0	250.2
10	Chronic lower respiratory diseases (J40-J47)	311	2.0	248.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	159	1.0	127.1
12	Pneumonitis due to solids and liquids (J69)	146	0.9	116.7
13	Atherosclerosis (I70)	110	0.7	87.9
14	Aortic aneurysm and dissection (I71)	79	0.5	63.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.5	58.3
...	All other causes (Residual)	2,212	14.3	1,767.9
Black, female, 90-94 years				
...	All causes	10,431	100.0	15,957.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,006	38.4	6,128.3
2	Cerebrovascular diseases (I60-I69)	1,145	11.0	1,751.6
3	Malignant neoplasms (C00-C97)	1,042	10.0	1,594.0
4	Influenza and pneumonia (J10-J18)	438	4.2	670.0
5	Diabetes mellitus (E10-E14)	398	3.8	608.9
6	Alzheimer's disease (G30)	365	3.5	558.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	287	2.8	439.0
8	Septicemia (A40-A41)	271	2.6	414.6
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	208	2.0	318.2
10	Chronic lower respiratory diseases (J40-J47)	187	1.8	286.1
11	Pneumonitis due to solids and liquids (J69)	113	1.1	172.9
12	Atherosclerosis (I70)	112	1.1	171.3
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	1.1	171.3
14	Nutritional deficiencies (E40-E64)	49	0.5	75.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.4	68.8
...	All other causes (Residual)	1,653	15.8	2,528.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 95-99 years				
...	All causes	4,875	100.0	17,601.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,955	40.1	7,058.5
2	Cerebrovascular diseases (I60-I69)	543	11.1	1,960.5
3	Malignant neoplasms (C00-C97)	372	7.6	1,343.1
4	Influenza and pneumonia (J10-J18)	229	4.7	826.8
5	Alzheimer's disease (G30)	183	3.8	660.7
6	Septicemia (A40-A41)	153	3.1	552.4
7	Diabetes mellitus (E10-E14)	139	2.9	501.9
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	115	2.4	415.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	2.3	400.8
10	Chronic lower respiratory diseases (J40-J47)	73	1.5	263.6
11	Atherosclerosis (I70)	57	1.2	205.8
12	Pneumonitis due to solids and liquids (J69)	53	1.1	191.4
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	1.0	173.3
14	Nutritional deficiencies (E40-E64)	30	0.6	108.3
15	Anemias (D50-D64)	23	0.5	83.0
...	All other causes (Residual)	791	16.2	2,855.9
Black, female, 100 years and over				
...	All causes	1,681	100.0	25,085.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	681	40.5	10,162.7
2	Cerebrovascular diseases (I60-I69)	176	10.5	2,626.5
3	Influenza and pneumonia (J10-J18)	97	5.8	1,447.5
4	Malignant neoplasms (C00-C97)	80	4.8	1,193.9
5	Alzheimer's disease (G30)	52	3.1	776.0
6	Diabetes mellitus (E10-E14)	47	2.8	701.4
7	Septicemia (A40-A41)	43	2.6	641.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	2.3	582.0
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	36	2.1	537.2
10	Atherosclerosis (I70)	35	2.1	522.3
11	Chronic lower respiratory diseases (J40-J47)	23	1.4	343.2
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	1.2	313.4
13	Pneumonitis due to solids and liquids (J69)	17	1.0	*
14	Nutritional deficiencies (E40-E64)	15	0.9	*
15	Anemias (D50-D64)	9	0.5	*
...	All other causes (Residual)	310	18.4	4,626.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, all ages				
...	All causes	11,363	100.0	466.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,417	21.3	99.2
2	Malignant neoplasms (C00-C97)	1,914	16.8	78.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,353	11.9	55.5
4	Diabetes mellitus (E10-E14)	616	5.4	25.3
5	Cerebrovascular diseases (I60-I69)	572	5.0	23.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	534	4.7	21.9
7	Chronic lower respiratory diseases (J40-J47)	429	3.8	17.6
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	297	2.6	12.2
9	Influenza and pneumonia (J10-J18)	289	2.5	11.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	215	1.9	8.8
11	Assault (homicide) (X85-Y09,Y87.1)	203	1.8	8.3
12	Septicemia (A40-A41)	162	1.4	6.6
13	Certain conditions originating in the perinatal period (P00-P96)	124	1.1	5.1
14	Alzheimer's disease (G30)	95	0.8	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	0.8	3.9
...	All other causes (Residual)	2,048	18.0	84.1
American Indian, both sexes, 1-4 years				
...	All causes	91	100.0	55.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	42.9	23.9
2	Assault (homicide) (X85-Y09,Y87.1)	8	8.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	4.4	*
4	Septicemia (A40-A41)	3	3.3	*
4	Malignant neoplasms (C00-C97)	3	3.3	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.2	*
6	Certain conditions originating in the perinatal period (P00-P96)	2	2.2	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1	1.1	*
8	Parkinson's disease (G20-G21)	1	1.1	*
8	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.1	*
8	Cerebrovascular diseases (I60-I69)	1	1.1	*
8	Diseases of appendix (K35-K38)	1	1.1	*
...	All other causes (Residual)	25	27.5	15.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 5-9 years				
...	All causes	50	100.0	23.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	62.0	14.6
2	Malignant neoplasms (C00-C97)	5	10.0	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.0	*
4	Influenza and pneumonia (J10-J18)	2	4.0	*
5	Septicemia (A40-A41)	1	2.0	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.0	*
5	Cerebrovascular diseases (I60-I69)	1	2.0	*
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	1	2.0	*
5	Assault (homicide) (X85-Y09,Y87.1)	1	2.0	*
...	All other causes (Residual)	3	6.0	*
American Indian, both sexes, 10-14 years				
...	All causes	63	100.0	24.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	61.9	15.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	6	9.5	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	6.3	*
4	Malignant neoplasms (C00-C97)	3	4.8	*
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1	1.6	*
5	Cerebrovascular diseases (I60-I69)	1	1.6	*
5	Influenza and pneumonia (J10-J18)	1	1.6	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.6	*
...	All other causes (Residual)	7	11.1	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-19 years				
...	All causes	251	100.0	105.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	59.4	62.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	44	17.5	18.4
3	Assault (homicide) (X85-Y09,Y87.1)	25	10.0	10.5
4	Malignant neoplasms (C00-C97)	6	2.4	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	0.8	*
7	Meningococcal infection (A39)	1	0.4	*
7	Septicemia (A40-A41)	1	0.4	*
7	Diabetes mellitus (E10-E14)	1	0.4	*
7	Influenza and pneumonia (J10-J18)	1	0.4	*
7	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
7	Pregnancy, childbirth and the puerperium (O00-099)	1	0.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
7	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (Residual)	13	5.2	*
American Indian, both sexes, 20-24 years				
...	All causes	268	100.0	133.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	56.0	74.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	45	16.8	22.3
3	Assault (homicide) (X85-Y09,Y87.1)	26	9.7	12.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.4	*
5	Malignant neoplasms (C00-C97)	7	2.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.7	*
6	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.7	*
9	Septicemia (A40-A41)	1	0.4	*
9	Diabetes mellitus (E10-E14)	1	0.4	*
9	Cerebrovascular diseases (I60-I69)	1	0.4	*
9	Influenza and pneumonia (J10-J18)	1	0.4	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
9	Pregnancy, childbirth and the puerperium (O00-099)	1	0.4	*
...	All other causes (Residual)	19	7.1	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 25-29 years				
...	All causes	272	100.0	140.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	45.2	63.7
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	43	15.8	22.3
3	Assault (homicide) (X85-Y09,Y87.1)	28	10.3	14.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	10	3.7	*
5	Malignant neoplasms (C00-C97)	9	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.9	*
7	Influenza and pneumonia (J10-J18)	4	1.5	*
7	Chronic lower respiratory diseases (J40-J47)	4	1.5	*
9	Cerebrovascular diseases (I60-I69)	3	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
12	Septicemia (A40-A41)	1	0.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.4	*
12	Pregnancy, childbirth and the puerperium (O00-099)	1	0.4	*
12	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (Residual)	31	11.4	16.0
American Indian, both sexes, 30-34 years				
...	All causes	324	100.0	177.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	35.8	63.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	35	10.8	19.1
3	Assault (homicide) (X85-Y09,Y87.1)	31	9.6	16.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	29	9.0	15.8
5	Malignant neoplasms (C00-C97)	17	5.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	4.9	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	7	2.2	*
7	Diabetes mellitus (E10-E14)	7	2.2	*
9	Viral hepatitis (B15-B19)	4	1.2	*
10	Pregnancy, childbirth and the puerperium (O00-099)	3	0.9	*
11	Septicemia (A40-A41)	2	0.6	*
11	Cerebrovascular diseases (I60-I69)	2	0.6	*
13	Influenza and pneumonia (J10-J18)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.3	*
...	All other causes (Residual)	50	15.4	27.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, both sexes, 35-39 years			
...	All causes	472	100.0	255.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	27.3	69.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	56	11.9	30.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	9.3	23.8
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	37	7.8	20.0
5	Malignant neoplasms (C00-C97)	34	7.2	18.4
6	Assault (homicide) (X85-Y09,Y87.1)	24	5.1	13.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	12	2.5	*
8	Influenza and pneumonia (J10-J18)	11	2.3	*
9	Diabetes mellitus (E10-E14)	7	1.5	*
9	Cerebrovascular diseases (I60-I69)	7	1.5	*
11	Septicemia (A40-A41)	6	1.3	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.8	*
13	Viral hepatitis (B15-B19)	3	0.6	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.4	*
...	All other causes (Residual)	93	19.7	50.3
	American Indian, both sexes, 40-44 years			
...	All causes	609	100.0	345.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	20.9	72.0
2	Chronic liver disease and cirrhosis (K70,K73-K74)	81	13.3	45.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	12.8	44.2
4	Malignant neoplasms (C00-C97)	71	11.7	40.2
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	30	4.9	17.0
6	Assault (homicide) (X85-Y09,Y87.1)	22	3.6	12.5
7	Influenza and pneumonia (J10-J18)	17	2.8	*
8	Diabetes mellitus (E10-E14)	16	2.6	*
9	Cerebrovascular diseases (I60-I69)	15	2.5	*
10	Viral hepatitis (B15-B19)	12	2.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	12	2.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.8	*
13	Septicemia (A40-A41)	7	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	6	1.0	*
15	Anemias (D50-D64)	2	0.3	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.3	*
...	All other causes (Residual)	98	16.1	55.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 45-49 years				
...	All causes	611	100.0	413.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	17.7	73.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	16.9	69.6
3	Malignant neoplasms (C00-C97)	88	14.4	59.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	72	11.8	48.7
5	Diabetes mellitus (E10-E14)	25	4.1	16.9
6	Cerebrovascular diseases (I60-I69)	22	3.6	14.9
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	16	2.6	*
8	Septicemia (A40-A41)	12	2.0	*
8	Viral hepatitis (B15-B19)	12	2.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	12	2.0	*
11	Chronic lower respiratory diseases (J40-J47)	10	1.6	*
12	Assault (homicide) (X85-Y09,Y87.1)	9	1.5	*
13	Influenza and pneumonia (J10-J18)	8	1.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
15	Tuberculosis (A16-A19)	2	0.3	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.3	*
15	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	Peptic ulcer (K25-K28)	2	0.3	*
...	All other causes (Residual)	101	16.5	68.3
American Indian, both sexes, 50-54 years				
...	All causes	743	100.0	628.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	146	19.7	123.6
2	Malignant neoplasms (C00-C97)	132	17.8	111.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	12.0	75.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	75	10.1	63.5
5	Diabetes mellitus (E10-E14)	53	7.1	44.9
6	Cerebrovascular diseases (I60-I69)	25	3.4	21.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.3	*
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	15	2.0	*
9	Viral hepatitis (B15-B19)	13	1.7	*
9	Influenza and pneumonia (J10-J18)	13	1.7	*
11	Septicemia (A40-A41)	9	1.2	*
11	Assault (homicide) (X85-Y09,Y87.1)	9	1.2	*
13	Chronic lower respiratory diseases (J40-J47)	7	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.4	*
15	Peptic ulcer (K25-K28)	3	0.4	*
...	All other causes (Residual)	128	17.2	108.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-59 years				
...	All causes	813	100.0	941.7
1	Malignant neoplasms (C00-C97)	190	23.4	220.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	22.1	208.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	69	8.5	79.9
4	Diabetes mellitus (E10-E14)	59	7.3	68.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	6.8	63.7
6	Chronic lower respiratory diseases (J40-J47)	36	4.4	41.7
7	Cerebrovascular diseases (I60-I69)	32	3.9	37.1
8	Septicemia (A40-A41)	18	2.2	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	2.1	*
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	13	1.6	*
11	Influenza and pneumonia (J10-J18)	6	0.7	*
12	Viral hepatitis (B15-B19)	4	0.5	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.5	*
12	Pneumonitis due to solids and liquids (J69)	4	0.5	*
15	Tuberculosis (A16-A19)	3	0.4	*
...	All other causes (Residual)	123	15.1	142.5
American Indian, both sexes, 60-64 years				
...	All causes	885	100.0	1,337.6
1	Malignant neoplasms (C00-C97)	227	25.6	343.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	214	24.2	323.4
3	Diabetes mellitus (E10-E14)	75	8.5	113.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	54	6.1	81.6
5	Cerebrovascular diseases (I60-I69)	47	5.3	71.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	4.4	58.9
7	Chronic lower respiratory diseases (J40-J47)	35	4.0	52.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.5	33.3
9	Influenza and pneumonia (J10-J18)	18	2.0	*
10	Septicemia (A40-A41)	11	1.2	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	1.1	*
12	Viral hepatitis (B15-B19)	6	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
13	Assault (homicide) (X85-Y09,Y87.1)	4	0.5	*
15	Tuberculosis (A16-A19)	3	0.3	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
15	Peptic ulcer (K25-K28)	3	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.3	*
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	3	0.3	*
...	All other causes (Residual)	104	11.8	157.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 65-69 years				
...	All causes	1,049	100.0	2,042.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	291	27.7	566.6
2	Malignant neoplasms (C00-C97)	251	23.9	488.7
3	Diabetes mellitus (E10-E14)	77	7.3	149.9
4	Chronic lower respiratory diseases (J40-J47)	58	5.5	112.9
5	Cerebrovascular diseases (I60-I69)	44	4.2	85.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	39	3.7	75.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	3.4	70.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	3.0	60.4
9	Septicemia (A40-A41)	22	2.1	42.8
10	Influenza and pneumonia (J10-J18)	20	1.9	38.9
11	Aortic aneurysm and dissection (I71)	13	1.2	*
12	Atherosclerosis (I70)	8	0.8	*
13	Tuberculosis (A16-A19)	4	0.4	*
13	Viral hepatitis (B15-B19)	4	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.4	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.4	*
13	Pneumonitis due to solids and liquids (J69)	4	0.4	*
13	Peptic ulcer (K25-K28)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	131	12.5	255.1
American Indian, both sexes, 70-74 years				
...	All causes	1,092	100.0	2,654.8
1	Malignant neoplasms (C00-C97)	288	26.4	700.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	257	23.5	624.8
3	Diabetes mellitus (E10-E14)	109	10.0	265.0
4	Chronic lower respiratory diseases (J40-J47)	66	6.0	160.5
5	Cerebrovascular diseases (I60-I69)	64	5.9	155.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	2.9	77.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	2.7	70.5
8	Influenza and pneumonia (J10-J18)	25	2.3	60.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	20	1.8	48.6
10	Alzheimer's disease (G30)	12	1.1	*
11	Septicemia (A40-A41)	11	1.0	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	0.8	*
13	Atherosclerosis (I70)	8	0.7	*
13	Aortic aneurysm and dissection (I71)	8	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
15	Parkinson's disease (G20-G21)	6	0.5	*
...	All other causes (Residual)	142	13.0	345.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-79 years				
...	All causes	1,130	100.0	3,460.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	326	28.8	998.4
2	Malignant neoplasms (C00-C97)	248	21.9	759.5
3	Cerebrovascular diseases (I60-I69)	86	7.6	263.4
4	Diabetes mellitus (E10-E14)	81	7.2	248.1
5	Chronic lower respiratory diseases (J40-J47)	66	5.8	202.1
6	Influenza and pneumonia (J10-J18)	37	3.3	113.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	2.6	88.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	1.9	67.4
9	Alzheimer's disease (G30)	21	1.9	64.3
10	Septicemia (A40-A41)	19	1.7	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.7	*
12	Parkinson's disease (G20-G21)	9	0.8	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	0.8	*
12	Aortic aneurysm and dissection (I71)	9	0.8	*
15	Anemias (D50-D64)	6	0.5	*
15	Pneumonitis due to solids and liquids (J69)	6	0.5	*
...	All other causes (Residual)	137	12.1	419.6
American Indian, both sexes, 80-84 years				
...	All causes	932	100.0	4,689.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	271	29.1	1,363.6
2	Malignant neoplasms (C00-C97)	164	17.6	825.2
3	Cerebrovascular diseases (I60-I69)	92	9.9	462.9
4	Chronic lower respiratory diseases (J40-J47)	60	6.4	301.9
5	Diabetes mellitus (E10-E14)	52	5.6	261.6
6	Influenza and pneumonia (J10-J18)	34	3.6	171.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	2.6	120.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	2.4	110.7
9	Alzheimer's disease (G30)	18	1.9	*
10	Septicemia (A40-A41)	15	1.6	*
11	Parkinson's disease (G20-G21)	11	1.2	*
12	Atherosclerosis (I70)	10	1.1	*
13	Pneumonitis due to solids and liquids (J69)	8	0.9	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	0.8	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.6	*
...	All other causes (Residual)	138	14.8	694.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, both sexes, 85-89 years			
...	All causes	728	100.0	6,056.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231	31.7	1,921.8
2	Malignant neoplasms (C00-C97)	103	14.1	856.9
3	Cerebrovascular diseases (I60-I69)	66	9.1	549.1
4	Chronic lower respiratory diseases (J40-J47)	49	6.7	407.7
5	Diabetes mellitus (E10-E14)	36	4.9	299.5
5	Influenza and pneumonia (J10-J18)	36	4.9	299.5
7	Alzheimer's disease (G30)	19	2.6	*
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13	1.8	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	1.8	*
10	Septicemia (A40-A41)	12	1.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.5	*
12	Nutritional deficiencies (E40-E64)	8	1.1	*
12	Atherosclerosis (I70)	8	1.1	*
12	Pneumonitis due to solids and liquids (J69)	8	1.1	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
15	Parkinson's disease (G20-G21)	6	0.8	*
...	All other causes (Residual)	103	14.1	856.9
	American Indian, both sexes, 90-94 years			
...	All causes	442	100.0	7,101.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	150	33.9	2,410.0
2	Cerebrovascular diseases (I60-I69)	47	10.6	755.1
3	Malignant neoplasms (C00-C97)	46	10.4	739.1
4	Influenza and pneumonia (J10-J18)	36	8.1	578.4
5	Alzheimer's disease (G30)	15	3.4	*
5	Chronic lower respiratory diseases (J40-J47)	15	3.4	*
7	Diabetes mellitus (E10-E14)	13	2.9	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	2.9	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.4	*
10	Parkinson's disease (G20-G21)	5	1.1	*
10	Pneumonitis due to solids and liquids (J69)	5	1.1	*
12	Septicemia (A40-A41)	4	0.9	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.9	*
14	Anemias (D50-D64)	3	0.7	*
14	Nutritional deficiencies (E40-E64)	3	0.7	*
14	Atherosclerosis (I70)	3	0.7	*
14	Aortic aneurysm and dissection (I71)	3	0.7	*
...	All other causes (Residual)	71	16.1	1,140.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 95-99 years				
...	All causes	169	100.0	6,273.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	38.5	2,412.8
2	Malignant neoplasms (C00-C97)	16	9.5	*
2	Cerebrovascular diseases (I60-I69)	16	9.5	*
4	Influenza and pneumonia (J10-J18)	9	5.3	*
5	Septicemia (A40-A41)	6	3.6	*
6	Alzheimer's disease (G30)	5	3.0	*
7	Atherosclerosis (I70)	4	2.4	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.8	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3	1.8	*
10	Diabetes mellitus (E10-E14)	2	1.2	*
10	Parkinson's disease (G20-G21)	2	1.2	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	1.2	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
14	Nutritional deficiencies (E40-E64)	1	0.6	*
14	Aortic aneurysm and dissection (I71)	1	0.6	*
14	Pneumonitis due to solids and liquids (J69)	1	0.6	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.6	*
14	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.6	*
...	All other causes (Residual)	28	16.6	1,039.3
American Indian, both sexes, 100 years and over				
...	All causes	43	100.0	5,858.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	30.2	*
2	Malignant neoplasms (C00-C97)	4	9.3	*
2	Chronic lower respiratory diseases (J40-J47)	4	9.3	*
4	Nutritional deficiencies (E40-E64)	3	7.0	*
4	Influenza and pneumonia (J10-J18)	3	7.0	*
6	Diabetes mellitus (E10-E14)	2	4.7	*
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	4.7	*
6	Atherosclerosis (I70)	2	4.7	*
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	4.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.3	*
10	Pneumonitis due to solids and liquids (J69)	1	2.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.3	*
10	Infections of kidney (N10-N12,N13.6,N15.1)	1	2.3	*
...	All other causes (Residual)	4	9.3	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	6,185	100.0	512.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,341	21.7	111.2
2	Malignant neoplasms (C00-C97)	997	16.1	82.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	900	14.6	74.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	298	4.8	24.7
5	Diabetes mellitus (E10-E14)	275	4.4	22.8
6	Cerebrovascular diseases (I60-I69)	250	4.0	20.7
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	237	3.8	19.6
8	Chronic lower respiratory diseases (J40-J47)	228	3.7	18.9
9	Assault (homicide) (X85-Y09,Y87.1)	159	2.6	13.2
10	Influenza and pneumonia (J10-J18)	149	2.4	12.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	1.6	8.1
12	Septicemia (A40-A41)	74	1.2	6.1
13	Certain conditions originating in the perinatal period (P00-P96)	72	1.2	6.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.9	4.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	43	0.7	3.6
...	All other causes (Residual)	1,008	16.3	83.6
American Indian, male, 1-4 years				
...	All causes	49	100.0	59.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	51.0	30.3
2	Assault (homicide) (X85-Y09,Y87.1)	6	12.2	*
3	Septicemia (A40-A41)	2	4.1	*
3	Malignant neoplasms (C00-C97)	2	4.1	*
5	Cerebrovascular diseases (I60-I69)	1	2.0	*
5	Chronic lower respiratory diseases (J40-J47)	1	2.0	*
5	Certain conditions originating in the perinatal period (P00-P96)	1	2.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.0	*
...	All other causes (Residual)	10	20.4	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 5-9 years				
...	All causes	28	100.0	26.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16	57.1	*
2	Malignant neoplasms (C00-C97)	3	10.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	7.1	*
4	Septicemia (A40-A41)	1	3.6	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	3.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	3.6	*
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	1	3.6	*
4	Assault (homicide) (X85-Y09,Y87.1)	1	3.6	*
...	All other causes (Residual)	2	7.1	*
American Indian, male, 10-14 years				
...	All causes	33	100.0	25.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	63.6	16.3
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	5	15.2	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	6.1	*
4	Malignant neoplasms (C00-C97)	1	3.0	*
...	All other causes (Residual)	4	12.1	*
American Indian, male, 15-19 years				
...	All causes	178	100.0	148.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	57.3	85.1
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	34	19.1	28.4
3	Assault (homicide) (X85-Y09,Y87.1)	21	11.8	17.5
4	Malignant neoplasms (C00-C97)	4	2.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.1	*
6	Septicemia (A40-A41)	1	0.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
6	Diabetes mellitus (E10-E14)	1	0.6	*
6	Influenza and pneumonia (J10-J18)	1	0.6	*
6	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
6	Legal intervention (Y35,Y89.0)	1	0.6	*
...	All other causes (Residual)	8	4.5	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 20-24 years				
...	All causes	191	100.0	189.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	52.4	99.0
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	37	19.4	36.6
3	Assault (homicide) (X85-Y09,Y87.1)	25	13.1	24.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	3.1	*
5	Malignant neoplasms (C00-C97)	3	1.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.0	*
6	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.0	*
9	Diabetes mellitus (E10-E14)	1	0.5	*
...	All other causes (Residual)	13	6.8	*
American Indian, male, 25-29 years				
...	All causes	192	100.0	193.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	49.0	94.8
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	33	17.2	33.3
3	Assault (homicide) (X85-Y09,Y87.1)	22	11.5	22.2
4	Malignant neoplasms (C00-C97)	5	2.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	2.6	*
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.6	*
8	Influenza and pneumonia (J10-J18)	2	1.0	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	1.0	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
10	Legal intervention (Y35,Y89.0)	1	0.5	*
...	All other causes (Residual)	18	9.4	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 30-34 years				
...	All causes	218	100.0	232.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	35.8	83.1
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	26	11.9	27.7
3	Assault (homicide) (X85-Y09,Y87.1)	24	11.0	25.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	18	8.3	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	5.0	*
6	Malignant neoplasms (C00-C97)	8	3.7	*
7	Diabetes mellitus (E10-E14)	6	2.8	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.8	*
9	Septicemia (A40-A41)	2	0.9	*
9	Viral hepatitis (B15-B19)	2	0.9	*
11	Influenza and pneumonia (J10-J18)	1	0.5	*
11	Pneumonitis due to solids and liquids (J69)	1	0.5	*
...	All other causes (Residual)	37	17.0	39.4
American Indian, male, 35-39 years				
...	All causes	284	100.0	305.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	28.5	87.1
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	31	10.9	33.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	27	9.5	29.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	9.2	28.0
5	Malignant neoplasms (C00-C97)	16	5.6	*
5	Assault (homicide) (X85-Y09,Y87.1)	16	5.6	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	9	3.2	*
8	Diabetes mellitus (E10-E14)	6	2.1	*
8	Influenza and pneumonia (J10-J18)	6	2.1	*
10	Cerebrovascular diseases (I60-I69)	4	1.4	*
11	Septicemia (A40-A41)	2	0.7	*
11	Viral hepatitis (B15-B19)	2	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
13	Meningitis (G00,G03)	1	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
13	Peptic ulcer (K25-K28)	1	0.4	*
13	Hernia (K40-K46)	1	0.4	*
...	All other causes (Residual)	53	18.7	57.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 40-44 years				
...	All causes	391	100.0	449.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	21.7	97.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	14.6	65.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	51	13.0	58.7
4	Malignant neoplasms (C00-C97)	33	8.4	38.0
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	22	5.6	25.3
6	Assault (homicide) (X85-Y09,Y87.1)	16	4.1	*
7	Influenza and pneumonia (J10-J18)	12	3.1	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	11	2.8	*
9	Diabetes mellitus (E10-E14)	10	2.6	*
10	Cerebrovascular diseases (I60-I69)	7	1.8	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.8	*
12	Viral hepatitis (B15-B19)	6	1.5	*
13	Septicemia (A40-A41)	5	1.3	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.5	*
...	All other causes (Residual)	64	16.4	73.6
American Indian, male, 45-49 years				
...	All causes	389	100.0	542.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	20.6	111.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	19.8	107.3
3	Malignant neoplasms (C00-C97)	44	11.3	61.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	42	10.8	58.5
5	Diabetes mellitus (E10-E14)	18	4.6	*
6	Human immunodeficiency virus (HIV) disease (B20-B24)	11	2.8	*
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	11	2.8	*
8	Cerebrovascular diseases (I60-I69)	10	2.6	*
9	Septicemia (A40-A41)	7	1.8	*
9	Viral hepatitis (B15-B19)	7	1.8	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.8	*
9	Assault (homicide) (X85-Y09,Y87.1)	7	1.8	*
13	Influenza and pneumonia (J10-J18)	5	1.3	*
14	Tuberculosis (A16-A19)	2	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
...	All other causes (Residual)	59	15.2	82.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 50-54 years				
...	All causes	469	100.0	824.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	102	21.7	179.3
2	Malignant neoplasms (C00-C97)	72	15.4	126.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	14.9	123.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	49	10.4	86.1
5	Diabetes mellitus (E10-E14)	23	4.9	40.4
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	13	2.8	*
7	Cerebrovascular diseases (I60-I69)	12	2.6	*
8	Influenza and pneumonia (J10-J18)	11	2.3	*
9	Viral hepatitis (B15-B19)	9	1.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.9	*
11	Assault (homicide) (X85-Y09,Y87.1)	7	1.5	*
12	Septicemia (A40-A41)	5	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.6	*
14	Peptic ulcer (K25-K28)	3	0.6	*
...	All other causes (Residual)	76	16.2	133.6
American Indian, male, 55-59 years				
...	All causes	459	100.0	1,123.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	118	25.7	288.8
2	Malignant neoplasms (C00-C97)	106	23.1	259.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	39	8.5	95.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	7.4	83.2
5	Diabetes mellitus (E10-E14)	26	5.7	63.6
6	Chronic lower respiratory diseases (J40-J47)	19	4.1	*
7	Cerebrovascular diseases (I60-I69)	17	3.7	*
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	12	2.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	2.0	*
10	Septicemia (A40-A41)	8	1.7	*
11	Influenza and pneumonia (J10-J18)	3	0.7	*
12	Tuberculosis (A16-A19)	2	0.4	*
12	Pneumonitis due to solids and liquids (J69)	2	0.4	*
14	Viral hepatitis (B15-B19)	1	0.2	*
14	Anemias (D50-D64)	1	0.2	*
14	Parkinson's disease (G20-G21)	1	0.2	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.2	*
14	Atherosclerosis (I70)	1	0.2	*
14	Aortic aneurysm and dissection (I71)	1	0.2	*
14	Peptic ulcer (K25-K28)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.2	*
14	Assault (homicide) (X85-Y09,Y87.1)	1	0.2	*
...	All other causes (Residual)	53	11.5	129.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 60-64 years				
...	All causes	505	100.0	1,645.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	134	26.5	436.6
2	Malignant neoplasms (C00-C97)	120	23.8	390.9
3	Diabetes mellitus (E10-E14)	42	8.3	136.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	32	6.3	104.3
5	Cerebrovascular diseases (I60-I69)	26	5.1	84.7
6	Chronic lower respiratory diseases (J40-J47)	22	4.4	71.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	4.4	71.7
8	Influenza and pneumonia (J10-J18)	13	2.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	2.4	*
10	Septicemia (A40-A41)	6	1.2	*
11	Viral hepatitis (B15-B19)	4	0.8	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	3	0.6	*
13	Assault (homicide) (X85-Y09,Y87.1)	3	0.6	*
...	All other causes (Residual)	56	11.1	182.4
American Indian, male, 65-69 years				
...	All causes	556	100.0	2,402.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	175	31.5	756.1
2	Malignant neoplasms (C00-C97)	136	24.5	587.6
3	Diabetes mellitus (E10-E14)	33	5.9	142.6
4	Cerebrovascular diseases (I60-I69)	27	4.9	116.7
5	Chronic lower respiratory diseases (J40-J47)	24	4.3	103.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	3.2	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	3.1	*
8	Influenza and pneumonia (J10-J18)	15	2.7	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	15	2.7	*
10	Septicemia (A40-A41)	10	1.8	*
11	Aortic aneurysm and dissection (I71)	7	1.3	*
12	Pneumonitis due to solids and liquids (J69)	4	0.7	*
13	Atherosclerosis (I70)	3	0.5	*
13	Peptic ulcer (K25-K28)	3	0.5	*
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	3	0.5	*
...	All other causes (Residual)	66	11.9	285.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 70-74 years				
...	All causes	557	100.0	3,020.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	161	28.9	873.1
2	Malignant neoplasms (C00-C97)	155	27.8	840.6
3	Cerebrovascular diseases (I60-I69)	38	6.8	206.1
4	Diabetes mellitus (E10-E14)	37	6.6	200.7
5	Chronic lower respiratory diseases (J40-J47)	26	4.7	141.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	3.4	*
7	Influenza and pneumonia (J10-J18)	14	2.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.8	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.1	*
10	Septicemia (A40-A41)	5	0.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	5	0.9	*
13	Aortic aneurysm and dissection (I71)	4	0.7	*
14	Parkinson's disease (G20-G21)	3	0.5	*
14	Alzheimer's disease (G30)	3	0.5	*
14	Atherosclerosis (I70)	3	0.5	*
14	Pneumonitis due to solids and liquids (J69)	3	0.5	*
...	All other causes (Residual)	60	10.8	325.4
American Indian, male, 75-79 years				
...	All causes	567	100.0	3,999.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	31.7	1,269.8
2	Malignant neoplasms (C00-C97)	133	23.5	938.2
3	Chronic lower respiratory diseases (J40-J47)	39	6.9	275.1
4	Diabetes mellitus (E10-E14)	37	6.5	261.0
5	Cerebrovascular diseases (I60-I69)	30	5.3	211.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	3.5	141.1
7	Influenza and pneumonia (J10-J18)	18	3.2	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	1.2	*
10	Septicemia (A40-A41)	6	1.1	*
10	Alzheimer's disease (G30)	6	1.1	*
12	Anemias (D50-D64)	5	0.9	*
12	Parkinson's disease (G20-G21)	5	0.9	*
12	Aortic aneurysm and dissection (I71)	5	0.9	*
12	Pneumonitis due to solids and liquids (J69)	5	0.9	*
...	All other causes (Residual)	60	10.6	423.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 80-84 years				
...	All causes	422	100.0	5,217.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	29.9	1,557.9
2	Malignant neoplasms (C00-C97)	80	19.0	989.1
3	Cerebrovascular diseases (I60-I69)	39	9.2	482.2
4	Chronic lower respiratory diseases (J40-J47)	38	9.0	469.8
5	Influenza and pneumonia (J10-J18)	15	3.6	*
6	Diabetes mellitus (E10-E14)	13	3.1	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	2.8	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	2.1	*
9	Septicemia (A40-A41)	8	1.9	*
10	Parkinson's disease (G20-G21)	7	1.7	*
10	Alzheimer's disease (G30)	7	1.7	*
12	Atherosclerosis (I70)	5	1.2	*
13	Pneumonitis due to solids and liquids (J69)	3	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	0.5	*
14	Aortic aneurysm and dissection (I71)	2	0.5	*
14	Peptic ulcer (K25-K28)	2	0.5	*
...	All other causes (Residual)	52	12.3	642.9
American Indian, male, 85-89 years				
...	All causes	286	100.0	6,833.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	31.8	2,174.4
2	Malignant neoplasms (C00-C97)	44	15.4	1,051.4
3	Cerebrovascular diseases (I60-I69)	24	8.4	573.5
3	Chronic lower respiratory diseases (J40-J47)	24	8.4	573.5
5	Diabetes mellitus (E10-E14)	15	5.2	*
6	Influenza and pneumonia (J10-J18)	11	3.8	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	3.1	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.7	*
9	Parkinson's disease (G20-G21)	4	1.4	*
9	Alzheimer's disease (G30)	4	1.4	*
9	Pneumonitis due to solids and liquids (J69)	4	1.4	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	1.0	*
12	Atherosclerosis (I70)	3	1.0	*
14	Septicemia (A40-A41)	2	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.7	*
14	Assault (homicide) (X85-Y09,Y87.1)	2	0.7	*
...	All other causes (Residual)	39	13.6	931.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 90-94 years				
...	All causes	155	100.0	8,630.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	29.7	2,561.2
2	Malignant neoplasms (C00-C97)	20	12.9	1,113.6
3	Influenza and pneumonia (J10-J18)	16	10.3	*
4	Cerebrovascular diseases (I60-I69)	12	7.7	*
5	Chronic lower respiratory diseases (J40-J47)	8	5.2	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	3.9	*
7	Diabetes mellitus (E10-E14)	5	3.2	*
8	Parkinson's disease (G20-G21)	4	2.6	*
8	Alzheimer's disease (G30)	4	2.6	*
10	Septicemia (A40-A41)	2	1.3	*
10	Pneumonitis due to solids and liquids (J69)	2	1.3	*
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	1.3	*
13	Nutritional deficiencies (E40-E64)	1	0.6	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.6	*
13	Aortic aneurysm and dissection (I71)	1	0.6	*
13	Pneumoconioses and chemical effects (J60-J66,J68)	1	0.6	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.6	*
...	All other causes (Residual)	23	14.8	1,280.6
American Indian, male, 95-99 years				
...	All causes	50	100.0	7,052.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	36.0	*
2	Malignant neoplasms (C00-C97)	9	18.0	*
3	Influenza and pneumonia (J10-J18)	4	8.0	*
4	Cerebrovascular diseases (I60-I69)	3	6.0	*
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	4.0	*
6	Septicemia (A40-A41)	1	2.0	*
6	Diabetes mellitus (E10-E14)	1	2.0	*
6	Alzheimer's disease (G30)	1	2.0	*
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	2.0	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.0	*
6	Pneumonitis due to solids and liquids (J69)	1	2.0	*
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	2.0	*
...	All other causes (Residual)	7	14.0	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 100 years and over				
...	All causes	10	100.0	*
1	Chronic lower respiratory diseases (J40-J47)	3	30.0	*
2	Malignant neoplasms (C00-C97)	1	10.0	*
2	Diabetes mellitus (E10-E14)	1	10.0	*
2	Nutritional deficiencies (E40-E64)	1	10.0	*
2	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	10.0	*
2	Atherosclerosis (I70)	1	10.0	*
...	All other causes (Residual)	2	20.0	*
American Indian, female, all ages				
...	All causes	5,178	100.0	421.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,076	20.8	87.5
2	Malignant neoplasms (C00-C97)	917	17.7	74.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	453	8.7	36.8
4	Diabetes mellitus (E10-E14)	341	6.6	27.7
5	Cerebrovascular diseases (I60-I69)	322	6.2	26.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	236	4.6	19.2
7	Chronic lower respiratory diseases (J40-J47)	201	3.9	16.3
8	Influenza and pneumonia (J10-J18)	140	2.7	11.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	2.3	9.5
10	Septicemia (A40-A41)	88	1.7	7.2
11	Alzheimer's disease (G30)	69	1.3	5.6
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	60	1.2	4.9
13	Certain conditions originating in the perinatal period (P00-P96)	52	1.0	4.2
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	49	0.9	4.0
15	Assault (homicide) (X85-Y09,Y87.1)	44	0.8	3.6
...	All other causes (Residual)	1,013	19.6	82.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 1-4 years				
...	All causes	42	100.0	52.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	33.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.1	*
3	Assault (homicide) (X85-Y09,Y87.1)	2	4.8	*
4	Septicemia (A40-A41)	1	2.4	*
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1	2.4	*
4	Malignant neoplasms (C00-C97)	1	2.4	*
4	Parkinson's disease (G20-G21)	1	2.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.4	*
4	Chronic lower respiratory diseases (J40-J47)	1	2.4	*
4	Diseases of appendix (K35-K38)	1	2.4	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	2.4	*
...	All other causes (Residual)	15	35.7	*
American Indian, female, 5-9 years				
...	All causes	22	100.0	21.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15	68.2	*
2	Malignant neoplasms (C00-C97)	2	9.1	*
2	Influenza and pneumonia (J10-J18)	2	9.1	*
4	Cerebrovascular diseases (I60-I69)	1	4.5	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.5	*
...	All other causes (Residual)	1	4.5	*
American Indian, female, 10-14 years				
...	All causes	30	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	60.0	*
2	Malignant neoplasms (C00-C97)	2	6.7	*
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	6.7	*
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1	3.3	*
4	Cerebrovascular diseases (I60-I69)	1	3.3	*
4	Influenza and pneumonia (J10-J18)	1	3.3	*
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	1	3.3	*
4	Complications of medical and surgical care (Y40-Y84,Y88)	1	3.3	*
...	All other causes (Residual)	3	10.0	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 15-19 years				
...	All causes	73	100.0	61.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	64.4	39.6
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	10	13.7	*
3	Assault (homicide) (X85-Y09,Y87.1)	4	5.5	*
4	Malignant neoplasms (C00-C97)	2	2.7	*
5	Meningococcal infection (A39)	1	1.4	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.4	*
5	Pregnancy, childbirth and the puerperium (O00-099)	1	1.4	*
5	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.4	*
...	All other causes (Residual)	5	6.8	*
American Indian, female, 20-24 years				
...	All causes	77	100.0	76.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	64.9	49.7
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	8	10.4	*
3	Malignant neoplasms (C00-C97)	4	5.2	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.9	*
5	Septicemia (A40-A41)	1	1.3	*
5	Cerebrovascular diseases (I60-I69)	1	1.3	*
5	Influenza and pneumonia (J10-J18)	1	1.3	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1	1.3	*
5	Pregnancy, childbirth and the puerperium (O00-099)	1	1.3	*
5	Assault (homicide) (X85-Y09,Y87.1)	1	1.3	*
...	All other causes (Residual)	6	7.8	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 25-29 years			
...	All causes	80	100.0	85.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	36.3	30.8
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	10	12.5	*
3	Assault (homicide) (X85-Y09,Y87.1)	6	7.5	*
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5	6.3	*
5	Malignant neoplasms (C00-C97)	4	5.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.8	*
6	Cerebrovascular diseases (I60-I69)	3	3.8	*
8	Influenza and pneumonia (J10-J18)	2	2.5	*
9	Septicemia (A40-A41)	1	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.3	*
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.3	*
9	Pregnancy, childbirth and the puerperium (O00-099)	1	1.3	*
...	All other causes (Residual)	13	16.3	*
	American Indian, female, 30-34 years			
...	All causes	106	100.0	118.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	35.8	42.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	11	10.4	*
3	Malignant neoplasms (C00-C97)	9	8.5	*
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	9	8.5	*
5	Assault (homicide) (X85-Y09,Y87.1)	7	6.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	4.7	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	3	2.8	*
7	Pregnancy, childbirth and the puerperium (O00-099)	3	2.8	*
9	Viral hepatitis (B15-B19)	2	1.9	*
9	Cerebrovascular diseases (I60-I69)	2	1.9	*
11	Diabetes mellitus (E10-E14)	1	0.9	*
11	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
11	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.9	*
...	All other causes (Residual)	13	12.3	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 35-39 years			
...	All causes	188	100.0	204.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48	25.5	52.3
2	Chronic liver disease and cirrhosis (K70,K73-K74)	29	15.4	31.6
3	Malignant neoplasms (C00-C97)	18	9.6	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	9.6	*
5	Assault (homicide) (X85-Y09,Y87.1)	8	4.3	*
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	6	3.2	*
7	Influenza and pneumonia (J10-J18)	5	2.7	*
8	Septicemia (A40-A41)	4	2.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.6	*
9	Cerebrovascular diseases (I60-I69)	3	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	3	1.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.1	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
14	Viral hepatitis (B15-B19)	1	0.5	*
14	Diabetes mellitus (E10-E14)	1	0.5	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
14	Legal intervention (Y35,Y89.0)	1	0.5	*
...	All other causes (Residual)	34	18.1	37.1
	American Indian, female, 40-44 years			
...	All causes	218	100.0	243.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	19.3	46.9
2	Malignant neoplasms (C00-C97)	38	17.4	42.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	30	13.8	33.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	9.6	23.5
5	Cerebrovascular diseases (I60-I69)	8	3.7	*
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	8	3.7	*
7	Viral hepatitis (B15-B19)	6	2.8	*
7	Diabetes mellitus (E10-E14)	6	2.8	*
7	Assault (homicide) (X85-Y09,Y87.1)	6	2.8	*
10	Influenza and pneumonia (J10-J18)	5	2.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.8	*
12	Chronic lower respiratory diseases (J40-J47)	3	1.4	*
13	Septicemia (A40-A41)	2	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
14	Anemias (D50-D64)	1	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	34	15.6	38.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 45-49 years			
...	All causes	222	100.0	291.4
1	Malignant neoplasms (C00-C97)	44	19.8	57.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	30	13.5	39.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	12.6	36.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	11.7	34.1
5	Cerebrovascular diseases (I60-I69)	12	5.4	*
6	Diabetes mellitus (E10-E14)	7	3.2	*
7	Septicemia (A40-A41)	5	2.3	*
7	Viral hepatitis (B15-B19)	5	2.3	*
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	5	2.3	*
10	Influenza and pneumonia (J10-J18)	3	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	3	1.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.4	*
13	Pneumonitis due to solids and liquids (J69)	2	0.9	*
13	Peptic ulcer (K25-K28)	2	0.9	*
13	Assault (homicide) (X85-Y09,Y87.1)	2	0.9	*
...	All other causes (Residual)	45	20.3	59.1
	American Indian, female, 50-54 years			
...	All causes	274	100.0	447.4
1	Malignant neoplasms (C00-C97)	60	21.9	98.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	16.1	71.8
3	Diabetes mellitus (E10-E14)	30	10.9	49.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	26	9.5	42.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	6.9	*
6	Cerebrovascular diseases (I60-I69)	13	4.7	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	2.9	*
8	Septicemia (A40-A41)	4	1.5	*
8	Viral hepatitis (B15-B19)	4	1.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.1	*
11	Influenza and pneumonia (J10-J18)	2	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	2	0.7	*
11	Assault (homicide) (X85-Y09,Y87.1)	2	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
15	Nutritional deficiencies (E40-E64)	1	0.4	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.4	*
15	Pneumonitis due to solids and liquids (J69)	1	0.4	*
15	Hernia (K40-K46)	1	0.4	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	*
...	All other causes (Residual)	49	17.9	80.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 55-59 years			
...	All causes	354	100.0	778.4
1	Malignant neoplasms (C00-C97)	84	23.7	184.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	17.5	136.3
3	Diabetes mellitus (E10-E14)	33	9.3	72.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	30	8.5	66.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	5.9	46.2
6	Chronic lower respiratory diseases (J40-J47)	17	4.8	*
7	Cerebrovascular diseases (I60-I69)	15	4.2	*
8	Septicemia (A40-A41)	10	2.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	2.3	*
10	Viral hepatitis (B15-B19)	3	0.8	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.8	*
10	Influenza and pneumonia (J10-J18)	3	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
13	Pneumonitis due to solids and liquids (J69)	2	0.6	*
15	Tuberculosis (A16-A19)	1	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
15	Alzheimer's disease (G30)	1	0.3	*
15	Atherosclerosis (I70)	1	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	1	0.3	*
15	Assault (homicide) (X85-Y09,Y87.1)	1	0.3	*
...	All other causes (Residual)	54	15.3	118.7
	American Indian, female, 60-64 years			
...	All causes	380	100.0	1,071.4
1	Malignant neoplasms (C00-C97)	107	28.2	301.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	21.1	225.5
3	Diabetes mellitus (E10-E14)	33	8.7	93.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	22	5.8	62.0
5	Cerebrovascular diseases (I60-I69)	21	5.5	59.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	4.5	*
7	Chronic lower respiratory diseases (J40-J47)	13	3.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	2.6	*
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	1.6	*
10	Septicemia (A40-A41)	5	1.3	*
10	Influenza and pneumonia (J10-J18)	5	1.3	*
12	Viral hepatitis (B15-B19)	2	0.5	*
12	Alzheimer's disease (G30)	2	0.5	*
12	Aortic aneurysm and dissection (I71)	2	0.5	*
12	Peptic ulcer (K25-K28)	2	0.5	*
...	All other causes (Residual)	53	13.9	149.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 65-69 years			
...	All causes	493	100.0	1,747.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	23.5	411.1
2	Malignant neoplasms (C00-C97)	115	23.3	407.5
3	Diabetes mellitus (E10-E14)	44	8.9	155.9
4	Chronic lower respiratory diseases (J40-J47)	34	6.9	120.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	24	4.9	85.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	3.7	*
7	Cerebrovascular diseases (I60-I69)	17	3.4	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	2.8	*
9	Septicemia (A40-A41)	12	2.4	*
10	Aortic aneurysm and dissection (I71)	6	1.2	*
11	Atherosclerosis (I70)	5	1.0	*
11	Influenza and pneumonia (J10-J18)	5	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.8	*
14	Viral hepatitis (B15-B19)	3	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.6	*
...	All other causes (Residual)	70	14.2	248.1
	American Indian, female, 70-74 years			
...	All causes	535	100.0	2,357.5
1	Malignant neoplasms (C00-C97)	133	24.9	586.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	17.9	423.0
3	Diabetes mellitus (E10-E14)	72	13.5	317.3
4	Chronic lower respiratory diseases (J40-J47)	40	7.5	176.3
5	Cerebrovascular diseases (I60-I69)	26	4.9	114.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	3.6	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	14	2.6	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	2.4	*
9	Influenza and pneumonia (J10-J18)	11	2.1	*
10	Alzheimer's disease (G30)	9	1.7	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	1.3	*
12	Septicemia (A40-A41)	6	1.1	*
13	Atherosclerosis (I70)	5	0.9	*
14	Aortic aneurysm and dissection (I71)	4	0.7	*
15	Nutritional deficiencies (E40-E64)	3	0.6	*
15	Parkinson's disease (G20-G21)	3	0.6	*
...	All other causes (Residual)	74	13.8	326.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 75-79 years			
...	All causes	563	100.0	3,047.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	146	25.9	790.2
2	Malignant neoplasms (C00-C97)	115	20.4	622.4
3	Cerebrovascular diseases (I60-I69)	56	9.9	303.1
4	Diabetes mellitus (E10-E14)	44	7.8	238.1
5	Chronic lower respiratory diseases (J40-J47)	27	4.8	146.1
6	Influenza and pneumonia (J10-J18)	19	3.4	*
7	Alzheimer's disease (G30)	15	2.7	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	2.7	*
9	Septicemia (A40-A41)	13	2.3	*
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9	1.6	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.4	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	6	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
13	Parkinson's disease (G20-G21)	4	0.7	*
13	Aortic aneurysm and dissection (I71)	4	0.7	*
...	All other causes (Residual)	78	13.9	422.2
	American Indian, female, 80-84 years			
...	All causes	510	100.0	4,327.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	145	28.4	1,230.3
2	Malignant neoplasms (C00-C97)	84	16.5	712.7
3	Cerebrovascular diseases (I60-I69)	53	10.4	449.7
4	Diabetes mellitus (E10-E14)	39	7.6	330.9
5	Chronic lower respiratory diseases (J40-J47)	22	4.3	186.7
6	Influenza and pneumonia (J10-J18)	19	3.7	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	2.9	*
8	Alzheimer's disease (G30)	11	2.2	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	2.0	*
10	Septicemia (A40-A41)	7	1.4	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	1.0	*
11	Atherosclerosis (I70)	5	1.0	*
11	Pneumonitis due to solids and liquids (J69)	5	1.0	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	5	1.0	*
15	Parkinson's disease (G20-G21)	4	0.8	*
...	All other causes (Residual)	81	15.9	687.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85-89 years				
...	All causes	442	100.0	5,641.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	140	31.7	1,786.9
2	Malignant neoplasms (C00-C97)	59	13.3	753.0
3	Cerebrovascular diseases (I60-I69)	42	9.5	536.1
4	Influenza and pneumonia (J10-J18)	25	5.7	319.1
4	Chronic lower respiratory diseases (J40-J47)	25	5.7	319.1
6	Diabetes mellitus (E10-E14)	21	4.8	268.0
7	Alzheimer's disease (G30)	15	3.4	*
8	Septicemia (A40-A41)	10	2.3	*
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	2.3	*
10	Nutritional deficiencies (E40-E64)	8	1.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.4	*
12	Atherosclerosis (I70)	5	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Pneumonitis due to solids and liquids (J69)	4	0.9	*
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	0.9	*
...	All other causes (Residual)	64	14.5	816.8
American Indian, female, 90-94 years				
...	All causes	287	100.0	6,481.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	104	36.2	2,348.7
2	Cerebrovascular diseases (I60-I69)	35	12.2	790.4
3	Malignant neoplasms (C00-C97)	26	9.1	587.2
4	Influenza and pneumonia (J10-J18)	20	7.0	451.7
5	Alzheimer's disease (G30)	11	3.8	*
6	Diabetes mellitus (E10-E14)	8	2.8	*
7	Chronic lower respiratory diseases (J40-J47)	7	2.4	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	2.4	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	1.4	*
10	Anemias (D50-D64)	3	1.0	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	1.0	*
10	Atherosclerosis (I70)	3	1.0	*
10	Pneumonitis due to solids and liquids (J69)	3	1.0	*
14	Septicemia (A40-A41)	2	0.7	*
14	Nutritional deficiencies (E40-E64)	2	0.7	*
14	Aortic aneurysm and dissection (I71)	2	0.7	*
...	All other causes (Residual)	47	16.4	1,061.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	American Indian, female, 95-99 years			
...	All causes	119	100.0	5,995.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	39.5	2,367.8
2	Cerebrovascular diseases (I60-I69)	13	10.9	*
3	Malignant neoplasms (C00-C97)	7	5.9	*
4	Septicemia (A40-A41)	5	4.2	*
4	Influenza and pneumonia (J10-J18)	5	4.2	*
6	Alzheimer's disease (G30)	4	3.4	*
6	Atherosclerosis (I70)	4	3.4	*
8	Parkinson's disease (G20-G21)	2	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	2	1.7	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
11	Diabetes mellitus (E10-E14)	1	0.8	*
11	Nutritional deficiencies (E40-E64)	1	0.8	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.8	*
11	Aortic aneurysm and dissection (I71)	1	0.8	*
11	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.8	*
11	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.8	*
...	All other causes (Residual)	21	17.6	1,057.9
	American Indian, female, 100 years and over			
...	All causes	33	100.0	5,892.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	39.4	*
2	Malignant neoplasms (C00-C97)	3	9.1	*
2	Influenza and pneumonia (J10-J18)	3	9.1	*
4	Nutritional deficiencies (E40-E64)	2	6.1	*
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	6.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	3.0	*
6	Diabetes mellitus (E10-E14)	1	3.0	*
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	3.0	*
6	Atherosclerosis (I70)	1	3.0	*
6	Chronic lower respiratory diseases (J40-J47)	1	3.0	*
6	Pneumonitis due to solids and liquids (J69)	1	3.0	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	3.0	*
6	Infections of kidney (N10-N12,N13.6,N15.1)	1	3.0	*
...	All other causes (Residual)	2	6.1	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, all ages				
...	All causes	34,875	100.0	309.4
1	Malignant neoplasms (C00-C97)	9,221	26.4	81.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,101	26.1	80.7
3	Cerebrovascular diseases (I60-I69)	3,288	9.4	29.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,678	4.8	14.9
5	Chronic lower respiratory diseases (J40-J47)	1,133	3.2	10.1
6	Influenza and pneumonia (J10-J18)	1,120	3.2	9.9
7	Diabetes mellitus (E10-E14)	1,103	3.2	9.8
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	616	1.8	5.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	527	1.5	4.7
10	Certain conditions originating in the perinatal period (P00-P96)	392	1.1	3.5
11	Assault (homicide) (X85-Y09,Y87.1)	356	1.0	3.2
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	338	1.0	3.0
13	Septicemia (A40-A41)	330	0.9	2.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	299	0.9	2.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	297	0.9	2.6
...	All other causes (Residual)	5,076	14.6	45.0
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	146	100.0	19.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	32.2	6.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	12.3	*
3	Malignant neoplasms (C00-C97)	17	11.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	3.4	*
5	Septicemia (A40-A41)	4	2.7	*
5	Influenza and pneumonia (J10-J18)	4	2.7	*
5	Certain conditions originating in the perinatal period (P00-P96)	4	2.7	*
5	Assault (homicide) (X85-Y09,Y87.1)	4	2.7	*
9	Cerebrovascular diseases (I60-I69)	2	1.4	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	2	1.4	*
11	Viral hepatitis (B15-B19)	1	0.7	*
11	Anemias (D50-D64)	1	0.7	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
...	All other causes (Residual)	36	24.7	4.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	88	100.0	9.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	29.5	2.9
2	Malignant neoplasms (C00-C97)	15	17.0	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	4.5	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	4.5	*
5	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.3	*
5	Influenza and pneumonia (J10-J18)	2	2.3	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	2	2.3	*
8	Septicemia (A40-A41)	1	1.1	*
8	Cerebrovascular diseases (I60-I69)	1	1.1	*
8	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.1	*
8	Assault (homicide) (X85-Y09,Y87.1)	1	1.1	*
...	All other causes (Residual)	27	30.7	3.0
Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	119	100.0	14.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	27.7	3.9
2	Malignant neoplasms (C00-C97)	23	19.3	2.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	7.6	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	7.6	*
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	6	5.0	*
6	Chronic lower respiratory diseases (J40-J47)	5	4.2	*
6	Assault (homicide) (X85-Y09,Y87.1)	5	4.2	*
8	Cerebrovascular diseases (I60-I69)	3	2.5	*
9	Septicemia (A40-A41)	1	0.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
9	Influenza and pneumonia (J10-J18)	1	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.8	*
...	All other causes (Residual)	21	17.6	2.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	322	100.0	38.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	42.9	16.4
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	51	15.8	6.1
3	Assault (homicide) (X85-Y09,Y87.1)	38	11.8	4.5
4	Malignant neoplasms (C00-C97)	27	8.4	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	2.8	*
7	Septicemia (A40-A41)	3	0.9	*
7	Cerebrovascular diseases (I60-I69)	3	0.9	*
9	Influenza and pneumonia (J10-J18)	2	0.6	*
9	Pregnancy, childbirth and the puerperium (O00-099)	2	0.6	*
11	Tuberculosis (A16-A19)	1	0.3	*
11	Meningococcal infection (A39)	1	0.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
11	Anemias (D50-D64)	1	0.3	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
...	All other causes (Residual)	32	9.9	3.8
Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	399	100.0	50.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	158	39.6	20.1
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	59	14.8	7.5
3	Assault (homicide) (X85-Y09,Y87.1)	50	12.5	6.3
4	Malignant neoplasms (C00-C97)	39	9.8	5.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	4.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.5	*
7	Cerebrovascular diseases (I60-I69)	4	1.0	*
8	Influenza and pneumonia (J10-J18)	3	0.8	*
9	Tuberculosis (A16-A19)	2	0.5	*
9	Meningococcal infection (A39)	2	0.5	*
9	Septicemia (A40-A41)	2	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
13	Anemias (D50-D64)	1	0.3	*
13	Aortic aneurysm and dissection (I71)	1	0.3	*
13	Pneumonitis due to solids and liquids (J69)	1	0.3	*
13	Diseases of appendix (K35-K38)	1	0.3	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	44	11.0	5.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	441	100.0	47.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	33.3	15.7
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	91	20.6	9.7
3	Malignant neoplasms (C00-C97)	54	12.2	5.8
4	Assault (homicide) (X85-Y09,Y87.1)	38	8.6	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	5.0	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.6	*
7	Cerebrovascular diseases (I60-I69)	5	1.1	*
8	Diabetes mellitus (E10-E14)	3	0.7	*
8	Aortic aneurysm and dissection (I71)	3	0.7	*
8	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
8	Legal intervention (Y35,Y89.0)	3	0.7	*
12	Septicemia (A40-A41)	2	0.5	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.5	*
12	Influenza and pneumonia (J10-J18)	2	0.5	*
12	Pregnancy, childbirth and the puerperium (O00-099)	2	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	55	12.5	5.9
Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	456	100.0	44.8
1	Malignant neoplasms (C00-C97)	115	25.2	11.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	20.6	9.2
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	63	13.8	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	8.3	3.7
5	Assault (homicide) (X85-Y09,Y87.1)	29	6.4	2.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	12	2.6	*
6	Cerebrovascular diseases (I60-I69)	12	2.6	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.5	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.5	*
10	Diabetes mellitus (E10-E14)	4	0.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.9	*
12	Influenza and pneumonia (J10-J18)	3	0.7	*
12	Pregnancy, childbirth and the puerperium (O00-099)	3	0.7	*
14	Tuberculosis (A16-A19)	2	0.4	*
14	Septicemia (A40-A41)	2	0.4	*
14	Viral hepatitis (B15-B19)	2	0.4	*
14	Aortic aneurysm and dissection (I71)	2	0.4	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.4	*
...	All other causes (Residual)	55	12.1	5.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	683	100.0	68.6
1	Malignant neoplasms (C00-C97)	198	29.0	19.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	16.5	11.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	11.6	7.9
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	65	9.5	6.5
5	Assault (homicide) (X85-Y09,Y87.1)	35	5.1	3.5
6	Cerebrovascular diseases (I60-I69)	32	4.7	3.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.8	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.2	*
10	Diabetes mellitus (E10-E14)	7	1.0	*
11	Viral hepatitis (B15-B19)	6	0.9	*
12	Septicemia (A40-A41)	5	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
13	Influenza and pneumonia (J10-J18)	4	0.6	*
13	Pregnancy, childbirth and the puerperium (O00-099)	4	0.6	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.6	*
...	All other causes (Residual)	96	14.1	9.6
Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	981	100.0	104.5
1	Malignant neoplasms (C00-C97)	345	35.2	36.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	14.6	15.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	9.6	10.0
4	Cerebrovascular diseases (I60-I69)	64	6.5	6.8
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	50	5.1	5.3
6	Assault (homicide) (X85-Y09,Y87.1)	37	3.8	3.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	27	2.8	2.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	18	1.8	*
9	Viral hepatitis (B15-B19)	15	1.5	*
9	Diabetes mellitus (E10-E14)	15	1.5	*
11	Chronic lower respiratory diseases (J40-J47)	13	1.3	*
12	Influenza and pneumonia (J10-J18)	12	1.2	*
13	Septicemia (A40-A41)	9	0.9	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	0.9	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
...	All other causes (Residual)	116	11.8	12.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	1,355	100.0	168.9
1	Malignant neoplasms (C00-C97)	535	39.5	66.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	209	15.4	26.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99	7.3	12.3
4	Cerebrovascular diseases (I60-I69)	98	7.2	12.2
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	53	3.9	6.6
6	Assault (homicide) (X85-Y09,Y87.1)	37	2.7	4.6
7	Viral hepatitis (B15-B19)	33	2.4	4.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	32	2.4	4.0
9	Diabetes mellitus (E10-E14)	31	2.3	3.9
10	Chronic lower respiratory diseases (J40-J47)	23	1.7	2.9
11	Influenza and pneumonia (J10-J18)	17	1.3	*
12	Septicemia (A40-A41)	15	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.8	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	0.7	*
15	Aortic aneurysm and dissection (I71)	6	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
...	All other causes (Residual)	140	10.3	17.4
Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	1,669	100.0	258.6
1	Malignant neoplasms (C00-C97)	654	39.2	101.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	351	21.0	54.4
3	Cerebrovascular diseases (I60-I69)	116	7.0	18.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	5.6	14.4
5	Diabetes mellitus (E10-E14)	52	3.1	8.1
6	Intentional self-harm (suicide) (X60-X84,Y87.0)	43	2.6	6.7
7	Viral hepatitis (B15-B19)	38	2.3	5.9
8	Chronic lower respiratory diseases (J40-J47)	28	1.7	4.3
9	Assault (homicide) (X85-Y09,Y87.1)	24	1.4	3.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	23	1.4	3.6
11	Aortic aneurysm and dissection (I71)	22	1.3	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.1	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	16	1.0	*
14	Septicemia (A40-A41)	15	0.9	*
15	Influenza and pneumonia (J10-J18)	14	0.8	*
...	All other causes (Residual)	162	9.7	25.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	1,937	100.0	431.1
1	Malignant neoplasms (C00-C97)	801	41.4	178.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	436	22.5	97.0
3	Cerebrovascular diseases (I60-I69)	140	7.2	31.2
4	Diabetes mellitus (E10-E14)	72	3.7	16.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	66	3.4	14.7
6	Viral hepatitis (B15-B19)	45	2.3	10.0
7	Chronic lower respiratory diseases (J40-J47)	35	1.8	7.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	32	1.7	7.1
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	29	1.5	6.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.4	6.0
11	Assault (homicide) (X85-Y09,Y87.1)	21	1.1	4.7
12	Influenza and pneumonia (J10-J18)	20	1.0	4.5
13	Septicemia (A40-A41)	17	0.9	*
14	Aortic aneurysm and dissection (I71)	11	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.5	*
...	All other causes (Residual)	175	9.0	39.0
Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	2,389	100.0	678.3
1	Malignant neoplasms (C00-C97)	909	38.0	258.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	603	25.2	171.2
3	Cerebrovascular diseases (I60-I69)	207	8.7	58.8
4	Diabetes mellitus (E10-E14)	86	3.6	24.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	3.1	20.7
6	Viral hepatitis (B15-B19)	41	1.7	11.6
7	Influenza and pneumonia (J10-J18)	39	1.6	11.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.6	10.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	1.4	9.7
10	Chronic lower respiratory diseases (J40-J47)	28	1.2	7.9
11	Aortic aneurysm and dissection (I71)	22	0.9	6.2
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	22	0.9	6.2
13	Septicemia (A40-A41)	19	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.8	*
15	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	0.4	*
15	Assault (homicide) (X85-Y09,Y87.1)	10	0.4	*
...	All other causes (Residual)	230	9.6	65.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	3,029	100.0	1,082.6
1	Malignant neoplasms (C00-C97)	1,122	37.0	401.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	769	25.4	274.8
3	Cerebrovascular diseases (I60-I69)	270	8.9	96.5
4	Diabetes mellitus (E10-E14)	143	4.7	51.1
5	Chronic lower respiratory diseases (J40-J47)	101	3.3	36.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	2.4	25.7
7	Influenza and pneumonia (J10-J18)	67	2.2	23.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	1.7	18.9
9	Viral hepatitis (B15-B19)	39	1.3	13.9
10	Aortic aneurysm and dissection (I71)	29	1.0	10.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	29	1.0	10.4
12	Septicemia (A40-A41)	25	0.8	8.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.6	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	19	0.6	*
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	14	0.5	*
...	All other causes (Residual)	258	8.5	92.2
Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	3,858	100.0	1,710.9
1	Malignant neoplasms (C00-C97)	1,264	32.8	560.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,049	27.2	465.2
3	Cerebrovascular diseases (I60-I69)	358	9.3	158.8
4	Chronic lower respiratory diseases (J40-J47)	153	4.0	67.8
5	Diabetes mellitus (E10-E14)	152	3.9	67.4
6	Influenza and pneumonia (J10-J18)	111	2.9	49.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	2.0	34.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	75	1.9	33.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	40	1.0	17.7
10	Septicemia (A40-A41)	39	1.0	17.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	35	0.9	15.5
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.8	13.3
12	Aortic aneurysm and dissection (I71)	30	0.8	13.3
14	Viral hepatitis (B15-B19)	29	0.8	12.9
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	17	0.4	*
...	All other causes (Residual)	398	10.3	176.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	4,772	100.0	2,916.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,461	30.6	893.0
2	Malignant neoplasms (C00-C97)	1,264	26.5	772.6
3	Cerebrovascular diseases (I60-I69)	533	11.2	325.8
4	Chronic lower respiratory diseases (J40-J47)	220	4.6	134.5
5	Diabetes mellitus (E10-E14)	184	3.9	112.5
6	Influenza and pneumonia (J10-J18)	147	3.1	89.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	2.2	63.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	1.7	50.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	53	1.1	32.4
10	Aortic aneurysm and dissection (I71)	49	1.0	29.9
11	Septicemia (A40-A41)	43	0.9	26.3
12	Alzheimer's disease (G30)	34	0.7	20.8
13	Parkinson's disease (G20-G21)	32	0.7	19.6
14	Pneumonitis due to solids and liquids (J69)	27	0.6	16.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.5	15.3
...	All other causes (Residual)	513	10.8	313.5
Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	4,438	100.0	4,838.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,367	30.8	1,490.4
2	Malignant neoplasms (C00-C97)	936	21.1	1,020.5
3	Cerebrovascular diseases (I60-I69)	543	12.2	592.0
4	Influenza and pneumonia (J10-J18)	223	5.0	243.1
5	Chronic lower respiratory diseases (J40-J47)	199	4.5	217.0
6	Diabetes mellitus (E10-E14)	174	3.9	189.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	85	1.9	92.7
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	74	1.7	80.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	1.6	76.3
10	Septicemia (A40-A41)	48	1.1	52.3
10	Alzheimer's disease (G30)	48	1.1	52.3
12	Pneumonitis due to solids and liquids (J69)	44	1.0	48.0
13	Parkinson's disease (G20-G21)	43	1.0	46.9
13	Aortic aneurysm and dissection (I71)	43	1.0	46.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.6	29.4
...	All other causes (Residual)	514	11.6	560.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	3,671	100.0	7,897.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,224	33.3	2,633.3
2	Malignant neoplasms (C00-C97)	589	16.0	1,267.2
3	Cerebrovascular diseases (I60-I69)	474	12.9	1,019.7
4	Influenza and pneumonia (J10-J18)	212	5.8	456.1
5	Chronic lower respiratory diseases (J40-J47)	179	4.9	385.1
6	Diabetes mellitus (E10-E14)	99	2.7	213.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	2.1	167.8
8	Alzheimer's disease (G30)	74	2.0	159.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	70	1.9	150.6
10	Pneumonitis due to solids and liquids (J69)	49	1.3	105.4
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	48	1.3	103.3
12	Septicemia (A40-A41)	47	1.3	101.1
13	Parkinson's disease (G20-G21)	41	1.1	88.2
14	Aortic aneurysm and dissection (I71)	32	0.9	68.8
15	Atherosclerosis (I70)	23	0.6	49.5
...	All other causes (Residual)	432	11.8	929.4
Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	2,219	100.0	11,178.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	825	37.2	4,156.0
2	Cerebrovascular diseases (I60-I69)	288	13.0	1,450.8
3	Malignant neoplasms (C00-C97)	246	11.1	1,239.2
4	Influenza and pneumonia (J10-J18)	145	6.5	730.4
5	Chronic lower respiratory diseases (J40-J47)	88	4.0	443.3
6	Diabetes mellitus (E10-E14)	59	2.7	297.2
7	Alzheimer's disease (G30)	55	2.5	277.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	49	2.2	246.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	1.9	211.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	32	1.4	161.2
10	Pneumonitis due to solids and liquids (J69)	32	1.4	161.2
12	Atherosclerosis (I70)	21	0.9	105.8
13	Septicemia (A40-A41)	19	0.9	*
14	Aortic aneurysm and dissection (I71)	18	0.8	*
15	Parkinson's disease (G20-G21)	15	0.7	*
...	All other causes (Residual)	285	12.8	1,435.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	898	100.0	13,256.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	371	41.3	5,476.8
2	Cerebrovascular diseases (I60-I69)	115	12.8	1,697.7
3	Influenza and pneumonia (J10-J18)	64	7.1	944.8
4	Malignant neoplasms (C00-C97)	51	5.7	752.9
5	Chronic lower respiratory diseases (J40-J47)	32	3.6	472.4
6	Alzheimer's disease (G30)	25	2.8	369.1
7	Diabetes mellitus (E10-E14)	21	2.3	310.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	2.1	*
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	17	1.9	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.3	*
11	Septicemia (A40-A41)	11	1.2	*
12	Pneumonitis due to solids and liquids (J69)	10	1.1	*
13	Atherosclerosis (I70)	7	0.8	*
14	Nutritional deficiencies (E40-E64)	5	0.6	*
14	Peptic ulcer (K25-K28)	5	0.6	*
...	All other causes (Residual)	133	14.8	1,963.4
Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	202	100.0	14,037.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	41.6	5,837.4
2	Influenza and pneumonia (J10-J18)	20	9.9	1,389.9
3	Cerebrovascular diseases (I60-I69)	17	8.4	*
4	Malignant neoplasms (C00-C97)	12	5.9	*
5	Alzheimer's disease (G30)	10	5.0	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	4.0	*
7	Pneumonitis due to solids and liquids (J69)	6	3.0	*
8	Chronic lower respiratory diseases (J40-J47)	5	2.5	*
9	Atherosclerosis (I70)	4	2.0	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	1.5	*
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	1.0	*
12	Diabetes mellitus (E10-E14)	1	0.5	*
12	Aortic aneurysm and dissection (I71)	1	0.5	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.5	*
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	1	0.5	*
...	All other causes (Residual)	27	13.4	1,876.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	19,018	100.0	349.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,175	27.2	95.0
2	Malignant neoplasms (C00-C97)	4,865	25.6	89.3
3	Cerebrovascular diseases (I60-I69)	1,555	8.2	28.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,057	5.6	19.4
5	Chronic lower respiratory diseases (J40-J47)	722	3.8	13.3
6	Influenza and pneumonia (J10-J18)	592	3.1	10.9
7	Diabetes mellitus (E10-E14)	547	2.9	10.0
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	452	2.4	8.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	254	1.3	4.7
9	Assault (homicide) (X85-Y09,Y87.1)	254	1.3	4.7
11	Certain conditions originating in the perinatal period (P00-P96)	224	1.2	4.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	184	1.0	3.4
13	Aortic aneurysm and dissection (I71)	172	0.9	3.2
14	Viral hepatitis (B15-B19)	161	0.8	3.0
15	Septicemia (A40-A41)	160	0.8	2.9
...	All other causes (Residual)	2,644	13.9	48.5
Asian or Pacific Islander, male, 1-4 years				
...	All causes	79	100.0	21.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	32.9	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	10.1	*
3	Malignant neoplasms (C00-C97)	7	8.9	*
4	Septicemia (A40-A41)	3	3.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.8	*
4	Influenza and pneumonia (J10-J18)	3	3.8	*
4	Certain conditions originating in the perinatal period (P00-P96)	3	3.8	*
4	Assault (homicide) (X85-Y09,Y87.1)	3	3.8	*
9	Viral hepatitis (B15-B19)	1	1.3	*
9	Cerebrovascular diseases (I60-I69)	1	1.3	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.3	*
...	All other causes (Residual)	20	25.3	5.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-9 years				
...	All causes	43	100.0	9.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	27.9	*
2	Malignant neoplasms (C00-C97)	9	20.9	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	7.0	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.7	*
4	Complications of medical and surgical care (Y40-Y84,Y88)	2	4.7	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.3	*
6	Influenza and pneumonia (J10-J18)	1	2.3	*
6	Assault (homicide) (X85-Y09,Y87.1)	1	2.3	*
...	All other causes (Residual)	12	27.9	*
Asian or Pacific Islander, male, 10-14 years				
...	All causes	68	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	29.4	4.6
2	Malignant neoplasms (C00-C97)	13	19.1	*
3	Assault (homicide) (X85-Y09,Y87.1)	5	7.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	5.9	*
4	Chronic lower respiratory diseases (J40-J47)	4	5.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	5.9	*
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	3	4.4	*
8	Cerebrovascular diseases (I60-I69)	2	2.9	*
9	Septicemia (A40-A41)	1	1.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	1.5	*
9	Influenza and pneumonia (J10-J18)	1	1.5	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.5	*
...	All other causes (Residual)	9	13.2	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 15-19 years				
...	All causes	232	100.0	54.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	106	45.7	25.0
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	37	15.9	8.7
3	Assault (homicide) (X85-Y09,Y87.1)	34	14.7	8.0
4	Malignant neoplasms (C00-C97)	16	6.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	4.3	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.2	*
7	Tuberculosis (A16-A19)	1	0.4	*
7	Meningococcal infection (A39)	1	0.4	*
7	Septicemia (A40-A41)	1	0.4	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
7	Cerebrovascular diseases (I60-I69)	1	0.4	*
7	Influenza and pneumonia (J10-J18)	1	0.4	*
...	All other causes (Residual)	18	7.8	*
Asian or Pacific Islander, male, 20-24 years				
...	All causes	284	100.0	73.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	42.6	31.2
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	48	16.9	12.4
3	Assault (homicide) (X85-Y09,Y87.1)	39	13.7	10.0
4	Malignant neoplasms (C00-C97)	22	7.7	5.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	4.9	*
6	Cerebrovascular diseases (I60-I69)	2	0.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.7	*
8	Tuberculosis (A16-A19)	1	0.4	*
8	Meningococcal infection (A39)	1	0.4	*
8	Septicemia (A40-A41)	1	0.4	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
8	Anemias (D50-D64)	1	0.4	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
8	Pneumonitis due to solids and liquids (J69)	1	0.4	*
8	Diseases of appendix (K35-K38)	1	0.4	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	25	8.8	6.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-29 years				
...	All causes	321	100.0	71.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	35.8	25.8
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	68	21.2	15.2
3	Malignant neoplasms (C00-C97)	34	10.6	7.6
4	Assault (homicide) (X85-Y09,Y87.1)	29	9.0	6.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	5.0	*
6	Cerebrovascular diseases (I60-I69)	4	1.2	*
7	Aortic aneurysm and dissection (I71)	3	0.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.9	*
7	Legal intervention (Y35,Y89.0)	3	0.9	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.6	*
10	Diabetes mellitus (E10-E14)	2	0.6	*
10	Influenza and pneumonia (J10-J18)	2	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	2	0.6	*
14	Meningococcal infection (A39)	1	0.3	*
14	Septicemia (A40-A41)	1	0.3	*
14	Viral hepatitis (B15-B19)	1	0.3	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	32	10.0	7.2
Asian or Pacific Islander, male, 30-34 years				
...	All causes	269	100.0	54.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	23.4	12.8
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	46	17.1	9.4
3	Malignant neoplasms (C00-C97)	45	16.7	9.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	10.0	5.5
5	Assault (homicide) (X85-Y09,Y87.1)	19	7.1	*
6	Human immunodeficiency virus (HIV) disease (B20-B24)	11	4.1	*
7	Cerebrovascular diseases (I60-I69)	8	3.0	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	6	2.2	*
9	Chronic lower respiratory diseases (J40-J47)	4	1.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.1	*
11	Tuberculosis (A16-A19)	2	0.7	*
11	Viral hepatitis (B15-B19)	2	0.7	*
11	Diabetes mellitus (E10-E14)	2	0.7	*
11	Influenza and pneumonia (J10-J18)	2	0.7	*
15	Septicemia (A40-A41)	1	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
15	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (Residual)	26	9.7	5.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 35-39 years				
...	All causes	424	100.0	88.0
1	Malignant neoplasms (C00-C97)	96	22.6	19.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76	17.9	15.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	14.2	12.5
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	49	11.6	10.2
5	Cerebrovascular diseases (I60-I69)	17	4.0	*
6	Assault (homicide) (X85-Y09,Y87.1)	15	3.5	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	10	2.4	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	9	2.1	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.7	*
10	Diabetes mellitus (E10-E14)	6	1.4	*
11	Viral hepatitis (B15-B19)	4	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Septicemia (A40-A41)	2	0.5	*
13	Aortic aneurysm and dissection (I71)	2	0.5	*
13	Influenza and pneumonia (J10-J18)	2	0.5	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
...	All other causes (Residual)	63	14.9	13.1
Asian or Pacific Islander, male, 40-44 years				
...	All causes	569	100.0	127.0
1	Malignant neoplasms (C00-C97)	151	26.5	33.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	114	20.0	25.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	10.9	13.8
4	Intentional self-harm (suicide) (X60-X84,Y87.0)	36	6.3	8.0
5	Cerebrovascular diseases (I60-I69)	30	5.3	6.7
5	Assault (homicide) (X85-Y09,Y87.1)	30	5.3	6.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	22	3.9	4.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	15	2.6	*
9	Viral hepatitis (B15-B19)	11	1.9	*
10	Diabetes mellitus (E10-E14)	9	1.6	*
11	Septicemia (A40-A41)	6	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.7	*
12	Aortic aneurysm and dissection (I71)	4	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
...	All other causes (Residual)	67	11.8	15.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	Asian or Pacific Islander, male, 45-49 years			
...	All causes	783	100.0	207.9
1	Malignant neoplasms (C00-C97)	243	31.0	64.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	161	20.6	42.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	8.0	16.7
4	Cerebrovascular diseases (I60-I69)	48	6.1	12.7
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	39	5.0	10.4
6	Assault (homicide) (X85-Y09,Y87.1)	28	3.6	7.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	24	3.1	6.4
8	Viral hepatitis (B15-B19)	23	2.9	6.1
9	Diabetes mellitus (E10-E14)	18	2.3	*
10	Chronic lower respiratory diseases (J40-J47)	13	1.7	*
11	Influenza and pneumonia (J10-J18)	10	1.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.1	*
13	Septicemia (A40-A41)	7	0.9	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	7	0.9	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.5	*
15	Aortic aneurysm and dissection (I71)	4	0.5	*
...	All other causes (Residual)	82	10.5	21.8
	Asian or Pacific Islander, male, 50-54 years			
...	All causes	978	100.0	327.0
1	Malignant neoplasms (C00-C97)	311	31.8	104.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	270	27.6	90.3
3	Cerebrovascular diseases (I60-I69)	58	5.9	19.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	5.4	17.7
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	33	3.4	11.0
6	Diabetes mellitus (E10-E14)	31	3.2	10.4
7	Viral hepatitis (B15-B19)	22	2.2	7.4
8	Chronic lower respiratory diseases (J40-J47)	21	2.1	7.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.6	*
9	Assault (homicide) (X85-Y09,Y87.1)	16	1.6	*
11	Aortic aneurysm and dissection (I71)	12	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.2	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	11	1.1	*
14	Influenza and pneumonia (J10-J18)	9	0.9	*
15	Septicemia (A40-A41)	8	0.8	*
...	All other causes (Residual)	95	9.7	31.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 55-59 years				
...	All causes	1,131	100.0	536.2
1	Malignant neoplasms (C00-C97)	417	36.9	197.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	307	27.1	145.5
3	Cerebrovascular diseases (I60-I69)	76	6.7	36.0
4	Diabetes mellitus (E10-E14)	47	4.2	22.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	3.4	18.5
6	Viral hepatitis (B15-B19)	28	2.5	13.3
7	Chronic lower respiratory diseases (J40-J47)	22	1.9	10.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	18	1.6	*
8	Intentional self-harm (suicide) (X60-X84,Y87.0)	18	1.6	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.5	*
11	Assault (homicide) (X85-Y09,Y87.1)	16	1.4	*
12	Influenza and pneumonia (J10-J18)	14	1.2	*
13	Aortic aneurysm and dissection (I71)	10	0.9	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.8	*
15	Septicemia (A40-A41)	6	0.5	*
...	All other causes (Residual)	87	7.7	41.2
Asian or Pacific Islander, male, 60-64 years				
...	All causes	1,434	100.0	875.2
1	Malignant neoplasms (C00-C97)	498	34.7	303.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	422	29.4	257.5
3	Cerebrovascular diseases (I60-I69)	121	8.4	73.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	3.2	28.1
5	Diabetes mellitus (E10-E14)	43	3.0	26.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	27	1.9	16.5
7	Viral hepatitis (B15-B19)	24	1.7	14.6
8	Influenza and pneumonia (J10-J18)	21	1.5	12.8
9	Aortic aneurysm and dissection (I71)	19	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.3	*
11	Intentional self-harm (suicide) (X60-X84,Y87.0)	17	1.2	*
12	Chronic lower respiratory diseases (J40-J47)	13	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.8	*
14	Septicemia (A40-A41)	9	0.6	*
15	Assault (homicide) (X85-Y09,Y87.1)	7	0.5	*
...	All other causes (Residual)	138	9.6	84.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-69 years				
...	All causes	1,692	100.0	1,401.1
1	Malignant neoplasms (C00-C97)	589	34.8	487.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	496	29.3	410.7
3	Cerebrovascular diseases (I60-I69)	128	7.6	106.0
4	Diabetes mellitus (E10-E14)	81	4.8	67.1
5	Chronic lower respiratory diseases (J40-J47)	70	4.1	58.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	2.7	37.3
7	Influenza and pneumonia (J10-J18)	38	2.2	31.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.2	17.4
9	Viral hepatitis (B15-B19)	19	1.1	*
10	Aortic aneurysm and dissection (I71)	16	0.9	*
11	Septicemia (A40-A41)	15	0.9	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.5	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	9	0.5	*
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	8	0.5	*
...	All other causes (Residual)	133	7.9	110.1
Asian or Pacific Islander, male, 70-74 years				
...	All causes	2,153	100.0	2,319.7
1	Malignant neoplasms (C00-C97)	706	32.8	760.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	621	28.8	669.1
3	Cerebrovascular diseases (I60-I69)	186	8.6	200.4
4	Chronic lower respiratory diseases (J40-J47)	95	4.4	102.4
5	Diabetes mellitus (E10-E14)	72	3.3	77.6
6	Influenza and pneumonia (J10-J18)	61	2.8	65.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	1.8	40.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	1.4	32.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	25	1.2	26.9
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.0	23.7
11	Aortic aneurysm and dissection (I71)	19	0.9	*
12	Septicemia (A40-A41)	18	0.8	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	17	0.8	*
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	12	0.6	*
15	Viral hepatitis (B15-B19)	11	0.5	*
15	Pneumonitis due to solids and liquids (J69)	11	0.5	*
...	All other causes (Residual)	209	9.7	225.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 75-79 years				
...	All causes	2,627	100.0	3,826.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	820	31.2	1,194.4
2	Malignant neoplasms (C00-C97)	718	27.3	1,045.8
3	Cerebrovascular diseases (I60-I69)	263	10.0	383.1
4	Chronic lower respiratory diseases (J40-J47)	157	6.0	228.7
5	Diabetes mellitus (E10-E14)	84	3.2	122.4
6	Influenza and pneumonia (J10-J18)	78	3.0	113.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	2.1	78.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	1.7	64.1
9	Aortic aneurysm and dissection (I71)	32	1.2	46.6
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	1.0	37.9
11	Parkinson's disease (G20-G21)	24	0.9	35.0
12	Septicemia (A40-A41)	23	0.9	33.5
13	Alzheimer's disease (G30)	19	0.7	*
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	16	0.6	*
15	Pneumonitis due to solids and liquids (J69)	15	0.6	*
...	All other causes (Residual)	254	9.7	370.0
Asian or Pacific Islander, male, 80-84 years				
...	All causes	2,207	100.0	5,729.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	693	31.4	1,799.2
2	Malignant neoplasms (C00-C97)	495	22.4	1,285.1
3	Cerebrovascular diseases (I60-I69)	227	10.3	589.4
4	Chronic lower respiratory diseases (J40-J47)	127	5.8	329.7
5	Influenza and pneumonia (J10-J18)	125	5.7	324.5
6	Diabetes mellitus (E10-E14)	72	3.3	186.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	46	2.1	119.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.6	90.9
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	30	1.4	77.9
10	Parkinson's disease (G20-G21)	29	1.3	75.3
11	Septicemia (A40-A41)	24	1.1	62.3
12	Pneumonitis due to solids and liquids (J69)	23	1.0	59.7
13	Aortic aneurysm and dissection (I71)	22	1.0	57.1
14	Alzheimer's disease (G30)	20	0.9	51.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.7	*
...	All other causes (Residual)	224	10.1	581.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85-89 years				
...	All causes	1,842	100.0	9,717.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	578	31.4	3,049.2
2	Malignant neoplasms (C00-C97)	335	18.2	1,767.3
3	Cerebrovascular diseases (I60-I69)	218	11.8	1,150.0
4	Chronic lower respiratory diseases (J40-J47)	119	6.5	627.8
5	Influenza and pneumonia (J10-J18)	111	6.0	585.6
6	Diabetes mellitus (E10-E14)	41	2.2	216.3
7	Alzheimer's disease (G30)	36	2.0	189.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.9	184.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	1.8	174.1
10	Parkinson's disease (G20-G21)	27	1.5	142.4
11	Pneumonitis due to solids and liquids (J69)	25	1.4	131.9
12	Septicemia (A40-A41)	22	1.2	116.1
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	22	1.2	116.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.9	*
15	Aortic aneurysm and dissection (I71)	13	0.7	*
...	All other causes (Residual)	211	11.5	1,113.1
Asian or Pacific Islander, male, 90-94 years				
...	All causes	1,026	100.0	12,627.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	364	35.5	4,480.0
2	Malignant neoplasms (C00-C97)	133	13.0	1,636.9
3	Cerebrovascular diseases (I60-I69)	112	10.9	1,378.5
4	Influenza and pneumonia (J10-J18)	74	7.2	910.8
5	Chronic lower respiratory diseases (J40-J47)	46	4.5	566.2
6	Diabetes mellitus (E10-E14)	28	2.7	344.6
7	Alzheimer's disease (G30)	24	2.3	295.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.0	258.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20	1.9	246.2
10	Pneumonitis due to solids and liquids (J69)	19	1.9	*
11	Aortic aneurysm and dissection (I71)	14	1.4	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	1.0	*
12	Atherosclerosis (I70)	10	1.0	*
14	Septicemia (A40-A41)	9	0.9	*
14	Parkinson's disease (G20-G21)	9	0.9	*
...	All other causes (Residual)	133	13.0	1,636.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
	Asian or Pacific Islander, male, 95-99 years			
...	All causes	346	100.0	13,616.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	45.1	6,139.3
2	Cerebrovascular diseases (I60-I69)	49	14.2	1,928.4
3	Malignant neoplasms (C00-C97)	22	6.4	865.8
3	Influenza and pneumonia (J10-J18)	22	6.4	865.8
5	Chronic lower respiratory diseases (J40-J47)	13	3.8	*
6	Diabetes mellitus (E10-E14)	10	2.9	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7	2.0	*
8	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	1.2	*
8	Pneumonitis due to solids and liquids (J69)	4	1.2	*
8	Peptic ulcer (K25-K28)	4	1.2	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.2	*
12	Septicemia (A40-A41)	3	0.9	*
12	Atherosclerosis (I70)	3	0.9	*
14	Alzheimer's disease (G30)	2	0.6	*
14	Aortic aneurysm and dissection (I71)	2	0.6	*
14	Hernia (K40-K46)	2	0.6	*
14	Intentional self-harm (suicide) (X60-X84,Y87.0)	2	0.6	*
...	All other causes (Residual)	37	10.7	1,456.1
	Asian or Pacific Islander, male, 100 years and over			
...	All causes	58	100.0	14,043.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	39.7	5,569.0
2	Influenza and pneumonia (J10-J18)	7	12.1	*
3	Alzheimer's disease (G30)	3	5.2	*
3	Cerebrovascular diseases (I60-I69)	3	5.2	*
3	Chronic lower respiratory diseases (J40-J47)	3	5.2	*
6	Malignant neoplasms (C00-C97)	2	3.4	*
6	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	2	3.4	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	3.4	*
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	3.4	*
10	Diabetes mellitus (E10-E14)	1	1.7	*
10	Atherosclerosis (I70)	1	1.7	*
10	Pneumonitis due to solids and liquids (J69)	1	1.7	*
10	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	1.7	*
10	Intentional self-harm (suicide) (X60-X84,Y87.0)	1	1.7	*
...	All other causes (Residual)	6	10.3	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, all ages				
...	All causes	15,857	100.0	272.2
1	Malignant neoplasms (C00-C97)	4,356	27.5	74.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,926	24.8	67.4
3	Cerebrovascular diseases (I60-I69)	1,733	10.9	29.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	621	3.9	10.7
5	Diabetes mellitus (E10-E14)	556	3.5	9.5
6	Influenza and pneumonia (J10-J18)	528	3.3	9.1
7	Chronic lower respiratory diseases (J40-J47)	411	2.6	7.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	273	1.7	4.7
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	179	1.1	3.1
10	Septicemia (A40-A41)	170	1.1	2.9
11	Certain conditions originating in the perinatal period (P00-P96)	168	1.1	2.9
12	Intentional self-harm (suicide) (X60-X84,Y87.0)	164	1.0	2.8
13	Alzheimer's disease (G30)	163	1.0	2.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	148	0.9	2.5
15	Viral hepatitis (B15-B19)	124	0.8	2.1
...	All other causes (Residual)	2,337	14.7	40.1
Asian or Pacific Islander, female, 1-4 years				
...	All causes	67	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	31.3	5.8
2	Malignant neoplasms (C00-C97)	10	14.9	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	14.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.0	*
5	Septicemia (A40-A41)	1	1.5	*
5	Anemias (D50-D64)	1	1.5	*
5	Cerebrovascular diseases (I60-I69)	1	1.5	*
5	Influenza and pneumonia (J10-J18)	1	1.5	*
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.5	*
5	Certain conditions originating in the perinatal period (P00-P96)	1	1.5	*
5	Assault (homicide) (X85-Y09,Y87.1)	1	1.5	*
5	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.5	*
...	All other causes (Residual)	16	23.9	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 5-9 years				
...	All causes	45	100.0	10.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	31.1	*
2	Malignant neoplasms (C00-C97)	6	13.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.4	*
4	Septicemia (A40-A41)	1	2.2	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.2	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.2	*
4	Cerebrovascular diseases (I60-I69)	1	2.2	*
4	Influenza and pneumonia (J10-J18)	1	2.2	*
4	Acute bronchitis and bronchiolitis (J20-J21)	1	2.2	*
4	Chronic lower respiratory diseases (J40-J47)	1	2.2	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	2.2	*
...	All other causes (Residual)	15	33.3	*
Asian or Pacific Islander, female, 10-14 years				
...	All causes	51	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	25.5	*
2	Malignant neoplasms (C00-C97)	10	19.6	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	9.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	9.8	*
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	3	5.9	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.0	*
6	Cerebrovascular diseases (I60-I69)	1	2.0	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.0	*
...	All other causes (Residual)	12	23.5	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-19 years				
...	All causes	90	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	35.6	7.7
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	14	15.6	*
3	Malignant neoplasms (C00-C97)	11	12.2	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	4.4	*
4	Assault (homicide) (X85-Y09,Y87.1)	4	4.4	*
6	Septicemia (A40-A41)	2	2.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.2	*
6	Cerebrovascular diseases (I60-I69)	2	2.2	*
6	Pregnancy, childbirth and the puerperium (O00-099)	2	2.2	*
10	Anemias (D50-D64)	1	1.1	*
10	Influenza and pneumonia (J10-J18)	1	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	1	1.1	*
...	All other causes (Residual)	14	15.6	*
Asian or Pacific Islander, female, 20-24 years				
...	All causes	115	100.0	28.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	32.2	9.3
2	Malignant neoplasms (C00-C97)	17	14.8	*
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	11	9.6	*
3	Assault (homicide) (X85-Y09,Y87.1)	11	9.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	4.3	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.5	*
7	Influenza and pneumonia (J10-J18)	3	2.6	*
8	Cerebrovascular diseases (I60-I69)	2	1.7	*
9	Tuberculosis (A16-A19)	1	0.9	*
9	Meningococcal infection (A39)	1	0.9	*
9	Septicemia (A40-A41)	1	0.9	*
9	Aortic aneurysm and dissection (I71)	1	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.9	*
...	All other causes (Residual)	19	16.5	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 25-29 years				
...	All causes	120	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	26.7	6.6
2	Intentional self-harm (suicide) (X60-X84,Y87.0)	23	19.2	4.7
3	Malignant neoplasms (C00-C97)	20	16.7	4.1
4	Assault (homicide) (X85-Y09,Y87.1)	9	7.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	5.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.3	*
7	Pregnancy, childbirth and the puerperium (O00-099)	2	1.7	*
8	Septicemia (A40-A41)	1	0.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
8	Diabetes mellitus (E10-E14)	1	0.8	*
8	Cerebrovascular diseases (I60-I69)	1	0.8	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.8	*
...	All other causes (Residual)	18	15.0	*
Asian or Pacific Islander, female, 30-34 years				
...	All causes	187	100.0	35.5
1	Malignant neoplasms (C00-C97)	70	37.4	13.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	16.6	5.9
3	Intentional self-harm (suicide) (X60-X84,Y87.0)	17	9.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	5.9	*
5	Assault (homicide) (X85-Y09,Y87.1)	10	5.3	*
6	Cerebrovascular diseases (I60-I69)	4	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.6	*
7	Pregnancy, childbirth and the puerperium (O00-099)	3	1.6	*
9	Diabetes mellitus (E10-E14)	2	1.1	*
9	Aortic aneurysm and dissection (I71)	2	1.1	*
11	Septicemia (A40-A41)	1	0.5	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
11	Nutritional deficiencies (E40-E64)	1	0.5	*
11	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.5	*
11	Influenza and pneumonia (J10-J18)	1	0.5	*
11	Acute bronchitis and bronchiolitis (J20-J21)	1	0.5	*
11	Peptic ulcer (K25-K28)	1	0.5	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
...	All other causes (Residual)	24	12.8	4.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-39 years				
...	All causes	259	100.0	50.4
1	Malignant neoplasms (C00-C97)	102	39.4	19.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	14.3	7.2
3	Assault (homicide) (X85-Y09,Y87.1)	20	7.7	3.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	7.3	*
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	16	6.2	*
6	Cerebrovascular diseases (I60-I69)	15	5.8	*
7	Pregnancy, childbirth and the puerperium (O00-099)	4	1.5	*
8	Septicemia (A40-A41)	3	1.2	*
9	Viral hepatitis (B15-B19)	2	0.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.8	*
9	Anemias (D50-D64)	2	0.8	*
9	Influenza and pneumonia (J10-J18)	2	0.8	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.8	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.8	*
...	All other causes (Residual)	29	11.2	5.6
Asian or Pacific Islander, female, 40-44 years				
...	All causes	412	100.0	83.9
1	Malignant neoplasms (C00-C97)	194	47.1	39.5
2	Cerebrovascular diseases (I60-I69)	34	8.3	6.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	7.8	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	7.0	5.9
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	14	3.4	*
6	Influenza and pneumonia (J10-J18)	9	2.2	*
6	Chronic lower respiratory diseases (J40-J47)	9	2.2	*
8	Assault (homicide) (X85-Y09,Y87.1)	7	1.7	*
9	Diabetes mellitus (E10-E14)	6	1.5	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.5	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	5	1.2	*
12	Viral hepatitis (B15-B19)	4	1.0	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.0	*
14	Septicemia (A40-A41)	3	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.7	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.7	*
...	All other causes (Residual)	50	12.1	10.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 45-49 years				
...	All causes	572	100.0	134.3
1	Malignant neoplasms (C00-C97)	292	51.0	68.6
2	Cerebrovascular diseases (I60-I69)	50	8.7	11.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	8.4	11.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	6.3	8.5
5	Intentional self-harm (suicide) (X60-X84,Y87.0)	14	2.4	*
6	Diabetes mellitus (E10-E14)	13	2.3	*
7	Viral hepatitis (B15-B19)	10	1.7	*
7	Chronic lower respiratory diseases (J40-J47)	10	1.7	*
9	Assault (homicide) (X85-Y09,Y87.1)	9	1.6	*
10	Septicemia (A40-A41)	8	1.4	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	8	1.4	*
12	Influenza and pneumonia (J10-J18)	7	1.2	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	3	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
15	Aortic aneurysm and dissection (I71)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	55	9.6	12.9
Asian or Pacific Islander, female, 50-54 years				
...	All causes	691	100.0	199.5
1	Malignant neoplasms (C00-C97)	343	49.6	99.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	11.7	23.4
3	Cerebrovascular diseases (I60-I69)	58	8.4	16.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	5.8	11.5
5	Diabetes mellitus (E10-E14)	21	3.0	6.1
6	Viral hepatitis (B15-B19)	16	2.3	*
7	Aortic aneurysm and dissection (I71)	10	1.4	*
7	Intentional self-harm (suicide) (X60-X84,Y87.0)	10	1.4	*
9	Assault (homicide) (X85-Y09,Y87.1)	8	1.2	*
10	Septicemia (A40-A41)	7	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	7	1.0	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	7	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.9	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	0.7	*
14	Influenza and pneumonia (J10-J18)	5	0.7	*
...	All other causes (Residual)	67	9.7	19.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-59 years				
...	All causes	806	100.0	338.1
1	Malignant neoplasms (C00-C97)	384	47.6	161.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	16.0	54.1
3	Cerebrovascular diseases (I60-I69)	64	7.9	26.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	3.3	11.3
5	Diabetes mellitus (E10-E14)	25	3.1	10.5
6	Viral hepatitis (B15-B19)	17	2.1	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.6	*
9	Septicemia (A40-A41)	11	1.4	*
9	Intentional self-harm (suicide) (X60-X84,Y87.0)	11	1.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.2	*
12	Influenza and pneumonia (J10-J18)	6	0.7	*
13	Assault (homicide) (X85-Y09,Y87.1)	5	0.6	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
14	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	4	0.5	*
...	All other causes (Residual)	82	10.2	34.4
Asian or Pacific Islander, female, 60-64 years				
...	All causes	955	100.0	507.0
1	Malignant neoplasms (C00-C97)	411	43.0	218.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	181	19.0	96.1
3	Cerebrovascular diseases (I60-I69)	86	9.0	45.7
4	Diabetes mellitus (E10-E14)	43	4.5	22.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	2.8	14.3
6	Influenza and pneumonia (J10-J18)	18	1.9	*
7	Viral hepatitis (B15-B19)	17	1.8	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	15	1.6	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	11	1.2	*
11	Septicemia (A40-A41)	10	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
13	Alzheimer's disease (G30)	5	0.5	*
13	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	5	0.5	*
13	Intentional self-harm (suicide) (X60-X84,Y87.0)	5	0.5	*
...	All other causes (Residual)	98	10.3	52.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 65-69 years				
...	All causes	1,337	100.0	840.7
1	Malignant neoplasms (C00-C97)	533	39.9	335.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	273	20.4	171.7
3	Cerebrovascular diseases (I60-I69)	142	10.6	89.3
4	Diabetes mellitus (E10-E14)	62	4.6	39.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	2.4	20.1
6	Chronic lower respiratory diseases (J40-J47)	31	2.3	19.5
7	Influenza and pneumonia (J10-J18)	29	2.2	18.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	2.0	17.0
9	Viral hepatitis (B15-B19)	20	1.5	12.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.0	*
11	Aortic aneurysm and dissection (I71)	13	1.0	*
12	Septicemia (A40-A41)	10	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.7	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	10	0.7	*
15	Intentional self-harm (suicide) (X60-X84,Y87.0)	6	0.4	*
...	All other causes (Residual)	125	9.3	78.6
Asian or Pacific Islander, female, 70-74 years				
...	All causes	1,705	100.0	1,285.0
1	Malignant neoplasms (C00-C97)	558	32.7	420.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	428	25.1	322.6
3	Cerebrovascular diseases (I60-I69)	172	10.1	129.6
4	Diabetes mellitus (E10-E14)	80	4.7	60.3
5	Chronic lower respiratory diseases (J40-J47)	58	3.4	43.7
6	Influenza and pneumonia (J10-J18)	50	2.9	37.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	2.6	33.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	2.3	30.1
9	Septicemia (A40-A41)	21	1.2	15.8
10	Viral hepatitis (B15-B19)	18	1.1	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18	1.1	*
12	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	15	0.9	*
13	Aortic aneurysm and dissection (I71)	11	0.6	*
14	Alzheimer's disease (G30)	9	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.5	*
...	All other causes (Residual)	174	10.2	131.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-79 years				
...	All causes	2,145	100.0	2,258.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	641	29.9	675.0
2	Malignant neoplasms (C00-C97)	546	25.5	575.0
3	Cerebrovascular diseases (I60-I69)	270	12.6	284.3
4	Diabetes mellitus (E10-E14)	100	4.7	105.3
5	Influenza and pneumonia (J10-J18)	69	3.2	72.7
6	Chronic lower respiratory diseases (J40-J47)	63	2.9	66.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	2.3	52.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.8	41.1
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	27	1.3	28.4
10	Septicemia (A40-A41)	20	0.9	21.1
11	Aortic aneurysm and dissection (I71)	17	0.8	*
12	Alzheimer's disease (G30)	15	0.7	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	13	0.6	*
14	Pneumonitis due to solids and liquids (J69)	12	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.5	*
...	All other causes (Residual)	252	11.7	265.4
Asian or Pacific Islander, female, 80-84 years				
...	All causes	2,231	100.0	4,193.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	674	30.2	1,266.8
2	Malignant neoplasms (C00-C97)	441	19.8	828.9
3	Cerebrovascular diseases (I60-I69)	316	14.2	593.9
4	Diabetes mellitus (E10-E14)	102	4.6	191.7
5	Influenza and pneumonia (J10-J18)	98	4.4	184.2
6	Chronic lower respiratory diseases (J40-J47)	72	3.2	135.3
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	44	2.0	82.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	1.7	73.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.6	65.8
10	Alzheimer's disease (G30)	28	1.3	52.6
11	Septicemia (A40-A41)	24	1.1	45.1
12	Aortic aneurysm and dissection (I71)	21	0.9	39.5
12	Pneumonitis due to solids and liquids (J69)	21	0.9	39.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	16	0.7	*
15	Parkinson's disease (G20-G21)	14	0.6	*
...	All other causes (Residual)	286	12.8	537.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 85-89 years				
...	All causes	1,829	100.0	6,644.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	646	35.3	2,346.9
2	Cerebrovascular diseases (I60-I69)	256	14.0	930.0
3	Malignant neoplasms (C00-C97)	254	13.9	922.8
4	Influenza and pneumonia (J10-J18)	101	5.5	366.9
5	Chronic lower respiratory diseases (J40-J47)	60	3.3	218.0
6	Diabetes mellitus (E10-E14)	58	3.2	210.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	2.4	156.2
8	Alzheimer's disease (G30)	38	2.1	138.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	2.0	134.4
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	26	1.4	94.5
11	Septicemia (A40-A41)	25	1.4	90.8
12	Pneumonitis due to solids and liquids (J69)	24	1.3	87.2
13	Aortic aneurysm and dissection (I71)	19	1.0	*
14	Parkinson's disease (G20-G21)	14	0.8	*
14	Atherosclerosis (I70)	14	0.8	*
...	All other causes (Residual)	214	11.7	777.4
Asian or Pacific Islander, female, 90-94 years				
...	All causes	1,193	100.0	10,174.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	461	38.6	3,931.4
2	Cerebrovascular diseases (I60-I69)	176	14.8	1,500.9
3	Malignant neoplasms (C00-C97)	113	9.5	963.7
4	Influenza and pneumonia (J10-J18)	71	6.0	605.5
5	Chronic lower respiratory diseases (J40-J47)	42	3.5	358.2
6	Diabetes mellitus (E10-E14)	31	2.6	264.4
6	Alzheimer's disease (G30)	31	2.6	264.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	2.4	247.3
9	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	22	1.8	187.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.8	179.1
11	Pneumonitis due to solids and liquids (J69)	13	1.1	*
12	Atherosclerosis (I70)	11	0.9	*
13	Septicemia (A40-A41)	10	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.8	*
15	Parkinson's disease (G20-G21)	6	0.5	*
...	All other causes (Residual)	146	12.2	1,245.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and sex: United States, 2000

[Rates per 100,000 population in specified group. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys. Population denominators for 2000 death rates are based on the 1990 census population.]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, 1992), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 95-99 years				
...	All causes	552	100.0	13,040.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	215	38.9	5,079.1
2	Cerebrovascular diseases (I60-I69)	66	12.0	1,559.2
3	Influenza and pneumonia (J10-J18)	42	7.6	992.2
4	Malignant neoplasms (C00-C97)	29	5.3	685.1
5	Alzheimer's disease (G30)	23	4.2	543.3
6	Chronic lower respiratory diseases (J40-J47)	19	3.4	*
7	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	13	2.4	*
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	2.2	*
9	Diabetes mellitus (E10-E14)	11	2.0	*
10	Septicemia (A40-A41)	8	1.4	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.4	*
12	Pneumonitis due to solids and liquids (J69)	6	1.1	*
13	Nutritional deficiencies (E40-E64)	5	0.9	*
14	Atherosclerosis (I70)	4	0.7	*
15	Parkinson's disease (G20-G21)	2	0.4	*
15	Aortic aneurysm and dissection (I71)	2	0.4	*
...	All other causes (Residual)	87	15.8	2,055.3
Asian or Pacific Islander, female, 100 years and over				
...	All causes	144	100.0	14,035.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	42.4	5,945.4
2	Cerebrovascular diseases (I60-I69)	14	9.7	*
3	Influenza and pneumonia (J10-J18)	13	9.0	*
4	Malignant neoplasms (C00-C97)	10	6.9	*
5	Alzheimer's disease (G30)	7	4.9	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	4.2	*
7	Pneumonitis due to solids and liquids (J69)	5	3.5	*
8	Atherosclerosis (I70)	3	2.1	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.4	*
10	Essential (primary) hypertension and hypertensive renal disease (I10,I12)	1	0.7	*
10	Aortic aneurysm and dissection (I71)	1	0.7	*
...	All other causes (Residual)	21	14.6	2,046.8
5	Chronic lower respiratory diseases (J40-J47)	4	2.9	*
8	Pneumonitis due to solids and liquids (J69)	3	2.2	*
9	Diabetes mellitus (E10-E14)	2	1.5	*
9	Alzheimer's disease (G30)	2	1.5	*
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	1.5	*
12	Peptic ulcer (K25-K28)	1	0.7	*
...	All other causes (Residual)	15	10.9	*

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable