

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, all ages				
...	All causes	2,543,279	100.0	810.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	599,711	23.6	191.0
2	Malignant neoplasms (C00-C97)	582,623	22.9	185.6
3	Chronic lower respiratory diseases (J40-J47)	143,489	5.6	45.7
4	Cerebrovascular diseases (I60-I69)	128,546	5.1	40.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127,792	5.0	40.7
6	Alzheimer's disease (G30)	83,637	3.3	26.6
7	Diabetes mellitus (E10-E14)	73,932	2.9	23.6
8	Influenza and pneumonia (J09-J18)	50,636	2.0	16.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45,622	1.8	14.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40,600	1.6	12.9
11	Septicemia (A40-A41)	35,842	1.4	11.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	34,979	1.4	11.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29,115	1.1	9.3
14	Parkinson's disease (G20-G21)	23,818	0.9	7.6
15	Pneumonitis due to solids and liquids (J69)	17,897	0.7	5.7
...	All other causes (Residual)	525,040	20.6	167.3
All races, both sexes, 1-4 years				
...	All causes	4,218	100.0	26.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,353	32.1	8.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	501	11.9	3.1
3	Malignant neoplasms (C00-C97)	392	9.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	8.0	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	154	3.7	1.0
6	Influenza and pneumonia (J09-J18)	93	2.2	0.6
7	Septicemia (A40-A41)	62	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	56	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	1.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.2	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	42	1.0	0.3
12	Acute bronchitis and bronchiolitis (J20-J21)	17	0.4	*
13	Anemias (D50-D64)	15	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.3	*
...	All other causes (Residual)	1,065	25.2	6.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 5-14 years				
...	All causes	5,200	100.0	12.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,550	29.8	3.8
2	Malignant neoplasms (C00-C97)	912	17.5	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	327	6.3	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	311	6.0	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	311	6.0	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	175	3.4	0.4
7	Chronic lower respiratory diseases (J40-J47)	119	2.3	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	1.7	0.2
9	Cerebrovascular diseases (I60-I69)	85	1.6	0.2
9	Influenza and pneumonia (J09-J18)	85	1.6	0.2
11	Septicemia (A40-A41)	52	1.0	0.1
12	Diabetes mellitus (E10-E14)	34	0.7	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	34	0.7	0.1
14	Anemias (D50-D64)	28	0.5	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	22	0.4	0.1
...	All other causes (Residual)	1,068	20.5	2.6
All races, both sexes, 15-24 years				
...	All causes	29,182	100.0	66.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,908	40.8	27.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,872	16.7	11.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,614	15.8	10.5
4	Malignant neoplasms (C00-C97)	1,574	5.4	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	956	3.3	2.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	423	1.4	1.0
7	Diabetes mellitus (E10-E14)	196	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	183	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	169	0.6	0.4
10	Influenza and pneumonia (J09-J18)	147	0.5	0.3
11	Chronic lower respiratory diseases (J40-J47)	143	0.5	0.3
12	Septicemia (A40-A41)	120	0.4	0.3
13	Legal intervention (Y35,Y89.0)	118	0.4	0.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	116	0.4	0.3
15	Anemias (D50-D64)	89	0.3	0.2
...	All other causes (Residual)	3,554	12.2	8.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 25-34 years				
...	All causes	44,591	100.0	105.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,851	35.5	37.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,216	13.9	14.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,342	9.7	10.3
4	Malignant neoplasms (C00-C97)	3,674	8.2	8.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,231	7.2	7.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	652	1.5	1.5
7	Diabetes mellitus (E10-E14)	646	1.4	1.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	597	1.3	1.4
9	Cerebrovascular diseases (I60-I69)	535	1.2	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	401	0.9	0.9
11	Pregnancy, childbirth and the puerperium (O00-O99)	390	0.9	0.9
12	Influenza and pneumonia (J09-J18)	356	0.8	0.8
13	Septicemia (A40-A41)	325	0.7	0.8
14	Chronic lower respiratory diseases (J40-J47)	289	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	232	0.5	0.5
...	All other causes (Residual)	6,854	15.4	16.2
All races, both sexes, 35-44 years				
...	All causes	69,162	100.0	170.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,034	21.7	37.1
2	Malignant neoplasms (C00-C97)	11,337	16.4	28.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,489	15.2	25.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,758	9.8	16.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,705	3.9	6.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,469	3.6	6.1
7	Diabetes mellitus (E10-E14)	1,867	2.7	4.6
8	Cerebrovascular diseases (I60-I69)	1,730	2.5	4.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,345	1.9	3.3
10	Septicemia (A40-A41)	757	1.1	1.9
11	Chronic lower respiratory diseases (J40-J47)	743	1.1	1.8
12	Influenza and pneumonia (J09-J18)	685	1.0	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	649	0.9	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	396	0.6	1.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	336	0.5	0.8
...	All other causes (Residual)	11,862	17.2	29.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 45-54 years				
...	All causes	179,463	100.0	405.4
1	Malignant neoplasms (C00-C97)	48,028	26.8	108.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,265	19.7	79.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,394	11.4	46.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	8,877	4.9	20.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,862	4.9	20.0
6	Diabetes mellitus (E10-E14)	5,747	3.2	13.0
7	Cerebrovascular diseases (I60-I69)	5,654	3.2	12.8
8	Chronic lower respiratory diseases (J40-J47)	4,533	2.5	10.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2,582	1.4	5.8
10	Septicemia (A40-A41)	2,340	1.3	5.3
11	Viral hepatitis (B15-B19)	2,067	1.2	4.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,064	1.2	4.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,055	1.1	4.6
14	Influenza and pneumonia (J09-J18)	1,825	1.0	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,335	0.7	3.0
...	All other causes (Residual)	27,835	15.5	62.9
All races, both sexes, 55-64 years				
...	All causes	329,606	100.0	854.2
1	Malignant neoplasms (C00-C97)	113,130	34.3	293.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,228	21.6	184.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,822	4.8	41.0
4	Chronic lower respiratory diseases (J40-J47)	15,212	4.6	39.4
5	Diabetes mellitus (E10-E14)	12,553	3.8	32.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	11,230	3.4	29.1
7	Cerebrovascular diseases (I60-I69)	11,070	3.4	28.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,929	2.1	18.0
9	Septicemia (A40-A41)	4,982	1.5	12.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,765	1.4	12.3
11	Influenza and pneumonia (J09-J18)	3,929	1.2	10.2
12	Viral hepatitis (B15-B19)	3,877	1.2	10.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,006	0.9	7.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,793	0.5	4.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,503	0.5	3.9
...	All other causes (Residual)	48,577	14.7	125.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 65-74 years				
...	All causes	432,346	100.0	1,802.5
1	Malignant neoplasms (C00-C97)	151,639	35.1	632.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	93,143	21.5	388.3
3	Chronic lower respiratory diseases (J40-J47)	33,575	7.8	140.0
4	Cerebrovascular diseases (I60-I69)	18,168	4.2	75.7
5	Diabetes mellitus (E10-E14)	16,713	3.9	69.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,558	2.4	44.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,999	1.9	33.3
8	Septicemia (A40-A41)	7,013	1.6	29.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	6,616	1.5	27.6
10	Influenza and pneumonia (J09-J18)	6,269	1.4	26.1
11	Alzheimer's disease (G30)	4,300	1.0	17.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,864	0.9	16.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,367	0.8	14.0
14	Parkinson's disease (G20-G21)	2,943	0.7	12.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,658	0.6	11.1
...	All other causes (Residual)	63,521	14.7	264.8
All races, both sexes, 75-84 years				
...	All causes	620,428	100.0	4,674.5
1	Malignant neoplasms (C00-C97)	154,192	24.9	1,161.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	146,491	23.6	1,103.7
3	Chronic lower respiratory diseases (J40-J47)	48,307	7.8	364.0
4	Cerebrovascular diseases (I60-I69)	36,134	5.8	272.2
5	Alzheimer's disease (G30)	23,281	3.8	175.4
6	Diabetes mellitus (E10-E14)	19,346	3.1	145.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,303	2.3	107.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13,257	2.1	99.9
9	Influenza and pneumonia (J09-J18)	13,040	2.1	98.2
10	Parkinson's disease (G20-G21)	10,117	1.6	76.2
11	Septicemia (A40-A41)	9,803	1.6	73.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,868	1.1	51.7
13	Pneumonitis due to solids and liquids (J69)	4,856	0.8	36.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,619	0.7	34.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,894	0.6	29.3
...	All other causes (Residual)	111,920	18.0	843.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, both sexes, 85 years and over				
...	All causes	805,307	100.0	13,678.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	238,206	29.6	4,046.1
2	Malignant neoplasms (C00-C97)	97,666	12.1	1,658.9
3	Alzheimer's disease (G30)	55,109	6.8	936.1
4	Cerebrovascular diseases (I60-I69)	54,825	6.8	931.2
5	Chronic lower respiratory diseases (J40-J47)	40,493	5.0	687.8
6	Influenza and pneumonia (J09-J18)	24,046	3.0	408.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,837	2.5	336.9
8	Diabetes mellitus (E10-E14)	16,822	2.1	285.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,484	2.0	280.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,580	1.7	230.7
11	Septicemia (A40-A41)	10,206	1.3	173.4
12	Parkinson's disease (G20-G21)	10,143	1.3	172.3
13	Pneumonitis due to solids and liquids (J69)	9,000	1.1	152.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,969	0.6	84.4
15	Atherosclerosis (I70)	3,940	0.5	66.9
...	All other causes (Residual)	189,981	23.6	3,226.9
All races, male, all ages				
...	All causes	1,273,722	100.0	824.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	312,491	24.5	202.3
2	Malignant neoplasms (C00-C97)	305,670	24.0	197.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80,010	6.3	51.8
4	Chronic lower respiratory diseases (J40-J47)	67,673	5.3	43.8
5	Cerebrovascular diseases (I60-I69)	52,638	4.1	34.1
6	Diabetes mellitus (E10-E14)	38,584	3.0	25.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31,780	2.5	20.6
8	Alzheimer's disease (G30)	25,653	2.0	16.6
9	Influenza and pneumonia (J09-J18)	24,013	1.9	15.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,731	1.8	14.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	22,623	1.8	14.6
12	Septicemia (A40-A41)	16,789	1.3	10.9
13	Parkinson's disease (G20-G21)	14,121	1.1	9.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13,208	1.0	8.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,859	0.9	7.7
...	All other causes (Residual)	233,879	18.4	151.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 1-4 years				
...	All causes	2,395	100.0	29.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	806	33.7	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	275	11.5	3.4
3	Malignant neoplasms (C00-C97)	218	9.1	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	197	8.2	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.3	1.0
6	Influenza and pneumonia (J09-J18)	55	2.3	0.7
7	Cerebrovascular diseases (I60-I69)	36	1.5	0.4
8	Chronic lower respiratory diseases (J40-J47)	35	1.5	0.4
9	Septicemia (A40-A41)	29	1.2	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	27	1.1	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.8	0.2
12	Acute bronchitis and bronchiolitis (J20-J21)	13	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
15	Anemias (D50-D64)	5	0.2	*
15	Pneumonitis due to solids and liquids (J69)	5	0.2	*
...	All other causes (Residual)	580	24.2	7.1
All races, male, 5-14 years				
...	All causes	3,036	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	982	32.3	4.7
2	Malignant neoplasms (C00-C97)	513	16.9	2.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	226	7.4	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	181	6.0	0.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	170	5.6	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	3.2	0.5
7	Chronic lower respiratory diseases (J40-J47)	62	2.0	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	41	1.4	0.2
10	Influenza and pneumonia (J09-J18)	40	1.3	0.2
11	Septicemia (A40-A41)	32	1.1	0.2
12	Diabetes mellitus (E10-E14)	18	0.6	*
13	Anemias (D50-D64)	16	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	11	0.4	*
...	All other causes (Residual)	588	19.4	2.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 15-24 years				
...	All causes	21,456	100.0	95.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,849	41.2	39.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,995	18.6	17.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,918	18.3	17.4
4	Malignant neoplasms (C00-C97)	924	4.3	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	644	3.0	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	250	1.2	1.1
7	Legal intervention (Y35,Y89.0)	112	0.5	0.5
8	Cerebrovascular diseases (I60-I69)	108	0.5	0.5
9	Diabetes mellitus (E10-E14)	106	0.5	0.5
10	Chronic lower respiratory diseases (J40-J47)	98	0.5	0.4
11	Influenza and pneumonia (J09-J18)	90	0.4	0.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	78	0.4	0.3
13	Septicemia (A40-A41)	54	0.3	0.2
14	Anemias (D50-D64)	50	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.2	0.2
...	All other causes (Residual)	2,131	9.9	9.5
All races, male, 25-34 years				
...	All causes	30,783	100.0	144.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,675	37.9	54.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,985	16.2	23.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,651	11.9	17.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,214	7.2	10.4
5	Malignant neoplasms (C00-C97)	1,802	5.9	8.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	456	1.5	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	379	1.2	1.8
8	Diabetes mellitus (E10-E14)	359	1.2	1.7
9	Cerebrovascular diseases (I60-I69)	297	1.0	1.4
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	0.8	1.1
11	Influenza and pneumonia (J09-J18)	205	0.7	1.0
12	Septicemia (A40-A41)	181	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	170	0.6	0.8
14	Legal intervention (Y35,Y89.0)	141	0.5	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	122	0.4	0.6
...	All other causes (Residual)	3,902	12.7	18.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 35-44 years				
...	All causes	42,929	100.0	212.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,473	24.4	51.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,231	16.8	35.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,185	12.1	25.7
4	Malignant neoplasms (C00-C97)	4,849	11.3	24.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,138	5.0	10.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,601	3.7	7.9
7	Diabetes mellitus (E10-E14)	1,156	2.7	5.7
8	Cerebrovascular diseases (I60-I69)	962	2.2	4.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	853	2.0	4.2
10	Septicemia (A40-A41)	386	0.9	1.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	376	0.9	1.9
12	Influenza and pneumonia (J09-J18)	358	0.8	1.8
13	Chronic lower respiratory diseases (J40-J47)	341	0.8	1.7
14	Aortic aneurysm and dissection (I71)	225	0.5	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	222	0.5	1.1
...	All other causes (Residual)	6,573	15.3	32.6
All races, male, 45-54 years				
...	All causes	109,214	100.0	500.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,927	22.8	114.3
2	Malignant neoplasms (C00-C97)	24,003	22.0	110.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,591	12.4	62.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,578	6.0	30.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,979	5.5	27.4
6	Diabetes mellitus (E10-E14)	3,604	3.3	16.5
7	Cerebrovascular diseases (I60-I69)	3,101	2.8	14.2
8	Chronic lower respiratory diseases (J40-J47)	2,133	2.0	9.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,894	1.7	8.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,530	1.4	7.0
11	Viral hepatitis (B15-B19)	1,472	1.3	6.8
12	Septicemia (A40-A41)	1,250	1.1	5.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,200	1.1	5.5
14	Influenza and pneumonia (J09-J18)	1,020	0.9	4.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	868	0.8	4.0
...	All other causes (Residual)	16,064	14.7	73.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 55-64 years				
...	All causes	200,951	100.0	1,080.2
1	Malignant neoplasms (C00-C97)	62,670	31.2	336.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,557	24.7	266.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,804	5.4	58.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	7,939	4.0	42.7
5	Chronic lower respiratory diseases (J40-J47)	7,881	3.9	42.4
6	Diabetes mellitus (E10-E14)	7,634	3.8	41.0
7	Cerebrovascular diseases (I60-I69)	6,410	3.2	34.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,332	2.7	28.7
9	Viral hepatitis (B15-B19)	2,754	1.4	14.8
10	Septicemia (A40-A41)	2,703	1.3	14.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,700	1.3	14.5
12	Influenza and pneumonia (J09-J18)	2,301	1.1	12.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,870	0.9	10.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,371	0.7	7.4
15	Aortic aneurysm and dissection (I71)	892	0.4	4.8
...	All other causes (Residual)	28,133	14.0	151.2
All races, male, 65-74 years				
...	All causes	244,935	100.0	2,186.4
1	Malignant neoplasms (C00-C97)	83,653	34.2	746.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	59,113	24.1	527.7
3	Chronic lower respiratory diseases (J40-J47)	16,901	6.9	150.9
4	Cerebrovascular diseases (I60-I69)	9,600	3.9	85.7
5	Diabetes mellitus (E10-E14)	9,489	3.9	84.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,705	2.7	59.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,313	1.8	38.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	4,108	1.7	36.7
9	Septicemia (A40-A41)	3,562	1.5	31.8
10	Influenza and pneumonia (J09-J18)	3,507	1.4	31.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,700	1.1	24.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,069	0.8	18.5
13	Parkinson's disease (G20-G21)	1,954	0.8	17.4
14	Alzheimer's disease (G30)	1,854	0.8	16.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,576	0.6	14.1
...	All other causes (Residual)	33,831	13.8	302.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, male, 75-84 years				
...	All causes	310,681	100.0	5,500.6
1	Malignant neoplasms (C00-C97)	81,761	26.3	1,447.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,401	25.2	1,388.1
3	Chronic lower respiratory diseases (J40-J47)	23,365	7.5	413.7
4	Cerebrovascular diseases (I60-I69)	15,686	5.0	277.7
5	Diabetes mellitus (E10-E14)	9,706	3.1	171.8
6	Alzheimer's disease (G30)	8,705	2.8	154.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,573	2.4	134.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,831	2.2	120.9
9	Influenza and pneumonia (J09-J18)	6,681	2.2	118.3
10	Parkinson's disease (G20-G21)	6,430	2.1	113.8
11	Septicemia (A40-A41)	4,565	1.5	80.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,879	0.9	51.0
13	Pneumonitis due to solids and liquids (J69)	2,791	0.9	49.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,546	0.8	45.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,028	0.7	35.9
...	All other causes (Residual)	50,733	16.3	898.2
All races, male, 85 years and over				
...	All causes	294,103	100.0	14,974.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	90,005	30.6	4,582.7
2	Malignant neoplasms (C00-C97)	45,234	15.4	2,303.1
3	Chronic lower respiratory diseases (J40-J47)	16,672	5.7	848.9
4	Cerebrovascular diseases (I60-I69)	16,342	5.6	832.1
5	Alzheimer's disease (G30)	14,676	5.0	747.2
6	Influenza and pneumonia (J09-J18)	9,659	3.3	491.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,866	2.7	400.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,098	2.4	361.4
9	Diabetes mellitus (E10-E14)	6,507	2.2	331.3
10	Parkinson's disease (G20-G21)	5,355	1.8	272.7
11	Pneumonitis due to solids and liquids (J69)	4,441	1.5	226.1
12	Septicemia (A40-A41)	3,921	1.3	199.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,895	1.3	198.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,210	0.8	112.5
15	Atherosclerosis (I70)	1,274	0.4	64.9
...	All other causes (Residual)	58,948	20.0	3,001.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, all ages				
...	All causes	1,269,557	100.0	796.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	287,220	22.6	180.2
2	Malignant neoplasms (C00-C97)	276,953	21.8	173.7
3	Cerebrovascular diseases (I60-I69)	75,908	6.0	47.6
4	Chronic lower respiratory diseases (J40-J47)	75,816	6.0	47.6
5	Alzheimer's disease (G30)	57,984	4.6	36.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,782	3.8	30.0
7	Diabetes mellitus (E10-E14)	35,348	2.8	22.2
8	Influenza and pneumonia (J09-J18)	26,623	2.1	16.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22,891	1.8	14.4
10	Septicemia (A40-A41)	19,053	1.5	12.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,256	1.4	10.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	12,356	1.0	7.8
13	Parkinson's disease (G20-G21)	9,697	0.8	6.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,820	0.7	5.5
15	Pneumonitis due to solids and liquids (J69)	8,184	0.6	5.1
...	All other causes (Residual)	287,666	22.7	180.4
All races, female, 1-4 years				
...	All causes	1,823	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	547	30.0	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	226	12.4	2.9
3	Malignant neoplasms (C00-C97)	174	9.5	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	7.8	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	4.1	0.9
6	Influenza and pneumonia (J09-J18)	38	2.1	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.9	0.4
8	Septicemia (A40-A41)	33	1.8	0.4
9	Cerebrovascular diseases (I60-I69)	20	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	16	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	15	0.8	*
12	Anemias (D50-D64)	10	0.5	*
13	Meningitis (G00,G03)	6	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
15	Acute bronchitis and bronchiolitis (J20-J21)	4	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.2	*
...	All other causes (Residual)	474	26.0	6.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 5-14 years				
...	All causes	2,164	100.0	10.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	568	26.2	2.8
2	Malignant neoplasms (C00-C97)	399	18.4	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	7.3	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	6.0	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	3.9	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	3.7	0.4
7	Chronic lower respiratory diseases (J40-J47)	57	2.6	0.3
8	Influenza and pneumonia (J09-J18)	45	2.1	0.2
9	Cerebrovascular diseases (I60-I69)	44	2.0	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	1.8	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	23	1.1	0.1
12	Septicemia (A40-A41)	20	0.9	0.1
13	Diabetes mellitus (E10-E14)	16	0.7	*
14	Anemias (D50-D64)	12	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.5	*
...	All other causes (Residual)	480	22.2	2.4
All races, female, 15-24 years				
...	All causes	7,726	100.0	36.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,059	39.6	14.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	954	12.3	4.5
3	Malignant neoplasms (C00-C97)	650	8.4	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	619	8.0	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	312	4.0	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	173	2.2	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	169	2.2	0.8
8	Diabetes mellitus (E10-E14)	90	1.2	0.4
9	Cerebrovascular diseases (I60-I69)	75	1.0	0.3
10	Septicemia (A40-A41)	66	0.9	0.3
11	Influenza and pneumonia (J09-J18)	57	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	45	0.6	0.2
13	Anemias (D50-D64)	39	0.5	0.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	38	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.5	0.2
...	All other causes (Residual)	1,342	17.4	6.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 25-34 years				
...	All causes	13,808	100.0	65.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,176	30.2	19.9
2	Malignant neoplasms (C00-C97)	1,872	13.6	8.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,231	8.9	5.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,017	7.4	4.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	691	5.0	3.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	390	2.8	1.9
7	Diabetes mellitus (E10-E14)	287	2.1	1.4
8	Cerebrovascular diseases (I60-I69)	238	1.7	1.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	218	1.6	1.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	196	1.4	0.9
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	157	1.1	0.7
12	Influenza and pneumonia (J09-J18)	151	1.1	0.7
13	Septicemia (A40-A41)	144	1.0	0.7
14	Chronic lower respiratory diseases (J40-J47)	119	0.9	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	0.8	0.5
...	All other causes (Residual)	2,811	20.4	13.4
All races, female, 35-44 years				
...	All causes	26,233	100.0	129.0
1	Malignant neoplasms (C00-C97)	6,488	24.7	31.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,561	17.4	22.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,258	12.4	16.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,573	6.0	7.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	868	3.3	4.3
6	Cerebrovascular diseases (I60-I69)	768	2.9	3.8
7	Diabetes mellitus (E10-E14)	711	2.7	3.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	567	2.2	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	492	1.9	2.4
10	Chronic lower respiratory diseases (J40-J47)	402	1.5	2.0
11	Septicemia (A40-A41)	371	1.4	1.8
12	Influenza and pneumonia (J09-J18)	327	1.2	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	273	1.0	1.3
13	Pregnancy, childbirth and the puerperium (O00-O99)	273	1.0	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	174	0.7	0.9
...	All other causes (Residual)	5,127	19.5	25.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 45-54 years				
...	All causes	70,249	100.0	312.7
1	Malignant neoplasms (C00-C97)	24,025	34.2	107.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,338	14.7	46.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,803	9.7	30.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,898	4.1	12.9
5	Cerebrovascular diseases (I60-I69)	2,553	3.6	11.4
6	Chronic lower respiratory diseases (J40-J47)	2,400	3.4	10.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,284	3.3	10.2
8	Diabetes mellitus (E10-E14)	2,143	3.1	9.5
9	Septicemia (A40-A41)	1,090	1.6	4.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	864	1.2	3.8
11	Influenza and pneumonia (J09-J18)	805	1.1	3.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	688	1.0	3.1
13	Viral hepatitis (B15-B19)	595	0.8	2.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	525	0.7	2.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	467	0.7	2.1
...	All other causes (Residual)	11,771	16.8	52.4
All races, female, 55-64 years				
...	All causes	128,655	100.0	643.8
1	Malignant neoplasms (C00-C97)	50,460	39.2	252.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,671	16.8	108.4
3	Chronic lower respiratory diseases (J40-J47)	7,331	5.7	36.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,018	3.9	25.1
5	Diabetes mellitus (E10-E14)	4,919	3.8	24.6
6	Cerebrovascular diseases (I60-I69)	4,660	3.6	23.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,291	2.6	16.5
8	Septicemia (A40-A41)	2,279	1.8	11.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,065	1.6	10.3
10	Influenza and pneumonia (J09-J18)	1,628	1.3	8.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,597	1.2	8.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,136	0.9	5.7
13	Viral hepatitis (B15-B19)	1,123	0.9	5.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	654	0.5	3.3
15	Alzheimer's disease (G30)	467	0.4	2.3
...	All other causes (Residual)	20,356	15.8	101.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 65-74 years				
...	All causes	187,411	100.0	1,466.1
1	Malignant neoplasms (C00-C97)	67,986	36.3	531.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,030	18.2	266.2
3	Chronic lower respiratory diseases (J40-J47)	16,674	8.9	130.4
4	Cerebrovascular diseases (I60-I69)	8,568	4.6	67.0
5	Diabetes mellitus (E10-E14)	7,224	3.9	56.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,853	2.1	30.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,686	2.0	28.8
8	Septicemia (A40-A41)	3,451	1.8	27.0
9	Influenza and pneumonia (J09-J18)	2,762	1.5	21.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,508	1.3	19.6
11	Alzheimer's disease (G30)	2,446	1.3	19.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,795	1.0	14.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,082	0.6	8.5
14	Parkinson's disease (G20-G21)	989	0.5	7.7
15	Pneumonitis due to solids and liquids (J69)	778	0.4	6.1
...	All other causes (Residual)	29,579	15.8	231.4
All races, female, 75-84 years				
...	All causes	309,747	100.0	4,062.5
1	Malignant neoplasms (C00-C97)	72,431	23.4	950.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,090	22.0	893.0
3	Chronic lower respiratory diseases (J40-J47)	24,942	8.1	327.1
4	Cerebrovascular diseases (I60-I69)	20,448	6.6	268.2
5	Alzheimer's disease (G30)	14,576	4.7	191.2
6	Diabetes mellitus (E10-E14)	9,640	3.1	126.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,730	2.2	88.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,426	2.1	84.3
9	Influenza and pneumonia (J09-J18)	6,359	2.1	83.4
10	Septicemia (A40-A41)	5,238	1.7	68.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,989	1.3	52.3
12	Parkinson's disease (G20-G21)	3,687	1.2	48.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,073	0.7	27.2
14	Pneumonitis due to solids and liquids (J69)	2,065	0.7	27.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,866	0.6	24.5
...	All other causes (Residual)	61,187	19.8	802.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
All races, female, 85 years and over				
...	All causes	511,204	100.0	13,030.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	148,201	29.0	3,777.5
2	Malignant neoplasms (C00-C97)	52,432	10.3	1,336.4
3	Alzheimer's disease (G30)	40,433	7.9	1,030.6
4	Cerebrovascular diseases (I60-I69)	38,483	7.5	980.9
5	Chronic lower respiratory diseases (J40-J47)	23,821	4.7	607.2
6	Influenza and pneumonia (J09-J18)	14,387	2.8	366.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,971	2.3	305.1
8	Diabetes mellitus (E10-E14)	10,315	2.0	262.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,685	1.9	246.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,386	1.8	239.2
11	Septicemia (A40-A41)	6,285	1.2	160.2
12	Parkinson's disease (G20-G21)	4,788	0.9	122.0
13	Pneumonitis due to solids and liquids (J69)	4,559	0.9	116.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,759	0.5	70.3
15	Atherosclerosis (I70)	2,666	0.5	68.0
...	All other causes (Residual)	131,033	25.6	3,339.9
White, both sexes, all ages				
...	All causes	2,175,178	100.0	876.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	514,372	23.6	207.2
2	Malignant neoplasms (C00-C97)	496,885	22.8	200.2
3	Chronic lower respiratory diseases (J40-J47)	131,782	6.1	53.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110,789	5.1	44.6
5	Cerebrovascular diseases (I60-I69)	107,964	5.0	43.5
6	Alzheimer's disease (G30)	76,590	3.5	30.9
7	Diabetes mellitus (E10-E14)	57,806	2.7	23.3
8	Influenza and pneumonia (J09-J18)	43,585	2.0	17.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,606	1.7	14.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36,066	1.7	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	30,747	1.4	12.4
12	Septicemia (A40-A41)	28,945	1.3	11.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22,908	1.1	9.2
14	Parkinson's disease (G20-G21)	22,352	1.0	9.0
15	Pneumonitis due to solids and liquids (J69)	15,938	0.7	6.4
...	All other causes (Residual)	441,843	20.3	178.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 1-4 years				
...	All causes	2,958	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,004	33.9	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	366	12.4	3.0
3	Malignant neoplasms (C00-C97)	301	10.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	189	6.4	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	3.4	0.8
6	Influenza and pneumonia (J09-J18)	67	2.3	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.5	0.4
8	Septicemia (A40-A41)	37	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	35	1.2	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	24	0.8	0.2
11	Chronic lower respiratory diseases (J40-J47)	20	0.7	0.2
12	Acute bronchitis and bronchiolitis (J20-J21)	11	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
14	Pneumonitis due to solids and liquids (J69)	7	0.2	*
15	Anemias (D50-D64)	6	0.2	*
15	Diabetes mellitus (E10-E14)	6	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.2	*
...	All other causes (Residual)	726	24.5	6.0
White, both sexes, 5-14 years				
...	All causes	3,711	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,087	29.3	3.5
2	Malignant neoplasms (C00-C97)	712	19.2	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	248	6.7	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	229	6.2	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	5.3	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	3.1	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	2.0	0.2
8	Cerebrovascular diseases (I60-I69)	67	1.8	0.2
9	Influenza and pneumonia (J09-J18)	63	1.7	0.2
10	Septicemia (A40-A41)	43	1.2	0.1
11	Chronic lower respiratory diseases (J40-J47)	41	1.1	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	24	0.6	0.1
13	Diabetes mellitus (E10-E14)	17	0.5	*
14	Meningitis (G00,G03)	14	0.4	*
15	Anemias (D50-D64)	11	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.3	*
...	All other causes (Residual)	758	20.4	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 15-24 years				
...	All causes	21,101	100.0	63.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,001	47.4	30.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,001	19.0	12.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,579	7.5	4.7
4	Malignant neoplasms (C00-C97)	1,218	5.8	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	613	2.9	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	321	1.5	1.0
7	Cerebrovascular diseases (I60-I69)	133	0.6	0.4
8	Diabetes mellitus (E10-E14)	131	0.6	0.4
9	Influenza and pneumonia (J09-J18)	110	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	98	0.5	0.3
11	Septicemia (A40-A41)	81	0.4	0.2
12	Chronic lower respiratory diseases (J40-J47)	75	0.4	0.2
13	Legal intervention (Y35,Y89.0)	65	0.3	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.2	0.1
...	All other causes (Residual)	2,572	12.2	7.7
White, both sexes, 25-34 years				
...	All causes	33,370	100.0	102.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,486	40.4	41.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,302	15.9	16.3
3	Malignant neoplasms (C00-C97)	2,791	8.4	8.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,169	6.5	6.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,744	5.2	5.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	484	1.5	1.5
7	Diabetes mellitus (E10-E14)	416	1.2	1.3
8	Cerebrovascular diseases (I60-I69)	374	1.1	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	327	1.0	1.0
10	Influenza and pneumonia (J09-J18)	259	0.8	0.8
11	Pregnancy, childbirth and the puerperium (O00-O99)	229	0.7	0.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	218	0.7	0.7
13	Septicemia (A40-A41)	216	0.6	0.7
14	Chronic lower respiratory diseases (J40-J47)	168	0.5	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	131	0.4	0.4
...	All other causes (Residual)	5,056	15.2	15.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 35-44 years				
...	All causes	52,228	100.0	166.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,770	24.5	40.6
2	Malignant neoplasms (C00-C97)	8,676	16.6	27.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,248	13.9	23.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,979	11.4	19.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,065	4.0	6.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,316	2.5	4.2
7	Diabetes mellitus (E10-E14)	1,276	2.4	4.1
8	Cerebrovascular diseases (I60-I69)	1,140	2.2	3.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	550	1.1	1.7
10	Chronic lower respiratory diseases (J40-J47)	526	1.0	1.7
11	Septicemia (A40-A41)	513	1.0	1.6
12	Influenza and pneumonia (J09-J18)	504	1.0	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	364	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	320	0.6	1.0
15	Viral hepatitis (B15-B19)	235	0.4	0.7
...	All other causes (Residual)	8,746	16.7	27.8
White, both sexes, 45-54 years				
...	All causes	140,139	100.0	394.2
1	Malignant neoplasms (C00-C97)	37,815	27.0	106.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,435	18.9	74.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,362	12.4	48.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,232	5.9	23.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	7,765	5.5	21.8
6	Diabetes mellitus (E10-E14)	4,064	2.9	11.4
7	Chronic lower respiratory diseases (J40-J47)	3,772	2.7	10.6
8	Cerebrovascular diseases (I60-I69)	3,672	2.6	10.3
9	Viral hepatitis (B15-B19)	1,749	1.2	4.9
10	Septicemia (A40-A41)	1,679	1.2	4.7
11	Influenza and pneumonia (J09-J18)	1,369	1.0	3.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,243	0.9	3.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,228	0.9	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,218	0.9	3.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	756	0.5	2.1
...	All other causes (Residual)	21,780	15.5	61.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 55-64 years				
...	All causes	263,657	100.0	826.8
1	Malignant neoplasms (C00-C97)	92,018	34.9	288.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,349	21.0	173.6
3	Chronic lower respiratory diseases (J40-J47)	13,390	5.1	42.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,245	5.0	41.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	9,783	3.7	30.7
6	Diabetes mellitus (E10-E14)	9,308	3.5	29.2
7	Cerebrovascular diseases (I60-I69)	7,803	3.0	24.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,528	2.5	20.5
9	Septicemia (A40-A41)	3,714	1.4	11.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,145	1.2	9.9
11	Viral hepatitis (B15-B19)	3,103	1.2	9.7
12	Influenza and pneumonia (J09-J18)	3,072	1.2	9.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,989	0.8	6.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,264	0.5	4.0
15	Pneumonitis due to solids and liquids (J69)	982	0.4	3.1
...	All other causes (Residual)	38,964	14.8	122.2
White, both sexes, 65-74 years				
...	All causes	364,706	100.0	1,781.2
1	Malignant neoplasms (C00-C97)	129,683	35.6	633.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,835	21.1	375.3
3	Chronic lower respiratory diseases (J40-J47)	30,730	8.4	150.1
4	Cerebrovascular diseases (I60-I69)	14,283	3.9	69.8
5	Diabetes mellitus (E10-E14)	12,859	3.5	62.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,211	2.5	45.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,932	1.6	29.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	5,895	1.6	28.8
9	Septicemia (A40-A41)	5,547	1.5	27.1
10	Influenza and pneumonia (J09-J18)	5,155	1.4	25.2
11	Alzheimer's disease (G30)	3,809	1.0	18.6
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,173	0.9	15.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,791	0.8	13.6
14	Parkinson's disease (G20-G21)	2,697	0.7	13.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,322	0.6	11.3
...	All other causes (Residual)	53,784	14.7	262.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, both sexes, 75-84 years				
...	All causes	545,479	100.0	4,721.9
1	Malignant neoplasms (C00-C97)	135,578	24.9	1,173.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	127,703	23.4	1,105.4
3	Chronic lower respiratory diseases (J40-J47)	45,058	8.3	390.0
4	Cerebrovascular diseases (I60-I69)	30,932	5.7	267.8
5	Alzheimer's disease (G30)	21,160	3.9	183.2
6	Diabetes mellitus (E10-E14)	15,512	2.8	134.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,068	2.4	113.1
8	Influenza and pneumonia (J09-J18)	11,315	2.1	97.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10,821	2.0	93.7
10	Parkinson's disease (G20-G21)	9,497	1.7	82.2
11	Septicemia (A40-A41)	8,175	1.5	70.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,448	1.0	47.2
13	Pneumonitis due to solids and liquids (J69)	4,306	0.8	37.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,191	0.8	36.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,564	0.7	30.9
...	All other causes (Residual)	99,151	18.2	858.3
White, both sexes, 85 years and over				
...	All causes	732,474	100.0	13,951.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	217,566	29.7	4,143.8
2	Malignant neoplasms (C00-C97)	88,033	12.0	1,676.7
3	Alzheimer's disease (G30)	50,764	6.9	966.9
4	Cerebrovascular diseases (I60-I69)	49,462	6.8	942.1
5	Chronic lower respiratory diseases (J40-J47)	37,986	5.2	723.5
6	Influenza and pneumonia (J09-J18)	21,578	2.9	411.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,837	2.6	358.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,338	2.0	273.1
9	Diabetes mellitus (E10-E14)	14,216	1.9	270.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,694	1.6	222.7
11	Parkinson's disease (G20-G21)	9,609	1.3	183.0
12	Septicemia (A40-A41)	8,843	1.2	168.4
13	Pneumonitis due to solids and liquids (J69)	8,256	1.1	157.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,621	0.6	88.0
15	Atherosclerosis (I70)	3,678	0.5	70.1
...	All other causes (Residual)	172,993	23.6	3,294.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, all ages				
...	All causes	1,085,250	100.0	882.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	267,970	24.7	218.0
2	Malignant neoplasms (C00-C97)	261,975	24.1	213.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68,677	6.3	55.9
4	Chronic lower respiratory diseases (J40-J47)	61,452	5.7	50.0
5	Cerebrovascular diseases (I60-I69)	43,435	4.0	35.3
6	Diabetes mellitus (E10-E14)	30,832	2.8	25.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,681	2.6	23.3
8	Alzheimer's disease (G30)	23,607	2.2	19.2
9	Influenza and pneumonia (J09-J18)	20,575	1.9	16.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	19,973	1.8	16.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,269	1.7	14.9
12	Septicemia (A40-A41)	13,654	1.3	11.1
13	Parkinson's disease (G20-G21)	13,281	1.2	10.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,098	0.8	7.4
15	Pneumonitis due to solids and liquids (J69)	8,662	0.8	7.0
...	All other causes (Residual)	195,109	18.0	158.7
White, male, 1-4 years				
...	All causes	1,683	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	591	35.1	9.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	12.4	3.4
3	Malignant neoplasms (C00-C97)	165	9.8	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	6.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	2.8	0.8
6	Influenza and pneumonia (J09-J18)	42	2.5	0.7
7	Cerebrovascular diseases (I60-I69)	25	1.5	0.4
8	Septicemia (A40-A41)	19	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	16	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.2	*
14	Pneumonitis due to solids and liquids (J69)	4	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4	0.2	*
...	All other causes (Residual)	408	24.2	6.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 5-14 years				
...	All causes	2,132	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	676	31.7	4.2
2	Malignant neoplasms (C00-C97)	399	18.7	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	178	8.3	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	5.3	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	106	5.0	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	2.9	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	1.9	0.3
8	Cerebrovascular diseases (I60-I69)	33	1.5	0.2
9	Influenza and pneumonia (J09-J18)	29	1.4	0.2
10	Septicemia (A40-A41)	28	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	20	0.9	0.1
12	Diabetes mellitus (E10-E14)	10	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.4	*
14	Meningitis (G00,G03)	7	0.3	*
15	Anemias (D50-D64)	5	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (Residual)	412	19.3	2.6
White, male, 15-24 years				
...	All causes	15,362	100.0	89.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,452	48.5	43.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,253	21.2	19.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,279	8.3	7.5
4	Malignant neoplasms (C00-C97)	705	4.6	4.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	2.7	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	186	1.2	1.1
7	Cerebrovascular diseases (I60-I69)	77	0.5	0.4
8	Diabetes mellitus (E10-E14)	76	0.5	0.4
9	Influenza and pneumonia (J09-J18)	68	0.4	0.4
10	Legal intervention (Y35,Y89.0)	62	0.4	0.4
11	Chronic lower respiratory diseases (J40-J47)	49	0.3	0.3
12	Septicemia (A40-A41)	38	0.2	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	23	0.1	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.1	0.1
...	All other causes (Residual)	1,626	10.6	9.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 25-34 years				
...	All causes	23,107	100.0	139.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,940	43.0	59.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,239	18.3	25.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,525	6.6	9.2
4	Malignant neoplasms (C00-C97)	1,409	6.1	8.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,340	5.8	8.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	322	1.4	1.9
7	Diabetes mellitus (E10-E14)	249	1.1	1.5
8	Cerebrovascular diseases (I60-I69)	208	0.9	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	190	0.8	1.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	179	0.8	1.1
11	Influenza and pneumonia (J09-J18)	152	0.7	0.9
12	Septicemia (A40-A41)	124	0.5	0.7
13	Legal intervention (Y35,Y89.0)	94	0.4	0.6
14	Chronic lower respiratory diseases (J40-J47)	92	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	0.3	0.5
...	All other causes (Residual)	2,968	12.8	17.9
White, male, 35-44 years				
...	All causes	32,909	100.0	207.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,887	27.0	56.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,095	15.5	32.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,591	14.0	28.9
4	Malignant neoplasms (C00-C97)	3,843	11.7	24.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,348	4.1	8.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	953	2.9	6.0
7	Diabetes mellitus (E10-E14)	790	2.4	5.0
8	Cerebrovascular diseases (I60-I69)	642	2.0	4.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	433	1.3	2.7
10	Septicemia (A40-A41)	273	0.8	1.7
11	Influenza and pneumonia (J09-J18)	266	0.8	1.7
12	Chronic lower respiratory diseases (J40-J47)	236	0.7	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	214	0.7	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	0.6	1.2
15	Aortic aneurysm and dissection (I71)	153	0.5	1.0
...	All other causes (Residual)	5,002	15.2	31.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 45-54 years				
...	All causes	86,567	100.0	489.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,169	22.1	108.3
2	Malignant neoplasms (C00-C97)	19,155	22.1	108.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,554	13.3	65.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,108	7.1	34.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,317	6.1	30.0
6	Diabetes mellitus (E10-E14)	2,614	3.0	14.8
7	Cerebrovascular diseases (I60-I69)	2,020	2.3	11.4
8	Chronic lower respiratory diseases (J40-J47)	1,753	2.0	9.9
9	Viral hepatitis (B15-B19)	1,253	1.4	7.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,020	1.2	5.8
11	Septicemia (A40-A41)	909	1.1	5.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	868	1.0	4.9
13	Influenza and pneumonia (J09-J18)	768	0.9	4.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	750	0.9	4.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	528	0.6	3.0
...	All other causes (Residual)	12,781	14.8	72.2
White, male, 55-64 years				
...	All causes	162,259	100.0	1,043.5
1	Malignant neoplasms (C00-C97)	51,210	31.6	329.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,373	24.3	253.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,985	5.5	57.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	6,929	4.3	44.6
5	Chronic lower respiratory diseases (J40-J47)	6,911	4.3	44.4
6	Diabetes mellitus (E10-E14)	5,812	3.6	37.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,020	3.1	32.3
8	Cerebrovascular diseases (I60-I69)	4,569	2.8	29.4
9	Viral hepatitis (B15-B19)	2,247	1.4	14.5
10	Septicemia (A40-A41)	2,045	1.3	13.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,807	1.1	11.6
12	Influenza and pneumonia (J09-J18)	1,798	1.1	11.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,267	0.8	8.1
14	Aortic aneurysm and dissection (I71)	732	0.5	4.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	718	0.4	4.6
...	All other causes (Residual)	22,836	14.1	146.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 65-74 years				
...	All causes	207,886	100.0	2,150.5
1	Malignant neoplasms (C00-C97)	71,804	34.5	742.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,634	23.9	513.5
3	Chronic lower respiratory diseases (J40-J47)	15,293	7.4	158.2
4	Cerebrovascular diseases (I60-I69)	7,545	3.6	78.1
5	Diabetes mellitus (E10-E14)	7,503	3.6	77.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,839	2.8	60.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,652	1.8	37.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,262	1.6	33.7
9	Influenza and pneumonia (J09-J18)	2,893	1.4	29.9
10	Septicemia (A40-A41)	2,856	1.4	29.5
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,554	1.2	26.4
12	Parkinson's disease (G20-G21)	1,801	0.9	18.6
13	Alzheimer's disease (G30)	1,649	0.8	17.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,511	0.7	15.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,411	0.7	14.6
...	All other causes (Residual)	28,679	13.8	296.7
White, male, 75-84 years				
...	All causes	275,179	100.0	5,529.5
1	Malignant neoplasms (C00-C97)	72,311	26.3	1,453.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,303	25.2	1,392.6
3	Chronic lower respiratory diseases (J40-J47)	21,597	7.8	434.0
4	Cerebrovascular diseases (I60-I69)	13,483	4.9	270.9
5	Diabetes mellitus (E10-E14)	8,036	2.9	161.5
6	Alzheimer's disease (G30)	7,923	2.9	159.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,905	2.5	138.8
8	Parkinson's disease (G20-G21)	6,045	2.2	121.5
9	Influenza and pneumonia (J09-J18)	5,813	2.1	116.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,742	2.1	115.4
11	Septicemia (A40-A41)	3,866	1.4	77.7
12	Pneumonitis due to solids and liquids (J69)	2,480	0.9	49.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,344	0.9	47.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,291	0.8	46.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,859	0.7	37.4
...	All other causes (Residual)	45,181	16.4	907.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, male, 85 years and over				
...	All causes	269,654	100.0	15,271.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	83,208	30.9	4,712.4
2	Malignant neoplasms (C00-C97)	40,942	15.2	2,318.7
3	Chronic lower respiratory diseases (J40-J47)	15,480	5.7	876.7
4	Cerebrovascular diseases (I60-I69)	14,801	5.5	838.2
5	Alzheimer's disease (G30)	13,658	5.1	773.5
6	Influenza and pneumonia (J09-J18)	8,690	3.2	492.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,437	2.8	421.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,355	2.4	359.9
9	Diabetes mellitus (E10-E14)	5,738	2.1	325.0
10	Parkinson's disease (G20-G21)	5,090	1.9	288.3
11	Pneumonitis due to solids and liquids (J69)	4,117	1.5	233.2
12	Septicemia (A40-A41)	3,443	1.3	195.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,353	1.2	189.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,071	0.8	117.3
15	Atherosclerosis (I70)	1,184	0.4	67.1
...	All other causes (Residual)	54,087	20.1	3,063.1
White, female, all ages				
...	All causes	1,089,928	100.0	869.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	246,402	22.6	196.7
2	Malignant neoplasms (C00-C97)	234,910	21.6	187.5
3	Chronic lower respiratory diseases (J40-J47)	70,330	6.5	56.1
4	Cerebrovascular diseases (I60-I69)	64,529	5.9	51.5
5	Alzheimer's disease (G30)	52,983	4.9	42.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42,112	3.9	33.6
7	Diabetes mellitus (E10-E14)	26,974	2.5	21.5
8	Influenza and pneumonia (J09-J18)	23,010	2.1	18.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,797	1.6	14.2
10	Septicemia (A40-A41)	15,291	1.4	12.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,810	1.3	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,774	1.0	8.6
13	Parkinson's disease (G20-G21)	9,071	0.8	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,925	0.7	6.3
15	Pneumonitis due to solids and liquids (J69)	7,276	0.7	5.8
...	All other causes (Residual)	246,734	22.6	196.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 1-4 years				
...	All causes	1,275	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	32.4	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	158	12.4	2.7
3	Malignant neoplasms (C00-C97)	136	10.7	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	6.3	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	4.2	0.9
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	2.2	0.5
7	Influenza and pneumonia (J09-J18)	25	2.0	0.4
8	Septicemia (A40-A41)	18	1.4	*
9	Cerebrovascular diseases (I60-I69)	10	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	9	0.7	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
13	Anemias (D50-D64)	3	0.2	*
13	Meningitis (G00,G03)	3	0.2	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
13	Pneumonitis due to solids and liquids (J69)	3	0.2	*
...	All other causes (Residual)	319	25.0	5.4
White, female, 5-14 years				
...	All causes	1,579	100.0	10.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	411	26.0	2.7
2	Malignant neoplasms (C00-C97)	313	19.8	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	7.3	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	5.7	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	4.4	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	3.5	0.4
7	Cerebrovascular diseases (I60-I69)	34	2.2	0.2
7	Influenza and pneumonia (J09-J18)	34	2.2	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	33	2.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	21	1.3	0.1
11	Certain conditions originating in the perinatal period (P00-P96)	16	1.0	*
12	Septicemia (A40-A41)	15	0.9	*
13	Diabetes mellitus (E10-E14)	7	0.4	*
13	Meningitis (G00,G03)	7	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
...	All other causes (Residual)	351	22.2	2.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 15-24 years				
...	All causes	5,739	100.0	35.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,549	44.4	15.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	748	13.0	4.6
3	Malignant neoplasms (C00-C97)	513	8.9	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	300	5.2	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	203	3.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	135	2.4	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	98	1.7	0.6
8	Cerebrovascular diseases (I60-I69)	56	1.0	0.3
9	Diabetes mellitus (E10-E14)	55	1.0	0.3
10	Septicemia (A40-A41)	43	0.7	0.3
11	Influenza and pneumonia (J09-J18)	42	0.7	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.5	0.2
12	Chronic lower respiratory diseases (J40-J47)	26	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.2	*
...	All other causes (Residual)	914	15.9	5.7
White, female, 25-34 years				
...	All causes	10,263	100.0	64.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,546	34.6	22.3
2	Malignant neoplasms (C00-C97)	1,382	13.5	8.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,063	10.4	6.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	644	6.3	4.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	404	3.9	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	229	2.2	1.4
7	Diabetes mellitus (E10-E14)	167	1.6	1.1
8	Cerebrovascular diseases (I60-I69)	166	1.6	1.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	162	1.6	1.0
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	1.3	0.9
11	Influenza and pneumonia (J09-J18)	107	1.0	0.7
12	Septicemia (A40-A41)	92	0.9	0.6
13	Chronic lower respiratory diseases (J40-J47)	76	0.7	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	0.5	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.5	0.3
...	All other causes (Residual)	1,986	19.4	12.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 35-44 years				
...	All causes	19,319	100.0	124.1
1	Malignant neoplasms (C00-C97)	4,833	25.0	31.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,883	20.1	24.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,153	11.1	13.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,388	7.2	8.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	717	3.7	4.6
6	Cerebrovascular diseases (I60-I69)	498	2.6	3.2
7	Diabetes mellitus (E10-E14)	486	2.5	3.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	363	1.9	2.3
9	Chronic lower respiratory diseases (J40-J47)	290	1.5	1.9
10	Septicemia (A40-A41)	240	1.2	1.5
11	Influenza and pneumonia (J09-J18)	238	1.2	1.5
12	Pregnancy, childbirth and the puerperium (O00-O99)	156	0.8	1.0
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	150	0.8	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	137	0.7	0.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	117	0.6	0.8
...	All other causes (Residual)	3,670	19.0	23.6
White, female, 45-54 years				
...	All causes	53,572	100.0	300.1
1	Malignant neoplasms (C00-C97)	18,660	34.8	104.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,266	13.6	40.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,808	10.8	32.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,448	4.6	13.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,124	4.0	11.9
6	Chronic lower respiratory diseases (J40-J47)	2,019	3.8	11.3
7	Cerebrovascular diseases (I60-I69)	1,652	3.1	9.3
8	Diabetes mellitus (E10-E14)	1,450	2.7	8.1
9	Septicemia (A40-A41)	770	1.4	4.3
10	Influenza and pneumonia (J09-J18)	601	1.1	3.4
11	Viral hepatitis (B15-B19)	496	0.9	2.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	478	0.9	2.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	350	0.7	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	254	0.5	1.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	241	0.4	1.3
...	All other causes (Residual)	8,955	16.7	50.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 55-64 years				
...	All causes	101,398	100.0	620.5
1	Malignant neoplasms (C00-C97)	40,808	40.2	249.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,976	15.8	97.8
3	Chronic lower respiratory diseases (J40-J47)	6,479	6.4	39.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,260	4.2	26.1
5	Diabetes mellitus (E10-E14)	3,496	3.4	21.4
6	Cerebrovascular diseases (I60-I69)	3,234	3.2	19.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,854	2.8	17.5
8	Septicemia (A40-A41)	1,669	1.6	10.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,508	1.5	9.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,338	1.3	8.2
11	Influenza and pneumonia (J09-J18)	1,274	1.3	7.8
12	Viral hepatitis (B15-B19)	856	0.8	5.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	722	0.7	4.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	546	0.5	3.3
15	Alzheimer's disease (G30)	422	0.4	2.6
...	All other causes (Residual)	15,956	15.7	97.6
White, female, 65-74 years				
...	All causes	156,820	100.0	1,450.9
1	Malignant neoplasms (C00-C97)	57,879	36.9	535.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,201	17.3	251.7
3	Chronic lower respiratory diseases (J40-J47)	15,437	9.8	142.8
4	Cerebrovascular diseases (I60-I69)	6,738	4.3	62.3
5	Diabetes mellitus (E10-E14)	5,356	3.4	49.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,372	2.2	31.2
7	Septicemia (A40-A41)	2,691	1.7	24.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,670	1.7	24.7
9	Influenza and pneumonia (J09-J18)	2,262	1.4	20.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,243	1.4	20.8
11	Alzheimer's disease (G30)	2,160	1.4	20.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,280	0.8	11.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	911	0.6	8.4
14	Parkinson's disease (G20-G21)	896	0.6	8.3
15	Pneumonitis due to solids and liquids (J69)	681	0.4	6.3
...	All other causes (Residual)	25,043	16.0	231.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
White, female, 75-84 years				
...	All causes	270,300	100.0	4,110.6
1	Malignant neoplasms (C00-C97)	63,267	23.4	962.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,400	21.6	888.1
3	Chronic lower respiratory diseases (J40-J47)	23,461	8.7	356.8
4	Cerebrovascular diseases (I60-I69)	17,449	6.5	265.4
5	Alzheimer's disease (G30)	13,237	4.9	201.3
6	Diabetes mellitus (E10-E14)	7,476	2.8	113.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,163	2.3	93.7
8	Influenza and pneumonia (J09-J18)	5,502	2.0	83.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,079	1.9	77.2
10	Septicemia (A40-A41)	4,309	1.6	65.5
11	Parkinson's disease (G20-G21)	3,452	1.3	52.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,157	1.2	48.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,847	0.7	28.1
14	Pneumonitis due to solids and liquids (J69)	1,826	0.7	27.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,705	0.6	25.9
...	All other causes (Residual)	53,970	20.0	820.8
White, female, 85 years and over				
...	All causes	462,820	100.0	13,281.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	134,358	29.0	3,855.8
2	Malignant neoplasms (C00-C97)	47,091	10.2	1,351.4
3	Alzheimer's disease (G30)	37,106	8.0	1,064.9
4	Cerebrovascular diseases (I60-I69)	34,661	7.5	994.7
5	Chronic lower respiratory diseases (J40-J47)	22,506	4.9	645.9
6	Influenza and pneumonia (J09-J18)	12,888	2.8	369.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,400	2.5	327.2
8	Diabetes mellitus (E10-E14)	8,478	1.8	243.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,341	1.8	239.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,983	1.7	229.1
11	Septicemia (A40-A41)	5,400	1.2	155.0
12	Parkinson's disease (G20-G21)	4,519	1.0	129.7
13	Pneumonitis due to solids and liquids (J69)	4,139	0.9	118.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,550	0.6	73.2
15	Atherosclerosis (I70)	2,494	0.5	71.6
...	All other causes (Residual)	118,906	25.7	3,412.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, all ages				
...	All causes	295,222	100.0	683.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	70,123	23.8	162.3
2	Malignant neoplasms (C00-C97)	67,379	22.8	156.0
3	Cerebrovascular diseases (I60-I69)	15,894	5.4	36.8
4	Diabetes mellitus (E10-E14)	12,983	4.4	30.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,709	4.3	29.4
6	Chronic lower respiratory diseases (J40-J47)	9,375	3.2	21.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,241	2.8	19.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,207	2.8	19.0
9	Septicemia (A40-A41)	5,983	2.0	13.8
10	Alzheimer's disease (G30)	5,451	1.8	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,200	1.8	12.0
12	Influenza and pneumonia (J09-J18)	5,000	1.7	11.6
13	Certain conditions originating in the perinatal period (P00-P96)	3,958	1.3	9.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3,921	1.3	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,791	0.9	6.5
...	All other causes (Residual)	58,007	19.6	134.3
Black, both sexes, 1-4 years				
...	All causes	1,021	100.0	37.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	280	27.4	10.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	13.0	4.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	10.6	3.9
4	Malignant neoplasms (C00-C97)	65	6.4	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	4.3	1.6
6	Chronic lower respiratory diseases (J40-J47)	30	2.9	1.1
7	Influenza and pneumonia (J09-J18)	21	2.1	0.8
8	Septicemia (A40-A41)	18	1.8	*
8	Cerebrovascular diseases (I60-I69)	18	1.8	*
10	Certain conditions originating in the perinatal period (P00-P96)	17	1.7	*
11	Anemias (D50-D64)	7	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
14	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	261	25.6	9.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 5-14 years				
...	All causes	1,196	100.0	17.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	30.0	5.3
2	Malignant neoplasms (C00-C97)	148	12.4	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	110	9.2	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	6.4	1.1
5	Chronic lower respiratory diseases (J40-J47)	75	6.3	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	3.7	0.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.6	0.6
8	Diabetes mellitus (E10-E14)	17	1.4	*
9	Influenza and pneumonia (J09-J18)	16	1.3	*
10	Cerebrovascular diseases (I60-I69)	15	1.3	*
11	Anemias (D50-D64)	13	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
14	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
15	Septicemia (A40-A41)	7	0.6	*
...	All other causes (Residual)	244	20.4	3.6
Black, both sexes, 15-24 years				
...	All causes	6,705	100.0	91.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,906	43.3	39.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,396	20.8	19.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	524	7.8	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	289	4.3	3.9
5	Malignant neoplasms (C00-C97)	250	3.7	3.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	87	1.3	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	85	1.3	1.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	68	1.0	0.9
9	Anemias (D50-D64)	65	1.0	0.9
10	Chronic lower respiratory diseases (J40-J47)	62	0.9	0.8
11	Diabetes mellitus (E10-E14)	58	0.9	0.8
12	Legal intervention (Y35,Y89.0)	45	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	43	0.6	0.6
14	Septicemia (A40-A41)	33	0.5	0.5
15	Influenza and pneumonia (J09-J18)	32	0.5	0.4
...	All other causes (Residual)	762	11.4	10.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 25-34 years				
...	All causes	9,105	100.0	148.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,433	26.7	39.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,734	19.0	28.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	897	9.9	14.7
4	Malignant neoplasms (C00-C97)	653	7.2	10.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	562	6.2	9.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	423	4.6	6.9
7	Diabetes mellitus (E10-E14)	204	2.2	3.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	135	1.5	2.2
9	Cerebrovascular diseases (I60-I69)	124	1.4	2.0
10	Anemias (D50-D64)	113	1.2	1.8
10	Chronic lower respiratory diseases (J40-J47)	113	1.2	1.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	92	1.0	1.5
13	Septicemia (A40-A41)	90	1.0	1.5
14	Influenza and pneumonia (J09-J18)	81	0.9	1.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	0.6	0.9
...	All other causes (Residual)	1,393	15.3	22.8
Black, both sexes, 35-44 years				
...	All causes	13,693	100.0	246.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,799	20.4	50.3
2	Malignant neoplasms (C00-C97)	1,948	14.2	35.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,685	12.3	30.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,290	9.4	23.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	760	5.6	13.7
6	Diabetes mellitus (E10-E14)	508	3.7	9.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	478	3.5	8.6
8	Cerebrovascular diseases (I60-I69)	470	3.4	8.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	256	1.9	4.6
10	Septicemia (A40-A41)	216	1.6	3.9
11	Chronic lower respiratory diseases (J40-J47)	196	1.4	3.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	196	1.4	3.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	160	1.2	2.9
14	Influenza and pneumonia (J09-J18)	152	1.1	2.7
15	Anemias (D50-D64)	146	1.1	2.6
...	All other causes (Residual)	2,433	17.8	43.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 45-54 years				
...	All causes	32,955	100.0	571.8
1	Malignant neoplasms (C00-C97)	8,209	24.9	142.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,767	23.6	134.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,379	7.2	41.3
4	Cerebrovascular diseases (I60-I69)	1,667	5.1	28.9
5	Diabetes mellitus (E10-E14)	1,410	4.3	24.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,307	4.0	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	760	2.3	13.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	758	2.3	13.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	720	2.2	12.5
10	Chronic lower respiratory diseases (J40-J47)	691	2.1	12.0
11	Septicemia (A40-A41)	588	1.8	10.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	535	1.6	9.3
13	Influenza and pneumonia (J09-J18)	393	1.2	6.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	373	1.1	6.5
15	Viral hepatitis (B15-B19)	240	0.7	4.2
...	All other causes (Residual)	5,158	15.7	89.5
Black, both sexes, 55-64 years				
...	All causes	55,658	100.0	1,253.3
1	Malignant neoplasms (C00-C97)	17,242	31.0	388.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,920	25.0	313.4
3	Cerebrovascular diseases (I60-I69)	2,718	4.9	61.2
4	Diabetes mellitus (E10-E14)	2,692	4.8	60.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,097	3.8	47.2
6	Chronic lower respiratory diseases (J40-J47)	1,567	2.8	35.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,484	2.7	33.4
8	Septicemia (A40-A41)	1,144	2.1	25.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,110	2.0	25.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	966	1.7	21.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	912	1.6	20.5
12	Influenza and pneumonia (J09-J18)	708	1.3	15.9
13	Viral hepatitis (B15-B19)	636	1.1	14.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	344	0.6	7.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	210	0.4	4.7
...	All other causes (Residual)	7,908	14.2	178.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 65-74 years				
...	All causes	55,151	100.0	2,415.5
1	Malignant neoplasms (C00-C97)	17,554	31.8	768.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,744	24.9	602.0
3	Diabetes mellitus (E10-E14)	3,116	5.6	136.5
4	Cerebrovascular diseases (I60-I69)	3,065	5.6	134.2
5	Chronic lower respiratory diseases (J40-J47)	2,384	4.3	104.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,788	3.2	78.3
7	Septicemia (A40-A41)	1,296	2.3	56.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	991	1.8	43.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	941	1.7	41.2
10	Influenza and pneumonia (J09-J18)	868	1.6	38.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	486	0.9	21.3
12	Alzheimer's disease (G30)	409	0.7	17.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	301	0.5	13.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	247	0.4	10.8
15	Pneumonitis due to solids and liquids (J69)	231	0.4	10.1
...	All other causes (Residual)	7,730	14.0	338.6
Black, both sexes, 75-84 years				
...	All causes	58,256	100.0	5,154.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,987	25.7	1,326.0
2	Malignant neoplasms (C00-C97)	14,109	24.2	1,248.3
3	Cerebrovascular diseases (I60-I69)	3,904	6.7	345.4
4	Diabetes mellitus (E10-E14)	2,977	5.1	263.4
5	Chronic lower respiratory diseases (J40-J47)	2,502	4.3	221.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,040	3.5	180.5
7	Alzheimer's disease (G30)	1,705	2.9	150.9
8	Septicemia (A40-A41)	1,391	2.4	123.1
9	Influenza and pneumonia (J09-J18)	1,190	2.0	105.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,175	2.0	104.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	808	1.4	71.5
12	Pneumonitis due to solids and liquids (J69)	419	0.7	37.1
13	Parkinson's disease (G20-G21)	396	0.7	35.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	320	0.5	28.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	186	0.3	16.5
...	All other causes (Residual)	10,147	17.4	897.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, both sexes, 85 years and over				
...	All causes	54,344	100.0	12,615.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,516	28.6	3,602.0
2	Malignant neoplasms (C00-C97)	7,188	13.2	1,668.7
3	Cerebrovascular diseases (I60-I69)	3,833	7.1	889.8
4	Alzheimer's disease (G30)	3,266	6.0	758.2
5	Diabetes mellitus (E10-E14)	2,000	3.7	464.3
6	Chronic lower respiratory diseases (J40-J47)	1,748	3.2	405.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,730	3.2	401.6
8	Influenza and pneumonia (J09-J18)	1,483	2.7	344.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,421	2.6	329.9
10	Septicemia (A40-A41)	1,126	2.1	261.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	579	1.1	134.4
12	Pneumonitis due to solids and liquids (J69)	515	0.9	119.6
13	Parkinson's disease (G20-G21)	301	0.6	69.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	246	0.5	57.1
15	Atherosclerosis (I70)	200	0.4	46.4
...	All other causes (Residual)	13,192	24.3	3,062.5
Black, male, all ages				
...	All causes	150,586	100.0	728.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,140	24.0	174.7
2	Malignant neoplasms (C00-C97)	34,427	22.9	166.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,583	5.7	41.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,129	4.7	34.5
5	Cerebrovascular diseases (I60-I69)	7,061	4.7	34.1
6	Diabetes mellitus (E10-E14)	6,145	4.1	29.7
7	Chronic lower respiratory diseases (J40-J47)	4,885	3.2	23.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,818	2.5	18.5
9	Septicemia (A40-A41)	2,709	1.8	13.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,546	1.7	12.3
11	Influenza and pneumonia (J09-J18)	2,404	1.6	11.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,310	1.5	11.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,235	1.5	10.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,908	1.3	9.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,807	1.2	8.7
...	All other causes (Residual)	26,479	17.6	128.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 1-4 years				
...	All causes	578	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	29.2	12.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	13.7	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	8.8	3.7
4	Malignant neoplasms (C00-C97)	39	6.7	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.8	2.0
6	Chronic lower respiratory diseases (J40-J47)	24	4.2	1.7
7	Influenza and pneumonia (J09-J18)	12	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	1.9	*
9	Septicemia (A40-A41)	9	1.6	*
9	Cerebrovascular diseases (I60-I69)	9	1.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Anemias (D50-D64)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	130	22.5	9.3
Black, male, 5-14 years				
...	All causes	731	100.0	21.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	32.6	6.9
2	Malignant neoplasms (C00-C97)	83	11.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	10.0	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	6.3	1.3
5	Chronic lower respiratory diseases (J40-J47)	40	5.5	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	4.5	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	3.3	0.7
8	Anemias (D50-D64)	8	1.1	*
8	Diabetes mellitus (E10-E14)	8	1.1	*
8	Influenza and pneumonia (J09-J18)	8	1.1	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	8	1.1	*
12	Cerebrovascular diseases (I60-I69)	7	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
14	Septicemia (A40-A41)	4	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	143	19.6	4.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 15-24 years				
...	All causes	5,127	100.0	138.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,617	51.0	70.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,020	19.9	27.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	423	8.3	11.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	193	3.8	5.2
5	Malignant neoplasms (C00-C97)	148	2.9	4.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.2	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	1.1	1.5
8	Chronic lower respiratory diseases (J40-J47)	46	0.9	1.2
9	Legal intervention (Y35,Y89.0)	43	0.8	1.2
10	Anemias (D50-D64)	34	0.7	0.9
11	Diabetes mellitus (E10-E14)	27	0.5	0.7
12	Cerebrovascular diseases (I60-I69)	24	0.5	0.6
13	Influenza and pneumonia (J09-J18)	20	0.4	0.5
14	Septicemia (A40-A41)	13	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.1	*
...	All other causes (Residual)	396	7.7	10.7
Black, male, 25-34 years				
...	All causes	6,267	100.0	212.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,177	34.7	73.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,277	20.4	43.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	567	9.0	19.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	482	7.7	16.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	268	4.3	9.1
5	Malignant neoplasms (C00-C97)	268	4.3	9.1
7	Diabetes mellitus (E10-E14)	101	1.6	3.4
8	Cerebrovascular diseases (I60-I69)	71	1.1	2.4
8	Chronic lower respiratory diseases (J40-J47)	71	1.1	2.4
10	Anemias (D50-D64)	59	0.9	2.0
11	Septicemia (A40-A41)	48	0.8	1.6
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	0.7	1.4
13	Influenza and pneumonia (J09-J18)	41	0.7	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	0.7	1.4
15	Legal intervention (Y35,Y89.0)	40	0.6	1.4
...	All other causes (Residual)	714	11.4	24.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 35-44 years				
...	All causes	8,019	100.0	306.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,796	22.4	68.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,184	14.8	45.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,115	13.9	42.6
4	Malignant neoplasms (C00-C97)	697	8.7	26.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	394	4.9	15.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	368	4.6	14.0
7	Diabetes mellitus (E10-E14)	302	3.8	11.5
8	Cerebrovascular diseases (I60-I69)	259	3.2	9.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	141	1.8	5.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	119	1.5	4.5
11	Septicemia (A40-A41)	99	1.2	3.8
11	Chronic lower respiratory diseases (J40-J47)	99	1.2	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	88	1.1	3.4
14	Influenza and pneumonia (J09-J18)	74	0.9	2.8
15	Anemias (D50-D64)	65	0.8	2.5
...	All other causes (Residual)	1,219	15.2	46.5
Black, male, 45-54 years				
...	All causes	18,852	100.0	696.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,938	26.2	182.4
2	Malignant neoplasms (C00-C97)	3,926	20.8	145.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,607	8.5	59.4
4	Cerebrovascular diseases (I60-I69)	903	4.8	33.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	848	4.5	31.3
6	Diabetes mellitus (E10-E14)	814	4.3	30.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	601	3.2	22.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	439	2.3	16.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	410	2.2	15.1
10	Chronic lower respiratory diseases (J40-J47)	342	1.8	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	308	1.6	11.4
12	Septicemia (A40-A41)	305	1.6	11.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	286	1.5	10.6
14	Influenza and pneumonia (J09-J18)	221	1.2	8.2
15	Viral hepatitis (B15-B19)	167	0.9	6.2
...	All other causes (Residual)	2,737	14.5	101.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 55-64 years				
...	All causes	32,662	100.0	1,615.8
1	Malignant neoplasms (C00-C97)	9,477	29.0	468.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,812	27.0	435.9
3	Cerebrovascular diseases (I60-I69)	1,537	4.7	76.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,504	4.6	74.4
5	Diabetes mellitus (E10-E14)	1,492	4.6	73.8
6	Chronic lower respiratory diseases (J40-J47)	826	2.5	40.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	815	2.5	40.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	783	2.4	38.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	683	2.1	33.8
10	Septicemia (A40-A41)	588	1.8	29.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	536	1.6	26.5
12	Viral hepatitis (B15-B19)	425	1.3	21.0
13	Influenza and pneumonia (J09-J18)	417	1.3	20.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	290	0.9	14.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	167	0.5	8.3
...	All other causes (Residual)	4,310	13.2	213.2
Black, male, 65-74 years				
...	All causes	30,118	100.0	3,077.9
1	Malignant neoplasms (C00-C97)	9,494	31.5	970.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,882	26.2	805.5
3	Cerebrovascular diseases (I60-I69)	1,614	5.4	164.9
4	Diabetes mellitus (E10-E14)	1,607	5.3	164.2
5	Chronic lower respiratory diseases (J40-J47)	1,353	4.5	138.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	908	3.0	92.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	645	2.1	65.9
8	Septicemia (A40-A41)	618	2.1	63.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	487	1.6	49.8
10	Influenza and pneumonia (J09-J18)	469	1.6	47.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	328	1.1	33.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	232	0.8	23.7
13	Alzheimer's disease (G30)	163	0.5	16.7
14	Pneumonitis due to solids and liquids (J69)	154	0.5	15.7
15	Parkinson's disease (G20-G21)	124	0.4	12.7
...	All other causes (Residual)	4,040	13.4	412.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, male, 75-84 years				
...	All causes	27,217	100.0	6,416.8
1	Malignant neoplasms (C00-C97)	7,150	26.3	1,685.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,116	26.1	1,677.7
3	Cerebrovascular diseases (I60-I69)	1,633	6.0	385.0
4	Chronic lower respiratory diseases (J40-J47)	1,307	4.8	308.1
5	Diabetes mellitus (E10-E14)	1,249	4.6	294.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	906	3.3	213.6
7	Alzheimer's disease (G30)	645	2.4	152.1
8	Septicemia (A40-A41)	586	2.2	138.2
9	Influenza and pneumonia (J09-J18)	575	2.1	135.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	490	1.8	115.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	463	1.7	109.2
12	Parkinson's disease (G20-G21)	252	0.9	59.4
13	Pneumonitis due to solids and liquids (J69)	237	0.9	55.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	150	0.6	35.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	104	0.4	24.5
...	All other causes (Residual)	4,354	16.0	1,026.5
Black, male, 85 years and over				
...	All causes	17,012	100.0	13,775.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,720	27.7	3,822.1
2	Malignant neoplasms (C00-C97)	3,137	18.4	2,540.2
3	Cerebrovascular diseases (I60-I69)	984	5.8	796.8
4	Chronic lower respiratory diseases (J40-J47)	773	4.5	625.9
5	Alzheimer's disease (G30)	712	4.2	576.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	573	3.4	464.0
7	Diabetes mellitus (E10-E14)	544	3.2	440.5
8	Influenza and pneumonia (J09-J18)	536	3.2	434.0
9	Septicemia (A40-A41)	389	2.3	315.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	371	2.2	300.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	1.4	192.7
12	Pneumonitis due to solids and liquids (J69)	207	1.2	167.6
13	Parkinson's disease (G20-G21)	146	0.9	118.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.5	75.3
15	Atherosclerosis (I70)	63	0.4	51.0
...	All other causes (Residual)	3,526	20.7	2,855.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, all ages				
...	All causes	144,636	100.0	642.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,983	23.5	150.9
2	Malignant neoplasms (C00-C97)	32,952	22.8	146.3
3	Cerebrovascular diseases (I60-I69)	8,833	6.1	39.2
4	Diabetes mellitus (E10-E14)	6,838	4.7	30.4
5	Chronic lower respiratory diseases (J40-J47)	4,490	3.1	19.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,389	3.0	19.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,126	2.9	18.3
8	Alzheimer's disease (G30)	3,899	2.7	17.3
9	Septicemia (A40-A41)	3,274	2.3	14.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,890	2.0	12.8
11	Influenza and pneumonia (J09-J18)	2,596	1.8	11.5
12	Certain conditions originating in the perinatal period (P00-P96)	1,723	1.2	7.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,375	1.0	6.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,112	0.8	4.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	984	0.7	4.4
...	All other causes (Residual)	31,172	21.6	138.4
Black, female, 1-4 years				
...	All causes	443	100.0	33.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	25.1	8.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	12.9	4.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	12.2	4.0
4	Malignant neoplasms (C00-C97)	26	5.9	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.6	*
6	Septicemia (A40-A41)	9	2.0	*
6	Cerebrovascular diseases (I60-I69)	9	2.0	*
6	Influenza and pneumonia (J09-J18)	9	2.0	*
9	Anemias (D50-D64)	6	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Meningitis (G00,G03)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	122	27.5	9.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 5-14 years				
...	All causes	465	100.0	14.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	26.0	3.7
2	Malignant neoplasms (C00-C97)	65	14.0	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	8.0	1.1
4	Chronic lower respiratory diseases (J40-J47)	35	7.5	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	6.7	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	4.3	0.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	2.2	*
8	Diabetes mellitus (E10-E14)	9	1.9	*
9	Cerebrovascular diseases (I60-I69)	8	1.7	*
9	Influenza and pneumonia (J09-J18)	8	1.7	*
11	Anemias (D50-D64)	5	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
14	Septicemia (A40-A41)	3	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
...	All other causes (Residual)	101	21.7	3.0
Black, female, 15-24 years				
...	All causes	1,578	100.0	43.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	376	23.8	10.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	289	18.3	8.0
3	Malignant neoplasms (C00-C97)	102	6.5	2.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	6.4	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	6.1	2.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	68	4.3	1.9
7	Anemias (D50-D64)	31	2.0	0.9
7	Diabetes mellitus (E10-E14)	31	2.0	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	1.9	0.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.6	0.7
11	Septicemia (A40-A41)	20	1.3	0.6
12	Cerebrovascular diseases (I60-I69)	19	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.2	*
14	Chronic lower respiratory diseases (J40-J47)	16	1.0	*
15	Influenza and pneumonia (J09-J18)	12	0.8	*
...	All other causes (Residual)	342	21.7	9.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 25-34 years				
...	All causes	2,838	100.0	89.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	457	16.1	14.4
2	Malignant neoplasms (C00-C97)	385	13.6	12.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	330	11.6	10.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	256	9.0	8.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	155	5.5	4.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	135	4.8	4.3
7	Diabetes mellitus (E10-E14)	103	3.6	3.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	2.8	2.5
9	Anemias (D50-D64)	54	1.9	1.7
10	Cerebrovascular diseases (I60-I69)	53	1.9	1.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	1.8	1.6
12	Septicemia (A40-A41)	42	1.5	1.3
12	Chronic lower respiratory diseases (J40-J47)	42	1.5	1.3
14	Influenza and pneumonia (J09-J18)	40	1.4	1.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	25	0.9	0.8
...	All other causes (Residual)	630	22.2	19.9
Black, female, 35-44 years				
...	All causes	5,674	100.0	192.9
1	Malignant neoplasms (C00-C97)	1,251	22.0	42.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,003	17.7	34.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	501	8.8	17.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	366	6.5	12.4
5	Cerebrovascular diseases (I60-I69)	211	3.7	7.2
6	Diabetes mellitus (E10-E14)	206	3.6	7.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	175	3.1	5.9
8	Septicemia (A40-A41)	117	2.1	4.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	2.0	3.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	110	1.9	3.7
11	Chronic lower respiratory diseases (J40-J47)	97	1.7	3.3
12	Pregnancy, childbirth and the puerperium (O00-O99)	95	1.7	3.2
13	Anemias (D50-D64)	81	1.4	2.8
14	Influenza and pneumonia (J09-J18)	78	1.4	2.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	77	1.4	2.6
...	All other causes (Residual)	1,191	21.0	40.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 45-54 years				
...	All causes	14,103	100.0	461.5
1	Malignant neoplasms (C00-C97)	4,283	30.4	140.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,829	20.1	92.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	772	5.5	25.3
4	Cerebrovascular diseases (I60-I69)	764	5.4	25.0
5	Diabetes mellitus (E10-E14)	596	4.2	19.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	459	3.3	15.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	350	2.5	11.5
8	Chronic lower respiratory diseases (J40-J47)	349	2.5	11.4
9	Septicemia (A40-A41)	283	2.0	9.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	281	2.0	9.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	227	1.6	7.4
12	Influenza and pneumonia (J09-J18)	172	1.2	5.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	157	1.1	5.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	0.6	2.8
15	Viral hepatitis (B15-B19)	73	0.5	2.4
15	Anemias (D50-D64)	73	0.5	2.4
...	All other causes (Residual)	2,348	16.6	76.8
Black, female, 55-64 years				
...	All causes	22,996	100.0	950.4
1	Malignant neoplasms (C00-C97)	7,765	33.8	320.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,108	22.2	211.1
3	Diabetes mellitus (E10-E14)	1,200	5.2	49.6
4	Cerebrovascular diseases (I60-I69)	1,181	5.1	48.8
5	Chronic lower respiratory diseases (J40-J47)	741	3.2	30.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	669	2.9	27.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	593	2.6	24.5
8	Septicemia (A40-A41)	556	2.4	23.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	376	1.6	15.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	327	1.4	13.5
11	Influenza and pneumonia (J09-J18)	291	1.3	12.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	283	1.2	11.7
13	Viral hepatitis (B15-B19)	211	0.9	8.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	92	0.4	3.8
15	Anemias (D50-D64)	83	0.4	3.4
...	All other causes (Residual)	3,520	15.3	145.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 65-74 years				
...	All causes	25,033	100.0	1,918.7
1	Malignant neoplasms (C00-C97)	8,060	32.2	617.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,862	23.4	449.3
3	Diabetes mellitus (E10-E14)	1,509	6.0	115.7
4	Cerebrovascular diseases (I60-I69)	1,451	5.8	111.2
5	Chronic lower respiratory diseases (J40-J47)	1,031	4.1	79.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	880	3.5	67.4
7	Septicemia (A40-A41)	678	2.7	52.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	454	1.8	34.8
9	Influenza and pneumonia (J09-J18)	399	1.6	30.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	346	1.4	26.5
11	Alzheimer's disease (G30)	246	1.0	18.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	158	0.6	12.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	126	0.5	9.7
14	Anemias (D50-D64)	83	0.3	6.4
15	Aortic aneurysm and dissection (I71)	80	0.3	6.1
...	All other causes (Residual)	3,670	14.7	281.3
Black, female, 75-84 years				
...	All causes	31,039	100.0	4,396.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,871	25.4	1,114.8
2	Malignant neoplasms (C00-C97)	6,959	22.4	985.6
3	Cerebrovascular diseases (I60-I69)	2,271	7.3	321.6
4	Diabetes mellitus (E10-E14)	1,728	5.6	244.7
5	Chronic lower respiratory diseases (J40-J47)	1,195	3.8	169.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,134	3.7	160.6
7	Alzheimer's disease (G30)	1,060	3.4	150.1
8	Septicemia (A40-A41)	805	2.6	114.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	685	2.2	97.0
10	Influenza and pneumonia (J09-J18)	615	2.0	87.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	345	1.1	48.9
12	Pneumonitis due to solids and liquids (J69)	182	0.6	25.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	170	0.5	24.1
14	Parkinson's disease (G20-G21)	144	0.5	20.4
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	98	0.3	13.9
...	All other causes (Residual)	5,777	18.6	818.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Black, female, 85 years and over				
...	All causes	37,332	100.0	12,149.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,796	28.9	3,513.5
2	Malignant neoplasms (C00-C97)	4,051	10.9	1,318.4
3	Cerebrovascular diseases (I60-I69)	2,849	7.6	927.2
4	Alzheimer's disease (G30)	2,554	6.8	831.2
5	Diabetes mellitus (E10-E14)	1,456	3.9	473.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,157	3.1	376.5
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,050	2.8	341.7
8	Chronic lower respiratory diseases (J40-J47)	975	2.6	317.3
9	Influenza and pneumonia (J09-J18)	947	2.5	308.2
10	Septicemia (A40-A41)	737	2.0	239.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	341	0.9	111.0
12	Pneumonitis due to solids and liquids (J69)	308	0.8	100.2
13	Parkinson's disease (G20-G21)	155	0.4	50.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	153	0.4	49.8
15	Atherosclerosis (I70)	137	0.4	44.6
...	All other causes (Residual)	9,666	25.9	3,145.8
American Indian, both sexes, all ages				
...	All causes	16,527	100.0	375.7
1	Malignant neoplasms (C00-C97)	3,019	18.3	68.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,950	17.8	67.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,922	11.6	43.7
4	Diabetes mellitus (E10-E14)	985	6.0	22.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	918	5.6	20.9
6	Chronic lower respiratory diseases (J40-J47)	708	4.3	16.1
7	Cerebrovascular diseases (I60-I69)	580	3.5	13.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	485	2.9	11.0
9	Influenza and pneumonia (J09-J18)	306	1.9	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	295	1.8	6.7
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	263	1.6	6.0
12	Septicemia (A40-A41)	255	1.5	5.8
13	Alzheimer's disease (G30)	217	1.3	4.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	150	0.9	3.4
15	Certain conditions originating in the perinatal period (P00-P96)	129	0.8	2.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	129	0.8	2.9
...	All other causes (Residual)	3,216	19.5	73.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 1-4 years				
...	All causes	89	100.0	28.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	43.8	12.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	12.4	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	6.7	*
4	Malignant neoplasms (C00-C97)	5	5.6	*
5	Septicemia (A40-A41)	3	3.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.1	*
6	Cerebrovascular diseases (I60-I69)	1	1.1	*
6	Influenza and pneumonia (J09-J18)	1	1.1	*
6	Hernia (K40-K46)	1	1.1	*
...	All other causes (Residual)	21	23.6	6.7
American Indian, both sexes, 5-14 years				
...	All causes	101	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	50.5	6.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	9.9	*
3	Malignant neoplasms (C00-C97)	6	5.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	5.0	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	4.0	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.0	*
7	Influenza and pneumonia (J09-J18)	2	2.0	*
9	Anemias (D50-D64)	1	1.0	*
9	Cerebrovascular diseases (I60-I69)	1	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (Residual)	15	14.9	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 15-24 years				
...	All causes	604	100.0	78.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	44.5	35.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	25.2	19.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	10.6	8.3
4	Malignant neoplasms (C00-C97)	15	2.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.3	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.8	*
8	Diabetes mellitus (E10-E14)	4	0.7	*
8	Legal intervention (Y35,Y89.0)	4	0.7	*
10	Septicemia (A40-A41)	3	0.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.3	*
12	Pneumonitis due to solids and liquids (J69)	2	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	2	0.3	*
...	All other causes (Residual)	57	9.4	7.4
American Indian, both sexes, 25-34 years				
...	All causes	934	100.0	138.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	354	37.9	52.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	118	12.6	17.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	8.8	12.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	65	7.0	9.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	6.6	9.2
6	Malignant neoplasms (C00-C97)	34	3.6	5.0
7	Diabetes mellitus (E10-E14)	13	1.4	*
8	Septicemia (A40-A41)	11	1.2	*
8	Influenza and pneumonia (J09-J18)	11	1.2	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	10	1.1	*
11	Cerebrovascular diseases (I60-I69)	8	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.5	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.5	*
14	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Legal intervention (Y35,Y89.0)	4	0.4	*
...	All other causes (Residual)	148	15.8	21.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 35-44 years				
...	All causes	1,198	100.0	206.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	330	27.5	56.8
2	Chronic liver disease and cirrhosis (K70,K73-K74)	170	14.2	29.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	136	11.4	23.4
4	Malignant neoplasms (C00-C97)	88	7.3	15.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	6.6	13.6
6	Diabetes mellitus (E10-E14)	40	3.3	6.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	3.3	6.7
8	Septicemia (A40-A41)	16	1.3	*
9	Cerebrovascular diseases (I60-I69)	14	1.2	*
10	Viral hepatitis (B15-B19)	13	1.1	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.1	*
13	Influenza and pneumonia (J09-J18)	10	0.8	*
14	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	7	0.6	*
...	All other causes (Residual)	223	18.6	38.4
American Indian, both sexes, 45-54 years				
...	All causes	2,264	100.0	414.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	381	16.8	69.7
2	Malignant neoplasms (C00-C97)	351	15.5	64.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	345	15.2	63.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	302	13.3	55.2
5	Diabetes mellitus (E10-E14)	130	5.7	23.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	70	3.1	12.8
7	Cerebrovascular diseases (I60-I69)	49	2.2	9.0
8	Chronic lower respiratory diseases (J40-J47)	40	1.8	7.3
9	Septicemia (A40-A41)	37	1.6	6.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	1.6	6.6
11	Viral hepatitis (B15-B19)	34	1.5	6.2
12	Influenza and pneumonia (J09-J18)	31	1.4	5.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.1	4.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.6	*
...	All other causes (Residual)	404	17.8	73.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 55-64 years				
...	All causes	2,922	100.0	756.6
1	Malignant neoplasms (C00-C97)	755	25.8	195.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	537	18.4	139.0
3	Diabetes mellitus (E10-E14)	233	8.0	60.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	230	7.9	59.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	214	7.3	55.4
6	Chronic lower respiratory diseases (J40-J47)	141	4.8	36.5
7	Cerebrovascular diseases (I60-I69)	90	3.1	23.3
8	Influenza and pneumonia (J09-J18)	50	1.7	12.9
9	Viral hepatitis (B15-B19)	48	1.6	12.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.5	11.7
11	Septicemia (A40-A41)	43	1.5	11.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	1.2	9.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	1.1	8.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	0.6	*
15	Pneumonitis due to solids and liquids (J69)	11	0.4	*
...	All other causes (Residual)	439	15.0	113.7
American Indian, both sexes, 65-74 years				
...	All causes	3,127	100.0	1,652.5
1	Malignant neoplasms (C00-C97)	892	28.5	471.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	647	20.7	341.9
3	Diabetes mellitus (E10-E14)	261	8.3	137.9
4	Chronic lower respiratory diseases (J40-J47)	208	6.7	109.9
5	Cerebrovascular diseases (I60-I69)	134	4.3	70.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	3.6	59.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	108	3.5	57.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	71	2.3	37.5
9	Septicemia (A40-A41)	51	1.6	27.0
10	Influenza and pneumonia (J09-J18)	46	1.5	24.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	1.0	15.9
12	Alzheimer's disease (G30)	21	0.7	11.1
13	Pneumonitis due to solids and liquids (J69)	18	0.6	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	0.5	*
15	Viral hepatitis (B15-B19)	14	0.4	*
...	All other causes (Residual)	497	15.9	262.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, both sexes, 75-84 years				
...	All causes	2,861	100.0	3,651.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	693	24.2	884.4
2	Malignant neoplasms (C00-C97)	634	22.2	809.1
3	Diabetes mellitus (E10-E14)	202	7.1	257.8
4	Chronic lower respiratory diseases (J40-J47)	198	6.9	252.7
5	Cerebrovascular diseases (I60-I69)	150	5.2	191.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	2.7	97.0
7	Influenza and pneumonia (J09-J18)	72	2.5	91.9
8	Alzheimer's disease (G30)	71	2.5	90.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	2.4	86.8
10	Septicemia (A40-A41)	57	2.0	72.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	42	1.5	53.6
12	Parkinson's disease (G20-G21)	27	0.9	34.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	0.9	33.2
14	Pneumonitis due to solids and liquids (J69)	21	0.7	26.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.6	*
...	All other causes (Residual)	508	17.8	648.3
American Indian, both sexes, 85 years and over				
...	All causes	2,057	100.0	8,206.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	505	24.6	2,014.6
2	Malignant neoplasms (C00-C97)	237	11.5	945.5
3	Cerebrovascular diseases (I60-I69)	131	6.4	522.6
4	Alzheimer's disease (G30)	120	5.8	478.7
5	Chronic lower respiratory diseases (J40-J47)	110	5.3	438.8
6	Diabetes mellitus (E10-E14)	102	5.0	406.9
7	Influenza and pneumonia (J09-J18)	78	3.8	311.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	2.9	235.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	2.9	235.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	2.0	163.6
11	Septicemia (A40-A41)	31	1.5	123.7
12	Pneumonitis due to solids and liquids (J69)	26	1.3	103.7
13	Parkinson's disease (G20-G21)	23	1.1	91.8
14	Anemias (D50-D64)	11	0.5	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	11	0.5	*
...	All other causes (Residual)	513	24.9	2,046.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, all ages				
...	All causes	9,067	100.0	410.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,731	19.1	78.3
2	Malignant neoplasms (C00-C97)	1,559	17.2	70.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,278	14.1	57.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	517	5.7	23.4
5	Diabetes mellitus (E10-E14)	514	5.7	23.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	386	4.3	17.5
7	Chronic lower respiratory diseases (J40-J47)	370	4.1	16.7
8	Cerebrovascular diseases (I60-I69)	266	2.9	12.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	207	2.3	9.4
10	Influenza and pneumonia (J09-J18)	144	1.6	6.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	1.6	6.5
12	Septicemia (A40-A41)	114	1.3	5.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	0.8	3.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	0.8	3.3
15	Certain conditions originating in the perinatal period (P00-P96)	72	0.8	3.3
...	All other causes (Residual)	1,619	17.9	73.2
American Indian, male, 1-4 years				
...	All causes	54	100.0	34.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	28	51.9	17.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	11.1	*
3	Malignant neoplasms (C00-C97)	4	7.4	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	5.6	*
5	Cerebrovascular diseases (I60-I69)	1	1.9	*
...	All other causes (Residual)	12	22.2	*
American Indian, male, 5-14 years				
...	All causes	65	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	56.9	9.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	12.3	*
3	Malignant neoplasms (C00-C97)	4	6.2	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	6.2	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.1	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.5	*
6	Influenza and pneumonia (J09-J18)	1	1.5	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.5	*
...	All other causes (Residual)	7	10.8	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 15-24 years				
...	All causes	427	100.0	107.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	43.8	47.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	27.2	29.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	11.5	12.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	2.3	*
5	Malignant neoplasms (C00-C97)	9	2.1	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	5	1.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
7	Legal intervention (Y35,Y89.0)	3	0.7	*
10	Diabetes mellitus (E10-E14)	2	0.5	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
13	Septicemia (A40-A41)	1	0.2	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
13	Cerebrovascular diseases (I60-I69)	1	0.2	*
13	Influenza and pneumonia (J09-J18)	1	0.2	*
13	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	31	7.3	7.8
American Indian, male, 25-34 years				
...	All causes	613	100.0	175.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	39.6	69.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	90	14.7	25.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	11.1	19.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	7.3	12.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	37	6.0	10.6
6	Malignant neoplasms (C00-C97)	18	2.9	*
7	Influenza and pneumonia (J09-J18)	8	1.3	*
8	Diabetes mellitus (E10-E14)	6	1.0	*
9	Septicemia (A40-A41)	4	0.7	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.7	*
9	Legal intervention (Y35,Y89.0)	4	0.7	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
12	Pneumonitis due to solids and liquids (J69)	3	0.5	*
15	Meningitis (G00,G03)	2	0.3	*
15	Cerebrovascular diseases (I60-I69)	2	0.3	*
...	All other causes (Residual)	73	11.9	20.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 35-44 years				
...	All causes	735	100.0	248.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	29.5	73.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	101	13.7	34.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	12.0	29.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	67	9.1	22.7
5	Malignant neoplasms (C00-C97)	42	5.7	14.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	4.2	10.5
7	Diabetes mellitus (E10-E14)	28	3.8	9.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.2	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.1	*
10	Septicemia (A40-A41)	7	1.0	*
11	Cerebrovascular diseases (I60-I69)	6	0.8	*
12	Influenza and pneumonia (J09-J18)	5	0.7	*
13	Viral hepatitis (B15-B19)	4	0.5	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.4	*
14	Legal intervention (Y35,Y89.0)	3	0.4	*
...	All other causes (Residual)	116	15.8	39.2
American Indian, male, 45-54 years				
...	All causes	1,352	100.0	500.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	248	18.3	91.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	236	17.5	87.4
3	Malignant neoplasms (C00-C97)	178	13.2	65.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	159	11.8	58.9
5	Diabetes mellitus (E10-E14)	77	5.7	28.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	57	4.2	21.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	2.4	11.9
8	Cerebrovascular diseases (I60-I69)	29	2.1	10.7
9	Viral hepatitis (B15-B19)	22	1.6	8.1
10	Chronic lower respiratory diseases (J40-J47)	18	1.3	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.3	*
12	Septicemia (A40-A41)	16	1.2	*
13	Influenza and pneumonia (J09-J18)	14	1.0	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.9	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10	0.7	*
...	All other causes (Residual)	227	16.8	84.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 55-64 years				
...	All causes	1,746	100.0	939.2
1	Malignant neoplasms (C00-C97)	402	23.0	216.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	369	21.1	198.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	166	9.5	89.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	137	7.8	73.7
5	Diabetes mellitus (E10-E14)	133	7.6	71.5
6	Chronic lower respiratory diseases (J40-J47)	77	4.4	41.4
7	Cerebrovascular diseases (I60-I69)	50	2.9	26.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	1.7	16.1
9	Influenza and pneumonia (J09-J18)	29	1.7	15.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	1.6	15.1
11	Viral hepatitis (B15-B19)	27	1.5	14.5
12	Septicemia (A40-A41)	24	1.4	12.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.1	10.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.7	*
15	Pneumonitis due to solids and liquids (J69)	8	0.5	*
...	All other causes (Residual)	234	13.4	125.9
American Indian, male, 65-74 years				
...	All causes	1,731	100.0	1,948.9
1	Malignant neoplasms (C00-C97)	496	28.7	558.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	401	23.2	451.5
3	Diabetes mellitus (E10-E14)	137	7.9	154.2
4	Chronic lower respiratory diseases (J40-J47)	99	5.7	111.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	4.6	90.1
6	Cerebrovascular diseases (I60-I69)	75	4.3	84.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	51	2.9	57.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	1.9	37.2
9	Septicemia (A40-A41)	29	1.7	32.7
10	Influenza and pneumonia (J09-J18)	22	1.3	24.8
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.8	*
13	Pneumonitis due to solids and liquids (J69)	11	0.6	*
14	Alzheimer's disease (G30)	9	0.5	*
15	Viral hepatitis (B15-B19)	8	0.5	*
...	All other causes (Residual)	253	14.6	284.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, male, 75-84 years				
...	All causes	1,387	100.0	4,190.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	377	27.2	1,138.9
2	Malignant neoplasms (C00-C97)	293	21.1	885.2
3	Chronic lower respiratory diseases (J40-J47)	116	8.4	350.4
4	Diabetes mellitus (E10-E14)	97	7.0	293.0
5	Cerebrovascular diseases (I60-I69)	57	4.1	172.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	2.6	108.8
7	Influenza and pneumonia (J09-J18)	35	2.5	105.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	2.4	99.7
9	Alzheimer's disease (G30)	25	1.8	75.5
10	Septicemia (A40-A41)	24	1.7	72.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	23	1.7	69.5
12	Parkinson's disease (G20-G21)	15	1.1	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.8	*
14	Pneumonitis due to solids and liquids (J69)	9	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	*
...	All other causes (Residual)	231	16.7	697.9
American Indian, male, 85 years and over				
...	All causes	749	100.0	8,618.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	198	26.4	2,278.2
2	Malignant neoplasms (C00-C97)	112	15.0	1,288.7
3	Chronic lower respiratory diseases (J40-J47)	55	7.3	632.8
4	Cerebrovascular diseases (I60-I69)	44	5.9	506.3
5	Diabetes mellitus (E10-E14)	34	4.5	391.2
6	Influenza and pneumonia (J09-J18)	26	3.5	299.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.8	241.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	2.8	241.6
9	Alzheimer's disease (G30)	19	2.5	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	2.0	*
11	Parkinson's disease (G20-G21)	13	1.7	*
12	Pneumonitis due to solids and liquids (J69)	12	1.6	*
13	Septicemia (A40-A41)	9	1.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.5	*
...	All other causes (Residual)	161	21.5	1,852.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, all ages				
...	All causes	7,460	100.0	340.9
1	Malignant neoplasms (C00-C97)	1,460	19.6	66.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,219	16.3	55.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	644	8.6	29.4
4	Diabetes mellitus (E10-E14)	471	6.3	21.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	401	5.4	18.3
6	Chronic lower respiratory diseases (J40-J47)	338	4.5	15.4
7	Cerebrovascular diseases (I60-I69)	314	4.2	14.4
8	Alzheimer's disease (G30)	162	2.2	7.4
8	Influenza and pneumonia (J09-J18)	162	2.2	7.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	2.0	6.9
11	Septicemia (A40-A41)	141	1.9	6.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	1.3	4.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	1.0	3.5
14	Certain conditions originating in the perinatal period (P00-P96)	57	0.8	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	0.8	2.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	0.8	2.6
...	All other causes (Residual)	1,652	22.1	75.5
American Indian, female, 1-4 years				
...	All causes	35	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	31.4	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	14.3	*
3	Septicemia (A40-A41)	3	8.6	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	8.6	*
5	Malignant neoplasms (C00-C97)	1	2.9	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.9	*
5	Influenza and pneumonia (J09-J18)	1	2.9	*
5	Hernia (K40-K46)	1	2.9	*
...	All other causes (Residual)	9	25.7	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 5-14 years				
...	All causes	36	100.0	9.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	38.9	*
2	Malignant neoplasms (C00-C97)	2	5.6	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	5.6	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	5.6	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	5.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	2.8	*
6	Anemias (D50-D64)	1	2.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.8	*
6	Cerebrovascular diseases (I60-I69)	1	2.8	*
6	Influenza and pneumonia (J09-J18)	1	2.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.8	*
...	All other causes (Residual)	8	22.2	*
American Indian, female, 15-24 years				
...	All causes	177	100.0	47.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	46.3	22.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	20.3	9.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	8.5	*
4	Malignant neoplasms (C00-C97)	6	3.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.3	*
6	Septicemia (A40-A41)	2	1.1	*
6	Diabetes mellitus (E10-E14)	2	1.1	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
10	Pneumonitis due to solids and liquids (J69)	1	0.6	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
10	Legal intervention (Y35,Y89.0)	1	0.6	*
...	All other causes (Residual)	22	12.4	5.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 25-34 years				
...	All causes	321	100.0	98.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	34.6	34.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	28	8.7	8.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	8.7	8.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	5.3	*
5	Malignant neoplasms (C00-C97)	16	5.0	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	4.4	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	10	3.1	*
8	Septicemia (A40-A41)	7	2.2	*
8	Diabetes mellitus (E10-E14)	7	2.2	*
10	Cerebrovascular diseases (I60-I69)	6	1.9	*
11	Influenza and pneumonia (J09-J18)	3	0.9	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
15	Viral hepatitis (B15-B19)	1	0.3	*
15	Anemias (D50-D64)	1	0.3	*
15	Pneumonitis due to solids and liquids (J69)	1	0.3	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (Residual)	63	19.6	19.4
American Indian, female, 35-44 years				
...	All causes	463	100.0	162.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	113	24.4	39.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	69	14.9	24.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	10.4	16.8
4	Malignant neoplasms (C00-C97)	46	9.9	16.1
5	Diabetes mellitus (E10-E14)	12	2.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.6	*
7	Septicemia (A40-A41)	9	1.9	*
7	Viral hepatitis (B15-B19)	9	1.9	*
9	Cerebrovascular diseases (I60-I69)	8	1.7	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	1.7	*
11	Pregnancy, childbirth and the puerperium (O00-O99)	7	1.5	*
12	Influenza and pneumonia (J09-J18)	5	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.1	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.9	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.9	*
...	All other causes (Residual)	99	21.4	34.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 45-54 years				
...	All causes	912	100.0	329.6
1	Malignant neoplasms (C00-C97)	173	19.0	62.5
2	Chronic liver disease and cirrhosis (K70,K73-K74)	143	15.7	51.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	133	14.6	48.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	109	12.0	39.4
5	Diabetes mellitus (E10-E14)	53	5.8	19.2
6	Chronic lower respiratory diseases (J40-J47)	22	2.4	8.0
7	Septicemia (A40-A41)	21	2.3	7.6
8	Cerebrovascular diseases (I60-I69)	20	2.2	7.2
9	Influenza and pneumonia (J09-J18)	17	1.9	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	1.4	*
11	Viral hepatitis (B15-B19)	12	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	5	0.5	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.4	*
...	All other causes (Residual)	173	19.0	62.5
American Indian, female, 55-64 years				
...	All causes	1,176	100.0	587.2
1	Malignant neoplasms (C00-C97)	353	30.0	176.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	168	14.3	83.9
3	Diabetes mellitus (E10-E14)	100	8.5	49.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	77	6.5	38.4
5	Chronic lower respiratory diseases (J40-J47)	64	5.4	32.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	5.4	32.0
7	Cerebrovascular diseases (I60-I69)	40	3.4	20.0
8	Viral hepatitis (B15-B19)	21	1.8	10.5
8	Influenza and pneumonia (J09-J18)	21	1.8	10.5
10	Septicemia (A40-A41)	19	1.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.4	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.1	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	0.5	*
14	Atherosclerosis (I70)	5	0.4	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.4	*
...	All other causes (Residual)	203	17.3	101.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 65-74 years				
...	All causes	1,396	100.0	1,390.3
1	Malignant neoplasms (C00-C97)	396	28.4	394.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	246	17.6	245.0
3	Diabetes mellitus (E10-E14)	124	8.9	123.5
4	Chronic lower respiratory diseases (J40-J47)	109	7.8	108.6
5	Cerebrovascular diseases (I60-I69)	59	4.2	58.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	57	4.1	56.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.7	37.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	2.4	32.9
9	Influenza and pneumonia (J09-J18)	24	1.7	23.9
10	Septicemia (A40-A41)	22	1.6	21.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	1.2	*
12	Alzheimer's disease (G30)	12	0.9	*
13	Parkinson's disease (G20-G21)	7	0.5	*
13	Pneumonitis due to solids and liquids (J69)	7	0.5	*
15	Viral hepatitis (B15-B19)	6	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	*
...	All other causes (Residual)	233	16.7	232.1
American Indian, female, 75-84 years				
...	All causes	1,474	100.0	3,256.9
1	Malignant neoplasms (C00-C97)	341	23.1	753.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	316	21.4	698.2
3	Diabetes mellitus (E10-E14)	105	7.1	232.0
4	Cerebrovascular diseases (I60-I69)	93	6.3	205.5
5	Chronic lower respiratory diseases (J40-J47)	82	5.6	181.2
6	Alzheimer's disease (G30)	46	3.1	101.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	2.9	95.0
8	Influenza and pneumonia (J09-J18)	37	2.5	81.8
9	Septicemia (A40-A41)	33	2.2	72.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	2.2	70.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	19	1.3	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	1.0	*
13	Parkinson's disease (G20-G21)	12	0.8	*
13	Pneumonitis due to solids and liquids (J69)	12	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
...	All other causes (Residual)	277	18.8	612.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
American Indian, female, 85 years and over				
...	All causes	1,308	100.0	7,987.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	307	23.5	1,874.7
2	Malignant neoplasms (C00-C97)	125	9.6	763.3
3	Alzheimer's disease (G30)	101	7.7	616.8
4	Cerebrovascular diseases (I60-I69)	87	6.7	531.3
5	Diabetes mellitus (E10-E14)	68	5.2	415.2
6	Chronic lower respiratory diseases (J40-J47)	55	4.2	335.9
7	Influenza and pneumonia (J09-J18)	52	4.0	317.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	2.9	232.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	2.9	232.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	2.0	158.8
11	Septicemia (A40-A41)	22	1.7	134.3
12	Pneumonitis due to solids and liquids (J69)	14	1.1	*
13	Parkinson's disease (G20-G21)	10	0.8	*
14	Anemias (D50-D64)	9	0.7	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.5	*
...	All other causes (Residual)	349	26.7	2,131.2
Asian or Pacific Islander, both sexes, all ages				
...	All causes	56,352	100.0	311.5
1	Malignant neoplasms (C00-C97)	15,340	27.2	84.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,266	21.8	67.8
3	Cerebrovascular diseases (I60-I69)	4,108	7.3	22.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,372	4.2	13.1
5	Diabetes mellitus (E10-E14)	2,158	3.8	11.9
6	Influenza and pneumonia (J09-J18)	1,745	3.1	9.6
7	Chronic lower respiratory diseases (J40-J47)	1,624	2.9	9.0
8	Alzheimer's disease (G30)	1,379	2.4	7.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,152	2.0	6.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,054	1.9	5.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	857	1.5	4.7
12	Septicemia (A40-A41)	659	1.2	3.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	523	0.9	2.9
14	Certain conditions originating in the perinatal period (P00-P96)	476	0.8	2.6
15	Parkinson's disease (G20-G21)	472	0.8	2.6
...	All other causes (Residual)	10,167	18.0	56.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	150	100.0	15.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	30	20.0	3.1
2	Malignant neoplasms (C00-C97)	21	14.0	2.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	14.0	2.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	5.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	4.0	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	3.3	*
7	Septicemia (A40-A41)	4	2.7	*
7	Influenza and pneumonia (J09-J18)	4	2.7	*
9	Anemias (D50-D64)	2	1.3	*
9	Cerebrovascular diseases (I60-I69)	2	1.3	*
9	Acute bronchitis and bronchiolitis (J20-J21)	2	1.3	*
12	Tuberculosis (A16-A19)	1	0.7	*
12	Meningitis (G00,G03)	1	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.7	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.7	*
...	All other causes (Residual)	40	26.7	4.1
Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	192	100.0	8.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	27.6	2.2
2	Malignant neoplasms (C00-C97)	46	24.0	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	8.9	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	5.2	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10	5.2	*
6	Influenza and pneumonia (J09-J18)	4	2.1	*
7	Anemias (D50-D64)	3	1.6	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.6	*
9	Septicemia (A40-A41)	2	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.0	*
9	Cerebrovascular diseases (I60-I69)	2	1.0	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	1.0	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.5	*
...	All other causes (Residual)	37	19.3	1.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	772	100.0	30.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	31.3	9.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	195	25.3	7.6
3	Malignant neoplasms (C00-C97)	91	11.8	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	8.4	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	5.2	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
7	Cerebrovascular diseases (I60-I69)	6	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
10	Influenza and pneumonia (J09-J18)	4	0.5	*
10	Legal intervention (Y35,Y89.0)	4	0.5	*
12	Septicemia (A40-A41)	3	0.4	*
12	Diabetes mellitus (E10-E14)	3	0.4	*
14	Tuberculosis (A16-A19)	2	0.3	*
14	Anemias (D50-D64)	2	0.3	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	88	11.4	3.4
Asian or Pacific Islander, both sexes, 25-34 years				
...	All causes	1,182	100.0	39.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	277	23.4	9.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	234	19.8	7.7
3	Malignant neoplasms (C00-C97)	196	16.6	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	8.7	3.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	7.0	2.7
6	Cerebrovascular diseases (I60-I69)	29	2.5	1.0
7	Pregnancy, childbirth and the puerperium (O00-O99)	16	1.4	*
8	Diabetes mellitus (E10-E14)	13	1.1	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.9	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.8	*
11	Septicemia (A40-A41)	8	0.7	*
12	Aortic aneurysm and dissection (I71)	7	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
...	All other causes (Residual)	178	15.1	5.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 35-44 years				
...	All causes	2,043	100.0	70.0
1	Malignant neoplasms (C00-C97)	625	30.6	21.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	306	15.0	10.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	249	12.2	8.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	222	10.9	7.6
5	Cerebrovascular diseases (I60-I69)	106	5.2	3.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	2.9	2.1
7	Diabetes mellitus (E10-E14)	43	2.1	1.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	38	1.9	1.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	22	1.1	0.8
10	Influenza and pneumonia (J09-J18)	19	0.9	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.9	*
12	Viral hepatitis (B15-B19)	16	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.8	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	15	0.7	*
...	All other causes (Residual)	272	13.3	9.3
Asian or Pacific Islander, both sexes, 45-54 years				
...	All causes	4,105	100.0	170.7
1	Malignant neoplasms (C00-C97)	1,653	40.3	68.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	718	17.5	29.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	6.6	11.3
4	Cerebrovascular diseases (I60-I69)	266	6.5	11.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	4.6	7.8
6	Diabetes mellitus (E10-E14)	143	3.5	5.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	90	2.2	3.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.2	2.1
9	Viral hepatitis (B15-B19)	44	1.1	1.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	1.0	1.8
11	Septicemia (A40-A41)	36	0.9	1.5
12	Influenza and pneumonia (J09-J18)	32	0.8	1.3
13	Chronic lower respiratory diseases (J40-J47)	30	0.7	1.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.7	1.2
15	Aortic aneurysm and dissection (I71)	27	0.7	1.1
...	All other causes (Residual)	485	11.8	20.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 55-64 years				
...	All causes	7,369	100.0	394.0
1	Malignant neoplasms (C00-C97)	3,115	42.3	166.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,422	19.3	76.0
3	Cerebrovascular diseases (I60-I69)	459	6.2	24.5
4	Diabetes mellitus (E10-E14)	320	4.3	17.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	250	3.4	13.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	2.1	8.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	123	1.7	6.6
8	Chronic lower respiratory diseases (J40-J47)	114	1.5	6.1
9	Influenza and pneumonia (J09-J18)	99	1.3	5.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	91	1.2	4.9
11	Viral hepatitis (B15-B19)	90	1.2	4.8
12	Septicemia (A40-A41)	81	1.1	4.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.0	3.9
14	Aortic aneurysm and dissection (I71)	52	0.7	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	0.7	2.7
...	All other causes (Residual)	875	11.9	46.8
Asian or Pacific Islander, both sexes, 65-74 years				
...	All causes	9,362	100.0	902.4
1	Malignant neoplasms (C00-C97)	3,510	37.5	338.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,917	20.5	184.8
3	Cerebrovascular diseases (I60-I69)	686	7.3	66.1
4	Diabetes mellitus (E10-E14)	477	5.1	46.0
5	Chronic lower respiratory diseases (J40-J47)	253	2.7	24.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	2.6	23.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	2.2	20.0
8	Influenza and pneumonia (J09-J18)	200	2.1	19.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	127	1.4	12.2
10	Septicemia (A40-A41)	119	1.3	11.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	102	1.1	9.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	0.9	8.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.8	7.4
14	Viral hepatitis (B15-B19)	76	0.8	7.3
15	Alzheimer's disease (G30)	61	0.7	5.9
...	All other causes (Residual)	1,221	13.0	117.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, both sexes, 75-84 years				
...	All causes	13,832	100.0	2,702.2
1	Malignant neoplasms (C00-C97)	3,871	28.0	756.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,108	22.5	607.2
3	Cerebrovascular diseases (I60-I69)	1,148	8.3	224.3
4	Diabetes mellitus (E10-E14)	655	4.7	128.0
5	Chronic lower respiratory diseases (J40-J47)	549	4.0	107.3
6	Influenza and pneumonia (J09-J18)	463	3.3	90.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	359	2.6	70.1
8	Alzheimer's disease (G30)	345	2.5	67.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	320	2.3	62.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	219	1.6	42.8
11	Parkinson's disease (G20-G21)	197	1.4	38.5
12	Septicemia (A40-A41)	180	1.3	35.2
13	Pneumonitis due to solids and liquids (J69)	110	0.8	21.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	102	0.7	19.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	92	0.7	18.0
...	All other causes (Residual)	2,114	15.3	413.0
Asian or Pacific Islander, both sexes, 85 years and over				
...	All causes	16,432	100.0	9,070.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,619	28.1	2,549.6
2	Malignant neoplasms (C00-C97)	2,208	13.4	1,218.8
3	Cerebrovascular diseases (I60-I69)	1,399	8.5	772.2
4	Alzheimer's disease (G30)	959	5.8	529.4
5	Influenza and pneumonia (J09-J18)	907	5.5	500.6
6	Chronic lower respiratory diseases (J40-J47)	649	3.9	358.2
7	Diabetes mellitus (E10-E14)	504	3.1	278.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	424	2.6	234.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	362	2.2	199.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	357	2.2	197.1
11	Parkinson's disease (G20-G21)	210	1.3	115.9
12	Septicemia (A40-A41)	206	1.3	113.7
13	Pneumonitis due to solids and liquids (J69)	203	1.2	112.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	93	0.6	51.3
15	Aortic aneurysm and dissection (I71)	71	0.4	39.2
...	All other causes (Residual)	3,261	19.8	1,800.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, all ages				
...	All causes	28,819	100.0	332.8
1	Malignant neoplasms (C00-C97)	7,709	26.7	89.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,650	23.1	76.8
3	Cerebrovascular diseases (I60-I69)	1,876	6.5	21.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,472	5.1	17.0
5	Diabetes mellitus (E10-E14)	1,093	3.8	12.6
6	Chronic lower respiratory diseases (J40-J47)	966	3.4	11.2
7	Influenza and pneumonia (J09-J18)	890	3.1	10.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	805	2.8	9.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	501	1.7	5.8
10	Alzheimer's disease (G30)	439	1.5	5.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	377	1.3	4.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	326	1.1	3.8
13	Septicemia (A40-A41)	312	1.1	3.6
14	Certain conditions originating in the perinatal period (P00-P96)	262	0.9	3.0
15	Parkinson's disease (G20-G21)	252	0.9	2.9
...	All other causes (Residual)	4,889	17.0	56.5
Asian or Pacific Islander, male, 1-4 years				
...	All causes	80	100.0	16.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	22.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	16.3	*
3	Malignant neoplasms (C00-C97)	10	12.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	6.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	3.8	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.5	*
7	Tuberculosis (A16-A19)	1	1.3	*
7	Septicemia (A40-A41)	1	1.3	*
7	Anemias (D50-D64)	1	1.3	*
7	Cerebrovascular diseases (I60-I69)	1	1.3	*
7	Influenza and pneumonia (J09-J18)	1	1.3	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	1.3	*
...	All other causes (Residual)	23	28.8	4.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 5-14 years				
...	All causes	108	100.0	9.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	28.7	2.6
2	Malignant neoplasms (C00-C97)	27	25.0	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	7.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	6.5	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	6.5	*
6	Anemias (D50-D64)	3	2.8	*
7	Influenza and pneumonia (J09-J18)	2	1.9	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
9	Cerebrovascular diseases (I60-I69)	1	0.9	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	0.9	*
...	All other causes (Residual)	18	16.7	*
Asian or Pacific Islander, male, 15-24 years				
...	All causes	540	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	35.2	14.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	126	23.3	9.7
3	Malignant neoplasms (C00-C97)	62	11.5	4.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	9.3	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	5.7	2.4
6	Cerebrovascular diseases (I60-I69)	6	1.1	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.1	*
8	Legal intervention (Y35,Y89.0)	4	0.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
11	Septicemia (A40-A41)	2	0.4	*
12	Anemias (D50-D64)	1	0.2	*
12	Diabetes mellitus (E10-E14)	1	0.2	*
12	Aortic aneurysm and dissection (I71)	1	0.2	*
12	Influenza and pneumonia (J09-J18)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	52	9.6	4.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 25-34 years				
...	All causes	796	100.0	54.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	27.0	14.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	174	21.9	12.0
3	Malignant neoplasms (C00-C97)	107	13.4	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	9.7	5.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	8.3	4.5
6	Cerebrovascular diseases (I60-I69)	16	2.0	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	1.0	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
8	Aortic aneurysm and dissection (I71)	6	0.8	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.8	*
11	Septicemia (A40-A41)	5	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
12	Influenza and pneumonia (J09-J18)	4	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.5	*
...	All other causes (Residual)	94	11.8	6.5
Asian or Pacific Islander, male, 35-44 years				
...	All causes	1,266	100.0	92.0
1	Malignant neoplasms (C00-C97)	267	21.1	19.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	252	19.9	18.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	14.6	13.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	12.6	11.6
5	Cerebrovascular diseases (I60-I69)	55	4.3	4.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	3.1	2.8
7	Diabetes mellitus (E10-E14)	36	2.8	2.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	33	2.6	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	17	1.3	*
10	Viral hepatitis (B15-B19)	14	1.1	*
11	Influenza and pneumonia (J09-J18)	13	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.0	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.9	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.8	*
15	Aortic aneurysm and dissection (I71)	8	0.6	*
...	All other causes (Residual)	154	12.2	11.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 45-54 years				
...	All causes	2,443	100.0	216.5
1	Malignant neoplasms (C00-C97)	744	30.5	65.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	584	23.9	51.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	7.4	16.1
4	Cerebrovascular diseases (I60-I69)	149	6.1	13.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	127	5.2	11.3
6	Diabetes mellitus (E10-E14)	99	4.1	8.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	64	2.6	5.7
8	Viral hepatitis (B15-B19)	30	1.2	2.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	1.2	2.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.9	2.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.9	1.9
12	Aortic aneurysm and dissection (I71)	21	0.9	1.9
13	Septicemia (A40-A41)	20	0.8	1.8
13	Chronic lower respiratory diseases (J40-J47)	20	0.8	1.8
15	Influenza and pneumonia (J09-J18)	17	0.7	*
...	All other causes (Residual)	312	12.8	27.7
Asian or Pacific Islander, male, 55-64 years				
...	All causes	4,284	100.0	505.9
1	Malignant neoplasms (C00-C97)	1,581	36.9	186.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,003	23.4	118.4
3	Cerebrovascular diseases (I60-I69)	254	5.9	30.0
4	Diabetes mellitus (E10-E14)	197	4.6	23.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	3.5	17.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	2.7	13.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	90	2.1	10.6
8	Chronic lower respiratory diseases (J40-J47)	67	1.6	7.9
9	Influenza and pneumonia (J09-J18)	57	1.3	6.7
10	Viral hepatitis (B15-B19)	55	1.3	6.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.2	5.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	47	1.1	5.6
13	Septicemia (A40-A41)	46	1.1	5.4
14	Aortic aneurysm and dissection (I71)	36	0.8	4.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	0.7	3.3
...	All other causes (Residual)	509	11.9	60.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 65-74 years				
...	All causes	5,200	100.0	1,109.2
1	Malignant neoplasms (C00-C97)	1,859	35.8	396.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,196	23.0	255.1
3	Cerebrovascular diseases (I60-I69)	366	7.0	78.1
4	Diabetes mellitus (E10-E14)	242	4.7	51.6
5	Chronic lower respiratory diseases (J40-J47)	156	3.0	33.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	141	2.7	30.1
7	Influenza and pneumonia (J09-J18)	123	2.4	26.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	2.1	23.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	77	1.5	16.4
10	Septicemia (A40-A41)	59	1.1	12.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	1.1	12.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	1.0	10.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.7	8.1
14	Viral hepatitis (B15-B19)	37	0.7	7.9
15	Aortic aneurysm and dissection (I71)	36	0.7	7.7
...	All other causes (Residual)	652	12.5	139.1
Asian or Pacific Islander, male, 75-84 years				
...	All causes	6,898	100.0	3,218.0
1	Malignant neoplasms (C00-C97)	2,007	29.1	936.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,605	23.3	748.7
3	Cerebrovascular diseases (I60-I69)	513	7.4	239.3
4	Chronic lower respiratory diseases (J40-J47)	345	5.0	160.9
5	Diabetes mellitus (E10-E14)	324	4.7	151.1
6	Influenza and pneumonia (J09-J18)	258	3.7	120.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	2.4	78.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	150	2.2	70.0
9	Parkinson's disease (G20-G21)	118	1.7	55.0
10	Alzheimer's disease (G30)	112	1.6	52.2
11	Septicemia (A40-A41)	89	1.3	41.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	87	1.3	40.6
13	Pneumonitis due to solids and liquids (J69)	65	0.9	30.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	47	0.7	21.9
14	Aortic aneurysm and dissection (I71)	47	0.7	21.9
...	All other causes (Residual)	962	13.9	448.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, male, 85 years and over				
...	All causes	6,688	100.0	10,116.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,879	28.1	2,842.2
2	Malignant neoplasms (C00-C97)	1,043	15.6	1,577.7
3	Cerebrovascular diseases (I60-I69)	513	7.7	776.0
4	Influenza and pneumonia (J09-J18)	407	6.1	615.6
5	Chronic lower respiratory diseases (J40-J47)	364	5.4	550.6
6	Alzheimer's disease (G30)	287	4.3	434.1
7	Diabetes mellitus (E10-E14)	191	2.9	288.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	2.5	257.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	156	2.3	236.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	149	2.2	225.4
11	Parkinson's disease (G20-G21)	106	1.6	160.3
12	Pneumonitis due to solids and liquids (J69)	105	1.6	158.8
13	Septicemia (A40-A41)	80	1.2	121.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.6	62.0
15	Aortic aneurysm and dissection (I71)	29	0.4	43.9
...	All other causes (Residual)	1,168	17.5	1,766.8
Asian or Pacific Islander, female, all ages				
...	All causes	27,533	100.0	292.0
1	Malignant neoplasms (C00-C97)	7,631	27.7	80.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,616	20.4	59.6
3	Cerebrovascular diseases (I60-I69)	2,232	8.1	23.7
4	Diabetes mellitus (E10-E14)	1,065	3.9	11.3
5	Alzheimer's disease (G30)	940	3.4	10.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	900	3.3	9.5
7	Influenza and pneumonia (J09-J18)	855	3.1	9.1
8	Chronic lower respiratory diseases (J40-J47)	658	2.4	7.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	553	2.0	5.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	480	1.7	5.1
11	Septicemia (A40-A41)	347	1.3	3.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	347	1.3	3.7
13	Parkinson's disease (G20-G21)	220	0.8	2.3
14	Certain conditions originating in the perinatal period (P00-P96)	214	0.8	2.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	197	0.7	2.1
...	All other causes (Residual)	5,278	19.2	56.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 1-4 years				
...	All causes	70	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	17.1	*
2	Malignant neoplasms (C00-C97)	11	15.7	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	11.4	*
4	Septicemia (A40-A41)	3	4.3	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	4.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	4.3	*
4	Influenza and pneumonia (J09-J18)	3	4.3	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	4.3	*
9	Anemias (D50-D64)	1	1.4	*
9	Meningitis (G00,G03)	1	1.4	*
9	Cerebrovascular diseases (I60-I69)	1	1.4	*
9	Acute bronchitis and bronchiolitis (J20-J21)	1	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
9	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.4	*
...	All other causes (Residual)	17	24.3	*
Asian or Pacific Islander, female, 5-14 years				
...	All causes	84	100.0	7.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	26.2	1.9
2	Malignant neoplasms (C00-C97)	19	22.6	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	10.7	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.6	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	3.6	*
6	Septicemia (A40-A41)	2	2.4	*
6	Influenza and pneumonia (J09-J18)	2	2.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.2	*
8	Cerebrovascular diseases (I60-I69)	1	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	1	1.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.2	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.2	*
...	All other causes (Residual)	19	22.6	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 15-24 years				
...	All causes	232	100.0	18.4
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	29.7	5.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	22.4	4.1
3	Malignant neoplasms (C00-C97)	29	12.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	6.5	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.9	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	2.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.3	*
7	Influenza and pneumonia (J09-J18)	3	1.3	*
9	Tuberculosis (A16-A19)	2	0.9	*
9	Diabetes mellitus (E10-E14)	2	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.9	*
12	Septicemia (A40-A41)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
12	Aortic aneurysm and dissection (I71)	1	0.4	*
12	Pneumonitis due to solids and liquids (J69)	1	0.4	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	34	14.7	2.7
Asian or Pacific Islander, female, 25-34 years				
...	All causes	386	100.0	24.4
1	Malignant neoplasms (C00-C97)	89	23.1	5.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	62	16.1	3.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	15.5	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	6.7	1.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	4.4	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	16	4.1	*
7	Cerebrovascular diseases (I60-I69)	13	3.4	*
8	Diabetes mellitus (E10-E14)	10	2.6	*
9	Septicemia (A40-A41)	3	0.8	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.8	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.8	*
12	Tuberculosis (A16-A19)	2	0.5	*
12	Meningococcal infection (A39)	2	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	*
12	Anemias (D50-D64)	2	0.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
12	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.5	*
...	All other causes (Residual)	70	18.1	4.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 35-44 years				
...	All causes	777	100.0	50.4
1	Malignant neoplasms (C00-C97)	358	46.1	23.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	8.2	4.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	63	8.1	4.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	6.9	3.5
5	Cerebrovascular diseases (I60-I69)	51	6.6	3.3
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	2.7	1.4
7	Pregnancy, childbirth and the puerperium (O00-O99)	15	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.3	*
9	Diabetes mellitus (E10-E14)	7	0.9	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
11	Influenza and pneumonia (J09-J18)	6	0.8	*
13	Septicemia (A40-A41)	5	0.6	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
13	Aortic aneurysm and dissection (I71)	5	0.6	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	5	0.6	*
...	All other causes (Residual)	95	12.2	6.2
Asian or Pacific Islander, female, 45-54 years				
...	All causes	1,662	100.0	130.3
1	Malignant neoplasms (C00-C97)	909	54.7	71.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	134	8.1	10.5
3	Cerebrovascular diseases (I60-I69)	117	7.0	9.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	90	5.4	7.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	60	3.6	4.7
6	Diabetes mellitus (E10-E14)	44	2.6	3.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.6	2.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	26	1.6	2.0
9	Septicemia (A40-A41)	16	1.0	*
10	Influenza and pneumonia (J09-J18)	15	0.9	*
11	Viral hepatitis (B15-B19)	14	0.8	*
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.7	*
14	Chronic lower respiratory diseases (J40-J47)	10	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.4	*
...	All other causes (Residual)	160	9.6	12.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 55-64 years				
...	All causes	3,085	100.0	301.5
1	Malignant neoplasms (C00-C97)	1,534	49.7	149.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	419	13.6	40.9
3	Cerebrovascular diseases (I60-I69)	205	6.6	20.0
4	Diabetes mellitus (E10-E14)	123	4.0	12.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	101	3.3	9.9
6	Chronic lower respiratory diseases (J40-J47)	47	1.5	4.6
7	Influenza and pneumonia (J09-J18)	42	1.4	4.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.3	4.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	1.3	3.9
10	Septicemia (A40-A41)	35	1.1	3.4
10	Viral hepatitis (B15-B19)	35	1.1	3.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	33	1.1	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25	0.8	2.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	0.7	2.2
15	Aortic aneurysm and dissection (I71)	16	0.5	*
...	All other causes (Residual)	366	11.9	35.8
Asian or Pacific Islander, female, 65-74 years				
...	All causes	4,162	100.0	731.9
1	Malignant neoplasms (C00-C97)	1,651	39.7	290.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	721	17.3	126.8
3	Cerebrovascular diseases (I60-I69)	320	7.7	56.3
4	Diabetes mellitus (E10-E14)	235	5.6	41.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	102	2.5	17.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	98	2.4	17.2
7	Chronic lower respiratory diseases (J40-J47)	97	2.3	17.1
8	Influenza and pneumonia (J09-J18)	77	1.9	13.5
9	Septicemia (A40-A41)	60	1.4	10.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	50	1.2	8.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.1	7.7
12	Viral hepatitis (B15-B19)	39	0.9	6.9
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.9	6.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	0.8	6.2
15	Parkinson's disease (G20-G21)	32	0.8	5.6
...	All other causes (Residual)	562	13.5	98.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and sex:
United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Data for races other than white and black should be interpreted with caution because of inconsistencies between the reporting of race on death certificates and on censuses and surveys]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race, sex, and age	Number	Percent of total deaths	Rate
Asian or Pacific Islander, female, 75-84 years				
...	All causes	6,934	100.0	2,330.6
1	Malignant neoplasms (C00-C97)	1,864	26.9	626.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,503	21.7	505.2
3	Cerebrovascular diseases (I60-I69)	635	9.2	213.4
4	Diabetes mellitus (E10-E14)	331	4.8	111.3
5	Alzheimer's disease (G30)	233	3.4	78.3
6	Influenza and pneumonia (J09-J18)	205	3.0	68.9
7	Chronic lower respiratory diseases (J40-J47)	204	2.9	68.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	2.7	63.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	170	2.5	57.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	132	1.9	44.4
11	Septicemia (A40-A41)	91	1.3	30.6
12	Parkinson's disease (G20-G21)	79	1.1	26.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	60	0.9	20.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.6	15.1
14	Pneumonitis due to solids and liquids (J69)	45	0.6	15.1
...	All other causes (Residual)	1,147	16.5	385.5
Asian or Pacific Islander, female, 85 years and over				
...	All causes	9,744	100.0	8,469.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,740	28.1	2,381.5
2	Malignant neoplasms (C00-C97)	1,165	12.0	1,012.6
3	Cerebrovascular diseases (I60-I69)	886	9.1	770.1
4	Alzheimer's disease (G30)	672	6.9	584.1
5	Influenza and pneumonia (J09-J18)	500	5.1	434.6
6	Diabetes mellitus (E10-E14)	313	3.2	272.0
7	Chronic lower respiratory diseases (J40-J47)	285	2.9	247.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	268	2.8	232.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	2.1	180.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	2.0	166.9
11	Septicemia (A40-A41)	126	1.3	109.5
12	Parkinson's disease (G20-G21)	104	1.1	90.4
13	Pneumonitis due to solids and liquids (J69)	98	1.0	85.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	45.2
15	Aortic aneurysm and dissection (I71)	42	0.4	36.5
...	All other causes (Residual)	2,093	21.5	1,819.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2012