

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	133,004	100.0	300.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,921	21.7	65.3
2	Malignant neoplasms (C00-C97)	26,633	20.0	60.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,052	9.1	27.2
4	Cerebrovascular diseases (I60-I69)	7,005	5.3	15.8
5	Diabetes mellitus (E10-E14)	6,287	4.7	14.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,592	2.7	8.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,524	2.6	8.0
8	Chronic lower respiratory diseases (J40-J47)	3,310	2.5	7.5
9	Influenza and pneumonia (J10-J18)	2,966	2.2	6.7
10	Certain conditions originating in the perinatal period (P00-P96)	2,804	2.1	6.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,592	1.9	5.8
12	Alzheimer's disease (G30)	2,399	1.8	5.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,177	1.6	4.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,952	1.5	4.4
15	Septicemia (A40-A41)	1,829	1.4	4.1
...	All other causes (Residual)	24,961	18.8	56.3
Hispanic, both sexes, 1-4 years				
...	All causes	987	100.0	26.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	361	36.6	9.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	12.7	3.3
3	Malignant neoplasms (C00-C97)	82	8.3	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	6.3	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	3.1	0.8
6	Influenza and pneumonia (J10-J18)	28	2.8	0.7
7	Septicemia (A40-A41)	18	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.1	*
10	Cerebrovascular diseases (I60-I69)	9	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
12	Meningitis (G00,G03)	6	0.6	*
13	Anemias (D50-D64)	4	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	223	22.6	6.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	484	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	36.6	4.3
2	Malignant neoplasms (C00-C97)	90	18.6	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	6.0	0.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	5.8	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.7	*
6	Influenza and pneumonia (J10-J18)	12	2.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	2.3	*
8	Septicemia (A40-A41)	9	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.9	*
10	Cerebrovascular diseases (I60-I69)	7	1.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	1.2	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
13	Tuberculosis (A16-A19)	1	0.2	*
13	Meningococcal infection (A39)	1	0.2	*
13	Anemias (D50-D64)	1	0.2	*
13	Nutritional deficiencies (E40-E64)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	79	16.3	1.9
Hispanic, both sexes, 10-14 years				
...	All causes	653	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	36.0	6.0
2	Malignant neoplasms (C00-C97)	93	14.2	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	10.1	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	5.1	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.1	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	3.7	0.6
7	Cerebrovascular diseases (I60-I69)	10	1.5	*
8	Septicemia (A40-A41)	7	1.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
8	Chronic lower respiratory diseases (J40-J47)	7	1.1	*
11	Diabetes mellitus (E10-E14)	4	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
13	Influenza and pneumonia (J10-J18)	3	0.5	*
13	Pneumonitis due to solids and liquids (J69)	3	0.5	*
13	Diseases of appendix (K35-K38)	3	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	124	19.0	3.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	2,366	100.0	65.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,058	44.7	29.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	574	24.3	15.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	198	8.4	5.5
4	Malignant neoplasms (C00-C97)	122	5.2	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	2.0	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	1.7	1.1
7	Cerebrovascular diseases (I60-I69)	20	0.8	0.6
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.4	*
10	Influenza and pneumonia (J10-J18)	8	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.3	*
10	Pregnancy, childbirth and the puerperium (O00-099)	8	0.3	*
13	Septicemia (A40-A41)	7	0.3	*
14	Legal intervention (Y35,Y89.0)	6	0.3	*
15	Meningitis (G00,G03)	5	0.2	*
...	All other causes (Residual)	243	10.3	6.7
Hispanic, both sexes, 20-24 years				
...	All causes	3,566	100.0	95.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,604	45.0	42.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	769	21.6	20.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	345	9.7	9.2
4	Malignant neoplasms (C00-C97)	199	5.6	5.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	2.3	2.2
6	Cerebrovascular diseases (I60-I69)	32	0.9	0.9
7	Pregnancy, childbirth and the puerperium (O00-099)	31	0.9	0.8
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	0.8	0.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.6	0.6
10	Legal intervention (Y35,Y89.0)	19	0.5	*
11	Influenza and pneumonia (J10-J18)	16	0.4	*
12	Septicemia (A40-A41)	15	0.4	*
13	Diabetes mellitus (E10-E14)	14	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.2	*
...	All other causes (Residual)	369	10.3	9.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	3,305	100.0	79.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,285	38.9	30.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	605	18.3	14.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	315	9.5	7.6
4	Malignant neoplasms (C00-C97)	243	7.4	5.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	143	4.3	3.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	62	1.9	1.5
7	Cerebrovascular diseases (I60-I69)	39	1.2	0.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	1.1	0.8
9	Pregnancy, childbirth and the puerperium (O00-099)	34	1.0	0.8
10	Diabetes mellitus (E10-E14)	31	0.9	0.7
11	Influenza and pneumonia (J10-J18)	23	0.7	0.6
12	Septicemia (A40-A41)	21	0.6	0.5
13	Legal intervention (Y35,Y89.0)	19	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	16	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.4	*
...	All other causes (Residual)	422	12.8	10.1
Hispanic, both sexes, 30-34 years				
...	All causes	3,287	100.0	83.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,115	33.9	28.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	390	11.9	9.9
3	Malignant neoplasms (C00-C97)	336	10.2	8.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	259	7.9	6.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	231	7.0	5.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	133	4.0	3.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	74	2.3	1.9
8	Cerebrovascular diseases (I60-I69)	55	1.7	1.4
9	Diabetes mellitus (E10-E14)	43	1.3	1.1
10	Pregnancy, childbirth and the puerperium (O00-099)	36	1.1	0.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.9	0.7
12	Influenza and pneumonia (J10-J18)	26	0.8	0.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.7	0.6
14	Septicemia (A40-A41)	20	0.6	0.5
14	Legal intervention (Y35,Y89.0)	20	0.6	0.5
...	All other causes (Residual)	499	15.2	12.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,822	100.0	108.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,021	26.7	28.9
2	Malignant neoplasms (C00-C97)	546	14.3	15.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	332	8.7	9.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	325	8.5	9.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	253	6.6	7.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	226	5.9	6.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	149	3.9	4.2
8	Cerebrovascular diseases (I60-I69)	100	2.6	2.8
9	Diabetes mellitus (E10-E14)	66	1.7	1.9
10	Septicemia (A40-A41)	35	0.9	1.0
11	Influenza and pneumonia (J10-J18)	29	0.8	0.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	0.6	0.7
13	Viral hepatitis (B15-B19)	23	0.6	0.7
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.6	0.7
15	Pregnancy, childbirth and the puerperium (O00-099)	22	0.6	0.6
...	All other causes (Residual)	648	17.0	18.3
Hispanic, both sexes, 40-44 years				
...	All causes	5,129	100.0	164.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,081	21.1	34.6
2	Malignant neoplasms (C00-C97)	894	17.4	28.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	598	11.7	19.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	331	6.5	10.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	295	5.8	9.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	215	4.2	6.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	212	4.1	6.8
8	Cerebrovascular diseases (I60-I69)	208	4.1	6.7
9	Diabetes mellitus (E10-E14)	119	2.3	3.8
10	Viral hepatitis (B15-B19)	81	1.6	2.6
11	Influenza and pneumonia (J10-J18)	48	0.9	1.5
12	Septicemia (A40-A41)	44	0.9	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	43	0.8	1.4
14	Chronic lower respiratory diseases (J40-J47)	29	0.6	0.9
15	Aortic aneurysm and dissection (I71)	24	0.5	0.8
...	All other causes (Residual)	907	17.7	29.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	6,486	100.0	253.6
1	Malignant neoplasms (C00-C97)	1,388	21.4	54.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	982	15.1	38.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	926	14.3	36.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	495	7.6	19.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	302	4.7	11.8
6	Cerebrovascular diseases (I60-I69)	295	4.5	11.5
7	Diabetes mellitus (E10-E14)	240	3.7	9.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	183	2.8	7.2
9	Viral hepatitis (B15-B19)	178	2.7	7.0
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	164	2.5	6.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	90	1.4	3.5
12	Influenza and pneumonia (J10-J18)	85	1.3	3.3
13	Septicemia (A40-A41)	76	1.2	3.0
14	Chronic lower respiratory diseases (J40-J47)	46	0.7	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	40	0.6	1.6
...	All other causes (Residual)	996	15.4	38.9
Hispanic, both sexes, 50-54 years				
...	All causes	7,549	100.0	385.2
1	Malignant neoplasms (C00-C97)	1,986	26.3	101.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,355	17.9	69.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	683	9.0	34.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	577	7.6	29.4
5	Cerebrovascular diseases (I60-I69)	357	4.7	18.2
6	Diabetes mellitus (E10-E14)	355	4.7	18.1
7	Viral hepatitis (B15-B19)	245	3.2	12.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	225	3.0	11.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	132	1.7	6.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	1.5	5.7
11	Septicemia (A40-A41)	103	1.4	5.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	99	1.3	5.1
13	Influenza and pneumonia (J10-J18)	80	1.1	4.1
14	Chronic lower respiratory diseases (J40-J47)	72	1.0	3.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	0.9	3.4
...	All other causes (Residual)	1,103	14.6	56.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	8,241	100.0	549.3
1	Malignant neoplasms (C00-C97)	2,390	29.0	159.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,695	20.6	113.0
3	Diabetes mellitus (E10-E14)	518	6.3	34.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	512	6.2	34.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	495	6.0	33.0
6	Cerebrovascular diseases (I60-I69)	438	5.3	29.2
7	Viral hepatitis (B15-B19)	218	2.6	14.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	2.1	11.3
9	Chronic lower respiratory diseases (J40-J47)	121	1.5	8.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	117	1.4	7.8
11	Septicemia (A40-A41)	110	1.3	7.3
12	Influenza and pneumonia (J10-J18)	103	1.2	6.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	92	1.1	6.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	81	1.0	5.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	0.6	3.5
...	All other causes (Residual)	1,129	13.7	75.3
Hispanic, both sexes, 60-64 years				
...	All causes	8,464	100.0	813.9
1	Malignant neoplasms (C00-C97)	2,654	31.4	255.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,917	22.6	184.3
3	Diabetes mellitus (E10-E14)	563	6.7	54.1
4	Cerebrovascular diseases (I60-I69)	431	5.1	41.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	375	4.4	36.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	332	3.9	31.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	197	2.3	18.9
8	Chronic lower respiratory diseases (J40-J47)	184	2.2	17.7
9	Influenza and pneumonia (J10-J18)	146	1.7	14.0
10	Septicemia (A40-A41)	141	1.7	13.6
11	Viral hepatitis (B15-B19)	111	1.3	10.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.1	8.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	77	0.9	7.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	0.9	6.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.6	5.3
...	All other causes (Residual)	1,120	13.2	107.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	9,550	100.0	1,229.6
1	Malignant neoplasms (C00-C97)	2,999	31.4	386.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,254	23.6	290.2
3	Diabetes mellitus (E10-E14)	748	7.8	96.3
4	Cerebrovascular diseases (I60-I69)	522	5.5	67.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	320	3.4	41.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	272	2.8	35.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	269	2.8	34.6
8	Chronic lower respiratory diseases (J40-J47)	249	2.6	32.1
9	Influenza and pneumonia (J10-J18)	154	1.6	19.8
10	Septicemia (A40-A41)	148	1.5	19.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	101	1.1	13.0
12	Viral hepatitis (B15-B19)	83	0.9	10.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	53	0.6	6.8
13	Alzheimer's disease (G30)	53	0.6	6.8
15	Aortic aneurysm and dissection (I71)	46	0.5	5.9
...	All other causes (Residual)	1,279	13.4	164.7
Hispanic, both sexes, 70-74 years				
...	All causes	11,511	100.0	1,922.0
1	Malignant neoplasms (C00-C97)	3,355	29.1	560.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,935	25.5	490.1
3	Diabetes mellitus (E10-E14)	875	7.6	146.1
4	Cerebrovascular diseases (I60-I69)	679	5.9	113.4
5	Chronic lower respiratory diseases (J40-J47)	377	3.3	62.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	285	2.5	47.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	271	2.4	45.2
8	Influenza and pneumonia (J10-J18)	269	2.3	44.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	245	2.1	40.9
10	Septicemia (A40-A41)	194	1.7	32.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	132	1.1	22.0
12	Alzheimer's disease (G30)	119	1.0	19.9
13	Parkinson's disease (G20-G21)	92	0.8	15.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.6	11.5
15	Viral hepatitis (B15-B19)	65	0.6	10.9
...	All other causes (Residual)	1,549	13.5	258.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	14,119	100.0	3,052.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,852	27.3	832.8
2	Malignant neoplasms (C00-C97)	3,577	25.3	773.4
3	Diabetes mellitus (E10-E14)	951	6.7	205.6
4	Cerebrovascular diseases (I60-I69)	909	6.4	196.5
5	Chronic lower respiratory diseases (J40-J47)	558	4.0	120.6
6	Influenza and pneumonia (J10-J18)	405	2.9	87.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	365	2.6	78.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	2.1	63.8
9	Alzheimer's disease (G30)	292	2.1	63.1
10	Septicemia (A40-A41)	251	1.8	54.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	242	1.7	52.3
12	Parkinson's disease (G20-G21)	140	1.0	30.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	128	0.9	27.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	89	0.6	19.2
15	Pneumonitis due to solids and liquids (J69)	86	0.6	18.6
...	All other causes (Residual)	1,979	14.0	427.9
Hispanic, both sexes, 80-84 years				
...	All causes	14,653	100.0	4,794.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,328	29.5	1,416.2
2	Malignant neoplasms (C00-C97)	2,906	19.8	950.9
3	Cerebrovascular diseases (I60-I69)	1,104	7.5	361.2
4	Diabetes mellitus (E10-E14)	807	5.5	264.1
5	Chronic lower respiratory diseases (J40-J47)	621	4.2	203.2
6	Alzheimer's disease (G30)	556	3.8	181.9
7	Influenza and pneumonia (J10-J18)	472	3.2	154.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	399	2.7	130.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	280	1.9	91.6
10	Septicemia (A40-A41)	249	1.7	81.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	180	1.2	58.9
12	Parkinson's disease (G20-G21)	179	1.2	58.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	176	1.2	57.6
14	Pneumonitis due to solids and liquids (J69)	113	0.8	37.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	103	0.7	33.7
...	All other causes (Residual)	2,180	14.9	713.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	11,831	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,801	32.1	---
2	Malignant neoplasms (C00-C97)	1,778	15.0	---
3	Cerebrovascular diseases (I60-I69)	912	7.7	---
4	Alzheimer's disease (G30)	608	5.1	---
5	Diabetes mellitus (E10-E14)	581	4.9	---
6	Chronic lower respiratory diseases (J40-J47)	510	4.3	---
7	Influenza and pneumonia (J10-J18)	483	4.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	290	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	188	1.6	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	1.6	---
11	Septicemia (A40-A41)	180	1.5	---
12	Parkinson's disease (G20-G21)	156	1.3	---
13	Pneumonitis due to solids and liquids (J69)	95	0.8	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	80	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.6	---
...	All other causes (Residual)	1,913	16.2	---
Hispanic, both sexes, 90-94 years				
...	All causes	7,521	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,702	35.9	---
2	Malignant neoplasms (C00-C97)	734	9.8	---
3	Cerebrovascular diseases (I60-I69)	583	7.8	---
4	Alzheimer's disease (G30)	493	6.6	---
5	Influenza and pneumonia (J10-J18)	337	4.5	---
6	Chronic lower respiratory diseases (J40-J47)	328	4.4	---
7	Diabetes mellitus (E10-E14)	264	3.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	156	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	126	1.7	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	1.5	---
11	Septicemia (A40-A41)	101	1.3	---
12	Parkinson's disease (G20-G21)	74	1.0	---
13	Pneumonitis due to solids and liquids (J69)	54	0.7	---
14	Atherosclerosis (I70)	51	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	0.5	---
...	All other causes (Residual)	1,364	18.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	2,960	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,201	40.6	---
2	Cerebrovascular diseases (I60-I69)	209	7.1	---
3	Malignant neoplasms (C00-C97)	206	7.0	---
4	Alzheimer's disease (G30)	173	5.8	---
5	Influenza and pneumonia (J10-J18)	145	4.9	---
6	Chronic lower respiratory diseases (J40-J47)	93	3.1	---
7	Diabetes mellitus (E10-E14)	86	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	2.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	50	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	1.6	---
11	Pneumonitis due to solids and liquids (J69)	41	1.4	---
12	Septicemia (A40-A41)	26	0.9	---
13	Atherosclerosis (I70)	24	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.5	---
15	Nutritional deficiencies (E40-E64)	12	0.4	---
...	All other causes (Residual)	575	19.4	---
Hispanic, both sexes, 100 years and over				
...	All causes	749	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	323	43.1	---
2	Cerebrovascular diseases (I60-I69)	52	6.9	---
3	Alzheimer's disease (G30)	41	5.5	---
4	Malignant neoplasms (C00-C97)	36	4.8	---
5	Influenza and pneumonia (J10-J18)	34	4.5	---
6	Chronic lower respiratory diseases (J40-J47)	26	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	2.1	---
9	Diabetes mellitus (E10-E14)	14	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	1.7	---
11	Pneumonitis due to solids and liquids (J69)	8	1.1	---
12	Septicemia (A40-A41)	7	0.9	---
13	Atherosclerosis (I70)	4	0.5	---
14	Nutritional deficiencies (E40-E64)	3	0.4	---
15	Parkinson's disease (G20-G21)	2	0.3	---
15	Peptic ulcer (K25-K28)	2	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.3	---
...	All other causes (Residual)	143	19.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	74,250	100.0	323.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,518	20.9	67.7
2	Malignant neoplasms (C00-C97)	13,856	18.7	60.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,102	12.3	39.7
4	Cerebrovascular diseases (I60-I69)	3,269	4.4	14.3
5	Diabetes mellitus (E10-E14)	3,140	4.2	13.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,004	4.0	13.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,527	3.4	11.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,813	2.4	7.9
9	Chronic lower respiratory diseases (J40-J47)	1,708	2.3	7.5
10	Certain conditions originating in the perinatal period (P00-P96)	1,565	2.1	6.8
11	Influenza and pneumonia (J10-J18)	1,439	1.9	6.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,295	1.7	5.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,275	1.7	5.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,046	1.4	4.6
15	Septicemia (A40-A41)	932	1.3	4.1
...	All other causes (Residual)	12,761	17.2	55.7
Hispanic, male, 1-4 years				
...	All causes	549	100.0	28.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	39.2	11.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	12.6	3.6
3	Malignant neoplasms (C00-C97)	49	8.9	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.4	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.5	*
6	Influenza and pneumonia (J10-J18)	13	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
8	Septicemia (A40-A41)	6	1.1	*
8	Cerebrovascular diseases (I60-I69)	6	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
12	Meningococcal infection (A39)	2	0.4	*
12	Anemias (D50-D64)	2	0.4	*
12	Meningitis (G00,G03)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	111	20.2	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	283	100.0	13.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	39.6	5.4
2	Malignant neoplasms (C00-C97)	51	18.0	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	5.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	4.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	4.2	*
6	Chronic lower respiratory diseases (J40-J47)	8	2.8	*
7	Septicemia (A40-A41)	6	2.1	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.1	*
9	Cerebrovascular diseases (I60-I69)	4	1.4	*
9	Influenza and pneumonia (J10-J18)	4	1.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	1.1	*
12	Meningitis (G00,G03)	1	0.4	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.4	*
...	All other causes (Residual)	46	16.3	2.2
Hispanic, male, 10-14 years				
...	All causes	390	100.0	19.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	37.7	7.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	12.8	2.5
3	Malignant neoplasms (C00-C97)	44	11.3	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	4.6	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	4.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.8	*
7	Cerebrovascular diseases (I60-I69)	8	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.3	*
9	Septicemia (A40-A41)	4	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	4	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.8	*
11	Diabetes mellitus (E10-E14)	3	0.8	*
11	Pneumonitis due to solids and liquids (J69)	3	0.8	*
14	Influenza and pneumonia (J10-J18)	2	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
...	All other causes (Residual)	65	16.7	3.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,831	100.0	98.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	818	44.7	43.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	522	28.5	28.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	147	8.0	7.9
4	Malignant neoplasms (C00-C97)	71	3.9	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	1.9	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	1.5	1.5
7	Cerebrovascular diseases (I60-I69)	12	0.7	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.3	*
8	Chronic lower respiratory diseases (J40-J47)	6	0.3	*
8	Legal intervention (Y35,Y89.0)	6	0.3	*
11	Meningitis (G00,G03)	5	0.3	*
11	Influenza and pneumonia (J10-J18)	5	0.3	*
13	Septicemia (A40-A41)	4	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.2	*
14	Diabetes mellitus (E10-E14)	3	0.2	*
14	Aortic aneurysm and dissection (I71)	3	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
...	All other causes (Residual)	154	8.4	8.2
Hispanic, male, 20-24 years				
...	All causes	2,881	100.0	141.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,351	46.9	66.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	688	23.9	33.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	305	10.6	15.0
4	Malignant neoplasms (C00-C97)	130	4.5	6.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	2.2	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.7	1.0
7	Cerebrovascular diseases (I60-I69)	19	0.7	*
7	Legal intervention (Y35,Y89.0)	19	0.7	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.5	*
10	Diabetes mellitus (E10-E14)	11	0.4	*
11	Influenza and pneumonia (J10-J18)	8	0.3	*
12	Septicemia (A40-A41)	7	0.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.1	*
...	All other causes (Residual)	231	8.0	11.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,593	100.0	111.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,079	41.6	46.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	545	21.0	23.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	266	10.3	11.5
4	Malignant neoplasms (C00-C97)	131	5.1	5.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	4.3	4.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	47	1.8	2.0
7	Cerebrovascular diseases (I60-I69)	27	1.0	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.8	0.9
9	Legal intervention (Y35,Y89.0)	19	0.7	*
10	Influenza and pneumonia (J10-J18)	16	0.6	*
11	Diabetes mellitus (E10-E14)	12	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	10	0.4	*
13	Septicemia (A40-A41)	9	0.3	*
13	Aortic aneurysm and dissection (I71)	9	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.3	*
...	All other causes (Residual)	283	10.9	12.2
Hispanic, male, 30-34 years				
...	All causes	2,431	100.0	113.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	921	37.9	43.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	335	13.8	15.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	9.0	10.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	7.4	8.4
5	Malignant neoplasms (C00-C97)	139	5.7	6.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	105	4.3	4.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	56	2.3	2.6
8	Cerebrovascular diseases (I60-I69)	34	1.4	1.6
9	Diabetes mellitus (E10-E14)	24	1.0	1.1
10	Legal intervention (Y35,Y89.0)	19	0.8	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.7	*
12	Influenza and pneumonia (J10-J18)	17	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.6	*
14	Septicemia (A40-A41)	13	0.5	*
15	Viral hepatitis (B15-B19)	8	0.3	*
...	All other causes (Residual)	328	13.5	15.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,685	100.0	142.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	825	30.7	43.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	269	10.0	14.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	250	9.3	13.3
4	Malignant neoplasms (C00-C97)	207	7.7	11.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	198	7.4	10.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	7.2	10.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	119	4.4	6.3
8	Cerebrovascular diseases (I60-I69)	68	2.5	3.6
9	Diabetes mellitus (E10-E14)	41	1.5	2.2
10	Viral hepatitis (B15-B19)	20	0.7	1.1
10	Influenza and pneumonia (J10-J18)	20	0.7	1.1
12	Septicemia (A40-A41)	19	0.7	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	14	0.5	*
15	Aortic aneurysm and dissection (I71)	13	0.5	*
...	All other causes (Residual)	415	15.5	22.0
Hispanic, male, 40-44 years				
...	All causes	3,538	100.0	215.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	866	24.5	52.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	459	13.0	28.0
3	Malignant neoplasms (C00-C97)	358	10.1	21.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	258	7.3	15.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	231	6.5	14.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	181	5.1	11.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	171	4.8	10.4
8	Cerebrovascular diseases (I60-I69)	125	3.5	7.6
9	Diabetes mellitus (E10-E14)	77	2.2	4.7
10	Viral hepatitis (B15-B19)	68	1.9	4.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	0.9	1.9
12	Influenza and pneumonia (J10-J18)	30	0.8	1.8
13	Septicemia (A40-A41)	24	0.7	1.5
14	Aortic aneurysm and dissection (I71)	21	0.6	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	0.4	*
...	All other causes (Residual)	623	17.6	38.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,277	100.0	326.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	705	16.5	53.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	702	16.4	53.6
3	Malignant neoplasms (C00-C97)	643	15.0	49.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	402	9.4	30.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	248	5.8	18.9
6	Cerebrovascular diseases (I60-I69)	168	3.9	12.8
7	Diabetes mellitus (E10-E14)	155	3.6	11.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	3.4	11.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	3.2	10.3
10	Viral hepatitis (B15-B19)	131	3.1	10.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.4	4.7
12	Influenza and pneumonia (J10-J18)	48	1.1	3.7
13	Septicemia (A40-A41)	46	1.1	3.5
14	Chronic lower respiratory diseases (J40-J47)	24	0.6	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.5	1.7
...	All other causes (Residual)	642	15.0	49.0
Hispanic, male, 50-54 years				
...	All causes	4,957	100.0	507.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,024	20.7	104.7
2	Malignant neoplasms (C00-C97)	990	20.0	101.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	534	10.8	54.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	453	9.1	46.3
5	Cerebrovascular diseases (I60-I69)	221	4.5	22.6
6	Diabetes mellitus (E10-E14)	215	4.3	22.0
7	Viral hepatitis (B15-B19)	188	3.8	19.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	178	3.6	18.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	1.8	8.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	1.6	8.2
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	1.5	7.6
12	Septicemia (A40-A41)	66	1.3	6.8
13	Influenza and pneumonia (J10-J18)	54	1.1	5.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	41	0.8	4.2
15	Chronic lower respiratory diseases (J40-J47)	39	0.8	4.0
...	All other causes (Residual)	713	14.4	72.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	5,239	100.0	718.7
1	Malignant neoplasms (C00-C97)	1,334	25.5	183.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,194	22.8	163.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	384	7.3	52.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	382	7.3	52.4
5	Diabetes mellitus (E10-E14)	305	5.8	41.8
6	Cerebrovascular diseases (I60-I69)	252	4.8	34.6
7	Viral hepatitis (B15-B19)	147	2.8	20.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	93	1.8	12.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	1.7	12.2
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	1.3	9.5
11	Influenza and pneumonia (J10-J18)	68	1.3	9.3
12	Septicemia (A40-A41)	66	1.3	9.1
13	Chronic lower respiratory diseases (J40-J47)	65	1.2	8.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	0.9	6.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	0.9	6.3
...	All other causes (Residual)	697	13.3	95.6
Hispanic, male, 60-64 years				
...	All causes	5,032	100.0	1,030.0
1	Malignant neoplasms (C00-C97)	1,403	27.9	287.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,269	25.2	259.7
3	Diabetes mellitus (E10-E14)	303	6.0	62.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	275	5.5	56.3
5	Cerebrovascular diseases (I60-I69)	249	4.9	51.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	4.8	49.3
7	Chronic lower respiratory diseases (J40-J47)	106	2.1	21.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	2.0	21.1
9	Influenza and pneumonia (J10-J18)	85	1.7	17.4
10	Septicemia (A40-A41)	72	1.4	14.7
11	Viral hepatitis (B15-B19)	67	1.3	13.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	64	1.3	13.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	1.2	12.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	1.1	11.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	0.7	7.4
...	All other causes (Residual)	646	12.8	132.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	5,505	100.0	1,553.4
1	Malignant neoplasms (C00-C97)	1,688	30.7	476.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,424	25.9	401.8
3	Diabetes mellitus (E10-E14)	391	7.1	110.3
4	Cerebrovascular diseases (I60-I69)	270	4.9	76.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	197	3.6	55.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	184	3.3	51.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	147	2.7	41.5
8	Chronic lower respiratory diseases (J40-J47)	140	2.5	39.5
9	Influenza and pneumonia (J10-J18)	88	1.6	24.8
10	Septicemia (A40-A41)	87	1.6	24.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	1.0	16.1
12	Viral hepatitis (B15-B19)	43	0.8	12.1
13	Aortic aneurysm and dissection (I71)	33	0.6	9.3
14	Alzheimer's disease (G30)	31	0.6	8.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	29	0.5	8.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	0.5	8.2
...	All other causes (Residual)	667	12.1	188.2
Hispanic, male, 70-74 years				
...	All causes	6,275	100.0	2,393.6
1	Malignant neoplasms (C00-C97)	1,856	29.6	708.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,693	27.0	645.8
3	Diabetes mellitus (E10-E14)	445	7.1	169.7
4	Cerebrovascular diseases (I60-I69)	347	5.5	132.4
5	Chronic lower respiratory diseases (J40-J47)	209	3.3	79.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	154	2.5	58.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	150	2.4	57.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	150	2.4	57.2
9	Influenza and pneumonia (J10-J18)	138	2.2	52.6
10	Septicemia (A40-A41)	101	1.6	38.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	1.0	22.9
12	Parkinson's disease (G20-G21)	56	0.9	21.4
13	Alzheimer's disease (G30)	50	0.8	19.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.7	16.0
15	Aortic aneurysm and dissection (I71)	34	0.5	13.0
...	All other causes (Residual)	790	12.6	301.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	7,224	100.0	3,720.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,052	28.4	1,056.9
2	Malignant neoplasms (C00-C97)	1,904	26.4	980.6
3	Cerebrovascular diseases (I60-I69)	475	6.6	244.6
4	Diabetes mellitus (E10-E14)	437	6.0	225.1
5	Chronic lower respiratory diseases (J40-J47)	301	4.2	155.0
6	Influenza and pneumonia (J10-J18)	203	2.8	104.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	173	2.4	89.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	2.3	84.5
9	Septicemia (A40-A41)	125	1.7	64.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	121	1.7	62.3
11	Alzheimer's disease (G30)	109	1.5	56.1
12	Parkinson's disease (G20-G21)	92	1.3	47.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	59	0.8	30.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.7	24.7
15	Pneumonitis due to solids and liquids (J69)	47	0.7	24.2
...	All other causes (Residual)	914	12.7	470.7
Hispanic, male, 80-84 years				
...	All causes	6,985	100.0	5,719.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,086	29.9	1,708.2
2	Malignant neoplasms (C00-C97)	1,570	22.5	1,285.7
3	Cerebrovascular diseases (I60-I69)	451	6.5	369.3
4	Diabetes mellitus (E10-E14)	378	5.4	309.5
5	Chronic lower respiratory diseases (J40-J47)	320	4.6	262.0
6	Alzheimer's disease (G30)	230	3.3	188.3
7	Influenza and pneumonia (J10-J18)	213	3.0	174.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	187	2.7	153.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	2.0	113.8
10	Septicemia (A40-A41)	117	1.7	95.8
11	Parkinson's disease (G20-G21)	110	1.6	90.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	72	1.0	59.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	64	0.9	52.4
14	Pneumonitis due to solids and liquids (J69)	54	0.8	44.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.7	39.3
...	All other causes (Residual)	946	13.5	774.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	4,813	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,573	32.7	---
2	Malignant neoplasms (C00-C97)	850	17.7	---
3	Cerebrovascular diseases (I60-I69)	293	6.1	---
4	Chronic lower respiratory diseases (J40-J47)	243	5.0	---
5	Diabetes mellitus (E10-E14)	232	4.8	---
6	Influenza and pneumonia (J10-J18)	207	4.3	---
7	Alzheimer's disease (G30)	189	3.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	117	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	1.9	---
10	Parkinson's disease (G20-G21)	92	1.9	---
11	Septicemia (A40-A41)	77	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	1.4	---
13	Pneumonitis due to solids and liquids (J69)	46	1.0	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	37	0.8	---
15	Aortic aneurysm and dissection (I71)	30	0.6	---
...	All other causes (Residual)	668	13.9	---
Hispanic, male, 90-94 years				
...	All causes	2,526	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	919	36.4	---
2	Malignant neoplasms (C00-C97)	322	12.7	---
3	Cerebrovascular diseases (I60-I69)	156	6.2	---
4	Influenza and pneumonia (J10-J18)	137	5.4	---
5	Chronic lower respiratory diseases (J40-J47)	134	5.3	---
6	Alzheimer's disease (G30)	118	4.7	---
7	Diabetes mellitus (E10-E14)	81	3.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	1.7	---
10	Septicemia (A40-A41)	37	1.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.5	---
12	Parkinson's disease (G20-G21)	35	1.4	---
13	Pneumonitis due to solids and liquids (J69)	17	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.6	---
15	Atherosclerosis (I70)	14	0.6	---
...	All other causes (Residual)	402	15.9	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	833	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	315	37.8	---
2	Malignant neoplasms (C00-C97)	88	10.6	---
3	Cerebrovascular diseases (I60-I69)	59	7.1	---
4	Alzheimer's disease (G30)	44	5.3	---
5	Chronic lower respiratory diseases (J40-J47)	40	4.8	---
6	Influenza and pneumonia (J10-J18)	35	4.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	3.2	---
8	Diabetes mellitus (E10-E14)	23	2.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	2.0	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	1.4	---
10	Pneumonitis due to solids and liquids (J69)	12	1.4	---
12	Septicemia (A40-A41)	7	0.8	---
13	Parkinson's disease (G20-G21)	5	0.6	---
13	Atherosclerosis (I70)	5	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	---
15	Aortic aneurysm and dissection (I71)	4	0.5	---
...	All other causes (Residual)	136	16.3	---
Hispanic, male, 100 years and over				
...	All causes	195	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	42.1	---
2	Malignant neoplasms (C00-C97)	17	8.7	---
3	Chronic lower respiratory diseases (J40-J47)	13	6.7	---
4	Influenza and pneumonia (J10-J18)	12	6.2	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	5.6	---
6	Alzheimer's disease (G30)	8	4.1	---
6	Cerebrovascular diseases (I60-I69)	8	4.1	---
8	Atherosclerosis (I70)	3	1.5	---
9	Septicemia (A40-A41)	2	1.0	---
9	Diabetes mellitus (E10-E14)	2	1.0	---
9	Parkinson's disease (G20-G21)	2	1.0	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	1.0	---
9	Pneumonitis due to solids and liquids (J69)	2	1.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	1.0	---
15	Peptic ulcer (K25-K28)	1	0.5	---
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	---
...	All other causes (Residual)	27	13.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	58,754	100.0	274.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,403	22.8	62.6
2	Malignant neoplasms (C00-C97)	12,777	21.7	59.7
3	Cerebrovascular diseases (I60-I69)	3,736	6.4	17.5
4	Diabetes mellitus (E10-E14)	3,147	5.4	14.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,950	5.0	13.8
6	Chronic lower respiratory diseases (J40-J47)	1,602	2.7	7.5
7	Alzheimer's disease (G30)	1,587	2.7	7.4
8	Influenza and pneumonia (J10-J18)	1,527	2.6	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,297	2.2	6.1
10	Certain conditions originating in the perinatal period (P00-P96)	1,239	2.1	5.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,065	1.8	5.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	906	1.5	4.2
13	Septicemia (A40-A41)	897	1.5	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	697	1.2	3.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	520	0.9	2.4
...	All other causes (Residual)	11,404	19.4	53.3
Hispanic, female, 1-4 years				
...	All causes	438	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	33.3	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	12.8	3.1
3	Malignant neoplasms (C00-C97)	33	7.5	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	6.2	1.5
5	Influenza and pneumonia (J10-J18)	15	3.4	*
6	Septicemia (A40-A41)	12	2.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
9	Meningitis (G00,G03)	4	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
12	Cerebrovascular diseases (I60-I69)	3	0.7	*
13	Anemias (D50-D64)	2	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	108	24.7	5.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	201	100.0	10.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	32.3	3.3
2	Malignant neoplasms (C00-C97)	39	19.4	2.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	8.0	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	6.0	*
5	Influenza and pneumonia (J10-J18)	8	4.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	3.0	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	2.5	*
8	Septicemia (A40-A41)	3	1.5	*
8	Cerebrovascular diseases (I60-I69)	3	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	3	1.5	*
11	Diabetes mellitus (E10-E14)	2	1.0	*
12	Tuberculosis (A16-A19)	1	0.5	*
12	Meningococcal infection (A39)	1	0.5	*
12	Anemias (D50-D64)	1	0.5	*
12	Nutritional deficiencies (E40-E64)	1	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (Residual)	33	16.4	1.7
Hispanic, female, 10-14 years				
...	All causes	263	100.0	13.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	33.5	4.6
2	Malignant neoplasms (C00-C97)	49	18.6	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	6.1	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	5.7	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	4.6	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	2.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.5	*
8	Septicemia (A40-A41)	3	1.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.8	*
9	Cerebrovascular diseases (I60-I69)	2	0.8	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.8	*
9	Diseases of appendix (K35-K38)	2	0.8	*
13	Meningococcal infection (A39)	1	0.4	*
13	Anemias (D50-D64)	1	0.4	*
13	Diabetes mellitus (E10-E14)	1	0.4	*
13	Influenza and pneumonia (J10-J18)	1	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
13	Pregnancy, childbirth and the puerperium (O00-099)	1	0.4	*
...	All other causes (Residual)	55	20.9	2.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	535	100.0	30.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	240	44.9	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	9.7	3.0
3	Malignant neoplasms (C00-C97)	51	9.5	2.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	9.5	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	2.4	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.5	*
7	Cerebrovascular diseases (I60-I69)	8	1.5	*
7	Pregnancy, childbirth and the puerperium (O00-099)	8	1.5	*
10	Septicemia (A40-A41)	3	0.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.6	*
10	Influenza and pneumonia (J10-J18)	3	0.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
14	Anemias (D50-D64)	2	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
...	All other causes (Residual)	75	14.0	4.3
Hispanic, female, 20-24 years				
...	All causes	685	100.0	40.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	253	36.9	14.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	11.8	4.7
3	Malignant neoplasms (C00-C97)	69	10.1	4.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	5.8	2.3
5	Pregnancy, childbirth and the puerperium (O00-099)	31	4.5	1.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	2.6	*
7	Cerebrovascular diseases (I60-I69)	13	1.9	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.5	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.3	*
10	Septicemia (A40-A41)	8	1.2	*
10	Influenza and pneumonia (J10-J18)	8	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
14	Diabetes mellitus (E10-E14)	3	0.4	*
14	Aortic aneurysm and dissection (I71)	3	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
...	All other causes (Residual)	127	18.5	7.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	712	100.0	38.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	206	28.9	11.2
2	Malignant neoplasms (C00-C97)	112	15.7	6.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	8.4	3.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	6.9	2.7
5	Pregnancy, childbirth and the puerperium (O00-099)	34	4.8	1.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	4.5	1.7
7	Diabetes mellitus (E10-E14)	19	2.7	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	15	2.1	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	2.0	*
10	Septicemia (A40-A41)	12	1.7	*
10	Cerebrovascular diseases (I60-I69)	12	1.7	*
12	Influenza and pneumonia (J10-J18)	7	1.0	*
13	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
...	All other causes (Residual)	125	17.6	6.8
Hispanic, female, 30-34 years				
...	All causes	856	100.0	47.7
1	Malignant neoplasms (C00-C97)	197	23.0	11.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	194	22.7	10.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	6.4	3.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	51	6.0	2.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	4.6	2.2
6	Pregnancy, childbirth and the puerperium (O00-099)	36	4.2	2.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	28	3.3	1.6
8	Cerebrovascular diseases (I60-I69)	21	2.5	1.2
9	Diabetes mellitus (E10-E14)	19	2.2	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18	2.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.6	*
12	Influenza and pneumonia (J10-J18)	9	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
15	Septicemia (A40-A41)	7	0.8	*
...	All other causes (Residual)	152	17.8	8.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,137	100.0	69.0
1	Malignant neoplasms (C00-C97)	339	29.8	20.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	17.2	11.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	7.2	5.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	4.9	3.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	55	4.8	3.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	3.0	2.1
7	Cerebrovascular diseases (I60-I69)	32	2.8	1.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	30	2.6	1.8
9	Diabetes mellitus (E10-E14)	25	2.2	1.5
10	Pregnancy, childbirth and the puerperium (O00-099)	22	1.9	1.3
11	Septicemia (A40-A41)	16	1.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
14	Influenza and pneumonia (J10-J18)	9	0.8	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	0.7	*
...	All other causes (Residual)	210	18.5	12.8
Hispanic, female, 40-44 years				
...	All causes	1,591	100.0	107.0
1	Malignant neoplasms (C00-C97)	536	33.7	36.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	13.5	14.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	8.7	9.4
4	Cerebrovascular diseases (I60-I69)	83	5.2	5.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	73	4.6	4.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	64	4.0	4.3
7	Diabetes mellitus (E10-E14)	42	2.6	2.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	2.6	2.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	2.1	2.3
10	Septicemia (A40-A41)	20	1.3	1.3
11	Influenza and pneumonia (J10-J18)	18	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	17	1.1	*
13	Viral hepatitis (B15-B19)	13	0.8	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.8	*
14	Pregnancy, childbirth and the puerperium (O00-099)	12	0.8	*
...	All other causes (Residual)	272	17.1	18.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,209	100.0	177.0
1	Malignant neoplasms (C00-C97)	745	33.7	59.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	277	12.5	22.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	224	10.1	17.9
4	Cerebrovascular diseases (I60-I69)	127	5.7	10.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	93	4.2	7.5
6	Diabetes mellitus (E10-E14)	85	3.8	6.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	54	2.4	4.3
8	Viral hepatitis (B15-B19)	47	2.1	3.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	1.7	3.0
10	Influenza and pneumonia (J10-J18)	37	1.7	3.0
11	Septicemia (A40-A41)	30	1.4	2.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.3	2.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	1.3	2.3
14	Chronic lower respiratory diseases (J40-J47)	22	1.0	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	0.8	*
...	All other causes (Residual)	354	16.0	28.4
Hispanic, female, 50-54 years				
...	All causes	2,592	100.0	263.9
1	Malignant neoplasms (C00-C97)	996	38.4	101.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	331	12.8	33.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	5.7	15.2
4	Diabetes mellitus (E10-E14)	140	5.4	14.3
5	Cerebrovascular diseases (I60-I69)	136	5.2	13.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	124	4.8	12.6
7	Viral hepatitis (B15-B19)	57	2.2	5.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	2.0	5.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	47	1.8	4.8
10	Septicemia (A40-A41)	37	1.4	3.8
11	Chronic lower respiratory diseases (J40-J47)	33	1.3	3.4
12	Influenza and pneumonia (J10-J18)	26	1.0	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	25	1.0	2.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	1.0	2.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24	0.9	2.4
...	All other causes (Residual)	390	15.0	39.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	3,002	100.0	389.3
1	Malignant neoplasms (C00-C97)	1,056	35.2	136.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	501	16.7	65.0
3	Diabetes mellitus (E10-E14)	213	7.1	27.6
4	Cerebrovascular diseases (I60-I69)	186	6.2	24.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	4.3	16.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	111	3.7	14.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	80	2.7	10.4
8	Viral hepatitis (B15-B19)	71	2.4	9.2
9	Chronic lower respiratory diseases (J40-J47)	56	1.9	7.3
10	Septicemia (A40-A41)	44	1.5	5.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.5	5.7
12	Influenza and pneumonia (J10-J18)	35	1.2	4.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	24	0.8	3.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.5	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	0.4	*
...	All other causes (Residual)	424	14.1	55.0
Hispanic, female, 60-64 years				
...	All causes	3,432	100.0	622.4
1	Malignant neoplasms (C00-C97)	1,251	36.5	226.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	648	18.9	117.5
3	Diabetes mellitus (E10-E14)	260	7.6	47.1
4	Cerebrovascular diseases (I60-I69)	182	5.3	33.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	100	2.9	18.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	2.7	17.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	2.7	16.5
8	Chronic lower respiratory diseases (J40-J47)	78	2.3	14.1
9	Septicemia (A40-A41)	69	2.0	12.5
10	Influenza and pneumonia (J10-J18)	61	1.8	11.1
11	Viral hepatitis (B15-B19)	44	1.3	8.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35	1.0	6.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.7	4.5
14	Alzheimer's disease (G30)	15	0.4	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	0.4	*
...	All other causes (Residual)	453	13.2	82.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	4,045	100.0	957.8
1	Malignant neoplasms (C00-C97)	1,311	32.4	310.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	830	20.5	196.5
3	Diabetes mellitus (E10-E14)	357	8.8	84.5
4	Cerebrovascular diseases (I60-I69)	252	6.2	59.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	123	3.0	29.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	122	3.0	28.9
7	Chronic lower respiratory diseases (J40-J47)	109	2.7	25.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	2.2	20.8
9	Influenza and pneumonia (J10-J18)	66	1.6	15.6
10	Septicemia (A40-A41)	61	1.5	14.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.1	10.4
12	Viral hepatitis (B15-B19)	40	1.0	9.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.6	6.2
14	Alzheimer's disease (G30)	22	0.5	5.2
15	Parkinson's disease (G20-G21)	14	0.3	*
...	All other causes (Residual)	580	14.3	137.3
Hispanic, female, 70-74 years				
...	All causes	5,236	100.0	1,554.8
1	Malignant neoplasms (C00-C97)	1,499	28.6	445.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,242	23.7	368.8
3	Diabetes mellitus (E10-E14)	430	8.2	127.7
4	Cerebrovascular diseases (I60-I69)	332	6.3	98.6
5	Chronic lower respiratory diseases (J40-J47)	168	3.2	49.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	135	2.6	40.1
7	Influenza and pneumonia (J10-J18)	131	2.5	38.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	117	2.2	34.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	1.8	28.2
10	Septicemia (A40-A41)	93	1.8	27.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	1.4	21.4
12	Alzheimer's disease (G30)	69	1.3	20.5
13	Parkinson's disease (G20-G21)	36	0.7	10.7
14	Viral hepatitis (B15-B19)	34	0.6	10.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.5	8.0
...	All other causes (Residual)	756	14.4	224.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	6,895	100.0	2,569.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,800	26.1	670.7
2	Malignant neoplasms (C00-C97)	1,673	24.3	623.4
3	Diabetes mellitus (E10-E14)	514	7.5	191.5
4	Cerebrovascular diseases (I60-I69)	434	6.3	161.7
5	Chronic lower respiratory diseases (J40-J47)	257	3.7	95.8
6	Influenza and pneumonia (J10-J18)	202	2.9	75.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	2.8	71.5
8	Alzheimer's disease (G30)	183	2.7	68.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	131	1.9	48.8
10	Septicemia (A40-A41)	126	1.8	47.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	121	1.8	45.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.0	25.7
13	Parkinson's disease (G20-G21)	48	0.7	17.9
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	42	0.6	15.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.6	15.3
...	All other causes (Residual)	1,062	15.4	395.7
Hispanic, female, 80-84 years				
...	All causes	7,668	100.0	4,178.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,242	29.2	1,221.8
2	Malignant neoplasms (C00-C97)	1,336	17.4	728.1
3	Cerebrovascular diseases (I60-I69)	653	8.5	355.9
4	Diabetes mellitus (E10-E14)	429	5.6	233.8
5	Alzheimer's disease (G30)	326	4.3	177.7
6	Chronic lower respiratory diseases (J40-J47)	301	3.9	164.0
7	Influenza and pneumonia (J10-J18)	259	3.4	141.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	212	2.8	115.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	141	1.8	76.8
10	Septicemia (A40-A41)	132	1.7	71.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	116	1.5	63.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	104	1.4	56.7
13	Parkinson's disease (G20-G21)	69	0.9	37.6
14	Pneumonitis due to solids and liquids (J69)	59	0.8	32.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.7	30.0
...	All other causes (Residual)	1,234	16.1	672.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	7,018	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,228	31.7	---
2	Malignant neoplasms (C00-C97)	928	13.2	---
3	Cerebrovascular diseases (I60-I69)	619	8.8	---
4	Alzheimer's disease (G30)	419	6.0	---
5	Diabetes mellitus (E10-E14)	349	5.0	---
6	Influenza and pneumonia (J10-J18)	276	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	267	3.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	173	2.5	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	122	1.7	---
10	Septicemia (A40-A41)	103	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	1.3	---
12	Parkinson's disease (G20-G21)	64	0.9	---
13	Pneumonitis due to solids and liquids (J69)	49	0.7	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	43	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	41	0.6	---
...	All other causes (Residual)	1,243	17.7	---
Hispanic, female, 90-94 years				
...	All causes	4,995	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,783	35.7	---
2	Cerebrovascular diseases (I60-I69)	427	8.5	---
3	Malignant neoplasms (C00-C97)	412	8.2	---
4	Alzheimer's disease (G30)	375	7.5	---
5	Influenza and pneumonia (J10-J18)	200	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	194	3.9	---
7	Diabetes mellitus (E10-E14)	183	3.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	97	1.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	89	1.8	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	72	1.4	---
11	Septicemia (A40-A41)	64	1.3	---
12	Parkinson's disease (G20-G21)	39	0.8	---
13	Atherosclerosis (I70)	37	0.7	---
13	Pneumonitis due to solids and liquids (J69)	37	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.5	---
...	All other causes (Residual)	962	19.3	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	2,127	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	886	41.7	---
2	Cerebrovascular diseases (I60-I69)	150	7.1	---
3	Alzheimer's disease (G30)	129	6.1	---
4	Malignant neoplasms (C00-C97)	118	5.5	---
5	Influenza and pneumonia (J10-J18)	110	5.2	---
6	Diabetes mellitus (E10-E14)	63	3.0	---
7	Chronic lower respiratory diseases (J40-J47)	53	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	1.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	1.6	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.5	---
11	Pneumonitis due to solids and liquids (J69)	29	1.4	---
12	Septicemia (A40-A41)	19	0.9	---
12	Atherosclerosis (I70)	19	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.5	---
15	Nutritional deficiencies (E40-E64)	10	0.5	---
...	All other causes (Residual)	432	20.3	---
Hispanic, female, 100 years and over				
...	All causes	554	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	241	43.5	---
2	Cerebrovascular diseases (I60-I69)	44	7.9	---
3	Alzheimer's disease (G30)	33	6.0	---
4	Influenza and pneumonia (J10-J18)	22	4.0	---
5	Malignant neoplasms (C00-C97)	19	3.4	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	2.5	---
7	Chronic lower respiratory diseases (J40-J47)	13	2.3	---
8	Diabetes mellitus (E10-E14)	12	2.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	2.0	---
11	Pneumonitis due to solids and liquids (J69)	6	1.1	---
12	Septicemia (A40-A41)	5	0.9	---
13	Nutritional deficiencies (E40-E64)	3	0.5	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	---
15	Anemias (D50-D64)	1	0.2	---
15	Atherosclerosis (I70)	1	0.2	---
15	Peptic ulcer (K25-K28)	1	0.2	---
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	---
...	All other causes (Residual)	111	20.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, all ages				
...	All causes	1,944,617	100.0	968.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	516,883	26.6	257.4
2	Malignant neoplasms (C00-C97)	455,978	23.4	227.1
3	Chronic lower respiratory diseases (J40-J47)	111,559	5.7	55.6
4	Cerebrovascular diseases (I60-I69)	108,886	5.6	54.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91,830	4.7	45.7
6	Alzheimer's disease (G30)	64,660	3.3	32.2
7	Diabetes mellitus (E10-E14)	50,950	2.6	25.4
8	Influenza and pneumonia (J10-J18)	46,419	2.4	23.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,208	1.7	16.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27,952	1.4	13.9
11	Septicemia (A40-A41)	25,556	1.3	12.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	20,605	1.1	10.3
13	Parkinson's disease (G20-G21)	17,806	0.9	8.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,034	0.9	8.5
15	Pneumonitis due to solids and liquids (J69)	14,563	0.7	7.3
...	All other causes (Residual)	340,728	17.5	169.7
White, non-Hispanic, both sexes, 1-4 years				
...	All causes	2,279	100.0	25.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	814	35.7	8.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	253	11.1	2.8
3	Malignant neoplasms (C00-C97)	209	9.2	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	5.8	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	3.5	0.9
6	Influenza and pneumonia (J10-J18)	54	2.4	0.6
7	Septicemia (A40-A41)	44	1.9	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.9	0.5
9	Cerebrovascular diseases (I60-I69)	32	1.4	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	29	1.3	0.3
11	Chronic lower respiratory diseases (J40-J47)	15	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.6	*
13	Meningococcal infection (A39)	10	0.4	*
13	Meningitis (G00,G03)	10	0.4	*
15	Anemias (D50-D64)	9	0.4	*
...	All other causes (Residual)	531	23.3	5.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 5-9 years				
...	All causes	1,495	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	554	37.1	4.8
2	Malignant neoplasms (C00-C97)	282	18.9	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	7.5	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	4.3	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	3.0	0.4
6	Cerebrovascular diseases (I60-I69)	24	1.6	0.2
7	Septicemia (A40-A41)	20	1.3	0.2
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.3	0.2
7	Influenza and pneumonia (J10-J18)	20	1.3	0.2
10	Chronic lower respiratory diseases (J40-J47)	16	1.1	*
11	Anemias (D50-D64)	6	0.4	*
12	Meningitis (G00,G03)	5	0.3	*
12	Pneumonitis due to solids and liquids (J69)	5	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
...	All other causes (Residual)	312	20.9	2.7
White, non-Hispanic, both sexes, 10-14 years				
...	All causes	1,858	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	713	38.4	5.8
2	Malignant neoplasms (C00-C97)	258	13.9	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	141	7.6	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	4.6	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	4.5	0.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	3.8	0.6
7	Septicemia (A40-A41)	27	1.5	0.2
7	Cerebrovascular diseases (I60-I69)	27	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.1	0.2
10	Influenza and pneumonia (J10-J18)	19	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.8	*
12	Diabetes mellitus (E10-E14)	9	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
14	Aortic aneurysm and dissection (I71)	6	0.3	*
15	Anemias (D50-D64)	5	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (Residual)	365	19.6	3.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 15-19 years				
...	All causes	7,837	100.0	59.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,577	58.4	34.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,098	14.0	8.3
3	Malignant neoplasms (C00-C97)	411	5.2	3.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	338	4.3	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	2.5	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	127	1.6	1.0
7	Influenza and pneumonia (J10-J18)	51	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	37	0.5	0.3
9	Septicemia (A40-A41)	31	0.4	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.3	0.2
11	Diabetes mellitus (E10-E14)	22	0.3	0.2
12	Legal intervention (Y35,Y89.0)	16	0.2	*
13	Pregnancy, childbirth and the puerperium (O00-099)	15	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	14	0.2	*
...	All other causes (Residual)	866	11.1	6.5
White, non-Hispanic, both sexes, 20-24 years				
...	All causes	12,218	100.0	93.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,626	54.2	50.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,846	15.1	14.1
3	Malignant neoplasms (C00-C97)	569	4.7	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	562	4.6	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	414	3.4	3.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	1.2	1.1
7	Cerebrovascular diseases (I60-I69)	68	0.6	0.5
8	Diabetes mellitus (E10-E14)	67	0.5	0.5
9	Influenza and pneumonia (J10-J18)	63	0.5	0.5
9	Pregnancy, childbirth and the puerperium (O00-099)	63	0.5	0.5
11	Chronic lower respiratory diseases (J40-J47)	41	0.3	0.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	40	0.3	0.3
13	Septicemia (A40-A41)	38	0.3	0.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.2	0.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	0.2	0.2
...	All other causes (Residual)	1,625	13.3	12.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 25-29 years				
...	All causes	11,942	100.0	96.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,477	45.9	44.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,870	15.7	15.1
3	Malignant neoplasms (C00-C97)	826	6.9	6.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	641	5.4	5.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	513	4.3	4.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	147	1.2	1.2
7	Diabetes mellitus (E10-E14)	114	1.0	0.9
8	Cerebrovascular diseases (I60-I69)	85	0.7	0.7
9	Influenza and pneumonia (J10-J18)	81	0.7	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	60	0.5	0.5
10	Pregnancy, childbirth and the puerperium (O00-099)	60	0.5	0.5
12	Septicemia (A40-A41)	59	0.5	0.5
13	Chronic lower respiratory diseases (J40-J47)	49	0.4	0.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.3	0.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	38	0.3	0.3
...	All other causes (Residual)	1,882	15.8	15.2
White, non-Hispanic, both sexes, 30-34 years				
...	All causes	12,664	100.0	107.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,608	36.4	39.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,819	14.4	15.5
3	Malignant neoplasms (C00-C97)	1,400	11.1	11.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,133	8.9	9.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	418	3.3	3.6
6	Diabetes mellitus (E10-E14)	220	1.7	1.9
7	Cerebrovascular diseases (I60-I69)	166	1.3	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	149	1.2	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	1.1	1.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	128	1.0	1.1
11	Influenza and pneumonia (J10-J18)	108	0.9	0.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	0.7	0.7
13	Septicemia (A40-A41)	86	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	66	0.5	0.6
15	Pregnancy, childbirth and the puerperium (O00-099)	53	0.4	0.5
...	All other causes (Residual)	2,077	16.4	17.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 35-39 years				
...	All causes	19,465	100.0	143.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,488	28.2	40.5
2	Malignant neoplasms (C00-C97)	2,858	14.7	21.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,554	13.1	18.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,361	12.1	17.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	479	2.5	3.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	437	2.2	3.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	431	2.2	3.2
8	Diabetes mellitus (E10-E14)	410	2.1	3.0
9	Cerebrovascular diseases (I60-I69)	334	1.7	2.5
10	Influenza and pneumonia (J10-J18)	185	1.0	1.4
11	Septicemia (A40-A41)	173	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	159	0.8	1.2
13	Chronic lower respiratory diseases (J40-J47)	124	0.6	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	120	0.6	0.9
15	Aortic aneurysm and dissection (I71)	79	0.4	0.6
...	All other causes (Residual)	3,273	16.8	24.1
White, non-Hispanic, both sexes, 40-44 years				
...	All causes	33,733	100.0	221.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,071	21.0	46.4
2	Malignant neoplasms (C00-C97)	6,469	19.2	42.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,599	16.6	36.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,954	8.8	19.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,256	3.7	8.2
6	Diabetes mellitus (E10-E14)	847	2.5	5.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	762	2.3	5.0
8	Cerebrovascular diseases (I60-I69)	752	2.2	4.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	560	1.7	3.7
10	Chronic lower respiratory diseases (J40-J47)	415	1.2	2.7
11	Influenza and pneumonia (J10-J18)	364	1.1	2.4
12	Septicemia (A40-A41)	317	0.9	2.1
13	Viral hepatitis (B15-B19)	252	0.7	1.7
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	227	0.7	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	181	0.5	1.2
...	All other causes (Residual)	5,707	16.9	37.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 45-49 years				
...	All causes	54,044	100.0	331.4
1	Malignant neoplasms (C00-C97)	13,629	25.2	83.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,410	19.3	63.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,945	14.7	48.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,424	6.3	21.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,401	4.4	14.7
6	Cerebrovascular diseases (I60-I69)	1,379	2.6	8.5
7	Diabetes mellitus (E10-E14)	1,328	2.5	8.1
8	Chronic lower respiratory diseases (J40-J47)	1,027	1.9	6.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	758	1.4	4.6
10	Viral hepatitis (B15-B19)	737	1.4	4.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	573	1.1	3.5
12	Influenza and pneumonia (J10-J18)	520	1.0	3.2
13	Septicemia (A40-A41)	518	1.0	3.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	442	0.8	2.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	255	0.5	1.6
...	All other causes (Residual)	8,698	16.1	53.3
White, non-Hispanic, both sexes, 50-54 years				
...	All causes	73,925	100.0	487.4
1	Malignant neoplasms (C00-C97)	22,829	30.9	150.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,728	21.3	103.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,674	9.0	44.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,190	4.3	21.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,140	4.2	20.7
6	Diabetes mellitus (E10-E14)	2,137	2.9	14.1
7	Cerebrovascular diseases (I60-I69)	2,116	2.9	14.0
8	Chronic lower respiratory diseases (J40-J47)	2,086	2.8	13.8
9	Viral hepatitis (B15-B19)	1,254	1.7	8.3
10	Septicemia (A40-A41)	819	1.1	5.4
11	Influenza and pneumonia (J10-J18)	807	1.1	5.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	671	0.9	4.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	599	0.8	3.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	448	0.6	3.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	372	0.5	2.5
...	All other causes (Residual)	11,055	15.0	72.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 55-59 years				
...	All causes	97,191	100.0	696.1
1	Malignant neoplasms (C00-C97)	35,381	36.4	253.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,987	22.6	157.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,061	5.2	36.2
4	Chronic lower respiratory diseases (J40-J47)	3,967	4.1	28.4
5	Diabetes mellitus (E10-E14)	3,318	3.4	23.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,088	3.2	22.1
7	Cerebrovascular diseases (I60-I69)	2,842	2.9	20.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,551	2.6	18.3
9	Septicemia (A40-A41)	1,211	1.2	8.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,080	1.1	7.7
11	Influenza and pneumonia (J10-J18)	1,005	1.0	7.2
12	Viral hepatitis (B15-B19)	898	0.9	6.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	513	0.5	3.7
14	Aortic aneurysm and dissection (I71)	457	0.5	3.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	453	0.5	3.2
...	All other causes (Residual)	13,379	13.8	95.8
White, non-Hispanic, both sexes, 60-64 years				
...	All causes	113,127	100.0	1,081.7
1	Malignant neoplasms (C00-C97)	44,000	38.9	420.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,066	23.0	249.2
3	Chronic lower respiratory diseases (J40-J47)	6,699	5.9	64.1
4	Diabetes mellitus (E10-E14)	4,080	3.6	39.0
5	Cerebrovascular diseases (I60-I69)	3,713	3.3	35.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,592	3.2	34.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,320	2.1	22.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,616	1.4	15.5
9	Septicemia (A40-A41)	1,468	1.3	14.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,432	1.3	13.7
11	Influenza and pneumonia (J10-J18)	1,308	1.2	12.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	672	0.6	6.4
13	Aortic aneurysm and dissection (I71)	604	0.5	5.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	567	0.5	5.4
15	Viral hepatitis (B15-B19)	407	0.4	3.9
...	All other causes (Residual)	14,583	12.9	139.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 65-69 years				
...	All causes	133,248	100.0	1,634.0
1	Malignant neoplasms (C00-C97)	51,411	38.6	630.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,242	22.7	370.9
3	Chronic lower respiratory diseases (J40-J47)	10,346	7.8	126.9
4	Cerebrovascular diseases (I60-I69)	5,024	3.8	61.6
5	Diabetes mellitus (E10-E14)	4,614	3.5	56.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,237	2.4	39.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,963	1.5	24.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,918	1.4	23.5
9	Septicemia (A40-A41)	1,853	1.4	22.7
10	Influenza and pneumonia (J10-J18)	1,769	1.3	21.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,183	0.9	14.5
12	Aortic aneurysm and dissection (I71)	887	0.7	10.9
13	Alzheimer's disease (G30)	858	0.6	10.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	801	0.6	9.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	766	0.6	9.4
...	All other causes (Residual)	16,376	12.3	200.8
White, non-Hispanic, both sexes, 70-74 years				
...	All causes	174,227	100.0	2,559.2
1	Malignant neoplasms (C00-C97)	60,534	34.7	889.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,904	23.5	600.8
3	Chronic lower respiratory diseases (J40-J47)	15,051	8.6	221.1
4	Cerebrovascular diseases (I60-I69)	8,013	4.6	117.7
5	Diabetes mellitus (E10-E14)	5,691	3.3	83.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,508	2.0	51.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,889	1.7	42.4
8	Influenza and pneumonia (J10-J18)	2,860	1.6	42.0
9	Septicemia (A40-A41)	2,482	1.4	36.5
10	Alzheimer's disease (G30)	2,395	1.4	35.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,876	1.1	27.6
12	Parkinson's disease (G20-G21)	1,470	0.8	21.6
13	Aortic aneurysm and dissection (I71)	1,316	0.8	19.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,203	0.7	17.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,152	0.7	16.9
...	All other causes (Residual)	22,883	13.1	336.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 75-79 years				
...	All causes	247,476	100.0	4,077.7
1	Malignant neoplasms (C00-C97)	71,385	28.8	1,176.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	62,329	25.2	1,027.0
3	Chronic lower respiratory diseases (J40-J47)	20,666	8.4	340.5
4	Cerebrovascular diseases (I60-I69)	14,162	5.7	233.4
5	Diabetes mellitus (E10-E14)	7,551	3.1	124.4
6	Alzheimer's disease (G30)	6,524	2.6	107.5
7	Influenza and pneumonia (J10-J18)	5,159	2.1	85.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,104	2.1	84.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,789	1.9	78.9
10	Septicemia (A40-A41)	3,777	1.5	62.2
11	Parkinson's disease (G20-G21)	3,346	1.4	55.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,929	0.8	31.8
13	Aortic aneurysm and dissection (I71)	1,914	0.8	31.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,828	0.7	30.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,761	0.7	29.0
...	All other causes (Residual)	35,252	14.2	580.9
White, non-Hispanic, both sexes, 80-84 years				
...	All causes	316,637	100.0	6,640.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	88,800	28.0	1,862.2
2	Malignant neoplasms (C00-C97)	69,261	21.9	1,452.4
3	Chronic lower respiratory diseases (J40-J47)	22,507	7.1	472.0
4	Cerebrovascular diseases (I60-I69)	21,806	6.9	457.3
5	Alzheimer's disease (G30)	13,749	4.3	288.3
6	Influenza and pneumonia (J10-J18)	8,711	2.8	182.7
7	Diabetes mellitus (E10-E14)	8,596	2.7	180.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,839	2.2	143.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,505	2.1	136.4
10	Parkinson's disease (G20-G21)	4,976	1.6	104.3
11	Septicemia (A40-A41)	4,718	1.5	98.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,032	1.0	63.6
13	Pneumonitis due to solids and liquids (J69)	2,926	0.9	61.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,310	0.7	48.4
15	Aortic aneurysm and dissection (I71)	2,173	0.7	45.6
...	All other causes (Residual)	49,728	15.7	1,042.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 85-89 years				
...	All causes	309,862	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	98,057	31.6	---
2	Malignant neoplasms (C00-C97)	47,001	15.2	---
3	Cerebrovascular diseases (I60-I69)	24,125	7.8	---
4	Alzheimer's disease (G30)	18,429	5.9	---
5	Chronic lower respiratory diseases (J40-J47)	17,165	5.5	---
6	Influenza and pneumonia (J10-J18)	10,157	3.3	---
7	Diabetes mellitus (E10-E14)	7,077	2.3	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,745	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,739	2.2	---
10	Parkinson's disease (G20-G21)	4,477	1.4	---
11	Septicemia (A40-A41)	4,386	1.4	---
12	Pneumonitis due to solids and liquids (J69)	3,650	1.2	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,619	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,068	0.7	---
15	Aortic aneurysm and dissection (I71)	1,790	0.6	---
...	All other causes (Residual)	54,377	17.5	---
White, non-Hispanic, both sexes, 90-94 years				
...	All causes	211,119	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,335	35.2	---
2	Malignant neoplasms (C00-C97)	21,246	10.1	---
3	Cerebrovascular diseases (I60-I69)	16,752	7.9	---
4	Alzheimer's disease (G30)	14,970	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	8,509	4.0	---
6	Influenza and pneumonia (J10-J18)	8,355	4.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,635	2.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,383	2.1	---
9	Diabetes mellitus (E10-E14)	3,606	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,826	1.3	---
11	Pneumonitis due to solids and liquids (J69)	2,611	1.2	---
12	Septicemia (A40-A41)	2,527	1.2	---
13	Parkinson's disease (G20-G21)	2,095	1.0	---
14	Atherosclerosis (I70)	1,608	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,226	0.6	---
...	All other causes (Residual)	41,435	19.6	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, both sexes, 95-99 years				
...	All causes	81,174	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,777	37.9	---
2	Cerebrovascular diseases (I60-I69)	6,296	7.8	---
3	Alzheimer's disease (G30)	6,049	7.5	---
4	Malignant neoplasms (C00-C97)	5,332	6.6	---
5	Influenza and pneumonia (J10-J18)	3,794	4.7	---
6	Chronic lower respiratory diseases (J40-J47)	2,440	3.0	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,657	2.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,503	1.9	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,372	1.7	---
10	Diabetes mellitus (E10-E14)	1,075	1.3	---
11	Pneumonitis due to solids and liquids (J69)	968	1.2	---
12	Septicemia (A40-A41)	802	1.0	---
13	Atherosclerosis (I70)	798	1.0	---
14	Parkinson's disease (G20-G21)	467	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	347	0.4	---
...	All other causes (Residual)	17,497	21.6	---
White, non-Hispanic, both sexes, 100 years and over				
...	All causes	16,098	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,526	40.5	---
2	Cerebrovascular diseases (I60-I69)	1,077	6.7	---
3	Alzheimer's disease (G30)	1,046	6.5	---
4	Influenza and pneumonia (J10-J18)	926	5.8	---
5	Malignant neoplasms (C00-C97)	634	3.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	348	2.2	---
7	Chronic lower respiratory diseases (J40-J47)	332	2.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	326	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	238	1.5	---
10	Atherosclerosis (I70)	234	1.5	---
11	Pneumonitis due to solids and liquids (J69)	182	1.1	---
12	Diabetes mellitus (E10-E14)	175	1.1	---
13	Septicemia (A40-A41)	109	0.7	---
14	Anemias (D50-D64)	54	0.3	---
15	Parkinson's disease (G20-G21)	51	0.3	---
...	All other causes (Residual)	3,840	23.9	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, all ages				
...	All causes	947,966	100.0	962.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	256,459	27.1	260.3
2	Malignant neoplasms (C00-C97)	236,393	24.9	239.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57,758	6.1	58.6
4	Chronic lower respiratory diseases (J40-J47)	52,270	5.5	53.0
5	Cerebrovascular diseases (I60-I69)	41,941	4.4	42.6
6	Diabetes mellitus (E10-E14)	25,931	2.7	26.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21,952	2.3	22.3
8	Influenza and pneumonia (J10-J18)	20,863	2.2	21.2
9	Alzheimer's disease (G30)	18,844	2.0	19.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,415	1.7	16.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	13,212	1.4	13.4
12	Septicemia (A40-A41)	11,526	1.2	11.7
13	Parkinson's disease (G20-G21)	10,277	1.1	10.4
14	Pneumonitis due to solids and liquids (J69)	7,694	0.8	7.8
15	Aortic aneurysm and dissection (I71)	6,512	0.7	6.6
...	All other causes (Residual)	149,919	15.8	152.1
White, non-Hispanic, male, 1-4 years				
...	All causes	1,247	100.0	26.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	458	36.7	9.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	122	9.8	2.6
3	Malignant neoplasms (C00-C97)	116	9.3	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	6.2	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.3	0.9
6	Influenza and pneumonia (J10-J18)	29	2.3	0.6
7	Septicemia (A40-A41)	24	1.9	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	21	1.7	0.4
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.3	*
10	Cerebrovascular diseases (I60-I69)	12	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.8	*
12	Meningococcal infection (A39)	7	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Anemias (D50-D64)	3	0.2	*
15	Meningitis (G00,G03)	3	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
...	All other causes (Residual)	293	23.5	6.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 5-9 years				
...	All causes	839	100.0	14.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	320	38.1	5.4
2	Malignant neoplasms (C00-C97)	160	19.1	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	7.2	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	3.2	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.7	0.4
6	Cerebrovascular diseases (I60-I69)	15	1.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.5	*
8	Influenza and pneumonia (J10-J18)	12	1.4	*
9	Septicemia (A40-A41)	11	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	11	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.5	*
12	Anemias (D50-D64)	3	0.4	*
12	Diabetes mellitus (E10-E14)	3	0.4	*
12	Meningitis (G00,G03)	3	0.4	*
12	Pneumonitis due to solids and liquids (J69)	3	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	168	20.0	2.8
White, non-Hispanic, male, 10-14 years				
...	All causes	1,144	100.0	18.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	446	39.0	7.0
2	Malignant neoplasms (C00-C97)	149	13.0	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	8.7	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	4.6	0.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	4.4	0.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	3.6	0.6
7	Cerebrovascular diseases (I60-I69)	19	1.7	*
8	Septicemia (A40-A41)	17	1.5	*
9	Influenza and pneumonia (J10-J18)	13	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	11	1.0	*
12	Aortic aneurysm and dissection (I71)	6	0.5	*
13	Diabetes mellitus (E10-E14)	5	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	214	18.7	3.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 15-19 years				
...	All causes	5,378	100.0	79.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,118	58.0	45.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	905	16.8	13.3
3	Malignant neoplasms (C00-C97)	257	4.8	3.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	230	4.3	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	130	2.4	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	75	1.4	1.1
7	Influenza and pneumonia (J10-J18)	23	0.4	0.3
8	Cerebrovascular diseases (I60-I69)	19	0.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.3	*
10	Septicemia (A40-A41)	14	0.3	*
10	Legal intervention (Y35,Y89.0)	14	0.3	*
12	Diabetes mellitus (E10-E14)	12	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
15	Meningococcal infection (A39)	6	0.1	*
15	Anemias (D50-D64)	6	0.1	*
15	Chronic lower respiratory diseases (J40-J47)	6	0.1	*
...	All other causes (Residual)	527	9.8	7.7
White, non-Hispanic, male, 20-24 years				
...	All causes	9,175	100.0	136.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,172	56.4	76.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,591	17.3	23.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	408	4.4	6.1
4	Malignant neoplasms (C00-C97)	351	3.8	5.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	280	3.1	4.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	0.9	1.2
7	Cerebrovascular diseases (I60-I69)	41	0.4	0.6
8	Influenza and pneumonia (J10-J18)	39	0.4	0.6
9	Diabetes mellitus (E10-E14)	35	0.4	0.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.3	0.4
11	Chronic lower respiratory diseases (J40-J47)	24	0.3	0.4
12	Septicemia (A40-A41)	19	0.2	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.2	*
13	Aortic aneurysm and dissection (I71)	17	0.2	*
15	Legal intervention (Y35,Y89.0)	14	0.2	*
...	All other causes (Residual)	1,056	11.5	15.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 25-29 years				
...	All causes	8,542	100.0	137.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,195	49.1	67.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,504	17.6	24.2
3	Malignant neoplasms (C00-C97)	461	5.4	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	442	5.2	7.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	361	4.2	5.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	1.1	1.5
7	Diabetes mellitus (E10-E14)	67	0.8	1.1
8	Cerebrovascular diseases (I60-I69)	45	0.5	0.7
9	Influenza and pneumonia (J10-J18)	44	0.5	0.7
10	Human immunodeficiency virus (HIV) disease (B20-B24)	43	0.5	0.7
11	Septicemia (A40-A41)	33	0.4	0.5
12	Chronic lower respiratory diseases (J40-J47)	28	0.3	0.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	25	0.3	0.4
14	Legal intervention (Y35,Y89.0)	23	0.3	0.4
15	Aortic aneurysm and dissection (I71)	21	0.2	0.3
...	All other causes (Residual)	1,155	13.5	18.6
White, non-Hispanic, male, 30-34 years				
...	All causes	8,573	100.0	145.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,383	39.5	57.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,456	17.0	24.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	808	9.4	13.7
4	Malignant neoplasms (C00-C97)	665	7.8	11.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	303	3.5	5.1
6	Diabetes mellitus (E10-E14)	122	1.4	2.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	109	1.3	1.8
8	Cerebrovascular diseases (I60-I69)	89	1.0	1.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	85	1.0	1.4
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	78	0.9	1.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	0.7	1.0
12	Influenza and pneumonia (J10-J18)	52	0.6	0.9
13	Septicemia (A40-A41)	44	0.5	0.7
14	Chronic lower respiratory diseases (J40-J47)	28	0.3	0.5
14	Legal intervention (Y35,Y89.0)	28	0.3	0.5
...	All other causes (Residual)	1,262	14.7	21.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 35-39 years				
...	All causes	12,463	100.0	183.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,851	30.9	56.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,955	15.7	28.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,691	13.6	24.8
4	Malignant neoplasms (C00-C97)	1,250	10.0	18.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	336	2.7	4.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	314	2.5	4.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	275	2.2	4.0
8	Diabetes mellitus (E10-E14)	242	1.9	3.6
9	Cerebrovascular diseases (I60-I69)	169	1.4	2.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	0.8	1.4
11	Influenza and pneumonia (J10-J18)	93	0.7	1.4
12	Septicemia (A40-A41)	90	0.7	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	0.6	1.0
14	Aortic aneurysm and dissection (I71)	58	0.5	0.9
15	Chronic lower respiratory diseases (J40-J47)	54	0.4	0.8
...	All other causes (Residual)	1,919	15.4	28.2
White, non-Hispanic, male, 40-44 years				
...	All causes	21,146	100.0	278.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,863	23.0	63.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,047	19.1	53.2
3	Malignant neoplasms (C00-C97)	2,915	13.8	38.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,224	10.5	29.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	797	3.8	10.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	645	3.1	8.5
7	Diabetes mellitus (E10-E14)	535	2.5	7.0
8	Cerebrovascular diseases (I60-I69)	389	1.8	5.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	386	1.8	5.1
10	Influenza and pneumonia (J10-J18)	204	1.0	2.7
11	Chronic lower respiratory diseases (J40-J47)	200	0.9	2.6
12	Septicemia (A40-A41)	172	0.8	2.3
13	Viral hepatitis (B15-B19)	165	0.8	2.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	140	0.7	1.8
15	Aortic aneurysm and dissection (I71)	102	0.5	1.3
...	All other causes (Residual)	3,362	15.9	44.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 45-49 years				
...	All causes	33,828	100.0	417.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,693	22.7	94.8
2	Malignant neoplasms (C00-C97)	6,562	19.4	80.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,509	16.3	67.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,561	7.6	31.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,629	4.8	20.1
6	Diabetes mellitus (E10-E14)	866	2.6	10.7
7	Cerebrovascular diseases (I60-I69)	734	2.2	9.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	657	1.9	8.1
9	Viral hepatitis (B15-B19)	521	1.5	6.4
10	Chronic lower respiratory diseases (J40-J47)	472	1.4	5.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	397	1.2	4.9
12	Influenza and pneumonia (J10-J18)	298	0.9	3.7
13	Septicemia (A40-A41)	284	0.8	3.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	278	0.8	3.4
15	Aortic aneurysm and dissection (I71)	187	0.6	2.3
...	All other causes (Residual)	5,180	15.3	63.9
White, non-Hispanic, male, 50-54 years				
...	All causes	46,599	100.0	621.0
1	Malignant neoplasms (C00-C97)	11,970	25.7	159.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,746	25.2	156.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,615	9.9	61.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,388	5.1	31.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,295	4.9	30.6
6	Diabetes mellitus (E10-E14)	1,345	2.9	17.9
7	Cerebrovascular diseases (I60-I69)	1,159	2.5	15.4
8	Chronic lower respiratory diseases (J40-J47)	1,066	2.3	14.2
9	Viral hepatitis (B15-B19)	929	2.0	12.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	498	1.1	6.6
11	Influenza and pneumonia (J10-J18)	473	1.0	6.3
12	Septicemia (A40-A41)	462	1.0	6.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	401	0.9	5.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	306	0.7	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	251	0.5	3.3
...	All other causes (Residual)	6,695	14.4	89.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 55-59 years				
...	All causes	59,651	100.0	869.7
1	Malignant neoplasms (C00-C97)	19,131	32.1	278.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,966	26.8	232.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,448	5.8	50.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,202	3.7	32.1
5	Chronic lower respiratory diseases (J40-J47)	2,033	3.4	29.6
6	Diabetes mellitus (E10-E14)	1,980	3.3	28.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,917	3.2	27.9
8	Cerebrovascular diseases (I60-I69)	1,600	2.7	23.3
9	Viral hepatitis (B15-B19)	669	1.1	9.8
10	Septicemia (A40-A41)	667	1.1	9.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	621	1.0	9.1
12	Influenza and pneumonia (J10-J18)	604	1.0	8.8
13	Aortic aneurysm and dissection (I71)	354	0.6	5.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	312	0.5	4.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	312	0.5	4.5
...	All other causes (Residual)	7,835	13.1	114.2
White, non-Hispanic, male, 60-64 years				
...	All causes	67,128	100.0	1,327.5
1	Malignant neoplasms (C00-C97)	24,169	36.0	478.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,935	26.7	354.7
3	Chronic lower respiratory diseases (J40-J47)	3,374	5.0	66.7
4	Diabetes mellitus (E10-E14)	2,506	3.7	49.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,399	3.6	47.4
6	Cerebrovascular diseases (I60-I69)	2,012	3.0	39.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,597	2.4	31.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,212	1.8	24.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	844	1.3	16.7
10	Influenza and pneumonia (J10-J18)	769	1.1	15.2
11	Septicemia (A40-A41)	745	1.1	14.7
12	Aortic aneurysm and dissection (I71)	442	0.7	8.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	418	0.6	8.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	312	0.5	6.2
15	Viral hepatitis (B15-B19)	278	0.4	5.5
...	All other causes (Residual)	8,116	12.1	160.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 65-69 years				
...	All causes	77,056	100.0	2,000.1
1	Malignant neoplasms (C00-C97)	28,490	37.0	739.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,874	25.8	515.9
3	Chronic lower respiratory diseases (J40-J47)	5,084	6.6	132.0
4	Cerebrovascular diseases (I60-I69)	2,735	3.5	71.0
5	Diabetes mellitus (E10-E14)	2,684	3.5	69.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,103	2.7	54.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,263	1.6	32.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,018	1.3	26.4
9	Influenza and pneumonia (J10-J18)	1,005	1.3	26.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	966	1.3	25.1
11	Septicemia (A40-A41)	954	1.2	24.8
12	Aortic aneurysm and dissection (I71)	603	0.8	15.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	445	0.6	11.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	435	0.6	11.3
15	Alzheimer's disease (G30)	425	0.6	11.0
...	All other causes (Residual)	8,972	11.6	232.9
White, non-Hispanic, male, 70-74 years				
...	All causes	96,000	100.0	3,100.3
1	Malignant neoplasms (C00-C97)	32,869	34.2	1,061.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,898	25.9	804.1
3	Chronic lower respiratory diseases (J40-J47)	7,462	7.8	241.0
4	Cerebrovascular diseases (I60-I69)	3,969	4.1	128.2
5	Diabetes mellitus (E10-E14)	3,146	3.3	101.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,061	2.1	66.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,604	1.7	51.8
8	Influenza and pneumonia (J10-J18)	1,571	1.6	50.7
9	Septicemia (A40-A41)	1,237	1.3	39.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,096	1.1	35.4
11	Alzheimer's disease (G30)	1,037	1.1	33.5
12	Parkinson's disease (G20-G21)	973	1.0	31.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	837	0.9	27.0
14	Aortic aneurysm and dissection (I71)	828	0.9	26.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	702	0.7	22.7
...	All other causes (Residual)	11,710	12.2	378.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 75-79 years				
...	All causes	129,406	100.0	4,983.5
1	Malignant neoplasms (C00-C97)	38,129	29.5	1,468.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,822	26.9	1,341.0
3	Chronic lower respiratory diseases (J40-J47)	10,217	7.9	393.5
4	Cerebrovascular diseases (I60-I69)	6,470	5.0	249.2
5	Diabetes mellitus (E10-E14)	3,971	3.1	152.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,837	2.2	109.3
7	Influenza and pneumonia (J10-J18)	2,758	2.1	106.2
8	Alzheimer's disease (G30)	2,624	2.0	101.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,549	2.0	98.2
10	Parkinson's disease (G20-G21)	2,149	1.7	82.8
11	Septicemia (A40-A41)	1,845	1.4	71.1
12	Aortic aneurysm and dissection (I71)	1,148	0.9	44.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,049	0.8	40.4
14	Pneumonitis due to solids and liquids (J69)	966	0.7	37.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	934	0.7	36.0
...	All other causes (Residual)	16,938	13.1	652.3
White, non-Hispanic, male, 80-84 years				
...	All causes	149,284	100.0	8,103.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,633	29.2	2,368.5
2	Malignant neoplasms (C00-C97)	35,687	23.9	1,937.1
3	Chronic lower respiratory diseases (J40-J47)	10,633	7.1	577.2
4	Cerebrovascular diseases (I60-I69)	8,643	5.8	469.2
5	Alzheimer's disease (G30)	4,739	3.2	257.2
6	Influenza and pneumonia (J10-J18)	4,208	2.8	228.4
7	Diabetes mellitus (E10-E14)	4,000	2.7	217.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,432	2.3	186.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,242	2.2	176.0
10	Parkinson's disease (G20-G21)	2,911	1.9	158.0
11	Septicemia (A40-A41)	2,084	1.4	113.1
12	Pneumonitis due to solids and liquids (J69)	1,660	1.1	90.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,211	0.8	65.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,173	0.8	63.7
15	Aortic aneurysm and dissection (I71)	1,160	0.8	63.0
...	All other causes (Residual)	20,868	14.0	1,132.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 85-89 years				
...	All causes	125,456	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,429	32.2	---
2	Malignant neoplasms (C00-C97)	22,365	17.8	---
3	Cerebrovascular diseases (I60-I69)	8,117	6.5	---
4	Chronic lower respiratory diseases (J40-J47)	7,459	5.9	---
5	Alzheimer's disease (G30)	5,455	4.3	---
6	Influenza and pneumonia (J10-J18)	4,416	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,153	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,019	2.4	---
9	Diabetes mellitus (E10-E14)	2,943	2.3	---
10	Parkinson's disease (G20-G21)	2,447	2.0	---
11	Pneumonitis due to solids and liquids (J69)	1,865	1.5	---
12	Septicemia (A40-A41)	1,752	1.4	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,214	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	851	0.7	---
15	Aortic aneurysm and dissection (I71)	850	0.7	---
...	All other causes (Residual)	19,121	15.2	---
White, non-Hispanic, male, 90-94 years				
...	All causes	66,870	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,837	35.6	---
2	Malignant neoplasms (C00-C97)	8,735	13.1	---
3	Cerebrovascular diseases (I60-I69)	4,390	6.6	---
4	Chronic lower respiratory diseases (J40-J47)	3,286	4.9	---
5	Alzheimer's disease (G30)	3,265	4.9	---
6	Influenza and pneumonia (J10-J18)	2,969	4.4	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,745	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,688	2.5	---
9	Diabetes mellitus (E10-E14)	1,192	1.8	---
10	Pneumonitis due to solids and liquids (J69)	1,184	1.8	---
11	Parkinson's disease (G20-G21)	994	1.5	---
12	Septicemia (A40-A41)	806	1.2	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	707	1.1	---
14	Atherosclerosis (I70)	453	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	432	0.6	---
...	All other causes (Residual)	11,187	16.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, male, 95-99 years				
...	All causes	18,370	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,026	38.2	---
2	Malignant neoplasms (C00-C97)	1,775	9.7	---
3	Cerebrovascular diseases (I60-I69)	1,149	6.3	---
4	Influenza and pneumonia (J10-J18)	1,037	5.6	---
5	Alzheimer's disease (G30)	894	4.9	---
6	Chronic lower respiratory diseases (J40-J47)	720	3.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	506	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	439	2.4	---
9	Pneumonitis due to solids and liquids (J69)	342	1.9	---
10	Diabetes mellitus (E10-E14)	247	1.3	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	223	1.2	---
12	Septicemia (A40-A41)	197	1.1	---
13	Atherosclerosis (I70)	177	1.0	---
14	Parkinson's disease (G20-G21)	170	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	88	0.5	---
...	All other causes (Residual)	3,380	18.4	---
White, non-Hispanic, male, 100 years and over				
...	All causes	2,455	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	990	40.3	---
2	Influenza and pneumonia (J10-J18)	178	7.3	---
3	Malignant neoplasms (C00-C97)	161	6.6	---
4	Cerebrovascular diseases (I60-I69)	134	5.5	---
5	Alzheimer's disease (G30)	101	4.1	---
6	Chronic lower respiratory diseases (J40-J47)	85	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	2.3	---
9	Pneumonitis due to solids and liquids (J69)	54	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	39	1.6	---
11	Atherosclerosis (I70)	29	1.2	---
12	Diabetes mellitus (E10-E14)	24	1.0	---
13	Septicemia (A40-A41)	21	0.9	---
14	Parkinson's disease (G20-G21)	14	0.6	---
15	Anemias (D50-D64)	9	0.4	---
...	All other causes (Residual)	496	20.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, all ages				
...	All causes	996,651	100.0	974.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	260,424	26.1	254.7
2	Malignant neoplasms (C00-C97)	219,585	22.0	214.7
3	Cerebrovascular diseases (I60-I69)	66,945	6.7	65.5
4	Chronic lower respiratory diseases (J40-J47)	59,289	5.9	58.0
5	Alzheimer's disease (G30)	45,816	4.6	44.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,072	3.4	33.3
7	Influenza and pneumonia (J10-J18)	25,556	2.6	25.0
8	Diabetes mellitus (E10-E14)	25,019	2.5	24.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,793	1.7	16.4
10	Septicemia (A40-A41)	14,030	1.4	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,583	1.1	10.3
12	Parkinson's disease (G20-G21)	7,529	0.8	7.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7,393	0.7	7.2
14	Pneumonitis due to solids and liquids (J69)	6,869	0.7	6.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,000	0.6	5.9
...	All other causes (Residual)	190,748	19.1	186.5
White, non-Hispanic, female, 1-4 years				
...	All causes	1,032	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	356	34.5	8.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	12.7	2.9
3	Malignant neoplasms (C00-C97)	93	9.0	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	5.4	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.7	0.9
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	2.7	0.6
7	Influenza and pneumonia (J10-J18)	25	2.4	0.6
8	Septicemia (A40-A41)	20	1.9	0.4
8	Cerebrovascular diseases (I60-I69)	20	1.9	0.4
10	Certain conditions originating in the perinatal period (P00-P96)	8	0.8	*
11	Meningitis (G00,G03)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
13	Anemias (D50-D64)	6	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
15	Pneumonitis due to solids and liquids (J69)	4	0.4	*
...	All other causes (Residual)	228	22.1	5.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 5-9 years				
...	All causes	656	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	35.7	4.2
2	Malignant neoplasms (C00-C97)	122	18.6	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	7.9	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	5.6	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.4	0.4
6	Septicemia (A40-A41)	9	1.4	*
6	Cerebrovascular diseases (I60-I69)	9	1.4	*
8	Influenza and pneumonia (J10-J18)	8	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.1	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
12	Anemias (D50-D64)	3	0.5	*
13	Meningitis (G00,G03)	2	0.3	*
13	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	Meningococcal infection (A39)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	133	20.3	2.4
White, non-Hispanic, female, 10-14 years				
...	All causes	714	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	267	37.4	4.4
2	Malignant neoplasms (C00-C97)	109	15.3	1.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	5.9	0.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	4.6	0.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	4.6	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	4.1	0.5
7	Septicemia (A40-A41)	10	1.4	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.4	*
9	Cerebrovascular diseases (I60-I69)	8	1.1	*
10	Influenza and pneumonia (J10-J18)	6	0.8	*
11	Diabetes mellitus (E10-E14)	4	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
13	Anemias (D50-D64)	3	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (Residual)	151	21.1	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 15-19 years				
...	All causes	2,459	100.0	38.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,459	59.3	22.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	193	7.8	3.0
3	Malignant neoplasms (C00-C97)	154	6.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	108	4.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	2.6	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	2.1	0.8
7	Influenza and pneumonia (J10-J18)	28	1.1	0.4
8	Cerebrovascular diseases (I60-I69)	18	0.7	*
9	Septicemia (A40-A41)	17	0.7	*
10	Pregnancy, childbirth and the puerperium (O00-099)	15	0.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.4	*
11	Diabetes mellitus (E10-E14)	10	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	7	0.3	*
14	Pneumonitis due to solids and liquids (J69)	5	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
...	All other causes (Residual)	314	12.8	4.9
White, non-Hispanic, female, 20-24 years				
...	All causes	3,043	100.0	47.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,454	47.8	22.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	8.4	4.0
3	Malignant neoplasms (C00-C97)	218	7.2	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	5.1	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	134	4.4	2.1
6	Pregnancy, childbirth and the puerperium (O00-099)	63	2.1	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	2.0	1.0
8	Diabetes mellitus (E10-E14)	32	1.1	0.5
9	Cerebrovascular diseases (I60-I69)	27	0.9	0.4
10	Influenza and pneumonia (J10-J18)	24	0.8	0.4
11	Septicemia (A40-A41)	19	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	17	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.3	*
...	All other causes (Residual)	551	18.1	8.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 25-29 years				
...	All causes	3,400	100.0	55.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,282	37.7	20.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	366	10.8	6.0
3	Malignant neoplasms (C00-C97)	365	10.7	5.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	199	5.9	3.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	152	4.5	2.5
6	Pregnancy, childbirth and the puerperium (O00-099)	60	1.8	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	1.5	0.8
8	Diabetes mellitus (E10-E14)	47	1.4	0.8
9	Cerebrovascular diseases (I60-I69)	40	1.2	0.7
10	Influenza and pneumonia (J10-J18)	37	1.1	0.6
11	Septicemia (A40-A41)	26	0.8	0.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.7	0.4
13	Chronic lower respiratory diseases (J40-J47)	21	0.6	0.3
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	18	0.5	*
...	All other causes (Residual)	693	20.4	11.3
White, non-Hispanic, female, 30-34 years				
...	All causes	4,091	100.0	70.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,225	29.9	21.0
2	Malignant neoplasms (C00-C97)	735	18.0	12.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	363	8.9	6.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	325	7.9	5.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	115	2.8	2.0
6	Diabetes mellitus (E10-E14)	98	2.4	1.7
7	Cerebrovascular diseases (I60-I69)	77	1.9	1.3
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	1.6	1.1
9	Influenza and pneumonia (J10-J18)	56	1.4	1.0
10	Pregnancy, childbirth and the puerperium (O00-099)	53	1.3	0.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	43	1.1	0.7
12	Septicemia (A40-A41)	42	1.0	0.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	40	1.0	0.7
14	Chronic lower respiratory diseases (J40-J47)	38	0.9	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.7	0.5
...	All other causes (Residual)	787	19.2	13.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 35-39 years				
...	All causes	7,002	100.0	103.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,637	23.4	24.3
2	Malignant neoplasms (C00-C97)	1,608	23.0	23.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	670	9.6	9.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	599	8.6	8.9
5	Diabetes mellitus (E10-E14)	168	2.4	2.5
6	Cerebrovascular diseases (I60-I69)	165	2.4	2.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	165	2.4	2.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	162	2.3	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	95	1.4	1.4
10	Influenza and pneumonia (J10-J18)	92	1.3	1.4
11	Septicemia (A40-A41)	83	1.2	1.2
12	Chronic lower respiratory diseases (J40-J47)	70	1.0	1.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	0.9	0.9
14	Pregnancy, childbirth and the puerperium (O00-099)	54	0.8	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	0.7	0.7
...	All other causes (Residual)	1,321	18.9	19.6
White, non-Hispanic, female, 40-44 years				
...	All causes	12,587	100.0	165.1
1	Malignant neoplasms (C00-C97)	3,554	28.2	46.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,208	17.5	29.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,552	12.3	20.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	730	5.8	9.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	459	3.6	6.0
6	Cerebrovascular diseases (I60-I69)	363	2.9	4.8
7	Diabetes mellitus (E10-E14)	312	2.5	4.1
8	Chronic lower respiratory diseases (J40-J47)	215	1.7	2.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	174	1.4	2.3
10	Influenza and pneumonia (J10-J18)	160	1.3	2.1
11	Septicemia (A40-A41)	145	1.2	1.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	117	0.9	1.5
13	Viral hepatitis (B15-B19)	87	0.7	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	0.7	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	0.7	1.1
...	All other causes (Residual)	2,342	18.6	30.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 45-49 years				
...	All causes	20,216	100.0	246.7
1	Malignant neoplasms (C00-C97)	7,067	35.0	86.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,717	13.4	33.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,436	12.0	29.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	863	4.3	10.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	772	3.8	9.4
6	Cerebrovascular diseases (I60-I69)	645	3.2	7.9
7	Chronic lower respiratory diseases (J40-J47)	555	2.7	6.8
8	Diabetes mellitus (E10-E14)	462	2.3	5.6
9	Septicemia (A40-A41)	234	1.2	2.9
10	Influenza and pneumonia (J10-J18)	222	1.1	2.7
11	Viral hepatitis (B15-B19)	216	1.1	2.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	176	0.9	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	0.8	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	0.6	1.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	110	0.5	1.3
...	All other causes (Residual)	3,463	17.1	42.3
White, non-Hispanic, female, 50-54 years				
...	All causes	27,326	100.0	356.7
1	Malignant neoplasms (C00-C97)	10,859	39.7	141.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,982	14.6	52.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,059	7.5	26.9
4	Chronic lower respiratory diseases (J40-J47)	1,020	3.7	13.3
5	Cerebrovascular diseases (I60-I69)	957	3.5	12.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	845	3.1	11.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	802	2.9	10.5
8	Diabetes mellitus (E10-E14)	792	2.9	10.3
9	Septicemia (A40-A41)	357	1.3	4.7
10	Influenza and pneumonia (J10-J18)	334	1.2	4.4
11	Viral hepatitis (B15-B19)	325	1.2	4.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	270	1.0	3.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	147	0.5	1.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	0.5	1.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	0.4	1.6
...	All other causes (Residual)	4,314	15.8	56.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 55-59 years				
...	All causes	37,540	100.0	528.4
1	Malignant neoplasms (C00-C97)	16,250	43.3	228.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,021	16.0	84.8
3	Chronic lower respiratory diseases (J40-J47)	1,934	5.2	27.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,613	4.3	22.7
5	Diabetes mellitus (E10-E14)	1,338	3.6	18.8
6	Cerebrovascular diseases (I60-I69)	1,242	3.3	17.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	886	2.4	12.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	634	1.7	8.9
9	Septicemia (A40-A41)	544	1.4	7.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	459	1.2	6.5
11	Influenza and pneumonia (J10-J18)	401	1.1	5.6
12	Viral hepatitis (B15-B19)	229	0.6	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	201	0.5	2.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	186	0.5	2.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	154	0.4	2.2
...	All other causes (Residual)	5,448	14.5	76.7
White, non-Hispanic, female, 60-64 years				
...	All causes	45,999	100.0	851.6
1	Malignant neoplasms (C00-C97)	19,831	43.1	367.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,131	17.7	150.5
3	Chronic lower respiratory diseases (J40-J47)	3,325	7.2	61.6
4	Cerebrovascular diseases (I60-I69)	1,701	3.7	31.5
5	Diabetes mellitus (E10-E14)	1,574	3.4	29.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,193	2.6	22.1
7	Septicemia (A40-A41)	723	1.6	13.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	723	1.6	13.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	588	1.3	10.9
10	Influenza and pneumonia (J10-J18)	539	1.2	10.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	404	0.9	7.5
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	255	0.6	4.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	254	0.6	4.7
14	Alzheimer's disease (G30)	195	0.4	3.6
15	Aortic aneurysm and dissection (I71)	162	0.4	3.0
...	All other causes (Residual)	6,401	13.9	118.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 65-69 years				
...	All causes	56,192	100.0	1,306.2
1	Malignant neoplasms (C00-C97)	22,921	40.8	532.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,368	18.5	241.0
3	Chronic lower respiratory diseases (J40-J47)	5,262	9.4	122.3
4	Cerebrovascular diseases (I60-I69)	2,289	4.1	53.2
5	Diabetes mellitus (E10-E14)	1,930	3.4	44.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,134	2.0	26.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	900	1.6	20.9
8	Septicemia (A40-A41)	899	1.6	20.9
9	Influenza and pneumonia (J10-J18)	764	1.4	17.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	700	1.2	16.3
11	Alzheimer's disease (G30)	433	0.8	10.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	366	0.7	8.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	321	0.6	7.5
14	Aortic aneurysm and dissection (I71)	284	0.5	6.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	217	0.4	5.0
...	All other causes (Residual)	7,404	13.2	172.1
White, non-Hispanic, female, 70-74 years				
...	All causes	78,227	100.0	2,107.8
1	Malignant neoplasms (C00-C97)	27,665	35.4	745.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,006	20.5	431.3
3	Chronic lower respiratory diseases (J40-J47)	7,589	9.7	204.5
4	Cerebrovascular diseases (I60-I69)	4,044	5.2	109.0
5	Diabetes mellitus (E10-E14)	2,545	3.3	68.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,447	1.8	39.0
7	Alzheimer's disease (G30)	1,358	1.7	36.6
8	Influenza and pneumonia (J10-J18)	1,289	1.6	34.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,285	1.6	34.6
10	Septicemia (A40-A41)	1,245	1.6	33.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	780	1.0	21.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	601	0.8	16.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	501	0.6	13.5
14	Parkinson's disease (G20-G21)	497	0.6	13.4
15	Aortic aneurysm and dissection (I71)	488	0.6	13.1
...	All other causes (Residual)	10,887	13.9	293.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 75-79 years				
...	All causes	118,070	100.0	3,400.4
1	Malignant neoplasms (C00-C97)	33,256	28.2	957.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,507	23.3	792.2
3	Chronic lower respiratory diseases (J40-J47)	10,449	8.8	300.9
4	Cerebrovascular diseases (I60-I69)	7,692	6.5	221.5
5	Alzheimer's disease (G30)	3,900	3.3	112.3
6	Diabetes mellitus (E10-E14)	3,580	3.0	103.1
7	Influenza and pneumonia (J10-J18)	2,401	2.0	69.1
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,267	1.9	65.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,240	1.9	64.5
10	Septicemia (A40-A41)	1,932	1.6	55.6
11	Parkinson's disease (G20-G21)	1,197	1.0	34.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,101	0.9	31.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	827	0.7	23.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	779	0.7	22.4
15	Aortic aneurysm and dissection (I71)	766	0.6	22.1
...	All other causes (Residual)	18,176	15.4	523.5
White, non-Hispanic, female, 80-84 years				
...	All causes	167,353	100.0	5,718.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,167	27.0	1,543.5
2	Malignant neoplasms (C00-C97)	33,574	20.1	1,147.3
3	Cerebrovascular diseases (I60-I69)	13,163	7.9	449.8
4	Chronic lower respiratory diseases (J40-J47)	11,874	7.1	405.8
5	Alzheimer's disease (G30)	9,010	5.4	307.9
6	Diabetes mellitus (E10-E14)	4,596	2.7	157.1
7	Influenza and pneumonia (J10-J18)	4,503	2.7	153.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,407	2.0	116.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,263	1.9	111.5
10	Septicemia (A40-A41)	2,634	1.6	90.0
11	Parkinson's disease (G20-G21)	2,065	1.2	70.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,821	1.1	62.2
13	Pneumonitis due to solids and liquids (J69)	1,266	0.8	43.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,137	0.7	38.9
15	Aortic aneurysm and dissection (I71)	1,013	0.6	34.6
...	All other causes (Residual)	28,860	17.2	986.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 85-89 years				
...	All causes	184,406	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,628	31.3	---
2	Malignant neoplasms (C00-C97)	24,636	13.4	---
3	Cerebrovascular diseases (I60-I69)	16,008	8.7	---
4	Alzheimer's disease (G30)	12,974	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	9,706	5.3	---
6	Influenza and pneumonia (J10-J18)	5,741	3.1	---
7	Diabetes mellitus (E10-E14)	4,134	2.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,726	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,586	1.9	---
10	Septicemia (A40-A41)	2,634	1.4	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,405	1.3	---
12	Parkinson's disease (G20-G21)	2,030	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,785	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,217	0.7	---
15	Atherosclerosis (I70)	1,024	0.6	---
...	All other causes (Residual)	35,172	19.1	---
White, non-Hispanic, female, 90-94 years				
...	All causes	144,249	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,498	35.0	---
2	Malignant neoplasms (C00-C97)	12,511	8.7	---
3	Cerebrovascular diseases (I60-I69)	12,362	8.6	---
4	Alzheimer's disease (G30)	11,705	8.1	---
5	Influenza and pneumonia (J10-J18)	5,386	3.7	---
6	Chronic lower respiratory diseases (J40-J47)	5,223	3.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,947	2.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,638	1.8	---
9	Diabetes mellitus (E10-E14)	2,414	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,119	1.5	---
11	Septicemia (A40-A41)	1,721	1.2	---
12	Pneumonitis due to solids and liquids (J69)	1,427	1.0	---
13	Atherosclerosis (I70)	1,155	0.8	---
14	Parkinson's disease (G20-G21)	1,101	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	794	0.6	---
...	All other causes (Residual)	30,248	21.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
White, non-Hispanic, female, 95-99 years				
...	All causes	62,804	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,751	37.8	---
2	Alzheimer's disease (G30)	5,155	8.2	---
3	Cerebrovascular diseases (I60-I69)	5,147	8.2	---
4	Malignant neoplasms (C00-C97)	3,557	5.7	---
5	Influenza and pneumonia (J10-J18)	2,757	4.4	---
6	Chronic lower respiratory diseases (J40-J47)	1,720	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,218	1.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,149	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	997	1.6	---
10	Diabetes mellitus (E10-E14)	828	1.3	---
11	Pneumonitis due to solids and liquids (J69)	626	1.0	---
12	Atherosclerosis (I70)	621	1.0	---
13	Septicemia (A40-A41)	605	1.0	---
14	Parkinson's disease (G20-G21)	297	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	259	0.4	---
...	All other causes (Residual)	14,117	22.5	---
White, non-Hispanic, female, 100 years and over				
...	All causes	13,643	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,536	40.6	---
2	Alzheimer's disease (G30)	945	6.9	---
3	Cerebrovascular diseases (I60-I69)	943	6.9	---
4	Influenza and pneumonia (J10-J18)	748	5.5	---
5	Malignant neoplasms (C00-C97)	473	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	309	2.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	2.0	---
8	Chronic lower respiratory diseases (J40-J47)	247	1.8	---
9	Atherosclerosis (I70)	205	1.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	1.3	---
11	Diabetes mellitus (E10-E14)	151	1.1	---
12	Pneumonitis due to solids and liquids (J69)	128	0.9	---
13	Septicemia (A40-A41)	88	0.6	---
14	Anemias (D50-D64)	45	0.3	---
15	Nutritional deficiencies (E40-E64)	41	0.3	---
...	All other causes (Residual)	3,340	24.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, all ages				
...	All causes	286,581	100.0	759.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	71,461	24.9	189.3
2	Malignant neoplasms (C00-C97)	62,475	21.8	165.5
3	Cerebrovascular diseases (I60-I69)	16,882	5.9	44.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,684	4.8	36.2
5	Diabetes mellitus (E10-E14)	12,671	4.4	33.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,902	3.1	23.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,334	2.9	22.1
8	Chronic lower respiratory diseases (J40-J47)	7,657	2.7	20.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	6,767	2.4	17.9
10	Septicemia (A40-A41)	6,045	2.1	16.0
11	Influenza and pneumonia (J10-J18)	5,242	1.8	13.9
12	Certain conditions originating in the perinatal period (P00-P96)	4,906	1.7	13.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,851	1.7	12.8
14	Alzheimer's disease (G30)	4,422	1.5	11.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,310	0.8	6.1
...	All other causes (Residual)	49,972	17.4	132.4
Black, non-Hispanic, both sexes, 1-4 years				
...	All causes	1,108	100.0	44.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	355	32.0	14.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	13.5	6.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	9.6	4.2
4	Malignant neoplasms (C00-C97)	63	5.7	2.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	4.0	1.8
6	Influenza and pneumonia (J10-J18)	34	3.1	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	22	2.0	0.9
8	Septicemia (A40-A41)	19	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	15	1.4	*
10	Anemias (D50-D64)	12	1.1	*
11	Cerebrovascular diseases (I60-I69)	11	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	256	23.1	10.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 5-9 years				
...	All causes	616	100.0	20.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	250	40.6	8.3
2	Malignant neoplasms (C00-C97)	70	11.4	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	8.1	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	5.7	1.2
5	Chronic lower respiratory diseases (J40-J47)	27	4.4	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.1	*
7	Cerebrovascular diseases (I60-I69)	11	1.8	*
8	Septicemia (A40-A41)	10	1.6	*
9	Anemias (D50-D64)	8	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.6	*
12	Meningitis (G00,G03)	3	0.5	*
12	Influenza and pneumonia (J10-J18)	3	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
15	Meningococcal infection (A39)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Hernia (K40-K46)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	110	17.9	3.7
Black, non-Hispanic, both sexes, 10-14 years				
...	All causes	745	100.0	23.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	210	28.2	6.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	13.0	3.0
3	Malignant neoplasms (C00-C97)	70	9.4	2.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	5.9	1.4
5	Chronic lower respiratory diseases (J40-J47)	39	5.2	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	4.7	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	4.3	1.0
8	Anemias (D50-D64)	11	1.5	*
9	Septicemia (A40-A41)	10	1.3	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.2	*
10	Cerebrovascular diseases (I60-I69)	9	1.2	*
13	Diabetes mellitus (E10-E14)	8	1.1	*
14	Influenza and pneumonia (J10-J18)	6	0.8	*
15	Meningitis (G00,G03)	5	0.7	*
...	All other causes (Residual)	151	20.3	4.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 15-19 years				
...	All causes	2,890	100.0	87.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,287	44.5	38.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	734	25.4	22.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	143	4.9	4.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	3.7	3.3
5	Malignant neoplasms (C00-C97)	104	3.6	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	1.6	1.4
7	Chronic lower respiratory diseases (J40-J47)	36	1.2	1.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	30	1.0	0.9
9	Anemias (D50-D64)	22	0.8	0.7
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.5	*
10	Diabetes mellitus (E10-E14)	15	0.5	*
12	Cerebrovascular diseases (I60-I69)	13	0.4	*
13	Septicemia (A40-A41)	11	0.4	*
13	Influenza and pneumonia (J10-J18)	11	0.4	*
15	Legal intervention (Y35,Y89.0)	10	0.3	*
...	All other causes (Residual)	304	10.5	9.2
Black, non-Hispanic, both sexes, 20-24 years				
...	All causes	4,457	100.0	146.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,966	44.1	64.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	969	21.7	31.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	264	5.9	8.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	4.4	6.4
5	Malignant neoplasms (C00-C97)	155	3.5	5.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	113	2.5	3.7
7	Pregnancy, childbirth and the puerperium (O00-099)	48	1.1	1.6
8	Anemias (D50-D64)	42	0.9	1.4
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	1.3
10	Diabetes mellitus (E10-E14)	39	0.9	1.3
11	Chronic lower respiratory diseases (J40-J47)	38	0.9	1.3
12	Cerebrovascular diseases (I60-I69)	31	0.7	1.0
13	Septicemia (A40-A41)	30	0.7	1.0
13	Influenza and pneumonia (J10-J18)	30	0.7	1.0
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.5	0.8
...	All other causes (Residual)	475	10.7	15.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 25-29 years				
...	All causes	4,814	100.0	171.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,586	32.9	56.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	981	20.4	34.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	375	7.8	13.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	293	6.1	10.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	251	5.2	8.9
6	Malignant neoplasms (C00-C97)	237	4.9	8.4
7	Diabetes mellitus (E10-E14)	97	2.0	3.5
8	Pregnancy, childbirth and the puerperium (O00-099)	66	1.4	2.3
9	Anemias (D50-D64)	57	1.2	2.0
10	Cerebrovascular diseases (I60-I69)	51	1.1	1.8
11	Chronic lower respiratory diseases (J40-J47)	45	0.9	1.6
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	0.9	1.6
13	Septicemia (A40-A41)	41	0.9	1.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	0.8	1.4
15	Influenza and pneumonia (J10-J18)	31	0.6	1.1
...	All other causes (Residual)	619	12.9	22.0
Black, non-Hispanic, both sexes, 30-34 years				
...	All causes	5,106	100.0	201.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,049	20.5	41.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	922	18.1	36.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	645	12.6	25.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	472	9.2	18.6
5	Malignant neoplasms (C00-C97)	399	7.8	15.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	215	4.2	8.5
7	Diabetes mellitus (E10-E14)	140	2.7	5.5
8	Cerebrovascular diseases (I60-I69)	98	1.9	3.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	1.3	2.6
10	Chronic lower respiratory diseases (J40-J47)	61	1.2	2.4
11	Anemias (D50-D64)	54	1.1	2.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	50	1.0	2.0
13	Pregnancy, childbirth and the puerperium (O00-099)	49	1.0	1.9
14	Septicemia (A40-A41)	47	0.9	1.9
15	Influenza and pneumonia (J10-J18)	46	0.9	1.8
...	All other causes (Residual)	792	15.5	31.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 35-39 years				
...	All causes	6,954	100.0	262.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,124	16.2	42.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,039	14.9	39.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	877	12.6	33.1
4	Malignant neoplasms (C00-C97)	776	11.2	29.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	739	10.6	27.9
6	Cerebrovascular diseases (I60-I69)	233	3.4	8.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	224	3.2	8.5
8	Diabetes mellitus (E10-E14)	216	3.1	8.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	111	1.6	4.2
10	Septicemia (A40-A41)	105	1.5	4.0
11	Chronic lower respiratory diseases (J40-J47)	72	1.0	2.7
12	Influenza and pneumonia (J10-J18)	66	0.9	2.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	66	0.9	2.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	61	0.9	2.3
15	Anemias (D50-D64)	60	0.9	2.3
...	All other causes (Residual)	1,185	17.0	44.7
Black, non-Hispanic, both sexes, 40-44 years				
...	All causes	10,746	100.0	384.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,913	17.8	68.5
2	Malignant neoplasms (C00-C97)	1,719	16.0	61.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	1,282	11.9	45.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,268	11.8	45.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	592	5.5	21.2
6	Cerebrovascular diseases (I60-I69)	458	4.3	16.4
7	Diabetes mellitus (E10-E14)	362	3.4	13.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	222	2.1	7.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	185	1.7	6.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	182	1.7	6.5
11	Septicemia (A40-A41)	165	1.5	5.9
12	Chronic lower respiratory diseases (J40-J47)	152	1.4	5.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	147	1.4	5.3
14	Influenza and pneumonia (J10-J18)	117	1.1	4.2
15	Anemias (D50-D64)	83	0.8	3.0
...	All other causes (Residual)	1,899	17.7	68.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 45-49 years				
...	All causes	16,058	100.0	590.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,515	21.9	129.3
2	Malignant neoplasms (C00-C97)	3,451	21.5	126.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,497	9.3	55.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,371	8.5	50.4
5	Cerebrovascular diseases (I60-I69)	761	4.7	28.0
6	Diabetes mellitus (E10-E14)	532	3.3	19.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	475	3.0	17.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	308	1.9	11.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	304	1.9	11.2
10	Septicemia (A40-A41)	280	1.7	10.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	215	1.3	7.9
12	Chronic lower respiratory diseases (J40-J47)	213	1.3	7.8
13	Influenza and pneumonia (J10-J18)	198	1.2	7.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	1.0	6.2
15	Viral hepatitis (B15-B19)	132	0.8	4.9
...	All other causes (Residual)	2,638	16.4	97.0
Black, non-Hispanic, both sexes, 50-54 years				
...	All causes	20,932	100.0	914.1
1	Malignant neoplasms (C00-C97)	5,262	25.1	229.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,005	23.9	218.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,320	6.3	57.6
4	Cerebrovascular diseases (I60-I69)	1,137	5.4	49.7
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,046	5.0	45.7
6	Diabetes mellitus (E10-E14)	870	4.2	38.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	528	2.5	23.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	488	2.3	21.3
9	Chronic lower respiratory diseases (J40-J47)	401	1.9	17.5
10	Septicemia (A40-A41)	393	1.9	17.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	350	1.7	15.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	337	1.6	14.7
13	Viral hepatitis (B15-B19)	259	1.2	11.3
14	Influenza and pneumonia (J10-J18)	251	1.2	11.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	0.6	5.9
...	All other causes (Residual)	3,150	15.0	137.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 55-59 years				
...	All causes	23,553	100.0	1,260.3
1	Malignant neoplasms (C00-C97)	6,860	29.1	367.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,985	25.4	320.2
3	Cerebrovascular diseases (I60-I69)	1,293	5.5	69.2
4	Diabetes mellitus (E10-E14)	1,178	5.0	63.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	980	4.2	52.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	686	2.9	36.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	624	2.6	33.4
8	Chronic lower respiratory diseases (J40-J47)	527	2.2	28.2
9	Septicemia (A40-A41)	494	2.1	26.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	439	1.9	23.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	364	1.5	19.5
12	Viral hepatitis (B15-B19)	276	1.2	14.8
13	Influenza and pneumonia (J10-J18)	246	1.0	13.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	0.9	10.9
15	Aortic aneurysm and dissection (I71)	115	0.5	6.2
...	All other causes (Residual)	3,282	13.9	175.6
Black, non-Hispanic, both sexes, 60-64 years				
...	All causes	22,408	100.0	1,789.0
1	Malignant neoplasms (C00-C97)	7,188	32.1	573.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,957	26.6	475.6
3	Cerebrovascular diseases (I60-I69)	1,299	5.8	103.7
4	Diabetes mellitus (E10-E14)	1,297	5.8	103.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	710	3.2	56.7
6	Chronic lower respiratory diseases (J40-J47)	663	3.0	52.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	564	2.5	45.0
8	Septicemia (A40-A41)	492	2.2	39.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	366	1.6	29.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	276	1.2	22.0
11	Influenza and pneumonia (J10-J18)	261	1.2	20.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	261	1.2	20.8
13	Viral hepatitis (B15-B19)	115	0.5	9.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	101	0.5	8.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	0.4	8.0
...	All other causes (Residual)	2,758	12.3	220.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 65-69 years				
...	All causes	23,929	100.0	2,454.3
1	Malignant neoplasms (C00-C97)	7,664	32.0	786.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,390	26.7	655.4
3	Diabetes mellitus (E10-E14)	1,447	6.0	148.4
4	Cerebrovascular diseases (I60-I69)	1,423	5.9	145.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	860	3.6	88.2
6	Chronic lower respiratory diseases (J40-J47)	831	3.5	85.2
7	Septicemia (A40-A41)	553	2.3	56.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	449	1.9	46.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	425	1.8	43.6
10	Influenza and pneumonia (J10-J18)	395	1.7	40.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	206	0.9	21.1
12	Human immunodeficiency virus (HIV) disease (B20-B24)	166	0.7	17.0
13	Aortic aneurysm and dissection (I71)	129	0.5	13.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.4	9.3
14	Pneumonitis due to solids and liquids (J69)	91	0.4	9.3
...	All other causes (Residual)	2,809	11.7	288.1
Black, non-Hispanic, both sexes, 70-74 years				
...	All causes	26,645	100.0	3,432.1
1	Malignant neoplasms (C00-C97)	7,933	29.8	1,021.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,378	27.7	950.3
3	Cerebrovascular diseases (I60-I69)	1,753	6.6	225.8
4	Diabetes mellitus (E10-E14)	1,530	5.7	197.1
5	Chronic lower respiratory diseases (J40-J47)	971	3.6	125.1
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	971	3.6	125.1
7	Septicemia (A40-A41)	677	2.5	87.2
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	453	1.7	58.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	404	1.5	52.0
10	Influenza and pneumonia (J10-J18)	401	1.5	51.7
11	Alzheimer's disease (G30)	259	1.0	33.4
12	Pneumonitis due to solids and liquids (J69)	156	0.6	20.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	127	0.5	16.4
14	Aortic aneurysm and dissection (I71)	126	0.5	16.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	117	0.4	15.1
...	All other causes (Residual)	3,389	12.7	436.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 75-79 years				
...	All causes	28,976	100.0	4,987.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,153	28.1	1,403.3
2	Malignant neoplasms (C00-C97)	7,589	26.2	1,306.2
3	Cerebrovascular diseases (I60-I69)	2,068	7.1	355.9
4	Diabetes mellitus (E10-E14)	1,612	5.6	277.5
5	Chronic lower respiratory diseases (J40-J47)	1,184	4.1	203.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,053	3.6	181.2
7	Septicemia (A40-A41)	742	2.6	127.7
8	Influenza and pneumonia (J10-J18)	589	2.0	101.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	556	1.9	95.7
10	Alzheimer's disease (G30)	537	1.9	92.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	421	1.5	72.5
12	Pneumonitis due to solids and liquids (J69)	195	0.7	33.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	155	0.5	26.7
14	Parkinson's disease (G20-G21)	148	0.5	25.5
15	Aortic aneurysm and dissection (I71)	140	0.5	24.1
...	All other causes (Residual)	3,834	13.2	659.9
Black, non-Hispanic, both sexes, 80-84 years				
...	All causes	30,440	100.0	7,380.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,031	29.7	2,189.6
2	Malignant neoplasms (C00-C97)	6,521	21.4	1,581.1
3	Cerebrovascular diseases (I60-I69)	2,330	7.7	564.9
4	Diabetes mellitus (E10-E14)	1,531	5.0	371.2
5	Chronic lower respiratory diseases (J40-J47)	1,100	3.6	266.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,071	3.5	259.7
7	Alzheimer's disease (G30)	946	3.1	229.4
8	Influenza and pneumonia (J10-J18)	814	2.7	197.4
9	Septicemia (A40-A41)	781	2.6	189.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	625	2.1	151.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	394	1.3	95.5
12	Pneumonitis due to solids and liquids (J69)	231	0.8	56.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	165	0.5	40.0
14	Parkinson's disease (G20-G21)	156	0.5	37.8
15	Aortic aneurysm and dissection (I71)	141	0.5	34.2
...	All other causes (Residual)	4,603	15.1	1,116.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 85-89 years				
...	All causes	23,671	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,327	31.0	---
2	Malignant neoplasms (C00-C97)	3,936	16.6	---
3	Cerebrovascular diseases (I60-I69)	1,943	8.2	---
4	Alzheimer's disease (G30)	1,075	4.5	---
5	Diabetes mellitus (E10-E14)	1,065	4.5	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	868	3.7	---
7	Chronic lower respiratory diseases (J40-J47)	739	3.1	---
8	Influenza and pneumonia (J10-J18)	707	3.0	---
9	Septicemia (A40-A41)	576	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	530	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	285	1.2	---
12	Pneumonitis due to solids and liquids (J69)	233	1.0	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	131	0.6	---
14	Parkinson's disease (G20-G21)	121	0.5	---
14	Atherosclerosis (I70)	121	0.5	---
...	All other causes (Residual)	4,014	17.0	---
Black, non-Hispanic, both sexes, 90-94 years				
...	All causes	15,413	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,056	32.8	---
2	Malignant neoplasms (C00-C97)	1,829	11.9	---
3	Cerebrovascular diseases (I60-I69)	1,259	8.2	---
4	Alzheimer's disease (G30)	925	6.0	---
5	Influenza and pneumonia (J10-J18)	576	3.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	564	3.7	---
7	Diabetes mellitus (E10-E14)	547	3.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	395	2.6	---
9	Septicemia (A40-A41)	360	2.3	---
10	Chronic lower respiratory diseases (J40-J47)	358	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	1.2	---
12	Pneumonitis due to solids and liquids (J69)	174	1.1	---
13	Atherosclerosis (I70)	88	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.5	---
15	Aortic aneurysm and dissection (I71)	59	0.4	---
...	All other causes (Residual)	2,967	19.2	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, both sexes, 95-99 years				
...	All causes	6,555	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,283	34.8	---
2	Malignant neoplasms (C00-C97)	550	8.4	---
3	Cerebrovascular diseases (I60-I69)	505	7.7	---
4	Alzheimer's disease (G30)	437	6.7	---
5	Influenza and pneumonia (J10-J18)	276	4.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	219	3.3	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	194	3.0	---
8	Diabetes mellitus (E10-E14)	148	2.3	---
9	Chronic lower respiratory diseases (J40-J47)	138	2.1	---
10	Septicemia (A40-A41)	129	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	89	1.4	---
12	Pneumonitis due to solids and liquids (J69)	62	0.9	---
13	Atherosclerosis (I70)	51	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.5	---
15	Anemias (D50-D64)	29	0.4	---
...	All other causes (Residual)	1,415	21.6	---
Black, non-Hispanic, both sexes, 100 years and over				
...	All causes	2,039	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	795	39.0	---
2	Cerebrovascular diseases (I60-I69)	147	7.2	---
3	Alzheimer's disease (G30)	104	5.1	---
4	Influenza and pneumonia (J10-J18)	97	4.8	---
5	Malignant neoplasms (C00-C97)	91	4.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	3.3	---
8	Chronic lower respiratory diseases (J40-J47)	36	1.8	---
9	Diabetes mellitus (E10-E14)	33	1.6	---
10	Septicemia (A40-A41)	31	1.5	---
11	Pneumonitis due to solids and liquids (J69)	24	1.2	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	21	1.0	---
13	Atherosclerosis (I70)	14	0.7	---
14	Nutritional deficiencies (E40-E64)	10	0.5	---
15	Anemias (D50-D64)	8	0.4	---
...	All other causes (Residual)	489	24.0	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, all ages				
...	All causes	146,729	100.0	815.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,787	24.4	198.8
2	Malignant neoplasms (C00-C97)	32,253	22.0	179.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,443	6.4	52.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,566	5.2	42.0
5	Cerebrovascular diseases (I60-I69)	7,349	5.0	40.8
6	Diabetes mellitus (E10-E14)	5,701	3.9	31.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	4,384	3.0	24.4
8	Chronic lower respiratory diseases (J40-J47)	4,091	2.8	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,782	2.6	21.0
10	Certain conditions originating in the perinatal period (P00-P96)	2,706	1.8	15.0
11	Septicemia (A40-A41)	2,688	1.8	14.9
12	Influenza and pneumonia (J10-J18)	2,449	1.7	13.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,100	1.4	11.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,630	1.1	9.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,485	1.0	8.3
...	All other causes (Residual)	23,315	15.9	129.5
Black, non-Hispanic, male, 1-4 years				
...	All causes	612	100.0	48.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	33.8	16.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	13.2	6.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	6.7	3.2
4	Malignant neoplasms (C00-C97)	34	5.6	2.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	4.4	2.1
6	Influenza and pneumonia (J10-J18)	16	2.6	*
7	Chronic lower respiratory diseases (J40-J47)	12	2.0	*
8	Septicemia (A40-A41)	11	1.8	*
9	Anemias (D50-D64)	9	1.5	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.5	*
11	Cerebrovascular diseases (I60-I69)	7	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
15	Meningococcal infection (A39)	1	0.2	*
15	Syphilis (A50-A53)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	143	23.4	11.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 5-9 years				
...	All causes	361	100.0	23.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	42.1	10.0
2	Malignant neoplasms (C00-C97)	39	10.8	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	9.4	2.2
4	Chronic lower respiratory diseases (J40-J47)	14	3.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	3.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.5	*
7	Septicemia (A40-A41)	7	1.9	*
8	Anemias (D50-D64)	5	1.4	*
8	Cerebrovascular diseases (I60-I69)	5	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.1	*
11	Meningitis (G00,G03)	2	0.6	*
12	Meningococcal infection (A39)	1	0.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
12	Diabetes mellitus (E10-E14)	1	0.3	*
12	Pneumonitis due to solids and liquids (J69)	1	0.3	*
12	Diseases of appendix (K35-K38)	1	0.3	*
12	Hernia (K40-K46)	1	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
...	All other causes (Residual)	68	18.8	4.5
Black, non-Hispanic, male, 10-14 years				
...	All causes	453	100.0	27.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	30.0	8.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	15.5	4.3
3	Malignant neoplasms (C00-C97)	41	9.1	2.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	6.2	1.7
5	Chronic lower respiratory diseases (J40-J47)	27	6.0	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	5.5	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	3.3	*
8	Septicemia (A40-A41)	6	1.3	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.1	*
10	Anemias (D50-D64)	4	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
11	Influenza and pneumonia (J10-J18)	3	0.7	*
13	Cerebrovascular diseases (I60-I69)	2	0.4	*
13	Pneumonitis due to solids and liquids (J69)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	84	18.5	5.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 15-19 years				
...	All causes	2,265	100.0	134.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,157	51.1	68.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	564	24.9	33.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	121	5.3	7.2
4	Malignant neoplasms (C00-C97)	66	2.9	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	65	2.9	3.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	1.2	1.7
7	Chronic lower respiratory diseases (J40-J47)	24	1.1	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.8	*
9	Anemias (D50-D64)	12	0.5	*
10	Legal intervention (Y35,Y89.0)	10	0.4	*
11	Diabetes mellitus (E10-E14)	8	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.3	*
13	Septicemia (A40-A41)	6	0.3	*
14	Cerebrovascular diseases (I60-I69)	5	0.2	*
14	Influenza and pneumonia (J10-J18)	5	0.2	*
...	All other causes (Residual)	170	7.5	10.1
Black, non-Hispanic, male, 20-24 years				
...	All causes	3,426	100.0	223.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,789	52.2	116.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	729	21.3	47.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	230	6.7	15.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	115	3.4	7.5
5	Malignant neoplasms (C00-C97)	85	2.5	5.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	66	1.9	4.3
7	Anemias (D50-D64)	25	0.7	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.7	1.6
9	Diabetes mellitus (E10-E14)	23	0.7	1.5
10	Chronic lower respiratory diseases (J40-J47)	22	0.6	1.4
11	Influenza and pneumonia (J10-J18)	18	0.5	*
12	Septicemia (A40-A41)	15	0.4	*
12	Cerebrovascular diseases (I60-I69)	15	0.4	*
14	Legal intervention (Y35,Y89.0)	14	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.3	*
...	All other causes (Residual)	247	7.2	16.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 25-29 years				
...	All causes	3,463	100.0	253.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,391	40.2	101.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	764	22.1	56.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	242	7.0	17.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	222	6.4	16.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	163	4.7	11.9
6	Malignant neoplasms (C00-C97)	109	3.1	8.0
7	Diabetes mellitus (E10-E14)	50	1.4	3.7
8	Anemias (D50-D64)	30	0.9	2.2
9	Cerebrovascular diseases (I60-I69)	28	0.8	2.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	0.8	1.9
11	Chronic lower respiratory diseases (J40-J47)	25	0.7	1.8
11	Legal intervention (Y35,Y89.0)	25	0.7	1.8
13	Influenza and pneumonia (J10-J18)	21	0.6	1.5
14	Septicemia (A40-A41)	18	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.5	*
...	All other causes (Residual)	331	9.6	24.2
Black, non-Hispanic, male, 30-34 years				
...	All causes	3,365	100.0	279.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	924	27.5	76.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	716	21.3	59.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	411	12.2	34.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	253	7.5	21.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	5.7	16.0
6	Malignant neoplasms (C00-C97)	162	4.8	13.5
7	Diabetes mellitus (E10-E14)	85	2.5	7.1
8	Cerebrovascular diseases (I60-I69)	47	1.4	3.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	1.1	3.2
10	Chronic lower respiratory diseases (J40-J47)	36	1.1	3.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	0.8	2.2
12	Septicemia (A40-A41)	26	0.8	2.2
13	Anemias (D50-D64)	25	0.7	2.1
14	Legal intervention (Y35,Y89.0)	19	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.5	*
...	All other causes (Residual)	387	11.5	32.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 35-39 years				
...	All causes	4,202	100.0	336.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	758	18.0	60.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	711	16.9	57.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	598	14.2	47.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	521	12.4	41.8
5	Malignant neoplasms (C00-C97)	287	6.8	23.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	4.6	15.4
7	Cerebrovascular diseases (I60-I69)	129	3.1	10.3
8	Diabetes mellitus (E10-E14)	128	3.0	10.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	58	1.4	4.6
10	Septicemia (A40-A41)	46	1.1	3.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	39	0.9	3.1
12	Influenza and pneumonia (J10-J18)	37	0.9	3.0
13	Chronic lower respiratory diseases (J40-J47)	33	0.8	2.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	32	0.8	2.6
15	Anemias (D50-D64)	25	0.6	2.0
...	All other causes (Residual)	608	14.5	48.7
Black, non-Hispanic, male, 40-44 years				
...	All causes	6,161	100.0	471.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,153	18.7	88.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	879	14.3	67.2
3	Human immunodeficiency virus (HIV) disease (B20-B24)	815	13.2	62.3
4	Malignant neoplasms (C00-C97)	686	11.1	52.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	454	7.4	34.7
6	Diabetes mellitus (E10-E14)	215	3.5	16.4
7	Cerebrovascular diseases (I60-I69)	210	3.4	16.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	150	2.4	11.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	123	2.0	9.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	119	1.9	9.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	87	1.4	6.7
12	Septicemia (A40-A41)	86	1.4	6.6
13	Chronic lower respiratory diseases (J40-J47)	69	1.1	5.3
14	Influenza and pneumonia (J10-J18)	58	0.9	4.4
15	Aortic aneurysm and dissection (I71)	43	0.7	3.3
...	All other causes (Residual)	1,014	16.5	77.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 45-49 years				
...	All causes	9,265	100.0	732.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,222	24.0	175.8
2	Malignant neoplasms (C00-C97)	1,563	16.9	123.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,059	11.4	83.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	880	9.5	69.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	380	4.1	30.1
6	Cerebrovascular diseases (I60-I69)	377	4.1	29.8
7	Diabetes mellitus (E10-E14)	307	3.3	24.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	193	2.1	15.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	178	1.9	14.1
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	1.5	10.9
11	Septicemia (A40-A41)	131	1.4	10.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	120	1.3	9.5
13	Influenza and pneumonia (J10-J18)	106	1.1	8.4
14	Chronic lower respiratory diseases (J40-J47)	101	1.1	8.0
15	Viral hepatitis (B15-B19)	81	0.9	6.4
...	All other causes (Residual)	1,429	15.4	113.0
Black, non-Hispanic, male, 50-54 years				
...	All causes	12,564	100.0	1,199.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,212	25.6	306.5
2	Malignant neoplasms (C00-C97)	2,760	22.0	263.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	978	7.8	93.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	745	5.9	71.1
5	Cerebrovascular diseases (I60-I69)	655	5.2	62.5
6	Diabetes mellitus (E10-E14)	485	3.9	46.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	337	2.7	32.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	299	2.4	28.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	260	2.1	24.8
10	Septicemia (A40-A41)	219	1.7	20.9
11	Chronic lower respiratory diseases (J40-J47)	213	1.7	20.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	1.7	19.9
13	Viral hepatitis (B15-B19)	175	1.4	16.7
14	Influenza and pneumonia (J10-J18)	164	1.3	15.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	106	0.8	10.1
...	All other causes (Residual)	1,748	13.9	166.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 55-59 years				
...	All causes	14,020	100.0	1,662.0
1	Malignant neoplasms (C00-C97)	3,804	27.1	450.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,784	27.0	448.6
3	Cerebrovascular diseases (I60-I69)	768	5.5	91.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	707	5.0	83.8
5	Diabetes mellitus (E10-E14)	621	4.4	73.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	496	3.5	58.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	348	2.5	41.3
8	Chronic liver disease and cirrhosis (K70,K73-K74)	307	2.2	36.4
9	Chronic lower respiratory diseases (J40-J47)	287	2.0	34.0
10	Septicemia (A40-A41)	272	1.9	32.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	214	1.5	25.4
12	Viral hepatitis (B15-B19)	201	1.4	23.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	166	1.2	19.7
14	Influenza and pneumonia (J10-J18)	138	1.0	16.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	0.6	9.5
...	All other causes (Residual)	1,827	13.0	216.6
Black, non-Hispanic, male, 60-64 years				
...	All causes	12,909	100.0	2,355.8
1	Malignant neoplasms (C00-C97)	3,949	30.6	720.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,701	28.7	675.4
3	Cerebrovascular diseases (I60-I69)	743	5.8	135.6
4	Diabetes mellitus (E10-E14)	668	5.2	121.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	403	3.1	73.5
6	Chronic lower respiratory diseases (J40-J47)	392	3.0	71.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	362	2.8	66.1
8	Septicemia (A40-A41)	255	2.0	46.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	204	1.6	37.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	185	1.4	33.8
11	Influenza and pneumonia (J10-J18)	166	1.3	30.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	165	1.3	30.1
13	Viral hepatitis (B15-B19)	78	0.6	14.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	74	0.6	13.5
15	Aortic aneurysm and dissection (I71)	57	0.4	10.4
...	All other causes (Residual)	1,507	11.7	275.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 65-69 years				
...	All causes	13,202	100.0	3,190.8
1	Malignant neoplasms (C00-C97)	4,252	32.2	1,027.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,739	28.3	903.7
3	Cerebrovascular diseases (I60-I69)	756	5.7	182.7
4	Diabetes mellitus (E10-E14)	685	5.2	165.6
5	Chronic lower respiratory diseases (J40-J47)	474	3.6	114.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	419	3.2	101.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	292	2.2	70.6
8	Septicemia (A40-A41)	259	2.0	62.6
9	Influenza and pneumonia (J10-J18)	213	1.6	51.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	209	1.6	50.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	143	1.1	34.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	115	0.9	27.8
13	Aortic aneurysm and dissection (I71)	77	0.6	18.6
14	Pneumonitis due to solids and liquids (J69)	64	0.5	15.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.4	12.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	0.4	12.3
...	All other causes (Residual)	1,403	10.6	339.1
Black, non-Hispanic, male, 70-74 years				
...	All causes	14,063	100.0	4,451.9
1	Malignant neoplasms (C00-C97)	4,350	30.9	1,377.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,015	28.6	1,271.0
3	Cerebrovascular diseases (I60-I69)	869	6.2	275.1
4	Diabetes mellitus (E10-E14)	711	5.1	225.1
5	Chronic lower respiratory diseases (J40-J47)	545	3.9	172.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	438	3.1	138.7
7	Septicemia (A40-A41)	304	2.2	96.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	247	1.8	78.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	233	1.7	73.8
10	Influenza and pneumonia (J10-J18)	221	1.6	70.0
11	Alzheimer's disease (G30)	106	0.8	33.6
12	Pneumonitis due to solids and liquids (J69)	91	0.6	28.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	71	0.5	22.5
14	Aortic aneurysm and dissection (I71)	64	0.5	20.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	61	0.4	19.3
...	All other causes (Residual)	1,737	12.4	549.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 75-79 years				
...	All causes	13,993	100.0	6,441.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,989	28.5	1,836.4
2	Malignant neoplasms (C00-C97)	3,946	28.2	1,816.6
3	Cerebrovascular diseases (I60-I69)	884	6.3	407.0
4	Diabetes mellitus (E10-E14)	660	4.7	303.8
5	Chronic lower respiratory diseases (J40-J47)	639	4.6	294.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	467	3.3	215.0
7	Septicemia (A40-A41)	341	2.4	157.0
8	Influenza and pneumonia (J10-J18)	308	2.2	141.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	1.7	110.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	216	1.5	99.4
11	Alzheimer's disease (G30)	197	1.4	90.7
12	Pneumonitis due to solids and liquids (J69)	122	0.9	56.2
13	Parkinson's disease (G20-G21)	94	0.7	43.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.5	33.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	70	0.5	32.2
...	All other causes (Residual)	1,746	12.5	803.8
Black, non-Hispanic, male, 80-84 years				
...	All causes	13,052	100.0	9,165.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,794	29.1	2,664.4
2	Malignant neoplasms (C00-C97)	3,303	25.3	2,319.6
3	Cerebrovascular diseases (I60-I69)	840	6.4	589.9
4	Chronic lower respiratory diseases (J40-J47)	602	4.6	422.8
5	Diabetes mellitus (E10-E14)	542	4.2	380.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	454	3.5	318.8
7	Influenza and pneumonia (J10-J18)	347	2.7	243.7
8	Septicemia (A40-A41)	305	2.3	214.2
9	Alzheimer's disease (G30)	288	2.2	202.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	248	1.9	174.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	1.4	127.1
12	Pneumonitis due to solids and liquids (J69)	124	1.0	87.1
13	Parkinson's disease (G20-G21)	91	0.7	63.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	79	0.6	55.5
15	Atherosclerosis (I70)	57	0.4	40.0
...	All other causes (Residual)	1,797	13.8	1,262.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 85-89 years				
...	All causes	8,523	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,534	29.7	---
2	Malignant neoplasms (C00-C97)	1,846	21.7	---
3	Cerebrovascular diseases (I60-I69)	571	6.7	---
4	Chronic lower respiratory diseases (J40-J47)	367	4.3	---
5	Diabetes mellitus (E10-E14)	337	4.0	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	297	3.5	---
7	Alzheimer's disease (G30)	279	3.3	---
8	Influenza and pneumonia (J10-J18)	272	3.2	---
9	Septicemia (A40-A41)	195	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	163	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	124	1.5	---
12	Pneumonitis due to solids and liquids (J69)	110	1.3	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.7	---
14	Parkinson's disease (G20-G21)	52	0.6	---
15	Atherosclerosis (I70)	42	0.5	---
...	All other causes (Residual)	1,277	15.0	---
Black, non-Hispanic, male, 90-94 years				
...	All causes	4,364	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,380	31.6	---
2	Malignant neoplasms (C00-C97)	738	16.9	---
3	Cerebrovascular diseases (I60-I69)	302	6.9	---
4	Influenza and pneumonia (J10-J18)	188	4.3	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	175	4.0	---
6	Alzheimer's disease (G30)	163	3.7	---
7	Chronic lower respiratory diseases (J40-J47)	142	3.3	---
8	Diabetes mellitus (E10-E14)	139	3.2	---
9	Septicemia (A40-A41)	101	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	93	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	71	1.6	---
12	Pneumonitis due to solids and liquids (J69)	65	1.5	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.6	---
13	Atherosclerosis (I70)	25	0.6	---
15	Parkinson's disease (G20-G21)	23	0.5	---
...	All other causes (Residual)	734	16.8	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, male, 95-99 years				
...	All causes	1,422	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	474	33.3	---
2	Malignant neoplasms (C00-C97)	200	14.1	---
3	Cerebrovascular diseases (I60-I69)	87	6.1	---
4	Influenza and pneumonia (J10-J18)	76	5.3	---
5	Alzheimer's disease (G30)	72	5.1	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	3.9	---
7	Chronic lower respiratory diseases (J40-J47)	50	3.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	2.7	---
9	Diabetes mellitus (E10-E14)	31	2.2	---
10	Septicemia (A40-A41)	26	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	1.5	---
12	Atherosclerosis (I70)	14	1.0	---
13	Pneumonitis due to solids and liquids (J69)	13	0.9	---
14	Aortic aneurysm and dissection (I71)	7	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.4	---
15	Anemias (D50-D64)	6	0.4	---
...	All other causes (Residual)	245	17.2	---
Black, non-Hispanic, male, 100 years and over				
...	All causes	319	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	119	37.3	---
2	Malignant neoplasms (C00-C97)	29	9.1	---
3	Influenza and pneumonia (J10-J18)	23	7.2	---
4	Cerebrovascular diseases (I60-I69)	16	5.0	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	4.7	---
6	Chronic lower respiratory diseases (J40-J47)	11	3.4	---
7	Alzheimer's disease (G30)	8	2.5	---
8	Septicemia (A40-A41)	6	1.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	1.9	---
8	Pneumonitis due to solids and liquids (J69)	6	1.9	---
11	Diabetes mellitus (E10-E14)	4	1.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	1.3	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.6	---
13	Nutritional deficiencies (E40-E64)	2	0.6	---
13	Atherosclerosis (I70)	2	0.6	---
...	All other causes (Residual)	66	20.7	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, all ages				
...	All causes	139,852	100.0	708.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,674	25.5	180.6
2	Malignant neoplasms (C00-C97)	30,222	21.6	153.0
3	Cerebrovascular diseases (I60-I69)	9,533	6.8	48.3
4	Diabetes mellitus (E10-E14)	6,970	5.0	35.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,552	3.3	23.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,241	3.0	21.5
7	Chronic lower respiratory diseases (J40-J47)	3,566	2.5	18.1
8	Septicemia (A40-A41)	3,357	2.4	17.0
9	Alzheimer's disease (G30)	3,246	2.3	16.4
10	Influenza and pneumonia (J10-J18)	2,793	2.0	14.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,751	2.0	13.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	2,383	1.7	12.1
13	Certain conditions originating in the perinatal period (P00-P96)	2,200	1.6	11.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,336	1.0	6.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	828	0.6	4.2
...	All other causes (Residual)	26,200	18.7	132.6
Black, non-Hispanic, female, 1-4 years				
...	All causes	496	100.0	40.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	29.8	12.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	13.9	5.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	13.1	5.3
4	Malignant neoplasms (C00-C97)	29	5.8	2.4
5	Influenza and pneumonia (J10-J18)	18	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.4	*
7	Certain conditions originating in the perinatal period (P00-P96)	13	2.6	*
8	Septicemia (A40-A41)	8	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
9	Cerebrovascular diseases (I60-I69)	4	0.8	*
11	Anemias (D50-D64)	3	0.6	*
11	Meningitis (G00,G03)	3	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	3	0.6	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (Residual)	108	21.8	8.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 5-9 years				
...	All causes	255	100.0	17.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98	38.4	6.6
2	Malignant neoplasms (C00-C97)	31	12.2	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	8.2	1.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	6.3	*
5	Chronic lower respiratory diseases (J40-J47)	13	5.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	3.9	*
7	Cerebrovascular diseases (I60-I69)	6	2.4	*
8	Septicemia (A40-A41)	3	1.2	*
8	Anemias (D50-D64)	3	1.2	*
8	Influenza and pneumonia (J10-J18)	3	1.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	3	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.4	*
13	Meningitis (G00,G03)	1	0.4	*
13	Peptic ulcer (K25-K28)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	42	16.5	2.8
Black, non-Hispanic, female, 10-14 years				
...	All causes	292	100.0	18.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	25.3	4.7
2	Malignant neoplasms (C00-C97)	29	9.9	1.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	9.2	1.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	6.5	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	5.8	*
6	Chronic lower respiratory diseases (J40-J47)	12	4.1	*
7	Anemias (D50-D64)	7	2.4	*
7	Diabetes mellitus (E10-E14)	7	2.4	*
7	Cerebrovascular diseases (I60-I69)	7	2.4	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	2.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.1	*
12	Septicemia (A40-A41)	4	1.4	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	4	1.4	*
12	Meningitis (G00,G03)	4	1.4	*
15	Influenza and pneumonia (J10-J18)	3	1.0	*
...	All other causes (Residual)	65	22.3	4.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 15-19 years				
...	All causes	625	100.0	38.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	27.2	10.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	20.8	7.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	6.9	2.6
4	Malignant neoplasms (C00-C97)	38	6.1	2.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	3.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	3.0	*
7	Human immunodeficiency virus (HIV) disease (B20-B24)	13	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	12	1.9	*
9	Anemias (D50-D64)	10	1.6	*
10	Pregnancy, childbirth and the puerperium (O00-099)	9	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.3	*
11	Cerebrovascular diseases (I60-I69)	8	1.3	*
13	Diabetes mellitus (E10-E14)	7	1.1	*
14	Influenza and pneumonia (J10-J18)	6	1.0	*
15	Septicemia (A40-A41)	5	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.8	*
...	All other causes (Residual)	120	19.2	7.3
Black, non-Hispanic, female, 20-24 years				
...	All causes	1,031	100.0	68.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	240	23.3	16.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	177	17.2	11.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	79	7.7	5.3
4	Malignant neoplasms (C00-C97)	70	6.8	4.7
5	Pregnancy, childbirth and the puerperium (O00-099)	48	4.7	3.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	47	4.6	3.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34	3.3	2.3
8	Anemias (D50-D64)	17	1.6	*
9	Diabetes mellitus (E10-E14)	16	1.6	*
9	Cerebrovascular diseases (I60-I69)	16	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	16	1.6	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	1.6	*
13	Septicemia (A40-A41)	15	1.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.4	*
15	Influenza and pneumonia (J10-J18)	12	1.2	*
...	All other causes (Residual)	214	20.8	14.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 25-29 years				
...	All causes	1,351	100.0	93.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	16.1	15.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	195	14.4	13.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	133	9.8	9.2
4	Human immunodeficiency virus (HIV) disease (B20-B24)	130	9.6	9.0
5	Malignant neoplasms (C00-C97)	128	9.5	8.9
6	Pregnancy, childbirth and the puerperium (O00-099)	66	4.9	4.6
7	Diabetes mellitus (E10-E14)	47	3.5	3.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	2.1	2.0
9	Anemias (D50-D64)	27	2.0	1.9
10	Septicemia (A40-A41)	23	1.7	1.6
10	Cerebrovascular diseases (I60-I69)	23	1.7	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	1.6	1.5
13	Chronic lower respiratory diseases (J40-J47)	20	1.5	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.4	*
15	Influenza and pneumonia (J10-J18)	10	0.7	*
...	All other causes (Residual)	263	19.5	18.2
Black, non-Hispanic, female, 30-34 years				
...	All causes	1,741	100.0	130.9
1	Malignant neoplasms (C00-C97)	237	13.6	17.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	234	13.4	17.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	219	12.6	16.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	206	11.8	15.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	7.2	9.4
6	Diabetes mellitus (E10-E14)	55	3.2	4.1
7	Cerebrovascular diseases (I60-I69)	51	2.9	3.8
8	Pregnancy, childbirth and the puerperium (O00-099)	49	2.8	3.7
9	Influenza and pneumonia (J10-J18)	30	1.7	2.3
10	Anemias (D50-D64)	29	1.7	2.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.7	2.2
12	Chronic lower respiratory diseases (J40-J47)	25	1.4	1.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	1.3	1.7
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23	1.3	1.7
15	Septicemia (A40-A41)	21	1.2	1.6
...	All other causes (Residual)	385	22.1	28.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 35-39 years				
...	All causes	2,752	100.0	196.5
1	Malignant neoplasms (C00-C97)	489	17.8	34.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	413	15.0	29.5
3	Human immunodeficiency virus (HIV) disease (B20-B24)	356	12.9	25.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	281	10.2	20.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	5.1	10.1
6	Cerebrovascular diseases (I60-I69)	104	3.8	7.4
7	Diabetes mellitus (E10-E14)	88	3.2	6.3
8	Septicemia (A40-A41)	59	2.1	4.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	1.9	3.8
10	Pregnancy, childbirth and the puerperium (O00-099)	51	1.9	3.6
11	Chronic lower respiratory diseases (J40-J47)	39	1.4	2.8
12	Anemias (D50-D64)	35	1.3	2.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	34	1.2	2.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	1.2	2.3
15	Influenza and pneumonia (J10-J18)	29	1.1	2.1
...	All other causes (Residual)	548	19.9	39.1
Black, non-Hispanic, female, 40-44 years				
...	All causes	4,585	100.0	308.4
1	Malignant neoplasms (C00-C97)	1,033	22.5	69.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	760	16.6	51.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	467	10.2	31.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	389	8.5	26.2
5	Cerebrovascular diseases (I60-I69)	248	5.4	16.7
6	Diabetes mellitus (E10-E14)	147	3.2	9.9
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	3.0	9.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	2.2	6.7
9	Chronic lower respiratory diseases (J40-J47)	83	1.8	5.6
10	Septicemia (A40-A41)	79	1.7	5.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	66	1.4	4.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	60	1.3	4.0
13	Influenza and pneumonia (J10-J18)	59	1.3	4.0
14	Anemias (D50-D64)	44	1.0	3.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	0.7	2.2
...	All other causes (Residual)	881	19.2	59.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 45-49 years				
...	All causes	6,793	100.0	467.0
1	Malignant neoplasms (C00-C97)	1,888	27.8	129.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,293	19.0	88.9
3	Human immunodeficiency virus (HIV) disease (B20-B24)	491	7.2	33.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	438	6.4	30.1
5	Cerebrovascular diseases (I60-I69)	384	5.7	26.4
6	Diabetes mellitus (E10-E14)	225	3.3	15.5
7	Septicemia (A40-A41)	149	2.2	10.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	130	1.9	8.9
9	Chronic lower respiratory diseases (J40-J47)	112	1.6	7.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	111	1.6	7.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	95	1.4	6.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	1.4	6.5
13	Influenza and pneumonia (J10-J18)	92	1.4	6.3
14	Viral hepatitis (B15-B19)	51	0.8	3.5
15	Anemias (D50-D64)	41	0.6	2.8
...	All other causes (Residual)	1,198	17.6	82.4
Black, non-Hispanic, female, 50-54 years				
...	All causes	8,368	100.0	673.7
1	Malignant neoplasms (C00-C97)	2,502	29.9	201.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,793	21.4	144.3
3	Cerebrovascular diseases (I60-I69)	482	5.8	38.8
4	Diabetes mellitus (E10-E14)	385	4.6	31.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	342	4.1	27.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	301	3.6	24.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	229	2.7	18.4
8	Chronic lower respiratory diseases (J40-J47)	188	2.2	15.1
9	Septicemia (A40-A41)	174	2.1	14.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	151	1.8	12.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	142	1.7	11.4
12	Influenza and pneumonia (J10-J18)	87	1.0	7.0
13	Viral hepatitis (B15-B19)	84	1.0	6.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	0.9	6.2
15	Aortic aneurysm and dissection (I71)	39	0.5	3.1
...	All other causes (Residual)	1,392	16.6	112.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 55-59 years				
...	All causes	9,533	100.0	929.8
1	Malignant neoplasms (C00-C97)	3,056	32.1	298.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,201	23.1	214.7
3	Diabetes mellitus (E10-E14)	557	5.8	54.3
4	Cerebrovascular diseases (I60-I69)	525	5.5	51.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	276	2.9	26.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	2.9	26.6
7	Chronic lower respiratory diseases (J40-J47)	240	2.5	23.4
8	Septicemia (A40-A41)	222	2.3	21.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	190	2.0	18.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	150	1.6	14.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	132	1.4	12.9
12	Influenza and pneumonia (J10-J18)	108	1.1	10.5
13	Viral hepatitis (B15-B19)	75	0.8	7.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	0.4	3.7
15	Aortic aneurysm and dissection (I71)	36	0.4	3.5
...	All other causes (Residual)	1,454	15.3	141.8
Black, non-Hispanic, female, 60-64 years				
...	All causes	9,499	100.0	1,348.2
1	Malignant neoplasms (C00-C97)	3,239	34.1	459.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,256	23.7	320.2
3	Diabetes mellitus (E10-E14)	629	6.6	89.3
4	Cerebrovascular diseases (I60-I69)	556	5.9	78.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	348	3.7	49.4
6	Chronic lower respiratory diseases (J40-J47)	271	2.9	38.5
7	Septicemia (A40-A41)	237	2.5	33.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	181	1.9	25.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	1.7	22.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	96	1.0	13.6
11	Influenza and pneumonia (J10-J18)	95	1.0	13.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	72	0.8	10.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	7.4
14	Aortic aneurysm and dissection (I71)	38	0.4	5.4
15	Viral hepatitis (B15-B19)	37	0.4	5.3
...	All other causes (Residual)	1,231	13.0	174.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 65-69 years				
...	All causes	10,727	100.0	1,911.3
1	Malignant neoplasms (C00-C97)	3,412	31.8	607.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,651	24.7	472.3
3	Diabetes mellitus (E10-E14)	762	7.1	135.8
4	Cerebrovascular diseases (I60-I69)	667	6.2	118.8
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	441	4.1	78.6
6	Chronic lower respiratory diseases (J40-J47)	357	3.3	63.6
7	Septicemia (A40-A41)	294	2.7	52.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	216	2.0	38.5
9	Influenza and pneumonia (J10-J18)	182	1.7	32.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	1.5	28.0
11	Chronic liver disease and cirrhosis (K70,K73-K74)	63	0.6	11.2
12	Aortic aneurysm and dissection (I71)	52	0.5	9.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	51	0.5	9.1
14	Viral hepatitis (B15-B19)	41	0.4	7.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.4	7.1
15	Alzheimer's disease (G30)	40	0.4	7.1
...	All other causes (Residual)	1,301	12.1	231.8
Black, non-Hispanic, female, 70-74 years				
...	All causes	12,582	100.0	2,732.5
1	Malignant neoplasms (C00-C97)	3,583	28.5	778.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,363	26.7	730.4
3	Cerebrovascular diseases (I60-I69)	884	7.0	192.0
4	Diabetes mellitus (E10-E14)	819	6.5	177.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	533	4.2	115.8
6	Chronic lower respiratory diseases (J40-J47)	426	3.4	92.5
7	Septicemia (A40-A41)	373	3.0	81.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	220	1.7	47.8
9	Influenza and pneumonia (J10-J18)	180	1.4	39.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	157	1.2	34.1
11	Alzheimer's disease (G30)	153	1.2	33.2
12	Pneumonitis due to solids and liquids (J69)	65	0.5	14.1
13	Aortic aneurysm and dissection (I71)	62	0.5	13.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.5	12.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	56	0.4	12.2
...	All other causes (Residual)	1,650	13.1	358.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 75-79 years				
...	All causes	14,983	100.0	4,118.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,164	27.8	1,144.6
2	Malignant neoplasms (C00-C97)	3,643	24.3	1,001.4
3	Cerebrovascular diseases (I60-I69)	1,184	7.9	325.5
4	Diabetes mellitus (E10-E14)	952	6.4	261.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	586	3.9	161.1
6	Chronic lower respiratory diseases (J40-J47)	545	3.6	149.8
7	Septicemia (A40-A41)	401	2.7	110.2
8	Alzheimer's disease (G30)	340	2.3	93.5
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	340	2.3	93.5
10	Influenza and pneumonia (J10-J18)	281	1.9	77.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	180	1.2	49.5
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	82	0.5	22.5
13	Aortic aneurysm and dissection (I71)	81	0.5	22.3
14	Pneumonitis due to solids and liquids (J69)	73	0.5	20.1
15	Parkinson's disease (G20-G21)	54	0.4	14.8
...	All other causes (Residual)	2,077	13.9	570.9
Black, non-Hispanic, female, 80-84 years				
...	All causes	17,388	100.0	6,438.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,237	30.1	1,939.3
2	Malignant neoplasms (C00-C97)	3,218	18.5	1,191.6
3	Cerebrovascular diseases (I60-I69)	1,490	8.6	551.8
4	Diabetes mellitus (E10-E14)	989	5.7	366.2
5	Alzheimer's disease (G30)	658	3.8	243.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	617	3.5	228.5
7	Chronic lower respiratory diseases (J40-J47)	498	2.9	184.4
8	Septicemia (A40-A41)	476	2.7	176.3
9	Influenza and pneumonia (J10-J18)	467	2.7	172.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	377	2.2	139.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	213	1.2	78.9
12	Pneumonitis due to solids and liquids (J69)	107	0.6	39.6
13	Aortic aneurysm and dissection (I71)	91	0.5	33.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	86	0.5	31.8
15	Atherosclerosis (I70)	66	0.4	24.4
...	All other causes (Residual)	2,798	16.1	1,036.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 85-89 years				
...	All causes	15,148	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,793	31.6	---
2	Malignant neoplasms (C00-C97)	2,090	13.8	---
3	Cerebrovascular diseases (I60-I69)	1,372	9.1	---
4	Alzheimer's disease (G30)	796	5.3	---
5	Diabetes mellitus (E10-E14)	728	4.8	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	571	3.8	---
7	Influenza and pneumonia (J10-J18)	435	2.9	---
8	Septicemia (A40-A41)	381	2.5	---
9	Chronic lower respiratory diseases (J40-J47)	372	2.5	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	367	2.4	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	1.1	---
12	Pneumonitis due to solids and liquids (J69)	123	0.8	---
13	Atherosclerosis (I70)	79	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	74	0.5	---
15	Parkinson's disease (G20-G21)	69	0.5	---
...	All other causes (Residual)	2,737	18.1	---
Black, non-Hispanic, female, 90-94 years				
...	All causes	11,049	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,676	33.3	---
2	Malignant neoplasms (C00-C97)	1,091	9.9	---
3	Cerebrovascular diseases (I60-I69)	957	8.7	---
4	Alzheimer's disease (G30)	762	6.9	---
5	Diabetes mellitus (E10-E14)	408	3.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	389	3.5	---
7	Influenza and pneumonia (J10-J18)	388	3.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	302	2.7	---
9	Septicemia (A40-A41)	259	2.3	---
10	Chronic lower respiratory diseases (J40-J47)	216	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	1.0	---
12	Pneumonitis due to solids and liquids (J69)	109	1.0	---
13	Atherosclerosis (I70)	63	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	0.4	---
15	Aortic aneurysm and dissection (I71)	47	0.4	---
...	All other causes (Residual)	2,222	20.1	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years because population figures are not available for these age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title for Essential hypertension and hypertensive renal disease was changed in 2006 to reflect the addition of Secondary hypertension (ICD code I15); see the "Technical Notes" of *Deaths: Final Data for 2006*]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Black, non-Hispanic, female, 95-99 years				
...	All causes	5,133	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,809	35.2	---
2	Cerebrovascular diseases (I60-I69)	418	8.1	---
3	Alzheimer's disease (G30)	365	7.1	---
4	Malignant neoplasms (C00-C97)	350	6.8	---
5	Influenza and pneumonia (J10-J18)	200	3.9	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	181	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	2.7	---
8	Diabetes mellitus (E10-E14)	117	2.3	---
9	Septicemia (A40-A41)	103	2.0	---
10	Chronic lower respiratory diseases (J40-J47)	88	1.7	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	1.3	---
12	Pneumonitis due to solids and liquids (J69)	49	1.0	---
13	Atherosclerosis (I70)	37	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	0.5	---
15	Anemias (D50-D64)	23	0.4	---
...	All other causes (Residual)	1,163	22.7	---
Black, non-Hispanic, female, 100 years and over				
...	All causes	1,720	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	676	39.3	---
2	Cerebrovascular diseases (I60-I69)	131	7.6	---
3	Alzheimer's disease (G30)	96	5.6	---
4	Influenza and pneumonia (J10-J18)	74	4.3	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	3.8	---
6	Malignant neoplasms (C00-C97)	62	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	3.0	---
8	Diabetes mellitus (E10-E14)	29	1.7	---
9	Septicemia (A40-A41)	25	1.5	---
9	Chronic lower respiratory diseases (J40-J47)	25	1.5	---
11	Pneumonitis due to solids and liquids (J69)	18	1.0	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.0	---
13	Atherosclerosis (I70)	12	0.7	---
14	Nutritional deficiencies (E40-E64)	8	0.5	---
15	Anemias (D50-D64)	7	0.4	---
...	All other causes (Residual)	422	24.5	---

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available