

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	141,576	100.0	292.4
1	Malignant neoplasms (C00-C97)	29,935	21.1	61.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,611	20.9	61.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,654	7.5	22.0
4	Cerebrovascular diseases (I60-I69)	7,065	5.0	14.6
5	Diabetes mellitus (E10-E14)	6,311	4.5	13.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,303	3.0	8.9
7	Chronic lower respiratory diseases (J40-J47)	4,026	2.8	8.3
8	Influenza and pneumonia (J09-J18)	3,679	2.6	7.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,179	2.2	6.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,107	2.2	6.4
11	Alzheimer's disease (G30)	2,986	2.1	6.2
12	Certain conditions originating in the perinatal period (P00-P96)	2,618	1.8	5.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,573	1.8	5.3
14	Septicemia (A40-A41)	2,077	1.5	4.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,851	1.3	3.8
...	All other causes (Residual)	27,601	19.5	57.0
Hispanic, both sexes, 1-4 years				
...	All causes	1,004	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	32.3	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	12.4	2.8
3	Malignant neoplasms (C00-C97)	99	9.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	7.5	1.7
5	Influenza and pneumonia (J09-J18)	34	3.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	2.6	0.6
7	Septicemia (A40-A41)	17	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.1	*
9	Cerebrovascular diseases (I60-I69)	11	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	1.0	*
12	Meningitis (G00,G03)	4	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
...	All other causes (Residual)	247	24.6	5.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	553	100.0	11.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	29.1	3.4
2	Malignant neoplasms (C00-C97)	115	20.8	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.9	0.8
4	Influenza and pneumonia (J09-J18)	30	5.4	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	4.9	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.6	0.4
7	Chronic lower respiratory diseases (J40-J47)	12	2.2	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.8	*
9	Septicemia (A40-A41)	7	1.3	*
9	Cerebrovascular diseases (I60-I69)	7	1.3	*
11	Anemias (D50-D64)	2	0.4	*
11	Diabetes mellitus (E10-E14)	2	0.4	*
11	Diseases of appendix (K35-K38)	2	0.4	*
14	Scarlet fever and erysipelas (A38,A46)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Aortic aneurysm and dissection (I71)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.2	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.2	*
...	All other causes (Residual)	114	20.6	2.4
Hispanic, both sexes, 10-14 years				
...	All causes	619	100.0	15.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	171	27.6	4.2
2	Malignant neoplasms (C00-C97)	101	16.3	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	7.6	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	7.1	1.1
5	Influenza and pneumonia (J09-J18)	33	5.3	0.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	5.0	0.8
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.6	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	2.3	*
9	Septicemia (A40-A41)	8	1.3	*
10	Chronic lower respiratory diseases (J40-J47)	7	1.1	*
11	Diabetes mellitus (E10-E14)	5	0.8	*
12	Anemias (D50-D64)	4	0.6	*
12	Meningitis (G00,G03)	4	0.6	*
14	Cerebrovascular diseases (I60-I69)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
...	All other causes (Residual)	119	19.2	2.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 15-19 years				
...	All causes	2,087	100.0	51.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	782	37.5	19.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	512	24.5	12.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	242	11.6	6.0
4	Malignant neoplasms (C00-C97)	131	6.3	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	2.4	1.2
6	Influenza and pneumonia (J09-J18)	38	1.8	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	1.6	0.8
8	Cerebrovascular diseases (I60-I69)	18	0.9	*
9	Septicemia (A40-A41)	17	0.8	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	12	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	10	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.3	*
12	Diabetes mellitus (E10-E14)	7	0.3	*
14	Anemias (D50-D64)	5	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.2	*
15	Diseases of appendix (K35-K38)	4	0.2	*
...	All other causes (Residual)	214	10.3	5.3
Hispanic, both sexes, 20-24 years				
...	All causes	3,100	100.0	79.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,217	39.3	31.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	656	21.2	16.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	341	11.0	8.8
4	Malignant neoplasms (C00-C97)	189	6.1	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	3.2	2.5
6	Influenza and pneumonia (J09-J18)	50	1.6	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.2	1.0
8	Pregnancy, childbirth and the puerperium (O00-O99)	29	0.9	0.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	26	0.8	0.7
10	Cerebrovascular diseases (I60-I69)	24	0.8	0.6
11	Septicemia (A40-A41)	19	0.6	*
12	Diabetes mellitus (E10-E14)	17	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.5	*
14	Legal intervention (Y35,Y89.0)	13	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	11	0.4	*
...	All other causes (Residual)	359	11.6	9.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-29 years				
...	All causes	3,151	100.0	75.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,083	34.4	26.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	483	15.3	11.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	305	9.7	7.3
4	Malignant neoplasms (C00-C97)	270	8.6	6.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	5.0	3.8
6	Influenza and pneumonia (J09-J18)	99	3.1	2.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	56	1.8	1.3
8	Cerebrovascular diseases (I60-I69)	45	1.4	1.1
9	Chronic liver disease and cirrhosis (K70,K73-K74)	42	1.3	1.0
9	Pregnancy, childbirth and the puerperium (O00-O99)	42	1.3	1.0
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	1.2	0.9
12	Diabetes mellitus (E10-E14)	32	1.0	0.8
13	Septicemia (A40-A41)	28	0.9	0.7
14	Chronic lower respiratory diseases (J40-J47)	17	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.5	*
...	All other causes (Residual)	438	13.9	10.6
Hispanic, both sexes, 30-34 years				
...	All causes	3,218	100.0	79.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	971	30.2	24.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	362	11.2	9.0
3	Malignant neoplasms (C00-C97)	358	11.1	8.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	293	9.1	7.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	227	7.1	5.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	82	2.5	2.0
7	Influenza and pneumonia (J09-J18)	79	2.5	2.0
8	Cerebrovascular diseases (I60-I69)	72	2.2	1.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	67	2.1	1.7
10	Diabetes mellitus (E10-E14)	58	1.8	1.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	42	1.3	1.0
12	Septicemia (A40-A41)	31	1.0	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	0.9	0.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	0.8	0.7
15	Chronic lower respiratory diseases (J40-J47)	23	0.7	0.6
...	All other causes (Residual)	498	15.5	12.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, both sexes, 35-39 years				
...	All causes	3,742	100.0	99.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	868	23.2	23.1
2	Malignant neoplasms (C00-C97)	581	15.5	15.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	326	8.7	8.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	299	8.0	8.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	260	6.9	6.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	158	4.2	4.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	154	4.1	4.1
8	Cerebrovascular diseases (I60-I69)	127	3.4	3.4
9	Influenza and pneumonia (J09-J18)	107	2.9	2.8
10	Diabetes mellitus (E10-E14)	83	2.2	2.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.1	1.1
12	Septicemia (A40-A41)	35	0.9	0.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	0.9	0.9
14	Pregnancy, childbirth and the puerperium (O00-O99)	32	0.9	0.9
15	Chronic lower respiratory diseases (J40-J47)	26	0.7	0.7
...	All other causes (Residual)	612	16.4	16.3
Hispanic, both sexes, 40-44 years				
...	All causes	4,898	100.0	148.1
1	Malignant neoplasms (C00-C97)	932	19.0	28.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	888	18.1	26.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	598	12.2	18.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	328	6.7	9.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	246	5.0	7.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	217	4.4	6.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	202	4.1	6.1
8	Cerebrovascular diseases (I60-I69)	158	3.2	4.8
9	Influenza and pneumonia (J09-J18)	156	3.2	4.7
10	Diabetes mellitus (E10-E14)	132	2.7	4.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	1.6	2.4
12	Viral hepatitis (B15-B19)	67	1.4	2.0
13	Septicemia (A40-A41)	60	1.2	1.8
14	Chronic lower respiratory diseases (J40-J47)	41	0.8	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	0.5	0.7
...	All other causes (Residual)	771	15.7	23.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-49 years				
...	All causes	6,875	100.0	237.6
1	Malignant neoplasms (C00-C97)	1,540	22.4	53.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,044	15.2	36.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	871	12.7	30.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	549	8.0	19.0
5	Cerebrovascular diseases (I60-I69)	289	4.2	10.0
6	Diabetes mellitus (E10-E14)	258	3.8	8.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	249	3.6	8.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	228	3.3	7.9
9	Influenza and pneumonia (J09-J18)	174	2.5	6.0
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	156	2.3	5.4
11	Viral hepatitis (B15-B19)	149	2.2	5.1
12	Septicemia (A40-A41)	95	1.4	3.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	1.2	2.9
14	Chronic lower respiratory diseases (J40-J47)	59	0.9	2.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	0.7	1.6
...	All other causes (Residual)	1,085	15.8	37.5
Hispanic, both sexes, 50-54 years				
...	All causes	8,351	100.0	367.3
1	Malignant neoplasms (C00-C97)	2,192	26.2	96.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,456	17.4	64.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	689	8.3	30.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	642	7.7	28.2
5	Diabetes mellitus (E10-E14)	354	4.2	15.6
6	Cerebrovascular diseases (I60-I69)	339	4.1	14.9
7	Viral hepatitis (B15-B19)	230	2.8	10.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	202	2.4	8.9
9	Influenza and pneumonia (J09-J18)	194	2.3	8.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	2.1	7.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	169	2.0	7.4
12	Septicemia (A40-A41)	133	1.6	5.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	109	1.3	4.8
14	Chronic lower respiratory diseases (J40-J47)	106	1.3	4.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	63	0.8	2.8
...	All other causes (Residual)	1,298	15.5	57.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 55-59 years				
...	All causes	9,259	100.0	538.3
1	Malignant neoplasms (C00-C97)	2,754	29.7	160.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,849	20.0	107.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	672	7.3	39.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	520	5.6	30.2
5	Diabetes mellitus (E10-E14)	481	5.2	28.0
6	Cerebrovascular diseases (I60-I69)	378	4.1	22.0
7	Viral hepatitis (B15-B19)	256	2.8	14.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	202	2.2	11.7
9	Septicemia (A40-A41)	165	1.8	9.6
10	Influenza and pneumonia (J09-J18)	155	1.7	9.0
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	140	1.5	8.1
12	Chronic lower respiratory diseases (J40-J47)	132	1.4	7.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	131	1.4	7.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	85	0.9	4.9
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	0.6	3.1
...	All other causes (Residual)	1,285	13.9	74.7
Hispanic, both sexes, 60-64 years				
...	All causes	9,992	100.0	784.2
1	Malignant neoplasms (C00-C97)	3,218	32.2	252.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,187	21.9	171.6
3	Diabetes mellitus (E10-E14)	687	6.9	53.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	497	5.0	39.0
5	Cerebrovascular diseases (I60-I69)	480	4.8	37.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	364	3.6	28.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	253	2.5	19.9
8	Chronic lower respiratory diseases (J40-J47)	192	1.9	15.1
9	Septicemia (A40-A41)	179	1.8	14.0
10	Influenza and pneumonia (J09-J18)	160	1.6	12.6
11	Viral hepatitis (B15-B19)	135	1.4	10.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	106	1.1	8.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	77	0.8	6.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	0.8	6.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	0.5	3.6
...	All other causes (Residual)	1,334	13.4	104.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-69 years				
...	All causes	10,437	100.0	1,171.6
1	Malignant neoplasms (C00-C97)	3,401	32.6	381.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,332	22.3	261.8
3	Diabetes mellitus (E10-E14)	698	6.7	78.4
4	Cerebrovascular diseases (I60-I69)	515	4.9	57.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	407	3.9	45.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	298	2.9	33.5
7	Chronic lower respiratory diseases (J40-J47)	271	2.6	30.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	248	2.4	27.8
9	Influenza and pneumonia (J09-J18)	233	2.2	26.2
10	Septicemia (A40-A41)	179	1.7	20.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	122	1.2	13.7
12	Viral hepatitis (B15-B19)	93	0.9	10.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	72	0.7	8.1
14	Alzheimer's disease (G30)	55	0.5	6.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	46	0.4	5.2
...	All other causes (Residual)	1,467	14.1	164.7
Hispanic, both sexes, 70-74 years				
...	All causes	11,942	100.0	1,767.3
1	Malignant neoplasms (C00-C97)	3,550	29.7	525.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,836	23.7	419.7
3	Diabetes mellitus (E10-E14)	766	6.4	113.4
4	Cerebrovascular diseases (I60-I69)	653	5.5	96.6
5	Chronic lower respiratory diseases (J40-J47)	422	3.5	62.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	368	3.1	54.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	329	2.8	48.7
8	Influenza and pneumonia (J09-J18)	280	2.3	41.4
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	257	2.2	38.0
10	Septicemia (A40-A41)	205	1.7	30.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	162	1.4	24.0
12	Alzheimer's disease (G30)	145	1.2	21.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	0.7	12.6
14	Parkinson's disease (G20-G21)	77	0.6	11.4
15	Aortic aneurysm and dissection (I71)	63	0.5	9.3
...	All other causes (Residual)	1,744	14.6	258.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 75-79 years				
...	All causes	14,345	100.0	2,819.8
1	Malignant neoplasms (C00-C97)	3,761	26.2	739.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,574	24.9	702.5
3	Cerebrovascular diseases (I60-I69)	907	6.3	178.3
4	Diabetes mellitus (E10-E14)	867	6.0	170.4
5	Chronic lower respiratory diseases (J40-J47)	639	4.5	125.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	410	2.9	80.6
7	Influenza and pneumonia (J09-J18)	374	2.6	73.5
8	Alzheimer's disease (G30)	329	2.3	64.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	2.0	56.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	281	2.0	55.2
11	Septicemia (A40-A41)	235	1.6	46.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	178	1.2	35.0
13	Parkinson's disease (G20-G21)	168	1.2	33.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	83	0.6	16.3
15	Pneumonitis due to solids and liquids (J69)	72	0.5	14.2
...	All other causes (Residual)	2,179	15.2	428.3
Hispanic, both sexes, 80-84 years				
...	All causes	16,153	100.0	4,466.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,451	27.6	1,230.8
2	Malignant neoplasms (C00-C97)	3,349	20.7	926.1
3	Cerebrovascular diseases (I60-I69)	1,105	6.8	305.6
4	Diabetes mellitus (E10-E14)	879	5.4	243.1
5	Chronic lower respiratory diseases (J40-J47)	784	4.9	216.8
6	Alzheimer's disease (G30)	646	4.0	178.6
7	Influenza and pneumonia (J09-J18)	473	2.9	130.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	471	2.9	130.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	1.7	78.0
10	Septicemia (A40-A41)	263	1.6	72.7
11	Parkinson's disease (G20-G21)	260	1.6	71.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	213	1.3	58.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	174	1.1	48.1
14	Pneumonitis due to solids and liquids (J69)	120	0.7	33.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.6	26.0
...	All other causes (Residual)	2,589	16.0	715.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, both sexes, 85-89 years				
...	All causes	14,113	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,123	29.2	---
2	Malignant neoplasms (C00-C97)	2,226	15.8	---
3	Cerebrovascular diseases (I60-I69)	1,017	7.2	---
4	Alzheimer's disease (G30)	841	6.0	---
5	Chronic lower respiratory diseases (J40-J47)	667	4.7	---
6	Diabetes mellitus (E10-E14)	648	4.6	---
7	Influenza and pneumonia (J09-J18)	475	3.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	357	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	263	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	240	1.7	---
11	Septicemia (A40-A41)	209	1.5	---
12	Parkinson's disease (G20-G21)	189	1.3	---
13	Pneumonitis due to solids and liquids (J69)	138	1.0	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	105	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.7	---
...	All other causes (Residual)	2,521	17.9	---
Hispanic, both sexes, 90-94 years				
...	All causes	8,117	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,699	33.3	---
2	Malignant neoplasms (C00-C97)	860	10.6	---
3	Alzheimer's disease (G30)	598	7.4	---
4	Cerebrovascular diseases (I60-I69)	596	7.3	---
5	Chronic lower respiratory diseases (J40-J47)	404	5.0	---
6	Influenza and pneumonia (J09-J18)	302	3.7	---
7	Diabetes mellitus (E10-E14)	248	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	2.4	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	155	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	1.8	---
11	Septicemia (A40-A41)	91	1.1	---
12	Pneumonitis due to solids and liquids (J69)	78	1.0	---
13	Parkinson's disease (G20-G21)	69	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	56	0.7	---
15	Atherosclerosis (I70)	41	0.5	---
...	All other causes (Residual)	1,582	19.5	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, both sexes, 95-99 years				
...	All causes	3,358	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,145	34.1	---
2	Alzheimer's disease (G30)	263	7.8	---
3	Malignant neoplasms (C00-C97)	260	7.7	---
4	Cerebrovascular diseases (I60-I69)	238	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	151	4.5	---
6	Influenza and pneumonia (J09-J18)	139	4.1	---
7	Diabetes mellitus (E10-E14)	77	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	76	2.3	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	72	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	53	1.6	---
11	Septicemia (A40-A41)	45	1.3	---
12	Pneumonitis due to solids and liquids (J69)	39	1.2	---
13	Atherosclerosis (I70)	22	0.7	---
14	Parkinson's disease (G20-G21)	20	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.5	---
...	All other causes (Residual)	741	22.1	---
Hispanic, both sexes, 100 years and over				
...	All causes	809	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	295	36.5	---
2	Alzheimer's disease (G30)	62	7.7	---
3	Cerebrovascular diseases (I60-I69)	50	6.2	---
4	Influenza and pneumonia (J09-J18)	40	4.9	---
5	Chronic lower respiratory diseases (J40-J47)	32	4.0	---
6	Malignant neoplasms (C00-C97)	31	3.8	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	2.2	---
9	Diabetes mellitus (E10-E14)	12	1.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	1.5	---
11	Atherosclerosis (I70)	7	0.9	---
11	Pneumonitis due to solids and liquids (J69)	7	0.9	---
13	Septicemia (A40-A41)	6	0.7	---
14	Anemias (D50-D64)	3	0.4	---
15	Parkinson's disease (G20-G21)	2	0.2	---
15	Aortic aneurysm and dissection (I71)	2	0.2	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.2	---
...	All other causes (Residual)	207	25.6	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	78,157	100.0	311.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,301	20.9	65.1
2	Malignant neoplasms (C00-C97)	15,772	20.2	62.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,841	10.0	31.3
4	Cerebrovascular diseases (I60-I69)	3,288	4.2	13.1
5	Diabetes mellitus (E10-E14)	3,212	4.1	12.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,969	3.8	11.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,635	3.4	10.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,135	2.7	8.5
9	Chronic lower respiratory diseases (J40-J47)	2,024	2.6	8.1
10	Influenza and pneumonia (J09-J18)	1,882	2.4	7.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,593	2.0	6.4
12	Certain conditions originating in the perinatal period (P00-P96)	1,462	1.9	5.8
13	Septicemia (A40-A41)	1,041	1.3	4.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	985	1.3	3.9
15	Alzheimer's disease (G30)	973	1.2	3.9
...	All other causes (Residual)	14,044	18.0	56.0
Hispanic, male, 1-4 years				
...	All causes	539	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	38.6	9.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	10.8	2.6
3	Malignant neoplasms (C00-C97)	44	8.2	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	5.9	1.4
5	Influenza and pneumonia (J09-J18)	16	3.0	*
6	Septicemia (A40-A41)	12	2.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.0	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
9	Cerebrovascular diseases (I60-I69)	5	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
12	Meningococcal infection (A39)	1	0.2	*
12	Anemias (D50-D64)	1	0.2	*
12	Meningitis (G00,G03)	1	0.2	*
12	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
12	Hernia (K40-K46)	1	0.2	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	131	24.3	5.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 5-9 years				
...	All causes	288	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	83	28.8	3.4
2	Malignant neoplasms (C00-C97)	65	22.6	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	7.3	0.9
4	Influenza and pneumonia (J09-J18)	16	5.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	3.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.8	*
6	Chronic lower respiratory diseases (J40-J47)	8	2.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.7	*
9	Cerebrovascular diseases (I60-I69)	4	1.4	*
10	Septicemia (A40-A41)	1	0.3	*
10	Anemias (D50-D64)	1	0.3	*
10	Diabetes mellitus (E10-E14)	1	0.3	*
10	Aortic aneurysm and dissection (I71)	1	0.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
...	All other causes (Residual)	62	21.5	2.5
Hispanic, male, 10-14 years				
...	All causes	362	100.0	17.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	29.6	5.1
2	Malignant neoplasms (C00-C97)	53	14.6	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	9.9	1.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	8.6	1.5
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	5.0	*
6	Influenza and pneumonia (J09-J18)	17	4.7	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	2.5	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.7	*
10	Diabetes mellitus (E10-E14)	3	0.8	*
11	Septicemia (A40-A41)	2	0.6	*
11	Meningitis (G00,G03)	2	0.6	*
11	Cerebrovascular diseases (I60-I69)	2	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
...	All other causes (Residual)	60	16.6	2.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, male, 15-19 years				
...	All causes	1,538	100.0	73.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	577	37.5	27.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	457	29.7	21.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	188	12.2	9.0
4	Malignant neoplasms (C00-C97)	71	4.6	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	2.1	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	1.3	1.0
7	Influenza and pneumonia (J09-J18)	19	1.2	*
8	Cerebrovascular diseases (I60-I69)	12	0.8	*
9	Septicemia (A40-A41)	10	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.2	*
10	Anemias (D50-D64)	3	0.2	*
10	Diabetes mellitus (E10-E14)	3	0.2	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.2	*
10	Diseases of appendix (K35-K38)	3	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.1	*
15	Aortic aneurysm and dissection (I71)	2	0.1	*
15	Pneumonitis due to solids and liquids (J69)	2	0.1	*
15	Legal intervention (Y35,Y89.0)	2	0.1	*
...	All other causes (Residual)	128	8.3	6.1
Hispanic, male, 20-24 years				
...	All causes	2,454	100.0	118.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,013	41.3	49.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	587	23.9	28.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	300	12.2	14.5
4	Malignant neoplasms (C00-C97)	133	5.4	6.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	68	2.8	3.3
6	Influenza and pneumonia (J09-J18)	21	0.9	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	0.8	1.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.7	*
9	Cerebrovascular diseases (I60-I69)	13	0.5	*
9	Legal intervention (Y35,Y89.0)	13	0.5	*
11	Septicemia (A40-A41)	11	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	7	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.2	*
14	Diabetes mellitus (E10-E14)	6	0.2	*
...	All other causes (Residual)	232	9.5	11.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-29 years				
...	All causes	2,363	100.0	103.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	891	37.7	39.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	417	17.6	18.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	251	10.6	11.0
4	Malignant neoplasms (C00-C97)	163	6.9	7.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	112	4.7	4.9
6	Influenza and pneumonia (J09-J18)	61	2.6	2.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	49	2.1	2.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	35	1.5	1.5
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	1.0	1.0
10	Cerebrovascular diseases (I60-I69)	21	0.9	0.9
11	Septicemia (A40-A41)	18	0.8	*
12	Diabetes mellitus (E10-E14)	16	0.7	*
13	Legal intervention (Y35,Y89.0)	14	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	9	0.4	*
...	All other causes (Residual)	263	11.1	11.5
Hispanic, male, 30-34 years				
...	All causes	2,325	100.0	104.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	783	33.7	35.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	304	13.1	13.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	250	10.8	11.3
4	Malignant neoplasms (C00-C97)	173	7.4	7.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	160	6.9	7.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	64	2.8	2.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	58	2.5	2.6
8	Influenza and pneumonia (J09-J18)	51	2.2	2.3
9	Cerebrovascular diseases (I60-I69)	46	2.0	2.1
10	Diabetes mellitus (E10-E14)	39	1.7	1.8
11	Septicemia (A40-A41)	22	0.9	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	0.7	*
14	Legal intervention (Y35,Y89.0)	16	0.7	*
15	Chronic lower respiratory diseases (J40-J47)	13	0.6	*
...	All other causes (Residual)	311	13.4	14.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 35-39 years				
...	All causes	2,573	100.0	127.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	698	27.1	34.5
2	Malignant neoplasms (C00-C97)	258	10.0	12.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	246	9.6	12.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	243	9.4	12.0
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	220	8.6	10.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	129	5.0	6.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	121	4.7	6.0
8	Cerebrovascular diseases (I60-I69)	77	3.0	3.8
9	Influenza and pneumonia (J09-J18)	68	2.6	3.4
10	Diabetes mellitus (E10-E14)	50	1.9	2.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.9	1.1
12	Septicemia (A40-A41)	22	0.9	1.1
13	Chronic lower respiratory diseases (J40-J47)	15	0.6	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.5	*
15	Legal intervention (Y35,Y89.0)	14	0.5	*
...	All other causes (Residual)	360	14.0	17.8
Hispanic, male, 40-44 years				
...	All causes	3,238	100.0	184.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	698	21.6	39.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	468	14.5	26.6
3	Malignant neoplasms (C00-C97)	402	12.4	22.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	262	8.1	14.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	196	6.1	11.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	177	5.5	10.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	157	4.8	8.9
8	Influenza and pneumonia (J09-J18)	102	3.2	5.8
9	Cerebrovascular diseases (I60-I69)	93	2.9	5.3
10	Diabetes mellitus (E10-E14)	81	2.5	4.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	1.6	2.9
12	Viral hepatitis (B15-B19)	44	1.4	2.5
13	Septicemia (A40-A41)	33	1.0	1.9
14	Chronic lower respiratory diseases (J40-J47)	16	0.5	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.4	*
...	All other causes (Residual)	445	13.7	25.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, male, 45-49 years				
...	All causes	4,447	100.0	296.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	779	17.5	52.0
2	Malignant neoplasms (C00-C97)	708	15.9	47.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	645	14.5	43.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	429	9.6	28.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	202	4.5	13.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	4.2	12.5
7	Cerebrovascular diseases (I60-I69)	175	3.9	11.7
8	Diabetes mellitus (E10-E14)	171	3.8	11.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	119	2.7	7.9
10	Viral hepatitis (B15-B19)	107	2.4	7.1
11	Influenza and pneumonia (J09-J18)	97	2.2	6.5
12	Septicemia (A40-A41)	54	1.2	3.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	53	1.2	3.5
14	Chronic lower respiratory diseases (J40-J47)	31	0.7	2.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.7	1.9
...	All other causes (Residual)	661	14.9	44.1
Hispanic, male, 50-54 years				
...	All causes	5,508	100.0	478.9
1	Malignant neoplasms (C00-C97)	1,159	21.0	100.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,095	19.9	95.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	9.5	45.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	502	9.1	43.7
5	Diabetes mellitus (E10-E14)	221	4.0	19.2
6	Cerebrovascular diseases (I60-I69)	212	3.8	18.4
7	Viral hepatitis (B15-B19)	168	3.1	14.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	160	2.9	13.9
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	2.7	12.9
10	Influenza and pneumonia (J09-J18)	123	2.2	10.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	1.9	9.1
12	Septicemia (A40-A41)	93	1.7	8.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	1.6	7.7
14	Chronic lower respiratory diseases (J40-J47)	52	0.9	4.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	0.7	3.2
...	All other causes (Residual)	824	15.0	71.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, male, 55-59 years				
...	All causes	5,871	100.0	695.4
1	Malignant neoplasms (C00-C97)	1,477	25.2	175.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,361	23.2	161.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	495	8.4	58.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	399	6.8	47.3
5	Diabetes mellitus (E10-E14)	288	4.9	34.1
6	Cerebrovascular diseases (I60-I69)	218	3.7	25.8
7	Viral hepatitis (B15-B19)	180	3.1	21.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	121	2.1	14.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	112	1.9	13.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	96	1.6	11.4
11	Septicemia (A40-A41)	95	1.6	11.3
12	Influenza and pneumonia (J09-J18)	89	1.5	10.5
13	Chronic lower respiratory diseases (J40-J47)	73	1.2	8.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	0.9	6.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	0.6	4.5
...	All other causes (Residual)	777	13.2	92.0
Hispanic, male, 60-64 years				
...	All causes	6,070	100.0	999.1
1	Malignant neoplasms (C00-C97)	1,774	29.2	292.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,490	24.5	245.2
3	Diabetes mellitus (E10-E14)	410	6.8	67.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	351	5.8	57.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	4.6	46.4
6	Cerebrovascular diseases (I60-I69)	274	4.5	45.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	144	2.4	23.7
8	Chronic lower respiratory diseases (J40-J47)	104	1.7	17.1
9	Influenza and pneumonia (J09-J18)	96	1.6	15.8
10	Septicemia (A40-A41)	87	1.4	14.3
10	Viral hepatitis (B15-B19)	87	1.4	14.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	1.1	11.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	59	1.0	9.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	58	1.0	9.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	0.6	5.9
...	All other causes (Residual)	749	12.3	123.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65-69 years				
...	All causes	6,009	100.0	1,472.3
1	Malignant neoplasms (C00-C97)	1,888	31.4	462.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,489	24.8	364.8
3	Diabetes mellitus (E10-E14)	393	6.5	96.3
4	Cerebrovascular diseases (I60-I69)	303	5.0	74.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	277	4.6	67.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	148	2.5	36.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	146	2.4	35.8
8	Chronic lower respiratory diseases (J40-J47)	144	2.4	35.3
9	Influenza and pneumonia (J09-J18)	117	1.9	28.7
10	Septicemia (A40-A41)	92	1.5	22.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	69	1.1	16.9
12	Viral hepatitis (B15-B19)	57	0.9	14.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	0.9	12.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	33	0.5	8.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.4	6.4
15	Pneumonitis due to solids and liquids (J69)	26	0.4	6.4
...	All other causes (Residual)	749	12.5	183.5
Hispanic, male, 70-74 years				
...	All causes	6,614	100.0	2,205.8
1	Malignant neoplasms (C00-C97)	1,963	29.7	654.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,700	25.7	567.0
3	Diabetes mellitus (E10-E14)	406	6.1	135.4
4	Cerebrovascular diseases (I60-I69)	350	5.3	116.7
5	Chronic lower respiratory diseases (J40-J47)	248	3.7	82.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	197	3.0	65.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	179	2.7	59.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	2.4	54.0
9	Influenza and pneumonia (J09-J18)	153	2.3	51.0
10	Septicemia (A40-A41)	100	1.5	33.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	95	1.4	31.7
12	Alzheimer's disease (G30)	62	0.9	20.7
13	Parkinson's disease (G20-G21)	52	0.8	17.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	40	0.6	13.3
15	Aortic aneurysm and dissection (I71)	38	0.6	12.7
...	All other causes (Residual)	869	13.1	289.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, male, 75-79 years				
...	All causes	7,312	100.0	3,374.1
1	Malignant neoplasms (C00-C97)	2,065	28.2	952.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,916	26.2	884.1
3	Cerebrovascular diseases (I60-I69)	418	5.7	192.9
4	Diabetes mellitus (E10-E14)	379	5.2	174.9
5	Chronic lower respiratory diseases (J40-J47)	337	4.6	155.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	204	2.8	94.1
7	Influenza and pneumonia (J09-J18)	200	2.7	92.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	155	2.1	71.5
9	Alzheimer's disease (G30)	146	2.0	67.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	120	1.6	55.4
11	Septicemia (A40-A41)	112	1.5	51.7
12	Parkinson's disease (G20-G21)	98	1.3	45.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	73	1.0	33.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.6	20.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	0.5	18.5
...	All other causes (Residual)	1,004	13.7	463.3
Hispanic, male, 80-84 years				
...	All causes	7,599	100.0	5,201.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,194	28.9	1,501.7
2	Malignant neoplasms (C00-C97)	1,763	23.2	1,206.7
3	Cerebrovascular diseases (I60-I69)	440	5.8	301.2
4	Chronic lower respiratory diseases (J40-J47)	393	5.2	269.0
5	Diabetes mellitus (E10-E14)	379	5.0	259.4
6	Alzheimer's disease (G30)	236	3.1	161.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	217	2.9	148.5
8	Influenza and pneumonia (J09-J18)	204	2.7	139.6
9	Parkinson's disease (G20-G21)	152	2.0	104.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	1.9	97.9
11	Septicemia (A40-A41)	111	1.5	76.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	92	1.2	63.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	68	0.9	46.5
14	Pneumonitis due to solids and liquids (J69)	61	0.8	41.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.6	30.8
...	All other causes (Residual)	1,101	14.5	753.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, male, 85-89 years				
...	All causes	6,057	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,799	29.7	---
2	Malignant neoplasms (C00-C97)	1,094	18.1	---
3	Cerebrovascular diseases (I60-I69)	370	6.1	---
4	Chronic lower respiratory diseases (J40-J47)	324	5.3	---
5	Alzheimer's disease (G30)	270	4.5	---
6	Diabetes mellitus (E10-E14)	261	4.3	---
7	Influenza and pneumonia (J09-J18)	218	3.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	2.7	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	2.1	---
10	Parkinson's disease (G20-G21)	96	1.6	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	95	1.6	---
12	Septicemia (A40-A41)	82	1.4	---
13	Pneumonitis due to solids and liquids (J69)	69	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.8	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	41	0.7	---
...	All other causes (Residual)	997	16.5	---
Hispanic, male, 90-94 years				
...	All causes	2,849	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	922	32.4	---
2	Malignant neoplasms (C00-C97)	393	13.8	---
3	Cerebrovascular diseases (I60-I69)	171	6.0	---
4	Chronic lower respiratory diseases (J40-J47)	169	5.9	---
5	Alzheimer's disease (G30)	150	5.3	---
6	Influenza and pneumonia (J09-J18)	119	4.2	---
7	Diabetes mellitus (E10-E14)	89	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	2.7	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	2.0	---
11	Septicemia (A40-A41)	41	1.4	---
12	Pneumonitis due to solids and liquids (J69)	37	1.3	---
13	Parkinson's disease (G20-G21)	36	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.8	---
15	Atherosclerosis (I70)	13	0.5	---
...	All other causes (Residual)	490	17.2	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 95-99 years				
...	All causes	948	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	319	33.6	---
2	Malignant neoplasms (C00-C97)	108	11.4	---
3	Cerebrovascular diseases (I60-I69)	55	5.8	---
3	Chronic lower respiratory diseases (J40-J47)	55	5.8	---
5	Alzheimer's disease (G30)	51	5.4	---
6	Influenza and pneumonia (J09-J18)	47	5.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	3.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	1.8	---
10	Diabetes mellitus (E10-E14)	16	1.7	---
11	Pneumonitis due to solids and liquids (J69)	15	1.6	---
12	Septicemia (A40-A41)	14	1.5	---
13	Parkinson's disease (G20-G21)	9	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	5	0.5	---
...	All other causes (Residual)	176	18.6	---
Hispanic, male, 100 years and over				
...	All causes	185	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78	42.2	---
2	Cerebrovascular diseases (I60-I69)	11	5.9	---
2	Influenza and pneumonia (J09-J18)	11	5.9	---
4	Malignant neoplasms (C00-C97)	10	5.4	---
4	Alzheimer's disease (G30)	10	5.4	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	3.8	---
6	Chronic lower respiratory diseases (J40-J47)	7	3.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	3.2	---
9	Atherosclerosis (I70)	3	1.6	---
10	Aortic aneurysm and dissection (I71)	2	1.1	---
11	Septicemia (A40-A41)	1	0.5	---
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	---
11	Anemias (D50-D64)	1	0.5	---
11	Parkinson's disease (G20-G21)	1	0.5	---
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.5	---
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1	0.5	---
...	All other causes (Residual)	33	17.8	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, female, all ages				
...	All causes	63,419	100.0	271.5
1	Malignant neoplasms (C00-C97)	14,163	22.3	60.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,310	21.0	57.0
3	Cerebrovascular diseases (I60-I69)	3,777	6.0	16.2
4	Diabetes mellitus (E10-E14)	3,099	4.9	13.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,813	4.4	12.0
6	Alzheimer's disease (G30)	2,013	3.2	8.6
7	Chronic lower respiratory diseases (J40-J47)	2,002	3.2	8.6
8	Influenza and pneumonia (J09-J18)	1,797	2.8	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,514	2.4	6.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,334	2.1	5.7
11	Certain conditions originating in the perinatal period (P00-P96)	1,156	1.8	4.9
12	Septicemia (A40-A41)	1,036	1.6	4.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	882	1.4	3.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	791	1.2	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	544	0.9	2.3
...	All other causes (Residual)	13,188	20.8	56.4
Hispanic, female, 1-4 years				
...	All causes	465	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	24.9	5.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	14.2	3.1
3	Malignant neoplasms (C00-C97)	55	11.8	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	9.2	2.0
5	Influenza and pneumonia (J09-J18)	18	3.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	7	1.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
8	Cerebrovascular diseases (I60-I69)	6	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.3	*
11	Septicemia (A40-A41)	5	1.1	*
12	Meningitis (G00,G03)	3	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	3	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Anemias (D50-D64)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	111	23.9	5.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 5-9 years				
...	All causes	265	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	78	29.4	3.3
2	Malignant neoplasms (C00-C97)	50	18.9	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	6.4	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	6.4	*
5	Influenza and pneumonia (J09-J18)	14	5.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	4.5	*
7	Septicemia (A40-A41)	6	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.9	*
9	Chronic lower respiratory diseases (J40-J47)	4	1.5	*
10	Cerebrovascular diseases (I60-I69)	3	1.1	*
11	Diseases of appendix (K35-K38)	2	0.8	*
12	Scarlet fever and erysipelas (A38,A46)	1	0.4	*
12	Anemias (D50-D64)	1	0.4	*
12	Diabetes mellitus (E10-E14)	1	0.4	*
12	Meningitis (G00,G03)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
...	All other causes (Residual)	52	19.6	2.2
Hispanic, female, 10-14 years				
...	All causes	257	100.0	13.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	24.9	3.2
2	Malignant neoplasms (C00-C97)	48	18.7	2.4
3	Influenza and pneumonia (J09-J18)	16	6.2	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	6.2	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	13	5.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	3.9	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	3.1	*
8	Septicemia (A40-A41)	6	2.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.9	*
10	Anemias (D50-D64)	3	1.2	*
11	Diabetes mellitus (E10-E14)	2	0.8	*
11	Meningitis (G00,G03)	2	0.8	*
13	Meningococcal infection (A39)	1	0.4	*
13	Viral hepatitis (B15-B19)	1	0.4	*
13	Cerebrovascular diseases (I60-I69)	1	0.4	*
13	Aortic aneurysm and dissection (I71)	1	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
13	Pneumonitis due to solids and liquids (J69)	1	0.4	*
13	Diseases of appendix (K35-K38)	1	0.4	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	53	20.6	2.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, female, 15-19 years				
...	All causes	549	100.0	28.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	205	37.3	10.5
2	Malignant neoplasms (C00-C97)	60	10.9	3.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	10.0	2.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	9.8	2.8
5	Influenza and pneumonia (J09-J18)	19	3.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.1	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	2.6	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	12	2.2	*
9	Septicemia (A40-A41)	7	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	7	1.3	*
11	Cerebrovascular diseases (I60-I69)	6	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
12	Diabetes mellitus (E10-E14)	4	0.7	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.5	*
15	Anemias (D50-D64)	2	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (Residual)	78	14.2	4.0
Hispanic, female, 20-24 years				
...	All causes	646	100.0	35.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	31.6	11.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	10.7	3.8
3	Malignant neoplasms (C00-C97)	56	8.7	3.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	6.3	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	4.6	1.7
6	Influenza and pneumonia (J09-J18)	29	4.5	1.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	29	4.5	1.6
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	2.6	*
9	Diabetes mellitus (E10-E14)	11	1.7	*
9	Cerebrovascular diseases (I60-I69)	11	1.7	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	10	1.5	*
12	Septicemia (A40-A41)	8	1.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.9	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
...	All other causes (Residual)	118	18.3	6.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-29 years				
...	All causes	788	100.0	42.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	24.4	10.3
2	Malignant neoplasms (C00-C97)	107	13.6	5.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	8.4	3.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	6.9	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	5.6	2.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	42	5.3	2.3
7	Influenza and pneumonia (J09-J18)	38	4.8	2.0
8	Cerebrovascular diseases (I60-I69)	24	3.0	1.3
9	Diabetes mellitus (E10-E14)	16	2.0	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	2.0	*
11	Septicemia (A40-A41)	10	1.3	*
12	Chronic lower respiratory diseases (J40-J47)	8	1.0	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.9	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	7	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
...	All other causes (Residual)	152	19.3	8.1
Hispanic, female, 30-34 years				
...	All causes	893	100.0	49.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	188	21.1	10.4
2	Malignant neoplasms (C00-C97)	185	20.7	10.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	7.5	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	58	6.5	3.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	4.8	2.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	42	4.7	2.3
7	Influenza and pneumonia (J09-J18)	28	3.1	1.5
8	Cerebrovascular diseases (I60-I69)	26	2.9	1.4
9	Diabetes mellitus (E10-E14)	19	2.1	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18	2.0	*
11	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.1	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	1.1	*
14	Septicemia (A40-A41)	9	1.0	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	9	1.0	*
...	All other causes (Residual)	171	19.1	9.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 35-39 years				
...	All causes	1,169	100.0	67.5
1	Malignant neoplasms (C00-C97)	323	27.6	18.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	14.5	9.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	7.1	4.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	4.5	3.1
5	Cerebrovascular diseases (I60-I69)	50	4.3	2.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	3.4	2.3
7	Influenza and pneumonia (J09-J18)	39	3.3	2.3
8	Human immunodeficiency virus (HIV) disease (B20-B24)	37	3.2	2.1
9	Diabetes mellitus (E10-E14)	33	2.8	1.9
10	Pregnancy, childbirth and the puerperium (O00-O99)	32	2.7	1.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	25	2.1	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.5	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.5	*
14	Septicemia (A40-A41)	13	1.1	*
15	Chronic lower respiratory diseases (J40-J47)	11	0.9	*
...	All other causes (Residual)	224	19.2	12.9
Hispanic, female, 40-44 years				
...	All causes	1,660	100.0	107.1
1	Malignant neoplasms (C00-C97)	530	31.9	34.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	11.4	12.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	130	7.8	8.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	66	4.0	4.3
5	Cerebrovascular diseases (I60-I69)	65	3.9	4.2
6	Influenza and pneumonia (J09-J18)	54	3.3	3.5
7	Diabetes mellitus (E10-E14)	51	3.1	3.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	50	3.0	3.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	45	2.7	2.9
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	2.4	2.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	1.7	1.8
12	Septicemia (A40-A41)	27	1.6	1.7
13	Chronic lower respiratory diseases (J40-J47)	25	1.5	1.6
14	Viral hepatitis (B15-B19)	23	1.4	1.5
15	Pregnancy, childbirth and the puerperium (O00-O99)	12	0.7	*
...	All other causes (Residual)	324	19.5	20.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-49 years				
...	All causes	2,428	100.0	174.1
1	Malignant neoplasms (C00-C97)	832	34.3	59.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	265	10.9	19.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	226	9.3	16.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	120	4.9	8.6
5	Cerebrovascular diseases (I60-I69)	114	4.7	8.2
6	Diabetes mellitus (E10-E14)	87	3.6	6.2
7	Influenza and pneumonia (J09-J18)	77	3.2	5.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	47	1.9	3.4
9	Viral hepatitis (B15-B19)	42	1.7	3.0
10	Septicemia (A40-A41)	41	1.7	2.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	41	1.7	2.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	37	1.5	2.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.3	2.2
14	Chronic lower respiratory diseases (J40-J47)	28	1.2	2.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.9	1.5
...	All other causes (Residual)	419	17.3	30.0
Hispanic, female, 50-54 years				
...	All causes	2,843	100.0	253.0
1	Malignant neoplasms (C00-C97)	1,033	36.3	91.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	361	12.7	32.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	168	5.9	14.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	140	4.9	12.5
5	Diabetes mellitus (E10-E14)	133	4.7	11.8
6	Cerebrovascular diseases (I60-I69)	127	4.5	11.3
7	Influenza and pneumonia (J09-J18)	71	2.5	6.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	2.3	5.7
9	Viral hepatitis (B15-B19)	62	2.2	5.5
10	Chronic lower respiratory diseases (J40-J47)	54	1.9	4.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	42	1.5	3.7
12	Septicemia (A40-A41)	40	1.4	3.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	0.9	2.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	0.9	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	0.7	1.9
...	All other causes (Residual)	474	16.7	42.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, female, 55-59 years				
...	All causes	3,388	100.0	386.8
1	Malignant neoplasms (C00-C97)	1,277	37.7	145.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	488	14.4	55.7
3	Diabetes mellitus (E10-E14)	193	5.7	22.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	177	5.2	20.2
5	Cerebrovascular diseases (I60-I69)	160	4.7	18.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	3.6	13.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	81	2.4	9.2
8	Viral hepatitis (B15-B19)	76	2.2	8.7
9	Septicemia (A40-A41)	70	2.1	8.0
10	Influenza and pneumonia (J09-J18)	66	1.9	7.5
11	Chronic lower respiratory diseases (J40-J47)	59	1.7	6.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	35	1.0	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33	1.0	3.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	0.8	3.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.6	2.4
...	All other causes (Residual)	503	14.8	57.4
Hispanic, female, 60-64 years				
...	All causes	3,922	100.0	588.3
1	Malignant neoplasms (C00-C97)	1,444	36.8	216.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	697	17.8	104.6
3	Diabetes mellitus (E10-E14)	277	7.1	41.6
4	Cerebrovascular diseases (I60-I69)	206	5.3	30.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	146	3.7	21.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	109	2.8	16.4
7	Septicemia (A40-A41)	92	2.3	13.8
8	Chronic lower respiratory diseases (J40-J47)	88	2.2	13.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	2.1	12.3
10	Influenza and pneumonia (J09-J18)	64	1.6	9.6
11	Viral hepatitis (B15-B19)	48	1.2	7.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	47	1.2	7.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.5	3.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.5	*
15	Alzheimer's disease (G30)	12	0.3	*
...	All other causes (Residual)	570	14.5	85.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Hispanic, female, 65-69 years				
...	All causes	4,428	100.0	917.4
1	Malignant neoplasms (C00-C97)	1,513	34.2	313.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	843	19.0	174.6
3	Diabetes mellitus (E10-E14)	305	6.9	63.2
4	Cerebrovascular diseases (I60-I69)	212	4.8	43.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	3.4	31.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	130	2.9	26.9
7	Chronic lower respiratory diseases (J40-J47)	127	2.9	26.3
8	Influenza and pneumonia (J09-J18)	116	2.6	24.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	2.3	20.7
10	Septicemia (A40-A41)	87	2.0	18.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	53	1.2	11.0
12	Viral hepatitis (B15-B19)	36	0.8	7.5
13	Alzheimer's disease (G30)	33	0.7	6.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	0.5	4.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.3	*
...	All other causes (Residual)	687	15.5	142.3
Hispanic, female, 70-74 years				
...	All causes	5,328	100.0	1,417.6
1	Malignant neoplasms (C00-C97)	1,587	29.8	422.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,136	21.3	302.2
3	Diabetes mellitus (E10-E14)	360	6.8	95.8
4	Cerebrovascular diseases (I60-I69)	303	5.7	80.6
5	Chronic lower respiratory diseases (J40-J47)	174	3.3	46.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	171	3.2	45.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	150	2.8	39.9
8	Influenza and pneumonia (J09-J18)	127	2.4	33.8
9	Septicemia (A40-A41)	105	2.0	27.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	1.8	25.3
11	Alzheimer's disease (G30)	83	1.6	22.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	1.3	17.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.8	12.0
14	Viral hepatitis (B15-B19)	32	0.6	8.5
15	Parkinson's disease (G20-G21)	25	0.5	6.7
15	Aortic aneurysm and dissection (I71)	25	0.5	6.7
...	All other causes (Residual)	843	15.8	224.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Hispanic, female, 75-79 years				
...	All causes	7,033	100.0	2,408.4
1	Malignant neoplasms (C00-C97)	1,696	24.1	580.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,658	23.6	567.8
3	Cerebrovascular diseases (I60-I69)	489	7.0	167.5
4	Diabetes mellitus (E10-E14)	488	6.9	167.1
5	Chronic lower respiratory diseases (J40-J47)	302	4.3	103.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	206	2.9	70.5
7	Alzheimer's disease (G30)	183	2.6	62.7
8	Influenza and pneumonia (J09-J18)	174	2.5	59.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	161	2.3	55.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	133	1.9	45.5
11	Septicemia (A40-A41)	123	1.7	42.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	105	1.5	36.0
13	Parkinson's disease (G20-G21)	70	1.0	24.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.5	13.0
15	Viral hepatitis (B15-B19)	36	0.5	12.3
...	All other causes (Residual)	1,171	16.7	401.0
Hispanic, female, 80-84 years				
...	All causes	8,554	100.0	3,968.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,257	26.4	1,047.2
2	Malignant neoplasms (C00-C97)	1,586	18.5	735.8
3	Cerebrovascular diseases (I60-I69)	665	7.8	308.5
4	Diabetes mellitus (E10-E14)	500	5.8	232.0
5	Alzheimer's disease (G30)	410	4.8	190.2
6	Chronic lower respiratory diseases (J40-J47)	391	4.6	181.4
7	Influenza and pneumonia (J09-J18)	269	3.1	124.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	254	3.0	117.8
9	Septicemia (A40-A41)	152	1.8	70.5
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	1.6	64.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.4	56.1
12	Parkinson's disease (G20-G21)	108	1.3	50.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	106	1.2	49.2
14	Pneumonitis due to solids and liquids (J69)	59	0.7	27.4
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	54	0.6	25.1
...	All other causes (Residual)	1,483	17.3	688.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85-89 years				
...	All causes	8,056	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,324	28.8	---
2	Malignant neoplasms (C00-C97)	1,132	14.1	---
3	Cerebrovascular diseases (I60-I69)	647	8.0	---
4	Alzheimer's disease (G30)	571	7.1	---
5	Diabetes mellitus (E10-E14)	387	4.8	---
6	Chronic lower respiratory diseases (J40-J47)	343	4.3	---
7	Influenza and pneumonia (J09-J18)	257	3.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	191	2.4	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	145	1.8	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	136	1.7	---
11	Septicemia (A40-A41)	127	1.6	---
12	Parkinson's disease (G20-G21)	93	1.2	---
13	Pneumonitis due to solids and liquids (J69)	69	0.9	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	64	0.8	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.6	---
...	All other causes (Residual)	1,524	18.9	---
Hispanic, female, 90-94 years				
...	All causes	5,268	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,777	33.7	---
2	Malignant neoplasms (C00-C97)	467	8.9	---
3	Alzheimer's disease (G30)	448	8.5	---
4	Cerebrovascular diseases (I60-I69)	425	8.1	---
5	Chronic lower respiratory diseases (J40-J47)	235	4.5	---
6	Influenza and pneumonia (J09-J18)	183	3.5	---
7	Diabetes mellitus (E10-E14)	159	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	99	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	82	1.6	---
11	Septicemia (A40-A41)	50	0.9	---
12	Pneumonitis due to solids and liquids (J69)	41	0.8	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.6	---
14	Parkinson's disease (G20-G21)	33	0.6	---
15	Atherosclerosis (I70)	28	0.5	---
...	All other causes (Residual)	1,092	20.7	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 95-99 years				
...	All causes	2,410	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	826	34.3	---
2	Alzheimer's disease (G30)	212	8.8	---
3	Cerebrovascular diseases (I60-I69)	183	7.6	---
4	Malignant neoplasms (C00-C97)	152	6.3	---
5	Chronic lower respiratory diseases (J40-J47)	96	4.0	---
6	Influenza and pneumonia (J09-J18)	92	3.8	---
7	Diabetes mellitus (E10-E14)	61	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	1.5	---
11	Septicemia (A40-A41)	31	1.3	---
12	Pneumonitis due to solids and liquids (J69)	24	1.0	---
13	Atherosclerosis (I70)	19	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.5	---
14	Parkinson's disease (G20-G21)	11	0.5	---
...	All other causes (Residual)	563	23.4	---
Hispanic, female, 100 years and over				
...	All causes	624	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	217	34.8	---
2	Alzheimer's disease (G30)	52	8.3	---
3	Cerebrovascular diseases (I60-I69)	39	6.3	---
4	Influenza and pneumonia (J09-J18)	29	4.6	---
5	Chronic lower respiratory diseases (J40-J47)	25	4.0	---
6	Malignant neoplasms (C00-C97)	21	3.4	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	2.2	---
8	Diabetes mellitus (E10-E14)	12	1.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.9	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	1.8	---
11	Pneumonitis due to solids and liquids (J69)	7	1.1	---
12	Septicemia (A40-A41)	5	0.8	---
13	Atherosclerosis (I70)	4	0.6	---
14	Anemias (D50-D64)	2	0.3	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.3	---
...	All other causes (Residual)	172	27.6	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	1,944,606	100.0	961.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	485,779	25.0	240.3
2	Malignant neoplasms (C00-C97)	457,189	23.5	226.2
3	Chronic lower respiratory diseases (J40-J47)	122,605	6.3	60.6
4	Cerebrovascular diseases (I60-I69)	101,703	5.2	50.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91,416	4.7	45.2
6	Alzheimer's disease (G30)	69,695	3.6	34.5
7	Diabetes mellitus (E10-E14)	47,851	2.5	23.7
8	Influenza and pneumonia (J09-J18)	42,752	2.2	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,670	1.8	17.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,813	1.6	15.2
11	Septicemia (A40-A41)	26,533	1.4	13.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	22,520	1.2	11.1
13	Parkinson's disease (G20-G21)	18,554	1.0	9.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,552	1.0	9.2
15	Pneumonitis due to solids and liquids (J69)	13,489	0.7	6.7
...	All other causes (Residual)	359,485	18.5	177.8
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,181	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	773	35.4	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	230	10.5	2.5
3	Malignant neoplasms (C00-C97)	180	8.3	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	6.5	1.5
5	Influenza and pneumonia (J09-J18)	75	3.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.2	0.8
7	Septicemia (A40-A41)	30	1.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	27	1.2	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	27	1.2	0.3
11	Chronic lower respiratory diseases (J40-J47)	19	0.9	*
12	Meningitis (G00,G03)	11	0.5	*
13	Anemias (D50-D64)	10	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
15	Meningococcal infection (A39)	7	0.3	*
...	All other causes (Residual)	545	25.0	6.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,236	100.0	10.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	377	30.5	3.3
2	Malignant neoplasms (C00-C97)	255	20.6	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	8.9	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	4.0	0.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	3.6	0.4
6	Influenza and pneumonia (J09-J18)	44	3.6	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	17	1.4	*
9	Septicemia (A40-A41)	14	1.1	*
9	Cerebrovascular diseases (I60-I69)	14	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	8	0.6	*
12	Diabetes mellitus (E10-E14)	6	0.5	*
13	Anemias (D50-D64)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
15	Meningitis (G00,G03)	3	0.2	*
15	Pneumonitis due to solids and liquids (J69)	3	0.2	*
15	Diseases of appendix (K35-K38)	3	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
...	All other causes (Residual)	256	20.7	2.2
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,631	100.0	13.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	518	31.8	4.4
2	Malignant neoplasms (C00-C97)	238	14.6	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	155	9.5	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	5.0	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	4.1	0.6
6	Influenza and pneumonia (J09-J18)	60	3.7	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	46	2.8	0.4
8	Cerebrovascular diseases (I60-I69)	29	1.8	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.2	0.2
10	Septicemia (A40-A41)	14	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	9	0.6	*
12	Diabetes mellitus (E10-E14)	8	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.5	*
14	Meningitis (G00,G03)	7	0.4	*
15	Anemias (D50-D64)	5	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.3	*
...	All other causes (Residual)	360	22.1	3.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	6,363	100.0	49.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,237	50.9	25.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,152	18.1	8.9
3	Malignant neoplasms (C00-C97)	360	5.7	2.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	280	4.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	162	2.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	2.0	1.0
7	Influenza and pneumonia (J09-J18)	87	1.4	0.7
8	Chronic lower respiratory diseases (J40-J47)	37	0.6	0.3
9	Septicemia (A40-A41)	27	0.4	0.2
9	Cerebrovascular diseases (I60-I69)	27	0.4	0.2
11	Pregnancy, childbirth and the puerperium (O00-O99)	24	0.4	0.2
12	Diabetes mellitus (E10-E14)	22	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.2	*
15	Pneumonitis due to solids and liquids (J69)	9	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.1	*
...	All other causes (Residual)	776	12.2	6.0
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	10,787	100.0	81.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,327	49.4	40.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,894	17.6	14.3
3	Malignant neoplasms (C00-C97)	585	5.4	4.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	475	4.4	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	337	3.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	144	1.3	1.1
7	Influenza and pneumonia (J09-J18)	136	1.3	1.0
8	Cerebrovascular diseases (I60-I69)	70	0.6	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	69	0.6	0.5
10	Diabetes mellitus (E10-E14)	67	0.6	0.5
11	Septicemia (A40-A41)	46	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	34	0.3	0.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	24	0.2	0.2
15	Legal intervention (Y35,Y89.0)	21	0.2	0.2
...	All other causes (Residual)	1,532	14.2	11.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	12,125	100.0	92.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,262	43.4	40.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,994	16.4	15.2
3	Malignant neoplasms (C00-C97)	820	6.8	6.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	611	5.0	4.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	498	4.1	3.8
6	Influenza and pneumonia (J09-J18)	218	1.8	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	1.1	1.0
8	Diabetes mellitus (E10-E14)	135	1.1	1.0
9	Cerebrovascular diseases (I60-I69)	95	0.8	0.7
10	Septicemia (A40-A41)	88	0.7	0.7
11	Pregnancy, childbirth and the puerperium (O00-O99)	83	0.7	0.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	66	0.5	0.5
13	Chronic lower respiratory diseases (J40-J47)	57	0.5	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.4	0.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	40	0.3	0.3
...	All other causes (Residual)	1,977	16.3	15.1
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	13,078	100.0	111.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,625	35.4	39.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,989	15.2	16.9
3	Malignant neoplasms (C00-C97)	1,386	10.6	11.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,077	8.2	9.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	432	3.3	3.7
6	Influenza and pneumonia (J09-J18)	216	1.7	1.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	176	1.3	1.5
8	Diabetes mellitus (E10-E14)	168	1.3	1.4
9	Cerebrovascular diseases (I60-I69)	150	1.1	1.3
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	114	0.9	1.0
11	Septicemia (A40-A41)	104	0.8	0.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	100	0.8	0.8
13	Pregnancy, childbirth and the puerperium (O00-O99)	82	0.6	0.7
14	Chronic lower respiratory diseases (J40-J47)	79	0.6	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	0.5	0.6
...	All other causes (Residual)	2,312	17.7	19.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	18,287	100.0	144.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,954	27.1	39.2
2	Malignant neoplasms (C00-C97)	2,680	14.7	21.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,488	13.6	19.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,241	12.3	17.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	463	2.5	3.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	454	2.5	3.6
7	Diabetes mellitus (E10-E14)	357	2.0	2.8
8	Cerebrovascular diseases (I60-I69)	310	1.7	2.5
9	Influenza and pneumonia (J09-J18)	267	1.5	2.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	227	1.2	1.8
11	Septicemia (A40-A41)	182	1.0	1.4
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	0.8	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	0.7	1.0
14	Chronic lower respiratory diseases (J40-J47)	124	0.7	1.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.4	0.5
...	All other causes (Residual)	3,190	17.4	25.2
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	29,449	100.0	214.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,048	20.5	44.1
2	Malignant neoplasms (C00-C97)	5,533	18.8	40.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,606	15.6	33.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,020	10.3	22.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,155	3.9	8.4
6	Diabetes mellitus (E10-E14)	688	2.3	5.0
7	Cerebrovascular diseases (I60-I69)	652	2.2	4.8
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	489	1.7	3.6
9	Influenza and pneumonia (J09-J18)	450	1.5	3.3
10	Human immunodeficiency virus (HIV) disease (B20-B24)	412	1.4	3.0
11	Chronic lower respiratory diseases (J40-J47)	345	1.2	2.5
12	Septicemia (A40-A41)	337	1.1	2.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	233	0.8	1.7
14	Viral hepatitis (B15-B19)	195	0.7	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	194	0.7	1.4
...	All other causes (Residual)	5,092	17.3	37.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	52,481	100.0	330.1
1	Malignant neoplasms (C00-C97)	12,897	24.6	81.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,501	18.1	59.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,920	15.1	49.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,627	6.9	22.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,421	4.6	15.2
6	Diabetes mellitus (E10-E14)	1,359	2.6	8.5
7	Cerebrovascular diseases (I60-I69)	1,347	2.6	8.5
8	Chronic lower respiratory diseases (J40-J47)	1,129	2.2	7.1
9	Influenza and pneumonia (J09-J18)	766	1.5	4.8
10	Viral hepatitis (B15-B19)	593	1.1	3.7
11	Septicemia (A40-A41)	556	1.1	3.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	551	1.0	3.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	543	1.0	3.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	462	0.9	2.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	251	0.5	1.6
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	251	0.5	1.6
...	All other causes (Residual)	8,307	15.8	52.2
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	78,259	100.0	494.4
1	Malignant neoplasms (C00-C97)	23,588	30.1	149.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,668	20.0	99.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,441	9.5	47.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,937	5.0	24.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,601	4.6	22.8
6	Chronic lower respiratory diseases (J40-J47)	2,570	3.3	16.2
7	Diabetes mellitus (E10-E14)	2,195	2.8	13.9
8	Cerebrovascular diseases (I60-I69)	2,077	2.7	13.1
9	Viral hepatitis (B15-B19)	1,178	1.5	7.4
10	Influenza and pneumonia (J09-J18)	1,133	1.4	7.2
11	Septicemia (A40-A41)	948	1.2	6.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	689	0.9	4.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	472	0.6	3.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	436	0.6	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	432	0.6	2.7
...	All other causes (Residual)	11,894	15.2	75.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	99,131	100.0	696.7
1	Malignant neoplasms (C00-C97)	34,413	34.7	241.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,015	21.2	147.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,706	5.8	40.1
4	Chronic lower respiratory diseases (J40-J47)	4,416	4.5	31.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,721	3.8	26.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,121	3.1	21.9
7	Diabetes mellitus (E10-E14)	3,000	3.0	21.1
8	Cerebrovascular diseases (I60-I69)	2,747	2.8	19.3
9	Influenza and pneumonia (J09-J18)	1,403	1.4	9.9
10	Viral hepatitis (B15-B19)	1,331	1.3	9.4
11	Septicemia (A40-A41)	1,310	1.3	9.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,106	1.1	7.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	565	0.6	4.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	462	0.5	3.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	391	0.4	2.7
...	All other causes (Residual)	14,424	14.6	101.4
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	125,244	100.0	1,021.0
1	Malignant neoplasms (C00-C97)	47,398	37.8	386.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,381	21.9	223.2
3	Chronic lower respiratory diseases (J40-J47)	7,842	6.3	63.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,305	3.4	35.1
5	Diabetes mellitus (E10-E14)	4,289	3.4	35.0
6	Cerebrovascular diseases (I60-I69)	3,823	3.1	31.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,049	2.4	24.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,100	1.7	17.1
9	Septicemia (A40-A41)	1,781	1.4	14.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,665	1.3	13.6
11	Influenza and pneumonia (J09-J18)	1,618	1.3	13.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	846	0.7	6.9
13	Viral hepatitis (B15-B19)	726	0.6	5.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	656	0.5	5.3
15	Aortic aneurysm and dissection (I71)	556	0.4	4.5
...	All other causes (Residual)	17,209	13.7	140.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	144,557	100.0	1,556.9
1	Malignant neoplasms (C00-C97)	55,213	38.2	594.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,952	21.4	333.4
3	Chronic lower respiratory diseases (J40-J47)	12,091	8.4	130.2
4	Cerebrovascular diseases (I60-I69)	5,079	3.5	54.7
5	Diabetes mellitus (E10-E14)	4,647	3.2	50.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,690	2.6	39.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,284	1.6	24.6
8	Septicemia (A40-A41)	2,199	1.5	23.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,185	1.5	23.5
10	Influenza and pneumonia (J09-J18)	1,988	1.4	21.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,508	1.0	16.2
12	Alzheimer's disease (G30)	948	0.7	10.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	934	0.6	10.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	873	0.6	9.4
15	Aortic aneurysm and dissection (I71)	806	0.6	8.7
...	All other causes (Residual)	19,160	13.3	206.4
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	170,975	100.0	2,407.8
1	Malignant neoplasms (C00-C97)	59,477	34.8	837.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,514	21.9	528.3
3	Chronic lower respiratory diseases (J40-J47)	16,009	9.4	225.4
4	Cerebrovascular diseases (I60-I69)	7,626	4.5	107.4
5	Diabetes mellitus (E10-E14)	5,304	3.1	74.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,621	2.1	51.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,153	1.8	44.4
8	Influenza and pneumonia (J09-J18)	2,705	1.6	38.1
9	Septicemia (A40-A41)	2,618	1.5	36.9
10	Alzheimer's disease (G30)	2,521	1.5	35.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,820	1.1	25.6
12	Parkinson's disease (G20-G21)	1,465	0.9	20.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,247	0.7	17.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,161	0.7	16.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,131	0.7	15.9
...	All other causes (Residual)	23,603	13.8	332.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	225,470	100.0	3,830.2
1	Malignant neoplasms (C00-C97)	65,847	29.2	1,118.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	52,624	23.3	894.0
3	Chronic lower respiratory diseases (J40-J47)	20,518	9.1	348.6
4	Cerebrovascular diseases (I60-I69)	12,197	5.4	207.2
5	Alzheimer's disease (G30)	6,426	2.9	109.2
6	Diabetes mellitus (E10-E14)	6,268	2.8	106.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,865	2.2	82.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,718	2.1	80.1
9	Influenza and pneumonia (J09-J18)	4,365	1.9	74.2
10	Septicemia (A40-A41)	3,591	1.6	61.0
11	Parkinson's disease (G20-G21)	3,199	1.4	54.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,872	0.8	31.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,657	0.7	28.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,596	0.7	27.1
15	Pneumonitis due to solids and liquids (J69)	1,445	0.6	24.5
...	All other causes (Residual)	34,282	15.2	582.4
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	299,194	100.0	6,218.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	77,762	26.0	1,616.2
2	Malignant neoplasms (C00-C97)	67,135	22.4	1,395.4
3	Chronic lower respiratory diseases (J40-J47)	24,011	8.0	499.1
4	Cerebrovascular diseases (I60-I69)	18,939	6.3	393.6
5	Alzheimer's disease (G30)	13,879	4.6	288.5
6	Diabetes mellitus (E10-E14)	7,491	2.5	155.7
7	Influenza and pneumonia (J09-J18)	6,992	2.3	145.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,938	2.3	144.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,677	2.2	138.8
10	Parkinson's disease (G20-G21)	5,080	1.7	105.6
11	Septicemia (A40-A41)	4,538	1.5	94.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,070	1.0	63.8
13	Pneumonitis due to solids and liquids (J69)	2,491	0.8	51.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,323	0.8	48.3
15	Aortic aneurysm and dissection (I71)	1,681	0.6	34.9
...	All other causes (Residual)	50,187	16.8	1,043.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	322,231	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	95,182	29.5	---
2	Malignant neoplasms (C00-C97)	50,609	15.7	---
3	Cerebrovascular diseases (I60-I69)	23,152	7.2	---
4	Alzheimer's disease (G30)	20,452	6.3	---
5	Chronic lower respiratory diseases (J40-J47)	19,935	6.2	---
6	Influenza and pneumonia (J09-J18)	8,786	2.7	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,632	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,467	2.3	---
9	Diabetes mellitus (E10-E14)	6,897	2.1	---
10	Parkinson's disease (G20-G21)	4,915	1.5	---
11	Septicemia (A40-A41)	4,475	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,968	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,285	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,416	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,588	0.5	---
...	All other causes (Residual)	61,472	19.1	---
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	217,564	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,016	33.1	---
2	Malignant neoplasms (C00-C97)	22,205	10.2	---
3	Alzheimer's disease (G30)	16,535	7.6	---
4	Cerebrovascular diseases (I60-I69)	16,087	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	9,992	4.6	---
6	Influenza and pneumonia (J09-J18)	7,205	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,235	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,830	2.2	---
9	Diabetes mellitus (E10-E14)	3,662	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,269	1.5	---
11	Septicemia (A40-A41)	2,607	1.2	---
12	Pneumonitis due to solids and liquids (J69)	2,409	1.1	---
13	Parkinson's disease (G20-G21)	2,283	1.0	---
14	Atherosclerosis (I70)	1,341	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,287	0.6	---
...	All other causes (Residual)	46,601	21.4	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	85,674	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,447	35.5	---
2	Alzheimer's disease (G30)	6,971	8.1	---
3	Cerebrovascular diseases (I60-I69)	6,119	7.1	---
4	Malignant neoplasms (C00-C97)	5,663	6.6	---
5	Influenza and pneumonia (J09-J18)	3,358	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	2,963	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,012	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,814	2.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,504	1.8	---
10	Diabetes mellitus (E10-E14)	1,127	1.3	---
11	Pneumonitis due to solids and liquids (J69)	1,027	1.2	---
12	Septicemia (A40-A41)	874	1.0	---
13	Atherosclerosis (I70)	683	0.8	---
14	Parkinson's disease (G20-G21)	507	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	386	0.5	---
...	All other causes (Residual)	20,219	23.6	---
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	17,004	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,343	37.3	---
2	Alzheimer's disease (G30)	1,261	7.4	---
3	Cerebrovascular diseases (I60-I69)	1,068	6.3	---
4	Influenza and pneumonia (J09-J18)	783	4.6	---
5	Malignant neoplasms (C00-C97)	660	3.9	---
6	Chronic lower respiratory diseases (J40-J47)	396	2.3	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	370	2.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	357	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	330	1.9	---
10	Atherosclerosis (I70)	217	1.3	---
11	Pneumonitis due to solids and liquids (J69)	210	1.2	---
12	Diabetes mellitus (E10-E14)	159	0.9	---
13	Septicemia (A40-A41)	116	0.7	---
14	Anemias (D50-D64)	73	0.4	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	59	0.3	---
...	All other causes (Residual)	4,602	27.1	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	959,014	100.0	965.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,489	25.8	249.2
2	Malignant neoplasms (C00-C97)	239,301	25.0	241.0
3	Chronic lower respiratory diseases (J40-J47)	57,370	6.0	57.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56,607	5.9	57.0
5	Cerebrovascular diseases (I60-I69)	39,912	4.2	40.2
6	Diabetes mellitus (E10-E14)	24,993	2.6	25.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24,248	2.5	24.4
8	Alzheimer's disease (G30)	21,087	2.2	21.2
9	Influenza and pneumonia (J09-J18)	19,736	2.1	19.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,735	1.8	17.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14,494	1.5	14.6
12	Septicemia (A40-A41)	12,167	1.3	12.3
13	Parkinson's disease (G20-G21)	10,919	1.1	11.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,316	0.8	7.4
15	Pneumonitis due to solids and liquids (J69)	7,068	0.7	7.1
...	All other causes (Residual)	158,572	16.5	159.7
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,262	100.0	27.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	500	39.6	10.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	8.8	2.4
3	Malignant neoplasms (C00-C97)	106	8.4	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	5.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.2	0.9
6	Influenza and pneumonia (J09-J18)	32	2.5	0.7
7	Certain conditions originating in the perinatal period (P00-P96)	18	1.4	*
8	Septicemia (A40-A41)	17	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.3	*
10	Cerebrovascular diseases (I60-I69)	15	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
12	Anemias (D50-D64)	8	0.6	*
12	Meningitis (G00,G03)	8	0.6	*
14	Meningococcal infection (A39)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
...	All other causes (Residual)	298	23.6	6.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-9 years				
...	All causes	719	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	33.1	4.0
2	Malignant neoplasms (C00-C97)	149	20.7	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	9.0	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	3.9	0.5
5	Influenza and pneumonia (J09-J18)	25	3.5	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	2.2	*
7	Chronic lower respiratory diseases (J40-J47)	12	1.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.1	*
8	Cerebrovascular diseases (I60-I69)	8	1.1	*
10	Septicemia (A40-A41)	6	0.8	*
11	Anemias (D50-D64)	4	0.6	*
11	Diabetes mellitus (E10-E14)	4	0.6	*
13	Meningitis (G00,G03)	3	0.4	*
14	Meningococcal infection (A39)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Diseases of appendix (K35-K38)	2	0.3	*
14	Hernia (K40-K46)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	139	19.3	2.3
Non-Hispanic, white, male, 10-14 years				
...	All causes	965	100.0	16.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	336	34.8	5.6
2	Malignant neoplasms (C00-C97)	128	13.3	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	100	10.4	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	4.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	4.1	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	3.2	0.5
7	Influenza and pneumonia (J09-J18)	26	2.7	0.4
8	Cerebrovascular diseases (I60-I69)	18	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.5	*
10	Chronic lower respiratory diseases (J40-J47)	6	0.6	*
11	Septicemia (A40-A41)	5	0.5	*
12	Meningitis (G00,G03)	4	0.4	*
13	Aortic aneurysm and dissection (I71)	3	0.3	*
13	Diseases of appendix (K35-K38)	3	0.3	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
...	All other causes (Residual)	203	21.0	3.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 15-19 years				
...	All causes	4,355	100.0	65.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,219	51.0	33.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	930	21.4	14.0
3	Malignant neoplasms (C00-C97)	211	4.8	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	190	4.4	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	2.4	1.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	1.5	1.0
7	Influenza and pneumonia (J09-J18)	50	1.1	0.8
8	Chronic lower respiratory diseases (J40-J47)	21	0.5	0.3
9	Septicemia (A40-A41)	19	0.4	*
10	Cerebrovascular diseases (I60-I69)	18	0.4	*
11	Diabetes mellitus (E10-E14)	16	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.1	*
15	Pneumonitis due to solids and liquids (J69)	5	0.1	*
...	All other causes (Residual)	480	11.0	7.2
Non-Hispanic, white, male, 20-24 years				
...	All causes	7,963	100.0	117.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,082	51.3	60.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,603	20.1	23.6
3	Malignant neoplasms (C00-C97)	367	4.6	5.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	346	4.3	5.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	224	2.8	3.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	1.1	1.3
7	Influenza and pneumonia (J09-J18)	79	1.0	1.2
8	Diabetes mellitus (E10-E14)	37	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	36	0.5	0.5
10	Septicemia (A40-A41)	26	0.3	0.4
11	Legal intervention (Y35,Y89.0)	21	0.3	0.3
12	Chronic lower respiratory diseases (J40-J47)	20	0.3	0.3
13	Pneumonitis due to solids and liquids (J69)	16	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.2	*
...	All other causes (Residual)	985	12.4	14.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-29 years				
...	All causes	8,514	100.0	128.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,954	46.4	59.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,631	19.2	24.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	439	5.2	6.6
4	Malignant neoplasms (C00-C97)	434	5.1	6.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	332	3.9	5.0
6	Influenza and pneumonia (J09-J18)	119	1.4	1.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	0.9	1.2
8	Diabetes mellitus (E10-E14)	68	0.8	1.0
9	Cerebrovascular diseases (I60-I69)	51	0.6	0.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	42	0.5	0.6
11	Septicemia (A40-A41)	41	0.5	0.6
12	Chronic lower respiratory diseases (J40-J47)	33	0.4	0.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	26	0.3	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.2	0.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.2	*
15	Pneumonitis due to solids and liquids (J69)	19	0.2	*
...	All other causes (Residual)	1,206	14.2	18.2
Non-Hispanic, white, male, 30-34 years				
...	All causes	8,651	100.0	146.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,332	38.5	56.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,587	18.3	26.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	755	8.7	12.8
4	Malignant neoplasms (C00-C97)	661	7.6	11.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	294	3.4	5.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	109	1.3	1.8
7	Influenza and pneumonia (J09-J18)	106	1.2	1.8
8	Diabetes mellitus (E10-E14)	105	1.2	1.8
9	Cerebrovascular diseases (I60-I69)	85	1.0	1.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	72	0.8	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	0.8	1.1
12	Septicemia (A40-A41)	51	0.6	0.9
13	Aortic aneurysm and dissection (I71)	33	0.4	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.4	0.6
15	Chronic lower respiratory diseases (J40-J47)	30	0.3	0.5
...	All other causes (Residual)	1,330	15.4	22.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 35-39 years				
...	All causes	11,621	100.0	183.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,475	29.9	54.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,929	16.6	30.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,566	13.5	24.7
4	Malignant neoplasms (C00-C97)	1,173	10.1	18.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	307	2.6	4.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	286	2.5	4.5
7	Diabetes mellitus (E10-E14)	224	1.9	3.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	174	1.5	2.7
9	Cerebrovascular diseases (I60-I69)	164	1.4	2.6
10	Influenza and pneumonia (J09-J18)	156	1.3	2.5
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	94	0.8	1.5
12	Septicemia (A40-A41)	83	0.7	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	0.6	1.1
14	Aortic aneurysm and dissection (I71)	55	0.5	0.9
15	Chronic lower respiratory diseases (J40-J47)	51	0.4	0.8
...	All other causes (Residual)	1,812	15.6	28.5
Non-Hispanic, white, male, 40-44 years				
...	All causes	18,213	100.0	265.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,114	22.6	59.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,280	18.0	47.8
3	Malignant neoplasms (C00-C97)	2,442	13.4	35.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,280	12.5	33.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	738	4.1	10.7
6	Diabetes mellitus (E10-E14)	463	2.5	6.7
7	Cerebrovascular diseases (I60-I69)	352	1.9	5.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	351	1.9	5.1
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	309	1.7	4.5
10	Influenza and pneumonia (J09-J18)	236	1.3	3.4
11	Septicemia (A40-A41)	168	0.9	2.4
12	Chronic lower respiratory diseases (J40-J47)	152	0.8	2.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	139	0.8	2.0
14	Viral hepatitis (B15-B19)	124	0.7	1.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	0.6	1.7
...	All other causes (Residual)	2,950	16.2	43.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-49 years				
...	All causes	32,242	100.0	407.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,986	21.7	88.4
2	Malignant neoplasms (C00-C97)	6,173	19.1	78.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,281	16.4	66.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,732	8.5	34.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,581	4.9	20.0
6	Diabetes mellitus (E10-E14)	873	2.7	11.0
7	Cerebrovascular diseases (I60-I69)	730	2.3	9.2
8	Chronic lower respiratory diseases (J40-J47)	526	1.6	6.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	459	1.4	5.8
10	Influenza and pneumonia (J09-J18)	397	1.2	5.0
11	Viral hepatitis (B15-B19)	392	1.2	5.0
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	355	1.1	4.5
13	Septicemia (A40-A41)	285	0.9	3.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	255	0.8	3.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	171	0.5	2.2
...	All other causes (Residual)	5,046	15.7	63.8
Non-Hispanic, white, male, 50-54 years				
...	All causes	48,585	100.0	620.2
1	Malignant neoplasms (C00-C97)	12,314	25.3	157.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,579	23.8	147.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,934	10.2	63.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,948	6.1	37.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,496	5.1	31.9
6	Diabetes mellitus (E10-E14)	1,411	2.9	18.0
7	Chronic lower respiratory diseases (J40-J47)	1,280	2.6	16.3
8	Cerebrovascular diseases (I60-I69)	1,173	2.4	15.0
9	Viral hepatitis (B15-B19)	850	1.7	10.9
10	Influenza and pneumonia (J09-J18)	605	1.2	7.7
11	Septicemia (A40-A41)	528	1.1	6.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	426	0.9	5.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	394	0.8	5.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	313	0.6	4.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	297	0.6	3.8
...	All other causes (Residual)	7,037	14.5	89.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 55-59 years				
...	All causes	61,732	100.0	883.5
1	Malignant neoplasms (C00-C97)	19,039	30.8	272.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,318	24.8	219.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,866	6.3	55.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,664	4.3	38.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,365	3.8	33.8
6	Chronic lower respiratory diseases (J40-J47)	2,319	3.8	33.2
7	Diabetes mellitus (E10-E14)	1,869	3.0	26.7
8	Cerebrovascular diseases (I60-I69)	1,599	2.6	22.9
9	Viral hepatitis (B15-B19)	1,004	1.6	14.4
10	Influenza and pneumonia (J09-J18)	788	1.3	11.3
11	Septicemia (A40-A41)	680	1.1	9.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	663	1.1	9.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	391	0.6	5.6
14	Human immunodeficiency virus (HIV) disease (B20-B24)	342	0.6	4.9
15	Aortic aneurysm and dissection (I71)	280	0.5	4.0
...	All other causes (Residual)	8,545	13.8	122.3
Non-Hispanic, white, male, 60-64 years				
...	All causes	75,143	100.0	1,261.6
1	Malignant neoplasms (C00-C97)	26,197	34.9	439.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,137	25.5	321.3
3	Chronic lower respiratory diseases (J40-J47)	3,987	5.3	66.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,975	4.0	49.9
5	Diabetes mellitus (E10-E14)	2,669	3.6	44.8
6	Cerebrovascular diseases (I60-I69)	2,141	2.8	35.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,133	2.8	35.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,621	2.2	27.2
9	Septicemia (A40-A41)	953	1.3	16.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	939	1.2	15.8
11	Influenza and pneumonia (J09-J18)	901	1.2	15.1
12	Viral hepatitis (B15-B19)	519	0.7	8.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	515	0.7	8.6
14	Aortic aneurysm and dissection (I71)	405	0.5	6.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	381	0.5	6.4
...	All other causes (Residual)	9,670	12.9	162.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-69 years				
...	All causes	83,839	100.0	1,904.6
1	Malignant neoplasms (C00-C97)	30,617	36.5	695.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,496	24.4	465.6
3	Chronic lower respiratory diseases (J40-J47)	6,034	7.2	137.1
4	Diabetes mellitus (E10-E14)	2,800	3.3	63.6
5	Cerebrovascular diseases (I60-I69)	2,787	3.3	63.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,412	2.9	54.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,498	1.8	34.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,234	1.5	28.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,206	1.4	27.4
10	Septicemia (A40-A41)	1,130	1.3	25.7
11	Influenza and pneumonia (J09-J18)	1,116	1.3	25.4
12	Aortic aneurysm and dissection (I71)	550	0.7	12.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	527	0.6	12.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	526	0.6	11.9
15	Alzheimer's disease (G30)	434	0.5	9.9
...	All other causes (Residual)	10,472	12.5	237.9
Non-Hispanic, white, male, 70-74 years				
...	All causes	94,883	100.0	2,910.5
1	Malignant neoplasms (C00-C97)	32,407	34.2	994.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,231	24.5	712.6
3	Chronic lower respiratory diseases (J40-J47)	7,892	8.3	242.1
4	Cerebrovascular diseases (I60-I69)	3,910	4.1	119.9
5	Diabetes mellitus (E10-E14)	3,002	3.2	92.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,176	2.3	66.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,736	1.8	53.3
8	Influenza and pneumonia (J09-J18)	1,479	1.6	45.4
9	Septicemia (A40-A41)	1,329	1.4	40.8
10	Alzheimer's disease (G30)	1,105	1.2	33.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,055	1.1	32.4
12	Parkinson's disease (G20-G21)	991	1.0	30.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	940	1.0	28.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	668	0.7	20.5
15	Aortic aneurysm and dissection (I71)	648	0.7	19.9
...	All other causes (Residual)	12,314	13.0	377.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 75-79 years				
...	All causes	118,658	100.0	4,627.4
1	Malignant neoplasms (C00-C97)	35,566	30.0	1,387.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,873	25.2	1,165.0
3	Chronic lower respiratory diseases (J40-J47)	9,925	8.4	387.1
4	Cerebrovascular diseases (I60-I69)	5,644	4.8	220.1
5	Diabetes mellitus (E10-E14)	3,307	2.8	129.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,797	2.4	109.1
7	Alzheimer's disease (G30)	2,556	2.2	99.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,498	2.1	97.4
9	Influenza and pneumonia (J09-J18)	2,320	2.0	90.5
10	Parkinson's disease (G20-G21)	2,062	1.7	80.4
11	Septicemia (A40-A41)	1,749	1.5	68.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	949	0.8	37.0
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	862	0.7	33.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	852	0.7	33.2
15	Pneumonitis due to solids and liquids (J69)	850	0.7	33.1
...	All other causes (Residual)	16,848	14.2	657.0
Non-Hispanic, white, male, 80-84 years				
...	All causes	143,895	100.0	7,501.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,346	27.3	2,051.3
2	Malignant neoplasms (C00-C97)	35,012	24.3	1,825.3
3	Chronic lower respiratory diseases (J40-J47)	11,385	7.9	593.6
4	Cerebrovascular diseases (I60-I69)	7,672	5.3	400.0
5	Alzheimer's disease (G30)	4,993	3.5	260.3
6	Diabetes mellitus (E10-E14)	3,604	2.5	187.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,579	2.5	186.6
8	Influenza and pneumonia (J09-J18)	3,443	2.4	179.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,412	2.4	177.9
10	Parkinson's disease (G20-G21)	3,043	2.1	158.6
11	Septicemia (A40-A41)	2,097	1.5	109.3
12	Pneumonitis due to solids and liquids (J69)	1,380	1.0	71.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,249	0.9	65.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,249	0.9	65.1
15	Aortic aneurysm and dissection (I71)	915	0.6	47.7
...	All other causes (Residual)	21,516	15.0	1,121.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85-89 years				
...	All causes	133,842	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,043	30.7	---
2	Malignant neoplasms (C00-C97)	24,420	18.2	---
3	Chronic lower respiratory diseases (J40-J47)	8,873	6.6	---
4	Cerebrovascular diseases (I60-I69)	7,798	5.8	---
5	Alzheimer's disease (G30)	6,338	4.7	---
6	Influenza and pneumonia (J09-J18)	3,853	2.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,507	2.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,414	2.6	---
9	Diabetes mellitus (E10-E14)	2,937	2.2	---
10	Parkinson's disease (G20-G21)	2,751	2.1	---
11	Septicemia (A40-A41)	1,810	1.4	---
12	Pneumonitis due to solids and liquids (J69)	1,744	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,280	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,154	0.9	---
15	Aortic aneurysm and dissection (I71)	689	0.5	---
...	All other causes (Residual)	22,231	16.6	---
Non-Hispanic, white, male, 90-94 years				
...	All causes	73,568	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,095	34.1	---
2	Malignant neoplasms (C00-C97)	9,749	13.3	---
3	Cerebrovascular diseases (I60-I69)	4,329	5.9	---
4	Alzheimer's disease (G30)	4,095	5.6	---
5	Chronic lower respiratory diseases (J40-J47)	3,818	5.2	---
6	Influenza and pneumonia (J09-J18)	2,739	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,062	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,953	2.7	---
9	Diabetes mellitus (E10-E14)	1,262	1.7	---
10	Parkinson's disease (G20-G21)	1,172	1.6	---
11	Pneumonitis due to solids and liquids (J69)	1,111	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	906	1.2	---
13	Septicemia (A40-A41)	892	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	530	0.7	---
15	Atherosclerosis (I70)	402	0.5	---
...	All other causes (Residual)	13,453	18.3	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, white, male, 95-99 years				
...	All causes	21,051	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,802	37.1	---
2	Malignant neoplasms (C00-C97)	1,944	9.2	---
3	Cerebrovascular diseases (I60-I69)	1,203	5.7	---
4	Alzheimer's disease (G30)	1,130	5.4	---
5	Influenza and pneumonia (J09-J18)	1,018	4.8	---
6	Chronic lower respiratory diseases (J40-J47)	891	4.2	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	619	2.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	555	2.6	---
9	Pneumonitis due to solids and liquids (J69)	376	1.8	---
10	Diabetes mellitus (E10-E14)	307	1.5	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	290	1.4	---
12	Septicemia (A40-A41)	228	1.1	---
13	Parkinson's disease (G20-G21)	190	0.9	---
14	Atherosclerosis (I70)	180	0.9	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	125	0.6	---
...	All other causes (Residual)	4,193	19.9	---
Non-Hispanic, white, male, 100 years and over				
...	All causes	2,737	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,022	37.3	---
2	Influenza and pneumonia (J09-J18)	189	6.9	---
3	Malignant neoplasms (C00-C97)	165	6.0	---
4	Cerebrovascular diseases (I60-I69)	141	5.2	---
5	Alzheimer's disease (G30)	127	4.6	---
6	Chronic lower respiratory diseases (J40-J47)	95	3.5	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	87	3.2	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	51	1.9	---
10	Pneumonitis due to solids and liquids (J69)	48	1.8	---
11	Diabetes mellitus (E10-E14)	31	1.1	---
12	Atherosclerosis (I70)	28	1.0	---
13	Septicemia (A40-A41)	27	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.6	---
15	Parkinson's disease (G20-G21)	12	0.4	---
...	All other causes (Residual)	623	22.8	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, white, female, all ages				
...	All causes	985,592	100.0	958.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	238,290	24.2	231.7
2	Malignant neoplasms (C00-C97)	217,888	22.1	211.9
3	Chronic lower respiratory diseases (J40-J47)	65,235	6.6	63.4
4	Cerebrovascular diseases (I60-I69)	61,791	6.3	60.1
5	Alzheimer's disease (G30)	48,608	4.9	47.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,809	3.5	33.8
7	Influenza and pneumonia (J09-J18)	23,016	2.3	22.4
8	Diabetes mellitus (E10-E14)	22,858	2.3	22.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,935	1.8	17.4
10	Septicemia (A40-A41)	14,366	1.5	14.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,236	1.1	10.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,026	0.8	7.8
13	Parkinson's disease (G20-G21)	7,635	0.8	7.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,565	0.7	6.4
15	Pneumonitis due to solids and liquids (J69)	6,421	0.7	6.2
...	All other causes (Residual)	200,913	20.4	195.4
Non-Hispanic, white, female, 1-4 years				
...	All causes	919	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	29.7	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	12.9	2.7
3	Malignant neoplasms (C00-C97)	74	8.1	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	7.9	1.6
5	Influenza and pneumonia (J09-J18)	43	4.7	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.2	0.7
7	Septicemia (A40-A41)	13	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.3	*
8	Cerebrovascular diseases (I60-I69)	12	1.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	9	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Meningitis (G00,G03)	3	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
...	All other causes (Residual)	240	26.1	5.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 5-9 years				
...	All causes	517	100.0	9.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	26.9	2.5
2	Malignant neoplasms (C00-C97)	106	20.5	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	8.7	0.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.6	0.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.3	0.4
6	Influenza and pneumonia (J09-J18)	19	3.7	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	2.1	*
8	Septicemia (A40-A41)	8	1.5	*
9	Cerebrovascular diseases (I60-I69)	6	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
12	Diabetes mellitus (E10-E14)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
14	Anemias (D50-D64)	1	0.2	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.2	*
...	All other causes (Residual)	111	21.5	2.0
Non-Hispanic, white, female, 10-14 years				
...	All causes	666	100.0	11.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	27.3	3.2
2	Malignant neoplasms (C00-C97)	110	16.5	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	8.3	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	5.6	0.6
5	Influenza and pneumonia (J09-J18)	34	5.1	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	4.1	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	2.3	*
8	Cerebrovascular diseases (I60-I69)	11	1.7	*
9	Septicemia (A40-A41)	9	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.9	*
10	Diabetes mellitus (E10-E14)	6	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.8	*
13	Anemias (D50-D64)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
15	Meningitis (G00,G03)	3	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
...	All other causes (Residual)	155	23.3	2.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-19 years				
...	All causes	2,008	100.0	31.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,018	50.7	16.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	222	11.1	3.5
3	Malignant neoplasms (C00-C97)	149	7.4	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	4.5	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	3.0	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	2.8	0.9
7	Influenza and pneumonia (J09-J18)	37	1.8	0.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	24	1.2	0.4
9	Chronic lower respiratory diseases (J40-J47)	16	0.8	*
10	Cerebrovascular diseases (I60-I69)	9	0.4	*
11	Septicemia (A40-A41)	8	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.3	*
13	Diabetes mellitus (E10-E14)	6	0.3	*
14	Anemias (D50-D64)	5	0.2	*
15	Pneumonitis due to solids and liquids (J69)	4	0.2	*
...	All other causes (Residual)	296	14.7	4.7
Non-Hispanic, white, female, 20-24 years				
...	All causes	2,824	100.0	43.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,245	44.1	19.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	291	10.3	4.5
3	Malignant neoplasms (C00-C97)	218	7.7	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	4.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	4.0	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	69	2.4	1.1
7	Influenza and pneumonia (J09-J18)	57	2.0	0.9
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	1.9	0.8
9	Cerebrovascular diseases (I60-I69)	34	1.2	0.5
10	Diabetes mellitus (E10-E14)	30	1.1	0.5
11	Septicemia (A40-A41)	20	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	14	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.4	*
14	Meningitis (G00,G03)	8	0.3	*
14	Pneumonitis due to solids and liquids (J69)	8	0.3	*
...	All other causes (Residual)	524	18.6	8.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 25-29 years				
...	All causes	3,611	100.0	55.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,308	36.2	20.3
2	Malignant neoplasms (C00-C97)	386	10.7	6.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	363	10.1	5.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	172	4.8	2.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	166	4.6	2.6
6	Influenza and pneumonia (J09-J18)	99	2.7	1.5
7	Pregnancy, childbirth and the puerperium (O00-O99)	83	2.3	1.3
8	Diabetes mellitus (E10-E14)	67	1.9	1.0
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.6	0.9
10	Septicemia (A40-A41)	47	1.3	0.7
11	Cerebrovascular diseases (I60-I69)	44	1.2	0.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	0.7	0.4
13	Chronic lower respiratory diseases (J40-J47)	24	0.7	0.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.7	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.5	*
15	Pneumonitis due to solids and liquids (J69)	17	0.5	*
...	All other causes (Residual)	713	19.7	11.0
Non-Hispanic, white, female, 30-34 years				
...	All causes	4,427	100.0	75.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,293	29.2	22.1
2	Malignant neoplasms (C00-C97)	725	16.4	12.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	402	9.1	6.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	322	7.3	5.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	3.1	2.4
6	Influenza and pneumonia (J09-J18)	110	2.5	1.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	82	1.9	1.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	67	1.5	1.1
9	Cerebrovascular diseases (I60-I69)	65	1.5	1.1
10	Diabetes mellitus (E10-E14)	63	1.4	1.1
11	Septicemia (A40-A41)	53	1.2	0.9
12	Chronic lower respiratory diseases (J40-J47)	49	1.1	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	1.0	0.8
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	0.8	0.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	28	0.6	0.5
...	All other causes (Residual)	949	21.4	16.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-39 years				
...	All causes	6,666	100.0	105.9
1	Malignant neoplasms (C00-C97)	1,507	22.6	23.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,479	22.2	23.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	675	10.1	10.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	559	8.4	8.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	177	2.7	2.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	2.2	2.3
7	Cerebrovascular diseases (I60-I69)	146	2.2	2.3
8	Diabetes mellitus (E10-E14)	133	2.0	2.1
9	Influenza and pneumonia (J09-J18)	111	1.7	1.8
10	Septicemia (A40-A41)	99	1.5	1.6
11	Chronic lower respiratory diseases (J40-J47)	73	1.1	1.2
12	Pregnancy, childbirth and the puerperium (O00-O99)	68	1.0	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	0.9	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	0.8	0.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	53	0.8	0.8
...	All other causes (Residual)	1,324	19.9	21.0
Non-Hispanic, white, female, 40-44 years				
...	All causes	11,236	100.0	164.1
1	Malignant neoplasms (C00-C97)	3,091	27.5	45.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,934	17.2	28.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,326	11.8	19.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	740	6.6	10.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	417	3.7	6.1
6	Cerebrovascular diseases (I60-I69)	300	2.7	4.4
7	Diabetes mellitus (E10-E14)	225	2.0	3.3
8	Influenza and pneumonia (J09-J18)	214	1.9	3.1
9	Chronic lower respiratory diseases (J40-J47)	193	1.7	2.8
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	180	1.6	2.6
11	Septicemia (A40-A41)	169	1.5	2.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	0.8	1.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	0.7	1.2
14	Viral hepatitis (B15-B19)	71	0.6	1.0
15	Human immunodeficiency virus (HIV) disease (B20-B24)	61	0.5	0.9
...	All other causes (Residual)	2,142	19.1	31.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-49 years				
...	All causes	20,239	100.0	253.1
1	Malignant neoplasms (C00-C97)	6,724	33.2	84.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,639	13.0	33.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,515	12.4	31.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	895	4.4	11.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	840	4.2	10.5
6	Cerebrovascular diseases (I60-I69)	617	3.0	7.7
7	Chronic lower respiratory diseases (J40-J47)	603	3.0	7.5
8	Diabetes mellitus (E10-E14)	486	2.4	6.1
9	Influenza and pneumonia (J09-J18)	369	1.8	4.6
10	Septicemia (A40-A41)	271	1.3	3.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	207	1.0	2.6
12	Viral hepatitis (B15-B19)	201	1.0	2.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	188	0.9	2.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	109	0.5	1.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	98	0.5	1.2
...	All other causes (Residual)	3,477	17.2	43.5
Non-Hispanic, white, female, 50-54 years				
...	All causes	29,674	100.0	371.2
1	Malignant neoplasms (C00-C97)	11,274	38.0	141.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,089	13.8	51.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,507	8.4	31.4
4	Chronic lower respiratory diseases (J40-J47)	1,290	4.3	16.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,105	3.7	13.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	989	3.3	12.4
7	Cerebrovascular diseases (I60-I69)	904	3.0	11.3
8	Diabetes mellitus (E10-E14)	784	2.6	9.8
9	Influenza and pneumonia (J09-J18)	528	1.8	6.6
10	Septicemia (A40-A41)	420	1.4	5.3
11	Viral hepatitis (B15-B19)	328	1.1	4.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	263	0.9	3.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	135	0.5	1.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	132	0.4	1.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	0.4	1.5
...	All other causes (Residual)	4,803	16.2	60.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-59 years				
...	All causes	37,399	100.0	516.4
1	Malignant neoplasms (C00-C97)	15,374	41.1	212.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,697	15.2	78.7
3	Chronic lower respiratory diseases (J40-J47)	2,097	5.6	29.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,840	4.9	25.4
5	Cerebrovascular diseases (I60-I69)	1,148	3.1	15.9
6	Diabetes mellitus (E10-E14)	1,131	3.0	15.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,057	2.8	14.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	756	2.0	10.4
9	Septicemia (A40-A41)	630	1.7	8.7
10	Influenza and pneumonia (J09-J18)	615	1.6	8.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	443	1.2	6.1
12	Viral hepatitis (B15-B19)	327	0.9	4.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	185	0.5	2.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	174	0.5	2.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	171	0.5	2.4
...	All other causes (Residual)	5,754	15.4	79.5
Non-Hispanic, white, female, 60-64 years				
...	All causes	50,101	100.0	793.9
1	Malignant neoplasms (C00-C97)	21,201	42.3	336.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,244	16.5	130.6
3	Chronic lower respiratory diseases (J40-J47)	3,855	7.7	61.1
4	Cerebrovascular diseases (I60-I69)	1,682	3.4	26.7
5	Diabetes mellitus (E10-E14)	1,620	3.2	25.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,330	2.7	21.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	916	1.8	14.5
8	Septicemia (A40-A41)	828	1.7	13.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	726	1.4	11.5
10	Influenza and pneumonia (J09-J18)	717	1.4	11.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	479	1.0	7.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	331	0.7	5.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	275	0.5	4.4
14	Alzheimer's disease (G30)	229	0.5	3.6
15	Viral hepatitis (B15-B19)	207	0.4	3.3
...	All other causes (Residual)	7,461	14.9	118.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 65-69 years				
...	All causes	60,718	100.0	1,243.5
1	Malignant neoplasms (C00-C97)	24,596	40.5	503.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,456	17.2	214.1
3	Chronic lower respiratory diseases (J40-J47)	6,057	10.0	124.0
4	Cerebrovascular diseases (I60-I69)	2,292	3.8	46.9
5	Diabetes mellitus (E10-E14)	1,847	3.0	37.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,278	2.1	26.2
7	Septicemia (A40-A41)	1,069	1.8	21.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	979	1.6	20.1
9	Influenza and pneumonia (J09-J18)	872	1.4	17.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	786	1.3	16.1
11	Alzheimer's disease (G30)	514	0.8	10.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	407	0.7	8.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	347	0.6	7.1
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	274	0.5	5.6
15	Aortic aneurysm and dissection (I71)	256	0.4	5.2
...	All other causes (Residual)	8,688	14.3	177.9
Non-Hispanic, white, female, 70-74 years				
...	All causes	76,092	100.0	1,981.1
1	Malignant neoplasms (C00-C97)	27,070	35.6	704.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,283	18.8	371.9
3	Chronic lower respiratory diseases (J40-J47)	8,117	10.7	211.3
4	Cerebrovascular diseases (I60-I69)	3,716	4.9	96.7
5	Diabetes mellitus (E10-E14)	2,302	3.0	59.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,445	1.9	37.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,417	1.9	36.9
8	Alzheimer's disease (G30)	1,416	1.9	36.9
9	Septicemia (A40-A41)	1,289	1.7	33.6
10	Influenza and pneumonia (J09-J18)	1,226	1.6	31.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	765	1.0	19.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	607	0.8	15.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	493	0.6	12.8
14	Parkinson's disease (G20-G21)	474	0.6	12.3
15	Aortic aneurysm and dissection (I71)	365	0.5	9.5
...	All other causes (Residual)	11,107	14.6	289.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-79 years				
...	All causes	106,812	100.0	3,215.0
1	Malignant neoplasms (C00-C97)	30,281	28.3	911.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,751	21.3	684.8
3	Chronic lower respiratory diseases (J40-J47)	10,593	9.9	318.8
4	Cerebrovascular diseases (I60-I69)	6,553	6.1	197.2
5	Alzheimer's disease (G30)	3,870	3.6	116.5
6	Diabetes mellitus (E10-E14)	2,961	2.8	89.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,220	2.1	66.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,068	1.9	62.2
9	Influenza and pneumonia (J09-J18)	2,045	1.9	61.6
10	Septicemia (A40-A41)	1,842	1.7	55.4
11	Parkinson's disease (G20-G21)	1,137	1.1	34.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,026	1.0	30.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	744	0.7	22.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	708	0.7	21.3
15	Pneumonitis due to solids and liquids (J69)	595	0.6	17.9
...	All other causes (Residual)	17,418	16.3	524.3
Non-Hispanic, white, female, 80-84 years				
...	All causes	155,299	100.0	5,367.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,416	24.7	1,327.8
2	Malignant neoplasms (C00-C97)	32,123	20.7	1,110.3
3	Chronic lower respiratory diseases (J40-J47)	12,626	8.1	436.4
4	Cerebrovascular diseases (I60-I69)	11,267	7.3	389.4
5	Alzheimer's disease (G30)	8,886	5.7	307.1
6	Diabetes mellitus (E10-E14)	3,887	2.5	134.4
7	Influenza and pneumonia (J09-J18)	3,549	2.3	122.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,359	2.2	116.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,265	2.1	112.9
10	Septicemia (A40-A41)	2,441	1.6	84.4
11	Parkinson's disease (G20-G21)	2,037	1.3	70.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,821	1.2	62.9
13	Pneumonitis due to solids and liquids (J69)	1,111	0.7	38.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,074	0.7	37.1
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	830	0.5	28.7
...	All other causes (Residual)	28,607	18.4	988.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, white, female, 85-89 years				
...	All causes	188,389	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,139	28.7	---
2	Malignant neoplasms (C00-C97)	26,189	13.9	---
3	Cerebrovascular diseases (I60-I69)	15,354	8.2	---
4	Alzheimer's disease (G30)	14,114	7.5	---
5	Chronic lower respiratory diseases (J40-J47)	11,062	5.9	---
6	Influenza and pneumonia (J09-J18)	4,933	2.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,218	2.2	---
8	Diabetes mellitus (E10-E14)	3,960	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,960	2.1	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,688	1.4	---
11	Septicemia (A40-A41)	2,665	1.4	---
12	Parkinson's disease (G20-G21)	2,164	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,541	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,262	0.7	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1,008	0.5	---
...	All other causes (Residual)	39,132	20.8	---
Non-Hispanic, white, female, 90-94 years				
...	All causes	143,996	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	46,921	32.6	---
2	Malignant neoplasms (C00-C97)	12,456	8.7	---
3	Alzheimer's disease (G30)	12,440	8.6	---
4	Cerebrovascular diseases (I60-I69)	11,758	8.2	---
5	Chronic lower respiratory diseases (J40-J47)	6,174	4.3	---
6	Influenza and pneumonia (J09-J18)	4,466	3.1	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,282	2.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,768	1.9	---
9	Diabetes mellitus (E10-E14)	2,400	1.7	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,363	1.6	---
11	Septicemia (A40-A41)	1,715	1.2	---
12	Pneumonitis due to solids and liquids (J69)	1,298	0.9	---
13	Parkinson's disease (G20-G21)	1,111	0.8	---
14	Atherosclerosis (I70)	939	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	757	0.5	---
...	All other causes (Residual)	33,148	23.0	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 95-99 years				
...	All causes	64,623	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,645	35.0	---
2	Alzheimer's disease (G30)	5,841	9.0	---
3	Cerebrovascular diseases (I60-I69)	4,916	7.6	---
4	Malignant neoplasms (C00-C97)	3,719	5.8	---
5	Influenza and pneumonia (J09-J18)	2,340	3.6	---
6	Chronic lower respiratory diseases (J40-J47)	2,072	3.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,457	2.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,214	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,195	1.8	---
10	Diabetes mellitus (E10-E14)	820	1.3	---
11	Pneumonitis due to solids and liquids (J69)	651	1.0	---
12	Septicemia (A40-A41)	646	1.0	---
13	Atherosclerosis (I70)	503	0.8	---
14	Parkinson's disease (G20-G21)	317	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	261	0.4	---
...	All other causes (Residual)	16,026	24.8	---
Non-Hispanic, white, female, 100 years and over				
...	All causes	14,267	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,321	37.3	---
2	Alzheimer's disease (G30)	1,134	7.9	---
3	Cerebrovascular diseases (I60-I69)	927	6.5	---
4	Influenza and pneumonia (J09-J18)	594	4.2	---
5	Malignant neoplasms (C00-C97)	495	3.5	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	319	2.2	---
7	Chronic lower respiratory diseases (J40-J47)	301	2.1	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	243	1.7	---
10	Atherosclerosis (I70)	189	1.3	---
11	Pneumonitis due to solids and liquids (J69)	162	1.1	---
12	Diabetes mellitus (E10-E14)	128	0.9	---
13	Septicemia (A40-A41)	89	0.6	---
14	Anemias (D50-D64)	62	0.4	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	46	0.3	---
...	All other causes (Residual)	3,975	27.9	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	282,982	100.0	728.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,811	24.3	177.1
2	Malignant neoplasms (C00-C97)	63,967	22.6	164.6
3	Cerebrovascular diseases (I60-I69)	15,718	5.6	40.4
4	Diabetes mellitus (E10-E14)	11,833	4.2	30.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,810	4.2	30.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,727	3.1	22.5
7	Chronic lower respiratory diseases (J40-J47)	8,444	3.0	21.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,733	2.7	19.9
9	Septicemia (A40-A41)	6,131	2.2	15.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5,307	1.9	13.7
11	Influenza and pneumonia (J09-J18)	5,214	1.8	13.4
12	Alzheimer's disease (G30)	5,042	1.8	13.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,819	1.7	12.4
14	Certain conditions originating in the perinatal period (P00-P96)	4,456	1.6	11.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,518	0.9	6.5
...	All other causes (Residual)	52,452	18.5	135.0
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,039	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	27.7	11.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	13.8	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	8.4	3.5
4	Malignant neoplasms (C00-C97)	56	5.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	4.8	2.0
6	Chronic lower respiratory diseases (J40-J47)	33	3.2	1.3
7	Influenza and pneumonia (J09-J18)	29	2.8	1.2
8	Septicemia (A40-A41)	19	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	19	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	12	1.2	*
12	Cerebrovascular diseases (I60-I69)	9	0.9	*
13	Anemias (D50-D64)	8	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	8	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
...	All other causes (Residual)	260	25.0	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	563	100.0	18.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	32.9	6.2
2	Malignant neoplasms (C00-C97)	76	13.5	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	6.7	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	6.6	1.2
5	Chronic lower respiratory diseases (J40-J47)	29	5.2	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	4.1	0.8
6	Influenza and pneumonia (J09-J18)	23	4.1	0.8
8	Septicemia (A40-A41)	10	1.8	*
8	Cerebrovascular diseases (I60-I69)	10	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.6	*
11	Anemias (D50-D64)	5	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
13	Diabetes mellitus (E10-E14)	3	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
15	Meningitis (G00,G03)	2	0.4	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
15	Peptic ulcer (K25-K28)	2	0.4	*
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	98	17.4	3.3
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	703	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	192	27.3	6.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	9.5	2.2
3	Malignant neoplasms (C00-C97)	52	7.4	1.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	45	6.4	1.5
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	6.0	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	5.8	1.4
6	Chronic lower respiratory diseases (J40-J47)	41	5.8	1.4
8	Influenza and pneumonia (J09-J18)	22	3.1	0.7
9	Anemias (D50-D64)	14	2.0	*
10	Septicemia (A40-A41)	12	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.6	*
12	Cerebrovascular diseases (I60-I69)	8	1.1	*
13	Diabetes mellitus (E10-E14)	5	0.7	*
14	Diseases of appendix (K35-K38)	4	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	135	19.2	4.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,506	100.0	73.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,073	42.8	31.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	563	22.5	16.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	6.1	4.4
4	Malignant neoplasms (C00-C97)	115	4.6	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	4.2	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	2.2	1.6
7	Chronic lower respiratory diseases (J40-J47)	34	1.4	1.0
8	Influenza and pneumonia (J09-J18)	28	1.1	0.8
9	Anemias (D50-D64)	23	0.9	0.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	21	0.8	0.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	15	0.6	*
12	Septicemia (A40-A41)	14	0.6	*
12	Cerebrovascular diseases (I60-I69)	14	0.6	*
14	Diabetes mellitus (E10-E14)	13	0.5	*
15	Legal intervention (Y35,Y89.0)	12	0.5	*
...	All other causes (Residual)	269	10.7	7.9
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	4,096	100.0	128.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,687	41.2	52.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	746	18.2	23.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	281	6.9	8.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	234	5.7	7.3
5	Malignant neoplasms (C00-C97)	169	4.1	5.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	91	2.2	2.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	67	1.6	2.1
8	Influenza and pneumonia (J09-J18)	57	1.4	1.8
9	Chronic lower respiratory diseases (J40-J47)	51	1.2	1.6
10	Anemias (D50-D64)	43	1.0	1.3
11	Diabetes mellitus (E10-E14)	37	0.9	1.2
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.9	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	0.7	0.8
14	Cerebrovascular diseases (I60-I69)	25	0.6	0.8
15	Legal intervention (Y35,Y89.0)	18	0.4	*
...	All other causes (Residual)	526	12.8	16.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Percent of		
		Number	total deaths	Rate
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	4,385	100.0	145.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,315	30.0	43.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	867	19.8	28.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	339	7.7	11.2
4	Malignant neoplasms (C00-C97)	253	5.8	8.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	253	5.8	8.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	230	5.2	7.6
7	Diabetes mellitus (E10-E14)	82	1.9	2.7
8	Influenza and pneumonia (J09-J18)	71	1.6	2.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	66	1.5	2.2
10	Anemias (D50-D64)	54	1.2	1.8
11	Septicemia (A40-A41)	51	1.2	1.7
11	Cerebrovascular diseases (I60-I69)	51	1.2	1.7
13	Chronic lower respiratory diseases (J40-J47)	49	1.1	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	0.9	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	0.8	1.2
...	All other causes (Residual)	626	14.3	20.7
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	4,646	100.0	177.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	976	21.0	37.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	699	15.0	26.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	610	13.1	23.3
4	Malignant neoplasms (C00-C97)	373	8.0	14.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	366	7.9	14.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	210	4.5	8.0
7	Diabetes mellitus (E10-E14)	107	2.3	4.1
8	Cerebrovascular diseases (I60-I69)	92	2.0	3.5
9	Influenza and pneumonia (J09-J18)	72	1.5	2.7
10	Pregnancy, childbirth and the puerperium (O00-O99)	63	1.4	2.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	62	1.3	2.4
12	Anemias (D50-D64)	57	1.2	2.2
13	Septicemia (A40-A41)	56	1.2	2.1
14	Chronic lower respiratory diseases (J40-J47)	51	1.1	1.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	0.9	1.6
...	All other causes (Residual)	810	17.4	30.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	6,025	100.0	229.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,033	17.1	39.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	791	13.1	30.1
3	Malignant neoplasms (C00-C97)	719	11.9	27.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	683	11.3	26.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	561	9.3	21.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	197	3.3	7.5
7	Cerebrovascular diseases (I60-I69)	182	3.0	6.9
8	Diabetes mellitus (E10-E14)	177	2.9	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	124	2.1	4.7
10	Septicemia (A40-A41)	101	1.7	3.8
10	Influenza and pneumonia (J09-J18)	101	1.7	3.8
12	Chronic lower respiratory diseases (J40-J47)	83	1.4	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	1.1	2.5
14	Anemias (D50-D64)	56	0.9	2.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	56	0.9	2.1
...	All other causes (Residual)	1,094	18.2	41.6
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	8,926	100.0	339.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,830	20.5	69.6
2	Malignant neoplasms (C00-C97)	1,414	15.8	53.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	945	10.6	36.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	807	9.0	30.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	492	5.5	18.7
6	Cerebrovascular diseases (I60-I69)	361	4.0	13.7
7	Diabetes mellitus (E10-E14)	347	3.9	13.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	201	2.3	7.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	187	2.1	7.1
10	Septicemia (A40-A41)	170	1.9	6.5
11	Influenza and pneumonia (J09-J18)	151	1.7	5.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	134	1.5	5.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	128	1.4	4.9
14	Chronic lower respiratory diseases (J40-J47)	96	1.1	3.7
15	Anemias (D50-D64)	75	0.8	2.9
...	All other causes (Residual)	1,588	17.8	60.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	14,258	100.0	516.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,202	22.5	116.0
2	Malignant neoplasms (C00-C97)	3,065	21.5	111.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,189	8.3	43.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	938	6.6	34.0
5	Cerebrovascular diseases (I60-I69)	724	5.1	26.2
6	Diabetes mellitus (E10-E14)	479	3.4	17.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	418	2.9	15.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	329	2.3	11.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	302	2.1	10.9
10	Chronic lower respiratory diseases (J40-J47)	247	1.7	8.9
11	Septicemia (A40-A41)	236	1.7	8.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	215	1.5	7.8
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	1.5	7.7
14	Influenza and pneumonia (J09-J18)	204	1.4	7.4
15	Viral hepatitis (B15-B19)	91	0.6	3.3
...	All other causes (Residual)	2,407	16.9	87.2
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	20,804	100.0	826.5
1	Malignant neoplasms (C00-C97)	5,463	26.3	217.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,886	23.5	194.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,209	5.8	48.0
4	Cerebrovascular diseases (I60-I69)	1,056	5.1	42.0
5	Human immunodeficiency virus (HIV) disease (B20-B24)	907	4.4	36.0
6	Diabetes mellitus (E10-E14)	849	4.1	33.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	527	2.5	20.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	497	2.4	19.7
9	Chronic lower respiratory diseases (J40-J47)	455	2.2	18.1
10	Septicemia (A40-A41)	394	1.9	15.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	339	1.6	13.5
12	Influenza and pneumonia (J09-J18)	315	1.5	12.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	296	1.4	11.8
14	Viral hepatitis (B15-B19)	243	1.2	9.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	154	0.7	6.1
...	All other causes (Residual)	3,214	15.4	127.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	24,828	100.0	1,209.5
1	Malignant neoplasms (C00-C97)	7,412	29.9	361.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,131	24.7	298.7
3	Cerebrovascular diseases (I60-I69)	1,255	5.1	61.1
4	Diabetes mellitus (E10-E14)	1,179	4.7	57.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	941	3.8	45.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	675	2.7	32.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	663	2.7	32.3
8	Chronic lower respiratory diseases (J40-J47)	607	2.4	29.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	558	2.2	27.2
10	Septicemia (A40-A41)	536	2.2	26.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	383	1.5	18.7
12	Viral hepatitis (B15-B19)	331	1.3	16.1
13	Influenza and pneumonia (J09-J18)	318	1.3	15.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	174	0.7	8.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	109	0.4	5.3
...	All other causes (Residual)	3,556	14.3	173.2
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	25,261	100.0	1,669.0
1	Malignant neoplasms (C00-C97)	8,132	32.2	537.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,645	26.3	439.0
3	Cerebrovascular diseases (I60-I69)	1,324	5.2	87.5
4	Diabetes mellitus (E10-E14)	1,249	4.9	82.5
5	Chronic lower respiratory diseases (J40-J47)	756	3.0	49.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	740	2.9	48.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	667	2.6	44.1
8	Septicemia (A40-A41)	540	2.1	35.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	441	1.7	29.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	370	1.5	24.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	370	1.5	24.4
12	Influenza and pneumonia (J09-J18)	322	1.3	21.3
13	Viral hepatitis (B15-B19)	183	0.7	12.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	116	0.5	7.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	95	0.4	6.3
...	All other causes (Residual)	3,311	13.1	218.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	25,187	100.0	2,347.1
1	Malignant neoplasms (C00-C97)	8,136	32.3	758.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,601	26.2	615.1
3	Cerebrovascular diseases (I60-I69)	1,384	5.5	129.0
4	Diabetes mellitus (E10-E14)	1,322	5.2	123.2
5	Chronic lower respiratory diseases (J40-J47)	958	3.8	89.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	911	3.6	84.9
7	Septicemia (A40-A41)	605	2.4	56.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	439	1.7	40.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	421	1.7	39.2
10	Influenza and pneumonia (J09-J18)	366	1.5	34.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	229	0.9	21.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	190	0.8	17.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	118	0.5	11.0
14	Alzheimer's disease (G30)	95	0.4	8.9
14	Aortic aneurysm and dissection (I71)	95	0.4	8.9
...	All other causes (Residual)	3,317	13.2	309.1
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	26,148	100.0	3,154.6
1	Malignant neoplasms (C00-C97)	8,094	31.0	976.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,791	26.0	819.3
3	Cerebrovascular diseases (I60-I69)	1,595	6.1	192.4
4	Diabetes mellitus (E10-E14)	1,426	5.5	172.0
5	Chronic lower respiratory diseases (J40-J47)	1,099	4.2	132.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,008	3.9	121.6
7	Septicemia (A40-A41)	656	2.5	79.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	444	1.7	53.6
9	Influenza and pneumonia (J09-J18)	442	1.7	53.3
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	1.4	44.8
11	Alzheimer's disease (G30)	266	1.0	32.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	138	0.5	16.6
13	Pneumonitis due to solids and liquids (J69)	132	0.5	15.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	102	0.4	12.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	84	0.3	10.1
15	Parkinson's disease (G20-G21)	84	0.3	10.1
...	All other causes (Residual)	3,416	13.1	412.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	27,777	100.0	4,377.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,407	26.7	1,167.3
2	Malignant neoplasms (C00-C97)	7,399	26.6	1,166.0
3	Cerebrovascular diseases (I60-I69)	1,889	6.8	297.7
4	Diabetes mellitus (E10-E14)	1,438	5.2	226.6
5	Chronic lower respiratory diseases (J40-J47)	1,143	4.1	180.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,131	4.1	178.2
7	Septicemia (A40-A41)	712	2.6	112.2
8	Alzheimer's disease (G30)	625	2.3	98.5
9	Influenza and pneumonia (J09-J18)	575	2.1	90.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	478	1.7	75.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	375	1.4	59.1
12	Pneumonitis due to solids and liquids (J69)	183	0.7	28.8
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	140	0.5	22.1
14	Parkinson's disease (G20-G21)	134	0.5	21.1
15	Aortic aneurysm and dissection (I71)	92	0.3	14.5
...	All other causes (Residual)	4,056	14.6	639.2
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	29,008	100.0	6,521.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,931	27.3	1,783.0
2	Malignant neoplasms (C00-C97)	6,440	22.2	1,447.8
3	Cerebrovascular diseases (I60-I69)	2,076	7.2	466.7
4	Diabetes mellitus (E10-E14)	1,307	4.5	293.8
5	Chronic lower respiratory diseases (J40-J47)	1,174	4.0	263.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,159	4.0	260.6
7	Alzheimer's disease (G30)	1,031	3.6	231.8
8	Septicemia (A40-A41)	737	2.5	165.7
9	Influenza and pneumonia (J09-J18)	659	2.3	148.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	593	2.0	133.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	338	1.2	76.0
12	Pneumonitis due to solids and liquids (J69)	233	0.8	52.4
13	Parkinson's disease (G20-G21)	185	0.6	41.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	151	0.5	33.9
15	Aortic aneurysm and dissection (I71)	106	0.4	23.8
...	All other causes (Residual)	4,888	16.9	1,098.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	24,840	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,216	29.0	---
2	Malignant neoplasms (C00-C97)	4,147	16.7	---
3	Cerebrovascular diseases (I60-I69)	1,859	7.5	---
4	Alzheimer's disease (G30)	1,309	5.3	---
5	Diabetes mellitus (E10-E14)	1,061	4.3	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	947	3.8	---
7	Chronic lower respiratory diseases (J40-J47)	898	3.6	---
8	Septicemia (A40-A41)	670	2.7	---
9	Influenza and pneumonia (J09-J18)	657	2.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	566	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	300	1.2	---
12	Pneumonitis due to solids and liquids (J69)	230	0.9	---
13	Parkinson's disease (G20-G21)	146	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	119	0.5	---
15	Aortic aneurysm and dissection (I71)	95	0.4	---
...	All other causes (Residual)	4,620	18.6	---
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	14,862	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,598	30.9	---
2	Malignant neoplasms (C00-C97)	1,773	11.9	---
3	Cerebrovascular diseases (I60-I69)	1,138	7.7	---
4	Alzheimer's disease (G30)	958	6.4	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	520	3.5	---
6	Diabetes mellitus (E10-E14)	516	3.5	---
7	Chronic lower respiratory diseases (J40-J47)	433	2.9	---
8	Influenza and pneumonia (J09-J18)	408	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	393	2.6	---
10	Septicemia (A40-A41)	325	2.2	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	202	1.4	---
12	Pneumonitis due to solids and liquids (J69)	153	1.0	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	87	0.6	---
14	Parkinson's disease (G20-G21)	73	0.5	---
15	Atherosclerosis (I70)	68	0.5	---
...	All other causes (Residual)	3,217	21.6	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	6,994	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,246	32.1	---
2	Malignant neoplasms (C00-C97)	564	8.1	---
3	Alzheimer's disease (G30)	511	7.3	---
4	Cerebrovascular diseases (I60-I69)	504	7.2	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	257	3.7	---
6	Influenza and pneumonia (J09-J18)	222	3.2	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	3.0	---
8	Diabetes mellitus (E10-E14)	191	2.7	---
9	Chronic lower respiratory diseases (J40-J47)	165	2.4	---
10	Septicemia (A40-A41)	142	2.0	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	1.4	---
12	Pneumonitis due to solids and liquids (J69)	73	1.0	---
13	Atherosclerosis (I70)	38	0.5	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.5	---
15	Anemias (D50-D64)	34	0.5	---
...	All other causes (Residual)	1,707	24.4	---
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,136	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	773	36.2	---
2	Alzheimer's disease (G30)	167	7.8	---
3	Cerebrovascular diseases (I60-I69)	118	5.5	---
4	Malignant neoplasms (C00-C97)	100	4.7	---
5	Influenza and pneumonia (J09-J18)	91	4.3	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	3.1	---
7	Diabetes mellitus (E10-E14)	44	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	2.0	---
9	Septicemia (A40-A41)	40	1.9	---
10	Chronic lower respiratory diseases (J40-J47)	33	1.5	---
11	Atherosclerosis (I70)	19	0.9	---
11	Pneumonitis due to solids and liquids (J69)	19	0.9	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	0.8	---
14	Nutritional deficiencies (E40-E64)	8	0.4	---
15	Anemias (D50-D64)	6	0.3	---
...	All other causes (Residual)	593	27.8	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, all ages				
...	All causes	144,197	100.0	777.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,935	24.2	188.3
2	Malignant neoplasms (C00-C97)	33,065	22.9	178.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,894	5.5	42.5
4	Cerebrovascular diseases (I60-I69)	6,892	4.8	37.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,605	4.6	35.6
6	Diabetes mellitus (E10-E14)	5,430	3.8	29.3
7	Chronic lower respiratory diseases (J40-J47)	4,457	3.1	24.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,930	2.7	21.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,476	2.4	18.7
10	Septicemia (A40-A41)	2,721	1.9	14.7
11	Certain conditions originating in the perinatal period (P00-P96)	2,533	1.8	13.7
12	Influenza and pneumonia (J09-J18)	2,463	1.7	13.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,045	1.4	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,649	1.1	8.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,636	1.1	8.8
...	All other causes (Residual)	24,466	17.0	131.9
Non-Hispanic, black, male, 1-4 years				
...	All causes	561	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	30.1	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	14.1	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	8.6	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.9	1.7
5	Malignant neoplasms (C00-C97)	20	3.6	1.6
6	Chronic lower respiratory diseases (J40-J47)	18	3.2	*
7	Influenza and pneumonia (J09-J18)	13	2.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	10	1.8	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
11	Septicemia (A40-A41)	7	1.2	*
12	Anemias (D50-D64)	6	1.1	*
13	Cerebrovascular diseases (I60-I69)	4	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	141	25.1	11.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-9 years				
...	All causes	315	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	35.2	7.3
2	Malignant neoplasms (C00-C97)	36	11.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	6.3	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	6.0	*
5	Chronic lower respiratory diseases (J40-J47)	17	5.4	*
6	Influenza and pneumonia (J09-J18)	15	4.8	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.5	*
9	Septicemia (A40-A41)	6	1.9	*
9	Cerebrovascular diseases (I60-I69)	6	1.9	*
11	Anemias (D50-D64)	2	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	2	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
15	Meningitis (G00,G03)	1	0.3	*
15	Pneumonitis due to solids and liquids (J69)	1	0.3	*
15	Peptic ulcer (K25-K28)	1	0.3	*
15	Hernia (K40-K46)	1	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
15	Legal intervention (Y35,Y89.0)	1	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (Residual)	49	15.6	3.2
Non-Hispanic, black, male, 10-14 years				
...	All causes	420	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	30.2	8.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	12.6	3.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	7.9	2.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	6.7	1.8
5	Chronic lower respiratory diseases (J40-J47)	26	6.2	1.7
6	Malignant neoplasms (C00-C97)	24	5.7	1.6
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	5.5	1.5
8	Influenza and pneumonia (J09-J18)	7	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.2	*
9	Anemias (D50-D64)	5	1.2	*
9	Cerebrovascular diseases (I60-I69)	5	1.2	*
12	Septicemia (A40-A41)	4	1.0	*
13	Diseases of appendix (K35-K38)	3	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
15	Diabetes mellitus (E10-E14)	2	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
...	All other causes (Residual)	70	16.7	4.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-19 years				
...	All causes	1,896	100.0	109.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	959	50.6	55.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	20.6	22.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	121	6.4	7.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	71	3.7	4.1
5	Malignant neoplasms (C00-C97)	69	3.6	4.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	1.7	1.9
7	Chronic lower respiratory diseases (J40-J47)	17	0.9	*
8	Anemias (D50-D64)	12	0.6	*
9	Influenza and pneumonia (J09-J18)	11	0.6	*
9	Legal intervention (Y35,Y89.0)	11	0.6	*
11	Cerebrovascular diseases (I60-I69)	9	0.5	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	8	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
14	Diabetes mellitus (E10-E14)	5	0.3	*
15	Septicemia (A40-A41)	4	0.2	*
...	All other causes (Residual)	169	8.9	9.7
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,089	100.0	190.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,524	49.3	94.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	546	17.7	33.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	239	7.7	14.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	153	5.0	9.4
5	Malignant neoplasms (C00-C97)	89	2.9	5.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	57	1.8	3.5
7	Chronic lower respiratory diseases (J40-J47)	36	1.2	2.2
8	Influenza and pneumonia (J09-J18)	29	0.9	1.8
9	Anemias (D50-D64)	24	0.8	1.5
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.7	1.4
11	Diabetes mellitus (E10-E14)	20	0.6	1.2
12	Legal intervention (Y35,Y89.0)	18	0.6	*
13	Cerebrovascular diseases (I60-I69)	12	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.4	*
15	Septicemia (A40-A41)	8	0.3	*
...	All other causes (Residual)	299	9.7	18.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-29 years				
...	All causes	3,077	100.0	204.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,177	38.3	78.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	666	21.6	44.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	201	6.5	13.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	198	6.4	13.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	142	4.6	9.4
6	Malignant neoplasms (C00-C97)	124	4.0	8.2
7	Diabetes mellitus (E10-E14)	46	1.5	3.1
8	Anemias (D50-D64)	32	1.0	2.1
9	Cerebrovascular diseases (I60-I69)	29	0.9	1.9
10	Chronic lower respiratory diseases (J40-J47)	27	0.9	1.8
11	Influenza and pneumonia (J09-J18)	26	0.8	1.7
12	Septicemia (A40-A41)	21	0.7	1.4
13	Legal intervention (Y35,Y89.0)	20	0.6	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	0.6	*
...	All other causes (Residual)	331	10.8	22.0
Non-Hispanic, black, male, 30-34 years				
...	All causes	3,035	100.0	241.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	837	27.6	66.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	17.2	41.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	375	12.4	29.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	200	6.6	15.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	175	5.8	13.9
6	Malignant neoplasms (C00-C97)	156	5.1	12.4
7	Diabetes mellitus (E10-E14)	63	2.1	5.0
8	Cerebrovascular diseases (I60-I69)	50	1.6	4.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	1.3	3.0
10	Influenza and pneumonia (J09-J18)	36	1.2	2.9
11	Septicemia (A40-A41)	29	1.0	2.3
12	Anemias (D50-D64)	28	0.9	2.2
13	Chronic lower respiratory diseases (J40-J47)	25	0.8	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	0.7	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	0.7	1.6
...	All other causes (Residual)	461	15.2	36.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-39 years				
...	All causes	3,665	100.0	295.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	680	18.6	54.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	582	15.9	46.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	572	15.6	46.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	342	9.3	27.5
5	Malignant neoplasms (C00-C97)	279	7.6	22.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	164	4.5	13.2
7	Cerebrovascular diseases (I60-I69)	109	3.0	8.8
8	Diabetes mellitus (E10-E14)	100	2.7	8.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	54	1.5	4.3
10	Septicemia (A40-A41)	50	1.4	4.0
11	Chronic lower respiratory diseases (J40-J47)	48	1.3	3.9
12	Influenza and pneumonia (J09-J18)	45	1.2	3.6
13	Chronic liver disease and cirrhosis (K70,K73-K74)	42	1.1	3.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	37	1.0	3.0
15	Anemias (D50-D64)	29	0.8	2.3
...	All other causes (Residual)	532	14.5	42.8
Non-Hispanic, black, male, 40-44 years				
...	All causes	5,161	100.0	418.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,159	22.5	94.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	666	12.9	54.0
3	Malignant neoplasms (C00-C97)	541	10.5	43.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	502	9.7	40.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	412	8.0	33.4
6	Diabetes mellitus (E10-E14)	227	4.4	18.4
7	Cerebrovascular diseases (I60-I69)	179	3.5	14.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	165	3.2	13.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	2.0	8.5
10	Septicemia (A40-A41)	80	1.6	6.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	1.5	6.2
12	Influenza and pneumonia (J09-J18)	72	1.4	5.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	69	1.3	5.6
14	Chronic lower respiratory diseases (J40-J47)	58	1.1	4.7
15	Anemias (D50-D64)	33	0.6	2.7
...	All other causes (Residual)	817	15.8	66.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-49 years				
...	All causes	7,994	100.0	624.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,999	25.0	156.2
2	Malignant neoplasms (C00-C97)	1,358	17.0	106.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	780	9.8	60.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	595	7.4	46.5
5	Cerebrovascular diseases (I60-I69)	352	4.4	27.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	330	4.1	25.8
7	Diabetes mellitus (E10-E14)	278	3.5	21.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	195	2.4	15.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	2.1	13.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	163	2.0	12.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	1.6	9.9
12	Septicemia (A40-A41)	123	1.5	9.6
13	Influenza and pneumonia (J09-J18)	112	1.4	8.7
14	Chronic lower respiratory diseases (J40-J47)	106	1.3	8.3
15	Viral hepatitis (B15-B19)	59	0.7	4.6
...	All other causes (Residual)	1,251	15.6	97.7
Non-Hispanic, black, male, 50-54 years				
...	All causes	12,072	100.0	1,044.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,111	25.8	269.2
2	Malignant neoplasms (C00-C97)	2,798	23.2	242.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	830	6.9	71.8
4	Human immunodeficiency virus (HIV) disease (B20-B24)	618	5.1	53.5
5	Cerebrovascular diseases (I60-I69)	564	4.7	48.8
6	Diabetes mellitus (E10-E14)	494	4.1	42.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	334	2.8	28.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	292	2.4	25.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	230	1.9	19.9
10	Chronic lower respiratory diseases (J40-J47)	227	1.9	19.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	199	1.6	17.2
12	Septicemia (A40-A41)	197	1.6	17.0
13	Influenza and pneumonia (J09-J18)	163	1.4	14.1
14	Viral hepatitis (B15-B19)	153	1.3	13.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	112	0.9	9.7
...	All other causes (Residual)	1,750	14.5	151.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-59 years				
...	All causes	14,842	100.0	1,608.7
1	Malignant neoplasms (C00-C97)	4,124	27.8	447.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,901	26.3	422.8
3	Cerebrovascular diseases (I60-I69)	740	5.0	80.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	660	4.4	71.5
5	Diabetes mellitus (E10-E14)	638	4.3	69.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	488	3.3	52.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	393	2.6	42.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	374	2.5	40.5
9	Chronic lower respiratory diseases (J40-J47)	326	2.2	35.3
10	Septicemia (A40-A41)	285	1.9	30.9
11	Viral hepatitis (B15-B19)	236	1.6	25.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	223	1.5	24.2
13	Influenza and pneumonia (J09-J18)	186	1.3	20.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	148	1.0	16.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	0.6	9.1
...	All other causes (Residual)	2,036	13.7	220.7
Non-Hispanic, black, male, 60-64 years				
...	All causes	14,685	100.0	2,207.2
1	Malignant neoplasms (C00-C97)	4,575	31.2	687.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,104	27.9	616.9
3	Cerebrovascular diseases (I60-I69)	722	4.9	108.5
4	Diabetes mellitus (E10-E14)	655	4.5	98.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	476	3.2	71.5
6	Chronic lower respiratory diseases (J40-J47)	423	2.9	63.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	383	2.6	57.6
8	Septicemia (A40-A41)	280	1.9	42.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	268	1.8	40.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	245	1.7	36.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	240	1.6	36.1
12	Influenza and pneumonia (J09-J18)	167	1.1	25.1
13	Viral hepatitis (B15-B19)	136	0.9	20.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	0.5	11.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	64	0.4	9.6
...	All other causes (Residual)	1,872	12.7	281.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65-69 years				
...	All causes	14,083	100.0	3,103.0
1	Malignant neoplasms (C00-C97)	4,474	31.8	985.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,930	27.9	865.9
3	Cerebrovascular diseases (I60-I69)	782	5.6	172.3
4	Diabetes mellitus (E10-E14)	686	4.9	151.2
5	Chronic lower respiratory diseases (J40-J47)	530	3.8	116.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	445	3.2	98.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	308	2.2	67.9
8	Septicemia (A40-A41)	293	2.1	64.6
9	Influenza and pneumonia (J09-J18)	225	1.6	49.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	207	1.5	45.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	152	1.1	33.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	135	1.0	29.7
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.5	14.5
14	Viral hepatitis (B15-B19)	60	0.4	13.2
15	Aortic aneurysm and dissection (I71)	57	0.4	12.6
...	All other causes (Residual)	1,733	12.3	381.8
Non-Hispanic, black, male, 70-74 years				
...	All causes	13,685	100.0	4,027.5
1	Malignant neoplasms (C00-C97)	4,403	32.2	1,295.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,698	27.0	1,088.3
3	Cerebrovascular diseases (I60-I69)	783	5.7	230.4
4	Diabetes mellitus (E10-E14)	645	4.7	189.8
5	Chronic lower respiratory diseases (J40-J47)	606	4.4	178.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	479	3.5	141.0
7	Septicemia (A40-A41)	304	2.2	89.5
8	Influenza and pneumonia (J09-J18)	240	1.8	70.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	229	1.7	67.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	197	1.4	58.0
11	Alzheimer's disease (G30)	115	0.8	33.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	82	0.6	24.1
13	Pneumonitis due to solids and liquids (J69)	66	0.5	19.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	62	0.5	18.2
15	Parkinson's disease (G20-G21)	59	0.4	17.4
...	All other causes (Residual)	1,717	12.5	505.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-79 years				
...	All causes	13,471	100.0	5,522.6
1	Malignant neoplasms (C00-C97)	3,888	28.9	1,593.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,598	26.7	1,475.0
3	Cerebrovascular diseases (I60-I69)	822	6.1	337.0
4	Chronic lower respiratory diseases (J40-J47)	647	4.8	265.2
5	Diabetes mellitus (E10-E14)	593	4.4	243.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	510	3.8	209.1
7	Septicemia (A40-A41)	340	2.5	139.4
8	Influenza and pneumonia (J09-J18)	300	2.2	123.0
9	Alzheimer's disease (G30)	253	1.9	103.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	1.6	91.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	192	1.4	78.7
12	Pneumonitis due to solids and liquids (J69)	99	0.7	40.6
13	Parkinson's disease (G20-G21)	79	0.6	32.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.4	24.6
15	Aortic aneurysm and dissection (I71)	48	0.4	19.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	48	0.4	19.7
...	All other causes (Residual)	1,772	13.2	726.5
Non-Hispanic, black, male, 80-84 years				
...	All causes	12,319	100.0	8,040.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,353	27.2	2,188.5
2	Malignant neoplasms (C00-C97)	3,162	25.7	2,063.8
3	Cerebrovascular diseases (I60-I69)	764	6.2	498.7
4	Chronic lower respiratory diseases (J40-J47)	626	5.1	408.6
5	Diabetes mellitus (E10-E14)	489	4.0	319.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	469	3.8	306.1
7	Alzheimer's disease (G30)	336	2.7	219.3
8	Influenza and pneumonia (J09-J18)	291	2.4	189.9
9	Septicemia (A40-A41)	259	2.1	169.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	208	1.7	135.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	1.4	116.2
12	Pneumonitis due to solids and liquids (J69)	109	0.9	71.1
13	Parkinson's disease (G20-G21)	104	0.8	67.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.6	47.0
15	Aortic aneurysm and dissection (I71)	40	0.3	26.1
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	40	0.3	26.1
...	All other causes (Residual)	1,819	14.8	1,187.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 85-89 years				
...	All causes	9,088	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,620	28.8	---
2	Malignant neoplasms (C00-C97)	1,936	21.3	---
3	Cerebrovascular diseases (I60-I69)	563	6.2	---
4	Chronic lower respiratory diseases (J40-J47)	432	4.8	---
5	Alzheimer's disease (G30)	359	4.0	---
6	Diabetes mellitus (E10-E14)	327	3.6	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	313	3.4	---
8	Influenza and pneumonia (J09-J18)	269	3.0	---
9	Septicemia (A40-A41)	235	2.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	149	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	1.4	---
12	Pneumonitis due to solids and liquids (J69)	103	1.1	---
13	Parkinson's disease (G20-G21)	80	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	0.5	---
15	Aortic aneurysm and dissection (I71)	28	0.3	---
...	All other causes (Residual)	1,507	16.6	---
Non-Hispanic, black, male, 90-94 years				
...	All causes	4,382	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,277	29.1	---
2	Malignant neoplasms (C00-C97)	780	17.8	---
3	Cerebrovascular diseases (I60-I69)	273	6.2	---
4	Chronic lower respiratory diseases (J40-J47)	189	4.3	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	179	4.1	---
6	Alzheimer's disease (G30)	172	3.9	---
7	Influenza and pneumonia (J09-J18)	136	3.1	---
8	Diabetes mellitus (E10-E14)	125	2.9	---
9	Septicemia (A40-A41)	104	2.4	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	100	2.3	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	1.5	---
12	Pneumonitis due to solids and liquids (J69)	58	1.3	---
13	Parkinson's disease (G20-G21)	39	0.9	---
14	Atherosclerosis (I70)	22	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.4	---
...	All other causes (Residual)	845	19.3	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, black, male, 95-99 years				
...	All causes	1,493	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	450	30.1	---
2	Malignant neoplasms (C00-C97)	195	13.1	---
3	Alzheimer's disease (G30)	86	5.8	---
3	Cerebrovascular diseases (I60-I69)	86	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	61	4.1	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	4.1	---
7	Influenza and pneumonia (J09-J18)	60	4.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	46	3.1	---
9	Septicemia (A40-A41)	36	2.4	---
10	Diabetes mellitus (E10-E14)	32	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	1.5	---
12	Pneumonitis due to solids and liquids (J69)	19	1.3	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.7	---
14	Anemias (D50-D64)	7	0.5	---
14	Parkinson's disease (G20-G21)	7	0.5	---
...	All other causes (Residual)	315	21.1	---
Non-Hispanic, black, male, 100 years and over				
...	All causes	326	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	38.7	---
2	Malignant neoplasms (C00-C97)	25	7.7	---
3	Alzheimer's disease (G30)	16	4.9	---
4	Influenza and pneumonia (J09-J18)	14	4.3	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	3.7	---
6	Cerebrovascular diseases (I60-I69)	10	3.1	---
7	Septicemia (A40-A41)	9	2.8	---
8	Chronic lower respiratory diseases (J40-J47)	7	2.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	1.8	---
11	Pneumonitis due to solids and liquids (J69)	5	1.5	---
12	Diabetes mellitus (E10-E14)	4	1.2	---
13	Atherosclerosis (I70)	3	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	---
14	Anemias (D50-D64)	1	0.3	---
14	Peptic ulcer (K25-K28)	1	0.3	---
14	Hyperplasia of prostate (N40)	1	0.3	---
...	All other causes (Residual)	78	23.9	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	138,785	100.0	683.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,876	24.4	166.8
2	Malignant neoplasms (C00-C97)	30,902	22.3	152.2
3	Cerebrovascular diseases (I60-I69)	8,826	6.4	43.5
4	Diabetes mellitus (E10-E14)	6,403	4.6	31.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,797	3.5	23.6
6	Chronic lower respiratory diseases (J40-J47)	3,987	2.9	19.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,916	2.8	19.3
8	Alzheimer's disease (G30)	3,612	2.6	17.8
9	Septicemia (A40-A41)	3,410	2.5	16.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,774	2.0	13.7
11	Influenza and pneumonia (J09-J18)	2,751	2.0	13.5
12	Certain conditions originating in the perinatal period (P00-P96)	1,923	1.4	9.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,831	1.3	9.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,128	0.8	5.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	882	0.6	4.3
...	All other causes (Residual)	27,767	20.0	136.7
Non-Hispanic, black, female, 1-4 years				
...	All causes	478	100.0	38.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	24.9	9.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	13.4	5.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	8.2	3.2
4	Malignant neoplasms (C00-C97)	36	7.5	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.9	2.3
6	Influenza and pneumonia (J09-J18)	16	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	15	3.1	*
8	Septicemia (A40-A41)	12	2.5	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.9	*
10	Cerebrovascular diseases (I60-I69)	5	1.0	*
10	Acute bronchitis and bronchiolitis (J20-J21)	5	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Anemias (D50-D64)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	119	24.9	9.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-9 years				
...	All causes	248	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	29.8	5.0
2	Malignant neoplasms (C00-C97)	40	16.1	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	7.3	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	7.3	*
5	Chronic lower respiratory diseases (J40-J47)	12	4.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.4	*
7	Influenza and pneumonia (J09-J18)	8	3.2	*
8	Septicemia (A40-A41)	4	1.6	*
8	Cerebrovascular diseases (I60-I69)	4	1.6	*
10	Anemias (D50-D64)	3	1.2	*
10	Diabetes mellitus (E10-E14)	3	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
14	Meningitis (G00,G03)	1	0.4	*
14	Aortic aneurysm and dissection (I71)	1	0.4	*
14	Peptic ulcer (K25-K28)	1	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (Residual)	44	17.7	3.0
Non-Hispanic, black, female, 10-14 years				
...	All causes	283	100.0	19.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	65	23.0	4.4
2	Malignant neoplasms (C00-C97)	28	9.9	1.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	6.4	*
4	Influenza and pneumonia (J09-J18)	15	5.3	*
4	Chronic lower respiratory diseases (J40-J47)	15	5.3	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	4.9	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	4.9	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	4.2	*
9	Anemias (D50-D64)	9	3.2	*
10	Septicemia (A40-A41)	8	2.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	2.1	*
12	Diabetes mellitus (E10-E14)	3	1.1	*
12	Cerebrovascular diseases (I60-I69)	3	1.1	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.1	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.7	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	2	0.7	*
...	All other causes (Residual)	66	23.3	4.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age agroups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-19 years				
...	All causes	610	100.0	36.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	28.2	10.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	114	18.7	6.8
3	Malignant neoplasms (C00-C97)	46	7.5	2.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	5.7	2.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	5.1	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	21	3.4	1.2
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	3.4	1.2
8	Influenza and pneumonia (J09-J18)	17	2.8	*
8	Chronic lower respiratory diseases (J40-J47)	17	2.8	*
10	Anemias (D50-D64)	11	1.8	*
11	Septicemia (A40-A41)	10	1.6	*
12	Diabetes mellitus (E10-E14)	8	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	7	1.1	*
14	Cerebrovascular diseases (I60-I69)	5	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
...	All other causes (Residual)	91	14.9	5.4
Non-Hispanic, black, female, 20-24 years				
...	All causes	1,007	100.0	63.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	19.9	12.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	16.2	10.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	81	8.0	5.1
4	Malignant neoplasms (C00-C97)	80	7.9	5.1
5	Pregnancy, childbirth and the puerperium (O00-O99)	67	6.7	4.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	4.2	2.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	34	3.4	2.2
8	Influenza and pneumonia (J09-J18)	28	2.8	1.8
9	Anemias (D50-D64)	19	1.9	*
10	Diabetes mellitus (E10-E14)	17	1.7	*
11	Chronic lower respiratory diseases (J40-J47)	15	1.5	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.5	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.4	*
14	Cerebrovascular diseases (I60-I69)	13	1.3	*
15	Septicemia (A40-A41)	9	0.9	*
...	All other causes (Residual)	210	20.9	13.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,308	100.0	86.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	201	15.4	13.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	141	10.8	9.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	138	10.6	9.1
4	Malignant neoplasms (C00-C97)	129	9.9	8.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	88	6.7	5.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	66	5.0	4.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	4.0	3.4
8	Influenza and pneumonia (J09-J18)	45	3.4	3.0
9	Diabetes mellitus (E10-E14)	36	2.8	2.4
10	Septicemia (A40-A41)	30	2.3	2.0
11	Anemias (D50-D64)	22	1.7	1.4
11	Cerebrovascular diseases (I60-I69)	22	1.7	1.4
11	Chronic lower respiratory diseases (J40-J47)	22	1.7	1.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	1.7	1.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.5	*
...	All other causes (Residual)	275	21.0	18.1
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,611	100.0	118.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	235	14.6	17.2
2	Malignant neoplasms (C00-C97)	217	13.5	15.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	11.0	13.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	166	10.3	12.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	8.6	10.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	63	3.9	4.6
7	Diabetes mellitus (E10-E14)	44	2.7	3.2
8	Cerebrovascular diseases (I60-I69)	42	2.6	3.1
9	Influenza and pneumonia (J09-J18)	36	2.2	2.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	2.2	2.6
11	Anemias (D50-D64)	29	1.8	2.1
12	Septicemia (A40-A41)	27	1.7	2.0
13	Chronic lower respiratory diseases (J40-J47)	26	1.6	1.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	1.5	1.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	1.4	1.6
...	All other causes (Residual)	328	20.4	24.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,360	100.0	170.0
1	Malignant neoplasms (C00-C97)	440	18.6	31.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	353	15.0	25.4
3	Human immunodeficiency virus (HIV) disease (B20-B24)	219	9.3	15.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	219	9.3	15.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	101	4.3	7.3
6	Diabetes mellitus (E10-E14)	77	3.3	5.5
7	Cerebrovascular diseases (I60-I69)	73	3.1	5.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	3.0	5.0
9	Influenza and pneumonia (J09-J18)	56	2.4	4.0
10	Septicemia (A40-A41)	51	2.2	3.7
11	Pregnancy, childbirth and the puerperium (O00-O99)	39	1.7	2.8
12	Chronic lower respiratory diseases (J40-J47)	35	1.5	2.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	1.4	2.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	1.3	2.2
15	Anemias (D50-D64)	27	1.1	1.9
...	All other causes (Residual)	537	22.8	38.7
Non-Hispanic, black, female, 40-44 years				
...	All causes	3,765	100.0	269.7
1	Malignant neoplasms (C00-C97)	873	23.2	62.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	671	17.8	48.1
3	Human immunodeficiency virus (HIV) disease (B20-B24)	305	8.1	21.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	7.4	20.0
5	Cerebrovascular diseases (I60-I69)	182	4.8	13.0
6	Diabetes mellitus (E10-E14)	120	3.2	8.6
7	Septicemia (A40-A41)	90	2.4	6.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	2.2	5.9
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	80	2.1	5.7
10	Influenza and pneumonia (J09-J18)	79	2.1	5.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	59	1.6	4.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	58	1.5	4.2
13	Anemias (D50-D64)	42	1.1	3.0
14	Chronic lower respiratory diseases (J40-J47)	38	1.0	2.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	1.0	2.6
...	All other causes (Residual)	771	20.5	55.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-49 years				
...	All causes	6,264	100.0	423.0
1	Malignant neoplasms (C00-C97)	1,707	27.3	115.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,203	19.2	81.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	6.5	27.6
4	Cerebrovascular diseases (I60-I69)	372	5.9	25.1
5	Human immunodeficiency virus (HIV) disease (B20-B24)	343	5.5	23.2
6	Diabetes mellitus (E10-E14)	201	3.2	13.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	2.6	11.0
8	Chronic lower respiratory diseases (J40-J47)	141	2.3	9.5
9	Septicemia (A40-A41)	113	1.8	7.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	107	1.7	7.2
11	Influenza and pneumonia (J09-J18)	92	1.5	6.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	88	1.4	5.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	88	1.4	5.9
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	0.8	3.3
15	Anemias (D50-D64)	41	0.7	2.8
...	All other causes (Residual)	1,147	18.3	77.5
Non-Hispanic, black, female, 50-54 years				
...	All causes	8,732	100.0	641.3
1	Malignant neoplasms (C00-C97)	2,665	30.5	195.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,775	20.3	130.4
3	Cerebrovascular diseases (I60-I69)	492	5.6	36.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	379	4.3	27.8
5	Diabetes mellitus (E10-E14)	355	4.1	26.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	289	3.3	21.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	235	2.7	17.3
8	Chronic lower respiratory diseases (J40-J47)	228	2.6	16.7
9	Septicemia (A40-A41)	197	2.3	14.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	163	1.9	12.0
11	Influenza and pneumonia (J09-J18)	152	1.7	11.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	140	1.6	10.3
13	Viral hepatitis (B15-B19)	90	1.0	6.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	66	0.8	4.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	0.5	3.1
...	All other causes (Residual)	1,464	16.8	107.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, black, female, 55-59 years				
...	All causes	9,986	100.0	883.6
1	Malignant neoplasms (C00-C97)	3,288	32.9	290.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,230	22.3	197.3
3	Diabetes mellitus (E10-E14)	541	5.4	47.9
4	Cerebrovascular diseases (I60-I69)	515	5.2	45.6
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	301	3.0	26.6
6	Chronic lower respiratory diseases (J40-J47)	281	2.8	24.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	281	2.8	24.9
8	Septicemia (A40-A41)	251	2.5	22.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	175	1.8	15.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	165	1.7	14.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	160	1.6	14.2
12	Influenza and pneumonia (J09-J18)	132	1.3	11.7
13	Viral hepatitis (B15-B19)	95	1.0	8.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	4.6
15	Pneumonitis due to solids and liquids (J69)	38	0.4	3.4
...	All other causes (Residual)	1,481	14.8	131.0
Non-Hispanic, black, female, 60-64 years				
...	All causes	10,576	100.0	1,246.8
1	Malignant neoplasms (C00-C97)	3,557	33.6	419.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,541	24.0	299.6
3	Cerebrovascular diseases (I60-I69)	602	5.7	71.0
4	Diabetes mellitus (E10-E14)	594	5.6	70.0
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	357	3.4	42.1
6	Chronic lower respiratory diseases (J40-J47)	333	3.1	39.3
7	Septicemia (A40-A41)	260	2.5	30.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	201	1.9	23.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	191	1.8	22.5
10	Influenza and pneumonia (J09-J18)	155	1.5	18.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	125	1.2	14.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	102	1.0	12.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	6.1
14	Viral hepatitis (B15-B19)	47	0.4	5.5
15	Pneumonitis due to solids and liquids (J69)	38	0.4	4.5
...	All other causes (Residual)	1,421	13.4	167.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of	
			total deaths	Rate
Non-Hispanic, black, female, 65-69 years				
...	All causes	11,104	100.0	1,793.0
1	Malignant neoplasms (C00-C97)	3,662	33.0	591.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,671	24.1	431.3
3	Diabetes mellitus (E10-E14)	636	5.7	102.7
4	Cerebrovascular diseases (I60-I69)	602	5.4	97.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	466	4.2	75.2
6	Chronic lower respiratory diseases (J40-J47)	428	3.9	69.1
7	Septicemia (A40-A41)	312	2.8	50.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	214	1.9	34.6
9	Influenza and pneumonia (J09-J18)	141	1.3	22.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	131	1.2	21.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	77	0.7	12.4
12	Human immunodeficiency virus (HIV) disease (B20-B24)	55	0.5	8.9
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	52	0.5	8.4
14	Alzheimer's disease (G30)	47	0.4	7.6
15	Pneumonitis due to solids and liquids (J69)	39	0.4	6.3
...	All other causes (Residual)	1,571	14.1	253.7
Non-Hispanic, black, female, 70-74 years				
...	All causes	12,463	100.0	2,548.2
1	Malignant neoplasms (C00-C97)	3,691	29.6	754.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,093	24.8	632.4
3	Cerebrovascular diseases (I60-I69)	812	6.5	166.0
4	Diabetes mellitus (E10-E14)	781	6.3	159.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	529	4.2	108.2
6	Chronic lower respiratory diseases (J40-J47)	493	4.0	100.8
7	Septicemia (A40-A41)	352	2.8	72.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	247	2.0	50.5
9	Influenza and pneumonia (J09-J18)	202	1.6	41.3
10	Alzheimer's disease (G30)	151	1.2	30.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	1.1	29.0
12	Pneumonitis due to solids and liquids (J69)	66	0.5	13.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	58	0.5	11.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	56	0.4	11.4
15	Aortic aneurysm and dissection (I71)	40	0.3	8.2
...	All other causes (Residual)	1,750	14.0	357.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 75-79 years				
...	All causes	14,306	100.0	3,662.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,809	26.6	975.1
2	Malignant neoplasms (C00-C97)	3,511	24.5	898.8
3	Cerebrovascular diseases (I60-I69)	1,067	7.5	273.2
4	Diabetes mellitus (E10-E14)	845	5.9	216.3
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	621	4.3	159.0
6	Chronic lower respiratory diseases (J40-J47)	496	3.5	127.0
7	Septicemia (A40-A41)	372	2.6	95.2
7	Alzheimer's disease (G30)	372	2.6	95.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	286	2.0	73.2
10	Influenza and pneumonia (J09-J18)	275	1.9	70.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	1.1	39.2
12	Pneumonitis due to solids and liquids (J69)	84	0.6	21.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.6	20.5
14	Parkinson's disease (G20-G21)	55	0.4	14.1
15	Aortic aneurysm and dissection (I71)	44	0.3	11.3
...	All other causes (Residual)	2,236	15.6	572.4
Non-Hispanic, black, female, 80-84 years				
...	All causes	16,689	100.0	5,723.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,578	27.4	1,569.9
2	Malignant neoplasms (C00-C97)	3,278	19.6	1,124.1
3	Cerebrovascular diseases (I60-I69)	1,312	7.9	449.9
4	Diabetes mellitus (E10-E14)	818	4.9	280.5
5	Alzheimer's disease (G30)	695	4.2	238.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	690	4.1	236.6
7	Chronic lower respiratory diseases (J40-J47)	548	3.3	187.9
8	Septicemia (A40-A41)	478	2.9	163.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	385	2.3	132.0
10	Influenza and pneumonia (J09-J18)	368	2.2	126.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	1.0	54.9
12	Pneumonitis due to solids and liquids (J69)	124	0.7	42.5
13	Parkinson's disease (G20-G21)	81	0.5	27.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	79	0.5	27.1
15	Atherosclerosis (I70)	70	0.4	24.0
...	All other causes (Residual)	3,025	18.1	1,037.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85-89 years				
...	All causes	15,752	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,596	29.2	---
2	Malignant neoplasms (C00-C97)	2,211	14.0	---
3	Cerebrovascular diseases (I60-I69)	1,296	8.2	---
4	Alzheimer's disease (G30)	950	6.0	---
5	Diabetes mellitus (E10-E14)	734	4.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	634	4.0	---
7	Chronic lower respiratory diseases (J40-J47)	466	3.0	---
8	Septicemia (A40-A41)	435	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	417	2.6	---
10	Influenza and pneumonia (J09-J18)	388	2.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	1.1	---
12	Pneumonitis due to solids and liquids (J69)	127	0.8	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.5	---
14	Aortic aneurysm and dissection (I71)	67	0.4	---
15	Parkinson's disease (G20-G21)	66	0.4	---
...	All other causes (Residual)	3,113	19.8	---
Non-Hispanic, black, female, 90-94 years				
...	All causes	10,480	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,321	31.7	---
2	Malignant neoplasms (C00-C97)	993	9.5	---
3	Cerebrovascular diseases (I60-I69)	865	8.3	---
4	Alzheimer's disease (G30)	786	7.5	---
5	Diabetes mellitus (E10-E14)	391	3.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	341	3.3	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	293	2.8	---
8	Influenza and pneumonia (J09-J18)	272	2.6	---
9	Chronic lower respiratory diseases (J40-J47)	244	2.3	---
10	Septicemia (A40-A41)	221	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	1.3	---
12	Pneumonitis due to solids and liquids (J69)	95	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.6	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	49	0.5	---
15	Atherosclerosis (I70)	46	0.4	---
...	All other causes (Residual)	2,357	22.5	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available

LCWK4. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 95-99 years				
...	All causes	5,501	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,796	32.6	---
2	Alzheimer's disease (G30)	425	7.7	---
3	Cerebrovascular diseases (I60-I69)	418	7.6	---
4	Malignant neoplasms (C00-C97)	369	6.7	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	196	3.6	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	162	2.9	---
6	Influenza and pneumonia (J09-J18)	162	2.9	---
8	Diabetes mellitus (E10-E14)	159	2.9	---
9	Septicemia (A40-A41)	106	1.9	---
10	Chronic lower respiratory diseases (J40-J47)	104	1.9	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	1.3	---
12	Pneumonitis due to solids and liquids (J69)	54	1.0	---
13	Atherosclerosis (I70)	32	0.6	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.5	---
14	Anemias (D50-D64)	27	0.5	---
...	All other causes (Residual)	1,391	25.3	---
Non-Hispanic, black, female, 100 years and over				
...	All causes	1,810	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	647	35.7	---
2	Alzheimer's disease (G30)	151	8.3	---
3	Cerebrovascular diseases (I60-I69)	108	6.0	---
4	Influenza and pneumonia (J09-J18)	77	4.3	---
5	Malignant neoplasms (C00-C97)	75	4.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	3.0	---
7	Diabetes mellitus (E10-E14)	40	2.2	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	1.9	---
9	Septicemia (A40-A41)	31	1.7	---
10	Chronic lower respiratory diseases (J40-J47)	26	1.4	---
11	Atherosclerosis (I70)	16	0.9	---
12	Pneumonitis due to solids and liquids (J69)	14	0.8	---
13	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	0.6	---
14	Nutritional deficiencies (E40-E64)	8	0.4	---
15	Anemias (D50-D64)	5	0.3	---
...	All other causes (Residual)	512	28.3	---

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

--- Data not available