

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	139,241	100.0	296.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,951	20.8	61.7
2	Malignant neoplasms (C00-C97)	28,851	20.7	61.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,080	8.0	23.6
4	Cerebrovascular diseases (I60-I69)	7,121	5.1	15.2
5	Diabetes mellitus (E10-E14)	6,544	4.7	13.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,091	2.9	8.7
7	Chronic lower respiratory diseases (J40-J47)	3,949	2.8	8.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,331	2.4	7.1
9	Influenza and pneumonia (J09-J18)	3,176	2.3	6.8
10	Alzheimer's disease (G30)	3,005	2.2	6.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,903	2.1	6.2
12	Certain conditions originating in the perinatal period (P00-P96)	2,873	2.1	6.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,345	1.7	5.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,011	1.4	4.3
15	Septicemia (A40-A41)	2,004	1.4	4.3
...	All other causes (Residual)	27,006	19.4	57.5
Hispanic, both sexes, 1-4 years				
...	All causes	1,029	100.0	24.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	329	32.0	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	130	12.6	3.1
3	Malignant neoplasms (C00-C97)	105	10.2	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	7.7	1.9
5	Influenza and pneumonia (J09-J18)	31	3.0	0.7
6	Septicemia (A40-A41)	28	2.7	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	2.6	0.6
8	Certain conditions originating in the perinatal period (P00-P96)	17	1.7	*
9	Cerebrovascular diseases (I60-I69)	15	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	5	0.5	*
12	Anemias (D50-D64)	4	0.4	*
13	Meningitis (G00,G03)	3	0.3	*
13	Pneumonitis due to solids and liquids (J69)	3	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	240	23.3	5.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,020	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	29.8	3.6
2	Malignant neoplasms (C00-C97)	186	18.2	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	6.7	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	6.2	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.7	0.4
6	Cerebrovascular diseases (I60-I69)	26	2.5	0.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	2.2	0.3
8	Influenza and pneumonia (J09-J18)	16	1.6	*
8	Chronic lower respiratory diseases (J40-J47)	16	1.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.4	*
11	Septicemia (A40-A41)	11	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
13	Anemias (D50-D64)	6	0.6	*
13	Diabetes mellitus (E10-E14)	6	0.6	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
...	All other causes (Residual)	232	22.7	2.7
Hispanic, both sexes, 15-24 years				
...	All causes	5,203	100.0	69.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,152	41.4	28.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,217	23.4	16.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	511	9.8	6.8
4	Malignant neoplasms (C00-C97)	350	6.7	4.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	2.4	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	1.2	0.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	39	0.7	0.5
8	Cerebrovascular diseases (I60-I69)	33	0.6	0.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	32	0.6	0.4
10	Influenza and pneumonia (J09-J18)	29	0.6	0.4
11	Diabetes mellitus (E10-E14)	21	0.4	0.3
11	Legal intervention (Y35,Y89.0)	21	0.4	0.3
13	Chronic lower respiratory diseases (J40-J47)	20	0.4	0.3
14	Septicemia (A40-A41)	19	0.4	*
15	Anemias (D50-D64)	17	0.3	*
...	All other causes (Residual)	551	10.6	7.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-34 years				
...	All causes	6,396	100.0	78.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,260	35.3	27.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	941	14.7	11.5
3	Malignant neoplasms (C00-C97)	643	10.1	7.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	551	8.6	6.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	362	5.7	4.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	175	2.7	2.1
7	Cerebrovascular diseases (I60-I69)	117	1.8	1.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	102	1.6	1.2
9	Pregnancy, childbirth and the puerperium (O00-O99)	74	1.2	0.9
10	Diabetes mellitus (E10-E14)	62	1.0	0.8
11	Influenza and pneumonia (J09-J18)	55	0.9	0.7
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	0.8	0.7
13	Septicemia (A40-A41)	37	0.6	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.5	0.4
15	Chronic lower respiratory diseases (J40-J47)	29	0.5	0.4
...	All other causes (Residual)	901	14.1	11.0
Hispanic, both sexes, 35-44 years				
...	All causes	8,621	100.0	123.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,833	21.3	26.2
2	Malignant neoplasms (C00-C97)	1,500	17.4	21.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	885	10.3	12.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	535	6.2	7.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	499	5.8	7.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	489	5.7	7.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	405	4.7	5.8
8	Cerebrovascular diseases (I60-I69)	301	3.5	4.3
9	Diabetes mellitus (E10-E14)	225	2.6	3.2
10	Viral hepatitis (B15-B19)	109	1.3	1.6
11	Influenza and pneumonia (J09-J18)	102	1.2	1.5
12	Septicemia (A40-A41)	93	1.1	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	79	0.9	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	60	0.7	0.9
15	Chronic lower respiratory diseases (J40-J47)	56	0.6	0.8
...	All other causes (Residual)	1,450	16.8	20.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-54 years				
...	All causes	14,370	100.0	288.4
1	Malignant neoplasms (C00-C97)	3,550	24.7	71.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,311	16.1	46.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,541	10.7	30.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,182	8.2	23.7
5	Cerebrovascular diseases (I60-I69)	638	4.4	12.8
6	Diabetes mellitus (E10-E14)	601	4.2	12.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	467	3.2	9.4
8	Viral hepatitis (B15-B19)	412	2.9	8.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	385	2.7	7.7
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	267	1.9	5.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	209	1.5	4.2
12	Influenza and pneumonia (J09-J18)	185	1.3	3.7
13	Septicemia (A40-A41)	179	1.2	3.6
14	Chronic lower respiratory diseases (J40-J47)	139	1.0	2.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	112	0.8	2.2
...	All other causes (Residual)	2,192	15.3	44.0
Hispanic, both sexes, 55-64 years				
...	All causes	18,502	100.0	648.3
1	Malignant neoplasms (C00-C97)	5,774	31.2	202.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,867	20.9	135.5
3	Diabetes mellitus (E10-E14)	1,194	6.5	41.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,026	5.5	35.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	863	4.7	30.2
6	Cerebrovascular diseases (I60-I69)	795	4.3	27.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	474	2.6	16.6
8	Viral hepatitis (B15-B19)	408	2.2	14.3
9	Chronic lower respiratory diseases (J40-J47)	352	1.9	12.3
10	Influenza and pneumonia (J09-J18)	290	1.6	10.2
11	Septicemia (A40-A41)	264	1.4	9.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	200	1.1	7.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	195	1.1	6.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	174	0.9	6.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	0.6	3.6
...	All other causes (Residual)	2,522	13.6	88.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65-74 years				
...	All causes	21,943	100.0	1,457.4
1	Malignant neoplasms (C00-C97)	6,686	30.5	444.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,103	23.3	338.9
3	Diabetes mellitus (E10-E14)	1,585	7.2	105.3
4	Cerebrovascular diseases (I60-I69)	1,159	5.3	77.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	739	3.4	49.1
6	Chronic lower respiratory diseases (J40-J47)	737	3.4	48.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	618	2.8	41.0
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	2.4	34.6
9	Influenza and pneumonia (J09-J18)	448	2.0	29.8
10	Septicemia (A40-A41)	392	1.8	26.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	251	1.1	16.7
12	Alzheimer's disease (G30)	233	1.1	15.5
13	Viral hepatitis (B15-B19)	144	0.7	9.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	133	0.6	8.8
15	Parkinson's disease (G20-G21)	132	0.6	8.8
...	All other causes (Residual)	3,062	14.0	203.4
Hispanic, both sexes, 75-84 years				
...	All causes	30,262	100.0	3,592.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,769	25.7	922.2
2	Malignant neoplasms (C00-C97)	6,953	23.0	825.3
3	Cerebrovascular diseases (I60-I69)	2,133	7.0	253.2
4	Diabetes mellitus (E10-E14)	1,814	6.0	215.3
5	Chronic lower respiratory diseases (J40-J47)	1,410	4.7	167.4
6	Alzheimer's disease (G30)	992	3.3	117.8
7	Influenza and pneumonia (J09-J18)	908	3.0	107.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	835	2.8	99.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	598	2.0	71.0
10	Septicemia (A40-A41)	523	1.7	62.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	432	1.4	51.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	412	1.4	48.9
13	Parkinson's disease (G20-G21)	394	1.3	46.8
14	Pneumonitis due to solids and liquids (J69)	194	0.6	23.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	178	0.6	21.1
...	All other causes (Residual)	4,717	15.6	559.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 85 years and over				
...	All causes	25,996	100.0	8,299.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,370	32.2	2,672.1
2	Malignant neoplasms (C00-C97)	3,086	11.9	985.2
3	Cerebrovascular diseases (I60-I69)	1,877	7.2	599.2
4	Alzheimer's disease (G30)	1,733	6.7	553.3
5	Chronic lower respiratory diseases (J40-J47)	1,177	4.5	375.8
6	Influenza and pneumonia (J09-J18)	1,066	4.1	340.3
7	Diabetes mellitus (E10-E14)	1,035	4.0	330.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	608	2.3	194.1
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	492	1.9	157.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	482	1.9	153.9
11	Septicemia (A40-A41)	389	1.5	124.2
12	Parkinson's disease (G20-G21)	284	1.1	90.7
13	Pneumonitis due to solids and liquids (J69)	246	0.9	78.5
14	Atherosclerosis (I70)	126	0.5	40.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	123	0.5	39.3
...	All other causes (Residual)	4,902	18.9	1,565.0
Hispanic, male, all ages				
...	All causes	76,861	100.0	316.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,498	20.2	63.9
2	Malignant neoplasms (C00-C97)	15,283	19.9	63.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,363	10.9	34.5
4	Cerebrovascular diseases (I60-I69)	3,370	4.4	13.9
5	Diabetes mellitus (E10-E14)	3,314	4.3	13.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,850	3.7	11.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,777	3.6	11.4
8	Chronic lower respiratory diseases (J40-J47)	2,019	2.6	8.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,955	2.5	8.1
10	Certain conditions originating in the perinatal period (P00-P96)	1,645	2.1	6.8
11	Influenza and pneumonia (J09-J18)	1,544	2.0	6.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,456	1.9	6.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,061	1.4	4.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,037	1.3	4.3
15	Septicemia (A40-A41)	973	1.3	4.0
...	All other causes (Residual)	13,716	17.8	56.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	587	100.0	27.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	36.1	9.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	11.1	3.0
3	Malignant neoplasms (C00-C97)	52	8.9	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	6.0	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.1	*
6	Septicemia (A40-A41)	14	2.4	*
6	Influenza and pneumonia (J09-J18)	14	2.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	10	1.7	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
12	Anemias (D50-D64)	2	0.3	*
12	Meningitis (G00,G03)	2	0.3	*
12	Acute bronchitis and bronchiolitis (J20-J21)	2	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	145	24.7	6.8
Hispanic, male, 5-14 years				
...	All causes	548	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	34.7	4.4
2	Malignant neoplasms (C00-C97)	95	17.3	2.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	7.5	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	5.3	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.1	*
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	2.4	*
7	Cerebrovascular diseases (I60-I69)	11	2.0	*
8	Influenza and pneumonia (J09-J18)	7	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.1	*
10	Septicemia (A40-A41)	4	0.7	*
10	Anemias (D50-D64)	4	0.7	*
12	Diseases of appendix (K35-K38)	3	0.5	*
13	Meningococcal infection (A39)	2	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
13	Diabetes mellitus (E10-E14)	2	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	120	21.9	2.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	4,079	100.0	103.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,740	42.7	44.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,087	26.6	27.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	424	10.4	10.8
4	Malignant neoplasms (C00-C97)	217	5.3	5.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	2.3	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.0	1.0
7	Legal intervention (Y35,Y89.0)	21	0.5	0.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.4	*
8	Cerebrovascular diseases (I60-I69)	17	0.4	*
10	Influenza and pneumonia (J09-J18)	16	0.4	*
11	Chronic lower respiratory diseases (J40-J47)	14	0.3	*
12	Diabetes mellitus (E10-E14)	13	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.3	*
14	Septicemia (A40-A41)	8	0.2	*
14	Anemias (D50-D64)	8	0.2	*
...	All other causes (Residual)	351	8.6	8.9
Hispanic, male, 25-34 years				
...	All causes	4,839	100.0	107.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,903	39.3	42.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	808	16.7	17.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	468	9.7	10.4
4	Malignant neoplasms (C00-C97)	344	7.1	7.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	264	5.5	5.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	134	2.8	3.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	85	1.8	1.9
8	Cerebrovascular diseases (I60-I69)	81	1.7	1.8
9	Diabetes mellitus (E10-E14)	37	0.8	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	0.6	0.7
11	Influenza and pneumonia (J09-J18)	28	0.6	0.6
12	Legal intervention (Y35,Y89.0)	24	0.5	0.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	0.4	0.4
14	Septicemia (A40-A41)	18	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	15	0.3	*
...	All other causes (Residual)	579	12.0	12.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 35-44 years				
...	All causes	5,924	100.0	158.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,494	25.2	40.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	672	11.3	18.0
3	Malignant neoplasms (C00-C97)	650	11.0	17.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	442	7.5	11.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	419	7.1	11.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	375	6.3	10.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	302	5.1	8.1
8	Cerebrovascular diseases (I60-I69)	188	3.2	5.0
9	Diabetes mellitus (E10-E14)	152	2.6	4.1
10	Viral hepatitis (B15-B19)	78	1.3	2.1
11	Influenza and pneumonia (J09-J18)	65	1.1	1.7
12	Septicemia (A40-A41)	57	1.0	1.5
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	0.8	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	0.6	1.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.5	0.8
15	Chronic lower respiratory diseases (J40-J47)	29	0.5	0.8
...	All other causes (Residual)	886	15.0	23.8
Hispanic, male, 45-54 years				
...	All causes	9,469	100.0	372.5
1	Malignant neoplasms (C00-C97)	1,799	19.0	70.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,717	18.1	67.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,194	12.6	47.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	930	9.8	36.6
5	Cerebrovascular diseases (I60-I69)	386	4.1	15.2
6	Diabetes mellitus (E10-E14)	376	4.0	14.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	363	3.8	14.3
8	Viral hepatitis (B15-B19)	318	3.4	12.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	306	3.2	12.0
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	208	2.2	8.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	125	1.3	4.9
12	Influenza and pneumonia (J09-J18)	114	1.2	4.5
13	Septicemia (A40-A41)	104	1.1	4.1
14	Chronic lower respiratory diseases (J40-J47)	85	0.9	3.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	74	0.8	2.9
...	All other causes (Residual)	1,370	14.5	53.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 55-64 years				
...	All causes	11,461	100.0	831.3
1	Malignant neoplasms (C00-C97)	3,133	27.3	227.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,692	23.5	195.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	782	6.8	56.7
4	Diabetes mellitus (E10-E14)	683	6.0	49.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	673	5.9	48.8
6	Cerebrovascular diseases (I60-I69)	483	4.2	35.0
7	Viral hepatitis (B15-B19)	276	2.4	20.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	249	2.2	18.1
9	Chronic lower respiratory diseases (J40-J47)	203	1.8	14.7
10	Influenza and pneumonia (J09-J18)	174	1.5	12.6
11	Human immunodeficiency virus (HIV) disease (B20-B24)	155	1.4	11.2
12	Septicemia (A40-A41)	146	1.3	10.6
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	1.3	10.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	0.8	7.0
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	82	0.7	5.9
...	All other causes (Residual)	1,488	13.0	107.9
Hispanic, male, 65-74 years				
...	All causes	12,421	100.0	1,826.9
1	Malignant neoplasms (C00-C97)	3,787	30.5	557.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,074	24.7	452.1
3	Diabetes mellitus (E10-E14)	831	6.7	122.2
4	Cerebrovascular diseases (I60-I69)	607	4.9	89.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	443	3.6	65.2
6	Chronic lower respiratory diseases (J40-J47)	405	3.3	59.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	336	2.7	49.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	312	2.5	45.9
9	Influenza and pneumonia (J09-J18)	253	2.0	37.2
10	Septicemia (A40-A41)	220	1.8	32.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	131	1.1	19.3
12	Alzheimer's disease (G30)	118	1.0	17.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	94	0.8	13.8
14	Parkinson's disease (G20-G21)	79	0.6	11.6
15	Aortic aneurysm and dissection (I71)	69	0.6	10.1
...	All other causes (Residual)	1,662	13.4	244.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 75-84 years				
...	All causes	14,851	100.0	4,264.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,970	26.7	1,139.8
2	Malignant neoplasms (C00-C97)	3,702	24.9	1,062.9
3	Cerebrovascular diseases (I60-I69)	951	6.4	273.0
4	Diabetes mellitus (E10-E14)	843	5.7	242.0
5	Chronic lower respiratory diseases (J40-J47)	738	5.0	211.9
6	Influenza and pneumonia (J09-J18)	450	3.0	129.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	410	2.8	117.7
8	Alzheimer's disease (G30)	379	2.6	108.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	2.1	88.7
10	Parkinson's disease (G20-G21)	229	1.5	65.7
11	Septicemia (A40-A41)	221	1.5	63.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	180	1.2	51.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	176	1.2	50.5
14	Pneumonitis due to solids and liquids (J69)	105	0.7	30.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	97	0.7	27.9
...	All other causes (Residual)	2,091	14.1	600.4
Hispanic, male, 85 years and over				
...	All causes	9,394	100.0	8,379.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,928	31.2	2,611.7
2	Malignant neoplasms (C00-C97)	1,493	15.9	1,331.7
3	Cerebrovascular diseases (I60-I69)	622	6.6	554.8
4	Chronic lower respiratory diseases (J40-J47)	516	5.5	460.2
5	Alzheimer's disease (G30)	447	4.8	398.7
6	Influenza and pneumonia (J09-J18)	400	4.3	356.8
7	Diabetes mellitus (E10-E14)	377	4.0	336.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	268	2.9	239.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	198	2.1	176.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	146	1.6	130.2
11	Parkinson's disease (G20-G21)	145	1.5	129.3
12	Septicemia (A40-A41)	144	1.5	128.4
13	Pneumonitis due to solids and liquids (J69)	99	1.1	88.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	50	0.5	44.6
15	Aortic aneurysm and dissection (I71)	44	0.5	39.2
...	All other causes (Residual)	1,517	16.1	1,353.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	62,380	100.0	274.9
1	Malignant neoplasms (C00-C97)	13,568	21.8	59.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,453	21.6	59.3
3	Cerebrovascular diseases (I60-I69)	3,751	6.0	16.5
4	Diabetes mellitus (E10-E14)	3,230	5.2	14.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,717	4.4	12.0
6	Alzheimer's disease (G30)	2,039	3.3	9.0
7	Chronic lower respiratory diseases (J40-J47)	1,930	3.1	8.5
8	Influenza and pneumonia (J09-J18)	1,632	2.6	7.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,447	2.3	6.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,241	2.0	5.5
11	Certain conditions originating in the perinatal period (P00-P96)	1,228	2.0	5.4
12	Septicemia (A40-A41)	1,031	1.7	4.5
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	950	1.5	4.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	856	1.4	3.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	554	0.9	2.4
...	All other causes (Residual)	12,753	20.4	56.2
Hispanic, female, 1-4 years				
...	All causes	442	100.0	21.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	26.5	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	14.7	3.2
3	Malignant neoplasms (C00-C97)	53	12.0	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	10.0	2.2
5	Influenza and pneumonia (J09-J18)	17	3.8	*
6	Septicemia (A40-A41)	14	3.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.0	*
8	Cerebrovascular diseases (I60-I69)	8	1.8	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	7	1.6	*
11	Anemias (D50-D64)	2	0.5	*
11	Pneumonitis due to solids and liquids (J69)	2	0.5	*
13	Tuberculosis (A16-A19)	1	0.2	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
13	Peptic ulcer (K25-K28)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	89	20.1	4.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	472	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	24.2	2.8
2	Malignant neoplasms (C00-C97)	91	19.3	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	8.3	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.7	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	4.4	0.5
6	Cerebrovascular diseases (I60-I69)	15	3.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.5	*
8	Chronic lower respiratory diseases (J40-J47)	10	2.1	*
9	Influenza and pneumonia (J09-J18)	9	1.9	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	1.9	*
11	Septicemia (A40-A41)	7	1.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	1.1	*
14	Diabetes mellitus (E10-E14)	4	0.8	*
14	Meningitis (G00,G03)	4	0.8	*
...	All other causes (Residual)	105	22.2	2.5
Hispanic, female, 15-24 years				
...	All causes	1,124	100.0	31.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	412	36.7	11.5
2	Malignant neoplasms (C00-C97)	133	11.8	3.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	130	11.6	3.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	7.7	2.4
5	Pregnancy, childbirth and the puerperium (O00-O99)	39	3.5	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.8	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	2.2	0.7
8	Cerebrovascular diseases (I60-I69)	16	1.4	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.3	*
10	Influenza and pneumonia (J09-J18)	13	1.2	*
11	Septicemia (A40-A41)	11	1.0	*
12	Anemias (D50-D64)	9	0.8	*
13	Diabetes mellitus (E10-E14)	8	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	6	0.5	*
...	All other causes (Residual)	182	16.2	5.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-34 years				
...	All causes	1,557	100.0	42.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	357	22.9	9.7
2	Malignant neoplasms (C00-C97)	299	19.2	8.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	8.5	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	6.3	2.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	5.3	2.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	74	4.8	2.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	41	2.6	1.1
8	Cerebrovascular diseases (I60-I69)	36	2.3	1.0
9	Influenza and pneumonia (J09-J18)	27	1.7	0.7
10	Diabetes mellitus (E10-E14)	25	1.6	0.7
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	1.5	0.6
12	Septicemia (A40-A41)	19	1.2	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	17	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	14	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.8	*
...	All other causes (Residual)	298	19.1	8.1
Hispanic, female, 35-44 years				
...	All causes	2,697	100.0	82.2
1	Malignant neoplasms (C00-C97)	850	31.5	25.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	339	12.6	10.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	213	7.9	6.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	114	4.2	3.5
5	Cerebrovascular diseases (I60-I69)	113	4.2	3.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	103	3.8	3.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	3.4	2.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	3.0	2.4
9	Diabetes mellitus (E10-E14)	73	2.7	2.2
10	Influenza and pneumonia (J09-J18)	37	1.4	1.1
11	Septicemia (A40-A41)	36	1.3	1.1
12	Pregnancy, childbirth and the puerperium (O00-O99)	32	1.2	1.0
13	Viral hepatitis (B15-B19)	31	1.1	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	1.1	0.9
15	Chronic lower respiratory diseases (J40-J47)	27	1.0	0.8
...	All other causes (Residual)	527	19.5	16.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-54 years				
...	All causes	4,901	100.0	200.8
1	Malignant neoplasms (C00-C97)	1,751	35.7	71.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	594	12.1	24.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	347	7.1	14.2
4	Cerebrovascular diseases (I60-I69)	252	5.1	10.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	252	5.1	10.3
6	Diabetes mellitus (E10-E14)	225	4.6	9.2
7	Human immunodeficiency virus (HIV) disease (B20-B24)	104	2.1	4.3
8	Viral hepatitis (B15-B19)	94	1.9	3.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	1.7	3.4
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	1.6	3.2
11	Septicemia (A40-A41)	75	1.5	3.1
12	Influenza and pneumonia (J09-J18)	71	1.4	2.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	1.2	2.4
14	Chronic lower respiratory diseases (J40-J47)	54	1.1	2.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	38	0.8	1.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	0.8	1.6
...	All other causes (Residual)	784	16.0	32.1
Hispanic, female, 55-64 years				
...	All causes	7,041	100.0	477.2
1	Malignant neoplasms (C00-C97)	2,641	37.5	179.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,175	16.7	79.6
3	Diabetes mellitus (E10-E14)	511	7.3	34.6
4	Cerebrovascular diseases (I60-I69)	312	4.4	21.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	244	3.5	16.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	225	3.2	15.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	2.7	12.9
8	Chronic lower respiratory diseases (J40-J47)	149	2.1	10.1
9	Viral hepatitis (B15-B19)	132	1.9	8.9
10	Septicemia (A40-A41)	118	1.7	8.0
11	Influenza and pneumonia (J09-J18)	116	1.6	7.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	98	1.4	6.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	45	0.6	3.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	0.4	2.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.4	1.8
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	26	0.4	1.8
...	All other causes (Residual)	1,004	14.3	68.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65-74 years				
...	All causes	9,522	100.0	1,153.1
1	Malignant neoplasms (C00-C97)	2,899	30.4	351.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,029	21.3	245.7
3	Diabetes mellitus (E10-E14)	754	7.9	91.3
4	Cerebrovascular diseases (I60-I69)	552	5.8	66.8
5	Chronic lower respiratory diseases (J40-J47)	332	3.5	40.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	306	3.2	37.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	296	3.1	35.8
8	Influenza and pneumonia (J09-J18)	195	2.0	23.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	1.9	22.4
10	Septicemia (A40-A41)	172	1.8	20.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	120	1.3	14.5
12	Alzheimer's disease (G30)	115	1.2	13.9
13	Viral hepatitis (B15-B19)	77	0.8	9.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.7	8.2
15	Parkinson's disease (G20-G21)	53	0.6	6.4
...	All other causes (Residual)	1,369	14.4	165.8
Hispanic, female, 75-84 years				
...	All causes	15,411	100.0	3,118.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,799	24.7	768.8
2	Malignant neoplasms (C00-C97)	3,251	21.1	657.9
3	Cerebrovascular diseases (I60-I69)	1,182	7.7	239.2
4	Diabetes mellitus (E10-E14)	971	6.3	196.5
5	Chronic lower respiratory diseases (J40-J47)	672	4.4	136.0
6	Alzheimer's disease (G30)	613	4.0	124.1
7	Influenza and pneumonia (J09-J18)	458	3.0	92.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	425	2.8	86.0
9	Septicemia (A40-A41)	302	2.0	61.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	1.9	58.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	252	1.6	51.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	236	1.5	47.8
13	Parkinson's disease (G20-G21)	165	1.1	33.4
14	Pneumonitis due to solids and liquids (J69)	89	0.6	18.0
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	89	0.6	18.0
...	All other causes (Residual)	2,618	17.0	529.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 85 years and over				
...	All causes	16,602	100.0	8,254.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,442	32.8	2,705.9
2	Malignant neoplasms (C00-C97)	1,593	9.6	792.1
3	Alzheimer's disease (G30)	1,286	7.7	639.4
4	Cerebrovascular diseases (I60-I69)	1,255	7.6	624.0
5	Influenza and pneumonia (J09-J18)	666	4.0	331.1
6	Chronic lower respiratory diseases (J40-J47)	661	4.0	328.7
7	Diabetes mellitus (E10-E14)	658	4.0	327.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	340	2.0	169.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	336	2.0	167.1
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	294	1.8	146.2
11	Septicemia (A40-A41)	245	1.5	121.8
12	Pneumonitis due to solids and liquids (J69)	147	0.9	73.1
13	Parkinson's disease (G20-G21)	139	0.8	69.1
14	Atherosclerosis (I70)	90	0.5	44.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	85	0.5	42.3
...	All other causes (Residual)	3,365	20.3	1,673.1
Non-Hispanic, white, both sexes, all ages				
...	All causes	1,981,034	100.0	982.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	503,096	25.4	249.4
2	Malignant neoplasms (C00-C97)	457,084	23.1	226.6
3	Chronic lower respiratory diseases (J40-J47)	126,146	6.4	62.5
4	Cerebrovascular diseases (I60-I69)	106,134	5.4	52.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94,722	4.8	47.0
6	Alzheimer's disease (G30)	73,219	3.7	36.3
7	Diabetes mellitus (E10-E14)	49,388	2.5	24.5
8	Influenza and pneumonia (J09-J18)	45,780	2.3	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,439	1.8	17.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,281	1.5	15.0
11	Septicemia (A40-A41)	26,692	1.3	13.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	22,210	1.1	11.0
13	Parkinson's disease (G20-G21)	18,474	0.9	9.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,351	0.9	9.1
15	Pneumonitis due to solids and liquids (J69)	14,135	0.7	7.0
...	All other causes (Residual)	359,883	18.2	178.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,373	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	788	33.2	8.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	256	10.8	2.8
3	Malignant neoplasms (C00-C97)	225	9.5	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	7.1	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	80	3.4	0.9
6	Influenza and pneumonia (J09-J18)	60	2.5	0.7
7	Septicemia (A40-A41)	41	1.7	0.5
8	Cerebrovascular diseases (I60-I69)	30	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	22	0.9	0.2
11	Certain conditions originating in the perinatal period (P00-P96)	16	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	12	0.5	*
13	Anemias (D50-D64)	10	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.4	*
15	Meningococcal infection (A39)	8	0.3	*
15	Meningitis (G00,G03)	8	0.3	*
...	All other causes (Residual)	612	25.8	6.7
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	3,070	100.0	13.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,091	35.5	4.7
2	Malignant neoplasms (C00-C97)	509	16.6	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	169	5.5	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	148	4.8	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	3.8	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	3.4	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	1.6	0.2
7	Influenza and pneumonia (J09-J18)	49	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	47	1.5	0.2
10	Septicemia (A40-A41)	32	1.0	0.1
11	Chronic lower respiratory diseases (J40-J47)	29	0.9	0.1
12	Diabetes mellitus (E10-E14)	18	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	18	0.6	*
14	Anemias (D50-D64)	13	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.4	*
...	All other causes (Residual)	666	21.7	2.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	18,560	100.0	70.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,911	53.4	37.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,025	16.3	11.5
3	Malignant neoplasms (C00-C97)	949	5.1	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	914	4.9	3.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	511	2.8	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	266	1.4	1.0
7	Influenza and pneumonia (J09-J18)	125	0.7	0.5
8	Diabetes mellitus (E10-E14)	110	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	107	0.6	0.4
10	Septicemia (A40-A41)	73	0.4	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	63	0.3	0.2
12	Chronic lower respiratory diseases (J40-J47)	62	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.2	0.1
15	Pneumonitis due to solids and liquids (J69)	28	0.2	0.1
...	All other causes (Residual)	2,327	12.5	8.9
Non-Hispanic, white, both sexes, 25-34 years				
...	All causes	24,852	100.0	101.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,206	41.1	41.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,977	16.0	16.3
3	Malignant neoplasms (C00-C97)	2,084	8.4	8.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,723	6.9	7.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,084	4.4	4.4
6	Diabetes mellitus (E10-E14)	313	1.3	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	258	1.0	1.1
8	Cerebrovascular diseases (I60-I69)	248	1.0	1.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	233	0.9	1.0
10	Influenza and pneumonia (J09-J18)	211	0.8	0.9
11	Septicemia (A40-A41)	192	0.8	0.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	167	0.7	0.7
13	Pregnancy, childbirth and the puerperium (O00-O99)	147	0.6	0.6
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	116	0.5	0.5
15	Chronic lower respiratory diseases (J40-J47)	115	0.5	0.5
...	All other causes (Residual)	3,778	15.2	15.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 35-44 years				
...	All causes	49,091	100.0	180.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,865	24.2	43.5
2	Malignant neoplasms (C00-C97)	8,246	16.8	30.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,190	14.6	26.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,571	11.3	20.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,687	3.4	6.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,058	2.2	3.9
7	Diabetes mellitus (E10-E14)	1,036	2.1	3.8
8	Cerebrovascular diseases (I60-I69)	1,005	2.0	3.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	823	1.7	3.0
10	Influenza and pneumonia (J09-J18)	545	1.1	2.0
11	Chronic lower respiratory diseases (J40-J47)	526	1.1	1.9
12	Septicemia (A40-A41)	514	1.0	1.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	373	0.8	1.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	311	0.6	1.1
15	Viral hepatitis (B15-B19)	256	0.5	0.9
...	All other causes (Residual)	8,085	16.5	29.6
Non-Hispanic, white, both sexes, 45-54 years				
...	All causes	130,423	100.0	410.3
1	Malignant neoplasms (C00-C97)	36,300	27.8	114.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,243	20.1	82.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,697	12.0	49.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,342	5.6	23.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,893	4.5	18.5
6	Chronic lower respiratory diseases (J40-J47)	3,489	2.7	11.0
7	Diabetes mellitus (E10-E14)	3,478	2.7	10.9
8	Cerebrovascular diseases (I60-I69)	3,249	2.5	10.2
9	Viral hepatitis (B15-B19)	1,862	1.4	5.9
10	Septicemia (A40-A41)	1,548	1.2	4.9
11	Influenza and pneumonia (J09-J18)	1,526	1.2	4.8
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,175	0.9	3.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,121	0.9	3.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,034	0.8	3.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	597	0.5	1.9
...	All other causes (Residual)	19,869	15.2	62.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 55-64 years				
...	All causes	220,255	100.0	855.0
1	Malignant neoplasms (C00-C97)	80,232	36.4	311.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	48,535	22.0	188.4
3	Chronic lower respiratory diseases (J40-J47)	12,097	5.5	47.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,936	4.5	38.6
5	Diabetes mellitus (E10-E14)	7,422	3.4	28.8
6	Cerebrovascular diseases (I60-I69)	6,606	3.0	25.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,351	2.9	24.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,993	2.3	19.4
9	Septicemia (A40-A41)	3,080	1.4	12.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,648	1.2	10.3
11	Influenza and pneumonia (J09-J18)	2,610	1.2	10.1
12	Viral hepatitis (B15-B19)	1,769	0.8	6.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,351	0.6	5.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,102	0.5	4.3
15	Aortic aneurysm and dissection (I71)	929	0.4	3.6
...	All other causes (Residual)	30,594	13.9	118.8
Non-Hispanic, white, both sexes, 65-74 years				
...	All causes	316,682	100.0	1,994.4
1	Malignant neoplasms (C00-C97)	114,417	36.1	720.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,415	21.9	437.2
3	Chronic lower respiratory diseases (J40-J47)	28,687	9.1	180.7
4	Cerebrovascular diseases (I60-I69)	12,910	4.1	81.3
5	Diabetes mellitus (E10-E14)	10,223	3.2	64.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,257	2.3	45.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,355	1.7	33.7
8	Influenza and pneumonia (J09-J18)	4,681	1.5	29.5
9	Septicemia (A40-A41)	4,633	1.5	29.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4,118	1.3	25.9
11	Alzheimer's disease (G30)	3,621	1.1	22.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,501	0.8	15.8
13	Parkinson's disease (G20-G21)	2,185	0.7	13.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,103	0.7	13.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,085	0.7	13.1
...	All other causes (Residual)	42,491	13.4	267.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 75-84 years				
...	All causes	549,086	100.0	5,143.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	138,405	25.2	1,296.4
2	Malignant neoplasms (C00-C97)	136,202	24.8	1,275.7
3	Chronic lower respiratory diseases (J40-J47)	46,984	8.6	440.1
4	Cerebrovascular diseases (I60-I69)	33,298	6.1	311.9
5	Alzheimer's disease (G30)	22,140	4.0	207.4
6	Diabetes mellitus (E10-E14)	14,578	2.7	136.5
7	Influenza and pneumonia (J09-J18)	12,630	2.3	118.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,153	2.2	113.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11,481	2.1	107.5
10	Septicemia (A40-A41)	8,411	1.5	78.8
11	Parkinson's disease (G20-G21)	8,365	1.5	78.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,963	0.9	46.5
13	Pneumonitis due to solids and liquids (J69)	4,282	0.8	40.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,171	0.8	39.1
15	Aortic aneurysm and dissection (I71)	3,310	0.6	31.0
...	All other causes (Residual)	87,713	16.0	821.6
Non-Hispanic, white, both sexes, 85 years and over				
...	All causes	654,044	100.0	13,556.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	210,693	32.2	4,367.1
2	Malignant neoplasms (C00-C97)	77,880	11.9	1,614.3
3	Cerebrovascular diseases (I60-I69)	48,563	7.4	1,006.6
4	Alzheimer's disease (G30)	46,725	7.1	968.5
5	Chronic lower respiratory diseases (J40-J47)	34,124	5.2	707.3
6	Influenza and pneumonia (J09-J18)	23,236	3.6	481.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,133	2.3	313.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,232	2.2	295.0
9	Diabetes mellitus (E10-E14)	12,205	1.9	253.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,110	1.4	188.8
11	Septicemia (A40-A41)	8,066	1.2	167.2
12	Parkinson's disease (G20-G21)	7,522	1.2	155.9
13	Pneumonitis due to solids and liquids (J69)	7,271	1.1	150.7
14	Atherosclerosis (I70)	3,979	0.6	82.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3,961	0.6	82.1
...	All other causes (Residual)	131,344	20.1	2,722.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	969,288	100.0	978.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	252,572	26.1	254.9
2	Malignant neoplasms (C00-C97)	238,846	24.6	241.1
3	Chronic lower respiratory diseases (J40-J47)	59,284	6.1	59.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59,173	6.1	59.7
5	Cerebrovascular diseases (I60-I69)	41,092	4.2	41.5
6	Diabetes mellitus (E10-E14)	25,303	2.6	25.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	23,835	2.5	24.1
8	Alzheimer's disease (G30)	21,780	2.2	22.0
9	Influenza and pneumonia (J09-J18)	20,516	2.1	20.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,535	1.8	17.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14,452	1.5	14.6
12	Septicemia (A40-A41)	12,092	1.2	12.2
13	Parkinson's disease (G20-G21)	10,845	1.1	10.9
14	Pneumonitis due to solids and liquids (J69)	7,369	0.8	7.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,088	0.7	7.2
...	All other causes (Residual)	157,506	16.2	159.0
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,365	100.0	29.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	479	35.1	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	10.0	2.9
3	Malignant neoplasms (C00-C97)	127	9.3	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	100	7.3	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	2.9	0.8
6	Influenza and pneumonia (J09-J18)	27	2.0	0.6
7	Septicemia (A40-A41)	17	1.2	*
8	Chronic lower respiratory diseases (J40-J47)	14	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.0	*
9	Cerebrovascular diseases (I60-I69)	13	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	7	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.4	*
14	Anemias (D50-D64)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	367	26.9	7.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,806	100.0	15.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	705	39.0	5.9
2	Malignant neoplasms (C00-C97)	264	14.6	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	108	6.0	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	4.8	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	3.8	0.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	3.4	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.6	0.2
8	Influenza and pneumonia (J09-J18)	27	1.5	0.2
9	Cerebrovascular diseases (I60-I69)	21	1.2	0.2
10	Septicemia (A40-A41)	17	0.9	*
10	Chronic lower respiratory diseases (J40-J47)	17	0.9	*
12	Diabetes mellitus (E10-E14)	11	0.6	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.4	*
14	Anemias (D50-D64)	7	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
...	All other causes (Residual)	370	20.5	3.1
Non-Hispanic, white, male, 15-24 years				
...	All causes	13,575	100.0	100.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,382	54.4	54.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,530	18.6	18.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	689	5.1	5.1
4	Malignant neoplasms (C00-C97)	584	4.3	4.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	351	2.6	2.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	163	1.2	1.2
7	Influenza and pneumonia (J09-J18)	84	0.6	0.6
8	Diabetes mellitus (E10-E14)	69	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	65	0.5	0.5
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.3	0.3
10	Chronic lower respiratory diseases (J40-J47)	34	0.3	0.3
12	Septicemia (A40-A41)	29	0.2	0.2
13	Legal intervention (Y35,Y89.0)	25	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	21	0.2	0.2
...	All other causes (Residual)	1,492	11.0	11.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-34 years				
...	All causes	17,330	100.0	140.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,732	44.6	62.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,202	18.5	25.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,188	6.9	9.6
4	Malignant neoplasms (C00-C97)	1,063	6.1	8.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	783	4.5	6.3
6	Diabetes mellitus (E10-E14)	182	1.1	1.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	150	0.9	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	140	0.8	1.1
9	Cerebrovascular diseases (I60-I69)	134	0.8	1.1
10	Influenza and pneumonia (J09-J18)	117	0.7	0.9
11	Human immunodeficiency virus (HIV) disease (B20-B24)	116	0.7	0.9
12	Septicemia (A40-A41)	105	0.6	0.9
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	0.4	0.6
14	Chronic lower respiratory diseases (J40-J47)	55	0.3	0.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.3	0.4
...	All other causes (Residual)	2,240	12.9	18.1
Non-Hispanic, white, male, 35-44 years				
...	All causes	30,810	100.0	225.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,158	26.5	59.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,108	16.6	37.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,215	13.7	30.8
4	Malignant neoplasms (C00-C97)	3,663	11.9	26.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,089	3.5	8.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	722	2.3	5.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	662	2.1	4.8
8	Diabetes mellitus (E10-E14)	645	2.1	4.7
9	Cerebrovascular diseases (I60-I69)	525	1.7	3.8
10	Influenza and pneumonia (J09-J18)	288	0.9	2.1
11	Septicemia (A40-A41)	252	0.8	1.8
12	Chronic lower respiratory diseases (J40-J47)	239	0.8	1.7
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	214	0.7	1.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	180	0.6	1.3
15	Viral hepatitis (B15-B19)	160	0.5	1.2
...	All other causes (Residual)	4,690	15.2	34.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-54 years				
...	All causes	81,331	100.0	515.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,308	23.7	122.4
2	Malignant neoplasms (C00-C97)	18,717	23.0	118.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,515	12.9	66.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,512	6.8	35.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,084	5.0	25.9
6	Diabetes mellitus (E10-E14)	2,237	2.8	14.2
7	Cerebrovascular diseases (I60-I69)	1,768	2.2	11.2
8	Chronic lower respiratory diseases (J40-J47)	1,736	2.1	11.0
9	Viral hepatitis (B15-B19)	1,323	1.6	8.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	961	1.2	6.1
11	Influenza and pneumonia (J09-J18)	834	1.0	5.3
12	Septicemia (A40-A41)	829	1.0	5.3
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	713	0.9	4.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	693	0.9	4.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	400	0.5	2.5
...	All other causes (Residual)	11,701	14.4	74.2
Non-Hispanic, white, male, 55-64 years				
...	All causes	134,290	100.0	1,067.2
1	Malignant neoplasms (C00-C97)	44,417	33.1	353.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,524	25.7	274.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,789	5.1	54.0
4	Chronic lower respiratory diseases (J40-J47)	6,284	4.7	49.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,529	3.4	36.0
6	Diabetes mellitus (E10-E14)	4,514	3.4	35.9
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,859	2.9	30.7
8	Cerebrovascular diseases (I60-I69)	3,700	2.8	29.4
9	Septicemia (A40-A41)	1,630	1.2	13.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,520	1.1	12.1
11	Influenza and pneumonia (J09-J18)	1,513	1.1	12.0
12	Viral hepatitis (B15-B19)	1,270	0.9	10.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	826	0.6	6.6
14	Aortic aneurysm and dissection (I71)	684	0.5	5.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	648	0.5	5.1
...	All other causes (Residual)	17,583	13.1	139.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65-74 years				
...	All causes	178,730	100.0	2,413.0
1	Malignant neoplasms (C00-C97)	62,924	35.2	849.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,081	24.7	595.1
3	Chronic lower respiratory diseases (J40-J47)	14,313	8.0	193.2
4	Cerebrovascular diseases (I60-I69)	6,648	3.7	89.8
5	Diabetes mellitus (E10-E14)	5,834	3.3	78.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,593	2.6	62.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,921	1.6	39.4
8	Influenza and pneumonia (J09-J18)	2,610	1.5	35.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	2,595	1.5	35.0
10	Septicemia (A40-A41)	2,327	1.3	31.4
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,067	1.2	27.9
12	Alzheimer's disease (G30)	1,542	0.9	20.8
13	Parkinson's disease (G20-G21)	1,443	0.8	19.5
14	Aortic aneurysm and dissection (I71)	1,267	0.7	17.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,197	0.7	16.2
...	All other causes (Residual)	22,368	12.5	302.0
Non-Hispanic, white, male, 75-84 years				
...	All causes	272,482	100.0	6,166.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	72,515	26.6	1,641.1
2	Malignant neoplasms (C00-C97)	71,808	26.4	1,625.1
3	Chronic lower respiratory diseases (J40-J47)	22,596	8.3	511.4
4	Cerebrovascular diseases (I60-I69)	14,218	5.2	321.8
5	Alzheimer's disease (G30)	8,143	3.0	184.3
6	Diabetes mellitus (E10-E14)	7,181	2.6	162.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,430	2.4	145.5
8	Influenza and pneumonia (J09-J18)	6,297	2.3	142.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,002	2.2	135.8
10	Parkinson's disease (G20-G21)	5,243	1.9	118.7
11	Septicemia (A40-A41)	3,913	1.4	88.6
12	Pneumonitis due to solids and liquids (J69)	2,411	0.9	54.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,333	0.9	52.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,042	0.7	46.2
15	Aortic aneurysm and dissection (I71)	1,884	0.7	42.6
...	All other causes (Residual)	39,466	14.5	893.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 85 years and over				
...	All causes	230,495	100.0	14,721.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,291	32.7	4,808.9
2	Malignant neoplasms (C00-C97)	35,252	15.3	2,251.6
3	Chronic lower respiratory diseases (J40-J47)	13,992	6.1	893.7
4	Cerebrovascular diseases (I60-I69)	13,970	6.1	892.3
5	Alzheimer's disease (G30)	11,767	5.1	751.6
6	Influenza and pneumonia (J09-J18)	8,658	3.8	553.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,049	2.6	386.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,003	2.6	383.4
9	Diabetes mellitus (E10-E14)	4,628	2.0	295.6
10	Parkinson's disease (G20-G21)	3,903	1.7	249.3
11	Pneumonitis due to solids and liquids (J69)	3,379	1.5	215.8
12	Septicemia (A40-A41)	2,925	1.3	186.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,525	1.1	161.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,702	0.7	108.7
15	Atherosclerosis (I70)	1,185	0.5	75.7
...	All other causes (Residual)	39,266	17.0	2,507.9
Non-Hispanic, white, female, all ages				
...	All causes	1,011,746	100.0	985.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250,524	24.8	244.0
2	Malignant neoplasms (C00-C97)	218,238	21.6	212.6
3	Chronic lower respiratory diseases (J40-J47)	66,862	6.6	65.1
4	Cerebrovascular diseases (I60-I69)	65,042	6.4	63.4
5	Alzheimer's disease (G30)	51,439	5.1	50.1
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,549	3.5	34.6
7	Influenza and pneumonia (J09-J18)	25,264	2.5	24.6
8	Diabetes mellitus (E10-E14)	24,085	2.4	23.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,904	1.8	17.4
10	Septicemia (A40-A41)	14,600	1.4	14.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,263	1.1	11.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	7,758	0.8	7.6
13	Parkinson's disease (G20-G21)	7,629	0.8	7.4
14	Pneumonitis due to solids and liquids (J69)	6,766	0.7	6.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,446	0.6	6.3
...	All other causes (Residual)	202,377	20.0	197.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	1,008	100.0	22.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	309	30.7	7.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	11.9	2.7
3	Malignant neoplasms (C00-C97)	98	9.7	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	6.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	4.1	0.9
6	Influenza and pneumonia (J09-J18)	33	3.3	0.7
7	Septicemia (A40-A41)	24	2.4	0.5
8	Cerebrovascular diseases (I60-I69)	17	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.8	*
11	Certain conditions originating in the perinatal period (P00-P96)	6	0.6	*
12	Meningococcal infection (A39)	5	0.5	*
12	Anemias (D50-D64)	5	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
...	All other causes (Residual)	250	24.8	5.6
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,264	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	386	30.5	3.4
2	Malignant neoplasms (C00-C97)	245	19.4	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	6.6	0.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	3.8	0.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	3.4	0.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	3.2	0.4
7	Cerebrovascular diseases (I60-I69)	26	2.1	0.2
8	Influenza and pneumonia (J09-J18)	22	1.7	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.7	0.2
10	Septicemia (A40-A41)	15	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	12	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	10	0.8	*
13	Diabetes mellitus (E10-E14)	7	0.6	*
14	Anemias (D50-D64)	6	0.5	*
15	Meningitis (G00,G03)	5	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	290	22.9	2.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,985	100.0	39.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,529	50.7	19.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	495	9.9	3.9
3	Malignant neoplasms (C00-C97)	365	7.3	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	225	4.5	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	160	3.2	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	2.1	0.8
7	Pregnancy, childbirth and the puerperium (O00-O99)	63	1.3	0.5
8	Septicemia (A40-A41)	44	0.9	0.3
9	Cerebrovascular diseases (I60-I69)	42	0.8	0.3
10	Diabetes mellitus (E10-E14)	41	0.8	0.3
10	Influenza and pneumonia (J09-J18)	41	0.8	0.3
12	Chronic lower respiratory diseases (J40-J47)	28	0.6	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	10	0.2	*
...	All other causes (Residual)	797	16.0	6.2
Non-Hispanic, white, female, 25-34 years				
...	All causes	7,522	100.0	62.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,474	32.9	20.4
2	Malignant neoplasms (C00-C97)	1,021	13.6	8.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	775	10.3	6.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	535	7.1	4.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	301	4.0	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	147	2.0	1.2
7	Diabetes mellitus (E10-E14)	131	1.7	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	1.6	1.0
9	Cerebrovascular diseases (I60-I69)	114	1.5	0.9
10	Influenza and pneumonia (J09-J18)	94	1.2	0.8
11	Septicemia (A40-A41)	87	1.2	0.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	83	1.1	0.7
13	Chronic lower respiratory diseases (J40-J47)	60	0.8	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	51	0.7	0.4
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.6	0.4
...	All other causes (Residual)	1,487	19.8	12.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 35-44 years				
...	All causes	18,281	100.0	134.4
1	Malignant neoplasms (C00-C97)	4,583	25.1	33.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,707	20.3	27.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,082	11.4	15.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,356	7.4	10.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	598	3.3	4.4
6	Cerebrovascular diseases (I60-I69)	480	2.6	3.5
7	Diabetes mellitus (E10-E14)	391	2.1	2.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	336	1.8	2.5
9	Chronic lower respiratory diseases (J40-J47)	287	1.6	2.1
10	Septicemia (A40-A41)	262	1.4	1.9
11	Influenza and pneumonia (J09-J18)	257	1.4	1.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	161	0.9	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	0.9	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	131	0.7	1.0
15	Viral hepatitis (B15-B19)	96	0.5	0.7
...	All other causes (Residual)	3,395	18.6	25.0
Non-Hispanic, white, female, 45-54 years				
...	All causes	49,092	100.0	306.5
1	Malignant neoplasms (C00-C97)	17,583	35.8	109.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,935	14.1	43.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,182	10.6	32.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,830	3.7	11.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,809	3.7	11.3
6	Chronic lower respiratory diseases (J40-J47)	1,753	3.6	10.9
7	Cerebrovascular diseases (I60-I69)	1,481	3.0	9.2
8	Diabetes mellitus (E10-E14)	1,241	2.5	7.7
9	Septicemia (A40-A41)	719	1.5	4.5
10	Influenza and pneumonia (J09-J18)	692	1.4	4.3
11	Viral hepatitis (B15-B19)	539	1.1	3.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	428	0.9	2.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	321	0.7	2.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	273	0.6	1.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	220	0.4	1.4
...	All other causes (Residual)	8,086	16.5	50.5

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 55-64 years				
...	All causes	85,965	100.0	652.3
1	Malignant neoplasms (C00-C97)	35,815	41.7	271.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,011	16.3	106.3
3	Chronic lower respiratory diseases (J40-J47)	5,813	6.8	44.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,147	3.7	23.9
5	Diabetes mellitus (E10-E14)	2,908	3.4	22.1
6	Cerebrovascular diseases (I60-I69)	2,906	3.4	22.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,822	2.1	13.8
8	Septicemia (A40-A41)	1,450	1.7	11.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,134	1.3	8.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,128	1.3	8.6
11	Influenza and pneumonia (J09-J18)	1,097	1.3	8.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	525	0.6	4.0
13	Viral hepatitis (B15-B19)	499	0.6	3.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	454	0.5	3.4
15	Alzheimer's disease (G30)	350	0.4	2.7
...	All other causes (Residual)	12,906	15.0	97.9
Non-Hispanic, white, female, 65-74 years				
...	All causes	137,952	100.0	1,628.5
1	Malignant neoplasms (C00-C97)	51,493	37.3	607.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,334	18.4	299.1
3	Chronic lower respiratory diseases (J40-J47)	14,374	10.4	169.7
4	Cerebrovascular diseases (I60-I69)	6,262	4.5	73.9
5	Diabetes mellitus (E10-E14)	4,389	3.2	51.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,664	1.9	31.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,434	1.8	28.7
8	Septicemia (A40-A41)	2,306	1.7	27.2
9	Alzheimer's disease (G30)	2,079	1.5	24.5
10	Influenza and pneumonia (J09-J18)	2,071	1.5	24.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,523	1.1	18.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	969	0.7	11.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	888	0.6	10.5
14	Parkinson's disease (G20-G21)	742	0.5	8.8
15	Aortic aneurysm and dissection (I71)	638	0.5	7.5
...	All other causes (Residual)	19,786	14.3	233.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 75-84 years				
...	All causes	276,604	100.0	4,420.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	65,890	23.8	1,052.9
2	Malignant neoplasms (C00-C97)	64,394	23.3	1,029.0
3	Chronic lower respiratory diseases (J40-J47)	24,388	8.8	389.7
4	Cerebrovascular diseases (I60-I69)	19,080	6.9	304.9
5	Alzheimer's disease (G30)	13,997	5.1	223.7
6	Diabetes mellitus (E10-E14)	7,397	2.7	118.2
7	Influenza and pneumonia (J09-J18)	6,333	2.3	101.2
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,723	2.1	91.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,479	2.0	87.6
10	Septicemia (A40-A41)	4,498	1.6	71.9
11	Parkinson's disease (G20-G21)	3,122	1.1	49.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,921	1.1	46.7
13	Pneumonitis due to solids and liquids (J69)	1,871	0.7	29.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,838	0.7	29.4
15	Aortic aneurysm and dissection (I71)	1,426	0.5	22.8
...	All other causes (Residual)	48,247	17.4	771.0
Non-Hispanic, white, female, 85 years and over				
...	All causes	423,549	100.0	12,996.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	135,402	32.0	4,154.9
2	Malignant neoplasms (C00-C97)	42,628	10.1	1,308.1
3	Alzheimer's disease (G30)	34,958	8.3	1,072.7
4	Cerebrovascular diseases (I60-I69)	34,593	8.2	1,061.5
5	Chronic lower respiratory diseases (J40-J47)	20,132	4.8	617.8
6	Influenza and pneumonia (J09-J18)	14,578	3.4	447.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,130	2.2	280.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,183	1.9	251.1
9	Diabetes mellitus (E10-E14)	7,577	1.8	232.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,585	1.6	202.1
11	Septicemia (A40-A41)	5,141	1.2	157.8
12	Pneumonitis due to solids and liquids (J69)	3,892	0.9	119.4
13	Parkinson's disease (G20-G21)	3,619	0.9	111.1
14	Atherosclerosis (I70)	2,794	0.7	85.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,259	0.5	69.3
...	All other causes (Residual)	92,078	21.7	2,825.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	285,522	100.0	745.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,918	24.5	182.5
2	Malignant neoplasms (C00-C97)	63,279	22.2	165.2
3	Cerebrovascular diseases (I60-I69)	16,527	5.8	43.1
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,215	4.3	31.9
5	Diabetes mellitus (E10-E14)	11,934	4.2	31.1
6	Chronic lower respiratory diseases (J40-J47)	8,679	3.0	22.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,564	3.0	22.4
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,206	2.9	21.4
9	Septicemia (A40-A41)	6,359	2.2	16.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5,686	2.0	14.8
11	Influenza and pneumonia (J09-J18)	5,396	1.9	14.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,028	1.8	13.1
13	Alzheimer's disease (G30)	5,024	1.8	13.1
14	Certain conditions originating in the perinatal period (P00-P96)	4,627	1.6	12.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,464	0.9	6.4
...	All other causes (Residual)	51,616	18.1	134.7
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,076	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	282	26.2	11.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	151	14.0	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	10.4	4.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	5.9	2.6
5	Malignant neoplasms (C00-C97)	51	4.7	2.1
6	Influenza and pneumonia (J09-J18)	44	4.1	1.8
7	Chronic lower respiratory diseases (J40-J47)	22	2.0	0.9
8	Septicemia (A40-A41)	21	2.0	0.9
9	Anemias (D50-D64)	18	1.7	*
10	Certain conditions originating in the perinatal period (P00-P96)	15	1.4	*
11	Cerebrovascular diseases (I60-I69)	13	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	258	24.0	10.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,258	100.0	20.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	380	30.2	6.3
2	Malignant neoplasms (C00-C97)	148	11.8	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	133	10.6	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	73	5.8	1.2
5	Chronic lower respiratory diseases (J40-J47)	72	5.7	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	5.1	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	2.4	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.7	0.3
9	Cerebrovascular diseases (I60-I69)	19	1.5	*
10	Influenza and pneumonia (J09-J18)	16	1.3	*
11	Anemias (D50-D64)	14	1.1	*
12	Septicemia (A40-A41)	13	1.0	*
13	Diabetes mellitus (E10-E14)	12	1.0	*
14	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	252	20.0	4.2
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	7,092	100.0	109.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,992	42.2	46.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,480	20.9	22.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	501	7.1	7.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	372	5.2	5.7
5	Malignant neoplasms (C00-C97)	280	3.9	4.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	113	1.6	1.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	1.5	1.6
8	Chronic lower respiratory diseases (J40-J47)	75	1.1	1.2
9	Diabetes mellitus (E10-E14)	67	0.9	1.0
10	Anemias (D50-D64)	64	0.9	1.0
11	Pregnancy, childbirth and the puerperium (O00-O99)	56	0.8	0.9
12	Septicemia (A40-A41)	45	0.6	0.7
13	Cerebrovascular diseases (I60-I69)	41	0.6	0.6
13	Influenza and pneumonia (J09-J18)	41	0.6	0.6
15	Legal intervention (Y35,Y89.0)	37	0.5	0.6
...	All other causes (Residual)	825	11.6	12.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-34 years				
...	All causes	9,116	100.0	166.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,439	26.8	44.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,543	16.9	28.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,043	11.4	19.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	603	6.6	11.0
5	Malignant neoplasms (C00-C97)	567	6.2	10.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	491	5.4	8.9
7	Diabetes mellitus (E10-E14)	173	1.9	3.1
8	Cerebrovascular diseases (I60-I69)	142	1.6	2.6
9	Anemias (D50-D64)	125	1.4	2.3
10	Septicemia (A40-A41)	113	1.2	2.1
11	Pregnancy, childbirth and the puerperium (O00-O99)	105	1.2	1.9
12	Chronic lower respiratory diseases (J40-J47)	99	1.1	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	1.1	1.7
14	Influenza and pneumonia (J09-J18)	82	0.9	1.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	0.7	1.2
...	All other causes (Residual)	1,429	15.7	26.0
Non-Hispanic, black, both sexes, 35-44 years				
...	All causes	15,579	100.0	291.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,872	18.4	53.7
2	Malignant neoplasms (C00-C97)	2,313	14.8	43.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,838	11.8	34.4
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,542	9.9	28.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,203	7.7	22.5
6	Cerebrovascular diseases (I60-I69)	613	3.9	11.5
7	Diabetes mellitus (E10-E14)	522	3.4	9.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	392	2.5	7.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	301	1.9	5.6
10	Septicemia (A40-A41)	247	1.6	4.6
11	Chronic lower respiratory diseases (J40-J47)	205	1.3	3.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	196	1.3	3.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	189	1.2	3.5
14	Influenza and pneumonia (J09-J18)	177	1.1	3.3
15	Anemias (D50-D64)	151	1.0	2.8
...	All other causes (Residual)	2,818	18.1	52.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-54 years				
...	All causes	35,527	100.0	682.0
1	Malignant neoplasms (C00-C97)	8,654	24.4	166.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,200	23.1	157.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,504	7.0	48.0
4	Human immunodeficiency virus (HIV) disease (B20-B24)	2,018	5.7	38.7
5	Cerebrovascular diseases (I60-I69)	1,898	5.3	36.4
6	Diabetes mellitus (E10-E14)	1,344	3.8	25.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	820	2.3	15.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	801	2.3	15.4
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	760	2.1	14.6
10	Septicemia (A40-A41)	705	2.0	13.5
11	Chronic lower respiratory diseases (J40-J47)	662	1.9	12.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	588	1.7	11.3
13	Influenza and pneumonia (J09-J18)	463	1.3	8.9
14	Viral hepatitis (B15-B19)	368	1.0	7.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	325	0.9	6.2
...	All other causes (Residual)	5,417	15.2	104.0
Non-Hispanic, black, both sexes, 55-64 years				
...	All causes	48,228	100.0	1,416.8
1	Malignant neoplasms (C00-C97)	14,841	30.8	436.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,295	25.5	361.2
3	Cerebrovascular diseases (I60-I69)	2,542	5.3	74.7
4	Diabetes mellitus (E10-E14)	2,280	4.7	67.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,591	3.3	46.7
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,501	3.1	44.1
7	Chronic lower respiratory diseases (J40-J47)	1,343	2.8	39.5
8	Septicemia (A40-A41)	1,068	2.2	31.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,066	2.2	31.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	849	1.8	24.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	837	1.7	24.6
12	Influenza and pneumonia (J09-J18)	696	1.4	20.4
13	Viral hepatitis (B15-B19)	461	1.0	13.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	261	0.5	7.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	161	0.3	4.7
...	All other causes (Residual)	6,436	13.3	189.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65-74 years				
...	All causes	51,220	100.0	2,787.1
1	Malignant neoplasms (C00-C97)	15,965	31.2	868.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,395	26.2	728.9
3	Cerebrovascular diseases (I60-I69)	3,032	5.9	165.0
4	Diabetes mellitus (E10-E14)	2,851	5.6	155.1
5	Chronic lower respiratory diseases (J40-J47)	2,100	4.1	114.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,938	3.8	105.5
7	Septicemia (A40-A41)	1,264	2.5	68.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	909	1.8	49.5
9	Influenza and pneumonia (J09-J18)	885	1.7	48.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	876	1.7	47.7
11	Alzheimer's disease (G30)	396	0.8	21.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	381	0.7	20.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	275	0.5	15.0
14	Pneumonitis due to solids and liquids (J69)	236	0.5	12.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	217	0.4	11.8
...	All other causes (Residual)	6,500	12.7	353.7
Non-Hispanic, black, both sexes, 75-84 years				
...	All causes	58,481	100.0	5,746.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,155	27.6	1,587.4
2	Malignant neoplasms (C00-C97)	13,772	23.5	1,353.3
3	Cerebrovascular diseases (I60-I69)	4,302	7.4	422.7
4	Diabetes mellitus (E10-E14)	2,923	5.0	287.2
5	Chronic lower respiratory diseases (J40-J47)	2,485	4.2	244.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,112	3.6	207.5
7	Alzheimer's disease (G30)	1,685	2.9	165.6
8	Septicemia (A40-A41)	1,549	2.6	152.2
9	Influenza and pneumonia (J09-J18)	1,401	2.4	137.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,172	2.0	115.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	735	1.3	72.2
12	Pneumonitis due to solids and liquids (J69)	445	0.8	43.7
13	Parkinson's disease (G20-G21)	337	0.6	33.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	279	0.5	27.4
15	Aortic aneurysm and dissection (I71)	205	0.4	20.1
...	All other causes (Residual)	8,924	15.3	876.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 85 years and over				
...	All causes	49,738	100.0	12,239.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,343	30.8	3,775.4
2	Malignant neoplasms (C00-C97)	6,672	13.4	1,641.8
3	Cerebrovascular diseases (I60-I69)	3,885	7.8	956.0
4	Alzheimer's disease (G30)	2,875	5.8	707.5
5	Diabetes mellitus (E10-E14)	1,758	3.5	432.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,730	3.5	425.7
7	Chronic lower respiratory diseases (J40-J47)	1,604	3.2	394.7
8	Influenza and pneumonia (J09-J18)	1,530	3.1	376.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,241	2.5	305.4
10	Septicemia (A40-A41)	1,230	2.5	302.7
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	584	1.2	143.7
12	Pneumonitis due to solids and liquids (J69)	495	1.0	121.8
13	Atherosclerosis (I70)	264	0.5	65.0
14	Parkinson's disease (G20-G21)	240	0.5	59.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	215	0.4	52.9
...	All other causes (Residual)	10,072	20.3	2,478.4
Non-Hispanic, black, male, all ages				
...	All causes	145,168	100.0	794.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,965	24.1	191.3
2	Malignant neoplasms (C00-C97)	32,654	22.5	178.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,282	5.7	45.3
4	Cerebrovascular diseases (I60-I69)	7,135	4.9	39.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,047	4.9	38.6
6	Diabetes mellitus (E10-E14)	5,378	3.7	29.4
7	Chronic lower respiratory diseases (J40-J47)	4,497	3.1	24.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,893	2.7	21.3
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,724	2.6	20.4
10	Septicemia (A40-A41)	2,841	2.0	15.5
11	Certain conditions originating in the perinatal period (P00-P96)	2,608	1.8	14.3
12	Influenza and pneumonia (J09-J18)	2,542	1.8	13.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,194	1.5	12.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,721	1.2	9.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,653	1.1	9.0
...	All other causes (Residual)	24,034	16.6	131.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	619	100.0	50.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	170	27.5	13.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	14.5	7.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	10.3	5.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	5.7	2.8
5	Malignant neoplasms (C00-C97)	32	5.2	2.6
6	Influenza and pneumonia (J09-J18)	26	4.2	2.1
7	Septicemia (A40-A41)	11	1.8	*
7	Chronic lower respiratory diseases (J40-J47)	11	1.8	*
9	Anemias (D50-D64)	9	1.5	*
10	Cerebrovascular diseases (I60-I69)	8	1.3	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (Residual)	144	23.3	11.6
Non-Hispanic, black, male, 5-14 years				
...	All causes	750	100.0	24.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	242	32.3	7.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	83	11.1	2.7
3	Malignant neoplasms (C00-C97)	68	9.1	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	6.1	1.5
5	Chronic lower respiratory diseases (J40-J47)	44	5.9	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.7	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	2.9	0.7
8	Cerebrovascular diseases (I60-I69)	14	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.7	*
10	Influenza and pneumonia (J09-J18)	10	1.3	*
11	Diabetes mellitus (E10-E14)	7	0.9	*
12	Septicemia (A40-A41)	6	0.8	*
12	Anemias (D50-D64)	6	0.8	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	147	19.6	4.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,423	100.0	164.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,672	49.3	81.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,101	20.3	33.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	417	7.7	12.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	255	4.7	7.7
5	Malignant neoplasms (C00-C97)	163	3.0	4.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	63	1.2	1.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	59	1.1	1.8
8	Anemias (D50-D64)	43	0.8	1.3
9	Diabetes mellitus (E10-E14)	41	0.8	1.2
9	Chronic lower respiratory diseases (J40-J47)	41	0.8	1.2
11	Legal intervention (Y35,Y89.0)	36	0.7	1.1
12	Cerebrovascular diseases (I60-I69)	23	0.4	0.7
13	Septicemia (A40-A41)	22	0.4	0.7
14	Influenza and pneumonia (J09-J18)	18	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.3	*
...	All other causes (Residual)	454	8.4	13.8
Non-Hispanic, black, male, 25-34 years				
...	All causes	6,248	100.0	234.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,190	35.1	82.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,171	18.7	43.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	634	10.1	23.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	413	6.6	15.5
5	Human immunodeficiency virus (HIV) disease (B20-B24)	334	5.3	12.5
6	Malignant neoplasms (C00-C97)	260	4.2	9.7
7	Diabetes mellitus (E10-E14)	84	1.3	3.1
8	Cerebrovascular diseases (I60-I69)	66	1.1	2.5
9	Chronic lower respiratory diseases (J40-J47)	59	0.9	2.2
10	Anemias (D50-D64)	56	0.9	2.1
11	Septicemia (A40-A41)	54	0.9	2.0
12	Influenza and pneumonia (J09-J18)	47	0.8	1.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	0.7	1.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	43	0.7	1.6
15	Legal intervention (Y35,Y89.0)	42	0.7	1.6
...	All other causes (Residual)	749	12.0	28.1

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 35-44 years				
...	All causes	9,070	100.0	361.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,780	19.6	70.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,297	14.3	51.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,002	11.0	39.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	934	10.3	37.2
5	Malignant neoplasms (C00-C97)	891	9.8	35.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	323	3.6	12.9
7	Cerebrovascular diseases (I60-I69)	302	3.3	12.0
8	Diabetes mellitus (E10-E14)	293	3.2	11.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	168	1.9	6.7
10	Septicemia (A40-A41)	137	1.5	5.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	113	1.2	4.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	112	1.2	4.5
13	Chronic lower respiratory diseases (J40-J47)	106	1.2	4.2
14	Influenza and pneumonia (J09-J18)	102	1.1	4.1
15	Anemias (D50-D64)	70	0.8	2.8
...	All other causes (Residual)	1,440	15.9	57.3
Non-Hispanic, black, male, 45-54 years				
...	All causes	20,404	100.0	848.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,971	24.4	206.7
2	Malignant neoplasms (C00-C97)	4,272	20.9	177.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,739	8.5	72.3
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,377	6.7	57.3
5	Cerebrovascular diseases (I60-I69)	1,015	5.0	42.2
6	Diabetes mellitus (E10-E14)	753	3.7	31.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	631	3.1	26.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	547	2.7	22.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	436	2.1	18.1
10	Septicemia (A40-A41)	358	1.8	14.9
11	Chronic lower respiratory diseases (J40-J47)	326	1.6	13.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	325	1.6	13.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	269	1.3	11.2
14	Viral hepatitis (B15-B19)	255	1.2	10.6
15	Influenza and pneumonia (J09-J18)	254	1.2	10.6
...	All other causes (Residual)	2,876	14.1	119.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 55-64 years				
...	All causes	28,344	100.0	1,869.1
1	Malignant neoplasms (C00-C97)	8,269	29.2	545.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,647	27.0	504.3
3	Cerebrovascular diseases (I60-I69)	1,400	4.9	92.3
4	Diabetes mellitus (E10-E14)	1,200	4.2	79.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,162	4.1	76.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	813	2.9	53.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	785	2.8	51.8
8	Chronic lower respiratory diseases (J40-J47)	737	2.6	48.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	615	2.2	40.6
10	Septicemia (A40-A41)	551	1.9	36.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	481	1.7	31.7
12	Influenza and pneumonia (J09-J18)	413	1.5	27.2
13	Viral hepatitis (B15-B19)	331	1.2	21.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	210	0.7	13.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	138	0.5	9.1
...	All other causes (Residual)	3,592	12.7	236.9
Non-Hispanic, black, male, 65-74 years				
...	All causes	27,653	100.0	3,610.9
1	Malignant neoplasms (C00-C97)	8,650	31.3	1,129.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,651	27.7	999.1
3	Cerebrovascular diseases (I60-I69)	1,609	5.8	210.1
4	Diabetes mellitus (E10-E14)	1,350	4.9	176.3
5	Chronic lower respiratory diseases (J40-J47)	1,152	4.2	150.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	934	3.4	122.0
7	Septicemia (A40-A41)	600	2.2	78.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	579	2.1	75.6
9	Influenza and pneumonia (J09-J18)	458	1.7	59.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	442	1.6	57.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	250	0.9	32.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	189	0.7	24.7
13	Alzheimer's disease (G30)	175	0.6	22.9
14	Pneumonitis due to solids and liquids (J69)	145	0.5	18.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	108	0.4	14.1
...	All other causes (Residual)	3,361	12.2	438.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 75-84 years				
...	All causes	26,761	100.0	7,234.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,386	27.6	1,996.6
2	Malignant neoplasms (C00-C97)	7,088	26.5	1,916.0
3	Cerebrovascular diseases (I60-I69)	1,676	6.3	453.1
4	Chronic lower respiratory diseases (J40-J47)	1,328	5.0	359.0
5	Diabetes mellitus (E10-E14)	1,176	4.4	317.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	898	3.4	242.7
7	Influenza and pneumonia (J09-J18)	678	2.5	183.3
8	Septicemia (A40-A41)	653	2.4	176.5
9	Alzheimer's disease (G30)	574	2.1	155.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	460	1.7	124.3
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	401	1.5	108.4
12	Pneumonitis due to solids and liquids (J69)	231	0.9	62.4
13	Parkinson's disease (G20-G21)	177	0.7	47.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	130	0.5	35.1
15	Aortic aneurysm and dissection (I71)	87	0.3	23.5
...	All other causes (Residual)	3,818	14.3	1,032.1
Non-Hispanic, black, male, 85 years and over				
...	All causes	15,336	100.0	12,742.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,506	29.4	3,744.0
2	Malignant neoplasms (C00-C97)	2,949	19.2	2,450.3
3	Cerebrovascular diseases (I60-I69)	1,001	6.5	831.7
4	Chronic lower respiratory diseases (J40-J47)	682	4.4	566.7
5	Alzheimer's disease (G30)	614	4.0	510.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	554	3.6	460.3
7	Influenza and pneumonia (J09-J18)	503	3.3	417.9
8	Diabetes mellitus (E10-E14)	471	3.1	391.4
9	Septicemia (A40-A41)	395	2.6	328.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	319	2.1	265.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	207	1.3	172.0
12	Pneumonitis due to solids and liquids (J69)	198	1.3	164.5
13	Parkinson's disease (G20-G21)	111	0.7	92.2
14	Atherosclerosis (I70)	77	0.5	64.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	72	0.5	59.8
...	All other causes (Residual)	2,677	17.5	2,224.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	140,354	100.0	700.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,953	24.9	174.4
2	Malignant neoplasms (C00-C97)	30,625	21.8	152.8
3	Cerebrovascular diseases (I60-I69)	9,392	6.7	46.9
4	Diabetes mellitus (E10-E14)	6,556	4.7	32.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,671	3.3	23.3
6	Chronic lower respiratory diseases (J40-J47)	4,182	3.0	20.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,933	2.8	19.6
8	Alzheimer's disease (G30)	3,630	2.6	18.1
9	Septicemia (A40-A41)	3,518	2.5	17.6
10	Influenza and pneumonia (J09-J18)	2,854	2.0	14.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,834	2.0	14.1
12	Certain conditions originating in the perinatal period (P00-P96)	2,019	1.4	10.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,962	1.4	9.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,159	0.8	5.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	811	0.6	4.0
...	All other causes (Residual)	27,255	19.4	136.0
Non-Hispanic, black, female, 1-4 years				
...	All causes	457	100.0	38.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	24.5	9.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	13.3	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	10.5	4.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	6.3	2.4
5	Malignant neoplasms (C00-C97)	19	4.2	*
6	Influenza and pneumonia (J09-J18)	18	3.9	*
7	Chronic lower respiratory diseases (J40-J47)	11	2.4	*
8	Septicemia (A40-A41)	10	2.2	*
9	Anemias (D50-D64)	9	2.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	8	1.8	*
11	Cerebrovascular diseases (I60-I69)	5	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	5	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
15	Salmonella infections (A01-A02)	1	0.2	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	111	24.3	9.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	508	100.0	17.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	138	27.2	4.7
2	Malignant neoplasms (C00-C97)	80	15.7	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	9.8	1.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.7	1.0
5	Chronic lower respiratory diseases (J40-J47)	28	5.5	0.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	5.3	0.9
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.6	*
7	Anemias (D50-D64)	8	1.6	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	1.6	*
10	Septicemia (A40-A41)	7	1.4	*
11	Influenza and pneumonia (J09-J18)	6	1.2	*
12	Diabetes mellitus (E10-E14)	5	1.0	*
12	Cerebrovascular diseases (I60-I69)	5	1.0	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
15	Meningococcal infection (A39)	1	0.2	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Diseases of appendix (K35-K38)	1	0.2	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	97	19.1	3.3
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,669	100.0	52.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	379	22.7	11.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	320	19.2	10.0
3	Malignant neoplasms (C00-C97)	117	7.0	3.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	117	7.0	3.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	5.0	2.6
6	Pregnancy, childbirth and the puerperium (O00-O99)	56	3.4	1.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	54	3.2	1.7
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	2.4	1.2
9	Chronic lower respiratory diseases (J40-J47)	34	2.0	1.1
10	Diabetes mellitus (E10-E14)	26	1.6	0.8
11	Septicemia (A40-A41)	23	1.4	0.7
11	Influenza and pneumonia (J09-J18)	23	1.4	0.7
13	Anemias (D50-D64)	21	1.3	0.7
14	Cerebrovascular diseases (I60-I69)	18	1.1	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.8	*
...	All other causes (Residual)	331	19.8	10.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-34 years				
...	All causes	2,868	100.0	101.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	409	14.3	14.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	372	13.0	13.2
3	Malignant neoplasms (C00-C97)	307	10.7	10.9
4	Human immunodeficiency virus (HIV) disease (B20-B24)	269	9.4	9.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	249	8.7	8.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	105	3.7	3.7
7	Diabetes mellitus (E10-E14)	89	3.1	3.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	78	2.7	2.8
9	Cerebrovascular diseases (I60-I69)	76	2.6	2.7
10	Anemias (D50-D64)	69	2.4	2.4
11	Septicemia (A40-A41)	59	2.1	2.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	1.7	1.8
13	Chronic lower respiratory diseases (J40-J47)	40	1.4	1.4
14	Influenza and pneumonia (J09-J18)	35	1.2	1.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.8	0.8
...	All other causes (Residual)	637	22.2	22.6
Non-Hispanic, black, female, 35-44 years				
...	All causes	6,509	100.0	229.8
1	Malignant neoplasms (C00-C97)	1,422	21.8	50.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,092	16.8	38.6
3	Human immunodeficiency virus (HIV) disease (B20-B24)	608	9.3	21.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	541	8.3	19.1
5	Cerebrovascular diseases (I60-I69)	311	4.8	11.0
6	Diabetes mellitus (E10-E14)	229	3.5	8.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	201	3.1	7.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	133	2.0	4.7
9	Septicemia (A40-A41)	110	1.7	3.9
10	Chronic lower respiratory diseases (J40-J47)	99	1.5	3.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	83	1.3	2.9
12	Anemias (D50-D64)	81	1.2	2.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	77	1.2	2.7
14	Influenza and pneumonia (J09-J18)	75	1.2	2.6
15	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.1	2.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	69	1.1	2.4
...	All other causes (Residual)	1,309	20.1	46.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-54 years				
...	All causes	15,123	100.0	539.2
1	Malignant neoplasms (C00-C97)	4,382	29.0	156.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,229	21.4	115.1
3	Cerebrovascular diseases (I60-I69)	883	5.8	31.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	765	5.1	27.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	641	4.2	22.9
6	Diabetes mellitus (E10-E14)	591	3.9	21.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	365	2.4	13.0
8	Septicemia (A40-A41)	347	2.3	12.4
9	Chronic lower respiratory diseases (J40-J47)	336	2.2	12.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	273	1.8	9.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	263	1.7	9.4
12	Influenza and pneumonia (J09-J18)	209	1.4	7.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	0.9	4.6
14	Viral hepatitis (B15-B19)	113	0.7	4.0
15	Anemias (D50-D64)	61	0.4	2.2
...	All other causes (Residual)	2,536	16.8	90.4
Non-Hispanic, black, female, 55-64 years				
...	All causes	19,884	100.0	1,053.4
1	Malignant neoplasms (C00-C97)	6,572	33.1	348.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,648	23.4	246.2
3	Cerebrovascular diseases (I60-I69)	1,142	5.7	60.5
4	Diabetes mellitus (E10-E14)	1,080	5.4	57.2
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	688	3.5	36.4
6	Chronic lower respiratory diseases (J40-J47)	606	3.0	32.1
7	Septicemia (A40-A41)	517	2.6	27.4
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	429	2.2	22.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	356	1.8	18.9
10	Influenza and pneumonia (J09-J18)	283	1.4	15.0
11	Human immunodeficiency virus (HIV) disease (B20-B24)	281	1.4	14.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	234	1.2	12.4
13	Viral hepatitis (B15-B19)	130	0.7	6.9
14	Anemias (D50-D64)	68	0.3	3.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	0.3	3.3
...	All other causes (Residual)	2,788	14.0	147.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65-74 years				
...	All causes	23,567	100.0	2,198.5
1	Malignant neoplasms (C00-C97)	7,315	31.0	682.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,744	24.4	535.9
3	Diabetes mellitus (E10-E14)	1,501	6.4	140.0
4	Cerebrovascular diseases (I60-I69)	1,423	6.0	132.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,004	4.3	93.7
6	Chronic lower respiratory diseases (J40-J47)	948	4.0	88.4
7	Septicemia (A40-A41)	664	2.8	61.9
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	467	2.0	43.6
9	Influenza and pneumonia (J09-J18)	427	1.8	39.8
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	297	1.3	27.7
11	Alzheimer's disease (G30)	221	0.9	20.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	131	0.6	12.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	109	0.5	10.2
14	Pneumonitis due to solids and liquids (J69)	91	0.4	8.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	86	0.4	8.0
...	All other causes (Residual)	3,139	13.3	292.8
Non-Hispanic, black, female, 75-84 years				
...	All causes	31,720	100.0	4,896.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,769	27.6	1,353.7
2	Malignant neoplasms (C00-C97)	6,684	21.1	1,031.9
3	Cerebrovascular diseases (I60-I69)	2,626	8.3	405.4
4	Diabetes mellitus (E10-E14)	1,747	5.5	269.7
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,214	3.8	187.4
6	Chronic lower respiratory diseases (J40-J47)	1,157	3.6	178.6
7	Alzheimer's disease (G30)	1,111	3.5	171.5
8	Septicemia (A40-A41)	896	2.8	138.3
9	Influenza and pneumonia (J09-J18)	723	2.3	111.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	712	2.2	109.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	334	1.1	51.6
12	Pneumonitis due to solids and liquids (J69)	214	0.7	33.0
13	Parkinson's disease (G20-G21)	160	0.5	24.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	149	0.5	23.0
15	Aortic aneurysm and dissection (I71)	118	0.4	18.2
...	All other causes (Residual)	5,106	16.1	788.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK5. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2008

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. New ICD-10 code J09 (Influenza due to identified avian influenza virus) was added to the category Influenza and pneumonia in 2007]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 85 years and over				
...	All causes	34,402	100.0	12,027.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,837	31.5	3,788.7
2	Malignant neoplasms (C00-C97)	3,723	10.8	1,301.6
3	Cerebrovascular diseases (I60-I69)	2,884	8.4	1,008.3
4	Alzheimer's disease (G30)	2,261	6.6	790.5
5	Diabetes mellitus (E10-E14)	1,287	3.7	449.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,176	3.4	411.1
7	Influenza and pneumonia (J09-J18)	1,027	3.0	359.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	922	2.7	322.3
8	Chronic lower respiratory diseases (J40-J47)	922	2.7	322.3
10	Septicemia (A40-A41)	835	2.4	291.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	377	1.1	131.8
12	Pneumonitis due to solids and liquids (J69)	297	0.9	103.8
13	Atherosclerosis (I70)	187	0.5	65.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	143	0.4	50.0
15	Parkinson's disease (G20-G21)	129	0.4	45.1
...	All other causes (Residual)	7,395	21.5	2,585.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable