

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	141,576	100.0	292.4
1	Malignant neoplasms (C00-C97)	29,935	21.1	61.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,611	20.9	61.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,654	7.5	22.0
4	Cerebrovascular diseases (I60-I69)	7,065	5.0	14.6
5	Diabetes mellitus (E10-E14)	6,311	4.5	13.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,303	3.0	8.9
7	Chronic lower respiratory diseases (J40-J47)	4,026	2.8	8.3
8	Influenza and pneumonia (J09-J18)	3,679	2.6	7.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,179	2.2	6.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,107	2.2	6.4
11	Alzheimer's disease (G30)	2,986	2.1	6.2
12	Certain conditions originating in the perinatal period (P00-P96)	2,618	1.8	5.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,573	1.8	5.3
14	Septicemia (A40-A41)	2,077	1.5	4.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,851	1.3	3.8
...	All other causes (Residual)	27,601	19.5	57.0
Hispanic, both sexes, 1-4 years				
...	All causes	1,004	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	324	32.3	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	12.4	2.8
3	Malignant neoplasms (C00-C97)	99	9.9	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	75	7.5	1.7
5	Influenza and pneumonia (J09-J18)	34	3.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	2.6	0.6
7	Septicemia (A40-A41)	17	1.7	*
8	Chronic lower respiratory diseases (J40-J47)	13	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.1	*
9	Cerebrovascular diseases (I60-I69)	11	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	10	1.0	*
12	Meningitis (G00,G03)	4	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
...	All other causes (Residual)	247	24.6	5.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,172	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	332	28.3	3.8
2	Malignant neoplasms (C00-C97)	216	18.4	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	6.1	0.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	5.9	0.8
5	Influenza and pneumonia (J09-J18)	63	5.4	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	4.1	0.5
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	42	3.6	0.5
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	2.0	0.3
9	Chronic lower respiratory diseases (J40-J47)	19	1.6	*
10	Septicemia (A40-A41)	15	1.3	*
11	Cerebrovascular diseases (I60-I69)	10	0.9	*
12	Diabetes mellitus (E10-E14)	7	0.6	*
13	Anemias (D50-D64)	6	0.5	*
14	Meningitis (G00,G03)	5	0.4	*
15	Diseases of appendix (K35-K38)	4	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
...	All other causes (Residual)	233	19.9	2.6
Hispanic, both sexes, 15-24 years				
...	All causes	5,187	100.0	65.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,999	38.5	25.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,168	22.5	14.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	583	11.2	7.4
4	Malignant neoplasms (C00-C97)	320	6.2	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	148	2.9	1.9
6	Influenza and pneumonia (J09-J18)	88	1.7	1.1
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	1.4	0.9
8	Cerebrovascular diseases (I60-I69)	42	0.8	0.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	41	0.8	0.5
10	Septicemia (A40-A41)	36	0.7	0.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	30	0.6	0.4
12	Diabetes mellitus (E10-E14)	24	0.5	0.3
13	Chronic lower respiratory diseases (J40-J47)	21	0.4	0.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.3	*
...	All other causes (Residual)	584	11.3	7.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	15,009	100.0	98.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,810	25.4	25.0
2	Malignant neoplasms (C00-C97)	2,141	14.3	14.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,361	9.1	8.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,307	8.7	8.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,104	7.4	7.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	606	4.0	4.0
7	Human immunodeficiency virus (HIV) disease (B20-B24)	483	3.2	3.2
8	Influenza and pneumonia (J09-J18)	441	2.9	2.9
9	Cerebrovascular diseases (I60-I69)	402	2.7	2.6
10	Diabetes mellitus (E10-E14)	305	2.0	2.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	1.1	1.1
12	Septicemia (A40-A41)	154	1.0	1.0
13	Pregnancy, childbirth and the puerperium (O00-O99)	128	0.9	0.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	113	0.8	0.7
15	Chronic lower respiratory diseases (J40-J47)	107	0.7	0.7
...	All other causes (Residual)	2,383	15.9	15.6
Hispanic, both sexes, 45-64 years				
...	All causes	34,477	100.0	422.4
1	Malignant neoplasms (C00-C97)	9,704	28.1	118.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,536	19.0	80.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,444	7.1	29.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,360	6.8	28.9
5	Diabetes mellitus (E10-E14)	1,780	5.2	21.8
6	Cerebrovascular diseases (I60-I69)	1,486	4.3	18.2
7	Viral hepatitis (B15-B19)	770	2.2	9.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	708	2.1	8.7
9	Influenza and pneumonia (J09-J18)	683	2.0	8.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	659	1.9	8.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	620	1.8	7.6
12	Septicemia (A40-A41)	572	1.7	7.0
13	Chronic lower respiratory diseases (J40-J47)	489	1.4	6.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	365	1.1	4.5
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	299	0.9	3.7
...	All other causes (Residual)	5,002	14.5	61.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent	
			of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	79,274	100.0	2,871.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,455	27.1	777.1
2	Malignant neoplasms (C00-C97)	17,438	22.0	631.6
3	Cerebrovascular diseases (I60-I69)	5,081	6.4	184.0
4	Diabetes mellitus (E10-E14)	4,195	5.3	151.9
5	Chronic lower respiratory diseases (J40-J47)	3,370	4.3	122.1
6	Alzheimer's disease (G30)	2,939	3.7	106.4
7	Influenza and pneumonia (J09-J18)	2,316	2.9	83.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,190	2.8	79.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,549	2.0	56.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,328	1.7	48.1
11	Septicemia (A40-A41)	1,233	1.6	44.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,163	1.5	42.1
13	Parkinson's disease (G20-G21)	823	1.0	29.8
14	Pneumonitis due to solids and liquids (J69)	527	0.7	19.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	470	0.6	17.0
...	All other causes (Residual)	13,197	16.6	478.0
Hispanic, male, all ages				
...	All causes	78,157	100.0	311.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,301	20.9	65.1
2	Malignant neoplasms (C00-C97)	15,772	20.2	62.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,841	10.0	31.3
4	Cerebrovascular diseases (I60-I69)	3,288	4.2	13.1
5	Diabetes mellitus (E10-E14)	3,212	4.1	12.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,969	3.8	11.8
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,635	3.4	10.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,135	2.7	8.5
9	Chronic lower respiratory diseases (J40-J47)	2,024	2.6	8.1
10	Influenza and pneumonia (J09-J18)	1,882	2.4	7.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,593	2.0	6.4
12	Certain conditions originating in the perinatal period (P00-P96)	1,462	1.9	5.8
13	Septicemia (A40-A41)	1,041	1.3	4.2
14	Human immunodeficiency virus (HIV) disease (B20-B24)	985	1.3	3.9
15	Alzheimer's disease (G30)	973	1.2	3.9
...	All other causes (Residual)	14,044	18.0	56.0

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	539	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	208	38.6	9.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	10.8	2.6
3	Malignant neoplasms (C00-C97)	44	8.2	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	5.9	1.4
5	Influenza and pneumonia (J09-J18)	16	3.0	*
6	Septicemia (A40-A41)	12	2.2	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	2.0	*
8	Chronic lower respiratory diseases (J40-J47)	6	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.9	*
9	Cerebrovascular diseases (I60-I69)	5	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
12	Meningococcal infection (A39)	1	0.2	*
12	Anemias (D50-D64)	1	0.2	*
12	Meningitis (G00,G03)	1	0.2	*
12	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
12	Hernia (K40-K46)	1	0.2	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (Residual)	131	24.3	5.9
Hispanic, male, 5-14 years				
...	All causes	650	100.0	14.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	29.2	4.2
2	Malignant neoplasms (C00-C97)	118	18.2	2.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	46	7.1	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	6.0	0.9
5	Influenza and pneumonia (J09-J18)	33	5.1	0.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	4.9	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.1	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	2.2	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.2	*
10	Cerebrovascular diseases (I60-I69)	6	0.9	*
11	Diabetes mellitus (E10-E14)	4	0.6	*
12	Septicemia (A40-A41)	3	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
14	Anemias (D50-D64)	2	0.3	*
14	Meningitis (G00,G03)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	122	18.8	2.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,992	100.0	96.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,590	39.8	38.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,044	26.2	25.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	488	12.2	11.7
4	Malignant neoplasms (C00-C97)	204	5.1	4.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	2.5	2.4
6	Influenza and pneumonia (J09-J18)	40	1.0	1.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.0	1.0
8	Cerebrovascular diseases (I60-I69)	25	0.6	0.6
9	Septicemia (A40-A41)	21	0.5	0.5
10	Human immunodeficiency virus (HIV) disease (B20-B24)	17	0.4	*
11	Legal intervention (Y35,Y89.0)	15	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	10	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.2	*
13	Diabetes mellitus (E10-E14)	9	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.2	*
...	All other causes (Residual)	371	9.3	8.9
Hispanic, male, 25-44 years				
...	All causes	10,499	100.0	126.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,070	29.2	37.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,144	10.9	13.8
3	Malignant neoplasms (C00-C97)	996	9.5	12.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	983	9.4	11.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	917	8.7	11.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	490	4.7	5.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	385	3.7	4.6
8	Influenza and pneumonia (J09-J18)	282	2.7	3.4
9	Cerebrovascular diseases (I60-I69)	237	2.3	2.9
10	Diabetes mellitus (E10-E14)	186	1.8	2.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	103	1.0	1.2
12	Septicemia (A40-A41)	95	0.9	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	0.6	0.8
14	Viral hepatitis (B15-B19)	65	0.6	0.8
15	Chronic lower respiratory diseases (J40-J47)	53	0.5	0.6
...	All other causes (Residual)	1,427	13.6	17.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	21,896	100.0	533.9
1	Malignant neoplasms (C00-C97)	5,118	23.4	124.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,725	21.6	115.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,847	8.4	45.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,777	8.1	43.3
5	Diabetes mellitus (E10-E14)	1,090	5.0	26.6
6	Cerebrovascular diseases (I60-I69)	879	4.0	21.4
7	Viral hepatitis (B15-B19)	542	2.5	13.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	516	2.4	12.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	516	2.4	12.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	423	1.9	10.3
11	Influenza and pneumonia (J09-J18)	405	1.8	9.9
12	Septicemia (A40-A41)	329	1.5	8.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	281	1.3	6.9
14	Chronic lower respiratory diseases (J40-J47)	260	1.2	6.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	0.8	4.3
...	All other causes (Residual)	3,011	13.8	73.4
Hispanic, male, 65 years and over				
...	All causes	37,573	100.0	3,170.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,417	27.7	878.9
2	Malignant neoplasms (C00-C97)	9,284	24.7	783.3
3	Cerebrovascular diseases (I60-I69)	2,118	5.6	178.7
4	Diabetes mellitus (E10-E14)	1,923	5.1	162.2
5	Chronic lower respiratory diseases (J40-J47)	1,677	4.5	141.5
6	Influenza and pneumonia (J09-J18)	1,069	2.8	90.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,044	2.8	88.1
8	Alzheimer's disease (G30)	947	2.5	79.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	817	2.2	68.9
10	Chronic liver disease and cirrhosis (K70,K73-K74)	697	1.9	58.8
11	Septicemia (A40-A41)	553	1.5	46.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	511	1.4	43.1
13	Parkinson's disease (G20-G21)	469	1.2	39.6
14	Pneumonitis due to solids and liquids (J69)	270	0.7	22.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	233	0.6	19.7
...	All other causes (Residual)	5,544	14.8	467.7

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	63,419	100.0	271.5
1	Malignant neoplasms (C00-C97)	14,163	22.3	60.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,310	21.0	57.0
3	Cerebrovascular diseases (I60-I69)	3,777	6.0	16.2
4	Diabetes mellitus (E10-E14)	3,099	4.9	13.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,813	4.4	12.0
6	Alzheimer's disease (G30)	2,013	3.2	8.6
7	Chronic lower respiratory diseases (J40-J47)	2,002	3.2	8.6
8	Influenza and pneumonia (J09-J18)	1,797	2.8	7.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,514	2.4	6.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,334	2.1	5.7
11	Certain conditions originating in the perinatal period (P00-P96)	1,156	1.8	4.9
12	Septicemia (A40-A41)	1,036	1.6	4.4
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	882	1.4	3.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	791	1.2	3.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	544	0.9	2.3
...	All other causes (Residual)	13,188	20.8	56.4
Hispanic, female, 1-4 years				
...	All causes	465	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	24.9	5.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	14.2	3.1
3	Malignant neoplasms (C00-C97)	55	11.8	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	43	9.2	2.0
5	Influenza and pneumonia (J09-J18)	18	3.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	7	1.5	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
8	Cerebrovascular diseases (I60-I69)	6	1.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	6	1.3	*
11	Septicemia (A40-A41)	5	1.1	*
12	Meningitis (G00,G03)	3	0.6	*
12	Acute bronchitis and bronchiolitis (J20-J21)	3	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Anemias (D50-D64)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	111	23.9	5.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	522	100.0	12.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	27.2	3.3
2	Malignant neoplasms (C00-C97)	98	18.8	2.3
3	Influenza and pneumonia (J09-J18)	30	5.7	0.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	5.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	4.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	4.2	0.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16	3.1	*
8	Septicemia (A40-A41)	12	2.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.9	*
10	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
11	Anemias (D50-D64)	4	0.8	*
11	Cerebrovascular diseases (I60-I69)	4	0.8	*
13	Diabetes mellitus (E10-E14)	3	0.6	*
13	Meningitis (G00,G03)	3	0.6	*
13	Diseases of appendix (K35-K38)	3	0.6	*
...	All other causes (Residual)	115	22.0	2.7
Hispanic, female, 15-24 years				
...	All causes	1,195	100.0	31.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	34.2	10.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	124	10.4	3.3
3	Malignant neoplasms (C00-C97)	116	9.7	3.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	95	7.9	2.5
5	Influenza and pneumonia (J09-J18)	48	4.0	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	3.9	1.2
7	Pregnancy, childbirth and the puerperium (O00-O99)	41	3.4	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	2.6	0.8
9	Cerebrovascular diseases (I60-I69)	17	1.4	*
10	Septicemia (A40-A41)	15	1.3	*
10	Diabetes mellitus (E10-E14)	15	1.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	13	1.1	*
13	Chronic lower respiratory diseases (J40-J47)	11	0.9	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
...	All other causes (Residual)	198	16.6	5.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	4,510	100.0	64.8
1	Malignant neoplasms (C00-C97)	1,145	25.4	16.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	740	16.4	10.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	324	7.2	4.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	217	4.8	3.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	4.1	2.7
6	Cerebrovascular diseases (I60-I69)	165	3.7	2.4
7	Influenza and pneumonia (J09-J18)	159	3.5	2.3
8	Pregnancy, childbirth and the puerperium (O00-O99)	128	2.8	1.8
9	Diabetes mellitus (E10-E14)	119	2.6	1.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	116	2.6	1.7
11	Human immunodeficiency virus (HIV) disease (B20-B24)	98	2.2	1.4
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	61	1.4	0.9
13	Septicemia (A40-A41)	59	1.3	0.8
14	Chronic lower respiratory diseases (J40-J47)	54	1.2	0.8
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	1.0	0.7
...	All other causes (Residual)	891	19.8	12.8
Hispanic, female, 45-64 years				
...	All causes	12,581	100.0	309.8
1	Malignant neoplasms (C00-C97)	4,586	36.5	112.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,811	14.4	44.6
3	Diabetes mellitus (E10-E14)	690	5.5	17.0
4	Cerebrovascular diseases (I60-I69)	607	4.8	14.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	597	4.7	14.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	583	4.6	14.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	285	2.3	7.0
8	Influenza and pneumonia (J09-J18)	278	2.2	6.8
9	Septicemia (A40-A41)	243	1.9	6.0
10	Chronic lower respiratory diseases (J40-J47)	229	1.8	5.6
11	Viral hepatitis (B15-B19)	228	1.8	5.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	143	1.1	3.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	122	1.0	3.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	104	0.8	2.6
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	0.7	2.1
...	All other causes (Residual)	1,991	15.8	49.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	41,701	100.0	2,646.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,038	26.5	700.5
2	Malignant neoplasms (C00-C97)	8,154	19.6	517.5
3	Cerebrovascular diseases (I60-I69)	2,963	7.1	188.0
4	Diabetes mellitus (E10-E14)	2,272	5.4	144.2
5	Alzheimer's disease (G30)	1,992	4.8	126.4
6	Chronic lower respiratory diseases (J40-J47)	1,693	4.1	107.4
7	Influenza and pneumonia (J09-J18)	1,247	3.0	79.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,146	2.7	72.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	732	1.8	46.5
10	Septicemia (A40-A41)	680	1.6	43.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	652	1.6	41.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	631	1.5	40.0
13	Parkinson's disease (G20-G21)	354	0.8	22.5
14	Pneumonitis due to solids and liquids (J69)	257	0.6	16.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	237	0.6	15.0
...	All other causes (Residual)	7,653	18.4	485.7
Non-Hispanic, white, both sexes, all ages				
...	All causes	1,944,606	100.0	961.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	485,779	25.0	240.3
2	Malignant neoplasms (C00-C97)	457,189	23.5	226.2
3	Chronic lower respiratory diseases (J40-J47)	122,605	6.3	60.6
4	Cerebrovascular diseases (I60-I69)	101,703	5.2	50.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91,416	4.7	45.2
6	Alzheimer's disease (G30)	69,695	3.6	34.5
7	Diabetes mellitus (E10-E14)	47,851	2.5	23.7
8	Influenza and pneumonia (J09-J18)	42,752	2.2	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,670	1.8	17.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30,813	1.6	15.2
11	Septicemia (A40-A41)	26,533	1.4	13.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	22,520	1.2	11.1
13	Parkinson's disease (G20-G21)	18,554	1.0	9.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,552	1.0	9.2
15	Pneumonitis due to solids and liquids (J69)	13,489	0.7	6.7
...	All other causes (Residual)	359,485	18.5	177.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,181	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	773	35.4	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	230	10.5	2.5
3	Malignant neoplasms (C00-C97)	180	8.3	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	6.5	1.5
5	Influenza and pneumonia (J09-J18)	75	3.4	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	70	3.2	0.8
7	Septicemia (A40-A41)	30	1.4	0.3
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.3	0.3
9	Cerebrovascular diseases (I60-I69)	27	1.2	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	27	1.2	0.3
11	Chronic lower respiratory diseases (J40-J47)	19	0.9	*
12	Meningitis (G00,G03)	11	0.5	*
13	Anemias (D50-D64)	10	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.4	*
15	Meningococcal infection (A39)	7	0.3	*
...	All other causes (Residual)	545	25.0	6.0
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,867	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	895	31.2	3.8
2	Malignant neoplasms (C00-C97)	493	17.2	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	192	6.7	0.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	157	5.5	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	117	4.1	0.5
6	Influenza and pneumonia (J09-J18)	104	3.6	0.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	91	3.2	0.4
8	Cerebrovascular diseases (I60-I69)	43	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	39	1.4	0.2
10	Septicemia (A40-A41)	28	1.0	0.1
11	Chronic lower respiratory diseases (J40-J47)	26	0.9	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.6	*
13	Diabetes mellitus (E10-E14)	14	0.5	*
14	Anemias (D50-D64)	10	0.3	*
14	Meningitis (G00,G03)	10	0.3	*
...	All other causes (Residual)	632	22.0	2.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	17,150	100.0	65.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,564	49.9	32.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,046	17.8	11.6
3	Malignant neoplasms (C00-C97)	945	5.5	3.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	755	4.4	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	499	2.9	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	270	1.6	1.0
7	Influenza and pneumonia (J09-J18)	223	1.3	0.9
8	Cerebrovascular diseases (I60-I69)	97	0.6	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	93	0.5	0.4
10	Diabetes mellitus (E10-E14)	89	0.5	0.3
11	Septicemia (A40-A41)	73	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	71	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.2	0.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	0.2	0.1
15	Pneumonitis due to solids and liquids (J69)	33	0.2	0.1
...	All other causes (Residual)	2,321	13.5	8.9
Non-Hispanic, white, both sexes, 25-44 years				
...	All causes	72,939	100.0	142.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	20,889	28.6	40.8
2	Malignant neoplasms (C00-C97)	10,419	14.3	20.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,491	13.0	18.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,535	11.7	16.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,873	2.6	3.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,860	2.6	3.6
7	Diabetes mellitus (E10-E14)	1,348	1.8	2.6
8	Cerebrovascular diseases (I60-I69)	1,207	1.7	2.4
9	Influenza and pneumonia (J09-J18)	1,151	1.6	2.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	779	1.1	1.5
11	Septicemia (A40-A41)	711	1.0	1.4
12	Chronic lower respiratory diseases (J40-J47)	605	0.8	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	599	0.8	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	472	0.6	0.9
15	Viral hepatitis (B15-B19)	271	0.4	0.5
...	All other causes (Residual)	12,729	17.5	24.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-64 years				
...	All causes	355,115	100.0	609.9
1	Malignant neoplasms (C00-C97)	118,296	33.3	203.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	73,565	20.7	126.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25,372	7.1	43.6
4	Chronic lower respiratory diseases (J40-J47)	15,957	4.5	27.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	12,792	3.6	22.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12,785	3.6	22.0
7	Diabetes mellitus (E10-E14)	10,843	3.1	18.6
8	Cerebrovascular diseases (I60-I69)	9,994	2.8	17.2
9	Influenza and pneumonia (J09-J18)	4,920	1.4	8.5
10	Septicemia (A40-A41)	4,595	1.3	7.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,922	1.1	6.7
12	Viral hepatitis (B15-B19)	3,828	1.1	6.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,094	0.6	3.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,651	0.5	2.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,601	0.5	2.7
...	All other causes (Residual)	52,900	14.9	90.9
Non-Hispanic, white, both sexes, 65 years and over				
...	All causes	1,482,669	100.0	4,655.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	402,840	27.2	1,264.8
2	Malignant neoplasms (C00-C97)	326,809	22.0	1,026.1
3	Chronic lower respiratory diseases (J40-J47)	105,915	7.1	332.5
4	Cerebrovascular diseases (I60-I69)	90,267	6.1	283.4
5	Alzheimer's disease (G30)	68,993	4.7	216.6
6	Influenza and pneumonia (J09-J18)	36,182	2.4	113.6
7	Diabetes mellitus (E10-E14)	35,555	2.4	111.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,350	2.3	107.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31,174	2.1	97.9
10	Septicemia (A40-A41)	21,018	1.4	66.0
11	Parkinson's disease (G20-G21)	18,104	1.2	56.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16,234	1.1	51.0
13	Pneumonitis due to solids and liquids (J69)	12,149	0.8	38.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10,162	0.7	31.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	7,858	0.5	24.7
...	All other causes (Residual)	265,059	17.9	832.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	959,014	100.0	965.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	247,489	25.8	249.2
2	Malignant neoplasms (C00-C97)	239,301	25.0	241.0
3	Chronic lower respiratory diseases (J40-J47)	57,370	6.0	57.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	56,607	5.9	57.0
5	Cerebrovascular diseases (I60-I69)	39,912	4.2	40.2
6	Diabetes mellitus (E10-E14)	24,993	2.6	25.2
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	24,248	2.5	24.4
8	Alzheimer's disease (G30)	21,087	2.2	21.2
9	Influenza and pneumonia (J09-J18)	19,736	2.1	19.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,735	1.8	17.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	14,494	1.5	14.6
12	Septicemia (A40-A41)	12,167	1.3	12.3
13	Parkinson's disease (G20-G21)	10,919	1.1	11.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,316	0.8	7.4
15	Pneumonitis due to solids and liquids (J69)	7,068	0.7	7.1
...	All other causes (Residual)	158,572	16.5	159.7
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,262	100.0	27.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	500	39.6	10.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	8.8	2.4
3	Malignant neoplasms (C00-C97)	106	8.4	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	5.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.2	0.9
6	Influenza and pneumonia (J09-J18)	32	2.5	0.7
7	Certain conditions originating in the perinatal period (P00-P96)	18	1.4	*
8	Septicemia (A40-A41)	17	1.3	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.3	*
10	Cerebrovascular diseases (I60-I69)	15	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
12	Anemias (D50-D64)	8	0.6	*
12	Meningitis (G00,G03)	8	0.6	*
14	Meningococcal infection (A39)	6	0.5	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
...	All other causes (Residual)	298	23.6	6.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,684	100.0	14.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	574	34.1	4.8
2	Malignant neoplasms (C00-C97)	277	16.4	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	6.5	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	6.0	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	4.0	0.6
6	Influenza and pneumonia (J09-J18)	51	3.0	0.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	2.8	0.4
8	Cerebrovascular diseases (I60-I69)	26	1.5	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.3	0.2
10	Chronic lower respiratory diseases (J40-J47)	18	1.1	*
11	Septicemia (A40-A41)	11	0.7	*
12	Meningitis (G00,G03)	7	0.4	*
13	Diabetes mellitus (E10-E14)	6	0.4	*
14	Anemias (D50-D64)	5	0.3	*
14	Diseases of appendix (K35-K38)	5	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
...	All other causes (Residual)	351	20.8	2.9
Non-Hispanic, white, male, 15-24 years				
...	All causes	12,318	100.0	91.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,301	51.2	46.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,533	20.6	18.9
3	Malignant neoplasms (C00-C97)	578	4.7	4.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	536	4.4	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	330	2.7	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	155	1.3	1.2
7	Influenza and pneumonia (J09-J18)	129	1.0	1.0
8	Cerebrovascular diseases (I60-I69)	54	0.4	0.4
9	Diabetes mellitus (E10-E14)	53	0.4	0.4
10	Septicemia (A40-A41)	45	0.4	0.3
11	Chronic lower respiratory diseases (J40-J47)	41	0.3	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	0.2	0.2
12	Legal intervention (Y35,Y89.0)	25	0.2	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	21	0.2	0.2
15	Complications of medical and surgical care (Y40-Y84,Y88)	21	0.2	0.2
...	All other causes (Residual)	1,449	11.8	10.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-44 years				
...	All causes	46,999	100.0	182.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,875	31.6	57.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,427	15.8	28.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,040	12.9	23.4
4	Malignant neoplasms (C00-C97)	4,710	10.0	18.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,242	2.6	4.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,175	2.5	4.6
7	Diabetes mellitus (E10-E14)	860	1.8	3.3
8	Cerebrovascular diseases (I60-I69)	652	1.4	2.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	623	1.3	2.4
10	Influenza and pneumonia (J09-J18)	617	1.3	2.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	357	0.8	1.4
12	Septicemia (A40-A41)	343	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	266	0.6	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	264	0.6	1.0
15	Aortic aneurysm and dissection (I71)	187	0.4	0.7
...	All other causes (Residual)	7,361	15.7	28.6
Non-Hispanic, white, male, 45-64 years				
...	All causes	217,702	100.0	759.0
1	Malignant neoplasms (C00-C97)	63,723	29.3	222.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,020	24.4	184.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,056	7.8	59.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,666	4.4	33.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	8,874	4.1	30.9
6	Chronic lower respiratory diseases (J40-J47)	8,112	3.7	28.3
7	Diabetes mellitus (E10-E14)	6,822	3.1	23.8
8	Cerebrovascular diseases (I60-I69)	5,643	2.6	19.7
9	Viral hepatitis (B15-B19)	2,765	1.3	9.6
10	Influenza and pneumonia (J09-J18)	2,691	1.2	9.4
11	Septicemia (A40-A41)	2,446	1.1	8.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,283	1.0	8.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,374	0.6	4.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,358	0.6	4.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,051	0.5	3.7
...	All other causes (Residual)	30,818	14.2	107.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65 years and over				
...	All causes	672,473	100.0	4,926.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	187,908	27.9	1,376.7
2	Malignant neoplasms (C00-C97)	169,880	25.3	1,244.6
3	Chronic lower respiratory diseases (J40-J47)	48,913	7.3	358.4
4	Cerebrovascular diseases (I60-I69)	33,484	5.0	245.3
5	Alzheimer's disease (G30)	20,778	3.1	152.2
6	Diabetes mellitus (E10-E14)	17,250	2.6	126.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,961	2.5	124.3
8	Influenza and pneumonia (J09-J18)	16,157	2.4	118.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,127	2.2	110.8
10	Parkinson's disease (G20-G21)	10,628	1.6	77.9
11	Septicemia (A40-A41)	9,262	1.4	67.9
12	Pneumonitis due to solids and liquids (J69)	6,246	0.9	45.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,789	0.9	42.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,217	0.8	38.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,517	0.7	33.1
...	All other causes (Residual)	104,356	15.5	764.5
Non-Hispanic, white, female, all ages				
...	All causes	985,592	100.0	958.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	238,290	24.2	231.7
2	Malignant neoplasms (C00-C97)	217,888	22.1	211.9
3	Chronic lower respiratory diseases (J40-J47)	65,235	6.6	63.4
4	Cerebrovascular diseases (I60-I69)	61,791	6.3	60.1
5	Alzheimer's disease (G30)	48,608	4.9	47.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,809	3.5	33.8
7	Influenza and pneumonia (J09-J18)	23,016	2.3	22.4
8	Diabetes mellitus (E10-E14)	22,858	2.3	22.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,935	1.8	17.4
10	Septicemia (A40-A41)	14,366	1.5	14.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,236	1.1	10.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,026	0.8	7.8
13	Parkinson's disease (G20-G21)	7,635	0.8	7.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,565	0.7	6.4
15	Pneumonitis due to solids and liquids (J69)	6,421	0.7	6.2
...	All other causes (Residual)	200,913	20.4	195.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	919	100.0	20.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	273	29.7	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	119	12.9	2.7
3	Malignant neoplasms (C00-C97)	74	8.1	1.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	7.9	1.6
5	Influenza and pneumonia (J09-J18)	43	4.7	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.2	0.7
7	Septicemia (A40-A41)	13	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.3	*
8	Cerebrovascular diseases (I60-I69)	12	1.3	*
10	Certain conditions originating in the perinatal period (P00-P96)	9	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
13	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Meningitis (G00,G03)	3	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
...	All other causes (Residual)	240	26.1	5.4
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,183	100.0	10.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	27.1	2.8
2	Malignant neoplasms (C00-C97)	216	18.3	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	6.9	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	4.7	0.5
5	Influenza and pneumonia (J09-J18)	53	4.5	0.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	4.1	0.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	3.7	0.4
8	Septicemia (A40-A41)	17	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.4	*
8	Cerebrovascular diseases (I60-I69)	17	1.4	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	0.9	*
12	Diabetes mellitus (E10-E14)	8	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	8	0.7	*
14	Anemias (D50-D64)	5	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (Residual)	274	23.2	2.4

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,832	100.0	37.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,263	46.8	17.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	513	10.6	4.0
3	Malignant neoplasms (C00-C97)	367	7.6	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	219	4.5	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	169	3.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	115	2.4	0.9
7	Influenza and pneumonia (J09-J18)	94	1.9	0.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	93	1.9	0.7
9	Cerebrovascular diseases (I60-I69)	43	0.9	0.3
10	Diabetes mellitus (E10-E14)	36	0.7	0.3
11	Chronic lower respiratory diseases (J40-J47)	30	0.6	0.2
12	Septicemia (A40-A41)	28	0.6	0.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
14	Pneumonitis due to solids and liquids (J69)	12	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.2	*
15	Meningitis (G00,G03)	11	0.2	*
...	All other causes (Residual)	815	16.9	6.4
Non-Hispanic, white, female, 25-44 years				
...	All causes	25,940	100.0	101.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,014	23.2	23.6
2	Malignant neoplasms (C00-C97)	5,709	22.0	22.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,495	9.6	9.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,064	8.0	8.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	685	2.6	2.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	631	2.4	2.5
7	Cerebrovascular diseases (I60-I69)	555	2.1	2.2
8	Influenza and pneumonia (J09-J18)	534	2.1	2.1
9	Diabetes mellitus (E10-E14)	488	1.9	1.9
10	Septicemia (A40-A41)	368	1.4	1.4
11	Chronic lower respiratory diseases (J40-J47)	339	1.3	1.3
12	Pregnancy, childbirth and the puerperium (O00-O99)	270	1.0	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	242	0.9	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	0.8	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	156	0.6	0.6
...	All other causes (Residual)	5,182	20.0	20.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-64 years				
...	All causes	137,413	100.0	465.1
1	Malignant neoplasms (C00-C97)	54,573	39.7	184.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,545	15.0	69.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,316	6.1	28.1
4	Chronic lower respiratory diseases (J40-J47)	7,845	5.7	26.6
5	Cerebrovascular diseases (I60-I69)	4,351	3.2	14.7
6	Diabetes mellitus (E10-E14)	4,021	2.9	13.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,918	2.9	13.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,119	2.3	10.6
9	Influenza and pneumonia (J09-J18)	2,229	1.6	7.5
10	Septicemia (A40-A41)	2,149	1.6	7.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,639	1.2	5.5
12	Viral hepatitis (B15-B19)	1,063	0.8	3.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	720	0.5	2.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	701	0.5	2.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	532	0.4	1.8
...	All other causes (Residual)	21,692	15.8	73.4
Non-Hispanic, white, female, 65 years and over				
...	All causes	810,196	100.0	4,451.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	214,932	26.5	1,180.9
2	Malignant neoplasms (C00-C97)	156,929	19.4	862.2
3	Chronic lower respiratory diseases (J40-J47)	57,002	7.0	313.2
4	Cerebrovascular diseases (I60-I69)	56,783	7.0	312.0
5	Alzheimer's disease (G30)	48,215	6.0	264.9
6	Influenza and pneumonia (J09-J18)	20,025	2.5	110.0
7	Diabetes mellitus (E10-E14)	18,305	2.3	100.6
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17,389	2.1	95.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,047	2.0	88.2
10	Septicemia (A40-A41)	11,756	1.5	64.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,445	1.3	57.4
12	Parkinson's disease (G20-G21)	7,476	0.9	41.1
13	Pneumonitis due to solids and liquids (J69)	5,903	0.7	32.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4,945	0.6	27.2
15	Atherosclerosis (I70)	3,722	0.5	20.4
...	All other causes (Residual)	160,322	19.8	880.8

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	282,982	100.0	728.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	68,811	24.3	177.1
2	Malignant neoplasms (C00-C97)	63,967	22.6	164.6
3	Cerebrovascular diseases (I60-I69)	15,718	5.6	40.4
4	Diabetes mellitus (E10-E14)	11,833	4.2	30.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,810	4.2	30.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,727	3.1	22.5
7	Chronic lower respiratory diseases (J40-J47)	8,444	3.0	21.7
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,733	2.7	19.9
9	Septicemia (A40-A41)	6,131	2.2	15.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	5,307	1.9	13.7
11	Influenza and pneumonia (J09-J18)	5,214	1.8	13.4
12	Alzheimer's disease (G30)	5,042	1.8	13.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,819	1.7	12.4
14	Certain conditions originating in the perinatal period (P00-P96)	4,456	1.6	11.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,518	0.9	6.5
...	All other causes (Residual)	52,452	18.5	135.0
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,039	100.0	41.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	288	27.7	11.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	13.8	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	87	8.4	3.5
4	Malignant neoplasms (C00-C97)	56	5.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	50	4.8	2.0
6	Chronic lower respiratory diseases (J40-J47)	33	3.2	1.3
7	Influenza and pneumonia (J09-J18)	29	2.8	1.2
8	Septicemia (A40-A41)	19	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	19	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	12	1.2	*
12	Cerebrovascular diseases (I60-I69)	9	0.9	*
13	Anemias (D50-D64)	8	0.8	*
13	Acute bronchitis and bronchiolitis (J20-J21)	8	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.6	*
...	All other causes (Residual)	260	25.0	10.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,266	100.0	21.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	377	29.8	6.3
2	Malignant neoplasms (C00-C97)	128	10.1	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	105	8.3	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	79	6.2	1.3
5	Chronic lower respiratory diseases (J40-J47)	70	5.5	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	5.1	1.1
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	3.7	0.8
8	Influenza and pneumonia (J09-J18)	45	3.6	0.7
9	Septicemia (A40-A41)	22	1.7	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.6	0.3
11	Anemias (D50-D64)	19	1.5	*
12	Cerebrovascular diseases (I60-I69)	18	1.4	*
13	Diabetes mellitus (E10-E14)	8	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
15	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (Residual)	248	19.6	4.1
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	6,602	100.0	99.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,760	41.8	41.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,309	19.8	19.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	433	6.6	6.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	340	5.1	5.1
5	Malignant neoplasms (C00-C97)	284	4.3	4.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	106	1.6	1.6
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	1.4	1.4
8	Pregnancy, childbirth and the puerperium (O00-O99)	88	1.3	1.3
9	Influenza and pneumonia (J09-J18)	85	1.3	1.3
9	Chronic lower respiratory diseases (J40-J47)	85	1.3	1.3
11	Anemias (D50-D64)	66	1.0	1.0
12	Diabetes mellitus (E10-E14)	50	0.8	0.8
13	Cerebrovascular diseases (I60-I69)	39	0.6	0.6
14	Septicemia (A40-A41)	31	0.5	0.5
15	Legal intervention (Y35,Y89.0)	30	0.5	0.5
...	All other causes (Residual)	805	12.2	12.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-44 years				
...	All causes	23,982	100.0	219.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,812	15.9	35.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,466	14.5	31.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,302	13.8	30.3
4	Malignant neoplasms (C00-C97)	2,759	11.5	25.3
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,964	8.2	18.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	861	3.6	7.9
7	Diabetes mellitus (E10-E14)	713	3.0	6.5
8	Cerebrovascular diseases (I60-I69)	686	2.9	6.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	410	1.7	3.8
10	Influenza and pneumonia (J09-J18)	395	1.6	3.6
11	Septicemia (A40-A41)	378	1.6	3.5
12	Chronic lower respiratory diseases (J40-J47)	279	1.2	2.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	263	1.1	2.4
14	Anemias (D50-D64)	242	1.0	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	224	0.9	2.1
...	All other causes (Residual)	4,228	17.6	38.8
Non-Hispanic, black, both sexes, 45-64 years				
...	All causes	85,151	100.0	962.8
1	Malignant neoplasms (C00-C97)	24,072	28.3	272.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,864	24.5	235.9
3	Cerebrovascular diseases (I60-I69)	4,359	5.1	49.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,006	4.7	45.3
5	Diabetes mellitus (E10-E14)	3,756	4.4	42.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,878	3.4	32.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,271	2.7	25.7
8	Chronic lower respiratory diseases (J40-J47)	2,065	2.4	23.3
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,727	2.0	19.5
10	Septicemia (A40-A41)	1,706	2.0	19.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,378	1.6	15.6
12	Influenza and pneumonia (J09-J18)	1,159	1.4	13.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	983	1.2	11.1
14	Viral hepatitis (B15-B19)	848	1.0	9.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	541	0.6	6.1
...	All other causes (Residual)	12,538	14.7	141.8

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65 years and over				
...	All causes	156,952	100.0	4,703.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,563	27.8	1,305.5
2	Malignant neoplasms (C00-C97)	36,653	23.4	1,098.4
3	Cerebrovascular diseases (I60-I69)	10,563	6.7	316.5
4	Diabetes mellitus (E10-E14)	7,305	4.7	218.9
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,975	3.8	179.1
6	Chronic lower respiratory diseases (J40-J47)	5,903	3.8	176.9
7	Alzheimer's disease (G30)	4,962	3.2	148.7
8	Septicemia (A40-A41)	3,887	2.5	116.5
9	Influenza and pneumonia (J09-J18)	3,420	2.2	102.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,169	2.0	95.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,137	1.4	64.0
12	Pneumonitis due to solids and liquids (J69)	1,110	0.7	33.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	759	0.5	22.7
14	Parkinson's disease (G20-G21)	687	0.4	20.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	562	0.4	16.8
...	All other causes (Residual)	26,297	16.8	788.1
Non-Hispanic, black, male, all ages				
...	All causes	144,197	100.0	777.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,935	24.2	188.3
2	Malignant neoplasms (C00-C97)	33,065	22.9	178.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,894	5.5	42.5
4	Cerebrovascular diseases (I60-I69)	6,892	4.8	37.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,605	4.6	35.6
6	Diabetes mellitus (E10-E14)	5,430	3.8	29.3
7	Chronic lower respiratory diseases (J40-J47)	4,457	3.1	24.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,930	2.7	21.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	3,476	2.4	18.7
10	Septicemia (A40-A41)	2,721	1.9	14.7
11	Certain conditions originating in the perinatal period (P00-P96)	2,533	1.8	13.7
12	Influenza and pneumonia (J09-J18)	2,463	1.7	13.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,045	1.4	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,649	1.1	8.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,636	1.1	8.8
...	All other causes (Residual)	24,466	17.0	131.9

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	561	100.0	44.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	30.1	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	14.1	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	8.6	3.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.9	1.7
5	Malignant neoplasms (C00-C97)	20	3.6	1.6
6	Chronic lower respiratory diseases (J40-J47)	18	3.2	*
7	Influenza and pneumonia (J09-J18)	13	2.3	*
8	Certain conditions originating in the perinatal period (P00-P96)	10	1.8	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
11	Septicemia (A40-A41)	7	1.2	*
12	Anemias (D50-D64)	6	1.1	*
13	Cerebrovascular diseases (I60-I69)	4	0.7	*
14	Acute bronchitis and bronchiolitis (J20-J21)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
...	All other causes (Residual)	141	25.1	11.1
Non-Hispanic, black, male, 5-14 years				
...	All causes	735	100.0	24.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	32.4	7.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	9.9	2.4
3	Malignant neoplasms (C00-C97)	60	8.2	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	6.4	1.5
5	Chronic lower respiratory diseases (J40-J47)	43	5.9	1.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	4.8	1.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	35	4.8	1.1
8	Influenza and pneumonia (J09-J18)	22	3.0	0.7
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.8	*
10	Cerebrovascular diseases (I60-I69)	11	1.5	*
11	Septicemia (A40-A41)	10	1.4	*
12	Anemias (D50-D64)	7	1.0	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
14	Diseases of appendix (K35-K38)	3	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	128	17.4	4.2

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	4,985	100.0	148.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,483	49.8	74.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	937	18.8	27.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	360	7.2	10.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	224	4.5	6.7
5	Malignant neoplasms (C00-C97)	158	3.2	4.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	65	1.3	1.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.1	1.7
8	Chronic lower respiratory diseases (J40-J47)	53	1.1	1.6
9	Influenza and pneumonia (J09-J18)	40	0.8	1.2
10	Anemias (D50-D64)	36	0.7	1.1
11	Legal intervention (Y35,Y89.0)	29	0.6	0.9
12	Diabetes mellitus (E10-E14)	25	0.5	0.7
13	Cerebrovascular diseases (I60-I69)	21	0.4	0.6
14	Septicemia (A40-A41)	12	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	0.2	*
...	All other causes (Residual)	474	9.5	14.1
Non-Hispanic, black, male, 25-44 years				
...	All causes	14,938	100.0	285.3
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,008	20.1	57.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,425	16.2	46.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,412	16.1	46.1
4	Human immunodeficiency virus (HIV) disease (B20-B24)	1,186	7.9	22.6
5	Malignant neoplasms (C00-C97)	1,100	7.4	21.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	705	4.7	13.5
7	Diabetes mellitus (E10-E14)	436	2.9	8.3
8	Cerebrovascular diseases (I60-I69)	367	2.5	7.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	215	1.4	4.1
10	Septicemia (A40-A41)	180	1.2	3.4
11	Influenza and pneumonia (J09-J18)	179	1.2	3.4
12	Chronic lower respiratory diseases (J40-J47)	158	1.1	3.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	142	1.0	2.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	134	0.9	2.6
15	Anemias (D50-D64)	122	0.8	2.3
...	All other causes (Residual)	2,169	14.5	41.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-64 years				
...	All causes	49,593	100.0	1,232.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,115	26.4	326.0
2	Malignant neoplasms (C00-C97)	12,855	25.9	319.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,746	5.5	68.2
4	Cerebrovascular diseases (I60-I69)	2,378	4.8	59.1
5	Diabetes mellitus (E10-E14)	2,065	4.2	51.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,969	4.0	48.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,215	2.4	30.2
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,167	2.4	29.0
9	Chronic lower respiratory diseases (J40-J47)	1,082	2.2	26.9
10	Septicemia (A40-A41)	885	1.8	22.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	789	1.6	19.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	783	1.6	19.5
13	Influenza and pneumonia (J09-J18)	628	1.3	15.6
14	Viral hepatitis (B15-B19)	584	1.2	14.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	417	0.8	10.4
...	All other causes (Residual)	6,915	13.9	171.9
Non-Hispanic, black, male, 65 years and over				
...	All causes	68,847	100.0	5,347.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,052	27.7	1,479.9
2	Malignant neoplasms (C00-C97)	18,863	27.4	1,465.2
3	Cerebrovascular diseases (I60-I69)	4,083	5.9	317.2
4	Chronic lower respiratory diseases (J40-J47)	3,098	4.5	240.6
5	Diabetes mellitus (E10-E14)	2,901	4.2	225.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,463	3.6	191.3
7	Septicemia (A40-A41)	1,580	2.3	122.7
8	Influenza and pneumonia (J09-J18)	1,535	2.2	119.2
9	Alzheimer's disease (G30)	1,385	2.0	107.6
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,152	1.7	89.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,111	1.6	86.3
12	Pneumonitis due to solids and liquids (J69)	507	0.7	39.4
13	Parkinson's disease (G20-G21)	390	0.6	30.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	330	0.5	25.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	316	0.5	24.5
...	All other causes (Residual)	10,081	14.6	783.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	138,785	100.0	683.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,876	24.4	166.8
2	Malignant neoplasms (C00-C97)	30,902	22.3	152.2
3	Cerebrovascular diseases (I60-I69)	8,826	6.4	43.5
4	Diabetes mellitus (E10-E14)	6,403	4.6	31.5
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,797	3.5	23.6
6	Chronic lower respiratory diseases (J40-J47)	3,987	2.9	19.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,916	2.8	19.3
8	Alzheimer's disease (G30)	3,612	2.6	17.8
9	Septicemia (A40-A41)	3,410	2.5	16.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,774	2.0	13.7
11	Influenza and pneumonia (J09-J18)	2,751	2.0	13.5
12	Certain conditions originating in the perinatal period (P00-P96)	1,923	1.4	9.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,831	1.3	9.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,128	0.8	5.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	882	0.6	4.3
...	All other causes (Residual)	27,767	20.0	136.7
Non-Hispanic, black, female, 1-4 years				
...	All causes	478	100.0	38.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	24.9	9.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	13.4	5.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	8.2	3.2
4	Malignant neoplasms (C00-C97)	36	7.5	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.9	2.3
6	Influenza and pneumonia (J09-J18)	16	3.3	*
7	Chronic lower respiratory diseases (J40-J47)	15	3.1	*
8	Septicemia (A40-A41)	12	2.5	*
9	Certain conditions originating in the perinatal period (P00-P96)	9	1.9	*
10	Cerebrovascular diseases (I60-I69)	5	1.0	*
10	Acute bronchitis and bronchiolitis (J20-J21)	5	1.0	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Anemias (D50-D64)	2	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
...	All other causes (Residual)	119	24.9	9.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	531	100.0	17.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	139	26.2	4.7
2	Malignant neoplasms (C00-C97)	68	12.8	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	6.0	1.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	6.0	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	5.5	1.0
6	Chronic lower respiratory diseases (J40-J47)	27	5.1	0.9
7	Influenza and pneumonia (J09-J18)	23	4.3	0.8
8	Septicemia (A40-A41)	12	2.3	*
8	Anemias (D50-D64)	12	2.3	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	2.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.3	*
11	Cerebrovascular diseases (I60-I69)	7	1.3	*
13	Diabetes mellitus (E10-E14)	6	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.6	*
...	All other causes (Residual)	119	22.4	4.0
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,617	100.0	49.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	372	23.0	11.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	277	17.1	8.5
3	Malignant neoplasms (C00-C97)	126	7.8	3.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	116	7.2	3.6
5	Pregnancy, childbirth and the puerperium (O00-O99)	88	5.4	2.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	73	4.5	2.2
7	Influenza and pneumonia (J09-J18)	45	2.8	1.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	41	2.5	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	2.2	1.1
10	Chronic lower respiratory diseases (J40-J47)	32	2.0	1.0
11	Anemias (D50-D64)	30	1.9	0.9
12	Diabetes mellitus (E10-E14)	25	1.5	0.8
13	Septicemia (A40-A41)	19	1.2	*
14	Cerebrovascular diseases (I60-I69)	18	1.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	1.1	*
...	All other causes (Residual)	303	18.7	9.3

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-44 years				
...	All causes	9,044	100.0	159.6
1	Malignant neoplasms (C00-C97)	1,659	18.3	29.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,400	15.5	24.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	877	9.7	15.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	778	8.6	13.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	458	5.1	8.1
6	Cerebrovascular diseases (I60-I69)	319	3.5	5.6
7	Diabetes mellitus (E10-E14)	277	3.1	4.9
8	Influenza and pneumonia (J09-J18)	216	2.4	3.8
9	Septicemia (A40-A41)	198	2.2	3.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	195	2.2	3.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	194	2.1	3.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	156	1.7	2.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.3	2.1
13	Chronic lower respiratory diseases (J40-J47)	121	1.3	2.1
15	Anemias (D50-D64)	120	1.3	2.1
...	All other causes (Residual)	1,955	21.6	34.5
Non-Hispanic, black, female, 45-64 years				
...	All causes	35,558	100.0	737.6
1	Malignant neoplasms (C00-C97)	11,217	31.5	232.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,749	21.8	160.7
3	Cerebrovascular diseases (I60-I69)	1,981	5.6	41.1
4	Diabetes mellitus (E10-E14)	1,691	4.8	35.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,260	3.5	26.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,056	3.0	21.9
7	Chronic lower respiratory diseases (J40-J47)	983	2.8	20.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	909	2.6	18.9
9	Septicemia (A40-A41)	821	2.3	17.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	589	1.7	12.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	560	1.6	11.6
12	Influenza and pneumonia (J09-J18)	531	1.5	11.0
13	Viral hepatitis (B15-B19)	264	0.7	5.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	200	0.6	4.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	173	0.5	3.6
...	All other causes (Residual)	5,574	15.7	115.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2009

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

Rank ¹	Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65 years and over				
...	All causes	88,105	100.0	4,298.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,511	27.8	1,195.9
2	Malignant neoplasms (C00-C97)	17,790	20.2	868.0
3	Cerebrovascular diseases (I60-I69)	6,480	7.4	316.2
4	Diabetes mellitus (E10-E14)	4,404	5.0	214.9
5	Alzheimer's disease (G30)	3,577	4.1	174.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,512	4.0	171.4
7	Chronic lower respiratory diseases (J40-J47)	2,805	3.2	136.9
8	Septicemia (A40-A41)	2,307	2.6	112.6
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,058	2.3	100.4
10	Influenza and pneumonia (J09-J18)	1,885	2.1	92.0
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	985	1.1	48.1
12	Pneumonitis due to solids and liquids (J69)	603	0.7	29.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	443	0.5	21.6
14	Atherosclerosis (I70)	305	0.3	14.9
15	Parkinson's disease (G20-G21)	297	0.3	14.5
...	All other causes (Residual)	16,143	18.3	787.6

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable