

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	149,635	100.0	287.5
1	Malignant neoplasms (C00-C97)	32,381	21.6	62.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,385	20.3	58.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,166	7.5	21.5
4	Cerebrovascular diseases (I60-I69)	7,552	5.0	14.5
5	Diabetes mellitus (E10-E14)	7,054	4.7	13.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,774	3.2	9.2
7	Chronic lower respiratory diseases (J40-J47)	4,355	2.9	8.4
8	Alzheimer's disease (G30)	3,599	2.4	6.9
9	Influenza and pneumonia (J09-J18)	3,256	2.2	6.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,799	1.9	5.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,759	1.8	5.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,720	1.8	5.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,387	1.6	4.6
14	Septicemia (A40-A41)	2,168	1.4	4.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,803	1.2	3.5
...	All other causes (Residual)	30,477	20.4	58.6
Hispanic, both sexes, 1-4 years				
...	All causes	972	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	317	32.6	7.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	13.0	3.0
3	Malignant neoplasms (C00-C97)	97	10.0	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	94	9.7	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.6	0.8
6	Influenza and pneumonia (J09-J18)	21	2.2	0.5
7	Septicemia (A40-A41)	16	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
8	Cerebrovascular diseases (I60-I69)	10	1.0	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	217	22.3	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,052	100.0	11.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	278	26.4	2.9
2	Malignant neoplasms (C00-C97)	200	19.0	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	8.7	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	5.5	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	5.1	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	4.3	0.5
7	Chronic lower respiratory diseases (J40-J47)	23	2.2	0.2
8	Cerebrovascular diseases (I60-I69)	22	2.1	0.2
8	Influenza and pneumonia (J09-J18)	22	2.1	0.2
10	Septicemia (A40-A41)	10	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
12	Anemias (D50-D64)	6	0.6	*
12	Diabetes mellitus (E10-E14)	6	0.6	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.5	*
15	Meningitis (G00,G03)	4	0.4	*
...	All other causes (Residual)	220	20.9	2.3
Hispanic, both sexes, 15-24 years				
...	All causes	4,826	100.0	53.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,911	39.6	21.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	933	19.3	10.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	635	13.2	7.0
4	Malignant neoplasms (C00-C97)	383	7.9	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	3.2	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	1.2	0.6
7	Cerebrovascular diseases (I60-I69)	53	1.1	0.6
8	Legal intervention (Y35,Y89.0)	34	0.7	0.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	31	0.6	0.3
10	Influenza and pneumonia (J09-J18)	28	0.6	0.3
11	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.4	0.2
12	Diabetes mellitus (E10-E14)	19	0.4	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.4	*
14	Chronic lower respiratory diseases (J40-J47)	17	0.4	*
15	Septicemia (A40-A41)	16	0.3	*
...	All other causes (Residual)	515	10.7	5.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	14,253	100.0	88.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,903	27.4	24.2
2	Malignant neoplasms (C00-C97)	2,104	14.8	13.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,331	9.3	8.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,226	8.6	7.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,177	8.3	7.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	575	4.0	3.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	390	2.7	2.4
8	Cerebrovascular diseases (I60-I69)	365	2.6	2.3
9	Diabetes mellitus (E10-E14)	305	2.1	1.9
10	Influenza and pneumonia (J09-J18)	187	1.3	1.2
11	Septicemia (A40-A41)	145	1.0	0.9
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	0.7	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	94	0.7	0.6
14	Pregnancy, childbirth and the puerperium (O00-O99)	87	0.6	0.5
15	Viral hepatitis (B15-B19)	77	0.5	0.5
...	All other causes (Residual)	2,185	15.3	13.6
Hispanic, both sexes, 45-64 years				
...	All causes	36,507	100.0	397.9
1	Malignant neoplasms (C00-C97)	10,497	28.8	114.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,670	18.3	72.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,778	7.6	30.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,713	7.4	29.6
5	Diabetes mellitus (E10-E14)	1,938	5.3	21.1
6	Cerebrovascular diseases (I60-I69)	1,630	4.5	17.8
7	Viral hepatitis (B15-B19)	841	2.3	9.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	649	1.8	7.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	643	1.8	7.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	611	1.7	6.7
11	Septicemia (A40-A41)	580	1.6	6.3
12	Chronic lower respiratory diseases (J40-J47)	542	1.5	5.9
13	Influenza and pneumonia (J09-J18)	476	1.3	5.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	0.9	3.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	307	0.8	3.3
...	All other causes (Residual)	5,293	14.5	57.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	87,193	100.0	2,936.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,083	25.3	743.8
2	Malignant neoplasms (C00-C97)	19,085	21.9	642.8
3	Cerebrovascular diseases (I60-I69)	5,451	6.3	183.6
4	Diabetes mellitus (E10-E14)	4,784	5.5	161.1
5	Chronic lower respiratory diseases (J40-J47)	3,691	4.2	124.3
6	Alzheimer's disease (G30)	3,524	4.0	118.7
7	Influenza and pneumonia (J09-J18)	2,492	2.9	83.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,028	2.3	68.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,817	2.1	61.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,473	1.7	49.6
11	Septicemia (A40-A41)	1,368	1.6	46.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,334	1.5	44.9
13	Parkinson's disease (G20-G21)	1,025	1.2	34.5
14	Pneumonitis due to solids and liquids (J69)	640	0.7	21.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	529	0.6	17.8
...	All other causes (Residual)	15,869	18.2	534.5
Hispanic, male, all ages				
...	All causes	81,887	100.0	309.7
1	Malignant neoplasms (C00-C97)	17,185	21.0	65.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,526	20.2	62.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,098	9.9	30.6
4	Diabetes mellitus (E10-E14)	3,726	4.6	14.1
5	Cerebrovascular diseases (I60-I69)	3,528	4.3	13.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,311	4.0	12.5
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,279	2.8	8.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,224	2.7	8.4
9	Chronic lower respiratory diseases (J40-J47)	2,222	2.7	8.4
10	Influenza and pneumonia (J09-J18)	1,606	2.0	6.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,388	1.7	5.2
12	Certain conditions originating in the perinatal period (P00-P96)	1,297	1.6	4.9
13	Alzheimer's disease (G30)	1,151	1.4	4.4
14	Septicemia (A40-A41)	1,048	1.3	4.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	958	1.2	3.6
...	All other causes (Residual)	15,340	18.7	58.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	539	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	187	34.7	8.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	13.0	3.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	10.2	2.6
4	Malignant neoplasms (C00-C97)	51	9.5	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.7	0.9
6	Influenza and pneumonia (J09-J18)	10	1.9	*
7	Septicemia (A40-A41)	8	1.5	*
8	Cerebrovascular diseases (I60-I69)	7	1.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
10	Meningitis (G00,G03)	3	0.6	*
10	Acute bronchitis and bronchiolitis (J20-J21)	3	0.6	*
10	Certain conditions originating in the perinatal period (P00-P96)	3	0.6	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
...	All other causes (Residual)	110	20.4	5.2
Hispanic, male, 5-14 years				
...	All causes	608	100.0	12.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	182	29.9	3.7
2	Malignant neoplasms (C00-C97)	108	17.8	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	7.9	1.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	6.3	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	5.9	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	2.8	*
7	Cerebrovascular diseases (I60-I69)	12	2.0	*
7	Chronic lower respiratory diseases (J40-J47)	12	2.0	*
9	Influenza and pneumonia (J09-J18)	9	1.5	*
10	Septicemia (A40-A41)	5	0.8	*
10	Anemias (D50-D64)	5	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
13	Meningitis (G00,G03)	3	0.5	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Diseases of appendix (K35-K38)	2	0.3	*
...	All other causes (Residual)	125	20.6	2.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,723	100.0	78.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,517	40.7	32.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	835	22.4	17.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	510	13.7	10.7
4	Malignant neoplasms (C00-C97)	250	6.7	5.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	2.6	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	1.0	0.8
7	Legal intervention (Y35,Y89.0)	33	0.9	0.7
8	Cerebrovascular diseases (I60-I69)	32	0.9	0.7
9	Influenza and pneumonia (J09-J18)	17	0.5	*
10	Chronic lower respiratory diseases (J40-J47)	13	0.3	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	11	0.3	*
12	Anemias (D50-D64)	9	0.2	*
12	Diabetes mellitus (E10-E14)	9	0.2	*
14	Septicemia (A40-A41)	8	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.2	*
...	All other causes (Residual)	336	9.0	7.1
Hispanic, male, 25-44 years				
...	All causes	9,945	100.0	118.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,077	30.9	36.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,009	10.1	12.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	984	9.9	11.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	979	9.8	11.7
5	Malignant neoplasms (C00-C97)	935	9.4	11.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	452	4.5	5.4
7	Human immunodeficiency virus (HIV) disease (B20-B24)	307	3.1	3.7
8	Cerebrovascular diseases (I60-I69)	227	2.3	2.7
9	Diabetes mellitus (E10-E14)	191	1.9	2.3
10	Influenza and pneumonia (J09-J18)	116	1.2	1.4
11	Septicemia (A40-A41)	81	0.8	1.0
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	0.7	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	60	0.6	0.7
14	Viral hepatitis (B15-B19)	59	0.6	0.7
15	Legal intervention (Y35,Y89.0)	49	0.5	0.6
...	All other causes (Residual)	1,354	13.6	16.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	23,383	100.0	516.0
1	Malignant neoplasms (C00-C97)	5,707	24.4	125.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,753	20.3	104.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,086	8.9	46.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,048	8.8	45.2
5	Diabetes mellitus (E10-E14)	1,201	5.1	26.5
6	Cerebrovascular diseases (I60-I69)	982	4.2	21.7
7	Viral hepatitis (B15-B19)	629	2.7	13.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	524	2.2	11.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	456	2.0	10.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	388	1.7	8.6
11	Septicemia (A40-A41)	332	1.4	7.3
12	Influenza and pneumonia (J09-J18)	295	1.3	6.5
13	Chronic lower respiratory diseases (J40-J47)	294	1.3	6.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	269	1.2	5.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	193	0.8	4.3
...	All other causes (Residual)	3,226	13.8	71.2
Hispanic, male, 65 years and over				
...	All causes	41,063	100.0	3,241.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,616	25.9	838.1
2	Malignant neoplasms (C00-C97)	10,126	24.7	799.4
3	Diabetes mellitus (E10-E14)	2,322	5.7	183.3
4	Cerebrovascular diseases (I60-I69)	2,260	5.5	178.4
5	Chronic lower respiratory diseases (J40-J47)	1,866	4.5	147.3
6	Influenza and pneumonia (J09-J18)	1,138	2.8	89.8
7	Alzheimer's disease (G30)	1,121	2.7	88.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	957	2.3	75.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	922	2.2	72.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	801	2.0	63.2
11	Septicemia (A40-A41)	599	1.5	47.3
12	Parkinson's disease (G20-G21)	594	1.4	46.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	544	1.3	42.9
14	Pneumonitis due to solids and liquids (J69)	332	0.8	26.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	274	0.7	21.6
...	All other causes (Residual)	6,591	16.1	520.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	67,748	100.0	264.6
1	Malignant neoplasms (C00-C97)	15,196	22.4	59.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,859	20.5	54.1
3	Cerebrovascular diseases (I60-I69)	4,024	5.9	15.7
4	Diabetes mellitus (E10-E14)	3,328	4.9	13.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,068	4.5	12.0
6	Alzheimer's disease (G30)	2,448	3.6	9.6
7	Chronic lower respiratory diseases (J40-J47)	2,133	3.1	8.3
8	Influenza and pneumonia (J09-J18)	1,650	2.4	6.4
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,463	2.2	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,411	2.1	5.5
11	Septicemia (A40-A41)	1,120	1.7	4.4
12	Certain conditions originating in the perinatal period (P00-P96)	1,090	1.6	4.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	927	1.4	3.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	845	1.2	3.3
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	496	0.7	1.9
...	All other causes (Residual)	14,690	21.7	57.4
Hispanic, female, 1-4 years				
...	All causes	433	100.0	21.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	30.0	6.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	56	12.9	2.8
3	Malignant neoplasms (C00-C97)	46	10.6	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	39	9.0	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.5	*
6	Influenza and pneumonia (J09-J18)	11	2.5	*
7	Septicemia (A40-A41)	8	1.8	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.6	*
9	Cerebrovascular diseases (I60-I69)	3	0.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
12	Diseases of appendix (K35-K38)	2	0.5	*
12	Hernia (K40-K46)	2	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.5	*
...	All other causes (Residual)	104	24.0	5.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	444	100.0	9.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	96	21.6	2.1
2	Malignant neoplasms (C00-C97)	92	20.7	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	9.9	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	6.3	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	4.5	0.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	4.1	*
7	Influenza and pneumonia (J09-J18)	13	2.9	*
8	Chronic lower respiratory diseases (J40-J47)	11	2.5	*
9	Cerebrovascular diseases (I60-I69)	10	2.3	*
10	Septicemia (A40-A41)	5	1.1	*
11	Diabetes mellitus (E10-E14)	4	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
14	Meningococcal infection (A39)	2	0.5	*
14	Pneumonitis due to solids and liquids (J69)	2	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
...	All other causes (Residual)	90	20.3	1.9
Hispanic, female, 15-24 years				
...	All causes	1,103	100.0	25.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	394	35.7	9.1
2	Malignant neoplasms (C00-C97)	133	12.1	3.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	11.3	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	8.9	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	5.3	1.3
6	Pregnancy, childbirth and the puerperium (O00-O99)	31	2.8	0.7
7	Cerebrovascular diseases (I60-I69)	21	1.9	0.5
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	1.7	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.2	*
10	Influenza and pneumonia (J09-J18)	11	1.0	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.9	*
11	Diabetes mellitus (E10-E14)	10	0.9	*
13	Septicemia (A40-A41)	8	0.7	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.5	*
15	Anemias (D50-D64)	4	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	4	0.4	*
...	All other causes (Residual)	158	14.3	3.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	4,308	100.0	55.7
1	Malignant neoplasms (C00-C97)	1,169	27.1	15.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	826	19.2	10.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	347	8.1	4.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	217	5.0	2.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	198	4.6	2.6
6	Cerebrovascular diseases (I60-I69)	138	3.2	1.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	123	2.9	1.6
8	Diabetes mellitus (E10-E14)	114	2.6	1.5
9	Pregnancy, childbirth and the puerperium (O00-O99)	87	2.0	1.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	83	1.9	1.1
11	Influenza and pneumonia (J09-J18)	71	1.6	0.9
12	Septicemia (A40-A41)	64	1.5	0.8
13	Chronic lower respiratory diseases (J40-J47)	39	0.9	0.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	0.9	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	34	0.8	0.4
...	All other causes (Residual)	761	17.7	9.8
Hispanic, female, 45-64 years				
...	All causes	13,124	100.0	282.7
1	Malignant neoplasms (C00-C97)	4,790	36.5	103.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,917	14.6	41.3
3	Diabetes mellitus (E10-E14)	737	5.6	15.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	692	5.3	14.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	665	5.1	14.3
6	Cerebrovascular diseases (I60-I69)	648	4.9	14.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	255	1.9	5.5
8	Septicemia (A40-A41)	248	1.9	5.3
8	Chronic lower respiratory diseases (J40-J47)	248	1.9	5.3
10	Viral hepatitis (B15-B19)	212	1.6	4.6
11	Influenza and pneumonia (J09-J18)	181	1.4	3.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	155	1.2	3.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	125	1.0	2.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	0.9	2.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	79	0.6	1.7
...	All other causes (Residual)	2,058	15.7	44.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	46,130	100.0	2,709.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,467	24.9	673.6
2	Malignant neoplasms (C00-C97)	8,959	19.4	526.3
3	Cerebrovascular diseases (I60-I69)	3,191	6.9	187.4
4	Diabetes mellitus (E10-E14)	2,462	5.3	144.6
5	Alzheimer's disease (G30)	2,403	5.2	141.2
6	Chronic lower respiratory diseases (J40-J47)	1,825	4.0	107.2
7	Influenza and pneumonia (J09-J18)	1,354	2.9	79.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,106	2.4	65.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	860	1.9	50.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	790	1.7	46.4
11	Septicemia (A40-A41)	769	1.7	45.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	672	1.5	39.5
13	Parkinson's disease (G20-G21)	431	0.9	25.3
14	Pneumonitis due to solids and liquids (J69)	308	0.7	18.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	255	0.6	15.0
...	All other causes (Residual)	9,278	20.1	545.0
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,006,319	100.0	1,001.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	482,979	24.1	241.0
2	Malignant neoplasms (C00-C97)	460,026	22.9	229.5
3	Chronic lower respiratory diseases (J40-J47)	127,035	6.3	63.4
4	Cerebrovascular diseases (I60-I69)	101,043	5.0	50.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	98,605	4.9	49.2
6	Alzheimer's disease (G30)	74,448	3.7	37.1
7	Diabetes mellitus (E10-E14)	50,964	2.5	25.4
8	Influenza and pneumonia (J09-J18)	43,350	2.2	21.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,492	1.7	16.7
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,031	1.6	16.5
11	Septicemia (A40-A41)	26,669	1.3	13.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	24,752	1.2	12.3
13	Parkinson's disease (G20-G21)	20,625	1.0	10.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20,153	1.0	10.1
15	Pneumonitis due to solids and liquids (J69)	15,513	0.8	7.7
...	All other causes (Residual)	393,634	19.6	196.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,055	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	724	35.2	8.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	236	11.5	2.8
3	Malignant neoplasms (C00-C97)	183	8.9	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	7.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	74	3.6	0.9
6	Influenza and pneumonia (J09-J18)	45	2.2	0.5
7	Septicemia (A40-A41)	30	1.5	0.4
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	24	1.2	0.3
9	Chronic lower respiratory diseases (J40-J47)	19	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	17	0.8	*
11	Cerebrovascular diseases (I60-I69)	14	0.7	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
14	Anemias (D50-D64)	6	0.3	*
15	Meningitis (G00,G03)	5	0.2	*
...	All other causes (Residual)	513	25.0	6.0
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,908	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	973	33.5	4.3
2	Malignant neoplasms (C00-C97)	481	16.5	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	168	5.8	0.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	5.8	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	4.3	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	97	3.3	0.4
7	Influenza and pneumonia (J09-J18)	61	2.1	0.3
8	Septicemia (A40-A41)	46	1.6	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	1.5	0.2
10	Cerebrovascular diseases (I60-I69)	38	1.3	0.2
11	Chronic lower respiratory diseases (J40-J47)	29	1.0	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.6	*
13	Diabetes mellitus (E10-E14)	15	0.5	*
14	Pneumonitis due to solids and liquids (J69)	10	0.3	*
15	Anemias (D50-D64)	9	0.3	*
...	All other causes (Residual)	629	21.6	2.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	16,961	100.0	67.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,528	50.3	33.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,413	20.1	13.5
3	Malignant neoplasms (C00-C97)	891	5.3	3.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	701	4.1	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	493	2.9	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	252	1.5	1.0
7	Influenza and pneumonia (J09-J18)	138	0.8	0.5
8	Diabetes mellitus (E10-E14)	97	0.6	0.4
9	Cerebrovascular diseases (I60-I69)	90	0.5	0.4
10	Septicemia (A40-A41)	77	0.5	0.3
11	Pregnancy, childbirth and the puerperium (O00-O99)	73	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	64	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.3	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.2	0.1
15	Pneumonitis due to solids and liquids (J69)	28	0.2	0.1
15	Complications of medical and surgical care (Y40-Y84,Y88)	28	0.2	0.1
...	All other causes (Residual)	2,004	11.8	7.9
Non-Hispanic, white, both sexes, 25-44 years				
...	All causes	71,645	100.0	145.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,384	31.2	45.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9,945	13.9	20.1
3	Malignant neoplasms (C00-C97)	9,520	13.3	19.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,395	11.7	17.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,880	2.6	3.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,783	2.5	3.6
7	Diabetes mellitus (E10-E14)	1,393	1.9	2.8
8	Cerebrovascular diseases (I60-I69)	1,077	1.5	2.2
9	Influenza and pneumonia (J09-J18)	831	1.2	1.7
10	Septicemia (A40-A41)	637	0.9	1.3
11	Chronic lower respiratory diseases (J40-J47)	618	0.9	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	577	0.8	1.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	539	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	384	0.5	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	299	0.4	0.6
...	All other causes (Residual)	11,383	15.9	23.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-64 years				
...	All causes	366,320	100.0	617.6
1	Malignant neoplasms (C00-C97)	119,916	32.7	202.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	75,098	20.5	126.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,568	7.5	46.5
4	Chronic lower respiratory diseases (J40-J47)	16,602	4.5	28.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	14,256	3.9	24.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13,713	3.7	23.1
7	Diabetes mellitus (E10-E14)	11,797	3.2	19.9
8	Cerebrovascular diseases (I60-I69)	10,061	2.7	17.0
9	Septicemia (A40-A41)	4,843	1.3	8.2
10	Influenza and pneumonia (J09-J18)	4,583	1.3	7.7
11	Viral hepatitis (B15-B19)	3,886	1.1	6.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,744	1.0	6.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,271	0.6	3.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,615	0.4	2.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,595	0.4	2.7
...	All other causes (Residual)	54,772	15.0	92.3
Non-Hispanic, white, both sexes, 65 years and over				
...	All causes	1,535,530	100.0	4,642.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	398,682	26.0	1,205.4
2	Malignant neoplasms (C00-C97)	328,995	21.4	994.7
3	Chronic lower respiratory diseases (J40-J47)	109,687	7.1	331.6
4	Cerebrovascular diseases (I60-I69)	89,696	5.8	271.2
5	Alzheimer's disease (G30)	73,674	4.8	222.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37,840	2.5	114.4
7	Diabetes mellitus (E10-E14)	37,657	2.5	113.9
8	Influenza and pneumonia (J09-J18)	37,601	2.4	113.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29,291	1.9	88.6
10	Septicemia (A40-A41)	20,969	1.4	63.4
11	Parkinson's disease (G20-G21)	20,161	1.3	61.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,682	1.2	53.5
13	Pneumonitis due to solids and liquids (J69)	14,019	0.9	42.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10,488	0.7	31.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	8,599	0.6	26.0
...	All other causes (Residual)	300,489	19.6	908.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	989,835	100.0	1,004.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	248,787	25.1	252.4
2	Malignant neoplasms (C00-C97)	242,140	24.5	245.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60,043	6.1	60.9
4	Chronic lower respiratory diseases (J40-J47)	59,223	6.0	60.1
5	Cerebrovascular diseases (I60-I69)	39,753	4.0	40.3
6	Diabetes mellitus (E10-E14)	27,060	2.7	27.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25,858	2.6	26.2
8	Alzheimer's disease (G30)	22,482	2.3	22.8
9	Influenza and pneumonia (J09-J18)	20,316	2.1	20.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,898	1.7	17.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	15,764	1.6	16.0
12	Septicemia (A40-A41)	12,319	1.2	12.5
13	Parkinson's disease (G20-G21)	12,134	1.2	12.3
14	Pneumonitis due to solids and liquids (J69)	8,242	0.8	8.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7,926	0.8	8.0
...	All other causes (Residual)	170,890	17.3	173.4
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,177	100.0	27.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	454	38.6	10.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	124	10.5	2.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	93	7.9	2.1
4	Malignant neoplasms (C00-C97)	89	7.6	2.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	35	3.0	0.8
6	Influenza and pneumonia (J09-J18)	22	1.9	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	15	1.3	*
9	Septicemia (A40-A41)	12	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	9	0.8	*
11	Cerebrovascular diseases (I60-I69)	8	0.7	*
12	Anemias (D50-D64)	4	0.3	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
15	Meningitis (G00,G03)	3	0.3	*
...	All other causes (Residual)	285	24.2	6.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,727	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	613	35.5	5.2
2	Malignant neoplasms (C00-C97)	274	15.9	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	121	7.0	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	94	5.4	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	4.2	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	48	2.8	0.4
7	Influenza and pneumonia (J09-J18)	35	2.0	0.3
8	Septicemia (A40-A41)	24	1.4	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.3	0.2
10	Cerebrovascular diseases (I60-I69)	21	1.2	0.2
11	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
12	Pneumonitis due to solids and liquids (J69)	8	0.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.5	*
14	Anemias (D50-D64)	6	0.3	*
15	Diabetes mellitus (E10-E14)	5	0.3	*
...	All other causes (Residual)	358	20.7	3.1
Non-Hispanic, white, male, 15-24 years				
...	All causes	12,222	100.0	94.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,249	51.1	48.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,817	23.0	21.9
3	Malignant neoplasms (C00-C97)	517	4.2	4.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	497	4.1	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	356	2.9	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	164	1.3	1.3
7	Influenza and pneumonia (J09-J18)	81	0.7	0.6
8	Diabetes mellitus (E10-E14)	67	0.5	0.5
9	Cerebrovascular diseases (I60-I69)	51	0.4	0.4
10	Septicemia (A40-A41)	48	0.4	0.4
11	Chronic lower respiratory diseases (J40-J47)	47	0.4	0.4
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.2	0.2
13	Pneumonitis due to solids and liquids (J69)	20	0.2	0.2
14	Legal intervention (Y35,Y89.0)	18	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.1	*
...	All other causes (Residual)	1,246	10.2	9.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-44 years				
...	All causes	46,254	100.0	186.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,614	33.8	62.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,709	16.7	31.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,905	12.8	23.8
4	Malignant neoplasms (C00-C97)	4,439	9.6	17.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,223	2.6	4.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,152	2.5	4.6
7	Diabetes mellitus (E10-E14)	855	1.8	3.4
8	Cerebrovascular diseases (I60-I69)	550	1.2	2.2
9	Influenza and pneumonia (J09-J18)	445	1.0	1.8
10	Human immunodeficiency virus (HIV) disease (B20-B24)	402	0.9	1.6
11	Septicemia (A40-A41)	326	0.7	1.3
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	326	0.7	1.3
13	Chronic lower respiratory diseases (J40-J47)	300	0.6	1.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	231	0.5	0.9
15	Aortic aneurysm and dissection (I71)	170	0.4	0.7
...	All other causes (Residual)	6,607	14.3	26.6
Non-Hispanic, white, male, 45-64 years				
...	All causes	224,488	100.0	768.2
1	Malignant neoplasms (C00-C97)	65,056	29.0	222.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,812	24.0	184.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,400	8.2	63.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,362	4.6	35.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	9,690	4.3	33.2
6	Chronic lower respiratory diseases (J40-J47)	8,450	3.8	28.9
7	Diabetes mellitus (E10-E14)	7,360	3.3	25.2
8	Cerebrovascular diseases (I60-I69)	5,651	2.5	19.3
9	Viral hepatitis (B15-B19)	2,830	1.3	9.7
10	Influenza and pneumonia (J09-J18)	2,658	1.2	9.1
11	Septicemia (A40-A41)	2,563	1.1	8.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,168	1.0	7.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,477	0.7	5.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,217	0.5	4.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,074	0.5	3.7
...	All other causes (Residual)	31,720	14.1	108.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65 years and over				
...	All causes	697,841	100.0	4,814.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	188,556	27.0	1,300.9
2	Malignant neoplasms (C00-C97)	171,742	24.6	1,184.9
3	Chronic lower respiratory diseases (J40-J47)	50,383	7.2	347.6
4	Cerebrovascular diseases (I60-I69)	33,430	4.8	230.6
5	Alzheimer's disease (G30)	22,139	3.2	152.7
6	Diabetes mellitus (E10-E14)	18,771	2.7	129.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,381	2.6	126.8
8	Influenza and pneumonia (J09-J18)	17,020	2.4	117.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,460	2.1	99.8
10	Parkinson's disease (G20-G21)	11,834	1.7	81.6
11	Septicemia (A40-A41)	9,313	1.3	64.3
12	Pneumonitis due to solids and liquids (J69)	7,342	1.1	50.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,316	0.9	43.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,457	0.8	37.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,911	0.7	33.9
...	All other causes (Residual)	117,786	16.9	812.7
Non-Hispanic, white, female, all ages				
...	All causes	1,016,484	100.0	998.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	234,192	23.0	230.0
2	Malignant neoplasms (C00-C97)	217,886	21.4	213.9
3	Chronic lower respiratory diseases (J40-J47)	67,812	6.7	66.6
4	Cerebrovascular diseases (I60-I69)	61,290	6.0	60.2
5	Alzheimer's disease (G30)	51,966	5.1	51.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38,562	3.8	37.9
7	Diabetes mellitus (E10-E14)	23,904	2.4	23.5
8	Influenza and pneumonia (J09-J18)	23,034	2.3	22.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,594	1.6	16.3
10	Septicemia (A40-A41)	14,350	1.4	14.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,227	1.2	12.0
12	Chronic liver disease and cirrhosis (K70,K73-K74)	8,988	0.9	8.8
13	Parkinson's disease (G20-G21)	8,491	0.8	8.3
14	Pneumonitis due to solids and liquids (J69)	7,271	0.7	7.1
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,173	0.7	7.0
...	All other causes (Residual)	222,744	21.9	218.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	878	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	270	30.8	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	12.8	2.7
3	Malignant neoplasms (C00-C97)	94	10.7	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	6.5	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	4.4	0.9
6	Influenza and pneumonia (J09-J18)	23	2.6	0.6
7	Septicemia (A40-A41)	18	2.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.9	*
8	Certain conditions originating in the perinatal period (P00-P96)	8	0.9	*
10	Cerebrovascular diseases (I60-I69)	6	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
11	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
13	Diabetes mellitus (E10-E14)	3	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	223	25.4	5.4
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,181	100.0	10.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	360	30.5	3.2
2	Malignant neoplasms (C00-C97)	207	17.5	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	6.3	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	4.5	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.1	0.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47	4.0	0.4
7	Influenza and pneumonia (J09-J18)	26	2.2	0.2
8	Septicemia (A40-A41)	22	1.9	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.7	0.2
10	Cerebrovascular diseases (I60-I69)	17	1.4	*
11	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
12	Diabetes mellitus (E10-E14)	10	0.8	*
13	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
14	Meningitis (G00,G03)	4	0.3	*
15	Anemias (D50-D64)	3	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (Residual)	263	22.3	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,739	100.0	38.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,279	48.1	18.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	596	12.6	4.8
3	Malignant neoplasms (C00-C97)	374	7.9	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	204	4.3	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	137	2.9	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	1.9	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	73	1.5	0.6
8	Influenza and pneumonia (J09-J18)	57	1.2	0.5
9	Cerebrovascular diseases (I60-I69)	39	0.8	0.3
10	Diabetes mellitus (E10-E14)	30	0.6	0.2
11	Septicemia (A40-A41)	29	0.6	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.6	0.2
13	Chronic lower respiratory diseases (J40-J47)	17	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
...	All other causes (Residual)	759	16.0	6.1
Non-Hispanic, white, female, 25-44 years				
...	All causes	25,391	100.0	103.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,770	26.7	27.6
2	Malignant neoplasms (C00-C97)	5,081	20.0	20.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,490	9.8	10.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,236	8.8	9.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	728	2.9	3.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	560	2.2	2.3
7	Diabetes mellitus (E10-E14)	538	2.1	2.2
8	Cerebrovascular diseases (I60-I69)	527	2.1	2.1
9	Influenza and pneumonia (J09-J18)	386	1.5	1.6
10	Chronic lower respiratory diseases (J40-J47)	318	1.3	1.3
11	Septicemia (A40-A41)	311	1.2	1.3
12	Pregnancy, childbirth and the puerperium (O00-O99)	299	1.2	1.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	251	1.0	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	0.6	0.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	137	0.5	0.6
...	All other causes (Residual)	4,606	18.1	18.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-64 years				
...	All causes	141,832	100.0	471.3
1	Malignant neoplasms (C00-C97)	54,860	38.7	182.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,286	15.0	70.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,168	6.5	30.5
4	Chronic lower respiratory diseases (J40-J47)	8,152	5.7	27.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,566	3.2	15.2
6	Diabetes mellitus (E10-E14)	4,437	3.1	14.7
7	Cerebrovascular diseases (I60-I69)	4,410	3.1	14.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,351	2.4	11.1
9	Septicemia (A40-A41)	2,280	1.6	7.6
10	Influenza and pneumonia (J09-J18)	1,925	1.4	6.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,576	1.1	5.2
12	Viral hepatitis (B15-B19)	1,056	0.7	3.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	794	0.6	2.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	702	0.5	2.3
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	608	0.4	2.0
...	All other causes (Residual)	22,661	16.0	75.3
Non-Hispanic, white, female, 65 years and over				
...	All causes	837,689	100.0	4,508.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	210,126	25.1	1,130.9
2	Malignant neoplasms (C00-C97)	157,253	18.8	846.3
3	Chronic lower respiratory diseases (J40-J47)	59,304	7.1	319.2
4	Cerebrovascular diseases (I60-I69)	56,266	6.7	302.8
5	Alzheimer's disease (G30)	51,535	6.2	277.4
6	Influenza and pneumonia (J09-J18)	20,581	2.5	110.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,459	2.3	104.7
8	Diabetes mellitus (E10-E14)	18,886	2.3	101.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,831	1.8	79.8
10	Septicemia (A40-A41)	11,656	1.4	62.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,366	1.4	61.2
12	Parkinson's disease (G20-G21)	8,327	1.0	44.8
13	Pneumonitis due to solids and liquids (J69)	6,677	0.8	35.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,031	0.6	27.1
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	4,006	0.5	21.6
...	All other causes (Residual)	182,385	21.8	981.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	286,797	100.0	718.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	67,595	23.6	169.2
2	Malignant neoplasms (C00-C97)	66,158	23.1	165.6
3	Cerebrovascular diseases (I60-I69)	15,722	5.5	39.4
4	Diabetes mellitus (E10-E14)	12,771	4.5	32.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,299	4.3	30.8
6	Chronic lower respiratory diseases (J40-J47)	8,994	3.1	22.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,044	2.8	20.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,764	2.7	19.4
9	Septicemia (A40-A41)	5,963	2.1	14.9
10	Alzheimer's disease (G30)	5,446	1.9	13.6
11	Influenza and pneumonia (J09-J18)	5,098	1.8	12.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,965	1.7	12.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	4,138	1.4	10.4
14	Certain conditions originating in the perinatal period (P00-P96)	3,890	1.4	9.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,783	1.0	7.0
...	All other causes (Residual)	55,167	19.2	138.1
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,011	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	286	28.3	11.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	154	15.2	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	10.4	4.2
4	Malignant neoplasms (C00-C97)	53	5.2	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	52	5.1	2.1
6	Influenza and pneumonia (J09-J18)	32	3.2	1.3
7	Chronic lower respiratory diseases (J40-J47)	24	2.4	1.0
8	Cerebrovascular diseases (I60-I69)	17	1.7	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.2	*
10	Septicemia (A40-A41)	11	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.8	*
13	Anemias (D50-D64)	6	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
15	Meningitis (G00,G03)	2	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	2	0.2	*
15	Pneumonitis due to solids and liquids (J69)	2	0.2	*
...	All other causes (Residual)	232	22.9	9.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,150	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	313	27.2	5.1
2	Malignant neoplasms (C00-C97)	127	11.0	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	89	7.7	1.4
4	Chronic lower respiratory diseases (J40-J47)	81	7.0	1.3
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	78	6.8	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	4.3	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	3.7	0.7
8	Influenza and pneumonia (J09-J18)	25	2.2	0.4
9	Anemias (D50-D64)	22	1.9	0.4
10	Cerebrovascular diseases (I60-I69)	20	1.7	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	1.7	*
12	Septicemia (A40-A41)	11	1.0	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	*
14	Pneumonitis due to solids and liquids (J69)	5	0.4	*
15	Diabetes mellitus (E10-E14)	4	0.3	*
15	Meningitis (G00,G03)	4	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (Residual)	243	21.1	3.9
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	6,576	100.0	97.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,797	42.5	41.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,357	20.6	20.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	487	7.4	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	303	4.6	4.5
5	Malignant neoplasms (C00-C97)	257	3.9	3.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	1.6	1.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	100	1.5	1.5
8	Chronic lower respiratory diseases (J40-J47)	81	1.2	1.2
9	Anemias (D50-D64)	71	1.1	1.1
10	Pregnancy, childbirth and the puerperium (O00-O99)	62	0.9	0.9
11	Diabetes mellitus (E10-E14)	47	0.7	0.7
12	Legal intervention (Y35,Y89.0)	46	0.7	0.7
13	Influenza and pneumonia (J09-J18)	44	0.7	0.7
14	Septicemia (A40-A41)	35	0.5	0.5
15	Cerebrovascular diseases (I60-I69)	31	0.5	0.5
...	All other causes (Residual)	756	11.5	11.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-44 years				
...	All causes	22,568	100.0	209.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,622	16.0	33.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,425	15.2	31.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,291	14.6	30.6
4	Malignant neoplasms (C00-C97)	2,729	12.1	25.4
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,301	5.8	12.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	965	4.3	9.0
7	Diabetes mellitus (E10-E14)	713	3.2	6.6
8	Cerebrovascular diseases (I60-I69)	644	2.9	6.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	364	1.6	3.4
10	Septicemia (A40-A41)	312	1.4	2.9
11	Influenza and pneumonia (J09-J18)	285	1.3	2.7
12	Chronic lower respiratory diseases (J40-J47)	281	1.2	2.6
13	Anemias (D50-D64)	243	1.1	2.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	236	1.0	2.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	225	1.0	2.1
...	All other causes (Residual)	3,932	17.4	36.6
Non-Hispanic, black, both sexes, 45-64 years				
...	All causes	86,447	100.0	896.0
1	Malignant neoplasms (C00-C97)	25,117	29.1	260.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,771	24.0	215.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,359	5.0	45.2
4	Cerebrovascular diseases (I60-I69)	4,321	5.0	44.8
5	Diabetes mellitus (E10-E14)	4,181	4.8	43.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	2,408	2.8	25.0
7	Chronic lower respiratory diseases (J40-J47)	2,162	2.5	22.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,142	2.5	22.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,934	2.2	20.0
10	Septicemia (A40-A41)	1,742	2.0	18.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,339	1.5	13.9
12	Influenza and pneumonia (J09-J18)	1,096	1.3	11.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,058	1.2	11.0
14	Viral hepatitis (B15-B19)	872	1.0	9.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	542	0.6	5.6
...	All other causes (Residual)	12,403	14.3	128.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65 years and over				
...	All causes	162,044	100.0	4,550.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,698	26.3	1,199.1
2	Malignant neoplasms (C00-C97)	37,859	23.4	1,063.2
3	Cerebrovascular diseases (I60-I69)	10,644	6.6	298.9
4	Diabetes mellitus (E10-E14)	7,826	4.8	219.8
5	Chronic lower respiratory diseases (J40-J47)	6,353	3.9	178.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,475	3.4	153.8
7	Alzheimer's disease (G30)	5,369	3.3	150.8
8	Septicemia (A40-A41)	3,791	2.3	106.5
9	Influenza and pneumonia (J09-J18)	3,536	2.2	99.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,381	2.1	94.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,326	1.4	65.3
12	Pneumonitis due to solids and liquids (J69)	1,164	0.7	32.7
13	Parkinson's disease (G20-G21)	828	0.5	23.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	791	0.5	22.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	616	0.4	17.3
...	All other causes (Residual)	29,387	18.1	825.3
Non-Hispanic, black, male, all ages				
...	All causes	145,052	100.0	760.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,476	23.8	180.7
2	Malignant neoplasms (C00-C97)	33,628	23.2	176.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,219	5.7	43.1
4	Cerebrovascular diseases (I60-I69)	6,979	4.8	36.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6,658	4.6	34.9
6	Diabetes mellitus (E10-E14)	5,986	4.1	31.4
7	Chronic lower respiratory diseases (J40-J47)	4,648	3.2	24.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,734	2.6	19.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	2,693	1.9	14.1
10	Septicemia (A40-A41)	2,666	1.8	14.0
11	Influenza and pneumonia (J09-J18)	2,386	1.6	12.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,180	1.5	11.4
13	Certain conditions originating in the perinatal period (P00-P96)	2,171	1.5	11.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,790	1.2	9.4
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,750	1.2	9.2
...	All other causes (Residual)	25,088	17.3	131.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	575	100.0	45.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	29.0	13.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	84	14.6	6.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	9.9	4.5
4	Malignant neoplasms (C00-C97)	30	5.2	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	4.9	2.2
6	Influenza and pneumonia (J09-J18)	18	3.1	*
7	Chronic lower respiratory diseases (J40-J47)	16	2.8	*
8	Cerebrovascular diseases (I60-I69)	14	2.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	8	1.4	*
10	Septicemia (A40-A41)	6	1.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.0	*
13	Anemias (D50-D64)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Meningitis (G00,G03)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
...	All other causes (Residual)	127	22.1	10.1
Non-Hispanic, black, male, 5-14 years				
...	All causes	684	100.0	21.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	28.7	6.3
2	Malignant neoplasms (C00-C97)	73	10.7	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	8.9	1.9
4	Chronic lower respiratory diseases (J40-J47)	57	8.3	1.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	43	6.3	1.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	4.7	1.0
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	3.7	0.8
8	Influenza and pneumonia (J09-J18)	15	2.2	*
9	Anemias (D50-D64)	14	2.0	*
10	Cerebrovascular diseases (I60-I69)	11	1.6	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.5	*
12	Pneumonitis due to solids and liquids (J69)	5	0.7	*
13	Septicemia (A40-A41)	4	0.6	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
15	Diabetes mellitus (E10-E14)	3	0.4	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
...	All other causes (Residual)	125	18.3	4.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,040	100.0	148.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,508	49.8	73.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,012	20.1	29.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	408	8.1	12.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	197	3.9	5.8
5	Malignant neoplasms (C00-C97)	151	3.0	4.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	62	1.2	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	1.2	1.8
8	Chronic lower respiratory diseases (J40-J47)	54	1.1	1.6
9	Legal intervention (Y35,Y89.0)	45	0.9	1.3
10	Anemias (D50-D64)	43	0.9	1.3
11	Diabetes mellitus (E10-E14)	23	0.5	0.7
12	Influenza and pneumonia (J09-J18)	20	0.4	0.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.3	*
14	Cerebrovascular diseases (I60-I69)	16	0.3	*
15	Septicemia (A40-A41)	12	0.2	*
...	All other causes (Residual)	410	8.1	12.1
Non-Hispanic, black, male, 25-44 years				
...	All causes	13,887	100.0	272.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,006	21.6	58.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,350	16.9	46.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,280	16.4	44.7
4	Malignant neoplasms (C00-C97)	1,007	7.3	19.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	782	5.6	15.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	742	5.3	14.5
7	Diabetes mellitus (E10-E14)	386	2.8	7.6
8	Cerebrovascular diseases (I60-I69)	329	2.4	6.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	1.4	3.9
10	Septicemia (A40-A41)	156	1.1	3.1
11	Influenza and pneumonia (J09-J18)	148	1.1	2.9
12	Chronic lower respiratory diseases (J40-J47)	143	1.0	2.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	1.0	2.7
14	Anemias (D50-D64)	129	0.9	2.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	121	0.9	2.4
...	All other causes (Residual)	1,970	14.2	38.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-64 years				
...	All causes	50,209	100.0	1,123.5
1	Malignant neoplasms (C00-C97)	13,137	26.2	294.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,136	26.2	293.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,032	6.0	67.8
4	Cerebrovascular diseases (I60-I69)	2,482	4.9	55.5
5	Diabetes mellitus (E10-E14)	2,326	4.6	52.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,657	3.3	37.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,260	2.5	28.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,183	2.4	26.5
9	Chronic lower respiratory diseases (J40-J47)	1,116	2.2	25.0
10	Septicemia (A40-A41)	922	1.8	20.6
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	847	1.7	19.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	773	1.5	17.3
13	Influenza and pneumonia (J09-J18)	611	1.2	13.7
14	Viral hepatitis (B15-B19)	588	1.2	13.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	437	0.9	9.8
...	All other causes (Residual)	6,702	13.3	150.0
Non-Hispanic, black, male, 65 years and over				
...	All causes	70,756	100.0	5,037.3
1	Malignant neoplasms (C00-C97)	19,222	27.2	1,368.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,758	26.5	1,335.4
3	Cerebrovascular diseases (I60-I69)	4,102	5.8	292.0
4	Chronic lower respiratory diseases (J40-J47)	3,253	4.6	231.6
5	Diabetes mellitus (E10-E14)	3,248	4.6	231.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,314	3.3	164.7
7	Influenza and pneumonia (J09-J18)	1,529	2.2	108.9
8	Septicemia (A40-A41)	1,528	2.2	108.8
9	Alzheimer's disease (G30)	1,518	2.1	108.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,263	1.8	89.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,260	1.8	89.7
12	Pneumonitis due to solids and liquids (J69)	571	0.8	40.7
13	Parkinson's disease (G20-G21)	462	0.7	32.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	366	0.5	26.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	306	0.4	21.8
...	All other causes (Residual)	11,056	15.6	787.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	141,745	100.0	679.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,119	23.4	158.7
2	Malignant neoplasms (C00-C97)	32,530	22.9	155.9
3	Cerebrovascular diseases (I60-I69)	8,743	6.2	41.9
4	Diabetes mellitus (E10-E14)	6,785	4.8	32.5
5	Chronic lower respiratory diseases (J40-J47)	4,346	3.1	20.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,310	3.0	20.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,080	2.9	19.5
8	Alzheimer's disease (G30)	3,889	2.7	18.6
9	Septicemia (A40-A41)	3,297	2.3	15.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,785	2.0	13.3
11	Influenza and pneumonia (J09-J18)	2,712	1.9	13.0
12	Certain conditions originating in the perinatal period (P00-P96)	1,719	1.2	8.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,445	1.0	6.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,106	0.8	5.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,033	0.7	4.9
...	All other causes (Residual)	29,846	21.1	143.0
Non-Hispanic, black, female, 1-4 years				
...	All causes	436	100.0	35.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	27.3	9.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	16.1	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	11.0	3.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.5	2.0
5	Malignant neoplasms (C00-C97)	23	5.3	1.9
6	Influenza and pneumonia (J09-J18)	14	3.2	*
7	Chronic lower respiratory diseases (J40-J47)	8	1.8	*
8	Septicemia (A40-A41)	5	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	4	0.9	*
11	Anemias (D50-D64)	3	0.7	*
11	Cerebrovascular diseases (I60-I69)	3	0.7	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
14	Meningitis (G00,G03)	1	0.2	*
14	Atherosclerosis (I70)	1	0.2	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Peptic ulcer (K25-K28)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	102	23.4	8.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	466	100.0	15.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	117	25.1	3.9
2	Malignant neoplasms (C00-C97)	54	11.6	1.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	7.5	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	6.0	0.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	5.2	0.8
5	Chronic lower respiratory diseases (J40-J47)	24	5.2	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	2.4	*
8	Influenza and pneumonia (J09-J18)	10	2.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.9	*
9	Cerebrovascular diseases (I60-I69)	9	1.9	*
11	Anemias (D50-D64)	8	1.7	*
12	Septicemia (A40-A41)	7	1.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	1.1	*
14	Meningitis (G00,G03)	3	0.6	*
15	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (Residual)	120	25.8	4.0
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,536	100.0	45.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	345	22.5	10.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	289	18.8	8.6
3	Malignant neoplasms (C00-C97)	106	6.9	3.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	106	6.9	3.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	5.1	2.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	62	4.0	1.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	2.6	1.2
8	Human immunodeficiency virus (HIV) disease (B20-B24)	38	2.5	1.1
9	Anemias (D50-D64)	28	1.8	0.8
10	Chronic lower respiratory diseases (J40-J47)	27	1.8	0.8
11	Diabetes mellitus (E10-E14)	24	1.6	0.7
11	Influenza and pneumonia (J09-J18)	24	1.6	0.7
13	Septicemia (A40-A41)	23	1.5	0.7
14	Cerebrovascular diseases (I60-I69)	15	1.0	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.7	*
...	All other causes (Residual)	319	20.8	9.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-44 years				
...	All causes	8,681	100.0	153.8
1	Malignant neoplasms (C00-C97)	1,722	19.8	30.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,342	15.5	23.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	941	10.8	16.7
4	Human immunodeficiency virus (HIV) disease (B20-B24)	559	6.4	9.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	419	4.8	7.4
6	Diabetes mellitus (E10-E14)	327	3.8	5.8
7	Cerebrovascular diseases (I60-I69)	315	3.6	5.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	209	2.4	3.7
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	183	2.1	3.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	163	1.9	2.9
11	Septicemia (A40-A41)	156	1.8	2.8
12	Chronic lower respiratory diseases (J40-J47)	138	1.6	2.4
13	Influenza and pneumonia (J09-J18)	137	1.6	2.4
14	Anemias (D50-D64)	114	1.3	2.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	104	1.2	1.8
...	All other causes (Residual)	1,852	21.3	32.8
Non-Hispanic, black, female, 45-64 years				
...	All causes	36,238	100.0	699.8
1	Malignant neoplasms (C00-C97)	11,980	33.1	231.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,635	21.1	147.4
3	Diabetes mellitus (E10-E14)	1,855	5.1	35.8
4	Cerebrovascular diseases (I60-I69)	1,839	5.1	35.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,327	3.7	25.6
6	Chronic lower respiratory diseases (J40-J47)	1,046	2.9	20.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	959	2.6	18.5
8	Septicemia (A40-A41)	820	2.3	15.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	751	2.1	14.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	674	1.9	13.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	566	1.6	10.9
12	Influenza and pneumonia (J09-J18)	485	1.3	9.4
13	Viral hepatitis (B15-B19)	284	0.8	5.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	211	0.6	4.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	156	0.4	3.0
...	All other causes (Residual)	5,650	15.6	109.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2011

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> , Second Edition, 2004), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65 years and over				
...	All causes	91,288	100.0	4,233.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,940	26.2	1,110.3
2	Malignant neoplasms (C00-C97)	18,637	20.4	864.3
3	Cerebrovascular diseases (I60-I69)	6,542	7.2	303.4
4	Diabetes mellitus (E10-E14)	4,578	5.0	212.3
5	Alzheimer's disease (G30)	3,851	4.2	178.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,161	3.5	146.6
7	Chronic lower respiratory diseases (J40-J47)	3,100	3.4	143.8
8	Septicemia (A40-A41)	2,263	2.5	105.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,118	2.3	98.2
10	Influenza and pneumonia (J09-J18)	2,007	2.2	93.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,066	1.2	49.4
12	Pneumonitis due to solids and liquids (J69)	593	0.6	27.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	485	0.5	22.5
14	Parkinson's disease (G20-G21)	366	0.4	17.0
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	298	0.3	13.8
...	All other causes (Residual)	18,283	20.0	847.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2011