

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	156,419	100.0	295.0
1	Malignant neoplasms (C00-C97)	34,147	21.8	64.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,595	20.2	59.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,447	7.3	21.6
4	Cerebrovascular diseases (I60-I69)	7,767	5.0	14.6
5	Diabetes mellitus (E10-E14)	7,363	4.7	13.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,988	3.2	9.4
7	Chronic lower respiratory diseases (J40-J47)	4,477	2.9	8.4
8	Alzheimer's disease (G30)	3,772	2.4	7.1
9	Influenza and pneumonia (J09-J18)	3,049	1.9	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,964	1.9	5.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,837	1.8	5.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,740	1.8	5.2
13	Certain conditions originating in the perinatal period (P00-P96)	2,429	1.6	4.6
14	Septicemia (A40-A41)	2,304	1.5	4.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,859	1.2	3.5
...	All other causes (Residual)	32,681	20.9	61.6
Hispanic, both sexes, 1-4 years				
...	All causes	899	100.0	21.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	269	29.9	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	14.0	3.1
3	Malignant neoplasms (C00-C97)	99	11.0	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	6.7	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.2	0.7
6	Influenza and pneumonia (J09-J18)	22	2.4	0.5
7	Septicemia (A40-A41)	14	1.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.6	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.3	*
10	Cerebrovascular diseases (I60-I69)	11	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	9	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.6	*
13	Pneumonitis due to solids and liquids (J69)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
15	Anemias (D50-D64)	2	0.2	*
15	Diabetes mellitus (E10-E14)	2	0.2	*
15	Meningitis (G00,G03)	2	0.2	*
...	All other causes (Residual)	216	24.0	5.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-14 years				
...	All causes	1,083	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	307	28.3	3.2
2	Malignant neoplasms (C00-C97)	219	20.2	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	7.1	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	5.3	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	4.2	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	2.9	0.3
7	Cerebrovascular diseases (I60-I69)	24	2.2	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	1.8	0.2
9	Influenza and pneumonia (J09-J18)	18	1.7	*
9	Chronic lower respiratory diseases (J40-J47)	18	1.7	*
11	Septicemia (A40-A41)	16	1.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
13	Meningitis (G00,G03)	5	0.5	*
14	Anemias (D50-D64)	4	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	225	20.8	2.3
Hispanic, both sexes, 15-24 years				
...	All causes	4,773	100.0	52.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,871	39.2	20.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	915	19.2	10.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	675	14.1	7.4
4	Malignant neoplasms (C00-C97)	370	7.8	4.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	139	2.9	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	68	1.4	0.7
7	Cerebrovascular diseases (I60-I69)	36	0.8	0.4
8	Legal intervention (Y35,Y89.0)	30	0.6	0.3
9	Influenza and pneumonia (J09-J18)	26	0.5	0.3
9	Pregnancy, childbirth and the puerperium (O00-O99)	26	0.5	0.3
11	Diabetes mellitus (E10-E14)	23	0.5	0.3
12	Human immunodeficiency virus (HIV) disease (B20-B24)	19	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	19	0.4	*
14	Septicemia (A40-A41)	15	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	*
...	All other causes (Residual)	526	11.0	5.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 25-44 years				
...	All causes	14,554	100.0	89.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,972	27.3	24.3
2	Malignant neoplasms (C00-C97)	2,269	15.6	13.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,296	8.9	7.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,234	8.5	7.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,179	8.1	7.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	594	4.1	3.6
7	Cerebrovascular diseases (I60-I69)	386	2.7	2.4
8	Diabetes mellitus (E10-E14)	318	2.2	1.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	291	2.0	1.8
10	Influenza and pneumonia (J09-J18)	146	1.0	0.9
11	Septicemia (A40-A41)	140	1.0	0.9
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	0.8	0.7
13	Pregnancy, childbirth and the puerperium (O00-O99)	110	0.8	0.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	0.7	0.7
15	Chronic lower respiratory diseases (J40-J47)	76	0.5	0.5
...	All other causes (Residual)	2,317	15.9	14.2
Hispanic, both sexes, 45-64 years				
...	All causes	38,398	100.0	403.2
1	Malignant neoplasms (C00-C97)	11,190	29.1	117.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,932	18.1	72.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,868	7.5	30.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,849	7.4	29.9
5	Diabetes mellitus (E10-E14)	2,044	5.3	21.5
6	Cerebrovascular diseases (I60-I69)	1,615	4.2	17.0
7	Viral hepatitis (B15-B19)	873	2.3	9.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	718	1.9	7.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	692	1.8	7.3
10	Septicemia (A40-A41)	614	1.6	6.4
11	Chronic lower respiratory diseases (J40-J47)	567	1.5	6.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	543	1.4	5.7
13	Influenza and pneumonia (J09-J18)	479	1.2	5.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	365	1.0	3.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	327	0.9	3.4
...	All other causes (Residual)	5,722	14.9	60.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 65 years and over				
...	All causes	91,892	100.0	2,923.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,091	25.1	734.5
2	Malignant neoplasms (C00-C97)	19,985	21.7	635.7
3	Cerebrovascular diseases (I60-I69)	5,673	6.2	180.5
4	Diabetes mellitus (E10-E14)	4,972	5.4	158.2
5	Chronic lower respiratory diseases (J40-J47)	3,784	4.1	120.4
6	Alzheimer's disease (G30)	3,705	4.0	117.9
7	Influenza and pneumonia (J09-J18)	2,330	2.5	74.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,126	2.3	67.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,991	2.2	63.3
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,538	1.7	48.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,488	1.6	47.3
12	Septicemia (A40-A41)	1,478	1.6	47.0
13	Parkinson's disease (G20-G21)	1,092	1.2	34.7
14	Pneumonitis due to solids and liquids (J69)	706	0.8	22.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	528	0.6	16.8
...	All other causes (Residual)	17,405	18.9	553.6
Hispanic, male, all ages				
...	All causes	85,238	100.0	316.5
1	Malignant neoplasms (C00-C97)	17,849	20.9	66.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,329	20.3	64.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,255	9.7	30.7
4	Diabetes mellitus (E10-E14)	3,763	4.4	14.0
5	Cerebrovascular diseases (I60-I69)	3,636	4.3	13.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,422	4.0	12.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,292	2.7	8.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,277	2.7	8.5
9	Chronic lower respiratory diseases (J40-J47)	2,268	2.7	8.4
10	Influenza and pneumonia (J09-J18)	1,557	1.8	5.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,552	1.8	5.8
12	Certain conditions originating in the perinatal period (P00-P96)	1,373	1.6	5.1
13	Alzheimer's disease (G30)	1,193	1.4	4.4
14	Septicemia (A40-A41)	1,175	1.4	4.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	952	1.1	3.5
...	All other causes (Residual)	16,345	19.2	60.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	494	100.0	23.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	154	31.2	7.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	15.0	3.5
3	Malignant neoplasms (C00-C97)	56	11.3	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	6.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.6	*
6	Influenza and pneumonia (J09-J18)	11	2.2	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	1.8	*
8	Septicemia (A40-A41)	7	1.4	*
8	Cerebrovascular diseases (I60-I69)	7	1.4	*
10	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.8	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.6	*
12	Pneumonitis due to solids and liquids (J69)	3	0.6	*
14	Anemias (D50-D64)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (Residual)	114	23.1	5.4
Hispanic, male, 5-14 years				
...	All causes	608	100.0	12.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	32.1	3.9
2	Malignant neoplasms (C00-C97)	115	18.9	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	6.9	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	5.8	0.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	4.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	3.0	*
7	Cerebrovascular diseases (I60-I69)	14	2.3	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	2.0	*
9	Septicemia (A40-A41)	9	1.5	*
9	Influenza and pneumonia (J09-J18)	9	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	9	1.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
13	Anemias (D50-D64)	2	0.3	*
13	Diabetes mellitus (E10-E14)	2	0.3	*
13	Diseases of appendix (K35-K38)	2	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
...	All other causes (Residual)	110	18.1	2.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 15-24 years				
...	All causes	3,655	100.0	76.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,438	39.3	30.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	818	22.4	17.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	548	15.0	11.5
4	Malignant neoplasms (C00-C97)	228	6.2	4.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	98	2.7	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	1.1	0.9
7	Legal intervention (Y35,Y89.0)	29	0.8	0.6
8	Cerebrovascular diseases (I60-I69)	23	0.6	0.5
9	Influenza and pneumonia (J09-J18)	17	0.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.3	*
10	Diabetes mellitus (E10-E14)	12	0.3	*
10	Chronic lower respiratory diseases (J40-J47)	12	0.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.2	*
14	Septicemia (A40-A41)	7	0.2	*
15	Anemias (D50-D64)	6	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.2	*
...	All other causes (Residual)	351	9.6	7.3
Hispanic, male, 25-44 years				
...	All causes	10,171	100.0	119.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,151	31.0	37.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,044	10.3	12.3
3	Malignant neoplasms (C00-C97)	1,025	10.1	12.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	984	9.7	11.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	967	9.5	11.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	455	4.5	5.4
7	Cerebrovascular diseases (I60-I69)	250	2.5	2.9
8	Human immunodeficiency virus (HIV) disease (B20-B24)	226	2.2	2.7
9	Diabetes mellitus (E10-E14)	206	2.0	2.4
10	Influenza and pneumonia (J09-J18)	100	1.0	1.2
11	Septicemia (A40-A41)	83	0.8	1.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	72	0.7	0.8
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	0.6	0.8
14	Legal intervention (Y35,Y89.0)	62	0.6	0.7
15	Aortic aneurysm and dissection (I71)	56	0.6	0.7
...	All other causes (Residual)	1,424	14.0	16.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 45-64 years				
...	All causes	24,427	100.0	518.2
1	Malignant neoplasms (C00-C97)	5,932	24.3	125.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,933	20.2	104.6
3	Chronic liver disease and cirrhosis (K70,K73-K74)	2,152	8.8	45.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,148	8.8	45.6
5	Diabetes mellitus (E10-E14)	1,240	5.1	26.3
6	Cerebrovascular diseases (I60-I69)	982	4.0	20.8
7	Viral hepatitis (B15-B19)	661	2.7	14.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	566	2.3	12.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	421	1.7	8.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	419	1.7	8.9
11	Septicemia (A40-A41)	371	1.5	7.9
12	Chronic lower respiratory diseases (J40-J47)	320	1.3	6.8
13	Influenza and pneumonia (J09-J18)	296	1.2	6.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	279	1.1	5.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	214	0.9	4.5
...	All other causes (Residual)	3,493	14.3	74.1
Hispanic, male, 65 years and over				
...	All causes	43,214	100.0	3,211.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,234	26.0	834.9
2	Malignant neoplasms (C00-C97)	10,484	24.3	779.2
3	Cerebrovascular diseases (I60-I69)	2,350	5.4	174.7
4	Diabetes mellitus (E10-E14)	2,303	5.3	171.2
5	Chronic lower respiratory diseases (J40-J47)	1,875	4.3	139.3
6	Alzheimer's disease (G30)	1,161	2.7	86.3
7	Influenza and pneumonia (J09-J18)	1,107	2.6	82.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,082	2.5	80.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,044	2.4	77.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	812	1.9	60.3
11	Septicemia (A40-A41)	683	1.6	50.8
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	599	1.4	44.5
13	Parkinson's disease (G20-G21)	591	1.4	43.9
14	Pneumonitis due to solids and liquids (J69)	379	0.9	28.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	260	0.6	19.3
...	All other causes (Residual)	7,250	16.8	538.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, all ages				
...	All causes	71,181	100.0	272.7
1	Malignant neoplasms (C00-C97)	16,298	22.9	62.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,266	20.0	54.7
3	Cerebrovascular diseases (I60-I69)	4,131	5.8	15.8
4	Diabetes mellitus (E10-E14)	3,600	5.1	13.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,192	4.5	12.2
6	Alzheimer's disease (G30)	2,579	3.6	9.9
7	Chronic lower respiratory diseases (J40-J47)	2,209	3.1	8.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,566	2.2	6.0
9	Influenza and pneumonia (J09-J18)	1,492	2.1	5.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,412	2.0	5.4
11	Septicemia (A40-A41)	1,129	1.6	4.3
12	Certain conditions originating in the perinatal period (P00-P96)	1,056	1.5	4.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,017	1.4	3.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	838	1.2	3.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	545	0.8	2.1
...	All other causes (Residual)	15,851	22.3	60.7
Hispanic, female, 1-4 years				
...	All causes	405	100.0	20.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	28.4	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	12.8	2.6
3	Malignant neoplasms (C00-C97)	43	10.6	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	7.4	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	4.0	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	2.7	*
6	Influenza and pneumonia (J09-J18)	11	2.7	*
8	Septicemia (A40-A41)	7	1.7	*
9	Cerebrovascular diseases (I60-I69)	4	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	4	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	3	0.7	*
12	Diabetes mellitus (E10-E14)	2	0.5	*
12	Meningitis (G00,G03)	2	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.5	*
15	Anemias (D50-D64)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Hernia (K40-K46)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
...	All other causes (Residual)	98	24.2	4.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 5-14 years				
...	All causes	475	100.0	10.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	23.6	2.4
2	Malignant neoplasms (C00-C97)	104	21.9	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	7.4	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.6	0.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	17	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	2.7	*
7	Cerebrovascular diseases (I60-I69)	10	2.1	*
8	Influenza and pneumonia (J09-J18)	9	1.9	*
8	Chronic lower respiratory diseases (J40-J47)	9	1.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.7	*
11	Septicemia (A40-A41)	7	1.5	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
13	Meningitis (G00,G03)	4	0.8	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.6	*
15	Anemias (D50-D64)	2	0.4	*
15	Diabetes mellitus (E10-E14)	2	0.4	*
...	All other causes (Residual)	113	23.8	2.4
Hispanic, female, 15-24 years				
...	All causes	1,118	100.0	25.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	433	38.7	9.9
2	Malignant neoplasms (C00-C97)	142	12.7	3.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	127	11.4	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	97	8.7	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.7	0.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	26	2.3	0.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	2.3	0.6
8	Cerebrovascular diseases (I60-I69)	13	1.2	*
9	Diabetes mellitus (E10-E14)	11	1.0	*
10	Influenza and pneumonia (J09-J18)	9	0.8	*
11	Septicemia (A40-A41)	8	0.7	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	7	0.6	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.6	*
15	Anemias (D50-D64)	4	0.4	*
...	All other causes (Residual)	160	14.3	3.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 25-44 years				
...	All causes	4,383	100.0	56.0
1	Malignant neoplasms (C00-C97)	1,244	28.4	15.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	821	18.7	10.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	312	7.1	4.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	212	4.8	2.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	190	4.3	2.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	139	3.2	1.8
7	Cerebrovascular diseases (I60-I69)	136	3.1	1.7
8	Diabetes mellitus (E10-E14)	112	2.6	1.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	110	2.5	1.4
10	Human immunodeficiency virus (HIV) disease (B20-B24)	65	1.5	0.8
11	Septicemia (A40-A41)	57	1.3	0.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	1.1	0.6
13	Influenza and pneumonia (J09-J18)	46	1.0	0.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	41	0.9	0.5
15	Chronic lower respiratory diseases (J40-J47)	32	0.7	0.4
...	All other causes (Residual)	819	18.7	10.5
Hispanic, female, 45-64 years				
...	All causes	13,971	100.0	290.4
1	Malignant neoplasms (C00-C97)	5,258	37.6	109.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,999	14.3	41.6
3	Diabetes mellitus (E10-E14)	804	5.8	16.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	720	5.2	15.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	697	5.0	14.5
6	Cerebrovascular diseases (I60-I69)	633	4.5	13.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	273	2.0	5.7
8	Chronic lower respiratory diseases (J40-J47)	247	1.8	5.1
9	Septicemia (A40-A41)	243	1.7	5.1
10	Viral hepatitis (B15-B19)	212	1.5	4.4
11	Influenza and pneumonia (J09-J18)	183	1.3	3.8
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	152	1.1	3.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	122	0.9	2.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	113	0.8	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	86	0.6	1.8
...	All other causes (Residual)	2,229	16.0	46.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 65 years and over				
...	All causes	48,678	100.0	2,707.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,857	24.4	659.4
2	Malignant neoplasms (C00-C97)	9,501	19.5	528.4
3	Cerebrovascular diseases (I60-I69)	3,323	6.8	184.8
4	Diabetes mellitus (E10-E14)	2,669	5.5	148.4
5	Alzheimer's disease (G30)	2,544	5.2	141.5
6	Chronic lower respiratory diseases (J40-J47)	1,909	3.9	106.2
7	Influenza and pneumonia (J09-J18)	1,223	2.5	68.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,082	2.2	60.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	909	1.9	50.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	889	1.8	49.4
11	Septicemia (A40-A41)	795	1.6	44.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	726	1.5	40.4
13	Parkinson's disease (G20-G21)	501	1.0	27.9
14	Pneumonitis due to solids and liquids (J69)	327	0.7	18.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	268	0.6	14.9
...	All other causes (Residual)	10,155	20.9	564.7
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,016,896	100.0	1,004.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	481,991	23.9	240.2
2	Malignant neoplasms (C00-C97)	462,499	22.9	230.4
3	Chronic lower respiratory diseases (J40-J47)	127,116	6.3	63.3
4	Cerebrovascular diseases (I60-I69)	100,154	5.0	49.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	99,288	4.9	49.5
6	Alzheimer's disease (G30)	72,772	3.6	36.3
7	Diabetes mellitus (E10-E14)	50,443	2.5	25.1
8	Influenza and pneumonia (J09-J18)	40,460	2.0	20.2
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33,727	1.7	16.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33,105	1.6	16.5
11	Septicemia (A40-A41)	26,592	1.3	13.2
12	Chronic liver disease and cirrhosis (K70,K73-K74)	25,720	1.3	12.8
13	Parkinson's disease (G20-G21)	21,201	1.1	10.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,056	1.0	10.5
15	Pneumonitis due to solids and liquids (J69)	15,073	0.7	7.5
...	All other causes (Residual)	405,699	20.1	202.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	2,094	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	745	35.6	8.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	244	11.7	2.9
3	Malignant neoplasms (C00-C97)	202	9.6	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	6.5	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	73	3.5	0.9
6	Influenza and pneumonia (J09-J18)	46	2.2	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	1.5	0.4
8	Cerebrovascular diseases (I60-I69)	23	1.1	0.3
9	Septicemia (A40-A41)	22	1.1	0.3
10	Chronic lower respiratory diseases (J40-J47)	11	0.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	11	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.3	*
14	Anemias (D50-D64)	4	0.2	*
14	Diabetes mellitus (E10-E14)	4	0.2	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.2	*
...	All other causes (Residual)	522	24.9	6.2
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,667	100.0	11.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	798	29.9	3.5
2	Malignant neoplasms (C00-C97)	501	18.8	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	203	7.6	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	151	5.7	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	143	5.4	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	87	3.3	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	2.0	0.2
8	Influenza and pneumonia (J09-J18)	45	1.7	0.2
9	Cerebrovascular diseases (I60-I69)	44	1.6	0.2
10	Septicemia (A40-A41)	27	1.0	0.1
11	Chronic lower respiratory diseases (J40-J47)	25	0.9	0.1
12	Certain conditions originating in the perinatal period (P00-P96)	16	0.6	*
13	Diabetes mellitus (E10-E14)	12	0.4	*
14	Meningitis (G00,G03)	9	0.3	*
15	Anemias (D50-D64)	7	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.3	*
...	All other causes (Residual)	538	20.2	2.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	16,475	100.0	65.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,173	49.6	32.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,347	20.3	13.3
3	Malignant neoplasms (C00-C97)	858	5.2	3.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	706	4.3	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	480	2.9	1.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	254	1.5	1.0
7	Diabetes mellitus (E10-E14)	109	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	97	0.6	0.4
9	Influenza and pneumonia (J09-J18)	85	0.5	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	73	0.4	0.3
11	Septicemia (A40-A41)	66	0.4	0.3
12	Chronic lower respiratory diseases (J40-J47)	56	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.3	0.2
14	Legal intervention (Y35,Y89.0)	36	0.2	0.1
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	33	0.2	0.1
...	All other causes (Residual)	2,054	12.5	8.2
Non-Hispanic, white, both sexes, 25-44 years				
...	All causes	71,206	100.0	144.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22,321	31.3	45.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,113	14.2	20.5
3	Malignant neoplasms (C00-C97)	9,226	13.0	18.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,135	11.4	16.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,964	2.8	4.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,856	2.6	3.8
7	Diabetes mellitus (E10-E14)	1,376	1.9	2.8
8	Cerebrovascular diseases (I60-I69)	1,136	1.6	2.3
9	Chronic lower respiratory diseases (J40-J47)	619	0.9	1.3
10	Influenza and pneumonia (J09-J18)	618	0.9	1.3
11	Septicemia (A40-A41)	592	0.8	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	540	0.8	1.1
13	Human immunodeficiency virus (HIV) disease (B20-B24)	481	0.7	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	378	0.5	0.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	277	0.4	0.6
...	All other causes (Residual)	11,574	16.3	23.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 45-64 years				
...	All causes	364,732	100.0	620.5
1	Malignant neoplasms (C00-C97)	118,619	32.5	201.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	74,661	20.5	127.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27,624	7.6	47.0
4	Chronic lower respiratory diseases (J40-J47)	16,544	4.5	28.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	14,651	4.0	24.9
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13,986	3.8	23.8
7	Diabetes mellitus (E10-E14)	11,307	3.1	19.2
8	Cerebrovascular diseases (I60-I69)	9,864	2.7	16.8
9	Septicemia (A40-A41)	4,764	1.3	8.1
10	Viral hepatitis (B15-B19)	3,968	1.1	6.8
11	Influenza and pneumonia (J09-J18)	3,949	1.1	6.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,682	1.0	6.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,421	0.7	4.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,646	0.5	2.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,593	0.4	2.7
...	All other causes (Residual)	55,453	15.2	94.3
Non-Hispanic, white, both sexes, 65 years and over				
...	All causes	1,549,055	100.0	4,508.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	398,396	25.7	1,159.4
2	Malignant neoplasms (C00-C97)	333,050	21.5	969.2
3	Chronic lower respiratory diseases (J40-J47)	109,849	7.1	319.7
4	Cerebrovascular diseases (I60-I69)	88,947	5.7	258.9
5	Alzheimer's disease (G30)	71,983	4.6	209.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39,061	2.5	113.7
7	Diabetes mellitus (E10-E14)	37,634	2.4	109.5
8	Influenza and pneumonia (J09-J18)	35,648	2.3	103.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28,964	1.9	84.3
10	Septicemia (A40-A41)	21,050	1.4	61.3
11	Parkinson's disease (G20-G21)	20,685	1.3	60.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,447	1.2	53.7
13	Pneumonitis due to solids and liquids (J69)	13,538	0.9	39.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10,603	0.7	30.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	9,086	0.6	26.4
...	All other causes (Residual)	312,114	20.1	908.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, all ages				
...	All causes	998,832	100.0	1,011.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	250,116	25.0	253.2
2	Malignant neoplasms (C00-C97)	243,925	24.4	247.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60,388	6.0	61.1
4	Chronic lower respiratory diseases (J40-J47)	59,084	5.9	59.8
5	Cerebrovascular diseases (I60-I69)	39,786	4.0	40.3
6	Diabetes mellitus (E10-E14)	27,043	2.7	27.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26,356	2.6	26.7
8	Alzheimer's disease (G30)	22,403	2.2	22.7
9	Influenza and pneumonia (J09-J18)	18,960	1.9	19.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,720	1.7	16.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	16,518	1.7	16.7
12	Parkinson's disease (G20-G21)	12,654	1.3	12.8
13	Septicemia (A40-A41)	12,446	1.2	12.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,263	0.8	8.4
15	Pneumonitis due to solids and liquids (J69)	8,172	0.8	8.3
...	All other causes (Residual)	175,998	17.6	178.2
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,202	100.0	27.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	442	36.8	10.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	11.0	3.1
3	Malignant neoplasms (C00-C97)	108	9.0	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	81	6.7	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	2.8	0.8
6	Influenza and pneumonia (J09-J18)	31	2.6	0.7
7	Cerebrovascular diseases (I60-I69)	17	1.4	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.2	*
9	Septicemia (A40-A41)	11	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Diabetes mellitus (E10-E14)	4	0.3	*
15	Meningococcal infection (A39)	3	0.2	*
...	All other causes (Residual)	302	25.1	7.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,548	100.0	13.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	494	31.9	4.3
2	Malignant neoplasms (C00-C97)	288	18.6	2.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	149	9.6	1.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	72	4.7	0.6
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	4.6	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	2.8	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	1.9	0.2
8	Cerebrovascular diseases (I60-I69)	20	1.3	0.2
8	Influenza and pneumonia (J09-J18)	20	1.3	0.2
10	Septicemia (A40-A41)	19	1.2	*
11	Chronic lower respiratory diseases (J40-J47)	12	0.8	*
12	Diabetes mellitus (E10-E14)	8	0.5	*
13	Meningitis (G00,G03)	6	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
15	Anemias (D50-D64)	3	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	3	0.2	*
15	Pneumonitis due to solids and liquids (J69)	3	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.2	*
...	All other causes (Residual)	297	19.2	2.6
Non-Hispanic, white, male, 15-24 years				
...	All causes	11,826	100.0	92.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,050	51.2	47.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,723	23.0	21.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	500	4.2	3.9
4	Malignant neoplasms (C00-C97)	485	4.1	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	316	2.7	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	145	1.2	1.1
7	Diabetes mellitus (E10-E14)	65	0.5	0.5
8	Cerebrovascular diseases (I60-I69)	54	0.5	0.4
9	Influenza and pneumonia (J09-J18)	52	0.4	0.4
10	Chronic lower respiratory diseases (J40-J47)	37	0.3	0.3
11	Legal intervention (Y35,Y89.0)	33	0.3	0.3
12	Septicemia (A40-A41)	31	0.3	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.2	0.2
14	Pneumonitis due to solids and liquids (J69)	21	0.2	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	0.1	*
...	All other causes (Residual)	1,269	10.7	9.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 25-44 years				
...	All causes	45,942	100.0	185.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,701	34.2	63.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,871	17.1	31.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,640	12.3	22.7
4	Malignant neoplasms (C00-C97)	4,233	9.2	17.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,278	2.8	5.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,220	2.7	4.9
7	Diabetes mellitus (E10-E14)	833	1.8	3.4
8	Cerebrovascular diseases (I60-I69)	605	1.3	2.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	389	0.8	1.6
10	Influenza and pneumonia (J09-J18)	319	0.7	1.3
11	Septicemia (A40-A41)	314	0.7	1.3
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	307	0.7	1.2
13	Chronic lower respiratory diseases (J40-J47)	283	0.6	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	219	0.5	0.9
15	Aortic aneurysm and dissection (I71)	164	0.4	0.7
...	All other causes (Residual)	6,566	14.3	26.5
Non-Hispanic, white, male, 45-64 years				
...	All causes	223,841	100.0	773.1
1	Malignant neoplasms (C00-C97)	64,382	28.8	222.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,445	23.9	184.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,303	8.2	63.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,518	4.7	36.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	10,055	4.5	34.7
6	Chronic lower respiratory diseases (J40-J47)	8,323	3.7	28.7
7	Diabetes mellitus (E10-E14)	7,163	3.2	24.7
8	Cerebrovascular diseases (I60-I69)	5,602	2.5	19.3
9	Viral hepatitis (B15-B19)	2,831	1.3	9.8
10	Septicemia (A40-A41)	2,577	1.2	8.9
11	Influenza and pneumonia (J09-J18)	2,260	1.0	7.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,141	1.0	7.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,584	0.7	5.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,270	0.6	4.4
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,085	0.5	3.7
...	All other causes (Residual)	32,302	14.4	111.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 65 years and over				
...	All causes	708,546	100.0	4,673.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	190,543	26.9	1,256.9
2	Malignant neoplasms (C00-C97)	174,408	24.6	1,150.4
3	Chronic lower respiratory diseases (J40-J47)	50,415	7.1	332.6
4	Cerebrovascular diseases (I60-I69)	33,464	4.7	220.7
5	Alzheimer's disease (G30)	22,058	3.1	145.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,066	2.7	125.8
7	Diabetes mellitus (E10-E14)	18,970	2.7	125.1
8	Influenza and pneumonia (J09-J18)	16,238	2.3	107.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,314	2.0	94.4
10	Parkinson's disease (G20-G21)	12,328	1.7	81.3
11	Septicemia (A40-A41)	9,456	1.3	62.4
12	Pneumonitis due to solids and liquids (J69)	7,226	1.0	47.7
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,558	0.9	43.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,567	0.8	36.7
15	Chronic liver disease and cirrhosis (K70,K73-K74)	5,228	0.7	34.5
...	All other causes (Residual)	122,707	17.3	809.4
Non-Hispanic, white, female, all ages				
...	All causes	1,018,064	100.0	998.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	231,875	22.8	227.5
2	Malignant neoplasms (C00-C97)	218,574	21.5	214.4
3	Chronic lower respiratory diseases (J40-J47)	68,032	6.7	66.7
4	Cerebrovascular diseases (I60-I69)	60,368	5.9	59.2
5	Alzheimer's disease (G30)	50,369	4.9	49.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38,900	3.8	38.2
7	Diabetes mellitus (E10-E14)	23,400	2.3	23.0
8	Influenza and pneumonia (J09-J18)	21,500	2.1	21.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16,385	1.6	16.1
10	Septicemia (A40-A41)	14,146	1.4	13.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,793	1.3	12.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	9,202	0.9	9.0
13	Parkinson's disease (G20-G21)	8,547	0.8	8.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,371	0.7	7.2
15	Pneumonitis due to solids and liquids (J69)	6,901	0.7	6.8
...	All other causes (Residual)	229,701	22.6	225.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	892	100.0	21.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	303	34.0	7.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	12.6	2.7
3	Malignant neoplasms (C00-C97)	94	10.5	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	6.2	1.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	4.4	0.9
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	2.0	*
7	Influenza and pneumonia (J09-J18)	15	1.7	*
8	Septicemia (A40-A41)	11	1.2	*
9	Cerebrovascular diseases (I60-I69)	6	0.7	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
12	Acute bronchitis and bronchiolitis (J20-J21)	3	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
14	Anemias (D50-D64)	2	0.2	*
14	Pneumonitis due to solids and liquids (J69)	2	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
...	All other causes (Residual)	218	24.4	5.3
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,119	100.0	10.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	304	27.2	2.8
2	Malignant neoplasms (C00-C97)	213	19.0	1.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	7.1	0.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	71	6.3	0.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	54	4.8	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	44	3.9	0.4
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	2.2	0.2
7	Influenza and pneumonia (J09-J18)	25	2.2	0.2
9	Cerebrovascular diseases (I60-I69)	24	2.1	0.2
10	Chronic lower respiratory diseases (J40-J47)	13	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	11	1.0	*
12	Septicemia (A40-A41)	8	0.7	*
13	Anemias (D50-D64)	4	0.4	*
13	Diabetes mellitus (E10-E14)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
...	All other causes (Residual)	235	21.0	2.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,649	100.0	37.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,123	45.7	17.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	624	13.4	5.1
3	Malignant neoplasms (C00-C97)	373	8.0	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	206	4.4	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	164	3.5	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	109	2.3	0.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	73	1.6	0.6
8	Diabetes mellitus (E10-E14)	44	0.9	0.4
9	Cerebrovascular diseases (I60-I69)	43	0.9	0.3
10	Septicemia (A40-A41)	35	0.8	0.3
11	Influenza and pneumonia (J09-J18)	33	0.7	0.3
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	19	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17	0.4	*
15	Pneumonitis due to solids and liquids (J69)	8	0.2	*
...	All other causes (Residual)	759	16.3	6.2
Non-Hispanic, white, female, 25-44 years				
...	All causes	25,264	100.0	103.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,620	26.2	27.1
2	Malignant neoplasms (C00-C97)	4,993	19.8	20.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,495	9.9	10.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,242	8.9	9.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	744	2.9	3.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	578	2.3	2.4
7	Diabetes mellitus (E10-E14)	543	2.1	2.2
8	Cerebrovascular diseases (I60-I69)	531	2.1	2.2
9	Chronic lower respiratory diseases (J40-J47)	336	1.3	1.4
10	Influenza and pneumonia (J09-J18)	299	1.2	1.2
11	Septicemia (A40-A41)	278	1.1	1.1
12	Pregnancy, childbirth and the puerperium (O00-O99)	277	1.1	1.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	233	0.9	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	159	0.6	0.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	106	0.4	0.4
...	All other causes (Residual)	4,830	19.1	19.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 45-64 years				
...	All causes	140,891	100.0	472.4
1	Malignant neoplasms (C00-C97)	54,237	38.5	181.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,216	15.1	71.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,321	6.6	31.3
4	Chronic lower respiratory diseases (J40-J47)	8,221	5.8	27.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,596	3.3	15.4
6	Cerebrovascular diseases (I60-I69)	4,262	3.0	14.3
7	Diabetes mellitus (E10-E14)	4,144	2.9	13.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,468	2.5	11.6
9	Septicemia (A40-A41)	2,187	1.6	7.3
10	Influenza and pneumonia (J09-J18)	1,689	1.2	5.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,541	1.1	5.2
12	Viral hepatitis (B15-B19)	1,137	0.8	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	837	0.6	2.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	730	0.5	2.4
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	564	0.4	1.9
...	All other causes (Residual)	22,741	16.1	76.2
Non-Hispanic, white, female, 65 years and over				
...	All causes	840,509	100.0	4,377.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	207,853	24.7	1,082.5
2	Malignant neoplasms (C00-C97)	158,642	18.9	826.2
3	Chronic lower respiratory diseases (J40-J47)	59,434	7.1	309.5
4	Cerebrovascular diseases (I60-I69)	55,483	6.6	288.9
5	Alzheimer's disease (G30)	49,925	5.9	260.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,995	2.4	104.1
7	Influenza and pneumonia (J09-J18)	19,410	2.3	101.1
8	Diabetes mellitus (E10-E14)	18,664	2.2	97.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14,650	1.7	76.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,889	1.4	61.9
11	Septicemia (A40-A41)	11,594	1.4	60.4
12	Parkinson's disease (G20-G21)	8,357	1.0	43.5
13	Pneumonitis due to solids and liquids (J69)	6,312	0.8	32.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,036	0.6	26.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	3,858	0.5	20.1
...	All other causes (Residual)	189,407	22.5	986.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, all ages				
...	All causes	291,179	100.0	720.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	69,147	23.7	171.2
2	Malignant neoplasms (C00-C97)	66,565	22.9	164.8
3	Cerebrovascular diseases (I60-I69)	15,714	5.4	38.9
4	Diabetes mellitus (E10-E14)	12,836	4.4	31.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,451	4.3	30.8
6	Chronic lower respiratory diseases (J40-J47)	9,264	3.2	22.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,129	2.8	20.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,123	2.8	20.1
9	Septicemia (A40-A41)	5,918	2.0	14.7
10	Alzheimer's disease (G30)	5,392	1.9	13.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,140	1.8	12.7
12	Influenza and pneumonia (J09-J18)	4,923	1.7	12.2
13	Human immunodeficiency virus (HIV) disease (B20-B24)	3,848	1.3	9.5
14	Certain conditions originating in the perinatal period (P00-P96)	3,767	1.3	9.3
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,739	0.9	6.8
...	All other causes (Residual)	57,223	19.7	141.7
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	989	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	271	27.4	11.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	129	13.0	5.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	10.5	4.2
4	Malignant neoplasms (C00-C97)	64	6.5	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	4.3	1.7
6	Chronic lower respiratory diseases (J40-J47)	30	3.0	1.2
7	Influenza and pneumonia (J09-J18)	20	2.0	0.8
8	Septicemia (A40-A41)	17	1.7	*
8	Cerebrovascular diseases (I60-I69)	17	1.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	17	1.7	*
11	Anemias (D50-D64)	7	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.5	*
14	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.3	*
...	All other causes (Residual)	252	25.5	10.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,155	100.0	18.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	346	30.0	5.6
2	Malignant neoplasms (C00-C97)	140	12.1	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	104	9.0	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	77	6.7	1.2
5	Chronic lower respiratory diseases (J40-J47)	74	6.4	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	42	3.6	0.7
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	3.5	0.7
8	Diabetes mellitus (E10-E14)	17	1.5	*
9	Influenza and pneumonia (J09-J18)	15	1.3	*
10	Cerebrovascular diseases (I60-I69)	14	1.2	*
11	Anemias (D50-D64)	13	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	11	1.0	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
14	Certain conditions originating in the perinatal period (P00-P96)	8	0.7	*
15	Septicemia (A40-A41)	7	0.6	*
...	All other causes (Residual)	237	20.5	3.8
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	6,581	100.0	96.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,869	43.6	42.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,357	20.6	20.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	510	7.7	7.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	282	4.3	4.1
5	Malignant neoplasms (C00-C97)	244	3.7	3.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	87	1.3	1.3
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	83	1.3	1.2
8	Pregnancy, childbirth and the puerperium (O00-O99)	67	1.0	1.0
9	Anemias (D50-D64)	65	1.0	1.0
10	Chronic lower respiratory diseases (J40-J47)	62	0.9	0.9
11	Diabetes mellitus (E10-E14)	56	0.9	0.8
12	Legal intervention (Y35,Y89.0)	44	0.7	0.6
13	Cerebrovascular diseases (I60-I69)	42	0.6	0.6
14	Septicemia (A40-A41)	33	0.5	0.5
15	Influenza and pneumonia (J09-J18)	32	0.5	0.5
...	All other causes (Residual)	748	11.4	11.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 25-44 years				
...	All causes	22,450	100.0	206.8
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,685	16.4	33.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,647	16.2	33.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,350	14.9	30.9
4	Malignant neoplasms (C00-C97)	2,557	11.4	23.6
5	Human immunodeficiency virus (HIV) disease (B20-B24)	1,161	5.2	10.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,010	4.5	9.3
7	Diabetes mellitus (E10-E14)	705	3.1	6.5
8	Cerebrovascular diseases (I60-I69)	585	2.6	5.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	347	1.5	3.2
10	Chronic lower respiratory diseases (J40-J47)	303	1.3	2.8
11	Septicemia (A40-A41)	301	1.3	2.8
12	Anemias (D50-D64)	256	1.1	2.4
13	Influenza and pneumonia (J09-J18)	231	1.0	2.1
14	Pregnancy, childbirth and the puerperium (O00-O99)	228	1.0	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	227	1.0	2.1
...	All other causes (Residual)	3,857	17.2	35.5
Non-Hispanic, black, both sexes, 45-64 years				
...	All causes	87,505	100.0	895.5
1	Malignant neoplasms (C00-C97)	25,128	28.7	257.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,431	24.5	219.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,401	5.0	45.0
4	Cerebrovascular diseases (I60-I69)	4,337	5.0	44.4
5	Diabetes mellitus (E10-E14)	4,057	4.6	41.5
6	Chronic lower respiratory diseases (J40-J47)	2,235	2.6	22.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	2,231	2.5	22.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,224	2.5	22.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,804	2.1	18.5
10	Septicemia (A40-A41)	1,711	2.0	17.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,433	1.6	14.7
12	Influenza and pneumonia (J09-J18)	1,084	1.2	11.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,084	1.2	11.1
14	Viral hepatitis (B15-B19)	869	1.0	8.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	574	0.7	5.9
...	All other causes (Residual)	12,902	14.7	132.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 65 years and over				
...	All causes	165,703	100.0	4,456.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,596	26.3	1,172.4
2	Malignant neoplasms (C00-C97)	38,421	23.2	1,033.2
3	Cerebrovascular diseases (I60-I69)	10,683	6.4	287.3
4	Diabetes mellitus (E10-E14)	8,000	4.8	215.1
5	Chronic lower respiratory diseases (J40-J47)	6,553	4.0	176.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,502	3.3	148.0
7	Alzheimer's disease (G30)	5,322	3.2	143.1
8	Septicemia (A40-A41)	3,775	2.3	101.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,494	2.1	94.0
10	Influenza and pneumonia (J09-J18)	3,487	2.1	93.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,345	1.4	63.1
12	Pneumonitis due to solids and liquids (J69)	1,159	0.7	31.2
13	Parkinson's disease (G20-G21)	864	0.5	23.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	803	0.5	21.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	704	0.4	18.9
...	All other causes (Residual)	30,995	18.7	833.5
Non-Hispanic, black, male, all ages				
...	All causes	148,344	100.0	768.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,586	24.0	184.3
2	Malignant neoplasms (C00-C97)	34,005	22.9	176.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,402	5.7	43.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,039	4.7	36.5
5	Cerebrovascular diseases (I60-I69)	6,975	4.7	36.1
6	Diabetes mellitus (E10-E14)	6,072	4.1	31.5
7	Chronic lower respiratory diseases (J40-J47)	4,826	3.3	25.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,776	2.5	19.6
9	Septicemia (A40-A41)	2,677	1.8	13.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	2,491	1.7	12.9
11	Influenza and pneumonia (J09-J18)	2,371	1.6	12.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,288	1.5	11.9
13	Certain conditions originating in the perinatal period (P00-P96)	2,131	1.4	11.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,861	1.3	9.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,770	1.2	9.2
...	All other causes (Residual)	26,074	17.6	135.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	563	100.0	45.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	164	29.1	13.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	13.9	6.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	9.1	4.1
4	Malignant neoplasms (C00-C97)	39	6.9	3.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	28	5.0	2.2
6	Chronic lower respiratory diseases (J40-J47)	24	4.3	1.9
7	Influenza and pneumonia (J09-J18)	12	2.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	11	2.0	*
9	Septicemia (A40-A41)	8	1.4	*
9	Cerebrovascular diseases (I60-I69)	8	1.4	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.9	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
14	Anemias (D50-D64)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Diseases of appendix (K35-K38)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (Residual)	124	22.0	9.9
Non-Hispanic, black, male, 5-14 years				
...	All causes	707	100.0	22.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	228	32.2	7.3
2	Malignant neoplasms (C00-C97)	80	11.3	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	9.9	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	46	6.5	1.5
5	Chronic lower respiratory diseases (J40-J47)	40	5.7	1.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	4.7	1.1
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	3.3	0.7
8	Anemias (D50-D64)	8	1.1	*
8	Diabetes mellitus (E10-E14)	8	1.1	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	8	1.1	*
11	Influenza and pneumonia (J09-J18)	7	1.0	*
12	Cerebrovascular diseases (I60-I69)	6	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.7	*
14	Septicemia (A40-A41)	4	0.6	*
15	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
...	All other causes (Residual)	138	19.5	4.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,028	100.0	146.5
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,586	51.4	75.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	986	19.6	28.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	411	8.2	12.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	188	3.7	5.5
5	Malignant neoplasms (C00-C97)	142	2.8	4.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	61	1.2	1.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	1.1	1.6
8	Chronic lower respiratory diseases (J40-J47)	46	0.9	1.3
9	Legal intervention (Y35,Y89.0)	42	0.8	1.2
10	Anemias (D50-D64)	34	0.7	1.0
11	Diabetes mellitus (E10-E14)	26	0.5	0.8
12	Cerebrovascular diseases (I60-I69)	23	0.5	0.7
13	Influenza and pneumonia (J09-J18)	20	0.4	0.6
14	Septicemia (A40-A41)	13	0.3	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.1	*
...	All other causes (Residual)	389	7.7	11.3
Non-Hispanic, black, male, 25-44 years				
...	All causes	14,060	100.0	272.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,260	23.2	63.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,409	17.1	46.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,329	16.6	45.1
4	Malignant neoplasms (C00-C97)	951	6.8	18.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	823	5.9	15.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	649	4.6	12.6
7	Diabetes mellitus (E10-E14)	400	2.8	7.7
8	Cerebrovascular diseases (I60-I69)	325	2.3	6.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	182	1.3	3.5
10	Chronic lower respiratory diseases (J40-J47)	167	1.2	3.2
11	Septicemia (A40-A41)	145	1.0	2.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	129	0.9	2.5
13	Anemias (D50-D64)	123	0.9	2.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	114	0.8	2.2
14	Influenza and pneumonia (J09-J18)	114	0.8	2.2
...	All other causes (Residual)	1,940	13.8	37.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 45-64 years				
...	All causes	50,820	100.0	1,122.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,582	26.7	300.1
2	Malignant neoplasms (C00-C97)	13,221	26.0	292.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,063	6.0	67.7
4	Cerebrovascular diseases (I60-I69)	2,411	4.7	53.3
5	Diabetes mellitus (E10-E14)	2,283	4.5	50.4
6	Human immunodeficiency virus (HIV) disease (B20-B24)	1,497	2.9	33.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,213	2.4	26.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,203	2.4	26.6
9	Chronic lower respiratory diseases (J40-J47)	1,150	2.3	25.4
10	Septicemia (A40-A41)	880	1.7	19.4
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	877	1.7	19.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	838	1.6	18.5
13	Influenza and pneumonia (J09-J18)	627	1.2	13.9
14	Viral hepatitis (B15-B19)	588	1.2	13.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	446	0.9	9.9
...	All other causes (Residual)	6,941	13.7	153.4
Non-Hispanic, black, male, 65 years and over				
...	All causes	73,353	100.0	4,974.5
1	Malignant neoplasms (C00-C97)	19,566	26.7	1,326.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	19,380	26.4	1,314.3
3	Cerebrovascular diseases (I60-I69)	4,183	5.7	283.7
4	Chronic lower respiratory diseases (J40-J47)	3,395	4.6	230.2
5	Diabetes mellitus (E10-E14)	3,354	4.6	227.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,358	3.2	159.9
7	Septicemia (A40-A41)	1,577	2.1	106.9
8	Influenza and pneumonia (J09-J18)	1,560	2.1	105.8
9	Alzheimer's disease (G30)	1,499	2.0	101.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,333	1.8	90.4
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,326	1.8	89.9
12	Pneumonitis due to solids and liquids (J69)	593	0.8	40.2
13	Parkinson's disease (G20-G21)	516	0.7	35.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	435	0.6	29.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	361	0.5	24.5
...	All other causes (Residual)	11,917	16.2	808.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, all ages				
...	All causes	142,835	100.0	677.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	33,561	23.5	159.2
2	Malignant neoplasms (C00-C97)	32,560	22.8	154.4
3	Cerebrovascular diseases (I60-I69)	8,739	6.1	41.4
4	Diabetes mellitus (E10-E14)	6,764	4.7	32.1
5	Chronic lower respiratory diseases (J40-J47)	4,438	3.1	21.0
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,353	3.0	20.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,049	2.8	19.2
8	Alzheimer's disease (G30)	3,861	2.7	18.3
9	Septicemia (A40-A41)	3,241	2.3	15.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,852	2.0	13.5
11	Influenza and pneumonia (J09-J18)	2,552	1.8	12.1
12	Certain conditions originating in the perinatal period (P00-P96)	1,636	1.1	7.8
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1,357	1.0	6.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,084	0.8	5.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	969	0.7	4.6
...	All other causes (Residual)	30,819	21.6	146.1
Non-Hispanic, black, female, 1-4 years				
...	All causes	426	100.0	35.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	107	25.1	8.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	53	12.4	4.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	51	12.0	4.2
4	Malignant neoplasms (C00-C97)	25	5.9	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	3.5	*
6	Septicemia (A40-A41)	9	2.1	*
6	Cerebrovascular diseases (I60-I69)	9	2.1	*
8	Influenza and pneumonia (J09-J18)	8	1.9	*
9	Anemias (D50-D64)	6	1.4	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
13	Meningitis (G00,G03)	2	0.5	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.5	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Nutritional deficiencies (E40-E64)	1	0.2	*
15	Aortic aneurysm and dissection (I71)	1	0.2	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
...	All other causes (Residual)	119	27.9	9.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 5-14 years				
...	All causes	448	100.0	14.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	26.3	3.9
2	Malignant neoplasms (C00-C97)	60	13.4	2.0
3	Chronic lower respiratory diseases (J40-J47)	34	7.6	1.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	7.6	1.1
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	6.9	1.0
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	4.0	*
7	Diabetes mellitus (E10-E14)	9	2.0	*
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	2.0	*
9	Cerebrovascular diseases (I60-I69)	8	1.8	*
9	Influenza and pneumonia (J09-J18)	8	1.8	*
11	Anemias (D50-D64)	5	1.1	*
11	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.9	*
14	Septicemia (A40-A41)	3	0.7	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.7	*
...	All other causes (Residual)	99	22.1	3.3
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,553	100.0	46.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	371	23.9	11.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	283	18.2	8.4
3	Malignant neoplasms (C00-C97)	102	6.6	3.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	6.4	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	94	6.1	2.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	67	4.3	2.0
7	Anemias (D50-D64)	31	2.0	0.9
8	Diabetes mellitus (E10-E14)	30	1.9	0.9
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	1.9	0.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	26	1.7	0.8
11	Septicemia (A40-A41)	20	1.3	0.6
12	Cerebrovascular diseases (I60-I69)	19	1.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.2	*
14	Chronic lower respiratory diseases (J40-J47)	16	1.0	*
15	Influenza and pneumonia (J09-J18)	12	0.8	*
...	All other causes (Residual)	335	21.6	9.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 25-44 years				
...	All causes	8,390	100.0	147.5
1	Malignant neoplasms (C00-C97)	1,606	19.1	28.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,318	15.7	23.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	941	11.2	16.5
4	Human immunodeficiency virus (HIV) disease (B20-B24)	512	6.1	9.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	425	5.1	7.5
6	Diabetes mellitus (E10-E14)	305	3.6	5.4
7	Cerebrovascular diseases (I60-I69)	260	3.1	4.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	228	2.7	4.0
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	187	2.2	3.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	165	2.0	2.9
11	Septicemia (A40-A41)	156	1.9	2.7
12	Chronic lower respiratory diseases (J40-J47)	136	1.6	2.4
13	Anemias (D50-D64)	133	1.6	2.3
14	Influenza and pneumonia (J09-J18)	117	1.4	2.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	98	1.2	1.7
...	All other causes (Residual)	1,803	21.5	31.7
Non-Hispanic, black, female, 45-64 years				
...	All causes	36,685	100.0	699.3
1	Malignant neoplasms (C00-C97)	11,907	32.5	227.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,849	21.4	149.6
3	Cerebrovascular diseases (I60-I69)	1,926	5.3	36.7
4	Diabetes mellitus (E10-E14)	1,774	4.8	33.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,338	3.6	25.5
6	Chronic lower respiratory diseases (J40-J47)	1,085	3.0	20.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,011	2.8	19.3
8	Septicemia (A40-A41)	831	2.3	15.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	734	2.0	14.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	601	1.6	11.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	595	1.6	11.3
12	Influenza and pneumonia (J09-J18)	457	1.2	8.7
13	Viral hepatitis (B15-B19)	281	0.8	5.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	207	0.6	3.9
15	Anemias (D50-D64)	155	0.4	3.0
...	All other causes (Residual)	5,934	16.2	113.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012

LCWK6. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by Hispanic origin, race for non-Hispanic population and sex: United States, 2012

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 65 years and over				
...	All causes	92,350	100.0	4,115.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	24,216	26.2	1,079.2
2	Malignant neoplasms (C00-C97)	18,855	20.4	840.3
3	Cerebrovascular diseases (I60-I69)	6,500	7.0	289.7
4	Diabetes mellitus (E10-E14)	4,646	5.0	207.0
5	Alzheimer's disease (G30)	3,823	4.1	170.4
6	Chronic lower respiratory diseases (J40-J47)	3,158	3.4	140.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,144	3.4	140.1
8	Septicemia (A40-A41)	2,198	2.4	98.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,161	2.3	96.3
10	Influenza and pneumonia (J09-J18)	1,927	2.1	85.9
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,019	1.1	45.4
12	Pneumonitis due to solids and liquids (J69)	566	0.6	25.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	442	0.5	19.7
14	Parkinson's disease (G20-G21)	348	0.4	15.5
15	Anemias (D50-D64)	284	0.3	12.7
...	All other causes (Residual)	19,063	20.6	849.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality, 2012