

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Infants, all races, both sexes				
...	All causes	24,586	100.0	614.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5,107	20.8	127.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	4,148	16.9	103.7
3	Sudden infant death syndrome (R95)	2,063	8.4	51.6
4	Newborn affected by maternal complications of pregnancy (P01)	1,561	6.3	39.0
5	Accidents (unintentional injuries) (V01-X59)	1,110	4.5	27.8
6	Newborn affected by complications of placenta, cord and membranes (P02)	1,030	4.2	25.8
7	Bacterial sepsis of newborn (P36)	583	2.4	14.6
8	Respiratory distress of newborn (P22)	514	2.1	12.9
9	Diseases of the circulatory system (I00-I99)	507	2.1	12.7
10	Necrotizing enterocolitis of newborn (P77)	472	1.9	11.8
11	Neonatal hemorrhage (P50-P52,P54)	469	1.9	11.7
12	Diarrhea and gastroenteritis of infectious origin (A09)	316	1.3	7.9
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	314	1.3	7.9
14	Assault (homicide) (*U01,X85-Y09)	311	1.3	7.8
15	Atelectasis (P28.0-P28.1)	248	1.0	6.2
...	All other causes (Residual)	5,833	23.7	145.8
Infants, all races, male				
...	All causes	13,702	100.0	669.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,618	19.1	127.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,319	16.9	113.3
3	Sudden infant death syndrome (R95)	1,211	8.8	59.2
4	Newborn affected by maternal complications of pregnancy (P01)	856	6.2	41.8
5	Accidents (unintentional injuries) (V01-X59)	637	4.6	31.1
6	Newborn affected by complications of placenta, cord and membranes (P02)	553	4.0	27.0
7	Bacterial sepsis of newborn (P36)	325	2.4	15.9
8	Respiratory distress of newborn (P22)	312	2.3	15.2
9	Diseases of the circulatory system (I00-I99)	286	2.1	14.0
9	Neonatal hemorrhage (P50-P52,P54)	286	2.1	14.0
11	Necrotizing enterocolitis of newborn (P77)	278	2.0	13.6
12	Diarrhea and gastroenteritis of infectious origin (A09)	187	1.4	9.1
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	178	1.3	8.7
14	Assault (homicide) (*U01,X85-Y09)	177	1.3	8.6
15	Atelectasis (P28.0-P28.1)	142	1.0	6.9
...	All other causes (Residual)	3,337	24.4	163.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Infants, all races, female				
...	All causes	10,884	100.0	557.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,489	22.9	127.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,829	16.8	93.7
3	Sudden infant death syndrome (R95)	852	7.8	43.6
4	Newborn affected by maternal complications of pregnancy (P01)	705	6.5	36.1
5	Newborn affected by complications of placenta, cord and membranes (P02)	477	4.4	24.4
6	Accidents (unintentional injuries) (V01-X59)	473	4.3	24.2
7	Bacterial sepsis of newborn (P36)	258	2.4	13.2
8	Diseases of the circulatory system (I00-I99)	221	2.0	11.3
9	Respiratory distress of newborn (P22)	202	1.9	10.3
10	Necrotizing enterocolitis of newborn (P77)	194	1.8	9.9
11	Neonatal hemorrhage (P50-P52,P54)	183	1.7	9.4
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	136	1.2	7.0
13	Assault (homicide) (*U01,X85-Y09)	134	1.2	6.9
14	Diarrhea and gastroenteritis of infectious origin (A09)	129	1.2	6.6
15	Atelectasis (P28.0-P28.1)	106	1.0	5.4
...	All other causes (Residual)	2,496	22.9	127.8
Infants, white, both sexes				
...	All causes	15,954	100.0	519.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3,822	24.0	124.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,314	14.5	75.4
3	Sudden infant death syndrome (R95)	1,311	8.2	42.7
4	Newborn affected by maternal complications of pregnancy (P01)	991	6.2	32.3
5	Accidents (unintentional injuries) (V01-X59)	703	4.4	22.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	643	4.0	20.9
7	Bacterial sepsis of newborn (P36)	382	2.4	12.4
8	Neonatal hemorrhage (P50-P52,P54)	337	2.1	11.0
9	Diseases of the circulatory system (I00-I99)	326	2.0	10.6
10	Respiratory distress of newborn (P22)	318	2.0	10.4
11	Necrotizing enterocolitis of newborn (P77)	282	1.8	9.2
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	214	1.3	7.0
13	Assault (homicide) (*U01,X85-Y09)	213	1.3	6.9
14	Atelectasis (P28.0-P28.1)	180	1.1	5.9
15	Diarrhea and gastroenteritis of infectious origin (A09)	170	1.1	5.5
...	All other causes (Residual)	3,748	23.5	122.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Infants, white, male				
...	All causes	8,871	100.0	564.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,941	21.9	123.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,294	14.6	82.3
3	Sudden infant death syndrome (R95)	771	8.7	49.1
4	Newborn affected by maternal complications of pregnancy (P01)	554	6.2	35.3
5	Accidents (unintentional injuries) (V01-X59)	402	4.5	25.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	353	4.0	22.5
7	Bacterial sepsis of newborn (P36)	213	2.4	13.6
8	Neonatal hemorrhage (P50-P52,P54)	202	2.3	12.9
9	Respiratory distress of newborn (P22)	194	2.2	12.3
10	Diseases of the circulatory system (I00-I99)	181	2.0	11.5
11	Necrotizing enterocolitis of newborn (P77)	169	1.9	10.8
12	Assault (homicide) (*U01,X85-Y09)	129	1.5	8.2
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	121	1.4	7.7
14	Atelectasis (P28.0-P28.1)	105	1.2	6.7
15	Diarrhea and gastroenteritis of infectious origin (A09)	99	1.1	6.3
...	All other causes (Residual)	2,143	24.2	136.4
Infants, white, female				
...	All causes	7,083	100.0	472.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,881	26.6	125.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,020	14.4	68.1
3	Sudden infant death syndrome (R95)	540	7.6	36.1
4	Newborn affected by maternal complications of pregnancy (P01)	437	6.2	29.2
5	Accidents (unintentional injuries) (V01-X59)	301	4.2	20.1
6	Newborn affected by complications of placenta, cord and membranes (P02)	290	4.1	19.4
7	Bacterial sepsis of newborn (P36)	169	2.4	11.3
8	Diseases of the circulatory system (I00-I99)	145	2.0	9.7
9	Neonatal hemorrhage (P50-P52,P54)	135	1.9	9.0
10	Respiratory distress of newborn (P22)	124	1.8	8.3
11	Necrotizing enterocolitis of newborn (P77)	113	1.6	7.5
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	93	1.3	6.2
13	Assault (homicide) (*U01,X85-Y09)	84	1.2	5.6
14	Atelectasis (P28.0-P28.1)	75	1.1	5.0
15	Diarrhea and gastroenteritis of infectious origin (A09)	71	1.0	4.7
...	All other causes (Residual)	1,605	22.7	107.2

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Infants, black, both sexes				
...	All causes	7,401	100.0	1,162.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,635	22.1	256.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,010	13.6	158.7
3	Sudden infant death syndrome (R95)	668	9.0	105.0
4	Newborn affected by maternal complications of pregnancy (P01)	486	6.6	76.4
5	Accidents (unintentional injuries) (V01-X59)	354	4.8	55.6
6	Newborn affected by complications of placenta, cord and membranes (P02)	327	4.4	51.4
7	Respiratory distress of newborn (P22)	176	2.4	27.7
7	Bacterial sepsis of newborn (P36)	176	2.4	27.7
9	Necrotizing enterocolitis of newborn (P77)	174	2.4	27.3
10	Diseases of the circulatory system (I00-I99)	152	2.1	23.9
11	Diarrhea and gastroenteritis of infectious origin (A09)	133	1.8	20.9
12	Neonatal hemorrhage (P50-P52,P54)	118	1.6	18.5
13	Septicemia (A40-A41)	90	1.2	14.1
14	Assault (homicide) (*U01,X85-Y09)	88	1.2	13.8
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	76	1.0	11.9
...	All other causes (Residual)	1,738	23.5	273.1
Infants, black, male				
...	All causes	4,116	100.0	1,270.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	907	22.0	280.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	530	12.9	163.6
3	Sudden infant death syndrome (R95)	395	9.6	121.9
4	Newborn affected by maternal complications of pregnancy (P01)	255	6.2	78.7
5	Accidents (unintentional injuries) (V01-X59)	202	4.9	62.4
6	Newborn affected by complications of placenta, cord and membranes (P02)	159	3.9	49.1
7	Respiratory distress of newborn (P22)	106	2.6	32.7
8	Bacterial sepsis of newborn (P36)	99	2.4	30.6
8	Necrotizing enterocolitis of newborn (P77)	99	2.4	30.6
10	Diseases of the circulatory system (I00-I99)	88	2.1	27.2
11	Diarrhea and gastroenteritis of infectious origin (A09)	80	1.9	24.7
12	Neonatal hemorrhage (P50-P52,P54)	74	1.8	22.8
13	Septicemia (A40-A41)	47	1.1	14.5
14	Influenza and pneumonia (J09-J18)	43	1.0	13.3
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	42	1.0	13.0
15	Assault (homicide) (*U01,X85-Y09)	42	1.0	13.0
...	All other causes (Residual)	948	23.0	292.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Infants, black, female				
...	All causes	3,285	100.0	1,051.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	728	22.2	233.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	480	14.6	153.6
3	Sudden infant death syndrome (R95)	273	8.3	87.4
4	Newborn affected by maternal complications of pregnancy (P01)	231	7.0	73.9
5	Newborn affected by complications of placenta, cord and membranes (P02)	168	5.1	53.8
6	Accidents (unintentional injuries) (V01-X59)	152	4.6	48.6
7	Bacterial sepsis of newborn (P36)	77	2.3	24.6
8	Necrotizing enterocolitis of newborn (P77)	75	2.3	24.0
9	Respiratory distress of newborn (P22)	70	2.1	22.4
10	Diseases of the circulatory system (I00-I99)	64	1.9	20.5
11	Diarrhea and gastroenteritis of infectious origin (A09)	53	1.6	17.0
12	Assault (homicide) (*U01,X85-Y09)	46	1.4	14.7
13	Neonatal hemorrhage (P50-P52,P54)	44	1.3	14.1
14	Septicemia (A40-A41)	43	1.3	13.8
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	34	1.0	10.9
...	All other causes (Residual)	747	22.7	239.1
Infants, American Indian, both sexes				
...	All causes	354	100.0	757.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	23.7	179.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	44	12.4	94.1
3	Sudden infant death syndrome (R95)	38	10.7	81.3
4	Accidents (unintentional injuries) (V01-X59)	26	7.3	55.6
5	Newborn affected by complications of placenta, cord and membranes (P02)	16	4.5	*
6	Newborn affected by maternal complications of pregnancy (P01)	11	3.1	*
7	Influenza and pneumonia (J09-J18)	9	2.5	*
8	Diseases of the circulatory system (I00-I99)	8	2.3	*
9	Diarrhea and gastroenteritis of infectious origin (A09)	6	1.7	*
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	6	1.7	*
11	Bacterial sepsis of newborn (P36)	5	1.4	*
11	Assault (homicide) (*U01,X85-Y09)	5	1.4	*
13	Bronchitis, chronic and unspecified (J40-J42)	4	1.1	*
13	Respiratory distress of newborn (P22)	4	1.1	*
13	Neonatal hemorrhage (P50-P52,P54)	4	1.1	*
...	All other causes (Residual)	84	23.7	179.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Infants, American Indian, male				
...	All causes	213	100.0	899.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	22.1	198.5
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	23	10.8	97.2
3	Sudden infant death syndrome (R95)	21	9.9	88.7
4	Accidents (unintentional injuries) (V01-X59)	16	7.5	*
5	Newborn affected by complications of placenta, cord and membranes (P02)	13	6.1	*
6	Newborn affected by maternal complications of pregnancy (P01)	8	3.8	*
7	Diseases of the circulatory system (I00-I99)	6	2.8	*
8	Influenza and pneumonia (J09-J18)	5	2.3	*
9	Bronchitis, chronic and unspecified (J40-J42)	4	1.9	*
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	1.9	*
9	Bacterial sepsis of newborn (P36)	4	1.9	*
12	Diarrhea and gastroenteritis of infectious origin (A09)	3	1.4	*
12	Assault (homicide) (*U01,X85-Y09)	3	1.4	*
14	Septicemia (A40-A41)	2	0.9	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	2	0.9	*
14	Newborn affected by other complications of labor and delivery (P03)	2	0.9	*
14	Respiratory distress of newborn (P22)	2	0.9	*
14	Neonatal hemorrhage (P50-P52,P54)	2	0.9	*
14	Necrotizing enterocolitis of newborn (P77)	2	0.9	*
...	All other causes (Residual)	44	20.7	185.9
Infants, American Indian, female				
...	All causes	141	100.0	610.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	26.2	160.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	21	14.9	91.0
3	Sudden infant death syndrome (R95)	17	12.1	*
4	Accidents (unintentional injuries) (V01-X59)	10	7.1	*
5	Influenza and pneumonia (J09-J18)	4	2.8	*
6	Diarrhea and gastroenteritis of infectious origin (A09)	3	2.1	*
6	Newborn affected by maternal complications of pregnancy (P01)	3	2.1	*
6	Newborn affected by complications of placenta, cord and membranes (P02)	3	2.1	*
9	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	2	1.4	*
9	Diseases of the circulatory system (I00-I99)	2	1.4	*
9	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	2	1.4	*
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	2	1.4	*
9	Respiratory distress of newborn (P22)	2	1.4	*
9	Neonatal hemorrhage (P50-P52,P54)	2	1.4	*
9	Assault (homicide) (*U01,X85-Y09)	2	1.4	*
...	All other causes (Residual)	29	20.6	125.6

Source: CDC/NCHS, National Vital Statistics System

<sup>1</sup> Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Infants, Asian or Pacific Islander, both sexes				
...	All causes	877	100.0	355.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	21.8	77.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	155	17.7	62.8
3	Newborn affected by maternal complications of pregnancy (P01)	73	8.3	29.6
4	Sudden infant death syndrome (R95)	46	5.2	18.6
5	Newborn affected by complications of placenta, cord and membranes (P02)	44	5.0	17.8
6	Accidents (unintentional injuries) (V01-X59)	27	3.1	10.9
7	Diseases of the circulatory system (I00-I99)	21	2.4	8.5
8	Bacterial sepsis of newborn (P36)	20	2.3	8.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	18	2.1	*
10	Respiratory distress of newborn (P22)	16	1.8	*
11	Necrotizing enterocolitis of newborn (P77)	13	1.5	*
12	Influenza and pneumonia (J09-J18)	12	1.4	*
13	Hydrops fetalis not due to hemolytic disease (P83.2)	11	1.3	*
14	Neonatal hemorrhage (P50-P52,P54)	10	1.1	*
15	Pulmonary hemorrhage originating in the perinatal period (P26)	9	1.0	*
...	All other causes (Residual)	211	24.1	85.5
Infants, Asian or Pacific Islander, male				
...	All causes	502	100.0	392.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	100	19.9	78.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	95	18.9	74.3
3	Newborn affected by maternal complications of pregnancy (P01)	39	7.8	30.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	28	5.6	21.9
5	Sudden infant death syndrome (R95)	24	4.8	18.8
6	Accidents (unintentional injuries) (V01-X59)	17	3.4	*
7	Diseases of the circulatory system (I00-I99)	11	2.2	*
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	11	2.2	*
9	Respiratory distress of newborn (P22)	10	2.0	*
10	Bacterial sepsis of newborn (P36)	9	1.8	*
11	Neonatal hemorrhage (P50-P52,P54)	8	1.6	*
11	Necrotizing enterocolitis of newborn (P77)	8	1.6	*
13	Influenza and pneumonia (J09-J18)	7	1.4	*
14	Pulmonary hemorrhage originating in the perinatal period (P26)	6	1.2	*
14	Atelectasis (P28.0-P28.1)	6	1.2	*
14	Hydrops fetalis not due to hemolytic disease (P83.2)	6	1.2	*
...	All other causes (Residual)	117	23.3	91.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Infants, Asian or Pacific Islander, female				
...	All causes	375	100.0	315.0
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	91	24.3	76.4
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	60	16.0	50.4
3	Newborn affected by maternal complications of pregnancy (P01)	34	9.1	28.6
4	Sudden infant death syndrome (R95)	22	5.9	18.5
5	Newborn affected by complications of placenta, cord and membranes (P02)	16	4.3	*
6	Bacterial sepsis of newborn (P36)	11	2.9	*
7	Diseases of the circulatory system (I00-I99)	10	2.7	*
7	Accidents (unintentional injuries) (V01-X59)	10	2.7	*
9	Septicemia (A40-A41)	7	1.9	*
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	7	1.9	*
11	Respiratory distress of newborn (P22)	6	1.6	*
12	Influenza and pneumonia (J09-J18)	5	1.3	*
12	Necrotizing enterocolitis of newborn (P77)	5	1.3	*
12	Hydrops fetalis not due to hemolytic disease (P83.2)	5	1.3	*
15	Malignant neoplasms (C00-C97)	3	0.8	*
15	Pulmonary hemorrhage originating in the perinatal period (P26)	3	0.8	*
15	Chronic respiratory disease originating in the perinatal period (P27)	3	0.8	*
...	All other causes (Residual)	77	20.5	64.7
Neonates, all races, both sexes				
...	All causes	16,188	100.0	404.8
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	4,063	25.1	101.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3,605	22.3	90.1
3	Newborn affected by maternal complications of pregnancy (P01)	1,547	9.6	38.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	1,015	6.3	25.4
5	Bacterial sepsis of newborn (P36)	552	3.4	13.8
6	Respiratory distress of newborn (P22)	497	3.1	12.4
7	Neonatal hemorrhage (P50-P52,P54)	458	2.8	11.5
8	Necrotizing enterocolitis of newborn (P77)	409	2.5	10.2
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	300	1.9	7.5
10	Atelectasis (P28.0-P28.1)	231	1.4	5.8
11	Sudden infant death syndrome (R95)	197	1.2	4.9
12	Pulmonary hemorrhage originating in the perinatal period (P26)	167	1.0	4.2
13	Hydrops fetalis not due to hemolytic disease (P83.2)	142	0.9	3.6
14	Accidents (unintentional injuries) (V01-X59)	132	0.8	3.3
15	Newborn affected by other complications of labor and delivery (P03)	108	0.7	2.7
...	All other causes (Residual)	2,765	17.1	69.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, all races, male				
...	All causes	8,953	100.0	437.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,279	25.5	111.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,880	21.0	91.8
3	Newborn affected by maternal complications of pregnancy (P01)	850	9.5	41.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	543	6.1	26.5
5	Bacterial sepsis of newborn (P36)	310	3.5	15.1
6	Respiratory distress of newborn (P22)	297	3.3	14.5
7	Neonatal hemorrhage (P50-P52,P54)	280	3.1	13.7
8	Necrotizing enterocolitis of newborn (P77)	246	2.7	12.0
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	170	1.9	8.3
10	Atelectasis (P28.0-P28.1)	129	1.4	6.3
11	Sudden infant death syndrome (R95)	108	1.2	5.3
12	Pulmonary hemorrhage originating in the perinatal period (P26)	100	1.1	4.9
13	Accidents (unintentional injuries) (V01-X59)	75	0.8	3.7
14	Hydrops fetalis not due to hemolytic disease (P83.2)	73	0.8	3.6
15	Diseases of the circulatory system (I00-I99)	68	0.8	3.3
...	All other causes (Residual)	1,545	17.3	75.5
Neonates, all races, female				
...	All causes	7,235	100.0	370.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,784	24.7	91.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,725	23.8	88.4
3	Newborn affected by maternal complications of pregnancy (P01)	697	9.6	35.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	472	6.5	24.2
5	Bacterial sepsis of newborn (P36)	242	3.3	12.4
6	Respiratory distress of newborn (P22)	200	2.8	10.2
7	Neonatal hemorrhage (P50-P52,P54)	178	2.5	9.1
8	Necrotizing enterocolitis of newborn (P77)	163	2.3	8.3
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	130	1.8	6.7
10	Atelectasis (P28.0-P28.1)	102	1.4	5.2
11	Sudden infant death syndrome (R95)	89	1.2	4.6
12	Hydrops fetalis not due to hemolytic disease (P83.2)	69	1.0	3.5
13	Pulmonary hemorrhage originating in the perinatal period (P26)	67	0.9	3.4
14	Accidents (unintentional injuries) (V01-X59)	57	0.8	2.9
15	Newborn affected by other complications of labor and delivery (P03)	47	0.6	2.4
...	All other causes (Residual)	1,213	16.8	62.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, both sexes				
...	All causes	10,612	100.0	345.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,760	26.0	89.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,267	21.4	73.9
3	Newborn affected by maternal complications of pregnancy (P01)	981	9.2	32.0
4	Newborn affected by complications of placenta, cord and membranes (P02)	633	6.0	20.6
5	Bacterial sepsis of newborn (P36)	360	3.4	11.7
6	Neonatal hemorrhage (P50-P52,P54)	333	3.1	10.8
7	Respiratory distress of newborn (P22)	308	2.9	10.0
8	Necrotizing enterocolitis of newborn (P77)	250	2.4	8.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	208	2.0	6.8
10	Atelectasis (P28.0-P28.1)	169	1.6	5.5
11	Sudden infant death syndrome (R95)	127	1.2	4.1
12	Hydrops fetalis not due to hemolytic disease (P83.2)	116	1.1	3.8
13	Pulmonary hemorrhage originating in the perinatal period (P26)	98	0.9	3.2
14	Accidents (unintentional injuries) (V01-X59)	91	0.9	3.0
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	83	0.8	2.7
...	All other causes (Residual)	1,828	17.2	59.6
Neonates, white, male				
...	All causes	5,856	100.0	372.6
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,426	24.4	90.7
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,267	21.6	80.6
3	Newborn affected by maternal complications of pregnancy (P01)	551	9.4	35.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	345	5.9	22.0
5	Bacterial sepsis of newborn (P36)	202	3.4	12.9
6	Neonatal hemorrhage (P50-P52,P54)	200	3.4	12.7
7	Respiratory distress of newborn (P22)	186	3.2	11.8
8	Necrotizing enterocolitis of newborn (P77)	154	2.6	9.8
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	117	2.0	7.4
10	Atelectasis (P28.0-P28.1)	96	1.6	6.1
11	Sudden infant death syndrome (R95)	73	1.2	4.6
12	Pulmonary hemorrhage originating in the perinatal period (P26)	61	1.0	3.9
13	Hydrops fetalis not due to hemolytic disease (P83.2)	56	1.0	3.6
14	Accidents (unintentional injuries) (V01-X59)	52	0.9	3.3
15	Diseases of the circulatory system (I00-I99)	49	0.8	3.1
...	All other causes (Residual)	1,021	17.4	65.0

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, female				
...	All causes	4,756	100.0	317.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,334	28.0	89.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,000	21.0	66.8
3	Newborn affected by maternal complications of pregnancy (P01)	430	9.0	28.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	288	6.1	19.2
5	Bacterial sepsis of newborn (P36)	158	3.3	10.5
6	Neonatal hemorrhage (P50-P52,P54)	133	2.8	8.9
7	Respiratory distress of newborn (P22)	122	2.6	8.1
8	Necrotizing enterocolitis of newborn (P77)	96	2.0	6.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	91	1.9	6.1
10	Atelectasis (P28.0-P28.1)	73	1.5	4.9
11	Hydrops fetalis not due to hemolytic disease (P83.2)	60	1.3	4.0
12	Sudden infant death syndrome (R95)	54	1.1	3.6
13	Accidents (unintentional injuries) (V01-X59)	39	0.8	2.6
14	Interstitial emphysema and related conditions originating in the perinatal period (P25)	37	0.8	2.5
14	Pulmonary hemorrhage originating in the perinatal period (P26)	37	0.8	2.5
...	All other causes (Residual)	804	16.9	53.7
Neonates, black, both sexes				
...	All causes	4,769	100.0	749.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,600	33.6	251.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	658	13.8	103.4
3	Newborn affected by maternal complications of pregnancy (P01)	483	10.1	75.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	322	6.8	50.6
5	Respiratory distress of newborn (P22)	170	3.6	26.7
6	Bacterial sepsis of newborn (P36)	167	3.5	26.2
7	Necrotizing enterocolitis of newborn (P77)	144	3.0	22.6
8	Neonatal hemorrhage (P50-P52,P54)	112	2.3	17.6
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	68	1.4	10.7
10	Pulmonary hemorrhage originating in the perinatal period (P26)	60	1.3	9.4
11	Atelectasis (P28.0-P28.1)	56	1.2	8.8
11	Sudden infant death syndrome (R95)	56	1.2	8.8
13	Accidents (unintentional injuries) (V01-X59)	35	0.7	5.5
14	Newborn affected by maternal hypertensive disorders (P00.0)	32	0.7	5.0
14	Newborn affected by other complications of labor and delivery (P03)	32	0.7	5.0
...	All other causes (Residual)	774	16.2	121.6

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, black, male				
...	All causes	2,616	100.0	807.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	897	34.3	276.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	345	13.2	106.5
3	Newborn affected by maternal complications of pregnancy (P01)	253	9.7	78.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	157	6.0	48.5
5	Respiratory distress of newborn (P22)	100	3.8	30.9
6	Bacterial sepsis of newborn (P36)	95	3.6	29.3
7	Necrotizing enterocolitis of newborn (P77)	83	3.2	25.6
8	Neonatal hemorrhage (P50-P52,P54)	71	2.7	21.9
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	38	1.5	11.7
10	Pulmonary hemorrhage originating in the perinatal period (P26)	33	1.3	10.2
11	Atelectasis (P28.0-P28.1)	30	1.1	9.3
12	Sudden infant death syndrome (R95)	28	1.1	8.6
13	Newborn affected by maternal hypertensive disorders (P00.0)	19	0.7	*
13	Accidents (unintentional injuries) (V01-X59)	19	0.7	*
15	Newborn affected by other complications of labor and delivery (P03)	16	0.6	*
15	Slow fetal growth and fetal malnutrition (P05)	16	0.6	*
...	All other causes (Residual)	416	15.9	128.4
Neonates, black, female				
...	All causes	2,153	100.0	689.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	703	32.7	225.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	313	14.5	100.2
3	Newborn affected by maternal complications of pregnancy (P01)	230	10.7	73.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	165	7.7	52.8
5	Bacterial sepsis of newborn (P36)	72	3.3	23.0
6	Respiratory distress of newborn (P22)	70	3.3	22.4
7	Necrotizing enterocolitis of newborn (P77)	61	2.8	19.5
8	Neonatal hemorrhage (P50-P52,P54)	41	1.9	13.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	30	1.4	9.6
10	Sudden infant death syndrome (R95)	28	1.3	9.0
11	Pulmonary hemorrhage originating in the perinatal period (P26)	27	1.3	8.6
12	Atelectasis (P28.0-P28.1)	26	1.2	8.3
13	Newborn affected by other complications of labor and delivery (P03)	16	0.7	*
13	Accidents (unintentional injuries) (V01-X59)	16	0.7	*
15	Newborn affected by maternal hypertensive disorders (P00.0)	13	0.6	*
15	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	13	0.6	*
15	Slow fetal growth and fetal malnutrition (P05)	13	0.6	*
...	All other causes (Residual)	316	14.7	101.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, American Indian, both sexes				
...	All causes	189	100.0	404.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	29.1	117.6
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	44	23.3	94.1
3	Newborn affected by complications of placenta, cord and membranes (P02)	16	8.5	*
4	Newborn affected by maternal complications of pregnancy (P01)	11	5.8	*
5	Intrauterine hypoxia and birth asphyxia (P20-P21)	6	3.2	*
5	Sudden infant death syndrome (R95)	6	3.2	*
7	Bacterial sepsis of newborn (P36)	5	2.6	*
8	Respiratory distress of newborn (P22)	4	2.1	*
9	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	3	1.6	*
9	Neonatal hemorrhage (P50-P52,P54)	3	1.6	*
9	Necrotizing enterocolitis of newborn (P77)	3	1.6	*
12	Newborn affected by other complications of labor and delivery (P03)	2	1.1	*
12	Hydrops fetalis not due to hemolytic disease (P83.2)	2	1.1	*
12	Accidents (unintentional injuries) (V01-X59)	2	1.1	*
15	Diarrhea and gastroenteritis of infectious origin (A09)	1	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
15	Diseases of the circulatory system (I00-I99)	1	0.5	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	0.5	*
15	Newborn affected by maternal hypertensive disorders (P00.0)	1	0.5	*
15	Newborn affected by noxious influences transmitted via placenta or breast milk (P04)	1	0.5	*
15	Birth trauma (P10-P15)	1	0.5	*
15	Neonatal aspiration syndromes (P24)	1	0.5	*
15	Atelectasis (P28.0-P28.1)	1	0.5	*
...	All other causes (Residual)	18	9.5	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, American Indian, male				
...	All causes	120	100.0	506.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	29.2	147.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	23	19.2	97.2
3	Newborn affected by complications of placenta, cord and membranes (P02)	13	10.8	*
4	Newborn affected by maternal complications of pregnancy (P01)	8	6.7	*
5	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	3.3	*
5	Bacterial sepsis of newborn (P36)	4	3.3	*
7	Newborn affected by other complications of labor and delivery (P03)	2	1.7	*
7	Respiratory distress of newborn (P22)	2	1.7	*
7	Necrotizing enterocolitis of newborn (P77)	2	1.7	*
7	Sudden infant death syndrome (R95)	2	1.7	*
11	Diarrhea and gastroenteritis of infectious origin (A09)	1	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
11	Diseases of the circulatory system (I00-I99)	1	0.8	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	0.8	*
11	Newborn affected by maternal hypertensive disorders (P00.0)	1	0.8	*
11	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	1	0.8	*
11	Neonatal aspiration syndromes (P24)	1	0.8	*
11	Neonatal hemorrhage (P50-P52,P54)	1	0.8	*
11	Hydrops fetalis not due to hemolytic disease (P83.2)	1	0.8	*
11	Accidents (unintentional injuries) (V01-X59)	1	0.8	*
...	All other causes (Residual)	15	12.5	*
Neonates, American Indian, female				
...	All causes	69	100.0	298.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	21	30.4	91.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	29.0	86.6
3	Sudden infant death syndrome (R95)	4	5.8	*
4	Newborn affected by maternal complications of pregnancy (P01)	3	4.3	*
4	Newborn affected by complications of placenta, cord and membranes (P02)	3	4.3	*
6	Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9)	2	2.9	*
6	Intrauterine hypoxia and birth asphyxia (P20-P21)	2	2.9	*
6	Respiratory distress of newborn (P22)	2	2.9	*
6	Neonatal hemorrhage (P50-P52,P54)	2	2.9	*
10	Newborn affected by noxious influences transmitted via placenta or breast milk (P04)	1	1.4	*
10	Birth trauma (P10-P15)	1	1.4	*
10	Atelectasis (P28.0-P28.1)	1	1.4	*
10	Bacterial sepsis of newborn (P36)	1	1.4	*
10	Necrotizing enterocolitis of newborn (P77)	1	1.4	*
10	Hydrops fetalis not due to hemolytic disease (P83.2)	1	1.4	*
10	Accidents (unintentional injuries) (V01-X59)	1	1.4	*
...	All other causes (Residual)	3	4.3	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, Asian or Pacific Islander, both sexes				
...	All causes	618	100.0	250.3
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	152	24.6	61.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	132	21.4	53.5
3	Newborn affected by maternal complications of pregnancy (P01)	72	11.7	29.2
4	Newborn affected by complications of placenta, cord and membranes (P02)	44	7.1	17.8
5	Bacterial sepsis of newborn (P36)	20	3.2	8.1
6	Intrauterine hypoxia and birth asphyxia (P20-P21)	18	2.9	*
7	Respiratory distress of newborn (P22)	15	2.4	*
8	Necrotizing enterocolitis of newborn (P77)	12	1.9	*
9	Neonatal hemorrhage (P50-P52,P54)	10	1.6	*
9	Hydrops fetalis not due to hemolytic disease (P83.2)	10	1.6	*
11	Pulmonary hemorrhage originating in the perinatal period (P26)	9	1.5	*
12	Sudden infant death syndrome (R95)	8	1.3	*
13	Diseases of the circulatory system (I00-I99)	5	0.8	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	5	0.8	*
13	Atelectasis (P28.0-P28.1)	5	0.8	*
...	All other causes (Residual)	101	16.3	40.9
Neonates, Asian or Pacific Islander, male				
...	All causes	361	100.0	282.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	92	25.5	72.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	20.5	57.9
3	Newborn affected by maternal complications of pregnancy (P01)	38	10.5	29.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	28	7.8	21.9
5	Intrauterine hypoxia and birth asphyxia (P20-P21)	11	3.0	*
6	Respiratory distress of newborn (P22)	9	2.5	*
6	Bacterial sepsis of newborn (P36)	9	2.5	*
8	Neonatal hemorrhage (P50-P52,P54)	8	2.2	*
9	Necrotizing enterocolitis of newborn (P77)	7	1.9	*
10	Pulmonary hemorrhage originating in the perinatal period (P26)	6	1.7	*
11	Hydrops fetalis not due to hemolytic disease (P83.2)	5	1.4	*
11	Sudden infant death syndrome (R95)	5	1.4	*
13	Diseases of the circulatory system (I00-I99)	4	1.1	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	4	1.1	*
15	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	3	0.8	*
15	Atelectasis (P28.0-P28.1)	3	0.8	*
15	Hematological disorders (P60-P61)	3	0.8	*
15	Accidents (unintentional injuries) (V01-X59)	3	0.8	*
...	All other causes (Residual)	49	13.6	38.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, Asian or Pacific Islander, female				
...	All causes	257	100.0	215.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	60	23.3	50.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	22.6	48.7
3	Newborn affected by maternal complications of pregnancy (P01)	34	13.2	28.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	16	6.2	*
5	Bacterial sepsis of newborn (P36)	11	4.3	*
6	Intrauterine hypoxia and birth asphyxia (P20-P21)	7	2.7	*
7	Respiratory distress of newborn (P22)	6	2.3	*
8	Necrotizing enterocolitis of newborn (P77)	5	1.9	*
8	Hydrops fetalis not due to hemolytic disease (P83.2)	5	1.9	*
10	Pulmonary hemorrhage originating in the perinatal period (P26)	3	1.2	*
10	Sudden infant death syndrome (R95)	3	1.2	*
12	Newborn affected by other complications of labor and delivery (P03)	2	0.8	*
12	Slow fetal growth and fetal malnutrition (P05)	2	0.8	*
12	Atelectasis (P28.0-P28.1)	2	0.8	*
12	Neonatal hemorrhage (P50-P52,P54)	2	0.8	*
...	All other causes (Residual)	41	16.0	34.4
Postneonates, all races, both sexes				
...	All causes	8,398	100.0	210.0
1	Sudden infant death syndrome (R95)	1,866	22.2	46.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,502	17.9	37.6
3	Accidents (unintentional injuries) (V01-X59)	978	11.6	24.5
4	Diseases of the circulatory system (I00-I99)	402	4.8	10.1
5	Diarrhea and gastroenteritis of infectious origin (A09)	300	3.6	7.5
6	Assault (homicide) (*U01,X85-Y09)	281	3.3	7.0
7	Septicemia (A40-A41)	212	2.5	5.3
8	Influenza and pneumonia (J09-J18)	193	2.3	4.8
9	Chronic respiratory disease originating in the perinatal period (P27)	90	1.1	2.3
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	85	1.0	2.1
11	Necrotizing enterocolitis of newborn (P77)	63	0.8	1.6
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	57	0.7	1.4
13	Malignant neoplasms (C00-C97)	50	0.6	1.3
14	Meningitis (G00,G03)	48	0.6	1.2
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	44	0.5	1.1
...	All other causes (Residual)	2,227	26.5	55.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable



LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, all races, male				
...	All causes	4,749	100.0	232.0
1	Sudden infant death syndrome (R95)	1,103	23.2	53.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	738	15.5	36.1
3	Accidents (unintentional injuries) (V01-X59)	562	11.8	27.5
4	Diseases of the circulatory system (I00-I99)	218	4.6	10.7
5	Diarrhea and gastroenteritis of infectious origin (A09)	177	3.7	8.6
6	Assault (homicide) (*U01,X85-Y09)	164	3.5	8.0
7	Septicemia (A40-A41)	118	2.5	5.8
8	Influenza and pneumonia (J09-J18)	110	2.3	5.4
9	Chronic respiratory disease originating in the perinatal period (P27)	49	1.0	2.4
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	40	0.8	2.0
11	Necrotizing enterocolitis of newborn (P77)	32	0.7	1.6
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	29	0.6	1.4
13	Malignant neoplasms (C00-C97)	26	0.5	1.3
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	25	0.5	1.2
15	Meningitis (G00,G03)	23	0.5	1.1
...	All other causes (Residual)	1,335	28.1	65.2
Postneonates, all races, female				
...	All causes	3,649	100.0	186.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	764	20.9	39.1
2	Sudden infant death syndrome (R95)	763	20.9	39.1
3	Accidents (unintentional injuries) (V01-X59)	416	11.4	21.3
4	Diseases of the circulatory system (I00-I99)	184	5.0	9.4
5	Diarrhea and gastroenteritis of infectious origin (A09)	123	3.4	6.3
6	Assault (homicide) (*U01,X85-Y09)	117	3.2	6.0
7	Septicemia (A40-A41)	94	2.6	4.8
8	Influenza and pneumonia (J09-J18)	83	2.3	4.3
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	45	1.2	2.3
10	Chronic respiratory disease originating in the perinatal period (P27)	41	1.1	2.1
11	Necrotizing enterocolitis of newborn (P77)	31	0.8	1.6
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	28	0.8	1.4
13	Meningitis (G00,G03)	25	0.7	1.3
14	Malignant neoplasms (C00-C97)	24	0.7	1.2
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	19	0.5	*
...	All other causes (Residual)	892	24.4	45.7

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, both sexes				
...	All causes	5,342	100.0	174.0
1	Sudden infant death syndrome (R95)	1,184	22.2	38.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,062	19.9	34.6
3	Accidents (unintentional injuries) (V01-X59)	612	11.5	19.9
4	Diseases of the circulatory system (I00-I99)	249	4.7	8.1
5	Assault (homicide) (*U01,X85-Y09)	191	3.6	6.2
6	Diarrhea and gastroenteritis of infectious origin (A09)	163	3.1	5.3
7	Septicemia (A40-A41)	114	2.1	3.7
8	Influenza and pneumonia (J09-J18)	99	1.9	3.2
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	47	0.9	1.5
10	Chronic respiratory disease originating in the perinatal period (P27)	46	0.9	1.5
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	42	0.8	1.4
12	Malignant neoplasms (C00-C97)	37	0.7	1.2
13	Necrotizing enterocolitis of newborn (P77)	32	0.6	1.0
14	Meningitis (G00,G03)	26	0.5	0.8
15	Anoxic brain damage, not elsewhere classified (G93.1)	24	0.4	0.8
...	All other causes (Residual)	1,414	26.5	46.1
Postneonates, white, male				
...	All causes	3,015	100.0	191.9
1	Sudden infant death syndrome (R95)	698	23.2	44.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	515	17.1	32.8
3	Accidents (unintentional injuries) (V01-X59)	350	11.6	22.3
4	Diseases of the circulatory system (I00-I99)	132	4.4	8.4
5	Assault (homicide) (*U01,X85-Y09)	121	4.0	7.7
6	Diarrhea and gastroenteritis of infectious origin (A09)	96	3.2	6.1
7	Septicemia (A40-A41)	69	2.3	4.4
8	Influenza and pneumonia (J09-J18)	55	1.8	3.5
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	27	0.9	1.7
10	Chronic respiratory disease originating in the perinatal period (P27)	23	0.8	1.5
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	21	0.7	1.3
12	Malignant neoplasms (C00-C97)	20	0.7	1.3
13	Necrotizing enterocolitis of newborn (P77)	15	0.5	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	13	0.4	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	13	0.4	*
...	All other causes (Residual)	847	28.1	53.9

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, female				
...	All causes	2,327	100.0	155.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	547	23.5	36.5
2	Sudden infant death syndrome (R95)	486	20.9	32.4
3	Accidents (unintentional injuries) (V01-X59)	262	11.3	17.5
4	Diseases of the circulatory system (I00-I99)	117	5.0	7.8
5	Assault (homicide) (*U01,X85-Y09)	70	3.0	4.7
6	Diarrhea and gastroenteritis of infectious origin (A09)	67	2.9	4.5
7	Septicemia (A40-A41)	45	1.9	3.0
8	Influenza and pneumonia (J09-J18)	44	1.9	2.9
9	Chronic respiratory disease originating in the perinatal period (P27)	23	1.0	1.5
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	21	0.9	1.4
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	20	0.9	1.3
12	Malignant neoplasms (C00-C97)	17	0.7	*
12	Necrotizing enterocolitis of newborn (P77)	17	0.7	*
14	Meningitis (G00,G03)	14	0.6	*
15	Anoxic brain damage, not elsewhere classified (G93.1)	13	0.6	*
...	All other causes (Residual)	564	24.2	37.7
Postneonates, black, both sexes				
...	All causes	2,632	100.0	413.6
1	Sudden infant death syndrome (R95)	612	23.3	96.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	352	13.4	55.3
3	Accidents (unintentional injuries) (V01-X59)	319	12.1	50.1
4	Diseases of the circulatory system (I00-I99)	130	4.9	20.4
5	Diarrhea and gastroenteritis of infectious origin (A09)	125	4.7	19.6
6	Septicemia (A40-A41)	87	3.3	13.7
7	Assault (homicide) (*U01,X85-Y09)	80	3.0	12.6
8	Influenza and pneumonia (J09-J18)	73	2.8	11.5
9	Chronic respiratory disease originating in the perinatal period (P27)	38	1.4	6.0
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	35	1.3	5.5
11	Necrotizing enterocolitis of newborn (P77)	30	1.1	4.7
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	18	0.7	*
13	Meningitis (G00,G03)	17	0.6	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	15	0.6	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	12	0.5	*
...	All other causes (Residual)	689	26.2	108.3

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, male				
...	All causes	1,500	100.0	463.0
1	Sudden infant death syndrome (R95)	367	24.5	113.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	185	12.3	57.1
3	Accidents (unintentional injuries) (V01-X59)	183	12.2	56.5
4	Diarrhea and gastroenteritis of infectious origin (A09)	74	4.9	22.8
4	Diseases of the circulatory system (I00-I99)	74	4.9	22.8
6	Septicemia (A40-A41)	46	3.1	14.2
7	Influenza and pneumonia (J09-J18)	43	2.9	13.3
8	Assault (homicide) (*U01,X85-Y09)	37	2.5	11.4
9	Chronic respiratory disease originating in the perinatal period (P27)	22	1.5	6.8
10	Necrotizing enterocolitis of newborn (P77)	16	1.1	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	0.7	*
11	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	10	0.7	*
13	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	9	0.6	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	8	0.5	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	7	0.5	*
15	Meningitis (G00,G03)	7	0.5	*
...	All other causes (Residual)	402	26.8	124.1
Postneonates, black, female				
...	All causes	1,132	100.0	362.3
1	Sudden infant death syndrome (R95)	245	21.6	78.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	167	14.8	53.4
3	Accidents (unintentional injuries) (V01-X59)	136	12.0	43.5
4	Diseases of the circulatory system (I00-I99)	56	4.9	17.9
5	Diarrhea and gastroenteritis of infectious origin (A09)	51	4.5	16.3
6	Assault (homicide) (*U01,X85-Y09)	43	3.8	13.8
7	Septicemia (A40-A41)	41	3.6	13.1
8	Influenza and pneumonia (J09-J18)	30	2.7	9.6
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	25	2.2	8.0
10	Chronic respiratory disease originating in the perinatal period (P27)	16	1.4	*
11	Necrotizing enterocolitis of newborn (P77)	14	1.2	*
12	Meningitis (G00,G03)	10	0.9	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	8	0.7	*
14	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	7	0.6	*
15	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	5	0.4	*
15	Bacterial sepsis of newborn (P36)	5	0.4	*
...	All other causes (Residual)	273	24.1	87.4

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, American Indian, both sexes				
...	All causes	165	100.0	352.9
1	Sudden infant death syndrome (R95)	32	19.4	68.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	17.6	62.0
3	Accidents (unintentional injuries) (V01-X59)	24	14.5	51.3
4	Influenza and pneumonia (J09-J18)	9	5.5	*
5	Diseases of the circulatory system (I00-I99)	7	4.2	*
6	Diarrhea and gastroenteritis of infectious origin (A09)	5	3.0	*
6	Assault (homicide) (*U01,X85-Y09)	5	3.0	*
8	Bronchitis, chronic and unspecified (J40-J42)	4	2.4	*
9	Septicemia (A40-A41)	3	1.8	*
10	Malignant neoplasms (C00-C97)	2	1.2	*
10	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	2	1.2	*
12	Meningitis (G00,G03)	1	0.6	*
12	Diseases of the ear and mastoid process (H60-H93)	1	0.6	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	1	0.6	*
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	1	0.6	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	0.6	*
12	Chronic respiratory disease originating in the perinatal period (P27)	1	0.6	*
12	Atelectasis (P28.0-P28.1)	1	0.6	*
12	Neonatal hemorrhage (P50-P52,P54)	1	0.6	*
12	Complications of medical and surgical care (Y40-Y84)	1	0.6	*
...	All other causes (Residual)	34	20.6	72.7
Postneonates, American Indian, male				
...	All causes	93	100.0	392.9
1	Sudden infant death syndrome (R95)	19	20.4	*
2	Accidents (unintentional injuries) (V01-X59)	15	16.1	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	12.9	*
4	Diseases of the circulatory system (I00-I99)	5	5.4	*
4	Influenza and pneumonia (J09-J18)	5	5.4	*
6	Bronchitis, chronic and unspecified (J40-J42)	4	4.3	*
7	Assault (homicide) (*U01,X85-Y09)	3	3.2	*
8	Diarrhea and gastroenteritis of infectious origin (A09)	2	2.2	*
8	Septicemia (A40-A41)	2	2.2	*
10	Malignant neoplasms (C00-C97)	1	1.1	*
10	Meningitis (G00,G03)	1	1.1	*
10	Diseases of the ear and mastoid process (H60-H93)	1	1.1	*
10	Acute bronchitis and acute bronchiolitis (J20-J21)	1	1.1	*
10	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	1.1	*
10	Chronic respiratory disease originating in the perinatal period (P27)	1	1.1	*
10	Atelectasis (P28.0-P28.1)	1	1.1	*
10	Neonatal hemorrhage (P50-P52,P54)	1	1.1	*
...	All other causes (Residual)	18	19.4	*

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, American Indian, female				
...	All causes	72	100.0	311.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	23.6	*
2	Sudden infant death syndrome (R95)	13	18.1	*
3	Accidents (unintentional injuries) (V01-X59)	9	12.5	*
4	Influenza and pneumonia (J09-J18)	4	5.6	*
5	Diarrhea and gastroenteritis of infectious origin (A09)	3	4.2	*
6	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	2	2.8	*
6	Diseases of the circulatory system (I00-I99)	2	2.8	*
6	Assault (homicide) (*U01,X85-Y09)	2	2.8	*
9	Septicemia (A40-A41)	1	1.4	*
9	Malignant neoplasms (C00-C97)	1	1.4	*
9	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	1	1.4	*
9	Complications of medical and surgical care (Y40-Y84)	1	1.4	*
...	All other causes (Residual)	16	22.2	*
Postneonates, Asian or Pacific Islander, both sexes				
...	All causes	259	100.0	104.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	22.8	23.9
2	Sudden infant death syndrome (R95)	38	14.7	15.4
3	Accidents (unintentional injuries) (V01-X59)	23	8.9	9.3
4	Diseases of the circulatory system (I00-I99)	16	6.2	*
5	Influenza and pneumonia (J09-J18)	12	4.6	*
6	Septicemia (A40-A41)	8	3.1	*
7	Diarrhea and gastroenteritis of infectious origin (A09)	7	2.7	*
8	Malignant neoplasms (C00-C97)	5	1.9	*
8	Chronic respiratory disease originating in the perinatal period (P27)	5	1.9	*
8	Assault (homicide) (*U01,X85-Y09)	5	1.9	*
11	Meningitis (G00,G03)	4	1.5	*
12	Anoxic brain damage, not elsewhere classified (G93.1)	3	1.2	*
12	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	3	1.2	*
12	Atelectasis (P28.0-P28.1)	3	1.2	*
15	Acute bronchitis and acute bronchiolitis (J20-J21)	2	0.8	*
15	Bronchitis, chronic and unspecified (J40-J42)	2	0.8	*
15	Renal failure and other disorders of kidney (N17-N19,N25,N27)	2	0.8	*
...	All other causes (Residual)	62	23.9	25.1

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2010

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision, Second Edition, 2004</i> ), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Asian or Pacific Islander, male				
...	All causes	141	100.0	110.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	26	18.4	20.3
2	Sudden infant death syndrome (R95)	19	13.5	*
3	Accidents (unintentional injuries) (V01-X59)	14	9.9	*
4	Diseases of the circulatory system (I00-I99)	7	5.0	*
4	Influenza and pneumonia (J09-J18)	7	5.0	*
6	Diarrhea and gastroenteritis of infectious origin (A09)	5	3.5	*
7	Meningitis (G00,G03)	3	2.1	*
7	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	3	2.1	*
7	Chronic respiratory disease originating in the perinatal period (P27)	3	2.1	*
7	Atelectasis (P28.0-P28.1)	3	2.1	*
7	Assault (homicide) (*U01,X85-Y09)	3	2.1	*
12	Malignant neoplasms (C00-C97)	2	1.4	*
12	Anoxic brain damage, not elsewhere classified (G93.1)	2	1.4	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	2	1.4	*
12	Bronchitis, chronic and unspecified (J40-J42)	2	1.4	*
...	All other causes (Residual)	40	28.4	31.3
Postneonates, Asian or Pacific Islander, female				
...	All causes	118	100.0	99.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	28.0	27.7
2	Sudden infant death syndrome (R95)	19	16.1	*
3	Diseases of the circulatory system (I00-I99)	9	7.6	*
3	Accidents (unintentional injuries) (V01-X59)	9	7.6	*
5	Septicemia (A40-A41)	7	5.9	*
6	Influenza and pneumonia (J09-J18)	5	4.2	*
7	Malignant neoplasms (C00-C97)	3	2.5	*
8	Diarrhea and gastroenteritis of infectious origin (A09)	2	1.7	*
8	Chronic respiratory disease originating in the perinatal period (P27)	2	1.7	*
8	Assault (homicide) (*U01,X85-Y09)	2	1.7	*
11	Meningitis (G00,G03)	1	0.8	*
11	Anoxic brain damage, not elsewhere classified (G93.1)	1	0.8	*
11	Pneumonitis due to solids and liquids (J69)	1	0.8	*
11	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	0.8	*
11	Complications of medical and surgical care (Y40-Y84)	1	0.8	*
...	All other causes (Residual)	22	18.6	18.5

Source: CDC/NCHS, National Vital Statistics System

1 Rank based on number of deaths

\* Figure does not meet standards of reliability or precision

... Category not applicable