

A revised mortality data file for 2014 was released on April 3, 2017 to include corrections from Tennessee and Massachusetts affecting 125 records previously coded to Accidental discharge of firearms (W32-W34). After the revision, 50% of the changed records are now classified as Assault (homicide) by discharge of firearms (*U01.4 and X93-X95); 42% as Intentional self-harm (suicide) by discharge of firearms (X72-X74); 4% as Discharge of firearms, undetermined intent (Y22-Y24); and the remaining 4% to various other causes. For more information, please see "[Deaths: Final Data for 2014](#)." This table was updated on May 3, 2017 to correct errors. The shaded areas in this table indicate the changes that were made.

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Infants, all races, both sexes				
...	All causes	23,215	100.0	582.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4,746	20.4	119.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	4,173	18.0	104.6
3	Newborn affected by maternal complications of pregnancy (P01)	1,574	6.8	39.5
4	Sudden infant death syndrome (R95)	1,545	6.7	38.7
5	Accidents (unintentional injuries) (V01-X59)	1,160	5.0	29.1
6	Newborn affected by complications of placenta, cord and membranes (P02)	965	4.2	24.2
7	Bacterial sepsis of newborn (P36)	544	2.3	13.6
8	Respiratory distress of newborn (P22)	460	2.0	11.5
9	Diseases of the circulatory system (I00-I99)	444	1.9	11.1
10	Neonatal hemorrhage (P50-P52,P54)	441	1.9	11.1
11	Necrotizing enterocolitis of newborn (P77)	380	1.6	9.5
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	324	1.4	8.1
13	Assault (homicide) (*U01,X85-Y09)	249	1.1	6.2
14	Atelectasis (P28.0-P28.1)	241	1.0	6.0
15	Diarrhea and gastroenteritis of infectious origin (A09)	214	0.9	5.4
...	All other causes (residual)	5,755	24.8	144.3
Infants, all races, male				
...	All causes	12,886	100.0	631.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,444	19.0	119.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,370	18.4	116.1
3	Sudden infant death syndrome (R95)	924	7.2	45.3
4	Newborn affected by maternal complications of pregnancy (P01)	916	7.1	44.9
5	Accidents (unintentional injuries) (V01-X59)	663	5.1	32.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	499	3.9	24.5
7	Bacterial sepsis of newborn (P36)	298	2.3	14.6
8	Neonatal hemorrhage (P50-P52,P54)	271	2.1	13.3
9	Respiratory distress of newborn (P22)	269	2.1	13.2
10	Diseases of the circulatory system (I00-I99)	247	1.9	12.1
11	Necrotizing enterocolitis of newborn (P77)	218	1.7	10.7
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	178	1.4	8.7
13	Atelectasis (P28.0-P28.1)	149	1.2	7.3
14	Assault (homicide) (*U01,X85-Y09)	144	1.1	7.1
15	Diarrhea and gastroenteritis of infectious origin (A09)	112	0.9	5.5
...	All other causes (residual)	3,184	24.7	156.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Infants, all races, female				
...	All causes	10,329	100.0	530.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,302	22.3	118.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,803	17.5	92.6
3	Newborn affected by maternal complications of pregnancy (P01)	658	6.4	33.8
4	Sudden infant death syndrome (R95)	621	6.0	31.9
5	Accidents (unintentional injuries) (V01-X59)	497	4.8	25.5
6	Newborn affected by complications of placenta, cord and membranes (P02)	466	4.5	23.9
7	Bacterial sepsis of newborn (P36)	246	2.4	12.6
8	Diseases of the circulatory system (I00-I99)	197	1.9	10.1
9	Respiratory distress of newborn (P22)	191	1.8	9.8
10	Neonatal hemorrhage (P50-P52,P54)	170	1.6	8.7
11	Necrotizing enterocolitis of newborn (P77)	162	1.6	8.3
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	146	1.4	7.5
13	Assault (homicide) (*U01,X85-Y09)	105	1.0	5.4
14	Diarrhea and gastroenteritis of infectious origin (A09)	102	1.0	5.2
15	Influenza and pneumonia (J09-J18)	97	0.9	5.0
...	All other causes (residual)	2,566	24.8	131.8
Infants, white, both sexes				
...	All causes	14,883	100.0	492.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3,556	23.9	117.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,342	15.7	77.6
3	Sudden infant death syndrome (R95)	997	6.7	33.0
4	Newborn affected by maternal complications of pregnancy (P01)	912	6.1	30.2
5	Accidents (unintentional injuries) (V01-X59)	699	4.7	23.1
6	Newborn affected by complications of placenta, cord and membranes (P02)	605	4.1	20.0
7	Bacterial sepsis of newborn (P36)	345	2.3	11.4
8	Neonatal hemorrhage (P50-P52,P54)	305	2.0	10.1
9	Respiratory distress of newborn (P22)	283	1.9	9.4
10	Diseases of the circulatory system (I00-I99)	265	1.8	8.8
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	220	1.5	7.3
12	Necrotizing enterocolitis of newborn (P77)	218	1.5	7.2
13	Atelectasis (P28.0-P28.1)	152	1.0	5.0
13	Assault (homicide) (*U01,X85-Y09)	152	1.0	5.0
15	Hydrops fetalis not due to hemolytic disease (P83.2)	137	0.9	4.5
...	All other causes (residual)	3,695	24.8	122.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Infants, white, male				
...	All causes	8,297	100.0	536.2
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,846	22.2	119.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,354	16.3	87.5
3	Sudden infant death syndrome (R95)	591	7.1	38.2
4	Newborn affected by maternal complications of pregnancy (P01)	521	6.3	33.7
5	Accidents (unintentional injuries) (V01-X59)	407	4.9	26.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	309	3.7	20.0
7	Bacterial sepsis of newborn (P36)	195	2.4	12.6
8	Neonatal hemorrhage (P50-P52,P54)	193	2.3	12.5
9	Respiratory distress of newborn (P22)	173	2.1	11.2
10	Diseases of the circulatory system (I00-I99)	151	1.8	9.8
11	Necrotizing enterocolitis of newborn (P77)	122	1.5	7.9
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	113	1.4	7.3
13	Atelectasis (P28.0-P28.1)	96	1.2	6.2
14	Assault (homicide) (*U01,X85-Y09)	87	1.0	5.6
15	Pulmonary hemorrhage originating in the perinatal period (P26)	66	0.8	4.3
15	Hydrops fetalis not due to hemolytic disease (P83.2)	66	0.8	4.3
...	All other causes (residual)	2,007	24.2	129.7
Infants, white, female				
...	All causes	6,586	100.0	447.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,710	26.0	116.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	988	15.0	67.1
3	Sudden infant death syndrome (R95)	406	6.2	27.6
4	Newborn affected by maternal complications of pregnancy (P01)	391	5.9	26.6
5	Newborn affected by complications of placenta, cord and membranes (P02)	296	4.5	20.1
6	Accidents (unintentional injuries) (V01-X59)	292	4.4	19.8
7	Bacterial sepsis of newborn (P36)	150	2.3	10.2
8	Diseases of the circulatory system (I00-I99)	114	1.7	7.7
9	Neonatal hemorrhage (P50-P52,P54)	112	1.7	7.6
10	Respiratory distress of newborn (P22)	110	1.7	7.5
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	107	1.6	7.3
12	Necrotizing enterocolitis of newborn (P77)	96	1.5	6.5
13	Hydrops fetalis not due to hemolytic disease (P83.2)	71	1.1	4.8
14	Assault (homicide) (*U01,X85-Y09)	65	1.0	4.4
15	Atelectasis (P28.0-P28.1)	56	0.9	3.8
...	All other causes (residual)	1,622	24.6	110.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Infants, black, both sexes				
...	All causes	7,076	100.0	1,104.7
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,611	22.8	251.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	931	13.2	145.3
3	Newborn affected by maternal complications of pregnancy (P01)	566	8.0	88.4
4	Sudden infant death syndrome (R95)	474	6.7	74.0
5	Accidents (unintentional injuries) (V01-X59)	421	5.9	65.7
6	Newborn affected by complications of placenta, cord and membranes (P02)	306	4.3	47.8
7	Bacterial sepsis of newborn (P36)	174	2.5	27.2
8	Respiratory distress of newborn (P22)	159	2.2	24.8
9	Diseases of the circulatory system (I00-I99)	149	2.1	23.3
10	Necrotizing enterocolitis of newborn (P77)	145	2.0	22.6
11	Neonatal hemorrhage (P50-P52,P54)	117	1.7	18.3
12	Diarrhea and gastroenteritis of infectious origin (A09)	96	1.4	15.0
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	86	1.2	13.4
14	Assault (homicide) (*U01,X85-Y09)	84	1.2	13.1
15	Influenza and pneumonia (J09-J18)	74	1.0	11.6
15	Atelectasis (P28.0-P28.1)	74	1.0	11.6
...	All other causes (residual)	1,609	22.7	251.2
Infants, black, male				
...	All causes	3,900	100.0	1,200.7
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	896	23.0	275.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	471	12.1	145.0
3	Newborn affected by maternal complications of pregnancy (P01)	339	8.7	104.4
4	Sudden infant death syndrome (R95)	277	7.1	85.3
5	Accidents (unintentional injuries) (V01-X59)	235	6.0	72.3
6	Newborn affected by complications of placenta, cord and membranes (P02)	166	4.3	51.1
7	Respiratory distress of newborn (P22)	88	2.3	27.1
7	Bacterial sepsis of newborn (P36)	88	2.3	27.1
9	Necrotizing enterocolitis of newborn (P77)	84	2.2	25.9
10	Diseases of the circulatory system (I00-I99)	78	2.0	24.0
11	Neonatal hemorrhage (P50-P52,P54)	65	1.7	20.0
12	Diarrhea and gastroenteritis of infectious origin (A09)	51	1.3	15.7
12	Intrauterine hypoxia and birth asphyxia (P20-P21)	51	1.3	15.7
14	Assault (homicide) (*U01,X85-Y09)	48	1.2	14.8
15	Atelectasis (P28.0-P28.1)	44	1.1	13.5
...	All other causes (residual)	919	23.6	282.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Infants, black, female				
...	All causes	3,176	100.0	1,005.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	715	22.5	226.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	460	14.5	145.7
3	Newborn affected by maternal complications of pregnancy (P01)	227	7.1	71.9
4	Sudden infant death syndrome (R95)	197	6.2	62.4
5	Accidents (unintentional injuries) (V01-X59)	186	5.9	58.9
6	Newborn affected by complications of placenta, cord and membranes (P02)	140	4.4	44.3
7	Bacterial sepsis of newborn (P36)	86	2.7	27.2
8	Diseases of the circulatory system (I00-I99)	71	2.2	22.5
8	Respiratory distress of newborn (P22)	71	2.2	22.5
10	Necrotizing enterocolitis of newborn (P77)	61	1.9	19.3
11	Neonatal hemorrhage (P50-P52,P54)	52	1.6	16.5
12	Influenza and pneumonia (J09-J18)	46	1.4	14.6
13	Diarrhea and gastroenteritis of infectious origin (A09)	45	1.4	14.3
14	Assault (homicide) (*U01,X85-Y09)	36	1.1	11.4
15	Intrauterine hypoxia and birth asphyxia (P20-P21)	35	1.1	11.1
...	All other causes (residual)	748	23.6	236.9
Infants, American Indian, both sexes				
...	All causes	360	100.0	801.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	20.0	160.3
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	54	15.0	120.2
3	Sudden infant death syndrome (R95)	39	10.8	86.8
4	Accidents (unintentional injuries) (V01-X59)	24	6.7	53.4
5	Newborn affected by maternal complications of pregnancy (P01)	16	4.4	*
6	Newborn affected by complications of placenta, cord and membranes (P02)	15	4.2	*
7	Necrotizing enterocolitis of newborn (P77)	9	2.5	*
8	Assault (homicide) (*U01,X85-Y09)	8	2.2	*
9	Influenza and pneumonia (J09-J18)	5	1.4	*
9	Respiratory distress of newborn (P22)	5	1.4	*
11	Diseases of the circulatory system (I00-I99)	4	1.1	*
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	1.1	*
11	Atelectasis (P28.0-P28.1)	4	1.1	*
11	Bacterial sepsis of newborn (P36)	4	1.1	*
15	Diarrhea and gastroenteritis of infectious origin (A09)	3	0.8	*
15	Septicemia (A40-A41)	3	0.8	*
...	All other causes (residual)	91	25.3	202.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Infants, American Indian, male				
...	All causes	202	100.0	885.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	37	18.3	162.2
2	Sudden infant death syndrome (R95)	30	14.9	131.5
3	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	27	13.4	118.4
4	Accidents (unintentional injuries) (V01-X59)	15	7.4	*
5	Newborn affected by maternal complications of pregnancy (P01)	10	5.0	*
6	Newborn affected by complications of placenta, cord and membranes (P02)	9	4.5	*
7	Assault (homicide) (*U01,X85-Y09)	7	3.5	*
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	2.0	*
8	Respiratory distress of newborn (P22)	4	2.0	*
8	Necrotizing enterocolitis of newborn (P77)	4	2.0	*
11	Diseases of the circulatory system (I00-I99)	3	1.5	*
11	Atelectasis (P28.0-P28.1)	3	1.5	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	2	1.0	*
13	Influenza and pneumonia (J09-J18)	2	1.0	*
13	Newborn affected by other complications of labor and delivery (P03)	2	1.0	*
13	Bacterial sepsis of newborn (P36)	2	1.0	*
13	Neonatal hemorrhage (P50-P52,P54)	2	1.0	*
...	All other causes (residual)	39	19.3	171.0
Infants, American Indian, female				
...	All causes	158	100.0	714.3
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	22.2	158.2
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	27	17.1	122.1
3	Sudden infant death syndrome (R95)	9	5.7	*
3	Accidents (unintentional injuries) (V01-X59)	9	5.7	*
5	Newborn affected by maternal complications of pregnancy (P01)	6	3.8	*
5	Newborn affected by complications of placenta, cord and membranes (P02)	6	3.8	*
7	Necrotizing enterocolitis of newborn (P77)	5	3.2	*
8	Influenza and pneumonia (J09-J18)	3	1.9	*
9	Diarrhea and gastroenteritis of infectious origin (A09)	2	1.3	*
9	Septicemia (A40-A41)	2	1.3	*
9	Bacterial sepsis of newborn (P36)	2	1.3	*
12	Whooping cough (A37)	1	0.6	*
12	Anoxic brain damage, not elsewhere classified (G93.1)	1	0.6	*
12	Diseases of the circulatory system (I00-I99)	1	0.6	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	1	0.6	*
12	Bronchitis, chronic and unspecified (J40-J42)	1	0.6	*
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	1	0.6	*
12	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	0.6	*
12	Newborn affected by noxious influences transmitted via placenta or breast milk (P04)	1	0.6	*
12	Respiratory distress of newborn (P22)	1	0.6	*
12	Neonatal aspiration syndromes (P24)	1	0.6	*
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	1	0.6	*
12	Atelectasis (P28.0-P28.1)	1	0.6	*
12	Assault (homicide) (*U01,X85-Y09)	1	0.6	*
...	All other causes (residual)	39	24.7	176.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Infants, Asian or Pacific Islander, both sexes				
...	All causes	896	100.0	316.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	187	20.9	66.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	166	18.5	58.7
3	Newborn affected by maternal complications of pregnancy (P01)	80	8.9	28.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	39	4.4	13.8
5	Sudden infant death syndrome (R95)	35	3.9	12.4
6	Diseases of the circulatory system (I00-I99)	26	2.9	9.2
7	Bacterial sepsis of newborn (P36)	21	2.3	7.4
8	Neonatal hemorrhage (P50-P52,P54)	17	1.9	*
8	Hydrops fetalis not due to hemolytic disease (P83.2)	17	1.9	*
10	Accidents (unintentional injuries) (V01-X59)	16	1.8	*
11	Intrauterine hypoxia and birth asphyxia (P20-P21)	14	1.6	*
12	Respiratory distress of newborn (P22)	13	1.5	*
13	Atelectasis (P28.0-P28.1)	11	1.2	*
14	Newborn affected by other complications of labor and delivery (P03)	9	1.0	*
15	Septicemia (A40-A41)	8	0.9	*
15	Hematological disorders (P60-P61)	8	0.9	*
15	Necrotizing enterocolitis of newborn (P77)	8	0.9	*
...	All other causes (residual)	221	24.7	78.2
Infants, Asian or Pacific Islander, male				
...	All causes	487	100.0	334.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	93	19.1	63.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	18.5	61.8
3	Newborn affected by maternal complications of pregnancy (P01)	46	9.4	31.6
4	Sudden infant death syndrome (R95)	26	5.3	17.9
5	Diseases of the circulatory system (I00-I99)	15	3.1	*
5	Newborn affected by complications of placenta, cord and membranes (P02)	15	3.1	*
7	Bacterial sepsis of newborn (P36)	13	2.7	*
8	Neonatal hemorrhage (P50-P52,P54)	11	2.3	*
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	10	2.1	*
9	Hydrops fetalis not due to hemolytic disease (P83.2)	10	2.1	*
11	Necrotizing enterocolitis of newborn (P77)	8	1.6	*
12	Atelectasis (P28.0-P28.1)	6	1.2	*
12	Accidents (unintentional injuries) (V01-X59)	6	1.2	*
14	Influenza and pneumonia (J09-J18)	4	0.8	*
14	Respiratory distress of newborn (P22)	4	0.8	*
14	Neonatal aspiration syndromes (P24)	4	0.8	*
...	All other causes (residual)	126	25.9	86.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Infants, Asian or Pacific Islander, female				
...	All causes	409	100.0	298.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	97	23.7	70.8
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	73	17.8	53.3
3	Newborn affected by maternal complications of pregnancy (P01)	34	8.3	24.8
4	Newborn affected by complications of placenta, cord and membranes (P02)	24	5.9	17.5
5	Diseases of the circulatory system (I00-I99)	11	2.7	*
6	Accidents (unintentional injuries) (V01-X59)	10	2.4	*
7	Respiratory distress of newborn (P22)	9	2.2	*
7	Sudden infant death syndrome (R95)	9	2.2	*
9	Bacterial sepsis of newborn (P36)	8	2.0	*
10	Hydrops fetalis not due to hemolytic disease (P83.2)	7	1.7	*
11	Newborn affected by other complications of labor and delivery (P03)	6	1.5	*
11	Neonatal hemorrhage (P50-P52,P54)	6	1.5	*
13	Septicemia (A40-A41)	5	1.2	*
13	Slow fetal growth and fetal malnutrition (P05)	5	1.2	*
13	Atelectasis (P28.0-P28.1)	5	1.2	*
13	Hematological disorders (P60-P61)	5	1.2	*
...	All other causes (residual)	95	23.2	69.3
Neonates, all races, both sexes				
...	All causes	15,720	100.0	394.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	4,078	25.9	102.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3,338	21.2	83.7
3	Newborn affected by maternal complications of pregnancy (P01)	1,566	10.0	39.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	955	6.1	23.9
5	Bacterial sepsis of newborn (P36)	527	3.4	13.2
6	Respiratory distress of newborn (P22)	450	2.9	11.3
7	Neonatal hemorrhage (P50-P52,P54)	428	2.7	10.7
8	Necrotizing enterocolitis of newborn (P77)	341	2.2	8.6
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	308	2.0	7.7
10	Atelectasis (P28.0-P28.1)	225	1.4	5.6
11	Hydrops fetalis not due to hemolytic disease (P83.2)	169	1.1	4.2
12	Pulmonary hemorrhage originating in the perinatal period (P26)	156	1.0	3.9
13	Sudden infant death syndrome (R95)	141	0.9	3.5
14	Accidents (unintentional injuries) (V01-X59)	122	0.8	3.1
15	Newborn affected by other complications of labor and delivery (P03)	113	0.7	2.8
...	All other causes (residual)	2,803	17.8	70.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, all races, male				
...	All causes	8,671	100.0	424.9
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,315	26.7	113.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,738	20.0	85.2
3	Newborn affected by maternal complications of pregnancy (P01)	913	10.5	44.7
4	Newborn affected by complications of placenta, cord and membranes (P02)	496	5.7	24.3
5	Bacterial sepsis of newborn (P36)	286	3.3	14.0
6	Respiratory distress of newborn (P22)	264	3.0	12.9
6	Neonatal hemorrhage (P50-P52,P54)	264	3.0	12.9
8	Necrotizing enterocolitis of newborn (P77)	196	2.3	9.6
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	169	1.9	8.3
10	Atelectasis (P28.0-P28.1)	135	1.6	6.6
11	Pulmonary hemorrhage originating in the perinatal period (P26)	103	1.2	5.0
12	Hydrops fetalis not due to hemolytic disease (P83.2)	85	1.0	4.2
13	Sudden infant death syndrome (R95)	75	0.9	3.7
14	Newborn affected by other complications of labor and delivery (P03)	71	0.8	3.5
15	Accidents (unintentional injuries) (V01-X59)	56	0.6	2.7
...	All other causes (residual)	1,505	17.4	73.7
Neonates, all races, female				
...	All causes	7,049	100.0	362.0
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,763	25.0	90.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,600	22.7	82.2
3	Newborn affected by maternal complications of pregnancy (P01)	653	9.3	33.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	459	6.5	23.6
5	Bacterial sepsis of newborn (P36)	241	3.4	12.4
6	Respiratory distress of newborn (P22)	186	2.6	9.6
7	Neonatal hemorrhage (P50-P52,P54)	164	2.3	8.4
8	Necrotizing enterocolitis of newborn (P77)	145	2.1	7.4
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	139	2.0	7.1
10	Atelectasis (P28.0-P28.1)	90	1.3	4.6
11	Hydrops fetalis not due to hemolytic disease (P83.2)	84	1.2	4.3
12	Sudden infant death syndrome (R95)	66	0.9	3.4
12	Accidents (unintentional injuries) (V01-X59)	66	0.9	3.4
14	Slow fetal growth and fetal malnutrition (P05)	57	0.8	2.9
15	Pulmonary hemorrhage originating in the perinatal period (P26)	53	0.8	2.7
...	All other causes (residual)	1,283	18.2	65.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, both sexes				
...	All causes	10,170	100.0	336.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2,568	25.3	85.0
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2,289	22.5	75.8
3	Newborn affected by maternal complications of pregnancy (P01)	909	8.9	30.1
4	Newborn affected by complications of placenta, cord and membranes (P02)	598	5.9	19.8
5	Bacterial sepsis of newborn (P36)	333	3.3	11.0
6	Neonatal hemorrhage (P50-P52,P54)	297	2.9	9.8
7	Respiratory distress of newborn (P22)	278	2.7	9.2
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	208	2.0	6.9
9	Necrotizing enterocolitis of newborn (P77)	199	2.0	6.6
10	Atelectasis (P28.0-P28.1)	140	1.4	4.6
11	Hydrops fetalis not due to hemolytic disease (P83.2)	129	1.3	4.3
12	Sudden infant death syndrome (R95)	99	1.0	3.3
13	Pulmonary hemorrhage originating in the perinatal period (P26)	96	0.9	3.2
14	Accidents (unintentional injuries) (V01-X59)	88	0.9	2.9
15	Hematological disorders (P60-P61)	70	0.7	2.3
...	All other causes (residual)	1,869	18.4	61.9
Neonates, white, male				
...	All causes	5,624	100.0	363.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,348	24.0	87.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,325	23.6	85.6
3	Newborn affected by maternal complications of pregnancy (P01)	519	9.2	33.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	307	5.5	19.8
5	Neonatal hemorrhage (P50-P52,P54)	189	3.4	12.2
6	Bacterial sepsis of newborn (P36)	186	3.3	12.0
7	Respiratory distress of newborn (P22)	171	3.0	11.1
8	Necrotizing enterocolitis of newborn (P77)	112	2.0	7.2
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	107	1.9	6.9
10	Atelectasis (P28.0-P28.1)	86	1.5	5.6
11	Pulmonary hemorrhage originating in the perinatal period (P26)	66	1.2	4.3
12	Hydrops fetalis not due to hemolytic disease (P83.2)	63	1.1	4.1
13	Sudden infant death syndrome (R95)	55	1.0	3.6
14	Newborn affected by other complications of labor and delivery (P03)	41	0.7	2.6
15	Accidents (unintentional injuries) (V01-X59)	39	0.7	2.5
...	All other causes (residual)	1,010	18.0	65.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, white, female				
...	All causes	4,546	100.0	308.7
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,220	26.8	82.9
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	964	21.2	65.5
3	Newborn affected by maternal complications of pregnancy (P01)	390	8.6	26.5
4	Newborn affected by complications of placenta, cord and membranes (P02)	291	6.4	19.8
5	Bacterial sepsis of newborn (P36)	147	3.2	10.0
6	Neonatal hemorrhage (P50-P52,P54)	108	2.4	7.3
7	Respiratory distress of newborn (P22)	107	2.4	7.3
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	101	2.2	6.9
9	Necrotizing enterocolitis of newborn (P77)	87	1.9	5.9
10	Hydrops fetalis not due to hemolytic disease (P83.2)	66	1.5	4.5
11	Atelectasis (P28.0-P28.1)	54	1.2	3.7
12	Accidents (unintentional injuries) (V01-X59)	49	1.1	3.3
13	Sudden infant death syndrome (R95)	44	1.0	3.0
14	Hematological disorders (P60-P61)	33	0.7	2.2
15	Slow fetal growth and fetal malnutrition (P05)	30	0.7	2.0
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	30	0.7	2.0
15	Pulmonary hemorrhage originating in the perinatal period (P26)	30	0.7	2.0
...	All other causes (residual)	795	17.5	54.0
Neonates, black, both sexes				
...	All causes	4,686	100.0	731.5
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1,574	33.6	245.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	595	12.7	92.9
3	Newborn affected by maternal complications of pregnancy (P01)	563	12.0	87.9
4	Newborn affected by complications of placenta, cord and membranes (P02)	304	6.5	47.5
5	Bacterial sepsis of newborn (P36)	169	3.6	26.4
6	Respiratory distress of newborn (P22)	154	3.3	24.0
7	Necrotizing enterocolitis of newborn (P77)	126	2.7	19.7
8	Neonatal hemorrhage (P50-P52,P54)	112	2.4	17.5
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	82	1.7	12.8
10	Atelectasis (P28.0-P28.1)	72	1.5	11.2
11	Pulmonary hemorrhage originating in the perinatal period (P26)	57	1.2	8.9
12	Slow fetal growth and fetal malnutrition (P05)	39	0.8	6.1
13	Sudden infant death syndrome (R95)	36	0.8	5.6
14	Newborn affected by other complications of labor and delivery (P03)	34	0.7	5.3
15	Accidents (unintentional injuries) (V01-X59)	33	0.7	5.2
...	All other causes (residual)	736	15.7	114.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, black, male				
...	All causes	2,578	100.0	793.7
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	873	33.9	268.8
2	Newborn affected by maternal complications of pregnancy (P01)	338	13.1	104.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	300	11.6	92.4
4	Newborn affected by complications of placenta, cord and membranes (P02)	165	6.4	50.8
5	Respiratory distress of newborn (P22)	85	3.3	26.2
5	Bacterial sepsis of newborn (P36)	85	3.3	26.2
7	Necrotizing enterocolitis of newborn (P77)	73	2.8	22.5
8	Neonatal hemorrhage (P50-P52,P54)	62	2.4	19.1
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	48	1.9	14.8
10	Atelectasis (P28.0-P28.1)	42	1.6	12.9
11	Pulmonary hemorrhage originating in the perinatal period (P26)	35	1.4	10.8
12	Newborn affected by other complications of labor and delivery (P03)	25	1.0	7.7
13	Slow fetal growth and fetal malnutrition (P05)	17	0.7	*
13	Sudden infant death syndrome (R95)	17	0.7	*
13	Accidents (unintentional injuries) (V01-X59)	17	0.7	*
...	All other causes (residual)	396	15.4	121.9
Neonates, black, female				
...	All causes	2,108	100.0	667.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	701	33.3	222.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	295	14.0	93.4
3	Newborn affected by maternal complications of pregnancy (P01)	225	10.7	71.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	139	6.6	44.0
5	Bacterial sepsis of newborn (P36)	84	4.0	26.6
6	Respiratory distress of newborn (P22)	69	3.3	21.9
7	Necrotizing enterocolitis of newborn (P77)	53	2.5	16.8
8	Neonatal hemorrhage (P50-P52,P54)	50	2.4	15.8
9	Intrauterine hypoxia and birth asphyxia (P20-P21)	34	1.6	10.8
10	Atelectasis (P28.0-P28.1)	30	1.4	9.5
11	Slow fetal growth and fetal malnutrition (P05)	22	1.0	7.0
11	Pulmonary hemorrhage originating in the perinatal period (P26)	22	1.0	7.0
13	Sudden infant death syndrome (R95)	19	0.9	*
14	Accidents (unintentional injuries) (V01-X59)	16	0.8	*
15	Congenital pneumonia (P23)	14	0.7	*
15	Hematological disorders (P60-P61)	14	0.7	*
...	All other causes (residual)	321	15.2	101.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, American Indian, both sexes				
...	All causes	202	100.0	449.6
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	52	25.7	115.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	24.8	111.3
3	Newborn affected by maternal complications of pregnancy (P01)	16	7.9	*
4	Newborn affected by complications of placenta, cord and membranes (P02)	15	7.4	*
5	Necrotizing enterocolitis of newborn (P77)	8	4.0	*
6	Respiratory distress of newborn (P22)	5	2.5	*
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	2.0	*
7	Atelectasis (P28.0-P28.1)	4	2.0	*
7	Bacterial sepsis of newborn (P36)	4	2.0	*
10	Sudden infant death syndrome (R95)	3	1.5	*
10	Assault (homicide) (*U01,X85-Y09)	3	1.5	*
12	Diseases of the circulatory system (I00-I99)	2	1.0	*
12	Newborn affected by other complications of labor and delivery (P03)	2	1.0	*
12	Interstitial emphysema and related conditions originating in the perinatal period (P25)	2	1.0	*
12	Neonatal hemorrhage (P50-P52,P54)	2	1.0	*
...	All other causes (residual)	30	14.9	66.8
Neonates, American Indian, male				
...	All causes	113	100.0	495.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	25.7	127.1
2	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	26	23.0	114.0
3	Newborn affected by maternal complications of pregnancy (P01)	10	8.8	*
4	Newborn affected by complications of placenta, cord and membranes (P02)	9	8.0	*
5	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	3.5	*
5	Respiratory distress of newborn (P22)	4	3.5	*
7	Atelectasis (P28.0-P28.1)	3	2.7	*
7	Necrotizing enterocolitis of newborn (P77)	3	2.7	*
7	Assault (homicide) (*U01,X85-Y09)	3	2.7	*
10	Diseases of the circulatory system (I00-I99)	2	1.8	*
10	Newborn affected by other complications of labor and delivery (P03)	2	1.8	*
10	Bacterial sepsis of newborn (P36)	2	1.8	*
10	Neonatal hemorrhage (P50-P52,P54)	2	1.8	*
10	Sudden infant death syndrome (R95)	2	1.8	*
15	Slow fetal growth and fetal malnutrition (P05)	1	0.9	*
15	Interstitial emphysema and related conditions originating in the perinatal period (P25)	1	0.9	*
...	All other causes (residual)	10	8.8	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, American Indian, female				
...	All causes	89	100.0	402.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	26	29.2	117.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	23.6	94.9
3	Newborn affected by maternal complications of pregnancy (P01)	6	6.7	*
3	Newborn affected by complications of placenta, cord and membranes (P02)	6	6.7	*
5	Necrotizing enterocolitis of newborn (P77)	5	5.6	*
6	Bacterial sepsis of newborn (P36)	2	2.2	*
7	Newborn affected by noxious influences transmitted via placenta or breast milk (P04)	1	1.1	*
7	Respiratory distress of newborn (P22)	1	1.1	*
7	Neonatal aspiration syndromes (P24)	1	1.1	*
7	Interstitial emphysema and related conditions originating in the perinatal period (P25)	1	1.1	*
7	Atelectasis (P28.0-P28.1)	1	1.1	*
7	Sudden infant death syndrome (R95)	1	1.1	*
7	Accidents (unintentional injuries) (V01-X59)	1	1.1	*
...	All other causes (residual)	16	18.0	*
Neonates, Asian or Pacific Islander, both sexes				
...	All causes	662	100.0	234.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	163	24.6	57.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	125	18.9	44.2
3	Newborn affected by maternal complications of pregnancy (P01)	78	11.8	27.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	38	5.7	13.4
5	Bacterial sepsis of newborn (P36)	21	3.2	7.4
6	Neonatal hemorrhage (P50-P52,P54)	17	2.6	*
6	Hydrops fetalis not due to hemolytic disease (P83.2)	17	2.6	*
8	Intrauterine hypoxia and birth asphyxia (P20-P21)	14	2.1	*
9	Respiratory distress of newborn (P22)	13	2.0	*
10	Atelectasis (P28.0-P28.1)	9	1.4	*
11	Newborn affected by other complications of labor and delivery (P03)	8	1.2	*
11	Hematological disorders (P60-P61)	8	1.2	*
11	Necrotizing enterocolitis of newborn (P77)	8	1.2	*
14	Slow fetal growth and fetal malnutrition (P05)	7	1.1	*
15	Neonatal aspiration syndromes (P24)	6	0.9	*
...	All other causes (residual)	130	19.6	46.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Neonates, Asian or Pacific Islander, male				
...	All causes	356	100.0	244.4
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	91	25.6	62.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	61	17.1	41.9
3	Newborn affected by maternal complications of pregnancy (P01)	46	12.9	31.6
4	Newborn affected by complications of placenta, cord and membranes (P02)	15	4.2	*
5	Bacterial sepsis of newborn (P36)	13	3.7	*
6	Neonatal hemorrhage (P50-P52,P54)	11	3.1	*
7	Intrauterine hypoxia and birth asphyxia (P20-P21)	10	2.8	*
7	Hydrops fetalis not due to hemolytic disease (P83.2)	10	2.8	*
9	Necrotizing enterocolitis of newborn (P77)	8	2.2	*
10	Respiratory distress of newborn (P22)	4	1.1	*
10	Neonatal aspiration syndromes (P24)	4	1.1	*
10	Atelectasis (P28.0-P28.1)	4	1.1	*
13	Newborn affected by other complications of labor and delivery (P03)	3	0.8	*
13	Interstitial emphysema and related conditions originating in the perinatal period (P25)	3	0.8	*
13	Hematological disorders (P60-P61)	3	0.8	*
...	All other causes (residual)	70	19.7	48.1
Neonates, Asian or Pacific Islander, female				
...	All causes	306	100.0	223.2
1	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	72	23.5	52.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	64	20.9	46.7
3	Newborn affected by maternal complications of pregnancy (P01)	32	10.5	23.3
4	Newborn affected by complications of placenta, cord and membranes (P02)	23	7.5	16.8
5	Respiratory distress of newborn (P22)	9	2.9	*
6	Bacterial sepsis of newborn (P36)	8	2.6	*
7	Hydrops fetalis not due to hemolytic disease (P83.2)	7	2.3	*
8	Neonatal hemorrhage (P50-P52,P54)	6	2.0	*
9	Newborn affected by other complications of labor and delivery (P03)	5	1.6	*
9	Slow fetal growth and fetal malnutrition (P05)	5	1.6	*
9	Atelectasis (P28.0-P28.1)	5	1.6	*
9	Hematological disorders (P60-P61)	5	1.6	*
13	Diseases of the circulatory system (I00-I99)	4	1.3	*
13	Intrauterine hypoxia and birth asphyxia (P20-P21)	4	1.3	*
15	Newborn affected by maternal hypertensive disorders (P00.0)	3	1.0	*
...	All other causes (residual)	54	17.6	39.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, all races, both sexes				
...	All causes	7,495	100.0	187.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1,408	18.8	35.3
2	Sudden infant death syndrome (R95)	1,404	18.7	35.2
3	Accidents (unintentional injuries) (V01-X59)	1,038	13.8	26.0
4	Diseases of the circulatory system (I00-I99)	351	4.7	8.8
5	Assault (homicide) (*U01,X85-Y09)	229	3.1	5.7
6	Diarrhea and gastroenteritis of infectious origin (A09)	210	2.8	5.3
7	Influenza and pneumonia (J09-J18)	183	2.4	4.6
8	Septicemia (A40-A41)	157	2.1	3.9
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	95	1.3	2.4
10	Chronic respiratory disease originating in the perinatal period (P27)	87	1.2	2.2
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	48	0.6	1.2
12	Malignant neoplasms (C00-C97)	41	0.5	1.0
13	Necrotizing enterocolitis of newborn (P77)	39	0.5	1.0
14	Acute bronchitis and acute bronchiolitis (J20-J21)	36	0.5	0.9
15	Meningitis (G00,G03)	31	0.4	0.8
...	All other causes (residual)	2,138	28.5	53.6
Postneonates, all races, male				
...	All causes	4,215	100.0	206.5
1	Sudden infant death syndrome (R95)	849	20.1	41.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	706	16.7	34.6
3	Accidents (unintentional injuries) (V01-X59)	607	14.4	29.7
4	Diseases of the circulatory system (I00-I99)	198	4.7	9.7
5	Assault (homicide) (*U01,X85-Y09)	136	3.2	6.7
6	Diarrhea and gastroenteritis of infectious origin (A09)	109	2.6	5.3
7	Septicemia (A40-A41)	86	2.0	4.2
7	Influenza and pneumonia (J09-J18)	86	2.0	4.2
9	Chronic respiratory disease originating in the perinatal period (P27)	56	1.3	2.7
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	55	1.3	2.7
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	29	0.7	1.4
12	Necrotizing enterocolitis of newborn (P77)	22	0.5	1.1
13	Acute bronchitis and acute bronchiolitis (J20-J21)	20	0.5	1.0
14	Malignant neoplasms (C00-C97)	18	0.4	*
14	Anoxic brain damage, not elsewhere classified (G93.1)	18	0.4	*
...	All other causes (residual)	1,220	28.9	59.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, all races, female				
...	All causes	3,280	100.0	168.4
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	702	21.4	36.0
2	Sudden infant death syndrome (R95)	555	16.9	28.5
3	Accidents (unintentional injuries) (V01-X59)	431	13.1	22.1
4	Diseases of the circulatory system (I00-I99)	153	4.7	7.9
5	Diarrhea and gastroenteritis of infectious origin (A09)	101	3.1	5.2
6	Influenza and pneumonia (J09-J18)	97	3.0	5.0
7	Assault (homicide) (*U01,X85-Y09)	93	2.8	4.8
8	Septicemia (A40-A41)	71	2.2	3.6
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	40	1.2	2.1
10	Chronic respiratory disease originating in the perinatal period (P27)	31	0.9	1.6
11	Malignant neoplasms (C00-C97)	23	0.7	1.2
12	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	19	0.6	*
13	Meningitis (G00,G03)	18	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.5	*
14	Necrotizing enterocolitis of newborn (P77)	17	0.5	*
...	All other causes (residual)	912	27.8	46.8
Postneonates, white, both sexes				
...	All causes	4,713	100.0	156.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	988	21.0	32.7
2	Sudden infant death syndrome (R95)	898	19.1	29.7
3	Accidents (unintentional injuries) (V01-X59)	611	13.0	20.2
4	Diseases of the circulatory system (I00-I99)	203	4.3	6.7
5	Assault (homicide) (*U01,X85-Y09)	140	3.0	4.6
6	Diarrhea and gastroenteritis of infectious origin (A09)	105	2.2	3.5
7	Influenza and pneumonia (J09-J18)	99	2.1	3.3
8	Septicemia (A40-A41)	93	2.0	3.1
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	53	1.1	1.8
10	Chronic respiratory disease originating in the perinatal period (P27)	48	1.0	1.6
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	37	0.8	1.2
12	Malignant neoplasms (C00-C97)	32	0.7	1.1
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	20	0.4	0.7
14	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	19	0.4	*
14	Necrotizing enterocolitis of newborn (P77)	19	0.4	*
...	All other causes (residual)	1,348	28.6	44.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, white, male				
...	All causes	2,673	100.0	172.7
1	Sudden infant death syndrome (R95)	536	20.1	34.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	498	18.6	32.2
3	Accidents (unintentional injuries) (V01-X59)	368	13.8	23.8
4	Diseases of the circulatory system (I00-I99)	118	4.4	7.6
5	Assault (homicide) (*U01,X85-Y09)	83	3.1	5.4
6	Diarrhea and gastroenteritis of infectious origin (A09)	55	2.1	3.6
6	Septicemia (A40-A41)	55	2.1	3.6
8	Influenza and pneumonia (J09-J18)	54	2.0	3.5
9	Chronic respiratory disease originating in the perinatal period (P27)	32	1.2	2.1
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	29	1.1	1.9
11	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	20	0.7	1.3
12	Malignant neoplasms (C00-C97)	18	0.7	*
13	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	10	0.4	*
13	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	0.4	*
13	Atelectasis (P28.0-P28.1)	10	0.4	*
13	Necrotizing enterocolitis of newborn (P77)	10	0.4	*
...	All other causes (residual)	767	28.7	49.6
Postneonates, white, female				
...	All causes	2,040	100.0	138.5
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	490	24.0	33.3
2	Sudden infant death syndrome (R95)	362	17.7	24.6
3	Accidents (unintentional injuries) (V01-X59)	243	11.9	16.5
4	Diseases of the circulatory system (I00-I99)	85	4.2	5.8
5	Assault (homicide) (*U01,X85-Y09)	57	2.8	3.9
6	Diarrhea and gastroenteritis of infectious origin (A09)	50	2.5	3.4
7	Influenza and pneumonia (J09-J18)	45	2.2	3.1
8	Septicemia (A40-A41)	38	1.9	2.6
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	24	1.2	1.6
10	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	17	0.8	*
11	Chronic respiratory disease originating in the perinatal period (P27)	16	0.8	*
12	Malignant neoplasms (C00-C97)	14	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.5	*
14	Acute bronchitis and acute bronchiolitis (J20-J21)	10	0.5	*
14	Renal failure and other disorders of kidney (N17-N19,N25,N27)	10	0.5	*
...	All other causes (residual)	568	27.8	38.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, both sexes				
...	All causes	2,390	100.0	373.1
1	Sudden infant death syndrome (R95)	438	18.3	68.4
2	Accidents (unintentional injuries) (V01-X59)	388	16.2	60.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	336	14.1	52.5
4	Diseases of the circulatory system (I00-I99)	125	5.2	19.5
5	Diarrhea and gastroenteritis of infectious origin (A09)	95	4.0	14.8
6	Assault (homicide) (*U01,X85-Y09)	81	3.4	12.6
7	Influenza and pneumonia (J09-J18)	73	3.1	11.4
8	Septicemia (A40-A41)	53	2.2	8.3
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	37	1.5	5.8
10	Chronic respiratory disease originating in the perinatal period (P27)	35	1.5	5.5
11	Necrotizing enterocolitis of newborn (P77)	19	0.8	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	14	0.6	*
13	Meningitis (G00,G03)	13	0.5	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	10	0.4	*
14	Anoxic brain damage, not elsewhere classified (G93.1)	10	0.4	*
...	All other causes (residual)	663	27.7	103.5
Postneonates, black, male				
...	All causes	1,322	100.0	407.0
1	Sudden infant death syndrome (R95)	260	19.7	80.0
2	Accidents (unintentional injuries) (V01-X59)	218	16.5	67.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	171	12.9	52.6
4	Diseases of the circulatory system (I00-I99)	65	4.9	20.0
5	Diarrhea and gastroenteritis of infectious origin (A09)	50	3.8	15.4
6	Assault (homicide) (*U01,X85-Y09)	47	3.6	14.5
7	Septicemia (A40-A41)	27	2.0	8.3
7	Influenza and pneumonia (J09-J18)	27	2.0	8.3
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	23	1.7	7.1
10	Chronic respiratory disease originating in the perinatal period (P27)	22	1.7	6.8
11	Necrotizing enterocolitis of newborn (P77)	11	0.8	*
12	Acute bronchitis and acute bronchiolitis (J20-J21)	10	0.8	*
13	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	7	0.5	*
13	Anoxic brain damage, not elsewhere classified (G93.1)	7	0.5	*
15	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	5	0.4	*
15	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	5	0.4	*
...	All other causes (residual)	367	27.8	113.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, black, female				
...	All causes	1,068	100.0	338.3
1	Sudden infant death syndrome (R95)	178	16.7	56.4
2	Accidents (unintentional injuries) (V01-X59)	170	15.9	53.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	165	15.4	52.3
4	Diseases of the circulatory system (I00-I99)	60	5.6	19.0
5	Influenza and pneumonia (J09-J18)	46	4.3	14.6
6	Diarrhea and gastroenteritis of infectious origin (A09)	45	4.2	14.3
7	Assault (homicide) (*U01,X85-Y09)	34	3.2	10.8
8	Septicemia (A40-A41)	26	2.4	8.2
9	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	14	1.3	*
10	Chronic respiratory disease originating in the perinatal period (P27)	13	1.2	*
11	Malignant neoplasms (C00-C97)	9	0.8	*
11	Meningitis (G00,G03)	9	0.8	*
13	Necrotizing enterocolitis of newborn (P77)	8	0.7	*
14	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	7	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
...	All other causes (residual)	279	26.1	88.4
Postneonates, American Indian, both sexes				
...	All causes	158	100.0	351.7
1	Sudden infant death syndrome (R95)	36	22.8	80.1
2	Accidents (unintentional injuries) (V01-X59)	23	14.6	51.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	13.9	49.0
4	Influenza and pneumonia (J09-J18)	5	3.2	*
4	Assault (homicide) (*U01,X85-Y09)	5	3.2	*
6	Diarrhea and gastroenteritis of infectious origin (A09)	3	1.9	*
6	Septicemia (A40-A41)	3	1.9	*
8	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	2	1.3	*
8	Diseases of the circulatory system (I00-I99)	2	1.3	*
8	Acute bronchitis and acute bronchiolitis (J20-J21)	2	1.3	*
8	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2	1.3	*
12	Whooping cough (A37)	1	0.6	*
12	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	1	0.6	*
12	Anoxic brain damage, not elsewhere classified (G93.1)	1	0.6	*
12	Acute upper respiratory infections (J00-J06)	1	0.6	*
12	Bronchitis, chronic and unspecified (J40-J42)	1	0.6	*
12	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	1	0.6	*
12	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	0.6	*
12	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	0.6	*
12	Chronic respiratory disease originating in the perinatal period (P27)	1	0.6	*
12	Necrotizing enterocolitis of newborn (P77)	1	0.6	*
...	All other causes (residual)	43	27.2	95.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, American Indian, male				
...	All causes	89	100.0	390.2
1	Sudden infant death syndrome (R95)	28	31.5	122.8
2	Accidents (unintentional injuries) (V01-X59)	15	16.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	9.0	*
4	Assault (homicide) (*U01,X85-Y09)	4	4.5	*
5	Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)	2	2.2	*
5	Influenza and pneumonia (J09-J18)	2	2.2	*
7	Diarrhea and gastroenteritis of infectious origin (A09)	1	1.1	*
7	Septicemia (A40-A41)	1	1.1	*
7	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	1	1.1	*
7	Diseases of the circulatory system (I00-I99)	1	1.1	*
7	Acute upper respiratory infections (J00-J06)	1	1.1	*
7	Acute bronchitis and acute bronchiolitis (J20-J21)	1	1.1	*
7	Renal failure and other disorders of kidney (N17-N19,N25,N27)	1	1.1	*
7	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1	1.1	*
7	Chronic respiratory disease originating in the perinatal period (P27)	1	1.1	*
7	Necrotizing enterocolitis of newborn (P77)	1	1.1	*
...	All other causes (residual)	20	22.5	87.7
Postneonates, American Indian, female				
...	All causes	69	100.0	311.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	20.3	*
2	Sudden infant death syndrome (R95)	8	11.6	*
2	Accidents (unintentional injuries) (V01-X59)	8	11.6	*
4	Influenza and pneumonia (J09-J18)	3	4.3	*
5	Diarrhea and gastroenteritis of infectious origin (A09)	2	2.9	*
5	Septicemia (A40-A41)	2	2.9	*
7	Whooping cough (A37)	1	1.4	*
7	Anoxic brain damage, not elsewhere classified (G93.1)	1	1.4	*
7	Diseases of the circulatory system (I00-I99)	1	1.4	*
7	Acute bronchitis and acute bronchiolitis (J20-J21)	1	1.4	*
7	Bronchitis, chronic and unspecified (J40-J42)	1	1.4	*
7	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	1	1.4	*
7	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	1.4	*
7	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1	1.4	*
7	Assault (homicide) (*U01,X85-Y09)	1	1.4	*
...	All other causes (residual)	23	33.3	104.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Asian or Pacific Islander, both sexes				
...	All causes	234	100.0	82.8
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	62	26.5	21.9
2	Sudden infant death syndrome (R95)	32	13.7	11.3
3	Diseases of the circulatory system (I00-I99)	21	9.0	7.4
4	Accidents (unintentional injuries) (V01-X59)	16	6.8	*
5	Septicemia (A40-A41)	8	3.4	*
6	Diarrhea and gastroenteritis of infectious origin (A09)	7	3.0	*
7	Influenza and pneumonia (J09-J18)	6	2.6	*
8	Anoxic brain damage, not elsewhere classified (G93.1)	3	1.3	*
8	Acute bronchitis and acute bronchiolitis (J20-J21)	3	1.3	*
8	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	3	1.3	*
8	Chronic respiratory disease originating in the perinatal period (P27)	3	1.3	*
8	Assault (homicide) (*U01,X85-Y09)	3	1.3	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	*
13	Meningitis (G00,G03)	2	0.9	*
13	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	2	0.9	*
13	Newborn affected by maternal complications of pregnancy (P01)	2	0.9	*
13	Atelectasis (P28.0-P28.1)	2	0.9	*
...	All other causes (residual)	57	24.4	20.2
Postneonates, Asian or Pacific Islander, male				
...	All causes	131	100.0	89.9
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	22.1	19.9
2	Sudden infant death syndrome (R95)	25	19.1	17.2
3	Diseases of the circulatory system (I00-I99)	14	10.7	*
4	Accidents (unintentional injuries) (V01-X59)	6	4.6	*
5	Diarrhea and gastroenteritis of infectious origin (A09)	3	2.3	*
5	Septicemia (A40-A41)	3	2.3	*
5	Influenza and pneumonia (J09-J18)	3	2.3	*
8	Anoxic brain damage, not elsewhere classified (G93.1)	2	1.5	*
8	Acute bronchitis and acute bronchiolitis (J20-J21)	2	1.5	*
8	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	2	1.5	*
8	Atelectasis (P28.0-P28.1)	2	1.5	*
8	Assault (homicide) (*U01,X85-Y09)	2	1.5	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
13	Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)	1	0.8	*
13	Meningitis (G00,G03)	1	0.8	*
13	Gastritis, duodenitis, and noninfective enteritis and colitis (K29,K50-K55)	1	0.8	*
13	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	0.8	*
13	Chronic respiratory disease originating in the perinatal period (P27)	1	0.8	*
...	All other causes (residual)	32	24.4	22.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014

LCWK7. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by race and sex: United States, 2014

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days through 11 months) deaths per 100,000 live births in specified group. Data for races other than white and black should be interpreted with caution. Deaths based on race of decedent; live births based on race of mother]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), race and sex	Number	Percent of total deaths	Mortality rate
Postneonates, Asian or Pacific Islander, female				
...	All causes	103	100.0	75.1
1	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	32.0	24.1
2	Accidents (unintentional injuries) (V01-X59)	10	9.7	*
3	Diseases of the circulatory system (I00-I99)	7	6.8	*
3	Sudden infant death syndrome (R95)	7	6.8	*
5	Septicemia (A40-A41)	5	4.9	*
6	Diarrhea and gastroenteritis of infectious origin (A09)	4	3.9	*
7	Influenza and pneumonia (J09-J18)	3	2.9	*
8	Newborn affected by maternal complications of pregnancy (P01)	2	1.9	*
8	Chronic respiratory disease originating in the perinatal period (P27)	2	1.9	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
10	Cystic fibrosis (E84)	1	1.0	*
10	Meningitis (G00,G03)	1	1.0	*
10	Anoxic brain damage, not elsewhere classified (G93.1)	1	1.0	*
10	Acute bronchitis and acute bronchiolitis (J20-J21)	1	1.0	*
10	Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56)	1	1.0	*
10	Newborn affected by complications of placenta, cord and membranes (P02)	1	1.0	*
10	Newborn affected by other complications of labor and delivery (P03)	1	1.0	*
10	Disorders related to short gestation and low birth weight, not elsewhere classified (P07)	1	1.0	*
10	Congenital pneumonia (P23)	1	1.0	*
10	Assault (homicide) (*U01,X85-Y09)	1	1.0	*
...	All other causes (residual)	19	18.4	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2014