

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|-------------------------------|---|--------|-------------------------|----------------|
| Infants, Hispanic, both sexes | | | | |
| ... | All causes | 5,424 | 100.0 | 542.6 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 1,348 | 24.9 | 134.9 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 859 | 15.8 | 85.9 |
| 3 | Sudden infant death syndrome (R95) | 326 | 6.0 | 32.6 |
| 4 | Newborn affected by maternal complications of pregnancy (P01) | 322 | 5.9 | 32.2 |
| 5 | Newborn affected by complications of placenta, cord and membranes (P02) | 216 | 4.0 | 21.6 |
| 6 | Accidents (unintentional injuries) (V01-X59) | 187 | 3.4 | 18.7 |
| 7 | Diseases of the circulatory system (I00-I99) | 142 | 2.6 | 14.2 |
| 8 | Bacterial sepsis of newborn (P36) | 119 | 2.2 | 11.9 |
| 9 | Respiratory distress of newborn (P22) | 117 | 2.2 | 11.7 |
| 10 | Neonatal hemorrhage (P50-P52,P54) | 104 | 1.9 | 10.4 |
| 11 | Necrotizing enterocolitis of newborn (P77) | 103 | 1.9 | 10.3 |
| 12 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 71 | 1.3 | 7.1 |
| 13 | Diarrhea and gastroenteritis of infectious origin (A09) | 69 | 1.3 | 6.9 |
| 13 | Atelectasis (P28.0-P28.1) | 69 | 1.3 | 6.9 |
| 15 | Hydrops fetalis not due to hemolytic disease (P83.2) | 61 | 1.1 | 6.1 |
| 15 | Assault (homicide) (*U01,X85-Y09) | 61 | 1.1 | 6.1 |
| ... | All other causes (Residual) | 1,250 | 23.0 | 125.1 |
| Infants, Hispanic, male | | | | |
| ... | All causes | 2,989 | 100.0 | 585.5 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 702 | 23.5 | 137.5 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 485 | 16.2 | 95.0 |
| 3 | Sudden infant death syndrome (R95) | 183 | 6.1 | 35.8 |
| 4 | Newborn affected by maternal complications of pregnancy (P01) | 158 | 5.3 | 31.0 |
| 5 | Newborn affected by complications of placenta, cord and membranes (P02) | 125 | 4.2 | 24.5 |
| 6 | Accidents (unintentional injuries) (V01-X59) | 112 | 3.7 | 21.9 |
| 7 | Respiratory distress of newborn (P22) | 77 | 2.6 | 15.1 |
| 8 | Bacterial sepsis of newborn (P36) | 68 | 2.3 | 13.3 |
| 9 | Diseases of the circulatory system (I00-I99) | 67 | 2.2 | 13.1 |
| 10 | Necrotizing enterocolitis of newborn (P77) | 63 | 2.1 | 12.3 |
| 11 | Neonatal hemorrhage (P50-P52,P54) | 60 | 2.0 | 11.8 |
| 12 | Diarrhea and gastroenteritis of infectious origin (A09) | 47 | 1.6 | 9.2 |
| 13 | Atelectasis (P28.0-P28.1) | 45 | 1.5 | 8.8 |
| 14 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 43 | 1.4 | 8.4 |
| 15 | Influenza and pneumonia (J09-J18) | 37 | 1.2 | 7.2 |
| ... | All other causes (Residual) | 717 | 24.0 | 140.5 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|--|---|--------|-------------------------|----------------|
| Infants, Hispanic, female | | | | |
| ... | All causes | 2,435 | 100.0 | 497.9 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 646 | 26.5 | 132.1 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 374 | 15.4 | 76.5 |
| 3 | Newborn affected by maternal complications of pregnancy (P01) | 164 | 6.7 | 33.5 |
| 4 | Sudden infant death syndrome (R95) | 143 | 5.9 | 29.2 |
| 5 | Newborn affected by complications of placenta, cord and membranes (P02) | 91 | 3.7 | 18.6 |
| 6 | Diseases of the circulatory system (I00-I99) | 75 | 3.1 | 15.3 |
| 6 | Accidents (unintentional injuries) (V01-X59) | 75 | 3.1 | 15.3 |
| 8 | Bacterial sepsis of newborn (P36) | 51 | 2.1 | 10.4 |
| 9 | Neonatal hemorrhage (P50-P52,P54) | 44 | 1.8 | 9.0 |
| 10 | Respiratory distress of newborn (P22) | 40 | 1.6 | 8.2 |
| 10 | Necrotizing enterocolitis of newborn (P77) | 40 | 1.6 | 8.2 |
| 12 | Hydrops fetalis not due to hemolytic disease (P83.2) | 32 | 1.3 | 6.5 |
| 13 | Assault (homicide) (*U01,X85-Y09) | 31 | 1.3 | 6.3 |
| 14 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 28 | 1.1 | 5.7 |
| 15 | Atelectasis (P28.0-P28.1) | 24 | 1.0 | 4.9 |
| ... | All other causes (Residual) | 577 | 23.7 | 118.0 |
| Infants, Non-Hispanic, white, both sexes | | | | |
| ... | All causes | 11,608 | 100.0 | 524.6 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 2,621 | 22.6 | 118.5 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 1,660 | 14.3 | 75.0 |
| 3 | Sudden infant death syndrome (R95) | 1,121 | 9.7 | 50.7 |
| 4 | Newborn affected by maternal complications of pregnancy (P01) | 664 | 5.7 | 30.0 |
| 5 | Accidents (unintentional injuries) (V01-X59) | 562 | 4.8 | 25.4 |
| 6 | Newborn affected by complications of placenta, cord and membranes (P02) | 482 | 4.2 | 21.8 |
| 7 | Bacterial sepsis of newborn (P36) | 265 | 2.3 | 12.0 |
| 8 | Diseases of the circulatory system (I00-I99) | 251 | 2.2 | 11.3 |
| 9 | Respiratory distress of newborn (P22) | 250 | 2.2 | 11.3 |
| 10 | Neonatal hemorrhage (P50-P52,P54) | 249 | 2.1 | 11.3 |
| 11 | Necrotizing enterocolitis of newborn (P77) | 192 | 1.7 | 8.7 |
| 12 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 154 | 1.3 | 7.0 |
| 13 | Assault (homicide) (*U01,X85-Y09) | 122 | 1.1 | 5.5 |
| 14 | Atelectasis (P28.0-P28.1) | 119 | 1.0 | 5.4 |
| 15 | Diarrhea and gastroenteritis of infectious origin (A09) | 107 | 0.9 | 4.8 |
| ... | All other causes (Residual) | 2,789 | 24.0 | 126.1 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|--------------------------------------|---|--------|-------------------------|----------------|
| Infants, Non-Hispanic, white, male | | | | |
| ... | All causes | 6,530 | 100.0 | 575.5 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 1,373 | 21.0 | 121.0 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 924 | 14.2 | 81.4 |
| 3 | Sudden infant death syndrome (R95) | 675 | 10.3 | 59.5 |
| 4 | Newborn affected by maternal complications of pregnancy (P01) | 376 | 5.8 | 33.1 |
| 5 | Accidents (unintentional injuries) (V01-X59) | 334 | 5.1 | 29.4 |
| 6 | Newborn affected by complications of placenta, cord and membranes (P02) | 251 | 3.8 | 22.1 |
| 7 | Neonatal hemorrhage (P50-P52,P54) | 157 | 2.4 | 13.8 |
| 8 | Bacterial sepsis of newborn (P36) | 149 | 2.3 | 13.1 |
| 9 | Respiratory distress of newborn (P22) | 145 | 2.2 | 12.8 |
| 10 | Diseases of the circulatory system (I00-I99) | 142 | 2.2 | 12.5 |
| 11 | Necrotizing enterocolitis of newborn (P77) | 107 | 1.6 | 9.4 |
| 12 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 88 | 1.3 | 7.8 |
| 13 | Assault (homicide) (*U01,X85-Y09) | 79 | 1.2 | 7.0 |
| 14 | Atelectasis (P28.0-P28.1) | 67 | 1.0 | 5.9 |
| 15 | Influenza and pneumonia (J09-J18) | 58 | 0.9 | 5.1 |
| ... | All other causes (Residual) | 1,605 | 24.6 | 141.5 |
| Infants, Non-Hispanic, white, female | | | | |
| ... | All causes | 5,078 | 100.0 | 471.1 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 1,248 | 24.6 | 115.8 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 736 | 14.5 | 68.3 |
| 3 | Sudden infant death syndrome (R95) | 446 | 8.8 | 41.4 |
| 4 | Newborn affected by maternal complications of pregnancy (P01) | 288 | 5.7 | 26.7 |
| 5 | Newborn affected by complications of placenta, cord and membranes (P02) | 231 | 4.5 | 21.4 |
| 6 | Accidents (unintentional injuries) (V01-X59) | 228 | 4.5 | 21.2 |
| 7 | Bacterial sepsis of newborn (P36) | 116 | 2.3 | 10.8 |
| 8 | Diseases of the circulatory system (I00-I99) | 109 | 2.1 | 10.1 |
| 9 | Respiratory distress of newborn (P22) | 105 | 2.1 | 9.7 |
| 10 | Neonatal hemorrhage (P50-P52,P54) | 92 | 1.8 | 8.5 |
| 11 | Necrotizing enterocolitis of newborn (P77) | 85 | 1.7 | 7.9 |
| 12 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 66 | 1.3 | 6.1 |
| 13 | Atelectasis (P28.0-P28.1) | 52 | 1.0 | 4.8 |
| 14 | Diarrhea and gastroenteritis of infectious origin (A09) | 50 | 1.0 | 4.6 |
| 15 | Assault (homicide) (*U01,X85-Y09) | 43 | 0.8 | 4.0 |
| ... | All other causes (Residual) | 1,183 | 23.3 | 109.8 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|--|---|--------|-------------------------|----------------|
| Infants, Non-Hispanic, black, both sexes | | | | |
| ... | All causes | 7,967 | 100.0 | 1,307.0 |
| 1 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 1,781 | 22.4 | 292.2 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 1,028 | 12.9 | 168.6 |
| 3 | Sudden infant death syndrome (R95) | 665 | 8.3 | 109.1 |
| 4 | Newborn affected by maternal complications of pregnancy (P01) | 531 | 6.7 | 87.1 |
| 5 | Accidents (unintentional injuries) (V01-X59) | 387 | 4.9 | 63.5 |
| 6 | Newborn affected by complications of placenta, cord and membranes (P02) | 308 | 3.9 | 50.5 |
| 7 | Bacterial sepsis of newborn (P36) | 233 | 2.9 | 38.2 |
| 8 | Respiratory distress of newborn (P22) | 204 | 2.6 | 33.5 |
| 9 | Necrotizing enterocolitis of newborn (P77) | 167 | 2.1 | 27.4 |
| 10 | Diseases of the circulatory system (I00-I99) | 165 | 2.1 | 27.1 |
| 11 | Diarrhea and gastroenteritis of infectious origin (A09) | 140 | 1.8 | 23.0 |
| 12 | Neonatal hemorrhage (P50-P52,P54) | 136 | 1.7 | 22.3 |
| 13 | Assault (homicide) (*U01,X85-Y09) | 112 | 1.4 | 18.4 |
| 14 | Septicemia (A40-A41) | 87 | 1.1 | 14.3 |
| 15 | Chronic respiratory disease originating in the perinatal period (P27) | 84 | 1.1 | 13.8 |
| ... | All other causes (Residual) | 1,939 | 24.3 | 318.1 |
| Infants, Non-Hispanic, black, male | | | | |
| ... | All causes | 4,522 | 100.0 | 1,459.9 |
| 1 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 1,005 | 22.2 | 324.5 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 530 | 11.7 | 171.1 |
| 3 | Sudden infant death syndrome (R95) | 381 | 8.4 | 123.0 |
| 4 | Newborn affected by maternal complications of pregnancy (P01) | 324 | 7.2 | 104.6 |
| 5 | Accidents (unintentional injuries) (V01-X59) | 223 | 4.9 | 72.0 |
| 6 | Newborn affected by complications of placenta, cord and membranes (P02) | 162 | 3.6 | 52.3 |
| 7 | Bacterial sepsis of newborn (P36) | 134 | 3.0 | 43.3 |
| 8 | Respiratory distress of newborn (P22) | 115 | 2.5 | 37.1 |
| 9 | Diseases of the circulatory system (I00-I99) | 103 | 2.3 | 33.3 |
| 10 | Necrotizing enterocolitis of newborn (P77) | 93 | 2.1 | 30.0 |
| 11 | Diarrhea and gastroenteritis of infectious origin (A09) | 85 | 1.9 | 27.4 |
| 12 | Neonatal hemorrhage (P50-P52,P54) | 83 | 1.8 | 26.8 |
| 13 | Assault (homicide) (*U01,X85-Y09) | 62 | 1.4 | 20.0 |
| 14 | Chronic respiratory disease originating in the perinatal period (P27) | 50 | 1.1 | 16.1 |
| 15 | Septicemia (A40-A41) | 47 | 1.0 | 15.2 |
| ... | All other causes (Residual) | 1,125 | 24.9 | 363.2 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|--------------------------------------|---|--------|-------------------------|----------------|
| Infants, Non-Hispanic, black, female | | | | |
| ... | All causes | 3,445 | 100.0 | 1,149.0 |
| 1 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 776 | 22.5 | 258.8 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 498 | 14.5 | 166.1 |
| 3 | Sudden infant death syndrome (R95) | 284 | 8.2 | 94.7 |
| 4 | Newborn affected by maternal complications of pregnancy (P01) | 207 | 6.0 | 69.0 |
| 5 | Accidents (unintentional injuries) (V01-X59) | 164 | 4.8 | 54.7 |
| 6 | Newborn affected by complications of placenta, cord and membranes (P02) | 146 | 4.2 | 48.7 |
| 7 | Bacterial sepsis of newborn (P36) | 99 | 2.9 | 33.0 |
| 8 | Respiratory distress of newborn (P22) | 89 | 2.6 | 29.7 |
| 9 | Necrotizing enterocolitis of newborn (P77) | 74 | 2.1 | 24.7 |
| 10 | Diseases of the circulatory system (I00-I99) | 62 | 1.8 | 20.7 |
| 11 | Diarrhea and gastroenteritis of infectious origin (A09) | 55 | 1.6 | 18.3 |
| 12 | Neonatal hemorrhage (P50-P52,P54) | 53 | 1.5 | 17.7 |
| 13 | Assault (homicide) (*U01,X85-Y09) | 50 | 1.5 | 16.7 |
| 14 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 41 | 1.2 | 13.7 |
| 15 | Septicemia (A40-A41) | 40 | 1.2 | 13.3 |
| ... | All other causes (Residual) | 807 | 23.4 | 269.1 |
| Neonates, Hispanic, both sexes | | | | |
| ... | All causes | 3,629 | 100.0 | 363.1 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 931 | 25.7 | 93.1 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 842 | 23.2 | 84.2 |
| 3 | Newborn affected by maternal complications of pregnancy (P01) | 322 | 8.9 | 32.2 |
| 4 | Newborn affected by complications of placenta, cord and membranes (P02) | 214 | 5.9 | 21.4 |
| 5 | Respiratory distress of newborn (P22) | 115 | 3.2 | 11.5 |
| 5 | Bacterial sepsis of newborn (P36) | 115 | 3.2 | 11.5 |
| 7 | Neonatal hemorrhage (P50-P52,P54) | 102 | 2.8 | 10.2 |
| 8 | Necrotizing enterocolitis of newborn (P77) | 87 | 2.4 | 8.7 |
| 9 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 67 | 1.8 | 6.7 |
| 9 | Atelectasis (P28.0-P28.1) | 67 | 1.8 | 6.7 |
| 11 | Hydrops fetalis not due to hemolytic disease (P83.2) | 58 | 1.6 | 5.8 |
| 12 | Pulmonary hemorrhage originating in the perinatal period (P26) | 34 | 0.9 | 3.4 |
| 13 | Interstitial emphysema and related conditions originating in the perinatal period (P25) | 29 | 0.8 | 2.9 |
| 14 | Diseases of the circulatory system (I00-I99) | 27 | 0.7 | 2.7 |
| 15 | Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9) | 26 | 0.7 | 2.6 |
| ... | All other causes (Residual) | 593 | 16.3 | 59.3 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|----------------------------|---|--------|-------------------------|----------------|
| Neonates, Hispanic, male | | | | |
| ... | All causes | 1,988 | 100.0 | 389.4 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 487 | 24.5 | 95.4 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 478 | 24.0 | 93.6 |
| 3 | Newborn affected by maternal complications of pregnancy (P01) | 158 | 7.9 | 31.0 |
| 4 | Newborn affected by complications of placenta, cord and membranes (P02) | 123 | 6.2 | 24.1 |
| 5 | Respiratory distress of newborn (P22) | 75 | 3.8 | 14.7 |
| 6 | Bacterial sepsis of newborn (P36) | 66 | 3.3 | 12.9 |
| 7 | Neonatal hemorrhage (P50-P52,P54) | 58 | 2.9 | 11.4 |
| 8 | Necrotizing enterocolitis of newborn (P77) | 57 | 2.9 | 11.2 |
| 9 | Atelectasis (P28.0-P28.1) | 43 | 2.2 | 8.4 |
| 10 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 41 | 2.1 | 8.0 |
| 11 | Hydrops fetalis not due to hemolytic disease (P83.2) | 28 | 1.4 | 5.5 |
| 12 | Interstitial emphysema and related conditions originating in the perinatal period (P25) | 18 | 0.9 | * |
| 12 | Pulmonary hemorrhage originating in the perinatal period (P26) | 18 | 0.9 | * |
| 14 | Diseases of the circulatory system (I00-I99) | 17 | 0.9 | * |
| 15 | Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9) | 15 | 0.8 | * |
| ... | All other causes (Residual) | 306 | 15.4 | 59.9 |
| Neonates, Hispanic, female | | | | |
| ... | All causes | 1,641 | 100.0 | 335.5 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 444 | 27.1 | 90.8 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 364 | 22.2 | 74.4 |
| 3 | Newborn affected by maternal complications of pregnancy (P01) | 164 | 10.0 | 33.5 |
| 4 | Newborn affected by complications of placenta, cord and membranes (P02) | 91 | 5.5 | 18.6 |
| 5 | Bacterial sepsis of newborn (P36) | 49 | 3.0 | 10.0 |
| 6 | Neonatal hemorrhage (P50-P52,P54) | 44 | 2.7 | 9.0 |
| 7 | Respiratory distress of newborn (P22) | 40 | 2.4 | 8.2 |
| 8 | Necrotizing enterocolitis of newborn (P77) | 30 | 1.8 | 6.1 |
| 8 | Hydrops fetalis not due to hemolytic disease (P83.2) | 30 | 1.8 | 6.1 |
| 10 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 26 | 1.6 | 5.3 |
| 11 | Atelectasis (P28.0-P28.1) | 24 | 1.5 | 4.9 |
| 12 | Pulmonary hemorrhage originating in the perinatal period (P26) | 16 | 1.0 | * |
| 13 | Newborn affected by other maternal conditions which may be unrelated to present pregnancy (P00.1-P00.9) | 11 | 0.7 | * |
| 13 | Interstitial emphysema and related conditions originating in the perinatal period (P25) | 11 | 0.7 | * |
| 13 | Accidents (unintentional injuries) (V01-X59) | 11 | 0.7 | * |
| ... | All other causes (Residual) | 286 | 17.4 | 58.5 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|---|---|--------|-------------------------|----------------|
| Neonates, Non-Hispanic, white, both sexes | | | | |
| ... | All causes | 7,562 | 100.0 | 341.8 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 1,920 | 25.4 | 86.8 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 1,630 | 21.6 | 73.7 |
| 3 | Newborn affected by maternal complications of pregnancy (P01) | 658 | 8.7 | 29.7 |
| 4 | Newborn affected by complications of placenta, cord and membranes (P02) | 479 | 6.3 | 21.6 |
| 5 | Bacterial sepsis of newborn (P36) | 256 | 3.4 | 11.6 |
| 6 | Neonatal hemorrhage (P50-P52,P54) | 246 | 3.3 | 11.1 |
| 7 | Respiratory distress of newborn (P22) | 245 | 3.2 | 11.1 |
| 8 | Necrotizing enterocolitis of newborn (P77) | 170 | 2.2 | 7.7 |
| 9 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 145 | 1.9 | 6.6 |
| 10 | Atelectasis (P28.0-P28.1) | 112 | 1.5 | 5.1 |
| 11 | Sudden infant death syndrome (R95) | 105 | 1.4 | 4.7 |
| 12 | Hydrops fetalis not due to hemolytic disease (P83.2) | 88 | 1.2 | 4.0 |
| 13 | Accidents (unintentional injuries) (V01-X59) | 67 | 0.9 | 3.0 |
| 14 | Pulmonary hemorrhage originating in the perinatal period (P26) | 59 | 0.8 | 2.7 |
| 15 | Diseases of the circulatory system (I00-I99) | 57 | 0.8 | 2.6 |
| ... | All other causes (Residual) | 1,325 | 17.5 | 59.9 |
| Neonates, Non-Hispanic, white, male | | | | |
| ... | All causes | 4,180 | 100.0 | 368.4 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 1,028 | 24.6 | 90.6 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 904 | 21.6 | 79.7 |
| 3 | Newborn affected by maternal complications of pregnancy (P01) | 371 | 8.9 | 32.7 |
| 4 | Newborn affected by complications of placenta, cord and membranes (P02) | 249 | 6.0 | 21.9 |
| 5 | Neonatal hemorrhage (P50-P52,P54) | 156 | 3.7 | 13.7 |
| 6 | Bacterial sepsis of newborn (P36) | 145 | 3.5 | 12.8 |
| 7 | Respiratory distress of newborn (P22) | 143 | 3.4 | 12.6 |
| 8 | Necrotizing enterocolitis of newborn (P77) | 93 | 2.2 | 8.2 |
| 9 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 80 | 1.9 | 7.1 |
| 10 | Atelectasis (P28.0-P28.1) | 62 | 1.5 | 5.5 |
| 11 | Sudden infant death syndrome (R95) | 58 | 1.4 | 5.1 |
| 12 | Hydrops fetalis not due to hemolytic disease (P83.2) | 49 | 1.2 | 4.3 |
| 13 | Pulmonary hemorrhage originating in the perinatal period (P26) | 38 | 0.9 | 3.3 |
| 14 | Diseases of the circulatory system (I00-I99) | 35 | 0.8 | 3.1 |
| 15 | Interstitial emphysema and related conditions originating in the perinatal period (P25) | 30 | 0.7 | 2.6 |
| ... | All other causes (Residual) | 739 | 17.7 | 65.1 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|---|---|--------|-------------------------|----------------|
| Neonates, Non-Hispanic, white, female | | | | |
| ... | All causes | 3,382 | 100.0 | 313.8 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 892 | 26.4 | 82.8 |
| 2 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 726 | 21.5 | 67.4 |
| 3 | Newborn affected by maternal complications of pregnancy (P01) | 287 | 8.5 | 26.6 |
| 4 | Newborn affected by complications of placenta, cord and membranes (P02) | 230 | 6.8 | 21.3 |
| 5 | Bacterial sepsis of newborn (P36) | 111 | 3.3 | 10.3 |
| 6 | Respiratory distress of newborn (P22) | 102 | 3.0 | 9.5 |
| 7 | Neonatal hemorrhage (P50-P52,P54) | 90 | 2.7 | 8.3 |
| 8 | Necrotizing enterocolitis of newborn (P77) | 77 | 2.3 | 7.1 |
| 9 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 65 | 1.9 | 6.0 |
| 10 | Atelectasis (P28.0-P28.1) | 50 | 1.5 | 4.6 |
| 11 | Sudden infant death syndrome (R95) | 47 | 1.4 | 4.4 |
| 12 | Accidents (unintentional injuries) (V01-X59) | 42 | 1.2 | 3.9 |
| 13 | Hydrops fetalis not due to hemolytic disease (P83.2) | 39 | 1.2 | 3.6 |
| 14 | Newborn affected by other complications of labor and delivery (P03) | 27 | 0.8 | 2.5 |
| 15 | Hematological disorders (P60-P61) | 26 | 0.8 | 2.4 |
| ... | All other causes (Residual) | 571 | 16.9 | 53.0 |
| Neonates, Non-Hispanic, black, both sexes | | | | |
| ... | All causes | 5,131 | 100.0 | 841.7 |
| 1 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 1,727 | 33.7 | 283.3 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 676 | 13.2 | 110.9 |
| 3 | Newborn affected by maternal complications of pregnancy (P01) | 528 | 10.3 | 86.6 |
| 4 | Newborn affected by complications of placenta, cord and membranes (P02) | 305 | 5.9 | 50.0 |
| 5 | Bacterial sepsis of newborn (P36) | 219 | 4.3 | 35.9 |
| 6 | Respiratory distress of newborn (P22) | 203 | 4.0 | 33.3 |
| 7 | Necrotizing enterocolitis of newborn (P77) | 144 | 2.8 | 23.6 |
| 8 | Neonatal hemorrhage (P50-P52,P54) | 134 | 2.6 | 22.0 |
| 9 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 73 | 1.4 | 12.0 |
| 10 | Atelectasis (P28.0-P28.1) | 71 | 1.4 | 11.6 |
| 11 | Pulmonary hemorrhage originating in the perinatal period (P26) | 59 | 1.1 | 9.7 |
| 11 | Sudden infant death syndrome (R95) | 59 | 1.1 | 9.7 |
| 13 | Newborn affected by other complications of labor and delivery (P03) | 37 | 0.7 | 6.1 |
| 14 | Newborn affected by maternal hypertensive disorders (P00.0) | 35 | 0.7 | 5.7 |
| 14 | Hydrops fetalis not due to hemolytic disease (P83.2) | 35 | 0.7 | 5.7 |
| ... | All other causes (Residual) | 826 | 16.1 | 135.5 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|---------------------------------------|---|--------|-------------------------|----------------|
| Neonates, Non-Hispanic, black, male | | | | |
| ... | All causes | 2,893 | 100.0 | 934.0 |
| 1 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 978 | 33.8 | 315.7 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 366 | 12.7 | 118.2 |
| 3 | Newborn affected by maternal complications of pregnancy (P01) | 322 | 11.1 | 104.0 |
| 4 | Newborn affected by complications of placenta, cord and membranes (P02) | 160 | 5.5 | 51.7 |
| 5 | Bacterial sepsis of newborn (P36) | 126 | 4.4 | 40.7 |
| 6 | Respiratory distress of newborn (P22) | 115 | 4.0 | 37.1 |
| 7 | Neonatal hemorrhage (P50-P52,P54) | 82 | 2.8 | 26.5 |
| 8 | Necrotizing enterocolitis of newborn (P77) | 78 | 2.7 | 25.2 |
| 9 | Pulmonary hemorrhage originating in the perinatal period (P26) | 42 | 1.5 | 13.6 |
| 10 | Atelectasis (P28.0-P28.1) | 35 | 1.2 | 11.3 |
| 11 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 33 | 1.1 | 10.7 |
| 12 | Sudden infant death syndrome (R95) | 30 | 1.0 | 9.7 |
| 13 | Diseases of the circulatory system (I00-I99) | 22 | 0.8 | 7.1 |
| 14 | Newborn affected by other complications of labor and delivery (P03) | 21 | 0.7 | 6.8 |
| 14 | Hydrops fetalis not due to hemolytic disease (P83.2) | 21 | 0.7 | 6.8 |
| ... | All other causes (Residual) | 462 | 16.0 | 149.2 |
| Neonates, Non-Hispanic, black, female | | | | |
| ... | All causes | 2,238 | 100.0 | 746.4 |
| 1 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 749 | 33.5 | 249.8 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 310 | 13.9 | 103.4 |
| 3 | Newborn affected by maternal complications of pregnancy (P01) | 206 | 9.2 | 68.7 |
| 4 | Newborn affected by complications of placenta, cord and membranes (P02) | 145 | 6.5 | 48.4 |
| 5 | Bacterial sepsis of newborn (P36) | 93 | 4.2 | 31.0 |
| 6 | Respiratory distress of newborn (P22) | 88 | 3.9 | 29.3 |
| 7 | Necrotizing enterocolitis of newborn (P77) | 66 | 2.9 | 22.0 |
| 8 | Neonatal hemorrhage (P50-P52,P54) | 52 | 2.3 | 17.3 |
| 9 | Intrauterine hypoxia and birth asphyxia (P20-P21) | 40 | 1.8 | 13.3 |
| 10 | Atelectasis (P28.0-P28.1) | 36 | 1.6 | 12.0 |
| 11 | Sudden infant death syndrome (R95) | 29 | 1.3 | 9.7 |
| 12 | Newborn affected by maternal hypertensive disorders (P00.0) | 23 | 1.0 | 7.7 |
| 13 | Pulmonary hemorrhage originating in the perinatal period (P26) | 17 | 0.8 | * |
| 13 | Accidents (unintentional injuries) (V01-X59) | 17 | 0.8 | * |
| 15 | Newborn affected by other complications of labor and delivery (P03) | 16 | 0.7 | * |
| ... | All other causes (Residual) | 351 | 15.7 | 117.1 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|------------------------------------|---|--------|-------------------------|----------------|
| Postneonates, Hispanic, both sexes | | | | |
| ... | All causes | 1,795 | 100.0 | 179.6 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 417 | 23.2 | 41.7 |
| 2 | Sudden infant death syndrome (R95) | 303 | 16.9 | 30.3 |
| 3 | Accidents (unintentional injuries) (V01-X59) | 166 | 9.2 | 16.6 |
| 4 | Diseases of the circulatory system (I00-I99) | 115 | 6.4 | 11.5 |
| 5 | Diarrhea and gastroenteritis of infectious origin (A09) | 68 | 3.8 | 6.8 |
| 6 | Assault (homicide) (*U01,X85-Y09) | 57 | 3.2 | 5.7 |
| 7 | Influenza and pneumonia (J09-J18) | 52 | 2.9 | 5.2 |
| 8 | Septicemia (A40-A41) | 49 | 2.7 | 4.9 |
| 9 | Chronic respiratory disease originating in the perinatal period (P27) | 24 | 1.3 | 2.4 |
| 10 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 17 | 0.9 | * |
| 11 | Necrotizing enterocolitis of newborn (P77) | 16 | 0.9 | * |
| 12 | Malignant neoplasms (C00-C97) | 15 | 0.8 | * |
| 13 | Renal failure and other disorders of kidney (N17-N19,N25,N27) | 14 | 0.8 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 9 | 0.5 | * |
| 14 | Meningitis (G00,G03) | 9 | 0.5 | * |
| ... | All other causes (Residual) | 464 | 25.8 | 46.4 |
| Postneonates, Hispanic, male | | | | |
| ... | All causes | 1,001 | 100.0 | 196.1 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 215 | 21.5 | 42.1 |
| 2 | Sudden infant death syndrome (R95) | 170 | 17.0 | 33.3 |
| 3 | Accidents (unintentional injuries) (V01-X59) | 102 | 10.2 | 20.0 |
| 4 | Diseases of the circulatory system (I00-I99) | 50 | 5.0 | 9.8 |
| 5 | Diarrhea and gastroenteritis of infectious origin (A09) | 47 | 4.7 | 9.2 |
| 6 | Influenza and pneumonia (J09-J18) | 36 | 3.6 | 7.1 |
| 7 | Assault (homicide) (*U01,X85-Y09) | 29 | 2.9 | 5.7 |
| 8 | Septicemia (A40-A41) | 27 | 2.7 | 5.3 |
| 9 | Chronic respiratory disease originating in the perinatal period (P27) | 16 | 1.6 | * |
| 10 | Renal failure and other disorders of kidney (N17-N19,N25,N27) | 8 | 0.8 | * |
| 11 | Malignant neoplasms (C00-C97) | 7 | 0.7 | * |
| 11 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 7 | 0.7 | * |
| 13 | Necrotizing enterocolitis of newborn (P77) | 6 | 0.6 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4 | 0.4 | * |
| 14 | Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87) | 4 | 0.4 | * |
| 14 | Anoxic brain damage, not elsewhere classified (G93.1) | 4 | 0.4 | * |
| 14 | Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56) | 4 | 0.4 | * |
| ... | All other causes (Residual) | 265 | 26.5 | 51.9 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|---|---|--------|-------------------------|----------------|
| Postneonates, Hispanic, female | | | | |
| ... | All causes | 794 | 100.0 | 162.3 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 202 | 25.4 | 41.3 |
| 2 | Sudden infant death syndrome (R95) | 133 | 16.8 | 27.2 |
| 3 | Diseases of the circulatory system (I00-I99) | 65 | 8.2 | 13.3 |
| 4 | Accidents (unintentional injuries) (V01-X59) | 64 | 8.1 | 13.1 |
| 5 | Assault (homicide) (*U01,X85-Y09) | 28 | 3.5 | 5.7 |
| 6 | Septicemia (A40-A41) | 22 | 2.8 | 4.5 |
| 7 | Diarrhea and gastroenteritis of infectious origin (A09) | 21 | 2.6 | 4.3 |
| 8 | Influenza and pneumonia (J09-J18) | 16 | 2.0 | * |
| 9 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 10 | 1.3 | * |
| 9 | Necrotizing enterocolitis of newborn (P77) | 10 | 1.3 | * |
| 11 | Malignant neoplasms (C00-C97) | 8 | 1.0 | * |
| 11 | Chronic respiratory disease originating in the perinatal period (P27) | 8 | 1.0 | * |
| 13 | Meningitis (G00,G03) | 6 | 0.8 | * |
| 13 | Renal failure and other disorders of kidney (N17-N19,N25,N27) | 6 | 0.8 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 5 | 0.6 | * |
| 15 | Acute bronchitis and acute bronchiolitis (J20-J21) | 5 | 0.6 | * |
| ... | All other causes (Residual) | 185 | 23.3 | 37.8 |
| Postneonates, Non-Hispanic, white, both sexes | | | | |
| ... | All causes | 4,046 | 100.0 | 182.9 |
| 1 | Sudden infant death syndrome (R95) | 1,016 | 25.1 | 45.9 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 701 | 17.3 | 31.7 |
| 3 | Accidents (unintentional injuries) (V01-X59) | 495 | 12.2 | 22.4 |
| 4 | Diseases of the circulatory system (I00-I99) | 194 | 4.8 | 8.8 |
| 5 | Assault (homicide) (*U01,X85-Y09) | 112 | 2.8 | 5.1 |
| 6 | Diarrhea and gastroenteritis of infectious origin (A09) | 99 | 2.4 | 4.5 |
| 7 | Influenza and pneumonia (J09-J18) | 95 | 2.3 | 4.3 |
| 8 | Septicemia (A40-A41) | 78 | 1.9 | 3.5 |
| 9 | Chronic respiratory disease originating in the perinatal period (P27) | 51 | 1.3 | 2.3 |
| 10 | Malignant neoplasms (C00-C97) | 31 | 0.8 | 1.4 |
| 11 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 30 | 0.7 | 1.4 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 24 | 0.6 | 1.1 |
| 13 | Necrotizing enterocolitis of newborn (P77) | 22 | 0.5 | 1.0 |
| 14 | Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89) | 20 | 0.5 | 0.9 |
| 14 | Meningitis (G00,G03) | 20 | 0.5 | 0.9 |
| 14 | Renal failure and other disorders of kidney (N17-N19,N25,N27) | 20 | 0.5 | 0.9 |
| ... | All other causes (Residual) | 1,038 | 25.7 | 46.9 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|---|---|--------|-------------------------|----------------|
| Postneonates, Non-Hispanic, white, male | | | | |
| ... | All causes | 2,350 | 100.0 | 207.1 |
| 1 | Sudden infant death syndrome (R95) | 617 | 26.3 | 54.4 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 345 | 14.7 | 30.4 |
| 3 | Accidents (unintentional injuries) (V01-X59) | 309 | 13.1 | 27.2 |
| 4 | Diseases of the circulatory system (I00-I99) | 107 | 4.6 | 9.4 |
| 5 | Assault (homicide) (*U01,X85-Y09) | 75 | 3.2 | 6.6 |
| 6 | Influenza and pneumonia (J09-J18) | 58 | 2.5 | 5.1 |
| 7 | Diarrhea and gastroenteritis of infectious origin (A09) | 53 | 2.3 | 4.7 |
| 8 | Septicemia (A40-A41) | 43 | 1.8 | 3.8 |
| 9 | Chronic respiratory disease originating in the perinatal period (P27) | 33 | 1.4 | 2.9 |
| 10 | Malignant neoplasms (C00-C97) | 21 | 0.9 | 1.9 |
| 11 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 20 | 0.9 | 1.8 |
| 12 | Renal failure and other disorders of kidney (N17-N19,N25,N27) | 15 | 0.6 | * |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 14 | 0.6 | * |
| 13 | Meningitis (G00,G03) | 14 | 0.6 | * |
| 13 | Necrotizing enterocolitis of newborn (P77) | 14 | 0.6 | * |
| ... | All other causes (Residual) | 612 | 26.0 | 53.9 |
| Postneonates, Non-Hispanic, white, female | | | | |
| ... | All causes | 1,696 | 100.0 | 157.3 |
| 1 | Sudden infant death syndrome (R95) | 399 | 23.5 | 37.0 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 356 | 21.0 | 33.0 |
| 3 | Accidents (unintentional injuries) (V01-X59) | 186 | 11.0 | 17.3 |
| 4 | Diseases of the circulatory system (I00-I99) | 87 | 5.1 | 8.1 |
| 5 | Diarrhea and gastroenteritis of infectious origin (A09) | 46 | 2.7 | 4.3 |
| 6 | Influenza and pneumonia (J09-J18) | 37 | 2.2 | 3.4 |
| 6 | Assault (homicide) (*U01,X85-Y09) | 37 | 2.2 | 3.4 |
| 8 | Septicemia (A40-A41) | 35 | 2.1 | 3.2 |
| 9 | Chronic respiratory disease originating in the perinatal period (P27) | 18 | 1.1 | * |
| 10 | Malignant neoplasms (C00-C97) | 10 | 0.6 | * |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 10 | 0.6 | * |
| 10 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 10 | 0.6 | * |
| 13 | Anoxic brain damage, not elsewhere classified (G93.1) | 9 | 0.5 | * |
| 14 | Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89) | 8 | 0.5 | * |
| 14 | Necrotizing enterocolitis of newborn (P77) | 8 | 0.5 | * |
| ... | All other causes (Residual) | 440 | 25.9 | 40.8 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|---|---|--------|-------------------------|----------------|
| Postneonates, Non-Hispanic, black, both sexes | | | | |
| ... | All causes | 2,836 | 100.0 | 465.2 |
| 1 | Sudden infant death syndrome (R95) | 606 | 21.4 | 99.4 |
| 2 | Accidents (unintentional injuries) (V01-X59) | 353 | 12.4 | 57.9 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 352 | 12.4 | 57.7 |
| 4 | Diarrhea and gastroenteritis of infectious origin (A09) | 137 | 4.8 | 22.5 |
| 5 | Diseases of the circulatory system (I00-I99) | 135 | 4.8 | 22.1 |
| 6 | Assault (homicide) (*U01,X85-Y09) | 105 | 3.7 | 17.2 |
| 7 | Septicemia (A40-A41) | 85 | 3.0 | 13.9 |
| 8 | Influenza and pneumonia (J09-J18) | 80 | 2.8 | 13.1 |
| 9 | Chronic respiratory disease originating in the perinatal period (P27) | 79 | 2.8 | 13.0 |
| 10 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 54 | 1.9 | 8.9 |
| 11 | Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89) | 24 | 0.8 | 3.9 |
| 12 | Necrotizing enterocolitis of newborn (P77) | 23 | 0.8 | 3.8 |
| 13 | Acute bronchitis and acute bronchiolitis (J20-J21) | 17 | 0.6 | * |
| 14 | Renal failure and other disorders of kidney (N17-N19,N25,N27) | 14 | 0.5 | * |
| 14 | Bacterial sepsis of newborn (P36) | 14 | 0.5 | * |
| ... | All other causes (Residual) | 758 | 26.7 | 124.3 |
| Postneonates, Non-Hispanic, black, male | | | | |
| ... | All causes | 1,629 | 100.0 | 525.9 |
| 1 | Sudden infant death syndrome (R95) | 351 | 21.5 | 113.3 |
| 2 | Accidents (unintentional injuries) (V01-X59) | 206 | 12.6 | 66.5 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 164 | 10.1 | 52.9 |
| 4 | Diarrhea and gastroenteritis of infectious origin (A09) | 84 | 5.2 | 27.1 |
| 5 | Diseases of the circulatory system (I00-I99) | 81 | 5.0 | 26.2 |
| 6 | Assault (homicide) (*U01,X85-Y09) | 60 | 3.7 | 19.4 |
| 7 | Chronic respiratory disease originating in the perinatal period (P27) | 47 | 2.9 | 15.2 |
| 8 | Septicemia (A40-A41) | 46 | 2.8 | 14.9 |
| 9 | Influenza and pneumonia (J09-J18) | 45 | 2.8 | 14.5 |
| 10 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 27 | 1.7 | 8.7 |
| 11 | Necrotizing enterocolitis of newborn (P77) | 15 | 0.9 | * |
| 12 | Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89) | 12 | 0.7 | * |
| 13 | Renal failure and other disorders of kidney (N17-N19,N25,N27) | 11 | 0.7 | * |
| 14 | Hernia of abdominal cavity and intestinal obstruction without hernia (K40-K46,K56) | 10 | 0.6 | * |
| 15 | Meningitis (G00,G03) | 8 | 0.5 | * |
| 15 | Acute bronchitis and acute bronchiolitis (J20-J21) | 8 | 0.5 | * |
| 15 | Bacterial sepsis of newborn (P36) | 8 | 0.5 | * |
| ... | All other causes (Residual) | 446 | 27.4 | 144.0 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable

LCWK8. Infant, neonatal, and postneonatal deaths, percent of total deaths, and mortality rates for the 15 leading causes of infant death by Hispanic origin, races for non-Hispanic population, and sex: United States, 2009

[Rates are infant (under 1 year), neonatal (under 28 days), and postneonatal (28 days-11 months) deaths per 100,000 live births in specified group. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on birth and death certificates. The title of ICD-10 code J09 was amended in 2009 to reflect the inclusion of Influenza A/H1N1 pandemic 2009 [swine flu]. The new J09 title is Influenza due to certain identified influenza virus]

| Rank ¹ | Cause of death (Based on the Tenth Revision, International Classification of Diseases, Second Edition, 2004), Hispanic origin and sex | Number | Percent of total deaths | Mortality rate |
|---|---|--------|-------------------------|----------------|
| Postneonates, Non-Hispanic, black, female | | | | |
| ... | All causes | 1,207 | 100.0 | 402.6 |
| 1 | Sudden infant death syndrome (R95) | 255 | 21.1 | 85.0 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 188 | 15.6 | 62.7 |
| 3 | Accidents (unintentional injuries) (V01-X59) | 147 | 12.2 | 49.0 |
| 4 | Diseases of the circulatory system (I00-I99) | 54 | 4.5 | 18.0 |
| 5 | Diarrhea and gastroenteritis of infectious origin (A09) | 53 | 4.4 | 17.7 |
| 6 | Assault (homicide) (*U01,X85-Y09) | 45 | 3.7 | 15.0 |
| 7 | Septicemia (A40-A41) | 39 | 3.2 | 13.0 |
| 8 | Influenza and pneumonia (J09-J18) | 35 | 2.9 | 11.7 |
| 9 | Chronic respiratory disease originating in the perinatal period (P27) | 32 | 2.7 | 10.7 |
| 10 | Disorders related to short gestation and low birth weight, not elsewhere classified (P07) | 27 | 2.2 | 9.0 |
| 11 | Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89) | 12 | 1.0 | * |
| 12 | Acute bronchitis and acute bronchiolitis (J20-J21) | 9 | 0.7 | * |
| 13 | Necrotizing enterocolitis of newborn (P77) | 8 | 0.7 | * |
| 14 | Anoxic brain damage, not elsewhere classified (G93.1) | 6 | 0.5 | * |
| 14 | Bacterial sepsis of newborn (P36) | 6 | 0.5 | * |
| ... | All other causes (Residual) | 291 | 24.1 | 97.1 |

Source: CDC/NCHS, National Vital Statistics System

¹ Rank based on number of deaths

* Figure does not meet standards of reliability or precision

... Category not applicable