

Person with disabilities experiencing problems accessing
NCHS birth and death tables should contact
nhsed@cdc.gov, or call 301-458-4688

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
All races 3/, both sexes										
2006.....	810.4	690.7	28.4	13.9	16.6	64.4	100.2	100.9	111.9	148.7
2005.....	825.9	692.5	29.4	14.5	18.1	65.1	97.6	97.5	111.4	149.6
2004.....	816.5	685.2	29.9	14.7	18.7	66.1	94.0	96.0	107.9	151.7
2003.....	841.9	700.0	31.5	14.7	19.1	66.4	96.4	95.2	111.3	159.1
2002.....	847.3	695.0	31.2	15.2	19.5	67.8	95.2	94.7	111.6	161.3
2001.....	848.5	683.4	33.3	15.3	19.2	66.9	95.0	96.2	113.5	165.9
2000.....	854.0	736.7	32.4	15.8	20.3	67.1	93.6	91.2	111.0	159.2
1999.....	857.0	736.0	34.2	16.9	20.4	68.6	90.8	90.4	113.4	160.8
All races 3/, male										
2006.....	814.8	756.3	30.5	15.4	19.6	90.7	148.0	143.4	150.4	189.0
2005.....	827.2	762.3	33.4	15.6	21.5	91.6	143.9	138.7	148.1	189.4
2004.....	817.6	753.7	32.4	16.4	21.7	91.0	138.0	135.6	143.2	193.8
2003.....	840.3	777.4	35.1	16.2	23.1	92.3	140.3	133.5	148.9	202.7
2002.....	846.6	761.5	35.2	16.7	23.1	94.0	140.8	134.4	149.4	206.6
2001.....	846.4	749.8	37.0	16.7	22.8	93.7	141.1	135.6	151.2	213.7
2000.....	853.0	806.5	35.9	17.6	24.2	93.3	138.0	128.8	147.9	205.4
1999.....	859.2	805.0	37.9	18.6	24.4	94.7	133.1	127.0	152.0	207.6
All races 3/, female										
2006.....	806.1	622.0	26.3	12.3	13.3	36.8	49.0	56.5	72.5	107.9
2005.....	824.6	619.4	25.1	13.4	14.4	37.2	48.2	54.4	73.7	109.3
2004.....	815.4	613.4	27.3	13.0	15.4	39.9	47.3	54.5	72.0	109.3
2003.....	843.4	619.1	27.8	13.0	15.0	39.0	49.8	55.4	73.0	115.3
2002.....	848.0	625.3	27.0	13.5	15.9	40.1	47.3	53.6	73.3	116.1
2001.....	850.4	613.9	29.5	13.9	15.3	38.5	46.8	55.8	75.2	118.2
2000.....	855.0	663.4	28.7	14.0	16.1	39.3	47.1	52.8	73.6	113.2
1999.....	854.9	663.6	30.3	15.0	16.2	41.0	46.6	53.2	74.4	114.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
White, both sexes										
2006 4/.....	858.1	576.0	25.5	12.8	15.6	61.1	94.8	93.6	103.3	137.8
2005 4/.....	873.7	579.1	27.0	13.3	16.7	62.5	91.5	89.3	101.7	138.3
2004 4/.....	863.2	574.0	27.0	13.4	17.4	64.4	88.3	87.3	98.3	140.6
2003 4/.....	890.1	591.3	28.5	13.7	17.8	64.7	89.4	86.1	101.9	145.5
2002.....	895.7	586.7	28.1	14.0	18.1	66.3	88.0	85.0	101.1	146.8
2001.....	895.1	570.9	30.7	14.4	17.9	64.2	87.1	87.0	102.7	151.2
2000.....	900.2	610.6	29.2	14.8	19.2	64.5	85.1	81.0	99.0	144.0
1999.....	901.4	606.1	30.8	15.3	19.2	65.3	81.6	80.8	101.4	144.5
White, male										
2006 4/.....	852.3	632.7	27.5	14.1	18.6	84.1	139.3	132.3	138.7	176.1
2005 4/.....	864.5	640.0	30.9	14.2	19.8	86.5	133.8	126.5	135.1	176.1
2004 4/.....	854.2	631.6	29.4	15.1	20.6	87.1	129.1	123.2	130.8	180.9
2003 4/.....	877.6	658.7	31.5	15.3	21.4	88.3	129.1	120.0	137.0	187.6
2002.....	884.0	650.9	31.5	15.4	21.2	90.5	128.9	119.7	136.0	188.7
2001.....	881.9	627.6	34.2	15.3	21.3	88.3	128.2	122.4	137.4	196.7
2000.....	887.8	667.6	32.6	16.5	23.1	88.1	124.6	113.9	133.6	188.2
1999.....	892.1	667.0	33.9	16.8	22.7	88.3	118.5	113.5	136.5	188.8
White, female										
2006 4/.....	863.9	516.5	23.5	11.4	12.5	36.9	46.6	52.2	65.9	98.1
2005 4/.....	882.8	515.3	22.9	12.3	13.4	37.2	45.8	49.4	66.6	99.0
2004 4/.....	871.9	513.6	24.4	11.6	14.2	40.4	44.3	48.8	64.3	98.9
2003 4/.....	902.3	520.7	25.5	12.1	14.0	39.6	46.7	50.0	65.2	102.2
2002.....	907.0	519.4	24.5	12.6	14.8	40.4	44.4	48.1	64.8	103.9
2001.....	907.9	511.5	27.1	13.3	14.4	38.5	43.2	49.6	66.6	104.9
2000.....	912.3	550.5	25.5	13.0	15.2	39.3	43.0	46.5	63.0	99.2
1999.....	910.4	542.0	27.5	13.8	15.4	40.8	42.3	46.7	65.0	99.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All causes										
Black, both sexes										
2006 4/.....	733.0	1,303.1	43.3	19.9	22.4	84.7	142.0	163.9	192.9	252.6
2005 4/.....	749.4	1,311.2	41.8	21.1	25.2	83.6	143.1	162.0	197.3	254.1
2004 4/.....	744.3	1,284.7	44.8	21.6	25.3	81.4	138.8	167.2	191.3	254.4
2003 4/.....	763.6	1,273.8	46.8	19.7	25.7	80.0	147.3	168.6	194.1	275.2
2002.....	768.4	1,263.6	47.1	21.3	27.5	82.0	146.8	172.1	200.7	284.0
2001.....	773.5	1,304.5	47.5	20.7	25.8	86.3	151.3	168.8	200.5	286.0
2000.....	781.1	1,426.1	49.9	21.8	26.6	87.4	155.0	168.4	206.9	283.1
1999.....	788.1	1,457.3	52.6	25.2	28.0	91.9	154.4	163.5	204.9	293.5
Black, male										
2006 4/.....	786.7	1,407.1	47.1	22.9	26.7	130.8	215.7	242.0	268.0	324.7
2005 4/.....	799.2	1,437.2	46.7	22.6	30.9	127.2	220.4	239.1	270.7	323.9
2004 4/.....	792.6	1,414.2	48.6	23.1	28.6	121.0	210.1	244.7	259.9	323.6
2003 4/.....	813.7	1,410.0	53.7	21.6	31.4	120.9	224.5	249.6	262.5	344.8
2002.....	816.7	1,351.5	54.4	24.1	33.4	121.7	227.6	255.7	272.9	366.5
2001.....	823.9	1,426.4	53.2	23.8	31.6	130.4	236.3	247.0	271.0	364.1
2000.....	834.1	1,567.6	54.5	24.5	32.1	130.1	240.6	246.8	275.0	361.6
1999.....	847.4	1,592.8	59.0	29.3	35.0	137.7	239.9	238.0	278.9	372.8
Black, female										
2006 4/.....	684.0	1,194.6	39.4	16.8	17.9	37.4	66.5	89.8	124.9	188.2
2005 4/.....	703.9	1,179.7	36.7	19.6	19.3	38.8	64.5	89.1	131.0	191.5
2004 4/.....	700.3	1,149.9	40.9	20.1	21.9	40.7	67.0	94.7	129.5	192.6
2003 4/.....	717.9	1,132.2	39.7	17.8	19.8	37.8	70.6	93.6	132.5	213.1
2002.....	724.4	1,172.0	39.5	18.5	21.3	41.0	68.0	95.7	135.7	210.6
2001.....	727.7	1,176.3	41.7	17.5	19.8	40.8	69.4	97.5	137.3	216.6
2000.....	733.0	1,279.8	45.3	19.0	21.0	43.7	73.9	97.3	145.6	213.4
1999.....	734.3	1,317.4	46.1	21.0	20.7	45.2	73.2	96.1	138.6	223.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
I. Certain infectious and parasitic diseases (A00-B99)										
All races 3/, both sexes										
2006.....	22.4	11.6	1.1	0.4	0.5	0.7	1.6	3.1	5.4	10.2
2005.....	22.4	12.9	1.1	0.4	0.5	0.8	1.4	3.2	6.1	11.0
2004.....	22.0	11.8	1.2	0.4	0.4	0.7	1.6	3.3	6.8	12.5
2003.....	22.2	12.3	1.2	0.5	0.5	0.8	1.5	3.4	7.2	14.4
2002.....	22.2	14.4	1.1	0.5	0.6	0.8	1.5	3.9	8.2	15.3
2001.....	21.2	14.0	1.5	0.5	0.5	1.0	1.7	4.1	9.0	15.5
2000.....	21.0	13.4	1.4	0.5	0.5	0.8	1.6	4.4	10.4	16.0
1999.....	21.5	14.8	1.5	0.7	0.6	0.7	1.8	5.1	11.6	16.5
All races 3/, male										
2006.....	23.8	12.3	1.2	0.5	0.5	0.8	1.6	3.5	6.4	13.2
2005.....	24.0	14.7	1.2	0.5	0.5	0.8	1.5	3.7	7.2	14.5
2004.....	23.7	13.8	1.1	0.4	0.4	0.7	1.8	3.6	8.4	16.8
2003.....	24.0	12.6	1.4	0.4	0.4	0.9	1.6	3.9	9.3	19.7
2002.....	24.2	14.8	1.2	0.5	0.6	0.7	1.6	4.4	10.5	21.6
2001.....	23.3	14.7	1.6	0.5	0.6	1.0	1.7	4.7	12.0	22.2
2000.....	23.0	14.1	1.5	0.5	0.5	0.9	1.8	4.9	13.8	23.1
1999.....	23.7	17.0	1.8	0.7	0.6	0.7	1.9	6.3	15.9	23.7
All races 3/, female										
2006.....	21.0	10.9	1.0	0.3	0.4	0.7	1.5	2.6	4.3	7.2
2005.....	20.9	11.1	1.0	0.4	0.4	0.8	1.3	2.7	5.0	7.4
2004.....	20.3	9.6	1.2	0.4	0.4	0.7	1.3	3.1	5.2	8.2
2003.....	20.6	12.1	1.1	0.5	0.5	0.8	1.5	3.0	5.0	9.0
2002.....	20.2	14.0	0.9	0.4	0.6	0.9	1.4	3.5	5.8	8.9
2001.....	19.2	13.2	1.3	0.5	0.5	0.9	1.7	3.5	6.0	8.9
2000.....	19.0	12.7	1.4	0.4	0.5	0.7	1.4	3.8	6.9	9.0
1999.....	19.3	12.5	1.2	0.7	0.5	0.7	1.8	3.9	7.2	9.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
I. Certain infectious and parasitic diseases (A00-B99)										
White, both sexes										
2006 4/.....	20.6	8.6	1.0	0.4	0.4	0.6	0.9	1.7	3.1	6.2
2005 4/.....	20.5	10.0	1.0	0.4	0.5	0.6	0.9	1.8	3.4	6.8
2004 4/.....	19.9	9.9	1.1	0.4	0.3	0.5	0.9	1.8	3.7	8.2
2003 4/.....	20.0	9.0	1.0	0.4	0.4	0.7	1.0	1.8	4.3	8.9
2002.....	19.7	12.4	1.0	0.4	0.5	0.7	0.9	2.3	4.5	9.6
2001.....	18.6	10.6	1.3	0.5	0.4	0.7	1.0	2.1	5.1	9.8
2000.....	18.2	10.6	1.2	0.4	0.4	0.6	1.0	2.3	5.8	10.3
1999.....	18.6	11.9	1.3	0.6	0.4	0.6	1.1	2.9	7.1	10.8
White, male										
2006 4/.....	21.4	9.5	1.1	0.4	0.5	0.6	1.0	2.0	4.1	8.7
2005 4/.....	21.3	11.6	1.1	0.3	0.4	0.5	1.1	2.2	4.3	9.7
2004 4/.....	20.9	11.6	1.1	0.4	0.3	0.5	1.2	2.0	5.2	11.6
2003 4/.....	20.9	8.6	1.2	0.3	0.3	0.8	1.0	2.3	5.9	13.1
2002.....	20.9	12.7	1.2	0.5	0.6	0.6	1.1	2.9	6.3	14.5
2001.....	19.8	11.3	1.5	0.5	0.5	0.7	1.1	2.7	7.3	15.0
2000.....	19.3	11.2	1.2	0.5	0.4	0.8	1.2	2.9	8.5	15.9
1999.....	19.8	14.3	1.6	0.6	0.5	0.6	1.3	3.8	10.6	16.7
White, female										
2006 4/.....	19.8	7.6	0.9	0.4	0.4	0.5	0.9	1.3	2.0	3.6
2005 4/.....	19.7	8.5	1.0	0.4	0.5	0.6	0.7	1.3	2.3	3.8
2004 4/.....	19.0	8.1	1.1	0.4	0.3	0.5	0.7	1.5	2.2	4.7
2003 4/.....	19.1	9.4	0.9	0.4	0.4	0.5	1.0	1.4	2.5	4.5
2002.....	18.6	12.1	0.8	0.4	0.5	0.8	0.8	1.6	2.6	4.6
2001.....	17.4	9.9	1.2	0.5	0.4	0.6	0.9	1.5	2.8	4.5
2000.....	17.2	10.0	1.2	0.4	0.5	0.5	0.7	1.8	2.9	4.6
1999.....	17.4	9.4	1.1	0.6	0.4	0.6	0.8	1.9	3.5	4.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
I. Certain infectious and parasitic diseases (A00-B99)										
Black, both sexes										
2006 4/.....	39.2	26.6	1.6	*	0.8	1.5	5.0	12.0	21.0	37.8
2005 4/.....	40.0	27.7	1.4	0.7	0.6	1.7	4.3	12.1	24.5	40.0
2004 4/.....	40.4	21.8	1.5	*	0.8	1.6	5.3	13.2	27.2	42.5
2003 4/.....	41.7	28.2	2.3	0.9	0.9	1.6	4.5	13.5	26.6	51.9
2002.....	42.9	24.6	1.7	*	1.2	1.5	4.6	14.8	32.8	54.9
2001.....	42.3	31.2	2.2	0.6	1.1	2.4	5.9	16.6	35.7	55.4
2000.....	43.2	29.9	2.4	0.8	0.7	1.6	5.4	17.6	41.6	56.7
1999.....	44.5	31.1	2.5	1.3	1.1	1.3	6.0	19.6	42.4	57.5
Black, male										
2006 4/.....	45.4	26.5	1.9	*	*	1.6	5.4	13.5	23.6	46.5
2005 4/.....	47.1	30.4	1.7	*	*	1.8	4.4	13.4	27.7	50.9
2004 4/.....	47.8	24.3	*	*	*	1.7	5.6	13.9	31.5	55.4
2003 4/.....	50.0	31.8	2.5	*	*	1.3	4.6	14.8	33.6	68.5
2002.....	51.7	25.3	1.6	*	*	1.4	4.6	14.6	40.9	75.9
2001.....	51.2	33.2	2.6	*	*	2.4	5.6	18.6	46.4	76.4
2000.....	52.7	31.3	2.8	*	*	1.7	5.4	19.4	52.7	79.0
1999.....	54.6	33.2	3.2	1.3	1.4	1.3	5.1	23.8	56.2	78.3
Black, female										
2006 4/.....	33.4	26.8	*	*	*	1.4	4.6	10.6	18.6	30.1
2005 4/.....	33.5	25.0	*	*	*	1.6	4.1	10.9	21.7	30.2
2004 4/.....	33.7	19.1	*	*	*	1.5	5.1	12.5	23.4	31.0
2003 4/.....	34.2	24.4	2.0	*	*	2.0	4.5	12.3	20.4	37.2
2002.....	35.0	23.9	1.7	*	1.4	1.7	4.6	14.9	25.5	36.4
2001.....	34.2	29.0	1.8	*	*	2.3	6.2	14.8	26.2	36.6
2000.....	34.5	28.4	2.1	*	*	1.4	5.4	16.1	31.7	36.9
1999.....	35.3	28.9	1.9	1.3	*	*	6.9	15.8	30.1	39.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Zoonotic and other bacterial diseases (A20-A49)										
All races 3/, both sexes										
2006.....	12.0	7.5	0.7	0.2	0.2	0.3	0.5	0.7	0.9	1.6
2005.....	12.1	8.7	0.7	0.2	0.3	0.4	0.5	0.7	1.0	1.6
2004.....	12.0	7.5	0.7	0.2	0.2	0.3	0.5	0.7	1.1	1.6
2003.....	12.3	7.9	0.7	0.2	0.2	0.4	0.6	0.7	1.0	1.6
2002.....	12.3	8.6	0.6	0.3	0.3	0.4	0.4	0.7	1.0	1.6
2001.....	11.9	8.9	0.9	0.2	0.3	0.5	0.5	0.6	1.0	1.5
2000.....	11.6	8.6	0.8	0.3	0.2	0.4	0.5	0.6	0.9	1.6
1999.....	11.7	9.1	0.8	0.3	0.2	0.3	0.6	0.7	1.0	1.6
All races 3/, male										
2006.....	11.1	8.0	0.7	0.3	0.3	0.3	0.5	0.7	1.0	1.6
2005.....	11.1	9.7	0.7	0.3	0.3	0.4	0.6	0.8	0.9	1.5
2004.....	11.0	8.9	0.7	0.2	*	0.3	0.5	0.6	1.2	1.7
2003.....	11.1	8.4	0.8	*	0.2	0.4	0.5	0.6	1.1	1.7
2002.....	11.1	8.6	0.7	0.3	0.4	0.4	0.5	0.8	1.0	1.6
2001.....	10.8	9.5	0.9	0.2	0.3	0.5	0.5	0.6	1.0	1.7
2000.....	10.4	8.9	0.9	0.3	*	0.5	0.5	0.7	1.0	1.6
1999.....	10.5	10.4	0.9	0.3	0.2	0.3	0.7	0.7	1.1	1.7
All races 3/, female										
2006.....	12.9	7.0	0.6	*	0.2	0.3	0.5	0.7	0.9	1.7
2005.....	13.1	7.6	0.6	*	0.2	0.4	0.5	0.6	1.1	1.7
2004.....	12.9	6.1	0.7	0.3	0.2	0.3	0.5	0.7	1.1	1.5
2003.....	13.5	7.5	0.5	0.3	0.3	0.4	0.6	0.8	0.8	1.6
2002.....	13.5	8.5	0.6	0.2	0.3	0.4	0.4	0.7	0.9	1.6
2001.....	12.9	8.3	0.8	*	0.3	0.5	0.4	0.6	1.0	1.4
2000.....	12.9	8.3	0.7	0.2	*	0.3	0.4	0.5	0.9	1.5
1999.....	12.9	7.7	0.7	0.3	0.3	0.3	0.5	0.6	0.9	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Zoonotic and other bacterial diseases (A20-A49)										
White, both sexes										
2006 4/.....	11.9	5.6	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.3
2005 4/.....	12.0	6.3	0.6	0.2	0.3	0.3	0.4	0.6	0.8	1.3
2004 4/.....	11.8	6.1	0.6	0.3	0.2	0.3	0.4	0.6	0.9	1.4
2003 4/.....	12.1	5.6	0.5	0.2	0.2	0.3	0.5	0.6	0.8	1.3
2002.....	12.1	7.3	0.6	0.3	0.3	0.4	0.4	0.6	0.8	1.3
2001.....	11.7	6.8	0.8	0.2	0.3	0.5	0.3	0.5	0.7	1.2
2000.....	11.3	7.1	0.7	0.2	0.2	0.4	0.4	0.5	0.7	1.2
1999.....	11.3	7.2	0.8	0.3	0.2	0.3	0.5	0.5	0.8	1.2
White, male										
2006 4/.....	11.0	6.1	0.6	0.3	0.3	0.3	0.4	0.5	0.8	1.4
2005 4/.....	10.9	7.2	0.7	*	0.3	0.3	0.5	0.6	0.7	1.2
2004 4/.....	10.8	7.3	0.6	0.3	*	0.3	0.4	0.5	0.9	1.5
2003 4/.....	10.8	5.6	0.7	*	*	0.4	0.5	0.5	0.8	1.4
2002.....	10.9	7.3	0.7	0.4	0.3	0.3	0.4	0.7	0.8	1.3
2001.....	10.5	7.5	0.9	*	0.3	0.5	0.4	0.5	0.7	1.4
2000.....	10.0	7.2	0.8	0.3	*	0.5	0.5	0.5	0.8	1.3
1999.....	10.0	8.5	0.8	0.3	*	0.3	0.6	0.6	0.9	1.4
White, female										
2006 4/.....	12.8	5.2	0.6	*	*	0.3	0.4	0.5	0.7	1.3
2005 4/.....	13.0	5.3	0.5	*	*	0.3	0.4	0.5	0.9	1.4
2004 4/.....	12.8	4.9	0.6	*	*	0.3	0.4	0.6	0.8	1.3
2003 4/.....	13.4	5.6	0.4	*	*	0.3	0.6	0.6	0.7	1.2
2002.....	13.4	7.4	0.5	*	*	0.4	0.4	0.5	0.8	1.3
2001.....	12.8	6.1	0.7	*	*	0.4	0.3	0.5	0.8	1.0
2000.....	12.6	7.0	0.7	*	*	0.3	0.3	0.5	0.6	1.1
1999.....	12.6	5.9	0.7	0.3	*	0.3	0.4	0.5	0.8	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Zoonotic and other bacterial diseases (A20-A49)										
Black, both sexes										
2006 4/.....	16.1	17.1	0.9	*	*	*	1.2	1.5	2.1	4.1
2005 4/.....	16.6	21.1	0.9	*	*	0.7	0.9	1.6	2.6	3.9
2004 4/.....	16.2	15.1	0.9	*	*	*	1.2	1.4	2.7	2.8
2003 4/.....	16.9	19.6	1.2	*	*	0.7	0.9	1.7	2.3	3.9
2002.....	16.9	14.8	0.8	*	0.6	*	0.7	1.8	2.2	3.8
2001.....	16.5	20.3	1.1	*	*	0.9	1.0	1.2	2.8	3.9
2000.....	16.8	18.4	1.2	*	*	*	0.9	1.6	2.7	4.2
1999.....	17.0	19.0	0.9	*	*	*	1.2	1.7	2.2	4.2
Black, male										
2006 4/.....	15.1	18.1	*	*	*	*	*	1.5	2.4	4.0
2005 4/.....	15.7	22.4	*	*	*	*	*	1.8	2.3	4.0
2004 4/.....	15.3	16.9	*	*	*	*	*	*	3.1	2.6
2003 4/.....	16.0	22.0	*	*	*	*	*	1.6	2.9	3.9
2002.....	15.7	15.7	*	*	*	*	*	*	2.6	4.1
2001.....	15.6	20.7	*	*	*	*	1.4	*	2.8	4.0
2000.....	15.6	19.8	*	*	*	*	*	2.1	2.8	4.2
1999.....	16.3	20.5	*	*	*	*	*	*	2.8	4.5
Black, female										
2006 4/.....	17.0	15.9	*	*	*	*	*	1.6	1.7	4.1
2005 4/.....	17.4	19.7	*	*	*	*	*	1.5	2.8	3.9
2004 4/.....	17.1	13.3	*	*	*	*	*	1.5	2.3	3.0
2003 4/.....	17.7	17.0	*	*	*	*	*	1.9	1.6	3.9
2002.....	18.0	13.9	*	*	*	*	*	2.2	2.0	3.5
2001.....	17.3	19.8	*	*	*	*	*	*	2.8	3.8
2000.....	18.0	16.9	*	*	*	*	*	*	2.6	4.2
1999.....	17.7	17.5	*	*	*	*	*	1.9	1.6	3.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
All races 3/, both sexes										
2006.....	11.4	6.5	0.5	0.2	0.2	0.2	0.4	0.6	0.8	1.5
2005.....	11.5	7.4	0.5	0.2	0.2	0.3	0.4	0.6	0.9	1.5
2004.....	11.4	6.6	0.5	0.2	0.1	0.2	0.4	0.6	1.0	1.5
2003.....	11.7	6.9	0.5	0.2	0.2	0.3	0.5	0.6	0.9	1.5
2002.....	11.7	7.3	0.5	0.2	0.3	0.2	0.3	0.6	0.9	1.5
2001.....	11.3	7.7	0.7	0.1	0.2	0.3	0.4	0.5	0.9	1.4
2000.....	11.1	7.2	0.6	0.2	0.1	0.2	0.3	0.5	0.9	1.4
1999.....	11.0	7.5	0.6	0.2	0.1	0.2	0.5	0.5	0.8	1.4
All races 3/, male										
2006.....	10.5	6.8	0.5	0.2	0.3	0.2	0.4	0.6	0.9	1.5
2005.....	10.5	8.1	0.6	0.2	0.2	0.3	0.5	0.7	0.8	1.4
2004.....	10.4	8.0	0.5	*	*	0.2	0.4	0.5	1.1	1.5
2003.....	10.5	7.4	0.7	*	*	0.3	0.4	0.6	1.0	1.6
2002.....	10.6	7.7	0.5	0.3	0.2	0.2	0.4	0.7	1.0	1.5
2001.....	10.2	8.4	0.8	*	0.2	0.3	0.4	0.5	0.9	1.5
2000.....	9.8	7.1	0.7	0.2	*	0.2	0.4	0.6	0.9	1.4
1999.....	9.8	9.0	0.6	0.2	*	*	0.5	0.5	0.9	1.5
All races 3/, female										
2006.....	12.3	6.2	0.6	*	*	0.3	0.4	0.6	0.8	1.6
2005.....	12.5	6.5	0.5	*	0.2	0.3	0.4	0.6	1.0	1.6
2004.....	12.3	5.2	0.6	0.2	*	0.2	0.4	0.7	1.0	1.4
2003.....	12.9	6.5	0.4	0.2	*	0.2	0.5	0.7	0.8	1.4
2002.....	12.9	7.0	0.5	*	0.3	0.3	0.3	0.6	0.9	1.5
2001.....	12.4	7.0	0.6	*	*	0.3	0.3	0.5	0.9	1.2
2000.....	12.3	7.3	0.6	*	*	*	0.2	0.4	0.8	1.4
1999.....	12.2	5.8	0.5	0.2	*	*	0.4	0.5	0.7	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
White, both sexes										
2006 4/.....	11.3	4.9	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.2
2005 4/.....	11.3	5.0	0.5	0.2	0.2	0.2	0.4	0.5	0.7	1.2
2004 4/.....	11.2	5.2	0.5	0.2	*	0.2	0.3	0.5	0.8	1.3
2003 4/.....	11.5	4.6	0.4	0.2	0.2	0.2	0.4	0.5	0.7	1.2
2002.....	11.5	6.1	0.5	0.2	0.2	0.2	0.3	0.5	0.8	1.2
2001.....	11.1	5.4	0.7	0.2	0.2	0.2	0.3	0.4	0.7	1.1
2000.....	10.8	5.7	0.6	0.2	0.1	0.2	0.2	0.4	0.6	1.1
1999.....	10.6	5.5	0.5	0.2	0.1	0.2	0.4	0.4	0.7	1.0
White, male										
2006 4/.....	10.4	5.1	0.4	*	0.3	*	0.3	0.5	0.7	1.3
2005 4/.....	10.2	5.9	0.5	*	*	*	0.4	0.6	0.6	1.1
2004 4/.....	10.2	6.3	0.5	*	*	*	0.3	0.4	0.8	1.4
2003 4/.....	10.2	4.6	0.6	*	*	0.3	0.4	0.4	0.8	1.3
2002.....	10.3	6.2	0.5	0.3	0.2	*	0.4	0.6	0.8	1.2
2001.....	9.9	6.2	0.8	*	*	0.3	0.3	0.4	0.6	1.3
2000.....	9.4	5.8	0.6	*	*	*	0.3	0.4	0.7	1.2
1999.....	9.3	6.9	0.5	*	*	*	0.5	0.5	0.8	1.2
White, female										
2006 4/.....	12.2	4.6	0.5	*	*	0.2	0.3	0.5	0.7	1.2
2005 4/.....	12.4	4.1	0.4	*	*	*	0.3	0.4	0.8	1.3
2004 4/.....	12.2	4.0	0.5	*	*	*	0.3	0.6	0.8	1.2
2003 4/.....	12.8	4.5	*	*	*	*	0.4	0.6	0.7	1.1
2002.....	12.8	6.0	0.4	*	*	0.3	*	0.4	0.8	1.2
2001.....	12.3	4.6	0.6	*	*	*	0.3	0.4	0.7	0.9
2000.....	12.1	5.6	0.5	*	*	*	*	0.4	0.5	1.0
1999.....	11.9	4.0	0.5	*	*	*	*	0.4	0.6	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Septicemia (A40-A41)										
Black, both sexes										
2006 4/.....	15.4	15.0	0.8	*	*	*	0.9	1.4	1.8	3.8
2005 4/.....	15.9	19.8	0.8	*	*	0.6	0.8	1.5	2.3	3.6
2004 4/.....	15.6	14.5	*	*	*	*	1.0	1.3	2.5	2.7
2003 4/.....	16.3	18.6	0.9	*	*	*	0.7	1.6	2.2	3.6
2002.....	16.3	13.5	*	*	*	*	*	1.6	2.2	3.3
2001.....	15.8	19.3	0.9	*	*	*	0.7	1.1	2.5	3.7
2000.....	16.2	16.4	1.0	*	*	*	0.8	1.5	2.5	3.7
1999.....	16.1	17.7	*	*	*	*	1.0	1.4	1.7	3.7
Black, male										
2006 4/.....	14.4	15.6	*	*	*	*	*	*	2.1	3.6
2005 4/.....	14.9	20.4	*	*	*	*	*	1.6	2.1	3.6
2004 4/.....	14.6	16.3	*	*	*	*	*	*	2.8	2.5
2003 4/.....	15.3	21.1	*	*	*	*	*	1.6	2.7	3.6
2002.....	15.0	14.8	*	*	*	*	*	*	2.5	3.7
2001.....	14.9	20.1	*	*	*	*	*	*	2.5	3.7
2000.....	14.9	16.0	*	*	*	*	*	2.1	2.5	3.5
1999.....	15.3	19.8	*	*	*	*	*	*	2.1	4.0
Black, female										
2006 4/.....	16.4	14.4	*	*	*	*	*	1.5	1.5	4.0
2005 4/.....	16.8	19.1	*	*	*	*	*	1.4	2.6	3.7
2004 4/.....	16.4	12.7	*	*	*	*	*	1.4	2.1	2.9
2003 4/.....	17.2	16.1	*	*	*	*	*	1.6	1.6	3.5
2002.....	17.4	12.1	*	*	*	*	*	1.9	2.0	3.1
2001.....	16.7	18.6	*	*	*	*	*	*	2.5	3.7
2000.....	17.4	16.9	*	*	*	*	*	*	2.5	3.9
1999.....	16.9	15.4	*	*	*	*	*	1.6	1.4	3.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral diseases (A80-B34)										
All races 3/, both sexes										
2006.....	6.9	2.9	0.3	*	0.2	0.3	0.9	2.2	4.1	8.2
2005.....	6.6	2.9	0.3	0.1	0.1	0.3	0.7	2.3	4.8	8.8
2004.....	6.8	2.9	0.3	0.1	0.2	0.3	0.9	2.4	5.3	10.3
2003.....	7.1	2.9	0.4	0.2	0.2	0.3	0.8	2.5	5.8	12.1
2002.....	7.4	4.1	0.3	0.1	0.2	0.3	0.9	2.9	6.8	13.1
2001.....	7.4	3.0	0.4	0.2	0.2	0.4	1.1	3.3	7.7	13.4
2000.....	7.5	2.9	0.4	0.2	0.2	0.3	1.0	3.6	9.1	13.9
1999.....	7.6	3.5	0.5	0.3	0.2	0.3	1.0	4.1	10.0	14.1
All races 3/, male										
2006.....	9.6	3.0	0.3	*	*	0.4	1.0	2.6	5.1	11.1
2005.....	9.2	3.3	0.3	*	*	0.2	0.8	2.7	5.8	12.2
2004.....	9.5	3.2	0.4	*	*	0.3	1.0	2.7	6.7	14.3
2003.....	10.1	2.5	0.4	*	*	0.3	0.9	3.0	7.7	17.2
2002.....	10.6	4.6	0.4	*	0.2	0.3	1.0	3.3	9.1	19.3
2001.....	10.5	3.1	0.5	0.2	*	0.4	1.0	3.8	10.7	19.7
2000.....	10.8	2.9	0.4	*	0.2	0.3	1.1	4.1	12.3	20.7
1999.....	10.9	4.0	0.6	0.3	0.3	0.3	1.0	5.3	14.2	21.0
All races 3/, female										
2006.....	4.3	2.8	*	*	*	0.3	0.8	1.8	3.2	5.3
2005.....	4.0	2.5	0.3	*	*	0.3	0.7	1.8	3.8	5.3
2004.....	4.1	2.6	0.3	*	*	0.3	0.8	2.1	3.9	6.3
2003.....	4.2	3.3	0.4	*	*	0.3	0.8	2.0	3.9	7.0
2002.....	4.4	3.6	*	*	0.2	0.4	0.8	2.6	4.5	6.9
2001.....	4.3	2.8	0.3	0.2	*	0.3	1.2	2.7	4.6	7.1
2000.....	4.4	2.9	0.4	*	0.3	0.3	0.9	3.1	5.8	7.1
1999.....	4.3	3.1	0.4	0.3	*	0.3	1.0	2.9	5.8	7.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral diseases (A80-B34)										
White, both sexes										
2006 4/.....	5.0	2.2	0.3	*	0.1	0.2	0.4	1.0	2.0	4.6
2005 4/.....	4.6	2.6	0.3	*	*	0.1	0.4	1.0	2.3	5.0
2004 4/.....	4.6	2.6	0.3	*	*	0.2	0.4	1.0	2.6	6.2
2003 4/.....	4.9	2.3	0.4	*	*	0.2	0.4	1.2	3.2	7.1
2002.....	5.1	3.7	0.3	*	0.2	0.2	0.4	1.5	3.4	7.9
2001.....	5.0	2.3	0.3	0.2	*	0.2	0.5	1.4	4.1	8.1
2000.....	5.1	2.1	0.3	0.2	0.2	0.2	0.4	1.7	4.8	8.7
1999.....	5.2	2.8	0.4	0.2	0.1	0.2	0.5	2.1	5.9	8.9
White, male										
2006 4/.....	7.2	2.5	0.3	*	*	0.2	0.5	1.3	3.0	7.0
2005 4/.....	6.7	2.8	0.3	*	*	*	0.4	1.4	3.2	7.8
2004 4/.....	6.8	2.8	0.3	*	*	*	0.5	1.3	4.0	9.4
2003 4/.....	7.2	1.9	0.3	*	*	0.3	0.5	1.6	4.8	11.0
2002.....	7.5	4.1	0.4	*	*	*	0.5	2.0	5.2	12.8
2001.....	7.4	2.4	0.4	*	*	*	0.6	1.9	6.3	13.0
2000.....	7.5	2.2	*	*	*	0.2	0.6	2.2	7.4	14.0
1999.....	7.6	3.4	0.6	*	*	*	0.6	2.9	9.2	14.5
White, female										
2006 4/.....	2.8	1.9	*	*	*	*	0.4	0.6	1.0	2.2
2005 4/.....	2.6	2.4	*	*	*	*	0.3	0.6	1.4	2.0
2004 4/.....	2.5	2.4	*	*	*	*	*	0.6	1.2	3.0
2003 4/.....	2.7	2.7	0.4	*	*	*	0.3	0.7	1.6	3.0
2002.....	2.8	3.3	*	*	*	0.3	0.3	0.9	1.6	2.9
2001.....	2.7	2.2	*	*	*	*	0.4	0.9	1.8	3.2
2000.....	2.8	2.1	*	*	*	*	*	1.2	2.2	3.3
1999.....	2.8	2.1	*	*	*	*	0.4	1.2	2.4	3.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral diseases (A80-B34)										
Black, both sexes										
2006 4/.....	20.5	6.2	*	*	*	0.9	3.7	10.0	18.2	32.7
2005 4/.....	20.6	4.4	*	*	*	0.8	2.9	9.9	21.2	35.1
2004 4/.....	21.5	4.2	*	*	0.6	0.9	4.0	11.3	23.6	38.5
2003 4/.....	22.3	5.0	*	*	*	0.8	3.4	11.3	23.6	46.9
2002.....	23.5	5.8	*	*	*	1.0	3.5	12.3	29.5	49.7
2001.....	23.7	6.6	0.8	*	0.6	1.3	4.7	15.0	31.8	50.1
2000.....	24.1	7.3	0.9	*	*	0.9	4.2	15.7	37.9	50.9
1999.....	24.3	7.3	1.1	0.9	0.6	0.8	4.3	17.0	38.3	51.0
Black, male										
2006 4/.....	27.6	*	*	*	*	*	4.3	11.3	20.6	41.2
2005 4/.....	28.3	*	*	*	*	*	3.1	11.1	24.2	45.5
2004 4/.....	29.4	*	*	*	*	*	4.1	12.2	27.1	51.1
2003 4/.....	31.1	*	*	*	*	*	3.6	12.8	29.3	63.3
2002.....	33.0	6.4	*	*	*	*	3.7	12.4	37.1	69.8
2001.....	33.1	7.5	*	*	*	1.4	3.9	17.1	42.7	70.6
2000.....	34.3	7.0	*	*	*	*	4.0	16.9	48.7	72.9
1999.....	35.0	7.5	*	*	*	*	3.8	21.5	51.2	71.2
Black, female										
2006 4/.....	14.0	7.2	*	*	*	*	3.2	8.8	16.1	25.1
2005 4/.....	13.5	*	*	*	*	*	2.7	8.8	18.5	25.8
2004 4/.....	14.2	*	*	*	*	*	3.9	10.4	20.6	27.2
2003 4/.....	14.2	*	*	*	*	*	3.3	9.9	18.4	32.3
2002.....	14.9	*	*	*	*	*	3.3	12.2	22.7	31.9
2001.....	15.0	*	*	*	*	*	5.4	13.0	22.1	31.9
2000.....	14.8	7.6	*	*	*	*	4.3	14.6	28.3	31.5
1999.....	14.6	7.0	*	*	*	*	4.9	13.0	26.7	33.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral hepatitis (B15-B19)										
All races 3/, both sexes										
2006.....	2.4	*	*	*	*	*	*	*	0.2	0.6
2005.....	1.9	*	*	*	*	*	*	0.1	0.2	0.6
2004.....	1.8	*	*	*	*	*	*	0.1	0.2	0.7
2003.....	1.9	*	*	*	*	*	*	0.1	0.2	0.8
2002.....	2.0	*	*	*	*	*	*	0.2	0.3	1.0
2001.....	2.0	*	*	*	*	*	*	*	0.3	0.9
2000.....	1.9	*	*	*	*	*	*	0.1	0.3	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9
All races 3/, male										
2006.....	3.2	*	*	*	*	*	*	*	0.3	0.8
2005.....	2.5	*	*	*	*	*	*	*	*	0.8
2004.....	2.4	*	*	*	*	*	*	*	0.2	0.8
2003.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2002.....	2.7	*	*	*	*	*	*	0.2	0.4	1.4
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.2
All races 3/, female										
2006.....	1.6	*	*	*	*	*	*	*	*	0.4
2005.....	1.3	*	*	*	*	*	*	*	*	0.4
2004.....	1.3	*	*	*	*	*	*	*	*	0.5
2003.....	1.3	*	*	*	*	*	*	*	*	0.4
2002.....	1.4	*	*	*	*	*	*	*	*	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.6
2000.....	1.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Viral hepatitis (B15-B19)										
White, both sexes										
2006 4/.....	2.4	*	*	*	*	*	*	*	0.2	0.5
2005 4/.....	1.8	*	*	*	*	*	*	*	0.1	0.6
2004 4/.....	1.8	*	*	*	*	*	*	*	0.2	0.6
2003 4/.....	1.8	*	*	*	*	*	*	*	0.2	0.8
2002.....	2.0	*	*	*	*	*	*	0.1	0.3	1.0
2001.....	1.9	*	*	*	*	*	*	*	0.2	0.9
2000.....	1.8	*	*	*	*	*	*	*	0.2	1.0
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.9
White, male										
2006 4/.....	3.2	*	*	*	*	*	*	*	0.3	0.7
2005 4/.....	2.5	*	*	*	*	*	*	*	*	0.9
2004 4/.....	2.4	*	*	*	*	*	*	*	*	0.8
2003 4/.....	2.4	*	*	*	*	*	*	*	0.2	1.2
2002.....	2.6	*	*	*	*	*	*	*	0.4	1.5
2001.....	2.5	*	*	*	*	*	*	*	0.3	1.2
2000.....	2.4	*	*	*	*	*	*	*	0.3	1.3
1999.....	2.2	*	*	*	*	*	*	*	0.4	1.3
White, female										
2006 4/.....	1.6	*	*	*	*	*	*	*	*	0.4
2005 4/.....	1.2	*	*	*	*	*	*	*	*	0.3
2004 4/.....	1.2	*	*	*	*	*	*	*	*	0.5
2003 4/.....	1.2	*	*	*	*	*	*	*	*	0.5
2002.....	1.3	*	*	*	*	*	*	*	*	0.5
2001.....	1.3	*	*	*	*	*	*	*	*	0.6
2000.....	1.3	*	*	*	*	*	*	*	*	0.6
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
All races 3/, both sexes										
2006.....	4.0	*	*	*	*	0.2	0.8	2.0	3.9	7.5
2005.....	4.2	*	*	*	*	0.1	0.6	2.1	4.5	8.1
2004.....	4.4	*	*	*	*	0.1	0.8	2.2	5.1	9.5
2003.....	4.7	*	*	*	0.1	0.2	0.7	2.3	5.5	11.2
2002.....	4.9	*	*	*	0.1	0.2	0.7	2.6	6.4	11.9
2001.....	5.0	*	*	*	0.1	0.2	0.9	3.1	7.3	12.3
2000.....	5.1	*	*	*	0.1	0.2	0.8	3.3	8.7	12.8
1999.....	5.3	*	0.2	0.2	0.1	0.2	0.9	3.9	9.5	13.0
All races 3/, male										
2006.....	5.9	*	*	*	*	0.2	0.9	2.4	4.7	10.1
2005.....	6.3	*	*	*	*	*	0.7	2.4	5.5	11.2
2004.....	6.6	*	*	*	*	*	0.9	2.6	6.4	13.3
2003.....	7.1	*	*	*	*	*	0.8	2.8	7.3	15.9
2002.....	7.4	*	*	*	*	*	0.7	3.0	8.5	17.7
2001.....	7.6	*	*	*	*	0.2	0.8	3.7	10.2	18.4
2000.....	7.9	*	*	*	*	*	0.8	3.8	11.9	19.2
1999.....	8.2	*	*	0.2	*	*	0.8	5.1	13.6	19.5
All races 3/, female										
2006.....	2.2	*	*	*	*	*	0.7	1.6	3.0	4.9
2005.....	2.2	*	*	*	*	*	0.5	1.7	3.5	4.9
2004.....	2.4	*	*	*	*	*	0.7	1.8	3.7	5.6
2003.....	2.4	*	*	*	*	*	0.7	1.8	3.7	6.5
2002.....	2.5	*	*	*	*	0.2	0.6	2.3	4.2	6.2
2001.....	2.5	*	*	*	*	0.2	1.0	2.5	4.4	6.3
2000.....	2.5	*	*	*	*	*	0.7	2.9	5.5	6.4
1999.....	2.5	*	*	*	*	*	1.0	2.7	5.4	6.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
White, both sexes										
2006 4/.....	2.1	*	*	*	*	*	0.3	0.8	1.8	4.0
2005 4/.....	2.2	*	*	*	*	*	0.3	0.8	2.1	4.2
2004 4/.....	2.4	*	*	*	*	*	0.2	0.8	2.3	5.4
2003 4/.....	2.5	*	*	*	*	*	0.3	0.9	2.9	6.1
2002.....	2.6	*	*	*	*	*	0.3	1.2	3.0	6.8
2001.....	2.7	*	*	*	*	*	0.3	1.3	3.8	7.1
2000.....	2.8	*	*	*	*	*	0.3	1.5	4.5	7.6
1999.....	3.0	*	*	*	*	*	0.4	2.0	5.4	7.9
White, male										
2006 4/.....	3.5	*	*	*	*	*	0.3	1.1	2.7	6.2
2005 4/.....	3.7	*	*	*	*	*	0.4	1.1	3.0	6.8
2004 4/.....	3.9	*	*	*	*	*	0.4	1.2	3.6	8.4
2003 4/.....	4.2	*	*	*	*	*	0.3	1.4	4.5	9.7
2002.....	4.3	*	*	*	*	*	0.3	1.8	4.7	11.1
2001.....	4.5	*	*	*	*	*	0.4	1.8	5.8	11.6
2000.....	4.7	*	*	*	*	*	0.4	2.0	7.1	12.5
1999.....	5.0	*	*	*	*	*	0.4	2.8	8.6	13.0
White, female										
2006 4/.....	0.7	*	*	*	*	*	0.3	0.4	0.9	1.8
2005 4/.....	0.8	*	*	*	*	*	*	0.5	1.2	1.6
2004 4/.....	0.8	*	*	*	*	*	*	0.4	1.0	2.4
2003 4/.....	0.9	*	*	*	*	*	*	0.5	1.3	2.4
2002.....	0.9	*	*	*	*	*	*	0.7	1.3	2.3
2001.....	0.9	*	*	*	*	*	0.3	0.8	1.6	2.4
2000.....	1.0	*	*	*	*	*	*	1.0	1.9	2.6
1999.....	1.0	*	*	*	*	*	0.3	1.0	2.1	2.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease (B20-B24)										
Black, both sexes										
2006 4/.....	17.3	*	*	*	*	0.9	3.6	9.9	17.7	31.8
2005 4/.....	18.0	*	*	*	*	0.6	2.7	9.7	20.4	34.3
2004 4/.....	18.8	*	*	*	*	0.7	3.7	10.8	23.2	37.4
2003 4/.....	19.6	*	*	*	*	0.6	3.2	11.2	23.2	46.0
2002.....	20.8	*	*	*	*	0.8	3.3	11.7	29.1	48.2
2001.....	21.1	*	*	*	*	1.1	4.4	14.5	31.5	48.8
2000.....	21.4	*	*	*	*	0.7	3.7	15.2	37.1	49.8
1999.....	21.8	*	*	0.7	*	0.7	4.1	16.6	37.8	49.9
Black, male										
2006 4/.....	23.5	*	*	*	*	*	4.1	11.3	20.0	40.2
2005 4/.....	25.1	*	*	*	*	*	2.9	11.0	23.5	44.5
2004 4/.....	25.9	*	*	*	*	*	3.8	11.9	26.5	50.1
2003 4/.....	27.7	*	*	*	*	*	3.2	12.7	28.6	61.9
2002.....	29.5	*	*	*	*	*	3.3	11.7	36.6	68.4
2001.....	30.1	*	*	*	*	*	3.5	16.8	42.3	69.2
2000.....	31.0	*	*	*	*	*	3.6	16.4	47.9	71.2
1999.....	31.9	*	*	*	*	*	3.5	21.1	50.6	69.9
Black, female										
2006 4/.....	11.7	*	*	*	*	*	3.0	8.7	15.6	24.3
2005 4/.....	11.5	*	*	*	*	*	2.5	8.5	17.6	25.1
2004 4/.....	12.4	*	*	*	*	*	3.7	9.8	20.3	25.9
2003 4/.....	12.3	*	*	*	*	*	3.1	9.8	18.3	31.9
2002.....	12.8	*	*	*	*	*	3.3	11.7	22.2	30.3
2001.....	12.9	*	*	*	*	*	5.2	12.4	21.8	30.7
2000.....	12.8	*	*	*	*	*	3.8	14.1	27.4	30.8
1999.....	12.6	*	*	*	*	*	4.7	12.5	26.3	32.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)										
All races 3/, both sexes										
2006.....	1.5	*	*	*	*	*	0.3	0.8	1.6	2.9
2005.....	1.6	*	*	*	*	*	0.2	0.8	1.8	3.2
2004.....	1.7	*	*	*	*	*	0.3	0.9	2.1	3.7
2003.....	1.8	*	*	*	*	*	0.3	1.0	2.2	4.3
2002.....	1.9	*	*	*	*	*	0.3	1.1	2.5	4.9
2001.....	1.9	*	*	*	*	*	0.4	1.3	2.9	4.8
2000.....	2.0	*	*	*	*	*	0.4	1.4	3.4	5.2
1999.....	2.1	*	*	*	*	*	0.4	1.7	3.8	5.4
All races 3/, male										
2006.....	2.2	*	*	*	*	*	0.4	1.1	1.9	3.7
2005.....	2.4	*	*	*	*	*	0.3	1.0	2.1	4.5
2004.....	2.6	*	*	*	*	*	0.4	1.1	2.7	5.2
2003.....	2.7	*	*	*	*	*	0.3	1.1	3.0	5.9
2002.....	2.9	*	*	*	*	*	0.4	1.2	3.3	7.1
2001.....	2.9	*	*	*	*	*	0.4	1.6	3.9	7.1
2000.....	3.1	*	*	*	*	*	0.4	1.7	4.7	7.8
1999.....	3.3	*	*	*	*	*	0.4	2.2	5.3	8.0
All races 3/, female										
2006.....	0.9	*	*	*	*	*	0.3	0.6	1.2	2.0
2005.....	0.9	*	*	*	*	*	0.2	0.6	1.4	2.0
2004.....	1.0	*	*	*	*	*	0.3	0.8	1.6	2.1
2003.....	0.9	*	*	*	*	*	0.2	0.8	1.4	2.6
2002.....	1.0	*	*	*	*	*	0.2	1.0	1.8	2.7
2001.....	1.0	*	*	*	*	*	0.4	1.0	1.9	2.5
2000.....	1.0	*	*	*	*	*	0.4	1.1	2.1	2.6
1999.....	1.0	*	*	*	*	*	0.5	1.2	2.4	2.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

White, both sexes

2006 4/.....	0.8	*	*	*	*	*	0.2	0.4	0.8	1.6
2005 4/.....	0.9	*	*	*	*	*	0.1	0.3	0.9	1.8
2004 4/.....	1.0	*	*	*	*	*	*	0.4	1.0	2.3
2003 4/.....	1.0	*	*	*	*	*	*	0.4	1.2	2.5
2002.....	1.1	*	*	*	*	*	0.1	0.5	1.2	2.9
2001.....	1.1	*	*	*	*	*	0.2	0.6	1.6	2.9
2000.....	1.1	*	*	*	*	*	0.2	0.7	1.8	3.1
1999.....	1.2	*	*	*	*	*	0.1	0.9	2.2	3.3

White, male

2006 4/.....	1.3	*	*	*	*	*	*	0.6	1.1	2.5
2005 4/.....	1.4	*	*	*	*	*	*	0.5	1.3	2.8
2004 4/.....	1.6	*	*	*	*	*	*	0.6	1.5	3.5
2003 4/.....	1.7	*	*	*	*	*	*	0.6	1.9	3.8
2002.....	1.7	*	*	*	*	*	*	0.8	1.7	4.6
2001.....	1.8	*	*	*	*	*	*	0.9	2.4	4.6
2000.....	1.9	*	*	*	*	*	*	0.9	2.9	5.0
1999.....	2.0	*	*	*	*	*	*	1.3	3.3	5.3

White, female

2006 4/.....	0.3	*	*	*	*	*	*	*	0.4	0.8
2005 4/.....	0.3	*	*	*	*	*	*	*	0.5	0.7
2004 4/.....	0.4	*	*	*	*	*	*	*	0.5	1.0
2003 4/.....	0.4	*	*	*	*	*	*	*	0.5	1.1
2002.....	0.4	*	*	*	*	*	*	0.3	0.7	1.2
2001.....	0.4	*	*	*	*	*	*	0.3	0.7	1.2
2000.....	0.4	*	*	*	*	*	*	0.5	0.8	1.2
1999.....	0.4	*	*	*	*	*	*	0.5	1.0	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)										
Black, both sexes										
2006 4/.....	6.3	*	*	*	*	*	1.3	3.8	7.1	11.2
2005 4/.....	6.7	*	*	*	*	*	1.0	4.0	7.5	13.3
2004 4/.....	7.2	*	*	*	*	*	1.8	4.1	9.8	13.5
2003 4/.....	7.1	*	*	*	*	*	1.1	4.4	8.9	16.5
2002.....	8.0	*	*	*	*	*	1.4	4.5	11.4	18.6
2001.....	8.0	*	*	*	*	*	1.8	5.6	12.0	18.1
2000.....	8.3	*	*	*	*	*	1.6	6.1	14.2	20.2
1999.....	8.6	*	*	*	*	*	2.1	6.9	15.1	20.1
Black, male										
2006 4/.....	8.4	*	*	*	*	*	1.5	4.7	7.7	12.8
2005 4/.....	9.3	*	*	*	*	*	*	4.4	8.2	17.0
2004 4/.....	9.7	*	*	*	*	*	1.7	4.0	11.3	18.1
2003 4/.....	10.2	*	*	*	*	*	*	4.9	11.4	21.5
2002.....	11.4	*	*	*	*	*	1.6	4.0	14.8	25.8
2001.....	11.3	*	*	*	*	*	1.9	6.1	15.2	26.2
2000.....	11.9	*	*	*	*	*	1.5	7.1	18.7	29.5
1999.....	12.6	*	*	*	*	*	1.7	8.3	19.7	28.5
Black, female										
2006 4/.....	4.3	*	*	*	*	*	*	2.9	6.5	9.8
2005 4/.....	4.4	*	*	*	*	*	*	3.7	6.9	9.9
2004 4/.....	4.9	*	*	*	*	*	1.8	4.3	8.4	9.4
2003 4/.....	4.3	*	*	*	*	*	*	3.9	6.6	12.0
2002.....	4.9	*	*	*	*	*	*	5.0	8.3	12.3
2001.....	4.9	*	*	*	*	*	1.7	5.2	9.1	11.0
2000.....	5.0	*	*	*	*	*	1.8	5.2	10.2	11.9
1999.....	5.0	*	*	*	*	*	2.5	5.6	11.0	12.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in other specified diseases (B22)

All races 3/, both sexes

2006.....	0.4	*	*	*	*	*	0.1	0.2	0.4	0.8
2005.....	0.4	*	*	*	*	*	*	0.2	0.4	0.8
2004.....	0.4	*	*	*	*	*	*	0.2	0.5	0.9
2003.....	0.5	*	*	*	*	*	*	0.2	0.5	1.2
2002.....	0.5	*	*	*	*	*	*	0.3	0.7	1.0
2001.....	0.5	*	*	*	*	*	*	0.3	0.8	1.3
2000.....	0.6	*	*	*	*	*	*	0.3	1.1	1.5
1999.....	0.6	*	*	*	*	*	*	0.4	1.1	1.5

All races 3/, male

2006.....	0.6	*	*	*	*	*	*	0.2	0.6	1.2
2005.....	0.6	*	*	*	*	*	*	0.2	0.4	1.1
2004.....	0.7	*	*	*	*	*	*	0.3	0.6	1.3
2003.....	0.7	*	*	*	*	*	*	0.3	0.6	1.8
2002.....	0.8	*	*	*	*	*	*	0.4	1.1	1.7
2001.....	0.8	*	*	*	*	*	*	0.4	1.2	2.2
2000.....	0.9	*	*	*	*	*	*	0.4	1.6	2.3
1999.....	0.9	*	*	*	*	*	*	0.6	1.6	2.5

All races 3/, female

2006.....	0.2	*	*	*	*	*	*	*	0.2	0.3
2005.....	0.2	*	*	*	*	*	*	*	0.3	0.5
2004.....	0.2	*	*	*	*	*	*	*	0.4	0.4
2003.....	0.2	*	*	*	*	*	*	*	0.3	0.6
2002.....	0.2	*	*	*	*	*	*	0.3	0.4	0.4
2001.....	0.2	*	*	*	*	*	*	*	0.4	0.5
2000.....	0.2	*	*	*	*	*	*	0.2	0.6	0.7
1999.....	0.2	*	*	*	*	*	*	0.2	0.5	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

All races 3/, both sexes

2006.....	0.7	*	*	*	*	*	*	0.3	0.5	1.2
2005.....	0.7	*	*	*	*	*	*	0.3	0.7	1.1
2004.....	0.7	*	*	*	*	*	*	0.3	0.8	1.4
2003.....	1.1	*	*	*	*	*	0.2	0.5	1.2	2.4
2002.....	1.2	*	*	*	*	*	0.1	0.6	1.4	2.6
2001.....	1.2	*	*	*	*	*	0.3	0.8	1.6	2.9
2000.....	1.2	*	*	*	*	*	0.1	0.7	1.9	2.9
1999.....	1.2	*	*	*	*	*	0.2	0.8	2.0	2.8

All races 3/, male

2006.....	1.0	*	*	*	*	*	*	0.4	0.5	1.6
2005.....	1.0	*	*	*	*	*	*	0.3	0.8	1.4
2004.....	1.1	*	*	*	*	*	*	0.4	1.0	1.9
2003.....	1.6	*	*	*	*	*	*	0.5	1.5	3.3
2002.....	1.7	*	*	*	*	*	*	0.6	1.9	3.9
2001.....	1.8	*	*	*	*	*	*	0.8	2.2	4.2
2000.....	1.8	*	*	*	*	*	*	0.7	2.4	4.1
1999.....	1.8	*	*	*	*	*	*	1.1	2.7	3.9

All races 3/, female

2006.....	0.4	*	*	*	*	*	*	0.2	0.6	0.8
2005.....	0.4	*	*	*	*	*	*	0.3	0.6	0.8
2004.....	0.4	*	*	*	*	*	*	0.3	0.5	0.9
2003.....	0.6	*	*	*	*	*	0.2	0.5	0.9	1.5
2002.....	0.7	*	*	*	*	*	*	0.5	0.9	1.4
2001.....	0.7	*	*	*	*	*	0.4	0.7	1.0	1.7
2000.....	0.7	*	*	*	*	*	*	0.7	1.4	1.6
1999.....	0.7	*	*	*	*	*	0.2	0.6	1.3	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)										
White, both sexes										
2006 4/.....	0.3	*	*	*	*	*	*	*	0.2	0.5
2005 4/.....	0.3	*	*	*	*	*	*	*	0.2	0.5
2004 4/.....	0.3	*	*	*	*	*	*	*	0.3	0.6
2003 4/.....	0.5	*	*	*	*	*	*	0.1	0.5	1.1
2002.....	0.5	*	*	*	*	*	*	0.2	0.5	1.3
2001.....	0.6	*	*	*	*	*	*	0.2	0.7	1.5
2000.....	0.6	*	*	*	*	*	*	0.3	0.8	1.5
1999.....	0.6	*	*	*	*	*	*	0.3	0.9	1.4
White, male										
2006 4/.....	0.5	*	*	*	*	*	*	*	*	0.7
2005 4/.....	0.5	*	*	*	*	*	*	*	0.3	0.7
2004 4/.....	0.5	*	*	*	*	*	*	*	0.4	0.9
2003 4/.....	0.8	*	*	*	*	*	*	*	0.7	1.7
2002.....	0.9	*	*	*	*	*	*	*	0.8	2.2
2001.....	0.9	*	*	*	*	*	*	*	1.1	2.4
2000.....	0.9	*	*	*	*	*	*	0.3	1.2	2.5
1999.....	0.9	*	*	*	*	*	*	0.5	1.4	2.2
White, female										
2006 4/.....	0.1	*	*	*	*	*	*	*	*	0.3
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*	*	*	0.3
2003 4/.....	0.2	*	*	*	*	*	*	*	0.3	0.5
2002.....	0.2	*	*	*	*	*	*	*	*	0.5
2001.....	0.2	*	*	*	*	*	*	*	0.4	0.6
2000.....	0.2	*	*	*	*	*	*	*	0.5	0.6
1999.....	0.2	*	*	*	*	*	*	*	0.4	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)										
Black, both sexes										
2006 4/.....	3.5	*	*	*	*	*	*	1.8	3.0	6.4
2005 4/.....	3.4	*	*	*	*	*	*	1.6	3.7	5.2
2004 4/.....	3.5	*	*	*	*	*	*	1.8	3.9	6.9
2003 4/.....	5.3	*	*	*	*	*	1.0	2.8	6.0	11.5
2002.....	5.7	*	*	*	*	*	*	3.0	7.1	11.7
2001.....	5.7	*	*	*	*	*	1.2	4.1	7.6	12.8
2000.....	5.5	*	*	*	*	*	*	3.9	9.1	12.3
1999.....	5.8	*	*	*	*	*	0.9	4.1	9.2	12.7
Black, male										
2006 4/.....	4.7	*	*	*	*	*	*	2.2	2.8	8.8
2005 4/.....	4.8	*	*	*	*	*	*	1.6	4.0	6.2
2004 4/.....	5.0	*	*	*	*	*	*	1.9	5.2	9.3
2003 4/.....	7.3	*	*	*	*	*	*	2.5	7.2	15.1
2002.....	8.0	*	*	*	*	*	*	3.3	9.2	17.0
2001.....	8.0	*	*	*	*	*	*	4.7	10.4	17.9
2000.....	7.8	*	*	*	*	*	*	4.1	11.0	16.7
1999.....	8.3	*	*	*	*	*	*	5.4	11.9	17.0
Black, female										
2006 4/.....	2.4	*	*	*	*	*	*	1.4	3.2	4.1
2005 4/.....	2.1	*	*	*	*	*	*	1.6	3.3	4.3
2004 4/.....	2.2	*	*	*	*	*	*	1.8	2.7	4.8
2003 4/.....	3.6	*	*	*	*	*	*	3.0	4.9	8.2
2002.....	3.6	*	*	*	*	*	*	2.6	5.2	7.0
2001.....	3.6	*	*	*	*	*	1.9	3.6	5.1	8.2
2000.....	3.5	*	*	*	*	*	*	3.7	7.3	8.4
1999.....	3.6	*	*	*	*	*	*	2.9	6.7	8.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Unspecified human immunodeficiency virus (HIV) disease (B24)										
All races 3/, both sexes										
2006.....	1.1	*	*	*	*	*	0.2	0.6	1.1	2.2
2005.....	1.2	*	*	*	*	*	0.2	0.6	1.4	2.3
2004.....	1.2	*	*	*	*	*	0.2	0.6	1.4	2.8
2003.....	1.0	*	*	*	*	*	0.2	0.5	1.4	2.7
2002.....	1.0	*	*	*	*	*	0.2	0.6	1.4	2.7
2001.....	1.0	*	*	*	*	*	0.2	0.6	1.5	2.5
2000.....	1.0	*	*	*	*	*	0.1	0.7	1.8	2.4
1999.....	1.0	*	*	*	*	*	0.2	0.8	1.9	2.3
All races 3/, male										
2006.....	1.6	*	*	*	*	*	0.2	0.6	1.3	2.8
2005.....	1.7	*	*	*	*	*	0.2	0.7	1.7	3.2
2004.....	1.7	*	*	*	*	*	0.3	0.8	1.7	3.8
2003.....	1.5	*	*	*	*	*	*	0.7	1.8	3.8
2002.....	1.5	*	*	*	*	*	*	0.7	1.7	3.9
2001.....	1.5	*	*	*	*	*	*	0.7	2.2	3.6
2000.....	1.5	*	*	*	*	*	*	0.7	2.4	3.6
1999.....	1.5	*	*	*	*	*	*	0.9	2.7	3.5
All races 3/, female										
2006.....	0.6	*	*	*	*	*	0.2	0.6	0.9	1.5
2005.....	0.6	*	*	*	*	*	*	0.5	1.0	1.4
2004.....	0.7	*	*	*	*	*	*	0.5	1.2	1.9
2003.....	0.5	*	*	*	*	*	*	0.4	1.0	1.5
2002.....	0.5	*	*	*	*	*	*	0.5	1.1	1.5
2001.....	0.5	*	*	*	*	*	0.2	0.4	0.9	1.4
2000.....	0.5	*	*	*	*	*	*	0.6	1.2	1.2
1999.....	0.5	*	*	*	*	*	*	0.6	1.1	1.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Unspecified human immunodeficiency virus (HIV) disease (B24)										
White, both sexes										
2006 4/.....	0.5	*	*	*	*	*	*	0.2	0.5	1.0
2005 4/.....	0.6	*	*	*	*	*	*	0.2	0.6	1.1
2004 4/.....	0.6	*	*	*	*	*	*	0.2	0.6	1.5
2003 4/.....	0.5	*	*	*	*	*	*	0.2	0.7	1.2
2002.....	0.5	*	*	*	*	*	*	0.3	0.6	1.3
2001.....	0.5	*	*	*	*	*	*	0.2	0.7	1.2
2000.....	0.5	*	*	*	*	*	*	0.3	0.8	1.3
1999.....	0.5	*	*	*	*	*	*	0.4	1.1	1.3
White, male										
2006 4/.....	0.9	*	*	*	*	*	*	0.3	0.6	1.6
2005 4/.....	1.0	*	*	*	*	*	*	0.3	0.9	1.8
2004 4/.....	1.0	*	*	*	*	*	*	*	0.9	2.2
2003 4/.....	0.9	*	*	*	*	*	*	0.3	1.1	2.0
2002.....	0.8	*	*	*	*	*	*	0.4	0.9	2.2
2001.....	0.8	*	*	*	*	*	*	0.3	1.1	2.1
2000.....	0.8	*	*	*	*	*	*	0.5	1.3	2.1
1999.....	0.9	*	*	*	*	*	*	0.5	1.7	2.2
White, female										
2006 4/.....	0.2	*	*	*	*	*	*	*	0.3	0.5
2005 4/.....	0.2	*	*	*	*	*	*	*	0.4	0.4
2004 4/.....	0.2	*	*	*	*	*	*	*	0.3	0.8
2003 4/.....	0.2	*	*	*	*	*	*	*	0.3	0.5
2002.....	0.2	*	*	*	*	*	*	*	0.3	0.4
2001.....	0.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	0.2	*	*	*	*	*	*	*	0.4	0.4
1999.....	0.2	*	*	*	*	*	*	*	0.4	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Unspecified human immunodeficiency virus (HIV) disease (B24)										
Black, both sexes										
2006 4/.....	5.0	*	*	*	*	*	1.2	2.9	5.4	10.0
2005 4/.....	5.1	*	*	*	*	*	0.9	2.9	6.2	10.5
2004 4/.....	5.4	*	*	*	*	*	1.1	3.4	6.8	12.0
2003 4/.....	4.5	*	*	*	*	*	0.7	2.5	5.8	12.6
2002.....	4.3	*	*	*	*	*	0.8	2.4	6.9	12.2
2001.....	4.4	*	*	*	*	*	0.9	2.8	7.2	11.1
2000.....	4.3	*	*	*	*	*	0.8	2.9	8.4	10.1
1999.....	4.1	*	*	*	*	*	0.7	3.3	7.5	9.5
Black, male										
2006 4/.....	6.6	*	*	*	*	*	*	2.8	6.3	12.2
2005 4/.....	7.1	*	*	*	*	*	*	3.4	7.6	14.0
2004 4/.....	7.3	*	*	*	*	*	*	4.2	7.1	15.6
2003 4/.....	6.3	*	*	*	*	*	*	3.2	6.8	17.5
2002.....	6.0	*	*	*	*	*	*	2.8	7.8	16.7
2001.....	6.3	*	*	*	*	*	*	3.5	9.9	14.5
2000.....	6.3	*	*	*	*	*	*	2.5	10.6	14.6
1999.....	5.9	*	*	*	*	*	*	4.0	10.1	13.1
Black, female										
2006 4/.....	3.5	*	*	*	*	*	1.3	3.1	4.5	8.0
2005 4/.....	3.3	*	*	*	*	*	*	2.4	4.9	7.5
2004 4/.....	3.7	*	*	*	*	*	*	2.7	6.5	8.8
2003 4/.....	2.8	*	*	*	*	*	*	1.9	4.8	8.2
2002.....	2.8	*	*	*	*	*	*	2.1	6.2	8.2
2001.....	2.7	*	*	*	*	*	*	2.2	4.8	8.1
2000.....	2.6	*	*	*	*	*	*	3.3	6.3	6.1
1999.....	2.4	*	*	*	*	*	*	2.7	5.2	6.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

All races 3/, both sexes

2006.....	1.0	0.8	*	*	*	*	*	0.1	0.2	0.2
2005.....	1.6	1.0	*	*	*	*	0.1	0.1	0.2	0.4
2004.....	1.5	1.0	*	*	*	*	0.1	0.2	0.2	0.5
2003.....	1.5	1.1	*	*	*	*	*	0.2	0.3	0.5
2002.....	1.3	1.3	*	*	*	*	0.1	0.2	0.3	0.4
2001.....	1.1	1.6	*	*	*	*	0.1	0.2	0.3	0.5
2000.....	1.0	1.3	*	*	*	*	*	0.1	0.3	0.4
1999.....	1.4	1.6	0.1	*	*	*	0.1	0.3	0.5	0.7

All races 3/, male

2006.....	1.0	0.9	*	*	*	*	*	*	0.2	0.3
2005.....	1.8	1.2	*	*	*	*	*	*	0.4	0.6
2004.....	1.8	1.3	*	*	*	*	0.2	*	0.3	0.6
2003.....	1.7	1.3	*	*	*	*	*	0.2	0.3	0.6
2002.....	1.5	1.5	*	*	*	*	*	0.2	0.4	0.6
2001.....	1.2	1.6	*	*	*	*	*	0.2	0.3	0.6
2000.....	1.1	1.5	*	*	*	*	*	*	0.3	0.6
1999.....	1.6	1.9	*	*	*	*	*	0.4	0.5	0.8

All races 3/, female

2006.....	0.9	*	*	*	*	*	*	*	0.2	*
2005.....	1.3	*	*	*	*	*	*	*	*	0.3
2004.....	1.3	*	*	*	*	*	*	*	*	0.4
2003.....	1.3	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.0	1.2	*	*	*	*	*	*	0.3	0.3
2001.....	1.0	1.6	*	*	*	*	*	*	0.2	0.4
2000.....	0.9	1.1	*	*	*	*	*	*	0.2	0.3
1999.....	1.3	1.2	*	*	*	*	*	0.3	0.4	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

Black, both sexes

2006 4/.....	1.1	*	*	*	*	*	*	*	*	*	0.7
2005 4/.....	1.5	*	*	*	*	*	*	*	*	*	0.7
2004 4/.....	1.5	*	*	*	*	*	*	*	*	*	0.8
2003 4/.....	1.5	*	*	*	*	*	*	*	*	*	0.9
2002.....	1.5	*	*	*	*	*	*	*	*	0.8	1.0
2001.....	1.3	3.7	*	*	*	*	*	*	*	0.8	1.0
2000.....	1.3	3.3	*	*	*	*	*	*	*	0.8	1.2
1999.....	2.2	3.8	*	*	*	*	*	0.9	1.7	1.9	1.9

Black, male

2006 4/.....	1.2	*	*	*	*	*	*	*	*	*	*
2005 4/.....	1.8	*	*	*	*	*	*	*	*	*	*
2004 4/.....	1.8	*	*	*	*	*	*	*	*	*	*
2003 4/.....	1.8	*	*	*	*	*	*	*	*	*	*
2002.....	1.8	*	*	*	*	*	*	*	*	*	*
2001.....	1.5	*	*	*	*	*	*	*	*	*	*
2000.....	1.5	*	*	*	*	*	*	*	*	*	1.7
1999.....	2.3	*	*	*	*	*	*	*	1.9	1.9	1.9

Black, female

2006 4/.....	1.0	*	*	*	*	*	*	*	*	*	*
2005 4/.....	1.3	*	*	*	*	*	*	*	*	*	*
2004 4/.....	1.3	*	*	*	*	*	*	*	*	*	*
2003 4/.....	1.4	*	*	*	*	*	*	*	*	*	*
2002.....	1.1	*	*	*	*	*	*	*	*	*	*
2001.....	1.1	*	*	*	*	*	*	*	*	*	*
2000.....	1.1	*	*	*	*	*	*	*	*	*	*
1999.....	2.2	*	*	*	*	*	*	*	1.4	1.9	1.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
II. Neoplasms (C00-D48)										
All races 3/, both sexes										
2006.....	191.7	3.4	2.7	2.5	2.4	3.4	4.8	7.0	11.9	21.3
2005.....	193.3	3.3	2.6	2.7	2.6	3.7	4.9	7.0	11.7	21.5
2004.....	193.2	3.6	2.8	2.9	2.5	3.8	4.9	6.9	12.0	22.1
2003.....	196.2	3.4	2.8	2.8	2.8	3.6	4.9	7.2	12.3	23.6
2002.....	197.9	3.5	3.0	2.9	2.7	3.7	5.2	7.0	13.0	24.1
2001.....	199.2	2.9	3.1	2.7	2.7	3.8	5.2	7.7	13.0	25.3
2000.....	201.3	4.0	3.1	2.7	2.7	3.9	5.4	7.4	12.9	25.2
1999.....	201.8	3.3	3.1	2.8	2.7	4.0	5.5	7.6	13.1	25.8
All races 3/, male										
2006.....	201.4	3.4	2.8	2.8	2.5	4.0	5.6	7.3	10.7	17.8
2005.....	203.6	3.2	3.0	2.8	2.9	4.4	5.7	7.7	10.9	18.1
2004.....	203.1	3.5	2.9	3.2	2.7	4.3	5.8	7.1	10.9	18.7
2003.....	206.0	3.2	3.2	3.1	2.9	4.4	5.4	7.2	11.5	20.0
2002.....	208.5	3.5	3.1	3.1	3.0	4.2	6.0	7.4	11.7	20.5
2001.....	210.1	2.8	3.3	2.7	2.9	4.5	6.1	7.6	11.6	21.7
2000.....	212.0	4.2	3.4	2.9	3.0	4.6	6.3	7.4	11.8	22.0
1999.....	213.7	3.4	3.3	2.9	2.8	4.6	6.4	8.0	12.2	22.3
All races 3/, female										
2006.....	182.3	3.4	2.6	2.2	2.2	2.8	3.9	6.8	13.1	24.9
2005.....	183.4	3.3	2.3	2.5	2.4	3.0	4.0	6.3	12.6	25.0
2004.....	183.7	3.7	2.7	2.5	2.3	3.2	4.0	6.6	13.1	25.5
2003.....	186.6	3.6	2.4	2.5	2.7	2.7	4.4	7.2	13.1	27.3
2002.....	187.6	3.6	2.8	2.7	2.4	3.2	4.4	6.5	14.2	27.8
2001.....	188.7	3.0	2.9	2.7	2.6	3.2	4.3	7.7	14.5	28.8
2000.....	191.1	3.8	2.8	2.5	2.4	3.2	4.4	7.4	14.1	28.4
1999.....	190.3	3.2	2.9	2.6	2.5	3.3	4.6	7.2	13.9	29.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
II. Neoplasms (C00-D48)										
White, both sexes										
2006 4/.....	204.5	3.6	2.7	2.6	2.4	3.4	4.8	6.9	11.6	20.7
2005 4/.....	205.9	3.3	2.7	2.7	2.7	3.8	4.9	6.6	11.3	20.8
2004 4/.....	205.7	3.8	2.8	2.9	2.5	3.8	4.8	6.7	11.6	21.4
2003 4/.....	208.9	3.6	2.9	2.9	2.8	3.7	5.0	7.1	11.9	22.8
2002.....	210.6	3.7	2.9	3.0	2.7	3.8	5.4	6.8	12.6	23.3
2001.....	211.7	2.9	3.4	2.7	2.7	3.9	5.2	7.5	12.6	24.4
2000.....	213.9	4.1	3.1	2.8	2.8	3.8	5.3	7.4	12.6	24.0
1999.....	213.9	3.2	3.3	2.8	2.6	3.8	5.4	7.6	12.6	24.6
White, male										
2006 4/.....	213.9	3.6	2.8	2.9	2.5	4.1	5.8	7.3	10.5	17.6
2005 4/.....	215.8	3.3	3.0	2.8	2.9	4.5	5.7	7.4	10.5	17.8
2004 4/.....	215.0	3.8	2.9	3.3	2.7	4.4	5.6	7.2	11.0	18.6
2003 4/.....	218.3	3.4	3.2	3.3	2.9	4.6	5.4	7.4	11.2	20.0
2002.....	220.6	3.9	3.0	3.1	3.0	4.4	6.2	7.2	11.6	19.9
2001.....	221.7	2.7	3.5	2.7	2.6	4.5	6.1	7.5	11.6	21.2
2000.....	223.3	4.4	3.5	2.9	3.1	4.4	6.3	7.4	11.8	21.2
1999.....	224.8	3.4	3.5	3.0	2.8	4.3	6.2	7.9	11.9	21.7
White, female										
2006 4/.....	195.3	3.5	2.5	2.3	2.2	2.7	3.8	6.4	12.7	23.9
2005 4/.....	196.2	3.3	2.3	2.6	2.4	3.1	4.0	5.8	12.1	23.9
2004 4/.....	196.7	3.9	2.8	2.5	2.2	3.1	3.9	6.2	12.4	24.3
2003 4/.....	199.7	3.8	2.5	2.6	2.7	2.8	4.4	6.8	12.6	25.7
2002.....	200.9	3.5	2.7	2.9	2.5	3.2	4.6	6.4	13.6	26.9
2001.....	201.9	3.1	3.2	2.8	2.7	3.2	4.3	7.6	13.7	27.8
2000.....	204.8	3.9	2.7	2.6	2.5	3.1	4.4	7.3	13.4	26.9
1999.....	203.3	2.9	3.2	2.6	2.4	3.3	4.5	7.2	13.4	27.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
II. Neoplasms (C00-D48)										
Black, both sexes										
2006 4/.....	162.6	3.2	2.8	2.5	2.4	3.5	5.2	8.5	15.6	29.0
2005 4/.....	164.6	3.8	2.6	2.8	2.8	3.3	5.2	9.6	16.1	29.2
2004 4/.....	165.0	*	2.8	2.8	2.9	4.2	6.1	9.4	16.1	29.4
2003 4/.....	167.2	*	2.6	2.6	3.0	3.2	5.2	9.1	16.2	32.3
2002.....	168.9	3.1	3.5	2.8	2.8	3.8	4.8	8.5	17.0	32.4
2001.....	170.2	*	2.1	2.8	3.2	4.0	5.4	9.4	16.8	33.6
2000.....	172.4	3.3	3.2	2.4	2.5	4.6	6.2	8.6	16.3	35.3
1999.....	174.5	4.3	2.5	2.8	3.2	4.9	7.0	8.9	16.6	36.9
Black, male										
2006 4/.....	175.4	*	3.0	2.8	2.6	4.1	5.6	7.9	13.3	22.7
2005 4/.....	178.1	*	2.9	3.1	3.3	3.5	6.2	9.8	15.3	22.6
2004 4/.....	179.2	*	3.2	2.7	2.9	4.4	7.2	8.3	12.9	21.0
2003 4/.....	181.2	*	3.0	2.8	3.3	3.7	5.8	7.7	14.2	22.8
2002.....	184.3	*	3.3	3.6	3.0	4.3	5.4	8.6	14.3	26.2
2001.....	187.5	*	2.6	2.9	3.8	4.7	6.3	8.7	13.0	27.8
2000.....	191.5	*	3.3	2.7	2.8	5.7	7.3	7.7	13.3	28.6
1999.....	194.1	*	2.8	2.6	3.5	6.1	7.8	9.4	14.9	28.8
Black, female										
2006 4/.....	150.9	*	2.6	2.2	2.1	2.9	4.8	9.0	17.6	34.6
2005 4/.....	152.3	*	2.3	2.6	2.3	3.0	4.1	9.5	16.8	35.1
2004 4/.....	152.1	*	2.4	2.8	2.9	3.9	5.1	10.4	18.9	37.0
2003 4/.....	154.4	*	2.1	2.3	2.8	2.6	4.6	10.3	18.1	40.6
2002.....	155.0	*	3.7	1.9	2.5	3.3	4.2	8.5	19.3	37.9
2001.....	154.6	*	1.6	2.6	2.5	3.2	4.5	10.0	20.3	38.8
2000.....	155.0	*	3.0	2.2	2.3	3.6	5.2	9.5	19.0	41.2
1999.....	156.8	*	2.1	2.9	2.8	3.7	6.3	8.4	18.1	44.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
All races 3/, both sexes										
2006.....	187.0	1.8	2.3	2.3	2.2	3.2	4.6	6.7	11.5	20.8
2005.....	188.7	1.8	2.3	2.5	2.5	3.5	4.7	6.7	11.3	20.8
2004.....	188.6	1.8	2.5	2.7	2.3	3.5	4.7	6.6	11.5	21.5
2003.....	191.5	1.9	2.5	2.6	2.6	3.4	4.6	6.8	11.8	23.0
2002.....	193.2	1.8	2.6	2.7	2.5	3.5	5.0	6.6	12.5	23.5
2001.....	194.4	1.6	2.7	2.4	2.5	3.6	4.9	7.3	12.6	24.6
2000.....	196.5	2.4	2.7	2.4	2.6	3.7	5.1	7.0	12.5	24.5
1999.....	197.0	1.8	2.7	2.5	2.5	3.7	5.3	7.3	12.5	25.2
All races 3/, male										
2006.....	196.6	1.8	2.5	2.6	2.4	3.8	5.4	7.0	10.3	17.2
2005.....	198.9	2.1	2.6	2.6	2.8	4.1	5.5	7.3	10.4	17.4
2004.....	198.4	1.8	2.6	3.0	2.5	4.1	5.5	6.8	10.4	18.1
2003.....	201.3	1.7	2.8	2.9	2.7	4.2	5.1	6.8	10.9	19.3
2002.....	203.8	2.0	2.7	2.9	2.8	4.0	5.7	7.0	11.3	19.8
2001.....	205.3	1.5	2.9	2.4	2.6	4.3	5.7	7.2	11.2	21.0
2000.....	207.2	2.6	3.0	2.6	2.9	4.3	5.9	7.1	11.3	21.2
1999.....	208.9	1.9	2.9	2.6	2.7	4.3	6.1	7.6	11.6	21.6
All races 3/, female										
2006.....	177.6	1.8	2.1	2.1	2.0	2.5	3.7	6.4	12.7	24.4
2005.....	178.8	1.5	2.0	2.3	2.2	2.8	3.9	6.0	12.2	24.3
2004.....	179.1	1.9	2.4	2.3	2.2	2.9	3.8	6.3	12.6	24.9
2003.....	182.0	2.1	2.1	2.3	2.5	2.5	4.2	6.8	12.6	26.7
2002.....	183.0	1.6	2.4	2.5	2.2	3.0	4.2	6.2	13.7	27.2
2001.....	183.9	1.8	2.5	2.4	2.3	2.9	4.1	7.4	14.0	28.1
2000.....	186.2	2.3	2.5	2.2	2.2	3.0	4.2	7.0	13.7	27.8
1999.....	185.6	1.7	2.5	2.3	2.3	3.0	4.4	7.0	13.4	28.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
White, both sexes										
2006 4/.....	199.3	2.0	2.3	2.4	2.2	3.2	4.6	6.6	11.2	20.1
2005 4/.....	200.8	2.0	2.3	2.5	2.5	3.6	4.7	6.3	10.8	20.1
2004 4/.....	200.7	2.0	2.5	2.7	2.3	3.6	4.5	6.4	11.1	20.8
2003 4/.....	203.7	2.1	2.6	2.8	2.6	3.5	4.7	6.7	11.3	22.2
2002.....	205.5	1.8	2.5	2.8	2.5	3.6	5.2	6.5	12.1	22.7
2001.....	206.4	1.7	3.0	2.5	2.4	3.7	5.0	7.2	12.2	23.7
2000.....	208.6	2.5	2.8	2.5	2.6	3.6	5.1	7.0	12.1	23.3
1999.....	208.7	1.8	2.9	2.5	2.4	3.6	5.2	7.3	12.1	24.0
White, male										
2006 4/.....	208.7	2.0	2.5	2.6	2.4	3.9	5.5	7.1	10.2	16.9
2005 4/.....	210.6	2.3	2.6	2.6	2.8	4.2	5.4	7.0	10.0	17.1
2004 4/.....	209.9	1.9	2.5	3.1	2.5	4.2	5.3	6.8	10.4	18.0
2003 4/.....	213.1	1.8	2.9	3.1	2.7	4.3	5.1	7.0	10.6	19.3
2002.....	215.5	2.2	2.7	2.9	2.7	4.2	5.9	6.9	11.1	19.2
2001.....	216.4	1.5	3.1	2.4	2.4	4.4	5.8	7.2	11.2	20.5
2000.....	218.1	2.8	3.1	2.6	2.9	4.2	5.9	7.1	11.2	20.5
1999.....	219.5	2.0	3.2	2.7	2.6	4.1	5.9	7.6	11.3	21.1
White, female										
2006 4/.....	190.1	2.0	2.0	2.1	2.0	2.5	3.6	6.1	12.4	23.4
2005 4/.....	191.1	1.7	2.0	2.4	2.3	2.9	3.9	5.5	11.7	23.3
2004 4/.....	191.7	2.1	2.5	2.3	2.1	2.9	3.7	6.0	11.9	23.7
2003 4/.....	194.6	2.3	2.3	2.4	2.5	2.6	4.2	6.5	12.0	25.2
2002.....	195.8	1.4	2.3	2.6	2.4	3.0	4.4	6.1	13.2	26.3
2001.....	196.7	2.0	2.8	2.5	2.4	2.9	4.1	7.2	13.3	27.1
2000.....	199.4	2.3	2.4	2.3	2.3	3.0	4.2	7.0	13.0	26.3
1999.....	198.2	1.5	2.7	2.2	2.2	3.0	4.3	7.0	12.9	27.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms (C00-C97)										
Black, both sexes										
2006 4/.....	159.5	*	2.5	2.3	2.1	3.1	4.9	8.0	15.0	28.1
2005 4/.....	161.7	*	2.4	2.6	2.6	3.0	5.1	9.2	15.6	28.1
2004 4/.....	161.9	*	2.5	2.5	2.6	3.8	5.9	8.9	15.1	28.4
2003 4/.....	164.3	*	2.2	2.3	2.8	2.9	4.9	8.5	15.7	31.0
2002.....	165.9	*	2.9	2.6	2.5	3.6	4.5	8.0	16.3	31.7
2001.....	167.1	*	1.9	2.4	2.8	3.6	5.1	9.1	16.4	32.7
2000.....	169.3	*	2.8	2.1	2.3	4.4	5.9	8.0	15.8	34.5
1999.....	171.3	*	2.0	2.4	3.0	4.5	6.6	8.6	15.8	35.8
Black, male										
2006 4/.....	172.3	*	2.6	2.5	2.4	3.7	5.4	7.5	12.8	21.9
2005 4/.....	175.4	*	2.8	2.9	3.2	3.4	6.1	9.3	14.7	21.7
2004 4/.....	176.2	*	2.8	2.5	2.4	4.1	6.8	7.9	12.2	19.7
2003 4/.....	178.3	*	2.4	2.4	3.1	3.5	5.6	7.0	13.5	21.7
2002.....	181.5	*	2.6	3.3	2.9	4.1	5.0	8.2	14.1	25.6
2001.....	184.5	*	2.4	2.5	3.4	4.3	5.9	8.5	12.4	26.8
2000.....	188.5	*	2.7	2.4	2.7	5.2	7.1	7.3	12.9	27.6
1999.....	191.0	*	2.1	2.3	3.3	5.6	7.3	8.8	14.0	27.8
Black, female										
2006 4/.....	147.7	*	2.3	2.0	1.8	2.4	4.4	8.5	16.9	33.6
2005 4/.....	149.1	*	2.1	2.3	2.0	2.7	4.1	9.0	16.3	33.8
2004 4/.....	148.9	*	2.2	2.5	2.7	3.5	5.0	9.9	17.8	36.2
2003 4/.....	151.4	*	2.1	2.1	2.5	2.3	4.3	10.0	17.7	39.2
2002.....	151.7	*	3.2	1.9	2.1	3.1	4.0	7.9	18.3	37.1
2001.....	151.3	*	*	2.2	2.2	2.9	4.3	9.7	20.0	37.9
2000.....	151.8	*	2.9	1.8	2.0	3.5	4.7	8.5	18.3	40.5
1999.....	153.3	*	1.9	2.6	2.7	3.4	5.9	8.3	17.5	43.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of digestive organs (C15-C26)										
All races 3/, both sexes										
2006.....	44.8	*	*	*	*	0.1	0.4	1.0	2.2	4.4
2005.....	44.8	*	0.1	*	*	0.1	0.4	0.9	2.2	4.2
2004.....	44.7	*	0.1	*	*	0.2	0.5	0.9	2.1	4.3
2003.....	45.5	*	0.1	*	*	0.1	0.4	1.0	2.1	4.6
2002.....	45.7	*	0.2	*	*	0.1	0.5	1.0	2.0	4.6
2001.....	46.0	*	*	*	*	0.2	0.4	0.9	2.1	5.0
2000.....	46.4	*	0.2	*	*	0.2	0.3	0.9	2.1	4.5
1999.....	46.4	*	0.2	*	*	0.2	0.4	1.0	2.0	4.6
All races 3/, male										
2006.....	50.2	*	*	*	*	*	0.4	1.0	2.3	5.1
2005.....	50.2	*	*	*	*	*	0.4	1.1	2.5	5.2
2004.....	49.8	*	*	*	*	0.2	0.5	1.1	2.4	5.0
2003.....	50.4	*	*	*	*	*	0.4	1.1	2.3	5.5
2002.....	50.7	*	0.3	*	*	*	0.5	1.2	2.3	5.3
2001.....	50.5	*	*	*	*	*	0.4	0.9	2.4	6.0
2000.....	50.8	*	*	*	*	0.2	0.4	1.1	2.5	5.5
1999.....	50.7	*	0.3	*	*	0.2	0.4	1.2	2.3	5.3
All races 3/, female										
2006.....	39.6	*	*	*	*	*	0.4	1.0	2.0	3.7
2005.....	39.5	*	*	*	*	*	0.3	0.8	1.8	3.2
2004.....	39.8	*	*	*	*	*	0.4	0.7	1.8	3.6
2003.....	40.7	*	*	*	*	*	0.4	0.9	1.9	3.8
2002.....	40.8	*	*	*	*	*	0.4	0.8	1.7	3.8
2001.....	41.7	*	*	*	*	*	0.3	0.8	1.8	4.0
2000.....	42.3	*	*	*	*	*	0.3	0.8	1.6	3.4
1999.....	42.3	*	*	*	*	*	0.4	0.8	1.7	3.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of digestive organs (C15-C26)										
White, both sexes										
2006 4/.....	46.4	*	*	*	*	0.1	0.4	0.9	2.0	4.1
2005 4/.....	46.3	*	*	*	*	*	0.3	0.8	2.0	3.8
2004 4/.....	46.3	*	*	*	*	0.1	0.4	0.8	1.9	4.0
2003 4/.....	47.0	*	*	*	*	0.1	0.4	0.9	1.8	4.3
2002.....	47.3	*	0.2	*	*	0.1	0.4	0.9	1.8	4.3
2001.....	47.6	*	*	*	*	0.1	0.3	0.8	1.9	4.5
2000.....	48.0	*	*	*	*	0.2	0.3	0.8	1.8	4.0
1999.....	48.0	*	0.2	*	*	0.1	0.4	0.8	1.7	4.1
White, male										
2006 4/.....	51.9	*	*	*	*	*	0.4	0.9	2.1	4.7
2005 4/.....	51.7	*	*	*	*	*	0.3	1.0	2.2	4.6
2004 4/.....	51.3	*	*	*	*	*	0.4	0.9	2.2	4.8
2003 4/.....	52.0	*	*	*	*	*	0.4	1.0	1.9	5.1
2002.....	52.3	*	*	*	*	*	0.4	1.0	2.0	4.8
2001.....	52.1	*	*	*	*	*	0.3	0.8	2.1	5.4
2000.....	52.3	*	*	*	*	*	0.3	0.9	2.2	5.0
1999.....	52.2	*	*	*	*	*	0.4	1.0	2.0	4.8
White, female										
2006 4/.....	41.1	*	*	*	*	*	0.4	1.0	1.9	3.4
2005 4/.....	40.9	*	*	*	*	*	0.3	0.7	1.7	3.0
2004 4/.....	41.3	*	*	*	*	*	0.4	0.7	1.6	3.2
2003 4/.....	42.2	*	*	*	*	*	0.3	0.8	1.7	3.4
2002.....	42.4	*	*	*	*	*	0.5	0.8	1.6	3.7
2001.....	43.2	*	*	*	*	*	0.4	0.8	1.6	3.5
2000.....	43.9	*	*	*	*	*	*	0.6	1.4	3.1
1999.....	43.9	*	*	*	*	*	0.4	0.6	1.5	3.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of digestive organs (C15-C26)										
Black, both sexes										
2006 4/.....	42.8	*	*	*	*	*	*	1.3	3.2	6.2
2005 4/.....	43.1	*	*	*	*	*	*	1.5	3.3	6.7
2004 4/.....	42.7	*	*	*	*	*	0.9	1.4	3.1	5.4
2003 4/.....	43.7	*	*	*	*	*	*	1.2	3.1	6.8
2002.....	43.6	*	*	*	*	*	*	1.4	2.7	5.9
2001.....	44.1	*	*	*	*	*	*	1.2	3.1	7.7
2000.....	44.5	*	*	*	*	*	*	1.3	3.3	6.9
1999.....	44.6	*	*	*	*	*	*	1.8	3.1	7.2
Black, male										
2006 4/.....	48.3	*	*	*	*	*	*	1.7	3.7	7.5
2005 4/.....	48.7	*	*	*	*	*	*	1.8	4.3	8.9
2004 4/.....	47.7	*	*	*	*	*	*	1.6	3.4	5.7
2003 4/.....	48.4	*	*	*	*	*	*	*	3.6	7.4
2002.....	48.5	*	*	*	*	*	*	1.9	3.5	7.6
2001.....	48.1	*	*	*	*	*	*	*	3.7	9.5
2000.....	49.2	*	*	*	*	*	*	*	4.1	8.5
1999.....	49.4	*	*	*	*	*	*	2.3	3.7	7.9
Black, female										
2006 4/.....	37.9	*	*	*	*	*	*	*	2.8	5.2
2005 4/.....	38.0	*	*	*	*	*	*	*	2.3	4.7
2004 4/.....	38.2	*	*	*	*	*	*	*	2.8	5.2
2003 4/.....	39.3	*	*	*	*	*	*	*	2.6	6.3
2002.....	39.2	*	*	*	*	*	*	*	2.0	4.4
2001.....	40.5	*	*	*	*	*	*	*	2.6	6.1
2000.....	40.3	*	*	*	*	*	*	1.4	2.5	5.5
1999.....	40.4	*	*	*	*	*	*	*	2.5	6.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of stomach (C16)										
All races 3/, both sexes										
2006.....	3.8	*	*	*	*	*	*	0.2	0.4	0.8
2005.....	3.9	*	*	*	*	*	*	0.1	0.4	0.7
2004.....	4.0	*	*	*	*	*	*	0.1	0.5	0.7
2003.....	4.2	*	*	*	*	*	*	0.2	0.4	0.8
2002.....	4.2	*	*	*	*	*	0.1	0.3	0.4	0.7
2001.....	4.3	*	*	*	*	*	*	0.2	0.4	0.8
2000.....	4.5	*	*	*	*	*	*	0.2	0.4	0.7
1999.....	4.6	*	*	*	*	*	*	0.2	0.4	0.8
All races 3/, male										
2006.....	4.5	*	*	*	*	*	*	*	0.4	0.7
2005.....	4.6	*	*	*	*	*	*	*	0.4	0.8
2004.....	4.8	*	*	*	*	*	*	*	0.4	0.7
2003.....	4.9	*	*	*	*	*	*	*	0.4	0.8
2002.....	5.0	*	*	*	*	*	*	0.3	0.5	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.5	0.9
2000.....	5.3	*	*	*	*	*	*	0.2	0.5	0.9
1999.....	5.5	*	*	*	*	*	*	0.2	0.5	0.9
All races 3/, female										
2006.....	3.1	*	*	*	*	*	*	0.2	0.4	0.9
2005.....	3.2	*	*	*	*	*	*	*	0.4	0.5
2004.....	3.3	*	*	*	*	*	*	*	0.5	0.6
2003.....	3.4	*	*	*	*	*	*	*	0.4	0.8
2002.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2001.....	3.5	*	*	*	*	*	*	*	0.4	0.7
2000.....	3.7	*	*	*	*	*	*	*	0.4	0.6
1999.....	3.7	*	*	*	*	*	*	0.3	0.3	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of colon, rectum and anus (C18-C21)										
All races 3/, both sexes										
2006.....	17.9	*	*	*	*	*	0.2	0.5	1.1	2.0
2005.....	18.0	*	*	*	*	*	0.2	0.4	1.1	1.9
2004.....	18.3	*	*	*	*	*	0.2	0.4	0.9	2.0
2003.....	19.2	*	*	*	*	*	0.2	0.5	0.9	2.1
2002.....	19.7	*	*	*	*	*	0.2	0.5	1.0	2.0
2001.....	20.0	*	*	*	*	*	0.2	0.4	1.1	2.3
2000.....	20.4	*	*	*	*	*	0.1	0.4	0.9	2.1
1999.....	20.5	*	*	*	*	*	0.2	0.4	1.0	1.9
All races 3/, male										
2006.....	18.3	*	*	*	*	*	*	0.5	1.1	2.1
2005.....	18.4	*	*	*	*	*	0.2	0.5	1.1	2.1
2004.....	18.6	*	*	*	*	*	0.2	0.5	1.0	2.1
2003.....	19.6	*	*	*	*	*	*	0.6	0.9	2.2
2002.....	20.1	*	*	*	*	*	0.3	0.5	1.1	2.1
2001.....	20.2	*	*	*	*	*	*	0.4	1.1	2.6
2000.....	20.6	*	*	*	*	*	*	0.4	1.1	2.4
1999.....	20.6	*	*	*	*	*	0.2	0.6	1.0	2.0
All races 3/, female										
2006.....	17.5	*	*	*	*	*	0.2	0.5	1.1	1.9
2005.....	17.6	*	*	*	*	*	*	0.4	1.0	1.7
2004.....	18.0	*	*	*	*	*	*	0.4	0.9	2.0
2003.....	18.9	*	*	*	*	*	*	0.4	0.9	2.0
2002.....	19.2	*	*	*	*	*	*	0.4	0.9	1.9
2001.....	19.8	*	*	*	*	*	*	0.5	1.0	1.9
2000.....	20.2	*	*	*	*	*	*	0.4	0.8	1.8
1999.....	20.3	*	*	*	*	*	*	0.3	0.9	1.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of colon, rectum and anus (C18-C21)										
White, both sexes										
2006 4/.....	18.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2005 4/.....	18.7	*	*	*	*	*	0.2	0.4	1.0	1.8
2004 4/.....	19.2	*	*	*	*	*	0.2	0.4	0.9	1.9
2003 4/.....	20.2	*	*	*	*	*	0.1	0.5	0.9	1.9
2002.....	20.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2001.....	21.0	*	*	*	*	*	0.2	0.4	1.0	2.1
2000.....	21.5	*	*	*	*	*	*	0.4	0.9	1.9
1999.....	21.6	*	*	*	*	*	0.2	0.4	0.9	1.8
White, male										
2006 4/.....	19.0	*	*	*	*	*	*	0.5	1.0	2.0
2005 4/.....	19.2	*	*	*	*	*	*	0.5	1.0	2.1
2004 4/.....	19.5	*	*	*	*	*	*	0.5	0.9	2.1
2003 4/.....	20.5	*	*	*	*	*	*	0.6	0.9	2.0
2002.....	21.1	*	*	*	*	*	0.2	0.5	1.0	2.0
2001.....	21.2	*	*	*	*	*	*	0.4	1.1	2.5
2000.....	21.6	*	*	*	*	*	*	0.4	1.0	2.2
1999.....	21.8	*	*	*	*	*	*	0.5	0.9	1.9
White, female										
2006 4/.....	18.4	*	*	*	*	*	*	0.5	0.9	1.8
2005 4/.....	18.3	*	*	*	*	*	*	0.3	1.1	1.5
2004 4/.....	18.9	*	*	*	*	*	*	0.4	0.8	1.7
2003 4/.....	19.8	*	*	*	*	*	*	0.4	0.9	1.8
2002.....	20.3	*	*	*	*	*	*	0.4	0.9	1.9
2001.....	20.7	*	*	*	*	*	*	0.4	0.9	1.8
2000.....	21.4	*	*	*	*	*	*	0.3	0.7	1.6
1999.....	21.4	*	*	*	*	*	*	0.3	0.9	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Colon (C18)										
All races 3/, both sexes										
2006.....	14.7	*	*	*	*	*	0.1	0.4	0.8	1.5
2005.....	14.8	*	*	*	*	*	0.1	0.4	0.8	1.3
2004.....	15.2	*	*	*	*	*	0.1	0.3	0.7	1.4
2003.....	16.1	*	*	*	*	*	0.1	0.4	0.7	1.7
2002.....	16.5	*	*	*	*	*	0.2	0.3	0.7	1.5
2001.....	16.8	*	*	*	*	*	0.1	0.3	0.8	1.7
2000.....	17.3	*	*	*	*	*	*	0.3	0.8	1.6
1999.....	17.4	*	*	*	*	*	0.2	0.3	0.7	1.5
All races 3/, male										
2006.....	14.8	*	*	*	*	*	*	0.4	0.8	1.6
2005.....	14.9	*	*	*	*	*	*	0.4	0.8	1.5
2004.....	15.1	*	*	*	*	*	*	0.4	0.6	1.4
2003.....	16.1	*	*	*	*	*	*	0.5	0.7	1.7
2002.....	16.6	*	*	*	*	*	0.2	0.3	0.8	1.5
2001.....	16.7	*	*	*	*	*	*	0.3	0.9	1.9
2000.....	17.1	*	*	*	*	*	*	0.3	0.9	1.7
1999.....	17.2	*	*	*	*	*	*	0.4	0.8	1.5
All races 3/, female										
2006.....	14.7	*	*	*	*	*	*	0.4	0.8	1.3
2005.....	14.8	*	*	*	*	*	*	0.3	0.8	1.2
2004.....	15.2	*	*	*	*	*	*	0.3	0.7	1.4
2003.....	16.1	*	*	*	*	*	*	0.3	0.7	1.7
2002.....	16.4	*	*	*	*	*	*	0.3	0.7	1.4
2001.....	16.9	*	*	*	*	*	*	0.3	0.8	1.5
2000.....	17.4	*	*	*	*	*	*	0.3	0.6	1.4
1999.....	17.5	*	*	*	*	*	*	0.3	0.7	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Colon (C18)										
White, both sexes										
2006 4/.....	15.3	*	*	*	*	*	0.1	0.4	0.7	1.3
2005 4/.....	15.4	*	*	*	*	*	0.1	0.3	0.8	1.3
2004 4/.....	15.9	*	*	*	*	*	*	0.3	0.6	1.3
2003 4/.....	16.9	*	*	*	*	*	*	0.4	0.6	1.6
2002.....	17.3	*	*	*	*	*	0.2	0.3	0.7	1.4
2001.....	17.6	*	*	*	*	*	0.1	0.3	0.8	1.6
2000.....	18.1	*	*	*	*	*	*	0.3	0.7	1.4
1999.....	18.2	*	*	*	*	*	*	0.3	0.7	1.4
White, male										
2006 4/.....	15.3	*	*	*	*	*	*	0.4	0.8	1.5
2005 4/.....	15.5	*	*	*	*	*	*	0.4	0.7	1.5
2004 4/.....	15.8	*	*	*	*	*	*	0.3	0.6	1.4
2003 4/.....	16.9	*	*	*	*	*	*	0.5	0.6	1.5
2002.....	17.3	*	*	*	*	*	*	0.4	0.8	1.4
2001.....	17.5	*	*	*	*	*	*	*	0.9	1.8
2000.....	17.9	*	*	*	*	*	*	0.3	0.9	1.7
1999.....	18.2	*	*	*	*	*	*	0.3	0.7	1.5
White, female										
2006 4/.....	15.4	*	*	*	*	*	*	0.4	0.7	1.2
2005 4/.....	15.3	*	*	*	*	*	*	0.3	0.9	1.1
2004 4/.....	16.0	*	*	*	*	*	*	0.3	0.6	1.2
2003 4/.....	16.8	*	*	*	*	*	*	0.3	0.7	1.6
2002.....	17.3	*	*	*	*	*	*	0.3	0.7	1.4
2001.....	17.7	*	*	*	*	*	*	0.3	0.7	1.4
2000.....	18.3	*	*	*	*	*	*	0.3	0.6	1.2
1999.....	18.3	*	*	*	*	*	*	*	0.7	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of pancreas (C25)										
All races 3/, both sexes										
2006.....	11.2	*	*	*	*	*	*	*	0.2	0.6
2005.....	11.1	*	*	*	*	*	*	0.1	0.2	0.6
2004.....	10.8	*	*	*	*	*	*	*	0.2	0.6
2003.....	10.6	*	*	*	*	*	*	0.1	0.2	0.7
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.7
2001.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2000.....	10.4	*	*	*	*	*	*	*	0.2	0.6
1999.....	10.4	*	*	*	*	*	*	*	0.2	0.8
All races 3/, male										
2006.....	11.2	*	*	*	*	*	*	*	*	0.8
2005.....	11.1	*	*	*	*	*	*	*	0.2	0.8
2004.....	10.9	*	*	*	*	*	*	*	0.3	0.7
2003.....	10.5	*	*	*	*	*	*	*	0.2	0.9
2002.....	10.5	*	*	*	*	*	*	*	0.2	0.8
2001.....	10.3	*	*	*	*	*	*	*	0.2	0.9
2000.....	10.3	*	*	*	*	*	*	*	0.2	0.8
1999.....	10.4	*	*	*	*	*	*	*	0.3	0.9
All races 3/, female										
2006.....	11.1	*	*	*	*	*	*	*	*	0.3
2005.....	11.0	*	*	*	*	*	*	*	*	0.5
2004.....	10.7	*	*	*	*	*	*	*	*	0.5
2003.....	10.6	*	*	*	*	*	*	*	*	0.5
2002.....	10.5	*	*	*	*	*	*	*	*	0.6
2001.....	10.6	*	*	*	*	*	*	*	*	0.6
2000.....	10.5	*	*	*	*	*	*	*	*	0.5
1999.....	10.5	*	*	*	*	*	*	*	0.2	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)										
All races 3/, both sexes										
2006.....	54.6	*	*	*	*	*	0.1	0.2	0.6	2.1
2005.....	55.3	*	*	*	*	*	0.1	0.2	0.5	2.3
2004.....	55.4	*	*	*	*	*	0.1	0.3	0.6	2.4
2003.....	56.0	*	*	*	*	*	0.1	0.2	0.7	2.6
2002.....	56.3	*	*	*	*	*	0.2	0.2	0.7	2.9
2001.....	56.5	*	*	*	*	*	0.1	0.2	0.7	3.1
2000.....	57.0	*	*	*	*	*	0.1	0.2	0.9	3.1
1999.....	56.2	*	*	*	*	*	0.1	0.3	0.8	3.1
All races 3/, male										
2006.....	62.9	*	*	*	*	*	*	0.3	0.7	2.2
2005.....	64.2	*	*	*	*	*	*	0.3	0.6	2.4
2004.....	64.4	*	*	*	*	*	*	0.4	0.7	2.4
2003.....	65.4	*	*	*	*	*	*	0.2	0.7	2.6
2002.....	66.1	*	*	*	*	*	0.3	0.3	0.8	3.0
2001.....	67.2	*	*	*	*	*	*	0.2	0.7	3.1
2000.....	68.1	*	*	*	*	*	*	0.4	0.8	3.5
1999.....	68.0	*	*	*	*	*	*	0.3	0.8	3.2
All races 3/, female										
2006.....	46.5	*	*	*	*	*	*	0.2	0.6	2.0
2005.....	46.8	*	*	*	*	*	*	0.2	0.5	2.1
2004.....	46.7	*	*	*	*	*	*	*	0.5	2.4
2003.....	46.9	*	*	*	*	*	*	0.3	0.6	2.6
2002.....	46.8	*	*	*	*	*	*	*	0.7	2.8
2001.....	46.1	*	*	*	*	*	*	0.3	0.7	3.1
2000.....	46.2	*	*	*	*	*	*	*	0.9	2.7
1999.....	45.0	*	*	*	*	*	*	0.2	0.7	3.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)										
White, both sexes										
2006 4/.....	59.0	*	*	*	*	*	*	0.2	0.6	2.1
2005 4/.....	59.7	*	*	*	*	*	*	0.2	0.5	2.3
2004 4/.....	59.7	*	*	*	*	*	*	0.3	0.6	2.4
2003 4/.....	60.3	*	*	*	*	*	0.1	0.2	0.7	2.5
2002.....	60.7	*	*	*	*	*	0.1	0.2	0.7	2.8
2001.....	60.8	*	*	*	*	*	0.1	0.2	0.7	3.1
2000.....	61.2	*	*	*	*	*	0.1	0.2	0.9	2.9
1999.....	60.2	*	*	*	*	*	*	0.3	0.8	2.9
White, male										
2006 4/.....	67.0	*	*	*	*	*	*	*	0.7	2.1
2005 4/.....	68.3	*	*	*	*	*	*	0.2	0.6	2.3
2004 4/.....	68.3	*	*	*	*	*	*	0.4	0.6	2.3
2003 4/.....	69.4	*	*	*	*	*	*	*	0.7	2.5
2002.....	70.1	*	*	*	*	*	*	0.3	0.8	2.8
2001.....	71.1	*	*	*	*	*	*	*	0.7	3.0
2000.....	72.0	*	*	*	*	*	*	0.4	0.8	3.2
1999.....	71.6	*	*	*	*	*	*	0.3	0.9	2.9
White, female										
2006 4/.....	51.1	*	*	*	*	*	*	*	0.5	2.0
2005 4/.....	51.2	*	*	*	*	*	*	*	0.4	2.2
2004 4/.....	51.2	*	*	*	*	*	*	*	0.5	2.5
2003 4/.....	51.5	*	*	*	*	*	*	*	0.6	2.5
2002.....	51.5	*	*	*	*	*	*	*	0.7	2.9
2001.....	50.7	*	*	*	*	*	*	*	0.7	3.1
2000.....	50.7	*	*	*	*	*	*	*	0.9	2.5
1999.....	49.1	*	*	*	*	*	*	*	0.7	2.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of trachea, bronchus and lung (C33-C34)										
All races 3/, both sexes										
2006.....	53.0	*	*	*	*	*	*	0.2	0.6	2.0
2005.....	53.7	*	*	*	*	*	*	0.2	0.5	2.1
2004.....	53.8	*	*	*	*	*	*	0.2	0.5	2.2
2003.....	54.4	*	*	*	*	*	*	0.2	0.6	2.4
2002.....	54.7	*	*	*	*	*	*	0.2	0.6	2.7
2001.....	54.8	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	55.3	*	*	*	*	*	*	0.2	0.8	2.9
1999.....	54.5	*	*	*	*	*	*	0.2	0.7	2.9
All races 3/, male										
2006.....	60.5	*	*	*	*	*	*	0.2	0.6	2.0
2005.....	61.8	*	*	*	*	*	*	*	0.6	2.2
2004.....	62.0	*	*	*	*	*	*	0.3	0.6	2.2
2003.....	62.9	*	*	*	*	*	*	*	0.7	2.4
2002.....	63.7	*	*	*	*	*	*	*	0.7	2.6
2001.....	64.7	*	*	*	*	*	*	*	0.6	2.8
2000.....	65.5	*	*	*	*	*	*	0.3	0.7	3.3
1999.....	65.4	*	*	*	*	*	*	0.2	0.7	2.8
All races 3/, female										
2006.....	45.7	*	*	*	*	*	*	*	0.5	2.0
2005.....	45.9	*	*	*	*	*	*	0.2	0.4	2.0
2004.....	45.9	*	*	*	*	*	*	*	0.5	2.2
2003.....	46.1	*	*	*	*	*	*	0.2	0.5	2.5
2002.....	46.0	*	*	*	*	*	*	*	0.6	2.7
2001.....	45.3	*	*	*	*	*	*	*	0.6	2.9
2000.....	45.4	*	*	*	*	*	*	*	0.8	2.6
1999.....	44.1	*	*	*	*	*	*	*	0.6	3.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of trachea, bronchus and lung (C33-C34)										
White, both sexes										
2006 4/.....	57.3	*	*	*	*	*	*	0.2	0.5	2.0
2005 4/.....	58.1	*	*	*	*	*	*	0.2	0.4	2.1
2004 4/.....	58.1	*	*	*	*	*	*	0.2	0.5	2.2
2003 4/.....	58.7	*	*	*	*	*	*	0.2	0.6	2.4
2002.....	59.1	*	*	*	*	*	*	0.2	0.6	2.6
2001.....	59.1	*	*	*	*	*	*	0.2	0.6	2.9
2000.....	59.5	*	*	*	*	*	*	0.2	0.8	2.7
1999.....	58.5	*	*	*	*	*	*	0.2	0.7	2.7
White, male										
2006 4/.....	64.6	*	*	*	*	*	*	*	0.6	2.0
2005 4/.....	65.9	*	*	*	*	*	*	*	0.6	2.2
2004 4/.....	65.9	*	*	*	*	*	*	0.3	0.5	2.1
2003 4/.....	66.9	*	*	*	*	*	*	*	0.6	2.3
2002.....	67.7	*	*	*	*	*	*	*	0.7	2.5
2001.....	68.6	*	*	*	*	*	*	*	0.6	2.8
2000.....	69.4	*	*	*	*	*	*	0.3	0.7	3.0
1999.....	69.1	*	*	*	*	*	*	*	0.8	2.6
White, female										
2006 4/.....	50.2	*	*	*	*	*	*	*	0.5	2.0
2005 4/.....	50.4	*	*	*	*	*	*	*	0.3	2.0
2004 4/.....	50.4	*	*	*	*	*	*	*	0.4	2.3
2003 4/.....	50.7	*	*	*	*	*	*	*	0.6	2.4
2002.....	50.7	*	*	*	*	*	*	*	0.5	2.8
2001.....	49.8	*	*	*	*	*	*	*	0.6	3.0
2000.....	49.9	*	*	*	*	*	*	*	0.8	2.4
1999.....	48.2	*	*	*	*	*	*	*	0.6	2.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of bone and articular cartilage (C40-C41)										
All races 3/, both sexes										
2006.....	0.4	*	*	*	0.2	0.5	0.4	0.3	0.2	0.2
2005.....	0.5	*	*	*	0.3	0.6	0.5	0.3	0.2	0.2
2004.....	0.4	*	*	*	0.3	0.5	0.4	0.3	0.2	0.2
2003.....	0.4	*	*	*	0.3	0.5	0.4	0.3	0.2	0.2
2002.....	0.4	*	*	0.1	0.3	0.5	0.4	0.3	0.2	0.2
2001.....	0.5	*	*	*	0.3	0.5	0.5	0.3	0.2	0.2
2000.....	0.4	*	*	*	0.3	0.5	0.4	0.2	0.2	0.2
1999.....	0.4	*	*	0.1	0.3	0.6	0.4	0.2	0.2	0.2
All races 3/, male										
2006.....	0.5	*	*	*	*	0.7	0.6	0.4	0.3	0.2
2005.....	0.5	*	*	*	0.3	0.7	0.6	0.4	0.3	0.2
2004.....	0.5	*	*	*	0.3	0.7	0.5	0.3	0.2	0.2
2003.....	0.5	*	*	*	0.3	0.7	0.5	0.3	0.2	0.3
2002.....	0.5	*	*	*	0.3	0.5	0.5	0.4	0.2	0.2
2001.....	0.5	*	*	*	0.4	0.6	0.7	0.3	0.2	0.2
2000.....	0.5	*	*	*	0.4	0.6	0.5	0.3	0.3	0.2
1999.....	0.5	*	*	*	0.3	0.7	0.5	0.3	0.2	0.2
All races 3/, female										
2006.....	0.4	*	*	*	*	0.4	0.3	0.2	*	*
2005.....	0.4	*	*	*	0.3	0.5	0.4	*	*	0.2
2004.....	0.4	*	*	*	0.3	0.4	0.3	0.2	*	*
2003.....	0.4	*	*	*	0.3	0.3	0.3	0.3	*	*
2002.....	0.4	*	*	*	0.3	0.4	0.3	*	*	*
2001.....	0.4	*	*	*	0.3	0.4	0.3	0.2	0.2	0.2
2000.....	0.4	*	*	*	0.2	0.4	0.3	*	*	*
1999.....	0.4	*	*	*	0.3	0.5	0.3	*	0.2	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of bone and articular cartilage (C40-C41)

White, both sexes

2006 4/.....	0.5	*	*	*	0.2	0.5	0.4	0.3	0.2	0.2
2005 4/.....	0.5	*	*	*	0.3	0.6	0.5	0.3	0.2	0.2
2004 4/.....	0.5	*	*	*	0.3	0.5	0.4	0.3	0.2	0.2
2003 4/.....	0.5	*	*	*	0.3	0.6	0.4	0.3	0.2	0.2
2002.....	0.4	*	*	*	0.3	0.5	0.4	0.3	0.2	0.2
2001.....	0.5	*	*	*	0.4	0.6	0.5	0.3	0.2	0.2
2000.....	0.5	*	*	*	0.3	0.5	0.4	0.2	0.2	0.2
1999.....	0.5	*	*	*	0.3	0.7	0.4	0.2	0.2	0.2

White, male

2006 4/.....	0.5	*	*	*	*	0.7	0.6	0.4	0.3	0.2
2005 4/.....	0.6	*	*	*	0.3	0.7	0.7	0.4	0.3	*
2004 4/.....	0.5	*	*	*	0.3	0.7	0.5	0.4	*	*
2003 4/.....	0.5	*	*	*	0.3	0.8	0.4	0.3	0.3	0.4
2002.....	0.5	*	*	*	0.3	0.6	0.5	0.5	*	0.3
2001.....	0.5	*	*	*	0.4	0.7	0.7	0.3	*	0.2
2000.....	0.5	*	*	*	0.4	0.5	0.6	0.3	0.3	*
1999.....	0.5	*	*	*	0.3	0.7	0.5	0.3	0.3	0.2

White, female

2006 4/.....	0.4	*	*	*	*	0.4	0.3	*	*	*
2005 4/.....	0.4	*	*	*	0.3	0.6	0.4	*	*	*
2004 4/.....	0.4	*	*	*	0.3	0.4	0.3	*	*	*
2003 4/.....	0.4	*	*	*	0.3	0.3	0.4	*	*	*
2002.....	0.4	*	*	*	0.4	0.5	0.3	*	*	*
2001.....	0.4	*	*	*	0.3	0.4	0.3	0.3	*	*
2000.....	0.4	*	*	*	0.3	0.4	0.3	*	*	*
1999.....	0.4	*	*	*	*	0.6	0.3	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Melanoma and other malignant neoplasms of skin (C43-C44)										
All races 3/, both sexes										
2006.....	3.7	*	*	*	*	*	0.2	0.3	0.7	1.0
2005.....	3.6	*	*	*	*	*	0.2	0.5	0.6	0.9
2004.....	3.5	*	*	*	*	*	0.1	0.4	0.6	1.0
2003.....	3.5	*	*	*	*	*	0.1	0.4	0.7	0.9
2002.....	3.4	*	*	*	*	*	0.2	0.4	0.6	1.1
2001.....	3.5	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	3.4	*	*	*	*	*	0.2	0.3	0.6	1.1
1999.....	3.4	*	*	*	*	*	0.2	0.3	0.7	1.2
All races 3/, male										
2006.....	5.0	*	*	*	*	*	0.2	0.4	0.8	1.2
2005.....	4.8	*	*	*	*	*	0.2	0.6	0.8	1.0
2004.....	4.7	*	*	*	*	*	*	0.4	0.7	1.1
2003.....	4.6	*	*	*	*	*	*	0.5	0.8	1.2
2002.....	4.5	*	*	*	*	*	*	0.4	0.8	1.2
2001.....	4.6	*	*	*	*	*	*	0.5	0.6	1.3
2000.....	4.5	*	*	*	*	*	*	0.4	0.7	1.1
1999.....	4.5	*	*	*	*	*	*	0.4	0.8	1.5
All races 3/, female										
2006.....	2.4	*	*	*	*	*	*	*	0.6	0.7
2005.....	2.5	*	*	*	*	*	*	0.3	0.5	0.9
2004.....	2.4	*	*	*	*	*	*	0.3	0.5	0.8
2003.....	2.4	*	*	*	*	*	*	0.3	0.5	0.7
2002.....	2.4	*	*	*	*	*	*	0.4	0.4	0.9
2001.....	2.5	*	*	*	*	*	*	0.3	0.5	0.8
2000.....	2.5	*	*	*	*	*	*	0.3	0.5	1.1
1999.....	2.4	*	*	*	*	*	*	0.2	0.6	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Melanoma and other malignant neoplasms of skin (C43-C44)										
White, both sexes										
2006 4/.....	4.4	*	*	*	*	*	0.2	0.3	0.9	1.2
2005 4/.....	4.3	*	*	*	*	*	0.2	0.6	0.8	1.1
2004 4/.....	4.2	*	*	*	*	*	0.2	0.4	0.8	1.2
2003 4/.....	4.2	*	*	*	*	*	0.2	0.4	0.8	1.1
2002.....	4.1	*	*	*	*	*	0.2	0.5	0.8	1.3
2001.....	4.2	*	*	*	*	*	0.2	0.5	0.7	1.2
2000.....	4.0	*	*	*	*	*	0.2	0.4	0.8	1.3
1999.....	4.0	*	*	*	*	*	0.2	0.4	0.8	1.4
White, male										
2006 4/.....	5.9	*	*	*	*	*	0.2	0.5	0.9	1.5
2005 4/.....	5.7	*	*	*	*	*	*	0.7	1.0	1.1
2004 4/.....	5.6	*	*	*	*	*	*	0.5	0.9	1.4
2003 4/.....	5.5	*	*	*	*	*	*	0.6	1.0	1.4
2002.....	5.3	*	*	*	*	*	*	0.5	1.0	1.4
2001.....	5.4	*	*	*	*	*	*	0.5	0.8	1.5
2000.....	5.2	*	*	*	*	*	*	0.4	0.9	1.3
1999.....	5.3	*	*	*	*	*	*	0.5	0.9	1.7
White, female										
2006 4/.....	2.9	*	*	*	*	*	*	*	0.8	0.8
2005 4/.....	3.0	*	*	*	*	*	*	0.4	0.6	1.1
2004 4/.....	2.8	*	*	*	*	*	*	0.4	0.6	1.0
2003 4/.....	2.9	*	*	*	*	*	*	0.3	0.6	0.9
2002.....	2.9	*	*	*	*	*	*	0.5	0.5	1.1
2001.....	2.9	*	*	*	*	*	*	0.4	0.6	1.0
2000.....	2.9	*	*	*	*	*	*	0.3	0.6	1.3
1999.....	2.9	*	*	*	*	*	*	0.3	0.7	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant melanoma of skin (C43)										
All races 3/, both sexes										
2006.....	2.8	*	*	*	*	*	0.2	0.3	0.7	0.9
2005.....	2.8	*	*	*	*	*	0.2	0.5	0.6	0.9
2004.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9
2003.....	2.7	*	*	*	*	*	0.1	0.3	0.6	0.9
2002.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2001.....	2.6	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
1999.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.0
All races 3/, male										
2006.....	3.7	*	*	*	*	*	0.2	0.3	0.7	1.2
2005.....	3.6	*	*	*	*	*	*	0.6	0.8	0.9
2004.....	3.5	*	*	*	*	*	*	0.4	0.7	1.0
2003.....	3.5	*	*	*	*	*	*	0.4	0.8	1.1
2002.....	3.3	*	*	*	*	*	*	0.4	0.8	1.1
2001.....	3.4	*	*	*	*	*	*	0.4	0.6	1.2
2000.....	3.3	*	*	*	*	*	*	0.3	0.7	1.0
1999.....	3.3	*	*	*	*	*	*	0.4	0.7	1.3
All races 3/, female										
2006.....	2.0	*	*	*	*	*	*	*	0.6	0.6
2005.....	2.0	*	*	*	*	*	*	0.3	0.5	0.8
2004.....	1.9	*	*	*	*	*	*	0.3	0.5	0.8
2003.....	1.9	*	*	*	*	*	*	0.2	0.5	0.7
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.9
2001.....	1.9	*	*	*	*	*	*	0.3	0.5	0.7
2000.....	2.0	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	1.9	*	*	*	*	*	*	0.2	0.6	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant melanoma of skin (C43)										
White, both sexes										
2006 4/.....	3.4	*	*	*	*	*	0.2	0.3	0.8	1.1
2005 4/.....	3.4	*	*	*	*	*	0.2	0.6	0.8	1.0
2004 4/.....	3.3	*	*	*	*	*	0.2	0.4	0.8	1.1
2003 4/.....	3.2	*	*	*	*	*	0.2	0.4	0.8	1.1
2002.....	3.1	*	*	*	*	*	0.2	0.5	0.7	1.2
2001.....	3.2	*	*	*	*	*	0.2	0.5	0.7	1.2
2000.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.2
1999.....	3.1	*	*	*	*	*	0.2	0.4	0.8	1.3
White, male										
2006 4/.....	4.5	*	*	*	*	*	0.2	0.4	0.9	1.4
2005 4/.....	4.4	*	*	*	*	*	*	0.7	0.9	1.1
2004 4/.....	4.3	*	*	*	*	*	*	0.5	0.9	1.3
2003 4/.....	4.2	*	*	*	*	*	*	0.5	0.9	1.3
2002.....	4.0	*	*	*	*	*	*	0.5	1.0	1.3
2001.....	4.1	*	*	*	*	*	*	0.5	0.7	1.4
2000.....	4.0	*	*	*	*	*	*	0.4	0.9	1.2
1999.....	4.0	*	*	*	*	*	*	0.5	0.8	1.6
White, female										
2006 4/.....	2.3	*	*	*	*	*	*	*	0.8	0.8
2005 4/.....	2.4	*	*	*	*	*	*	0.4	0.6	1.0
2004 4/.....	2.3	*	*	*	*	*	*	0.4	0.6	1.0
2003 4/.....	2.3	*	*	*	*	*	*	0.3	0.6	0.9
2002.....	2.3	*	*	*	*	*	*	0.4	0.5	1.1
2001.....	2.3	*	*	*	*	*	*	0.4	0.6	0.9
2000.....	2.3	*	*	*	*	*	*	0.3	0.6	1.2
1999.....	2.3	*	*	*	*	*	*	0.3	0.7	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of mesothelial and soft tissue (C45-C49)										
All races 3/, both sexes										
2006.....	2.4	*	0.1	0.2	0.1	0.3	0.3	0.4	0.5	0.5
2005.....	2.4	*	0.1	0.1	0.2	0.4	0.4	0.4	0.5	0.5
2004.....	2.4	*	0.2	0.1	0.2	0.3	0.4	0.4	0.5	0.7
2003.....	2.4	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.6
2002.....	2.4	*	0.1	0.1	0.2	0.3	0.4	0.4	0.4	0.7
2001.....	2.4	*	*	0.1	0.2	0.3	0.3	0.4	0.6	0.6
2000.....	2.4	*	0.1	*	0.1	0.4	0.4	0.5	0.6	0.7
1999.....	2.4	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.7
All races 3/, male										
2006.....	2.8	*	*	0.2	*	0.3	0.3	0.5	0.6	0.6
2005.....	2.8	*	*	*	0.2	0.3	0.5	0.5	0.6	0.6
2004.....	2.8	*	*	*	0.2	0.3	0.4	0.5	0.6	0.8
2003.....	2.8	*	*	*	*	0.4	0.4	0.5	0.6	0.7
2002.....	2.8	*	*	*	0.2	0.3	0.4	0.5	0.5	0.8
2001.....	2.8	*	*	*	*	0.4	0.4	0.5	0.7	0.6
2000.....	2.9	*	*	*	*	0.4	0.5	0.5	0.6	0.8
1999.....	2.8	*	*	*	0.2	0.4	0.4	0.5	0.7	0.9
All races 3/, female										
2006.....	2.1	*	*	*	*	0.3	0.3	0.4	0.4	0.4
2005.....	2.1	*	*	*	*	0.4	0.3	0.4	0.5	0.4
2004.....	2.0	*	*	*	0.2	0.3	0.3	0.2	0.4	0.5
2003.....	2.0	*	*	*	0.2	0.3	0.3	0.4	0.4	0.6
2002.....	1.9	*	*	*	0.2	0.3	0.3	0.4	0.3	0.5
2001.....	2.0	*	*	*	*	0.3	0.2	0.4	0.4	0.6
2000.....	2.0	*	*	*	*	0.4	0.3	0.4	0.6	0.6
1999.....	1.9	*	*	*	*	*	0.4	0.4	0.4	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of mesothelial and soft tissue (C45-C49)										
White, both sexes										
2006 4/.....	2.7	*	*	0.2	0.1	0.3	0.3	0.4	0.5	0.5
2005 4/.....	2.7	*	*	*	0.2	0.4	0.4	0.4	0.5	0.5
2004 4/.....	2.7	*	*	0.2	0.2	0.3	0.3	0.4	0.5	0.6
2003 4/.....	2.6	*	*	0.1	0.2	0.4	0.4	0.4	0.5	0.7
2002.....	2.6	*	*	0.1	0.2	0.3	0.4	0.4	0.4	0.7
2001.....	2.6	*	*	*	0.2	0.3	0.3	0.4	0.5	0.6
2000.....	2.7	*	*	*	0.1	0.4	0.3	0.5	0.6	0.6
1999.....	2.6	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.7
White, male										
2006 4/.....	3.1	*	*	*	*	0.4	0.4	0.4	0.6	0.6
2005 4/.....	3.1	*	*	*	*	0.3	0.6	0.5	0.6	0.6
2004 4/.....	3.2	*	*	*	*	0.3	0.4	0.5	0.6	0.8
2003 4/.....	3.1	*	*	*	*	0.4	0.5	0.5	0.6	0.7
2002.....	3.1	*	*	*	*	0.3	0.4	0.5	0.5	0.8
2001.....	3.1	*	*	*	*	0.3	0.4	0.4	0.7	0.7
2000.....	3.2	*	*	*	*	0.4	0.4	0.6	0.6	0.7
1999.....	3.1	*	*	*	*	0.4	0.3	0.5	0.6	0.8
White, female										
2006 4/.....	2.2	*	*	*	*	0.3	*	0.3	0.4	0.5
2005 4/.....	2.2	*	*	*	*	0.5	0.3	0.3	0.4	0.4
2004 4/.....	2.2	*	*	*	*	0.3	0.3	*	0.4	0.5
2003 4/.....	2.2	*	*	*	*	0.3	0.4	0.3	0.4	0.6
2002.....	2.1	*	*	*	*	0.3	0.3	0.3	0.4	0.5
2001.....	2.1	*	*	*	*	0.3	0.3	0.4	0.4	0.6
2000.....	2.2	*	*	*	*	0.3	0.3	0.4	0.6	0.5
1999.....	2.1	*	*	*	*	*	0.4	0.4	0.4	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of other mesothelial and soft tissue (C47-C49)										
All races 3/, both sexes										
2006.....	1.6	*	0.1	0.2	0.1	0.3	0.3	0.4	0.5	0.4
2005.....	1.6	*	0.1	0.1	0.2	0.4	0.4	0.4	0.5	0.5
2004.....	1.6	*	0.2	0.1	0.2	0.3	0.4	0.3	0.5	0.6
2003.....	1.5	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.6
2002.....	1.5	*	0.1	0.1	0.2	0.3	0.4	0.4	0.4	0.6
2001.....	1.5	*	*	0.1	0.2	0.3	0.3	0.4	0.5	0.5
2000.....	1.6	*	0.1	*	0.1	0.4	0.4	0.4	0.5	0.6
1999.....	1.5	*	*	0.1	0.2	0.3	0.4	0.4	0.5	0.7
All races 3/, male										
2006.....	1.4	*	*	0.2	*	0.3	0.3	0.4	0.6	0.5
2005.....	1.4	*	*	*	0.2	0.3	0.5	0.5	0.6	0.5
2004.....	1.4	*	*	*	0.2	0.3	0.4	0.5	0.6	0.7
2003.....	1.4	*	*	*	*	0.4	0.4	0.4	0.6	0.6
2002.....	1.4	*	*	*	0.2	0.3	0.4	0.5	0.5	0.7
2001.....	1.4	*	*	*	*	0.4	0.4	0.4	0.6	0.5
2000.....	1.4	*	*	*	*	0.4	0.5	0.5	0.5	0.7
1999.....	1.5	*	*	*	0.2	0.4	0.4	0.5	0.6	0.7
All races 3/, female										
2006.....	1.7	*	*	*	*	0.3	0.3	0.4	0.4	0.4
2005.....	1.7	*	*	*	*	0.4	0.3	0.4	0.4	0.4
2004.....	1.7	*	*	*	0.2	0.3	0.3	0.2	0.4	0.4
2003.....	1.6	*	*	*	0.2	0.3	0.3	0.3	0.4	0.5
2002.....	1.6	*	*	*	0.2	0.3	0.3	0.4	0.3	0.5
2001.....	1.6	*	*	*	*	0.3	0.2	0.4	0.4	0.5
2000.....	1.7	*	*	*	*	0.4	0.3	0.4	0.5	0.5
1999.....	1.6	*	*	*	*	*	0.4	0.4	0.4	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of other mesothelial and soft tissue (C47-C49)										
White, both sexes										
2006 4/.....	1.7	*	*	0.2	0.1	0.3	0.3	0.4	0.5	0.5
2005 4/.....	1.7	*	*	*	0.2	0.4	0.4	0.4	0.5	0.4
2004 4/.....	1.7	*	*	0.2	0.2	0.3	0.3	0.4	0.5	0.6
2003 4/.....	1.6	*	*	0.1	0.2	0.4	0.4	0.4	0.4	0.6
2002.....	1.6	*	*	0.1	0.2	0.3	0.4	0.4	0.4	0.6
2001.....	1.6	*	*	*	0.2	0.3	0.3	0.4	0.5	0.6
2000.....	1.7	*	*	*	0.1	0.3	0.3	0.5	0.6	0.6
1999.....	1.6	*	*	0.1	0.2	0.3	0.3	0.4	0.4	0.6
White, male										
2006 4/.....	1.6	*	*	*	*	0.3	0.4	0.4	0.6	0.5
2005 4/.....	1.5	*	*	*	*	0.3	0.6	0.4	0.5	0.5
2004 4/.....	1.5	*	*	*	*	0.3	0.4	0.5	0.5	0.7
2003 4/.....	1.4	*	*	*	*	0.4	0.5	0.4	0.5	0.7
2002.....	1.4	*	*	*	*	0.3	0.4	0.5	0.4	0.7
2001.....	1.5	*	*	*	*	0.3	0.4	0.4	0.6	0.6
2000.....	1.5	*	*	*	*	0.4	0.4	0.5	0.6	0.6
1999.....	1.5	*	*	*	*	0.4	0.3	0.5	0.5	0.7
White, female										
2006 4/.....	1.9	*	*	*	*	0.3	*	0.3	0.4	0.4
2005 4/.....	1.8	*	*	*	*	0.5	0.3	0.3	0.4	0.4
2004 4/.....	1.8	*	*	*	*	0.3	0.3	*	0.4	0.4
2003 4/.....	1.8	*	*	*	*	0.3	0.4	0.3	0.3	0.5
2002.....	1.8	*	*	*	*	0.3	0.3	0.3	0.3	0.5
2001.....	1.7	*	*	*	*	0.3	0.3	0.4	0.4	0.6
2000.....	1.8	*	*	*	*	0.3	*	0.4	0.5	0.5
1999.....	1.7	*	*	*	*	*	0.4	0.3	0.4	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of breast (C50)										
All races 3/, both sexes										
2006.....	13.8	*	*	*	*	*	*	0.4	1.5	3.8
2005.....	14.0	*	*	*	*	*	*	0.4	1.4	3.8
2004.....	14.1	*	*	*	*	*	*	0.4	1.6	3.9
2003.....	14.4	*	*	*	*	*	*	0.3	1.7	4.3
2002.....	14.5	*	*	*	*	*	*	0.3	1.7	4.1
2001.....	14.7	*	*	*	*	*	*	0.5	1.8	4.3
2000.....	15.0	*	*	*	*	*	*	0.4	1.9	4.3
1999.....	14.9	*	*	*	*	*	*	0.4	1.7	4.3
All races 3/, male										
2006.....	0.3	*	*	*	*	*	*	*	*	*
2005.....	0.3	*	*	*	*	*	*	*	*	*
2004.....	0.3	*	*	*	*	*	*	*	*	*
2003.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*
All races 3/, female										
2006.....	26.9	*	*	*	*	*	*	0.7	3.0	7.5
2005.....	27.3	*	*	*	*	*	*	0.8	2.8	7.7
2004.....	27.5	*	*	*	*	*	*	0.8	3.2	7.8
2003.....	28.2	*	*	*	*	*	*	0.6	3.4	8.5
2002.....	28.3	*	*	*	*	*	*	0.7	3.5	8.2
2001.....	28.6	*	*	*	*	*	*	0.9	3.6	8.6
2000.....	29.2	*	*	*	*	*	*	0.8	3.8	8.5
1999.....	28.9	*	*	*	*	*	*	0.9	3.4	8.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of breast (C50)										
White, both sexes										
2006 4/.....	14.2	*	*	*	*	*	*	0.3	1.3	3.3
2005 4/.....	14.4	*	*	*	*	*	*	0.3	1.2	3.3
2004 4/.....	14.6	*	*	*	*	*	*	0.3	1.3	3.4
2003 4/.....	15.0	*	*	*	*	*	*	0.2	1.5	3.8
2002.....	15.1	*	*	*	*	*	*	0.3	1.5	3.6
2001.....	15.2	*	*	*	*	*	*	0.4	1.5	3.9
2000.....	15.7	*	*	*	*	*	*	0.4	1.6	3.9
1999.....	15.5	*	*	*	*	*	*	0.4	1.5	3.7
White, male										
2006 4/.....	0.3	*	*	*	*	*	*	*	*	*
2005 4/.....	0.3	*	*	*	*	*	*	*	*	*
2004 4/.....	0.3	*	*	*	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*
White, female										
2006 4/.....	27.9	*	*	*	*	*	*	0.6	2.6	6.6
2005 4/.....	28.3	*	*	*	*	*	*	0.6	2.4	6.7
2004 4/.....	28.6	*	*	*	*	*	*	0.6	2.7	6.9
2003 4/.....	29.3	*	*	*	*	*	*	0.4	3.0	7.7
2002.....	29.5	*	*	*	*	*	*	0.5	3.0	7.2
2001.....	29.8	*	*	*	*	*	*	0.8	3.0	7.8
2000.....	30.7	*	*	*	*	*	*	0.8	3.3	7.7
1999.....	30.2	*	*	*	*	*	*	0.8	3.0	7.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasm of breast (C50)										
Black, both sexes										
2006 4/.....	14.5	*	*	*	*	*	*	0.8	3.2	7.6
2005 4/.....	15.0	*	*	*	*	*	*	0.8	2.9	7.8
2004 4/.....	14.5	*	*	*	*	*	*	0.9	3.2	8.1
2003 4/.....	15.1	*	*	*	*	*	*	0.9	3.4	8.0
2002.....	14.9	*	*	*	*	*	*	0.8	3.7	8.1
2001.....	15.0	*	*	*	*	*	*	1.1	4.0	7.5
2000.....	14.8	*	*	*	*	*	*	0.7	3.7	7.7
1999.....	15.0	*	*	*	*	*	*	0.9	3.7	8.8
Black, male										
2006 4/.....	0.3	*	*	*	*	*	*	*	*	*
2005 4/.....	0.3	*	*	*	*	*	*	*	*	*
2004 4/.....	0.3	*	*	*	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	27.5	*	*	*	*	*	*	1.5	6.0	14.3
2005 4/.....	28.4	*	*	*	*	*	*	1.6	5.6	14.7
2004 4/.....	27.5	*	*	*	*	*	*	1.8	6.2	15.1
2003 4/.....	28.5	*	*	*	*	*	*	1.6	6.5	15.1
2002.....	28.2	*	*	*	*	*	*	1.6	6.9	15.1
2001.....	28.3	*	*	*	*	*	*	2.2	7.6	14.2
2000.....	27.9	*	*	*	*	*	*	*	7.1	14.3
1999.....	28.2	*	*	*	*	*	*	1.8	6.9	16.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of female genital organs (C51-C58)										
All races 3/, both sexes										
2006.....	9.3	*	*	*	*	*	0.1	0.4	1.1	2.0
2005.....	9.2	*	*	*	*	*	0.1	0.4	1.0	1.9
2004.....	9.2	*	*	*	*	*	0.2	0.6	1.1	2.0
2003.....	9.3	*	*	*	*	*	0.2	0.5	1.1	2.0
2002.....	9.4	*	*	*	*	*	0.2	0.4	1.2	2.2
2001.....	9.4	*	*	*	*	*	0.2	0.5	1.2	2.3
2000.....	9.4	*	*	*	*	*	0.2	0.6	1.2	2.2
1999.....	9.3	*	*	*	*	*	0.2	0.5	1.1	2.2
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	18.3	*	*	*	*	*	0.3	0.8	2.1	4.1
2005.....	18.1	*	*	*	*	*	0.3	0.8	2.1	3.9
2004.....	18.1	*	*	*	*	*	0.4	1.1	2.2	4.1
2003.....	18.3	*	*	*	*	*	0.4	1.1	2.1	4.1
2002.....	18.5	*	*	*	*	*	0.5	0.9	2.5	4.5
2001.....	18.5	*	*	*	*	*	0.3	1.1	2.5	4.5
2000.....	18.4	*	*	*	*	*	0.4	1.2	2.4	4.4
1999.....	18.2	*	*	*	*	*	0.4	1.1	2.2	4.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of female genital organs (C51-C58)										
White, both sexes										
2006 4/.....	9.8	*	*	*	*	*	0.1	0.4	1.0	2.1
2005 4/.....	9.7	*	*	*	*	*	0.2	0.4	1.1	1.9
2004 4/.....	9.7	*	*	*	*	*	0.2	0.5	1.1	2.0
2003 4/.....	9.8	*	*	*	*	*	0.2	0.5	1.0	1.9
2002.....	9.9	*	*	*	*	*	0.2	0.4	1.2	2.2
2001.....	9.9	*	*	*	*	*	0.1	0.5	1.2	2.2
2000.....	9.9	*	*	*	*	*	0.2	0.7	1.1	2.1
1999.....	9.8	*	*	*	*	*	0.2	0.5	1.1	2.1
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	19.4	*	*	*	*	*	0.3	0.8	2.1	4.2
2005 4/.....	19.2	*	*	*	*	*	0.3	0.7	2.2	3.8
2004 4/.....	19.2	*	*	*	*	*	0.4	1.0	2.2	4.0
2003 4/.....	19.3	*	*	*	*	*	0.4	1.0	2.0	3.8
2002.....	19.5	*	*	*	*	*	0.5	0.9	2.4	4.4
2001.....	19.6	*	*	*	*	*	0.3	1.1	2.4	4.4
2000.....	19.4	*	*	*	*	*	0.3	1.3	2.3	4.3
1999.....	19.2	*	*	*	*	*	0.4	1.1	2.2	4.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of female genital organs (C51-C58)										
Black, both sexes										
2006 4/.....	8.5	*	*	*	*	*	*	*	1.5	2.3
2005 4/.....	8.5	*	*	*	*	*	*	0.8	1.3	2.4
2004 4/.....	8.6	*	*	*	*	*	*	1.1	1.4	2.8
2003 4/.....	8.5	*	*	*	*	*	*	1.0	1.6	3.2
2002.....	8.6	*	*	*	*	*	*	*	1.7	3.3
2001.....	8.5	*	*	*	*	*	*	0.9	1.7	3.0
2000.....	8.6	*	*	*	*	*	*	*	1.9	3.1
1999.....	8.4	*	*	*	*	*	*	*	1.3	2.9
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	16.3	*	*	*	*	*	*	*	2.8	4.3
2005 4/.....	16.2	*	*	*	*	*	*	1.5	2.5	4.6
2004 4/.....	16.4	*	*	*	*	*	*	2.2	2.7	5.4
2003 4/.....	16.3	*	*	*	*	*	*	2.0	3.1	6.0
2002.....	16.4	*	*	*	*	*	*	*	3.3	6.3
2001.....	16.2	*	*	*	*	*	*	1.7	3.2	5.6
2000.....	16.4	*	*	*	*	*	*	*	3.7	5.8
1999.....	16.1	*	*	*	*	*	*	*	2.5	5.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of cervix uteri (C53)										
All races 3/, both sexes										
2006.....	1.3	*	*	*	*	*	*	0.2	0.7	1.2
2005.....	1.3	*	*	*	*	*	*	0.2	0.6	1.1
2004.....	1.3	*	*	*	*	*	*	0.4	0.7	1.1
2003.....	1.3	*	*	*	*	*	*	0.3	0.7	1.2
2002.....	1.4	*	*	*	*	*	*	0.2	0.8	1.3
2001.....	1.4	*	*	*	*	*	*	0.3	0.8	1.3
2000.....	1.5	*	*	*	*	*	*	0.4	0.7	1.3
1999.....	1.5	*	*	*	*	*	*	0.3	0.8	1.3
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	2.6	*	*	*	*	*	*	0.5	1.4	2.4
2005.....	2.6	*	*	*	*	*	*	0.4	1.3	2.3
2004.....	2.6	*	*	*	*	*	*	0.7	1.4	2.2
2003.....	2.7	*	*	*	*	*	*	0.6	1.4	2.3
2002.....	2.7	*	*	*	*	*	*	0.5	1.5	2.5
2001.....	2.8	*	*	*	*	*	*	0.5	1.5	2.6
2000.....	2.9	*	*	*	*	*	*	0.8	1.5	2.6
1999.....	3.0	*	*	*	*	*	*	0.6	1.5	2.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of cervix uteri (C53)										
White, both sexes										
2006 4/.....	1.2	*	*	*	*	*	*	0.2	0.6	1.2
2005 4/.....	1.2	*	*	*	*	*	*	0.2	0.7	1.1
2004 4/.....	1.2	*	*	*	*	*	*	0.3	0.7	1.1
2003 4/.....	1.2	*	*	*	*	*	*	0.3	0.6	1.1
2002.....	1.3	*	*	*	*	*	*	0.2	0.8	1.2
2001.....	1.3	*	*	*	*	*	*	0.3	0.7	1.3
2000.....	1.4	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	1.4	*	*	*	*	*	*	0.3	0.7	1.2
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	2.5	*	*	*	*	*	*	0.5	1.2	2.4
2005 4/.....	2.5	*	*	*	*	*	*	0.4	1.4	2.2
2004 4/.....	2.4	*	*	*	*	*	*	0.6	1.4	2.2
2003 4/.....	2.5	*	*	*	*	*	*	0.5	1.2	2.1
2002.....	2.5	*	*	*	*	*	*	0.5	1.5	2.4
2001.....	2.7	*	*	*	*	*	*	0.6	1.5	2.5
2000.....	2.7	*	*	*	*	*	*	0.8	1.4	2.4
1999.....	2.8	*	*	*	*	*	*	0.7	1.5	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of cervix uteri (C53)										
Black, both sexes										
2006 4/.....	2.0	*	*	*	*	*	*	*	1.2	1.6
2005 4/.....	2.0	*	*	*	*	*	*	*	0.8	1.6
2004 4/.....	2.0	*	*	*	*	*	*	0.7	0.9	1.6
2003 4/.....	2.1	*	*	*	*	*	*	*	1.2	2.0
2002.....	2.2	*	*	*	*	*	*	*	1.1	2.1
2001.....	2.1	*	*	*	*	*	*	*	1.1	1.8
2000.....	2.4	*	*	*	*	*	*	*	1.4	2.1
1999.....	2.3	*	*	*	*	*	*	*	1.0	2.1
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	3.8	*	*	*	*	*	*	*	2.2	3.0
2005 4/.....	3.8	*	*	*	*	*	*	*	1.5	3.0
2004 4/.....	3.9	*	*	*	*	*	*	1.4	1.7	3.0
2003 4/.....	4.0	*	*	*	*	*	*	*	2.2	3.8
2002.....	4.2	*	*	*	*	*	*	*	2.0	3.9
2001.....	4.0	*	*	*	*	*	*	*	2.1	3.5
2000.....	4.5	*	*	*	*	*	*	*	2.7	4.0
1999.....	4.4	*	*	*	*	*	*	*	1.9	3.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of ovary (C56)										
All races 3/, both sexes										
2006.....	5.0	*	*	*	*	*	*	0.1	0.3	0.5
2005.....	5.0	*	*	*	*	*	*	0.2	0.3	0.5
2004.....	5.0	*	*	*	*	*	0.1	0.2	0.3	0.6
2003.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	5.1	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.1	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.0	*	*	*	*	*	*	0.2	0.3	0.6
1999.....	4.9	*	*	*	*	*	*	0.1	0.2	0.7
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	9.8	*	*	*	*	*	*	0.3	0.5	1.1
2005.....	9.8	*	*	*	*	*	*	0.3	0.5	1.0
2004.....	9.9	*	*	*	*	*	0.2	0.3	0.6	1.2
2003.....	9.9	*	*	*	*	*	*	0.3	0.5	1.3
2002.....	10.0	*	*	*	*	*	0.3	0.4	0.6	1.4
2001.....	9.9	*	*	*	*	*	*	0.4	0.6	1.4
2000.....	9.8	*	*	*	*	*	*	0.4	0.6	1.3
1999.....	9.6	*	*	*	*	*	*	0.3	0.5	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of ovary (C56)										
White, both sexes										
2006 4/.....	5.5	*	*	*	*	*	*	0.1	0.3	0.6
2005 4/.....	5.5	*	*	*	*	*	*	0.2	0.3	0.5
2004 4/.....	5.5	*	*	*	*	*	*	0.1	0.3	0.6
2003 4/.....	5.6	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	5.6	*	*	*	*	*	0.1	0.2	0.3	0.7
2001.....	5.6	*	*	*	*	*	*	0.2	0.3	0.7
2000.....	5.5	*	*	*	*	*	*	0.2	0.3	0.6
1999.....	5.4	*	*	*	*	*	*	0.1	0.2	0.7
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	10.9	*	*	*	*	*	*	0.3	0.6	1.2
2005 4/.....	10.9	*	*	*	*	*	*	0.3	0.5	1.1
2004 4/.....	10.9	*	*	*	*	*	*	0.3	0.6	1.2
2003 4/.....	11.0	*	*	*	*	*	*	0.3	0.6	1.2
2002.....	11.1	*	*	*	*	*	0.3	0.4	0.6	1.4
2001.....	11.0	*	*	*	*	*	*	0.4	0.6	1.3
2000.....	10.9	*	*	*	*	*	*	0.4	0.6	1.3
1999.....	10.6	*	*	*	*	*	*	0.3	0.5	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes

2006.....	4.3	*	0.6	0.9	0.6	0.5	0.5	0.8	1.1	1.6
2005.....	4.4	*	0.5	1.0	0.7	0.6	0.6	0.8	1.1	1.6
2004.....	4.4	*	0.6	0.9	0.7	0.5	0.5	0.7	1.1	1.7
2003.....	4.4	*	0.7	0.9	0.8	0.6	0.6	0.9	1.2	1.9
2002.....	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7
2001.....	4.4	*	0.7	0.8	0.6	0.5	0.6	0.7	1.4	1.8
2000.....	4.5	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0
1999.....	4.6	*	0.7	0.9	0.6	0.6	0.6	0.9	1.3	1.9

All races 3/, male

2006.....	4.9	*	0.7	0.9	0.7	0.5	0.5	0.9	1.2	1.9
2005.....	5.0	*	0.5	0.9	0.7	0.7	0.6	1.1	1.3	2.0
2004.....	4.9	*	0.7	1.0	0.7	0.6	0.7	0.8	1.3	2.1
2003.....	5.0	*	0.7	0.9	0.8	0.6	0.6	0.9	1.4	2.3
2002.....	5.0	*	0.8	1.1	0.7	0.6	0.7	0.7	1.5	2.0
2001.....	5.0	*	0.7	0.8	0.6	0.6	0.7	0.9	1.7	2.2
2000.....	5.1	*	0.7	0.8	0.8	0.6	0.7	1.0	1.6	2.6
1999.....	5.1	*	0.7	0.9	0.6	0.7	0.7	1.1	1.7	2.3

All races 3/, female

2006.....	3.7	*	0.6	0.8	0.6	0.4	0.5	0.7	1.0	1.2
2005.....	3.9	*	0.6	1.0	0.6	0.4	0.5	0.5	0.8	1.2
2004.....	3.9	*	0.6	0.9	0.7	0.5	0.4	0.6	0.8	1.2
2003.....	3.9	*	0.6	0.9	0.7	0.5	0.5	0.9	0.9	1.5
2002.....	3.9	*	0.7	0.9	0.5	0.4	0.3	0.7	1.0	1.4
2001.....	3.9	*	0.6	0.9	0.6	0.4	0.6	0.5	1.1	1.4
2000.....	3.9	*	0.6	0.8	0.6	0.4	0.5	0.7	0.8	1.4
1999.....	4.1	*	0.8	0.9	0.7	0.5	0.5	0.8	0.9	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)										
White, both sexes										
2006 4/.....	4.9	*	0.6	0.9	0.7	0.5	0.5	0.8	1.3	1.8
2005 4/.....	5.0	*	0.5	0.9	0.7	0.6	0.6	0.9	1.2	1.8
2004 4/.....	4.9	*	0.7	1.0	0.6	0.6	0.6	0.7	1.2	1.9
2003 4/.....	5.0	*	0.7	0.9	0.8	0.6	0.6	1.0	1.3	2.1
2002.....	5.0	*	0.7	1.0	0.6	0.5	0.5	0.7	1.4	1.9
2001.....	5.0	*	0.7	0.9	0.6	0.6	0.6	0.8	1.5	2.1
2000.....	5.1	*	0.7	0.8	0.7	0.5	0.7	0.9	1.3	2.2
1999.....	5.2	*	0.8	0.9	0.6	0.5	0.6	1.1	1.4	2.2
White, male										
2006 4/.....	5.5	*	0.7	1.0	0.7	0.5	0.5	1.0	1.4	2.1
2005 4/.....	5.6	*	0.4	0.9	0.8	0.8	0.7	1.2	1.5	2.2
2004 4/.....	5.5	*	0.7	1.0	0.7	0.6	0.7	0.8	1.5	2.4
2003 4/.....	5.6	*	0.8	0.9	0.8	0.7	0.7	1.0	1.6	2.5
2002.....	5.7	*	0.7	1.1	0.7	0.7	0.8	0.7	1.6	2.3
2001.....	5.7	*	0.8	0.8	0.6	0.7	0.7	1.1	1.9	2.6
2000.....	5.7	*	0.7	0.9	0.8	0.6	0.7	1.0	1.7	2.8
1999.....	5.8	*	0.8	0.9	0.6	0.6	0.7	1.2	1.8	2.6
White, female										
2006 4/.....	4.3	*	0.5	0.8	0.6	0.4	0.5	0.6	1.1	1.4
2005 4/.....	4.4	*	0.6	1.0	0.6	0.4	0.5	0.6	0.9	1.3
2004 4/.....	4.4	*	0.6	0.9	0.6	0.5	0.4	0.7	0.9	1.4
2003 4/.....	4.4	*	0.6	0.9	0.8	0.6	0.5	0.9	1.0	1.8
2002.....	4.4	*	0.6	1.0	0.6	0.4	0.3	0.7	1.2	1.6
2001.....	4.4	*	0.7	1.0	0.7	0.4	0.6	0.5	1.2	1.6
2000.....	4.4	*	0.6	0.8	0.7	0.4	0.6	0.8	0.9	1.5
1999.....	4.6	*	0.8	0.8	0.7	0.5	0.5	0.9	1.0	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of brain (C71)										
All races 3/, both sexes										
2006.....	4.2	*	0.6	0.9	0.6	0.5	0.5	0.7	1.1	1.6
2005.....	4.4	*	0.5	1.0	0.6	0.5	0.5	0.8	1.1	1.6
2004.....	4.3	*	0.6	0.9	0.6	0.5	0.5	0.7	1.1	1.6
2003.....	4.4	*	0.6	0.9	0.7	0.6	0.5	0.8	1.1	1.9
2002.....	4.4	*	0.7	1.0	0.6	0.5	0.5	0.7	1.2	1.7
2001.....	4.3	*	0.6	0.8	0.6	0.5	0.6	0.7	1.3	1.8
2000.....	4.4	0.5	0.7	0.8	0.7	0.5	0.6	0.8	1.2	2.0
1999.....	4.5	*	0.7	0.9	0.6	0.5	0.5	0.9	1.3	1.9
All races 3/, male										
2006.....	4.8	*	0.7	0.9	0.7	0.5	0.5	0.9	1.2	1.9
2005.....	4.9	*	0.5	0.9	0.7	0.7	0.6	1.1	1.3	1.9
2004.....	4.8	*	0.7	1.0	0.6	0.6	0.6	0.8	1.3	2.1
2003.....	4.9	*	0.7	0.8	0.8	0.6	0.6	0.9	1.4	2.3
2002.....	4.9	*	0.8	1.1	0.7	0.6	0.7	0.7	1.4	2.0
2001.....	4.9	*	0.6	0.8	0.6	0.6	0.6	0.9	1.6	2.2
2000.....	5.0	*	0.7	0.8	0.8	0.6	0.7	0.9	1.6	2.6
1999.....	5.0	*	0.7	0.9	0.6	0.6	0.6	1.0	1.6	2.2
All races 3/, female										
2006.....	3.7	*	0.6	0.8	0.5	0.4	0.5	0.6	1.0	1.2
2005.....	3.8	*	0.6	1.0	0.5	0.4	0.5	0.5	0.8	1.2
2004.....	3.8	*	0.5	0.9	0.6	0.4	0.4	0.6	0.8	1.2
2003.....	3.8	*	0.6	0.9	0.7	0.5	0.5	0.8	0.9	1.5
2002.....	3.8	*	0.6	0.9	0.5	0.4	0.3	0.7	1.0	1.4
2001.....	3.8	*	0.6	0.9	0.6	0.4	0.6	0.5	1.0	1.4
2000.....	3.9	*	0.6	0.7	0.6	0.4	0.5	0.7	0.7	1.4
1999.....	4.0	*	0.7	0.9	0.6	0.4	0.5	0.8	0.9	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of brain (C71)										
White, both sexes										
2006 4/.....	4.8	*	0.6	0.9	0.6	0.5	0.5	0.8	1.2	1.7
2005 4/.....	5.0	*	0.5	0.9	0.7	0.6	0.6	0.9	1.2	1.7
2004 4/.....	4.9	*	0.6	0.9	0.6	0.5	0.6	0.7	1.2	1.9
2003 4/.....	4.9	*	0.7	0.9	0.8	0.6	0.6	0.9	1.3	2.1
2002.....	4.9	*	0.6	1.0	0.6	0.5	0.5	0.7	1.3	1.9
2001.....	4.9	*	0.7	0.9	0.6	0.5	0.6	0.8	1.5	2.1
2000.....	5.0	*	0.7	0.8	0.7	0.5	0.6	0.9	1.3	2.2
1999.....	5.1	*	0.8	0.8	0.6	0.5	0.6	1.0	1.4	2.1
White, male										
2006 4/.....	5.5	*	0.7	1.0	0.7	0.5	0.5	1.0	1.4	2.1
2005 4/.....	5.6	*	0.4	0.9	0.7	0.7	0.6	1.2	1.5	2.1
2004 4/.....	5.4	*	0.7	1.0	0.6	0.6	0.7	0.8	1.4	2.3
2003 4/.....	5.5	*	0.7	0.8	0.8	0.7	0.7	1.0	1.6	2.4
2002.....	5.5	*	0.7	1.0	0.7	0.6	0.7	0.7	1.5	2.2
2001.....	5.6	*	0.7	0.8	0.6	0.6	0.6	1.0	1.8	2.5
2000.....	5.6	*	0.7	0.9	0.8	0.6	0.7	1.0	1.7	2.8
1999.....	5.6	*	0.8	0.9	0.6	0.6	0.6	1.2	1.8	2.5
White, female										
2006 4/.....	4.2	*	0.5	0.8	0.5	0.4	0.5	0.6	1.1	1.4
2005 4/.....	4.4	*	0.6	1.0	0.6	0.4	0.5	0.5	0.9	1.3
2004 4/.....	4.3	*	0.6	0.9	0.6	0.4	0.4	0.6	0.9	1.4
2003 4/.....	4.3	*	0.6	0.9	0.7	0.6	0.5	0.9	0.9	1.7
2002.....	4.3	*	0.5	1.0	0.6	0.4	0.3	0.7	1.1	1.6
2001.....	4.3	*	0.6	1.0	0.7	0.4	0.6	0.5	1.2	1.6
2000.....	4.4	*	0.6	0.8	0.6	0.4	0.6	0.8	0.9	1.5
1999.....	4.5	*	0.8	0.8	0.6	0.4	0.5	0.9	1.0	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)										
All races 3/, both sexes										
2006.....	0.7	*	0.4	0.4	0.1	*	*	*	0.1	0.1
2005.....	0.7	*	0.4	0.3	0.1	*	*	*	0.1	0.2
2004.....	0.7	*	0.5	0.4	0.1	*	0.1	*	*	0.2
2003.....	0.7	*	0.4	0.4	0.1	*	*	*	0.1	0.2
2002.....	0.7	*	0.5	0.4	0.1	*	*	*	0.1	0.1
2001.....	0.7	*	0.5	0.3	0.1	*	*	0.1	*	0.2
2000.....	0.7	*	0.5	0.3	0.1	*	*	*	0.1	0.2
1999.....	0.7	*	0.5	0.4	0.1	*	*	*	0.1	0.2
All races 3/, male										
2006.....	0.6	*	0.4	0.4	*	*	*	*	*	*
2005.....	0.7	*	0.5	0.4	*	*	*	*	*	*
2004.....	0.6	*	0.5	0.4	*	*	*	*	*	*
2003.....	0.6	*	0.5	0.5	*	*	*	*	*	0.2
2002.....	0.6	*	0.4	0.4	*	*	*	*	*	*
2001.....	0.7	*	0.5	0.3	*	*	*	*	*	*
2000.....	0.7	*	0.5	0.3	*	*	*	*	*	*
1999.....	0.7	*	0.5	0.4	*	*	*	*	*	0.2
All races 3/, female										
2006.....	0.8	*	0.4	0.3	*	*	*	*	*	*
2005.....	0.8	*	0.4	0.3	*	*	*	*	*	*
2004.....	0.8	*	0.5	0.3	*	*	*	*	*	*
2003.....	0.7	*	0.4	0.4	*	*	*	*	*	0.2
2002.....	0.8	*	0.5	0.3	*	*	*	*	*	*
2001.....	0.8	*	0.4	0.2	*	*	*	*	*	*
2000.....	0.8	*	0.5	0.3	*	*	*	*	*	0.2
1999.....	0.8	*	0.4	0.4	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Malignant neoplasms of thyroid and other endocrine glands (C73-C75)

White, both sexes

2006 4/.....	0.8	*	0.4	0.4	*	*	*	*	0.1	0.2
2005 4/.....	0.8	*	0.4	0.3	*	*	*	*	0.2	0.2
2004 4/.....	0.8	*	0.5	0.3	*	*	*	*	*	0.2
2003 4/.....	0.7	*	0.4	0.4	*	*	*	*	*	0.2
2002.....	0.8	*	0.4	0.3	0.1	*	*	*	*	0.2
2001.....	0.8	*	0.5	0.3	*	*	*	*	*	0.1
2000.....	0.8	*	0.5	0.4	*	*	*	*	*	0.2
1999.....	0.8	*	0.5	0.4	*	*	*	*	*	0.2

White, male

2006 4/.....	0.7	*	0.4	0.4	*	*	*	*	*	*
2005 4/.....	0.7	*	0.5	0.4	*	*	*	*	*	*
2004 4/.....	0.7	*	0.5	0.4	*	*	*	*	*	*
2003 4/.....	0.7	*	0.5	0.5	*	*	*	*	*	0.2
2002.....	0.7	*	0.4	0.4	*	*	*	*	*	*
2001.....	0.7	*	0.6	0.3	*	*	*	*	*	*
2000.....	0.7	*	0.6	0.4	*	*	*	*	*	*
1999.....	0.7	*	0.6	0.4	*	*	*	*	*	0.2

White, female

2006 4/.....	0.9	*	0.3	0.3	*	*	*	*	*	*
2005 4/.....	0.8	*	0.3	*	*	*	*	*	*	*
2004 4/.....	0.8	*	0.5	*	*	*	*	*	*	*
2003 4/.....	0.8	*	0.4	0.4	*	*	*	*	*	*
2002.....	0.8	*	0.4	0.3	*	*	*	*	*	*
2001.....	0.8	*	0.5	*	*	*	*	*	*	*
2000.....	0.8	*	0.4	0.3	*	*	*	*	*	*
1999.....	0.8	*	0.4	0.4	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

All races 3/, both sexes

2006.....	13.3	*	*	*	*	0.2	0.3	0.5	0.6	1.2
2005.....	13.5	*	0.1	*	0.1	0.2	0.3	0.4	0.7	1.4
2004.....	13.5	*	*	*	*	0.2	0.3	0.5	0.8	1.4
2003.....	13.8	*	*	*	0.1	0.2	0.3	0.4	0.6	1.4
2002.....	14.1	*	0.1	*	*	0.3	0.3	0.4	0.9	1.5
2001.....	14.1	*	*	*	0.1	0.2	0.3	0.5	0.8	1.6
2000.....	14.5	*	0.1	*	0.1	0.2	0.3	0.4	0.7	1.6
1999.....	15.7	*	*	*	0.1	0.2	0.4	0.5	0.9	1.7

All races 3/, male

2006.....	13.9	*	*	*	*	0.2	0.4	0.6	0.6	1.1
2005.....	14.1	*	*	*	*	0.2	0.3	0.5	0.7	1.4
2004.....	14.1	*	*	*	*	0.3	0.4	0.5	0.8	1.5
2003.....	14.3	*	*	*	*	0.3	0.3	0.4	0.7	1.3
2002.....	14.7	*	*	*	*	0.3	0.4	0.4	0.9	1.6
2001.....	14.5	*	*	*	*	0.3	0.3	0.5	0.9	1.5
2000.....	14.8	*	*	*	*	0.2	0.3	0.5	0.7	1.5
1999.....	16.2	*	*	*	*	0.2	0.5	0.6	1.1	1.7

All races 3/, female

2006.....	12.7	*	*	*	*	*	0.2	0.5	0.6	1.3
2005.....	13.0	*	*	*	*	0.2	0.2	0.3	0.6	1.5
2004.....	13.0	*	*	*	*	*	*	0.5	0.8	1.3
2003.....	13.4	*	*	*	*	*	0.3	0.4	0.6	1.5
2002.....	13.5	*	*	*	*	0.2	0.2	0.3	0.8	1.5
2001.....	13.6	*	*	*	*	*	0.2	0.5	0.6	1.6
2000.....	14.1	*	*	*	*	0.2	0.3	0.4	0.7	1.7
1999.....	15.3	*	*	*	*	*	0.3	0.4	0.8	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other malignant neoplasms of other and unspecified sites (C76-C80,C97)										
White, both sexes										
2006 4/.....	14.1	*	*	*	*	0.2	0.3	0.5	0.6	1.1
2005 4/.....	14.4	*	*	*	0.1	0.2	0.2	0.3	0.6	1.4
2004 4/.....	14.4	*	*	*	*	0.2	0.3	0.5	0.8	1.4
2003 4/.....	14.7	*	*	*	*	0.2	0.3	0.4	0.6	1.3
2002.....	14.9	*	*	*	*	0.3	0.3	0.3	0.9	1.4
2001.....	14.8	*	*	*	*	0.2	0.3	0.4	0.7	1.5
2000.....	15.3	*	*	*	*	0.2	0.3	0.5	0.7	1.5
1999.....	16.6	*	*	*	*	0.2	0.4	0.5	0.9	1.6
White, male										
2006 4/.....	14.8	*	*	*	*	*	0.4	0.5	0.6	1.1
2005 4/.....	14.9	*	*	*	*	*	0.3	0.4	0.7	1.4
2004 4/.....	14.8	*	*	*	*	0.3	0.4	0.5	0.9	1.5
2003 4/.....	15.1	*	*	*	*	0.3	0.3	0.5	0.6	1.3
2002.....	15.5	*	*	*	*	0.3	0.4	0.4	1.0	1.4
2001.....	15.1	*	*	*	*	0.3	0.4	0.5	0.9	1.4
2000.....	15.5	*	*	*	*	*	0.3	0.5	0.7	1.4
1999.....	17.0	*	*	*	*	*	0.5	0.7	1.0	1.6
White, female										
2006 4/.....	13.5	*	*	*	*	*	*	0.4	0.6	1.2
2005 4/.....	14.0	*	*	*	*	*	*	0.3	0.6	1.4
2004 4/.....	13.9	*	*	*	*	*	*	0.5	0.8	1.3
2003 4/.....	14.4	*	*	*	*	*	0.3	0.4	0.5	1.4
2002.....	14.4	*	*	*	*	*	*	0.3	0.7	1.5
2001.....	14.6	*	*	*	*	*	*	0.4	0.6	1.5
2000.....	15.2	*	*	*	*	*	0.3	0.4	0.6	1.6
1999.....	16.2	*	*	*	*	*	*	0.4	0.8	1.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes

2006.....	18.4	0.8	0.8	0.6	0.9	1.3	1.9	1.9	2.2	2.8
2005.....	18.6	0.6	0.8	0.8	0.9	1.3	1.8	1.8	2.2	2.7
2004.....	18.6	0.6	0.8	0.9	0.9	1.4	1.7	1.8	2.3	2.7
2003.....	19.1	0.8	0.9	0.9	1.0	1.2	1.7	1.9	2.4	3.1
2002.....	19.5	0.7	0.8	0.8	1.0	1.5	1.9	1.9	2.6	3.2
2001.....	19.8	0.7	1.0	0.8	1.0	1.5	2.0	2.3	2.7	3.5
2000.....	20.1	1.0	0.9	0.8	1.0	1.6	2.1	2.1	2.5	3.4
1999.....	20.2	0.8	0.9	0.7	1.0	1.5	2.2	2.2	2.6	3.7

All races 3/, male

2006.....	20.3	*	0.9	0.7	1.0	1.6	2.3	2.3	2.6	3.1
2005.....	20.6	*	1.0	0.9	1.0	1.6	2.2	2.1	2.4	3.0
2004.....	20.4	*	0.9	1.1	1.0	1.6	2.1	2.1	2.7	3.2
2003.....	21.0	*	1.0	1.0	1.1	1.6	2.0	2.3	3.0	3.5
2002.....	21.4	*	0.8	0.9	1.2	1.8	2.2	2.3	3.0	3.7
2001.....	21.5	*	1.0	0.9	1.1	1.8	2.3	2.6	3.1	4.2
2000.....	21.6	1.1	1.0	0.9	1.2	1.9	2.6	2.3	3.0	3.9
1999.....	21.7	*	1.0	0.8	1.2	1.7	2.7	2.4	3.1	4.4

All races 3/, female

2006.....	16.5	*	0.8	0.5	0.7	0.9	1.3	1.5	1.8	2.4
2005.....	16.6	*	0.6	0.6	0.7	1.0	1.3	1.5	2.0	2.3
2004.....	16.8	*	0.6	0.7	0.7	1.1	1.3	1.5	1.9	2.2
2003.....	17.3	1.1	0.8	0.7	0.9	0.9	1.4	1.6	1.7	2.6
2002.....	17.7	*	0.7	0.8	0.8	1.2	1.7	1.5	2.2	2.6
2001.....	18.1	*	1.0	0.7	0.8	1.2	1.7	2.0	2.2	2.7
2000.....	18.6	*	0.9	0.7	0.8	1.2	1.5	1.9	1.9	3.0
1999.....	18.7	*	0.8	0.6	0.9	1.3	1.8	1.9	2.2	3.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
White, both sexes										
2006 4/.....	20.1	0.8	0.9	0.6	0.9	1.3	1.8	2.0	2.1	2.6
2005 4/.....	20.3	0.6	0.7	0.8	0.9	1.4	1.8	1.6	2.1	2.7
2004 4/.....	20.2	0.6	0.8	1.0	0.9	1.4	1.7	1.7	2.2	2.6
2003 4/.....	20.9	0.9	0.9	0.9	1.0	1.3	1.7	1.9	2.3	3.0
2002.....	21.3	*	0.8	0.9	1.0	1.5	2.1	1.9	2.5	3.1
2001.....	21.5	0.7	1.1	0.8	1.0	1.6	2.0	2.1	2.6	3.4
2000.....	21.9	1.2	1.0	0.9	1.1	1.6	2.0	2.1	2.4	3.3
1999.....	21.9	0.8	1.0	0.7	1.0	1.5	2.3	2.1	2.5	3.6
White, male										
2006 4/.....	22.1	*	0.9	0.6	1.0	1.7	2.4	2.4	2.5	3.0
2005 4/.....	22.5	*	0.9	0.9	1.1	1.7	2.2	1.9	2.2	3.1
2004 4/.....	22.2	*	0.8	1.2	1.0	1.7	2.1	2.1	2.6	3.1
2003 4/.....	23.1	*	1.1	1.2	1.1	1.6	2.0	2.2	2.9	3.4
2002.....	23.4	*	0.9	0.9	1.2	1.9	2.4	2.3	2.9	3.5
2001.....	23.3	*	1.1	0.9	1.0	1.9	2.3	2.4	3.2	4.0
2000.....	23.6	*	1.0	0.9	1.2	1.9	2.5	2.3	2.9	3.8
1999.....	23.6	*	1.0	0.9	1.1	1.7	2.8	2.3	2.9	4.3
White, female										
2006 4/.....	18.0	*	0.8	0.6	0.7	0.9	1.3	1.5	1.7	2.3
2005 4/.....	18.1	*	0.5	0.7	0.7	1.0	1.4	1.3	1.9	2.3
2004 4/.....	18.3	*	0.7	0.7	0.8	1.1	1.3	1.3	1.8	2.1
2003 4/.....	18.8	1.4	0.8	0.7	0.9	0.8	1.5	1.6	1.7	2.5
2002.....	19.2	*	0.8	0.9	0.8	1.1	1.7	1.5	2.2	2.6
2001.....	19.7	*	1.1	0.7	0.9	1.2	1.6	1.9	2.0	2.8
2000.....	20.3	*	1.0	0.8	0.9	1.2	1.4	1.8	1.9	2.9
1999.....	20.4	*	0.9	0.6	0.9	1.3	1.7	1.9	2.1	2.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
Black, both sexes										
2006 4/.....	13.3	*	*	*	0.9	1.2	2.0	1.9	2.9	4.1
2005 4/.....	12.9	*	1.0	*	0.8	1.3	1.9	2.7	3.8	3.2
2004 4/.....	13.6	*	*	0.7	0.8	1.4	1.9	2.8	3.1	3.7
2003 4/.....	13.4	*	*	0.6	1.0	1.1	1.7	2.4	3.3	4.1
2002.....	13.6	*	*	*	1.0	1.4	1.5	2.3	3.4	3.8
2001.....	14.1	*	*	0.9	1.0	1.5	2.1	3.2	3.5	4.5
2000.....	13.9	*	0.8	0.6	0.7	1.9	2.6	2.8	3.1	4.4
1999.....	14.1	*	*	0.6	1.3	1.7	2.9	2.6	3.2	4.6
Black, male										
2006 4/.....	14.9	*	*	*	*	1.5	2.4	2.1	3.5	4.9
2005 4/.....	14.1	*	*	*	*	1.4	2.5	2.8	4.7	3.7
2004 4/.....	14.4	*	*	*	*	1.4	2.5	2.9	3.6	4.3
2003 4/.....	13.9	*	*	*	1.2	*	2.1	2.5	4.2	4.7
2002.....	14.6	*	*	*	1.1	1.6	1.3	2.8	4.4	5.1
2001.....	15.1	*	*	*	1.7	1.8	2.2	3.8	4.0	6.5
2000.....	14.7	*	*	*	*	2.3	3.2	2.6	4.4	4.8
1999.....	15.1	*	*	*	1.7	2.1	3.1	3.3	4.2	5.3
Black, female										
2006 4/.....	11.9	*	*	*	*	*	1.7	1.8	2.3	3.3
2005 4/.....	11.8	*	*	*	*	1.3	*	2.6	3.0	2.8
2004 4/.....	12.8	*	*	*	*	1.4	1.3	2.7	2.7	3.1
2003 4/.....	12.9	*	*	*	*	*	1.3	2.4	2.4	3.6
2002.....	12.8	*	*	*	*	1.3	1.6	2.0	2.5	2.7
2001.....	13.1	*	*	*	*	*	2.1	2.7	3.1	2.8
2000.....	13.1	*	*	*	*	1.4	2.1	3.0	1.9	4.1
1999.....	13.2	*	*	*	*	1.4	2.8	2.0	2.3	3.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hodgkin's disease (C81)										
All races 3/, both sexes										
2006.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3
2005.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3
2004.....	0.4	*	*	*	*	*	0.2	0.3	0.3	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.3
1999.....	0.5	*	*	*	*	0.2	0.3	0.4	0.3	0.4
All races 3/, male										
2006.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4
2005.....	0.5	*	*	*	*	*	0.2	0.3	0.3	0.3
2004.....	0.5	*	*	*	*	*	0.3	0.3	0.4	0.3
2003.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.5	*	*	*	*	*	0.3	0.3	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.4	0.4	0.4	0.5
All races 3/, female										
2006.....	0.4	*	*	*	*	*	*	0.3	0.2	0.3
2005.....	0.4	*	*	*	*	*	0.3	0.2	0.3	0.2
2004.....	0.4	*	*	*	*	*	*	0.3	0.3	0.2
2003.....	0.4	*	*	*	*	*	0.2	0.4	0.4	0.3
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.3
2001.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.3
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.4	0.3
1999.....	0.4	*	*	*	*	*	0.3	0.4	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hodgkin's disease (C81)										
White, both sexes										
2006 4/.....	0.5	*	*	*	*	*	0.2	0.4	0.3	0.4
2005 4/.....	0.5	*	*	*	*	*	0.2	0.2	0.2	0.3
2004 4/.....	0.5	*	*	*	*	*	0.2	0.3	0.3	0.3
2003 4/.....	0.5	*	*	*	*	*	0.2	0.4	0.5	0.4
2002.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2001.....	0.5	*	*	*	*	0.1	0.3	0.4	0.4	0.4
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999.....	0.6	*	*	*	*	0.2	0.3	0.4	0.4	0.4
White, male										
2006 4/.....	0.6	*	*	*	*	*	0.3	0.4	0.4	0.4
2005 4/.....	0.5	*	*	*	*	*	*	0.3	*	0.3
2004 4/.....	0.5	*	*	*	*	*	0.3	0.3	0.4	0.3
2003 4/.....	0.5	*	*	*	*	*	*	0.4	0.5	0.4
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.6	*	*	*	*	*	0.3	0.4	0.5	0.5
2000.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.4
1999.....	0.6	*	*	*	*	*	0.4	0.4	0.5	0.5
White, female										
2006 4/.....	0.4	*	*	*	*	*	*	0.3	*	0.3
2005 4/.....	0.4	*	*	*	*	*	0.3	*	0.3	*
2004 4/.....	0.4	*	*	*	*	*	*	0.3	0.3	*
2003 4/.....	0.5	*	*	*	*	*	*	0.4	0.4	0.3
2002.....	0.5	*	*	*	*	*	0.4	0.3	0.5	0.3
2001.....	0.4	*	*	*	*	*	*	0.4	0.3	0.3
2000.....	0.5	*	*	*	*	*	*	0.3	0.4	0.3
1999.....	0.5	*	*	*	*	*	0.3	0.4	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

All races 3/, both sexes

2006.....	6.9	*	*	*	0.1	0.2	0.4	0.5	0.6	0.9
2005.....	7.1	*	*	0.1	0.1	0.2	0.4	0.4	0.7	0.9
2004.....	7.2	*	*	0.1	*	0.3	0.4	0.6	0.7	0.9
2003.....	7.4	*	*	*	0.1	0.3	0.4	0.5	0.8	1.1
2002.....	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2
2001.....	7.9	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4
2000.....	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3
1999.....	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4

All races 3/, male

2006.....	7.4	*	*	*	*	0.3	0.5	0.6	0.7	1.0
2005.....	7.7	*	*	*	0.2	0.3	0.5	0.5	0.8	1.1
2004.....	7.7	*	*	*	*	0.3	0.5	0.9	0.8	1.2
2003.....	8.0	*	*	*	*	0.3	0.5	0.6	1.0	1.4
2002.....	8.2	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5
2001.....	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8
2000.....	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7
1999.....	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8

All races 3/, female

2006.....	6.4	*	*	*	*	*	0.3	0.4	0.5	0.7
2005.....	6.5	*	*	*	*	*	0.3	0.4	0.5	0.7
2004.....	6.7	*	*	*	*	*	0.3	0.4	0.6	0.6
2003.....	6.8	*	*	*	*	*	0.3	0.4	0.5	0.7
2002.....	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8
2001.....	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0
2000.....	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9
1999.....	7.8	*	*	*	*	0.2	0.4	0.4	0.7	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)										
White, both sexes										
2006 4/.....	7.7	*	*	*	*	0.2	0.4	0.5	0.6	0.7
2005 4/.....	7.9	*	*	*	0.2	0.2	0.3	0.4	0.6	0.9
2004 4/.....	8.0	*	*	*	*	0.3	0.4	0.6	0.6	0.9
2003 4/.....	8.3	*	*	*	*	0.2	0.4	0.4	0.7	1.1
2002.....	8.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1
2001.....	8.8	*	*	*	0.1	0.3	0.4	0.6	0.8	1.3
2000.....	9.1	*	*	*	0.1	0.3	0.6	0.6	0.8	1.2
1999.....	9.2	*	*	*	*	0.2	0.5	0.5	0.8	1.4
White, male										
2006 4/.....	8.3	*	*	*	*	0.3	0.5	0.6	0.7	0.9
2005 4/.....	8.5	*	*	*	0.3	0.4	0.5	0.4	0.7	1.1
2004 4/.....	8.5	*	*	*	*	0.4	0.5	0.8	0.7	1.3
2003 4/.....	9.0	*	*	*	*	0.3	0.5	0.6	0.9	1.4
2002.....	9.1	*	*	*	0.3	0.3	0.6	0.8	0.9	1.4
2001.....	9.3	*	*	*	*	0.4	0.5	0.7	1.0	1.6
2000.....	9.6	*	*	*	*	0.5	0.8	0.6	1.0	1.6
1999.....	9.6	*	*	*	*	0.3	0.6	0.6	1.0	1.7
White, female										
2006 4/.....	7.2	*	*	*	*	*	*	0.4	0.4	0.6
2005 4/.....	7.3	*	*	*	*	*	*	0.4	0.5	0.6
2004 4/.....	7.5	*	*	*	*	*	0.3	0.3	0.5	0.6
2003 4/.....	7.7	*	*	*	*	*	0.4	0.3	0.4	0.7
2002.....	8.0	*	*	*	*	*	0.3	0.4	0.6	0.8
2001.....	8.3	*	*	*	*	*	0.4	0.5	0.5	1.0
2000.....	8.7	*	*	*	*	*	0.4	0.5	0.6	0.8
1999.....	8.8	*	*	*	*	*	0.4	0.4	0.6	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Non-Hodgkin's lymphoma (C82-C85)										
All races 3/, both sexes										
2006.....	6.9	*	*	*	0.1	0.2	0.4	0.5	0.6	0.8
2005.....	7.0	*	*	*	0.1	0.2	0.4	0.4	0.7	0.9
2004.....	7.1	*	*	0.1	*	0.3	0.4	0.6	0.7	0.9
2003.....	7.4	*	*	*	0.1	0.3	0.4	0.5	0.8	1.1
2002.....	7.6	*	*	*	0.2	0.2	0.5	0.6	0.8	1.2
2001.....	7.8	*	*	*	0.1	0.3	0.4	0.7	0.8	1.4
2000.....	8.1	*	*	0.1	0.1	0.3	0.6	0.6	0.8	1.3
1999.....	8.2	*	*	*	0.1	0.3	0.5	0.6	0.9	1.4
All races 3/, male										
2006.....	7.4	*	*	*	*	0.3	0.5	0.6	0.7	1.0
2005.....	7.6	*	*	*	0.2	0.3	0.5	0.5	0.8	1.1
2004.....	7.6	*	*	*	*	0.3	0.5	0.8	0.8	1.2
2003.....	8.0	*	*	*	*	0.3	0.5	0.6	1.0	1.4
2002.....	8.1	*	*	*	0.3	0.3	0.6	0.7	1.0	1.5
2001.....	8.4	*	*	*	0.2	0.4	0.5	0.8	1.0	1.8
2000.....	8.6	*	*	*	*	0.4	0.8	0.7	1.0	1.7
1999.....	8.6	*	*	*	0.2	0.3	0.6	0.7	1.1	1.8
All races 3/, female										
2006.....	6.4	*	*	*	*	*	0.3	0.4	0.5	0.7
2005.....	6.5	*	*	*	*	*	0.3	0.4	0.5	0.7
2004.....	6.6	*	*	*	*	*	0.3	0.4	0.6	0.6
2003.....	6.8	*	*	*	*	*	0.3	0.4	0.5	0.7
2002.....	7.1	*	*	*	*	*	0.4	0.4	0.6	0.8
2001.....	7.3	*	*	*	*	*	0.4	0.6	0.6	1.0
2000.....	7.6	*	*	*	*	*	0.4	0.6	0.6	0.9
1999.....	7.7	*	*	*	*	0.2	0.4	0.4	0.7	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Non-Hodgkin's lymphoma (C82-C85)										
White, both sexes										
2006 4/.....	7.7	*	*	*	*	0.2	0.4	0.5	0.6	0.7
2005 4/.....	7.9	*	*	*	0.2	0.2	0.3	0.4	0.6	0.8
2004 4/.....	8.0	*	*	*	*	0.3	0.4	0.5	0.6	0.9
2003 4/.....	8.3	*	*	*	*	0.2	0.4	0.4	0.7	1.1
2002.....	8.5	*	*	*	0.2	0.2	0.5	0.6	0.7	1.1
2001.....	8.8	*	*	*	0.1	0.3	0.4	0.6	0.8	1.3
2000.....	9.1	*	*	*	0.1	0.3	0.6	0.6	0.8	1.2
1999.....	9.2	*	*	*	*	0.2	0.5	0.5	0.8	1.4
White, male										
2006 4/.....	8.2	*	*	*	*	0.3	0.5	0.6	0.7	0.9
2005 4/.....	8.5	*	*	*	0.3	0.4	0.5	0.4	0.7	1.1
2004 4/.....	8.5	*	*	*	*	0.4	0.5	0.8	0.7	1.2
2003 4/.....	8.9	*	*	*	*	0.3	0.5	0.6	0.9	1.4
2002.....	9.1	*	*	*	0.3	0.3	0.6	0.8	0.9	1.4
2001.....	9.3	*	*	*	*	0.4	0.5	0.7	1.0	1.6
2000.....	9.6	*	*	*	*	0.5	0.8	0.6	1.0	1.6
1999.....	9.6	*	*	*	*	0.3	0.6	0.6	1.0	1.7
White, female										
2006 4/.....	7.2	*	*	*	*	*	*	0.4	0.4	0.6
2005 4/.....	7.3	*	*	*	*	*	*	0.3	0.5	0.6
2004 4/.....	7.5	*	*	*	*	*	0.3	0.3	0.5	0.6
2003 4/.....	7.7	*	*	*	*	*	0.4	0.3	0.4	0.7
2002.....	8.0	*	*	*	*	*	0.3	0.4	0.6	0.8
2001.....	8.2	*	*	*	*	*	0.4	0.5	0.5	1.0
2000.....	8.6	*	*	*	*	*	0.4	0.5	0.6	0.8
1999.....	8.8	*	*	*	*	*	0.4	0.4	0.6	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
All races 3/, both sexes										
2006.....	7.3	0.8	0.8	0.5	0.7	1.0	1.2	1.1	1.2	1.4
2005.....	7.3	0.5	0.7	0.6	0.7	1.1	1.2	1.1	1.2	1.4
2004.....	7.3	0.6	0.8	0.8	0.7	1.0	1.1	0.9	1.2	1.4
2003.....	7.4	0.7	0.8	0.8	0.9	0.9	1.1	1.1	1.1	1.5
2002.....	7.5	0.5	0.7	0.7	0.8	1.2	1.2	1.0	1.3	1.5
2001.....	7.5	0.6	1.0	0.8	0.8	1.1	1.3	1.2	1.4	1.6
2000.....	7.6	1.0	0.9	0.7	0.8	1.2	1.2	1.2	1.2	1.6
1999.....	7.5	0.7	0.8	0.6	0.9	1.1	1.4	1.2	1.3	1.6
All races 3/, male										
2006.....	8.4	*	0.8	0.6	0.8	1.2	1.5	1.3	1.4	1.6
2005.....	8.4	*	1.0	0.8	0.8	1.2	1.5	1.3	1.2	1.4
2004.....	8.3	*	0.9	1.0	0.8	1.2	1.3	1.0	1.4	1.5
2003.....	8.4	*	0.9	0.9	1.0	1.2	1.2	1.3	1.4	1.6
2002.....	8.5	*	0.7	0.7	0.9	1.4	1.4	1.2	1.5	1.6
2001.....	8.5	*	1.0	0.8	0.9	1.3	1.5	1.4	1.6	1.8
2000.....	8.5	1.1	0.9	0.8	1.0	1.4	1.5	1.3	1.5	1.6
1999.....	8.4	*	0.8	0.7	0.9	1.3	1.7	1.3	1.4	1.8
All races 3/, female										
2006.....	6.3	*	0.8	0.5	0.6	0.7	0.9	0.8	1.0	1.3
2005.....	6.2	*	0.5	0.5	0.7	0.9	0.8	0.9	1.2	1.3
2004.....	6.3	*	0.6	0.6	0.7	0.8	0.8	0.8	0.9	1.2
2003.....	6.4	1.1	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4
2002.....	6.5	*	0.6	0.8	0.7	1.0	1.0	0.8	1.1	1.3
2001.....	6.6	*	1.0	0.7	0.7	0.9	1.0	1.0	1.2	1.3
2000.....	6.7	*	0.9	0.6	0.6	0.9	0.9	1.0	0.9	1.6
1999.....	6.7	*	0.7	0.5	0.8	0.9	1.1	1.1	1.2	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
White, both sexes										
2006 4/.....	8.1	0.8	0.8	0.5	0.8	1.0	1.2	1.1	1.2	1.4
2005 4/.....	8.1	*	0.7	0.7	0.7	1.1	1.2	1.0	1.2	1.5
2004 4/.....	8.0	*	0.7	0.8	0.8	1.0	1.1	0.9	1.2	1.3
2003 4/.....	8.2	0.9	0.9	0.9	0.9	0.9	1.1	1.1	1.1	1.4
2002.....	8.2	*	0.7	0.8	0.8	1.2	1.3	1.0	1.3	1.5
2001.....	8.3	0.6	1.1	0.7	0.8	1.2	1.3	1.1	1.4	1.5
2000.....	8.3	1.2	1.0	0.7	0.9	1.2	1.2	1.1	1.1	1.6
1999.....	8.3	0.8	0.9	0.7	0.8	1.1	1.4	1.2	1.3	1.6
White, male										
2006 4/.....	9.3	*	0.8	0.5	0.9	1.3	1.6	1.5	1.3	1.6
2005 4/.....	9.3	*	0.9	0.8	0.8	1.3	1.5	1.2	1.3	1.5
2004 4/.....	9.1	*	0.8	1.0	0.8	1.2	1.4	1.0	1.4	1.4
2003 4/.....	9.4	*	1.0	1.0	1.0	1.2	1.2	1.3	1.4	1.4
2002.....	9.3	*	0.8	0.7	0.9	1.4	1.5	1.2	1.5	1.6
2001.....	9.3	*	1.1	0.8	0.8	1.4	1.6	1.3	1.6	1.7
2000.....	9.4	*	0.9	0.8	1.1	1.4	1.5	1.3	1.4	1.7
1999.....	9.2	*	0.9	0.8	0.9	1.2	1.8	1.3	1.4	1.8
White, female										
2006 4/.....	7.0	*	0.8	0.5	0.7	0.8	0.9	0.8	1.0	1.3
2005 4/.....	6.9	*	0.5	0.6	0.7	0.9	0.9	0.8	1.2	1.4
2004 4/.....	6.9	*	0.7	0.7	0.7	0.8	0.9	0.8	1.0	1.2
2003 4/.....	7.0	1.4	0.8	0.7	0.8	0.6	0.9	0.9	0.8	1.4
2002.....	7.1	*	0.6	0.8	0.7	1.0	1.1	0.8	1.1	1.4
2001.....	7.3	*	1.1	0.7	0.8	0.9	1.0	0.9	1.2	1.4
2000.....	7.4	*	1.0	0.7	0.7	0.9	0.9	0.9	0.9	1.6
1999.....	7.3	*	0.8	0.6	0.8	1.0	1.0	1.1	1.2	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Leukemia (C91-C95)										
Black, both sexes										
2006 4/.....	4.6	*	*	*	0.7	0.9	1.2	0.9	1.3	1.5
2005 4/.....	4.4	*	1.0	*	0.7	1.0	0.9	1.5	1.5	1.1
2004 4/.....	4.8	*	*	*	0.6	1.1	0.9	1.1	1.5	1.8
2003 4/.....	4.6	*	*	*	0.8	0.7	1.0	1.0	1.4	2.0
2002.....	4.7	*	*	*	0.7	0.9	0.8	1.1	1.4	1.5
2001.....	4.7	*	*	0.8	0.9	0.9	1.2	1.4	1.4	1.8
2000.....	4.8	*	*	*	0.6	1.5	1.3	1.4	1.8	1.7
1999.....	4.7	*	*	*	1.1	1.2	1.7	1.1	1.4	1.8
Black, male										
2006 4/.....	5.3	*	*	*	*	*	1.5	*	1.6	1.7
2005 4/.....	4.9	*	*	*	*	*	1.6	1.5	*	*
2004 4/.....	5.3	*	*	*	*	*	*	*	1.8	2.1
2003 4/.....	5.0	*	*	*	*	*	1.3	*	1.8	2.5
2002.....	5.1	*	*	*	*	*	*	*	1.6	1.9
2001.....	5.2	*	*	*	1.5	*	*	1.7	1.8	2.3
2000.....	5.3	*	*	*	*	2.0	1.8	*	2.5	1.4
1999.....	5.2	*	*	*	1.5	1.5	1.7	*	1.7	1.8
Black, female										
2006 4/.....	3.9	*	*	*	*	*	*	*	*	*
2005 4/.....	4.0	*	*	*	*	*	*	1.4	1.6	*
2004 4/.....	4.4	*	*	*	*	*	*	*	*	1.5
2003 4/.....	4.2	*	*	*	*	*	*	*	*	1.7
2002.....	4.3	*	*	*	*	*	*	*	*	*
2001.....	4.3	*	*	*	*	*	*	*	*	1.4
2000.....	4.2	*	*	*	*	*	*	1.5	*	1.9
1999.....	4.3	*	*	*	*	*	1.8	*	*	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Lymphoid leukemia (C91)										
All races 3/, both sexes										
2006.....	2.1	*	0.3	0.3	0.4	0.5	0.5	0.3	0.4	0.3
2005.....	2.1	*	0.3	0.3	0.4	0.4	0.5	0.3	0.3	0.3
2004.....	2.1	*	0.3	0.5	0.3	0.5	0.4	0.3	0.3	0.3
2003.....	2.2	*	0.4	0.4	0.4	0.4	0.5	0.3	0.3	0.4
2002.....	2.2	*	0.2	0.4	0.4	0.5	0.5	0.3	0.4	0.4
2001.....	2.2	*	0.3	0.4	0.4	0.5	0.5	0.4	0.4	0.4
2000.....	2.2	*	0.3	0.4	0.5	0.5	0.4	0.3	0.2	0.3
1999.....	2.2	*	0.3	0.3	0.4	0.5	0.5	0.4	0.3	0.3
All races 3/, male										
2006.....	2.5	*	0.3	0.3	0.4	0.5	0.8	0.4	0.5	0.4
2005.....	2.5	*	0.4	0.5	0.4	0.5	0.7	0.5	0.4	0.4
2004.....	2.5	*	0.4	0.5	0.4	0.6	0.6	0.4	0.4	0.4
2003.....	2.6	*	0.4	0.5	0.6	0.6	0.6	0.4	0.4	0.4
2002.....	2.6	*	*	0.4	0.5	0.7	0.6	0.4	0.4	0.5
2001.....	2.5	*	0.3	0.4	0.5	0.7	0.7	0.4	0.5	0.5
2000.....	2.6	*	0.3	0.4	0.6	0.7	0.7	0.4	0.3	0.4
1999.....	2.6	*	0.3	0.4	0.4	0.6	0.7	0.4	0.4	0.4
All races 3/, female										
2006.....	1.7	*	*	0.2	0.3	0.3	0.3	0.3	0.3	0.3
2005.....	1.7	*	*	0.2	0.4	0.3	0.3	0.2	0.3	0.3
2004.....	1.7	*	*	0.4	0.3	0.4	0.3	0.2	0.2	0.3
2003.....	1.8	*	0.4	0.3	0.3	0.3	0.3	0.3	*	0.3
2002.....	1.8	*	*	0.4	0.4	0.4	0.4	*	0.3	0.3
2001.....	1.8	*	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.3
2000.....	1.8	*	0.3	0.3	0.3	0.4	0.2	0.2	*	0.2
1999.....	1.9	*	0.3	0.2	0.3	0.3	0.4	0.3	0.2	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Lymphoid leukemia (C91)										
White, both sexes										
2006 4/.....	2.4	*	0.3	0.3	0.4	0.5	0.6	0.4	0.4	0.3
2005 4/.....	2.4	*	0.3	0.4	0.4	0.5	0.6	0.3	0.4	0.4
2004 4/.....	2.3	*	0.3	0.5	0.4	0.5	0.4	0.3	0.3	0.3
2003 4/.....	2.4	*	0.4	0.5	0.4	0.5	0.5	0.4	0.3	0.3
2002.....	2.4	*	0.2	0.4	0.5	0.5	0.6	0.3	0.4	0.4
2001.....	2.4	*	0.3	0.4	0.4	0.6	0.6	0.4	0.4	0.3
2000.....	2.4	*	0.4	0.4	0.5	0.5	0.5	0.3	0.2	0.3
1999.....	2.5	*	0.4	0.4	0.4	0.5	0.6	0.4	0.3	0.3
White, male										
2006 4/.....	2.8	*	0.3	0.3	0.5	0.6	0.8	0.5	0.5	0.4
2005 4/.....	2.8	*	0.4	0.5	0.5	0.6	0.8	0.4	0.4	0.4
2004 4/.....	2.7	*	0.3	0.5	0.4	0.7	0.6	0.4	0.5	0.4
2003 4/.....	2.9	*	0.5	0.7	0.6	0.6	0.6	0.4	0.4	0.4
2002.....	2.8	*	*	0.4	0.5	0.7	0.7	0.4	0.4	0.5
2001.....	2.8	*	*	0.4	0.5	0.8	0.8	0.4	0.5	0.4
2000.....	2.8	*	0.3	0.5	0.7	0.7	0.7	0.4	0.3	0.4
1999.....	2.9	*	0.4	0.5	0.4	0.6	0.8	0.5	0.4	0.4
White, female										
2006 4/.....	2.0	*	*	0.3	0.3	0.4	0.3	0.3	0.3	0.3
2005 4/.....	1.9	*	*	0.3	0.4	0.3	0.4	*	0.3	0.3
2004 4/.....	1.9	*	*	0.4	0.4	0.4	0.3	*	*	0.3
2003 4/.....	2.0	*	0.4	0.4	0.3	0.3	0.3	0.3	*	0.3
2002.....	2.0	*	*	0.5	0.5	0.3	0.5	*	0.3	0.3
2001.....	2.0	*	*	0.4	0.3	0.3	0.4	0.3	0.3	0.2
2000.....	2.1	*	0.4	0.3	0.3	0.4	*	*	*	*
1999.....	2.1	*	*	0.3	0.3	0.3	0.4	0.3	0.3	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Myeloid leukemia (C92)										
All races 3/, both sexes										
2006.....	3.4	*	0.3	0.1	0.2	0.4	0.5	0.5	0.6	0.9
2005.....	3.4	*	0.4	0.2	0.2	0.4	0.4	0.5	0.6	0.7
2004.....	3.4	*	0.4	0.2	0.3	0.4	0.5	0.5	0.6	0.8
2003.....	3.4	*	0.3	0.2	0.3	0.4	0.4	0.5	0.6	0.8
2002.....	3.4	*	0.3	0.2	0.2	0.4	0.4	0.5	0.7	0.8
2001.....	3.5	*	0.5	0.2	0.2	0.4	0.5	0.6	0.8	1.0
2000.....	3.5	*	0.4	0.2	0.3	0.5	0.5	0.6	0.8	1.1
1999.....	3.3	*	0.4	0.2	0.3	0.4	0.6	0.6	0.7	1.0
All races 3/, male										
2006.....	3.9	*	0.4	*	0.2	0.5	0.5	0.7	0.7	0.9
2005.....	3.8	*	0.4	*	0.3	0.4	0.5	0.6	0.6	0.7
2004.....	3.8	*	0.4	0.3	0.2	0.4	0.5	0.4	0.7	0.8
2003.....	3.8	*	0.3	0.2	0.3	0.5	0.4	0.6	0.7	0.8
2002.....	3.9	*	0.4	*	0.2	0.4	0.4	0.6	0.8	0.9
2001.....	3.9	*	0.5	0.2	0.2	0.4	0.5	0.7	0.8	1.0
2000.....	3.9	*	0.4	*	0.3	0.5	0.6	0.6	0.9	0.9
1999.....	3.7	*	0.4	*	0.3	0.4	0.7	0.6	0.7	1.0
All races 3/, female										
2006.....	3.0	*	0.3	*	0.2	0.3	0.4	0.4	0.6	0.8
2005.....	2.9	*	0.3	*	0.2	0.5	0.3	0.5	0.6	0.8
2004.....	3.0	*	0.3	*	0.3	0.3	0.5	0.5	0.6	0.7
2003.....	3.0	*	*	*	0.4	0.3	0.4	0.4	0.5	0.8
2002.....	3.0	*	*	0.2	0.2	0.4	0.4	0.4	0.6	0.8
2001.....	3.1	*	0.5	*	0.3	0.4	0.5	0.5	0.7	0.9
2000.....	3.1	*	0.3	*	0.2	0.4	0.4	0.6	0.6	1.2
1999.....	3.0	*	0.3	*	0.3	0.4	0.5	0.6	0.6	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Myeloid leukemia (C92)										
White, both sexes										
2006 4/.....	3.8	*	0.3	*	0.2	0.4	0.4	0.5	0.6	0.9
2005 4/.....	3.7	*	0.3	0.2	0.2	0.4	0.4	0.5	0.6	0.8
2004 4/.....	3.7	*	0.3	0.2	0.2	0.3	0.5	0.5	0.6	0.7
2003 4/.....	3.7	*	0.3	0.2	0.3	0.3	0.4	0.6	0.6	0.8
2002.....	3.8	*	0.3	0.2	0.2	0.4	0.4	0.5	0.7	0.8
2001.....	3.8	*	0.5	0.2	0.2	0.4	0.5	0.6	0.8	0.9
2000.....	3.8	*	0.4	0.2	0.3	0.4	0.5	0.5	0.7	1.1
1999.....	3.6	*	0.4	0.1	0.3	0.3	0.6	0.6	0.7	1.0
White, male										
2006 4/.....	4.2	*	0.4	*	*	0.5	0.5	0.7	0.7	0.9
2005 4/.....	4.3	*	0.3	*	0.3	0.4	0.5	0.6	0.6	0.7
2004 4/.....	4.2	*	0.3	0.3	0.2	0.4	0.5	0.4	0.6	0.8
2003 4/.....	4.3	*	*	*	0.2	0.4	0.4	0.6	0.7	0.7
2002.....	4.3	*	0.4	*	0.2	0.4	0.4	0.6	0.8	0.8
2001.....	4.3	*	0.5	*	*	0.4	0.5	0.7	0.9	1.0
2000.....	4.3	*	0.4	*	0.3	0.5	0.5	0.6	0.8	1.0
1999.....	4.0	*	0.3	*	0.3	0.3	0.7	0.6	0.7	1.1
White, female										
2006 4/.....	3.3	*	*	*	*	0.3	0.4	0.3	0.5	0.8
2005 4/.....	3.2	*	*	*	*	0.4	0.4	0.4	0.6	0.8
2004 4/.....	3.3	*	0.3	*	0.3	0.3	0.5	0.5	0.6	0.7
2003 4/.....	3.3	*	*	*	0.4	*	0.4	0.5	0.5	0.9
2002.....	3.3	*	*	*	*	0.4	0.4	0.4	0.6	0.8
2001.....	3.4	*	0.6	*	0.3	0.4	0.5	0.5	0.7	0.9
2000.....	3.3	*	*	*	*	0.3	0.4	0.5	0.6	1.2
1999.....	3.3	*	0.4	*	0.3	0.3	0.5	0.7	0.6	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified leukemia (C94-C95)										
All races 3/, both sexes										
2006.....	1.7	*	0.1	0.1	0.1	0.1	0.2	0.2	0.2	0.2
2005.....	1.8	*	0.1	0.1	0.1	0.2	0.2	0.2	0.2	0.3
2004.....	1.8	*	0.1	0.1	0.1	0.2	0.2	0.1	0.2	0.2
2003.....	1.8	*	0.2	0.1	0.1	0.1	0.2	0.2	0.2	0.3
2002.....	1.8	*	*	0.2	0.1	0.2	0.2	0.2	0.2	0.2
2001.....	1.8	*	0.2	0.1	0.2	0.2	0.2	0.2	0.2	0.2
2000.....	1.9	*	0.2	0.1	0.1	0.2	0.2	0.3	0.2	0.3
1999.....	1.9	*	*	0.2	0.2	0.2	0.3	0.2	0.3	0.3
All races 3/, male										
2006.....	1.9	*	*	*	*	0.2	0.3	0.2	0.2	0.3
2005.....	2.0	*	*	*	*	0.3	0.3	0.3	0.3	0.3
2004.....	2.0	*	*	*	*	0.2	0.2	*	0.3	0.3
2003.....	1.9	*	*	*	*	*	0.2	0.2	0.2	0.3
2002.....	2.0	*	*	*	*	0.3	0.3	0.2	0.3	0.2
2001.....	2.0	*	0.3	*	0.2	*	0.3	0.3	0.3	0.3
2000.....	2.0	*	*	*	*	0.2	0.2	0.4	0.2	0.3
1999.....	2.1	*	*	0.2	*	0.3	0.3	0.2	0.3	0.3
All races 3/, female										
2006.....	1.5	*	*	*	*	*	*	*	*	*
2005.....	1.6	*	*	*	*	*	*	*	0.2	0.2
2004.....	1.6	*	*	*	*	*	*	*	*	0.2
2003.....	1.6	*	*	*	*	*	*	*	*	0.3
2002.....	1.6	*	*	*	*	*	*	*	0.2	0.2
2001.....	1.6	*	*	*	*	0.2	*	*	*	0.2
2000.....	1.7	*	*	*	*	0.2	0.2	0.2	*	0.2
1999.....	1.8	*	*	*	*	0.2	*	*	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other and unspecified leukemia (C94-C95)										
White, both sexes										
2006 4/.....	1.9	*	*	0.1	0.1	0.1	0.2	0.2	0.2	0.2
2005 4/.....	2.0	*	*	*	*	0.2	0.2	0.2	0.2	0.3
2004 4/.....	1.9	*	*	0.1	0.1	0.2	0.2	*	0.2	0.2
2003 4/.....	2.0	*	0.2	0.1	0.1	0.1	0.2	0.2	0.2	0.3
2002.....	2.0	*	*	0.1	*	0.2	0.2	0.2	0.2	0.2
2001.....	2.0	*	0.3	0.1	0.2	0.2	0.2	0.2	0.2	0.3
2000.....	2.1	*	0.2	0.1	*	0.2	0.2	0.3	0.2	0.3
1999.....	2.1	*	*	0.2	0.1	0.3	0.3	0.2	0.3	0.3
White, male										
2006 4/.....	2.2	*	*	*	*	*	0.3	0.2	*	0.3
2005 4/.....	2.1	*	*	*	*	0.3	0.3	*	0.3	0.3
2004 4/.....	2.2	*	*	*	*	*	0.2	*	0.3	0.3
2003 4/.....	2.1	*	*	*	*	*	*	*	0.2	0.3
2002.....	2.2	*	*	*	*	0.3	0.3	*	0.3	0.2
2001.....	2.2	*	*	*	*	*	0.3	0.3	0.3	0.3
2000.....	2.2	*	*	*	*	*	*	0.3	0.2	0.3
1999.....	2.3	*	*	*	*	0.2	0.4	*	0.3	0.3
White, female										
2006 4/.....	1.7	*	*	*	*	*	*	*	*	*
2005 4/.....	1.8	*	*	*	*	*	*	*	*	0.3
2004 4/.....	1.7	*	*	*	*	*	*	*	*	*
2003 4/.....	1.8	*	*	*	*	*	*	*	*	0.3
2002.....	1.8	*	*	*	*	*	*	*	*	0.2
2001.....	1.8	*	*	*	*	*	*	*	*	*
2000.....	1.9	*	*	*	*	*	*	*	*	0.2
1999.....	1.9	*	*	*	*	0.3	*	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)										
All races 3/, both sexes										
2006.....	3.9	0.6	0.3	0.2	0.1	0.1	0.1	0.2	0.3	0.4
2005.....	3.9	0.9	0.2	0.2	0.1	0.2	0.1	0.2	0.3	0.5
2004.....	3.8	0.8	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.4
2003.....	3.9	0.9	0.2	0.2	0.1	0.1	0.2	0.2	0.4	0.5
2002.....	3.8	0.9	0.3	0.2	0.2	0.1	0.2	0.2	0.3	0.4
2001.....	4.0	0.5	0.3	0.2	0.2	0.1	0.2	0.2	0.3	0.5
2000.....	4.0	0.7	0.3	0.3	0.2	0.1	0.2	0.2	0.3	0.5
1999.....	4.0	0.8	0.4	0.3	0.2	0.2	0.2	0.2	0.4	0.4
All races 3/, male										
2006.....	4.1	*	0.2	0.2	*	*	*	0.2	0.3	0.5
2005.....	4.1	*	0.3	*	*	0.2	0.2	0.3	0.4	0.5
2004.....	4.1	*	0.3	*	*	*	0.2	0.3	0.4	0.4
2003.....	4.1	*	0.3	*	*	*	*	0.2	0.4	0.6
2002.....	4.0	*	0.3	*	0.2	*	0.2	0.2	0.3	0.4
2001.....	4.2	*	0.3	0.2	0.2	*	*	0.2	0.3	0.6
2000.....	4.2	*	0.4	0.3	*	*	0.3	0.3	0.3	0.6
1999.....	4.2	*	0.4	0.2	*	*	0.3	0.2	0.5	0.5
All races 3/, female										
2006.....	3.8	*	0.3	*	*	*	*	0.2	0.3	0.3
2005.....	3.7	1.1	*	0.2	*	*	*	0.2	0.3	0.4
2004.....	3.6	*	*	*	*	*	*	*	0.3	0.4
2003.....	3.8	*	*	*	*	*	*	*	0.3	0.3
2002.....	3.7	*	0.3	*	*	*	*	*	0.3	0.3
2001.....	3.9	*	0.3	0.2	0.2	*	*	*	0.3	0.5
2000.....	3.9	*	*	0.2	*	*	*	0.2	0.3	0.4
1999.....	3.8	*	0.3	0.3	*	0.2	*	*	0.3	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

White, both sexes

2006 4/.....	4.4	0.7	0.3	0.2	0.1	0.1	0.1	0.2	0.3	0.4
2005 4/.....	4.3	0.7	0.2	0.2	0.1	0.2	0.2	0.3	0.3	0.5
2004 4/.....	4.3	0.8	0.3	0.2	0.1	0.1	0.2	0.3	0.3	0.4
2003 4/.....	4.4	0.8	0.2	0.1	0.1	0.1	0.2	0.2	0.4	0.4
2002.....	4.3	1.0	0.3	0.2	0.2	0.2	0.2	0.2	0.3	0.4
2001.....	4.5	*	0.3	0.2	0.2	0.1	0.1	0.2	0.3	0.5
2000.....	4.5	0.7	0.3	0.3	0.2	*	0.2	0.2	0.4	0.5
1999.....	4.4	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.4	0.4

White, male

2006 4/.....	4.6	*	*	*	*	*	*	*	0.3	0.5
2005 4/.....	4.6	*	*	*	*	*	*	0.3	0.4	0.5
2004 4/.....	4.5	*	0.3	*	*	*	*	0.3	0.4	0.4
2003 4/.....	4.5	*	*	*	*	*	*	*	0.4	0.6
2002.....	4.5	*	*	*	*	*	*	*	0.4	0.5
2001.....	4.7	*	*	*	*	*	*	0.3	0.3	0.6
2000.....	4.6	*	0.4	0.3	*	*	0.3	0.3	0.4	0.6
1999.....	4.6	*	0.3	0.2	*	*	*	*	0.4	0.5

White, female

2006 4/.....	4.2	*	0.4	*	*	*	*	*	*	0.3
2005 4/.....	4.1	*	*	*	*	*	*	*	0.3	0.4
2004 4/.....	4.0	*	*	*	*	*	*	*	*	0.4
2003 4/.....	4.2	*	*	*	*	*	*	*	0.3	0.3
2002.....	4.1	*	*	*	*	*	*	*	0.3	0.3
2001.....	4.3	*	*	*	*	*	*	*	0.2	0.5
2000.....	4.4	*	*	0.3	*	*	*	*	0.3	0.4
1999.....	4.2	*	0.4	0.3	*	*	*	*	0.3	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

All races 3/, both sexes

2006.....	3.0	2.5	0.4	0.2	0.2	0.3	0.5	0.7	0.7	1.0
2005.....	3.2	2.3	0.5	0.1	0.2	0.3	0.5	0.7	0.7	1.2
2004.....	3.2	2.3	0.4	0.3	0.2	0.4	0.5	0.7	0.8	1.1
2003.....	3.3	2.6	0.5	0.2	0.3	0.3	0.5	0.6	0.9	1.2
2002.....	3.4	2.0	0.5	0.2	0.2	0.4	0.5	0.7	0.9	1.2
2001.....	3.3	2.6	0.5	0.2	0.2	0.4	0.5	0.6	1.1	1.2
2000.....	3.3	1.8	0.5	0.2	0.2	0.3	0.5	0.6	0.9	1.3
1999.....	3.2	2.5	0.5	0.2	0.2	0.4	0.6	0.6	1.0	1.2

All races 3/, male

2006.....	2.6	2.7	0.6	*	*	0.3	0.5	0.8	0.7	0.9
2005.....	2.8	2.7	0.5	*	0.2	0.3	0.5	0.7	0.7	1.3
2004.....	2.8	2.8	0.5	0.2	0.2	0.4	0.5	0.7	0.9	1.2
2003.....	2.9	2.9	0.5	*	0.3	0.3	0.6	0.6	0.9	1.2
2002.....	2.9	2.5	0.5	*	0.2	0.4	0.6	0.9	1.0	1.2
2001.....	2.8	2.9	0.6	0.2	0.3	0.4	0.5	0.7	1.2	1.3
2000.....	2.8	2.1	0.5	0.3	0.2	0.3	0.5	0.6	0.9	1.2
1999.....	2.8	3.0	0.6	0.2	0.3	0.4	0.7	0.7	1.1	1.4

All races 3/, female

2006.....	3.3	2.2	0.3	*	*	0.3	0.4	0.6	0.8	1.1
2005.....	3.6	1.8	0.4	*	*	0.3	0.5	0.6	0.7	1.2
2004.....	3.6	1.8	0.4	0.3	0.2	0.4	0.5	0.8	0.7	1.0
2003.....	3.7	2.3	0.4	*	0.3	0.3	0.5	0.7	0.9	1.2
2002.....	3.8	1.5	0.5	0.2	*	0.3	0.5	0.6	0.8	1.2
2001.....	3.9	2.4	0.3	0.2	*	0.4	0.5	0.5	1.0	1.1
2000.....	3.8	1.6	0.6	*	0.2	0.3	0.5	0.6	1.0	1.3
1999.....	3.7	1.9	0.4	0.3	*	0.4	0.6	0.5	1.0	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

White, both sexes

2006 4/.....	2.7	2.2	0.4	*	*	0.2	0.2	0.3	0.3	0.5
2005 4/.....	3.0	1.9	0.4	*	*	0.2	0.2	0.3	0.3	0.6
2004 4/.....	3.0	2.0	0.3	0.2	0.2	0.3	0.2	0.4	0.3	0.5
2003 4/.....	3.0	2.3	0.4	*	0.2	0.2	0.3	0.3	0.4	0.6
2002.....	3.1	1.9	0.4	0.2	*	0.2	0.3	0.3	0.5	0.6
2001.....	3.1	2.3	0.4	0.2	0.2	0.2	0.2	0.3	0.5	0.5
2000.....	3.1	1.7	0.4	0.1	0.2	0.2	0.2	0.2	0.4	0.6
1999.....	3.0	1.8	0.4	0.2	0.2	0.3	0.2	0.2	0.4	0.6

White, male

2006 4/.....	2.3	2.4	0.4	*	*	*	0.3	0.4	0.4	0.5
2005 4/.....	2.5	2.2	0.5	*	*	*	*	0.3	0.3	0.6
2004 4/.....	2.6	2.2	0.3	*	*	0.3	0.3	0.4	0.3	0.6
2003 4/.....	2.6	2.8	0.3	*	0.2	*	0.3	*	0.5	0.7
2002.....	2.6	2.5	0.5	*	*	0.3	0.3	0.4	0.6	0.7
2001.....	2.6	2.4	0.6	*	0.3	*	0.2	0.4	0.5	0.6
2000.....	2.5	1.7	0.4	*	*	*	*	*	0.4	0.6
1999.....	2.5	2.2	0.5	*	*	0.3	*	0.3	0.4	0.7

White, female

2006 4/.....	3.0	2.0	*	*	*	*	*	*	*	0.5
2005 4/.....	3.4	1.6	0.4	*	*	*	0.3	*	0.3	0.6
2004 4/.....	3.3	1.7	*	*	*	0.3	*	0.4	0.3	0.4
2003 4/.....	3.5	1.7	0.4	*	*	*	*	0.3	0.4	0.5
2002.....	3.6	*	0.4	*	*	*	0.3	0.3	0.4	0.5
2001.....	3.6	2.1	*	*	*	*	*	*	0.5	0.4
2000.....	3.6	1.7	0.4	*	*	*	*	*	0.3	0.6
1999.....	3.4	1.4	*	*	*	0.3	*	*	0.4	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

Black, both sexes

2006 4/.....	5.4	3.1	0.9	*	*	1.0	1.7	2.6	3.4	4.5
2005 4/.....	5.8	3.8	*	*	*	1.1	2.1	3.0	3.5	5.4
2004 4/.....	5.7	3.5	0.9	*	*	1.1	2.2	2.9	4.1	5.1
2003 4/.....	6.0	3.0	0.9	*	0.6	0.7	2.3	2.8	4.0	5.2
2002.....	6.0	*	0.8	*	*	0.9	2.1	3.2	3.8	5.5
2001.....	5.9	3.5	*	*	*	1.1	2.0	2.4	4.7	5.5
2000.....	5.8	*	1.2	0.6	0.6	0.7	2.1	3.0	4.7	5.7
1999.....	5.7	5.5	1.4	0.6	0.6	0.9	2.8	2.9	5.1	5.8

Black, male

2006 4/.....	4.8	*	*	*	*	*	1.7	2.8	2.8	4.1
2005 4/.....	5.3	*	*	*	*	1.2	2.2	3.2	3.6	6.1
2004 4/.....	5.3	*	*	*	*	1.4	2.0	3.0	4.7	5.6
2003 4/.....	5.6	*	*	*	*	*	2.2	2.7	4.1	5.3
2002.....	5.5	*	*	*	*	*	2.7	4.1	4.3	5.2
2001.....	5.2	*	*	*	*	1.3	2.2	2.9	5.6	5.9
2000.....	5.1	*	*	*	*	*	2.1	3.0	4.4	5.8
1999.....	5.3	6.5	*	*	*	*	3.3	3.3	5.4	6.4

Black, female

2006 4/.....	6.0	*	*	*	*	*	1.7	2.4	3.8	4.9
2005 4/.....	6.2	*	*	*	*	*	1.9	2.8	3.4	4.8
2004 4/.....	6.0	*	*	*	*	*	2.4	2.9	3.5	4.7
2003 4/.....	6.3	*	*	*	*	*	2.4	2.9	3.9	5.2
2002.....	6.4	*	*	*	*	*	1.6	2.3	3.3	5.7
2001.....	6.4	*	*	*	*	*	1.8	1.9	3.8	5.0
2000.....	6.4	*	*	*	*	*	2.2	3.0	5.0	5.6
1999.....	6.2	*	*	*	*	*	2.3	2.5	4.7	5.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Anemias (D50-D64)										
All races 3/, both sexes										
2006.....	1.3	*	0.2	*	*	0.2	0.3	0.4	0.3	0.4
2005.....	1.6	*	0.1	*	*	0.2	0.3	0.4	0.4	0.4
2004.....	1.6	*	0.2	0.1	0.1	0.2	0.4	0.4	0.4	0.4
2003.....	1.6	0.5	0.2	*	0.1	0.2	0.3	0.4	0.4	0.4
2002.....	1.6	*	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.5
2001.....	1.6	*	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.4
2000.....	1.6	*	0.2	0.1	0.1	0.1	0.3	0.3	0.4	0.4
1999.....	1.6	*	0.2	0.1	0.1	0.2	0.4	0.3	0.5	0.4
All races 3/, male										
2006.....	1.1	*	*	*	*	0.2	0.3	0.4	0.3	0.3
2005.....	1.2	*	*	*	*	0.2	0.3	0.4	0.4	0.4
2004.....	1.3	*	*	*	*	0.3	0.4	0.4	0.5	0.5
2003.....	1.3	*	*	*	*	*	0.3	0.4	0.4	0.3
2002.....	1.3	*	*	*	*	0.2	0.4	0.5	0.5	0.5
2001.....	1.3	*	*	*	*	0.2	0.3	0.4	0.5	0.4
2000.....	1.2	*	*	*	*	*	0.3	0.3	0.4	0.3
1999.....	1.3	*	*	*	*	0.2	0.5	0.3	0.5	0.4
All races 3/, female										
2006.....	1.6	*	*	*	*	*	0.2	0.3	0.4	0.5
2005.....	1.9	*	*	*	*	*	0.3	0.4	0.3	0.4
2004.....	1.8	*	*	*	*	*	0.4	0.5	0.3	0.3
2003.....	1.9	*	*	*	*	*	0.3	0.3	0.4	0.4
2002.....	1.9	*	*	*	*	*	0.3	0.3	0.5	0.4
2001.....	2.0	*	*	*	*	0.2	0.3	0.3	0.4	0.4
2000.....	2.0	*	*	*	*	*	0.3	0.3	0.5	0.4
1999.....	1.9	*	*	*	*	*	0.4	0.3	0.5	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Anemias (D50-D64)										
Black, both sexes										
2006 4/.....	2.3	*	*	*	*	0.6	1.3	1.9	2.1	2.2
2005 4/.....	2.6	*	*	*	*	0.9	1.5	2.3	2.4	2.2
2004 4/.....	2.7	*	*	*	*	0.9	2.0	2.2	2.4	2.4
2003 4/.....	2.7	*	*	*	*	*	1.9	1.9	2.2	2.2
2002.....	2.8	*	*	*	*	0.9	1.7	2.5	2.3	2.6
2001.....	2.7	*	*	*	*	0.9	1.5	1.9	2.5	2.1
2000.....	2.6	*	*	*	*	*	1.6	1.7	2.8	2.2
1999.....	2.8	*	0.8	*	*	0.7	2.2	2.1	3.0	2.2
Black, male										
2006 4/.....	2.1	*	*	*	*	*	1.5	2.1	2.0	1.9
2005 4/.....	2.4	*	*	*	*	*	1.7	2.4	2.7	2.7
2004 4/.....	2.6	*	*	*	*	1.2	1.9	2.3	3.0	3.0
2003 4/.....	2.7	*	*	*	*	*	1.8	2.2	2.3	2.2
2002.....	2.7	*	*	*	*	*	2.3	3.2	2.5	2.7
2001.....	2.5	*	*	*	*	*	1.6	2.5	3.3	2.2
2000.....	2.3	*	*	*	*	*	1.7	1.6	2.7	1.9
1999.....	2.7	*	*	*	*	*	2.8	2.3	3.3	2.5
Black, female										
2006 4/.....	2.6	*	*	*	*	*	*	1.8	2.1	2.4
2005 4/.....	2.8	*	*	*	*	*	1.3	2.1	2.2	1.8
2004 4/.....	2.8	*	*	*	*	*	2.1	2.2	1.9	1.9
2003 4/.....	2.8	*	*	*	*	*	2.1	1.7	2.2	2.2
2002.....	3.0	*	*	*	*	*	*	1.8	2.2	2.4
2001.....	2.8	*	*	*	*	*	1.5	*	1.8	2.1
2000.....	3.0	*	*	*	*	*	1.6	1.9	2.9	2.5
1999.....	3.0	*	*	*	*	*	1.7	1.8	2.8	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/ year 2/	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hemolytic anemias (D55-D59)										
All races 3/, both sexes										
2006.....	0.3	*	*	*	*	0.1	0.2	0.3	0.3	0.3
2005.....	0.3	*	*	*	*	0.1	0.2	0.3	0.3	0.3
2004.....	0.3	*	*	*	*	0.1	0.3	0.3	0.3	0.3
2003.....	0.3	*	*	*	*	*	0.3	0.3	0.3	0.3
2002.....	0.3	*	*	*	*	0.1	0.3	0.4	0.4	0.3
2001.....	0.3	*	*	*	*	0.1	0.2	0.3	0.4	0.3
2000.....	0.3	*	0.2	*	*	*	0.2	0.3	0.3	0.3
1999.....	0.3	*	0.1	*	*	0.1	0.3	0.3	0.4	0.3
All races 3/, male										
2006.....	0.2	*	*	*	*	*	0.3	0.3	0.2	0.2
2005.....	0.3	*	*	*	*	*	0.3	0.4	0.3	0.3
2004.....	0.3	*	*	*	*	0.2	0.3	0.3	0.4	0.3
2003.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.3
2002.....	0.3	*	*	*	*	*	0.3	0.4	0.4	0.4
2001.....	0.3	*	*	*	*	*	0.2	0.3	0.5	0.3
2000.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.2
1999.....	0.3	*	*	*	*	*	0.4	0.3	0.4	0.3
All races 3/, female										
2006.....	0.3	*	*	*	*	*	*	0.3	0.3	0.4
2005.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.3
2004.....	0.3	*	*	*	*	*	0.3	0.4	0.3	0.2
2003.....	0.3	*	*	*	*	*	0.3	0.3	0.3	0.3
2002.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.3
2001.....	0.3	*	*	*	*	*	*	0.2	0.3	0.3
2000.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.3
1999.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hemolytic anemias (D55-D59)										
Black, both sexes										
2006 4/.....	1.2	*	*	*	*	0.6	1.2	1.8	1.8	1.9
2005 4/.....	1.4	*	*	*	*	0.8	1.4	2.2	2.2	2.1
2004 4/.....	1.5	*	*	*	*	0.8	1.9	2.2	2.3	2.1
2003 4/.....	1.3	*	*	*	*	*	1.7	1.9	2.0	1.9
2002.....	1.4	*	*	*	*	0.7	1.6	2.2	2.3	2.3
2001.....	1.4	*	*	*	*	0.7	1.4	1.6	2.3	1.8
2000.....	1.4	*	*	*	*	*	1.5	1.6	2.4	2.0
1999.....	1.5	*	0.8	*	*	0.6	1.9	1.9	2.8	2.1
Black, male										
2006 4/.....	1.2	*	*	*	*	*	1.5	2.0	1.7	1.6
2005 4/.....	1.4	*	*	*	*	*	1.6	2.4	2.5	2.7
2004 4/.....	1.5	*	*	*	*	*	1.8	2.2	3.0	2.5
2003 4/.....	1.4	*	*	*	*	*	1.5	2.1	2.0	1.9
2002.....	1.4	*	*	*	*	*	2.0	2.9	2.4	2.5
2001.....	1.4	*	*	*	*	*	1.5	2.0	3.1	1.9
2000.....	1.3	*	*	*	*	*	1.5	1.6	2.1	1.9
1999.....	1.7	*	*	*	*	*	2.6	2.2	3.1	2.3
Black, female										
2006 4/.....	1.3	*	*	*	*	*	*	1.6	1.8	2.1
2005 4/.....	1.3	*	*	*	*	*	1.3	1.9	2.0	1.6
2004 4/.....	1.4	*	*	*	*	*	2.0	2.1	1.8	1.8
2003 4/.....	1.3	*	*	*	*	*	2.0	1.7	2.0	2.0
2002.....	1.4	*	*	*	*	*	*	1.7	2.2	2.1
2001.....	1.3	*	*	*	*	*	*	*	1.5	1.8
2000.....	1.4	*	*	*	*	*	1.5	1.6	2.6	2.0
1999.....	1.3	*	*	*	*	*	*	1.8	2.5	1.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IV. Endocrine, nutritional and metabolic diseases (E00-E88)										
All races 3/, both sexes										
2006.....	33.1	5.0	1.0	0.4	0.5	0.8	1.6	2.5	3.7	5.4
2005.....	34.8	5.5	0.9	0.5	0.7	1.0	1.7	2.3	3.7	5.8
2004.....	33.7	5.7	0.9	0.5	0.6	0.9	1.6	2.2	3.6	5.5
2003.....	34.5	6.2	1.2	0.5	0.6	1.0	1.6	2.6	3.6	5.5
2002.....	34.5	6.6	0.9	0.5	0.6	0.9	1.7	2.5	3.7	5.6
2001.....	34.0	7.2	1.2	0.5	0.6	1.0	1.5	2.3	3.2	5.3
2000.....	33.5	6.8	1.0	0.4	0.7	1.0	1.6	2.3	3.4	4.9
1999.....	33.3	6.9	1.0	0.5	0.6	1.1	1.6	2.2	3.1	4.9
All races 3/, male										
2006.....	32.3	6.0	0.8	0.4	0.5	0.9	1.7	2.6	4.2	6.4
2005.....	33.3	5.9	1.0	0.6	0.7	1.1	1.8	2.5	4.3	6.7
2004.....	32.2	7.1	1.0	0.5	0.6	0.9	1.7	2.4	3.9	6.4
2003.....	32.5	6.8	1.4	0.6	0.6	1.1	1.7	2.8	4.3	6.0
2002.....	31.9	7.4	0.8	0.6	0.7	0.9	1.9	2.7	4.4	6.3
2001.....	30.9	8.0	1.2	0.4	0.7	1.0	1.5	2.4	3.6	6.1
2000.....	30.2	7.4	1.1	0.4	0.7	1.0	1.7	2.6	4.1	5.6
1999.....	29.8	7.3	1.1	0.5	0.6	1.1	1.5	2.2	3.6	5.8
All races 3/, female										
2006.....	33.9	4.0	1.1	0.3	0.5	0.8	1.6	2.3	3.3	4.4
2005.....	36.2	5.0	0.9	0.4	0.7	0.9	1.7	2.1	3.1	4.9
2004.....	35.2	4.2	0.8	0.5	0.5	0.8	1.6	2.0	3.3	4.5
2003.....	36.5	5.5	1.1	0.3	0.5	0.9	1.5	2.4	2.9	4.9
2002.....	37.0	5.8	1.0	0.3	0.5	0.9	1.6	2.3	2.9	4.8
2001.....	37.0	6.4	1.2	0.5	0.5	1.0	1.6	2.2	2.8	4.5
2000.....	36.7	6.2	1.0	0.3	0.6	1.0	1.4	2.0	2.7	4.3
1999.....	36.6	6.5	0.9	0.5	0.6	1.1	1.7	2.1	2.7	4.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IV. Endocrine, nutritional and metabolic diseases (E00-E88)										
White, both sexes										
2006 4/.....	33.1	4.9	1.0	0.4	0.5	0.9	1.6	2.2	3.1	4.6
2005 4/.....	34.9	5.0	1.0	0.5	0.7	1.0	1.6	1.9	3.1	5.0
2004 4/.....	33.7	5.8	0.9	0.5	0.6	0.8	1.6	2.0	3.2	4.7
2003 4/.....	34.5	5.4	1.2	0.5	0.6	1.0	1.6	2.2	3.2	4.8
2002.....	34.4	6.2	1.0	0.5	0.6	0.9	1.6	2.3	3.2	5.0
2001.....	34.0	6.8	1.2	0.5	0.6	0.9	1.4	2.0	2.8	4.7
2000.....	33.5	6.3	1.1	0.4	0.7	1.0	1.5	1.9	3.0	4.4
1999.....	33.1	6.0	1.1	0.4	0.6	1.1	1.6	2.0	2.7	4.3
White, male										
2006 4/.....	32.6	5.9	0.9	0.5	0.5	0.9	1.6	2.3	3.6	5.3
2005 4/.....	33.7	5.4	1.1	0.6	0.7	1.0	1.7	2.1	3.6	5.7
2004 4/.....	32.5	7.2	1.0	0.5	0.6	0.8	1.7	2.2	3.4	5.5
2003 4/.....	32.9	6.3	1.4	0.7	0.7	1.1	1.7	2.3	3.9	5.5
2002.....	32.2	7.6	0.8	0.7	0.7	0.9	1.6	2.5	3.9	5.7
2001.....	31.2	7.2	1.3	0.5	0.7	1.0	1.4	2.0	3.2	5.3
2000.....	30.6	6.8	1.2	0.4	0.8	1.0	1.7	2.3	3.7	5.0
1999.....	29.9	6.3	1.2	0.5	0.6	1.2	1.4	2.1	3.0	5.1
White, female										
2006 4/.....	33.5	3.8	1.1	0.4	0.4	0.8	1.6	2.0	2.7	3.8
2005 4/.....	36.0	4.6	0.9	0.4	0.6	0.9	1.5	1.7	2.6	4.3
2004 4/.....	34.9	4.3	0.9	0.6	0.5	0.9	1.5	1.9	2.9	3.8
2003 4/.....	36.1	4.5	1.1	0.4	0.5	0.9	1.5	2.2	2.5	4.1
2002.....	36.6	4.7	1.1	0.3	0.5	1.0	1.5	2.1	2.6	4.2
2001.....	36.8	6.3	1.2	0.5	0.5	0.9	1.5	2.0	2.5	4.0
2000.....	36.3	5.7	1.0	0.4	0.6	1.0	1.4	1.6	2.3	3.8
1999.....	36.2	5.7	0.9	0.4	0.6	1.1	1.7	2.0	2.4	3.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IV. Endocrine, nutritional and metabolic diseases (E00-E88)										
Black, both sexes										
2006 4/.....	40.9	6.2	0.9	*	0.6	0.8	2.4	5.0	8.4	12.2
2005 4/.....	42.2	7.5	0.8	*	0.8	1.4	2.7	4.7	8.3	12.1
2004 4/.....	41.8	5.9	*	*	0.6	1.1	2.4	4.2	7.7	12.2
2003 4/.....	42.3	9.9	1.0	*	*	1.2	2.2	5.5	7.1	11.0
2002.....	42.6	8.3	0.8	*	0.7	1.0	2.9	4.9	7.5	10.7
2001.....	41.6	10.3	0.8	*	0.6	1.1	2.5	4.5	6.4	10.5
2000.....	41.4	8.9	0.8	*	*	1.1	2.5	5.3	6.8	9.6
1999.....	41.4	11.4	0.8	0.6	0.6	1.1	2.2	3.6	6.4	10.2
Black, male										
2006 4/.....	38.1	6.9	*	*	*	*	2.4	5.5	9.6	15.5
2005 4/.....	38.4	7.7	*	*	*	1.7	2.3	4.8	10.1	14.9
2004 4/.....	37.7	7.7	*	*	*	1.4	2.2	4.8	8.8	14.2
2003 4/.....	37.1	8.9	*	*	*	*	2.1	6.7	8.5	11.3
2002.....	36.6	7.0	*	*	*	1.2	3.5	5.5	9.3	12.0
2001.....	35.9	11.7	*	*	*	*	2.4	5.2	7.8	12.8
2000.....	34.6	9.9	*	*	*	*	2.7	5.5	8.0	10.7
1999.....	35.2	12.7	*	*	*	*	2.3	3.4	7.8	11.7
Black, female										
2006 4/.....	43.4	*	*	*	*	*	2.3	4.5	7.4	9.2
2005 4/.....	45.7	7.4	*	*	*	*	3.0	4.7	6.7	9.6
2004 4/.....	45.5	*	*	*	*	*	2.5	3.6	6.7	10.3
2003 4/.....	47.0	10.8	*	*	*	1.3	2.2	4.4	5.9	10.6
2002.....	48.0	9.7	*	*	*	*	2.4	4.3	5.8	9.6
2001.....	46.8	8.8	*	*	*	*	2.5	3.9	5.2	8.6
2000.....	47.6	7.9	*	*	*	*	2.2	5.1	5.7	8.6
1999.....	47.1	10.1	*	*	*	*	2.1	3.9	5.1	8.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Endocrine diseases (E00-E34)										
All races 3/, both sexes										
2006.....	25.1	0.8	*	*	0.1	0.2	0.7	1.3	2.3	3.5
2005.....	26.3	0.8	*	*	0.2	0.4	0.8	1.2	2.2	3.8
2004.....	25.8	0.6	0.1	*	0.2	0.3	0.7	1.1	2.2	3.5
2003.....	26.5	0.7	0.1	*	0.1	0.3	0.7	1.3	2.2	3.5
2002.....	26.4	1.0	*	*	0.1	0.3	0.7	1.3	2.1	3.7
2001.....	26.1	1.1	*	*	0.2	0.3	0.6	1.2	2.0	3.5
2000.....	25.6	1.2	0.1	*	0.1	0.3	0.6	1.2	2.2	3.2
1999.....	25.5	1.0	0.1	*	0.1	0.3	0.7	1.1	2.0	3.4
All races 3/, male										
2006.....	25.0	1.0	*	*	*	0.3	0.8	1.4	2.5	4.1
2005.....	25.6	*	*	*	*	0.4	0.8	1.3	2.6	4.5
2004.....	25.0	*	*	*	0.2	0.3	0.6	1.3	2.4	4.4
2003.....	25.3	*	*	*	*	0.3	0.8	1.4	2.6	4.0
2002.....	24.8	1.1	*	*	0.2	0.3	0.8	1.4	2.6	4.2
2001.....	24.1	1.3	*	*	0.2	0.3	0.7	1.2	2.3	4.1
2000.....	23.4	1.3	*	*	*	0.3	0.7	1.3	2.7	3.7
1999.....	23.3	*	*	*	*	0.3	0.7	1.1	2.3	4.0
All races 3/, female										
2006.....	25.3	*	*	*	*	0.2	0.6	1.3	2.0	2.9
2005.....	27.0	*	*	*	0.2	0.3	0.7	1.1	1.7	3.1
2004.....	26.7	*	*	*	*	0.2	0.7	0.9	1.9	2.7
2003.....	27.7	*	*	*	*	0.2	0.6	1.3	1.8	2.9
2002.....	28.0	*	*	*	*	0.3	0.6	1.2	1.7	3.1
2001.....	28.0	*	*	*	*	0.3	0.5	1.2	1.6	2.9
2000.....	27.7	1.1	*	*	*	0.2	0.6	1.1	1.7	2.8
1999.....	27.5	1.3	*	*	*	0.3	0.7	1.1	1.8	2.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Endocrine diseases (E00-E34)										
White, both sexes										
2006 4/.....	24.6	0.7	*	*	*	0.2	0.6	1.0	1.8	2.9
2005 4/.....	25.9	0.6	*	*	0.2	0.3	0.6	0.9	1.7	3.2
2004 4/.....	25.3	0.7	*	*	*	0.2	0.5	0.9	1.8	3.0
2003 4/.....	26.1	*	*	*	*	0.2	0.6	1.0	1.9	3.0
2002.....	26.0	1.1	*	*	*	0.3	0.6	1.1	1.9	3.2
2001.....	25.7	1.1	*	*	0.1	0.2	0.4	0.9	1.7	3.0
2000.....	25.1	1.0	*	*	*	0.3	0.5	0.9	1.9	2.8
1999.....	24.9	1.0	*	*	*	0.3	0.6	1.0	1.7	2.9
White, male										
2006 4/.....	24.8	*	*	*	*	0.2	0.6	1.0	1.9	3.4
2005 4/.....	25.5	*	*	*	*	0.3	0.7	1.0	2.0	3.7
2004 4/.....	24.9	*	*	*	*	*	0.6	1.0	2.0	3.7
2003 4/.....	25.3	*	*	*	*	0.3	0.7	0.9	2.3	3.4
2002.....	24.8	1.3	*	*	*	0.3	0.6	1.2	2.2	3.7
2001.....	24.1	*	*	*	*	0.3	0.5	0.9	1.9	3.5
2000.....	23.5	*	*	*	*	0.3	0.6	1.0	2.4	3.2
1999.....	23.2	*	*	*	*	0.3	0.6	1.0	1.9	3.5
White, female										
2006 4/.....	24.4	*	*	*	*	*	0.5	0.9	1.6	2.5
2005 4/.....	26.3	*	*	*	*	0.3	0.5	0.7	1.3	2.7
2004 4/.....	25.8	*	*	*	*	*	0.5	0.8	1.6	2.3
2003 4/.....	26.8	*	*	*	*	*	0.5	1.1	1.5	2.5
2002.....	27.1	*	*	*	*	0.3	0.5	1.0	1.5	2.7
2001.....	27.2	*	*	*	*	*	0.4	0.9	1.5	2.5
2000.....	26.8	*	*	*	*	*	0.4	0.8	1.3	2.5
1999.....	26.5	1.4	*	*	*	0.3	0.5	1.0	1.5	2.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Endocrine diseases (E00-E34)										
Black, both sexes										
2006 4/.....	33.4	*	*	*	*	*	1.5	3.6	5.6	8.3
2005 4/.....	34.2	*	*	*	*	0.8	1.7	3.1	5.5	8.6
2004 4/.....	34.2	*	*	*	*	0.7	1.5	2.3	4.9	8.0
2003 4/.....	34.7	*	*	*	*	*	1.3	3.5	4.7	7.5
2002.....	34.5	*	*	*	*	*	1.6	3.1	4.6	7.6
2001.....	34.0	*	*	*	*	0.6	1.4	3.4	4.4	7.3
2000.....	33.8	*	*	*	*	*	1.5	3.3	4.5	6.7
1999.....	34.0	*	*	*	*	*	1.2	2.1	4.4	7.1
Black, male										
2006 4/.....	31.4	*	*	*	*	*	1.7	3.7	7.0	10.3
2005 4/.....	31.5	*	*	*	*	*	1.5	3.2	6.9	11.3
2004 4/.....	31.0	*	*	*	*	*	1.3	2.8	5.8	9.9
2003 4/.....	30.5	*	*	*	*	*	1.4	4.4	5.4	8.7
2002.....	29.5	*	*	*	*	*	1.8	3.3	6.0	8.7
2001.....	29.2	*	*	*	*	*	1.5	4.0	5.8	9.3
2000.....	28.1	*	*	*	*	*	1.5	3.5	5.0	7.6
1999.....	28.5	*	*	*	*	*	*	1.8	5.2	8.2
Black, female										
2006 4/.....	35.1	*	*	*	*	*	1.3	3.4	4.4	6.5
2005 4/.....	36.7	*	*	*	*	*	1.8	3.0	4.2	6.2
2004 4/.....	37.2	*	*	*	*	*	1.8	1.8	4.2	6.2
2003 4/.....	38.6	*	*	*	*	*	*	2.7	4.1	6.5
2002.....	39.0	*	*	*	*	*	1.4	2.8	3.4	6.6
2001.....	38.3	*	*	*	*	*	1.3	2.8	3.1	5.5
2000.....	39.1	*	*	*	*	*	1.5	3.2	4.1	5.8
1999.....	39.0	*	*	*	*	*	*	2.5	3.8	6.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
All races 3/, both sexes										
2006.....	24.2	*	*	*	0.1	0.2	0.6	1.2	2.1	3.3
2005.....	25.3	*	*	*	0.2	0.3	0.6	1.0	2.0	3.6
2004.....	24.9	*	*	*	0.1	0.2	0.5	0.9	2.0	3.4
2003.....	25.5	*	*	*	0.1	0.2	0.5	1.2	2.1	3.3
2002.....	25.4	*	*	*	0.1	0.3	0.6	1.2	2.0	3.5
2001.....	25.1	*	*	*	0.2	0.2	0.5	1.1	1.8	3.3
2000.....	24.6	*	*	*	*	0.2	0.6	1.1	2.0	3.0
1999.....	24.5	*	*	*	0.1	0.3	0.6	1.0	1.9	3.2
All races 3/, male										
2006.....	24.4	*	*	*	*	0.2	0.7	1.3	2.4	4.0
2005.....	25.0	*	*	*	*	0.4	0.7	1.1	2.5	4.3
2004.....	24.4	*	*	*	*	0.3	0.5	1.0	2.3	4.2
2003.....	24.8	*	*	*	*	0.3	0.6	1.2	2.5	3.9
2002.....	24.2	*	*	*	*	0.3	0.7	1.3	2.4	4.1
2001.....	23.5	*	*	*	0.2	0.3	0.6	1.2	2.2	3.9
2000.....	22.9	*	*	*	*	0.3	0.7	1.2	2.5	3.5
1999.....	22.8	*	*	*	*	0.2	0.6	1.0	2.2	3.8
All races 3/, female										
2006.....	24.0	*	*	*	*	*	0.5	1.1	1.9	2.7
2005.....	25.7	*	*	*	0.2	0.3	0.6	0.9	1.5	2.9
2004.....	25.4	*	*	*	*	*	0.6	0.8	1.8	2.5
2003.....	26.2	*	*	*	*	*	0.5	1.1	1.7	2.8
2002.....	26.5	*	*	*	*	0.3	0.5	1.1	1.5	3.0
2001.....	26.6	*	*	*	*	0.2	0.5	1.1	1.5	2.7
2000.....	26.3	*	*	*	*	*	0.5	0.9	1.6	2.5
1999.....	26.2	*	*	*	*	0.3	0.5	1.0	1.6	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
White, both sexes										
2006 4/.....	23.6	*	*	*	*	0.2	0.5	0.9	1.7	2.8
2005 4/.....	24.9	*	*	*	0.1	0.2	0.5	0.8	1.6	3.0
2004 4/.....	24.4	*	*	*	*	0.2	0.4	0.8	1.7	2.9
2003 4/.....	25.0	*	*	*	*	0.2	0.5	0.9	1.8	2.8
2002.....	24.9	*	*	*	*	0.2	0.5	1.0	1.7	3.1
2001.....	24.6	*	*	*	*	0.2	0.4	0.8	1.6	2.8
2000.....	24.1	*	*	*	*	0.2	0.5	0.8	1.8	2.7
1999.....	23.9	*	*	*	*	0.2	0.5	0.9	1.6	2.8
White, male										
2006 4/.....	24.2	*	*	*	*	*	0.5	0.9	1.8	3.3
2005 4/.....	24.9	*	*	*	*	0.3	0.6	0.9	1.9	3.5
2004 4/.....	24.3	*	*	*	*	*	0.4	0.8	2.0	3.6
2003 4/.....	24.8	*	*	*	*	*	0.6	0.8	2.2	3.4
2002.....	24.2	*	*	*	*	*	0.5	1.1	2.1	3.7
2001.....	23.5	*	*	*	*	*	0.4	0.8	1.8	3.3
2000.....	22.9	*	*	*	*	0.3	0.5	1.0	2.3	3.1
1999.....	22.7	*	*	*	*	0.3	0.5	0.9	1.8	3.4
White, female										
2006 4/.....	23.0	*	*	*	*	*	0.4	0.8	1.6	2.3
2005 4/.....	24.9	*	*	*	*	*	0.4	0.7	1.2	2.5
2004 4/.....	24.5	*	*	*	*	*	0.4	0.7	1.5	2.1
2003 4/.....	25.2	*	*	*	*	*	0.4	1.0	1.4	2.3
2002.....	25.6	*	*	*	*	*	0.4	0.9	1.3	2.5
2001.....	25.7	*	*	*	*	*	0.3	0.8	1.4	2.3
2000.....	25.3	*	*	*	*	*	0.4	0.7	1.2	2.3
1999.....	25.0	*	*	*	*	*	0.4	0.9	1.4	2.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diabetes mellitus (E10-E14)										
Black, both sexes										
2006 4/.....	32.4	*	*	*	*	*	1.3	3.4	5.2	7.8
2005 4/.....	33.2	*	*	*	*	0.7	1.5	2.8	5.1	8.1
2004 4/.....	33.2	*	*	*	*	0.6	1.3	2.0	4.6	7.5
2003 4/.....	33.8	*	*	*	*	*	1.0	3.2	4.4	7.3
2002.....	33.6	*	*	*	*	*	1.4	2.8	4.3	7.3
2001.....	33.1	*	*	*	*	*	1.4	3.2	4.2	6.9
2000.....	32.8	*	*	*	*	*	1.5	2.8	4.1	6.0
1999.....	33.0	*	*	*	*	*	1.1	1.8	4.1	6.6
Black, male										
2006 4/.....	30.6	*	*	*	*	*	1.5	3.6	6.7	9.8
2005 4/.....	30.7	*	*	*	*	*	1.4	2.9	6.7	10.8
2004 4/.....	30.2	*	*	*	*	*	*	2.5	5.4	9.5
2003 4/.....	29.8	*	*	*	*	*	*	4.1	5.2	8.4
2002.....	29.0	*	*	*	*	*	1.7	3.0	5.7	8.5
2001.....	28.5	*	*	*	*	*	1.5	3.9	5.6	8.7
2000.....	27.4	*	*	*	*	*	1.5	3.1	4.5	7.1
1999.....	27.7	*	*	*	*	*	*	1.6	5.0	7.7
Black, female										
2006 4/.....	34.1	*	*	*	*	*	*	3.1	3.9	6.0
2005 4/.....	35.5	*	*	*	*	*	1.7	2.6	3.6	5.7
2004 4/.....	36.0	*	*	*	*	*	1.6	1.6	3.8	5.6
2003 4/.....	37.4	*	*	*	*	*	*	2.3	3.7	6.3
2002.....	37.8	*	*	*	*	*	*	2.6	3.1	6.3
2001.....	37.2	*	*	*	*	*	1.3	2.7	2.9	5.3
2000.....	37.8	*	*	*	*	*	1.5	2.5	3.8	5.0
1999.....	37.8	*	*	*	*	*	*	2.0	3.3	5.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nutritional diseases (E40-E68)										
All races 3/, both sexes										
2006.....	2.4	*	*	*	*	*	0.3	0.6	0.9	1.3
2005.....	2.6	*	*	*	*	0.1	0.3	0.6	1.0	1.3
2004.....	2.5	*	*	*	*	*	0.2	0.6	0.9	1.2
2003.....	2.5	*	*	*	*	0.1	0.3	0.6	0.8	1.3
2002.....	2.6	*	*	*	*	*	0.3	0.6	0.9	1.2
2001.....	2.4	*	*	*	*	*	0.3	0.4	0.7	1.2
2000.....	2.6	*	*	*	*	*	0.2	0.5	0.7	1.1
1999.....	2.5	*	*	*	*	0.1	0.2	0.4	0.6	0.9
All races 3/, male										
2006.....	2.1	*	*	*	*	*	0.3	0.7	1.0	1.4
2005.....	2.3	*	*	*	*	*	0.3	0.7	1.2	1.3
2004.....	2.1	*	*	*	*	*	0.3	0.6	0.9	1.3
2003.....	2.1	*	*	*	*	*	0.3	0.8	1.1	1.2
2002.....	2.2	*	*	*	*	*	0.4	0.6	1.1	1.4
2001.....	1.9	*	*	*	*	*	0.3	0.6	0.8	1.2
2000.....	2.0	*	*	*	*	*	0.2	0.6	0.9	1.2
1999.....	2.0	*	*	*	*	*	*	0.4	0.8	1.1
All races 3/, female										
2006.....	2.7	*	*	*	*	*	0.3	0.4	0.8	1.1
2005.....	3.0	*	*	*	*	*	0.3	0.5	0.9	1.3
2004.....	2.8	*	*	*	*	*	*	0.6	0.9	1.2
2003.....	2.9	*	*	*	*	*	0.2	0.5	0.6	1.3
2002.....	3.0	*	*	*	*	*	0.3	0.5	0.8	1.1
2001.....	2.9	*	*	*	*	*	0.3	0.3	0.7	1.2
2000.....	3.1	*	*	*	*	*	*	0.4	0.5	0.9
1999.....	2.9	*	*	*	*	*	0.3	0.4	0.4	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Nutritional diseases (E40-E68)										
White, both sexes										
2006 4/.....	2.5	*	*	*	*	*	0.2	0.5	0.8	1.0
2005 4/.....	2.6	*	*	*	*	*	0.2	0.5	0.8	1.1
2004 4/.....	2.4	*	*	*	*	*	0.2	0.5	0.7	1.0
2003 4/.....	2.6	*	*	*	*	*	0.2	0.5	0.7	1.1
2002.....	2.6	*	*	*	*	*	0.2	0.5	0.7	1.1
2001.....	2.4	*	*	*	*	*	0.2	0.4	0.6	1.0
2000.....	2.6	*	*	*	*	*	0.1	0.3	0.5	0.9
1999.....	2.5	*	*	*	*	*	0.2	0.3	0.5	0.8
White, male										
2006 4/.....	2.2	*	*	*	*	*	0.3	0.6	0.9	1.1
2005 4/.....	2.3	*	*	*	*	*	0.2	0.6	1.0	1.2
2004 4/.....	2.1	*	*	*	*	*	*	0.5	0.7	1.1
2003 4/.....	2.2	*	*	*	*	*	0.3	0.6	1.0	1.2
2002.....	2.2	*	*	*	*	*	0.3	0.5	0.8	1.3
2001.....	1.9	*	*	*	*	*	*	0.5	0.7	1.0
2000.....	2.1	*	*	*	*	*	*	0.4	0.7	1.2
1999.....	2.0	*	*	*	*	*	*	0.3	0.7	0.9
White, female										
2006 4/.....	2.8	*	*	*	*	*	*	0.4	0.6	1.0
2005 4/.....	3.0	*	*	*	*	*	*	0.4	0.7	1.1
2004 4/.....	2.8	*	*	*	*	*	*	0.5	0.7	0.8
2003 4/.....	3.0	*	*	*	*	*	*	0.4	0.4	1.0
2002.....	3.0	*	*	*	*	*	*	0.4	0.6	0.9
2001.....	2.9	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	3.1	*	*	*	*	*	*	*	0.3	0.7
1999.....	3.0	*	*	*	*	*	*	0.3	0.3	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Obesity and other hyperalimentation (E65-E68)

All races 3/, both sexes

2006.....	1.6	*	*	*	*	*	0.3	0.5	0.9	1.2
2005.....	1.6	*	*	*	*	*	0.3	0.6	1.0	1.3
2004.....	1.4	*	*	*	*	*	0.2	0.6	0.9	1.2
2003.....	1.4	*	*	*	*	0.1	0.2	0.6	0.8	1.2
2002.....	1.3	*	*	*	*	*	0.3	0.5	0.9	1.1
2001.....	1.1	*	*	*	*	*	0.3	0.4	0.7	1.1
2000.....	1.1	*	*	*	*	*	0.2	0.4	0.6	0.9
1999.....	0.9	*	*	*	*	*	0.2	0.4	0.6	0.8

All races 3/, male

2006.....	1.5	*	*	*	*	*	0.3	0.7	0.9	1.4
2005.....	1.5	*	*	*	*	*	0.3	0.7	1.1	1.3
2004.....	1.4	*	*	*	*	*	0.2	0.6	0.9	1.2
2003.....	1.3	*	*	*	*	*	0.2	0.7	1.0	1.2
2002.....	1.2	*	*	*	*	*	0.3	0.5	1.1	1.3
2001.....	1.0	*	*	*	*	*	0.2	0.5	0.7	1.1
2000.....	1.0	*	*	*	*	*	0.2	0.5	0.8	1.1
1999.....	0.9	*	*	*	*	*	*	0.4	0.8	1.0

All races 3/, female

2006.....	1.7	*	*	*	*	*	0.3	0.4	0.8	1.1
2005.....	1.6	*	*	*	*	*	0.3	0.4	0.8	1.2
2004.....	1.5	*	*	*	*	*	*	0.6	0.9	1.1
2003.....	1.5	*	*	*	*	*	0.2	0.5	0.6	1.2
2002.....	1.3	*	*	*	*	*	0.2	0.5	0.7	0.9
2001.....	1.2	*	*	*	*	*	0.3	0.3	0.6	1.1
2000.....	1.1	*	*	*	*	*	*	0.3	0.4	0.7
1999.....	1.0	*	*	*	*	*	0.2	0.4	0.4	0.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Obesity and other hyperalimentation (E65-E68)										
White, both sexes										
2006 4/.....	1.6	*	*	*	*	*	0.2	0.5	0.7	1.0
2005 4/.....	1.5	*	*	*	*	*	0.2	0.4	0.8	1.1
2004 4/.....	1.4	*	*	*	*	*	0.1	0.5	0.7	0.9
2003 4/.....	1.4	*	*	*	*	*	0.2	0.5	0.7	1.0
2002.....	1.2	*	*	*	*	*	0.2	0.4	0.7	1.0
2001.....	1.0	*	*	*	*	*	0.2	0.4	0.6	0.9
2000.....	1.0	*	*	*	*	*	*	0.3	0.5	0.8
1999.....	0.9	*	*	*	*	*	0.1	0.3	0.5	0.7
White, male										
2006 4/.....	1.5	*	*	*	*	*	0.3	0.6	0.9	1.1
2005 4/.....	1.6	*	*	*	*	*	0.2	0.5	0.9	1.1
2004 4/.....	1.4	*	*	*	*	*	*	0.5	0.7	1.1
2003 4/.....	1.4	*	*	*	*	*	*	0.6	0.9	1.1
2002.....	1.2	*	*	*	*	*	*	0.4	0.8	1.2
2001.....	1.0	*	*	*	*	*	*	0.5	0.7	1.0
2000.....	1.0	*	*	*	*	*	*	0.4	0.7	1.1
1999.....	0.9	*	*	*	*	*	*	0.3	0.6	0.8
White, female										
2006 4/.....	1.6	*	*	*	*	*	*	0.3	0.6	0.9
2005 4/.....	1.5	*	*	*	*	*	*	0.3	0.7	1.0
2004 4/.....	1.4	*	*	*	*	*	*	0.4	0.7	0.8
2003 4/.....	1.4	*	*	*	*	*	*	0.4	0.4	1.0
2002.....	1.2	*	*	*	*	*	*	0.4	0.5	0.8
2001.....	1.1	*	*	*	*	*	*	*	0.5	0.9
2000.....	1.0	*	*	*	*	*	*	*	0.3	0.6
1999.....	0.9	*	*	*	*	*	*	*	0.3	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Metabolic disorders (E70-E88)										
All races 3/, both sexes										
2006.....	5.6	4.0	0.9	0.3	0.3	0.5	0.6	0.6	0.6	0.6
2005.....	5.8	4.6	0.9	0.4	0.5	0.5	0.7	0.5	0.5	0.6
2004.....	5.4	4.9	0.8	0.4	0.4	0.5	0.8	0.5	0.6	0.7
2003.....	5.5	5.2	1.1	0.4	0.4	0.6	0.7	0.7	0.5	0.7
2002.....	5.5	5.5	0.8	0.4	0.4	0.5	0.7	0.6	0.6	0.6
2001.....	5.5	6.0	1.0	0.4	0.4	0.6	0.7	0.6	0.5	0.7
2000.....	5.4	5.3	0.9	0.3	0.5	0.6	0.8	0.6	0.6	0.7
1999.....	5.3	5.6	0.8	0.4	0.5	0.7	0.7	0.6	0.5	0.6
All races 3/, male										
2006.....	5.2	4.6	0.8	0.3	0.3	0.5	0.6	0.6	0.7	0.8
2005.....	5.4	5.0	1.0	0.5	0.5	0.5	0.6	0.5	0.5	0.8
2004.....	5.1	6.0	0.8	0.4	0.4	0.5	0.8	0.5	0.7	0.7
2003.....	5.1	5.8	1.1	0.5	0.5	0.7	0.7	0.7	0.6	0.8
2002.....	4.9	6.1	0.8	0.5	0.5	0.5	0.7	0.7	0.7	0.7
2001.....	4.9	6.5	1.1	0.4	0.4	0.6	0.6	0.6	0.6	0.8
2000.....	4.8	5.5	0.9	0.3	0.6	0.6	0.8	0.8	0.6	0.7
1999.....	4.5	6.3	0.9	0.4	0.5	0.7	0.7	0.7	0.5	0.7
All races 3/, female										
2006.....	5.9	3.5	1.0	0.3	0.3	0.5	0.7	0.7	0.5	0.4
2005.....	6.2	4.2	0.8	0.3	0.4	0.5	0.7	0.6	0.5	0.4
2004.....	5.8	3.8	0.7	0.4	0.3	0.5	0.8	0.5	0.5	0.7
2003.....	5.9	4.5	1.0	0.3	0.3	0.6	0.7	0.7	0.4	0.7
2002.....	6.0	4.9	0.9	0.3	0.4	0.6	0.7	0.6	0.5	0.5
2001.....	6.1	5.5	1.0	0.4	0.4	0.6	0.7	0.6	0.5	0.5
2000.....	6.0	5.0	0.8	0.3	0.4	0.7	0.7	0.5	0.5	0.6
1999.....	6.1	4.9	0.7	0.4	0.4	0.7	0.8	0.6	0.5	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Metabolic disorders (E70-E88)										
White, both sexes										
2006 4/.....	6.0	4.0	0.9	0.3	0.4	0.6	0.8	0.7	0.6	0.6
2005 4/.....	6.3	4.3	0.9	0.4	0.5	0.6	0.7	0.6	0.6	0.7
2004 4/.....	5.9	5.0	0.8	0.4	0.4	0.6	0.9	0.6	0.6	0.7
2003 4/.....	5.9	4.6	1.1	0.4	0.5	0.7	0.8	0.7	0.6	0.7
2002.....	5.9	5.0	0.9	0.4	0.5	0.6	0.8	0.7	0.7	0.7
2001.....	5.9	5.6	1.1	0.4	0.4	0.7	0.8	0.7	0.6	0.7
2000.....	5.8	5.0	0.9	0.3	0.6	0.7	0.9	0.7	0.6	0.7
1999.....	5.7	4.9	0.9	0.4	0.5	0.8	0.9	0.8	0.5	0.6
White, male										
2006 4/.....	5.7	4.6	0.8	0.4	0.4	0.6	0.7	0.7	0.7	0.8
2005 4/.....	5.8	4.6	1.0	0.6	0.5	0.6	0.7	0.6	0.6	0.9
2004 4/.....	5.5	6.2	0.9	0.4	0.5	0.5	0.9	0.6	0.7	0.7
2003 4/.....	5.4	5.6	1.1	0.5	0.6	0.7	0.8	0.7	0.6	0.8
2002.....	5.2	6.2	0.8	0.6	0.5	0.6	0.8	0.8	0.8	0.7
2001.....	5.2	5.9	1.1	0.4	0.4	0.7	0.7	0.7	0.6	0.8
2000.....	5.1	5.2	1.0	0.4	0.7	0.6	0.9	0.8	0.6	0.7
1999.....	4.7	5.6	1.0	0.4	0.5	0.7	0.7	0.8	0.5	0.7
White, female										
2006 4/.....	6.3	3.4	1.0	0.3	0.4	0.6	0.9	0.8	0.5	0.4
2005 4/.....	6.7	4.0	0.8	0.3	0.4	0.6	0.8	0.6	0.6	0.5
2004 4/.....	6.3	3.7	0.8	0.5	0.4	0.6	0.9	0.6	0.6	0.7
2003 4/.....	6.3	3.7	1.0	0.3	0.3	0.6	0.8	0.7	0.5	0.6
2002.....	6.5	3.9	1.0	0.3	0.5	0.7	0.8	0.7	0.5	0.6
2001.....	6.6	5.3	1.1	0.5	0.4	0.8	0.9	0.7	0.5	0.6
2000.....	6.4	4.8	0.8	0.3	0.5	0.8	0.8	0.5	0.6	0.7
1999.....	6.7	4.1	0.8	0.4	0.5	0.8	1.0	0.7	0.6	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cystic fibrosis (E84)										
All races 3/, both sexes										
2006.....	0.1	*	*	*	0.1	0.3	0.4	0.4	0.3	0.2
2005.....	0.1	*	*	*	0.1	0.3	0.4	0.3	0.2	0.2
2004.....	0.2	*	*	*	0.1	0.3	0.5	0.3	0.3	0.2
2003.....	0.2	*	*	*	0.1	0.3	0.4	0.4	0.2	0.2
2002.....	0.2	*	*	*	0.1	0.3	0.5	0.4	0.3	0.2
2001.....	0.2	*	*	*	0.1	0.4	0.5	0.4	0.2	0.2
2000.....	0.2	*	*	*	0.2	0.4	0.5	0.4	0.3	0.3
1999.....	0.2	*	*	*	0.2	0.4	0.5	0.4	0.3	0.2
All races 3/, male										
2006.....	0.1	*	*	*	*	0.3	0.3	0.3	0.3	*
2005.....	0.1	*	*	*	*	0.2	0.4	0.3	0.2	0.2
2004.....	0.2	*	*	*	*	0.3	0.5	0.3	0.3	0.2
2003.....	0.2	*	*	*	*	0.3	0.4	0.4	0.3	0.2
2002.....	0.2	*	*	*	*	0.2	0.4	0.4	0.4	0.2
2001.....	0.2	*	*	*	*	0.3	0.4	0.4	0.3	0.2
2000.....	0.2	*	*	*	*	0.3	0.5	0.4	0.3	0.2
1999.....	0.2	*	*	*	*	0.3	0.5	0.4	0.3	*
All races 3/, female										
2006.....	0.1	*	*	*	*	0.3	0.5	0.5	0.2	*
2005.....	0.1	*	*	*	*	0.3	0.5	0.3	*	*
2004.....	0.2	*	*	*	*	0.3	0.6	0.3	0.3	0.2
2003.....	0.2	*	*	*	*	0.3	0.5	0.4	*	*
2002.....	0.2	*	*	*	*	0.4	0.5	0.3	0.2	0.2
2001.....	0.2	*	*	*	*	0.4	0.6	0.4	0.2	*
2000.....	0.2	*	*	*	0.2	0.4	0.5	0.3	0.4	0.3
1999.....	0.2	*	*	*	*	0.5	0.6	0.4	0.2	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cystic fibrosis (E84)										
White, both sexes										
2006 4/.....	0.2	*	*	*	0.2	0.3	0.5	0.4	0.3	0.2
2005 4/.....	0.2	*	*	*	*	0.3	0.5	0.4	0.2	0.2
2004 4/.....	0.2	*	*	*	0.2	0.3	0.6	0.4	0.3	0.3
2003 4/.....	0.2	*	*	*	*	0.4	0.5	0.5	0.3	0.2
2002.....	0.2	*	*	*	0.2	0.4	0.5	0.5	0.4	0.2
2001.....	0.2	*	*	*	0.2	0.5	0.6	0.5	0.3	0.2
2000.....	0.2	*	*	*	0.2	0.4	0.6	0.4	0.4	0.3
1999.....	0.2	*	*	*	0.2	0.5	0.7	0.5	0.3	0.2
White, male										
2006 4/.....	0.2	*	*	*	*	0.3	0.4	0.3	0.4	*
2005 4/.....	0.2	*	*	*	*	0.3	0.4	0.4	0.3	0.2
2004 4/.....	0.2	*	*	*	*	0.3	0.6	0.4	0.4	0.2
2003 4/.....	0.2	*	*	*	*	0.3	0.4	0.5	0.3	0.2
2002.....	0.2	*	*	*	*	0.3	0.5	0.5	0.5	0.3
2001.....	0.2	*	*	*	*	0.4	0.5	0.4	0.3	0.3
2000.....	0.2	*	*	*	*	0.3	0.6	0.5	0.3	0.3
1999.....	0.2	*	*	*	*	0.4	0.6	0.5	0.3	*
White, female										
2006 4/.....	0.2	*	*	*	*	0.4	0.7	0.5	0.3	*
2005 4/.....	0.2	*	*	*	*	0.4	0.6	0.4	*	*
2004 4/.....	0.2	*	*	*	*	0.4	0.7	0.4	0.3	0.3
2003 4/.....	0.2	*	*	*	*	0.4	0.6	0.5	*	*
2002.....	0.2	*	*	*	*	0.5	0.6	0.4	0.3	0.2
2001.....	0.2	*	*	*	*	0.6	0.7	0.5	0.3	*
2000.....	0.2	*	*	*	0.3	0.5	0.6	0.4	0.5	0.3
1999.....	0.2	*	*	*	*	0.6	0.7	0.5	0.3	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other metabolic disorders (E70-E83,E85,E88)										
All races 3/, both sexes										
2006.....	3.4	2.5	0.7	0.2	0.2	0.2	0.2	0.2	0.2	0.3
2005.....	3.4	2.9	0.7	0.3	0.3	0.2	0.2	0.1	0.2	0.3
2004.....	3.0	3.1	0.7	0.3	0.2	0.2	0.2	0.1	0.2	0.3
2003.....	2.9	3.1	0.8	0.3	0.3	0.3	0.2	0.2	0.2	0.3
2002.....	2.8	3.4	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.3
2001.....	2.6	3.7	0.8	0.3	0.2	0.2	0.1	0.1	0.1	0.3
2000.....	2.4	3.2	0.8	0.2	0.3	0.2	0.2	0.2	0.2	0.2
1999.....	2.2	3.4	0.7	0.3	0.3	0.2	0.1	0.1	0.2	0.3
All races 3/, male										
2006.....	3.6	2.6	0.6	0.2	0.2	0.2	0.2	0.2	0.3	0.5
2005.....	3.6	3.3	0.7	0.4	0.4	0.2	0.2	*	0.2	0.5
2004.....	3.3	3.8	0.7	0.3	0.3	*	0.2	*	0.2	0.4
2003.....	3.2	3.5	1.0	0.4	0.4	0.3	0.2	*	*	0.4
2002.....	3.0	3.6	0.7	0.4	0.4	0.3	0.3	0.2	0.2	0.4
2001.....	2.9	4.1	0.9	0.3	0.3	0.3	*	*	*	0.4
2000.....	2.6	3.6	0.8	0.3	0.4	0.2	0.3	0.3	0.3	0.3
1999.....	2.4	3.7	0.8	0.3	0.3	0.3	*	*	*	0.3
All races 3/, female										
2006.....	3.1	2.4	0.9	0.2	*	*	*	*	*	*
2005.....	3.1	2.6	0.6	0.2	0.3	*	*	*	*	*
2004.....	2.8	2.4	0.6	0.3	*	*	*	*	*	0.2
2003.....	2.6	2.8	0.7	0.2	*	0.2	*	*	*	0.2
2002.....	2.5	3.2	0.7	*	0.2	*	*	*	*	*
2001.....	2.3	3.2	0.7	0.3	*	*	*	*	*	0.2
2000.....	2.3	2.8	0.7	0.2	*	*	*	*	*	0.2
1999.....	2.1	3.1	0.6	0.3	0.2	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other metabolic disorders (E70-E83,E85,E88)										
White, both sexes										
2006 4/.....	3.7	2.5	0.8	0.2	0.2	0.2	0.2	0.2	0.2	0.3
2005 4/.....	3.7	3.1	0.8	0.4	0.3	0.2	0.2	0.1	0.2	0.3
2004 4/.....	3.3	3.4	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.3
2003 4/.....	3.2	3.1	0.9	0.4	0.3	0.3	0.2	0.1	0.2	0.3
2002.....	3.0	3.3	0.7	0.3	0.3	0.2	0.2	0.2	0.2	0.3
2001.....	2.8	3.7	0.8	0.4	0.3	0.2	0.1	0.1	0.1	0.3
2000.....	2.7	3.4	0.8	0.3	0.3	0.2	0.2	0.2	0.1	0.3
1999.....	2.4	3.1	0.8	0.3	0.3	0.2	0.1	0.1	0.1	0.2
White, male										
2006 4/.....	4.0	2.6	0.6	*	*	0.3	0.2	*	0.3	0.5
2005 4/.....	4.0	3.5	0.8	0.5	0.4	0.2	*	*	*	0.5
2004 4/.....	3.6	4.2	0.8	0.4	0.3	*	0.3	*	*	0.4
2003 4/.....	3.5	3.8	1.0	0.5	0.5	0.3	0.2	*	*	0.4
2002.....	3.3	4.0	0.7	0.4	0.4	0.3	0.3	*	*	0.4
2001.....	3.1	4.0	0.9	0.4	0.3	0.3	*	*	*	0.4
2000.....	2.9	3.8	0.9	0.3	0.4	*	*	*	*	0.3
1999.....	2.6	3.5	0.9	0.3	0.3	0.3	*	*	*	0.4
White, female										
2006 4/.....	3.3	2.4	0.9	*	*	*	*	*	*	*
2005 4/.....	3.4	2.6	0.7	*	0.3	*	*	*	*	*
2004 4/.....	3.0	2.5	0.7	0.3	*	*	*	*	*	0.3
2003 4/.....	2.9	2.4	0.8	0.3	*	*	*	*	*	*
2002.....	2.7	2.6	0.8	*	0.3	*	*	*	*	*
2001.....	2.5	3.3	0.8	0.4	*	*	*	*	*	0.2
2000.....	2.4	3.0	0.7	0.3	*	*	*	*	*	0.2
1999.....	2.3	2.7	0.7	*	0.3	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

V. Mental and behavioral disorders (F01-F99)

All races 3/, both sexes

2006.....	30.6	*	*	*	*	0.4	1.2	1.7	2.3	3.5
2005.....	24.1	*	*	*	*	0.4	1.2	1.6	2.0	3.5
2004.....	21.0	*	*	*	0.1	0.4	1.0	1.6	2.1	3.8
2003.....	20.7	*	*	*	*	0.4	1.0	1.5	2.1	4.0
2002.....	19.3	*	*	*	0.1	0.3	0.8	1.3	2.0	4.0
2001.....	17.8	*	*	*	*	0.4	0.8	1.4	2.2	3.7
2000.....	16.4	*	*	*	*	0.4	0.9	1.3	2.3	3.9
1999.....	15.1	*	*	*	0.1	0.4	0.8	1.3	2.3	4.1

All races 3/, male

2006.....	22.7	*	*	*	*	0.6	1.6	2.5	3.3	4.5
2005.....	18.2	*	*	*	*	0.6	1.6	2.3	2.8	4.8
2004.....	16.2	*	*	*	*	0.5	1.4	2.4	3.0	5.5
2003.....	15.7	*	*	*	*	0.5	1.4	2.3	2.9	5.6
2002.....	14.8	*	*	*	*	0.4	1.1	1.8	2.7	5.6
2001.....	13.8	*	*	*	*	0.5	1.2	2.1	3.2	5.2
2000.....	12.8	*	*	*	*	0.5	1.3	1.9	3.3	6.0
1999.....	12.3	*	*	*	*	0.5	1.1	2.0	3.3	6.0

All races 3/, female

2006.....	38.3	*	*	*	*	0.2	0.7	1.0	1.3	2.4
2005.....	29.9	*	*	*	*	0.2	0.7	0.8	1.1	2.2
2004.....	25.6	*	*	*	*	0.4	0.7	0.8	1.3	2.1
2003.....	25.5	*	*	*	*	0.3	0.5	0.7	1.2	2.3
2002.....	23.7	*	*	*	*	0.2	0.5	0.7	1.3	2.4
2001.....	21.7	*	*	*	*	0.3	0.3	0.8	1.2	2.2
2000.....	19.8	*	*	*	*	0.3	0.5	0.7	1.3	1.8
1999.....	17.8	*	*	*	*	*	0.5	0.6	1.4	2.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
V. Mental and behavioral disorders (F01-F99)										
White, both sexes										
2006 4/.....	33.8	*	*	*	*	0.5	1.3	1.9	2.3	3.4
2005 4/.....	26.6	*	*	*	*	0.5	1.3	1.7	2.1	3.5
2004 4/.....	23.1	*	*	*	*	0.5	1.1	1.7	2.1	3.8
2003 4/.....	22.8	*	*	*	*	0.4	1.0	1.6	2.1	3.9
2002.....	21.3	*	*	*	*	0.4	0.9	1.3	2.1	3.9
2001.....	19.6	*	*	*	*	0.4	0.8	1.4	2.2	3.6
2000.....	17.9	*	*	*	*	0.4	0.9	1.4	2.2	3.7
1999.....	16.5	*	*	*	*	0.4	0.8	1.4	2.4	3.7
White, male										
2006 4/.....	24.4	*	*	*	*	0.7	1.8	2.7	3.4	4.5
2005 4/.....	19.5	*	*	*	*	0.7	1.8	2.4	3.1	4.9
2004 4/.....	17.2	*	*	*	*	0.5	1.6	2.4	2.9	5.5
2003 4/.....	16.7	*	*	*	*	0.5	1.5	2.4	2.9	5.6
2002.....	15.7	*	*	*	*	0.5	1.2	1.9	2.8	5.5
2001.....	14.5	*	*	*	*	0.5	1.2	2.0	3.2	5.1
2000.....	13.4	*	*	*	*	0.5	1.3	2.0	3.1	5.7
1999.....	12.8	*	*	*	*	0.6	1.1	2.1	3.4	5.6
White, female										
2006 4/.....	43.0	*	*	*	*	0.2	0.8	1.1	1.2	2.3
2005 4/.....	33.5	*	*	*	*	*	0.8	1.0	1.1	2.1
2004 4/.....	28.8	*	*	*	*	0.4	0.7	0.8	1.3	2.1
2003 4/.....	28.7	*	*	*	*	0.4	0.5	0.7	1.2	2.2
2002.....	26.7	*	*	*	*	0.3	0.5	0.7	1.3	2.3
2001.....	24.4	*	*	*	*	0.3	0.4	0.8	1.2	2.1
2000.....	22.2	*	*	*	*	0.3	0.6	0.8	1.2	1.7
1999.....	20.1	*	*	*	*	*	0.4	0.6	1.3	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
V. Mental and behavioral disorders (F01-F99)										
Black, both sexes										
2006 4/.....	20.8	*	*	*	*	*	0.7	1.1	2.3	3.5
2005 4/.....	16.5	*	*	*	*	*	0.6	1.4	1.6	3.7
2004 4/.....	14.4	*	*	*	*	*	*	1.7	2.5	4.1
2003 4/.....	14.0	*	*	*	*	*	0.7	1.6	2.2	5.2
2002.....	13.2	*	*	*	*	*	0.7	1.2	2.3	4.9
2001.....	12.6	*	*	*	*	*	0.8	1.4	2.3	4.9
2000.....	11.8	*	*	*	*	*	1.1	1.4	3.3	5.3
1999.....	10.7	*	*	*	*	*	0.9	1.3	2.4	6.8
Black, male										
2006 4/.....	18.3	*	*	*	*	*	*	1.6	3.0	4.3
2005 4/.....	14.7	*	*	*	*	*	*	2.4	2.2	4.6
2004 4/.....	14.0	*	*	*	*	*	*	3.0	4.1	5.8
2003 4/.....	13.4	*	*	*	*	*	*	2.2	2.7	7.4
2002.....	13.1	*	*	*	*	*	*	1.8	2.9	6.9
2001.....	13.0	*	*	*	*	*	1.4	2.5	3.4	7.1
2000.....	12.5	*	*	*	*	*	1.8	2.3	4.6	8.0
1999.....	11.9	*	*	*	*	*	*	2.0	3.1	9.8
Black, female										
2006 4/.....	23.1	*	*	*	*	*	*	*	1.6	2.9
2005 4/.....	18.2	*	*	*	*	*	*	*	*	2.9
2004 4/.....	14.8	*	*	*	*	*	*	*	*	2.6
2003 4/.....	14.5	*	*	*	*	*	*	*	1.8	3.3
2002.....	13.3	*	*	*	*	*	*	*	1.8	3.1
2001.....	12.1	*	*	*	*	*	*	*	1.4	3.0
2000.....	11.2	*	*	*	*	*	*	*	2.2	2.8
1999.....	9.6	*	*	*	*	*	*	*	1.8	4.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

All races 3/, both sexes

2006.....	3.9	*	*	*	*	0.3	1.0	1.5	2.1	3.1
2005.....	3.8	*	*	*	*	0.3	1.0	1.5	1.7	3.3
2004.....	3.6	*	*	*	*	0.3	0.9	1.4	2.0	3.5
2003.....	3.5	*	*	*	*	0.2	0.8	1.3	1.9	3.7
2002.....	3.3	*	*	*	*	0.2	0.7	1.1	1.8	3.7
2001.....	3.2	*	*	*	*	0.3	0.6	1.2	2.0	3.4
2000.....	3.1	*	*	*	*	0.2	0.7	1.1	2.1	3.6
1999.....	3.1	*	*	*	*	0.3	0.6	1.1	2.1	3.8

All races 3/, male

2006.....	5.9	*	*	*	*	0.5	1.4	2.3	3.1	4.2
2005.....	5.7	*	*	*	*	0.5	1.4	2.2	2.6	4.5
2004.....	5.6	*	*	*	*	0.4	1.3	2.2	2.8	5.2
2003.....	5.3	*	*	*	*	0.3	1.2	2.0	2.7	5.3
2002.....	5.1	*	*	*	*	0.3	1.0	1.7	2.5	5.3
2001.....	4.9	*	*	*	*	0.4	1.1	1.8	2.9	4.9
2000.....	4.8	*	*	*	*	0.3	1.2	1.7	3.0	5.7
1999.....	4.8	*	*	*	*	0.4	1.0	1.7	3.0	5.6

All races 3/, female

2006.....	2.1	*	*	*	*	*	0.5	0.7	1.0	2.0
2005.....	2.0	*	*	*	*	*	0.5	0.7	0.9	2.0
2004.....	1.8	*	*	*	*	*	0.4	0.6	1.1	1.8
2003.....	1.7	*	*	*	*	*	0.3	0.5	1.1	2.1
2002.....	1.6	*	*	*	*	*	0.3	0.5	1.0	2.2
2001.....	1.5	*	*	*	*	*	0.2	0.6	1.0	1.9
2000.....	1.5	*	*	*	*	*	0.3	0.6	1.1	1.6
1999.....	1.4	*	*	*	*	*	0.3	0.4	1.2	1.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

White, both sexes

2006 4/.....	4.0	*	*	*	*	0.4	1.1	1.7	2.1	3.1
2005 4/.....	3.9	*	*	*	*	0.4	1.1	1.6	1.9	3.2
2004 4/.....	3.6	*	*	*	*	0.3	1.0	1.5	1.9	3.5
2003 4/.....	3.4	*	*	*	*	0.3	0.8	1.4	1.9	3.6
2002.....	3.2	*	*	*	*	0.3	0.7	1.1	1.8	3.6
2001.....	3.0	*	*	*	*	0.3	0.7	1.2	1.9	3.3
2000.....	3.0	*	*	*	*	0.3	0.8	1.2	2.0	3.5
1999.....	2.9	*	*	*	*	0.3	0.7	1.1	2.1	3.4

White, male

2006 4/.....	5.9	*	*	*	*	0.6	1.7	2.5	3.2	4.2
2005 4/.....	5.8	*	*	*	*	0.6	1.7	2.3	2.8	4.6
2004 4/.....	5.5	*	*	*	*	0.5	1.4	2.3	2.8	5.2
2003 4/.....	5.2	*	*	*	*	0.3	1.3	2.2	2.7	5.2
2002.....	4.9	*	*	*	*	0.4	1.1	1.7	2.7	5.2
2001.....	4.6	*	*	*	*	0.5	1.1	1.8	2.9	4.7
2000.....	4.6	*	*	*	*	0.4	1.2	1.8	2.9	5.5
1999.....	4.6	*	*	*	*	0.4	1.0	1.7	3.1	5.2

White, female

2006 4/.....	2.1	*	*	*	*	*	0.6	0.8	0.9	1.9
2005 4/.....	2.0	*	*	*	*	*	0.5	0.9	0.9	1.8
2004 4/.....	1.8	*	*	*	*	*	0.5	0.6	1.1	1.8
2003 4/.....	1.7	*	*	*	*	*	0.3	0.5	1.0	1.9
2002.....	1.5	*	*	*	*	*	0.3	0.5	1.0	2.0
2001.....	1.5	*	*	*	*	*	0.3	0.6	1.0	1.8
2000.....	1.4	*	*	*	*	*	0.3	0.6	1.0	1.5
1999.....	1.3	*	*	*	*	*	0.3	0.4	1.1	1.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mental and behavioral disorders due to psychoactive substance use (F10-F19)										
Black, both sexes										
2006 4/.....	4.5	*	*	*	*	*	*	0.7	1.9	3.1
2005 4/.....	4.2	*	*	*	*	*	*	1.0	1.4	3.4
2004 4/.....	4.3	*	*	*	*	*	*	1.2	2.2	3.8
2003 4/.....	4.3	*	*	*	*	*	*	1.4	2.0	4.8
2002.....	4.4	*	*	*	*	*	*	1.1	2.0	4.7
2001.....	4.6	*	*	*	*	*	*	1.2	2.1	4.6
2000.....	4.6	*	*	*	*	*	0.9	1.1	3.1	4.8
1999.....	4.7	*	*	*	*	*	*	1.1	2.2	6.2
Black, male										
2006 4/.....	6.9	*	*	*	*	*	*	*	2.5	3.9
2005 4/.....	6.5	*	*	*	*	*	*	1.9	1.7	4.2
2004 4/.....	6.9	*	*	*	*	*	*	2.1	3.6	5.5
2003 4/.....	6.7	*	*	*	*	*	*	1.9	2.5	6.8
2002.....	7.0	*	*	*	*	*	*	1.6	2.7	6.6
2001.....	7.4	*	*	*	*	*	*	2.0	3.0	6.7
2000.....	7.3	*	*	*	*	*	1.5	2.0	4.3	7.2
1999.....	7.4	*	*	*	*	*	*	1.8	2.8	9.1
Black, female										
2006 4/.....	2.3	*	*	*	*	*	*	*	1.4	2.4
2005 4/.....	2.2	*	*	*	*	*	*	*	*	2.7
2004 4/.....	2.0	*	*	*	*	*	*	*	*	2.3
2003 4/.....	2.0	*	*	*	*	*	*	*	1.5	3.1
2002.....	2.1	*	*	*	*	*	*	*	1.4	3.0
2001.....	2.1	*	*	*	*	*	*	*	*	2.7
2000.....	2.1	*	*	*	*	*	*	*	2.0	2.6
1999.....	2.3	*	*	*	*	*	*	*	1.7	3.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mental and behavioral disorders due to use of alcohol (F10)										
All races 3/, both sexes										
2006.....	2.6	*	*	*	*	0.2	0.3	0.5	0.9	1.7
2005.....	2.5	*	*	*	*	0.1	0.3	0.5	0.7	1.8
2004.....	2.5	*	*	*	*	0.1	0.3	0.5	1.0	2.0
2003.....	2.4	*	*	*	*	0.1	0.3	0.4	1.0	2.0
2002.....	2.4	*	*	*	*	0.1	0.2	0.5	0.8	2.2
2001.....	2.3	*	*	*	*	0.1	0.2	0.5	1.0	2.2
2000.....	2.2	*	*	*	*	*	0.2	0.4	1.0	2.2
1999.....	2.2	*	*	*	*	*	0.2	0.4	1.0	2.3
All races 3/, male										
2006.....	4.0	*	*	*	*	0.3	0.4	0.9	1.3	2.5
2005.....	3.9	*	*	*	*	0.2	0.5	0.8	1.1	2.6
2004.....	3.9	*	*	*	*	*	0.5	0.8	1.4	3.0
2003.....	3.8	*	*	*	*	*	0.5	0.7	1.5	3.0
2002.....	3.8	*	*	*	*	*	0.3	0.8	1.3	3.3
2001.....	3.7	*	*	*	*	0.2	0.4	0.7	1.5	3.3
2000.....	3.6	*	*	*	*	*	0.3	0.7	1.5	3.6
1999.....	3.6	*	*	*	*	*	0.4	0.7	1.4	3.4
All races 3/, female										
2006.....	1.1	*	*	*	*	*	*	*	0.4	0.9
2005.....	1.1	*	*	*	*	*	*	0.2	0.3	0.9
2004.....	1.0	*	*	*	*	*	*	*	0.5	1.0
2003.....	1.1	*	*	*	*	*	*	*	0.4	1.0
2002.....	1.0	*	*	*	*	*	*	*	0.4	1.1
2001.....	1.0	*	*	*	*	*	*	0.2	0.5	1.0
2000.....	0.9	*	*	*	*	*	*	*	0.4	0.9
1999.....	0.9	*	*	*	*	*	*	*	0.6	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Mental and behavioral disorders due to use of alcohol (F10)										
White, both sexes										
2006 4/.....	2.6	*	*	*	*	0.2	0.3	0.6	0.9	1.7
2005 4/.....	2.6	*	*	*	*	0.1	0.3	0.6	0.7	1.8
2004 4/.....	2.5	*	*	*	*	*	0.3	0.5	0.9	2.0
2003 4/.....	2.4	*	*	*	*	0.1	0.2	0.5	1.0	2.1
2002.....	2.4	*	*	*	*	0.1	0.2	0.5	0.8	2.2
2001.....	2.3	*	*	*	*	0.1	0.2	0.5	1.0	2.2
2000.....	2.2	*	*	*	*	*	0.2	0.4	0.9	2.2
1999.....	2.1	*	*	*	*	*	0.3	0.4	1.0	2.1
White, male										
2006 4/.....	4.1	*	*	*	*	0.3	0.5	0.9	1.4	2.5
2005 4/.....	4.1	*	*	*	*	0.2	0.6	0.8	1.1	2.7
2004 4/.....	4.0	*	*	*	*	*	0.5	0.8	1.3	3.0
2003 4/.....	3.8	*	*	*	*	*	0.4	0.8	1.5	3.1
2002.....	3.8	*	*	*	*	*	0.3	0.8	1.3	3.4
2001.....	3.6	*	*	*	*	*	0.4	0.7	1.4	3.2
2000.....	3.5	*	*	*	*	*	0.3	0.7	1.4	3.6
1999.....	3.4	*	*	*	*	*	0.4	0.8	1.5	3.3
White, female										
2006 4/.....	1.2	*	*	*	*	*	*	*	0.3	0.9
2005 4/.....	1.1	*	*	*	*	*	*	0.3	0.3	0.9
2004 4/.....	1.1	*	*	*	*	*	*	*	0.5	1.0
2003 4/.....	1.1	*	*	*	*	*	*	*	0.4	1.1
2002.....	1.0	*	*	*	*	*	*	*	0.4	1.1
2001.....	1.0	*	*	*	*	*	*	*	0.5	1.1
2000.....	0.9	*	*	*	*	*	*	*	0.5	0.8
1999.....	0.9	*	*	*	*	*	*	*	0.6	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

All races 3/, both sexes

2006.....	1.4	*	*	*	*	0.2	0.7	1.0	1.2	1.4
2005.....	1.3	*	*	*	*	0.2	0.7	0.9	1.0	1.5
2004.....	1.2	*	*	*	*	0.2	0.6	0.9	1.0	1.5
2003.....	1.1	*	*	*	*	0.1	0.5	0.9	0.9	1.6
2002.....	0.9	*	*	*	*	0.1	0.5	0.6	1.0	1.5
2001.....	0.9	*	*	*	*	0.1	0.4	0.7	0.9	1.3
2000.....	0.9	*	*	*	*	0.1	0.5	0.7	1.1	1.4
1999.....	0.9	*	*	*	*	0.2	0.4	0.6	1.1	1.5

All races 3/, male

2006.....	1.9	*	*	*	*	0.3	1.0	1.4	1.7	1.7
2005.....	1.8	*	*	*	*	0.3	0.9	1.3	1.5	1.9
2004.....	1.6	*	*	*	*	0.3	0.8	1.4	1.4	2.2
2003.....	1.5	*	*	*	*	*	0.8	1.3	1.2	2.2
2002.....	1.3	*	*	*	*	*	0.7	0.9	1.3	1.9
2001.....	1.2	*	*	*	*	0.2	0.6	1.1	1.4	1.6
2000.....	1.2	*	*	*	*	*	0.8	1.0	1.5	2.1
1999.....	1.3	*	*	*	*	0.3	0.6	1.0	1.5	2.2

All races 3/, female

2006.....	0.9	*	*	*	*	*	0.4	0.5	0.6	1.1
2005.....	0.9	*	*	*	*	*	0.4	0.5	0.6	1.0
2004.....	0.7	*	*	*	*	*	0.3	0.4	0.6	0.9
2003.....	0.7	*	*	*	*	*	0.2	0.4	0.6	1.0
2002.....	0.6	*	*	*	*	*	0.3	0.4	0.6	1.0
2001.....	0.5	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	0.6	*	*	*	*	*	0.2	0.4	0.7	0.8
1999.....	0.5	*	*	*	*	*	*	0.3	0.6	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

White, both sexes

2006 4/.....	1.3	*	*	*	*	0.2	0.8	1.2	1.3	1.4
2005 4/.....	1.3	*	*	*	*	0.2	0.8	1.0	1.1	1.4
2004 4/.....	1.1	*	*	*	*	0.2	0.7	1.0	1.0	1.5
2003 4/.....	1.0	*	*	*	*	0.1	0.6	0.9	0.9	1.5
2002.....	0.9	*	*	*	*	0.1	0.5	0.7	1.0	1.4
2001.....	0.8	*	*	*	*	0.2	0.5	0.7	1.0	1.1
2000.....	0.8	*	*	*	*	0.2	0.6	0.7	1.0	1.3
1999.....	0.8	*	*	*	*	0.2	0.4	0.6	1.1	1.3

White, male

2006 4/.....	1.8	*	*	*	*	0.3	1.2	1.6	1.8	1.7
2005 4/.....	1.7	*	*	*	*	0.4	1.1	1.4	1.7	1.9
2004 4/.....	1.5	*	*	*	*	0.3	0.9	1.5	1.4	2.2
2003 4/.....	1.4	*	*	*	*	*	0.9	1.4	1.3	2.1
2002.....	1.2	*	*	*	*	*	0.7	1.0	1.4	1.8
2001.....	1.1	*	*	*	*	0.3	0.7	1.1	1.4	1.5
2000.....	1.1	*	*	*	*	*	0.8	1.0	1.4	1.9
1999.....	1.1	*	*	*	*	0.3	0.6	1.0	1.6	2.0

White, female

2006 4/.....	0.9	*	*	*	*	*	0.4	0.6	0.6	1.1
2005 4/.....	0.9	*	*	*	*	*	0.5	0.6	0.6	0.9
2004 4/.....	0.7	*	*	*	*	*	0.4	0.5	0.6	0.8
2003 4/.....	0.6	*	*	*	*	*	0.3	0.3	0.6	0.8
2002.....	0.6	*	*	*	*	*	*	0.3	0.6	1.0
2001.....	0.5	*	*	*	*	*	*	0.4	0.5	0.8
2000.....	0.5	*	*	*	*	*	*	0.5	0.6	0.7
1999.....	0.4	*	*	*	*	*	*	0.3	0.5	0.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

VI. Diseases of the nervous system (G00-G98)

All races 3/, both sexes

2006.....	41.1	9.0	1.6	1.0	1.3	2.1	2.5	2.3	2.6	3.4
2005.....	40.7	8.6	1.8	1.1	1.3	2.1	2.2	2.2	2.3	3.6
2004.....	38.1	9.2	2.0	1.1	1.4	1.9	2.1	2.2	2.4	3.2
2003.....	37.7	10.0	1.8	1.3	1.4	1.9	2.4	2.3	2.4	3.3
2002.....	35.7	9.5	2.1	1.1	1.3	1.8	2.2	2.1	2.3	3.3
2001.....	33.9	8.9	2.1	1.1	1.3	2.0	2.2	2.1	2.3	3.2
2000.....	32.3	10.2	2.1	1.1	1.4	1.8	2.0	1.9	2.3	3.2
1999.....	30.1	11.6	2.4	1.3	1.3	1.9	2.2	1.9	2.2	3.2

All races 3/, male

2006.....	32.7	9.9	1.7	1.2	1.5	2.7	3.2	2.8	2.8	3.9
2005.....	32.0	10.3	2.0	1.2	1.4	2.9	2.8	2.7	2.7	4.2
2004.....	30.0	10.5	2.0	1.2	1.7	2.6	2.6	2.6	2.7	3.6
2003.....	29.9	11.0	1.8	1.4	1.6	2.6	3.0	2.7	2.7	3.6
2002.....	28.5	11.0	2.4	1.2	1.5	2.3	2.8	2.6	2.8	4.0
2001.....	27.4	10.0	2.2	1.1	1.4	2.7	2.9	2.5	2.3	3.8
2000.....	26.2	11.5	2.1	1.1	1.6	2.2	2.6	2.3	2.6	3.5
1999.....	25.0	12.2	2.6	1.4	1.4	2.5	3.0	2.3	2.5	3.7

All races 3/, female

2006.....	49.2	8.1	1.5	0.9	1.1	1.5	1.7	1.9	2.3	2.8
2005.....	49.2	6.9	1.6	1.0	1.2	1.3	1.6	1.8	1.9	2.9
2004.....	45.9	8.0	2.0	1.0	1.2	1.1	1.6	1.7	2.1	2.9
2003.....	45.3	9.0	1.8	1.1	1.1	1.3	1.7	1.9	2.1	3.0
2002.....	42.8	7.8	1.7	1.0	1.1	1.2	1.6	1.7	1.8	2.5
2001.....	40.2	7.8	1.9	1.2	1.1	1.2	1.5	1.6	2.3	2.5
2000.....	38.2	8.8	2.0	1.1	1.2	1.3	1.3	1.5	2.1	2.8
1999.....	35.1	11.0	2.1	1.2	1.1	1.2	1.4	1.6	1.8	2.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
VI. Diseases of the nervous system (G00-G98)										
White, both sexes										
2006 4/.....	46.3	8.6	1.4	1.0	1.2	1.9	2.5	2.2	2.4	3.2
2005 4/.....	45.7	8.1	1.5	1.0	1.1	2.1	2.2	2.1	2.1	3.3
2004 4/.....	42.7	9.1	1.8	1.0	1.4	1.8	2.0	2.1	2.2	3.1
2003 4/.....	42.4	9.2	1.6	1.2	1.2	1.9	2.4	2.2	2.2	3.0
2002.....	40.2	9.3	1.8	1.1	1.2	1.7	2.0	2.0	2.2	3.1
2001.....	38.0	7.9	1.8	1.1	1.2	1.9	2.1	1.8	2.1	3.0
2000.....	36.3	9.4	1.8	1.1	1.3	1.7	2.0	1.8	2.2	3.0
1999.....	33.7	10.4	2.2	1.2	1.1	1.8	2.1	1.9	2.0	3.1
White, male										
2006 4/.....	36.6	9.4	1.5	1.1	1.5	2.5	3.3	2.8	2.7	3.8
2005 4/.....	35.6	9.9	1.7	1.1	1.3	2.9	2.8	2.5	2.6	4.0
2004 4/.....	33.4	10.6	1.8	1.2	1.6	2.4	2.5	2.7	2.5	3.5
2003 4/.....	33.3	11.1	1.5	1.4	1.4	2.6	3.1	2.6	2.4	3.3
2002.....	31.7	11.0	2.1	1.2	1.3	2.4	2.5	2.5	2.6	3.8
2001.....	30.5	8.7	2.1	1.1	1.4	2.6	2.8	2.2	2.1	3.7
2000.....	29.1	10.3	2.0	1.1	1.5	2.3	2.6	2.3	2.5	3.3
1999.....	27.7	11.8	2.5	1.2	1.2	2.5	2.8	2.3	2.3	3.6
White, female										
2006 4/.....	55.7	7.7	1.3	0.8	0.9	1.3	1.6	1.7	2.1	2.6
2005 4/.....	55.6	6.2	1.3	0.8	1.0	1.2	1.5	1.7	1.6	2.6
2004 4/.....	51.8	7.5	1.7	0.9	1.1	1.1	1.5	1.4	1.8	2.7
2003 4/.....	51.4	7.1	1.7	1.0	1.0	1.2	1.6	1.8	2.0	2.7
2002.....	48.6	7.5	1.5	0.9	1.0	1.0	1.5	1.5	1.7	2.3
2001.....	45.4	7.1	1.6	1.1	1.0	1.1	1.3	1.4	2.1	2.3
2000.....	43.2	8.4	1.6	1.0	1.1	1.1	1.3	1.3	1.9	2.6
1999.....	39.5	9.0	1.8	1.1	1.1	1.0	1.3	1.5	1.7	2.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
VI. Diseases of the nervous system (G00-G98)										
Black, both sexes										
2006 4/.....	22.9	12.2	2.5	1.4	1.9	3.2	2.9	3.6	4.3	5.0
2005 4/.....	23.5	10.7	2.9	2.0	2.2	2.6	3.0	3.5	4.2	5.9
2004 4/.....	21.9	10.4	3.2	1.7	1.9	2.8	3.2	3.3	4.8	5.0
2003 4/.....	20.8	14.6	2.8	1.7	2.2	2.3	3.0	3.6	4.2	6.1
2002.....	19.5	11.6	3.3	1.3	2.0	2.3	3.5	3.6	4.1	5.5
2001.....	19.0	14.4	3.1	1.5	1.8	3.0	3.0	3.9	4.2	5.4
2000.....	17.9	14.3	3.4	1.4	2.2	2.5	2.5	3.2	3.8	5.4
1999.....	16.7	17.8	3.3	1.9	1.8	2.5	3.5	2.8	3.5	5.3
Black, male										
2006 4/.....	18.2	13.8	2.6	1.7	1.9	4.1	3.3	3.5	4.6	5.6
2005 4/.....	19.0	12.1	3.1	2.1	2.5	3.5	3.8	4.4	4.6	6.6
2004 4/.....	17.3	10.4	3.2	1.2	2.1	3.9	3.5	2.8	4.9	5.0
2003 4/.....	17.5	12.5	3.2	1.3	2.8	2.8	3.3	4.0	5.2	6.6
2002.....	16.7	11.9	4.0	1.3	2.4	2.5	4.8	4.3	4.8	6.7
2001.....	16.1	17.1	2.9	1.2	1.7	4.0	3.5	5.0	4.5	6.5
2000.....	15.4	17.9	3.1	1.1	2.7	2.3	3.0	3.5	3.8	5.8
1999.....	14.6	15.9	3.2	2.3	2.4	2.9	5.0	3.3	3.9	5.6
Black, female										
2006 4/.....	27.2	10.5	2.4	*	1.9	2.3	2.6	3.7	4.1	4.5
2005 4/.....	27.5	9.2	2.6	1.9	1.9	1.7	2.1	2.7	4.0	5.3
2004 4/.....	26.1	10.5	3.3	2.2	1.7	1.7	2.9	3.8	4.7	5.0
2003 4/.....	23.9	16.7	2.3	2.1	1.6	1.8	2.6	3.2	3.3	5.7
2002.....	22.2	11.2	2.7	1.3	1.6	2.1	2.4	2.9	3.5	4.5
2001.....	21.5	11.7	3.4	1.8	1.8	1.9	2.5	2.9	4.0	4.4
2000.....	20.2	10.6	3.6	1.8	1.8	2.7	2.1	2.8	3.8	4.9
1999.....	18.5	19.8	3.4	1.5	1.2	2.0	2.1	2.3	3.1	5.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Multiple sclerosis (G35)										
All races 3/, both sexes										
2006.....	1.2	*	*	*	*	*	*	*	0.1	0.3
2005.....	1.2	*	*	*	*	*	*	0.1	0.1	0.4
2004.....	1.1	*	*	*	*	*	*	*	0.2	0.4
2003.....	1.1	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.1	*	*	*	*	*	*	*	0.1	0.4
2001.....	1.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	1.0	*	*	*	*	*	*	0.1	0.2	0.4
1999.....	1.0	*	*	*	*	*	*	*	0.2	0.4
All races 3/, male										
2006.....	0.8	*	*	*	*	*	*	*	*	0.2
2005.....	0.8	*	*	*	*	*	*	*	*	0.3
2004.....	0.8	*	*	*	*	*	*	*	*	0.3
2003.....	0.8	*	*	*	*	*	*	*	*	0.3
2002.....	0.8	*	*	*	*	*	*	*	*	0.4
2001.....	0.7	*	*	*	*	*	*	*	*	0.3
2000.....	0.7	*	*	*	*	*	*	*	*	0.3
1999.....	0.6	*	*	*	*	*	*	*	*	0.3
All races 3/, female										
2006.....	1.5	*	*	*	*	*	*	*	*	0.4
2005.....	1.6	*	*	*	*	*	*	*	*	0.4
2004.....	1.5	*	*	*	*	*	*	*	0.2	0.5
2003.....	1.4	*	*	*	*	*	*	*	0.2	0.6
2002.....	1.4	*	*	*	*	*	*	*	*	0.5
2001.....	1.4	*	*	*	*	*	*	*	0.2	0.5
2000.....	1.3	*	*	*	*	*	*	*	0.3	0.5
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Epilepsy (G40-G41)										
All races 3/, both sexes										
2006.....	0.5	*	0.2	*	*	0.3	0.4	0.5	0.4	0.4
2005.....	0.5	*	0.1	*	0.1	0.2	0.4	0.4	0.5	0.4
2004.....	0.5	*	0.2	*	0.1	0.2	0.3	0.4	0.5	0.5
2003.....	0.5	*	0.2	*	0.2	0.2	0.4	0.4	0.4	0.5
2002.....	0.5	*	0.1	0.1	0.1	0.2	0.4	0.4	0.4	0.5
2001.....	0.5	*	*	0.1	*	0.2	0.3	0.4	0.4	0.5
2000.....	0.5	*	0.2	*	0.1	0.2	0.3	0.3	0.4	0.5
1999.....	0.5	*	0.2	0.1	0.1	0.2	0.3	0.4	0.4	0.5
All races 3/, male										
2006.....	0.5	*	*	*	*	0.3	0.4	0.6	0.5	0.5
2005.....	0.5	*	*	*	*	0.3	0.5	0.4	0.6	0.5
2004.....	0.5	*	*	*	*	0.2	0.3	0.5	0.6	0.5
2003.....	0.5	*	*	*	0.2	0.2	0.4	0.5	0.5	0.6
2002.....	0.5	*	*	*	*	*	0.4	0.5	0.5	0.6
2001.....	0.5	*	*	*	*	0.2	0.4	0.5	0.4	0.7
2000.....	0.5	*	*	*	*	*	0.4	0.4	0.5	0.5
1999.....	0.5	*	0.3	*	*	0.3	0.4	0.6	0.5	0.7
All races 3/, female										
2006.....	0.4	*	*	*	*	0.3	0.3	0.4	0.3	0.3
2005.....	0.5	*	*	*	*	0.2	0.3	0.4	0.4	0.3
2004.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.4
2003.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.4
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.4	0.4
2001.....	0.4	*	*	*	*	*	*	0.3	0.4	0.3
2000.....	0.4	*	*	*	*	0.2	*	0.2	0.4	0.4
1999.....	0.4	*	*	*	*	*	0.3	0.3	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Epilepsy (G40-G41)										
White, both sexes										
2006 4/.....	0.4	*	0.2	*	*	0.2	0.3	0.5	0.4	0.4
2005 4/.....	0.5	*	*	*	*	0.2	0.4	0.3	0.4	0.4
2004 4/.....	0.4	*	0.2	*	0.1	0.2	0.3	0.4	0.4	0.4
2003 4/.....	0.4	*	0.2	*	0.1	0.2	0.4	0.4	0.4	0.5
2002.....	0.4	*	*	*	*	0.2	0.3	0.4	0.4	0.5
2001.....	0.4	*	*	0.1	*	0.2	0.2	0.3	0.4	0.4
2000.....	0.4	*	0.2	*	*	0.1	0.3	0.3	0.4	0.4
1999.....	0.4	*	0.2	*	*	0.2	0.3	0.4	0.4	0.5
White, male										
2006 4/.....	0.5	*	*	*	*	*	0.4	0.6	0.5	0.5
2005 4/.....	0.5	*	*	*	*	*	0.5	0.3	0.5	0.5
2004 4/.....	0.5	*	*	*	*	*	0.2	0.5	0.6	0.5
2003 4/.....	0.5	*	*	*	*	*	0.4	0.5	0.4	0.5
2002.....	0.5	*	*	*	*	*	0.3	0.4	0.5	0.6
2001.....	0.5	*	*	*	*	*	0.4	0.4	0.4	0.6
2000.....	0.5	*	*	*	*	*	0.3	0.5	0.4	0.4
1999.....	0.5	*	0.3	*	*	0.2	0.3	0.5	0.5	0.6
White, female										
2006 4/.....	0.4	*	*	*	*	0.3	0.3	0.4	*	0.3
2005 4/.....	0.4	*	*	*	*	*	0.3	0.4	0.3	0.3
2004 4/.....	0.4	*	*	*	*	*	0.3	0.3	0.3	0.3
2003 4/.....	0.4	*	*	*	*	*	0.3	0.3	0.3	0.4
2002.....	0.4	*	*	*	*	*	0.3	0.3	0.3	0.4
2001.....	0.4	*	*	*	*	*	*	0.3	0.4	0.3
2000.....	0.4	*	*	*	*	*	*	*	0.3	0.4
1999.....	0.4	*	*	*	*	*	0.3	*	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Infantile cerebral palsy (G80)										
All races 3/, both sexes										
2006.....	0.5	*	0.3	0.4	0.7	0.7	0.8	0.6	0.4	0.4
2005.....	0.5	*	0.4	0.6	0.5	0.7	0.6	0.5	0.4	0.4
2004.....	0.5	*	0.5	0.5	0.7	0.8	0.7	0.6	0.3	0.3
2003.....	0.5	*	0.4	0.6	0.6	0.7	0.7	0.6	0.4	0.3
2002.....	0.4	*	0.6	0.5	0.5	0.6	0.7	0.5	0.2	0.2
2001.....	0.4	*	0.5	0.4	0.5	0.7	0.6	0.4	0.3	0.2
2000.....	0.4	*	0.4	0.5	0.6	0.6	0.6	0.3	0.3	0.2
1999.....	0.4	*	0.5	0.4	0.5	0.5	0.5	0.4	0.3	0.3
All races 3/, male										
2006.....	0.6	*	0.4	0.5	0.7	0.9	1.0	0.6	0.4	0.4
2005.....	0.5	*	0.4	0.6	0.5	0.9	0.7	0.5	0.4	0.4
2004.....	0.6	*	0.4	0.5	0.7	1.0	0.8	0.7	0.3	0.3
2003.....	0.5	*	0.5	0.7	0.7	0.8	0.7	0.6	0.4	0.3
2002.....	0.5	*	0.7	0.6	0.5	0.8	0.8	0.5	0.3	0.3
2001.....	0.5	*	0.5	0.4	0.6	0.8	0.7	0.4	0.4	0.2
2000.....	0.4	*	0.3	0.5	0.7	0.7	0.7	0.4	0.4	0.2
1999.....	0.4	*	0.6	0.5	0.5	0.6	0.6	0.4	0.3	0.2
All races 3/, female										
2006.....	0.4	*	*	0.3	0.6	0.6	0.5	0.5	0.4	0.4
2005.....	0.4	*	0.4	0.6	0.5	0.5	0.4	0.4	0.3	0.4
2004.....	0.4	*	0.5	0.4	0.6	0.5	0.6	0.4	0.3	0.3
2003.....	0.4	*	0.4	0.5	0.5	0.6	0.6	0.5	0.3	0.3
2002.....	0.4	*	0.5	0.5	0.5	0.5	0.5	0.4	0.2	0.2
2001.....	0.3	*	0.4	0.5	0.5	0.6	0.5	0.3	0.3	0.2
2000.....	0.3	*	0.4	0.6	0.4	0.5	0.5	0.3	0.3	0.2
1999.....	0.3	*	0.4	0.4	0.4	0.4	0.3	0.4	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Infantile cerebral palsy (G80)										
White, both sexes										
2006 4/.....	0.5	*	0.2	0.4	0.6	0.6	0.7	0.5	0.4	0.4
2005 4/.....	0.5	*	0.3	0.4	0.4	0.7	0.5	0.4	0.4	0.4
2004 4/.....	0.5	*	0.3	0.4	0.6	0.7	0.7	0.6	0.3	0.3
2003 4/.....	0.4	*	0.3	0.5	0.5	0.6	0.6	0.6	0.3	0.3
2002.....	0.4	*	0.5	0.5	0.4	0.6	0.5	0.4	0.2	0.2
2001.....	0.4	*	0.4	0.4	0.5	0.6	0.6	0.3	0.3	0.2
2000.....	0.4	*	0.3	0.5	0.5	0.6	0.5	0.3	0.3	0.2
1999.....	0.3	*	0.5	0.4	0.4	0.5	0.4	0.4	0.3	0.3
White, male										
2006 4/.....	0.5	*	*	0.5	0.7	0.8	1.0	0.6	0.4	0.3
2005 4/.....	0.5	*	0.3	0.5	0.5	0.8	0.7	0.5	0.4	0.4
2004 4/.....	0.5	*	*	0.5	0.7	0.9	0.8	0.8	0.3	0.3
2003 4/.....	0.5	*	0.4	0.7	0.6	0.8	0.7	0.5	0.4	0.3
2002.....	0.4	*	0.6	0.5	0.4	0.8	0.6	0.4	*	0.3
2001.....	0.5	*	0.5	0.3	0.5	0.7	0.7	0.4	0.3	0.2
2000.....	0.4	*	*	0.5	0.7	0.7	0.6	0.4	0.3	0.2
1999.....	0.4	*	0.6	0.4	0.4	0.6	0.5	0.4	0.3	0.3
White, female										
2006 4/.....	0.4	*	*	0.3	0.5	0.5	0.5	0.4	0.4	0.4
2005 4/.....	0.4	*	*	0.4	0.4	0.5	0.4	0.3	0.4	0.4
2004 4/.....	0.4	*	0.4	0.3	0.5	0.5	0.5	0.4	0.3	0.3
2003 4/.....	0.4	*	*	0.4	0.4	0.5	0.5	0.6	0.3	0.3
2002.....	0.3	*	0.4	0.4	0.4	0.4	0.4	0.4	*	*
2001.....	0.3	*	*	0.4	0.4	0.5	0.4	0.3	0.3	0.3
2000.....	0.3	*	*	0.5	0.4	0.5	0.4	0.3	0.2	0.2
1999.....	0.3	*	*	0.4	0.4	0.4	0.3	0.4	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)										
All races 3/, both sexes										
2006.....	7.7	6.8	0.9	0.4	0.5	0.9	1.1	1.0	1.4	1.9
2005.....	7.3	6.7	1.0	0.4	0.6	1.0	1.0	1.1	1.1	2.0
2004.....	6.9	6.7	1.1	0.5	0.5	0.8	0.9	0.9	1.2	1.9
2003.....	7.1	7.3	0.9	0.5	0.6	0.9	1.2	1.1	1.2	1.8
2002.....	6.9	6.9	1.1	0.3	0.6	0.8	1.0	1.0	1.3	1.8
2001.....	6.7	6.1	1.2	0.5	0.6	0.9	1.1	1.0	1.1	1.8
2000.....	6.7	7.1	1.2	0.4	0.5	0.9	1.0	1.0	1.2	1.8
1999.....	6.5	7.7	1.2	0.5	0.6	0.9	1.2	0.9	1.1	1.7
All races 3/, male										
2006.....	8.3	7.7	1.0	0.5	0.6	1.4	1.6	1.3	1.6	2.4
2005.....	7.9	8.6	1.2	0.4	0.6	1.6	1.4	1.5	1.3	2.6
2004.....	7.3	7.8	1.1	0.5	0.6	1.2	1.3	1.2	1.4	2.2
2003.....	7.5	8.3	0.9	0.5	0.7	1.3	1.7	1.4	1.4	2.1
2002.....	7.3	8.4	1.3	0.4	0.7	1.2	1.4	1.4	1.7	2.4
2001.....	7.1	7.1	1.2	0.5	0.7	1.5	1.6	1.4	1.1	2.3
2000.....	7.2	8.1	1.3	0.4	0.6	1.3	1.4	1.3	1.5	2.1
1999.....	7.0	8.6	1.2	0.6	0.6	1.5	1.7	1.1	1.3	2.2
All races 3/, female										
2006.....	7.2	5.8	0.8	0.4	0.3	0.4	0.7	0.7	1.2	1.5
2005.....	6.8	4.7	0.9	0.3	0.5	0.4	0.6	0.7	0.8	1.5
2004.....	6.5	5.6	1.1	0.4	0.4	0.4	0.5	0.7	1.0	1.6
2003.....	6.6	6.2	1.0	0.5	0.4	0.5	0.6	0.7	1.0	1.4
2002.....	6.5	5.4	0.9	0.3	0.4	0.4	0.6	0.7	0.9	1.3
2001.....	6.3	5.0	1.2	0.5	0.5	0.3	0.7	0.7	1.1	1.2
2000.....	6.3	6.0	1.1	0.4	0.5	0.5	0.6	0.6	1.0	1.4
1999.....	6.1	6.8	1.2	0.5	0.5	0.4	0.6	0.6	0.8	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

White, both sexes

2006 4/.....	8.2	6.5	0.8	0.4	0.5	0.9	1.2	1.0	1.4	1.9
2005 4/.....	7.8	6.5	0.9	0.3	0.5	1.1	1.0	1.1	1.0	1.9
2004 4/.....	7.3	6.8	1.0	0.5	0.5	0.8	0.9	0.9	1.1	1.9
2003 4/.....	7.5	6.8	0.9	0.5	0.5	1.0	1.2	1.1	1.2	1.7
2002.....	7.3	6.9	1.0	0.3	0.6	0.8	1.0	1.0	1.3	1.8
2001.....	7.0	5.8	1.1	0.5	0.6	0.9	1.1	0.9	1.0	1.8
2000.....	7.1	6.9	1.1	0.4	0.5	0.9	1.0	0.9	1.2	1.7
1999.....	6.9	7.5	1.1	0.5	0.5	1.0	1.2	0.9	1.0	1.7

White, male

2006 4/.....	8.9	7.2	0.9	0.4	0.6	1.4	1.7	1.3	1.6	2.5
2005 4/.....	8.4	8.6	1.1	0.4	0.6	1.7	1.4	1.6	1.3	2.5
2004 4/.....	7.8	8.0	1.0	0.5	0.7	1.3	1.3	1.2	1.3	2.2
2003 4/.....	8.0	8.2	0.7	0.5	0.6	1.5	1.8	1.4	1.3	1.9
2002.....	7.8	8.5	1.1	0.4	0.7	1.3	1.4	1.4	1.7	2.4
2001.....	7.5	6.6	1.1	0.5	0.7	1.5	1.6	1.2	1.0	2.4
2000.....	7.6	7.4	1.2	0.4	0.5	1.4	1.5	1.3	1.5	2.1
1999.....	7.4	8.4	1.1	0.5	0.5	1.6	1.8	1.2	1.2	2.1

White, female

2006 4/.....	7.6	5.7	0.7	0.3	0.3	0.4	0.7	0.6	1.2	1.3
2005 4/.....	7.2	4.4	0.7	0.3	0.5	0.4	0.7	0.7	0.7	1.3
2004 4/.....	6.8	5.6	1.0	0.4	0.4	0.3	0.4	0.5	0.9	1.5
2003 4/.....	7.0	5.2	1.0	0.4	0.4	0.5	0.6	0.7	1.0	1.4
2002.....	6.9	5.2	0.8	0.3	0.4	0.3	0.5	0.6	0.8	1.3
2001.....	6.6	4.9	1.0	0.5	0.4	0.3	0.6	0.6	0.9	1.2
2000.....	6.6	6.3	1.0	0.5	0.4	0.4	0.6	0.5	0.9	1.3
1999.....	6.4	6.5	1.1	0.5	0.5	0.3	0.6	0.6	0.8	1.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IX. Diseases of the circulatory system (I00-I99)										
All races 3/, both sexes										
2006.....	276.5	13.1	1.4	0.7	1.1	2.2	4.5	7.4	13.6	24.0
2005.....	290.4	12.9	1.4	0.9	1.0	2.4	4.4	7.6	13.2	24.7
2004.....	294.9	14.5	1.5	0.6	1.0	2.2	4.3	7.5	13.0	24.8
2003.....	311.9	14.8	1.5	0.7	1.0	2.6	4.7	7.3	13.6	25.7
2002.....	320.2	16.5	1.5	0.6	1.1	2.5	4.0	7.3	13.1	25.9
2001.....	325.5	15.4	1.9	0.7	1.1	2.2	4.4	7.4	13.4	25.5
2000.....	334.6	17.4	1.5	0.7	1.1	2.5	4.4	7.0	12.5	24.7
1999.....	342.0	17.6	1.5	0.7	1.0	2.8	4.2	6.6	12.9	26.3
All races 3/, male										
2006.....	268.9	14.2	1.4	0.8	1.2	2.7	5.5	9.7	18.0	32.1
2005.....	279.4	14.0	1.6	0.8	1.1	3.0	5.6	9.7	17.0	32.5
2004.....	282.7	15.6	1.5	0.5	1.1	2.8	5.4	9.5	16.7	32.6
2003.....	297.7	16.2	1.5	0.8	1.2	3.1	5.5	8.7	17.4	34.0
2002.....	304.7	17.3	1.5	0.6	1.3	3.2	5.1	9.1	16.6	33.5
2001.....	307.6	15.8	2.0	0.6	1.1	2.7	5.4	8.9	16.8	33.5
2000.....	317.2	18.3	1.7	0.8	1.2	2.9	5.1	8.3	15.7	31.9
1999.....	324.2	18.5	1.6	0.8	1.1	3.4	4.9	7.9	16.4	34.6
All races 3/, female										
2006.....	283.8	12.0	1.3	0.6	1.0	1.8	3.3	4.9	9.2	15.8
2005.....	301.1	11.7	1.2	0.9	0.8	1.7	3.2	5.3	9.4	16.8
2004.....	306.7	13.5	1.5	0.7	1.0	1.6	3.2	5.5	9.3	16.9
2003.....	325.8	13.3	1.5	0.6	0.8	2.1	3.8	5.9	9.8	17.4
2002.....	335.2	15.7	1.4	0.7	0.9	1.7	2.9	5.4	9.5	18.2
2001.....	342.8	15.0	1.8	0.8	1.1	1.7	3.4	5.8	9.9	17.5
2000.....	351.3	16.5	1.4	0.6	1.0	2.1	3.7	5.6	9.4	17.6
1999.....	359.1	16.6	1.4	0.6	1.0	2.1	3.6	5.3	9.4	18.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IX. Diseases of the circulatory system (I00-I99)										
White, both sexes										
2006 4/.....	293.5	10.7	1.2	0.6	1.0	2.0	3.9	6.1	11.2	20.1
2005 4/.....	308.7	10.6	1.1	0.7	0.8	2.0	3.7	6.1	10.8	20.9
2004 4/.....	313.3	12.6	1.2	0.5	0.9	1.9	3.7	6.0	10.8	21.5
2003 4/.....	331.7	11.9	1.2	0.6	0.9	2.3	3.8	5.9	11.3	21.8
2002.....	341.0	14.1	1.3	0.6	0.9	2.2	3.4	5.9	10.6	22.1
2001.....	346.4	12.7	1.6	0.6	1.0	2.0	3.5	6.1	11.2	21.8
2000.....	356.4	15.3	1.4	0.6	0.9	2.1	3.6	5.7	10.4	21.1
1999.....	364.2	15.2	1.3	0.7	0.9	2.4	3.5	5.5	10.9	22.4
White, male										
2006 4/.....	282.7	11.4	1.2	0.8	1.2	2.5	5.0	8.0	15.1	27.3
2005 4/.....	294.4	11.5	1.4	0.7	1.0	2.5	4.7	8.1	14.0	28.0
2004 4/.....	298.1	13.7	1.3	0.5	1.0	2.4	4.7	7.8	14.1	28.6
2003 4/.....	314.0	12.5	1.1	0.6	1.1	2.7	4.5	7.1	14.7	29.9
2002.....	322.2	15.2	1.5	0.5	1.1	2.8	4.4	7.6	13.7	29.0
2001.....	324.9	13.7	1.6	0.6	1.0	2.3	4.4	7.5	14.3	29.2
2000.....	335.6	16.1	1.6	0.7	1.1	2.5	4.3	7.0	13.5	28.1
1999.....	342.5	15.9	1.5	0.8	0.9	2.9	3.9	6.6	14.2	30.2
White, female										
2006 4/.....	304.2	10.1	1.2	0.5	0.7	1.5	2.7	4.0	7.1	12.5
2005 4/.....	322.6	9.7	0.9	0.7	0.7	1.5	2.6	4.0	7.4	13.5
2004 4/.....	328.3	11.4	1.1	0.6	0.8	1.4	2.7	4.1	7.3	14.1
2003 4/.....	348.9	11.2	1.4	0.5	0.7	1.8	3.0	4.7	7.7	13.6
2002.....	359.5	13.0	1.2	0.7	0.8	1.5	2.3	4.1	7.4	14.9
2001.....	367.4	11.7	1.5	0.7	0.9	1.6	2.6	4.6	7.9	14.2
2000.....	376.6	14.4	1.1	0.5	0.8	1.7	2.9	4.4	7.2	13.9
1999.....	385.2	14.4	1.2	0.6	0.9	1.9	3.0	4.3	7.4	14.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
IX. Diseases of the circulatory system (I00-I99)										
Black, both sexes										
2006 4/.....	248.0	25.6	2.2	1.1	1.7	3.9	8.0	16.0	31.8	54.0
2005 4/.....	258.2	23.7	2.4	1.6	1.5	4.3	8.7	17.2	30.5	53.8
2004 4/.....	262.4	23.0	2.9	1.1	1.8	4.2	8.0	17.6	29.6	51.1
2003 4/.....	275.3	27.1	2.8	1.2	1.6	4.5	10.0	16.8	30.5	54.2
2002.....	278.2	28.6	2.0	0.8	2.0	4.1	8.1	17.2	31.0	54.4
2001.....	282.4	29.3	3.6	1.0	1.6	4.0	9.7	17.0	29.5	52.7
2000.....	287.0	28.6	2.6	1.0	1.9	4.8	9.4	15.9	28.7	52.4
1999.....	291.1	29.9	2.5	1.0	1.8	5.0	9.3	14.8	27.5	56.2
Black, male										
2006 4/.....	252.0	28.8	2.6	*	1.6	4.5	9.1	21.1	40.6	70.9
2005 4/.....	257.1	26.9	2.2	1.4	1.7	5.7	11.0	21.2	38.6	70.0
2004 4/.....	259.1	24.6	2.8	*	1.8	5.6	10.2	21.1	36.5	66.9
2003 4/.....	270.6	32.7	3.3	1.4	1.7	5.8	12.0	20.5	37.1	66.6
2002.....	270.5	28.8	2.0	*	2.3	5.3	9.8	21.1	38.5	68.3
2001.....	274.3	29.0	3.8	*	1.5	5.1	11.6	20.8	36.0	66.9
2000.....	277.6	29.7	2.2	*	1.8	5.3	10.5	18.7	33.7	63.2
1999.....	284.2	31.6	2.6	*	2.1	6.4	11.1	18.0	33.8	70.7
Black, female										
2006 4/.....	244.3	22.3	1.7	*	1.8	3.3	6.8	11.3	23.7	38.9
2005 4/.....	259.2	20.3	2.5	1.7	1.4	2.9	6.3	13.3	23.2	39.2
2004 4/.....	265.3	21.3	3.0	1.3	1.8	2.7	5.7	14.2	23.4	36.9
2003 4/.....	279.5	21.3	2.4	*	1.5	3.3	8.0	13.3	24.5	43.2
2002.....	285.2	28.5	2.1	*	1.6	2.8	6.5	13.6	24.2	42.0
2001.....	289.9	29.6	3.4	1.3	1.6	2.9	8.0	13.4	23.5	40.1
2000.....	295.6	27.4	2.9	*	2.0	4.4	8.4	13.5	24.1	42.8
1999.....	297.4	28.2	2.4	*	1.5	3.5	7.6	11.8	21.9	43.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive diseases (I10-I15)										
All races 3/, both sexes										
2006.....	18.9	*	*	*	*	*	0.3	0.7	1.5	2.9
2005.....	19.4	*	*	*	*	*	0.3	0.7	1.5	2.9
2004.....	18.6	*	*	*	*	0.1	0.2	0.7	1.4	2.7
2003.....	18.4	*	*	*	*	*	0.3	0.7	1.5	2.7
2002.....	17.2	*	*	*	*	*	0.2	0.7	1.3	2.5
2001.....	16.4	*	*	*	*	*	0.2	0.6	1.3	2.2
2000.....	15.9	*	*	*	*	*	0.2	0.5	1.0	2.0
1999.....	15.4	*	*	*	*	*	0.2	0.4	1.1	2.2
All races 3/, male										
2006.....	16.5	*	*	*	*	*	0.5	1.0	2.0	4.1
2005.....	16.5	*	*	*	*	*	0.3	0.9	1.9	4.1
2004.....	16.0	*	*	*	*	*	0.3	1.0	1.9	3.9
2003.....	15.4	*	*	*	*	*	0.4	0.8	1.9	3.6
2002.....	14.5	*	*	*	*	*	0.3	1.0	1.8	3.5
2001.....	13.5	*	*	*	*	*	0.3	0.7	1.7	2.8
2000.....	13.0	*	*	*	*	*	0.3	0.6	1.3	2.5
1999.....	12.6	*	*	*	*	*	0.3	0.6	1.5	2.8
All races 3/, female										
2006.....	21.2	*	*	*	*	*	*	0.4	0.9	1.6
2005.....	22.1	*	*	*	*	*	*	0.6	1.0	1.7
2004.....	21.2	*	*	*	*	*	*	0.5	0.9	1.5
2003.....	21.2	*	*	*	*	*	*	0.5	1.0	1.8
2002.....	19.9	*	*	*	*	*	*	0.4	0.9	1.6
2001.....	19.2	*	*	*	*	*	*	0.5	0.9	1.6
2000.....	18.6	*	*	*	*	*	*	0.4	0.7	1.5
1999.....	18.1	*	*	*	*	*	*	0.2	0.7	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive diseases (I10-I15)										
White, both sexes										
2006 4/.....	17.5	*	*	*	*	*	0.2	0.4	0.9	2.0
2005 4/.....	18.0	*	*	*	*	*	0.1	0.4	0.9	1.9
2004 4/.....	17.2	*	*	*	*	*	0.2	0.4	0.9	1.7
2003 4/.....	16.9	*	*	*	*	*	0.2	0.4	0.9	1.7
2002.....	15.8	*	*	*	*	*	0.2	0.4	0.8	1.5
2001.....	15.0	*	*	*	*	*	*	0.3	0.8	1.4
2000.....	14.4	*	*	*	*	*	0.1	0.3	0.5	1.2
1999.....	13.9	*	*	*	*	*	*	0.2	0.7	1.2
White, male										
2006 4/.....	14.7	*	*	*	*	*	0.3	0.5	1.3	3.0
2005 4/.....	14.6	*	*	*	*	*	*	0.6	1.2	2.9
2004 4/.....	14.2	*	*	*	*	*	0.2	0.6	1.3	2.5
2003 4/.....	13.6	*	*	*	*	*	0.3	0.6	1.2	2.6
2002.....	12.7	*	*	*	*	*	0.2	0.6	1.0	2.2
2001.....	11.8	*	*	*	*	*	*	0.4	1.1	2.0
2000.....	11.3	*	*	*	*	*	*	0.4	0.8	1.7
1999.....	10.8	*	*	*	*	*	*	0.4	1.1	1.7
White, female										
2006 4/.....	20.4	*	*	*	*	*	*	*	0.5	0.9
2005 4/.....	21.3	*	*	*	*	*	*	*	0.5	1.0
2004 4/.....	20.1	*	*	*	*	*	*	*	0.4	0.8
2003 4/.....	20.1	*	*	*	*	*	*	*	0.6	0.8
2002.....	18.8	*	*	*	*	*	*	*	0.5	0.9
2001.....	18.2	*	*	*	*	*	*	*	0.4	0.8
2000.....	17.4	*	*	*	*	*	*	*	0.3	0.7
1999.....	16.9	*	*	*	*	*	*	*	0.3	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive diseases (I10-I15)										
Black, both sexes										
2006 4/.....	31.8	*	*	*	*	*	0.8	2.3	5.6	9.3
2005 4/.....	32.7	*	*	*	*	*	1.1	2.4	5.5	9.7
2004 4/.....	32.2	*	*	*	*	*	0.7	3.0	4.8	9.7
2003 4/.....	32.1	*	*	*	*	*	1.0	2.2	5.4	9.1
2002.....	30.7	*	*	*	*	*	0.7	2.4	5.3	9.3
2001.....	29.1	*	*	*	*	*	0.7	2.0	4.9	7.9
2000.....	28.9	*	*	*	*	*	0.8	1.8	4.4	7.5
1999.....	28.6	*	*	*	*	*	1.0	1.5	3.9	8.9
Black, male										
2006 4/.....	32.2	*	*	*	*	*	1.2	3.4	7.4	12.9
2005 4/.....	32.3	*	*	*	*	*	1.3	2.3	6.9	12.7
2004 4/.....	31.3	*	*	*	*	*	*	3.7	6.0	14.1
2003 4/.....	30.8	*	*	*	*	*	1.4	2.2	7.2	11.0
2002.....	29.3	*	*	*	*	*	*	3.6	7.0	13.1
2001.....	27.7	*	*	*	*	*	*	2.4	6.2	9.1
2000.....	26.8	*	*	*	*	*	*	2.0	5.2	8.5
1999.....	26.8	*	*	*	*	*	*	1.9	4.4	11.2
Black, female										
2006 4/.....	31.4	*	*	*	*	*	*	1.3	4.0	6.1
2005 4/.....	33.0	*	*	*	*	*	*	2.4	4.1	6.9
2004 4/.....	33.0	*	*	*	*	*	*	2.4	3.8	5.8
2003 4/.....	33.3	*	*	*	*	*	*	2.2	3.8	7.5
2002.....	31.9	*	*	*	*	*	*	*	3.8	6.0
2001.....	30.5	*	*	*	*	*	*	1.7	3.8	6.8
2000.....	30.8	*	*	*	*	*	*	1.6	3.6	6.5
1999.....	30.3	*	*	*	*	*	*	*	3.6	6.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/ year 2/	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
All races 3/, both sexes										
2006.....	9.9	*	*	*	*	*	0.2	0.5	0.9	2.2
2005.....	9.9	*	*	*	*	*	0.2	0.6	1.1	2.2
2004.....	9.7	*	*	*	*	*	0.1	0.5	1.0	1.9
2003.....	9.7	*	*	*	*	*	0.2	0.5	1.0	1.9
2002.....	9.2	*	*	*	*	*	0.1	0.4	1.0	1.8
2001.....	8.7	*	*	*	*	*	*	0.3	0.9	1.6
2000.....	8.4	*	*	*	*	*	0.1	0.4	0.7	1.4
1999.....	8.1	*	*	*	*	*	0.1	0.3	0.8	1.5
All races 3/, male										
2006.....	9.3	*	*	*	*	*	0.3	0.7	1.3	3.2
2005.....	9.1	*	*	*	*	*	0.2	0.7	1.4	3.2
2004.....	8.9	*	*	*	*	*	*	0.7	1.5	2.8
2003.....	8.8	*	*	*	*	*	0.3	0.7	1.4	2.7
2002.....	8.2	*	*	*	*	*	0.2	0.7	1.4	2.6
2001.....	7.5	*	*	*	*	*	*	0.4	1.2	2.1
2000.....	7.2	*	*	*	*	*	0.2	0.4	0.9	1.9
1999.....	6.8	*	*	*	*	*	*	0.4	1.0	2.0
All races 3/, female										
2006.....	10.6	*	*	*	*	*	*	0.3	0.5	1.1
2005.....	10.7	*	*	*	*	*	*	0.4	0.7	1.2
2004.....	10.5	*	*	*	*	*	*	0.3	0.5	1.0
2003.....	10.7	*	*	*	*	*	*	0.3	0.7	1.2
2002.....	10.2	*	*	*	*	*	*	*	0.6	1.0
2001.....	9.8	*	*	*	*	*	*	0.3	0.5	1.1
2000.....	9.7	*	*	*	*	*	*	0.3	0.5	1.0
1999.....	9.4	*	*	*	*	*	*	*	0.5	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
Black, both sexes										
2006 4/.....	17.3	*	*	*	*	*	*	1.7	3.3	6.4
2005 4/.....	17.7	*	*	*	*	*	0.7	1.6	3.4	6.7
2004 4/.....	17.6	*	*	*	*	*	*	2.1	3.2	6.1
2003 4/.....	17.9	*	*	*	*	*	0.6	1.5	3.5	6.0
2002.....	17.2	*	*	*	*	*	*	1.5	3.9	5.8
2001.....	16.0	*	*	*	*	*	*	1.1	3.1	5.1
2000.....	16.1	*	*	*	*	*	*	1.4	2.8	4.9
1999.....	15.5	*	*	*	*	*	*	1.0	2.6	5.7
Black, male										
2006 4/.....	18.8	*	*	*	*	*	*	2.6	4.6	9.0
2005 4/.....	18.5	*	*	*	*	*	*	1.7	4.2	9.2
2004 4/.....	18.4	*	*	*	*	*	*	2.5	4.6	8.8
2003 4/.....	18.1	*	*	*	*	*	*	1.7	4.3	7.3
2002.....	17.5	*	*	*	*	*	*	2.4	5.4	8.5
2001.....	15.9	*	*	*	*	*	*	*	4.0	5.9
2000.....	15.6	*	*	*	*	*	*	1.6	3.5	5.8
1999.....	15.2	*	*	*	*	*	*	*	2.8	7.3
Black, female										
2006 4/.....	15.9	*	*	*	*	*	*	*	2.1	4.0
2005 4/.....	17.0	*	*	*	*	*	*	1.6	2.7	4.4
2004 4/.....	16.9	*	*	*	*	*	*	1.8	2.1	3.6
2003 4/.....	17.8	*	*	*	*	*	*	*	2.7	4.8
2002.....	16.9	*	*	*	*	*	*	*	2.6	3.4
2001.....	16.2	*	*	*	*	*	*	*	2.3	4.4
2000.....	16.5	*	*	*	*	*	*	*	2.1	4.1
1999.....	15.9	*	*	*	*	*	*	*	2.5	4.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes										
2006.....	142.1	0.5	*	*	*	0.2	0.5	1.4	3.9	8.8
2005.....	150.4	*	*	*	*	0.1	0.6	1.4	3.6	9.1
2004.....	153.7	*	*	*	*	0.1	0.5	1.5	3.9	9.4
2003.....	165.1	0.5	*	*	*	0.2	0.6	1.4	3.9	10.1
2002.....	171.4	0.6	*	*	*	0.1	0.5	1.3	3.6	10.1
2001.....	176.3	0.6	*	*	*	0.2	0.7	1.5	4.0	9.9
2000.....	183.1	0.8	*	*	*	0.2	0.5	1.2	3.4	9.8
1999.....	189.8	0.7	*	*	*	0.3	0.5	1.3	3.7	10.4
All races 3/, male										
2006.....	152.2	*	*	*	*	0.3	0.7	2.1	6.1	13.2
2005.....	159.0	*	*	*	*	0.2	0.9	2.2	5.4	13.6
2004.....	161.6	*	*	*	*	0.2	0.7	2.1	6.0	13.9
2003.....	172.2	*	*	*	*	0.2	0.8	2.0	5.9	15.2
2002.....	178.4	*	*	*	*	*	0.7	2.0	5.2	15.2
2001.....	181.7	*	*	*	*	0.2	0.9	2.2	5.9	15.0
2000.....	188.7	*	*	*	*	0.2	0.6	1.8	5.1	14.7
1999.....	195.4	*	*	*	*	0.3	0.6	1.7	5.5	15.8
All races 3/, female										
2006.....	132.3	*	*	*	*	*	0.2	0.6	1.7	4.4
2005.....	142.0	*	*	*	*	*	0.2	0.6	1.9	4.6
2004.....	146.1	*	*	*	*	*	0.2	0.9	1.8	4.9
2003.....	158.1	*	*	*	*	*	0.4	0.8	1.9	4.9
2002.....	164.7	*	*	*	*	*	0.3	0.6	1.9	4.9
2001.....	171.2	*	*	*	*	*	0.4	0.7	2.1	4.9
2000.....	177.6	*	*	*	*	*	0.3	0.6	1.7	5.0
1999.....	184.5	*	*	*	*	*	0.3	0.8	1.9	5.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/ year 2/	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
White, both sexes										
2006 4/.....	153.4	*	*	*	*	0.2	0.5	1.3	3.7	8.3
2005 4/.....	162.6	*	*	*	*	0.1	0.5	1.2	3.5	8.5
2004 4/.....	166.1	*	*	*	*	*	0.4	1.3	3.6	9.2
2003 4/.....	178.3	*	*	*	*	0.2	0.5	1.3	3.7	9.8
2002.....	185.4	*	*	*	*	0.1	0.4	1.2	3.3	9.6
2001.....	190.5	*	*	*	*	0.1	0.5	1.3	3.8	9.5
2000.....	197.9	0.8	*	*	*	0.2	0.4	1.0	3.1	9.4
1999.....	205.1	*	*	*	*	0.2	0.3	1.2	3.5	9.9
White, male										
2006 4/.....	163.7	*	*	*	*	0.3	0.7	2.0	5.7	12.6
2005 4/.....	171.5	*	*	*	*	*	0.8	1.9	5.0	12.7
2004 4/.....	174.3	*	*	*	*	*	0.5	1.9	5.6	13.6
2003 4/.....	185.5	*	*	*	*	*	0.7	1.8	5.6	15.0
2002.....	192.5	*	*	*	*	*	0.6	1.9	5.0	14.6
2001.....	195.8	*	*	*	*	*	0.8	2.0	5.5	14.5
2000.....	203.6	*	*	*	*	*	0.6	1.6	4.8	14.3
1999.....	210.5	*	*	*	*	0.3	0.5	1.6	5.1	15.1
White, female										
2006 4/.....	143.3	*	*	*	*	*	*	0.5	1.5	3.8
2005 4/.....	153.9	*	*	*	*	*	*	0.5	1.8	4.1
2004 4/.....	158.1	*	*	*	*	*	*	0.7	1.6	4.7
2003 4/.....	171.4	*	*	*	*	*	0.4	0.7	1.7	4.4
2002.....	178.4	*	*	*	*	*	*	0.5	1.6	4.5
2001.....	185.3	*	*	*	*	*	0.3	0.6	1.9	4.5
2000.....	192.4	*	*	*	*	*	0.3	0.5	1.4	4.5
1999.....	200.0	*	*	*	*	*	*	0.7	1.7	4.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
Black, both sexes										
2006 4/.....	112.6	*	*	*	*	*	0.7	2.3	6.2	14.7
2005 4/.....	117.8	*	*	*	*	*	0.9	2.6	5.3	15.5
2004 4/.....	121.0	*	*	*	*	*	1.2	2.9	6.5	12.7
2003 4/.....	129.2	*	*	*	*	*	1.2	2.4	6.1	14.0
2002.....	132.9	*	*	*	*	*	0.9	2.1	5.6	15.1
2001.....	136.9	*	*	*	*	*	1.5	2.8	6.4	14.9
2000.....	140.1	*	*	*	*	*	0.7	2.4	6.0	14.6
1999.....	143.7	*	*	*	*	*	1.2	1.9	5.8	16.1
Black, male										
2006 4/.....	120.2	*	*	*	*	*	*	3.2	9.3	20.7
2005 4/.....	122.9	*	*	*	*	*	1.3	4.2	7.9	23.0
2004 4/.....	125.2	*	*	*	*	*	1.8	3.4	9.6	18.4
2003 4/.....	133.4	*	*	*	*	*	1.6	3.1	9.2	19.6
2002.....	135.3	*	*	*	*	*	*	2.7	7.5	22.0
2001.....	138.9	*	*	*	*	*	1.9	3.9	9.3	21.6
2000.....	141.5	*	*	*	*	*	*	3.7	8.0	20.3
1999.....	146.0	*	*	*	*	*	*	2.9	8.4	23.3
Black, female										
2006 4/.....	105.6	*	*	*	*	*	*	1.4	3.3	9.4
2005 4/.....	113.1	*	*	*	*	*	*	*	3.0	8.8
2004 4/.....	117.1	*	*	*	*	*	*	2.4	3.7	7.6
2003 4/.....	125.3	*	*	*	*	*	*	1.7	3.3	9.1
2002.....	130.8	*	*	*	*	*	*	1.5	4.0	9.0
2001.....	135.2	*	*	*	*	*	*	1.7	3.8	9.0
2000.....	138.8	*	*	*	*	*	*	*	4.3	9.7
1999.....	141.7	*	*	*	*	*	*	*	3.4	9.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes										
2006.....	47.2	*	*	*	*	0.1	0.2	0.6	1.5	3.6
2005.....	50.9	*	*	*	*	*	0.3	0.6	1.3	3.5
2004.....	53.4	*	*	*	*	*	0.2	0.6	1.6	3.8
2003.....	58.7	*	*	*	*	*	0.3	0.5	1.5	4.1
2002.....	62.3	*	*	*	*	*	0.2	0.5	1.5	4.0
2001.....	64.9	*	*	*	*	*	0.2	0.6	1.8	4.2
2000.....	68.5	0.5	*	*	*	*	0.2	0.6	1.3	4.0
1999.....	71.5	*	*	*	*	0.1	0.2	0.5	1.6	4.1
All races 3/, male										
2006.....	51.6	*	*	*	*	*	0.2	0.9	2.3	5.2
2005.....	54.8	*	*	*	*	*	0.4	1.0	2.0	5.1
2004.....	57.4	*	*	*	*	*	0.3	0.8	2.3	5.7
2003.....	62.6	*	*	*	*	*	0.3	0.8	2.2	6.2
2002.....	66.2	*	*	*	*	*	0.3	0.7	2.1	6.0
2001.....	68.6	*	*	*	*	*	0.3	0.7	2.5	6.2
2000.....	72.7	*	*	*	*	*	0.2	0.8	1.9	5.8
1999.....	76.1	*	*	*	*	*	0.3	0.7	2.3	6.2
All races 3/, female										
2006.....	43.0	*	*	*	*	*	*	0.3	0.8	1.9
2005.....	47.2	*	*	*	*	*	*	0.3	0.7	2.0
2004.....	49.6	*	*	*	*	*	*	0.4	0.8	1.9
2003.....	54.8	*	*	*	*	*	0.2	0.3	0.8	2.0
2002.....	58.4	*	*	*	*	*	*	0.3	0.8	2.1
2001.....	61.3	*	*	*	*	*	*	0.4	1.0	2.1
2000.....	64.6	*	*	*	*	*	*	0.3	0.7	2.2
1999.....	67.0	*	*	*	*	*	*	0.3	0.9	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
White, both sexes										
2006 4/.....	51.0	*	*	*	*	*	0.2	0.6	1.5	3.5
2005 4/.....	55.1	*	*	*	*	*	0.2	0.5	1.3	3.4
2004 4/.....	57.8	*	*	*	*	*	0.1	0.5	1.6	3.8
2003 4/.....	63.4	*	*	*	*	*	0.2	0.5	1.4	4.1
2002.....	67.3	*	*	*	*	*	0.2	0.4	1.4	4.0
2001.....	70.1	*	*	*	*	*	0.2	0.5	1.7	4.1
2000.....	74.1	*	*	*	*	*	0.2	0.5	1.2	4.0
1999.....	77.4	*	*	*	*	*	0.1	0.5	1.5	4.0
White, male										
2006 4/.....	55.8	*	*	*	*	*	0.3	0.8	2.2	5.3
2005 4/.....	59.5	*	*	*	*	*	0.4	0.8	1.9	4.9
2004 4/.....	62.3	*	*	*	*	*	*	0.7	2.4	5.6
2003 4/.....	67.7	*	*	*	*	*	0.3	0.7	2.0	6.3
2002.....	71.8	*	*	*	*	*	0.3	0.6	2.1	5.9
2001.....	74.4	*	*	*	*	*	0.3	0.6	2.5	6.1
2000.....	78.8	*	*	*	*	*	*	0.6	1.8	6.0
1999.....	82.7	*	*	*	*	*	*	0.6	2.1	6.2
White, female										
2006 4/.....	46.3	*	*	*	*	*	*	0.3	0.7	1.7
2005 4/.....	50.8	*	*	*	*	*	*	*	0.7	1.8
2004 4/.....	53.3	*	*	*	*	*	*	0.4	0.7	1.9
2003 4/.....	59.2	*	*	*	*	*	*	*	0.8	1.9
2002.....	62.8	*	*	*	*	*	*	*	0.7	1.9
2001.....	66.0	*	*	*	*	*	*	0.3	0.9	2.0
2000.....	69.6	*	*	*	*	*	*	0.3	0.6	2.0
1999.....	72.3	*	*	*	*	*	*	0.3	0.9	1.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Acute myocardial infarction (I21-I22)										
Black, both sexes										
2006 4/.....	37.4	*	*	*	*	*	*	1.0	2.4	5.2
2005 4/.....	39.8	*	*	*	*	*	*	1.4	2.0	5.4
2004 4/.....	42.0	*	*	*	*	*	*	1.2	2.2	4.9
2003 4/.....	45.7	*	*	*	*	*	*	0.9	2.3	5.6
2002.....	48.5	*	*	*	*	*	*	0.9	2.3	5.5
2001.....	50.0	*	*	*	*	*	*	1.1	2.5	5.8
2000.....	52.2	*	*	*	*	*	*	1.2	2.3	4.9
1999.....	53.3	*	*	*	*	*	*	0.8	2.4	5.5
Black, male										
2006 4/.....	39.1	*	*	*	*	*	*	1.4	3.4	6.9
2005 4/.....	40.3	*	*	*	*	*	*	2.1	2.9	7.5
2004 4/.....	42.4	*	*	*	*	*	*	*	2.7	7.7
2003 4/.....	46.6	*	*	*	*	*	*	*	3.5	7.5
2002.....	48.3	*	*	*	*	*	*	*	3.0	7.6
2001.....	49.2	*	*	*	*	*	*	*	3.4	8.0
2000.....	52.0	*	*	*	*	*	*	1.6	2.7	6.5
1999.....	53.4	*	*	*	*	*	*	*	3.4	7.7
Black, female										
2006 4/.....	35.7	*	*	*	*	*	*	*	1.4	3.7
2005 4/.....	39.2	*	*	*	*	*	*	*	*	3.6
2004 4/.....	41.6	*	*	*	*	*	*	*	1.8	2.4
2003 4/.....	44.9	*	*	*	*	*	*	*	*	3.9
2002.....	48.8	*	*	*	*	*	*	*	1.6	3.6
2001.....	50.6	*	*	*	*	*	*	*	1.7	3.8
2000.....	52.5	*	*	*	*	*	*	*	1.9	3.6
1999.....	53.2	*	*	*	*	*	*	*	1.4	3.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes

2006.....	93.5	*	*	*	*	*	0.3	0.8	2.3	5.1
2005.....	98.2	*	*	*	*	*	0.3	0.8	2.3	5.5
2004.....	99.5	*	*	*	*	*	0.3	0.9	2.3	5.5
2003.....	105.3	*	*	*	*	*	0.3	0.9	2.4	5.8
2002.....	108.0	*	*	*	*	*	0.3	0.8	2.0	6.0
2001.....	110.3	*	*	*	*	*	0.4	0.9	2.2	5.6
2000.....	113.3	*	*	*	*	*	0.3	0.6	2.0	5.7
1999.....	117.2	*	*	*	*	0.1	0.3	0.7	2.1	6.2

All races 3/, male

2006.....	99.2	*	*	*	*	*	0.4	1.2	3.7	7.9
2005.....	102.9	*	*	*	*	*	0.5	1.2	3.4	8.4
2004.....	103.3	*	*	*	*	*	0.4	1.3	3.6	8.1
2003.....	108.5	*	*	*	*	*	0.4	1.2	3.7	8.8
2002.....	110.9	*	*	*	*	*	0.4	1.3	3.0	9.2
2001.....	111.9	*	*	*	*	*	0.5	1.5	3.3	8.6
2000.....	114.8	*	*	*	*	*	0.4	1.1	3.1	8.7
1999.....	118.0	*	*	*	*	0.2	0.3	1.0	3.2	9.4

All races 3/, female

2006.....	88.0	*	*	*	*	*	*	0.3	0.9	2.3
2005.....	93.7	*	*	*	*	*	*	0.3	1.2	2.5
2004.....	95.7	*	*	*	*	*	*	0.5	0.9	3.0
2003.....	102.3	*	*	*	*	*	0.2	0.5	1.0	2.8
2002.....	105.2	*	*	*	*	*	*	0.4	1.0	2.8
2001.....	108.8	*	*	*	*	*	0.2	0.3	1.1	2.7
2000.....	111.9	*	*	*	*	*	*	0.2	1.0	2.8
1999.....	116.4	*	*	*	*	*	*	0.4	1.0	3.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
White, both sexes										
2006 4/.....	101.0	*	*	*	*	*	0.3	0.7	2.1	4.7
2005 4/.....	106.2	*	*	*	*	*	0.3	0.7	2.2	5.0
2004 4/.....	107.5	*	*	*	*	*	0.2	0.8	2.0	5.4
2003 4/.....	113.8	*	*	*	*	*	0.3	0.8	2.2	5.6
2002.....	116.9	*	*	*	*	*	0.2	0.8	1.8	5.6
2001.....	119.1	*	*	*	*	*	0.3	0.8	2.0	5.3
2000.....	122.6	*	*	*	*	*	0.2	0.6	1.9	5.3
1999.....	126.5	*	*	*	*	*	0.2	0.7	1.9	5.8
White, male										
2006 4/.....	106.5	*	*	*	*	*	0.4	1.1	3.4	7.3
2005 4/.....	110.7	*	*	*	*	*	0.4	1.1	3.1	7.6
2004 4/.....	111.1	*	*	*	*	*	0.3	1.2	3.1	7.9
2003 4/.....	116.6	*	*	*	*	*	0.4	1.1	3.5	8.5
2002.....	119.4	*	*	*	*	*	0.4	1.3	2.8	8.6
2001.....	120.1	*	*	*	*	*	0.4	1.4	3.0	8.2
2000.....	123.5	*	*	*	*	*	0.3	0.9	2.9	8.2
1999.....	126.5	*	*	*	*	*	*	0.9	3.0	8.8
White, female										
2006 4/.....	95.6	*	*	*	*	*	*	*	0.8	2.0
2005 4/.....	101.9	*	*	*	*	*	*	0.3	1.1	2.3
2004 4/.....	103.9	*	*	*	*	*	*	0.3	0.9	2.8
2003 4/.....	111.0	*	*	*	*	*	*	0.4	0.9	2.5
2002.....	114.4	*	*	*	*	*	*	0.3	0.8	2.5
2001.....	118.1	*	*	*	*	*	*	*	0.9	2.4
2000.....	121.6	*	*	*	*	*	*	*	0.8	2.4
1999.....	126.5	*	*	*	*	*	*	0.4	0.9	2.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other forms of chronic ischemic heart disease (I20,I25)										
Black, both sexes										
2006 4/.....	73.9	*	*	*	*	*	*	1.3	3.6	9.0
2005 4/.....	76.6	*	*	*	*	*	*	1.1	3.3	9.8
2004 4/.....	78.0	*	*	*	*	*	*	1.6	4.1	7.7
2003 4/.....	82.3	*	*	*	*	*	*	1.5	3.7	8.2
2002.....	83.2	*	*	*	*	*	*	1.1	3.2	9.6
2001.....	85.8	*	*	*	*	*	0.8	1.7	3.8	8.9
2000.....	86.6	*	*	*	*	*	*	1.3	3.6	9.5
1999.....	89.3	*	*	*	*	*	0.7	1.1	3.4	10.1
Black, male										
2006 4/.....	79.6	*	*	*	*	*	*	1.8	5.7	13.2
2005 4/.....	81.0	*	*	*	*	*	*	2.0	4.9	15.3
2004 4/.....	81.5	*	*	*	*	*	*	1.9	6.7	10.5
2003 4/.....	85.5	*	*	*	*	*	*	2.1	5.7	11.7
2002.....	85.7	*	*	*	*	*	*	*	4.1	14.4
2001.....	88.2	*	*	*	*	*	*	2.4	5.9	13.3
2000.....	88.0	*	*	*	*	*	*	2.1	5.0	13.4
1999.....	91.1	*	*	*	*	*	*	1.7	5.0	14.9
Black, female										
2006 4/.....	68.7	*	*	*	*	*	*	*	1.8	5.2
2005 4/.....	72.6	*	*	*	*	*	*	*	1.8	4.9
2004 4/.....	74.7	*	*	*	*	*	*	*	1.7	5.2
2003 4/.....	79.4	*	*	*	*	*	*	*	2.0	5.0
2002.....	81.0	*	*	*	*	*	*	*	2.3	5.3
2001.....	83.5	*	*	*	*	*	*	*	1.9	4.9
2000.....	85.3	*	*	*	*	*	*	*	2.4	5.9
1999.....	87.6	*	*	*	*	*	*	*	1.9	5.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes										
2006.....	20.4	*	*	*	*	*	*	0.2	0.8	1.9
2005.....	21.2	*	*	*	*	*	*	0.2	0.7	1.9
2004.....	21.7	*	*	*	*	*	*	0.2	0.8	1.7
2003.....	23.1	*	*	*	*	*	*	0.3	0.8	1.9
2002.....	23.6	*	*	*	*	*	*	0.2	0.7	1.9
2001.....	23.9	*	*	*	*	*	*	0.2	0.7	1.8
2000.....	24.7	*	*	*	*	*	*	0.2	0.5	1.7
1999.....	25.8	*	*	*	*	*	*	0.2	0.6	2.0
All races 3/, male										
2006.....	23.2	*	*	*	*	*	*	0.4	1.4	3.1
2005.....	23.6	*	*	*	*	*	*	0.4	1.1	3.0
2004.....	23.9	*	*	*	*	*	*	0.3	1.2	2.5
2003.....	25.2	*	*	*	*	*	*	0.4	1.3	2.9
2002.....	25.4	*	*	*	*	*	*	0.3	1.1	3.0
2001.....	25.3	*	*	*	*	*	*	0.3	1.0	2.8
2000.....	26.0	*	*	*	*	*	*	0.3	0.9	2.8
1999.....	26.8	*	*	*	*	*	*	0.2	0.9	3.1
All races 3/, female										
2006.....	17.7	*	*	*	*	*	*	*	0.3	0.7
2005.....	18.8	*	*	*	*	*	*	*	0.3	0.8
2004.....	19.5	*	*	*	*	*	*	*	0.3	0.8
2003.....	21.1	*	*	*	*	*	*	*	0.2	0.9
2002.....	21.9	*	*	*	*	*	*	*	0.3	0.8
2001.....	22.5	*	*	*	*	*	*	*	0.4	0.8
2000.....	23.5	*	*	*	*	*	*	*	0.2	0.7
1999.....	24.8	*	*	*	*	*	*	*	0.3	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
All races 3/, both sexes										
2006.....	73.1	*	*	*	*	*	0.2	0.5	1.5	3.2
2005.....	77.0	*	*	*	*	*	0.2	0.5	1.6	3.6
2004.....	77.8	*	*	*	*	*	0.2	0.7	1.5	3.9
2003.....	82.2	*	*	*	*	*	0.3	0.6	1.6	3.9
2002.....	84.4	*	*	*	*	*	0.2	0.6	1.3	4.1
2001.....	86.4	*	*	*	*	*	0.3	0.7	1.5	3.8
2000.....	88.6	*	*	*	*	*	0.2	0.5	1.5	4.0
1999.....	91.4	*	*	*	*	0.1	0.2	0.6	1.5	4.3
All races 3/, male										
2006.....	76.1	*	*	*	*	*	0.3	0.9	2.3	4.8
2005.....	79.3	*	*	*	*	*	0.4	0.9	2.3	5.4
2004.....	79.4	*	*	*	*	*	0.3	1.0	2.3	5.6
2003.....	83.3	*	*	*	*	*	0.3	0.8	2.3	5.9
2002.....	85.5	*	*	*	*	*	0.3	0.9	1.9	6.2
2001.....	86.5	*	*	*	*	*	0.4	1.1	2.3	5.8
2000.....	88.8	*	*	*	*	*	0.2	0.8	2.2	5.9
1999.....	91.2	*	*	*	*	0.2	0.2	0.8	2.2	6.3
All races 3/, female										
2006.....	70.3	*	*	*	*	*	*	*	0.6	1.6
2005.....	74.8	*	*	*	*	*	*	0.2	0.9	1.7
2004.....	76.2	*	*	*	*	*	*	0.4	0.7	2.1
2003.....	81.1	*	*	*	*	*	*	0.4	0.8	1.9
2002.....	83.3	*	*	*	*	*	*	0.3	0.8	2.0
2001.....	86.2	*	*	*	*	*	*	0.2	0.7	1.9
2000.....	88.4	*	*	*	*	*	*	*	0.8	2.1
1999.....	91.6	*	*	*	*	*	*	0.3	0.7	2.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
White, both sexes										
2006 4/.....	80.3	*	*	*	*	*	0.2	0.5	1.4	2.9
2005 4/.....	84.7	*	*	*	*	*	0.2	0.5	1.6	3.3
2004 4/.....	85.4	*	*	*	*	*	0.2	0.6	1.4	3.7
2003 4/.....	90.2	*	*	*	*	*	0.2	0.5	1.5	3.7
2002.....	92.8	*	*	*	*	*	0.2	0.6	1.2	3.9
2001.....	94.8	*	*	*	*	*	0.2	0.6	1.4	3.6
2000.....	97.4	*	*	*	*	*	0.1	0.4	1.4	3.7
1999.....	100.4	*	*	*	*	*	0.1	0.5	1.4	4.0
White, male										
2006 4/.....	83.4	*	*	*	*	*	0.3	0.8	2.2	4.4
2005 4/.....	87.0	*	*	*	*	*	0.3	0.8	2.3	5.0
2004 4/.....	87.0	*	*	*	*	*	0.2	0.9	2.1	5.5
2003 4/.....	91.3	*	*	*	*	*	0.3	0.7	2.2	5.7
2002.....	93.9	*	*	*	*	*	0.3	0.9	1.8	5.9
2001.....	94.8	*	*	*	*	*	0.4	1.1	2.1	5.5
2000.....	97.5	*	*	*	*	*	*	0.7	2.1	5.6
1999.....	99.9	*	*	*	*	*	*	0.7	2.1	6.0
White, female										
2006 4/.....	77.3	*	*	*	*	*	*	*	0.6	1.3
2005 4/.....	82.4	*	*	*	*	*	*	*	0.9	1.6
2004 4/.....	83.8	*	*	*	*	*	*	0.3	0.6	2.0
2003 4/.....	89.1	*	*	*	*	*	*	0.3	0.7	1.6
2002.....	91.7	*	*	*	*	*	*	*	0.6	1.9
2001.....	94.9	*	*	*	*	*	*	*	0.6	1.7
2000.....	97.3	*	*	*	*	*	*	*	0.6	1.8
1999.....	100.8	*	*	*	*	*	*	0.3	0.6	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
All races 3/, both sexes										
2006.....	56.9	7.7	0.9	0.4	0.7	1.5	2.7	3.9	5.6	7.4
2005.....	57.6	8.3	0.9	0.5	0.6	1.7	2.7	3.8	5.4	7.8
2004.....	56.6	10.0	1.1	0.4	0.7	1.5	2.5	3.6	5.0	7.8
2003.....	58.5	10.4	1.1	0.5	0.7	1.7	2.7	3.6	5.4	7.8
2002.....	58.8	11.7	1.0	0.4	0.7	1.8	2.3	3.8	5.2	8.1
2001.....	58.6	11.1	1.4	0.5	0.7	1.5	2.5	3.6	5.1	8.0
2000.....	58.8	12.0	1.1	0.5	0.7	1.8	2.6	3.5	5.2	7.5
1999.....	59.4	12.9	1.2	0.5	0.7	2.0	2.6	3.5	5.3	8.2
All races 3/, male										
2006.....	51.0	8.0	1.0	0.5	0.8	1.8	3.3	4.9	7.0	9.2
2005.....	51.4	8.9	0.9	0.4	0.7	2.3	3.4	4.9	6.7	9.6
2004.....	50.6	10.5	1.1	0.3	0.8	1.9	3.2	4.5	6.1	9.4
2003.....	52.3	11.4	1.0	0.5	0.8	2.1	3.2	4.1	6.5	9.5
2002.....	52.4	12.1	1.0	0.4	0.8	2.4	3.0	4.6	6.6	9.6
2001.....	51.8	10.8	1.5	0.4	0.8	1.9	3.1	4.3	6.0	9.9
2000.....	52.2	12.2	1.2	0.6	0.9	2.1	3.1	4.2	6.2	9.1
1999.....	53.1	13.1	1.3	0.5	0.8	2.4	3.0	4.0	6.4	10.3
All races 3/, female										
2006.....	62.6	7.5	0.9	0.4	0.6	1.1	2.0	2.8	4.2	5.6
2005.....	63.5	7.7	0.8	0.5	0.6	1.1	1.9	2.7	4.1	6.0
2004.....	62.4	9.4	1.1	0.5	0.6	1.2	1.8	2.6	3.9	6.1
2003.....	64.5	9.3	1.2	0.4	0.5	1.3	2.1	3.1	4.3	6.1
2002.....	65.0	11.4	1.0	0.5	0.5	1.1	1.7	3.0	3.8	6.6
2001.....	65.2	11.5	1.3	0.5	0.7	1.1	1.9	2.9	4.1	6.0
2000.....	65.1	11.8	0.9	0.4	0.6	1.4	2.1	2.9	4.1	5.9
1999.....	65.5	12.7	1.0	0.4	0.7	1.5	2.2	2.9	4.2	6.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
White, both sexes										
2006 4/.....	60.9	6.0	0.8	0.4	0.6	1.2	2.3	3.2	4.4	5.9
2005 4/.....	61.4	6.9	0.7	0.4	0.5	1.5	2.3	3.2	4.1	6.4
2004 4/.....	60.4	8.7	0.9	0.3	0.6	1.3	2.2	2.8	4.0	6.5
2003 4/.....	62.2	8.2	0.9	0.4	0.6	1.5	2.2	2.8	4.4	6.4
2002.....	62.6	10.0	0.9	0.4	0.6	1.5	1.9	3.0	4.1	6.6
2001.....	62.3	9.3	1.1	0.4	0.7	1.3	2.0	2.9	4.1	6.5
2000.....	62.5	10.6	1.0	0.4	0.6	1.4	2.1	2.8	4.3	6.1
1999.....	62.9	11.0	1.0	0.4	0.6	1.8	2.1	2.7	4.4	6.7
White, male										
2006 4/.....	53.6	6.1	0.9	0.4	0.7	1.6	2.9	4.1	5.6	7.3
2005 4/.....	54.0	7.2	0.8	0.3	0.6	1.9	2.9	4.2	5.2	8.1
2004 4/.....	53.0	9.1	0.9	*	0.7	1.6	2.9	3.6	5.0	8.0
2003 4/.....	54.7	8.8	0.8	0.4	0.8	1.8	2.5	3.2	5.5	7.9
2002.....	54.9	10.5	1.0	0.4	0.7	2.1	2.5	3.6	5.2	7.9
2001.....	54.1	9.6	1.2	0.4	0.7	1.5	2.4	3.5	4.8	8.1
2000.....	54.6	10.8	1.2	0.5	0.8	1.7	2.5	3.4	5.2	7.5
1999.....	55.1	11.4	1.1	0.5	0.7	2.1	2.4	3.3	5.4	8.5
White, female										
2006 4/.....	68.1	5.9	0.8	0.3	0.5	0.9	1.6	2.2	3.2	4.5
2005 4/.....	68.7	6.5	0.6	0.4	0.5	1.0	1.5	2.1	3.0	4.6
2004 4/.....	67.6	8.3	0.8	0.4	0.5	1.0	1.5	2.0	3.1	4.9
2003 4/.....	69.6	7.6	1.0	0.3	0.4	1.1	1.8	2.4	3.2	5.0
2002.....	70.2	9.6	0.8	0.4	0.5	1.0	1.3	2.3	2.9	5.3
2001.....	70.4	8.9	1.1	0.5	0.6	1.0	1.5	2.3	3.4	4.8
2000.....	70.2	10.4	0.7	0.3	0.4	1.2	1.6	2.2	3.3	4.6
1999.....	70.5	10.7	0.8	0.4	0.6	1.4	1.9	2.2	3.4	4.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other heart diseases (I26-I51)										
Black, both sexes										
2006 4/.....	50.1	16.0	1.6	0.6	1.2	2.8	4.9	8.6	14.3	18.7
2005 4/.....	51.4	15.2	1.7	1.1	1.1	3.2	5.1	8.6	14.0	18.5
2004 4/.....	50.8	14.8	2.3	0.8	1.3	3.0	4.5	8.0	11.9	17.9
2003 4/.....	52.5	19.6	2.0	0.9	1.0	3.0	5.9	8.9	12.8	18.1
2002.....	52.5	20.2	1.5	*	1.2	3.1	5.1	9.5	13.0	18.9
2001.....	52.8	20.1	2.9	0.8	1.1	2.9	5.5	8.7	12.3	18.4
2000.....	52.7	19.0	1.9	0.8	1.4	3.6	6.0	8.8	11.9	17.8
1999.....	54.4	22.3	2.0	0.7	1.2	3.5	5.6	8.6	12.4	19.5
Black, male										
2006 4/.....	50.1	17.0	1.9	*	1.4	3.4	5.3	10.8	18.0	24.0
2005 4/.....	50.6	17.7	*	*	1.2	4.6	6.6	10.4	17.8	22.6
2004 4/.....	50.4	15.7	2.3	*	1.3	4.0	5.6	10.4	13.9	21.5
2003 4/.....	51.5	23.2	2.2	*	1.2	3.9	7.5	10.5	14.3	22.5
2002.....	50.8	20.9	*	*	1.3	4.0	6.2	11.9	16.4	22.8
2001.....	51.7	17.7	3.2	*	1.2	3.8	7.0	10.7	14.9	23.1
2000.....	51.5	18.9	1.9	*	1.4	4.3	6.7	10.1	14.3	21.6
1999.....	54.3	21.8	2.1	*	1.5	4.8	6.8	10.2	15.2	24.3
Black, female										
2006 4/.....	50.0	15.0	*	*	*	2.2	4.4	6.4	10.9	13.9
2005 4/.....	52.2	12.6	2.0	1.3	*	1.7	3.7	6.9	10.6	14.8
2004 4/.....	51.1	13.9	2.3	*	1.3	2.0	3.4	5.8	10.1	14.7
2003 4/.....	53.4	15.8	1.8	*	*	2.1	4.2	7.3	11.4	14.3
2002.....	54.0	19.4	1.8	*	*	2.2	4.0	7.4	9.8	15.5
2001.....	53.8	22.7	2.5	*	*	2.0	4.1	6.9	10.0	14.1
2000.....	53.9	19.2	2.0	*	1.3	2.9	5.3	7.6	9.8	14.5
1999.....	54.5	22.8	1.8	*	*	2.2	4.5	7.1	9.9	15.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
All races 3/, both sexes										
2006.....	4.1	2.0	0.1	*	*	0.2	0.3	0.6	0.9	1.3
2005.....	4.6	2.4	0.1	*	*	0.2	0.4	0.6	1.0	1.4
2004.....	4.5	3.8	0.2	*	*	0.2	0.4	0.7	0.9	1.4
2003.....	4.8	3.3	0.2	*	*	0.2	0.4	0.7	1.1	1.5
2002.....	4.8	4.6	0.2	*	0.1	0.2	0.3	0.7	0.9	1.4
2001.....	4.7	4.2	0.1	*	*	0.2	0.4	0.8	1.0	1.4
2000.....	4.6	4.9	0.2	*	*	0.2	0.4	0.6	1.1	1.4
1999.....	4.7	5.7	*	0.1	*	0.3	0.5	0.7	1.0	1.5
All races 3/, male										
2006.....	3.2	2.0	*	*	*	*	0.3	0.5	0.7	1.1
2005.....	3.7	2.6	*	*	*	*	0.3	0.4	0.7	1.2
2004.....	3.6	4.1	*	*	*	*	0.3	0.5	0.7	1.1
2003.....	3.9	3.7	*	*	*	0.2	0.3	0.4	0.8	1.3
2002.....	3.8	4.7	*	*	*	*	0.2	0.7	0.7	1.3
2001.....	3.8	3.9	*	*	*	*	0.3	0.6	0.7	1.1
2000.....	3.8	5.1	*	*	*	*	0.3	0.5	0.9	1.2
1999.....	3.9	5.9	*	*	*	0.3	0.3	0.5	0.8	1.3
All races 3/, female										
2006.....	5.0	1.9	*	*	*	0.2	0.4	0.7	1.2	1.4
2005.....	5.4	2.1	*	*	*	0.3	0.4	0.9	1.2	1.5
2004.....	5.3	3.5	*	*	*	0.3	0.5	0.8	1.1	1.8
2003.....	5.7	2.9	*	*	*	0.3	0.6	0.9	1.3	1.8
2002.....	5.7	4.5	*	*	*	0.2	0.5	0.8	1.1	1.6
2001.....	5.5	4.6	*	*	*	*	0.6	0.9	1.3	1.7
2000.....	5.5	4.6	*	*	*	0.3	0.6	0.8	1.2	1.6
1999.....	5.5	5.6	*	*	*	0.3	0.7	0.9	1.2	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
White, both sexes										
2006 4/.....	4.2	1.3	*	*	*	0.1	0.3	0.5	0.7	1.0
2005 4/.....	4.6	2.1	*	*	*	0.2	0.3	0.5	0.7	1.2
2004 4/.....	4.5	3.2	*	*	*	0.2	0.3	0.5	0.7	1.2
2003 4/.....	4.8	2.5	*	*	*	0.2	0.3	0.5	0.9	1.2
2002.....	4.8	3.9	*	*	*	0.2	0.3	0.5	0.7	1.3
2001.....	4.7	3.6	*	*	*	0.1	0.4	0.6	0.8	1.1
2000.....	4.7	4.1	0.2	*	*	0.2	0.4	0.5	0.9	1.2
1999.....	4.7	5.2	*	*	*	0.3	0.4	0.5	0.8	1.3
White, male										
2006 4/.....	3.3	1.3	*	*	*	*	0.2	0.4	0.4	0.9
2005 4/.....	3.7	2.4	*	*	*	*	0.3	0.3	0.5	1.1
2004 4/.....	3.6	3.4	*	*	*	*	*	0.4	0.6	0.8
2003 4/.....	3.9	2.9	*	*	*	*	*	0.3	0.7	1.0
2002.....	3.9	4.1	*	*	*	*	*	0.5	0.6	1.1
2001.....	3.8	3.4	*	*	*	*	0.3	0.5	0.6	0.9
2000.....	3.8	4.3	*	*	*	*	0.3	0.5	0.7	1.1
1999.....	3.9	5.4	*	*	*	0.2	*	0.4	0.6	1.2
White, female										
2006 4/.....	5.1	1.4	*	*	*	*	0.4	0.6	0.9	1.1
2005 4/.....	5.5	1.8	*	*	*	0.3	0.4	0.7	0.9	1.3
2004 4/.....	5.4	3.0	*	*	*	0.3	0.4	0.6	0.9	1.6
2003 4/.....	5.7	2.0	*	*	*	0.3	0.5	0.8	1.1	1.4
2002.....	5.8	3.7	*	*	*	*	0.4	0.5	0.8	1.4
2001.....	5.6	3.9	*	*	*	*	0.5	0.7	1.1	1.4
2000.....	5.5	3.9	*	*	*	0.3	0.4	0.6	1.0	1.3
1999.....	5.5	4.9	*	*	*	0.4	0.6	0.6	1.0	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
Black, both sexes										
2006 4/.....	5.2	5.1	*	*	*	*	0.7	1.4	2.6	3.2
2005 4/.....	5.6	4.2	*	*	*	*	0.7	1.6	2.8	3.0
2004 4/.....	5.6	5.9	*	*	*	*	1.1	1.6	2.2	3.4
2003 4/.....	6.1	6.5	*	*	*	*	1.2	1.4	2.5	3.8
2002.....	5.9	8.2	*	*	*	*	0.8	2.2	2.5	3.1
2001.....	5.8	6.8	*	*	*	*	0.8	1.9	2.3	3.2
2000.....	5.7	8.0	*	*	*	*	1.0	1.6	2.5	3.1
1999.....	6.2	9.4	*	*	*	*	1.0	1.8	2.6	3.3
Black, male										
2006 4/.....	4.2	*	*	*	*	*	*	*	2.4	2.6
2005 4/.....	4.6	*	*	*	*	*	*	*	2.0	2.5
2004 4/.....	4.6	6.5	*	*	*	*	*	1.6	1.5	3.3
2003 4/.....	5.2	7.1	*	*	*	*	*	*	2.0	3.4
2002.....	4.5	9.0	*	*	*	*	*	2.2	1.8	2.9
2001.....	4.8	6.9	*	*	*	*	*	*	1.7	2.8
2000.....	4.6	7.7	*	*	*	*	*	*	2.5	2.5
1999.....	5.1	8.8	*	*	*	*	*	*	2.5	2.4
Black, female										
2006 4/.....	6.1	*	*	*	*	*	*	1.4	2.8	3.7
2005 4/.....	6.6	*	*	*	*	*	*	2.0	3.6	3.4
2004 4/.....	6.5	*	*	*	*	*	*	1.5	2.9	3.5
2003 4/.....	6.9	*	*	*	*	*	1.3	1.6	2.9	4.2
2002.....	7.2	7.3	*	*	*	*	*	2.2	3.1	3.3
2001.....	6.8	6.6	*	*	*	*	*	2.4	2.9	3.5
2000.....	6.8	8.3	*	*	*	*	1.5	2.2	2.6	3.6
1999.....	7.2	10.1	*	*	*	*	*	2.3	2.7	4.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary embolism (I26)										
All races 3/, both sexes										
2006.....	2.3	*	*	*	*	*	0.2	0.4	0.7	0.8
2005.....	2.8	*	*	*	*	0.1	0.3	0.4	0.6	1.0
2004.....	2.8	*	*	*	*	0.1	0.3	0.4	0.6	1.0
2003.....	3.0	*	*	*	*	0.1	0.3	0.4	0.7	1.1
2002.....	3.0	*	*	*	*	0.1	0.3	0.5	0.6	1.0
2001.....	3.0	*	*	*	*	*	0.3	0.5	0.7	0.9
2000.....	3.0	*	*	*	*	0.1	0.3	0.4	0.7	1.0
1999.....	3.2	*	*	*	*	0.2	0.3	0.4	0.7	0.9
All races 3/, male										
2006.....	2.0	*	*	*	*	*	*	0.3	0.5	0.8
2005.....	2.4	*	*	*	*	*	0.2	0.3	0.5	1.0
2004.....	2.3	*	*	*	*	*	0.2	0.4	0.5	0.7
2003.....	2.6	*	*	*	*	*	0.2	0.3	0.6	0.9
2002.....	2.5	*	*	*	*	*	*	0.5	0.5	0.9
2001.....	2.6	*	*	*	*	*	0.2	0.4	0.5	0.8
2000.....	2.6	*	*	*	*	*	*	0.3	0.6	0.8
1999.....	2.7	*	*	*	*	*	0.3	0.3	0.6	0.9
All races 3/, female										
2006.....	2.6	*	*	*	*	*	0.2	0.5	0.8	0.9
2005.....	3.2	*	*	*	*	*	0.3	0.5	0.8	0.9
2004.....	3.2	*	*	*	*	*	0.3	0.5	0.8	1.3
2003.....	3.4	*	*	*	*	*	0.4	0.5	0.8	1.2
2002.....	3.5	*	*	*	*	*	0.3	0.5	0.7	1.1
2001.....	3.5	*	*	*	*	*	0.4	0.6	0.9	1.1
2000.....	3.4	*	*	*	*	*	0.4	0.5	0.7	1.1
1999.....	3.7	*	*	*	*	*	0.5	0.6	0.8	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary embolism (I26)										
White, both sexes										
2006 4/.....	2.3	*	*	*	*	*	0.2	0.4	0.5	0.6
2005 4/.....	2.8	*	*	*	*	0.1	0.2	0.3	0.5	0.9
2004 4/.....	2.8	*	*	*	*	*	0.2	0.4	0.6	0.8
2003 4/.....	3.0	*	*	*	*	*	0.2	0.3	0.6	0.8
2002.....	3.0	*	*	*	*	*	0.2	0.3	0.5	0.9
2001.....	3.1	*	*	*	*	*	0.3	0.4	0.6	0.7
2000.....	3.0	*	*	*	*	*	0.2	0.4	0.6	0.8
1999.....	3.2	*	*	*	*	0.1	0.3	0.3	0.5	0.8
White, male										
2006 4/.....	2.0	*	*	*	*	*	*	0.3	0.3	0.6
2005 4/.....	2.4	*	*	*	*	*	*	0.2	0.4	0.9
2004 4/.....	2.3	*	*	*	*	*	*	0.3	0.5	0.5
2003 4/.....	2.6	*	*	*	*	*	*	*	0.5	0.7
2002.....	2.5	*	*	*	*	*	*	0.3	0.4	0.7
2001.....	2.6	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	2.6	*	*	*	*	*	*	0.4	0.5	0.7
1999.....	2.7	*	*	*	*	*	*	*	0.5	0.8
White, female										
2006 4/.....	2.7	*	*	*	*	*	*	0.4	0.7	0.7
2005 4/.....	3.2	*	*	*	*	*	*	0.4	0.5	0.8
2004 4/.....	3.2	*	*	*	*	*	0.3	0.4	0.7	1.2
2003 4/.....	3.4	*	*	*	*	*	0.3	0.5	0.6	0.9
2002.....	3.5	*	*	*	*	*	0.3	0.4	0.6	1.0
2001.....	3.6	*	*	*	*	*	0.3	0.4	0.8	0.8
2000.....	3.4	*	*	*	*	*	0.3	0.3	0.6	0.8
1999.....	3.7	*	*	*	*	*	0.4	0.4	0.6	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pulmonary embolism (I26)										
Black, both sexes										
2006 4/.....	3.0	*	*	*	*	*	*	0.9	1.9	2.2
2005 4/.....	3.6	*	*	*	*	*	*	1.3	1.9	2.1
2004 4/.....	3.6	*	*	*	*	*	*	0.9	1.4	2.4
2003 4/.....	4.0	*	*	*	*	*	1.0	1.0	1.9	2.9
2002.....	3.8	*	*	*	*	*	*	1.6	1.5	2.1
2001.....	3.8	*	*	*	*	*	*	1.4	1.7	2.4
2000.....	3.9	*	*	*	*	*	0.9	0.9	1.6	2.3
1999.....	4.3	*	*	*	*	*	0.8	1.3	2.0	2.0
Black, male										
2006 4/.....	2.7	*	*	*	*	*	*	*	2.0	1.8
2005 4/.....	3.2	*	*	*	*	*	*	*	*	2.0
2004 4/.....	3.2	*	*	*	*	*	*	*	*	2.4
2003 4/.....	3.6	*	*	*	*	*	*	*	1.6	2.7
2002.....	3.0	*	*	*	*	*	*	1.6	*	2.0
2001.....	3.3	*	*	*	*	*	*	*	*	2.1
2000.....	3.3	*	*	*	*	*	*	*	1.5	2.0
1999.....	3.8	*	*	*	*	*	*	*	2.0	1.4
Black, female										
2006 4/.....	3.3	*	*	*	*	*	*	*	1.8	2.4
2005 4/.....	3.9	*	*	*	*	*	*	1.6	2.6	2.1
2004 4/.....	3.9	*	*	*	*	*	*	*	1.6	2.4
2003 4/.....	4.3	*	*	*	*	*	*	*	2.2	3.1
2002.....	4.4	*	*	*	*	*	*	1.5	1.6	2.2
2001.....	4.3	*	*	*	*	*	*	1.7	2.0	2.7
2000.....	4.4	*	*	*	*	*	*	1.4	1.6	2.6
1999.....	4.8	*	*	*	*	*	*	1.5	2.1	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

All races 3/, both sexes

2006.....	1.8	1.9	0.1	*	*	0.1	0.1	0.2	0.3	0.4
2005.....	1.8	2.3	0.1	*	*	*	0.1	0.2	0.3	0.4
2004.....	1.7	3.7	0.1	*	*	0.1	0.1	0.2	0.3	0.4
2003.....	1.8	3.2	0.1	*	*	0.1	0.1	0.2	0.4	0.5
2002.....	1.8	4.6	0.2	*	*	*	*	0.2	0.3	0.4
2001.....	1.6	4.2	0.1	*	*	*	0.2	0.3	0.3	0.5
2000.....	1.6	4.7	0.2	*	*	0.1	0.1	0.2	0.4	0.5
1999.....	1.5	5.7	*	0.1	*	0.1	0.2	0.2	0.3	0.5

All races 3/, male

2006.....	1.3	2.0	*	*	*	*	*	0.2	*	0.3
2005.....	1.2	2.5	*	*	*	*	*	*	0.2	0.2
2004.....	1.2	4.0	*	*	*	*	*	*	*	0.4
2003.....	1.3	3.6	*	*	*	*	*	*	0.2	0.4
2002.....	1.3	4.7	*	*	*	*	*	*	0.2	0.4
2001.....	1.2	3.9	*	*	*	*	*	*	0.2	0.3
2000.....	1.2	4.9	*	*	*	*	*	*	0.3	0.4
1999.....	1.2	5.9	*	*	*	*	*	*	*	0.4

All races 3/, female

2006.....	2.4	1.8	*	*	*	*	*	0.2	0.4	0.5
2005.....	2.3	2.1	*	*	*	*	*	0.3	0.4	0.6
2004.....	2.2	3.5	*	*	*	*	*	0.3	0.4	0.5
2003.....	2.3	2.8	*	*	*	0.2	*	0.4	0.5	0.6
2002.....	2.2	4.4	*	*	*	*	*	0.3	0.4	0.5
2001.....	2.0	4.5	*	*	*	*	0.2	0.4	0.4	0.6
2000.....	2.1	4.5	*	*	*	*	*	0.3	0.5	0.5
1999.....	1.8	5.5	*	*	*	*	0.3	0.3	0.5	0.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)										
White, both sexes										
2006 4/.....	1.9	1.3	*	*	*	*	0.1	0.2	0.2	0.3
2005 4/.....	1.8	2.0	*	*	*	*	*	0.2	0.2	0.3
2004 4/.....	1.7	3.2	*	*	*	*	*	0.2	0.2	0.4
2003 4/.....	1.8	2.5	*	*	*	0.1	*	0.2	0.3	0.4
2002.....	1.8	3.8	*	*	*	*	*	0.1	0.2	0.4
2001.....	1.7	3.6	*	*	*	*	*	0.2	0.3	0.4
2000.....	1.7	4.0	0.2	*	*	*	*	0.2	0.3	0.4
1999.....	1.5	5.1	*	*	*	0.2	0.2	0.2	0.3	0.4
White, male										
2006 4/.....	1.3	1.3	*	*	*	*	*	*	*	0.3
2005 4/.....	1.3	2.2	*	*	*	*	*	*	*	*
2004 4/.....	1.2	3.3	*	*	*	*	*	*	*	0.3
2003 4/.....	1.4	2.9	*	*	*	*	*	*	*	0.3
2002.....	1.3	4.0	*	*	*	*	*	*	*	0.3
2001.....	1.2	3.4	*	*	*	*	*	*	*	0.3
2000.....	1.2	4.2	*	*	*	*	*	*	*	0.4
1999.....	1.2	5.4	*	*	*	*	*	*	*	0.4
White, female										
2006 4/.....	2.4	1.3	*	*	*	*	*	*	*	0.4
2005 4/.....	2.3	1.7	*	*	*	*	*	0.3	0.4	0.4
2004 4/.....	2.2	3.0	*	*	*	*	*	*	*	0.4
2003 4/.....	2.3	2.0	*	*	*	*	*	0.3	0.4	0.5
2002.....	2.2	3.6	*	*	*	*	*	*	0.2	0.4
2001.....	2.1	3.8	*	*	*	*	*	0.3	0.4	0.6
2000.....	2.1	3.8	*	*	*	*	*	*	0.4	0.4
1999.....	1.8	4.8	*	*	*	*	*	*	0.4	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other non-pulmonary forms of heart disease (I30-I51)										
All races 3/, both sexes										
2006.....	52.7	5.8	0.8	0.4	0.6	1.3	2.3	3.3	4.7	6.1
2005.....	53.0	5.9	0.8	0.5	0.6	1.5	2.3	3.2	4.4	6.4
2004.....	52.2	6.2	0.9	0.3	0.6	1.3	2.2	2.9	4.1	6.3
2003.....	53.6	7.1	0.9	0.4	0.6	1.5	2.2	3.0	4.3	6.3
2002.....	54.0	7.1	0.8	0.4	0.6	1.6	2.0	3.1	4.3	6.6
2001.....	53.9	6.9	1.3	0.4	0.6	1.3	2.0	2.9	4.1	6.6
2000.....	54.1	7.1	0.9	0.4	0.7	1.5	2.2	2.9	4.1	6.1
1999.....	54.7	7.2	1.0	0.4	0.7	1.7	2.1	2.8	4.3	6.7
All races 3/, male										
2006.....	47.7	6.0	0.9	0.4	0.7	1.7	3.0	4.4	6.3	8.1
2005.....	47.8	6.3	0.8	0.4	0.7	2.1	3.1	4.4	6.0	8.4
2004.....	47.1	6.4	1.0	0.2	0.7	1.7	2.9	3.9	5.4	8.3
2003.....	48.3	7.7	0.8	0.4	0.8	1.9	2.9	3.7	5.7	8.2
2002.....	48.6	7.3	0.9	0.3	0.7	2.2	2.7	4.0	5.9	8.3
2001.....	48.0	6.9	1.3	0.3	0.7	1.7	2.7	3.7	5.3	8.8
2000.....	48.5	7.1	1.0	0.4	0.8	2.0	2.8	3.6	5.3	7.9
1999.....	49.1	7.3	1.1	0.4	0.8	2.2	2.6	3.6	5.6	9.0
All races 3/, female										
2006.....	57.6	5.6	0.7	0.3	0.5	0.9	1.6	2.1	3.0	4.2
2005.....	58.1	5.5	0.7	0.5	0.6	0.8	1.4	1.9	2.8	4.5
2004.....	57.1	5.9	0.9	0.4	0.5	0.9	1.3	1.8	2.8	4.4
2003.....	58.8	6.4	1.0	0.3	0.4	1.0	1.5	2.2	3.0	4.4
2002.....	59.3	6.9	0.8	0.4	0.4	0.9	1.2	2.2	2.7	4.9
2001.....	59.7	7.0	1.2	0.5	0.6	1.0	1.3	2.0	2.8	4.4
2000.....	59.6	7.2	0.7	0.3	0.5	1.0	1.5	2.1	2.9	4.3
1999.....	60.0	7.1	0.9	0.4	0.6	1.2	1.5	2.0	3.0	4.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other non-pulmonary forms of heart disease (I30-I51)										
White, both sexes										
2006 4/.....	56.7	4.7	0.7	0.3	0.6	1.1	2.0	2.7	3.7	4.9
2005 4/.....	56.8	4.8	0.6	0.3	0.5	1.2	1.9	2.7	3.5	5.2
2004 4/.....	55.9	5.4	0.7	0.3	0.5	1.1	2.0	2.3	3.3	5.3
2003 4/.....	57.4	5.7	0.8	0.3	0.5	1.2	1.9	2.3	3.5	5.2
2002.....	57.8	6.2	0.8	0.3	0.5	1.4	1.6	2.5	3.4	5.4
2001.....	57.6	5.7	1.0	0.4	0.6	1.1	1.6	2.3	3.3	5.3
2000.....	57.8	6.5	0.8	0.3	0.5	1.2	1.7	2.3	3.4	4.9
1999.....	58.1	5.9	0.9	0.3	0.6	1.5	1.7	2.2	3.6	5.4
White, male										
2006 4/.....	50.3	4.8	0.8	0.4	0.7	1.5	2.7	3.6	5.2	6.4
2005 4/.....	50.2	4.8	0.7	0.3	0.6	1.7	2.7	3.9	4.7	7.0
2004 4/.....	49.4	5.7	0.8	*	0.6	1.4	2.7	3.2	4.4	7.1
2003 4/.....	50.8	5.8	0.7	0.4	0.7	1.6	2.4	3.0	4.8	6.8
2002.....	51.1	6.4	0.9	0.3	0.6	1.9	2.3	3.2	4.6	6.9
2001.....	50.3	6.3	1.0	0.3	0.6	1.4	2.1	3.0	4.3	7.2
2000.....	50.8	6.4	1.0	0.4	0.7	1.6	2.2	2.9	4.5	6.4
1999.....	51.1	6.0	1.0	0.4	0.6	1.8	2.2	2.9	4.8	7.4
White, female										
2006 4/.....	63.0	4.5	0.6	0.3	0.5	0.7	1.3	1.6	2.2	3.4
2005 4/.....	63.2	4.7	0.4	0.3	0.4	0.7	1.2	1.4	2.1	3.4
2004 4/.....	62.2	5.2	0.7	0.3	0.4	0.7	1.2	1.4	2.2	3.3
2003 4/.....	63.9	5.6	0.9	0.3	0.3	0.8	1.3	1.5	2.2	3.6
2002.....	64.4	5.9	0.7	0.4	0.4	0.8	0.9	1.8	2.1	3.9
2001.....	64.8	5.0	1.0	0.4	0.5	0.8	1.0	1.6	2.2	3.4
2000.....	64.6	6.6	0.6	0.3	0.4	0.9	1.1	1.6	2.3	3.3
1999.....	64.9	5.8	0.8	0.3	0.5	1.1	1.2	1.6	2.4	3.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other non-pulmonary forms of heart disease (I30-I51)										
Black, both sexes										
2006 4/.....	44.9	10.9	1.2	*	1.1	2.4	4.2	7.2	11.6	15.5
2005 4/.....	45.8	11.0	1.5	1.0	1.1	2.9	4.4	6.9	11.2	15.5
2004 4/.....	45.2	8.9	2.1	0.8	1.2	2.7	3.4	6.5	9.7	14.5
2003 4/.....	46.4	13.0	1.7	0.7	1.0	2.7	4.7	7.5	10.3	14.3
2002.....	46.6	12.0	1.1	*	1.1	2.7	4.3	7.3	10.5	15.8
2001.....	47.0	13.4	2.7	0.7	1.0	2.5	4.7	6.8	10.0	15.2
2000.....	47.0	11.1	1.6	0.6	1.3	3.2	4.9	7.2	9.4	14.7
1999.....	48.2	12.9	1.8	0.7	1.2	3.2	4.6	6.8	9.8	16.2
Black, male										
2006 4/.....	46.0	11.5	*	*	1.3	3.1	4.7	9.5	15.6	21.5
2005 4/.....	46.0	13.0	*	*	1.1	4.4	5.9	9.2	15.8	20.1
2004 4/.....	45.8	9.2	2.2	*	1.2	3.8	4.6	8.8	12.4	18.1
2003 4/.....	46.4	16.1	1.8	*	1.2	3.6	6.5	9.4	12.2	19.0
2002.....	46.3	11.9	*	*	1.3	3.7	5.6	9.7	14.6	19.9
2001.....	46.9	10.8	3.0	*	*	3.3	6.4	9.3	13.2	20.4
2000.....	46.9	11.2	*	*	1.4	4.1	6.2	9.2	11.8	19.1
1999.....	49.2	13.0	2.0	*	1.4	4.5	5.9	9.0	12.7	21.9
Black, female										
2006 4/.....	43.9	10.2	*	*	*	1.8	3.6	5.0	8.0	10.2
2005 4/.....	45.6	8.9	1.9	*	*	1.4	2.9	4.8	7.0	11.4
2004 4/.....	44.6	8.6	2.0	*	1.2	1.6	2.2	4.3	7.2	11.3
2003 4/.....	46.5	9.9	1.6	*	*	1.8	3.0	5.7	8.5	10.1
2002.....	46.8	12.1	*	*	*	1.7	3.0	5.1	6.8	12.2
2001.....	47.1	16.1	2.3	*	*	1.8	3.1	4.5	7.1	10.6
2000.....	47.1	10.9	1.6	*	1.2	2.1	3.8	5.4	7.2	10.8
1999.....	47.3	12.8	1.6	*	*	1.8	3.4	4.8	7.2	11.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of endocardium (I34-I38)										
All races 3/, both sexes										
2006.....	7.1	*	*	*	*	*	0.2	0.2	0.4	0.4
2005.....	7.0	*	*	*	*	*	0.2	0.2	0.4	0.5
2004.....	6.9	*	*	*	*	0.1	0.1	0.2	0.4	0.5
2003.....	6.9	*	*	*	*	*	0.1	0.2	0.4	0.5
2002.....	6.9	*	*	*	*	*	0.1	0.3	0.3	0.6
2001.....	6.9	*	*	*	*	*	0.1	0.3	0.3	0.6
2000.....	7.0	*	*	*	*	*	0.2	0.2	0.3	0.6
1999.....	7.0	*	*	*	*	0.1	0.2	0.2	0.5	0.6
All races 3/, male										
2006.....	5.7	*	*	*	*	*	*	0.3	0.5	0.5
2005.....	5.5	*	*	*	*	*	0.2	0.4	0.4	0.6
2004.....	5.5	*	*	*	*	*	*	*	0.4	0.6
2003.....	5.4	*	*	*	*	*	0.2	*	0.4	0.6
2002.....	5.4	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	5.4	*	*	*	*	*	*	0.3	0.3	0.7
2000.....	5.5	*	*	*	*	*	*	0.3	0.3	0.7
1999.....	5.5	*	*	*	*	*	0.2	*	0.5	0.8
All races 3/, female										
2006.....	8.5	*	*	*	*	*	*	0.2	0.3	0.4
2005.....	8.6	*	*	*	*	*	*	*	0.4	0.4
2004.....	8.3	*	*	*	*	*	*	*	0.3	0.5
2003.....	8.3	*	*	*	*	*	*	0.3	0.4	0.4
2002.....	8.4	*	*	*	*	*	*	0.4	0.2	0.5
2001.....	8.2	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	8.5	*	*	*	*	*	*	0.2	0.4	0.6
1999.....	8.5	*	*	*	*	*	*	0.3	0.5	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of endocardium (I34-I38)										
White, both sexes										
2006 4/.....	8.2	*	*	*	*	*	0.2	0.2	0.4	0.4
2005 4/.....	8.0	*	*	*	*	*	0.2	0.2	0.4	0.4
2004 4/.....	7.8	*	*	*	*	*	0.1	0.1	0.4	0.5
2003 4/.....	7.8	*	*	*	*	*	*	0.2	0.3	0.5
2002.....	7.9	*	*	*	*	*	*	0.3	0.3	0.5
2001.....	7.8	*	*	*	*	*	*	0.2	0.3	0.5
2000.....	7.9	*	*	*	*	*	*	0.2	0.3	0.5
1999.....	7.9	*	*	*	*	0.1	0.2	0.2	0.5	0.6
White, male										
2006 4/.....	6.5	*	*	*	*	*	*	0.2	0.5	0.5
2005 4/.....	6.2	*	*	*	*	*	*	0.3	0.4	0.6
2004 4/.....	6.1	*	*	*	*	*	*	*	0.5	0.5
2003 4/.....	6.1	*	*	*	*	*	*	*	0.4	0.6
2002.....	6.1	*	*	*	*	*	*	*	0.4	0.6
2001.....	6.1	*	*	*	*	*	*	0.3	0.3	0.6
2000.....	6.1	*	*	*	*	*	*	*	0.3	0.6
1999.....	6.2	*	*	*	*	*	*	*	0.5	0.7
White, female										
2006 4/.....	9.8	*	*	*	*	*	*	*	0.3	0.3
2005 4/.....	9.8	*	*	*	*	*	*	*	0.3	0.3
2004 4/.....	9.5	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	9.5	*	*	*	*	*	*	*	0.3	0.4
2002.....	9.6	*	*	*	*	*	*	0.3	*	0.4
2001.....	9.5	*	*	*	*	*	*	*	0.3	0.4
2000.....	9.7	*	*	*	*	*	*	*	0.4	0.5
1999.....	9.6	*	*	*	*	*	*	*	0.5	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases of endocardium (I36-I38)										
All races 3/, both sexes										
2006.....	1.7	*	*	*	*	*	*	*	0.2	0.2
2005.....	1.7	*	*	*	*	*	*	0.1	0.2	0.2
2004.....	1.7	*	*	*	*	*	*	*	0.1	0.2
2003.....	1.7	*	*	*	*	*	*	0.1	0.1	0.2
2002.....	1.7	*	*	*	*	*	*	0.1	0.1	0.2
2001.....	1.6	*	*	*	*	*	*	*	*	0.2
2000.....	1.6	*	*	*	*	*	*	*	0.1	0.3
1999.....	1.6	*	*	*	*	*	*	*	0.2	0.2
All races 3/, male										
2006.....	1.4	*	*	*	*	*	*	*	0.2	0.2
2005.....	1.4	*	*	*	*	*	*	*	*	0.3
2004.....	1.4	*	*	*	*	*	*	*	*	0.3
2003.....	1.3	*	*	*	*	*	*	*	*	0.2
2002.....	1.3	*	*	*	*	*	*	*	*	0.3
2001.....	1.3	*	*	*	*	*	*	*	*	0.2
2000.....	1.2	*	*	*	*	*	*	*	*	0.3
1999.....	1.3	*	*	*	*	*	*	*	0.2	0.2
All races 3/, female										
2006.....	2.0	*	*	*	*	*	*	*	*	*
2005.....	2.1	*	*	*	*	*	*	*	*	*
2004.....	2.0	*	*	*	*	*	*	*	*	0.2
2003.....	2.1	*	*	*	*	*	*	*	*	*
2002.....	2.0	*	*	*	*	*	*	*	*	0.2
2001.....	1.9	*	*	*	*	*	*	*	*	0.2
2000.....	1.9	*	*	*	*	*	*	*	*	0.2
1999.....	2.0	*	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cardiomyopathy (I42)										
All races 3/, both sexes										
2006.....	8.3	2.8	0.3	*	0.2	0.4	0.8	1.3	1.9	2.4
2005.....	8.6	2.0	0.2	0.1	0.2	0.6	0.9	1.3	1.8	2.6
2004.....	8.7	2.8	0.2	0.1	0.2	0.5	0.8	1.2	1.6	2.7
2003.....	9.2	3.1	0.4	0.1	0.2	0.5	0.8	1.3	1.7	2.6
2002.....	9.6	2.9	0.2	0.1	0.2	0.6	0.8	1.4	1.8	2.8
2001.....	9.4	3.4	0.4	0.1	0.2	0.6	0.8	1.1	1.7	2.6
2000.....	9.7	3.1	0.3	0.1	0.3	0.5	0.9	1.2	1.8	2.6
1999.....	9.8	3.5	0.2	0.1	0.2	0.6	0.7	1.2	1.8	3.0
All races 3/, male										
2006.....	9.8	3.1	0.3	*	0.3	0.6	1.1	1.9	2.7	3.4
2005.....	10.2	1.8	*	*	0.3	0.8	1.3	1.9	2.6	3.5
2004.....	10.3	2.7	0.3	*	0.2	0.7	1.2	1.7	2.4	3.8
2003.....	10.8	3.4	0.3	*	0.2	0.6	1.1	1.8	2.4	3.7
2002.....	11.3	2.7	*	*	0.3	0.8	1.1	2.0	2.6	3.7
2001.....	11.0	2.9	0.5	*	0.2	0.7	1.1	1.5	2.4	3.8
2000.....	11.4	2.8	0.3	*	0.4	0.7	1.2	1.6	2.6	3.6
1999.....	11.5	3.1	*	*	0.3	0.9	0.9	1.6	2.4	4.3
All races 3/, female										
2006.....	6.8	2.5	*	*	*	0.2	0.5	0.6	1.0	1.4
2005.....	7.0	2.2	*	*	*	0.3	0.5	0.7	1.0	1.6
2004.....	7.2	3.0	*	*	0.2	0.3	0.4	0.6	0.9	1.7
2003.....	7.7	2.9	0.4	*	*	0.4	0.5	0.7	1.1	1.5
2002.....	8.0	3.1	*	*	*	0.3	0.4	0.8	1.0	2.0
2001.....	8.0	3.9	0.3	*	*	0.4	0.4	0.7	0.9	1.5
2000.....	8.0	3.4	*	*	*	0.3	0.6	0.8	1.1	1.6
1999.....	8.1	3.8	*	*	*	0.3	0.5	0.8	1.2	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cardiomyopathy (I42)										
White, both sexes										
2006 4/.....	8.2	2.5	0.2	*	0.1	0.4	0.7	1.0	1.4	1.8
2005 4/.....	8.6	1.8	0.2	*	0.2	0.4	0.7	1.0	1.3	1.9
2004 4/.....	8.7	2.7	0.2	*	0.2	0.4	0.7	0.9	1.2	2.1
2003 4/.....	9.2	3.0	0.3	*	0.1	0.4	0.7	0.9	1.3	1.9
2002.....	9.6	2.8	0.2	*	0.2	0.4	0.6	1.0	1.3	2.1
2001.....	9.3	2.8	0.3	*	0.2	0.4	0.6	0.8	1.3	2.0
2000.....	9.5	3.0	0.2	*	0.2	0.3	0.6	0.8	1.4	1.9
1999.....	9.6	3.1	0.2	*	0.2	0.4	0.5	0.9	1.4	2.3
White, male										
2006 4/.....	9.6	2.6	*	*	*	0.6	1.0	1.5	2.0	2.6
2005 4/.....	10.1	1.6	*	*	*	0.7	1.0	1.5	2.0	2.7
2004 4/.....	10.2	2.6	*	*	*	0.5	1.1	1.3	1.8	3.0
2003 4/.....	10.7	2.9	*	*	*	0.5	0.9	1.3	1.9	2.8
2002.....	11.2	2.4	*	*	0.3	0.6	0.9	1.4	1.9	2.8
2001.....	10.8	2.7	0.4	*	*	0.5	0.8	1.2	1.8	2.9
2000.....	11.1	2.7	*	*	0.3	0.5	0.9	1.1	1.9	2.7
1999.....	11.1	2.6	*	*	*	0.6	0.7	1.2	2.0	3.2
White, female										
2006 4/.....	6.8	2.3	*	*	*	*	0.4	0.4	0.7	1.1
2005 4/.....	7.0	2.0	*	*	*	*	0.4	0.4	0.7	1.1
2004 4/.....	7.2	2.8	*	*	*	0.3	0.3	0.4	0.6	1.2
2003 4/.....	7.8	3.1	*	*	*	*	0.4	0.4	0.7	1.1
2002.....	8.0	3.2	*	*	*	0.3	*	0.5	0.7	1.4
2001.....	7.9	2.9	*	*	*	0.3	0.3	0.5	0.7	1.1
2000.....	8.0	3.4	*	*	*	*	0.3	0.5	0.8	1.1
1999.....	8.1	3.5	*	*	*	0.3	0.3	0.5	0.8	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/ year 2/	Under 1 year 2/ years	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cardiomyopathy (I42)										
Black, both sexes										
2006 4/.....	10.8	4.3	*	*	*	0.7	1.6	3.1	5.3	6.5
2005 4/.....	11.3	*	*	*	*	1.2	2.3	3.2	5.2	7.1
2004 4/.....	11.2	*	*	*	*	1.1	1.5	2.8	4.6	7.1
2003 4/.....	11.9	3.5	*	*	*	1.1	1.9	3.8	4.7	7.0
2002.....	12.2	3.4	*	*	*	1.2	2.1	4.0	5.2	7.9
2001.....	12.4	6.8	0.9	*	*	1.3	2.1	3.2	4.6	7.1
2000.....	12.9	3.6	*	*	*	1.4	2.3	3.6	4.8	7.3
1999.....	13.2	5.1	*	*	0.6	1.7	1.9	3.2	4.4	8.0
Black, male										
2006 4/.....	13.3	*	*	*	*	*	2.0	4.6	7.5	9.3
2005 4/.....	14.1	*	*	*	*	1.7	3.1	4.4	7.9	9.6
2004 4/.....	13.7	*	*	*	*	1.8	2.2	3.8	6.8	9.7
2003 4/.....	14.5	*	*	*	*	*	2.6	5.3	6.3	9.9
2002.....	14.9	*	*	*	*	1.7	2.6	5.9	7.6	10.3
2001.....	14.9	*	*	*	*	1.7	2.9	4.1	6.7	9.9
2000.....	15.8	*	*	*	*	2.0	2.9	4.7	7.0	10.0
1999.....	16.4	*	*	*	*	2.7	2.5	4.5	5.8	11.9
Black, female										
2006 4/.....	8.4	*	*	*	*	*	*	1.7	3.4	3.9
2005 4/.....	8.7	*	*	*	*	*	1.4	2.0	2.8	4.8
2004 4/.....	8.9	*	*	*	*	*	*	1.8	2.7	4.7
2003 4/.....	9.5	*	*	*	*	*	*	2.4	3.2	4.5
2002.....	9.8	*	*	*	*	*	1.6	2.2	3.1	5.8
2001.....	10.2	8.8	*	*	*	*	*	2.3	2.7	4.5
2000.....	10.3	*	*	*	*	*	1.7	2.7	2.9	4.9
1999.....	10.3	*	*	*	*	*	*	2.0	3.1	4.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Conduction disorders and cardiac dysrhythmias (I44-I49)										
All races 3/, both sexes										
2006.....	12.6	0.9	*	*	0.2	0.3	0.6	0.7	1.0	1.3
2005.....	13.0	1.0	0.1	*	0.1	0.4	0.5	0.7	0.9	1.5
2004.....	12.8	0.8	0.1	*	0.2	0.3	0.6	0.7	0.9	1.3
2003.....	13.4	0.8	0.1	*	0.2	0.4	0.5	0.6	1.0	1.4
2002.....	13.6	1.4	0.1	0.1	0.1	0.4	0.5	0.7	0.9	1.3
2001.....	13.5	0.9	0.2	*	0.2	0.3	0.5	0.6	0.8	1.3
2000.....	13.6	1.0	0.2	*	0.2	0.4	0.5	0.6	0.8	1.3
1999.....	14.3	1.0	0.3	*	0.2	0.4	0.5	0.7	0.9	1.3
All races 3/, male										
2006.....	11.3	*	*	*	0.2	0.4	0.7	0.9	1.2	1.7
2005.....	11.6	1.1	*	*	0.2	0.6	0.7	0.9	1.1	1.8
2004.....	11.5	*	*	*	*	0.4	0.7	0.8	1.1	1.6
2003.....	12.0	*	*	*	0.2	0.5	0.6	0.6	1.1	1.7
2002.....	12.2	1.6	*	*	*	0.6	0.7	0.8	1.1	1.6
2001.....	12.0	*	*	*	0.2	0.4	0.7	0.7	1.0	1.6
2000.....	12.1	1.1	*	*	0.2	0.5	0.6	0.8	0.9	1.6
1999.....	12.9	1.2	0.3	*	0.2	0.5	0.6	0.8	1.1	1.5
All races 3/, female										
2006.....	13.9	1.0	*	*	*	0.2	0.5	0.5	0.9	1.0
2005.....	14.3	*	*	*	*	0.2	0.3	0.5	0.6	1.2
2004.....	14.1	*	*	*	*	0.2	0.4	0.5	0.8	1.0
2003.....	14.8	*	*	*	*	0.3	0.4	0.5	0.8	1.2
2002.....	15.1	1.2	*	*	*	0.2	0.3	0.5	0.7	1.1
2001.....	15.1	*	*	*	0.2	*	0.3	0.4	0.7	1.0
2000.....	15.1	*	*	*	*	0.3	0.4	0.4	0.7	1.0
1999.....	15.7	*	*	*	*	0.4	0.4	0.5	0.7	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Conduction disorders and cardiac dysrhythmias (I44-I49)										
White, both sexes										
2006 4/.....	13.3	0.7	*	*	0.2	0.3	0.5	0.6	0.8	1.1
2005 4/.....	13.7	0.8	*	*	*	0.4	0.5	0.7	0.7	1.4
2004 4/.....	13.5	0.7	*	*	0.1	0.3	0.5	0.5	0.8	1.1
2003 4/.....	14.2	*	*	*	0.1	0.3	0.4	0.5	0.8	1.3
2002.....	14.4	1.1	*	*	0.1	0.4	0.4	0.6	0.7	1.1
2001.....	14.3	0.9	0.2	*	0.2	0.3	0.4	0.5	0.7	1.0
2000.....	14.4	1.0	0.2	*	0.1	0.4	0.5	0.5	0.7	1.2
1999.....	15.2	0.8	0.3	*	0.2	0.4	0.4	0.6	0.8	1.1
White, male										
2006 4/.....	11.7	*	*	*	*	0.4	0.7	0.8	0.9	1.4
2005 4/.....	12.0	*	*	*	*	0.5	0.6	0.9	0.9	1.6
2004 4/.....	11.9	*	*	*	*	0.4	0.6	0.6	0.9	1.3
2003 4/.....	12.5	*	*	*	*	0.4	0.5	0.5	1.1	1.5
2002.....	12.7	1.6	*	*	*	0.7	0.6	0.7	0.9	1.3
2001.....	12.4	*	*	*	*	0.4	0.6	0.6	0.9	1.3
2000.....	12.7	*	*	*	*	0.5	0.6	0.7	0.8	1.5
1999.....	13.5	*	0.3	*	*	0.5	0.5	0.7	1.0	1.3
White, female										
2006 4/.....	14.9	*	*	*	*	*	0.3	0.4	0.7	0.7
2005 4/.....	15.4	*	*	*	*	0.3	0.3	0.4	0.5	1.1
2004 4/.....	15.0	*	*	*	*	*	0.4	0.4	0.7	0.9
2003 4/.....	15.9	*	*	*	*	0.3	0.3	0.4	0.6	1.1
2002.....	16.1	*	*	*	*	*	*	0.4	0.6	0.9
2001.....	16.1	*	*	*	*	*	*	0.4	0.5	0.8
2000.....	16.1	*	*	*	*	*	0.4	0.3	0.5	0.9
1999.....	16.9	*	*	*	*	0.4	0.4	0.4	0.5	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Conduction disorders and cardiac dysrhythmias (I44-I49)										
Black, both sexes										
2006 4/.....	12.1	*	*	*	*	0.6	1.1	1.5	2.5	3.5
2005 4/.....	12.0	*	*	*	*	0.7	0.9	1.2	2.1	3.0
2004 4/.....	12.1	*	*	*	*	*	0.8	1.4	1.9	2.9
2003 4/.....	12.4	*	*	*	*	0.6	0.8	1.4	2.0	2.4
2002.....	12.5	*	*	*	*	*	0.9	1.3	2.0	2.9
2001.....	12.6	*	*	*	*	*	1.0	1.0	1.9	3.2
2000.....	12.3	*	*	*	*	*	0.8	1.1	1.7	2.5
1999.....	13.0	*	*	*	*	*	1.1	1.4	2.0	3.1
Black, male										
2006 4/.....	12.1	*	*	*	*	*	*	2.0	2.9	4.3
2005 4/.....	11.6	*	*	*	*	1.2	*	*	2.8	3.7
2004 4/.....	11.9	*	*	*	*	*	*	1.5	1.9	3.6
2003 4/.....	12.0	*	*	*	*	*	*	1.6	1.7	2.6
2002.....	11.9	*	*	*	*	*	*	*	2.5	3.4
2001.....	12.2	*	*	*	*	*	1.5	*	2.1	3.8
2000.....	11.7	*	*	*	*	*	*	*	1.7	3.3
1999.....	12.9	*	*	*	*	*	*	1.6	2.2	3.3
Black, female										
2006 4/.....	12.2	*	*	*	*	*	*	*	2.1	2.9
2005 4/.....	12.4	*	*	*	*	*	*	*	1.5	2.4
2004 4/.....	12.3	*	*	*	*	*	*	*	1.9	2.4
2003 4/.....	12.8	*	*	*	*	*	*	*	2.2	2.2
2002.....	13.1	*	*	*	*	*	*	*	1.5	2.4
2001.....	12.9	*	*	*	*	*	*	*	1.6	2.7
2000.....	12.8	*	*	*	*	*	*	*	1.7	1.8
1999.....	13.1	*	*	*	*	*	*	*	1.8	2.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
All races 3/, both sexes										
2006.....	45.8	3.4	0.3	0.2	0.2	0.3	0.6	0.9	1.7	3.4
2005.....	48.4	3.1	0.4	0.3	0.2	0.4	0.6	1.0	1.8	3.3
2004.....	51.1	3.1	0.3	0.2	0.2	0.3	0.7	1.1	1.8	3.4
2003.....	54.2	2.5	0.3	0.1	0.2	0.4	0.6	1.0	1.9	3.6
2002.....	56.4	2.9	0.3	0.2	0.3	0.3	0.6	1.0	1.8	3.6
2001.....	57.4	2.7	0.4	0.2	0.2	0.3	0.7	1.1	1.9	3.8
2000.....	59.6	3.3	0.3	0.1	0.2	0.3	0.7	1.1	1.9	3.9
1999.....	60.0	2.7	0.3	0.1	0.2	0.3	0.6	1.0	1.8	4.0
All races 3/, male										
2006.....	37.0	3.9	0.3	0.3	0.3	0.3	0.7	1.0	1.8	3.7
2005.....	38.8	3.5	0.5	0.3	0.3	0.3	0.6	1.1	1.9	3.4
2004.....	40.7	3.4	0.3	*	0.2	0.4	0.7	1.1	1.7	3.7
2003.....	42.9	2.8	0.3	*	0.2	0.5	0.6	1.0	2.0	3.8
2002.....	44.2	3.2	0.4	*	0.3	0.3	0.7	1.0	1.7	3.6
2001.....	45.2	3.1	0.3	0.2	0.2	0.3	0.7	1.1	2.0	3.9
2000.....	46.9	3.8	*	*	0.2	0.4	0.7	1.1	1.9	3.8
1999.....	47.1	3.3	0.3	*	*	0.3	0.6	1.0	1.9	3.9
All races 3/, female										
2006.....	54.4	2.9	0.4	*	0.2	0.4	0.6	0.8	1.6	3.1
2005.....	57.8	2.6	0.3	0.3	*	0.4	0.6	0.8	1.6	3.2
2004.....	61.2	2.8	*	*	0.2	0.3	0.7	1.0	1.8	3.1
2003.....	65.1	2.2	0.3	*	*	0.4	0.7	0.9	1.8	3.3
2002.....	68.2	2.5	0.3	*	0.3	*	0.5	0.9	1.9	3.7
2001.....	69.2	2.3	0.4	*	0.2	0.3	0.6	1.0	1.9	3.6
2000.....	71.8	2.7	0.4	*	0.3	0.3	0.7	1.0	1.9	4.0
1999.....	72.3	2.1	0.3	*	0.2	0.3	0.6	1.0	1.8	4.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
White, both sexes										
2006 4/.....	47.9	2.8	0.3	0.2	0.2	0.4	0.6	0.8	1.4	2.6
2005 4/.....	50.7	2.5	0.4	0.2	0.2	0.3	0.5	0.7	1.5	2.7
2004 4/.....	53.7	2.5	0.2	0.2	0.2	0.3	0.6	0.9	1.4	2.9
2003 4/.....	57.0	2.0	0.2	0.1	0.2	0.4	0.5	0.8	1.6	2.7
2002.....	59.5	2.3	0.3	0.2	0.2	0.2	0.6	0.8	1.5	2.9
2001.....	60.5	1.9	0.3	0.2	0.2	0.3	0.5	0.9	1.7	3.1
2000.....	62.8	2.5	0.3	0.1	0.2	0.3	0.6	1.0	1.6	3.0
1999.....	63.3	2.3	0.3	0.1	0.2	0.3	0.6	0.9	1.5	3.2
White, male										
2006 4/.....	37.7	2.9	*	*	0.3	0.4	0.7	0.9	1.6	2.8
2005 4/.....	39.7	2.8	0.4	*	*	0.2	0.5	0.8	1.6	2.8
2004 4/.....	41.8	2.8	*	*	*	0.3	0.6	1.1	1.4	3.1
2003 4/.....	44.2	2.0	*	*	*	0.4	0.6	0.8	1.7	2.8
2002.....	45.7	2.6	0.4	*	*	0.3	0.7	1.0	1.4	2.8
2001.....	46.6	2.2	*	*	*	0.3	0.7	0.9	1.8	3.1
2000.....	48.4	3.0	*	*	*	0.4	0.7	1.1	1.6	3.0
1999.....	48.7	2.8	*	*	*	0.3	0.6	0.9	1.7	3.2
White, female										
2006 4/.....	57.9	2.6	0.4	*	*	0.3	0.5	0.7	1.3	2.4
2005 4/.....	61.6	2.1	*	*	*	0.4	0.5	0.7	1.4	2.7
2004 4/.....	65.3	2.2	*	*	*	0.3	0.6	0.7	1.5	2.7
2003 4/.....	69.5	2.0	*	*	*	0.3	0.5	0.8	1.5	2.5
2002.....	73.0	2.0	*	*	0.3	*	0.5	0.7	1.6	2.9
2001.....	74.0	1.6	0.4	*	*	0.3	0.4	0.9	1.6	3.0
2000.....	76.9	2.0	*	*	0.3	*	0.6	0.9	1.5	3.0
1999.....	77.6	1.9	*	*	*	*	0.6	0.9	1.4	3.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Cerebrovascular diseases (I60-I69)										
Black, both sexes										
2006 4/.....	43.1	7.2	*	*	*	*	1.0	1.7	3.7	8.4
2005 4/.....	44.9	6.0	*	*	*	*	1.0	2.2	3.6	6.8
2004 4/.....	46.9	6.0	*	*	*	*	1.0	2.2	4.1	7.0
2003 4/.....	49.3	4.9	*	*	*	0.7	1.2	2.0	4.1	8.8
2002.....	50.0	5.5	*	*	*	*	0.8	1.9	4.2	8.2
2001.....	51.1	6.6	*	*	*	*	1.3	2.2	3.6	8.1
2000.....	52.5	7.5	*	*	*	*	1.1	1.5	4.3	9.6
1999.....	52.2	4.6	*	*	*	*	0.8	1.8	3.6	9.1
Black, male										
2006 4/.....	39.3	9.5	*	*	*	*	*	1.9	3.7	9.8
2005 4/.....	40.3	6.8	*	*	*	*	*	2.6	4.1	7.1
2004 4/.....	41.5	6.8	*	*	*	*	*	1.7	4.1	8.5
2003 4/.....	43.2	5.9	*	*	*	*	*	2.6	4.1	9.7
2002.....	43.5	5.8	*	*	*	*	*	1.7	4.0	7.5
2001.....	44.6	8.1	*	*	*	*	*	2.8	3.4	8.8
2000.....	46.1	8.6	*	*	*	*	*	*	4.1	9.2
1999.....	45.9	*	*	*	*	*	*	1.9	3.7	8.9
Black, female										
2006 4/.....	46.5	*	*	*	*	*	*	1.5	3.8	7.1
2005 4/.....	49.1	*	*	*	*	*	*	1.9	3.2	6.5
2004 4/.....	51.9	*	*	*	*	*	*	2.6	4.0	5.7
2003 4/.....	54.8	*	*	*	*	*	1.6	1.5	4.1	8.0
2002.....	55.8	*	*	*	*	*	*	2.2	4.3	8.9
2001.....	56.9	*	*	*	*	*	1.7	1.7	3.8	7.4
2000.....	58.3	*	*	*	*	*	*	1.6	4.6	10.0
1999.....	57.9	*	*	*	*	*	*	1.8	3.6	9.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Subarachnoid hemorrhage (I60)										
All races 3/, both sexes										
2006.....	2.0	*	*	*	*	0.1	0.2	0.3	0.6	1.0
2005.....	2.1	*	*	*	*	0.1	0.3	0.3	0.7	1.1
2004.....	2.1	*	*	*	*	*	0.2	0.3	0.6	1.2
2003.....	2.1	*	*	*	*	*	0.2	0.4	0.6	1.2
2002.....	2.1	*	*	*	*	*	0.2	0.3	0.7	1.3
2001.....	2.2	*	*	*	*	0.1	0.2	0.4	0.7	1.3
2000.....	2.3	*	*	*	*	*	0.3	0.4	0.7	1.3
1999.....	2.3	*	*	*	*	0.1	0.2	0.4	0.7	1.6
All races 3/, male										
2006.....	1.4	*	*	*	*	*	0.2	0.3	0.6	0.9
2005.....	1.5	*	*	*	*	*	0.2	0.4	0.6	0.9
2004.....	1.5	*	*	*	*	*	0.2	0.3	0.6	1.0
2003.....	1.5	*	*	*	*	*	0.2	0.4	0.7	1.1
2002.....	1.5	*	*	*	*	*	0.3	0.3	0.6	1.1
2001.....	1.5	*	*	*	*	*	0.2	0.4	0.7	1.0
2000.....	1.6	*	*	*	*	*	0.3	0.4	0.7	1.0
1999.....	1.6	*	*	*	*	*	*	0.4	0.7	1.3
All races 3/, female										
2006.....	2.5	*	*	*	*	*	0.2	0.3	0.5	1.1
2005.....	2.6	*	*	*	*	*	0.3	0.2	0.7	1.4
2004.....	2.6	*	*	*	*	*	0.3	0.3	0.6	1.4
2003.....	2.6	*	*	*	*	*	*	0.4	0.6	1.3
2002.....	2.8	*	*	*	*	*	0.2	0.4	0.8	1.5
2001.....	2.8	*	*	*	*	*	0.2	0.3	0.7	1.5
2000.....	3.0	*	*	*	*	*	0.3	0.4	0.8	1.6
1999.....	3.0	*	*	*	*	*	0.2	0.4	0.8	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Subarachnoid hemorrhage (I60)										
White, both sexes										
2006 4/.....	1.9	*	*	*	*	0.1	0.2	0.3	0.5	0.8
2005 4/.....	2.0	*	*	*	*	0.1	0.2	0.2	0.6	1.0
2004 4/.....	2.1	*	*	*	*	*	0.2	0.3	0.6	1.1
2003 4/.....	2.0	*	*	*	*	*	0.2	0.3	0.6	1.0
2002.....	2.1	*	*	*	*	*	0.2	0.3	0.5	1.1
2001.....	2.1	*	*	*	*	*	0.2	0.3	0.6	1.1
2000.....	2.3	*	*	*	*	*	0.3	0.5	0.7	1.2
1999.....	2.3	*	*	*	*	*	0.2	0.4	0.7	1.4
White, male										
2006 4/.....	1.4	*	*	*	*	*	0.3	0.4	0.6	0.7
2005 4/.....	1.5	*	*	*	*	*	0.2	0.3	0.5	0.8
2004 4/.....	1.5	*	*	*	*	*	*	0.3	0.6	0.9
2003 4/.....	1.5	*	*	*	*	*	0.2	0.3	0.7	0.9
2002.....	1.5	*	*	*	*	*	0.3	0.3	0.4	1.0
2001.....	1.5	*	*	*	*	*	0.3	0.4	0.6	0.9
2000.....	1.6	*	*	*	*	*	0.3	0.5	0.7	1.0
1999.....	1.6	*	*	*	*	*	*	0.4	0.7	1.2
White, female										
2006 4/.....	2.4	*	*	*	*	*	*	0.3	0.5	1.0
2005 4/.....	2.6	*	*	*	*	*	*	*	0.7	1.2
2004 4/.....	2.6	*	*	*	*	*	*	*	0.5	1.4
2003 4/.....	2.6	*	*	*	*	*	*	0.4	0.6	1.1
2002.....	2.7	*	*	*	*	*	*	0.3	0.6	1.3
2001.....	2.7	*	*	*	*	*	*	*	0.7	1.3
2000.....	2.9	*	*	*	*	*	0.3	0.4	0.7	1.4
1999.....	2.9	*	*	*	*	*	*	0.3	0.7	1.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Subarachnoid hemorrhage (I60)										
Black, both sexes										
2006 4/.....	2.1	*	*	*	*	*	*	*	0.9	1.9
2005 4/.....	2.3	*	*	*	*	*	*	*	1.2	1.8
2004 4/.....	2.2	*	*	*	*	*	*	*	1.1	1.9
2003 4/.....	2.3	*	*	*	*	*	*	0.8	0.9	2.6
2002.....	2.4	*	*	*	*	*	*	*	1.4	2.2
2001.....	2.5	*	*	*	*	*	*	*	1.0	2.4
2000.....	2.7	*	*	*	*	*	*	*	1.3	2.5
1999.....	2.6	*	*	*	*	*	*	*	1.2	2.7
Black, male										
2006 4/.....	1.5	*	*	*	*	*	*	*	*	2.0
2005 4/.....	1.6	*	*	*	*	*	*	*	*	*
2004 4/.....	1.6	*	*	*	*	*	*	*	*	2.0
2003 4/.....	1.6	*	*	*	*	*	*	*	*	2.2
2002.....	1.6	*	*	*	*	*	*	*	*	1.5
2001.....	1.7	*	*	*	*	*	*	*	*	1.7
2000.....	1.7	*	*	*	*	*	*	*	*	1.6
1999.....	1.8	*	*	*	*	*	*	*	*	2.2
Black, female										
2006 4/.....	2.7	*	*	*	*	*	*	*	*	1.8
2005 4/.....	2.9	*	*	*	*	*	*	*	*	2.3
2004 4/.....	2.8	*	*	*	*	*	*	*	*	1.8
2003 4/.....	2.9	*	*	*	*	*	*	*	*	2.9
2002.....	3.1	*	*	*	*	*	*	*	1.6	2.9
2001.....	3.2	*	*	*	*	*	*	*	*	3.0
2000.....	3.5	*	*	*	*	*	*	*	1.9	3.3
1999.....	3.3	*	*	*	*	*	*	*	1.4	3.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intracerebral and other intracranial hemorrhage (I61-I62)										
All races 3/, both sexes										
2006.....	9.1	1.6	*	*	*	0.1	0.2	0.3	0.7	1.3
2005.....	9.4	1.1	*	*	*	*	0.2	0.3	0.6	1.2
2004.....	9.4	1.3	*	*	*	0.1	0.3	0.4	0.6	1.2
2003.....	9.4	0.8	*	*	*	0.2	0.3	0.4	0.6	1.3
2002.....	9.4	1.0	*	*	*	0.1	0.2	0.3	0.6	1.3
2001.....	9.4	1.0	*	*	*	0.1	0.3	0.4	0.7	1.5
2000.....	9.7	1.1	*	*	*	0.1	0.3	0.3	0.7	1.5
1999.....	9.1	1.1	*	*	*	0.1	0.2	0.3	0.6	1.5
All races 3/, male										
2006.....	8.7	1.8	*	*	*	*	0.3	0.3	0.7	1.7
2005.....	9.0	1.3	*	*	*	*	0.2	0.3	0.8	1.4
2004.....	8.9	1.4	*	*	*	*	0.3	0.5	0.6	1.6
2003.....	8.9	*	*	*	*	0.2	0.3	0.4	0.7	1.6
2002.....	8.7	1.1	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	8.8	1.2	*	*	*	0.2	0.3	0.5	0.8	1.8
2000.....	9.1	1.3	*	*	*	*	0.3	0.4	0.8	1.7
1999.....	8.5	1.3	*	*	*	*	0.2	0.3	0.7	1.7
All races 3/, female										
2006.....	9.5	1.4	*	*	*	*	0.2	0.3	0.6	1.0
2005.....	9.8	*	*	*	*	*	0.2	0.3	0.4	1.0
2004.....	9.8	1.3	*	*	*	*	0.2	0.4	0.6	0.8
2003.....	9.8	*	*	*	*	*	0.3	0.3	0.6	1.0
2002.....	10.0	1.0	*	*	*	*	*	0.3	0.6	1.3
2001.....	9.9	*	*	*	*	*	0.3	0.4	0.7	1.1
2000.....	10.4	*	*	*	*	*	0.3	0.2	0.6	1.3
1999.....	9.8	*	*	*	*	*	*	0.4	0.5	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intracerebral and other intracranial hemorrhage (I61-I62)										
White, both sexes										
2006 4/.....	9.1	1.4	*	*	*	0.1	0.2	0.3	0.5	0.9
2005 4/.....	9.4	1.0	*	*	*	*	0.2	0.2	0.5	0.9
2004 4/.....	9.4	1.0	*	*	*	*	0.2	0.3	0.5	0.9
2003 4/.....	9.3	*	*	*	*	0.1	0.2	0.3	0.5	0.9
2002.....	9.5	0.9	*	*	*	*	0.2	0.3	0.5	1.0
2001.....	9.4	0.7	*	*	*	0.1	0.3	0.3	0.6	1.1
2000.....	9.8	0.7	*	*	*	*	0.3	0.3	0.5	1.0
1999.....	9.2	0.8	*	*	*	*	0.2	0.3	0.4	1.1
White, male										
2006 4/.....	8.5	1.4	*	*	*	*	*	0.3	0.5	1.2
2005 4/.....	8.8	1.2	*	*	*	*	*	0.2	0.6	1.1
2004 4/.....	8.8	*	*	*	*	*	0.3	0.4	0.5	1.2
2003 4/.....	8.6	*	*	*	*	*	0.2	0.3	0.5	1.1
2002.....	8.7	*	*	*	*	*	0.3	0.4	0.5	1.0
2001.....	8.6	*	*	*	*	*	0.3	0.4	0.7	1.4
2000.....	8.9	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	8.3	*	*	*	*	*	*	0.3	0.6	1.2
White, female										
2006 4/.....	9.7	1.3	*	*	*	*	*	0.3	0.5	0.7
2005 4/.....	10.0	*	*	*	*	*	*	*	0.3	0.8
2004 4/.....	10.1	*	*	*	*	*	*	*	0.4	0.6
2003 4/.....	10.0	*	*	*	*	*	*	*	0.5	0.8
2002.....	10.2	*	*	*	*	*	*	*	0.5	0.9
2001.....	10.1	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	10.7	*	*	*	*	*	*	*	0.4	0.9
1999.....	10.0	*	*	*	*	*	*	0.3	0.3	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intracerebral and other intracranial hemorrhage (I61-I62)										
Black, both sexes										
2006 4/.....	9.9	3.1	*	*	*	*	*	0.7	1.5	3.7
2005 4/.....	10.1	*	*	*	*	*	*	0.8	1.5	2.8
2004 4/.....	10.2	*	*	*	*	*	*	1.1	1.6	3.0
2003 4/.....	10.5	*	*	*	*	*	*	0.9	1.5	3.2
2002.....	9.7	*	*	*	*	*	*	*	1.7	3.1
2001.....	10.0	*	*	*	*	*	*	1.1	1.6	3.3
2000.....	10.3	*	*	*	*	*	*	*	2.0	4.4
1999.....	9.5	*	*	*	*	*	*	*	1.4	4.4
Black, male										
2006 4/.....	10.5	*	*	*	*	*	*	*	1.7	4.9
2005 4/.....	10.8	*	*	*	*	*	*	*	1.9	3.6
2004 4/.....	10.7	*	*	*	*	*	*	*	1.7	4.6
2003 4/.....	11.0	*	*	*	*	*	*	*	1.6	4.0
2002.....	10.1	*	*	*	*	*	*	*	2.0	3.1
2001.....	10.3	*	*	*	*	*	*	*	1.5	4.2
2000.....	10.9	*	*	*	*	*	*	*	2.2	5.2
1999.....	9.8	*	*	*	*	*	*	*	1.5	5.0
Black, female										
2006 4/.....	9.4	*	*	*	*	*	*	*	*	2.6
2005 4/.....	9.6	*	*	*	*	*	*	*	*	2.1
2004 4/.....	9.8	*	*	*	*	*	*	*	1.4	1.6
2003 4/.....	10.1	*	*	*	*	*	*	*	1.4	2.5
2002.....	9.4	*	*	*	*	*	*	*	1.4	3.2
2001.....	9.8	*	*	*	*	*	*	*	1.6	2.4
2000.....	9.8	*	*	*	*	*	*	*	1.7	3.8
1999.....	9.2	*	*	*	*	*	*	*	1.4	4.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Stroke, not specified as hemorrhage or infarction (I64)										
All races 3/, both sexes										
2006.....	24.2	*	*	*	*	*	*	0.1	0.3	0.6
2005.....	25.1	*	*	*	*	*	*	0.1	0.2	0.5
2004.....	26.0	*	*	*	*	*	*	0.1	0.3	0.6
2003.....	28.4	*	*	*	*	*	*	0.1	0.4	0.6
2002.....	30.1	*	*	*	*	*	*	0.1	0.3	0.6
2001.....	31.1	*	*	*	*	*	*	0.1	0.2	0.5
2000.....	32.1	*	*	*	*	*	*	0.2	0.2	0.5
1999.....	32.6	*	*	*	*	*	*	0.1	0.2	0.4
All races 3/, male										
2006.....	18.5	*	*	*	*	*	*	*	0.3	0.7
2005.....	19.2	*	*	*	*	*	*	*	0.2	0.5
2004.....	19.7	*	*	*	*	*	*	*	0.2	0.6
2003.....	21.5	*	*	*	*	*	*	*	0.4	0.7
2002.....	22.6	*	*	*	*	*	*	*	*	0.7
2001.....	23.5	*	*	*	*	*	*	*	*	0.5
2000.....	24.2	*	*	*	*	*	*	*	0.3	0.6
1999.....	24.7	*	*	*	*	*	*	*	0.2	0.4
All races 3/, female										
2006.....	29.8	*	*	*	*	*	*	*	0.3	0.5
2005.....	30.9	*	*	*	*	*	*	*	0.2	0.5
2004.....	32.2	*	*	*	*	*	*	*	0.4	0.5
2003.....	35.0	*	*	*	*	*	*	*	0.3	0.5
2002.....	37.4	*	*	*	*	*	*	*	0.4	0.5
2001.....	38.5	*	*	*	*	*	*	*	0.3	0.5
2000.....	39.7	*	*	*	*	*	*	*	*	0.5
1999.....	40.3	*	*	*	*	*	*	*	0.3	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Stroke, not specified as hemorrhage or infarction (I64)										
White, both sexes										
2006 4/.....	25.6	*	*	*	*	*	*	*	0.2	0.5
2005 4/.....	26.6	*	*	*	*	*	*	*	0.2	0.4
2004 4/.....	27.6	*	*	*	*	*	*	*	0.2	0.4
2003 4/.....	30.2	*	*	*	*	*	*	*	0.2	0.4
2002.....	32.1	*	*	*	*	*	*	*	0.2	0.4
2001.....	33.1	*	*	*	*	*	*	*	0.2	0.4
2000.....	34.2	*	*	*	*	*	*	*	0.2	0.4
1999.....	34.8	*	*	*	*	*	*	*	0.2	0.3
White, male										
2006 4/.....	19.0	*	*	*	*	*	*	*	0.3	0.6
2005 4/.....	19.8	*	*	*	*	*	*	*	*	0.4
2004 4/.....	20.5	*	*	*	*	*	*	*	*	0.5
2003 4/.....	22.4	*	*	*	*	*	*	*	0.3	0.5
2002.....	23.5	*	*	*	*	*	*	*	*	0.5
2001.....	24.4	*	*	*	*	*	*	*	*	0.4
2000.....	25.2	*	*	*	*	*	*	*	0.3	0.4
1999.....	25.8	*	*	*	*	*	*	*	*	0.3
White, female										
2006 4/.....	32.1	*	*	*	*	*	*	*	*	0.3
2005 4/.....	33.3	*	*	*	*	*	*	*	*	0.4
2004 4/.....	34.7	*	*	*	*	*	*	*	0.3	0.3
2003 4/.....	37.8	*	*	*	*	*	*	*	*	0.3
2002.....	40.4	*	*	*	*	*	*	*	0.3	0.3
2001.....	41.6	*	*	*	*	*	*	*	*	0.4
2000.....	43.0	*	*	*	*	*	*	*	*	0.3
1999.....	43.6	*	*	*	*	*	*	*	0.3	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of arteries, arterioles and capillaries (I70-I78)										
All races 3/, both sexes										
2006.....	10.4	0.5	*	*	*	0.1	0.2	0.3	0.4	0.8
2005.....	12.0	*	*	*	*	*	0.2	0.3	0.5	0.8
2004.....	12.1	*	*	*	*	*	0.2	0.3	0.5	0.9
2003.....	13.0	*	*	*	*	0.2	0.2	0.3	0.5	0.8
2002.....	13.4	*	*	*	*	0.2	0.2	0.3	0.6	0.8
2001.....	13.8	*	*	*	*	0.1	0.2	0.4	0.5	0.9
2000.....	14.4	*	*	*	*	*	0.2	0.3	0.5	0.8
1999.....	14.6	*	*	*	*	*	0.2	0.3	0.5	0.8
All races 3/, male										
2006.....	10.3	*	*	*	*	*	0.3	0.5	0.6	1.2
2005.....	11.6	*	*	*	*	*	0.2	0.4	0.7	1.2
2004.....	11.6	*	*	*	*	*	0.3	0.5	0.7	1.2
2003.....	12.6	*	*	*	*	*	0.3	0.4	0.6	1.2
2002.....	13.0	*	*	*	*	*	0.3	0.3	0.7	0.8
2001.....	13.3	*	*	*	*	*	0.2	0.4	0.7	1.3
2000.....	14.1	*	*	*	*	*	0.3	0.4	0.7	0.9
1999.....	14.1	*	*	*	*	*	0.2	0.4	0.7	1.1
All races 3/, female										
2006.....	10.5	*	*	*	*	*	*	*	0.3	0.5
2005.....	12.3	*	*	*	*	*	*	0.3	0.3	0.4
2004.....	12.6	*	*	*	*	*	*	*	0.4	0.6
2003.....	13.3	*	*	*	*	*	*	*	0.3	0.5
2002.....	13.8	*	*	*	*	*	*	0.2	0.4	0.7
2001.....	14.4	*	*	*	*	*	*	0.3	0.3	0.6
2000.....	14.6	*	*	*	*	*	*	*	0.4	0.6
1999.....	15.2	*	*	*	*	*	*	*	0.3	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other disorders of circulatory system (I80-I99)										
All races 3/, both sexes										
2006.....	1.3	0.8	*	*	*	*	0.2	0.2	0.4	0.5
2005.....	1.6	0.6	*	*	*	*	0.1	0.2	0.4	0.6
2004.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.5
2003.....	1.6	0.9	*	*	*	*	0.2	0.3	0.4	0.6
2002.....	1.6	0.7	*	*	*	*	0.1	0.2	0.5	0.6
2001.....	1.6	0.5	*	*	*	*	0.1	0.2	0.4	0.4
2000.....	1.6	0.7	*	*	*	*	0.1	0.3	0.4	0.6
1999.....	1.4	0.6	*	*	*	*	*	0.2	0.3	0.4
All races 3/, male										
2006.....	1.2	*	*	*	*	*	*	*	0.4	0.6
2005.....	1.5	*	*	*	*	*	*	0.2	0.4	0.5
2004.....	1.4	*	*	*	*	*	*	0.3	0.4	0.5
2003.....	1.4	1.0	*	*	*	*	*	0.2	0.4	0.5
2002.....	1.4	*	*	*	*	*	*	*	0.5	0.6
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.5
2000.....	1.4	*	*	*	*	*	*	*	0.4	0.6
1999.....	1.3	*	*	*	*	*	*	*	0.3	0.5
All races 3/, female										
2006.....	1.5	*	*	*	*	*	0.2	*	0.4	0.5
2005.....	1.8	*	*	*	*	*	*	0.3	0.4	0.7
2004.....	1.7	*	*	*	*	*	*	0.3	0.4	0.5
2003.....	1.9	*	*	*	*	*	0.2	0.3	0.5	0.6
2002.....	1.9	*	*	*	*	*	*	0.3	0.4	0.6
2001.....	1.9	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.8	*	*	*	*	*	*	0.4	0.4	0.5
1999.....	1.6	*	*	*	*	*	*	*	0.3	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other disorders of circulatory system (I80-I99)

White, both sexes

2006 4/.....	1.3	0.9	*	*	*	*	*	0.1	0.3	0.5
2005 4/.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5
2004 4/.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.4
2003 4/.....	1.6	0.9	*	*	*	*	0.1	0.2	0.3	0.5
2002.....	1.6	0.6	*	*	*	*	*	0.2	0.4	0.5
2001.....	1.7	*	*	*	*	*	*	0.2	0.3	0.3
2000.....	1.6	0.7	*	*	*	*	*	0.2	0.3	0.5
1999.....	1.5	*	*	*	*	*	*	0.1	0.3	0.4

White, male

2006 4/.....	1.2	*	*	*	*	*	*	*	0.3	0.6
2005 4/.....	1.4	*	*	*	*	*	*	*	0.3	0.5
2004 4/.....	1.4	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	1.3	*	*	*	*	*	*	*	0.3	0.5
2002.....	1.3	*	*	*	*	*	*	*	0.4	0.5
2001.....	1.4	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.4	*	*	*	*	*	*	*	0.3	0.5
1999.....	1.2	*	*	*	*	*	*	*	0.2	0.4

White, female

2006 4/.....	1.5	*	*	*	*	*	*	*	0.3	0.4
2005 4/.....	1.8	*	*	*	*	*	*	0.3	0.3	0.6
2004 4/.....	1.8	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	1.9	*	*	*	*	*	*	0.3	0.3	0.4
2002.....	1.9	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.9	*	*	*	*	*	*	*	0.3	0.3
2000.....	1.9	*	*	*	*	*	*	0.3	0.3	0.4
1999.....	1.7	*	*	*	*	*	*	*	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)										
All races 3/, both sexes										
2006.....	1.0	*	*	*	*	*	0.2	0.2	0.4	0.5
2005.....	1.2	*	*	*	*	*	0.1	0.2	0.4	0.4
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.4	0.4
2003.....	1.2	*	*	*	*	*	0.2	0.3	0.4	0.5
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.4	0.5
2001.....	1.2	*	*	*	*	*	0.1	0.2	0.3	0.4
2000.....	1.2	*	*	*	*	*	0.1	0.3	0.3	0.5
1999.....	0.9	*	*	*	*	*	*	0.1	0.3	0.3
All races 3/, male										
2006.....	0.8	*	*	*	*	*	*	*	0.4	0.5
2005.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2004.....	1.0	*	*	*	*	*	*	0.3	0.4	0.4
2003.....	1.0	*	*	*	*	*	*	0.2	0.3	0.5
2002.....	1.0	*	*	*	*	*	*	*	0.5	0.5
2001.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.0	*	*	*	*	*	*	*	0.3	0.5
1999.....	0.8	*	*	*	*	*	*	*	0.3	0.4
All races 3/, female										
2006.....	1.1	*	*	*	*	*	0.2	*	0.4	0.5
2005.....	1.3	*	*	*	*	*	*	0.3	0.4	0.5
2004.....	1.4	*	*	*	*	*	*	0.2	0.4	0.5
2003.....	1.5	*	*	*	*	*	0.2	0.3	0.4	0.5
2002.....	1.5	*	*	*	*	*	*	0.3	0.4	0.5
2001.....	1.5	*	*	*	*	*	*	0.3	0.4	0.4
2000.....	1.4	*	*	*	*	*	*	0.4	0.3	0.4
1999.....	1.1	*	*	*	*	*	*	*	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)										
White, both sexes										
2006 4/.....	0.9	*	*	*	*	*	*	*	0.3	0.4
2005 4/.....	1.2	*	*	*	*	*	*	0.2	0.3	0.4
2004 4/.....	1.2	*	*	*	*	*	*	0.2	0.3	0.3
2003 4/.....	1.2	*	*	*	*	*	*	0.2	0.3	0.4
2002.....	1.2	*	*	*	*	*	*	0.1	0.3	0.4
2001.....	1.2	*	*	*	*	*	*	0.2	0.3	0.3
2000.....	1.2	*	*	*	*	*	*	0.2	0.3	0.4
1999.....	0.9	*	*	*	*	*	*	*	0.2	0.3
White, male										
2006 4/.....	0.8	*	*	*	*	*	*	*	0.3	0.5
2005 4/.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2004 4/.....	1.0	*	*	*	*	*	*	*	0.3	0.3
2003 4/.....	1.0	*	*	*	*	*	*	*	0.3	0.5
2002.....	0.9	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.0	*	*	*	*	*	*	*	*	0.3
2000.....	1.0	*	*	*	*	*	*	*	0.3	0.4
1999.....	0.7	*	*	*	*	*	*	*	*	0.3
White, female										
2006 4/.....	1.1	*	*	*	*	*	*	*	0.3	0.4
2005 4/.....	1.4	*	*	*	*	*	*	0.3	0.3	0.4
2004 4/.....	1.4	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	1.5	*	*	*	*	*	*	0.3	0.3	0.4
2002.....	1.5	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.5	*	*	*	*	*	*	*	0.3	0.3
2000.....	1.4	*	*	*	*	*	*	*	0.3	0.4
1999.....	1.1	*	*	*	*	*	*	*	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
X. Diseases of the respiratory system (J00-J98)										
All races 3/, both sexes										
2006.....	75.7	16.8	1.8	0.7	0.6	0.9	1.3	1.8	2.5	3.6
2005.....	80.9	16.3	1.8	0.8	0.7	0.9	1.3	1.7	2.6	3.9
2004.....	77.3	15.8	1.9	0.6	0.8	1.1	1.5	1.7	2.3	4.0
2003.....	81.1	17.5	2.2	0.7	1.0	1.2	1.8	1.9	2.7	4.4
2002.....	81.4	15.8	1.9	0.7	1.0	1.2	1.5	1.9	2.6	4.4
2001.....	80.8	17.3	1.8	0.6	0.8	1.0	1.6	1.9	2.8	4.7
2000.....	82.1	17.1	1.7	0.7	0.9	1.1	1.7	2.0	2.9	4.5
1999.....	82.4	18.1	2.1	0.7	0.9	1.3	1.6	2.1	2.9	4.6
All races 3/, male										
2006.....	73.6	19.6	2.0	0.7	0.8	1.0	1.5	2.0	2.5	3.8
2005.....	77.8	18.8	1.9	0.8	0.8	1.2	1.5	1.9	2.6	4.0
2004.....	75.2	17.6	1.8	0.7	0.9	1.2	1.7	1.9	2.8	4.2
2003.....	78.2	20.6	2.5	0.8	1.0	1.3	2.0	2.3	3.1	4.6
2002.....	78.9	18.8	2.2	0.7	0.9	1.4	1.6	1.9	2.8	4.6
2001.....	78.2	19.7	1.9	0.7	0.8	1.1	1.9	1.9	2.9	4.9
2000.....	80.2	18.9	1.7	0.8	0.9	1.3	1.9	2.2	3.1	4.7
1999.....	81.3	19.6	2.2	0.7	1.1	1.4	1.7	2.2	3.0	4.9
All races 3/, female										
2006.....	77.8	13.7	1.6	0.6	0.5	0.8	1.0	1.6	2.6	3.4
2005.....	83.9	13.7	1.6	0.7	0.6	0.6	1.1	1.5	2.5	3.7
2004.....	79.4	14.1	1.9	0.5	0.6	0.9	1.2	1.4	1.9	3.7
2003.....	84.0	14.2	1.8	0.6	0.9	1.0	1.6	1.6	2.4	4.2
2002.....	83.7	12.6	1.5	0.7	1.0	0.9	1.3	1.8	2.5	4.3
2001.....	83.2	14.7	1.8	0.6	0.7	0.9	1.4	1.9	2.6	4.4
2000.....	84.0	15.2	1.7	0.6	0.8	0.9	1.5	1.8	2.7	4.2
1999.....	83.4	16.6	1.9	0.6	0.8	1.1	1.4	2.1	2.8	4.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
X. Diseases of the respiratory system (J00-J98)										
White, both sexes										
2006 4/.....	84.7	12.6	1.5	0.6	0.4	0.8	1.0	1.6	2.2	3.2
2005 4/.....	90.3	12.9	1.5	0.7	0.5	0.8	1.1	1.6	2.2	3.5
2004 4/.....	86.4	12.2	1.5	0.4	0.6	0.9	1.3	1.4	2.1	3.4
2003 4/.....	90.6	14.4	1.8	0.6	0.8	0.9	1.5	1.6	2.4	3.9
2002.....	90.8	12.0	1.4	0.6	0.7	0.9	1.3	1.7	2.3	3.8
2001.....	89.7	12.3	1.5	0.5	0.7	0.8	1.3	1.5	2.2	3.8
2000.....	91.2	13.3	1.5	0.6	0.7	0.9	1.5	1.7	2.5	3.8
1999.....	91.2	13.3	1.6	0.6	0.8	1.1	1.2	1.6	2.3	3.9
White, male										
2006 4/.....	81.2	14.4	1.7	0.7	0.5	0.8	1.2	1.7	2.1	3.3
2005 4/.....	85.5	15.2	1.5	0.7	0.6	1.1	1.2	1.7	2.2	3.6
2004 4/.....	82.8	13.5	1.5	0.4	0.7	1.0	1.6	1.5	2.4	3.5
2003 4/.....	85.8	16.9	2.0	0.6	0.8	0.9	1.7	1.9	2.6	4.1
2002.....	86.7	14.6	1.7	0.6	0.6	1.1	1.4	1.6	2.5	3.7
2001.....	85.7	14.3	1.5	0.6	0.7	0.8	1.6	1.4	2.3	4.1
2000.....	87.7	15.1	1.3	0.7	0.7	1.0	1.7	1.7	2.5	3.9
1999.....	88.8	14.7	1.5	0.6	0.9	1.2	1.4	1.7	2.5	4.1
White, female										
2006 4/.....	88.1	10.6	1.3	0.5	0.3	0.7	0.9	1.5	2.3	3.1
2005 4/.....	95.0	10.4	1.4	0.6	0.5	0.5	1.0	1.4	2.3	3.3
2004 4/.....	89.9	10.9	1.4	0.4	0.5	0.8	1.0	1.3	1.7	3.3
2003 4/.....	95.2	11.8	1.6	0.5	0.8	0.9	1.3	1.3	2.1	3.6
2002.....	94.7	9.2	1.2	0.5	0.8	0.8	1.1	1.7	2.1	3.9
2001.....	93.7	10.2	1.6	0.4	0.6	0.8	1.0	1.6	2.2	3.5
2000.....	94.6	11.3	1.6	0.5	0.7	0.8	1.2	1.7	2.4	3.6
1999.....	93.6	11.8	1.7	0.6	0.6	1.0	1.1	1.5	2.2	3.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
X. Diseases of the respiratory system (J00-J98)										
Black, both sexes										
2006 4/.....	43.0	37.1	2.9	1.2	1.8	1.8	2.9	3.4	5.4	6.9
2005 4/.....	46.6	35.1	3.4	1.3	1.4	1.7	2.7	3.0	5.6	7.2
2004 4/.....	44.2	33.4	3.7	1.5	1.5	2.2	2.6	3.5	4.8	8.5
2003 4/.....	45.9	32.7	3.5	1.4	1.7	2.4	3.4	4.4	5.7	8.5
2002.....	46.7	34.1	4.1	1.2	2.3	2.5	2.9	3.5	5.6	9.4
2001.....	47.2	39.6	3.2	1.2	1.5	2.0	3.6	4.3	6.8	11.1
2000.....	48.0	37.7	3.1	1.2	1.9	2.5	3.4	4.4	6.1	9.6
1999.....	49.2	41.6	4.0	1.1	2.1	2.0	3.6	5.7	6.9	10.2
Black, male										
2006 4/.....	45.6	44.4	3.4	1.2	2.3	2.2	3.5	4.3	5.6	7.8
2005 4/.....	49.8	39.0	4.2	1.6	1.8	2.2	3.3	3.4	6.3	7.9
2004 4/.....	47.9	37.1	3.5	2.2	1.8	2.6	2.6	4.9	6.1	9.9
2003 4/.....	50.0	38.7	4.5	1.8	2.0	3.1	3.9	5.5	7.4	8.8
2002.....	50.2	39.2	4.7	*	2.4	3.4	3.2	4.0	6.3	10.7
2001.....	50.5	43.7	3.8	1.2	1.7	2.6	3.5	4.6	7.8	11.3
2000.....	52.4	39.3	3.9	1.4	2.2	3.1	3.7	5.9	7.7	10.7
1999.....	54.0	44.6	5.1	1.4	2.5	2.3	4.3	5.6	7.3	10.9
Black, female										
2006 4/.....	40.6	29.5	2.3	*	1.3	1.5	2.3	2.6	5.3	6.2
2005 4/.....	43.6	31.1	2.6	*	*	*	2.2	2.5	4.9	6.5
2004 4/.....	40.9	29.6	3.8	*	1.2	1.8	2.5	2.2	3.6	7.3
2003 4/.....	42.1	26.6	2.4	*	1.3	1.7	2.9	3.3	4.2	8.2
2002.....	43.5	28.8	3.5	1.4	2.2	1.5	2.7	3.1	4.9	8.3
2001.....	44.1	35.3	2.5	*	1.4	1.4	3.6	4.0	5.9	10.9
2000.....	44.1	36.0	2.2	*	1.5	1.9	3.2	3.0	4.6	8.6
1999.....	44.8	38.6	2.9	*	1.6	1.8	2.9	5.7	6.5	9.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of the respiratory system (J10-J98)										
All races 3/, both sexes										
2006.....	75.6	16.5	1.7	0.7	0.6	0.9	1.3	1.8	2.5	3.6
2005.....	80.8	16.0	1.7	0.7	0.7	0.9	1.3	1.7	2.6	3.9
2004.....	77.3	15.7	1.8	0.6	0.8	1.0	1.5	1.6	2.3	4.0
2003.....	81.1	17.1	2.1	0.7	0.9	1.1	1.8	1.9	2.7	4.3
2002.....	81.3	15.5	1.8	0.7	0.9	1.1	1.4	1.9	2.6	4.4
2001.....	80.7	17.0	1.7	0.6	0.8	1.0	1.6	1.9	2.8	4.7
2000.....	82.0	16.9	1.7	0.7	0.9	1.1	1.7	2.0	2.9	4.5
1999.....	82.3	17.8	2.0	0.7	0.9	1.2	1.6	2.1	2.9	4.6
All races 3/, male										
2006.....	73.5	19.2	1.9	0.7	0.8	1.0	1.5	2.0	2.5	3.8
2005.....	77.7	18.5	1.8	0.8	0.8	1.2	1.5	1.9	2.6	4.0
2004.....	75.1	17.4	1.8	0.7	0.9	1.2	1.7	1.9	2.7	4.2
2003.....	78.1	20.1	2.4	0.8	1.0	1.3	2.0	2.3	3.0	4.5
2002.....	78.9	18.6	2.1	0.7	0.9	1.4	1.6	1.9	2.8	4.5
2001.....	78.1	19.4	1.8	0.6	0.8	1.1	1.9	1.8	2.9	4.9
2000.....	80.1	18.7	1.7	0.8	0.9	1.3	1.9	2.2	3.1	4.7
1999.....	81.3	19.2	2.1	0.7	1.1	1.3	1.7	2.1	3.0	4.8
All races 3/, female										
2006.....	77.7	13.6	1.5	0.6	0.5	0.8	1.0	1.6	2.6	3.4
2005.....	83.8	13.5	1.5	0.7	0.6	0.6	1.1	1.5	2.5	3.7
2004.....	79.4	14.0	1.8	0.4	0.6	0.9	1.2	1.4	1.9	3.7
2003.....	83.9	13.9	1.8	0.6	0.9	1.0	1.5	1.6	2.3	4.2
2002.....	83.7	12.4	1.4	0.6	1.0	0.9	1.3	1.8	2.5	4.3
2001.....	83.2	14.6	1.7	0.6	0.7	0.9	1.4	1.9	2.6	4.4
2000.....	83.9	15.1	1.6	0.6	0.8	0.9	1.5	1.8	2.7	4.2
1999.....	83.3	16.4	1.9	0.6	0.7	1.1	1.4	2.0	2.8	4.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of the respiratory system (J10-J98)										
White, both sexes										
2006 4/.....	84.7	12.3	1.4	0.5	0.4	0.7	1.0	1.6	2.1	3.2
2005 4/.....	90.3	12.7	1.4	0.6	0.5	0.8	1.1	1.5	2.2	3.5
2004 4/.....	86.3	12.1	1.4	0.3	0.6	0.9	1.3	1.4	2.1	3.4
2003 4/.....	90.5	14.1	1.7	0.6	0.8	0.9	1.5	1.6	2.3	3.9
2002.....	90.7	11.9	1.3	0.6	0.7	0.9	1.2	1.7	2.3	3.8
2001.....	89.7	12.1	1.4	0.5	0.7	0.8	1.3	1.5	2.2	3.8
2000.....	91.1	13.1	1.4	0.6	0.7	0.9	1.4	1.7	2.4	3.8
1999.....	91.2	13.1	1.5	0.6	0.7	1.1	1.2	1.6	2.3	3.9
White, male										
2006 4/.....	81.2	14.1	1.6	0.6	0.5	0.8	1.2	1.7	2.0	3.3
2005 4/.....	85.5	15.0	1.4	0.6	0.6	1.1	1.2	1.7	2.2	3.6
2004 4/.....	82.7	13.2	1.4	0.4	0.7	1.0	1.6	1.5	2.4	3.5
2003 4/.....	85.8	16.4	1.9	0.6	0.8	0.9	1.7	1.8	2.5	4.1
2002.....	86.7	14.5	1.6	0.6	0.6	1.1	1.4	1.6	2.5	3.7
2001.....	85.6	14.1	1.4	0.5	0.7	0.8	1.6	1.4	2.3	4.1
2000.....	87.6	14.8	1.3	0.7	0.7	1.0	1.7	1.7	2.4	3.9
1999.....	88.7	14.5	1.4	0.6	0.9	1.2	1.4	1.7	2.5	4.1
White, female										
2006 4/.....	88.1	10.4	1.3	0.5	0.3	0.7	0.9	1.5	2.3	3.1
2005 4/.....	94.9	10.3	1.3	0.6	0.5	0.5	0.9	1.4	2.3	3.3
2004 4/.....	89.9	10.8	1.3	0.3	0.5	0.7	1.0	1.3	1.7	3.3
2003 4/.....	95.2	11.6	1.5	0.5	0.8	0.8	1.3	1.3	2.1	3.6
2002.....	94.7	9.0	1.1	0.5	0.8	0.8	1.1	1.7	2.1	3.9
2001.....	93.6	10.0	1.5	0.4	0.6	0.8	1.0	1.6	2.2	3.5
2000.....	94.5	11.3	1.6	0.5	0.7	0.7	1.2	1.7	2.4	3.6
1999.....	93.5	11.7	1.6	0.6	0.6	1.0	1.1	1.5	2.2	3.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of the respiratory system (J10-J98)										
Black, both sexes										
2006 4/.....	42.9	36.6	2.8	1.2	1.8	1.8	2.9	3.4	5.4	6.9
2005 4/.....	46.5	34.7	3.4	1.3	1.3	1.7	2.7	2.9	5.6	7.2
2004 4/.....	44.2	33.4	3.7	1.5	1.5	2.2	2.6	3.5	4.8	8.4
2003 4/.....	45.8	32.3	3.4	1.4	1.7	2.4	3.4	4.4	5.7	8.4
2002.....	46.6	33.5	4.0	1.2	2.3	2.5	2.9	3.5	5.6	9.4
2001.....	47.1	39.1	3.0	1.2	1.5	1.9	3.6	4.3	6.8	11.1
2000.....	48.0	37.6	3.0	1.2	1.8	2.5	3.3	4.3	6.1	9.6
1999.....	49.1	41.0	3.9	1.0	2.1	2.0	3.6	5.7	6.7	10.1
Black, male										
2006 4/.....	45.5	43.5	3.3	1.2	2.3	2.2	3.5	4.3	5.6	7.8
2005 4/.....	49.7	38.7	4.1	1.4	1.8	2.2	3.2	3.4	6.3	7.9
2004 4/.....	47.8	37.1	3.5	2.2	1.8	2.5	2.6	4.9	6.1	9.8
2003 4/.....	49.9	38.4	4.4	1.8	2.0	3.1	3.8	5.5	7.4	8.7
2002.....	50.2	38.3	4.7	*	2.4	3.4	3.1	4.0	6.3	10.7
2001.....	50.4	42.8	3.6	1.2	1.7	2.5	3.5	4.6	7.8	11.2
2000.....	52.4	39.3	3.9	1.4	2.2	3.1	3.5	5.8	7.7	10.7
1999.....	53.9	43.9	4.8	1.3	2.5	2.2	4.3	5.6	7.0	10.9
Black, female										
2006 4/.....	40.6	29.5	2.3	*	1.3	1.3	2.3	2.6	5.3	6.2
2005 4/.....	43.6	30.5	2.6	*	*	*	2.2	2.4	4.9	6.5
2004 4/.....	40.9	29.6	3.8	*	1.2	1.8	2.5	2.2	3.6	7.2
2003 4/.....	42.1	26.0	2.3	*	1.3	1.7	2.9	3.3	4.2	8.1
2002.....	43.4	28.5	3.4	1.4	2.2	1.5	2.6	3.1	4.9	8.3
2001.....	44.1	35.3	2.5	*	1.3	1.4	3.6	4.0	5.9	10.9
2000.....	44.0	35.7	2.2	*	1.5	1.9	3.1	3.0	4.6	8.6
1999.....	44.7	37.9	2.9	*	1.6	1.8	2.9	5.7	6.4	9.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
All races 3/, both sexes										
2006.....	18.5	5.9	0.7	0.1	*	0.3	0.5	0.7	1.0	1.4
2005.....	20.6	6.0	0.6	0.2	0.2	0.3	0.5	0.7	1.0	1.5
2004.....	19.9	6.3	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.4
2003.....	21.8	7.3	0.7	0.2	0.3	0.3	0.6	0.8	1.0	1.7
2002.....	22.5	6.3	0.7	0.2	0.2	0.3	0.4	0.7	1.0	1.7
2001.....	21.7	7.2	0.7	0.2	0.2	0.3	0.6	0.6	1.0	1.8
2000.....	22.6	7.4	0.6	0.2	0.2	0.3	0.6	0.7	1.1	1.7
1999.....	22.2	8.1	0.8	0.2	0.2	0.3	0.5	0.7	1.0	2.0
All races 3/, male										
2006.....	17.1	7.2	0.6	*	*	0.3	0.6	0.8	0.9	1.5
2005.....	18.8	6.6	0.6	0.2	0.2	0.4	0.6	0.8	1.1	1.8
2004.....	18.3	7.5	0.5	*	0.2	0.4	0.7	0.7	1.2	1.7
2003.....	19.6	8.5	0.7	0.2	0.2	0.4	0.7	0.9	1.3	1.9
2002.....	20.2	7.6	0.8	*	0.2	0.4	0.5	0.7	1.2	1.8
2001.....	19.5	7.8	0.7	0.2	0.2	0.3	0.7	0.6	1.2	2.0
2000.....	20.3	8.1	0.6	0.2	*	0.3	0.7	0.8	1.2	1.8
1999.....	19.8	8.5	0.7	0.2	0.2	0.3	0.6	0.8	1.1	2.2
All races 3/, female										
2006.....	19.9	4.6	0.7	*	*	0.3	0.4	0.6	1.0	1.3
2005.....	22.5	5.3	0.6	0.2	0.2	*	0.3	0.7	0.9	1.2
2004.....	21.5	5.0	0.8	*	*	0.2	0.4	0.4	0.7	1.2
2003.....	23.9	6.0	0.6	0.2	0.3	0.2	0.6	0.6	0.8	1.5
2002.....	24.8	5.0	0.6	*	0.2	0.3	0.4	0.7	0.8	1.5
2001.....	23.8	6.7	0.7	*	0.2	0.3	0.5	0.6	0.8	1.6
2000.....	24.8	6.6	0.6	*	*	0.3	0.6	0.6	0.9	1.7
1999.....	24.6	7.7	0.8	*	*	0.3	0.5	0.6	0.8	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
White, both sexes										
2006 4/.....	20.1	4.6	0.6	*	*	0.3	0.4	0.6	0.9	1.3
2005 4/.....	22.4	4.5	0.5	0.2	0.2	0.3	0.5	0.7	0.9	1.4
2004 4/.....	21.6	4.7	0.5	*	0.2	0.3	0.5	0.5	0.9	1.2
2003 4/.....	23.7	6.0	0.6	0.2	0.3	0.3	0.7	0.6	0.9	1.5
2002.....	24.6	5.0	0.5	0.2	0.2	0.3	0.4	0.7	0.9	1.4
2001.....	23.5	5.1	0.6	0.2	0.2	0.3	0.5	0.5	0.9	1.5
2000.....	24.4	5.6	0.5	0.2	0.2	0.3	0.6	0.6	1.0	1.6
1999.....	24.1	5.9	0.6	0.2	0.2	0.3	0.4	0.5	0.8	1.7
White, male										
2006 4/.....	18.3	5.7	0.5	*	*	0.3	0.5	0.7	0.8	1.3
2005 4/.....	20.0	5.3	0.4	*	*	0.4	0.6	0.7	1.0	1.6
2004 4/.....	19.5	5.8	0.4	*	*	0.4	0.7	0.6	1.1	1.4
2003 4/.....	20.9	7.1	0.7	*	0.3	0.4	0.7	0.8	1.1	1.7
2002.....	21.7	6.4	0.5	*	*	0.4	0.5	0.6	1.0	1.5
2001.....	20.6	5.8	0.6	*	*	0.3	0.7	0.5	1.0	1.6
2000.....	21.5	6.1	0.4	0.3	*	0.3	0.6	0.6	1.0	1.7
1999.....	21.0	6.7	0.4	*	0.2	0.3	0.5	0.6	0.9	1.9
White, female										
2006 4/.....	21.8	3.5	0.6	*	*	0.3	0.4	0.6	0.9	1.2
2005 4/.....	24.8	3.7	0.5	*	*	*	0.3	0.7	0.8	1.1
2004 4/.....	23.6	3.5	0.6	*	*	*	0.3	0.4	0.7	1.0
2003 4/.....	26.5	4.8	0.5	*	0.3	*	0.6	0.5	0.7	1.3
2002.....	27.4	3.6	0.5	*	*	*	0.4	0.7	0.7	1.4
2001.....	26.2	4.4	0.6	*	*	0.3	0.3	0.6	0.7	1.4
2000.....	27.3	5.0	0.6	*	*	*	0.5	0.6	0.9	1.5
1999.....	27.1	5.2	0.7	*	*	0.3	0.4	0.4	0.6	1.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia (J12-J18)										
Black, both sexes										
2006 4/.....	13.3	12.5	1.1	*	*	*	0.9	1.0	1.6	2.4
2005 4/.....	14.6	14.0	1.0	*	*	*	*	0.9	1.8	2.5
2004 4/.....	14.6	13.5	1.2	*	*	*	0.8	0.9	1.5	3.0
2003 4/.....	15.1	13.0	0.9	*	*	*	0.7	1.6	2.0	3.3
2002.....	15.5	12.7	1.5	*	*	*	0.7	1.0	2.0	3.5
2001.....	15.5	16.3	1.0	*	*	*	1.0	1.2	2.0	3.7
2000.....	16.1	16.4	1.2	*	*	*	1.0	1.4	2.0	2.9
1999.....	16.1	18.3	1.7	*	*	*	1.3	1.7	2.2	4.0
Black, male										
2006 4/.....	13.0	15.3	*	*	*	*	*	1.4	*	2.8
2005 4/.....	14.5	13.9	*	*	*	*	*	*	2.2	3.2
2004 4/.....	14.9	15.1	*	*	*	*	*	*	1.9	3.7
2003 4/.....	15.8	15.2	*	*	*	*	*	1.6	2.8	3.5
2002.....	15.3	13.7	1.6	*	*	*	*	*	2.4	4.0
2001.....	15.8	15.9	*	*	*	*	*	*	2.7	4.5
2000.....	16.5	17.9	1.7	*	*	*	*	2.0	2.5	3.0
1999.....	16.3	17.9	1.9	*	*	*	*	1.9	2.5	4.4
Black, female										
2006 4/.....	13.6	9.6	*	*	*	*	*	*	2.1	2.0
2005 4/.....	14.8	14.2	*	*	*	*	*	*	1.5	1.8
2004 4/.....	14.4	11.7	1.7	*	*	*	*	*	*	2.3
2003 4/.....	14.6	10.8	*	*	*	*	*	1.5	1.4	3.1
2002.....	15.6	11.8	*	*	*	*	*	*	1.6	3.1
2001.....	15.1	16.7	*	*	*	*	*	*	1.4	3.0
2000.....	15.8	14.9	*	*	*	*	*	*	1.6	2.7
1999.....	15.9	18.8	*	*	*	*	*	1.5	1.9	3.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia due to other or unspecified organisms (J16,J18)										
All races 3/, both sexes										
2006.....	16.9	3.5	0.4	*	*	0.2	0.4	0.6	0.8	1.2
2005.....	18.9	3.7	0.4	0.1	0.2	0.2	0.4	0.6	0.8	1.2
2004.....	18.4	4.1	0.4	0.1	0.1	0.3	0.5	0.4	0.8	1.2
2003.....	20.0	4.6	0.4	0.1	0.2	0.2	0.6	0.7	0.9	1.4
2002.....	20.8	3.9	0.5	0.1	0.2	0.3	0.4	0.6	0.8	1.4
2001.....	20.0	4.6	0.5	0.2	0.2	0.3	0.5	0.5	0.9	1.5
2000.....	20.8	5.1	0.4	0.2	0.1	0.2	0.5	0.6	0.9	1.4
1999.....	20.7	4.7	0.5	0.1	0.2	0.3	0.5	0.6	0.9	1.6
All races 3/, male										
2006.....	15.4	4.4	0.4	*	*	0.3	0.4	0.7	0.8	1.3
2005.....	17.0	3.9	0.4	*	*	0.3	0.5	0.6	0.9	1.5
2004.....	16.7	4.9	0.3	*	*	0.4	0.6	0.6	1.0	1.5
2003.....	17.8	5.5	0.4	*	0.2	0.3	0.6	0.8	1.1	1.6
2002.....	18.5	4.6	0.5	*	*	0.3	0.5	0.5	1.0	1.6
2001.....	17.8	4.9	0.5	*	0.2	0.3	0.6	0.5	1.0	1.7
2000.....	18.5	5.3	0.4	*	*	0.3	0.6	0.7	1.0	1.5
1999.....	18.3	5.0	0.5	*	*	0.3	0.5	0.6	1.0	1.8
All races 3/, female										
2006.....	18.4	2.5	0.4	*	*	0.2	0.3	0.5	0.9	1.1
2005.....	20.8	3.5	0.4	*	*	*	0.3	0.6	0.7	1.0
2004.....	20.0	3.3	0.5	*	*	*	0.3	0.3	0.6	0.9
2003.....	22.2	3.8	0.5	*	*	*	0.5	0.5	0.7	1.3
2002.....	23.0	3.1	0.4	*	*	0.3	0.3	0.6	0.7	1.3
2001.....	22.2	4.2	0.5	*	*	0.3	0.4	0.5	0.7	1.3
2000.....	23.1	4.9	0.4	*	*	0.2	0.5	0.4	0.8	1.4
1999.....	23.0	4.3	0.5	*	*	0.3	0.4	0.5	0.7	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia due to other or unspecified organisms (J16,J18)										
White, both sexes										
2006 4/.....	18.4	2.7	0.3	*	*	0.3	0.3	0.5	0.7	1.0
2005 4/.....	20.6	2.7	0.3	0.1	0.2	0.2	0.4	0.6	0.7	1.2
2004 4/.....	19.9	2.9	0.3	*	*	0.3	0.5	0.4	0.8	1.0
2003 4/.....	21.8	3.8	0.4	*	0.2	0.2	0.6	0.6	0.8	1.3
2002.....	22.7	2.9	0.4	0.1	0.2	0.3	0.4	0.5	0.7	1.2
2001.....	21.7	3.0	0.5	0.2	0.2	0.2	0.5	0.4	0.7	1.3
2000.....	22.6	3.9	0.3	0.2	0.1	0.2	0.5	0.5	0.8	1.3
1999.....	22.5	3.3	0.4	0.1	0.2	0.3	0.4	0.5	0.7	1.4
White, male										
2006 4/.....	16.5	3.4	0.3	*	*	0.3	0.4	0.6	0.8	1.1
2005 4/.....	18.2	3.3	0.3	*	*	0.3	0.5	0.6	0.8	1.4
2004 4/.....	17.8	3.4	*	*	*	0.4	0.7	0.5	0.9	1.2
2003 4/.....	19.0	4.5	0.4	*	*	0.3	0.6	0.7	0.9	1.4
2002.....	19.9	3.7	0.4	*	*	0.3	0.5	0.5	0.9	1.3
2001.....	18.9	3.5	0.5	*	*	*	0.6	0.4	0.8	1.4
2000.....	19.6	4.3	*	*	*	0.2	0.6	0.6	0.9	1.4
1999.....	19.5	3.8	*	*	*	0.3	0.4	0.5	0.8	1.6
White, female										
2006 4/.....	20.2	1.9	*	*	*	*	*	0.5	0.7	1.0
2005 4/.....	23.0	2.1	0.4	*	*	*	0.3	0.6	0.7	0.9
2004 4/.....	22.0	2.5	0.3	*	*	*	*	0.3	0.6	0.8
2003 4/.....	24.6	3.0	0.4	*	*	*	0.5	0.4	0.6	1.1
2002.....	25.5	2.0	0.3	*	*	*	0.3	0.5	0.6	1.2
2001.....	24.5	2.5	0.4	*	*	0.3	0.3	0.5	0.6	1.1
2000.....	25.5	3.5	0.4	*	*	*	0.4	0.4	0.7	1.2
1999.....	25.4	2.8	0.4	*	*	0.3	0.3	0.4	0.6	1.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pneumonia due to other or unspecified organisms (J16,J18)										
Black, both sexes										
2006 4/.....	12.0	7.9	*	*	*	*	0.8	0.9	1.4	2.2
2005 4/.....	13.3	9.3	0.8	*	*	*	*	0.8	1.5	2.1
2004 4/.....	13.2	10.3	1.0	*	*	*	0.8	0.7	1.2	2.6
2003 4/.....	13.7	8.8	*	*	*	*	0.7	1.3	1.9	2.9
2002.....	14.2	8.2	1.0	*	*	*	*	0.9	1.6	3.0
2001.....	14.1	11.8	0.8	*	*	*	1.0	1.1	1.7	3.3
2000.....	14.7	11.2	0.8	*	*	*	0.9	1.2	1.9	2.4
1999.....	14.7	11.4	1.1	*	*	*	1.1	1.4	1.9	3.4
Black, male										
2006 4/.....	11.6	9.8	*	*	*	*	*	*	*	2.7
2005 4/.....	13.0	7.4	*	*	*	*	*	*	1.9	2.7
2004 4/.....	13.4	12.5	*	*	*	*	*	*	*	3.5
2003 4/.....	14.2	10.1	*	*	*	*	*	*	2.6	3.1
2002.....	13.9	8.1	*	*	*	*	*	*	2.2	3.3
2001.....	14.3	11.1	*	*	*	*	*	*	2.4	3.8
2000.....	14.9	10.6	*	*	*	*	*	1.8	2.3	2.5
1999.....	14.8	12.0	*	*	*	*	*	1.6	2.4	3.7
Black, female										
2006 4/.....	12.4	6.0	*	*	*	*	*	*	1.7	1.8
2005 4/.....	13.5	11.4	*	*	*	*	*	*	*	1.6
2004 4/.....	13.0	8.0	*	*	*	*	*	*	*	1.8
2003 4/.....	13.3	7.4	*	*	*	*	*	*	1.4	2.7
2002.....	14.4	8.2	*	*	*	*	*	*	*	2.8
2001.....	14.0	12.6	*	*	*	*	*	*	*	2.8
2000.....	14.6	11.9	*	*	*	*	*	*	1.5	2.4
1999.....	14.6	10.7	*	*	*	*	*	*	1.5	3.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
All races 3/, both sexes										
2006.....	41.6	0.7	0.3	0.3	0.3	0.3	0.4	0.5	0.7	1.0
2005.....	44.2	0.8	0.3	0.3	0.3	0.3	0.4	0.5	0.8	1.2
2004.....	41.5	0.9	0.3	0.2	0.3	0.4	0.4	0.6	0.7	1.3
2003.....	43.5	0.8	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
2002.....	43.3	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.6
2001.....	43.2	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6
2000.....	43.4	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
1999.....	44.5	0.9	0.4	0.2	0.4	0.5	0.5	0.7	0.9	1.3
All races 3/, male										
2006.....	40.2	0.9	0.3	0.3	0.4	0.4	0.5	0.6	0.7	1.0
2005.....	42.8	*	0.4	0.3	0.3	0.4	0.5	0.6	0.7	1.2
2004.....	40.6	1.2	0.4	0.3	0.4	0.5	0.5	0.6	0.9	1.2
2003.....	42.4	1.1	0.5	0.2	0.5	0.5	0.6	0.7	0.9	1.3
2002.....	42.9	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.5
2001.....	42.7	1.1	0.4	0.2	0.3	0.4	0.6	0.6	0.9	1.5
2000.....	43.5	1.2	0.4	0.2	0.5	0.6	0.6	0.7	0.9	1.3
1999.....	45.6	*	0.4	0.3	0.6	0.6	0.7	0.7	0.8	1.2
All races 3/, female										
2006.....	43.0	*	*	*	*	0.2	0.4	0.5	0.7	1.1
2005.....	45.5	*	0.3	0.2	0.2	*	0.4	0.4	0.9	1.3
2004.....	42.5	*	*	*	0.3	0.3	0.4	0.5	0.5	1.4
2003.....	44.4	*	*	*	0.3	0.3	0.4	0.5	0.7	1.5
2002.....	43.7	*	0.3	*	0.4	0.3	0.5	0.5	0.9	1.7
2001.....	43.7	*	*	*	0.2	0.3	0.4	0.5	0.9	1.7
2000.....	43.2	*	0.3	0.2	0.4	0.3	0.5	0.6	0.8	1.4
1999.....	43.4	*	0.3	0.2	0.3	0.5	0.4	0.8	1.0	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
White, both sexes										
2006 4/.....	47.5	*	0.2	0.2	*	0.1	0.3	0.4	0.5	0.9
2005 4/.....	50.3	0.6	0.2	0.2	0.1	0.2	0.2	0.4	0.6	1.1
2004 4/.....	47.4	0.7	0.2	*	0.2	0.2	0.3	0.4	0.5	1.1
2003 4/.....	49.5	0.7	0.2	*	0.2	0.2	0.4	0.5	0.6	1.2
2002.....	49.2	0.8	0.2	*	0.2	0.3	0.4	0.5	0.7	1.3
2001.....	49.0	0.7	0.2	*	0.2	0.3	0.3	0.4	0.6	1.2
2000.....	49.0	0.8	0.2	0.1	0.2	0.3	0.4	0.4	0.6	1.0
1999.....	50.2	*	0.3	0.2	0.3	0.4	0.4	0.5	0.6	1.1
White, male										
2006 4/.....	45.1	*	*	*	*	*	0.3	0.5	0.4	0.8
2005 4/.....	47.9	*	*	*	*	0.3	0.3	0.4	0.4	1.0
2004 4/.....	45.5	*	*	*	0.3	0.3	0.4	0.4	0.6	1.0
2003 4/.....	47.4	1.3	*	*	0.3	0.2	0.5	0.4	0.5	1.1
2002.....	47.8	*	0.3	*	*	0.4	0.4	0.5	0.8	1.1
2001.....	47.6	*	*	*	*	*	0.3	0.4	0.5	1.2
2000.....	48.3	*	*	*	0.3	0.3	0.5	0.5	0.6	0.9
1999.....	50.6	*	*	*	0.4	0.4	0.4	0.4	0.5	1.0
White, female										
2006 4/.....	49.9	*	*	*	*	*	0.3	0.3	0.6	0.9
2005 4/.....	52.8	*	*	*	*	*	*	0.4	0.7	1.2
2004 4/.....	49.3	*	*	*	*	*	0.3	0.4	0.4	1.2
2003 4/.....	51.5	*	*	*	*	*	0.3	0.5	0.7	1.3
2002.....	50.5	*	*	*	*	*	0.4	0.4	0.7	1.5
2001.....	50.3	*	*	*	*	0.3	*	0.3	0.7	1.2
2000.....	49.7	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	49.8	*	*	*	*	0.4	0.4	0.5	0.7	1.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Chronic lower respiratory diseases (J40-J47)										
Black, both sexes										
2006 4/.....	19.5	*	*	0.9	1.2	1.1	1.2	1.5	2.3	2.7
2005 4/.....	21.1	*	1.2	0.7	0.9	0.7	1.6	1.2	2.3	2.6
2004 4/.....	19.2	*	0.8	0.9	1.0	1.3	1.2	1.7	1.9	3.0
2003 4/.....	20.2	*	1.0	0.7	1.1	1.3	1.3	1.7	2.1	2.8
2002.....	20.7	*	1.2	0.6	1.6	1.4	1.2	1.6	2.2	3.8
2001.....	20.4	*	0.9	0.7	0.9	1.0	1.6	1.8	3.0	4.5
2000.....	20.8	*	0.8	0.7	1.4	1.4	1.3	1.9	2.4	3.5
1999.....	21.9	*	*	0.6	1.5	1.2	1.5	2.5	2.7	3.2
Black, male										
2006 4/.....	21.9	*	*	*	1.7	1.4	1.4	1.7	2.8	2.6
2005 4/.....	23.9	*	1.6	*	1.2	1.2	1.8	1.5	2.5	2.7
2004 4/.....	22.0	*	*	1.3	1.3	1.6	1.3	2.4	2.6	3.1
2003 4/.....	23.3	*	1.6	*	1.5	1.6	1.5	2.4	3.1	2.8
2002.....	24.1	*	1.6	*	1.7	2.2	1.5	2.2	2.3	4.5
2001.....	23.6	*	*	*	*	1.4	1.8	2.1	3.4	4.3
2000.....	24.3	*	*	*	1.7	2.0	1.5	2.4	3.1	3.7
1999.....	26.2	*	*	*	2.1	1.5	2.2	2.6	2.7	3.3
Black, female										
2006 4/.....	17.4	*	*	*	*	*	*	1.4	1.8	2.7
2005 4/.....	18.4	*	*	*	*	*	1.4	*	2.2	2.5
2004 4/.....	16.6	*	*	*	*	*	*	*	1.4	2.9
2003 4/.....	17.3	*	*	*	*	*	*	*	*	2.9
2002.....	17.7	*	*	*	1.4	*	*	*	2.1	3.2
2001.....	17.5	*	*	*	*	*	1.4	1.5	2.6	4.7
2000.....	17.6	*	*	*	*	*	*	1.4	1.7	3.3
1999.....	18.0	*	*	*	*	*	*	2.4	2.8	3.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
All races 3/, both sexes										
2006.....	41.3	0.7	0.3	0.3	0.3	0.3	0.4	0.5	0.7	1.0
2005.....	43.8	0.8	0.3	0.3	0.3	0.3	0.4	0.5	0.8	1.2
2004.....	41.2	0.9	0.3	0.2	0.3	0.4	0.4	0.6	0.7	1.3
2003.....	43.1	0.8	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.4
2002.....	42.9	1.0	0.4	0.2	0.4	0.5	0.5	0.6	0.9	1.5
2001.....	42.9	1.0	0.3	0.2	0.3	0.4	0.5	0.5	0.9	1.6
2000.....	43.0	0.9	0.3	0.2	0.4	0.4	0.5	0.6	0.8	1.3
1999.....	44.2	0.9	0.3	0.2	0.4	0.5	0.5	0.7	0.9	1.3
All races 3/, male										
2006.....	40.0	0.9	0.3	0.3	0.4	0.4	0.5	0.6	0.7	1.0
2005.....	42.6	*	0.4	0.3	0.3	0.4	0.5	0.6	0.7	1.2
2004.....	40.4	1.2	0.4	0.3	0.4	0.5	0.5	0.6	0.8	1.2
2003.....	42.3	1.1	0.5	0.2	0.5	0.4	0.6	0.7	0.8	1.3
2002.....	42.7	1.1	0.6	0.2	0.5	0.6	0.5	0.7	0.9	1.4
2001.....	42.5	1.1	0.4	0.2	0.3	0.4	0.5	0.6	0.9	1.5
2000.....	43.3	1.2	0.4	0.2	0.5	0.5	0.6	0.7	0.9	1.3
1999.....	45.4	*	0.4	0.3	0.6	0.6	0.6	0.7	0.8	1.2
All races 3/, female										
2006.....	42.6	*	*	*	*	0.2	0.4	0.5	0.7	1.1
2005.....	45.1	*	0.3	0.2	*	*	0.4	0.4	0.9	1.3
2004.....	42.0	*	*	*	0.3	0.3	0.4	0.5	0.5	1.3
2003.....	44.0	*	*	*	0.3	0.3	0.4	0.5	0.7	1.5
2002.....	43.2	*	0.3	*	0.4	0.3	0.5	0.5	0.8	1.6
2001.....	43.2	*	*	*	0.2	0.3	0.4	0.5	0.9	1.6
2000.....	42.8	*	0.3	0.2	0.4	0.3	0.4	0.5	0.7	1.4
1999.....	43.0	*	0.3	0.2	0.3	0.5	0.4	0.7	1.0	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
White, both sexes										
2006 4/.....	47.1	*	0.2	0.2	*	0.1	0.3	0.4	0.5	0.9
2005 4/.....	50.0	0.6	0.2	0.2	*	0.2	0.2	0.4	0.6	1.0
2004 4/.....	47.0	0.7	0.2	*	0.2	0.2	0.3	0.4	0.5	1.1
2003 4/.....	49.1	0.7	0.2	*	0.2	0.2	0.4	0.5	0.6	1.2
2002.....	48.8	0.8	0.2	*	0.2	0.3	0.4	0.4	0.7	1.3
2001.....	48.6	0.7	0.2	*	0.2	0.2	0.3	0.3	0.6	1.2
2000.....	48.7	0.8	0.2	0.1	0.2	0.2	0.4	0.4	0.6	1.0
1999.....	49.8	*	0.3	0.2	0.3	0.4	0.4	0.4	0.6	1.1
White, male										
2006 4/.....	44.9	*	*	*	*	*	0.3	0.5	0.4	0.8
2005 4/.....	47.6	*	*	*	*	0.2	0.3	0.4	0.4	1.0
2004 4/.....	45.3	*	*	*	0.3	0.3	0.3	0.4	0.6	1.0
2003 4/.....	47.2	1.3	*	*	0.3	*	0.5	0.4	0.5	1.1
2002.....	47.6	*	0.3	*	*	0.4	0.3	0.5	0.7	1.1
2001.....	47.4	*	*	*	*	*	0.3	0.4	0.5	1.2
2000.....	48.1	*	*	*	0.3	0.3	0.5	0.4	0.6	0.9
1999.....	50.3	*	*	*	0.4	0.4	0.4	0.4	0.5	0.9
White, female										
2006 4/.....	49.4	*	*	*	*	*	0.3	0.3	0.6	0.9
2005 4/.....	52.2	*	*	*	*	*	*	0.4	0.7	1.1
2004 4/.....	48.7	*	*	*	*	*	0.3	0.4	0.4	1.2
2003 4/.....	51.0	*	*	*	*	*	0.3	0.5	0.6	1.3
2002.....	49.9	*	*	*	*	*	0.4	0.4	0.7	1.5
2001.....	49.8	*	*	*	*	0.3	*	0.3	0.7	1.1
2000.....	49.2	*	*	*	*	*	0.3	0.4	0.6	1.2
1999.....	49.3	*	*	*	*	0.4	0.4	0.5	0.7	1.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/ year 2/	Under 1 year 2/ year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
Black, both sexes										
2006 4/.....	19.5	*	*	0.9	1.2	1.1	1.2	1.5	2.3	2.7
2005 4/.....	21.0	*	1.2	0.7	0.9	0.7	1.6	1.2	2.3	2.6
2004 4/.....	19.1	*	0.8	0.9	1.0	1.3	1.2	1.7	1.9	2.9
2003 4/.....	20.1	*	1.0	0.7	1.1	1.3	1.3	1.6	2.1	2.8
2002.....	20.7	*	1.2	0.6	1.6	1.4	1.2	1.6	2.2	3.8
2001.....	20.3	*	0.9	0.7	0.9	1.0	1.6	1.7	2.9	4.5
2000.....	20.7	*	0.8	0.7	1.4	1.4	1.3	1.9	2.3	3.5
1999.....	21.8	*	*	0.6	1.5	1.2	1.5	2.4	2.7	3.2
Black, male										
2006 4/.....	21.8	*	*	*	1.7	1.4	1.4	1.7	2.8	2.6
2005 4/.....	23.9	*	1.6	*	1.2	1.2	1.8	1.5	2.5	2.7
2004 4/.....	21.9	*	*	1.3	1.3	1.6	1.3	2.4	2.6	3.0
2003 4/.....	23.3	*	1.6	*	1.5	1.6	1.5	2.3	3.1	2.8
2002.....	24.1	*	1.6	*	1.7	2.2	1.5	2.2	2.3	4.5
2001.....	23.6	*	*	*	*	1.4	1.8	2.1	3.4	4.3
2000.....	24.3	*	*	*	1.7	1.9	1.5	2.4	3.1	3.7
1999.....	26.1	*	*	*	2.1	1.5	2.2	2.6	2.7	3.3
Black, female										
2006 4/.....	17.3	*	*	*	*	*	*	1.4	1.8	2.7
2005 4/.....	18.4	*	*	*	*	*	1.4	*	2.2	2.5
2004 4/.....	16.5	*	*	*	*	*	*	*	1.4	2.9
2003 4/.....	17.3	*	*	*	*	*	*	*	*	2.9
2002.....	17.6	*	*	*	1.4	*	*	*	2.1	3.2
2001.....	17.4	*	*	*	*	*	1.4	1.4	2.5	4.6
2000.....	17.5	*	*	*	*	*	*	1.4	1.6	3.3
1999.....	17.9	*	*	*	*	*	*	2.2	2.8	3.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
All races 3/, both sexes										
2006.....	1.2	*	0.2	0.2	0.3	0.3	0.4	0.4	0.5	0.7
2005.....	1.3	*	0.2	0.2	0.3	0.2	0.4	0.5	0.6	0.7
2004.....	1.3	*	0.2	0.2	0.3	0.4	0.4	0.5	0.5	0.7
2003.....	1.4	*	0.2	0.2	0.4	0.3	0.4	0.5	0.6	0.7
2002.....	1.5	*	0.3	0.2	0.4	0.4	0.4	0.5	0.6	0.9
2001.....	1.5	*	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9
2000.....	1.6	*	0.2	0.2	0.4	0.4	0.5	0.5	0.7	0.8
1999.....	1.7	*	0.2	0.2	0.4	0.5	0.5	0.6	0.7	0.9
All races 3/, male										
2006.....	0.9	*	*	0.3	0.4	0.3	0.4	0.5	0.5	0.6
2005.....	0.9	*	0.3	0.3	0.3	0.3	0.4	0.5	0.5	0.5
2004.....	1.0	*	*	0.2	0.4	0.5	0.5	0.5	0.6	0.6
2003.....	1.0	*	0.4	0.2	0.4	0.4	0.5	0.6	0.7	0.6
2002.....	1.1	*	0.4	0.2	0.5	0.6	0.5	0.7	0.6	0.8
2001.....	1.1	*	0.3	0.2	0.3	0.3	0.4	0.5	0.6	0.8
2000.....	1.2	*	0.3	0.2	0.5	0.5	0.5	0.7	0.8	0.7
1999.....	1.2	*	*	0.2	0.6	0.5	0.6	0.6	0.6	0.8
All races 3/, female										
2006.....	1.5	*	*	*	*	0.2	0.3	0.4	0.5	0.7
2005.....	1.7	*	*	*	*	*	0.3	0.4	0.7	0.8
2004.....	1.6	*	*	*	0.2	0.3	0.3	0.5	0.4	0.9
2003.....	1.8	*	*	*	0.3	0.3	0.4	0.4	0.6	0.9
2002.....	1.8	*	*	*	0.4	0.2	0.4	0.4	0.6	1.0
2001.....	1.9	*	*	*	0.2	0.3	0.4	0.4	0.8	1.0
2000.....	2.0	*	*	0.2	0.3	0.3	0.4	0.4	0.6	0.9
1999.....	2.1	*	*	*	0.3	0.4	0.4	0.6	0.8	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
White, both sexes										
2006 4/.....	1.0	*	*	*	*	*	0.2	0.3	0.3	0.5
2005 4/.....	1.1	*	*	0.1	*	0.2	0.2	0.4	0.4	0.5
2004 4/.....	1.1	*	*	*	0.2	0.2	0.3	0.3	0.3	0.6
2003 4/.....	1.2	*	*	*	0.2	0.2	0.3	0.4	0.4	0.5
2002.....	1.3	*	*	*	0.2	0.3	0.3	0.4	0.5	0.7
2001.....	1.3	*	*	*	0.2	0.2	0.2	0.3	0.4	0.5
2000.....	1.4	*	*	*	0.2	0.2	0.4	0.3	0.5	0.5
1999.....	1.5	*	*	0.1	0.2	0.4	0.3	0.4	0.4	0.7
White, male										
2006 4/.....	0.7	*	*	*	*	*	0.3	0.4	*	0.4
2005 4/.....	0.7	*	*	*	*	*	0.2	0.4	0.3	0.4
2004 4/.....	0.7	*	*	*	*	0.3	0.3	0.3	0.4	0.4
2003 4/.....	0.8	*	*	*	0.3	*	0.3	0.4	0.4	0.4
2002.....	0.9	*	*	*	*	0.3	0.3	0.4	0.5	0.5
2001.....	0.8	*	*	*	*	*	0.2	0.3	0.3	0.5
2000.....	0.9	*	*	*	0.2	0.3	0.4	0.4	0.5	0.4
1999.....	0.9	*	*	*	0.3	0.4	0.4	0.4	0.4	0.6
White, female										
2006 4/.....	1.4	*	*	*	*	*	*	*	0.4	0.5
2005 4/.....	1.5	*	*	*	*	*	*	0.4	0.5	0.6
2004 4/.....	1.5	*	*	*	*	*	*	0.4	0.3	0.7
2003 4/.....	1.6	*	*	*	*	*	*	0.4	0.5	0.7
2002.....	1.7	*	*	*	*	*	0.3	0.4	0.6	0.9
2001.....	1.7	*	*	*	*	*	*	*	0.6	0.6
2000.....	1.8	*	*	*	*	*	0.3	0.3	0.5	0.6
1999.....	2.0	*	*	*	*	0.4	*	0.4	0.5	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Asthma (J45-J46)										
Black, both sexes										
2006 4/.....	2.4	*	*	0.8	1.1	1.1	1.2	1.3	1.9	2.2
2005 4/.....	2.6	*	1.1	0.7	0.9	0.7	1.5	1.2	1.9	1.8
2004 4/.....	2.6	*	*	0.7	1.0	1.2	1.1	1.6	1.7	2.0
2003 4/.....	2.7	*	0.9	0.7	1.0	1.1	1.3	1.4	1.9	2.0
2002.....	2.9	*	0.9	*	1.5	1.3	1.2	1.6	1.6	2.9
2001.....	3.0	*	0.8	0.7	0.8	0.9	1.4	1.7	2.6	3.4
2000.....	3.2	*	*	0.7	1.4	1.3	1.2	1.8	2.0	2.7
1999.....	3.2	*	*	0.6	1.4	1.1	1.4	2.3	2.4	2.6
Black, male										
2006 4/.....	2.2	*	*	*	1.6	1.4	1.3	1.5	2.4	2.0
2005 4/.....	2.2	*	1.6	*	1.2	*	1.6	1.5	1.9	1.7
2004 4/.....	2.5	*	*	*	1.2	1.5	1.3	2.2	2.3	1.8
2003 4/.....	2.4	*	1.5	*	1.4	1.3	1.4	2.1	2.7	1.9
2002.....	2.8	*	*	*	1.7	2.1	1.5	2.2	1.7	3.4
2001.....	2.6	*	*	*	*	*	1.5	1.9	3.0	3.0
2000.....	2.8	*	*	*	1.7	1.7	*	2.3	2.6	2.7
1999.....	2.8	*	*	*	2.0	1.3	2.0	2.6	2.6	2.6
Black, female										
2006 4/.....	2.6	*	*	*	*	*	*	*	1.6	2.4
2005 4/.....	3.0	*	*	*	*	*	1.3	*	1.9	1.9
2004 4/.....	2.7	*	*	*	*	*	*	*	*	2.2
2003 4/.....	2.9	*	*	*	*	*	*	*	*	2.1
2002.....	3.0	*	*	*	1.4	*	*	*	1.4	2.3
2001.....	3.3	*	*	*	*	*	*	1.4	2.3	3.7
2000.....	3.5	*	*	*	*	*	*	*	1.4	2.6
1999.....	3.5	*	*	*	*	*	*	2.0	2.3	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Lung diseases due to external agents (J60-J70)										
All races 3/, both sexes										
2006.....	6.0	*	*	*	*	*	*	0.2	0.2	0.3
2005.....	6.2	*	*	*	*	0.1	0.1	0.1	0.2	0.4
2004.....	6.1	*	*	*	*	*	0.1	0.2	0.2	0.3
2003.....	6.4	*	*	*	*	*	0.2	0.1	0.2	0.3
2002.....	6.5	*	*	*	*	*	0.1	0.2	0.2	0.3
2001.....	6.5	*	*	*	*	*	0.1	0.2	0.2	0.3
2000.....	6.3	*	*	*	*	*	0.1	0.1	0.2	0.3
1999.....	5.9	*	*	*	*	*	0.1	0.2	0.2	0.3
All races 3/, male										
2006.....	6.7	*	*	*	*	*	*	0.2	0.3	0.3
2005.....	6.7	*	*	*	*	*	*	*	0.2	0.4
2004.....	6.7	*	*	*	*	*	*	0.2	*	0.3
2003.....	6.9	*	*	*	*	*	0.3	*	0.3	0.4
2002.....	7.0	*	*	*	*	*	*	*	0.2	0.4
2001.....	7.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	6.9	*	*	*	*	*	*	*	0.3	0.4
1999.....	6.5	*	*	*	*	*	*	0.2	0.3	0.3
All races 3/, female										
2006.....	5.3	*	*	*	*	*	*	*	*	*
2005.....	5.7	*	*	*	*	*	*	*	*	0.3
2004.....	5.5	*	*	*	*	*	*	*	*	0.2
2003.....	5.9	*	*	*	*	*	*	*	*	0.3
2002.....	6.0	*	*	*	*	*	*	*	*	0.3
2001.....	6.1	*	*	*	*	*	*	*	0.2	0.2
2000.....	5.8	*	*	*	*	*	*	*	*	0.2
1999.....	5.3	*	*	*	*	*	*	*	0.2	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases of respiratory system (J80-J84,J93-J98)										
All races 3/, both sexes										
2006.....	8.4	7.1	0.4	0.1	0.1	0.2	0.2	0.3	0.5	0.7
2005.....	8.3	6.6	0.4	0.1	*	0.2	0.3	0.3	0.5	0.6
2004.....	8.5	5.9	0.5	0.1	0.1	0.2	0.3	0.3	0.4	0.8
2003.....	8.0	6.2	0.4	*	0.1	0.2	0.3	0.3	0.5	0.7
2002.....	7.9	6.1	0.4	0.1	0.1	0.2	0.3	0.3	0.5	0.7
2001.....	8.3	6.5	0.4	0.1	0.2	0.2	0.4	0.4	0.5	0.8
2000.....	8.3	6.4	0.4	0.1	0.2	0.3	0.3	0.4	0.6	0.9
1999.....	8.1	5.8	0.4	0.2	0.1	0.2	0.2	0.4	0.6	0.8
All races 3/, male										
2006.....	8.5	8.1	0.6	*	*	0.2	0.2	0.3	0.4	0.8
2005.....	8.2	8.0	0.5	*	*	0.2	0.2	0.3	0.4	0.5
2004.....	8.5	5.9	0.6	*	*	*	0.2	0.3	0.4	0.8
2003.....	7.8	7.1	0.4	*	*	0.2	0.3	0.3	0.5	0.7
2002.....	7.7	7.3	0.5	*	*	0.2	0.3	0.3	0.4	0.7
2001.....	8.1	8.1	0.4	*	0.2	0.2	0.4	0.3	0.4	0.8
2000.....	8.1	7.1	0.4	*	*	0.2	0.4	0.4	0.6	1.0
1999.....	7.9	6.6	0.4	*	*	0.3	0.3	0.3	0.6	0.8
All races 3/, female										
2006.....	8.3	6.1	0.3	*	*	*	*	0.4	0.6	0.7
2005.....	8.4	5.1	0.3	*	*	*	0.3	0.2	0.5	0.8
2004.....	8.6	5.8	0.3	*	*	0.2	0.3	0.3	0.4	0.8
2003.....	8.1	5.3	0.4	*	*	*	*	0.3	0.6	0.7
2002.....	8.0	4.9	0.3	*	*	0.2	0.3	0.4	0.5	0.8
2001.....	8.6	4.9	0.4	*	*	*	0.3	0.5	0.5	0.9
2000.....	8.4	5.7	0.3	*	*	0.3	0.2	0.4	0.6	0.8
1999.....	8.3	4.9	0.3	*	*	*	*	0.4	0.7	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases of respiratory system (J80-J84,J93-J98)										
White, both sexes										
2006 4/.....	9.2	4.9	0.4	0.1	*	0.2	0.1	0.4	0.5	0.7
2005 4/.....	9.1	5.3	0.4	0.1	*	0.2	0.2	0.3	0.5	0.6
2004 4/.....	9.3	4.2	0.4	*	*	0.2	0.3	0.3	0.3	0.6
2003 4/.....	8.7	4.6	0.3	*	*	0.2	0.2	0.3	0.5	0.6
2002.....	8.6	4.2	0.3	*	*	0.2	0.2	0.3	0.4	0.6
2001.....	9.1	4.5	0.4	*	0.2	0.2	0.3	0.3	0.4	0.6
2000.....	9.0	4.9	0.4	0.1	0.2	0.2	0.3	0.4	0.5	0.7
1999.....	8.8	4.1	0.3	0.1	0.1	0.2	0.3	0.3	0.6	0.7
White, male										
2006 4/.....	9.3	5.3	0.5	*	*	*	*	0.3	0.4	0.7
2005 4/.....	8.9	6.2	0.4	*	*	*	*	0.3	0.4	0.5
2004 4/.....	9.3	4.2	0.5	*	*	*	0.3	*	0.3	0.5
2003 4/.....	8.6	4.9	0.3	*	*	*	0.3	0.3	0.5	0.6
2002.....	8.4	5.0	0.5	*	*	*	0.3	0.3	0.4	0.6
2001.....	8.8	5.3	0.3	*	*	*	0.3	*	0.3	0.7
2000.....	8.8	5.8	0.4	*	*	*	0.3	0.3	0.5	0.8
1999.....	8.6	4.8	0.4	*	*	0.3	0.3	0.4	0.5	0.7
White, female										
2006 4/.....	9.2	4.5	*	*	*	*	*	0.4	0.6	0.6
2005 4/.....	9.2	4.4	*	*	*	*	0.3	*	0.5	0.6
2004 4/.....	9.4	4.2	*	*	*	*	0.3	0.3	0.4	0.7
2003 4/.....	8.9	4.4	*	*	*	*	*	*	0.5	0.6
2002.....	8.8	3.5	*	*	*	*	*	0.3	0.4	0.7
2001.....	9.3	3.5	0.4	*	*	*	0.3	0.4	0.4	0.6
2000.....	9.2	4.1	*	*	*	*	*	0.4	0.5	0.6
1999.....	9.0	3.4	*	*	*	*	*	0.3	0.6	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other diseases of respiratory system (J80-J84,J93-J98)										
Black, both sexes										
2006 4/.....	5.4	17.5	*	*	*	*	*	*	0.7	1.2
2005 4/.....	5.6	14.0	*	*	*	*	*	*	0.9	1.3
2004 4/.....	5.5	14.1	0.8	*	*	*	*	*	0.9	1.9
2003 4/.....	5.1	14.1	*	*	*	*	0.7	*	1.0	1.6
2002.....	5.3	14.8	0.9	*	*	*	0.7	*	1.0	1.5
2001.....	5.8	16.6	*	*	*	*	*	0.9	1.2	2.1
2000.....	5.7	15.1	*	*	*	*	*	*	1.1	2.1
1999.....	5.6	14.5	*	*	*	*	*	0.8	1.4	1.8
Black, male										
2006 4/.....	5.4	20.7	*	*	*	*	*	*	*	*
2005 4/.....	5.6	18.6	*	*	*	*	*	*	*	*
2004 4/.....	5.6	14.8	*	*	*	*	*	*	*	2.4
2003 4/.....	5.2	17.6	*	*	*	*	*	*	*	1.6
2002.....	5.2	17.7	*	*	*	*	*	*	*	1.5
2001.....	5.4	21.3	*	*	*	*	*	*	*	1.5
2000.....	5.6	15.0	*	*	*	*	*	*	*	2.6
1999.....	5.3	16.9	*	*	*	*	*	*	*	1.7
Black, female										
2006 4/.....	5.4	14.1	*	*	*	*	*	*	*	*
2005 4/.....	5.7	9.2	*	*	*	*	*	*	*	1.7
2004 4/.....	5.5	13.3	*	*	*	*	*	*	*	1.5
2003 4/.....	5.0	10.5	*	*	*	*	*	*	*	1.6
2002.....	5.4	11.8	*	*	*	*	*	*	*	1.6
2001.....	6.2	11.7	*	*	*	*	*	*	*	2.6
2000.....	5.7	15.2	*	*	*	*	*	*	*	1.7
1999.....	5.9	12.1	*	*	*	*	*	*	1.5	1.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XI. Diseases of the digestive system (K00-K92)

All races 3/, both sexes

2006.....	29.3	14.1	0.7	0.2	0.3	0.4	0.9	1.5	2.9	6.4
2005.....	29.5	15.2	0.8	0.3	0.3	0.4	0.8	1.5	2.8	6.2
2004.....	29.4	14.2	0.8	0.2	0.3	0.4	0.9	1.3	2.9	7.0
2003.....	30.2	13.8	0.8	0.3	0.3	0.5	0.8	1.6	3.2	7.3
2002.....	30.1	12.6	0.8	0.2	0.3	0.4	0.9	1.4	3.2	7.3
2001.....	30.1	11.9	0.8	0.3	0.3	0.4	0.7	1.4	3.2	8.1
2000.....	29.9	13.1	0.7	0.3	0.3	0.4	1.0	1.4	3.3	8.0
1999.....	29.7	13.2	0.7	0.3	0.4	0.5	0.7	1.3	3.5	7.9

All races 3/, male

2006.....	30.1	15.8	0.7	0.2	0.3	0.4	0.9	1.9	3.5	8.0
2005.....	30.4	18.5	1.0	0.3	0.3	0.4	0.8	1.7	3.3	7.8
2004.....	30.0	15.6	0.9	0.3	0.2	0.4	0.9	1.6	3.9	9.0
2003.....	30.9	16.6	0.9	0.3	0.3	0.6	0.8	1.8	4.0	9.0
2002.....	30.7	15.1	0.9	0.3	0.3	0.4	0.9	1.5	4.1	9.1
2001.....	30.6	13.4	0.9	0.3	0.4	0.4	0.8	1.6	4.0	10.3
2000.....	30.3	14.8	0.9	0.3	0.3	0.4	1.1	1.8	4.1	10.2
1999.....	30.4	16.5	0.7	0.2	0.4	0.5	0.7	1.5	4.3	9.9

All races 3/, female

2006.....	28.4	12.3	0.7	0.2	0.3	0.3	0.8	1.1	2.4	4.7
2005.....	28.6	11.9	0.6	0.3	0.2	0.4	0.7	1.2	2.2	4.7
2004.....	28.8	12.8	0.6	*	0.3	0.5	0.8	1.0	2.0	5.0
2003.....	29.4	10.8	0.7	0.2	0.3	0.4	0.7	1.3	2.3	5.6
2002.....	29.5	10.0	0.7	*	0.4	0.4	0.8	1.3	2.3	5.6
2001.....	29.7	10.3	0.6	0.3	0.3	0.3	0.7	1.2	2.4	6.0
2000.....	29.4	11.4	0.5	0.3	0.4	0.3	0.8	1.0	2.5	5.8
1999.....	29.1	9.7	0.8	0.3	0.4	0.4	0.8	1.1	2.7	5.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XI. Diseases of the digestive system (K00-K92)										
White, both sexes										
2006 4/.....	31.5	10.3	0.6	0.2	0.3	0.3	0.8	1.4	2.9	6.4
2005 4/.....	31.6	12.2	0.7	0.3	0.2	0.4	0.6	1.4	2.6	6.2
2004 4/.....	31.5	10.7	0.6	0.2	0.2	0.4	0.7	1.1	2.8	6.8
2003 4/.....	32.3	11.0	0.7	0.3	0.3	0.5	0.6	1.3	3.1	7.1
2002.....	32.2	10.1	0.7	0.2	0.3	0.4	0.8	1.2	3.0	7.2
2001.....	32.1	8.7	0.7	0.3	0.3	0.3	0.6	1.2	3.0	7.9
2000.....	31.7	10.0	0.5	0.2	0.3	0.3	0.9	1.2	3.1	7.6
1999.....	31.5	10.1	0.7	0.2	0.4	0.4	0.6	1.2	3.3	7.6
White, male										
2006 4/.....	32.1	11.4	0.6	*	0.3	0.4	0.9	1.8	3.5	8.1
2005 4/.....	32.2	14.6	0.8	*	*	0.4	0.7	1.7	3.2	7.9
2004 4/.....	31.6	11.4	0.7	0.3	*	0.4	0.7	1.4	3.8	8.6
2003 4/.....	32.6	13.9	0.9	0.3	0.3	0.6	0.7	1.5	4.0	8.9
2002.....	32.4	12.1	0.7	0.3	0.3	0.4	0.9	1.2	4.0	9.0
2001.....	31.9	9.3	0.8	0.3	0.3	0.4	0.6	1.4	3.8	9.9
2000.....	31.7	11.2	0.6	0.3	0.3	0.4	1.0	1.7	3.9	9.8
1999.....	31.7	12.5	0.5	*	0.3	0.5	0.6	1.4	4.0	9.7
White, female										
2006 4/.....	31.0	9.0	0.6	*	0.3	0.2	0.7	1.1	2.3	4.6
2005 4/.....	31.0	9.7	0.6	0.3	*	0.4	0.6	1.2	2.0	4.6
2004 4/.....	31.3	9.9	0.5	*	*	0.4	0.6	0.8	1.9	4.9
2003 4/.....	32.0	8.0	0.6	*	0.3	0.3	0.5	1.1	2.2	5.3
2002.....	32.1	7.9	0.7	*	0.3	0.4	0.6	1.1	1.9	5.4
2001.....	32.2	8.1	0.5	0.3	0.3	0.3	0.5	1.1	2.3	5.9
2000.....	31.6	8.8	0.4	*	0.3	0.3	0.7	0.7	2.2	5.3
1999.....	31.2	7.5	0.8	0.3	0.4	0.4	0.6	1.0	2.5	5.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XI. Diseases of the digestive system (K00-K92)										
Black, both sexes										
2006 4/.....	22.2	33.2	0.9	*	*	0.6	1.3	1.9	2.9	6.7
2005 4/.....	23.3	32.1	1.1	*	0.7	*	1.6	1.7	3.6	6.5
2004 4/.....	23.7	31.6	1.4	*	*	0.7	1.7	2.2	3.6	8.6
2003 4/.....	24.2	29.0	1.2	*	*	0.7	1.4	3.1	3.6	8.9
2002.....	23.7	25.9	1.4	*	*	0.6	1.5	2.5	4.3	8.0
2001.....	25.1	28.6	1.2	*	0.6	0.6	1.5	2.6	4.3	10.1
2000.....	25.2	29.6	1.5	*	*	*	1.6	2.9	4.7	10.8
1999.....	25.1	29.6	1.1	*	*	0.7	1.8	2.1	4.6	10.1
Black, male										
2006 4/.....	24.3	37.2	*	*	*	*	1.2	2.4	3.1	8.1
2005 4/.....	25.7	41.3	1.5	*	*	*	1.8	1.6	3.7	7.7
2004 4/.....	26.4	36.2	1.7	*	*	*	1.6	2.5	5.0	11.4
2003 4/.....	27.1	32.4	*	*	*	*	1.6	3.6	4.5	10.0
2002.....	26.0	31.4	1.7	*	*	*	*	2.9	4.9	9.1
2001.....	28.4	33.5	*	*	*	*	*	3.3	5.3	13.2
2000.....	27.7	33.9	2.0	*	*	*	2.1	3.0	5.1	13.6
1999.....	27.8	36.4	*	*	*	*	1.6	2.2	5.7	12.2
Black, female										
2006 4/.....	20.2	29.2	*	*	*	*	1.4	1.4	2.8	5.6
2005 4/.....	21.0	22.5	*	*	*	*	1.4	1.9	3.5	5.3
2004 4/.....	21.1	26.9	*	*	*	*	1.7	2.0	2.3	6.2
2003 4/.....	21.6	25.4	*	*	*	*	1.3	2.7	2.9	7.9
2002.....	21.5	20.3	*	*	*	*	1.8	2.1	3.8	7.0
2001.....	22.1	23.3	*	*	*	*	1.7	2.0	3.4	7.3
2000.....	23.0	25.1	*	*	*	*	*	2.8	4.3	8.4
1999.....	22.7	22.5	*	*	*	*	1.9	2.0	3.6	8.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of other parts of digestive system (K20-K92)										
All races 3/, both sexes										
2006.....	29.2	14.0	0.7	0.2	0.3	0.4	0.9	1.5	2.9	6.4
2005.....	29.4	15.2	0.8	0.3	0.3	0.4	0.7	1.5	2.8	6.2
2004.....	29.3	14.1	0.8	0.2	0.3	0.4	0.9	1.3	2.9	7.0
2003.....	30.1	13.7	0.8	0.3	0.3	0.5	0.7	1.5	3.2	7.3
2002.....	30.0	12.5	0.8	0.2	0.3	0.4	0.9	1.4	3.2	7.3
2001.....	30.1	11.8	0.8	0.3	0.3	0.4	0.7	1.4	3.2	8.1
2000.....	29.8	13.0	0.7	0.3	0.3	0.4	1.0	1.4	3.3	8.0
1999.....	29.7	13.1	0.7	0.3	0.4	0.5	0.7	1.3	3.5	7.9
All races 3/, male										
2006.....	30.1	15.7	0.7	0.2	0.3	0.4	0.9	1.9	3.5	8.0
2005.....	30.3	18.4	1.0	0.3	0.3	0.4	0.8	1.7	3.3	7.7
2004.....	29.9	15.5	0.9	0.3	0.2	0.4	0.9	1.6	3.9	8.9
2003.....	30.9	16.5	0.9	0.3	0.3	0.6	0.8	1.8	4.0	9.0
2002.....	30.6	15.0	0.9	0.3	0.3	0.4	0.9	1.5	4.1	9.1
2001.....	30.5	13.3	0.9	0.3	0.4	0.4	0.8	1.6	4.0	10.3
2000.....	30.3	14.8	0.9	0.3	0.3	0.4	1.1	1.8	4.1	10.2
1999.....	30.3	16.4	0.7	0.2	0.4	0.5	0.7	1.5	4.3	9.9
All races 3/, female										
2006.....	28.3	12.2	0.7	0.2	0.3	0.3	0.8	1.1	2.3	4.7
2005.....	28.5	11.8	0.6	0.2	0.2	0.4	0.7	1.2	2.2	4.7
2004.....	28.7	12.6	0.6	*	0.3	0.5	0.8	1.0	2.0	5.0
2003.....	29.4	10.8	0.7	0.2	0.3	0.4	0.6	1.3	2.3	5.6
2002.....	29.4	9.9	0.7	*	0.4	0.4	0.8	1.3	2.3	5.6
2001.....	29.7	10.2	0.6	0.3	0.3	0.3	0.7	1.2	2.4	6.0
2000.....	29.4	11.2	0.5	0.3	0.4	0.3	0.8	1.0	2.5	5.8
1999.....	29.0	9.7	0.8	0.3	0.4	0.4	0.7	1.1	2.7	5.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of other parts of digestive system (K20-K92)										
White, both sexes										
2006 4/.....	31.5	10.1	0.6	0.2	0.3	0.3	0.8	1.4	2.9	6.4
2005 4/.....	31.5	12.1	0.7	0.3	0.2	0.4	0.6	1.4	2.6	6.2
2004 4/.....	31.4	10.5	0.6	0.2	0.2	0.4	0.7	1.1	2.8	6.8
2003 4/.....	32.2	11.0	0.7	0.3	0.3	0.5	0.6	1.3	3.1	7.1
2002.....	32.2	10.0	0.7	0.2	0.3	0.4	0.8	1.2	3.0	7.2
2001.....	32.0	8.6	0.7	0.3	0.3	0.3	0.5	1.2	3.0	7.9
2000.....	31.6	9.9	0.5	0.2	0.3	0.3	0.9	1.2	3.0	7.6
1999.....	31.4	10.0	0.7	0.2	0.4	0.4	0.6	1.2	3.3	7.6
White, male										
2006 4/.....	32.0	11.3	0.6	*	0.3	0.4	0.9	1.8	3.4	8.1
2005 4/.....	32.1	14.5	0.8	*	*	0.4	0.7	1.7	3.1	7.9
2004 4/.....	31.6	11.3	0.7	0.3	*	0.4	0.7	1.4	3.7	8.6
2003 4/.....	32.6	13.8	0.9	0.3	0.3	0.6	0.7	1.5	4.0	8.9
2002.....	32.4	12.0	0.7	0.3	0.3	0.4	0.9	1.2	4.0	9.0
2001.....	31.9	9.3	0.8	0.3	0.3	0.4	0.6	1.4	3.8	9.9
2000.....	31.7	11.2	0.6	0.3	0.3	0.4	1.0	1.6	3.9	9.8
1999.....	31.7	12.4	0.5	*	0.3	0.5	0.6	1.4	4.0	9.7
White, female										
2006 4/.....	30.9	8.9	0.6	*	0.3	0.2	0.7	1.1	2.3	4.6
2005 4/.....	30.9	9.6	0.6	0.3	*	0.4	0.6	1.2	2.0	4.6
2004 4/.....	31.2	9.7	0.5	*	*	0.4	0.6	0.8	1.9	4.9
2003 4/.....	31.9	8.0	0.6	*	0.3	0.3	0.5	1.1	2.2	5.3
2002.....	32.0	7.9	0.7	*	0.3	0.4	0.6	1.1	1.9	5.4
2001.....	32.1	8.0	0.5	0.3	0.3	0.3	0.5	1.0	2.3	5.9
2000.....	31.6	8.6	0.4	*	0.3	0.3	0.7	0.7	2.2	5.3
1999.....	31.2	7.5	0.8	0.3	0.4	0.4	0.6	1.0	2.5	5.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of other parts of digestive system (K20-K92)										
Black, both sexes										
2006 4/.....	22.1	33.2	0.9	*	*	0.6	1.3	1.9	2.9	6.7
2005 4/.....	23.2	32.1	1.1	*	0.7	*	1.6	1.7	3.6	6.5
2004 4/.....	23.6	31.6	1.4	*	*	0.7	1.7	2.1	3.5	8.6
2003 4/.....	24.2	28.8	1.2	*	*	0.7	1.4	3.1	3.6	8.9
2002.....	23.6	25.9	1.4	*	*	0.6	1.4	2.5	4.2	7.9
2001.....	25.0	28.6	1.2	*	0.6	0.6	1.5	2.6	4.3	10.1
2000.....	25.2	29.4	1.5	*	*	*	1.6	2.9	4.6	10.8
1999.....	25.1	29.6	1.1	*	*	0.7	1.8	2.1	4.6	10.1
Black, male										
2006 4/.....	24.3	37.2	*	*	*	*	1.2	2.4	3.1	8.1
2005 4/.....	25.7	41.3	1.5	*	*	*	1.8	1.6	3.7	7.7
2004 4/.....	26.4	36.2	1.7	*	*	*	1.6	2.5	4.9	11.3
2003 4/.....	27.1	32.4	*	*	*	*	1.5	3.5	4.5	10.0
2002.....	25.9	31.4	1.7	*	*	*	*	2.9	4.8	9.0
2001.....	28.3	33.5	*	*	*	*	*	3.2	5.3	13.2
2000.....	27.7	33.9	2.0	*	*	*	2.1	3.0	5.1	13.6
1999.....	27.7	36.4	*	*	*	*	1.6	2.2	5.7	12.2
Black, female										
2006 4/.....	20.2	29.2	*	*	*	*	1.3	1.4	2.8	5.6
2005 4/.....	20.9	22.5	*	*	*	*	1.4	1.8	3.5	5.3
2004 4/.....	21.1	26.9	*	*	*	*	1.7	1.8	2.3	6.1
2003 4/.....	21.5	25.0	*	*	*	*	1.3	2.7	2.9	7.9
2002.....	21.5	20.3	*	*	*	*	1.7	2.1	3.7	7.0
2001.....	22.0	23.3	*	*	*	*	1.7	2.0	3.4	7.3
2000.....	22.9	24.8	*	*	*	*	*	2.8	4.3	8.4
1999.....	22.6	22.5	*	*	*	*	1.9	2.0	3.6	8.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Vascular disorders and obstruction of intestine without hernia (K55-K56)										
All races 3/, both sexes										
2006.....	4.8	1.7	*	*	*	*	0.1	0.1	0.1	0.3
2005.....	4.9	3.1	0.2	*	*	*	0.1	0.1	0.1	0.2
2004.....	4.9	6.9	0.2	*	*	*	*	*	0.2	0.3
2003.....	5.1	6.8	0.1	*	*	*	0.1	0.1	0.1	0.2
2002.....	5.0	5.8	0.1	*	*	*	0.1	0.1	0.1	0.3
2001.....	5.0	5.8	0.1	*	*	*	*	0.1	0.1	0.2
2000.....	5.1	6.0	0.1	*	*	*	0.1	0.1	0.1	0.3
1999.....	5.0	6.1	0.1	*	*	*	*	0.1	0.2	0.2
All races 3/, male										
2006.....	3.5	1.6	*	*	*	*	*	*	*	0.3
2005.....	3.5	3.9	*	*	*	*	*	*	*	0.2
2004.....	3.5	7.7	0.2	*	*	*	*	*	*	0.3
2003.....	3.7	8.2	*	*	*	*	*	*	*	0.2
2002.....	3.6	6.5	*	*	*	*	*	*	0.2	0.3
2001.....	3.6	6.5	*	*	*	*	*	*	*	0.2
2000.....	3.6	6.3	*	*	*	*	*	*	*	0.3
1999.....	3.5	7.6	*	*	*	*	*	*	*	0.2
All races 3/, female										
2006.....	6.1	1.8	*	*	*	*	*	*	*	0.3
2005.....	6.2	2.3	*	*	*	*	*	*	*	0.3
2004.....	6.3	6.0	*	*	*	*	*	*	0.2	0.2
2003.....	6.5	5.3	*	*	*	*	*	*	*	0.2
2002.....	6.4	5.0	*	*	*	*	*	*	*	0.2
2001.....	6.4	5.1	*	*	*	*	*	*	*	0.2
2000.....	6.5	5.8	*	*	*	*	*	*	*	0.2
1999.....	6.4	4.4	*	*	*	*	*	*	0.2	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of liver (K70-K76)										
All races 3/, both sexes										
2006.....	12.4	2.0	0.2	*	*	0.1	0.3	0.7	1.9	4.4
2005.....	12.4	2.3	0.2	*	*	0.1	0.3	0.7	1.7	4.4
2004.....	12.3	2.5	0.2	*	*	0.1	0.3	0.6	1.8	4.9
2003.....	12.6	1.5	0.2	*	*	0.2	0.2	0.7	2.0	5.3
2002.....	12.4	2.2	0.2	*	*	0.1	0.3	0.7	1.9	5.3
2001.....	12.2	1.9	0.2	*	*	0.1	0.3	0.6	2.0	6.1
2000.....	12.0	2.0	0.2	*	*	*	0.4	0.6	2.1	6.0
1999.....	12.0	2.3	0.1	*	*	0.2	0.3	0.6	2.2	6.0
All races 3/, male										
2006.....	15.8	2.0	*	*	*	*	0.3	0.8	2.2	5.5
2005.....	15.9	2.9	*	*	*	*	0.3	0.8	2.1	5.6
2004.....	15.7	2.5	*	*	*	*	0.3	0.7	2.5	6.3
2003.....	16.1	2.2	0.2	*	*	*	0.2	0.9	2.5	6.7
2002.....	15.8	3.1	*	*	*	*	0.3	0.8	2.4	6.8
2001.....	15.7	2.3	*	*	*	*	0.3	0.7	2.5	7.8
2000.....	15.5	2.7	*	*	*	*	0.4	0.8	2.6	7.8
1999.....	15.6	2.9	*	*	*	*	*	0.6	2.8	7.6
All races 3/, female										
2006.....	9.1	1.9	*	*	*	*	0.4	0.5	1.5	3.3
2005.....	9.0	1.7	*	*	*	0.2	0.3	0.6	1.3	3.2
2004.....	9.0	2.4	*	*	*	*	0.3	0.4	1.2	3.5
2003.....	9.1	*	*	*	*	*	0.3	0.6	1.4	4.0
2002.....	9.0	1.3	*	*	*	*	0.3	0.6	1.4	3.9
2001.....	8.9	1.6	*	*	*	*	0.3	0.5	1.5	4.4
2000.....	8.7	1.3	*	*	*	*	0.3	0.4	1.6	4.2
1999.....	8.6	1.6	*	*	*	*	0.3	0.5	1.6	4.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of liver (K70-K76)										
White, both sexes										
2006 4/.....	13.3	1.7	0.2	*	*	*	0.3	0.6	1.9	4.5
2005 4/.....	13.2	2.1	*	*	*	*	0.2	0.7	1.6	4.6
2004 4/.....	13.0	1.9	0.2	*	*	*	0.2	0.5	1.9	4.9
2003 4/.....	13.3	1.2	0.2	*	*	*	0.2	0.6	2.0	5.5
2002.....	13.1	1.9	*	*	*	*	0.2	0.6	1.9	5.4
2001.....	12.9	1.6	0.2	*	*	*	0.2	0.5	2.0	6.1
2000.....	12.6	1.6	*	*	*	*	0.3	0.5	2.0	5.8
1999.....	12.5	1.7	*	*	*	0.1	0.2	0.5	2.1	5.9
White, male										
2006 4/.....	16.9	2.0	*	*	*	*	0.3	0.7	2.3	5.8
2005 4/.....	17.0	2.4	*	*	*	*	0.2	0.8	2.1	6.0
2004 4/.....	16.6	1.8	*	*	*	*	0.3	0.6	2.6	6.2
2003 4/.....	17.1	1.8	*	*	*	*	*	0.7	2.6	6.9
2002.....	16.8	2.7	*	*	*	*	0.3	0.7	2.4	6.9
2001.....	16.4	1.9	*	*	*	*	*	0.6	2.5	7.7
2000.....	16.2	2.0	*	*	*	*	0.3	0.7	2.6	7.7
1999.....	16.3	2.0	*	*	*	*	*	0.7	2.8	7.6
White, female										
2006 4/.....	9.7	1.4	*	*	*	*	0.3	0.5	1.5	3.2
2005 4/.....	9.5	1.7	*	*	*	*	*	0.5	1.1	3.2
2004 4/.....	9.5	1.9	*	*	*	*	*	0.4	1.2	3.5
2003 4/.....	9.7	*	*	*	*	*	*	0.5	1.3	4.0
2002.....	9.5	*	*	*	*	*	*	0.5	1.3	3.9
2001.....	9.4	1.3	*	*	*	*	*	0.4	1.5	4.5
2000.....	9.1	*	*	*	*	*	*	0.3	1.4	3.9
1999.....	8.9	1.4	*	*	*	*	*	0.4	1.5	4.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of liver (K70-K76)										
Black, both sexes										
2006 4/.....	8.9	3.4	*	*	*	*	*	0.7	1.1	3.5
2005 4/.....	9.6	3.9	*	*	*	*	*	0.7	1.9	3.2
2004 4/.....	9.8	5.4	*	*	*	*	0.6	0.7	1.4	4.8
2003 4/.....	10.0	3.0	*	*	*	*	0.6	1.4	1.8	5.0
2002.....	9.9	3.7	*	*	*	*	*	1.0	1.9	4.5
2001.....	10.4	3.4	*	*	*	*	*	0.9	1.9	5.8
2000.....	10.4	4.2	*	*	*	*	0.7	1.2	2.3	6.9
1999.....	10.8	5.0	*	*	*	*	*	0.9	2.5	6.3
Black, male										
2006 4/.....	11.5	*	*	*	*	*	*	*	*	3.7
2005 4/.....	12.4	*	*	*	*	*	*	*	1.6	3.6
2004 4/.....	12.9	*	*	*	*	*	*	*	1.8	6.1
2003 4/.....	13.3	*	*	*	*	*	*	1.6	1.9	5.8
2002.....	12.6	*	*	*	*	*	*	*	2.2	4.8
2001.....	14.0	*	*	*	*	*	*	*	2.4	7.8
2000.....	13.5	6.4	*	*	*	*	*	*	2.4	8.9
1999.....	14.1	7.2	*	*	*	*	*	*	2.7	7.6
Black, female										
2006 4/.....	6.6	*	*	*	*	*	*	*	*	3.3
2005 4/.....	6.9	*	*	*	*	*	*	*	2.2	2.9
2004 4/.....	7.0	*	*	*	*	*	*	*	*	3.6
2003 4/.....	7.1	*	*	*	*	*	*	*	1.7	4.3
2002.....	7.4	*	*	*	*	*	*	*	1.7	4.2
2001.....	7.2	*	*	*	*	*	*	*	1.5	4.0
2000.....	7.7	*	*	*	*	*	*	*	2.3	5.1
1999.....	7.9	*	*	*	*	*	*	*	2.3	5.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcoholic liver disease (K70)										
All races 3/, both sexes										
2006.....	4.4	*	*	*	*	*	*	0.3	1.0	2.4
2005.....	4.4	*	*	*	*	*	*	0.3	0.8	2.4
2004.....	4.3	*	*	*	*	*	*	0.2	0.9	2.5
2003.....	4.3	*	*	*	*	*	*	0.3	0.9	2.9
2002.....	4.2	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3
2000.....	4.3	*	*	*	*	*	*	0.3	1.1	3.3
1999.....	4.3	*	*	*	*	*	*	0.2	1.1	3.3
All races 3/, male										
2006.....	6.4	*	*	*	*	*	*	0.4	1.4	3.1
2005.....	6.5	*	*	*	*	*	*	0.4	1.0	3.2
2004.....	6.3	*	*	*	*	*	*	0.3	1.3	3.3
2003.....	6.4	*	*	*	*	*	*	0.4	1.4	3.7
2002.....	6.2	*	*	*	*	*	*	0.3	1.2	3.6
2001.....	6.4	*	*	*	*	*	*	0.3	1.4	4.4
2000.....	6.5	*	*	*	*	*	*	0.5	1.5	4.3
1999.....	6.5	*	*	*	*	*	*	0.2	1.4	4.3
All races 3/, female										
2006.....	2.4	*	*	*	*	*	*	*	0.6	1.7
2005.....	2.3	*	*	*	*	*	*	*	0.6	1.6
2004.....	2.3	*	*	*	*	*	*	*	0.5	1.8
2003.....	2.2	*	*	*	*	*	*	*	0.5	2.0
2002.....	2.3	*	*	*	*	*	*	*	0.7	2.0
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3
2000.....	2.2	*	*	*	*	*	*	*	0.6	2.2
1999.....	2.1	*	*	*	*	*	*	*	0.8	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Alcoholic liver disease (K70)										
White, both sexes										
2006 4/.....	4.7	*	*	*	*	*	*	0.2	1.0	2.5
2005 4/.....	4.7	*	*	*	*	*	*	0.3	0.8	2.6
2004 4/.....	4.6	*	*	*	*	*	*	0.2	1.0	2.6
2003 4/.....	4.5	*	*	*	*	*	*	0.2	1.0	2.9
2002.....	4.4	*	*	*	*	*	*	0.2	0.9	2.8
2001.....	4.5	*	*	*	*	*	*	0.2	1.1	3.4
2000.....	4.5	*	*	*	*	*	*	0.3	1.1	3.2
1999.....	4.4	*	*	*	*	*	*	0.2	1.0	3.3
White, male										
2006 4/.....	7.0	*	*	*	*	*	*	0.3	1.4	3.3
2005 4/.....	6.9	*	*	*	*	*	*	0.4	1.0	3.4
2004 4/.....	6.8	*	*	*	*	*	*	0.3	1.5	3.3
2003 4/.....	6.8	*	*	*	*	*	*	0.4	1.4	3.9
2002.....	6.6	*	*	*	*	*	*	0.3	1.3	3.7
2001.....	6.7	*	*	*	*	*	*	0.3	1.5	4.5
2000.....	6.8	*	*	*	*	*	*	0.5	1.6	4.3
1999.....	6.7	*	*	*	*	*	*	0.3	1.4	4.3
White, female										
2006 4/.....	2.5	*	*	*	*	*	*	*	0.6	1.7
2005 4/.....	2.4	*	*	*	*	*	*	*	0.5	1.7
2004 4/.....	2.4	*	*	*	*	*	*	*	0.5	1.9
2003 4/.....	2.3	*	*	*	*	*	*	*	0.5	1.9
2002.....	2.3	*	*	*	*	*	*	*	0.6	2.0
2001.....	2.3	*	*	*	*	*	*	*	0.7	2.3
2000.....	2.1	*	*	*	*	*	*	*	0.5	2.2
1999.....	2.1	*	*	*	*	*	*	*	0.7	2.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Fibrosis and cirrhosis of liver (K74)										
All races 3/, both sexes										
2006.....	4.8	*	*	*	*	*	*	*	0.3	0.9
2005.....	4.9	*	*	*	*	*	*	*	0.4	1.0
2004.....	4.9	*	*	*	*	*	*	*	0.3	1.2
2003.....	5.2	*	*	*	*	*	*	0.2	0.4	1.2
2002.....	5.1	*	*	*	*	*	*	0.1	0.4	1.4
2001.....	5.0	*	*	*	*	*	*	0.1	0.4	1.6
2000.....	5.0	*	*	*	*	*	*	0.1	0.5	1.5
1999.....	5.0	*	*	*	*	*	*	0.1	0.5	1.4
All races 3/, male										
2006.....	5.7	*	*	*	*	*	*	*	0.3	1.2
2005.....	5.8	*	*	*	*	*	*	*	0.4	1.3
2004.....	5.8	*	*	*	*	*	*	*	0.5	1.7
2003.....	6.1	*	*	*	*	*	*	*	0.5	1.6
2002.....	6.0	*	*	*	*	*	*	*	0.5	1.9
2001.....	6.0	*	*	*	*	*	*	*	0.5	1.9
2000.....	5.9	*	*	*	*	*	*	*	0.6	2.0
1999.....	5.9	*	*	*	*	*	*	*	0.7	2.0
All races 3/, female										
2006.....	4.0	*	*	*	*	*	*	*	0.2	0.7
2005.....	4.0	*	*	*	*	*	*	*	0.3	0.7
2004.....	4.0	*	*	*	*	*	*	*	0.2	0.7
2003.....	4.2	*	*	*	*	*	*	*	0.3	0.9
2002.....	4.2	*	*	*	*	*	*	*	0.3	0.9
2001.....	4.2	*	*	*	*	*	*	*	0.3	1.2
2000.....	4.1	*	*	*	*	*	*	*	0.4	0.9
1999.....	4.1	*	*	*	*	*	*	*	0.4	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Fibrosis and cirrhosis of liver (K74)										
White, both sexes										
2006 4/.....	5.2	*	*	*	*	*	*	*	0.3	1.0
2005 4/.....	5.3	*	*	*	*	*	*	*	0.3	1.1
2004 4/.....	5.3	*	*	*	*	*	*	*	0.3	1.2
2003 4/.....	5.6	*	*	*	*	*	*	*	0.4	1.3
2002.....	5.5	*	*	*	*	*	*	*	0.4	1.4
2001.....	5.4	*	*	*	*	*	*	*	0.4	1.6
2000.....	5.4	*	*	*	*	*	*	*	0.4	1.5
1999.....	5.3	*	*	*	*	*	*	*	0.6	1.5
White, male										
2006 4/.....	6.1	*	*	*	*	*	*	*	0.4	1.3
2005 4/.....	6.2	*	*	*	*	*	*	*	0.4	1.4
2004 4/.....	6.2	*	*	*	*	*	*	*	0.5	1.7
2003 4/.....	6.6	*	*	*	*	*	*	*	0.6	1.7
2002.....	6.4	*	*	*	*	*	*	*	0.6	2.0
2001.....	6.3	*	*	*	*	*	*	*	0.5	2.0
2000.....	6.3	*	*	*	*	*	*	*	0.5	2.0
1999.....	6.3	*	*	*	*	*	*	*	0.7	2.0
White, female										
2006 4/.....	4.4	*	*	*	*	*	*	*	0.3	0.6
2005 4/.....	4.4	*	*	*	*	*	*	*	0.3	0.7
2004 4/.....	4.4	*	*	*	*	*	*	*	*	0.7
2003 4/.....	4.7	*	*	*	*	*	*	*	0.3	0.9
2002.....	4.6	*	*	*	*	*	*	*	0.3	0.9
2001.....	4.6	*	*	*	*	*	*	*	0.3	1.3
2000.....	4.5	*	*	*	*	*	*	*	0.3	0.9
1999.....	4.4	*	*	*	*	*	*	*	0.4	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of liver (K71-K72,K75-K76)										
All races 3/, both sexes										
2006.....	3.2	1.8	0.1	*	*	0.1	0.2	0.3	0.6	1.1
2005.....	3.1	2.1	0.2	*	*	0.1	0.2	0.3	0.5	1.0
2004.....	3.1	2.3	0.2	*	*	0.1	0.2	0.3	0.6	1.2
2003.....	3.1	1.5	0.2	*	*	0.1	0.2	0.3	0.6	1.2
2002.....	2.9	2.1	0.2	*	*	*	0.2	0.3	0.5	1.1
2001.....	2.7	1.9	0.2	*	*	*	0.2	0.2	0.5	1.1
2000.....	2.6	1.9	0.2	*	*	*	0.2	0.2	0.5	1.2
1999.....	2.6	2.1	0.1	*	*	0.1	0.2	0.2	0.6	1.2
All races 3/, male										
2006.....	3.6	1.9	*	*	*	*	*	0.3	0.5	1.2
2005.....	3.7	2.6	*	*	*	*	*	0.3	0.7	1.1
2004.....	3.5	2.3	*	*	*	*	0.2	0.4	0.7	1.3
2003.....	3.6	2.1	0.2	*	*	*	*	0.3	0.6	1.3
2002.....	3.5	2.8	*	*	*	*	*	0.3	0.6	1.2
2001.....	3.2	2.2	*	*	*	*	*	*	0.5	1.4
2000.....	3.0	2.5	*	*	*	*	0.2	*	0.5	1.5
1999.....	3.1	2.7	*	*	*	*	*	0.2	0.7	1.4
All races 3/, female										
2006.....	2.7	1.8	*	*	*	*	0.3	0.3	0.7	1.0
2005.....	2.6	1.5	*	*	*	*	0.2	0.3	0.4	0.8
2004.....	2.6	2.3	*	*	*	*	*	0.3	0.5	1.0
2003.....	2.6	*	*	*	*	*	0.2	0.3	0.6	1.1
2002.....	2.3	1.3	*	*	*	*	*	0.3	0.4	1.0
2001.....	2.3	1.5	*	*	*	*	*	0.2	0.5	0.9
2000.....	2.2	1.3	*	*	*	*	*	*	0.5	0.9
1999.....	2.2	1.5	*	*	*	*	0.2	*	0.4	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other diseases of liver (K71-K72,K75-K76)										
White, both sexes										
2006 4/.....	3.3	1.6	*	*	*	*	0.2	0.3	0.6	1.0
2005 4/.....	3.2	1.8	*	*	*	*	0.2	0.3	0.5	1.0
2004 4/.....	3.1	1.6	0.2	*	*	*	0.2	0.2	0.6	1.1
2003 4/.....	3.2	1.2	0.2	*	*	*	0.1	0.3	0.6	1.2
2002.....	3.0	1.8	*	*	*	*	*	0.3	0.5	1.1
2001.....	2.8	1.5	0.2	*	*	*	0.1	0.2	0.4	1.0
2000.....	2.6	1.6	*	*	*	*	0.2	0.1	0.5	1.1
1999.....	2.6	1.5	*	*	*	*	0.1	0.2	0.5	1.0
White, male										
2006 4/.....	3.8	1.9	*	*	*	*	*	0.3	0.5	1.2
2005 4/.....	3.8	2.1	*	*	*	*	*	0.2	0.7	1.1
2004 4/.....	3.6	1.6	*	*	*	*	*	0.3	0.6	1.2
2003 4/.....	3.7	1.7	*	*	*	*	*	*	0.6	1.3
2002.....	3.6	2.4	*	*	*	*	*	*	0.6	1.3
2001.....	3.3	1.9	*	*	*	*	*	*	0.4	1.2
2000.....	3.0	2.0	*	*	*	*	*	*	0.5	1.4
1999.....	3.1	1.7	*	*	*	*	*	*	0.7	1.3
White, female										
2006 4/.....	2.8	1.3	*	*	*	*	*	0.3	0.7	0.9
2005 4/.....	2.6	1.5	*	*	*	*	*	0.3	0.3	0.8
2004 4/.....	2.6	1.7	*	*	*	*	*	*	0.5	0.9
2003 4/.....	2.7	*	*	*	*	*	*	0.3	0.5	1.1
2002.....	2.4	*	*	*	*	*	*	*	0.4	0.9
2001.....	2.3	*	*	*	*	*	*	*	0.5	0.8
2000.....	2.2	*	*	*	*	*	*	*	0.5	0.8
1999.....	2.2	1.4	*	*	*	*	*	*	0.3	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Disorders of biliary tract and pancreas (K83-K86)										
All races 3/, both sexes										
2006.....	1.7	*	*	*	*	*	*	0.3	0.3	0.6
2005.....	1.6	*	*	*	*	*	0.1	0.2	0.3	0.6
2004.....	1.7	*	*	*	*	*	0.1	0.3	0.4	0.7
2003.....	1.7	*	*	*	*	*	*	0.2	0.4	0.6
2002.....	1.7	*	*	*	*	*	0.2	0.2	0.4	0.6
2001.....	1.7	*	*	*	*	*	0.1	0.2	0.5	0.7
2000.....	1.7	*	*	*	*	*	0.2	0.3	0.4	0.7
1999.....	1.6	*	*	*	*	*	0.2	0.3	0.4	0.6
All races 3/, male										
2006.....	1.8	*	*	*	*	*	*	0.4	0.5	0.8
2005.....	1.8	*	*	*	*	*	*	0.3	0.4	0.8
2004.....	1.8	*	*	*	*	*	*	0.4	0.6	1.0
2003.....	1.8	*	*	*	*	*	*	0.3	0.6	0.8
2002.....	1.9	*	*	*	*	*	0.2	0.3	0.6	0.8
2001.....	1.8	*	*	*	*	*	*	0.3	0.6	1.0
2000.....	1.8	*	*	*	*	*	0.2	0.4	0.6	0.9
1999.....	1.7	*	*	*	*	*	*	0.4	0.6	0.8
All races 3/, female										
2006.....	1.6	*	*	*	*	*	*	*	*	0.4
2005.....	1.5	*	*	*	*	*	*	*	0.2	0.4
2004.....	1.6	*	*	*	*	*	*	*	*	0.3
2003.....	1.6	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.5	*	*	*	*	*	*	*	0.3	0.4
2001.....	1.7	*	*	*	*	*	*	*	0.3	0.5
2000.....	1.6	*	*	*	*	*	*	*	0.2	0.4
1999.....	1.5	*	*	*	*	*	*	*	0.3	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Disorders of biliary tract and pancreas (K83-K86)										
White, both sexes										
2006 4/.....	1.7	*	*	*	*	*	*	0.3	0.3	0.6
2005 4/.....	1.7	*	*	*	*	*	*	0.2	0.3	0.6
2004 4/.....	1.7	*	*	*	*	*	*	0.2	0.3	0.6
2003 4/.....	1.7	*	*	*	*	*	*	0.2	0.4	0.5
2002.....	1.8	*	*	*	*	*	0.1	0.2	0.4	0.6
2001.....	1.8	*	*	*	*	*	*	0.2	0.4	0.6
2000.....	1.7	*	*	*	*	*	0.2	0.2	0.4	0.6
1999.....	1.6	*	*	*	*	*	*	0.2	0.4	0.5
White, male										
2006 4/.....	1.8	*	*	*	*	*	*	0.4	0.4	0.8
2005 4/.....	1.8	*	*	*	*	*	*	0.3	0.4	0.7
2004 4/.....	1.8	*	*	*	*	*	*	0.3	0.5	0.9
2003 4/.....	1.8	*	*	*	*	*	*	*	0.6	0.7
2002.....	1.9	*	*	*	*	*	*	0.3	0.5	0.7
2001.....	1.8	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	1.8	*	*	*	*	*	*	0.3	0.5	0.8
1999.....	1.7	*	*	*	*	*	*	0.3	0.5	0.6
White, female										
2006 4/.....	1.7	*	*	*	*	*	*	*	*	0.4
2005 4/.....	1.6	*	*	*	*	*	*	*	*	0.4
2004 4/.....	1.6	*	*	*	*	*	*	*	*	0.3
2003 4/.....	1.7	*	*	*	*	*	*	*	*	0.4
2002.....	1.6	*	*	*	*	*	*	*	*	0.4
2001.....	1.8	*	*	*	*	*	*	*	*	0.4
2000.....	1.6	*	*	*	*	*	*	*	*	0.4
1999.....	1.6	*	*	*	*	*	*	*	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pancreatitis (K85,K86.0-K86.1)										
All races 3/, both sexes										
2006.....	1.2	*	*	*	*	*	*	0.3	0.3	0.5
2005.....	1.1	*	*	*	*	*	0.1	0.2	0.3	0.5
2004.....	1.2	*	*	*	*	*	0.1	0.2	0.3	0.6
2003.....	1.2	*	*	*	*	*	*	0.2	0.3	0.6
2002.....	1.2	*	*	*	*	*	0.2	0.2	0.4	0.6
2001.....	1.3	*	*	*	*	*	0.1	0.2	0.4	0.6
2000.....	1.2	*	*	*	*	*	0.2	0.2	0.4	0.6
1999.....	1.2	*	*	*	*	*	0.1	0.2	0.4	0.5
All races 3/, male										
2006.....	1.3	*	*	*	*	*	*	0.4	0.5	0.7
2005.....	1.3	*	*	*	*	*	*	0.3	0.4	0.7
2004.....	1.3	*	*	*	*	*	*	0.4	0.6	0.9
2003.....	1.3	*	*	*	*	*	*	0.2	0.5	0.8
2002.....	1.4	*	*	*	*	*	0.2	0.3	0.6	0.8
2001.....	1.4	*	*	*	*	*	*	0.3	0.5	0.9
2000.....	1.3	*	*	*	*	*	0.2	0.3	0.6	0.9
1999.....	1.3	*	*	*	*	*	*	0.3	0.6	0.7
All races 3/, female										
2006.....	1.0	*	*	*	*	*	*	*	*	0.3
2005.....	1.0	*	*	*	*	*	*	*	0.2	0.3
2004.....	1.0	*	*	*	*	*	*	*	*	0.2
2003.....	1.1	*	*	*	*	*	*	*	*	0.4
2002.....	1.1	*	*	*	*	*	*	*	*	0.4
2001.....	1.2	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.1	*	*	*	*	*	*	*	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pancreatitis (K85,K86.0-K86.1)										
White, both sexes										
2006 4/.....	1.2	*	*	*	*	*	*	0.3	0.3	0.6
2005 4/.....	1.2	*	*	*	*	*	*	0.2	0.3	0.5
2004 4/.....	1.2	*	*	*	*	*	*	0.2	0.3	0.5
2003 4/.....	1.2	*	*	*	*	*	*	0.1	0.3	0.5
2002.....	1.3	*	*	*	*	*	0.1	0.2	0.3	0.5
2001.....	1.3	*	*	*	*	*	*	0.2	0.3	0.6
2000.....	1.2	*	*	*	*	*	0.2	0.2	0.3	0.6
1999.....	1.2	*	*	*	*	*	*	0.2	0.4	0.4
White, male										
2006 4/.....	1.4	*	*	*	*	*	*	0.3	0.4	0.7
2005 4/.....	1.3	*	*	*	*	*	*	0.3	0.3	0.6
2004 4/.....	1.3	*	*	*	*	*	*	0.3	0.5	0.8
2003 4/.....	1.3	*	*	*	*	*	*	*	0.5	0.6
2002.....	1.4	*	*	*	*	*	*	0.3	0.5	0.7
2001.....	1.4	*	*	*	*	*	*	*	0.4	0.8
2000.....	1.3	*	*	*	*	*	*	0.3	0.5	0.8
1999.....	1.3	*	*	*	*	*	*	0.3	0.5	0.6
White, female										
2006 4/.....	1.1	*	*	*	*	*	*	*	*	0.4
2005 4/.....	1.0	*	*	*	*	*	*	*	*	0.3
2004 4/.....	1.1	*	*	*	*	*	*	*	*	*
2003 4/.....	1.1	*	*	*	*	*	*	*	*	0.3
2002.....	1.1	*	*	*	*	*	*	*	*	0.4
2001.....	1.2	*	*	*	*	*	*	*	*	0.4
2000.....	1.1	*	*	*	*	*	*	*	*	0.3
1999.....	1.1	*	*	*	*	*	*	*	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

All races 3/, both sexes

2006.....	4.7	*	*	*	0.1	0.3	0.5	0.7	0.8	1.1
2005.....	4.9	*	*	*	0.1	0.3	0.5	0.6	0.8	1.1
2004.....	4.9	*	*	*	0.2	0.3	0.5	0.7	0.7	1.1
2003.....	5.0	*	*	*	0.1	0.3	0.5	0.7	0.9	1.1
2002.....	5.0	*	*	*	0.1	0.2	0.5	0.7	0.7	1.2
2001.....	4.9	*	*	*	0.1	0.3	0.6	0.6	0.8	1.3
2000.....	4.9	*	*	*	0.1	0.2	0.6	0.7	0.8	1.1
1999.....	4.8	*	*	*	0.1	0.3	0.5	0.6	0.8	1.2

All races 3/, male

2006.....	2.9	*	*	*	*	*	0.3	0.4	0.4	0.6
2005.....	3.0	*	*	*	*	0.2	0.3	0.3	0.5	0.6
2004.....	2.9	*	*	*	*	0.2	0.3	0.3	0.4	0.7
2003.....	3.0	*	*	*	*	0.2	0.2	0.4	0.4	0.8
2002.....	3.0	*	*	*	*	*	0.2	0.4	0.5	0.7
2001.....	2.9	*	*	*	*	*	0.2	0.3	0.5	1.0
2000.....	2.9	*	*	*	*	*	0.4	0.4	0.5	0.6
1999.....	2.8	*	*	*	*	*	0.2	0.3	0.5	0.8

All races 3/, female

2006.....	6.4	*	*	*	*	0.4	0.7	1.1	1.1	1.5
2005.....	6.7	*	*	*	0.2	0.3	0.8	1.0	1.2	1.5
2004.....	6.7	*	*	*	0.2	0.3	0.7	1.1	1.0	1.6
2003.....	6.9	*	*	*	*	0.5	0.8	1.1	1.4	1.4
2002.....	7.0	*	*	*	*	0.4	0.8	1.0	1.0	1.6
2001.....	6.9	*	*	*	*	0.5	0.9	1.0	1.2	1.7
2000.....	6.8	*	*	*	*	0.3	0.8	1.1	1.2	1.6
1999.....	6.7	*	*	*	0.2	0.4	0.7	1.0	1.2	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

White, both sexes

2006 4/.....	4.9	*	*	*	*	0.2	0.3	0.4	0.5	0.7
2005 4/.....	5.1	*	*	*	0.1	0.2	0.3	0.4	0.5	0.7
2004 4/.....	5.0	*	*	*	0.1	0.2	0.3	0.3	0.4	0.8
2003 4/.....	5.2	*	*	*	*	0.3	0.3	0.4	0.6	0.7
2002.....	5.2	*	*	*	*	0.2	0.3	0.4	0.4	0.8
2001.....	5.1	*	*	*	*	0.3	0.4	0.3	0.6	1.0
2000.....	5.1	*	*	*	*	*	0.3	0.5	0.6	0.7
1999.....	5.0	*	*	*	*	0.2	0.3	0.4	0.6	0.7

White, male

2006 4/.....	3.0	*	*	*	*	*	0.3	*	0.3	0.5
2005 4/.....	3.1	*	*	*	*	*	0.2	*	0.3	0.5
2004 4/.....	3.0	*	*	*	*	*	*	*	*	0.5
2003 4/.....	3.1	*	*	*	*	*	0.2	0.3	0.3	0.6
2002.....	3.0	*	*	*	*	*	*	0.3	0.3	0.6
2001.....	3.0	*	*	*	*	*	*	*	0.3	0.9
2000.....	3.0	*	*	*	*	*	*	0.3	0.4	0.4
1999.....	2.9	*	*	*	*	*	*	*	0.4	0.5

White, female

2006 4/.....	6.7	*	*	*	*	0.3	0.4	0.6	0.7	1.0
2005 4/.....	7.0	*	*	*	*	*	0.4	0.6	0.7	0.9
2004 4/.....	7.0	*	*	*	*	*	0.4	0.6	0.6	1.0
2003 4/.....	7.3	*	*	*	*	0.3	0.5	0.6	0.9	0.9
2002.....	7.4	*	*	*	*	0.3	0.4	0.5	0.6	0.9
2001.....	7.2	*	*	*	*	0.3	0.5	0.5	0.8	1.2
2000.....	7.1	*	*	*	*	*	0.4	0.7	0.8	1.1
1999.....	7.0	*	*	*	*	*	0.4	0.6	0.8	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

Black, both sexes

2006 4/.....	4.6	*	*	*	*	0.7	1.6	2.6	2.2	3.2
2005 4/.....	4.7	*	*	*	*	0.6	1.5	1.9	2.7	3.4
2004 4/.....	5.0	*	*	*	*	*	1.7	2.6	2.5	3.5
2003 4/.....	4.9	*	*	*	*	0.9	1.5	2.3	2.7	3.4
2002.....	4.8	*	*	*	*	*	1.9	2.3	2.6	3.9
2001.....	4.8	*	*	*	*	0.7	1.6	2.3	2.5	3.5
2000.....	4.9	*	*	*	*	0.7	1.9	2.3	2.5	3.7
1999.....	4.7	*	*	*	*	0.7	1.7	2.0	2.2	4.1

Black, male

2006 4/.....	3.0	*	*	*	*	*	*	*	*	*
2005 4/.....	2.9	*	*	*	*	*	*	*	*	1.8
2004 4/.....	3.3	*	*	*	*	*	*	*	1.7	1.7
2003 4/.....	3.3	*	*	*	*	*	*	*	*	2.6
2002.....	3.1	*	*	*	*	*	*	*	1.8	2.0
2001.....	3.1	*	*	*	*	*	*	*	*	2.1
2000.....	3.4	*	*	*	*	*	*	*	*	2.4
1999.....	2.9	*	*	*	*	*	*	*	*	2.4

Black, female

2006 4/.....	6.0	*	*	*	*	*	2.5	3.8	3.2	4.8
2005 4/.....	6.4	*	*	*	*	*	2.4	3.0	3.8	4.9
2004 4/.....	6.5	*	*	*	*	*	2.3	3.7	3.2	5.1
2003 4/.....	6.4	*	*	*	*	1.4	2.6	3.5	4.2	4.1
2002.....	6.3	*	*	*	*	*	3.0	3.4	3.4	5.6
2001.....	6.3	*	*	*	*	*	2.7	3.6	3.6	4.8
2000.....	6.2	*	*	*	*	*	2.5	3.2	3.4	4.9
1999.....	6.2	*	*	*	*	1.4	2.4	3.1	3.4	5.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Systemic lupus erythematosus (M32)										
All races 3/, both sexes										
2006.....	0.4	*	*	*	*	0.2	0.3	0.4	0.4	0.4
2005.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.4
2004.....	0.4	*	*	*	*	*	0.3	0.4	0.4	0.5
2003.....	0.5	*	*	*	*	0.2	0.3	0.3	0.4	0.4
2002.....	0.4	*	*	*	*	0.1	0.3	0.3	0.3	0.5
2001.....	0.5	*	*	*	*	0.2	0.4	0.4	0.4	0.5
2000.....	0.5	*	*	*	*	0.1	0.3	0.4	0.3	0.5
1999.....	0.5	*	*	*	*	0.1	0.3	0.3	0.4	0.5
All races 3/, male										
2006.....	0.1	*	*	*	*	*	*	*	*	*
2005.....	0.1	*	*	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	0.2
2003.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
All races 3/, female										
2006.....	0.7	*	*	*	*	0.3	0.5	0.8	0.7	0.7
2005.....	0.8	*	*	*	*	*	0.5	0.6	0.7	0.7
2004.....	0.7	*	*	*	*	*	0.5	0.7	0.6	0.7
2003.....	0.8	*	*	*	*	0.3	0.5	0.6	0.7	0.7
2002.....	0.8	*	*	*	*	0.2	0.5	0.5	0.5	0.9
2001.....	0.8	*	*	*	*	0.3	0.6	0.7	0.6	0.8
2000.....	0.8	*	*	*	*	*	0.5	0.6	0.6	0.9
1999.....	0.8	*	*	*	*	0.2	0.5	0.6	0.6	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Systemic lupus erythematosus (M32)										
White, both sexes										
2006 4/.....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.2
2005 4/.....	0.4	*	*	*	*	*	0.2	0.2	0.2	0.2
2004 4/.....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.3
2003 4/.....	0.4	*	*	*	*	*	0.2	0.2	0.2	0.3
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.2
2001.....	0.4	*	*	*	*	*	0.2	0.1	0.2	0.2
2000.....	0.4	*	*	*	*	*	0.2	0.2	0.2	0.3
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.2	0.3
White, male										
2006 4/.....	0.1	*	*	*	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
White, female										
2006 4/.....	0.5	*	*	*	*	*	*	0.3	0.4	0.4
2005 4/.....	0.6	*	*	*	*	*	*	0.3	0.3	0.4
2004 4/.....	0.5	*	*	*	*	*	*	0.3	0.3	0.4
2003 4/.....	0.6	*	*	*	*	*	0.3	0.3	0.4	0.4
2002.....	0.6	*	*	*	*	*	*	*	0.3	0.4
2001.....	0.6	*	*	*	*	*	0.3	*	0.3	0.4
2000.....	0.6	*	*	*	*	*	0.3	0.3	0.3	0.5
1999.....	0.6	*	*	*	*	*	*	0.4	0.3	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Systemic lupus erythematosus (M32)										
Black, both sexes										
2006 4/.....	1.1	*	*	*	*	*	1.1	1.8	1.5	1.5
2005 4/.....	1.2	*	*	*	*	*	1.1	1.2	1.3	1.5
2004 4/.....	1.2	*	*	*	*	*	1.1	1.5	1.4	1.6
2003 4/.....	1.1	*	*	*	*	0.6	1.0	1.1	1.3	1.4
2002.....	1.2	*	*	*	*	*	1.2	1.3	1.3	2.1
2001.....	1.2	*	*	*	*	*	1.1	1.6	1.4	1.9
2000.....	1.2	*	*	*	*	*	1.1	1.4	1.2	1.9
1999.....	1.2	*	*	*	*	*	1.2	1.1	1.3	2.0
Black, male										
2006 4/.....	0.4	*	*	*	*	*	*	*	*	*
2005 4/.....	0.3	*	*	*	*	*	*	*	*	*
2004 4/.....	0.4	*	*	*	*	*	*	*	*	*
2003 4/.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	1.8	*	*	*	*	*	1.8	3.0	2.4	2.4
2005 4/.....	2.0	*	*	*	*	*	1.8	2.0	2.3	2.5
2004 4/.....	2.0	*	*	*	*	*	1.7	2.4	2.4	2.4
2003 4/.....	1.9	*	*	*	*	*	1.9	1.9	2.3	2.4
2002.....	2.1	*	*	*	*	*	2.1	2.3	2.0	3.8
2001.....	2.0	*	*	*	*	*	2.0	2.7	2.2	3.2
2000.....	2.0	*	*	*	*	*	1.8	2.5	2.0	2.8
1999.....	2.0	*	*	*	*	*	1.9	1.9	2.1	3.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)										
All races 3/, both sexes										
2006.....	1.5	*	*	*	*	*	0.1	0.2	0.2	0.4
2005.....	1.6	*	*	*	*	*	0.1	0.2	0.3	0.4
2004.....	1.6	*	*	*	*	*	0.1	0.2	0.2	0.4
2003.....	1.6	*	*	*	*	0.1	0.1	0.2	0.3	0.4
2002.....	1.7	*	*	*	*	*	0.1	0.2	0.3	0.5
2001.....	1.7	*	*	*	*	*	0.1	0.2	0.3	0.5
2000.....	1.7	*	*	*	*	*	0.2	0.2	0.4	0.4
1999.....	1.7	*	*	*	*	0.1	*	0.2	0.3	0.5
All races 3/, male										
2006.....	0.9	*	*	*	*	*	*	*	0.2	0.3
2005.....	1.0	*	*	*	*	*	*	*	0.2	0.3
2004.....	1.0	*	*	*	*	*	*	*	*	0.2
2003.....	1.0	*	*	*	*	*	*	*	*	0.4
2002.....	1.0	*	*	*	*	*	*	*	0.2	0.4
2001.....	1.0	*	*	*	*	*	*	*	*	0.4
2000.....	1.1	*	*	*	*	*	*	0.2	0.3	0.2
1999.....	1.0	*	*	*	*	*	*	*	*	0.4
All races 3/, female										
2006.....	2.1	*	*	*	*	*	*	0.2	0.3	0.6
2005.....	2.2	*	*	*	*	*	*	0.3	0.4	0.6
2004.....	2.2	*	*	*	*	*	*	0.3	0.2	0.6
2003.....	2.2	*	*	*	*	*	*	0.4	0.4	0.5
2002.....	2.3	*	*	*	*	*	0.2	0.3	0.4	0.5
2001.....	2.3	*	*	*	*	*	*	0.3	0.4	0.6
2000.....	2.3	*	*	*	*	*	*	0.3	0.5	0.6
1999.....	2.3	*	*	*	*	*	*	0.2	0.4	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)										
White, both sexes										
2006 4/.....	1.6	*	*	*	*	*	*	*	0.2	0.2
2005 4/.....	1.7	*	*	*	*	*	*	*	0.2	0.3
2004 4/.....	1.7	*	*	*	*	*	*	*	0.1	0.3
2003 4/.....	1.7	*	*	*	*	*	*	*	0.1	0.3
2002.....	1.7	*	*	*	*	*	*	0.2	0.2	0.3
2001.....	1.8	*	*	*	*	*	*	0.1	0.2	0.4
2000.....	1.7	*	*	*	*	*	*	0.2	0.3	0.2
1999.....	1.7	*	*	*	*	*	*	*	0.2	0.3
White, male										
2006 4/.....	1.0	*	*	*	*	*	*	*	*	*
2005 4/.....	1.0	*	*	*	*	*	*	*	*	*
2004 4/.....	1.0	*	*	*	*	*	*	*	*	*
2003 4/.....	1.0	*	*	*	*	*	*	*	*	*
2002.....	1.0	*	*	*	*	*	*	*	*	0.3
2001.....	1.1	*	*	*	*	*	*	*	*	0.3
2000.....	1.0	*	*	*	*	*	*	*	*	*
1999.....	1.0	*	*	*	*	*	*	*	*	0.3
White, female										
2006 4/.....	2.2	*	*	*	*	*	*	*	*	0.3
2005 4/.....	2.3	*	*	*	*	*	*	*	*	0.4
2004 4/.....	2.3	*	*	*	*	*	*	*	*	0.4
2003 4/.....	2.3	*	*	*	*	*	*	*	*	0.3
2002.....	2.4	*	*	*	*	*	*	*	0.2	0.3
2001.....	2.4	*	*	*	*	*	*	*	0.3	0.5
2000.....	2.4	*	*	*	*	*	*	*	0.3	0.4
1999.....	2.3	*	*	*	*	*	*	*	0.3	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XIV. Diseases of the genitourinary system (N00-N98)										
All races 3/, both sexes										
2006.....	21.2	4.4	*	*	0.1	0.1	0.4	0.6	1.1	1.6
2005.....	21.2	4.4	0.1	*	0.1	0.2	0.4	0.7	1.1	1.6
2004.....	20.5	4.9	0.2	*	*	0.2	0.4	0.6	1.0	1.7
2003.....	20.7	5.1	0.1	*	0.1	0.2	0.4	0.7	1.1	1.6
2002.....	20.2	4.6	*	*	*	0.2	0.4	0.7	1.0	1.6
2001.....	20.0	3.9	0.2	*	*	0.1	0.4	0.6	0.9	1.7
2000.....	19.4	5.0	0.1	*	*	0.2	0.4	0.5	1.0	1.5
1999.....	19.0	5.1	0.2	0.1	0.1	0.2	0.3	0.7	0.9	1.5
All races 3/, male										
2006.....	19.4	5.5	*	*	*	*	0.4	0.5	1.2	1.7
2005.....	19.1	4.8	*	*	*	0.2	0.4	0.7	1.1	1.6
2004.....	18.5	5.8	*	*	*	*	0.4	0.6	0.8	1.8
2003.....	18.6	6.0	*	*	*	0.2	0.4	0.6	1.1	1.6
2002.....	18.1	5.6	*	*	*	*	0.4	0.6	1.0	1.6
2001.....	17.7	4.2	*	*	*	*	0.4	0.6	0.9	1.8
2000.....	17.2	6.1	*	*	*	0.2	0.3	0.5	1.0	1.5
1999.....	16.9	5.0	*	*	*	*	0.3	0.6	0.9	1.6
All races 3/, female										
2006.....	23.0	3.2	*	*	*	*	0.4	0.6	1.0	1.4
2005.....	23.2	3.9	*	*	*	0.2	0.5	0.8	1.0	1.5
2004.....	22.5	4.0	*	*	*	*	0.3	0.7	1.1	1.7
2003.....	22.6	4.3	*	*	*	0.2	0.5	0.7	1.1	1.5
2002.....	22.3	3.6	*	*	*	*	0.4	0.9	1.0	1.6
2001.....	22.2	3.6	*	*	*	*	0.5	0.6	1.0	1.6
2000.....	21.5	3.9	*	*	*	*	0.5	0.5	1.0	1.5
1999.....	21.1	5.1	*	*	*	0.2	0.4	0.8	0.9	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XIV. Diseases of the genitourinary system (N00-N98)										
White, both sexes										
2006 4/.....	21.4	3.9	*	*	*	0.1	0.4	0.4	0.9	1.1
2005 4/.....	21.3	3.5	0.2	*	*	0.2	0.4	0.5	0.7	1.2
2004 4/.....	20.6	3.6	*	*	*	0.2	0.3	0.5	0.7	1.3
2003 4/.....	20.7	4.1	*	*	*	0.2	0.3	0.5	0.8	1.1
2002.....	20.3	4.0	*	*	*	*	0.3	0.5	0.8	1.1
2001.....	20.0	3.0	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	19.4	3.6	*	*	*	0.2	0.4	0.4	0.7	1.0
1999.....	18.9	4.0	*	*	*	0.1	0.2	0.5	0.7	1.0
White, male										
2006 4/.....	19.5	5.1	*	*	*	*	0.4	0.3	1.0	1.2
2005 4/.....	19.3	3.6	*	*	*	*	0.3	0.4	0.8	1.2
2004 4/.....	18.5	4.2	*	*	*	*	0.3	0.4	0.4	1.2
2003 4/.....	18.6	4.8	*	*	*	*	0.3	0.5	0.8	1.0
2002.....	18.1	4.7	*	*	*	*	0.3	0.4	0.8	1.1
2001.....	17.7	3.4	*	*	*	*	0.3	0.4	0.6	1.5
2000.....	17.2	4.6	*	*	*	*	0.3	0.4	0.7	1.0
1999.....	16.6	4.5	*	*	*	*	*	0.4	0.7	1.1
White, female										
2006 4/.....	23.2	2.7	*	*	*	*	0.3	0.4	0.7	1.0
2005 4/.....	23.4	3.3	*	*	*	*	0.4	0.6	0.7	1.1
2004 4/.....	22.7	2.8	*	*	*	*	0.3	0.5	0.9	1.3
2003 4/.....	22.8	3.4	*	*	*	*	0.3	0.6	0.8	1.1
2002.....	22.5	3.2	*	*	*	*	0.3	0.6	0.7	1.1
2001.....	22.3	2.6	*	*	*	*	0.5	0.5	0.8	1.0
2000.....	21.5	2.6	*	*	*	*	0.4	0.5	0.7	1.0
1999.....	21.2	3.4	*	*	*	*	0.3	0.6	0.6	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XIV. Diseases of the genitourinary system (N00-N98)										
Black, both sexes										
2006 4/.....	26.2	6.8	*	*	*	*	0.8	1.5	2.8	4.8
2005 4/.....	26.0	9.7	*	*	*	*	0.8	2.0	3.2	4.2
2004 4/.....	25.7	11.6	*	*	*	*	0.9	1.6	2.8	4.9
2003 4/.....	26.0	11.1	*	*	*	*	1.0	1.6	3.4	5.1
2002.....	25.2	8.3	*	*	*	*	1.1	2.3	2.6	4.8
2001.....	25.1	9.2	*	*	*	*	0.8	1.6	2.6	4.7
2000.....	24.4	12.7	*	*	*	*	*	1.2	2.9	4.9
1999.....	24.3	11.1	*	*	*	*	1.0	1.8	2.5	4.6
Black, male										
2006 4/.....	24.1	7.5	*	*	*	*	*	1.4	3.1	5.3
2005 4/.....	23.7	12.1	*	*	*	*	*	1.9	3.5	4.0
2004 4/.....	23.4	13.0	*	*	*	*	*	*	3.1	5.8
2003 4/.....	24.0	12.5	*	*	*	*	*	*	3.9	6.0
2002.....	23.2	10.5	*	*	*	*	*	1.9	2.4	4.6
2001.....	22.2	9.3	*	*	*	*	*	2.1	3.1	3.8
2000.....	21.9	14.1	*	*	*	*	*	*	3.1	5.6
1999.....	22.4	8.1	*	*	*	*	*	*	2.4	5.3
Black, female										
2006 4/.....	28.1	6.0	*	*	*	*	*	1.6	2.4	4.3
2005 4/.....	28.2	7.1	*	*	*	*	*	2.0	3.0	4.4
2004 4/.....	27.7	10.2	*	*	*	*	*	1.8	2.6	4.0
2003 4/.....	27.7	9.6	*	*	*	*	*	1.8	2.9	4.3
2002.....	27.0	6.1	*	*	*	*	1.3	2.7	2.8	5.0
2001.....	27.8	9.1	*	*	*	*	*	*	2.1	5.5
2000.....	26.6	11.2	*	*	*	*	*	*	2.8	4.3
1999.....	26.1	14.1	*	*	*	*	*	2.2	2.5	4.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of urinary system (N00-N39)										
All races 3/, both sexes										
2006.....	20.8	4.3	*	*	0.1	0.1	0.4	0.5	1.1	1.5
2005.....	20.8	4.4	0.1	*	0.1	0.2	0.4	0.7	1.0	1.5
2004.....	20.2	4.8	0.2	*	*	0.2	0.4	0.6	0.9	1.7
2003.....	20.3	5.0	0.1	*	0.1	0.2	0.4	0.7	1.1	1.5
2002.....	19.9	4.6	*	*	*	0.1	0.4	0.7	0.9	1.5
2001.....	19.7	3.7	0.2	*	*	0.1	0.4	0.6	0.9	1.7
2000.....	19.1	4.8	0.1	*	*	0.2	0.4	0.5	0.9	1.4
1999.....	18.7	4.7	0.1	0.1	0.1	0.2	0.3	0.7	0.9	1.4
All races 3/, male										
2006.....	18.9	5.5	*	*	*	*	0.4	0.5	1.2	1.7
2005.....	18.6	4.8	*	*	*	0.2	0.4	0.6	1.1	1.6
2004.....	18.0	5.8	*	*	*	*	0.4	0.6	0.8	1.8
2003.....	18.2	5.9	*	*	*	0.2	0.4	0.6	1.1	1.6
2002.....	17.7	5.6	*	*	*	*	0.4	0.6	1.0	1.5
2001.....	17.3	4.2	*	*	*	*	0.4	0.6	0.9	1.7
2000.....	16.8	6.0	*	*	*	0.2	0.3	0.5	1.0	1.5
1999.....	16.4	5.0	*	*	*	*	0.3	0.6	0.9	1.6
All races 3/, female										
2006.....	22.8	3.1	*	*	*	*	0.4	0.6	0.9	1.4
2005.....	22.9	3.9	*	*	*	0.2	0.4	0.8	0.9	1.4
2004.....	22.2	3.8	*	*	*	*	0.3	0.7	1.1	1.6
2003.....	22.4	4.1	*	*	*	*	0.4	0.7	1.0	1.4
2002.....	22.0	3.5	*	*	*	*	0.4	0.9	0.9	1.4
2001.....	22.0	3.2	*	*	*	*	0.5	0.6	1.0	1.6
2000.....	21.2	3.6	*	*	*	*	0.4	0.5	0.9	1.3
1999.....	20.9	4.3	*	*	*	*	0.3	0.7	0.8	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of urinary system (N00-N39)										
White, both sexes										
2006 4/.....	21.0	3.9	*	*	*	*	0.4	0.4	0.9	1.1
2005 4/.....	20.9	3.4	0.2	*	*	0.2	0.4	0.5	0.7	1.1
2004 4/.....	20.2	3.5	*	*	*	0.1	0.3	0.4	0.6	1.2
2003 4/.....	20.4	4.1	*	*	*	0.2	0.3	0.5	0.8	1.0
2002.....	20.0	3.9	*	*	*	*	0.3	0.5	0.7	1.0
2001.....	19.7	2.8	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	19.0	3.5	*	*	*	0.2	0.4	0.4	0.7	1.0
1999.....	18.6	3.7	*	*	*	0.1	0.2	0.5	0.6	1.0
White, male										
2006 4/.....	18.9	5.1	*	*	*	*	0.4	0.3	1.0	1.2
2005 4/.....	18.7	3.6	*	*	*	*	0.3	0.4	0.8	1.2
2004 4/.....	18.0	4.2	*	*	*	*	0.3	0.4	0.4	1.2
2003 4/.....	18.1	4.8	*	*	*	*	0.3	0.5	0.8	1.0
2002.....	17.7	4.7	*	*	*	*	0.3	0.4	0.8	1.1
2001.....	17.3	3.3	*	*	*	*	0.3	0.4	0.6	1.5
2000.....	16.7	4.5	*	*	*	*	0.3	0.4	0.7	1.0
1999.....	16.2	4.5	*	*	*	*	*	0.4	0.7	1.1
White, female										
2006 4/.....	22.9	2.6	*	*	*	*	0.3	0.4	0.7	1.0
2005 4/.....	23.1	3.3	*	*	*	*	0.4	0.6	0.6	1.0
2004 4/.....	22.4	2.8	*	*	*	*	0.3	0.5	0.8	1.3
2003 4/.....	22.5	3.3	*	*	*	*	0.3	0.5	0.8	1.0
2002.....	22.2	3.0	*	*	*	*	*	0.6	0.6	1.0
2001.....	22.0	2.3	*	*	*	*	0.4	0.5	0.8	1.0
2000.....	21.3	2.4	*	*	*	*	0.4	0.4	0.7	0.9
1999.....	20.9	2.8	*	*	*	*	*	0.5	0.6	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Diseases of urinary system (N00-N39)										
Black, both sexes										
2006 4/.....	25.9	6.6	*	*	*	*	0.8	1.4	2.7	4.7
2005 4/.....	25.7	9.7	*	*	*	*	0.7	1.8	3.1	4.1
2004 4/.....	25.3	11.2	*	*	*	*	0.8	1.6	2.8	4.8
2003 4/.....	25.7	10.6	*	*	*	*	0.9	1.5	3.3	5.0
2002.....	24.8	8.3	*	*	*	*	1.0	2.3	2.5	4.5
2001.....	24.8	8.9	*	*	*	*	0.8	1.6	2.6	4.4
2000.....	24.1	12.2	*	*	*	*	*	1.2	2.8	4.7
1999.....	23.9	9.9	*	*	*	*	0.9	1.7	2.4	4.5
Black, male										
2006 4/.....	23.7	7.5	*	*	*	*	*	1.4	3.1	5.2
2005 4/.....	23.3	12.1	*	*	*	*	*	1.8	3.3	4.0
2004 4/.....	23.0	13.0	*	*	*	*	*	*	3.0	5.8
2003 4/.....	23.7	12.2	*	*	*	*	*	*	3.9	5.9
2002.....	22.8	10.5	*	*	*	*	*	1.9	2.4	4.5
2001.....	21.8	9.3	*	*	*	*	*	2.1	3.1	3.6
2000.....	21.6	14.1	*	*	*	*	*	*	3.0	5.5
1999.....	22.0	8.1	*	*	*	*	*	*	2.4	5.3
Black, female										
2006 4/.....	27.8	*	*	*	*	*	*	1.4	2.3	4.1
2005 4/.....	27.8	7.1	*	*	*	*	*	1.9	2.9	4.1
2004 4/.....	27.4	9.3	*	*	*	*	*	1.8	2.5	3.8
2003 4/.....	27.4	9.0	*	*	*	*	*	1.6	2.7	4.1
2002.....	26.6	6.1	*	*	*	*	*	2.6	2.6	4.4
2001.....	27.5	8.5	*	*	*	*	*	*	2.1	5.2
2000.....	26.3	10.3	*	*	*	*	*	*	2.6	4.0
1999.....	25.7	11.7	*	*	*	*	*	2.0	2.4	3.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
All races 3/, both sexes										
2006.....	14.5	3.7	*	*	*	0.1	0.3	0.4	0.9	1.2
2005.....	14.5	3.7	*	*	*	0.1	0.2	0.5	0.8	1.2
2004.....	14.2	4.0	*	*	*	0.1	0.3	0.4	0.7	1.3
2003.....	14.4	4.2	*	*	*	0.1	0.3	0.5	0.8	1.2
2002.....	13.9	4.1	*	*	*	*	0.3	0.5	0.7	1.2
2001.....	13.6	3.1	*	*	*	*	0.3	0.4	0.6	1.2
2000.....	13.0	4.0	*	*	*	0.1	0.3	0.4	0.7	1.2
1999.....	12.4	4.3	*	*	*	0.1	0.2	0.4	0.7	1.1
All races 3/, male										
2006.....	14.3	4.4	*	*	*	*	0.3	0.3	1.0	1.3
2005.....	14.2	4.1	*	*	*	*	0.3	0.5	0.9	1.2
2004.....	13.8	5.1	*	*	*	*	0.3	0.4	0.6	1.4
2003.....	14.1	4.8	*	*	*	*	0.3	0.5	0.9	1.3
2002.....	13.7	5.1	*	*	*	*	0.3	0.4	0.7	1.3
2001.....	13.2	3.5	*	*	*	*	0.3	0.4	0.6	1.3
2000.....	12.6	5.1	*	*	*	*	0.2	0.4	0.7	1.3
1999.....	12.1	4.5	*	*	*	*	*	0.3	0.7	1.3
All races 3/, female										
2006.....	14.6	3.0	*	*	*	*	0.3	0.4	0.7	1.1
2005.....	14.7	3.2	*	*	*	*	0.2	0.5	0.7	1.1
2004.....	14.6	3.0	*	*	*	*	0.2	0.4	0.8	1.2
2003.....	14.6	3.7	*	*	*	*	0.2	0.5	0.8	1.1
2002.....	14.2	3.0	*	*	*	*	0.3	0.6	0.7	1.1
2001.....	14.0	2.7	*	*	*	*	0.4	0.4	0.7	1.2
2000.....	13.3	2.8	*	*	*	*	0.3	0.3	0.7	1.0
1999.....	12.7	4.1	*	*	*	*	0.3	0.5	0.7	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
White, both sexes										
2006 4/.....	14.1	3.3	*	*	*	*	0.3	0.3	0.7	0.8
2005 4/.....	14.2	2.8	*	*	*	*	0.2	0.4	0.5	0.8
2004 4/.....	13.9	2.9	*	*	*	*	0.2	0.3	0.4	0.9
2003 4/.....	14.0	3.5	*	*	*	*	0.2	0.4	0.5	0.8
2002.....	13.6	3.4	*	*	*	*	0.2	0.3	0.5	0.8
2001.....	13.2	2.2	*	*	*	*	0.3	0.3	0.4	0.8
2000.....	12.6	2.7	*	*	*	*	0.2	0.3	0.5	0.7
1999.....	12.0	3.3	*	*	*	*	0.2	0.3	0.5	0.7
White, male										
2006 4/.....	14.1	4.0	*	*	*	*	0.3	*	0.8	0.9
2005 4/.....	14.1	3.1	*	*	*	*	0.2	0.3	0.6	0.9
2004 4/.....	13.6	3.6	*	*	*	*	*	0.3	0.3	0.9
2003 4/.....	13.8	4.1	*	*	*	*	*	0.4	0.5	0.8
2002.....	13.4	4.2	*	*	*	*	*	*	0.5	0.9
2001.....	13.0	2.6	*	*	*	*	*	0.3	0.4	1.0
2000.....	12.4	3.6	*	*	*	*	0.3	0.3	0.5	0.8
1999.....	11.8	3.9	*	*	*	*	*	*	0.5	0.8
White, female										
2006 4/.....	14.2	2.6	*	*	*	*	*	0.3	0.5	0.7
2005 4/.....	14.2	2.6	*	*	*	*	*	0.4	0.4	0.7
2004 4/.....	14.1	2.1	*	*	*	*	*	0.3	0.5	1.0
2003 4/.....	14.2	2.8	*	*	*	*	*	0.4	0.5	0.7
2002.....	13.8	2.6	*	*	*	*	*	0.4	0.5	0.7
2001.....	13.4	1.8	*	*	*	*	0.3	0.3	0.5	0.7
2000.....	12.8	1.7	*	*	*	*	*	*	0.5	0.7
1999.....	12.2	2.8	*	*	*	*	*	0.3	0.5	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Renal failure (N17-N19)										
Black, both sexes										
2006 4/.....	20.3	6.2	*	*	*	*	0.7	1.2	2.3	3.8
2005 4/.....	20.2	8.4	*	*	*	*	*	1.4	2.6	3.5
2004 4/.....	20.0	9.8	*	*	*	*	0.7	1.2	2.3	4.0
2003 4/.....	20.3	8.9	*	*	*	*	0.6	1.3	2.8	4.4
2002.....	19.5	7.7	*	*	*	*	0.8	1.7	2.0	3.9
2001.....	19.3	8.1	*	*	*	*	0.7	1.2	2.1	3.9
2000.....	18.5	11.1	*	*	*	*	*	1.0	2.5	4.1
1999.....	18.2	9.6	*	*	*	*	*	1.4	2.2	3.8
Black, male										
2006 4/.....	19.2	6.6	*	*	*	*	*	*	2.6	4.3
2005 4/.....	19.1	10.6	*	*	*	*	*	1.4	2.7	3.5
2004 4/.....	18.9	11.9	*	*	*	*	*	*	2.7	5.1
2003 4/.....	19.7	9.2	*	*	*	*	*	*	3.4	5.4
2002.....	18.8	9.6	*	*	*	*	*	*	2.0	4.4
2001.....	17.7	8.7	*	*	*	*	*	1.6	2.6	3.3
2000.....	17.3	13.4	*	*	*	*	*	*	2.7	4.9
1999.....	17.1	8.1	*	*	*	*	*	*	2.2	4.5
Black, female										
2006 4/.....	21.3	*	*	*	*	*	*	1.3	2.0	3.4
2005 4/.....	21.2	6.2	*	*	*	*	*	1.4	2.5	3.4
2004 4/.....	21.0	7.7	*	*	*	*	*	*	1.9	3.0
2003 4/.....	20.9	8.7	*	*	*	*	*	*	2.3	3.5
2002.....	20.2	*	*	*	*	*	*	1.9	2.0	3.5
2001.....	20.7	7.6	*	*	*	*	*	*	1.7	4.4
2000.....	19.6	8.6	*	*	*	*	*	*	2.3	3.4
1999.....	19.1	11.1	*	*	*	*	*	1.6	2.2	3.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XV. Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes

2006.....	0.3	*	0.2	0.7	0.8	0.8	0.7
2005.....	0.3	*	0.2	0.7	0.8	0.8	0.6
2004.....	0.2	*	0.2	0.6	0.7	0.7	0.5
2003.....	0.2	*	0.1	0.4	0.6	0.6	0.5
2002.....	0.1	*	0.1	0.3	0.4	0.4	0.4
2001.....	0.1	*	0.2	0.4	0.5	0.5	0.3
2000.....	0.1	*	*	0.4	0.4	0.5	0.4
1999.....	0.1	*	0.2	0.3	0.5	0.4	0.4

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.5	*	0.3	1.4	1.7	1.6	1.3
2005.....	0.5	*	0.4	1.4	1.6	1.6	1.3
2004.....	0.5	*	0.4	1.3	1.5	1.4	0.9
2003.....	0.4	*	0.3	0.9	1.2	1.2	1.0
2002.....	0.3	*	0.3	0.6	0.9	0.9	0.7
2001.....	0.3	*	0.4	0.8	1.0	0.9	0.7
2000.....	0.3	*	*	0.8	0.9	0.9	0.8
1999.....	0.3	*	0.3	0.7	0.9	0.9	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XV. Pregnancy, childbirth and the puerperium (000-099)

White, both sexes

2006 4/.....	0.2	*	0.1	0.6	0.6	0.6	0.5
2005 4/.....	0.2	*	0.1	0.5	0.5	0.6	0.5
2004 4/.....	0.2	*	0.1	0.4	0.5	0.6	0.4
2003 4/.....	0.1	*	*	0.3	0.5	0.4	0.3
2002.....	0.1	*	*	0.2	0.3	0.3	0.2
2001.....	0.1	*	0.1	0.3	0.3	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.3	0.3
1999.....	0.1	*	*	0.2	0.3	0.3	0.3

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	0.4	*	0.3	1.2	1.2	1.2	0.9
2005 4/.....	0.4	*	0.3	1.1	1.0	1.3	1.0
2004 4/.....	0.3	*	0.3	0.8	1.0	1.2	0.7
2003 4/.....	0.3	*	*	0.6	1.0	0.7	0.7
2002.....	0.2	*	*	0.4	0.6	0.7	0.5
2001.....	0.2	*	0.3	0.6	0.7	0.8	0.4
2000.....	0.2	*	*	0.6	0.7	0.7	0.6
1999.....	0.2	*	*	0.4	0.6	0.6	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XV. Pregnancy, childbirth and the puerperium (000-099)										
Black, both sexes										
2006 4/.....	0.7	*	*	1.5	2.3	1.9	1.8
2005 4/.....	0.7	*	0.6	1.5	2.3	1.9	1.5
2004 4/.....	0.7	*	*	2.0	2.2	1.7	1.1
2003 4/.....	0.5	*	*	1.0	1.2	1.8	1.4
2002.....	0.4	*	*	1.0	1.4	1.0	1.2
2001.....	0.4	*	*	1.1	1.4	1.1	1.0
2000.....	0.4	*	*	1.1	1.1	1.2	0.9
1999.....	0.4	*	*	1.1	1.4	1.2	0.9
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	1.3	*	*	3.1	4.5	3.6	3.5
2005 4/.....	1.3	*	1.2	3.1	4.5	3.6	2.8
2004 4/.....	1.2	*	*	4.0	4.3	3.2	2.0
2003 4/.....	1.0	*	*	2.1	2.3	3.5	2.7
2002.....	0.8	*	*	2.0	2.8	1.9	2.3
2001.....	0.8	*	*	2.1	2.6	2.1	1.9
2000.....	0.7	*	*	2.2	2.2	2.3	1.7
1999.....	0.9	*	*	2.1	2.7	2.3	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy with abortive outcome (000-007)										
All races 3/, both sexes										
2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy with abortive outcome (000-007)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy with abortive outcome (000-007)										
Black, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ectopic pregnancy (000)										
All races 3/, both sexes										
2006.....	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ectopic pregnancy (000)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ectopic pregnancy (000)										
Black, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Spontaneous abortion (003)

All races 3/, both sexes

2006.....	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Spontaneous abortion (003)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Spontaneous abortion (003)										
Black, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Medical abortion (O04)										
All races 3/, both sexes										
2006.....	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Medical abortion (O04)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Medical abortion (O04)										
Black, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other abortion (005)										
All races 3/, both sexes										
2006.....	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other abortion (005)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other abortion (005)										
Black, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

All races 3/, both sexes

2006.....	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

Black, both sexes

2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other direct obstetric deaths (O10-O92)

All races 3/, both sexes

2006.....	0.1	*	*	0.4	0.5	0.4	0.3
2005.....	0.1	*	*	0.3	0.5	0.4	0.4
2004.....	0.1	*	0.1	0.3	0.4	0.4	0.2
2003.....	0.1	*	*	0.3	0.3	0.3	0.3
2002.....	0.1	*	0.1	0.2	0.3	0.3	0.3
2001.....	0.1	*	0.1	0.3	0.4	0.4	0.2
2000.....	0.1	*	*	0.3	0.3	0.4	0.3
1999.....	0.1	*	0.1	0.2	0.3	0.3	0.3

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.3	*	*	0.8	1.0	0.9	0.7
2005.....	0.3	*	*	0.7	0.9	0.7	0.7
2004.....	0.2	*	0.2	0.6	0.8	0.8	0.5
2003.....	0.2	*	*	0.5	0.7	0.7	0.7
2002.....	0.2	*	0.2	0.4	0.7	0.7	0.6
2001.....	0.2	*	0.3	0.5	0.8	0.7	0.5
2000.....	0.2	*	*	0.6	0.6	0.8	0.6
1999.....	0.2	*	0.3	0.5	0.7	0.7	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other direct obstetric deaths (O10-O92)										
White, both sexes										
2006 4/.....	0.1	*	*	0.3	0.3	0.3	0.2
2005 4/.....	0.1	*	*	0.3	0.3	0.3	0.3
2004 4/.....	0.1	*	*	0.2	0.2	0.3	0.2
2003 4/.....	0.1	*	*	0.2	0.3	0.2	0.2
2002.....	0.1	*	*	*	0.2	0.3	0.2
2001.....	0.1	*	*	0.2	0.3	0.3	0.1
2000.....	0.1	*	*	0.2	0.2	0.3	0.2
1999.....	0.1	*	*	0.2	0.2	0.2	0.2
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	0.2	*	*	0.6	0.6	0.6	0.4
2005 4/.....	0.2	*	*	0.6	0.6	0.5	0.6
2004 4/.....	0.2	*	*	0.3	0.5	0.6	0.4
2003 4/.....	0.2	*	*	0.4	0.5	0.4	0.4
2002.....	0.1	*	*	*	0.4	0.5	0.4
2001.....	0.2	*	*	0.4	0.6	0.6	0.3
2000.....	0.2	*	*	0.5	0.5	0.6	0.5
1999.....	0.1	*	*	0.3	0.5	0.5	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other direct obstetric deaths (O10-O92)										
Black, both sexes										
2006 4/.....	0.4	*	*	1.0	1.5	1.2	1.1
2005 4/.....	0.4	*	*	0.7	1.5	1.1	0.9
2004 4/.....	0.4	*	*	1.0	1.5	1.0	*
2003 4/.....	0.4	*	*	0.6	0.9	1.3	1.1
2002.....	0.3	*	*	0.8	1.0	0.9	0.9
2001.....	0.3	*	*	0.7	1.1	0.8	*
2000.....	0.3	*	*	0.8	0.8	0.9	*
1999.....	0.3	*	*	*	1.1	0.9	0.7
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	0.8	*	*	2.0	3.0	2.3	2.1
2005 4/.....	0.7	*	*	1.4	2.9	2.0	1.7
2004 4/.....	0.7	*	*	2.0	2.9	2.0	*
2003 4/.....	0.7	*	*	1.3	1.7	2.5	2.1
2002.....	0.6	*	*	1.5	1.9	1.6	1.7
2001.....	0.6	*	*	1.5	2.0	1.6	*
2000.....	0.5	*	*	1.6	1.5	1.7	*
1999.....	0.6	*	*	*	2.0	1.7	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Eclampsia and pre-eclampsia (O11,O13-O16)										
All races 3/, both sexes										
2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	0.1	*
1999.....	0.0	*	*	*	0.1	*	*
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.2	*
1999.....	0.1	*	*	*	0.2	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Eclampsia and pre-eclampsia (O11,O13-O16)										
White, both sexes										
2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Eclampsia and pre-eclampsia (O11,O13-O16)										
Black, both sexes										
2006 4/.....	0.1	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	0.1	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hemorrhage of pregnancy and childbirth and placenta previa (020,044-046,067,072)

All races 3/, both sexes

2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Hemorrhage of pregnancy and childbirth and placenta previa (020,044-046,067,072)

White, both sexes

2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hemorrhage of pregnancy and childbirth and placenta previa (020,044-046,067,072)										
Black, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Complications predominately related to the puerperium (085-092)

All races 3/, both sexes

2006.....	0.0	*	*	0.1	0.1	*	0.1
2005.....	0.0	*	*	0.2	0.1	*	0.1
2004.....	0.0	*	*	*	0.1	0.1	0.1
2003.....	0.0	*	*	*	0.1	*	0.1
2002.....	0.0	*	*	*	*	0.1	*
2001.....	0.0	*	*	*	0.1	0.1	*
2000.....	0.0	*	*	0.1	*	0.1	*
1999.....	0.0	*	*	*	*	0.1	*

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.1	*	*	0.3	0.3	*	0.2
2005.....	0.1	*	*	0.4	0.3	*	0.2
2004.....	0.1	*	*	*	0.2	0.2	0.2
2003.....	0.1	*	*	*	0.2	*	0.2
2002.....	0.1	*	*	*	*	0.2	*
2001.....	0.1	*	*	*	0.3	0.2	*
2000.....	0.1	*	*	0.2	*	0.3	*
1999.....	0.1	*	*	*	*	0.2	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Complications predominately related to the puerperium (085-092)										
White, both sexes										
2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.0	*	*	0.1	*	*	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.1	*	*	0.3	*	*	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Complications predominately related to the puerperium (085-092)

Black, both sexes

2006 4/.....	0.1	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	0.2	*	*	*	*	*	*
2005 4/.....	0.2	*	*	*	*	*	*
2004 4/.....	0.2	*	*	*	*	*	*
2003 4/.....	0.2	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Obstetric embolism (O88)										
All races 3/, both sexes										
2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	0.1	*
1999.....	0.0	*	*	*	*	*	*
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	0.2	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Obstetric embolism (O88)										
White, both sexes										
2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Obstetric embolism (O88)										
Black, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications predominately related to the puerperium (O85-O87,089-092)

All races 3/, both sexes

2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	0.1	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	0.2	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications predominately related to the puerperium (O85-O87,089-092)

White, both sexes

2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications predominately related to the puerperium (O85-O87,089-092)

Black, both sexes

2006 4/.....	0.1	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	0.2	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other direct obstetric causes (010,012,021-043,047-066,068-071,073-075)

All races 3/, both sexes

2006.....	0.1	*	*	0.2	0.2	0.2	0.1
2005.....	0.1	*	*	0.1	0.2	0.2	0.2
2004.....	0.1	*	*	0.1	0.2	0.2	0.1
2003.....	0.0	*	*	0.1	*	0.2	0.1
2002.....	0.0	*	*	0.1	0.1	0.1	0.1
2001.....	0.0	*	*	*	0.1	0.1	*
2000.....	0.0	*	*	0.1	*	0.1	0.1
1999.....	0.0	*	*	*	0.1	*	*

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.1	*	*	0.3	0.5	0.4	0.3
2005.....	0.1	*	*	0.2	0.4	0.4	0.3
2004.....	0.1	*	*	0.3	0.4	0.3	0.2
2003.....	0.1	*	*	0.2	*	0.3	0.2
2002.....	0.1	*	*	0.2	0.3	0.3	0.3
2001.....	0.1	*	*	*	0.3	0.2	*
2000.....	0.1	*	*	0.2	*	0.2	0.2
1999.....	0.1	*	*	*	0.2	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other direct obstetric causes (010,012,021-043,047-066,068-071,073-075)										
White, both sexes										
2006 4/.....	0.0	*	*	0.1	0.2	0.1	*
2005 4/.....	0.0	*	*	*	0.1	0.1	0.1
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	0.1	*	*	0.3	0.3	0.3	*
2005 4/.....	0.1	*	*	*	0.3	0.3	0.2
2004 4/.....	0.1	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other direct obstetric causes (010,012,021-043,047-066,068-071,073-075)

Black, both sexes

2006 4/.....	0.2	*	*	*	0.7	*	*
2005 4/.....	0.2	*	*	*	0.7	*	*
2004 4/.....	0.2	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	0.4	*	*	*	1.4	*	*
2005 4/.....	0.3	*	*	*	1.4	*	*
2004 4/.....	0.3	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Obstetric death of unspecified cause (O95)										
All races 3/, both sexes										
2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	0.0	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Obstetric death of unspecified cause (O95)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Obstetric death of unspecified cause (O95)										
Black, both sexes										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

All races 3/, both sexes

2006.....	0.1	*	*	0.2	0.2	0.2	0.2
2005.....	0.0	*	*	0.1	0.1	0.2	*
2004.....	0.1	*	*	*	0.1	0.2	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.1	*	*	0.4	0.3	0.3	0.4
2005.....	0.1	*	*	0.3	0.2	0.4	*
2004.....	0.1	*	*	*	0.2	0.3	*
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other deaths related to pregnancy, childbirth and the puerperium (096-097)										
White, both sexes										
2006 4/.....	0.1	*	*	0.2	0.1	0.2	0.1
2005 4/.....	0.0	*	*	*	*	0.2	*
2004 4/.....	0.0	*	*	*	*	0.1	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	0.1	*	*	0.3	0.3	0.3	0.3
2005 4/.....	0.1	*	*	*	*	0.4	*
2004 4/.....	0.1	*	*	*	*	0.3	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other deaths related to pregnancy, childbirth and the puerperium (096-097)										
Black, both sexes										
2006 4/.....	0.1	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	0.2	*	*	*	*	*	*
2005 4/.....	0.2	*	*	*	*	*	*
2004 4/.....	0.2	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Indirect obstetric deaths (O98-O99)										
All races 3/, both sexes										
2006.....	0.0	*	*	0.1	0.1	0.1	0.1
2005.....	0.1	*	*	0.1	0.1	0.2	0.1
2004.....	0.0	*	*	0.1	0.1	0.1	0.1
2003.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	0.1	*	*	0.2	0.2	0.2	0.2
2005.....	0.1	*	*	0.3	0.3	0.4	0.2
2004.....	0.1	*	*	0.3	0.3	0.2	0.2
2003.....	0.1	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Indirect obstetric deaths (O98-O99)										
White, both sexes										
2006 4/.....	0.0	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	0.1	*
2004 4/.....	0.0	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	0.1	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	0.3	*
2004 4/.....	0.1	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Indirect obstetric deaths (O98-O99)										
Black, both sexes										
2006 4/.....	0.1	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	0.2	*	*	*	*	*	*
2005 4/.....	0.3	*	*	*	*	*	*
2004 4/.....	0.2	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVI. Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes

2006.....	4.8	346.7	0.4	*	*	*	*	*	*	*
2005.....	4.9	351.2	0.4	*	*	*	*	*	*	*
2004.....	4.8	345.3	0.4	0.1	*	*	*	*	*	*
2003.....	4.9	356.0	0.5	*	*	*	*	*	*	*
2002.....	4.9	349.7	0.4	*	*	*	*	*	*	*
2001.....	4.9	340.5	0.5	0.1	*	*	*	*	*	*
2000.....	5.0	365.7	0.5	0.1	*	*	*	*	*	*
1999.....	5.1	371.0	0.6	0.1	*	*	*	*	*	*

All races 3/, male

2006.....	5.5	379.8	0.4	*	*	*	*	*	*	*
2005.....	5.6	386.1	0.4	*	*	*	*	*	*	*
2004.....	5.6	384.2	0.5	*	*	*	*	*	*	*
2003.....	5.7	397.4	0.5	*	*	*	*	*	*	*
2002.....	5.7	385.7	0.5	*	*	*	*	*	*	*
2001.....	5.6	376.6	0.6	*	*	*	*	*	*	*
2000.....	5.8	403.9	0.7	*	*	*	*	*	*	*
1999.....	5.9	408.6	0.7	*	*	*	*	*	*	*

All races 3/, female

2006.....	4.2	312.1	0.4	*	*	*	*	*	*	*
2005.....	4.2	314.6	0.3	*	*	*	*	*	*	*
2004.....	4.1	304.6	0.3	*	*	*	*	*	*	*
2003.....	4.2	312.8	0.5	*	*	*	*	*	*	*
2002.....	4.2	312.0	0.3	*	*	*	*	*	*	*
2001.....	4.2	302.6	0.4	*	*	*	*	*	*	*
2000.....	4.3	325.6	0.4	*	*	*	*	*	*	*
1999.....	4.4	331.7	0.5	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2006.....	3.5	140.9	3.2	0.9	0.8	1.1	1.1	1.1	1.0	1.1
2005.....	3.5	135.2	3.2	1.0	1.0	1.2	1.2	1.2	1.0	1.1
2004.....	3.6	137.9	3.6	1.0	0.9	1.2	1.1	1.1	1.0	1.0
2003.....	3.6	140.4	3.4	0.9	1.0	1.1	1.1	1.0	1.1	1.1
2002.....	3.7	139.4	3.4	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.7	136.7	3.6	0.9	0.9	1.3	1.3	1.2	1.1	1.2
2000.....	3.8	150.9	3.2	1.0	1.0	1.1	1.1	1.1	1.3	1.2
1999.....	3.7	144.2	3.6	1.0	1.1	1.1	1.1	1.1	1.2	1.2

All races 3/, male

2006.....	3.8	146.6	2.9	0.9	0.9	1.3	1.3	1.4	1.2	1.3
2005.....	3.7	141.1	3.2	0.9	1.0	1.6	1.4	1.3	1.1	1.2
2004.....	3.8	139.0	3.9	1.1	0.9	1.4	1.3	1.2	1.1	1.2
2003.....	3.8	147.6	3.4	0.9	1.0	1.4	1.2	1.2	1.1	1.2
2002.....	3.9	142.6	3.5	1.0	1.1	1.4	1.3	1.4	1.4	1.2
2001.....	3.9	143.1	3.6	1.0	1.0	1.5	1.4	1.3	1.2	1.2
2000.....	4.0	155.4	3.3	1.0	1.0	1.3	1.4	1.2	1.5	1.4
1999.....	3.9	147.9	4.0	1.0	1.1	1.3	1.3	1.1	1.3	1.3

All races 3/, female

2006.....	3.2	134.9	3.4	1.0	0.7	0.8	0.9	0.9	0.9	0.9
2005.....	3.3	129.0	3.3	1.2	0.9	0.8	1.0	1.0	0.9	1.0
2004.....	3.4	136.8	3.2	1.0	0.8	1.1	0.9	0.9	1.0	0.8
2003.....	3.4	132.8	3.5	0.9	1.0	0.8	1.0	0.8	1.1	1.1
2002.....	3.5	136.1	3.3	1.0	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	129.9	3.7	0.8	0.8	1.0	1.1	1.1	1.0	1.1
2000.....	3.5	146.2	3.1	0.9	1.0	0.9	0.9	0.9	1.1	1.1
1999.....	3.5	140.3	3.2	1.0	1.1	0.9	1.0	1.1	1.2	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2006 4/.....	3.4	137.2	3.0	0.9	0.8	1.0	1.0	1.1	1.1	1.1
2005 4/.....	3.4	131.0	3.1	0.9	0.9	1.1	1.2	1.1	1.0	1.1
2004 4/.....	3.5	134.5	3.2	1.0	0.9	1.1	1.0	1.1	1.1	1.0
2003 4/.....	3.5	137.6	3.2	0.9	1.0	1.1	1.0	1.1	1.1	1.2
2002.....	3.6	138.3	3.3	1.0	1.0	1.2	1.2	1.1	1.3	1.2
2001.....	3.6	135.0	3.2	0.9	0.9	1.2	1.2	1.2	1.2	1.1
2000.....	3.6	147.1	2.9	0.9	0.9	1.0	1.2	1.1	1.4	1.2
1999.....	3.6	141.3	3.2	0.9	1.0	1.1	1.1	1.1	1.3	1.2

White, male

2006 4/.....	3.7	142.3	2.9	1.0	0.9	1.2	1.2	1.4	1.2	1.3
2005 4/.....	3.6	136.9	3.1	0.8	1.0	1.5	1.4	1.2	1.1	1.2
2004 4/.....	3.6	134.3	3.5	1.1	0.9	1.2	1.1	1.3	1.1	1.1
2003 4/.....	3.7	143.2	3.1	0.8	0.9	1.3	1.1	1.2	1.1	1.2
2002.....	3.8	143.3	3.4	0.9	1.0	1.3	1.3	1.4	1.5	1.2
2001.....	3.8	143.0	3.2	1.0	1.1	1.4	1.3	1.4	1.3	1.2
2000.....	3.9	151.9	2.9	0.9	0.9	1.1	1.4	1.2	1.6	1.4
1999.....	3.8	146.5	3.5	0.8	1.0	1.3	1.3	1.2	1.3	1.4

White, female

2006 4/.....	3.1	131.8	3.0	0.8	0.6	0.8	0.9	0.8	0.9	0.9
2005 4/.....	3.2	124.8	3.0	1.0	0.8	0.7	0.9	1.0	0.9	1.1
2004 4/.....	3.4	134.7	2.9	0.8	0.8	1.0	0.9	1.0	1.1	0.8
2003 4/.....	3.4	131.8	3.2	0.9	1.0	0.8	1.0	0.9	1.1	1.1
2002.....	3.5	133.1	3.1	1.1	1.0	1.0	1.1	0.9	1.1	1.2
2001.....	3.4	126.7	3.3	0.8	0.7	0.9	1.1	1.0	1.1	1.0
2000.....	3.5	142.0	2.9	1.0	0.9	0.9	0.9	0.9	1.1	1.0
1999.....	3.5	135.7	2.9	1.0	1.0	0.8	0.9	1.1	1.2	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, both sexes

2006 4/.....	4.6	170.8	4.1	1.1	0.9	1.3	1.3	1.5	1.0	1.6
2005 4/.....	4.4	162.9	4.0	1.3	1.3	1.6	1.4	1.6	1.3	1.3
2004 4/.....	4.5	162.9	5.1	1.5	1.0	1.8	1.7	1.1	1.1	1.2
2003 4/.....	4.5	157.5	4.7	1.1	1.1	1.2	1.7	1.2	1.5	1.2
2002.....	4.7	155.2	4.5	1.2	1.2	1.7	1.5	1.3	1.4	1.4
2001.....	4.6	158.4	5.3	1.0	1.2	1.9	1.8	1.4	1.2	1.5
2000.....	4.9	179.2	5.0	1.3	1.4	1.8	1.3	1.4	1.4	1.6
1999.....	4.8	169.3	5.5	1.4	1.7	1.4	1.5	1.0	1.2	1.4

Black, male

2006 4/.....	5.0	176.8	3.1	*	*	1.6	1.5	1.8	*	1.7
2005 4/.....	4.8	166.5	3.6	*	1.3	2.1	1.8	2.1	*	1.8
2004 4/.....	5.1	169.9	5.8	1.2	*	2.0	2.5	*	1.6	1.7
2003 4/.....	5.1	170.7	4.8	1.2	*	1.7	1.8	*	*	*
2002.....	5.1	151.7	4.6	1.4	1.4	2.1	1.7	1.7	1.6	*
2001.....	5.0	160.8	5.5	*	*	2.6	2.2	*	*	*
2000.....	5.4	181.4	5.5	1.6	1.4	2.5	*	1.9	1.7	1.8
1999.....	5.2	172.7	6.2	1.4	1.8	1.7	1.8	*	*	*

Black, female

2006 4/.....	4.2	164.5	5.0	1.3	*	*	*	*	*	1.5
2005 4/.....	4.1	159.0	4.5	1.5	1.4	*	*	*	1.5	*
2004 4/.....	4.0	155.6	4.3	1.8	*	1.6	*	*	*	*
2003 4/.....	3.9	143.8	4.6	*	*	*	1.5	*	1.6	*
2002.....	4.3	158.9	4.4	*	*	1.3	1.3	*	*	1.4
2001.....	4.3	155.9	5.1	*	1.4	1.3	1.5	1.7	*	1.7
2000.....	4.4	176.9	4.6	*	1.5	*	*	*	*	1.5
1999.....	4.3	165.8	4.8	1.5	1.7	*	*	*	*	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)

All races 3/, both sexes

2006.....	0.3	11.6	0.8	0.3	0.2	0.1	0.1	*	*	*
2005.....	0.3	9.8	0.8	0.3	0.2	0.2	0.1	0.1	*	0.1
2004.....	0.3	10.8	0.8	0.3	0.2	0.2	*	*	*	*
2003.....	0.3	10.5	0.6	0.3	0.2	0.2	0.1	*	0.1	*
2002.....	0.3	9.3	0.8	0.3	0.2	0.2	0.1	*	*	0.1
2001.....	0.3	9.7	0.8	0.2	0.2	0.2	0.1	0.1	*	*
2000.....	0.3	10.7	0.6	0.3	0.2	0.1	0.1	*	*	*
1999.....	0.3	9.4	0.7	0.3	0.2	0.1	*	0.1	0.1	*

All races 3/, male

2006.....	0.3	11.6	0.6	0.3	0.3	*	*	*	*	*
2005.....	0.3	9.8	0.8	0.3	0.2	0.3	*	*	*	*
2004.....	0.3	9.2	0.9	0.2	0.2	0.2	*	*	*	*
2003.....	0.3	10.8	0.7	0.2	*	0.2	*	*	*	*
2002.....	0.3	9.0	0.7	0.3	*	*	*	*	*	*
2001.....	0.3	9.4	0.8	0.2	0.2	*	*	*	*	*
2000.....	0.3	10.6	0.6	0.3	0.2	*	*	*	*	*
1999.....	0.3	8.6	0.7	0.3	0.2	*	*	*	*	*

All races 3/, female

2006.....	0.3	11.5	0.9	0.2	*	*	*	*	*	*
2005.....	0.3	9.9	0.7	0.3	0.2	*	*	*	*	*
2004.....	0.3	12.4	0.8	0.3	0.2	0.2	*	*	*	*
2003.....	0.3	10.2	0.6	0.3	*	*	*	*	*	*
2002.....	0.3	9.6	0.9	0.3	0.3	*	*	*	*	*
2001.....	0.3	10.0	0.9	*	*	*	*	*	*	*
2000.....	0.3	10.9	0.7	0.3	0.2	*	*	*	*	*
1999.....	0.3	10.2	0.7	0.3	0.2	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other congenital malformations of nervous system (Q01-Q04,Q06-Q07)										
White, both sexes										
2006 4/.....	0.3	11.5	0.7	0.3	0.2	0.1	*	*	*	*
2005 4/.....	0.3	9.7	0.7	0.3	0.3	0.2	*	*	*	*
2004 4/.....	0.3	10.7	0.8	0.3	0.2	0.2	*	*	*	*
2003 4/.....	0.3	10.2	0.6	0.2	0.2	0.2	0.1	*	*	*
2002.....	0.3	9.4	0.7	0.3	0.2	0.2	0.2	*	*	*
2001.....	0.3	9.6	0.7	0.2	0.2	0.2	*	*	*	*
2000.....	0.3	10.5	0.6	0.3	0.2	*	*	*	*	*
1999.....	0.2	8.8	0.6	0.2	0.2	*	*	*	*	*
White, male										
2006 4/.....	0.3	11.7	0.6	0.4	0.3	*	*	*	*	*
2005 4/.....	0.3	9.7	0.8	*	0.3	0.3	*	*	*	*
2004 4/.....	0.3	9.1	0.8	0.3	*	*	*	*	*	*
2003 4/.....	0.3	10.2	0.7	*	*	0.3	*	*	*	*
2002.....	0.3	9.0	0.6	0.3	*	*	*	*	*	*
2001.....	0.3	9.6	0.7	*	0.2	*	*	*	*	*
2000.....	0.3	10.4	0.6	*	*	*	*	*	*	*
1999.....	0.2	8.1	0.6	*	*	*	*	*	*	*
White, female										
2006 4/.....	0.3	11.3	0.8	*	*	*	*	*	*	*
2005 4/.....	0.3	9.8	0.7	0.3	0.3	*	*	*	*	*
2004 4/.....	0.3	12.3	0.7	*	*	*	*	*	*	*
2003 4/.....	0.3	10.1	0.6	0.3	*	*	*	*	*	*
2002.....	0.3	9.8	0.8	0.3	*	*	*	*	*	*
2001.....	0.2	9.7	0.8	*	*	*	*	*	*	*
2000.....	0.2	10.6	0.7	0.3	*	*	*	*	*	*
1999.....	0.2	9.4	0.6	0.3	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Congenital malformations of heart (Q20-Q24)										
All races 3/, both sexes										
2006.....	1.0	33.8	1.1	0.2	0.2	0.5	0.5	0.4	0.4	0.4
2005.....	1.0	33.5	1.1	0.3	0.3	0.6	0.5	0.6	0.4	0.5
2004.....	1.0	34.1	1.2	0.4	0.3	0.6	0.5	0.5	0.5	0.4
2003.....	1.1	36.1	1.4	0.3	0.4	0.4	0.4	0.5	0.5	0.5
2002.....	1.1	37.4	1.2	0.3	0.4	0.6	0.5	0.5	0.6	0.5
2001.....	1.2	38.6	1.3	0.2	0.4	0.6	0.6	0.5	0.5	0.5
2000.....	1.2	42.9	1.2	0.3	0.4	0.6	0.5	0.5	0.7	0.5
1999.....	1.3	42.1	1.3	0.3	0.5	0.6	0.5	0.6	0.6	0.6
All races 3/, male										
2006.....	1.1	36.7	1.1	0.2	0.2	0.6	0.6	0.6	0.5	0.5
2005.....	1.1	35.9	1.2	0.3	0.4	0.8	0.6	0.7	0.5	0.6
2004.....	1.1	36.7	1.4	0.4	0.3	0.7	0.6	0.6	0.6	0.5
2003.....	1.2	40.1	1.5	0.3	0.4	0.6	0.5	0.6	0.6	0.6
2002.....	1.2	40.4	1.5	0.3	0.5	0.7	0.6	0.7	0.8	0.6
2001.....	1.3	41.8	1.4	0.3	0.4	0.9	0.7	0.6	0.7	0.6
2000.....	1.3	44.5	1.3	0.3	0.4	0.7	0.6	0.6	0.9	0.6
1999.....	1.4	45.7	1.5	0.3	0.6	0.8	0.7	0.6	0.7	0.7
All races 3/, female										
2006.....	0.8	30.8	1.0	0.3	0.2	0.3	0.3	0.3	0.3	0.4
2005.....	0.9	31.0	1.1	0.4	0.3	0.3	0.4	0.5	0.4	0.4
2004.....	0.9	31.4	1.0	0.3	0.3	0.5	0.4	0.4	0.4	0.3
2003.....	1.0	31.9	1.4	0.3	0.4	0.2	0.3	0.4	0.5	0.5
2002.....	1.0	34.4	1.0	0.3	0.3	0.4	0.3	0.4	0.5	0.5
2001.....	1.1	35.2	1.2	0.2	0.3	0.3	0.5	0.5	0.4	0.5
2000.....	1.1	41.3	1.2	0.3	0.3	0.4	0.4	0.4	0.5	0.4
1999.....	1.2	38.3	1.1	0.3	0.4	0.4	0.4	0.5	0.6	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Congenital malformations of heart (Q20-Q24)										
White, both sexes										
2006 4/.....	0.9	32.6	1.0	0.2	0.2	0.4	0.4	0.4	0.4	0.5
2005 4/.....	1.0	33.6	1.1	0.3	0.3	0.5	0.5	0.6	0.4	0.5
2004 4/.....	1.0	33.2	1.1	0.3	0.3	0.5	0.4	0.5	0.5	0.4
2003 4/.....	1.0	35.0	1.3	0.3	0.4	0.4	0.3	0.5	0.5	0.5
2002.....	1.1	36.4	1.3	0.3	0.4	0.6	0.4	0.5	0.7	0.6
2001.....	1.1	37.4	1.2	0.2	0.3	0.5	0.6	0.5	0.6	0.5
2000.....	1.2	41.5	1.1	0.3	0.3	0.5	0.5	0.5	0.7	0.5
1999.....	1.3	42.0	1.1	0.3	0.4	0.6	0.5	0.6	0.7	0.6
White, male										
2006 4/.....	1.0	35.6	1.1	*	*	0.6	0.6	0.5	0.5	0.6
2005 4/.....	1.1	37.2	1.2	0.3	0.3	0.7	0.6	0.6	0.5	0.6
2004 4/.....	1.1	35.9	1.3	0.4	0.4	0.6	0.4	0.7	0.6	0.5
2003 4/.....	1.1	39.3	1.3	0.3	0.4	0.5	0.4	0.6	0.6	0.6
2002.....	1.2	40.0	1.5	0.3	0.4	0.7	0.6	0.6	0.8	0.6
2001.....	1.2	41.3	1.3	0.3	0.4	0.7	0.6	0.6	0.7	0.6
2000.....	1.3	43.6	1.2	0.3	0.4	0.6	0.6	0.6	0.9	0.6
1999.....	1.3	46.2	1.3	0.3	0.5	0.8	0.6	0.6	0.7	0.7
White, female										
2006 4/.....	0.8	29.5	0.8	*	*	0.3	0.3	0.3	0.3	0.3
2005 4/.....	0.9	29.9	1.0	*	*	0.3	0.3	0.5	0.4	0.5
2004 4/.....	0.9	30.5	0.9	0.3	0.3	0.4	0.4	0.4	0.4	0.3
2003 4/.....	1.0	30.5	1.3	0.3	0.4	*	*	0.4	0.5	0.5
2002.....	1.0	32.7	1.0	0.3	0.3	0.4	0.3	0.4	0.6	0.5
2001.....	1.0	33.2	1.0	*	0.3	0.3	0.5	0.4	0.4	0.5
2000.....	1.1	39.3	1.1	0.3	0.3	0.4	0.4	0.4	0.6	0.4
1999.....	1.2	37.6	1.0	0.3	0.3	0.4	0.4	0.6	0.6	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Congenital malformations of heart (Q20-Q24)										
Black, both sexes										
2006 4/.....	1.3	42.5	1.5	*	*	0.6	0.7	0.7	*	*
2005 4/.....	1.2	36.3	1.3	*	0.6	0.9	0.6	0.8	*	*
2004 4/.....	1.3	42.2	1.7	0.7	*	0.9	0.7	*	*	*
2003 4/.....	1.4	41.7	2.1	*	0.6	0.7	0.8	*	0.8	*
2002.....	1.4	44.3	1.3	*	*	0.8	0.7	*	0.7	*
2001.....	1.6	47.7	1.9	*	*	1.2	1.0	*	*	*
2000.....	1.6	51.5	1.8	*	0.7	1.1	0.7	*	*	0.7
1999.....	1.6	47.9	2.2	*	0.8	0.7	0.9	*	*	0.8
Black, male										
2006 4/.....	1.5	43.5	*	*	*	*	*	*	*	*
2005 4/.....	1.2	33.4	*	*	*	1.2	*	*	*	*
2004 4/.....	1.5	44.8	2.0	*	*	*	*	*	*	*
2003 4/.....	1.6	44.9	2.2	*	*	*	*	*	*	*
2002.....	1.6	43.0	*	*	*	*	*	*	*	*
2001.....	1.7	48.8	1.7	*	*	1.9	*	*	*	*
2000.....	1.8	49.9	2.1	*	*	1.6	*	*	*	*
1999.....	1.8	50.1	2.5	*	*	*	*	*	*	*
Black, female										
2006 4/.....	1.2	41.5	1.6	*	*	*	*	*	*	*
2005 4/.....	1.2	39.4	1.7	*	*	*	*	*	*	*
2004 4/.....	1.1	39.5	*	*	*	*	*	*	*	*
2003 4/.....	1.2	38.3	2.1	*	*	*	*	*	*	*
2002.....	1.3	45.7	*	*	*	*	*	*	*	*
2001.....	1.4	46.6	2.0	*	*	*	*	*	*	*
2000.....	1.4	53.2	*	*	*	*	*	*	*	*
1999.....	1.4	45.6	2.0	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other congenital malformations and deformations (Q10-Q18,Q35-Q89)

All races 3/, both sexes

2006.....	1.0	43.2	0.8	0.2	0.1	0.3	0.2	0.3	0.2	0.3
2005.....	0.9	38.1	0.7	0.2	0.2	0.2	0.3	0.2	0.2	0.2
2004.....	0.9	36.4	0.8	0.2	0.1	0.2	0.2	0.2	0.2	0.2
2003.....	0.9	37.5	0.7	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2002.....	0.9	36.2	0.7	0.2	0.2	0.3	0.3	0.2	0.2	0.2
2001.....	0.9	36.3	0.8	0.2	0.2	0.2	0.2	0.2	0.2	0.3
2000.....	0.9	37.9	0.7	0.2	0.2	0.2	0.2	0.2	0.2	0.3
1999.....	0.9	38.3	0.9	0.2	0.1	0.1	0.2	0.2	0.2	0.2

All races 3/, male

2006.....	1.1	47.6	0.7	*	*	0.3	0.2	0.4	0.2	0.3
2005.....	1.1	42.5	0.7	0.2	0.2	0.3	0.4	*	0.3	0.3
2004.....	1.0	38.7	0.9	*	*	0.3	0.3	0.2	0.2	0.3
2003.....	1.0	42.8	0.7	0.2	*	0.3	0.3	0.3	0.2	0.2
2002.....	1.0	39.2	0.6	0.2	0.2	0.3	0.3	0.2	*	0.2
2001.....	1.0	41.1	0.8	*	*	0.2	0.3	0.2	*	0.3
2000.....	1.0	42.1	0.7	0.2	*	0.2	0.2	0.2	0.3	0.3
1999.....	1.0	42.5	1.1	0.2	*	*	0.3	*	0.2	0.2

All races 3/, female

2006.....	0.9	38.5	0.8	0.2	*	*	0.2	0.2	0.2	0.2
2005.....	0.8	33.5	0.7	0.3	*	*	*	*	*	0.2
2004.....	0.9	34.0	0.7	*	*	0.2	0.2	*	0.2	*
2003.....	0.8	31.9	0.7	*	*	*	0.2	*	0.2	0.2
2002.....	0.8	33.1	0.8	*	*	0.2	0.3	*	*	*
2001.....	0.8	31.3	0.8	*	*	*	*	0.2	*	0.3
2000.....	0.8	33.6	0.6	*	0.2	*	0.2	*	*	0.2
1999.....	0.8	33.8	0.8	*	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other congenital malformations and deformations (Q10-Q18,Q35-Q89)										
White, both sexes										
2006 4/.....	1.0	41.6	0.8	0.2	0.1	0.3	0.2	0.3	0.2	0.2
2005 4/.....	0.9	35.9	0.6	0.2	0.2	0.2	0.3	0.2	0.2	0.2
2004 4/.....	0.9	35.6	0.8	0.2	0.1	0.2	0.2	0.2	0.2	0.2
2003 4/.....	0.9	36.4	0.7	0.1	0.1	0.2	0.2	0.2	0.2	0.3
2002.....	0.9	37.0	0.7	0.2	0.2	0.2	0.3	0.2	0.2	0.2
2001.....	0.9	36.0	0.7	0.2	0.1	0.2	0.2	0.2	0.2	0.3
2000.....	0.9	37.4	0.6	0.2	0.2	0.2	0.3	0.2	0.2	0.3
1999.....	0.9	36.8	0.9	0.2	0.1	0.2	0.2	0.2	0.2	0.2
White, male										
2006 4/.....	1.1	45.8	0.7	*	*	0.3	0.2	0.4	*	0.3
2005 4/.....	1.0	38.9	0.6	*	*	0.3	0.4	*	0.3	0.2
2004 4/.....	0.9	37.5	0.8	*	*	0.3	0.3	*	*	0.2
2003 4/.....	1.0	41.0	0.7	*	*	0.3	0.3	0.3	0.3	0.2
2002.....	1.0	40.4	0.6	*	*	0.2	0.3	0.3	*	*
2001.....	1.0	40.6	0.6	*	*	0.3	0.3	*	*	0.3
2000.....	1.0	41.3	0.7	*	*	*	0.3	*	0.2	0.3
1999.....	1.0	42.1	1.0	*	*	*	0.3	*	*	0.2
White, female										
2006 4/.....	0.9	37.2	0.8	*	*	*	*	*	*	*
2005 4/.....	0.8	32.7	0.7	0.3	*	*	*	*	*	0.3
2004 4/.....	0.9	33.7	0.7	*	*	*	*	*	0.3	*
2003 4/.....	0.8	31.6	0.7	*	*	*	*	*	*	0.3
2002.....	0.8	33.4	0.7	*	*	*	0.3	*	*	*
2001.....	0.8	31.1	0.7	*	*	*	*	*	*	0.3
2000.....	0.8	33.2	0.6	*	*	*	*	*	*	0.2
1999.....	0.8	31.2	0.7	*	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
All other chromosomal abnormalities, not elsewhere classified (Q91-Q99)										
All races 3/, both sexes										
2006.....	0.4	25.0	0.3	*	*	*	*	*	*	*
2005.....	0.3	22.1	0.2	*	*	*	*	*	*	*
2004.....	0.4	24.4	0.3	*	*	*	*	*	*	*
2003.....	0.4	23.5	0.3	*	*	*	*	*	*	*
2002.....	0.4	24.3	0.2	*	*	*	*	*	*	*
2001.....	0.4	22.7	0.3	*	*	*	*	*	*	*
2000.....	0.4	24.2	0.3	*	*	*	*	*	*	*
1999.....	0.3	22.5	0.3	*	*	*	*	*	*	*
All races 3/, male										
2006.....	0.3	21.6	*	*	*	*	*	*	*	*
2005.....	0.3	18.6	*	*	*	*	*	*	*	*
2004.....	0.3	20.2	*	*	*	*	*	*	*	*
2003.....	0.3	20.0	*	*	*	*	*	*	*	*
2002.....	0.3	20.7	*	*	*	*	*	*	*	*
2001.....	0.3	20.1	*	*	*	*	*	*	*	*
2000.....	0.3	21.4	*	*	*	*	*	*	*	*
1999.....	0.3	18.7	*	*	*	*	*	*	*	*
All races 3/, female										
2006.....	0.4	28.6	0.4	*	*	*	*	*	*	*
2005.....	0.4	25.7	0.3	*	*	*	*	*	*	*
2004.....	0.4	28.8	0.4	*	*	*	*	*	*	*
2003.....	0.4	27.1	0.4	*	*	*	*	*	*	*
2002.....	0.4	28.0	*	*	*	*	*	*	*	*
2001.....	0.4	25.4	0.5	*	*	*	*	*	*	*
2000.....	0.4	27.2	0.3	*	*	*	*	*	*	*
1999.....	0.4	26.5	0.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2006.....	10.6	83.8	1.5	0.3	0.3	0.9	2.2	2.4	2.6	3.2
2005.....	10.8	87.4	1.4	0.3	0.5	1.2	2.5	2.8	3.1	4.1
2004.....	10.3	82.4	1.2	0.3	0.3	1.0	2.1	2.7	2.8	3.6
2003.....	10.8	82.9	1.6	0.4	0.5	1.5	2.6	3.2	3.4	4.8
2002.....	10.4	85.7	1.6	0.3	0.4	1.3	2.4	3.0	3.6	4.6
2001.....	11.4	86.5	1.8	0.4	0.5	1.8	3.0	3.5	4.2	6.2
2000.....	11.3	95.6	1.6	0.4	0.4	1.6	2.9	3.4	4.3	5.7
1999.....	9.6	93.2	1.5	0.3	0.4	1.1	2.0	2.3	3.1	4.1

All races 3/, male

2006.....	9.5	95.2	1.6	0.4	0.4	1.1	2.9	3.2	3.7	4.0
2005.....	10.0	101.8	1.6	0.3	0.6	1.6	3.4	3.9	4.0	5.1
2004.....	9.5	96.9	1.2	0.3	0.2	1.3	3.0	3.6	3.6	4.6
2003.....	10.2	96.5	1.8	0.3	0.6	1.9	3.4	4.4	4.5	6.4
2002.....	10.1	98.5	1.8	0.4	0.4	1.6	3.2	4.1	4.9	6.1
2001.....	11.3	98.9	2.1	0.4	0.5	2.4	4.0	4.8	5.6	8.0
2000.....	11.5	110.6	1.7	0.5	0.5	1.9	4.0	4.7	5.9	7.6
1999.....	9.4	106.6	1.7	0.3	0.4	1.5	2.8	2.9	4.2	5.3

All races 3/, female

2006.....	11.7	71.9	1.3	0.3	0.2	0.7	1.5	1.5	1.5	2.4
2005.....	11.5	72.3	1.3	0.3	0.3	0.8	1.4	1.7	2.1	3.1
2004.....	11.0	67.3	1.1	0.2	0.4	0.7	1.3	1.7	2.0	2.7
2003.....	11.4	68.6	1.5	0.5	0.4	1.0	1.7	1.9	2.3	3.3
2002.....	10.7	72.3	1.4	0.2	0.3	0.8	1.6	1.8	2.3	3.1
2001.....	11.4	73.6	1.5	0.4	0.5	1.1	1.9	2.2	2.8	4.4
2000.....	11.2	79.9	1.5	0.4	0.4	1.2	1.8	2.2	2.8	3.7
1999.....	9.7	79.2	1.3	0.3	0.3	0.8	1.2	1.6	2.1	3.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes

2006 4/.....	11.0	70.2	1.2	0.3	0.3	0.9	2.2	2.4	2.6	3.2
2005 4/.....	11.1	72.9	1.2	0.2	0.4	1.1	2.4	2.7	3.1	4.0
2004 4/.....	10.5	65.8	1.0	0.2	0.2	1.0	2.2	2.6	2.7	3.6
2003 4/.....	11.1	68.1	1.4	0.4	0.4	1.5	2.5	3.1	3.4	4.7
2002.....	10.5	71.0	1.3	0.2	0.3	1.2	2.3	2.8	3.5	4.4
2001.....	11.6	70.9	1.5	0.3	0.4	1.8	3.0	3.5	4.1	6.0
2000.....	11.4	77.2	1.5	0.4	0.4	1.5	2.8	3.2	4.2	5.6
1999.....	9.4	74.7	1.3	0.3	0.3	1.0	1.8	2.0	2.9	3.7

White, male

2006 4/.....	9.5	81.8	1.3	0.3	0.3	1.1	2.9	3.3	3.6	4.0
2005 4/.....	10.0	87.7	1.4	*	0.5	1.6	3.4	3.8	4.1	4.9
2004 4/.....	9.4	76.9	1.1	0.3	*	1.2	3.0	3.6	3.5	4.6
2003 4/.....	10.1	82.4	1.4	0.3	0.4	2.0	3.4	4.2	4.5	6.1
2002.....	9.8	82.5	1.4	0.3	0.4	1.6	3.2	3.9	4.7	5.8
2001.....	11.2	82.0	1.7	0.3	0.5	2.4	4.0	4.9	5.5	7.7
2000.....	11.3	89.4	1.7	0.4	0.4	1.9	3.9	4.4	5.7	7.7
1999.....	9.0	86.4	1.6	0.3	0.3	1.3	2.6	2.7	3.8	4.9

White, female

2006 4/.....	12.5	58.1	1.1	*	*	0.6	1.5	1.5	1.5	2.4
2005 4/.....	12.2	57.4	1.1	*	*	0.7	1.4	1.5	2.0	3.0
2004 4/.....	11.6	54.2	1.0	*	0.3	0.7	1.3	1.5	1.9	2.5
2003 4/.....	12.0	53.3	1.3	0.4	0.3	1.0	1.6	1.9	2.1	3.2
2002.....	11.2	59.0	1.2	*	*	0.8	1.4	1.6	2.2	3.0
2001.....	11.9	59.3	1.4	0.4	0.4	1.2	1.9	1.9	2.7	4.3
2000.....	11.5	64.3	1.3	0.3	0.4	1.1	1.6	1.9	2.6	3.5
1999.....	9.9	62.4	1.1	0.3	*	0.7	1.0	1.3	2.1	2.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2006 4/.....	11.1	155.8	2.8	0.6	*	1.1	2.5	2.9	3.9	4.3
2005 4/.....	12.3	169.0	2.5	0.8	0.9	1.9	3.1	4.2	4.2	6.0
2004 4/.....	11.6	171.7	1.9	*	0.6	1.3	2.3	3.9	4.2	5.2
2003 4/.....	12.3	159.3	3.1	*	1.0	1.5	3.4	4.9	5.0	7.1
2002.....	12.3	162.3	2.6	0.7	0.7	1.8	3.2	4.6	5.6	7.5
2001.....	12.6	169.5	2.8	0.6	0.6	1.8	3.8	4.6	5.3	8.2
2000.....	13.7	197.1	2.3	0.6	0.8	2.2	4.3	5.3	6.3	7.3
1999.....	12.7	192.8	2.7	*	*	1.7	3.7	4.3	5.4	7.8

Black, male

2006 4/.....	11.7	165.9	3.1	*	*	1.4	3.1	3.7	5.7	5.2
2005 4/.....	13.3	186.6	2.7	*	*	2.4	4.5	5.7	5.2	7.1
2004 4/.....	12.6	204.3	2.1	*	*	1.6	3.1	4.8	5.5	6.3
2003 4/.....	13.5	170.7	3.4	*	1.3	1.8	4.2	7.3	6.2	9.6
2002.....	14.1	183.0	3.5	*	*	2.4	3.6	5.8	7.8	10.3
2001.....	13.9	187.4	3.5	*	*	2.4	5.2	5.5	6.5	10.5
2000.....	15.5	227.1	2.3	*	*	2.0	5.6	7.1	8.2	8.3
1999.....	14.6	217.6	2.8	*	*	2.4	4.7	5.2	8.3	9.4

Black, female

2006 4/.....	10.5	145.3	2.5	*	*	*	1.9	2.1	2.3	3.5
2005 4/.....	11.4	150.7	2.2	*	*	1.3	1.7	2.8	3.3	5.0
2004 4/.....	10.8	137.7	1.7	*	*	*	1.5	3.0	2.9	4.3
2003 4/.....	11.3	147.5	2.9	*	*	*	2.7	2.7	3.9	4.9
2002.....	10.6	140.8	1.8	*	*	*	2.7	3.5	3.7	5.1
2001.....	11.4	150.6	2.0	*	*	*	2.4	3.9	4.2	6.2
2000.....	12.0	166.0	2.3	*	*	2.3	3.0	3.6	4.6	6.3
1999.....	10.9	167.1	2.7	*	*	*	2.6	3.4	2.8	6.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ill-defined and unknown causes of mortality (R95-R99)										
All races 3/, both sexes										
2006.....	4.8	82.0	1.3	0.2	0.2	0.6	1.6	1.7	2.1	2.5
2005.....	5.7	85.8	1.2	0.2	0.3	1.0	2.0	2.2	2.6	3.4
2004.....	5.3	80.7	1.0	0.2	0.2	0.8	1.7	2.1	2.2	3.0
2003.....	6.0	81.2	1.3	0.3	0.4	1.2	2.0	2.5	2.7	4.0
2002.....	6.0	83.8	1.3	0.2	0.3	1.0	1.9	2.3	2.8	3.7
2001.....	7.1	84.8	1.5	0.3	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	7.3	93.9	1.2	0.3	0.3	1.3	2.4	2.7	3.7	4.8
1999.....	5.9	91.3	1.3	0.2	0.3	0.9	1.7	1.7	2.6	3.3
All races 3/, male										
2006.....	5.3	93.1	1.4	0.2	0.2	0.8	2.1	2.4	3.0	3.1
2005.....	6.5	100.1	1.3	0.2	0.4	1.3	2.8	3.2	3.4	4.3
2004.....	6.0	94.9	1.1	0.2	*	1.0	2.3	2.9	2.9	3.8
2003.....	6.8	94.3	1.4	0.2	0.4	1.6	2.7	3.4	3.7	5.3
2002.....	6.8	96.5	1.5	0.3	0.3	1.3	2.6	3.2	3.8	5.1
2001.....	8.2	96.5	1.8	0.3	0.3	2.0	3.4	4.1	4.9	6.9
2000.....	8.5	108.9	1.4	0.3	0.4	1.6	3.3	3.7	5.1	6.5
1999.....	6.7	104.3	1.4	0.2	0.3	1.2	2.4	2.2	3.5	4.2
All races 3/, female										
2006.....	4.2	70.2	1.1	*	*	0.4	1.0	1.0	1.2	1.8
2005.....	5.0	70.8	1.1	0.2	*	0.6	1.1	1.2	1.6	2.5
2004.....	4.6	65.8	0.9	*	0.2	0.6	1.0	1.2	1.5	2.2
2003.....	5.2	67.6	1.2	0.4	0.3	0.9	1.2	1.5	1.7	2.7
2002.....	5.1	70.4	1.1	*	0.2	0.6	1.2	1.4	1.8	2.4
2001.....	6.1	72.4	1.2	0.2	0.4	1.0	1.5	1.7	2.3	3.8
2000.....	6.2	78.0	1.1	0.2	0.3	1.0	1.4	1.7	2.2	3.2
1999.....	5.2	77.7	1.1	0.3	0.2	0.6	0.9	1.2	1.6	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ill-defined and unknown causes of mortality (R95-R99)										
White, both sexes										
2006 4/.....	4.7	68.6	1.0	0.1	0.2	0.6	1.7	1.9	2.1	2.5
2005 4/.....	5.6	71.6	1.0	0.2	0.2	0.9	2.0	2.3	2.6	3.4
2004 4/.....	5.2	64.4	0.8	0.2	0.1	0.8	1.8	2.1	2.2	3.0
2003 4/.....	5.9	66.4	1.1	0.3	0.3	1.3	2.0	2.4	2.7	4.0
2002.....	5.8	69.5	1.1	0.2	0.2	1.0	2.0	2.2	2.8	3.6
2001.....	7.1	69.3	1.3	0.2	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	7.2	75.6	1.2	0.3	0.3	1.2	2.3	2.6	3.7	4.9
1999.....	5.6	73.1	1.1	0.2	0.2	0.8	1.6	1.6	2.4	3.1
White, male										
2006 4/.....	5.2	79.8	1.1	*	*	0.8	2.2	2.6	3.0	3.2
2005 4/.....	6.3	86.2	1.1	*	0.4	1.3	2.8	3.3	3.5	4.3
2004 4/.....	5.8	75.7	0.9	*	*	1.0	2.5	3.1	3.0	3.9
2003 4/.....	6.6	80.2	1.1	*	0.3	1.7	2.8	3.3	3.8	5.2
2002.....	6.5	81.1	1.1	*	0.2	1.3	2.7	3.2	3.8	4.9
2001.....	8.1	79.9	1.5	*	0.3	2.0	3.5	4.3	4.8	6.9
2000.....	8.2	87.8	1.4	0.3	0.3	1.6	3.3	3.6	5.2	6.6
1999.....	6.2	84.5	1.3	*	0.2	1.1	2.3	2.1	3.3	4.0
White, female										
2006 4/.....	4.2	56.8	0.9	*	*	0.4	1.1	1.1	1.2	1.8
2005 4/.....	5.0	56.2	0.9	*	*	0.5	1.1	1.2	1.6	2.5
2004 4/.....	4.6	52.7	0.7	*	*	0.6	1.0	1.1	1.5	2.1
2003 4/.....	5.2	51.9	1.1	0.3	0.2	0.8	1.2	1.5	1.6	2.7
2002.....	5.1	57.4	1.0	*	*	0.6	1.2	1.2	1.8	2.4
2001.....	6.1	58.2	1.1	*	0.3	1.0	1.5	1.6	2.2	3.8
2000.....	6.2	62.7	0.9	*	0.3	0.9	1.3	1.6	2.1	3.1
1999.....	5.1	61.2	0.8	0.3	*	0.5	0.7	1.0	1.6	2.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ill-defined and unknown causes of mortality (R95-R99)										
Black, both sexes										
2006 4/.....	6.0	153.4	2.6	*	*	0.8	1.5	1.6	2.8	2.9
2005 4/.....	8.0	165.7	2.1	*	*	1.3	2.3	2.8	3.1	4.3
2004 4/.....	7.3	168.5	1.7	*	*	0.9	1.4	2.5	2.6	3.8
2003 4/.....	8.2	158.1	2.5	*	0.7	1.2	2.5	3.4	3.8	5.4
2002.....	8.4	158.0	2.3	*	0.6	1.1	2.0	3.4	3.6	5.6
2001.....	8.8	167.3	2.3	*	*	1.2	2.5	3.4	4.4	6.3
2000.....	9.9	194.0	1.8	*	0.7	1.7	3.3	3.8	4.6	5.8
1999.....	9.1	189.1	2.3	*	*	1.3	2.7	2.8	4.0	5.8
Black, male										
2006 4/.....	7.3	163.9	2.9	*	*	*	1.9	2.5	4.0	3.6
2005 4/.....	9.5	183.1	2.2	*	*	1.8	3.1	3.8	4.0	5.0
2004 4/.....	8.8	199.0	1.7	*	*	*	1.8	2.9	3.5	4.2
2003 4/.....	9.9	168.7	2.8	*	*	1.4	3.2	5.2	4.7	7.2
2002.....	10.4	178.1	2.9	*	*	1.4	2.5	4.0	5.1	8.0
2001.....	10.4	184.7	2.9	*	*	1.7	3.5	4.2	5.3	7.9
2000.....	12.0	224.2	1.9	*	*	1.7	4.2	5.2	6.3	6.5
1999.....	11.2	212.7	2.3	*	*	1.7	3.6	3.7	6.2	7.3
Black, female										
2006 4/.....	4.9	142.6	2.2	*	*	*	*	*	1.6	2.4
2005 4/.....	6.6	147.6	2.0	*	*	*	1.4	1.9	2.2	3.7
2004 4/.....	6.0	136.8	1.6	*	*	*	*	2.1	1.7	3.5
2003 4/.....	6.7	147.2	2.1	*	*	*	1.9	1.7	2.9	3.8
2002.....	6.5	137.1	1.7	*	*	*	1.6	2.8	2.2	3.5
2001.....	7.4	149.0	1.6	*	*	*	1.5	2.6	3.5	4.9
2000.....	7.9	162.7	1.6	*	*	1.6	2.4	2.5	3.1	5.2
1999.....	7.2	164.8	2.4	*	*	*	1.9	2.0	2.1	4.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other ill-defined and unknown causes of mortality (R96-R99)										
All races 3/, both sexes										
2006.....	4.0	25.7	1.3	0.2	0.2	0.6	1.6	1.7	2.1	2.5
2005.....	5.0	31.5	1.2	0.2	0.3	1.0	2.0	2.2	2.6	3.4
2004.....	4.5	25.6	1.0	0.2	0.2	0.8	1.7	2.1	2.2	3.0
2003.....	5.2	27.2	1.3	0.3	0.4	1.2	2.0	2.5	2.7	4.0
2002.....	5.2	26.9	1.3	0.2	0.3	1.0	1.9	2.3	2.8	3.7
2001.....	6.3	29.4	1.5	0.3	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	6.4	27.6	1.2	0.3	0.3	1.3	2.4	2.7	3.7	4.8
1999.....	5.0	21.6	1.3	0.2	0.3	0.9	1.7	1.7	2.6	3.3
All races 3/, male										
2006.....	4.4	28.9	1.4	0.2	0.2	0.8	2.1	2.4	3.0	3.1
2005.....	5.6	34.8	1.3	0.2	0.4	1.3	2.8	3.2	3.4	4.3
2004.....	5.1	31.2	1.1	0.2	*	1.0	2.3	2.9	2.9	3.8
2003.....	5.9	31.7	1.4	0.2	0.4	1.6	2.7	3.4	3.7	5.3
2002.....	5.9	31.1	1.5	0.3	0.3	1.3	2.6	3.2	3.8	5.1
2001.....	7.3	32.7	1.8	0.3	0.3	2.0	3.4	4.1	4.9	6.9
2000.....	7.4	31.5	1.4	0.3	0.4	1.6	3.3	3.7	5.1	6.5
1999.....	5.5	24.4	1.4	0.2	0.3	1.2	2.4	2.2	3.5	4.2
All races 3/, female										
2006.....	3.5	22.4	1.1	*	*	0.4	1.0	1.0	1.2	1.8
2005.....	4.4	28.0	1.1	0.2	*	0.6	1.1	1.2	1.6	2.5
2004.....	4.0	19.7	0.9	*	0.2	0.6	1.0	1.2	1.5	2.2
2003.....	4.6	22.5	1.2	0.4	0.3	0.9	1.2	1.5	1.7	2.7
2002.....	4.5	22.5	1.1	*	0.2	0.6	1.2	1.4	1.8	2.4
2001.....	5.5	25.8	1.2	0.2	0.4	1.0	1.5	1.7	2.3	3.8
2000.....	5.5	23.4	1.1	0.2	0.3	1.0	1.4	1.7	2.2	3.2
1999.....	4.5	18.6	1.1	0.3	0.2	0.6	0.9	1.2	1.6	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other ill-defined and unknown causes of mortality (R96-R99)

White, both sexes

2006 4/.....	4.1	21.1	1.0	0.1	0.2	0.6	1.7	1.9	2.1	2.5
2005 4/.....	5.0	24.9	1.0	0.2	0.2	0.9	2.0	2.3	2.6	3.4
2004 4/.....	4.6	19.1	0.8	0.2	0.1	0.8	1.8	2.1	2.2	3.0
2003 4/.....	5.3	22.5	1.1	0.3	0.3	1.3	2.0	2.4	2.7	4.0
2002.....	5.2	21.8	1.1	0.2	0.2	1.0	2.0	2.2	2.8	3.6
2001.....	6.5	24.6	1.3	0.2	0.3	1.5	2.5	3.0	3.6	5.4
2000.....	6.5	21.5	1.2	0.3	0.3	1.2	2.3	2.6	3.7	4.9
1999.....	4.9	15.7	1.1	0.2	0.2	0.8	1.6	1.6	2.4	3.1

White, male

2006 4/.....	4.5	24.4	1.1	*	*	0.8	2.2	2.6	3.0	3.2
2005 4/.....	5.5	27.6	1.1	*	0.4	1.3	2.8	3.3	3.5	4.3
2004 4/.....	5.1	23.5	0.9	*	*	1.0	2.5	3.1	3.0	3.9
2003 4/.....	5.9	27.3	1.1	*	0.3	1.7	2.8	3.3	3.8	5.2
2002.....	5.8	25.6	1.1	*	0.2	1.3	2.7	3.2	3.8	4.9
2001.....	7.3	27.6	1.5	*	0.3	2.0	3.5	4.3	4.8	6.9
2000.....	7.4	24.4	1.4	0.3	0.3	1.6	3.3	3.6	5.2	6.6
1999.....	5.3	18.2	1.3	*	0.2	1.1	2.3	2.1	3.3	4.0

White, female

2006 4/.....	3.7	17.7	0.9	*	*	0.4	1.1	1.1	1.2	1.8
2005 4/.....	4.5	22.0	0.9	*	*	0.5	1.1	1.2	1.6	2.5
2004 4/.....	4.1	14.4	0.7	*	*	0.6	1.0	1.1	1.5	2.1
2003 4/.....	4.7	17.6	1.1	0.3	0.2	0.8	1.2	1.5	1.6	2.7
2002.....	4.6	17.9	1.0	*	*	0.6	1.2	1.2	1.8	2.4
2001.....	5.6	21.4	1.1	*	0.3	1.0	1.5	1.6	2.2	3.8
2000.....	5.7	18.4	0.9	*	0.3	0.9	1.3	1.6	2.1	3.1
1999.....	4.5	13.1	0.8	0.3	*	0.5	0.7	1.0	1.6	2.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other ill-defined and unknown causes of mortality (R96-R99)

Black, both sexes

2006 4/.....	4.2	48.0	2.6	*	*	0.8	1.5	1.6	2.8	2.9
2005 4/.....	6.3	65.7	2.1	*	*	1.3	2.3	2.8	3.1	4.3
2004 4/.....	5.4	58.7	1.7	*	*	0.9	1.4	2.5	2.6	3.8
2003 4/.....	6.4	53.4	2.5	*	0.7	1.2	2.5	3.4	3.8	5.4
2002.....	6.5	53.8	2.3	*	0.6	1.1	2.0	3.4	3.6	5.6
2001.....	6.8	53.0	2.3	*	*	1.2	2.5	3.4	4.4	6.3
2000.....	7.6	59.7	1.8	*	0.7	1.7	3.3	3.8	4.6	5.8
1999.....	6.8	50.5	2.3	*	*	1.3	2.7	2.8	4.0	5.8

Black, male

2006 4/.....	5.2	51.3	2.9	*	*	*	1.9	2.5	4.0	3.6
2005 4/.....	7.6	74.4	2.2	*	*	1.8	3.1	3.8	4.0	5.0
2004 4/.....	6.4	69.7	1.7	*	*	*	1.8	2.9	3.5	4.2
2003 4/.....	7.9	59.5	2.8	*	*	1.4	3.2	5.2	4.7	7.2
2002.....	8.2	60.7	2.9	*	*	1.4	2.5	4.0	5.1	8.0
2001.....	8.0	58.4	2.9	*	*	1.7	3.5	4.2	5.3	7.9
2000.....	9.3	70.0	1.9	*	*	1.7	4.2	5.2	6.3	6.5
1999.....	8.5	57.6	2.3	*	*	1.7	3.6	3.7	6.2	7.3

Black, female

2006 4/.....	3.3	44.5	2.2	*	*	*	*	*	1.6	2.4
2005 4/.....	5.2	56.7	2.0	*	*	*	1.4	1.9	2.2	3.7
2004 4/.....	4.6	47.2	1.6	*	*	*	*	2.1	1.7	3.5
2003 4/.....	5.1	47.0	2.1	*	*	*	1.9	1.7	2.9	3.8
2002.....	5.0	46.6	1.7	*	*	*	1.6	2.8	2.2	3.5
2001.....	5.7	47.3	1.6	*	*	*	1.5	2.6	3.5	4.9
2000.....	6.1	48.9	1.6	*	*	1.6	2.4	2.5	3.1	5.2
1999.....	5.3	43.3	2.4	*	*	*	1.9	2.0	2.1	4.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

All races 3/, both sexes

2006.....	4.2	1.9	0.2	0.2	0.1	0.3	0.6	0.6	0.6	0.7
2005.....	3.5	1.6	0.2	*	0.2	0.3	0.5	0.6	0.5	0.7
2004.....	3.3	1.7	0.2	*	0.1	0.2	0.5	0.6	0.6	0.7
2003.....	3.2	1.6	0.3	0.1	0.1	0.2	0.6	0.7	0.7	0.8
2002.....	2.9	1.9	0.3	*	0.1	0.3	0.5	0.6	0.8	0.9
2001.....	2.7	1.8	0.3	0.1	0.1	0.3	0.5	0.5	0.6	0.8
2000.....	2.6	1.8	0.4	0.1	0.1	0.3	0.5	0.7	0.7	0.8
1999.....	2.5	1.9	0.3	*	*	0.3	0.4	0.6	0.6	0.8

All races 3/, male

2006.....	3.2	2.1	*	*	*	0.3	0.8	0.8	0.8	0.9
2005.....	2.6	1.7	0.3	*	0.2	0.3	0.7	0.7	0.6	0.8
2004.....	2.5	2.0	*	*	*	0.3	0.6	0.7	0.7	0.8
2003.....	2.5	2.2	0.4	*	*	0.3	0.7	1.0	0.8	1.1
2002.....	2.4	1.9	0.4	*	*	0.4	0.5	0.8	1.0	1.1
2001.....	2.3	2.3	0.3	*	0.2	0.4	0.6	0.7	0.7	1.0
2000.....	2.3	1.7	0.3	*	*	0.3	0.7	0.9	0.7	1.1
1999.....	2.1	2.3	0.3	*	*	0.3	0.4	0.7	0.7	1.0

All races 3/, female

2006.....	5.2	1.6	*	*	*	0.2	0.5	0.5	0.3	0.6
2005.....	4.3	1.5	*	*	*	0.2	0.3	0.4	0.5	0.6
2004.....	4.1	1.5	*	*	*	*	0.3	0.5	0.5	0.5
2003.....	3.9	1.1	0.3	*	*	*	0.5	0.4	0.6	0.6
2002.....	3.5	1.9	*	*	*	0.2	0.4	0.4	0.6	0.7
2001.....	3.2	1.2	0.3	*	*	*	0.4	0.4	0.5	0.6
2000.....	3.0	1.8	0.4	*	*	0.3	0.4	0.5	0.6	0.5
1999.....	2.9	1.5	*	*	*	*	0.3	0.5	0.5	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

White, both sexes

2006 4/.....	4.4	1.7	0.2	*	*	0.3	0.6	0.5	0.5	0.7
2005 4/.....	3.6	1.3	0.2	*	0.1	0.2	0.5	0.4	0.5	0.6
2004 4/.....	3.5	1.4	0.2	*	*	0.2	0.4	0.5	0.5	0.6
2003 4/.....	3.3	1.8	0.3	*	*	0.2	0.5	0.6	0.6	0.7
2002.....	3.0	1.5	0.3	*	*	0.2	0.4	0.6	0.7	0.8
2001.....	2.8	1.7	0.2	*	0.1	0.3	0.4	0.5	0.6	0.7
2000.....	2.7	1.6	0.4	*	0.1	0.3	0.5	0.6	0.5	0.7
1999.....	2.5	1.6	0.3	*	*	0.2	0.3	0.5	0.5	0.7

White, male

2006 4/.....	3.2	2.0	*	*	*	0.3	0.7	0.7	0.6	0.8
2005 4/.....	2.6	1.5	*	*	*	0.3	0.6	0.5	0.5	0.6
2004 4/.....	2.5	1.2	*	*	*	0.3	0.5	0.5	0.5	0.7
2003 4/.....	2.5	2.2	0.4	*	*	0.3	0.7	0.9	0.7	0.9
2002.....	2.4	1.4	*	*	*	0.2	0.4	0.8	0.8	0.9
2001.....	2.2	2.2	*	*	*	0.4	0.5	0.6	0.6	0.8
2000.....	2.2	1.6	0.3	*	*	0.3	0.6	0.8	0.6	1.0
1999.....	2.1	1.9	*	*	*	0.3	0.3	0.6	0.5	0.9

White, female

2006 4/.....	5.6	1.3	*	*	*	*	0.4	0.4	0.3	0.6
2005 4/.....	4.6	*	*	*	*	*	0.3	0.4	0.4	0.5
2004 4/.....	4.4	1.5	*	*	*	*	0.3	0.4	0.4	0.4
2003 4/.....	4.2	1.3	*	*	*	*	0.4	0.3	0.5	0.5
2002.....	3.7	1.6	*	*	*	*	*	0.4	0.5	0.6
2001.....	3.4	*	*	*	*	*	0.4	0.3	0.5	0.5
2000.....	3.2	1.6	0.4	*	*	*	0.3	0.4	0.4	0.4
1999.....	3.0	*	*	*	*	*	*	0.3	0.5	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

Black, both sexes

2006 4/.....	4.2	*	*	*	*	*	1.0	1.2	1.2	1.3
2005 4/.....	3.4	3.3	*	*	*	*	0.9	1.4	1.1	1.7
2004 4/.....	3.5	3.2	*	*	*	*	1.0	1.4	1.6	1.4
2003 4/.....	3.4	*	*	*	*	*	0.9	1.5	1.2	1.7
2002.....	3.2	4.3	*	*	*	0.7	1.1	1.2	2.1	1.9
2001.....	3.1	*	*	*	*	*	1.3	1.3	0.9	1.9
2000.....	3.0	*	*	*	*	*	1.0	1.5	1.7	1.4
1999.....	3.0	3.6	*	*	*	*	1.0	1.4	1.4	2.0

Black, male

2006 4/.....	4.0	*	*	*	*	*	*	*	1.7	1.6
2005 4/.....	3.2	*	*	*	*	*	1.4	1.9	*	2.1
2004 4/.....	3.3	*	*	*	*	*	1.3	1.9	2.1	2.1
2003 4/.....	3.2	*	*	*	*	*	*	2.1	1.5	2.4
2002.....	3.3	*	*	*	*	*	*	1.7	2.8	2.3
2001.....	3.1	*	*	*	*	*	1.7	*	*	2.6
2000.....	3.0	*	*	*	*	*	*	2.0	1.9	1.8
1999.....	3.0	*	*	*	*	*	*	*	2.1	2.2

Black, female

2006 4/.....	4.5	*	*	*	*	*	*	*	*	*
2005 4/.....	3.5	*	*	*	*	*	*	*	*	*
2004 4/.....	3.7	*	*	*	*	*	*	*	*	*
2003 4/.....	3.6	*	*	*	*	*	*	*	*	*
2002.....	3.2	*	*	*	*	*	*	*	1.4	1.6
2001.....	3.0	*	*	*	*	*	*	*	*	1.3
2000.....	3.0	*	*	*	*	*	*	*	1.5	*
1999.....	3.0	*	*	*	*	*	*	1.4	*	1.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XX. External causes of mortality (*U01-*U03,V01-Y89)										
All races 3/, both sexes										
2006.....	60.7	38.7	12.6	6.2	8.3	50.4	76.4	67.2	59.8	62.1
2005.....	59.5	36.8	13.1	6.3	9.0	49.9	73.9	63.6	59.3	59.6
2004.....	57.9	36.7	13.0	6.5	9.8	51.4	70.7	62.5	55.5	59.5
2003.....	57.4	34.4	13.6	6.3	9.5	50.7	72.1	60.3	56.0	60.3
2002.....	56.9	33.8	13.7	6.8	9.9	52.6	71.9	60.6	55.5	61.1
2001 5/.....	56.2	35.1	14.3	7.2	9.9	51.0	70.6	60.9	55.6	62.4
2000.....	53.8	34.5	14.6	7.7	10.6	51.7	68.9	56.5	51.9	57.4
1999.....	54.2	33.9	15.3	8.1	10.7	52.7	67.2	56.4	54.0	57.6
All races 3/, male										
2006.....	84.6	44.3	14.1	7.1	10.6	74.4	120.9	104.9	91.6	90.7
2005.....	83.0	39.7	15.6	7.2	11.6	73.2	116.5	99.8	89.7	86.8
2004.....	80.3	40.4	14.7	7.6	12.3	74.0	111.2	98.0	84.0	87.2
2003.....	80.2	39.2	15.9	7.3	12.7	73.8	113.1	94.6	85.5	88.7
2002.....	79.8	37.6	16.3	7.9	12.7	76.5	114.0	95.6	84.8	90.3
2001 5/.....	79.3	38.9	16.8	8.5	12.9	75.1	113.0	96.1	85.3	92.4
2000.....	75.8	38.1	17.0	8.8	14.0	75.5	109.5	89.6	79.6	86.0
1999.....	76.2	38.3	17.3	9.5	14.1	76.3	106.5	88.9	82.6	86.0
All races 3/, female										
2006.....	37.4	32.8	11.0	5.3	6.0	25.2	28.9	27.8	27.1	33.3
2005.....	36.7	33.8	10.5	5.4	6.3	25.3	28.4	25.7	28.1	32.1
2004.....	36.2	32.9	11.3	5.4	7.2	27.6	27.7	25.5	26.4	31.5
2003.....	35.3	29.3	11.2	5.3	6.1	26.3	28.6	24.7	25.9	31.9
2002.....	34.8	29.8	11.1	5.7	6.9	27.5	27.6	24.4	25.7	31.8
2001 5/.....	34.0	31.2	11.8	5.8	6.7	25.4	26.2	24.7	25.4	32.5
2000.....	32.5	30.8	12.1	6.4	7.1	26.5	26.5	22.6	23.8	29.0
1999.....	33.0	29.2	13.3	6.6	7.2	27.8	26.2	23.3	25.2	29.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

XX. External causes of mortality (*U01-*U03,V01-Y89)

White, both sexes

2006 4/.....	61.7	32.2	11.1	5.5	8.0	48.2	73.1	64.4	58.4	62.8
2005 4/.....	60.4	31.5	12.2	5.7	8.4	48.4	70.0	60.5	57.7	59.9
2004 4/.....	59.0	31.3	11.9	5.8	9.3	50.9	67.7	59.0	54.1	59.9
2003 4/.....	58.2	28.9	12.5	5.7	8.9	49.9	67.7	56.4	54.7	60.5
2002.....	57.4	26.1	12.3	5.9	9.3	52.1	67.3	56.1	53.8	60.3
2001 5/.....	56.5	28.8	13.3	6.5	9.3	49.5	65.6	57.1	53.9	62.0
2000.....	53.6	27.6	13.1	7.0	10.3	50.4	63.2	51.7	49.6	56.6
1999.....	53.7	27.9	13.6	7.1	10.5	50.6	61.4	52.1	51.7	56.4

White, male

2006 4/.....	84.2	36.2	12.4	6.1	10.0	68.7	113.7	98.0	87.0	89.7
2005 4/.....	82.5	33.6	14.7	6.5	10.7	69.2	108.5	92.5	84.9	85.9
2004 4/.....	80.3	35.3	13.7	6.7	11.7	71.4	104.7	90.1	80.0	87.0
2003 4/.....	79.7	32.8	14.7	6.5	11.9	70.6	104.0	86.0	82.1	87.5
2002.....	79.0	30.7	14.5	6.8	11.8	74.1	104.5	86.0	80.8	87.9
2001 5/.....	78.3	31.8	15.8	7.4	12.1	71.1	102.8	88.2	81.6	91.2
2000.....	74.1	31.8	15.4	8.1	13.4	71.7	98.4	80.1	75.1	84.0
1999.....	74.2	32.6	15.4	8.2	13.6	71.1	95.2	80.8	77.8	83.3

White, female

2006 4/.....	39.7	28.0	9.7	4.8	5.9	26.5	29.3	28.4	28.3	34.8
2005 4/.....	38.7	29.3	9.5	4.8	6.0	26.4	28.5	26.2	29.1	33.0
2004 4/.....	38.1	27.0	10.2	4.9	6.7	29.2	27.9	25.7	26.9	32.0
2003 4/.....	37.1	24.8	10.3	4.9	5.8	27.9	28.6	24.7	26.1	32.8
2002.....	36.4	21.3	10.1	5.0	6.6	28.9	27.7	24.3	25.6	32.2
2001 5/.....	35.2	25.7	10.5	5.6	6.5	26.5	26.0	24.3	25.2	32.2
2000.....	33.6	23.2	10.7	5.9	7.0	27.7	25.7	22.0	23.2	28.7
1999.....	33.9	23.1	11.7	5.9	7.1	28.9	25.4	22.2	24.6	29.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
XX. External causes of mortality (*U01-*U03,V01-Y89)										
Black, both sexes										
2006 4/.....	66.1	73.1	20.9	10.0	10.6	64.2	104.0	98.6	85.5	76.7
2005 4/.....	65.4	67.6	18.7	9.2	12.0	61.5	103.9	93.4	85.7	73.5
2004 4/.....	62.3	66.5	19.4	10.4	12.7	58.8	97.9	97.1	78.2	71.1
2003 4/.....	62.8	63.5	20.2	9.2	12.3	58.5	105.7	96.5	78.6	72.6
2002.....	63.7	72.8	21.0	11.2	12.9	59.9	106.7	100.3	78.6	78.5
2001 5/.....	64.2	68.9	21.2	10.6	12.6	62.5	107.7	94.8	76.8	76.9
2000.....	63.9	70.6	22.8	11.3	13.0	63.3	111.4	94.0	76.1	73.4
1999.....	65.2	67.1	24.8	13.3	13.2	68.4	108.3	91.2	76.9	75.1
Black, male										
2006 4/.....	104.6	86.7	23.5	12.3	14.3	107.7	176.1	170.9	150.3	125.4
2005 4/.....	102.4	74.7	21.4	10.2	16.1	101.3	177.2	164.2	147.0	117.4
2004 4/.....	96.9	68.2	21.4	12.4	15.7	94.8	166.8	171.2	133.1	112.1
2003 4/.....	98.2	72.0	23.9	11.0	16.5	96.9	180.8	170.4	132.1	117.2
2002.....	99.9	73.5	26.0	13.1	17.9	96.4	184.7	178.2	132.9	126.6
2001 5/.....	100.2	77.2	24.1	14.2	17.7	102.8	189.7	165.4	128.5	117.9
2000.....	100.2	72.3	25.5	13.3	18.2	103.1	193.6	165.9	126.7	117.9
1999.....	101.7	71.2	27.7	16.7	18.0	110.9	189.6	158.4	127.6	119.4
Black, female										
2006 4/.....	31.0	58.9	18.1	7.6	6.8	19.5	30.0	30.0	26.9	33.3
2005 4/.....	31.5	60.1	15.9	8.2	7.7	20.7	29.3	26.5	30.2	34.0
2004 4/.....	30.8	64.8	17.4	8.3	9.5	21.8	28.6	27.8	28.7	34.5
2003 4/.....	30.5	54.7	16.4	7.4	7.9	18.9	31.0	28.2	30.3	32.9
2002.....	30.9	72.0	15.8	9.1	7.9	22.2	30.7	29.1	29.9	35.7
2001 5/.....	31.5	60.2	18.3	6.9	7.4	20.9	28.8	30.4	30.3	40.5
2000.....	30.9	68.8	19.9	9.2	7.6	22.6	33.5	29.0	30.5	34.0
1999.....	32.2	62.8	21.9	9.9	8.2	25.0	31.1	30.4	31.3	35.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All races 3/, both sexes										
2006.....	40.6	27.8	9.9	5.3	5.9	31.2	45.3	38.8	35.1	36.9
2005.....	39.7	26.4	10.3	5.5	6.4	31.4	43.4	36.0	33.8	35.3
2004.....	38.1	25.8	10.3	5.7	7.3	32.9	41.1	34.6	30.6	34.8
2003.....	37.6	23.6	10.9	5.5	7.2	33.0	41.1	32.8	30.2	35.0
2002.....	37.0	23.5	10.5	5.9	7.3	35.0	40.9	32.3	30.8	34.9
2001.....	35.7	24.2	11.2	6.4	7.4	32.8	39.5	31.4	28.5	34.1
2000.....	34.8	23.1	11.9	6.8	7.7	33.4	38.8	30.4	28.6	32.9
1999.....	35.1	22.3	12.4	7.1	8.1	33.3	37.5	30.0	29.2	32.2
All races 3/, male										
2006.....	53.5	31.6	11.1	6.1	7.3	43.0	69.1	59.2	52.1	52.7
2005.....	52.3	27.5	12.4	6.3	8.1	43.0	66.2	54.8	49.7	50.6
2004.....	49.8	29.0	11.8	6.8	9.0	44.6	61.8	52.2	45.0	49.7
2003.....	49.3	26.4	13.0	6.4	9.3	44.7	61.8	49.4	45.1	50.2
2002.....	48.9	26.8	12.9	7.1	9.2	47.6	62.4	49.5	46.1	50.5
2001.....	47.2	26.2	13.3	7.6	9.5	45.3	60.7	48.0	43.1	49.5
2000.....	46.2	25.7	14.0	7.8	9.9	45.8	58.9	47.0	43.1	48.9
1999.....	46.4	25.4	14.4	8.5	10.4	44.8	57.1	46.1	43.9	47.4
All races 3/, female										
2006.....	28.1	23.8	8.6	4.4	4.4	18.9	19.9	17.6	17.6	20.8
2005.....	27.5	25.2	8.0	4.7	4.7	19.2	19.2	16.3	17.5	19.8
2004.....	26.8	22.5	8.7	4.6	5.4	20.6	19.1	16.2	15.9	19.7
2003.....	26.2	20.7	8.7	4.6	4.9	20.6	19.2	15.5	15.1	19.7
2002.....	25.6	20.0	8.1	4.7	5.3	21.8	18.4	14.5	15.2	19.2
2001.....	24.5	22.1	8.9	5.0	5.3	19.5	17.2	14.3	13.8	18.7
2000.....	23.8	20.5	9.7	5.7	5.5	20.3	17.8	13.5	13.9	17.0
1999.....	24.1	19.0	10.3	5.6	5.6	21.1	17.1	13.7	14.3	17.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, both sexes

2006 4/.....	42.9	22.8	9.2	4.7	5.9	33.8	49.5	41.6	37.1	38.5
2005 4/.....	41.8	23.0	10.0	5.1	6.3	33.9	46.9	38.5	35.3	36.8
2004 4/.....	40.2	22.2	10.0	5.2	7.1	36.0	44.4	36.5	32.1	36.1
2003 4/.....	39.5	20.0	10.6	5.2	7.0	35.9	44.3	34.6	31.6	36.2
2002.....	38.7	18.6	9.9	5.2	7.0	37.9	43.8	33.5	32.2	35.5
2001.....	37.0	20.6	11.0	5.8	7.1	34.9	41.7	32.7	29.3	34.8
2000.....	35.9	19.1	11.1	6.2	7.6	36.0	40.7	30.9	29.1	33.4
1999.....	36.0	18.9	11.5	6.3	7.9	35.6	38.9	30.8	29.3	32.4

White, male

2006 4/.....	55.7	25.8	10.4	5.4	7.2	45.9	75.5	62.8	54.3	54.4
2005 4/.....	54.3	23.7	12.3	5.9	7.8	46.1	71.6	58.1	51.2	52.1
2004 4/.....	52.0	25.5	11.5	6.1	8.9	48.5	67.0	54.6	46.5	51.4
2003 4/.....	51.3	22.5	12.8	6.0	9.0	48.4	66.2	51.9	46.8	51.4
2002.....	50.4	22.3	11.8	6.3	8.8	51.0	66.4	50.7	47.9	50.9
2001.....	48.4	22.1	13.3	6.7	9.0	47.8	64.2	49.6	44.0	50.3
2000.....	47.0	21.7	13.2	7.2	9.6	48.8	61.7	47.2	43.7	49.5
1999.....	46.9	21.8	13.2	7.4	10.1	47.2	58.9	47.1	43.8	47.5

White, female

2006 4/.....	30.3	19.8	8.0	4.0	4.5	21.0	21.4	18.9	19.0	22.1
2005 4/.....	29.5	22.2	7.6	4.3	4.7	21.0	20.3	17.4	18.6	20.9
2004 4/.....	28.8	18.8	8.4	4.3	5.2	22.9	20.1	17.2	16.9	20.3
2003 4/.....	28.0	17.4	8.3	4.4	4.8	22.7	20.7	16.1	15.7	20.6
2002.....	27.3	14.7	7.8	4.2	5.2	23.9	19.6	15.1	15.9	19.7
2001.....	25.9	18.9	8.5	4.8	5.1	21.1	17.8	14.8	14.0	19.0
2000.....	25.1	16.4	9.0	5.3	5.5	22.3	18.3	13.8	13.8	17.0
1999.....	25.3	15.8	9.6	5.1	5.6	23.3	17.6	13.8	14.3	17.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
Black, both sexes										
2006 4/.....	35.2	52.7	14.0	8.1	6.3	21.8	31.1	33.4	35.0	37.9
2005 4/.....	34.9	47.4	12.5	7.5	7.6	22.8	31.8	29.4	34.9	36.1
2004 4/.....	32.8	45.8	12.5	8.5	8.5	21.6	30.1	32.2	30.6	34.9
2003 4/.....	32.4	43.5	13.4	7.3	8.1	20.7	30.6	29.8	29.9	35.5
2002.....	33.1	48.9	14.1	9.3	9.0	23.9	32.0	32.8	30.1	38.1
2001.....	33.5	43.1	13.6	9.4	9.4	24.7	33.0	30.5	30.0	35.8
2000.....	33.5	44.5	16.2	9.7	8.8	23.7	34.3	33.1	31.9	35.9
1999.....	35.2	41.6	17.7	11.3	9.6	25.0	34.6	31.5	32.9	36.6
Black, male										
2006 4/.....	50.8	61.3	15.9	9.7	8.0	33.1	46.2	53.4	57.2	58.8
2005 4/.....	50.0	50.8	13.8	8.3	10.3	32.5	48.5	47.2	54.8	55.7
2004 4/.....	46.7	48.3	14.2	10.4	10.1	31.2	44.8	51.7	48.9	51.5
2003 4/.....	46.1	47.6	15.7	8.7	10.6	30.0	47.6	46.6	46.9	54.9
2002.....	47.9	51.1	18.5	11.1	12.1	34.6	49.5	53.6	47.7	58.9
2001.....	48.2	48.5	15.4	12.6	12.9	36.1	51.0	48.3	46.6	53.3
2000.....	49.0	47.7	18.8	11.9	12.1	34.5	52.9	54.1	49.6	55.7
1999.....	51.0	45.9	21.0	14.5	13.0	36.6	54.7	51.0	51.7	55.4
Black, female										
2006 4/.....	20.9	43.6	12.1	6.3	4.6	10.1	15.6	14.4	14.9	19.2
2005 4/.....	21.2	43.8	11.1	6.7	4.8	12.7	14.9	12.6	16.8	18.5
2004 4/.....	20.1	43.2	10.6	6.5	6.7	11.8	15.4	14.1	14.1	20.0
2003 4/.....	19.9	39.3	11.0	5.9	5.5	11.2	13.7	14.3	14.7	18.3
2002.....	19.7	46.6	9.6	7.5	5.8	12.8	14.9	13.8	14.4	19.7
2001.....	20.1	37.5	11.9	6.2	5.8	12.9	15.6	14.2	15.1	20.2
2000.....	19.5	41.3	13.5	7.6	5.4	12.7	16.7	14.1	15.9	18.3
1999.....	20.8	37.3	14.3	8.0	6.1	13.1	15.6	13.8	16.0	19.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
All races 3/, both sexes										
2006.....	16.2	3.4	3.8	3.1	4.1	23.9	29.8	20.9	17.4	16.4
2005.....	16.3	3.6	4.0	3.3	4.3	24.4	29.3	20.5	17.6	16.3
2004.....	16.4	3.6	4.1	3.5	5.1	26.0	28.5	20.9	16.7	16.4
2003.....	16.5	3.7	4.1	3.5	5.1	26.7	28.8	20.1	16.5	16.7
2002.....	16.8	3.1	4.1	3.5	4.9	28.4	29.9	20.5	17.1	17.1
2001.....	16.6	3.7	4.4	3.8	5.0	26.6	29.2	20.9	16.9	17.4
2000.....	16.6	4.6	4.5	4.1	5.3	27.1	29.3	20.4	16.8	17.1
1999.....	16.6	5.0	4.6	4.4	5.3	27.1	28.2	20.0	17.0	16.9
All races 3/, male										
2006.....	23.1	3.3	4.0	3.5	4.7	31.2	44.6	31.8	25.9	23.7
2005.....	23.3	3.5	4.5	3.7	5.1	31.8	44.0	31.2	26.0	23.5
2004.....	23.0	3.9	4.4	3.9	6.0	33.6	42.0	31.2	24.4	24.0
2003.....	23.3	4.0	4.5	4.0	6.3	34.7	42.4	29.8	24.8	24.1
2002.....	23.7	3.5	4.8	4.0	5.9	37.0	44.4	31.0	25.4	24.7
2001.....	23.5	3.3	5.0	4.3	6.2	35.4	43.9	31.4	25.4	24.7
2000.....	23.3	4.8	5.0	4.4	6.4	35.5	43.3	30.8	24.7	24.8
1999.....	23.1	5.2	5.1	5.0	6.5	34.8	41.8	29.9	24.7	24.0
All races 3/, female										
2006.....	9.4	3.6	3.6	2.8	3.3	16.2	14.0	9.5	8.6	9.1
2005.....	9.5	3.6	3.5	3.0	3.5	16.5	13.6	9.3	8.9	8.9
2004.....	9.9	3.2	3.8	3.0	4.1	18.0	14.2	10.2	8.9	8.8
2003.....	10.0	3.4	3.7	3.0	3.8	18.3	14.3	10.0	8.1	9.3
2002.....	10.1	2.8	3.4	3.1	3.9	19.3	14.6	9.7	8.7	9.6
2001.....	10.0	4.2	3.8	3.3	3.8	17.3	13.8	10.1	8.3	10.2
2000.....	10.2	4.3	4.0	3.8	4.1	18.1	14.6	9.8	8.7	9.5
1999.....	10.4	4.8	4.2	3.6	4.1	19.0	14.0	10.0	9.3	9.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
White, both sexes										
2006 4/.....	16.7	3.1	3.5	3.0	4.2	26.0	31.9	21.4	17.6	16.8
2005 4/.....	16.9	3.1	3.9	3.3	4.3	26.4	31.1	21.2	17.9	16.6
2004 4/.....	16.9	3.3	4.0	3.4	5.2	28.6	30.3	21.4	17.0	16.7
2003 4/.....	17.0	3.6	4.0	3.5	5.2	29.2	30.4	20.7	16.8	17.0
2002.....	17.3	2.7	3.9	3.4	5.0	31.0	31.5	20.7	17.6	17.2
2001.....	17.0	3.7	4.3	3.6	5.0	28.6	30.4	21.6	17.1	17.6
2000.....	17.0	4.1	4.3	4.1	5.4	29.6	30.5	20.3	16.8	17.3
1999.....	17.0	4.6	4.1	4.1	5.5	29.4	29.2	20.4	17.2	16.8
White, male										
2006 4/.....	23.7	3.2	3.6	3.2	4.8	33.7	47.7	32.4	25.8	23.9
2005 4/.....	24.0	3.3	4.3	3.7	5.1	34.2	46.7	32.0	26.0	23.5
2004 4/.....	23.7	3.3	4.2	3.8	6.3	36.7	44.8	31.5	24.3	24.3
2003 4/.....	23.9	4.0	4.5	3.9	6.5	37.6	44.4	30.6	25.1	24.2
2002.....	24.2	3.1	4.5	3.8	6.0	40.2	46.7	31.0	25.9	24.5
2001.....	23.9	3.2	4.9	4.0	6.2	37.8	45.8	32.3	25.4	24.7
2000.....	23.6	4.5	4.7	4.4	6.5	38.4	45.0	30.3	24.6	24.8
1999.....	23.4	5.2	4.4	4.7	6.7	37.3	43.1	30.2	24.6	23.7
White, female										
2006 4/.....	9.8	3.1	3.4	2.7	3.5	17.9	14.8	9.6	9.1	9.5
2005 4/.....	10.0	2.9	3.5	3.0	3.6	18.2	14.2	9.7	9.5	9.4
2004 4/.....	10.3	3.2	3.8	2.9	4.1	20.0	14.7	10.5	9.3	8.9
2003 4/.....	10.3	3.1	3.4	3.1	3.9	20.2	15.3	10.2	8.2	9.6
2002.....	10.5	2.2	3.4	2.9	3.9	21.3	15.3	9.8	9.0	9.7
2001.....	10.3	4.1	3.6	3.1	3.8	18.9	14.0	10.4	8.4	10.4
2000.....	10.5	3.7	3.8	3.7	4.3	20.2	15.1	9.9	8.7	9.8
1999.....	10.7	3.9	3.8	3.5	4.3	21.0	14.4	10.1	9.6	9.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Transport accidents (V01-V99,Y85)										
Black, both sexes										
2006 4/.....	15.1	4.6	5.2	4.0	3.8	15.7	23.0	21.7	21.0	17.7
2005 4/.....	15.0	5.7	4.5	3.6	4.3	17.1	23.6	19.2	20.0	17.9
2004 4/.....	15.2	4.8	5.1	4.0	4.8	16.6	22.7	22.4	19.3	17.6
2003 4/.....	15.1	4.2	5.0	3.6	4.5	16.2	23.2	20.3	18.3	17.9
2002.....	15.4	5.3	5.0	4.6	4.6	17.9	24.6	23.0	18.0	19.2
2001.....	15.9	*	5.2	5.3	5.3	18.8	25.8	21.0	18.9	18.1
2000.....	16.0	6.5	5.6	4.5	4.8	17.3	26.6	23.5	19.4	17.9
1999.....	16.3	7.3	7.1	5.8	5.0	18.7	25.8	21.3	18.5	19.9
Black, male										
2006 4/.....	22.9	*	5.7	4.8	4.6	22.5	33.9	34.4	35.2	28.9
2005 4/.....	22.7	*	4.9	3.6	5.5	23.5	35.8	31.2	33.6	29.4
2004 4/.....	22.8	5.9	5.4	4.4	5.3	22.9	33.2	36.1	31.5	27.9
2003 4/.....	22.6	*	5.1	4.5	5.7	22.7	36.3	31.8	29.4	28.4
2002.....	23.1	*	6.4	5.0	5.6	24.5	37.1	37.5	28.9	29.8
2001.....	23.8	*	5.7	6.6	7.3	26.5	39.1	33.2	30.0	27.2
2000.....	24.1	6.7	6.2	4.9	6.5	24.2	40.3	38.2	30.7	28.6
1999.....	24.2	*	8.0	7.2	6.4	25.8	39.2	34.3	29.4	29.7
Black, female										
2006 4/.....	8.0	*	4.6	3.1	3.0	8.7	11.8	9.7	8.2	7.7
2005 4/.....	7.9	6.8	4.2	3.5	3.0	10.6	11.2	7.9	7.8	7.7
2004 4/.....	8.2	*	4.8	3.6	4.2	10.0	12.1	9.6	8.3	8.5
2003 4/.....	8.3	*	5.0	2.8	3.4	9.5	10.2	9.8	8.2	8.4
2002.....	8.3	*	3.4	4.2	3.5	11.2	12.5	9.8	8.2	9.8
2001.....	8.7	*	4.7	4.0	3.4	10.8	13.0	9.9	8.9	10.0
2000.....	8.5	*	5.0	4.1	3.0	10.3	13.7	10.2	9.2	8.4
1999.....	9.2	8.7	6.2	4.4	3.6	11.4	13.1	9.5	8.6	11.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
All races 3/, both sexes										
2006.....	15.1	3.4	3.6	2.9	3.7	23.2	28.8	19.9	16.4	15.2
2005.....	15.3	3.6	3.8	3.2	4.0	23.6	28.2	19.6	16.5	15.1
2004.....	15.3	3.5	4.0	3.3	4.8	25.2	27.5	19.7	15.6	15.2
2003.....	15.4	3.6	3.9	3.3	4.7	25.7	27.5	19.0	15.4	15.4
2002.....	15.7	3.0	3.9	3.3	4.5	27.6	28.8	19.7	16.0	15.8
2001.....	15.4	3.6	4.1	3.6	4.6	25.7	28.0	19.6	15.6	15.9
2000.....	15.4	4.4	4.2	3.8	4.8	26.0	28.0	19.2	15.5	15.6
1999.....	15.2	4.8	4.2	3.9	4.8	25.9	26.5	18.4	15.4	15.1
All races 3/, male										
2006.....	21.4	3.3	3.8	3.2	4.2	30.2	42.9	30.1	24.4	21.7
2005.....	21.7	3.5	4.2	3.5	4.7	30.7	42.3	29.6	24.2	21.8
2004.....	21.3	3.8	4.2	3.7	5.6	32.5	40.4	29.3	22.4	21.9
2003.....	21.4	3.9	4.4	3.7	5.9	33.3	40.4	28.2	23.0	21.9
2002.....	21.9	3.3	4.5	3.8	5.3	36.0	42.6	29.6	23.6	22.4
2001.....	21.5	3.2	4.5	4.0	5.7	34.0	42.1	29.5	23.1	22.3
2000.....	21.3	4.6	4.7	4.1	5.8	33.9	41.2	28.6	22.6	22.1
1999.....	20.9	5.0	4.7	4.4	5.7	33.1	39.1	27.2	22.2	21.2
All races 3/, female										
2006.....	9.0	3.5	3.4	2.6	3.2	15.7	13.7	9.2	8.1	8.7
2005.....	9.1	3.6	3.4	2.9	3.2	16.2	13.2	9.1	8.6	8.4
2004.....	9.5	3.2	3.7	2.8	3.9	17.5	13.8	9.7	8.5	8.3
2003.....	9.5	3.3	3.5	2.8	3.5	17.8	13.9	9.5	7.7	8.8
2002.....	9.8	2.8	3.3	2.9	3.6	18.9	14.3	9.4	8.3	9.2
2001.....	9.5	4.0	3.6	3.0	3.5	16.9	13.4	9.5	8.0	9.6
2000.....	9.7	4.2	3.8	3.5	3.8	17.5	14.2	9.5	8.2	9.1
1999.....	9.7	4.7	3.8	3.3	3.8	18.3	13.4	9.4	8.5	9.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, both sexes

2006 4/.....	15.6	3.1	3.4	2.7	3.8	25.2	30.7	20.4	16.6	15.6
2005 4/.....	15.8	3.1	3.7	3.2	4.0	25.6	30.0	20.3	16.8	15.4
2004 4/.....	15.8	3.2	3.9	3.2	4.9	27.8	29.2	20.0	15.7	15.3
2003 4/.....	15.8	3.5	3.8	3.3	4.8	28.1	29.1	19.5	15.7	15.6
2002.....	16.2	2.6	3.7	3.2	4.6	30.2	30.4	19.8	16.4	15.8
2001.....	15.7	3.5	3.9	3.3	4.6	27.6	29.1	20.2	15.7	16.1
2000.....	15.7	3.9	4.0	3.7	5.0	28.4	29.1	19.0	15.4	15.8
1999.....	15.5	4.5	3.8	3.7	5.0	28.1	27.4	18.6	15.5	14.9

White, male

2006 4/.....	22.0	3.2	3.5	3.0	4.2	32.6	45.8	30.7	24.2	21.9
2005 4/.....	22.3	3.3	4.1	3.5	4.6	33.0	45.1	30.4	24.1	21.7
2004 4/.....	21.9	3.3	4.0	3.6	5.9	35.6	43.2	29.5	22.2	22.1
2003 4/.....	22.0	3.9	4.3	3.6	6.0	36.1	42.3	28.8	23.3	21.9
2002.....	22.4	2.9	4.2	3.6	5.5	39.0	44.8	29.5	23.9	22.2
2001.....	21.9	3.1	4.4	3.7	5.6	36.2	43.8	30.1	23.1	22.3
2000.....	21.6	4.2	4.5	4.0	5.9	36.8	42.6	27.9	22.4	22.1
1999.....	21.1	5.0	4.1	4.2	5.9	35.5	40.4	27.4	22.1	20.8

White, female

2006 4/.....	9.4	3.0	3.3	2.5	3.3	17.4	14.3	9.3	8.5	9.1
2005 4/.....	9.5	2.9	3.4	2.8	3.2	17.7	13.8	9.4	9.1	8.8
2004 4/.....	9.8	3.2	3.7	2.7	3.9	19.5	14.1	9.9	8.9	8.3
2003 4/.....	9.8	3.0	3.2	2.9	3.6	19.6	14.8	9.6	7.7	9.0
2002.....	10.1	2.2	3.2	2.7	3.7	20.9	15.0	9.5	8.6	9.3
2001.....	9.8	3.8	3.4	2.9	3.5	18.5	13.5	9.7	8.0	9.8
2000.....	10.0	3.5	3.6	3.4	4.0	19.5	14.6	9.6	8.2	9.3
1999.....	10.0	3.9	3.5	3.2	4.0	20.2	13.7	9.5	8.7	8.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2006 4/.....	14.3	4.4	5.0	3.8	3.6	15.3	22.5	20.6	20.1	16.5
2005 4/.....	14.1	5.7	4.3	3.4	4.0	16.7	22.4	18.4	19.0	17.0
2004 4/.....	14.3	4.7	4.9	3.8	4.5	15.8	21.8	21.2	18.4	16.6
2003 4/.....	14.2	4.2	4.9	3.4	4.3	15.5	22.3	19.5	17.3	16.7
2002.....	14.5	5.2	4.8	4.3	4.2	17.4	24.0	22.4	17.2	17.9
2001.....	14.8	*	5.0	4.9	4.8	18.1	25.2	20.1	17.7	16.7
2000.....	15.0	6.5	5.4	4.3	4.6	16.5	25.8	22.8	18.0	16.4
1999.....	15.1	7.1	6.4	5.3	4.6	17.8	24.8	20.1	17.0	18.2

Black, male

2006 4/.....	21.5	*	5.7	4.6	4.3	21.8	33.1	32.4	33.6	26.6
2005 4/.....	21.2	*	4.7	3.4	5.1	22.7	33.8	29.9	31.9	27.6
2004 4/.....	21.2	*	5.2	4.2	5.0	21.6	31.5	33.8	29.8	26.0
2003 4/.....	21.1	*	5.0	4.1	5.2	21.6	34.6	30.5	27.7	26.3
2002.....	21.5	*	6.3	4.8	5.0	23.7	36.1	36.2	27.4	27.6
2001.....	22.0	*	5.3	6.2	6.5	25.5	38.0	31.9	28.1	24.9
2000.....	22.5	6.7	5.8	4.5	6.2	22.8	38.8	36.7	28.5	26.0
1999.....	22.2	*	7.0	6.3	5.7	24.5	37.7	32.4	27.0	27.0

Black, female

2006 4/.....	7.7	*	4.4	3.1	2.9	8.6	11.6	9.4	7.9	7.5
2005 4/.....	7.5	6.8	3.9	3.4	3.0	10.6	10.8	7.6	7.5	7.5
2004 4/.....	7.9	*	4.7	3.5	4.0	9.8	12.0	9.4	8.1	8.2
2003 4/.....	8.0	*	4.8	2.7	3.4	9.2	10.1	9.4	8.0	8.2
2002.....	8.0	*	3.4	3.8	3.3	11.0	12.2	9.7	7.9	9.3
2001.....	8.2	*	4.6	3.6	3.1	10.5	12.9	9.5	8.4	9.5
2000.....	8.2	*	4.9	4.0	2.9	10.1	13.5	10.2	8.6	7.9
1999.....	8.7	8.7	5.8	4.2	3.5	11.0	12.6	9.0	8.1	10.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pedestrian involved in collision with motor vehicle (V02-V04)										
All races 3/, both sexes										
2006.....	1.1	*	0.9	0.4	0.4	0.8	1.0	1.0	1.0	1.0
2005.....	1.1	*	1.1	0.5	0.4	0.8	1.1	0.9	1.0	1.1
2004.....	1.1	*	1.0	0.5	0.5	0.8	1.0	1.0	0.9	1.1
2003.....	1.2	*	1.1	0.5	0.5	1.0	1.1	1.1	0.9	1.2
2002.....	1.3	*	1.0	0.6	0.5	1.0	1.1	1.0	1.0	1.3
2001.....	1.2	*	1.1	0.6	0.7	0.9	1.1	1.0	1.0	1.4
2000.....	1.2	*	1.2	0.7	0.6	0.8	1.0	1.0	1.0	1.2
1999.....	1.2	*	1.1	0.8	0.5	0.8	1.1	1.0	1.0	1.3
All races 3/, male										
2006.....	1.6	*	1.2	0.6	0.6	1.1	1.3	1.7	1.4	1.5
2005.....	1.6	*	1.4	0.5	0.6	0.9	1.6	1.3	1.5	1.6
2004.....	1.6	*	1.2	0.6	0.6	1.1	1.5	1.4	1.4	1.6
2003.....	1.7	*	1.4	0.7	0.7	1.3	1.6	1.6	1.4	1.7
2002.....	1.7	*	1.3	0.7	0.6	1.4	1.7	1.6	1.6	1.8
2001.....	1.8	*	1.4	0.8	1.0	1.2	1.8	1.6	1.5	1.9
2000.....	1.7	*	1.4	0.8	0.7	1.0	1.5	1.5	1.5	1.7
1999.....	1.7	*	1.2	0.9	0.7	1.2	1.6	1.5	1.5	1.9
All races 3/, female										
2006.....	0.7	*	0.7	*	0.3	0.6	0.6	0.4	0.5	0.4
2005.....	0.7	*	0.9	0.4	0.3	0.6	0.4	0.5	0.6	0.6
2004.....	0.7	*	0.8	0.4	0.3	0.4	0.5	0.5	0.5	0.7
2003.....	0.7	*	0.8	0.4	0.4	0.6	0.6	0.5	0.4	0.6
2002.....	0.8	*	0.6	0.5	0.4	0.6	0.5	0.5	0.5	0.8
2001.....	0.8	*	0.8	0.4	0.4	0.5	0.5	0.3	0.5	0.8
2000.....	0.8	*	0.9	0.6	0.4	0.6	0.4	0.4	0.5	0.7
1999.....	0.7	*	1.0	0.7	0.4	0.5	0.5	0.5	0.4	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pedestrian involved in collision with motor vehicle (V02-V04)										
White, both sexes										
2006 4/.....	1.1	*	0.9	0.3	0.4	0.9	1.0	1.0	1.0	0.9
2005 4/.....	1.1	*	1.2	0.4	0.4	0.7	1.1	0.9	1.0	1.0
2004 4/.....	1.1	*	1.0	0.5	0.5	0.8	1.0	1.0	0.8	1.1
2003 4/.....	1.2	*	1.1	0.4	0.5	1.0	1.1	1.0	0.9	1.1
2002.....	1.2	*	0.9	0.5	0.5	0.9	1.1	1.0	1.0	1.1
2001.....	1.2	*	1.0	0.5	0.7	0.8	1.2	1.0	1.0	1.2
2000.....	1.1	*	1.0	0.6	0.6	0.8	1.0	0.9	0.9	1.1
1999.....	1.1	*	1.0	0.6	0.5	0.8	1.0	0.9	0.9	1.1
White, male										
2006 4/.....	1.5	*	1.1	0.4	0.5	1.1	1.3	1.7	1.5	1.4
2005 4/.....	1.5	*	1.3	0.4	0.4	0.8	1.7	1.3	1.3	1.5
2004 4/.....	1.5	*	1.2	0.5	0.6	1.2	1.5	1.5	1.3	1.5
2003 4/.....	1.6	*	1.4	0.6	0.6	1.4	1.7	1.7	1.3	1.6
2002.....	1.6	*	1.2	0.7	0.6	1.3	1.8	1.5	1.5	1.5
2001.....	1.7	*	1.4	0.6	1.0	1.2	1.8	1.7	1.4	1.8
2000.....	1.6	*	1.2	0.7	0.7	1.0	1.6	1.3	1.4	1.5
1999.....	1.5	*	1.1	0.7	0.7	1.2	1.6	1.3	1.4	1.6
White, female										
2006 4/.....	0.6	*	0.6	*	*	0.6	0.6	0.3	0.5	0.4
2005 4/.....	0.7	*	1.0	0.4	0.3	0.6	0.4	0.5	0.6	0.5
2004 4/.....	0.7	*	0.8	0.4	0.3	0.4	0.5	0.4	0.4	0.7
2003 4/.....	0.7	*	0.8	0.3	0.3	0.6	0.5	0.4	0.5	0.6
2002.....	0.7	*	0.6	0.4	0.4	0.5	0.5	0.4	0.5	0.6
2001.....	0.7	*	0.7	0.3	0.4	0.5	0.5	0.3	0.5	0.7
2000.....	0.7	*	0.8	0.5	0.4	0.6	0.5	0.4	0.4	0.6
1999.....	0.7	*	1.0	0.5	0.4	0.5	0.5	0.4	0.4	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pedestrian involved in collision with motor vehicle (V02-V04)										
Black, both sexes										
2006 4/.....	1.4	*	1.5	0.9	0.6	0.7	0.9	1.2	1.1	1.2
2005 4/.....	1.4	*	1.0	0.7	0.9	0.9	1.0	1.1	1.4	1.6
2004 4/.....	1.4	*	1.1	0.9	*	*	1.1	1.0	1.2	1.3
2003 4/.....	1.5	*	1.5	1.0	0.8	1.0	1.0	1.1	1.3	1.4
2002.....	1.6	*	1.3	1.0	0.6	1.3	1.0	1.4	1.4	2.4
2001.....	1.7	*	1.5	1.2	0.9	1.1	1.0	1.0	1.2	2.2
2000.....	1.7	*	1.9	1.2	0.7	0.8	0.8	1.3	1.1	2.1
1999.....	1.8	*	1.6	1.6	0.7	0.8	1.1	1.7	1.5	2.4
Black, male										
2006 4/.....	2.1	*	1.9	1.6	*	*	1.3	1.8	1.8	1.7
2005 4/.....	2.1	*	*	*	1.2	*	1.4	1.6	2.4	2.4
2004 4/.....	2.1	*	*	*	*	*	1.6	*	2.1	1.9
2003 4/.....	2.3	*	1.9	1.3	*	1.2	1.4	*	2.3	2.5
2002.....	2.4	*	1.7	*	*	1.7	1.5	2.1	2.2	3.6
2001.....	2.4	*	1.7	1.8	1.4	1.3	1.6	1.6	1.7	2.8
2000.....	2.5	*	2.3	1.5	*	*	*	2.1	1.6	3.4
1999.....	2.8	*	2.0	1.8	*	*	1.8	2.6	2.7	3.9
Black, female										
2006 4/.....	0.9	*	*	*	*	*	*	*	*	*
2005 4/.....	0.7	*	*	*	*	*	*	*	*	*
2004 4/.....	0.8	*	*	*	*	*	*	*	*	*
2003 4/.....	0.8	*	*	*	*	*	*	*	*	*
2002.....	0.9	*	*	*	*	*	*	*	*	1.4
2001.....	1.0	*	*	*	*	*	*	*	*	1.7
2000.....	0.8	*	*	*	*	*	*	*	*	*
1999.....	0.9	*	*	1.3	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pedalcyclist involved in collision with motor vehicle (V12-V14,V19.0-V19.2,V19.4-V19.6)										
All races 3/, both sexes										
2006.....	0.2	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.2
2005.....	0.2	*	*	0.2	0.3	0.2	0.2	0.2	0.2	0.3
2004.....	0.2	*	*	0.2	0.3	0.3	0.2	0.1	0.1	0.3
2003.....	0.2	*	*	0.2	0.4	0.3	0.1	0.1	0.1	0.2
2002.....	0.2	*	*	0.2	0.3	0.2	0.1	0.1	0.2	0.2
2001.....	0.2	*	*	0.2	0.3	0.3	0.2	0.2	0.2	0.2
2000.....	0.2	*	*	0.2	0.5	0.2	0.1	0.1	0.2	0.2
1999.....	0.2	*	*	0.3	0.4	0.3	0.2	0.2	0.2	0.2
All races 3/, male										
2006.....	0.4	*	*	0.3	0.4	0.4	0.3	0.3	0.3	0.3
2005.....	0.4	*	*	0.3	0.5	0.4	0.4	0.3	0.3	0.5
2004.....	0.4	*	*	0.3	0.5	0.4	0.3	0.3	0.2	0.5
2003.....	0.4	*	*	0.2	0.6	0.4	0.3	0.2	*	0.4
2002.....	0.4	*	*	0.3	0.6	0.4	0.2	0.2	0.4	0.3
2001.....	0.4	*	*	0.3	0.5	0.5	0.3	0.4	0.3	0.4
2000.....	0.4	*	*	0.4	0.8	0.4	0.3	0.2	0.3	0.3
1999.....	0.4	*	*	0.5	0.6	0.6	0.4	0.3	0.4	0.4
All races 3/, female										
2006.....	0.1	*	*	*	*	*	*	*	*	*
2005.....	0.1	*	*	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	0.2	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

All races 3/, both sexes

2006.....	1.6	*	*	*	0.1	1.1	3.5	2.6	2.4	2.4
2005.....	1.5	*	*	*	0.2	1.1	3.1	2.5	2.4	2.2
2004.....	1.4	*	*	*	0.2	1.1	2.7	2.3	2.1	2.0
2003.....	1.3	*	*	*	0.2	1.0	2.5	2.1	2.0	1.9
2002.....	1.1	*	*	*	0.1	0.8	2.3	2.0	1.8	1.7
2001.....	1.1	*	*	*	0.1	0.8	2.3	2.2	1.9	1.7
2000.....	1.0	*	*	*	0.1	0.8	2.1	2.1	1.6	1.5
1999.....	0.8	*	*	*	0.1	0.6	2.0	1.8	1.3	1.3

All races 3/, male

2006.....	3.0	*	*	*	0.2	1.7	6.5	4.8	4.4	4.3
2005.....	2.7	*	*	*	0.3	1.9	5.6	4.6	4.3	3.8
2004.....	2.5	*	*	*	0.4	2.0	5.0	4.2	3.9	3.5
2003.....	2.3	*	*	*	0.3	1.7	4.4	3.9	3.5	3.4
2002.....	2.1	*	*	*	*	1.5	4.1	3.7	3.3	3.0
2001.....	2.0	*	*	*	0.3	1.5	4.2	4.1	3.4	2.9
2000.....	1.8	*	*	*	0.2	1.5	3.8	3.9	2.9	2.7
1999.....	1.6	*	*	*	0.2	1.1	3.6	3.3	2.4	2.4

All races 3/, female

2006.....	0.3	*	*	*	*	0.3	0.4	0.2	0.4	0.5
2005.....	0.3	*	*	*	*	0.2	0.4	0.3	0.5	0.5
2004.....	0.3	*	*	*	*	0.2	0.4	0.3	0.3	0.4
2003.....	0.3	*	*	*	*	0.2	0.4	0.3	0.4	0.4
2002.....	0.2	*	*	*	*	*	0.3	0.3	0.3	0.5
2001.....	0.2	*	*	*	*	*	0.3	0.3	0.4	0.5
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.4
1999.....	0.2	*	*	*	*	*	0.2	0.2	*	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Motorcyclist involved in any accident except collision with railway train (V20-V24,V26-V34,V36-V39)

White, both sexes

2006 4/.....	1.7	*	*	*	0.1	1.2	3.8	2.6	2.4	2.5
2005 4/.....	1.6	*	*	*	0.2	1.2	3.4	2.5	2.3	2.2
2004 4/.....	1.5	*	*	*	0.2	1.2	3.0	2.3	2.1	2.0
2003 4/.....	1.4	*	*	*	0.2	1.1	2.7	2.1	1.9	1.9
2002.....	1.2	*	*	*	*	0.9	2.3	2.0	1.8	1.9
2001.....	1.1	*	*	*	0.2	1.0	2.4	2.2	1.8	1.8
2000.....	1.0	*	*	*	0.1	0.9	2.1	2.0	1.5	1.6
1999.....	0.9	*	*	*	0.2	0.7	2.0	1.6	1.3	1.4

White, male

2006 4/.....	3.2	*	*	*	*	2.0	7.1	4.8	4.3	4.3
2005 4/.....	2.9	*	*	*	0.3	2.2	6.1	4.5	4.0	3.7
2004 4/.....	2.7	*	*	*	0.4	2.2	5.4	4.1	3.7	3.5
2003 4/.....	2.4	*	*	*	0.3	2.0	4.8	3.7	3.3	3.2
2002.....	2.2	*	*	*	*	1.7	4.2	3.6	3.1	3.1
2001.....	2.1	*	*	*	0.3	1.8	4.4	4.0	3.2	3.0
2000.....	1.9	*	*	*	*	1.6	3.9	3.6	2.6	2.8
1999.....	1.6	*	*	*	0.2	1.2	3.7	2.9	2.4	2.5

White, female

2006 4/.....	0.3	*	*	*	*	0.4	0.4	0.3	0.5	0.5
2005 4/.....	0.3	*	*	*	*	*	0.4	0.4	0.5	0.6
2004 4/.....	0.3	*	*	*	*	*	0.4	0.4	0.3	0.5
2003 4/.....	0.3	*	*	*	*	*	0.5	0.3	0.4	0.4
2002.....	0.2	*	*	*	*	*	0.3	0.3	0.4	0.6
2001.....	0.2	*	*	*	*	*	0.3	0.3	0.4	0.6
2000.....	0.2	*	*	*	*	*	*	0.3	0.3	0.4
1999.....	0.2	*	*	*	*	*	*	0.3	*	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

All races 3/, both sexes

2006.....	1.4	*	0.2	0.1	0.3	3.6	3.9	2.2	1.6	1.5
2005.....	1.5	*	0.1	0.1	0.2	3.6	3.8	2.3	1.7	1.5
2004.....	1.6	*	0.2	0.2	0.3	3.9	4.4	2.5	1.7	1.6
2003.....	1.7	*	0.2	0.2	0.4	4.0	4.4	2.8	1.9	1.6
2002.....	1.8	*	0.2	0.2	0.4	4.5	4.6	2.8	1.9	1.8
2001.....	1.5	*	0.1	0.2	0.2	3.7	4.0	2.2	1.8	1.5
2000.....	1.4	*	0.2	0.2	0.3	3.4	3.5	2.0	1.5	1.6
1999.....	1.4	*	0.1	0.1	0.3	3.3	3.3	2.0	1.6	1.4

All races 3/, male

2006.....	2.1	*	*	*	0.3	5.1	5.9	3.3	2.4	2.2
2005.....	2.2	*	*	*	0.3	5.0	6.0	3.7	2.5	2.2
2004.....	2.4	*	*	0.2	0.3	5.3	6.6	3.8	2.5	2.6
2003.....	2.6	*	*	0.2	0.6	5.7	6.8	4.3	2.9	2.4
2002.....	2.6	*	*	0.2	0.4	6.3	7.0	4.4	2.9	2.7
2001.....	2.2	*	*	*	0.3	5.2	6.1	3.4	2.7	2.1
2000.....	2.0	*	*	*	0.3	4.8	5.7	3.2	2.2	2.3
1999.....	2.0	*	*	*	0.4	4.4	5.3	3.2	2.5	2.1

All races 3/, female

2006.....	0.8	*	*	*	0.2	2.0	1.7	1.0	0.6	0.9
2005.....	0.7	*	*	*	0.2	2.1	1.5	0.9	0.8	0.8
2004.....	0.8	*	*	*	0.3	2.3	1.9	1.1	0.8	0.7
2003.....	0.9	*	0.3	*	0.3	2.2	1.8	1.3	0.9	0.8
2002.....	0.9	*	*	*	0.3	2.6	1.9	1.1	0.8	0.9
2001.....	0.8	*	*	*	0.2	2.1	1.7	1.0	0.8	0.8
2000.....	0.8	*	*	*	0.3	1.9	1.3	0.8	0.8	1.0
1999.....	0.7	*	*	*	0.2	2.2	1.3	0.8	0.7	0.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

White, both sexes

2006 4/.....	1.5	*	*	0.1	0.3	3.9	4.1	2.1	1.6	1.5
2005 4/.....	1.5	*	*	0.2	0.3	3.9	4.1	2.4	1.8	1.5
2004 4/.....	1.6	*	*	0.2	0.3	4.3	4.6	2.5	1.6	1.7
2003 4/.....	1.8	*	0.2	0.2	0.4	4.4	4.8	2.8	2.0	1.7
2002.....	1.8	*	0.2	0.2	0.3	4.9	4.8	2.8	1.9	1.8
2001.....	1.5	*	*	0.1	0.3	3.8	3.9	2.2	1.7	1.5
2000.....	1.4	*	*	0.2	0.3	3.6	3.5	1.9	1.5	1.7
1999.....	1.3	*	*	0.1	0.3	3.6	3.3	2.0	1.6	1.3

White, male

2006 4/.....	2.2	*	*	*	0.3	5.5	6.2	3.2	2.5	2.1
2005 4/.....	2.3	*	*	*	0.3	5.3	6.4	3.7	2.6	2.2
2004 4/.....	2.4	*	*	*	0.3	6.0	7.1	3.8	2.3	2.7
2003 4/.....	2.6	*	*	0.3	0.6	6.2	7.3	4.2	3.0	2.5
2002.....	2.6	*	*	*	0.3	6.7	7.3	4.4	3.0	2.5
2001.....	2.2	*	*	*	0.3	5.3	6.1	3.4	2.6	2.1
2000.....	2.0	*	*	*	0.3	5.0	5.6	3.0	2.2	2.3
1999.....	2.0	*	*	*	0.4	4.7	5.3	3.1	2.4	2.0

White, female

2006 4/.....	0.8	*	*	*	0.3	2.1	1.8	0.9	0.7	1.0
2005 4/.....	0.8	*	*	*	*	2.3	1.6	1.0	0.9	0.8
2004 4/.....	0.9	*	*	*	0.3	2.6	1.9	1.2	0.9	0.7
2003 4/.....	0.9	*	*	*	0.3	2.5	2.0	1.4	0.9	0.9
2002.....	1.0	*	*	*	0.3	2.9	2.0	1.2	0.8	1.0
2001.....	0.8	*	*	*	*	2.3	1.5	1.0	0.8	0.8
2000.....	0.8	*	*	*	0.3	2.1	1.3	0.8	0.9	1.0
1999.....	0.7	*	*	*	*	2.4	1.2	0.8	0.7	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

Black, both sexes

2006 4/.....	1.5	*	*	*	*	2.4	3.4	2.9	1.8	1.9
2005 4/.....	1.5	*	*	*	*	2.6	3.2	2.4	1.7	1.9
2004 4/.....	1.7	*	*	*	*	2.2	3.8	3.0	2.3	1.8
2003 4/.....	1.6	*	*	*	*	2.8	3.4	3.2	1.7	1.3
2002.....	1.9	*	*	*	*	3.3	4.3	3.1	2.1	2.4
2001.....	1.8	*	*	*	*	3.6	4.7	2.6	2.3	1.7
2000.....	1.6	*	*	*	*	2.6	3.7	2.7	1.4	1.7
1999.....	1.6	*	*	*	*	2.8	4.1	2.5	2.1	1.8

Black, male

2006 4/.....	2.4	*	*	*	*	3.3	5.2	4.3	3.3	3.3
2005 4/.....	2.4	*	*	*	*	4.0	5.1	4.3	3.0	3.0
2004 4/.....	2.5	*	*	*	*	3.1	5.3	4.9	4.1	2.6
2003 4/.....	2.6	*	*	*	*	4.0	5.9	5.7	2.7	2.3
2002.....	3.0	*	*	*	*	5.1	6.7	5.8	3.3	4.0
2001.....	2.8	*	*	*	*	5.3	7.4	4.2	3.7	2.6
2000.....	2.5	*	*	*	*	3.9	5.9	4.8	2.6	2.7
1999.....	2.6	*	*	*	*	4.0	6.4	4.3	3.6	2.7

Black, female

2006 4/.....	0.8	*	*	*	*	1.5	1.5	1.6	*	*
2005 4/.....	0.6	*	*	*	*	*	1.3	*	*	*
2004 4/.....	0.9	*	*	*	*	1.2	2.4	*	*	*
2003 4/.....	0.7	*	*	*	*	1.6	*	*	*	*
2002.....	0.9	*	*	*	*	1.5	1.9	*	*	*
2001.....	0.8	*	*	*	*	1.7	2.1	*	*	*
2000.....	0.7	*	*	*	*	*	1.7	*	*	*
1999.....	0.8	*	*	*	*	1.6	1.9	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

All races 3/, both sexes

2006.....	3.1	1.2	0.7	0.6	0.7	4.9	5.4	3.8	3.1	2.6
2005.....	3.4	1.4	0.8	0.7	0.8	5.5	5.9	3.9	3.3	2.9
2004.....	3.6	1.5	0.8	0.8	1.0	6.2	5.7	4.0	3.2	3.0
2003.....	3.7	1.5	0.9	0.8	0.8	6.2	6.0	4.1	3.4	3.2
2002.....	3.9	1.1	1.0	0.8	0.9	6.9	6.8	4.2	3.4	3.4
2001.....	3.7	1.6	1.0	0.8	1.0	6.3	5.9	4.2	3.1	3.3
2000.....	3.7	1.5	0.9	0.9	0.9	6.2	6.0	4.1	3.2	3.3
1999.....	3.7	1.6	0.9	0.9	1.0	6.0	5.7	4.0	3.5	3.0

All races 3/, male

2006.....	3.8	1.1	0.5	0.6	0.6	5.8	7.4	4.9	3.8	3.1
2005.....	4.2	1.5	0.7	0.7	0.7	6.5	7.8	5.4	4.3	3.6
2004.....	4.4	1.9	0.8	0.8	1.1	7.3	7.5	5.4	4.1	4.0
2003.....	4.5	1.4	0.9	0.7	0.9	7.0	7.9	5.4	4.5	4.1
2002.....	4.8	1.5	1.0	0.9	0.9	8.3	9.1	6.0	4.3	4.2
2001.....	4.5	1.6	0.9	0.8	1.0	7.6	8.1	5.7	4.3	4.2
2000.....	4.5	1.5	0.9	0.9	0.9	7.1	8.0	5.3	4.3	4.1
1999.....	4.4	1.7	1.0	0.9	1.0	7.1	7.5	5.2	4.6	3.5

All races 3/, female

2006.....	2.5	1.2	0.8	0.6	0.7	4.0	3.3	2.6	2.3	2.1
2005.....	2.7	1.2	0.9	0.8	0.8	4.5	4.0	2.4	2.2	2.2
2004.....	2.9	1.1	0.9	0.8	1.0	5.0	3.9	2.6	2.2	2.1
2003.....	2.9	1.5	0.9	0.9	0.7	5.3	4.1	2.7	2.3	2.3
2002.....	3.0	*	0.9	0.8	0.9	5.3	4.4	2.4	2.4	2.5
2001.....	2.9	1.6	1.1	0.9	1.0	4.9	3.6	2.7	2.0	2.4
2000.....	2.9	1.5	0.9	0.9	0.8	5.2	3.8	3.0	2.2	2.4
1999.....	2.9	1.4	0.7	0.9	1.1	4.8	3.8	2.7	2.4	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

White, both sexes

2006 4/.....	3.3	1.1	0.6	0.6	0.7	5.4	5.8	4.0	3.2	2.7
2005 4/.....	3.6	1.4	0.8	0.7	0.7	6.2	6.3	4.2	3.4	3.0
2004 4/.....	3.8	1.5	0.8	0.8	1.1	6.8	6.2	4.1	3.2	3.1
2003 4/.....	3.9	1.4	0.9	0.8	0.8	6.9	6.3	4.3	3.6	3.2
2002.....	4.1	1.0	0.9	0.8	1.0	7.6	7.2	4.2	3.4	3.4
2001.....	3.8	1.5	1.0	0.8	1.1	6.8	6.3	4.4	3.2	3.3
2000.....	3.8	1.5	0.9	0.9	0.9	6.8	6.2	4.2	3.2	3.3
1999.....	3.8	1.5	0.8	0.9	1.1	6.6	5.9	4.1	3.5	2.8

White, male

2006 4/.....	3.9	*	0.5	0.6	0.6	6.3	8.0	5.1	3.9	3.3
2005 4/.....	4.4	1.5	0.8	0.7	0.7	7.1	8.3	5.7	4.5	3.7
2004 4/.....	4.5	1.9	0.8	0.9	1.2	7.9	8.2	5.5	3.9	4.0
2003 4/.....	4.6	1.5	0.9	0.7	1.0	7.8	8.2	5.7	4.7	4.2
2002.....	5.0	1.4	1.0	0.9	1.0	9.0	9.7	5.9	4.3	4.2
2001.....	4.7	1.6	0.9	0.7	1.1	8.1	8.5	5.9	4.3	4.1
2000.....	4.6	1.5	0.9	1.0	0.9	7.8	8.1	5.3	4.2	4.1
1999.....	4.5	1.8	0.9	0.9	1.0	7.6	7.7	5.4	4.6	3.2

White, female

2006 4/.....	2.6	*	0.7	0.6	0.8	4.4	3.4	2.8	2.3	2.2
2005 4/.....	2.8	*	0.9	0.8	0.8	5.1	4.2	2.6	2.3	2.3
2004 4/.....	3.0	*	0.8	0.8	1.0	5.7	4.1	2.6	2.3	2.1
2003 4/.....	3.1	1.4	0.9	0.9	0.7	6.0	4.3	2.9	2.4	2.2
2002.....	3.2	*	0.9	0.7	1.0	6.1	4.6	2.3	2.4	2.6
2001.....	3.0	1.5	1.1	0.9	1.0	5.3	3.9	2.9	2.1	2.4
2000.....	3.1	1.5	0.9	0.9	0.9	5.7	4.1	2.9	2.3	2.4
1999.....	3.1	*	0.7	0.9	1.2	5.4	3.9	2.7	2.4	2.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

Black, both sexes

2006 4/.....	2.8	*	0.9	0.7	0.6	3.1	4.3	3.6	3.4	2.5
2005 4/.....	2.9	*	1.0	0.7	0.7	3.4	4.8	3.3	3.3	3.3
2004 4/.....	3.1	*	0.9	0.8	0.8	3.7	4.2	4.5	3.9	3.1
2003 4/.....	3.2	*	0.9	0.8	0.7	3.5	5.2	3.8	3.4	3.3
2002.....	3.3	*	1.2	0.8	0.7	4.1	5.5	5.5	4.1	3.4
2001.....	3.3	*	1.1	0.9	0.8	4.4	4.9	4.3	3.5	3.7
2000.....	3.3	*	1.0	0.7	0.8	3.7	5.5	4.7	3.7	3.3
1999.....	3.4	*	1.1	0.9	1.0	3.6	5.1	4.2	4.0	4.2

Black, male

2006 4/.....	3.7	*	*	*	*	4.1	5.6	4.8	4.3	3.2
2005 4/.....	3.8	*	*	*	*	4.6	6.3	4.6	4.6	4.4
2004 4/.....	4.0	*	*	*	*	5.0	5.1	6.3	6.2	4.2
2003 4/.....	4.1	*	*	*	*	4.6	7.2	5.2	4.8	3.7
2002.....	4.4	*	1.6	*	*	5.7	7.0	7.8	5.7	4.5
2001.....	4.4	*	*	*	*	5.6	6.9	5.9	5.2	4.8
2000.....	4.4	*	*	*	*	4.5	8.0	6.0	5.7	4.0
1999.....	4.5	*	*	*	*	5.0	6.9	5.7	5.5	5.7

Black, female

2006 4/.....	2.0	*	*	*	*	2.1	3.0	2.4	2.6	1.9
2005 4/.....	2.1	*	*	*	*	2.1	3.3	2.2	2.2	2.3
2004 4/.....	2.2	*	*	*	*	2.3	3.3	2.8	1.7	2.1
2003 4/.....	2.4	*	*	*	*	2.3	3.2	2.6	2.2	3.1
2002.....	2.4	*	*	*	*	2.4	4.1	3.4	2.6	2.5
2001.....	2.3	*	*	*	*	3.2	3.1	2.8	2.1	2.8
2000.....	2.2	*	*	*	*	2.9	3.2	3.5	1.9	2.6
1999.....	2.3	*	*	*	*	2.2	3.4	2.9	2.7	2.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)										
All races 3/, both sexes										
2006.....	1.3	0.6	0.3	0.2	0.4	2.6	3.1	1.9	1.7	1.5
2005.....	1.3	0.6	0.2	0.2	0.4	2.8	3.0	2.2	1.7	1.4
2004.....	1.4	*	0.3	0.4	0.4	2.9	3.2	1.9	1.5	1.6
2003.....	1.5	*	0.3	0.3	0.5	3.4	3.2	2.2	1.6	1.6
2002.....	1.5	*	0.3	0.3	0.5	3.7	3.4	2.3	1.8	1.7
2001.....	1.4	*	0.2	0.2	0.5	2.8	3.3	1.9	1.7	1.6
2000.....	1.3	0.6	0.2	0.2	0.5	2.9	3.2	2.2	1.5	1.5
1999.....	1.3	*	0.3	0.3	0.3	3.0	3.1	2.0	1.5	1.5
All races 3/, male										
2006.....	1.9	*	0.3	*	0.4	3.3	4.6	2.9	2.6	2.1
2005.....	1.9	*	*	0.3	0.4	3.5	4.5	3.3	2.3	1.9
2004.....	1.9	*	0.3	0.4	0.4	3.6	4.7	2.9	2.1	2.3
2003.....	2.1	*	0.3	0.3	0.6	4.3	4.8	3.2	2.4	2.2
2002.....	2.2	*	0.4	0.3	0.5	4.6	5.0	3.4	2.6	2.4
2001.....	2.0	*	*	0.3	0.5	3.4	5.0	2.8	2.6	2.3
2000.....	1.9	*	*	0.2	0.6	3.9	4.7	3.4	2.2	2.1
1999.....	1.9	*	0.3	0.3	0.3	3.9	4.6	3.0	2.1	2.1
All races 3/, female										
2006.....	0.8	*	*	0.2	0.3	1.9	1.6	0.9	0.8	0.8
2005.....	0.8	*	*	0.2	0.4	2.1	1.4	1.0	1.1	0.8
2004.....	0.8	*	0.3	0.3	0.4	2.3	1.6	0.9	0.9	0.8
2003.....	0.9	*	0.4	0.3	0.4	2.5	1.6	1.1	0.8	1.0
2002.....	0.9	*	*	0.3	0.4	2.7	1.7	1.1	1.0	1.1
2001.....	0.8	*	*	0.2	0.5	2.1	1.5	1.0	0.8	0.9
2000.....	0.7	*	*	0.3	0.4	1.9	1.6	0.9	0.8	0.9
1999.....	0.8	*	0.3	0.3	0.3	2.2	1.5	0.9	0.8	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)										
White, both sexes										
2006 4/.....	1.4	*	0.2	0.2	0.4	2.8	3.4	2.0	1.8	1.5
2005 4/.....	1.4	*	0.2	0.2	0.4	3.0	3.2	2.3	1.7	1.4
2004 4/.....	1.4	*	0.2	0.4	0.4	3.3	3.3	2.0	1.6	1.6
2003 4/.....	1.5	*	0.3	0.3	0.5	3.8	3.5	2.3	1.7	1.6
2002.....	1.6	*	0.3	0.3	0.5	4.0	3.6	2.5	1.9	1.8
2001.....	1.4	*	0.2	0.2	0.5	3.1	3.5	1.9	1.7	1.7
2000.....	1.4	*	0.2	0.2	0.4	3.2	3.4	2.3	1.6	1.5
1999.....	1.3	*	0.3	0.3	0.3	3.3	3.3	2.0	1.5	1.5
White, male										
2006 4/.....	1.9	*	*	*	0.4	3.6	5.0	3.0	2.7	2.1
2005 4/.....	2.0	*	*	0.3	0.4	3.9	4.9	3.4	2.4	2.0
2004 4/.....	2.0	*	*	0.4	0.4	4.0	5.0	3.0	2.2	2.4
2003 4/.....	2.1	*	*	*	0.5	4.8	5.1	3.5	2.5	2.1
2002.....	2.2	*	0.3	0.3	0.6	5.1	5.3	3.5	2.7	2.4
2001.....	2.0	*	*	*	0.5	3.7	5.5	2.8	2.6	2.4
2000.....	2.0	*	*	*	0.5	4.3	5.0	3.5	2.4	2.2
1999.....	1.9	*	*	0.3	0.3	4.2	4.9	3.1	2.2	2.1
White, female										
2006 4/.....	0.8	*	*	*	0.4	2.0	1.7	0.8	0.9	0.8
2005 4/.....	0.8	*	*	*	0.4	2.1	1.4	1.0	1.1	0.9
2004 4/.....	0.8	*	*	0.3	0.4	2.6	1.5	0.9	0.8	0.8
2003 4/.....	0.9	*	0.3	0.4	0.4	2.8	1.7	1.1	0.7	1.2
2002.....	0.9	*	*	0.3	0.4	2.8	1.7	1.3	1.1	1.1
2001.....	0.8	*	*	*	0.5	2.4	1.5	0.9	0.8	1.0
2000.....	0.7	*	*	0.3	0.3	2.1	1.6	1.0	0.7	0.9
1999.....	0.8	*	*	0.3	0.3	2.4	1.5	1.0	0.8	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

Black, both sexes

2006 4/.....	1.1	*	*	*	*	1.7	1.7	1.6	1.6	1.6
2005 4/.....	1.1	*	*	*	*	1.7	1.8	2.0	1.8	1.2
2004 4/.....	1.1	*	*	*	*	1.4	2.2	1.5	1.7	1.3
2003 4/.....	1.2	*	*	*	*	1.2	2.1	1.6	1.8	1.5
2002.....	1.2	*	*	*	*	1.8	2.2	1.7	1.2	1.4
2001.....	1.1	*	*	*	*	1.3	2.0	1.8	1.7	1.0
2000.....	1.1	*	*	*	0.6	1.4	2.5	1.6	1.1	1.3
1999.....	1.2	*	*	*	*	1.7	2.2	1.6	1.4	1.6

Black, male

2006 4/.....	1.6	*	*	*	*	2.0	2.5	2.3	2.5	2.3
2005 4/.....	1.6	*	*	*	*	2.0	2.8	2.8	2.6	2.1
2004 4/.....	1.7	*	*	*	*	2.0	3.0	2.4	2.0	1.9
2003 4/.....	1.7	*	*	*	*	1.7	3.1	2.0	2.3	2.6
2002.....	1.8	*	*	*	*	2.2	3.2	3.0	2.2	1.8
2001.....	1.6	*	*	*	*	1.8	2.4	2.5	3.2	1.5
2000.....	1.6	*	*	*	*	1.8	3.5	3.0	1.7	2.3
1999.....	1.7	*	*	*	*	2.2	2.9	2.8	2.0	2.4

Black, female

2006 4/.....	0.6	*	*	*	*	1.3	*	*	*	*
2005 4/.....	0.7	*	*	*	*	1.4	*	*	*	*
2004 4/.....	0.6	*	*	*	*	*	1.3	*	1.4	*
2003 4/.....	0.7	*	*	*	*	*	*	*	*	*
2002.....	0.6	*	*	*	*	1.4	*	*	*	*
2001.....	0.6	*	*	*	*	*	1.6	*	*	*
2000.....	0.6	*	*	*	*	*	1.5	*	*	*
1999.....	0.7	*	*	*	*	*	1.5	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

All races 3/, both sexes

2006.....	5.7	1.3	1.5	1.2	1.1	9.2	10.9	7.7	5.8	5.5
2005.....	5.6	1.3	1.4	1.1	1.2	8.8	10.4	7.0	5.7	5.1
2004.....	5.5	1.2	1.6	1.0	1.4	9.3	9.6	7.2	5.5	5.0
2003.....	5.3	1.3	1.3	1.1	1.4	9.2	9.4	6.0	4.9	5.2
2002.....	5.5	1.1	1.3	1.0	1.3	9.9	9.9	6.6	5.5	5.1
2001.....	5.8	1.3	1.5	1.2	1.3	10.2	10.6	7.2	5.4	5.8
2000.....	6.0	1.8	1.6	1.3	1.6	10.7	11.4	7.2	5.9	5.8
1999.....	6.0	2.3	1.6	1.3	1.7	11.1	10.5	7.0	5.7	6.0

All races 3/, male

2006.....	7.7	1.4	1.6	1.2	1.1	11.6	15.5	11.4	8.2	7.3
2005.....	7.6	1.1	1.6	1.2	1.3	11.1	15.1	10.0	8.1	7.0
2004.....	7.3	1.2	1.6	1.0	1.4	11.6	13.6	10.2	7.3	6.7
2003.....	7.0	1.6	1.5	1.2	1.5	11.6	13.3	8.6	7.1	7.0
2002.....	7.3	1.1	1.4	1.1	1.4	12.4	14.3	9.3	7.7	7.1
2001.....	7.7	1.1	1.6	1.3	1.4	13.3	15.5	10.4	7.5	7.7
2000.....	8.0	1.9	1.8	1.3	1.7	13.7	16.0	10.2	8.2	8.0
1999.....	7.9	2.1	1.8	1.4	1.8	13.8	15.0	9.9	7.6	8.0

All races 3/, female

2006.....	3.8	1.2	1.4	1.2	1.1	6.6	6.0	3.9	3.3	3.7
2005.....	3.7	1.5	1.2	1.0	1.1	6.4	5.4	3.8	3.3	3.3
2004.....	3.7	1.3	1.5	1.0	1.5	6.8	5.3	4.0	3.6	3.4
2003.....	3.6	*	1.1	0.9	1.2	6.6	5.2	3.3	2.6	3.4
2002.....	3.7	1.1	1.3	0.9	1.1	7.3	5.4	3.8	3.1	3.2
2001.....	3.9	1.5	1.3	1.1	1.1	6.9	5.5	4.0	3.4	4.0
2000.....	4.1	1.7	1.4	1.3	1.4	7.5	6.6	4.0	3.6	3.6
1999.....	4.2	2.5	1.5	1.2	1.5	8.2	5.8	4.1	3.8	4.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

White, both sexes

2006 4/.....	5.9	1.2	1.5	1.1	1.1	9.9	11.5	7.9	5.6	5.7
2005 4/.....	5.8	1.0	1.3	1.1	1.2	9.4	10.9	7.2	5.8	5.2
2004 4/.....	5.6	1.1	1.5	0.9	1.3	10.0	10.1	7.2	5.6	5.0
2003 4/.....	5.3	1.2	1.2	1.1	1.4	9.8	9.7	6.1	4.9	5.2
2002.....	5.6	0.8	1.3	0.9	1.3	10.9	10.3	6.6	5.6	5.1
2001.....	5.9	1.1	1.4	1.1	1.2	11.0	10.9	7.4	5.5	5.8
2000.....	6.2	1.5	1.6	1.3	1.6	11.8	11.9	7.1	5.8	5.9
1999.....	6.2	2.2	1.4	1.2	1.7	12.0	11.0	7.3	5.7	6.1

White, male

2006 4/.....	7.8	1.3	1.5	1.1	1.1	12.3	16.3	11.6	7.8	7.5
2005 4/.....	7.8	*	1.4	1.2	1.3	11.8	15.9	10.3	8.0	6.8
2004 4/.....	7.3	*	1.5	0.9	1.2	12.4	14.4	10.1	7.4	6.5
2003 4/.....	7.1	1.6	1.4	1.2	1.5	12.3	13.6	8.9	7.0	6.9
2002.....	7.4	*	1.3	0.9	1.4	13.5	14.7	9.2	7.8	7.0
2001.....	7.8	*	1.5	1.2	1.2	14.2	15.9	10.6	7.6	7.4
2000.....	8.2	1.6	1.8	1.2	1.8	15.0	16.8	10.0	8.2	8.1
1999.....	8.0	2.2	1.4	1.3	1.8	14.8	15.6	10.3	7.5	8.1

White, female

2006 4/.....	4.0	*	1.5	1.1	1.2	7.3	6.3	4.1	3.3	3.9
2005 4/.....	3.8	*	1.2	0.9	1.0	6.9	5.5	3.9	3.5	3.5
2004 4/.....	3.9	1.4	1.5	0.9	1.4	7.4	5.5	4.1	3.8	3.4
2003 4/.....	3.6	*	1.0	0.9	1.3	7.2	5.5	3.2	2.6	3.5
2002.....	3.9	*	1.3	0.9	1.1	8.0	5.6	3.8	3.2	3.1
2001.....	4.0	1.3	1.3	1.0	1.1	7.6	5.5	4.1	3.3	4.2
2000.....	4.2	1.4	1.4	1.3	1.5	8.4	6.6	4.0	3.4	3.7
1999.....	4.3	2.3	1.4	1.2	1.5	9.0	6.0	4.1	3.9	4.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

Black, both sexes

2006 4/.....	5.6	*	1.7	1.6	1.2	6.0	9.4	7.7	8.1	5.6
2005 4/.....	5.5	3.0	1.8	1.3	1.3	6.7	9.1	6.0	6.2	5.6
2004 4/.....	5.4	*	2.1	1.3	2.0	6.7	8.1	7.6	5.7	6.1
2003 4/.....	5.2	*	1.6	1.0	1.3	6.0	8.6	6.1	5.5	5.8
2002.....	5.1	*	1.5	1.4	1.2	6.3	8.7	7.7	5.6	5.9
2001.....	5.6	*	1.7	1.8	1.7	6.7	10.0	7.4	5.9	6.4
2000.....	5.9	3.6	1.9	1.5	1.3	7.0	10.8	8.7	7.4	6.0
1999.....	5.8	*	2.6	1.7	1.6	7.9	9.4	6.9	6.0	6.4

Black, male

2006 4/.....	8.4	*	2.0	1.6	1.5	8.9	13.3	12.0	13.2	8.4
2005 4/.....	8.0	*	2.2	1.3	1.3	8.7	13.8	9.5	9.9	8.8
2004 4/.....	7.8	*	2.3	1.2	2.3	8.6	12.0	11.7	8.3	9.3
2003 4/.....	7.4	*	1.6	*	1.3	8.3	13.4	9.1	8.4	8.6
2002.....	7.3	*	1.8	1.9	1.4	7.9	13.1	11.7	8.3	8.8
2001.....	8.0	*	1.8	2.2	2.2	9.3	14.9	11.3	8.2	9.7
2000.....	8.5	*	2.0	1.4	1.5	9.4	15.3	13.0	10.2	9.2
1999.....	8.2	*	3.1	2.0	1.9	10.5	14.1	10.2	9.0	8.7

Black, female

2006 4/.....	3.2	*	*	1.5	*	3.0	5.5	3.5	3.4	3.0
2005 4/.....	3.1	*	*	1.4	1.3	4.7	4.3	2.8	2.9	2.7
2004 4/.....	3.2	*	1.9	1.3	1.7	4.8	4.1	3.9	3.4	3.2
2003 4/.....	3.2	*	*	*	1.2	3.6	3.9	3.4	2.9	3.4
2002.....	3.1	*	*	*	*	4.7	4.4	4.0	3.3	3.4
2001.....	3.4	*	*	1.3	1.2	3.9	5.4	3.8	3.8	3.5
2000.....	3.6	*	1.8	1.6	*	4.5	6.6	4.8	5.0	3.2
1999.....	3.7	*	2.1	1.4	1.4	5.4	4.9	3.9	3.2	4.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2006.....	0.2	*	*	*	0.1	0.2	0.2	0.2	0.1	0.2
2005.....	0.3	*	0.2	*	0.1	0.3	0.3	0.2	0.2	0.2
2004.....	0.3	*	*	0.1	0.1	0.3	0.3	0.2	0.2	0.2
2003.....	0.3	*	*	0.1	0.1	0.5	0.5	0.3	0.3	0.2
2002.....	0.2	*	*	*	0.2	0.1	0.2	0.1	0.2	0.2
2001.....	0.3	*	0.2	0.1	0.2	0.3	0.3	0.2	0.2	0.4
2000.....	0.4	*	0.2	0.2	0.2	0.4	0.4	0.3	0.4	0.3
1999.....	0.6	*	0.3	0.3	0.2	0.7	0.7	0.5	0.6	0.6

All races 3/, male

2006.....	0.4	*	*	*	*	0.3	0.3	0.2	*	0.3
2005.....	0.4	*	*	*	*	0.4	0.4	0.4	0.4	0.4
2004.....	0.4	*	*	*	*	0.4	0.4	0.4	0.3	0.4
2003.....	0.5	*	*	*	*	0.6	0.7	0.5	0.4	0.3
2002.....	0.3	*	*	*	0.2	0.2	0.4	*	*	0.4
2001.....	0.5	*	*	*	0.2	0.4	0.5	0.3	0.4	0.6
2000.....	0.5	*	*	*	0.3	0.5	0.6	0.5	0.5	0.5
1999.....	0.8	*	0.3	0.3	0.3	0.9	1.1	0.9	0.7	0.8

All races 3/, female

2006.....	0.1	*	*	*	*	*	*	*	*	*
2005.....	0.1	*	*	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	0.3	0.2	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	0.2	*	*	0.2	*
1999.....	0.3	*	0.3	*	*	0.5	0.3	*	0.4	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, both sexes

2006 4/.....	0.2	*	*	*	0.1	0.2	0.2	*	0.1	0.2
2005 4/.....	0.3	*	*	*	0.1	0.2	0.3	0.2	0.2	0.3
2004 4/.....	0.3	*	*	*	*	0.3	0.3	0.2	0.2	0.2
2003 4/.....	0.3	*	*	*	0.1	0.5	0.5	0.4	0.3	0.2
2002.....	0.2	*	*	*	0.2	0.1	0.2	*	0.2	0.2
2001.....	0.3	*	0.2	*	0.2	0.3	0.3	0.2	0.2	0.3
2000.....	0.4	*	*	0.2	0.2	0.4	0.4	0.3	0.4	0.3
1999.....	0.6	*	0.3	0.2	0.3	0.7	0.8	0.6	0.6	0.6

White, male

2006 4/.....	0.4	*	*	*	*	0.3	0.3	*	*	0.3
2005 4/.....	0.4	*	*	*	*	0.3	0.4	0.4	0.3	0.4
2004 4/.....	0.4	*	*	*	*	0.4	0.3	0.3	0.3	0.4
2003 4/.....	0.5	*	*	*	*	0.7	0.8	0.6	0.4	0.3
2002.....	0.3	*	*	*	0.2	*	0.4	*	*	0.4
2001.....	0.5	*	*	*	*	0.4	0.5	0.3	0.3	0.5
2000.....	0.5	*	*	*	0.3	0.6	0.6	0.5	0.6	0.5
1999.....	0.8	*	*	0.3	0.3	0.9	1.1	0.9	0.7	0.8

White, female

2006 4/.....	0.1	*	*	*	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*	*	*	*
2003 4/.....	0.2	*	*	*	*	0.3	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	0.3	*	*	*	*
1999.....	0.4	*	*	*	*	0.5	0.3	*	0.4	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Falls (W00-W19)										
All races 3/, both sexes										
2006.....	7.0	0.6	0.2	*	0.1	0.4	0.8	0.7	0.9	1.1
2005.....	6.6	*	0.2	*	*	0.4	0.8	0.7	0.8	1.1
2004.....	6.4	0.6	0.3	*	0.1	0.4	0.7	0.8	0.8	1.1
2003.....	5.9	*	0.3	*	0.1	0.4	0.7	0.7	0.7	1.2
2002.....	5.6	*	0.2	*	0.1	0.4	0.8	0.7	0.8	1.2
2001.....	5.3	0.6	0.2	0.2	0.2	0.4	0.9	0.8	0.9	1.2
2000.....	4.7	*	0.2	*	0.1	0.5	0.7	0.7	0.8	1.0
1999.....	4.7	*	0.4	0.1	0.1	0.6	0.7	0.8	0.9	1.2
All races 3/, male										
2006.....	7.3	*	0.3	*	*	0.6	1.3	1.1	1.5	1.8
2005.....	7.0	*	0.2	*	*	0.6	1.2	1.2	1.3	1.8
2004.....	6.8	*	0.4	*	*	0.7	1.3	1.3	1.4	1.8
2003.....	6.2	*	0.5	*	0.2	0.7	1.1	1.3	1.1	2.0
2002.....	6.0	*	0.4	*	*	0.6	1.4	1.2	1.4	1.7
2001.....	5.8	*	0.3	0.2	0.2	0.8	1.5	1.5	1.5	2.1
2000.....	5.2	*	0.3	*	*	0.8	1.2	1.2	1.5	1.8
1999.....	5.2	*	0.5	*	0.2	0.9	1.2	1.4	1.5	2.0
All races 3/, female										
2006.....	6.6	*	*	*	*	*	*	*	0.3	0.4
2005.....	6.3	*	*	*	*	*	0.3	*	0.3	0.4
2004.....	6.0	*	*	*	*	*	*	0.2	0.2	0.5
2003.....	5.6	*	*	*	*	*	0.2	*	0.3	0.3
2002.....	5.3	*	*	*	*	*	*	*	0.3	0.6
2001.....	4.8	*	*	*	*	*	*	*	0.3	0.4
2000.....	4.3	*	*	*	*	*	0.2	*	*	0.3
1999.....	4.3	*	*	*	*	*	*	*	0.3	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Falls (W00-W19)										
White, both sexes										
2006 4/.....	7.9	*	0.2	*	*	0.4	0.9	0.7	1.0	1.2
2005 4/.....	7.5	*	*	*	*	0.4	0.9	0.8	0.8	1.1
2004 4/.....	7.3	*	0.3	*	0.1	0.5	0.8	0.8	0.9	1.2
2003 4/.....	6.7	*	0.3	*	0.2	0.5	0.8	0.8	0.8	1.2
2002.....	6.4	*	0.2	*	*	0.4	0.9	0.8	0.9	1.2
2001.....	6.0	*	0.2	0.2	0.2	0.5	1.0	0.9	0.9	1.3
2000.....	5.3	*	0.2	*	*	0.6	0.8	0.8	0.9	1.0
1999.....	5.2	*	0.4	*	0.2	0.6	0.8	0.9	0.9	1.3
White, male										
2006 4/.....	8.2	*	*	*	*	0.7	1.5	1.2	1.7	1.9
2005 4/.....	7.8	*	*	*	*	0.7	1.4	1.4	1.3	1.9
2004 4/.....	7.6	*	0.4	*	*	0.8	1.4	1.4	1.5	2.0
2003 4/.....	6.9	*	0.5	*	*	0.8	1.3	1.4	1.2	2.1
2002.....	6.6	*	*	*	*	0.7	1.6	1.4	1.5	1.8
2001.....	6.4	*	0.3	0.2	0.2	0.9	1.8	1.6	1.6	2.1
2000.....	5.7	*	*	*	*	1.0	1.4	1.4	1.5	1.7
1999.....	5.6	*	0.5	*	*	1.1	1.3	1.5	1.5	2.0
White, female										
2006 4/.....	7.7	*	*	*	*	*	*	*	0.3	0.4
2005 4/.....	7.3	*	*	*	*	*	0.3	*	*	0.4
2004 4/.....	7.0	*	*	*	*	*	*	*	0.3	0.4
2003 4/.....	6.5	*	*	*	*	*	0.3	*	0.3	0.3
2002.....	6.2	*	*	*	*	*	*	*	0.3	0.7
2001.....	5.5	*	*	*	*	*	*	*	0.3	0.4
2000.....	5.0	*	*	*	*	*	0.3	*	*	0.3
1999.....	4.9	*	*	*	*	*	*	*	0.3	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes										
2006.....	1.2	1.2	2.8	0.7	0.6	1.5	1.4	1.2	0.9	1.0
2005.....	1.2	1.6	3.0	0.6	0.6	1.5	1.6	1.0	0.9	1.0
2004.....	1.1	1.5	2.7	0.7	0.7	1.5	1.3	1.1	0.8	0.9
2003.....	1.1	1.4	2.9	0.6	0.7	1.4	1.4	1.0	0.8	1.0
2002.....	1.2	1.6	2.9	0.8	0.8	1.6	1.5	1.2	1.0	1.2
2001.....	1.2	1.7	3.0	0.8	0.8	1.6	1.4	1.1	0.8	1.0
2000.....	1.2	2.0	3.2	1.0	0.8	1.8	1.5	1.1	1.0	1.1
1999.....	1.3	1.8	3.2	0.9	0.9	1.8	1.5	1.1	1.1	1.1
All races 3/, male										
2006.....	1.9	1.3	3.5	1.0	0.8	2.6	2.5	2.2	1.6	1.6
2005.....	1.9	1.3	4.1	0.8	1.0	2.6	2.8	1.7	1.5	1.7
2004.....	1.8	1.8	3.2	0.9	1.0	2.6	2.3	1.9	1.4	1.6
2003.....	1.8	1.7	3.7	0.9	1.0	2.5	2.5	1.8	1.4	1.7
2002.....	1.9	1.9	3.9	1.2	1.1	2.8	2.8	2.1	1.7	2.0
2001.....	1.8	1.5	3.7	1.2	1.1	2.8	2.5	1.9	1.4	1.6
2000.....	2.0	2.2	4.0	1.3	1.2	3.3	2.5	1.9	1.7	1.9
1999.....	2.0	2.0	4.0	1.3	1.3	3.2	2.8	2.0	2.0	1.9
All races 3/, female										
2006.....	0.5	1.2	2.1	0.4	0.2	0.3	0.4	0.3	*	0.4
2005.....	0.5	1.8	2.0	0.4	0.3	0.3	0.3	0.3	0.3	0.3
2004.....	0.5	1.2	2.1	0.4	0.3	0.3	0.2	0.3	0.2	0.2
2003.....	0.5	1.2	2.0	0.4	0.3	*	0.2	*	*	0.3
2002.....	0.5	1.2	1.9	0.3	0.4	0.3	*	0.3	0.3	0.4
2001.....	0.5	1.9	2.2	0.5	0.5	0.3	0.3	0.3	0.2	0.4
2000.....	0.5	1.7	2.4	0.6	0.4	0.3	0.3	0.3	0.3	0.3
1999.....	0.5	1.6	2.4	0.5	0.4	0.3	0.3	*	0.3	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
White, both sexes										
2006 4/.....	1.2	1.4	2.9	0.5	0.5	1.3	1.5	1.3	0.8	1.0
2005 4/.....	1.2	1.8	3.3	0.5	0.5	1.3	1.6	1.0	0.9	0.9
2004 4/.....	1.1	1.6	3.0	0.5	0.4	1.2	1.3	1.1	0.8	0.9
2003 4/.....	1.1	1.4	3.1	0.6	0.5	1.3	1.3	1.0	0.8	1.0
2002.....	1.1	1.6	3.0	0.6	0.5	1.4	1.5	1.1	1.0	1.1
2001.....	1.1	1.7	3.3	0.6	0.5	1.3	1.4	1.0	0.7	1.0
2000.....	1.2	1.6	3.5	0.8	0.6	1.6	1.4	1.1	1.0	1.1
1999.....	1.2	1.5	3.5	0.7	0.5	1.6	1.5	1.0	1.1	1.1
White, male										
2006 4/.....	1.8	1.4	3.6	0.8	0.7	2.1	2.4	2.2	1.5	1.6
2005 4/.....	1.8	1.3	4.5	0.6	0.6	2.2	2.8	1.8	1.4	1.5
2004 4/.....	1.7	1.8	3.5	0.7	0.6	2.1	2.2	1.8	1.3	1.5
2003 4/.....	1.8	1.5	4.0	0.8	0.7	2.3	2.4	1.7	1.3	1.7
2002.....	1.8	1.9	4.0	0.9	0.6	2.4	2.7	1.9	1.7	1.9
2001.....	1.7	1.6	4.2	0.8	0.6	2.3	2.4	1.7	1.2	1.5
2000.....	1.9	1.8	4.4	1.0	0.8	2.8	2.4	1.8	1.7	1.8
1999.....	1.9	1.7	4.3	0.9	0.7	2.8	2.5	1.7	1.9	1.8
White, female										
2006 4/.....	0.5	1.4	2.3	0.3	*	0.3	0.4	0.3	*	0.4
2005 4/.....	0.5	2.2	2.1	0.4	0.3	0.3	0.3	0.3	0.3	0.3
2004 4/.....	0.5	1.4	2.4	0.3	*	*	0.3	0.3	0.3	0.3
2003 4/.....	0.5	1.3	2.2	0.3	*	*	*	*	*	0.3
2002.....	0.4	*	2.0	0.3	0.3	0.3	*	0.3	*	0.4
2001.....	0.5	1.8	2.4	0.5	0.3	*	0.3	0.3	*	0.4
2000.....	0.5	*	2.5	0.5	0.3	0.3	0.3	0.3	*	0.4
1999.....	0.5	1.4	2.6	0.4	*	0.3	0.3	*	0.3	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental drowning and submersion (W65-W74)

Black, both sexes

2006 4/.....	1.4	*	2.6	1.4	1.0	2.4	1.4	1.1	1.2	1.0
2005 4/.....	1.4	*	2.2	1.2	1.4	2.2	1.9	1.1	1.1	1.5
2004 4/.....	1.4	*	1.9	1.3	1.8	2.4	1.2	1.3	0.9	1.1
2003 4/.....	1.3	*	2.1	1.0	1.6	1.8	1.4	1.1	1.1	0.8
2002.....	1.6	*	2.5	1.8	2.2	2.3	1.9	1.6	1.2	1.5
2001.....	1.6	*	1.9	1.7	2.3	2.9	1.5	1.2	1.1	1.2
2000.....	1.6	4.2	2.2	1.9	2.1	2.8	1.7	1.3	0.9	1.2
1999.....	1.8	3.5	2.4	2.2	2.6	2.7	1.9	1.8	1.4	1.1

Black, male

2006 4/.....	2.4	*	3.6	2.1	1.6	4.7	2.5	2.3	2.2	2.1
2005 4/.....	2.5	*	2.7	1.6	2.6	4.2	3.3	1.9	2.1	2.7
2004 4/.....	2.3	*	2.8	1.6	2.7	4.3	2.3	2.2	1.6	2.2
2003 4/.....	2.2	*	3.0	1.4	2.4	3.4	2.6	2.1	2.2	1.7
2002.....	2.8	*	3.4	2.8	3.5	4.3	3.7	3.1	2.0	2.8
2001.....	2.7	*	2.2	2.7	3.2	5.2	3.1	2.6	2.2	2.2
2000.....	2.7	*	2.8	2.7	3.2	5.2	3.3	2.3	1.8	2.3
1999.....	3.1	*	3.2	3.4	3.9	5.0	3.8	3.5	2.7	2.2

Black, female

2006 4/.....	0.4	*	1.5	*	*	*	*	*	*	*
2005 4/.....	0.4	*	1.7	*	*	*	*	*	*	*
2004 4/.....	0.5	*	*	*	*	*	*	*	*	*
2003 4/.....	0.4	*	*	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*	*	*
2001.....	0.5	*	1.6	*	1.3	*	*	*	*	*
2000.....	0.6	*	1.7	*	*	*	*	*	*	*
1999.....	0.5	*	1.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

All races 3/, both sexes

2006.....	1.4	1.5	0.4	0.1	*	0.1	0.2	0.2	0.2	0.3
2005.....	1.5	1.2	0.4	*	*	0.1	0.2	0.3	0.2	0.3
2004.....	1.5	1.4	0.4	*	*	0.1	0.2	0.2	0.2	0.3
2003.....	1.5	1.5	0.6	*	*	0.1	0.2	0.2	0.2	0.3
2002.....	1.4	1.5	0.5	*	*	*	0.2	0.2	0.2	0.3
2001.....	1.5	1.5	0.5	*	*	*	0.2	0.1	0.2	0.3
2000.....	1.5	2.0	0.6	*	*	0.1	0.1	0.2	0.3	0.3
1999.....	1.4	1.7	0.6	0.1	0.1	*	0.1	0.1	0.3	0.3

All races 3/, male

2006.....	1.6	1.8	0.4	*	*	0.2	0.3	0.3	0.3	0.5
2005.....	1.5	1.0	0.5	*	*	*	0.3	0.4	0.3	0.4
2004.....	1.6	2.3	0.6	*	*	0.2	0.2	0.3	0.3	0.3
2003.....	1.6	1.6	0.7	*	*	*	0.2	0.3	0.3	0.5
2002.....	1.5	2.0	0.6	*	*	*	0.2	0.3	0.3	0.4
2001.....	1.5	2.0	0.6	*	*	*	0.2	*	0.2	0.4
2000.....	1.6	2.3	0.7	*	*	*	*	0.2	0.4	0.5
1999.....	1.4	1.9	0.8	*	*	*	*	*	0.3	0.3

All races 3/, female

2006.....	1.3	1.2	0.5	*	*	*	*	*	*	0.2
2005.....	1.4	1.3	0.3	*	*	*	*	*	*	0.2
2004.....	1.5	*	0.3	*	*	*	*	*	0.2	0.3
2003.....	1.4	1.5	0.4	*	*	*	*	*	*	0.2
2002.....	1.4	1.0	0.3	*	*	*	*	*	*	0.3
2001.....	1.4	1.0	0.4	*	*	*	*	*	*	0.3
2000.....	1.5	1.8	0.5	*	*	*	*	*	0.2	0.2
1999.....	1.4	1.4	0.4	*	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

All races 3/, both sexes

2006.....	0.5	18.9	0.4	0.1	0.2	0.2	0.3	0.2	0.2	0.3
2005.....	0.5	17.0	0.4	0.1	0.2	0.2	0.3	0.2	0.3	0.3
2004.....	0.5	16.4	0.3	0.1	0.3	0.2	0.2	0.2	0.3	0.3
2003.....	0.4	13.9	0.5	0.1	0.2	0.1	0.2	0.3	0.2	0.4
2002.....	0.5	14.3	0.4	0.1	0.2	0.2	0.3	0.2	0.3	0.3
2001.....	0.5	13.7	0.4	0.1	0.2	0.2	0.2	0.2	0.3	0.4
2000.....	0.5	11.8	0.4	0.2	0.3	0.2	0.3	0.2	0.3	0.4
1999.....	0.6	10.8	0.5	0.1	0.3	0.2	0.3	0.3	0.2	0.3

All races 3/, male

2006.....	0.7	22.1	0.4	*	0.4	0.4	0.4	0.4	0.4	0.4
2005.....	0.7	18.9	0.5	*	0.4	0.4	0.5	0.4	0.5	0.4
2004.....	0.7	17.7	0.4	0.2	0.5	0.3	0.4	0.3	0.4	0.5
2003.....	0.6	16.3	0.5	*	0.3	0.3	0.4	0.5	0.4	0.6
2002.....	0.7	15.7	0.5	*	0.4	0.4	0.5	0.4	0.5	0.5
2001.....	0.7	15.3	0.5	0.2	0.4	0.4	0.4	0.4	0.5	0.6
2000.....	0.7	13.2	0.5	*	0.5	0.4	0.5	0.4	0.5	0.7
1999.....	0.8	12.6	0.5	0.2	0.5	0.4	0.5	0.5	0.4	0.5

All races 3/, female

2006.....	0.3	15.5	0.4	*	*	*	*	*	*	*
2005.....	0.3	15.1	0.3	*	*	*	*	*	*	*
2004.....	0.3	14.9	*	*	*	*	*	*	*	*
2003.....	0.3	11.4	0.4	*	*	*	*	*	*	*
2002.....	0.3	12.8	0.4	*	*	*	*	*	*	*
2001.....	0.3	12.0	0.3	*	*	*	*	*	*	*
2000.....	0.3	10.3	0.3	*	*	*	*	*	*	*
1999.....	0.4	8.9	0.5	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)										
White, both sexes										
2006 4/.....	0.5	14.5	0.4	0.1	0.2	0.2	0.3	0.3	0.2	0.3
2005 4/.....	0.5	14.3	0.4	*	0.2	0.2	0.3	0.2	0.3	0.3
2004 4/.....	0.5	13.2	0.4	0.1	0.3	0.2	0.3	0.2	0.3	0.4
2003 4/.....	0.4	11.1	0.5	*	0.2	0.2	0.3	0.3	0.3	0.4
2002.....	0.4	10.7	0.4	*	0.3	0.3	0.3	0.3	0.3	0.4
2001.....	0.5	10.8	0.4	0.2	0.3	0.2	0.3	0.3	0.3	0.4
2000.....	0.5	9.4	0.4	0.2	0.3	0.3	0.3	0.3	0.3	0.4
1999.....	0.6	8.7	0.5	0.1	0.3	0.3	0.3	0.3	0.3	0.3
White, male										
2006 4/.....	0.7	16.9	0.4	*	0.4	0.4	0.5	0.5	0.4	0.5
2005 4/.....	0.7	15.9	0.5	*	0.3	0.4	0.5	0.4	0.6	0.4
2004 4/.....	0.7	15.0	0.5	*	0.5	0.4	0.5	0.3	0.5	0.6
2003 4/.....	0.6	13.2	0.6	*	0.3	0.3	0.5	0.6	0.4	0.7
2002.....	0.6	12.3	0.5	*	0.5	0.5	0.5	0.4	0.6	0.6
2001.....	0.6	11.9	0.4	*	0.5	0.4	0.5	0.5	0.5	0.7
2000.....	0.7	10.8	0.5	*	0.5	0.5	0.5	0.4	0.5	0.8
1999.....	0.8	9.7	0.5	0.2	0.5	0.5	0.5	0.5	0.4	0.6
White, female										
2006 4/.....	0.3	12.0	0.3	*	*	*	*	*	*	*
2005 4/.....	0.3	12.7	0.3	*	*	*	*	*	*	*
2004 4/.....	0.3	11.4	*	*	*	*	*	*	*	*
2003 4/.....	0.2	8.9	0.4	*	*	*	*	*	*	*
2002.....	0.3	9.0	0.4	*	*	*	*	*	*	*
2001.....	0.3	9.7	*	*	*	*	*	*	*	*
2000.....	0.3	7.9	0.3	*	*	*	*	*	*	*
1999.....	0.4	7.5	0.5	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes										
2006.....	1.0	0.7	1.2	0.6	0.3	0.4	0.5	0.5	0.5	0.6
2005.....	1.1	0.8	1.3	0.7	0.4	0.3	0.5	0.6	0.5	0.7
2004.....	1.1	0.6	1.4	0.9	0.4	0.3	0.6	0.7	0.5	0.7
2003.....	1.2	0.7	1.4	0.7	0.4	0.4	0.6	0.6	0.6	0.8
2002.....	1.1	0.9	1.4	0.8	0.5	0.4	0.5	0.6	0.7	0.7
2001.....	1.2	1.2	1.5	0.8	0.4	0.4	0.7	0.6	0.6	0.9
2000.....	1.2	1.0	1.9	0.9	0.4	0.4	0.6	0.6	0.6	0.8
1999.....	1.2	1.1	2.0	0.8	0.5	0.4	0.6	0.7	0.7	0.9
All races 3/, male										
2006.....	1.2	*	1.4	0.6	0.3	0.4	0.6	0.7	0.6	0.8
2005.....	1.3	*	1.5	0.8	0.4	0.4	0.6	0.7	0.6	0.9
2004.....	1.3	*	1.5	1.1	0.4	0.4	0.6	0.9	0.7	0.9
2003.....	1.4	*	1.6	0.7	0.4	0.4	0.7	0.7	0.8	1.1
2002.....	1.4	1.0	1.6	0.9	0.6	0.6	0.7	0.8	0.9	0.9
2001.....	1.4	1.5	1.8	1.0	0.4	0.4	0.9	0.8	0.8	1.2
2000.....	1.5	1.0	2.2	1.0	0.3	0.4	0.8	0.8	0.8	1.2
1999.....	1.5	1.1	2.2	0.9	0.5	0.6	0.8	0.9	1.0	1.2
All races 3/, female										
2006.....	0.8	*	1.0	0.6	0.3	0.3	0.3	0.3	0.4	0.5
2005.....	0.9	*	1.0	0.6	0.4	0.2	0.4	0.4	0.5	0.4
2004.....	0.9	*	1.3	0.6	0.4	0.2	0.5	0.4	0.3	0.4
2003.....	1.0	1.1	1.2	0.7	0.4	0.3	0.5	0.5	0.5	0.5
2002.....	0.8	*	1.2	0.6	0.4	0.3	0.4	0.4	0.6	0.4
2001.....	0.9	1.0	1.1	0.6	0.4	0.3	0.4	0.4	0.4	0.5
2000.....	0.9	*	1.5	0.7	0.5	0.4	0.4	0.4	0.4	0.5
1999.....	0.9	1.1	1.7	0.7	0.4	0.3	0.4	0.5	0.4	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
White, both sexes										
2006 4/.....	1.0	*	0.9	0.4	0.3	0.4	0.5	0.4	0.5	0.6
2005 4/.....	1.0	0.7	1.0	0.5	0.3	0.2	0.5	0.5	0.5	0.6
2004 4/.....	1.0	*	1.1	0.6	0.3	0.3	0.5	0.6	0.5	0.7
2003 4/.....	1.1	0.7	1.2	0.5	0.2	0.3	0.6	0.6	0.7	0.8
2002.....	1.0	*	1.0	0.5	0.4	0.4	0.5	0.5	0.7	0.6
2001.....	1.0	1.1	1.2	0.7	0.4	0.3	0.7	0.5	0.6	0.8
2000.....	1.0	0.8	1.3	0.7	0.3	0.4	0.5	0.5	0.5	0.7
1999.....	1.0	0.9	1.5	0.6	0.4	0.4	0.6	0.7	0.7	0.8
White, male										
2006 4/.....	1.2	*	1.2	0.4	0.3	0.5	0.6	0.6	0.6	0.7
2005 4/.....	1.2	*	1.3	0.6	0.3	0.3	0.6	0.7	0.6	0.8
2004 4/.....	1.2	*	1.2	0.7	0.3	0.4	0.6	0.8	0.7	0.8
2003 4/.....	1.3	*	1.4	0.5	*	0.4	0.8	0.7	0.8	1.1
2002.....	1.2	*	1.1	0.6	0.5	0.5	0.6	0.7	0.9	0.8
2001.....	1.3	*	1.6	0.8	0.4	0.4	0.9	0.7	0.8	1.0
2000.....	1.3	*	1.6	0.7	0.3	0.4	0.8	0.7	0.7	1.0
1999.....	1.3	*	1.7	0.6	0.4	0.6	0.7	0.9	1.0	1.0
White, female										
2006 4/.....	0.8	*	0.7	0.5	*	0.3	0.3	0.3	0.4	0.5
2005 4/.....	0.8	*	0.7	0.4	0.3	*	0.4	0.4	0.3	0.4
2004 4/.....	0.8	*	1.0	0.5	0.3	0.3	0.5	0.4	0.4	0.5
2003 4/.....	0.9	*	1.0	0.5	0.3	0.3	0.5	0.5	0.5	0.5
2002.....	0.7	*	1.0	0.4	*	*	0.3	0.3	0.5	0.4
2001.....	0.8	*	0.8	0.5	0.3	0.3	0.4	*	0.4	0.5
2000.....	0.8	*	1.1	0.6	0.4	0.4	0.3	0.4	0.3	0.4
1999.....	0.8	*	1.3	0.6	0.3	0.3	0.4	0.4	0.3	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental exposure to smoke, fire and flames (X00-X09)										
Black, both sexes										
2006 4/.....	1.8	*	3.0	1.5	*	*	0.7	0.8	0.9	1.1
2005 4/.....	1.9	*	2.8	1.6	0.9	*	*	0.9	1.0	1.1
2004 4/.....	2.1	*	2.8	2.3	0.9	*	0.8	1.2	0.8	0.9
2003 4/.....	2.0	*	3.0	1.9	1.0	0.6	*	0.9	0.8	1.0
2002.....	2.1	*	3.5	1.9	0.9	0.7	0.9	1.2	1.1	1.0
2001.....	2.3	*	3.3	1.8	0.8	*	1.0	1.4	1.0	1.5
2000.....	2.5	*	4.8	2.1	0.8	*	1.0	1.1	1.2	1.5
1999.....	2.5	*	4.5	2.1	0.9	*	0.9	1.1	0.9	1.9
Black, male										
2006 4/.....	2.1	*	3.0	1.5	*	*	*	*	*	*
2005 4/.....	2.4	*	2.8	1.7	*	*	*	*	*	*
2004 4/.....	2.6	*	2.9	3.1	*	*	*	1.6	*	*
2003 4/.....	2.4	*	3.2	1.9	*	*	*	*	*	*
2002.....	2.6	*	4.1	2.2	*	*	*	*	*	1.5
2001.....	2.9	*	3.6	2.3	*	*	*	*	*	2.5
2000.....	3.2	*	5.6	2.6	*	*	*	*	1.5	2.2
1999.....	3.0	*	5.0	2.4	*	*	*	*	*	2.8
Black, female										
2006 4/.....	1.5	*	2.9	1.5	*	*	*	*	*	*
2005 4/.....	1.5	*	2.7	1.6	*	*	*	*	*	*
2004 4/.....	1.6	*	2.7	1.4	*	*	*	*	*	*
2003 4/.....	1.7	*	2.7	1.8	*	*	*	*	*	*
2002.....	1.6	*	2.8	1.7	*	*	*	*	*	*
2001.....	1.7	*	2.9	1.2	*	*	*	1.5	*	*
2000.....	2.0	*	4.1	1.6	*	*	*	*	*	*
1999.....	2.0	*	3.9	1.8	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
All races 3/, both sexes										
2006.....	9.2	*	0.2	*	0.2	3.5	10.4	13.0	13.0	15.1
2005.....	8.0	0.5	0.1	*	0.2	3.0	8.8	10.6	11.3	13.5
2004.....	7.1	*	*	*	0.2	3.1	7.7	8.9	9.3	12.8
2003.....	6.7	0.5	0.3	*	0.2	2.5	7.1	7.9	9.3	12.5
2002.....	6.1	0.6	0.2	*	0.1	2.4	5.9	6.9	8.6	11.9
2001.....	4.9	*	0.2	*	0.2	2.0	4.9	5.7	6.9	10.4
2000.....	4.5	*	0.2	*	0.1	1.7	4.3	5.1	6.8	9.5
1999.....	4.4	*	0.2	*	0.1	1.3	3.8	4.9	6.8	9.1
All races 3/, male										
2006.....	12.6	*	*	*	0.2	5.2	16.0	19.2	18.6	20.5
2005.....	10.9	*	*	*	0.2	4.5	13.4	15.5	15.7	18.0
2004.....	9.6	*	*	*	0.2	4.6	11.8	13.3	13.0	16.8
2003.....	9.2	*	0.3	*	0.2	3.8	10.9	11.7	13.1	16.8
2002.....	8.5	1.0	*	*	*	3.5	9.1	10.4	12.5	16.5
2001.....	7.1	*	0.3	*	*	3.0	7.6	8.7	9.9	14.6
2000.....	6.6	*	0.3	*	*	2.6	6.7	7.9	10.1	13.7
1999.....	6.5	*	*	*	*	1.9	5.9	7.7	10.3	13.3
All races 3/, female										
2006.....	5.9	*	*	*	*	1.6	4.4	6.5	7.3	9.6
2005.....	5.1	*	*	*	*	1.5	3.9	5.4	6.7	8.9
2004.....	4.7	*	*	*	0.2	1.5	3.4	4.4	5.4	8.7
2003.....	4.3	*	0.3	*	*	1.2	3.1	4.0	5.4	8.1
2002.....	3.7	*	*	*	*	1.2	2.6	3.3	4.7	7.2
2001.....	2.9	*	*	*	*	1.0	1.9	2.6	3.9	6.1
2000.....	2.5	*	*	*	*	0.8	1.7	2.1	3.5	5.3
1999.....	2.3	*	*	*	*	0.7	1.6	2.0	3.2	4.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
White, both sexes										
2006 4/.....	9.7	*	*	*	0.2	4.2	12.3	15.1	14.6	16.2
2005 4/.....	8.4	*	*	*	0.2	3.6	10.4	12.1	12.5	14.6
2004 4/.....	7.6	*	*	*	0.2	3.7	9.1	10.4	10.3	13.8
2003 4/.....	7.1	*	0.3	*	0.1	3.0	8.5	9.1	10.2	13.3
2002.....	6.3	*	*	*	0.2	2.8	6.9	7.8	9.5	12.5
2001.....	5.0	*	0.2	*	0.2	2.3	5.6	6.4	7.4	10.8
2000.....	4.6	*	*	*	0.1	2.0	4.9	5.5	7.2	9.8
1999.....	4.3	*	0.2	*	0.1	1.5	4.2	5.2	6.9	9.3
White, male										
2006 4/.....	13.2	*	*	*	*	6.4	19.0	22.0	20.6	21.7
2005 4/.....	11.4	*	*	*	*	5.5	15.8	17.7	17.4	19.5
2004 4/.....	10.2	*	*	*	0.3	5.6	13.8	15.3	14.5	18.2
2003 4/.....	9.7	*	*	*	*	4.6	12.9	13.4	14.4	17.7
2002.....	8.8	*	*	*	*	4.1	10.6	11.6	13.6	17.3
2001.....	7.2	*	*	*	*	3.5	8.7	9.5	10.7	15.2
2000.....	6.7	*	*	*	*	3.1	7.8	8.6	10.7	14.3
1999.....	6.4	*	*	*	*	2.1	6.5	8.3	10.6	13.7
White, female										
2006 4/.....	6.3	*	*	*	*	1.9	5.2	7.8	8.3	10.4
2005 4/.....	5.4	*	*	*	*	1.7	4.4	6.2	7.3	9.5
2004 4/.....	5.0	*	*	*	*	1.7	4.0	5.1	6.0	9.3
2003 4/.....	4.5	*	*	*	*	1.4	3.8	4.5	5.8	8.7
2002.....	3.9	*	*	*	*	1.4	3.0	3.8	5.1	7.5
2001.....	2.9	*	*	*	*	1.0	2.3	3.0	3.9	6.2
2000.....	2.5	*	*	*	*	0.9	1.8	2.3	3.6	5.2
1999.....	2.3	*	*	*	*	0.8	1.7	2.1	3.1	4.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
Black, both sexes										
2006 4/.....	8.9	*	*	*	*	0.9	3.5	6.8	9.3	14.5
2005 4/.....	7.6	*	*	*	*	0.7	3.0	5.6	8.8	11.6
2004 4/.....	6.5	*	*	*	*	0.9	2.9	4.5	6.2	11.0
2003 4/.....	6.3	*	*	*	*	*	2.3	4.0	7.3	11.8
2002.....	6.3	*	*	*	*	*	2.3	4.4	6.8	11.9
2001.....	5.8	*	*	*	*	0.9	2.3	4.0	6.3	11.1
2000.....	5.5	*	*	*	*	0.7	2.1	4.3	6.8	10.7
1999.....	5.7	*	*	*	*	*	2.8	4.2	8.0	10.3
Black, male										
2006 4/.....	12.9	*	*	*	*	1.2	5.0	11.0	14.3	20.6
2005 4/.....	10.8	*	*	*	*	*	4.2	8.6	12.0	15.5
2004 4/.....	8.9	*	*	*	*	*	4.4	6.5	8.7	13.5
2003 4/.....	8.8	*	*	*	*	*	3.5	5.7	10.2	16.7
2002.....	9.1	*	*	*	*	*	3.5	7.0	9.9	17.0
2001.....	8.4	*	*	*	*	*	3.8	6.7	8.2	15.4
2000.....	8.2	*	*	*	*	*	2.7	7.1	9.6	14.9
1999.....	8.7	*	*	*	*	*	4.7	6.7	11.5	14.9
Black, female										
2006 4/.....	5.3	*	*	*	*	*	2.1	2.8	4.8	9.0
2005 4/.....	4.8	*	*	*	*	*	1.7	2.8	5.9	8.1
2004 4/.....	4.3	*	*	*	*	*	1.4	2.5	4.0	8.8
2003 4/.....	3.9	*	*	*	*	*	*	2.4	4.8	7.4
2002.....	3.7	*	*	*	*	*	*	2.0	4.0	7.4
2001.....	3.5	*	*	*	*	*	*	1.6	4.6	7.3
2000.....	3.1	*	*	*	*	*	1.5	1.7	4.3	7.1
1999.....	3.0	*	*	*	*	*	*	2.0	4.9	6.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

All races 3/, both sexes

2006.....	8.8	*	*	*	0.1	3.2	10.1	12.6	12.7	14.6
2005.....	7.6	*	*	*	0.1	2.8	8.4	10.3	11.0	13.0
2004.....	6.8	*	*	*	0.1	2.9	7.5	8.6	8.9	12.3
2003.....	6.3	*	0.2	*	*	2.3	6.8	7.6	8.9	11.9
2002.....	5.7	*	*	*	*	2.2	5.6	6.6	8.2	11.4
2001.....	4.6	*	*	*	*	1.7	4.5	5.4	6.6	9.8
2000.....	4.2	*	*	*	*	1.4	4.0	4.7	6.5	9.1
1999.....	4.0	*	*	*	*	1.0	3.4	4.6	6.4	8.6

All races 3/, male

2006.....	12.0	*	*	*	*	4.9	15.4	18.6	18.1	19.7
2005.....	10.3	*	*	*	*	4.2	12.8	15.0	15.2	17.2
2004.....	9.1	*	*	*	*	4.3	11.3	12.8	12.5	16.0
2003.....	8.6	*	*	*	*	3.4	10.4	11.2	12.5	16.0
2002.....	7.9	*	*	*	*	3.2	8.6	9.9	11.8	15.7
2001.....	6.5	*	*	*	*	2.6	7.1	8.2	9.5	13.8
2000.....	6.1	*	*	*	*	2.1	6.2	7.5	9.4	13.0
1999.....	5.9	*	*	*	*	1.5	5.4	7.2	9.8	12.5

All races 3/, female

2006.....	5.7	*	*	*	*	1.5	4.3	6.4	7.2	9.5
2005.....	4.9	*	*	*	*	1.3	3.7	5.3	6.6	8.7
2004.....	4.5	*	*	*	*	1.3	3.3	4.3	5.3	8.5
2003.....	4.1	*	*	*	*	1.2	3.0	3.8	5.3	7.9
2002.....	3.5	*	*	*	*	1.1	2.5	3.2	4.6	7.0
2001.....	2.7	*	*	*	*	0.8	1.8	2.5	3.7	5.8
2000.....	2.3	*	*	*	*	0.7	1.6	2.0	3.4	5.2
1999.....	2.1	*	*	*	*	0.5	1.5	2.0	3.0	4.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)										
White, both sexes										
2006 4/.....	9.3	*	*	*	0.1	3.9	12.0	14.7	14.3	15.7
2005 4/.....	8.0	*	*	*	*	3.4	10.0	11.8	12.1	14.0
2004 4/.....	7.2	*	*	*	0.1	3.4	8.8	10.0	10.0	13.2
2003 4/.....	6.7	*	*	*	*	2.8	8.1	8.7	9.8	12.7
2002.....	5.9	*	*	*	*	2.6	6.6	7.4	9.1	12.0
2001.....	4.7	*	*	*	*	2.0	5.2	6.0	7.1	10.2
2000.....	4.2	*	*	*	*	1.7	4.6	5.2	6.8	9.3
1999.....	4.0	*	*	*	*	1.2	3.8	5.0	6.6	8.8
White, male										
2006 4/.....	12.6	*	*	*	*	5.9	18.4	21.4	20.0	21.0
2005 4/.....	10.8	*	*	*	*	5.1	15.2	17.1	16.8	18.6
2004 4/.....	9.6	*	*	*	*	5.2	13.3	14.7	13.9	17.3
2003 4/.....	9.1	*	*	*	*	4.2	12.4	12.8	13.8	16.9
2002.....	8.2	*	*	*	*	3.7	10.0	11.0	13.0	16.5
2001.....	6.6	*	*	*	*	3.1	8.1	9.1	10.3	14.4
2000.....	6.1	*	*	*	*	2.5	7.3	8.1	10.0	13.5
1999.....	5.9	*	*	*	*	1.7	5.9	7.8	10.1	12.9
White, female										
2006 4/.....	6.1	*	*	*	*	1.8	5.1	7.6	8.2	10.3
2005 4/.....	5.3	*	*	*	*	1.6	4.3	6.1	7.2	9.3
2004 4/.....	4.8	*	*	*	*	1.6	3.9	5.0	5.8	9.0
2003 4/.....	4.3	*	*	*	*	1.4	3.6	4.3	5.7	8.4
2002.....	3.7	*	*	*	*	1.3	2.9	3.7	5.0	7.3
2001.....	2.7	*	*	*	*	0.9	2.2	2.8	3.8	6.0
2000.....	2.3	*	*	*	*	0.8	1.7	2.2	3.5	5.1
1999.....	2.1	*	*	*	*	0.7	1.6	2.1	3.0	4.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Black, both sexes

2006 4/.....	8.6	*	*	*	*	0.8	3.3	6.5	9.1	14.1
2005 4/.....	7.2	*	*	*	*	*	2.6	5.5	8.6	11.2
2004 4/.....	6.1	*	*	*	*	0.7	2.7	4.3	6.0	10.7
2003 4/.....	5.9	*	*	*	*	*	2.1	3.7	7.1	11.5
2002.....	5.8	*	*	*	*	*	2.2	4.2	6.4	11.2
2001.....	5.4	*	*	*	*	*	2.1	3.8	6.1	10.5
2000.....	5.2	*	*	*	*	*	2.0	3.9	6.6	10.3
1999.....	5.3	*	*	*	*	*	2.5	4.2	7.6	9.8

Black, male

2006 4/.....	12.4	*	*	*	*	*	4.4	10.4	14.1	20.2
2005 4/.....	10.2	*	*	*	*	*	3.7	8.4	11.7	15.0
2004 4/.....	8.4	*	*	*	*	*	4.1	6.4	8.4	13.1
2003 4/.....	8.2	*	*	*	*	*	3.2	5.1	9.7	16.3
2002.....	8.4	*	*	*	*	*	3.4	6.7	9.3	15.8
2001.....	7.8	*	*	*	*	*	3.5	6.3	8.0	14.5
2000.....	7.7	*	*	*	*	*	2.6	6.6	9.3	14.0
1999.....	8.1	*	*	*	*	*	4.2	6.7	10.9	13.9

Black, female

2006 4/.....	5.1	*	*	*	*	*	2.1	2.7	4.7	8.7
2005 4/.....	4.5	*	*	*	*	*	1.5	2.7	5.8	7.7
2004 4/.....	4.1	*	*	*	*	*	1.3	2.4	3.8	8.6
2003 4/.....	3.8	*	*	*	*	*	*	2.4	4.7	7.3
2002.....	3.5	*	*	*	*	*	*	2.0	3.9	7.1
2001.....	3.2	*	*	*	*	*	*	1.4	4.4	6.9
2000.....	2.9	*	*	*	*	*	1.4	1.5	4.2	7.1
1999.....	2.8	*	*	*	*	*	*	2.0	4.6	6.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

All races 3/, both sexes

2006.....	3.6	1.0	0.7	0.3	0.2	0.7	1.1	1.3	1.3	1.6
2005.....	3.7	1.2	0.6	0.3	0.3	1.0	1.2	1.4	1.5	1.6
2004.....	3.5	1.3	0.7	0.3	0.3	0.7	1.1	1.2	1.4	1.7
2003.....	3.7	1.1	0.7	0.2	0.3	0.7	1.3	1.1	1.2	1.5
2002.....	3.8	0.9	0.6	0.3	0.4	0.8	1.0	1.1	1.3	1.6
2001.....	4.0	1.3	0.8	0.3	0.3	0.8	1.2	1.1	1.3	1.8
2000.....	3.9	1.1	0.7	0.3	0.3	0.8	1.2	1.2	1.4	1.9
1999.....	4.3	1.2	0.8	0.4	0.4	0.9	1.2	1.3	1.4	1.7

All races 3/, male

2006.....	4.3	1.3	0.8	0.4	0.3	1.1	1.9	2.2	2.2	2.8
2005.....	4.5	1.2	0.7	0.4	0.4	1.5	2.0	2.2	2.5	2.6
2004.....	4.2	1.6	0.9	0.4	0.4	1.2	1.9	2.0	2.3	2.8
2003.....	4.3	1.1	0.8	0.3	0.5	1.2	2.0	1.8	2.1	2.4
2002.....	4.3	1.2	0.7	0.4	0.4	1.2	1.8	1.8	2.3	2.6
2001.....	4.5	1.6	0.9	0.4	0.4	1.2	2.1	2.0	2.3	3.0
2000.....	4.5	1.2	0.8	0.4	0.4	1.3	2.2	2.2	2.4	3.3
1999.....	4.8	1.4	1.0	0.5	0.5	1.4	2.0	2.3	2.3	2.9

All races 3/, female

2006.....	3.0	*	0.6	*	*	*	0.4	0.4	0.5	0.5
2005.....	3.0	1.3	0.6	0.2	*	0.4	0.4	0.4	0.5	0.5
2004.....	2.9	*	0.4	*	*	0.3	0.2	0.3	0.4	0.5
2003.....	3.1	1.1	0.5	*	*	0.2	0.4	0.4	0.4	0.6
2002.....	3.2	*	0.5	*	0.3	0.3	0.2	0.4	0.3	0.6
2001.....	3.5	1.1	0.7	0.2	*	0.4	0.3	0.3	0.4	0.6
2000.....	3.3	*	0.5	0.2	*	0.3	0.2	0.2	0.4	0.5
1999.....	3.7	*	0.5	0.3	0.2	0.4	0.3	0.4	0.4	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

White, both sexes

2006 4/.....	3.9	1.0	0.7	0.3	0.2	0.6	1.2	1.4	1.4	1.6
2005 4/.....	4.0	1.0	0.7	0.4	0.3	1.0	1.3	1.5	1.5	1.6
2004 4/.....	3.8	1.2	0.7	0.3	0.2	0.8	1.1	1.3	1.4	1.6
2003 4/.....	4.0	1.0	0.6	0.2	0.3	0.7	1.3	1.2	1.3	1.4
2002.....	4.1	0.8	0.5	0.3	0.3	0.8	1.1	1.2	1.3	1.5
2001.....	4.3	1.3	0.8	0.3	0.3	0.8	1.3	1.2	1.3	1.8
2000.....	4.2	1.0	0.6	0.3	0.3	0.8	1.3	1.3	1.4	1.9
1999.....	4.5	1.0	0.8	0.4	0.4	0.9	1.2	1.3	1.4	1.8

White, male

2006 4/.....	4.5	*	0.8	0.4	0.3	1.1	2.1	2.3	2.3	2.8
2005 4/.....	4.7	*	0.7	0.5	0.5	1.5	2.1	2.4	2.6	2.7
2004 4/.....	4.4	1.5	1.0	0.4	0.3	1.3	1.9	2.1	2.4	2.7
2003 4/.....	4.5	*	0.8	0.3	0.5	1.2	2.1	2.0	2.1	2.3
2002.....	4.5	*	0.6	0.5	0.3	1.2	2.0	2.0	2.3	2.5
2001.....	4.8	1.5	0.9	0.3	0.4	1.3	2.2	2.0	2.3	3.1
2000.....	4.6	*	0.7	0.4	0.4	1.3	2.2	2.3	2.5	3.2
1999.....	5.0	*	1.0	0.5	0.5	1.4	2.1	2.3	2.3	2.9

White, female

2006 4/.....	3.3	*	0.6	*	*	*	0.4	0.3	0.5	0.5
2005 4/.....	3.3	*	0.6	*	*	0.4	0.4	0.5	0.4	0.4
2004 4/.....	3.3	*	0.4	*	*	0.3	*	0.4	0.3	0.5
2003 4/.....	3.5	*	0.5	*	*	*	0.3	0.4	0.4	0.6
2002.....	3.6	*	0.4	*	0.3	0.3	*	0.4	0.3	0.5
2001.....	3.9	*	0.6	*	*	0.4	0.3	*	0.4	0.5
2000.....	3.7	*	0.5	*	*	0.3	*	0.3	0.4	0.5
1999.....	4.1	*	0.5	0.3	0.3	0.4	0.3	0.3	0.4	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
All races 3/, both sexes										
2006.....	11.1	*	1.0	7.3	12.5	12.4	12.3	14.7
2005.....	11.0	*	1.3	7.7	12.4	11.8	13.0	13.8
2004.....	11.0	*	1.3	8.2	12.5	12.3	13.0	14.0
2003.....	10.8	*	1.2	7.3	12.1	11.9	13.5	13.8
2002.....	11.0	*	1.2	7.4	12.4	12.8	12.5	14.3
2001 5/.....	10.8	*	1.3	7.9	12.0	12.6	13.0	14.3
2000.....	10.4	*	1.5	8.0	12.5	12.1	12.0	13.7
1999.....	10.5	*	1.2	8.0	12.3	12.3	13.1	14.4
All races 3/, male										
2006.....	17.8	*	1.4	11.5	20.8	19.7	19.7	22.7
2005.....	17.7	*	1.9	12.1	20.2	19.2	20.6	21.4
2004.....	17.7	*	1.7	12.6	20.8	20.2	20.6	21.7
2003.....	17.6	*	1.7	11.6	20.2	19.5	21.6	21.9
2002.....	17.9	*	1.8	12.2	20.8	20.8	20.2	22.7
2001 5/.....	17.6	*	1.9	12.9	20.5	21.0	21.0	22.7
2000.....	17.1	*	2.3	13.0	21.4	20.0	19.2	21.7
1999.....	17.1	*	1.9	13.0	20.8	20.3	21.1	22.7
All races 3/, female										
2006.....	4.6	*	0.6	2.8	3.6	4.7	4.7	6.6
2005.....	4.5	*	0.7	3.0	4.0	4.1	5.3	6.1
2004.....	4.6	*	1.0	3.5	3.6	4.2	5.3	6.3
2003.....	4.3	*	0.5	2.7	3.4	4.0	5.2	5.7
2002.....	4.3	*	0.6	2.4	3.5	4.5	4.7	6.0
2001 5/.....	4.1	*	0.6	2.7	3.1	4.0	4.8	5.9
2000.....	4.0	*	0.6	2.7	3.2	4.0	4.6	5.8
1999.....	4.0	*	0.5	2.7	3.4	4.1	5.1	6.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

White, both sexes

2006 4/.....	12.4	*	1.0	7.8	13.2	13.4	13.5	16.5
2005 4/.....	12.3	*	1.3	8.3	13.1	12.5	14.3	15.4
2004 4/.....	12.3	*	1.3	8.8	13.2	13.0	14.2	15.6
2003 4/.....	12.1	*	1.1	7.9	12.5	12.6	14.7	15.3
2002.....	12.2	*	1.3	8.2	13.2	13.5	13.7	15.9
2001 5/.....	11.9	*	1.3	8.6	12.6	13.5	14.2	15.8
2000.....	11.5	*	1.5	8.5	13.1	12.7	13.0	15.1
1999.....	11.5	*	1.3	8.5	12.7	12.8	14.2	15.7

White, male

2006 4/.....	19.8	*	1.4	12.3	21.9	21.0	21.2	25.0
2005 4/.....	19.7	*	1.9	13.3	21.3	20.3	22.2	23.7
2004 4/.....	19.6	*	1.7	13.6	22.0	21.2	22.3	24.0
2003 4/.....	19.5	*	1.7	12.7	21.1	20.3	23.3	23.9
2002.....	19.9	*	1.9	13.4	22.0	21.4	21.7	24.7
2001 5/.....	19.5	*	2.0	14.0	21.3	22.2	22.6	24.7
2000.....	18.8	*	2.4	13.7	22.3	20.8	20.7	23.6
1999.....	18.7	*	2.0	13.7	21.5	21.0	22.5	24.3

White, female

2006 4/.....	5.2	*	0.6	3.0	3.7	5.3	5.4	7.7
2005 4/.....	5.0	*	0.6	3.1	4.3	4.2	6.0	6.8
2004 4/.....	5.1	*	0.9	3.7	3.8	4.3	5.6	7.0
2003 4/.....	4.7	*	0.5	2.9	3.3	4.3	5.7	6.5
2002.....	4.8	*	0.7	2.6	3.7	5.1	5.3	6.8
2001 5/.....	4.6	*	0.6	2.9	3.3	4.3	5.4	6.7
2000.....	4.4	*	0.7	2.9	3.3	4.1	5.1	6.5
1999.....	4.5	*	0.5	2.9	3.5	4.3	5.5	6.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
Black, both sexes										
2006 4/.....	4.9	*	1.1	4.2	8.5	8.6	8.1	8.2
2005 4/.....	5.1	*	1.1	4.4	9.1	8.8	8.5	7.2
2004 4/.....	5.2	*	1.4	4.7	9.8	10.1	8.7	7.0
2003 4/.....	5.1	*	1.2	3.8	10.5	9.5	9.1	7.5
2002.....	5.1	*	1.0	4.0	9.1	10.3	8.2	8.1
2001 5/.....	5.3	*	1.1	4.3	10.1	9.6	8.6	7.8
2000.....	5.4	*	1.2	5.5	11.0	9.9	7.6	7.3
1999.....	5.4	*	0.9	5.8	10.7	9.6	8.7	8.4
Black, male										
2006 4/.....	8.8	*	1.7	7.0	14.5	15.7	15.2	14.9
2005 4/.....	8.7	*	1.5	7.2	16.2	14.9	15.0	12.0
2004 4/.....	9.0	*	1.6	7.4	17.2	18.0	14.4	11.7
2003 4/.....	8.8	*	1.9	6.6	18.0	17.4	15.7	12.9
2002.....	9.1	*	1.6	6.9	16.1	19.5	15.2	14.5
2001 5/.....	9.2	*	1.7	7.3	19.4	17.7	15.4	13.8
2000.....	9.4	*	2.1	9.5	19.5	17.5	13.6	13.0
1999.....	9.6	*	1.3	9.9	19.5	17.9	15.7	15.6
Black, female										
2006 4/.....	1.4	*	*	1.3	2.3	1.9	1.7	2.2
2005 4/.....	1.8	*	*	1.4	2.0	3.0	2.7	2.8
2004 4/.....	1.8	*	*	1.9	2.4	2.8	3.6	2.7
2003 4/.....	1.8	*	*	*	3.1	2.2	3.1	2.6
2002.....	1.5	*	*	*	2.3	2.0	2.0	2.5
2001 5/.....	1.7	*	*	1.3	*	2.2	2.5	2.5
2000.....	1.7	*	*	1.4	3.0	3.0	2.2	2.2
1999.....	1.6	*	*	1.6	2.3	2.1	2.5	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

All races 3/, both sexes

2006.....	1.5	*	*	0.3	0.9	1.4	1.7	2.3
2005.....	1.4	*	*	0.4	0.9	1.0	1.7	2.2
2004.....	1.4	*	*	0.5	0.8	1.2	1.6	2.3
2003.....	1.3	*	*	0.4	0.7	1.1	1.6	2.1
2002.....	1.3	*	*	0.4	0.7	1.1	1.6	2.2
2001.....	1.2	*	*	0.4	0.7	1.0	1.5	2.1
2000.....	1.2	*	*	0.4	0.7	1.0	1.3	1.8
1999.....	1.1	*	*	0.3	0.7	1.0	1.3	1.9

All races 3/, male

2006.....	1.4	*	*	0.3	1.0	1.4	1.7	2.3
2005.....	1.3	*	*	0.4	0.8	1.0	1.6	2.1
2004.....	1.4	*	*	0.4	1.0	1.3	1.7	2.4
2003.....	1.3	*	*	0.4	0.7	1.1	1.6	2.2
2002.....	1.3	*	*	0.4	0.7	1.3	1.7	2.2
2001.....	1.2	*	*	0.4	0.7	1.1	1.6	2.4
2000.....	1.1	*	*	0.4	0.7	0.9	1.5	1.9
1999.....	1.1	*	*	0.3	0.7	1.0	1.2	2.0

All races 3/, female

2006.....	1.6	*	*	0.3	0.8	1.4	1.7	2.3
2005.....	1.5	*	*	0.4	0.9	1.0	1.7	2.3
2004.....	1.5	*	*	0.5	0.7	1.1	1.6	2.2
2003.....	1.4	*	*	0.4	0.7	1.0	1.7	2.1
2002.....	1.4	*	*	0.4	0.8	1.0	1.4	2.1
2001.....	1.3	*	*	0.5	0.8	0.9	1.4	1.8
2000.....	1.2	*	*	0.4	0.6	1.1	1.2	1.8
1999.....	1.2	*	*	0.4	0.6	1.1	1.4	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)										
White, both sexes										
2006 4/.....	1.8	*	*	0.3	1.1	1.6	1.9	2.6
2005 4/.....	1.6	*	*	0.4	1.0	1.0	1.9	2.5
2004 4/.....	1.6	*	*	0.5	1.0	1.4	1.9	2.6
2003 4/.....	1.5	*	*	0.4	0.8	1.2	1.9	2.4
2002.....	1.5	*	*	0.4	0.9	1.3	1.7	2.5
2001.....	1.4	*	*	0.5	0.8	1.1	1.7	2.4
2000.....	1.3	*	*	0.4	0.7	1.1	1.5	2.1
1999.....	1.3	*	*	0.4	0.7	1.1	1.4	2.2
White, male										
2006 4/.....	1.6	*	*	0.3	1.2	1.6	1.8	2.5
2005 4/.....	1.5	*	*	0.5	1.0	1.1	1.9	2.4
2004 4/.....	1.6	*	*	0.5	1.1	1.5	1.9	2.7
2003 4/.....	1.5	*	*	0.5	0.8	1.3	1.8	2.5
2002.....	1.5	*	*	0.5	0.8	1.4	1.9	2.5
2001.....	1.4	*	*	0.5	0.8	1.2	1.8	2.7
2000.....	1.2	*	*	0.4	0.7	1.0	1.7	2.1
1999.....	1.3	*	*	0.3	0.8	1.1	1.3	2.2
White, female										
2006 4/.....	1.9	*	*	0.4	0.9	1.7	2.0	2.7
2005 4/.....	1.7	*	*	0.4	1.0	1.0	1.9	2.6
2004 4/.....	1.7	*	*	0.6	0.8	1.3	1.8	2.5
2003 4/.....	1.6	*	*	0.4	0.7	1.1	2.0	2.4
2002.....	1.6	*	*	0.4	0.9	1.2	1.5	2.4
2001.....	1.5	*	*	0.5	0.9	1.0	1.6	2.1
2000.....	1.4	*	*	0.4	0.7	1.2	1.4	2.0
1999.....	1.3	*	*	0.5	0.7	1.2	1.5	2.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

Black, both sexes

2006 4/.....	0.4	*	*	*	*	*	*	0.9
2005 4/.....	0.5	*	*	*	*	*	0.9	0.8
2004 4/.....	0.4	*	*	*	*	*	0.8	0.9
2003 4/.....	0.4	*	*	*	*	*	*	0.9
2002.....	0.4	*	*	*	*	*	1.0	0.8
2001.....	0.4	*	*	*	*	*	*	1.0
2000.....	0.4	*	*	*	*	*	*	0.8
1999.....	0.4	*	*	*	*	*	0.8	0.7

Black, male

2006 4/.....	0.5	*	*	*	*	*	*	*
2005 4/.....	0.4	*	*	*	*	*	*	*
2004 4/.....	0.4	*	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*

Black, female

2006 4/.....	0.4	*	*	*	*	*	*	*
2005 4/.....	0.5	*	*	*	*	*	*	*
2004 4/.....	0.5	*	*	*	*	*	*	*
2003 4/.....	0.5	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

All races 3/, both sexes

2006.....	0.1	*	*	*	*	*	*	0.1
2005.....	0.1	*	*	*	*	*	0.1	0.1
2004.....	0.1	*	*	*	*	0.1	0.1	0.1
2003.....	0.1	*	*	*	*	*	0.1	0.1
2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	0.1	*	0.1
2000.....	0.1	*	*	*	*	*	*	0.1
1999.....	0.1	*	*	*	*	*	*	0.1

All races 3/, male

2006.....	0.1	*	*	*	*	*	*	*
2005.....	0.1	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	0.2	*
2003.....	0.1	*	*	*	*	*	*	0.2
2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	0.2
1999.....	0.1	*	*	*	*	*	*	*

All races 3/, female

2006.....	0.0	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

White, both sexes

2006 4/.....	0.1	*	*	*	*	*	*	0.1
2005 4/.....	0.1	*	*	*	*	*	*	0.1
2004 4/.....	0.1	*	*	*	*	*	0.1	*
2003 4/.....	0.1	*	*	*	*	*	*	0.1
2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	0.1
1999.....	0.1	*	*	*	*	*	*	*

White, male

2006 4/.....	0.1	*	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*	0.2
2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	0.2
1999.....	0.1	*	*	*	*	*	*	*

White, female

2006 4/.....	0.0	*	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other and unspecified solid or liquid substances and their vapors (X65-X66,X68-X69)

Black, both sexes

2006 4/.....	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Black, male

2006 4/.....	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

All races 3/, both sexes

2006.....	0.4	*	*	*	0.3	0.3	0.4	0.8
2005.....	0.4	*	*	0.1	0.2	0.4	0.6	0.5
2004.....	0.5	*	*	0.1	0.2	0.4	0.5	0.7
2003.....	0.5	*	*	*	0.2	0.3	0.7	0.9
2002.....	0.5	*	*	0.2	0.2	0.4	0.7	0.8
2001.....	0.5	*	*	0.1	0.3	0.4	0.7	0.8
2000.....	0.5	*	*	0.2	0.3	0.6	0.7	0.9
1999.....	0.6	*	*	0.2	0.5	0.5	0.9	0.9

All races 3/, male

2006.....	0.7	*	*	*	0.4	0.5	0.7	1.2
2005.....	0.7	*	*	0.2	0.4	0.6	0.9	0.9
2004.....	0.7	*	*	0.2	0.4	0.7	0.9	1.1
2003.....	0.7	*	*	*	0.4	0.6	1.2	1.4
2002.....	0.8	*	*	0.3	0.4	0.7	1.0	1.2
2001.....	0.8	*	*	0.2	0.6	0.7	1.1	1.3
2000.....	0.8	*	*	0.3	0.5	0.9	1.1	1.4
1999.....	0.9	*	*	0.3	0.9	0.9	1.3	1.4

All races 3/, female

2006.....	0.2	*	*	*	*	*	*	0.3
2005.....	0.2	*	*	*	*	*	0.2	0.2
2004.....	0.2	*	*	*	*	*	*	0.3
2003.....	0.2	*	*	*	*	*	0.2	0.3
2002.....	0.2	*	*	*	*	*	0.3	0.3
2001.....	0.2	*	*	*	*	*	0.2	0.3
2000.....	0.2	*	*	*	*	0.2	0.3	0.4
1999.....	0.2	*	*	*	*	*	0.4	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

White, both sexes

2006 4/.....	0.5	*	*	*	0.3	0.4	0.5	0.9
2005 4/.....	0.5	*	*	0.2	0.3	0.4	0.7	0.6
2004 4/.....	0.5	*	*	0.2	0.3	0.5	0.6	0.9
2003 4/.....	0.6	*	*	*	0.3	0.4	0.8	1.0
2002.....	0.6	*	*	0.2	0.3	0.5	0.8	0.9
2001.....	0.6	*	*	0.2	0.4	0.5	0.8	0.9
2000.....	0.6	*	*	0.2	0.4	0.6	0.8	1.0
1999.....	0.6	*	*	0.2	0.6	0.6	1.0	1.0

White, male

2006 4/.....	0.8	*	*	*	0.4	0.5	0.8	1.4
2005 4/.....	0.8	*	*	0.3	0.4	0.6	1.1	1.0
2004 4/.....	0.8	*	*	0.3	0.4	0.8	1.0	1.3
2003 4/.....	0.9	*	*	*	0.4	0.7	1.4	1.6
2002.....	0.9	*	*	0.3	0.4	0.8	1.2	1.4
2001.....	0.9	*	*	0.2	0.7	0.8	1.3	1.5
2000.....	1.0	*	*	0.3	0.7	1.0	1.3	1.6
1999.....	1.0	*	*	0.3	1.1	0.9	1.6	1.6

White, female

2006 4/.....	0.2	*	*	*	*	*	*	0.4
2005 4/.....	0.2	*	*	*	*	*	0.3	0.3
2004 4/.....	0.2	*	*	*	*	*	*	0.4
2003 4/.....	0.2	*	*	*	*	*	*	0.4
2002.....	0.3	*	*	*	*	*	0.4	0.4
2001.....	0.3	*	*	*	*	*	*	0.4
2000.....	0.3	*	*	*	*	*	0.3	0.4
1999.....	0.3	*	*	*	*	*	0.5	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)										
Black, both sexes										
2006 4/.....	0.1	*	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*
Black, male										
2006 4/.....	0.1	*	*	*	*	*	*	*
2005 4/.....	0.2	*	*	*	*	*	*	*
2004 4/.....	0.2	*	*	*	*	*	*	*
2003 4/.....	0.2	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*
Black, female										
2006 4/.....	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

All races 3/, both sexes

2006.....	2.5	*	0.7	3.1	4.0	4.1	3.8	4.1
2005.....	2.4	*	0.8	3.1	4.4	3.8	3.8	3.8
2004.....	2.5	*	1.0	3.3	4.0	3.9	4.1	3.9
2003.....	2.3	*	0.7	2.8	3.8	3.6	3.8	3.5
2002.....	2.2	*	0.7	2.7	3.6	3.6	3.5	3.4
2001.....	2.2	*	0.8	2.7	3.5	3.5	3.5	3.6
2000.....	2.0	*	0.8	2.5	3.3	3.3	3.0	3.0
1999.....	1.9	*	0.6	2.2	2.9	3.1	3.4	3.2

All races 3/, male

2006.....	4.2	*	0.9	4.5	6.7	6.8	6.4	6.9
2005.....	4.0	*	1.2	4.4	7.1	6.3	6.3	6.4
2004.....	4.1	*	1.2	4.7	6.6	6.6	6.8	6.5
2003.....	3.8	*	1.1	4.2	6.3	6.0	6.4	6.0
2002.....	3.8	*	1.1	4.3	6.2	6.0	6.0	5.9
2001.....	3.7	*	1.2	4.3	6.0	6.0	5.9	6.2
2000.....	3.4	*	1.3	4.0	5.7	5.6	5.2	5.2
1999.....	3.3	*	1.0	3.4	4.8	5.3	5.9	5.6

All races 3/, female

2006.....	0.9	*	0.5	1.6	1.2	1.3	1.1	1.3
2005.....	0.9	*	0.5	1.6	1.5	1.2	1.2	1.2
2004.....	0.9	*	0.7	1.7	1.2	1.1	1.2	1.2
2003.....	0.8	*	0.3	1.2	1.1	1.1	1.2	1.0
2002.....	0.8	*	0.3	1.0	1.0	1.2	0.8	1.0
2001.....	0.7	*	0.3	1.0	0.9	0.9	1.0	1.1
2000.....	0.7	*	0.3	1.0	0.8	0.9	0.8	0.8
1999.....	0.7	*	0.2	0.9	0.8	0.9	0.9	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)										
White, both sexes										
2006 4/.....	2.7	*	0.6	3.3	4.3	4.5	4.2	4.5
2005 4/.....	2.6	*	0.8	3.3	4.7	4.0	4.1	4.2
2004 4/.....	2.6	*	0.9	3.3	4.3	4.1	4.4	4.3
2003 4/.....	2.4	*	0.7	3.0	3.9	3.8	4.2	3.7
2002.....	2.4	*	0.7	3.0	4.0	3.8	3.7	3.7
2001.....	2.3	*	0.8	2.9	3.8	3.6	3.6	4.0
2000.....	2.1	*	0.8	2.7	3.4	3.4	3.1	3.2
1999.....	2.0	*	0.6	2.2	3.0	3.3	3.6	3.4
White, male										
2006 4/.....	4.5	*	0.7	4.8	7.1	7.4	7.0	7.5
2005 4/.....	4.3	*	1.1	5.0	7.5	6.7	6.9	7.1
2004 4/.....	4.4	*	1.2	4.8	7.1	7.0	7.3	7.2
2003 4/.....	4.1	*	1.0	4.6	6.6	6.3	7.0	6.3
2002.....	4.0	*	1.1	4.9	6.7	6.3	6.3	6.2
2001.....	4.0	*	1.2	4.8	6.6	6.2	6.1	6.6
2000.....	3.6	*	1.2	4.3	6.0	6.0	5.4	5.5
1999.....	3.4	*	1.0	3.5	5.1	5.6	6.2	5.9
White, female										
2006 4/.....	0.9	*	0.4	1.7	1.2	1.4	1.3	1.4
2005 4/.....	0.9	*	0.4	1.6	1.7	1.1	1.3	1.3
2004 4/.....	0.9	*	0.7	1.8	1.3	1.0	1.3	1.3
2003 4/.....	0.8	*	0.2	1.3	1.1	1.2	1.3	1.0
2002.....	0.8	*	0.4	1.1	1.0	1.2	0.9	1.1
2001.....	0.7	*	0.3	1.0	0.9	0.9	1.0	1.2
2000.....	0.7	*	0.4	1.1	0.7	0.8	0.8	0.9
1999.....	0.7	*	*	0.9	0.8	0.9	0.9	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

Black, both sexes

2006 4/.....	1.2	*	0.9	1.5	1.8	2.2	2.1	2.3
2005 4/.....	1.3	*	0.9	1.4	2.7	2.4	2.1	1.8
2004 4/.....	1.4	*	1.0	2.2	2.6	2.9	2.5	1.5
2003 4/.....	1.3	*	1.0	1.3	2.7	2.3	2.0	2.0
2002.....	1.1	*	0.7	1.0	2.0	2.3	2.3	1.9
2001.....	1.1	*	0.6	1.1	1.8	2.5	2.6	1.8
2000.....	1.2	*	1.0	1.3	2.3	2.3	1.9	1.7
1999.....	1.0	*	0.7	1.3	1.8	2.0	2.2	1.9

Black, male

2006 4/.....	2.2	*	1.5	2.4	3.0	4.1	4.1	4.6
2005 4/.....	2.1	*	1.2	2.0	4.5	3.9	3.7	3.1
2004 4/.....	2.4	*	1.2	3.4	4.4	5.0	4.4	2.8
2003 4/.....	2.3	*	1.6	2.2	4.2	4.2	3.6	3.7
2002.....	2.0	*	1.1	1.6	3.1	4.0	4.3	3.6
2001.....	2.0	*	*	1.9	3.1	4.7	4.8	3.3
2000.....	2.1	*	1.8	2.1	4.1	3.9	3.7	3.0
1999.....	1.9	*	*	1.9	3.3	3.8	4.2	4.0

Black, female

2006 4/.....	0.3	*	*	*	*	*	*	*
2005 4/.....	0.5	*	*	*	*	*	*	*
2004 4/.....	0.4	*	*	*	*	*	*	*
2003 4/.....	0.4	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes

2006.....	5.6	*	0.3	3.3	6.0	5.4	5.2	6.2
2005.....	5.7	*	0.4	3.5	5.8	5.6	5.7	5.9
2004.....	5.7	*	0.3	3.8	6.3	5.7	5.7	5.9
2003.....	5.8	*	0.3	3.6	6.5	5.8	6.1	6.0
2002.....	5.9	*	0.4	3.6	6.7	6.4	5.6	6.6
2001.....	5.9	*	0.4	4.1	6.6	6.7	6.3	6.4
2000.....	5.9	*	0.5	4.4	7.2	6.3	6.0	6.7
1999.....	5.9	*	0.5	4.9	7.2	6.6	6.5	7.1

All races 3/, male

2006.....	10.0	*	0.5	5.8	10.6	9.2	9.2	10.4
2005.....	10.2	*	0.7	6.1	10.3	9.7	9.7	10.1
2004.....	10.0	*	0.5	6.5	11.1	9.9	9.6	9.8
2003.....	10.4	*	0.6	6.3	11.4	10.1	10.6	10.2
2002.....	10.6	*	0.6	6.4	11.9	11.1	9.6	11.3
2001.....	10.6	*	0.6	7.1	11.8	11.7	10.8	10.8
2000.....	10.5	*	0.9	7.6	12.9	11.1	10.1	11.1
1999.....	10.6	*	0.8	8.4	12.8	11.5	10.9	11.8

All races 3/, female

2006.....	1.4	*	*	0.6	1.2	1.4	1.2	1.9
2005.....	1.4	*	*	0.8	1.1	1.3	1.6	1.7
2004.....	1.5	*	*	1.0	1.1	1.3	1.7	1.9
2003.....	1.4	*	*	0.8	1.2	1.3	1.5	1.8
2002.....	1.4	*	*	0.7	1.2	1.6	1.7	2.0
2001.....	1.5	*	0.2	1.0	1.0	1.5	1.7	2.1
2000.....	1.5	*	0.2	1.1	1.3	1.5	1.9	2.3
1999.....	1.5	*	0.2	1.1	1.4	1.6	1.9	2.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by discharge of firearms (X72-X74)										
White, both sexes										
2006 4/.....	6.4	*	0.4	3.5	6.3	5.7	5.7	7.0
2005 4/.....	6.5	*	0.5	3.8	6.1	6.0	6.3	6.7
2004 4/.....	6.5	*	0.3	4.2	6.6	6.0	6.3	6.6
2003 4/.....	6.6	*	0.4	4.0	6.7	6.1	6.7	6.8
2002.....	6.7	*	0.4	3.9	7.0	6.8	6.2	7.4
2001.....	6.7	*	0.4	4.5	6.7	7.2	7.0	7.2
2000.....	6.6	*	0.6	4.6	7.6	6.6	6.6	7.5
1999.....	6.6	*	0.6	5.2	7.4	7.0	7.1	7.8
White, male										
2006 4/.....	11.3	*	0.6	6.3	11.0	9.6	9.9	11.6
2005 4/.....	11.5	*	0.8	6.6	10.7	10.3	10.5	11.2
2004 4/.....	11.3	*	0.5	7.1	11.7	10.2	10.4	11.0
2003 4/.....	11.7	*	0.7	6.9	11.8	10.5	11.3	11.3
2002.....	11.9	*	0.7	6.9	12.3	11.3	10.4	12.4
2001.....	11.8	*	0.7	7.6	11.9	12.3	11.7	11.9
2000.....	11.6	*	1.0	7.9	13.5	11.4	11.0	12.2
1999.....	11.7	*	0.9	8.9	13.0	11.9	11.8	12.7
White, female										
2006 4/.....	1.6	*	*	0.6	1.2	1.6	1.4	2.3
2005 4/.....	1.6	*	*	0.9	1.2	1.4	1.9	2.0
2004 4/.....	1.7	*	*	1.1	1.2	1.5	2.0	2.1
2003 4/.....	1.6	*	*	0.9	1.2	1.5	1.8	2.1
2002.....	1.6	*	*	0.8	1.3	2.0	1.9	2.3
2001.....	1.7	*	*	1.1	1.1	1.7	2.0	2.4
2000.....	1.7	*	*	1.1	1.3	1.6	2.0	2.7
1999.....	1.7	*	0.3	1.2	1.4	1.8	2.2	2.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Black, both sexes

2006 4/.....	2.6	*	*	2.4	5.3	5.0	4.1	3.4
2005 4/.....	2.6	*	*	2.4	5.3	4.6	4.2	3.5
2004 4/.....	2.6	*	*	2.2	5.9	5.4	4.2	2.9
2003 4/.....	2.6	*	*	2.1	6.5	5.5	4.8	2.8
2002.....	2.8	*	*	2.5	5.8	6.1	4.0	4.0
2001.....	2.9	*	*	2.8	7.1	5.8	4.1	3.7
2000.....	3.0	*	*	3.8	6.8	6.0	4.0	3.4
1999.....	3.1	*	*	3.9	7.6	5.7	4.2	4.2

Black, male

2006 4/.....	4.9	*	*	4.2	9.5	9.5	7.9	6.6
2005 4/.....	4.9	*	*	4.3	9.8	8.4	8.3	6.7
2004 4/.....	4.9	*	*	3.7	10.7	10.2	7.8	5.5
2003 4/.....	5.0	*	*	3.9	11.9	11.0	9.1	5.5
2002.....	5.4	*	*	4.3	11.0	12.4	7.7	7.4
2001.....	5.5	*	*	5.0	13.9	10.9	8.0	7.1
2000.....	5.5	*	*	6.9	12.6	11.3	7.3	6.4
1999.....	5.9	*	*	7.0	14.2	11.2	8.1	8.1

Black, female

2006 4/.....	0.4	*	*	*	*	*	*	*
2005 4/.....	0.5	*	*	*	*	*	*	*
2004 4/.....	0.5	*	*	*	*	*	*	*
2003 4/.....	0.5	*	*	*	*	*	*	*
2002.....	0.5	*	*	*	*	*	*	*
2001.....	0.6	*	*	*	*	*	*	*
2000.....	0.7	*	*	*	*	*	*	*
1999.....	0.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by jumping from a high place (X80)

All races 3/, both sexes

2006.....	0.2	*	*	0.1	0.4	0.3	0.3	0.3
2005.....	0.2	*	*	0.2	0.3	0.3	0.3	0.3
2004.....	0.2	*	*	0.2	0.3	0.3	0.3	0.3
2003.....	0.2	*	*	0.2	0.3	0.3	0.3	0.4
2002.....	0.3	*	*	0.2	0.4	0.4	0.4	0.3
2001.....	0.2	*	*	0.2	0.2	0.3	0.3	0.3
2000.....	0.2	*	*	0.2	0.4	0.3	0.3	0.3
1999.....	0.2	*	*	0.2	0.4	0.3	0.3	0.4

All races 3/, male

2006.....	0.4	*	*	0.2	0.8	0.6	0.4	0.4
2005.....	0.3	*	*	0.3	0.5	0.5	0.5	0.4
2004.....	0.3	*	*	0.3	0.5	0.5	0.5	0.4
2003.....	0.4	*	*	0.2	0.5	0.4	0.5	0.5
2002.....	0.4	*	*	0.2	0.6	0.6	0.6	0.5
2001.....	0.3	*	*	0.3	0.3	0.4	0.4	0.4
2000.....	0.3	*	*	0.3	0.5	0.4	0.3	0.5
1999.....	0.4	*	*	0.3	0.6	0.5	0.4	0.6

All races 3/, female

2006.....	0.1	*	*	*	*	*	*	0.2
2005.....	0.1	*	*	*	*	*	*	0.2
2004.....	0.1	*	*	*	*	*	*	0.2
2003.....	0.1	*	*	*	*	*	*	0.2
2002.....	0.2	*	*	*	*	*	0.3	0.2
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.2	*	*	*
1999.....	0.1	*	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by jumping from a high place (X80)

White, both sexes

2006 4/.....	0.2	*	*	0.1	0.4	0.4	0.3	0.3
2005 4/.....	0.2	*	*	0.2	0.4	0.3	0.3	0.3
2004 4/.....	0.2	*	*	0.2	0.3	0.3	0.3	0.3
2003 4/.....	0.3	*	*	0.1	0.3	0.3	0.3	0.4
2002.....	0.3	*	*	0.1	0.4	0.4	0.4	0.4
2001.....	0.2	*	*	0.2	0.3	0.2	0.3	0.3
2000.....	0.2	*	*	0.2	0.4	0.2	0.2	0.3
1999.....	0.2	*	*	0.2	0.4	0.3	0.3	0.4

White, male

2006 4/.....	0.4	*	*	0.2	0.7	0.6	0.4	0.3
2005 4/.....	0.4	*	*	0.4	0.6	0.5	0.5	0.4
2004 4/.....	0.3	*	*	0.3	0.5	0.4	0.4	0.4
2003 4/.....	0.4	*	*	*	0.5	0.4	0.5	0.5
2002.....	0.4	*	*	*	0.6	0.5	0.5	0.5
2001.....	0.3	*	*	0.3	0.4	0.3	0.4	0.4
2000.....	0.3	*	*	0.3	0.5	0.4	0.3	0.5
1999.....	0.4	*	*	0.2	0.6	0.4	0.4	0.5

White, female

2006 4/.....	0.1	*	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	0.3	*
2001.....	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	0.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by jumping from a high place (X80)

Black, both sexes

2006 4/.....	0.2	*	*	*	*	*	*	*
2005 4/.....	0.2	*	*	*	*	*	*	*
2004 4/.....	0.2	*	*	*	*	*	*	*
2003 4/.....	0.2	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*
2001.....	0.2	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*
1999.....	0.2	*	*	*	*	*	*	*

Black, male

2006 4/.....	0.3	*	*	*	*	*	*	*
2005 4/.....	0.2	*	*	*	*	*	*	*
2004 4/.....	0.3	*	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*	*
2002.....	0.3	*	*	*	*	*	*	*
2001.....	0.3	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*

Black, female

2006 4/.....	*	*	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,Y87.0)

All races 3/, both sexes

2006.....	0.7	*	*	0.4	0.7	0.7	0.8	0.9
2005.....	0.7	*	*	0.3	0.6	0.7	0.9	0.9
2004.....	0.6	*	*	0.4	0.7	0.6	0.6	0.9
2003.....	0.6	*	*	0.2	0.5	0.7	0.7	0.9
2002.....	0.6	*	*	0.3	0.6	0.7	0.7	0.9
2001 5/.....	0.6	*	*	0.3	0.6	0.7	0.7	0.9
2000.....	0.6	*	*	0.3	0.6	0.6	0.6	0.9
1999.....	0.6	*	*	0.3	0.6	0.6	0.7	0.8

All races 3/, male

2006.....	1.1	*	*	0.6	1.2	1.2	1.3	1.4
2005.....	1.0	*	*	0.6	1.0	1.0	1.3	1.4
2004.....	1.0	*	*	0.5	1.1	1.0	0.9	1.3
2003.....	0.9	*	*	0.3	0.8	1.1	1.1	1.4
2002.....	1.0	*	*	0.6	1.0	1.0	1.2	1.5
2001 5/.....	0.9	*	*	0.5	1.0	1.1	1.0	1.4
2000.....	0.9	*	*	0.4	1.0	0.9	0.9	1.4
1999.....	0.8	*	*	0.5	1.0	0.9	1.2	1.1

All races 3/, female

2006.....	0.3	*	*	0.2	*	0.3	0.4	0.5
2005.....	0.3	*	*	*	0.2	0.3	0.4	0.4
2004.....	0.3	*	*	*	0.3	0.2	0.4	0.5
2003.....	0.3	*	*	*	0.2	0.2	0.4	0.3
2002.....	0.3	*	*	*	0.3	0.4	0.3	0.4
2001 5/.....	0.3	*	*	*	0.2	0.2	0.3	0.4
2000.....	0.3	*	*	*	0.2	*	0.3	0.3
1999.....	0.3	*	*	*	0.2	*	0.3	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,Y87.0)										
White, both sexes										
2006 4/.....	0.7	*	*	0.4	0.7	0.8	0.8	1.0
2005 4/.....	0.7	*	*	0.4	0.7	0.7	0.9	0.9
2004 4/.....	0.7	*	*	0.4	0.7	0.7	0.7	0.9
2003 4/.....	0.6	*	*	0.2	0.5	0.6	0.7	0.9
2002.....	0.7	*	*	0.4	0.6	0.7	0.8	1.0
2001 5/.....	0.6	*	*	0.4	0.6	0.7	0.7	0.9
2000.....	0.6	*	*	0.3	0.6	0.6	0.6	0.9
1999.....	0.6	*	*	0.3	0.6	0.5	0.7	0.8
White, male										
2006 4/.....	1.1	*	*	0.6	1.3	1.2	1.2	1.4
2005 4/.....	1.1	*	*	0.6	1.0	1.0	1.3	1.5
2004 4/.....	1.0	*	*	0.6	1.1	1.0	1.0	1.3
2003 4/.....	1.0	*	*	0.3	0.8	1.1	1.1	1.4
2002.....	1.0	*	*	0.6	1.0	1.0	1.2	1.5
2001 5/.....	1.0	*	*	0.6	0.9	1.1	1.1	1.4
2000.....	0.9	*	*	0.4	0.9	1.0	0.9	1.4
1999.....	0.9	*	*	0.5	1.0	0.9	1.1	1.1
White, female										
2006 4/.....	0.4	*	*	*	*	0.3	0.4	0.5
2005 4/.....	0.3	*	*	*	*	0.3	0.4	0.4
2004 4/.....	0.3	*	*	*	0.3	*	0.3	0.5
2003 4/.....	0.3	*	*	*	*	*	0.3	0.4
2002.....	0.3	*	*	*	0.3	0.4	0.3	0.4
2001 5/.....	0.3	*	*	*	*	*	0.4	0.4
2000.....	0.3	*	*	*	*	*	0.3	0.3
1999.....	0.3	*	*	*	*	*	0.3	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,Y87.0)

Black, both sexes

2006 4/.....	0.4	*	*	*	*	*	0.7	0.7
2005 4/.....	0.5	*	*	*	0.7	*	0.9	*
2004 4/.....	0.5	*	*	*	0.7	*	*	1.0
2003 4/.....	0.5	*	*	*	*	0.8	1.0	0.9
2002.....	0.5	*	*	*	0.7	0.8	*	1.1
2001 5/.....	0.5	*	*	*	*	*	*	0.9
2000.....	0.5	*	*	*	1.0	*	*	0.9
1999.....	0.5	*	*	*	*	*	0.9	0.8

Black, male

2006 4/.....	0.7	*	*	*	*	*	*	*
2005 4/.....	0.8	*	*	*	1.3	*	1.5	*
2004 4/.....	0.7	*	*	*	*	*	*	1.5
2003 4/.....	0.7	*	*	*	*	*	1.5	1.6
2002.....	0.8	*	*	*	*	*	*	1.8
2001 5/.....	0.7	*	*	*	*	*	*	1.5
2000.....	0.8	*	*	*	*	*	*	1.9
1999.....	0.7	*	*	*	*	*	1.5	1.5

Black, female

2006 4/.....	0.2	*	*	*	*	*	*	*
2005 4/.....	0.2	*	*	*	*	*	*	*
2004 4/.....	0.2	*	*	*	*	*	*	*
2003 4/.....	0.2	*	*	*	*	*	*	*
2002.....	0.2	*	*	*	*	*	*	*
2001 5/.....	0.2	*	*	*	*	*	*	*
2000.....	0.2	*	*	*	*	*	*	*
1999.....	0.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
All races 3/, both sexes										
2006.....	6.2	8.1	2.2	0.8	1.2	10.7	16.2	13.5	9.8	7.6
2005.....	6.1	7.5	2.3	0.6	1.1	9.9	16.1	13.6	10.0	7.5
2004.....	5.9	8.0	2.4	0.6	1.0	9.3	15.0	13.3	9.3	7.3
2003.....	6.1	8.5	2.4	0.6	1.0	9.5	16.5	13.4	9.4	7.6
2002.....	6.1	7.5	2.7	0.7	1.0	9.3	16.5	13.3	9.4	7.9
2001 5/.....	7.1	8.2	2.7	0.7	0.9	9.4	17.3	14.9	11.5	10.4
2000.....	6.0	9.2	2.3	0.7	1.1	9.5	16.0	11.9	9.0	7.6
1999.....	6.1	8.7	2.5	0.9	1.2	10.4	15.6	12.1	9.1	7.7
All races 3/, male										
2006.....	10.0	9.4	2.5	0.8	1.6	18.2	27.5	22.4	16.2	11.4
2005.....	9.8	8.2	2.6	0.7	1.3	16.8	27.2	22.8	16.4	11.2
2004.....	9.4	8.0	2.5	0.6	1.3	15.3	25.4	22.3	14.9	11.2
2003.....	9.7	10.0	2.5	0.7	1.3	15.9	27.6	22.4	14.8	11.4
2002.....	9.6	7.9	2.9	0.6	1.2	15.3	27.7	22.0	14.7	11.8
2001 5/.....	11.1	9.5	3.0	0.7	1.1	15.7	28.9	24.1	17.7	15.3
2000.....	9.3	10.4	2.5	0.7	1.4	15.5	26.7	19.6	14.0	10.9
1999.....	9.3	9.6	2.5	0.9	1.5	16.9	25.7	19.5	13.8	11.0
All races 3/, female										
2006.....	2.5	6.8	2.0	0.7	0.7	2.9	4.2	4.2	3.3	3.6
2005.....	2.5	6.6	2.0	0.6	0.8	2.5	4.3	4.0	3.5	3.9
2004.....	2.5	7.9	2.2	0.6	0.7	3.0	4.0	3.9	3.6	3.3
2003.....	2.6	6.9	2.3	0.5	0.6	2.6	4.8	4.0	3.9	3.9
2002.....	2.7	7.1	2.5	0.8	0.8	2.9	4.6	4.2	4.1	4.1
2001 5/.....	3.3	6.9	2.4	0.6	0.7	2.7	5.1	5.5	5.2	5.5
2000.....	2.8	7.9	2.1	0.6	0.8	3.1	4.7	4.1	4.0	4.4
1999.....	2.9	7.8	2.4	0.9	0.9	3.5	5.1	4.5	4.3	4.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
White, both sexes										
2006 4/.....	3.7	6.9	1.5	0.6	0.8	5.4	7.9	6.8	5.2	4.7
2005 4/.....	3.7	6.1	1.7	0.4	0.6	5.3	7.9	7.2	5.6	4.7
2004 4/.....	3.6	6.5	1.6	0.5	0.7	5.1	7.9	7.0	5.2	4.6
2003 4/.....	3.7	6.9	1.7	0.4	0.6	5.0	8.4	6.9	5.3	5.0
2002.....	3.7	5.4	2.0	0.5	0.7	5.2	8.2	6.8	5.1	5.0
2001 5/.....	4.9	6.2	1.9	0.7	0.7	5.1	9.4	8.9	7.8	7.8
2000.....	3.6	6.6	1.7	0.6	0.9	5.2	7.7	6.2	5.3	4.9
1999.....	3.8	6.7	1.7	0.7	1.0	5.6	8.0	6.5	5.5	5.1
White, male										
2006 4/.....	5.4	7.4	1.7	0.5	1.1	8.7	12.6	10.6	7.9	6.7
2005 4/.....	5.4	6.7	1.8	0.5	0.7	8.6	12.5	11.1	8.4	6.5
2004 4/.....	5.3	6.9	1.8	0.5	0.9	7.9	12.5	10.9	7.7	6.8
2003 4/.....	5.4	8.1	1.5	0.4	0.9	8.0	13.3	10.7	7.8	6.9
2002.....	5.4	6.2	2.1	0.4	0.7	8.2	12.9	10.5	7.3	7.0
2001 5/.....	7.2	7.3	2.2	0.6	0.8	7.9	14.5	13.5	11.3	11.6
2000.....	5.2	8.2	1.9	0.7	1.1	8.0	12.0	9.2	7.6	6.6
1999.....	5.5	7.9	1.8	0.7	1.1	8.6	12.1	9.9	7.8	7.0
White, female										
2006 4/.....	1.9	6.3	1.3	0.7	0.6	1.9	2.9	2.7	2.3	2.7
2005 4/.....	1.9	5.5	1.6	0.4	0.6	1.8	2.9	3.0	2.7	2.9
2004 4/.....	1.9	6.2	1.4	0.5	0.5	2.1	3.0	2.9	2.6	2.4
2003 4/.....	2.0	5.7	1.8	0.4	0.3	1.8	3.2	2.9	2.6	3.0
2002.....	2.0	4.6	1.8	0.6	0.6	1.9	3.1	2.8	2.8	3.1
2001 5/.....	2.6	5.1	1.6	0.7	0.6	2.0	3.9	4.1	4.2	3.9
2000.....	2.1	5.0	1.5	0.5	0.7	2.1	3.2	3.0	2.9	3.2
1999.....	2.1	5.4	1.7	0.7	0.8	2.3	3.7	2.9	3.2	3.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
Black, both sexes										
2006 4/.....	22.8	16.0	5.8	1.6	2.9	37.5	62.4	54.0	39.7	26.9
2005 4/.....	22.2	14.3	5.3	1.4	3.0	33.5	61.2	53.0	39.2	26.5
2004 4/.....	21.1	15.6	6.3	1.4	2.4	31.6	55.9	52.8	35.6	25.2
2003 4/.....	22.0	16.1	6.1	1.7	2.7	33.2	62.4	54.5	36.3	25.4
2002.....	22.0	17.3	6.3	1.4	2.6	31.0	63.4	54.2	37.0	26.9
2001 5/.....	22.1	18.9	6.8	0.9	1.9	32.7	62.8	52.6	35.3	28.2
2000.....	21.5	22.8	5.8	1.2	2.5	33.1	64.1	48.2	33.5	25.9
1999.....	21.1	19.5	6.1	1.9	2.5	36.5	60.4	47.0	31.8	25.1
Black, male										
2006 4/.....	40.6	20.7	6.3	2.1	4.1	66.3	112.2	97.3	73.6	46.3
2005 4/.....	39.7	15.9	6.6	1.6	4.1	60.0	110.1	98.6	72.8	44.7
2004 4/.....	37.1	14.2	6.3	1.4	3.3	54.8	101.8	98.4	64.7	43.1
2003 4/.....	38.9	17.8	7.4	2.0	3.7	58.9	111.6	101.6	63.9	43.7
2002.....	38.4	16.3	7.1	1.4	3.5	53.3	115.3	99.9	65.4	45.2
2001 5/.....	38.3	21.0	8.0	1.2	2.7	58.2	116.3	96.0	62.3	42.8
2000.....	37.2	23.3	5.9	*	3.3	57.2	117.8	89.4	58.6	42.5
1999.....	36.1	19.5	5.9	2.0	3.3	62.4	110.9	84.0	54.7	40.1
Black, female										
2006 4/.....	6.6	11.1	5.4	*	1.7	7.9	11.3	12.9	9.0	9.6
2005 4/.....	6.2	12.6	4.0	1.3	1.9	6.3	11.5	9.9	8.8	10.1
2004 4/.....	6.4	17.0	6.3	1.4	1.5	7.8	9.8	10.2	9.3	9.2
2003 4/.....	6.6	14.2	4.8	1.3	1.7	6.7	13.6	10.9	11.4	9.0
2002.....	7.0	18.5	5.5	1.3	1.6	8.0	12.7	12.3	11.5	10.6
2001 5/.....	7.4	16.7	5.6	*	*	6.4	11.4	13.1	11.1	15.3
2000.....	7.2	22.2	5.7	1.4	1.7	8.4	13.2	10.9	10.9	11.1
1999.....	7.6	19.5	6.3	1.8	1.6	10.0	12.5	13.6	11.2	11.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Assault (homicide) by hanging, strangulation and suffocation (X91)

All races 3/, both sexes

2006.....	0.2	0.8	0.1	*	*	0.2	0.3	0.3	0.2	0.3
2005.....	0.2	0.7	0.1	*	0.1	0.2	0.3	0.3	0.2	0.3
2004.....	0.2	1.0	0.2	*	*	0.1	0.4	0.3	0.3	0.3
2003.....	0.2	1.0	*	*	*	0.2	0.3	0.3	0.3	0.4
2002.....	0.2	0.8	0.2	*	*	0.2	0.3	0.3	0.3	0.4
2001.....	0.2	1.0	*	*	*	0.2	0.4	0.3	0.3	0.3
2000.....	0.2	0.7	0.2	*	*	0.2	0.4	0.3	0.4	0.3
1999.....	0.3	0.9	0.1	*	*	0.2	0.4	0.4	0.4	0.4

All races 3/, male

2006.....	0.2	1.0	*	*	*	*	*	*	*	0.2
2005.....	0.1	*	*	*	*	*	*	0.2	*	*
2004.....	0.1	*	*	*	*	*	0.2	*	0.2	*
2003.....	0.2	1.0	*	*	*	*	*	*	*	0.3
2002.....	0.2	*	*	*	*	*	0.2	0.2	*	0.2
2001.....	0.2	1.2	*	*	*	*	0.2	*	*	0.2
2000.....	0.2	*	*	*	*	*	*	*	0.3	*
1999.....	0.2	1.1	*	*	*	*	*	0.2	0.3	0.2

All races 3/, female

2006.....	0.3	*	*	*	*	0.3	0.4	0.4	0.3	0.3
2005.....	0.3	*	*	*	*	0.3	0.5	0.5	0.4	0.4
2004.....	0.3	1.3	*	*	*	0.2	0.5	0.6	0.4	0.4
2003.....	0.3	*	*	*	*	0.4	0.5	0.5	0.5	0.4
2002.....	0.3	*	*	*	*	0.3	0.5	0.5	0.4	0.6
2001.....	0.3	*	*	*	*	0.3	0.6	0.5	0.5	0.4
2000.....	0.3	*	*	*	*	0.3	0.6	0.5	0.4	0.4
1999.....	0.3	*	*	*	*	0.4	0.6	0.6	0.6	0.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by hanging, strangulation and suffocation (X91)										
White, both sexes										
2006 4/.....	0.2	0.6	*	*	*	*	0.2	0.2	0.2	0.2
2005 4/.....	0.2	*	*	*	*	0.2	0.3	0.3	0.2	0.2
2004 4/.....	0.2	0.9	*	*	*	*	0.3	0.3	0.2	0.2
2003 4/.....	0.2	0.9	*	*	*	0.2	0.3	0.3	0.3	0.3
2002.....	0.2	0.6	*	*	*	0.1	0.3	0.3	0.2	0.3
2001.....	0.2	0.7	*	*	*	0.2	0.3	0.3	0.3	0.3
2000.....	0.2	*	0.2	*	*	0.2	0.3	0.3	0.2	0.2
1999.....	0.2	0.7	*	*	*	0.2	0.3	0.3	0.3	0.3
White, male										
2006 4/.....	0.1	*	*	*	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*	*	*	0.2
2002.....	0.1	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*
White, female										
2006 4/.....	0.2	*	*	*	*	*	0.4	0.3	0.3	*
2005 4/.....	0.2	*	*	*	*	0.3	0.4	0.4	0.3	0.3
2004 4/.....	0.2	1.4	*	*	*	*	0.4	0.5	0.3	0.3
2003 4/.....	0.2	*	*	*	*	0.3	0.4	0.5	0.3	0.3
2002.....	0.2	*	*	*	*	*	0.5	0.3	0.3	0.4
2001.....	0.2	*	*	*	*	0.3	0.4	0.4	0.4	0.3
2000.....	0.2	*	*	*	*	*	0.5	0.4	0.3	0.3
1999.....	0.2	*	*	*	*	0.3	0.5	0.4	0.4	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by hanging, strangulation and suffocation (X91)

Black, both sexes

2006 4/.....	0.5	*	*	*	*	*	*	0.8	*	0.7
2005 4/.....	0.5	*	*	*	*	*	0.7	*	0.9	0.8
2004 4/.....	0.5	*	*	*	*	*	0.7	0.8	*	1.0
2003 4/.....	0.5	*	*	*	*	0.7	0.7	*	*	0.9
2002.....	0.6	*	*	*	*	*	0.7	0.9	0.9	1.1
2001.....	0.6	*	*	*	*	*	1.0	*	0.7	0.9
2000.....	0.6	*	*	*	*	*	1.0	0.9	1.0	0.8
1999.....	0.6	*	*	*	*	*	0.8	1.1	1.1	1.1

Black, male

2006 4/.....	0.4	*	*	*	*	*	*	*	*	*
2005 4/.....	0.4	*	*	*	*	*	*	*	*	*
2004 4/.....	0.3	*	*	*	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*
2000.....	0.3	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	0.6	*	*	*	*	*	*	*	*	*
2005 4/.....	0.6	*	*	*	*	*	*	*	*	*
2004 4/.....	0.7	*	*	*	*	*	*	*	*	*
2003 4/.....	0.7	*	*	*	*	*	*	*	*	*
2002.....	0.7	*	*	*	*	*	*	*	*	1.6
2001.....	0.7	*	*	*	*	*	*	*	*	1.4
2000.....	0.8	*	*	*	*	*	1.4	1.4	1.4	*
1999.....	0.8	*	*	*	*	*	*	1.6	1.6	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All races 3/, both sexes

2006.....	4.3	*	0.3	0.3	0.8	9.1	13.7	10.8	7.7	5.2
2005.....	4.2	*	0.2	0.2	0.7	8.3	13.1	11.0	7.8	5.1
2004.....	4.0	*	0.2	0.2	0.7	7.6	12.2	10.5	7.0	4.9
2003.....	4.1	*	0.3	0.2	0.7	7.7	13.6	10.8	7.1	5.0
2002.....	4.1	*	0.3	0.3	0.7	7.7	13.6	10.6	6.9	5.2
2001.....	4.0	*	0.4	0.3	0.6	7.5	13.6	10.3	6.6	5.1
2000.....	3.8	*	0.2	0.2	0.7	7.7	12.7	9.2	6.3	4.7
1999.....	3.9	*	0.3	0.3	0.8	8.5	12.5	9.2	6.1	4.6

All races 3/, male

2006.....	7.4	*	0.3	0.3	1.3	16.0	24.1	18.7	13.4	8.4
2005.....	7.2	*	0.3	0.3	1.0	14.7	23.1	19.3	13.5	8.1
2004.....	6.9	*	*	0.2	1.0	13.1	21.5	18.8	11.9	8.3
2003.....	7.1	*	0.2	0.3	1.0	13.7	23.8	19.1	12.0	8.0
2002.....	7.0	*	0.4	0.3	0.9	13.2	23.8	18.6	11.6	8.4
2001.....	6.8	*	0.3	0.3	0.8	13.2	24.0	17.8	11.1	8.0
2000.....	6.5	*	*	0.3	1.0	13.3	22.4	16.1	10.4	7.4
1999.....	6.5	*	0.3	0.4	1.1	14.5	21.8	15.9	10.1	7.1

All races 3/, female

2006.....	1.3	*	*	0.3	0.4	1.8	2.6	2.5	1.9	2.0
2005.....	1.2	*	*	*	0.4	1.5	2.4	2.3	2.1	2.1
2004.....	1.1	*	*	0.2	0.3	1.8	2.2	2.0	2.0	1.4
2003.....	1.2	*	0.3	0.2	0.3	1.5	2.8	2.2	2.1	1.9
2002.....	1.3	*	*	0.3	0.5	1.8	2.9	2.4	2.1	2.0
2001.....	1.3	*	0.4	0.2	0.3	1.5	2.7	2.5	2.1	2.2
2000.....	1.3	*	*	0.2	0.3	1.7	2.7	2.2	2.1	1.9
1999.....	1.3	*	0.3	0.2	0.5	2.2	2.8	2.3	2.1	2.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
White, both sexes										
2006 4/.....	2.2	*	0.2	0.3	0.6	4.2	5.9	4.8	3.6	3.0
2005 4/.....	2.2	*	0.2	0.2	0.4	4.1	5.5	5.1	4.0	3.0
2004 4/.....	2.1	*	*	0.2	0.4	3.8	5.5	4.9	3.7	2.9
2003 4/.....	2.2	*	0.2	0.1	0.4	3.8	6.3	4.9	3.6	3.0
2002.....	2.2	*	0.3	0.2	0.4	4.1	6.0	4.9	3.4	3.0
2001.....	2.2	*	0.3	0.3	0.4	3.7	6.3	5.0	3.5	3.2
2000.....	2.1	*	*	0.2	0.5	3.9	5.5	4.2	3.3	2.8
1999.....	2.2	*	0.2	0.2	0.6	4.2	5.6	4.3	3.4	3.0
White, male										
2006 4/.....	3.5	*	*	*	0.8	7.2	9.9	7.9	5.9	4.4
2005 4/.....	3.5	*	*	*	0.5	7.0	9.3	8.4	6.3	4.2
2004 4/.....	3.5	*	*	*	0.6	6.3	9.3	8.3	5.8	4.7
2003 4/.....	3.5	*	*	*	0.6	6.4	10.6	8.2	5.7	4.4
2002.....	3.5	*	0.3	*	0.5	6.7	10.0	8.1	5.3	4.5
2001.....	3.5	*	*	0.3	0.6	6.2	10.5	8.2	5.4	4.9
2000.....	3.2	*	*	0.3	0.7	6.5	9.0	6.7	5.1	4.1
1999.....	3.4	*	*	0.2	0.8	6.9	9.1	7.1	5.2	4.3
White, female										
2006 4/.....	0.9	*	*	0.3	0.3	1.1	1.5	1.5	1.3	1.5
2005 4/.....	0.9	*	*	*	*	1.0	1.4	1.6	1.5	1.7
2004 4/.....	0.9	*	*	*	*	1.2	1.5	1.2	1.4	1.0
2003 4/.....	0.9	*	*	*	*	1.0	1.7	1.4	1.4	1.6
2002.....	1.0	*	*	*	0.3	1.2	1.6	1.6	1.4	1.5
2001.....	1.0	*	*	*	0.3	1.1	1.9	1.6	1.6	1.6
2000.....	1.0	*	*	*	*	1.1	1.7	1.6	1.5	1.5
1999.....	0.9	*	*	*	0.4	1.2	1.9	1.4	1.5	1.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by discharge of firearms (*U01.4,X93-X95)										
Black, both sexes										
2006 4/.....	18.0	*	0.8	*	2.4	33.5	57.1	47.4	34.4	20.6
2005 4/.....	17.2	*	*	*	2.1	30.0	55.1	46.9	33.5	19.4
2004 4/.....	16.1	*	0.8	*	1.8	27.8	50.4	46.3	29.3	18.5
2003 4/.....	16.8	*	*	0.7	2.0	29.0	55.3	47.9	30.1	18.1
2002.....	16.7	*	*	*	1.9	27.3	57.0	46.9	30.1	19.5
2001.....	15.8	*	*	*	1.3	28.7	55.7	43.6	26.9	18.4
2000.....	15.6	*	*	*	1.7	28.6	55.3	40.7	26.0	17.3
1999.....	15.4	*	*	0.7	1.8	32.0	53.4	39.3	23.7	15.9
Black, male										
2006 4/.....	33.7	*	*	*	3.7	60.8	104.4	87.9	65.8	37.7
2005 4/.....	32.5	*	*	*	3.1	54.9	101.6	89.5	64.2	35.7
2004 4/.....	30.3	*	*	*	2.9	49.8	93.9	88.9	55.2	34.6
2003 4/.....	31.6	*	*	*	2.9	53.2	101.7	92.4	56.4	33.9
2002.....	31.0	*	*	*	2.7	48.4	105.9	90.2	56.3	35.7
2001.....	29.8	*	*	*	2.1	52.8	106.5	83.2	51.3	31.5
2000.....	29.2	*	*	*	2.8	51.7	105.2	78.7	48.1	31.7
1999.....	28.4	*	*	*	2.8	56.3	101.0	74.4	44.0	27.3
Black, female										
2006 4/.....	3.6	*	*	*	*	5.5	8.6	9.0	5.9	5.4
2005 4/.....	3.1	*	*	*	*	4.4	7.6	6.6	5.8	4.7
2004 4/.....	3.0	*	*	*	*	5.1	6.6	6.4	5.9	4.0
2003 4/.....	3.3	*	*	*	1.1	4.0	9.1	6.8	6.3	4.1
2002.....	3.6	*	*	*	1.2	5.4	9.2	7.3	6.5	5.0
2001.....	3.1	*	*	*	*	3.9	7.0	7.6	5.0	6.8
2000.....	3.2	*	*	*	*	4.9	8.0	6.2	6.0	4.5
1999.....	3.7	*	*	*	*	7.0	8.2	7.5	5.5	5.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by sharp object (X99)										
All races 3/, both sexes										
2006.....	0.7	*	*	*	*	1.0	1.3	1.4	0.9	1.0
2005.....	0.7	*	*	*	*	0.8	1.7	1.3	1.0	1.0
2004.....	0.7	*	*	*	0.1	0.8	1.5	1.3	1.1	1.0
2003.....	0.7	*	*	*	*	0.8	1.5	1.3	1.1	1.1
2002.....	0.7	*	*	0.1	*	0.7	1.4	1.3	1.2	1.1
2001.....	0.7	*	*	*	*	0.8	1.6	1.4	1.0	1.1
2000.....	0.6	*	*	*	0.1	0.8	1.3	1.1	1.0	1.1
1999.....	0.7	*	*	*	0.1	0.8	1.4	1.1	1.2	1.1
All races 3/, male										
2006.....	1.0	*	*	*	*	1.5	1.9	2.0	1.5	1.5
2005.....	1.0	*	*	*	*	1.2	2.6	2.0	1.5	1.3
2004.....	1.0	*	*	*	*	1.3	2.3	1.9	1.5	1.4
2003.....	1.0	*	*	*	*	1.3	2.1	1.8	1.5	1.6
2002.....	1.0	*	*	*	*	1.1	2.2	1.9	1.5	1.6
2001.....	1.0	*	*	*	*	1.3	2.4	2.0	1.4	1.4
2000.....	0.9	*	*	*	*	1.1	2.1	1.6	1.5	1.4
1999.....	1.0	*	*	*	*	1.2	2.0	1.6	1.7	1.6
All races 3/, female										
2006.....	0.4	*	*	*	*	0.4	0.6	0.7	0.4	0.6
2005.....	0.4	*	*	*	*	0.4	0.7	0.7	0.5	0.7
2004.....	0.4	*	*	*	*	0.4	0.6	0.8	0.6	0.7
2003.....	0.4	*	*	*	*	0.3	0.7	0.7	0.7	0.6
2002.....	0.4	*	*	*	*	0.3	0.5	0.8	0.9	0.7
2001.....	0.4	*	*	*	*	0.3	0.8	0.7	0.6	0.8
2000.....	0.4	*	*	*	*	0.4	0.5	0.5	0.6	0.7
1999.....	0.4	*	*	*	*	0.4	0.7	0.6	0.7	0.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by sharp object (X99)										
White, both sexes										
2006 4/.....	0.5	*	*	*	*	0.6	1.0	1.0	0.7	0.7
2005 4/.....	0.5	*	*	*	*	0.6	1.2	1.0	0.7	0.6
2004 4/.....	0.5	*	*	*	*	0.6	1.2	1.0	0.7	0.7
2003 4/.....	0.5	*	*	*	*	0.5	0.9	0.9	0.8	0.7
2002.....	0.5	*	*	*	*	0.4	1.0	0.9	0.7	0.8
2001.....	0.5	*	*	*	*	0.5	1.2	1.0	0.7	0.7
2000.....	0.4	*	*	*	*	0.6	0.9	0.8	0.8	0.7
1999.....	0.5	*	*	*	*	0.5	1.0	0.9	0.8	0.6
White, male										
2006 4/.....	0.7	*	*	*	*	0.9	1.5	1.4	1.0	1.0
2005 4/.....	0.7	*	*	*	*	0.9	1.9	1.5	1.0	0.8
2004 4/.....	0.7	*	*	*	*	0.9	1.8	1.3	0.8	1.0
2003 4/.....	0.7	*	*	*	*	0.8	1.3	1.3	1.1	1.0
2002.....	0.7	*	*	*	*	0.7	1.6	1.3	0.9	1.2
2001.....	0.7	*	*	*	*	0.8	1.9	1.4	0.9	0.9
2000.....	0.6	*	*	*	*	0.9	1.5	1.2	1.0	1.0
1999.....	0.7	*	*	*	*	0.8	1.4	1.3	1.1	0.9
White, female										
2006 4/.....	0.3	*	*	*	*	0.3	0.5	0.5	0.3	0.4
2005 4/.....	0.3	*	*	*	*	0.3	0.5	0.5	0.5	0.4
2004 4/.....	0.3	*	*	*	*	*	0.5	0.7	0.5	0.5
2003 4/.....	0.3	*	*	*	*	*	0.5	0.5	0.4	0.4
2002.....	0.3	*	*	*	*	*	0.4	0.5	0.6	0.4
2001.....	0.3	*	*	*	*	*	0.5	0.5	0.4	0.5
2000.....	0.3	*	*	*	*	0.3	0.3	0.4	0.5	0.5
1999.....	0.3	*	*	*	*	0.3	0.5	0.4	0.5	0.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Assault (homicide) by sharp object (X99)										
Black, both sexes										
2006 4/.....	2.0	*	*	*	*	2.6	2.9	3.7	2.8	3.1
2005 4/.....	2.1	*	*	*	*	2.0	4.0	3.4	2.9	3.5
2004 4/.....	2.1	*	*	*	*	2.1	3.3	3.4	3.3	3.0
2003 4/.....	2.2	*	*	*	*	2.4	4.2	3.5	3.1	3.7
2002.....	2.2	*	*	*	*	2.0	3.3	4.2	3.4	3.2
2001.....	2.0	*	*	*	*	2.2	3.7	4.1	2.8	3.5
2000.....	1.9	*	*	*	*	1.6	3.5	2.9	2.7	3.4
1999.....	2.0	*	*	*	*	2.0	3.7	2.6	3.6	3.9
Black, male										
2006 4/.....	3.1	*	*	*	*	4.0	4.5	5.6	4.6	4.9
2005 4/.....	3.4	*	*	*	*	3.0	6.1	5.5	5.1	4.7
2004 4/.....	3.2	*	*	*	*	2.9	5.1	5.4	5.4	4.0
2003 4/.....	3.4	*	*	*	*	3.7	6.5	5.6	4.5	5.4
2002.....	3.2	*	*	*	*	3.0	5.4	6.3	4.9	4.1
2001.....	3.0	*	*	*	*	3.3	5.4	6.3	4.2	4.7
2000.....	2.8	*	*	*	*	2.0	5.7	4.5	4.4	4.6
1999.....	3.1	*	*	*	*	3.1	5.3	3.6	5.5	6.4
Black, female										
2006 4/.....	1.0	*	*	*	*	*	1.3	1.8	*	1.4
2005 4/.....	1.0	*	*	*	*	*	1.9	1.4	*	2.5
2004 4/.....	1.1	*	*	*	*	1.3	1.4	1.5	1.4	2.2
2003 4/.....	1.1	*	*	*	*	*	2.0	1.6	1.9	2.1
2002.....	1.2	*	*	*	*	*	1.3	2.2	2.0	2.3
2001.....	1.1	*	*	*	*	*	2.0	2.1	1.6	2.5
2000.....	1.1	*	*	*	*	*	1.5	1.4	*	2.3
1999.....	1.0	*	*	*	*	*	2.2	1.7	1.9	1.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Neglect, abandonment and other maltreatment syndromes (Y06-Y07)

All races 3/, both sexes

2006.....	0.1	1.8	0.3	*	*	*	*	*	*	*
2005.....	0.1	2.4	0.5	*	*	*	*	*	*	*
2004.....	0.1	2.5	0.4	*	*	*	*	*	*	*
2003.....	0.1	2.5	0.5	*	*	*	*	*	*	*
2002.....	0.1	2.4	0.6	*	*	*	*	*	*	*
2001.....	0.1	2.9	0.5	*	*	*	*	*	*	*
2000.....	0.1	3.4	0.5	*	*	*	*	*	*	*
1999.....	0.1	3.3	0.4	*	*	*	*	*	*	*

All races 3/, male

2006.....	0.1	2.1	0.3	*	*	*	*	*	*	*
2005.....	0.1	2.5	0.4	*	*	*	*	*	*	*
2004.....	0.1	2.9	0.5	*	*	*	*	*	*	*
2003.....	0.1	3.3	0.5	*	*	*	*	*	*	*
2002.....	0.1	2.5	0.6	*	*	*	*	*	*	*
2001.....	0.1	3.2	0.7	*	*	*	*	*	*	*
2000.....	0.1	4.0	0.5	*	*	*	*	*	*	*
1999.....	0.1	3.7	0.4	*	*	*	*	*	*	*

All races 3/, female

2006.....	0.1	1.5	0.3	*	*	*	*	*	*	*
2005.....	0.1	2.3	0.5	*	*	*	*	*	*	*
2004.....	0.1	2.0	0.4	*	*	*	*	*	*	*
2003.....	0.1	1.7	0.6	*	*	*	*	*	*	*
2002.....	0.1	2.4	0.6	*	*	*	*	*	*	*
2001.....	0.1	2.5	0.4	*	*	*	*	*	*	*
2000.....	0.1	2.7	0.4	*	*	*	*	*	*	*
1999.....	0.1	2.9	0.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

All races 3/, both sexes

2006.....	0.9	4.8	1.3	0.2	0.1	0.5	0.9	0.9	0.7	0.9
2005.....	0.9	3.8	1.3	0.2	0.2	0.5	0.9	0.9	0.9	1.1
2004.....	0.9	3.8	1.3	0.2	0.1	0.6	0.9	1.0	0.8	1.0
2003.....	0.9	4.4	1.3	0.2	0.1	0.5	1.0	0.9	0.8	1.1
2002.....	0.9	3.7	1.4	0.2	0.1	0.6	0.9	0.8	0.9	1.0
2001 5/.....	2.0	3.5	1.6	0.2	0.2	0.7	1.5	2.8	3.4	3.6
2000.....	1.0	4.3	1.3	0.3	0.2	0.7	1.3	1.1	1.2	1.4
1999.....	1.0	3.8	1.3	0.4	0.2	0.8	1.1	1.1	1.1	1.3

All races 3/, male

2006.....	1.2	5.6	1.5	0.2	*	0.6	1.2	1.3	1.0	1.2
2005.....	1.2	4.1	1.5	0.2	*	0.7	1.1	1.2	1.2	1.6
2004.....	1.2	3.8	1.3	0.2	*	0.7	1.3	1.4	1.2	1.3
2003.....	1.2	4.7	1.5	0.2	*	0.8	1.3	1.2	1.0	1.4
2002.....	1.2	4.0	1.6	*	*	0.7	1.2	1.1	1.2	1.3
2001 5/.....	2.9	4.0	1.7	0.3	*	0.9	2.0	3.8	4.9	5.5
2000.....	1.4	4.7	1.5	0.3	0.2	0.9	1.8	1.6	1.5	1.7
1999.....	1.3	4.0	1.4	0.3	0.2	1.0	1.4	1.4	1.4	1.8

All races 3/, female

2006.....	0.5	4.0	1.2	0.2	*	0.3	0.5	0.5	0.5	0.6
2005.....	0.5	3.5	1.1	*	*	0.3	0.6	0.5	0.5	0.7
2004.....	0.6	3.8	1.3	*	*	0.6	0.5	0.5	0.5	0.7
2003.....	0.6	4.0	1.2	*	*	0.3	0.7	0.6	0.6	0.8
2002.....	0.5	3.3	1.2	*	*	0.4	0.6	0.5	0.6	0.7
2001 5/.....	1.1	3.1	1.4	*	0.2	0.5	1.0	1.7	1.9	1.8
2000.....	0.7	3.9	1.1	0.3	0.3	0.6	0.8	0.7	0.8	1.1
1999.....	0.7	3.6	1.2	0.4	*	0.6	0.7	0.8	0.9	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

White, both sexes

2006 4/.....	0.7	4.2	0.9	0.2	*	0.4	0.8	0.7	0.6	0.7
2005 4/.....	0.7	3.0	1.0	*	*	0.4	0.7	0.7	0.7	0.9
2004 4/.....	0.7	3.1	0.9	0.2	*	0.5	0.8	0.8	0.6	0.7
2003 4/.....	0.7	3.5	0.9	*	*	0.5	0.8	0.7	0.6	0.9
2002.....	0.7	2.5	0.9	*	*	0.4	0.7	0.6	0.7	0.8
2001 5/.....	1.8	2.7	1.1	0.2	0.2	0.6	1.4	2.6	3.3	3.4
2000.....	0.7	3.0	0.8	0.2	0.2	0.5	0.9	0.8	0.8	1.0
1999.....	0.8	2.9	1.0	0.2	0.2	0.6	0.9	0.8	0.8	1.0

White, male

2006 4/.....	1.0	4.7	1.0	*	*	0.5	1.0	1.0	0.8	1.0
2005 4/.....	0.9	3.2	1.1	*	*	0.5	0.9	1.1	1.0	1.3
2004 4/.....	0.9	3.5	1.0	*	*	0.6	1.1	1.0	0.8	0.9
2003 4/.....	0.9	3.8	0.9	*	*	0.6	1.1	0.8	0.8	1.1
2002.....	0.9	2.9	1.1	*	*	0.5	0.9	0.9	0.9	1.0
2001 5/.....	2.6	3.1	1.2	*	*	0.7	1.8	3.5	4.7	5.5
2000.....	0.9	3.5	1.0	*	*	0.5	1.2	1.0	1.0	1.2
1999.....	1.0	3.1	1.1	*	*	0.7	1.2	1.0	1.0	1.3

White, female

2006 4/.....	0.4	3.7	0.8	*	*	0.3	0.5	0.4	0.4	0.4
2005 4/.....	0.4	2.8	0.8	*	*	*	0.5	0.4	0.4	0.5
2004 4/.....	0.5	2.6	0.8	*	*	0.4	0.5	0.5	0.4	0.5
2003 4/.....	0.5	3.1	0.9	*	*	0.3	0.6	0.5	0.4	0.6
2002.....	0.4	2.2	0.8	*	*	0.4	0.5	0.4	0.4	0.6
2001 5/.....	1.0	2.3	0.9	*	*	0.4	0.9	1.6	1.7	1.3
2000.....	0.5	2.4	0.7	*	*	0.5	0.6	0.6	0.6	0.7
1999.....	0.5	2.5	0.9	*	*	0.5	0.7	0.6	0.7	0.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

Black, both sexes

2006 4/.....	2.0	8.7	3.4	0.6	*	0.9	1.6	1.9	1.7	2.3
2005 4/.....	2.0	7.7	2.9	*	*	1.1	1.4	1.8	1.7	2.6
2004 4/.....	2.0	7.7	3.7	*	*	1.3	1.5	2.2	2.2	2.5
2003 4/.....	2.1	8.8	3.6	0.6	*	0.9	2.1	2.1	2.2	2.5
2002.....	2.1	9.2	3.8	*	*	1.0	2.0	1.9	2.3	2.6
2001 5/.....	3.1	8.0	4.1	*	*	1.3	2.2	3.8	4.5	5.1
2000.....	2.9	11.2	3.8	0.7	*	2.2	3.9	3.3	3.3	4.0
1999.....	2.5	8.9	3.2	0.9	*	1.7	2.0	3.1	3.0	3.6

Black, male

2006 4/.....	2.9	10.7	3.7	*	*	1.3	2.5	3.0	2.3	2.8
2005 4/.....	3.0	8.3	3.5	*	*	1.7	1.9	2.5	2.8	3.4
2004 4/.....	2.8	5.9	3.4	*	*	1.4	2.3	3.5	3.6	3.7
2003 4/.....	3.1	8.6	4.5	*	*	1.3	2.7	3.3	2.5	3.5
2002.....	3.1	9.3	4.3	*	*	1.5	2.9	2.5	3.2	3.7
2001 5/.....	4.3	8.4	4.8	*	*	1.6	3.4	5.7	5.9	6.1
2000.....	4.2	11.2	3.9	*	*	3.0	5.7	5.2	4.9	5.3
1999.....	3.4	9.1	3.6	*	*	2.5	3.3	4.2	3.9	4.7

Black, female

2006 4/.....	1.1	6.6	3.2	*	*	*	*	*	*	1.8
2005 4/.....	1.2	7.1	2.3	*	*	*	*	*	*	1.8
2004 4/.....	1.3	9.6	4.0	*	*	*	*	*	*	1.5
2003 4/.....	1.2	9.0	2.6	*	*	*	1.4	*	1.8	1.7
2002.....	1.2	9.1	3.3	*	*	*	*	*	1.6	1.6
2001 5/.....	2.1	7.6	3.3	*	*	*	*	2.0	3.1	4.3
2000.....	1.7	11.2	3.7	*	*	1.3	2.2	1.7	1.9	2.8
1999.....	1.6	8.7	2.9	*	*	*	*	2.1	2.1	2.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2006.....	1.7	2.2	0.3	0.1	0.2	0.8	2.0	2.0	2.1	2.6
2005.....	1.6	2.5	0.4	0.1	0.2	0.7	1.6	1.8	1.9	2.3
2004.....	1.7	2.6	0.3	0.1	0.2	0.7	1.7	1.9	2.1	2.9
2003.....	1.7	1.9	0.2	*	0.2	0.8	1.9	1.8	2.3	3.2
2002.....	1.7	2.5	0.4	0.1	0.2	0.7	1.7	1.9	2.3	3.3
2001.....	1.5	2.1	0.3	*	0.2	0.7	1.4	1.6	2.0	3.0
2000.....	1.4	1.7	0.3	*	0.2	0.6	1.3	1.6	1.8	2.6
1999.....	1.4	1.4	0.3	*	0.2	0.8	1.5	1.6	2.1	2.8
All races 3/, male										
2006.....	2.2	2.7	0.3	*	0.3	1.2	2.9	2.8	2.7	3.2
2005.....	2.0	3.2	0.5	*	0.3	1.0	2.3	2.4	2.3	2.7
2004.....	2.2	3.1	0.3	*	0.2	1.1	2.4	2.6	2.9	3.8
2003.....	2.3	2.4	*	*	0.3	1.2	2.7	2.5	3.2	4.2
2002.....	2.2	2.5	0.4	*	0.2	1.0	2.4	2.7	3.1	4.5
2001.....	2.0	2.4	*	*	0.3	1.0	2.1	2.4	2.7	3.9
2000.....	1.9	1.4	0.3	*	0.4	0.9	2.0	2.3	2.6	3.7
1999.....	2.0	1.7	0.3	*	0.3	1.3	2.2	2.3	2.8	3.9
All races 3/, female										
2006.....	1.2	1.7	0.3	*	*	0.5	1.0	1.1	1.4	2.0
2005.....	1.3	1.8	0.3	*	*	0.4	0.9	1.2	1.5	1.9
2004.....	1.2	2.0	0.3	*	*	0.4	0.8	1.1	1.4	1.9
2003.....	1.2	1.3	*	*	*	0.5	1.1	1.1	1.5	2.2
2002.....	1.2	2.5	0.4	*	*	0.4	1.0	1.0	1.4	2.1
2001.....	0.9	1.7	0.3	*	*	0.3	0.7	0.9	1.3	2.1
2000.....	0.8	2.0	*	*	*	0.3	0.6	0.9	1.1	1.6
1999.....	0.9	1.2	0.3	*	*	0.3	0.6	0.9	1.3	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
White, both sexes										
2006 4/.....	1.7	2.0	0.2	*	0.2	1.0	2.2	2.2	2.2	2.7
2005 4/.....	1.6	2.1	0.3	*	0.2	0.7	1.7	2.0	2.0	2.4
2004 4/.....	1.7	2.1	0.2	*	0.2	0.8	1.9	2.1	2.2	3.0
2003 4/.....	1.8	1.7	0.2	*	0.2	0.9	2.1	2.0	2.6	3.4
2002.....	1.7	1.8	0.4	*	0.2	0.7	1.9	1.9	2.3	3.4
2001.....	1.5	1.4	0.3	*	0.2	0.7	1.5	1.7	2.1	3.0
2000.....	1.3	1.4	0.2	*	0.2	0.6	1.3	1.6	1.8	2.7
1999.....	1.4	1.2	0.2	*	0.2	0.8	1.5	1.6	2.1	2.8
White, male										
2006 4/.....	2.2	2.5	*	*	0.3	1.4	3.2	3.0	2.9	3.2
2005 4/.....	2.0	2.8	0.4	*	0.3	1.0	2.5	2.5	2.4	2.8
2004 4/.....	2.2	2.7	*	*	*	1.1	2.7	2.9	2.9	3.9
2003 4/.....	2.3	1.9	*	*	0.3	1.3	2.9	2.7	3.3	4.4
2002.....	2.2	1.8	0.4	*	0.2	1.1	2.6	2.7	3.1	4.5
2001.....	2.0	1.7	*	*	0.3	1.1	2.2	2.5	2.9	3.8
2000.....	1.8	*	*	*	0.3	1.0	1.9	2.3	2.5	3.7
1999.....	1.9	1.4	*	*	0.3	1.3	2.3	2.3	2.8	3.7
White, female										
2006 4/.....	1.3	1.4	*	*	*	0.5	1.1	1.3	1.5	2.2
2005 4/.....	1.3	1.5	*	*	*	0.4	0.9	1.4	1.6	2.0
2004 4/.....	1.3	1.5	*	*	*	0.4	0.9	1.2	1.5	2.1
2003 4/.....	1.3	1.4	*	*	*	0.5	1.2	1.2	1.7	2.3
2002.....	1.2	1.8	0.4	*	*	0.4	1.1	1.1	1.5	2.3
2001.....	1.0	*	*	*	*	0.3	0.8	0.9	1.4	2.3
2000.....	0.8	1.6	*	*	*	0.3	0.7	0.9	1.1	1.6
1999.....	0.9	*	*	*	*	*	0.6	1.0	1.4	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
Black, both sexes										
2006 4/.....	1.8	3.4	0.8	*	*	*	1.3	1.5	1.6	2.8
2005 4/.....	1.8	4.7	0.9	*	*	0.6	1.1	1.4	1.9	2.5
2004 4/.....	1.8	4.8	*	*	*	0.7	1.1	1.0	2.6	3.1
2003 4/.....	1.8	3.2	*	*	*	*	1.1	1.4	2.1	3.2
2002.....	2.0	5.5	*	*	*	*	1.3	2.0	2.5	4.1
2001.....	1.7	5.5	*	*	*	*	1.1	1.5	1.9	3.8
2000.....	1.8	*	*	*	*	*	1.4	1.7	2.2	3.3
1999.....	1.9	*	*	*	*	0.8	1.7	1.9	2.5	3.9
Black, male										
2006 4/.....	2.7	*	*	*	*	*	2.0	2.6	2.3	4.0
2005 4/.....	2.4	*	*	*	*	*	1.6	2.3	2.3	3.1
2004 4/.....	2.6	*	*	*	*	*	1.5	1.5	4.0	4.6
2003 4/.....	2.7	*	*	*	*	*	1.9	2.5	3.4	4.4
2002.....	2.9	*	*	*	*	*	1.9	3.2	3.7	6.5
2001.....	2.7	6.0	*	*	*	*	1.9	2.5	2.7	5.9
2000.....	2.8	*	*	*	*	*	2.5	2.7	3.4	4.8
1999.....	3.0	*	*	*	*	1.3	2.7	3.4	3.7	6.3
Black, female										
2006 4/.....	1.0	*	*	*	*	*	*	*	*	1.8
2005 4/.....	1.1	*	*	*	*	*	*	*	1.5	2.0
2004 4/.....	1.1	*	*	*	*	*	*	*	1.4	1.8
2003 4/.....	0.9	*	*	*	*	*	*	*	*	2.2
2002.....	1.2	*	*	*	*	*	*	*	1.5	1.9
2001.....	0.9	*	*	*	*	*	*	*	*	2.0
2000.....	0.9	*	*	*	*	*	*	*	*	2.0
1999.....	0.9	*	*	*	*	*	*	*	1.4	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)										
All races 3/, both sexes										
2006.....	1.1	*	*	*	*	0.4	1.4	1.4	1.6	1.8
2005.....	1.0	*	*	*	*	0.3	1.0	1.3	1.4	1.7
2004.....	1.1	*	*	*	*	0.4	1.1	1.3	1.6	2.3
2003.....	1.2	*	*	*	*	0.4	1.4	1.4	1.8	2.6
2002.....	1.1	*	*	*	*	0.4	1.0	1.4	1.7	2.5
2001.....	1.0	*	*	*	*	0.3	0.8	1.1	1.5	2.3
2000.....	0.9	*	*	*	*	0.3	0.7	1.0	1.2	2.1
1999.....	0.9	*	*	*	*	0.3	0.8	1.1	1.5	2.1
All races 3/, male										
2006.....	1.3	*	*	*	*	0.5	2.0	1.9	2.0	2.1
2005.....	1.2	*	*	*	*	0.5	1.4	1.6	1.6	1.8
2004.....	1.4	*	*	*	*	0.6	1.5	1.8	2.0	2.9
2003.....	1.6	*	*	*	*	0.6	1.9	1.8	2.4	3.3
2002.....	1.4	*	*	*	*	0.5	1.4	1.9	2.1	3.3
2001.....	1.3	*	*	*	*	0.5	1.2	1.5	1.9	2.9
2000.....	1.2	*	*	*	*	0.4	1.0	1.4	1.7	2.8
1999.....	1.2	*	*	*	*	0.4	1.1	1.5	2.0	2.9
All races 3/, female										
2006.....	0.9	*	*	*	*	0.3	0.8	0.9	1.1	1.6
2005.....	0.9	*	*	*	*	0.2	0.6	1.0	1.3	1.6
2004.....	0.9	*	*	*	*	*	0.6	0.9	1.1	1.6
2003.....	0.9	*	*	*	*	0.3	0.7	0.9	1.2	1.8
2002.....	0.8	*	*	*	*	0.2	0.6	0.8	1.2	1.7
2001.....	0.7	*	*	*	*	*	0.4	0.7	1.1	1.7
2000.....	0.5	*	*	*	*	*	0.4	0.6	0.7	1.3
1999.....	0.6	*	*	*	*	*	0.4	0.6	1.0	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

White, both sexes

2006 4/.....	1.2	*	*	*	*	0.5	1.6	1.6	1.8	2.0
2005 4/.....	1.1	*	*	*	*	0.4	1.2	1.5	1.6	1.8
2004 4/.....	1.2	*	*	*	*	0.4	1.3	1.6	1.7	2.4
2003 4/.....	1.3	*	*	*	*	0.5	1.6	1.6	2.0	2.8
2002.....	1.2	*	*	*	*	0.4	1.2	1.5	1.8	2.6
2001.....	1.0	*	*	*	*	0.4	1.0	1.2	1.6	2.4
2000.....	0.9	*	*	*	*	0.3	0.8	1.1	1.3	2.1
1999.....	0.9	*	*	*	*	0.3	0.9	1.1	1.7	2.1

White, male

2006 4/.....	1.4	*	*	*	*	0.7	2.3	2.1	2.2	2.2
2005 4/.....	1.2	*	*	*	*	0.6	1.6	1.8	1.8	1.9
2004 4/.....	1.5	*	*	*	*	0.7	1.9	2.2	2.1	3.0
2003 4/.....	1.6	*	*	*	*	0.7	2.3	2.1	2.6	3.5
2002.....	1.4	*	*	*	*	0.5	1.6	2.1	2.2	3.3
2001.....	1.3	*	*	*	*	0.6	1.4	1.6	2.1	2.9
2000.....	1.2	*	*	*	*	0.4	1.2	1.5	1.8	2.9
1999.....	1.1	*	*	*	*	0.5	1.2	1.5	2.2	2.8

White, female

2006 4/.....	1.0	*	*	*	*	0.3	0.9	1.1	1.3	1.7
2005 4/.....	1.0	*	*	*	*	0.3	0.7	1.1	1.4	1.7
2004 4/.....	1.0	*	*	*	*	*	0.8	1.0	1.2	1.8
2003 4/.....	1.0	*	*	*	*	0.3	0.9	1.1	1.4	2.0
2002.....	0.9	*	*	*	*	0.3	0.8	0.9	1.2	1.9
2001.....	0.7	*	*	*	*	*	0.6	0.8	1.1	1.8
2000.....	0.6	*	*	*	*	*	0.5	0.7	0.8	1.3
1999.....	0.6	*	*	*	*	*	0.5	0.7	1.1	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
---	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2006.....	0.8	0.6	0.1	*	*	0.1	0.1	0.1	0.2	0.2
2005.....	0.9	*	*	*	*	*	*	0.2	0.2	0.3
2004.....	1.0	*	*	*	*	0.1	0.1	0.2	0.2	0.3
2003.....	1.0	*	0.2	*	*	0.1	0.1	0.1	0.2	0.4
2002.....	1.0	*	*	*	0.1	*	*	0.1	0.2	0.3
2001.....	1.1	0.6	0.2	*	*	*	0.1	0.1	0.3	0.3
2000.....	1.1	*	0.1	*	*	*	0.1	0.2	0.2	0.3
1999.....	1.0	1.4	0.2	*	*	*	*	0.1	0.2	0.3

All races 3/, male

2006.....	0.8	*	*	*	*	*	*	*	*	*
2005.....	0.8	*	*	*	*	*	*	*	*	0.2
2004.....	0.9	*	*	*	*	*	*	*	*	0.2
2003.....	0.9	*	*	*	*	*	*	*	0.2	0.3
2002.....	0.9	*	*	*	*	*	*	*	0.3	0.3
2001.....	1.0	*	*	*	*	*	*	*	0.3	0.4
2000.....	1.0	*	*	*	*	*	*	0.2	0.2	0.3
1999.....	1.0	1.6	*	*	*	*	*	*	0.3	0.3

All races 3/, female

2006.....	0.9	*	*	*	*	*	*	0.2	*	0.2
2005.....	1.0	*	*	*	*	*	*	*	0.3	0.3
2004.....	1.0	*	*	*	*	*	*	*	0.3	0.3
2003.....	1.0	*	*	*	*	*	*	*	0.2	0.4
2002.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2001.....	1.1	*	*	*	*	*	*	*	0.2	0.3
2000.....	1.1	*	*	*	*	*	*	*	0.2	0.3
1999.....	1.1	1.2	*	*	*	*	*	0.2	*	0.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
All races 3/, both sexes										
2006.....	229.3	347.7	516.4	730.1	1,110.2	1,656.6	2,554.8	4,033.4	6,524.0	13,253.1
2005.....	233.4	353.1	520.8	734.6	1,136.9	1,700.0	2,657.6	4,154.0	6,712.9	13,798.6
2004.....	231.7	352.3	511.7	734.8	1,140.1	1,727.4	2,676.4	4,193.2	6,717.2	13,823.5
2003.....	241.3	359.2	517.8	755.6	1,182.7	1,793.7	2,778.3	4,363.4	6,976.2	14,593.3
2002.....	242.6	357.1	512.8	770.1	1,187.7	1,832.7	2,845.9	4,449.1	7,103.8	14,828.3
2001.....	240.5	354.8	512.4	771.8	1,210.7	1,869.7	2,878.3	4,494.0	7,151.9	15,112.8
2000.....	239.1	352.5	509.0	792.6	1,241.0	1,906.3	2,929.4	4,553.8	7,335.0	15,524.4
1999.....	236.3	346.6	501.4	801.6	1,254.0	1,956.7	2,995.3	4,608.0	7,378.5	15,554.6
All races 3/, male										
2006.....	285.9	435.3	659.7	920.0	1,373.6	2,040.2	3,117.5	4,944.6	7,942.7	14,309.1
2005.....	292.7	443.7	666.0	925.7	1,410.0	2,084.2	3,267.0	5,103.4	8,147.4	14,889.4
2004.....	289.5	444.4	657.0	917.3	1,411.0	2,125.1	3,279.2	5,173.5	8,179.0	15,031.1
2003.....	304.1	457.9	660.8	942.3	1,462.2	2,216.6	3,428.7	5,400.3	8,513.7	15,794.0
2002.....	306.4	454.7	653.5	961.8	1,476.0	2,265.9	3,533.0	5,522.7	8,652.3	16,254.5
2001.....	304.8	455.4	646.8	958.4	1,496.8	2,315.6	3,587.4	5,603.2	8,781.9	16,744.8
2000.....	305.9	456.3	642.2	989.8	1,535.9	2,362.0	3,676.1	5,693.2	9,095.4	17,501.4
1999.....	304.0	448.0	632.4	1,003.8	1,560.8	2,444.1	3,785.1	5,806.4	9,216.8	17,597.2
All races 3/, female										
2006.....	173.1	262.2	378.9	551.0	869.7	1,321.4	2,097.0	3,366.6	5,637.2	12,759.0
2005.....	174.7	264.6	381.6	554.3	887.9	1,364.7	2,164.0	3,464.7	5,822.0	13,297.7
2004.....	174.5	262.5	372.7	562.7	893.6	1,381.2	2,191.0	3,488.7	5,816.6	13,280.3
2003.....	179.2	263.2	380.9	579.7	928.8	1,427.1	2,256.6	3,625.3	6,036.0	14,062.5
2002.....	179.7	262.4	378.2	590.0	926.0	1,458.7	2,296.7	3,690.9	6,162.2	14,209.6
2001.....	177.2	257.2	383.7	596.9	951.3	1,485.9	2,314.8	3,717.6	6,175.1	14,429.9
2000.....	173.4	252.0	381.3	608.1	973.7	1,515.7	2,341.2	3,760.2	6,296.6	14,719.2
1999.....	169.8	248.5	375.9	612.6	976.4	1,539.6	2,378.2	3,774.8	6,302.8	14,731.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
White, both sexes										
2006 4/.....	214.0	323.7	479.5	686.5	1,064.0	1,607.4	2,519.6	4,020.7	6,550.9	13,480.4
2005 4/.....	217.8	326.6	480.7	689.7	1,085.7	1,649.5	2,618.1	4,128.9	6,731.9	14,021.1
2004 4/.....	215.0	325.0	471.5	688.5	1,090.5	1,675.5	2,634.1	4,160.5	6,736.0	14,010.1
2003 4/.....	222.6	329.4	474.5	710.6	1,127.8	1,739.2	2,728.6	4,329.5	6,992.1	14,792.3
2002.....	224.0	327.4	471.3	723.5	1,133.0	1,777.6	2,796.3	4,407.8	7,111.3	15,015.9
2001.....	218.8	322.8	471.4	723.2	1,156.1	1,805.8	2,823.2	4,446.8	7,149.4	15,287.5
2000.....	215.9	317.9	467.4	743.9	1,184.9	1,844.5	2,866.8	4,496.6	7,333.2	15,707.0
1999.....	210.8	310.5	460.0	747.2	1,193.5	1,895.5	2,923.9	4,542.8	7,370.9	15,718.0
White, male										
2006 4/.....	269.8	408.3	612.8	861.2	1,309.5	1,973.1	3,060.0	4,914.2	7,982.5	14,576.8
2005 4/.....	276.4	413.4	616.6	864.0	1,341.8	2,016.7	3,208.8	5,065.8	8,169.0	15,156.5
2004 4/.....	272.8	412.3	607.8	855.8	1,343.7	2,055.9	3,219.3	5,123.2	8,214.4	15,250.7
2003 4/.....	284.0	423.5	606.9	884.2	1,390.6	2,146.2	3,359.5	5,349.4	8,522.6	16,037.9
2002.....	287.2	420.4	601.2	902.0	1,402.7	2,196.1	3,469.7	5,469.3	8,660.8	16,473.2
2001.....	281.4	417.1	595.5	896.0	1,427.9	2,237.6	3,513.9	5,543.5	8,771.1	16,935.4
2000.....	279.4	416.1	588.3	925.1	1,465.1	2,285.1	3,593.1	5,620.3	9,092.5	17,716.4
1999.....	274.0	402.1	579.3	934.6	1,484.0	2,367.8	3,698.3	5,727.2	9,211.2	17,752.9
White, female										
2006 4/.....	157.3	239.3	348.7	518.0	835.1	1,281.6	2,071.0	3,353.8	5,646.4	12,965.7
2005 4/.....	158.6	239.9	347.5	521.9	847.6	1,322.9	2,129.8	3,435.6	5,831.3	13,498.3
2004 4/.....	156.6	238.0	338.1	527.6	855.6	1,338.1	2,153.2	3,456.3	5,817.7	13,450.9
2003 4/.....	160.7	235.8	344.9	543.9	884.5	1,379.6	2,212.3	3,592.1	6,048.6	14,240.6
2002.....	160.5	235.1	344.3	552.6	883.7	1,409.1	2,247.4	3,648.0	6,161.6	14,382.8
2001.....	155.9	229.4	350.1	558.4	905.3	1,426.7	2,263.9	3,669.9	6,169.3	14,597.6
2000.....	152.2	220.7	349.3	571.5	926.5	1,459.3	2,284.0	3,706.2	6,286.6	14,890.7
1999.....	147.4	219.9	343.6	569.0	925.9	1,483.5	2,308.5	3,711.2	6,284.5	14,900.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All causes										
Black, both sexes										
2006 4/.....	373.1	576.3	892.9	1,232.7	1,750.2	2,409.9	3,371.9	4,902.4	7,275.3	12,489.7
2005 4/.....	379.3	602.7	914.9	1,254.6	1,812.2	2,466.3	3,514.3	5,161.7	7,547.7	13,083.1
2004 4/.....	386.6	608.3	910.1	1,271.1	1,813.5	2,509.7	3,582.0	5,266.0	7,493.0	13,338.5
2003 4/.....	410.4	632.9	940.3	1,295.1	1,889.9	2,595.8	3,754.9	5,421.2	7,745.3	13,976.7
2002.....	412.9	634.1	925.9	1,321.6	1,880.7	2,632.0	3,824.9	5,571.3	7,962.2	14,260.8
2001.....	428.4	644.1	928.5	1,328.9	1,891.7	2,738.8	3,904.2	5,660.2	8,028.9	14,550.8
2000.....	434.5	666.6	938.7	1,347.0	1,929.0	2,769.0	4,019.7	5,782.6	8,161.0	14,714.3
1999.....	448.6	668.3	929.9	1,399.3	1,969.2	2,813.7	4,148.0	5,878.3	8,169.4	14,757.9
Black, male										
2006 4/.....	457.3	715.6	1,171.2	1,625.1	2,302.6	3,132.8	4,373.3	6,329.2	9,021.0	13,206.0
2005 4/.....	463.8	755.9	1,184.8	1,667.1	2,380.7	3,174.7	4,509.2	6,579.2	9,376.2	13,809.8
2004 4/.....	467.1	766.8	1,186.6	1,668.7	2,374.4	3,242.7	4,587.3	6,747.0	9,229.2	14,452.5
2003 4/.....	507.1	800.8	1,227.5	1,688.7	2,454.5	3,334.8	4,843.9	6,978.2	9,827.8	14,903.4
2002.....	502.9	801.4	1,206.6	1,719.3	2,464.7	3,357.1	4,918.1	7,088.6	9,968.4	15,635.5
2001.....	527.8	830.2	1,201.4	1,718.2	2,424.6	3,497.3	5,053.2	7,255.7	10,256.3	16,439.9
2000.....	549.1	853.0	1,227.4	1,776.2	2,462.1	3,513.8	5,203.9	7,410.0	10,405.7	16,791.0
1999.....	572.7	885.6	1,220.1	1,822.6	2,549.7	3,599.1	5,336.3	7,538.1	10,433.8	17,304.5
Black, female										
2006 4/.....	298.9	455.0	657.4	909.1	1,319.8	1,876.0	2,683.8	4,048.5	6,352.9	12,196.7
2005 4/.....	304.8	469.0	686.6	914.1	1,369.2	1,940.8	2,833.5	4,316.0	6,581.9	12,789.9
2004 4/.....	315.7	470.1	676.4	943.8	1,376.3	1,965.5	2,899.8	4,388.5	6,578.2	12,896.9
2003 4/.....	325.1	486.7	697.2	971.3	1,449.4	2,047.3	3,020.5	4,495.8	6,654.1	13,616.7
2002.....	333.6	489.0	688.0	995.2	1,423.7	2,093.9	3,091.2	4,664.6	6,917.8	13,734.2
2001.....	340.6	482.8	696.2	1,009.5	1,472.5	2,178.4	3,138.8	4,707.8	6,894.3	13,849.9
2000.....	333.3	505.0	692.6	995.2	1,508.3	2,219.3	3,233.3	4,801.6	7,030.5	13,941.3
1999.....	339.3	480.9	683.9	1,053.5	1,514.0	2,231.6	3,359.6	4,878.6	7,047.2	13,805.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

All races 3/, both sexes

2006.....	16.0	21.0	27.1	28.2	30.6	41.5	61.2	96.4	150.3	253.9
2005.....	17.0	22.3	28.0	27.0	29.8	41.0	62.0	93.3	147.9	259.6
2004.....	18.2	22.8	27.2	26.5	29.9	40.2	60.3	91.6	142.7	249.5
2003.....	18.9	24.0	26.4	25.9	29.7	39.6	60.2	94.7	141.6	255.2
2002.....	19.6	24.7	25.1	24.0	28.8	41.5	60.8	94.0	140.5	252.8
2001.....	19.7	23.2	22.6	22.9	28.7	39.4	57.2	87.5	130.0	246.3
2000.....	19.5	23.5	22.2	21.9	27.5	37.5	54.7	85.1	131.5	256.0
1999.....	20.9	22.9	21.1	22.2	26.5	40.3	56.4	86.9	132.5	267.9

All races 3/, male

2006.....	22.1	29.0	38.4	38.2	37.5	48.0	67.3	109.4	168.0	276.6
2005.....	23.7	31.5	39.3	36.3	36.5	47.5	70.5	105.2	168.2	278.2
2004.....	24.9	32.1	38.0	35.8	35.8	46.6	68.8	103.7	163.7	273.1
2003.....	26.4	34.3	37.6	34.6	35.3	46.3	68.0	106.2	162.9	269.1
2002.....	27.5	35.6	36.6	31.2	35.3	47.9	69.6	109.1	164.0	270.9
2001.....	28.1	33.9	32.3	30.4	34.0	44.8	65.5	102.4	153.8	269.7
2000.....	27.8	34.9	32.3	28.8	33.4	43.8	62.7	99.8	154.8	273.0
1999.....	30.6	33.9	30.2	29.1	32.0	47.4	65.4	102.6	157.8	290.1

All races 3/, female

2006.....	9.9	13.3	16.3	18.7	24.4	35.8	56.2	86.9	139.2	243.3
2005.....	10.4	13.2	17.3	18.1	23.6	35.3	55.1	84.7	135.3	251.0
2004.....	11.5	13.6	16.8	17.7	24.6	34.7	53.5	82.9	129.7	238.8
2003.....	11.6	13.9	15.7	17.8	24.6	33.8	54.0	86.5	128.6	249.1
2002.....	11.8	14.0	14.1	17.2	22.9	36.0	53.8	83.4	126.2	245.0
2001.....	11.4	12.8	13.2	15.8	23.9	34.8	50.6	77.1	115.8	236.5
2000.....	11.3	12.5	12.6	15.4	22.1	32.1	48.4	74.9	117.8	249.1
1999.....	11.3	12.4	12.4	15.8	21.5	34.2	49.3	75.9	117.8	258.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
I. Certain infectious and parasitic diseases (A00-B99)										
White, both sexes										
2006 4/.....	10.7	14.9	21.0	22.1	25.9	36.5	55.7	91.9	145.3	250.0
2005 4/.....	11.6	16.0	21.8	21.6	24.8	36.3	56.8	87.6	141.7	251.5
2004 4/.....	12.0	16.4	20.8	21.1	24.7	34.8	55.2	86.4	136.4	242.1
2003 4/.....	12.9	17.6	19.7	20.4	24.4	33.9	55.1	88.6	136.1	245.0
2002.....	12.8	17.8	18.4	18.7	23.5	35.8	55.1	87.3	133.9	242.3
2001.....	12.8	16.3	16.3	17.9	23.2	34.0	51.5	81.4	123.5	235.4
2000.....	12.8	16.2	15.9	16.9	22.0	31.9	49.0	77.9	123.0	242.9
1999.....	13.2	15.3	15.3	16.6	21.6	34.7	49.5	78.9	124.1	251.9
White, male										
2006 4/.....	15.6	21.7	29.9	29.6	30.8	42.1	61.2	103.6	162.7	275.1
2005 4/.....	17.0	23.5	30.6	28.5	30.1	41.3	63.9	99.0	160.7	269.2
2004 4/.....	17.5	23.6	29.4	28.0	28.8	40.0	62.3	98.1	155.4	266.6
2003 4/.....	18.8	26.0	28.4	26.7	28.5	39.0	61.5	98.9	156.5	259.4
2002.....	18.9	26.5	26.6	24.1	28.5	40.7	63.0	100.0	155.3	260.3
2001.....	19.5	24.4	23.5	23.3	27.2	38.3	58.3	95.3	146.1	256.7
2000.....	18.9	24.6	22.9	21.9	26.3	36.8	56.4	90.4	145.0	258.5
1999.....	20.3	22.6	21.8	21.2	25.5	40.2	57.2	94.2	146.7	270.4
White, female										
2006 4/.....	5.6	8.1	12.3	14.9	21.3	31.5	51.2	83.3	134.4	238.3
2005 4/.....	6.1	8.6	13.2	15.0	19.8	31.8	50.9	79.2	129.8	243.4
2004 4/.....	6.5	9.2	12.3	14.6	20.8	30.2	49.4	77.9	124.5	231.1
2003 4/.....	6.9	9.2	11.3	14.5	20.6	29.4	49.9	81.1	123.6	238.6
2002.....	6.6	9.3	10.4	13.5	18.9	31.5	48.6	78.2	120.8	234.5
2001.....	6.1	8.4	9.2	12.6	19.5	30.3	46.0	71.6	109.8	226.4
2000.....	6.6	7.9	9.0	12.1	18.0	27.7	43.1	69.1	110.0	236.5
1999.....	6.2	8.1	8.9	12.3	18.1	29.8	43.4	68.2	110.7	244.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Black, both sexes

2006 4/.....	54.1	67.4	77.5	82.6	77.9	92.2	119.1	158.7	229.3	343.8
2005 4/.....	56.6	70.1	80.4	76.0	79.1	88.1	116.3	161.9	244.9	392.6
2004 4/.....	62.4	71.4	80.3	76.5	80.1	91.4	115.8	161.6	240.1	375.3
2003 4/.....	64.0	73.9	81.9	76.6	80.0	92.2	115.9	171.4	230.0	416.6
2002.....	68.7	78.3	81.7	73.6	76.1	94.7	122.7	181.9	238.1	413.1
2001.....	69.7	77.3	75.6	69.7	79.2	89.6	119.4	166.2	226.7	404.7
2000.....	68.7	80.8	77.5	66.3	74.5	89.9	119.3	177.5	253.3	436.7
1999.....	76.9	83.1	71.9	72.0	69.0	93.0	131.6	181.7	249.4	469.0

Black, male

2006 4/.....	72.2	88.5	113.3	121.2	108.5	112.5	140.3	204.1	258.8	350.3
2005 4/.....	76.7	97.8	117.2	113.1	106.3	117.4	145.8	190.9	290.7	425.7
2004 4/.....	82.4	102.3	115.2	114.8	108.6	115.1	147.6	192.4	299.2	408.3
2003 4/.....	87.5	105.3	121.3	115.4	105.7	122.6	144.5	205.4	285.0	456.7
2002.....	93.7	112.7	126.8	104.1	101.6	123.4	151.2	232.9	297.3	448.1
2001.....	96.3	114.3	111.6	101.8	102.5	111.6	151.1	204.0	278.5	470.9
2000.....	97.4	121.5	120.5	96.7	100.2	116.0	144.1	229.2	311.1	486.7
1999.....	112.3	128.3	108.8	103.4	95.7	120.0	166.2	213.7	325.2	537.8

Black, female

2006 4/.....	38.2	49.1	47.3	50.8	54.1	77.2	104.6	131.5	213.8	341.1
2005 4/.....	38.9	46.0	49.2	45.3	57.8	66.3	96.1	144.5	220.7	379.2
2004 4/.....	44.7	44.5	50.7	45.0	58.0	73.8	94.2	143.4	209.0	362.3
2003 4/.....	43.4	46.6	48.5	44.7	60.0	69.6	96.6	151.1	201.2	401.0
2002.....	46.7	48.4	43.5	48.5	56.2	73.4	103.5	151.5	207.3	399.6
2001.....	46.1	45.2	44.9	43.4	60.8	73.4	98.2	143.7	200.3	380.2
2000.....	43.5	45.6	40.9	41.4	54.3	70.7	102.8	146.4	224.2	418.1
1999.....	45.7	44.0	40.7	46.4	48.1	73.1	108.7	162.5	211.7	443.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intestinal infectious diseases (A00-A09)										
All races 3/, both sexes										
2006.....	0.1	0.2	0.4	0.9	1.6	3.3	6.8	13.6	26.3	50.7
2005.....	0.1	0.2	0.4	0.6	1.3	2.8	6.5	11.6	21.6	45.9
2004.....	0.1	0.2	0.3	0.6	1.3	2.2	4.7	9.4	17.5	34.4
2003.....	*	0.1	0.2	0.5	0.9	1.7	3.4	7.2	12.4	25.0
2002.....	*	0.1	0.2	0.4	0.7	1.4	2.7	5.1	9.8	21.0
2001.....	*	*	0.2	0.2	0.5	0.8	1.9	3.3	6.6	13.1
2000.....	*	*	0.1	0.3	0.4	0.9	1.6	2.8	5.4	11.9
1999.....	*	*	*	*	0.3	0.8	1.3	2.4	4.6	10.6
All races 3/, male										
2006.....	*	0.3	0.4	0.9	1.8	3.7	6.5	14.0	25.5	50.2
2005.....	*	0.2	0.4	0.6	1.5	2.9	6.3	12.5	22.2	46.2
2004.....	*	0.2	0.3	0.6	1.4	2.2	4.3	9.4	17.9	32.2
2003.....	*	*	0.2	0.5	0.9	1.5	3.1	7.0	12.9	21.9
2002.....	*	*	*	0.4	0.8	1.6	3.1	5.1	10.3	20.4
2001.....	*	*	*	*	0.4	0.8	2.0	3.6	6.5	12.2
2000.....	*	*	*	0.4	0.4	1.0	1.7	3.0	5.9	11.5
1999.....	*	*	*	*	*	0.8	1.3	2.7	5.1	9.7
All races 3/, female										
2006.....	*	*	0.4	0.8	1.4	3.0	7.0	13.4	26.8	50.9
2005.....	*	*	0.4	0.6	1.1	2.7	6.6	11.0	21.3	45.7
2004.....	*	*	0.3	0.6	1.2	2.2	4.9	9.3	17.3	35.4
2003.....	*	*	0.2	0.5	1.0	1.8	3.6	7.2	12.1	26.4
2002.....	*	*	*	0.5	0.6	1.3	2.3	5.1	9.5	21.3
2001.....	*	*	0.2	*	0.6	0.8	1.8	3.1	6.6	13.4
2000.....	*	*	*	*	*	0.8	1.6	2.7	5.1	12.0
1999.....	*	*	*	*	*	0.8	1.3	2.2	4.4	10.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intestinal infectious diseases (A00-A09)										
White, both sexes										
2006 4/.....	0.1	0.2	0.3	0.9	1.5	3.5	7.0	14.4	28.0	53.7
2005 4/.....	*	0.2	0.4	0.5	1.3	3.0	6.7	12.3	22.8	48.7
2004 4/.....	*	0.1	0.3	0.6	1.3	2.3	4.9	9.9	18.4	36.7
2003 4/.....	*	0.1	0.2	0.5	0.9	1.8	3.6	7.3	13.2	26.3
2002.....	*	0.1	0.2	0.4	0.6	1.6	2.7	5.3	10.4	22.1
2001.....	*	*	0.2	0.2	0.5	0.8	2.0	3.5	6.7	13.8
2000.....	*	*	*	0.3	0.4	0.9	1.7	3.0	5.8	12.3
1999.....	*	*	*	*	0.3	0.8	1.3	2.5	4.8	11.2
White, male										
2006 4/.....	*	0.2	0.3	0.9	1.7	3.9	6.7	14.6	26.8	53.2
2005 4/.....	*	*	0.4	0.5	1.5	3.2	6.5	13.0	22.8	49.4
2004 4/.....	*	*	0.3	0.5	1.4	2.2	4.6	9.8	18.9	34.2
2003 4/.....	*	*	*	0.5	0.8	1.7	3.3	7.3	13.8	23.2
2002.....	*	*	*	0.4	0.7	1.7	3.0	5.3	10.9	21.5
2001.....	*	*	*	*	*	0.8	2.0	3.7	6.5	12.8
2000.....	*	*	*	0.4	*	1.1	1.8	3.2	6.2	12.0
1999.....	*	*	*	*	*	0.8	1.4	2.8	5.3	10.2
White, female										
2006 4/.....	*	*	0.4	0.8	1.4	3.1	7.3	14.3	28.8	54.0
2005 4/.....	*	*	0.4	0.5	1.0	2.8	6.9	11.7	22.8	48.3
2004 4/.....	*	*	0.3	0.6	1.2	2.3	5.2	9.9	18.2	37.9
2003 4/.....	*	*	*	0.5	0.9	1.9	3.9	7.3	12.9	27.7
2002.....	*	*	*	0.4	0.6	1.5	2.4	5.4	10.1	22.3
2001.....	*	*	0.3	*	0.5	0.8	1.9	3.3	6.9	14.2
2000.....	*	*	*	*	*	0.8	1.6	2.8	5.6	12.5
1999.....	*	*	*	*	*	0.8	1.3	2.2	4.5	11.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other intestinal infectious diseases (A01-A08)

All races 3/, both sexes

2006.....	0.1	0.2	0.4	0.9	1.5	3.3	6.6	13.3	25.7	49.7
2005.....	0.1	0.2	0.4	0.6	1.2	2.7	6.3	11.4	21.2	44.8
2004.....	0.1	0.2	0.3	0.6	1.2	2.2	4.5	9.1	17.1	33.6
2003.....	*	0.1	0.2	0.5	0.9	1.6	3.3	6.9	12.1	24.2
2002.....	*	0.1	0.2	0.4	0.7	1.4	2.6	5.0	9.4	20.2
2001.....	*	*	0.2	0.2	0.5	0.8	1.8	3.2	6.4	12.6
2000.....	*	*	0.1	0.3	0.3	0.8	1.5	2.7	5.3	11.3
1999.....	*	*	*	*	0.3	0.7	1.1	2.2	4.3	9.9

All races 3/, male

2006.....	*	0.2	0.4	0.9	1.7	3.7	6.4	13.6	24.9	49.3
2005.....	*	*	0.4	0.5	1.4	2.9	6.1	12.1	21.7	45.5
2004.....	*	0.2	0.3	0.6	1.3	2.2	4.3	9.2	17.5	31.7
2003.....	*	*	0.2	0.5	0.8	1.5	3.1	6.7	12.6	21.1
2002.....	*	*	*	0.4	0.7	1.5	3.0	5.0	9.9	19.7
2001.....	*	*	*	*	0.4	0.8	2.0	3.4	6.4	11.9
2000.....	*	*	*	0.4	*	1.0	1.6	2.9	5.8	10.6
1999.....	*	*	*	*	*	0.7	1.1	2.6	4.9	9.3

All races 3/, female

2006.....	*	*	0.4	0.8	1.3	2.9	6.8	13.0	26.1	49.8
2005.....	*	*	0.4	0.6	1.0	2.6	6.4	10.9	20.9	44.5
2004.....	*	*	0.3	0.6	1.2	2.1	4.7	9.0	16.9	34.5
2003.....	*	*	0.2	0.5	0.9	1.7	3.4	7.0	11.8	25.5
2002.....	*	*	*	0.4	0.6	1.3	2.2	5.0	9.1	20.4
2001.....	*	*	*	*	0.6	0.8	1.7	3.0	6.4	13.0
2000.....	*	*	*	*	*	0.7	1.5	2.6	5.0	11.5
1999.....	*	*	*	*	*	0.7	1.2	1.9	4.0	10.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other intestinal infectious diseases (A01-A08)										
White, both sexes										
2006 4/.....	0.1	0.2	0.3	0.8	1.5	3.4	6.9	14.1	27.3	52.7
2005 4/.....	*	0.2	0.4	0.5	1.2	2.9	6.5	12.0	22.3	47.6
2004 4/.....	*	0.1	0.3	0.6	1.3	2.2	4.8	9.6	18.0	35.9
2003 4/.....	*	0.1	0.2	0.5	0.9	1.7	3.5	7.0	12.9	25.4
2002.....	*	0.1	0.2	0.4	0.6	1.5	2.6	5.2	10.1	21.1
2001.....	*	*	0.2	0.2	0.5	0.7	1.9	3.3	6.5	13.3
2000.....	*	*	*	0.3	0.3	0.8	1.6	2.8	5.6	11.7
1999.....	*	*	*	*	0.3	0.7	1.2	2.2	4.5	10.5
White, male										
2006 4/.....	*	0.2	0.3	0.9	1.7	3.8	6.7	14.2	26.2	52.2
2005 4/.....	*	*	0.4	0.5	1.5	3.1	6.3	12.6	22.3	48.7
2004 4/.....	*	*	0.3	0.5	1.4	2.2	4.5	9.7	18.4	33.7
2003 4/.....	*	*	*	0.4	0.8	1.7	3.3	7.0	13.4	22.4
2002.....	*	*	*	0.4	0.7	1.6	2.9	5.2	10.6	20.7
2001.....	*	*	*	*	*	0.7	2.0	3.5	6.4	12.4
2000.....	*	*	*	0.4	*	1.0	1.6	3.0	6.0	11.1
1999.....	*	*	*	*	*	0.7	1.2	2.6	5.1	9.7
White, female										
2006 4/.....	*	*	0.4	0.8	1.3	3.0	7.1	14.0	28.0	53.0
2005 4/.....	*	*	0.4	0.5	1.0	2.7	6.7	11.6	22.4	47.1
2004 4/.....	*	*	0.3	0.6	1.2	2.2	5.1	9.6	17.7	36.9
2003 4/.....	*	*	*	0.5	0.9	1.8	3.7	7.1	12.6	26.7
2002.....	*	*	*	0.4	0.6	1.4	2.3	5.3	9.7	21.4
2001.....	*	*	*	*	0.5	0.7	1.9	3.2	6.6	13.7
2000.....	*	*	*	*	*	0.7	1.5	2.7	5.4	11.9
1999.....	*	*	*	*	*	0.7	1.2	2.0	4.1	10.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intestinal infections due to other specified organisms (A04,A07-A08)

All races 3/, both sexes

2006.....	0.1	0.2	0.4	0.8	1.5	3.2	6.5	13.2	25.6	49.5
2005.....	0.1	0.1	0.4	0.5	1.2	2.7	6.3	11.3	21.0	44.7
2004.....	0.1	0.2	0.3	0.6	1.2	2.1	4.5	9.1	17.0	33.5
2003.....	*	0.1	0.2	0.4	0.8	1.6	3.2	6.8	12.0	23.9
2002.....	*	0.1	0.2	0.4	0.7	1.3	2.5	4.9	9.3	20.2
2001.....	*	*	0.2	0.2	0.5	0.7	1.8	3.1	6.2	12.4
2000.....	*	*	*	0.2	0.3	0.8	1.5	2.6	5.1	11.1
1999.....	*	*	*	*	0.3	0.6	1.1	2.0	4.2	9.7

All races 3/, male

2006.....	*	0.2	0.4	0.9	1.7	3.6	6.3	13.5	24.7	49.1
2005.....	*	*	0.4	0.5	1.4	2.8	6.1	12.1	21.6	45.4
2004.....	*	*	0.3	0.5	1.3	2.1	4.2	9.1	17.4	31.4
2003.....	*	*	0.2	0.4	0.8	1.5	3.0	6.6	12.6	20.6
2002.....	*	*	*	0.4	0.7	1.4	2.9	5.0	9.7	19.6
2001.....	*	*	*	*	0.4	0.7	1.9	3.3	6.1	11.8
2000.....	*	*	*	0.3	*	0.9	1.5	2.8	5.7	10.5
1999.....	*	*	*	*	*	0.6	1.0	2.5	4.7	9.1

All races 3/, female

2006.....	*	*	0.4	0.8	1.3	2.9	6.8	13.0	26.1	49.7
2005.....	*	*	0.4	0.5	1.0	2.6	6.4	10.8	20.7	44.4
2004.....	*	*	0.3	0.6	1.2	2.1	4.7	9.0	16.8	34.4
2003.....	*	*	0.2	0.4	0.9	1.7	3.3	6.9	11.7	25.4
2002.....	*	*	*	0.4	0.6	1.2	2.2	4.9	9.1	20.4
2001.....	*	*	*	*	0.5	0.8	1.7	2.9	6.2	12.7
2000.....	*	*	*	*	*	0.7	1.5	2.5	4.8	11.4
1999.....	*	*	*	*	*	0.7	1.1	1.7	4.0	9.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intestinal infections due to other specified organisms (A04,A07-A08)

White, both sexes

2006 4/.....	*	0.2	0.3	0.8	1.5	3.4	6.8	14.0	27.2	52.6
2005 4/.....	*	0.2	0.3	0.5	1.2	2.9	6.5	12.0	22.2	47.5
2004 4/.....	*	0.1	0.3	0.6	1.3	2.2	4.8	9.6	17.9	35.8
2003 4/.....	*	*	0.2	0.4	0.8	1.7	3.4	7.0	12.8	25.2
2002.....	*	*	0.2	0.4	0.6	1.4	2.6	5.2	9.9	21.1
2001.....	*	*	0.2	0.2	0.4	0.7	1.9	3.2	6.3	13.1
2000.....	*	*	*	0.3	0.3	0.8	1.5	2.7	5.4	11.5
1999.....	*	*	*	*	0.3	0.7	1.1	2.1	4.4	10.2

White, male

2006 4/.....	*	*	0.3	0.9	1.7	3.8	6.5	14.1	26.0	52.1
2005 4/.....	*	*	0.3	0.4	1.4	3.1	6.3	12.6	22.3	48.6
2004 4/.....	*	*	0.3	0.5	1.4	2.1	4.5	9.5	18.3	33.5
2003 4/.....	*	*	*	0.4	0.8	1.6	3.2	6.9	13.3	22.0
2002.....	*	*	*	0.4	0.7	1.5	2.9	5.1	10.4	20.6
2001.....	*	*	*	*	*	0.7	1.9	3.3	6.1	12.3
2000.....	*	*	*	*	*	1.0	1.5	2.9	5.9	11.0
1999.....	*	*	*	*	*	0.6	1.1	2.6	5.0	9.5

White, female

2006 4/.....	*	*	0.4	0.8	1.3	3.0	7.1	14.0	28.0	52.9
2005 4/.....	*	*	0.4	0.5	1.0	2.7	6.7	11.5	22.1	47.0
2004 4/.....	*	*	0.3	0.6	1.2	2.2	5.0	9.6	17.7	36.8
2003 4/.....	*	*	*	0.5	0.9	1.8	3.6	7.0	12.5	26.6
2002.....	*	*	*	0.4	0.6	1.4	2.3	5.3	9.7	21.3
2001.....	*	*	*	*	0.5	0.7	1.9	3.1	6.5	13.4
2000.....	*	*	*	*	*	0.7	1.5	2.6	5.2	11.8
1999.....	*	*	*	*	*	0.7	1.1	1.8	4.1	10.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Tuberculosis (A16-A19)										
All races 3/, both sexes										
2006.....	0.1	0.2	0.2	0.3	0.4	0.6	0.7	0.9	1.7	2.1
2005.....	*	0.2	0.2	0.2	0.4	0.5	0.7	1.3	1.3	2.2
2004.....	0.1	0.1	0.2	0.2	0.3	0.6	1.0	1.1	1.5	2.4
2003.....	0.1	0.2	0.3	0.4	0.3	0.5	0.8	1.4	2.0	2.3
2002.....	0.1	0.2	0.2	0.3	0.5	0.9	0.9	1.5	2.3	2.6
2001.....	0.1	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.0	2.8
2000.....	0.1	0.2	0.3	0.2	0.5	0.7	1.0	1.8	2.4	2.7
1999.....	0.1	0.3	0.3	0.5	0.6	0.7	1.2	1.8	2.4	4.2
All races 3/, male										
2006.....	0.2	0.3	0.4	0.4	0.7	0.8	0.9	1.4	2.5	3.1
2005.....	*	0.2	0.3	0.3	0.6	0.7	1.1	1.7	1.7	3.3
2004.....	0.2	0.2	0.3	0.3	0.6	0.8	1.4	1.2	2.2	3.8
2003.....	*	0.4	0.4	0.6	0.4	0.6	1.2	2.0	2.7	3.1
2002.....	0.2	0.2	0.4	0.5	0.7	1.2	1.2	1.8	3.2	4.0
2001.....	*	0.3	0.4	0.7	0.7	0.9	1.5	1.7	2.4	4.4
2000.....	*	0.4	0.5	0.4	0.7	1.0	1.1	2.3	3.8	4.6
1999.....	0.2	0.4	0.5	0.8	0.9	1.1	1.7	2.4	3.7	7.1
All races 3/, female										
2006.....	*	*	*	*	*	*	0.5	0.6	1.3	1.6
2005.....	*	*	*	*	*	0.4	0.4	0.9	1.0	1.7
2004.....	*	*	*	*	*	*	0.6	1.0	1.1	1.7
2003.....	*	*	*	*	*	0.4	0.5	1.0	1.7	1.9
2002.....	*	*	*	*	0.4	0.5	0.6	1.3	1.7	2.1
2001.....	*	*	*	*	*	0.4	0.7	1.1	1.8	2.2
2000.....	*	*	*	*	*	0.5	0.9	1.4	1.6	2.0
1999.....	*	*	*	*	*	0.4	0.7	1.5	1.7	3.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Tuberculosis (A16-A19)										
White, both sexes										
2006 4/.....	*	0.1	0.1	0.2	0.3	0.4	0.4	0.8	1.2	1.6
2005 4/.....	*	0.1	0.1	*	0.2	0.4	0.4	0.8	0.9	1.7
2004 4/.....	*	*	0.1	*	0.2	0.3	0.7	0.8	1.2	1.7
2003 4/.....	*	0.1	0.1	0.3	0.2	0.3	0.7	1.0	1.7	1.8
2002.....	*	*	0.1	0.2	0.4	0.5	0.6	1.2	1.8	2.1
2001.....	*	0.1	0.1	0.3	0.3	0.5	0.7	1.1	1.4	2.2
2000.....	*	*	0.2	*	0.3	0.5	0.6	1.2	1.7	2.2
1999.....	*	0.2	0.2	0.3	0.4	0.6	0.8	1.3	2.0	3.2
White, male										
2006 4/.....	*	*	0.2	*	0.4	0.5	*	1.0	1.7	2.1
2005 4/.....	*	*	*	*	0.4	0.5	*	1.0	1.0	2.4
2004 4/.....	*	*	*	*	*	*	1.0	1.0	1.6	2.8
2003 4/.....	*	*	*	0.4	*	*	1.0	1.3	2.2	2.4
2002.....	*	*	*	*	0.5	0.8	1.0	1.3	2.5	3.2
2001.....	*	*	*	0.5	0.5	0.6	0.8	1.4	1.8	3.1
2000.....	*	*	0.3	*	*	0.7	0.8	1.5	2.3	3.8
1999.....	*	0.3	0.4	0.5	0.6	0.9	1.1	1.5	3.0	5.4
White, female										
2006 4/.....	*	*	*	*	*	*	*	0.6	0.9	1.4
2005 4/.....	*	*	*	*	*	*	*	0.6	0.8	1.4
2004 4/.....	*	*	*	*	*	*	*	0.7	0.9	1.3
2003 4/.....	*	*	*	*	*	*	*	0.9	1.3	1.6
2002.....	*	*	*	*	*	*	*	1.1	1.3	1.6
2001.....	*	*	*	*	*	*	0.5	0.9	1.2	1.8
2000.....	*	*	*	*	*	*	0.5	1.0	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	1.1	1.4	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Respiratory tuberculosis (A16)										
All races 3/, both sexes										
2006.....	*	0.1	0.2	0.2	0.4	0.4	0.5	0.7	1.4	1.6
2005.....	*	0.1	0.2	0.2	0.3	0.3	0.6	1.0	1.0	1.9
2004.....	0.1	0.1	0.2	0.2	0.3	0.4	0.8	0.8	1.1	2.2
2003.....	0.1	0.2	0.2	0.3	0.2	0.4	0.6	1.2	1.6	1.8
2002.....	0.1	0.1	0.2	0.2	0.4	0.6	0.7	1.2	2.0	2.3
2001.....	*	0.1	0.2	0.4	0.4	0.5	0.8	1.1	1.6	2.4
2000.....	0.1	0.1	0.2	0.2	0.3	0.5	0.8	1.4	2.1	2.3
1999.....	0.1	0.2	0.2	0.4	0.5	0.6	0.9	1.5	2.1	3.7
All races 3/, male										
2006.....	*	0.2	0.3	0.3	0.6	0.6	0.7	1.2	2.0	2.4
2005.....	*	*	0.3	0.2	0.4	0.5	0.9	1.2	1.4	2.7
2004.....	*	*	0.3	*	0.4	0.6	1.2	0.9	1.7	3.7
2003.....	*	0.3	0.3	0.5	*	0.5	1.0	1.7	2.2	2.5
2002.....	0.2	*	0.3	0.4	0.5	1.0	0.9	1.3	2.8	3.5
2001.....	*	0.2	0.3	0.6	0.7	0.7	1.2	1.5	2.0	3.8
2000.....	*	0.3	0.4	0.3	0.6	0.9	0.9	1.9	3.2	4.1
1999.....	*	0.3	0.4	0.6	0.9	0.9	1.4	2.0	3.3	6.4
All races 3/, female										
2006.....	*	*	*	*	*	*	*	*	1.0	1.2
2005.....	*	*	*	*	*	*	*	0.8	0.8	1.5
2004.....	*	*	*	*	*	*	0.5	0.7	0.8	1.5
2003.....	*	*	*	*	*	*	*	0.8	1.3	1.5
2002.....	*	*	*	*	*	*	0.5	1.1	1.5	1.8
2001.....	*	*	*	*	*	*	0.5	0.8	1.4	1.8
2000.....	*	*	*	*	*	*	0.6	1.1	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	1.2	1.4	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Respiratory tuberculosis (A16)										
White, both sexes										
2006 4/.....	*	*	*	*	0.2	0.3	0.3	0.6	1.0	1.2
2005 4/.....	*	*	*	*	0.2	0.3	0.3	0.6	0.7	1.3
2004 4/.....	*	*	*	*	*	0.3	0.5	0.6	0.9	1.6
2003 4/.....	*	*	*	0.2	*	*	0.5	0.9	1.4	1.4
2002.....	*	*	*	*	0.3	0.4	0.5	0.9	1.5	1.8
2001.....	*	*	*	0.2	0.3	0.4	0.5	0.9	1.2	1.8
2000.....	*	*	0.2	*	*	0.3	0.5	1.0	1.5	1.9
1999.....	*	0.1	0.1	0.2	0.4	0.5	0.6	1.1	1.7	3.0
White, male										
2006 4/.....	*	*	*	*	0.4	*	*	0.9	1.5	1.5
2005 4/.....	*	*	*	*	*	*	*	*	*	1.9
2004 4/.....	*	*	*	*	*	*	0.8	0.8	1.2	2.6
2003 4/.....	*	*	*	0.3	*	*	0.9	1.1	1.8	1.8
2002.....	*	*	*	*	*	0.7	0.8	0.9	2.0	2.9
2001.....	*	*	*	0.4	0.5	*	0.7	1.2	1.4	2.6
2000.....	*	*	0.3	*	*	0.5	0.7	1.3	1.9	3.4
1999.....	*	*	*	0.4	0.5	0.7	1.0	1.3	2.6	5.2
White, female										
2006 4/.....	*	*	*	*	*	*	*	*	0.7	1.0
2005 4/.....	*	*	*	*	*	*	*	*	0.7	1.1
2004 4/.....	*	*	*	*	*	*	*	*	0.7	1.1
2003 4/.....	*	*	*	*	*	*	*	*	0.8	1.2
2002.....	*	*	*	*	*	*	*	0.9	1.1	1.3
2001.....	*	*	*	*	*	*	*	0.6	1.0	1.4
2000.....	*	*	*	*	*	*	*	0.7	1.2	1.2
1999.....	*	*	*	*	*	*	*	0.9	1.1	2.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Zoonotic and other bacterial diseases (A20-A49)										
All races 3/, both sexes										
2006.....	2.6	4.2	7.1	10.9	17.1	26.5	42.0	69.0	108.3	184.6
2005.....	2.5	4.3	7.1	11.1	16.9	26.8	43.0	67.0	109.7	194.3
2004.....	2.5	4.4	7.3	11.0	17.2	26.0	43.1	68.3	108.7	194.8
2003.....	2.7	4.5	6.9	11.1	17.6	26.5	43.0	71.4	112.4	209.5
2002.....	2.5	4.5	6.7	10.6	16.8	28.1	45.1	73.3	113.9	210.5
2001.....	2.4	4.3	6.6	10.4	16.3	26.6	42.6	70.3	108.3	213.7
2000.....	2.6	4.2	6.5	9.9	15.9	25.2	40.0	66.9	109.2	222.5
1999.....	2.5	4.1	6.1	9.6	15.2	26.5	40.5	67.5	108.5	231.2
All races 3/, male										
2006.....	2.8	4.5	8.2	12.6	18.4	29.1	45.8	78.8	123.2	204.8
2005.....	2.6	4.8	7.7	12.3	18.6	30.0	48.7	76.1	125.5	210.8
2004.....	2.5	4.8	8.0	12.2	18.0	28.5	49.0	77.9	126.8	214.7
2003.....	2.7	5.0	7.5	12.2	18.8	30.2	47.7	80.4	128.2	223.2
2002.....	2.7	5.1	7.4	11.3	18.6	30.2	50.7	84.9	132.3	222.5
2001.....	2.6	4.7	7.2	11.3	17.6	29.4	48.7	82.3	127.8	233.5
2000.....	2.7	4.8	7.6	10.7	17.4	28.2	45.0	77.3	127.0	233.1
1999.....	2.9	4.6	6.7	10.3	16.9	30.0	46.3	79.3	127.7	246.7
All races 3/, female										
2006.....	2.3	3.9	5.9	9.4	15.8	24.3	38.9	61.9	99.0	175.1
2005.....	2.4	3.9	6.5	10.0	15.3	24.1	38.4	60.3	99.9	186.7
2004.....	2.5	3.9	6.7	9.9	16.4	23.9	38.3	61.4	97.6	185.8
2003.....	2.7	4.1	6.3	10.0	16.5	23.2	39.2	65.0	102.8	203.5
2002.....	2.3	4.0	6.0	10.0	15.2	26.3	40.7	65.1	102.7	205.3
2001.....	2.2	3.9	6.0	9.5	15.1	24.1	37.8	62.0	96.5	205.4
2000.....	2.5	3.6	5.5	9.1	14.6	22.7	36.1	59.6	98.7	218.2
1999.....	2.1	3.6	5.5	9.0	13.7	23.6	35.9	59.3	97.2	225.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Zoonotic and other bacterial diseases (A20-A49)										
White, both sexes										
2006 4/.....	2.1	3.4	5.8	9.1	14.9	23.7	37.9	64.9	102.9	178.3
2005 4/.....	2.0	3.5	5.9	9.6	14.8	24.0	39.6	62.3	103.4	184.0
2004 4/.....	1.9	3.5	6.0	9.5	14.7	22.9	39.4	63.7	102.6	186.1
2003 4/.....	2.2	3.7	5.5	9.4	15.2	22.9	39.0	66.6	107.3	198.2
2002.....	2.1	3.8	5.3	8.9	14.1	24.7	41.0	67.6	107.5	199.4
2001.....	1.9	3.4	5.3	8.5	14.0	23.4	38.4	65.0	102.8	202.9
2000.....	2.2	3.2	5.3	8.2	13.3	21.8	36.0	60.8	101.6	209.6
1999.....	2.0	3.1	4.9	7.8	13.0	23.1	35.6	60.9	100.8	216.1
White, male										
2006 4/.....	2.4	3.8	6.8	10.4	15.8	26.3	41.8	74.1	117.9	201.2
2005 4/.....	2.1	4.0	6.3	10.5	16.5	26.5	44.4	71.1	118.8	199.5
2004 4/.....	2.0	3.8	6.7	10.4	15.2	25.2	44.5	72.9	119.0	207.4
2003 4/.....	2.2	4.2	6.0	10.1	16.2	25.9	43.3	74.9	122.5	212.7
2002.....	2.2	4.3	5.9	9.6	15.9	26.3	46.3	77.5	124.3	211.9
2001.....	2.2	3.7	5.8	9.1	15.3	25.7	43.5	76.2	121.5	221.0
2000.....	2.3	3.7	6.0	9.0	14.8	24.4	41.1	70.0	119.3	219.5
1999.....	2.3	3.5	5.4	8.2	13.9	25.9	40.6	72.4	117.9	230.1
White, female										
2006 4/.....	1.9	3.0	4.8	7.9	14.1	21.3	34.8	58.0	93.4	167.6
2005 4/.....	1.9	3.1	5.5	8.8	13.1	21.7	35.6	55.7	93.7	176.9
2004 4/.....	1.9	3.1	5.4	8.6	14.1	20.9	35.2	56.9	92.4	176.5
2003 4/.....	2.2	3.3	5.0	8.7	14.2	20.1	35.5	60.7	97.9	191.9
2002.....	1.9	3.3	4.8	8.3	12.4	23.3	36.6	60.5	97.3	194.0
2001.....	1.7	3.2	4.8	8.0	12.7	21.4	34.4	57.1	91.5	195.4
2000.....	2.0	2.7	4.6	7.5	12.0	19.5	31.9	54.3	91.0	205.6
1999.....	1.7	2.8	4.5	7.4	12.1	20.6	31.7	52.8	90.6	210.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Zoonotic and other bacterial diseases (A20-A49)										
Black, both sexes										
2006 4/.....	6.0	10.8	17.4	27.6	39.7	57.8	87.9	129.0	191.5	296.3
2005 4/.....	5.8	10.7	17.5	26.2	39.8	57.6	84.7	128.6	209.0	346.4
2004 4/.....	6.7	11.0	18.2	26.6	41.7	59.5	88.2	134.2	202.7	328.8
2003 4/.....	6.6	11.2	18.6	26.9	43.1	62.8	90.6	137.8	197.7	379.6
2002.....	6.1	10.1	17.8	28.1	42.6	63.2	94.6	151.5	212.7	376.3
2001.....	5.9	11.0	17.9	28.6	40.5	59.8	92.3	143.8	198.0	373.1
2000.....	6.1	11.5	17.9	25.4	41.3	61.3	89.6	150.3	221.7	402.3
1999.....	6.6	11.7	16.6	26.7	37.7	61.3	98.2	151.0	219.6	428.2
Black, male										
2006 4/.....	6.7	11.0	21.5	33.8	47.2	63.3	97.5	158.9	215.6	302.9
2005 4/.....	6.0	11.8	20.6	31.6	44.9	70.8	102.2	148.9	244.4	384.8
2004 4/.....	7.2	12.6	20.0	31.8	46.7	67.8	108.1	160.6	254.0	352.7
2003 4/.....	6.9	12.9	21.2	34.1	48.2	78.1	105.1	164.8	241.4	416.0
2002.....	6.8	10.9	21.3	30.2	47.1	73.1	110.5	191.1	265.2	402.0
2001.....	6.1	12.2	20.1	35.2	43.7	69.9	114.1	175.6	243.2	435.7
2000.....	6.1	13.1	23.0	29.0	46.2	71.3	101.0	188.1	256.3	441.7
1999.....	8.1	13.7	19.5	31.2	48.4	75.6	120.1	176.7	286.0	472.2
Black, female										
2006 4/.....	5.4	10.6	14.0	22.5	33.9	53.8	81.4	111.1	178.7	293.5
2005 4/.....	5.6	9.8	14.9	21.7	35.8	47.8	72.7	116.5	190.3	330.9
2004 4/.....	6.3	9.6	16.6	22.4	37.7	53.2	74.7	118.6	175.8	319.4
2003 4/.....	6.4	9.8	16.3	21.0	39.1	51.5	80.9	121.8	174.9	365.4
2002.....	5.5	9.4	14.9	26.3	39.0	55.9	84.0	127.9	185.4	366.4
2001.....	5.7	9.9	16.0	23.2	37.9	52.4	77.8	124.8	175.0	349.8
2000.....	6.1	10.1	13.6	22.4	37.4	54.0	82.1	127.6	204.2	387.6
1999.....	5.3	9.9	14.1	23.0	29.3	50.8	83.7	135.6	186.7	411.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
All races 3/, both sexes										
2006.....	2.4	3.9	6.6	10.3	16.2	25.3	40.3	66.0	103.7	177.3
2005.....	2.3	4.0	6.6	10.5	16.1	25.7	40.8	63.6	104.8	187.3
2004.....	2.3	4.1	6.9	10.3	16.2	24.9	41.2	65.1	103.6	186.7
2003.....	2.6	4.2	6.5	10.3	16.7	25.2	40.9	68.4	107.8	202.5
2002.....	2.3	4.2	6.2	10.0	16.0	27.0	43.1	70.0	109.4	203.0
2001.....	2.3	4.0	6.2	9.8	15.5	25.3	40.9	67.3	103.9	205.9
2000.....	2.5	3.9	6.1	9.4	15.2	24.2	38.4	64.0	105.0	215.7
1999.....	2.2	3.7	5.6	9.0	14.4	24.7	38.3	63.8	102.9	220.7
All races 3/, male										
2006.....	2.6	4.2	7.7	11.7	17.4	27.6	43.9	75.3	117.7	194.8
2005.....	2.4	4.4	7.1	11.5	17.8	28.6	45.9	71.7	119.8	201.9
2004.....	2.4	4.5	7.6	11.4	17.1	26.9	46.6	74.5	120.9	203.9
2003.....	2.5	4.6	7.1	11.3	17.7	28.7	45.1	76.7	122.5	213.8
2002.....	2.5	4.7	6.8	10.6	17.8	29.1	48.2	80.6	127.0	213.9
2001.....	2.5	4.3	6.7	10.6	16.6	27.8	46.7	78.6	122.3	224.3
2000.....	2.6	4.4	7.0	10.1	16.7	26.9	43.2	73.9	121.7	223.0
1999.....	2.6	4.2	6.2	9.7	15.7	27.7	43.7	74.8	120.6	233.5
All races 3/, female										
2006.....	2.2	3.7	5.6	8.9	15.1	23.2	37.4	59.2	94.9	169.2
2005.....	2.3	3.6	6.1	9.5	14.5	23.2	36.7	57.8	95.4	180.6
2004.....	2.3	3.7	6.4	9.3	15.5	23.2	36.8	58.4	92.9	179.0
2003.....	2.7	3.8	6.0	9.4	15.7	22.2	37.5	62.5	98.8	197.5
2002.....	2.2	3.8	5.6	9.5	14.5	25.2	39.0	62.5	98.7	198.3
2001.....	2.1	3.8	5.7	9.1	14.4	23.1	36.2	59.5	92.9	198.3
2000.....	2.3	3.4	5.2	8.7	13.8	21.8	34.6	57.1	95.2	212.7
1999.....	1.9	3.3	5.1	8.4	13.1	22.1	34.1	56.2	92.5	215.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
White, both sexes										
2006 4/.....	2.0	3.2	5.4	8.6	14.1	22.5	36.2	61.9	98.3	171.0
2005 4/.....	1.9	3.3	5.4	9.0	14.0	22.9	37.5	58.9	98.3	177.3
2004 4/.....	1.8	3.2	5.7	8.8	13.8	21.9	37.6	60.7	97.5	178.1
2003 4/.....	2.1	3.4	5.2	8.8	14.3	21.7	36.9	63.7	102.5	191.1
2002.....	1.9	3.6	5.0	8.4	13.5	23.6	39.2	64.4	103.1	191.9
2001.....	1.8	3.2	4.9	8.1	13.2	22.2	36.8	62.1	98.5	195.3
2000.....	2.0	3.0	4.9	7.8	12.6	20.7	34.5	58.0	97.4	203.0
1999.....	1.7	2.8	4.5	7.2	12.1	21.3	33.6	57.6	95.5	205.9
White, male										
2006 4/.....	2.2	3.5	6.3	9.7	14.8	24.8	40.0	70.8	112.6	190.7
2005 4/.....	2.0	3.6	5.7	9.7	15.8	25.2	41.8	66.7	113.0	190.7
2004 4/.....	1.8	3.5	6.4	9.7	14.4	23.7	42.5	69.8	113.4	196.5
2003 4/.....	2.0	3.8	5.7	9.3	15.3	24.6	40.8	71.3	116.7	203.3
2002.....	2.0	4.0	5.4	8.9	15.3	25.2	44.0	73.4	119.0	202.9
2001.....	2.1	3.3	5.3	8.5	14.4	24.4	41.6	72.5	116.0	211.8
2000.....	2.2	3.5	5.5	8.4	14.0	23.1	39.4	66.7	114.0	209.8
1999.....	2.0	3.1	4.9	7.6	12.8	23.7	38.2	68.3	111.6	217.4
White, female										
2006 4/.....	1.8	2.8	4.6	7.5	13.4	20.4	33.2	55.2	89.3	161.8
2005 4/.....	1.8	2.9	5.1	8.3	12.4	20.8	33.9	53.1	89.2	171.0
2004 4/.....	1.8	2.9	5.1	8.0	13.3	20.2	33.5	54.1	87.7	169.9
2003 4/.....	2.2	3.0	4.7	8.2	13.4	19.2	33.7	58.2	93.7	185.8
2002.....	1.9	3.1	4.5	7.8	11.8	22.3	35.2	58.0	93.4	187.1
2001.....	1.6	3.0	4.5	7.7	12.2	20.4	32.9	54.8	87.9	188.3
2000.....	1.9	2.5	4.3	7.2	11.3	18.7	30.6	51.9	87.6	200.2
1999.....	1.4	2.4	4.1	6.9	11.5	19.2	29.9	50.1	86.1	201.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Septicemia (A40-A41)										
Black, both sexes										
2006 4/.....	5.7	10.0	16.8	26.0	38.4	55.7	85.8	124.8	186.2	287.5
2005 4/.....	5.5	10.0	16.6	25.5	38.4	55.8	81.2	125.4	203.7	336.8
2004 4/.....	6.2	10.3	17.3	25.5	39.9	57.9	85.1	128.0	196.1	320.3
2003 4/.....	6.3	10.5	17.5	25.3	41.5	61.0	88.0	133.4	193.4	372.0
2002.....	5.5	9.5	17.0	27.2	40.9	62.0	90.6	146.5	208.8	367.0
2001.....	5.6	10.6	16.9	27.1	38.4	57.4	89.6	139.1	193.1	364.8
2000.....	5.8	11.0	16.9	24.3	40.2	59.7	86.6	146.4	216.3	392.6
1999.....	6.2	11.0	15.5	25.4	36.3	57.9	94.6	142.8	209.5	412.3
Black, male										
2006 4/.....	6.4	10.1	20.4	31.8	45.4	61.6	94.4	153.0	209.5	296.7
2005 4/.....	5.5	10.9	19.5	30.5	43.4	68.2	97.1	143.5	238.8	374.6
2004 4/.....	6.7	12.0	19.0	30.4	44.5	65.2	102.3	153.8	244.6	341.7
2003 4/.....	6.4	11.8	19.7	31.8	46.1	74.9	100.2	160.2	235.5	404.5
2002.....	6.1	10.3	20.0	29.0	45.1	71.6	105.5	183.7	259.8	393.5
2001.....	5.7	11.6	19.1	33.3	41.5	66.3	111.1	171.0	236.8	424.4
2000.....	5.8	12.4	21.3	27.8	45.1	68.9	97.0	182.9	250.4	427.8
1999.....	7.7	12.8	18.1	29.7	46.5	71.4	115.0	166.2	266.8	453.5
Black, female										
2006 4/.....	5.2	9.9	13.7	21.2	32.9	51.4	79.9	107.9	174.0	283.7
2005 4/.....	5.5	9.2	14.1	21.3	34.4	46.5	70.3	114.7	185.1	321.5
2004 4/.....	5.7	8.9	15.8	21.5	36.4	52.5	73.4	112.7	170.5	311.9
2003 4/.....	6.3	9.4	15.6	20.0	37.9	50.8	79.8	117.5	171.4	359.3
2002.....	5.0	8.9	14.4	25.7	37.6	54.9	80.7	124.3	182.2	356.9
2001.....	5.4	9.7	15.0	22.1	36.0	50.9	75.3	120.1	170.9	342.7
2000.....	5.8	9.7	13.2	21.5	36.3	52.8	79.7	124.5	199.1	379.5
1999.....	5.0	9.4	13.3	21.9	28.3	47.9	81.0	128.7	181.1	396.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

All races 3/, both sexes

2006.....	0.1	0.3	0.4	0.6	0.9	1.3	1.6	3.0	4.6	7.2
2005.....	0.1	0.3	0.5	0.6	0.8	1.1	2.2	3.3	4.9	6.8
2004.....	0.1	0.3	0.3	0.7	0.9	1.1	1.9	3.1	5.1	7.8
2003.....	0.1	0.3	0.3	0.7	0.9	1.2	2.1	3.0	4.6	7.0
2002.....	0.2	0.2	0.4	0.5	0.7	1.0	2.0	3.2	4.3	7.4
2001.....	0.1	0.2	0.4	0.5	0.8	1.2	1.6	2.9	4.3	7.5
2000.....	0.1	0.2	0.4	0.5	0.7	1.0	1.5	2.8	4.0	6.7
1999.....	0.2	0.3	0.4	0.6	0.8	1.8	2.2	3.5	5.4	10.3

All races 3/, male

2006.....	0.2	0.3	0.5	0.8	1.0	1.5	1.8	3.5	5.5	10.0
2005.....	0.2	0.4	0.6	0.8	0.8	1.4	2.8	4.5	5.7	8.8
2004.....	*	0.3	0.4	0.8	0.9	1.5	2.3	3.4	5.9	10.8
2003.....	*	0.4	0.4	0.9	1.0	1.5	2.6	3.7	5.6	9.3
2002.....	0.2	0.3	0.5	0.7	0.8	1.1	2.4	4.2	5.2	8.5
2001.....	*	0.3	0.5	0.6	1.0	1.5	1.9	3.6	5.4	8.9
2000.....	*	0.3	0.6	0.6	0.8	1.4	1.8	3.3	5.2	9.9
1999.....	0.2	0.4	0.4	0.6	1.1	2.2	2.6	4.4	7.0	13.1

All races 3/, female

2006.....	*	0.3	0.2	0.4	0.7	1.1	1.5	2.7	4.0	5.9
2005.....	*	0.2	0.4	0.4	0.7	0.9	1.6	2.4	4.4	6.0
2004.....	*	0.2	0.3	0.5	0.9	0.7	1.5	2.9	4.7	6.4
2003.....	*	0.2	0.3	0.5	0.8	0.9	1.6	2.5	4.0	5.9
2002.....	*	0.2	0.3	0.4	0.7	1.0	1.6	2.5	3.8	6.9
2001.....	*	*	0.3	0.3	0.7	1.0	1.5	2.4	3.6	6.9
2000.....	*	*	*	0.3	0.7	0.8	1.3	2.4	3.4	5.3
1999.....	0.2	0.3	0.4	0.6	0.6	1.4	1.8	2.9	4.5	9.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

White, both sexes

2006 4/.....	0.1	0.2	0.4	0.5	0.8	1.2	1.6	3.0	4.5	7.2
2005 4/.....	0.1	0.2	0.4	0.6	0.7	1.1	2.0	3.4	5.0	6.6
2004 4/.....	*	0.2	0.3	0.6	0.9	1.0	1.8	2.9	5.1	7.7
2003 4/.....	*	0.3	0.2	0.6	0.8	1.1	2.0	2.9	4.7	7.0
2002.....	0.1	0.2	0.3	0.5	0.6	1.0	1.8	3.1	4.3	7.3
2001.....	*	0.2	0.3	0.4	0.7	1.1	1.5	2.8	4.2	7.4
2000.....	*	0.2	0.3	0.4	0.7	1.0	1.4	2.7	4.0	6.4
1999.....	0.2	0.3	0.4	0.5	0.8	1.6	2.0	3.1	5.1	9.9

White, male

2006 4/.....	*	*	0.5	0.7	0.9	1.5	1.7	3.3	5.3	10.5
2005 4/.....	*	0.3	0.5	0.7	0.7	1.3	2.6	4.5	5.9	8.7
2004 4/.....	*	0.3	0.3	0.7	0.8	1.4	2.0	3.1	5.6	10.9
2003 4/.....	*	0.3	*	0.8	0.9	1.3	2.4	3.6	5.7	9.3
2002.....	*	0.3	0.4	0.6	0.6	1.1	2.2	4.1	5.1	8.7
2001.....	*	0.3	0.4	0.5	0.9	1.3	1.8	3.6	5.4	8.9
2000.....	*	0.2	0.5	0.6	0.7	1.2	1.6	3.1	5.2	9.5
1999.....	*	0.3	0.4	0.5	1.1	2.0	2.4	4.0	6.2	12.6

White, female

2006 4/.....	*	0.2	0.2	0.4	0.8	0.9	1.5	2.7	4.0	5.7
2005 4/.....	*	*	0.4	0.5	0.7	0.9	1.6	2.6	4.4	5.7
2004 4/.....	*	*	*	0.5	0.9	0.7	1.6	2.7	4.7	6.3
2003 4/.....	*	0.2	0.2	0.5	0.8	0.9	1.7	2.4	4.1	6.0
2002.....	*	*	0.3	0.4	0.6	1.0	1.4	2.4	3.7	6.7
2001.....	*	*	*	*	0.6	0.9	1.4	2.1	3.5	6.8
2000.....	*	*	*	*	0.6	0.7	1.2	2.4	3.2	5.1
1999.....	*	*	0.3	0.5	0.6	1.3	1.8	2.5	4.5	8.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

Black, both sexes

2006 4/.....	*	*	*	1.6	*	2.1	*	4.2	5.0	8.8
2005 4/.....	*	0.7	0.9	*	*	*	3.5	*	5.3	9.6
2004 4/.....	*	*	*	*	1.7	*	3.0	6.3	6.7	8.2
2003 4/.....	*	*	*	1.3	*	*	2.6	4.4	*	7.6
2002.....	*	*	*	*	*	*	4.0	5.0	*	9.2
2001.....	*	*	1.0	1.4	*	2.3	2.7	4.5	*	8.0
2000.....	*	*	*	*	*	*	2.9	3.9	*	9.7
1999.....	*	*	*	*	*	3.4	3.5	8.0	10.1	15.9

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	10.0	19.2	*

Black, female

2006 4/.....	*	*	*	*	*	*	*	*	*	9.9
2005 4/.....	*	*	*	*	*	*	*	*	*	9.4
2004 4/.....	*	*	*	*	*	*	*	5.9	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	9.4
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	6.9	*	14.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral diseases (A80-B34)										
All races 3/, both sexes										
2006.....	12.7	15.8	18.5	14.9	9.8	8.4	7.8	8.0	7.4	7.8
2005.....	13.5	15.8	16.9	12.1	8.4	7.2	7.0	6.6	7.3	7.9
2004.....	14.5	16.1	16.4	11.7	8.4	7.5	6.5	6.5	6.6	7.8
2003.....	15.1	17.2	16.2	11.4	8.1	7.4	8.0	7.6	7.5	9.1
2002.....	15.9	18.4	15.9	10.6	8.3	7.9	7.8	7.8	7.7	10.1
2001.....	16.5	17.5	14.2	10.3	9.1	8.1	7.6	7.3	7.0	8.3
2000.....	16.1	18.1	14.2	10.1	8.7	7.9	8.0	8.1	8.1	9.6
1999.....	17.2	17.1	12.9	9.9	7.5	8.2	8.1	8.0	8.7	11.6
All races 3/, male										
2006.....	18.3	23.1	28.1	22.8	14.6	11.2	9.3	9.5	8.6	8.1
2005.....	19.7	23.8	25.8	19.1	12.3	9.7	8.8	7.6	8.3	7.7
2004.....	20.9	24.1	25.1	18.6	12.5	10.5	8.1	7.7	6.7	9.0
2003.....	22.2	26.1	25.4	17.8	11.8	9.7	10.5	9.2	9.6	10.2
2002.....	23.2	28.2	25.5	16.3	12.1	10.7	9.2	10.0	9.1	11.7
2001.....	24.4	27.3	22.9	16.1	12.5	10.0	9.1	8.3	8.9	9.3
2000.....	24.0	28.4	22.7	15.7	12.4	10.2	9.5	10.7	9.9	10.8
1999.....	26.2	27.0	20.8	15.5	10.5	10.7	9.8	9.3	10.9	13.3
All races 3/, female										
2006.....	7.1	8.7	9.3	7.5	5.4	6.0	6.6	6.9	6.6	7.7
2005.....	7.3	8.0	8.4	5.5	4.7	5.0	5.5	5.8	6.7	8.0
2004.....	8.2	8.3	8.0	5.3	4.7	4.9	5.2	5.5	6.5	7.2
2003.....	8.1	8.4	7.3	5.4	4.7	5.4	6.0	6.4	6.2	8.7
2002.....	8.8	9.0	6.7	5.2	4.8	5.4	6.7	6.3	6.9	9.3
2001.....	8.7	8.1	5.9	4.8	6.0	6.4	6.5	6.7	5.9	7.9
2000.....	8.3	8.1	6.0	4.8	5.3	6.0	6.9	6.3	7.0	9.1
1999.....	8.3	7.5	5.3	4.7	4.7	6.0	6.7	7.1	7.4	10.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral diseases (A80-B34)										
White, both sexes										
2006 4/.....	7.9	10.7	14.0	10.8	7.4	6.3	6.6	7.1	6.8	7.7
2005 4/.....	8.6	10.4	12.0	8.6	5.9	5.4	5.6	5.8	6.9	8.1
2004 4/.....	9.0	10.7	11.3	8.2	5.9	5.7	5.3	5.9	6.2	7.8
2003 4/.....	9.6	11.8	11.2	7.9	5.6	5.5	6.9	6.9	7.0	9.3
2002.....	9.8	12.5	10.8	7.5	6.0	5.9	6.5	7.0	7.4	10.3
2001.....	10.2	11.7	9.5	7.4	6.4	6.3	6.5	6.8	6.6	8.4
2000.....	10.0	12.0	9.3	6.9	6.2	6.2	6.8	7.6	7.8	9.6
1999.....	10.4	10.9	8.7	6.6	5.5	6.4	6.8	7.4	8.6	11.6
White, male										
2006 4/.....	12.5	16.8	21.6	16.7	10.9	8.2	7.6	8.2	8.3	8.2
2005 4/.....	13.6	16.6	18.6	13.6	8.5	7.1	7.0	6.9	7.8	8.0
2004 4/.....	14.2	16.7	17.9	13.0	8.6	7.9	6.5	7.2	6.2	8.8
2003 4/.....	15.2	18.8	18.0	12.3	8.1	7.0	8.8	8.1	8.9	10.3
2002.....	15.4	19.9	17.4	11.6	8.4	7.8	7.5	8.9	8.6	11.7
2001.....	16.5	19.0	15.9	11.7	8.5	7.8	7.9	7.7	8.3	9.7
2000.....	15.8	19.6	15.3	10.8	8.6	7.6	7.6	9.9	9.3	10.9
1999.....	16.8	17.4	14.3	10.2	7.9	8.3	8.1	8.6	10.9	12.7
White, female										
2006 4/.....	3.3	4.5	6.5	5.2	4.0	4.7	5.8	6.3	5.9	7.5
2005 4/.....	3.5	4.2	5.5	3.8	3.4	3.9	4.5	5.0	6.3	8.2
2004 4/.....	3.8	4.7	4.8	3.5	3.4	3.8	4.4	5.0	6.2	7.3
2003 4/.....	4.0	4.8	4.4	3.6	3.3	4.2	5.3	6.1	5.8	8.9
2002.....	4.1	5.1	4.3	3.6	3.8	4.1	5.8	5.7	6.7	9.7
2001.....	4.0	4.5	3.2	3.3	4.3	5.0	5.4	6.1	5.6	7.9
2000.....	4.3	4.5	3.5	3.2	4.0	5.0	6.1	5.9	6.8	9.1
1999.....	4.0	4.4	3.1	3.2	3.4	4.9	5.7	6.6	7.2	11.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral diseases (A80-B34)										
Black, both sexes										
2006 4/.....	46.6	54.6	56.1	50.8	31.9	26.3	16.4	13.6	12.3	8.3
2005 4/.....	49.1	56.4	58.0	43.1	31.8	22.9	17.5	13.5	10.9	6.4
2004 4/.....	53.7	57.6	58.0	44.1	31.7	22.5	15.2	10.7	9.9	7.3
2003 4/.....	55.5	58.9	58.0	43.7	30.4	22.9	17.6	13.1	12.7	7.9
2002.....	60.5	65.2	59.0	38.4	27.5	24.2	18.4	16.3	9.4	7.4
2001.....	62.3	63.5	54.4	36.3	32.7	21.8	15.9	12.9	10.0	6.1
2000.....	60.6	66.4	55.8	36.7	27.8	21.0	19.3	13.7	11.3	8.8
1999.....	66.5	66.5	49.7	39.3	23.0	22.5	19.6	12.7	9.0	11.5
Black, male										
2006 4/.....	63.3	74.7	86.5	81.2	52.3	39.8	25.8	23.3	13.7	*
2005 4/.....	68.4	82.1	90.1	72.4	52.2	36.6	26.9	16.5	*	*
2004 4/.....	72.4	85.6	88.9	73.4	52.5	36.8	24.0	12.7	*	*
2003 4/.....	78.3	87.4	92.8	72.9	49.4	36.4	28.6	18.7	16.2	*
2002.....	83.9	97.6	98.4	63.1	45.7	39.6	27.7	24.3	*	*
2001.....	87.8	97.7	86.4	60.2	51.5	31.5	21.9	17.2	*	*
2000.....	88.6	104.1	91.6	61.9	46.0	33.8	30.8	19.8	18.5	*
1999.....	100.2	108.7	82.0	66.3	35.9	33.2	28.8	15.7	*	*
Black, female										
2006 4/.....	31.9	37.1	30.5	25.7	16.0	16.3	9.9	7.8	11.6	8.8
2005 4/.....	32.1	34.1	30.9	19.0	15.9	12.7	11.0	11.7	10.0	7.5
2004 4/.....	37.1	33.3	31.9	20.0	15.4	11.9	9.2	9.4	8.7	*
2003 4/.....	35.5	34.1	28.6	19.7	15.6	12.8	10.2	9.8	10.8	*
2002.....	39.9	37.1	25.6	18.1	13.2	12.8	12.1	11.5	*	*
2001.....	39.7	33.9	27.1	16.7	17.9	14.6	11.9	10.3	8.2	*
2000.....	35.9	33.7	25.3	16.1	13.4	11.5	11.8	10.0	*	8.6
1999.....	36.8	30.1	22.2	17.3	13.0	14.6	13.5	10.9	8.6	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)										
All races 3/, both sexes										
2006.....	1.8	4.8	8.9	8.0	5.0	4.5	4.5	5.0	4.0	3.1
2005.....	1.7	4.2	6.8	5.3	3.5	3.2	3.8	3.6	3.8	2.4
2004.....	2.0	4.3	6.5	4.9	3.5	3.3	3.6	3.6	3.6	2.7
2003.....	2.1	4.8	6.3	4.5	3.1	3.4	4.3	4.3	4.1	2.4
2002.....	2.3	5.6	6.1	4.3	3.7	3.9	4.5	4.7	4.4	3.9
2001.....	2.6	5.3	5.2	4.1	3.8	4.3	4.9	5.1	4.6	3.8
2000.....	2.5	5.2	4.8	3.7	3.8	4.3	4.9	5.4	5.2	4.0
1999.....	2.3	4.4	3.8	3.3	3.3	4.5	5.1	5.3	5.4	4.7
All races 3/, male										
2006.....	2.5	6.8	13.4	11.9	7.0	5.2	4.7	5.1	3.8	3.0
2005.....	2.3	6.3	10.2	8.2	4.7	3.5	4.0	3.3	4.2	2.6
2004.....	2.8	6.4	9.9	7.3	4.5	4.1	3.7	3.8	3.3	3.0
2003.....	2.9	7.1	9.7	6.6	3.9	3.7	4.8	4.7	5.0	2.8
2002.....	3.2	8.5	9.5	6.2	4.6	4.3	4.7	5.7	4.8	5.0
2001.....	3.7	8.1	8.2	5.9	4.5	4.7	4.9	5.1	5.5	4.4
2000.....	3.5	8.0	7.4	5.2	4.7	4.6	5.0	6.5	6.2	5.7
1999.....	3.5	6.7	5.9	4.7	3.8	4.9	5.4	6.1	7.3	6.9
All races 3/, female										
2006.....	1.1	2.8	4.6	4.2	3.3	3.8	4.3	5.0	4.1	3.1
2005.....	1.1	2.2	3.6	2.7	2.3	2.9	3.7	3.8	3.6	2.3
2004.....	1.3	2.3	3.3	2.5	2.6	2.6	3.5	3.4	3.7	2.5
2003.....	1.2	2.6	3.1	2.5	2.4	3.1	3.9	4.0	3.5	2.3
2002.....	1.4	2.8	2.8	2.5	2.9	3.4	4.4	4.1	4.2	3.3
2001.....	1.5	2.5	2.4	2.5	3.3	3.9	4.9	5.0	4.1	3.5
2000.....	1.5	2.5	2.4	2.2	3.1	4.1	4.9	4.6	4.7	3.3
1999.....	1.2	2.2	1.7	2.1	2.8	4.1	4.9	4.7	4.2	3.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)										
White, both sexes										
2006 4/.....	1.8	4.9	8.8	7.2	4.5	3.8	4.1	4.5	3.5	2.8
2005 4/.....	1.7	4.3	6.7	4.9	3.0	2.7	3.2	3.3	3.6	2.2
2004 4/.....	2.1	4.3	6.2	4.2	3.1	3.0	3.2	3.3	3.1	2.5
2003 4/.....	2.1	4.9	6.0	3.8	2.7	2.9	3.8	4.0	3.8	2.3
2002.....	2.2	5.5	5.8	3.8	3.2	3.4	4.1	4.5	4.3	3.7
2001.....	2.6	5.2	4.7	3.7	3.5	3.8	4.4	4.9	4.4	3.6
2000.....	2.4	4.9	4.3	3.0	3.4	3.9	4.6	5.3	5.0	3.9
1999.....	2.2	4.2	3.3	2.9	2.9	4.0	4.6	4.9	5.3	4.5
White, male										
2006 4/.....	2.5	7.0	13.3	10.8	6.2	4.4	4.3	4.6	3.6	3.1
2005 4/.....	2.4	6.4	10.0	7.6	4.0	3.1	3.4	3.3	3.9	2.4
2004 4/.....	2.9	6.4	9.4	6.4	4.0	3.7	3.4	3.6	2.8	2.7
2003 4/.....	2.9	7.2	9.2	5.7	3.4	3.3	4.3	4.2	4.8	2.8
2002.....	3.1	8.3	9.1	5.5	3.9	4.0	4.3	5.5	4.8	4.9
2001.....	3.7	8.1	7.6	5.4	4.1	4.4	4.6	5.2	5.6	4.5
2000.....	3.4	7.6	6.6	4.4	4.2	4.0	4.6	6.4	5.9	5.7
1999.....	3.3	6.1	5.2	4.1	3.6	4.6	5.1	5.8	7.7	6.5
White, female										
2006 4/.....	1.1	2.8	4.5	3.8	2.9	3.2	3.9	4.3	3.5	2.7
2005 4/.....	1.1	2.2	3.4	2.4	2.0	2.4	3.0	3.2	3.4	2.2
2004 4/.....	1.3	2.3	3.0	2.2	2.4	2.4	3.0	3.1	3.3	2.4
2003 4/.....	1.2	2.6	2.9	2.0	2.0	2.6	3.5	3.8	3.2	2.1
2002.....	1.3	2.6	2.7	2.2	2.6	2.8	3.9	3.8	4.0	3.2
2001.....	1.4	2.4	2.0	2.1	2.9	3.3	4.2	4.6	3.7	3.2
2000.....	1.4	2.3	2.0	1.8	2.7	3.7	4.5	4.5	4.4	3.2
1999.....	1.1	2.2	1.5	1.7	2.3	3.5	4.3	4.3	3.9	3.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Viral hepatitis (B15-B19)										
Black, both sexes										
2006 4/.....	1.8	4.8	11.0	14.5	9.0	8.9	5.0	7.0	6.9	*
2005 4/.....	1.7	4.7	9.7	9.2	7.2	5.8	6.4	5.5	4.9	*
2004 4/.....	2.0	5.2	10.3	10.2	6.5	4.2	5.1	3.9	6.7	*
2003 4/.....	2.7	5.2	10.1	9.4	6.3	5.1	5.7	6.5	5.3	*
2002.....	2.8	7.2	9.0	7.7	6.3	6.1	6.2	6.0	*	*
2001.....	3.3	6.8	9.4	7.3	6.0	5.1	6.2	6.1	*	*
2000.....	3.1	7.3	8.8	6.9	4.6	5.2	6.3	5.7	7.1	*
1999.....	3.5	7.0	6.7	5.7	4.2	6.6	7.2	6.8	*	6.4
Black, male										
2006 4/.....	2.3	6.3	16.2	23.5	13.8	11.2	6.5	9.4	*	*
2005 4/.....	2.2	6.7	14.5	14.3	10.5	6.7	7.9	*	*	*
2004 4/.....	2.2	8.3	16.0	16.6	9.9	6.8	6.5	*	*	*
2003 4/.....	3.6	7.6	16.2	14.9	7.5	5.4	6.9	*	*	*
2002.....	3.8	10.5	14.6	12.4	8.1	8.0	*	*	*	*
2001.....	4.7	10.1	13.9	10.2	7.3	*	*	*	*	*
2000.....	4.2	11.0	13.9	10.4	5.2	6.8	7.4	*	*	*
1999.....	5.1	11.9	11.2	7.6	4.5	5.5	*	*	*	*
Black, female										
2006 4/.....	1.4	3.4	6.6	7.1	5.2	7.1	*	5.6	7.6	*
2005 4/.....	*	3.0	5.7	5.0	4.7	5.1	5.4	7.5	*	*
2004 4/.....	1.8	2.5	5.5	4.9	3.9	*	*	*	*	*
2003 4/.....	1.9	3.2	5.0	4.8	5.3	4.9	4.9	5.7	*	*
2002.....	1.9	4.4	4.2	3.9	4.8	4.6	6.3	5.8	*	*
2001.....	2.0	3.9	5.5	4.9	5.0	5.4	6.7	7.2	*	*
2000.....	2.1	4.0	4.4	4.0	4.1	4.0	5.5	*	*	*
1999.....	2.0	2.8	2.8	4.2	4.0	7.5	7.7	6.6	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)										
All races 3/, both sexes										
2006.....	10.8	10.9	9.3	6.5	4.1	3.1	1.8	1.0	0.6	*
2005.....	11.7	11.3	9.9	6.3	4.0	2.8	1.8	1.0	0.5	*
2004.....	12.3	11.5	9.6	6.3	4.2	3.1	1.5	1.1	0.5	*
2003.....	12.8	12.1	9.5	6.4	4.1	2.8	1.9	1.0	0.4	*
2002.....	13.4	12.6	9.6	5.9	4.0	2.9	1.4	1.0	0.5	*
2001.....	13.7	12.1	8.7	5.8	4.5	2.7	1.4	0.9	0.4	*
2000.....	13.5	12.7	9.0	5.9	4.1	2.6	1.7	0.9	*	*
1999.....	14.6	12.4	8.7	5.9	3.5	2.7	1.5	0.7	*	*
All races 3/, male										
2006.....	15.6	16.1	14.4	10.3	6.9	5.0	3.1	1.9	1.1	*
2005.....	17.2	17.2	15.4	10.5	6.6	4.8	3.1	1.8	*	*
2004.....	17.9	17.5	14.9	10.5	7.0	5.1	2.6	1.7	*	*
2003.....	19.1	18.8	15.3	10.7	6.9	4.6	3.3	1.8	1.0	*
2002.....	19.8	19.4	15.7	9.8	6.8	5.0	2.6	1.7	*	*
2001.....	20.5	19.0	14.4	9.8	7.1	4.2	2.7	1.8	*	*
2000.....	20.4	20.3	15.0	10.1	7.0	4.5	2.9	1.8	*	*
1999.....	22.6	20.0	14.5	10.2	5.9	4.8	2.7	1.3	*	*
All races 3/, female										
2006.....	5.9	5.7	4.3	2.8	1.5	1.4	0.8	*	*	*
2005.....	6.2	5.6	4.6	2.4	1.6	1.1	0.7	*	*	*
2004.....	6.7	5.7	4.5	2.3	1.5	1.4	0.6	0.6	*	*
2003.....	6.6	5.6	3.9	2.5	1.6	1.3	0.7	*	*	*
2002.....	7.2	5.9	3.6	2.3	1.5	1.0	0.5	*	*	*
2001.....	7.0	5.4	3.3	2.0	2.0	1.4	0.4	*	*	*
2000.....	6.7	5.3	3.4	2.0	1.5	1.0	0.7	*	*	*
1999.....	6.9	5.0	3.1	1.9	1.2	1.0	0.6	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)										
White, both sexes										
2006 4/.....	6.0	5.6	4.8	3.1	2.2	1.6	0.9	0.5	0.4	*
2005 4/.....	6.7	5.9	5.1	3.2	1.9	1.4	0.9	0.6	*	*
2004 4/.....	6.8	6.1	4.8	3.3	2.0	1.5	0.8	0.6	*	*
2003 4/.....	7.4	6.7	4.9	3.5	2.1	1.3	1.0	0.6	*	*
2002.....	7.3	6.8	4.6	3.3	2.2	1.3	0.6	0.5	*	*
2001.....	7.5	6.4	4.5	3.3	2.0	1.2	0.8	0.5	*	*
2000.....	7.5	6.9	4.7	3.4	2.1	1.3	0.7	0.5	*	*
1999.....	8.1	6.5	4.9	3.1	1.9	1.4	0.7	0.4	*	*
White, male										
2006 4/.....	9.8	9.7	8.0	5.4	4.0	2.8	1.6	1.0	*	*
2005 4/.....	11.0	10.0	8.4	5.6	3.4	2.5	1.7	1.1	*	*
2004 4/.....	11.2	10.1	8.2	5.9	3.7	2.7	1.5	1.2	*	*
2003 4/.....	12.1	11.3	8.4	6.0	3.6	2.2	2.0	1.3	*	*
2002.....	12.0	11.4	8.0	5.8	3.8	2.4	1.2	0.8	*	*
2001.....	12.6	10.8	8.0	5.9	3.4	2.1	1.7	1.0	*	*
2000.....	12.3	11.7	8.3	6.0	3.7	2.5	1.3	1.0	*	*
1999.....	13.4	11.0	8.6	5.6	3.5	2.7	1.2	0.8	*	*
White, female										
2006 4/.....	2.1	1.6	1.7	0.9	0.5	0.5	*	*	*	*
2005 4/.....	2.4	1.9	1.9	0.9	0.5	*	*	*	*	*
2004 4/.....	2.4	2.1	1.5	0.8	0.4	0.5	*	*	*	*
2003 4/.....	2.6	2.1	1.3	1.1	0.6	0.5	*	*	*	*
2002.....	2.6	2.2	1.3	0.9	0.7	*	*	*	*	*
2001.....	2.4	2.0	1.0	0.9	0.7	0.5	*	*	*	*
2000.....	2.7	2.1	1.2	0.9	0.7	*	*	*	*	*
1999.....	2.8	2.0	1.2	0.7	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease (B20-B24)										
Black, both sexes										
2006 4/.....	44.6	49.5	44.4	35.8	22.1	16.7	10.8	5.9	*	*
2005 4/.....	47.2	51.2	47.8	33.8	24.0	16.6	10.5	6.0	*	*
2004 4/.....	51.4	52.1	47.4	33.4	24.6	17.6	8.5	6.3	*	*
2003 4/.....	52.3	53.4	47.2	33.9	23.8	17.2	11.0	4.9	*	*
2002.....	57.3	57.4	49.7	30.2	20.7	17.5	10.5	7.0	*	*
2001.....	58.8	56.5	44.7	28.9	26.2	16.3	8.5	6.4	*	*
2000.....	57.3	58.7	46.6	29.3	22.0	15.1	11.6	6.2	*	*
1999.....	62.8	59.0	42.6	32.7	18.3	15.4	11.2	4.8	*	*
Black, male										
2006 4/.....	60.6	68.2	69.1	57.1	37.3	27.2	19.4	12.5	*	*
2005 4/.....	65.8	74.8	75.4	57.8	40.7	29.7	18.7	11.6	*	*
2004 4/.....	69.9	76.9	72.4	56.1	42.0	29.6	15.2	9.6	*	*
2003 4/.....	74.0	79.5	75.7	57.6	41.3	30.0	20.0	*	*	*
2002.....	79.6	86.3	83.5	50.0	37.0	31.4	19.4	14.7	*	*
2001.....	82.8	87.5	72.2	49.9	43.5	26.4	15.5	13.0	*	*
2000.....	84.1	92.7	77.4	51.0	39.5	26.3	22.4	11.8	*	*
1999.....	94.9	96.1	70.6	57.5	30.6	27.3	21.4	*	*	*
Black, female										
2006 4/.....	30.5	33.2	23.5	18.2	10.3	9.0	4.9	*	*	*
2005 4/.....	30.8	30.6	24.6	14.0	10.9	6.9	5.0	*	*	*
2004 4/.....	35.0	30.5	26.3	14.7	10.9	8.6	*	*	*	*
2003 4/.....	33.1	30.7	23.1	14.3	10.2	7.7	4.9	*	*	*
2002.....	37.6	32.4	21.0	13.9	7.9	7.2	4.5	*	*	*
2001.....	37.5	29.7	21.2	11.7	12.6	8.8	*	*	*	*
2000.....	33.7	29.2	20.4	11.5	8.2	6.9	4.4	*	*	*
1999.....	34.6	26.9	18.8	12.5	8.7	6.5	4.5	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

All races 3/, both sexes

2006.....	4.0	4.2	3.8	2.3	1.5	1.0	0.4	0.3	*	*
2005.....	4.4	4.3	3.9	2.5	1.3	1.0	0.5	0.3	*	*
2004.....	5.0	4.6	3.7	2.4	1.3	1.2	0.5	0.4	*	*
2003.....	4.8	4.7	3.8	2.4	1.5	0.9	0.7	*	*	*
2002.....	5.3	5.1	3.9	2.3	1.4	1.0	0.4	0.4	*	*
2001.....	5.3	4.9	3.4	2.2	1.5	0.9	0.6	0.3	*	*
2000.....	5.4	5.2	3.5	2.0	1.4	0.9	0.6	0.4	*	*
1999.....	5.8	5.2	3.6	1.9	1.1	0.9	0.6	0.4	*	*

All races 3/, male

2006.....	5.8	6.0	5.9	3.7	2.5	1.5	0.7	*	*	*
2005.....	6.4	6.7	5.7	4.2	2.3	1.6	0.7	*	*	*
2004.....	7.3	6.9	5.5	3.9	2.1	2.0	0.9	0.7	*	*
2003.....	7.1	7.5	6.2	3.8	2.5	1.5	1.0	*	*	*
2002.....	7.9	7.8	6.5	3.7	2.2	1.7	0.9	0.7	*	*
2001.....	8.0	7.6	5.6	3.7	2.3	1.2	1.1	*	*	*
2000.....	7.9	8.2	5.8	3.3	2.4	1.7	1.1	*	*	*
1999.....	8.9	8.2	5.9	3.4	1.9	1.6	1.0	0.7	*	*

All races 3/, female

2006.....	2.2	2.4	1.7	0.9	0.6	0.5	*	*	*	*
2005.....	2.4	2.0	2.1	0.9	0.5	0.4	*	*	*	*
2004.....	2.7	2.4	1.9	0.9	0.5	0.6	*	*	*	*
2003.....	2.4	2.0	1.6	1.1	0.6	0.4	*	*	*	*
2002.....	2.7	2.4	1.4	0.9	0.7	0.4	*	*	*	*
2001.....	2.7	2.3	1.3	0.8	0.8	0.5	*	*	*	*
2000.....	2.9	2.2	1.2	0.9	0.5	*	*	*	*	*
1999.....	2.7	2.2	1.3	0.6	0.5	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

White, both sexes

2006 4/.....	2.4	2.2	2.0	1.1	0.8	0.4	0.3	*	*	*
2005 4/.....	2.7	2.3	2.1	1.3	0.7	0.4	*	*	*	*
2004 4/.....	2.9	2.6	1.9	1.2	0.6	0.6	*	*	*	*
2003 4/.....	3.0	2.8	1.9	1.3	0.8	0.5	0.3	*	*	*
2002.....	3.0	2.7	1.9	1.3	0.9	0.5	*	*	*	*
2001.....	3.1	2.7	1.7	1.3	0.7	0.4	0.3	*	*	*
2000.....	3.0	2.9	1.9	1.2	0.8	0.5	*	*	*	*
1999.....	3.2	2.6	2.1	1.1	0.7	0.5	*	*	*	*

White, male

2006 4/.....	3.8	3.6	3.3	2.0	1.5	0.7	*	*	*	*
2005 4/.....	4.3	3.8	3.2	2.3	1.3	0.7	*	*	*	*
2004 4/.....	4.8	4.2	3.0	2.2	1.1	1.1	*	*	*	*
2003 4/.....	4.9	4.7	3.3	2.1	1.3	0.8	0.6	*	*	*
2002.....	4.8	4.6	3.2	2.3	1.3	0.9	*	*	*	*
2001.....	5.0	4.6	3.0	2.3	1.2	0.6	0.6	*	*	*
2000.....	4.9	4.8	3.4	1.9	1.3	1.0	*	*	*	*
1999.....	5.2	4.4	3.6	1.9	1.2	0.9	*	*	*	*

White, female

2006 4/.....	1.0	0.8	0.8	0.3	*	*	*	*	*	*
2005 4/.....	1.0	0.7	1.0	0.4	*	*	*	*	*	*
2004 4/.....	1.0	1.0	0.8	*	*	*	*	*	*	*
2003 4/.....	1.1	0.9	0.6	0.6	*	*	*	*	*	*
2002.....	1.2	0.9	0.6	0.4	0.5	*	*	*	*	*
2001.....	1.1	0.9	0.5	0.4	*	*	*	*	*	*
2000.....	1.2	0.9	0.5	0.4	*	*	*	*	*	*
1999.....	1.2	0.9	0.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in infectious and parasitic diseases (B20)

Black, both sexes

2006 4/.....	15.4	18.8	17.1	12.4	8.0	5.7	*	*	*	*
2005 4/.....	16.8	19.7	18.1	13.1	7.7	6.2	3.7	*	*	*
2004 4/.....	20.2	19.8	17.5	12.9	8.1	6.8	3.5	*	*	*
2003 4/.....	17.8	19.0	19.2	12.5	8.5	5.0	4.1	*	*	*
2002.....	21.6	23.0	20.3	10.9	6.6	6.1	2.7	*	*	*
2001.....	21.9	22.2	17.5	10.7	8.5	5.2	3.6	*	*	*
2000.....	22.4	23.5	16.5	9.9	7.4	5.1	4.4	*	*	*
1999.....	24.5	25.1	16.8	10.1	5.8	4.6	4.3	*	*	*

Black, male

2006 4/.....	21.4	25.1	27.2	20.3	12.7	9.1	*	*	*	*
2005 4/.....	23.3	29.5	27.4	22.3	13.1	10.9	*	*	*	*
2004 4/.....	27.6	28.5	26.1	20.8	13.5	11.1	*	*	*	*
2003 4/.....	25.3	29.8	31.2	21.0	14.7	9.1	6.6	*	*	*
2002.....	31.3	34.8	35.1	17.7	11.1	9.5	*	*	*	*
2001.....	31.5	33.3	28.3	18.3	13.8	8.1	6.7	*	*	*
2000.....	31.4	36.9	28.0	16.3	13.3	9.4	9.0	*	*	*
1999.....	37.7	40.3	28.2	18.3	8.9	8.8	7.8	*	*	*

Black, female

2006 4/.....	10.1	13.3	8.6	6.0	4.3	*	*	*	*	*
2005 4/.....	11.1	11.1	10.3	5.5	3.5	*	*	*	*	*
2004 4/.....	13.7	12.3	10.3	6.3	3.9	3.6	*	*	*	*
2003 4/.....	11.2	9.5	9.1	5.4	3.8	*	*	*	*	*
2002.....	13.1	12.7	7.7	5.3	3.1	*	*	*	*	*
2001.....	13.5	12.6	8.3	4.5	4.3	*	*	*	*	*
2000.....	14.4	11.9	6.7	4.6	*	*	*	*	*	*
1999.....	12.8	12.0	7.2	3.3	3.3	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Human immunodeficiency virus (HIV) disease resulting in malignant neoplasms (B21)										
All races 3/, both sexes										
2006.....	0.8	0.7	0.6	0.5	0.4	0.3	*	*	*	*
2005.....	0.9	0.8	0.7	0.5	0.4	0.3	*	*	*	*
2004.....	0.7	0.8	0.8	0.6	0.4	0.2	*	*	*	*
2003.....	0.9	0.8	0.6	0.6	0.3	0.3	*	*	*	*
2002.....	0.9	0.9	0.7	0.4	0.3	*	*	*	*	*
2001.....	1.0	0.8	0.6	0.6	0.4	0.2	*	*	*	*
2000.....	1.0	0.9	0.7	0.5	0.4	0.2	*	*	*	*
1999.....	1.1	0.9	0.7	0.6	0.3	*	*	*	*	*
All races 3/, male										
2006.....	1.2	1.1	1.1	0.9	0.7	0.5	*	*	*	*
2005.....	1.4	1.3	1.3	0.9	0.6	0.5	*	*	*	*
2004.....	1.2	1.4	1.3	1.0	0.7	*	*	*	*	*
2003.....	1.5	1.2	1.0	0.9	0.5	0.4	*	*	*	*
2002.....	1.5	1.5	1.2	0.7	0.6	*	*	*	*	*
2001.....	1.7	1.2	1.1	0.9	0.8	*	*	*	*	*
2000.....	1.7	1.6	1.2	1.0	0.6	*	*	*	*	*
1999.....	1.8	1.5	1.2	1.1	0.5	*	*	*	*	*
All races 3/, female										
2006.....	0.3	0.3	*	*	*	*	*	*	*	*
2005.....	0.3	0.3	0.2	*	*	*	*	*	*	*
2004.....	0.3	0.2	0.2	*	*	*	*	*	*	*
2003.....	0.3	0.4	0.3	*	*	*	*	*	*	*
2002.....	0.3	0.3	0.2	*	*	*	*	*	*	*
2001.....	0.3	0.3	*	*	*	*	*	*	*	*
2000.....	0.3	0.3	*	*	*	*	*	*	*	*
1999.....	0.3	0.2	0.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

All races 3/, both sexes

2006.....	1.8	1.8	1.7	1.4	0.7	0.7	0.5	*	*	*
2005.....	1.9	1.8	1.8	1.2	0.8	0.5	0.4	0.3	*	*
2004.....	2.0	1.8	1.7	1.1	1.0	0.7	0.3	*	*	*
2003.....	3.0	3.0	2.3	1.6	1.1	0.8	0.5	0.3	*	*
2002.....	3.3	3.1	2.4	1.6	1.1	0.8	0.4	0.3	*	*
2001.....	3.3	3.0	2.2	1.4	1.3	0.7	0.4	0.3	*	*
2000.....	3.0	2.9	2.3	1.7	1.2	0.7	0.4	0.3	*	*
1999.....	3.3	2.9	2.2	1.7	1.1	0.9	0.4	*	*	*

All races 3/, male

2006.....	2.6	2.7	2.5	2.2	1.1	1.2	0.7	*	*	*
2005.....	2.7	2.7	2.8	2.0	1.3	0.8	0.7	*	*	*
2004.....	2.8	2.8	2.6	1.7	1.6	1.1	*	*	*	*
2003.....	4.3	4.4	3.6	2.7	1.7	1.2	1.0	*	*	*
2002.....	4.4	4.6	3.8	2.6	2.0	1.5	0.5	*	*	*
2001.....	4.6	4.5	3.6	2.5	2.0	1.1	0.7	*	*	*
2000.....	4.4	4.4	3.6	2.8	1.8	1.1	0.7	0.7	*	*
1999.....	4.7	4.5	3.5	2.7	1.9	1.6	0.7	*	*	*

All races 3/, female

2006.....	1.0	0.9	1.0	0.6	0.4	0.4	*	*	*	*
2005.....	1.0	0.9	0.8	0.5	0.4	*	*	*	*	*
2004.....	1.2	0.9	0.8	0.4	0.4	*	*	*	*	*
2003.....	1.8	1.7	1.0	0.6	0.5	0.4	*	*	*	*
2002.....	2.1	1.7	1.1	0.7	0.4	*	*	*	*	*
2001.....	2.1	1.6	0.9	0.5	0.6	*	*	*	*	*
2000.....	1.7	1.4	1.0	0.6	0.6	*	*	*	*	*
1999.....	1.9	1.3	0.9	0.7	0.4	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease resulting in other conditions (B23)

Black, both sexes

2006 4/.....	8.5	8.7	9.3	8.8	4.6	4.6	3.4	*	*	*
2005 4/.....	8.9	9.3	9.5	7.5	5.5	3.0	*	*	*	*
2004 4/.....	9.4	9.8	9.4	6.6	6.1	3.4	*	*	*	*
2003 4/.....	14.3	15.7	12.5	9.0	6.8	5.5	3.6	*	*	*
2002.....	15.7	16.8	13.7	9.6	7.0	5.8	3.2	*	*	*
2001.....	15.7	15.5	12.9	7.8	8.0	4.1	*	*	*	*
2000.....	14.0	14.7	13.4	9.0	7.2	4.6	2.9	*	*	*
1999.....	16.2	15.5	13.1	10.4	5.7	5.6	3.2	*	*	*

Black, male

2006 4/.....	11.5	12.3	12.6	14.0	6.7	7.5	*	*	*	*
2005 4/.....	13.2	13.4	15.7	12.5	9.3	5.5	*	*	*	*
2004 4/.....	12.6	15.5	14.6	10.6	9.9	6.1	*	*	*	*
2003 4/.....	19.4	22.2	19.8	15.4	11.2	8.6	6.6	*	*	*
2002.....	20.1	25.2	21.6	15.5	12.5	11.2	*	*	*	*
2001.....	20.6	23.5	20.9	14.2	12.4	7.4	*	*	*	*
2000.....	20.1	22.9	21.4	15.1	11.6	8.1	*	*	*	*
1999.....	22.8	25.3	21.1	18.0	9.1	10.4	6.8	*	*	*

Black, female

2006 4/.....	5.9	5.4	6.5	4.4	2.9	*	*	*	*	*
2005 4/.....	5.1	5.7	4.3	3.4	*	*	*	*	*	*
2004 4/.....	6.6	4.8	5.0	3.3	3.1	*	*	*	*	*
2003 4/.....	9.7	10.1	6.4	3.7	3.3	*	*	*	*	*
2002.....	11.9	9.4	7.0	4.8	*	*	*	*	*	*
2001.....	11.4	8.6	6.1	2.6	4.5	*	*	*	*	*
2000.....	8.5	7.7	6.5	4.0	3.8	*	*	*	*	*
1999.....	10.4	7.2	6.3	4.3	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified human immunodeficiency virus (HIV) disease (B24)										
All races 3/, both sexes										
2006.....	3.1	3.1	2.3	1.6	1.0	0.7	0.6	0.3	*	*
2005.....	3.4	3.2	2.5	1.4	1.1	0.7	0.4	0.3	*	*
2004.....	3.4	3.1	2.5	1.6	1.1	0.7	0.3	*	*	*
2003.....	2.8	2.4	1.9	1.3	0.8	0.6	0.3	0.3	*	*
2002.....	2.6	2.4	1.8	1.0	0.7	0.5	0.3	*	*	*
2001.....	2.9	2.3	1.7	0.9	0.7	0.6	0.2	*	*	*
2000.....	2.6	2.4	1.6	1.0	0.7	0.5	0.3	*	*	*
1999.....	2.9	2.3	1.3	1.0	0.6	0.5	0.3	*	*	*
All races 3/, male										
2006.....	4.4	4.6	3.6	2.5	1.8	1.2	1.1	*	*	*
2005.....	4.9	4.8	4.0	2.4	1.8	1.2	0.8	*	*	*
2004.....	4.8	4.5	3.8	2.7	1.9	1.1	0.6	*	*	*
2003.....	4.1	3.8	3.0	2.2	1.4	0.9	0.6	*	*	*
2002.....	3.9	3.7	2.9	1.7	1.2	0.8	0.6	*	*	*
2001.....	4.3	3.8	2.8	1.6	1.2	0.9	0.5	*	*	*
2000.....	4.0	3.9	2.6	1.7	1.4	0.9	0.5	*	*	*
1999.....	4.5	3.7	2.2	1.8	1.0	0.8	0.5	*	*	*
All races 3/, female										
2006.....	1.9	1.6	1.0	0.7	0.4	*	*	*	*	*
2005.....	1.9	1.7	1.1	0.5	0.4	*	*	*	*	*
2004.....	1.9	1.7	1.2	0.7	0.3	*	*	*	*	*
2003.....	1.6	1.1	0.8	0.5	*	*	*	*	*	*
2002.....	1.4	1.2	0.7	0.4	*	*	*	*	*	*
2001.....	1.5	0.8	0.6	0.3	*	*	*	*	*	*
2000.....	1.3	1.0	0.7	*	*	*	*	*	*	*
1999.....	1.4	0.9	0.4	0.3	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified human immunodeficiency virus (HIV) disease (B24)										
Black, both sexes										
2006 4/.....	14.0	14.7	11.3	8.3	5.7	3.7	3.9	*	*	*
2005 4/.....	14.4	14.4	12.8	7.5	6.5	4.4	*	*	*	*
2004 4/.....	14.7	15.5	13.0	8.3	6.5	4.5	*	*	*	*
2003 4/.....	12.7	11.4	9.3	6.7	4.1	3.9	*	*	*	*
2002.....	11.8	10.7	9.2	5.3	3.5	3.0	*	*	*	*
2001.....	12.8	11.7	8.6	4.8	4.7	3.7	*	*	*	*
2000.....	11.9	11.7	9.4	5.0	3.8	3.1	*	*	*	*
1999.....	12.2	11.3	6.2	5.9	3.5	2.2	*	*	*	*
Black, male										
2006 4/.....	18.1	20.5	18.3	13.1	10.6	5.9	7.4	*	*	*
2005 4/.....	19.3	20.7	20.3	13.3	11.5	7.8	*	*	*	*
2004 4/.....	19.5	22.4	19.5	14.3	11.8	7.5	*	*	*	*
2003 4/.....	17.3	17.2	14.9	11.2	7.5	7.4	*	*	*	*
2002.....	16.3	15.8	15.8	9.1	6.5	5.2	*	*	*	*
2001.....	17.8	19.7	13.7	8.1	8.7	5.8	*	*	*	*
2000.....	17.8	18.5	15.2	9.4	7.9	5.2	*	*	*	*
1999.....	18.2	18.5	10.7	10.5	5.7	*	*	*	*	*
Black, female										
2006 4/.....	10.4	9.6	5.3	4.4	*	*	*	*	*	*
2005 4/.....	10.1	8.8	6.4	2.7	*	*	*	*	*	*
2004 4/.....	10.4	9.5	7.5	3.5	*	*	*	*	*	*
2003 4/.....	8.6	6.4	4.6	3.0	*	*	*	*	*	*
2002.....	7.8	6.3	3.6	*	*	*	*	*	*	*
2001.....	8.4	4.8	4.1	*	*	*	*	*	*	*
2000.....	6.6	5.7	4.5	*	*	*	*	*	*	*
1999.....	6.9	5.1	2.4	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

All races 3/, both sexes

2006.....	0.1	0.1	0.3	0.4	0.5	0.6	1.0	1.0	1.4	1.8
2005.....	0.1	0.2	0.2	0.3	0.7	0.9	0.9	1.2	1.7	2.7
2004.....	0.1	0.2	0.2	0.4	0.6	0.8	0.8	1.1	1.4	2.0
2003.....	0.2	0.2	0.2	0.4	0.6	0.8	1.2	1.2	1.4	2.8
2002.....	0.2	0.2	0.3	0.3	0.5	0.8	1.0	1.2	1.5	2.9
2001.....	0.1	0.1	0.2	0.3	0.6	0.9	0.9	0.9	1.2	2.2
2000.....	0.1	0.2	0.3	0.4	0.6	0.9	1.1	1.2	1.7	3.3
1999.....	0.2	0.2	0.4	0.6	0.6	0.8	1.1	1.5	2.1	4.0

All races 3/, male

2006.....	*	*	0.3	0.4	0.5	0.6	1.1	1.3	1.9	2.1
2005.....	*	0.2	*	0.3	0.8	1.1	0.9	1.6	1.8	2.7
2004.....	*	*	0.3	0.5	0.7	0.8	0.9	1.5	1.6	2.3
2003.....	0.2	0.2	0.3	0.4	0.8	0.9	1.4	1.3	1.7	3.0
2002.....	0.2	0.2	0.3	0.3	0.5	0.9	1.0	1.4	1.8	2.8
2001.....	0.2	*	0.2	0.3	0.7	0.9	1.0	1.0	1.5	2.4
2000.....	*	*	0.3	0.4	0.7	1.0	1.4	1.7	2.0	3.1
1999.....	*	0.2	0.4	0.5	0.7	0.8	1.4	1.4	2.0	3.9

All races 3/, female

2006.....	*	*	0.3	0.4	0.4	0.6	1.0	0.8	1.2	1.7
2005.....	*	*	0.2	0.3	0.6	0.8	0.8	1.0	1.6	2.7
2004.....	*	0.2	*	0.4	0.5	0.7	0.7	0.8	1.3	1.8
2003.....	0.2	*	*	0.4	0.4	0.7	0.9	1.2	1.3	2.7
2002.....	0.2	0.2	0.2	0.3	0.5	0.8	1.0	1.1	1.3	2.9
2001.....	*	*	0.2	0.3	0.5	0.9	0.8	0.8	1.0	2.1
2000.....	*	0.2	0.3	0.4	0.6	0.8	1.0	0.8	1.6	3.4
1999.....	*	*	0.4	0.6	0.6	0.8	0.9	1.5	2.1	4.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified viral diseases (A81,A85.0-A85.1,A85.8,A86-A89,B01,B04,B07-B09,B25-B34)

White, both sexes

2006 4/.....	*	0.1	0.3	0.4	0.4	0.7	1.1	1.1	1.5	1.9
2005 4/.....	0.1	0.2	0.2	0.4	0.7	1.1	0.9	1.3	1.6	3.0
2004 4/.....	*	0.2	0.2	0.4	0.6	0.8	0.9	1.2	1.5	2.0
2003 4/.....	0.1	0.2	0.2	0.4	0.7	0.8	1.2	1.3	1.4	3.0
2002.....	0.2	0.2	0.3	0.3	0.5	0.9	1.1	1.2	1.6	3.1
2001.....	0.1	*	0.2	0.3	0.7	1.0	0.9	0.9	1.2	2.3
2000.....	0.1	0.2	0.3	0.5	0.6	1.0	1.2	1.2	1.8	3.4
1999.....	0.2	0.2	0.4	0.6	0.7	0.8	1.1	1.5	2.1	4.2

White, male

2006 4/.....	*	*	0.2	0.5	0.4	0.6	1.2	1.4	2.0	2.2
2005 4/.....	*	*	*	0.4	0.8	1.2	1.0	1.6	1.7	3.0
2004 4/.....	*	*	0.2	0.5	0.8	0.9	1.0	1.6	1.8	2.4
2003 4/.....	*	*	0.3	0.5	0.8	0.9	1.5	1.4	1.7	3.2
2002.....	0.2	*	0.3	*	0.5	1.0	1.1	1.5	1.9	2.9
2001.....	*	*	*	0.3	0.7	0.9	1.0	1.1	1.6	2.6
2000.....	*	*	0.3	0.4	0.7	1.0	1.5	1.7	2.0	3.3
1999.....	*	*	0.4	0.5	0.7	0.8	1.4	1.5	1.9	3.9

White, female

2006 4/.....	*	*	0.3	0.4	0.4	0.7	1.0	0.9	1.1	1.8
2005 4/.....	*	*	*	0.4	0.6	0.9	0.9	1.0	1.6	3.0
2004 4/.....	*	*	*	0.4	0.4	0.7	0.7	0.9	1.3	1.8
2003 4/.....	*	*	*	0.4	0.5	0.8	1.0	1.3	1.3	2.8
2002.....	*	*	*	0.4	0.5	0.8	1.0	1.0	1.4	3.1
2001.....	*	*	*	0.4	0.6	1.0	0.9	0.8	1.0	2.2
2000.....	*	*	*	0.5	0.5	0.9	1.0	0.8	1.7	3.5
1999.....	*	*	0.4	0.6	0.6	0.9	0.9	1.6	2.2	4.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

All races 3/, both sexes

2006.....	0.5	0.6	0.9	1.2	1.8	2.6	3.9	4.8	6.4	8.6
2005.....	0.9	1.9	3.4	2.9	2.8	3.6	4.7	6.8	7.9	9.0
2004.....	0.9	2.0	2.9	2.9	2.7	3.9	5.0	6.2	8.2	9.7
2003.....	0.9	1.9	2.8	2.6	2.8	3.5	5.0	7.0	7.1	9.0
2002.....	0.9	1.4	2.1	2.0	2.5	3.2	4.3	6.2	6.7	8.3
2001.....	0.6	1.1	1.2	1.6	2.3	3.3	4.0	5.1	5.9	8.2
2000.....	0.6	0.9	1.0	1.4	2.1	2.7	4.0	5.4	6.1	9.0
1999.....	1.0	1.4	1.7	2.0	2.8	4.1	5.3	7.0	8.2	10.0

All races 3/, male

2006.....	0.6	0.8	1.2	1.5	2.0	3.0	4.8	5.7	8.0	10.3
2005.....	1.2	2.6	5.0	4.0	3.4	4.2	5.5	7.2	10.4	9.7
2004.....	1.2	2.8	4.2	4.1	3.4	4.4	5.8	7.3	10.0	12.9
2003.....	1.2	2.7	3.9	3.5	3.4	4.1	5.4	7.6	9.2	10.4
2002.....	1.2	2.1	3.0	2.7	3.2	4.1	5.3	7.1	9.0	11.9
2001.....	0.9	1.6	1.6	2.1	2.6	3.6	4.1	6.5	8.0	10.1
2000.....	0.8	1.2	1.3	1.6	2.5	3.3	5.4	6.3	7.7	12.6
1999.....	1.3	1.7	2.1	2.2	3.2	4.8	6.2	8.9	10.3	13.1

All races 3/, female

2006.....	0.4	0.4	0.6	0.9	1.6	2.3	3.2	4.1	5.4	7.7
2005.....	0.5	1.1	1.8	1.9	2.3	3.1	4.1	6.5	6.3	8.7
2004.....	0.7	1.1	1.7	1.9	2.1	3.3	4.4	5.5	7.2	8.3
2003.....	0.6	1.2	1.7	1.7	2.2	3.1	4.8	6.6	5.8	8.3
2002.....	0.5	0.7	1.1	1.4	1.8	2.4	3.5	5.6	5.3	6.8
2001.....	0.4	0.6	0.9	1.1	2.0	3.1	3.9	4.2	4.7	7.4
2000.....	0.4	0.7	0.8	1.2	1.7	2.2	2.9	4.7	5.2	7.6
1999.....	0.8	1.1	1.4	1.8	2.4	3.5	4.6	5.6	6.9	8.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

White, both sexes

2006 4/.....	0.4	0.5	0.8	1.1	1.7	2.6	3.8	4.7	6.3	8.5
2005 4/.....	0.8	1.8	3.4	2.7	2.6	3.5	4.4	6.4	7.6	8.8
2004 4/.....	0.9	2.0	3.0	2.7	2.5	3.6	4.8	6.0	7.9	9.6
2003 4/.....	0.9	1.9	2.7	2.4	2.5	3.5	4.9	6.6	6.9	8.9
2002.....	0.8	1.3	1.9	1.7	2.3	3.1	4.2	6.2	6.6	8.3
2001.....	0.5	1.0	1.1	1.4	2.0	3.1	3.8	5.1	5.8	7.9
2000.....	0.5	0.9	0.9	1.3	1.8	2.5	3.8	5.3	6.0	8.9
1999.....	0.7	1.1	1.4	1.8	2.3	3.7	5.0	6.8	7.8	9.6

White, male

2006 4/.....	0.4	0.7	1.1	1.3	1.8	3.2	4.6	5.5	7.9	10.3
2005 4/.....	1.1	2.5	5.1	3.7	3.2	4.1	5.4	6.9	10.1	9.6
2004 4/.....	1.1	2.9	4.2	3.7	3.2	4.3	5.7	7.0	9.8	13.2
2003 4/.....	1.2	2.7	3.8	3.3	3.1	4.0	5.1	7.2	8.9	10.5
2002.....	1.1	2.0	2.8	2.3	3.0	4.0	5.2	6.9	8.9	11.8
2001.....	0.7	1.5	1.4	1.8	2.4	3.4	4.0	6.3	8.0	9.9
2000.....	0.7	1.1	1.2	1.6	2.1	3.1	5.2	5.9	7.5	12.0
1999.....	1.0	1.4	1.7	2.1	2.7	4.4	5.9	8.7	9.4	12.0

White, female

2006 4/.....	0.3	0.4	0.5	0.9	1.6	2.1	3.1	4.1	5.4	7.6
2005 4/.....	0.5	1.1	1.7	1.8	2.1	3.0	3.6	6.1	6.0	8.5
2004 4/.....	0.7	1.1	1.7	1.8	2.0	3.0	4.1	5.3	6.7	7.9
2003 4/.....	0.6	1.1	1.6	1.5	2.0	3.0	4.8	6.1	5.6	8.2
2002.....	0.4	0.7	1.0	1.2	1.7	2.3	3.4	5.6	5.2	6.7
2001.....	0.4	0.6	0.9	1.0	1.8	2.9	3.7	4.2	4.4	7.1
2000.....	0.3	0.6	0.7	1.1	1.5	2.1	2.8	4.9	5.0	7.6
1999.....	0.4	0.8	1.1	1.4	2.1	3.2	4.3	5.4	6.9	8.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (B35-B99)

Black, both sexes

2006 4/.....	1.1	1.0	2.0	2.0	2.5	3.6	5.1	5.9	7.6	11.2
2005 4/.....	1.2	2.2	3.7	4.6	4.6	4.0	5.7	8.2	9.2	11.0
2004 4/.....	1.5	2.0	2.8	4.2	4.4	5.3	5.6	8.0	11.1	13.3
2003 4/.....	1.3	2.7	3.8	4.1	4.3	3.5	4.8	10.0	8.9	9.0
2002.....	1.5	2.2	3.8	4.8	3.3	3.5	4.0	7.0	5.5	9.5
2001.....	1.3	2.0	2.0	2.9	3.9	4.4	5.9	4.9	7.0	11.4
2000.....	1.3	1.9	2.2	2.5	3.7	3.9	5.3	5.9	9.0	10.9
1999.....	3.0	3.9	4.4	4.1	6.2	6.9	8.4	9.5	11.0	14.3

Black, male

2006 4/.....	1.5	*	2.9	3.0	*	*	6.5	*	*	*
2005 4/.....	1.5	3.0	4.8	6.5	5.8	5.7	*	10.7	*	*
2004 4/.....	2.0	3.1	4.5	6.7	6.4	*	7.1	11.4	*	*
2003 4/.....	1.5	3.3	5.0	5.3	5.8	*	6.6	9.6	*	*
2002.....	2.1	3.2	5.4	7.3	4.9	*	*	*	*	*
2001.....	2.1	3.0	2.9	3.6	4.7	5.6	*	*	*	*
2000.....	1.7	2.3	3.0	*	5.0	5.5	7.4	9.9	*	*
1999.....	3.0	4.2	5.1	*	7.6	7.8	9.8	10.9	18.3	*

Black, female

2006 4/.....	*	*	*	*	*	4.2	4.2	*	*	10.6
2005 4/.....	*	1.6	2.6	3.0	3.7	*	5.8	6.7	7.8	10.5
2004 4/.....	*	*	*	*	2.9	5.9	4.6	5.9	10.6	13.0
2003 4/.....	*	2.1	2.7	3.1	3.2	*	*	10.3	*	8.9
2002.....	*	*	2.5	2.7	*	*	*	6.6	*	*
2001.....	*	*	*	*	3.2	*	5.6	*	*	11.8
2000.....	*	*	*	*	*	*	*	*	*	9.0
1999.....	3.1	3.7	3.9	5.0	5.2	6.2	7.4	8.6	*	11.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mycoses (B35-B49)										
All races 3/, both sexes										
2006.....	0.2	0.2	0.4	0.4	0.7	1.1	1.6	1.5	2.1	2.3
2005.....	0.2	0.3	0.4	0.5	0.7	1.0	1.2	1.9	2.1	2.1
2004.....	0.2	0.3	0.3	0.6	0.8	1.0	1.3	1.7	2.4	2.5
2003.....	0.2	0.2	0.4	0.5	0.7	1.1	1.5	2.1	2.1	2.4
2002.....	0.2	0.3	0.5	0.6	0.8	1.1	1.7	2.3	2.7	2.5
2001.....	0.2	0.3	0.3	0.5	0.8	1.2	1.4	2.3	2.0	3.0
2000.....	0.2	0.3	0.5	0.7	1.0	1.2	1.7	2.3	2.1	3.2
1999.....	0.4	0.6	0.8	1.0	1.4	2.1	2.6	3.2	3.3	3.9
All races 3/, male										
2006.....	0.3	0.3	0.4	0.6	0.8	1.3	2.1	2.0	2.7	3.4
2005.....	0.3	0.3	0.4	0.4	0.8	1.3	1.7	2.3	3.3	2.2
2004.....	0.2	0.4	0.4	0.8	0.8	1.1	1.5	2.2	3.4	3.8
2003.....	0.3	0.3	0.4	0.6	0.9	1.1	1.7	2.7	3.2	3.3
2002.....	0.3	0.4	0.7	0.9	1.1	1.7	2.3	3.1	4.1	4.2
2001.....	0.3	0.5	0.4	0.7	1.0	1.4	1.5	3.2	2.9	3.8
2000.....	0.3	0.3	0.6	0.7	1.2	1.6	2.4	3.0	2.8	4.9
1999.....	0.5	0.6	0.8	1.2	1.7	2.4	3.4	4.4	4.9	6.6
All races 3/, female										
2006.....	0.2	*	0.3	0.4	0.5	0.8	1.3	1.2	1.7	1.9
2005.....	*	0.2	0.3	0.5	0.6	0.7	0.8	1.6	1.3	2.0
2004.....	*	*	0.3	0.4	0.7	0.8	1.2	1.3	1.8	1.9
2003.....	*	0.2	0.3	0.4	0.6	1.1	1.3	1.7	1.5	2.0
2002.....	*	*	0.3	0.4	0.6	0.7	1.2	1.6	1.9	1.7
2001.....	*	0.2	0.3	0.4	0.7	1.0	1.3	1.7	1.5	2.6
2000.....	0.2	0.3	0.4	0.6	0.8	1.0	1.2	1.8	1.7	2.6
1999.....	0.3	0.5	0.7	0.9	1.2	1.9	1.9	2.4	2.3	2.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mycoses (B35-B49)										
White, both sexes										
2006 4/.....	0.2	0.2	0.3	0.4	0.6	1.0	1.6	1.4	2.0	2.2
2005 4/.....	0.1	0.2	0.3	0.4	0.6	0.9	1.1	1.8	2.0	1.9
2004 4/.....	0.1	0.2	0.2	0.5	0.7	0.9	1.2	1.5	2.2	2.4
2003 4/.....	0.1	0.2	0.3	0.4	0.6	1.1	1.5	2.0	2.0	2.3
2002.....	0.2	0.2	0.3	0.5	0.7	1.0	1.6	2.1	2.7	2.4
2001.....	0.2	0.3	0.3	0.5	0.7	1.0	1.2	2.2	1.8	2.6
2000.....	0.2	0.3	0.4	0.6	0.9	1.2	1.6	2.2	2.1	3.0
1999.....	0.3	0.5	0.7	0.9	1.2	1.9	2.4	3.1	3.0	3.5
White, male										
2006 4/.....	*	0.3	0.3	0.4	0.7	1.4	1.9	1.8	2.6	3.4
2005 4/.....	*	0.2	0.3	0.3	0.8	1.2	1.6	2.3	3.3	2.1
2004 4/.....	*	0.3	0.3	0.6	0.7	1.0	1.4	1.9	3.2	4.0
2003 4/.....	0.2	0.2	0.4	0.4	0.8	1.1	1.5	2.6	3.1	3.4
2002.....	0.3	0.4	0.4	0.7	1.0	1.5	2.2	2.9	4.1	4.1
2001.....	*	0.4	0.4	0.6	0.8	1.2	1.3	3.0	2.8	3.4
2000.....	*	0.3	0.5	0.7	1.1	1.5	2.2	2.7	2.6	4.4
1999.....	0.3	0.5	0.7	1.2	1.4	2.2	3.3	4.4	4.4	5.7
White, female										
2006 4/.....	*	*	*	0.3	0.4	0.6	1.2	1.1	1.6	1.6
2005 4/.....	*	*	*	0.4	0.5	0.7	0.6	1.4	1.2	1.8
2004 4/.....	*	*	*	0.3	0.6	0.7	1.1	1.2	1.5	1.7
2003 4/.....	*	*	0.3	0.4	0.5	1.0	1.4	1.5	1.4	1.7
2002.....	*	*	*	0.4	0.5	0.6	1.1	1.5	1.9	1.7
2001.....	*	*	0.3	0.4	0.6	0.9	1.2	1.6	1.2	2.2
2000.....	*	0.3	0.3	0.5	0.7	0.9	1.1	1.8	1.8	2.4
1999.....	*	0.4	0.6	0.6	1.0	1.7	1.7	2.2	2.1	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Sequelae of tuberculosis (B90)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	0.5
2002.....	*	*	*	*	*	*	*	*	*	0.9
2001.....	*	*	*	*	*	*	*	*	0.5	0.6
2000.....	*	*	*	*	*	*	*	*	0.5	0.7
1999.....	*	*	*	*	*	*	*	0.4	0.9	0.8
White, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, female										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	0.9
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	1.0	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Sequelae of poliomyelitis (B91)										
All races 3/, both sexes										
2006.....	*	*	*	*	0.3	0.4	0.5	0.9	1.1	1.3
2005.....	*	*	*	0.1	0.3	0.3	0.5	0.9	1.4	1.2
2004.....	*	*	*	*	0.2	0.4	0.6	0.7	1.0	1.4
2003.....	*	*	*	0.1	0.2	0.3	0.5	0.7	0.8	1.2
2002.....	*	*	*	0.1	0.2	0.4	0.6	0.9	0.9	1.3
2001.....	*	*	*	*	0.3	0.4	0.6	0.8	0.9	1.2
2000.....	*	*	*	0.2	0.2	0.3	0.5	0.8	0.8	1.2
1999.....	*	*	*	*	0.2	0.3	0.5	0.6	0.7	1.3
All races 3/, male										
2006.....	*	*	*	*	*	0.5	0.6	1.1	1.2	1.3
2005.....	*	*	*	*	*	*	0.7	0.9	2.0	*
2004.....	*	*	*	*	*	*	0.6	0.8	1.4	1.9
2003.....	*	*	*	*	*	*	0.6	*	1.3	1.5
2002.....	*	*	*	*	*	0.5	0.5	0.7	1.1	1.9
2001.....	*	*	*	*	*	*	0.7	1.2	*	1.8
2000.....	*	*	*	*	*	*	0.6	0.8	*	*
1999.....	*	*	*	*	*	*	*	0.8	*	*
All races 3/, female										
2006.....	*	*	*	*	0.3	0.4	*	0.8	1.1	1.2
2005.....	*	*	*	*	*	*	*	1.0	1.1	1.2
2004.....	*	*	*	*	*	*	0.5	0.7	0.8	1.1
2003.....	*	*	*	*	*	*	0.5	0.8	*	1.1
2002.....	*	*	*	*	*	0.4	0.6	1.0	0.8	1.0
2001.....	*	*	*	*	*	0.4	0.6	0.6	0.9	1.0
2000.....	*	*	*	*	*	*	0.5	0.8	0.8	1.1
1999.....	*	*	*	*	*	*	0.5	0.5	0.7	1.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Sequelae of poliomyelitis (B91)										
White, both sexes										
2006 4/.....	*	*	*	*	0.3	0.5	0.5	1.1	1.2	1.4
2005 4/.....	*	*	*	*	0.3	0.3	0.6	1.0	1.5	1.3
2004 4/.....	*	*	*	*	0.2	0.4	0.6	0.8	1.1	1.5
2003 4/.....	*	*	*	0.2	0.3	0.3	0.6	0.8	0.9	1.3
2002.....	*	*	*	0.2	0.2	0.5	0.6	1.0	1.0	1.4
2001.....	*	*	*	*	0.4	0.5	0.7	0.9	1.0	1.4
2000.....	*	*	*	0.2	0.2	0.4	0.6	0.9	0.9	1.4
1999.....	*	*	*	*	0.2	0.3	0.5	0.7	0.8	1.4
White, male										
2006 4/.....	*	*	*	*	*	0.5	0.6	1.2	1.3	1.4
2005 4/.....	*	*	*	*	*	*	0.8	1.0	2.2	*
2004 4/.....	*	*	*	*	*	*	0.7	0.9	1.6	2.1
2003 4/.....	*	*	*	*	*	*	0.6	*	1.5	1.5
2002.....	*	*	*	*	*	0.5	0.6	0.8	1.3	2.1
2001.....	*	*	*	*	*	*	0.8	1.3	*	2.0
2000.....	*	*	*	*	*	*	0.7	0.9	*	*
1999.....	*	*	*	*	*	*	*	0.8	*	*
White, female										
2006 4/.....	*	*	*	*	0.4	0.4	*	0.9	1.2	1.4
2005 4/.....	*	*	*	*	*	*	*	1.1	1.2	1.4
2004 4/.....	*	*	*	*	*	*	0.6	0.7	0.8	1.3
2003 4/.....	*	*	*	*	*	*	0.6	0.9	*	1.2
2002.....	*	*	*	*	*	*	0.6	1.1	0.8	1.1
2001.....	*	*	*	*	*	0.5	0.6	0.7	0.9	1.1
2000.....	*	*	*	*	*	*	0.5	0.9	0.9	1.2
1999.....	*	*	*	*	*	*	0.6	0.5	0.8	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

All races 3/, both sexes

2006.....	0.1	0.2	0.4	0.5	0.7	1.0	1.5	2.1	2.9	4.5
2005.....	0.6	1.5	2.9	2.2	1.8	2.2	2.8	3.6	3.8	5.3
2004.....	0.7	1.6	2.4	2.1	1.6	2.2	2.8	3.4	4.3	5.3
2003.....	0.7	1.5	2.3	1.8	1.6	1.9	2.7	3.9	3.7	4.8
2002.....	0.5	1.0	1.4	1.1	1.3	1.4	1.7	2.6	2.6	3.6
2001.....	0.3	0.6	0.7	0.7	0.8	1.3	1.4	1.5	2.3	3.2
2000.....	0.2	0.4	0.4	0.4	0.6	0.8	1.2	1.7	2.3	3.7
1999.....	0.5	0.7	0.7	0.7	1.0	1.4	1.6	2.4	3.0	3.9

All races 3/, male

2006.....	*	0.3	0.6	0.7	0.7	1.1	1.8	2.1	3.8	4.9
2005.....	0.8	2.1	4.4	3.3	2.2	2.4	2.7	3.6	4.3	5.7
2004.....	0.8	2.3	3.6	3.0	2.1	2.4	3.3	3.8	4.4	6.3
2003.....	0.9	2.2	3.4	2.6	1.9	2.4	2.8	4.1	4.3	4.8
2002.....	0.8	1.4	2.1	1.5	1.6	1.7	2.0	2.8	3.2	4.7
2001.....	0.5	0.9	1.0	1.0	0.9	1.3	1.3	1.5	3.2	3.4
2000.....	0.4	0.6	0.5	0.5	0.7	1.0	1.5	2.0	2.9	4.3
1999.....	0.6	0.9	0.9	0.7	1.0	1.7	1.6	2.9	3.7	3.8

All races 3/, female

2006.....	*	0.2	0.2	0.4	0.7	0.9	1.3	2.0	2.4	4.3
2005.....	0.4	0.9	1.5	1.2	1.3	2.0	2.8	3.6	3.6	5.1
2004.....	0.5	0.9	1.3	1.3	1.2	2.1	2.4	3.2	4.2	4.8
2003.....	0.5	0.9	1.2	1.0	1.4	1.5	2.7	3.8	3.3	4.8
2002.....	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.5	2.2	3.1
2001.....	0.2	0.3	0.4	0.4	0.7	1.2	1.5	1.5	1.8	3.2
2000.....	*	0.3	0.3	0.3	0.5	0.6	0.9	1.5	2.0	3.4
1999.....	0.3	0.5	0.5	0.7	0.9	1.1	1.6	2.1	2.6	3.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

White, both sexes

2006 4/.....	0.1	0.2	0.4	0.5	0.6	1.0	1.5	2.0	2.9	4.4
2005 4/.....	0.6	1.5	3.0	2.1	1.6	2.1	2.6	3.3	3.5	5.1
2004 4/.....	0.7	1.7	2.6	2.1	1.6	2.1	2.7	3.4	4.1	5.1
2003 4/.....	0.7	1.6	2.2	1.7	1.5	1.9	2.6	3.5	3.5	4.8
2002.....	0.5	1.0	1.4	0.9	1.2	1.4	1.8	2.6	2.6	3.5
2001.....	0.3	0.6	0.6	0.6	0.7	1.2	1.3	1.5	2.3	3.3
2000.....	0.2	0.5	0.4	0.3	0.5	0.7	1.2	1.8	2.2	3.6
1999.....	0.3	0.5	0.5	0.6	0.7	1.2	1.5	2.3	2.9	3.8

White, male

2006 4/.....	*	0.3	0.6	0.6	0.6	1.1	1.7	2.1	3.7	4.7
2005 4/.....	0.9	2.2	4.6	3.1	2.0	2.3	2.7	3.3	4.0	5.5
2004 4/.....	0.8	2.4	3.8	2.9	2.1	2.3	3.3	3.7	4.4	6.2
2003 4/.....	1.0	2.3	3.3	2.5	1.8	2.3	2.6	3.8	3.9	4.8
2002.....	0.7	1.5	2.2	1.3	1.5	1.7	2.0	2.8	3.2	4.7
2001.....	0.4	0.9	0.9	0.9	0.8	1.4	1.3	1.5	3.1	3.5
2000.....	0.4	0.6	0.5	0.4	0.5	0.8	1.5	2.0	2.9	4.1
1999.....	0.5	0.7	0.8	0.6	0.8	1.4	1.5	2.8	3.3	3.7

White, female

2006 4/.....	*	*	0.2	0.4	0.6	0.9	1.3	2.0	2.4	4.3
2005 4/.....	0.4	0.9	1.4	1.2	1.2	1.9	2.5	3.2	3.2	4.9
2004 4/.....	0.6	0.9	1.4	1.3	1.1	1.8	2.3	3.1	4.0	4.6
2003 4/.....	0.5	0.9	1.2	0.9	1.2	1.6	2.6	3.4	3.2	4.8
2002.....	0.3	0.5	0.6	0.5	0.9	1.1	1.5	2.5	2.2	2.9
2001.....	*	0.3	0.4	*	0.7	1.1	1.4	1.5	1.7	3.2
2000.....	*	0.3	0.3	*	0.5	0.7	1.0	1.6	1.7	3.4
1999.....	*	*	0.3	0.5	0.7	1.0	1.5	2.0	2.7	3.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified infectious and parasitic diseases and their sequelae (B85-B89,B92-B99)

Black, both sexes

2006 4/.....	*	*	*	*	*	*	*	*	*	6.0
2005 4/.....	*	1.3	2.4	3.3	2.9	2.2	3.1	4.5	5.6	6.9
2004 4/.....	*	1.1	1.7	2.4	2.2	3.4	2.9	3.9	5.0	8.2
2003 4/.....	*	1.6	2.8	2.6	2.3	*	*	5.6	5.3	*
2002.....	*	1.1	1.7	2.3	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	1.3	2.1	2.1	2.0	2.9	2.4	3.2	3.9	*	*

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	1.7	3.0	5.0	4.2	*	*	*	*	*
2004 4/.....	*	1.8	3.2	3.8	*	*	*	*	*	*
2003 4/.....	*	2.2	3.9	3.5	*	*	*	*	*	*
2002.....	*	*	2.4	3.4	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	2.0	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	1.8	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	4.1	*	*	*	7.9
2003 4/.....	*	*	1.8	*	*	*	*	5.7	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	1.4	2.1	2.2	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
II. Neoplasms (C00-D48)										
All races 3/, both sexes										
2006.....	43.2	85.6	154.4	255.9	420.1	624.1	880.0	1,173.6	1,465.2	1,684.1
2005.....	45.6	87.9	156.9	258.9	427.8	631.7	902.6	1,186.3	1,472.5	1,714.4
2004.....	45.1	88.1	157.6	265.0	433.4	647.1	909.1	1,199.4	1,472.0	1,731.0
2003.....	47.0	91.2	161.1	271.1	447.0	657.7	925.5	1,223.9	1,497.3	1,778.8
2002.....	48.5	92.1	163.6	280.5	452.5	673.5	949.9	1,235.8	1,504.6	1,802.6
2001.....	49.7	93.9	167.1	284.6	459.6	683.7	961.4	1,241.2	1,513.7	1,848.3
2000.....	49.8	95.0	168.5	292.8	470.0	701.2	969.0	1,251.3	1,555.9	1,904.3
1999.....	50.3	95.1	169.3	298.8	477.9	711.0	982.0	1,253.3	1,540.6	1,888.7
All races 3/, male										
2006.....	37.9	82.3	164.6	286.0	482.6	742.7	1,067.2	1,480.5	1,962.8	2,341.0
2005.....	40.5	84.9	167.3	288.4	491.7	755.7	1,112.1	1,511.4	1,961.1	2,412.0
2004.....	40.2	87.5	170.5	293.9	499.6	768.6	1,112.1	1,538.1	1,955.7	2,448.4
2003.....	42.7	92.0	172.2	299.6	514.6	786.0	1,139.0	1,569.2	2,001.3	2,515.6
2002.....	43.9	91.4	173.9	313.2	526.0	809.1	1,177.8	1,591.4	2,010.6	2,592.2
2001.....	45.0	93.4	176.6	315.6	534.1	826.9	1,200.6	1,607.6	2,041.1	2,721.3
2000.....	45.3	95.1	176.6	325.4	543.2	848.0	1,211.6	1,622.8	2,115.8	2,819.6
1999.....	46.6	96.1	177.6	334.6	558.0	861.6	1,236.8	1,637.2	2,107.0	2,831.4
All races 3/, female										
2006.....	48.4	88.9	144.5	227.4	363.1	520.5	727.8	949.0	1,154.1	1,376.8
2005.....	50.5	90.8	146.9	231.1	369.5	523.5	732.8	950.3	1,169.0	1,394.0
2004.....	49.9	88.8	145.2	237.8	373.1	541.3	745.6	956.0	1,174.0	1,408.2
2003.....	51.2	90.4	150.5	244.3	385.5	546.4	754.1	978.0	1,189.2	1,453.0
2002.....	53.1	92.8	153.8	249.8	385.7	556.4	767.8	984.7	1,196.9	1,460.0
2001.....	54.3	94.4	158.0	255.5	391.9	560.4	771.5	984.7	1,197.7	1,483.0
2000.....	54.1	95.0	160.7	262.4	403.7	575.3	777.8	992.7	1,225.7	1,531.6
1999.....	54.0	94.1	161.4	265.4	405.4	582.2	783.0	986.4	1,209.2	1,508.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
II. Neoplasms (C00-D48)										
White, both sexes										
2006 4/.....	41.3	81.6	148.2	248.9	413.6	621.6	883.6	1,181.4	1,475.0	1,698.1
2005 4/.....	43.9	83.2	149.9	250.6	419.7	628.2	905.5	1,193.6	1,479.0	1,725.0
2004 4/.....	42.7	83.6	150.1	257.2	427.4	642.9	908.6	1,201.8	1,477.8	1,739.9
2003 4/.....	44.8	86.1	153.3	264.3	440.7	654.6	925.3	1,224.7	1,499.4	1,784.4
2002.....	46.5	87.7	156.7	272.3	445.0	669.1	948.1	1,234.8	1,504.1	1,806.4
2001.....	47.3	89.3	160.7	276.2	453.3	676.3	956.5	1,237.9	1,507.6	1,849.1
2000.....	46.7	89.6	161.8	285.0	463.7	695.8	961.4	1,243.6	1,555.2	1,903.1
1999.....	47.7	89.8	162.1	289.3	468.4	703.7	974.3	1,242.7	1,537.1	1,884.1
White, male										
2006 4/.....	36.6	78.7	156.6	275.3	470.3	732.3	1,060.7	1,478.9	1,966.3	2,352.0
2005 4/.....	39.7	80.6	159.0	275.4	477.3	744.2	1,104.4	1,509.7	1,956.8	2,419.6
2004 4/.....	38.2	82.5	160.9	282.0	486.7	757.4	1,098.9	1,526.1	1,950.8	2,452.1
2003 4/.....	40.8	86.4	162.0	288.9	501.8	775.0	1,129.8	1,556.3	1,993.4	2,517.5
2002.....	42.3	87.0	164.9	300.9	509.8	798.0	1,165.9	1,575.7	1,994.7	2,590.9
2001.....	42.9	88.3	167.6	302.3	521.3	810.7	1,183.4	1,588.2	2,014.4	2,707.4
2000.....	42.3	89.2	166.6	311.3	529.3	833.1	1,187.6	1,596.8	2,094.9	2,806.3
1999.....	44.2	89.4	167.0	320.0	540.8	844.6	1,216.4	1,608.8	2,087.0	2,810.7
White, female										
2006 4/.....	46.1	84.6	139.9	223.5	360.7	522.9	736.5	959.3	1,164.6	1,391.1
2005 4/.....	48.3	85.9	141.0	226.7	366.2	525.0	741.1	959.6	1,179.5	1,405.2
2004 4/.....	47.3	84.7	139.4	233.3	372.3	541.4	752.2	964.6	1,184.0	1,418.8
2003 4/.....	49.0	85.8	144.8	240.7	384.1	548.3	757.9	984.9	1,194.9	1,459.6
2002.....	50.6	88.4	148.7	245.0	385.2	555.6	770.6	990.7	1,203.4	1,465.6
2001.....	51.8	90.3	154.0	251.2	390.6	558.3	772.7	989.8	1,201.4	1,489.8
2000.....	51.1	89.9	157.1	260.0	403.2	575.6	779.9	995.3	1,234.1	1,536.2
1999.....	51.2	90.3	157.3	260.1	401.8	580.9	781.9	985.6	1,212.5	1,511.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
II. Neoplasms (C00-D48)										
Black, both sexes										
2006 4/.....	60.7	126.0	226.8	362.4	569.2	780.8	1,018.9	1,309.0	1,596.5	1,740.5
2005 4/.....	62.2	131.5	233.9	378.3	580.8	792.6	1,042.6	1,320.7	1,619.0	1,799.2
2004 4/.....	65.5	131.3	238.7	382.7	577.4	813.2	1,075.2	1,381.7	1,617.5	1,831.0
2003 4/.....	67.5	140.6	245.9	386.7	595.7	813.9	1,095.5	1,408.1	1,680.6	1,910.9
2002.....	70.0	137.1	241.3	404.8	602.9	843.8	1,141.1	1,450.2	1,707.0	1,949.3
2001.....	71.6	139.2	244.2	409.2	597.0	869.3	1,175.3	1,457.5	1,759.5	2,005.1
2000.....	75.8	148.0	255.0	412.1	611.8	872.0	1,199.5	1,508.3	1,738.5	2,063.0
1999.....	74.6	146.6	258.7	434.4	641.8	892.8	1,209.9	1,531.6	1,742.9	2,070.3
Black, male										
2006 4/.....	51.7	122.9	258.9	443.6	711.8	1,019.8	1,370.0	1,816.3	2,332.9	2,623.7
2005 4/.....	51.8	129.6	261.4	464.0	738.5	1,035.7	1,413.4	1,831.4	2,396.9	2,738.5
2004 4/.....	57.5	135.6	278.2	462.5	738.9	1,047.9	1,461.9	1,960.9	2,387.5	2,810.4
2003 4/.....	62.4	147.7	285.9	470.3	755.8	1,058.2	1,477.1	2,002.0	2,470.2	2,892.3
2002.....	61.7	138.8	274.3	491.8	784.4	1,092.0	1,552.3	2,077.8	2,567.7	3,055.5
2001.....	64.1	145.0	283.1	499.5	755.4	1,148.3	1,623.1	2,115.9	2,703.9	3,298.4
2000.....	71.7	154.0	297.3	517.2	779.4	1,152.9	1,678.6	2,200.5	2,722.0	3,340.9
1999.....	69.5	162.3	304.1	539.8	824.0	1,176.7	1,665.8	2,241.5	2,684.7	3,404.0
Black, female										
2006 4/.....	68.7	128.6	199.6	295.4	458.0	604.2	777.7	1,005.4	1,207.4	1,379.3
2005 4/.....	71.4	133.2	210.7	307.5	457.8	612.2	788.9	1,016.0	1,208.1	1,420.2
2004 4/.....	72.6	127.5	205.3	316.9	451.5	639.0	812.7	1,038.6	1,211.9	1,442.8
2003 4/.....	72.0	134.4	212.1	318.0	470.8	632.5	838.1	1,055.1	1,266.9	1,529.6
2002.....	77.3	135.7	213.2	333.3	460.9	659.6	865.2	1,075.3	1,258.9	1,525.5
2001.....	78.3	134.3	211.1	335.1	472.5	663.3	877.1	1,064.5	1,278.4	1,525.3
2000.....	79.5	142.8	218.9	325.9	479.5	664.7	881.3	1,091.1	1,243.1	1,587.4
1999.....	79.1	133.2	220.2	348.4	498.8	682.4	907.4	1,103.9	1,276.1	1,571.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
All races 3/, both sexes										
2006.....	42.3	84.1	152.1	252.7	414.6	615.1	863.3	1,144.9	1,418.8	1,606.1
2005.....	44.6	86.5	154.8	255.6	422.1	622.2	886.3	1,158.4	1,427.7	1,637.7
2004.....	44.3	86.7	155.6	261.5	427.6	637.8	892.4	1,172.1	1,424.8	1,653.3
2003.....	46.1	89.8	159.1	267.9	440.9	647.9	909.2	1,195.4	1,450.0	1,698.2
2002.....	47.5	90.6	161.5	276.8	447.0	663.8	933.6	1,207.0	1,458.3	1,723.9
2001.....	48.7	92.4	164.8	280.7	453.3	673.2	943.5	1,211.7	1,465.9	1,765.6
2000.....	48.8	93.6	166.2	289.0	463.4	690.6	951.6	1,222.5	1,505.3	1,819.4
1999.....	49.4	93.7	167.0	295.3	471.6	700.2	963.4	1,224.0	1,493.1	1,805.8
All races 3/, male										
2006.....	37.1	80.7	162.1	282.3	476.3	731.6	1,045.7	1,442.3	1,901.9	2,248.7
2005.....	39.5	83.4	164.9	284.8	484.5	743.6	1,091.8	1,475.1	1,901.6	2,319.3
2004.....	39.2	86.1	168.2	289.9	492.5	756.9	1,091.6	1,503.3	1,894.3	2,349.5
2003.....	41.7	90.6	169.8	295.9	507.6	774.4	1,117.9	1,533.3	1,939.7	2,413.8
2002.....	42.8	89.8	171.7	308.6	519.7	796.8	1,157.8	1,554.1	1,951.5	2,491.1
2001.....	44.0	91.7	174.2	311.6	526.7	814.1	1,178.0	1,568.4	1,978.7	2,613.6
2000.....	44.4	93.5	173.9	321.3	535.7	834.7	1,190.4	1,586.0	2,050.4	2,710.7
1999.....	45.5	94.7	174.9	330.6	550.1	848.2	1,212.9	1,600.1	2,046.0	2,722.2
All races 3/, female										
2006.....	47.6	87.5	142.6	224.8	358.2	513.3	714.8	927.2	1,116.9	1,305.5
2005.....	49.6	89.6	145.1	228.1	365.2	516.2	719.8	928.5	1,133.4	1,324.6
2004.....	49.2	87.4	143.5	234.8	368.5	534.2	732.0	934.2	1,135.5	1,340.1
2003.....	50.5	89.1	148.8	241.5	380.3	538.1	741.8	954.8	1,150.5	1,381.9
2002.....	52.2	91.4	151.8	246.8	381.1	548.9	754.4	961.9	1,158.4	1,391.1
2001.....	53.5	93.2	155.9	251.7	386.7	551.9	757.2	962.0	1,158.6	1,410.7
2000.....	53.2	93.7	158.8	258.9	398.0	567.2	763.5	969.3	1,183.8	1,456.4
1999.....	53.3	92.7	159.5	262.3	400.6	573.7	768.4	962.5	1,169.5	1,436.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
White, both sexes										
2006 4/.....	40.6	80.2	146.0	245.8	408.2	612.3	866.4	1,151.9	1,427.2	1,618.1
2005 4/.....	43.0	82.0	147.8	247.2	414.2	618.5	888.9	1,164.9	1,432.7	1,645.4
2004 4/.....	42.0	82.2	148.2	253.8	421.6	633.6	891.5	1,174.1	1,429.3	1,659.7
2003 4/.....	44.0	84.8	151.3	261.2	434.5	644.8	908.5	1,195.1	1,450.7	1,701.4
2002.....	45.5	86.2	154.7	268.7	439.5	659.2	931.4	1,205.1	1,456.9	1,725.7
2001.....	46.4	87.8	158.4	272.2	447.1	665.6	938.4	1,207.5	1,458.8	1,765.2
2000.....	45.8	88.2	159.5	281.1	457.1	685.0	944.1	1,214.2	1,503.0	1,815.7
1999.....	46.8	88.5	159.9	285.8	462.2	693.1	955.3	1,213.0	1,488.4	1,798.8
White, male										
2006 4/.....	35.8	77.2	154.0	271.5	464.1	721.0	1,038.6	1,439.4	1,903.8	2,258.3
2005 4/.....	38.7	79.2	156.6	271.7	470.2	731.9	1,083.4	1,472.2	1,894.9	2,322.7
2004 4/.....	37.4	81.1	158.7	278.1	479.7	745.4	1,077.9	1,490.7	1,887.6	2,348.9
2003 4/.....	39.8	85.1	159.7	285.4	494.7	763.2	1,108.1	1,518.9	1,930.1	2,412.1
2002.....	41.2	85.4	162.8	296.5	503.4	785.6	1,145.2	1,537.3	1,934.6	2,486.8
2001.....	41.9	86.5	165.1	298.2	513.9	797.6	1,160.4	1,547.8	1,950.7	2,597.6
2000.....	41.4	87.7	164.0	307.0	522.0	819.7	1,166.6	1,559.2	2,026.9	2,693.7
1999.....	43.1	88.0	164.3	316.0	532.7	831.3	1,191.9	1,571.1	2,024.1	2,696.3
White, female										
2006 4/.....	45.4	83.3	138.1	220.9	356.0	515.5	723.4	937.3	1,126.1	1,317.5
2005 4/.....	47.3	84.8	139.2	223.6	362.1	517.7	728.2	937.5	1,143.0	1,333.6
2004 4/.....	46.6	83.4	137.9	230.5	367.7	534.4	738.3	942.5	1,144.7	1,348.9
2003 4/.....	48.3	84.6	143.1	237.8	378.9	540.3	745.2	961.0	1,155.2	1,386.5
2002.....	49.9	86.9	146.8	242.0	380.4	548.0	757.1	967.4	1,164.2	1,395.1
2001.....	51.0	89.0	152.0	247.4	385.4	549.7	758.6	966.4	1,161.4	1,416.7
2000.....	50.1	88.7	155.2	256.5	397.2	567.2	765.6	971.5	1,191.3	1,459.1
1999.....	50.5	89.1	155.5	257.1	397.2	572.5	767.3	961.5	1,172.1	1,438.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms (C00-C97)										
Black, both sexes										
2006 4/.....	59.3	123.7	224.1	358.6	561.4	771.6	1,004.1	1,283.1	1,557.0	1,678.3
2005 4/.....	60.6	129.1	231.4	374.3	573.3	782.6	1,026.2	1,296.6	1,585.5	1,744.7
2004 4/.....	64.2	129.6	235.7	377.1	570.2	803.0	1,059.1	1,355.6	1,578.2	1,774.0
2003 4/.....	66.5	138.3	243.2	382.6	589.1	802.7	1,080.1	1,385.5	1,640.8	1,853.1
2002.....	68.3	135.1	238.2	399.6	597.1	833.2	1,126.1	1,425.8	1,665.1	1,885.7
2001.....	70.4	137.8	241.5	404.7	589.4	858.4	1,156.8	1,433.0	1,716.9	1,931.7
2000.....	74.4	145.6	251.9	407.9	604.6	861.4	1,179.2	1,481.7	1,700.6	1,998.9
1999.....	73.3	144.5	255.6	429.5	633.9	879.1	1,191.7	1,502.1	1,704.2	2,008.4
Black, male										
2006 4/.....	50.6	120.6	256.5	438.9	703.1	1,007.6	1,351.8	1,783.7	2,278.2	2,543.3
2005 4/.....	50.0	127.0	258.6	459.7	729.5	1,024.8	1,395.3	1,803.3	2,354.0	2,683.7
2004 4/.....	56.2	134.0	274.7	455.9	729.5	1,035.5	1,442.5	1,926.3	2,340.8	2,748.8
2003 4/.....	61.2	145.6	283.0	464.8	747.3	1,045.9	1,455.8	1,975.5	2,417.1	2,825.5
2002.....	60.4	137.1	271.5	484.4	777.8	1,079.3	1,536.2	2,047.9	2,514.3	2,976.1
2001.....	62.8	143.6	280.7	494.9	746.2	1,135.3	1,600.3	2,084.3	2,645.3	3,209.9
2000.....	70.3	151.8	293.7	512.8	769.2	1,137.3	1,654.9	2,166.0	2,683.2	3,264.8
1999.....	68.3	160.5	300.8	533.7	815.7	1,159.8	1,644.1	2,205.4	2,642.0	3,340.7
Black, female										
2006 4/.....	67.0	126.3	196.7	292.4	450.9	597.3	765.2	983.5	1,175.9	1,324.4
2005 4/.....	70.0	131.0	208.4	303.9	451.6	603.0	773.5	994.4	1,179.6	1,365.8
2004 4/.....	71.2	125.9	202.6	312.3	446.1	630.3	798.9	1,017.5	1,176.3	1,387.7
2003 4/.....	71.1	132.0	209.5	314.9	465.7	622.1	826.8	1,034.8	1,234.0	1,475.3
2002.....	75.2	133.5	210.0	330.0	455.6	650.5	850.9	1,054.1	1,223.1	1,468.0
2001.....	77.1	132.8	208.1	330.6	466.1	653.9	861.4	1,044.2	1,244.0	1,457.5
2000.....	78.1	140.2	216.2	321.9	474.8	657.8	863.3	1,069.2	1,205.8	1,527.7
1999.....	77.7	130.6	217.2	344.4	491.4	671.1	891.6	1,078.5	1,239.4	1,510.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
All races 3/, both sexes										
2006.....	0.9	1.8	3.5	5.2	6.6	8.7	10.6	13.0	14.1	16.3
2005.....	0.8	1.9	3.5	5.6	7.5	9.2	10.6	11.8	14.2	17.0
2004.....	0.9	2.0	3.8	5.1	7.5	9.4	11.0	12.7	14.9	18.0
2003.....	0.9	2.0	3.6	5.5	7.9	9.7	10.8	12.8	13.6	18.6
2002.....	0.8	2.0	3.6	5.8	8.1	8.8	12.1	13.1	14.1	18.7
2001.....	0.8	2.1	3.5	6.1	7.9	9.4	11.6	13.5	15.2	18.5
2000.....	0.9	2.0	3.8	5.7	8.1	9.9	11.5	12.6	14.0	19.4
1999.....	0.9	2.3	3.6	5.8	8.1	9.3	11.3	13.5	15.4	19.3
All races 3/, male										
2006.....	1.2	2.9	5.6	8.4	10.6	13.9	16.0	19.7	21.4	21.9
2005.....	1.2	2.8	5.7	9.1	11.9	14.8	16.0	17.9	20.4	22.0
2004.....	1.2	3.2	6.2	8.6	11.9	14.9	16.4	19.3	21.7	24.9
2003.....	1.4	3.2	6.2	9.1	12.6	15.0	16.4	19.0	19.9	26.9
2002.....	1.2	3.2	5.8	9.4	12.9	13.9	18.1	19.2	20.3	26.9
2001.....	1.2	3.2	5.8	10.0	13.0	14.3	17.9	19.6	22.1	25.8
2000.....	1.4	3.1	6.0	9.5	12.7	15.2	16.9	18.7	20.2	27.1
1999.....	1.3	3.6	5.7	9.0	13.1	14.3	17.4	20.1	22.3	27.1
All races 3/, female										
2006.....	0.5	0.8	1.4	2.1	2.9	4.2	6.3	8.1	9.6	13.7
2005.....	0.4	1.0	1.4	2.3	3.4	4.3	6.2	7.3	10.4	14.7
2004.....	0.5	0.9	1.5	1.7	3.6	4.7	6.7	8.0	10.8	14.9
2003.....	0.5	0.9	1.1	2.1	3.6	5.2	6.3	8.4	9.7	15.0
2002.....	0.4	0.9	1.4	2.5	3.7	4.4	7.3	8.8	10.3	15.1
2001.....	0.4	1.0	1.4	2.4	3.3	5.1	6.6	9.3	11.1	15.4
2000.....	0.4	1.0	1.6	2.2	3.9	5.4	7.2	8.3	10.4	16.3
1999.....	0.4	1.0	1.6	2.7	3.6	5.1	6.6	9.0	11.3	16.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
White, both sexes										
2006 4/.....	0.8	1.7	3.2	4.8	6.2	8.3	10.4	13.0	14.3	16.9
2005 4/.....	0.7	1.7	3.1	5.1	7.1	8.7	10.5	11.7	14.2	17.3
2004 4/.....	0.7	1.7	3.4	4.6	7.0	9.0	11.0	12.7	15.0	18.6
2003 4/.....	0.8	1.8	3.2	5.2	7.3	9.3	10.4	12.6	13.4	19.0
2002.....	0.7	1.7	3.1	5.3	7.5	8.4	12.3	13.3	14.2	18.9
2001.....	0.7	1.7	3.2	5.5	7.3	8.7	11.4	13.6	15.5	19.0
2000.....	0.7	1.7	3.2	5.0	7.3	9.4	11.1	12.5	14.0	19.7
1999.....	0.7	1.8	3.0	5.0	7.5	8.9	11.0	13.2	15.5	19.8
White, male										
2006 4/.....	1.1	2.6	5.2	7.8	9.8	12.9	15.3	19.5	21.3	22.4
2005 4/.....	1.0	2.4	5.1	8.1	11.2	13.8	15.3	17.3	20.0	22.1
2004 4/.....	0.9	2.6	5.4	7.8	10.9	14.0	15.8	18.8	21.5	24.8
2003 4/.....	1.3	2.7	5.5	8.5	11.3	14.1	15.2	18.2	19.2	27.2
2002.....	1.0	2.7	5.0	8.6	11.7	12.9	18.0	19.4	20.2	27.0
2001.....	1.0	2.6	5.2	8.9	11.8	13.2	17.1	19.2	22.3	26.4
2000.....	1.1	2.7	4.9	8.3	11.2	14.0	15.9	18.0	19.9	27.6
1999.....	1.1	2.7	4.7	7.6	12.0	13.5	16.8	19.3	22.0	27.3
White, female										
2006 4/.....	0.5	0.7	1.3	2.0	2.8	4.2	6.3	8.2	9.9	14.3
2005 4/.....	0.5	0.9	1.3	2.2	3.3	4.2	6.5	7.6	10.6	15.1
2004 4/.....	0.4	0.8	1.4	1.5	3.4	4.6	7.1	8.2	10.9	15.7
2003 4/.....	0.3	0.9	0.9	2.1	3.7	5.1	6.4	8.5	9.8	15.4
2002.....	0.4	0.7	1.3	2.2	3.6	4.3	7.6	8.8	10.5	15.3
2001.....	0.3	0.8	1.2	2.2	3.2	4.8	6.7	9.6	11.3	15.9
2000.....	0.4	0.8	1.6	1.9	3.6	5.3	7.2	8.6	10.6	16.5
1999.....	0.4	0.8	1.3	2.5	3.3	4.9	6.4	9.0	11.7	16.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)										
Black, both sexes										
2006 4/.....	1.2	3.2	5.6	8.4	11.1	13.4	14.3	14.2	12.5	10.4
2005 4/.....	1.1	3.6	6.6	9.2	12.6	14.1	13.7	13.0	14.3	12.6
2004 4/.....	1.9	4.6	7.8	9.5	13.3	13.8	13.7	14.9	15.8	12.4
2003 4/.....	1.7	3.9	6.5	8.8	14.4	15.5	15.3	14.8	16.0	15.5
2002.....	1.5	4.2	7.1	10.1	14.7	13.3	12.5	11.3	12.6	16.0
2001.....	1.7	4.3	6.2	12.2	13.1	15.3	15.7	14.2	13.3	12.9
2000.....	1.6	4.1	8.3	12.2	15.2	14.9	16.1	14.6	15.8	15.3
1999.....	1.5	5.9	8.8	12.6	14.6	14.4	16.2	17.7	15.6	14.3
Black, male										
2006 4/.....	1.7	5.1	9.1	15.0	19.8	25.5	24.9	25.1	25.3	*
2005 4/.....	2.1	6.1	11.7	17.5	22.4	26.6	25.9	26.8	28.8	21.4
2004 4/.....	2.9	7.6	13.8	17.0	23.7	26.2	27.2	25.9	27.3	26.8
2003 4/.....	1.9	7.0	11.4	16.3	27.8	27.6	30.2	26.5	30.3	25.0
2002.....	2.4	6.7	12.9	18.2	26.9	24.2	21.4	16.5	21.4	26.8
2001.....	2.4	7.0	10.8	22.2	24.8	25.4	29.3	25.1	20.9	*
2000.....	2.7	6.3	15.9	21.9	27.3	27.6	29.1	28.8	28.7	*
1999.....	2.3	10.3	14.5	21.5	25.2	24.1	26.5	33.2	28.8	24.6
Black, female										
2006 4/.....	*	1.5	2.6	3.0	4.3	4.5	7.0	7.8	*	8.8
2005 4/.....	*	1.5	2.2	2.4	5.0	4.8	5.4	*	*	9.0
2004 4/.....	*	1.9	2.7	3.4	5.2	4.7	4.6	8.4	9.8	*
2003 4/.....	1.5	*	2.3	2.7	3.9	6.6	5.3	7.9	8.5	11.7
2002.....	*	1.9	2.2	3.4	5.1	5.2	6.5	8.2	8.0	11.9
2001.....	*	2.0	2.2	4.0	3.8	7.9	6.7	7.8	9.4	10.5
2000.....	*	2.1	*	4.2	5.6	5.6	7.5	6.0	9.3	14.2
1999.....	*	2.1	3.9	5.3	6.3	7.1	9.5	8.3	9.1	10.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of tongue (C01-C02)										
All races 3/, both sexes										
2006.....	0.2	0.5	0.9	1.3	1.6	2.2	2.7	2.9	3.4	3.6
2005.....	0.2	0.5	0.9	1.4	1.9	2.2	2.8	2.8	3.5	4.0
2004.....	0.2	0.5	1.0	1.3	1.7	2.5	2.5	2.5	3.4	4.1
2003.....	0.3	0.5	1.0	1.5	1.8	2.4	2.6	3.1	3.0	3.7
2002.....	0.2	0.5	0.9	1.4	2.1	2.1	2.8	3.2	3.4	4.0
2001.....	0.2	0.5	0.9	1.6	1.8	2.0	2.6	3.1	3.5	3.8
2000.....	0.3	0.5	0.9	1.4	2.1	2.3	2.4	2.8	3.2	4.0
1999.....	0.3	0.6	0.9	1.3	1.8	2.2	2.6	3.0	3.4	4.0
All races 3/, male										
2006.....	0.3	0.8	1.5	2.1	2.6	3.4	3.8	4.1	4.4	4.0
2005.....	0.3	0.8	1.4	2.1	3.0	3.6	4.1	4.2	4.4	4.5
2004.....	0.3	0.7	1.6	2.3	2.8	3.8	3.4	3.7	4.5	4.8
2003.....	0.4	0.8	1.6	2.5	2.6	3.6	3.7	4.2	4.3	5.0
2002.....	0.3	0.7	1.4	2.3	3.1	3.5	3.8	4.5	4.3	5.2
2001.....	0.3	0.8	1.5	2.6	3.0	3.1	4.0	4.1	4.9	4.1
2000.....	0.3	0.8	1.5	2.2	3.4	3.5	3.3	3.6	3.8	5.1
1999.....	0.4	0.9	1.4	1.9	3.0	3.5	4.1	4.7	4.3	5.4
All races 3/, female										
2006.....	*	0.2	0.4	0.6	0.7	1.1	1.8	2.0	2.7	3.4
2005.....	*	0.2	0.3	0.7	1.0	0.9	1.8	1.7	2.9	3.8
2004.....	0.2	0.3	0.4	0.4	0.8	1.3	1.7	1.7	2.8	3.8
2003.....	*	0.3	0.4	0.5	1.1	1.3	1.8	2.3	2.2	3.2
2002.....	*	0.2	0.4	0.6	1.2	0.9	2.0	2.3	2.8	3.5
2001.....	*	0.3	0.4	0.6	0.7	1.1	1.4	2.4	2.7	3.6
2000.....	0.2	0.3	0.4	0.6	1.0	1.2	1.7	2.3	2.8	3.6
1999.....	0.2	0.3	0.4	0.7	0.8	1.1	1.4	1.8	2.8	3.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of tongue (C01-C02)										
White, both sexes										
2006 4/.....	0.3	0.5	0.9	1.3	1.7	2.1	2.8	2.9	3.4	3.8
2005 4/.....	0.2	0.5	0.8	1.4	1.9	2.2	2.8	2.8	3.6	4.2
2004 4/.....	0.2	0.5	1.0	1.3	1.8	2.4	2.5	2.7	3.5	4.2
2003 4/.....	0.2	0.5	0.9	1.4	1.8	2.3	2.6	3.1	3.0	3.8
2002.....	0.2	0.4	0.9	1.4	2.0	2.0	2.9	3.3	3.5	4.1
2001.....	0.2	0.5	0.9	1.6	1.8	1.9	2.6	3.2	3.5	3.9
2000.....	0.2	0.5	0.8	1.3	2.0	2.3	2.4	2.9	3.3	4.1
1999.....	0.3	0.5	0.8	1.1	1.8	2.2	2.6	3.0	3.5	4.1
White, male										
2006 4/.....	0.3	0.9	1.5	2.1	2.7	3.2	3.8	4.2	4.5	4.3
2005 4/.....	0.3	0.7	1.3	2.1	2.9	3.5	3.8	4.1	4.5	4.6
2004 4/.....	0.2	0.6	1.5	2.2	2.8	3.7	3.4	3.9	4.4	4.8
2003 4/.....	0.4	0.8	1.5	2.5	2.6	3.4	3.7	4.1	4.3	5.1
2002.....	0.3	0.6	1.4	2.2	3.0	3.2	3.9	4.6	4.4	5.2
2001.....	0.3	0.7	1.5	2.5	2.9	3.0	4.1	4.2	4.8	4.3
2000.....	0.3	0.7	1.3	2.2	3.3	3.5	3.3	3.5	3.8	5.1
1999.....	0.4	0.7	1.3	1.6	2.9	3.5	4.0	4.7	4.5	5.5
White, female										
2006 4/.....	*	*	0.3	0.5	0.7	1.1	2.0	2.0	2.8	3.6
2005 4/.....	*	0.3	0.3	0.7	1.0	1.0	1.9	1.9	3.1	4.0
2004 4/.....	*	0.3	0.4	0.5	0.9	1.3	1.9	1.8	2.9	3.9
2003 4/.....	*	0.3	0.3	0.5	1.1	1.3	1.8	2.4	2.2	3.3
2002.....	*	*	0.4	0.6	1.2	0.9	2.1	2.4	3.0	3.6
2001.....	*	0.3	0.4	0.7	0.7	1.0	1.4	2.5	2.8	3.8
2000.....	*	0.3	0.5	0.5	0.9	1.3	1.7	2.4	3.0	3.7
1999.....	*	0.3	0.4	0.7	0.8	1.0	1.4	1.9	3.0	3.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pharynx (C10-C13,C14.0)										
All races 3/, both sexes										
2006.....	0.4	0.8	1.4	2.4	2.9	3.9	4.6	5.5	5.8	4.6
2005.....	0.3	0.7	1.5	2.5	3.3	4.3	4.3	4.7	5.2	5.0
2004.....	0.4	1.0	1.6	2.2	3.6	4.2	4.9	5.6	5.6	5.1
2003.....	0.3	0.8	1.5	2.3	3.6	4.5	4.8	5.1	5.0	6.0
2002.....	0.3	0.8	1.6	2.5	3.5	3.8	5.2	5.2	4.9	5.4
2001.....	0.4	0.9	1.5	2.6	3.7	4.1	4.9	5.5	5.7	5.9
2000.....	0.4	0.8	1.6	2.7	3.7	4.3	5.1	5.2	5.3	5.6
1999.....	0.4	1.0	1.6	2.7	3.6	4.1	5.1	5.5	5.6	5.2
All races 3/, male										
2006.....	0.6	1.2	2.3	3.9	4.8	6.4	7.2	9.0	10.0	7.4
2005.....	0.5	1.1	2.5	4.3	5.5	6.8	7.0	7.3	8.6	8.0
2004.....	0.6	1.6	2.7	3.8	5.8	7.0	7.9	9.2	8.9	8.7
2003.....	0.5	1.3	2.7	3.9	5.9	7.2	7.6	8.2	8.7	10.8
2002.....	0.5	1.4	2.6	4.0	5.9	6.0	8.2	8.2	8.2	9.8
2001.....	0.6	1.5	2.6	4.4	6.1	6.5	7.9	8.8	9.0	10.0
2000.....	0.6	1.3	2.6	4.6	5.9	6.8	8.2	8.6	8.6	9.9
1999.....	0.6	1.6	2.6	4.5	5.9	6.5	8.0	8.7	9.6	9.9
All races 3/, female										
2006.....	0.2	0.4	0.5	0.9	1.1	1.6	2.5	2.9	3.1	3.4
2005.....	*	0.4	0.5	0.8	1.4	2.0	2.2	2.8	3.0	3.6
2004.....	0.2	0.4	0.6	0.8	1.6	1.9	2.5	3.1	3.5	3.5
2003.....	*	0.3	0.5	0.9	1.5	2.2	2.5	2.8	2.8	3.9
2002.....	*	0.3	0.6	1.2	1.4	1.8	2.9	3.1	2.8	3.5
2001.....	*	0.4	0.6	0.9	1.6	2.1	2.5	3.1	3.6	4.2
2000.....	*	0.3	0.6	1.0	1.6	2.2	2.8	2.9	3.3	3.9
1999.....	*	0.4	0.7	1.1	1.6	2.1	2.8	3.3	3.2	3.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pharynx (C10-C13,C14.0)										
White, both sexes										
2006 4/.....	0.3	0.6	1.2	2.0	2.6	3.6	4.2	5.3	5.6	4.5
2005 4/.....	0.3	0.6	1.2	2.2	3.0	3.9	4.1	4.5	4.9	4.8
2004 4/.....	0.3	0.8	1.3	1.8	3.0	3.8	4.7	5.3	5.4	5.1
2003 4/.....	0.3	0.6	1.2	2.1	3.1	4.2	4.4	4.8	4.6	5.8
2002.....	0.2	0.6	1.2	2.2	3.0	3.5	5.2	5.0	4.6	5.2
2001.....	0.2	0.7	1.2	2.2	3.3	3.7	4.6	5.2	5.5	6.0
2000.....	0.2	0.6	1.3	2.2	3.1	3.9	4.7	5.0	5.2	5.3
1999.....	0.2	0.6	1.2	2.3	3.2	3.8	4.7	5.1	5.3	5.3
White, male										
2006 4/.....	0.4	0.9	2.0	3.3	4.2	5.8	6.5	8.5	9.4	7.0
2005 4/.....	0.4	0.8	2.0	3.6	4.9	6.1	6.4	6.6	8.0	7.6
2004 4/.....	0.4	1.2	2.1	3.2	4.8	6.1	7.4	8.5	8.8	8.2
2003 4/.....	0.4	0.9	2.2	3.4	4.8	6.4	6.7	7.7	7.8	10.6
2002.....	0.3	1.0	2.0	3.4	4.9	5.4	8.1	8.0	7.6	9.5
2001.....	0.4	1.1	2.0	3.6	5.3	5.6	7.4	8.3	8.8	9.9
2000.....	0.3	0.9	2.0	3.7	4.8	6.1	7.3	7.8	8.4	9.5
1999.....	0.3	1.0	1.9	3.7	5.1	5.8	7.4	7.9	8.9	9.9
White, female										
2006 4/.....	*	0.3	0.5	0.8	1.0	1.6	2.3	3.0	3.2	3.3
2005 4/.....	*	0.3	0.5	0.8	1.2	1.9	2.2	2.9	3.0	3.6
2004 4/.....	*	0.3	0.5	0.6	1.4	1.7	2.5	3.0	3.4	3.7
2003 4/.....	*	0.3	0.3	0.8	1.5	2.1	2.5	2.8	2.7	3.7
2002.....	*	0.2	0.5	1.0	1.3	1.8	2.8	2.9	2.7	3.4
2001.....	*	*	0.4	0.8	1.5	2.1	2.4	3.1	3.5	4.3
2000.....	*	0.2	0.6	0.8	1.5	1.9	2.7	2.9	3.3	3.7
1999.....	*	0.3	0.5	0.9	1.4	1.9	2.6	3.2	3.1	3.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pharynx (C10-C13,C14.0)										
Black, both sexes										
2006 4/.....	*	1.8	2.7	4.8	5.6	6.4	8.6	7.4	7.8	5.4
2005 4/.....	*	1.6	3.4	5.0	7.0	7.5	6.8	7.5	7.5	6.7
2004 4/.....	1.0	2.3	4.3	5.4	9.3	7.8	8.2	9.1	6.7	5.6
2003 4/.....	0.7	2.0	3.3	4.2	8.4	8.6	8.5	7.7	8.1	7.3
2002.....	0.9	2.0	3.8	5.1	8.1	5.7	6.4	5.3	7.6	7.4
2001.....	0.9	2.2	3.7	6.5	6.4	7.4	7.8	7.3	7.6	*
2000.....	0.8	2.1	3.8	7.1	8.7	8.2	9.1	8.4	7.3	7.5
1999.....	0.8	3.0	4.6	6.5	7.5	7.5	8.9	10.2	9.0	*
Black, male										
2006 4/.....	*	3.0	4.6	9.2	10.6	12.4	15.4	15.2	18.5	*
2005 4/.....	*	2.7	6.5	9.7	12.5	14.3	13.6	16.1	17.6	*
2004 4/.....	1.5	4.0	7.6	9.2	16.8	14.8	16.2	16.8	*	*
2003 4/.....	*	3.7	6.0	8.0	16.4	16.5	17.1	14.6	17.7	*
2002.....	*	3.4	7.2	9.2	15.2	10.7	10.3	*	*	*
2001.....	*	3.4	6.5	11.7	11.6	14.5	14.5	13.5	*	*
2000.....	*	3.5	7.1	12.7	15.9	14.1	17.7	17.0	*	*
1999.....	*	5.4	8.1	11.2	12.9	13.0	14.6	19.9	17.4	*
Black, female										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	2.2	3.4	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	3.8	*	*	*	*
1999.....	*	*	*	*	*	*	5.2	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

All races 3/, both sexes

2006.....	0.3	0.5	1.2	1.5	2.1	2.7	3.2	4.5	4.9	7.7
2005.....	0.3	0.7	1.1	1.7	2.2	2.7	3.4	4.1	5.4	7.7
2004.....	0.2	0.5	1.2	1.5	2.1	2.7	3.6	4.4	5.6	8.4
2003.....	0.3	0.7	1.1	1.7	2.5	2.7	3.3	4.4	5.4	8.7
2002.....	0.2	0.7	1.1	1.9	2.4	2.9	4.0	4.5	5.7	8.7
2001.....	0.2	0.6	1.0	1.8	2.3	3.1	4.1	4.8	5.9	8.3
2000.....	0.3	0.7	1.2	1.6	2.2	3.3	3.8	4.4	5.2	9.5
1999.....	0.2	0.7	1.1	1.7	2.6	3.0	3.6	4.9	6.3	9.7

All races 3/, male

2006.....	0.4	0.8	1.8	2.4	3.2	4.1	4.8	6.3	6.9	9.7
2005.....	0.4	1.0	1.7	2.6	3.4	4.3	4.7	6.2	7.1	9.2
2004.....	0.4	0.9	1.9	2.5	3.2	4.0	5.0	6.1	7.8	10.8
2003.....	0.5	1.1	1.9	2.7	4.1	4.0	4.9	6.1	6.7	10.7
2002.....	0.3	1.1	1.8	3.1	3.8	4.4	5.9	6.2	7.5	11.1
2001.....	0.3	0.9	1.7	2.9	3.7	4.6	5.8	6.4	8.0	10.6
2000.....	0.4	1.1	1.9	2.6	3.3	4.8	5.2	6.3	7.4	11.4
1999.....	0.3	1.1	1.7	2.5	4.2	4.3	5.2	6.5	8.2	11.4

All races 3/, female

2006.....	*	0.3	0.5	0.6	1.1	1.5	1.9	3.2	3.7	6.8
2005.....	*	0.3	0.5	0.8	1.1	1.3	2.3	2.6	4.3	7.0
2004.....	*	0.2	0.5	0.5	1.2	1.5	2.5	3.2	4.3	7.3
2003.....	0.2	0.3	0.3	0.8	1.0	1.7	2.0	3.2	4.7	7.7
2002.....	*	0.3	0.4	0.7	1.1	1.6	2.4	3.4	4.6	7.7
2001.....	*	0.3	0.4	0.8	1.0	1.9	2.7	3.7	4.7	7.3
2000.....	*	0.4	0.5	0.6	1.3	1.9	2.8	3.1	4.0	8.8
1999.....	*	0.4	0.5	0.9	1.2	1.8	2.3	3.7	5.2	9.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites within the lip, oral cavity and pharynx (C03-C09,C14.2-C14.8)

White, both sexes

2006 4/.....	0.2	0.5	1.1	1.5	1.9	2.6	3.3	4.7	5.2	8.1
2005 4/.....	0.3	0.6	1.1	1.6	2.1	2.6	3.5	4.3	5.5	7.9
2004 4/.....	0.2	0.4	1.1	1.4	2.1	2.7	3.7	4.5	5.7	8.8
2003 4/.....	0.3	0.7	1.0	1.7	2.4	2.8	3.2	4.4	5.6	9.0
2002.....	0.2	0.7	1.0	1.8	2.4	2.9	4.1	4.7	5.9	9.1
2001.....	0.2	0.5	1.0	1.7	2.2	3.0	4.0	5.0	6.3	8.5
2000.....	0.3	0.6	1.1	1.4	2.1	3.2	3.9	4.6	5.2	9.9
1999.....	0.2	0.6	1.0	1.6	2.4	2.9	3.6	4.9	6.5	10.0

White, male

2006 4/.....	0.4	0.8	1.7	2.4	2.8	3.9	4.8	6.6	7.3	10.3
2005 4/.....	0.4	0.9	1.8	2.4	3.3	4.1	4.9	6.3	7.4	9.6
2004 4/.....	0.3	0.7	1.8	2.3	3.2	4.0	4.9	6.1	7.9	11.2
2003 4/.....	0.5	1.0	1.8	2.6	3.9	4.0	4.7	5.9	6.8	11.1
2002.....	0.3	1.0	1.7	2.9	3.8	4.4	5.9	6.4	7.8	11.5
2001.....	0.3	0.8	1.6	2.6	3.5	4.5	5.5	6.5	8.4	11.0
2000.....	0.4	1.0	1.6	2.3	3.1	4.4	5.2	6.4	7.2	12.2
1999.....	0.3	0.9	1.6	2.3	3.9	4.0	5.2	6.5	8.3	11.5

White, female

2006 4/.....	*	0.2	0.5	0.6	1.1	1.5	2.0	3.2	3.9	7.1
2005 4/.....	*	0.3	0.4	0.8	1.0	1.3	2.4	2.7	4.3	7.2
2004 4/.....	*	*	0.5	0.5	1.1	1.6	2.7	3.4	4.4	7.7
2003 4/.....	*	0.3	0.3	0.8	1.0	1.7	2.1	3.3	4.8	8.1
2002.....	*	0.3	0.4	0.7	1.1	1.6	2.6	3.5	4.7	8.0
2001.....	*	0.3	0.4	0.8	1.0	1.7	2.8	3.9	5.0	7.5
2000.....	*	0.3	0.5	0.6	1.3	2.1	2.8	3.2	4.1	8.9
1999.....	*	0.3	0.4	0.9	1.1	1.9	2.3	3.8	5.5	9.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of digestive organs (C15-C26)										
All races 3/, both sexes										
2006.....	9.5	20.3	39.8	64.3	98.2	141.0	197.1	266.4	343.1	420.0
2005.....	9.9	20.8	39.8	62.7	95.8	141.9	200.6	268.2	345.7	431.1
2004.....	9.6	20.5	39.3	62.7	99.2	142.7	202.9	270.8	345.7	445.3
2003.....	9.3	21.2	40.1	62.7	100.4	148.1	209.2	277.0	355.1	461.2
2002.....	10.1	21.1	39.5	64.8	101.0	152.4	212.5	281.9	354.0	467.4
2001.....	9.8	21.6	39.7	63.4	102.1	154.5	215.5	282.2	362.2	487.9
2000.....	10.0	21.7	38.8	65.0	105.6	153.7	217.5	290.1	374.6	504.9
1999.....	10.2	21.8	38.1	65.5	106.6	157.7	218.4	288.9	367.2	507.4
All races 3/, male										
2006.....	11.1	25.9	53.1	87.4	131.8	187.6	256.2	337.5	441.7	499.9
2005.....	11.7	26.5	53.9	85.7	129.0	188.8	263.5	347.7	437.3	513.8
2004.....	11.8	27.1	53.7	84.7	131.5	188.8	264.4	349.3	437.2	531.7
2003.....	11.4	28.0	53.7	84.2	133.1	196.2	271.8	356.3	455.5	546.7
2002.....	12.7	27.5	53.5	86.9	135.8	202.1	278.3	364.4	449.0	557.9
2001.....	11.6	28.1	52.7	84.4	136.1	203.6	278.0	366.3	458.9	593.2
2000.....	12.3	28.4	51.4	86.2	139.3	202.8	281.4	377.0	474.6	614.1
1999.....	12.5	29.0	50.4	86.9	138.8	205.8	285.0	374.7	472.3	620.8
All races 3/, female										
2006.....	7.9	14.8	27.0	42.5	67.6	100.2	149.0	214.3	281.5	382.6
2005.....	8.0	15.3	26.3	41.0	65.6	100.9	149.6	210.4	288.9	393.1
2004.....	7.4	14.1	25.5	42.1	69.8	102.7	153.3	214.4	289.4	406.4
2003.....	7.2	14.7	27.0	42.4	70.7	106.4	159.0	220.5	293.6	423.4
2002.....	7.5	14.8	26.1	44.1	69.4	109.5	159.9	223.5	296.1	428.1
2001.....	7.9	15.3	27.2	43.8	71.3	112.3	165.8	223.4	304.3	443.9
2000.....	7.7	15.2	26.6	45.1	75.1	111.6	167.2	229.6	315.5	460.4
1999.....	8.0	14.9	26.2	45.5	77.5	116.7	166.3	229.3	305.7	461.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of digestive organs (C15-C26)										
White, both sexes										
2006 4/.....	8.6	18.6	37.0	60.4	93.2	135.4	192.2	259.9	336.1	416.3
2005 4/.....	9.2	19.1	36.5	58.6	90.4	135.8	195.3	261.1	338.3	425.7
2004 4/.....	8.7	18.6	36.2	58.5	93.9	136.5	195.2	264.7	340.6	440.0
2003 4/.....	8.2	19.1	36.0	58.7	94.9	143.0	202.2	270.8	348.6	453.8
2002.....	9.1	19.2	36.3	60.5	95.8	145.8	205.6	274.7	346.3	460.4
2001.....	8.8	19.6	36.6	58.8	97.3	147.3	208.2	274.5	354.1	481.6
2000.....	8.8	19.6	35.6	60.3	100.2	147.0	210.3	281.8	365.9	497.0
1999.....	9.1	19.9	34.8	60.2	100.9	151.7	211.0	280.4	358.9	499.7
White, male										
2006 4/.....	10.0	24.0	49.2	82.1	125.1	180.7	249.9	330.4	434.5	496.7
2005 4/.....	11.0	24.5	49.4	79.8	121.8	180.3	257.3	339.1	429.1	510.8
2004 4/.....	10.7	24.7	49.8	78.7	124.7	180.8	253.8	341.7	432.2	528.5
2003 4/.....	10.2	25.1	48.3	79.4	125.7	189.7	263.8	348.6	450.2	539.4
2002.....	11.6	25.3	49.1	81.2	129.0	194.6	269.7	356.9	439.1	550.5
2001.....	10.5	25.5	48.6	78.4	130.0	194.4	269.8	358.3	449.9	588.1
2000.....	10.9	25.6	47.1	79.3	132.3	195.1	271.9	368.1	464.1	605.3
1999.....	11.3	26.3	45.8	79.5	131.0	198.2	277.1	363.3	463.6	614.4
White, female										
2006 4/.....	7.2	13.3	25.1	39.5	63.5	95.0	144.3	207.3	273.9	378.5
2005 4/.....	7.2	13.7	24.0	38.3	61.2	96.2	144.1	203.4	281.3	386.5
2004 4/.....	6.7	12.5	23.0	39.0	65.3	97.3	147.1	208.4	283.7	400.0
2003 4/.....	6.3	13.2	24.1	38.9	66.4	101.7	151.9	214.5	285.9	415.8
2002.....	6.7	13.1	23.9	40.6	65.2	102.9	153.3	215.9	289.5	421.3
2001.....	7.2	13.7	24.8	40.1	67.2	105.9	158.4	215.0	296.1	437.0
2000.....	6.8	13.6	24.4	42.3	70.6	105.0	160.8	221.1	307.4	453.1
1999.....	7.0	13.6	24.1	41.8	73.1	111.2	158.5	222.3	297.0	453.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of digestive organs (C15-C26)										
Black, both sexes										
2006 4/.....	14.7	31.9	64.2	101.0	151.8	200.1	256.4	347.3	440.9	486.7
2005 4/.....	13.9	32.9	66.1	102.0	149.6	204.6	260.5	356.9	442.7	515.6
2004 4/.....	14.8	33.7	63.3	102.7	151.9	210.1	284.1	352.2	418.9	524.2
2003 4/.....	15.5	35.7	71.8	102.0	156.0	201.4	287.0	359.2	450.1	567.1
2002.....	16.2	34.7	64.6	107.2	151.8	217.4	295.9	380.0	450.0	566.6
2001.....	15.7	36.7	65.8	107.4	150.7	224.7	297.8	380.2	475.1	585.6
2000.....	16.8	37.8	68.1	105.8	158.3	221.2	302.3	395.9	483.7	606.9
1999.....	17.0	36.1	66.2	112.9	160.5	222.0	306.6	395.8	484.5	613.1
Black, male										
2006 4/.....	18.0	40.0	89.4	144.4	207.1	270.4	346.0	453.7	569.5	571.0
2005 4/.....	16.1	41.8	92.5	147.8	211.0	282.3	341.6	474.7	560.4	589.4
2004 4/.....	18.2	44.3	87.2	143.2	206.1	286.1	390.4	458.6	536.6	575.2
2003 4/.....	19.6	49.0	101.0	138.7	214.4	268.9	386.3	467.4	547.8	654.8
2002.....	20.1	44.6	90.5	147.5	206.9	288.2	394.9	489.1	600.7	671.1
2001.....	19.1	48.4	89.0	146.3	199.9	305.9	385.4	489.7	602.7	676.3
2000.....	21.6	50.2	93.2	152.0	212.2	289.6	403.5	508.4	629.7	728.8
1999.....	21.1	50.3	92.5	159.0	216.8	292.9	397.5	543.8	615.6	715.9
Black, female										
2006 4/.....	11.9	24.9	42.9	65.2	108.7	148.2	194.8	283.6	372.9	452.2
2005 4/.....	12.0	25.2	43.7	64.2	101.7	147.0	205.0	286.7	380.6	485.8
2004 4/.....	11.8	24.5	43.1	69.4	109.5	153.6	212.0	289.2	356.8	504.0
2003 4/.....	11.9	24.1	47.1	71.8	110.4	151.3	220.0	294.9	398.8	533.1
2002.....	12.7	26.1	42.7	74.1	108.7	164.8	229.4	314.8	371.6	526.6
2001.....	12.7	26.5	46.1	75.6	112.0	164.6	239.4	314.8	410.0	551.9
2000.....	12.5	27.0	46.8	67.9	115.7	170.7	235.1	328.1	410.1	561.5
1999.....	13.4	23.9	44.0	75.2	116.3	169.5	246.2	306.7	419.6	574.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of esophagus (C15)										
All races 3/, both sexes										
2006.....	0.9	2.3	4.8	8.4	13.2	17.4	22.0	26.9	27.0	25.6
2005.....	1.1	2.3	4.9	8.4	12.6	17.5	22.4	27.6	27.3	24.9
2004.....	1.0	2.3	4.7	8.1	13.2	17.5	21.3	26.0	27.7	25.8
2003.....	0.9	2.4	4.6	8.8	12.8	17.6	22.2	25.4	27.6	25.0
2002.....	1.0	2.4	4.7	8.6	12.7	18.1	22.5	25.7	27.8	24.6
2001.....	1.1	2.4	5.0	7.9	13.0	18.5	22.9	26.1	26.5	25.8
2000.....	1.0	2.4	4.9	8.8	13.2	17.9	22.1	24.5	27.2	25.9
1999.....	1.0	2.4	4.7	8.7	13.0	18.2	22.0	24.3	24.6	26.8
All races 3/, male										
2006.....	1.5	4.1	8.1	15.0	23.2	30.8	38.7	47.0	51.3	48.5
2005.....	1.7	3.9	8.8	14.8	22.2	31.1	39.6	49.6	52.4	46.0
2004.....	1.6	4.0	8.1	14.1	22.9	30.5	37.0	46.8	51.7	48.5
2003.....	1.5	4.1	8.1	15.3	22.7	30.6	38.3	45.7	51.0	45.3
2002.....	1.6	4.1	8.3	15.1	22.1	31.4	40.3	46.8	50.6	43.4
2001.....	1.8	4.1	8.7	13.7	22.6	32.6	39.9	46.5	48.8	48.1
2000.....	1.8	4.2	8.7	15.2	23.0	30.7	38.0	43.9	48.8	47.8
1999.....	1.7	4.2	8.3	15.3	21.9	31.7	39.3	44.1	44.8	48.0
All races 3/, female										
2006.....	0.2	0.7	1.7	2.2	4.1	5.8	8.4	12.2	11.8	14.9
2005.....	0.4	0.7	1.2	2.5	3.9	5.6	8.5	11.6	11.6	15.3
2004.....	0.3	0.7	1.4	2.4	4.3	6.1	8.7	11.1	12.9	15.7
2003.....	0.3	0.7	1.3	2.6	3.8	6.3	9.2	11.0	13.3	16.1
2002.....	0.4	0.7	1.3	2.4	4.2	6.7	8.2	10.9	14.0	16.5
2001.....	0.3	0.7	1.4	2.4	4.2	6.4	9.4	11.9	13.2	16.5
2000.....	0.2	0.6	1.4	2.8	4.4	7.1	9.6	11.0	14.5	17.1
1999.....	0.3	0.6	1.3	2.5	5.0	6.7	8.5	10.5	12.8	18.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of esophagus (C15)										
White, both sexes										
2006 4/.....	0.9	2.3	4.7	8.3	13.3	17.5	22.5	27.4	27.9	26.4
2005 4/.....	1.1	2.3	4.9	8.3	12.5	17.6	22.9	28.0	28.0	25.9
2004 4/.....	1.0	2.2	4.6	7.9	12.8	17.4	21.2	26.5	28.0	26.2
2003 4/.....	0.9	2.3	4.3	8.5	12.3	17.0	22.5	25.8	27.8	25.8
2002.....	0.9	2.3	4.6	8.1	12.2	17.5	22.4	25.7	28.2	25.1
2001.....	1.0	2.3	4.6	7.1	12.3	17.7	22.1	26.2	26.9	26.3
2000.....	0.9	2.3	4.5	8.1	12.4	17.4	21.8	24.0	27.2	26.6
1999.....	0.9	2.2	4.3	8.0	12.1	17.5	21.5	23.7	24.9	27.2
White, male										
2006 4/.....	1.7	4.1	8.0	15.0	23.4	31.1	39.3	47.9	53.7	49.9
2005 4/.....	1.8	3.9	9.0	14.7	22.0	31.2	40.4	50.6	53.7	48.5
2004 4/.....	1.7	3.9	8.0	13.9	22.5	30.5	37.2	47.6	52.8	49.2
2003 4/.....	1.6	3.9	7.6	15.0	22.0	29.5	39.0	46.5	52.3	46.7
2002.....	1.6	4.0	8.1	14.5	21.5	30.7	40.2	47.2	51.3	44.6
2001.....	1.8	3.9	8.1	12.5	21.6	31.2	39.0	46.9	49.5	48.8
2000.....	1.8	4.1	8.0	14.1	22.0	30.0	37.5	43.2	49.7	49.3
1999.....	1.6	4.0	7.6	14.1	20.5	30.8	38.4	43.0	45.6	48.0
White, female										
2006 4/.....	0.2	0.6	1.5	1.8	3.8	5.5	8.5	12.1	11.6	15.3
2005 4/.....	0.3	0.6	1.0	2.2	3.6	5.5	8.4	11.3	11.8	15.5
2004 4/.....	0.3	0.5	1.1	2.1	3.8	5.9	8.1	11.1	12.6	15.9
2003 4/.....	0.3	0.6	1.0	2.3	3.4	6.0	9.0	10.9	12.7	16.5
2002.....	0.3	0.6	1.2	2.1	3.7	5.9	7.8	10.3	14.1	16.7
2001.....	0.3	0.6	1.2	1.9	3.7	5.8	8.5	11.6	13.2	16.8
2000.....	*	0.4	1.1	2.4	3.6	6.4	9.1	10.6	13.8	17.3
1999.....	*	0.5	1.0	2.1	4.4	5.8	8.0	10.1	12.6	18.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of esophagus (C15)										
Black, both sexes										
2006 4/.....	0.8	3.0	7.4	11.5	16.9	21.5	22.8	25.1	23.7	18.2
2005 4/.....	1.3	3.2	6.8	11.8	17.4	21.4	23.9	30.2	22.6	17.4
2004 4/.....	1.2	4.0	6.6	12.0	20.4	23.0	26.8	27.4	27.0	21.4
2003 4/.....	1.3	4.1	8.4	14.1	20.7	26.1	24.9	26.9	29.4	21.3
2002.....	1.6	3.6	6.9	14.9	20.4	27.5	29.8	31.8	26.2	20.8
2001.....	1.6	3.7	8.6	15.9	21.9	29.6	35.8	30.7	27.6	23.6
2000.....	1.5	4.1	9.8	16.9	23.3	27.0	29.6	36.1	29.9	22.2
1999.....	1.7	4.3	9.4	17.1	23.5	28.7	31.3	35.7	23.7	23.9
Black, male										
2006 4/.....	*	4.8	12.2	18.7	28.8	38.2	42.1	44.3	39.0	34.9
2005 4/.....	2.1	5.1	11.4	19.8	31.3	39.7	41.4	52.7	48.5	27.0
2004 4/.....	1.7	6.3	10.8	19.9	33.4	40.0	43.1	49.6	45.2	38.7
2003 4/.....	2.1	7.1	14.0	23.3	36.2	46.5	41.4	47.5	43.6	38.6
2002.....	2.5	5.7	11.5	26.8	34.4	45.4	54.4	54.2	48.1	37.5
2001.....	2.6	6.0	15.3	27.5	38.8	51.8	59.9	55.3	53.8	46.5
2000.....	2.4	6.3	17.4	30.5	38.5	44.8	49.8	65.7	45.5	38.1
1999.....	3.2	7.4	16.4	30.3	40.3	46.5	56.0	67.9	36.6	51.6
Black, female										
2006 4/.....	*	1.4	3.3	5.6	7.6	9.2	9.5	13.7	15.6	11.3
2005 4/.....	*	1.6	2.9	5.2	6.5	7.8	11.9	16.8	8.9	13.5
2004 4/.....	*	1.9	3.1	5.5	10.2	10.4	15.8	14.3	17.4	14.6
2003 4/.....	*	1.6	3.6	6.5	8.6	11.0	13.7	14.7	22.0	14.6
2002.....	*	1.8	3.0	5.1	9.4	14.2	13.2	18.4	14.7	14.4
2001.....	*	1.7	2.9	6.3	8.6	13.1	19.7	16.1	14.3	15.2
2000.....	*	2.1	3.4	5.8	11.3	13.8	16.2	18.2	22.1	16.3
1999.....	*	1.6	3.4	6.4	10.3	15.6	14.9	16.3	17.3	13.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of stomach (C16)										
All races 3/, both sexes										
2006.....	1.3	2.1	3.3	5.0	7.2	11.0	15.5	22.3	29.9	35.8
2005.....	1.3	2.1	3.1	4.8	7.3	11.8	16.7	23.5	30.6	38.5
2004.....	1.4	2.1	3.5	5.3	8.0	12.1	18.2	23.4	30.3	42.2
2003.....	1.3	2.3	3.8	5.3	8.6	12.4	18.4	24.2	32.2	43.6
2002.....	1.2	2.1	3.5	5.6	8.5	12.3	18.8	25.6	33.0	47.6
2001.....	1.3	2.2	3.7	5.3	9.2	13.7	19.5	27.0	34.7	45.4
2000.....	1.4	2.3	3.6	5.6	9.7	13.8	19.6	27.1	36.9	52.7
1999.....	1.4	2.3	3.8	6.2	9.8	14.2	20.2	28.1	36.8	50.1
All races 3/, male										
2006.....	1.4	2.6	4.4	7.0	10.1	15.4	21.4	31.3	44.7	50.2
2005.....	1.6	2.5	4.3	6.6	10.4	16.4	23.8	32.5	44.3	54.2
2004.....	1.7	2.6	4.6	7.5	11.2	17.0	24.9	33.6	42.5	62.9
2003.....	1.5	2.8	5.0	7.5	12.3	17.9	26.6	34.2	44.7	60.0
2002.....	1.4	2.7	4.8	7.7	11.9	18.1	27.5	36.4	47.1	67.2
2001.....	1.4	2.8	4.9	7.5	13.7	19.5	27.5	39.1	49.4	62.6
2000.....	1.7	2.9	4.8	7.8	14.1	19.5	28.3	39.4	51.4	75.6
1999.....	1.7	3.1	5.4	8.8	13.7	20.3	28.6	41.5	54.9	71.3
All races 3/, female										
2006.....	1.2	1.7	2.3	3.1	4.5	7.2	10.6	15.8	20.7	29.1
2005.....	1.1	1.8	2.0	3.0	4.5	7.7	11.0	17.0	22.1	31.2
2004.....	1.2	1.6	2.4	3.3	5.1	7.8	12.8	16.0	22.7	32.9
2003.....	1.1	1.7	2.6	3.1	5.3	7.7	11.9	17.1	24.5	36.4
2002.....	1.0	1.5	2.3	3.7	5.4	7.3	11.9	18.1	24.4	39.1
2001.....	1.1	1.6	2.5	3.2	5.1	8.8	13.1	18.5	25.9	38.3
2000.....	1.2	1.7	2.4	3.5	5.8	8.9	12.7	18.5	28.3	43.4
1999.....	1.2	1.5	2.4	3.7	6.4	9.1	13.7	18.8	26.2	41.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of stomach (C16)										
White, both sexes										
2006 4/.....	1.0	1.8	3.0	4.4	6.1	9.5	13.5	19.6	26.1	32.4
2005 4/.....	1.1	1.7	2.7	4.2	6.2	10.0	14.7	20.5	27.0	34.8
2004 4/.....	1.2	1.7	2.9	4.5	6.9	10.2	15.9	20.9	27.3	38.8
2003 4/.....	1.1	1.8	3.1	4.6	7.6	11.0	16.3	21.4	28.8	38.7
2002.....	1.0	1.8	2.9	4.9	7.3	10.5	16.5	22.6	29.3	42.2
2001.....	1.1	1.9	3.1	4.7	7.9	11.7	17.4	24.0	31.0	41.1
2000.....	1.1	1.9	3.1	4.9	8.5	11.9	17.2	23.6	32.7	47.3
1999.....	1.1	2.0	3.1	5.2	8.3	12.8	17.7	24.8	33.0	44.8
White, male										
2006 4/.....	1.1	2.2	3.9	6.2	8.6	13.4	18.8	27.8	39.2	45.8
2005 4/.....	1.3	2.0	3.7	5.8	8.6	13.7	20.7	28.7	39.1	49.0
2004 4/.....	1.4	2.1	3.9	6.5	9.5	14.4	21.3	29.9	38.0	58.4
2003 4/.....	1.2	2.3	4.1	6.6	10.6	15.5	23.5	30.3	40.3	53.5
2002.....	1.1	2.4	3.9	6.8	10.3	15.7	24.3	32.8	42.4	60.5
2001.....	1.2	2.4	4.1	6.5	11.7	16.3	24.5	35.4	44.0	57.7
2000.....	1.2	2.3	4.2	6.7	12.5	17.1	25.1	34.6	46.1	68.9
1999.....	1.2	2.7	4.4	7.6	11.5	18.4	25.2	36.9	49.0	63.4
White, female										
2006 4/.....	1.0	1.3	2.0	2.6	3.9	6.0	9.1	13.5	17.8	26.1
2005 4/.....	0.9	1.5	1.8	2.6	3.9	6.7	9.7	14.5	19.5	28.2
2004 4/.....	1.0	1.3	2.0	2.7	4.4	6.5	11.5	14.2	20.7	29.9
2003 4/.....	0.9	1.4	2.2	2.7	4.8	7.0	10.5	15.0	21.7	32.1
2002.....	0.8	1.2	1.8	3.0	4.6	5.9	10.1	15.4	21.3	34.2
2001.....	1.0	1.3	2.1	2.9	4.3	7.7	11.7	16.0	23.1	34.1
2000.....	1.0	1.5	2.0	3.1	4.9	7.4	10.9	15.8	24.7	38.6
1999.....	0.9	1.2	1.9	2.9	5.4	8.0	11.7	16.3	23.5	37.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of stomach (C16)										
Black, both sexes										
2006 4/.....	2.3	3.6	5.3	8.3	14.7	20.7	28.4	42.1	64.4	66.9
2005 4/.....	2.3	4.0	5.2	9.0	14.4	23.5	30.1	48.6	57.3	74.3
2004 4/.....	2.7	3.9	6.9	10.6	15.8	27.3	34.0	43.8	59.9	71.3
2003 4/.....	2.4	4.3	7.4	9.6	16.7	22.3	33.1	48.8	62.4	85.8
2002.....	2.2	3.6	7.5	10.4	16.1	24.6	38.9	52.7	63.1	100.4
2001.....	2.5	3.7	7.5	9.2	17.7	27.1	34.7	51.1	69.9	88.1
2000.....	2.9	4.7	7.0	11.1	17.3	26.8	38.2	56.9	75.1	101.7
1999.....	3.2	3.9	8.5	13.2	19.1	23.6	40.3	58.6	74.3	102.4
Black, male										
2006 4/.....	2.9	4.7	7.6	12.5	21.9	30.7	38.1	67.6	100.6	84.0
2005 4/.....	3.2	5.4	7.7	13.5	24.2	34.9	48.4	67.5	79.4	114.3
2004 4/.....	3.1	5.3	9.5	15.3	25.2	40.7	56.0	63.7	93.3	100.3
2003 4/.....	3.3	6.2	10.9	14.9	26.0	36.4	54.2	74.9	87.1	106.3
2002.....	2.4	5.1	11.0	15.2	24.0	36.9	59.8	73.5	90.2	137.2
2001.....	2.8	5.0	11.5	13.8	27.5	43.7	54.2	75.7	106.7	104.4
2000.....	3.9	6.6	9.6	16.5	25.9	38.5	57.2	83.2	105.4	136.1
1999.....	4.0	5.7	12.2	19.0	28.2	35.1	60.4	89.3	121.2	147.6
Black, female										
2006 4/.....	1.7	2.6	3.3	4.9	9.1	13.3	21.8	26.8	45.2	60.0
2005 4/.....	1.5	2.8	3.1	5.3	6.8	15.0	17.5	37.3	45.6	58.1
2004 4/.....	2.3	2.7	4.7	6.7	8.4	17.3	19.1	32.1	42.3	59.9
2003 4/.....	1.5	2.7	4.5	5.3	9.5	11.9	18.8	33.4	49.5	77.8
2002.....	2.1	2.3	4.6	6.5	9.9	15.5	24.9	40.3	48.9	86.3
2001.....	2.2	2.5	4.1	5.4	10.1	14.8	21.7	36.3	51.1	82.1
2000.....	1.9	3.0	4.7	6.6	10.5	18.2	25.5	41.0	59.9	88.9
1999.....	2.4	2.4	5.4	8.4	12.0	15.2	27.0	40.0	51.0	85.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of small intestine (C17)										
All races 3/, both sexes										
2006.....	0.2	0.2	0.4	0.5	0.8	1.1	1.8	2.2	2.7	2.9
2005.....	0.1	0.2	0.3	0.6	0.8	1.3	1.5	2.2	2.9	2.9
2004.....	0.1	0.2	0.4	0.5	0.9	1.4	1.6	2.1	2.7	3.1
2003.....	0.1	0.2	0.3	0.6	0.9	1.1	1.6	2.1	3.0	3.0
2002.....	0.1	0.2	0.4	0.6	0.7	1.2	1.6	2.1	2.6	2.8
2001.....	0.1	0.2	0.4	0.5	1.0	1.0	1.9	2.4	2.6	3.2
2000.....	0.1	0.3	0.4	0.6	1.0	1.2	1.7	2.2	2.8	3.0
1999.....	0.1	0.4	0.5	0.4	0.9	1.2	1.7	2.2	2.6	3.2
All races 3/, male										
2006.....	*	0.2	0.4	0.5	0.8	1.3	2.4	2.4	3.9	3.7
2005.....	*	0.3	0.3	0.6	1.0	1.6	2.1	3.0	3.4	2.7
2004.....	*	0.2	0.4	0.7	1.1	1.8	2.1	2.4	3.9	4.3
2003.....	*	0.3	0.3	0.9	1.0	1.5	2.0	2.2	3.5	3.5
2002.....	0.2	0.3	0.4	0.6	1.0	1.4	1.9	2.7	2.9	3.7
2001.....	*	0.3	0.5	0.6	1.3	1.3	2.5	2.7	3.3	4.1
2000.....	*	0.3	0.4	0.6	1.2	1.4	2.2	2.5	3.7	4.2
1999.....	0.2	0.4	0.5	0.6	1.4	1.5	2.0	3.0	3.6	4.4
All races 3/, female										
2006.....	*	*	0.3	0.5	0.7	0.9	1.3	2.1	1.9	2.5
2005.....	*	*	0.3	0.5	0.7	1.1	1.0	1.7	2.6	3.0
2004.....	*	0.2	0.4	0.4	0.8	1.1	1.2	1.9	2.0	2.5
2003.....	*	0.2	0.4	0.4	0.7	0.7	1.3	2.0	2.8	2.7
2002.....	*	*	0.3	0.5	0.5	1.1	1.3	1.7	2.5	2.4
2001.....	*	0.2	0.2	0.5	0.8	0.8	1.4	2.2	2.1	2.8
2000.....	*	0.2	0.3	0.5	0.8	1.1	1.4	1.9	2.3	2.5
1999.....	*	0.3	0.4	0.3	0.4	0.9	1.4	1.7	2.0	2.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of small intestine (C17)										
White, both sexes										
2006 4/.....	0.1	0.2	0.3	0.5	0.7	1.1	1.6	2.2	2.6	2.9
2005 4/.....	0.1	0.2	0.3	0.5	0.8	1.3	1.5	2.3	2.8	3.0
2004 4/.....	*	0.2	0.3	0.5	0.8	1.4	1.5	2.0	2.7	2.9
2003 4/.....	*	0.2	0.3	0.6	0.8	1.1	1.6	2.0	3.0	2.9
2002.....	0.1	0.2	0.4	0.5	0.6	1.2	1.6	2.0	2.5	2.6
2001.....	0.1	0.2	0.3	0.5	1.0	1.0	1.8	2.4	2.5	3.0
2000.....	0.1	0.2	0.3	0.6	1.0	1.2	1.7	2.1	2.7	3.1
1999.....	0.1	0.3	0.4	0.4	0.8	1.2	1.6	2.1	2.4	3.0
White, male										
2006 4/.....	*	0.2	0.4	0.5	0.8	1.3	2.2	2.4	3.8	3.6
2005 4/.....	*	0.2	0.3	0.5	0.9	1.6	2.1	3.0	3.4	2.9
2004 4/.....	*	0.2	0.4	0.7	0.9	1.7	1.9	2.3	3.8	4.1
2003 4/.....	*	0.2	0.3	0.8	1.0	1.4	1.8	2.1	3.4	3.3
2002.....	*	0.2	0.4	0.6	0.9	1.3	1.9	2.6	2.8	3.6
2001.....	*	*	0.4	0.5	1.2	1.2	2.5	2.7	3.3	3.7
2000.....	*	0.3	0.4	0.6	1.2	1.4	2.1	2.4	3.6	4.3
1999.....	*	0.4	0.5	0.5	1.3	1.5	1.9	2.8	3.3	4.4
White, female										
2006 4/.....	*	*	0.3	0.5	0.6	0.9	1.1	2.0	1.8	2.6
2005 4/.....	*	*	*	0.5	0.7	1.0	1.0	1.7	2.5	3.0
2004 4/.....	*	0.2	0.3	0.3	0.8	1.1	1.1	1.8	2.1	2.3
2003 4/.....	*	*	0.3	0.4	0.7	0.8	1.5	2.0	2.7	2.7
2002.....	*	*	0.3	0.5	0.4	1.0	1.4	1.6	2.4	2.2
2001.....	*	*	*	0.4	0.8	0.8	1.3	2.2	2.0	2.7
2000.....	*	*	0.3	0.6	0.8	0.9	1.4	2.0	2.2	2.6
1999.....	*	*	0.4	*	0.4	0.9	1.4	1.6	1.8	2.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of colon, rectum and anus (C18-C21)										
All races 3/, both sexes										
2006.....	4.0	7.6	13.5	21.8	34.8	52.0	73.9	104.4	146.8	207.5
2005.....	4.0	7.4	13.4	21.4	34.2	52.7	75.8	105.7	149.2	216.9
2004.....	3.9	7.4	13.5	22.4	35.9	53.2	78.3	111.3	152.3	225.6
2003.....	3.8	7.9	14.3	23.0	38.6	59.3	83.3	117.0	159.0	239.7
2002.....	4.3	8.2	15.0	24.6	40.7	61.6	86.2	118.9	163.2	241.4
2001.....	4.1	8.4	15.2	25.4	39.3	62.9	88.8	121.1	165.8	255.7
2000.....	4.0	8.2	14.7	25.2	42.0	62.0	92.4	126.9	174.8	270.4
1999.....	4.0	8.3	14.4	25.3	43.6	63.9	91.9	129.0	173.2	271.7
All races 3/, male										
2006.....	4.2	8.5	15.1	25.6	43.1	64.9	91.9	127.3	181.3	234.6
2005.....	4.2	8.2	15.2	25.9	42.4	67.7	94.4	131.8	180.2	245.9
2004.....	4.2	8.3	16.1	27.0	43.7	66.4	98.0	139.0	183.1	254.7
2003.....	4.2	9.0	16.6	27.6	47.1	75.0	103.4	143.0	196.5	273.3
2002.....	4.9	9.1	17.5	30.1	51.6	77.8	106.8	147.7	197.4	275.1
2001.....	4.3	9.5	17.1	30.8	48.0	78.9	110.4	149.4	201.3	298.3
2000.....	4.4	9.0	17.0	30.6	51.0	77.4	115.7	159.4	213.0	315.6
1999.....	4.1	9.1	16.6	30.6	52.9	78.0	114.3	161.3	216.2	324.0
All races 3/, female										
2006.....	3.8	6.7	11.9	18.2	27.2	40.8	59.2	87.6	125.3	194.9
2005.....	3.7	6.6	11.6	17.1	26.7	39.7	60.8	86.8	130.0	203.6
2004.....	3.6	6.5	11.1	18.0	28.7	41.8	62.4	91.3	133.3	212.5
2003.....	3.4	6.9	12.1	18.6	30.9	45.6	67.3	98.5	136.1	224.8
2002.....	3.6	7.4	12.6	19.5	30.8	47.6	69.7	98.6	142.3	226.8
2001.....	3.8	7.4	13.4	20.4	31.5	49.2	71.5	101.2	144.6	237.8
2000.....	3.5	7.5	12.4	20.2	33.9	48.8	74.1	104.3	152.2	251.9
1999.....	3.9	7.5	12.3	20.4	35.1	51.8	74.3	106.6	148.0	250.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of colon, rectum and anus (C18-C21)										
White, both sexes										
2006 4/.....	3.6	7.1	12.6	20.7	32.9	50.0	72.5	101.8	144.5	207.4
2005 4/.....	3.7	6.8	12.2	19.9	32.1	50.3	74.4	103.1	147.3	215.7
2004 4/.....	3.6	6.8	12.6	21.3	34.2	51.2	76.2	109.5	150.9	225.1
2003 4/.....	3.4	7.3	13.0	21.5	36.9	58.0	80.7	115.1	158.2	238.5
2002.....	3.9	7.5	14.2	23.3	38.9	59.9	84.6	116.9	160.3	240.7
2001.....	3.8	7.7	14.3	23.9	37.9	60.8	86.8	118.0	163.7	255.3
2000.....	3.6	7.5	13.8	23.7	40.5	60.3	90.6	125.0	172.1	269.2
1999.....	3.7	7.7	13.5	23.8	42.0	62.0	90.0	126.9	170.8	270.9
White, male										
2006 4/.....	3.8	7.9	14.0	24.6	40.8	62.7	89.6	124.4	177.1	233.0
2005 4/.....	4.1	7.6	13.9	24.1	40.0	64.6	92.9	128.0	179.5	245.3
2004 4/.....	3.9	7.8	15.1	26.0	41.9	64.0	95.3	135.9	181.7	255.4
2003 4/.....	3.8	8.3	15.2	26.2	45.0	73.6	100.8	140.7	196.0	272.9
2002.....	4.5	8.4	16.5	28.7	49.4	76.0	105.4	144.9	193.3	273.6
2001.....	4.0	8.7	16.4	29.3	46.7	76.7	108.7	146.7	198.9	297.3
2000.....	3.9	8.3	16.1	28.6	49.1	75.3	113.5	157.6	208.6	313.3
1999.....	3.9	8.5	15.8	28.8	51.1	76.1	113.3	158.1	214.4	324.5
White, female										
2006 4/.....	3.5	6.3	11.2	17.0	25.4	38.7	58.4	85.0	123.9	195.4
2005 4/.....	3.4	6.0	10.5	15.9	24.8	37.6	59.2	84.7	127.1	202.0
2004 4/.....	3.4	5.9	10.3	16.8	27.1	39.8	60.5	90.3	131.8	211.4
2003 4/.....	2.9	6.3	10.8	17.0	29.5	44.1	64.2	96.6	134.8	223.2
2002.....	3.3	6.5	12.0	18.2	29.3	45.7	67.6	96.9	140.1	226.4
2001.....	3.6	6.7	12.2	18.8	29.8	46.9	69.1	97.7	142.5	237.7
2000.....	3.3	6.6	11.5	19.1	32.6	47.2	72.3	102.1	150.4	251.3
1999.....	3.5	6.9	11.2	19.2	33.6	49.7	71.5	104.9	145.1	249.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of colon, rectum and anus (C18-C21)										
Black, both sexes										
2006 4/.....	6.9	12.6	22.4	33.8	57.3	79.1	101.6	147.8	197.8	240.8
2005 4/.....	5.7	12.6	23.8	36.5	57.5	82.9	101.3	148.3	197.6	261.7
2004 4/.....	5.9	12.2	21.9	36.1	56.7	79.9	113.2	148.6	189.1	260.8
2003 4/.....	6.6	13.2	25.5	39.4	59.7	81.2	124.6	154.7	197.7	281.5
2002.....	7.2	14.5	23.5	40.0	61.9	86.8	114.9	160.5	218.7	274.7
2001.....	6.5	14.5	24.6	43.1	58.3	90.9	120.3	172.6	211.0	286.5
2000.....	6.6	15.2	24.8	39.9	62.7	86.9	126.1	167.4	226.2	310.0
1999.....	6.6	13.3	23.4	42.0	65.7	90.5	125.4	172.3	226.8	306.1
Black, male										
2006 4/.....	7.3	14.9	26.4	39.5	70.5	99.8	137.8	188.4	266.3	309.2
2005 4/.....	5.7	13.7	27.3	46.0	74.2	106.7	124.9	197.1	231.8	295.6
2004 4/.....	6.6	13.2	26.3	42.0	69.7	100.5	147.0	198.8	239.6	278.2
2003 4/.....	7.1	15.2	30.3	46.1	76.4	102.7	156.7	194.9	242.2	317.0
2002.....	8.4	15.7	27.9	47.5	79.4	109.2	138.2	209.0	285.1	328.1
2001.....	7.6	16.4	26.4	51.3	67.7	114.1	146.8	208.2	263.2	347.2
2000.....	8.0	16.4	29.2	51.4	76.4	109.3	160.1	208.9	298.4	377.1
1999.....	6.3	15.0	26.9	54.1	80.4	110.6	147.6	228.0	274.7	350.4
Black, female										
2006 4/.....	6.4	10.7	19.1	29.1	47.0	63.9	76.7	123.5	161.7	212.8
2005 4/.....	5.6	11.6	20.9	28.7	44.5	65.2	85.1	119.2	179.5	248.0
2004 4/.....	5.3	11.3	18.2	31.3	46.6	64.6	90.3	118.9	162.5	254.0
2003 4/.....	6.1	11.4	21.4	33.8	46.8	65.2	103.0	130.8	174.5	267.8
2002.....	6.1	13.5	19.7	33.9	48.2	70.1	99.2	131.5	184.2	254.2
2001.....	5.6	12.8	23.1	36.3	50.9	73.8	102.7	151.4	184.4	264.0
2000.....	5.3	14.2	21.1	30.5	52.0	70.5	103.4	142.4	189.8	285.0
1999.....	6.7	11.9	20.5	32.1	54.2	75.6	110.7	138.7	203.1	289.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Colon (C18)										
All races 3/, both sexes										
2006.....	2.9	5.6	10.4	17.0	27.7	42.3	61.1	87.5	124.9	179.1
2005.....	3.0	5.5	10.2	16.6	27.2	42.8	62.6	89.0	127.6	188.1
2004.....	2.8	5.5	10.5	17.6	28.4	43.7	65.7	94.0	130.9	195.3
2003.....	2.8	6.1	11.2	17.9	31.1	49.4	70.1	99.8	137.6	208.4
2002.....	3.3	6.3	11.9	19.8	32.5	50.5	72.7	101.5	142.3	211.3
2001.....	3.1	6.4	11.8	20.4	32.4	52.4	75.6	103.9	143.5	221.6
2000.....	3.1	6.3	11.8	20.1	34.4	51.4	78.2	108.3	153.7	236.6
1999.....	3.1	6.4	11.6	20.4	35.9	53.3	77.7	110.7	151.6	238.1
All races 3/, male										
2006.....	3.0	6.3	11.3	19.6	33.9	52.1	75.2	105.0	152.5	200.5
2005.....	3.1	5.8	11.7	20.1	33.0	53.8	76.7	109.8	152.5	212.2
2004.....	2.9	6.2	12.4	20.5	34.2	53.7	80.4	115.7	155.2	219.9
2003.....	3.0	6.8	12.8	21.2	37.1	62.2	85.5	121.3	168.6	236.1
2002.....	3.7	6.8	13.5	23.8	40.2	63.2	89.8	124.7	169.8	240.2
2001.....	3.2	7.1	12.9	24.5	38.5	64.5	92.6	126.3	173.0	256.7
2000.....	3.4	6.7	13.6	24.0	40.9	62.7	96.6	135.2	185.0	275.8
1999.....	3.2	7.0	13.2	24.2	43.3	64.3	96.0	137.4	187.2	282.2
All races 3/, female										
2006.....	2.8	5.0	9.5	14.5	22.0	33.7	49.7	74.7	107.6	169.1
2005.....	2.9	5.1	8.9	13.4	21.9	33.2	51.2	74.0	112.2	177.0
2004.....	2.7	4.8	8.7	14.9	23.2	34.9	53.8	78.4	116.0	184.2
2003.....	2.5	5.4	9.6	14.8	25.7	38.3	57.6	84.5	118.7	196.1
2002.....	2.9	5.8	10.3	16.1	25.4	39.5	59.1	85.1	125.6	198.7
2001.....	3.0	5.7	10.7	16.7	26.9	42.0	62.1	88.3	125.8	206.9
2000.....	2.8	5.9	10.1	16.6	28.5	41.7	63.7	89.6	135.2	220.6
1999.....	3.0	5.9	10.1	16.8	29.3	43.8	63.4	92.1	130.8	220.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Colon (C18)										
White, both sexes										
2006 4/.....	2.7	5.1	9.5	15.9	25.8	40.2	59.7	85.1	123.0	179.0
2005 4/.....	2.8	4.9	9.2	15.3	25.2	40.7	61.1	86.5	125.7	186.8
2004 4/.....	2.6	5.1	9.7	16.5	26.8	41.6	63.3	92.0	129.3	194.8
2003 4/.....	2.4	5.5	10.0	16.6	29.5	48.1	67.5	98.1	136.6	206.9
2002.....	3.0	5.5	11.1	18.6	30.8	48.6	71.0	99.4	140.0	210.5
2001.....	2.8	5.8	10.9	19.1	30.9	50.1	73.7	100.7	141.6	221.2
2000.....	2.8	5.6	11.0	18.8	33.0	49.9	76.4	106.4	150.8	235.0
1999.....	2.8	5.9	10.8	19.0	34.5	51.6	75.8	108.5	149.3	237.1
White, male										
2006 4/.....	2.8	5.8	10.3	18.5	31.8	49.8	72.9	102.1	149.2	198.3
2005 4/.....	3.0	5.5	10.5	18.4	30.8	51.3	75.1	106.2	151.5	211.8
2004 4/.....	2.8	5.8	11.4	19.4	32.4	51.5	77.4	112.3	153.5	220.7
2003 4/.....	2.8	6.3	11.7	20.0	35.0	60.7	83.1	119.2	167.3	236.2
2002.....	3.4	6.1	12.6	22.5	38.1	61.2	88.2	121.7	166.9	238.7
2001.....	2.9	6.4	12.1	23.2	37.1	62.2	91.1	123.1	170.6	255.5
2000.....	3.0	6.1	12.7	22.3	39.0	61.2	94.6	133.3	180.4	272.5
1999.....	3.0	6.5	12.6	22.6	41.6	62.7	95.0	134.3	185.6	282.9
White, female										
2006 4/.....	2.5	4.5	8.8	13.3	20.2	31.7	48.6	72.4	106.5	169.9
2005 4/.....	2.6	4.4	7.8	12.2	20.0	31.2	49.5	72.0	109.6	175.3
2004 4/.....	2.4	4.3	7.9	13.7	21.7	32.8	51.7	77.2	114.2	183.1
2003 4/.....	2.1	4.8	8.5	13.4	24.5	36.9	54.8	82.8	117.7	193.9
2002.....	2.6	4.9	9.7	14.8	24.0	37.6	57.0	83.4	123.5	198.2
2001.....	2.7	5.1	9.6	15.1	25.1	39.5	59.7	84.8	124.1	206.9
2000.....	2.6	5.1	9.3	15.5	27.3	40.1	61.8	87.5	133.2	219.8
1999.....	2.6	5.3	9.2	15.5	27.9	41.8	60.6	90.5	127.8	218.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Colon (C18)										
Black, both sexes										
2006 4/.....	5.0	10.1	18.3	28.6	48.3	68.2	87.9	127.7	166.8	206.5
2005 4/.....	4.5	9.8	19.7	30.7	48.7	70.5	87.2	128.4	176.0	230.4
2004 4/.....	4.2	9.2	18.0	30.6	47.7	68.9	100.8	131.9	168.3	227.3
2003 4/.....	5.2	11.0	20.8	32.2	50.8	70.1	108.9	135.1	175.7	249.4
2002.....	5.7	12.1	19.7	33.6	52.1	74.8	100.3	142.8	190.2	244.1
2001.....	5.5	11.8	20.5	36.2	51.3	80.1	105.9	154.9	182.6	249.9
2000.....	5.4	12.5	21.0	33.5	53.6	73.0	109.7	148.6	206.7	279.7
1999.....	5.5	11.4	19.6	35.7	56.4	78.1	109.4	151.6	204.0	273.9
Black, male										
2006 4/.....	4.9	11.4	21.5	32.9	58.3	85.5	117.2	162.4	221.8	275.2
2005 4/.....	4.1	9.4	22.3	38.8	61.3	89.1	106.9	169.9	207.9	251.9
2004 4/.....	3.9	9.3	22.1	34.8	57.9	84.6	128.2	173.8	208.8	243.4
2003 4/.....	5.3	11.8	24.3	36.7	64.4	89.1	133.7	168.0	220.7	270.0
2002.....	6.3	12.1	23.0	39.3	66.3	94.2	119.2	186.9	238.4	292.7
2001.....	5.9	13.5	22.0	41.6	57.2	97.8	124.9	184.9	231.1	307.5
2000.....	6.6	12.9	24.3	41.5	64.4	86.9	136.1	184.3	271.4	346.0
1999.....	5.1	12.3	21.6	44.4	67.7	94.8	126.5	200.9	238.9	304.7
Black, female										
2006 4/.....	5.0	9.0	15.6	25.1	40.5	55.4	67.8	106.9	137.8	178.4
2005 4/.....	4.9	10.2	17.4	24.1	39.0	56.8	73.8	103.7	159.1	221.7
2004 4/.....	4.4	9.1	14.6	27.0	39.8	57.2	82.2	107.0	147.0	220.9
2003 4/.....	5.2	10.2	17.8	28.4	40.2	55.9	92.2	115.6	152.0	241.4
2002.....	5.1	12.0	16.9	29.0	41.0	60.3	87.6	116.4	165.1	225.5
2001.....	5.1	10.3	19.2	31.8	46.7	67.0	93.3	137.0	157.8	228.6
2000.....	4.4	12.3	18.1	27.0	45.1	62.8	92.1	127.0	174.1	255.0
1999.....	5.9	10.7	18.0	28.5	47.6	65.8	98.1	121.9	186.7	262.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rectosigmoid junction and rectum (C19-C20)										
All races 3/, both sexes										
2006.....	1.0	1.8	2.9	4.4	6.6	9.3	12.0	16.0	20.9	27.0
2005.....	0.8	1.8	2.8	4.3	6.6	9.3	12.6	15.9	20.5	27.6
2004.....	1.0	1.6	2.7	4.4	7.0	9.0	11.9	16.3	20.3	29.1
2003.....	0.9	1.6	2.8	4.7	7.0	9.3	12.6	16.3	20.4	29.9
2002.....	0.9	1.8	2.8	4.5	7.7	10.6	12.9	16.7	19.9	29.0
2001.....	0.8	1.8	3.2	4.7	6.6	10.0	12.5	16.3	21.4	32.6
2000.....	0.7	1.7	2.6	4.7	7.2	10.1	13.6	17.7	20.3	32.6
1999.....	0.8	1.6	2.5	4.6	7.3	10.1	13.5	17.6	20.6	32.1
All races 3/, male										
2006.....	1.1	2.0	3.6	5.7	8.9	12.4	15.9	21.4	28.3	32.7
2005.....	0.9	2.2	3.4	5.4	9.1	13.4	17.0	21.5	26.7	32.8
2004.....	1.1	2.0	3.4	6.1	9.1	12.0	17.0	22.5	27.5	33.5
2003.....	1.0	2.0	3.6	6.1	9.7	12.2	17.4	21.2	27.5	36.4
2002.....	1.1	2.2	3.6	6.1	10.8	14.2	16.6	22.4	26.9	33.6
2001.....	1.0	2.2	4.0	6.0	9.2	13.9	17.2	22.3	27.7	40.7
2000.....	0.9	2.2	3.2	6.3	9.7	14.2	18.7	23.5	27.0	38.9
1999.....	0.9	1.9	3.2	6.2	9.4	13.3	17.8	23.3	28.2	40.7
All races 3/, female										
2006.....	0.9	1.5	2.1	3.2	4.6	6.5	8.9	12.0	16.3	24.3
2005.....	0.7	1.3	2.3	3.2	4.3	5.7	8.9	11.8	16.6	25.2
2004.....	0.9	1.3	2.1	2.8	5.0	6.3	7.8	11.9	16.0	27.2
2003.....	0.7	1.2	2.1	3.3	4.6	6.8	8.8	12.8	16.0	27.0
2002.....	0.6	1.4	2.0	3.0	4.8	7.5	9.9	12.6	15.6	27.0
2001.....	0.7	1.5	2.4	3.3	4.2	6.6	8.8	12.0	17.6	29.2
2000.....	0.6	1.3	2.0	3.3	4.9	6.6	9.6	13.7	16.4	30.1
1999.....	0.8	1.4	1.9	3.1	5.3	7.4	10.1	13.6	16.2	28.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rectosigmoid junction and rectum (C19-C20)										
White, both sexes										
2006 4/.....	0.9	1.7	2.8	4.4	6.5	9.2	12.1	15.9	20.4	27.0
2005 4/.....	0.8	1.7	2.7	4.2	6.5	9.0	12.7	15.7	20.4	27.6
2004 4/.....	0.9	1.5	2.7	4.4	6.9	9.0	12.2	16.6	20.5	29.1
2003 4/.....	0.9	1.6	2.7	4.5	6.9	9.4	12.5	16.2	20.5	30.1
2002.....	0.8	1.7	2.8	4.4	7.6	10.7	13.0	16.8	19.3	29.0
2001.....	0.8	1.8	3.1	4.5	6.6	10.1	12.4	16.4	21.1	32.6
2000.....	0.7	1.7	2.5	4.6	7.1	9.8	13.5	17.8	20.5	33.0
1999.....	0.8	1.6	2.4	4.5	7.1	10.0	13.5	17.5	20.7	32.2
White, male										
2006 4/.....	0.9	1.9	3.5	5.7	8.6	12.5	15.8	21.4	27.3	33.3
2005 4/.....	0.9	1.9	3.2	5.3	8.8	12.8	17.2	21.2	26.9	32.6
2004 4/.....	0.9	1.8	3.4	6.1	9.0	11.9	17.4	22.7	27.6	33.6
2003 4/.....	1.0	1.9	3.4	5.9	9.6	12.3	17.2	20.9	28.3	36.0
2002.....	1.0	2.0	3.6	5.9	10.8	14.3	16.7	22.7	25.6	33.7
2001.....	0.9	2.1	4.0	5.8	9.2	14.0	16.9	22.8	27.8	41.0
2000.....	0.9	2.1	3.1	5.9	9.7	13.6	18.4	23.7	27.2	39.8
1999.....	0.8	1.8	3.0	5.9	9.3	13.0	17.8	23.2	28.0	40.6
White, female										
2006 4/.....	0.8	1.5	2.0	3.2	4.5	6.3	9.0	11.8	16.0	24.0
2005 4/.....	0.7	1.4	2.2	3.1	4.3	5.7	8.9	11.7	16.3	25.2
2004 4/.....	0.9	1.2	2.0	2.7	4.9	6.4	8.0	12.2	16.1	27.1
2003 4/.....	0.7	1.2	1.9	3.1	4.5	6.7	8.6	12.7	15.8	27.5
2002.....	0.6	1.4	1.9	2.9	4.7	7.4	9.9	12.5	15.4	27.0
2001.....	0.7	1.4	2.2	3.3	4.2	6.8	8.7	11.9	17.1	29.1
2000.....	0.6	1.3	1.9	3.3	4.6	6.5	9.7	13.7	16.5	30.3
1999.....	0.8	1.4	1.8	3.2	5.2	7.4	10.1	13.5	16.4	28.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rectosigmoid junction and rectum (C19-C20)										
Black, both sexes										
2006 4/.....	1.6	2.3	3.9	4.6	8.4	10.9	13.0	19.4	30.1	32.2
2005 4/.....	0.9	2.5	3.9	5.4	8.4	11.4	13.5	19.5	20.6	30.2
2004 4/.....	1.6	2.7	3.5	5.3	8.6	10.4	11.9	15.7	20.8	31.9
2003 4/.....	1.2	2.1	4.3	6.8	8.5	10.5	14.7	18.2	21.3	31.5
2002.....	1.3	2.3	3.5	6.0	9.0	11.6	13.9	17.2	27.7	29.7
2001.....	0.9	2.6	3.7	6.5	6.9	10.3	14.1	17.2	26.8	35.6
2000.....	1.0	2.5	3.4	6.1	8.8	13.6	15.6	17.6	19.2	28.8
1999.....	0.9	1.7	3.3	6.1	9.1	11.9	14.7	20.4	21.4	31.9
Black, male										
2006 4/.....	2.0	3.1	5.0	6.3	12.0	14.1	19.4	26.0	44.5	32.2
2005 4/.....	*	3.9	4.7	7.0	12.7	17.3	17.4	27.3	23.2	42.8
2004 4/.....	2.4	3.6	3.8	7.0	11.2	15.0	18.1	24.6	30.8	31.8
2003 4/.....	1.5	3.2	5.5	9.1	12.0	12.3	22.7	25.6	21.4	45.9
2002.....	1.7	3.5	4.4	7.9	12.5	14.5	18.7	21.1	45.9	34.3
2001.....	1.6	2.8	4.2	9.2	10.4	15.8	21.5	22.8	30.5	38.6
2000.....	*	3.5	4.2	9.6	11.4	21.9	23.4	22.7	27.0	30.0
1999.....	*	2.5	4.7	9.5	12.5	15.6	20.4	27.1	34.9	45.7
Black, female										
2006 4/.....	1.3	1.5	3.0	3.2	5.5	8.5	8.7	15.5	22.4	32.2
2005 4/.....	*	*	3.1	4.1	5.0	7.1	10.8	14.9	19.3	25.1
2004 4/.....	*	1.9	3.3	3.9	6.6	7.0	7.7	10.5	15.5	31.9
2003 4/.....	*	*	3.3	4.8	5.9	9.1	9.3	13.8	21.3	25.9
2002.....	*	*	2.7	4.5	6.3	9.4	10.8	14.8	18.3	27.9
2001.....	*	2.3	3.3	4.2	4.2	6.2	9.2	13.9	24.9	34.5
2000.....	*	1.7	2.8	3.2	6.7	7.5	10.4	14.5	15.3	28.3
1999.....	*	*	2.1	3.3	6.5	9.2	11.0	16.3	14.7	26.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anus and anus canal (C21)										
All races 3/, both sexes										
2006.....	0.1	0.2	0.3	0.4	0.5	0.5	0.7	0.9	1.1	1.4
2005.....	0.1	0.2	0.3	0.4	0.4	0.6	0.7	0.8	1.1	1.2
2004.....	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.9	1.0	1.2
2003.....	0.1	0.2	0.3	0.4	0.5	0.5	0.7	0.9	1.0	1.4
2002.....	0.1	0.2	0.3	0.3	0.6	0.5	0.6	0.8	1.0	1.2
2001.....	0.1	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.0	1.4
2000.....	0.1	0.2	0.3	0.3	0.5	0.5	0.7	0.9	0.8	1.2
1999.....	0.1	0.2	0.2	0.3	0.4	0.5	0.7	0.7	0.9	1.4
All races 3/, male										
2006.....	*	0.2	0.2	0.4	0.4	0.4	0.8	0.8	*	1.4
2005.....	*	0.2	0.2	0.4	0.3	0.5	0.6	*	1.0	*
2004.....	*	*	0.3	0.4	0.5	0.6	0.6	0.8	*	*
2003.....	*	*	*	0.3	*	0.6	*	*	*	*
2002.....	*	0.2	0.3	*	0.5	0.5	*	*	*	*
2001.....	*	*	0.2	0.3	*	0.5	0.6	0.8	*	*
2000.....	*	*	0.3	0.4	0.4	0.5	0.5	0.7	*	*
1999.....	*	0.2	*	*	*	*	*	*	*	*
All races 3/, female										
2006.....	*	0.2	0.3	0.5	0.7	0.6	0.7	0.9	1.4	1.5
2005.....	*	0.2	0.5	0.5	0.5	0.7	0.7	1.0	1.2	1.4
2004.....	*	0.3	0.4	0.3	0.5	0.6	0.8	1.0	1.4	1.1
2003.....	*	0.2	0.4	0.5	0.6	0.5	0.8	1.1	1.3	1.6
2002.....	*	0.2	0.3	0.4	0.6	0.6	0.7	0.9	1.1	1.2
2001.....	*	*	0.4	0.4	0.4	0.6	0.7	0.9	1.2	1.7
2000.....	*	0.3	0.3	0.3	0.5	0.5	0.8	1.0	0.7	1.3
1999.....	*	*	0.3	0.4	0.4	0.5	0.8	0.9	1.0	1.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anus and anus canal (C21)										
White, both sexes										
2006 4/.....	0.1	0.2	0.3	0.4	0.6	0.6	0.8	0.9	1.1	1.4
2005 4/.....	*	0.2	0.4	0.5	0.4	0.6	0.7	0.9	1.1	1.3
2004 4/.....	0.1	0.2	0.3	0.4	0.5	0.6	0.7	0.9	1.1	1.1
2003 4/.....	0.1	0.2	0.3	0.4	0.5	0.5	0.6	0.9	1.0	1.5
2002.....	0.1	0.2	0.4	0.4	0.6	0.6	0.6	0.8	1.0	1.2
2001.....	0.1	0.2	0.3	0.4	0.4	0.5	0.7	0.9	1.0	1.5
2000.....	0.1	0.2	0.3	0.3	0.5	0.5	0.7	0.8	0.8	1.1
1999.....	*	0.2	0.2	0.4	0.4	0.5	0.6	0.8	0.9	1.5
White, male										
2006 4/.....	*	*	*	0.4	0.4	*	0.8	0.9	*	1.4
2005 4/.....	*	*	*	0.4	*	0.5	0.6	*	1.0	*
2004 4/.....	*	*	0.3	0.5	0.5	0.5	*	0.8	*	*
2003 4/.....	*	*	*	0.3	*	0.5	*	*	*	*
2002.....	*	0.2	0.3	*	0.5	*	*	*	*	*
2001.....	*	*	*	*	*	*	0.7	0.8	*	*
2000.....	*	*	*	0.4	*	0.5	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, female										
2006 4/.....	*	0.2	0.3	0.4	0.7	0.7	0.8	0.9	1.4	1.5
2005 4/.....	*	0.2	0.5	0.5	0.5	0.7	0.7	1.1	1.2	1.4
2004 4/.....	*	0.3	0.4	0.4	0.5	0.6	0.8	0.9	1.5	1.2
2003 4/.....	*	0.3	0.4	0.5	0.5	0.5	0.8	1.1	1.3	1.8
2002.....	*	0.2	0.4	0.5	0.6	0.6	0.7	1.0	1.2	1.2
2001.....	*	*	0.3	0.4	0.5	0.6	0.7	1.0	1.2	1.7
2000.....	*	0.3	0.3	*	0.6	0.6	0.8	1.0	0.7	1.2
1999.....	*	*	*	0.5	0.5	0.5	0.8	0.9	0.9	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of liver and intrahepatic bile ducts (C22)										
All races 3/, both sexes										
2006.....	1.1	3.2	8.2	11.5	13.1	17.2	23.9	30.2	34.1	33.5
2005.....	1.2	3.7	8.5	10.2	12.9	16.6	24.3	30.2	34.4	32.5
2004.....	1.2	3.9	7.6	9.4	12.5	16.6	23.7	28.8	34.6	32.4
2003.....	1.2	3.7	7.4	8.5	11.7	16.5	23.5	29.6	32.1	33.4
2002.....	1.3	3.4	6.6	8.1	11.6	16.9	22.6	27.6	31.6	35.0
2001.....	1.2	3.8	5.8	7.5	11.6	16.3	21.5	27.0	32.0	34.1
2000.....	1.4	3.7	5.4	7.7	11.3	16.0	21.6	27.1	31.3	32.7
1999.....	1.4	3.4	4.9	7.1	10.9	16.3	21.8	26.1	30.9	30.6
All races 3/, male										
2006.....	1.6	5.0	13.3	18.5	19.6	25.7	33.7	42.2	48.0	43.8
2005.....	1.6	5.9	13.8	16.7	19.5	24.2	34.8	42.8	48.2	44.8
2004.....	1.8	6.2	12.8	15.1	18.6	23.6	34.0	38.8	48.4	41.5
2003.....	1.8	6.0	12.2	13.0	17.3	23.3	33.4	43.7	46.1	47.5
2002.....	2.0	5.7	11.1	12.4	17.4	24.8	32.8	39.5	44.8	46.2
2001.....	1.8	6.0	9.7	11.4	17.2	23.5	30.6	39.2	43.9	50.3
2000.....	2.0	6.0	8.5	11.7	16.5	23.6	30.9	39.3	46.7	44.7
1999.....	2.1	5.5	7.6	10.6	15.6	23.7	31.9	36.6	42.9	40.1
All races 3/, female										
2006.....	0.7	1.5	3.3	4.9	7.1	9.8	15.9	21.4	25.4	28.7
2005.....	0.8	1.5	3.4	4.1	6.8	9.9	15.8	21.0	25.9	26.8
2004.....	0.6	1.8	2.6	4.1	6.9	10.6	15.4	21.7	26.1	28.3
2003.....	0.7	1.6	2.8	4.3	6.6	10.7	15.6	19.6	23.6	27.2
2002.....	0.6	1.2	2.4	4.0	6.4	10.1	14.4	19.2	23.6	30.2
2001.....	0.5	1.7	2.1	3.9	6.5	10.2	14.2	18.4	24.8	27.2
2000.....	0.8	1.4	2.4	4.0	6.5	9.5	14.2	18.6	22.2	27.8
1999.....	0.8	1.3	2.3	3.8	6.6	10.1	13.8	18.7	23.9	26.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of liver and intrahepatic bile ducts (C22)										
White, both sexes										
2006 4/.....	0.9	2.7	7.2	9.9	11.7	15.8	22.7	28.9	32.8	33.1
2005 4/.....	1.0	3.3	7.4	8.9	11.3	15.3	22.5	29.1	32.8	31.7
2004 4/.....	1.0	3.3	6.6	8.0	11.2	15.1	21.9	27.3	33.7	31.2
2003 4/.....	1.0	3.1	6.3	7.5	10.3	15.2	21.7	28.3	31.0	32.1
2002.....	1.1	3.0	5.5	6.8	10.6	15.8	21.1	26.7	30.7	33.8
2001.....	0.8	3.2	4.9	6.4	10.5	15.1	20.2	26.1	30.4	33.3
2000.....	1.1	3.1	4.4	6.6	10.0	14.6	20.2	25.8	30.7	32.0
1999.....	1.1	2.7	4.2	5.9	9.7	15.2	20.5	25.2	30.2	29.9
White, male										
2006 4/.....	1.2	4.1	11.6	15.6	17.1	23.1	32.0	40.6	46.9	43.7
2005 4/.....	1.3	5.2	11.6	14.2	17.1	22.0	32.2	41.5	45.8	45.0
2004 4/.....	1.5	5.0	11.1	12.4	16.5	21.5	31.1	37.4	47.2	39.9
2003 4/.....	1.4	4.9	10.2	11.4	15.2	21.4	31.4	41.8	44.8	45.6
2002.....	1.6	4.9	9.0	10.4	15.6	23.0	30.4	38.0	42.9	45.1
2001.....	1.3	5.0	8.1	9.5	15.7	22.1	29.1	37.7	42.6	49.7
2000.....	1.5	4.8	6.8	9.6	14.4	21.4	28.9	38.1	45.8	43.1
1999.....	1.7	4.3	6.3	8.8	13.9	21.9	30.2	35.3	42.4	40.6
White, female										
2006 4/.....	0.7	1.3	2.9	4.4	6.7	9.2	15.0	20.2	23.9	28.1
2005 4/.....	0.7	1.3	3.2	3.7	6.0	9.3	14.5	19.9	24.7	25.6
2004 4/.....	0.5	1.5	2.2	3.7	6.3	9.5	14.3	19.8	25.3	27.3
2003 4/.....	0.6	1.3	2.5	3.8	5.7	9.7	13.9	18.6	22.5	26.2
2002.....	0.5	1.1	2.0	3.4	5.9	9.3	13.5	18.6	23.2	29.0
2001.....	0.4	1.4	1.9	3.4	5.7	8.9	13.0	17.8	23.1	26.5
2000.....	0.6	1.3	2.0	3.6	5.9	8.5	13.2	17.1	21.7	27.4
1999.....	0.6	1.2	2.1	3.1	5.8	9.3	12.8	18.1	23.0	25.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of liver and intrahepatic bile ducts (C22)										
Black, both sexes										
2006 4/.....	1.7	5.4	14.3	22.7	20.4	22.3	26.7	30.8	36.9	31.1
2005 4/.....	1.8	5.6	14.5	18.9	22.1	21.4	29.4	29.1	39.3	29.1
2004 4/.....	1.8	7.1	12.7	17.9	16.6	23.2	28.5	33.7	35.4	36.1
2003 4/.....	1.9	6.2	14.2	14.7	18.4	18.9	27.9	30.3	32.7	41.7
2002.....	2.1	5.8	12.3	15.5	14.0	19.5	27.3	30.2	31.9	38.0
2001.....	2.3	7.1	10.7	15.2	14.3	20.0	24.0	26.9	40.9	35.0
2000.....	2.7	7.0	10.6	12.7	15.8	20.9	26.9	28.8	28.5	31.6
1999.....	2.2	6.6	8.1	12.0	14.7	21.1	28.0	26.6	33.8	34.4
Black, male										
2006 4/.....	2.9	8.0	24.3	42.3	35.2	37.0	42.4	45.2	45.9	34.9
2005 4/.....	2.5	8.9	26.8	35.0	36.7	35.9	44.6	45.1	59.7	31.6
2004 4/.....	2.9	11.7	22.9	31.9	27.1	36.8	45.0	39.6	50.9	46.7
2003 4/.....	3.0	10.3	25.5	24.0	28.5	26.8	42.4	43.8	45.0	63.6
2002.....	3.3	9.8	21.7	25.3	22.8	32.2	45.1	44.6	51.2	41.8
2001.....	3.6	11.6	18.6	25.3	21.4	28.0	32.7	39.5	49.0	42.0
2000.....	3.8	12.6	17.7	22.1	25.3	34.1	41.5	36.4	45.5	48.4
1999.....	3.4	11.8	14.2	18.6	21.6	32.2	42.1	40.4	42.7	30.5
Black, female										
2006 4/.....	*	3.0	5.8	6.5	9.0	11.4	15.8	22.2	32.2	29.6
2005 4/.....	*	2.6	4.2	5.6	10.8	10.6	19.0	19.5	28.6	28.1
2004 4/.....	*	3.0	4.1	6.4	8.4	13.1	17.4	30.2	27.2	31.9
2003 4/.....	*	2.7	4.5	7.0	10.5	13.0	18.2	22.2	26.3	33.2
2002.....	*	2.3	4.3	7.5	7.1	10.2	15.5	21.7	21.9	36.6
2001.....	*	3.2	3.9	6.8	8.8	14.1	18.2	19.4	36.8	32.4
2000.....	1.8	2.2	4.7	5.0	8.3	11.1	17.3	24.2	20.0	25.3
1999.....	*	2.0	2.9	6.5	9.3	12.9	18.7	18.3	29.4	35.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Liver (C22.0,C22.2-C22.9)										
All races 3/, both sexes										
2006.....	0.9	2.6	7.1	9.7	10.1	13.3	17.9	22.9	24.9	23.6
2005.....	0.9	3.1	7.3	8.6	10.1	12.7	18.9	22.0	25.5	23.4
2004.....	0.9	3.3	6.6	7.8	9.4	12.7	18.2	21.3	25.1	23.2
2003.....	1.0	3.2	6.3	6.9	8.9	12.8	18.1	22.6	23.6	24.3
2002.....	1.0	3.0	5.7	6.4	9.1	13.5	18.1	21.2	23.6	24.7
2001.....	0.9	3.4	5.0	6.2	9.2	12.8	16.7	20.6	23.8	24.2
2000.....	1.2	3.2	4.5	6.4	8.9	12.7	17.1	20.7	24.1	23.4
1999.....	1.2	3.0	4.1	5.7	8.7	13.0	17.5	20.4	23.3	22.5
All races 3/, male										
2006.....	1.4	4.3	12.2	16.5	16.2	21.0	27.1	34.0	37.5	31.8
2005.....	1.4	5.2	12.5	14.9	16.3	19.7	28.2	33.1	37.7	33.5
2004.....	1.6	5.5	11.6	13.2	14.9	19.5	27.6	30.6	36.8	30.6
2003.....	1.6	5.4	11.1	11.1	14.1	19.0	26.6	35.1	35.9	35.2
2002.....	1.7	5.1	10.0	10.6	14.5	20.9	27.6	31.4	34.8	34.6
2001.....	1.5	5.6	8.7	10.0	14.5	19.5	24.9	31.3	33.6	38.5
2000.....	1.8	5.4	7.6	10.2	14.0	19.7	25.5	31.4	37.6	33.9
1999.....	1.9	5.1	6.9	8.9	13.4	19.9	26.7	29.9	34.2	31.2
All races 3/, female										
2006.....	0.4	1.0	2.2	3.3	4.5	6.5	10.4	14.8	17.0	19.8
2005.....	0.4	1.0	2.3	2.6	4.4	6.6	11.3	14.0	18.0	18.7
2004.....	0.3	1.2	1.8	2.7	4.4	6.8	10.6	14.6	17.9	19.8
2003.....	0.4	1.0	1.7	3.0	4.2	7.5	11.3	13.6	16.2	19.5
2002.....	0.4	0.9	1.6	2.5	4.2	7.1	10.5	14.0	16.8	20.4
2001.....	0.4	1.2	1.5	2.5	4.5	7.1	10.1	13.1	17.8	18.2
2000.....	0.6	1.1	1.5	2.7	4.4	6.6	10.4	13.3	16.1	19.1
1999.....	0.6	0.9	1.5	2.7	4.5	7.1	10.3	13.7	16.9	18.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Liver (C22.0,C22.2-C22.9)										
White, both sexes										
2006 4/.....	0.7	2.1	6.1	8.1	8.6	11.8	16.7	21.7	23.5	23.1
2005 4/.....	0.7	2.7	6.3	7.3	8.6	11.5	17.3	20.9	24.3	22.5
2004 4/.....	0.8	2.7	5.7	6.4	8.0	11.3	16.4	19.8	24.2	21.9
2003 4/.....	0.8	2.6	5.3	5.9	7.6	11.6	16.4	21.2	22.4	23.0
2002.....	0.8	2.6	4.6	5.3	8.1	12.4	16.6	20.4	22.8	23.7
2001.....	0.6	2.8	4.2	5.1	8.2	11.6	15.5	19.6	22.5	23.5
2000.....	0.8	2.6	3.5	5.3	7.7	11.2	15.6	19.5	23.4	22.5
1999.....	0.9	2.3	3.4	4.6	7.6	11.8	16.1	19.4	22.5	21.7
White, male										
2006 4/.....	1.0	3.5	10.5	13.6	13.6	18.4	25.3	32.4	36.3	31.6
2005 4/.....	1.1	4.5	10.5	12.5	13.9	17.7	25.9	32.1	35.9	33.2
2004 4/.....	1.2	4.4	10.0	10.6	12.8	17.5	24.8	29.2	35.2	28.8
2003 4/.....	1.2	4.4	9.2	9.4	12.2	17.3	24.8	33.1	34.4	33.1
2002.....	1.4	4.3	8.0	8.7	12.8	19.3	25.1	30.2	33.2	33.5
2001.....	1.0	4.6	7.1	8.2	13.2	18.1	23.6	29.9	32.4	37.8
2000.....	1.3	4.2	5.9	8.4	12.0	17.5	23.4	30.2	36.6	31.8
1999.....	1.5	3.8	5.6	7.2	11.6	18.2	24.9	28.7	33.8	31.4
White, female										
2006 4/.....	0.4	0.7	1.9	2.8	4.0	5.8	9.5	13.7	15.5	19.1
2005 4/.....	0.4	0.8	2.1	2.2	3.8	6.0	10.2	12.7	17.1	17.5
2004 4/.....	0.3	1.0	1.5	2.3	3.7	5.8	9.5	12.9	17.4	18.8
2003 4/.....	0.3	0.7	1.4	2.5	3.4	6.5	9.6	12.6	14.9	18.5
2002.....	0.3	0.8	1.3	2.0	3.8	6.2	9.7	13.3	16.5	19.4
2001.....	0.2	1.0	1.3	2.1	3.6	5.8	8.9	12.4	16.5	17.5
2000.....	0.4	0.9	1.2	2.3	3.8	5.7	9.3	11.9	15.5	18.7
1999.....	0.4	0.8	1.3	2.1	3.8	6.3	9.1	13.0	15.9	17.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Liver (C22.0,C22.2-C22.9)										
Black, both sexes										
2006 4/.....	1.6	4.8	13.2	21.0	18.0	18.4	21.2	23.3	30.1	24.9
2005 4/.....	1.4	5.0	13.3	17.4	19.1	17.6	23.4	21.2	31.6	23.8
2004 4/.....	1.4	6.4	11.8	16.6	14.4	19.0	23.9	27.1	28.7	29.0
2003 4/.....	1.8	5.4	12.8	13.3	15.5	14.8	23.4	24.0	27.2	33.0
2002.....	1.9	5.3	11.5	13.8	12.0	16.5	23.9	24.2	24.9	28.8
2001.....	2.1	6.5	9.9	13.4	12.1	16.7	20.5	22.8	31.4	27.0
2000.....	2.5	6.5	9.8	11.3	13.1	18.1	23.1	23.8	24.3	24.7
1999.....	1.9	6.3	7.6	10.4	13.1	17.9	24.6	22.7	26.9	28.4
Black, male										
2006 4/.....	2.9	7.5	23.5	39.7	32.7	31.9	36.0	36.7	40.4	27.7
2005 4/.....	2.1	8.4	25.0	33.6	33.3	31.6	37.0	33.1	49.9	26.0
2004 4/.....	2.5	11.0	21.8	30.6	24.1	31.5	40.1	32.8	46.6	36.8
2003 4/.....	2.9	9.6	23.9	22.0	25.5	21.7	35.8	37.9	36.2	51.1
2002.....	3.2	9.2	21.0	23.8	20.4	27.4	40.1	36.3	41.3	35.4
2001.....	3.4	11.2	17.8	23.2	17.9	23.9	27.9	32.1	40.9	34.0
2000.....	3.4	12.0	17.0	20.3	23.0	30.7	36.4	30.7	40.5	42.7
1999.....	3.1	11.5	13.8	17.5	20.4	28.6	37.3	34.7	34.9	24.6
Black, female										
2006 4/.....	*	2.5	4.5	5.7	6.6	8.5	11.0	15.3	24.6	23.8
2005 4/.....	*	2.1	3.4	4.0	8.1	7.2	14.1	14.1	21.9	22.9
2004 4/.....	*	2.4	3.2	5.1	6.8	9.7	13.0	23.7	19.3	26.0
2003 4/.....	*	1.6	3.4	6.1	7.7	9.7	15.1	15.7	22.4	25.9
2002.....	*	1.9	3.5	5.5	5.4	8.3	13.0	17.0	16.3	26.3
2001.....	*	2.3	3.1	5.4	7.5	11.5	15.5	17.2	26.6	24.4
2000.....	1.6	1.8	3.7	4.0	5.2	8.8	14.2	19.7	16.1	18.0
1999.....	*	1.9	2.3	4.7	7.3	10.0	16.2	15.4	22.9	29.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intrahepatic bile duct carcinoma (C22.1)										
All races 3/, both sexes										
2006.....	0.2	0.6	1.1	1.8	3.0	4.0	6.0	7.3	9.2	9.9
2005.....	0.3	0.6	1.1	1.6	2.8	3.9	5.4	8.1	8.9	9.1
2004.....	0.2	0.6	1.0	1.6	3.1	3.9	5.5	7.5	9.5	9.2
2003.....	0.2	0.6	1.1	1.6	2.8	3.7	5.4	7.1	8.5	9.1
2002.....	0.2	0.4	0.9	1.6	2.5	3.4	4.5	6.4	8.0	10.3
2001.....	0.2	0.4	0.8	1.3	2.3	3.5	4.8	6.4	8.2	9.9
2000.....	0.2	0.5	0.9	1.4	2.3	3.3	4.5	6.4	7.2	9.3
1999.....	0.2	0.4	0.7	1.4	2.2	3.3	4.3	5.7	7.7	8.1
All races 3/, male										
2006.....	0.2	0.7	1.1	2.1	3.4	4.7	6.6	8.1	10.5	12.0
2005.....	0.3	0.7	1.2	1.8	3.2	4.5	6.6	9.7	10.5	11.3
2004.....	0.3	0.7	1.1	1.9	3.7	4.1	6.4	8.2	11.5	10.9
2003.....	0.2	0.6	1.1	1.9	3.2	4.3	6.7	8.6	10.2	12.4
2002.....	0.3	0.5	1.0	1.8	2.8	3.9	5.3	8.0	10.0	11.6
2001.....	0.3	0.4	1.0	1.3	2.7	4.0	5.7	7.9	10.3	11.9
2000.....	0.3	0.6	0.9	1.4	2.5	3.9	5.5	7.9	9.0	10.8
1999.....	0.2	0.4	0.7	1.7	2.3	3.8	5.2	6.7	8.8	8.9
All races 3/, female										
2006.....	0.3	0.5	1.1	1.6	2.6	3.4	5.4	6.6	8.4	8.9
2005.....	0.4	0.4	1.0	1.4	2.4	3.3	4.4	7.0	7.9	8.1
2004.....	0.2	0.5	0.8	1.4	2.6	3.8	4.8	7.0	8.2	8.5
2003.....	0.3	0.6	1.1	1.3	2.4	3.2	4.3	6.0	7.5	7.6
2002.....	0.2	0.3	0.8	1.5	2.3	3.0	3.9	5.3	6.8	9.8
2001.....	0.2	0.4	0.6	1.3	2.0	3.0	4.1	5.3	7.0	9.0
2000.....	0.2	0.4	0.9	1.3	2.2	2.8	3.8	5.4	6.1	8.8
1999.....	0.2	0.4	0.8	1.1	2.1	3.0	3.6	5.0	7.0	7.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intrahepatic bile duct carcinoma (C22.1)										
White, both sexes										
2006 4/.....	0.3	0.6	1.1	1.8	3.1	4.0	6.0	7.2	9.2	10.0
2005 4/.....	0.3	0.6	1.1	1.6	2.7	3.8	5.2	8.1	8.5	9.3
2004 4/.....	0.2	0.6	0.9	1.6	3.2	3.8	5.5	7.5	9.5	9.3
2003 4/.....	0.3	0.5	1.0	1.6	2.7	3.6	5.3	7.1	8.7	9.2
2002.....	0.2	0.4	0.8	1.5	2.5	3.4	4.5	6.4	7.9	10.2
2001.....	0.2	0.4	0.8	1.3	2.3	3.5	4.7	6.4	7.9	9.8
2000.....	0.2	0.5	0.9	1.3	2.2	3.4	4.6	6.3	7.3	9.5
1999.....	0.2	0.4	0.7	1.3	2.2	3.3	4.4	5.7	7.7	8.3
White, male										
2006 4/.....	0.3	0.7	1.1	1.9	3.5	4.7	6.7	8.2	10.6	12.2
2005 4/.....	0.3	0.7	1.1	1.8	3.2	4.3	6.3	9.4	9.9	11.8
2004 4/.....	0.3	0.6	1.1	1.8	3.7	3.9	6.3	8.3	12.0	11.1
2003 4/.....	0.2	0.5	1.0	2.0	3.1	4.0	6.6	8.7	10.4	12.5
2002.....	0.3	0.5	1.0	1.7	2.8	3.7	5.3	7.9	9.7	11.6
2001.....	0.3	0.4	0.9	1.2	2.5	4.0	5.5	7.8	10.2	11.8
2000.....	0.2	0.6	0.9	1.3	2.5	3.9	5.5	7.9	9.3	11.3
1999.....	0.2	0.5	0.7	1.6	2.3	3.7	5.3	6.6	8.6	9.2
White, female										
2006 4/.....	0.3	0.5	1.0	1.7	2.7	3.4	5.5	6.5	8.4	9.0
2005 4/.....	0.3	0.4	1.0	1.5	2.3	3.3	4.3	7.2	7.7	8.1
2004 4/.....	0.2	0.5	0.8	1.4	2.7	3.7	4.8	6.9	7.9	8.5
2003 4/.....	0.3	0.5	1.1	1.3	2.3	3.1	4.3	6.0	7.6	7.7
2002.....	*	0.3	0.7	1.4	2.2	3.1	3.9	5.3	6.7	9.6
2001.....	*	0.4	0.6	1.3	2.1	3.0	4.1	5.5	6.6	9.0
2000.....	0.2	0.4	0.9	1.3	2.0	2.9	3.9	5.2	6.2	8.8
1999.....	0.2	0.4	0.8	1.0	2.0	3.0	3.7	5.1	7.1	7.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of gallbladder and extrahepatic bile ducts (C23-C24)										
All races 3/, both sexes										
2006.....	0.2	0.4	0.8	1.4	2.2	3.3	5.1	7.6	9.7	12.6
2005.....	0.2	0.5	0.7	1.3	2.3	3.4	5.6	7.5	10.1	13.1
2004.....	0.2	0.2	0.6	1.3	2.3	3.6	5.9	8.0	9.7	13.3
2003.....	0.1	0.4	0.8	1.2	2.4	3.3	5.3	8.1	10.9	14.3
2002.....	0.2	0.4	0.8	1.4	2.0	3.7	5.7	8.3	9.9	14.5
2001.....	0.2	0.5	0.8	1.4	2.3	3.7	6.0	8.3	12.1	15.8
2000.....	0.2	0.4	0.9	1.5	2.6	4.2	6.4	9.1	11.4	16.0
1999.....	0.2	0.4	0.8	1.6	2.5	4.2	6.0	8.8	11.9	16.2
All races 3/, male										
2006.....	*	0.2	0.7	1.0	2.0	2.7	4.9	7.1	9.2	12.0
2005.....	0.2	0.3	0.6	1.2	2.0	2.7	4.9	7.0	9.2	12.3
2004.....	*	0.2	0.5	1.2	1.7	3.2	5.2	8.1	9.4	12.9
2003.....	*	0.4	0.6	1.0	2.0	3.4	5.1	7.6	9.8	14.1
2002.....	*	0.2	0.6	1.2	1.8	3.3	5.5	8.2	9.1	15.0
2001.....	*	0.3	0.7	1.2	1.7	2.8	5.3	8.1	12.8	15.8
2000.....	*	0.3	0.6	1.1	2.2	3.9	6.2	8.6	11.0	15.8
1999.....	0.2	0.3	0.6	1.2	2.1	3.8	5.4	8.3	11.0	15.2
All races 3/, female										
2006.....	0.3	0.5	0.9	1.7	2.4	3.8	5.3	8.0	10.1	13.0
2005.....	0.2	0.6	0.8	1.5	2.6	4.0	6.2	7.9	10.7	13.5
2004.....	0.2	0.3	0.8	1.5	2.9	4.0	6.5	7.9	10.0	13.5
2003.....	*	0.5	0.9	1.4	2.7	3.1	5.5	8.5	11.5	14.4
2002.....	0.3	0.6	1.0	1.6	2.2	4.1	5.8	8.3	10.3	14.3
2001.....	0.3	0.6	0.9	1.5	2.8	4.5	6.5	8.5	11.7	15.8
2000.....	0.2	0.5	1.1	1.8	2.9	4.5	6.6	9.4	11.7	16.0
1999.....	0.3	0.4	1.0	2.0	2.9	4.6	6.5	9.1	12.4	16.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of gallbladder and extrahepatic bile ducts (C23-C24)										
White, both sexes										
2006 4/.....	0.2	0.3	0.8	1.4	2.2	3.2	5.2	7.4	9.7	12.6
2005 4/.....	0.2	0.4	0.6	1.3	2.3	3.4	5.6	7.4	9.8	13.0
2004 4/.....	0.2	0.2	0.6	1.3	2.2	3.4	5.8	7.9	9.6	13.4
2003 4/.....	*	0.4	0.7	1.2	2.4	3.3	5.3	8.0	10.7	14.4
2002.....	0.2	0.4	0.8	1.3	2.0	3.7	5.6	8.1	9.8	14.5
2001.....	0.2	0.5	0.8	1.4	2.2	3.5	5.9	8.3	12.2	15.8
2000.....	0.2	0.4	0.8	1.5	2.5	4.2	6.3	9.2	11.6	16.3
1999.....	0.2	0.3	0.8	1.6	2.5	4.2	6.0	8.7	11.5	16.4
White, male										
2006 4/.....	*	0.2	0.7	1.1	2.1	2.7	5.1	6.4	9.3	12.3
2005 4/.....	*	0.3	0.6	1.2	2.0	2.7	4.8	6.8	8.5	12.3
2004 4/.....	*	0.2	0.5	1.1	1.7	2.9	5.2	8.0	9.3	13.2
2003 4/.....	*	0.4	0.5	1.0	2.1	3.6	5.0	7.5	9.7	14.5
2002.....	*	*	0.6	1.1	1.8	3.3	5.4	8.1	9.0	15.3
2001.....	*	0.3	0.7	1.2	1.7	2.8	5.3	8.0	12.8	15.5
2000.....	*	0.3	0.6	1.1	2.2	4.1	6.0	8.7	11.3	16.2
1999.....	*	0.3	0.6	1.2	2.1	3.6	5.4	8.5	10.8	15.8
White, female										
2006 4/.....	0.3	0.4	0.9	1.6	2.3	3.7	5.2	8.1	9.9	12.8
2005 4/.....	*	0.5	0.7	1.4	2.6	3.9	6.2	7.9	10.7	13.4
2004 4/.....	*	0.2	0.7	1.5	2.7	3.8	6.3	7.7	9.9	13.5
2003 4/.....	*	0.4	0.9	1.4	2.8	3.1	5.4	8.5	11.2	14.4
2002.....	0.3	0.5	1.0	1.6	2.2	4.0	5.8	8.1	10.2	14.2
2001.....	0.3	0.6	0.9	1.5	2.7	4.1	6.4	8.5	11.9	15.9
2000.....	0.2	0.5	1.1	1.8	2.8	4.3	6.6	9.6	11.8	16.3
1999.....	0.2	0.4	1.0	2.0	2.9	4.7	6.5	8.9	12.0	16.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of gallbladder and extrahepatic bile ducts (C23-C24)

Black, both sexes

2006 4/.....	*	*	1.0	1.7	2.4	3.7	4.6	10.7	10.2	9.9
2005 4/.....	*	0.8	1.0	1.9	1.9	3.2	5.9	9.0	13.1	13.4
2004 4/.....	*	*	*	1.7	3.0	5.0	6.2	8.1	9.7	11.3
2003 4/.....	*	*	1.0	1.3	1.8	2.6	5.3	7.7	11.2	12.3
2002.....	*	*	1.1	2.2	2.5	4.7	5.9	10.0	9.4	14.3
2001.....	*	*	*	1.4	3.0	5.2	6.5	7.6	10.6	15.7
2000.....	*	*	1.3	*	3.0	5.1	6.7	7.6	8.2	11.6
1999.....	*	*	*	1.8	2.6	4.2	6.5	7.9	13.3	12.1

Black, male

2006 4/.....	*	*	*	*	*	*	*	14.8	*	*
2005 4/.....	*	*	*	*	*	*	*	8.9	14.7	*
2004 4/.....	*	*	*	*	*	*	*	9.6	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	7.7	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	*	*	*	2.3	2.8	4.8	5.5	8.3	11.9	11.7
2005 4/.....	*	*	*	2.3	*	4.1	5.8	9.1	12.2	13.5
2004 4/.....	*	*	*	*	4.5	5.8	6.8	7.3	10.2	14.2
2003 4/.....	*	*	*	*	*	*	5.5	7.9	12.8	14.2
2002.....	*	*	*	2.6	*	5.6	6.0	10.7	9.9	16.4
2001.....	*	*	*	*	4.2	6.4	6.5	7.5	10.2	16.0
2000.....	*	*	2.2	*	3.6	6.9	6.0	7.7	9.3	12.0
1999.....	*	*	*	2.9	*	4.0	6.5	8.3	13.8	14.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pancreas (C25)										
All races 3/, both sexes										
2006.....	1.7	4.4	8.5	15.3	26.2	37.7	53.4	70.2	89.7	96.9
2005.....	1.9	4.5	8.7	15.7	25.1	37.6	52.6	69.6	88.4	97.4
2004.....	1.7	4.2	8.8	15.2	25.8	37.3	52.4	68.9	85.4	97.7
2003.....	1.7	4.1	8.7	14.7	24.7	36.4	53.0	67.6	86.2	95.2
2002.....	1.9	4.2	8.2	15.5	24.0	37.2	53.2	70.2	81.8	94.5
2001.....	1.8	4.0	8.6	14.9	24.8	37.0	52.8	67.2	84.2	100.1
2000.....	1.8	4.2	8.7	15.1	24.8	37.2	51.4	70.4	85.7	96.4
1999.....	1.9	4.5	8.7	15.6	25.0	38.1	52.5	67.2	82.0	99.7
All races 3/, male										
2006.....	2.0	5.2	10.7	19.2	31.9	45.3	61.2	77.3	99.0	102.1
2005.....	2.2	5.2	10.6	19.6	30.6	43.9	61.6	78.7	96.5	102.8
2004.....	2.2	5.5	11.0	18.6	31.5	45.2	61.6	77.7	94.8	102.0
2003.....	2.0	5.2	10.7	18.2	29.8	42.5	60.9	76.1	99.5	95.6
2002.....	2.4	5.3	10.6	19.2	29.3	43.3	60.8	78.9	91.9	100.9
2001.....	2.0	4.9	10.7	18.5	30.3	43.6	59.0	77.2	94.9	105.1
2000.....	2.1	5.4	11.1	18.5	30.0	44.7	57.0	80.4	94.9	102.9
1999.....	2.4	5.9	11.0	19.1	29.9	44.9	60.9	75.7	92.0	107.5
All races 3/, female										
2006.....	1.4	3.5	6.3	11.7	21.0	31.1	47.0	64.9	83.9	94.4
2005.....	1.6	3.8	6.9	12.0	20.0	32.1	45.3	63.0	83.4	94.9
2004.....	1.3	2.9	6.7	12.0	20.6	30.5	45.1	62.6	79.6	95.8
2003.....	1.4	3.0	6.8	11.5	20.2	31.1	46.7	61.6	78.1	95.0
2002.....	1.4	3.1	6.0	12.0	19.2	31.9	47.1	64.1	75.6	91.7
2001.....	1.6	3.1	6.5	11.5	19.8	31.3	47.8	60.1	77.8	98.1
2000.....	1.5	3.0	6.4	11.8	20.0	30.8	47.1	63.5	80.2	93.8
1999.....	1.4	3.1	6.5	12.3	20.5	32.2	46.0	61.2	76.1	96.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pancreas (C25)										
White, both sexes										
2006 4/.....	1.6	4.2	8.2	14.9	25.6	37.3	52.6	70.0	89.5	96.1
2005 4/.....	1.9	4.4	8.2	15.2	24.6	37.1	52.2	68.8	87.7	96.8
2004 4/.....	1.6	4.0	8.4	14.6	25.1	36.8	51.3	68.3	85.1	97.1
2003 4/.....	1.6	3.9	8.1	14.3	23.8	36.0	52.2	67.2	85.2	94.4
2002.....	1.9	4.1	7.8	15.0	23.3	36.1	52.0	69.2	81.4	94.3
2001.....	1.7	3.8	8.3	14.4	24.7	36.3	51.7	66.3	83.3	99.0
2000.....	1.7	4.1	8.4	14.5	24.3	36.2	50.4	69.2	84.5	94.7
1999.....	1.8	4.4	8.2	14.8	24.3	37.2	51.5	65.8	80.6	98.6
White, male										
2006 4/.....	1.9	5.1	10.3	18.7	31.3	45.1	60.9	77.9	100.3	103.4
2005 4/.....	2.2	5.1	10.0	18.9	30.4	43.4	62.0	78.1	96.2	102.9
2004 4/.....	2.0	5.3	10.5	17.7	30.8	44.5	60.0	77.7	95.8	102.9
2003 4/.....	1.8	4.8	10.0	17.7	28.9	42.7	60.2	76.1	99.3	96.0
2002.....	2.4	5.1	10.1	18.6	28.8	42.6	59.7	78.9	92.2	101.6
2001.....	1.9	4.7	10.5	18.2	30.3	42.8	58.0	77.0	94.5	106.5
2000.....	2.0	5.3	10.7	17.8	29.6	44.3	55.9	79.9	94.0	102.4
1999.....	2.3	5.8	10.3	17.9	29.2	44.0	60.1	74.6	90.9	107.4
White, female										
2006 4/.....	1.3	3.3	6.1	11.3	20.3	30.3	45.7	64.1	82.7	92.7
2005 4/.....	1.5	3.6	6.4	11.7	19.3	31.5	44.0	61.9	82.4	94.0
2004 4/.....	1.3	2.8	6.2	11.5	19.7	30.0	44.1	61.5	78.5	94.5
2003 4/.....	1.4	2.9	6.3	11.0	19.1	30.0	45.8	60.7	76.6	93.8
2002.....	1.4	3.0	5.5	11.6	18.2	30.3	45.7	62.2	74.9	91.1
2001.....	1.6	2.9	6.3	10.8	19.6	30.6	46.5	58.7	76.5	95.9
2000.....	1.4	2.9	6.1	11.4	19.4	29.2	46.0	61.6	78.9	91.5
1999.....	1.3	3.1	6.3	11.9	19.9	31.3	44.6	59.5	74.5	95.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of pancreas (C25)										
Black, both sexes										
2006 4/.....	2.3	6.2	12.6	21.6	37.2	49.2	67.3	85.1	99.9	113.4
2005 4/.....	2.2	6.2	13.7	22.3	33.6	49.1	66.4	86.9	105.1	111.4
2004 4/.....	2.6	6.0	13.3	22.6	36.3	48.0	70.5	84.3	93.3	113.6
2003 4/.....	2.7	6.4	14.5	20.7	36.1	47.1	68.0	83.6	107.4	112.9
2002.....	2.5	5.6	12.3	22.4	34.4	50.4	73.7	88.6	91.8	106.6
2001.....	2.2	6.4	12.1	20.7	31.6	48.5	70.8	84.9	104.8	121.3
2000.....	2.6	5.8	13.2	22.8	33.3	50.3	69.8	93.8	104.2	119.8
1999.....	2.8	6.2	14.1	25.0	32.2	50.9	70.1	87.5	104.6	118.6
Black, male										
2006 4/.....	2.7	6.6	16.8	28.7	45.8	58.6	75.7	86.8	98.6	91.1
2005 4/.....	2.3	7.2	17.3	30.1	39.6	59.2	71.5	96.1	116.6	100.4
2004 4/.....	3.4	7.6	16.1	30.0	44.8	60.6	88.1	88.7	91.1	100.3
2003 4/.....	3.5	9.0	18.6	26.2	42.8	50.0	80.8	89.0	112.2	104.3
2002.....	2.9	7.2	16.5	29.0	40.8	56.6	84.1	90.5	105.5	100.8
2001.....	2.3	7.9	15.2	24.2	38.2	60.0	78.8	95.2	109.2	99.8
2000.....	3.0	7.3	17.5	29.3	40.4	55.7	78.9	101.6	116.3	109.6
1999.....	3.4	8.3	19.5	34.1	39.5	60.0	78.4	100.7	117.7	114.8
Black, female										
2006 4/.....	1.9	5.8	9.1	15.8	30.6	42.2	61.5	84.1	100.6	122.5
2005 4/.....	2.2	5.3	10.7	15.8	28.9	41.6	63.0	81.3	99.0	115.9
2004 4/.....	1.9	4.6	11.0	16.6	29.7	38.7	58.6	81.7	94.5	118.9
2003 4/.....	2.1	4.1	11.0	16.2	30.8	45.0	59.4	80.3	104.8	116.3
2002.....	2.1	4.3	8.8	17.0	29.4	45.9	66.8	87.5	84.7	108.8
2001.....	2.2	5.1	9.6	17.8	26.4	40.0	65.5	78.8	102.6	129.2
2000.....	2.3	4.4	9.4	17.4	27.8	46.3	63.7	89.1	98.1	123.6
1999.....	2.2	4.3	9.5	17.6	26.5	44.3	64.6	79.5	98.1	120.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and ill-defined digestive organs (C26)

All races 3/, both sexes

2006.....	*	0.1	0.3	0.4	0.8	1.2	1.6	2.6	3.1	5.2
2005.....	*	0.1	0.2	0.3	0.6	0.9	1.6	1.9	2.8	4.8
2004.....	0.1	0.1	0.2	0.4	0.7	1.0	1.4	2.3	3.1	5.1
2003.....	*	0.2	0.2	0.5	0.7	1.5	1.9	2.9	4.0	7.0
2002.....	*	0.1	0.2	0.5	0.7	1.3	2.0	3.4	4.1	6.9
2001.....	*	0.2	0.3	0.5	0.9	1.3	2.2	3.2	4.3	7.8
2000.....	*	0.2	0.3	0.5	1.0	1.3	2.1	2.8	4.5	7.8
1999.....	0.1	0.3	0.3	0.6	1.0	1.6	2.3	3.3	5.3	9.1

All races 3/, male

2006.....	*	*	0.3	0.5	1.0	1.4	2.0	3.0	4.3	5.1
2005.....	*	*	0.3	0.4	0.9	1.2	2.3	2.5	3.1	5.1
2004.....	*	*	0.2	0.4	0.9	1.3	1.7	2.9	3.5	5.0
2003.....	*	*	0.3	0.7	0.9	2.0	2.2	3.8	4.5	7.3
2002.....	*	*	0.3	0.7	0.8	2.0	2.6	4.3	5.3	6.4
2001.....	*	0.2	0.4	0.7	1.3	1.4	2.7	4.1	4.5	8.9
2000.....	*	*	0.3	0.7	1.3	1.6	3.1	3.4	5.2	7.5
1999.....	0.2	0.3	0.5	0.7	1.3	1.9	2.6	4.2	6.9	10.4

All races 3/, female

2006.....	*	*	0.2	0.3	0.5	0.9	1.2	2.3	2.4	5.2
2005.....	*	*	*	0.3	0.4	0.7	1.1	1.5	2.6	4.8
2004.....	*	*	*	0.4	0.5	0.8	1.3	1.9	2.8	5.2
2003.....	*	*	*	0.4	0.5	1.0	1.6	2.3	3.7	6.8
2002.....	*	*	*	0.3	0.7	0.8	1.5	2.7	3.4	7.2
2001.....	*	*	*	0.3	0.7	1.2	1.8	2.6	4.2	7.3
2000.....	*	*	0.2	0.4	0.7	1.0	1.4	2.3	4.1	8.0
1999.....	*	*	*	0.4	0.7	1.4	2.1	2.7	4.4	8.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and ill-defined digestive organs (C26)

White, both sexes

2006 4/.....	*	0.1	0.3	0.4	0.7	1.1	1.5	2.6	3.1	5.3
2005 4/.....	*	*	0.2	0.3	0.6	0.9	1.6	1.9	2.8	4.9
2004 4/.....	*	0.1	0.2	0.4	0.7	0.9	1.4	2.3	3.2	5.3
2003 4/.....	*	0.1	0.2	0.5	0.6	1.5	1.9	2.9	3.9	7.0
2002.....	*	*	0.2	0.5	0.7	1.3	1.9	3.5	4.0	7.0
2001.....	*	0.1	0.2	0.5	0.8	1.3	2.2	3.2	4.1	7.7
2000.....	*	0.2	0.2	0.6	1.0	1.2	2.0	2.8	4.3	7.9
1999.....	0.1	0.2	0.3	0.5	1.0	1.6	2.3	3.3	5.5	9.0

White, male

2006 4/.....	*	*	0.3	0.4	1.0	1.4	1.9	3.0	4.3	4.9
2005 4/.....	*	*	0.3	0.3	0.9	1.2	2.1	2.4	3.0	4.9
2004 4/.....	*	*	*	0.4	0.8	1.2	1.7	2.7	3.6	5.3
2003 4/.....	*	*	0.3	0.7	0.9	2.1	2.1	3.6	4.5	7.0
2002.....	*	*	0.3	0.7	0.7	2.0	2.4	4.4	5.3	6.2
2001.....	*	*	0.4	0.7	1.1	1.4	2.7	4.1	4.4	8.8
2000.....	*	*	0.3	0.7	1.3	1.5	2.8	3.6	5.1	7.7
1999.....	*	0.3	0.4	0.7	1.3	1.8	2.6	4.1	7.1	10.3

White, female

2006 4/.....	*	*	0.3	0.3	0.4	0.8	1.2	2.3	2.3	5.4
2005 4/.....	*	*	*	0.3	*	0.8	1.1	1.6	2.6	4.9
2004 4/.....	*	*	*	0.3	0.5	0.7	1.2	2.0	2.9	5.3
2003 4/.....	*	*	*	0.3	0.4	1.0	1.6	2.3	3.6	7.0
2002.....	*	*	*	*	0.8	0.7	1.4	2.8	3.3	7.4
2001.....	*	*	*	*	0.6	1.2	1.8	2.5	3.9	7.3
2000.....	*	*	*	0.4	0.7	1.0	1.3	2.2	3.8	8.0
1999.....	*	*	*	0.4	0.7	1.5	2.1	2.7	4.6	8.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)										
All races 3/, both sexes										
2006.....	7.5	20.9	41.1	74.8	140.9	223.1	304.4	375.4	391.5	309.1
2005.....	8.7	21.4	41.8	76.7	147.2	225.3	316.3	380.5	389.3	311.0
2004.....	8.8	21.6	42.2	81.0	149.6	232.7	318.6	381.1	383.2	306.3
2003.....	9.0	22.2	42.4	84.8	154.3	236.8	321.9	384.6	389.5	307.9
2002.....	9.6	21.5	43.4	89.5	157.8	239.7	328.9	386.5	387.5	305.1
2001.....	10.0	21.6	44.1	92.3	159.7	246.4	331.7	382.6	377.8	312.2
2000.....	9.8	22.0	45.6	95.9	165.1	252.9	333.8	377.8	385.1	310.8
1999.....	9.7	20.9	46.4	96.7	166.0	250.7	331.0	370.6	372.2	306.6
All races 3/, male										
2006.....	7.9	23.0	50.2	93.3	174.9	282.5	390.4	500.3	561.2	475.2
2005.....	9.1	24.2	51.5	95.7	184.8	286.8	412.0	511.6	557.6	486.7
2004.....	9.6	25.1	52.6	100.8	187.3	296.4	413.5	523.7	545.3	485.7
2003.....	10.2	26.3	52.9	106.1	192.7	303.6	424.2	529.4	559.7	493.3
2002.....	10.1	25.3	54.2	112.3	201.0	307.8	436.0	532.5	563.8	498.8
2001.....	11.2	25.9	55.2	115.5	204.0	323.8	449.3	538.5	558.1	534.1
2000.....	11.2	27.2	56.2	121.6	211.2	333.9	455.0	533.6	578.3	541.7
1999.....	11.2	25.9	57.7	124.3	216.4	334.1	453.2	529.0	569.3	550.6
All races 3/, female										
2006.....	7.2	18.9	32.4	57.4	109.9	171.1	234.4	284.0	285.4	231.3
2005.....	8.3	18.6	32.5	58.7	112.8	171.7	238.8	285.3	284.7	230.3
2004.....	8.1	18.3	32.3	62.3	115.3	177.3	242.2	278.7	283.2	225.6
2003.....	7.8	18.2	32.5	64.8	119.5	178.8	239.8	281.5	285.4	225.9
2002.....	9.0	17.8	32.9	68.0	118.6	180.9	243.4	283.4	280.3	221.1
2001.....	8.7	17.3	33.6	70.6	119.5	179.8	238.3	273.4	269.7	219.3
2000.....	8.4	17.0	35.5	71.8	123.3	183.5	238.3	269.3	271.1	216.8
1999.....	8.4	16.1	35.5	70.9	120.4	179.4	235.5	260.5	256.8	208.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)										
White, both sexes										
2006 4/.....	7.3	20.0	39.0	73.7	140.9	226.3	310.7	384.4	397.5	314.7
2005 4/.....	8.5	20.5	40.2	74.6	146.6	228.4	322.6	388.9	394.8	313.9
2004 4/.....	8.5	20.7	40.3	78.9	149.9	235.0	325.1	386.6	386.8	310.7
2003 4/.....	8.7	20.7	40.6	83.5	154.9	239.9	327.1	389.3	393.6	311.2
2002.....	9.1	20.4	41.6	87.8	157.2	242.4	334.8	391.2	393.1	309.0
2001.....	9.5	20.2	42.6	91.1	160.7	248.3	335.4	385.9	380.2	314.8
2000.....	9.0	20.4	43.8	94.4	165.8	254.9	335.8	379.8	391.0	313.4
1999.....	9.2	19.3	44.3	95.3	165.4	251.2	333.1	371.7	374.7	308.1
White, male										
2006 4/.....	7.6	21.6	46.6	89.5	170.7	280.5	391.3	504.4	565.7	481.6
2005 4/.....	8.8	22.7	48.9	90.4	179.7	285.5	413.1	517.1	560.5	487.8
2004 4/.....	9.1	23.3	48.6	95.8	182.5	294.8	414.0	523.5	544.6	490.5
2003 4/.....	9.5	24.2	49.3	101.6	188.7	301.6	424.3	527.2	559.1	496.2
2002.....	9.4	23.3	50.6	107.5	194.8	305.5	436.9	530.6	564.7	503.6
2001.....	10.4	23.6	51.5	110.8	201.1	319.8	446.3	534.5	554.0	535.5
2000.....	9.8	24.8	52.0	116.2	208.2	329.8	450.1	527.8	580.1	543.3
1999.....	10.2	23.2	53.4	119.7	210.9	328.7	448.4	524.2	565.4	551.9
White, female										
2006 4/.....	7.1	18.4	31.5	58.4	113.0	178.0	243.9	294.8	291.3	236.3
2005 4/.....	8.2	18.2	31.6	59.4	115.8	177.5	247.7	294.0	290.9	233.9
2004 4/.....	7.9	18.1	32.1	62.5	119.7	182.0	252.0	286.5	288.7	229.6
2003 4/.....	7.8	17.3	32.1	66.1	123.6	185.4	247.6	289.6	291.5	229.3
2002.....	8.9	17.5	32.7	68.9	122.4	186.8	251.5	291.5	287.9	224.5
2001.....	8.6	16.9	33.8	72.2	123.4	185.6	245.6	280.6	275.2	222.3
2000.....	8.2	16.0	35.7	73.7	126.7	189.5	244.1	275.7	278.5	220.1
1999.....	8.2	15.4	35.5	72.1	123.5	183.7	241.5	264.7	262.1	210.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of respiratory and intrathoracic organs (C30-C39)										
Black, both sexes										
2006 4/.....	10.2	32.0	65.9	103.4	180.9	258.4	316.7	365.8	390.0	274.0
2005 4/.....	11.5	32.6	64.1	114.3	190.1	260.6	329.2	372.5	386.4	301.5
2004 4/.....	13.3	32.6	66.9	120.0	188.5	267.0	322.3	395.4	400.8	287.6
2003 4/.....	13.2	37.4	65.8	121.3	193.3	268.4	343.1	401.4	391.2	293.2
2002.....	14.9	34.8	68.4	127.3	203.6	278.0	343.7	407.8	378.3	278.0
2001.....	15.4	35.0	67.8	127.4	191.9	284.8	363.2	410.1	381.9	290.2
2000.....	16.8	37.8	74.0	131.6	200.8	291.8	375.8	412.1	357.5	295.9
1999.....	15.7	37.5	75.3	133.2	209.9	294.7	364.5	414.0	375.3	300.4
Black, male										
2006 4/.....	11.4	38.9	88.1	150.1	265.6	388.5	474.3	566.1	618.8	461.1
2005 4/.....	12.7	41.1	84.9	167.9	282.7	386.4	493.4	564.6	625.7	527.1
2004 4/.....	15.5	43.4	96.6	172.9	287.8	388.4	492.0	626.9	656.4	508.6
2003 4/.....	16.7	48.1	93.7	179.7	286.1	401.1	522.9	650.0	662.2	530.7
2002.....	18.1	46.6	98.0	184.4	311.8	417.4	531.1	666.4	642.7	498.5
2001.....	19.6	48.9	100.0	189.0	285.8	440.1	582.0	691.4	685.4	571.9
2000.....	23.1	51.1	108.6	200.9	294.2	453.8	593.7	688.0	645.7	584.7
1999.....	20.7	53.3	111.5	201.4	321.8	451.3	581.4	680.1	694.9	599.9
Black, female										
2006 4/.....	9.2	26.1	47.0	64.9	114.8	162.3	208.4	245.9	269.1	197.4
2005 4/.....	10.4	25.1	46.6	70.1	118.0	167.3	216.9	257.9	260.0	210.4
2004 4/.....	11.4	23.3	41.8	76.5	111.2	176.8	207.1	258.2	266.1	200.0
2003 4/.....	10.1	28.1	42.3	73.3	120.8	170.0	221.8	253.7	249.1	200.9
2002.....	12.0	24.6	43.2	80.4	118.9	174.6	218.0	253.3	240.7	193.5
2001.....	11.7	23.0	40.3	77.0	118.1	170.1	217.5	242.1	227.3	185.7
2000.....	11.3	26.3	44.4	74.8	127.0	172.2	231.1	245.8	212.3	188.4
1999.....	11.3	23.9	44.7	77.5	122.1	178.7	220.5	253.7	216.9	188.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of nasal cavity, middle ear and accessory sinuses (C30-C31)

All races 3/, both sexes

2006.....	*	0.1	0.2	0.2	0.4	0.4	0.5	0.8	0.9	1.1
2005.....	*	0.1	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.4
2004.....	*	0.1	0.1	0.3	0.3	0.5	0.7	0.8	1.1	1.4
2003.....	0.1	0.1	0.2	0.2	0.4	0.5	0.6	1.0	0.9	1.2
2002.....	*	0.2	0.2	0.2	0.4	0.4	0.7	0.7	1.0	1.2
2001.....	0.1	0.1	0.2	0.3	0.4	0.4	0.6	0.8	1.4	1.7
2000.....	0.1	*	0.2	0.3	0.4	0.5	0.6	0.6	0.9	1.2
1999.....	0.1	0.1	0.3	0.3	0.3	0.5	0.6	0.8	1.0	1.5

All races 3/, male

2006.....	*	*	0.2	0.3	0.4	0.5	0.6	0.8	1.1	1.3
2005.....	*	0.2	0.3	0.4	0.6	0.6	0.8	1.0	1.7	1.7
2004.....	*	*	*	0.4	0.4	0.5	1.0	1.0	1.5	1.8
2003.....	*	*	0.3	0.3	0.7	0.7	0.9	1.4	1.3	*
2002.....	*	0.3	0.3	0.3	0.6	0.7	1.0	1.0	1.1	1.7
2001.....	*	0.2	0.3	0.4	0.5	0.5	0.8	0.9	1.9	1.8
2000.....	*	*	0.2	0.4	0.6	0.7	0.9	1.0	1.1	*
1999.....	*	*	0.4	0.5	0.4	0.7	0.6	1.2	1.1	1.8

All races 3/, female

2006.....	*	*	*	*	*	0.4	*	0.8	0.7	1.0
2005.....	*	*	*	*	*	*	0.4	0.6	0.8	1.2
2004.....	*	*	*	*	*	0.4	*	0.6	0.9	1.2
2003.....	*	*	*	*	*	*	*	0.7	0.6	1.1
2002.....	*	*	*	*	*	*	0.5	0.6	0.8	1.0
2001.....	*	*	*	*	*	*	0.4	0.8	1.1	1.6
2000.....	*	*	*	*	*	*	0.4	*	0.8	1.1
1999.....	*	*	*	*	*	*	0.6	0.6	0.9	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of nasal cavity, middle ear and accessory sinuses (C30-C31)										
White, both sexes										
2006 4/.....	*	*	0.2	0.2	0.3	0.5	0.5	0.8	0.9	1.2
2005 4/.....	*	0.1	0.2	0.2	0.3	0.4	0.6	0.7	1.2	1.4
2004 4/.....	*	*	0.1	0.3	0.3	0.4	0.7	0.8	1.1	1.4
2003 4/.....	0.1	*	0.2	0.2	0.5	0.4	0.7	0.9	0.9	1.2
2002.....	*	0.2	0.2	0.3	0.4	0.4	0.7	0.8	1.0	1.3
2001.....	*	0.2	0.2	0.3	0.4	0.4	0.5	0.8	1.4	1.7
2000.....	*	*	0.2	0.3	0.4	0.5	0.6	0.7	0.9	1.3
1999.....	0.1	*	0.3	0.3	0.3	0.5	0.7	0.9	1.0	1.5
White, male										
2006 4/.....	*	*	*	0.3	0.4	*	*	0.8	1.1	1.4
2005 4/.....	*	*	0.3	0.3	0.5	0.6	0.8	0.9	1.7	1.7
2004 4/.....	*	*	*	0.4	0.4	*	1.0	0.9	1.5	1.8
2003 4/.....	*	*	0.3	*	0.7	0.6	0.9	1.3	1.2	*
2002.....	*	0.3	0.3	0.4	0.6	0.6	0.9	1.0	1.2	1.7
2001.....	*	*	*	0.4	0.5	*	0.8	0.9	2.0	1.8
2000.....	*	*	*	0.4	0.6	0.6	0.9	1.0	1.3	*
1999.....	*	*	0.4	0.5	*	0.7	0.7	1.3	*	1.9
White, female										
2006 4/.....	*	*	*	*	*	0.5	*	0.8	0.7	1.0
2005 4/.....	*	*	*	*	*	*	0.5	0.6	0.9	1.2
2004 4/.....	*	*	*	*	*	*	*	0.6	0.9	1.2
2003 4/.....	*	*	*	*	*	*	*	0.7	*	1.2
2002.....	*	*	*	*	*	*	0.5	0.6	0.9	1.0
2001.....	*	*	*	*	*	*	*	0.8	1.1	1.7
2000.....	*	*	*	*	*	*	*	*	0.7	1.2
1999.....	*	*	*	*	*	*	0.7	0.6	0.9	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of larynx (C32)										
All races 3/, both sexes										
2006.....	0.3	0.7	1.4	2.6	3.8	5.2	6.2	6.9	7.4	6.3
2005.....	0.2	0.7	1.3	2.4	3.9	5.8	6.7	6.8	7.3	6.3
2004.....	0.2	0.7	1.4	2.6	4.1	5.3	6.2	7.1	6.0	6.5
2003.....	0.3	0.7	1.4	2.6	4.6	5.8	6.6	6.6	7.4	6.5
2002.....	0.2	0.7	1.4	2.7	4.2	5.3	6.6	7.9	7.2	5.5
2001.....	0.3	0.7	1.5	2.9	4.5	5.6	7.3	7.4	6.9	6.7
2000.....	0.2	0.8	1.5	2.9	4.5	6.0	7.3	8.0	8.1	6.5
1999.....	0.3	0.8	1.5	3.1	4.6	6.1	7.4	7.2	7.4	7.0
All races 3/, male										
2006.....	0.3	1.2	2.3	4.4	6.2	9.2	10.5	12.5	14.7	14.8
2005.....	0.4	1.0	2.1	4.0	6.7	10.2	11.4	12.2	14.5	15.7
2004.....	0.4	1.2	2.4	4.4	6.8	9.4	11.0	13.3	12.4	15.6
2003.....	0.5	1.2	2.4	4.2	7.9	10.3	11.4	12.6	15.4	15.3
2002.....	0.4	1.2	2.4	4.7	7.2	8.9	11.3	14.8	14.9	13.2
2001.....	0.5	1.2	2.5	4.9	7.5	9.5	12.9	14.5	13.9	18.3
2000.....	0.4	1.2	2.5	4.9	7.9	10.0	12.9	14.8	17.1	17.4
1999.....	0.4	1.3	2.6	5.4	7.8	10.4	12.6	13.2	15.8	17.8
All races 3/, female										
2006.....	*	0.3	0.5	1.0	1.5	1.7	2.7	2.8	2.8	2.3
2005.....	*	0.4	0.6	0.8	1.3	1.9	3.0	2.8	2.8	1.9
2004.....	*	0.2	0.5	0.9	1.6	1.7	2.2	2.6	2.1	2.4
2003.....	*	0.3	0.4	1.0	1.6	2.0	2.8	2.3	2.5	2.7
2002.....	*	0.2	0.5	0.9	1.5	2.2	2.9	3.1	2.5	2.2
2001.....	*	0.2	0.5	1.0	1.7	2.2	2.8	2.5	2.8	1.9
2000.....	*	0.3	0.6	0.9	1.4	2.6	2.8	3.3	2.8	2.1
1999.....	*	0.2	0.4	1.0	1.7	2.5	3.3	3.0	2.6	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of larynx (C32)										
White, both sexes										
2006 4/.....	0.2	0.7	1.2	2.4	3.6	4.8	6.1	6.8	7.2	6.2
2005 4/.....	0.2	0.6	1.1	2.1	3.4	5.4	6.3	6.8	7.2	6.0
2004 4/.....	0.2	0.7	1.2	2.3	3.8	5.0	5.9	6.9	5.7	6.3
2003 4/.....	0.3	0.6	1.1	2.3	4.2	5.5	6.2	6.3	7.1	6.2
2002.....	0.2	0.6	1.2	2.5	3.9	5.0	6.4	7.5	6.9	5.6
2001.....	0.3	0.6	1.2	2.5	4.1	5.2	6.7	7.0	6.8	6.6
2000.....	0.2	0.6	1.3	2.4	3.9	5.6	6.7	7.5	8.0	6.6
1999.....	0.2	0.6	1.3	2.7	4.2	5.7	7.1	7.1	7.3	6.7
White, male										
2006 4/.....	0.3	1.0	2.0	4.0	5.8	8.2	9.9	12.0	14.3	14.7
2005 4/.....	0.4	0.9	1.8	3.5	5.7	9.5	10.5	12.1	14.0	14.9
2004 4/.....	0.3	1.2	1.9	3.9	6.2	8.8	10.4	12.8	11.7	15.0
2003 4/.....	0.5	1.1	2.0	3.7	7.2	9.3	10.4	11.9	14.4	14.5
2002.....	0.3	1.0	2.0	4.1	6.3	8.3	10.7	13.8	14.6	13.3
2001.....	0.4	0.9	2.0	4.2	6.9	8.8	11.9	13.3	13.2	18.3
2000.....	0.3	1.0	2.1	4.1	6.8	9.2	11.7	13.9	16.8	17.6
1999.....	0.3	1.1	2.2	4.6	7.0	9.6	12.0	12.9	15.1	16.8
White, female										
2006 4/.....	*	0.3	0.5	0.9	1.5	1.7	2.9	3.0	2.8	2.2
2005 4/.....	*	0.3	0.5	0.8	1.3	1.7	2.9	2.9	2.9	1.9
2004 4/.....	*	0.2	0.5	0.9	1.6	1.7	2.1	2.6	2.0	2.5
2003 4/.....	*	*	0.3	0.9	1.4	2.1	2.7	2.3	2.5	2.5
2002.....	*	*	0.4	0.9	1.5	2.1	2.8	2.9	2.3	2.3
2001.....	*	*	0.4	0.9	1.5	2.1	2.6	2.5	2.9	1.7
2000.....	*	0.3	0.5	0.9	1.3	2.5	2.6	3.0	2.7	2.1
1999.....	*	*	0.3	1.0	1.5	2.3	3.3	3.0	2.7	2.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of larynx (C32)										
Black, both sexes										
2006 4/.....	*	1.6	2.8	5.4	6.6	10.9	9.6	10.2	9.5	8.0
2005 4/.....	*	1.3	3.3	5.5	9.1	11.5	11.4	9.2	10.9	10.2
2004 4/.....	*	1.2	3.5	5.5	8.0	9.3	11.6	11.2	11.1	8.2
2003 4/.....	*	1.7	3.6	5.8	9.7	10.6	12.7	11.1	12.7	11.1
2002.....	*	1.9	3.6	6.1	9.3	9.1	11.8	15.3	11.3	*
2001.....	*	1.9	3.8	7.1	8.9	9.5	13.9	13.4	9.2	8.3
2000.....	*	2.0	4.1	7.5	10.8	10.6	15.3	15.6	11.6	6.9
1999.....	0.9	2.1	3.9	7.5	10.0	11.4	11.8	10.2	9.8	8.3
Black, male										
2006 4/.....	*	2.8	5.2	9.5	12.2	21.6	19.4	22.4	21.2	18.8
2005 4/.....	*	2.1	5.0	10.3	18.5	22.1	22.1	18.3	26.0	28.8
2004 4/.....	*	2.1	6.7	11.0	14.9	18.7	22.7	24.1	25.8	23.8
2003 4/.....	*	2.7	6.8	10.7	17.5	22.7	25.3	25.6	31.7	27.1
2002.....	*	3.2	6.0	11.5	18.8	17.0	22.4	33.1	22.9	*
2001.....	*	3.6	7.0	13.0	15.9	17.5	26.3	30.7	22.5	*
2000.....	*	3.2	7.6	14.3	20.7	19.8	30.1	29.3	27.0	*
1999.....	1.6	3.9	6.9	14.4	17.8	21.0	22.7	19.9	27.9	24.6
Black, female										
2006 4/.....	*	*	*	2.0	*	*	*	*	*	*
2005 4/.....	*	*	1.8	*	*	3.7	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	3.6	*	*	*	*	*
2002.....	*	*	*	*	*	*	4.7	*	*	*
2001.....	*	*	*	*	3.4	*	5.6	*	*	*
2000.....	*	*	*	*	*	3.8	5.5	7.4	*	*
1999.....	*	*	*	*	3.8	4.2	4.5	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of trachea, bronchus and lung (C33-C34)										
All races 3/, both sexes										
2006.....	7.1	19.9	39.4	71.8	136.5	216.9	297.1	366.8	382.3	300.5
2005.....	8.2	20.4	40.1	73.8	142.7	218.7	308.3	372.3	380.0	302.3
2004.....	8.5	20.6	40.4	77.9	144.9	226.5	311.2	372.5	375.1	297.5
2003.....	8.5	21.2	40.7	81.8	149.0	230.0	314.0	376.1	380.1	298.9
2002.....	9.2	20.4	41.6	86.3	152.9	233.4	320.9	376.9	378.5	297.2
2001.....	9.5	20.5	42.3	88.8	154.5	239.7	323.2	373.4	368.6	302.7
2000.....	9.3	21.0	43.8	92.5	159.8	245.8	325.4	368.2	374.7	302.1
1999.....	9.3	19.9	44.4	93.0	160.7	243.6	322.3	361.7	362.6	297.0
All races 3/, male										
2006.....	7.3	21.6	47.5	88.3	167.8	272.4	378.8	486.0	544.1	457.7
2005.....	8.5	22.8	49.0	91.0	177.3	275.5	399.2	497.5	540.5	468.0
2004.....	9.1	23.5	49.8	95.8	179.7	285.8	400.8	508.4	530.2	466.8
2003.....	9.5	24.8	50.0	101.3	183.7	292.1	411.0	514.2	541.5	475.1
2002.....	9.6	23.6	51.4	107.0	192.7	297.5	423.0	515.5	546.8	482.2
2001.....	10.5	24.3	52.2	109.6	195.7	312.9	434.7	522.3	541.0	512.4
2000.....	10.5	25.7	53.2	115.9	202.3	322.5	440.5	516.5	558.1	521.2
1999.....	10.5	24.3	54.4	118.1	207.7	322.4	439.2	513.4	551.3	529.2
All races 3/, female										
2006.....	6.9	18.3	31.6	56.2	107.8	168.4	230.7	279.5	281.3	226.9
2005.....	8.0	18.1	31.7	57.6	111.2	169.1	234.7	281.3	280.3	226.2
2004.....	7.9	17.8	31.5	61.1	113.2	174.9	239.1	274.9	279.5	221.3
2003.....	7.5	17.8	31.9	63.5	117.4	176.2	236.1	277.9	281.4	221.0
2002.....	8.8	17.3	32.2	66.8	116.8	177.9	239.3	279.1	276.1	217.0
2001.....	8.4	16.8	32.8	69.2	117.2	176.7	234.5	269.1	265.2	215.0
2000.....	8.1	16.5	34.7	70.5	121.3	180.2	234.7	264.9	266.6	212.8
1999.....	8.1	15.7	34.9	69.5	118.1	176.1	230.9	256.2	252.2	203.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of trachea, bronchus and lung (C33-C34)										
White, both sexes										
2006 4/.....	7.0	19.1	37.5	70.9	136.6	220.6	303.6	375.7	388.5	306.1
2005 4/.....	8.1	19.6	38.7	72.0	142.7	222.2	315.0	380.7	385.6	305.6
2004 4/.....	8.1	19.7	38.8	76.1	145.5	229.2	318.0	378.3	378.9	302.0
2003 4/.....	8.2	19.9	39.1	80.7	149.9	233.5	319.7	381.3	384.5	302.4
2002.....	8.8	19.4	40.0	84.9	152.6	236.4	327.0	382.1	384.1	300.9
2001.....	9.1	19.3	40.9	87.9	155.9	242.0	327.4	377.2	371.2	305.4
2000.....	8.6	19.5	42.2	91.5	161.2	248.2	328.0	370.7	380.7	304.5
1999.....	8.8	18.4	42.6	92.0	160.6	244.5	324.6	362.9	365.2	298.8
White, male										
2006 4/.....	7.1	20.3	44.3	84.9	164.0	271.4	380.3	490.4	548.9	464.0
2005 4/.....	8.1	21.5	46.6	86.3	173.3	275.0	401.3	503.1	543.8	469.9
2004 4/.....	8.5	21.8	46.4	91.4	175.5	284.9	401.9	508.8	530.1	472.2
2003 4/.....	8.9	22.8	46.9	97.3	180.4	291.1	412.1	513.0	541.9	478.5
2002.....	8.9	21.8	48.1	102.8	187.3	295.9	424.5	514.5	547.8	486.6
2001.....	9.7	22.2	49.0	105.6	193.4	309.8	432.8	519.4	537.4	513.9
2000.....	9.3	23.5	49.5	111.5	200.5	319.1	436.8	511.6	560.0	522.4
1999.....	9.7	21.9	50.5	114.3	203.1	317.9	435.0	508.8	548.0	531.5
White, female										
2006 4/.....	6.8	17.9	30.8	57.3	111.0	175.3	240.0	290.2	287.2	231.9
2005 4/.....	8.0	17.7	30.9	58.3	114.2	175.2	243.6	290.1	286.4	229.9
2004 4/.....	7.7	17.7	31.3	61.3	117.7	179.7	249.0	282.8	285.0	225.3
2003 4/.....	7.5	17.0	31.5	64.8	121.6	182.6	244.0	286.0	287.4	224.4
2002.....	8.7	17.1	32.0	67.7	120.6	184.0	247.5	287.3	283.8	220.2
2001.....	8.4	16.4	33.1	71.0	121.3	182.5	242.1	276.4	270.7	218.1
2000.....	8.0	15.6	35.0	72.5	125.0	186.2	240.7	271.6	274.1	215.9
1999.....	7.9	15.0	35.0	70.8	121.6	180.5	236.9	260.4	257.3	205.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of trachea, bronchus and lung (C33-C34)										
Black, both sexes										
2006 4/.....	9.4	30.2	62.5	97.4	173.6	246.1	305.4	354.5	378.9	264.6
2005 4/.....	10.9	30.9	60.6	107.8	180.3	248.0	316.4	361.1	374.0	289.2
2004 4/.....	12.7	31.0	62.7	114.0	179.5	256.4	309.0	382.2	387.7	277.5
2003 4/.....	12.4	35.4	62.0	114.8	182.9	256.6	328.9	387.1	377.2	280.9
2002.....	14.1	32.4	64.4	120.6	193.7	267.2	330.7	391.2	366.3	272.3
2001.....	14.5	32.6	63.5	119.3	182.2	273.6	347.6	395.1	370.3	280.7
2000.....	15.7	35.5	69.3	123.1	188.7	279.9	359.1	394.9	343.3	288.1
1999.....	14.5	35.1	70.8	125.2	198.4	282.1	351.1	402.9	363.5	290.5
Black, male										
2006 4/.....	10.2	36.0	82.2	139.9	252.7	365.1	452.2	543.2	595.6	442.3
2005 4/.....	12.1	38.4	79.5	156.2	262.7	363.2	469.6	544.5	597.6	495.5
2004 4/.....	14.5	40.8	89.1	161.1	271.7	367.8	467.4	600.5	627.7	481.8
2003 4/.....	15.6	45.1	86.6	167.9	267.8	376.2	495.0	618.5	628.3	502.5
2002.....	17.0	42.6	91.6	172.1	292.0	398.4	506.4	630.6	619.0	484.6
2001.....	18.3	45.0	92.4	174.1	268.7	418.8	552.7	659.8	659.7	549.2
2000.....	21.4	47.4	100.4	185.2	271.7	432.2	561.6	656.8	617.1	562.8
1999.....	18.6	48.8	103.6	186.3	301.7	429.2	556.6	658.7	665.3	574.2
Black, female										
2006 4/.....	8.7	25.3	45.8	62.4	112.0	158.2	204.6	241.6	264.4	191.9
2005 4/.....	9.9	24.4	44.6	67.7	116.2	162.5	211.5	251.7	255.9	205.9
2004 4/.....	11.1	22.5	40.4	75.2	107.7	173.8	201.4	252.8	261.2	196.5
2003 4/.....	9.6	26.9	41.2	71.2	116.7	167.8	216.9	249.6	245.6	194.8
2002.....	11.5	23.6	41.5	78.3	116.8	169.8	212.8	248.1	234.7	191.0
2001.....	11.1	21.9	38.8	74.3	114.1	166.3	211.0	237.1	222.8	181.0
2000.....	10.7	25.1	42.7	72.2	123.2	167.6	224.7	237.0	205.5	185.9
1999.....	10.8	23.3	42.9	75.3	117.3	173.2	214.7	248.9	213.9	184.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of thymus, heart, mediastinum and pleura (C37-C38)

All races 3/, both sexes

2006.....	0.1	0.1	0.2	0.2	0.3	0.4	0.6	0.8	0.8	1.1
2005.....	0.1	0.1	0.1	0.2	0.2	0.4	0.6	0.7	0.8	1.0
2004.....	*	0.1	0.2	0.2	0.3	0.5	0.6	0.7	0.9	0.9
2003.....	0.1	0.1	*	0.2	0.3	0.4	0.6	0.8	0.9	1.3
2002.....	*	0.2	0.1	0.2	0.3	0.5	0.6	0.8	0.8	1.1
2001.....	0.1	0.2	0.2	0.3	0.3	0.6	0.6	0.8	0.9	1.0
2000.....	0.1	0.1	0.1	0.2	0.3	0.5	0.5	0.8	1.3	0.9
1999.....	*	0.1	0.2	0.3	0.3	0.4	0.7	0.8	1.0	1.0

All races 3/, male

2006.....	*	*	*	0.3	0.3	*	0.5	0.9	1.1	*
2005.....	*	*	*	0.3	*	0.5	0.6	0.9	*	*
2004.....	*	0.2	*	*	0.4	0.6	0.7	0.9	1.3	1.5
2003.....	*	*	*	0.3	0.4	0.5	0.7	1.0	1.2	1.7
2002.....	*	0.2	*	*	0.5	0.6	0.6	1.0	*	1.7
2001.....	*	0.2	0.2	0.5	*	0.7	0.8	0.8	1.2	1.5
2000.....	*	*	*	*	0.4	0.6	0.6	1.1	1.8	*
1999.....	*	*	0.3	0.3	0.4	*	0.7	1.0	*	*

All races 3/, female

2006.....	*	*	*	*	*	0.4	0.6	0.8	0.6	1.1
2005.....	*	*	*	*	*	0.4	0.7	0.6	0.7	0.9
2004.....	*	*	*	*	*	*	0.5	0.5	0.6	0.7
2003.....	*	*	*	*	*	*	0.5	0.7	0.7	1.1
2002.....	*	*	*	*	*	0.4	0.6	0.6	0.7	0.9
2001.....	*	*	*	*	*	0.4	0.5	0.8	0.7	0.8
2000.....	*	*	*	*	*	0.4	*	0.6	0.9	0.7
1999.....	*	*	*	*	*	0.4	0.6	0.7	1.0	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of thymus, heart, mediastinum and pleura (C37-C38)										
White, both sexes										
2006 4/.....	*	0.1	0.1	0.2	0.3	0.4	0.5	0.9	0.8	1.1
2005 4/.....	*	*	0.1	0.2	0.2	0.4	0.6	0.7	0.8	0.9
2004 4/.....	*	0.2	0.1	0.2	0.3	0.4	0.5	0.6	0.9	1.0
2003 4/.....	*	0.1	*	0.2	0.3	0.4	0.6	0.7	1.0	1.4
2002.....	*	0.2	0.1	*	0.3	0.5	0.7	0.8	0.8	1.3
2001.....	0.1	0.2	0.2	0.3	0.3	0.5	0.6	0.7	0.9	1.0
2000.....	*	*	*	0.2	0.3	0.5	0.5	0.8	1.3	0.9
1999.....	*	*	0.2	0.3	0.3	0.4	0.6	0.8	1.1	1.0
White, male										
2006 4/.....	*	*	*	0.3	*	*	*	0.9	1.1	*
2005 4/.....	*	*	*	0.3	*	*	*	1.0	*	*
2004 4/.....	*	*	*	*	0.4	0.6	0.7	0.9	1.4	1.6
2003 4/.....	*	*	*	*	*	*	0.7	0.8	1.3	1.8
2002.....	*	*	*	*	0.5	0.6	0.7	1.0	*	1.9
2001.....	*	*	*	0.5	*	0.6	0.8	0.8	1.3	*
2000.....	*	*	*	*	*	0.7	0.6	1.1	1.9	*
1999.....	*	*	*	0.4	*	*	0.7	1.0	*	*
White, female										
2006 4/.....	*	*	*	*	*	*	0.6	0.8	*	1.0
2005 4/.....	*	*	*	*	*	*	0.6	*	0.7	0.8
2004 4/.....	*	*	*	*	*	*	*	*	0.6	0.7
2003 4/.....	*	*	*	*	*	*	0.5	0.6	0.8	1.2
2002.....	*	*	*	*	*	*	0.7	0.6	0.7	1.0
2001.....	*	*	*	*	*	0.5	0.5	0.7	*	0.9
2000.....	*	*	*	*	*	*	*	0.6	0.9	*
1999.....	*	*	*	*	*	*	0.5	0.7	1.0	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of bone and articular cartilage (C40-C41)

All races 3/, both sexes

2006.....	0.2	0.3	0.3	0.4	0.6	0.8	1.4	1.4	2.1	3.0
2005.....	0.2	0.4	0.4	0.4	0.6	0.9	1.1	1.7	2.4	2.7
2004.....	0.2	0.2	0.3	0.5	0.7	0.9	1.1	1.5	2.2	2.7
2003.....	0.3	0.3	0.4	0.4	0.5	0.6	1.1	1.5	2.5	2.8
2002.....	0.2	0.3	0.3	0.5	0.7	0.6	1.0	1.3	2.0	2.9
2001.....	0.2	0.3	0.3	0.5	0.6	0.9	1.3	1.7	2.2	3.5
2000.....	0.2	0.3	0.3	0.5	0.6	0.9	1.0	1.8	2.1	3.4
1999.....	0.2	0.3	0.4	0.3	0.6	0.8	1.1	1.6	2.1	3.0

All races 3/, male

2006.....	0.3	0.3	0.4	0.6	0.9	1.1	1.7	1.6	2.2	3.5
2005.....	0.2	0.4	0.5	0.5	0.7	1.0	1.6	2.1	3.4	3.5
2004.....	0.2	0.4	0.5	0.6	0.8	1.1	1.1	1.6	2.7	3.2
2003.....	0.3	0.3	0.4	0.5	0.8	0.8	1.4	1.7	3.6	3.5
2002.....	0.3	0.4	0.4	0.6	0.7	0.7	1.1	1.8	2.4	3.7
2001.....	0.3	0.4	0.4	0.5	0.7	1.1	1.7	2.3	2.9	3.6
2000.....	0.2	0.4	0.5	0.7	0.7	1.2	1.2	2.2	2.7	4.2
1999.....	0.3	0.4	0.5	0.4	0.8	0.9	1.6	1.8	2.2	4.5

All races 3/, female

2006.....	0.2	0.3	0.3	0.3	0.4	0.6	1.0	1.2	2.0	2.8
2005.....	*	0.3	0.3	0.4	0.5	0.8	0.8	1.4	1.8	2.4
2004.....	*	*	0.2	0.4	0.7	0.8	1.1	1.5	1.8	2.5
2003.....	0.2	*	0.3	0.4	*	0.5	1.0	1.3	1.8	2.5
2002.....	*	*	*	0.4	0.6	0.6	1.0	1.0	1.8	2.5
2001.....	*	*	0.2	0.5	0.5	0.7	1.0	1.3	1.8	3.4
2000.....	*	*	*	0.4	0.5	0.7	0.8	1.5	1.8	3.1
1999.....	*	0.3	0.3	*	0.4	0.8	0.8	1.5	2.0	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of bone and articular cartilage (C40-C41)										
White, both sexes										
2006 4/.....	0.2	0.3	0.3	0.4	0.6	0.8	1.4	1.4	2.2	3.0
2005 4/.....	0.2	0.4	0.3	0.4	0.6	0.9	1.2	1.8	2.5	2.8
2004 4/.....	0.2	0.2	0.3	0.5	0.7	1.0	1.2	1.5	2.2	2.8
2003 4/.....	0.3	0.3	0.4	0.4	0.5	0.6	1.2	1.4	2.5	2.9
2002.....	0.2	0.3	0.3	0.5	0.6	0.6	1.0	1.3	2.1	3.0
2001.....	0.2	0.2	0.3	0.5	0.6	0.8	1.3	1.7	2.3	3.6
2000.....	0.2	0.3	0.3	0.5	0.6	1.0	1.0	1.8	2.2	3.5
1999.....	0.2	0.3	0.4	0.3	0.5	0.9	1.2	1.7	2.2	3.1
White, male										
2006 4/.....	0.3	0.3	0.4	0.6	0.8	1.0	1.8	1.6	2.3	3.4
2005 4/.....	0.2	0.4	0.4	0.5	0.8	1.1	1.6	2.3	3.5	3.5
2004 4/.....	0.3	0.4	0.4	0.6	0.8	1.1	1.2	1.7	2.7	3.2
2003 4/.....	0.3	0.3	0.4	0.5	0.8	0.8	1.4	1.8	3.7	3.5
2002.....	0.3	0.4	0.4	0.6	0.7	0.8	1.1	1.8	2.5	4.0
2001.....	0.3	0.3	0.3	0.5	0.7	1.0	1.7	2.3	3.1	3.6
2000.....	*	0.3	0.5	0.6	0.7	1.3	1.1	2.2	2.6	4.5
1999.....	0.2	0.4	0.5	0.4	0.7	1.0	1.6	1.9	2.3	4.8
White, female										
2006 4/.....	*	0.3	0.3	0.3	0.4	0.6	1.1	1.2	2.1	2.8
2005 4/.....	*	0.3	0.3	0.3	0.5	0.8	0.8	1.4	1.8	2.5
2004 4/.....	*	*	*	0.4	0.7	0.8	1.2	1.3	1.9	2.6
2003 4/.....	0.2	*	0.3	0.3	*	0.5	1.0	1.2	1.8	2.6
2002.....	*	*	*	0.4	0.6	0.5	1.0	1.0	1.8	2.6
2001.....	*	*	0.3	0.5	0.4	0.5	0.9	1.4	1.8	3.6
2000.....	*	*	*	0.4	0.5	0.7	0.9	1.4	1.9	3.1
1999.....	*	0.3	0.3	*	*	0.8	0.8	1.5	2.1	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Melanoma and other malignant neoplasms of skin (C43-C44)										
All races 3/, both sexes										
2006.....	1.5	2.6	3.8	5.6	8.1	10.5	13.9	18.4	24.7	33.8
2005.....	1.7	2.5	4.1	5.6	7.5	10.3	13.9	19.5	23.7	31.7
2004.....	1.6	2.6	4.0	5.8	7.8	10.1	13.7	17.9	22.3	30.2
2003.....	1.8	2.8	4.0	6.0	7.9	9.5	13.5	18.3	21.6	30.8
2002.....	1.8	2.6	4.1	5.4	7.7	10.0	13.2	17.8	21.8	29.8
2001.....	1.9	2.9	4.4	5.9	8.0	10.2	14.2	17.1	21.0	31.2
2000.....	1.9	3.0	4.5	5.8	7.8	10.5	13.3	16.6	20.3	31.8
1999.....	1.9	3.1	4.2	5.7	8.0	10.6	13.1	17.2	21.3	29.2
All races 3/, male										
2006.....	1.9	3.3	5.3	8.0	12.3	15.9	20.8	30.1	43.2	61.2
2005.....	2.2	3.2	5.5	8.1	10.7	15.3	21.5	30.8	39.4	56.4
2004.....	2.0	3.4	5.5	8.3	11.9	15.2	21.2	28.9	37.5	53.9
2003.....	2.2	3.6	5.6	8.6	11.6	14.0	20.4	29.8	35.4	53.5
2002.....	2.4	3.8	5.6	7.7	11.5	14.9	20.2	28.3	35.3	50.2
2001.....	2.4	3.8	5.9	8.3	11.9	15.6	22.2	27.1	34.5	54.0
2000.....	2.4	3.7	6.3	8.3	11.0	15.5	20.4	26.5	34.7	55.4
1999.....	2.5	4.2	5.7	8.4	11.6	15.5	20.5	26.9	35.2	50.6
All races 3/, female										
2006.....	1.2	1.8	2.4	3.3	4.3	5.8	8.2	9.9	13.2	21.0
2005.....	1.1	1.9	2.7	3.2	4.6	6.0	7.8	11.2	13.9	20.3
2004.....	1.2	1.8	2.5	3.3	4.2	5.6	7.7	9.9	12.9	19.5
2003.....	1.4	2.1	2.5	3.5	4.7	5.5	7.9	10.2	13.2	20.7
2002.....	1.3	1.5	2.7	3.2	4.3	5.7	7.6	10.5	13.6	21.0
2001.....	1.5	2.1	2.9	3.6	4.5	5.5	7.8	10.1	12.9	21.7
2000.....	1.4	2.2	2.7	3.4	4.9	6.2	7.8	9.7	11.8	22.2
1999.....	1.3	2.0	2.8	3.2	4.7	6.3	7.2	10.4	13.1	20.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Melanoma and other malignant neoplasms of skin (C43-C44)										
White, both sexes										
2006 4/.....	1.8	3.0	4.4	6.3	9.2	11.9	15.6	20.4	26.9	36.6
2005 4/.....	1.9	2.9	4.8	6.3	8.4	11.6	15.6	21.6	25.6	34.2
2004 4/.....	1.8	3.1	4.6	6.5	8.8	11.3	15.4	19.7	24.1	32.7
2003 4/.....	2.1	3.3	4.6	6.8	8.9	10.6	15.0	19.9	23.6	33.3
2002.....	2.2	3.1	4.7	6.0	8.6	11.1	14.5	19.6	23.8	32.2
2001.....	2.3	3.4	5.0	6.6	8.9	11.3	15.5	18.7	22.6	33.7
2000.....	2.2	3.4	5.1	6.4	8.7	11.6	14.5	17.9	21.8	34.3
1999.....	2.3	3.5	4.8	6.4	8.9	11.8	14.3	18.5	22.9	31.5
White, male										
2006 4/.....	2.2	3.8	6.0	9.0	13.7	17.8	23.2	33.0	46.9	66.5
2005 4/.....	2.5	3.5	6.4	9.0	11.9	17.1	24.0	33.9	42.8	61.7
2004 4/.....	2.3	3.9	6.3	9.3	13.1	17.1	23.7	31.7	40.5	58.9
2003 4/.....	2.6	4.2	6.4	9.7	12.9	15.6	22.7	32.4	38.6	58.2
2002.....	2.8	4.4	6.2	8.4	12.8	16.4	21.9	30.9	38.5	54.5
2001.....	2.8	4.3	6.7	9.3	13.1	17.1	24.1	29.6	37.4	58.7
2000.....	2.7	4.3	7.0	9.1	12.2	17.0	22.1	28.6	37.2	60.6
1999.....	3.0	4.7	6.3	9.3	12.8	17.2	22.3	28.7	37.9	55.0
White, female										
2006 4/.....	1.4	2.2	2.8	3.7	5.0	6.6	9.2	11.0	14.2	22.5
2005 4/.....	1.4	2.2	3.1	3.7	5.1	6.8	8.7	12.5	14.9	21.5
2004 4/.....	1.4	2.2	3.0	3.9	4.8	6.2	8.6	11.0	13.8	20.9
2003 4/.....	1.7	2.4	2.8	3.9	5.2	6.3	8.8	10.9	14.3	22.2
2002.....	1.6	1.8	3.2	3.7	4.8	6.4	8.4	11.5	14.7	22.5
2001.....	1.8	2.5	3.4	4.1	5.0	6.2	8.5	11.0	13.7	23.2
2000.....	1.7	2.6	3.2	3.7	5.4	6.9	8.4	10.4	12.6	23.6
1999.....	1.5	2.3	3.3	3.7	5.3	7.0	7.9	11.3	14.1	22.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant melanoma of skin (C43)										
All races 3/, both sexes										
2006.....	1.4	2.2	3.2	4.5	6.2	8.2	10.8	14.0	17.8	20.1
2005.....	1.5	2.2	3.5	4.6	5.9	8.1	10.9	15.1	16.2	19.8
2004.....	1.4	2.3	3.3	4.7	6.1	8.1	10.6	13.6	16.3	17.7
2003.....	1.6	2.5	3.4	4.9	6.4	7.3	10.3	13.9	15.5	17.6
2002.....	1.7	2.3	3.4	4.4	6.0	7.8	10.0	12.9	14.9	17.9
2001.....	1.8	2.6	3.6	4.6	6.1	7.8	10.7	12.4	14.7	17.8
2000.....	1.7	2.7	3.8	4.7	6.3	7.9	10.0	12.0	14.9	18.9
1999.....	1.7	2.6	3.6	4.7	6.1	8.3	9.7	12.9	14.2	16.4
All races 3/, male										
2006.....	1.6	2.8	4.3	6.2	9.0	11.9	15.9	22.5	29.9	36.7
2005.....	1.9	2.7	4.6	6.4	7.9	11.9	16.2	23.5	26.0	34.9
2004.....	1.8	2.9	4.5	6.6	8.9	11.8	16.1	21.8	26.7	31.4
2003.....	1.9	3.1	4.6	6.9	8.9	10.4	15.1	22.3	24.9	30.2
2002.....	2.1	3.3	4.4	6.1	8.5	11.3	14.9	19.8	23.3	29.2
2001.....	2.2	3.2	4.7	6.3	8.8	11.9	16.3	19.6	24.0	29.9
2000.....	2.1	3.2	5.2	6.5	8.7	11.2	14.7	18.4	24.7	31.8
1999.....	2.2	3.5	4.6	6.7	8.4	12.0	14.6	19.8	22.7	27.5
All races 3/, female										
2006.....	1.2	1.7	2.1	2.9	3.6	5.0	6.7	7.8	10.2	12.4
2005.....	1.1	1.8	2.4	2.9	4.2	4.8	6.6	9.0	10.1	12.9
2004.....	1.1	1.7	2.2	3.0	3.6	4.9	6.2	7.7	9.9	11.5
2003.....	1.3	1.9	2.2	3.1	4.0	4.6	6.4	7.8	9.8	12.1
2002.....	1.2	1.4	2.5	2.8	3.6	4.8	6.1	7.9	9.8	12.9
2001.....	1.4	1.9	2.6	3.0	3.7	4.3	6.4	7.3	9.2	12.8
2000.....	1.4	2.1	2.4	3.0	4.1	5.0	6.3	7.5	9.1	13.7
1999.....	1.3	1.8	2.6	2.8	4.0	5.2	5.8	8.1	9.2	11.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant melanoma of skin (C43)										
White, both sexes										
2006 4/.....	1.7	2.7	3.8	5.2	7.0	9.5	12.3	15.5	19.5	21.7
2005 4/.....	1.8	2.6	4.1	5.3	6.8	9.3	12.3	16.8	17.6	21.4
2004 4/.....	1.7	2.8	3.9	5.5	7.0	9.2	12.0	15.1	17.6	19.2
2003 4/.....	1.9	2.9	3.9	5.6	7.3	8.3	11.6	15.2	17.0	19.1
2002.....	2.0	2.8	4.0	5.0	6.8	8.9	11.0	14.2	16.3	19.2
2001.....	2.2	3.1	4.3	5.3	7.0	8.8	11.9	13.6	15.9	19.3
2000.....	2.0	3.1	4.4	5.3	7.2	8.9	11.0	13.1	16.0	20.3
1999.....	2.1	3.1	4.2	5.4	6.9	9.4	10.7	14.0	15.3	17.6
White, male										
2006 4/.....	2.0	3.3	5.1	7.1	10.2	13.7	18.0	24.8	32.8	39.7
2005 4/.....	2.2	3.1	5.4	7.3	9.0	13.6	18.3	26.0	28.3	38.1
2004 4/.....	2.1	3.5	5.3	7.5	10.1	13.4	18.1	24.0	28.9	34.2
2003 4/.....	2.3	3.6	5.3	7.9	10.1	11.7	17.1	24.4	27.1	32.9
2002.....	2.5	3.8	5.1	6.9	9.7	12.7	16.4	21.8	25.6	31.7
2001.....	2.6	3.8	5.5	7.2	10.0	13.3	18.0	21.5	26.1	32.7
2000.....	2.5	3.8	6.0	7.3	9.9	12.5	16.2	20.1	26.6	34.7
1999.....	2.7	4.1	5.3	7.7	9.5	13.4	16.2	21.5	24.5	29.7
White, female										
2006 4/.....	1.4	2.1	2.5	3.3	4.1	5.7	7.6	8.6	11.0	13.3
2005 4/.....	1.3	2.1	2.8	3.4	4.7	5.5	7.4	10.0	10.9	13.8
2004 4/.....	1.4	2.1	2.6	3.5	4.1	5.4	7.0	8.6	10.6	12.4
2003 4/.....	1.5	2.3	2.6	3.5	4.6	5.2	7.1	8.5	10.8	12.9
2002.....	1.5	1.7	2.9	3.2	4.2	5.5	6.7	8.8	10.7	13.8
2001.....	1.7	2.3	3.1	3.5	4.2	4.8	7.0	8.0	9.8	13.7
2000.....	1.6	2.5	2.8	3.4	4.6	5.7	6.9	8.2	9.7	14.5
1999.....	1.5	2.2	3.0	3.3	4.6	5.8	6.4	8.8	9.9	12.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other malignant neoplasms of skin (C44)										
All races 3/, both sexes										
2006.....	0.1	0.3	0.6	1.1	1.9	2.2	3.0	4.4	6.9	13.6
2005.....	0.2	0.3	0.6	1.0	1.6	2.2	3.1	4.4	7.5	11.8
2004.....	0.1	0.3	0.6	1.0	1.7	2.0	3.1	4.3	6.0	12.5
2003.....	0.2	0.3	0.7	1.0	1.6	2.2	3.2	4.4	6.1	13.1
2002.....	0.2	0.3	0.7	1.0	1.7	2.2	3.2	5.0	6.9	12.0
2001.....	0.1	0.3	0.7	1.3	1.9	2.4	3.5	4.8	6.3	13.4
2000.....	0.1	0.3	0.7	1.1	1.5	2.6	3.3	4.6	5.4	12.9
1999.....	0.2	0.4	0.7	1.0	1.9	2.3	3.4	4.3	7.0	12.8
All races 3/, male										
2006.....	0.2	0.5	1.0	1.8	3.3	3.9	4.9	7.6	13.3	24.6
2005.....	0.3	0.5	0.9	1.7	2.8	3.4	5.3	7.3	13.4	21.6
2004.....	0.2	0.4	1.0	1.7	3.0	3.5	5.1	7.2	10.9	22.5
2003.....	0.3	0.6	1.1	1.7	2.6	3.6	5.3	7.4	10.6	23.3
2002.....	0.2	0.5	1.2	1.6	3.0	3.7	5.3	8.5	12.0	21.0
2001.....	0.2	0.5	1.2	2.1	3.1	3.7	5.9	7.5	10.5	24.1
2000.....	0.2	0.5	1.1	1.9	2.4	4.2	5.7	8.1	9.9	23.6
1999.....	0.3	0.7	1.1	1.7	3.2	3.6	5.9	7.1	12.4	23.1
All races 3/, female										
2006.....	*	*	0.2	0.4	0.7	0.8	1.5	2.1	3.0	8.5
2005.....	*	*	0.3	0.3	0.4	1.2	1.2	2.3	3.8	7.4
2004.....	*	*	0.3	0.4	0.6	0.7	1.5	2.1	3.1	8.0
2003.....	*	*	0.3	0.4	0.6	0.9	1.5	2.3	3.3	8.6
2002.....	*	*	0.3	0.5	0.7	0.9	1.6	2.5	3.8	8.1
2001.....	*	*	0.2	0.5	0.8	1.2	1.5	2.8	3.7	8.9
2000.....	*	*	0.3	0.4	0.8	1.2	1.5	2.2	2.7	8.5
1999.....	*	*	0.3	0.4	0.7	1.1	1.4	2.4	3.9	8.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other malignant neoplasms of skin (C44)										
White, both sexes										
2006 4/.....	0.1	0.3	0.6	1.1	2.1	2.4	3.3	4.9	7.4	14.8
2005 4/.....	0.1	0.3	0.6	1.0	1.6	2.3	3.3	4.8	8.0	12.7
2004 4/.....	0.1	0.3	0.7	1.1	1.7	2.2	3.3	4.6	6.4	13.6
2003 4/.....	0.2	0.3	0.7	1.1	1.7	2.3	3.4	4.8	6.6	14.2
2002.....	0.2	0.3	0.7	1.0	1.8	2.2	3.4	5.4	7.4	13.0
2001.....	0.1	0.3	0.7	1.3	1.9	2.5	3.6	5.1	6.7	14.3
2000.....	0.1	0.3	0.7	1.1	1.5	2.7	3.5	4.9	5.8	14.0
1999.....	0.2	0.4	0.6	1.0	2.0	2.4	3.6	4.5	7.6	13.9
White, male										
2006 4/.....	0.2	0.5	0.9	1.8	3.5	4.1	5.3	8.2	14.1	26.8
2005 4/.....	0.2	0.4	1.0	1.8	2.9	3.5	5.7	7.9	14.5	23.6
2004 4/.....	*	0.4	1.0	1.8	2.9	3.7	5.5	7.7	11.6	24.7
2003 4/.....	0.3	0.6	1.1	1.9	2.7	3.8	5.6	7.9	11.5	25.3
2002.....	0.2	0.5	1.2	1.5	3.1	3.7	5.6	9.2	13.0	22.7
2001.....	*	0.5	1.2	2.1	3.1	3.7	6.2	8.1	11.3	25.9
2000.....	0.2	0.5	1.0	1.8	2.3	4.4	6.0	8.6	10.7	25.9
1999.....	0.3	0.7	1.0	1.7	3.3	3.8	6.2	7.3	13.3	25.3
White, female										
2006 4/.....	*	*	0.3	0.4	0.8	0.8	1.6	2.4	3.2	9.2
2005 4/.....	*	*	0.3	0.3	0.4	1.3	1.3	2.4	4.0	7.7
2004 4/.....	*	*	0.3	0.4	0.6	0.8	1.6	2.3	3.2	8.5
2003 4/.....	*	*	0.3	0.4	0.6	1.0	1.7	2.4	3.5	9.3
2002.....	*	*	*	0.5	0.6	0.9	1.7	2.7	4.1	8.7
2001.....	*	*	0.3	0.6	0.8	1.4	1.6	3.0	3.9	9.5
2000.....	*	*	0.3	0.4	0.8	1.2	1.5	2.3	2.8	9.2
1999.....	*	*	0.3	0.4	0.7	1.2	1.5	2.5	4.2	9.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

All races 3/, both sexes

2006.....	1.0	1.3	2.0	3.0	4.9	7.4	10.7	14.2	18.3	15.8
2005.....	0.8	1.4	2.1	3.2	5.0	7.5	10.5	14.7	17.9	16.1
2004.....	0.9	1.3	1.9	3.3	5.2	7.6	10.0	15.0	17.7	15.7
2003.....	0.8	1.3	2.1	3.1	5.5	7.6	10.1	14.8	16.9	14.9
2002.....	0.9	1.2	2.0	3.4	5.5	7.4	10.9	13.8	16.3	15.7
2001.....	0.9	1.4	2.3	3.4	5.3	7.7	10.6	14.7	15.7	14.9
2000.....	0.9	1.4	2.2	3.7	5.3	8.1	11.2	14.4	16.2	15.7
1999.....	1.0	1.4	2.3	3.5	5.6	7.7	11.0	14.0	15.9	14.8

All races 3/, male

2006.....	1.1	1.4	2.0	3.2	5.8	9.7	13.9	21.0	28.7	25.9
2005.....	0.9	1.4	2.0	3.5	6.1	9.5	14.2	20.9	28.3	26.3
2004.....	0.9	1.3	1.9	3.4	6.7	9.6	13.3	22.6	27.8	27.9
2003.....	0.9	1.5	2.2	3.3	6.3	9.7	13.9	22.5	27.1	25.1
2002.....	1.1	1.3	2.0	3.7	6.7	9.4	15.6	20.8	25.1	28.1
2001.....	1.0	1.5	2.6	3.6	6.3	10.0	15.3	21.3	24.9	25.9
2000.....	0.9	1.5	2.2	4.2	6.6	10.8	15.2	21.9	26.2	28.4
1999.....	1.3	1.4	2.5	4.3	6.9	10.4	15.6	20.8	25.7	24.6

All races 3/, female

2006.....	1.0	1.2	2.1	2.9	4.1	5.4	8.2	9.2	11.8	11.0
2005.....	0.7	1.3	2.2	3.0	3.9	5.7	7.5	10.3	11.5	11.4
2004.....	1.0	1.3	1.9	3.3	3.9	5.9	7.3	9.5	11.5	10.2
2003.....	0.8	1.2	2.0	3.0	4.7	5.8	7.1	9.3	10.6	10.3
2002.....	0.8	1.1	1.9	3.1	4.4	5.7	7.1	8.9	11.0	10.4
2001.....	0.9	1.2	2.0	3.2	4.3	5.7	6.8	10.1	10.2	10.3
2000.....	0.9	1.3	2.2	3.2	4.1	5.8	8.0	9.1	10.3	10.5
1999.....	0.7	1.5	2.0	2.7	4.4	5.3	7.4	9.3	10.1	10.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of mesothelial and soft tissue (C45-C49)										
White, both sexes										
2006 4/.....	1.0	1.3	2.1	3.1	5.1	7.9	11.3	14.8	19.5	16.6
2005 4/.....	0.8	1.3	2.0	3.3	5.1	7.7	11.1	15.4	19.0	16.8
2004 4/.....	0.9	1.3	1.9	3.4	5.6	8.1	10.6	15.8	18.5	16.5
2003 4/.....	0.8	1.3	2.1	3.3	5.5	8.0	10.7	15.4	17.9	15.6
2002.....	0.9	1.2	2.0	3.4	5.6	7.7	11.4	14.6	17.0	16.5
2001.....	0.9	1.3	2.3	3.4	5.6	8.3	11.1	15.4	16.5	15.4
2000.....	0.9	1.4	2.2	3.8	5.5	8.3	11.7	15.1	17.1	16.2
1999.....	0.9	1.5	2.3	3.6	5.7	8.1	11.4	14.4	16.3	15.3
White, male										
2006 4/.....	1.1	1.4	2.0	3.3	6.1	10.5	14.9	21.8	30.4	27.5
2005 4/.....	0.9	1.4	1.9	3.7	6.3	9.8	15.1	22.3	29.9	27.8
2004 4/.....	0.9	1.3	1.9	3.5	7.3	10.3	14.2	23.9	29.2	29.5
2003 4/.....	0.8	1.4	2.2	3.5	6.4	10.2	14.8	23.6	29.2	26.1
2002.....	1.0	1.3	2.1	3.8	7.0	10.0	16.4	22.0	26.5	29.8
2001.....	0.9	1.4	2.7	3.6	6.8	10.8	16.3	22.4	26.2	27.0
2000.....	0.9	1.6	2.3	4.5	6.9	11.1	16.0	23.1	27.6	29.5
1999.....	1.2	1.4	2.6	4.6	7.1	11.0	16.2	21.6	26.9	26.2
White, female										
2006 4/.....	1.0	1.1	2.1	2.9	4.2	5.5	8.4	9.6	12.6	11.5
2005 4/.....	0.7	1.3	2.1	3.0	4.0	5.9	7.8	10.3	12.2	11.7
2004 4/.....	0.9	1.3	1.9	3.3	4.1	6.1	7.7	9.9	11.9	10.6
2003 4/.....	0.8	1.2	2.0	3.1	4.7	6.1	7.4	9.5	10.9	10.9
2002.....	0.8	1.1	1.9	3.1	4.4	5.8	7.3	9.3	11.2	10.7
2001.....	0.8	1.2	1.9	3.2	4.5	6.0	6.9	10.4	10.7	10.5
2000.....	0.9	1.3	2.0	3.2	4.2	5.9	8.3	9.5	10.9	10.7
1999.....	0.6	1.5	2.0	2.7	4.4	5.5	7.7	9.4	10.1	10.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of mesothelial and soft tissue (C45-C49)

Black, both sexes

2006 4/.....	1.1	1.7	2.1	3.2	4.4	5.5	8.8	9.2	9.7	9.3
2005 4/.....	0.9	1.7	2.4	3.3	4.5	6.6	6.7	10.9	9.2	10.4
2004 4/.....	1.2	1.5	2.8	3.9	2.9	6.0	6.0	9.0	12.6	9.3
2003 4/.....	1.2	2.0	2.5	2.5	5.8	5.9	6.9	11.2	8.4	9.9
2002.....	1.3	1.7	2.0	3.8	4.7	6.6	7.5	9.1	11.8	7.7
2001.....	1.3	1.7	2.9	3.9	3.4	4.5	7.5	8.7	8.1	11.1
2000.....	1.4	1.9	2.6	3.8	4.8	7.7	8.8	8.5	7.9	10.9
1999.....	1.5	1.5	2.4	2.9	6.1	5.1	8.4	11.4	11.6	10.5

Black, male

2006 4/.....	*	*	1.9	2.8	4.9	5.4	8.6	14.8	16.4	*
2005 4/.....	*	1.9	2.6	2.6	4.9	8.6	8.2	9.4	14.7	*
2004 4/.....	*	*	2.7	3.8	*	6.5	7.8	12.7	18.7	*
2003 4/.....	1.5	2.9	2.8	2.9	6.9	6.4	7.6	15.5	*	*
2002.....	1.7	2.3	*	4.0	4.8	6.7	9.7	11.9	*	*
2001.....	1.5	*	2.9	4.4	*	5.1	8.1	10.7	*	*
2000.....	*	1.8	*	4.0	6.2	9.4	11.0	12.3	*	*
1999.....	1.9	*	2.5	3.6	7.2	6.0	10.5	14.7	*	*

Black, female

2006 4/.....	*	2.1	2.3	3.5	4.0	5.5	8.9	5.9	*	*
2005 4/.....	*	1.6	2.3	3.9	4.1	5.1	5.6	11.7	*	9.4
2004 4/.....	1.4	1.5	2.8	3.9	*	5.6	4.8	6.7	9.4	*
2003 4/.....	*	*	2.3	*	5.0	5.5	6.4	8.7	8.5	*
2002.....	*	*	2.1	3.5	4.6	6.5	6.0	7.4	10.3	*
2001.....	*	1.8	2.9	3.6	3.5	4.1	7.2	7.5	*	8.8
2000.....	1.3	1.9	3.7	3.6	3.8	6.5	7.3	6.3	*	8.6
1999.....	*	1.6	2.3	*	5.2	4.4	7.0	9.4	10.4	10.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mesothelioma (C45)										
All races 3/, both sexes										
2006.....	0.1	0.2	0.3	0.7	1.6	2.9	4.7	6.4	8.3	6.0
2005.....	0.1	0.2	0.4	0.9	2.0	3.2	4.4	6.8	8.2	6.4
2004.....	0.1	0.2	0.4	0.8	2.0	2.9	4.2	7.6	8.4	5.7
2003.....	0.1	0.2	0.5	0.8	1.9	3.2	4.4	7.1	8.1	5.7
2002.....	0.1	0.1	0.4	1.0	1.9	3.0	5.1	7.0	7.4	5.9
2001.....	0.1	0.2	0.5	0.9	1.9	3.1	4.8	7.2	6.9	5.5
2000.....	0.1	0.2	0.5	1.1	1.9	3.3	5.2	6.6	7.2	5.7
1999.....	*	0.2	0.5	1.2	2.0	3.4	5.1	6.6	7.1	4.8
All races 3/, male										
2006.....	*	0.2	0.4	1.0	2.8	5.1	8.6	13.1	18.0	13.9
2005.....	*	0.2	0.5	1.3	3.5	5.5	8.1	13.4	17.2	15.3
2004.....	*	0.2	0.6	1.2	3.5	5.1	7.6	15.5	17.4	14.4
2003.....	*	0.4	0.7	1.3	3.4	5.6	8.3	14.4	17.1	14.2
2002.....	*	*	0.6	1.6	3.4	5.6	9.8	14.0	15.6	15.9
2001.....	*	0.3	0.8	1.5	3.2	5.6	9.1	14.1	14.7	14.5
2000.....	*	0.2	0.7	1.9	3.4	5.9	9.8	13.4	15.5	15.5
1999.....	*	0.4	0.8	1.9	3.3	6.1	9.6	12.9	15.8	12.7
All races 3/, female										
2006.....	*	*	0.2	0.3	0.5	1.0	1.5	1.5	2.2	2.4
2005.....	*	*	0.3	0.5	0.6	1.2	1.4	1.9	2.6	2.3
2004.....	*	*	0.2	0.4	0.6	1.0	1.5	1.9	2.9	1.8
2003.....	*	*	0.2	0.4	0.6	1.1	1.2	1.9	2.6	1.9
2002.....	*	*	*	0.4	0.5	0.7	1.3	2.0	2.3	1.6
2001.....	*	*	0.3	0.4	0.7	0.9	1.4	2.4	2.1	1.7
2000.....	*	*	0.2	0.4	0.6	1.1	1.5	1.9	2.3	1.8
1999.....	*	*	0.3	0.5	0.7	1.0	1.6	2.2	2.1	1.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mesothelioma (C45)										
White, both sexes										
2006 4/.....	0.2	0.2	0.3	0.7	1.7	3.2	5.1	6.9	8.9	6.5
2005 4/.....	*	0.2	0.4	1.0	2.1	3.5	4.9	7.4	8.8	6.8
2004 4/.....	0.1	0.2	0.4	0.9	2.3	3.2	4.6	8.3	9.0	6.1
2003 4/.....	0.1	0.2	0.5	1.0	2.1	3.5	4.8	7.6	8.7	6.0
2002.....	0.1	0.2	0.4	1.1	2.1	3.3	5.5	7.5	7.9	6.4
2001.....	*	0.2	0.5	1.0	2.2	3.5	5.3	7.9	7.3	5.9
2000.....	*	0.2	0.5	1.2	2.1	3.5	5.6	7.1	7.8	6.1
1999.....	*	0.3	0.5	1.3	2.1	3.7	5.5	7.0	7.5	5.0
White, male										
2006 4/.....	*	0.2	0.4	1.1	3.1	5.7	9.3	13.9	19.1	14.7
2005 4/.....	*	0.2	0.5	1.4	3.7	5.9	8.8	14.6	18.4	16.4
2004 4/.....	*	0.2	0.7	1.4	3.9	5.6	8.3	16.7	18.2	15.6
2003 4/.....	*	0.3	0.7	1.5	3.6	6.1	9.0	15.3	18.5	14.9
2002.....	*	*	0.6	1.7	3.7	6.1	10.5	14.8	16.7	17.3
2001.....	*	0.3	0.8	1.6	3.6	6.3	9.9	15.3	15.6	15.6
2000.....	*	*	0.7	2.0	3.6	6.2	10.5	14.4	16.6	16.6
1999.....	*	0.4	0.8	2.1	3.5	6.7	10.3	13.6	16.7	13.6
White, female										
2006 4/.....	*	*	*	0.4	0.5	1.1	1.7	1.6	2.5	2.6
2005 4/.....	*	*	0.3	0.5	0.7	1.3	1.6	2.1	2.8	2.4
2004 4/.....	*	*	*	0.4	0.7	1.1	1.6	2.1	3.2	1.9
2003 4/.....	*	*	*	0.5	0.6	1.2	1.4	2.1	2.7	2.1
2002.....	*	*	*	0.5	0.5	0.8	1.4	2.2	2.5	1.7
2001.....	*	*	0.3	0.4	0.8	1.0	1.5	2.7	2.3	1.9
2000.....	*	*	0.3	0.4	0.6	1.2	1.6	2.0	2.6	1.8
1999.....	*	*	0.3	0.6	0.8	1.1	1.7	2.3	2.0	1.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other mesothelial and soft tissue (C47-C49)										
All races 3/, both sexes										
2006.....	0.9	1.1	1.7	2.4	3.3	4.5	6.0	7.7	9.9	9.4
2005.....	0.7	1.2	1.7	2.3	3.0	4.3	6.1	7.9	9.6	9.4
2004.....	0.8	1.1	1.5	2.5	3.2	4.7	5.8	7.3	9.2	9.6
2003.....	0.7	1.1	1.7	2.3	3.5	4.4	5.7	7.6	8.7	8.8
2002.....	0.8	1.1	1.6	2.4	3.6	4.4	5.8	6.8	8.8	9.3
2001.....	0.8	1.1	1.8	2.4	3.4	4.6	5.8	7.4	8.7	9.1
2000.....	0.8	1.2	1.7	2.6	3.4	4.8	5.9	7.7	8.8	9.6
1999.....	0.9	1.2	1.8	2.3	3.6	4.3	5.9	7.4	8.6	9.6
All races 3/, male										
2006.....	0.9	1.1	1.6	2.2	2.9	4.6	5.2	7.7	10.5	11.5
2005.....	0.8	1.2	1.6	2.2	2.7	4.0	6.1	7.4	10.7	10.5
2004.....	0.7	1.0	1.3	2.1	3.2	4.5	5.7	7.0	10.2	12.7
2003.....	0.7	1.1	1.5	2.0	2.9	4.0	5.5	7.9	9.9	10.7
2002.....	0.9	1.1	1.4	2.1	3.3	3.8	5.7	6.7	9.3	11.7
2001.....	0.9	1.1	1.8	2.1	3.1	4.4	6.1	7.1	10.0	10.7
2000.....	0.8	1.3	1.5	2.3	3.3	4.9	5.2	8.4	10.4	12.5
1999.....	1.1	1.0	1.7	2.4	3.6	4.4	5.9	7.8	9.6	11.3
All races 3/, female										
2006.....	0.8	1.1	1.9	2.6	3.6	4.4	6.6	7.8	9.6	8.5
2005.....	0.7	1.1	1.9	2.5	3.3	4.5	6.1	8.3	8.9	8.9
2004.....	0.9	1.1	1.7	2.9	3.3	4.9	5.8	7.6	8.5	8.2
2003.....	0.7	1.0	1.8	2.6	4.1	4.7	5.9	7.3	8.0	8.0
2002.....	0.8	1.0	1.7	2.7	3.8	4.9	5.8	6.8	8.6	8.3
2001.....	0.8	1.1	1.8	2.8	3.6	4.8	5.4	7.6	8.0	8.4
2000.....	0.8	1.1	1.9	2.8	3.5	4.7	6.5	7.1	7.8	8.4
1999.....	0.7	1.3	1.8	2.2	3.7	4.3	5.8	7.1	8.0	8.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other mesothelial and soft tissue (C47-C49)										
White, both sexes										
2006 4/.....	0.9	1.1	1.8	2.4	3.4	4.6	6.2	7.9	10.5	9.8
2005 4/.....	0.7	1.1	1.6	2.4	3.0	4.2	6.2	8.0	10.0	9.6
2004 4/.....	0.8	1.1	1.5	2.5	3.4	4.9	6.0	7.6	9.5	9.9
2003 4/.....	0.7	1.0	1.6	2.3	3.4	4.5	5.9	7.6	9.1	9.1
2002.....	0.8	1.0	1.6	2.4	3.6	4.4	5.9	7.1	9.0	9.6
2001.....	0.8	1.1	1.7	2.4	3.4	4.8	5.8	7.4	9.1	9.1
2000.....	0.8	1.2	1.7	2.6	3.4	4.8	6.1	7.9	9.1	9.7
1999.....	0.8	1.2	1.8	2.3	3.6	4.4	5.9	7.4	8.7	9.8
White, male										
2006 4/.....	0.9	1.2	1.6	2.3	3.0	4.8	5.6	7.8	11.0	12.2
2005 4/.....	0.8	1.1	1.5	2.3	2.7	3.9	6.3	7.7	11.0	10.9
2004 4/.....	0.7	1.0	1.3	2.1	3.4	4.6	5.9	7.2	10.8	13.0
2003 4/.....	0.6	1.0	1.5	2.0	2.8	4.0	5.7	8.1	10.5	10.9
2002.....	0.8	1.1	1.5	2.1	3.3	3.8	5.9	7.1	9.6	12.0
2001.....	0.8	1.0	1.8	2.0	3.2	4.5	6.4	7.1	10.4	10.6
2000.....	0.8	1.3	1.6	2.4	3.2	4.9	5.4	8.6	10.8	12.4
1999.....	1.1	1.0	1.8	2.4	3.6	4.3	5.9	7.9	9.8	12.0
White, female										
2006 4/.....	0.9	1.0	1.9	2.5	3.8	4.4	6.7	8.1	10.1	8.7
2005 4/.....	0.7	1.1	1.8	2.4	3.3	4.5	6.2	8.2	9.3	9.1
2004 4/.....	0.8	1.1	1.6	2.9	3.4	5.1	6.1	7.8	8.7	8.5
2003 4/.....	0.7	1.0	1.8	2.6	4.1	4.9	6.0	7.3	8.2	8.3
2002.....	0.7	1.0	1.8	2.6	3.8	4.9	5.9	7.1	8.6	8.5
2001.....	0.7	1.1	1.7	2.8	3.7	5.1	5.4	7.7	8.3	8.4
2000.....	0.8	1.1	1.7	2.8	3.6	4.7	6.6	7.4	8.2	8.5
1999.....	0.5	1.3	1.7	2.1	3.6	4.4	6.0	7.1	8.0	8.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of other mesothelial and soft tissue (C47-C49)										
Black, both sexes										
2006 4/.....	1.1	1.6	1.8	2.7	3.3	4.2	5.8	6.2	5.2	6.0
2005 4/.....	0.9	1.7	2.0	2.6	3.4	4.9	5.1	8.5	5.8	7.5
2004 4/.....	1.1	1.3	2.4	3.5	2.1	5.0	4.4	6.8	8.4	7.0
2003 4/.....	1.2	1.7	2.1	2.3	4.4	4.6	5.6	8.2	5.6	7.0
2002.....	1.2	1.6	1.7	3.2	4.1	5.3	5.5	5.7	8.9	5.9
2001.....	1.2	1.4	2.5	3.5	3.0	3.8	6.1	7.3	6.0	9.8
2000.....	1.2	1.6	2.4	2.9	3.7	5.2	5.9	6.0	*	9.1
1999.....	1.3	1.3	1.9	2.4	4.8	4.0	6.6	8.0	7.5	8.3
Black, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	1.8	2.0	*	*	5.2	*	*	*	*
2004 4/.....	*	*	2.1	3.2	*	5.1	*	*	*	*
2003 4/.....	*	2.2	*	*	4.2	*	*	*	*	*
2002.....	1.5	2.1	*	*	*	*	*	*	*	*
2001.....	*	*	2.3	3.4	*	*	*	*	*	*
2000.....	*	*	*	*	4.1	*	*	*	*	*
1999.....	1.5	*	*	*	4.5	*	7.1	*	*	*
Black, female										
2006 4/.....	*	1.9	1.9	3.3	3.3	4.7	7.6	5.4	*	*
2005 4/.....	*	1.6	2.1	3.4	4.0	4.6	5.4	10.7	*	7.5
2004 4/.....	1.3	1.4	2.7	3.7	*	4.9	4.4	6.7	9.4	*
2003 4/.....	*	*	2.2	*	4.5	5.1	6.4	8.4	*	*
2002.....	*	*	2.0	3.4	4.3	6.3	5.4	6.3	9.5	*
2001.....	*	1.7	2.7	3.5	3.5	3.8	6.7	7.2	*	8.8
2000.....	*	1.7	3.7	3.4	3.4	5.6	6.7	*	*	*
1999.....	*	*	2.2	*	5.0	*	6.3	8.0	*	9.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)										
All races 3/, both sexes										
2006.....	6.9	11.6	17.0	24.7	32.7	38.0	47.1	63.1	82.7	116.7
2005.....	7.4	11.8	17.9	25.0	33.1	38.7	49.3	63.5	82.9	122.7
2004.....	7.3	11.7	18.7	26.1	33.1	40.8	49.6	64.7	82.4	124.2
2003.....	7.9	12.6	19.1	26.6	33.5	39.9	51.8	66.0	87.3	132.5
2002.....	7.9	12.9	19.7	26.2	33.7	41.3	52.4	67.1	89.8	134.6
2001.....	8.2	13.4	20.6	27.1	34.1	41.7	54.0	69.1	88.0	134.2
2000.....	8.3	13.6	20.8	28.1	35.0	43.4	55.2	70.9	90.9	147.4
1999.....	7.9	13.3	21.0	28.1	34.8	43.5	55.9	70.0	92.4	144.4
All races 3/, male										
2006.....	*	*	0.3	0.4	0.8	0.8	1.2	1.8	2.6	3.0
2005.....	*	*	0.3	0.4	0.7	1.0	1.7	1.8	2.5	2.4
2004.....	*	*	0.2	0.4	0.6	0.9	1.3	2.1	2.1	3.3
2003.....	*	*	0.4	0.4	0.7	0.9	1.2	1.5	2.8	3.9
2002.....	*	*	0.3	0.4	0.7	0.8	1.3	1.9	2.7	3.4
2001.....	*	*	0.4	0.5	0.8	1.2	1.7	2.3	2.4	3.5
2000.....	*	*	0.3	0.7	0.8	1.1	1.7	2.1	2.4	4.2
1999.....	*	*	0.4	0.6	0.7	1.1	1.4	1.9	2.3	4.5
All races 3/, female										
2006.....	13.8	22.7	33.0	47.5	61.9	70.4	84.4	108.0	132.9	169.9
2005.....	14.7	23.2	34.9	48.3	62.6	71.6	87.9	108.3	132.8	177.9
2004.....	14.5	23.0	36.3	50.4	62.6	75.6	88.5	109.6	131.8	178.6
2003.....	15.6	24.7	37.0	51.3	63.4	73.7	92.4	111.9	139.0	189.4
2002.....	15.6	25.3	38.3	50.4	63.7	76.2	93.2	113.2	142.7	191.5
2001.....	16.2	26.3	40.1	52.0	64.3	76.5	95.6	115.9	139.4	188.9
2000.....	16.4	26.6	40.4	53.8	66.0	79.6	97.3	118.8	143.1	205.7
1999.....	15.6	26.1	40.8	53.9	65.6	79.7	98.4	117.4	145.1	200.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)										
White, both sexes										
2006 4/.....	6.3	10.4	15.7	23.3	31.9	37.7	46.9	63.3	83.0	116.9
2005 4/.....	6.6	10.4	16.4	23.4	32.0	38.1	49.2	63.6	83.6	123.4
2004 4/.....	6.5	10.5	17.1	25.1	32.0	40.6	49.5	65.0	83.4	124.8
2003 4/.....	7.1	11.4	17.8	25.3	32.6	39.2	51.7	66.0	87.5	132.5
2002.....	7.0	11.7	18.5	25.0	33.1	40.8	52.0	67.5	89.0	135.3
2001.....	7.3	12.4	19.4	25.8	33.3	40.9	53.6	69.4	88.2	134.8
2000.....	7.5	12.4	19.7	26.9	34.4	43.3	55.5	71.1	91.7	147.3
1999.....	7.1	12.1	19.8	26.8	33.8	42.8	55.5	69.7	92.8	146.0
White, male										
2006 4/.....	*	*	*	0.4	0.8	0.8	1.2	1.8	2.5	3.2
2005 4/.....	*	*	0.3	0.4	0.6	0.9	1.6	1.6	2.7	2.4
2004 4/.....	*	*	*	0.3	0.6	0.9	1.2	2.2	2.0	3.0
2003 4/.....	*	*	0.4	0.4	0.7	0.9	1.2	1.5	2.9	3.9
2002.....	*	*	0.3	0.4	0.6	0.6	1.3	1.7	2.7	3.3
2001.....	*	*	0.3	0.5	0.7	1.2	1.6	2.1	2.3	3.3
2000.....	*	*	0.3	0.6	0.7	1.1	1.7	2.2	2.3	3.9
1999.....	*	*	0.3	0.5	0.6	1.0	1.3	1.8	2.2	4.4
White, female										
2006 4/.....	12.6	20.6	31.0	45.4	60.8	70.5	84.8	109.2	133.8	170.3
2005 4/.....	13.3	20.7	32.3	45.7	61.1	71.1	88.6	109.5	134.4	179.1
2004 4/.....	12.9	20.9	33.7	48.9	61.1	75.9	89.2	111.0	134.0	179.7
2003 4/.....	14.2	22.7	34.8	49.2	62.0	72.9	93.0	112.7	139.7	189.4
2002.....	13.9	23.2	36.3	48.6	63.1	76.2	93.3	114.7	142.0	192.6
2001.....	14.7	24.5	38.1	50.1	63.4	75.7	95.8	117.0	140.0	189.9
2000.....	15.0	24.4	38.7	51.8	65.4	80.2	98.6	119.7	144.9	205.5
1999.....	14.2	23.9	38.9	51.7	64.4	79.2	98.6	117.4	146.3	202.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasm of breast (C50)										
Black, both sexes										
2006 4/.....	12.5	21.3	27.7	39.5	47.4	49.8	59.1	78.1	104.6	142.2
2005 4/.....	13.6	23.1	31.8	43.6	47.7	53.8	61.5	75.5	91.5	144.3
2004 4/.....	13.7	21.2	31.8	40.0	48.9	51.2	62.8	77.0	90.1	143.5
2003 4/.....	14.1	22.6	31.2	43.4	48.1	54.6	65.1	80.3	106.4	161.6
2002.....	15.3	22.9	32.1	40.5	46.3	54.5	67.2	78.5	117.5	152.6
2001.....	14.8	22.6	33.1	42.4	47.4	58.0	70.4	83.5	105.9	152.0
2000.....	14.8	23.9	32.8	43.9	47.3	52.7	65.6	84.2	100.5	176.4
1999.....	14.6	24.2	34.7	44.4	48.6	57.8	69.9	88.0	104.9	147.3
Black, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	23.5	39.6	50.4	71.3	83.6	85.5	98.3	123.7	157.7	199.2
2005 4/.....	25.4	43.0	58.4	79.1	83.9	91.8	101.5	117.3	139.5	201.8
2004 4/.....	25.6	39.5	58.1	72.4	85.8	88.1	103.7	121.6	135.7	196.9
2003 4/.....	26.3	42.0	57.3	78.3	84.7	94.1	107.4	127.2	160.5	222.4
2002.....	28.7	42.4	58.7	73.1	81.2	92.9	111.1	123.2	176.7	209.1
2001.....	27.5	41.9	60.5	76.7	83.7	99.5	115.0	130.9	157.8	205.9
2000.....	27.7	44.3	60.4	78.8	83.3	91.0	107.4	133.6	149.0	238.7
1999.....	27.5	44.9	63.3	79.6	85.2	99.5	114.1	139.3	154.7	199.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of female genital organs (C51-C58)

All races 3/, both sexes

2006.....	3.5	6.0	9.7	14.6	22.2	29.6	38.9	49.1	60.6	72.7
2005.....	3.6	6.1	9.8	15.3	21.8	29.6	38.1	46.9	61.5	72.7
2004.....	3.8	6.2	9.4	14.5	21.5	30.9	39.4	48.8	62.9	72.7
2003.....	3.7	5.9	9.9	15.2	22.9	30.9	38.2	49.9	63.8	75.2
2002.....	3.7	6.4	10.3	15.1	22.5	30.3	39.9	52.0	63.6	77.5
2001.....	3.8	6.4	10.6	14.7	22.8	30.1	39.6	52.1	66.9	79.9
2000.....	3.8	6.5	10.4	15.5	22.2	31.0	38.9	52.5	65.4	82.3
1999.....	3.9	6.6	10.3	15.1	22.7	30.8	39.4	51.7	64.6	79.1

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	7.0	11.8	19.1	28.5	42.4	55.5	70.5	85.0	98.5	106.7
2005.....	7.1	12.1	19.2	29.7	41.7	55.3	68.9	80.9	99.6	106.1
2004.....	7.5	12.1	18.3	28.3	41.0	57.8	71.1	83.8	101.6	105.4
2003.....	7.3	11.7	19.4	29.5	43.7	57.6	68.8	85.4	102.8	108.5
2002.....	7.3	12.6	20.2	29.2	42.9	56.4	71.8	88.8	102.3	111.2
2001.....	7.5	12.6	20.7	28.4	43.4	56.0	71.0	88.6	107.0	113.4
2000.....	7.5	12.7	20.3	30.1	42.4	57.5	69.6	89.1	103.9	115.8
1999.....	7.8	13.0	20.3	29.3	43.2	57.2	70.3	87.6	102.4	110.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of female genital organs (C51-C58)										
White, both sexes										
2006 4/.....	3.4	5.8	9.6	14.5	21.7	28.6	38.3	48.5	60.7	73.0
2005 4/.....	3.4	5.9	9.8	15.0	21.1	28.9	37.3	46.2	61.5	73.3
2004 4/.....	3.6	6.0	9.0	14.3	21.1	29.7	38.1	48.6	63.5	73.0
2003 4/.....	3.6	5.8	9.7	15.2	22.2	30.0	37.4	49.2	63.6	75.2
2002.....	3.6	6.2	10.3	14.9	22.1	29.2	38.9	51.3	63.3	76.9
2001.....	3.7	6.3	10.4	14.5	21.9	29.0	38.9	51.6	66.9	79.7
2000.....	3.6	6.4	10.2	15.4	21.7	29.8	37.7	52.0	65.0	82.1
1999.....	3.8	6.7	10.3	14.9	22.2	29.7	38.3	50.7	64.5	78.8
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	6.8	11.6	19.0	28.5	41.9	54.1	70.1	84.7	99.1	107.2
2005 4/.....	6.8	11.8	19.3	29.4	40.8	54.6	68.1	80.4	100.0	107.1
2004 4/.....	7.3	12.0	17.9	28.0	40.6	56.1	69.5	84.1	103.0	105.9
2003 4/.....	7.2	11.5	19.2	29.8	42.8	56.6	67.9	84.7	102.9	108.6
2002.....	7.2	12.3	20.3	29.1	42.5	55.0	70.7	88.1	102.1	110.3
2001.....	7.4	12.6	20.6	28.4	42.2	54.4	70.5	88.2	107.3	113.0
2000.....	7.1	12.7	20.1	30.1	41.7	55.8	68.0	88.6	103.7	115.5
1999.....	7.7	13.2	20.3	29.0	42.7	55.5	68.8	86.3	102.6	110.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of female genital organs (C51-C58)

Black, both sexes

2006 4/.....	5.2	7.6	11.1	17.7	31.0	44.8	52.1	63.7	69.8	83.8
2005 4/.....	5.3	7.9	10.7	18.9	31.9	41.3	52.9	63.3	74.0	81.8
2004 4/.....	4.9	7.9	12.4	18.3	29.2	46.9	59.0	62.5	68.8	80.9
2003 4/.....	4.5	7.7	12.3	17.9	32.1	43.7	52.9	65.2	76.4	89.3
2002.....	4.9	8.0	12.1	18.9	29.4	44.0	57.2	66.8	77.4	98.3
2001.....	5.2	7.0	12.4	17.9	31.8	44.6	55.0	67.0	78.0	94.9
2000.....	5.6	7.3	12.4	18.5	30.2	47.0	57.9	68.9	79.6	96.7
1999.....	5.1	6.5	11.9	20.0	31.3	46.0	60.9	70.7	74.0	91.5

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	9.7	14.1	20.5	32.3	55.2	77.9	87.9	101.8	106.7	118.1
2005 4/.....	10.1	14.8	19.8	34.5	56.8	71.9	89.2	101.1	113.1	114.8
2004 4/.....	9.1	14.8	22.9	33.5	52.0	81.7	99.1	99.5	105.1	113.0
2003 4/.....	8.5	14.5	22.6	32.7	57.2	76.2	88.6	103.9	116.4	124.0
2002.....	9.2	14.9	22.4	34.3	52.3	76.6	95.7	106.8	117.8	136.0
2001.....	9.8	13.1	23.0	32.6	56.8	77.5	91.7	107.1	117.7	130.1
2000.....	10.5	13.7	23.0	33.7	54.1	81.6	96.3	110.5	119.7	132.6
1999.....	9.6	12.0	21.9	36.4	55.9	80.0	101.3	113.3	110.6	125.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of vulva and vagina (C51-C52)										
All races 3/, both sexes										
2006.....	*	0.2	0.3	0.5	0.7	1.0	1.3	2.2	3.6	6.9
2005.....	0.1	0.2	0.3	0.4	0.5	0.9	1.3	2.0	4.1	6.6
2004.....	0.1	0.2	0.2	0.4	0.7	0.9	1.6	2.1	3.9	6.9
2003.....	*	0.2	0.2	0.3	0.6	1.0	1.4	2.4	3.6	7.2
2002.....	0.1	0.1	0.3	0.3	0.6	0.7	1.5	2.5	3.5	7.7
2001.....	0.1	0.2	0.3	0.3	0.5	0.9	1.4	2.5	3.9	7.3
2000.....	0.1	0.1	0.3	0.3	0.6	1.0	1.5	2.5	3.6	7.7
1999.....	0.1	0.1	0.3	0.3	0.6	0.9	1.5	2.5	3.9	8.6
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	*	0.3	0.5	1.0	1.2	2.0	2.3	3.9	5.9	10.1
2005.....	0.3	0.3	0.6	0.8	0.9	1.7	2.4	3.4	6.6	9.6
2004.....	0.3	0.4	0.5	0.8	1.3	1.7	2.9	3.7	6.3	10.1
2003.....	*	0.3	0.5	0.7	1.1	1.8	2.5	4.0	5.7	10.4
2002.....	0.2	0.3	0.5	0.7	1.2	1.3	2.7	4.3	5.7	11.0
2001.....	0.2	0.3	0.6	0.6	1.0	1.6	2.6	4.2	6.3	10.3
2000.....	0.2	0.3	0.6	0.6	1.1	1.9	2.8	4.3	5.7	10.9
1999.....	0.2	0.3	0.6	0.7	1.2	1.6	2.6	4.2	6.1	12.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of vulva and vagina (C51-C52)										
White, both sexes										
2006 4/.....	*	0.2	0.3	0.5	0.7	1.1	1.4	2.4	3.6	7.2
2005 4/.....	0.1	0.2	0.3	0.4	0.5	0.9	1.4	2.1	4.3	6.8
2004 4/.....	0.1	0.2	0.2	0.3	0.7	0.9	1.6	2.3	4.0	7.1
2003 4/.....	*	0.2	0.2	0.3	0.6	1.0	1.4	2.5	3.7	7.5
2002.....	0.1	0.1	0.3	0.4	0.7	0.7	1.5	2.6	3.6	8.0
2001.....	*	0.1	0.3	0.3	0.4	0.9	1.4	2.5	4.1	7.4
2000.....	*	0.1	0.3	0.3	0.6	1.1	1.6	2.5	3.7	7.9
1999.....	*	0.1	0.3	0.3	0.6	0.9	1.5	2.5	3.9	9.0
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	*	0.3	0.5	1.0	1.3	2.1	2.5	4.1	5.9	10.6
2005 4/.....	0.2	0.3	0.5	0.8	0.9	1.8	2.5	3.6	7.0	9.9
2004 4/.....	0.3	0.4	0.5	0.7	1.3	1.7	3.0	3.9	6.4	10.4
2003 4/.....	*	0.4	0.4	0.7	1.1	1.9	2.6	4.3	5.9	10.8
2002.....	0.2	0.3	0.6	0.7	1.3	1.4	2.8	4.4	5.8	11.4
2001.....	*	0.3	0.5	0.6	0.8	1.7	2.6	4.2	6.5	10.4
2000.....	*	0.3	0.6	0.6	1.2	2.0	2.9	4.3	5.9	11.1
1999.....	*	0.3	0.6	0.6	1.1	1.6	2.7	4.2	6.3	12.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of cervix uteri (C53)										
All races 3/, both sexes										
2006.....	1.7	2.1	2.4	2.5	2.7	3.1	3.4	3.6	4.1	5.2
2005.....	1.7	2.2	2.3	2.6	2.6	3.0	3.1	3.5	4.6	5.4
2004.....	1.7	2.1	2.0	2.7	2.9	3.2	3.3	3.3	4.3	5.1
2003.....	1.7	2.0	2.4	2.8	2.9	3.2	3.0	3.7	5.1	5.7
2002.....	1.7	2.1	2.4	2.7	2.8	3.3	3.6	4.1	4.2	6.2
2001.....	1.8	2.3	2.6	2.6	3.2	2.9	3.5	4.0	5.5	7.2
2000.....	1.8	2.2	2.7	3.1	3.1	3.6	3.9	4.3	5.6	7.3
1999.....	1.8	2.4	2.7	3.0	3.5	3.7	3.9	4.7	4.6	7.9
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	3.4	4.1	4.7	4.9	5.2	5.7	6.1	6.2	6.6	7.7
2005.....	3.4	4.3	4.6	5.1	5.0	5.7	5.7	6.0	7.5	7.8
2004.....	3.4	4.1	4.0	5.2	5.6	6.0	6.0	5.6	7.0	7.3
2003.....	3.3	4.0	4.6	5.5	5.5	5.9	5.4	6.3	8.3	8.3
2002.....	3.4	4.1	4.7	5.2	5.3	6.1	6.5	7.0	6.7	8.9
2001.....	3.7	4.5	5.1	5.1	6.0	5.4	6.3	6.8	8.9	10.1
2000.....	3.5	4.4	5.2	6.0	5.9	6.7	6.9	7.3	9.0	10.3
1999.....	3.6	4.6	5.2	5.8	6.7	6.8	6.9	8.0	7.2	11.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of cervix uteri (C53)										
White, both sexes										
2006 4/.....	1.5	1.9	2.2	2.4	2.5	2.6	2.8	3.0	3.7	4.2
2005 4/.....	1.6	1.9	2.2	2.3	2.3	2.7	2.8	3.0	4.0	4.8
2004 4/.....	1.5	1.9	1.8	2.5	2.6	2.7	2.8	2.9	3.8	4.1
2003 4/.....	1.6	1.9	2.1	2.6	2.6	2.8	2.4	2.8	4.6	4.9
2002.....	1.6	1.8	2.2	2.5	2.4	2.7	3.0	3.6	3.5	5.1
2001.....	1.7	2.2	2.3	2.3	2.9	2.4	3.0	3.4	4.7	6.2
2000.....	1.6	2.0	2.4	2.7	2.6	2.9	3.3	3.8	5.0	6.3
1999.....	1.7	2.3	2.5	2.7	3.0	3.1	3.1	4.0	4.1	6.8
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	3.0	3.8	4.4	4.7	4.8	5.0	5.2	5.3	6.0	6.2
2005 4/.....	3.1	3.8	4.3	4.5	4.5	5.1	5.2	5.2	6.6	6.9
2004 4/.....	3.1	3.8	3.5	4.9	5.1	5.1	5.1	5.0	6.1	5.9
2003 4/.....	3.2	3.7	4.2	5.1	4.9	5.2	4.4	4.9	7.4	7.1
2002.....	3.1	3.7	4.3	4.9	4.5	5.2	5.4	6.2	5.7	7.3
2001.....	3.5	4.3	4.6	4.6	5.5	4.4	5.5	5.8	7.6	8.8
2000.....	3.1	4.0	4.7	5.3	5.0	5.5	5.9	6.5	8.0	8.8
1999.....	3.4	4.5	5.0	5.2	5.7	5.9	5.5	6.8	6.5	9.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of cervix uteri (C53)										
Black, both sexes										
2006 4/.....	3.2	3.2	4.0	3.4	4.9	6.7	8.5	8.2	9.0	17.6
2005 4/.....	3.0	4.0	3.3	5.1	5.3	6.5	6.2	9.4	12.4	13.1
2004 4/.....	2.9	3.5	3.9	4.7	6.1	7.1	8.1	8.0	10.1	16.1
2003 4/.....	2.4	3.5	3.8	5.1	6.1	7.1	7.7	11.2	11.4	14.6
2002.....	3.0	3.7	4.6	4.2	6.0	8.2	10.1	9.6	12.6	17.5
2001.....	3.0	3.2	4.4	4.6	5.9	7.9	7.7	9.7	15.2	19.0
2000.....	3.1	3.7	4.9	6.2	6.4	9.4	9.7	10.8	13.6	19.7
1999.....	3.0	3.0	4.1	6.1	8.3	8.7	13.1	12.3	9.5	21.7
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	6.0	5.9	7.4	6.1	8.7	11.6	14.4	13.1	13.7	24.9
2005 4/.....	5.7	7.6	6.2	9.3	9.5	11.3	10.4	14.9	18.9	18.4
2004 4/.....	5.4	6.6	7.2	8.6	10.9	12.4	13.6	12.7	15.5	22.4
2003 4/.....	4.5	6.5	7.1	9.3	10.8	12.4	12.8	17.9	17.4	20.3
2002.....	5.6	7.0	8.4	7.6	10.7	14.2	16.8	15.4	19.1	24.2
2001.....	5.6	6.1	8.2	8.3	10.6	13.7	12.8	15.5	22.9	26.1
2000.....	5.9	7.0	9.1	11.2	11.4	16.3	16.2	17.4	20.4	27.0
1999.....	5.7	5.5	7.5	11.1	14.8	15.2	21.8	19.7	14.3	29.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of corpus uteri and uterus, part unspecified (C54-C55)

All races 3/, both sexes

2006.....	0.4	1.0	1.9	3.8	6.2	8.8	11.6	14.3	17.6	21.6
2005.....	0.4	0.9	1.8	3.7	6.4	8.7	10.8	13.1	17.4	22.1
2004.....	0.5	0.9	1.8	3.4	5.8	8.8	11.5	13.8	17.7	22.3
2003.....	0.5	0.8	1.9	3.5	6.1	9.0	11.3	13.8	17.1	22.8
2002.....	0.4	0.9	2.0	3.5	5.8	8.4	11.8	14.2	17.0	24.6
2001.....	0.4	0.9	2.0	3.2	5.8	8.2	11.4	14.5	19.2	24.7
2000.....	0.4	0.9	1.8	3.5	6.0	8.4	11.1	14.0	18.6	24.7
1999.....	0.5	0.9	1.7	3.3	5.5	8.8	10.9	14.4	19.4	24.0

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	0.9	1.9	3.6	7.4	11.9	16.4	21.0	24.7	28.6	31.6
2005.....	0.8	1.8	3.5	7.2	12.1	16.4	19.6	22.6	28.2	32.2
2004.....	1.0	1.8	3.5	6.7	11.0	16.5	20.8	23.8	28.7	32.3
2003.....	1.0	1.6	3.7	6.8	11.7	16.8	20.4	23.5	27.6	33.0
2002.....	0.9	1.7	3.9	6.9	11.0	15.7	21.2	24.2	27.4	35.2
2001.....	0.8	1.8	4.0	6.2	11.1	15.3	20.4	24.6	30.7	35.0
2000.....	0.9	1.9	3.5	6.9	11.4	15.6	19.8	23.8	29.5	34.8
1999.....	0.9	1.8	3.3	6.3	10.5	16.2	19.5	24.5	30.7	33.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of corpus uteri and uterus, part unspecified (C54-C55)

White, both sexes

2006 4/.....	0.4	0.9	1.8	3.5	5.5	7.6	10.3	13.3	16.9	21.7
2005 4/.....	0.4	0.9	1.7	3.4	5.5	7.9	9.8	12.0	16.9	21.8
2004 4/.....	0.5	0.9	1.6	3.2	5.2	7.7	10.2	13.0	17.4	22.4
2003 4/.....	0.4	0.7	1.9	3.4	5.4	7.9	10.5	13.0	16.2	22.7
2002.....	0.4	0.8	1.9	3.2	5.2	7.2	10.8	13.1	16.3	23.8
2001.....	0.4	0.9	1.9	3.1	5.2	7.2	10.5	13.7	18.6	24.5
2000.....	0.4	0.9	1.8	3.4	5.4	7.3	9.9	13.0	17.5	24.4
1999.....	0.4	0.9	1.6	3.0	5.2	7.6	10.2	13.7	18.8	23.4

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	0.9	1.8	3.6	6.8	10.6	14.3	18.8	23.3	27.6	31.8
2005 4/.....	0.8	1.8	3.4	6.7	10.6	15.0	18.0	20.8	27.5	31.8
2004 4/.....	1.0	1.8	3.2	6.3	10.0	14.5	18.6	22.6	28.2	32.6
2003 4/.....	0.8	1.5	3.7	6.6	10.4	14.9	19.1	22.4	26.2	32.8
2002.....	0.9	1.7	3.7	6.3	10.1	13.6	19.6	22.4	26.3	34.2
2001.....	0.8	1.7	3.8	6.1	9.9	13.6	18.9	23.5	29.8	34.7
2000.....	0.8	1.9	3.5	6.7	10.3	13.6	17.8	22.2	28.0	34.3
1999.....	0.9	1.8	3.2	5.9	9.9	14.2	18.4	23.3	30.0	32.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of corpus uteri and uterus, part unspecified (C54-C55)

Black, both sexes

2006 4/.....	*	1.4	2.1	6.9	13.9	20.8	25.8	27.6	30.8	25.9
2005 4/.....	*	1.3	2.5	6.1	15.1	18.0	22.7	27.6	27.4	31.3
2004 4/.....	0.7	1.0	3.0	5.2	11.9	20.2	26.4	26.6	25.5	26.2
2003 4/.....	0.9	1.1	2.5	5.3	13.9	19.9	22.4	25.5	31.2	30.0
2002.....	*	1.1	2.9	6.6	10.8	20.0	25.2	27.8	29.0	38.9
2001.....	*	1.4	2.9	4.6	11.6	18.8	24.2	26.6	31.1	32.2
2000.....	*	1.0	2.4	5.3	12.0	19.6	26.4	27.7	35.0	33.2
1999.....	0.8	*	2.2	5.8	10.5	20.9	21.8	25.7	29.8	35.4

Black, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

Black, female

2006 4/.....	*	2.7	3.8	12.7	24.8	36.2	43.5	44.2	47.0	36.6
2005 4/.....	*	2.5	4.7	11.2	26.9	31.4	38.3	44.0	41.9	43.9
2004 4/.....	1.4	1.9	5.5	9.4	21.1	35.3	44.4	42.3	38.9	36.6
2003 4/.....	1.6	2.1	4.7	9.6	24.8	34.7	37.4	40.7	47.6	41.7
2002.....	*	2.0	5.4	12.1	19.2	34.8	42.1	44.5	44.2	53.8
2001.....	*	2.6	5.4	8.3	20.8	32.7	40.4	42.4	47.0	44.2
2000.....	*	1.9	4.4	9.6	21.4	34.0	44.0	44.4	52.6	45.5
1999.....	1.6	*	4.1	10.5	18.8	36.4	36.2	41.2	44.5	48.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of ovary (C56)										
All races 3/, both sexes										
2006.....	1.3	2.8	5.1	7.7	12.4	16.3	22.0	28.3	34.8	37.8
2005.....	1.3	2.8	5.3	8.4	12.2	16.5	22.3	28.0	34.7	38.1
2004.....	1.4	2.9	5.2	7.9	11.9	17.8	22.5	28.9	36.4	37.7
2003.....	1.4	2.9	5.2	8.3	13.0	17.3	21.9	29.4	36.9	38.4
2002.....	1.4	3.2	5.5	8.3	12.9	17.3	22.3	30.3	37.8	38.1
2001.....	1.4	2.9	5.5	8.3	13.0	17.6	22.6	30.2	37.4	39.6
2000.....	1.5	3.1	5.5	8.3	12.2	17.5	21.8	30.9	36.5	40.9
1999.....	1.5	3.1	5.5	8.3	12.6	17.0	22.4	29.2	35.5	37.6
All races 3/, male										
2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....
All races 3/, female										
2006.....	2.5	5.5	9.9	14.9	23.7	30.6	39.9	49.1	56.5	55.5
2005.....	2.7	5.5	10.3	16.3	23.4	30.9	40.4	48.3	56.3	55.6
2004.....	2.8	5.8	10.1	15.3	22.7	33.3	40.7	49.7	58.8	54.6
2003.....	2.8	5.7	10.2	16.2	24.8	32.3	39.5	50.4	59.5	55.4
2002.....	2.8	6.4	10.7	16.2	24.6	32.3	40.1	51.7	60.8	54.6
2001.....	2.8	5.8	10.8	16.0	24.7	32.8	40.5	51.4	59.8	56.1
2000.....	2.9	6.0	10.8	16.1	23.3	32.5	39.0	52.5	58.0	57.6
1999.....	3.0	6.2	10.8	16.0	24.0	31.6	39.9	49.5	56.2	52.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of ovary (C56)										
White, both sexes										
2006 4/.....	1.3	2.8	5.2	7.9	12.8	16.9	23.2	29.1	36.0	38.8
2005 4/.....	1.3	2.9	5.5	8.8	12.7	16.9	22.8	28.8	35.5	39.4
2004 4/.....	1.5	3.0	5.3	8.0	12.4	18.2	23.1	29.8	37.8	38.6
2003 4/.....	1.4	3.0	5.4	8.7	13.5	17.9	22.4	30.3	38.1	39.1
2002.....	1.4	3.3	5.8	8.6	13.5	18.0	22.9	31.1	38.8	39.0
2001.....	1.4	3.1	5.7	8.5	13.2	18.1	23.3	31.1	38.6	40.4
2000.....	1.5	3.2	5.6	8.7	12.7	18.0	22.4	32.0	37.6	42.0
1999.....	1.6	3.3	5.7	8.6	13.0	17.5	22.9	29.6	36.4	38.7
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	2.6	5.7	10.3	15.6	24.8	32.0	42.4	50.8	58.7	57.0
2005 4/.....	2.7	5.7	10.9	17.2	24.4	32.0	41.6	50.1	57.8	57.5
2004 4/.....	2.9	5.9	10.4	15.8	23.8	34.4	42.1	51.6	61.3	56.0
2003 4/.....	2.9	6.0	10.6	17.1	25.9	33.7	40.7	52.1	61.5	56.4
2002.....	2.9	6.6	11.4	16.8	25.9	33.9	41.6	53.4	62.5	55.9
2001.....	2.9	6.2	11.4	16.6	25.3	33.9	42.2	53.1	61.9	57.3
2000.....	3.0	6.4	11.1	17.0	24.5	33.7	40.3	54.5	60.0	59.0
1999.....	3.2	6.5	11.2	16.8	25.1	32.8	41.0	50.5	57.9	54.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of ovary (C56)										
Black, both sexes										
2006 4/.....	1.3	2.7	4.4	6.6	11.3	15.5	16.3	25.3	24.8	33.5
2005 4/.....	1.4	2.3	4.2	6.9	11.1	15.6	22.4	25.2	31.1	31.3
2004 4/.....	1.1	3.0	4.9	7.5	10.2	18.3	22.5	26.2	29.0	32.7
2003 4/.....	1.2	2.8	5.1	6.8	11.0	15.3	20.6	26.2	29.4	36.8
2002.....	1.2	2.9	4.1	7.7	10.9	14.9	20.0	26.5	32.4	34.7
2001.....	1.4	1.9	4.4	7.9	12.6	16.3	20.7	26.2	27.9	36.5
2000.....	1.5	2.2	4.8	6.4	10.9	16.8	19.6	26.1	27.4	34.4
1999.....	1.1	2.4	5.0	7.2	11.4	15.0	22.7	28.9	28.3	28.7
Black, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
Black, female										
2006 4/.....	2.4	5.0	8.1	12.1	20.1	27.0	27.5	40.4	38.0	47.2
2005 4/.....	2.7	4.3	7.8	12.5	19.7	27.1	37.7	40.3	47.5	43.9
2004 4/.....	2.0	5.7	9.1	13.7	18.1	31.8	37.8	41.8	44.2	45.7
2003 4/.....	2.3	5.3	9.4	12.5	19.6	26.7	34.6	41.8	44.9	51.0
2002.....	2.3	5.3	7.7	13.9	19.5	25.9	33.4	42.3	49.3	48.1
2001.....	2.7	3.6	8.1	14.4	22.6	28.3	34.5	41.9	42.1	50.1
2000.....	2.9	4.0	8.8	11.6	19.4	29.2	32.6	41.9	41.2	47.2
1999.....	2.1	4.5	9.3	13.2	20.3	26.2	37.8	46.3	42.3	39.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified female genital organs (C57)

All races 3/, both sexes

2006.....	*	*	0.1	0.2	0.2	0.5	0.7	0.7	0.6	1.2
2005.....	*	*	0.1	0.1	0.2	0.4	0.4	0.4	0.6	0.6
2004.....	*	*	0.1	0.1	0.2	0.2	0.4	0.6	0.5	0.7
2003.....	*	*	0.1	0.2	0.3	0.4	0.6	0.6	1.1	1.1
2002.....	*	*	0.2	0.2	0.4	0.5	0.6	1.0	1.1	1.0
2001.....	*	*	0.1	0.3	0.3	0.4	0.7	0.9	0.9	1.2
2000.....	*	*	0.1	0.2	0.4	0.5	0.6	0.7	1.1	1.6
1999.....	*	*	0.2	0.3	0.4	0.5	0.7	0.8	1.3	0.9

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	*	*	0.3	0.3	0.4	0.8	1.2	1.2	0.9	1.8
2005.....	*	*	0.2	0.2	0.3	0.7	0.8	0.6	1.0	0.9
2004.....	*	*	0.2	0.3	0.4	0.4	0.7	1.0	0.8	1.1
2003.....	*	*	0.3	0.4	0.5	0.8	1.1	1.1	1.7	1.5
2002.....	*	*	0.3	0.4	0.8	1.0	1.2	1.6	1.7	1.5
2001.....	*	*	0.3	0.5	0.6	0.8	1.2	1.5	1.4	1.7
2000.....	*	*	0.2	0.5	0.8	0.8	1.1	1.3	1.7	2.2
1999.....	*	*	0.4	0.5	0.8	0.9	1.3	1.4	2.1	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified female genital organs (C57)

White, both sexes

2006 4/.....	*	*	0.1	0.2	0.2	0.4	0.7	0.7	0.6	1.2
2005 4/.....	*	*	*	*	*	0.4	0.4	0.4	0.7	0.7
2004 4/.....	*	*	*	0.1	0.2	*	0.4	0.6	0.5	0.8
2003 4/.....	*	*	0.1	0.2	0.2	0.5	0.6	0.6	1.1	1.0
2002.....	*	*	0.2	0.2	0.3	0.5	0.7	1.0	1.1	1.0
2001.....	*	*	0.1	0.3	0.3	0.4	0.7	0.9	0.9	1.3
2000.....	*	*	*	0.2	0.4	0.5	0.6	0.6	1.1	1.6
1999.....	*	*	0.2	0.3	0.4	0.5	0.7	0.8	1.2	0.9

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	*	*	0.2	0.3	0.4	0.7	1.3	1.2	1.0	1.7
2005 4/.....	*	*	*	*	*	0.8	0.8	0.7	1.1	1.0
2004 4/.....	*	*	*	0.3	0.4	*	0.8	1.1	0.9	1.1
2003 4/.....	*	*	0.3	0.4	0.4	0.9	1.1	1.1	1.8	1.5
2002.....	*	*	0.3	0.4	0.6	1.0	1.2	1.7	1.7	1.5
2001.....	*	*	0.3	0.5	0.6	0.8	1.3	1.5	1.5	1.8
2000.....	*	*	*	0.5	0.8	0.9	1.1	1.1	1.8	2.2
1999.....	*	*	0.4	0.5	0.9	0.9	1.2	1.4	2.0	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of urinary tract (C64-C68)										
All races 3/, both sexes										
2006.....	1.2	2.8	5.5	10.3	17.0	26.7	38.5	56.0	78.5	105.1
2005.....	1.3	3.0	5.8	10.1	17.9	26.3	39.7	55.7	79.4	103.8
2004.....	1.2	3.1	5.9	10.6	18.3	26.6	39.0	57.3	77.5	102.7
2003.....	1.4	2.9	5.9	10.8	18.6	26.7	39.2	56.5	77.3	101.4
2002.....	1.4	3.3	6.1	11.2	17.6	28.7	41.0	58.2	74.7	102.0
2001.....	1.7	3.2	6.3	10.7	19.3	27.3	40.9	55.2	76.1	103.9
2000.....	1.5	3.1	6.2	11.4	17.9	27.5	40.1	55.2	77.4	106.0
1999.....	1.4	2.9	6.3	11.3	18.2	26.9	40.6	54.4	73.1	103.0
All races 3/, male										
2006.....	1.7	4.0	8.2	15.4	25.5	40.2	57.9	88.6	130.4	189.3
2005.....	1.8	4.2	8.2	14.8	26.7	40.0	61.5	86.3	133.2	186.4
2004.....	1.6	4.3	8.9	15.7	27.9	40.1	59.9	89.6	128.3	182.8
2003.....	2.0	4.5	8.6	16.0	28.4	39.9	58.6	90.0	127.6	178.8
2002.....	1.9	4.7	9.0	16.8	26.8	42.7	61.3	92.6	119.6	183.6
2001.....	2.1	4.7	9.2	15.9	28.9	41.4	62.9	87.5	126.4	192.4
2000.....	2.0	4.3	9.2	17.1	26.7	41.5	61.0	87.4	129.7	198.9
1999.....	1.9	4.3	9.2	16.9	27.2	40.6	61.8	88.2	121.7	188.4
All races 3/, female										
2006.....	0.6	1.7	3.0	5.4	9.1	15.0	22.7	32.2	46.1	65.7
2005.....	0.9	1.9	3.5	5.6	9.8	14.4	22.0	33.4	46.0	65.8
2004.....	0.8	1.9	3.1	5.7	9.7	14.9	22.2	34.1	46.1	66.7
2003.....	0.8	1.4	3.3	5.8	9.6	15.3	23.6	32.7	46.6	67.2
2002.....	0.9	2.0	3.4	6.0	9.4	16.5	24.8	33.9	47.4	66.6
2001.....	1.2	1.8	3.4	5.8	10.5	15.2	23.3	32.7	46.0	66.9
2000.....	1.0	1.9	3.3	6.2	10.0	15.4	23.7	32.7	46.6	68.1
1999.....	0.9	1.6	3.5	6.1	10.0	15.1	23.9	30.8	44.6	68.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of urinary tract (C64-C68)										
White, both sexes										
2006 4/.....	1.1	2.9	5.6	10.6	17.7	27.8	39.9	58.3	81.1	108.2
2005 4/.....	1.4	3.0	5.9	10.4	18.3	27.4	41.3	57.5	82.3	106.1
2004 4/.....	1.2	3.1	6.1	11.0	19.3	27.9	40.3	59.3	80.4	105.5
2003 4/.....	1.3	2.9	6.1	11.0	19.2	27.6	40.9	58.3	79.3	104.1
2002.....	1.4	3.4	6.3	11.5	18.3	29.8	42.4	59.6	76.4	103.9
2001.....	1.6	3.3	6.3	10.9	19.7	28.6	42.2	56.6	77.4	106.4
2000.....	1.5	3.2	6.4	11.8	18.7	28.4	41.1	56.2	78.9	109.2
1999.....	1.4	2.8	6.4	11.5	18.8	27.6	41.4	55.5	74.7	104.6
White, male										
2006 4/.....	1.7	4.0	8.3	15.7	26.5	41.9	60.6	92.7	135.8	197.1
2005 4/.....	1.7	4.3	8.2	15.3	27.3	41.8	63.7	89.2	138.7	193.0
2004 4/.....	1.6	4.3	9.2	16.4	29.2	42.2	62.1	92.9	134.5	190.9
2003 4/.....	1.9	4.5	8.8	16.5	29.5	41.2	61.1	92.7	132.2	186.6
2002.....	1.9	4.9	9.3	17.4	27.8	44.6	63.7	95.5	122.7	190.7
2001.....	2.1	4.9	9.3	16.4	29.5	43.5	65.2	89.8	129.9	200.5
2000.....	2.1	4.4	9.4	17.5	28.0	42.9	62.8	89.8	132.5	206.8
1999.....	1.9	4.2	9.1	17.2	28.2	41.8	63.7	90.7	125.5	194.9
White, female										
2006 4/.....	0.6	1.8	2.9	5.6	9.4	15.3	22.7	32.6	46.5	66.4
2005 4/.....	1.0	1.8	3.6	5.5	10.0	14.7	22.8	34.1	47.0	66.1
2004 4/.....	0.7	1.8	3.1	5.8	10.1	15.3	22.3	34.7	46.7	67.0
2003 4/.....	0.7	1.2	3.4	5.7	9.7	15.6	24.4	33.3	46.7	67.6
2002.....	0.9	1.9	3.3	5.9	9.5	16.8	25.0	33.9	48.1	66.2
2001.....	1.1	1.7	3.5	5.7	10.6	15.5	23.6	33.0	45.7	67.0
2000.....	1.0	1.9	3.5	6.4	10.1	15.8	23.7	32.5	47.0	69.6
1999.....	0.9	1.5	3.7	6.1	10.1	15.2	23.8	30.8	44.7	68.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of urinary tract (C64-C68)										
Black, both sexes										
2006 4/.....	1.7	3.4	6.9	10.8	15.4	23.9	35.1	45.9	66.0	90.5
2005 4/.....	1.5	3.5	6.7	10.8	18.7	23.6	34.2	47.8	63.6	89.3
2004 4/.....	1.8	3.6	6.4	10.1	16.0	21.8	36.5	50.1	59.4	88.8
2003 4/.....	2.2	3.9	6.4	11.4	16.5	25.6	30.4	49.9	69.6	89.0
2002.....	1.3	3.5	6.4	11.7	16.5	25.7	36.1	54.1	64.9	93.2
2001.....	2.4	3.5	6.9	11.1	19.8	22.5	34.5	50.4	71.2	86.6
2000.....	1.5	3.4	6.0	10.9	15.9	25.1	37.5	53.5	74.0	80.4
1999.....	1.7	4.1	7.4	12.8	17.0	26.0	39.2	49.1	61.0	93.1
Black, male										
2006 4/.....	2.7	5.0	9.9	17.0	24.0	34.7	48.3	61.3	97.9	145.7
2005 4/.....	2.4	4.6	9.9	15.2	29.1	33.3	55.7	71.1	97.6	143.2
2004 4/.....	2.5	4.9	9.3	14.1	24.3	28.6	54.4	73.7	83.2	120.2
2003 4/.....	3.4	5.4	9.3	16.0	23.5	38.4	43.7	78.5	101.9	126.2
2002.....	1.7	4.4	8.3	16.5	24.5	34.4	47.4	77.2	98.6	135.1
2001.....	2.7	4.4	10.7	14.9	30.3	31.0	50.2	76.2	101.9	127.1
2000.....	1.7	4.2	9.1	16.3	22.6	38.0	53.5	76.5	122.2	140.7
1999.....	2.3	5.8	11.9	18.6	24.0	36.1	52.2	73.1	89.8	138.3
Black, female										
2006 4/.....	*	2.1	4.4	5.8	8.7	15.9	26.0	36.7	49.2	68.0
2005 4/.....	*	2.6	4.1	7.2	10.6	16.4	19.5	33.9	45.6	67.5
2004 4/.....	*	2.5	3.9	6.7	9.5	16.7	24.4	36.1	46.9	76.4
2003 4/.....	*	2.6	3.9	7.6	11.0	16.1	21.5	32.8	52.6	74.5
2002.....	*	2.8	4.8	7.8	10.2	19.2	28.5	40.3	47.3	77.2
2001.....	2.2	2.7	3.7	8.1	11.5	16.1	24.0	34.9	55.6	71.6
2000.....	*	2.7	3.5	6.5	10.6	15.6	26.9	39.6	49.7	57.9
1999.....	*	2.5	3.6	8.0	11.5	18.5	30.6	34.6	46.7	76.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of kidney and renal pelvis (C64-C65)										
All races 3/, both sexes										
2006.....	0.8	2.0	3.7	6.5	9.6	14.8	18.2	24.0	28.9	34.3
2005.....	0.9	2.0	4.1	6.4	10.8	14.4	19.9	24.3	30.3	32.6
2004.....	0.8	2.1	4.0	6.8	11.3	14.6	19.1	24.5	29.8	32.7
2003.....	1.0	2.0	4.0	7.1	11.4	14.5	19.6	24.8	30.0	33.2
2002.....	0.9	2.4	4.3	7.2	10.9	16.0	19.6	24.6	28.7	32.5
2001.....	1.2	2.4	4.4	7.1	11.9	14.6	20.8	24.4	29.1	32.3
2000.....	1.1	2.2	4.5	7.6	10.8	15.0	19.4	24.5	29.6	33.8
1999.....	1.1	2.2	4.3	7.3	10.8	14.4	19.2	23.1	26.9	31.4
All races 3/, male										
2006.....	1.2	2.8	5.4	9.5	14.0	21.6	25.0	34.5	40.8	52.1
2005.....	1.2	2.9	5.8	9.3	15.7	20.7	28.9	34.1	43.3	50.6
2004.....	1.2	2.9	5.9	9.8	16.6	20.7	27.8	34.2	44.0	47.0
2003.....	1.5	3.0	5.7	10.4	17.2	20.2	27.7	35.9	44.9	48.7
2002.....	1.3	3.4	6.4	10.6	16.1	23.0	27.6	36.6	39.9	50.1
2001.....	1.6	3.5	6.5	10.5	17.2	21.0	30.0	34.4	42.2	52.9
2000.....	1.6	3.1	6.6	11.2	15.6	21.4	27.4	34.6	44.2	55.1
1999.....	1.5	3.2	6.1	10.6	15.8	20.3	27.3	33.7	38.9	49.4
All races 3/, female										
2006.....	0.4	1.2	2.1	3.6	5.6	8.8	12.6	16.4	21.5	25.9
2005.....	0.7	1.2	2.4	3.6	6.4	8.8	12.5	17.2	22.2	24.3
2004.....	0.5	1.3	2.2	3.8	6.4	9.2	12.1	17.6	21.0	26.3
2003.....	0.6	1.0	2.3	4.0	6.1	9.4	13.0	16.9	20.9	26.4
2002.....	0.5	1.5	2.3	3.9	6.2	10.0	13.2	16.2	21.9	24.9
2001.....	0.9	1.4	2.4	3.9	7.0	9.1	13.5	17.4	21.2	23.7
2000.....	0.7	1.4	2.4	4.3	6.5	9.5	13.0	17.5	21.0	25.1
1999.....	0.6	1.2	2.5	4.3	6.2	9.3	12.8	15.7	19.9	24.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of kidney and renal pelvis (C64-C65)										
White, both sexes										
2006 4/.....	0.8	2.0	3.7	6.7	10.0	15.3	18.7	24.9	29.6	34.9
2005 4/.....	0.9	2.0	4.2	6.5	11.0	14.8	20.6	24.9	31.1	33.1
2004 4/.....	0.8	2.2	4.2	7.0	11.7	15.1	19.4	24.9	30.6	33.7
2003 4/.....	1.0	2.0	4.2	7.2	11.6	15.0	20.2	25.4	30.8	33.8
2002.....	1.0	2.4	4.4	7.4	11.2	16.6	20.2	25.0	29.2	33.2
2001.....	1.2	2.5	4.4	7.2	12.1	15.3	21.3	24.7	29.5	32.8
2000.....	1.1	2.3	4.6	7.8	11.1	15.4	19.7	24.9	30.0	34.5
1999.....	1.1	2.1	4.3	7.5	11.2	14.8	19.4	23.4	27.3	31.5
White, male										
2006 4/.....	1.2	2.8	5.4	9.5	14.4	22.0	25.7	35.7	41.8	53.4
2005 4/.....	1.2	2.8	5.9	9.5	15.7	21.3	29.6	34.8	44.2	52.0
2004 4/.....	1.1	3.0	6.2	10.2	17.2	21.3	28.2	34.5	45.4	48.6
2003 4/.....	1.5	3.1	5.9	10.7	17.5	20.8	28.2	36.5	46.2	50.0
2002.....	1.4	3.5	6.6	10.9	16.4	23.8	28.3	37.5	40.2	51.9
2001.....	1.6	3.6	6.5	10.8	17.2	21.9	30.6	34.4	42.8	54.4
2000.....	1.6	3.2	6.7	11.3	16.1	21.5	27.7	35.1	44.5	56.6
1999.....	1.5	3.1	6.0	10.7	16.4	20.6	27.6	34.1	40.1	49.8
White, female										
2006 4/.....	0.4	1.3	2.0	3.9	5.8	9.3	12.9	16.9	22.0	26.2
2005 4/.....	0.7	1.2	2.6	3.7	6.7	9.1	13.1	17.6	22.9	24.3
2004 4/.....	0.5	1.3	2.3	4.0	6.7	9.6	12.2	17.8	21.3	27.0
2003 4/.....	0.6	0.9	2.5	3.8	6.2	9.7	13.7	17.4	21.3	26.6
2002.....	0.5	1.4	2.2	4.1	6.4	10.3	13.6	16.1	22.5	25.1
2001.....	0.8	1.3	2.4	3.8	7.4	9.5	13.8	17.8	21.4	23.8
2000.....	0.7	1.4	2.5	4.5	6.5	10.0	13.3	17.7	21.4	25.5
1999.....	0.6	1.1	2.6	4.4	6.3	9.7	13.0	15.8	19.8	24.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of kidney and renal pelvis (C64-C65)										
Black, both sexes										
2006 4/.....	1.3	2.4	4.9	6.7	8.4	13.2	16.7	19.3	26.0	33.5
2005 4/.....	1.1	2.4	4.4	6.6	12.1	13.5	17.5	22.2	26.2	30.5
2004 4/.....	1.2	2.3	4.0	6.2	10.5	13.1	19.2	26.2	25.3	28.2
2003 4/.....	1.5	2.5	4.1	8.2	11.2	13.4	16.1	22.3	26.1	33.8
2002.....	0.9	2.8	4.4	7.0	10.6	14.3	16.8	25.8	26.7	28.2
2001.....	1.8	2.6	5.0	7.4	12.2	11.7	19.2	24.0	28.2	29.8
2000.....	1.1	2.5	4.7	7.4	10.4	14.8	18.5	23.5	29.1	29.7
1999.....	1.2	2.9	4.9	7.8	9.7	13.5	19.3	23.0	24.3	31.2
Black, male										
2006 4/.....	2.1	3.8	6.8	11.1	13.4	22.0	23.1	27.3	39.7	49.1
2005 4/.....	1.9	3.6	7.1	10.0	19.6	20.0	29.1	32.2	39.3	39.0
2004 4/.....	1.6	3.2	6.1	9.2	16.1	19.6	29.8	38.2	35.9	38.7
2003 4/.....	2.5	3.6	6.5	11.5	17.0	19.5	25.9	35.6	39.1	43.8
2002.....	*	3.5	6.4	11.8	16.8	19.9	23.7	34.4	39.0	39.7
2001.....	1.9	3.5	7.8	10.5	20.5	17.0	29.3	39.0	42.5	46.5
2000.....	*	3.2	7.1	11.4	15.3	25.0	28.1	32.1	48.9	49.6
1999.....	1.8	4.1	8.0	12.5	14.0	20.0	28.2	34.2	27.9	49.2
Black, female										
2006 4/.....	*	*	3.3	3.0	4.5	6.8	12.3	14.5	18.8	27.1
2005 4/.....	*	*	2.1	3.8	6.2	8.6	9.5	16.3	19.3	27.0
2004 4/.....	*	1.5	2.2	3.7	6.1	8.3	12.1	19.1	19.7	24.0
2003 4/.....	*	1.5	2.1	5.5	6.6	9.0	9.5	14.4	19.3	30.0
2002.....	*	2.3	2.6	3.1	5.7	10.2	12.1	20.6	20.3	23.8
2001.....	1.7	1.8	2.7	4.9	5.6	7.7	12.6	15.0	20.8	23.6
2000.....	*	1.8	2.6	4.2	6.5	7.3	12.2	18.2	19.1	22.3
1999.....	*	1.9	2.3	3.9	6.3	8.7	13.5	16.3	22.5	24.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of bladder (C67)										
All races 3/, both sexes										
2006.....	0.3	0.8	1.7	3.5	7.0	11.2	19.2	30.0	46.9	67.4
2005.....	0.4	0.9	1.6	3.5	6.8	11.5	18.9	30.0	46.7	68.3
2004.....	0.4	0.9	1.8	3.6	6.8	11.5	18.9	31.1	45.7	67.9
2003.....	0.3	0.8	1.8	3.5	6.9	11.8	18.5	30.0	45.1	64.8
2002.....	0.4	0.9	1.7	3.8	6.4	12.0	20.2	31.9	44.1	66.6
2001.....	0.4	0.8	1.7	3.4	7.0	12.0	19.0	29.3	44.8	68.9
2000.....	0.3	0.8	1.6	3.6	6.8	11.7	19.6	29.1	45.5	69.6
1999.....	0.3	0.7	1.9	3.8	7.0	11.8	20.2	29.7	43.8	68.5
All races 3/, male										
2006.....	0.5	1.1	2.6	5.5	11.1	17.7	31.4	51.4	85.8	131.6
2005.....	0.5	1.2	2.2	5.3	10.8	18.7	31.2	50.4	86.3	132.1
2004.....	0.5	1.3	2.8	5.7	10.8	18.5	30.6	53.1	81.6	133.0
2003.....	0.4	1.4	2.7	5.5	10.9	19.1	29.6	51.7	80.3	125.1
2002.....	0.5	1.2	2.4	5.9	10.1	19.0	32.2	54.0	76.6	128.9
2001.....	0.5	1.2	2.6	5.2	11.3	19.5	31.6	50.8	81.0	134.8
2000.....	0.4	1.1	2.5	5.7	10.6	19.0	32.2	50.6	82.4	139.4
1999.....	0.4	1.1	2.9	5.9	11.0	19.3	32.9	52.3	79.5	134.8
All races 3/, female										
2006.....	*	0.5	0.8	1.6	3.3	5.6	9.4	14.4	22.6	37.4
2005.....	0.3	0.6	1.0	1.7	3.1	5.2	8.9	15.2	22.1	39.0
2004.....	0.3	0.5	0.8	1.7	3.1	5.3	9.5	15.3	23.5	38.6
2003.....	0.2	0.3	0.9	1.7	3.3	5.4	9.5	14.6	23.5	38.1
2002.....	0.3	0.5	1.0	1.9	2.9	6.0	10.6	16.3	24.2	39.6
2001.....	0.3	0.4	0.9	1.7	3.2	5.6	9.0	14.3	23.2	41.4
2000.....	0.3	0.5	0.8	1.7	3.4	5.5	9.7	14.0	23.8	41.2
1999.....	0.2	0.3	0.9	1.7	3.4	5.4	10.3	13.9	22.9	41.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of bladder (C67)										
White, both sexes										
2006 4/.....	0.3	0.8	1.8	3.6	7.4	11.8	20.0	31.3	48.6	69.8
2005 4/.....	0.4	0.9	1.6	3.6	7.1	12.0	19.8	31.1	48.7	70.0
2004 4/.....	0.3	0.9	1.8	3.8	7.3	12.3	19.8	32.6	47.6	69.6
2003 4/.....	0.3	0.8	1.8	3.7	7.3	12.2	19.6	31.1	46.2	66.6
2002.....	0.4	0.9	1.8	3.9	6.7	12.5	20.9	32.9	45.2	67.7
2001.....	0.4	0.8	1.8	3.5	7.3	12.6	19.8	30.3	45.7	70.8
2000.....	0.3	0.8	1.7	3.8	7.2	12.3	20.2	29.7	46.4	72.0
1999.....	0.3	0.7	2.0	3.8	7.2	12.1	20.8	30.5	45.0	69.8
White, male										
2006 4/.....	0.5	1.2	2.7	5.7	11.6	18.8	33.1	54.2	90.0	137.9
2005 4/.....	0.5	1.4	2.3	5.6	11.3	19.9	32.8	52.5	90.5	137.4
2004 4/.....	0.4	1.3	2.9	6.0	11.6	20.2	32.3	55.9	86.2	139.3
2003 4/.....	0.4	1.4	2.8	5.7	11.7	19.8	31.6	53.8	83.5	131.0
2002.....	0.5	1.3	2.6	6.2	10.8	20.1	33.7	56.0	79.4	134.1
2001.....	0.5	1.2	2.7	5.4	12.0	20.7	33.1	53.1	83.7	141.1
2000.....	0.4	1.2	2.7	6.0	11.4	20.2	33.6	52.7	84.6	145.7
1999.....	0.4	1.1	3.0	6.2	11.4	20.1	34.4	54.3	82.1	140.7
White, female										
2006 4/.....	*	0.5	0.9	1.6	3.4	5.6	9.0	14.1	22.5	37.9
2005 4/.....	0.3	0.5	0.9	1.6	3.2	5.1	9.1	15.3	22.4	39.0
2004 4/.....	0.3	0.5	0.7	1.7	3.3	5.3	9.5	15.6	23.7	38.1
2003 4/.....	*	0.2	0.8	1.7	3.3	5.5	9.8	14.6	23.3	38.1
2002.....	0.3	0.5	1.0	1.7	2.9	5.9	10.4	16.4	24.3	38.9
2001.....	0.3	0.3	0.9	1.7	3.0	5.5	9.0	14.2	22.7	41.3
2000.....	0.3	0.5	0.8	1.8	3.4	5.5	9.6	13.6	23.7	42.1
1999.....	0.2	0.3	1.0	1.5	3.4	5.1	10.0	13.8	23.2	41.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of bladder (C67)										
Black, both sexes										
2006 4/.....	*	0.9	1.7	3.8	6.2	9.7	18.2	25.6	38.6	54.5
2005 4/.....	*	1.0	2.0	3.7	5.9	9.7	15.7	25.1	36.2	57.5
2004 4/.....	*	1.2	2.0	3.3	5.1	7.9	16.5	23.0	33.4	60.1
2003 4/.....	*	1.2	2.2	3.2	5.0	11.4	12.7	26.5	40.9	53.4
2002.....	*	*	1.6	4.0	5.2	10.9	18.6	27.0	37.2	63.3
2001.....	*	0.9	1.7	3.4	6.9	10.0	14.1	24.8	41.2	55.0
2000.....	*	0.8	1.3	3.0	5.4	9.5	17.9	28.4	44.6	49.7
1999.....	*	1.0	2.2	4.5	6.9	11.8	18.9	25.4	35.3	60.3
Black, male										
2006 4/.....	*	*	2.8	5.6	10.4	12.4	24.9	32.7	56.1	92.9
2005 4/.....	*	*	2.7	4.9	9.1	13.1	25.0	38.0	56.9	101.3
2004 4/.....	*	1.7	3.0	4.4	8.0	8.2	23.0	35.0	46.6	81.5
2003 4/.....	*	1.7	2.8	4.5	6.6	18.2	16.4	41.1	61.3	80.3
2002.....	*	*	*	4.5	7.1	14.2	23.7	41.8	57.3	91.1
2001.....	*	*	2.8	4.2	9.2	12.7	20.5	34.4	57.8	78.3
2000.....	*	*	*	4.6	7.2	12.5	24.1	41.1	73.3	88.8
1999.....	*	*	3.9	5.4	9.8	15.6	23.7	38.9	59.3	86.7
Black, female										
2006 4/.....	*	*	*	2.4	2.9	7.6	13.5	21.4	29.3	38.7
2005 4/.....	*	*	*	2.7	3.4	7.2	9.3	17.3	25.2	39.8
2004 4/.....	*	*	*	2.5	*	7.7	12.1	15.9	26.5	51.6
2003 4/.....	*	*	1.7	*	3.8	6.4	10.2	17.9	30.2	42.9
2002.....	*	*	*	3.7	3.7	8.5	15.2	18.1	26.7	52.6
2001.....	*	*	*	2.7	5.1	8.1	9.9	19.1	32.7	46.3
2000.....	*	*	*	*	3.9	7.3	13.8	20.8	30.1	35.2
1999.....	*	*	*	3.7	4.7	9.0	15.8	17.2	23.3	50.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites within the urinary tract (C66,C68)

All races 3/, both sexes

2006.....	*	*	0.1	0.3	0.4	0.7	1.1	2.0	2.7	3.4
2005.....	*	*	0.1	0.3	0.2	0.5	0.9	1.4	2.4	2.9
2004.....	*	*	0.1	0.2	0.3	0.6	1.0	1.6	2.0	2.1
2003.....	*	0.1	*	0.2	0.3	0.5	1.1	1.7	2.3	3.4
2002.....	*	*	0.1	0.2	0.4	0.7	1.2	1.6	1.9	2.9
2001.....	*	*	0.1	0.2	0.4	0.7	1.1	1.5	2.2	2.7
2000.....	*	*	*	0.2	0.3	0.7	1.2	1.6	2.3	2.6
1999.....	*	*	0.1	0.2	0.4	0.7	1.1	1.6	2.3	3.1

All races 3/, male

2006.....	*	*	*	0.4	0.5	0.9	1.6	2.8	3.8	5.6
2005.....	*	*	*	*	*	0.6	1.3	1.9	3.6	3.7
2004.....	*	*	*	*	0.4	0.8	1.5	2.3	2.7	2.8
2003.....	*	*	*	*	*	0.6	1.2	2.3	2.4	5.1
2002.....	*	*	*	0.3	0.6	0.7	1.5	2.0	3.1	4.7
2001.....	*	*	*	*	0.4	0.9	1.3	2.2	3.3	4.7
2000.....	*	*	*	*	0.4	1.1	1.5	2.1	3.1	4.4
1999.....	*	*	*	0.3	*	1.0	1.6	2.2	3.3	4.2

All races 3/, female

2006.....	*	*	*	0.2	*	0.6	0.7	1.5	2.0	2.4
2005.....	*	*	*	0.3	*	0.4	0.6	1.1	1.6	2.6
2004.....	*	*	*	*	*	0.4	0.6	1.2	1.6	1.8
2003.....	*	*	*	*	*	0.4	1.0	1.2	2.2	2.7
2002.....	*	*	*	*	*	0.6	1.0	1.4	1.2	2.2
2001.....	*	*	*	*	*	0.4	0.9	1.0	1.6	1.8
2000.....	*	*	*	*	*	*	0.9	1.2	1.8	1.8
1999.....	*	*	*	*	0.4	0.4	0.8	1.2	1.8	2.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites within the urinary tract (C66,C68)

White, both sexes

2006 4/.....	*	*	*	0.3	0.3	0.7	1.2	2.1	2.8	3.4
2005 4/.....	*	*	*	0.2	0.2	0.5	0.9	1.5	2.5	3.0
2004 4/.....	*	*	*	0.1	0.3	0.6	1.0	1.8	2.2	2.3
2003 4/.....	*	*	*	0.2	0.3	0.5	1.1	1.8	2.3	3.6
2002.....	*	*	*	0.2	0.4	0.7	1.3	1.7	2.0	3.0
2001.....	*	*	0.1	*	0.3	0.7	1.1	1.6	2.3	2.8
2000.....	*	*	*	0.2	0.3	0.7	1.2	1.6	2.5	2.7
1999.....	*	*	*	0.2	0.4	0.7	1.2	1.7	2.4	3.3

White, male

2006 4/.....	*	*	*	0.4	0.5	1.0	1.8	2.9	4.0	5.7
2005 4/.....	*	*	*	*	*	0.7	1.3	2.0	3.9	3.7
2004 4/.....	*	*	*	*	0.4	0.8	1.6	2.5	2.9	3.0
2003 4/.....	*	*	*	*	*	0.5	1.2	2.4	2.5	5.5
2002.....	*	*	*	*	0.6	0.8	1.7	2.1	3.1	4.7
2001.....	*	*	*	*	*	0.9	1.4	2.3	3.4	4.9
2000.....	*	*	*	*	*	1.2	1.5	2.1	3.4	4.6
1999.....	*	*	*	*	*	1.0	1.7	2.4	3.3	4.4

White, female

2006 4/.....	*	*	*	*	*	*	0.8	1.6	2.0	2.4
2005 4/.....	*	*	*	*	*	0.4	0.6	1.2	1.7	2.7
2004 4/.....	*	*	*	*	*	*	0.6	1.2	1.7	1.9
2003 4/.....	*	*	*	*	*	*	0.9	1.3	2.2	2.8
2002.....	*	*	*	*	*	0.6	1.0	1.4	1.3	2.2
2001.....	*	*	*	*	*	0.5	0.8	1.1	1.6	1.9
2000.....	*	*	*	*	*	*	0.9	1.3	1.9	1.9
1999.....	*	*	*	*	*	*	0.8	1.2	1.8	2.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of eye and adnexa (C69)

All races 3/, both sexes

2006.....	*	*	*	0.1	*	0.2	0.4	0.4	0.5	0.4
2005.....	*	*	0.1	0.1	*	0.3	0.3	0.4	0.6	0.8
2004.....	*	*	*	*	0.2	0.3	0.3	0.4	*	0.5
2003.....	*	*	*	0.2	0.2	*	0.3	0.4	0.5	0.7
2002.....	*	*	*	0.2	0.2	0.2	0.3	0.4	0.5	0.7
2001.....	*	*	*	*	*	0.3	0.3	0.4	0.6	0.6
2000.....	*	*	*	0.2	0.2	0.3	0.3	0.4	0.5	0.8
1999.....	*	*	*	*	*	0.2	0.3	0.4	0.6	0.8

All races 3/, male

2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	1.2
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	0.5	*	*	*

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	0.7
2001.....	*	*	*	*	*	*	*	*	*	0.6
2000.....	*	*	*	*	*	*	*	*	*	0.7
1999.....	*	*	*	*	*	*	*	*	0.8	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of eye and adnexa (C69)

White, both sexes

2006 4/.....	*	*	*	0.1	*	0.3	0.4	0.4	0.6	0.5
2005 4/.....	*	*	*	*	*	0.4	0.3	0.4	0.6	0.8
2004 4/.....	*	*	*	*	0.2	0.3	0.4	0.5	*	0.5
2003 4/.....	*	*	*	0.2	0.2	*	0.3	0.4	0.6	0.7
2002.....	*	*	*	0.2	0.2	*	0.3	0.5	0.6	0.8
2001.....	*	*	*	*	*	0.3	0.3	0.4	0.7	0.7
2000.....	*	*	*	*	0.2	0.3	0.4	0.5	0.5	0.8
1999.....	*	*	*	*	*	*	0.4	0.4	0.7	0.9

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	0.6	*	*	*

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	0.7
2001.....	*	*	*	*	*	*	*	*	*	0.7
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	0.9	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes

2006.....	2.3	3.5	5.7	8.1	10.8	13.0	16.8	18.5	19.6	16.7
2005.....	2.7	3.7	5.6	8.3	10.7	14.0	18.0	20.0	20.4	15.8
2004.....	2.4	3.8	5.5	8.4	10.9	14.0	17.7	19.0	20.4	16.6
2003.....	2.9	3.9	5.8	8.5	11.2	13.8	17.3	19.4	19.6	15.2
2002.....	2.8	4.2	6.2	8.5	11.1	14.7	17.8	19.6	18.5	16.4
2001.....	2.8	4.0	6.1	8.3	11.6	14.2	16.7	20.3	20.6	16.4
2000.....	2.9	4.1	6.3	8.2	11.4	14.8	18.3	19.4	21.1	16.9
1999.....	2.8	4.3	6.1	9.2	11.7	15.7	18.3	21.0	19.4	16.0

All races 3/, male

2006.....	2.9	4.3	6.8	9.9	13.5	16.4	20.4	22.9	25.6	21.2
2005.....	3.5	4.3	7.0	10.0	13.1	17.7	21.2	25.3	25.3	19.3
2004.....	2.9	4.7	7.1	10.4	13.7	16.0	21.3	23.1	24.5	19.4
2003.....	3.4	5.1	7.4	9.7	13.7	16.4	21.3	24.0	25.4	20.0
2002.....	3.5	5.3	7.2	10.6	13.1	18.4	22.2	24.4	22.9	20.1
2001.....	3.5	4.9	7.5	10.7	13.9	17.3	21.2	25.4	26.5	20.1
2000.....	3.7	4.9	7.7	10.1	14.0	17.9	22.8	23.2	26.3	22.4
1999.....	3.4	5.2	7.4	10.6	14.7	19.5	22.3	26.1	24.5	20.7

All races 3/, female

2006.....	1.8	2.7	4.6	6.4	8.4	10.0	13.9	15.3	15.8	14.6
2005.....	1.9	3.0	4.3	6.8	8.6	10.7	15.4	16.1	17.4	14.3
2004.....	1.9	2.8	4.0	6.4	8.4	12.2	14.7	16.0	17.9	15.4
2003.....	2.4	2.8	4.3	7.3	9.0	11.6	14.1	16.2	16.0	13.1
2002.....	2.1	3.1	5.1	6.5	9.2	11.4	14.2	16.2	15.8	14.8
2001.....	2.1	3.1	4.8	6.0	9.5	11.5	13.1	16.7	17.1	14.8
2000.....	2.0	3.3	4.9	6.5	9.0	12.1	14.8	16.7	18.0	14.7
1999.....	2.3	3.4	4.9	7.9	8.9	12.4	15.1	17.5	16.5	14.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, both sexes

2006 4/.....	2.6	3.9	6.3	8.7	11.9	14.0	18.1	20.0	20.6	17.4
2005 4/.....	3.0	4.0	6.2	9.1	11.6	15.2	19.4	21.1	21.5	16.5
2004 4/.....	2.6	4.1	6.0	9.2	11.9	15.2	19.1	20.2	21.7	17.3
2003 4/.....	3.3	4.3	6.3	9.1	12.2	14.9	18.5	20.6	20.7	15.7
2002.....	3.1	4.7	6.7	9.1	12.0	15.9	19.0	20.7	19.3	17.1
2001.....	3.0	4.4	6.6	8.8	12.6	15.2	17.8	21.6	21.6	17.0
2000.....	3.1	4.5	6.9	8.8	12.3	16.0	19.7	20.5	21.8	17.5
1999.....	3.0	4.7	6.6	10.0	12.5	16.9	19.6	22.2	20.3	16.6

White, male

2006 4/.....	3.3	4.7	7.5	10.5	14.8	17.7	21.8	24.6	26.6	22.1
2005 4/.....	3.9	4.6	7.6	10.7	14.0	19.3	22.5	26.3	26.5	20.2
2004 4/.....	3.1	5.1	7.6	11.3	14.8	17.4	23.1	24.2	25.8	20.6
2003 4/.....	3.8	5.5	8.0	10.3	14.8	17.4	22.6	25.3	26.6	20.7
2002.....	3.9	5.9	7.9	11.2	14.0	19.9	23.6	25.6	23.7	21.2
2001.....	3.8	5.4	8.1	11.4	14.9	18.6	22.6	26.7	27.8	20.5
2000.....	4.0	5.4	8.3	10.9	15.2	19.4	24.3	24.3	27.5	23.5
1999.....	3.6	5.7	7.9	11.4	15.7	20.9	23.9	27.4	25.8	21.8

White, female

2006 4/.....	1.9	3.0	5.1	7.0	9.3	10.8	15.0	16.6	16.8	15.3
2005 4/.....	2.1	3.3	4.8	7.4	9.4	11.5	16.7	17.3	18.4	14.8
2004 4/.....	2.1	3.1	4.4	7.2	9.1	13.2	15.7	17.2	19.2	15.8
2003 4/.....	2.7	3.0	4.8	7.9	9.8	12.7	15.2	17.3	17.0	13.4
2002.....	2.3	3.5	5.5	7.1	10.2	12.4	15.3	17.2	16.7	15.4
2001.....	2.3	3.4	5.2	6.4	10.5	12.2	13.9	18.0	17.9	15.6
2000.....	2.1	3.5	5.5	6.9	9.7	13.0	16.0	17.7	18.5	15.0
1999.....	2.5	3.8	5.4	8.7	9.7	13.4	16.2	18.5	17.1	14.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, both sexes

2006 4/.....	1.3	1.9	3.2	5.4	5.0	7.0	9.3	7.4	12.3	9.6
2005 4/.....	1.3	2.4	3.1	5.2	5.6	7.2	9.8	12.9	10.9	9.4
2004 4/.....	1.8	2.2	3.5	4.1	6.0	7.1	9.2	10.0	9.9	10.4
2003 4/.....	1.6	2.7	3.6	5.5	5.6	8.3	9.7	10.6	10.4	10.5
2002.....	1.3	1.9	3.7	5.7	5.6	8.0	9.5	12.2	11.3	9.5
2001.....	1.6	2.6	3.4	5.4	6.4	9.0	9.4	9.4	10.8	11.7
2000.....	2.0	2.2	3.7	5.4	6.0	7.5	8.8	10.3	14.4	10.6
1999.....	1.8	2.6	4.0	5.3	6.7	8.5	7.8	12.0	11.6	9.9

Black, male

2006 4/.....	1.5	2.4	4.0	7.3	6.5	8.4	12.3	*	19.2	*
2005 4/.....	1.5	2.9	4.5	6.6	7.5	8.1	13.3	19.2	14.7	*
2004 4/.....	2.8	2.9	5.0	5.9	7.8	7.8	8.1	14.1	14.3	*
2003 4/.....	2.4	3.1	5.1	6.6	6.7	11.8	12.2	16.0	*	*
2002.....	*	2.1	3.3	8.2	7.9	9.5	11.7	15.6	17.6	*
2001.....	1.9	3.0	4.4	6.9	9.2	9.1	11.4	12.5	*	*
2000.....	2.7	2.2	5.7	6.6	7.7	7.8	12.4	11.8	*	*
1999.....	2.6	3.7	5.3	5.9	8.7	10.9	8.8	14.2	*	*

Black, female

2006 4/.....	*	1.5	2.5	3.9	3.9	5.9	7.2	7.0	8.7	9.1
2005 4/.....	*	2.0	1.9	4.0	4.1	6.5	7.4	9.1	8.9	9.8
2004 4/.....	*	1.6	2.2	2.6	4.5	6.7	9.9	7.5	7.6	11.4
2003 4/.....	*	2.3	2.2	4.6	4.7	5.7	8.0	7.3	8.5	11.7
2002.....	*	1.8	4.0	3.7	3.7	6.8	8.1	10.2	8.0	10.3
2001.....	1.3	2.3	2.6	4.1	4.3	8.8	8.1	7.5	10.2	9.7
2000.....	1.5	2.2	2.1	4.5	4.7	7.3	6.4	9.4	14.0	10.7
1999.....	*	1.6	2.8	4.8	5.2	6.7	7.2	10.6	12.5	9.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of brain (C71)										
All races 3/, both sexes										
2006.....	2.3	3.4	5.6	8.0	10.7	12.8	16.5	18.2	19.3	16.2
2005.....	2.6	3.6	5.5	8.2	10.6	13.8	17.6	19.6	19.9	15.5
2004.....	2.4	3.7	5.4	8.2	10.7	13.9	17.4	18.7	20.1	16.1
2003.....	2.9	3.9	5.7	8.3	11.1	13.6	17.1	19.0	19.2	14.6
2002.....	2.7	4.2	6.0	8.4	10.9	14.4	17.3	19.2	18.0	16.0
2001.....	2.7	3.9	6.0	8.2	11.4	14.0	16.4	19.8	20.1	16.0
2000.....	2.8	4.0	6.2	8.1	11.2	14.5	18.0	19.1	20.6	16.3
1999.....	2.7	4.2	6.0	9.1	11.4	15.5	17.9	20.5	19.0	15.5
All races 3/, male										
2006.....	2.9	4.2	6.7	9.7	13.4	16.2	20.0	22.6	25.3	20.5
2005.....	3.4	4.2	6.9	9.9	12.9	17.5	20.9	24.8	24.6	19.0
2004.....	2.9	4.7	7.0	10.2	13.4	15.9	21.1	22.8	24.1	19.1
2003.....	3.4	5.0	7.3	9.5	13.6	16.2	21.1	23.7	25.2	19.2
2002.....	3.4	5.3	7.1	10.4	12.9	18.0	21.6	23.9	21.9	19.7
2001.....	3.4	4.8	7.4	10.5	13.7	17.0	20.8	24.9	26.0	19.7
2000.....	3.6	4.8	7.6	10.0	13.9	17.7	22.5	22.8	25.8	21.8
1999.....	3.3	5.2	7.3	10.5	14.4	19.2	21.9	25.5	24.0	20.1
All races 3/, female										
2006.....	1.7	2.7	4.5	6.3	8.3	9.8	13.5	15.0	15.5	14.2
2005.....	1.9	3.0	4.2	6.6	8.5	10.6	15.0	15.9	17.0	13.9
2004.....	1.9	2.8	3.9	6.4	8.2	12.1	14.5	15.8	17.6	14.7
2003.....	2.3	2.8	4.2	7.2	9.0	11.4	13.9	15.7	15.5	12.6
2002.....	2.0	3.1	5.0	6.4	9.1	11.2	13.9	15.9	15.5	14.4
2001.....	2.0	3.1	4.7	6.0	9.3	11.4	12.9	16.3	16.6	14.4
2000.....	2.0	3.2	4.8	6.3	8.8	11.9	14.5	16.5	17.6	14.1
1999.....	2.2	3.3	4.7	7.8	8.7	12.3	14.8	17.1	16.0	13.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of brain (C71)										
White, both sexes										
2006 4/.....	2.6	3.8	6.1	8.6	11.8	13.9	17.8	19.7	20.3	16.9
2005 4/.....	3.0	3.9	6.1	8.9	11.5	15.0	19.0	20.7	21.0	16.2
2004 4/.....	2.6	4.0	5.9	9.1	11.6	15.1	18.8	19.9	21.3	16.7
2003 4/.....	3.2	4.2	6.2	9.0	12.1	14.6	18.3	20.2	20.3	15.0
2002.....	3.0	4.6	6.6	9.0	11.8	15.6	18.6	20.3	18.8	16.7
2001.....	3.0	4.3	6.6	8.7	12.4	14.9	17.4	21.1	21.2	16.6
2000.....	3.0	4.4	6.8	8.7	12.2	15.7	19.4	20.2	21.4	16.9
1999.....	3.0	4.6	6.5	9.9	12.3	16.6	19.2	21.7	19.8	16.2
White, male										
2006 4/.....	3.3	4.6	7.3	10.4	14.6	17.4	21.5	24.3	26.3	21.4
2005 4/.....	3.9	4.6	7.5	10.6	13.9	19.0	22.3	25.8	25.8	19.9
2004 4/.....	3.0	5.0	7.5	11.1	14.5	17.3	22.9	23.9	25.3	20.4
2003 4/.....	3.7	5.5	7.8	10.1	14.7	17.2	22.4	24.9	26.4	19.8
2002.....	3.8	5.8	7.9	11.0	13.9	19.4	23.0	25.1	22.7	20.8
2001.....	3.8	5.3	8.0	11.2	14.6	18.3	22.1	26.1	27.4	20.1
2000.....	3.9	5.3	8.1	10.7	15.0	19.1	24.0	24.1	27.0	23.0
1999.....	3.5	5.6	7.8	11.3	15.4	20.6	23.5	26.8	25.3	21.2
White, female										
2006 4/.....	1.9	3.0	5.0	6.9	9.2	10.7	14.7	16.3	16.5	14.9
2005 4/.....	2.1	3.3	4.8	7.3	9.3	11.4	16.3	17.0	18.0	14.5
2004 4/.....	2.1	3.1	4.3	7.1	9.0	13.0	15.5	16.9	18.8	15.1
2003 4/.....	2.7	3.0	4.7	7.8	9.8	12.4	15.0	16.9	16.5	12.9
2002.....	2.2	3.5	5.4	7.0	10.0	12.2	15.0	16.9	16.4	14.9
2001.....	2.3	3.3	5.2	6.3	10.3	12.0	13.6	17.5	17.4	15.1
2000.....	2.1	3.4	5.4	6.7	9.5	12.8	15.7	17.5	18.1	14.5
1999.....	2.5	3.7	5.2	8.5	9.5	13.2	15.9	18.1	16.6	14.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of brain (C71)										
Black, both sexes										
2006 4/.....	1.3	1.8	3.1	5.3	5.0	6.7	8.8	7.4	12.1	9.3
2005 4/.....	1.3	2.3	3.1	4.9	5.5	7.0	9.3	12.5	10.2	8.8
2004 4/.....	1.7	2.2	3.3	3.9	5.7	7.0	8.9	9.8	9.7	10.4
2003 4/.....	1.6	2.7	3.5	5.2	5.5	8.1	9.0	10.0	10.2	10.2
2002.....	1.3	1.8	3.6	5.3	5.3	7.5	9.0	11.9	10.5	9.2
2001.....	1.5	2.6	3.3	5.3	6.2	8.9	9.4	9.2	10.6	11.7
2000.....	1.9	2.2	3.6	5.3	5.8	7.2	8.8	9.8	14.1	10.0
1999.....	1.8	2.6	3.7	5.0	6.4	8.5	7.4	11.4	10.7	9.2
Black, male										
2006 4/.....	1.5	2.4	3.9	7.2	6.4	8.2	11.4	*	18.5	*
2005 4/.....	1.5	2.8	4.5	6.2	7.3	7.8	12.3	18.3	*	*
2004 4/.....	2.8	2.9	4.9	5.5	7.7	7.8	7.4	13.6	*	*
2003 4/.....	2.4	3.1	5.0	6.2	6.7	11.6	11.5	16.0	*	*
2002.....	*	2.0	3.1	7.7	7.5	9.0	10.7	15.2	16.0	*
2001.....	1.8	2.9	4.2	6.7	8.7	9.1	11.4	12.5	*	*
2000.....	2.6	2.2	5.6	6.4	7.5	7.5	12.4	10.9	*	*
1999.....	2.5	3.7	4.9	5.8	8.5	10.9	7.8	13.8	*	*
Black, female										
2006 4/.....	*	1.3	2.5	3.7	3.9	5.5	7.0	7.0	8.7	8.8
2005 4/.....	*	2.0	1.9	3.7	4.1	6.3	7.1	9.1	8.5	9.0
2004 4/.....	*	1.6	1.9	2.6	4.2	6.5	9.9	7.5	7.6	11.4
2003 4/.....	*	2.3	2.2	4.4	4.5	5.5	7.3	6.5	8.1	11.3
2002.....	*	1.7	4.0	3.3	3.6	6.5	7.8	9.9	*	10.3
2001.....	*	2.3	2.6	4.1	4.2	8.6	8.1	7.2	9.8	9.7
2000.....	*	2.1	2.0	4.3	4.6	6.9	6.4	9.1	14.0	10.3
1999.....	*	1.6	2.7	4.4	4.8	6.7	7.2	10.0	11.7	8.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of thyroid and other endocrine glands (C73-C75)

All races 3/, both sexes

2006.....	0.2	0.3	0.6	1.0	1.4	2.1	3.0	3.7	4.9	5.8
2005.....	0.2	0.4	0.7	1.0	1.4	2.1	2.9	3.5	4.8	5.6
2004.....	0.2	0.4	0.5	0.9	1.4	2.1	2.8	3.4	4.7	6.0
2003.....	0.2	0.4	0.6	1.0	1.4	1.8	2.6	3.5	4.2	5.1
2002.....	0.2	0.5	0.6	1.0	1.4	1.9	2.7	3.9	4.7	5.3
2001.....	0.2	0.5	0.7	1.0	1.7	2.2	3.1	3.8	4.7	5.3
2000.....	0.2	0.4	0.5	1.0	1.6	2.3	2.9	3.9	4.5	6.2
1999.....	0.2	0.4	0.6	1.1	1.6	2.0	2.6	3.9	4.4	5.8

All races 3/, male

2006.....	0.2	0.4	0.6	0.9	1.4	2.1	3.3	3.3	4.7	4.7
2005.....	0.2	0.4	0.7	1.1	1.3	1.8	3.2	2.8	5.4	5.4
2004.....	0.2	0.3	0.5	1.0	1.4	2.2	3.3	3.3	4.2	5.5
2003.....	0.2	0.4	0.7	0.9	1.5	1.7	2.9	3.5	3.7	4.5
2002.....	0.2	0.5	0.7	1.0	1.6	1.9	2.6	4.1	4.6	4.2
2001.....	0.3	0.5	0.7	1.0	1.8	2.5	3.3	3.4	4.7	4.5
2000.....	0.2	0.5	0.5	1.0	1.8	2.5	3.0	3.9	5.0	5.3
1999.....	0.3	0.4	0.7	1.1	1.7	2.1	3.1	4.0	4.5	5.1

All races 3/, female

2006.....	0.2	0.3	0.7	1.0	1.4	2.2	2.8	3.9	5.0	6.3
2005.....	0.2	0.4	0.6	0.9	1.5	2.4	2.6	3.9	4.5	5.7
2004.....	*	0.5	0.5	0.9	1.4	2.1	2.4	3.5	4.9	6.3
2003.....	*	0.4	0.6	1.0	1.4	1.8	2.5	3.5	4.5	5.4
2002.....	0.2	0.4	0.6	0.9	1.3	1.9	2.8	3.8	4.8	5.8
2001.....	0.2	0.5	0.6	0.9	1.6	1.9	3.0	4.0	4.7	5.7
2000.....	0.3	0.3	0.6	1.1	1.5	2.1	2.8	3.8	4.3	6.5
1999.....	0.2	0.4	0.5	1.1	1.6	1.9	2.2	3.8	4.4	6.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of thyroid and other endocrine glands (C73-C75)										
White, both sexes										
2006 4/.....	0.2	0.3	0.7	1.0	1.4	2.2	2.9	3.8	4.9	5.7
2005 4/.....	0.2	0.4	0.7	0.9	1.4	2.1	2.9	3.4	4.9	5.4
2004 4/.....	0.2	0.4	0.5	0.9	1.4	2.1	2.9	3.5	4.7	5.9
2003 4/.....	0.2	0.4	0.7	1.0	1.5	1.9	2.7	3.5	4.1	5.2
2002.....	0.2	0.5	0.6	1.0	1.5	2.0	2.6	4.0	4.7	5.3
2001.....	0.3	0.5	0.7	1.0	1.7	2.3	3.3	3.7	4.5	5.3
2000.....	0.2	0.4	0.5	1.1	1.6	2.2	2.8	3.9	4.6	6.2
1999.....	0.2	0.5	0.6	1.1	1.6	2.0	2.6	3.8	4.5	5.8
White, male										
2006 4/.....	*	0.4	0.6	1.0	1.5	2.2	3.4	3.6	4.7	4.8
2005 4/.....	0.2	0.4	0.7	1.1	1.4	1.9	3.3	2.8	5.5	5.7
2004 4/.....	*	0.3	0.5	1.0	1.4	2.3	3.4	3.4	4.4	5.3
2003 4/.....	0.2	0.5	0.7	1.0	1.6	1.9	2.9	3.5	3.9	4.6
2002.....	0.2	0.4	0.7	1.0	1.7	2.0	2.6	4.2	4.8	4.1
2001.....	0.3	0.6	0.7	1.0	1.8	2.7	3.6	3.6	4.7	4.5
2000.....	*	0.5	0.5	1.1	1.6	2.5	3.1	4.1	5.2	5.6
1999.....	0.3	0.5	0.7	1.0	1.7	2.2	3.1	4.2	4.7	4.9
White, female										
2006 4/.....	*	0.3	0.7	0.9	1.3	2.3	2.5	4.0	5.0	6.1
2005 4/.....	*	0.3	0.6	0.8	1.4	2.3	2.5	3.8	4.6	5.2
2004 4/.....	*	0.5	0.4	0.9	1.4	1.9	2.4	3.6	4.8	6.2
2003 4/.....	*	0.4	0.6	1.0	1.4	1.8	2.5	3.5	4.3	5.5
2002.....	0.2	0.5	0.6	0.9	1.4	2.0	2.7	3.9	4.5	5.9
2001.....	0.2	0.4	0.6	0.9	1.6	1.9	3.0	3.8	4.4	5.7
2000.....	0.2	0.3	0.5	1.0	1.5	2.0	2.6	3.8	4.2	6.5
1999.....	*	0.5	0.5	1.2	1.5	1.9	2.2	3.6	4.4	6.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

All races 3/, both sexes

2006.....	2.9	5.8	10.2	17.3	27.5	40.5	58.7	79.0	105.2	135.8
2005.....	2.9	5.9	10.5	17.9	29.3	41.0	59.2	79.7	108.6	141.4
2004.....	2.9	6.0	10.4	18.0	27.9	42.6	60.1	82.9	109.6	141.2
2003.....	3.0	6.4	11.4	18.7	28.9	43.2	61.9	86.5	110.1	147.9
2002.....	3.2	6.5	11.3	18.8	31.0	44.0	64.4	85.4	112.6	154.5
2001.....	3.3	6.3	11.3	19.9	30.4	44.2	64.4	86.9	112.4	155.2
2000.....	3.4	6.5	11.8	19.6	32.0	46.9	66.7	90.0	119.6	159.0
1999.....	3.9	7.0	12.2	22.1	35.9	52.6	74.2	96.8	128.2	168.4

All races 3/, male

2006.....	2.9	6.6	12.2	21.2	34.1	48.7	70.3	98.1	132.3	168.1
2005.....	3.0	6.7	12.5	22.3	35.5	50.7	71.7	97.2	133.2	175.5
2004.....	2.9	6.7	12.7	22.0	33.6	53.0	72.7	103.5	137.1	170.3
2003.....	3.1	7.4	13.3	22.7	35.2	52.7	74.4	106.9	137.6	181.3
2002.....	3.3	7.2	14.0	21.9	38.3	54.3	79.5	105.3	138.8	196.9
2001.....	3.5	7.2	13.8	23.5	37.1	52.9	79.1	107.0	138.3	196.5
2000.....	3.6	7.5	13.9	23.3	38.6	56.6	81.7	110.4	148.1	200.1
1999.....	4.0	8.3	14.3	26.5	43.1	62.9	92.2	118.6	158.2	218.9

All races 3/, female

2006.....	3.0	5.0	8.4	13.5	21.4	33.4	49.3	64.9	88.2	120.7
2005.....	2.8	5.1	8.5	13.8	23.8	32.4	49.1	66.9	93.4	125.7
2004.....	2.9	5.3	8.3	14.2	22.6	33.6	49.9	68.2	92.7	128.1
2003.....	2.8	5.4	9.7	14.9	23.2	34.9	51.8	72.0	93.3	133.2
2002.....	3.2	5.8	8.6	15.9	24.4	35.1	52.3	71.4	96.6	136.0
2001.....	3.0	5.4	8.9	16.4	24.3	36.8	52.8	72.9	96.9	137.9
2000.....	3.1	5.5	9.7	16.2	26.0	38.6	54.8	75.8	102.8	142.3
1999.....	3.8	5.8	10.3	18.0	29.3	43.8	60.1	81.5	110.7	148.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

White, both sexes

2006 4/.....	2.8	5.5	9.6	16.5	26.5	39.8	59.2	79.2	107.0	137.3
2005 4/.....	2.8	5.5	9.9	17.3	28.6	40.2	59.1	80.3	109.2	144.0
2004 4/.....	2.8	5.4	9.8	17.2	27.1	41.7	60.0	83.0	109.8	143.3
2003 4/.....	2.7	5.9	10.7	17.7	28.3	42.5	61.5	87.0	111.1	149.3
2002.....	3.0	6.1	10.7	18.2	30.0	43.3	63.6	84.8	112.5	154.9
2001.....	3.0	5.8	10.6	18.8	29.3	43.2	62.9	86.8	111.6	157.1
2000.....	3.1	5.9	11.0	19.0	30.9	46.5	65.6	88.5	120.1	160.3
1999.....	3.6	6.5	11.5	20.9	34.6	52.1	72.9	96.0	128.0	168.0

White, male

2006 4/.....	2.8	6.1	11.1	20.1	33.1	47.9	70.6	98.5	134.1	170.5
2005 4/.....	2.9	6.1	11.8	21.0	34.4	49.2	70.6	98.1	133.7	178.6
2004 4/.....	2.6	5.9	11.6	20.7	32.2	51.2	71.6	103.6	138.0	172.2
2003 4/.....	2.8	6.7	12.3	21.3	34.2	51.4	73.5	107.2	138.3	184.1
2002.....	2.9	6.7	13.2	21.1	36.5	53.9	78.5	103.6	139.1	198.4
2001.....	3.2	6.6	12.5	21.8	35.2	51.0	76.9	106.2	136.9	199.0
2000.....	3.2	6.7	12.8	22.1	36.1	55.9	80.1	106.9	149.0	201.3
1999.....	3.7	7.3	13.3	24.8	41.4	61.7	90.2	117.7	158.1	218.7

White, female

2006 4/.....	2.8	5.0	8.1	13.0	20.4	32.6	49.7	64.7	90.0	121.7
2005 4/.....	2.7	4.9	8.0	13.7	23.3	32.2	49.7	67.2	93.8	128.1
2004 4/.....	2.9	4.9	8.1	13.8	22.3	33.3	50.5	67.9	92.3	130.3
2003 4/.....	2.7	5.2	9.2	14.2	22.8	34.6	51.7	72.4	94.3	133.9
2002.....	3.2	5.4	8.2	15.5	23.9	33.9	51.4	71.3	96.2	136.0
2001.....	2.9	5.0	8.7	16.0	23.9	36.4	51.5	73.0	96.3	139.6
2000.....	3.0	5.2	9.2	16.1	26.1	38.3	53.9	75.6	102.9	143.7
1999.....	3.5	5.6	9.8	17.1	28.3	43.8	59.2	80.8	110.2	147.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other malignant neoplasms of other and unspecified sites (C76-C80,C97)

Black, both sexes

2006 4/.....	4.1	8.3	16.5	27.9	41.6	55.0	68.4	92.2	102.7	136.5
2005 4/.....	4.2	9.3	17.1	27.7	40.8	54.9	72.1	91.5	123.1	126.4
2004 4/.....	4.5	10.6	16.9	28.4	40.9	60.1	73.4	98.3	124.5	135.9
2003 4/.....	4.7	10.4	19.4	30.8	42.0	58.1	76.9	97.5	118.5	150.0
2002.....	4.8	10.3	17.2	28.5	46.1	58.8	84.7	108.1	127.9	168.4
2001.....	4.8	11.1	18.8	31.3	45.0	61.9	88.7	101.9	134.1	152.9
2000.....	5.9	11.5	19.9	29.1	46.8	59.5	86.2	117.5	128.8	155.2
1999.....	6.3	12.4	20.1	36.2	53.4	65.5	95.9	117.3	143.9	184.3

Black, male

2006 4/.....	4.0	11.1	22.5	36.4	53.2	69.1	88.3	118.1	145.8	163.5
2005 4/.....	4.8	11.8	20.3	39.6	53.8	76.3	98.4	107.7	160.8	165.5
2004 4/.....	5.4	13.2	24.5	38.4	55.9	81.6	101.6	126.9	153.5	178.8
2003 4/.....	5.4	13.4	24.3	41.2	54.4	77.1	99.9	124.6	165.4	174.1
2002.....	5.9	11.7	22.7	35.6	63.4	71.6	108.2	145.6	162.0	212.3
2001.....	5.6	13.4	26.6	41.9	62.2	82.1	117.5	138.5	170.1	208.8
2000.....	7.5	15.1	26.0	38.7	67.5	76.0	110.7	169.6	160.2	204.1
1999.....	6.5	16.7	25.7	49.0	69.0	87.8	126.2	145.3	183.1	241.4

Black, female

2006 4/.....	4.3	5.8	11.5	20.9	32.5	44.5	54.7	76.6	79.9	125.4
2005 4/.....	3.7	7.2	14.4	17.9	30.7	39.0	54.1	81.9	103.1	110.7
2004 4/.....	3.8	8.4	10.5	20.1	29.3	44.1	54.3	81.4	109.2	118.9
2003 4/.....	4.1	7.7	15.2	22.2	32.3	44.0	61.4	81.4	94.0	140.6
2002.....	3.8	9.1	12.6	22.7	32.5	49.4	69.0	85.6	110.2	151.6
2001.....	4.1	9.2	12.1	22.6	31.5	46.9	69.5	80.2	115.7	132.2
2000.....	4.5	8.4	14.7	21.2	30.4	47.4	69.9	86.0	112.9	136.9
1999.....	6.1	8.7	15.2	25.8	41.3	49.1	75.8	100.4	124.5	163.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes

2006.....	4.2	6.3	11.0	18.5	32.4	51.7	81.7	119.0	162.2	186.1
2005.....	4.1	6.6	11.0	19.0	33.2	52.9	83.8	121.4	161.5	189.8
2004.....	4.1	6.7	11.6	20.2	33.1	54.5	84.2	122.7	162.6	189.4
2003.....	4.7	7.2	11.7	20.1	35.5	55.2	87.3	127.4	165.2	198.0
2002.....	4.6	7.5	12.6	21.5	36.9	59.2	90.1	125.6	168.8	197.5
2001.....	4.8	8.1	13.0	22.7	38.1	58.8	91.8	129.0	168.5	199.9
2000.....	4.8	8.3	13.4	23.4	38.2	62.2	91.6	131.4	176.0	204.6
1999.....	4.8	8.4	13.6	25.6	39.4	62.8	93.9	130.7	174.0	199.4

All races 3/, male

2006.....	5.1	7.4	14.0	23.4	41.2	65.7	103.4	157.6	219.4	244.7
2005.....	4.9	7.8	13.3	23.9	40.7	68.1	109.5	161.0	214.7	262.4
2004.....	5.0	8.1	14.4	24.9	41.7	70.0	108.7	158.6	214.5	254.9
2003.....	5.8	8.8	14.5	25.2	45.6	71.9	112.3	163.5	216.8	268.7
2002.....	5.5	9.1	15.1	26.8	45.7	76.4	116.8	164.3	224.2	269.1
2001.....	5.9	9.8	16.0	27.7	47.5	75.2	116.9	165.7	220.9	275.3
2000.....	5.7	10.3	16.1	28.4	45.8	78.3	117.7	169.8	230.6	281.8
1999.....	5.9	10.2	16.7	31.0	48.6	78.1	118.9	169.4	221.8	279.0

All races 3/, female

2006.....	3.3	5.2	8.2	13.9	24.3	39.4	64.0	90.8	126.3	158.7
2005.....	3.3	5.5	8.7	14.4	26.4	39.6	63.0	92.7	128.4	156.4
2004.....	3.2	5.3	9.0	15.7	25.2	40.9	64.5	96.8	130.6	160.0
2003.....	3.5	5.5	9.1	15.3	26.4	40.8	67.3	101.7	133.6	166.8
2002.....	3.7	6.0	10.1	16.4	28.9	44.4	68.8	98.2	135.1	166.5
2001.....	3.7	6.4	10.1	18.1	29.5	44.6	71.8	103.4	137.1	168.4
2000.....	3.9	6.5	10.8	18.8	31.3	48.4	71.0	104.7	143.8	173.1
1999.....	3.8	6.6	10.6	20.5	31.0	49.8	74.3	103.8	145.9	167.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
White, both sexes										
2006 4/.....	4.1	5.9	10.9	18.2	32.0	52.4	83.5	122.3	168.0	192.2
2005 4/.....	3.9	6.3	10.5	18.7	33.3	53.6	85.6	125.4	167.1	197.3
2004 4/.....	4.0	6.5	11.2	19.9	33.1	55.0	85.0	125.4	167.2	194.9
2003 4/.....	4.6	7.0	11.3	20.1	35.8	56.1	88.7	130.3	169.5	204.5
2002.....	4.6	7.2	12.1	21.0	36.9	60.3	91.7	128.7	173.6	204.0
2001.....	4.6	7.9	12.8	22.3	38.0	59.0	93.7	131.9	172.8	205.9
2000.....	4.7	8.0	13.2	23.3	38.5	63.5	93.4	134.4	181.5	208.7
1999.....	4.7	8.3	13.4	25.3	39.2	63.8	95.6	133.1	178.4	204.1
White, male										
2006 4/.....	4.9	6.9	13.7	22.9	40.7	66.4	105.6	161.6	227.7	252.0
2005 4/.....	4.7	7.4	12.7	23.5	41.1	69.4	111.8	165.6	221.9	275.3
2004 4/.....	5.0	7.8	13.9	24.8	41.9	70.4	109.7	162.3	221.1	264.0
2003 4/.....	5.7	8.7	13.9	25.1	46.3	73.6	115.2	168.7	224.3	279.5
2002.....	5.5	8.8	14.8	26.5	45.7	77.9	119.2	169.0	231.0	279.4
2001.....	5.7	9.7	15.8	27.1	47.3	75.6	119.2	169.8	226.4	284.8
2000.....	5.8	9.9	15.6	28.3	46.1	80.4	119.0	174.1	238.0	290.2
1999.....	5.6	10.0	16.5	30.6	48.6	79.4	121.0	172.1	228.5	285.8
White, female										
2006 4/.....	3.3	4.9	8.1	13.6	23.9	39.9	65.1	93.1	130.3	164.1
2005 4/.....	3.0	5.3	8.3	14.1	26.1	39.5	64.0	95.7	132.7	161.4
2004 4/.....	3.0	5.2	8.6	15.2	25.0	41.4	64.6	98.3	133.7	163.8
2003 4/.....	3.4	5.4	8.8	15.3	26.1	40.7	67.0	102.6	135.8	171.3
2002.....	3.7	5.7	9.6	15.8	28.8	44.8	69.4	99.9	138.5	171.2
2001.....	3.5	6.1	9.8	17.7	29.3	44.4	73.0	105.0	140.4	172.8
2000.....	3.6	6.1	10.7	18.6	31.4	48.7	72.8	106.5	147.9	175.5
1999.....	3.8	6.6	10.3	20.2	30.5	50.3	75.5	105.7	148.8	171.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)										
Black, both sexes										
2006 4/.....	6.0	10.0	14.6	26.0	42.9	57.3	83.4	109.2	127.8	143.5
2005 4/.....	5.8	9.2	16.0	24.9	38.7	58.5	81.2	103.1	127.4	136.8
2004 4/.....	5.2	9.1	16.8	26.3	40.8	60.7	93.3	115.8	134.7	154.0
2003 4/.....	6.3	9.5	17.3	23.5	41.1	59.4	90.5	117.4	139.1	150.5
2002.....	5.7	10.5	17.3	29.8	43.9	62.3	93.2	111.8	138.2	152.3
2001.....	6.6	10.4	16.5	30.2	44.9	67.8	90.7	116.7	142.7	155.7
2000.....	6.3	12.5	17.5	29.1	44.0	62.2	90.5	114.3	131.6	175.5
1999.....	6.5	10.4	18.2	33.2	45.9	64.6	90.4	119.1	139.3	162.0
Black, male										
2006 4/.....	8.0	12.2	19.2	34.4	56.0	76.2	110.7	144.5	167.7	194.8
2005 4/.....	6.8	11.9	19.9	33.1	46.7	73.4	108.2	139.0	172.1	164.5
2004 4/.....	5.9	11.3	21.4	31.5	51.2	80.4	122.7	139.2	168.6	193.7
2003 4/.....	8.0	11.9	22.5	32.0	51.1	73.4	105.1	137.8	166.8	184.5
2002.....	7.2	13.1	19.7	36.0	55.6	78.8	123.2	137.8	177.3	202.6
2001.....	8.2	11.9	21.0	37.5	56.9	87.9	114.1	141.2	191.0	206.5
2000.....	6.6	14.9	21.8	36.7	54.2	73.1	125.4	147.9	167.8	224.9
1999.....	8.4	13.3	22.5	41.2	55.2	80.7	114.6	159.1	164.8	220.3
Black, female										
2006 4/.....	4.2	8.0	10.7	19.0	32.6	43.4	64.7	88.1	106.7	122.5
2005 4/.....	5.0	6.7	12.8	18.2	32.4	47.4	62.8	81.6	103.9	125.7
2004 4/.....	4.6	7.2	12.9	21.9	32.8	46.1	73.4	101.9	116.8	138.2
2003 4/.....	4.8	7.4	12.9	16.6	33.2	49.0	80.6	105.3	124.6	137.3
2002.....	4.3	8.3	15.2	24.7	34.7	50.1	73.0	96.3	117.8	133.1
2001.....	5.3	9.2	12.7	24.2	35.4	52.9	75.1	102.1	118.1	136.8
2000.....	6.0	10.4	13.9	22.9	36.0	54.1	67.3	94.0	113.4	157.1
1999.....	4.8	8.0	14.5	26.6	38.6	52.7	74.3	95.0	126.6	140.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hodgkin's disease (C81)										
All races 3/, both sexes										
2006.....	0.4	0.3	0.4	0.5	0.7	0.9	1.3	2.1	2.3	2.6
2005.....	0.3	0.4	0.4	0.5	0.7	1.0	1.2	2.2	2.5	2.1
2004.....	0.3	0.4	0.4	0.7	0.8	1.0	1.4	1.8	2.1	2.2
2003.....	0.3	0.3	0.4	0.6	0.8	0.9	1.5	2.1	2.3	2.4
2002.....	0.3	0.4	0.5	0.5	0.8	1.0	1.6	1.7	2.3	2.7
2001.....	0.4	0.4	0.5	0.4	0.7	1.1	1.2	2.0	2.5	2.2
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	1.8	2.4	2.5
1999.....	0.3	0.4	0.5	0.7	0.9	1.3	1.8	2.0	2.3	2.4
All races 3/, male										
2006.....	0.6	0.3	0.6	0.8	0.9	1.2	1.6	2.3	3.0	3.7
2005.....	0.3	0.5	0.4	0.8	0.9	1.2	1.6	2.3	2.7	3.1
2004.....	0.3	0.5	0.5	0.9	1.0	1.4	1.9	2.0	3.2	3.2
2003.....	0.5	0.4	0.5	0.7	1.1	1.1	2.0	2.4	2.9	2.6
2002.....	0.4	0.5	0.6	0.7	1.0	1.4	2.2	2.0	3.2	3.6
2001.....	0.5	0.5	0.6	0.6	0.9	1.5	1.6	2.5	2.9	2.6
2000.....	0.4	0.5	0.6	0.5	1.1	1.3	1.9	2.3	2.7	3.7
1999.....	0.4	0.5	0.6	0.9	1.2	1.6	2.3	2.3	2.8	3.8
All races 3/, female										
2006.....	0.2	0.2	0.2	0.3	0.5	0.7	1.0	2.0	1.9	2.1
2005.....	0.2	0.4	0.3	0.3	0.5	0.8	1.0	2.1	2.3	1.6
2004.....	0.3	0.3	0.3	0.5	0.5	0.8	1.0	1.6	1.5	1.7
2003.....	0.2	0.2	0.4	0.5	0.6	0.6	1.2	1.8	1.9	2.3
2002.....	0.3	0.3	0.4	0.3	0.7	0.6	1.1	1.5	1.7	2.3
2001.....	0.3	0.4	0.4	0.3	0.6	0.7	0.9	1.7	2.2	2.0
2000.....	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.4	2.3	2.0
1999.....	0.3	0.3	0.4	0.5	0.5	0.9	1.5	1.8	2.0	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hodgkin's disease (C81)										
White, both sexes										
2006 4/.....	0.4	0.3	0.4	0.5	0.7	1.0	1.3	2.2	2.5	2.6
2005 4/.....	0.2	0.4	0.4	0.5	0.8	1.1	1.3	2.4	2.6	2.1
2004 4/.....	0.3	0.4	0.4	0.7	0.8	1.1	1.5	1.9	2.2	2.3
2003 4/.....	0.4	0.4	0.5	0.6	0.9	0.8	1.7	2.2	2.4	2.5
2002.....	0.4	0.4	0.5	0.5	0.8	1.0	1.7	1.8	2.4	2.9
2001.....	0.4	0.4	0.5	0.4	0.8	1.2	1.3	2.1	2.6	2.4
2000.....	0.3	0.4	0.5	0.5	0.9	1.0	1.6	2.0	2.5	2.6
1999.....	0.4	0.4	0.5	0.7	0.9	1.3	2.0	2.2	2.5	2.4
White, male										
2006 4/.....	0.6	0.3	0.6	0.7	1.0	1.2	1.6	2.4	3.1	3.7
2005 4/.....	0.3	0.5	0.4	0.7	1.0	1.3	1.6	2.4	2.9	3.4
2004 4/.....	0.4	0.5	0.6	0.9	1.0	1.3	2.0	2.1	3.4	3.5
2003 4/.....	0.5	0.4	0.5	0.7	1.1	1.1	2.0	2.4	3.0	2.7
2002.....	0.4	0.5	0.5	0.7	0.9	1.5	2.3	2.1	3.5	3.9
2001.....	0.5	0.5	0.7	0.6	0.9	1.6	1.8	2.5	3.0	2.7
2000.....	0.4	0.5	0.6	0.5	1.1	1.4	1.9	2.6	2.8	3.9
1999.....	0.5	0.5	0.6	0.9	1.3	1.7	2.4	2.5	3.0	3.8
White, female										
2006 4/.....	*	*	0.3	0.3	0.5	0.8	1.0	2.1	2.1	2.1
2005 4/.....	*	0.4	0.3	0.4	0.6	0.9	1.0	2.3	2.5	1.5
2004 4/.....	0.2	0.4	0.3	0.5	0.5	0.8	1.0	1.7	1.5	1.7
2003 4/.....	0.2	0.3	0.4	0.5	0.6	0.6	1.4	2.0	2.0	2.4
2002.....	0.3	0.3	0.5	*	0.7	0.6	1.2	1.6	1.7	2.5
2001.....	0.3	0.4	0.4	*	0.7	0.8	0.9	1.8	2.4	2.2
2000.....	0.3	0.3	0.4	0.5	0.7	0.7	1.3	1.6	2.4	2.1
1999.....	0.3	0.4	0.4	0.5	0.6	0.9	1.6	2.0	2.2	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

All races 3/, both sexes

2006.....	1.5	2.5	4.2	6.7	12.5	19.0	30.2	46.4	63.6	72.7
2005.....	1.6	2.4	4.3	7.6	12.4	19.2	31.6	47.0	64.6	75.4
2004.....	1.5	2.8	4.8	7.8	12.7	21.1	32.1	47.0	65.1	75.3
2003.....	1.7	3.0	4.7	8.1	14.3	21.6	32.9	49.4	65.5	79.0
2002.....	1.9	3.0	5.0	8.8	14.9	23.3	35.1	50.3	67.3	77.6
2001.....	2.0	3.4	5.6	9.6	15.6	23.1	36.2	51.0	70.1	80.1
2000.....	2.1	3.7	5.8	10.7	16.1	25.1	36.4	53.9	72.1	80.7
1999.....	2.1	3.6	6.2	10.9	16.3	25.7	38.5	54.8	72.8	78.0

All races 3/, male

2006.....	1.9	3.1	5.6	8.8	16.0	23.9	37.3	57.5	80.9	90.2
2005.....	2.2	3.1	5.7	9.7	15.0	24.6	40.1	59.2	79.7	100.9
2004.....	2.1	3.7	6.1	9.8	15.7	26.7	39.7	57.6	81.9	96.3
2003.....	2.2	3.9	6.0	10.1	18.7	27.9	41.2	61.0	81.8	104.0
2002.....	2.4	3.9	6.3	11.3	18.4	30.2	43.0	61.9	84.8	100.2
2001.....	2.7	4.4	7.4	11.9	19.1	29.3	45.0	61.4	87.8	107.1
2000.....	2.6	4.8	7.5	13.4	19.2	30.7	45.3	66.6	88.5	106.6
1999.....	2.8	4.8	7.9	13.5	19.6	30.9	46.8	68.4	85.9	104.7

All races 3/, female

2006.....	1.1	1.8	2.8	4.7	9.3	14.8	24.4	38.3	52.8	64.6
2005.....	1.1	1.7	3.0	5.7	10.0	14.6	24.8	38.1	55.2	63.7
2004.....	0.9	1.8	3.5	5.9	9.9	16.3	25.9	39.4	54.7	65.8
2003.....	1.2	2.1	3.5	6.1	10.4	16.2	26.2	41.3	55.5	68.0
2002.....	1.3	2.2	3.9	6.4	11.7	17.4	28.7	42.1	56.6	67.8
2001.....	1.3	2.5	3.8	7.5	12.4	17.8	29.2	43.7	59.5	68.8
2000.....	1.5	2.6	4.1	8.2	13.3	20.3	29.4	45.0	62.5	70.2
1999.....	1.4	2.4	4.6	8.3	13.3	21.2	31.9	45.3	65.2	67.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

White, both sexes

2006 4/.....	1.4	2.3	4.2	6.7	12.8	19.8	32.0	49.3	67.2	76.8
2005 4/.....	1.6	2.4	4.2	7.8	12.7	20.3	33.3	50.1	68.1	79.9
2004 4/.....	1.5	2.7	4.7	8.0	13.1	22.2	33.6	49.4	68.9	79.0
2003 4/.....	1.7	2.9	4.8	8.5	14.9	22.8	34.4	52.0	69.0	83.6
2002.....	1.8	3.0	5.0	9.0	15.6	24.5	36.6	53.0	70.6	82.4
2001.....	2.0	3.4	5.7	9.8	16.3	24.0	38.1	53.7	73.3	83.8
2000.....	2.1	3.6	5.9	11.2	17.0	26.4	38.3	56.7	76.2	84.8
1999.....	2.0	3.6	6.5	11.2	17.1	27.2	40.3	57.2	76.9	81.5

White, male

2006 4/.....	1.8	2.9	5.6	8.8	16.4	24.7	39.4	60.8	85.7	94.6
2005 4/.....	2.2	3.0	5.5	9.9	15.4	25.8	42.0	63.0	83.8	107.4
2004 4/.....	2.1	3.6	6.1	10.0	16.2	27.7	41.4	60.3	86.6	101.7
2003 4/.....	2.2	3.7	6.0	10.6	19.4	29.7	43.3	64.2	86.5	110.1
2002.....	2.3	4.0	6.2	11.6	19.1	31.7	44.8	65.2	88.9	106.1
2001.....	2.7	4.4	7.5	12.0	19.8	30.5	46.9	64.3	91.7	111.9
2000.....	2.7	4.7	7.6	13.8	20.1	32.4	46.9	70.1	93.1	112.5
1999.....	2.7	4.7	8.1	13.8	20.3	32.4	49.0	70.5	91.1	108.3

White, female

2006 4/.....	1.0	1.7	2.9	4.7	9.5	15.5	25.9	40.7	55.5	68.5
2005 4/.....	1.0	1.7	3.0	5.8	10.2	15.4	26.1	40.5	58.2	67.2
2004 4/.....	0.9	1.8	3.4	6.1	10.2	17.4	27.2	41.4	57.9	68.8
2003 4/.....	1.2	2.1	3.5	6.4	10.7	16.8	27.0	43.1	58.3	71.8
2002.....	1.3	2.1	3.7	6.5	12.3	18.2	29.9	44.3	59.5	72.1
2001.....	1.3	2.4	3.9	7.7	13.1	18.3	30.9	46.2	62.2	72.1
2000.....	1.4	2.6	4.3	8.8	14.1	21.2	31.3	47.2	66.1	73.5
1999.....	1.4	2.4	4.8	8.7	14.2	22.7	33.4	47.8	68.5	70.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other lymphoid, hematopoietic and related tissue (C82-C85,C96)

Black, both sexes

2006 4/.....	2.4	4.1	4.8	8.5	12.4	15.2	19.4	25.1	31.5	34.0
2005 4/.....	2.4	3.2	5.5	7.6	11.5	13.9	20.8	22.4	32.3	33.9
2004 4/.....	1.7	3.9	5.9	7.7	11.0	15.9	21.7	29.1	27.0	39.2
2003 4/.....	2.3	3.9	5.9	6.1	12.7	14.7	22.8	26.5	31.0	31.5
2002.....	2.7	3.5	6.1	9.3	12.8	17.7	25.7	27.1	35.6	31.2
2001.....	2.4	4.3	5.9	9.9	12.4	19.2	23.5	25.9	38.7	39.6
2000.....	2.3	4.6	5.5	8.9	12.6	16.6	22.7	26.3	28.5	36.9
1999.....	2.8	4.1	5.7	10.6	11.7	17.0	23.9	32.1	31.5	40.8

Black, male

2006 4/.....	3.4	5.9	7.1	11.9	16.4	20.2	24.0	30.0	34.9	42.0
2005 4/.....	2.8	4.4	8.2	10.5	14.7	17.8	29.7	25.9	37.2	40.0
2004 4/.....	2.2	5.4	7.8	10.0	14.4	19.6	28.5	31.8	34.4	42.7
2003 4/.....	3.0	5.8	7.5	7.7	16.0	17.2	24.6	28.8	34.7	34.4
2002.....	3.8	4.0	7.4	11.5	16.6	23.2	34.4	31.7	42.0	42.9
2001.....	3.4	5.1	8.7	12.4	15.5	24.9	31.6	31.1	48.2	55.6
2000.....	2.6	6.3	7.9	13.7	15.7	19.3	32.1	31.2	32.0	46.1
1999.....	3.9	5.6	8.0	13.9	15.7	23.4	28.5	47.5	28.8	59.8

Black, female

2006 4/.....	1.5	2.7	2.8	5.7	9.2	11.6	16.3	22.2	29.7	30.7
2005 4/.....	2.0	2.1	3.2	5.2	8.9	10.9	14.7	20.3	29.7	31.5
2004 4/.....	1.3	2.5	4.3	5.9	8.3	13.1	17.1	27.5	23.1	37.8
2003 4/.....	1.6	2.3	4.5	4.7	10.1	12.8	21.5	25.2	29.0	30.4
2002.....	1.7	3.1	5.0	7.6	9.8	13.7	19.9	24.4	32.2	26.7
2001.....	1.6	3.6	3.6	8.0	9.9	15.0	18.2	22.7	33.9	33.7
2000.....	2.1	3.2	3.5	5.0	10.1	14.6	16.4	23.4	26.7	33.5
1999.....	1.8	2.8	3.7	7.9	8.7	12.3	20.9	22.9	32.8	33.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
All races 3/, both sexes										
2006.....	1.5	2.5	4.2	6.7	12.5	19.0	30.1	46.2	63.4	72.3
2005.....	1.6	2.4	4.3	7.6	12.3	19.2	31.5	46.9	64.4	75.2
2004.....	1.5	2.8	4.8	7.7	12.6	21.1	32.0	46.9	64.8	75.0
2003.....	1.7	3.0	4.7	8.1	14.3	21.6	32.8	49.3	65.4	78.8
2002.....	1.9	3.0	5.0	8.8	14.9	23.2	35.0	50.2	67.0	77.3
2001.....	2.0	3.4	5.5	9.6	15.6	23.0	36.0	50.9	69.8	79.6
2000.....	2.1	3.6	5.8	10.6	16.1	25.0	36.3	53.7	71.8	80.2
1999.....	2.1	3.5	6.2	10.8	16.3	25.6	38.4	54.6	72.6	77.7
All races 3/, male										
2006.....	1.9	3.1	5.6	8.8	16.0	23.8	37.2	57.3	80.7	89.6
2005.....	2.2	3.1	5.7	9.7	15.0	24.5	39.9	59.0	79.4	100.5
2004.....	2.1	3.7	6.1	9.8	15.7	26.7	39.6	57.5	81.6	95.8
2003.....	2.2	3.9	6.0	10.1	18.7	27.8	41.0	60.7	81.8	103.6
2002.....	2.4	3.9	6.2	11.3	18.4	30.0	42.8	61.8	84.4	99.7
2001.....	2.7	4.4	7.4	11.9	19.1	29.2	44.7	61.3	87.3	106.6
2000.....	2.6	4.7	7.5	13.3	19.2	30.5	45.2	66.3	88.1	105.9
1999.....	2.8	4.7	7.9	13.5	19.6	30.8	46.8	68.1	85.7	104.4
All races 3/, female										
2006.....	1.1	1.8	2.8	4.7	9.3	14.8	24.3	38.2	52.6	64.2
2005.....	1.1	1.7	3.0	5.7	9.9	14.5	24.7	38.1	55.2	63.5
2004.....	0.9	1.8	3.5	5.8	9.8	16.2	25.8	39.3	54.5	65.7
2003.....	1.2	2.1	3.5	6.1	10.4	16.2	26.1	41.2	55.4	67.8
2002.....	1.3	2.2	3.8	6.4	11.7	17.3	28.7	41.9	56.4	67.6
2001.....	1.3	2.5	3.8	7.5	12.4	17.7	29.1	43.6	59.3	68.3
2000.....	1.5	2.6	4.1	8.2	13.3	20.3	29.2	45.0	62.2	69.8
1999.....	1.4	2.4	4.5	8.3	13.3	21.1	31.8	45.2	65.0	66.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
White, both sexes										
2006 4/.....	1.4	2.3	4.2	6.7	12.8	19.8	32.0	49.1	67.0	76.4
2005 4/.....	1.6	2.4	4.2	7.8	12.6	20.2	33.2	50.0	67.9	79.7
2004 4/.....	1.5	2.7	4.7	7.9	13.1	22.2	33.5	49.2	68.6	78.8
2003 4/.....	1.7	2.9	4.7	8.4	14.9	22.8	34.2	51.8	69.0	83.3
2002.....	1.8	3.0	4.9	9.0	15.6	24.5	36.5	52.9	70.4	82.1
2001.....	2.0	3.4	5.7	9.8	16.2	23.9	37.9	53.6	73.0	83.4
2000.....	2.1	3.6	5.9	11.2	16.9	26.3	38.1	56.5	75.8	84.3
1999.....	2.0	3.6	6.4	11.1	17.0	27.1	40.2	57.0	76.7	81.2
White, male										
2006 4/.....	1.8	2.9	5.6	8.7	16.3	24.6	39.3	60.6	85.5	94.1
2005 4/.....	2.2	3.0	5.5	9.8	15.3	25.7	41.9	62.8	83.6	107.0
2004 4/.....	2.1	3.6	6.1	9.9	16.2	27.7	41.3	60.1	86.2	101.2
2003 4/.....	2.1	3.7	6.0	10.6	19.4	29.6	43.1	63.9	86.5	109.7
2002.....	2.3	4.0	6.2	11.6	19.1	31.6	44.5	65.0	88.6	105.6
2001.....	2.6	4.4	7.4	12.0	19.7	30.3	46.6	64.2	91.2	111.5
2000.....	2.7	4.7	7.6	13.7	20.0	32.2	46.8	69.8	92.6	111.8
1999.....	2.7	4.7	8.1	13.7	20.2	32.3	48.9	70.2	90.8	108.0
White, female										
2006 4/.....	1.0	1.7	2.9	4.7	9.5	15.5	25.8	40.5	55.3	68.1
2005 4/.....	1.0	1.7	3.0	5.8	10.2	15.3	26.0	40.4	58.1	67.1
2004 4/.....	0.9	1.8	3.4	6.0	10.2	17.3	27.1	41.3	57.7	68.7
2003 4/.....	1.2	2.1	3.5	6.4	10.7	16.7	27.0	43.0	58.2	71.6
2002.....	1.3	2.1	3.7	6.5	12.3	18.2	29.9	44.2	59.3	71.8
2001.....	1.3	2.4	3.9	7.7	13.0	18.3	30.9	46.0	62.0	71.6
2000.....	1.4	2.6	4.3	8.8	14.1	21.2	31.1	47.2	65.8	73.1
1999.....	1.4	2.4	4.8	8.7	14.1	22.6	33.3	47.7	68.3	70.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Non-Hodgkin's lymphoma (C82-C85)										
Black, both sexes										
2006 4/.....	2.4	4.1	4.7	8.4	12.3	15.2	19.3	25.1	31.5	34.0
2005 4/.....	2.4	3.2	5.5	7.5	11.5	13.9	20.6	22.4	32.0	33.7
2004 4/.....	1.7	3.9	5.9	7.6	11.0	15.9	21.7	28.9	26.5	38.6
2003 4/.....	2.3	3.9	5.9	6.1	12.7	14.6	22.8	26.5	31.0	31.2
2002.....	2.7	3.5	6.1	9.3	12.8	17.6	25.7	26.8	35.3	30.6
2001.....	2.4	4.3	5.9	9.8	12.3	19.1	23.4	25.7	37.9	39.3
2000.....	2.3	4.5	5.5	8.8	12.6	16.6	22.7	25.9	28.2	36.3
1999.....	2.8	4.1	5.6	10.5	11.7	17.0	23.9	32.0	31.5	40.5
Black, male										
2006 4/.....	3.4	5.9	7.0	11.9	16.3	20.2	23.7	30.0	34.9	42.0
2005 4/.....	2.8	4.4	8.2	10.4	14.7	17.8	29.1	25.9	37.2	40.0
2004 4/.....	2.2	5.4	7.8	10.0	14.4	19.6	28.5	31.8	34.4	41.7
2003 4/.....	3.0	5.7	7.5	7.7	16.0	17.2	24.6	28.8	34.7	34.4
2002.....	3.8	4.0	7.4	11.5	16.6	22.9	34.4	31.7	41.3	42.9
2001.....	3.4	5.1	8.7	12.2	15.5	24.6	31.6	30.7	47.4	54.5
2000.....	2.6	6.1	7.9	13.5	15.7	19.3	32.1	30.7	32.0	43.8
1999.....	3.9	5.6	7.9	13.7	15.7	23.4	28.5	47.0	28.8	59.8
Black, female										
2006 4/.....	1.5	2.7	2.8	5.6	9.2	11.6	16.3	22.2	29.7	30.7
2005 4/.....	2.0	2.1	3.2	5.2	8.9	10.9	14.7	20.3	29.3	31.1
2004 4/.....	1.3	2.5	4.2	5.6	8.3	13.1	17.1	27.2	22.3	37.4
2003 4/.....	1.6	2.3	4.5	4.7	10.1	12.6	21.5	25.2	29.0	30.0
2002.....	1.7	3.1	5.0	7.6	9.8	13.7	19.9	23.9	32.2	25.9
2001.....	1.6	3.6	3.6	7.8	9.8	15.0	17.9	22.7	33.1	33.7
2000.....	2.1	3.2	3.5	5.0	10.1	14.6	16.4	23.1	26.3	33.5
1999.....	1.7	2.8	3.7	7.9	8.7	12.3	20.9	22.9	32.8	33.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant immunoproliferative diseases (C88)

All races 3/, both sexes

2006.....	*	*	*	*	*	0.3	0.3	0.9	1.2	1.9
2005.....	*	*	*	*	0.2	0.3	0.4	1.0	1.2	2.1
2004.....	*	*	*	*	*	0.3	0.6	0.9	1.1	1.5
2003.....	*	*	*	*	0.2	0.2	0.8	1.2	1.1	2.2
2002.....	*	*	*	*	0.2	0.3	0.6	1.2	1.7	2.0
2001.....	*	*	*	*	0.2	0.4	0.6	0.8	1.6	2.0
2000.....	*	*	*	*	*	0.3	0.6	0.9	1.7	2.1
1999.....	*	*	*	*	0.4	0.3	0.7	1.2	1.6	2.7

All races 3/, male

2006.....	*	*	*	*	*	0.4	*	1.3	1.9	3.0
2005.....	*	*	*	*	*	*	*	1.3	1.9	3.8
2004.....	*	*	*	*	*	*	0.8	1.2	1.6	2.5
2003.....	*	*	*	*	*	*	1.2	1.7	1.7	3.6
2002.....	*	*	*	*	0.4	0.5	0.9	1.7	2.9	2.8
2001.....	*	*	*	*	*	0.6	0.7	1.1	2.2	2.9
2000.....	*	*	*	*	*	*	0.8	1.3	2.4	3.4
1999.....	*	*	*	*	0.5	0.5	1.1	1.7	2.3	3.9

All races 3/, female

2006.....	*	*	*	*	*	*	*	0.5	0.8	1.4
2005.....	*	*	*	*	*	*	*	0.7	0.8	1.3
2004.....	*	*	*	*	*	*	*	0.7	0.8	1.1
2003.....	*	*	*	*	*	*	0.4	0.8	0.7	1.6
2002.....	*	*	*	*	*	*	*	0.8	1.0	1.6
2001.....	*	*	*	*	*	*	0.5	0.6	1.2	1.7
2000.....	*	*	*	*	*	*	0.4	0.5	1.4	1.6
1999.....	*	*	*	*	*	*	0.5	0.9	1.2	2.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malignant immunoproliferative diseases (C88)										
White, both sexes										
2006 4/.....	*	*	*	*	*	0.3	0.3	1.0	1.3	2.1
2005 4/.....	*	*	*	*	0.2	0.3	0.4	1.1	1.3	2.3
2004 4/.....	*	*	*	*	*	0.3	0.6	1.0	1.2	1.7
2003 4/.....	*	*	*	*	*	0.2	0.9	1.3	1.2	2.4
2002.....	*	*	*	*	0.3	0.3	0.6	1.3	1.8	2.1
2001.....	*	*	*	*	*	0.4	0.6	0.9	1.7	2.2
2000.....	*	*	*	*	*	0.3	0.6	0.9	1.8	2.2
1999.....	*	*	*	*	0.4	0.4	0.8	1.2	1.8	2.9
White, male										
2006 4/.....	*	*	*	*	*	*	*	1.4	2.0	3.3
2005 4/.....	*	*	*	*	*	*	*	1.4	2.0	4.2
2004 4/.....	*	*	*	*	*	*	0.9	1.3	1.6	2.8
2003 4/.....	*	*	*	*	*	*	1.3	1.8	1.9	4.0
2002.....	*	*	*	*	0.4	*	1.0	1.8	3.0	2.9
2001.....	*	*	*	*	*	0.6	0.7	1.3	2.4	3.1
2000.....	*	*	*	*	*	*	0.9	1.4	2.5	3.5
1999.....	*	*	*	*	0.5	0.6	1.1	1.7	2.4	4.3
White, female										
2006 4/.....	*	*	*	*	*	*	*	0.6	0.9	1.6
2005 4/.....	*	*	*	*	*	*	*	0.9	0.9	1.5
2004 4/.....	*	*	*	*	*	*	*	0.8	0.9	1.2
2003 4/.....	*	*	*	*	*	*	0.5	0.9	0.7	1.7
2002.....	*	*	*	*	*	*	*	0.8	1.0	1.7
2001.....	*	*	*	*	*	*	0.6	0.7	1.3	1.8
2000.....	*	*	*	*	*	*	*	0.6	1.4	1.7
1999.....	*	*	*	*	*	*	0.5	0.9	1.4	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and malignant plasma cell neoplasms (C90)										
All races 3/, both sexes										
2006.....	0.5	1.1	2.4	4.2	7.4	11.5	18.7	24.9	32.1	31.9
2005.....	0.4	1.1	2.2	4.3	8.2	12.6	19.2	25.5	32.2	31.5
2004.....	0.5	1.0	2.2	4.5	7.6	12.1	18.4	27.4	32.8	31.6
2003.....	0.5	1.1	2.4	4.3	7.8	12.9	19.6	27.4	33.4	34.1
2002.....	0.4	1.2	2.4	4.7	8.0	13.8	20.5	26.7	34.1	34.7
2001.....	0.5	1.2	2.3	5.0	8.3	13.6	20.5	27.4	33.1	34.2
2000.....	0.5	1.2	2.6	4.4	8.3	14.1	19.8	27.4	34.6	35.7
1999.....	0.5	1.2	2.4	5.4	8.7	13.8	19.8	26.3	34.5	33.9
All races 3/, male										
2006.....	0.6	1.2	2.8	5.0	9.0	13.9	22.6	31.7	41.9	42.8
2005.....	0.6	1.3	2.7	5.1	9.2	15.9	23.4	32.2	42.7	42.3
2004.....	0.5	1.1	2.7	5.1	9.1	14.9	22.3	33.8	41.2	42.1
2003.....	0.7	1.3	3.0	5.3	9.8	15.7	23.1	32.2	41.6	46.8
2002.....	0.5	1.5	2.9	5.5	9.8	16.4	25.0	33.8	43.3	48.0
2001.....	0.6	1.5	2.8	5.8	9.9	16.2	24.6	34.5	41.2	45.1
2000.....	0.6	1.4	2.8	4.8	9.2	16.6	23.4	34.4	43.7	48.3
1999.....	0.6	1.3	2.9	6.4	10.5	16.0	23.1	32.4	43.2	47.4
All races 3/, female										
2006.....	0.4	0.9	1.9	3.5	6.0	9.5	15.5	20.0	26.0	26.8
2005.....	0.3	0.9	1.7	3.5	7.2	9.7	15.8	20.6	25.7	26.5
2004.....	0.4	0.9	1.8	3.9	6.1	9.6	15.2	22.8	27.6	26.9
2003.....	0.4	0.9	1.9	3.4	5.9	10.5	16.8	23.9	28.4	28.5
2002.....	0.4	1.0	1.9	3.9	6.4	11.6	16.9	21.7	28.6	29.0
2001.....	0.4	0.8	1.9	4.2	7.0	11.3	17.2	22.5	28.3	29.7
2000.....	0.4	1.0	2.3	4.1	7.5	12.0	16.9	22.6	29.3	30.6
1999.....	0.4	1.1	2.0	4.6	7.1	11.8	17.3	22.1	29.4	28.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and malignant plasma cell neoplasms (C90)										
White, both sexes										
2006 4/.....	0.5	0.8	2.0	3.7	6.5	10.6	17.3	23.9	31.1	30.7
2005 4/.....	0.4	0.9	1.8	3.6	7.5	11.4	17.9	24.4	31.4	30.6
2004 4/.....	0.4	0.9	1.9	3.9	6.9	11.0	16.8	26.0	31.3	30.3
2003 4/.....	0.5	0.9	2.0	3.9	7.2	11.9	18.2	25.6	31.9	32.6
2002.....	0.4	1.1	2.1	4.1	7.2	12.7	19.2	25.6	33.1	33.6
2001.....	0.4	1.0	2.0	4.3	7.3	12.5	19.2	26.1	31.3	32.8
2000.....	0.4	1.0	2.2	3.7	7.6	13.1	18.5	25.6	33.4	33.0
1999.....	0.4	1.1	2.0	4.6	7.5	12.8	18.4	24.5	33.2	31.9
White, male										
2006 4/.....	0.6	1.0	2.5	4.5	7.8	12.9	21.0	30.6	40.8	41.7
2005 4/.....	0.5	1.0	2.3	4.3	8.7	14.8	22.0	30.8	41.9	41.7
2004 4/.....	0.5	1.0	2.3	4.6	8.5	13.8	20.8	32.5	39.6	40.7
2003 4/.....	0.6	1.1	2.5	4.8	9.1	14.7	22.1	31.0	40.0	45.8
2002.....	0.4	1.3	2.6	4.9	9.0	15.4	23.9	32.9	42.3	47.3
2001.....	0.5	1.3	2.5	5.0	8.7	15.0	23.4	33.4	39.3	44.0
2000.....	0.5	1.2	2.4	4.1	8.6	15.6	21.9	32.5	42.9	46.7
1999.....	0.4	1.2	2.6	5.5	9.5	15.2	21.4	30.8	41.9	45.4
White, female										
2006 4/.....	0.4	0.7	1.5	3.0	5.3	8.5	14.3	18.9	24.9	25.5
2005 4/.....	0.3	0.7	1.4	3.0	6.4	8.4	14.6	19.6	24.8	25.5
2004 4/.....	0.3	0.8	1.5	3.3	5.5	8.5	13.4	21.2	26.1	25.6
2003 4/.....	0.3	0.7	1.6	3.0	5.3	9.4	15.1	21.7	26.9	26.7
2002.....	0.3	0.8	1.5	3.4	5.6	10.4	15.4	20.4	27.4	27.7
2001.....	*	0.7	1.6	3.6	6.0	10.2	15.9	21.0	26.5	28.1
2000.....	0.2	0.8	2.0	3.3	6.7	10.9	15.8	20.8	27.8	27.4
1999.....	0.3	0.9	1.5	3.8	5.7	10.7	16.0	20.1	28.1	26.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple myeloma and malignant plasma cell neoplasms (C90)										
Black, both sexes										
2006 4/.....	0.9	2.6	5.4	9.4	17.1	23.0	34.1	43.7	52.5	51.6
2005 4/.....	0.9	2.3	5.0	9.7	16.0	25.4	34.7	44.1	51.2	49.2
2004 4/.....	1.0	2.0	5.2	10.2	15.3	24.3	38.0	48.2	58.9	53.9
2003 4/.....	1.2	2.7	5.7	9.3	15.0	25.3	37.3	51.9	58.6	59.8
2002.....	1.0	2.9	5.6	10.5	16.2	26.3	37.7	44.1	54.2	54.3
2001.....	1.5	2.6	4.8	11.8	19.0	25.5	38.1	47.6	60.9	57.1
2000.....	1.1	3.1	6.1	11.6	16.5	27.2	37.6	52.8	57.0	70.1
1999.....	1.4	2.4	6.4	13.6	19.6	24.1	37.5	50.2	56.1	62.5
Black, male										
2006 4/.....	*	3.0	6.3	11.5	22.3	27.2	46.8	54.6	67.8	65.2
2005 4/.....	*	3.0	6.2	12.6	17.1	30.4	42.4	57.7	67.4	61.4
2004 4/.....	*	2.1	6.1	11.2	18.1	30.5	44.0	56.9	72.5	71.5
2003 4/.....	1.5	3.1	7.5	12.1	19.5	29.1	39.4	56.6	73.1	73.0
2002.....	*	3.5	5.9	12.2	20.0	30.7	44.7	51.4	69.5	70.8
2001.....	1.6	3.4	5.3	14.5	23.0	29.7	45.1	57.1	79.5	69.2
2000.....	*	3.3	7.0	12.2	18.4	31.0	48.1	66.2	67.4	76.1
1999.....	1.9	2.6	6.4	15.9	20.8	26.0	46.5	58.4	73.2	80.9
Black, female										
2006 4/.....	*	2.3	4.7	7.8	13.1	19.9	25.4	37.2	44.5	46.1
2005 4/.....	*	1.7	4.0	7.4	15.2	21.7	29.4	36.0	42.7	44.3
2004 4/.....	*	1.9	4.4	9.3	13.1	19.6	33.8	43.1	51.8	46.9
2003 4/.....	*	2.4	4.3	7.0	11.4	22.5	35.9	49.1	51.1	54.7
2002.....	*	2.3	5.3	9.2	13.3	23.1	32.9	39.8	46.2	48.1
2001.....	1.5	1.9	4.3	9.5	15.8	22.3	33.4	41.9	51.5	52.6
2000.....	*	2.8	5.4	11.2	15.0	24.4	30.6	44.7	51.8	67.8
1999.....	*	2.3	6.4	11.8	18.6	22.7	31.5	45.2	47.5	55.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
All races 3/, both sexes										
2006.....	1.8	2.5	4.1	7.0	11.6	19.9	31.3	44.7	62.9	76.9
2005.....	1.7	2.7	4.1	6.5	11.8	19.8	31.3	45.7	61.0	78.7
2004.....	1.8	2.5	4.2	7.1	12.0	19.9	31.8	45.6	61.5	78.8
2003.....	2.1	2.7	4.0	7.0	12.4	19.6	32.5	47.4	62.9	80.3
2002.....	1.9	2.8	4.5	7.4	12.9	20.9	32.4	45.7	63.4	80.5
2001.....	1.9	3.0	4.6	7.6	13.1	20.6	33.3	47.7	61.3	81.4
2000.....	1.9	3.1	4.5	7.7	12.7	21.7	33.2	47.4	65.1	83.6
1999.....	1.9	3.2	4.4	8.5	13.1	21.8	33.1	46.3	62.7	82.4
All races 3/, male										
2006.....	2.0	2.7	4.9	8.7	15.1	26.3	41.6	64.8	91.8	105.1
2005.....	1.8	2.9	4.4	8.3	15.4	26.1	43.9	66.0	87.9	112.4
2004.....	2.1	2.8	5.0	9.0	15.7	26.7	43.9	64.1	86.6	110.7
2003.....	2.4	3.2	4.9	8.9	15.8	26.8	44.9	66.3	88.8	111.8
2002.....	2.2	3.2	5.3	9.2	16.1	28.0	45.8	65.0	90.0	114.5
2001.....	2.1	3.4	5.2	9.2	17.4	27.6	44.9	66.2	86.9	117.5
2000.....	2.1	3.6	5.1	9.7	16.0	29.2	46.2	65.1	93.4	119.8
1999.....	2.1	3.6	5.2	10.1	16.8	29.0	45.7	64.6	87.6	119.2
All races 3/, female										
2006.....	1.7	2.3	3.3	5.3	8.5	14.3	22.8	30.0	44.8	63.7
2005.....	1.6	2.5	3.7	4.9	8.5	14.4	21.1	31.0	44.3	63.3
2004.....	1.6	2.2	3.3	5.4	8.6	14.1	22.1	32.3	46.0	64.5
2003.....	1.7	2.2	3.2	5.2	9.4	13.3	22.7	33.9	47.0	66.4
2002.....	1.6	2.5	3.8	5.7	10.0	14.7	21.8	32.1	47.3	65.7
2001.....	1.7	2.7	4.0	6.0	9.3	14.6	24.1	34.8	45.9	66.3
2000.....	1.7	2.6	3.9	5.9	9.7	15.3	22.9	35.1	48.4	68.8
1999.....	1.7	2.8	3.6	7.0	9.8	15.6	23.2	33.6	48.1	67.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
White, both sexes										
2006 4/.....	1.9	2.5	4.2	7.1	11.8	20.7	32.5	46.0	66.0	79.9
2005 4/.....	1.7	2.7	4.0	6.7	12.1	20.5	32.7	47.5	63.7	82.4
2004 4/.....	1.8	2.5	4.2	7.3	12.2	20.5	32.5	47.1	63.6	81.6
2003 4/.....	2.1	2.8	4.0	7.1	12.7	20.3	33.6	49.3	65.0	83.4
2002.....	2.0	2.7	4.5	7.4	13.1	21.8	33.6	47.0	65.7	83.0
2001.....	1.9	3.1	4.5	7.6	13.5	20.9	34.5	49.1	63.8	84.7
2000.....	1.9	3.0	4.5	7.8	12.8	22.6	34.4	49.2	67.6	86.0
1999.....	1.9	3.2	4.3	8.7	13.3	22.2	34.2	48.0	64.1	85.4
White, male										
2006 4/.....	1.9	2.8	5.0	8.9	15.3	27.1	43.2	66.3	96.1	108.7
2005 4/.....	1.7	2.9	4.5	8.5	15.8	27.0	45.6	67.9	91.4	118.7
2004 4/.....	2.1	2.8	4.9	9.2	16.0	27.2	44.4	66.2	89.8	115.3
2003 4/.....	2.4	3.4	4.8	9.0	16.4	27.7	46.4	69.2	92.8	116.9
2002.....	2.3	3.0	5.3	9.3	16.3	29.0	47.2	67.0	93.3	119.1
2001.....	2.0	3.5	5.1	9.4	17.8	27.8	46.5	68.3	90.1	123.1
2000.....	2.2	3.5	5.0	9.8	16.1	30.5	47.4	67.5	96.7	123.6
1999.....	2.1	3.5	5.1	10.3	17.0	29.5	47.1	66.7	90.2	124.1
White, female										
2006 4/.....	1.8	2.3	3.4	5.5	8.5	15.0	23.6	30.8	46.9	66.4
2005 4/.....	1.6	2.5	3.6	4.9	8.7	14.7	22.0	32.4	46.4	65.7
2004 4/.....	1.6	2.3	3.4	5.4	8.6	14.5	22.6	33.2	47.3	66.4
2003 4/.....	1.7	2.3	3.2	5.4	9.3	13.8	23.1	34.9	47.9	68.6
2002.....	1.7	2.5	3.8	5.6	10.1	15.4	22.6	32.8	48.9	67.3
2001.....	1.7	2.7	3.9	6.0	9.5	14.8	24.7	35.4	47.9	68.6
2000.....	1.6	2.5	4.0	5.9	9.8	15.7	23.9	36.3	50.2	70.8
1999.....	1.8	2.9	3.6	7.2	9.8	15.8	23.9	34.9	48.7	69.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Leukemia (C91-C95)										
Black, both sexes										
2006 4/.....	2.2	2.6	4.1	7.3	12.8	17.9	28.6	39.2	41.7	55.5
2005 4/.....	2.0	3.1	5.0	6.8	10.7	18.7	24.3	35.1	43.2	52.1
2004 4/.....	2.2	2.9	5.3	7.7	13.8	19.4	32.5	37.1	47.0	60.1
2003 4/.....	2.5	2.5	5.1	7.2	12.8	17.9	29.6	37.3	48.0	59.2
2002.....	1.7	3.6	5.1	9.1	13.8	17.3	28.6	39.2	46.6	64.7
2001.....	2.3	3.0	5.6	7.9	12.3	22.3	28.0	41.3	42.0	57.1
2000.....	2.4	4.5	5.2	7.9	14.1	17.9	28.4	34.7	43.8	66.0
1999.....	2.0	3.5	5.4	8.1	13.6	22.2	27.9	35.0	51.1	55.8
Black, male										
2006 4/.....	3.0	2.6	5.2	9.7	16.6	26.9	38.4	58.6	61.6	84.9
2005 4/.....	2.2	3.7	5.1	8.3	14.0	24.5	34.2	52.7	66.7	62.3
2004 4/.....	2.3	3.1	7.1	9.4	17.6	28.3	48.6	48.2	58.8	79.5
2003 4/.....	3.1	2.7	6.8	10.7	14.5	24.9	39.4	50.2	58.3	77.2
2002.....	1.7	4.8	5.7	11.0	17.2	23.4	42.4	52.4	64.2	86.8
2001.....	2.7	2.8	6.6	9.9	17.1	32.3	36.4	51.6	61.0	77.2
2000.....	2.5	4.9	6.2	10.0	18.6	22.4	43.5	49.6	64.1	98.0
1999.....	2.2	4.4	6.9	10.3	17.2	29.1	38.0	51.3	61.0	77.3
Black, female										
2006 4/.....	1.4	2.6	3.2	5.3	9.8	11.3	21.8	27.6	31.1	43.5
2005 4/.....	1.8	2.6	4.9	5.5	8.2	14.5	17.5	24.5	30.8	48.0
2004 4/.....	2.0	2.6	3.8	6.3	10.9	12.8	21.5	30.5	40.8	52.4
2003 4/.....	2.1	2.4	3.8	4.3	11.4	12.8	23.0	29.6	42.6	52.3
2002.....	1.6	2.6	4.7	7.6	11.2	12.8	19.3	31.3	37.4	56.3
2001.....	2.0	3.3	4.7	6.3	8.5	15.0	22.4	35.2	32.3	49.7
2000.....	2.3	4.2	4.4	6.2	10.5	14.6	18.4	25.6	33.5	54.1
1999.....	1.7	2.6	4.2	6.2	10.8	17.1	21.2	25.2	46.2	47.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lymphoid leukemia (C91)										
All races 3/, both sexes										
2006.....	0.4	0.6	0.9	1.8	3.1	4.8	8.2	12.0	18.1	28.9
2005.....	0.4	0.7	0.9	1.5	3.1	5.2	8.1	12.3	17.5	29.2
2004.....	0.4	0.5	1.0	1.9	3.0	5.0	8.4	12.0	17.0	28.9
2003.....	0.4	0.6	0.9	1.8	3.0	5.2	8.5	12.9	18.8	29.6
2002.....	0.5	0.7	1.1	1.8	3.3	5.3	8.2	12.2	18.6	30.1
2001.....	0.4	0.6	1.1	1.9	3.3	5.4	8.7	12.9	18.2	31.2
2000.....	0.4	0.6	1.0	1.8	3.2	5.5	8.9	13.1	19.5	32.0
1999.....	0.3	0.6	1.1	2.2	3.5	6.0	9.2	13.4	19.7	30.8
All races 3/, male										
2006.....	0.5	0.7	1.2	2.4	4.4	6.4	11.8	18.6	27.5	40.2
2005.....	0.4	0.8	1.2	2.2	4.5	7.4	12.3	18.1	25.4	41.6
2004.....	0.5	0.7	1.3	2.7	4.6	6.8	12.0	17.0	24.7	41.7
2003.....	0.6	0.8	1.2	2.7	4.2	8.2	12.5	18.5	27.7	41.7
2002.....	0.6	0.8	1.5	2.4	4.4	7.8	12.5	17.7	26.9	43.0
2001.....	0.5	0.7	1.4	2.6	4.9	8.1	12.2	18.4	25.7	43.8
2000.....	0.5	0.8	1.2	2.7	4.6	8.2	12.8	18.9	28.4	43.8
1999.....	0.4	0.7	1.5	3.0	5.1	8.6	13.1	19.1	28.7	45.6
All races 3/, female										
2006.....	0.3	0.5	0.7	1.1	1.9	3.3	5.3	7.3	12.3	23.6
2005.....	0.3	0.5	0.7	0.9	1.7	3.4	4.7	8.1	12.6	23.6
2004.....	0.3	0.4	0.8	1.1	1.6	3.3	5.4	8.3	12.3	23.1
2003.....	0.3	0.5	0.5	1.0	1.9	2.7	5.3	8.9	13.3	24.3
2002.....	0.3	0.5	0.7	1.3	2.3	3.2	4.8	8.3	13.6	24.5
2001.....	0.3	0.5	0.7	1.1	1.9	3.1	5.8	9.0	13.8	25.9
2000.....	0.2	0.5	0.7	1.0	1.9	3.2	5.8	9.1	14.2	27.3
1999.....	0.3	0.5	0.6	1.4	2.1	3.7	6.2	9.4	14.4	24.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lymphoid leukemia (C91)										
White, both sexes										
2006 4/.....	0.4	0.6	1.0	1.8	3.1	4.8	8.4	12.5	19.1	30.2
2005 4/.....	0.4	0.6	1.0	1.6	3.1	5.1	8.6	12.7	18.3	31.0
2004 4/.....	0.4	0.6	1.0	2.0	3.0	5.1	8.5	12.3	17.6	30.2
2003 4/.....	0.4	0.6	0.9	1.9	3.1	5.4	8.7	13.4	19.7	31.0
2002.....	0.5	0.7	1.2	1.8	3.3	5.5	8.5	12.3	19.5	31.6
2001.....	0.4	0.6	1.0	1.9	3.4	5.3	9.0	13.2	19.1	32.8
2000.....	0.4	0.6	1.0	1.9	3.2	5.7	9.1	13.6	20.3	33.1
1999.....	0.4	0.6	1.1	2.1	3.5	6.1	9.6	13.9	20.3	32.0
White, male										
2006 4/.....	0.4	0.7	1.2	2.4	4.3	6.5	11.9	19.0	28.8	42.0
2005 4/.....	0.5	0.8	1.2	2.2	4.7	7.2	13.0	18.3	26.3	44.4
2004 4/.....	0.5	0.7	1.3	2.9	4.5	6.8	11.9	17.4	25.7	43.7
2003 4/.....	0.6	0.8	1.1	2.7	4.4	8.5	13.0	19.3	29.2	43.5
2002.....	0.7	0.8	1.6	2.4	4.5	8.2	12.8	17.8	28.3	45.1
2001.....	0.5	0.7	1.4	2.7	4.9	8.0	12.7	19.0	26.6	46.2
2000.....	0.5	0.8	1.2	2.8	4.6	8.4	13.1	19.5	29.5	45.0
1999.....	0.4	0.7	1.6	3.0	5.2	8.7	13.5	19.8	29.8	47.7
White, female										
2006 4/.....	0.4	0.5	0.7	1.1	1.9	3.4	5.4	7.6	12.9	24.7
2005 4/.....	0.3	0.4	0.7	0.9	1.7	3.3	4.9	8.5	13.2	24.9
2004 4/.....	0.3	0.4	0.8	1.1	1.6	3.6	5.6	8.5	12.5	24.1
2003 4/.....	0.3	0.5	0.6	1.0	1.9	2.7	5.2	9.1	13.9	25.4
2002.....	0.3	0.5	0.7	1.2	2.2	3.2	5.0	8.4	14.1	25.8
2001.....	0.3	0.4	0.6	1.1	2.0	2.9	5.9	9.0	14.5	27.3
2000.....	0.2	0.5	0.8	1.0	2.0	3.3	6.0	9.5	14.8	28.3
1999.....	0.3	0.5	0.6	1.4	2.1	3.8	6.5	9.8	14.7	25.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lymphoid leukemia (C91)										
Black, both sexes										
2006 4/.....	*	*	1.1	2.3	3.9	6.1	9.6	10.9	14.0	19.5
2005 4/.....	*	0.9	1.1	1.4	3.4	7.3	6.3	11.9	12.9	16.0
2004 4/.....	*	*	1.2	2.0	4.2	5.2	10.1	12.0	14.4	19.7
2003 4/.....	*	*	1.1	2.0	2.9	5.6	9.5	11.2	12.2	19.8
2002.....	*	*	1.0	2.6	4.3	5.2	7.9	14.3	13.1	18.4
2001.....	*	0.9	1.7	2.4	4.0	7.4	7.9	13.0	13.5	19.0
2000.....	*	1.0	1.1	1.5	3.5	4.5	9.1	10.5	13.6	25.0
1999.....	*	*	1.2	2.8	4.0	7.0	8.0	10.0	16.2	22.0
Black, male										
2006 4/.....	*	*	*	3.6	6.2	9.1	14.8	19.7	21.2	31.3
2005 4/.....	*	*	*	*	4.5	10.7	9.2	20.1	20.4	19.5
2004 4/.....	*	*	*	2.6	6.7	9.0	17.2	17.7	20.1	30.8
2003 4/.....	*	*	2.0	3.2	*	8.4	11.8	15.1	17.0	26.1
2002.....	*	*	*	3.3	5.7	7.0	11.7	21.6	18.3	31.1
2001.....	*	*	2.2	3.1	6.5	11.2	10.8	16.7	22.5	28.4
2000.....	*	*	*	*	6.0	7.3	13.7	16.5	21.1	40.4
1999.....	*	*	*	3.9	5.5	10.9	12.2	15.2	21.8	32.8
Black, female										
2006 4/.....	*	*	*	*	*	3.8	6.1	5.6	10.1	14.6
2005 4/.....	*	*	*	*	*	4.8	4.3	6.9	8.9	14.6
2004 4/.....	*	*	*	*	*	*	5.3	8.6	11.3	15.4
2003 4/.....	*	*	*	*	*	*	8.0	9.0	9.7	17.4
2002.....	*	*	*	*	3.1	3.9	5.4	9.9	10.3	13.6
2001.....	*	*	*	*	*	4.7	6.1	10.8	9.0	15.6
2000.....	*	*	*	*	*	*	6.0	6.8	9.8	19.3
1999.....	*	*	*	*	*	4.0	5.2	6.9	13.4	18.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Myeloid leukemia (C92)										
All races 3/, both sexes										
2006.....	1.1	1.4	2.4	3.9	6.1	10.8	15.6	20.7	28.1	24.8
2005.....	1.0	1.6	2.3	3.6	6.4	10.0	15.9	21.7	25.4	25.8
2004.....	1.1	1.6	2.4	3.7	6.6	10.7	15.5	21.7	26.9	25.6
2003.....	1.2	1.5	2.4	3.7	6.9	10.1	16.3	22.4	25.5	26.1
2002.....	1.1	1.6	2.5	4.0	7.1	10.9	16.1	21.6	26.9	25.7
2001.....	1.2	1.9	2.7	4.2	6.9	10.7	16.8	22.2	25.5	25.1
2000.....	1.2	1.8	2.7	4.3	6.8	11.1	16.0	22.0	26.7	25.3
1999.....	1.2	2.0	2.5	4.6	6.5	10.8	15.2	19.5	24.5	25.6
All races 3/, male										
2006.....	1.2	1.4	2.8	4.6	7.5	14.2	20.6	28.6	40.6	34.5
2005.....	1.0	1.6	2.4	4.4	8.1	12.9	21.3	31.3	37.6	38.1
2004.....	1.2	1.6	2.8	4.4	7.9	14.5	20.7	29.9	36.8	36.1
2003.....	1.4	1.8	2.7	4.5	8.4	13.2	21.7	31.0	35.8	37.9
2002.....	1.2	1.8	2.8	5.0	8.6	14.1	22.1	30.5	38.6	37.6
2001.....	1.3	2.0	2.8	4.9	9.0	13.4	22.2	29.9	37.0	37.2
2000.....	1.2	2.0	3.1	5.0	8.2	14.1	21.9	29.7	39.1	39.1
1999.....	1.2	2.1	2.8	5.2	8.0	13.8	20.7	26.3	34.6	37.7
All races 3/, female										
2006.....	1.0	1.5	2.0	3.3	4.9	7.8	11.5	14.9	20.2	20.3
2005.....	1.1	1.6	2.2	2.9	4.8	7.6	11.4	14.6	17.8	20.2
2004.....	1.0	1.5	2.0	3.1	5.3	7.4	11.3	15.8	20.8	20.8
2003.....	1.1	1.3	2.1	2.9	5.5	7.4	12.0	16.3	19.3	20.9
2002.....	1.0	1.5	2.3	3.1	5.7	8.2	11.3	15.3	19.9	20.5
2001.....	1.2	1.8	2.6	3.6	5.1	8.4	12.4	16.9	18.6	20.1
2000.....	1.2	1.6	2.3	3.7	5.6	8.5	11.3	16.6	19.4	19.7
1999.....	1.2	1.9	2.2	4.1	5.2	8.3	10.9	14.7	18.6	20.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Myeloid leukemia (C92)										
White, both sexes										
2006 4/.....	1.1	1.4	2.5	4.0	6.3	11.3	16.3	21.3	29.5	25.6
2005 4/.....	1.0	1.6	2.2	3.7	6.7	10.6	16.6	22.7	26.8	26.9
2004 4/.....	1.1	1.6	2.4	3.8	6.8	11.1	16.1	22.6	27.7	26.7
2003 4/.....	1.2	1.6	2.4	3.8	7.1	10.6	17.0	23.5	26.4	26.9
2002.....	1.2	1.6	2.5	4.0	7.3	11.4	16.9	22.5	27.9	26.5
2001.....	1.2	1.9	2.7	4.3	7.3	11.0	17.4	23.0	26.5	26.0
2000.....	1.1	1.7	2.7	4.3	6.8	11.6	16.8	23.0	27.7	25.9
1999.....	1.2	2.0	2.5	4.8	6.6	11.1	15.7	20.2	25.2	26.7
White, male										
2006 4/.....	1.2	1.4	2.7	4.7	7.8	14.7	21.6	29.5	42.5	35.1
2005 4/.....	1.0	1.5	2.4	4.5	8.4	13.7	22.3	32.7	39.1	40.7
2004 4/.....	1.2	1.6	2.7	4.6	8.3	15.0	21.6	30.9	38.0	38.3
2003 4/.....	1.4	1.9	2.7	4.5	8.8	13.7	22.6	32.6	37.6	39.6
2002.....	1.2	1.7	2.8	5.0	8.8	14.5	23.1	31.8	39.9	38.7
2001.....	1.2	2.0	2.8	4.9	9.4	13.7	22.9	30.9	38.7	39.1
2000.....	1.2	2.0	3.1	5.0	8.2	14.8	22.6	31.4	40.5	41.0
1999.....	1.2	2.1	2.7	5.4	8.0	14.0	21.4	27.2	35.8	39.5
White, female										
2006 4/.....	1.1	1.4	2.2	3.4	5.0	8.2	11.9	15.2	21.2	21.2
2005 4/.....	1.1	1.7	2.1	3.0	5.1	7.9	11.9	15.4	19.0	20.6
2004 4/.....	1.0	1.5	2.1	3.1	5.4	7.7	11.7	16.5	21.4	21.5
2003 4/.....	1.1	1.3	2.1	3.1	5.5	7.8	12.3	16.9	19.6	21.3
2002.....	1.1	1.5	2.2	3.0	5.9	8.6	11.8	15.9	20.6	21.2
2001.....	1.2	1.8	2.6	3.7	5.3	8.7	13.0	17.4	19.1	20.5
2000.....	1.1	1.5	2.4	3.7	5.5	8.8	12.1	17.2	20.1	19.8
1999.....	1.2	1.9	2.2	4.2	5.4	8.5	11.1	15.3	18.9	21.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Myeloid leukemia (C92)										
Black, both sexes										
2006 4/.....	1.3	1.7	2.2	3.7	6.0	9.3	11.8	16.9	16.3	17.4
2005 4/.....	1.2	1.8	3.1	3.7	4.9	7.6	11.6	12.9	15.5	17.4
2004 4/.....	1.1	1.7	3.1	3.7	6.8	9.4	11.1	14.7	20.3	17.8
2003 4/.....	1.5	1.6	2.9	3.0	7.0	7.8	13.6	15.5	18.3	21.0
2002.....	0.8	1.9	2.4	4.8	6.6	8.4	12.5	15.1	18.1	19.3
2001.....	1.6	1.6	2.8	4.2	4.6	10.4	12.8	17.2	15.7	17.2
2000.....	1.7	2.5	3.2	4.5	7.9	8.6	11.2	13.3	16.1	20.3
1999.....	1.4	2.1	2.8	3.4	6.7	9.6	12.4	14.1	17.9	14.3
Black, male										
2006 4/.....	1.5	*	2.8	4.7	6.9	13.1	14.5	22.4	24.0	29.5
2005 4/.....	*	2.2	2.9	4.3	6.4	8.8	14.2	18.3	26.7	*
2004 4/.....	*	*	4.2	4.2	7.5	12.8	14.6	17.7	25.1	19.9
2003 4/.....	1.5	*	3.2	4.8	7.1	11.6	17.7	22.4	19.9	31.3
2002.....	*	2.5	2.5	5.8	8.1	11.0	17.7	19.7	22.9	32.2
2001.....	1.7	*	3.1	5.3	6.3	14.0	17.8	20.9	19.3	*
2000.....	1.8	2.5	3.8	5.1	9.3	9.6	17.1	15.6	22.8	24.2
1999.....	*	2.5	3.7	4.4	8.7	12.5	16.6	19.9	20.9	*
Black, female										
2006 4/.....	*	2.0	1.6	2.9	5.4	6.4	9.9	13.7	12.3	12.4
2005 4/.....	*	1.4	3.3	3.1	3.7	6.7	9.7	9.6	9.6	18.0
2004 4/.....	*	1.9	2.2	3.3	6.3	6.8	8.8	12.9	17.8	16.9
2003 4/.....	1.4	1.8	2.6	*	6.9	4.9	10.9	11.4	17.4	17.0
2002.....	*	*	2.4	3.9	5.4	6.5	9.0	12.3	15.5	14.4
2001.....	1.4	1.8	2.5	3.3	3.2	7.7	9.4	15.0	13.9	16.4
2000.....	1.7	2.6	2.6	3.9	6.9	7.9	7.3	12.0	12.7	18.9
1999.....	1.4	1.6	*	*	5.2	7.5	9.7	10.6	16.4	12.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified leukemia (C94-C95)										
All races 3/, both sexes										
2006.....	0.3	0.5	0.7	1.3	2.3	4.2	7.4	11.6	16.3	22.7
2005.....	0.3	0.4	0.8	1.3	2.3	4.5	7.2	11.5	17.6	23.0
2004.....	0.4	0.4	0.8	1.5	2.4	4.1	7.9	11.5	17.1	23.8
2003.....	0.4	0.5	0.7	1.4	2.4	4.2	7.5	11.8	17.9	24.0
2002.....	0.4	0.5	0.9	1.5	2.4	4.5	7.9	11.6	17.5	24.1
2001.....	0.3	0.5	0.8	1.4	2.8	4.4	7.7	12.2	17.0	24.4
2000.....	0.4	0.6	0.8	1.6	2.6	5.0	8.0	12.0	18.5	25.6
1999.....	0.4	0.6	0.8	1.7	3.0	4.8	8.3	13.0	17.8	25.2
All races 3/, male										
2006.....	0.3	0.6	0.9	1.6	3.1	5.6	9.1	17.1	23.0	29.7
2005.....	0.3	0.5	0.8	1.7	2.7	5.7	10.1	16.1	24.3	31.5
2004.....	0.4	0.5	0.9	1.8	3.1	5.2	11.0	16.5	24.4	31.7
2003.....	0.5	0.7	0.9	1.7	3.0	5.3	10.4	16.4	24.6	31.1
2002.....	0.4	0.6	1.0	1.7	3.0	5.9	10.9	16.1	24.3	32.6
2001.....	0.4	0.7	0.9	1.7	3.4	6.0	10.2	17.3	23.6	35.3
2000.....	0.4	0.8	0.7	2.0	3.1	6.8	11.1	16.1	25.1	35.7
1999.....	0.4	0.7	0.8	1.8	3.7	6.3	11.5	18.4	23.4	34.6
All races 3/, female										
2006.....	0.3	0.3	0.5	0.9	1.6	3.1	6.0	7.6	12.2	19.5
2005.....	0.2	0.4	0.8	1.0	1.9	3.4	4.9	8.2	13.5	19.1
2004.....	0.4	0.3	0.6	1.2	1.7	3.2	5.4	7.9	12.7	20.3
2003.....	0.3	0.4	0.5	1.2	1.9	3.2	5.2	8.6	13.9	20.8
2002.....	0.3	0.5	0.8	1.3	2.0	3.2	5.5	8.4	13.4	20.4
2001.....	0.2	0.4	0.7	1.2	2.2	3.0	5.6	8.6	13.1	19.8
2000.....	0.3	0.5	0.9	1.2	2.1	3.5	5.6	9.2	14.6	21.4
1999.....	0.3	0.5	0.8	1.5	2.4	3.4	5.9	9.2	14.6	21.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified leukemia (C94-C95)										
White, both sexes										
2006 4/.....	0.3	0.5	0.7	1.3	2.4	4.5	7.7	11.8	17.1	23.5
2005 4/.....	0.2	0.4	0.8	1.3	2.3	4.7	7.4	11.8	18.2	23.7
2004 4/.....	0.4	0.4	0.7	1.5	2.4	4.1	7.7	11.9	17.7	24.1
2003 4/.....	0.4	0.6	0.7	1.4	2.5	4.2	7.7	12.2	18.2	24.8
2002.....	0.4	0.5	0.8	1.5	2.4	4.7	8.1	11.8	18.0	24.2
2001.....	0.3	0.6	0.8	1.5	2.7	4.4	7.8	12.5	17.6	25.1
2000.....	0.4	0.6	0.8	1.6	2.7	5.1	8.2	12.2	19.2	26.3
1999.....	0.4	0.6	0.7	1.7	3.1	4.8	8.6	13.4	18.0	25.7
White, male										
2006 4/.....	0.3	0.6	1.0	1.7	3.2	5.9	9.4	17.2	24.2	30.9
2005 4/.....	0.3	0.5	0.8	1.7	2.7	6.0	10.1	16.4	25.3	32.3
2004 4/.....	0.4	0.5	0.9	1.8	3.2	5.2	10.8	17.2	25.4	32.1
2003 4/.....	0.5	0.7	0.9	1.6	3.1	5.4	10.6	16.9	25.2	32.7
2002.....	0.4	0.5	0.9	1.8	3.0	6.0	10.9	16.6	24.9	34.0
2001.....	0.3	0.7	0.9	1.7	3.3	6.0	10.6	17.8	24.1	36.4
2000.....	0.4	0.7	0.7	1.9	3.2	7.0	11.3	16.1	26.0	36.4
1999.....	0.5	0.7	0.8	1.8	3.9	6.5	11.9	19.0	23.6	35.4
White, female										
2006 4/.....	0.3	0.3	0.5	0.9	1.6	3.3	6.3	7.7	12.6	20.0
2005 4/.....	0.2	0.3	0.8	1.0	1.9	3.5	5.1	8.4	13.8	19.8
2004 4/.....	0.3	0.3	0.6	1.2	1.7	3.2	5.2	8.0	13.0	20.5
2003 4/.....	0.3	0.5	0.5	1.2	1.9	3.1	5.4	8.8	13.9	21.4
2002.....	0.3	0.4	0.8	1.3	1.9	3.5	5.7	8.4	13.7	20.0
2001.....	0.2	0.4	0.6	1.2	2.1	3.0	5.6	8.8	13.7	20.3
2000.....	0.3	0.5	0.9	1.2	2.2	3.5	5.7	9.4	15.1	22.2
1999.....	0.3	0.5	0.7	1.5	2.4	3.2	6.0	9.5	14.7	21.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified leukemia (C94-C95)										
Black, both sexes										
2006 4/.....	*	*	0.8	1.2	2.8	2.5	7.0	11.4	11.4	18.2
2005 4/.....	*	*	*	1.6	2.5	3.7	6.4	10.4	14.3	18.7
2004 4/.....	0.7	*	1.0	2.0	2.9	4.7	11.3	10.2	12.4	22.3
2003 4/.....	*	*	1.2	2.1	2.9	4.5	6.2	10.4	17.5	18.4
2002.....	*	1.0	1.6	1.7	3.0	3.7	8.0	9.4	15.4	26.1
2001.....	*	*	1.0	*	3.6	4.5	7.3	10.9	12.5	20.6
2000.....	*	0.9	*	2.0	2.6	4.6	8.0	10.8	13.8	20.0
1999.....	*	*	1.5	1.8	2.9	5.4	7.2	10.5	16.5	19.4
Black, male										
2006 4/.....	*	*	*	*	3.5	*	8.9	16.6	16.4	23.2
2005 4/.....	*	*	*	*	*	5.0	10.8	14.3	19.0	27.0
2004 4/.....	*	*	*	*	*	6.5	16.8	12.3	*	27.8
2003 4/.....	*	*	*	*	*	*	9.5	12.8	21.4	*
2002.....	*	*	*	*	*	5.5	13.0	10.6	22.9	22.5
2001.....	*	*	*	*	4.3	7.1	7.7	13.9	18.5	29.5
2000.....	*	*	*	*	*	5.5	12.7	17.5	19.4	32.3
1999.....	*	*	*	*	*	5.7	8.8	15.2	18.3	24.6
Black, female										
2006 4/.....	*	*	*	*	*	*	5.7	8.3	8.7	16.1
2005 4/.....	*	*	*	*	*	*	*	8.0	11.9	15.4
2004 4/.....	*	*	*	*	*	*	7.5	8.9	11.7	20.1
2003 4/.....	*	*	*	*	*	4.4	*	9.0	15.5	17.8
2002.....	*	*	*	*	*	*	4.7	8.8	11.5	27.5
2001.....	*	*	*	*	*	*	7.0	9.2	9.4	17.3
2000.....	*	*	*	*	*	4.0	4.9	6.8	11.0	15.5
1999.....	*	*	*	*	*	5.2	6.1	7.7	15.6	17.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Benign neoplasms (D10-D36)										
All races 3/, both sexes										
2006.....	0.2	0.3	0.4	0.4	0.6	0.9	1.4	2.6	4.0	7.1
2005.....	0.2	0.2	0.3	0.5	0.6	0.9	1.5	2.7	3.9	6.4
2004.....	0.1	0.3	0.3	0.5	0.8	0.9	1.7	3.0	4.5	7.9
2003.....	0.2	0.3	0.4	0.5	0.8	1.1	1.7	2.8	4.9	7.4
2002.....	0.2	0.4	0.4	0.5	0.8	1.4	1.8	3.1	4.3	7.8
2001.....	0.2	0.2	0.4	0.6	0.9	1.3	1.7	2.9	4.6	8.7
2000.....	0.2	0.2	0.4	0.4	0.7	1.2	1.9	3.2	5.5	9.0
1999.....	0.1	0.2	0.3	0.6	0.9	1.3	1.8	3.4	4.8	8.4
All races 3/, male										
2006.....	*	0.2	0.4	0.4	0.4	1.1	1.5	2.5	4.1	6.1
2005.....	*	0.2	0.3	0.3	0.5	0.9	1.3	2.5	3.4	6.1
2004.....	*	0.2	0.3	0.5	0.6	0.8	1.5	2.9	4.2	7.4
2003.....	0.2	0.2	0.4	0.5	0.8	1.1	1.8	2.7	4.9	7.3
2002.....	*	0.3	0.3	0.5	0.8	1.4	1.9	3.1	3.8	6.3
2001.....	*	*	0.4	0.6	1.0	1.3	1.6	3.1	5.1	8.9
2000.....	*	0.3	0.4	0.4	0.5	1.3	1.7	3.2	4.9	8.7
1999.....	0.2	*	0.3	0.6	0.9	1.1	2.1	3.1	4.6	9.2
All races 3/, female										
2006.....	0.2	0.3	0.4	0.4	0.8	0.8	1.4	2.7	3.9	7.6
2005.....	0.3	0.2	0.4	0.6	0.8	0.9	1.7	2.8	4.2	6.6
2004.....	0.2	0.4	0.4	0.6	0.9	1.0	1.8	3.0	4.7	8.1
2003.....	0.2	0.4	0.3	0.5	0.7	1.1	1.5	2.8	4.9	7.4
2002.....	0.3	0.4	0.4	0.5	0.7	1.5	1.8	3.2	4.6	8.4
2001.....	0.2	0.3	0.4	0.6	0.8	1.2	1.7	2.7	4.3	8.6
2000.....	0.3	0.2	0.4	0.5	0.9	1.1	2.0	3.3	5.9	9.1
1999.....	*	0.2	0.3	0.6	0.9	1.5	1.6	3.7	4.9	8.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Benign neoplasms (D10-D36)										
White, both sexes										
2006 4/.....	0.2	0.2	0.3	0.4	0.6	0.9	1.4	2.5	3.9	7.1
2005 4/.....	0.2	0.2	0.3	0.4	0.5	0.9	1.4	2.5	3.9	6.2
2004 4/.....	0.1	0.3	0.3	0.5	0.8	0.8	1.5	2.8	4.4	7.8
2003 4/.....	0.2	0.3	0.4	0.5	0.8	1.0	1.5	2.8	4.8	7.4
2002.....	0.2	0.3	0.4	0.5	0.7	1.3	1.8	3.1	4.1	7.4
2001.....	0.2	0.2	0.4	0.5	0.7	1.1	1.6	2.8	4.3	8.2
2000.....	0.2	0.2	0.4	0.4	0.7	1.1	1.7	3.2	5.5	8.9
1999.....	0.1	0.2	0.3	0.5	0.8	1.2	1.7	3.2	4.7	8.4
White, male										
2006 4/.....	*	0.2	0.4	0.4	0.4	1.1	1.4	2.4	3.8	5.7
2005 4/.....	*	*	0.3	0.3	0.4	0.8	1.3	2.4	3.4	6.2
2004 4/.....	*	*	0.3	0.4	0.6	0.8	1.5	2.7	3.9	7.3
2003 4/.....	*	0.2	0.3	0.4	0.8	1.0	1.7	2.7	4.7	7.3
2002.....	*	0.3	0.4	0.5	0.7	1.3	1.9	3.0	3.6	5.7
2001.....	*	*	0.4	0.5	0.8	1.2	1.5	3.2	4.7	8.4
2000.....	*	0.3	0.4	0.4	*	1.1	1.5	3.2	5.0	8.7
1999.....	*	*	*	0.5	0.8	1.1	2.0	2.8	4.4	9.3
White, female										
2006 4/.....	*	*	0.3	0.3	0.7	0.8	1.3	2.5	3.9	7.8
2005 4/.....	*	*	0.4	0.5	0.6	0.9	1.6	2.5	4.2	6.2
2004 4/.....	*	0.3	0.3	0.5	0.9	0.8	1.5	2.9	4.7	8.0
2003 4/.....	*	0.3	0.4	0.5	0.7	1.0	1.4	3.0	4.9	7.4
2002.....	0.2	0.4	0.4	0.5	0.7	1.4	1.7	3.1	4.3	8.1
2001.....	*	*	0.4	0.5	0.7	1.0	1.8	2.5	4.1	8.0
2000.....	0.2	*	0.4	0.5	1.0	1.0	1.9	3.1	5.8	9.0
1999.....	*	*	0.3	0.5	0.8	1.3	1.5	3.6	4.8	8.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of eye, brain and other parts of central nervous system (D31-D33)

All races 3/, both sexes

2006.....	0.1	0.1	0.2	0.2	0.3	0.6	1.0	1.8	2.9	4.9
2005.....	*	0.1	0.2	0.2	0.3	0.5	1.1	1.8	2.7	4.7
2004.....	*	0.1	0.2	0.3	0.5	0.6	1.2	1.7	3.5	5.6
2003.....	0.1	0.1	0.2	0.3	0.5	0.8	1.1	1.8	3.4	5.4
2002.....	0.1	0.2	0.2	0.3	0.4	0.9	1.2	2.1	2.9	5.1
2001.....	*	0.1	0.2	0.3	0.5	0.8	1.2	1.8	3.0	5.9
2000.....	*	0.1	0.2	0.2	0.5	0.7	1.2	2.3	3.7	6.2
1999.....	*	*	0.2	0.4	0.6	0.9	1.1	2.2	3.5	5.5

All races 3/, male

2006.....	*	*	*	0.2	*	0.7	1.0	1.5	2.9	3.9
2005.....	*	*	*	*	*	0.5	1.0	1.6	2.0	3.9
2004.....	*	*	*	*	*	*	1.0	1.5	3.2	5.2
2003.....	*	*	*	0.3	0.4	0.7	1.1	1.6	3.4	5.2
2002.....	*	*	*	*	0.4	0.8	1.2	1.7	2.4	3.2
2001.....	*	*	*	0.3	0.4	0.8	1.0	1.8	2.7	5.4
2000.....	*	*	*	*	*	0.8	1.1	2.0	3.3	5.9
1999.....	*	*	*	0.3	0.4	0.8	1.2	1.9	3.2	5.5

All races 3/, female

2006.....	*	*	0.2	0.2	0.5	0.5	1.0	2.0	3.0	5.3
2005.....	*	*	0.2	0.4	0.4	0.6	1.2	1.9	3.2	5.0
2004.....	*	*	0.2	0.4	0.6	0.8	1.3	1.9	3.6	5.8
2003.....	*	*	*	0.3	0.5	0.8	1.1	2.0	3.3	5.5
2002.....	*	*	0.3	0.3	0.4	1.0	1.2	2.3	3.2	5.9
2001.....	*	*	0.2	0.3	0.5	0.8	1.3	1.9	3.1	6.2
2000.....	*	*	*	0.3	0.6	0.7	1.3	2.4	4.0	6.4
1999.....	*	*	*	0.4	0.7	1.1	1.1	2.5	3.7	5.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Of eye, brain and other parts of central nervous system (D31-D33)										
White, both sexes										
2006 4/.....	*	0.1	0.2	0.2	0.3	0.7	0.9	1.6	3.0	4.9
2005 4/.....	*	*	0.2	0.2	0.2	0.5	1.0	1.6	2.8	4.5
2004 4/.....	*	0.1	0.2	0.2	0.4	0.4	1.0	1.6	3.3	5.7
2003 4/.....	*	0.1	0.2	0.2	0.5	0.6	1.0	1.9	3.4	5.4
2002.....	*	0.2	0.2	0.3	0.4	0.8	1.2	2.0	2.8	4.9
2001.....	*	*	0.2	0.2	0.4	0.7	1.1	1.8	2.8	5.5
2000.....	*	*	0.2	0.2	0.5	0.7	1.1	2.2	3.7	6.3
1999.....	*	*	0.1	0.3	0.5	0.9	1.1	2.1	3.4	5.6
White, male										
2006 4/.....	*	*	*	*	*	0.8	1.0	1.4	3.0	3.7
2005 4/.....	*	*	*	*	*	0.5	1.0	1.5	2.1	3.8
2004 4/.....	*	*	*	*	*	*	1.0	1.4	2.9	5.2
2003 4/.....	*	*	*	*	0.4	0.6	1.0	1.6	3.4	5.3
2002.....	*	*	*	*	*	0.8	1.2	1.6	2.3	2.9
2001.....	*	*	*	*	*	0.8	1.0	1.9	2.5	4.9
2000.....	*	*	*	*	*	0.7	0.9	2.1	3.2	6.0
1999.....	*	*	*	*	*	0.8	1.2	1.7	3.0	5.5
White, female										
2006 4/.....	*	*	*	*	0.5	0.6	0.9	1.9	3.1	5.4
2005 4/.....	*	*	*	0.3	*	0.6	1.1	1.7	3.2	4.8
2004 4/.....	*	*	*	0.3	0.6	0.6	1.1	1.7	3.6	5.9
2003 4/.....	*	*	*	*	0.5	0.6	1.0	2.1	3.3	5.4
2002.....	*	*	*	0.3	0.5	0.9	1.1	2.3	3.1	5.8
2001.....	*	*	*	*	0.4	0.6	1.3	1.7	3.0	5.8
2000.....	*	*	*	*	0.7	0.7	1.3	2.3	4.0	6.4
1999.....	*	*	*	0.4	0.6	0.9	1.0	2.4	3.6	5.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites (D10-D24,D29-D30,D34-D36)

All races 3/, both sexes

2006.....	*	0.1	0.1	0.1	0.3	0.3	0.4	0.8	1.0	2.1
2005.....	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.9	1.1	1.7
2004.....	*	0.1	0.1	0.2	0.3	0.3	0.5	1.2	1.0	2.2
2003.....	0.1	0.1	0.2	0.2	0.3	0.4	0.5	0.9	1.4	1.9
2002.....	0.1	0.2	0.1	0.2	0.3	0.5	0.6	1.0	1.3	2.6
2001.....	0.1	0.1	0.2	0.3	0.4	0.5	0.5	1.0	1.6	2.5
2000.....	*	0.1	0.2	0.2	0.2	0.4	0.6	1.0	1.7	2.6
1999.....	*	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.3	2.8

All races 3/, male

2006.....	*	*	*	*	*	*	*	1.0	1.1	2.2
2005.....	*	*	*	0.2	*	*	*	1.0	1.3	2.2
2004.....	*	*	*	0.3	0.4	0.5	*	1.5	1.0	2.3
2003.....	*	*	*	*	0.4	*	0.7	1.2	1.5	2.1
2002.....	*	*	*	0.3	0.4	0.6	0.7	1.4	1.4	3.1
2001.....	*	*	0.2	0.3	0.5	0.5	0.5	1.3	2.4	3.5
2000.....	*	*	0.2	*	*	0.5	0.6	1.1	1.6	2.9
1999.....	*	*	*	*	0.5	*	0.9	1.2	1.4	3.7

All races 3/, female

2006.....	*	*	*	*	0.3	*	*	0.6	0.9	2.1
2005.....	*	*	*	*	0.4	*	0.5	0.8	1.0	1.5
2004.....	*	*	*	*	*	*	0.5	1.1	1.0	2.2
2003.....	*	*	*	*	*	*	*	0.7	1.4	1.8
2002.....	*	0.2	*	*	*	0.4	0.5	0.8	1.3	2.4
2001.....	*	*	*	*	*	0.4	*	0.8	1.1	2.1
2000.....	*	*	*	*	*	0.4	0.6	0.8	1.8	2.6
1999.....	*	*	*	*	*	0.4	0.5	1.2	1.2	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Of other and unspecified sites (D10-D24,D29-D30,D34-D36)

White, both sexes

2006 4/.....	*	*	0.1	0.1	0.2	0.3	0.4	0.8	0.8	2.1
2005 4/.....	*	*	0.1	0.2	0.3	0.3	0.4	0.8	1.1	1.7
2004 4/.....	*	*	*	0.2	0.4	0.4	0.4	1.1	1.1	2.0
2003 4/.....	*	0.1	0.2	0.2	0.3	0.4	0.5	0.9	1.4	1.9
2002.....	*	0.1	0.1	0.2	0.3	0.4	0.5	1.1	1.3	2.3
2001.....	*	*	0.2	0.3	0.4	0.4	0.5	1.0	1.4	2.5
2000.....	*	*	0.2	0.2	0.2	0.4	0.5	0.9	1.8	2.5
1999.....	*	*	*	0.2	0.3	0.3	0.6	1.1	1.3	2.8

White, male

2006 4/.....	*	*	*	*	*	*	*	1.1	*	1.9
2005 4/.....	*	*	*	*	*	*	*	0.9	1.3	2.4
2004 4/.....	*	*	*	*	0.4	0.5	*	1.3	1.0	2.1
2003 4/.....	*	*	*	*	*	*	0.7	1.1	1.3	2.0
2002.....	*	*	*	*	*	0.5	0.7	1.4	1.4	2.8
2001.....	*	*	*	*	0.5	*	*	1.3	2.2	3.5
2000.....	*	*	*	*	*	*	*	1.1	1.7	2.7
1999.....	*	*	*	*	*	*	0.8	1.1	1.4	3.8

White, female

2006 4/.....	*	*	*	*	*	*	*	0.5	0.8	2.3
2005 4/.....	*	*	*	*	0.4	*	*	0.7	1.0	1.3
2004 4/.....	*	*	*	*	*	*	*	1.0	1.1	2.0
2003 4/.....	*	*	*	*	*	*	*	0.8	1.5	1.8
2002.....	*	*	*	*	*	*	*	0.9	1.2	2.1
2001.....	*	*	*	*	*	*	*	0.8	1.0	2.1
2000.....	*	*	*	*	*	*	0.5	0.8	1.8	2.5
1999.....	*	*	*	*	*	*	0.5	1.1	1.2	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

All races 3/, both sexes

2006.....	0.6	1.0	1.7	2.4	4.5	7.5	14.4	24.7	40.2	66.9
2005.....	0.7	1.0	1.6	2.6	4.6	8.1	13.8	24.0	39.1	66.4
2004.....	0.6	1.0	1.5	2.7	4.7	7.9	14.3	23.1	40.9	65.6
2003.....	0.6	1.0	1.5	2.5	5.0	8.3	14.1	24.8	40.9	69.6
2002.....	0.7	1.0	1.5	3.0	4.4	7.9	14.0	24.7	40.0	67.0
2001.....	0.7	1.1	1.7	3.0	5.1	8.7	15.6	25.6	41.5	70.1
2000.....	0.7	1.1	1.8	3.2	5.5	9.0	14.9	24.6	43.2	72.0
1999.....	0.7	1.1	1.8	2.8	5.1	9.1	16.2	24.9	41.1	70.8

All races 3/, male

2006.....	0.7	1.2	2.0	3.1	5.3	9.4	19.0	34.2	54.6	83.5
2005.....	0.8	1.2	2.0	3.0	6.1	10.7	18.0	32.3	53.9	84.0
2004.....	0.8	1.1	1.8	3.2	6.1	10.5	18.1	30.8	55.3	87.9
2003.....	0.7	1.1	1.8	3.0	6.0	10.1	18.8	32.2	55.2	91.6
2002.....	0.8	1.2	1.7	3.9	5.3	10.5	17.5	33.2	53.7	91.9
2001.....	0.8	1.4	1.9	3.2	6.3	10.9	20.3	35.1	55.7	96.6
2000.....	0.8	1.2	2.1	3.6	6.7	11.6	18.9	32.6	58.8	97.6
1999.....	0.9	1.2	2.2	3.2	6.7	11.8	21.2	33.2	55.0	97.8

All races 3/, female

2006.....	0.5	0.9	1.3	1.8	3.7	5.8	10.7	17.8	31.2	59.1
2005.....	0.6	0.9	1.2	2.2	3.2	5.9	10.4	18.0	29.8	58.3
2004.....	0.5	0.9	1.1	2.2	3.4	5.6	11.2	17.6	32.0	55.5
2003.....	0.4	0.9	1.2	2.1	4.2	6.7	10.3	19.5	32.2	59.9
2002.....	0.6	0.9	1.3	2.3	3.5	5.6	11.1	18.7	31.7	56.2
2001.....	0.6	0.9	1.5	2.9	4.1	6.9	11.8	19.0	33.0	59.0
2000.....	0.6	1.0	1.4	2.8	4.4	6.8	11.7	19.0	33.9	61.5
1999.....	0.6	1.0	1.4	2.3	3.6	6.8	12.2	19.2	33.0	60.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

White, both sexes

2006 4/.....	0.6	1.0	1.7	2.5	4.5	7.7	15.0	25.6	41.8	68.8
2005 4/.....	0.7	1.0	1.6	2.6	4.6	8.3	14.2	25.0	40.6	69.3
2004 4/.....	0.6	1.0	1.4	2.6	4.6	8.1	14.8	23.8	42.4	68.2
2003 4/.....	0.6	1.0	1.5	2.5	5.1	8.5	14.6	25.8	42.2	72.1
2002.....	0.6	1.1	1.5	3.0	4.5	8.2	14.4	25.6	41.1	69.3
2001.....	0.7	1.2	1.7	3.2	5.3	9.2	15.9	26.5	42.8	71.6
2000.....	0.7	1.1	1.7	3.3	5.5	9.3	15.0	25.3	44.9	74.3
1999.....	0.7	1.1	1.8	2.8	5.1	9.2	16.7	25.6	42.5	73.4

White, male

2006 4/.....	0.6	1.1	2.0	3.1	5.3	9.6	19.9	35.5	56.8	85.3
2005 4/.....	0.7	1.1	2.0	3.1	6.1	11.0	18.7	33.6	56.3	88.0
2004 4/.....	0.8	1.0	1.7	3.1	6.0	10.9	18.6	31.6	57.6	92.1
2003 4/.....	0.8	1.1	1.8	2.9	6.0	10.5	19.5	33.7	56.9	95.6
2002.....	0.8	1.3	1.6	3.7	5.5	10.8	18.2	34.4	54.9	95.4
2001.....	0.8	1.4	2.0	3.4	6.4	11.4	21.0	36.1	57.3	99.2
2000.....	0.7	1.2	2.1	3.7	6.6	11.9	18.9	33.4	61.3	101.1
1999.....	0.9	1.2	2.2	3.3	6.9	11.8	21.9	34.2	57.1	102.7

White, female

2006 4/.....	0.5	0.9	1.3	1.9	3.6	6.0	10.9	18.2	32.4	61.1
2005 4/.....	0.6	0.8	1.2	2.2	3.2	6.0	10.5	18.7	30.8	60.7
2004 4/.....	0.4	0.9	1.1	2.2	3.3	5.7	11.6	18.0	32.9	57.4
2003 4/.....	0.5	0.9	1.2	2.2	4.3	6.7	10.7	20.0	33.1	61.7
2002.....	0.5	0.9	1.3	2.2	3.6	5.9	11.4	19.3	32.6	57.9
2001.....	0.6	0.9	1.5	2.9	4.3	7.2	11.8	19.7	34.1	60.1
2000.....	0.6	1.0	1.4	2.8	4.5	7.0	11.8	19.6	35.0	63.4
1999.....	0.6	1.0	1.4	2.3	3.5	6.9	12.5	19.6	33.9	61.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Neoplasms of uncertain or unknown behavior of specified sites (D37-D47)

Black, both sexes

2006 4/.....	0.8	1.3	2.0	2.6	5.7	7.2	11.8	20.4	30.3	50.1
2005 4/.....	0.9	1.5	1.8	2.8	4.7	7.8	12.3	17.5	27.4	42.8
2004 4/.....	0.8	1.0	2.1	3.8	5.9	7.4	12.0	19.6	30.7	42.3
2003 4/.....	*	1.5	2.1	3.2	5.2	7.8	11.9	19.1	32.5	46.7
2002.....	1.1	1.1	2.0	4.2	4.3	7.3	11.9	19.2	33.2	49.9
2001.....	1.0	0.9	1.9	2.8	5.1	6.9	14.8	19.8	33.3	55.6
2000.....	*	1.4	2.5	3.0	5.7	8.0	15.6	20.1	28.5	51.6
1999.....	1.0	1.4	2.1	2.9	5.6	10.5	14.2	21.6	30.3	48.8

Black, male

2006 4/.....	*	1.7	2.2	3.8	6.9	10.1	14.5	26.8	41.8	64.3
2005 4/.....	*	1.6	2.2	3.6	6.2	8.8	14.5	21.9	38.6	47.4
2004 4/.....	*	*	2.6	5.0	8.0	9.7	16.5	25.9	35.9	48.7
2003 4/.....	*	*	2.1	4.3	6.6	8.9	17.1	21.5	45.0	56.3
2002.....	*	*	2.4	6.3	4.2	9.5	13.4	24.8	45.1	66.5
2001.....	*	*	*	*	6.3	9.1	17.8	27.4	46.6	71.5
2000.....	*	*	2.9	3.5	7.9	11.4	19.4	27.9	32.9	64.6
1999.....	*	*	*	3.6	5.9	14.5	17.6	26.1	34.9	52.7

Black, female

2006 4/.....	*	*	1.9	*	4.8	5.0	9.9	16.6	24.2	44.2
2005 4/.....	*	1.4	*	2.1	3.5	7.1	10.8	14.9	21.5	40.9
2004 4/.....	*	*	*	2.7	4.2	5.8	9.0	15.9	28.0	39.8
2003 4/.....	*	1.4	2.0	*	4.2	6.9	8.4	17.6	25.9	42.9
2002.....	*	*	*	2.6	4.3	5.7	11.0	15.9	27.1	43.5
2001.....	*	*	2.1	3.0	4.2	5.3	12.8	15.3	26.6	49.7
2000.....	*	*	2.2	2.7	3.9	5.4	13.1	15.4	26.3	46.8
1999.....	*	*	*	*	5.3	7.5	11.9	18.9	28.1	47.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Neoplasms of uncertain or unknown behavior of unspecified sites (D48)

All races 3/, both sexes

2006.....	*	0.2	0.2	0.3	0.5	0.6	0.9	1.4	2.1	4.0
2005.....	0.1	0.1	0.2	0.3	0.4	0.5	1.0	1.2	1.7	3.8
2004.....	*	0.2	0.2	0.3	0.3	0.4	0.8	1.1	1.7	4.1
2003.....	*	*	0.1	0.2	0.3	0.4	0.5	0.9	1.5	3.4
2002.....	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.9	1.9	3.8
2001.....	*	0.1	0.2	0.2	0.3	0.5	0.6	0.9	1.6	3.8
2000.....	*	0.1	*	0.2	0.4	0.3	0.6	1.1	1.9	3.9
1999.....	*	*	0.2	*	0.3	0.3	0.6	0.9	1.5	3.5

All races 3/, male

2006.....	*	*	*	0.3	0.5	0.6	0.9	1.4	2.2	2.6
2005.....	*	*	*	0.3	0.6	0.5	0.9	1.5	2.1	2.6
2004.....	*	0.2	0.2	0.3	0.4	0.4	0.9	1.0	1.7	3.6
2003.....	*	*	*	*	*	*	0.5	1.0	1.5	2.6
2002.....	*	*	*	*	*	*	0.6	0.9	1.5	3.0
2001.....	*	*	*	*	*	0.5	0.6	0.9	1.4	1.8
2000.....	*	*	*	*	*	*	*	1.1	1.6	2.4
1999.....	*	*	*	*	*	0.5	0.6	0.9	1.2	2.1

All races 3/, female

2006.....	*	*	0.2	0.3	0.5	0.6	0.8	1.3	2.1	4.6
2005.....	*	*	0.2	0.3	0.3	0.5	1.0	1.0	1.5	4.4
2004.....	*	*	*	0.2	0.3	0.4	0.7	1.2	1.7	4.4
2003.....	*	*	*	*	*	0.4	0.5	0.8	1.5	3.8
2002.....	*	*	*	*	0.4	*	0.4	0.9	2.2	4.2
2001.....	*	*	*	0.3	*	0.4	0.7	0.9	1.7	4.6
2000.....	*	*	*	*	0.4	*	0.6	1.1	2.0	4.4
1999.....	*	*	*	*	*	*	0.6	0.9	1.7	4.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Neoplasms of uncertain or unknown behavior of unspecified sites (D48)

White, both sexes

2006 4/.....	*	0.2	0.2	0.3	0.4	0.6	0.8	1.4	2.0	4.0
2005 4/.....	0.1	0.1	0.2	0.3	0.4	0.5	0.9	1.2	1.7	4.0
2004 4/.....	*	0.2	0.2	0.3	0.3	0.4	0.8	1.2	1.7	4.2
2003 4/.....	*	*	0.1	0.2	0.3	0.4	0.6	0.9	1.6	3.5
2002.....	*	0.1	*	0.2	0.3	0.4	0.5	1.0	2.0	3.9
2001.....	*	0.1	0.2	0.2	0.3	0.4	0.5	0.9	1.6	4.0
2000.....	*	0.1	*	0.2	0.4	0.3	0.5	1.0	1.9	4.1
1999.....	*	*	0.1	*	0.3	0.3	0.6	0.8	1.5	3.4

White, male

2006 4/.....	*	*	*	0.3	0.5	0.6	0.8	1.4	1.8	2.6
2005 4/.....	*	*	*	0.3	0.5	0.5	1.0	1.5	2.1	2.8
2004 4/.....	*	*	*	0.4	*	*	0.9	1.1	1.6	3.7
2003 4/.....	*	*	*	*	*	*	0.6	1.0	1.6	2.5
2002.....	*	*	*	*	*	*	0.6	1.0	1.5	3.1
2001.....	*	*	*	*	*	0.5	*	0.9	1.5	2.0
2000.....	*	*	*	*	*	*	*	1.0	1.7	2.6
1999.....	*	*	*	*	*	*	0.6	0.8	1.3	2.2

White, female

2006 4/.....	*	*	*	0.3	0.3	0.6	0.8	1.3	2.1	4.7
2005 4/.....	*	*	0.2	0.3	*	0.5	0.9	1.0	1.5	4.6
2004 4/.....	*	*	*	*	*	*	0.7	1.2	1.8	4.4
2003 4/.....	*	*	*	*	*	*	0.6	0.9	1.6	3.9
2002.....	*	*	*	*	0.4	*	*	0.9	2.3	4.3
2001.....	*	*	*	*	*	*	0.6	1.0	1.7	4.8
2000.....	*	*	*	*	*	*	0.6	1.0	2.0	4.7
1999.....	*	*	*	*	*	*	0.6	0.9	1.7	3.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

All races 3/, both sexes

2006.....	1.5	1.8	2.4	3.0	4.1	5.4	8.1	12.3	20.7	42.4
2005.....	1.6	2.0	2.6	3.1	4.0	6.0	8.9	13.8	22.6	51.2
2004.....	1.6	2.0	2.4	3.0	4.2	5.7	8.6	13.2	22.8	52.9
2003.....	1.6	2.0	2.6	3.0	4.1	5.8	9.1	13.6	23.0	56.7
2002.....	1.7	2.2	2.6	3.0	4.3	6.0	9.4	13.9	24.7	56.8
2001.....	1.8	2.2	2.4	3.0	4.3	6.1	8.7	15.0	24.5	58.5
2000.....	1.6	2.1	2.5	3.1	4.3	6.4	8.8	14.1	25.9	58.9
1999.....	1.6	2.1	2.1	2.9	3.9	5.4	9.3	14.5	23.7	60.3

All races 3/, male

2006.....	1.6	1.7	2.3	3.1	4.0	5.2	7.9	13.4	22.1	43.8
2005.....	1.5	2.0	2.6	3.3	3.9	6.0	8.9	14.2	26.6	49.6
2004.....	1.7	2.1	2.4	3.3	4.4	6.0	8.9	13.6	24.6	52.1
2003.....	1.8	2.2	2.5	2.8	4.2	6.1	10.1	14.8	24.9	55.1
2002.....	1.6	2.1	2.5	3.1	4.5	5.5	10.2	14.8	25.1	57.2
2001.....	1.6	2.1	2.4	2.7	4.1	5.9	8.6	15.3	27.3	58.1
2000.....	1.5	2.2	2.4	2.9	4.2	6.5	9.8	14.2	27.3	60.4
1999.....	1.6	2.2	2.1	2.7	3.6	5.4	9.9	15.7	25.4	63.2

All races 3/, female

2006.....	1.5	2.0	2.6	3.0	4.1	5.6	8.4	11.5	19.8	41.7
2005.....	1.6	2.0	2.6	2.9	4.2	6.0	8.9	13.4	20.1	51.9
2004.....	1.6	1.9	2.4	2.7	4.1	5.5	8.3	13.0	21.6	53.3
2003.....	1.5	1.9	2.6	3.2	4.0	5.6	8.4	12.8	21.9	57.5
2002.....	1.8	2.3	2.6	3.0	4.1	6.4	8.8	13.3	24.4	56.6
2001.....	2.0	2.2	2.5	3.4	4.4	6.2	8.8	14.8	22.8	58.7
2000.....	1.6	1.9	2.6	3.3	4.4	6.3	8.1	14.1	25.0	58.3
1999.....	1.7	2.0	2.2	3.0	4.1	5.4	8.8	13.6	22.8	59.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

White, both sexes

2006 4/.....	0.7	1.0	1.5	2.2	3.3	4.7	7.6	11.9	20.5	42.2
2005 4/.....	0.8	1.1	1.7	2.3	3.4	4.9	8.1	13.1	21.9	51.3
2004 4/.....	0.9	1.2	1.5	2.1	3.4	5.1	8.0	13.0	22.7	52.6
2003 4/.....	0.8	1.2	1.5	2.0	3.3	5.0	8.6	13.0	22.7	56.4
2002.....	0.8	1.3	1.6	2.3	3.5	5.3	8.8	13.1	24.3	56.2
2001.....	0.9	1.1	1.5	2.3	3.6	5.5	8.1	14.5	23.8	58.3
2000.....	0.8	1.1	1.6	2.4	3.5	5.7	8.3	13.7	25.5	58.8
1999.....	0.8	1.2	1.4	2.0	3.2	4.7	8.7	14.0	23.5	59.6

White, male

2006 4/.....	0.7	1.0	1.5	2.3	3.2	4.9	7.5	12.9	22.3	44.4
2005 4/.....	0.8	1.2	1.8	2.5	3.4	4.9	8.3	13.6	25.4	50.3
2004 4/.....	1.0	1.4	1.7	2.3	3.8	5.4	8.5	13.3	24.4	52.5
2003 4/.....	0.9	1.4	1.5	1.9	3.4	5.4	9.5	14.1	25.0	55.1
2002.....	0.8	1.4	1.6	2.4	3.8	5.0	9.5	14.2	24.5	56.3
2001.....	0.9	1.2	1.6	2.1	3.6	5.4	7.9	14.9	27.0	58.9
2000.....	0.9	1.3	1.6	2.4	3.5	6.2	9.2	13.8	26.8	59.7
1999.....	0.9	1.4	1.4	1.9	3.0	4.9	9.2	15.2	24.8	63.6

White, female

2006 4/.....	0.6	1.0	1.6	2.1	3.3	4.5	7.7	11.1	19.5	41.1
2005 4/.....	0.8	1.1	1.6	2.1	3.3	5.0	8.0	12.6	19.8	51.8
2004 4/.....	0.8	1.0	1.3	1.8	3.1	4.8	7.6	12.7	21.7	52.6
2003 4/.....	0.7	1.0	1.5	2.2	3.2	4.6	7.9	12.2	21.2	57.0
2002.....	0.8	1.3	1.6	2.2	3.2	5.6	8.2	12.3	24.2	56.2
2001.....	0.9	1.0	1.5	2.5	3.7	5.5	8.3	14.3	21.9	58.0
2000.....	0.7	1.0	1.6	2.3	3.6	5.3	7.6	13.6	24.6	58.4
1999.....	0.7	1.0	1.3	2.0	3.5	4.6	8.2	13.1	22.7	58.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

III. Diseases of the blood and blood-forming organs and certain disorders involving the immune mechanism (D50-D89)

Black, both sexes

2006 4/.....	7.5	7.8	9.4	10.5	11.4	12.9	15.4	18.4	25.8	52.7
2005 4/.....	6.8	8.3	9.7	10.5	10.9	15.3	17.9	23.4	33.3	56.7
2004 4/.....	6.8	7.8	9.7	11.4	12.6	13.0	15.3	19.8	25.5	65.7
2003 4/.....	7.6	8.5	10.8	12.0	12.0	13.8	16.5	22.0	32.2	69.4
2002.....	7.6	9.0	10.7	10.5	12.1	13.4	16.9	25.6	31.9	71.9
2001.....	8.5	10.2	10.0	9.9	10.1	12.3	16.8	22.9	35.8	64.8
2000.....	7.0	9.2	10.5	10.3	11.8	13.0	14.7	21.0	35.9	66.6
1999.....	7.4	8.3	8.7	11.3	10.1	12.2	15.6	22.3	29.2	71.1

Black, male

2006 4/.....	7.8	7.6	8.7	10.4	12.2	10.3	13.2	20.1	23.3	46.5
2005 4/.....	6.6	8.3	8.8	11.0	9.1	16.4	15.5	24.6	43.5	50.2
2004 4/.....	6.9	7.7	9.0	13.2	11.8	14.5	15.2	20.9	26.5	61.6
2003 4/.....	8.1	9.3	10.9	12.1	12.5	12.3	20.0	26.5	30.3	66.7
2002.....	7.0	8.1	11.1	10.4	12.1	12.2	19.4	24.8	35.9	78.3
2001.....	7.6	9.6	9.7	8.6	9.6	11.2	18.2	24.6	36.1	52.2
2000.....	6.1	9.4	10.6	8.1	11.8	10.9	15.7	19.8	36.2	73.8
1999.....	6.8	8.2	8.3	10.2	10.4	11.7	16.6	23.3	34.9	63.3

Black, female

2006 4/.....	7.3	7.9	10.0	10.6	10.7	14.9	16.9	17.4	27.1	55.2
2005 4/.....	6.9	8.4	10.4	10.0	12.3	14.5	19.5	22.7	27.8	59.3
2004 4/.....	6.8	7.8	10.3	9.9	13.3	11.9	15.4	19.1	24.9	67.3
2003 4/.....	7.2	7.9	10.8	11.9	11.6	15.0	14.2	19.3	33.3	70.5
2002.....	8.0	9.8	10.4	10.5	12.1	14.2	15.2	26.1	29.8	69.4
2001.....	9.3	10.8	10.1	11.0	10.6	13.1	15.9	21.9	35.6	69.5
2000.....	7.7	9.0	10.5	12.1	11.8	14.6	14.0	21.6	35.7	64.0
1999.....	7.9	8.4	9.1	12.2	9.8	12.5	14.9	21.7	26.4	74.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
All races 3/, both sexes										
2006.....	0.5	0.5	0.5	0.8	1.2	1.4	3.0	5.1	10.5	30.0
2005.....	0.5	0.6	0.6	0.8	1.2	2.0	3.5	6.3	12.8	36.6
2004.....	0.5	0.5	0.6	0.8	1.3	1.8	3.4	5.9	12.2	38.1
2003.....	0.4	0.6	0.5	0.8	1.3	1.9	3.5	6.0	12.7	40.7
2002.....	0.4	0.6	0.6	0.8	1.2	1.9	3.7	6.4	12.9	40.3
2001.....	0.6	0.5	0.7	0.8	1.2	2.0	3.5	7.1	13.6	42.0
2000.....	0.5	0.5	0.7	0.9	1.2	2.2	3.4	6.5	13.9	42.4
1999.....	0.5	0.7	0.6	0.8	1.2	1.8	3.5	6.8	13.7	43.8
All races 3/, male										
2006.....	0.5	0.5	0.5	0.8	1.1	1.5	2.8	5.4	11.6	29.6
2005.....	0.5	0.5	0.5	0.7	1.3	1.9	3.5	6.4	14.1	33.5
2004.....	0.5	0.5	0.6	1.0	1.4	1.9	3.6	6.3	12.8	36.5
2003.....	0.5	0.7	0.5	0.9	1.5	2.3	3.8	6.6	13.3	38.0
2002.....	0.4	0.5	0.6	0.8	1.3	1.7	4.3	6.6	12.4	38.3
2001.....	0.4	0.6	0.6	0.8	1.1	2.1	3.8	7.2	14.3	40.0
2000.....	0.4	0.5	0.7	1.0	1.2	2.2	3.6	6.5	14.6	40.1
1999.....	0.5	0.7	0.6	0.7	1.2	2.1	3.4	7.2	14.8	43.4
All races 3/, female										
2006.....	0.5	0.6	0.6	0.8	1.2	1.3	3.2	5.0	9.8	30.2
2005.....	0.5	0.7	0.7	0.9	1.2	2.0	3.5	6.3	12.1	38.0
2004.....	0.5	0.6	0.6	0.7	1.3	1.7	3.3	5.7	11.8	38.8
2003.....	0.4	0.5	0.6	0.7	1.0	1.5	3.2	5.6	12.3	41.9
2002.....	0.5	0.7	0.7	0.8	1.1	2.1	3.2	6.2	13.2	41.2
2001.....	0.7	0.5	0.7	0.9	1.2	1.9	3.3	7.0	13.2	42.9
2000.....	0.5	0.6	0.7	0.9	1.2	2.2	3.3	6.5	13.6	43.3
1999.....	0.4	0.7	0.6	0.9	1.1	1.6	3.6	6.5	13.0	44.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
White, both sexes										
2006 4/.....	0.1	0.3	0.3	0.6	0.9	1.2	2.8	5.0	10.4	29.6
2005 4/.....	0.1	0.3	0.3	0.6	1.0	1.6	3.1	6.0	12.4	36.6
2004 4/.....	0.2	0.3	0.3	0.5	1.0	1.7	3.2	5.6	12.2	37.7
2003 4/.....	0.1	0.3	0.3	0.4	0.9	1.6	3.2	5.7	12.4	40.6
2002.....	0.1	0.3	0.3	0.6	0.9	1.7	3.4	5.9	12.6	39.7
2001.....	0.2	0.2	0.4	0.6	1.0	1.8	3.2	6.7	13.2	41.7
2000.....	0.1	0.2	0.4	0.7	0.9	1.9	3.2	6.2	13.8	42.0
1999.....	0.1	0.3	0.3	0.4	0.8	1.6	3.2	6.5	13.6	43.0
White, male										
2006 4/.....	*	*	0.3	0.5	0.9	1.4	2.6	5.1	11.5	29.9
2005 4/.....	*	0.3	0.3	0.6	1.0	1.5	3.3	5.9	13.5	33.7
2004 4/.....	0.2	0.3	0.3	0.5	1.2	1.7	3.5	5.7	12.7	36.7
2003 4/.....	*	0.3	0.3	0.5	1.1	2.0	3.3	6.3	13.4	37.9
2002.....	*	0.3	0.3	0.6	1.0	1.6	3.7	6.3	11.9	37.2
2001.....	*	*	0.4	0.5	1.0	1.9	3.3	7.0	14.1	40.4
2000.....	*	*	0.4	0.8	0.9	2.1	3.4	6.1	14.2	39.1
1999.....	*	0.4	0.3	*	0.7	1.9	3.1	7.0	14.8	43.4
White, female										
2006 4/.....	*	0.3	0.3	0.7	1.0	1.1	3.1	4.8	9.7	29.4
2005 4/.....	*	0.4	0.3	0.7	1.0	1.7	3.0	6.1	11.7	37.9
2004 4/.....	*	0.2	0.3	0.5	0.9	1.6	3.0	5.6	11.9	38.2
2003 4/.....	*	*	0.3	0.3	0.8	1.3	3.1	5.3	11.8	41.8
2002.....	*	0.3	0.3	0.5	0.8	1.8	3.1	5.6	13.0	40.8
2001.....	0.3	*	0.4	0.6	1.1	1.6	3.1	6.5	12.6	42.3
2000.....	*	0.3	0.4	0.7	0.9	1.8	3.0	6.3	13.5	43.2
1999.....	*	0.3	0.3	0.5	0.9	1.4	3.4	6.1	12.9	42.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Anemias (D50-D64)										
Black, both sexes										
2006 4/.....	2.9	2.6	2.4	2.9	3.4	3.0	5.5	8.0	13.5	38.9
2005 4/.....	2.9	2.5	2.9	2.5	3.7	5.3	7.3	11.0	20.1	42.5
2004 4/.....	2.4	2.5	3.0	3.9	4.2	3.7	6.2	10.7	13.4	48.2
2003 4/.....	2.5	3.3	2.7	4.1	4.4	4.3	7.1	10.9	19.8	49.3
2002.....	2.4	3.1	3.1	3.2	4.1	4.4	7.9	13.7	19.4	52.3
2001.....	3.2	3.2	2.9	3.2	2.6	3.9	7.4	12.5	21.4	48.5
2000.....	2.7	2.9	3.2	3.0	3.9	4.8	6.1	10.3	18.1	51.0
1999.....	2.8	3.2	2.9	4.2	4.3	3.6	6.6	11.4	16.5	52.9
Black, male										
2006 4/.....	2.9	2.4	2.1	3.3	3.9	*	*	9.4	15.7	29.5
2005 4/.....	3.1	2.4	2.6	*	*	6.7	*	13.4	23.2	37.2
2004 4/.....	2.5	2.1	2.9	5.3	4.5	*	*	13.6	14.3	42.7
2003 4/.....	2.9	3.9	2.6	4.9	6.2	5.7	10.8	12.8	18.5	49.0
2002.....	2.3	2.6	2.6	3.0	5.1	*	11.7	12.9	23.7	56.8
2001.....	2.7	3.9	3.2	3.3	*	*	10.4	12.1	20.1	42.0
2000.....	2.2	3.2	3.4	*	*	*	*	10.9	19.4	55.4
1999.....	2.9	3.3	3.2	4.2	6.2	*	7.8	10.9	*	44.5
Black, female										
2006 4/.....	3.0	2.7	2.7	2.6	3.0	*	5.7	7.2	12.3	42.8
2005 4/.....	2.7	2.6	3.2	2.9	4.0	4.2	8.2	9.6	18.5	44.6
2004 4/.....	2.4	2.8	3.1	2.7	3.9	*	6.2	8.9	12.9	50.4
2003 4/.....	2.1	2.8	2.7	3.3	3.0	*	4.7	9.8	20.5	49.4
2002.....	2.6	3.6	3.5	3.3	3.3	4.6	5.4	14.3	17.1	50.5
2001.....	3.7	2.6	2.6	3.2	*	3.9	5.4	12.8	22.1	50.9
2000.....	3.2	2.7	3.0	3.0	3.9	5.8	6.2	10.0	17.4	49.4
1999.....	2.6	3.2	2.7	4.2	*	*	5.9	11.7	16.4	56.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional anemias (D50-D53)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	*	*	0.7	2.7
2005 4/.....	*	*	*	*	*	*	*	0.4	0.8	3.2
2004 4/.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2003 4/.....	*	*	*	*	*	*	*	0.4	0.7	3.6
2002.....	*	*	*	*	*	*	*	0.3	0.5	3.6
2001.....	*	*	*	*	*	*	*	0.5	0.8	3.7
2000.....	*	*	*	*	*	*	*	0.3	0.8	4.4
1999.....	*	*	*	*	*	*	*	0.3	1.1	3.9
White, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	2.4
2005 4/.....	*	*	*	*	*	*	*	*	*	2.8
2004 4/.....	*	*	*	*	*	*	*	*	*	3.0
2003 4/.....	*	*	*	*	*	*	*	*	*	2.9
2002.....	*	*	*	*	*	*	*	*	*	2.9
2001.....	*	*	*	*	*	*	*	*	*	3.3
2000.....	*	*	*	*	*	*	*	*	*	5.0
1999.....	*	*	*	*	*	*	*	*	*	4.1
White, female										
2006 4/.....	*	*	*	*	*	*	*	*	0.8	2.8
2005 4/.....	*	*	*	*	*	*	*	0.6	0.7	3.4
2004 4/.....	*	*	*	*	*	*	*	0.6	0.8	2.6
2003 4/.....	*	*	*	*	*	*	*	*	0.7	3.9
2002.....	*	*	*	*	*	*	*	*	*	3.9
2001.....	*	*	*	*	*	*	*	*	0.7	3.8
2000.....	*	*	*	*	*	*	*	*	1.1	4.1
1999.....	*	*	*	*	*	*	*	*	1.2	3.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hemolytic anemias (D55-D59)										
All races 3/, both sexes										
2006.....	0.4	0.3	0.3	0.3	0.2	*	0.4	0.5	0.9	1.5
2005.....	0.3	0.3	0.3	0.2	0.3	0.3	0.5	0.5	1.1	1.8
2004.....	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.5	0.8	1.9
2003.....	0.3	0.3	0.3	0.3	0.2	*	0.3	0.6	1.1	1.9
2002.....	0.3	0.4	0.3	0.2	0.2	0.4	0.4	0.7	1.0	2.2
2001.....	0.4	0.3	0.3	0.3	0.2	0.3	0.4	0.7	1.2	2.2
2000.....	0.3	0.3	0.4	0.3	0.2	0.3	0.4	0.6	1.2	2.4
1999.....	0.3	0.4	0.2	0.3	0.2	0.3	0.4	0.7	1.2	2.2
All races 3/, male										
2006.....	0.4	0.2	*	0.3	*	*	*	*	1.0	1.8
2005.....	0.3	0.3	0.2	*	*	*	*	0.6	1.1	2.4
2004.....	0.3	0.2	0.3	0.3	*	*	*	0.8	1.0	2.1
2003.....	0.3	0.4	0.2	0.3	*	*	*	0.7	1.1	1.5
2002.....	0.3	0.3	0.3	*	*	*	*	*	*	2.2
2001.....	0.3	0.4	0.3	*	*	*	*	0.8	1.4	2.2
2000.....	0.3	0.3	0.3	*	*	*	*	*	1.2	2.5
1999.....	0.4	0.4	0.2	*	*	*	*	0.8	1.4	1.8
All races 3/, female										
2006.....	0.4	0.3	0.3	0.3	*	*	0.4	0.6	0.9	1.3
2005.....	0.3	0.3	0.4	0.3	0.3	0.4	0.6	*	1.1	1.5
2004.....	0.3	0.3	0.4	*	0.3	0.4	*	*	0.6	1.8
2003.....	0.3	0.3	0.3	*	*	*	*	0.5	1.1	2.1
2002.....	0.3	0.4	0.4	*	*	0.4	0.4	0.8	1.1	2.2
2001.....	0.5	0.3	0.3	0.3	*	*	*	0.7	1.1	2.2
2000.....	0.4	0.3	0.4	0.3	*	*	0.4	0.7	1.2	2.3
1999.....	0.3	0.4	*	0.4	*	*	0.5	0.7	1.0	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hemolytic anemias (D55-D59)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	0.3	0.5	0.9	1.6
2005 4/.....	*	*	*	*	*	*	0.4	0.5	1.1	1.9
2004 4/.....	*	*	*	*	*	*	*	0.5	0.9	1.9
2003 4/.....	*	*	*	*	*	*	0.3	0.6	1.2	2.0
2002.....	*	*	*	*	*	0.3	0.3	0.6	1.1	2.3
2001.....	*	*	*	*	*	0.3	0.3	0.7	1.2	2.2
2000.....	*	*	*	*	*	*	0.3	0.7	1.3	2.5
1999.....	*	*	*	*	*	0.2	0.4	0.8	1.2	2.4
White, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	2.0
2005 4/.....	*	*	*	*	*	*	*	*	*	2.6
2004 4/.....	*	*	*	*	*	*	*	0.8	1.1	2.0
2003 4/.....	*	*	*	*	*	*	*	0.7	1.2	*
2002.....	*	*	*	*	*	*	*	*	*	2.4
2001.....	*	*	*	*	*	*	*	0.8	1.5	2.2
2000.....	*	*	*	*	*	*	*	*	1.3	2.8
1999.....	*	*	*	*	*	*	*	0.9	1.5	*
White, female										
2006 4/.....	*	*	*	*	*	*	*	0.6	0.9	1.4
2005 4/.....	*	*	*	*	*	*	0.5	*	1.2	1.6
2004 4/.....	*	*	*	*	*	*	*	*	0.7	1.9
2003 4/.....	*	*	*	*	*	*	*	0.5	1.2	2.2
2002.....	*	*	*	*	*	*	*	0.7	1.2	2.2
2001.....	*	*	*	*	*	*	*	0.6	1.0	2.2
2000.....	*	*	*	*	*	*	*	0.7	1.3	2.4
1999.....	*	*	*	*	*	*	0.5	0.7	1.0	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hemolytic anemias (D55-D59)										
Black, both sexes										
2006 4/.....	2.7	1.9	1.8	1.9	*	*	*	*	*	*
2005 4/.....	2.4	2.1	2.0	1.2	*	*	*	*	*	*
2004 4/.....	2.1	1.9	2.4	2.2	1.6	*	*	*	*	*
2003 4/.....	2.2	2.5	1.9	2.2	*	*	*	*	*	*
2002.....	2.2	2.6	2.2	1.6	*	*	*	*	*	*
2001.....	2.8	2.7	2.0	1.9	*	*	*	*	*	*
2000.....	2.3	2.2	2.5	1.6	*	*	*	*	*	*
1999.....	2.3	2.5	1.8	2.7	*	*	*	*	*	*
Black, male										
2006 4/.....	2.9	1.8	*	*	*	*	*	*	*	*
2005 4/.....	2.6	2.1	*	*	*	*	*	*	*	*
2004 4/.....	2.2	1.7	2.1	3.0	*	*	*	*	*	*
2003 4/.....	2.6	2.8	*	*	*	*	*	*	*	*
2002.....	2.1	2.2	*	*	*	*	*	*	*	*
2001.....	2.4	3.1	2.2	*	*	*	*	*	*	*
2000.....	2.0	2.3	2.6	*	*	*	*	*	*	*
1999.....	2.6	2.4	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	2.6	2.1	2.2	*	*	*	*	*	*	*
2005 4/.....	2.2	2.0	2.4	*	*	*	*	*	*	*
2004 4/.....	2.0	2.1	2.6	*	*	*	*	*	*	*
2003 4/.....	1.9	2.3	2.0	*	*	*	*	*	*	*
2002.....	2.3	3.0	2.6	*	*	*	*	*	*	*
2001.....	3.1	2.3	1.9	*	*	*	*	*	*	*
2000.....	2.6	2.1	2.5	*	*	*	*	*	*	*
1999.....	2.0	2.6	*	2.8	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aplastic anemias (D60-D61)										
All races 3/, both sexes										
2006.....	*	0.1	0.1	0.2	0.4	0.5	0.9	1.5	2.6	5.4
2005.....	*	0.1	0.1	0.3	0.3	0.8	1.2	1.7	2.6	4.9
2004.....	0.1	0.1	0.1	0.3	0.5	0.6	1.2	1.5	2.6	5.6
2003.....	*	0.1	0.1	0.2	0.4	0.7	1.1	1.7	2.8	5.7
2002.....	*	0.1	0.1	0.2	0.4	0.6	1.1	1.8	3.2	5.3
2001.....	0.1	0.1	0.2	0.3	0.5	0.6	1.2	2.1	3.5	6.8
2000.....	*	0.1	0.2	0.3	0.5	0.8	1.1	1.8	3.6	7.4
1999.....	*	0.2	0.1	0.2	0.5	0.7	1.3	2.0	3.5	7.7
All races 3/, male										
2006.....	*	*	*	*	0.4	0.6	0.9	1.6	3.4	6.6
2005.....	*	*	*	0.3	0.4	0.8	1.4	1.7	3.4	5.2
2004.....	*	*	*	*	0.5	0.7	1.4	1.7	3.2	6.2
2003.....	*	*	*	*	0.4	1.0	1.3	2.0	3.4	7.4
2002.....	*	*	*	*	0.5	*	1.3	2.3	3.7	7.0
2001.....	*	*	0.2	*	0.5	0.6	1.1	2.3	3.9	7.8
2000.....	*	*	*	0.3	0.5	1.0	1.2	1.8	4.4	7.6
1999.....	*	*	*	*	0.5	1.0	1.2	2.4	4.3	10.1
All races 3/, female										
2006.....	*	*	*	*	0.5	0.4	0.9	1.5	2.1	4.8
2005.....	*	*	*	0.3	0.3	0.7	1.0	1.7	2.2	4.7
2004.....	*	*	*	0.3	0.5	0.6	1.0	1.4	2.2	5.4
2003.....	*	*	*	*	0.4	0.6	0.9	1.5	2.5	5.0
2002.....	*	*	*	0.3	*	0.8	0.9	1.5	2.8	4.5
2001.....	*	*	*	0.4	0.5	0.6	1.3	1.9	3.3	6.3
2000.....	*	*	*	*	0.5	0.7	1.0	1.8	3.1	7.3
1999.....	*	*	*	*	0.4	0.4	1.3	1.8	3.0	6.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aplastic anemias (D60-D61)										
White, both sexes										
2006 4/.....	*	0.1	*	0.2	0.4	0.5	0.9	1.5	2.7	5.5
2005 4/.....	*	0.1	*	0.3	0.3	0.7	1.2	1.7	2.7	5.0
2004 4/.....	*	0.1	*	0.3	0.5	0.6	1.1	1.5	2.7	5.6
2003 4/.....	*	0.1	*	0.2	0.4	0.7	1.0	1.8	2.8	5.8
2002.....	*	0.1	*	0.2	0.3	0.6	1.0	1.9	3.2	5.2
2001.....	*	0.1	0.2	0.3	0.5	0.6	1.2	2.2	3.6	7.1
2000.....	*	0.1	0.1	0.3	0.5	0.8	1.1	1.8	3.7	7.4
1999.....	*	0.1	0.1	*	0.4	0.7	1.2	2.0	3.6	7.8
White, male										
2006 4/.....	*	*	*	*	0.4	0.6	0.8	1.5	3.7	7.0
2005 4/.....	*	*	*	*	*	0.7	1.5	1.8	3.3	5.4
2004 4/.....	*	*	*	*	0.5	0.7	1.5	1.6	3.2	6.2
2003 4/.....	*	*	*	*	*	0.9	1.2	2.2	3.5	7.6
2002.....	*	*	*	*	0.4	*	1.2	2.5	3.6	6.9
2001.....	*	*	*	*	0.5	0.6	0.9	2.4	4.1	8.5
2000.....	*	*	*	*	0.5	1.0	1.3	1.7	4.2	7.0
1999.....	*	*	*	*	*	1.0	1.1	2.4	4.4	10.3
White, female										
2006 4/.....	*	*	*	*	0.5	*	1.1	1.5	2.0	4.8
2005 4/.....	*	*	*	0.3	*	0.7	0.9	1.7	2.3	4.8
2004 4/.....	*	*	*	0.3	0.4	0.6	0.9	1.4	2.3	5.3
2003 4/.....	*	*	*	*	0.4	0.6	0.9	1.5	2.3	5.0
2002.....	*	*	*	*	*	0.8	0.9	1.5	2.9	4.5
2001.....	*	*	*	0.4	0.5	0.5	1.4	2.0	3.3	6.4
2000.....	*	*	*	*	0.5	0.7	1.1	1.9	3.3	7.5
1999.....	*	*	*	*	*	*	1.3	1.8	3.1	6.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute posthemorrhagic and other anemias (D62,D64)

All races 3/, both sexes

2006.....	*	0.1	0.1	0.3	0.5	0.6	1.5	2.8	6.3	20.2
2005.....	0.1	0.1	0.2	0.3	0.6	0.7	1.7	3.8	8.3	26.7
2004.....	*	0.1	0.2	0.3	0.5	0.9	1.8	3.5	8.1	27.8
2003.....	*	0.1	0.2	0.3	0.6	0.9	1.9	3.3	8.1	29.6
2002.....	*	0.1	0.1	0.3	0.6	0.8	2.0	3.6	8.3	29.4
2001.....	*	*	0.1	0.2	0.4	0.9	1.7	3.8	8.1	29.5
2000.....	*	*	0.1	0.3	0.5	1.0	1.8	3.6	8.3	28.2
1999.....	*	0.1	0.2	0.2	0.4	0.8	1.6	3.6	7.9	29.9

All races 3/, male

2006.....	*	*	*	0.3	0.5	0.7	1.3	3.2	6.7	18.7
2005.....	*	*	*	0.3	0.7	0.7	1.5	3.8	8.8	23.2
2004.....	*	*	*	0.4	0.6	1.0	1.9	3.5	7.9	25.4
2003.....	*	*	*	0.4	0.8	1.1	1.9	3.2	8.2	26.2
2002.....	*	*	*	0.4	0.6	0.9	2.4	3.6	7.8	26.3
2001.....	*	*	*	*	*	1.0	2.1	3.7	8.0	26.9
2000.....	*	*	*	0.4	0.6	0.9	1.9	3.6	8.6	24.9
1999.....	*	*	*	*	0.4	0.8	1.7	3.5	8.0	27.7

All races 3/, female

2006.....	*	0.2	*	0.3	0.4	0.5	1.7	2.6	6.1	21.0
2005.....	*	*	*	0.3	0.5	0.8	1.7	3.7	8.0	28.3
2004.....	*	*	*	*	0.4	0.8	1.7	3.4	8.3	28.9
2003.....	*	*	*	*	0.4	0.7	1.8	3.4	8.0	31.1
2002.....	*	*	*	0.3	0.5	0.7	1.8	3.6	8.5	30.7
2001.....	*	*	*	*	0.4	0.9	1.4	3.9	8.1	30.5
2000.....	*	*	*	0.3	0.4	1.1	1.7	3.6	8.1	29.6
1999.....	*	*	*	*	0.5	0.8	1.6	3.7	7.8	30.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute posthemorrhagic and other anemias (D62,D64)

White, both sexes

2006 4/.....	*	*	*	0.3	0.3	0.6	1.4	2.7	6.1	19.8
2005 4/.....	*	*	0.1	0.2	0.5	0.6	1.4	3.4	7.9	26.4
2004 4/.....	*	*	*	0.2	0.4	0.8	1.7	3.2	8.0	27.4
2003 4/.....	*	*	*	0.2	0.4	0.7	1.7	3.0	7.8	29.2
2002.....	*	*	*	0.3	0.4	0.7	1.9	3.0	7.8	28.7
2001.....	*	*	*	0.2	0.3	0.8	1.5	3.4	7.6	28.8
2000.....	*	*	*	0.3	0.3	0.8	1.6	3.4	8.0	27.8
1999.....	*	*	*	*	0.3	0.7	1.4	3.4	7.8	28.9

White, male

2006 4/.....	*	*	*	0.3	*	0.6	1.3	3.1	6.3	18.5
2005 4/.....	*	*	*	*	0.6	*	1.3	3.3	8.5	23.0
2004 4/.....	*	*	*	*	0.4	0.8	1.8	3.1	7.6	25.4
2003 4/.....	*	*	*	*	0.5	0.9	1.6	2.8	8.0	25.9
2002.....	*	*	*	0.3	*	0.8	2.0	3.0	7.2	25.1
2001.....	*	*	*	*	*	0.9	1.8	3.4	7.6	26.4
2000.....	*	*	*	*	*	0.9	1.7	3.5	8.3	24.3
1999.....	*	*	*	*	*	0.7	1.5	3.3	8.0	27.2

White, female

2006 4/.....	*	*	*	0.3	*	0.5	1.4	2.4	6.0	20.3
2005 4/.....	*	*	*	*	0.4	0.7	1.5	3.4	7.5	28.0
2004 4/.....	*	*	*	*	*	0.7	1.6	3.2	8.2	28.3
2003 4/.....	*	*	*	*	*	0.5	1.7	3.1	7.7	30.7
2002.....	*	*	*	*	0.4	0.5	1.8	2.9	8.2	30.2
2001.....	*	*	*	*	*	0.8	1.3	3.4	7.6	29.8
2000.....	*	*	*	*	*	0.8	1.6	3.3	7.8	29.2
1999.....	*	*	*	*	*	0.6	1.3	3.4	7.6	29.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute posthemorrhagic and other anemias (D62,D64)

Black, both sexes

2006 4/.....	*	*	*	*	1.9	*	3.5	4.7	10.6	28.8
2005 4/.....	*	*	*	*	2.0	2.3	4.5	8.9	15.3	34.7
2004 4/.....	*	*	*	*	2.0	2.2	3.0	7.8	11.1	37.8
2003 4/.....	*	*	*	*	2.3	2.3	4.2	8.2	14.0	39.7
2002.....	*	*	*	*	2.3	2.3	4.3	11.2	15.4	41.6
2001.....	*	*	*	*	*	*	4.0	9.2	16.0	41.1
2000.....	*	*	*	*	*	2.8	4.0	7.1	13.3	37.2
1999.....	*	*	*	*	1.9	*	3.9	7.7	10.4	39.9

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	13.7	21.4
2005 4/.....	*	*	*	*	*	*	*	10.3	14.7	31.6
2004 4/.....	*	*	*	*	*	*	*	9.6	*	33.8
2003 4/.....	*	*	*	*	*	*	*	10.0	15.5	37.5
2002.....	*	*	*	*	*	*	7.0	11.0	18.3	45.0
2001.....	*	*	*	*	*	*	*	*	*	37.4
2000.....	*	*	*	*	*	*	*	*	*	33.4
1999.....	*	*	*	*	*	*	*	*	*	31.6

Black, female

2006 4/.....	*	*	*	*	*	*	4.4	*	9.0	31.8
2005 4/.....	*	*	*	*	*	*	4.5	8.0	15.6	36.0
2004 4/.....	*	*	*	*	*	*	*	6.7	11.3	39.4
2003 4/.....	*	*	*	*	*	*	*	7.1	13.2	40.5
2002.....	*	*	*	*	*	*	*	11.3	13.9	40.3
2001.....	*	*	*	*	*	*	*	9.7	16.4	42.5
2000.....	*	*	*	*	*	*	*	7.7	12.7	38.6
1999.....	*	*	*	*	*	*	*	7.7	10.8	42.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

All races 3/, both sexes

2006.....	0.4	0.5	0.8	0.9	1.4	2.1	2.9	4.3	6.9	8.5
2005.....	0.3	0.6	0.8	1.1	1.4	2.1	3.4	4.9	6.3	9.9
2004.....	0.4	0.6	0.8	1.0	1.5	2.0	3.0	4.9	7.5	10.1
2003.....	0.4	0.6	0.9	1.0	1.3	2.3	3.3	5.0	7.0	10.5
2002.....	0.4	0.7	0.7	0.9	1.4	2.2	3.5	4.8	7.7	11.3
2001.....	0.5	0.6	0.7	1.1	1.5	2.3	3.1	4.9	7.6	11.1
2000.....	0.4	0.5	0.8	1.1	1.6	2.6	3.3	4.8	7.9	11.6
1999.....	0.4	0.6	0.7	1.0	1.5	2.2	3.7	5.4	6.9	11.5

All races 3/, male

2006.....	0.4	0.5	0.7	1.0	1.4	2.4	3.0	5.0	7.0	10.1
2005.....	0.3	0.6	0.9	1.2	1.4	2.4	3.4	5.3	8.2	11.7
2004.....	0.4	0.6	0.8	1.2	1.6	2.2	3.5	5.3	8.5	10.9
2003.....	0.4	0.6	0.9	1.1	1.3	2.3	3.8	5.5	7.9	11.9
2002.....	0.4	0.6	0.7	0.9	1.6	2.1	3.5	5.6	8.4	13.9
2001.....	0.4	0.6	0.7	1.0	1.6	2.3	3.3	4.8	9.6	12.3
2000.....	0.4	0.7	0.8	1.0	1.5	2.7	3.8	4.9	8.3	14.6
1999.....	0.4	0.6	0.8	1.0	1.3	2.2	4.1	5.8	6.8	14.3

All races 3/, female

2006.....	0.4	0.5	0.8	0.8	1.5	1.9	2.9	3.8	6.8	7.7
2005.....	0.3	0.6	0.7	0.9	1.3	1.8	3.3	4.7	5.2	9.0
2004.....	0.4	0.5	0.8	0.8	1.3	1.9	2.6	4.7	6.8	9.7
2003.....	0.3	0.6	0.9	1.0	1.3	2.2	2.9	4.7	6.4	9.9
2002.....	0.4	0.8	0.7	0.9	1.2	2.3	3.6	4.3	7.2	10.1
2001.....	0.5	0.6	0.7	1.1	1.4	2.2	3.0	5.0	6.4	10.5
2000.....	0.3	0.4	0.8	1.3	1.7	2.5	2.9	4.7	7.7	10.4
1999.....	0.5	0.5	0.6	1.0	1.8	2.2	3.3	5.1	6.9	10.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

White, both sexes

2006 4/.....	0.3	0.4	0.7	0.8	1.3	2.0	2.8	4.1	6.8	8.5
2005 4/.....	0.3	0.5	0.8	0.9	1.3	1.9	3.2	4.7	6.3	10.0
2004 4/.....	0.3	0.5	0.7	0.8	1.4	1.9	2.9	4.9	7.5	10.0
2003 4/.....	0.3	0.5	0.7	0.9	1.2	2.1	3.3	4.8	6.9	10.3
2002.....	0.4	0.6	0.6	0.8	1.3	2.0	3.5	4.6	7.8	11.3
2001.....	0.4	0.5	0.6	1.0	1.3	2.2	3.1	4.8	7.4	11.0
2000.....	0.3	0.5	0.6	1.0	1.5	2.5	3.1	4.7	7.9	11.8
1999.....	0.4	0.5	0.6	0.8	1.5	1.9	3.5	5.2	6.7	11.4

White, male

2006 4/.....	0.3	0.4	0.6	0.9	1.2	2.2	2.9	4.7	7.1	10.3
2005 4/.....	0.3	0.5	0.9	1.0	1.3	2.2	3.3	5.1	8.0	12.0
2004 4/.....	0.3	0.6	0.7	1.1	1.5	2.0	3.4	5.3	8.5	11.0
2003 4/.....	0.4	0.6	0.7	0.9	1.2	2.1	3.8	5.3	8.0	11.8
2002.....	0.4	0.5	0.6	0.8	1.5	1.9	3.5	5.4	8.2	14.1
2001.....	0.4	0.4	0.6	0.9	1.4	2.2	3.1	4.6	9.6	12.5
2000.....	0.3	0.6	0.6	0.9	1.4	2.5	3.6	4.9	8.3	14.8
1999.....	0.4	0.5	0.7	0.8	1.2	1.9	3.9	5.4	6.3	14.6

White, female

2006 4/.....	0.3	0.4	0.7	0.7	1.4	1.8	2.8	3.6	6.5	7.7
2005 4/.....	0.3	0.5	0.7	0.8	1.2	1.6	3.1	4.4	5.3	9.1
2004 4/.....	0.3	0.5	0.7	0.6	1.3	1.8	2.4	4.7	6.9	9.5
2003 4/.....	0.3	0.4	0.8	0.9	1.2	2.2	2.9	4.4	6.3	9.7
2002.....	0.4	0.6	0.6	0.8	1.1	2.1	3.4	4.0	7.5	10.1
2001.....	0.4	0.5	0.6	1.1	1.2	2.2	3.0	5.0	6.1	10.4
2000.....	0.2	0.4	0.7	1.1	1.6	2.4	2.8	4.5	7.6	10.5
1999.....	0.3	0.4	0.5	0.7	1.8	2.0	3.1	5.0	7.0	10.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Coagulation defects, purpura and other hemorrhagic conditions (D65-D69)

Black, both sexes

2006 4/.....	1.1	1.2	1.6	2.0	2.5	3.6	4.4	6.2	8.3	10.1
2005 4/.....	*	1.4	1.5	2.3	2.2	4.0	5.5	7.5	8.0	9.6
2004 4/.....	1.1	1.1	2.0	2.3	2.4	3.4	4.7	6.1	7.4	13.0
2003 4/.....	0.8	1.5	2.1	2.2	2.3	3.9	4.8	7.5	7.9	13.7
2002.....	*	1.4	1.8	1.9	2.3	4.5	4.8	7.9	7.3	13.1
2001.....	1.0	1.9	1.8	1.6	2.9	3.0	4.2	6.4	10.3	12.3
2000.....	0.8	1.1	2.5	2.4	2.8	3.6	5.3	6.9	10.2	11.3
1999.....	1.1	1.1	1.8	2.8	2.3	4.6	5.4	8.2	8.7	13.7

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	1.6	*	3.1	*	5.0	*	*	14.0	*
2004 4/.....	*	*	2.0	*	*	*	*	*	*	*
2003 4/.....	*	*	2.3	*	*	*	*	*	*	*
2002.....	*	*	2.3	*	*	*	*	*	*	*
2001.....	*	1.9	*	*	*	*	*	*	*	*
2000.....	*	*	2.6	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	5.5	*	10.0	*	*

Black, female

2006 4/.....	*	*	1.6	1.9	*	*	4.6	5.9	10.1	9.1
2005 4/.....	*	*	*	*	*	*	5.8	7.7	*	9.8
2004 4/.....	1.3	*	2.1	*	*	*	4.4	5.9	*	12.6
2003 4/.....	*	1.7	1.9	*	*	*	*	7.1	8.1	13.8
2002.....	*	1.8	*	*	*	4.6	5.4	8.0	*	12.3
2001.....	*	1.8	*	*	*	*	*	5.8	9.8	13.9
2000.....	*	*	2.4	3.0	*	*	4.7	7.7	11.0	10.3
1999.....	*	*	*	2.9	*	4.0	5.4	7.2	*	13.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of blood and blood-forming organs (D70-D76)

All races 3/, both sexes

2006.....	*	0.1	0.2	0.3	0.4	0.7	1.0	1.4	1.8	2.7
2005.....	0.1	0.1	0.2	0.3	0.4	0.7	0.8	1.2	2.1	3.1
2004.....	0.1	0.1	0.2	0.3	0.4	0.6	1.1	1.3	1.9	3.0
2003.....	0.1	0.1	0.3	0.3	0.5	0.6	1.0	1.4	2.0	3.9
2002.....	0.1	0.1	0.3	0.4	0.5	0.7	1.1	1.6	2.6	3.4
2001.....	0.1	0.1	0.2	0.3	0.5	0.8	0.9	1.8	2.0	3.9
2000.....	0.1	0.2	0.3	0.3	0.5	0.8	1.0	1.6	2.2	3.4
1999.....	0.1	0.1	0.2	0.4	0.4	0.5	1.2	1.3	2.2	3.6

All races 3/, male

2006.....	*	*	0.3	0.3	0.5	0.8	1.4	1.8	2.2	2.8
2005.....	*	*	0.3	0.4	0.5	0.7	0.9	1.5	2.8	3.2
2004.....	*	*	0.3	0.3	0.5	0.7	1.2	1.5	2.3	3.2
2003.....	*	*	0.3	*	0.5	0.6	1.2	1.7	2.4	4.0
2002.....	*	*	0.3	0.4	0.6	0.8	1.4	1.8	2.7	3.5
2001.....	*	*	0.3	0.3	0.6	0.8	0.8	2.2	2.2	3.9
2000.....	*	0.2	*	*	0.5	1.0	1.2	1.8	2.7	4.1
1999.....	*	*	*	0.4	0.5	0.5	1.5	1.8	2.7	4.0

All races 3/, female

2006.....	*	*	0.2	0.3	0.4	0.6	0.8	1.1	1.5	2.7
2005.....	*	*	*	0.3	0.4	0.7	0.7	1.0	1.7	3.1
2004.....	*	*	*	*	0.3	0.5	1.0	1.1	1.7	2.9
2003.....	*	*	0.2	0.3	0.5	0.6	0.8	1.2	1.8	3.9
2002.....	*	*	0.2	0.3	0.5	0.6	0.8	1.4	2.5	3.3
2001.....	*	*	0.2	0.3	0.5	0.7	0.9	1.5	1.9	3.8
2000.....	*	*	0.3	*	0.6	0.7	0.8	1.4	1.9	3.1
1999.....	*	*	0.3	0.4	0.4	0.6	0.9	1.0	1.8	3.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of blood and blood-forming organs (D70-D76)

White, both sexes

2006 4/.....	*	0.1	0.2	0.3	0.5	0.7	1.0	1.5	1.9	2.9
2005 4/.....	0.1	0.1	0.2	0.3	0.4	0.7	0.8	1.2	2.1	3.3
2004 4/.....	*	0.1	0.2	0.2	0.4	0.6	1.1	1.4	2.0	3.2
2003 4/.....	*	0.1	0.2	0.2	0.6	0.6	1.1	1.4	2.1	4.1
2002.....	0.1	0.1	0.3	0.4	0.6	0.7	1.1	1.6	2.6	3.4
2001.....	0.1	0.1	0.2	0.3	0.5	0.7	0.9	1.9	2.0	3.9
2000.....	*	0.1	0.2	0.3	0.5	0.8	1.0	1.6	2.2	3.5
1999.....	0.1	*	0.2	0.4	0.4	0.5	1.2	1.4	2.2	3.7

White, male

2006 4/.....	*	*	*	0.3	0.6	0.8	1.5	1.8	2.4	2.9
2005 4/.....	*	*	0.3	0.3	0.5	0.6	0.9	1.5	2.7	3.3
2004 4/.....	*	*	0.3	*	0.5	0.7	1.2	1.6	2.2	3.4
2003 4/.....	*	*	0.3	*	0.6	0.6	1.2	1.6	2.5	4.4
2002.....	*	*	0.3	0.4	0.6	0.8	1.3	1.8	2.9	3.5
2001.....	*	*	*	*	0.6	0.7	0.9	2.4	2.1	4.1
2000.....	*	*	*	*	*	1.0	1.2	1.7	2.8	4.1
1999.....	*	*	*	0.4	0.5	*	1.6	1.9	2.7	4.0

White, female

2006 4/.....	*	*	*	0.3	0.4	0.6	0.7	1.2	1.6	2.8
2005 4/.....	*	*	*	0.3	0.4	0.7	0.7	1.0	1.7	3.3
2004 4/.....	*	*	*	*	*	0.6	1.0	1.2	1.8	3.0
2003 4/.....	*	*	*	0.3	0.5	0.6	0.9	1.2	1.9	4.0
2002.....	*	*	*	0.3	0.5	0.6	0.9	1.5	2.4	3.3
2001.....	*	*	*	*	0.4	0.7	1.0	1.5	1.9	3.8
2000.....	*	*	0.3	*	0.6	0.6	0.9	1.5	1.9	3.2
1999.....	*	*	0.3	*	*	0.6	0.9	1.0	1.9	3.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain disorders involving the immune mechanism (D80-D89)

All races 3/, both sexes

2006.....	0.6	0.7	0.9	1.0	1.0	1.3	1.2	1.4	1.5	1.2
2005.....	0.6	0.7	0.9	0.9	1.0	1.2	1.3	1.3	1.3	1.6
2004.....	0.6	0.7	0.8	0.9	1.0	1.3	1.1	1.1	1.2	1.7
2003.....	0.7	0.7	0.9	1.0	1.0	1.1	1.3	1.2	1.3	1.6
2002.....	0.7	0.8	1.0	1.0	1.1	1.1	1.0	1.1	1.5	1.8
2001.....	0.7	0.9	0.8	0.8	1.1	1.1	1.2	1.2	1.3	1.5
2000.....	0.6	0.8	0.8	0.8	0.9	0.8	1.1	1.3	1.8	1.6
1999.....	0.6	0.7	0.6	0.7	0.7	0.9	0.9	1.0	1.0	1.4

All races 3/, male

2006.....	0.6	0.7	0.8	0.9	1.0	0.6	0.7	1.3	1.2	1.4
2005.....	0.6	0.7	0.8	1.0	0.8	0.9	1.1	1.1	1.5	1.3
2004.....	0.7	0.8	0.8	0.8	0.8	1.2	0.7	0.6	1.0	1.5
2003.....	0.8	0.7	0.8	0.7	0.8	0.9	1.3	1.1	1.2	*
2002.....	0.7	0.9	0.9	1.0	1.0	0.9	1.0	0.8	1.5	1.5
2001.....	0.7	0.8	0.8	0.6	0.9	0.7	0.8	1.1	1.2	1.8
2000.....	0.6	0.8	0.7	0.7	0.9	0.7	1.1	1.0	1.7	1.6
1999.....	0.5	0.8	0.6	0.6	0.7	0.7	0.8	1.0	*	*

All races 3/, female

2006.....	0.5	0.7	1.0	1.1	1.1	1.8	1.5	1.6	1.7	1.2
2005.....	0.6	0.7	1.0	0.9	1.2	1.5	1.4	1.4	1.2	1.7
2004.....	0.5	0.6	0.8	1.0	1.2	1.3	1.5	1.5	1.2	1.9
2003.....	0.7	0.7	0.9	1.2	1.2	1.2	1.4	1.3	1.3	1.7
2002.....	0.8	0.7	1.0	1.0	1.2	1.4	1.1	1.3	1.5	1.9
2001.....	0.7	1.0	0.9	1.0	1.3	1.4	1.6	1.3	1.4	1.4
2000.....	0.6	0.8	0.8	0.9	0.9	0.9	1.0	1.5	1.9	1.5
1999.....	0.6	0.7	0.6	0.8	0.8	1.0	1.0	1.1	1.0	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain disorders involving the immune mechanism (D80-D89)

White, both sexes

2006 4/.....	0.2	0.2	0.4	0.5	0.6	0.8	0.9	1.3	1.4	1.2
2005 4/.....	0.3	0.2	0.4	0.5	0.7	0.8	1.0	1.1	1.1	1.5
2004 4/.....	0.3	0.3	0.4	0.5	0.6	0.9	0.9	1.0	1.0	1.8
2003 4/.....	0.2	0.3	0.3	0.5	0.6	0.7	1.1	1.1	1.2	1.5
2002.....	0.2	0.3	0.4	0.6	0.7	0.8	0.9	1.1	1.4	1.8
2001.....	0.2	0.3	0.4	0.4	0.8	0.8	0.9	1.1	1.2	1.6
2000.....	0.3	0.3	0.3	0.4	0.6	0.5	1.0	1.2	1.6	1.5
1999.....	0.2	0.3	0.3	0.4	0.5	0.7	0.7	1.0	0.9	1.4

White, male

2006 4/.....	0.2	0.2	0.4	0.5	0.6	0.5	*	1.2	1.2	*
2005 4/.....	0.3	0.3	0.4	0.6	0.6	0.6	0.9	1.0	1.2	*
2004 4/.....	0.3	0.4	0.4	0.5	0.6	0.9	*	*	*	1.5
2003 4/.....	0.3	0.4	0.3	0.3	0.6	0.8	1.2	0.9	1.1	*
2002.....	0.2	0.4	0.4	0.6	0.8	0.6	1.0	0.8	1.6	*
2001.....	0.2	0.4	0.4	*	0.7	0.6	0.6	0.9	1.1	2.0
2000.....	0.3	0.3	0.4	0.5	0.7	0.6	1.1	1.0	1.5	*
1999.....	0.3	0.4	0.3	0.4	0.6	0.6	0.6	1.0	*	*

White, female

2006 4/.....	*	0.3	0.4	0.5	0.6	1.1	1.2	1.4	1.5	1.2
2005 4/.....	0.2	*	0.5	0.4	0.7	1.0	1.1	1.2	1.0	1.7
2004 4/.....	*	*	0.3	0.5	0.6	0.9	1.2	1.2	1.0	1.9
2003 4/.....	*	0.3	0.3	0.6	0.7	0.6	1.0	1.3	1.2	1.7
2002.....	*	0.2	0.4	0.5	0.6	1.0	0.8	1.2	1.3	1.9
2001.....	*	0.3	0.3	0.5	0.9	1.0	1.1	1.2	1.3	1.5
2000.....	0.2	*	0.3	*	0.5	0.5	0.9	1.3	1.6	1.5
1999.....	*	*	*	0.4	0.5	0.7	0.8	1.0	0.9	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain disorders involving the immune mechanism (D80-D89)

Black, both sexes

2006 4/.....	3.4	3.8	4.8	5.2	5.1	5.8	4.1	*	*	*
2005 4/.....	3.2	4.3	5.0	4.9	4.5	5.4	4.2	3.7	*	*
2004 4/.....	3.2	3.9	4.5	4.8	5.3	5.0	3.8	*	*	*
2003 4/.....	4.2	3.5	5.8	5.4	4.8	4.9	4.1	*	*	*
2002.....	4.4	4.3	5.5	5.0	5.0	4.1	3.1	*	*	*
2001.....	4.0	5.0	4.8	4.8	4.0	4.0	4.6	*	*	*
2000.....	3.3	4.9	4.5	4.5	4.2	3.5	*	*	*	*
1999.....	3.4	3.7	3.6	3.3	2.7	3.1	*	*	*	*

Black, male

2006 4/.....	3.8	4.0	4.3	4.5	4.8	*	*	*	*	*
2005 4/.....	3.1	4.0	4.5	4.9	*	*	*	*	*	*
2004 4/.....	3.6	3.9	4.1	4.6	3.9	4.8	*	*	*	*
2003 4/.....	4.1	3.8	5.7	4.1	*	*	*	*	*	*
2002.....	4.0	4.3	5.8	4.8	4.0	*	*	*	*	*
2001.....	3.9	3.7	3.9	3.4	*	*	*	*	*	*
2000.....	3.1	4.7	4.3	*	*	*	*	*	*	*
1999.....	2.8	3.5	3.0	*	*	*	*	*	*	*

Black, female

2006 4/.....	3.1	3.6	5.1	5.8	5.4	8.7	5.1	*	*	*
2005 4/.....	3.2	4.5	5.5	5.0	5.5	6.2	4.3	*	*	*
2004 4/.....	2.8	4.0	4.9	4.9	6.4	5.0	*	*	*	*
2003 4/.....	4.3	3.3	5.9	6.5	5.9	7.1	5.1	*	*	*
2002.....	4.9	4.3	5.3	5.1	5.9	4.8	*	*	*	*
2001.....	4.2	6.2	5.6	5.9	4.6	5.3	5.8	*	*	*
2000.....	3.4	5.1	4.8	5.9	4.4	4.6	*	*	*	*
1999.....	4.0	3.9	4.1	4.0	3.5	4.2	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

All races 3/, both sexes

2006.....	9.3	14.4	23.9	38.2	59.1	85.7	123.3	179.9	264.5	433.0
2005.....	9.0	14.4	24.2	38.9	60.5	89.1	131.3	192.4	280.9	480.7
2004.....	8.6	14.6	23.5	37.9	59.1	87.8	130.2	187.8	278.7	470.6
2003.....	8.8	14.8	24.3	39.0	60.7	91.7	132.9	194.0	284.5	492.8
2002.....	9.2	14.0	24.2	39.1	59.2	91.3	133.0	197.1	288.8	504.8
2001.....	8.0	14.0	23.8	37.3	59.1	90.6	132.7	194.8	286.9	519.2
2000.....	8.4	13.3	22.9	37.3	58.9	89.4	131.9	195.7	286.5	521.0
1999.....	8.1	13.3	21.7	36.5	58.8	90.3	132.3	191.6	285.9	529.8

All races 3/, male

2006.....	11.1	17.9	29.2	45.4	71.3	100.4	143.5	207.4	302.0	450.1
2005.....	11.2	18.2	29.0	47.4	72.2	102.1	151.3	217.9	313.6	489.7
2004.....	10.3	17.6	28.6	44.5	70.2	102.7	148.6	213.4	315.6	481.0
2003.....	10.8	17.6	28.7	45.5	71.0	106.9	151.4	220.3	318.9	490.8
2002.....	11.4	16.3	28.7	46.4	68.9	102.1	147.7	217.5	313.8	504.5
2001.....	9.5	16.8	27.9	42.9	67.4	102.8	148.3	215.1	310.2	507.7
2000.....	9.8	16.0	26.4	42.5	66.6	98.5	146.0	215.1	311.8	517.4
1999.....	9.4	16.1	25.2	41.9	65.3	98.8	146.0	213.4	312.5	521.0

All races 3/, female

2006.....	7.5	11.0	18.7	31.3	48.0	72.8	106.9	159.7	241.0	425.0
2005.....	6.7	10.7	19.6	30.9	49.7	77.8	115.0	173.8	260.5	476.6
2004.....	6.9	11.7	18.6	31.7	48.9	74.9	115.5	169.4	255.9	465.9
2003.....	6.9	12.1	20.1	32.7	51.3	78.6	118.1	175.2	263.5	493.7
2002.....	7.0	11.7	19.8	32.2	50.4	81.9	121.2	182.7	273.5	504.9
2001.....	6.5	11.3	19.9	31.9	51.6	80.0	120.3	180.6	272.9	524.0
2000.....	7.0	10.7	19.4	32.4	52.0	81.6	120.7	182.3	271.5	522.5
1999.....	6.8	10.6	18.3	31.4	52.9	83.0	121.6	176.4	270.3	533.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

White, both sexes

2006 4/.....	8.2	13.0	21.1	34.0	53.1	77.2	113.7	168.5	250.2	418.8
2005 4/.....	8.1	12.7	21.7	34.7	54.7	80.2	120.4	180.2	267.8	465.3
2004 4/.....	7.6	13.2	20.7	33.8	52.8	79.4	118.7	174.8	264.7	453.5
2003 4/.....	8.0	13.1	21.4	34.6	54.2	82.9	121.8	181.0	269.3	475.4
2002.....	8.3	12.3	21.2	34.5	52.5	81.0	121.2	183.8	274.3	487.1
2001.....	7.2	12.2	21.0	32.8	52.6	81.3	121.4	181.3	272.1	502.4
2000.....	7.7	11.6	19.9	32.9	52.9	80.1	120.6	181.8	270.5	503.5
1999.....	7.2	11.4	18.7	31.1	51.4	80.8	119.6	178.3	269.9	511.4

White, male

2006 4/.....	9.9	16.5	26.3	41.1	65.5	93.1	134.7	198.0	292.3	441.5
2005 4/.....	10.4	16.6	26.6	42.9	66.6	94.1	141.6	207.6	304.5	481.0
2004 4/.....	9.3	16.1	26.1	40.3	64.4	94.5	138.5	202.5	307.0	470.5
2003 4/.....	9.9	15.9	26.0	41.6	64.9	98.9	141.8	210.2	308.1	484.1
2002.....	10.5	14.6	25.9	42.1	62.2	93.2	138.7	208.7	303.9	498.9
2001.....	8.9	14.8	25.1	38.7	61.2	95.1	139.9	204.3	297.8	500.5
2000.....	9.2	14.4	23.8	38.3	60.9	91.0	137.2	204.2	302.4	510.5
1999.....	8.5	14.2	22.5	36.8	58.9	90.4	136.6	203.3	301.4	504.7

White, female

2006 4/.....	6.6	9.6	16.1	27.2	41.7	63.0	96.3	146.4	223.6	408.1
2005 4/.....	5.8	8.9	16.9	26.9	43.7	67.9	102.9	159.9	244.8	458.1
2004 4/.....	5.9	10.3	15.4	27.6	42.1	66.0	102.5	154.5	238.5	445.8
2003 4/.....	6.0	10.2	16.9	27.9	44.4	68.9	105.4	159.9	245.4	471.5
2002.....	6.1	10.1	16.5	27.3	43.6	70.2	107.0	166.0	256.1	482.1
2001.....	5.4	9.6	17.0	27.1	44.6	69.2	106.5	165.0	256.6	503.2
2000.....	6.2	8.9	16.1	27.8	45.5	70.5	107.4	166.0	251.5	500.6
1999.....	5.8	8.7	15.0	25.6	44.5	72.4	106.1	160.7	251.2	514.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

IV. Endocrine, nutritional and metabolic diseases (E00-E88)

Black, both sexes

2006 4/.....	17.5	27.2	48.0	76.7	119.6	171.2	227.4	319.5	453.7	635.1
2005 4/.....	16.4	28.6	46.7	78.4	122.5	175.6	249.3	342.9	465.8	694.9
2004 4/.....	16.6	27.2	48.6	78.0	120.8	174.2	257.1	351.2	465.4	719.1
2003 4/.....	16.5	29.2	50.1	83.1	123.9	179.3	255.5	354.1	484.1	736.9
2002.....	17.3	28.3	51.2	83.3	125.4	186.0	261.4	356.3	491.9	757.0
2001.....	15.1	28.7	50.2	81.1	121.3	180.4	263.0	363.7	482.9	763.1
2000.....	15.2	28.3	49.9	79.6	117.6	183.3	258.5	375.1	503.2	766.1
1999.....	16.2	28.7	49.8	84.7	127.7	183.9	268.1	349.9	497.3	763.6

Black, male

2006 4/.....	21.6	32.1	57.6	87.2	138.4	187.9	258.7	349.0	471.0	610.3
2005 4/.....	19.0	34.2	51.3	93.4	139.8	189.7	272.3	375.5	481.1	636.8
2004 4/.....	19.2	31.5	54.5	88.4	134.5	199.6	277.1	370.3	476.3	683.5
2003 4/.....	19.3	32.9	55.0	91.3	138.4	197.7	272.0	372.0	497.6	619.3
2002.....	20.6	32.0	56.7	92.7	143.3	192.0	264.0	352.7	492.9	610.0
2001.....	16.2	33.8	55.1	88.9	133.2	189.6	265.6	375.4	511.2	658.1
2000.....	16.5	31.6	53.2	87.3	126.7	180.1	265.1	376.6	480.5	655.0
1999.....	18.0	33.5	53.0	92.9	130.5	193.2	261.9	347.6	485.7	746.4

Black, female

2006 4/.....	13.9	22.9	39.8	68.1	104.9	158.9	205.8	301.8	444.5	645.2
2005 4/.....	14.2	23.6	42.7	66.0	109.0	165.2	233.5	323.5	457.7	718.3
2004 4/.....	14.3	23.5	43.6	69.5	110.1	155.2	243.6	339.9	459.6	733.2
2003 4/.....	14.0	25.9	45.9	76.3	112.5	165.6	244.4	343.5	477.0	782.6
2002.....	14.4	25.0	46.5	75.5	111.4	181.5	259.7	358.4	491.4	813.3
2001.....	14.2	24.2	46.0	74.8	112.0	173.7	261.2	356.7	468.5	802.0
2000.....	14.0	25.4	47.1	73.3	110.5	185.7	254.2	374.2	514.6	807.4
1999.....	14.7	24.6	47.2	78.0	125.5	177.0	272.3	351.3	503.0	770.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Endocrine diseases (E00-E34)										
All races 3/, both sexes										
2006.....	6.4	9.9	17.6	29.3	47.3	69.7	100.5	145.6	207.2	307.0
2005.....	5.9	10.2	17.7	30.3	48.4	72.9	106.7	155.5	218.9	337.3
2004.....	6.0	10.4	17.6	29.9	48.2	72.6	107.7	153.4	220.9	330.4
2003.....	6.1	10.7	18.3	31.2	49.9	76.3	111.0	159.7	225.2	342.7
2002.....	6.3	10.2	18.5	30.7	48.4	76.9	111.3	161.9	226.5	347.6
2001.....	5.6	10.2	18.3	30.1	49.3	76.6	110.8	160.4	227.1	350.3
2000.....	5.8	9.8	17.7	30.3	48.9	75.2	110.7	160.9	222.4	345.2
1999.....	5.7	10.3	17.0	30.2	49.7	77.0	111.6	157.7	223.2	343.6
All races 3/, male										
2006.....	7.8	12.5	21.8	34.9	57.6	81.7	118.0	170.3	238.4	333.1
2005.....	7.4	13.1	21.4	36.8	57.7	83.4	122.7	177.2	249.6	359.1
2004.....	7.3	12.7	21.5	34.9	57.5	84.1	122.8	175.5	252.8	349.1
2003.....	7.6	12.9	21.7	36.1	58.4	87.9	126.2	182.8	255.1	354.4
2002.....	7.9	12.1	21.9	36.6	56.4	85.4	122.9	178.9	248.9	365.2
2001.....	6.8	12.3	21.8	34.9	55.3	85.7	123.0	175.6	247.7	362.2
2000.....	7.0	11.6	20.1	34.0	54.5	81.7	121.4	177.0	247.4	360.9
1999.....	6.6	12.5	19.8	34.1	54.6	83.6	121.3	175.9	248.6	361.8
All races 3/, female										
2006.....	5.0	7.3	13.6	24.0	38.0	59.1	86.2	127.5	187.8	294.8
2005.....	4.4	7.5	14.3	24.1	39.9	63.8	93.7	139.6	199.8	327.3
2004.....	4.6	8.0	13.9	25.2	39.6	62.6	95.5	137.6	201.2	322.0
2003.....	4.5	8.5	15.1	26.5	42.2	66.2	98.9	143.3	206.9	337.6
2002.....	4.7	8.3	15.3	25.2	41.2	69.6	102.0	149.9	212.9	339.9
2001.....	4.5	8.1	15.0	25.6	44.0	68.7	101.2	149.7	214.7	345.3
2000.....	4.7	8.1	15.4	26.9	43.8	69.5	102.3	149.7	207.7	338.8
1999.....	4.8	8.1	14.3	26.5	45.3	71.3	104.1	145.0	208.4	336.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Endocrine diseases (E00-E34)										
White, both sexes										
2006 4/.....	5.5	8.7	15.1	25.5	41.4	61.6	91.0	134.2	193.7	292.8
2005 4/.....	5.2	8.9	15.4	26.4	42.8	64.5	96.1	143.9	206.2	322.1
2004 4/.....	5.2	9.2	15.0	25.9	42.2	64.7	96.6	140.8	207.2	313.3
2003 4/.....	5.3	9.1	15.6	27.0	43.6	67.7	100.2	147.6	211.2	326.4
2002.....	5.5	8.7	15.7	26.5	42.2	67.2	100.5	149.5	213.6	331.6
2001.....	5.0	8.8	15.7	25.8	43.1	67.9	100.1	147.7	214.1	335.2
2000.....	5.2	8.3	15.0	26.4	43.1	66.2	100.1	147.8	207.5	329.3
1999.....	4.9	8.6	14.3	25.0	42.6	68.1	99.9	145.2	208.9	325.9
White, male										
2006 4/.....	6.8	11.2	19.0	30.7	51.5	74.4	109.0	161.2	229.2	324.2
2005 4/.....	6.7	11.6	18.8	32.4	52.1	75.7	112.7	167.6	240.9	350.8
2004 4/.....	6.5	11.3	19.0	30.8	51.7	76.2	113.0	165.2	244.2	338.1
2003 4/.....	6.9	11.3	19.0	32.1	52.1	80.1	117.0	173.8	246.1	347.0
2002.....	7.0	10.5	19.1	32.4	50.2	76.9	114.5	170.9	240.8	361.5
2001.....	6.3	10.7	19.1	30.8	49.6	78.5	115.3	165.5	237.4	356.8
2000.....	6.4	10.0	17.6	30.1	49.3	74.6	113.2	167.2	238.9	353.0
1999.....	5.7	10.8	17.2	29.3	48.4	75.9	113.0	167.1	240.0	349.4
White, female										
2006 4/.....	4.2	6.1	11.3	20.5	32.0	50.2	76.1	114.0	171.2	278.0
2005 4/.....	3.7	6.2	12.1	20.6	34.2	54.5	82.4	126.3	184.5	308.9
2004 4/.....	3.9	7.1	11.1	21.3	33.4	54.4	83.1	123.0	184.2	302.1
2003 4/.....	3.8	6.9	12.2	22.2	35.7	56.8	86.4	128.7	189.6	317.3
2002.....	4.0	6.9	12.5	20.8	34.7	58.7	89.1	134.2	196.9	318.5
2001.....	3.6	6.9	12.5	21.0	37.2	58.5	87.8	135.1	200.0	326.1
2000.....	4.0	6.7	12.4	22.9	37.4	58.8	89.6	134.2	188.8	319.7
1999.....	4.1	6.5	11.4	21.0	37.4	61.3	89.4	129.9	190.5	316.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Endocrine diseases (E00-E34)										
Black, both sexes										
2006 4/.....	13.2	19.9	38.3	63.1	103.8	148.1	197.0	278.8	377.5	490.3
2005 4/.....	11.5	21.3	37.1	65.2	104.9	152.2	216.1	290.3	387.9	530.3
2004 4/.....	11.9	20.1	39.8	66.1	104.0	150.5	224.4	304.5	394.4	555.3
2003 4/.....	12.1	23.0	40.6	71.4	108.8	158.0	225.2	304.4	402.1	552.8
2002.....	12.6	22.0	42.1	69.3	107.6	163.3	226.6	305.3	400.6	559.8
2001.....	11.4	21.6	41.1	70.7	106.2	159.8	230.8	312.3	394.1	555.5
2000.....	10.8	22.4	41.4	67.6	103.3	162.5	228.0	325.0	415.6	550.6
1999.....	11.8	23.7	41.5	74.8	113.5	162.3	233.8	301.0	408.8	552.6
Black, male										
2006 4/.....	16.7	24.6	47.0	73.1	122.3	165.2	227.0	302.9	386.8	476.3
2005 4/.....	13.2	26.6	42.9	79.7	120.3	164.2	241.0	316.0	398.9	484.3
2004 4/.....	14.3	25.1	45.0	75.5	118.1	172.0	242.8	315.7	399.6	525.5
2003 4/.....	14.7	26.5	44.6	79.0	122.8	172.1	235.2	312.7	397.9	473.4
2002.....	15.4	25.9	47.9	77.6	121.2	167.3	228.0	299.5	391.3	427.8
2001.....	12.3	26.2	46.3	77.7	113.5	164.4	226.5	317.3	409.3	469.7
2000.....	12.1	25.8	44.2	73.8	108.9	156.9	228.7	317.5	389.5	474.0
1999.....	13.6	28.3	45.1	80.5	115.2	168.2	220.8	288.7	388.0	523.8
Black, female										
2006 4/.....	10.1	15.8	30.9	54.8	89.4	135.5	176.5	264.3	372.6	496.1
2005 4/.....	10.1	16.8	32.3	53.2	92.9	143.3	199.1	274.9	382.1	548.8
2004 4/.....	9.8	15.7	35.5	58.3	93.1	134.6	212.0	297.9	391.6	567.0
2003 4/.....	9.9	19.9	37.2	65.2	97.8	147.5	218.4	299.5	404.2	583.7
2002.....	10.1	18.7	37.2	62.4	97.0	160.4	225.6	308.7	405.4	610.4
2001.....	10.6	17.6	36.7	64.9	100.5	156.4	233.6	309.2	386.3	587.3
2000.....	9.7	19.4	39.0	62.5	98.9	166.7	227.5	329.5	428.8	579.1
1999.....	10.3	19.7	38.5	70.1	112.1	158.0	242.4	308.4	419.2	563.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of thyroid gland (E00-E07)										
All races 3/, both sexes										
2006.....	0.1	0.1	0.2	0.2	0.3	0.7	1.2	2.1	5.1	18.3
2005.....	0.2	0.1	0.2	0.3	0.4	0.6	1.0	2.3	5.2	21.3
2004.....	0.1	0.2	0.2	0.2	0.5	0.5	1.0	2.2	5.3	19.2
2003.....	0.1	0.2	0.2	0.3	0.4	0.7	1.2	2.9	6.1	21.1
2002.....	0.1	0.1	0.2	0.3	0.4	0.5	1.4	2.5	6.0	22.7
2001.....	0.1	0.1	0.2	0.3	0.4	0.6	1.1	3.0	5.8	23.2
2000.....	0.1	*	0.2	0.3	0.4	0.5	1.3	2.7	5.9	21.1
1999.....	0.1	0.2	0.2	0.3	0.3	0.6	1.3	2.5	6.2	22.4
All races 3/, male										
2006.....	*	*	0.2	*	*	0.4	1.1	1.3	3.3	8.4
2005.....	*	*	*	*	0.5	0.4	0.6	1.4	3.2	10.8
2004.....	*	*	*	*	0.4	0.5	0.8	1.5	3.5	11.2
2003.....	*	*	*	*	*	0.6	0.6	1.9	2.9	10.5
2002.....	*	*	*	*	*	*	0.8	1.6	3.0	12.7
2001.....	*	*	*	*	*	*	0.7	2.3	2.8	14.1
2000.....	*	*	*	*	*	*	0.9	1.5	2.6	12.1
1999.....	*	*	*	*	*	0.5	0.8	1.5	4.4	12.9
All races 3/, female										
2006.....	*	*	0.2	0.3	0.4	1.0	1.2	2.7	6.3	22.9
2005.....	0.3	*	0.3	0.4	0.4	0.8	1.3	3.0	6.4	26.1
2004.....	*	0.2	0.2	*	0.5	0.6	1.3	2.7	6.4	22.8
2003.....	*	0.2	0.3	0.4	0.6	0.8	1.7	3.6	8.0	25.8
2002.....	*	0.2	0.3	0.4	0.6	0.6	1.8	3.1	7.8	27.0
2001.....	*	*	0.3	0.4	0.5	0.8	1.4	3.6	7.6	27.1
2000.....	0.2	*	0.3	0.4	0.6	0.7	1.5	3.5	7.8	24.8
1999.....	*	0.3	*	0.4	0.4	0.8	1.7	3.2	7.3	26.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of thyroid gland (E00-E07)										
White, both sexes										
2006 4/.....	*	*	0.2	0.2	0.3	0.7	1.1	2.1	5.2	18.9
2005 4/.....	0.1	*	0.2	0.2	0.4	0.6	0.9	2.2	5.3	22.1
2004 4/.....	*	0.1	0.2	0.2	0.4	0.5	0.9	2.0	5.3	19.8
2003 4/.....	*	0.1	0.2	0.3	0.4	0.7	1.2	3.0	6.1	21.7
2002.....	*	*	0.1	0.2	0.4	0.6	1.3	2.5	6.2	23.5
2001.....	*	*	0.2	0.3	0.4	0.6	1.1	3.0	5.5	23.9
2000.....	*	*	0.2	0.3	0.5	0.4	1.2	2.6	5.8	21.6
1999.....	*	0.1	0.2	0.2	0.3	0.6	1.3	2.5	6.1	23.0
White, male										
2006 4/.....	*	*	*	*	*	*	1.0	1.2	3.3	8.4
2005 4/.....	*	*	*	*	0.5	*	*	1.3	3.3	11.1
2004 4/.....	*	*	*	*	*	*	0.7	1.4	3.3	11.6
2003 4/.....	*	*	*	*	*	0.5	0.6	1.9	2.9	10.7
2002.....	*	*	*	*	*	*	0.8	1.6	3.2	13.1
2001.....	*	*	*	*	*	*	0.7	2.1	2.6	14.5
2000.....	*	*	*	*	*	*	0.8	1.6	2.8	12.0
1999.....	*	*	*	*	*	*	0.9	1.5	4.1	13.2
White, female										
2006 4/.....	*	*	*	0.3	0.4	1.0	1.3	2.7	6.4	23.8
2005 4/.....	0.2	*	0.2	0.3	0.4	0.9	1.2	2.9	6.6	27.2
2004 4/.....	*	*	*	*	0.5	0.5	1.1	2.5	6.5	23.5
2003 4/.....	*	*	0.2	0.4	0.6	0.8	1.6	3.8	8.0	26.5
2002.....	*	*	*	0.3	0.6	0.7	1.7	3.2	8.1	28.0
2001.....	*	*	*	0.4	0.4	0.8	1.4	3.6	7.3	27.8
2000.....	*	*	0.3	0.4	0.6	0.5	1.5	3.3	7.7	25.5
1999.....	*	*	*	0.4	*	0.9	1.6	3.1	7.2	27.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
All races 3/, both sexes										
2006.....	6.2	9.6	17.1	28.7	46.4	68.3	98.3	141.9	199.3	285.2
2005.....	5.6	9.9	17.3	29.7	47.3	71.7	104.9	151.4	211.2	312.1
2004.....	5.7	10.0	17.2	29.4	47.2	71.3	105.7	149.8	213.1	307.0
2003.....	5.8	10.3	17.9	30.4	48.9	75.0	108.8	155.3	216.5	317.5
2002.....	6.0	9.9	18.0	30.1	47.5	75.7	108.8	157.7	217.9	320.6
2001.....	5.4	9.9	17.8	29.5	48.4	75.5	108.6	155.5	218.7	321.8
2000.....	5.5	9.5	17.3	29.6	48.1	74.0	108.6	156.5	213.9	319.7
1999.....	5.5	9.9	16.5	29.5	48.9	75.6	109.2	153.5	214.8	317.2
All races 3/, male										
2006.....	7.6	12.3	21.3	34.5	56.8	80.7	115.9	167.5	232.2	321.0
2005.....	7.2	12.7	21.0	36.3	56.6	82.4	121.2	174.0	244.1	344.4
2004.....	7.1	12.4	21.2	34.5	56.6	82.7	121.2	172.2	246.9	333.9
2003.....	7.4	12.6	21.3	35.4	57.8	86.8	124.4	179.0	249.7	340.7
2002.....	7.7	11.9	21.5	36.0	55.5	84.3	121.1	175.9	243.5	348.5
2001.....	6.6	12.1	21.3	34.4	54.4	84.7	121.0	171.5	242.3	343.1
2000.....	6.8	11.4	19.7	33.4	53.9	80.8	119.7	173.7	242.5	345.1
1999.....	6.4	12.3	19.4	33.5	53.9	82.5	119.4	172.3	241.3	345.7
All races 3/, female										
2006.....	4.7	6.9	13.2	23.3	36.9	57.5	84.0	123.1	178.7	268.4
2005.....	4.1	7.2	13.8	23.4	38.9	62.3	91.6	134.9	190.8	297.2
2004.....	4.3	7.7	13.4	24.6	38.6	61.4	93.2	133.6	192.2	294.9
2003.....	4.2	8.1	14.7	25.7	40.9	64.8	96.2	138.4	196.2	307.2
2002.....	4.4	7.9	14.7	24.5	40.2	68.2	99.0	144.9	202.3	308.4
2001.....	4.2	7.9	14.5	24.9	42.9	67.6	98.8	144.3	204.5	312.9
2000.....	4.3	7.8	14.9	26.1	42.8	68.1	99.9	144.6	197.0	309.4
1999.....	4.6	7.5	13.8	25.8	44.5	69.7	101.2	140.4	199.3	305.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
White, both sexes										
2006 4/.....	5.3	8.4	14.7	25.0	40.6	60.2	89.0	130.6	185.8	270.4
2005 4/.....	4.9	8.6	15.0	25.9	41.8	63.3	94.6	139.9	198.7	296.3
2004 4/.....	5.0	8.9	14.6	25.5	41.3	63.6	94.8	137.4	199.5	289.4
2003 4/.....	5.1	8.8	15.2	26.4	42.7	66.5	98.1	143.1	202.7	300.8
2002.....	5.3	8.5	15.4	26.0	41.3	66.0	98.2	145.4	204.9	303.7
2001.....	4.8	8.6	15.3	25.3	42.3	66.8	98.0	143.0	205.9	306.0
2000.....	5.0	8.0	14.6	25.7	42.3	65.2	98.1	143.7	199.1	303.5
1999.....	4.7	8.3	13.8	24.5	41.9	66.9	97.6	141.2	200.8	298.9
White, male										
2006 4/.....	6.7	10.9	18.5	30.4	50.9	73.3	107.3	158.5	223.5	312.0
2005 4/.....	6.5	11.3	18.4	32.0	51.0	74.8	111.5	164.6	235.4	336.2
2004 4/.....	6.3	11.1	18.7	30.3	51.0	75.0	111.6	162.1	238.5	323.0
2003 4/.....	6.6	11.1	18.7	31.6	51.5	79.1	115.3	170.2	241.1	333.4
2002.....	6.9	10.3	18.8	32.0	49.5	75.9	112.9	167.9	235.5	344.3
2001.....	6.1	10.5	18.6	30.4	48.8	77.6	113.5	161.8	232.1	337.3
2000.....	6.3	9.7	17.3	29.5	48.7	73.8	111.6	164.0	234.1	337.4
1999.....	5.6	10.6	16.8	28.9	47.7	75.2	111.3	163.7	233.2	333.4
White, female										
2006 4/.....	3.9	5.8	10.8	19.9	31.0	48.5	73.9	109.8	162.0	250.8
2005 4/.....	3.3	5.9	11.6	20.0	33.2	53.1	80.7	121.7	175.7	278.0
2004 4/.....	3.6	6.7	10.6	20.8	32.4	53.4	81.1	119.4	175.3	274.3
2003 4/.....	3.5	6.5	11.8	21.4	34.5	55.4	84.0	123.5	179.1	286.3
2002.....	3.8	6.6	12.0	20.2	33.7	57.3	86.3	129.3	186.1	286.1
2001.....	3.4	6.7	12.0	20.4	36.3	57.4	85.4	129.7	190.1	293.0
2000.....	3.7	6.4	12.0	22.1	36.3	57.6	87.3	129.3	178.3	289.7
1999.....	3.9	6.1	11.0	20.4	36.6	59.6	86.7	125.4	181.7	285.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diabetes mellitus (E10-E14)										
Black, both sexes										
2006 4/.....	12.5	19.1	37.1	61.5	101.7	145.8	193.3	272.8	367.5	471.1
2005 4/.....	10.8	20.7	36.3	63.6	102.9	150.2	211.2	283.6	377.0	506.5
2004 4/.....	11.5	19.2	38.8	64.8	102.0	147.7	219.3	297.6	384.2	532.4
2003 4/.....	11.7	22.1	39.9	69.8	106.9	156.1	221.2	300.0	389.9	527.5
2002.....	12.0	21.4	40.7	67.8	105.5	161.6	221.6	300.3	392.7	537.8
2001.....	10.7	21.0	40.2	69.6	104.4	157.6	226.8	305.2	382.2	529.7
2000.....	10.1	21.9	40.7	66.3	102.4	159.2	223.7	316.5	404.9	524.9
1999.....	11.3	22.5	40.5	72.7	111.6	159.6	229.0	294.9	395.8	529.0
Black, male										
2006 4/.....	15.9	23.8	45.3	71.9	120.0	163.3	220.8	297.6	373.8	463.8
2005 4/.....	12.3	25.9	42.2	78.0	118.2	161.8	236.9	310.7	391.9	464.8
2004 4/.....	13.9	24.2	44.1	74.6	115.7	168.6	238.3	309.8	389.5	505.7
2003 4/.....	14.4	25.4	44.0	77.3	121.7	170.6	232.6	306.3	387.6	453.5
2002.....	14.9	25.4	46.4	76.5	118.8	165.3	223.3	295.8	386.7	417.0
2001.....	11.7	25.5	45.6	76.7	112.5	162.1	221.5	308.1	400.5	450.5
2000.....	11.8	25.5	43.5	72.1	108.5	154.3	224.3	311.4	382.7	452.1
1999.....	13.2	27.5	44.1	78.2	113.3	163.8	216.1	282.6	374.1	506.2
Black, female										
2006 4/.....	9.6	15.0	30.2	52.9	87.4	132.9	174.3	257.9	364.2	474.1
2005 4/.....	9.5	16.2	31.3	51.7	91.0	141.5	193.7	267.5	369.1	523.3
2004 4/.....	9.4	14.9	34.4	56.8	91.3	132.2	206.5	290.3	381.4	543.0
2003 4/.....	9.3	19.2	36.4	63.6	95.4	145.3	213.6	296.3	391.1	556.2
2002.....	9.4	17.9	35.8	60.6	95.1	158.9	220.4	303.0	395.9	584.1
2001.....	9.8	17.0	35.6	63.9	98.1	154.3	230.3	303.4	372.8	559.1
2000.....	8.7	18.7	38.4	61.5	97.6	162.8	223.3	319.6	416.1	552.0
1999.....	9.7	18.1	37.4	68.3	110.3	156.4	237.6	302.4	406.6	537.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of adrenal glands (E24-E27)										
All races 3/, both sexes										
2006.....	*	*	*	0.1	0.2	0.2	0.2	0.4	0.8	0.9
2005.....	*	*	*	0.1	0.2	*	*	0.5	0.7	1.0
2004.....	*	*	*	*	0.2	*	0.3	0.4	0.7	1.2
2003.....	*	*	*	0.1	0.2	0.2	0.3	0.6	0.8	1.2
2002.....	*	*	*	*	0.2	0.2	0.3	0.5	0.9	1.1
2001.....	*	*	*	*	*	*	0.5	0.7	1.0	1.8
2000.....	*	*	*	0.2	*	0.2	0.2	0.5	1.0	1.0
1999.....	*	*	0.1	*	*	0.3	0.3	0.5	0.4	1.0
All races 3/, male										
2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	1.3
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	0.6	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	0.7	1.2	2.1
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2006.....	*	*	*	*	*	*	*	*	1.0	0.9
2005.....	*	*	*	*	*	*	*	*	0.8	0.9
2004.....	*	*	*	*	*	*	*	*	0.8	1.1
2003.....	*	*	*	*	*	*	*	0.5	0.9	1.3
2002.....	*	*	*	*	*	*	0.4	0.6	0.9	1.1
2001.....	*	*	*	*	*	*	0.5	0.7	0.8	1.7
2000.....	*	*	*	*	*	*	*	0.5	1.0	1.1
1999.....	*	*	*	*	*	*	0.4	0.6	*	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of adrenal glands (E24-E27)										
White, both sexes										
2006 4/.....	*	*	*	*	*	0.2	*	0.4	0.9	1.0
2005 4/.....	*	*	*	*	0.2	*	*	0.5	0.7	1.0
2004 4/.....	*	*	*	*	*	*	*	0.4	0.8	1.2
2003 4/.....	*	*	*	*	*	*	0.3	0.6	0.8	1.2
2002.....	*	*	*	*	*	*	0.3	0.5	0.9	1.1
2001.....	*	*	*	*	*	*	0.5	0.7	1.0	1.8
2000.....	*	*	*	0.2	*	*	*	0.5	1.0	1.1
1999.....	*	*	0.1	*	*	0.3	0.3	0.6	0.5	1.0
White, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	1.3	2.1
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, female										
2006 4/.....	*	*	*	*	*	*	*	*	1.1	1.0
2005 4/.....	*	*	*	*	*	*	*	*	0.8	0.9
2004 4/.....	*	*	*	*	*	*	*	*	0.8	1.2
2003 4/.....	*	*	*	*	*	*	*	*	1.0	1.3
2002.....	*	*	*	*	*	*	*	0.6	0.9	1.1
2001.....	*	*	*	*	*	*	0.6	0.8	0.8	1.7
2000.....	*	*	*	*	*	*	*	*	1.1	1.1
1999.....	*	*	*	*	*	*	*	0.6	*	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other endocrine diseases (E15-E21,E28-E31,E34)										
All races 3/, both sexes										
2006.....	*	0.1	0.2	0.2	0.3	0.3	0.6	0.8	1.4	1.8
2005.....	*	0.1	0.1	0.1	0.3	0.2	0.4	0.8	1.2	2.2
2004.....	*	0.1	0.1	0.2	0.2	0.4	0.4	0.8	1.2	2.1
2003.....	*	*	*	0.2	0.2	0.3	0.5	0.6	1.2	2.1
2002.....	*	0.1	0.2	0.2	0.2	0.3	0.5	0.8	1.3	2.2
2001.....	*	*	0.1	0.1	0.3	0.2	0.5	0.7	1.0	2.5
2000.....	*	0.1	*	*	*	0.2	0.4	0.9	0.9	2.2
1999.....	*	*	0.2	0.2	0.2	0.2	0.6	0.7	1.2	1.9
All races 3/, male										
2006.....	*	*	*	*	0.3	*	0.5	0.7	1.7	1.8
2005.....	*	*	*	*	*	*	*	0.9	1.1	1.6
2004.....	*	*	*	*	*	0.5	*	1.0	1.3	1.8
2003.....	*	*	*	*	*	*	0.6	0.7	1.2	1.7
2002.....	*	*	*	*	*	*	*	0.7	1.1	1.8
2001.....	*	*	*	*	*	*	0.6	0.8	*	2.2
2000.....	*	*	*	*	*	*	*	0.9	*	1.7
1999.....	*	*	*	*	*	*	0.7	0.8	1.4	*
All races 3/, female										
2006.....	*	*	*	*	0.3	*	0.6	0.9	1.2	1.8
2005.....	*	*	*	*	*	*	*	0.8	1.3	2.5
2004.....	*	*	*	*	*	*	*	0.7	1.1	2.2
2003.....	*	*	*	0.3	*	*	0.4	0.6	1.2	2.2
2002.....	*	*	*	*	*	*	0.6	0.8	1.3	2.3
2001.....	*	*	*	*	*	*	*	0.7	1.2	2.6
2000.....	*	*	*	*	*	*	0.4	0.9	0.9	2.4
1999.....	*	*	*	*	*	*	0.5	0.6	1.1	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other endocrine diseases (E15-E21,E28-E31,E34)

White, both sexes

2006 4/.....	*	*	*	*	0.3	0.3	0.5	0.7	1.2	1.6
2005 4/.....	*	*	0.1	*	0.2	*	0.3	0.8	1.0	1.9
2004 4/.....	*	*	*	0.1	0.2	0.3	0.3	0.8	1.1	1.9
2003 4/.....	*	*	*	*	0.2	*	0.4	0.6	1.0	2.0
2002.....	*	*	0.1	*	0.2	0.3	0.4	0.7	1.2	2.1
2001.....	*	*	*	*	0.2	*	0.3	0.6	0.9	2.4
2000.....	*	*	*	*	*	*	0.3	0.7	0.7	2.0
1999.....	*	*	*	*	*	*	0.5	0.6	1.1	1.8

White, male

2006 4/.....	*	*	*	*	*	*	*	*	1.3	1.8
2005 4/.....	*	*	*	*	*	*	*	0.9	*	*
2004 4/.....	*	*	*	*	*	*	*	0.9	1.3	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	2.0
2001.....	*	*	*	*	*	*	*	*	*	2.1
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	0.8	1.4	*

White, female

2006 4/.....	*	*	*	*	*	*	0.5	0.8	1.2	1.5
2005 4/.....	*	*	*	*	*	*	*	0.7	1.0	2.3
2004 4/.....	*	*	*	*	*	*	*	0.6	1.0	2.1
2003 4/.....	*	*	*	*	*	*	*	0.6	1.0	2.2
2002.....	*	*	*	*	*	*	0.5	0.8	1.3	2.2
2001.....	*	*	*	*	*	*	*	0.6	1.0	2.5
2000.....	*	*	*	*	*	*	*	0.7	0.7	2.2
1999.....	*	*	*	*	*	*	*	0.5	0.9	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional diseases (E40-E68)										
All races 3/, both sexes										
2006.....	1.9	2.8	3.4	4.3	4.5	5.2	6.1	7.1	10.9	24.4
2005.....	1.9	2.5	3.5	4.1	5.1	5.3	6.6	7.9	12.8	32.6
2004.....	1.7	2.5	3.4	3.7	4.3	4.8	5.8	7.7	12.8	32.3
2003.....	1.7	2.5	3.1	3.7	4.1	5.0	5.8	8.3	12.9	37.5
2002.....	1.9	2.2	3.0	3.8	3.9	4.7	5.0	8.7	15.0	42.9
2001.....	1.4	2.1	2.8	3.1	3.5	3.9	5.4	8.1	14.5	44.5
2000.....	1.6	2.0	2.7	2.9	3.6	4.5	5.4	9.1	17.9	51.7
1999.....	1.5	1.7	2.3	2.8	2.9	3.8	5.6	8.6	17.9	55.3
All races 3/, male										
2006.....	2.1	3.1	3.4	4.1	4.2	4.9	5.7	6.3	10.5	19.4
2005.....	2.2	2.9	3.5	4.3	5.0	4.6	5.8	6.9	11.3	29.3
2004.....	1.8	2.5	3.5	3.8	4.0	4.7	5.0	7.6	12.3	28.8
2003.....	1.8	2.6	3.1	3.9	3.9	5.1	5.2	7.6	12.8	31.5
2002.....	2.2	2.2	3.1	3.7	3.6	4.3	4.9	8.2	13.5	37.8
2001.....	1.5	2.2	2.6	2.9	3.4	3.4	5.3	6.9	14.4	35.8
2000.....	1.7	2.2	2.8	3.3	3.2	4.0	5.3	8.5	15.8	44.6
1999.....	1.8	1.8	2.3	3.1	3.0	3.5	5.8	8.4	17.1	48.7
All races 3/, female										
2006.....	1.7	2.5	3.3	4.5	4.8	5.4	6.5	7.7	11.1	26.7
2005.....	1.5	2.2	3.5	3.9	5.1	5.9	7.3	8.7	13.7	34.2
2004.....	1.6	2.6	3.2	3.6	4.6	4.9	6.4	7.8	13.1	33.9
2003.....	1.7	2.5	3.1	3.6	4.4	5.0	6.3	8.8	12.9	40.1
2002.....	1.6	2.2	2.9	3.9	4.2	5.1	5.2	9.0	15.9	45.1
2001.....	1.3	2.0	3.0	3.3	3.7	4.3	5.6	9.0	14.6	48.1
2000.....	1.6	1.8	2.7	2.6	3.9	4.9	5.5	9.6	19.2	54.6
1999.....	1.3	1.6	2.4	2.4	2.9	4.0	5.5	8.8	18.4	58.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional diseases (E40-E68)										
White, both sexes										
2006 4/.....	1.7	2.6	3.2	4.1	4.4	5.1	6.1	7.1	10.1	23.7
2005 4/.....	1.6	2.4	3.3	4.0	5.0	5.0	6.4	7.5	12.2	31.9
2004 4/.....	1.5	2.3	3.2	3.6	4.1	4.5	5.5	7.2	12.4	31.0
2003 4/.....	1.6	2.4	3.0	3.7	4.0	5.1	5.5	8.1	12.3	36.1
2002.....	1.8	2.1	2.9	3.6	3.7	4.4	4.8	8.2	14.4	41.6
2001.....	1.2	1.9	2.6	3.0	3.4	3.6	5.0	7.6	13.6	43.8
2000.....	1.5	1.8	2.6	2.7	3.5	4.3	5.0	8.5	17.4	50.6
1999.....	1.4	1.5	2.2	2.6	2.8	3.6	4.9	8.0	16.8	54.2
White, male										
2006 4/.....	1.9	2.9	3.4	4.1	4.2	5.1	5.6	6.0	9.6	18.8
2005 4/.....	2.0	2.9	3.7	4.3	5.1	4.5	5.8	6.5	10.6	28.5
2004 4/.....	1.8	2.4	3.5	3.9	3.9	4.3	5.1	7.1	11.9	27.8
2003 4/.....	1.7	2.5	3.1	4.0	3.9	5.3	4.8	7.1	12.1	30.5
2002.....	2.1	2.3	3.2	3.6	3.5	4.1	4.6	8.0	12.5	37.1
2001.....	1.4	2.0	2.5	2.9	3.2	3.1	4.9	6.6	13.5	34.5
2000.....	1.6	2.2	2.8	3.1	3.0	3.8	4.9	7.9	15.3	43.7
1999.....	1.8	1.7	2.3	2.9	2.8	3.3	5.0	7.6	15.8	46.9
White, female										
2006 4/.....	1.6	2.3	3.1	4.2	4.6	5.2	6.5	7.8	10.5	26.0
2005 4/.....	1.2	1.8	3.0	3.7	4.9	5.5	6.9	8.2	13.2	33.4
2004 4/.....	1.3	2.2	2.9	3.4	4.3	4.6	5.9	7.4	12.7	32.5
2003 4/.....	1.5	2.3	3.0	3.4	4.1	5.0	6.1	8.8	12.3	38.6
2002.....	1.4	2.0	2.6	3.5	3.8	4.6	5.0	8.3	15.5	43.5
2001.....	1.1	1.7	2.7	3.1	3.6	4.1	5.1	8.3	13.8	47.7
2000.....	1.4	1.4	2.4	2.3	3.9	4.7	5.2	9.0	18.7	53.4
1999.....	1.1	1.4	2.2	2.2	2.7	3.8	4.8	8.3	17.4	57.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional diseases (E40-E68)										
Black, both sexes										
2006 4/.....	3.2	4.9	5.6	7.3	6.6	7.4	8.8	9.7	21.3	34.8
2005 4/.....	3.7	4.7	5.7	6.6	7.7	9.3	10.9	13.7	21.6	45.7
2004 4/.....	3.2	4.7	5.7	6.0	7.6	9.3	10.3	13.9	18.6	50.8
2003 4/.....	3.0	4.1	4.8	5.6	6.8	5.6	9.9	13.4	22.8	58.1
2002.....	3.4	3.7	5.2	7.5	7.0	8.7	8.8	16.3	24.9	62.1
2001.....	2.9	4.3	5.5	5.4	5.7	6.8	10.6	15.5	26.5	56.2
2000.....	3.0	3.8	4.9	6.2	5.3	7.5	10.4	16.9	26.5	68.5
1999.....	2.6	3.2	4.0	5.3	5.5	6.6	13.4	16.2	34.4	72.4
Black, male										
2006 4/.....	3.6	5.0	4.6	5.7	4.8	5.9	8.6	10.7	23.3	30.4
2005 4/.....	3.9	4.3	3.6	5.8	6.5	7.1	7.6	10.7	21.1	40.9
2004 4/.....	2.8	3.3	4.8	5.0	6.0	10.4	*	15.5	17.9	45.7
2003 4/.....	2.8	3.7	4.4	4.5	5.4	4.9	10.5	16.4	25.1	48.0
2002.....	3.7	2.6	3.8	6.7	5.7	7.0	8.7	12.4	29.8	48.2
2001.....	2.9	3.9	4.4	4.2	6.1	6.4	10.4	13.5	26.5	57.9
2000.....	2.8	2.8	4.0	6.4	5.6	6.5	12.0	18.0	25.3	65.7
1999.....	2.4	2.6	3.1	6.1	5.3	6.2	14.9	18.0	35.8	75.0
Black, female										
2006 4/.....	2.8	4.8	6.4	8.5	8.0	8.5	8.9	9.1	20.3	36.6
2005 4/.....	3.5	5.1	7.5	7.3	8.6	10.9	13.2	15.5	21.9	47.6
2004 4/.....	3.5	5.9	6.4	6.7	8.9	8.5	13.4	12.9	18.9	52.8
2003 4/.....	3.2	4.5	5.1	6.5	8.0	6.0	9.5	11.7	21.7	62.0
2002.....	3.2	4.6	6.4	8.2	8.1	10.0	9.0	18.7	22.3	67.4
2001.....	2.8	4.5	6.4	6.3	5.4	7.1	10.8	16.6	26.6	55.6
2000.....	3.1	4.7	5.6	6.1	5.1	8.3	9.3	16.2	27.2	69.5
1999.....	2.9	3.7	4.8	4.6	5.7	6.9	12.4	15.2	33.7	71.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
All races 3/, both sexes										
2006.....	0.1	0.2	0.3	0.4	0.7	1.0	1.8	3.7	7.0	20.9
2005.....	0.1	0.2	0.3	0.4	0.6	1.2	2.5	4.6	9.3	28.9
2004.....	0.1	0.2	0.4	0.4	0.7	1.2	2.0	4.4	9.6	28.3
2003.....	0.1	0.2	0.3	0.5	0.7	1.2	2.4	5.5	10.0	33.4
2002.....	0.1	0.2	0.3	0.5	0.7	1.4	2.5	5.6	12.2	38.4
2001.....	0.1	0.2	0.3	0.4	0.8	1.3	2.6	5.5	12.0	40.2
2000.....	0.1	0.2	0.3	0.5	0.9	1.9	2.8	6.4	14.6	46.3
1999.....	0.2	0.2	0.3	0.5	0.8	1.5	3.0	6.0	14.7	50.0
All races 3/, male										
2006.....	*	*	0.2	0.4	0.6	1.0	1.9	3.7	7.4	17.5
2005.....	*	0.2	0.3	0.5	0.6	1.2	2.4	4.4	8.8	26.3
2004.....	*	*	0.3	0.5	0.7	1.3	2.2	4.8	9.2	25.3
2003.....	*	*	0.3	0.5	0.7	1.3	2.5	5.4	10.4	28.2
2002.....	*	*	0.3	0.5	0.8	1.4	2.8	5.7	11.4	35.1
2001.....	*	0.2	0.3	0.5	0.8	1.3	2.6	5.1	12.6	33.6
2000.....	*	0.2	*	0.6	0.9	1.8	3.3	6.4	13.1	40.7
1999.....	*	0.2	0.3	0.6	1.1	1.5	3.6	6.2	14.8	44.4
All races 3/, female										
2006.....	*	0.2	0.3	0.5	0.7	1.0	1.8	3.8	6.8	22.4
2005.....	*	0.2	0.4	0.3	0.5	1.2	2.6	4.7	9.7	30.2
2004.....	*	0.2	0.4	0.3	0.7	1.1	1.9	4.2	9.9	29.6
2003.....	*	*	0.3	0.5	0.8	1.1	2.2	5.7	9.7	35.6
2002.....	*	*	0.3	0.4	0.6	1.4	2.2	5.6	12.6	39.8
2001.....	*	0.2	0.3	0.4	0.7	1.3	2.5	5.7	11.6	43.0
2000.....	*	0.2	0.3	0.3	0.9	1.9	2.4	6.5	15.5	48.6
1999.....	0.2	0.2	0.3	0.4	0.6	1.4	2.5	5.8	14.7	52.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
White, both sexes										
2006 4/.....	0.1	0.2	0.2	0.4	0.7	0.9	1.7	3.7	6.6	20.3
2005 4/.....	*	0.2	0.3	0.4	0.5	1.0	2.4	4.2	8.8	28.2
2004 4/.....	*	0.1	0.3	0.4	0.6	1.0	1.9	4.1	9.3	26.9
2003 4/.....	*	0.1	0.2	0.4	0.6	1.1	2.2	5.4	9.3	32.1
2002.....	0.1	0.2	0.3	0.4	0.6	1.3	2.4	5.4	11.7	37.0
2001.....	*	0.2	0.2	0.4	0.7	1.1	2.4	5.1	11.2	39.5
2000.....	*	0.2	0.2	0.4	0.8	1.8	2.4	6.1	14.2	45.2
1999.....	0.2	0.2	0.3	0.4	0.7	1.3	2.5	5.5	13.8	48.9
White, male										
2006 4/.....	*	*	*	0.3	0.6	0.8	1.8	3.4	6.7	16.9
2005 4/.....	*	*	0.3	0.5	0.5	1.0	2.3	4.0	8.1	25.7
2004 4/.....	*	*	0.3	0.4	0.6	1.0	2.0	4.3	8.7	24.1
2003 4/.....	*	*	*	0.4	0.5	1.2	2.2	4.9	9.5	27.2
2002.....	*	*	*	0.4	0.6	1.3	2.7	5.5	10.5	34.3
2001.....	*	*	*	0.4	0.7	1.0	2.3	4.7	11.7	32.2
2000.....	*	*	*	0.5	0.7	1.6	2.8	5.9	12.5	39.7
1999.....	*	*	0.3	0.4	0.9	1.2	2.8	5.6	13.4	42.2
White, female										
2006 4/.....	*	*	0.2	0.5	0.7	0.9	1.7	3.9	6.5	21.8
2005 4/.....	*	*	0.3	0.4	0.5	1.0	2.4	4.4	9.2	29.4
2004 4/.....	*	*	0.3	0.4	0.6	1.1	1.7	3.9	9.7	28.2
2003 4/.....	*	*	*	0.5	0.7	1.0	2.2	5.7	9.3	34.2
2002.....	*	*	0.3	0.4	0.6	1.2	2.1	5.3	12.4	38.2
2001.....	*	*	0.3	0.4	0.8	1.2	2.5	5.4	10.9	42.6
2000.....	*	*	0.3	*	0.9	1.9	2.1	6.2	15.1	47.4
1999.....	*	*	0.3	0.4	0.5	1.3	2.3	5.5	14.0	51.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Malnutrition (E40-E46)										
Black, both sexes										
2006 4/.....	*	*	*	*	*	2.6	3.4	4.4	13.0	29.3
2005 4/.....	*	*	*	*	*	2.7	4.2	8.7	16.7	40.4
2004 4/.....	*	*	0.9	*	*	2.5	4.2	8.1	13.6	46.0
2003 4/.....	*	*	1.0	*	1.8	*	4.4	8.8	19.3	53.4
2002.....	*	*	*	1.5	1.9	3.0	4.4	10.1	19.9	57.6
2001.....	*	*	*	*	*	2.8	4.0	10.1	22.2	51.6
2000.....	*	*	*	*	1.9	3.1	6.5	11.2	20.9	63.8
1999.....	*	*	*	1.6	2.0	3.2	8.0	11.1	28.9	66.0
Black, male										
2006 4/.....	*	*	*	*	*	*	*	*	17.1	28.6
2005 4/.....	*	*	*	*	*	*	*	*	17.6	34.4
2004 4/.....	*	*	*	*	*	*	*	10.5	14.3	43.7
2003 4/.....	*	*	*	*	*	*	6.9	13.2	23.6	45.9
2002.....	*	*	*	*	*	*	*	9.2	25.2	46.1
2001.....	*	*	*	*	*	*	*	11.2	24.1	55.6
2000.....	*	*	*	*	*	*	9.7	14.2	22.8	61.1
1999.....	*	*	*	*	*	*	12.2	13.3	33.1	73.8
Black, female										
2006 4/.....	*	*	*	*	*	*	*	*	10.9	29.6
2005 4/.....	*	*	*	*	*	*	4.8	9.1	16.3	42.8
2004 4/.....	*	*	*	*	*	*	*	6.7	13.2	46.9
2003 4/.....	*	*	*	*	*	*	*	6.2	17.0	56.3
2002.....	*	*	*	*	*	*	*	10.7	17.1	62.0
2001.....	*	*	*	*	*	*	*	9.4	21.3	50.1
2000.....	*	*	*	*	*	*	4.4	9.4	20.0	64.8
1999.....	*	*	*	*	*	*	5.2	9.7	26.8	63.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional marasmus (E41)										
All races 3/, both sexes										
2006.....	*	*	*	*	*	*	*	*	*	0.4
2005.....	*	*	*	*	*	*	*	*	0.4	1.1
2004.....	*	*	*	*	*	*	*	*	0.5	1.3
2003.....	*	*	*	*	*	*	*	0.3	0.6	1.6
2002.....	*	*	*	*	*	*	*	*	0.6	2.4
2001.....	*	*	*	*	*	*	*	*	0.6	2.5
2000.....	*	*	*	*	*	*	*	*	0.8	2.7
1999.....	*	*	*	*	*	*	*	0.3	0.8	2.5
All races 3/, male										
2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	2.2
2000.....	*	*	*	*	*	*	*	*	*	2.9
1999.....	*	*	*	*	*	*	*	*	*	3.0
All races 3/, female										
2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	1.3
2004.....	*	*	*	*	*	*	*	*	*	1.4
2003.....	*	*	*	*	*	*	*	*	0.6	1.7
2002.....	*	*	*	*	*	*	*	*	0.6	2.7
2001.....	*	*	*	*	*	*	*	*	*	2.6
2000.....	*	*	*	*	*	*	*	*	0.9	2.7
1999.....	*	*	*	*	*	*	*	*	0.6	2.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nutritional marasmus (E41)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	1.1
2004 4/.....	*	*	*	*	*	*	*	*	0.5	1.2
2003 4/.....	*	*	*	*	*	*	*	*	0.5	1.6
2002.....	*	*	*	*	*	*	*	*	0.6	2.4
2001.....	*	*	*	*	*	*	*	*	0.6	2.5
2000.....	*	*	*	*	*	*	*	*	0.9	2.7
1999.....	*	*	*	*	*	*	*	*	0.8	2.4
White, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	2.1
2000.....	*	*	*	*	*	*	*	*	*	2.7
1999.....	*	*	*	*	*	*	*	*	*	2.9
White, female										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	1.3
2004 4/.....	*	*	*	*	*	*	*	*	*	1.4
2003 4/.....	*	*	*	*	*	*	*	*	*	1.8
2002.....	*	*	*	*	*	*	*	*	*	2.7
2001.....	*	*	*	*	*	*	*	*	*	2.7
2000.....	*	*	*	*	*	*	*	*	1.0	2.7
1999.....	*	*	*	*	*	*	*	*	0.7	2.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified malnutrition (E40,E42-E46)

All races 3/, both sexes

2006.....	0.1	0.2	0.3	0.4	0.7	1.0	1.8	3.7	6.8	20.5
2005.....	0.1	0.2	0.3	0.4	0.6	1.1	2.4	4.4	8.9	27.8
2004.....	0.1	0.2	0.4	0.4	0.7	1.1	1.9	4.3	9.2	27.0
2003.....	0.1	0.2	0.3	0.5	0.7	1.1	2.3	5.3	9.4	31.8
2002.....	0.1	0.2	0.3	0.5	0.7	1.4	2.4	5.4	11.6	36.0
2001.....	0.1	0.2	0.3	0.4	0.7	1.2	2.4	5.2	11.4	37.7
2000.....	0.1	0.2	0.2	0.4	0.9	1.8	2.7	6.2	13.8	43.6
1999.....	0.2	0.2	0.3	0.5	0.8	1.4	2.8	5.7	14.0	47.5

All races 3/, male

2006.....	*	*	0.2	0.4	0.6	1.0	1.9	3.6	7.1	17.3
2005.....	*	0.2	0.3	0.5	0.6	1.2	2.3	4.3	8.4	25.8
2004.....	*	*	0.3	0.5	0.7	1.2	2.1	4.6	8.7	24.3
2003.....	*	*	0.3	0.5	0.6	1.2	2.5	5.1	9.9	27.1
2002.....	*	*	0.2	0.5	0.8	1.4	2.8	5.6	10.8	33.5
2001.....	*	0.2	0.3	0.5	0.7	1.2	2.5	4.8	11.9	31.3
2000.....	*	0.2	*	0.6	0.9	1.7	3.2	6.1	12.4	37.8
1999.....	*	0.2	0.3	0.6	1.0	1.5	3.3	6.0	13.8	41.4

All races 3/, female

2006.....	*	0.2	0.3	0.5	0.7	1.0	1.7	3.7	6.6	21.9
2005.....	*	0.2	0.4	0.3	0.5	1.1	2.4	4.5	9.3	28.8
2004.....	*	*	0.4	0.3	0.7	1.1	1.8	4.1	9.4	28.2
2003.....	*	*	0.3	0.5	0.7	1.1	2.1	5.4	9.1	33.9
2002.....	*	*	0.3	0.4	0.6	1.3	2.1	5.3	12.0	37.1
2001.....	*	0.2	0.3	0.4	0.7	1.2	2.4	5.5	11.1	40.4
2000.....	*	0.2	0.3	0.3	0.8	1.9	2.3	6.3	14.6	45.9
1999.....	0.2	*	0.3	0.4	0.6	1.4	2.4	5.5	14.1	50.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified malnutrition (E40,E42-E46)										
White, both sexes										
2006 4/.....	0.1	0.1	0.2	0.4	0.7	0.9	1.7	3.6	6.3	19.9
2005 4/.....	*	0.2	0.3	0.4	0.5	1.0	2.2	4.0	8.4	27.1
2004 4/.....	*	0.1	0.3	0.4	0.6	1.0	1.8	3.9	8.8	25.7
2003 4/.....	*	0.1	0.2	0.4	0.6	1.1	2.1	5.1	8.8	30.5
2002.....	0.1	0.2	0.2	0.4	0.6	1.2	2.3	5.2	11.1	34.7
2001.....	*	0.2	0.2	0.4	0.7	1.1	2.3	4.8	10.6	37.0
2000.....	*	0.2	0.2	0.4	0.7	1.7	2.3	5.9	13.3	42.5
1999.....	0.2	0.2	0.3	0.3	0.7	1.2	2.3	5.4	12.9	46.5
White, male										
2006 4/.....	*	*	*	0.3	0.6	0.8	1.7	3.3	6.4	16.6
2005 4/.....	*	*	0.3	0.4	0.5	1.0	2.1	3.9	7.8	25.1
2004 4/.....	*	*	0.3	0.4	0.5	0.9	1.9	4.2	8.2	23.2
2003 4/.....	*	*	*	0.3	0.5	1.1	2.1	4.6	9.0	26.0
2002.....	*	*	*	0.4	0.6	1.3	2.6	5.4	10.0	32.5
2001.....	*	*	*	0.4	0.6	1.0	2.3	4.4	11.0	30.1
2000.....	*	*	*	0.4	0.7	1.6	2.6	5.7	11.7	37.0
1999.....	*	*	0.3	*	0.9	1.2	2.6	5.4	12.5	39.4
White, female										
2006 4/.....	*	*	0.2	0.5	0.7	0.9	1.7	3.9	6.3	21.4
2005 4/.....	*	*	0.3	0.3	0.5	1.0	2.3	4.1	8.8	28.1
2004 4/.....	*	*	0.3	0.3	0.6	1.1	1.7	3.8	9.2	26.9
2003 4/.....	*	*	*	0.5	0.7	1.0	2.1	5.5	8.7	32.4
2002.....	*	*	0.3	0.4	0.6	1.2	2.0	5.0	11.8	35.6
2001.....	*	*	0.3	0.3	0.8	1.2	2.4	5.1	10.4	39.9
2000.....	*	*	*	*	0.8	1.8	2.0	6.0	14.2	44.7
1999.....	*	*	0.3	0.3	0.5	1.3	2.1	5.3	13.2	49.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified malnutrition (E40,E42-E46)

Black, both sexes

2006 4/.....	*	*	*	*	*	2.6	3.4	4.2	12.8	29.3
2005 4/.....	*	*	*	*	*	2.6	4.1	8.5	16.5	39.0
2004 4/.....	*	*	0.9	*	*	2.4	3.9	8.1	13.6	44.8
2003 4/.....	*	*	0.9	*	*	*	4.2	8.3	18.0	52.5
2002.....	*	*	*	1.4	1.8	2.9	4.3	9.3	18.8	55.8
2001.....	*	*	*	*	*	2.7	3.9	10.1	20.9	49.1
2000.....	*	*	*	*	1.9	3.0	6.3	10.7	20.9	59.7
1999.....	*	*	*	1.5	1.9	3.1	7.4	9.6	28.6	62.8

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	16.4	28.6
2005 4/.....	*	*	*	*	*	*	*	*	17.6	34.4
2004 4/.....	*	*	*	*	*	*	*	10.5	14.3	42.7
2003 4/.....	*	*	*	*	*	*	6.9	12.3	22.9	44.8
2002.....	*	*	*	*	*	*	*	*	22.9	45.0
2001.....	*	*	*	*	*	*	*	11.2	23.3	53.3
2000.....	*	*	*	*	*	*	9.4	12.8	22.8	55.4
1999.....	*	*	*	*	*	*	11.5	12.8	32.3	69.1

Black, female

2006 4/.....	*	*	*	*	*	*	*	*	10.9	29.6
2005 4/.....	*	*	*	*	*	*	4.5	8.8	16.0	40.9
2004 4/.....	*	*	*	*	*	*	*	6.7	13.2	45.7
2003 4/.....	*	*	*	*	*	*	*	6.0	15.5	55.5
2002.....	*	*	*	*	*	*	*	9.6	16.7	60.0
2001.....	*	*	*	*	*	*	*	9.4	19.6	47.6
2000.....	*	*	*	*	*	*	*	9.4	20.0	61.4
1999.....	*	*	*	*	*	*	4.7	7.7	26.8	60.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other nutritional deficiencies (E50-E64)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*	*	0.5	1.9
2005 4/.....	*	*	*	*	*	*	*	*	0.6	1.8
2004 4/.....	*	*	*	*	*	*	*	*	0.4	2.2
2003 4/.....	*	*	*	*	*	*	*	*	0.5	2.2
2002.....	*	*	*	*	*	*	*	0.5	0.6	3.1
2001.....	*	*	*	*	*	*	*	0.4	0.8	2.7
2000.....	*	*	*	*	*	*	0.3	0.3	1.2	3.7
1999.....	*	*	*	*	*	*	*	0.4	1.1	3.9

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	1.5
2004 4/.....	*	*	*	*	*	*	*	*	*	2.2
2003 4/.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	2.0
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	1.4	3.2
1999.....	*	*	*	*	*	*	*	*	1.3	4.3

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	2.2
2005 4/.....	*	*	*	*	*	*	*	*	0.7	1.9
2004 4/.....	*	*	*	*	*	*	*	*	*	2.2
2003 4/.....	*	*	*	*	*	*	*	*	*	2.1
2002.....	*	*	*	*	*	*	*	*	0.7	3.6
2001.....	*	*	*	*	*	*	*	*	0.9	3.1
2000.....	*	*	*	*	*	*	*	*	1.0	3.9
1999.....	*	*	*	*	*	*	*	*	1.0	3.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Obesity and other hyperalimentation (E65-E68)										
All races 3/, both sexes										
2006.....	1.7	2.6	3.1	3.9	3.7	4.1	4.2	3.2	3.3	1.7
2005.....	1.8	2.3	3.1	3.6	4.5	4.1	3.9	3.2	2.8	1.9
2004.....	1.6	2.3	3.0	3.3	3.6	3.6	3.7	3.1	2.7	2.0
2003.....	1.6	2.4	2.8	3.2	3.4	3.8	3.3	2.6	2.5	2.0
2002.....	1.8	2.0	2.7	3.3	3.1	3.1	2.4	2.6	2.2	1.4
2001.....	1.3	1.9	2.5	2.7	2.7	2.5	2.7	2.2	1.7	1.6
2000.....	1.5	1.7	2.4	2.4	2.6	2.5	2.4	2.3	2.2	1.7
1999.....	1.3	1.5	2.0	2.2	2.1	2.2	2.4	2.2	2.0	1.4
All races 3/, male										
2006.....	2.0	2.9	3.1	3.7	3.5	3.8	3.6	2.4	2.4	*
2005.....	2.1	2.7	3.2	3.7	4.3	3.4	3.2	2.4	2.0	1.4
2004.....	1.8	2.3	3.2	3.3	3.2	3.4	2.8	2.7	2.7	1.4
2003.....	1.7	2.4	2.8	3.4	3.2	3.7	2.5	1.9	1.9	*
2002.....	2.1	2.0	2.8	3.2	2.7	2.7	1.9	1.9	1.5	*
2001.....	1.4	1.9	2.3	2.4	2.5	1.9	2.4	1.3	1.2	*
2000.....	1.6	1.9	2.5	2.6	2.2	2.0	1.9	1.7	1.4	*
1999.....	1.6	1.5	1.9	2.5	1.9	1.9	1.9	1.6	*	*
All races 3/, female										
2006.....	1.5	2.3	3.0	4.0	4.0	4.4	4.7	3.7	3.9	2.2
2005.....	1.4	2.0	3.1	3.6	4.6	4.7	4.5	3.7	3.3	2.1
2004.....	1.4	2.3	2.8	3.3	3.9	3.8	4.5	3.5	2.8	2.2
2003.....	1.6	2.3	2.8	3.1	3.6	3.9	3.9	3.0	2.8	2.5
2002.....	1.5	2.1	2.6	3.4	3.5	3.5	2.8	3.1	2.6	1.6
2001.....	1.1	1.8	2.6	2.9	3.0	3.0	2.9	2.9	2.0	2.0
2000.....	1.4	1.6	2.3	2.2	3.0	2.9	2.8	2.7	2.6	2.0
1999.....	1.1	1.4	2.0	1.9	2.2	2.5	2.8	2.6	2.6	1.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Obesity and other hyperalimentation (E65-E68)										
White, both sexes										
2006 4/.....	1.6	2.5	3.0	3.7	3.7	4.2	4.3	3.1	3.0	1.6
2005 4/.....	1.6	2.2	3.0	3.5	4.5	4.0	3.8	3.1	2.8	1.8
2004 4/.....	1.4	2.1	2.9	3.2	3.5	3.4	3.6	3.0	2.6	1.9
2003 4/.....	1.5	2.3	2.8	3.2	3.3	3.9	3.2	2.5	2.4	1.9
2002.....	1.6	2.0	2.6	3.1	3.0	3.0	2.3	2.3	2.1	1.4
2001.....	1.1	1.7	2.3	2.6	2.6	2.4	2.4	2.1	1.7	1.6
2000.....	1.4	1.6	2.3	2.2	2.6	2.3	2.3	2.2	2.1	1.7
1999.....	1.2	1.4	1.9	2.2	2.0	2.2	2.2	2.0	2.0	1.4
White, male										
2006 4/.....	1.8	2.8	3.2	3.8	3.6	4.2	3.7	2.4	2.3	*
2005 4/.....	2.0	2.7	3.4	3.8	4.5	3.4	3.3	2.4	2.0	*
2004 4/.....	1.7	2.3	3.2	3.4	3.2	3.3	3.0	2.6	2.8	1.5
2003 4/.....	1.6	2.4	2.8	3.6	3.3	3.9	2.5	1.9	2.0	*
2002.....	2.0	2.1	2.9	3.2	2.9	2.7	1.8	1.9	1.5	*
2001.....	1.3	1.8	2.3	2.5	2.4	2.0	2.3	1.3	1.3	*
2000.....	1.6	1.9	2.6	2.5	2.2	2.0	1.9	1.6	1.4	*
1999.....	1.6	1.5	2.0	2.5	1.8	2.0	2.0	1.5	*	*
White, female										
2006 4/.....	1.4	2.2	2.8	3.7	3.8	4.2	4.7	3.7	3.5	2.0
2005 4/.....	1.2	1.6	2.6	3.3	4.4	4.4	4.3	3.6	3.3	2.0
2004 4/.....	1.2	2.0	2.5	3.0	3.7	3.5	4.1	3.3	2.6	2.0
2003 4/.....	1.4	2.1	2.7	2.9	3.3	4.0	3.8	2.9	2.7	2.3
2002.....	1.3	1.8	2.2	3.1	3.2	3.3	2.7	2.6	2.4	1.7
2001.....	0.9	1.5	2.3	2.8	2.8	2.8	2.4	2.6	1.9	2.0
2000.....	1.2	1.2	2.1	1.9	3.0	2.7	2.6	2.5	2.5	2.1
1999.....	0.9	1.2	1.8	1.8	2.1	2.4	2.4	2.4	2.5	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Obesity and other hyperalimentation (E65-E68)										
Black, both sexes										
2006 4/.....	3.0	4.5	4.9	6.3	5.3	4.8	5.4	4.7	7.8	*
2005 4/.....	3.5	4.3	5.0	5.8	6.1	6.6	6.6	5.0	*	*
2004 4/.....	3.0	4.3	4.8	5.2	5.9	6.6	6.2	5.8	*	*
2003 4/.....	2.9	3.7	3.8	4.5	5.0	3.6	5.3	4.4	*	*
2002.....	3.2	3.4	4.6	6.0	5.0	5.2	3.8	5.8	*	*
2001.....	2.5	3.8	4.4	4.2	4.6	3.8	6.2	4.7	*	*
2000.....	2.7	3.5	4.0	4.9	3.2	4.3	3.9	4.8	*	*
1999.....	2.3	2.5	3.3	3.3	3.5	3.4	5.1	3.9	*	*
Black, male										
2006 4/.....	3.4	4.7	3.8	4.2	3.5	*	*	*	*	*
2005 4/.....	3.8	3.8	2.8	4.4	4.4	*	*	*	*	*
2004 4/.....	2.7	2.8	3.9	3.7	3.9	5.8	*	*	*	*
2003 4/.....	2.6	3.1	3.4	2.8	*	*	*	*	*	*
2002.....	3.4	2.1	3.0	4.3	*	*	*	*	*	*
2001.....	2.4	3.0	2.9	*	4.1	*	*	*	*	*
2000.....	2.6	2.4	3.0	4.0	*	*	*	*	*	*
1999.....	2.0	2.2	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	2.7	4.4	5.8	7.9	6.7	6.6	6.3	5.6	9.4	*
2005 4/.....	3.3	4.7	6.8	7.0	7.5	8.5	8.2	6.4	*	*
2004 4/.....	3.3	5.5	5.5	6.4	7.4	7.2	9.2	6.2	*	*
2003 4/.....	3.1	4.2	4.2	5.9	6.5	4.2	6.6	5.4	*	*
2002.....	3.0	4.5	6.0	7.3	7.1	6.5	*	7.7	*	*
2001.....	2.6	4.5	5.7	5.6	5.0	5.3	7.8	6.4	*	*
2000.....	2.8	4.4	4.9	5.7	3.8	5.6	4.9	6.0	*	*
1999.....	2.5	2.8	4.3	3.5	4.2	4.6	7.2	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Metabolic disorders (E70-E88)										
All races 3/, both sexes										
2006.....	1.0	1.8	2.9	4.5	7.3	10.8	16.7	27.1	46.4	101.6
2005.....	1.2	1.6	3.0	4.6	7.0	10.9	17.9	29.0	49.2	110.7
2004.....	0.9	1.7	2.5	4.3	6.6	10.4	16.8	26.7	45.1	107.9
2003.....	1.0	1.6	2.9	4.1	6.6	10.4	16.1	25.9	46.5	112.6
2002.....	1.0	1.5	2.6	4.6	6.9	9.7	16.6	26.5	47.3	114.3
2001.....	1.0	1.7	2.7	4.1	6.3	10.1	16.4	26.3	45.2	124.4
2000.....	1.0	1.5	2.5	4.0	6.5	9.8	15.7	25.7	46.1	124.1
1999.....	0.9	1.3	2.3	3.5	6.2	9.5	15.1	25.3	44.7	130.8
All races 3/, male										
2006.....	1.2	2.3	4.0	6.4	9.6	13.7	19.9	30.7	53.1	97.6
2005.....	1.6	2.2	4.1	6.3	9.5	14.0	22.8	33.8	52.7	101.2
2004.....	1.1	2.4	3.6	5.8	8.7	13.9	20.7	30.3	50.5	103.1
2003.....	1.4	2.1	4.0	5.6	8.7	13.9	20.1	30.0	51.0	104.8
2002.....	1.3	2.0	3.7	6.1	8.9	12.5	20.0	30.4	51.5	101.5
2001.....	1.2	2.3	3.5	5.2	8.8	13.7	20.0	32.6	48.0	109.8
2000.....	1.2	2.2	3.6	5.3	8.9	12.8	19.2	29.6	48.6	112.0
1999.....	1.1	1.8	3.1	4.7	7.8	11.6	19.0	29.2	46.9	110.5
All races 3/, female										
2006.....	0.8	1.2	1.8	2.8	5.3	8.3	14.1	24.5	42.2	103.5
2005.....	0.8	1.0	1.9	2.9	4.7	8.1	14.0	25.5	47.1	115.1
2004.....	0.7	1.1	1.5	2.9	4.7	7.4	13.6	24.1	41.7	110.1
2003.....	0.7	1.1	1.9	2.7	4.7	7.4	12.9	23.0	43.8	116.1
2002.....	0.8	1.1	1.6	3.2	5.0	7.2	13.9	23.8	44.7	119.9
2001.....	0.7	1.1	1.9	3.0	3.9	7.0	13.6	22.0	43.5	130.6
2000.....	0.8	0.8	1.4	2.8	4.3	7.2	12.9	22.9	44.7	129.1
1999.....	0.7	0.9	1.7	2.4	4.7	7.7	12.1	22.7	43.4	139.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Metabolic disorders (E70-E88)										
White, both sexes										
2006 4/.....	1.0	1.7	2.8	4.4	7.3	10.5	16.6	27.2	46.4	102.3
2005 4/.....	1.2	1.5	3.0	4.4	6.9	10.7	17.9	28.8	49.4	111.3
2004 4/.....	0.9	1.7	2.5	4.2	6.5	10.2	16.7	26.7	45.2	109.2
2003 4/.....	1.0	1.6	2.8	3.9	6.6	10.1	16.1	25.3	45.9	112.9
2002.....	1.0	1.5	2.6	4.5	6.6	9.4	16.0	26.1	46.3	114.0
2001.....	1.0	1.5	2.7	4.0	6.1	9.9	16.3	26.0	44.4	123.4
2000.....	0.9	1.5	2.4	3.9	6.3	9.7	15.5	25.4	45.6	123.6
1999.....	0.8	1.3	2.2	3.5	6.0	9.1	14.8	25.0	44.2	131.3
White, male										
2006 4/.....	1.2	2.4	3.9	6.3	9.7	13.7	20.1	30.8	53.5	98.5
2005 4/.....	1.6	2.1	4.2	6.2	9.5	13.9	23.2	33.4	53.0	101.7
2004 4/.....	1.0	2.4	3.5	5.7	8.8	13.9	20.5	30.2	50.9	104.6
2003 4/.....	1.3	2.1	3.9	5.5	8.8	13.5	20.0	29.3	49.9	106.6
2002.....	1.3	1.8	3.7	6.0	8.4	12.1	19.6	29.8	50.6	100.2
2001.....	1.2	2.1	3.5	5.1	8.4	13.5	19.7	32.2	46.9	109.2
2000.....	1.1	2.2	3.5	5.2	8.6	12.6	19.1	29.1	48.1	113.9
1999.....	1.0	1.7	3.0	4.7	7.8	11.2	18.6	28.6	45.7	108.4
White, female										
2006 4/.....	0.8	1.1	1.8	2.6	5.1	7.6	13.7	24.6	42.0	104.1
2005 4/.....	0.9	1.0	1.8	2.7	4.6	7.9	13.6	25.4	47.1	115.7
2004 4/.....	0.7	1.0	1.5	2.8	4.4	7.0	13.5	24.1	41.6	111.2
2003 4/.....	0.7	1.1	1.7	2.4	4.6	7.0	12.9	22.5	43.4	115.6
2002.....	0.7	1.1	1.5	3.0	5.0	6.9	13.0	23.4	43.7	120.0
2001.....	0.8	1.0	1.9	3.0	3.8	6.7	13.6	21.6	42.9	129.4
2000.....	0.7	0.8	1.2	2.7	4.2	7.1	12.6	22.8	44.1	127.6
1999.....	0.6	0.9	1.4	2.4	4.4	7.4	11.9	22.5	43.3	140.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Metabolic disorders (E70-E88)										
Black, both sexes										
2006 4/.....	1.2	2.4	4.1	6.4	9.2	15.7	21.5	31.0	54.9	110.0
2005 4/.....	1.2	2.5	3.8	6.6	9.8	14.1	22.2	38.9	56.3	118.9
2004 4/.....	1.5	2.4	3.1	6.0	9.2	14.3	22.4	32.8	52.5	113.1
2003 4/.....	1.3	2.1	4.7	6.1	8.3	15.7	20.4	36.2	59.1	126.0
2002.....	1.3	2.5	3.8	6.5	10.8	13.9	26.0	34.7	66.5	135.1
2001.....	0.9	2.8	3.6	5.1	9.4	13.8	21.5	36.0	62.3	151.4
2000.....	1.4	2.1	3.6	5.8	9.0	13.3	20.1	33.2	61.0	147.0
1999.....	1.8	1.9	4.3	4.6	8.7	14.9	21.0	32.7	54.0	138.7
Black, male										
2006 4/.....	*	2.4	6.1	8.4	11.3	16.9	23.1	35.4	60.9	103.7
2005 4/.....	1.9	3.4	4.8	8.0	12.9	18.3	23.7	48.7	61.1	111.5
2004 4/.....	2.0	3.1	4.8	7.9	10.5	17.2	28.5	39.1	58.8	112.3
2003 4/.....	1.8	2.8	6.0	7.9	10.2	20.7	26.3	42.9	74.6	98.0
2002.....	1.5	3.5	5.0	8.5	16.4	17.7	27.4	40.9	71.8	134.0
2001.....	*	3.7	4.3	6.9	13.6	18.8	28.6	44.6	75.4	130.5
2000.....	1.7	2.9	5.1	7.1	12.2	16.7	24.4	41.1	65.8	115.3
1999.....	2.0	2.5	4.8	6.3	10.0	18.7	26.1	40.8	61.9	147.6
Black, female										
2006 4/.....	*	2.3	2.5	4.7	7.6	14.9	20.5	28.4	51.7	112.6
2005 4/.....	*	1.8	3.0	5.6	7.4	10.9	21.2	33.1	53.8	121.9
2004 4/.....	*	1.9	1.7	4.5	8.2	12.2	18.2	29.1	49.1	113.4
2003 4/.....	*	1.5	3.6	4.6	6.8	12.1	16.4	32.3	51.1	136.9
2002.....	*	1.8	2.8	4.9	6.3	11.1	25.1	31.0	63.7	135.5
2001.....	*	2.1	2.9	3.6	6.1	10.1	16.8	30.8	55.6	159.1
2000.....	*	*	2.4	4.7	6.5	10.8	17.3	28.5	58.6	158.8
1999.....	1.6	*	3.9	3.3	7.7	12.1	17.6	27.7	50.1	135.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

All races 3/, both sexes

2006.....	0.3	0.5	0.7	1.0	1.5	2.2	4.2	8.4	17.7	57.9
2005.....	0.3	0.4	0.7	0.8	1.5	2.3	4.6	9.2	21.4	68.7
2004.....	0.2	0.5	0.5	0.9	1.3	2.1	4.5	9.1	19.7	70.7
2003.....	0.2	0.4	0.7	0.8	1.5	2.3	4.6	9.9	22.4	77.3
2002.....	0.3	0.5	0.6	1.0	1.6	2.5	5.2	9.8	24.7	83.2
2001.....	0.3	0.5	0.6	0.8	1.7	2.6	5.6	11.0	25.1	93.9
2000.....	0.4	0.4	0.6	1.0	1.7	3.0	5.2	11.3	26.6	96.8
1999.....	0.3	0.4	0.7	0.8	1.4	3.0	5.5	11.9	27.7	106.4

All races 3/, male

2006.....	0.3	0.5	0.8	1.1	1.9	2.1	3.9	7.6	17.5	46.4
2005.....	0.3	0.5	0.7	1.0	1.6	2.6	4.7	10.0	20.9	56.7
2004.....	0.3	0.6	0.7	1.0	1.3	2.4	4.8	9.0	19.6	59.8
2003.....	0.2	0.5	0.7	0.9	1.5	2.5	4.6	10.2	22.2	65.6
2002.....	0.3	0.5	0.6	1.0	1.7	2.6	5.2	9.6	24.1	67.9
2001.....	0.3	0.5	0.6	0.8	2.0	2.8	5.5	12.2	24.0	76.5
2000.....	0.4	0.5	0.7	1.2	1.9	3.4	5.8	12.0	27.2	84.3
1999.....	0.4	0.5	0.8	1.1	1.4	3.1	6.2	13.2	27.2	85.0

All races 3/, female

2006.....	0.3	0.4	0.6	0.8	1.1	2.2	4.4	8.9	17.7	63.3
2005.....	0.3	0.4	0.6	0.7	1.4	2.0	4.4	8.6	21.7	74.2
2004.....	0.2	0.4	0.4	0.7	1.2	1.9	4.2	9.2	19.8	75.7
2003.....	0.2	0.4	0.7	0.6	1.5	2.2	4.6	9.6	22.5	82.5
2002.....	0.2	0.4	0.5	1.0	1.5	2.3	5.2	10.0	25.0	89.8
2001.....	0.3	0.4	0.6	0.8	1.4	2.5	5.7	10.2	25.7	101.2
2000.....	0.3	0.3	0.5	0.9	1.5	2.6	4.7	10.8	26.2	101.9
1999.....	0.2	0.3	0.5	0.7	1.5	2.9	5.0	11.1	27.9	115.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

White, both sexes

2006 4/.....	0.2	0.4	0.6	0.8	1.3	1.9	3.9	8.1	17.1	57.7
2005 4/.....	0.3	0.3	0.6	0.7	1.4	2.0	4.2	8.6	21.0	68.4
2004 4/.....	0.2	0.4	0.5	0.7	1.1	1.9	4.1	8.9	19.4	70.8
2003 4/.....	0.2	0.4	0.6	0.7	1.4	2.0	4.4	9.2	21.7	76.6
2002.....	0.2	0.4	0.5	0.9	1.4	2.2	4.7	9.2	23.5	82.4
2001.....	0.3	0.3	0.5	0.7	1.4	2.4	5.1	10.4	24.0	93.0
2000.....	0.3	0.3	0.5	0.9	1.5	2.7	4.8	10.9	25.9	96.0
1999.....	0.2	0.4	0.5	0.7	1.2	2.5	5.2	11.4	27.2	106.6

White, male

2006 4/.....	*	0.5	0.7	0.9	1.8	1.9	3.6	7.1	16.7	45.8
2005 4/.....	0.3	0.4	0.6	0.8	1.4	2.3	4.5	9.2	20.4	56.1
2004 4/.....	*	0.5	0.7	0.8	1.0	2.3	4.5	8.6	19.3	59.0
2003 4/.....	*	0.4	0.5	0.9	1.4	2.1	4.3	9.4	20.9	65.6
2002.....	0.2	0.4	0.5	0.8	1.5	2.2	4.8	8.9	22.7	66.1
2001.....	0.2	0.4	0.5	0.6	1.6	2.6	4.7	11.5	22.3	75.8
2000.....	0.3	0.4	0.6	1.0	1.6	3.0	5.3	11.2	26.3	84.9
1999.....	0.2	0.5	0.7	0.9	1.1	2.3	5.6	12.3	26.0	82.4

White, female

2006 4/.....	0.3	0.4	0.5	0.7	0.9	1.9	4.1	8.8	17.3	63.2
2005 4/.....	0.3	0.3	0.6	0.6	1.4	1.8	4.0	8.2	21.4	74.0
2004 4/.....	*	0.3	0.4	0.6	1.2	1.6	3.8	9.1	19.5	76.1
2003 4/.....	*	0.3	0.6	0.5	1.4	1.9	4.4	9.0	22.2	81.5
2002.....	*	0.4	0.5	1.0	1.4	2.2	4.7	9.4	24.0	89.5
2001.....	0.3	0.3	0.5	0.7	1.3	2.1	5.4	9.7	25.0	100.2
2000.....	0.3	0.3	0.4	0.8	1.4	2.3	4.4	10.7	25.7	100.5
1999.....	*	0.3	0.4	0.6	1.2	2.6	4.8	10.8	27.9	116.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Volume depletion, disorders of fluid, electrolyte and acid-base balance (E86-E87)

Black, both sexes

2006 4/.....	0.8	*	1.5	2.3	2.9	5.1	8.0	13.7	28.2	72.4
2005 4/.....	*	1.0	1.5	1.8	3.2	5.3	8.7	17.9	31.1	83.1
2004 4/.....	0.7	1.0	*	2.4	3.0	4.5	9.3	13.9	28.0	84.3
2003 4/.....	*	0.8	2.1	2.0	3.0	6.1	8.2	20.2	35.0	97.1
2002.....	0.8	1.2	1.3	2.0	3.1	5.4	11.4	18.9	42.9	104.2
2001.....	*	1.4	1.6	1.8	4.2	5.5	12.1	20.7	42.8	119.4
2000.....	0.8	1.3	1.6	2.2	3.4	6.6	10.1	18.5	40.9	119.8
1999.....	1.1	1.0	1.9	2.0	4.0	7.8	11.0	19.8	36.7	115.1

Black, male

2006 4/.....	*	*	2.0	3.2	3.5	4.9	7.7	14.8	34.2	65.2
2005 4/.....	*	*	*	*	4.2	6.4	8.2	22.4	35.8	72.5
2004 4/.....	*	*	*	3.6	4.1	5.3	10.4	16.4	31.6	86.4
2003 4/.....	*	*	2.7	*	*	7.4	9.5	23.3	48.0	77.2
2002.....	*	1.8	*	*	4.4	7.0	11.3	20.7	46.6	99.7
2001.....	*	*	*	*	5.9	5.8	15.8	24.2	54.6	97.6
2000.....	*	*	*	*	4.6	8.1	12.0	26.0	47.2	98.0
1999.....	1.5	*	*	*	*	11.4	14.9	26.6	44.5	125.4

Black, female

2006 4/.....	*	*	*	*	*	5.2	8.2	13.1	25.0	75.3
2005 4/.....	*	*	*	*	*	4.4	9.1	15.2	28.6	87.4
2004 4/.....	*	*	*	*	*	4.0	8.6	12.4	26.1	83.5
2003 4/.....	*	*	*	*	3.0	5.1	7.3	18.4	28.2	104.9
2002.....	*	*	*	*	*	4.3	11.4	17.8	41.0	106.0
2001.....	*	*	*	*	*	5.3	9.6	18.6	36.8	127.6
2000.....	*	*	*	*	*	5.6	8.9	14.0	37.8	127.9
1999.....	*	*	*	*	4.0	5.2	8.3	15.7	32.8	111.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other metabolic disorders (E70-E83,E85,E88)										
All races 3/, both sexes										
2006.....	0.6	1.2	2.1	3.5	5.8	8.6	12.5	18.8	28.7	43.6
2005.....	0.7	1.1	2.3	3.7	5.5	8.5	13.3	19.8	27.9	42.0
2004.....	0.6	1.2	1.9	3.4	5.3	8.2	12.3	17.6	25.3	37.1
2003.....	0.6	1.1	2.2	3.3	5.1	8.1	11.5	16.0	24.1	35.2
2002.....	0.6	1.0	2.0	3.6	5.2	7.2	11.4	16.7	22.6	31.1
2001.....	0.5	1.1	2.0	3.2	4.6	7.4	10.8	15.3	20.0	30.5
2000.....	0.4	1.0	1.9	2.9	4.8	6.8	10.5	14.3	19.6	27.3
1999.....	0.5	0.9	1.6	2.6	4.7	6.5	9.5	13.4	17.0	24.3
All races 3/, male										
2006.....	0.8	1.7	3.0	5.2	7.7	11.5	15.9	23.1	35.6	51.2
2005.....	1.1	1.6	3.3	5.4	7.9	11.2	18.0	23.7	31.9	44.5
2004.....	0.8	1.7	2.8	4.7	7.4	11.4	15.8	21.3	30.9	43.2
2003.....	0.9	1.5	3.2	4.6	7.2	11.4	15.5	19.7	28.8	39.2
2002.....	0.9	1.4	3.1	5.1	7.1	9.8	14.7	20.8	27.3	33.6
2001.....	0.7	1.6	2.8	4.4	6.8	10.9	14.5	20.3	23.8	33.3
2000.....	0.6	1.5	2.9	4.0	6.9	9.3	13.4	17.6	21.4	27.7
1999.....	0.6	1.3	2.2	3.6	6.4	8.6	12.7	15.9	19.6	25.5
All races 3/, female										
2006.....	0.4	0.7	1.2	2.0	4.1	6.1	9.7	15.6	24.4	40.1
2005.....	0.4	0.6	1.3	2.2	3.3	6.1	9.5	16.9	25.4	40.8
2004.....	0.4	0.6	1.0	2.2	3.4	5.5	9.4	14.9	21.8	34.4
2003.....	0.3	0.7	1.2	2.0	3.2	5.2	8.4	13.4	21.2	33.4
2002.....	0.4	0.7	1.0	2.2	3.5	4.9	8.7	13.8	19.7	30.0
2001.....	0.4	0.6	1.2	2.1	2.6	4.5	7.8	11.7	17.8	29.3
2000.....	0.3	0.5	0.9	1.9	2.8	4.6	8.2	12.1	18.5	27.1
1999.....	0.4	0.5	1.0	1.7	3.2	4.8	7.1	11.6	15.5	23.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other metabolic disorders (E70-E83,E85,E88)										
White, both sexes										
2006 4/.....	0.6	1.2	2.1	3.6	5.9	8.6	12.7	19.1	29.3	44.6
2005 4/.....	0.7	1.1	2.3	3.7	5.5	8.6	13.6	20.1	28.4	42.9
2004 4/.....	0.6	1.2	1.9	3.5	5.4	8.3	12.5	17.8	25.7	38.4
2003 4/.....	0.6	1.1	2.2	3.2	5.2	8.1	11.7	16.1	24.2	36.1
2002.....	0.6	1.0	2.0	3.6	5.2	7.2	11.2	16.9	22.7	31.6
2001.....	0.6	1.1	2.1	3.3	4.6	7.5	11.2	15.5	20.3	30.4
2000.....	0.4	1.1	1.9	3.0	4.8	7.0	10.7	14.5	19.7	27.5
1999.....	0.5	0.9	1.6	2.7	4.8	6.7	9.7	13.6	16.9	24.6
White, male										
2006 4/.....	0.8	1.8	3.0	5.4	7.9	11.8	16.4	23.6	36.7	52.7
2005 4/.....	1.1	1.6	3.5	5.4	8.1	11.5	18.6	24.2	32.6	45.6
2004 4/.....	0.8	1.8	2.8	4.9	7.7	11.6	15.9	21.5	31.6	45.5
2003 4/.....	0.9	1.5	3.3	4.6	7.4	11.4	15.7	19.8	29.0	40.9
2002.....	0.9	1.4	3.2	5.2	6.9	9.9	14.8	20.9	27.8	34.1
2001.....	0.8	1.6	2.9	4.4	6.8	10.9	15.0	20.7	24.4	33.3
2000.....	0.6	1.7	2.9	4.1	6.9	9.5	13.8	18.0	21.8	29.0
1999.....	0.6	1.2	2.2	3.7	6.6	8.9	13.0	16.2	19.6	25.9
White, female										
2006 4/.....	0.4	0.6	1.2	1.9	4.1	5.7	9.6	15.8	24.6	40.8
2005 4/.....	0.4	0.6	1.2	2.0	3.2	6.1	9.5	17.2	25.7	41.7
2004 4/.....	0.4	0.6	1.0	2.1	3.2	5.3	9.7	15.0	22.0	35.1
2003 4/.....	0.3	0.7	1.1	1.9	3.2	5.2	8.5	13.4	21.2	34.0
2002.....	0.4	0.7	0.9	2.1	3.6	4.7	8.3	14.0	19.6	30.4
2001.....	0.4	0.6	1.3	2.2	2.6	4.5	8.1	11.8	17.9	29.1
2000.....	0.2	0.5	0.9	1.8	2.8	4.7	8.2	12.1	18.4	26.9
1999.....	0.3	0.5	0.9	1.7	3.2	4.7	7.0	11.8	15.4	24.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other metabolic disorders (E70-E83,E85,E88)										
Black, both sexes										
2006 4/.....	*	1.7	2.6	4.0	6.3	10.6	13.5	17.3	26.7	37.6
2005 4/.....	0.8	1.5	2.3	4.8	6.6	8.8	13.5	21.0	25.2	35.8
2004 4/.....	0.7	1.4	2.4	3.6	6.1	9.8	13.1	19.0	24.5	28.8
2003 4/.....	0.7	1.3	2.5	4.1	5.3	9.7	12.2	15.8	24.1	28.9
2002.....	*	1.3	2.4	4.4	7.6	8.5	14.5	15.8	23.5	30.9
2001.....	*	1.5	1.9	3.2	5.2	8.3	9.4	15.3	19.5	31.9
2000.....	*	*	2.0	3.6	5.7	6.6	10.0	14.7	20.0	27.2
1999.....	0.7	0.9	2.3	2.7	4.7	7.1	10.0	12.9	17.3	23.6
Black, male										
2006 4/.....	*	1.8	4.1	5.2	7.8	11.7	15.4	20.6	26.7	38.4
2005 4/.....	*	2.0	3.1	6.0	8.7	11.9	15.5	26.4	25.3	39.0
2004 4/.....	*	1.9	3.6	4.4	6.4	11.9	18.1	22.7	27.3	25.8
2003 4/.....	*	1.6	3.2	5.8	7.3	13.3	16.8	19.6	26.6	20.9
2002.....	*	1.7	3.3	5.8	12.1	10.7	15.7	20.2	25.2	34.3
2001.....	*	2.1	2.5	4.7	7.7	13.0	12.8	20.4	20.9	32.9
2000.....	*	*	3.1	4.0	7.7	8.6	12.4	15.1	18.5	*
1999.....	*	*	2.5	*	5.9	7.3	11.2	14.2	17.4	*
Black, female										
2006 4/.....	*	1.5	*	3.1	5.1	9.7	12.3	15.3	26.8	37.3
2005 4/.....	*	*	1.6	3.8	5.0	6.5	12.1	17.9	25.2	34.5
2004 4/.....	*	*	*	3.0	6.0	8.3	9.7	16.7	23.1	29.9
2003 4/.....	*	*	2.0	2.7	3.8	6.9	9.1	13.6	22.8	32.0
2002.....	*	*	*	3.3	4.0	6.8	13.7	13.2	22.7	29.6
2001.....	*	*	*	*	3.2	4.9	7.2	12.2	18.8	31.6
2000.....	*	*	*	3.2	4.1	5.2	8.4	14.5	20.8	30.9
1999.....	*	*	2.2	*	3.7	6.9	9.2	12.0	17.3	24.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

V. Mental and behavioral disorders (F01-F99)

All races 3/, both sexes

2006.....	6.5	8.7	10.6	10.7	12.9	19.0	40.6	104.5	275.6	977.5
2005.....	6.4	8.7	10.1	9.8	11.6	15.3	31.4	77.8	211.9	766.4
2004.....	6.0	8.6	9.5	9.2	10.5	13.8	28.6	69.6	180.4	671.4
2003.....	6.3	8.4	8.9	8.3	10.4	13.6	27.2	69.8	179.2	679.3
2002.....	6.5	7.9	8.3	8.3	9.5	13.0	25.6	64.2	168.2	646.7
2001.....	6.0	7.9	7.7	7.7	9.4	13.0	23.8	60.6	154.3	604.8
2000.....	5.8	7.5	7.1	7.8	8.6	11.7	22.5	54.1	143.7	561.6
1999.....	6.1	7.4	6.8	7.3	8.9	11.2	21.2	49.9	128.6	515.3

All races 3/, male

2006.....	9.4	12.5	16.0	16.2	18.5	24.7	47.1	108.7	277.0	775.4
2005.....	9.4	12.6	15.5	15.1	17.1	19.7	35.5	81.7	208.4	597.0
2004.....	8.8	12.5	14.9	14.5	15.8	18.7	34.0	73.2	174.5	529.7
2003.....	9.2	12.7	14.0	12.8	15.0	18.5	32.1	73.2	175.7	523.8
2002.....	9.3	12.0	13.4	12.9	14.4	18.4	30.2	68.2	166.1	506.0
2001.....	8.7	12.2	12.3	11.7	14.3	18.1	29.3	63.3	151.3	488.0
2000.....	8.5	11.6	11.4	11.9	13.0	16.5	28.0	59.4	140.5	451.0
1999.....	9.4	11.5	11.1	11.7	13.8	16.1	26.9	56.9	133.4	415.8

All races 3/, female

2006.....	3.7	5.1	5.4	5.4	7.8	14.0	35.4	101.5	274.7	1,072.1
2005.....	3.3	4.9	4.9	4.9	6.6	11.4	28.0	74.9	214.1	844.2
2004.....	3.2	4.9	4.3	4.3	5.6	9.6	24.2	67.0	184.1	735.1
2003.....	3.5	4.2	4.1	4.1	6.2	9.4	23.4	67.4	181.4	748.1
2002.....	3.6	4.0	3.3	4.0	5.1	8.4	21.9	61.3	169.4	707.8
2001.....	3.3	3.8	3.3	3.9	4.9	8.6	19.5	58.8	156.0	653.7
2000.....	3.2	3.5	3.1	4.0	4.6	7.7	18.3	50.4	145.6	606.6
1999.....	2.8	3.4	2.8	3.3	4.5	7.0	16.7	45.1	125.8	555.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
V. Mental and behavioral disorders (F01-F99)										
White, both sexes										
2006 4/.....	6.5	8.5	10.1	10.0	12.3	18.5	39.5	104.5	276.2	998.9
2005 4/.....	6.4	8.5	9.5	9.5	11.5	14.9	30.5	77.3	214.0	783.9
2004 4/.....	6.0	8.4	9.0	8.9	10.0	13.0	28.1	69.0	183.4	687.9
2003 4/.....	6.1	8.1	8.3	7.8	9.8	13.5	27.1	69.8	180.9	697.4
2002.....	6.2	7.5	7.6	7.7	8.8	12.5	25.3	64.1	170.5	663.8
2001.....	5.5	7.1	6.8	6.9	8.9	12.5	23.5	61.1	157.4	621.6
2000.....	5.4	6.7	6.6	7.2	7.9	11.2	22.0	54.0	145.3	577.6
1999.....	5.5	6.5	6.1	6.5	8.3	10.7	20.7	50.0	130.3	534.6
White, male										
2006 4/.....	9.3	12.2	15.1	14.9	17.4	23.5	44.5	106.9	274.2	790.6
2005 4/.....	9.5	12.3	14.4	14.3	16.6	18.7	33.9	80.3	208.1	610.0
2004 4/.....	8.9	12.3	13.8	13.7	14.6	17.0	33.0	70.7	175.8	538.9
2003 4/.....	9.0	12.2	12.7	12.0	13.4	18.0	31.0	71.8	175.5	535.8
2002.....	9.2	11.3	12.3	11.6	13.0	16.9	29.0	66.5	167.4	514.9
2001.....	8.0	10.9	10.8	10.2	13.2	16.8	28.3	63.1	152.8	498.4
2000.....	7.9	10.3	10.4	10.8	11.6	15.2	26.7	58.9	140.3	459.4
1999.....	8.6	10.2	9.8	10.1	12.8	14.9	25.7	55.8	132.9	429.2
White, female										
2006 4/.....	3.6	4.8	5.2	5.3	7.5	14.1	35.3	102.8	277.4	1,096.6
2005 4/.....	3.2	4.7	4.6	4.9	6.7	11.4	27.7	75.1	217.7	864.0
2004 4/.....	3.1	4.5	4.2	4.3	5.7	9.5	24.0	67.8	188.1	755.1
2003 4/.....	3.1	4.0	3.9	3.8	6.4	9.4	23.9	68.2	184.2	769.0
2002.....	3.3	3.7	3.1	4.0	4.9	8.7	22.3	62.4	172.4	728.4
2001.....	2.9	3.4	2.9	3.8	4.9	8.7	19.6	59.7	160.2	673.1
2000.....	2.9	3.1	2.8	3.8	4.4	7.7	18.3	50.6	148.3	625.6
1999.....	2.3	2.8	2.4	3.1	4.2	6.9	16.7	45.9	128.8	576.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
V. Mental and behavioral disorders (F01-F99)										
Black, both sexes										
2006 4/.....	7.9	12.4	16.1	18.0	23.0	27.4	63.0	131.9	325.9	896.1
2005 4/.....	7.7	12.1	16.2	15.0	16.4	21.4	47.9	104.1	238.6	701.6
2004 4/.....	6.7	12.6	15.5	14.7	16.6	24.0	41.0	95.5	188.4	589.4
2003 4/.....	8.4	12.1	15.5	14.4	18.1	18.0	34.7	86.8	198.0	578.8
2002.....	9.5	12.6	15.0	15.9	16.8	19.8	34.5	78.3	173.2	540.8
2001.....	10.3	15.3	16.2	15.4	15.9	20.7	33.5	70.5	147.9	488.5
2000.....	9.4	15.2	13.7	14.2	16.2	18.7	33.4	68.4	145.4	442.6
1999.....	10.5	15.1	13.9	16.5	15.4	18.8	30.8	57.8	128.3	349.1
Black, male										
2006 4/.....	11.0	17.3	25.7	30.3	36.2	42.6	88.3	158.9	373.1	726.5
2005 4/.....	10.7	17.9	26.2	25.3	27.4	31.1	63.6	122.9	273.2	544.7
2004 4/.....	9.9	17.5	25.8	26.0	30.3	39.5	54.7	126.0	203.7	510.6
2003 4/.....	11.5	18.9	27.0	23.1	33.0	27.3	51.6	106.4	216.3	482.7
2002.....	12.8	19.5	25.7	28.0	28.9	35.4	51.1	103.3	189.5	457.8
2001.....	14.9	24.5	27.3	27.4	29.1	34.6	47.8	84.6	171.8	423.2
2000.....	13.6	24.4	23.6	24.4	27.7	32.5	48.8	82.2	169.4	415.2
1999.....	15.4	23.1	23.2	29.5	26.5	31.4	46.5	79.8	168.3	309.3
Black, female										
2006 4/.....	5.2	8.2	7.9	7.8	12.7	16.1	45.6	115.7	300.9	965.5
2005 4/.....	5.0	7.0	7.7	6.5	7.8	14.1	37.2	92.8	220.3	764.9
2004 4/.....	4.0	8.2	6.9	5.3	6.0	12.4	31.6	77.4	180.3	620.6
2003 4/.....	5.7	6.2	5.8	7.2	6.5	11.1	23.3	75.2	188.4	616.1
2002.....	6.7	6.7	5.8	6.0	7.3	8.1	23.3	63.4	164.7	572.6
2001.....	6.2	7.4	6.7	5.5	5.6	10.5	24.0	62.1	135.7	512.8
2000.....	5.6	7.2	5.4	5.9	7.0	8.4	23.1	60.1	133.3	452.9
1999.....	6.2	8.1	6.1	6.0	6.7	9.4	20.5	44.6	108.5	364.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Organic dementia (F01,F03)										
All races 3/, both sexes										
2006.....	*	0.2	0.5	1.5	4.1	10.7	31.8	92.9	258.4	939.9
2005.....	*	0.1	0.3	0.9	2.7	7.2	23.3	66.8	193.9	725.2
2004.....	*	0.1	0.2	0.7	2.2	5.7	19.7	58.5	163.4	628.2
2003.....	*	*	0.2	0.6	2.2	5.9	18.5	58.6	162.7	631.7
2002.....	*	0.1	0.2	0.5	1.8	5.4	17.0	53.8	150.9	598.3
2001.....	*	*	0.2	0.6	1.8	5.5	15.9	49.5	135.8	554.0
2000.....	*	0.1	0.1	0.6	1.4	4.5	14.4	43.0	125.3	505.6
1999.....	*	*	0.2	0.5	1.4	4.0	13.0	37.8	108.1	454.4
All races 3/, male										
2006.....	*	0.2	0.5	1.5	4.7	12.3	34.5	94.0	255.6	737.2
2005.....	*	*	0.3	0.9	3.1	7.6	24.2	66.2	187.8	557.7
2004.....	*	*	0.3	0.8	2.5	6.2	20.5	58.4	154.1	487.8
2003.....	*	*	0.2	0.7	2.4	6.7	20.1	58.0	155.7	480.4
2002.....	*	*	*	0.5	1.9	6.2	18.7	55.1	147.0	465.7
2001.....	*	*	*	0.5	2.0	6.2	17.5	49.7	130.8	442.4
2000.....	*	*	*	0.5	1.7	5.2	16.3	45.4	120.5	398.9
1999.....	*	*	*	0.6	1.7	4.9	15.3	40.5	109.7	361.7
All races 3/, female										
2006.....	*	0.2	0.4	1.5	3.5	9.3	29.6	92.2	260.1	1,034.7
2005.....	*	*	0.2	0.9	2.3	6.9	22.5	67.2	197.7	802.1
2004.....	*	*	*	0.6	2.0	5.3	19.0	58.6	169.2	691.3
2003.....	*	*	0.2	0.5	2.1	5.2	17.3	59.0	166.9	698.5
2002.....	*	*	*	0.6	1.6	4.6	15.7	52.9	153.3	655.9
2001.....	*	*	*	0.6	1.5	4.9	14.5	49.4	138.8	600.8
2000.....	*	*	*	0.7	1.2	4.0	12.8	41.4	128.2	549.1
1999.....	*	*	*	*	1.1	3.2	11.3	35.9	107.1	491.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Organic dementia (F01,F03)										
White, both sexes										
2006 4/.....	*	0.2	0.4	1.5	3.8	10.3	30.7	92.8	258.2	959.7
2005 4/.....	*	0.1	0.3	0.9	2.6	6.8	22.4	66.1	195.5	741.4
2004 4/.....	*	*	0.2	0.7	2.2	5.5	19.2	57.8	165.7	642.8
2003 4/.....	*	*	0.2	0.6	2.2	5.8	18.4	58.3	163.9	647.9
2002.....	*	*	0.2	0.5	1.8	5.2	16.9	53.6	153.0	613.9
2001.....	*	*	0.2	0.6	1.8	5.4	15.6	49.9	138.3	569.1
2000.....	*	*	0.1	0.6	1.3	4.4	13.9	42.9	126.6	519.6
1999.....	*	*	0.2	0.5	1.4	3.8	12.9	37.9	109.6	471.2
White, male										
2006 4/.....	*	*	0.5	1.4	4.5	11.6	32.4	92.0	252.5	751.1
2005 4/.....	*	*	0.3	0.9	3.0	7.0	22.8	64.9	187.5	569.5
2004 4/.....	*	*	0.3	0.8	2.4	5.9	19.9	56.2	154.3	496.5
2003 4/.....	*	*	*	0.7	2.3	6.6	19.4	56.6	155.3	491.1
2002.....	*	*	*	0.5	1.9	5.8	18.2	53.7	148.1	473.7
2001.....	*	*	*	0.5	2.0	5.9	16.9	49.8	132.0	452.2
2000.....	*	*	*	0.5	1.6	4.9	15.4	45.2	120.2	406.1
1999.....	*	*	*	0.6	1.7	4.8	14.9	40.2	109.4	373.7
White, female										
2006 4/.....	*	0.2	0.4	1.5	3.2	9.2	29.3	93.3	261.8	1,057.7
2005 4/.....	*	*	*	0.9	2.3	6.6	22.1	67.0	200.5	820.5
2004 4/.....	*	*	*	0.6	2.0	5.2	18.7	59.0	172.8	708.7
2003 4/.....	*	*	*	0.5	2.2	5.1	17.6	59.5	169.2	717.4
2002.....	*	*	*	0.6	1.7	4.8	15.9	53.6	155.9	674.7
2001.....	*	*	*	0.6	1.6	5.0	14.5	50.0	142.2	618.0
2000.....	*	*	*	0.7	1.1	3.9	12.6	41.3	130.4	565.7
1999.....	*	*	*	*	1.1	3.0	11.2	36.3	109.7	510.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Organic dementia (F01,F03)										
Black, both sexes										
2006 4/.....	*	*	*	1.8	7.8	16.8	51.6	119.3	313.3	869.6
2005 4/.....	*	*	*	*	4.0	12.0	37.6	91.5	221.4	667.6
2004 4/.....	*	*	*	*	2.9	9.6	29.3	83.3	175.3	560.3
2003 4/.....	*	*	*	*	3.2	8.3	24.3	75.0	182.0	545.3
2002.....	*	*	*	*	2.2	7.8	22.4	66.8	157.3	503.1
2001.....	*	*	*	*	2.2	8.2	23.0	57.5	132.7	452.3
2000.....	*	*	*	*	2.5	6.4	22.5	55.3	127.6	403.2
1999.....	*	*	*	*	*	6.0	17.7	42.8	107.5	309.3
Black, male										
2006 4/.....	*	*	*	*	8.3	23.2	67.1	141.9	351.2	700.6
2005 4/.....	*	*	*	*	5.3	13.5	46.2	101.0	247.9	509.4
2004 4/.....	*	*	*	*	3.9	12.6	32.7	104.2	192.3	468.9
2003 4/.....	*	*	*	*	4.6	9.6	33.5	88.1	194.9	445.2
2002.....	*	*	*	*	*	11.0	28.7	84.0	169.7	425.6
2001.....	*	*	*	*	*	11.2	29.3	62.7	150.9	382.4
2000.....	*	*	*	*	*	9.1	29.4	61.4	145.8	372.5
1999.....	*	*	*	*	*	7.5	23.1	51.3	136.0	266.0
Black, female										
2006 4/.....	*	*	*	*	7.4	12.1	41.0	105.8	293.3	938.8
2005 4/.....	*	*	*	*	3.0	10.9	31.8	85.9	207.4	731.5
2004 4/.....	*	*	*	*	*	7.4	27.0	70.9	166.3	596.6
2003 4/.....	*	*	*	*	*	7.3	18.2	67.3	175.2	584.1
2002.....	*	*	*	*	*	5.4	18.1	56.5	150.8	532.7
2001.....	*	*	*	*	*	6.0	18.8	54.4	123.5	478.2
2000.....	*	*	*	*	*	4.4	18.0	51.6	118.5	414.7
1999.....	*	*	*	*	*	4.8	14.2	37.8	93.3	325.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified organic mental disorders (F04-F09)

All races 3/, both sexes

2006.....	*	*	0.1	*	0.2	0.4	0.9	2.5	4.8	18.7
2005.....	*	*	*	*	0.2	0.4	0.7	2.4	5.5	20.1
2004.....	*	*	*	*	0.2	0.4	0.8	2.2	6.0	22.3
2003.....	*	*	*	*	0.2	0.3	1.0	2.6	6.0	24.5
2002.....	*	*	*	0.1	0.2	0.5	1.2	2.4	6.5	26.7
2001.....	*	*	*	*	0.2	0.5	1.1	2.9	7.8	29.3
2000.....	*	*	*	0.2	0.3	0.5	1.2	3.2	7.5	33.7
1999.....	*	*	*	*	0.2	0.4	1.2	3.3	8.7	37.7

All races 3/, male

2006.....	*	*	*	*	*	*	1.0	2.3	5.1	16.3
2005.....	*	*	*	*	*	0.4	0.8	2.8	4.9	16.2
2004.....	*	*	*	*	*	0.5	1.0	2.7	6.4	19.4
2003.....	*	*	*	*	*	0.5	1.0	3.1	6.7	20.8
2002.....	*	*	*	*	*	0.7	1.3	2.4	6.9	20.1
2001.....	*	*	*	*	*	0.7	1.3	3.0	7.7	23.8
2000.....	*	*	*	*	*	0.6	1.4	3.4	6.4	28.4
1999.....	*	*	*	*	*	0.5	1.3	3.6	9.0	29.8

All races 3/, female

2006.....	*	*	*	*	*	0.5	0.7	2.5	4.6	19.9
2005.....	*	*	*	*	*	*	0.7	2.0	6.0	21.8
2004.....	*	*	*	*	*	*	0.6	1.9	5.7	23.6
2003.....	*	*	*	*	*	*	1.0	2.3	5.5	26.2
2002.....	*	*	*	*	*	*	1.1	2.4	6.4	29.6
2001.....	*	*	*	*	*	*	1.0	2.9	7.9	31.6
2000.....	*	*	*	*	*	*	1.1	3.1	8.2	35.8
1999.....	*	*	*	*	*	*	1.1	3.1	8.6	40.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified organic mental disorders (F04-F09)										
White, both sexes										
2006 4/.....	*	*	*	*	0.2	0.4	0.9	2.5	5.1	19.6
2005 4/.....	*	*	*	*	0.2	0.4	0.8	2.3	5.7	20.6
2004 4/.....	*	*	*	*	0.2	0.4	0.9	2.3	6.1	23.3
2003 4/.....	*	*	*	*	0.2	0.3	1.0	2.7	6.1	25.1
2002.....	*	*	*	*	*	0.4	1.1	2.5	6.7	27.5
2001.....	*	*	*	*	*	0.5	1.1	2.9	8.0	30.2
2000.....	*	*	*	0.2	0.2	0.5	1.3	3.3	7.5	34.7
1999.....	*	*	*	*	0.2	0.4	1.1	3.2	8.8	38.9
White, male										
2006 4/.....	*	*	*	*	*	*	1.0	2.4	5.3	17.2
2005 4/.....	*	*	*	*	*	*	0.8	2.7	5.0	16.7
2004 4/.....	*	*	*	*	*	*	1.1	2.8	6.7	20.1
2003 4/.....	*	*	*	*	*	*	0.9	3.1	6.8	21.3
2002.....	*	*	*	*	*	0.6	1.3	2.5	7.1	20.6
2001.....	*	*	*	*	*	0.8	1.3	2.9	7.8	24.2
2000.....	*	*	*	*	*	0.6	1.4	3.5	6.4	28.5
1999.....	*	*	*	*	*	*	1.3	3.3	8.9	30.1
White, female										
2006 4/.....	*	*	*	*	*	0.5	0.8	2.6	5.0	20.8
2005 4/.....	*	*	*	*	*	*	0.7	2.0	6.2	22.3
2004 4/.....	*	*	*	*	*	*	0.7	1.9	5.8	24.7
2003 4/.....	*	*	*	*	*	*	1.0	2.3	5.7	26.8
2002.....	*	*	*	*	*	*	1.0	2.5	6.5	30.5
2001.....	*	*	*	*	*	*	1.0	2.9	8.2	32.6
2000.....	*	*	*	*	*	*	1.2	3.2	8.2	37.3
1999.....	*	*	*	*	*	*	1.0	3.2	8.7	42.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified organic mental disorders (F04-F09)

Black, both sexes

2006 4/.....	*	*	*	*	*	*	*	*	*	13.8
2005 4/.....	*	*	*	*	*	*	*	3.5	5.6	18.2
2004 4/.....	*	*	*	*	*	*	*	*	5.4	16.6
2003 4/.....	*	*	*	*	*	*	*	*	5.8	20.4
2002.....	*	*	*	*	*	*	*	*	5.5	22.9
2001.....	*	*	*	*	*	*	*	4.2	7.3	24.3
2000.....	*	*	*	*	*	*	*	*	9.6	25.3
1999.....	*	*	*	*	*	*	*	4.3	10.4	28.7

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	22.7
2000.....	*	*	*	*	*	*	*	*	*	25.4
1999.....	*	*	*	*	*	*	*	*	*	29.3

Black, female

2006 4/.....	*	*	*	*	*	*	*	*	*	15.7
2005 4/.....	*	*	*	*	*	*	*	*	*	19.9
2004 4/.....	*	*	*	*	*	*	*	*	*	16.5
2003 4/.....	*	*	*	*	*	*	*	*	*	20.7
2002.....	*	*	*	*	*	*	*	*	*	25.1
2001.....	*	*	*	*	*	*	*	*	*	24.8
2000.....	*	*	*	*	*	*	*	*	9.8	25.3
1999.....	*	*	*	*	*	*	*	*	8.6	28.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

All races 3/, both sexes

2006.....	6.1	8.0	9.5	8.4	7.8	6.8	6.3	6.1	6.6	6.7
2005.....	6.0	8.1	9.2	8.2	7.8	6.6	5.4	5.4	5.9	6.8
2004.....	5.6	8.0	8.6	7.7	7.2	6.7	6.1	5.5	5.3	4.6
2003.....	5.9	7.8	8.1	6.9	7.1	6.2	5.7	4.9	4.4	4.3
2002.....	6.0	7.3	7.6	6.9	6.7	6.1	5.3	4.7	3.8	3.7
2001.....	5.5	7.4	7.0	6.4	6.6	5.7	5.1	4.6	4.5	4.0
2000.....	5.5	7.0	6.5	6.4	6.0	5.8	5.1	4.7	4.5	4.1
1999.....	5.7	6.9	6.2	6.2	6.6	5.8	5.0	5.3	4.4	3.8

All races 3/, male

2006.....	8.9	11.7	14.7	13.7	12.6	10.9	10.0	9.2	10.8	11.4
2005.....	8.9	11.9	14.5	13.2	12.6	10.7	8.7	9.6	9.5	10.7
2004.....	8.4	11.7	13.8	12.8	12.2	10.8	10.1	9.0	8.8	8.2
2003.....	8.8	12.0	13.0	11.2	11.5	10.2	9.1	8.4	7.4	7.3
2002.....	8.8	11.2	12.6	11.5	11.2	10.4	8.1	7.8	6.2	7.0
2001.....	8.2	11.6	11.5	10.4	11.1	9.7	8.6	7.2	7.3	7.5
2000.....	8.1	11.1	10.7	10.4	10.1	9.7	8.3	8.0	7.9	7.3
1999.....	9.0	11.0	10.2	10.2	11.1	9.5	8.1	9.0	8.4	7.4

All races 3/, female

2006.....	3.4	4.5	4.4	3.3	3.3	3.3	3.2	3.8	4.0	4.5
2005.....	3.1	4.4	4.1	3.4	3.5	3.1	2.7	2.3	3.6	5.0
2004.....	2.8	4.3	3.6	2.9	2.6	3.1	2.9	2.9	3.1	3.0
2003.....	3.1	3.7	3.3	2.9	3.1	2.9	2.9	2.5	2.5	3.0
2002.....	3.2	3.4	2.8	2.6	2.5	2.4	3.1	2.5	2.4	2.2
2001.....	2.9	3.4	2.8	2.7	2.6	2.3	2.3	2.9	2.9	2.5
2000.....	2.9	3.0	2.5	2.7	2.3	2.4	2.6	2.5	2.6	2.8
1999.....	2.5	3.1	2.3	2.5	2.5	2.6	2.5	2.7	2.0	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

White, both sexes

2006 4/.....	6.0	7.8	9.0	7.8	7.4	6.7	6.1	6.1	6.7	6.7
2005 4/.....	6.0	7.8	8.5	7.9	7.7	6.6	5.4	5.4	5.9	6.9
2004 4/.....	5.6	7.7	8.0	7.3	6.7	6.2	5.8	5.4	5.5	4.7
2003 4/.....	5.7	7.5	7.4	6.4	6.4	6.2	5.6	5.0	4.5	4.5
2002.....	5.8	6.8	6.9	6.3	5.9	5.8	5.1	4.7	3.8	3.6
2001.....	5.0	6.7	6.2	5.7	6.1	5.3	5.0	4.6	4.6	4.0
2000.....	5.0	6.2	5.9	5.8	5.4	5.3	5.0	4.7	4.6	4.1
1999.....	5.1	6.0	5.4	5.4	6.0	5.4	4.7	5.1	4.3	3.9

White, male

2006 4/.....	8.8	11.5	13.9	12.5	11.7	10.3	9.4	9.3	10.9	11.5
2005 4/.....	9.0	11.5	13.4	12.6	12.1	10.4	8.5	9.6	9.2	10.9
2004 4/.....	8.4	11.5	12.7	12.0	11.1	9.6	9.3	8.7	9.3	8.2
2003 4/.....	8.6	11.5	11.7	10.4	10.0	9.9	8.8	8.3	7.4	7.4
2002.....	8.7	10.6	11.5	10.2	9.8	9.4	7.4	7.7	5.9	6.8
2001.....	7.5	10.4	10.1	9.0	10.0	8.7	8.1	6.9	7.2	7.3
2000.....	7.6	9.8	9.6	9.3	9.0	8.7	7.9	7.6	7.8	7.2
1999.....	8.2	9.7	9.0	8.6	10.1	8.4	7.4	8.4	8.3	7.4

White, female

2006 4/.....	3.2	4.2	4.2	3.2	3.4	3.4	3.4	3.7	4.1	4.5
2005 4/.....	2.9	4.2	3.8	3.4	3.5	3.3	2.8	2.3	3.8	5.1
2004 4/.....	2.7	3.9	3.5	2.8	2.6	3.2	3.0	3.0	3.2	3.1
2003 4/.....	2.8	3.6	3.2	2.6	3.1	2.9	2.9	2.6	2.7	3.2
2002.....	2.8	3.1	2.5	2.5	2.3	2.5	3.2	2.6	2.5	2.2
2001.....	2.6	3.0	2.3	2.6	2.5	2.4	2.4	3.0	2.9	2.6
2000.....	2.5	2.5	2.2	2.4	2.2	2.4	2.6	2.6	2.7	2.9
1999.....	2.0	2.5	1.8	2.3	2.3	2.7	2.5	2.7	2.0	2.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to psychoactive substance use (F10-F19)

Black, both sexes

2006 4/.....	7.5	11.1	14.5	15.1	14.0	9.5	8.4	6.4	5.4	5.4
2005 4/.....	7.2	11.4	14.9	12.8	11.5	7.7	7.7	6.3	6.3	6.7
2004 4/.....	6.0	11.8	14.6	12.7	12.5	11.7	9.9	5.9	*	*
2003 4/.....	7.9	11.2	14.2	12.6	13.8	8.4	7.4	5.3	*	*
2002.....	8.6	11.9	14.2	14.3	13.5	9.6	8.3	5.8	*	*
2001.....	9.6	14.5	15.0	13.2	12.7	10.0	7.4	6.4	*	*
2000.....	9.0	14.6	13.1	12.9	12.4	11.0	7.5	6.9	*	*
1999.....	10.2	14.3	13.1	14.9	12.4	10.8	9.5	8.0	6.1	*

Black, male

2006 4/.....	10.4	15.9	23.8	27.1	26.3	18.0	17.8	10.3	*	*
2005 4/.....	10.0	17.3	24.6	22.4	21.1	15.2	14.9	13.4	14.0	*
2004 4/.....	9.1	16.7	24.6	23.8	24.9	23.7	20.1	13.6	*	*
2003 4/.....	10.9	17.9	25.4	20.8	27.2	15.5	14.1	11.4	*	*
2002.....	11.5	18.5	24.8	26.5	25.3	20.7	16.7	11.5	*	*
2001.....	14.0	23.1	25.3	24.2	24.8	20.6	15.1	13.9	*	*
2000.....	12.7	23.6	22.9	23.1	22.8	21.6	15.0	15.6	*	*
1999.....	15.0	22.3	21.6	27.6	22.9	21.6	19.0	16.6	*	*

Black, female

2006 4/.....	4.9	6.9	6.5	5.2	4.4	*	*	*	*	*
2005 4/.....	4.7	6.2	6.7	4.9	4.0	*	*	*	*	*
2004 4/.....	3.3	7.6	6.2	3.7	2.9	*	*	*	*	*
2003 4/.....	5.2	5.3	4.7	5.9	3.3	*	*	*	*	*
2002.....	6.1	6.1	5.3	4.3	4.2	*	*	*	*	*
2001.....	5.7	7.0	6.3	4.1	3.2	*	*	*	*	*
2000.....	5.6	6.8	4.8	4.6	4.2	*	*	*	*	*
1999.....	5.9	7.5	5.8	4.6	4.2	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to use of alcohol (F10)

All races 3/, both sexes

2006.....	3.9	5.5	6.7	6.5	6.2	5.0	3.9	3.3	3.2	2.3
2005.....	3.7	5.5	6.7	6.3	6.3	5.1	3.6	3.0	2.8	2.6
2004.....	3.5	5.4	6.3	6.2	5.9	5.3	4.4	3.4	2.9	1.9
2003.....	3.9	5.6	6.1	5.7	6.0	4.9	4.1	3.1	2.6	2.0
2002.....	3.9	5.3	6.0	5.9	5.8	5.1	4.1	3.5	2.4	2.0
2001.....	3.7	5.4	5.6	5.5	5.9	5.1	4.1	3.3	3.0	2.1
2000.....	3.6	5.1	5.2	5.5	5.3	5.0	4.1	3.4	2.9	2.5
1999.....	3.6	5.1	5.0	5.3	5.7	4.9	3.8	3.8	2.9	2.3

All races 3/, male

2006.....	5.8	8.5	10.7	10.9	10.3	8.6	6.8	5.9	5.7	4.3
2005.....	5.8	8.2	10.8	10.5	10.3	8.9	6.2	6.0	5.3	4.7
2004.....	5.6	8.5	10.4	10.5	10.3	8.9	8.0	6.4	5.3	3.6
2003.....	6.1	8.7	10.1	9.4	10.0	8.2	7.1	5.6	5.4	3.5
2002.....	6.1	8.3	10.1	9.9	10.0	9.0	6.8	6.4	4.5	4.4
2001.....	5.7	8.6	9.2	9.0	10.0	8.8	7.3	5.8	5.4	4.7
2000.....	5.7	8.3	8.6	9.2	9.1	8.8	7.2	6.1	5.9	5.2
1999.....	5.9	8.1	8.4	8.9	9.7	8.4	6.7	7.2	6.2	4.9

All races 3/, female

2006.....	1.9	2.7	2.8	2.3	2.4	1.9	1.5	1.5	1.6	1.4
2005.....	1.6	2.8	2.7	2.4	2.6	1.8	1.5	0.8	1.2	1.6
2004.....	1.5	2.5	2.4	2.2	2.0	2.2	1.5	1.3	1.4	1.2
2003.....	1.8	2.5	2.4	2.2	2.5	2.0	1.7	1.3	0.9	1.3
2002.....	1.9	2.3	2.1	2.0	1.9	1.6	1.9	1.4	1.1	1.0
2001.....	1.8	2.3	2.1	2.2	2.1	1.9	1.5	1.6	1.6	1.1
2000.....	1.6	2.0	1.9	2.1	1.8	1.8	1.6	1.5	1.2	1.5
1999.....	1.5	2.1	1.7	2.0	2.1	2.0	1.6	1.5	1.0	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to use of alcohol (F10)

White, both sexes

2006 4/.....	4.0	5.8	6.8	6.2	6.0	4.9	3.7	3.4	3.2	2.3
2005 4/.....	3.8	5.5	6.6	6.4	6.2	5.0	3.4	2.9	2.7	2.5
2004 4/.....	3.7	5.6	6.3	6.1	5.6	4.8	4.0	3.4	3.0	2.0
2003 4/.....	4.0	5.6	5.9	5.5	5.6	4.7	3.9	3.0	2.5	2.0
2002.....	4.0	5.3	5.8	5.4	5.1	4.7	3.8	3.4	2.3	2.0
2001.....	3.5	5.2	5.1	4.9	5.5	4.6	3.9	3.2	3.0	2.1
2000.....	3.5	4.7	4.9	5.1	4.7	4.5	3.9	3.2	2.9	2.5
1999.....	3.5	4.7	4.5	4.7	5.2	4.5	3.5	3.6	2.8	2.3

White, male

2006 4/.....	6.0	8.8	10.7	10.3	9.8	8.3	6.3	5.9	5.6	4.3
2005 4/.....	6.0	8.3	10.6	10.4	10.0	8.4	5.8	5.8	5.1	4.6
2004 4/.....	5.8	8.6	10.2	10.2	9.5	7.7	7.2	6.1	5.6	3.7
2003 4/.....	6.2	8.6	9.4	9.1	8.9	7.9	6.7	5.3	5.2	3.4
2002.....	6.2	8.2	9.8	9.0	8.9	8.2	6.0	6.3	4.2	4.1
2001.....	5.3	8.2	8.4	7.9	9.1	7.7	6.8	5.5	5.4	4.4
2000.....	5.5	7.7	8.0	8.3	8.0	7.8	6.7	5.6	5.7	5.1
1999.....	5.6	7.6	7.7	7.7	8.7	7.3	5.9	6.7	5.9	4.7

White, female

2006 4/.....	1.9	2.7	2.9	2.3	2.5	1.9	1.6	1.5	1.6	1.3
2005 4/.....	1.5	2.8	2.7	2.5	2.6	1.9	1.5	0.8	1.3	1.5
2004 4/.....	1.5	2.6	2.5	2.2	2.0	2.2	1.4	1.4	1.4	1.2
2003 4/.....	1.8	2.6	2.5	2.1	2.5	2.0	1.6	1.4	0.9	1.4
2002.....	1.8	2.4	2.0	2.0	1.7	1.7	1.9	1.4	1.1	1.1
2001.....	1.7	2.2	1.9	2.1	2.0	1.9	1.5	1.6	1.6	1.1
2000.....	1.5	1.8	1.8	1.9	1.6	1.7	1.6	1.6	1.2	1.5
1999.....	1.3	1.9	1.4	1.9	1.9	2.0	1.6	1.4	0.9	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to use of alcohol (F10)

Black, both sexes

2006 4/.....	3.3	4.9	7.0	9.6	9.1	7.2	5.6	3.9	*	*
2005 4/.....	3.4	6.0	7.7	7.6	8.9	7.0	6.2	4.2	*	*
2004 4/.....	2.7	5.6	7.8	8.7	9.6	10.0	9.0	4.6	*	*
2003 4/.....	3.6	6.0	8.4	8.4	11.1	7.0	6.2	4.8	*	*
2002.....	4.1	6.2	8.4	10.9	11.2	8.3	7.9	5.0	*	*
2001.....	5.2	7.9	10.3	11.0	11.0	9.6	6.3	5.2	*	*
2000.....	4.7	8.8	8.7	10.3	11.1	10.2	6.8	6.0	*	*
1999.....	4.9	7.9	8.9	11.6	11.3	9.6	8.4	7.1	*	*

Black, male

2006 4/.....	4.6	7.4	12.2	17.9	18.2	13.6	12.9	*	*	*
2005 4/.....	5.2	9.7	13.8	13.8	15.8	14.0	12.3	9.8	*	*
2004 4/.....	4.4	9.4	14.0	16.8	19.2	20.4	17.8	10.5	*	*
2003 4/.....	5.6	10.1	16.1	14.3	22.4	12.6	12.5	10.5	*	*
2002.....	5.9	10.4	14.6	20.7	21.0	18.0	16.4	9.6	*	*
2001.....	8.4	13.4	17.4	20.6	21.6	19.6	13.5	11.6	*	*
2000.....	7.3	14.5	15.7	18.9	21.3	20.3	14.4	14.6	*	*
1999.....	7.5	12.9	15.2	21.9	21.0	19.5	17.3	14.2	*	*

Black, female

2006 4/.....	2.1	2.7	2.6	2.8	*	*	*	*	*	*
2005 4/.....	1.7	2.8	2.6	2.5	3.5	*	*	*	*	*
2004 4/.....	*	2.4	2.6	*	*	*	*	*	*	*
2003 4/.....	1.9	2.4	1.8	3.6	*	*	*	*	*	*
2002.....	2.4	2.5	3.2	2.8	3.6	*	*	*	*	*
2001.....	2.4	3.1	4.2	3.1	*	*	*	*	*	*
2000.....	2.3	3.8	2.8	3.2	*	*	*	*	*	*
1999.....	2.5	3.7	3.6	3.2	3.7	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

All races 3/, both sexes

2006.....	2.2	2.5	2.8	1.9	1.6	1.8	2.4	2.7	3.4	4.4
2005.....	2.3	2.6	2.5	1.8	1.5	1.5	1.8	2.3	3.1	4.2
2004.....	2.1	2.5	2.3	1.5	1.2	1.4	1.7	2.0	2.4	2.7
2003.....	2.0	2.2	1.9	1.2	1.0	1.4	1.6	1.8	1.8	2.3
2002.....	2.0	2.0	1.6	1.1	0.9	1.0	1.2	1.2	1.4	1.7
2001.....	1.8	2.1	1.4	0.9	0.7	0.7	1.0	1.3	1.5	1.8
2000.....	1.8	1.9	1.3	0.9	0.8	0.8	1.0	1.4	1.6	1.6
1999.....	2.1	1.9	1.2	0.9	0.9	0.9	1.2	1.4	1.5	1.5

All races 3/, male

2006.....	3.0	3.3	4.0	2.9	2.4	2.3	3.2	3.4	5.0	7.1
2005.....	3.1	3.6	3.6	2.7	2.3	1.8	2.5	3.6	4.2	6.0
2004.....	2.8	3.3	3.4	2.3	1.9	1.9	2.1	2.7	3.5	4.6
2003.....	2.7	3.3	3.0	1.8	1.5	2.0	2.0	2.8	2.0	3.8
2002.....	2.8	2.9	2.6	1.6	1.2	1.3	1.3	1.4	1.7	2.6
2001.....	2.5	3.0	2.3	1.3	1.0	0.9	1.3	1.4	1.9	2.8
2000.....	2.4	2.8	2.1	1.2	1.0	0.9	1.1	1.9	2.0	2.1
1999.....	3.1	2.8	1.8	1.3	1.4	1.1	1.5	1.8	2.3	2.5

All races 3/, female

2006.....	1.5	1.8	1.6	1.0	1.0	1.4	1.7	2.3	2.4	3.1
2005.....	1.5	1.6	1.4	1.0	0.8	1.3	1.2	1.4	2.4	3.4
2004.....	1.3	1.8	1.3	0.7	0.6	0.9	1.4	1.5	1.7	1.8
2003.....	1.3	1.2	1.0	0.7	0.6	0.8	1.2	1.1	1.6	1.7
2002.....	1.3	1.1	0.7	0.6	0.6	0.7	1.2	1.1	1.3	1.2
2001.....	1.2	1.1	0.6	0.5	0.5	0.4	0.9	1.3	1.3	1.4
2000.....	1.3	1.0	0.6	0.6	0.6	0.6	1.0	1.0	1.4	1.3
1999.....	1.0	1.0	0.6	0.5	0.4	0.7	0.9	1.2	1.0	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

White, both sexes

2006 4/.....	2.1	2.1	2.3	1.5	1.3	1.8	2.4	2.8	3.6	4.5
2005 4/.....	2.2	2.3	1.9	1.5	1.5	1.7	1.9	2.5	3.1	4.4
2004 4/.....	1.9	2.1	1.7	1.2	1.1	1.4	1.8	2.1	2.5	2.7
2003 4/.....	1.8	1.9	1.5	0.9	0.9	1.4	1.7	2.0	1.9	2.5
2002.....	1.8	1.6	1.1	0.9	0.8	1.0	1.3	1.3	1.5	1.6
2001.....	1.5	1.5	1.1	0.8	0.7	0.7	1.1	1.4	1.5	1.9
2000.....	1.6	1.4	1.0	0.7	0.8	0.8	1.1	1.4	1.7	1.6
1999.....	1.6	1.3	0.9	0.6	0.9	0.9	1.2	1.5	1.6	1.6

White, male

2006 4/.....	2.8	2.7	3.2	2.2	1.8	2.1	3.1	3.4	5.3	7.2
2005 4/.....	3.0	3.2	2.8	2.1	2.1	1.9	2.6	3.8	4.1	6.2
2004 4/.....	2.7	2.9	2.5	1.8	1.6	1.8	2.2	2.6	3.7	4.5
2003 4/.....	2.5	2.9	2.3	1.3	1.2	2.0	2.1	3.0	2.2	4.0
2002.....	2.5	2.4	1.7	1.2	0.9	1.2	1.4	1.4	1.8	2.7
2001.....	2.2	2.2	1.7	1.1	0.9	0.9	1.3	1.3	1.9	2.9
2000.....	2.1	2.1	1.6	0.9	1.0	0.9	1.1	2.0	2.1	2.1
1999.....	2.6	2.1	1.3	0.9	1.4	1.1	1.5	1.7	2.4	2.7

White, female

2006 4/.....	1.3	1.5	1.3	0.9	0.8	1.5	1.8	2.3	2.5	3.1
2005 4/.....	1.4	1.4	1.1	0.9	0.9	1.4	1.3	1.5	2.5	3.6
2004 4/.....	1.2	1.3	1.0	0.6	0.6	1.0	1.6	1.7	1.8	2.0
2003 4/.....	1.0	1.0	0.7	0.5	0.6	0.9	1.3	1.2	1.8	1.8
2002.....	1.0	0.7	0.5	0.5	0.6	0.8	1.3	1.2	1.4	1.1
2001.....	0.9	0.8	0.4	0.4	0.5	0.5	0.9	1.4	1.3	1.5
2000.....	1.0	0.8	0.4	0.5	0.5	0.7	1.0	1.0	1.5	1.4
1999.....	0.7	0.6	0.4	0.4	*	0.7	1.0	1.3	1.1	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Mental and behavioral disorders due to other psychoactive substance use (F11-F19)

Black, both sexes

2006 4/.....	4.2	6.2	7.4	5.4	4.9	2.3	2.8	*	*	*
2005 4/.....	3.8	5.4	7.2	5.1	2.5	*	*	*	*	*
2004 4/.....	3.4	6.2	6.8	4.1	2.9	*	*	*	*	*
2003 4/.....	4.3	5.2	5.8	4.2	2.7	*	*	*	*	*
2002.....	4.6	5.7	5.8	3.4	2.3	*	*	*	*	*
2001.....	4.4	6.6	4.7	2.2	*	*	*	*	*	*
2000.....	4.3	5.8	4.4	2.6	*	*	*	*	*	*
1999.....	5.3	6.4	4.2	3.4	*	*	*	*	*	*

Black, male

2006 4/.....	5.8	8.5	11.6	9.2	8.1	*	*	*	*	*
2005 4/.....	4.8	7.7	10.8	8.6	5.3	*	*	*	*	*
2004 4/.....	4.6	7.3	10.6	7.0	5.6	*	*	*	*	*
2003 4/.....	5.3	7.7	9.3	6.5	4.8	*	*	*	*	*
2002.....	5.6	8.1	10.2	5.8	4.4	*	*	*	*	*
2001.....	5.6	9.7	7.9	3.6	*	*	*	*	*	*
2000.....	5.5	9.1	7.2	4.1	*	*	*	*	*	*
1999.....	7.5	9.3	6.4	5.8	*	*	*	*	*	*

Black, female

2006 4/.....	2.8	4.2	3.9	2.4	*	*	*	*	*	*
2005 4/.....	2.9	3.4	4.2	2.3	*	*	*	*	*	*
2004 4/.....	2.2	5.2	3.6	*	*	*	*	*	*	*
2003 4/.....	3.3	2.9	2.9	2.3	*	*	*	*	*	*
2002.....	3.7	3.7	2.1	*	*	*	*	*	*	*
2001.....	3.3	3.9	2.0	*	*	*	*	*	*	*
2000.....	3.3	3.0	2.0	*	*	*	*	*	*	*
1999.....	3.4	3.8	2.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Schizophrenia, schizotypal and delusional disorders (F20-F29)										
All races 3/, both sexes										
2006.....	*	0.1	0.1	0.1	0.3	0.4	0.6	1.0	1.4	2.2
2005.....	*	0.1	*	0.2	0.3	0.3	0.6	1.0	1.6	2.4
2004.....	0.1	0.1	0.2	0.2	0.2	0.4	0.6	0.9	1.3	2.7
2003.....	*	0.1	0.2	*	0.2	0.3	0.8	0.9	1.5	2.7
2002.....	0.1	0.1	*	0.2	*	0.3	0.7	1.0	1.3	3.3
2001.....	*	0.1	*	0.1	0.2	0.3	0.4	1.1	1.3	2.7
2000.....	*	*	*	*	0.2	0.3	0.7	0.9	1.7	3.3
1999.....	*	*	*	*	0.2	0.3	0.6	1.0	2.1	4.9
All races 3/, male										
2006.....	*	*	*	*	0.3	0.4	0.6	1.0	1.2	1.5
2005.....	*	*	*	0.2	0.4	*	*	0.8	1.5	1.8
2004.....	*	0.2	*	0.2	*	*	0.7	0.9	1.1	2.3
2003.....	*	*	0.2	*	*	*	0.6	0.6	1.1	1.8
2002.....	*	*	*	*	*	*	0.7	0.9	*	1.9
2001.....	*	*	*	*	*	0.5	0.5	0.9	1.1	1.5
2000.....	*	*	*	*	*	*	0.7	0.7	1.6	2.4
1999.....	*	*	*	*	*	*	0.7	1.0	1.4	4.6
All races 3/, female										
2006.....	*	*	*	*	*	*	0.6	1.0	1.6	2.5
2005.....	*	*	*	*	*	0.4	0.7	1.2	1.6	2.7
2004.....	*	*	*	*	*	*	0.6	0.9	1.4	2.8
2003.....	*	*	*	*	*	*	0.9	1.0	1.7	3.0
2002.....	*	*	*	*	*	*	0.6	1.1	1.6	3.9
2001.....	*	*	*	*	*	*	*	1.2	1.4	3.1
2000.....	*	*	*	*	*	*	0.7	1.0	1.8	3.8
1999.....	*	*	*	*	*	*	0.5	1.0	2.5	5.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Schizophrenia, schizotypal and delusional disorders (F20-F29)

White, both sexes

2006 4/.....	*	*	*	*	0.2	0.4	0.6	1.0	1.5	2.3
2005 4/.....	*	*	*	0.2	0.3	0.3	0.6	1.0	1.5	2.5
2004 4/.....	*	0.1	0.1	0.2	0.2	0.3	0.6	0.9	1.3	2.6
2003 4/.....	*	*	0.1	*	*	0.4	0.8	0.9	1.4	2.8
2002.....	*	*	*	0.2	*	0.3	0.6	1.0	1.3	3.4
2001.....	*	*	*	*	0.2	0.3	0.4	1.1	1.3	2.8
2000.....	*	*	*	*	*	0.4	0.7	0.9	1.8	3.4
1999.....	*	*	*	*	*	0.3	0.6	1.0	2.1	5.0

White, male

2006 4/.....	*	*	*	*	*	0.5	0.6	1.1	1.1	1.6
2005 4/.....	*	*	*	*	*	*	*	0.7	1.4	1.9
2004 4/.....	*	*	*	*	*	*	0.7	0.8	1.1	1.9
2003 4/.....	*	*	*	*	*	*	0.6	*	1.1	1.9
2002.....	*	*	*	*	*	*	0.6	0.8	*	1.9
2001.....	*	*	*	*	*	*	*	0.9	1.1	*
2000.....	*	*	*	*	*	*	0.8	0.8	1.7	2.4
1999.....	*	*	*	*	*	*	0.7	0.8	1.4	4.7

White, female

2006 4/.....	*	*	*	*	*	*	0.6	0.9	1.7	2.7
2005 4/.....	*	*	*	*	*	*	0.7	1.2	1.6	2.8
2004 4/.....	*	*	*	*	*	*	0.6	1.0	1.5	3.0
2003 4/.....	*	*	*	*	*	*	0.9	1.1	1.6	3.2
2002.....	*	*	*	*	*	*	0.7	1.1	1.6	4.0
2001.....	*	*	*	*	*	*	*	1.2	1.4	3.3
2000.....	*	*	*	*	*	*	0.7	1.0	1.9	3.8
1999.....	*	*	*	*	*	*	0.5	1.0	2.5	5.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mood (affective) disorders (F30-F39)										
All races 3/, both sexes										
2006.....	*	*	*	*	0.1	*	0.5	1.2	2.5	6.5
2005.....	*	*	0.1	*	*	*	0.5	1.2	3.0	7.4
2004.....	*	*	*	0.1	*	0.2	0.7	1.4	2.6	8.4
2003.....	*	*	*	0.1	*	0.2	0.5	1.5	2.9	10.5
2002.....	*	*	*	*	0.2	0.2	0.7	1.3	3.5	9.1
2001.....	0.1	*	*	*	*	0.4	0.6	1.4	3.2	9.6
2000.....	*	*	*	*	*	0.3	0.5	1.2	2.9	9.8
1999.....	*	*	*	*	*	0.3	0.6	1.6	3.4	9.4
All races 3/, male										
2006.....	*	*	*	*	*	*	*	0.9	1.7	5.4
2005.....	*	*	*	*	*	*	*	1.1	2.4	6.0
2004.....	*	*	*	*	*	*	0.7	1.0	2.1	7.2
2003.....	*	*	*	*	*	*	*	1.5	2.4	8.0
2002.....	*	*	*	*	*	*	0.5	0.9	2.6	6.6
2001.....	*	*	*	*	*	*	0.6	1.3	2.4	8.0
2000.....	*	*	*	*	*	*	0.5	0.9	2.3	9.1
1999.....	*	*	*	*	*	*	*	1.6	2.7	8.8
All races 3/, female										
2006.....	*	*	*	*	*	*	0.5	1.5	3.0	7.0
2005.....	*	*	*	*	*	*	0.6	1.3	3.4	8.1
2004.....	*	*	*	*	*	*	0.6	1.6	3.0	9.0
2003.....	*	*	*	*	*	*	0.5	1.6	3.3	11.6
2002.....	*	*	*	*	*	*	0.7	1.6	4.0	10.1
2001.....	*	*	*	*	*	*	0.6	1.5	3.7	10.3
2000.....	*	*	*	*	*	*	0.5	1.5	3.2	10.1
1999.....	*	*	*	*	*	*	0.6	1.7	3.9	9.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Mood (affective) disorders (F30-F39)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	0.5	1.2	2.7	6.9
2005 4/.....	*	*	0.1	*	*	*	0.5	1.3	3.3	8.2
2004 4/.....	*	*	*	*	*	0.3	0.7	1.4	2.8	9.2
2003 4/.....	*	*	*	*	*	0.2	0.5	1.6	3.1	11.3
2002.....	*	*	*	*	0.2	0.2	0.7	1.4	3.6	9.8
2001.....	*	*	*	*	*	0.4	0.6	1.6	3.4	10.1
2000.....	*	*	*	*	*	0.3	0.6	1.3	3.0	10.6
1999.....	*	*	*	*	*	0.3	0.6	1.8	3.7	10.2
White, male										
2006 4/.....	*	*	*	*	*	*	*	0.8	1.9	5.8
2005 4/.....	*	*	*	*	*	*	*	1.1	2.6	6.6
2004 4/.....	*	*	*	*	*	*	0.8	1.1	2.2	7.7
2003 4/.....	*	*	*	*	*	*	*	1.5	2.6	8.6
2002.....	*	*	*	*	*	*	*	0.9	2.7	7.2
2001.....	*	*	*	*	*	*	0.7	1.4	2.6	8.3
2000.....	*	*	*	*	*	*	*	0.9	2.4	10.1
1999.....	*	*	*	*	*	*	*	1.7	2.9	9.5
White, female										
2006 4/.....	*	*	*	*	*	*	0.6	1.5	3.3	7.3
2005 4/.....	*	*	*	*	*	*	0.6	1.5	3.7	8.9
2004 4/.....	*	*	*	*	*	*	0.6	1.7	3.1	9.8
2003 4/.....	*	*	*	*	*	*	0.5	1.6	3.5	12.5
2002.....	*	*	*	*	*	*	0.8	1.8	4.1	10.9
2001.....	*	*	*	*	*	*	0.6	1.6	4.0	10.9
2000.....	*	*	*	*	*	*	0.6	1.6	3.4	10.7
1999.....	*	*	*	*	*	*	0.7	1.9	4.2	10.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)

All races 3/, both sexes

2006.....	*	*	*	0.1	*	*	0.3	0.4	1.0	2.4
2005.....	*	*	*	*	*	*	0.3	0.4	1.0	2.8
2004.....	*	*	*	*	*	*	*	0.5	1.0	3.9
2003.....	*	0.1	*	*	*	*	0.3	0.7	1.1	4.4
2002.....	*	*	*	*	*	*	0.3	0.4	1.2	4.4
2001.....	*	*	*	*	*	*	*	0.5	1.0	4.1
2000.....	*	*	*	*	*	*	*	0.4	1.2	3.9
1999.....	*	*	*	*	*	*	0.4	0.4	1.0	4.0

All races 3/, male

2006.....	*	*	*	*	*	*	*	0.7	1.3	2.5
2005.....	*	*	*	*	*	*	*	*	1.1	2.5
2004.....	*	*	*	*	*	*	*	*	1.1	3.7
2003.....	*	*	*	*	*	*	*	0.7	1.3	4.2
2002.....	*	*	*	*	*	*	*	*	1.6	3.6
2001.....	*	*	*	*	*	*	*	*	*	3.5
2000.....	*	*	*	*	*	*	*	*	1.3	3.3
1999.....	*	*	*	*	*	*	*	*	*	2.5

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	0.9	2.3
2005.....	*	*	*	*	*	*	*	*	0.9	2.9
2004.....	*	*	*	*	*	*	*	0.6	1.0	4.0
2003.....	*	*	*	*	*	*	*	0.7	1.0	4.5
2002.....	*	*	*	*	*	*	*	*	1.0	4.7
2001.....	*	*	*	*	*	*	*	0.5	1.0	4.4
2000.....	*	*	*	*	*	*	*	0.5	1.1	4.1
1999.....	*	*	*	*	*	*	*	*	1.1	4.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified mental and behavioral disorders (F50-F69,F80-F99)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	0.3	0.4	1.0	2.4
2005 4/.....	*	*	*	*	*	*	*	0.4	0.9	2.7
2004 4/.....	*	*	*	*	*	*	*	0.5	1.1	4.0
2003 4/.....	*	*	*	*	*	*	0.3	0.7	1.1	4.4
2002.....	*	*	*	*	*	*	*	0.4	1.3	4.4
2001.....	*	*	*	*	*	*	*	0.5	1.0	4.4
2000.....	*	*	*	*	*	*	*	0.4	1.2	4.0
1999.....	*	*	*	*	*	*	0.4	0.5	1.1	4.3
White, male										
2006 4/.....	*	*	*	*	*	*	*	*	1.2	2.4
2005 4/.....	*	*	*	*	*	*	*	*	1.0	2.4
2004 4/.....	*	*	*	*	*	*	*	*	1.2	3.5
2003 4/.....	*	*	*	*	*	*	*	0.7	1.3	4.2
2002.....	*	*	*	*	*	*	*	*	1.6	3.7
2001.....	*	*	*	*	*	*	*	*	*	3.7
2000.....	*	*	*	*	*	*	*	*	1.3	3.2
1999.....	*	*	*	*	*	*	*	*	*	2.7
White, female										
2006 4/.....	*	*	*	*	*	*	*	*	0.8	2.4
2005 4/.....	*	*	*	*	*	*	*	*	0.9	2.8
2004 4/.....	*	*	*	*	*	*	*	0.6	1.0	4.2
2003 4/.....	*	*	*	*	*	*	*	0.7	1.0	4.5
2002.....	*	*	*	*	*	*	*	*	1.0	4.7
2001.....	*	*	*	*	*	*	*	0.5	1.0	4.7
2000.....	*	*	*	*	*	*	*	*	1.1	4.3
1999.....	*	*	*	*	*	*	*	*	1.2	5.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

VI. Diseases of the nervous system (G00-G98)

All races 3/, both sexes

2006.....	5.4	7.5	11.5	17.0	26.5	43.2	90.3	200.1	424.8	1,062.1
2005.....	5.2	8.0	11.8	17.1	26.2	42.9	88.7	198.0	429.6	1,069.1
2004.....	4.9	7.6	11.0	16.3	25.0	42.3	83.4	190.8	405.4	1,014.7
2003.....	5.0	7.7	11.0	16.2	25.7	43.0	88.3	193.1	400.0	1,006.7
2002.....	5.1	7.8	10.5	15.7	24.9	42.7	83.2	185.2	388.8	951.0
2001.....	5.0	7.3	10.3	15.7	24.0	41.0	81.1	178.4	375.0	906.2
2000.....	5.0	7.3	11.0	15.6	23.8	40.2	79.5	172.9	361.9	864.9
1999.....	4.6	7.0	10.3	14.4	22.1	39.3	76.3	163.6	340.4	790.8

All races 3/, male

2006.....	5.9	8.1	12.4	18.1	28.9	48.4	105.8	230.1	460.0	954.3
2005.....	5.9	8.6	12.6	18.4	27.9	47.9	100.9	223.7	468.1	943.2
2004.....	5.6	8.4	12.0	17.5	26.9	47.0	95.4	211.4	441.6	910.7
2003.....	6.0	8.6	12.2	17.5	27.9	47.2	100.6	218.6	436.2	907.9
2002.....	5.7	8.5	11.2	16.5	27.4	48.2	95.7	210.0	422.2	867.4
2001.....	5.5	7.7	10.9	16.2	25.7	47.4	92.9	206.0	421.1	861.6
2000.....	5.7	8.3	11.8	16.5	25.9	46.5	92.8	198.3	407.5	826.2
1999.....	5.1	7.2	11.0	15.9	24.4	44.9	89.1	193.7	394.2	772.7

All races 3/, female

2006.....	4.8	6.8	10.7	15.9	24.4	38.7	77.8	178.2	402.8	1,112.4
2005.....	4.5	7.4	11.0	15.8	24.7	38.6	78.8	179.3	405.8	1,126.9
2004.....	4.2	6.8	10.1	15.2	23.3	38.3	73.8	176.0	383.1	1,061.5
2003.....	4.1	6.9	9.9	15.0	23.7	39.4	78.4	175.0	377.9	1,050.4
2002.....	4.4	7.0	9.9	14.9	22.6	37.8	73.2	167.7	368.5	987.3
2001.....	4.5	6.9	9.7	15.2	22.5	35.5	71.7	159.1	347.4	924.9
2000.....	4.4	6.4	10.3	14.8	21.8	34.9	69.0	155.1	335.0	880.7
1999.....	4.1	6.7	9.6	13.0	20.1	34.6	66.3	142.7	308.9	798.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

VI. Diseases of the nervous system (G00-G98)

White, both sexes

2006 4/.....	5.2	7.2	11.4	16.9	26.8	44.2	93.8	208.7	443.0	1,103.5
2005 4/.....	5.0	7.8	11.7	17.2	26.2	44.0	91.4	205.3	445.0	1,109.0
2004 4/.....	4.8	7.4	10.9	16.3	25.3	43.2	86.9	197.4	419.4	1,051.0
2003 4/.....	4.9	7.4	10.7	16.3	25.9	44.0	91.4	200.1	416.1	1,047.3
2002.....	4.9	7.5	10.2	15.7	25.4	43.8	85.9	192.1	405.4	989.7
2001.....	4.8	7.0	10.2	15.5	24.3	42.0	84.1	184.4	389.0	942.3
2000.....	4.8	7.0	10.9	15.6	24.1	41.2	81.7	179.1	375.6	900.8
1999.....	4.4	6.6	10.1	14.3	22.2	40.1	78.1	169.0	353.7	825.1

White, male

2006 4/.....	6.0	7.8	12.4	17.9	29.1	49.3	109.6	239.9	479.7	996.5
2005 4/.....	5.8	8.5	12.6	18.3	27.8	48.7	104.1	231.0	485.1	980.2
2004 4/.....	5.6	8.2	11.8	17.4	27.3	48.0	99.2	218.0	459.2	945.7
2003 4/.....	5.9	8.3	11.9	17.6	27.7	48.0	104.2	226.5	452.6	944.4
2002.....	5.5	8.2	10.7	16.4	28.0	49.0	98.4	217.7	438.9	904.3
2001.....	5.3	7.5	10.7	16.1	26.0	48.4	95.9	213.3	436.5	896.4
2000.....	5.3	8.1	11.9	16.3	26.0	47.8	95.3	205.0	422.6	859.9
1999.....	4.7	6.9	10.7	15.9	24.4	45.7	91.7	200.0	411.2	806.2

White, female

2006 4/.....	4.5	6.6	10.3	15.9	24.7	39.7	80.6	185.4	419.9	1,153.8
2005 4/.....	4.2	7.1	10.9	16.0	24.6	39.7	80.9	186.3	419.9	1,168.3
2004 4/.....	4.0	6.5	10.0	15.2	23.5	39.0	76.7	182.3	394.7	1,098.4
2003 4/.....	3.8	6.6	9.6	15.0	24.2	40.4	81.0	181.0	393.5	1,092.9
2002.....	4.3	6.9	9.7	15.1	22.9	39.2	75.8	173.7	384.8	1,026.8
2001.....	4.4	6.6	9.7	15.0	22.7	36.3	74.5	163.9	360.3	961.5
2000.....	4.3	5.9	10.0	14.9	22.3	35.4	70.7	160.9	347.7	917.4
1999.....	4.1	6.3	9.4	12.8	20.1	35.3	67.3	147.1	319.8	832.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

VI. Diseases of the nervous system (G00-G98)

Black, both sexes

2006 4/.....	8.1	11.1	16.1	23.6	31.5	44.0	84.1	163.2	317.6	779.1
2005 4/.....	7.7	11.4	16.1	20.9	33.8	46.7	88.4	166.0	347.6	811.1
2004 4/.....	6.9	11.1	14.8	21.4	30.2	45.2	72.9	166.6	327.0	776.9
2003 4/.....	7.3	11.9	16.1	20.5	30.8	46.5	78.6	159.4	295.2	699.3
2002.....	7.7	11.7	16.0	18.8	28.2	43.4	76.3	150.1	269.2	655.4
2001.....	7.4	10.7	14.9	21.3	27.3	42.2	69.6	152.0	273.6	624.9
2000.....	7.9	12.1	15.4	20.0	26.2	41.6	73.1	136.1	259.8	561.2
1999.....	7.1	11.7	15.6	18.6	26.3	40.1	73.7	131.6	225.9	485.3

Black, male

2006 4/.....	7.8	12.5	16.4	26.2	35.5	51.3	96.9	185.3	330.6	611.2
2005 4/.....	7.9	11.8	16.2	24.5	36.7	55.4	97.4	189.5	371.5	686.0
2004 4/.....	7.5	12.2	17.0	23.8	31.8	49.2	83.8	183.3	327.8	661.6
2003 4/.....	8.1	13.0	17.9	21.5	37.0	53.9	89.7	180.3	324.8	620.4
2002.....	8.5	13.5	17.1	20.5	30.7	52.1	93.8	168.1	303.4	570.4
2001.....	8.5	11.5	16.5	22.1	29.5	49.3	81.8	170.5	302.6	593.4
2000.....	9.6	12.2	15.3	23.9	30.6	46.6	86.9	159.7	285.8	537.4
1999.....	8.9	12.3	18.1	19.3	29.7	46.2	81.1	152.4	231.1	481.6

Black, female

2006 4/.....	8.4	9.9	15.9	21.6	28.4	38.6	75.2	150.0	310.7	847.7
2005 4/.....	7.6	10.9	16.1	18.0	31.5	40.2	82.2	152.0	335.0	861.6
2004 4/.....	6.3	10.3	13.0	19.6	29.0	42.3	65.5	156.6	326.6	822.6
2003 4/.....	6.6	10.9	14.7	19.7	25.9	40.9	71.1	147.0	279.7	730.0
2002.....	7.0	10.2	15.0	17.4	26.2	37.0	64.5	139.4	251.5	688.0
2001.....	6.4	10.0	13.6	20.7	25.6	37.0	61.4	140.9	258.8	636.5
2000.....	6.4	11.9	15.5	16.9	22.7	38.0	63.9	121.9	246.7	570.1
1999.....	5.6	11.2	13.5	18.0	23.6	35.6	68.9	119.0	223.4	486.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningitis (G00,G03)										
All races 3/, both sexes										
2006.....	0.2	0.2	0.3	0.2	0.2	0.3	0.5	0.6	0.8	0.9
2005.....	0.2	0.3	0.3	0.4	0.3	0.4	0.4	0.5	0.6	0.9
2004.....	0.2	0.3	0.3	0.3	0.3	0.5	0.5	0.6	0.7	0.9
2003.....	0.2	0.3	0.3	0.3	0.4	0.6	0.5	0.7	0.6	1.1
2002.....	0.2	0.3	0.3	0.3	0.3	0.5	0.7	0.6	0.7	1.2
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.5	0.6	1.0	0.9
2000.....	0.2	0.2	0.3	0.3	0.4	0.5	0.7	0.9	1.1	1.2
1999.....	0.2	0.3	0.3	0.3	0.5	0.5	0.7	0.9	1.2	1.5
All races 3/, male										
2006.....	*	0.3	0.3	0.2	*	*	0.5	*	0.9	*
2005.....	0.2	0.3	0.3	0.4	0.3	*	*	*	*	*
2004.....	0.3	0.3	0.4	0.3	0.3	*	*	*	*	*
2003.....	*	0.3	0.4	0.4	0.5	0.6	*	0.8	*	*
2002.....	0.2	0.4	0.3	0.4	*	0.6	0.9	0.7	1.0	*
2001.....	*	0.3	0.3	0.4	0.6	0.5	0.7	0.9	1.1	*
2000.....	0.2	0.3	0.4	0.4	0.5	0.6	0.9	1.3	1.1	*
1999.....	0.2	0.4	0.4	0.4	0.4	0.6	0.8	0.9	1.3	1.7
All races 3/, female										
2006.....	0.2	0.2	0.2	0.3	*	0.4	0.4	0.5	0.8	0.8
2005.....	0.2	0.2	0.3	0.4	0.3	*	*	0.5	0.6	0.9
2004.....	*	*	0.3	0.3	0.3	0.6	0.6	0.6	0.6	0.7
2003.....	*	0.2	0.2	0.3	0.3	0.6	0.4	0.7	0.6	1.0
2002.....	*	0.2	0.2	*	0.3	0.4	0.6	0.6	*	1.1
2001.....	*	0.2	0.2	0.4	0.4	0.4	0.4	0.5	0.9	0.8
2000.....	*	0.2	0.3	*	*	0.4	0.5	0.6	1.1	1.0
1999.....	*	0.2	*	*	0.5	0.4	0.7	0.9	1.1	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Meningitis (G00,G03)										
White, both sexes										
2006 4/.....	0.2	0.2	0.3	0.2	0.2	0.3	0.5	0.5	0.8	0.9
2005 4/.....	0.2	0.3	0.2	0.4	0.3	0.3	0.4	0.5	0.6	0.9
2004 4/.....	0.2	0.2	0.2	0.2	0.3	0.4	0.4	0.6	0.6	0.8
2003 4/.....	0.1	0.2	0.2	0.3	0.3	0.5	0.4	0.7	0.6	1.0
2002.....	0.1	0.2	0.2	0.3	0.3	0.5	0.7	0.6	0.6	1.2
2001.....	0.1	0.2	0.2	0.3	0.5	0.4	0.6	0.6	0.9	0.8
2000.....	0.1	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1	1.0
1999.....	0.1	0.2	0.2	0.3	0.4	0.5	0.6	0.7	1.1	1.3
White, male										
2006 4/.....	*	0.2	0.3	*	*	*	*	*	*	*
2005 4/.....	*	0.3	0.3	0.4	*	*	*	*	*	*
2004 4/.....	0.2	0.3	0.3	*	*	*	*	*	*	*
2003 4/.....	*	0.3	0.3	0.3	0.4	0.6	*	0.9	*	*
2002.....	*	0.3	*	0.3	*	0.6	0.8	*	*	*
2001.....	*	0.2	0.3	0.4	0.5	*	0.7	0.8	*	*
2000.....	*	0.3	0.3	0.4	0.5	*	0.8	1.2	*	*
1999.....	*	0.3	*	0.4	*	0.5	0.7	*	1.3	*
White, female										
2006 4/.....	*	*	0.2	*	*	0.4	*	*	0.7	0.8
2005 4/.....	*	*	*	0.3	*	*	*	*	*	0.9
2004 4/.....	*	*	*	*	*	0.5	0.5	0.6	*	*
2003 4/.....	*	*	*	*	*	0.5	*	0.6	*	0.9
2002.....	*	*	*	*	*	*	0.5	0.5	*	1.2
2001.....	*	*	*	*	0.4	*	*	*	0.8	0.8
2000.....	*	*	*	*	*	*	*	0.6	1.0	1.0
1999.....	*	*	*	*	0.5	*	0.6	0.8	1.0	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other inflammatory diseases of central nervous system (G04-G09)

All races 3/, both sexes

2006.....	0.1	0.2	0.2	0.3	0.4	0.5	0.8	1.0	1.0	1.2
2005.....	0.1	0.2	0.3	0.2	0.4	0.8	0.5	0.8	1.0	1.0
2004.....	0.2	0.2	0.2	0.3	0.4	0.4	0.8	1.0	1.3	1.1
2003.....	0.2	0.2	0.2	0.3	0.4	0.7	0.7	1.0	1.5	1.1
2002.....	0.2	0.2	0.2	0.3	0.4	0.5	0.8	1.1	1.4	1.2
2001.....	0.1	0.2	0.2	0.3	0.4	0.5	0.8	0.9	1.1	1.0
2000.....	0.2	0.1	0.2	0.3	0.3	0.5	0.8	1.1	1.1	1.0
1999.....	0.1	0.2	0.2	0.3	0.3	0.5	0.7	0.7	1.0	1.1

All races 3/, male

2006.....	*	0.2	0.3	0.4	0.5	0.6	1.0	1.3	1.3	1.5
2005.....	0.2	0.3	0.4	0.3	0.5	1.0	0.6	1.0	1.2	1.3
2004.....	0.2	0.2	0.2	0.4	0.5	0.6	1.0	1.2	1.8	1.5
2003.....	0.2	0.3	0.3	0.4	0.5	0.8	0.9	1.3	1.7	*
2002.....	0.2	0.3	0.3	0.4	0.5	0.8	1.0	1.3	2.0	*
2001.....	*	0.2	0.3	0.3	0.4	0.7	0.8	1.1	1.3	1.8
2000.....	0.2	*	0.3	0.3	0.4	0.6	1.0	1.6	1.4	*
1999.....	0.2	*	*	0.5	*	0.6	0.6	0.9	1.6	*

All races 3/, female

2006.....	*	*	0.2	0.2	0.4	0.4	0.7	0.8	0.8	1.1
2005.....	*	0.2	0.2	*	0.3	0.6	*	0.7	0.8	0.9
2004.....	*	*	0.2	*	0.3	*	0.6	0.8	0.9	1.0
2003.....	*	*	*	0.3	0.3	0.6	0.5	0.8	1.4	0.9
2002.....	*	0.2	0.2	0.3	0.3	*	0.6	0.9	1.1	1.1
2001.....	*	0.2	*	0.3	0.4	0.4	0.8	0.8	0.9	0.7
2000.....	*	*	*	0.3	*	0.5	0.6	0.8	0.9	0.9
1999.....	*	*	*	*	0.4	0.4	0.8	0.6	0.7	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other inflammatory diseases of central nervous system (G04-G09)

White, both sexes

2006 4/.....	0.1	0.2	0.2	0.3	0.4	0.5	0.8	1.0	1.0	1.2
2005 4/.....	0.1	0.2	0.2	0.2	0.3	0.7	0.4	0.9	1.0	1.0
2004 4/.....	0.1	0.2	0.2	0.3	0.4	0.4	0.8	1.0	1.2	1.1
2003 4/.....	0.2	0.2	0.2	0.3	0.4	0.7	0.7	1.0	1.5	1.1
2002.....	0.1	0.2	0.2	0.3	0.4	0.5	0.7	1.0	1.4	1.3
2001.....	0.1	0.2	0.2	0.3	0.4	0.5	0.7	0.9	1.0	1.0
2000.....	0.2	*	0.2	0.3	0.4	0.5	0.7	1.1	1.0	1.0
1999.....	0.1	0.2	0.2	0.3	0.3	0.5	0.6	0.7	1.1	1.1

White, male

2006 4/.....	*	*	0.3	0.5	0.5	0.5	1.0	1.3	1.3	1.5
2005 4/.....	*	0.2	0.3	*	0.4	1.0	*	1.0	1.3	*
2004 4/.....	*	0.2	0.2	0.4	0.6	0.6	1.0	1.2	1.8	*
2003 4/.....	0.2	0.2	0.3	0.4	0.5	0.7	0.9	1.4	1.7	*
2002.....	*	0.2	*	0.4	0.5	0.7	0.9	1.3	2.0	*
2001.....	*	*	*	0.4	*	0.6	0.8	0.9	1.3	1.8
2000.....	0.2	*	0.3	*	0.5	0.6	1.0	1.7	1.4	*
1999.....	*	*	*	0.4	*	0.5	0.6	1.0	1.6	*

White, female

2006 4/.....	*	*	*	*	*	*	0.7	0.8	0.7	1.1
2005 4/.....	*	*	*	*	*	0.5	*	0.8	0.8	0.9
2004 4/.....	*	*	*	*	*	*	0.6	0.8	0.9	1.0
2003 4/.....	*	*	*	*	*	0.7	0.5	0.8	1.4	1.0
2002.....	*	*	*	*	*	*	0.5	0.9	1.0	1.2
2001.....	*	*	*	*	*	*	0.7	0.8	0.9	*
2000.....	*	*	*	*	*	*	0.5	0.7	0.8	0.9
1999.....	*	*	*	*	*	*	0.7	0.5	0.8	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)										
All races 3/, both sexes										
2006.....	*	*	0.3	0.7	2.1	6.0	19.8	50.1	95.4	144.8
2005.....	*	0.1	0.2	0.8	2.3	6.8	20.3	50.2	98.7	143.7
2004.....	*	*	0.2	0.6	2.0	6.4	18.5	47.6	94.0	135.8
2003.....	*	*	0.2	0.7	2.0	6.4	19.9	49.3	93.3	138.2
2002.....	*	*	0.2	0.6	1.9	6.5	18.4	45.4	89.8	135.2
2001.....	*	*	0.2	0.7	1.8	5.8	18.1	46.4	90.9	134.2
2000.....	*	*	0.2	0.6	1.7	6.6	16.7	44.8	87.6	131.9
1999.....	*	*	0.2	0.6	1.6	5.6	16.7	41.8	82.8	124.4
All races 3/, male										
2006.....	*	*	0.4	0.9	2.8	8.5	29.0	76.0	145.6	231.7
2005.....	*	*	0.3	0.9	3.2	9.6	30.0	75.0	153.6	228.6
2004.....	*	*	0.4	0.9	2.6	9.1	27.3	70.6	147.1	220.0
2003.....	*	*	0.2	0.8	2.7	8.9	29.2	74.2	140.7	222.4
2002.....	*	*	0.3	0.8	2.5	9.8	27.3	67.2	136.6	220.1
2001.....	*	*	0.3	1.0	2.7	8.2	26.5	68.8	141.7	224.4
2000.....	*	*	*	0.8	2.2	9.0	24.1	67.4	135.0	222.3
1999.....	*	*	0.3	0.7	2.3	7.7	24.8	64.4	131.3	205.6
All races 3/, female										
2006.....	*	*	*	0.4	1.3	3.8	12.2	31.0	64.0	104.1
2005.....	*	*	*	0.6	1.5	4.3	12.5	32.3	64.6	104.7
2004.....	*	*	*	0.4	1.4	4.1	11.4	31.0	61.3	97.9
2003.....	*	*	0.2	0.6	1.4	4.3	12.4	31.5	64.3	101.0
2002.....	*	*	*	0.5	1.4	3.8	11.3	30.0	61.3	98.4
2001.....	*	*	*	0.4	1.0	3.7	11.5	30.7	60.4	96.4
2000.....	*	*	*	0.4	1.3	4.4	10.9	29.1	59.5	95.0
1999.....	*	*	*	0.4	0.9	3.9	10.4	26.0	54.4	91.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)										
White, both sexes										
2006 4/.....	*	*	0.2	0.7	2.1	6.3	21.2	53.5	101.9	154.1
2005 4/.....	*	0.1	0.2	0.8	2.4	7.1	21.8	53.7	104.4	152.9
2004 4/.....	*	*	0.2	0.7	2.1	6.8	20.2	50.6	99.9	143.1
2003 4/.....	*	*	0.2	0.7	2.0	6.8	21.2	52.4	99.0	146.0
2002.....	*	*	0.2	0.6	2.1	6.9	19.6	48.3	95.7	143.3
2001.....	*	*	0.2	0.7	1.9	6.2	19.4	49.6	96.4	141.8
2000.....	*	*	0.2	0.6	1.9	7.1	17.8	47.5	92.5	139.5
1999.....	*	*	0.2	0.6	1.6	6.0	17.7	44.1	87.4	131.2
White, male										
2006 4/.....	*	*	0.4	1.0	2.9	8.8	30.8	80.5	154.3	247.3
2005 4/.....	*	*	0.3	1.0	3.3	10.0	31.9	79.5	162.2	242.7
2004 4/.....	*	*	0.4	1.0	2.7	9.6	29.4	74.7	156.1	231.3
2003 4/.....	*	*	*	0.8	2.7	9.4	30.9	78.4	148.9	233.3
2002.....	*	*	0.3	0.8	2.7	10.2	28.8	71.2	145.0	232.7
2001.....	*	*	0.3	1.0	3.0	8.8	28.1	73.4	149.6	237.6
2000.....	*	*	*	0.9	2.5	9.6	25.5	70.7	141.5	235.3
1999.....	*	*	0.3	0.7	2.4	8.1	26.4	67.7	138.6	216.7
White, female										
2006 4/.....	*	*	*	0.5	1.4	4.0	13.2	33.4	68.8	110.3
2005 4/.....	*	*	*	0.7	1.6	4.6	13.4	34.6	68.2	111.5
2004 4/.....	*	*	*	0.4	1.5	4.3	12.6	33.0	65.0	103.3
2003 4/.....	*	*	*	0.6	1.4	4.5	13.3	33.6	68.2	107.4
2002.....	*	*	*	0.5	1.5	3.9	12.0	31.9	65.5	104.4
2001.....	*	*	*	0.5	0.9	4.0	12.3	32.7	64.2	101.7
2000.....	*	*	*	0.4	1.4	4.8	11.7	31.2	63.3	100.6
1999.....	*	*	*	0.4	0.9	4.2	10.8	27.6	57.3	96.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Parkinson's disease (G20-G21)										
Black, both sexes										
2006 4/.....	*	*	*	*	1.5	4.1	11.5	25.0	37.2	51.4
2005 4/.....	*	*	*	*	*	4.3	10.9	22.7	50.0	56.7
2004 4/.....	*	*	*	*	*	3.9	7.2	22.3	39.1	61.2
2003 4/.....	*	*	*	*	1.9	4.1	10.7	22.8	38.3	52.8
2002.....	*	*	*	*	*	4.6	9.5	20.6	32.7	51.7
2001.....	*	*	*	*	*	2.7	8.9	18.9	34.9	59.0
2000.....	*	*	*	*	*	3.1	8.8	21.0	37.6	51.6
1999.....	*	*	*	*	*	3.6	8.7	19.5	32.4	52.0
Black, male										
2006 4/.....	*	*	*	*	*	6.6	18.5	42.5	63.0	70.6
2005 4/.....	*	*	*	*	*	7.4	18.7	36.2	82.2	97.6
2004 4/.....	*	*	*	*	*	5.3	12.6	34.1	58.1	106.3
2003 4/.....	*	*	*	*	*	5.4	16.8	37.0	59.1	97.0
2002.....	*	*	*	*	*	7.5	16.0	30.3	55.0	89.0
2001.....	*	*	*	*	*	*	14.8	25.6	57.8	99.8
2000.....	*	*	*	*	*	*	13.7	37.8	65.8	86.5
1999.....	*	*	*	*	*	5.7	12.6	31.8	48.8	86.7
Black, female										
2006 4/.....	*	*	*	*	*	*	6.8	14.5	23.5	43.5
2005 4/.....	*	*	*	*	*	*	5.6	14.7	33.0	40.1
2004 4/.....	*	*	*	*	*	*	*	15.4	29.1	43.3
2003 4/.....	*	*	*	*	*	*	6.6	14.4	27.5	35.6
2002.....	*	*	*	*	*	*	5.2	14.8	21.1	37.4
2001.....	*	*	*	*	*	*	4.9	15.0	23.3	43.8
2000.....	*	*	*	*	*	*	5.5	10.8	23.4	38.6
1999.....	*	*	*	*	*	*	6.1	12.0	24.2	39.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
All races 3/, both sexes										
2006.....	*	0.1	0.4	1.1	3.4	9.6	32.9	101.2	272.6	848.3
2005.....	*	0.1	0.3	1.1	3.5	9.7	33.3	102.5	275.5	861.6
2004.....	*	0.2	0.3	0.9	3.1	10.1	31.0	99.4	261.0	818.8
2003.....	*	0.1	0.3	1.0	3.3	10.0	33.3	98.6	255.0	802.4
2002.....	*	*	0.2	0.9	3.2	9.8	30.6	95.4	245.7	752.3
2001.....	*	*	0.3	1.2	3.3	9.0	29.2	88.5	232.5	710.3
2000.....	*	*	0.3	1.1	3.2	9.1	28.9	85.2	221.1	667.7
1999.....	*	0.1	0.2	1.0	3.0	8.7	26.7	78.6	206.2	601.3
All races 3/, male										
2006.....	*	*	0.4	1.0	3.4	10.3	31.6	95.1	244.5	638.6
2005.....	*	*	0.2	1.0	3.2	9.6	31.4	95.9	246.9	639.9
2004.....	*	*	0.3	0.8	3.1	9.4	29.6	89.3	236.4	619.6
2003.....	*	*	0.3	1.0	2.9	10.1	32.1	92.1	231.2	606.5
2002.....	*	*	0.2	0.8	3.1	9.2	29.5	90.9	220.6	572.0
2001.....	*	*	0.3	0.9	3.2	9.0	27.7	85.6	216.6	556.1
2000.....	*	*	0.3	0.9	2.9	9.5	29.1	79.9	206.8	522.1
1999.....	*	*	*	1.1	3.0	8.8	26.9	77.4	198.2	485.7
All races 3/, female										
2006.....	*	*	0.3	1.3	3.3	9.0	34.0	105.7	290.1	946.4
2005.....	*	*	0.4	1.2	3.7	9.8	34.8	107.3	293.2	963.4
2004.....	*	0.2	0.4	1.0	3.1	10.7	32.1	106.7	276.2	908.4
2003.....	*	*	0.3	0.9	3.7	10.0	34.2	103.2	269.5	889.0
2002.....	*	*	*	1.0	3.3	10.3	31.5	98.5	260.9	830.5
2001.....	*	*	0.3	1.4	3.4	9.0	30.4	90.5	242.1	774.9
2000.....	*	*	0.3	1.2	3.5	8.7	28.8	88.9	229.5	727.0
1999.....	*	*	0.3	0.8	3.0	8.6	26.6	79.4	210.9	648.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
White, both sexes										
2006 4/.....	*	0.2	0.4	1.2	3.6	10.2	34.2	104.8	282.8	878.9
2005 4/.....	*	*	0.3	1.2	3.7	10.0	34.1	105.3	284.3	891.6
2004 4/.....	*	0.2	0.4	1.0	3.3	10.4	32.0	102.3	268.5	846.8
2003 4/.....	*	*	0.3	1.1	3.6	10.3	34.5	101.8	264.4	834.0
2002.....	*	*	0.2	1.0	3.4	10.3	31.9	99.1	255.3	781.8
2001.....	*	*	0.3	1.3	3.6	9.3	30.6	90.9	240.5	738.1
2000.....	*	*	0.3	1.1	3.4	9.6	29.9	88.1	229.1	695.3
1999.....	*	*	0.2	1.0	3.3	9.1	27.6	81.3	213.6	627.9
White, male										
2006 4/.....	*	*	0.4	1.1	3.6	10.9	32.6	98.3	253.6	662.5
2005 4/.....	*	*	0.2	1.1	3.5	9.8	31.9	97.5	254.3	661.6
2004 4/.....	*	*	0.4	0.9	3.3	9.7	30.2	91.6	243.9	640.9
2003 4/.....	*	*	0.3	1.1	3.0	10.3	33.1	94.9	238.5	630.3
2002.....	*	*	*	0.8	3.3	9.4	30.5	94.3	227.6	594.8
2001.....	*	*	0.3	0.9	3.4	9.2	28.8	88.1	222.7	577.2
2000.....	*	*	0.3	1.0	3.1	10.1	29.9	82.6	214.4	542.1
1999.....	*	*	*	1.1	3.3	9.2	27.6	80.1	205.6	506.9
White, female										
2006 4/.....	*	*	0.3	1.4	3.5	9.7	35.5	109.7	301.2	980.5
2005 4/.....	*	*	0.4	1.4	3.9	10.2	35.8	111.1	303.1	997.5
2004 4/.....	*	0.3	0.4	1.1	3.3	11.1	33.5	110.2	283.8	939.6
2003 4/.....	*	*	0.4	1.0	4.2	10.3	35.7	106.8	280.3	924.3
2002.....	*	*	*	1.2	3.5	11.0	33.1	102.6	272.2	863.0
2001.....	*	*	0.4	1.6	3.8	9.4	32.1	92.9	251.3	805.4
2000.....	*	*	0.3	1.3	3.7	9.2	29.9	92.0	237.7	757.6
1999.....	*	*	*	0.9	3.3	9.1	27.5	82.1	218.3	676.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alzheimer's disease (G30)										
Black, both sexes										
2006 4/.....	*	*	*	*	3.1	7.6	32.6	90.5	226.0	663.9
2005 4/.....	*	*	*	*	2.8	10.3	37.0	100.4	241.5	689.5
2004 4/.....	*	*	*	*	2.2	10.3	30.6	96.0	236.7	657.9
2003 4/.....	*	*	*	*	1.8	10.5	30.2	91.4	207.9	583.8
2002.....	*	*	*	*	2.5	8.5	27.2	81.3	186.8	543.7
2001.....	*	*	*	*	1.9	8.7	23.3	86.8	183.6	506.7
2000.....	*	*	*	*	2.1	7.4	25.9	71.1	171.7	445.1
1999.....	*	*	*	*	*	6.9	25.3	65.0	152.0	372.4
Black, male										
2006 4/.....	*	*	*	*	3.5	8.4	32.9	88.6	200.6	470.9
2005 4/.....	*	*	*	*	*	11.6	36.1	103.7	222.6	518.7
2004 4/.....	*	*	*	*	*	9.4	32.0	89.6	210.9	497.7
2003 4/.....	*	*	*	*	*	11.1	32.5	93.1	206.0	459.8
2002.....	*	*	*	*	*	9.5	29.4	82.2	189.5	412.8
2001.....	*	*	*	*	*	9.9	24.2	84.6	185.4	414.1
2000.....	*	*	*	*	*	7.8	27.7	64.7	156.0	371.3
1999.....	*	*	*	*	*	8.0	27.5	62.7	139.5	317.5
Black, female										
2006 4/.....	*	*	*	*	2.8	6.9	32.3	91.6	239.4	742.8
2005 4/.....	*	*	*	*	3.5	9.3	37.7	98.4	251.5	758.5
2004 4/.....	*	*	*	*	*	11.0	29.7	99.7	250.2	721.4
2003 4/.....	*	*	*	*	*	10.1	28.6	90.3	208.9	631.9
2002.....	*	*	*	*	*	7.8	25.8	80.7	185.4	593.9
2001.....	*	*	*	*	*	7.9	22.6	88.2	182.7	541.0
2000.....	*	*	*	*	*	7.1	24.6	74.9	179.6	472.6
1999.....	*	*	*	*	*	6.0	23.9	66.4	158.2	392.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple sclerosis (G35)										
All races 3/, both sexes										
2006.....	0.7	1.2	2.4	3.0	3.8	3.9	3.9	4.1	3.4	2.4
2005.....	0.8	1.5	2.5	3.1	3.8	3.6	3.7	4.0	4.0	2.2
2004.....	0.7	1.4	2.3	3.2	3.5	3.8	3.7	3.9	3.2	2.6
2003.....	0.8	1.5	2.4	3.1	3.3	3.6	3.6	4.0	3.0	2.2
2002.....	0.8	1.7	2.3	3.0	3.5	3.4	3.5	3.4	3.3	2.2
2001.....	0.9	1.6	2.3	3.0	3.3	3.4	3.3	3.6	3.4	2.1
2000.....	0.9	1.4	2.5	3.0	3.1	3.1	3.1	3.0	3.1	1.7
1999.....	0.9	1.4	2.4	2.4	2.7	3.0	2.8	3.2	2.8	2.1
All races 3/, male										
2006.....	0.5	0.8	1.7	2.2	2.6	2.8	3.3	3.6	2.4	1.8
2005.....	0.5	1.0	1.8	2.0	2.7	2.5	2.6	2.8	3.3	1.3
2004.....	0.5	1.0	1.5	2.4	2.4	2.9	2.9	2.8	2.4	1.7
2003.....	0.6	1.1	1.7	2.3	2.7	2.7	3.0	2.8	2.2	1.5
2002.....	0.5	1.1	1.6	2.2	3.1	2.4	2.6	2.7	2.4	1.9
2001.....	0.6	1.2	1.7	2.1	2.3	3.1	2.3	3.1	2.1	1.5
2000.....	0.7	1.1	1.8	2.2	2.3	2.3	2.4	2.4	2.2	1.8
1999.....	0.7	0.8	1.8	1.9	1.8	2.4	2.4	2.3	2.2	*
All races 3/, female										
2006.....	0.9	1.6	3.0	3.7	4.9	4.8	4.4	4.5	4.0	2.7
2005.....	1.0	2.0	3.1	4.1	4.9	4.6	4.7	4.8	4.4	2.7
2004.....	0.9	1.8	3.0	4.0	4.6	4.6	4.4	4.6	3.7	3.1
2003.....	0.9	1.9	3.1	3.8	3.8	4.4	4.1	4.8	3.4	2.5
2002.....	1.1	2.3	3.0	3.8	3.8	4.2	4.2	3.9	3.8	2.4
2001.....	1.2	2.0	3.0	3.7	4.1	3.6	4.1	4.0	4.1	2.3
2000.....	1.1	1.8	3.1	3.8	3.8	3.7	3.6	3.5	3.7	1.7
1999.....	1.2	2.0	3.0	2.9	3.5	3.5	3.1	3.8	3.1	2.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple sclerosis (G35)										
White, both sexes										
2006 4/.....	0.6	1.3	2.5	3.1	3.9	4.1	4.3	4.4	3.6	2.6
2005 4/.....	0.8	1.5	2.6	3.3	4.1	3.9	3.9	4.2	4.2	2.3
2004 4/.....	0.7	1.5	2.4	3.4	3.6	4.1	4.1	4.2	3.5	2.9
2003 4/.....	0.8	1.5	2.5	3.3	3.4	3.8	3.8	4.2	3.2	2.3
2002.....	0.8	1.7	2.3	3.2	3.6	3.6	3.8	3.6	3.5	2.4
2001.....	0.9	1.6	2.4	3.2	3.5	3.6	3.6	3.9	3.6	2.2
2000.....	0.9	1.4	2.6	3.1	3.2	3.3	3.3	3.3	3.3	1.9
1999.....	0.9	1.5	2.5	2.7	2.9	3.1	2.9	3.5	3.0	2.2
White, male										
2006 4/.....	0.5	0.9	1.8	2.4	2.7	3.1	3.6	3.9	2.5	1.9
2005 4/.....	0.5	1.0	1.9	2.1	2.9	2.7	2.7	2.8	3.5	1.4
2004 4/.....	0.5	1.1	1.6	2.6	2.6	3.0	3.1	2.8	2.5	1.8
2003 4/.....	0.7	1.2	1.7	2.5	2.8	2.9	3.2	3.0	2.4	*
2002.....	0.5	1.1	1.7	2.4	3.4	2.6	2.8	2.8	2.6	2.1
2001.....	0.6	1.2	1.8	2.3	2.5	3.4	2.5	3.3	2.2	1.7
2000.....	0.7	1.2	1.9	2.3	2.4	2.5	2.7	2.6	2.4	2.0
1999.....	0.7	0.9	1.8	2.1	1.9	2.6	2.5	2.5	2.4	*
White, female										
2006 4/.....	0.7	1.7	3.1	3.8	5.0	5.1	4.9	4.8	4.4	2.9
2005 4/.....	1.1	2.0	3.2	4.4	5.2	5.0	4.9	5.2	4.7	2.8
2004 4/.....	1.0	1.9	3.2	4.1	4.6	5.0	4.9	5.1	4.1	3.3
2003 4/.....	0.9	1.9	3.2	4.1	4.0	4.6	4.4	5.0	3.7	2.7
2002.....	1.1	2.3	3.0	4.0	3.9	4.4	4.6	4.1	4.1	2.5
2001.....	1.2	2.1	3.1	4.0	4.3	3.9	4.5	4.3	4.4	2.5
2000.....	1.1	1.7	3.2	4.0	4.0	4.0	3.8	3.8	3.9	1.9
1999.....	1.2	2.2	3.1	3.2	3.8	3.6	3.3	4.2	3.4	2.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Multiple sclerosis (G35)										
Black, both sexes										
2006 4/.....	1.4	1.3	2.7	3.2	4.9	2.8	*	*	*	*
2005 4/.....	1.1	1.9	2.8	2.7	3.3	2.3	3.6	*	*	*
2004 4/.....	1.0	1.7	2.2	3.2	4.3	3.2	*	*	*	*
2003 4/.....	1.1	2.2	2.7	2.2	3.1	3.4	2.6	*	*	*
2002.....	1.3	2.2	3.0	2.7	3.0	2.9	*	*	*	*
2001.....	1.2	2.0	2.4	2.6	3.0	2.5	*	*	*	*
2000.....	1.5	2.1	2.4	3.0	3.6	*	*	*	*	*
1999.....	1.3	1.4	2.5	1.5	2.2	2.5	*	*	*	*
Black, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	2.2	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	2.3	1.7	3.4	4.3	6.1	3.5	*	*	*	*
2005 4/.....	1.4	2.5	3.9	3.3	4.1	*	4.3	*	*	*
2004 4/.....	*	2.2	3.0	4.8	6.0	*	*	*	*	*
2003 4/.....	1.4	2.9	3.1	3.0	3.3	4.0	*	*	*	*
2002.....	2.1	2.9	4.1	3.5	4.0	4.3	*	*	*	*
2001.....	1.6	2.5	3.3	3.5	4.0	*	*	*	*	*
2000.....	1.7	3.1	3.1	3.4	3.9	*	*	*	*	*
1999.....	1.6	1.8	2.8	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Epilepsy (G40-G41)										
All races 3/, both sexes										
2006.....	0.6	0.6	0.7	0.6	0.7	0.6	0.8	1.0	1.3	1.5
2005.....	0.6	0.6	0.8	0.6	0.6	0.8	0.7	0.9	1.4	1.9
2004.....	0.6	0.6	0.6	0.6	0.6	0.7	0.6	1.1	1.1	1.5
2003.....	0.5	0.7	0.6	0.7	0.6	0.7	0.8	1.0	1.5	1.5
2002.....	0.6	0.6	0.6	0.5	0.6	0.7	0.9	1.3	1.4	1.8
2001.....	0.6	0.5	0.6	0.6	0.6	0.7	0.8	1.1	1.2	1.6
2000.....	0.6	0.6	0.6	0.7	0.6	0.7	1.0	1.0	1.6	2.0
1999.....	0.6	0.5	0.6	0.5	0.6	0.8	1.0	1.1	1.5	1.8
All races 3/, male										
2006.....	0.7	0.7	0.8	0.6	0.6	0.8	0.7	0.9	1.3	1.7
2005.....	0.8	0.8	0.8	0.8	0.6	0.8	0.7	0.7	1.2	1.9
2004.....	0.8	0.7	0.8	0.7	0.5	0.7	*	1.2	0.9	1.9
2003.....	0.7	0.8	0.8	0.7	0.6	0.6	0.9	1.0	1.8	1.5
2002.....	0.8	0.8	0.8	0.4	0.7	0.8	0.9	1.2	1.5	2.3
2001.....	0.8	0.6	0.7	0.7	0.7	0.8	0.8	1.0	*	2.0
2000.....	0.8	0.8	0.7	0.8	0.8	0.7	1.0	1.0	*	2.4
1999.....	0.7	0.7	0.6	0.6	0.7	0.9	1.0	0.9	1.3	2.3
All races 3/, female										
2006.....	0.5	0.5	0.6	0.5	0.7	0.5	0.9	1.0	1.2	1.4
2005.....	0.5	0.5	0.7	0.5	0.6	0.7	0.7	1.1	1.5	2.0
2004.....	0.4	0.4	0.3	0.6	0.7	0.6	0.7	1.1	1.2	1.4
2003.....	0.4	0.5	0.5	0.6	0.6	0.8	0.8	1.0	1.3	1.6
2002.....	0.4	0.5	0.4	0.5	0.6	0.7	0.9	1.3	1.4	1.5
2001.....	0.5	0.4	0.5	0.5	0.6	0.6	0.7	1.2	1.5	1.4
2000.....	0.4	0.4	0.5	0.5	0.5	0.6	1.0	1.0	2.0	1.9
1999.....	0.4	0.4	0.5	0.5	0.5	0.7	0.9	1.2	1.6	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Epilepsy (G40-G41)										
White, both sexes										
2006 4/.....	0.5	0.5	0.6	0.5	0.6	0.5	0.7	0.9	1.1	1.3
2005 4/.....	0.6	0.6	0.7	0.6	0.5	0.6	0.6	0.7	1.2	1.8
2004 4/.....	0.5	0.5	0.5	0.5	0.5	0.6	0.5	0.9	1.1	1.4
2003 4/.....	0.5	0.6	0.6	0.5	0.5	0.6	0.7	0.8	1.3	1.4
2002.....	0.6	0.6	0.5	0.4	0.6	0.6	0.7	1.1	1.3	1.6
2001.....	0.6	0.5	0.5	0.5	0.6	0.6	0.6	0.9	1.0	1.6
2000.....	0.6	0.5	0.6	0.6	0.5	0.6	0.8	0.8	1.5	1.7
1999.....	0.5	0.5	0.5	0.4	0.4	0.6	0.8	0.9	1.4	1.7
White, male										
2006 4/.....	0.6	0.6	0.7	0.6	0.5	0.7	0.7	0.9	1.2	1.6
2005 4/.....	0.8	0.7	0.8	0.7	0.4	0.7	0.7	*	1.2	1.8
2004 4/.....	0.7	0.7	0.8	0.5	0.4	0.6	*	0.9	*	1.8
2003 4/.....	0.6	0.7	0.7	0.6	0.5	*	0.7	0.8	1.5	*
2002.....	0.7	0.7	0.6	0.4	0.6	0.7	0.7	0.9	1.2	2.2
2001.....	0.7	0.6	0.6	0.6	0.6	0.7	0.7	0.7	*	2.0
2000.....	0.7	0.7	0.7	0.7	0.6	0.7	0.8	*	*	1.9
1999.....	0.6	0.6	0.5	0.4	*	0.7	0.9	*	1.3	2.3
White, female										
2006 4/.....	0.4	0.5	0.5	0.5	0.7	0.4	0.8	0.9	1.1	1.2
2005 4/.....	0.4	0.4	0.7	0.5	0.5	0.6	0.6	1.0	1.2	1.8
2004 4/.....	0.3	0.4	0.3	0.5	0.5	0.6	0.6	0.9	1.2	1.2
2003 4/.....	0.4	0.5	0.5	0.5	0.5	0.8	0.8	0.8	1.2	1.5
2002.....	0.4	0.4	0.3	0.4	0.5	0.6	0.7	1.2	1.3	1.3
2001.....	0.5	0.4	0.5	0.4	0.5	0.6	0.6	1.0	1.2	1.4
2000.....	0.5	0.3	0.5	0.4	0.4	0.5	0.9	0.9	1.9	1.6
1999.....	0.4	0.4	0.5	0.5	*	0.5	0.7	1.0	1.5	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transient cerebral ischemic attacks and related syndromes (G45)										
All races 3/, both sexes										
2006.....	*	*	*	*	*	*	*	*	*	1.5
2005.....	*	*	*	*	*	*	*	*	0.6	2.3
2004.....	*	*	*	*	*	*	*	0.3	0.6	2.1
2003.....	*	*	*	*	*	*	*	0.3	0.7	3.8
2002.....	*	*	*	*	*	*	*	0.4	0.8	2.9
2001.....	*	*	*	*	*	*	*	0.4	0.7	3.8
2000.....	*	*	*	*	*	*	*	0.5	1.0	3.9
1999.....	*	*	*	*	*	*	*	0.4	1.0	4.6
All races 3/, male										
2006.....	*	*	*	*	*	*	*	*	*	1.2
2005.....	*	*	*	*	*	*	*	*	*	1.8
2004.....	*	*	*	*	*	*	*	*	*	1.6
2003.....	*	*	*	*	*	*	*	*	*	2.8
2002.....	*	*	*	*	*	*	*	*	1.0	1.8
2001.....	*	*	*	*	*	*	*	*	*	4.2
2000.....	*	*	*	*	*	*	*	0.7	*	2.9
1999.....	*	*	*	*	*	*	*	0.7	1.2	3.8
All races 3/, female										
2006.....	*	*	*	*	*	*	*	*	*	1.6
2005.....	*	*	*	*	*	*	*	*	0.6	2.5
2004.....	*	*	*	*	*	*	*	*	0.7	2.4
2003.....	*	*	*	*	*	*	*	*	0.6	4.2
2002.....	*	*	*	*	*	*	*	*	0.6	3.4
2001.....	*	*	*	*	*	*	*	*	*	3.6
2000.....	*	*	*	*	*	*	*	*	1.1	4.2
1999.....	*	*	*	*	*	*	*	*	0.9	4.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Transient cerebral ischemic attacks and related syndromes (G45)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*	*	*	1.5
2005 4/.....	*	*	*	*	*	*	*	*	0.6	2.4
2004 4/.....	*	*	*	*	*	*	*	*	0.6	2.2
2003 4/.....	*	*	*	*	*	*	*	0.3	0.6	4.0
2002.....	*	*	*	*	*	*	*	0.4	0.8	3.0
2001.....	*	*	*	*	*	*	*	0.5	0.8	3.9
2000.....	*	*	*	*	*	*	*	0.5	1.1	4.0
1999.....	*	*	*	*	*	*	*	0.4	1.1	4.9

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	1.3
2005 4/.....	*	*	*	*	*	*	*	*	*	1.9
2004 4/.....	*	*	*	*	*	*	*	*	*	1.8
2003 4/.....	*	*	*	*	*	*	*	*	*	2.9
2002.....	*	*	*	*	*	*	*	*	*	1.8
2001.....	*	*	*	*	*	*	*	*	*	4.3
2000.....	*	*	*	*	*	*	*	0.7	*	3.0
1999.....	*	*	*	*	*	*	*	*	1.3	3.9

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	1.6
2005 4/.....	*	*	*	*	*	*	*	*	0.6	2.7
2004 4/.....	*	*	*	*	*	*	*	*	0.7	2.4
2003 4/.....	*	*	*	*	*	*	*	*	*	4.5
2002.....	*	*	*	*	*	*	*	*	0.7	3.6
2001.....	*	*	*	*	*	*	*	*	*	3.7
2000.....	*	*	*	*	*	*	*	*	1.1	4.4
1999.....	*	*	*	*	*	*	*	*	1.0	5.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

All races 3/, both sexes

2006.....	3.3	4.7	7.0	10.4	15.5	21.6	30.8	41.3	49.5	60.9
2005.....	3.1	4.8	7.0	10.3	14.8	20.3	29.1	38.2	47.4	55.1
2004.....	3.0	4.6	6.6	9.9	14.6	19.9	27.6	36.3	43.1	51.4
2003.....	3.0	4.5	6.5	9.7	15.2	20.6	29.0	37.8	43.9	56.0
2002.....	3.1	4.5	6.4	9.7	14.4	20.7	27.7	37.2	45.2	53.8
2001.....	2.9	4.3	6.1	9.2	13.8	20.7	27.6	36.3	43.9	51.9
2000.....	2.9	4.5	6.7	9.3	13.9	19.2	27.7	35.9	44.6	55.4
1999.....	2.5	4.1	6.2	9.1	13.1	19.8	27.1	36.6	43.4	53.6

All races 3/, male

2006.....	4.0	5.5	8.2	12.0	18.0	24.6	38.6	51.4	62.8	76.0
2005.....	3.7	5.6	8.2	12.4	16.7	23.3	34.4	47.0	59.9	67.3
2004.....	3.6	5.5	7.8	11.6	16.7	23.2	32.9	44.4	51.3	62.4
2003.....	3.9	5.5	7.9	11.4	17.4	23.1	33.3	45.4	56.3	70.0
2002.....	3.7	5.6	7.3	11.1	16.5	24.2	32.9	44.8	56.2	66.1
2001.....	3.4	5.0	7.0	10.4	15.3	24.6	33.0	44.4	56.2	69.7
2000.....	3.4	5.5	7.7	10.7	16.3	22.9	33.5	43.6	58.4	72.0
1999.....	3.0	4.6	7.2	10.5	15.3	23.3	32.0	45.6	56.5	70.3

All races 3/, female

2006.....	2.7	3.9	5.8	9.0	13.2	19.1	24.5	33.9	41.1	53.9
2005.....	2.4	3.9	5.9	8.3	13.0	17.8	24.9	31.8	39.6	49.5
2004.....	2.3	3.7	5.4	8.3	12.7	17.1	23.3	30.5	38.1	46.5
2003.....	2.2	3.5	5.1	8.2	13.1	18.4	25.5	32.3	36.3	49.9
2002.....	2.4	3.4	5.5	8.3	12.4	17.7	23.6	31.8	38.4	48.5
2001.....	2.3	3.7	5.2	8.0	12.4	17.3	23.4	30.6	36.6	44.5
2000.....	2.4	3.6	5.7	8.0	11.8	16.0	23.2	30.6	36.5	48.7
1999.....	2.0	3.6	5.1	7.8	11.0	16.8	23.2	30.4	35.7	46.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

White, both sexes

2006 4/.....	3.3	4.5	6.8	10.1	15.5	21.6	31.2	42.5	50.8	62.4
2005 4/.....	3.0	4.7	6.9	10.2	14.3	20.6	29.5	39.3	48.1	55.7
2004 4/.....	3.0	4.4	6.4	9.7	14.6	19.9	28.1	36.9	43.7	52.3
2003 4/.....	2.9	4.4	6.2	9.6	15.0	20.9	29.4	38.3	44.8	56.9
2002.....	3.0	4.4	6.2	9.5	14.4	20.9	28.0	37.5	46.1	54.7
2001.....	2.8	4.1	5.9	8.8	13.6	20.8	27.9	36.6	44.4	52.5
2000.....	2.7	4.3	6.5	9.2	13.9	19.1	27.8	36.6	45.4	56.2
1999.....	2.5	3.7	6.0	8.8	12.9	19.8	27.2	36.9	44.5	54.2

White, male

2006 4/.....	4.0	5.4	8.2	11.5	18.1	24.6	39.3	53.0	64.7	78.5
2005 4/.....	3.8	5.6	8.3	12.3	16.2	23.6	35.2	48.5	60.7	68.4
2004 4/.....	3.8	5.4	7.7	11.3	16.7	23.3	33.8	45.0	52.3	64.7
2003 4/.....	3.9	5.4	7.8	11.4	17.2	23.3	34.2	46.3	57.5	72.0
2002.....	3.7	5.5	7.0	10.9	16.6	24.1	33.0	45.5	57.7	67.5
2001.....	3.3	4.8	6.7	10.1	15.2	24.8	33.3	44.7	57.4	70.1
2000.....	3.2	5.4	7.7	10.5	16.1	23.0	33.8	44.5	59.3	73.1
1999.....	3.0	4.5	7.1	10.4	15.2	23.4	32.3	46.3	58.5	71.5

White, female

2006 4/.....	2.7	3.6	5.5	8.7	13.1	18.9	24.4	34.7	42.1	54.9
2005 4/.....	2.2	3.7	5.5	8.1	12.5	17.9	24.9	32.4	40.2	49.8
2004 4/.....	2.2	3.4	5.2	8.1	12.6	16.9	23.3	30.9	38.3	46.8
2003 4/.....	2.0	3.4	4.7	7.9	12.9	18.8	25.4	32.6	37.0	50.3
2002.....	2.4	3.3	5.3	8.1	12.4	18.1	23.9	31.8	39.0	49.2
2001.....	2.3	3.5	5.1	7.5	12.1	17.3	23.5	30.8	36.5	45.1
2000.....	2.3	3.3	5.4	8.0	11.8	15.7	23.1	31.1	37.1	49.3
1999.....	2.0	3.0	4.9	7.3	10.8	16.7	23.1	30.4	36.2	47.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of nervous system (G10-G12,G23-G25,G31,G36-G37,G43-G44,G47-G72,G81-G98)

Black, both sexes

2006 4/.....	4.4	7.4	10.1	16.2	18.6	26.2	33.9	40.2	46.1	55.5
2005 4/.....	4.3	6.6	10.1	14.3	22.4	24.8	33.0	34.6	46.8	55.3
2004 4/.....	3.7	7.2	9.5	14.2	18.9	24.5	29.2	39.4	44.6	49.6
2003 4/.....	4.4	6.5	9.9	13.0	19.9	24.3	30.7	37.1	39.9	52.5
2002.....	4.2	6.4	9.2	13.0	18.1	23.5	31.6	37.6	42.6	50.5
2001.....	4.0	6.5	9.7	14.6	17.7	24.3	30.7	38.4	45.8	48.8
2000.....	4.5	7.5	9.9	12.5	16.7	24.2	30.4	35.2	40.7	51.3
1999.....	3.5	7.6	9.4	13.1	16.5	22.5	31.4	38.0	34.7	50.7

Black, male

2006 4/.....	4.9	8.3	11.0	19.7	21.4	30.0	40.3	49.2	57.5	63.4
2005 4/.....	4.3	7.2	10.5	17.0	25.8	28.8	35.7	39.3	59.0	62.3
2004 4/.....	3.8	8.2	11.8	17.6	21.3	27.1	32.7	46.9	50.2	50.7
2003 4/.....	5.1	7.9	10.7	14.3	23.9	29.3	32.5	42.9	47.2	50.0
2002.....	5.1	7.9	10.5	15.6	20.4	29.2	40.1	42.3	49.7	57.9
2001.....	5.0	7.5	11.4	15.2	19.3	28.2	35.7	49.7	53.0	62.4
2000.....	5.6	8.3	10.2	16.0	20.1	27.6	37.4	44.9	56.5	63.4
1999.....	4.2	7.5	11.1	13.1	18.9	24.7	33.9	47.0	37.5	65.6

Black, female

2006 4/.....	3.9	6.6	9.3	13.3	16.4	23.4	29.6	34.8	40.1	52.3
2005 4/.....	4.2	6.0	9.7	12.0	19.7	21.9	31.2	31.7	40.4	52.5
2004 4/.....	3.6	6.4	7.5	11.4	17.0	22.5	26.8	35.0	41.6	49.2
2003 4/.....	3.8	5.3	9.3	12.0	16.7	20.7	29.5	33.6	36.0	53.5
2002.....	3.5	5.1	8.2	10.9	16.3	19.2	26.0	34.9	39.0	47.6
2001.....	3.1	5.5	8.2	14.1	16.5	21.4	27.4	31.6	42.1	43.8
2000.....	3.5	6.7	9.6	9.7	14.1	21.7	25.8	29.3	32.7	46.8
1999.....	2.9	7.6	7.9	13.2	14.6	21.0	29.7	32.6	33.3	45.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

IX. Diseases of the circulatory system (I00-I99)

All races 3/, both sexes

2006.....	47.3	84.4	138.4	207.3	325.4	492.8	802.2	1,373.9	2,477.4	5,946.6
2005.....	48.1	85.5	141.6	210.2	338.9	518.5	852.7	1,459.7	2,646.0	6,421.6
2004.....	48.7	87.2	140.6	213.0	345.4	540.6	889.4	1,521.5	2,741.8	6,639.9
2003.....	51.3	89.0	144.2	226.3	366.2	577.3	951.0	1,634.2	2,930.7	7,162.1
2002.....	50.4	89.0	145.7	235.5	376.5	605.9	997.5	1,707.6	3,047.6	7,414.0
2001.....	49.6	86.6	146.1	236.8	388.4	623.5	1,022.9	1,767.1	3,145.7	7,699.8
2000.....	50.5	89.8	148.2	250.1	410.1	651.3	1,071.3	1,832.2	3,265.6	8,055.8
1999.....	50.8	89.3	149.7	256.4	418.3	682.1	1,115.6	1,895.5	3,369.1	8,220.5

All races 3/, male

2006.....	63.4	116.8	197.5	293.0	446.1	660.3	1,031.1	1,719.4	2,990.5	6,168.0
2005.....	65.0	118.5	202.0	297.1	464.9	683.9	1,098.4	1,826.8	3,170.0	6,635.1
2004.....	65.6	121.7	200.3	298.4	474.9	718.1	1,139.5	1,908.7	3,295.5	6,837.7
2003.....	68.6	124.5	206.3	316.8	500.6	769.4	1,230.2	2,058.7	3,527.9	7,348.4
2002.....	68.7	125.2	207.7	329.0	515.2	805.6	1,291.7	2,147.8	3,646.7	7,659.3
2001.....	67.3	122.7	206.2	329.7	529.2	825.3	1,331.5	2,231.5	3,786.4	7,958.0
2000.....	69.1	128.3	210.8	352.5	559.6	861.1	1,409.4	2,313.7	3,969.1	8,470.3
1999.....	69.8	126.7	213.7	360.1	570.7	912.4	1,466.7	2,400.1	4,092.1	8,624.0

All races 3/, female

2006.....	31.2	52.7	81.7	126.5	215.1	346.5	616.0	1,121.1	2,156.6	5,843.1
2005.....	31.4	53.4	83.7	128.1	224.0	374.1	653.7	1,193.0	2,320.6	6,323.5
2004.....	32.0	53.7	83.4	132.5	227.5	386.2	688.0	1,243.2	2,400.8	6,551.0
2003.....	34.1	54.4	84.9	141.0	244.0	410.7	727.1	1,332.1	2,565.4	7,079.7
2002.....	32.4	53.9	86.3	147.7	250.7	433.5	762.4	1,396.8	2,683.2	7,307.6
2001.....	32.2	51.6	88.6	149.8	260.6	449.7	777.7	1,441.9	2,761.9	7,591.7
2000.....	32.1	52.4	88.3	154.3	274.6	471.5	804.9	1,496.7	2,850.6	7,887.0
1999.....	32.0	53.1	88.4	159.5	280.3	485.1	841.2	1,544.6	2,946.1	8,057.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
IX. Diseases of the circulatory system (I00-I99)										
White, both sexes										
2006 4/.....	41.7	74.3	121.9	186.6	299.9	459.9	768.1	1,346.2	2,465.7	6,047.9
2005 4/.....	42.1	74.9	123.9	189.0	311.6	486.2	819.0	1,427.3	2,634.3	6,531.2
2004 4/.....	42.6	76.0	123.3	191.3	318.5	506.9	855.3	1,486.6	2,733.2	6,728.0
2003 4/.....	44.2	77.3	126.2	204.9	335.4	540.6	911.7	1,600.0	2,921.5	7,256.7
2002.....	44.3	77.7	128.7	213.7	346.9	571.0	961.2	1,670.2	3,036.0	7,507.0
2001.....	42.3	75.2	128.8	214.1	357.9	583.9	985.3	1,728.3	3,134.6	7,796.7
2000.....	43.4	77.7	132.0	227.1	378.7	615.7	1,032.2	1,790.7	3,255.4	8,159.4
1999.....	43.1	77.5	133.3	231.7	387.3	647.6	1,072.5	1,854.7	3,357.7	8,322.3
White, male										
2006 4/.....	57.8	105.5	177.7	267.1	414.4	620.9	991.4	1,691.6	2,995.8	6,303.1
2005 4/.....	58.7	106.6	181.0	270.8	432.8	647.0	1,061.6	1,799.5	3,175.4	6,791.7
2004 4/.....	59.8	109.2	180.6	272.3	442.5	680.8	1,105.9	1,877.0	3,310.2	6,960.5
2003 4/.....	61.4	111.5	185.2	292.0	466.1	730.5	1,188.2	2,029.2	3,531.1	7,485.4
2002.....	62.6	113.0	188.2	303.5	481.4	768.5	1,257.1	2,119.7	3,654.4	7,786.5
2001.....	59.8	110.2	187.0	303.7	496.2	785.5	1,291.8	2,200.7	3,789.8	8,082.3
2000.....	61.7	115.8	192.1	325.2	526.1	826.1	1,371.9	2,281.0	3,975.9	8,607.0
1999.....	61.6	113.9	194.5	331.6	535.7	877.3	1,424.9	2,368.9	4,101.9	8,738.2
White, female										
2006 4/.....	25.2	43.2	67.2	109.1	193.3	316.5	582.7	1,088.4	2,130.8	5,928.1
2005 4/.....	25.3	43.4	67.8	110.1	198.9	343.1	618.4	1,151.9	2,295.3	6,411.3
2004 4/.....	25.2	43.0	67.3	113.3	203.5	352.6	649.5	1,201.0	2,374.8	6,623.2
2003 4/.....	26.8	43.2	68.4	121.2	214.4	372.9	685.4	1,289.6	2,545.7	7,155.4
2002.....	25.9	42.6	70.5	127.7	222.6	397.2	720.1	1,348.4	2,656.9	7,385.6
2001.....	24.6	40.5	72.0	128.6	230.3	406.9	737.1	1,393.7	2,738.6	7,677.1
2000.....	25.0	39.9	73.1	133.8	242.8	431.7	759.7	1,445.8	2,826.7	7,977.5
1999.....	24.5	41.5	73.6	136.8	250.6	447.2	792.4	1,493.6	2,918.3	8,155.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
IX. Diseases of the circulatory system (I00-I99)										
Black, both sexes										
2006 4/.....	92.4	168.4	287.7	414.7	615.4	862.1	1,258.7	1,899.5	2,993.2	5,612.9
2005 4/.....	96.0	174.8	298.8	426.3	645.8	887.7	1,309.0	2,052.8	3,185.8	6,041.2
2004 4/.....	100.0	181.9	297.1	438.1	655.7	928.1	1,383.6	2,144.9	3,231.9	6,397.4
2003 4/.....	106.5	186.5	308.3	455.3	700.9	989.8	1,498.2	2,263.5	3,423.0	6,895.6
2002.....	100.9	186.4	305.2	472.5	703.6	1,007.6	1,517.4	2,356.8	3,577.5	7,111.7
2001.....	107.5	184.1	309.6	475.0	719.8	1,064.4	1,557.8	2,436.9	3,638.1	7,346.8
2000.....	106.7	194.4	306.7	493.2	742.0	1,070.6	1,632.2	2,510.7	3,733.9	7,548.9
1999.....	111.7	190.9	313.3	511.4	747.2	1,100.0	1,711.0	2,564.7	3,805.3	7,610.9
Black, male										
2006 4/.....	113.9	219.4	395.7	574.4	852.6	1,162.1	1,646.5	2,388.2	3,511.0	5,441.3
2005 4/.....	120.0	228.1	406.0	590.6	880.3	1,164.6	1,677.4	2,520.7	3,687.0	5,732.7
2004 4/.....	121.0	236.6	398.3	597.9	896.8	1,216.2	1,737.7	2,639.2	3,693.8	6,222.9
2003 4/.....	131.6	243.1	416.3	613.6	931.7	1,281.3	1,910.4	2,805.4	4,056.0	6,580.1
2002.....	124.8	240.2	409.4	638.6	940.4	1,300.5	1,916.4	2,815.9	4,121.4	6,961.1
2001.....	132.1	239.0	408.8	632.4	942.9	1,350.6	1,998.5	2,960.2	4,299.5	7,393.4
2000.....	133.7	250.3	410.7	675.6	960.7	1,337.4	2,049.3	3,012.7	4,411.3	7,717.4
1999.....	140.9	249.2	426.4	688.3	996.7	1,409.3	2,153.2	3,061.2	4,428.7	7,943.3
Black, female										
2006 4/.....	73.4	124.1	196.4	282.9	430.5	640.6	992.2	1,607.0	2,719.6	5,683.1
2005 4/.....	74.9	128.2	208.1	290.6	462.9	682.4	1,056.8	1,773.6	2,921.1	6,165.8
2004 4/.....	81.6	134.1	211.5	306.5	467.8	714.2	1,143.3	1,852.1	2,988.4	6,466.6
2003 4/.....	84.4	137.1	216.9	325.0	520.9	773.4	1,220.2	1,941.4	3,091.2	7,018.2
2002.....	79.8	139.8	216.8	336.2	518.4	790.4	1,249.6	2,082.4	3,294.4	7,169.4
2001.....	85.8	136.5	225.1	345.8	544.3	853.0	1,264.3	2,124.5	3,301.2	7,329.5
2000.....	82.8	145.9	218.0	343.7	569.4	873.8	1,355.3	2,208.1	3,392.7	7,486.2
1999.....	86.0	140.6	217.4	367.0	551.6	870.8	1,417.5	2,265.6	3,496.4	7,486.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes

2006.....	0.2	0.4	0.4	0.9	1.5	2.5	4.1	6.8	11.0	16.7
2005.....	0.2	0.4	0.5	0.9	1.5	2.5	4.2	6.7	10.9	19.0
2004.....	0.3	0.4	0.6	0.9	1.4	2.6	4.2	7.3	10.9	17.1
2003.....	0.3	0.5	0.6	1.0	1.8	3.0	4.6	8.1	11.6	20.5
2002.....	0.2	0.5	0.7	1.3	2.1	2.8	5.0	8.4	11.4	17.8
2001.....	0.4	0.5	0.7	1.2	1.7	3.2	5.0	8.1	11.4	17.7
2000.....	0.4	0.6	0.8	1.4	2.2	3.4	5.6	8.1	11.6	16.8
1999.....	0.3	0.5	0.9	1.4	2.1	3.5	5.8	8.4	12.6	18.0

All races 3/, male

2006.....	*	0.4	0.4	0.7	1.2	1.9	3.0	5.1	7.7	12.0
2005.....	0.2	0.3	0.5	0.9	1.1	1.9	3.1	4.9	8.0	14.0
2004.....	0.2	0.4	0.5	0.7	0.9	2.1	2.8	5.7	9.5	11.5
2003.....	0.2	0.4	0.4	0.8	1.5	2.3	3.6	6.4	9.0	16.2
2002.....	0.2	0.4	0.6	1.0	1.5	1.9	4.2	5.8	8.2	11.8
2001.....	0.3	0.4	0.5	1.1	1.2	2.4	3.2	6.0	7.7	12.8
2000.....	0.3	0.5	0.6	1.1	1.8	2.2	3.6	5.8	7.8	12.2
1999.....	0.3	0.4	0.6	1.0	1.4	2.4	4.7	5.4	8.9	10.3

All races 3/, female

2006.....	0.2	0.4	0.5	1.0	1.7	3.0	5.0	8.0	13.0	18.9
2005.....	0.3	0.5	0.6	1.0	1.9	3.0	5.1	8.0	12.7	21.3
2004.....	0.3	0.4	0.7	1.1	1.9	3.0	5.3	8.3	11.8	19.5
2003.....	0.3	0.6	0.8	1.1	2.1	3.6	5.5	9.3	13.2	22.3
2002.....	0.3	0.5	0.9	1.5	2.6	3.7	5.6	10.2	13.4	20.4
2001.....	0.4	0.6	0.9	1.3	2.2	3.9	6.4	9.5	13.7	19.8
2000.....	0.5	0.7	0.9	1.6	2.5	4.5	7.1	9.7	13.9	18.7
1999.....	0.4	0.6	1.1	1.8	2.7	4.5	6.7	10.4	14.7	21.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

White, both sexes

2006 4/.....	0.2	0.3	0.4	0.8	1.3	2.5	4.2	7.1	11.6	18.0
2005 4/.....	0.2	0.3	0.4	0.8	1.4	2.5	4.4	7.0	11.5	20.4
2004 4/.....	0.2	0.3	0.6	0.7	1.4	2.6	4.4	7.6	11.6	18.2
2003 4/.....	0.2	0.4	0.5	0.8	1.6	3.1	4.7	8.6	12.4	21.8
2002.....	0.2	0.4	0.7	1.2	2.0	2.8	5.2	8.6	12.1	18.9
2001.....	0.3	0.4	0.6	1.0	1.7	3.2	5.0	8.4	12.2	18.7
2000.....	0.3	0.5	0.7	1.3	2.2	3.5	5.6	8.6	12.4	17.6
1999.....	0.3	0.5	0.8	1.4	2.0	3.6	6.0	8.8	13.1	18.9

White, male

2006 4/.....	*	0.3	0.3	0.6	1.1	1.9	3.0	5.3	8.3	12.9
2005 4/.....	*	0.3	0.4	0.8	1.1	1.9	3.3	5.2	8.3	14.7
2004 4/.....	*	0.3	0.4	0.6	0.9	2.0	3.1	6.1	10.1	12.4
2003 4/.....	*	0.3	0.3	0.7	1.3	2.3	3.7	6.9	9.8	17.3
2002.....	*	0.3	0.5	0.9	1.5	1.8	4.4	5.9	8.8	12.6
2001.....	0.3	0.4	0.4	1.0	1.2	2.3	3.3	6.3	8.0	13.3
2000.....	0.3	0.5	0.7	1.1	1.9	2.3	3.5	6.1	8.4	12.7
1999.....	0.2	0.4	0.6	1.0	1.3	2.6	4.9	5.5	9.4	11.1

White, female

2006 4/.....	*	0.3	0.4	1.0	1.6	3.0	5.2	8.4	13.6	20.4
2005 4/.....	*	0.4	0.5	0.9	1.8	3.1	5.3	8.4	13.6	23.1
2004 4/.....	0.2	0.3	0.7	0.9	1.9	3.0	5.4	8.6	12.5	20.8
2003 4/.....	0.2	0.5	0.6	1.0	1.9	3.8	5.5	9.8	14.0	23.8
2002.....	0.2	0.5	0.9	1.4	2.5	3.7	5.9	10.6	14.2	21.6
2001.....	0.3	0.4	0.7	1.1	2.3	3.9	6.4	9.9	14.7	21.0
2000.....	0.4	0.6	0.8	1.5	2.4	4.6	7.3	10.4	14.7	19.6
1999.....	0.3	0.6	0.9	1.8	2.7	4.6	6.9	11.0	15.4	22.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)										
Black, both sexes										
2006 4/.....	*	0.8	*	1.1	2.5	2.8	3.1	4.7	6.2	*
2005 4/.....	*	*	1.3	1.2	1.8	2.5	*	4.2	*	6.1
2004 4/.....	0.7	1.0	1.0	2.0	*	2.6	2.7	4.1	*	5.9
2003 4/.....	*	1.2	1.4	2.2	3.2	3.1	4.8	3.9	*	7.6
2002.....	*	0.8	1.1	2.0	2.9	3.2	*	4.8	*	*
2001.....	0.9	1.1	1.6	2.2	2.0	3.3	5.1	4.9	*	6.8
2000.....	0.9	1.0	1.1	1.9	2.2	3.4	3.6	3.7	*	8.4
1999.....	0.7	0.9	1.6	2.2	2.1	2.8	4.2	4.3	*	8.3
Black, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	*	*	*	*	*	*	*	*	8.0	*
2005 4/.....	*	*	*	*	*	*	*	5.3	*	*
2004 4/.....	*	*	*	*	*	*	*	5.4	*	*
2003 4/.....	*	1.6	1.7	*	3.3	*	6.2	*	*	8.1
2002.....	*	*	*	*	3.6	*	*	5.5	*	*
2001.....	*	*	2.2	2.7	*	3.8	7.2	6.4	*	*
2000.....	*	*	*	*	*	4.4	*	*	*	*
1999.....	*	*	*	2.8	*	*	4.7	*	*	11.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic rheumatic heart diseases (I05-I09)										
All races 3/, both sexes										
2006.....	0.2	0.4	0.4	0.9	1.4	2.5	4.1	6.8	10.9	16.6
2005.....	0.2	0.4	0.5	0.9	1.5	2.4	4.2	6.7	10.9	18.9
2004.....	0.3	0.4	0.6	0.9	1.4	2.5	4.1	7.2	10.8	17.0
2003.....	0.2	0.5	0.6	0.9	1.7	3.0	4.6	8.0	11.6	20.3
2002.....	0.2	0.4	0.7	1.3	2.1	2.8	4.9	8.2	11.3	17.5
2001.....	0.4	0.5	0.7	1.2	1.7	3.2	4.9	8.0	11.3	17.2
2000.....	0.4	0.6	0.7	1.4	2.1	3.4	5.5	8.0	11.6	16.7
1999.....	0.3	0.5	0.8	1.4	2.1	3.5	5.8	8.3	12.5	17.9
All races 3/, male										
2006.....	*	0.4	0.4	0.7	1.1	1.9	3.0	5.1	7.7	12.0
2005.....	0.2	0.3	0.5	0.9	1.1	1.9	3.0	4.8	8.0	13.9
2004.....	0.2	0.3	0.5	0.7	0.9	2.0	2.8	5.7	9.4	11.4
2003.....	0.2	0.4	0.4	0.8	1.4	2.3	3.5	6.4	9.0	16.1
2002.....	0.2	0.4	0.6	1.0	1.5	1.9	4.1	5.7	8.1	11.4
2001.....	0.3	0.4	0.4	1.0	1.2	2.4	3.2	5.9	7.5	12.3
2000.....	0.3	0.5	0.6	1.1	1.8	2.2	3.5	5.7	7.8	12.1
1999.....	0.3	0.4	0.6	1.0	1.3	2.4	4.6	5.4	8.8	10.3
All races 3/, female										
2006.....	0.2	0.4	0.5	1.0	1.7	3.0	5.0	8.0	12.9	18.8
2005.....	0.3	0.5	0.6	0.9	1.9	2.9	5.1	8.0	12.7	21.2
2004.....	0.3	0.4	0.7	1.1	1.8	3.0	5.2	8.3	11.7	19.5
2003.....	0.3	0.6	0.8	1.1	2.0	3.6	5.4	9.1	13.2	22.2
2002.....	0.3	0.5	0.9	1.5	2.6	3.6	5.5	10.0	13.3	20.2
2001.....	0.4	0.6	0.9	1.3	2.2	3.8	6.3	9.4	13.5	19.3
2000.....	0.5	0.7	0.8	1.6	2.5	4.4	7.1	9.7	13.8	18.5
1999.....	0.4	0.6	1.1	1.8	2.7	4.5	6.6	10.3	14.6	21.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic rheumatic heart diseases (I05-I09)										
White, both sexes										
2006 4/.....	0.2	0.3	0.4	0.8	1.3	2.5	4.2	7.0	11.5	18.0
2005 4/.....	0.2	0.3	0.4	0.8	1.4	2.5	4.4	7.0	11.5	20.3
2004 4/.....	0.2	0.3	0.6	0.7	1.4	2.5	4.3	7.5	11.5	18.1
2003 4/.....	0.2	0.4	0.5	0.8	1.6	3.1	4.6	8.5	12.4	21.7
2002.....	0.2	0.4	0.7	1.1	2.0	2.8	5.2	8.5	12.0	18.6
2001.....	0.3	0.4	0.6	1.0	1.7	3.1	4.9	8.3	12.0	18.3
2000.....	0.3	0.5	0.7	1.3	2.1	3.5	5.6	8.6	12.3	17.5
1999.....	0.3	0.5	0.7	1.4	2.0	3.6	6.0	8.7	13.1	18.9
White, male										
2006 4/.....	*	0.3	0.3	0.6	1.1	1.9	3.0	5.2	8.3	12.8
2005 4/.....	*	0.3	0.4	0.8	1.1	1.9	3.2	5.1	8.2	14.7
2004 4/.....	*	0.3	0.4	0.6	0.8	2.0	3.0	6.1	10.0	12.3
2003 4/.....	*	0.3	0.3	0.6	1.2	2.3	3.6	6.9	9.7	17.2
2002.....	*	0.3	0.5	0.9	1.5	1.8	4.3	5.8	8.7	12.2
2001.....	0.3	0.3	0.4	1.0	1.2	2.3	3.2	6.2	7.9	12.9
2000.....	0.3	0.5	0.7	1.1	1.8	2.3	3.4	6.0	8.4	12.6
1999.....	0.2	0.4	0.5	0.9	1.3	2.6	4.8	5.5	9.3	11.1
White, female										
2006 4/.....	*	0.3	0.4	1.0	1.6	3.0	5.1	8.4	13.5	20.4
2005 4/.....	*	0.4	0.4	0.9	1.7	3.0	5.3	8.3	13.6	22.9
2004 4/.....	0.2	0.3	0.7	0.9	1.9	3.0	5.4	8.6	12.4	20.7
2003 4/.....	0.2	0.5	0.6	1.0	1.9	3.8	5.4	9.6	14.0	23.7
2002.....	0.2	0.5	0.8	1.4	2.4	3.7	5.9	10.4	14.1	21.4
2001.....	0.3	0.4	0.7	1.1	2.2	3.9	6.3	9.8	14.5	20.5
2000.....	0.4	0.6	0.8	1.5	2.4	4.6	7.3	10.3	14.7	19.5
1999.....	0.3	0.6	0.9	1.8	2.7	4.6	6.9	11.0	15.3	22.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic rheumatic heart diseases (I05-I09)										
Black, both sexes										
2006 4/.....	*	0.7	*	1.1	2.3	2.7	3.1	4.7	6.2	*
2005 4/.....	*	*	1.3	1.2	1.8	2.5	*	4.0	*	6.1
2004 4/.....	0.7	1.0	1.0	1.9	*	2.5	2.7	4.1	*	5.9
2003 4/.....	*	1.2	1.3	2.2	3.0	3.1	4.6	3.6	*	7.6
2002.....	*	0.8	1.1	2.0	2.9	3.2	*	4.5	*	*
2001.....	0.9	1.1	1.6	2.2	1.9	3.0	5.1	4.9	*	*
2000.....	0.9	1.0	*	1.9	2.2	3.3	3.6	3.7	*	8.1
1999.....	*	0.9	1.6	2.1	2.1	2.8	4.2	4.1	*	7.7
Black, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	*	*	*	*	*	*	*	*	8.0	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	5.4	*	*
2003 4/.....	*	1.6	*	*	3.2	*	6.0	*	*	8.1
2002.....	*	*	*	*	3.6	*	*	*	*	*
2001.....	*	*	2.2	2.7	*	*	7.2	6.4	*	*
2000.....	*	*	*	*	*	4.2	*	*	*	*
1999.....	*	*	*	*	*	*	4.7	*	*	10.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rheumatic mitral valve diseases (I05)										
All races 3/, both sexes										
2006.....	0.1	0.2	0.2	0.5	0.8	1.3	2.0	3.3	4.9	6.4
2005.....	0.1	0.2	0.2	0.5	0.8	1.2	1.9	3.3	5.0	7.1
2004.....	0.1	0.2	0.3	0.4	0.7	1.2	2.1	3.3	4.7	6.6
2003.....	0.1	0.3	0.4	0.4	1.0	1.7	2.3	4.0	5.6	7.6
2002.....	0.2	0.2	0.5	0.8	1.3	1.4	2.6	4.4	5.7	7.4
2001.....	0.2	0.3	0.5	0.7	1.0	1.8	2.8	4.5	6.0	8.5
2000.....	0.2	0.4	0.5	0.8	1.3	2.0	3.4	4.4	6.8	8.5
1999.....	0.2	0.3	0.5	0.8	1.2	2.1	3.2	4.9	6.8	9.0
All races 3/, male										
2006.....	*	*	*	0.4	0.6	0.9	1.6	2.7	3.4	4.0
2005.....	*	0.2	*	0.5	0.6	1.0	1.4	2.3	4.2	4.4
2004.....	*	0.2	0.3	0.3	0.4	0.8	1.4	2.4	3.9	4.6
2003.....	*	0.2	0.2	0.4	1.0	1.4	1.6	3.1	4.2	5.9
2002.....	*	*	0.4	0.6	0.9	0.9	2.3	2.6	3.7	4.2
2001.....	0.2	0.3	0.3	0.6	0.5	1.4	1.8	2.9	3.7	6.2
2000.....	0.2	0.3	0.3	0.7	1.1	1.2	2.2	3.2	4.7	6.1
1999.....	*	0.3	0.4	0.6	0.7	1.5	2.4	3.2	4.4	4.9
All races 3/, female										
2006.....	*	0.2	0.2	0.6	0.9	1.6	2.3	3.7	5.8	7.5
2005.....	*	0.3	0.3	0.6	1.0	1.4	2.4	4.0	5.5	8.3
2004.....	0.2	0.2	0.4	0.5	1.0	1.4	2.7	3.9	5.3	7.5
2003.....	0.2	0.4	0.5	0.5	1.0	1.9	2.9	4.7	6.5	8.4
2002.....	*	0.3	0.6	0.9	1.6	1.8	2.9	5.7	7.0	8.7
2001.....	0.2	0.4	0.6	0.8	1.4	2.1	3.6	5.5	7.4	9.4
2000.....	0.3	0.4	0.6	0.9	1.5	2.7	4.3	5.2	8.0	9.5
1999.....	0.2	0.4	0.7	1.1	1.6	2.7	3.8	6.1	8.3	10.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rheumatic mitral valve diseases (I05)										
White, both sexes										
2006 4/.....	*	0.2	0.2	0.5	0.8	1.3	2.0	3.4	5.1	6.9
2005 4/.....	*	0.2	0.2	0.5	0.8	1.2	2.1	3.5	5.2	7.7
2004 4/.....	*	0.2	0.3	0.4	0.7	1.2	2.2	3.4	4.9	7.1
2003 4/.....	0.1	0.2	0.3	0.3	0.9	1.7	2.4	4.3	5.9	8.1
2002.....	0.1	0.2	0.5	0.7	1.2	1.3	2.8	4.6	6.2	7.9
2001.....	0.2	0.3	0.4	0.6	1.0	1.7	2.8	4.7	6.5	9.1
2000.....	0.2	0.4	0.5	0.8	1.3	2.2	3.4	4.6	7.3	9.1
1999.....	0.1	0.3	0.4	0.8	1.2	2.2	3.3	5.2	7.2	9.5
White, male										
2006 4/.....	*	*	*	0.3	0.6	0.9	1.6	2.8	3.7	4.1
2005 4/.....	*	*	*	0.5	0.6	0.9	1.6	2.5	4.3	4.8
2004 4/.....	*	*	*	0.3	*	0.9	1.6	2.5	4.1	5.0
2003 4/.....	*	*	*	*	0.8	1.4	1.7	3.3	4.5	6.4
2002.....	*	*	0.3	0.5	0.9	0.9	2.4	2.6	4.1	4.5
2001.....	*	0.2	*	0.5	0.6	1.2	1.8	3.1	4.0	6.5
2000.....	*	0.3	0.4	0.7	1.1	1.4	2.1	3.3	5.1	6.4
1999.....	*	0.3	0.4	0.5	0.7	1.5	2.5	3.2	4.6	5.3
White, female										
2006 4/.....	*	*	*	0.6	0.9	1.7	2.3	3.9	6.0	8.2
2005 4/.....	*	0.3	0.3	0.6	1.0	1.4	2.5	4.2	5.8	9.0
2004 4/.....	*	*	0.4	0.4	1.1	1.5	2.7	4.0	5.5	8.0
2003 4/.....	*	0.3	0.4	0.4	0.9	2.0	3.0	5.0	6.8	8.8
2002.....	*	0.3	0.6	0.9	1.5	1.7	3.1	5.9	7.5	9.3
2001.....	*	0.3	0.5	0.7	1.4	2.1	3.7	5.8	8.0	10.1
2000.....	0.3	0.4	0.5	0.9	1.4	2.9	4.4	5.6	8.6	10.3
1999.....	*	0.4	0.5	1.1	1.6	2.7	3.9	6.6	8.7	11.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Disorders of both mitral and aortic valves (I08.0)

All races 3/, both sexes

2006.....	*	*	*	0.1	0.2	0.3	0.8	1.4	2.4	4.9
2005.....	*	*	0.1	0.1	0.2	0.4	0.8	1.6	2.6	6.0
2004.....	*	*	*	0.1	0.2	0.5	0.7	1.5	2.6	5.2
2003.....	*	*	*	0.2	0.2	0.4	0.8	1.6	2.7	6.5
2002.....	*	*	*	*	*	0.3	0.7	1.3	1.8	3.5
2001.....	*	*	*	*	*	0.3	0.6	0.7	1.6	2.5
2000.....	*	*	*	*	0.2	0.2	0.5	0.8	1.0	2.0
1999.....	*	*	*	*	0.2	0.2	0.6	0.5	1.1	2.0

All races 3/, male

2006.....	*	*	*	*	*	*	0.5	1.2	2.2	4.2
2005.....	*	*	*	*	*	0.4	0.5	1.4	2.0	5.4
2004.....	*	*	*	*	*	0.6	*	1.5	3.0	3.8
2003.....	*	*	*	*	*	*	0.9	1.6	2.4	6.0
2002.....	*	*	*	*	*	*	0.7	1.4	1.5	2.4
2001.....	*	*	*	*	*	*	0.6	*	1.6	2.3
2000.....	*	*	*	*	*	*	*	*	*	1.9
1999.....	*	*	*	*	*	*	0.6	*	*	*

All races 3/, female

2006.....	*	*	*	*	*	0.4	1.1	1.5	2.5	5.2
2005.....	*	*	*	*	*	0.4	0.9	1.7	3.0	6.2
2004.....	*	*	*	*	*	0.4	0.9	1.5	2.3	5.9
2003.....	*	*	*	*	*	0.5	0.7	1.6	2.9	6.7
2002.....	*	*	*	*	*	*	0.6	1.2	2.1	4.0
2001.....	*	*	*	*	*	0.4	0.6	0.7	1.6	2.5
2000.....	*	*	*	*	*	*	0.5	0.9	1.2	2.1
1999.....	*	*	*	*	*	*	0.6	0.6	1.2	2.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of both mitral and aortic valves (I08.0)										
White, both sexes										
2006 4/.....	*	*	*	*	0.2	0.2	0.9	1.4	2.6	5.4
2005 4/.....	*	*	*	*	0.2	0.5	0.8	1.7	2.8	6.6
2004 4/.....	*	*	*	*	0.2	0.5	0.8	1.5	2.8	5.6
2003 4/.....	*	*	*	0.2	*	0.4	0.8	1.7	3.0	7.0
2002.....	*	*	*	*	*	0.4	0.7	1.4	1.9	3.8
2001.....	*	*	*	*	*	0.4	0.6	0.7	1.7	2.6
2000.....	*	*	*	*	*	*	0.5	0.8	1.1	2.0
1999.....	*	*	*	*	*	*	0.6	0.6	1.2	2.1
White, male										
2006 4/.....	*	*	*	*	*	*	*	1.2	2.4	4.7
2005 4/.....	*	*	*	*	*	*	*	1.5	2.1	6.0
2004 4/.....	*	*	*	*	*	0.6	*	1.5	3.2	4.1
2003 4/.....	*	*	*	*	*	*	0.9	1.8	2.7	6.4
2002.....	*	*	*	*	*	*	0.7	1.4	1.5	2.7
2001.....	*	*	*	*	*	*	0.6	*	1.7	2.5
2000.....	*	*	*	*	*	*	*	*	*	1.9
1999.....	*	*	*	*	*	*	0.7	*	*	*
White, female										
2006 4/.....	*	*	*	*	*	*	1.1	1.5	2.7	5.7
2005 4/.....	*	*	*	*	*	0.5	1.0	1.8	3.3	6.9
2004 4/.....	*	*	*	*	*	*	1.0	1.5	2.5	6.3
2003 4/.....	*	*	*	*	*	0.4	0.7	1.6	3.1	7.2
2002.....	*	*	*	*	*	*	0.6	1.4	2.2	4.2
2001.....	*	*	*	*	*	0.5	0.6	0.8	1.6	2.6
2000.....	*	*	*	*	*	*	0.6	1.0	1.3	2.1
1999.....	*	*	*	*	*	*	0.6	0.6	1.2	2.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)

All races 3/, both sexes

2006.....	*	0.1	0.2	0.2	0.4	0.8	1.2	2.0	3.2	4.6
2005.....	*	0.1	0.1	0.2	0.4	0.8	1.4	1.7	2.9	5.2
2004.....	0.1	0.1	0.2	0.2	0.4	0.8	1.2	2.2	3.1	4.4
2003.....	*	0.1	0.2	0.3	0.5	0.8	1.4	2.1	2.9	5.4
2002.....	*	0.1	0.2	0.4	0.6	1.0	1.5	2.2	3.2	5.6
2001.....	0.1	0.1	0.2	0.4	0.5	1.0	1.4	2.7	3.4	5.9
2000.....	0.1	0.2	0.2	0.4	0.7	1.1	1.6	2.9	3.7	5.8
1999.....	0.1	0.1	0.2	0.4	0.7	1.1	1.8	2.7	4.3	6.6

All races 3/, male

2006.....	*	*	*	*	*	0.6	0.8	1.1	1.6	3.0
2005.....	*	*	*	0.2	*	0.5	1.0	1.0	1.4	3.6
2004.....	*	*	*	*	*	0.6	0.7	1.6	2.1	2.5
2003.....	*	*	*	*	*	0.5	0.9	1.5	1.8	3.4
2002.....	*	*	*	0.4	0.4	0.6	1.0	1.5	2.4	3.8
2001.....	*	*	*	0.4	0.4	0.7	0.8	2.2	1.8	3.6
2000.....	*	*	*	*	0.5	0.7	0.9	1.9	2.2	4.0
1999.....	*	*	*	*	0.4	0.7	1.6	1.6	3.2	3.7

All races 3/, female

2006.....	*	*	0.2	0.3	0.6	0.9	1.5	2.7	4.2	5.3
2005.....	*	*	*	0.2	0.6	1.0	1.7	2.2	3.8	5.9
2004.....	*	*	0.2	0.3	0.5	1.1	1.6	2.7	3.7	5.3
2003.....	*	*	*	0.4	0.7	1.1	1.7	2.6	3.5	6.3
2002.....	*	*	*	0.4	0.8	1.3	1.9	2.7	3.8	6.3
2001.....	*	*	*	0.4	0.6	1.2	2.0	3.1	4.4	6.8
2000.....	*	0.2	0.3	0.5	0.8	1.4	2.2	3.5	4.6	6.6
1999.....	*	*	0.3	0.6	0.9	1.5	2.0	3.5	5.0	7.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic rheumatic heart diseases (I07,I08.1-I08.9,I09)

White, both sexes

2006 4/.....	*	*	0.1	0.2	0.4	0.8	1.2	2.1	3.3	4.9
2005 4/.....	*	*	*	0.2	0.4	0.8	1.4	1.7	3.0	5.4
2004 4/.....	*	*	0.2	0.2	0.4	0.8	1.2	2.4	3.3	4.7
2003 4/.....	*	0.1	0.1	0.3	0.5	0.9	1.3	2.3	3.1	5.7
2002.....	*	*	0.1	0.4	0.6	1.0	1.6	2.3	3.4	5.9
2001.....	*	*	*	0.3	0.5	1.0	1.5	2.8	3.6	6.3
2000.....	*	0.1	0.2	0.3	0.7	1.1	1.7	3.1	3.8	6.0
1999.....	*	*	0.2	0.4	0.6	1.2	1.9	2.8	4.6	6.9

White, male

2006 4/.....	*	*	*	*	*	0.6	0.7	1.2	1.7	3.2
2005 4/.....	*	*	*	*	*	*	1.0	1.0	1.5	3.5
2004 4/.....	*	*	*	*	*	0.5	0.7	1.8	2.2	2.6
2003 4/.....	*	*	*	*	*	0.5	1.0	1.6	2.0	3.5
2002.....	*	*	*	0.3	*	0.5	1.1	1.6	2.6	4.0
2001.....	*	*	*	0.4	*	0.7	0.8	2.3	1.8	3.7
2000.....	*	*	*	*	0.5	0.7	0.9	2.0	2.3	4.1
1999.....	*	*	*	*	*	0.8	1.7	1.7	3.4	3.9

White, female

2006 4/.....	*	*	*	0.3	0.5	1.0	1.6	2.9	4.4	5.7
2005 4/.....	*	*	*	*	0.6	1.1	1.7	2.3	4.0	6.2
2004 4/.....	*	*	*	0.3	0.5	1.0	1.7	2.8	4.0	5.6
2003 4/.....	*	*	*	0.4	0.7	1.2	1.6	2.8	3.7	6.7
2002.....	*	*	*	0.4	0.8	1.4	2.0	2.8	3.9	6.7
2001.....	*	*	*	*	0.6	1.3	2.0	3.2	4.7	7.3
2000.....	*	*	*	0.4	0.8	1.4	2.4	3.8	4.7	6.8
1999.....	*	*	0.3	0.6	0.9	1.6	2.2	3.6	5.2	8.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive diseases (I10-I15)										
All races 3/, both sexes										
2006.....	5.4	9.2	14.5	17.9	24.7	32.9	46.5	76.1	142.5	406.0
2005.....	5.5	9.1	13.1	17.4	24.1	33.3	48.9	81.3	150.8	433.9
2004.....	5.2	9.4	12.9	17.8	24.3	33.3	49.5	79.7	146.8	416.1
2003.....	5.2	8.9	13.3	17.6	24.0	34.0	49.3	83.1	146.3	409.5
2002.....	4.7	8.1	12.0	16.5	22.8	32.2	47.5	77.8	142.1	390.8
2001.....	4.5	7.2	11.1	14.9	21.2	30.3	45.3	77.2	140.5	384.7
2000.....	4.3	6.8	10.0	14.5	21.9	29.5	45.1	75.2	138.9	380.3
1999.....	4.0	6.6	9.4	14.6	20.6	30.4	43.8	75.7	136.9	366.0
All races 3/, male										
2006.....	7.0	12.7	19.6	23.9	32.3	40.7	53.2	81.6	145.3	326.0
2005.....	7.2	12.3	18.0	23.7	31.4	40.5	54.8	86.6	146.5	344.2
2004.....	6.8	12.4	17.6	23.3	31.9	41.4	55.6	84.3	144.0	335.7
2003.....	6.8	12.0	17.8	23.0	29.3	41.8	54.8	85.9	139.5	322.4
2002.....	6.4	10.8	16.0	21.9	28.6	37.8	52.5	82.0	136.9	311.1
2001.....	6.2	9.8	14.6	18.8	25.8	36.4	50.0	80.0	134.2	309.4
2000.....	5.6	9.0	13.8	18.4	27.5	33.8	51.2	76.7	135.5	309.3
1999.....	5.4	8.6	12.2	18.7	24.7	35.1	47.9	79.8	131.9	297.3
All races 3/, female										
2006.....	3.8	5.8	9.5	12.3	17.7	26.2	41.1	72.1	140.8	443.4
2005.....	3.7	5.8	8.3	11.5	17.3	27.1	44.2	77.5	153.5	475.1
2004.....	3.6	6.6	8.5	12.6	17.4	26.3	44.6	76.4	148.6	452.3
2003.....	3.7	5.8	9.0	12.4	19.1	27.3	44.8	81.1	150.4	448.0
2002.....	3.1	5.5	8.1	11.5	17.5	27.5	43.5	74.8	145.3	425.4
2001.....	2.9	4.7	7.7	11.3	17.0	25.2	41.5	75.2	144.3	416.3
2000.....	3.0	4.7	6.4	10.8	16.9	25.8	40.3	74.1	140.9	409.3
1999.....	2.7	4.7	6.7	10.8	16.9	26.3	40.6	72.9	139.8	393.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive diseases (I10-I15)										
White, both sexes										
2006 4/.....	3.7	6.5	10.1	13.3	18.6	25.1	36.8	64.5	129.1	387.3
2005 4/.....	3.6	6.1	8.9	12.7	18.4	25.1	39.0	69.3	135.4	413.1
2004 4/.....	3.5	6.3	8.8	12.5	17.8	25.9	39.8	67.5	132.1	394.8
2003 4/.....	3.4	5.7	9.0	12.1	17.8	25.4	38.9	70.4	133.7	389.7
2002.....	3.1	5.4	8.1	11.5	16.5	24.3	38.0	65.5	126.8	368.7
2001.....	2.7	4.5	7.3	10.2	15.2	22.3	35.8	65.7	126.0	364.3
2000.....	2.8	4.3	6.4	9.6	15.3	21.9	35.2	63.1	123.9	357.1
1999.....	2.3	4.0	6.0	9.6	14.4	22.3	33.9	62.7	121.1	343.4
White, male										
2006 4/.....	5.2	9.5	14.5	18.0	24.9	30.7	41.5	68.6	130.3	308.3
2005 4/.....	5.1	8.9	12.7	17.7	24.4	31.0	43.0	73.1	131.7	322.5
2004 4/.....	5.0	8.8	12.6	16.9	23.7	32.6	45.4	71.1	130.1	318.7
2003 4/.....	4.9	8.2	12.8	16.7	22.4	32.0	43.9	73.7	127.4	301.5
2002.....	4.6	7.7	11.4	16.0	21.1	28.7	42.7	69.5	121.3	290.6
2001.....	4.0	6.3	10.0	13.5	19.2	27.5	40.3	68.4	119.3	288.6
2000.....	3.8	6.2	9.3	12.8	19.7	25.8	40.8	65.4	120.1	286.2
1999.....	3.5	5.6	8.1	12.9	17.9	26.2	37.7	66.4	116.4	273.5
White, female										
2006 4/.....	2.2	3.5	5.8	8.7	12.8	20.1	32.9	61.4	128.4	424.4
2005 4/.....	2.1	3.3	5.1	7.9	12.8	19.9	35.7	66.4	137.8	454.8
2004 4/.....	2.1	3.7	5.1	8.3	12.3	20.0	35.2	64.8	133.4	429.1
2003 4/.....	1.9	3.1	5.3	7.7	13.6	19.5	34.8	68.0	137.5	428.7
2002.....	1.6	3.2	4.8	7.1	12.2	20.4	34.2	62.6	130.2	402.6
2001.....	1.4	2.7	4.6	7.1	11.6	17.8	32.2	63.7	130.1	396.0
2000.....	1.7	2.4	3.5	6.6	11.2	18.5	30.6	61.5	126.1	385.8
1999.....	1.2	2.5	3.9	6.5	11.2	19.0	30.9	60.1	123.9	371.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive diseases (I10-I15)										
Black, both sexes										
2006 4/.....	17.4	29.7	48.9	59.8	83.0	106.3	143.2	207.3	316.4	659.7
2005 4/.....	18.0	31.0	46.5	59.9	80.3	111.1	147.0	218.1	350.0	721.9
2004 4/.....	17.1	33.1	46.6	65.2	86.0	104.8	150.5	222.2	337.4	698.5
2003 4/.....	18.2	33.2	48.7	67.7	82.8	113.4	157.0	231.9	315.3	686.5
2002.....	16.3	28.9	44.5	64.1	81.0	105.7	150.4	224.0	341.7	685.4
2001.....	17.2	28.3	43.3	58.1	76.4	104.6	145.9	213.3	325.6	664.2
2000.....	15.3	26.9	40.6	59.6	82.1	103.2	152.4	221.1	326.4	673.8
1999.....	16.3	26.7	39.8	60.1	75.9	105.6	151.7	230.5	335.2	639.9
Black, male										
2006 4/.....	20.8	39.0	63.6	80.4	111.7	142.5	180.9	249.7	359.4	583.5
2005 4/.....	22.9	39.7	63.7	82.5	109.4	138.6	180.6	259.7	356.8	640.5
2004 4/.....	20.7	40.8	61.3	86.2	117.2	136.4	173.2	258.9	343.6	572.2
2003 4/.....	21.6	43.0	62.5	87.0	102.4	143.3	186.2	247.4	321.1	612.0
2002.....	20.3	36.6	57.8	82.5	104.1	126.9	170.9	247.6	354.6	602.5
2001.....	22.3	38.1	56.4	72.4	92.5	129.1	167.0	230.5	349.1	604.8
2000.....	19.5	33.2	54.3	75.6	103.7	119.4	173.5	224.0	349.8	606.6
1999.....	20.3	33.9	51.4	75.3	92.3	126.7	169.6	251.7	351.4	589.4
Black, female										
2006 4/.....	14.4	21.5	36.4	42.8	60.6	79.6	117.3	181.9	293.7	690.9
2005 4/.....	13.6	23.4	32.0	41.2	57.5	90.8	124.0	193.3	346.5	754.7
2004 4/.....	13.9	26.5	34.1	47.9	61.6	81.3	135.1	200.5	334.1	748.6
2003 4/.....	15.2	24.7	37.0	51.9	67.5	91.2	137.4	222.7	312.2	715.4
2002.....	12.8	22.2	33.2	49.0	62.9	89.9	136.7	209.9	335.0	717.1
2001.....	12.7	19.7	32.2	46.3	63.7	86.5	131.8	203.0	313.6	686.2
2000.....	11.5	21.5	28.9	46.6	65.0	91.2	138.3	219.3	314.6	698.8
1999.....	12.7	20.5	30.1	47.6	63.1	90.0	139.7	217.7	327.1	658.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension (I10)										
All races 3/, both sexes										
2006.....	0.6	1.2	2.4	3.2	5.2	7.7	11.7	19.9	38.4	119.3
2005.....	0.6	1.1	1.9	3.0	4.8	7.6	12.0	22.1	43.0	137.5
2004.....	0.5	1.2	2.0	2.8	4.6	6.9	11.6	20.8	38.9	127.4
2003.....	0.5	1.0	2.0	2.9	4.6	7.2	11.6	20.5	39.5	121.9
2002.....	0.5	1.0	1.6	2.5	4.2	6.6	10.9	18.5	37.1	112.0
2001.....	0.4	1.0	1.6	2.7	4.3	6.9	10.1	19.1	35.6	107.4
2000.....	0.5	1.1	1.7	2.6	4.3	6.1	10.4	18.2	33.1	101.1
1999.....	0.4	0.9	1.6	2.5	4.3	6.7	9.7	18.0	33.0	98.1
All races 3/, male										
2006.....	0.8	1.5	3.2	4.1	6.6	9.2	13.2	19.0	35.4	80.3
2005.....	0.7	1.4	2.5	3.8	6.3	9.1	13.0	22.0	37.8	88.0
2004.....	0.6	1.5	2.7	3.2	5.9	8.4	11.9	20.7	35.3	86.5
2003.....	0.6	1.3	2.6	3.6	5.6	8.9	12.3	18.1	33.5	77.4
2002.....	0.7	1.3	2.1	3.3	5.0	7.3	11.9	17.5	32.0	74.2
2001.....	0.5	1.2	2.2	3.4	5.2	7.9	10.9	18.8	30.1	74.8
2000.....	0.6	1.5	2.3	3.3	5.4	6.5	11.7	17.4	29.5	70.3
1999.....	0.5	1.1	1.9	3.2	5.1	7.8	10.7	17.2	29.7	68.1
All races 3/, female										
2006.....	0.4	0.9	1.6	2.3	3.9	6.3	10.5	20.5	40.2	137.6
2005.....	0.6	0.9	1.2	2.2	3.5	6.3	11.2	22.2	46.2	160.2
2004.....	0.3	1.0	1.3	2.4	3.5	5.6	11.3	20.8	41.1	145.8
2003.....	0.5	0.6	1.4	2.2	3.7	5.7	11.0	22.1	43.1	141.5
2002.....	0.4	0.7	1.2	1.8	3.5	6.0	10.2	19.1	40.2	128.3
2001.....	0.4	0.7	1.1	2.0	3.5	6.1	9.4	19.3	38.9	121.1
2000.....	0.4	0.6	1.2	1.9	3.4	5.7	9.4	18.7	35.2	113.6
1999.....	0.3	0.7	1.3	1.9	3.6	5.7	8.9	18.6	34.9	110.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension (I10)										
White, both sexes										
2006 4/.....	0.4	0.9	1.8	2.5	4.4	6.4	10.6	18.1	35.7	114.8
2005 4/.....	0.5	0.9	1.4	2.4	4.2	6.6	10.5	19.9	40.9	134.3
2004 4/.....	0.4	0.9	1.6	2.2	4.0	5.9	10.3	18.9	36.9	123.5
2003 4/.....	0.4	0.7	1.6	2.2	3.8	5.9	9.6	18.3	36.6	119.9
2002.....	0.4	0.7	1.3	2.1	3.7	5.7	9.5	16.4	35.1	108.6
2001.....	0.3	0.7	1.2	2.1	3.6	5.7	8.6	17.6	33.1	104.6
2000.....	0.4	0.8	1.2	2.1	3.5	5.3	9.2	16.2	31.2	97.5
1999.....	0.3	0.7	1.1	2.0	3.6	5.6	8.7	16.0	30.9	94.3
White, male										
2006 4/.....	0.6	1.3	2.6	3.3	5.9	7.7	11.5	17.2	32.2	76.9
2005 4/.....	0.6	1.2	2.0	3.1	5.4	7.9	11.1	19.1	35.3	83.8
2004 4/.....	0.5	1.2	2.3	2.5	5.2	7.4	10.6	18.4	33.5	84.0
2003 4/.....	0.5	1.0	2.2	3.0	4.5	7.6	10.5	15.9	29.7	74.3
2002.....	0.5	1.0	1.7	2.8	4.4	6.3	10.4	15.6	29.1	71.0
2001.....	0.4	0.9	1.6	2.6	4.4	6.8	9.2	16.9	26.9	71.1
2000.....	0.5	1.1	1.7	2.7	4.3	5.8	10.3	15.6	26.8	66.4
1999.....	0.4	0.9	1.4	2.6	4.3	6.7	9.5	15.3	27.0	62.7
White, female										
2006 4/.....	0.2	0.6	1.1	1.8	3.1	5.3	9.8	18.7	37.9	132.6
2005 4/.....	0.3	0.5	0.9	1.6	3.1	5.4	10.1	20.6	44.5	157.6
2004 4/.....	0.2	0.5	0.9	1.8	2.8	4.7	10.1	19.2	39.0	141.3
2003 4/.....	0.4	0.5	1.0	1.6	3.1	4.5	8.9	20.0	40.8	140.2
2002.....	0.3	0.4	0.9	1.4	3.0	5.1	8.7	17.1	38.7	124.9
2001.....	*	0.4	0.8	1.6	2.7	4.8	8.1	18.1	36.9	118.7
2000.....	0.3	0.4	0.8	1.5	2.8	4.8	8.4	16.7	33.8	110.2
1999.....	*	0.5	0.8	1.4	2.9	4.7	8.0	16.5	33.2	107.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Essential (primary) hypertension (I10)										
Black, both sexes										
2006 4/.....	1.8	3.3	7.0	8.9	13.2	19.6	24.4	40.5	73.6	179.0
2005 4/.....	1.8	3.1	5.2	8.7	12.0	18.6	27.5	45.9	72.1	188.4
2004 4/.....	1.3	3.8	5.4	8.5	11.5	16.5	25.0	44.5	64.9	183.6
2003 4/.....	1.4	2.7	5.3	8.9	12.9	19.6	32.3	47.0	78.9	157.5
2002.....	1.5	3.4	4.7	6.9	9.1	15.4	27.5	43.6	67.0	158.6
2001.....	1.6	3.4	4.9	8.3	11.9	18.2	25.8	39.4	66.6	150.8
2000.....	1.4	3.2	5.9	7.9	11.9	14.7	23.6	41.4	57.0	149.2
1999.....	1.4	2.4	6.0	8.0	11.2	16.1	22.2	42.3	58.9	143.2
Black, male										
2006 4/.....	2.3	3.7	9.0	11.8	15.9	24.6	34.4	42.1	80.1	119.7
2005 4/.....	1.7	3.2	6.5	11.1	17.1	22.3	34.2	59.5	73.0	135.7
2004 4/.....	1.5	3.8	7.0	10.2	13.8	19.4	26.5	53.2	62.4	124.2
2003 4/.....	1.5	3.7	6.8	10.5	17.2	24.4	32.8	48.4	87.1	119.9
2002.....	1.8	4.0	5.7	8.8	10.3	17.2	30.4	44.1	73.4	117.9
2001.....	1.7	3.8	6.9	11.3	13.6	20.3	31.3	46.0	72.2	132.8
2000.....	*	4.5	7.7	10.7	15.9	14.6	28.4	39.7	64.9	123.4
1999.....	1.7	2.6	6.3	9.5	13.6	19.0	26.5	41.3	66.3	124.2
Black, female										
2006 4/.....	1.4	3.0	5.2	6.5	11.2	15.9	17.5	39.6	70.2	203.3
2005 4/.....	2.0	3.1	4.0	6.7	8.1	15.9	22.9	37.9	71.6	209.7
2004 4/.....	*	3.7	4.1	7.1	9.6	14.4	23.9	39.4	66.1	207.1
2003 4/.....	1.4	1.8	4.0	7.6	9.6	16.1	31.9	46.1	74.7	172.2
2002.....	*	2.9	3.9	5.4	8.2	14.1	25.5	43.4	63.7	174.2
2001.....	1.5	3.0	3.1	5.9	10.6	16.7	22.2	35.5	63.8	157.5
2000.....	1.4	2.1	4.5	5.5	8.7	14.8	20.4	42.4	53.1	158.8
1999.....	*	2.1	5.7	6.8	9.3	14.0	19.4	42.9	55.3	150.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
All races 3/, both sexes										
2006.....	4.0	6.8	10.0	11.8	14.9	18.2	23.1	35.2	64.2	194.0
2005.....	4.0	6.5	9.2	11.7	14.7	17.6	23.8	35.8	65.9	198.6
2004.....	4.0	6.8	9.0	12.2	14.9	18.8	25.1	36.7	66.1	192.8
2003.....	3.9	6.6	9.4	11.7	14.9	19.4	25.1	39.5	66.8	195.0
2002.....	3.5	5.9	8.6	11.1	14.3	18.9	24.0	37.9	66.0	187.0
2001.....	3.2	5.1	7.7	9.7	12.5	16.3	23.5	36.6	66.0	189.5
2000.....	2.9	4.7	6.6	9.3	13.2	16.5	23.0	36.3	66.1	193.1
1999.....	2.8	4.5	6.2	9.6	11.9	16.3	22.1	36.4	64.7	184.9
All races 3/, male										
2006.....	5.2	9.8	13.8	16.2	20.4	24.0	27.8	40.2	63.1	143.0
2005.....	5.4	9.1	13.0	16.5	20.3	22.5	28.5	39.3	62.2	147.5
2004.....	5.2	9.2	12.4	17.0	20.4	24.7	30.5	39.1	60.9	145.4
2003.....	5.2	9.0	12.8	16.0	18.8	25.2	29.4	42.8	63.3	142.2
2002.....	4.9	8.0	11.9	15.2	19.0	23.6	27.0	41.0	62.2	134.6
2001.....	4.5	7.0	10.3	12.8	16.5	20.5	27.1	38.9	63.5	134.4
2000.....	3.9	6.3	9.4	12.1	17.4	20.7	27.2	37.3	61.8	139.1
1999.....	3.9	6.1	8.1	12.7	15.0	19.5	24.6	39.4	59.2	133.6
All races 3/, female										
2006.....	2.8	3.8	6.3	7.5	9.8	13.2	19.2	31.4	64.9	217.9
2005.....	2.5	3.9	5.5	7.2	9.7	13.3	20.0	33.2	68.2	222.0
2004.....	2.7	4.4	5.8	7.6	9.9	13.5	20.7	35.0	69.3	214.1
2003.....	2.6	4.4	6.1	7.7	11.2	14.4	21.6	37.2	69.0	218.3
2002.....	2.2	3.9	5.4	7.2	10.0	14.8	21.6	35.8	68.2	209.7
2001.....	2.0	3.2	5.1	6.8	8.9	12.7	20.6	35.1	67.5	212.6
2000.....	2.0	3.2	3.8	6.6	9.3	12.9	19.6	35.7	68.6	215.1
1999.....	1.7	3.0	4.3	6.8	9.2	13.5	20.2	34.3	67.9	205.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
White, both sexes										
2006 4/.....	3.0	5.0	7.3	9.0	11.5	14.2	17.9	29.8	58.1	186.9
2005 4/.....	2.8	4.6	6.5	8.9	11.7	13.5	19.4	30.4	58.0	188.9
2004 4/.....	2.8	4.8	6.4	9.0	11.2	15.1	20.4	31.0	59.1	183.7
2003 4/.....	2.7	4.4	6.6	8.5	11.4	14.7	20.2	33.2	61.2	185.3
2002.....	2.4	4.3	6.0	8.1	10.5	14.4	19.3	31.9	58.1	177.3
2001.....	2.1	3.3	5.2	6.9	9.1	12.1	18.5	30.9	59.2	179.9
2000.....	2.0	3.1	4.3	6.2	9.4	12.5	17.6	30.2	58.4	181.8
1999.....	1.8	2.9	4.1	6.5	8.7	12.0	16.9	30.0	57.3	175.3
White, male										
2006 4/.....	4.2	7.6	10.8	12.8	15.8	18.2	21.7	33.4	56.2	134.5
2005 4/.....	4.0	6.9	9.6	12.8	16.4	17.7	22.9	33.6	54.7	136.9
2004 4/.....	4.1	7.0	9.3	12.8	15.3	20.0	25.0	33.4	54.3	136.4
2003 4/.....	4.0	6.4	9.7	12.2	15.0	19.4	23.6	36.1	57.9	131.9
2002.....	3.7	6.1	8.9	11.6	14.0	18.1	22.0	34.5	54.8	124.8
2001.....	3.1	4.8	7.5	9.6	12.5	15.9	21.9	32.9	56.1	123.7
2000.....	2.8	4.6	6.5	8.6	12.7	16.0	21.5	31.1	53.4	129.3
1999.....	2.7	4.2	5.8	8.9	11.3	14.4	19.1	32.4	51.8	123.7
White, female										
2006 4/.....	1.7	2.4	3.9	5.3	7.5	10.5	14.7	27.1	59.3	211.5
2005 4/.....	1.5	2.3	3.6	5.2	7.3	9.7	16.5	28.0	60.1	212.9
2004 4/.....	1.6	2.7	3.5	5.3	7.4	10.8	16.6	29.3	62.1	205.0
2003 4/.....	1.3	2.4	3.5	5.0	8.1	10.7	17.4	31.1	63.3	209.0
2002.....	1.2	2.4	3.3	4.7	7.1	11.1	17.1	30.0	60.1	200.1
2001.....	1.0	1.8	3.0	4.4	6.0	8.8	15.8	29.5	61.1	203.5
2000.....	1.2	1.6	2.2	4.0	6.3	9.4	14.5	29.6	61.3	203.1
1999.....	0.8	1.6	2.5	4.2	6.3	10.0	15.1	28.2	60.6	196.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart disease (I11)										
Black, both sexes										
2006 4/.....	11.3	20.4	31.3	37.3	48.4	56.9	75.0	96.8	145.3	299.6
2005 4/.....	12.2	20.8	30.3	37.8	46.0	57.8	69.9	100.9	168.5	338.1
2004 4/.....	12.3	22.0	31.3	41.6	49.9	55.4	76.2	103.1	161.4	314.7
2003 4/.....	12.8	23.9	32.8	41.9	48.3	63.0	77.1	113.3	145.7	332.3
2002.....	11.4	19.3	30.3	39.7	50.5	61.2	76.5	110.3	167.7	320.4
2001.....	11.5	18.6	28.4	35.7	43.6	55.8	77.8	105.1	154.7	326.1
2000.....	9.9	17.8	25.4	37.6	47.9	54.6	81.6	111.1	164.9	340.0
1999.....	10.2	17.9	24.8	38.0	41.6	57.6	79.0	113.7	161.2	305.1
Black, male										
2006 4/.....	12.9	28.0	40.6	50.6	70.9	84.1	94.7	129.8	165.0	272.6
2005 4/.....	16.3	27.1	42.3	53.5	64.9	74.2	91.1	118.9	169.2	301.2
2004 4/.....	14.5	27.8	40.8	59.1	72.3	78.3	97.8	114.6	158.5	267.2
2003 4/.....	14.9	29.8	41.5	55.1	59.8	86.7	99.9	129.6	149.1	287.8
2002.....	14.6	24.4	39.9	53.1	71.1	79.0	89.8	125.4	167.4	267.0
2001.....	14.8	25.1	36.6	45.1	55.9	69.6	91.6	117.1	171.8	284.8
2000.....	12.7	21.4	35.8	47.9	63.8	70.0	95.9	117.2	178.7	273.3
1999.....	13.1	22.5	31.3	49.5	52.0	73.0	90.2	129.2	166.5	263.6
Black, female										
2006 4/.....	9.9	13.7	23.4	26.4	31.0	36.9	61.5	77.1	134.9	310.7
2005 4/.....	8.6	15.3	20.2	24.9	31.3	45.7	55.4	90.1	168.0	353.0
2004 4/.....	10.3	17.0	23.2	27.2	32.5	38.5	61.5	96.2	162.9	333.5
2003 4/.....	11.0	18.7	25.5	31.0	39.4	45.5	61.8	103.6	143.9	349.6
2002.....	8.6	14.9	22.3	28.7	34.4	47.9	67.7	101.3	167.9	340.9
2001.....	8.6	13.0	21.5	28.0	33.9	45.6	68.6	97.9	145.9	341.4
2000.....	7.5	14.6	16.5	29.2	35.3	43.2	72.1	107.4	157.9	364.9
1999.....	7.6	13.8	19.3	28.7	33.4	46.2	71.6	104.4	158.6	320.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive renal disease and secondary hypertension (I12,I15)										
All races 3/, both sexes										
2006.....	0.6	0.9	1.6	2.3	3.6	5.6	9.3	16.8	31.3	70.0
2005.....	0.6	1.1	1.5	2.1	3.5	6.3	10.3	18.4	32.5	72.5
2004.....	0.5	1.0	1.4	2.2	3.5	6.0	10.3	17.8	32.4	71.1
2003.....	0.5	0.9	1.3	2.1	3.4	5.9	9.8	18.0	30.4	67.0
2002.....	0.5	0.8	1.2	2.0	3.1	5.2	9.7	16.5	29.6	68.4
2001.....	0.6	0.9	1.3	1.7	3.2	5.3	8.9	16.4	29.6	64.5
2000.....	0.5	0.7	1.3	1.8	3.5	5.4	8.7	15.7	29.9	61.8
1999.....	0.5	0.8	1.2	1.6	3.0	5.3	8.9	14.7	26.9	53.9
All races 3/, male										
2006.....	0.7	1.1	1.9	2.6	4.0	6.0	9.6	17.6	36.8	79.0
2005.....	0.8	1.3	1.8	2.6	3.7	6.8	10.3	19.9	36.7	81.4
2004.....	0.6	1.2	1.7	2.3	3.9	6.5	10.4	19.4	37.4	79.5
2003.....	0.7	1.2	1.7	2.3	3.7	6.1	10.2	19.4	32.6	76.1
2002.....	0.6	0.9	1.4	2.2	3.2	5.4	10.3	18.4	32.7	79.2
2001.....	0.8	1.1	1.5	1.8	2.9	5.8	9.0	17.2	31.2	76.0
2000.....	0.7	0.8	1.6	2.1	3.7	5.0	8.9	17.0	33.4	72.0
1999.....	0.6	0.9	1.6	1.8	3.1	5.6	9.4	15.9	29.7	64.3
All races 3/, female										
2006.....	0.5	0.8	1.3	2.1	3.2	5.3	9.1	16.2	27.9	65.9
2005.....	0.4	0.9	1.1	1.6	3.3	5.9	10.2	17.2	29.8	68.5
2004.....	0.4	0.8	1.0	2.0	3.1	5.7	10.2	16.6	29.3	67.4
2003.....	0.4	0.6	1.0	1.8	3.2	5.7	9.4	17.0	29.1	63.0
2002.....	0.4	0.6	1.1	1.9	2.9	5.0	9.3	15.1	27.7	63.7
2001.....	0.4	0.7	1.2	1.7	3.4	5.0	8.8	15.9	28.6	59.7
2000.....	0.4	0.6	1.0	1.6	3.3	5.7	8.5	14.7	27.8	57.7
1999.....	0.4	0.7	0.8	1.4	2.9	5.1	8.5	13.8	25.3	49.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive renal disease and secondary hypertension (I12,I15)										
White, both sexes										
2006 4/.....	0.2	0.4	0.7	1.4	2.2	3.7	6.8	13.5	27.7	64.7
2005 4/.....	0.3	0.5	0.7	1.2	2.0	4.0	7.1	15.0	28.6	65.9
2004 4/.....	0.2	0.4	0.6	1.1	1.8	3.9	7.3	14.0	28.0	64.6
2003 4/.....	0.3	0.4	0.7	1.0	2.0	3.8	7.0	14.7	27.5	61.1
2002.....	0.2	0.3	0.6	1.0	1.7	3.3	7.3	13.3	25.3	61.9
2001.....	0.2	0.4	0.6	0.8	1.8	3.4	6.7	13.3	25.5	58.7
2000.....	0.2	0.2	0.6	0.9	1.9	3.3	6.1	12.7	26.3	55.7
1999.....	0.3	0.3	0.5	0.7	1.4	3.5	6.5	11.5	22.4	47.7
White, male										
2006 4/.....	0.2	0.5	0.9	1.5	2.6	3.9	6.8	14.7	32.9	74.4
2005 4/.....	0.4	0.6	0.8	1.4	2.1	4.3	6.9	16.3	33.6	76.3
2004 4/.....	0.3	0.5	0.7	1.3	2.0	4.2	7.6	15.3	33.1	75.4
2003 4/.....	0.4	0.6	0.7	1.1	2.1	4.0	7.5	16.8	30.6	71.0
2002.....	0.3	0.4	0.7	1.1	1.9	3.5	7.9	15.2	28.6	74.2
2001.....	0.3	0.5	0.6	0.9	1.5	3.5	7.2	14.5	27.6	71.1
2000.....	0.3	0.3	0.8	1.1	2.2	3.0	6.5	14.3	30.6	66.1
1999.....	0.3	0.4	0.6	0.9	1.5	3.7	7.1	12.8	25.5	58.3
White, female										
2006 4/.....	*	0.4	0.6	1.3	1.8	3.4	6.9	12.6	24.4	60.1
2005 4/.....	*	0.4	0.6	0.9	2.0	3.8	7.3	14.1	25.4	61.2
2004 4/.....	*	0.4	0.5	0.9	1.6	3.7	7.0	13.1	24.9	59.8
2003 4/.....	*	0.3	0.6	0.9	1.9	3.7	6.6	13.1	25.6	56.6
2002.....	*	0.3	0.5	0.9	1.5	3.2	6.9	12.0	23.3	56.5
2001.....	*	0.3	0.5	0.8	2.0	3.4	6.4	12.4	24.2	53.5
2000.....	*	*	0.3	0.7	1.7	3.5	5.8	11.5	23.6	51.4
1999.....	*	0.3	0.4	0.6	1.4	3.2	6.0	10.6	20.7	43.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive renal disease and secondary hypertension (I12,I15)										
Black, both sexes										
2006 4/.....	3.3	4.4	8.0	10.2	15.6	23.3	32.7	53.1	76.4	139.3
2005 4/.....	2.8	5.4	7.7	9.7	16.6	26.0	38.9	54.8	81.1	152.6
2004 4/.....	2.2	4.6	7.0	11.3	18.2	25.3	39.7	59.4	85.2	152.0
2003 4/.....	2.4	4.4	6.8	11.6	16.1	23.3	36.6	55.1	66.5	143.5
2002.....	2.3	4.0	6.1	11.5	15.2	21.1	34.1	52.7	84.0	150.3
2001.....	2.8	4.4	7.7	9.7	15.1	22.1	30.4	50.4	81.5	137.3
2000.....	2.8	4.0	6.8	10.1	17.1	25.6	35.2	51.0	73.4	134.2
1999.....	2.6	4.0	6.8	9.0	15.9	21.9	35.0	52.7	79.7	126.3
Black, male										
2006 4/.....	4.1	5.5	10.5	13.0	17.3	25.5	38.1	55.0	91.0	151.9
2005 4/.....	3.6	6.7	10.8	13.2	19.4	30.4	44.9	64.4	81.5	153.4
2004 4/.....	2.7	6.0	9.7	12.5	22.4	30.5	39.5	70.5	95.4	139.1
2003 4/.....	3.0	6.1	9.6	13.7	19.1	25.6	41.7	53.9	60.5	154.3
2002.....	2.9	5.1	7.6	13.0	15.4	23.4	36.7	61.1	90.9	156.5
2001.....	3.8	6.2	9.4	10.5	16.7	28.5	30.6	49.3	85.9	143.0
2000.....	3.9	4.6	7.9	11.9	18.4	25.5	35.4	53.4	72.5	146.5
1999.....	2.9	5.2	10.3	10.0	17.4	24.1	35.6	57.5	88.9	141.8
Black, female										
2006 4/.....	2.5	3.4	5.8	7.8	14.2	21.6	29.0	52.0	68.7	134.2
2005 4/.....	2.0	4.3	5.0	6.8	14.3	22.7	34.8	49.1	80.9	152.3
2004 4/.....	1.9	3.3	4.6	10.2	14.9	21.4	39.8	52.8	79.8	157.1
2003 4/.....	1.9	2.9	4.4	9.8	13.7	21.6	33.2	55.9	69.6	139.4
2002.....	1.8	2.9	4.9	10.3	15.0	19.4	32.3	47.8	80.4	147.9
2001.....	1.9	2.8	6.2	9.1	13.9	17.5	30.3	51.0	79.3	135.1
2000.....	1.7	3.6	5.8	8.6	16.0	25.7	35.1	49.6	73.9	129.6
1999.....	2.2	2.9	3.8	8.2	14.6	20.2	34.7	49.8	75.2	120.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
All races 3/, both sexes										
2006.....	0.2	0.3	0.5	0.7	1.0	1.4	2.4	4.3	8.6	22.6
2005.....	0.2	0.3	0.6	0.6	1.0	1.8	2.9	5.0	9.5	25.3
2004.....	0.3	0.5	0.5	0.7	1.3	1.6	2.6	4.5	9.4	24.8
2003.....	0.3	0.4	0.6	0.9	1.1	1.6	2.9	5.2	9.5	25.6
2002.....	0.2	0.4	0.5	0.9	1.2	1.6	2.8	4.9	9.5	23.5
2001.....	0.3	0.3	0.4	0.8	1.2	1.8	2.8	5.0	9.3	23.3
2000.....	0.3	0.3	0.4	0.8	1.0	1.6	3.0	5.1	9.8	24.3
1999.....	0.3	0.4	0.4	0.9	1.4	2.1	3.0	6.6	12.3	29.1
All races 3/, male										
2006.....	0.3	0.3	0.6	0.9	1.3	1.6	2.6	4.7	10.0	23.8
2005.....	0.3	0.5	0.7	0.8	1.2	2.1	2.9	5.3	9.8	27.4
2004.....	0.3	0.5	0.7	0.7	1.7	1.7	2.8	5.1	10.4	24.3
2003.....	0.3	0.6	0.7	1.1	1.3	1.7	3.0	5.6	10.1	26.7
2002.....	0.2	0.5	0.6	1.1	1.4	1.5	3.4	5.1	9.9	23.1
2001.....	0.4	0.4	0.6	0.9	1.2	2.1	2.9	5.2	9.4	24.2
2000.....	0.3	0.4	0.5	0.9	1.0	1.7	3.4	5.0	10.8	27.8
1999.....	0.4	0.5	0.6	1.0	1.6	2.3	3.2	7.3	13.3	31.3
All races 3/, female										
2006.....	*	0.3	0.4	0.4	0.8	1.3	2.3	4.0	7.8	22.1
2005.....	0.2	*	0.4	0.5	0.8	1.6	2.8	4.8	9.3	24.3
2004.....	*	0.4	0.4	0.7	0.9	1.5	2.3	4.0	8.8	25.0
2003.....	0.2	0.2	0.5	0.6	0.9	1.5	2.8	4.9	9.2	25.2
2002.....	0.2	0.2	0.4	0.7	1.0	1.7	2.4	4.7	9.2	23.7
2001.....	*	*	0.3	0.7	1.3	1.5	2.8	4.9	9.3	22.9
2000.....	0.2	0.2	0.4	0.7	1.0	1.5	2.7	5.1	9.3	22.9
1999.....	0.3	0.3	0.3	0.7	1.2	2.0	2.9	6.1	11.7	28.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
White, both sexes										
2006 4/.....	*	0.1	0.2	0.4	0.5	0.9	1.5	3.1	7.7	20.9
2005 4/.....	*	0.1	0.2	0.3	0.5	1.1	2.0	3.9	7.9	23.9
2004 4/.....	0.1	0.1	0.2	0.3	0.7	0.9	1.8	3.5	8.1	23.0
2003 4/.....	*	0.1	0.2	0.3	0.6	0.9	2.1	4.2	8.3	23.4
2002.....	*	*	0.1	0.3	0.6	0.9	1.9	3.8	8.4	21.0
2001.....	*	*	0.2	0.4	0.7	1.1	2.0	3.9	8.3	21.0
2000.....	0.2	0.1	0.2	0.4	0.5	0.9	2.2	4.0	8.1	22.0
1999.....	*	0.1	0.2	0.4	0.7	1.2	1.9	5.2	10.4	26.1
White, male										
2006 4/.....	*	*	0.3	0.5	0.6	0.8	1.6	3.3	9.0	22.5
2005 4/.....	*	*	0.3	0.4	0.5	1.1	2.1	4.2	8.2	25.5
2004 4/.....	*	*	*	0.3	1.0	1.0	2.1	3.9	9.2	22.9
2003 4/.....	*	*	*	0.5	0.8	1.1	2.2	4.9	9.3	24.3
2002.....	*	*	*	0.5	0.7	0.8	2.4	4.2	8.8	20.6
2001.....	*	*	0.3	0.4	0.7	1.3	2.0	4.1	8.7	22.7
2000.....	*	*	*	0.4	0.5	1.0	2.5	4.4	9.2	24.4
1999.....	*	*	*	0.5	0.8	1.4	2.0	5.8	12.1	28.8
White, female										
2006 4/.....	*	*	*	*	0.4	0.9	1.5	2.9	6.8	20.1
2005 4/.....	*	*	*	*	0.4	1.0	1.9	3.7	7.8	23.2
2004 4/.....	*	*	*	0.3	0.4	0.8	1.5	3.2	7.4	23.0
2003 4/.....	*	*	*	*	0.5	0.7	1.9	3.7	7.8	22.9
2002.....	*	*	*	*	0.5	0.9	1.5	3.6	8.1	21.1
2001.....	*	*	*	0.3	0.8	0.8	2.0	3.7	8.0	20.3
2000.....	*	*	*	0.4	0.5	0.8	2.0	3.7	7.4	21.1
1999.....	*	*	*	0.4	0.6	1.1	1.8	4.7	9.5	25.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hypertensive heart and renal disease (I13)										
Black, both sexes										
2006 4/.....	1.0	1.7	2.7	3.4	5.7	6.5	11.0	16.8	21.1	41.8
2005 4/.....	1.2	1.7	3.4	3.7	5.7	8.7	10.7	16.5	28.4	42.8
2004 4/.....	1.3	2.8	2.9	3.9	6.4	7.5	9.7	15.2	26.0	48.2
2003 4/.....	1.5	2.3	3.8	5.3	5.5	7.4	11.0	16.5	24.1	53.1
2002.....	1.1	2.2	3.3	6.0	6.2	8.0	12.3	17.4	23.0	56.1
2001.....	1.4	1.9	2.3	4.4	5.7	8.4	11.8	18.4	22.8	50.1
2000.....	1.2	1.9	2.5	4.1	5.3	8.3	11.9	17.6	31.1	50.4
1999.....	2.1	2.5	2.3	5.0	7.3	10.1	15.4	21.8	35.3	65.4
Black, male										
2006 4/.....	1.5	1.9	3.5	5.0	7.6	8.2	13.5	22.8	23.3	39.3
2005 4/.....	*	2.8	4.0	4.7	8.0	11.6	10.4	17.0	33.0	50.2
2004 4/.....	2.0	3.3	3.8	4.4	8.6	8.2	9.4	20.5	27.3	41.7
2003 4/.....	2.1	3.4	4.6	7.6	6.4	6.6	11.8	15.5	24.4	50.0
2002.....	*	3.0	4.7	7.6	7.3	7.2	14.0	17.0	22.9	61.1
2001.....	2.1	3.0	3.5	5.6	6.3	10.7	13.5	18.1	19.3	44.3
2000.....	1.5	2.7	2.9	5.1	5.6	9.4	13.7	13.7	33.7	63.4
1999.....	2.6	3.5	3.4	6.3	9.3	10.6	17.3	23.7	29.6	59.8
Black, female										
2006 4/.....	*	1.5	1.9	2.1	4.3	5.2	9.3	13.1	19.9	42.8
2005 4/.....	*	*	2.9	2.9	3.8	6.5	10.8	16.3	26.0	39.8
2004 4/.....	*	2.4	2.2	3.5	4.7	7.0	9.9	12.1	25.3	50.8
2003 4/.....	*	*	3.1	3.5	4.8	8.0	10.4	17.1	24.0	54.3
2002.....	*	1.5	2.1	4.6	5.3	8.5	11.2	17.6	23.1	54.2
2001.....	*	*	*	3.3	5.3	6.8	10.8	18.6	24.5	52.2
2000.....	*	*	2.1	3.2	5.1	7.5	10.7	19.9	29.7	45.5
1999.....	1.6	1.6	*	4.0	5.7	9.6	14.2	20.6	38.0	67.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
All races 3/, both sexes										
2006.....	21.5	42.2	75.0	118.1	188.5	279.1	441.7	726.4	1,263.9	2,904.7
2005.....	21.6	42.9	78.3	120.9	198.6	295.4	472.3	778.6	1,359.8	3,158.1
2004.....	22.4	44.0	77.9	123.9	203.1	310.5	495.9	813.5	1,418.0	3,287.4
2003.....	23.5	45.9	80.7	132.4	218.7	335.0	538.1	886.4	1,538.2	3,610.2
2002.....	23.8	46.4	83.5	140.1	226.2	358.4	569.8	940.2	1,611.2	3,775.0
2001.....	23.4	45.6	84.8	142.8	236.6	372.4	591.8	984.3	1,685.2	3,958.2
2000.....	23.6	48.0	86.7	153.9	250.0	390.8	628.0	1,032.5	1,761.7	4,182.5
1999.....	24.2	48.5	90.1	157.2	260.0	420.1	661.9	1,081.1	1,841.7	4,318.1
All races 3/, male										
2006.....	32.7	64.4	116.9	180.0	277.1	400.1	613.0	988.2	1,672.7	3,303.0
2005.....	33.0	66.2	121.3	183.8	291.3	415.2	653.7	1,054.3	1,780.2	3,576.8
2004.....	34.0	68.2	121.1	187.5	297.8	440.2	682.0	1,106.0	1,855.0	3,680.3
2003.....	35.5	71.4	126.9	200.3	319.7	476.1	744.1	1,200.1	2,013.5	4,005.2
2002.....	36.5	72.3	130.3	211.4	331.4	508.5	791.5	1,274.8	2,094.2	4,205.3
2001.....	35.8	71.6	131.1	215.5	344.4	525.7	820.3	1,337.5	2,180.0	4,395.8
2000.....	37.0	76.9	134.2	232.8	362.9	550.5	879.6	1,394.4	2,305.0	4,701.7
1999.....	37.9	77.0	140.3	237.7	376.1	595.2	924.6	1,462.5	2,400.4	4,837.5
All races 3/, female										
2006.....	10.3	20.6	34.8	59.7	107.5	173.3	302.4	534.8	1,008.4	2,718.3
2005.....	10.4	20.2	37.0	61.5	114.1	190.9	325.4	578.5	1,098.7	2,965.8
2004.....	10.8	20.4	36.6	64.0	116.8	197.7	346.0	603.2	1,148.8	3,110.6
2003.....	11.6	21.1	36.5	68.4	126.9	212.6	373.0	663.1	1,247.5	3,435.5
2002.....	11.2	21.4	38.7	73.1	130.7	228.8	392.6	704.0	1,317.5	3,588.3
2001.....	11.1	20.3	40.3	74.8	138.8	240.5	410.2	737.0	1,388.7	3,775.1
2000.....	10.3	20.0	41.3	80.0	147.7	253.9	429.9	780.5	1,441.2	3,971.0
1999.....	10.7	20.8	42.1	82.1	155.0	270.2	456.7	815.9	1,514.7	4,108.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
White, both sexes										
2006 4/.....	21.0	40.1	71.3	112.1	180.9	268.6	432.1	720.3	1,267.5	2,964.3
2005 4/.....	21.0	41.1	73.8	114.2	189.8	286.3	463.7	771.1	1,364.5	3,227.0
2004 4/.....	21.6	42.0	73.4	117.5	195.2	298.7	485.8	807.0	1,423.8	3,343.9
2003 4/.....	22.4	43.5	76.1	126.4	208.2	322.6	526.3	877.2	1,542.2	3,671.7
2002.....	22.9	44.2	79.1	133.3	216.1	346.6	559.2	928.9	1,616.4	3,836.4
2001.....	22.1	43.1	80.3	135.4	225.8	357.9	580.5	972.9	1,689.9	4,022.7
2000.....	22.2	45.3	82.4	146.2	238.5	378.4	615.0	1,019.7	1,768.5	4,257.7
1999.....	22.6	46.0	85.4	149.1	249.5	408.5	647.3	1,070.6	1,849.5	4,389.2
White, male										
2006 4/.....	32.6	62.2	112.7	172.6	267.5	388.4	603.3	983.8	1,689.7	3,396.0
2005 4/.....	32.7	64.1	116.2	175.7	281.5	406.5	646.8	1,053.1	1,798.9	3,687.1
2004 4/.....	33.7	66.3	116.1	180.0	289.4	428.5	673.9	1,105.8	1,880.8	3,768.8
2003 4/.....	34.4	68.8	121.6	194.2	308.7	464.4	731.5	1,195.8	2,028.8	4,106.1
2002.....	35.9	70.3	125.6	203.9	320.5	498.1	784.6	1,273.1	2,115.1	4,299.7
2001.....	34.6	69.5	126.4	207.2	333.9	512.8	810.3	1,335.1	2,201.7	4,494.0
2000.....	35.9	74.5	130.0	223.7	352.1	541.2	870.1	1,390.0	2,328.3	4,813.9
1999.....	36.2	74.7	135.0	228.2	364.4	586.0	913.4	1,463.9	2,429.7	4,933.9
White, female										
2006 4/.....	9.2	18.1	30.6	53.9	100.1	161.9	289.9	523.6	1,000.8	2,761.6
2005 4/.....	9.2	18.1	32.1	54.9	104.5	179.4	312.3	562.4	1,092.2	3,015.2
2004 4/.....	9.4	17.9	31.5	57.3	107.8	183.6	331.3	588.5	1,140.0	3,152.3
2003 4/.....	10.4	18.4	31.5	61.3	115.2	197.3	358.4	646.9	1,242.2	3,479.3
2002.....	9.8	18.2	33.6	65.7	119.6	213.2	375.5	682.5	1,310.6	3,635.2
2001.....	9.5	17.0	35.2	66.8	126.1	222.0	394.4	716.3	1,380.6	3,825.5
2000.....	8.5	16.4	36.0	72.5	133.8	236.0	410.4	759.2	1,435.5	4,031.8
1999.....	8.9	17.6	37.0	73.8	143.6	253.7	435.9	794.5	1,507.0	4,170.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ischemic heart diseases (I20-I25)										
Black, both sexes										
2006 4/.....	29.7	64.9	121.2	191.9	295.9	422.1	616.2	922.5	1,418.1	2,616.1
2005 4/.....	30.2	65.0	129.0	202.1	320.9	431.0	638.8	1,004.8	1,517.3	2,804.4
2004 4/.....	33.2	67.4	130.6	207.3	320.6	471.5	691.7	1,030.8	1,554.7	3,014.3
2003 4/.....	35.2	71.5	133.3	213.8	352.2	503.0	756.8	1,139.8	1,696.1	3,310.1
2002.....	35.1	72.8	136.5	230.9	362.0	522.5	779.6	1,196.0	1,768.0	3,462.9
2001.....	36.5	72.5	138.8	236.1	376.4	562.5	810.8	1,249.3	1,837.2	3,615.1
2000.....	37.1	78.2	140.3	250.9	392.9	566.1	866.5	1,296.9	1,882.8	3,684.7
1999.....	39.7	76.9	149.3	258.5	395.4	596.2	906.3	1,321.5	1,926.9	3,805.8
Black, male										
2006 4/.....	41.0	91.7	178.2	281.0	431.2	594.0	850.5	1,242.2	1,785.3	2,656.7
2005 4/.....	41.2	95.4	187.9	294.0	454.6	584.9	857.4	1,296.8	1,887.0	2,785.0
2004 4/.....	44.9	96.6	189.8	297.2	454.8	641.3	922.9	1,332.6	1,860.9	3,100.5
2003 4/.....	50.2	103.2	195.6	302.7	490.5	677.1	1,027.8	1,501.3	2,138.8	3,285.3
2002.....	48.4	101.6	196.9	330.3	509.6	703.3	1,024.8	1,495.0	2,171.1	3,556.2
2001.....	50.8	101.9	199.9	333.7	515.8	743.8	1,093.0	1,582.5	2,244.8	3,775.0
2000.....	51.8	111.6	197.6	366.7	530.3	727.5	1,139.7	1,630.6	2,316.5	3,885.2
1999.....	56.8	110.4	217.6	372.5	553.5	788.8	1,188.9	1,630.3	2,312.4	4,080.1
Black, female										
2006 4/.....	19.7	41.6	73.0	118.5	190.5	295.1	455.2	731.2	1,224.0	2,599.5
2005 4/.....	20.5	38.6	79.1	126.3	216.6	316.9	489.1	830.7	1,322.0	2,812.2
2004 4/.....	22.9	42.0	80.5	133.4	215.9	345.4	534.9	852.0	1,393.3	2,980.1
2003 4/.....	21.9	43.9	80.5	140.7	244.3	373.7	574.0	924.9	1,464.2	3,319.7
2002.....	23.5	47.9	85.4	149.3	246.6	388.3	615.0	1,017.3	1,558.1	3,427.2
2001.....	23.9	47.1	86.8	156.0	266.7	428.6	622.8	1,050.3	1,629.6	3,555.7
2000.....	24.1	49.2	91.4	156.0	284.5	447.0	685.1	1,095.7	1,664.3	3,610.0
1999.....	24.7	48.0	91.4	165.5	271.4	453.5	718.8	1,135.4	1,735.9	3,703.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
All races 3/, both sexes										
2006.....	8.5	16.5	29.2	46.7	73.9	107.0	160.4	252.0	406.4	825.8
2005.....	8.7	17.3	30.9	47.8	78.9	115.0	176.6	272.1	448.1	923.1
2004.....	8.9	17.7	31.6	50.6	82.7	124.1	187.7	291.1	476.0	992.9
2003.....	9.6	18.8	33.2	55.5	90.8	134.5	207.4	325.6	532.5	1,118.8
2002.....	9.9	19.0	34.9	58.4	96.2	148.5	227.5	352.5	572.7	1,191.3
2001.....	9.9	19.7	36.5	61.3	100.1	154.1	235.6	374.0	611.5	1,265.4
2000.....	9.9	21.1	37.8	67.0	110.9	166.8	254.7	405.3	650.1	1,338.5
1999.....	10.4	21.5	39.0	69.7	114.9	181.7	271.6	422.9	689.4	1,379.3
All races 3/, male										
2006.....	13.0	24.7	44.8	70.0	107.3	151.8	219.9	331.3	527.1	936.3
2005.....	13.1	26.5	47.2	72.2	114.0	159.3	240.5	361.8	567.1	1,045.0
2004.....	13.2	27.4	48.7	76.3	120.6	173.0	254.1	386.8	608.9	1,120.5
2003.....	14.4	29.2	52.1	83.4	131.6	188.5	281.4	430.0	682.4	1,263.5
2002.....	14.8	29.2	53.7	86.6	139.0	207.6	312.0	466.1	726.6	1,366.4
2001.....	15.0	30.8	55.6	91.3	143.5	214.2	322.0	495.6	776.1	1,461.3
2000.....	15.5	33.6	57.5	99.7	158.6	231.3	349.6	535.2	835.8	1,568.3
1999.....	16.0	33.7	60.9	105.3	164.9	253.5	373.8	564.2	885.1	1,622.1
All races 3/, female										
2006.....	4.1	8.4	14.3	24.7	43.4	67.8	112.0	194.0	330.9	774.1
2005.....	4.4	8.4	15.2	24.7	47.0	76.4	124.8	206.9	374.1	867.1
2004.....	4.5	8.3	15.3	26.4	48.2	81.5	134.3	222.3	394.1	935.5
2003.....	5.0	8.7	15.1	29.3	53.8	87.7	148.0	251.3	440.9	1,054.8
2002.....	5.1	9.1	17.0	32.0	57.4	97.6	159.8	272.2	479.1	1,115.3
2001.....	4.9	9.0	18.2	33.2	60.7	102.4	167.0	288.8	512.9	1,183.5
2000.....	4.4	8.9	18.9	36.5	67.7	111.5	179.9	314.9	540.6	1,244.8
1999.....	4.9	9.7	18.0	36.5	69.6	120.2	191.7	324.7	574.9	1,281.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
White, both sexes										
2006 4/.....	8.5	16.1	28.6	45.6	72.9	105.2	158.9	250.6	408.0	836.0
2005 4/.....	8.6	17.0	30.3	46.3	78.0	113.8	175.8	270.1	449.4	936.8
2004 4/.....	8.7	17.4	30.6	49.4	81.8	121.7	185.3	291.1	478.6	1,004.1
2003 4/.....	9.3	18.2	32.1	54.4	89.1	132.3	205.3	324.2	532.9	1,132.6
2002.....	9.7	18.4	34.2	57.3	94.3	146.4	225.0	349.5	572.7	1,201.2
2001.....	9.5	19.3	35.4	59.8	98.4	150.8	233.3	371.9	612.7	1,277.8
2000.....	9.5	20.6	36.7	65.6	108.9	164.1	251.8	401.7	652.6	1,353.5
1999.....	10.0	21.1	38.3	68.2	113.4	180.6	268.3	421.1	692.5	1,393.2
White, male										
2006 4/.....	13.1	24.6	44.6	69.2	106.3	151.2	220.0	331.9	532.5	955.4
2005 4/.....	13.1	26.3	47.2	70.7	114.2	160.0	241.8	363.3	572.8	1,070.1
2004 4/.....	13.4	27.2	48.0	75.4	121.2	171.7	252.9	391.6	618.6	1,143.7
2003 4/.....	14.1	28.7	51.2	83.0	130.8	188.3	280.0	430.3	685.7	1,289.3
2002.....	14.7	29.0	53.3	86.2	138.7	207.7	313.1	467.1	730.8	1,386.1
2001.....	14.9	30.9	55.1	90.6	143.7	213.3	321.9	497.6	784.0	1,481.1
2000.....	15.3	33.6	57.2	98.8	158.3	231.1	349.3	536.3	843.5	1,592.9
1999.....	15.8	33.8	60.8	104.0	165.1	254.3	374.0	568.8	896.6	1,644.0
White, female										
2006 4/.....	3.9	7.6	13.0	22.9	41.7	64.3	108.1	189.9	329.4	780.0
2005 4/.....	3.9	7.7	13.7	22.9	44.3	72.6	121.3	201.2	372.1	875.4
2004 4/.....	3.9	7.5	13.5	24.3	45.2	77.4	129.7	217.6	391.6	941.1
2003 4/.....	4.5	7.7	13.4	26.8	50.5	82.8	144.1	247.5	438.7	1,063.2
2002.....	4.6	8.0	15.4	29.7	53.3	92.5	153.2	265.3	475.8	1,120.8
2001.....	4.2	7.7	16.1	30.3	56.7	96.0	161.5	282.8	509.1	1,192.7
2000.....	3.7	7.7	16.6	34.1	63.5	105.5	173.6	307.0	539.1	1,256.2
1999.....	4.3	8.4	16.2	34.2	65.8	116.3	184.3	317.3	572.1	1,292.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute myocardial infarction (I21-I22)										
Black, both sexes										
2006 4/.....	11.1	23.1	41.6	67.0	100.9	144.9	207.1	311.5	448.2	804.0
2005 4/.....	11.8	24.2	43.3	69.9	107.8	150.2	217.0	345.4	499.1	892.1
2004 4/.....	12.4	25.0	47.9	75.3	112.7	169.2	249.7	351.1	520.9	979.9
2003 4/.....	14.4	27.5	49.5	79.4	124.3	180.3	270.2	401.4	597.0	1,094.9
2002.....	13.9	27.7	50.4	84.2	136.3	193.6	295.9	434.1	643.9	1,201.2
2001.....	14.3	27.8	53.2	89.2	135.9	211.1	304.3	452.3	671.2	1,239.9
2000.....	15.1	29.6	55.5	94.7	151.4	219.0	328.8	490.6	698.8	1,275.7
1999.....	15.4	30.0	54.9	96.9	150.2	224.0	347.5	495.1	729.6	1,318.8
Black, male										
2006 4/.....	15.9	31.5	59.0	95.0	144.9	195.4	271.9	391.1	552.4	814.1
2005 4/.....	15.8	34.4	59.5	100.2	145.2	190.8	278.3	425.6	598.3	885.9
2004 4/.....	15.4	36.3	67.6	104.8	150.1	223.1	322.7	419.5	591.1	1,009.3
2003 4/.....	20.0	38.6	70.9	108.1	167.9	230.2	353.4	516.2	736.1	1,113.5
2002.....	19.0	37.1	70.4	114.5	177.6	247.8	370.2	531.4	776.4	1,280.1
2001.....	18.5	36.4	71.2	120.1	174.5	264.8	386.4	548.7	792.2	1,382.0
2000.....	20.6	41.0	72.7	131.4	196.7	270.9	419.2	592.5	856.5	1,418.5
1999.....	21.1	41.1	76.9	138.8	198.8	290.8	435.5	591.7	858.0	1,498.7
Black, female										
2006 4/.....	7.0	15.8	26.9	43.9	66.7	107.7	162.5	263.8	393.2	799.9
2005 4/.....	8.4	15.3	29.6	44.9	78.6	120.0	175.1	297.6	446.6	894.6
2004 4/.....	9.7	15.2	31.2	51.0	83.6	129.2	200.1	310.5	483.8	968.3
2003 4/.....	9.5	18.0	31.5	55.7	90.3	143.3	214.0	333.2	524.2	1,087.7
2002.....	9.3	19.5	33.4	59.4	104.1	153.4	246.0	376.0	574.9	1,171.0
2001.....	10.5	20.3	37.8	63.8	105.6	171.4	249.5	394.7	609.5	1,187.2
2000.....	10.1	19.8	40.8	64.6	115.7	180.7	268.8	429.2	619.4	1,222.5
1999.....	10.4	20.5	36.2	62.6	112.1	174.5	289.2	436.8	665.9	1,251.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
All races 3/, both sexes										
2006.....	0.3	0.6	1.1	1.5	2.5	2.9	4.4	6.4	10.0	21.5
2005.....	0.3	0.6	1.1	1.4	2.1	2.4	4.2	5.4	9.5	20.8
2004.....	0.3	0.4	0.7	1.1	1.4	1.8	2.6	4.2	6.5	14.8
2003.....	0.3	0.5	0.9	1.3	1.7	2.7	3.6	5.2	8.3	20.0
2002.....	0.3	0.6	1.2	1.4	2.2	3.0	3.9	5.7	9.0	21.1
2001.....	0.3	0.6	1.0	1.4	2.4	2.9	4.1	5.3	9.2	21.5
2000.....	0.3	0.5	1.1	1.6	2.5	3.1	4.0	5.9	9.4	21.9
1999.....	0.3	0.5	0.9	1.6	2.2	3.1	4.3	5.5	9.2	21.9
All races 3/, male										
2006.....	0.4	0.8	1.6	1.9	3.6	3.6	5.9	8.0	12.1	21.8
2005.....	0.4	0.8	1.6	2.1	3.2	3.4	5.3	7.1	10.6	20.2
2004.....	0.4	0.5	1.0	1.6	2.0	2.7	3.4	5.0	8.3	16.3
2003.....	0.5	0.7	1.3	1.8	2.4	3.8	4.9	7.1	9.4	19.5
2002.....	0.4	0.8	1.8	2.1	3.0	4.3	5.1	7.0	11.3	21.4
2001.....	0.4	0.9	1.6	2.0	3.5	4.1	5.5	6.7	11.3	20.7
2000.....	0.4	0.8	1.6	2.5	3.5	4.4	5.2	7.6	11.7	20.7
1999.....	0.4	0.7	1.3	2.5	3.1	4.3	5.6	7.6	10.8	22.3
All races 3/, female										
2006.....	0.2	0.3	0.6	1.0	1.5	2.2	3.3	5.3	8.8	21.4
2005.....	*	0.4	0.6	0.7	1.2	1.6	3.3	4.2	8.8	21.1
2004.....	*	0.2	0.4	0.6	0.8	1.0	1.9	3.6	5.4	14.1
2003.....	*	0.3	0.5	0.8	1.0	1.8	2.6	3.9	7.6	20.2
2002.....	*	0.3	0.7	0.7	1.4	1.9	3.0	4.7	7.6	21.0
2001.....	*	0.3	0.5	0.9	1.3	2.0	3.1	4.4	8.0	21.8
2000.....	*	*	0.5	0.8	1.5	1.9	2.9	4.8	8.1	22.4
1999.....	*	0.2	0.6	0.7	1.5	2.0	3.2	4.1	8.3	21.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
White, both sexes										
2006 4/.....	0.2	0.5	1.0	1.4	2.3	2.7	4.2	6.2	9.8	21.6
2005 4/.....	0.2	0.5	0.9	1.2	1.9	2.3	3.9	5.1	9.2	20.5
2004 4/.....	0.3	0.3	0.6	0.9	1.3	1.6	2.4	3.9	6.3	15.0
2003 4/.....	0.3	0.4	0.8	1.1	1.6	2.6	3.4	5.0	8.1	20.2
2002.....	0.2	0.5	1.1	1.3	2.0	2.9	3.8	5.6	8.9	21.4
2001.....	0.2	0.5	0.9	1.3	2.1	2.7	4.0	5.2	9.2	21.5
2000.....	0.2	0.4	1.0	1.4	2.2	2.9	3.8	5.6	9.3	22.1
1999.....	0.2	0.3	0.8	1.4	2.1	3.0	4.0	5.4	8.9	22.4
White, male										
2006 4/.....	0.3	0.7	1.5	1.8	3.4	3.5	5.5	7.6	11.6	21.8
2005 4/.....	0.3	0.7	1.4	1.8	2.8	3.2	4.9	6.6	10.3	19.7
2004 4/.....	0.4	0.4	0.8	1.4	1.8	2.3	3.1	4.6	8.3	16.2
2003 4/.....	0.4	0.6	1.2	1.5	2.2	3.6	4.8	6.7	8.9	19.7
2002.....	0.4	0.7	1.6	2.0	2.8	4.1	4.8	7.0	11.1	21.9
2001.....	0.3	0.8	1.3	1.8	3.1	3.5	5.4	6.5	11.6	20.7
2000.....	0.4	0.7	1.5	2.2	3.2	4.0	4.9	7.2	11.6	21.0
1999.....	0.3	0.5	1.2	2.2	2.8	4.1	5.2	7.6	10.3	22.9
White, female										
2006 4/.....	*	0.2	0.5	0.9	1.4	2.0	3.1	5.2	8.7	21.6
2005 4/.....	*	0.2	0.4	0.6	1.0	1.6	3.1	4.0	8.5	20.8
2004 4/.....	*	*	0.3	0.5	0.8	1.0	1.9	3.4	5.1	14.4
2003 4/.....	*	0.2	0.4	0.7	0.9	1.6	2.4	3.8	7.7	20.5
2002.....	*	0.3	0.6	0.7	1.2	1.8	2.9	4.6	7.5	21.1
2001.....	*	0.2	0.4	0.8	1.2	1.9	2.9	4.3	7.7	21.7
2000.....	*	*	0.5	0.6	1.3	1.9	2.8	4.5	8.0	22.6
1999.....	*	*	0.5	0.7	1.5	2.0	3.0	3.8	8.1	22.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
Black, both sexes										
2006 4/.....	*	1.1	2.3	2.6	4.5	4.7	7.6	9.0	14.2	20.8
2005 4/.....	*	1.3	2.8	3.2	4.5	4.1	7.3	9.7	14.8	27.3
2004 4/.....	*	1.0	1.9	2.9	3.0	3.9	4.3	8.1	9.9	15.2
2003 4/.....	0.7	1.1	1.5	2.7	3.3	4.9	6.2	8.7	11.7	19.8
2002.....	*	1.1	2.4	2.1	4.1	5.0	7.1	7.7	11.8	21.1
2001.....	0.7	1.1	2.1	2.8	5.2	5.3	6.2	6.9	10.8	24.0
2000.....	*	1.0	2.1	3.8	4.7	5.2	6.1	10.0	12.1	21.9
1999.....	*	1.5	2.2	3.3	3.9	4.3	7.3	8.7	13.9	17.2
Black, male										
2006 4/.....	*	*	3.2	3.9	6.9	5.6	11.1	12.5	19.2	23.2
2005 4/.....	*	*	3.6	5.0	6.9	6.4	9.8	14.8	17.6	27.9
2004 4/.....	*	*	2.7	4.5	4.7	7.0	7.1	11.8	*	22.8
2003 4/.....	*	1.7	2.2	4.6	5.0	7.4	7.9	14.2	19.2	24.0
2002.....	*	1.9	3.6	3.1	5.7	8.0	10.7	9.2	16.0	21.4
2001.....	*	*	3.6	3.4	8.1	9.1	7.7	11.2	*	23.8
2000.....	*	*	3.6	5.8	6.6	8.3	9.0	13.2	*	*
1999.....	*	2.5	2.8	5.8	6.6	6.8	10.2	10.0	17.4	*
Black, female										
2006 4/.....	*	*	*	*	*	4.0	5.3	7.0	11.6	19.7
2005 4/.....	*	*	2.2	*	*	*	5.6	6.7	13.4	27.0
2004 4/.....	*	*	*	*	*	*	*	5.9	9.1	12.2
2003 4/.....	*	*	*	*	*	*	5.1	5.4	7.7	18.2
2002.....	*	*	*	*	*	*	4.7	6.9	9.5	20.9
2001.....	*	*	*	*	*	*	5.2	*	11.9	24.0
2000.....	*	*	*	*	3.3	*	*	8.0	10.6	22.3
1999.....	*	*	*	*	*	*	5.4	8.0	12.1	16.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes										
2006.....	12.7	25.2	44.7	70.0	112.1	169.3	276.9	468.0	847.5	2,057.4
2005.....	12.7	25.1	46.3	71.7	117.5	178.0	291.6	501.1	902.3	2,214.2
2004.....	13.2	25.9	45.6	72.2	119.0	184.6	305.6	518.2	935.6	2,279.7
2003.....	13.5	26.6	46.6	75.6	126.1	197.8	327.1	555.6	997.3	2,471.4
2002.....	13.6	26.8	47.3	80.2	127.8	206.8	338.4	582.1	1,029.5	2,562.5
2001.....	13.2	25.2	47.3	80.1	134.1	215.4	352.0	605.0	1,064.5	2,671.3
2000.....	13.4	26.5	47.9	85.2	136.6	221.0	369.4	621.2	1,102.2	2,822.1
1999.....	13.5	26.5	50.2	85.9	142.9	235.3	386.1	652.6	1,143.1	2,917.0
All races 3/, male										
2006.....	19.4	38.8	70.5	108.1	166.2	244.7	387.2	649.0	1,133.5	2,344.9
2005.....	19.5	39.0	72.5	109.5	174.1	252.6	407.9	685.4	1,202.5	2,511.6
2004.....	20.4	40.3	71.4	109.5	175.2	264.4	424.5	714.2	1,237.9	2,543.6
2003.....	20.7	41.5	73.5	115.1	185.6	283.8	457.8	763.1	1,321.6	2,722.2
2002.....	21.4	42.2	74.9	122.7	189.4	296.6	474.3	801.6	1,356.2	2,817.4
2001.....	20.4	40.0	74.0	122.2	197.4	307.4	492.9	835.3	1,392.6	2,913.8
2000.....	21.1	42.5	75.1	130.6	200.8	314.8	524.7	851.5	1,457.6	3,112.7
1999.....	21.5	42.6	78.1	129.9	208.1	337.3	545.2	890.7	1,504.6	3,193.1
All races 3/, female										
2006.....	6.0	11.8	19.9	34.1	62.7	103.3	187.1	335.6	668.6	1,922.9
2005.....	5.9	11.5	21.2	36.1	65.9	112.9	197.3	367.3	715.8	2,077.6
2004.....	6.1	11.9	20.9	37.0	67.8	115.1	209.8	377.3	749.3	2,161.0
2003.....	6.5	12.1	20.9	38.4	72.0	123.2	222.4	408.0	799.1	2,360.5
2002.....	6.0	11.9	21.0	40.3	72.0	129.4	229.7	427.1	830.8	2,451.9
2001.....	6.1	11.0	21.7	40.6	76.8	136.2	240.1	443.8	867.9	2,569.8
2000.....	5.8	10.9	21.8	42.7	78.5	140.5	247.0	460.8	892.5	2,703.8
1999.....	5.6	10.9	23.4	44.9	83.9	148.1	261.7	487.1	931.6	2,805.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
White, both sexes										
2006 4/.....	12.2	23.5	41.7	65.2	105.7	160.6	269.0	463.4	849.7	2,106.6
2005 4/.....	12.2	23.6	42.6	66.7	109.9	170.2	283.9	495.8	905.8	2,269.8
2004 4/.....	12.6	24.4	42.2	67.2	112.1	175.4	298.1	512.0	939.0	2,324.8
2003 4/.....	12.9	24.9	43.2	71.0	117.6	187.7	317.6	548.0	1,001.1	2,518.8
2002.....	13.0	25.2	43.9	74.7	119.8	197.3	330.5	573.8	1,034.7	2,613.9
2001.....	12.3	23.3	44.0	74.3	125.3	204.5	343.2	595.8	1,068.1	2,723.5
2000.....	12.5	24.3	44.8	79.2	127.3	211.4	359.4	612.4	1,106.6	2,882.1
1999.....	12.3	24.6	46.3	79.4	134.0	225.0	375.0	644.2	1,148.1	2,973.6
White, male										
2006 4/.....	19.2	36.9	66.7	101.7	157.9	233.6	377.8	644.3	1,145.6	2,418.9
2005 4/.....	19.2	37.1	67.6	103.3	164.5	243.3	400.0	683.2	1,215.8	2,597.3
2004 4/.....	19.9	38.7	67.2	103.3	166.4	254.5	417.9	709.7	1,254.0	2,608.9
2003 4/.....	19.9	39.5	69.2	109.6	175.8	272.6	446.7	758.8	1,334.2	2,797.1
2002.....	20.8	40.6	70.7	115.8	179.1	286.2	466.8	799.0	1,373.2	2,891.6
2001.....	19.4	37.7	69.9	114.8	187.1	296.0	483.0	831.0	1,406.1	2,992.2
2000.....	20.3	40.1	71.3	122.7	190.6	306.1	515.9	846.5	1,473.2	3,200.0
1999.....	20.1	40.4	72.9	122.0	196.5	327.6	534.1	887.5	1,522.9	3,267.1
White, female										
2006 4/.....	5.2	10.2	17.1	30.0	57.0	95.6	178.7	328.5	662.7	1,960.0
2005 4/.....	5.1	10.2	18.0	31.4	59.2	105.2	187.9	357.2	711.6	2,119.0
2004 4/.....	5.3	10.2	17.7	32.5	61.8	105.2	199.7	367.4	743.3	2,196.8
2003 4/.....	5.8	10.5	17.7	33.8	63.8	112.8	211.9	395.6	795.8	2,395.5
2002.....	5.1	10.0	17.6	35.3	65.1	118.9	219.3	412.6	827.2	2,493.2
2001.....	5.2	9.0	18.7	35.8	68.2	124.1	230.0	429.2	863.8	2,611.0
2000.....	4.7	8.6	18.8	37.8	69.0	128.7	233.9	447.7	888.4	2,753.0
1999.....	4.5	9.0	20.3	38.9	76.3	135.5	248.5	473.4	926.8	2,855.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other forms of chronic ischemic heart disease (I20,I25)										
Black, both sexes										
2006 4/.....	18.1	40.7	77.3	122.3	190.5	272.5	401.5	602.0	955.6	1,791.4
2005 4/.....	17.9	39.5	82.8	129.0	208.5	276.7	414.4	649.7	1,003.4	1,885.0
2004 4/.....	20.3	41.4	80.8	129.1	204.8	298.4	437.7	671.6	1,023.9	2,019.1
2003 4/.....	20.1	42.8	82.2	131.7	224.6	317.7	480.4	729.7	1,087.4	2,195.3
2002.....	20.7	44.0	83.7	144.6	221.6	323.9	476.6	754.2	1,112.3	2,240.6
2001.....	21.5	43.7	83.6	144.1	235.3	346.1	500.3	790.1	1,155.2	2,351.2
2000.....	21.5	47.6	82.7	152.4	236.7	341.9	531.5	796.3	1,171.8	2,387.1
1999.....	23.8	45.4	92.2	158.4	241.2	367.9	551.4	817.7	1,183.5	2,469.8
Black, male										
2006 4/.....	24.6	58.7	116.0	182.2	279.4	393.0	567.5	838.6	1,213.7	1,819.4
2005 4/.....	24.7	59.5	124.8	188.8	302.5	387.6	569.3	856.5	1,271.1	1,871.2
2004 4/.....	28.8	59.1	119.5	187.9	300.1	411.1	593.0	901.3	1,258.3	2,068.3
2003 4/.....	29.4	62.9	122.5	189.9	317.6	439.5	666.5	970.9	1,383.5	2,147.8
2002.....	29.0	62.5	122.8	212.7	326.3	447.5	643.9	954.4	1,378.6	2,254.6
2001.....	31.2	64.0	125.1	210.1	333.1	469.8	698.8	1,022.7	1,443.9	2,369.1
2000.....	30.3	69.2	121.3	229.5	326.9	448.3	711.4	1,024.9	1,444.9	2,446.0
1999.....	34.9	66.9	138.0	227.8	348.1	491.2	743.2	1,028.6	1,437.0	2,561.5
Black, female										
2006 4/.....	12.4	24.9	44.5	73.0	121.2	183.5	287.4	460.4	819.3	1,779.9
2005 4/.....	11.9	22.1	47.3	79.7	135.3	194.4	308.4	526.4	862.1	1,890.6
2004 4/.....	12.8	26.0	48.2	80.7	130.5	214.6	332.3	535.6	900.4	1,999.6
2003 4/.....	11.9	25.3	48.2	83.8	152.1	227.3	354.9	586.3	932.3	2,213.8
2002.....	13.3	28.0	50.5	88.6	139.7	232.2	364.3	634.5	973.6	2,235.2
2001.....	13.0	26.1	48.2	89.9	158.2	254.8	368.1	651.2	1,008.1	2,344.5
2000.....	13.7	28.8	49.7	89.2	165.6	263.4	412.0	658.5	1,034.3	2,365.2
1999.....	14.1	26.8	53.5	101.6	157.4	276.5	424.2	690.6	1,057.9	2,435.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
All races 3/, both sexes										
2006.....	4.6	9.4	17.9	26.1	38.5	47.4	64.4	89.1	147.1	345.5
2005.....	4.4	9.6	18.6	26.9	37.7	46.8	64.6	96.5	161.9	376.4
2004.....	4.8	9.5	18.2	26.6	39.1	49.8	69.0	100.8	166.6	400.4
2003.....	4.7	9.9	18.6	26.6	40.5	53.3	73.3	110.7	186.2	442.9
2002.....	4.8	9.8	17.5	27.9	39.7	53.9	76.5	116.6	187.6	474.5
2001.....	4.6	8.8	17.4	27.7	40.6	55.3	76.7	121.8	197.2	493.6
2000.....	4.4	9.4	17.5	28.5	40.7	55.7	81.9	125.0	207.8	534.6
1999.....	4.7	9.1	17.4	28.6	43.4	59.6	86.8	131.6	218.2	563.4
All races 3/, male										
2006.....	7.0	14.9	29.3	41.5	58.9	69.9	90.6	117.9	188.1	365.3
2005.....	7.0	15.3	30.0	42.5	57.7	68.3	91.5	127.0	204.6	390.7
2004.....	7.7	15.3	29.6	42.0	58.4	73.1	96.4	133.8	210.5	414.0
2003.....	7.4	16.0	30.5	41.8	60.6	77.1	102.8	146.6	233.1	462.1
2002.....	7.8	15.6	28.4	44.8	59.7	78.8	106.7	154.5	232.4	491.8
2001.....	7.2	14.6	28.3	43.4	61.8	79.0	107.6	160.2	247.4	502.0
2000.....	7.1	15.5	28.4	45.8	61.3	80.8	115.6	167.1	252.8	550.8
1999.....	7.6	14.9	28.0	45.0	64.6	87.3	120.8	172.7	264.3	578.6
All races 3/, female										
2006.....	2.2	4.1	7.1	11.5	20.0	27.7	43.1	67.9	121.4	336.1
2005.....	1.9	4.0	7.7	12.2	19.4	28.1	42.9	74.3	135.4	369.9
2004.....	2.0	3.9	7.4	12.0	21.5	29.5	47.0	77.0	139.6	394.3
2003.....	2.0	4.0	7.2	12.4	22.2	32.6	49.7	85.2	157.6	434.4
2002.....	1.8	4.1	7.1	12.1	21.4	32.4	52.5	89.7	160.4	467.1
2001.....	2.1	3.3	7.0	12.9	21.4	34.9	52.1	94.9	167.1	490.1
2000.....	1.8	3.5	7.2	12.4	22.1	34.1	55.4	95.7	181.2	528.0
1999.....	1.8	3.5	7.3	13.4	24.1	36.0	60.2	103.1	191.3	557.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
White, both sexes										
2006 4/.....	4.3	8.6	16.2	23.4	34.9	42.7	58.6	83.0	140.9	343.2
2005 4/.....	4.3	9.0	16.6	23.8	33.3	42.5	58.7	89.9	154.9	373.8
2004 4/.....	4.6	8.7	16.4	23.7	35.1	44.4	63.4	93.3	160.3	395.6
2003 4/.....	4.4	9.0	16.7	24.2	35.4	47.5	66.6	103.0	179.6	439.3
2002.....	4.4	9.1	15.6	24.7	35.3	47.3	69.7	107.8	180.5	470.6
2001.....	4.2	7.9	15.6	23.9	35.0	47.9	69.4	112.2	190.6	489.7
2000.....	4.0	8.3	15.7	24.8	35.2	49.5	73.7	116.0	198.5	529.5
1999.....	4.2	8.0	15.2	24.6	37.5	52.3	77.9	121.5	209.8	557.5
White, male										
2006 4/.....	6.7	13.8	26.8	37.6	53.1	62.8	82.2	109.7	180.2	362.7
2005 4/.....	7.0	14.6	27.2	38.2	51.2	62.2	82.8	118.8	194.8	392.6
2004 4/.....	7.5	14.4	27.3	37.8	52.4	66.1	89.3	123.9	203.3	408.8
2003 4/.....	6.9	14.8	27.7	38.3	53.4	69.2	92.5	136.9	224.4	458.7
2002.....	7.4	14.7	25.7	40.0	53.6	69.4	98.0	143.3	222.2	488.0
2001.....	6.6	13.3	25.7	38.3	54.0	69.3	97.6	148.7	239.9	498.6
2000.....	6.6	14.0	25.9	40.0	54.2	72.8	104.4	154.6	241.6	545.6
1999.....	7.0	13.2	24.8	39.4	56.2	77.6	109.0	159.8	253.9	571.2
White, female										
2006 4/.....	1.8	3.4	5.7	9.7	17.9	24.7	39.1	63.1	116.1	334.1
2005 4/.....	1.5	3.4	6.2	9.9	16.6	25.1	38.9	68.5	129.9	365.1
2004 4/.....	1.7	3.0	5.8	10.2	19.1	25.1	42.1	70.9	133.6	389.7
2003 4/.....	1.8	3.2	5.9	10.6	18.7	28.3	45.4	78.5	152.0	430.7
2002.....	1.5	3.5	5.6	10.2	18.4	27.8	46.6	82.5	154.9	463.0
2001.....	1.8	2.6	5.7	10.1	17.6	29.1	46.7	86.4	160.8	486.0
2000.....	1.4	2.7	5.8	10.2	17.6	29.0	49.0	88.8	172.9	522.9
1999.....	1.3	2.8	5.8	10.6	20.3	30.3	53.1	94.6	183.8	552.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerotic cardiovascular disease, so described (I25.0)										
Black, both sexes										
2006 4/.....	7.6	16.7	34.8	52.2	78.1	97.2	126.6	167.0	244.5	428.1
2005 4/.....	6.5	15.5	37.3	56.6	83.4	92.4	130.4	183.1	265.1	469.6
2004 4/.....	7.5	17.5	35.7	55.9	81.5	105.1	139.0	198.5	268.4	505.1
2003 4/.....	7.3	17.4	35.9	53.2	91.8	111.4	149.2	214.1	282.8	543.2
2002.....	8.1	17.0	35.9	60.5	85.5	119.5	157.5	231.9	300.4	575.5
2001.....	8.2	16.8	35.7	64.2	94.4	128.0	162.6	244.0	301.5	594.5
2000.....	7.4	19.4	36.4	65.4	94.0	119.6	176.8	239.7	340.0	646.6
1999.....	9.1	19.1	38.4	67.8	98.2	133.9	188.9	261.7	338.9	668.6
Black, male										
2006 4/.....	10.5	24.8	54.4	82.0	126.0	151.8	190.7	239.0	329.3	465.6
2005 4/.....	8.9	23.1	58.1	87.6	130.3	140.2	200.8	251.7	361.7	450.8
2004 4/.....	10.7	25.1	54.0	88.3	127.8	153.1	197.1	278.0	350.8	531.5
2003 4/.....	11.4	26.4	57.8	81.3	140.0	161.3	228.9	301.3	380.2	583.9
2002.....	11.8	25.5	55.0	97.1	130.5	180.0	220.3	321.0	417.3	599.3
2001.....	12.4	25.7	56.0	97.0	144.0	183.5	241.0	330.4	388.5	609.3
2000.....	11.0	30.3	56.8	106.4	137.1	168.6	257.1	338.3	425.7	701.2
1999.....	13.7	29.9	59.7	103.8	147.9	197.3	271.0	350.0	435.1	715.9
Black, female										
2006 4/.....	5.1	9.6	18.2	27.7	40.8	56.8	82.6	124.0	199.7	412.7
2005 4/.....	4.3	8.9	19.7	31.1	46.8	56.9	82.2	142.1	214.0	477.1
2004 4/.....	4.7	10.9	20.2	29.3	45.3	69.4	99.5	151.5	224.9	494.6
2003 4/.....	3.7	9.6	17.3	30.0	54.2	74.4	95.5	162.2	231.7	527.4
2002.....	4.8	9.7	19.7	30.4	50.3	74.6	115.4	178.7	239.5	566.4
2001.....	4.5	9.0	18.4	37.3	55.4	87.1	110.3	192.5	257.1	589.0
2000.....	4.2	10.0	19.0	31.8	60.0	83.5	123.4	180.3	296.8	626.3
1999.....	5.0	9.8	20.4	38.4	59.2	87.0	134.3	208.5	291.3	650.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
All races 3/, both sexes										
2006.....	8.1	15.7	26.8	43.9	73.6	121.9	212.5	379.0	700.4	1,712.0
2005.....	8.2	15.5	27.7	44.9	79.9	131.2	226.9	404.6	740.4	1,837.8
2004.....	8.4	16.4	27.4	45.7	79.9	134.8	236.6	417.4	769.0	1,879.3
2003.....	8.9	16.7	28.0	49.0	85.6	144.5	253.8	444.9	811.1	2,028.5
2002.....	8.8	17.1	29.9	52.3	88.2	152.9	261.8	465.6	841.9	2,088.0
2001.....	8.6	16.4	29.8	52.4	93.5	160.1	275.3	483.2	867.3	2,177.7
2000.....	9.0	17.0	30.3	56.7	95.9	165.3	287.4	496.2	894.4	2,287.5
1999.....	8.8	17.4	32.8	57.3	99.5	175.7	299.3	521.0	924.9	2,353.6
All races 3/, male										
2006.....	12.3	23.9	41.2	66.6	107.4	174.8	296.7	531.1	945.4	1,979.5
2005.....	12.5	23.7	42.4	67.0	116.4	184.3	316.4	558.3	997.9	2,120.9
2004.....	12.7	25.0	41.8	67.5	116.8	191.3	328.2	580.4	1,027.4	2,129.6
2003.....	13.3	25.5	43.0	73.3	125.0	206.7	355.0	616.5	1,088.5	2,260.1
2002.....	13.6	26.6	46.5	78.0	129.6	217.8	367.6	647.1	1,123.8	2,325.6
2001.....	13.2	25.4	45.7	78.8	135.6	228.4	385.2	675.1	1,145.1	2,411.8
2000.....	14.0	27.0	46.7	84.9	139.5	234.0	409.1	684.5	1,204.8	2,561.9
1999.....	13.9	27.7	50.2	84.9	143.5	250.0	424.4	718.0	1,240.3	2,614.4
All races 3/, female										
2006.....	3.8	7.7	12.9	22.6	42.7	75.6	144.0	267.7	547.2	1,586.8
2005.....	4.0	7.4	13.5	23.9	46.5	84.8	154.5	293.0	580.4	1,707.7
2004.....	4.1	8.0	13.6	25.0	46.3	85.6	162.8	300.3	609.8	1,766.7
2003.....	4.5	8.2	13.7	26.0	49.8	90.5	172.7	322.7	641.5	1,926.2
2002.....	4.2	7.8	13.9	28.2	50.5	97.0	177.3	337.3	670.4	1,984.9
2001.....	4.0	7.6	14.7	27.7	55.4	101.2	188.0	348.9	700.7	2,079.8
2000.....	4.0	7.3	14.6	30.3	56.4	106.3	191.6	365.1	711.3	2,175.8
1999.....	3.9	7.4	16.1	31.5	59.7	112.1	201.5	384.0	740.3	2,248.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)										
White, both sexes										
2006 4/.....	8.0	14.9	25.5	41.8	70.8	118.0	210.4	380.4	708.8	1,763.4
2005 4/.....	8.0	14.6	26.0	42.9	76.7	127.6	225.2	405.9	750.9	1,896.0
2004 4/.....	8.1	15.7	25.8	43.5	77.0	131.0	234.7	418.8	778.7	1,929.2
2003 4/.....	8.5	16.0	26.5	46.8	82.2	140.3	251.0	445.0	821.5	2,079.5
2002.....	8.6	16.1	28.3	49.9	84.5	150.0	260.7	466.0	854.2	2,143.3
2001.....	8.1	15.4	28.4	50.5	90.2	156.6	273.8	483.6	877.5	2,233.8
2000.....	8.5	16.0	29.1	54.4	92.2	162.0	285.8	496.4	908.0	2,352.7
1999.....	8.2	16.6	31.1	54.8	96.5	172.6	297.1	522.7	938.2	2,416.1
White, male										
2006 4/.....	12.5	23.1	39.9	64.1	104.8	170.8	295.6	534.6	965.4	2,056.2
2005 4/.....	12.3	22.5	40.4	65.0	113.3	181.1	317.3	564.5	1,021.0	2,204.7
2004 4/.....	12.4	24.3	40.0	65.5	114.0	188.4	328.6	585.8	1,050.7	2,200.2
2003 4/.....	12.9	24.7	41.5	71.3	122.4	203.4	354.1	622.0	1,109.8	2,338.4
2002.....	13.4	25.9	45.0	75.8	125.4	216.8	368.7	655.7	1,151.0	2,403.6
2001.....	12.8	24.4	44.2	76.5	133.1	226.8	385.4	682.4	1,166.2	2,493.6
2000.....	13.7	26.1	45.5	82.7	136.4	233.3	411.5	691.9	1,231.6	2,654.4
1999.....	13.1	27.2	48.1	82.6	140.3	249.9	425.1	727.7	1,269.0	2,695.9
White, female										
2006 4/.....	3.4	6.8	11.4	20.3	39.1	70.9	139.6	265.4	546.6	1,626.0
2005 4/.....	3.6	6.8	11.8	21.5	42.6	80.1	149.1	288.6	581.7	1,753.8
2004 4/.....	3.7	7.2	11.9	22.3	42.8	80.1	157.6	296.5	609.7	1,807.0
2003 4/.....	4.0	7.3	11.8	23.3	45.1	84.5	166.6	317.1	643.8	1,964.9
2002.....	3.6	6.5	12.0	25.2	46.7	91.1	172.7	330.1	672.3	2,030.2
2001.....	3.5	6.4	13.0	25.7	50.7	95.0	183.3	342.8	703.0	2,125.0
2000.....	3.3	5.9	13.0	27.5	51.4	99.6	184.9	358.9	715.6	2,230.1
1999.....	3.2	6.2	14.5	28.3	56.1	105.2	195.4	378.8	743.0	2,303.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Black, both sexes

2006 4/.....	10.5	24.0	42.5	70.1	112.4	175.3	274.8	435.0	711.2	1,363.3
2005 4/.....	11.4	24.0	45.5	72.4	125.2	184.3	284.0	466.7	738.4	1,415.4
2004 4/.....	12.8	23.9	45.2	73.2	123.4	193.3	298.8	473.1	755.5	1,514.0
2003 4/.....	12.8	25.4	46.4	78.5	132.8	206.3	331.2	515.6	804.7	1,652.1
2002.....	12.6	27.0	47.8	84.1	136.1	204.4	319.1	522.2	811.9	1,665.1
2001.....	13.3	26.9	47.9	79.9	140.9	218.1	337.8	546.0	853.7	1,756.7
2000.....	14.1	28.1	46.3	87.0	142.8	222.3	354.7	556.6	831.8	1,740.5
1999.....	14.7	26.3	53.8	90.6	143.1	233.9	362.6	555.9	844.6	1,801.2

Black, male

2006 4/.....	14.0	33.9	61.6	100.1	153.4	241.1	376.8	599.6	884.4	1,353.8
2005 4/.....	15.8	36.4	66.7	101.3	172.1	247.4	368.4	604.8	909.5	1,420.4
2004 4/.....	18.1	34.0	65.5	99.6	172.3	258.0	395.9	623.3	907.5	1,536.8
2003 4/.....	18.0	36.5	64.7	108.6	177.6	278.2	437.5	669.6	1,003.3	1,563.9
2002.....	17.2	37.0	67.8	115.7	195.8	267.5	423.6	633.4	961.4	1,655.3
2001.....	18.8	38.3	69.2	113.1	189.2	286.4	457.8	692.3	1,055.4	1,759.8
2000.....	19.3	39.0	64.5	123.2	189.9	279.7	454.3	686.6	1,019.2	1,744.8
1999.....	21.2	37.0	78.2	124.1	200.3	293.9	472.2	678.6	1,001.9	1,845.5

Black, female

2006 4/.....	7.3	15.3	26.3	45.3	80.4	126.7	204.8	336.4	619.6	1,367.2
2005 4/.....	7.6	13.2	27.6	48.5	88.6	137.5	226.2	384.3	648.0	1,413.4
2004 4/.....	8.1	15.1	28.0	51.4	85.2	145.2	232.8	384.1	675.5	1,505.0
2003 4/.....	8.3	15.7	30.9	53.8	97.9	153.0	259.4	424.0	700.6	1,686.4
2002.....	8.5	18.4	30.8	58.2	89.4	157.6	248.9	455.8	734.1	1,668.8
2001.....	8.5	17.0	29.8	52.6	102.9	167.6	257.8	458.7	751.0	1,755.5
2000.....	9.5	18.7	30.7	57.4	105.6	179.9	288.6	478.2	737.5	1,738.9
1999.....	9.1	17.1	33.1	63.3	98.2	189.5	289.9	482.0	766.6	1,784.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
All races 3/, both sexes										
2006.....	11.4	17.7	25.0	36.3	55.5	85.7	144.6	259.0	493.3	1,342.8
2005.....	11.6	17.9	25.5	36.8	57.4	88.2	150.8	264.6	500.7	1,377.4
2004.....	11.5	17.9	25.6	35.7	57.5	90.3	151.8	268.1	503.3	1,373.9
2003.....	12.6	17.9	25.3	39.1	59.7	97.3	160.2	279.1	525.5	1,427.2
2002.....	12.4	18.1	25.9	39.4	60.6	97.9	164.5	286.7	536.3	1,443.5
2001.....	11.9	17.9	25.9	39.7	62.0	100.7	164.7	287.2	537.0	1,475.5
2000.....	12.3	17.6	25.9	40.2	64.3	103.6	167.1	290.2	542.8	1,509.4
1999.....	12.5	17.8	25.9	41.8	66.5	105.2	175.0	295.4	553.5	1,512.9
All races 3/, male										
2006.....	14.0	22.6	32.4	46.6	69.5	106.9	173.2	302.8	566.0	1,338.1
2005.....	14.8	22.6	33.3	47.4	71.3	108.0	180.8	310.3	568.1	1,377.8
2004.....	14.4	23.2	32.8	46.4	72.7	111.3	181.2	316.2	573.7	1,377.7
2003.....	15.6	22.7	32.4	49.9	74.9	119.4	195.4	334.0	599.9	1,431.3
2002.....	15.9	23.5	33.6	49.4	75.7	118.8	199.8	340.7	612.4	1,448.6
2001.....	14.9	23.1	32.7	49.4	77.7	121.5	200.9	344.7	622.1	1,473.4
2000.....	15.4	23.0	33.4	52.0	80.8	126.9	205.9	345.1	626.8	1,549.1
1999.....	15.6	23.0	33.3	54.1	84.3	133.4	217.0	350.9	641.4	1,548.9
All races 3/, female										
2006.....	8.9	12.9	18.0	26.6	42.7	67.2	121.3	226.9	447.9	1,345.0
2005.....	8.5	13.3	18.0	26.9	44.8	70.9	126.4	231.4	458.9	1,377.3
2004.....	8.6	12.8	18.6	25.7	43.8	72.0	128.2	233.5	459.9	1,372.1
2003.....	9.5	13.2	18.6	28.8	45.9	78.1	131.9	240.0	480.0	1,425.4
2002.....	9.0	12.8	18.5	30.1	46.8	79.8	136.3	248.6	490.0	1,441.3
2001.....	8.9	12.8	19.5	30.6	47.8	82.8	135.9	247.0	486.0	1,476.4
2000.....	9.2	12.3	18.7	29.2	49.3	83.6	136.6	252.1	493.2	1,493.2
1999.....	9.4	12.8	18.9	30.2	50.5	81.0	142.2	256.8	502.0	1,498.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
White, both sexes										
2006 4/.....	9.9	15.2	21.4	32.0	50.8	79.9	139.1	256.9	496.7	1,386.4
2005 4/.....	10.0	15.2	21.6	32.4	51.6	81.9	145.2	260.8	504.5	1,419.0
2004 4/.....	9.7	15.0	21.8	31.3	52.7	84.7	146.5	262.9	507.6	1,408.8
2003 4/.....	10.5	15.2	21.4	34.8	53.8	90.7	153.2	276.4	528.9	1,461.4
2002.....	10.7	15.0	22.1	35.0	55.3	92.4	158.5	282.4	539.1	1,475.5
2001.....	9.7	14.9	22.3	34.8	57.2	94.1	159.0	281.6	538.8	1,509.6
2000.....	10.2	14.6	22.1	35.9	59.0	97.3	160.6	284.5	545.7	1,541.6
1999.....	10.2	14.6	22.2	36.3	60.4	98.6	167.6	288.5	553.7	1,543.3
White, male										
2006 4/.....	12.1	19.5	27.7	41.3	63.5	100.0	166.3	300.6	572.3	1,381.2
2005 4/.....	12.8	19.6	28.7	41.9	64.7	100.4	173.5	306.2	575.6	1,424.2
2004 4/.....	12.4	19.6	28.7	40.9	66.8	104.6	176.0	309.6	578.2	1,407.9
2003 4/.....	13.5	19.7	27.5	44.5	68.3	111.8	188.4	332.0	604.3	1,467.6
2002.....	14.1	19.9	29.2	44.1	69.9	112.5	193.0	336.5	617.5	1,482.1
2001.....	12.7	19.7	28.9	43.5	72.1	114.9	193.7	338.2	621.9	1,503.4
2000.....	13.0	19.7	28.6	46.9	74.8	119.7	199.2	339.4	630.8	1,580.5
1999.....	13.3	19.4	28.7	47.5	76.9	125.6	209.0	344.0	641.4	1,571.5
White, female										
2006 4/.....	7.7	11.0	15.2	23.0	38.9	62.0	116.6	224.3	449.0	1,388.9
2005 4/.....	7.1	10.8	14.5	23.3	39.5	65.4	121.9	227.1	460.0	1,416.6
2004 4/.....	6.9	10.5	15.1	22.0	39.7	66.9	122.2	228.7	463.8	1,409.2
2003 4/.....	7.5	10.8	15.4	25.5	40.4	72.1	124.3	236.1	482.4	1,458.6
2002.....	7.3	10.2	15.2	26.2	41.8	74.7	130.3	243.7	491.0	1,472.6
2001.....	6.6	10.1	15.8	26.5	43.5	75.9	130.9	241.5	488.5	1,512.2
2000.....	7.4	9.6	15.8	25.4	44.3	77.7	129.5	245.9	495.2	1,525.8
1999.....	7.2	9.8	15.8	25.7	45.1	75.0	134.8	249.5	501.9	1,531.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other heart diseases (I26-I51)										
Black, both sexes										
2006 4/.....	24.1	38.4	57.4	80.0	111.5	155.8	229.4	342.1	566.8	1,094.3
2005 4/.....	25.3	40.3	60.2	82.0	120.4	162.1	237.3	368.3	563.4	1,135.3
2004 4/.....	26.3	41.7	60.2	82.6	114.5	159.9	240.5	381.0	551.3	1,180.7
2003 4/.....	28.9	40.8	61.1	85.3	124.6	176.1	265.5	364.8	574.9	1,238.1
2002.....	26.6	43.6	60.8	87.1	121.2	169.4	255.8	386.7	594.7	1,270.4
2001.....	29.2	43.1	60.9	91.4	118.8	179.7	254.5	402.9	601.8	1,277.7
2000.....	28.5	43.3	62.0	86.2	122.8	181.5	266.8	406.9	590.7	1,306.0
1999.....	30.2	44.9	62.2	96.8	133.5	185.5	284.2	417.4	621.2	1,302.8
Black, male										
2006 4/.....	30.0	49.8	76.5	102.5	147.0	199.2	288.8	409.0	645.5	1,101.8
2005 4/.....	31.8	50.1	77.6	107.6	151.1	204.4	300.1	446.6	614.5	1,088.5
2004 4/.....	31.6	54.3	75.0	109.0	149.1	202.5	284.5	469.5	653.5	1,224.9
2003 4/.....	34.7	51.0	79.5	113.5	158.3	223.6	318.0	434.1	668.1	1,220.9
2002.....	32.0	56.7	77.1	110.7	150.7	208.7	317.5	454.2	675.6	1,238.3
2001.....	33.7	53.5	73.7	117.2	151.5	212.9	318.8	494.8	743.2	1,349.1
2000.....	34.8	54.3	82.1	112.1	153.6	224.8	318.9	480.5	696.3	1,402.3
1999.....	36.2	56.1	81.9	127.0	172.7	237.8	353.1	489.1	747.3	1,454.1
Black, female										
2006 4/.....	18.8	28.5	41.3	61.4	83.9	123.8	188.5	302.1	525.2	1,091.2
2005 4/.....	19.6	31.7	45.4	60.8	96.5	130.8	194.4	321.6	536.4	1,154.2
2004 4/.....	21.6	30.6	47.7	60.9	87.6	128.3	210.7	328.6	497.4	1,163.2
2003 4/.....	23.8	31.9	45.4	62.1	98.2	140.9	230.2	323.7	526.1	1,244.8
2002.....	21.8	32.3	47.0	67.7	98.2	140.2	214.4	346.3	552.6	1,282.7
2001.....	25.3	34.0	50.1	70.3	93.1	155.1	211.6	348.1	529.8	1,251.2
2000.....	22.9	33.9	44.8	64.9	98.4	149.6	232.2	362.6	537.5	1,270.2
1999.....	24.9	35.2	45.4	72.1	102.8	146.8	238.5	374.2	558.7	1,246.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
All races 3/, both sexes										
2006.....	1.7	2.5	3.6	4.8	6.7	9.5	13.9	21.2	30.4	50.3
2005.....	1.9	2.7	3.7	5.2	7.6	11.4	16.8	22.3	34.5	54.4
2004.....	1.9	2.9	3.7	4.7	7.1	11.0	16.3	23.3	33.1	54.8
2003.....	2.2	2.8	4.0	5.4	8.1	11.6	17.5	25.8	36.9	58.0
2002.....	1.9	2.9	3.8	5.9	8.1	12.7	17.5	25.7	35.5	56.3
2001.....	2.1	2.9	3.9	5.9	8.2	12.0	17.3	24.2	34.8	57.5
2000.....	2.1	2.8	4.2	5.2	8.5	12.3	16.8	24.1	34.7	57.4
1999.....	2.1	2.8	3.9	5.6	8.0	11.9	17.9	25.7	36.5	59.6
All races 3/, male										
2006.....	1.5	2.2	3.5	4.4	6.0	9.4	13.4	19.8	27.0	43.1
2005.....	1.7	2.3	3.7	5.0	7.1	11.6	16.0	21.9	33.1	47.8
2004.....	1.7	2.7	3.6	5.1	6.8	10.3	15.2	21.5	30.6	48.3
2003.....	1.8	2.7	4.0	5.1	7.7	11.4	17.5	26.5	35.0	54.4
2002.....	1.6	2.8	3.5	5.3	8.0	12.3	16.3	23.8	33.8	52.1
2001.....	1.9	2.6	3.4	5.1	8.2	11.7	17.6	23.1	35.5	55.6
2000.....	1.7	2.7	4.0	4.9	8.3	11.5	16.6	25.1	33.5	54.8
1999.....	1.8	2.5	3.6	5.3	8.6	12.0	18.5	25.5	35.2	60.3
All races 3/, female										
2006.....	2.0	2.7	3.7	5.1	7.2	9.6	14.3	22.3	32.6	53.7
2005.....	2.2	3.1	3.7	5.4	8.0	11.4	17.5	22.5	35.3	57.5
2004.....	2.0	3.1	3.7	4.5	7.3	11.7	17.2	24.5	34.7	57.7
2003.....	2.6	2.8	4.0	5.7	8.4	11.9	17.6	25.3	38.0	59.6
2002.....	2.2	3.1	4.1	6.4	8.3	13.0	18.4	27.1	36.6	58.1
2001.....	2.2	3.1	4.4	6.6	8.2	12.3	17.1	24.9	34.4	58.2
2000.....	2.5	3.0	4.4	5.5	8.8	12.9	17.1	23.5	35.3	58.5
1999.....	2.3	3.1	4.1	5.9	7.5	11.8	17.4	25.8	37.2	59.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
White, both sexes										
2006 4/.....	1.5	2.1	3.4	4.3	6.2	8.7	13.3	20.7	30.0	50.9
2005 4/.....	1.6	2.4	3.3	4.8	6.8	10.8	16.3	21.6	33.7	54.3
2004 4/.....	1.6	2.6	3.3	4.3	6.6	10.2	15.5	22.2	32.4	55.3
2003 4/.....	1.8	2.5	3.6	4.9	7.4	10.7	16.1	25.4	36.5	57.8
2002.....	1.7	2.6	3.5	5.3	7.7	12.0	16.7	24.8	35.1	56.1
2001.....	1.8	2.5	3.6	5.3	7.7	11.2	16.7	23.2	34.6	56.8
2000.....	1.8	2.5	3.7	4.9	7.9	11.7	16.1	23.3	34.2	56.4
1999.....	1.7	2.4	3.5	4.9	7.2	11.0	16.7	24.7	35.3	59.2
White, male										
2006 4/.....	1.3	1.9	3.4	4.1	5.7	8.6	13.1	19.4	26.9	43.0
2005 4/.....	1.4	2.1	3.4	4.8	6.4	11.0	15.7	21.3	32.7	46.9
2004 4/.....	1.6	2.5	3.3	4.5	6.5	9.9	14.6	20.5	28.6	48.0
2003 4/.....	1.5	2.4	3.6	4.6	7.3	10.6	16.4	26.3	34.8	54.0
2002.....	1.5	2.5	3.5	4.8	7.9	11.8	15.9	23.2	33.1	51.9
2001.....	1.7	2.3	3.3	4.5	7.7	11.2	17.1	22.2	34.8	53.6
2000.....	1.4	2.5	3.6	4.7	7.8	11.3	16.2	24.6	33.0	52.9
1999.....	1.6	2.3	3.3	4.9	7.8	11.2	17.4	24.7	33.8	60.3
White, female										
2006 4/.....	1.7	2.3	3.4	4.4	6.6	8.9	13.4	21.7	32.0	54.6
2005 4/.....	1.8	2.7	3.3	4.8	7.2	10.7	16.9	21.8	34.3	57.7
2004 4/.....	1.6	2.6	3.3	4.0	6.8	10.6	16.2	23.4	34.8	58.6
2003 4/.....	2.2	2.6	3.6	5.2	7.6	10.8	15.9	24.8	37.5	59.5
2002.....	1.9	2.7	3.5	5.8	7.4	12.1	17.4	25.9	36.3	58.0
2001.....	1.9	2.7	3.9	6.1	7.7	11.2	16.3	23.9	34.4	58.1
2000.....	2.1	2.6	3.8	5.2	8.1	12.1	16.0	22.5	35.0	57.8
1999.....	1.9	2.6	3.6	5.0	6.7	10.9	16.1	24.7	36.2	58.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary heart disease and diseases of pulmonary circulation (I26-I28)										
Black, both sexes										
2006 4/.....	3.9	5.4	6.4	10.5	13.5	18.9	24.8	31.3	43.5	55.3
2005 4/.....	4.3	5.7	7.6	9.9	16.2	20.5	25.8	36.6	51.7	66.3
2004 4/.....	4.2	5.9	7.6	10.2	13.3	21.0	29.1	42.8	46.8	62.0
2003 4/.....	5.2	5.6	8.1	11.0	16.0	23.5	36.6	36.9	50.0	72.9
2002.....	3.9	6.4	7.3	12.8	15.0	22.6	30.3	43.6	49.5	68.6
2001.....	4.5	6.3	7.4	12.4	15.3	23.8	28.5	39.8	45.2	77.7
2000.....	4.9	5.8	9.3	8.9	16.3	20.9	29.2	39.3	46.9	76.6
1999.....	4.9	6.2	8.6	12.6	17.4	23.0	35.9	43.7	57.8	72.1
Black, male										
2006 4/.....	3.7	4.9	5.9	8.8	12.2	20.4	21.2	26.8	38.3	57.2
2005 4/.....	3.9	4.9	6.9	8.7	15.3	21.4	24.4	38.4	48.5	67.9
2004 4/.....	3.2	4.9	7.1	11.3	11.8	17.7	24.9	41.9	62.4	64.6
2003 4/.....	4.7	5.9	8.2	11.5	14.5	23.1	34.5	37.0	50.2	76.1
2002.....	2.9	6.2	4.9	11.9	11.7	21.7	25.0	37.7	51.2	65.4
2001.....	3.8	5.5	5.2	11.9	16.3	21.1	28.6	38.1	54.6	95.3
2000.....	3.9	4.8	8.5	9.2	15.9	17.2	26.4	38.7	50.6	92.3
1999.....	3.9	5.0	7.9	10.3	18.9	23.9	38.3	43.2	62.8	71.5
Black, female										
2006 4/.....	4.1	5.9	6.9	11.8	14.5	17.8	27.3	34.0	46.3	54.5
2005 4/.....	4.7	6.4	8.1	10.8	17.0	19.9	26.8	35.5	53.4	65.6
2004 4/.....	5.2	6.9	8.0	9.2	14.4	23.4	31.9	43.4	38.6	61.0
2003 4/.....	5.7	5.3	8.0	10.5	17.1	23.8	38.1	36.9	49.9	71.7
2002.....	4.9	6.7	9.3	13.6	17.7	23.3	33.8	47.2	48.5	69.8
2001.....	5.2	7.0	9.3	12.8	14.6	25.7	28.5	40.8	40.5	71.1
2000.....	5.7	6.7	9.9	8.6	16.7	23.6	31.1	39.6	45.0	70.8
1999.....	5.7	7.3	9.2	14.4	16.3	22.3	34.2	44.1	55.3	72.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary embolism (I26)										
All races 3/, both sexes										
2006.....	1.2	1.5	2.2	2.8	3.9	5.1	7.5	11.7	16.2	26.7
2005.....	1.3	1.7	2.5	3.3	4.7	7.0	9.8	13.9	20.7	32.8
2004.....	1.2	2.0	2.4	3.0	4.5	6.8	9.7	14.4	20.7	33.4
2003.....	1.5	1.9	2.8	3.5	5.1	7.0	10.7	15.7	23.2	35.5
2002.....	1.3	2.0	2.5	3.9	5.3	8.1	11.0	15.8	22.2	35.4
2001.....	1.4	1.9	2.7	4.1	5.4	7.8	11.4	15.7	22.3	37.8
2000.....	1.4	1.9	2.8	3.5	5.8	7.9	10.8	15.7	22.5	38.4
1999.....	1.4	2.0	2.7	3.9	5.6	7.9	12.6	17.9	25.8	42.0
All races 3/, male										
2006.....	1.1	1.4	2.4	2.9	4.0	5.6	8.1	11.9	16.5	24.3
2005.....	1.3	1.6	2.7	3.5	4.8	7.5	10.1	14.9	20.7	30.2
2004.....	1.2	1.9	2.4	3.5	4.7	7.2	9.9	14.3	20.5	30.6
2003.....	1.2	2.0	2.8	3.8	5.4	7.7	11.4	16.9	23.5	34.1
2002.....	1.1	2.0	2.4	3.8	5.5	8.2	10.8	15.0	21.3	34.1
2001.....	1.3	1.8	2.4	3.6	5.6	8.4	11.8	16.1	23.5	38.0
2000.....	1.1	2.0	3.0	3.5	6.0	8.0	11.2	17.4	22.4	37.5
1999.....	1.2	1.9	2.6	4.1	6.3	8.3	13.7	18.1	24.2	42.0
All races 3/, female										
2006.....	1.3	1.6	2.1	2.7	3.9	4.6	7.1	11.5	16.1	27.8
2005.....	1.3	1.8	2.3	3.1	4.5	6.6	9.6	13.2	20.7	34.0
2004.....	1.3	2.2	2.4	2.4	4.4	6.6	9.6	14.5	20.8	34.7
2003.....	1.7	1.8	2.7	3.3	4.9	6.4	10.2	14.8	23.0	36.0
2002.....	1.4	1.9	2.6	4.0	5.1	8.0	11.1	16.4	22.8	36.0
2001.....	1.5	2.1	2.9	4.6	5.2	7.3	11.1	15.4	21.5	37.7
2000.....	1.6	1.9	2.6	3.5	5.6	7.8	10.6	14.5	22.5	38.8
1999.....	1.6	2.1	2.8	3.8	5.0	7.5	11.7	17.8	26.7	42.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary embolism (I26)										
White, both sexes										
2006 4/.....	1.0	1.3	2.2	2.6	3.6	4.7	7.3	11.2	15.8	26.7
2005 4/.....	1.1	1.5	2.3	3.1	4.4	6.6	9.5	13.2	20.0	32.7
2004 4/.....	1.0	1.8	2.1	2.7	4.3	6.3	9.1	13.7	20.2	33.4
2003 4/.....	1.2	1.7	2.5	3.3	4.8	6.3	9.8	15.5	22.8	34.8
2002.....	1.2	1.8	2.3	3.5	5.1	7.7	10.4	15.1	21.7	34.8
2001.....	1.2	1.7	2.5	3.8	5.1	7.4	10.9	15.0	22.1	37.0
2000.....	1.1	1.7	2.5	3.3	5.4	7.4	10.4	15.0	22.0	37.4
1999.....	1.2	1.7	2.4	3.5	5.1	7.3	11.6	17.0	25.1	41.7
White, male										
2006 4/.....	0.9	1.2	2.3	2.8	3.7	5.0	8.0	11.5	16.2	23.9
2005 4/.....	1.1	1.4	2.6	3.3	4.5	7.0	9.7	14.1	20.1	29.7
2004 4/.....	1.1	1.8	2.2	3.2	4.5	6.8	9.3	13.3	19.0	30.0
2003 4/.....	1.1	1.7	2.6	3.4	5.0	7.1	10.7	16.6	23.0	33.2
2002.....	1.1	1.8	2.4	3.4	5.4	7.8	10.6	14.6	20.6	33.3
2001.....	1.1	1.6	2.4	3.3	5.2	8.1	11.5	15.4	22.9	36.1
2000.....	0.9	1.8	2.7	3.2	5.7	7.8	11.0	16.8	21.6	35.8
1999.....	1.1	1.7	2.3	3.7	5.7	7.6	12.8	17.2	23.0	41.3
White, female										
2006 4/.....	1.1	1.4	2.0	2.4	3.5	4.3	6.7	11.0	15.5	28.1
2005 4/.....	1.1	1.6	2.0	2.9	4.3	6.2	9.3	12.6	19.9	34.1
2004 4/.....	1.0	1.8	2.1	2.3	4.1	5.8	9.0	13.9	21.0	34.9
2003 4/.....	1.4	1.7	2.4	3.1	4.5	5.6	9.1	14.7	22.6	35.6
2002.....	1.3	1.8	2.3	3.6	4.7	7.6	10.3	15.5	22.4	35.5
2001.....	1.3	1.8	2.5	4.4	5.0	6.8	10.5	14.7	21.6	37.4
2000.....	1.4	1.6	2.3	3.3	5.2	7.1	9.9	13.6	22.2	38.1
1999.....	1.3	1.7	2.5	3.4	4.4	7.1	10.6	16.8	26.4	41.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pulmonary embolism (I26)										
Black, both sexes										
2006 4/.....	2.9	3.2	3.6	5.2	8.4	10.4	12.9	19.9	26.5	33.2
2005 4/.....	3.1	3.4	4.9	6.0	8.8	12.9	16.3	25.7	35.2	40.6
2004 4/.....	3.0	4.4	4.9	5.8	8.2	13.8	18.1	27.4	30.7	42.6
2003 4/.....	3.3	3.6	5.8	7.3	9.6	15.3	23.6	23.0	34.8	51.1
2002.....	2.4	4.1	5.0	8.5	9.1	14.3	19.4	28.7	33.5	49.0
2001.....	3.1	4.4	5.0	7.9	9.9	14.3	19.2	27.3	30.3	55.9
2000.....	3.3	4.3	6.1	6.0	11.1	14.7	18.5	28.8	33.0	56.6
1999.....	3.2	4.4	6.0	8.8	12.1	15.7	26.4	33.4	40.7	53.2
Black, male										
2006 4/.....	2.9	3.1	4.1	5.4	8.3	13.6	12.3	19.2	28.1	38.4
2005 4/.....	3.2	3.6	4.8	6.4	9.8	14.5	18.0	30.4	35.1	43.7
2004 4/.....	2.4	3.6	4.6	7.9	7.8	12.8	17.8	31.8	45.2	45.7
2003 4/.....	3.0	4.5	5.8	8.6	10.4	16.7	22.7	27.4	39.9	57.3
2002.....	1.8	4.6	3.8	8.6	8.5	14.5	16.0	24.8	35.9	51.5
2001.....	2.9	4.0	3.4	8.4	12.0	14.2	18.2	28.8	40.1	72.6
2000.....	3.0	3.9	7.0	7.4	12.0	12.7	17.1	30.2	38.8	71.5
1999.....	2.6	3.8	5.9	8.8	13.4	18.7	29.2	35.6	49.7	60.9
Black, female										
2006 4/.....	2.9	3.2	3.2	5.1	8.4	8.0	13.3	20.4	25.7	31.1
2005 4/.....	3.0	3.3	4.9	5.7	7.9	11.6	15.2	22.9	35.2	39.4
2004 4/.....	3.5	5.1	5.1	4.1	8.4	14.6	18.2	24.8	23.1	41.3
2003 4/.....	3.6	2.8	5.8	6.2	9.0	14.3	24.1	20.3	32.1	48.6
2002.....	3.0	3.7	6.1	8.4	9.6	14.2	21.7	31.0	32.2	48.1
2001.....	3.2	4.8	6.4	7.4	8.3	14.3	20.0	26.3	25.3	49.7
2000.....	3.6	4.7	5.4	4.9	10.3	16.1	19.5	27.9	30.1	51.1
1999.....	3.7	5.0	6.1	8.7	11.1	13.5	24.5	32.0	36.3	50.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

All races 3/, both sexes

2006.....	0.5	1.0	1.4	2.0	2.8	4.4	6.4	9.6	14.2	23.6
2005.....	0.6	1.0	1.2	1.9	2.9	4.4	7.0	8.4	13.7	21.6
2004.....	0.6	0.9	1.3	1.8	2.6	4.2	6.6	8.9	12.4	21.4
2003.....	0.8	0.9	1.2	1.8	2.9	4.6	6.8	10.1	13.6	22.6
2002.....	0.6	1.0	1.3	2.0	2.9	4.6	6.5	9.9	13.3	20.9
2001.....	0.7	1.0	1.2	1.8	2.8	4.2	6.0	8.5	12.5	19.6
2000.....	0.7	0.9	1.4	1.7	2.7	4.4	6.0	8.4	12.2	19.0
1999.....	0.7	0.9	1.2	1.7	2.4	4.0	5.4	7.8	10.7	17.5

All races 3/, male

2006.....	0.4	0.8	1.1	1.4	2.1	3.7	5.3	8.0	10.5	18.8
2005.....	0.4	0.7	1.0	1.5	2.2	4.1	6.0	7.0	12.4	17.5
2004.....	0.5	0.8	1.2	1.5	2.2	3.1	5.3	7.2	10.0	17.7
2003.....	0.6	0.7	1.1	1.3	2.3	3.7	6.0	9.6	11.4	20.3
2002.....	0.4	0.7	1.1	1.5	2.5	4.1	5.5	8.8	12.5	18.0
2001.....	0.6	0.9	1.0	1.5	2.6	3.3	5.9	7.0	11.9	17.6
2000.....	0.5	0.7	1.0	1.4	2.3	3.5	5.4	7.7	11.1	17.3
1999.....	0.6	0.7	1.1	1.3	2.3	3.7	4.9	7.4	10.9	18.4

All races 3/, female

2006.....	0.6	1.1	1.6	2.5	3.4	5.0	7.2	10.7	16.5	25.9
2005.....	0.8	1.3	1.4	2.3	3.5	4.7	7.9	9.4	14.6	23.5
2004.....	0.7	0.9	1.4	2.0	3.0	5.1	7.6	10.0	13.9	23.1
2003.....	0.9	1.1	1.3	2.3	3.5	5.5	7.4	10.5	15.0	23.6
2002.....	0.8	1.2	1.5	2.4	3.2	4.9	7.3	10.7	13.8	22.1
2001.....	0.8	1.1	1.5	2.0	3.0	5.0	6.0	9.5	12.9	20.5
2000.....	0.9	1.1	1.8	2.0	3.2	5.2	6.5	8.9	12.8	19.7
1999.....	0.8	1.1	1.4	2.0	2.5	4.3	5.7	8.0	10.5	17.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)										
White, both sexes										
2006 4/.....	0.5	0.8	1.2	1.6	2.6	4.1	6.0	9.6	14.2	24.2
2005 4/.....	0.5	0.9	1.0	1.7	2.5	4.2	6.9	8.4	13.7	21.6
2004 4/.....	0.6	0.8	1.2	1.5	2.4	4.0	6.3	8.6	12.2	21.9
2003 4/.....	0.6	0.8	1.1	1.7	2.7	4.4	6.3	9.9	13.7	23.0
2002.....	0.5	0.8	1.2	1.7	2.6	4.3	6.3	9.7	13.3	21.3
2001.....	0.6	0.8	1.1	1.5	2.6	3.8	5.8	8.2	12.5	19.7
2000.....	0.6	0.9	1.2	1.7	2.5	4.3	5.7	8.4	12.3	19.0
1999.....	0.5	0.7	1.1	1.4	2.1	3.7	5.1	7.7	10.2	17.6
White, male										
2006 4/.....	0.4	0.7	1.1	1.3	2.0	3.6	5.1	7.9	10.7	19.1
2005 4/.....	0.3	0.7	0.9	1.5	2.0	4.0	6.0	7.1	12.6	17.2
2004 4/.....	0.5	0.8	1.1	1.4	2.0	3.1	5.2	7.2	9.6	18.0
2003 4/.....	0.5	0.6	1.0	1.2	2.3	3.5	5.7	9.7	11.8	20.8
2002.....	0.4	0.6	1.1	1.3	2.5	3.9	5.3	8.7	12.5	18.6
2001.....	0.5	0.8	0.9	1.3	2.5	3.1	5.6	6.8	11.9	17.4
2000.....	0.5	0.7	1.0	1.5	2.2	3.4	5.2	7.8	11.4	17.1
1999.....	0.4	0.6	1.0	1.3	2.0	3.5	4.6	7.5	10.8	19.0
White, female										
2006 4/.....	0.6	0.9	1.4	2.0	3.1	4.5	6.7	10.8	16.5	26.6
2005 4/.....	0.7	1.1	1.2	1.9	3.0	4.4	7.6	9.2	14.4	23.6
2004 4/.....	0.6	0.8	1.2	1.7	2.7	4.7	7.2	9.5	13.8	23.7
2003 4/.....	0.8	0.9	1.2	2.1	3.0	5.1	6.8	10.1	14.9	23.9
2002.....	0.7	1.0	1.3	2.1	2.7	4.6	7.1	10.4	13.9	22.5
2001.....	0.6	0.9	1.3	1.7	2.7	4.4	5.9	9.2	12.8	20.7
2000.....	0.7	1.0	1.5	1.9	2.8	5.0	6.1	8.8	12.8	19.8
1999.....	0.6	0.8	1.2	1.6	2.2	3.8	5.4	7.8	9.8	17.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other pulmonary heart disease and diseases of pulmonary circulation (I27-I28)

Black, both sexes

2006 4/.....	1.0	2.2	2.8	5.2	5.1	8.6	11.9	11.4	17.0	22.1
2005 4/.....	1.2	2.3	2.7	3.9	7.5	7.7	9.5	10.9	16.5	25.7
2004 4/.....	1.3	1.6	2.7	4.3	5.1	7.1	11.0	15.4	16.1	19.5
2003 4/.....	1.9	2.0	2.3	3.7	6.3	8.2	13.1	14.0	15.2	21.9
2002.....	1.5	2.3	2.2	4.3	5.9	8.3	10.9	14.9	16.0	19.6
2001.....	1.5	1.9	2.4	4.5	5.4	9.5	9.3	12.5	14.9	21.8
2000.....	1.6	1.5	3.1	2.9	5.3	6.2	10.7	10.5	13.8	20.0
1999.....	1.7	1.8	2.6	3.8	5.3	7.3	9.5	10.4	17.0	18.8

Black, male

2006 4/.....	*	1.8	1.8	3.4	3.9	6.8	8.9	*	*	18.8
2005 4/.....	*	*	2.1	*	5.5	6.9	6.3	*	*	24.2
2004 4/.....	*	*	2.5	3.4	3.9	4.8	7.1	10.0	17.2	*
2003 4/.....	1.7	*	2.4	2.9	4.0	6.4	11.8	9.6	*	*
2002.....	*	*	*	3.3	*	7.2	9.0	12.9	15.3	*
2001.....	*	*	*	3.4	4.3	6.9	10.4	9.3	*	22.7
2000.....	*	*	*	*	*	*	9.4	*	*	*
1999.....	*	*	*	*	5.5	5.2	9.2	*	*	*

Black, female

2006 4/.....	*	2.7	3.7	6.7	6.1	9.9	13.9	13.7	20.6	23.4
2005 4/.....	1.7	3.1	3.2	5.2	9.1	8.3	11.7	12.5	18.2	26.3
2004 4/.....	1.7	1.8	2.9	5.1	6.0	8.8	13.6	18.6	15.5	19.7
2003 4/.....	2.1	2.5	2.2	4.3	8.1	9.5	14.0	16.5	17.8	23.1
2002.....	1.8	3.0	3.3	5.1	8.1	9.1	12.1	16.2	16.3	21.8
2001.....	2.0	2.3	2.9	5.4	6.2	11.5	8.5	14.4	15.1	21.5
2000.....	2.1	2.1	4.5	3.8	6.4	7.5	11.5	11.7	14.9	19.7
1999.....	1.9	2.4	3.1	5.7	5.2	8.9	9.7	12.0	19.0	21.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other non-pulmonary forms of heart disease (I30-I51)

All races 3/, both sexes

2006.....	9.7	15.3	21.4	31.6	48.8	76.2	130.7	237.7	462.9	1,292.5
2005.....	9.7	15.2	21.8	31.6	49.9	76.8	133.9	242.3	466.3	1,323.0
2004.....	9.6	15.0	21.9	31.0	50.4	79.2	135.6	244.8	470.2	1,319.1
2003.....	10.3	15.1	21.4	33.7	51.6	85.6	142.7	253.3	488.6	1,369.2
2002.....	10.5	15.1	22.1	33.6	52.4	85.2	147.1	261.0	500.8	1,387.2
2001.....	9.8	15.0	22.0	33.8	53.8	88.6	147.4	263.0	502.3	1,418.1
2000.....	10.2	14.8	21.7	35.0	55.7	91.3	150.3	266.1	508.1	1,452.0
1999.....	10.4	15.0	22.0	36.2	58.5	93.3	157.1	269.7	517.0	1,453.4

All races 3/, male

2006.....	12.5	20.4	28.9	42.2	63.4	97.5	159.8	283.0	539.1	1,295.0
2005.....	13.1	20.3	29.5	42.4	64.2	96.5	164.8	288.4	535.0	1,330.0
2004.....	12.7	20.5	29.2	41.3	65.8	101.0	166.0	294.7	543.2	1,329.4
2003.....	13.8	20.1	28.5	44.9	67.1	108.0	178.0	307.6	564.9	1,376.9
2002.....	14.3	20.7	30.0	44.1	67.8	106.5	183.6	316.9	578.7	1,396.5
2001.....	13.0	20.5	29.3	44.3	69.5	109.7	183.3	321.6	586.6	1,417.8
2000.....	13.7	20.3	29.4	47.1	72.5	115.4	189.3	320.0	593.3	1,494.4
1999.....	13.8	20.5	29.7	48.8	75.7	121.4	198.5	325.4	606.3	1,488.5

All races 3/, female

2006.....	6.9	10.3	14.2	21.5	35.5	57.7	106.9	204.6	415.3	1,291.3
2005.....	6.4	10.2	14.3	21.5	36.8	59.5	108.9	208.9	423.6	1,319.8
2004.....	6.6	9.7	14.9	21.2	36.4	60.3	111.0	209.0	425.2	1,314.4
2003.....	6.9	10.3	14.6	23.2	37.5	66.3	114.4	214.7	442.0	1,365.8
2002.....	6.8	9.7	14.4	23.6	38.5	66.8	117.9	221.5	453.4	1,383.1
2001.....	6.7	9.6	15.1	24.0	39.6	70.4	118.8	222.0	451.7	1,418.2
2000.....	6.7	9.3	14.3	23.7	40.6	70.7	119.5	228.6	457.9	1,434.8
1999.....	7.0	9.7	14.7	24.4	42.9	69.2	124.8	231.0	464.8	1,439.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other non-pulmonary forms of heart disease (I30-I51)

White, both sexes

2006 4/.....	8.4	13.1	18.0	27.7	44.6	71.2	125.9	236.2	466.8	1,335.5
2005 4/.....	8.3	12.8	18.2	27.6	44.8	71.0	128.9	239.2	470.8	1,364.7
2004 4/.....	8.1	12.5	18.5	27.0	46.1	74.4	131.0	240.7	475.2	1,353.6
2003 4/.....	8.7	12.8	17.8	29.9	46.4	80.0	137.1	250.9	492.4	1,403.6
2002.....	9.0	12.4	18.6	29.7	47.6	80.4	141.8	257.7	504.0	1,419.3
2001.....	7.9	12.3	18.7	29.5	49.5	83.0	142.3	258.4	504.2	1,452.9
2000.....	8.5	12.1	18.4	30.9	51.0	85.6	144.5	261.2	511.5	1,485.2
1999.....	8.5	12.1	18.7	31.4	53.2	87.6	151.0	263.8	518.4	1,484.0

White, male

2006 4/.....	10.8	17.6	24.3	37.3	57.9	91.4	153.1	281.2	545.5	1,338.2
2005 4/.....	11.4	17.5	25.3	37.0	58.2	89.4	157.8	284.9	542.9	1,377.4
2004 4/.....	10.8	17.1	25.4	36.4	60.3	94.8	161.5	289.1	549.6	1,359.9
2003 4/.....	12.0	17.3	23.9	39.9	61.0	101.1	172.1	305.7	569.5	1,413.6
2002.....	12.6	17.4	25.7	39.3	62.0	100.7	177.1	313.3	584.4	1,430.2
2001.....	11.0	17.3	25.6	38.9	64.5	103.7	176.6	316.0	587.2	1,449.9
2000.....	11.6	17.2	25.0	42.2	67.0	108.4	183.0	314.8	597.8	1,527.6
1999.....	11.7	17.1	25.5	42.6	69.2	114.4	191.6	319.3	607.6	1,511.1

White, female

2006 4/.....	6.0	8.7	11.8	18.5	32.3	53.1	103.3	202.6	417.0	1,334.3
2005 4/.....	5.2	8.1	11.3	18.5	32.3	54.7	105.0	205.3	425.6	1,358.9
2004 4/.....	5.3	7.9	11.8	18.0	32.9	56.4	106.0	205.3	429.0	1,350.7
2003 4/.....	5.4	8.2	11.8	20.3	32.8	61.4	108.5	211.3	444.9	1,399.2
2002.....	5.4	7.5	11.7	20.5	34.3	62.5	112.9	217.8	454.8	1,414.6
2001.....	4.8	7.4	12.0	20.4	35.8	64.8	114.5	217.6	454.1	1,454.1
2000.....	5.3	7.0	11.9	20.2	36.3	65.6	113.5	223.4	460.2	1,468.0
1999.....	5.4	7.3	12.2	20.8	38.4	64.2	118.7	224.8	465.7	1,473.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other non-pulmonary forms of heart disease (I30-I51)

Black, both sexes

2006 4/.....	20.2	33.0	51.0	69.5	98.1	136.9	204.6	310.8	523.2	1,039.0
2005 4/.....	21.0	34.6	52.6	72.1	104.1	141.6	211.5	331.7	511.7	1,069.1
2004 4/.....	22.1	35.7	52.6	72.4	101.3	139.0	211.5	338.2	504.5	1,118.7
2003 4/.....	23.7	35.2	53.0	74.3	108.6	152.6	228.9	327.9	524.9	1,165.2
2002.....	22.7	37.2	53.5	74.3	106.2	146.8	225.5	343.1	545.3	1,201.8
2001.....	24.7	36.7	53.5	79.0	103.5	155.9	225.9	363.2	556.6	1,200.0
2000.....	23.6	37.5	52.7	77.3	106.4	160.6	237.6	367.7	543.8	1,229.4
1999.....	25.3	38.6	53.6	84.2	116.1	162.6	248.4	373.7	563.4	1,230.8

Black, male

2006 4/.....	26.3	44.9	70.6	93.7	134.8	178.8	267.6	382.1	607.2	1,044.6
2005 4/.....	27.9	45.2	70.7	98.9	135.8	183.0	275.8	408.1	566.0	1,020.7
2004 4/.....	28.4	49.5	67.9	97.6	137.3	184.9	259.6	427.7	591.1	1,160.3
2003 4/.....	30.0	45.2	71.4	102.0	143.8	200.4	283.5	397.1	617.9	1,144.8
2002.....	29.1	50.5	72.3	98.8	139.0	187.0	292.4	416.6	624.4	1,172.9
2001.....	29.9	48.0	68.5	105.3	135.3	191.8	290.2	456.7	688.6	1,253.8
2000.....	30.9	49.4	73.7	102.9	137.7	207.6	292.5	441.8	645.7	1,310.1
1999.....	32.3	51.1	74.0	116.6	153.8	213.9	314.8	445.9	684.5	1,382.7

Black, female

2006 4/.....	14.8	22.6	34.4	49.5	69.4	105.9	161.2	268.1	478.9	1,036.7
2005 4/.....	14.9	25.3	37.3	49.9	79.5	110.9	167.5	286.1	483.0	1,088.6
2004 4/.....	16.4	23.8	39.7	51.7	73.1	104.9	178.8	285.2	458.9	1,102.2
2003 4/.....	18.1	26.6	37.4	51.6	81.1	117.1	192.1	286.8	476.2	1,173.1
2002.....	17.0	25.6	37.7	54.1	80.5	116.9	180.6	299.1	504.1	1,212.9
2001.....	20.1	27.0	40.7	57.5	78.6	129.3	183.2	307.3	489.4	1,180.0
2000.....	17.2	27.1	34.9	56.3	81.7	126.0	201.1	323.0	492.5	1,199.3
1999.....	19.2	27.9	36.2	57.7	86.5	124.5	204.3	330.1	503.4	1,174.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and subacute endocarditis (I33)										
All races 3/, both sexes										
2006.....	0.2	0.4	0.5	0.7	0.8	1.0	1.4	2.2	2.7	2.2
2005.....	0.2	0.4	0.5	0.6	0.7	1.1	1.4	2.4	2.7	3.0
2004.....	0.2	0.4	0.5	0.5	0.7	1.0	1.5	2.1	2.9	3.2
2003.....	0.3	0.4	0.5	0.6	0.9	1.0	1.5	2.3	2.7	3.0
2002.....	0.3	0.4	0.5	0.5	0.8	1.1	1.5	2.2	2.7	2.6
2001.....	0.3	0.4	0.5	0.5	0.8	1.1	1.7	2.2	3.0	2.5
2000.....	0.3	0.4	0.6	0.7	0.7	1.1	1.3	2.2	2.8	2.8
1999.....	0.3	0.3	0.4	0.7	0.7	1.1	1.5	2.1	2.7	2.9
All races 3/, male										
2006.....	0.3	0.6	0.7	0.9	0.9	1.2	1.6	2.3	3.3	2.3
2005.....	0.2	0.4	0.7	0.8	0.9	1.2	1.8	3.1	3.5	5.1
2004.....	0.3	0.5	0.7	0.6	0.8	1.4	1.4	2.3	4.2	4.8
2003.....	0.4	0.6	0.7	0.8	1.1	1.3	1.6	2.9	3.7	4.1
2002.....	0.4	0.5	0.7	0.6	0.9	1.5	1.7	2.5	3.7	3.5
2001.....	0.3	0.7	0.7	0.6	1.1	1.1	2.1	3.1	4.3	3.8
2000.....	0.4	0.6	0.7	0.8	0.9	1.3	1.6	3.1	4.0	4.2
1999.....	0.4	0.5	0.7	1.0	0.8	1.2	1.7	2.8	3.9	4.7
All races 3/, female										
2006.....	0.2	0.3	0.3	0.4	0.7	0.7	1.2	2.2	2.3	2.1
2005.....	*	0.3	0.3	0.4	0.6	1.0	1.1	1.9	2.2	2.1
2004.....	*	0.3	0.4	0.3	0.6	0.7	1.7	2.0	2.1	2.5
2003.....	0.2	0.3	0.3	0.4	0.8	0.9	1.4	1.9	2.1	2.5
2002.....	0.2	0.3	0.3	0.4	0.8	0.8	1.3	1.9	2.1	2.2
2001.....	0.3	0.2	0.3	0.5	0.5	1.0	1.3	1.6	2.2	2.0
2000.....	0.2	0.3	0.4	0.5	0.6	0.9	1.1	1.6	2.0	2.3
1999.....	*	*	0.2	0.5	0.7	1.0	1.4	1.7	2.0	2.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute and subacute endocarditis (I33)										
White, both sexes										
2006 4/.....	0.2	0.4	0.4	0.5	0.7	0.9	1.4	2.1	2.8	2.1
2005 4/.....	0.1	0.3	0.3	0.4	0.7	1.0	1.5	2.4	2.6	3.1
2004 4/.....	0.2	0.3	0.4	0.4	0.6	1.0	1.5	2.0	2.9	3.2
2003 4/.....	0.3	0.3	0.4	0.5	0.8	1.0	1.4	2.3	2.8	3.1
2002.....	0.2	0.3	0.4	0.4	0.8	1.0	1.4	2.2	2.9	2.7
2001.....	0.2	0.4	0.4	0.4	0.7	0.9	1.5	2.0	3.0	2.6
2000.....	0.3	0.3	0.4	0.5	0.7	1.1	1.3	2.2	2.7	2.7
1999.....	0.2	0.3	0.3	0.5	0.6	1.0	1.4	2.2	2.8	3.0
White, male										
2006 4/.....	0.3	0.6	0.5	0.8	0.8	1.1	1.6	2.3	3.3	2.5
2005 4/.....	*	0.3	0.5	0.6	0.8	1.2	1.8	3.2	3.5	5.3
2004 4/.....	0.3	0.4	0.5	0.6	0.7	1.3	1.4	2.2	4.3	4.9
2003 4/.....	0.3	0.4	0.5	0.6	1.0	1.2	1.5	2.8	3.9	4.4
2002.....	0.3	0.4	0.6	0.4	0.9	1.3	1.7	2.7	3.9	3.6
2001.....	0.3	0.5	0.5	0.5	0.9	0.9	1.8	3.0	4.3	3.9
2000.....	0.4	0.5	0.6	0.7	0.9	1.2	1.6	3.2	4.0	4.2
1999.....	0.3	0.4	0.5	0.8	0.7	1.2	1.6	3.0	4.0	5.1
White, female										
2006 4/.....	*	0.2	0.2	0.3	0.5	0.8	1.1	2.0	2.5	1.9
2005 4/.....	*	0.2	*	0.3	0.5	0.9	1.2	1.7	2.0	2.1
2004 4/.....	*	*	0.2	*	0.5	0.6	1.6	1.8	2.1	2.4
2003 4/.....	*	0.2	0.2	0.4	0.6	0.8	1.3	1.8	2.1	2.6
2002.....	*	*	*	0.3	0.7	0.7	1.2	1.8	2.3	2.3
2001.....	*	*	*	0.4	0.4	0.9	1.3	1.3	2.2	2.1
2000.....	*	*	0.3	0.4	0.5	1.0	1.0	1.6	2.0	2.1
1999.....	*	*	*	*	0.6	0.8	1.3	1.6	2.0	2.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
All races 3/, both sexes										
2006.....	0.2	0.2	0.3	0.3	0.5	0.4	0.8	1.1	1.3	1.8
2005.....	0.2	0.3	0.3	0.3	0.4	0.7	0.8	1.1	1.3	1.9
2004.....	0.2	0.2	0.3	0.3	0.5	0.6	0.6	1.1	1.3	1.8
2003.....	0.2	0.3	0.3	0.4	0.4	0.8	0.8	0.9	1.5	1.6
2002.....	0.2	0.3	0.3	0.4	0.4	0.6	1.0	1.1	1.4	2.1
2001.....	0.2	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.1	1.3	1.8
1999.....	0.3	0.3	0.3	0.5	0.6	0.7	0.9	1.2	1.5	2.0
All races 3/, male										
2006.....	0.2	0.3	0.3	0.4	0.6	0.6	0.9	1.1	1.7	1.8
2005.....	0.3	0.4	0.4	0.3	0.5	0.7	1.0	1.4	1.5	1.9
2004.....	0.2	0.2	0.4	0.4	0.6	0.8	0.7	1.2	1.4	2.3
2003.....	0.3	0.4	0.3	0.4	0.4	1.0	1.0	1.1	1.5	1.5
2002.....	0.3	0.4	0.3	0.5	0.6	0.7	1.1	1.1	1.3	2.4
2001.....	*	0.3	0.3	0.4	0.6	0.6	0.8	1.3	1.1	1.8
2000.....	0.3	0.3	*	0.5	0.4	0.5	1.1	1.4	1.7	2.1
1999.....	0.3	0.3	0.3	0.6	0.6	0.8	0.9	1.3	1.5	2.5
All races 3/, female										
2006.....	*	*	0.2	0.2	0.4	*	0.7	1.1	1.0	1.8
2005.....	*	0.3	0.3	0.3	0.4	0.6	0.6	0.9	1.1	1.9
2004.....	*	0.2	0.2	0.3	0.4	0.5	0.6	1.1	1.3	1.6
2003.....	0.2	0.2	*	0.3	0.3	0.7	0.7	0.8	1.5	1.7
2002.....	*	*	0.2	0.4	*	0.5	1.0	1.1	1.4	2.0
2001.....	0.3	0.3	0.3	0.4	*	*	0.9	1.1	1.4	2.0
2000.....	*	*	0.2	*	0.4	0.5	0.7	0.9	1.0	1.7
1999.....	0.3	0.3	0.2	0.3	0.6	0.7	0.8	1.2	1.5	1.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of pericardium and acute myocarditis (I30-I31,I40)										
White, both sexes										
2006 4/.....	0.2	0.2	0.3	0.3	0.5	0.5	0.8	1.1	1.3	1.8
2005 4/.....	0.2	0.3	0.3	0.3	0.4	0.7	0.8	1.1	1.2	1.9
2004 4/.....	0.1	0.2	0.3	0.3	0.4	0.6	0.6	1.0	1.3	1.8
2003 4/.....	0.2	0.3	0.2	0.4	0.4	0.8	0.8	1.0	1.5	1.7
2002.....	0.2	0.2	0.2	0.4	0.4	0.6	1.0	1.0	1.3	2.2
2001.....	0.2	0.2	0.2	0.3	0.5	0.4	0.9	1.2	1.3	1.9
2000.....	0.2	0.2	0.2	0.3	0.4	0.5	0.9	1.0	1.4	1.8
1999.....	0.2	0.3	0.2	0.5	0.6	0.7	0.8	1.2	1.5	2.0
White, male										
2006 4/.....	*	0.3	0.3	0.4	0.7	0.6	0.9	1.2	1.8	1.8
2005 4/.....	0.2	0.3	0.4	0.3	0.4	0.7	1.0	1.3	1.5	1.9
2004 4/.....	*	*	0.3	0.4	0.6	0.6	0.7	1.2	1.4	2.2
2003 4/.....	0.2	0.3	0.3	0.4	0.5	1.0	1.1	1.1	1.6	1.7
2002.....	0.3	0.3	0.3	0.4	0.6	0.6	1.1	1.1	1.2	2.5
2001.....	*	0.2	0.3	0.4	0.6	0.5	0.9	1.3	1.1	2.0
2000.....	0.2	*	*	0.5	*	*	1.1	1.2	1.9	2.1
1999.....	0.3	0.3	*	0.6	0.6	0.7	0.9	1.3	1.5	2.5
White, female										
2006 4/.....	*	*	*	*	0.4	*	0.7	1.0	1.0	1.9
2005 4/.....	*	*	0.3	*	0.4	0.6	0.6	0.9	1.1	1.8
2004 4/.....	*	*	*	0.3	*	0.5	0.6	0.9	1.3	1.6
2003 4/.....	*	*	*	0.4	*	0.7	0.6	0.8	1.4	1.7
2002.....	*	*	*	0.4	*	0.5	0.9	1.0	1.4	2.0
2001.....	*	0.3	*	0.3	*	*	0.9	1.2	1.4	1.9
2000.....	*	*	*	*	*	0.6	0.7	0.9	1.1	1.6
1999.....	*	0.3	*	*	0.5	0.7	0.8	1.1	1.4	1.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of endocardium (I34-I38)										
All races 3/, both sexes										
2006.....	0.8	1.2	1.9	2.9	4.6	8.5	15.3	30.4	67.0	205.5
2005.....	0.8	1.4	2.0	2.9	4.7	8.0	14.5	31.2	67.0	204.7
2004.....	0.8	1.5	2.0	2.8	4.8	8.3	16.4	31.5	67.0	200.3
2003.....	0.9	1.2	1.9	3.1	4.5	9.1	15.9	32.4	68.7	201.9
2002.....	1.0	1.3	1.9	2.9	4.8	8.4	15.9	33.0	72.0	203.9
2001.....	0.8	1.2	2.1	2.8	4.7	9.0	17.3	33.9	70.3	204.2
2000.....	0.9	1.4	1.9	3.0	5.1	8.9	17.8	34.1	73.3	213.2
1999.....	1.1	1.3	2.0	2.9	5.1	9.2	17.6	35.2	76.6	209.8
All races 3/, male										
2006.....	1.0	1.6	2.2	3.8	5.2	9.7	16.9	31.7	74.4	199.6
2005.....	0.9	1.7	2.5	3.7	5.6	8.4	15.1	33.8	69.6	192.7
2004.....	1.0	2.0	2.4	3.6	5.2	9.1	17.6	34.1	70.3	194.2
2003.....	1.1	1.4	2.1	3.9	4.9	9.6	17.4	35.9	73.9	195.3
2002.....	1.1	1.7	2.3	3.3	5.2	8.7	17.3	36.4	74.9	198.5
2001.....	0.9	1.6	2.7	3.3	5.3	9.7	18.9	38.6	75.4	200.8
2000.....	1.0	1.9	2.2	3.8	5.9	9.3	19.2	35.1	78.9	215.8
1999.....	1.2	1.5	2.4	3.5	5.8	10.4	19.6	38.5	86.3	206.4
All races 3/, female										
2006.....	0.7	0.9	1.6	2.0	4.0	7.6	13.9	29.4	62.4	208.2
2005.....	0.6	1.1	1.6	2.2	4.0	7.7	14.0	29.3	65.4	210.2
2004.....	0.6	1.0	1.5	2.0	4.4	7.6	15.4	29.6	64.9	203.0
2003.....	0.7	1.0	1.7	2.4	4.1	8.6	14.7	29.8	65.6	204.8
2002.....	0.8	1.0	1.5	2.5	4.5	8.2	14.8	30.6	70.2	206.3
2001.....	0.6	0.9	1.5	2.4	4.2	8.4	16.1	30.6	67.2	205.6
2000.....	0.7	1.0	1.5	2.4	4.4	8.6	16.8	33.5	69.9	212.1
1999.....	0.9	1.2	1.7	2.4	4.4	8.2	16.0	32.9	70.9	211.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of endocardium (I34-I38)										
White, both sexes										
2006 4/.....	0.8	1.1	1.6	2.7	4.7	8.5	15.7	31.9	70.3	219.3
2005 4/.....	0.7	1.3	1.7	2.7	4.5	8.2	14.7	32.8	70.7	218.9
2004 4/.....	0.7	1.3	1.7	2.6	4.7	8.2	17.1	32.8	70.3	212.5
2003 4/.....	0.7	1.1	1.6	2.9	4.2	9.1	16.2	33.5	72.3	214.4
2002.....	0.9	1.1	1.7	2.7	4.8	8.2	16.4	34.7	75.8	216.2
2001.....	0.7	1.1	1.9	2.6	4.6	8.8	17.8	35.2	74.1	216.5
2000.....	0.8	1.3	1.7	2.7	5.0	8.9	18.1	35.1	77.0	225.8
1999.....	1.0	1.1	1.9	2.7	4.9	9.1	17.4	36.4	80.2	221.4
White, male										
2006 4/.....	1.0	1.4	1.9	3.6	5.1	9.8	17.3	33.2	78.5	213.0
2005 4/.....	1.0	1.6	2.2	3.3	5.4	8.8	15.4	35.5	73.6	206.9
2004 4/.....	0.9	1.7	2.1	3.4	5.1	9.2	18.5	35.2	73.7	205.2
2003 4/.....	0.9	1.2	1.8	3.5	4.7	9.7	17.9	37.3	77.6	207.8
2002.....	1.1	1.4	2.1	3.1	5.1	8.7	17.8	38.2	78.9	211.7
2001.....	0.8	1.4	2.5	3.1	5.2	9.5	19.4	39.9	79.8	212.0
2000.....	1.0	1.7	2.0	3.5	5.9	9.4	19.5	36.2	82.7	229.0
1999.....	1.1	1.1	2.3	3.3	5.6	10.5	19.4	39.9	90.2	218.2
White, female										
2006 4/.....	0.6	0.8	1.4	1.9	4.3	7.2	14.5	31.0	65.1	222.3
2005 4/.....	0.4	0.9	1.3	2.1	3.6	7.6	14.0	30.8	68.9	224.4
2004 4/.....	0.6	0.8	1.3	1.9	4.3	7.2	15.9	31.0	68.1	215.8
2003 4/.....	0.5	0.9	1.5	2.3	3.7	8.5	14.8	30.7	69.0	217.2
2002.....	0.6	0.7	1.3	2.3	4.4	7.8	15.2	32.1	73.9	218.2
2001.....	0.5	0.7	1.4	2.1	4.2	8.2	16.5	31.8	70.6	218.4
2000.....	0.6	0.8	1.4	2.0	4.2	8.4	17.0	34.3	73.6	224.5
1999.....	0.8	1.0	1.4	2.2	4.2	7.9	15.8	33.9	74.4	222.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of endocardium (I34-I38)										
Black, both sexes										
2006 4/.....	1.3	2.1	3.8	4.4	4.4	10.9	14.0	21.3	42.6	76.5
2005 4/.....	1.3	2.4	4.6	5.3	8.1	8.8	15.8	22.0	36.2	73.8
2004 4/.....	1.6	3.1	4.7	4.6	6.6	11.6	13.9	22.7	40.6	88.3
2003 4/.....	2.1	2.7	4.3	5.5	7.6	10.2	16.1	26.0	37.6	85.2
2002.....	1.8	3.6	3.8	4.8	6.7	12.1	13.8	22.2	37.2	86.4
2001.....	1.5	2.9	3.9	5.9	6.4	12.2	15.1	25.7	38.2	85.4
2000.....	1.9	3.0	3.6	5.9	7.2	10.9	18.9	29.9	39.5	92.9
1999.....	2.2	3.2	3.7	5.3	7.8	11.4	21.8	25.9	41.9	91.5
Black, male										
2006 4/.....	1.5	2.9	4.5	6.1	5.8	9.8	16.9	23.3	41.1	69.7
2005 4/.....	*	3.0	5.7	7.5	9.5	7.8	16.4	24.6	35.8	58.6
2004 4/.....	2.0	4.3	5.9	5.7	7.7	9.9	13.9	25.0	40.2	84.4
2003 4/.....	2.6	3.3	5.5	8.4	7.5	10.6	15.1	26.9	39.9	70.9
2002.....	1.5	4.5	4.0	6.1	7.1	10.7	14.0	24.3	35.2	67.5
2001.....	1.9	3.8	5.1	6.7	7.9	13.0	15.8	29.7	32.9	86.2
2000.....	1.8	4.1	4.5	6.6	7.5	10.4	19.7	29.8	43.0	86.5
1999.....	2.3	3.8	4.1	5.4	9.3	11.2	25.1	27.1	47.1	77.3
Black, female										
2006 4/.....	*	1.5	3.1	2.9	3.3	11.6	12.0	20.1	43.4	79.3
2005 4/.....	1.7	1.9	3.7	3.5	7.1	9.5	15.4	20.5	36.4	79.9
2004 4/.....	1.3	2.1	3.6	3.8	5.8	12.8	13.8	21.3	40.8	89.8
2003 4/.....	1.7	2.1	3.4	3.1	7.7	9.9	16.8	25.5	36.4	90.7
2002.....	2.0	2.8	3.6	3.8	6.3	13.1	13.7	20.9	38.2	93.6
2001.....	*	2.1	2.9	5.1	5.1	11.6	14.6	23.3	40.9	85.0
2000.....	1.9	2.0	2.8	5.3	7.0	11.3	18.4	29.9	37.8	95.3
1999.....	2.2	2.7	3.4	5.1	6.7	11.5	19.6	25.2	39.3	96.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nonrheumatic mitral valve disorders (I34)										
All races 3/, both sexes										
2006.....	0.2	0.3	0.3	0.4	0.7	1.4	2.3	3.9	8.1	19.8
2005.....	0.2	0.3	0.4	0.5	0.8	1.3	2.2	4.3	8.5	20.4
2004.....	0.2	0.3	0.3	0.4	0.7	1.3	2.5	4.7	8.4	20.1
2003.....	0.2	0.2	0.3	0.5	0.8	1.4	2.4	4.4	9.0	20.1
2002.....	0.2	0.3	0.3	0.5	1.0	1.5	2.8	5.0	9.8	22.1
2001.....	0.2	0.3	0.4	0.5	0.7	1.7	2.7	5.7	9.6	22.7
2000.....	0.3	0.3	0.3	0.5	1.0	1.6	3.0	5.8	10.6	24.3
1999.....	0.3	0.3	0.4	0.5	0.9	1.6	3.1	5.8	10.5	25.1
All races 3/, male										
2006.....	*	0.3	0.3	0.5	0.7	1.5	1.9	3.6	7.6	17.9
2005.....	0.2	0.2	0.4	0.5	0.7	0.9	1.8	4.0	7.4	17.6
2004.....	0.2	0.3	0.3	0.4	0.6	1.1	2.2	4.4	7.8	16.6
2003.....	0.2	0.2	0.3	0.5	0.8	1.3	2.4	4.5	8.2	16.4
2002.....	0.2	0.2	0.3	0.4	0.9	1.3	2.4	4.2	8.2	21.0
2001.....	0.2	0.3	0.4	0.5	0.7	1.7	2.8	4.7	8.1	19.0
2000.....	0.2	0.4	0.3	0.5	0.8	1.6	2.6	5.6	9.8	22.9
1999.....	0.3	0.3	0.3	0.4	0.9	1.3	3.1	5.6	10.6	22.2
All races 3/, female										
2006.....	0.2	0.3	0.4	0.4	0.7	1.3	2.6	4.2	8.5	20.6
2005.....	0.2	0.4	0.4	0.5	0.8	1.6	2.4	4.5	9.2	21.7
2004.....	0.3	0.3	0.3	0.4	0.8	1.4	2.7	5.0	8.7	21.7
2003.....	0.2	0.3	0.4	0.5	0.9	1.5	2.3	4.4	9.5	21.7
2002.....	0.2	0.3	0.3	0.5	1.1	1.8	3.1	5.6	10.9	22.6
2001.....	0.3	0.3	0.4	0.5	0.8	1.7	2.7	6.3	10.5	24.3
2000.....	0.3	0.2	0.4	0.6	1.1	1.5	3.2	5.9	11.1	24.9
1999.....	0.4	0.4	0.5	0.6	0.9	1.8	3.1	5.9	10.5	26.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nonrheumatic mitral valve disorders (I34)										
White, both sexes										
2006 4/.....	0.2	0.3	0.3	0.4	0.7	1.4	2.4	4.2	8.6	21.1
2005 4/.....	0.2	0.3	0.3	0.5	0.7	1.3	2.2	4.5	8.9	21.7
2004 4/.....	0.2	0.3	0.3	0.4	0.6	1.3	2.5	4.9	8.7	21.1
2003 4/.....	0.2	0.2	0.3	0.5	0.8	1.4	2.5	4.6	9.3	21.4
2002.....	0.2	0.2	0.3	0.4	1.0	1.5	2.9	5.2	10.4	23.8
2001.....	0.2	0.3	0.4	0.5	0.7	1.6	2.8	6.0	10.1	24.2
2000.....	0.3	0.3	0.3	0.5	1.0	1.6	3.0	6.0	11.3	25.5
1999.....	0.3	0.3	0.3	0.5	0.9	1.5	3.1	5.9	11.1	26.5
White, male										
2006 4/.....	*	0.3	0.3	0.5	0.6	1.5	1.9	3.9	8.2	19.3
2005 4/.....	*	*	0.3	0.5	0.7	1.0	1.9	4.1	7.9	19.1
2004 4/.....	0.2	0.3	0.3	0.4	0.5	1.2	2.3	4.4	8.0	17.5
2003 4/.....	*	*	0.3	0.5	0.8	1.3	2.6	4.6	8.5	17.6
2002.....	0.2	0.2	*	0.4	0.9	1.3	2.4	4.3	8.7	22.6
2001.....	*	0.3	0.4	0.6	0.7	1.6	2.8	5.1	8.7	20.5
2000.....	*	0.4	*	0.4	0.9	1.7	2.7	5.8	10.5	24.3
1999.....	0.3	0.2	0.3	0.4	0.9	1.3	3.0	5.7	11.4	23.7
White, female										
2006 4/.....	0.2	0.2	0.3	0.4	0.7	1.3	2.8	4.5	8.9	22.0
2005 4/.....	*	0.3	0.3	0.5	0.7	1.6	2.4	4.8	9.6	22.9
2004 4/.....	0.3	0.2	*	0.4	0.7	1.4	2.7	5.2	9.2	22.7
2003 4/.....	0.2	0.3	0.3	0.5	0.7	1.5	2.5	4.6	9.9	23.0
2002.....	*	0.3	0.3	0.5	1.2	1.7	3.3	5.8	11.4	24.3
2001.....	0.3	0.3	0.3	0.4	0.8	1.6	2.8	6.7	11.0	25.7
2000.....	0.3	*	0.3	0.6	1.1	1.5	3.3	6.1	11.8	25.9
1999.....	0.4	0.3	0.4	0.6	0.9	1.7	3.1	6.0	10.9	27.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nonrheumatic aortic valve disorders (I35)										
All races 3/, both sexes										
2006.....	0.3	0.4	0.7	1.1	2.1	4.2	8.4	18.0	43.6	149.5
2005.....	0.2	0.4	0.7	1.2	2.2	3.7	7.8	17.8	43.0	146.9
2004.....	0.3	0.5	0.8	1.2	2.2	3.8	8.9	18.4	42.7	142.5
2003.....	0.3	0.4	0.6	1.3	1.9	4.5	8.5	18.4	44.1	144.9
2002.....	0.3	0.5	0.8	1.2	2.2	4.0	8.8	19.2	45.4	144.8
2001.....	0.2	0.5	0.9	1.2	2.3	4.5	9.2	19.9	44.6	144.1
2000.....	0.3	0.5	0.8	1.4	2.3	4.5	9.9	20.0	47.1	151.3
1999.....	0.3	0.5	0.8	1.3	2.3	4.8	9.3	20.7	49.9	147.8
All races 3/, male										
2006.....	0.4	0.5	0.9	1.6	2.5	5.1	10.4	19.9	51.1	149.9
2005.....	0.4	0.6	0.9	1.7	2.9	4.6	8.9	20.8	47.3	142.5
2004.....	0.4	0.7	1.0	1.7	2.7	4.6	10.3	21.0	46.0	141.9
2003.....	0.4	0.6	0.8	1.8	2.4	5.5	9.9	21.9	50.4	144.7
2002.....	0.4	0.7	1.0	1.5	2.6	4.5	10.8	23.2	51.4	144.9
2001.....	0.3	0.8	1.3	1.5	2.9	5.3	11.0	24.7	50.7	145.9
2000.....	0.4	0.8	1.2	2.1	3.1	5.2	11.7	21.5	52.9	158.2
1999.....	0.4	0.7	1.2	1.7	3.0	5.9	11.7	24.3	58.9	150.3
All races 3/, female										
2006.....	0.2	0.3	0.4	0.7	1.8	3.5	6.7	16.7	38.9	149.3
2005.....	*	0.3	0.4	0.8	1.5	3.0	7.0	15.6	40.4	148.9
2004.....	*	0.2	0.5	0.8	1.8	3.0	7.8	16.5	40.6	142.7
2003.....	*	0.2	0.5	0.8	1.4	3.6	7.4	15.9	40.2	145.0
2002.....	0.2	0.2	0.6	0.8	1.9	3.5	7.1	16.4	41.7	144.8
2001.....	*	0.3	0.5	0.9	1.8	3.8	7.8	16.6	40.9	143.4
2000.....	*	0.2	0.4	0.8	1.6	3.8	8.5	19.0	43.7	148.5
1999.....	0.2	0.4	0.5	0.9	1.7	3.8	7.4	18.1	44.7	146.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nonrheumatic aortic valve disorders (I35)										
White, both sexes										
2006 4/.....	0.3	0.4	0.7	1.2	2.3	4.4	8.8	19.0	45.9	159.6
2005 4/.....	0.2	0.4	0.6	1.3	2.2	3.8	8.1	19.0	45.7	157.2
2004 4/.....	0.3	0.5	0.7	1.3	2.3	3.9	9.5	19.1	45.0	151.6
2003 4/.....	0.2	0.4	0.6	1.3	1.9	4.6	8.8	19.1	46.6	154.1
2002.....	0.3	0.4	0.7	1.2	2.3	3.9	9.1	20.3	47.9	153.2
2001.....	0.2	0.5	0.9	1.2	2.4	4.6	9.6	20.5	47.3	152.8
2000.....	0.3	0.5	0.8	1.4	2.3	4.4	10.2	20.9	49.5	160.7
1999.....	0.3	0.4	0.8	1.3	2.3	4.8	9.2	21.7	52.3	156.1
White, male										
2006 4/.....	0.5	0.5	0.9	1.7	2.5	5.4	10.9	20.8	53.9	159.8
2005 4/.....	0.4	0.6	0.9	1.7	2.9	4.8	9.2	22.1	50.2	152.7
2004 4/.....	0.4	0.7	0.9	1.7	2.7	4.8	10.9	21.7	48.5	150.1
2003 4/.....	0.3	0.5	0.7	1.7	2.5	5.8	10.3	22.7	53.4	154.4
2002.....	0.4	0.6	1.0	1.5	2.7	4.6	11.3	24.5	54.3	154.4
2001.....	0.3	0.7	1.2	1.5	3.0	5.4	11.4	25.2	53.8	153.5
2000.....	0.4	0.7	1.2	2.0	3.2	5.2	12.0	22.6	55.4	168.4
1999.....	0.4	0.6	1.1	1.7	3.0	6.1	11.5	25.6	61.2	159.0
White, female										
2006 4/.....	*	0.3	0.5	0.7	2.0	3.6	7.1	17.7	40.9	159.5
2005 4/.....	*	0.2	0.3	0.9	1.5	2.9	7.1	16.7	42.8	159.3
2004 4/.....	*	0.2	0.5	0.9	1.9	3.0	8.3	17.2	42.8	152.2
2003 4/.....	*	0.2	0.5	0.9	1.4	3.7	7.6	16.5	42.3	154.0
2002.....	*	*	0.5	0.9	1.9	3.4	7.4	17.3	44.0	152.7
2001.....	*	0.3	0.5	0.8	2.0	3.9	8.2	17.2	43.3	152.5
2000.....	*	*	0.5	0.7	1.5	3.8	8.7	19.7	46.1	157.6
1999.....	0.2	0.3	0.4	0.8	1.8	3.7	7.4	19.0	47.0	155.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nonrheumatic aortic valve disorders (I35)										
Black, both sexes										
2006 4/.....	*	*	1.0	*	*	3.6	6.4	10.9	27.5	56.0
2005 4/.....	*	*	1.2	1.4	2.7	3.5	7.3	10.4	20.9	53.5
2004 4/.....	*	*	1.3	1.4	2.3	4.0	5.8	13.2	23.3	59.2
2003 4/.....	*	*	0.9	1.7	1.8	3.3	7.8	14.1	21.1	58.6
2002.....	*	0.9	1.1	*	2.2	5.1	6.8	11.5	23.3	63.0
2001.....	*	*	1.3	1.7	2.0	4.4	6.9	16.3	21.4	59.6
2000.....	*	0.8	*	2.1	2.6	5.5	9.1	13.9	25.4	62.3
1999.....	*	1.0	1.4	1.7	2.8	5.0	10.7	12.0	26.9	64.4
Black, male										
2006 4/.....	*	*	*	*	*	*	8.6	13.9	30.1	50.9
2005 4/.....	*	*	*	2.4	4.2	*	7.9	13.0	22.5	47.4
2004 4/.....	*	*	*	*	4.1	*	6.5	14.1	23.7	58.6
2003 4/.....	*	*	*	2.8	*	*	7.9	16.4	19.9	48.0
2002.....	*	*	*	*	*	5.2	7.7	13.3	22.2	52.5
2001.....	*	*	2.3	*	*	*	8.1	22.8	19.3	68.1
2000.....	*	*	*	*	*	6.2	10.7	13.2	30.3	57.7
1999.....	*	*	*	*	*	5.7	13.9	13.3	36.6	58.6
Black, female										
2006 4/.....	*	*	*	*	*	3.6	4.9	9.1	26.0	58.1
2005 4/.....	*	*	*	*	*	*	6.9	8.8	20.0	55.9
2004 4/.....	*	*	*	*	*	4.1	5.3	12.7	23.1	59.5
2003 4/.....	*	*	*	*	*	*	7.8	12.8	21.7	62.8
2002.....	*	*	*	*	*	5.0	6.3	10.4	23.9	66.9
2001.....	*	*	*	*	*	4.1	6.1	12.5	22.5	56.4
2000.....	*	*	*	*	*	5.0	8.0	14.2	22.9	64.0
1999.....	*	*	*	*	*	4.4	8.6	11.2	22.0	66.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of endocardium (I36-I38)										
All races 3/, both sexes										
2006.....	0.4	0.5	0.8	1.3	1.7	2.9	4.6	8.4	15.3	36.2
2005.....	0.4	0.7	0.9	1.2	1.8	3.0	4.5	9.1	15.5	37.4
2004.....	0.3	0.7	0.9	1.1	1.9	3.3	5.0	8.4	15.9	37.7
2003.....	0.4	0.6	0.9	1.3	1.8	3.2	5.0	9.5	15.7	36.9
2002.....	0.4	0.6	0.8	1.2	1.6	2.9	4.4	8.7	16.7	37.0
2001.....	0.3	0.4	0.8	1.1	1.6	2.8	5.3	8.3	16.1	37.4
2000.....	0.3	0.6	0.8	1.1	1.8	2.9	5.0	8.3	15.6	37.6
1999.....	0.4	0.5	0.8	1.1	1.8	2.9	5.2	8.7	16.1	37.0
All races 3/, male										
2006.....	0.4	0.7	1.0	1.7	2.0	3.1	4.6	8.2	15.7	31.8
2005.....	0.4	0.9	1.1	1.5	2.0	2.9	4.4	9.0	14.9	32.6
2004.....	0.4	1.0	1.1	1.4	2.0	3.4	5.1	8.7	16.5	35.7
2003.....	0.5	0.7	1.0	1.6	1.7	2.9	5.0	9.5	15.4	34.2
2002.....	0.5	0.8	1.0	1.3	1.7	2.9	4.1	9.1	15.3	32.6
2001.....	0.4	0.5	1.0	1.3	1.7	2.7	5.1	9.2	16.5	35.9
2000.....	0.4	0.7	0.8	1.2	2.0	2.5	4.9	8.0	16.2	34.7
1999.....	0.5	0.5	0.9	1.3	1.9	3.1	4.9	8.6	16.8	33.9
All races 3/, female										
2006.....	0.3	0.3	0.7	0.9	1.5	2.7	4.6	8.6	15.1	38.3
2005.....	0.3	0.4	0.8	0.9	1.6	3.2	4.6	9.2	15.9	39.7
2004.....	0.2	0.5	0.7	0.8	1.8	3.2	5.0	8.1	15.6	38.5
2003.....	0.3	0.5	0.8	1.0	1.8	3.5	5.0	9.5	15.9	38.1
2002.....	0.4	0.5	0.6	1.1	1.5	3.0	4.6	8.5	17.6	38.9
2001.....	0.2	0.3	0.6	1.0	1.6	2.8	5.5	7.7	15.8	38.0
2000.....	0.2	0.5	0.7	0.9	1.7	3.2	5.0	8.6	15.2	38.7
1999.....	0.3	0.5	0.7	0.9	1.7	2.7	5.5	8.8	15.7	38.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of endocardium (I36-I38)										
White, both sexes										
2006 4/.....	0.3	0.5	0.7	1.1	1.7	2.6	4.6	8.7	15.8	38.6
2005 4/.....	0.3	0.6	0.8	1.0	1.6	3.0	4.4	9.3	16.1	40.0
2004 4/.....	0.2	0.5	0.7	0.9	1.8	3.0	5.1	8.8	16.5	39.9
2003 4/.....	0.3	0.4	0.7	1.1	1.5	3.0	4.9	9.8	16.4	38.9
2002.....	0.3	0.4	0.7	1.0	1.4	2.8	4.3	9.2	17.5	39.2
2001.....	0.3	0.3	0.7	0.9	1.5	2.6	5.4	8.6	16.7	39.5
2000.....	0.2	0.5	0.6	0.8	1.7	2.9	4.9	8.2	16.2	39.6
1999.....	0.3	0.3	0.7	0.9	1.6	2.8	5.1	8.8	16.8	38.8
White, male										
2006 4/.....	0.4	0.6	0.7	1.5	2.0	3.0	4.5	8.5	16.4	33.9
2005 4/.....	0.4	0.8	0.9	1.2	1.8	3.0	4.3	9.2	15.4	35.0
2004 4/.....	0.3	0.7	0.9	1.3	1.9	3.3	5.3	9.0	17.1	37.6
2003 4/.....	0.4	0.5	0.8	1.3	1.4	2.7	5.0	9.9	15.7	35.8
2002.....	0.4	0.5	0.9	1.1	1.5	2.9	4.1	9.4	15.9	34.8
2001.....	0.3	0.3	0.9	1.0	1.5	2.5	5.2	9.7	17.3	38.0
2000.....	0.4	0.6	0.6	1.0	1.8	2.5	4.8	7.8	16.9	36.4
1999.....	0.4	0.3	0.8	1.1	1.7	3.0	4.8	8.6	17.6	35.5
White, female										
2006 4/.....	0.2	0.3	0.6	0.8	1.5	2.3	4.6	8.8	15.4	40.8
2005 4/.....	*	0.4	0.6	0.7	1.4	3.1	4.5	9.3	16.5	42.3
2004 4/.....	*	0.4	0.5	0.6	1.6	2.8	4.9	8.6	16.1	40.9
2003 4/.....	*	0.4	0.7	0.9	1.6	3.3	4.7	9.6	16.8	40.2
2002.....	0.2	0.3	0.4	1.0	1.3	2.7	4.6	9.0	18.5	41.1
2001.....	*	*	0.5	0.8	1.4	2.7	5.5	7.9	16.3	40.2
2000.....	*	0.4	0.6	0.7	1.6	3.2	4.9	8.4	15.7	40.9
1999.....	0.2	0.3	0.6	0.8	1.6	2.5	5.3	8.9	16.4	40.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of endocardium (I36-I38)

Black, both sexes

2006 4/.....	0.8	1.1	2.2	2.8	1.9	5.9	6.3	8.5	11.4	14.5
2005 4/.....	0.9	1.3	2.4	3.1	3.9	3.7	6.6	9.2	9.7	12.3
2004 4/.....	0.9	2.1	2.9	2.6	2.9	5.9	5.9	5.8	12.1	19.2
2003 4/.....	1.2	1.7	2.7	3.0	4.3	5.3	6.9	8.5	11.2	18.7
2002.....	1.3	2.2	2.1	2.8	3.6	4.9	4.8	6.9	9.7	16.6
2001.....	*	1.6	1.9	3.3	3.5	5.3	5.4	6.9	11.6	18.1
2000.....	1.0	1.5	2.1	2.9	3.6	3.8	7.2	11.0	10.4	18.5
1999.....	1.3	1.6	1.7	2.8	3.7	4.5	7.7	9.5	10.7	16.9

Black, male

2006 4/.....	*	1.6	2.5	3.9	*	4.7	6.8	*	*	*
2005 4/.....	*	1.8	3.1	4.4	4.2	*	7.9	*	*	*
2004 4/.....	*	3.1	3.8	2.9	*	4.8	*	*	*	*
2003 4/.....	*	2.0	3.6	4.6	4.2	5.4	*	*	15.5	*
2002.....	*	3.0	*	3.7	*	*	*	*	*	*
2001.....	*	1.9	2.6	4.4	4.3	5.8	*	*	*	*
2000.....	*	1.9	2.4	*	*	*	7.0	12.8	*	*
1999.....	*	*	*	*	4.9	*	*	10.0	*	*

Black, female

2006 4/.....	*	*	1.9	1.9	*	6.8	5.9	8.6	13.0	13.9
2005 4/.....	1.3	*	1.8	2.0	3.7	4.2	5.6	9.6	9.6	13.5
2004 4/.....	*	*	2.2	2.3	3.4	6.7	6.2	*	12.1	19.3
2003 4/.....	*	1.4	2.0	*	4.4	5.3	7.8	9.2	8.9	19.0
2002.....	1.4	1.6	2.2	*	3.6	5.4	5.4	6.3	9.5	19.3
2001.....	*	*	*	*	*	4.9	5.6	7.8	11.9	18.9
2000.....	*	*	*	2.7	3.3	4.0	7.3	10.0	11.0	17.6
1999.....	*	*	*	*	*	4.6	8.6	9.2	11.2	18.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cardiomyopathy (I42)										
All races 3/, both sexes										
2006.....	3.6	5.1	6.5	8.6	11.2	16.3	25.1	38.7	62.8	122.0
2005.....	3.7	5.4	6.7	8.8	12.4	17.4	26.5	42.2	65.7	126.7
2004.....	3.5	5.3	6.9	9.0	13.0	18.3	28.0	43.8	68.9	127.6
2003.....	4.0	5.6	7.0	9.7	13.9	20.8	30.7	45.7	72.5	137.8
2002.....	4.1	5.9	7.2	10.5	13.8	21.1	33.6	49.5	77.3	142.0
2001.....	4.0	5.5	7.3	10.6	14.5	22.9	31.9	48.9	76.8	138.3
2000.....	4.1	5.6	8.0	10.8	15.8	24.1	33.7	51.5	76.2	142.3
1999.....	4.2	5.7	7.9	11.7	16.8	25.6	35.5	50.0	76.1	139.2
All races 3/, male										
2006.....	5.1	7.3	9.1	12.4	15.7	23.0	35.5	54.5	90.8	159.6
2005.....	5.5	7.8	9.9	12.6	16.9	24.7	37.1	58.9	97.1	169.6
2004.....	5.0	7.7	9.7	12.5	18.5	25.4	39.2	62.4	96.3	174.6
2003.....	5.7	8.0	10.0	13.5	19.6	28.6	42.9	65.5	104.5	178.8
2002.....	5.9	8.6	10.4	14.7	19.4	28.9	47.9	71.8	110.8	189.7
2001.....	5.8	7.9	10.0	14.8	20.5	31.0	44.7	69.9	108.8	184.7
2000.....	5.9	7.9	11.5	15.5	21.6	33.1	47.9	74.4	109.5	195.3
1999.....	6.1	8.2	11.3	16.8	23.0	36.1	50.7	70.9	106.8	188.2
All races 3/, female										
2006.....	2.2	3.0	3.9	5.1	7.1	10.4	16.6	27.1	45.3	104.4
2005.....	1.8	3.0	3.6	5.3	8.3	11.0	17.8	30.1	46.3	107.0
2004.....	2.0	2.9	4.3	5.7	8.0	12.2	19.0	30.5	52.1	106.4
2003.....	2.3	3.3	4.2	6.1	8.7	14.1	20.9	31.6	52.8	119.7
2002.....	2.4	3.2	4.1	6.5	8.7	14.3	22.1	33.7	56.9	121.3
2001.....	2.3	3.2	4.7	6.7	9.0	15.9	21.8	34.3	57.6	118.9
2000.....	2.2	3.4	4.7	6.4	10.5	16.4	22.5	35.6	56.5	120.7
1999.....	2.3	3.4	4.7	6.9	11.2	16.6	23.6	35.5	58.2	119.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cardiomyopathy (I42)										
White, both sexes										
2006 4/.....	3.0	4.2	5.2	7.1	9.6	14.7	23.2	37.3	61.9	123.3
2005 4/.....	3.0	4.3	5.3	7.4	10.7	15.4	24.9	41.1	64.9	128.5
2004 4/.....	2.8	4.1	5.6	7.7	11.3	16.3	26.2	42.2	68.7	129.3
2003 4/.....	3.3	4.6	5.6	8.1	11.9	18.8	28.8	44.3	71.3	139.8
2002.....	3.3	4.7	5.8	8.9	12.2	19.5	31.7	47.7	76.6	141.7
2001.....	3.0	4.2	5.9	8.8	12.7	21.0	30.3	46.9	75.3	139.2
2000.....	3.2	4.2	6.4	9.0	14.0	21.6	31.7	49.1	74.7	142.4
1999.....	3.1	4.4	6.3	9.6	14.2	23.4	33.1	48.0	74.3	138.6
White, male										
2006 4/.....	4.2	6.0	7.4	10.2	13.6	20.8	32.8	53.4	90.6	162.1
2005 4/.....	4.7	6.3	7.9	10.6	14.6	21.8	34.5	57.9	97.0	175.0
2004 4/.....	4.0	6.2	8.1	10.6	16.1	22.6	37.3	60.8	96.8	178.3
2003 4/.....	4.8	6.7	8.0	11.4	16.9	25.8	40.6	64.2	103.5	182.6
2002.....	4.8	7.0	8.5	12.6	17.4	26.7	45.3	70.3	110.8	190.9
2001.....	4.5	6.2	8.3	12.4	18.2	28.5	42.7	67.7	107.1	187.0
2000.....	4.6	6.3	9.2	13.0	19.4	29.5	45.4	71.2	108.2	197.5
1999.....	4.7	6.4	9.1	13.8	19.4	33.2	47.2	68.7	105.1	188.0
White, female										
2006 4/.....	1.8	2.5	3.1	4.2	5.8	9.2	15.3	25.3	43.8	105.0
2005 4/.....	1.3	2.2	2.7	4.2	7.0	9.7	17.0	28.6	44.8	107.1
2004 4/.....	1.5	2.1	3.2	4.9	6.8	10.7	17.0	28.6	51.2	107.2
2003 4/.....	1.7	2.5	3.2	4.9	7.2	12.7	19.2	30.0	51.4	120.8
2002.....	1.8	2.4	3.2	5.3	7.4	13.1	20.7	31.6	55.6	120.4
2001.....	1.5	2.2	3.5	5.4	7.6	14.3	20.2	32.2	56.1	119.2
2000.....	1.7	2.3	3.6	5.2	9.0	14.6	20.6	33.5	54.8	120.0
1999.....	1.5	2.4	3.6	5.6	9.4	14.9	21.9	33.4	56.1	118.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cardiomyopathy (I42)										
Black, both sexes										
2006 4/.....	8.3	12.2	17.3	21.9	28.2	33.3	46.1	59.8	86.9	125.8
2005 4/.....	8.8	14.4	18.1	22.3	29.3	38.3	45.7	66.8	89.1	125.9
2004 4/.....	9.1	13.8	18.2	23.1	31.7	40.0	51.2	69.1	85.4	124.4
2003 4/.....	9.8	14.1	19.0	25.7	34.1	42.5	55.0	68.6	99.5	135.7
2002.....	10.4	15.2	18.4	27.0	30.4	39.5	57.5	77.3	96.3	161.0
2001.....	11.8	15.5	19.8	27.8	33.3	44.7	54.5	80.9	104.5	141.2
2000.....	10.7	16.6	22.7	28.6	32.2	51.2	59.6	85.5	104.8	151.1
1999.....	12.1	16.6	22.7	30.9	40.5	49.4	64.3	79.1	103.7	153.0
Black, male										
2006 4/.....	12.1	18.2	25.1	32.9	40.3	48.0	68.0	78.8	119.1	155.5
2005 4/.....	12.9	20.5	27.8	32.4	41.4	57.0	70.2	92.1	126.4	145.9
2004 4/.....	13.1	20.6	24.3	33.8	46.5	58.1	69.3	96.0	112.6	158.0
2003 4/.....	13.5	19.8	27.0	36.3	49.0	62.8	77.5	94.5	141.0	172.0
2002.....	14.8	22.4	27.4	38.6	42.0	55.8	85.5	103.8	130.7	204.8
2001.....	16.3	21.4	26.6	40.0	47.6	62.5	73.4	112.9	153.3	180.4
2000.....	16.2	22.0	33.1	42.8	44.5	73.6	82.2	123.3	149.2	193.7
1999.....	16.8	23.5	33.6	47.1	58.1	70.6	96.3	105.0	140.4	215.6
Black, female										
2006 4/.....	5.0	6.9	10.6	12.8	18.7	22.5	31.1	48.5	69.8	113.7
2005 4/.....	5.1	9.0	9.9	14.0	19.8	24.3	29.0	51.7	69.4	117.8
2004 4/.....	5.6	7.8	13.1	14.2	20.1	26.4	38.9	53.1	71.1	111.0
2003 4/.....	6.5	9.2	12.1	17.0	22.6	27.4	39.9	53.2	77.8	121.5
2002.....	6.5	9.1	10.7	17.5	21.4	27.4	38.8	61.5	78.4	144.2
2001.....	7.8	10.3	14.0	17.8	22.1	31.5	41.9	61.8	79.7	126.7
2000.....	5.8	12.0	13.9	17.0	22.6	34.6	44.6	62.7	82.4	135.2
1999.....	7.9	10.7	13.4	17.7	26.6	33.7	43.0	63.5	85.6	129.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Conduction disorders and cardiac dysrhythmias (I44-I49)										
All races 3/, both sexes										
2006.....	2.1	4.0	6.0	9.2	15.0	21.8	35.1	61.2	109.9	274.0
2005.....	2.4	3.8	6.1	9.5	15.1	21.9	37.2	61.0	111.3	297.0
2004.....	2.4	3.9	6.2	9.1	15.6	22.4	36.3	62.1	114.4	295.3
2003.....	2.4	4.0	6.1	10.2	15.4	24.3	39.4	66.7	122.1	313.7
2002.....	2.6	3.6	6.4	10.1	15.8	24.7	39.0	70.1	126.0	323.2
2001.....	2.2	3.9	6.3	10.0	15.9	25.3	40.5	69.3	125.7	328.2
2000.....	2.5	3.8	5.9	10.5	16.2	26.1	41.0	68.8	127.3	340.2
1999.....	2.5	4.1	6.5	10.8	19.1	27.3	44.8	72.4	134.3	353.7
All races 3/, male										
2006.....	2.6	5.0	8.0	12.0	19.7	27.8	41.8	70.3	120.5	257.5
2005.....	2.8	4.9	8.2	13.0	20.0	28.0	46.8	69.1	118.0	284.0
2004.....	3.1	5.0	8.1	12.4	21.2	28.9	45.1	71.8	126.1	278.5
2003.....	3.1	5.1	8.4	13.8	20.4	31.5	50.2	77.3	134.0	292.7
2002.....	3.4	4.8	8.7	13.5	21.2	31.1	48.1	81.3	138.2	310.5
2001.....	2.9	5.1	8.5	13.1	20.9	31.0	49.7	82.2	140.8	311.6
2000.....	3.0	5.1	8.2	14.3	21.7	34.5	52.3	80.6	137.4	333.2
1999.....	3.1	5.7	8.7	14.7	24.9	36.8	55.1	84.9	154.2	353.4
All races 3/, female										
2006.....	1.6	3.0	4.0	6.5	10.7	16.6	29.7	54.6	103.3	281.6
2005.....	2.0	2.8	4.2	6.1	10.6	16.6	29.4	55.1	107.2	303.0
2004.....	1.8	2.8	4.3	6.1	10.5	16.7	29.3	55.2	107.3	302.9
2003.....	1.8	3.0	4.0	6.9	10.9	18.0	30.6	59.1	114.7	323.0
2002.....	1.8	2.4	4.3	6.9	10.9	19.1	31.8	62.2	118.6	328.8
2001.....	1.5	2.7	4.3	7.0	11.4	20.4	33.3	60.3	116.7	335.2
2000.....	1.9	2.5	3.7	7.0	11.2	18.8	32.2	60.6	121.4	343.0
1999.....	1.8	2.5	4.4	7.2	13.8	19.2	36.7	63.7	122.7	353.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Conduction disorders and cardiac dysrhythmias (I44-I49)										
White, both sexes										
2006 4/.....	1.8	3.7	5.3	8.3	13.8	20.1	33.0	59.6	108.1	277.9
2005 4/.....	2.2	3.4	5.4	8.5	13.8	20.1	35.1	59.1	111.3	299.8
2004 4/.....	2.1	3.4	5.6	8.0	14.6	21.1	34.5	59.8	113.5	295.0
2003 4/.....	2.2	3.6	5.4	9.4	14.3	22.2	37.4	65.0	120.9	314.5
2002.....	2.4	3.2	5.7	9.1	14.1	23.4	37.1	68.0	124.4	324.4
2001.....	1.9	3.5	5.6	8.9	14.7	23.7	38.6	66.9	123.8	329.7
2000.....	2.2	3.3	5.4	9.8	14.7	24.7	39.4	66.6	127.1	341.6
1999.....	2.3	3.6	5.9	9.8	17.9	26.0	43.0	70.0	133.1	355.9
White, male										
2006 4/.....	2.2	4.6	7.3	10.9	18.0	25.5	39.5	67.9	118.6	259.3
2005 4/.....	2.6	4.6	7.6	11.9	18.7	25.7	44.0	66.9	117.6	283.8
2004 4/.....	2.8	4.5	7.6	11.1	20.1	27.2	42.8	68.9	124.4	273.7
2003 4/.....	2.9	4.7	7.5	12.7	19.0	29.1	47.7	74.8	132.6	291.3
2002.....	3.4	4.4	7.8	12.4	19.3	29.9	45.7	78.2	136.9	308.9
2001.....	2.7	4.7	7.7	11.6	19.4	29.7	46.9	78.8	137.5	310.2
2000.....	2.8	4.6	7.4	13.3	20.0	32.7	50.4	78.4	138.4	332.8
1999.....	3.0	5.3	8.0	13.3	23.6	35.1	53.3	81.3	152.3	350.9
White, female										
2006 4/.....	1.3	2.7	3.4	5.8	9.8	15.3	27.7	53.5	101.5	286.6
2005 4/.....	1.8	2.3	3.3	5.3	9.4	15.2	27.7	53.2	107.4	307.1
2004 4/.....	1.4	2.4	3.6	5.0	9.5	15.7	27.8	53.1	106.7	304.7
2003 4/.....	1.5	2.5	3.3	6.1	9.9	16.1	28.9	58.0	113.7	324.7
2002.....	1.5	2.0	3.6	6.1	9.4	17.7	30.0	60.6	116.8	331.1
2001.....	1.2	2.2	3.6	6.2	10.4	18.4	31.8	58.5	115.5	337.9
2000.....	1.6	2.0	3.3	6.4	9.9	17.6	30.5	58.4	120.4	345.2
1999.....	1.6	2.0	3.9	6.5	12.6	18.1	34.9	62.1	121.8	357.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Conduction disorders and cardiac dysrhythmias (I44-I49)										
Black, both sexes										
2006 4/.....	4.6	7.2	12.3	18.2	30.0	40.0	61.8	91.3	149.3	275.3
2005 4/.....	4.3	7.4	12.7	19.4	28.9	39.7	64.4	89.0	128.9	303.9
2004 4/.....	5.2	7.8	12.4	20.6	28.4	37.5	61.8	96.0	140.6	315.6
2003 4/.....	4.8	8.2	13.3	19.6	28.8	46.5	66.2	95.0	145.2	327.3
2002.....	4.4	7.2	14.0	20.3	33.5	41.6	63.8	99.1	160.7	337.6
2001.....	4.0	8.0	13.6	21.1	29.8	44.7	65.7	105.2	162.5	340.2
2000.....	4.8	8.3	11.8	19.4	32.3	43.4	64.4	105.6	147.1	346.3
1999.....	4.2	8.9	12.8	21.7	33.6	44.6	70.6	109.4	170.2	359.0
Black, male										
2006 4/.....	5.6	9.1	16.0	24.1	41.9	52.3	76.6	115.0	167.7	284.2
2005 4/.....	5.0	8.4	14.8	25.7	37.4	51.8	85.1	101.5	136.2	325.4
2004 4/.....	5.4	9.9	15.5	26.8	38.9	49.9	81.3	117.8	166.4	332.8
2003 4/.....	5.4	9.8	17.6	26.1	38.2	60.3	88.4	121.0	160.9	307.6
2002.....	5.1	9.3	18.4	26.5	43.9	51.4	82.5	120.3	173.5	336.6
2001.....	4.4	9.4	17.6	28.5	40.5	50.6	86.2	135.7	198.2	357.4
2000.....	5.1	10.8	17.1	26.2	42.9	58.8	79.6	121.4	152.6	361.0
1999.....	4.7	11.1	16.5	31.0	43.5	61.0	85.8	135.3	215.4	411.3
Black, female										
2006 4/.....	3.7	5.5	9.2	13.3	20.8	31.0	51.6	77.1	139.6	271.6
2005 4/.....	3.7	6.5	11.0	14.3	22.2	30.7	50.2	81.6	125.0	295.2
2004 4/.....	5.0	5.9	9.9	15.5	20.2	28.2	48.5	83.0	127.0	308.7
2003 4/.....	4.2	6.9	9.7	14.2	21.5	36.2	51.2	79.5	136.9	335.0
2002.....	3.8	5.4	10.2	15.3	25.4	34.4	51.3	86.4	154.0	338.0
2001.....	3.7	6.7	10.2	15.0	21.4	40.4	52.0	87.1	144.3	333.8
2000.....	4.6	6.2	7.3	13.8	24.0	32.1	54.4	96.0	144.4	340.8
1999.....	3.7	6.9	9.6	14.1	25.8	32.5	60.5	93.8	147.8	339.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
All races 3/, both sexes										
2006.....	1.0	2.0	3.7	6.6	12.0	21.9	43.1	88.7	190.0	614.4
2005.....	1.0	1.7	3.5	6.2	11.8	21.5	43.6	88.8	190.5	618.6
2004.....	0.9	1.7	3.3	6.1	11.4	22.4	43.8	88.7	187.7	621.2
2003.....	1.0	1.6	3.3	6.4	12.1	23.6	44.7	89.9	192.7	641.1
2002.....	0.8	1.7	3.4	6.0	12.4	22.9	45.7	89.0	194.0	645.0
2001.....	0.8	1.7	3.3	6.5	13.0	23.8	45.6	92.6	199.2	674.6
2000.....	0.8	1.4	3.0	6.1	12.8	24.0	46.5	93.5	199.9	684.9
1999.....	0.8	1.5	2.9	6.5	11.7	23.5	48.3	94.3	201.6	681.9
All races 3/, male										
2006.....	1.2	2.6	4.9	8.2	14.5	26.6	50.0	103.3	213.5	599.0
2005.....	1.3	2.1	4.3	7.6	14.0	24.6	50.1	102.1	213.7	603.1
2004.....	1.0	2.2	4.0	7.5	13.7	26.4	51.1	103.2	211.1	605.5
2003.....	1.3	2.1	3.9	7.9	14.7	28.3	53.1	105.3	213.9	629.8
2002.....	1.0	2.0	4.4	7.2	14.8	27.0	53.2	103.9	218.0	621.4
2001.....	0.9	2.2	4.0	8.1	15.0	28.0	54.2	107.9	226.6	646.4
2000.....	1.1	1.9	3.5	7.4	15.3	28.2	55.6	107.9	229.5	672.2
1999.....	1.0	1.9	3.6	8.0	14.2	28.0	58.9	108.4	224.9	667.0
All races 3/, female										
2006.....	0.8	1.5	2.6	5.1	9.7	17.9	37.6	78.0	175.2	621.6
2005.....	0.6	1.4	2.7	4.9	9.8	18.7	38.4	79.1	176.1	625.6
2004.....	0.7	1.3	2.8	4.8	9.4	18.8	37.9	78.3	173.3	628.2
2003.....	0.7	1.2	2.7	4.9	9.7	19.6	37.8	78.9	179.8	646.1
2002.....	0.5	1.3	2.4	4.9	10.1	19.3	39.8	78.4	179.3	655.3
2001.....	0.7	1.2	2.6	5.1	11.3	20.2	38.7	81.9	182.7	686.4
2000.....	0.6	0.9	2.5	5.0	10.5	20.3	39.3	83.5	182.4	690.0
1999.....	0.6	1.1	2.1	5.1	9.5	19.6	39.9	84.5	188.0	687.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
White, both sexes										
2006 4/.....	0.8	1.5	2.8	5.6	11.0	20.5	42.2	89.0	193.3	637.2
2005 4/.....	0.7	1.2	2.7	5.3	10.5	19.8	42.3	87.6	192.4	640.6
2004 4/.....	0.7	1.4	2.5	5.1	10.3	21.5	42.9	87.9	191.0	641.5
2003 4/.....	0.7	1.3	2.6	5.6	10.7	22.4	43.2	89.5	195.3	660.4
2002.....	0.6	1.2	2.7	5.3	11.4	21.7	44.1	88.1	195.6	664.5
2001.....	0.6	1.4	2.7	5.7	12.1	22.5	44.3	92.1	201.1	695.1
2000.....	0.6	1.1	2.5	5.4	11.8	22.6	44.9	92.8	201.8	703.7
1999.....	0.6	1.1	2.3	5.6	10.9	22.0	46.8	92.5	202.5	699.2
White, male										
2006 4/.....	0.9	1.9	3.7	7.2	13.4	25.3	48.3	104.2	217.8	622.6
2005 4/.....	0.8	1.5	3.4	6.5	12.5	22.7	48.4	100.6	217.9	629.3
2004 4/.....	0.8	1.6	3.1	6.2	12.2	25.2	50.7	101.7	215.6	625.5
2003 4/.....	0.9	1.7	3.0	7.0	13.1	27.0	52.1	105.9	217.1	650.8
2002.....	0.8	1.5	3.6	6.4	13.6	25.5	51.4	103.4	221.4	641.9
2001.....	0.7	1.9	3.4	7.2	14.2	26.5	52.8	107.6	228.2	666.5
2000.....	0.9	1.5	3.0	6.8	14.1	27.2	54.3	107.7	231.7	690.0
1999.....	0.8	1.4	2.9	7.1	13.3	26.2	58.1	107.2	226.3	680.0
White, female										
2006 4/.....	0.6	1.0	2.0	4.1	8.8	16.3	37.1	77.6	177.8	644.0
2005 4/.....	0.5	1.0	2.1	4.2	8.6	17.3	37.2	78.0	176.5	645.8
2004 4/.....	0.6	1.1	2.0	4.0	8.6	18.1	36.5	77.8	175.7	648.7
2003 4/.....	0.4	0.9	2.1	4.2	8.5	18.3	35.9	77.6	181.9	664.7
2002.....	0.4	0.9	1.9	4.2	9.3	18.4	38.2	77.1	179.8	674.3
2001.....	0.5	0.8	2.0	4.2	10.1	18.9	37.4	81.0	184.7	707.1
2000.....	0.4	0.6	2.0	4.1	9.6	18.6	37.3	82.4	183.9	709.3
1999.....	0.4	0.8	1.7	4.2	8.8	18.4	37.8	82.1	188.5	707.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Heart failure (I50)										
Black, both sexes										
2006 4/.....	2.8	6.4	10.7	16.1	24.5	41.1	64.8	109.6	200.0	484.9
2005 4/.....	3.0	5.2	10.0	15.6	26.8	41.3	69.1	123.8	216.0	489.6
2004 4/.....	2.2	5.0	10.5	16.3	23.8	35.7	64.0	120.2	194.6	506.5
2003 4/.....	3.3	4.7	9.4	15.3	27.7	41.1	70.9	114.7	203.6	531.8
2002.....	2.2	5.5	9.3	13.9	24.6	38.6	71.7	118.5	215.3	528.0
2001.....	2.7	4.5	8.7	15.7	24.6	41.2	71.1	121.1	213.7	543.8
2000.....	2.2	4.2	7.6	14.0	24.3	41.2	72.6	119.8	209.5	558.4
1999.....	2.4	4.8	8.4	16.4	22.0	42.0	74.0	130.5	216.7	552.9
Black, male										
2006 4/.....	3.7	8.0	14.9	18.4	31.3	49.0	82.1	126.2	223.8	463.8
2005 4/.....	4.6	6.4	12.6	20.1	32.7	48.5	82.5	152.0	224.0	423.0
2004 4/.....	2.6	6.8	12.2	21.0	30.6	45.5	69.3	148.3	220.2	503.7
2003 4/.....	3.9	5.5	11.6	19.4	35.9	51.0	79.5	124.6	227.4	513.0
2002.....	2.7	6.6	12.0	16.5	31.3	47.6	87.5	136.9	236.9	479.2
2001.....	2.6	5.0	10.4	18.9	26.4	49.6	86.2	140.8	256.8	541.2
2000.....	2.5	5.5	9.1	14.8	29.6	45.5	82.9	134.7	242.8	581.2
1999.....	2.7	6.4	11.0	19.3	27.4	51.4	83.1	146.7	238.9	603.5
Black, female										
2006 4/.....	2.0	5.0	7.2	14.3	19.1	35.3	52.8	99.6	187.4	493.5
2005 4/.....	1.6	4.0	7.7	11.9	22.1	36.0	60.0	106.9	211.8	516.5
2004 4/.....	1.9	3.4	8.9	12.4	18.5	28.4	60.4	103.5	181.1	507.6
2003 4/.....	2.8	4.0	7.6	11.9	21.4	33.8	65.1	108.8	191.1	539.2
2002.....	1.7	4.5	7.0	11.7	19.4	31.8	61.2	107.6	204.1	546.7
2001.....	2.8	4.1	7.2	13.1	23.2	35.1	61.0	109.3	191.7	544.8
2000.....	1.9	3.0	6.4	13.4	20.1	38.0	65.7	110.8	192.8	549.9
1999.....	2.1	3.5	6.3	14.0	17.8	35.0	68.0	120.7	205.7	534.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congestive heart failure (I50.0)										
All races 3/, both sexes										
2006.....	0.8	1.6	3.1	5.6	10.6	19.6	39.1	81.6	176.9	579.3
2005.....	0.8	1.4	2.9	5.3	10.3	19.3	39.7	81.9	179.0	586.0
2004.....	0.7	1.5	2.8	5.1	9.9	19.8	40.0	82.1	175.8	588.9
2003.....	0.8	1.4	2.8	5.5	10.6	21.2	40.6	82.9	181.6	608.2
2002.....	0.6	1.3	2.8	5.1	10.7	20.3	41.4	82.3	181.6	609.0
2001.....	0.7	1.4	2.8	5.4	11.3	21.5	41.5	85.8	185.4	636.1
2000.....	0.7	1.1	2.5	5.4	11.1	20.9	41.8	86.1	186.9	644.4
1999.....	0.7	1.2	2.3	5.3	10.0	20.9	43.6	86.6	188.4	642.7
All races 3/, male										
2006.....	0.9	2.1	3.9	7.0	12.6	23.6	44.9	94.3	196.7	565.2
2005.....	1.1	1.7	3.5	6.4	12.0	21.9	45.1	94.5	200.5	571.0
2004.....	0.8	1.8	3.3	6.2	12.0	23.0	46.5	95.6	196.2	571.8
2003.....	1.0	1.7	3.2	6.7	12.6	25.5	48.1	96.9	200.7	594.6
2002.....	0.8	1.6	3.5	6.1	12.5	23.8	48.2	95.6	203.2	584.6
2001.....	0.8	1.9	3.3	6.6	12.8	24.9	49.1	98.7	208.5	609.3
2000.....	0.8	1.6	2.9	6.4	13.0	24.7	49.6	99.1	214.1	632.2
1999.....	0.8	1.5	2.9	6.5	11.9	24.5	53.2	99.2	209.4	628.2
All races 3/, female										
2006.....	0.6	1.2	2.3	4.4	8.7	16.1	34.4	72.4	164.5	585.9
2005.....	0.5	1.0	2.3	4.2	8.8	17.0	35.3	72.8	165.6	592.9
2004.....	0.6	1.2	2.4	4.1	8.0	16.9	34.8	72.4	163.3	596.6
2003.....	0.6	1.0	2.3	4.3	8.7	17.6	34.6	72.9	169.9	614.2
2002.....	0.5	1.0	2.1	4.2	9.0	17.2	36.0	72.8	168.4	619.6
2001.....	0.6	1.0	2.3	4.2	10.0	18.5	35.5	76.8	171.5	647.3
2000.....	0.5	0.7	2.1	4.4	9.4	17.7	35.7	77.0	170.9	649.4
1999.....	0.5	0.9	1.8	4.2	8.4	17.9	36.1	77.9	176.1	648.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congestive heart failure (I50.0)										
White, both sexes										
2006 4/.....	0.5	1.2	2.4	4.8	9.8	18.4	38.4	82.0	180.4	602.1
2005 4/.....	0.5	1.0	2.3	4.5	9.3	17.9	38.6	81.3	181.8	608.3
2004 4/.....	0.6	1.1	2.2	4.3	9.0	19.1	39.3	81.5	179.5	609.2
2003 4/.....	0.5	1.0	2.2	4.8	9.3	20.3	39.5	82.7	184.5	628.4
2002.....	0.5	0.9	2.2	4.5	9.8	19.3	40.1	81.7	183.6	628.8
2001.....	0.5	1.1	2.3	4.7	10.5	20.2	40.5	85.7	187.6	656.5
2000.....	0.5	0.8	2.1	4.8	10.2	19.7	40.6	85.6	189.3	663.6
1999.....	0.5	0.8	1.8	4.5	9.4	19.6	42.3	85.3	189.4	660.1
White, male										
2006 4/.....	0.6	1.5	3.0	6.2	11.8	22.6	43.7	95.5	201.4	588.7
2005 4/.....	0.7	1.2	2.8	5.5	10.8	20.2	43.7	93.3	205.6	597.3
2004 4/.....	0.6	1.4	2.6	5.2	10.6	22.1	46.3	94.3	201.1	591.7
2003 4/.....	0.7	1.4	2.6	5.9	11.2	24.5	47.2	97.8	204.3	616.2
2002.....	0.6	1.2	2.8	5.4	11.4	22.5	46.5	95.3	207.2	605.8
2001.....	0.6	1.6	2.8	6.0	12.1	23.5	47.9	99.1	210.8	629.0
2000.....	0.7	1.2	2.4	6.1	11.8	23.9	48.6	99.2	216.9	650.5
1999.....	0.7	1.1	2.3	5.7	11.0	22.9	52.5	98.4	210.7	641.7
White, female										
2006 4/.....	0.5	0.8	1.8	3.5	7.9	14.8	34.0	72.0	167.2	608.4
2005 4/.....	0.4	0.8	1.7	3.6	7.9	15.8	34.4	72.4	166.9	613.4
2004 4/.....	0.5	0.9	1.7	3.4	7.5	16.4	33.6	72.2	166.1	617.1
2003 4/.....	0.3	0.7	1.9	3.7	7.6	16.6	33.1	71.8	172.3	633.9
2002.....	0.3	0.7	1.6	3.6	8.2	16.5	35.0	71.9	169.1	638.8
2001.....	0.4	0.7	1.9	3.5	9.1	17.2	34.4	76.1	173.6	667.9
2000.....	0.4	0.5	1.7	3.5	8.7	16.0	34.1	76.1	172.8	668.8
1999.....	0.4	0.6	1.3	3.4	7.8	16.8	34.2	76.2	176.9	667.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congestive heart failure (I50.0)										
Black, both sexes										
2006 4/.....	2.3	5.2	9.1	13.7	21.1	36.0	56.7	98.7	182.0	444.2
2005 4/.....	2.7	4.0	8.0	12.9	22.4	36.6	61.5	110.1	192.0	448.2
2004 4/.....	1.7	4.2	8.5	13.8	20.1	30.7	57.3	109.5	174.8	468.7
2003 4/.....	2.9	4.1	7.6	12.9	24.3	35.8	63.5	102.9	185.8	484.3
2002.....	1.9	4.6	7.7	11.9	21.9	34.4	63.8	106.7	196.0	482.6
2001.....	2.2	4.0	7.3	13.3	21.0	37.6	63.1	108.5	194.2	502.4
2000.....	1.8	3.6	6.7	11.9	21.3	36.5	63.0	108.4	190.3	510.2
1999.....	1.8	4.2	7.3	13.6	18.7	37.2	66.6	116.4	200.2	509.5
Black, male										
2006 4/.....	2.9	6.6	12.2	15.8	26.2	42.9	69.5	109.2	199.2	419.1
2005 4/.....	3.9	5.2	9.9	16.3	26.9	43.0	73.1	137.7	197.3	381.1
2004 4/.....	1.9	5.5	9.6	17.9	26.7	38.5	61.2	135.6	194.4	466.9
2003 4/.....	3.3	4.8	9.1	16.4	30.7	44.6	71.6	110.5	203.8	466.1
2002.....	2.2	5.6	9.6	14.3	27.3	42.9	81.5	122.6	209.4	428.8
2001.....	2.3	4.5	8.8	15.2	22.6	44.2	75.4	123.1	224.7	500.4
2000.....	2.0	4.7	8.0	11.7	26.1	38.2	71.2	120.0	216.6	528.2
1999.....	2.1	5.6	9.2	15.4	23.1	44.9	74.6	131.5	224.1	554.2
Black, female										
2006 4/.....	1.7	4.1	6.5	11.9	17.1	31.0	48.0	92.4	172.9	454.4
2005 4/.....	1.5	2.9	6.4	10.1	18.8	31.9	53.7	93.6	189.2	475.3
2004 4/.....	1.5	3.1	7.5	10.4	15.0	24.8	54.7	94.1	164.4	469.4
2003 4/.....	2.5	3.5	6.3	9.9	19.4	29.2	58.0	98.5	176.4	491.4
2002.....	1.6	3.7	6.1	9.9	17.7	28.1	52.0	97.2	189.0	503.2
2001.....	2.2	3.6	6.0	11.7	19.7	32.7	54.9	99.8	178.7	503.1
2000.....	1.5	2.5	5.5	12.0	17.5	35.1	57.5	101.4	177.0	503.5
1999.....	1.6	3.0	5.7	12.0	15.3	31.6	61.2	107.3	188.4	492.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified heart failure (I50.1-I50.9)										
All races 3/, both sexes										
2006.....	0.2	0.4	0.6	0.9	1.4	2.3	4.1	7.1	13.1	35.1
2005.....	0.2	0.4	0.7	0.9	1.4	2.2	3.9	6.8	11.5	32.5
2004.....	0.2	0.3	0.5	1.0	1.6	2.6	3.8	6.6	11.9	32.3
2003.....	0.2	0.3	0.5	0.9	1.5	2.4	4.0	7.0	11.2	32.9
2002.....	0.1	0.4	0.6	0.9	1.7	2.6	4.3	6.7	12.4	36.1
2001.....	0.2	0.3	0.5	1.1	1.7	2.4	4.1	6.8	13.8	38.5
2000.....	0.2	0.3	0.5	0.8	1.7	3.0	4.7	7.4	12.9	40.5
1999.....	0.2	0.3	0.5	1.2	1.7	2.6	4.6	7.7	13.3	39.2
All races 3/, male										
2006.....	0.3	0.5	0.9	1.2	1.8	3.0	5.1	9.1	16.7	33.8
2005.....	0.2	0.4	0.8	1.2	2.0	2.7	4.9	7.7	13.2	32.1
2004.....	0.2	0.4	0.7	1.2	1.7	3.4	4.7	7.6	15.0	33.7
2003.....	0.3	0.4	0.7	1.2	2.1	2.8	5.0	8.4	13.2	35.2
2002.....	0.2	0.4	1.0	1.1	2.3	3.2	5.0	8.3	14.8	36.8
2001.....	0.2	0.3	0.7	1.5	2.1	3.1	5.2	9.2	18.1	37.2
2000.....	0.3	0.4	0.7	0.9	2.3	3.6	6.1	8.8	15.4	40.0
1999.....	0.2	0.4	0.7	1.6	2.4	3.5	5.7	9.2	15.5	38.8
All races 3/, female										
2006.....	*	0.3	0.3	0.7	1.0	1.8	3.2	5.6	10.8	35.7
2005.....	*	0.3	0.5	0.7	1.0	1.7	3.1	6.2	10.5	32.7
2004.....	*	0.2	0.4	0.7	1.4	1.9	3.1	5.8	10.0	31.6
2003.....	*	0.2	0.3	0.7	1.0	2.0	3.2	6.0	9.9	31.9
2002.....	*	0.3	0.3	0.7	1.1	2.1	3.8	5.6	10.9	35.7
2001.....	*	*	0.3	0.8	1.3	1.7	3.2	5.2	11.2	39.1
2000.....	*	*	0.4	0.6	1.1	2.5	3.6	6.5	11.5	40.6
1999.....	*	0.3	0.4	0.9	1.1	1.7	3.8	6.6	11.9	39.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified heart failure (I50.1-I50.9)										
White, both sexes										
2006 4/.....	0.2	0.3	0.5	0.8	1.2	2.1	3.8	6.9	12.9	35.1
2005 4/.....	0.1	0.2	0.5	0.8	1.2	1.9	3.7	6.4	10.6	32.3
2004 4/.....	0.1	0.2	0.3	0.8	1.4	2.4	3.6	6.4	11.5	32.3
2003 4/.....	0.2	0.3	0.3	0.8	1.3	2.1	3.7	6.7	10.9	32.0
2002.....	0.1	0.3	0.5	0.8	1.6	2.4	4.0	6.4	12.0	35.6
2001.....	0.1	0.2	0.4	1.0	1.5	2.3	3.8	6.4	13.5	38.7
2000.....	0.1	0.2	0.5	0.7	1.6	2.9	4.3	7.2	12.5	40.2
1999.....	0.1	0.3	0.5	1.1	1.6	2.4	4.5	7.2	13.1	39.2
White, male										
2006 4/.....	0.3	0.4	0.7	1.0	1.6	2.7	4.6	8.7	16.4	33.9
2005 4/.....	*	0.3	0.6	1.0	1.7	2.5	4.7	7.3	12.3	32.0
2004 4/.....	*	0.3	0.5	1.1	1.6	3.1	4.4	7.4	14.5	33.8
2003 4/.....	0.3	0.3	0.4	1.0	1.8	2.5	4.9	8.1	12.8	34.6
2002.....	*	0.3	0.8	1.0	2.2	3.0	4.9	8.0	14.2	36.0
2001.....	*	0.3	0.6	1.3	2.0	3.0	4.9	8.5	17.4	37.5
2000.....	0.2	0.3	0.6	0.7	2.2	3.2	5.7	8.4	14.8	39.5
1999.....	*	0.4	0.6	1.4	2.2	3.3	5.6	8.8	15.6	38.3
White, female										
2006 4/.....	*	*	0.2	0.5	0.9	1.5	3.2	5.6	10.6	35.6
2005 4/.....	*	*	0.4	0.6	0.7	1.5	2.8	5.7	9.6	32.4
2004 4/.....	*	*	*	0.6	1.1	1.7	3.0	5.6	9.7	31.6
2003 4/.....	*	*	*	0.5	0.9	1.7	2.8	5.7	9.7	30.8
2002.....	*	0.2	*	0.6	1.1	1.9	3.3	5.2	10.6	35.5
2001.....	*	*	*	0.8	1.1	1.7	3.0	4.9	11.1	39.2
2000.....	*	*	0.3	0.6	1.0	2.6	3.2	6.3	11.1	40.5
1999.....	*	*	0.3	0.8	1.0	1.6	3.6	6.0	11.6	39.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other and unspecified heart failure (I50.1-I50.9)										
Black, both sexes										
2006 4/.....	*	1.1	1.6	2.4	3.4	5.1	8.0	10.9	18.0	40.7
2005 4/.....	*	1.2	2.0	2.7	4.4	4.7	7.6	13.7	24.0	41.4
2004 4/.....	*	0.8	2.0	2.4	3.7	5.1	6.7	10.7	19.8	37.8
2003 4/.....	*	*	1.9	2.4	3.4	5.3	7.4	11.7	17.8	47.6
2002.....	*	0.9	1.6	2.0	2.7	4.1	7.9	11.9	19.4	45.4
2001.....	*	*	1.4	2.5	3.7	3.7	7.9	12.5	19.5	41.5
2000.....	*	*	*	2.2	3.0	4.8	9.6	11.4	19.2	48.2
1999.....	*	*	*	2.8	3.3	4.8	7.4	14.1	16.5	43.4
Black, male										
2006 4/.....	*	*	2.7	2.5	5.1	6.1	12.6	17.0	24.6	44.7
2005 4/.....	*	*	2.8	3.8	5.8	5.5	9.5	14.3	26.7	41.8
2004 4/.....	*	*	2.6	3.0	3.9	7.0	8.1	12.7	25.8	36.8
2003 4/.....	*	*	2.5	2.9	5.2	6.4	7.9	14.2	23.6	46.9
2002.....	*	*	2.5	*	4.0	*	*	14.2	27.5	50.4
2001.....	*	*	*	3.8	*	5.3	10.8	17.7	32.1	40.8
2000.....	*	*	*	*	*	7.3	11.7	14.6	26.1	53.0
1999.....	*	*	*	3.9	4.2	6.5	8.5	15.2	*	49.2
Black, female										
2006 4/.....	*	*	*	2.4	*	4.3	4.9	7.2	14.5	39.1
2005 4/.....	*	*	*	*	3.3	4.1	6.3	13.3	22.6	41.3
2004 4/.....	*	*	*	*	3.5	3.6	5.7	9.4	16.6	38.2
2003 4/.....	*	*	*	*	*	4.6	7.1	10.3	14.7	47.8
2002.....	*	*	*	*	*	3.7	9.2	10.4	15.1	43.5
2001.....	*	*	*	*	3.5	*	6.1	9.4	13.1	41.7
2000.....	*	*	*	*	*	*	8.2	9.4	15.7	46.4
1999.....	*	*	*	*	*	*	6.8	13.4	17.3	41.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Myocarditis, unspecified and myocardial degeneration (I51.4-I51.5)

All races 3/, both sexes

2006.....	0.2	0.3	0.3	0.2	0.2	*	0.2	0.3	0.4	1.5
2005.....	0.3	0.3	0.2	0.2	0.2	*	0.3	0.3	0.6	1.9
2004.....	0.3	0.3	0.2	0.2	*	0.3	0.3	0.3	0.4	2.0
2003.....	0.2	0.2	0.2	0.2	0.2	*	*	0.3	0.5	2.2
2002.....	0.3	0.3	0.3	0.4	0.2	0.4	0.6	0.8	1.1	3.2
2001.....	0.3	0.3	0.3	0.2	0.3	0.4	0.4	0.5	0.6	2.0
2000.....	0.3	0.3	0.1	0.3	0.3	*	0.4	0.4	0.7	2.4
1999.....	0.3	0.3	0.3	0.2	0.2	0.3	0.3	0.4	0.7	2.4

All races 3/, male

2006.....	0.2	0.3	0.3	*	*	*	*	*	*	1.7
2005.....	0.3	0.3	0.3	*	*	*	*	*	*	1.9
2004.....	0.3	0.3	0.3	*	*	*	*	*	*	1.7
2003.....	*	0.2	0.2	0.3	*	*	*	*	*	2.6
2002.....	0.4	0.3	0.3	0.3	*	*	0.8	0.8	1.0	3.8
2001.....	0.3	0.3	0.3	*	*	0.5	0.5	0.8	*	2.2
2000.....	0.3	0.2	*	0.4	*	*	*	*	*	2.6
1999.....	0.3	0.3	0.3	*	*	*	*	*	*	2.1

All races 3/, female

2006.....	0.2	0.3	0.3	0.2	*	*	*	*	*	1.4
2005.....	0.3	0.3	0.2	*	*	*	*	*	0.7	1.9
2004.....	0.2	0.3	0.2	*	*	*	*	*	*	2.2
2003.....	0.3	0.3	0.2	*	*	*	*	*	*	2.0
2002.....	0.3	0.3	0.3	0.4	*	*	0.5	0.8	1.1	3.0
2001.....	0.3	0.3	*	*	*	*	*	*	*	1.9
2000.....	0.2	0.3	*	0.3	*	*	*	*	*	2.4
1999.....	0.3	0.3	0.2	*	*	*	*	*	*	2.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

All races 3/, both sexes

2006.....	1.5	1.9	2.3	3.1	4.6	6.1	9.6	15.2	28.8	71.2
2005.....	1.3	1.9	2.4	3.2	4.4	6.0	9.6	15.3	27.1	69.2
2004.....	1.3	1.7	2.4	3.0	4.4	5.9	8.6	15.1	27.4	67.7
2003.....	1.3	1.7	2.1	3.1	4.2	5.8	9.6	15.2	27.9	67.8
2002.....	1.3	1.7	2.1	2.8	4.2	6.1	9.7	15.3	26.4	65.0
2001.....	1.2	1.7	2.0	2.8	4.1	5.7	9.0	14.4	25.4	66.3
2000.....	1.1	1.6	2.0	3.1	4.5	6.4	8.7	14.3	26.8	64.6
1999.....	1.0	1.4	1.8	2.8	4.3	5.6	8.3	14.1	23.4	61.5

All races 3/, male

2006.....	1.9	2.7	3.3	4.3	6.6	8.6	12.7	19.3	34.5	73.6
2005.....	1.8	2.7	3.4	4.2	6.0	8.6	12.6	19.5	31.0	71.8
2004.....	1.8	2.4	3.7	4.1	5.7	8.7	10.7	19.3	33.2	67.8
2003.....	1.9	2.3	2.9	4.3	5.8	7.5	11.5	19.3	32.5	72.1
2002.....	1.8	2.4	3.0	4.0	5.5	8.2	13.5	18.9	30.7	66.7
2001.....	1.7	2.4	2.8	3.9	5.8	7.8	12.3	17.8	28.8	66.4
2000.....	1.7	2.3	2.8	4.4	6.5	8.1	11.3	17.0	31.2	69.0
1999.....	1.5	2.1	2.5	4.0	6.2	7.8	11.2	18.3	27.7	64.3

All races 3/, female

2006.....	1.0	1.1	1.3	2.0	2.7	3.9	7.0	12.2	25.2	70.1
2005.....	0.8	1.1	1.4	2.2	3.0	3.8	7.3	12.3	24.6	68.0
2004.....	0.9	1.0	1.2	1.8	3.1	3.5	6.9	12.1	23.9	67.7
2003.....	0.7	1.1	1.3	2.0	2.7	4.3	8.1	12.3	25.0	66.0
2002.....	0.8	1.1	1.3	1.6	3.0	4.3	6.6	12.8	23.8	64.3
2001.....	0.7	0.9	1.2	1.9	2.6	3.9	6.5	12.0	23.3	66.3
2000.....	0.6	0.8	1.2	1.9	2.7	4.9	6.6	12.5	24.1	62.7
1999.....	0.6	0.8	1.1	1.7	2.7	3.7	6.0	11.2	20.8	60.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)										
White, both sexes										
2006 4/.....	1.4	1.8	2.1	2.9	4.2	5.9	9.3	14.9	28.6	72.5
2005 4/.....	1.2	1.7	2.2	2.9	4.1	5.7	9.5	14.9	27.0	70.0
2004 4/.....	1.2	1.5	2.3	2.7	4.1	5.6	8.0	14.6	27.1	68.1
2003 4/.....	1.2	1.5	1.8	2.9	3.9	5.5	9.1	15.1	27.8	67.5
2002.....	1.1	1.5	1.8	2.6	3.7	5.7	9.4	15.2	26.3	64.5
2001.....	1.0	1.4	1.7	2.5	4.0	5.4	8.6	13.7	25.1	65.7
2000.....	1.0	1.5	1.7	2.8	4.3	6.1	7.9	13.9	26.1	64.7
1999.....	0.9	1.2	1.6	2.4	3.9	5.1	8.1	13.3	23.2	61.4
White, male										
2006 4/.....	1.9	2.6	3.0	3.9	6.1	8.1	12.3	18.6	34.5	75.1
2005 4/.....	1.7	2.5	3.1	3.7	5.6	8.2	12.4	19.1	31.3	73.2
2004 4/.....	1.6	2.3	3.4	3.9	5.4	8.3	9.9	18.7	33.0	68.2
2003 4/.....	1.8	2.1	2.6	4.0	5.5	7.2	11.1	19.2	32.4	72.2
2002.....	1.6	2.1	2.6	3.7	4.9	7.6	13.3	18.7	30.2	66.9
2001.....	1.6	2.1	2.6	3.5	5.7	7.5	11.6	17.0	28.4	66.2
2000.....	1.5	2.2	2.4	3.9	6.3	7.7	10.3	16.4	30.0	69.2
1999.....	1.2	2.0	2.2	3.5	5.7	7.2	10.8	17.5	27.2	64.5
White, female										
2006 4/.....	0.9	1.0	1.2	1.8	2.6	3.9	6.8	12.1	24.8	71.2
2005 4/.....	0.6	1.0	1.2	2.0	2.6	3.4	7.1	11.8	24.3	68.5
2004 4/.....	0.8	0.8	1.1	1.6	2.8	3.2	6.4	11.7	23.5	68.1
2003 4/.....	0.6	0.9	1.1	1.8	2.5	4.0	7.5	12.2	25.0	65.4
2002.....	0.7	0.8	1.0	1.5	2.7	4.0	6.3	12.8	23.9	63.4
2001.....	0.5	0.7	0.9	1.6	2.5	3.5	6.2	11.3	23.1	65.5
2000.....	0.5	0.7	0.9	1.7	2.4	4.6	6.0	12.1	23.8	62.8
1999.....	0.5	0.5	0.9	1.4	2.2	3.2	5.9	10.3	20.8	60.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and ill-defined forms of heart disease (I50.0-I51.3,I51.6-I51.9)

Black, both sexes

2006 4/.....	2.3	3.2	4.8	6.3	8.4	9.3	14.8	23.3	40.9	70.3
2005 4/.....	2.4	3.1	5.0	6.3	7.9	10.6	13.5	23.9	34.2	69.2
2004 4/.....	2.6	3.6	4.0	5.8	8.2	10.8	17.0	24.5	37.9	76.4
2003 4/.....	2.4	3.2	4.7	5.7	7.5	9.8	16.7	19.4	34.0	80.5
2002.....	2.3	3.6	5.4	5.1	8.7	11.2	14.5	19.9	32.2	80.5
2001.....	2.6	3.7	4.5	6.1	6.4	9.4	15.2	23.8	32.2	83.8
2000.....	2.4	2.9	4.8	6.5	8.0	11.0	18.7	22.2	37.8	73.8
1999.....	2.6	3.1	3.8	6.7	9.1	11.3	13.3	24.8	26.9	68.9

Black, male

2006 4/.....	2.6	4.3	7.0	8.7	12.9	15.9	21.2	34.5	49.3	69.7
2005 4/.....	2.9	4.5	6.7	9.7	10.0	15.2	18.0	31.3	36.5	62.3
2004 4/.....	3.5	4.4	6.2	8.0	11.0	15.5	22.7	35.0	45.9	75.5
2003 4/.....	3.4	3.8	6.5	7.7	11.0	13.0	18.7	25.6	44.3	80.3
2002.....	3.2	4.6	7.2	7.3	12.1	16.0	18.7	26.6	42.8	77.2
2001.....	3.0	5.1	5.3	8.4	8.3	10.9	21.9	28.8	39.3	84.0
2000.....	3.3	4.1	7.1	9.7	10.6	14.1	25.4	26.9	52.3	80.7
1999.....	3.6	3.8	5.3	9.7	11.7	16.1	19.0	29.9	36.6	72.6

Black, female

2006 4/.....	1.9	2.3	3.0	4.3	5.0	4.3	10.4	16.6	36.5	70.6
2005 4/.....	1.9	2.0	3.5	3.5	6.2	7.2	10.4	19.5	33.0	72.0
2004 4/.....	1.8	2.9	2.2	3.9	6.0	7.4	13.2	18.3	33.6	76.8
2003 4/.....	1.5	2.7	3.2	4.0	4.8	7.3	15.3	15.7	28.6	80.6
2002.....	1.5	2.6	4.0	3.3	6.0	7.6	11.6	15.9	26.7	81.7
2001.....	2.3	2.5	3.9	4.2	4.8	8.3	10.8	20.8	28.6	83.8
2000.....	1.7	1.8	2.8	3.9	6.0	8.8	14.2	19.4	30.6	71.3
1999.....	1.6	2.6	2.6	4.3	7.2	7.7	9.5	21.7	22.0	67.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
All races 3/, both sexes										
2006.....	6.7	11.2	18.5	26.4	42.7	70.7	127.5	240.3	458.6	1,039.6
2005.....	6.9	11.7	18.8	25.8	42.7	73.8	133.6	253.6	497.3	1,141.8
2004.....	7.1	11.9	18.3	26.5	44.5	78.4	142.2	276.7	532.1	1,245.9
2003.....	7.4	11.8	18.6	27.8	45.9	79.7	150.7	294.1	571.1	1,370.1
2002.....	7.1	12.3	18.3	28.6	48.3	83.8	160.7	308.8	601.7	1,445.9
2001.....	7.2	12.1	18.5	28.8	49.6	85.7	164.3	320.2	622.4	1,500.2
2000.....	7.7	13.1	19.3	30.8	53.8	91.0	169.1	332.8	654.0	1,589.2
1999.....	7.5	12.5	18.4	31.7	51.4	91.1	173.5	340.6	664.0	1,614.8
All races 3/, male										
2006.....	6.9	12.0	21.2	31.2	49.2	81.9	140.9	259.6	468.5	932.4
2005.....	7.0	12.1	21.5	30.1	50.0	84.6	149.5	275.8	513.0	1,023.3
2004.....	7.4	12.8	21.1	29.9	52.2	90.2	158.8	301.2	551.6	1,118.1
2003.....	7.6	12.9	21.1	31.3	53.5	91.4	170.8	322.5	595.6	1,236.0
2002.....	6.9	13.4	20.4	32.7	55.8	97.0	178.7	334.8	615.6	1,317.9
2001.....	7.4	13.2	20.5	32.8	57.2	98.5	187.9	349.0	654.6	1,380.2
2000.....	7.8	14.2	21.3	35.2	62.3	102.6	192.9	367.6	695.1	1,484.3
1999.....	7.8	13.3	20.7	35.7	60.0	102.8	197.7	376.9	706.8	1,512.8
All races 3/, female										
2006.....	6.5	10.4	15.8	21.9	36.8	60.9	116.5	226.2	452.3	1,089.8
2005.....	6.7	11.3	16.2	21.8	36.0	64.3	120.7	237.6	487.5	1,196.1
2004.....	6.8	11.0	15.5	23.2	37.5	68.1	128.8	259.1	520.2	1,303.4
2003.....	7.2	10.8	16.3	24.6	38.9	69.5	134.5	273.9	556.1	1,429.4
2002.....	7.2	11.2	16.3	24.8	41.4	72.4	146.2	290.4	593.3	1,501.5
2001.....	7.1	10.9	16.5	25.1	42.7	74.7	145.6	300.0	603.2	1,550.4
2000.....	7.5	12.0	17.3	26.6	46.0	81.0	150.3	308.5	629.7	1,632.0
1999.....	7.1	11.7	16.1	28.0	43.7	81.1	154.5	315.3	639.0	1,656.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
White, both sexes										
2006 4/.....	5.3	8.9	14.5	21.3	36.3	62.4	117.8	231.6	453.0	1,050.8
2005 4/.....	5.3	9.1	14.5	21.1	36.2	65.6	123.8	244.2	491.2	1,153.3
2004 4/.....	5.7	9.1	14.1	21.8	37.5	70.0	132.6	264.8	526.2	1,257.8
2003 4/.....	5.6	9.0	14.2	22.6	38.3	70.8	140.3	283.2	565.9	1,382.2
2002.....	5.5	9.5	14.0	23.8	41.2	74.2	150.1	298.6	595.7	1,459.5
2001.....	5.6	9.3	13.8	23.9	41.7	75.2	153.2	309.7	617.1	1,511.2
2000.....	5.9	9.7	15.4	25.4	46.5	81.9	159.0	321.3	647.0	1,601.6
1999.....	5.7	9.4	14.4	26.2	43.5	82.9	161.3	329.3	658.4	1,631.6
White, male										
2006 4/.....	5.6	9.6	16.4	24.5	41.0	71.7	128.9	249.8	464.4	941.3
2005 4/.....	5.5	9.5	16.5	24.2	41.8	74.3	138.0	266.4	505.3	1,033.7
2004 4/.....	6.1	9.7	16.3	24.1	43.3	80.6	147.0	288.0	546.9	1,129.3
2003 4/.....	5.9	10.1	16.0	24.7	44.5	81.8	158.5	310.0	590.4	1,247.0
2002.....	5.5	10.6	15.5	26.9	47.2	85.1	167.3	323.4	610.1	1,327.4
2001.....	5.9	10.2	15.4	26.9	48.1	86.7	174.8	337.7	648.8	1,386.2
2000.....	6.1	10.8	16.9	28.6	53.8	91.7	180.5	354.6	686.1	1,490.7
1999.....	6.1	9.8	16.2	29.5	50.6	93.2	183.6	363.2	700.5	1,522.7
White, female										
2006 4/.....	4.9	8.2	12.7	18.2	31.8	54.1	108.6	218.0	445.7	1,102.2
2005 4/.....	5.2	8.6	12.5	18.2	31.1	57.8	112.1	227.7	482.4	1,208.5
2004 4/.....	5.2	8.5	11.9	19.5	32.2	60.5	120.7	247.9	513.3	1,315.7
2003 4/.....	5.3	7.9	12.3	20.5	32.5	61.1	125.4	263.8	550.7	1,442.1
2002.....	5.5	8.4	12.6	20.9	35.6	64.5	136.0	280.8	586.8	1,516.9
2001.....	5.4	8.3	12.2	21.1	35.8	65.1	135.7	289.8	598.0	1,563.5
2000.....	5.7	8.7	13.9	22.4	39.8	73.3	141.9	297.9	623.8	1,646.7
1999.....	5.3	8.9	12.7	23.0	37.0	73.9	143.7	305.5	633.6	1,675.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebrovascular diseases (I60-I69)										
Black, both sexes										
2006 4/.....	15.9	27.2	48.6	67.5	101.4	143.3	221.7	349.3	557.8	1,008.7
2005 4/.....	17.1	30.5	50.3	65.4	99.5	145.1	231.4	372.3	611.0	1,106.7
2004 4/.....	17.2	31.5	48.0	66.3	108.1	154.9	246.5	418.6	647.9	1,211.2
2003 4/.....	18.6	31.4	52.7	72.4	113.1	157.4	256.3	433.2	677.0	1,348.7
2002.....	17.4	32.1	51.2	71.1	109.9	169.6	272.5	447.2	712.5	1,383.9
2001.....	18.4	32.6	55.0	70.6	117.1	178.3	277.6	460.2	716.4	1,452.7
2000.....	19.7	37.2	51.2	77.4	114.0	174.8	281.1	475.2	769.4	1,530.0
1999.....	19.9	34.4	50.6	78.4	118.2	170.0	307.8	487.9	756.4	1,492.5
Black, male										
2006 4/.....	15.6	29.0	61.1	88.9	132.2	179.3	270.4	400.0	579.8	882.0
2005 4/.....	17.4	32.0	60.5	83.6	133.4	184.9	276.7	407.2	655.2	983.5
2004 4/.....	16.9	35.4	56.4	84.7	139.4	188.3	297.8	470.9	677.2	1,061.0
2003 4/.....	18.8	34.2	62.6	90.6	141.7	184.9	307.4	503.0	727.9	1,180.3
2002.....	16.9	34.3	61.6	88.9	138.8	213.7	328.8	502.5	729.8	1,252.2
2001.....	17.8	36.5	63.9	87.6	143.4	218.5	336.3	524.6	768.1	1,349.1
2000.....	20.5	41.0	60.4	97.8	137.5	212.8	340.0	544.8	863.2	1,458.8
1999.....	20.8	38.8	61.3	92.7	149.4	204.3	366.7	559.0	818.8	1,448.3
Black, female										
2006 4/.....	16.2	25.6	38.1	49.9	77.4	116.7	188.3	319.0	546.2	1,060.5
2005 4/.....	16.8	29.3	41.8	50.3	73.0	115.6	200.4	351.5	587.6	1,156.5
2004 4/.....	17.5	28.1	40.8	51.2	83.6	130.1	211.8	387.6	632.4	1,270.7
2003 4/.....	18.5	29.0	44.4	57.3	90.7	137.1	221.8	391.8	650.3	1,414.2
2002.....	17.9	30.1	42.4	56.5	87.3	136.9	234.8	414.1	703.4	1,434.3
2001.....	19.0	29.1	47.4	56.7	96.5	148.5	238.6	421.9	690.1	1,491.2
2000.....	19.0	33.9	43.4	60.6	95.4	146.7	242.0	433.2	722.2	1,556.5
1999.....	19.0	30.5	41.6	66.6	93.7	144.5	268.7	445.1	725.6	1,509.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Subarachnoid hemorrhage (I60)										
All races 3/, both sexes										
2006.....	1.9	2.6	3.2	3.4	3.9	4.0	5.2	7.4	9.3	12.0
2005.....	2.1	2.9	3.3	3.5	3.9	4.6	5.6	7.2	9.7	12.2
2004.....	2.0	2.8	3.4	3.7	4.3	4.7	5.9	7.7	9.2	12.2
2003.....	2.1	2.6	3.6	3.8	4.3	5.0	5.8	7.7	10.1	11.5
2002.....	2.3	3.0	3.1	3.8	4.8	5.5	6.2	7.9	10.3	12.2
2001.....	2.2	3.0	3.4	4.2	4.3	5.3	6.3	8.0	9.8	13.2
2000.....	2.5	3.3	3.8	4.7	5.0	5.2	6.4	8.6	10.7	14.1
1999.....	2.6	3.3	3.7	4.4	5.1	5.7	6.7	8.4	10.9	13.1
All races 3/, male										
2006.....	1.5	2.2	2.6	2.7	2.8	2.9	3.7	5.9	7.0	8.9
2005.....	1.7	2.1	2.7	2.9	3.1	3.4	4.1	5.2	6.9	10.2
2004.....	1.5	2.4	3.0	2.8	3.3	3.2	4.2	5.7	6.5	9.3
2003.....	1.6	2.1	2.9	2.8	3.4	4.0	4.0	5.4	7.4	9.1
2002.....	1.8	2.4	2.2	3.0	3.8	4.2	4.6	5.6	6.1	9.1
2001.....	1.9	2.4	2.8	3.2	3.1	3.5	5.2	5.7	7.5	9.5
2000.....	2.0	2.6	2.9	3.4	3.6	3.4	5.2	6.3	7.7	11.2
1999.....	2.2	2.7	2.9	3.3	3.6	4.2	4.4	6.1	7.7	9.0
All races 3/, female										
2006.....	2.3	3.0	3.7	4.1	4.9	5.0	6.4	8.5	10.7	13.4
2005.....	2.4	3.7	3.9	4.1	4.7	5.7	6.8	8.7	11.5	13.1
2004.....	2.6	3.3	3.7	4.5	5.2	6.1	7.4	9.2	10.9	13.5
2003.....	2.5	3.2	4.3	4.7	5.1	5.8	7.2	9.3	11.8	12.5
2002.....	2.7	3.6	4.0	4.6	5.6	6.6	7.4	9.6	12.9	13.6
2001.....	2.6	3.6	3.9	5.2	5.3	6.8	7.2	9.7	11.2	14.8
2000.....	2.9	4.0	4.7	6.0	6.4	6.7	7.4	10.3	12.4	15.3
1999.....	3.0	3.9	4.4	5.5	6.4	6.9	8.5	10.0	12.8	14.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Subarachnoid hemorrhage (I60)										
White, both sexes										
2006 4/.....	1.6	2.4	2.8	3.2	3.7	3.9	5.0	7.3	9.1	11.8
2005 4/.....	1.8	2.6	3.0	3.4	3.7	4.6	5.3	7.2	9.5	12.4
2004 4/.....	1.8	2.6	3.0	3.5	4.0	4.6	5.8	7.4	9.1	12.4
2003 4/.....	1.8	2.3	3.3	3.6	4.2	4.6	5.6	7.3	9.7	11.1
2002.....	2.0	2.8	2.8	3.5	4.6	5.3	5.7	7.7	10.1	12.0
2001.....	1.9	2.7	2.9	3.8	4.2	4.8	6.1	7.7	9.9	12.7
2000.....	2.1	2.9	3.5	4.6	4.8	4.8	6.0	8.4	10.4	13.6
1999.....	2.3	2.9	3.4	4.2	4.7	5.6	6.5	8.3	10.7	12.6
White, male										
2006 4/.....	1.3	2.1	2.4	2.6	2.6	2.7	3.6	6.0	6.9	9.0
2005 4/.....	1.6	1.9	2.4	2.9	2.8	3.2	3.9	5.2	6.9	10.7
2004 4/.....	1.4	2.1	2.7	2.6	3.0	3.1	4.2	5.5	6.5	9.9
2003 4/.....	1.4	1.9	2.7	2.6	3.2	3.7	4.2	5.1	7.2	8.6
2002.....	1.7	2.4	2.0	2.7	3.6	4.1	4.3	5.6	5.9	9.1
2001.....	1.7	2.1	2.5	2.8	3.1	3.3	5.0	5.5	7.9	9.2
2000.....	1.8	2.4	2.7	3.3	3.3	3.2	4.8	6.3	7.8	11.8
1999.....	2.1	2.5	2.7	3.2	3.4	4.0	4.4	6.3	7.6	9.0
White, female										
2006 4/.....	1.9	2.6	3.3	3.7	4.7	5.0	6.1	8.3	10.4	13.2
2005 4/.....	1.9	3.2	3.5	3.9	4.5	5.9	6.5	8.7	11.1	13.1
2004 4/.....	2.2	3.0	3.4	4.4	4.9	5.9	7.1	8.7	10.8	13.5
2003 4/.....	2.1	2.7	3.9	4.6	5.0	5.4	6.8	8.9	11.3	12.2
2002.....	2.4	3.1	3.6	4.3	5.5	6.4	6.8	9.3	12.7	13.3
2001.....	2.2	3.2	3.4	4.8	5.3	6.1	6.9	9.3	11.1	14.2
2000.....	2.5	3.4	4.3	5.9	6.2	6.1	7.0	10.0	12.0	14.3
1999.....	2.5	3.3	4.0	5.1	5.9	7.0	8.3	9.7	12.5	14.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Subarachnoid hemorrhage (I60)										
Black, both sexes										
2006 4/.....	3.5	4.0	5.6	4.9	5.0	5.3	6.6	7.2	10.6	11.4
2005 4/.....	4.1	5.2	5.7	4.7	5.3	4.4	7.2	7.3	10.7	10.2
2004 4/.....	3.9	4.6	5.3	5.2	6.8	4.9	5.8	8.0	10.1	9.0
2003 4/.....	3.8	4.4	5.7	5.3	4.8	6.6	5.4	8.7	11.7	14.6
2002.....	4.1	5.3	5.5	6.0	5.6	6.2	6.7	8.2	9.2	12.8
2001.....	4.4	5.1	6.5	7.2	4.8	8.2	6.3	9.7	6.8	13.8
2000.....	4.7	6.1	6.2	6.2	6.2	7.4	8.7	11.0	10.4	17.2
1999.....	5.0	6.0	6.3	6.7	6.7	5.5	8.7	7.5	12.4	14.7
Black, male										
2006 4/.....	2.3	2.7	4.3	3.7	4.1	5.6	*	*	*	*
2005 4/.....	2.7	3.1	4.4	3.7	5.3	*	7.0	*	*	*
2004 4/.....	2.5	3.7	4.6	4.5	6.0	*	*	*	*	*
2003 4/.....	2.8	2.8	3.9	5.0	5.8	5.7	*	*	*	*
2002.....	2.7	3.1	3.4	5.2	5.1	5.7	6.7	*	*	*
2001.....	3.4	3.9	5.3	5.8	4.1	*	*	*	*	*
2000.....	3.6	4.2	4.6	3.8	4.8	*	8.4	*	*	*
1999.....	3.5	4.5	5.3	5.1	4.9	5.7	*	*	*	*
Black, female										
2006 4/.....	4.5	5.2	6.6	6.0	5.8	5.0	8.5	8.3	13.0	12.4
2005 4/.....	5.3	7.0	6.8	5.5	5.4	4.6	7.4	8.3	13.7	12.0
2004 4/.....	5.1	5.4	6.0	5.9	7.4	5.8	7.0	9.7	12.1	11.8
2003 4/.....	4.8	5.8	7.2	5.5	4.1	7.3	7.8	10.3	12.8	15.0
2002.....	5.2	7.2	7.3	6.6	6.0	6.5	6.7	9.9	10.7	15.6
2001.....	5.3	6.1	7.6	8.3	5.4	10.9	7.2	10.8	8.6	16.4
2000.....	5.6	7.8	7.5	8.1	7.4	9.4	8.9	12.8	13.2	22.8
1999.....	6.4	7.3	7.2	8.0	8.2	5.4	10.8	10.3	13.0	18.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intracerebral and other intracranial hemorrhage (I61-I62)										
All races 3/, both sexes										
2006.....	2.7	4.6	7.9	9.9	13.1	18.7	31.3	52.9	85.0	120.8
2005.....	2.7	4.8	7.8	9.8	13.5	20.5	33.3	56.0	89.1	125.2
2004.....	2.9	5.1	7.5	9.8	13.9	21.2	34.1	57.5	89.6	124.3
2003.....	3.1	5.2	7.6	9.5	13.9	20.4	34.9	57.7	89.5	126.6
2002.....	2.8	5.0	7.2	9.7	14.3	20.9	36.0	59.1	90.6	127.2
2001.....	3.1	5.2	7.4	9.5	14.8	21.2	35.2	58.2	91.1	127.9
2000.....	3.2	5.6	7.6	10.6	15.9	23.5	35.5	60.1	95.1	137.9
1999.....	2.8	4.9	7.1	10.7	14.3	20.7	36.4	59.5	90.5	122.3
All races 3/, male										
2006.....	3.0	5.4	9.7	11.9	15.5	21.7	35.5	57.9	92.0	130.5
2005.....	3.1	5.6	9.4	12.2	16.0	24.2	38.5	63.2	96.6	133.7
2004.....	3.5	6.0	9.1	11.7	16.8	24.9	37.7	63.2	98.2	134.4
2003.....	3.4	6.1	9.4	11.0	16.4	23.6	39.9	63.2	99.8	137.4
2002.....	3.2	6.0	8.8	11.0	17.3	24.7	41.0	63.4	98.4	135.0
2001.....	3.4	6.3	8.8	11.2	17.3	23.7	41.5	65.5	99.5	134.2
2000.....	3.6	6.8	8.9	13.4	19.0	27.0	40.0	65.9	100.6	149.6
1999.....	3.4	5.6	8.6	12.7	17.3	23.5	41.7	65.1	94.2	130.7
All races 3/, female										
2006.....	2.4	3.9	6.1	8.1	10.9	16.0	27.8	49.2	80.7	116.3
2005.....	2.4	4.2	6.3	7.5	11.2	17.4	29.1	50.8	84.4	121.3
2004.....	2.3	4.2	6.1	8.0	11.3	18.0	31.2	53.4	84.3	119.8
2003.....	2.7	4.4	5.8	8.0	11.7	17.7	30.9	53.8	83.3	121.8
2002.....	2.4	3.9	5.7	8.4	11.7	17.6	32.1	56.1	85.9	123.8
2001.....	2.7	4.1	6.2	7.8	12.6	19.1	30.2	53.1	86.2	125.2
2000.....	2.8	4.4	6.4	7.9	13.0	20.4	32.0	56.1	91.8	133.1
1999.....	2.3	4.2	5.6	8.8	11.6	18.2	32.3	55.6	88.4	118.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intracerebral and other intracranial hemorrhage (I61-I62)										
White, both sexes										
2006 4/.....	2.0	3.4	5.7	7.4	11.2	16.7	29.9	52.7	85.1	122.9
2005 4/.....	2.0	3.4	5.6	7.5	11.4	18.8	32.1	55.6	89.5	127.1
2004 4/.....	2.1	3.5	5.3	7.6	11.9	19.4	32.9	56.7	89.8	126.5
2003 4/.....	2.1	3.6	5.0	7.3	11.3	18.5	33.6	57.1	89.5	127.7
2002.....	1.8	3.4	5.0	7.8	12.5	18.8	34.9	58.7	91.3	129.2
2001.....	2.2	3.5	5.1	7.8	12.5	19.1	33.7	57.8	91.7	128.9
2000.....	2.2	3.7	5.6	8.5	13.7	21.7	34.7	59.4	95.6	139.4
1999.....	1.9	3.3	5.1	8.5	12.2	19.4	35.4	58.9	90.7	123.2
White, male										
2006 4/.....	2.2	4.0	7.1	8.5	12.8	19.6	33.7	57.7	93.6	132.9
2005 4/.....	2.2	4.1	6.7	9.2	13.4	22.1	37.0	63.5	96.8	137.5
2004 4/.....	2.7	4.2	6.6	9.0	14.1	22.9	36.1	62.3	99.1	137.6
2003 4/.....	2.5	4.4	6.3	8.3	13.2	21.7	38.0	62.7	100.0	138.9
2002.....	2.1	4.2	6.1	8.7	15.1	22.2	40.0	63.5	99.5	136.9
2001.....	2.5	4.5	6.0	9.2	14.9	21.2	39.7	65.1	100.4	134.9
2000.....	2.5	4.7	6.6	10.7	16.1	24.5	38.9	65.5	101.3	150.5
1999.....	2.3	3.7	6.2	10.1	14.8	22.2	40.7	64.0	93.7	131.7
White, female										
2006 4/.....	1.7	2.8	4.4	6.4	9.6	14.0	26.7	49.1	79.7	118.3
2005 4/.....	1.7	2.8	4.5	5.9	9.6	15.8	28.1	49.8	85.0	122.3
2004 4/.....	1.5	2.8	4.1	6.4	9.9	16.3	30.3	52.5	84.0	121.5
2003 4/.....	1.7	2.8	3.8	6.2	9.6	15.7	30.1	53.0	83.0	122.8
2002.....	1.5	2.6	4.0	7.0	10.1	15.8	30.7	55.2	86.2	125.9
2001.....	1.9	2.6	4.1	6.5	10.2	17.3	28.8	52.7	86.4	126.3
2000.....	1.8	2.7	4.6	6.4	11.4	19.2	31.3	55.1	92.2	134.9
1999.....	1.5	2.8	4.1	7.1	9.7	16.9	31.2	55.3	89.0	119.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intracerebral and other intracranial hemorrhage (I61-I62)										
Black, both sexes										
2006 4/.....	7.0	12.8	23.3	28.8	28.6	31.5	41.1	53.3	81.4	94.7
2005 4/.....	7.6	14.5	24.0	27.4	29.5	35.4	42.5	56.5	80.8	102.1
2004 4/.....	7.9	15.9	22.9	26.1	31.4	35.6	44.6	61.1	88.4	95.6
2003 4/.....	8.9	16.4	25.8	27.3	34.1	33.8	43.4	64.2	86.1	107.4
2002.....	8.8	15.7	22.9	24.1	28.1	34.8	44.6	61.7	81.1	98.3
2001.....	8.7	16.6	25.5	22.6	32.0	37.0	43.7	59.2	83.2	106.6
2000.....	10.0	19.0	23.4	27.5	32.3	36.1	42.4	63.3	78.2	117.0
1999.....	9.4	16.1	22.3	26.6	29.6	30.5	44.4	62.5	77.4	96.9
Black, male										
2006 4/.....	7.7	14.3	30.4	39.1	39.8	36.3	52.0	58.6	78.0	93.8
2005 4/.....	8.9	16.7	31.0	37.0	40.5	43.7	52.8	57.2	91.3	93.0
2004 4/.....	8.6	19.0	27.1	35.1	42.0	42.6	53.1	64.1	96.1	100.3
2003 4/.....	9.2	18.7	32.9	34.9	44.2	37.7	54.5	75.3	98.9	116.8
2002.....	10.4	19.3	30.0	30.8	36.0	43.4	52.4	61.5	81.0	101.8
2001.....	9.3	20.3	31.3	29.1	36.4	46.0	54.2	62.7	85.9	112.3
2000.....	11.6	23.4	28.5	38.4	43.5	46.3	47.1	69.9	78.4	130.3
1999.....	11.6	19.2	28.6	33.2	37.8	34.0	48.2	70.8	81.1	109.0
Black, female										
2006 4/.....	6.3	11.5	17.3	20.2	20.0	27.9	33.6	50.1	83.2	95.0
2005 4/.....	6.5	12.6	18.0	19.4	20.8	29.3	35.5	56.0	75.3	105.8
2004 4/.....	7.2	13.2	19.3	18.7	23.0	30.4	38.9	59.3	84.3	93.7
2003 4/.....	8.6	14.3	19.7	21.1	26.2	30.9	35.9	57.5	79.3	103.7
2002.....	7.4	12.5	16.9	18.6	21.8	28.5	39.4	61.7	81.2	96.9
2001.....	8.1	13.4	20.6	17.2	28.5	30.4	36.8	57.1	81.8	104.4
2000.....	8.5	15.2	19.1	18.6	23.5	28.6	39.3	59.2	78.1	112.0
1999.....	7.4	13.3	16.9	21.2	23.1	27.9	41.9	57.5	75.6	92.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebral infarction (I63)										
All races 3/, both sexes										
2006.....	0.4	0.6	1.0	1.4	2.6	4.0	6.6	12.8	23.0	50.4
2005.....	0.4	0.6	1.0	1.6	2.8	4.5	7.5	13.0	25.3	56.8
2004.....	0.4	0.6	1.0	1.8	3.1	5.5	10.3	20.3	37.8	97.7
2003.....	0.4	0.8	1.1	1.8	3.0	6.2	10.9	21.6	44.5	112.2
2002.....	0.3	0.7	1.3	1.7	3.5	5.8	11.6	22.6	45.6	118.3
2001.....	0.4	0.8	1.1	1.9	3.1	6.1	13.0	24.3	47.6	124.8
2000.....	0.5	0.7	1.2	2.0	4.0	6.6	12.8	27.2	54.5	141.0
1999.....	0.5	0.7	1.2	2.4	4.0	8.1	14.9	29.7	58.5	153.1
All races 3/, male										
2006.....	0.5	0.7	1.2	1.9	3.2	4.8	8.0	14.4	24.0	45.8
2005.....	0.4	0.8	1.2	2.0	3.6	5.4	8.1	12.8	26.7	51.9
2004.....	0.4	0.7	1.4	2.3	4.1	6.8	11.6	23.5	39.7	88.5
2003.....	0.5	1.0	1.3	2.2	4.0	7.6	13.2	23.4	47.7	100.3
2002.....	0.3	0.9	1.6	2.3	4.2	7.2	12.4	25.8	50.1	108.7
2001.....	0.4	0.9	1.2	2.6	3.8	7.7	14.9	28.0	53.3	111.2
2000.....	0.5	0.9	1.4	2.5	5.2	7.8	15.4	31.2	60.8	133.3
1999.....	0.5	0.9	1.5	2.9	4.8	10.0	17.4	34.0	63.3	144.6
All races 3/, female										
2006.....	0.4	0.5	0.7	1.0	2.0	3.2	5.5	11.6	22.4	52.5
2005.....	0.4	0.4	0.7	1.2	2.0	3.7	7.0	13.2	24.4	59.1
2004.....	0.4	0.6	0.6	1.3	2.2	4.3	9.2	18.0	36.7	101.9
2003.....	0.3	0.5	0.9	1.5	2.1	5.1	9.0	20.4	42.6	117.5
2002.....	0.3	0.5	0.9	1.1	2.9	4.6	11.0	20.4	42.8	122.4
2001.....	0.3	0.7	0.9	1.3	2.5	4.6	11.4	21.8	44.2	130.4
2000.....	0.4	0.5	1.1	1.6	3.0	5.6	10.7	24.4	50.7	144.2
1999.....	0.4	0.5	0.9	1.8	3.3	6.4	12.9	26.7	55.7	156.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebral infarction (I63)										
White, both sexes										
2006 4/.....	0.4	0.5	0.8	1.3	2.3	3.7	6.4	12.6	22.8	51.5
2005 4/.....	0.4	0.4	0.9	1.3	2.4	4.0	7.2	12.3	25.6	57.4
2004 4/.....	0.3	0.5	0.8	1.5	2.6	5.0	9.6	19.5	37.5	97.6
2003 4/.....	0.3	0.6	0.9	1.5	2.5	5.9	10.1	21.0	44.4	113.7
2002.....	0.3	0.6	1.0	1.5	3.2	5.3	10.9	22.1	44.8	119.1
2001.....	0.3	0.6	0.9	1.6	2.7	5.4	12.3	23.6	47.5	126.0
2000.....	0.3	0.6	1.0	1.6	3.7	6.0	12.0	26.3	54.0	141.7
1999.....	0.4	0.6	1.0	2.0	3.5	7.5	13.8	28.5	58.6	154.4
White, male										
2006 4/.....	0.4	0.5	1.0	1.7	2.8	4.5	7.9	13.8	23.6	46.6
2005 4/.....	0.4	0.6	1.1	1.7	3.0	4.9	7.7	12.6	27.5	51.4
2004 4/.....	0.4	0.6	1.1	2.0	3.3	6.3	10.6	22.5	39.7	87.9
2003 4/.....	0.4	0.9	1.2	1.8	3.4	7.2	12.2	22.4	48.5	101.5
2002.....	0.3	0.9	1.3	2.0	3.7	6.5	11.7	25.3	49.3	108.6
2001.....	0.4	0.7	1.1	2.2	3.4	7.1	14.3	26.8	52.9	110.5
2000.....	0.4	0.8	1.0	2.0	4.7	7.0	14.6	29.8	59.8	133.1
1999.....	0.4	0.7	1.3	2.6	4.2	9.5	15.9	32.3	63.5	144.0
White, female										
2006 4/.....	0.3	0.5	0.6	0.9	1.8	3.0	5.2	11.7	22.3	53.8
2005 4/.....	0.3	0.3	0.7	1.0	1.8	3.3	6.8	12.2	24.4	60.2
2004 4/.....	0.3	0.5	0.5	1.1	2.0	3.9	8.7	17.3	36.2	102.0
2003 4/.....	0.2	0.3	0.6	1.3	1.7	4.7	8.5	19.9	41.8	119.1
2002.....	0.3	0.4	0.8	1.0	2.7	4.2	10.3	19.8	42.1	123.7
2001.....	0.2	0.5	0.8	1.1	2.1	4.0	10.7	21.3	44.2	132.5
2000.....	0.3	0.4	0.9	1.2	2.7	5.2	10.0	23.8	50.5	145.2
1999.....	0.3	0.5	0.8	1.5	2.9	5.7	12.1	25.9	55.7	158.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cerebral infarction (I63)										
Black, both sexes										
2006 4/.....	0.8	1.5	2.5	2.9	5.5	6.6	9.5	16.1	29.8	45.1
2005 4/.....	0.8	1.7	2.0	3.5	6.3	8.3	10.5	21.5	26.0	55.9
2004 4/.....	1.0	1.5	2.2	4.1	7.4	9.5	19.1	33.2	43.8	103.5
2003 4/.....	0.9	1.6	2.6	4.7	7.7	10.6	18.8	32.3	51.8	107.7
2002.....	*	1.3	3.1	3.6	6.7	11.0	20.0	32.3	56.8	113.1
2001.....	0.9	2.0	2.2	4.4	7.3	12.3	21.7	34.9	53.4	121.3
2000.....	1.2	1.3	3.1	5.4	7.8	13.0	21.1	37.7	67.5	142.6
1999.....	1.1	1.8	2.7	5.0	8.7	14.4	27.7	47.0	62.7	150.8
Black, male										
2006 4/.....	*	2.1	3.7	4.1	7.1	8.0	10.5	21.0	32.2	42.0
2005 4/.....	*	2.1	2.7	4.4	9.3	10.7	13.0	18.3	22.5	63.2
2004 4/.....	*	2.0	3.1	5.1	11.2	13.6	22.7	41.9	45.9	97.4
2003 4/.....	*	2.1	2.8	6.0	10.8	13.0	25.0	38.8	50.2	97.0
2002.....	*	*	4.4	4.9	8.9	14.7	22.7	35.8	63.4	114.7
2001.....	*	2.4	2.6	6.6	8.3	15.8	23.2	47.4	65.0	127.1
2000.....	*	*	4.5	7.1	10.1	17.7	24.7	46.8	83.5	137.2
1999.....	*	2.7	3.1	6.3	10.8	17.1	37.7	60.3	68.9	162.9
Black, female										
2006 4/.....	*	*	*	*	4.3	5.5	8.9	13.1	28.6	46.4
2005 4/.....	*	*	*	2.7	4.0	6.5	8.9	23.5	27.8	52.9
2004 4/.....	*	*	*	3.1	4.4	6.5	16.7	28.0	42.7	105.9
2003 4/.....	*	*	2.5	3.7	5.3	8.8	14.6	28.5	52.6	111.8
2002.....	*	*	2.0	2.4	5.0	8.3	18.1	30.2	53.3	112.5
2001.....	*	1.7	1.9	2.7	6.4	9.8	20.6	27.5	47.4	119.1
2000.....	1.3	*	*	4.0	5.9	9.6	18.6	32.2	59.4	144.7
1999.....	*	*	2.4	3.9	7.0	12.3	21.2	38.9	59.6	146.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Stroke, not specified as hemorrhage or infarction (I64)										
All races 3/, both sexes										
2006.....	1.3	2.6	5.0	9.2	18.1	34.0	64.1	126.8	257.2	628.5
2005.....	1.3	2.5	5.2	8.6	17.3	33.8	65.4	132.4	271.7	675.7
2004.....	1.3	2.6	4.9	8.6	17.9	36.0	70.1	141.8	288.1	713.4
2003.....	1.5	2.4	5.1	9.9	19.5	36.9	75.1	155.8	315.4	799.7
2002.....	1.2	2.8	5.3	10.7	20.5	40.3	82.7	165.7	338.0	859.4
2001.....	1.2	2.5	5.2	10.5	22.0	41.7	85.7	176.0	355.4	906.2
2000.....	1.2	2.6	5.2	10.7	23.0	44.6	89.8	181.1	372.2	954.7
1999.....	1.1	2.7	5.0	11.3	22.4	45.1	89.6	186.1	380.9	982.4
All races 3/, male										
2006.....	1.5	2.9	6.2	11.8	21.6	40.6	70.8	135.4	255.2	539.0
2005.....	1.4	2.9	6.6	10.5	21.4	39.4	74.1	145.0	274.5	584.7
2004.....	1.5	3.0	6.1	10.0	21.4	41.6	80.0	152.1	292.2	615.5
2003.....	1.7	2.9	6.1	12.1	23.3	43.1	85.5	171.8	321.2	699.0
2002.....	1.2	3.2	6.1	13.2	24.3	47.5	93.1	179.9	334.5	762.8
2001.....	1.3	3.0	6.2	12.6	26.5	49.4	97.9	188.5	363.9	819.7
2000.....	1.3	3.1	6.5	12.8	27.8	51.0	102.3	198.1	390.9	863.2
1999.....	1.3	3.2	6.2	13.1	27.2	51.7	103.2	206.0	404.7	896.4
All races 3/, female										
2006.....	1.1	2.2	3.9	6.8	15.0	28.3	58.6	120.4	258.5	670.4
2005.....	1.1	2.2	3.9	6.8	13.6	28.9	58.3	123.2	270.0	717.4
2004.....	1.1	2.2	3.8	7.3	14.8	31.1	62.2	134.3	285.5	757.4
2003.....	1.2	2.0	4.1	7.9	16.1	31.6	66.7	144.4	311.9	844.2
2002.....	1.2	2.4	4.4	8.4	16.9	34.1	74.3	155.8	340.2	901.3
2001.....	1.1	2.0	4.2	8.6	17.8	35.1	76.1	167.2	350.3	942.4
2000.....	1.1	2.2	3.9	8.6	18.7	39.2	80.0	169.2	361.2	992.0
1999.....	0.9	2.2	3.9	9.6	18.2	39.4	78.9	172.4	367.0	1,017.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Stroke, not specified as hemorrhage or infarction (I64)										
White, both sexes										
2006 4/.....	1.0	2.0	4.0	7.4	14.9	29.4	57.8	120.9	253.9	635.9
2005 4/.....	1.0	1.9	4.0	6.9	14.3	29.4	59.2	126.9	267.3	682.2
2004 4/.....	1.0	1.9	3.7	6.9	14.7	31.1	64.6	134.4	284.2	721.0
2003 4/.....	1.0	1.8	3.9	8.0	16.0	32.2	69.0	148.7	312.5	807.1
2002.....	1.0	2.1	4.0	8.6	16.6	34.9	76.1	159.4	333.5	867.3
2001.....	0.9	1.9	3.9	8.5	17.8	35.8	78.8	169.6	350.8	913.6
2000.....	1.0	1.9	4.1	8.4	19.4	39.5	83.1	174.3	366.9	963.5
1999.....	0.8	1.9	3.8	9.0	18.4	39.9	82.3	178.9	377.1	993.4
White, male										
2006 4/.....	1.2	2.3	4.8	9.3	17.6	34.9	63.1	129.0	251.5	543.1
2005 4/.....	1.1	2.3	5.2	8.3	17.7	33.8	67.0	138.6	268.4	589.5
2004 4/.....	1.1	2.3	4.7	7.9	17.6	36.1	73.3	144.3	288.8	622.4
2003 4/.....	1.2	2.2	4.7	9.5	19.5	37.9	78.3	164.2	317.7	704.0
2002.....	1.1	2.5	4.7	10.6	19.6	40.6	85.5	172.6	329.7	766.7
2001.....	1.1	2.3	4.7	10.0	21.2	42.3	89.6	181.7	357.5	823.5
2000.....	1.1	2.2	5.3	10.1	23.6	45.2	94.2	189.8	384.8	865.2
1999.....	1.0	2.3	4.8	10.7	22.3	45.4	95.3	197.4	401.4	903.1
White, female										
2006 4/.....	0.8	1.7	3.2	5.5	12.3	24.5	53.4	114.8	255.5	679.5
2005 4/.....	0.9	1.6	2.8	5.5	11.1	25.5	52.8	118.3	266.6	724.9
2004 4/.....	0.8	1.6	2.7	6.0	12.1	26.7	57.5	127.1	281.3	765.4
2003 4/.....	0.8	1.5	3.0	6.4	12.8	27.1	61.3	137.4	309.3	852.8
2002.....	0.9	1.6	3.3	6.7	13.8	30.0	68.4	149.9	335.9	910.9
2001.....	0.7	1.4	3.1	7.0	14.6	30.1	70.1	161.0	346.7	951.3
2000.....	0.8	1.5	2.9	6.8	15.5	34.6	74.2	163.4	356.3	1,003.5
1999.....	0.6	1.5	2.9	7.4	14.9	35.1	72.0	165.9	362.8	1,029.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Stroke, not specified as hemorrhage or infarction (I64)										
Black, both sexes										
2006 4/.....	3.6	6.9	13.3	24.9	49.1	78.1	128.3	204.2	323.3	612.5
2005 4/.....	3.6	7.1	15.0	24.0	45.6	73.1	128.5	206.8	351.0	669.0
2004 4/.....	3.7	7.4	14.4	24.1	48.5	82.1	132.2	236.6	367.9	706.1
2003 4/.....	4.3	7.1	14.9	27.8	52.6	81.9	142.9	250.5	386.3	797.6
2002.....	3.2	8.0	15.7	30.9	55.6	91.8	156.7	259.7	425.2	840.7
2001.....	3.6	7.1	16.3	29.3	59.5	96.7	163.1	268.9	435.0	883.4
2000.....	2.9	8.6	14.7	31.0	55.2	97.3	167.8	275.6	473.0	919.4
1999.....	3.4	8.5	16.0	32.5	59.9	97.2	174.4	287.1	454.8	917.6
Black, male										
2006 4/.....	3.7	8.0	18.2	34.3	62.4	98.4	157.2	235.8	336.8	521.0
2005 4/.....	4.3	8.2	18.2	32.0	60.7	95.3	154.0	237.4	377.1	581.0
2004 4/.....	4.1	8.5	17.2	30.9	61.5	98.6	159.6	257.5	375.9	589.1
2003 4/.....	5.3	8.1	18.3	36.2	63.0	95.5	170.8	285.3	410.5	681.9
2002.....	2.5	8.9	18.8	40.0	71.1	118.4	194.6	293.9	428.7	761.2
2001.....	3.6	8.2	19.8	37.9	77.9	122.7	199.6	302.9	470.3	816.9
2000.....	3.0	10.0	18.6	39.0	66.7	117.1	203.6	321.8	519.3	897.2
1999.....	3.8	10.1	19.8	37.8	77.2	120.2	202.2	328.6	489.2	877.6
Black, female										
2006 4/.....	3.5	6.0	9.2	17.2	38.7	63.2	108.4	185.3	316.1	650.0
2005 4/.....	3.0	6.2	12.3	17.5	33.9	56.6	111.0	188.5	337.2	704.5
2004 4/.....	3.3	6.4	12.0	18.5	38.3	69.8	113.6	224.3	363.6	752.5
2003 4/.....	3.5	6.2	12.0	20.9	44.4	71.8	124.1	229.8	373.7	842.6
2002.....	3.8	7.2	13.0	23.5	43.5	72.0	131.3	239.3	423.3	871.2
2001.....	3.7	6.1	13.3	22.2	45.0	77.5	138.8	248.5	417.0	908.1
2000.....	2.8	7.4	11.4	24.4	46.1	82.8	144.1	247.8	449.6	927.6
1999.....	3.1	7.2	12.7	28.1	46.3	80.2	156.0	262.0	437.8	932.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other cerebrovascular diseases and their sequelae (I67,I69)										
All races 3/, both sexes										
2006.....	0.4	0.8	1.5	2.4	5.1	10.0	20.3	40.5	84.0	228.0
2005.....	0.4	0.8	1.4	2.4	5.2	10.3	21.9	45.0	101.5	271.8
2004.....	0.5	0.7	1.4	2.6	5.2	11.0	21.7	49.4	107.4	298.3
2003.....	0.4	0.8	1.3	2.8	5.1	11.1	24.0	51.3	111.4	320.1
2002.....	0.5	0.8	1.5	2.7	5.2	11.3	24.2	53.4	117.2	328.9
2001.....	0.4	0.7	1.4	2.7	5.4	11.4	24.1	53.6	118.5	328.2
2000.....	0.4	0.8	1.4	2.8	5.8	11.0	24.6	55.7	121.5	341.4
1999.....	0.4	0.8	1.4	2.9	5.6	11.7	25.8	56.9	123.1	343.9
All races 3/, male										
2006.....	0.4	0.8	1.6	2.9	6.3	11.9	22.9	45.9	90.4	208.2
2005.....	0.3	0.8	1.5	2.6	5.9	12.2	24.7	49.6	108.3	242.8
2004.....	0.5	0.7	1.6	3.1	6.5	13.7	25.4	56.8	115.0	270.4
2003.....	0.3	0.9	1.4	3.2	6.3	13.2	28.1	58.7	119.4	290.2
2002.....	0.4	0.8	1.7	3.2	6.2	13.3	27.6	60.2	126.5	302.3
2001.....	0.4	0.7	1.5	3.2	6.5	14.2	28.3	61.4	130.6	305.5
2000.....	0.4	0.8	1.6	3.1	6.8	13.3	30.1	66.1	135.0	326.9
1999.....	0.4	0.8	1.5	3.6	7.1	13.4	31.0	65.8	136.9	332.1
All races 3/, female										
2006.....	0.4	0.8	1.5	2.0	4.0	8.3	18.3	36.5	80.0	237.2
2005.....	0.5	0.9	1.3	2.3	4.6	8.6	19.7	41.7	97.2	285.2
2004.....	0.4	0.7	1.3	2.2	4.1	8.7	18.8	44.1	102.8	310.8
2003.....	0.5	0.7	1.2	2.4	4.0	9.3	20.8	46.0	106.6	333.3
2002.....	0.6	0.8	1.2	2.4	4.3	9.5	21.5	48.6	111.5	340.4
2001.....	0.4	0.7	1.2	2.3	4.4	9.0	20.7	48.2	111.3	337.6
2000.....	0.4	0.9	1.3	2.4	5.0	9.0	20.3	48.5	113.6	347.4
1999.....	0.5	0.9	1.2	2.3	4.2	10.2	21.8	50.7	115.1	348.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other cerebrovascular diseases and their sequelae (I67,I69)										
White, both sexes										
2006 4/.....	0.3	0.7	1.2	2.0	4.3	8.7	18.7	38.1	82.1	228.7
2005 4/.....	0.3	0.7	1.1	2.1	4.4	8.8	19.9	42.1	99.3	274.3
2004 4/.....	0.4	0.6	1.2	2.2	4.3	9.8	19.7	46.9	105.6	300.3
2003 4/.....	0.3	0.6	1.0	2.3	4.2	9.6	22.0	49.2	109.8	322.5
2002.....	0.4	0.6	1.2	2.4	4.3	9.8	22.4	50.7	115.9	332.0
2001.....	0.3	0.6	1.0	2.2	4.5	10.0	22.3	51.0	117.3	329.9
2000.....	0.3	0.7	1.2	2.3	5.0	9.9	23.1	52.9	120.2	343.4
1999.....	0.4	0.7	1.1	2.4	4.7	10.5	23.3	54.7	121.3	348.1
White, male										
2006 4/.....	0.4	0.7	1.2	2.4	5.1	10.0	20.6	43.3	88.8	209.8
2005 4/.....	0.2	0.7	1.2	2.2	4.8	10.2	22.5	46.6	105.7	244.6
2004 4/.....	0.5	0.5	1.2	2.6	5.3	12.2	22.9	53.4	112.8	271.5
2003 4/.....	0.3	0.7	1.0	2.5	5.2	11.3	25.9	55.6	117.0	294.0
2002.....	0.3	0.7	1.4	2.7	5.1	11.7	25.8	56.5	125.7	306.3
2001.....	0.3	0.6	1.0	2.7	5.5	12.8	26.2	58.6	130.0	308.1
2000.....	0.3	0.7	1.3	2.5	6.2	11.8	27.9	63.3	132.5	330.0
1999.....	0.3	0.6	1.1	2.9	5.9	12.1	27.4	63.2	134.3	334.9
White, female										
2006 4/.....	0.3	0.7	1.2	1.6	3.5	7.6	17.1	34.2	77.9	237.5
2005 4/.....	0.3	0.7	1.0	2.0	4.0	7.4	17.8	38.7	95.2	288.0
2004 4/.....	0.4	0.6	1.1	1.8	3.4	7.7	17.1	42.2	101.1	313.3
2003 4/.....	0.4	0.5	1.0	2.0	3.3	8.2	18.8	44.5	105.3	335.2
2002.....	0.5	0.6	1.0	2.0	3.5	8.1	19.7	46.6	109.9	343.2
2001.....	0.3	0.6	0.9	1.8	3.6	7.6	19.2	45.6	109.6	339.1
2000.....	0.3	0.6	1.0	2.1	4.0	8.1	19.3	45.7	112.9	348.8
1999.....	0.4	0.7	1.1	1.9	3.6	9.2	20.2	48.7	113.6	353.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other cerebrovascular diseases and their sequelae (I67,I69)										
Black, both sexes										
2006 4/.....	1.0	1.9	4.0	6.1	13.2	21.8	36.2	68.5	112.6	244.9
2005 4/.....	1.0	2.1	3.7	5.9	12.7	24.0	42.7	80.2	142.5	269.7
2004 4/.....	0.7	2.0	3.1	6.9	14.1	22.8	44.9	79.7	137.6	296.9
2003 4/.....	0.7	2.0	3.7	7.2	13.9	24.5	45.8	77.6	141.1	321.5
2002.....	0.8	1.8	4.0	6.5	13.9	25.8	44.5	85.2	140.2	318.9
2001.....	0.9	1.7	4.4	7.2	13.6	24.0	42.8	87.5	138.1	327.6
2000.....	0.9	2.2	3.8	7.3	12.5	20.9	41.1	87.6	140.3	333.8
1999.....	0.9	2.0	3.3	7.6	13.3	22.3	52.6	83.9	149.1	312.5
Black, male										
2006 4/.....	*	1.8	4.5	7.7	18.9	30.9	46.8	79.2	126.6	216.3
2005 4/.....	*	1.9	4.2	6.6	17.6	31.1	50.0	88.5	159.4	240.8
2004 4/.....	*	2.1	4.3	9.1	18.7	29.8	58.6	102.4	152.8	272.2
2003 4/.....	*	2.4	4.6	8.4	17.9	33.0	55.2	97.7	158.7	271.1
2002.....	*	*	5.0	7.9	17.6	31.4	52.4	105.6	150.5	269.1
2001.....	*	*	4.9	8.3	16.7	29.5	54.2	103.6	143.7	285.9
2000.....	*	*	4.2	9.6	12.4	27.1	56.2	98.3	177.0	291.8
1999.....	*	2.3	4.6	10.3	18.7	27.3	73.3	96.4	168.3	294.1
Black, female										
2006 4/.....	*	1.9	3.5	4.7	8.7	15.1	29.0	62.1	105.3	256.6
2005 4/.....	*	2.2	3.2	5.3	8.9	18.7	37.7	75.2	133.5	281.3
2004 4/.....	*	1.9	2.2	5.0	10.5	17.6	35.6	66.3	129.7	306.8
2003 4/.....	*	1.6	3.0	6.1	10.8	18.3	39.4	65.7	131.9	341.1
2002.....	*	1.9	3.2	5.4	11.0	21.6	39.2	73.0	134.9	338.0
2001.....	*	1.8	4.1	6.3	11.2	19.9	35.2	77.9	135.3	343.1
2000.....	*	2.6	3.5	5.4	12.6	16.3	31.1	81.2	121.9	349.4
1999.....	*	1.7	2.3	5.4	9.2	18.7	38.9	76.4	139.6	319.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of arteries, arterioles and capillaries (I70-I78)										
All races 3/, both sexes										
2006.....	1.3	2.5	3.6	6.1	10.7	19.5	34.3	59.7	99.4	218.4
2005.....	1.4	2.3	3.8	6.4	12.3	22.0	38.4	67.8	115.7	268.4
2004.....	1.4	2.4	3.9	6.2	12.2	22.4	41.5	69.1	119.1	276.3
2003.....	1.4	2.7	4.2	6.8	13.8	25.0	43.2	76.4	126.2	300.6
2002.....	1.4	2.5	3.8	7.6	14.1	27.5	45.8	78.3	131.6	316.5
2001.....	1.4	2.3	3.9	7.4	14.6	27.3	46.5	82.5	137.2	337.3
2000.....	1.4	2.5	4.1	7.7	15.3	29.6	51.1	85.6	144.6	352.0
1999.....	1.5	2.4	4.0	8.0	15.4	28.6	51.3	87.6	147.8	367.2
All races 3/, male										
2006.....	1.9	3.4	5.2	8.8	14.6	26.5	44.3	76.1	122.0	240.2
2005.....	1.9	3.2	5.4	8.9	16.9	29.6	51.4	88.4	143.2	279.9
2004.....	1.8	3.4	5.4	8.4	16.6	29.6	54.4	87.3	149.4	294.7
2003.....	1.9	3.7	5.9	9.3	19.1	34.8	56.6	102.8	159.5	318.3
2002.....	2.0	3.6	5.2	10.3	19.6	38.1	60.7	102.7	166.1	344.0
2001.....	2.0	3.3	5.2	10.0	20.0	37.0	63.3	106.0	175.7	364.5
2000.....	1.9	3.3	5.8	11.0	21.6	41.2	70.6	116.1	186.4	391.4
1999.....	1.8	3.4	5.5	11.3	22.0	39.8	70.0	117.2	189.4	396.5
All races 3/, female										
2006.....	0.7	1.5	2.2	3.6	7.1	13.5	26.2	47.7	85.2	208.2
2005.....	1.0	1.3	2.4	3.9	8.1	15.3	27.8	52.9	98.6	263.1
2004.....	0.9	1.4	2.4	4.2	8.3	16.2	31.1	56.1	100.5	268.0
2003.....	0.9	1.8	2.5	4.4	8.9	16.4	32.4	57.7	105.8	292.8
2002.....	0.8	1.5	2.5	5.0	9.2	18.3	33.8	61.1	110.6	304.6
2001.....	0.9	1.4	2.7	5.0	9.7	18.9	33.1	66.0	114.2	325.9
2000.....	1.0	1.7	2.5	4.6	9.6	19.7	35.6	64.4	120.0	336.0
1999.....	1.2	1.5	2.7	5.0	9.5	19.1	36.7	67.0	123.5	355.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of arteries, arterioles and capillaries (I70-I78)										
White, both sexes										
2006 4/.....	1.1	2.2	3.1	5.6	10.4	19.1	34.6	60.3	99.4	222.9
2005 4/.....	1.3	2.0	3.3	5.9	12.0	21.6	38.6	68.3	116.7	274.8
2004 4/.....	1.2	2.1	3.5	5.8	11.6	22.1	42.3	69.9	120.5	281.3
2003 4/.....	1.2	2.4	3.7	6.5	13.4	25.0	43.6	77.7	126.9	306.4
2002.....	1.2	2.1	3.3	7.1	13.7	27.7	46.1	79.0	133.2	323.7
2001.....	1.2	2.1	3.5	7.0	14.0	27.5	46.6	82.8	138.7	344.1
2000.....	1.2	2.1	3.5	7.2	14.8	29.7	51.8	86.1	146.1	358.3
1999.....	1.3	2.1	3.7	7.6	15.4	28.7	52.1	88.3	149.4	372.8
White, male										
2006 4/.....	1.7	3.1	4.6	8.1	14.3	25.9	44.9	77.6	123.3	247.4
2005 4/.....	1.6	2.8	4.6	8.4	16.7	29.1	52.4	89.1	145.0	290.3
2004 4/.....	1.6	3.0	5.0	7.7	15.7	29.4	56.1	88.6	152.3	303.7
2003 4/.....	1.7	3.3	5.3	9.1	18.6	35.0	57.5	104.4	160.3	327.4
2002.....	1.8	3.1	4.4	9.6	19.0	39.0	61.2	104.5	169.2	354.1
2001.....	1.6	2.9	4.6	9.6	19.3	37.6	63.8	107.1	178.0	375.1
2000.....	1.5	2.9	5.0	10.5	21.3	42.0	72.3	117.6	190.0	401.0
1999.....	1.6	3.1	5.0	10.8	22.3	40.5	71.7	118.9	191.8	404.8
White, female										
2006 4/.....	0.6	1.2	1.7	3.3	6.7	13.0	26.0	47.4	84.3	211.3
2005 4/.....	0.9	1.1	2.0	3.6	7.7	14.9	27.2	52.9	98.9	267.7
2004 4/.....	0.8	1.2	2.0	3.9	7.8	15.6	30.9	56.2	100.7	271.1
2003 4/.....	0.8	1.4	2.1	4.0	8.6	16.2	32.2	58.5	106.4	297.1
2002.....	0.7	1.2	2.2	4.7	8.8	17.7	33.9	60.8	111.2	310.4
2001.....	0.8	1.3	2.4	4.5	9.0	18.6	32.7	65.6	115.0	331.0
2000.....	0.8	1.3	2.1	4.1	8.7	19.0	35.3	63.9	119.9	340.9
1999.....	1.0	1.2	2.4	4.5	9.2	18.4	36.5	66.8	124.4	360.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of arteries, arterioles and capillaries (I70-I78)										
Black, both sexes										
2006 4/.....	3.0	4.8	7.9	11.5	16.7	27.7	39.3	65.5	113.1	203.9
2005 4/.....	2.7	4.6	8.0	11.7	17.8	30.3	44.7	73.5	122.1	242.4
2004 4/.....	2.9	4.7	7.2	11.0	20.5	29.6	42.7	75.3	122.0	258.0
2003 4/.....	2.7	5.7	8.6	10.4	19.9	30.5	48.6	76.1	137.6	271.0
2002.....	3.1	5.9	8.1	13.2	20.7	30.9	49.3	85.2	134.7	273.8
2001.....	3.3	4.4	7.5	12.7	23.0	29.2	54.9	92.2	138.9	300.3
2000.....	3.5	5.3	8.3	13.3	22.9	32.9	52.8	92.4	146.3	315.9
1999.....	3.0	4.8	7.4	13.4	18.2	33.0	50.5	91.8	143.3	333.2
Black, male										
2006 4/.....	4.4	6.7	10.9	17.5	22.1	40.1	47.7	74.7	121.2	192.1
2005 4/.....	3.9	6.9	11.2	16.8	23.6	41.4	51.9	96.1	144.0	204.5
2004 4/.....	3.8	6.6	9.7	15.4	29.2	39.5	49.5	91.4	139.2	236.4
2003 4/.....	3.7	7.7	11.6	13.9	28.3	41.6	58.1	101.3	177.9	242.9
2002.....	4.4	7.4	12.0	19.1	29.3	36.9	64.1	100.6	159.7	276.6
2001.....	4.7	6.2	10.8	15.3	29.7	35.6	71.0	109.2	176.6	279.1
2000.....	4.6	6.5	12.0	17.5	28.6	41.4	65.9	116.2	171.1	325.2
1999.....	3.5	6.5	10.0	17.5	23.5	40.8	64.8	111.6	174.4	348.0
Black, female										
2006 4/.....	1.7	3.2	5.4	6.5	12.5	18.5	33.6	60.0	108.9	208.7
2005 4/.....	1.7	2.6	5.4	7.5	13.3	22.0	39.8	60.0	110.5	257.7
2004 4/.....	2.1	3.0	5.0	7.4	13.8	22.3	38.0	65.8	113.0	266.6
2003 4/.....	1.8	4.0	6.1	7.5	13.2	22.3	42.1	61.0	116.4	281.9
2002.....	1.9	4.5	4.8	8.3	13.9	26.5	39.4	76.0	121.7	272.7
2001.....	2.2	2.9	4.7	10.5	17.8	24.4	44.2	82.1	119.8	308.2
2000.....	2.5	4.4	5.2	9.9	18.5	26.7	44.2	78.0	133.7	312.5
1999.....	2.7	3.3	5.2	10.1	14.0	27.3	41.0	79.8	127.9	327.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
All races 3/, both sexes										
2006.....	0.1	0.2	0.4	0.9	1.7	3.2	6.1	12.6	27.2	89.9
2005.....	0.1	0.3	0.7	1.0	2.1	3.9	7.2	16.9	37.8	132.9
2004.....	0.1	0.3	0.5	1.0	2.3	4.1	8.5	17.5	38.0	137.0
2003.....	0.2	0.3	0.6	1.1	2.4	4.5	9.0	19.9	43.5	154.6
2002.....	0.1	0.3	0.6	1.4	2.4	5.2	9.3	20.5	47.8	168.1
2001.....	0.1	0.3	0.7	1.4	2.5	4.9	9.8	21.9	49.2	179.1
2000.....	0.1	0.4	0.6	1.3	2.6	5.2	10.5	22.7	52.4	188.8
1999.....	0.1	0.3	0.8	1.3	2.9	5.1	11.2	24.1	53.6	202.0
All races 3/, male										
2006.....	*	0.3	0.6	1.2	2.2	3.8	7.3	14.2	29.5	85.3
2005.....	*	0.4	1.0	1.2	2.7	4.8	8.7	20.0	42.0	121.7
2004.....	*	0.4	0.7	1.2	3.0	4.9	10.3	20.2	43.5	128.2
2003.....	0.2	0.5	0.8	1.5	3.2	5.4	10.6	23.9	48.3	138.6
2002.....	0.2	0.4	0.8	1.7	3.2	6.4	11.5	24.4	52.5	156.6
2001.....	*	0.4	0.8	1.7	3.2	6.2	12.1	25.0	53.2	167.2
2000.....	0.2	0.5	0.8	1.7	3.6	6.6	13.1	27.5	59.8	176.5
1999.....	0.2	0.4	1.0	1.8	3.8	6.3	13.3	28.1	60.0	189.0
All races 3/, female										
2006.....	*	*	0.2	0.5	1.3	2.6	5.1	11.4	25.7	92.1
2005.....	*	0.2	0.4	0.8	1.5	3.1	5.9	14.7	35.2	138.1
2004.....	*	*	0.3	0.8	1.7	3.4	7.1	15.5	34.6	141.0
2003.....	*	*	0.4	0.7	1.7	3.7	7.8	17.0	40.6	161.7
2002.....	*	*	0.5	1.0	1.7	4.1	7.5	17.8	45.0	173.1
2001.....	*	0.2	0.5	1.1	1.8	3.9	8.1	19.7	46.8	184.1
2000.....	*	0.2	0.4	0.9	1.7	4.0	8.4	19.4	48.1	193.8
1999.....	*	*	0.5	0.9	2.2	4.0	9.5	21.4	49.8	207.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
White, both sexes										
2006 4/.....	*	0.2	0.4	0.8	1.7	3.1	6.0	12.8	27.4	93.2
2005 4/.....	0.1	0.3	0.6	0.9	2.0	3.8	7.1	17.2	38.8	138.7
2004 4/.....	0.1	0.3	0.4	1.0	2.3	4.0	8.5	17.8	39.3	142.0
2003 4/.....	0.2	0.3	0.6	1.0	2.4	4.4	8.9	20.2	44.1	159.5
2002.....	0.1	0.2	0.5	1.3	2.3	5.3	9.1	20.4	48.4	174.7
2001.....	0.1	0.3	0.6	1.4	2.3	4.9	9.8	22.2	50.3	185.9
2000.....	0.1	0.3	0.5	1.2	2.4	5.2	10.4	23.1	53.0	195.2
1999.....	*	0.3	0.7	1.2	2.9	5.0	11.3	24.2	54.7	207.7
White, male										
2006 4/.....	*	0.3	0.6	1.1	2.2	3.6	7.1	14.3	29.3	87.9
2005 4/.....	*	0.4	0.9	1.2	2.6	4.8	8.5	20.6	42.9	128.3
2004 4/.....	*	0.4	0.7	1.2	2.9	4.9	10.4	20.5	44.8	134.4
2003 4/.....	0.2	0.5	0.8	1.4	3.1	5.2	10.1	23.8	48.4	143.3
2002.....	*	0.4	0.6	1.7	3.0	6.5	11.3	24.0	53.1	163.2
2001.....	*	0.4	0.8	1.7	3.0	6.2	12.1	25.2	54.0	173.9
2000.....	*	0.4	0.7	1.7	3.5	6.5	13.2	28.2	60.3	181.4
1999.....	*	0.4	1.0	1.6	3.7	6.2	13.3	27.7	60.8	193.7
White, female										
2006 4/.....	*	*	*	0.6	1.2	2.6	5.1	11.6	26.2	95.7
2005 4/.....	*	*	0.3	0.7	1.5	3.0	5.9	14.7	36.2	143.5
2004 4/.....	*	*	0.2	0.8	1.7	3.2	7.0	15.9	35.9	145.4
2003 4/.....	*	*	0.4	0.7	1.7	3.6	7.8	17.6	41.4	166.7
2002.....	*	*	0.4	1.0	1.6	4.2	7.3	17.7	45.5	179.7
2001.....	*	*	0.5	1.0	1.6	3.9	8.0	20.2	48.0	190.9
2000.....	*	*	0.3	0.8	1.5	4.1	8.2	19.5	48.7	200.8
1999.....	*	*	0.5	0.9	2.0	3.9	9.7	21.7	51.0	213.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Atherosclerosis (I70)										
Black, both sexes										
2006 4/.....	*	*	*	1.5	3.1	4.5	8.8	14.2	29.3	71.6
2005 4/.....	*	*	1.3	1.3	3.1	5.8	9.5	16.5	33.7	97.8
2004 4/.....	*	*	*	1.7	3.2	5.4	9.9	18.4	31.9	102.9
2003 4/.....	*	*	1.3	1.9	3.5	6.5	12.7	20.8	45.4	121.1
2002.....	*	*	1.5	1.9	4.2	5.2	11.5	26.8	51.3	117.0
2001.....	*	*	1.0	2.0	5.0	6.3	12.1	22.9	42.8	125.6
2000.....	*	*	1.5	2.1	4.7	5.6	12.9	23.5	53.9	138.3
1999.....	*	*	1.3	2.1	4.7	7.0	11.9	27.5	46.8	154.3
Black, male										
2006 4/.....	*	*	*	2.5	3.7	6.6	11.4	16.6	39.0	75.1
2005 4/.....	*	*	*	*	4.4	7.6	12.3	17.0	40.0	75.3
2004 4/.....	*	*	*	*	4.9	6.1	12.0	23.2	40.9	78.5
2003 4/.....	*	*	*	*	4.8	8.1	18.1	28.8	59.1	104.3
2002.....	*	*	2.3	*	6.3	6.2	15.4	34.9	59.6	110.4
2001.....	*	*	*	*	5.9	7.6	15.1	30.2	49.0	113.5
2000.....	*	*	*	*	6.0	8.6	14.0	25.5	66.6	147.6
1999.....	*	*	*	3.4	5.7	8.8	15.6	37.0	56.7	164.0
Black, female										
2006 4/.....	*	*	*	*	*	*	7.0	12.9	24.2	70.2
2005 4/.....	*	*	*	*	*	4.4	7.6	16.3	30.4	106.9
2004 4/.....	*	*	*	*	*	4.9	8.6	15.6	27.2	112.6
2003 4/.....	*	*	*	*	*	5.3	9.1	16.0	38.3	127.6
2002.....	*	*	*	*	*	4.4	9.0	22.0	46.9	119.5
2001.....	*	*	*	*	4.3	5.3	10.1	18.6	39.7	130.1
2000.....	*	*	*	*	3.6	*	12.2	22.2	47.6	134.8
1999.....	*	*	*	*	3.8	5.6	9.5	21.7	41.9	150.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
All races 3/, both sexes										
2006.....	1.0	1.6	2.1	3.4	5.6	10.6	18.0	29.5	43.2	62.7
2005.....	0.9	1.4	2.2	3.3	6.8	11.7	19.8	31.6	46.6	64.9
2004.....	0.8	1.4	2.2	3.1	6.4	11.4	21.1	32.6	48.8	64.8
2003.....	0.8	1.7	2.4	3.7	7.3	13.5	22.7	36.6	50.7	70.0
2002.....	0.9	1.5	2.0	4.0	7.4	14.7	23.9	37.5	51.5	69.4
2001.....	0.9	1.4	2.0	3.7	7.8	15.0	23.8	39.0	55.4	77.5
2000.....	0.9	1.3	2.2	4.3	8.4	16.3	27.8	41.0	56.5	79.2
1999.....	0.9	1.4	2.1	4.4	8.1	16.5	27.3	42.2	58.2	80.3
All races 3/, male										
2006.....	1.5	2.4	3.3	5.3	8.4	15.2	24.8	41.1	59.0	87.6
2005.....	1.5	2.3	3.3	5.3	10.0	17.1	29.0	43.8	65.4	91.0
2004.....	1.3	2.1	3.3	4.9	9.7	17.0	30.6	44.2	70.3	95.0
2003.....	1.3	2.5	3.7	5.8	11.4	20.2	33.0	53.8	74.2	105.5
2002.....	1.4	2.4	3.2	6.3	11.5	23.0	34.9	55.0	74.5	106.6
2001.....	1.4	2.1	3.2	5.8	12.0	22.2	36.0	56.3	82.5	114.8
2000.....	1.3	1.9	3.5	6.7	13.1	24.9	41.8	61.9	84.9	127.3
1999.....	1.2	2.2	3.3	6.8	13.0	25.3	41.0	63.3	88.6	125.3
All races 3/, female										
2006.....	0.4	0.8	1.0	1.6	3.1	6.5	12.5	21.0	33.3	51.0
2005.....	0.4	0.6	1.1	1.5	3.8	7.0	12.4	22.7	35.0	52.9
2004.....	0.4	0.7	1.0	1.5	3.3	6.6	13.5	24.3	35.6	51.2
2003.....	0.4	0.9	1.1	1.7	3.6	7.8	14.5	24.3	36.3	54.2
2002.....	0.3	0.7	1.0	1.8	3.7	7.5	15.1	25.1	37.6	53.3
2001.....	0.3	0.6	0.9	1.8	3.9	8.8	14.0	26.9	39.2	61.9
2000.....	0.5	0.8	1.0	2.1	4.1	8.9	16.7	26.5	39.7	59.6
1999.....	0.5	0.7	1.0	2.1	3.7	9.0	16.6	27.5	40.4	62.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
White, both sexes										
2006 4/.....	0.8	1.4	1.8	3.1	5.6	10.5	18.5	30.4	44.1	64.3
2005 4/.....	0.8	1.2	1.8	3.2	6.7	11.7	20.6	32.5	47.6	66.2
2004 4/.....	0.7	1.2	1.9	2.8	6.1	11.7	22.1	33.7	49.8	66.8
2003 4/.....	0.7	1.4	2.1	3.6	7.3	14.0	23.6	38.1	52.2	72.4
2002.....	0.7	1.2	1.7	3.8	7.5	15.3	24.8	39.0	53.1	71.2
2001.....	0.7	1.2	1.7	3.6	7.7	15.5	24.6	39.8	57.0	79.7
2000.....	0.7	1.2	2.0	4.1	8.4	16.7	29.0	42.0	58.1	80.8
1999.....	0.7	1.2	1.9	4.2	8.5	17.1	28.4	43.5	60.0	82.2
White, male										
2006 4/.....	1.3	2.2	2.9	5.0	8.5	15.2	25.8	42.8	61.0	90.4
2005 4/.....	1.2	1.9	2.8	5.1	10.0	17.2	30.6	45.1	67.1	94.2
2004 4/.....	1.1	1.8	3.0	4.5	9.3	17.4	32.1	45.8	72.1	98.7
2003 4/.....	1.2	2.1	3.4	5.7	11.4	21.1	34.6	56.2	76.5	110.0
2002.....	1.2	2.0	2.7	5.9	11.6	24.3	36.1	57.5	77.7	111.8
2001.....	1.1	1.9	2.8	5.6	11.9	23.2	37.3	57.7	85.5	120.3
2000.....	1.0	1.8	3.2	6.5	13.3	26.1	43.8	63.6	88.6	132.8
1999.....	1.0	1.9	2.9	6.7	13.6	26.5	43.0	66.3	91.5	129.8
White, female										
2006 4/.....	0.3	0.6	0.7	1.4	2.9	6.3	12.5	21.2	33.4	52.0
2005 4/.....	0.3	0.4	0.9	1.4	3.5	6.9	12.3	23.1	35.3	53.3
2004 4/.....	0.3	0.5	0.9	1.2	3.2	6.7	13.9	24.8	35.9	52.4
2003 4/.....	0.3	0.7	0.8	1.5	3.5	7.9	14.6	25.0	37.2	55.7
2002.....	0.3	0.5	0.8	1.7	3.7	7.4	15.5	25.8	38.1	53.5
2001.....	0.3	0.5	0.7	1.6	3.9	8.8	14.3	27.1	39.8	62.6
2000.....	0.3	0.5	0.9	1.9	3.9	8.5	17.1	26.8	40.0	59.6
1999.....	0.4	0.5	0.9	1.8	3.7	8.9	16.7	27.5	41.3	63.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Aortic aneurysm and dissection (I71)										
Black, both sexes										
2006 4/.....	2.2	3.1	5.1	6.0	7.5	13.0	16.0	23.6	34.1	47.2
2005 4/.....	1.8	3.3	4.9	5.6	8.8	13.3	15.9	25.6	38.6	50.2
2004 4/.....	1.8	3.2	4.0	5.8	10.0	11.6	14.9	26.1	40.6	43.7
2003 4/.....	1.8	3.8	4.7	5.7	9.0	11.7	17.9	25.9	36.0	46.4
2002.....	2.1	3.8	4.7	6.8	8.3	11.6	18.4	24.9	37.4	50.5
2001.....	2.1	2.7	4.4	6.2	9.3	11.6	18.4	34.4	42.5	55.3
2000.....	2.3	2.9	3.7	7.3	9.3	14.3	20.3	33.8	40.4	62.3
1999.....	2.0	3.5	4.0	6.5	6.5	14.1	19.8	29.6	40.5	60.3
Black, male										
2006 4/.....	3.1	4.5	7.5	9.2	10.2	18.5	19.7	26.4	34.9	50.9
2005 4/.....	3.1	5.3	7.1	8.8	12.0	18.8	17.1	34.0	47.1	51.1
2004 4/.....	2.8	5.1	6.0	8.8	15.9	17.9	18.5	29.6	48.1	56.6
2003 4/.....	2.8	5.8	7.1	8.0	14.7	16.3	20.4	32.9	49.5	51.1
2002.....	3.5	5.7	7.7	11.3	13.1	14.7	24.7	31.2	41.3	49.3
2001.....	3.4	3.9	6.9	8.6	14.6	15.0	26.6	43.2	54.6	48.8
2000.....	3.5	3.6	6.0	10.4	12.8	17.7	27.7	47.7	45.5	69.2
1999.....	2.7	5.4	6.4	9.0	9.3	19.2	25.4	29.9	54.1	76.2
Black, female										
2006 4/.....	1.4	1.9	3.0	3.4	5.4	9.0	13.5	22.0	33.6	45.7
2005 4/.....	*	1.6	3.1	3.0	6.4	9.2	15.2	20.5	34.1	49.9
2004 4/.....	*	1.6	2.4	3.3	5.4	6.8	12.5	24.0	36.7	38.6
2003 4/.....	*	2.1	2.7	3.8	4.7	8.4	16.2	21.7	29.0	44.6
2002.....	*	2.1	2.2	3.2	4.5	9.3	14.1	21.1	35.4	50.9
2001.....	*	1.7	2.2	4.2	5.1	9.0	13.0	29.1	36.4	57.7
2000.....	*	2.3	*	4.7	6.5	11.7	15.3	25.3	37.8	59.7
1999.....	1.4	2.0	*	4.4	4.3	10.4	16.0	29.5	33.7	54.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)										
All races 3/, both sexes										
2006.....	0.3	0.7	1.0	1.9	3.3	5.8	10.2	17.6	29.1	65.8
2005.....	0.4	0.5	1.0	2.1	3.5	6.3	11.4	19.3	31.3	70.6
2004.....	0.4	0.7	1.2	2.1	3.5	6.9	11.9	19.1	32.3	74.5
2003.....	0.4	0.7	1.2	2.0	4.0	7.0	11.4	20.0	32.0	76.0
2002.....	0.4	0.7	1.1	2.2	4.3	7.7	12.6	20.4	32.3	79.0
2001.....	0.5	0.6	1.2	2.2	4.3	7.3	12.9	21.5	32.7	80.7
2000.....	0.4	0.8	1.3	2.1	4.3	8.2	12.8	21.8	35.8	84.0
1999.....	0.5	0.7	1.2	2.3	4.4	7.0	12.9	21.3	36.1	84.9
All races 3/, male										
2006.....	0.3	0.7	1.2	2.3	4.0	7.5	12.1	20.8	33.5	67.3
2005.....	0.3	0.6	1.1	2.5	4.2	7.7	13.7	24.6	35.8	67.2
2004.....	0.4	0.8	1.4	2.3	3.9	7.7	13.6	22.9	35.5	71.6
2003.....	0.3	0.7	1.3	2.0	4.4	9.3	13.0	25.1	37.1	74.2
2002.....	0.4	0.8	1.2	2.3	4.9	8.7	14.2	23.4	39.2	80.8
2001.....	0.4	0.7	1.2	2.4	4.8	8.5	15.2	24.7	40.0	82.5
2000.....	0.3	0.8	1.4	2.6	4.9	9.7	15.7	26.6	41.7	87.6
1999.....	0.4	0.8	1.2	2.6	5.2	8.1	15.7	25.9	40.8	82.2
All races 3/, female										
2006.....	0.2	0.6	0.9	1.5	2.7	4.3	8.6	15.3	26.2	65.0
2005.....	0.5	0.5	0.9	1.7	2.8	5.2	9.6	15.5	28.4	72.2
2004.....	0.5	0.6	1.0	1.8	3.2	6.2	10.5	16.3	30.3	75.8
2003.....	0.5	0.7	1.0	1.9	3.6	5.0	10.1	16.3	28.9	76.9
2002.....	0.4	0.6	1.0	2.1	3.7	6.7	11.3	18.3	28.1	78.2
2001.....	0.5	0.6	1.3	2.0	3.9	6.3	11.1	19.3	28.3	79.9
2000.....	0.5	0.7	1.1	1.6	3.7	6.9	10.5	18.5	32.2	82.6
1999.....	0.5	0.6	1.2	2.1	3.6	6.0	10.6	18.2	33.3	86.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of arteries, arterioles and capillaries (I72-I78)										
White, both sexes										
2006 4/.....	0.2	0.6	0.9	1.7	3.1	5.5	10.0	17.1	27.8	65.4
2005 4/.....	0.4	0.5	1.0	1.8	3.3	6.0	11.0	18.6	30.3	69.9
2004 4/.....	0.4	0.7	1.1	1.9	3.2	6.4	11.7	18.4	31.4	72.5
2003 4/.....	0.3	0.6	1.0	1.9	3.7	6.6	11.1	19.5	30.7	74.6
2002.....	0.4	0.6	1.0	2.0	3.9	7.1	12.2	19.7	31.7	77.8
2001.....	0.4	0.6	1.2	2.1	4.0	7.1	12.2	20.8	31.4	78.5
2000.....	0.3	0.7	1.0	1.9	3.9	7.8	12.4	21.0	34.9	82.3
1999.....	0.5	0.7	1.1	2.1	4.1	6.6	12.5	20.6	34.8	82.9
White, male										
2006 4/.....	0.2	0.7	1.1	2.0	3.7	7.1	12.1	20.5	33.0	69.0
2005 4/.....	0.3	0.5	1.0	2.1	4.1	7.1	13.3	23.4	35.0	67.8
2004 4/.....	0.4	0.8	1.3	2.1	3.6	7.1	13.6	22.3	35.4	70.7
2003 4/.....	0.3	0.7	1.2	2.0	4.0	8.7	12.8	24.4	35.4	74.1
2002.....	0.4	0.8	1.1	2.0	4.4	8.2	13.7	23.0	38.4	79.1
2001.....	0.3	0.6	1.1	2.3	4.4	8.3	14.4	24.2	38.5	80.9
2000.....	0.3	0.7	1.1	2.4	4.5	9.4	15.3	25.9	41.1	86.9
1999.....	0.4	0.8	1.1	2.4	4.9	7.9	15.3	24.8	39.6	81.3
White, female										
2006 4/.....	0.3	0.5	0.8	1.4	2.6	4.1	8.3	14.5	24.6	63.7
2005 4/.....	0.5	0.5	0.9	1.5	2.6	4.9	9.0	15.1	27.4	70.8
2004 4/.....	0.4	0.5	0.9	1.8	2.8	5.7	10.1	15.6	28.9	73.3
2003 4/.....	0.4	0.6	0.9	1.9	3.4	4.8	9.8	15.9	27.8	74.8
2002.....	0.3	0.5	0.9	2.0	3.4	6.1	11.0	17.3	27.5	77.2
2001.....	0.5	0.6	1.2	1.8	3.5	6.0	10.4	18.4	27.1	77.5
2000.....	0.4	0.6	0.9	1.4	3.3	6.5	10.0	17.6	31.2	80.4
1999.....	0.5	0.5	1.0	1.8	3.4	5.6	10.2	17.6	32.0	83.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, both sexes

2006 4/.....	*	1.3	2.1	4.0	6.1	10.2	14.5	27.6	49.7	85.1
2005 4/.....	0.7	1.0	1.8	4.7	5.9	11.2	19.3	31.4	49.8	94.3
2004 4/.....	0.8	1.1	2.3	3.6	7.4	12.7	17.8	30.8	49.5	111.4
2003 4/.....	0.8	1.5	2.6	2.8	7.3	12.3	18.0	29.4	56.1	103.6
2002.....	0.7	1.5	2.0	4.4	8.3	14.1	19.4	33.5	46.1	106.3
2001.....	1.1	1.3	2.1	4.5	8.7	11.3	24.4	34.9	53.6	119.4
2000.....	0.9	1.7	3.1	3.9	9.0	13.0	19.6	35.2	52.0	115.4
1999.....	0.8	1.0	2.1	4.9	7.0	11.9	18.8	34.6	56.1	118.6

Black, male

2006 4/.....	*	1.6	2.6	5.8	8.1	15.0	16.6	31.8	47.2	66.1
2005 4/.....	*	*	2.4	6.2	7.3	15.0	22.5	45.1	56.9	78.1
2004 4/.....	*	*	2.9	4.9	8.4	15.5	19.1	38.7	50.2	101.3
2003 4/.....	*	*	2.8	3.2	8.9	17.2	19.7	39.7	69.4	87.6
2002.....	*	*	2.1	5.2	9.9	16.0	24.0	34.4	58.8	116.9
2001.....	*	*	2.5	4.5	9.2	13.0	29.3	35.8	73.0	116.9
2000.....	*	1.9	4.0	4.8	9.7	15.1	24.1	43.0	59.0	108.4
1999.....	*	*	*	5.1	8.5	12.7	23.7	44.6	63.7	107.8

Black, female

2006 4/.....	*	*	1.7	2.6	4.5	6.6	13.1	25.2	51.0	92.9
2005 4/.....	*	*	*	3.4	4.8	8.5	17.1	23.2	46.0	100.9
2004 4/.....	*	*	1.8	2.5	6.6	10.6	16.9	26.1	49.1	115.4
2003 4/.....	*	1.8	2.4	2.4	6.0	8.6	16.8	23.3	49.1	109.8
2002.....	*	1.9	1.8	3.7	7.0	12.8	16.4	32.9	39.4	102.3
2001.....	*	*	1.8	4.5	8.3	10.1	21.1	34.4	43.7	120.4
2000.....	*	*	2.4	3.2	8.3	11.5	16.6	30.5	48.4	118.0
1999.....	*	*	*	4.7	5.8	11.4	15.5	28.6	52.3	122.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes

2006.....	0.7	1.1	1.4	1.6	1.9	2.4	3.5	5.6	8.7	18.4
2005.....	0.9	1.3	1.6	1.9	2.4	3.3	4.4	7.0	10.8	23.0
2004.....	0.9	1.3	1.4	1.9	2.5	3.1	4.3	7.1	11.5	23.3
2003.....	0.9	1.3	1.5	1.7	2.5	3.3	4.9	7.0	11.9	24.0
2002.....	0.8	1.2	1.5	2.0	2.5	3.4	4.3	7.4	13.2	24.4
2001.....	0.8	1.1	1.3	1.9	2.6	3.8	5.4	7.7	11.9	26.1
2000.....	0.9	1.2	1.5	1.7	2.6	3.4	5.3	7.7	11.9	25.6
1999.....	0.8	1.0	1.0	1.6	2.2	3.3	4.3	6.8	12.7	23.4

All races 3/, male

2006.....	0.8	1.3	1.8	1.9	2.3	2.4	3.5	6.1	8.3	16.2
2005.....	1.0	1.6	2.1	2.3	3.0	4.0	5.0	6.6	11.1	19.2
2004.....	1.0	1.4	1.8	2.2	3.0	3.3	4.6	7.9	12.2	19.6
2003.....	1.0	1.4	1.8	2.2	2.7	3.5	4.9	6.9	10.9	19.1
2002.....	0.8	1.3	1.6	2.3	2.6	3.6	4.2	7.0	13.3	20.5
2001.....	0.9	1.2	1.5	2.2	2.9	3.9	5.9	8.3	12.1	22.0
2000.....	1.1	1.5	1.7	1.9	2.7	3.9	5.6	8.0	12.4	22.2
1999.....	0.9	1.1	1.2	1.7	2.3	3.7	4.8	7.5	13.2	20.8

All races 3/, female

2006.....	0.7	1.0	0.9	1.2	1.6	2.5	3.6	5.3	9.0	19.5
2005.....	0.9	1.0	1.2	1.5	1.8	2.7	4.0	7.2	10.7	24.8
2004.....	0.8	1.1	1.1	1.6	2.0	2.9	4.0	6.5	11.0	25.0
2003.....	0.9	1.1	1.2	1.3	2.3	3.0	5.0	7.0	12.5	26.2
2002.....	0.8	1.1	1.4	1.7	2.5	3.2	4.4	7.7	13.1	26.2
2001.....	0.8	0.9	1.0	1.7	2.3	3.7	5.0	7.3	11.7	27.8
2000.....	0.7	1.0	1.3	1.5	2.6	3.0	5.0	7.5	11.6	26.9
1999.....	0.6	1.0	0.9	1.5	2.1	2.9	3.9	6.3	12.4	24.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)										
White, both sexes										
2006 4/.....	0.6	1.0	1.2	1.5	1.7	2.3	3.4	5.5	8.4	18.1
2005 4/.....	0.7	1.2	1.4	1.7	2.1	3.2	4.2	6.7	10.5	23.5
2004 4/.....	0.7	1.1	1.2	1.8	2.3	3.0	3.9	6.8	11.3	23.2
2003 4/.....	0.8	1.1	1.4	1.6	2.2	3.0	4.7	6.5	11.6	23.6
2002.....	0.6	1.0	1.4	1.9	2.2	3.2	4.1	7.1	12.7	24.3
2001.....	0.7	0.9	1.1	1.8	2.3	3.6	5.1	7.2	11.8	26.0
2000.....	0.8	1.1	1.4	1.5	2.4	2.9	5.0	7.3	11.8	25.5
1999.....	0.6	0.9	0.9	1.6	2.1	2.9	4.1	6.5	12.4	23.2
White, male										
2006 4/.....	0.6	1.1	1.5	1.9	1.9	2.2	3.4	6.0	7.6	16.0
2005 4/.....	0.8	1.4	1.9	2.1	2.6	3.7	4.6	6.4	10.5	19.1
2004 4/.....	0.8	1.4	1.5	2.1	2.7	3.1	4.3	7.6	11.9	19.7
2003 4/.....	0.9	1.2	1.6	2.1	2.3	3.2	4.6	6.4	10.2	18.5
2002.....	0.6	1.1	1.5	2.1	2.3	3.3	4.0	6.7	12.4	19.9
2001.....	0.7	1.2	1.3	2.0	2.5	3.6	5.6	7.8	12.0	21.7
2000.....	1.0	1.3	1.6	1.7	2.4	3.4	5.5	7.8	12.3	22.0
1999.....	0.8	0.9	1.0	1.7	2.3	3.2	4.7	7.0	12.8	20.7
White, female										
2006 4/.....	0.5	0.8	0.8	1.1	1.4	2.3	3.5	5.2	8.9	19.1
2005 4/.....	0.7	1.0	1.0	1.4	1.6	2.7	3.9	6.9	10.5	25.5
2004 4/.....	0.6	0.9	0.9	1.4	1.8	2.8	3.6	6.2	11.0	24.8
2003 4/.....	0.7	1.1	1.2	1.1	2.2	2.9	4.7	6.5	12.4	25.8
2002.....	0.7	1.0	1.2	1.6	2.1	3.0	4.2	7.4	12.9	26.3
2001.....	0.7	0.7	1.0	1.5	2.1	3.5	4.8	6.8	11.7	27.9
2000.....	0.5	0.9	1.1	1.4	2.4	2.5	4.7	7.0	11.5	26.9
1999.....	0.5	0.9	0.9	1.5	1.9	2.7	3.7	6.1	12.2	24.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other disorders of circulatory system (I80-I99)										
Black, both sexes										
2006 4/.....	1.9	2.7	3.0	2.7	4.4	4.2	5.8	8.0	14.9	25.7
2005 4/.....	2.3	2.6	3.5	4.0	5.2	5.5	7.6	11.5	17.5	24.3
2004 4/.....	2.5	2.4	3.6	3.6	5.0	4.9	8.9	12.9	14.6	28.8
2003 4/.....	2.3	2.6	2.5	3.6	5.2	6.2	9.3	13.8	17.8	33.6
2002.....	1.8	2.4	2.9	4.2	5.9	6.4	7.2	12.9	22.2	30.0
2001.....	1.8	2.1	2.4	3.9	6.1	6.9	9.0	14.1	14.9	30.1
2000.....	1.8	2.5	3.2	3.9	5.1	8.7	9.1	14.6	14.1	30.0
1999.....	1.9	2.4	2.3	2.1	3.9	6.9	6.4	11.4	17.6	28.4
Black, male										
2006 4/.....	1.8	2.6	4.7	2.9	6.2	4.7	*	*	17.1	22.3
2005 4/.....	2.4	3.6	4.1	4.8	6.9	7.8	10.1	12.1	22.5	25.1
2004 4/.....	2.5	2.1	5.0	3.6	6.2	5.8	8.7	14.1	16.5	24.8
2003 4/.....	1.8	3.4	3.4	3.6	7.3	7.6	10.2	15.5	20.7	32.3
2002.....	2.1	3.0	2.9	5.4	5.9	7.7	7.7	12.4	28.3	32.2
2001.....	2.2	1.7	3.1	4.7	8.5	7.9	10.4	16.3	*	29.5
2000.....	2.0	3.1	3.7	4.8	5.6	9.4	8.7	13.2	*	30.0
1999.....	2.6	2.7	3.0	*	*	9.3	6.8	14.7	20.9	*
Black, female										
2006 4/.....	1.9	2.8	1.6	2.6	3.0	3.8	5.9	8.0	13.7	27.1
2005 4/.....	2.2	1.8	3.0	3.3	3.8	3.7	5.8	11.2	14.8	24.0
2004 4/.....	2.6	2.6	2.4	3.7	4.1	4.1	9.0	12.1	13.6	30.3
2003 4/.....	2.7	1.9	1.8	3.6	3.6	5.1	8.6	12.8	16.2	34.0
2002.....	1.6	1.8	2.9	3.2	5.9	5.4	6.9	13.2	19.1	29.2
2001.....	1.5	2.4	*	3.2	4.2	6.2	8.1	12.8	14.7	30.3
2000.....	1.7	1.9	2.8	3.2	4.7	8.3	9.3	15.4	14.4	30.0
1999.....	*	2.1	*	*	4.2	5.0	6.1	9.4	16.0	30.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)										
All races 3/, both sexes										
2006.....	0.6	0.9	1.0	1.1	1.4	1.7	2.4	3.9	6.1	12.0
2005.....	0.7	0.9	1.0	1.3	1.6	2.3	3.0	5.2	7.9	16.8
2004.....	0.8	0.9	1.0	1.4	1.8	2.2	3.1	5.4	8.8	17.3
2003.....	0.8	1.0	1.1	1.3	1.9	2.6	3.7	5.2	9.0	18.1
2002.....	0.7	0.9	1.2	1.6	1.9	2.6	3.1	5.4	10.1	18.6
2001.....	0.6	0.8	0.9	1.4	2.0	2.8	4.1	5.9	8.9	19.6
2000.....	0.6	0.9	1.1	1.3	1.8	2.5	3.9	5.8	8.8	18.7
1999.....	0.6	0.7	0.7	1.1	1.4	2.2	2.8	4.3	8.0	14.6
All races 3/, male										
2006.....	0.6	0.9	1.3	1.4	1.5	1.5	2.3	3.9	5.7	10.0
2005.....	0.6	1.0	1.1	1.5	1.9	2.7	3.4	5.0	7.9	13.5
2004.....	0.8	0.9	1.2	1.4	2.0	2.2	3.3	5.7	9.0	14.3
2003.....	0.7	1.0	1.2	1.5	2.0	2.7	3.5	4.8	7.8	13.8
2002.....	0.6	0.9	1.1	1.7	1.7	2.7	2.9	4.7	9.6	14.8
2001.....	0.6	0.8	1.0	1.5	2.1	2.7	4.4	5.9	8.3	15.2
2000.....	0.7	0.9	1.1	1.3	1.6	2.8	3.8	5.7	8.9	15.0
1999.....	0.6	0.6	0.6	1.0	1.2	2.3	3.1	4.4	7.9	11.2
All races 3/, female										
2006.....	0.6	0.9	0.7	0.8	1.3	1.9	2.5	3.9	6.3	12.9
2005.....	0.8	0.8	0.9	1.2	1.4	2.0	2.7	5.4	8.0	18.2
2004.....	0.8	1.0	0.8	1.4	1.5	2.1	3.0	5.2	8.7	18.6
2003.....	0.8	1.0	1.0	1.1	1.7	2.5	3.9	5.4	9.8	20.1
2002.....	0.7	0.9	1.2	1.5	2.0	2.6	3.2	5.9	10.4	20.2
2001.....	0.7	0.8	0.8	1.3	1.9	2.9	3.8	5.9	9.2	21.4
2000.....	0.5	0.9	1.1	1.2	1.9	2.3	4.0	5.9	8.7	20.2
1999.....	0.6	0.7	0.7	1.1	1.5	2.0	2.5	4.2	8.1	15.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)										
White, both sexes										
2006 4/.....	0.4	0.7	0.8	1.1	1.2	1.6	2.4	3.8	5.8	11.8
2005 4/.....	0.5	0.8	0.8	1.2	1.5	2.2	2.8	5.0	7.6	17.0
2004 4/.....	0.6	0.8	0.8	1.2	1.5	2.1	2.8	5.1	8.7	17.1
2003 4/.....	0.6	0.8	1.0	1.2	1.6	2.5	3.5	4.8	8.8	17.8
2002.....	0.5	0.8	1.0	1.5	1.6	2.5	2.9	5.2	9.6	18.4
2001.....	0.5	0.7	0.8	1.3	1.7	2.6	3.8	5.6	8.9	19.6
2000.....	0.6	0.8	1.0	1.1	1.6	2.1	3.7	5.5	8.8	18.8
1999.....	0.5	0.6	0.6	1.0	1.3	2.0	2.7	4.1	7.8	14.3
White, male										
2006 4/.....	0.4	0.8	1.0	1.5	1.2	1.4	2.2	3.9	5.0	9.9
2005 4/.....	0.5	0.9	1.0	1.4	1.7	2.6	3.0	4.7	7.4	13.6
2004 4/.....	0.6	0.8	1.0	1.4	1.8	2.1	3.0	5.5	8.7	14.3
2003 4/.....	0.7	0.8	1.1	1.5	1.7	2.5	3.3	4.3	7.5	13.3
2002.....	0.4	0.8	1.0	1.5	1.5	2.4	2.8	4.7	8.8	14.4
2001.....	0.5	0.7	0.9	1.4	1.7	2.6	4.1	5.7	8.3	15.0
2000.....	0.7	0.8	1.0	1.1	1.4	2.5	3.6	5.6	9.0	14.9
1999.....	0.5	0.6	0.5	1.0	1.2	2.0	3.0	4.2	7.7	11.4
White, female										
2006 4/.....	0.4	0.7	0.6	0.7	1.1	1.7	2.5	3.8	6.2	12.7
2005 4/.....	0.6	0.8	0.7	1.1	1.3	1.9	2.6	5.2	7.7	18.6
2004 4/.....	0.5	0.8	0.7	1.1	1.3	2.1	2.7	4.9	8.7	18.4
2003 4/.....	0.6	0.9	1.0	0.8	1.6	2.4	3.7	5.1	9.7	19.8
2002.....	0.6	0.9	1.0	1.4	1.8	2.5	3.0	5.6	10.1	20.2
2001.....	0.6	0.6	0.8	1.2	1.7	2.6	3.5	5.6	9.2	21.5
2000.....	0.5	0.8	0.9	1.1	1.8	1.8	3.8	5.4	8.7	20.4
1999.....	0.5	0.6	0.7	1.1	1.4	1.9	2.4	4.1	7.8	15.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Phlebitis, thrombophlebitis, venous embolism and thrombosis (I80-I82)

Black, both sexes

2006 4/.....	1.9	2.3	2.6	1.9	3.2	3.6	3.9	5.5	11.6	16.3
2005 4/.....	1.8	1.8	2.5	2.8	3.7	3.9	6.3	9.2	14.1	19.0
2004 4/.....	2.2	2.1	2.6	3.0	4.3	3.2	6.8	10.2	11.1	22.8
2003 4/.....	2.1	2.2	1.9	2.9	4.5	4.8	7.4	10.6	13.7	26.3
2002.....	1.6	1.8	2.5	3.2	4.4	4.9	5.6	9.3	18.3	23.8
2001.....	1.5	1.8	1.7	2.9	5.3	5.3	7.8	10.4	11.6	23.3
2000.....	1.3	1.9	2.5	2.9	3.9	6.8	7.1	11.7	9.3	21.9
1999.....	1.5	1.6	1.5	1.6	2.4	4.8	4.5	7.1	12.7	18.8

Black, male

2006 4/.....	1.8	2.1	3.9	*	4.6	*	*	*	15.1	*
2005 4/.....	*	2.2	2.5	2.8	4.5	5.2	8.5	10.3	16.9	*
2004 4/.....	2.0	1.7	3.3	2.6	5.0	*	7.4	10.0	*	19.9
2003 4/.....	1.6	2.8	2.3	*	6.2	5.9	7.6	11.9	*	25.0
2002.....	1.7	2.1	2.4	4.2	4.4	5.7	*	*	22.2	22.5
2001.....	1.7	*	2.4	3.3	6.9	5.1	8.8	11.2	*	*
2000.....	*	2.2	2.4	*	*	6.5	7.0	*	*	*
1999.....	1.7	*	*	*	*	6.8	*	*	*	*

Black, female

2006 4/.....	1.9	2.5	1.6	2.1	*	*	*	6.2	9.8	17.9
2005 4/.....	2.1	1.5	2.5	2.8	3.0	*	4.8	8.5	12.6	19.5
2004 4/.....	2.4	2.4	1.9	3.3	3.8	*	6.4	10.2	10.6	24.0
2003 4/.....	2.5	1.7	*	3.3	3.2	4.0	7.3	9.8	13.5	26.7
2002.....	1.6	1.6	2.6	*	4.5	4.3	5.8	11.0	16.3	24.2
2001.....	1.3	2.2	*	2.6	4.0	5.4	7.2	10.0	12.3	24.0
2000.....	*	1.7	2.6	2.7	3.9	7.1	7.1	13.4	10.6	22.3
1999.....	*	1.8	*	*	*	*	4.5	6.3	12.5	21.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified disorders of circulatory system (I83-I99)

All races 3/, both sexes

2006.....	0.1	0.2	0.4	0.4	0.5	0.7	1.1	1.7	2.6	6.5
2005.....	0.2	0.4	0.6	0.6	0.7	1.0	1.4	1.7	2.9	6.3
2004.....	0.1	0.3	0.4	0.5	0.7	0.9	1.2	1.7	2.7	6.0
2003.....	0.2	0.3	0.4	0.4	0.6	0.7	1.2	1.8	2.8	5.9
2002.....	0.1	0.3	0.3	0.4	0.7	0.7	1.2	2.1	3.1	5.9
2001.....	0.2	0.3	0.4	0.5	0.6	1.0	1.4	1.8	3.0	6.5
2000.....	0.2	0.3	0.4	0.4	0.9	0.9	1.4	1.9	3.1	6.8
1999.....	0.2	0.3	0.4	0.5	0.8	1.1	1.5	2.5	4.7	8.8

All races 3/, male

2006.....	*	0.3	0.5	0.5	0.8	0.8	1.2	2.2	2.6	6.3
2005.....	0.4	0.6	0.9	0.8	1.1	1.3	1.6	1.6	3.2	5.7
2004.....	0.2	0.5	0.6	0.8	0.9	1.1	1.3	2.3	3.3	5.4
2003.....	0.2	0.4	0.6	0.7	0.6	0.8	1.4	2.1	3.0	5.3
2002.....	0.2	0.4	0.5	0.6	0.8	0.9	1.3	2.3	3.7	5.7
2001.....	0.3	0.4	0.5	0.7	0.8	1.2	1.5	2.4	3.8	6.8
2000.....	0.3	0.5	0.6	0.6	1.1	1.0	1.8	2.4	3.5	7.3
1999.....	0.3	0.4	0.5	0.7	1.1	1.4	1.7	3.1	5.3	9.6

All races 3/, female

2006.....	*	*	0.2	0.4	0.3	0.6	1.0	1.4	2.7	6.5
2005.....	*	0.2	0.3	0.3	0.4	0.7	1.2	1.8	2.7	6.6
2004.....	*	*	0.2	0.3	0.5	0.8	1.0	1.3	2.3	6.4
2003.....	*	0.2	0.2	*	0.6	0.5	1.0	1.6	2.7	6.1
2002.....	*	*	0.2	0.3	0.5	0.6	1.2	1.9	2.7	5.9
2001.....	*	*	0.2	0.3	0.4	0.8	1.2	1.4	2.5	6.4
2000.....	*	*	0.2	*	0.7	0.8	1.0	1.6	2.9	6.7
1999.....	*	0.3	*	0.4	0.6	0.9	1.3	2.1	4.3	8.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified disorders of circulatory system (I83-I99)

White, both sexes

2006 4/.....	0.1	0.2	0.4	0.4	0.5	0.7	1.1	1.7	2.6	6.3
2005 4/.....	0.2	0.4	0.6	0.5	0.6	0.9	1.4	1.7	2.9	6.5
2004 4/.....	0.1	0.4	0.4	0.5	0.7	0.9	1.1	1.7	2.6	6.0
2003 4/.....	0.1	0.3	0.4	0.4	0.6	0.6	1.2	1.7	2.7	5.8
2002.....	0.1	0.2	0.3	0.4	0.6	0.7	1.2	2.0	3.1	5.9
2001.....	0.2	0.3	0.3	0.5	0.6	0.9	1.3	1.6	3.0	6.5
2000.....	0.2	0.3	0.4	0.4	0.8	0.8	1.3	1.9	3.0	6.7
1999.....	0.2	0.3	0.3	0.5	0.8	1.0	1.5	2.4	4.6	8.8

White, male

2006 4/.....	*	0.3	0.5	0.4	0.7	0.9	1.2	2.1	2.6	6.2
2005 4/.....	0.3	0.5	0.9	0.7	0.9	1.2	1.6	1.7	3.1	5.5
2004 4/.....	*	0.6	0.5	0.7	0.9	1.0	1.3	2.2	3.1	5.4
2003 4/.....	0.2	0.4	0.5	0.6	0.6	0.7	1.4	2.1	2.7	5.2
2002.....	*	0.4	0.5	0.5	0.8	0.9	1.2	2.1	3.6	5.5
2001.....	0.2	0.5	0.5	0.6	0.8	1.0	1.4	2.1	3.7	6.6
2000.....	0.3	0.5	0.6	0.5	1.0	0.9	1.9	2.2	3.3	7.1
1999.....	0.3	0.3	0.4	0.7	1.1	1.2	1.7	2.8	5.0	9.3

White, female

2006 4/.....	*	*	*	0.4	*	0.6	1.0	1.4	2.7	6.4
2005 4/.....	*	*	0.3	0.3	0.4	0.7	1.3	1.7	2.8	6.9
2004 4/.....	*	*	*	0.3	0.5	0.8	0.9	1.3	2.3	6.3
2003 4/.....	*	*	*	*	0.6	0.5	1.0	1.4	2.8	6.1
2002.....	*	*	*	*	*	0.5	1.2	1.9	2.8	6.0
2001.....	*	*	*	*	0.4	0.9	1.3	1.3	2.6	6.4
2000.....	*	*	*	*	0.6	0.7	0.9	1.6	2.8	6.5
1999.....	*	0.3	*	0.4	0.4	0.7	1.3	2.0	4.3	8.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

X. Diseases of the respiratory system (J00-J98)

All races 3/, both sexes

2006.....	7.2	12.6	23.8	42.2	86.5	163.3	288.7	482.9	772.0	1,391.2
2005.....	7.3	14.0	24.4	44.5	92.7	171.7	310.9	517.5	816.8	1,540.5
2004.....	7.1	12.6	22.5	43.2	90.4	166.1	302.7	503.4	791.7	1,497.1
2003.....	7.3	13.6	23.5	44.5	94.9	176.0	311.6	524.2	827.7	1,625.4
2002.....	7.6	13.1	22.8	45.7	91.4	172.4	315.3	528.4	849.0	1,663.7
2001.....	7.4	12.7	23.1	46.9	93.4	178.6	319.0	526.3	833.1	1,671.8
2000.....	8.0	13.2	23.3	48.3	94.2	181.4	324.3	529.4	874.7	1,734.0
1999.....	7.4	12.9	22.4	49.4	97.1	185.1	330.2	537.8	873.1	1,733.0

All races 3/, male

2006.....	7.5	13.1	26.5	47.3	97.8	183.8	337.6	591.2	982.5	1,752.9
2005.....	7.6	15.3	27.4	49.2	104.7	198.2	365.8	632.0	1,031.2	1,900.6
2004.....	7.5	13.6	25.8	47.6	101.4	192.5	358.3	620.1	1,012.1	1,915.3
2003.....	7.5	15.2	26.2	49.5	106.5	202.6	371.4	646.6	1,052.6	2,057.2
2002.....	8.3	14.2	25.5	50.2	102.9	197.5	379.7	661.7	1,094.2	2,150.0
2001.....	8.0	14.3	25.7	51.9	104.0	204.9	385.2	662.7	1,080.0	2,187.2
2000.....	8.9	14.8	25.7	54.2	106.1	210.7	394.4	666.6	1,158.0	2,344.4
1999.....	7.9	14.6	24.5	54.1	111.1	220.6	411.8	691.1	1,170.2	2,356.4

All races 3/, female

2006.....	7.0	12.2	21.1	37.4	76.2	145.4	248.8	403.6	640.4	1,222.0
2005.....	7.1	12.8	21.5	40.0	81.7	148.6	266.4	434.3	683.7	1,375.2
2004.....	6.8	11.7	19.3	39.1	80.4	143.2	257.9	419.5	655.9	1,309.0
2003.....	7.2	12.1	20.8	39.8	84.3	153.0	263.6	437.0	690.2	1,434.5
2002.....	6.8	12.0	20.2	41.5	81.1	150.7	263.8	434.3	699.8	1,452.7
2001.....	6.8	11.0	20.6	42.2	83.9	155.9	266.4	430.8	685.2	1,456.2
2000.....	7.1	11.6	21.0	42.9	83.5	156.3	269.0	433.8	707.6	1,485.4
1999.....	7.0	11.4	20.5	45.0	84.3	154.8	266.4	431.1	699.3	1,481.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

X. Diseases of the respiratory system (J00-J98)

White, both sexes

2006 4/.....	6.6	11.9	22.9	42.2	88.7	169.7	304.2	502.3	799.2	1,436.6
2005 4/.....	6.7	13.0	23.1	44.1	94.8	177.8	324.3	536.3	845.4	1,590.1
2004 4/.....	6.5	11.5	21.4	42.9	92.2	172.9	316.6	521.6	817.9	1,541.7
2003 4/.....	6.4	12.5	22.4	43.9	96.7	183.2	323.5	541.5	855.8	1,676.3
2002.....	6.7	12.1	21.6	45.3	93.0	179.4	326.6	545.8	873.3	1,713.3
2001.....	6.5	11.4	21.7	46.7	95.4	184.8	329.8	542.3	856.5	1,713.5
2000.....	7.0	11.6	21.8	47.7	95.6	185.4	333.7	544.9	898.9	1,783.5
1999.....	6.1	11.2	21.2	48.7	98.7	190.9	338.4	551.4	897.0	1,778.5

White, male

2006 4/.....	6.8	12.3	25.2	47.0	97.9	187.3	349.2	605.7	1,006.0	1,798.4
2005 4/.....	6.9	14.0	25.6	47.5	105.1	201.7	375.4	643.7	1,053.9	1,946.0
2004 4/.....	6.6	12.3	24.4	46.1	101.3	196.2	367.6	632.9	1,035.5	1,960.9
2003 4/.....	6.4	14.0	24.9	48.0	106.0	207.1	377.9	655.2	1,072.5	2,106.7
2002.....	7.3	13.0	23.9	48.8	102.3	202.3	386.7	672.5	1,114.4	2,201.7
2001.....	7.1	13.1	23.6	51.2	103.7	208.2	391.6	672.0	1,098.6	2,230.5
2000.....	7.8	12.9	23.5	52.6	105.9	210.3	398.5	676.6	1,179.6	2,402.2
1999.....	6.3	12.6	22.5	52.3	110.8	223.6	415.4	699.7	1,188.8	2,409.4

White, female

2006 4/.....	6.3	11.5	20.7	37.6	80.2	154.0	266.8	425.2	668.6	1,266.7
2005 4/.....	6.5	12.1	20.6	40.9	85.2	156.5	282.1	456.9	714.8	1,426.3
2004 4/.....	6.3	10.6	18.5	39.8	83.9	152.2	274.6	440.1	682.8	1,352.8
2003 4/.....	6.3	11.0	19.9	40.0	88.1	162.1	279.0	459.2	722.2	1,485.6
2002.....	6.0	11.3	19.3	42.0	84.4	159.3	277.5	455.0	725.5	1,501.2
2001.....	5.9	9.6	19.8	42.3	87.8	164.3	279.7	450.5	710.2	1,497.0
2000.....	6.2	10.3	20.2	43.1	86.1	163.6	281.7	452.2	731.9	1,532.2
1999.....	5.9	9.9	19.9	45.3	87.5	162.3	277.1	447.2	724.8	1,525.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
X. Diseases of the respiratory system (J00-J98)										
Black, both sexes										
2006 4/.....	13.1	21.0	36.7	54.8	93.3	152.7	226.6	377.8	579.3	993.1
2005 4/.....	13.2	24.4	40.3	60.2	99.1	166.7	262.9	431.8	603.7	1,099.0
2004 4/.....	13.7	23.1	36.8	57.7	100.1	154.3	244.8	404.7	595.6	1,082.3
2003 4/.....	15.1	24.7	37.8	60.3	102.7	160.3	264.1	427.8	606.9	1,135.5
2002.....	15.5	23.1	36.6	61.9	102.5	154.4	271.2	424.1	670.3	1,204.5
2001.....	14.9	24.9	39.3	60.7	99.0	162.5	275.3	430.7	647.1	1,258.0
2000.....	16.0	27.1	40.4	65.3	106.0	183.3	288.6	427.5	662.7	1,239.4
1999.....	17.6	28.0	38.2	68.0	105.5	173.1	302.0	456.5	655.6	1,298.0
Black, male										
2006 4/.....	13.5	22.2	44.6	64.3	127.2	201.8	312.2	543.2	835.1	1,316.3
2005 4/.....	14.2	28.4	48.0	76.9	129.8	224.1	356.7	625.8	876.4	1,475.2
2004 4/.....	15.2	26.0	44.3	74.2	131.2	215.1	344.8	585.5	873.1	1,501.1
2003 4/.....	15.4	27.4	43.2	76.0	137.5	215.7	378.7	640.4	937.6	1,577.5
2002.....	17.9	26.6	43.7	77.9	135.8	204.2	378.9	634.7	986.6	1,685.3
2001.....	16.2	27.3	47.7	72.0	130.6	218.5	390.5	653.7	981.6	1,775.7
2000.....	18.5	32.3	48.7	82.0	134.0	254.7	419.2	634.6	980.4	1,808.3
1999.....	19.8	33.3	47.4	83.9	141.1	237.6	438.2	682.9	1,034.1	1,900.6
Black, female										
2006 4/.....	12.8	20.0	30.0	47.0	66.9	116.5	167.8	278.8	444.2	860.9
2005 4/.....	12.3	21.0	33.8	46.5	75.1	124.1	198.7	316.0	459.6	947.2
2004 4/.....	12.3	20.6	30.5	44.1	75.9	109.2	177.0	297.6	449.4	916.3
2003 4/.....	14.9	22.4	33.3	47.4	75.5	119.2	186.8	301.4	433.6	963.7
2002.....	13.4	20.1	30.5	48.8	76.5	117.5	198.9	298.3	505.7	1,020.3
2001.....	13.8	22.8	32.1	51.4	74.1	121.1	198.6	297.6	476.7	1,066.0
2000.....	13.7	22.6	33.4	51.7	84.0	130.6	201.8	302.8	502.7	1,027.6
1999.....	15.6	23.5	30.4	55.0	77.5	125.3	211.5	320.1	468.0	1,072.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of the respiratory system (J10-J98)										
All races 3/, both sexes										
2006.....	7.2	12.6	23.7	42.1	86.5	163.2	288.6	482.7	771.8	1,390.0
2005.....	7.3	14.0	24.4	44.4	92.6	171.7	310.8	517.3	816.5	1,539.0
2004.....	7.1	12.6	22.5	43.2	90.3	166.1	302.6	503.2	791.4	1,496.3
2003.....	7.3	13.6	23.4	44.5	94.8	176.0	311.5	524.1	827.4	1,623.9
2002.....	7.5	13.1	22.7	45.6	91.4	172.3	315.1	528.3	848.8	1,662.6
2001.....	7.4	12.6	23.1	46.8	93.4	178.5	318.9	526.1	832.8	1,670.6
2000.....	8.0	13.2	23.2	48.3	94.2	181.3	324.1	529.1	874.3	1,732.2
1999.....	7.4	12.9	22.4	49.4	97.0	185.1	330.1	537.5	872.6	1,731.2
All races 3/, male										
2006.....	7.4	13.1	26.5	47.3	97.8	183.7	337.6	590.9	982.2	1,751.8
2005.....	7.5	15.3	27.4	49.2	104.6	198.1	365.7	631.8	1,030.8	1,899.9
2004.....	7.4	13.5	25.8	47.5	101.4	192.5	358.2	620.0	1,011.8	1,914.6
2003.....	7.4	15.1	26.2	49.5	106.4	202.5	371.4	646.6	1,052.2	2,055.5
2002.....	8.3	14.2	25.4	50.1	102.9	197.4	379.5	661.6	1,094.0	2,149.3
2001.....	7.9	14.3	25.7	51.9	104.0	204.9	385.2	662.6	1,079.8	2,186.6
2000.....	8.9	14.8	25.6	54.2	106.1	210.6	394.3	666.5	1,157.7	2,343.3
1999.....	7.8	14.5	24.5	54.1	111.0	220.5	411.7	690.9	1,169.5	2,355.3
All races 3/, female										
2006.....	7.0	12.1	21.1	37.3	76.2	145.3	248.8	403.5	640.3	1,220.8
2005.....	7.0	12.8	21.5	39.9	81.7	148.6	266.3	434.2	683.4	1,373.3
2004.....	6.8	11.7	19.3	39.1	80.3	143.1	257.9	419.3	655.6	1,308.1
2003.....	7.2	12.1	20.8	39.7	84.2	153.0	263.5	436.9	689.9	1,433.1
2002.....	6.8	11.9	20.1	41.4	81.0	150.6	263.7	434.1	699.7	1,451.5
2001.....	6.8	11.0	20.6	42.1	83.8	155.9	266.3	430.6	684.9	1,454.6
2000.....	7.1	11.6	20.9	42.8	83.5	156.2	268.8	433.5	707.2	1,483.4
1999.....	7.0	11.3	20.4	45.0	84.3	154.8	266.3	430.9	698.8	1,479.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of the respiratory system (J10-J98)										
White, both sexes										
2006 4/.....	6.6	11.9	22.9	42.2	88.7	169.6	304.1	502.1	799.0	1,435.3
2005 4/.....	6.7	13.0	23.1	44.1	94.7	177.8	324.2	536.1	845.0	1,588.5
2004 4/.....	6.4	11.4	21.4	42.9	92.2	172.8	316.5	521.4	817.6	1,540.9
2003 4/.....	6.3	12.5	22.4	43.9	96.7	183.2	323.5	541.4	855.4	1,674.8
2002.....	6.6	12.1	21.6	45.3	93.0	179.4	326.4	545.6	873.2	1,712.2
2001.....	6.5	11.3	21.7	46.6	95.4	184.8	329.7	542.2	856.2	1,712.1
2000.....	6.9	11.6	21.8	47.7	95.5	185.3	333.5	544.6	898.5	1,781.7
1999.....	6.1	11.2	21.2	48.7	98.6	190.8	338.3	551.1	896.5	1,776.7
White, male										
2006 4/.....	6.8	12.2	25.1	47.0	97.9	187.2	349.1	605.4	1,005.7	1,797.4
2005 4/.....	6.9	13.9	25.6	47.5	105.0	201.7	375.3	643.5	1,053.5	1,945.3
2004 4/.....	6.6	12.3	24.4	46.1	101.3	196.2	367.5	632.8	1,035.2	1,960.3
2003 4/.....	6.4	14.0	24.9	48.0	105.9	207.0	377.9	655.2	1,072.2	2,105.2
2002.....	7.3	12.9	23.9	48.8	102.3	202.2	386.6	672.4	1,114.3	2,201.0
2001.....	7.0	13.1	23.6	51.2	103.6	208.1	391.6	671.9	1,098.3	2,229.8
2000.....	7.8	12.9	23.5	52.6	105.8	210.3	398.3	676.5	1,179.3	2,401.1
1999.....	6.2	12.5	22.5	52.3	110.8	223.5	415.3	699.4	1,188.2	2,408.6
White, female										
2006 4/.....	6.3	11.5	20.7	37.5	80.1	154.0	266.7	425.1	668.4	1,265.3
2005 4/.....	6.5	12.1	20.6	40.8	85.2	156.5	282.0	456.7	714.4	1,424.2
2004 4/.....	6.3	10.6	18.4	39.8	83.8	152.1	274.6	439.9	682.5	1,351.9
2003 4/.....	6.3	11.0	19.9	40.0	88.1	162.1	279.0	459.1	721.8	1,484.2
2002.....	6.0	11.2	19.3	41.9	84.3	159.2	277.4	454.8	725.4	1,499.8
2001.....	5.9	9.6	19.8	42.3	87.8	164.2	279.6	450.3	709.9	1,495.4
2000.....	6.1	10.3	20.2	43.1	86.1	163.5	281.5	451.8	731.5	1,530.1
1999.....	5.9	9.9	19.8	45.3	87.4	162.3	277.0	447.0	724.3	1,522.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of the respiratory system (J10-J98)

Black, both sexes

2006 4/.....	13.1	20.9	36.6	54.8	93.2	152.6	226.6	377.7	579.3	992.6
2005 4/.....	13.0	24.4	40.2	60.1	99.1	166.5	262.8	431.4	603.4	1,098.2
2004 4/.....	13.5	23.1	36.7	57.6	100.1	154.2	244.7	404.7	595.4	1,081.5
2003 4/.....	15.0	24.6	37.7	60.3	102.4	160.1	264.1	427.8	606.9	1,134.0
2002.....	15.4	23.0	36.5	61.8	102.5	154.2	271.2	424.1	669.8	1,204.5
2001.....	14.9	24.8	39.2	60.6	98.8	162.5	275.2	430.5	647.1	1,257.1
2000.....	16.0	27.0	40.4	65.3	106.0	183.1	288.6	427.5	662.4	1,238.8
1999.....	17.6	28.0	38.1	67.9	105.3	173.1	301.6	456.5	655.3	1,296.4

Black, male

2006 4/.....	13.5	22.2	44.5	64.3	127.1	201.5	312.2	543.2	835.1	1,314.5
2005 4/.....	14.0	28.4	47.8	76.9	129.8	224.1	356.7	624.9	876.4	1,473.4
2004 4/.....	15.0	26.0	44.2	74.0	131.2	215.1	344.4	585.5	872.3	1,500.1
2003 4/.....	15.2	27.3	43.0	76.0	136.9	215.2	378.7	640.4	937.6	1,574.4
2002.....	17.8	26.6	43.5	77.9	135.8	203.9	378.9	634.7	985.8	1,685.3
2001.....	16.1	27.2	47.7	71.8	130.4	218.5	390.5	653.7	981.6	1,775.7
2000.....	18.5	32.3	48.7	82.0	134.0	254.2	419.2	634.6	979.5	1,806.0
1999.....	19.8	33.2	47.4	83.9	140.7	237.6	437.9	682.9	1,034.1	1,895.9

Black, female

2006 4/.....	12.8	19.8	30.0	47.0	66.8	116.5	167.8	278.5	444.2	860.9
2005 4/.....	12.1	21.0	33.8	46.3	75.1	123.7	198.5	316.0	459.2	946.8
2004 4/.....	12.2	20.6	30.5	44.1	75.9	109.0	177.0	297.6	449.4	915.5
2003 4/.....	14.8	22.3	33.1	47.4	75.5	119.2	186.8	301.4	433.6	962.9
2002.....	13.4	20.0	30.5	48.6	76.5	117.3	198.9	298.3	505.3	1,020.3
2001.....	13.8	22.8	32.0	51.4	73.9	121.1	198.4	297.3	476.7	1,064.7
2000.....	13.7	22.5	33.3	51.7	84.0	130.6	201.8	302.8	502.7	1,027.6
1999.....	15.6	23.5	30.3	54.8	77.5	125.3	211.1	320.1	467.6	1,072.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza (J10-J11)										
All races 3/, both sexes										
2006.....	*	*	*	0.1	0.2	0.3	0.6	1.3	2.5	7.0
2005.....	*	*	0.1	0.1	0.3	0.4	1.1	2.4	5.4	19.3
2004.....	*	*	*	0.1	0.3	0.3	0.8	1.6	3.2	11.4
2003.....	*	0.1	0.2	0.3	0.3	0.7	1.3	2.7	4.7	17.0
2002.....	*	*	*	*	*	*	0.4	0.8	2.2	8.8
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	0.1	0.1	0.3	0.5	0.8	1.4	2.5	6.1	20.2
1999.....	*	*	*	0.2	0.4	0.6	1.2	2.1	6.1	21.2
All races 3/, male										
2006.....	*	*	*	*	*	*	0.7	1.3	3.0	8.1
2005.....	*	*	*	*	0.4	*	1.1	2.5	5.3	18.2
2004.....	*	*	*	*	*	*	1.1	1.8	3.3	11.1
2003.....	*	*	*	0.3	*	0.8	1.3	2.9	4.9	17.0
2002.....	*	*	*	*	*	*	*	0.9	2.6	8.9
2001.....	*	*	*	*	*	*	*	*	*	2.2
2000.....	*	*	*	0.3	0.5	1.0	1.6	2.7	6.3	20.7
1999.....	*	*	*	*	0.5	0.6	1.2	2.4	7.3	19.9
All races 3/, female										
2006.....	*	*	*	*	*	*	0.5	1.3	2.2	6.5
2005.....	*	*	*	*	*	0.5	1.1	2.3	5.5	19.8
2004.....	*	*	*	*	*	*	0.6	1.4	3.1	11.5
2003.....	*	*	*	0.3	0.3	0.7	1.3	2.5	4.7	17.1
2002.....	*	*	*	*	*	*	*	0.7	2.0	8.7
2001.....	*	*	*	*	*	*	*	*	0.7	2.2
2000.....	*	*	*	*	0.5	0.6	1.2	2.5	6.0	19.9
1999.....	*	*	*	*	*	0.6	1.1	1.9	5.4	21.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Influenza (J10-J11)										
White, both sexes										
2006 4/.....	*	*	*	0.1	0.2	0.3	0.7	1.4	2.7	7.5
2005 4/.....	*	*	0.1	0.2	0.3	0.5	1.2	2.5	5.8	20.8
2004 4/.....	*	*	*	0.2	0.3	0.3	0.9	1.5	3.4	12.3
2003 4/.....	*	0.1	0.2	0.3	0.3	0.8	1.4	2.9	5.1	18.2
2002.....	*	*	*	*	*	*	0.4	0.9	2.4	9.4
2001.....	*	*	*	*	*	*	*	*	0.7	2.3
2000.....	*	*	0.1	0.3	0.5	0.8	1.4	2.7	6.5	21.4
1999.....	*	*	*	0.2	0.4	0.7	1.2	2.2	6.4	22.8
White, male										
2006 4/.....	*	*	*	*	*	*	0.8	1.3	3.1	8.6
2005 4/.....	*	*	*	*	*	*	1.2	2.7	5.6	19.2
2004 4/.....	*	*	*	*	*	*	1.1	1.7	3.6	12.1
2003 4/.....	*	*	*	0.3	*	0.9	1.4	3.1	5.2	18.3
2002.....	*	*	*	*	*	*	*	1.0	2.8	9.7
2001.....	*	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	*	0.5	1.0	1.7	2.7	6.7	21.6
1999.....	*	*	*	*	0.5	0.6	1.2	2.6	7.5	21.7
White, female										
2006 4/.....	*	*	*	*	*	*	0.5	1.4	2.4	7.1
2005 4/.....	*	*	*	*	*	0.5	1.2	2.4	5.9	21.5
2004 4/.....	*	*	*	*	*	*	0.7	1.4	3.3	12.4
2003 4/.....	*	*	*	*	0.4	0.7	1.4	2.8	5.0	18.1
2002.....	*	*	*	*	*	*	*	0.8	2.2	9.3
2001.....	*	*	*	*	*	*	*	*	0.7	2.3
2000.....	*	*	*	*	0.6	0.7	1.2	2.7	6.3	21.3
1999.....	*	*	*	*	*	0.7	1.1	2.0	5.7	23.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
All races 3/, both sexes										
2006.....	2.4	3.6	5.6	7.5	13.0	22.8	42.3	85.0	179.3	495.5
2005.....	2.6	4.2	6.0	8.6	14.3	24.1	47.5	93.7	197.3	574.7
2004.....	2.5	3.6	5.6	8.2	13.9	23.5	46.4	93.1	195.6	571.2
2003.....	2.6	4.1	6.2	7.9	14.9	25.8	48.1	100.6	212.1	649.1
2002.....	2.6	4.1	5.4	8.4	14.6	25.1	50.5	105.7	225.1	687.8
2001.....	2.6	3.9	5.3	8.1	13.8	24.3	49.2	100.9	216.1	683.4
2000.....	2.9	3.8	5.5	8.6	15.1	25.8	51.2	104.1	234.7	723.9
1999.....	2.7	4.0	5.2	8.2	13.9	24.3	49.3	103.8	227.7	730.6
All races 3/, male										
2006.....	2.7	4.2	7.0	9.3	16.3	27.8	51.8	106.9	223.8	578.3
2005.....	3.0	5.0	7.4	10.5	17.1	29.2	57.9	116.8	248.4	655.3
2004.....	2.7	4.2	7.0	9.7	17.0	29.4	57.6	116.5	249.1	666.1
2003.....	3.0	4.8	7.9	9.3	17.6	32.4	60.8	127.2	267.5	740.9
2002.....	3.0	4.9	6.7	10.1	17.8	30.9	63.4	133.0	285.0	796.9
2001.....	3.0	4.8	6.6	9.9	16.9	30.3	61.4	130.5	273.9	806.2
2000.....	3.7	4.6	6.8	10.6	18.4	31.9	64.1	131.5	302.6	871.3
1999.....	3.1	5.1	6.7	9.5	16.3	31.1	63.0	131.1	294.5	874.7
All races 3/, female										
2006.....	2.1	3.1	4.3	5.9	10.1	18.5	34.5	69.0	151.5	456.7
2005.....	2.3	3.4	4.7	6.8	11.7	19.6	39.1	77.0	165.6	537.6
2004.....	2.3	2.9	4.1	6.7	11.1	18.4	37.3	76.4	162.6	528.5
2003.....	2.2	3.4	4.7	6.5	12.4	20.1	37.9	81.7	178.2	608.5
2002.....	2.2	3.4	4.3	6.8	11.8	20.0	40.2	86.5	188.6	640.5
2001.....	2.2	2.9	4.1	6.3	11.0	19.1	39.4	80.2	181.5	632.0
2000.....	2.1	3.0	4.2	6.7	12.0	20.5	41.0	85.1	194.6	663.9
1999.....	2.3	2.9	3.9	7.0	11.7	18.6	38.6	84.8	188.7	672.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
White, both sexes										
2006 4/.....	2.2	3.2	5.1	7.1	12.5	21.4	41.9	84.2	178.8	501.7
2005 4/.....	2.4	3.8	5.4	7.8	13.4	22.8	45.8	92.6	198.2	585.6
2004 4/.....	2.2	3.2	4.9	7.5	13.1	22.0	45.3	91.5	194.9	578.4
2003 4/.....	2.3	3.6	5.6	7.1	13.7	24.4	46.5	99.5	213.3	661.1
2002.....	2.3	3.8	4.9	7.6	13.7	24.3	49.5	104.5	225.6	699.6
2001.....	2.1	3.3	4.8	7.4	12.9	23.2	48.0	98.3	215.8	692.7
2000.....	2.4	3.2	4.8	7.8	13.8	23.7	48.9	102.9	234.3	737.0
1999.....	2.1	3.3	4.6	7.4	12.6	23.4	47.6	103.3	229.2	741.6
White, male										
2006 4/.....	2.5	3.7	6.2	8.8	15.3	25.8	50.5	105.2	222.6	582.8
2005 4/.....	2.7	4.4	6.5	9.2	15.8	27.2	55.3	115.1	248.3	665.2
2004 4/.....	2.3	3.8	6.2	9.0	15.7	26.7	55.6	112.9	247.3	673.4
2003 4/.....	2.5	4.2	7.0	8.2	15.4	29.9	57.8	124.1	266.2	748.8
2002.....	2.7	4.3	6.0	9.0	16.3	29.6	61.9	130.4	283.7	806.2
2001.....	2.5	4.1	5.6	9.0	15.3	28.3	59.2	125.3	271.5	812.1
2000.....	3.1	3.8	5.7	9.2	16.7	28.6	60.6	129.1	299.7	884.4
1999.....	2.2	4.2	5.6	8.4	14.6	29.5	59.7	129.6	294.3	884.0
White, female										
2006 4/.....	1.9	2.8	4.1	5.5	9.9	17.4	34.8	68.5	151.2	463.7
2005 4/.....	2.1	3.1	4.3	6.5	11.1	19.0	38.0	75.9	166.9	548.9
2004 4/.....	2.1	2.6	3.7	6.1	10.6	17.8	36.9	75.9	162.3	535.6
2003 4/.....	2.0	3.0	4.2	6.0	12.1	19.5	37.2	81.7	180.7	622.3
2002.....	1.9	3.2	3.9	6.2	11.3	19.7	39.4	86.0	190.0	653.3
2001.....	1.8	2.4	3.9	5.9	10.6	18.8	38.9	79.3	182.2	642.8
2000.....	1.8	2.7	3.9	6.4	11.1	19.4	39.6	84.5	195.4	677.1
1999.....	1.9	2.5	3.5	6.5	10.8	18.0	37.9	84.8	190.7	684.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia (J12-J18)										
Black, both sexes										
2006 4/.....	4.1	7.1	10.7	12.9	20.1	39.5	50.9	99.0	195.0	432.7
2005 4/.....	4.9	7.8	12.0	16.7	24.1	39.0	67.8	114.9	197.1	467.2
2004 4/.....	4.9	6.9	11.5	15.5	23.5	39.5	61.9	115.6	214.1	502.5
2003 4/.....	5.4	8.3	12.4	15.4	27.0	41.0	69.6	121.7	203.6	519.6
2002.....	5.2	7.5	10.1	17.2	26.1	35.0	68.6	123.2	230.8	572.8
2001.....	5.8	9.0	10.7	15.4	23.1	34.1	65.7	133.7	222.9	579.7
2000.....	6.0	8.1	12.1	17.7	26.9	45.4	76.0	123.5	243.1	591.5
1999.....	6.8	9.7	11.6	16.0	26.4	36.5	73.9	117.8	217.0	629.1
Black, male										
2006 4/.....	4.5	8.3	15.3	16.1	29.2	49.9	68.9	138.7	241.6	502.2
2005 4/.....	6.1	10.2	15.7	23.6	32.5	53.7	89.8	143.9	257.7	524.3
2004 4/.....	5.9	8.4	15.8	19.1	32.9	61.1	86.4	165.6	297.0	597.1
2003 4/.....	7.0	10.0	16.5	20.8	39.5	60.8	100.2	176.6	283.5	638.1
2002.....	6.2	9.9	12.9	23.5	35.6	47.4	89.8	173.2	320.2	687.2
2001.....	7.0	11.4	15.7	20.6	33.2	49.6	94.9	203.5	309.0	712.6
2000.....	8.1	11.0	17.0	25.5	35.0	64.3	106.6	170.6	343.1	742.7
1999.....	8.9	13.3	17.1	21.7	33.9	51.4	108.2	159.6	307.8	800.3
Black, female										
2006 4/.....	3.7	6.0	6.9	10.3	13.1	31.8	38.5	75.3	170.4	404.3
2005 4/.....	3.8	5.7	8.8	10.9	17.6	28.0	52.8	97.6	165.1	444.1
2004 4/.....	4.1	5.7	7.9	12.6	16.2	23.6	45.3	86.0	170.5	465.0
2003 4/.....	4.1	6.7	8.9	11.0	17.1	26.3	49.0	89.0	161.7	473.5
2002.....	4.3	5.5	7.7	12.0	18.6	25.9	54.4	93.3	184.2	529.0
2001.....	4.7	6.9	6.5	11.2	15.2	22.7	46.2	92.1	179.1	530.5
2000.....	4.2	5.6	7.9	11.2	20.4	31.5	55.7	95.1	192.8	535.3
1999.....	5.0	6.6	7.0	11.4	20.5	25.4	51.1	92.7	172.0	565.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral pneumonia, not elsewhere classified (J12)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*	*	0.5	1.1
2005 4/.....	*	*	*	*	*	*	0.3	*	0.8	1.5
2004 4/.....	*	*	*	*	*	*	*	0.3	0.7	1.3
2003 4/.....	*	*	*	*	*	*	*	*	0.4	0.6
2002.....	*	*	*	*	*	*	0.3	0.4	0.8	1.8
2001.....	*	*	*	*	*	*	*	0.3	0.8	1.5
2000.....	*	*	*	*	*	0.3	0.3	0.6	1.0	3.0
1999.....	*	*	*	*	*	0.2	0.3	0.5	1.3	2.5

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	1.6
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	2.0
2001.....	*	*	*	*	*	*	*	*	1.4	2.0
2000.....	*	*	*	*	*	*	*	0.7	*	2.7
1999.....	*	*	*	*	*	*	*	0.8	1.8	1.9

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	1.1
2005 4/.....	*	*	*	*	*	*	*	*	0.7	1.5
2004 4/.....	*	*	*	*	*	*	*	*	*	1.5
2003 4/.....	*	*	*	*	*	*	*	*	*	2.2
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	1.4
2000.....	*	*	*	*	*	*	*	0.5	1.0	3.2
1999.....	*	*	*	*	*	*	*	*	1.0	2.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bacterial pneumonia (J13-J15)										
All races 3/, both sexes										
2006.....	0.4	0.5	1.0	1.1	1.8	2.8	4.4	8.3	14.1	27.7
2005.....	0.4	0.6	0.8	1.3	1.9	3.0	4.8	8.3	13.6	31.3
2004.....	0.3	0.5	0.8	1.2	1.7	2.6	4.3	7.9	13.9	30.1
2003.....	0.4	0.5	1.0	1.1	1.8	2.9	4.7	8.2	15.2	34.9
2002.....	0.4	0.6	0.7	1.2	1.6	2.6	4.7	8.3	15.5	35.7
2001.....	0.3	0.5	0.7	1.4	1.6	2.7	4.4	8.4	14.8	34.8
2000.....	0.5	0.6	0.8	1.3	1.9	2.7	4.9	8.1	15.3	35.7
1999.....	0.5	0.5	0.6	1.1	1.5	2.4	4.3	7.2	13.6	30.1
All races 3/, male										
2006.....	0.4	0.6	1.2	1.5	2.3	3.5	5.8	11.4	19.4	37.2
2005.....	0.4	0.7	1.1	1.6	2.0	3.7	5.9	10.4	19.0	43.1
2004.....	0.3	0.6	1.0	1.2	1.9	3.3	5.3	10.2	19.9	41.1
2003.....	0.5	0.6	1.3	1.3	2.2	3.8	6.3	10.6	20.5	45.7
2002.....	0.4	0.6	0.8	1.4	1.8	3.0	6.1	11.3	20.9	46.8
2001.....	0.5	0.6	0.9	1.6	2.0	3.6	5.6	11.7	20.6	46.8
2000.....	0.6	0.6	0.9	1.5	2.3	3.3	6.4	11.9	21.7	52.3
1999.....	0.5	0.6	0.8	1.2	1.6	3.0	5.5	9.0	18.4	40.4
All races 3/, female										
2006.....	0.4	0.5	0.7	0.7	1.4	2.1	3.2	6.0	10.7	23.3
2005.....	0.4	0.5	0.6	1.0	1.8	2.4	4.0	6.8	10.3	25.9
2004.....	0.4	0.4	0.6	1.1	1.6	2.0	3.4	6.2	10.3	25.2
2003.....	0.3	0.5	0.7	0.9	1.4	2.2	3.4	6.5	11.9	30.2
2002.....	0.4	0.5	0.6	0.9	1.4	2.3	3.6	6.2	12.2	31.0
2001.....	0.2	0.5	0.6	1.1	1.3	1.9	3.4	6.2	11.3	29.8
2000.....	0.4	0.5	0.6	1.1	1.6	2.1	3.7	5.4	11.6	29.0
1999.....	0.5	0.4	0.4	0.9	1.5	1.8	3.3	5.9	10.8	26.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bacterial pneumonia (J13-J15)										
White, both sexes										
2006 4/.....	0.4	0.4	0.9	1.1	1.8	2.6	4.5	8.2	13.9	27.8
2005 4/.....	0.4	0.6	0.8	1.1	1.8	3.0	4.7	8.2	13.7	31.7
2004 4/.....	0.3	0.5	0.7	1.1	1.6	2.5	4.2	7.5	13.8	29.9
2003 4/.....	0.4	0.5	0.9	1.0	1.6	2.9	4.6	8.0	15.3	34.9
2002.....	0.4	0.5	0.7	1.0	1.5	2.6	4.7	8.3	15.3	36.3
2001.....	0.3	0.5	0.7	1.4	1.5	2.6	4.2	8.1	14.8	35.0
2000.....	0.4	0.5	0.7	1.2	1.9	2.5	4.7	7.9	15.0	35.6
1999.....	0.4	0.5	0.5	1.0	1.4	2.3	4.1	7.1	13.6	30.3
White, male										
2006 4/.....	0.4	0.4	1.1	1.5	2.2	3.4	5.8	11.3	18.9	37.7
2005 4/.....	0.4	0.7	1.1	1.3	1.8	3.5	5.8	10.1	19.2	43.7
2004 4/.....	0.3	0.6	0.9	1.0	1.7	2.9	5.0	9.7	19.7	41.7
2003 4/.....	0.4	0.5	1.2	1.1	1.8	3.7	6.2	10.3	20.8	45.9
2002.....	0.4	0.5	0.7	1.2	1.7	2.9	5.9	11.1	20.1	47.9
2001.....	0.4	0.5	0.8	1.6	1.8	3.5	5.3	10.9	20.5	46.6
2000.....	0.6	0.6	0.9	1.4	2.1	3.0	6.1	11.4	21.7	51.1
1999.....	0.4	0.5	0.7	1.1	1.5	2.9	5.2	8.9	18.1	40.1
White, female										
2006 4/.....	0.3	0.5	0.7	0.7	1.4	1.9	3.5	5.9	10.6	23.2
2005 4/.....	0.3	0.5	0.6	0.8	1.8	2.5	3.9	6.8	10.3	26.2
2004 4/.....	0.3	0.4	0.6	1.1	1.6	2.0	3.5	5.9	10.2	24.7
2003 4/.....	0.3	0.5	0.7	0.9	1.4	2.2	3.4	6.4	11.8	30.1
2002.....	0.3	0.5	0.6	0.8	1.4	2.3	3.7	6.3	12.3	31.2
2001.....	*	0.5	0.7	1.1	1.2	1.8	3.4	6.1	11.3	30.2
2000.....	0.3	0.4	0.6	1.0	1.6	2.0	3.6	5.5	11.0	29.2
1999.....	0.4	0.4	0.4	0.9	1.3	1.8	3.2	5.9	11.0	26.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bacterial pneumonia (J13-J15)										
Black, both sexes										
2006 4/.....	*	1.4	1.7	1.1	2.0	4.6	3.6	8.9	15.6	29.1
2005 4/.....	*	0.7	1.0	3.2	2.9	3.5	6.8	9.2	13.1	26.2
2004 4/.....	*	0.9	1.7	2.1	2.8	4.3	5.4	11.8	15.6	35.0
2003 4/.....	0.8	1.0	1.6	2.2	3.4	3.4	4.9	10.7	12.9	35.0
2002.....	0.8	1.1	1.0	2.3	2.2	2.9	5.4	8.4	16.0	28.5
2001.....	0.7	*	1.2	1.6	3.0	3.3	6.1	12.3	14.1	32.5
2000.....	*	1.0	1.3	2.0	2.8	4.6	7.1	8.9	17.5	34.7
1999.....	1.1	1.0	*	2.1	2.6	2.8	6.9	8.2	12.1	25.2
Black, male										
2006 4/.....	*	1.8	2.6	*	*	4.7	*	13.9	19.9	36.6
2005 4/.....	*	*	*	4.0	3.6	5.7	8.2	11.2	15.5	33.5
2004 4/.....	*	*	2.4	3.3	3.9	7.0	8.1	16.4	22.2	41.7
2003 4/.....	*	*	2.2	3.2	5.4	*	7.6	16.0	*	41.7
2002.....	*	*	*	3.4	*	*	7.7	13.3	26.0	32.2
2001.....	*	*	*	*	*	5.1	10.1	21.4	18.5	49.9
2000.....	*	*	*	*	4.1	5.5	10.7	16.1	21.1	60.0
1999.....	*	*	*	*	*	*	9.8	10.9	20.9	38.7
Black, female										
2006 4/.....	*	*	*	*	*	4.5	*	5.9	13.4	26.0
2005 4/.....	*	*	*	2.4	*	*	5.8	8.0	11.9	23.3
2004 4/.....	*	*	*	*	*	*	*	9.2	12.1	32.3
2003 4/.....	*	*	*	*	*	*	*	7.6	12.4	32.4
2002.....	*	*	*	*	*	*	*	5.5	10.7	27.1
2001.....	*	*	*	*	*	*	*	6.9	11.9	26.1
2000.....	*	*	*	*	*	4.0	4.7	*	15.7	25.3
1999.....	*	*	*	*	*	*	5.0	6.6	*	20.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia due to other or unspecified organisms (J16,J18)										
All races 3/, both sexes										
2006.....	2.0	3.0	4.7	6.4	11.2	20.0	37.7	76.5	164.8	466.8
2005.....	2.2	3.5	5.1	7.3	12.3	21.0	42.4	85.2	183.0	541.9
2004.....	2.1	3.0	4.7	7.0	12.1	20.8	42.0	85.0	181.0	539.8
2003.....	2.2	3.5	5.2	6.7	13.0	22.8	43.2	92.0	196.3	612.1
2002.....	2.2	3.5	4.7	7.2	13.0	22.4	45.5	97.0	208.8	650.3
2001.....	2.2	3.3	4.6	6.7	12.1	21.5	44.7	92.1	200.6	647.1
2000.....	2.4	3.2	4.7	7.2	13.0	22.9	46.0	95.5	218.3	685.3
1999.....	2.2	3.5	4.6	7.0	12.2	21.7	44.8	96.1	212.9	698.1
All races 3/, male										
2006.....	2.3	3.6	5.8	7.7	13.9	24.2	45.8	95.2	203.9	540.2
2005.....	2.6	4.3	6.2	8.9	15.0	25.4	51.7	106.1	228.6	610.7
2004.....	2.3	3.6	6.0	8.5	15.1	26.0	52.2	106.1	228.3	623.9
2003.....	2.5	4.2	6.5	8.0	15.4	28.5	54.2	116.2	246.2	693.0
2002.....	2.6	4.2	5.8	8.6	15.9	27.8	56.9	121.3	263.1	748.2
2001.....	2.5	4.2	5.7	8.2	14.9	26.5	55.7	118.3	252.1	757.4
2000.....	3.1	3.9	5.8	9.0	16.0	28.3	57.5	118.9	279.9	816.5
1999.....	2.6	4.5	5.8	8.1	14.5	27.9	57.2	121.2	274.4	832.5
All races 3/, female										
2006.....	1.7	2.5	3.6	5.2	8.7	16.3	31.2	62.8	140.3	432.4
2005.....	1.9	2.8	4.1	5.8	9.8	17.1	35.0	70.0	154.6	510.3
2004.....	1.9	2.5	3.5	5.6	9.5	16.2	33.8	69.8	151.8	502.0
2003.....	1.9	2.9	3.9	5.6	10.8	17.8	34.4	74.8	165.9	576.3
2002.....	1.7	2.8	3.6	5.8	10.3	17.7	36.4	79.8	175.8	607.9
2001.....	1.9	2.4	3.5	5.2	9.6	17.1	36.0	73.8	169.8	601.0
2000.....	1.7	2.5	3.6	5.6	10.3	18.2	37.0	79.2	182.0	631.9
1999.....	1.8	2.5	3.4	6.0	10.1	16.5	35.0	78.7	177.0	643.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia due to other or unspecified organisms (J16,J18)										
White, both sexes										
2006 4/.....	1.8	2.8	4.2	6.0	10.7	18.7	37.3	75.7	164.5	472.8
2005 4/.....	2.0	3.2	4.5	6.7	11.5	19.7	40.8	84.1	183.7	552.3
2004 4/.....	1.9	2.7	4.2	6.4	11.4	19.4	41.1	83.8	180.3	547.1
2003 4/.....	1.9	3.1	4.6	6.1	12.0	21.4	41.6	91.1	197.4	624.0
2002.....	1.9	3.2	4.2	6.5	12.1	21.7	44.5	95.8	209.5	661.6
2001.....	1.8	2.7	4.0	6.0	11.3	20.5	43.6	89.9	200.3	656.2
2000.....	2.0	2.7	4.0	6.5	11.8	21.0	43.9	94.4	218.3	698.3
1999.....	1.7	2.9	4.0	6.4	11.1	20.8	43.2	95.8	214.3	708.8
White, male										
2006 4/.....	2.1	3.2	5.1	7.3	13.0	22.3	44.6	93.6	203.1	544.1
2005 4/.....	2.3	3.7	5.4	7.8	13.9	23.5	49.2	104.6	228.2	619.9
2004 4/.....	2.0	3.2	5.3	7.9	14.0	23.7	50.5	103.1	226.7	630.7
2003 4/.....	2.1	3.7	5.8	7.1	13.6	26.1	51.4	113.4	244.6	700.5
2002.....	2.3	3.8	5.2	7.7	14.5	26.6	55.6	118.8	262.5	756.4
2001.....	2.1	3.5	4.8	7.3	13.5	24.6	53.8	113.8	249.6	763.5
2000.....	2.5	3.3	4.8	7.7	14.5	25.3	54.2	117.0	276.9	830.6
1999.....	1.9	3.7	4.9	7.2	12.9	26.4	54.2	119.9	274.5	841.9
White, female										
2006 4/.....	1.6	2.3	3.3	4.7	8.5	15.4	31.2	62.4	140.1	439.3
2005 4/.....	1.8	2.7	3.7	5.6	9.2	16.4	33.9	68.9	155.9	521.2
2004 4/.....	1.7	2.2	3.1	5.0	9.0	15.7	33.3	69.6	151.5	509.4
2003 4/.....	1.6	2.5	3.5	5.1	10.5	17.3	33.6	75.0	168.3	590.1
2002.....	1.6	2.7	3.2	5.3	9.8	17.3	35.5	79.3	177.0	620.4
2001.....	1.6	2.0	3.2	4.7	9.3	16.9	35.4	73.0	170.5	611.3
2000.....	1.5	2.2	3.2	5.3	9.4	17.2	35.7	78.5	183.3	644.6
1999.....	1.5	2.1	3.1	5.6	9.4	15.9	34.5	78.8	178.7	655.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonia due to other or unspecified organisms (J16,J18)										
Black, both sexes										
2006 4/.....	3.6	5.7	9.0	11.7	18.1	34.8	47.0	90.1	179.1	403.4
2005 4/.....	4.3	7.0	11.0	13.5	21.0	35.3	60.8	105.7	183.7	440.5
2004 4/.....	4.3	6.0	9.8	13.3	20.6	35.1	56.5	103.8	198.5	467.3
2003 4/.....	4.6	7.2	10.7	13.2	23.4	37.7	64.7	110.9	190.1	484.0
2002.....	4.4	6.4	9.1	14.9	23.9	32.2	63.1	114.6	214.6	543.4
2001.....	5.1	8.2	9.5	13.8	20.1	30.6	59.5	121.4	208.6	546.9
2000.....	5.4	7.1	10.7	15.6	23.8	40.7	68.7	114.6	225.3	556.5
1999.....	5.7	8.6	10.6	13.9	23.8	33.5	67.0	109.1	204.3	602.6
Black, male										
2006 4/.....	4.1	6.6	12.7	14.1	26.0	45.0	62.7	124.8	221.8	464.7
2005 4/.....	5.5	9.6	14.4	19.5	28.5	47.8	81.0	132.8	241.6	490.8
2004 4/.....	5.3	7.5	13.4	15.7	29.0	54.0	78.3	149.2	274.8	554.3
2003 4/.....	5.6	8.8	14.2	17.5	33.7	56.4	92.6	160.7	268.0	595.3
2002.....	5.5	8.6	11.7	20.1	32.7	44.4	82.1	159.4	293.5	652.9
2001.....	5.9	10.1	13.7	18.6	29.3	44.2	84.5	182.1	290.5	661.5
2000.....	7.5	10.0	15.3	22.6	30.6	58.8	95.6	154.5	322.0	682.7
1999.....	7.7	11.8	15.4	18.8	31.6	47.5	98.4	147.2	285.1	760.5
Black, female										
2006 4/.....	3.2	4.9	5.9	9.7	12.0	27.3	36.1	69.4	156.6	378.4
2005 4/.....	3.3	4.8	8.1	8.5	15.2	26.1	47.0	89.6	153.2	420.1
2004 4/.....	3.4	4.8	6.8	11.4	14.1	21.0	41.7	76.8	158.4	432.8
2003 4/.....	3.7	5.8	7.7	9.6	15.3	23.8	45.9	81.4	149.3	440.7
2002.....	3.4	4.5	6.9	10.6	17.0	23.1	50.4	87.8	173.5	501.5
2001.....	4.3	6.4	5.9	9.9	12.8	20.5	42.8	85.1	166.8	504.4
2000.....	3.5	4.6	6.8	9.9	18.5	27.3	50.8	90.6	176.6	509.5
1999.....	4.0	5.9	6.6	9.8	17.6	23.1	46.1	86.1	164.2	543.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute lower respiratory infections (J20-J22)

All races 3/, both sexes

2006.....	*	*	*	*	*	*	*	0.3	0.6	2.0
2005.....	*	*	*	*	*	*	*	0.4	0.9	3.3
2004.....	*	*	*	*	*	*	*	0.3	0.6	2.6
2003.....	*	*	*	*	*	*	*	0.5	0.8	3.5
2002.....	*	*	*	*	*	*	0.3	0.3	0.8	3.3
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.3
2000.....	*	*	*	*	*	*	0.3	0.6	0.8	4.5
1999.....	*	*	*	*	*	*	0.2	0.6	1.3	6.0

All races 3/, male

2006.....	*	*	*	*	*	*	*	*	*	1.8
2005.....	*	*	*	*	*	*	*	*	1.3	3.1
2004.....	*	*	*	*	*	*	*	*	*	2.5
2003.....	*	*	*	*	*	*	*	*	*	2.4
2002.....	*	*	*	*	*	*	*	*	*	2.8
2001.....	*	*	*	*	*	*	*	*	*	2.3
2000.....	*	*	*	*	*	*	*	0.7	*	3.8
1999.....	*	*	*	*	*	*	*	0.7	1.6	4.0

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	0.7	2.1
2005.....	*	*	*	*	*	*	*	*	0.6	3.5
2004.....	*	*	*	*	*	*	*	*	0.7	2.7
2003.....	*	*	*	*	*	*	*	*	0.8	3.9
2002.....	*	*	*	*	*	*	*	*	0.9	3.5
2001.....	*	*	*	*	*	*	*	*	0.7	3.7
2000.....	*	*	*	*	*	*	*	0.5	0.9	4.8
1999.....	*	*	*	*	*	*	*	0.5	1.1	6.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute lower respiratory infections (J20-J22)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*	0.3	0.7	2.2
2005 4/.....	*	*	*	*	*	*	*	0.4	0.9	3.5
2004 4/.....	*	*	*	*	*	*	*	*	0.7	2.8
2003 4/.....	*	*	*	*	*	*	*	0.5	0.9	3.7
2002.....	*	*	*	*	*	*	*	*	0.8	3.5
2001.....	*	*	*	*	*	*	*	0.3	0.7	3.5
2000.....	*	*	*	*	*	*	0.3	0.6	0.9	4.7
1999.....	*	*	*	*	*	*	*	0.6	1.3	6.3

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	1.9
2005 4/.....	*	*	*	*	*	*	*	*	1.4	3.3
2004 4/.....	*	*	*	*	*	*	*	*	*	2.6
2003 4/.....	*	*	*	*	*	*	*	*	*	2.7
2002.....	*	*	*	*	*	*	*	*	*	2.9
2001.....	*	*	*	*	*	*	*	*	*	2.4
2000.....	*	*	*	*	*	*	*	0.8	*	3.8
1999.....	*	*	*	*	*	*	*	0.8	1.6	4.0

White, female

2006 4/.....	*	*	*	*	*	*	*	*	0.8	2.3
2005 4/.....	*	*	*	*	*	*	*	*	*	3.6
2004 4/.....	*	*	*	*	*	*	*	*	0.7	2.8
2003 4/.....	*	*	*	*	*	*	*	*	0.8	4.2
2002.....	*	*	*	*	*	*	*	*	0.9	3.7
2001.....	*	*	*	*	*	*	*	*	0.8	3.9
2000.....	*	*	*	*	*	*	*	*	1.0	5.1
1999.....	*	*	*	*	*	*	*	0.5	1.1	7.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Acute bronchitis and bronchiolitis (J20-J21)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	*	*	0.5	1.2
2005 4/.....	*	*	*	*	*	*	*	0.3	0.6	2.0
2004 4/.....	*	*	*	*	*	*	*	*	*	1.5
2003 4/.....	*	*	*	*	*	*	*	*	0.5	2.3
2002.....	*	*	*	*	*	*	*	*	0.5	2.1
2001.....	*	*	*	*	*	*	*	*	0.5	2.1
2000.....	*	*	*	*	*	*	*	0.4	0.5	2.7
1999.....	*	*	*	*	*	*	*	0.4	0.9	3.8
White, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	1.7
2004 4/.....	*	*	*	*	*	*	*	*	*	1.8
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	1.7
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	2.4
1999.....	*	*	*	*	*	*	*	*	*	2.6
White, female										
2006 4/.....	*	*	*	*	*	*	*	*	*	1.1
2005 4/.....	*	*	*	*	*	*	*	*	*	2.1
2004 4/.....	*	*	*	*	*	*	*	*	*	1.3
2003 4/.....	*	*	*	*	*	*	*	*	*	2.7
2002.....	*	*	*	*	*	*	*	*	*	2.3
2001.....	*	*	*	*	*	*	*	*	*	2.5
2000.....	*	*	*	*	*	*	*	*	*	2.9
1999.....	*	*	*	*	*	*	*	*	0.7	4.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of upper respiratory tract (J30-J39)

All races 3/, both sexes

2006.....	*	*	0.2	0.2	0.3	0.3	0.4	0.8	0.9	2.0
2005.....	*	0.1	0.2	0.2	0.2	0.3	0.4	0.6	1.2	2.2
2004.....	*	0.1	0.1	0.1	0.2	0.2	0.3	0.7	1.1	2.1
2003.....	*	*	*	0.2	0.3	0.3	0.4	0.7	1.0	2.3
2002.....	*	0.1	*	0.2	0.2	0.3	0.4	0.6	0.7	2.1
2001.....	*	0.1	0.1	*	0.2	0.3	0.4	0.6	1.3	2.4
2000.....	*	0.1	0.1	0.1	0.2	0.3	0.5	0.7	1.2	2.1
1999.....	*	*	*	*	0.3	0.4	0.4	0.9	1.1	2.4

All races 3/, male

2006.....	*	*	*	*	*	*	0.5	1.1	1.0	2.7
2005.....	*	*	*	*	*	*	0.6	0.7	1.2	3.1
2004.....	*	*	*	*	*	*	*	0.8	1.5	3.1
2003.....	*	*	*	*	*	0.4	*	0.9	1.5	3.0
2002.....	*	*	*	*	*	*	*	0.7	*	2.5
2001.....	*	*	*	*	*	*	*	0.7	1.9	3.4
2000.....	*	*	*	*	*	*	0.6	0.7	1.9	3.1
1999.....	*	*	*	*	*	0.5	0.5	1.1	1.3	3.5

All races 3/, female

2006.....	*	*	*	*	*	*	*	0.6	0.8	1.7
2005.....	*	*	0.2	*	*	*	*	0.5	1.1	1.8
2004.....	*	*	*	*	*	*	*	0.6	0.8	1.6
2003.....	*	*	*	*	*	*	*	0.5	0.7	2.0
2002.....	*	*	*	*	*	*	*	0.5	0.6	1.9
2001.....	*	*	*	*	*	*	*	0.6	1.0	1.9
2000.....	*	*	*	*	*	*	0.5	0.6	0.8	1.7
1999.....	*	*	*	*	*	*	*	0.7	1.0	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of upper respiratory tract (J30-J39)

White, both sexes

2006 4/.....	*	*	0.1	0.1	0.2	0.2	0.4	0.8	0.9	1.9
2005 4/.....	*	*	0.2	0.2	*	*	0.4	0.6	1.2	2.3
2004 4/.....	*	0.1	*	*	*	*	*	0.6	1.1	2.1
2003 4/.....	*	*	*	*	0.3	0.3	0.4	0.6	1.1	2.3
2002.....	*	*	*	0.2	*	0.2	0.4	0.6	0.6	2.2
2001.....	*	*	*	*	0.2	0.3	0.3	0.6	1.4	2.3
2000.....	*	*	0.1	*	0.2	0.3	0.5	0.6	1.3	2.2
1999.....	*	*	*	*	0.3	0.3	0.4	0.8	1.1	2.4

White, male

2006 4/.....	*	*	*	*	*	*	*	1.1	1.1	2.7
2005 4/.....	*	*	*	*	*	*	*	0.8	1.2	3.3
2004 4/.....	*	*	*	*	*	*	*	*	1.5	3.2
2003 4/.....	*	*	*	*	*	*	*	0.9	1.7	3.2
2002.....	*	*	*	*	*	*	*	0.7	*	2.7
2001.....	*	*	*	*	*	*	*	*	1.9	3.2
2000.....	*	*	*	*	*	*	0.6	*	1.9	3.0
1999.....	*	*	*	*	*	*	*	1.1	1.5	3.2

White, female

2006 4/.....	*	*	*	*	*	*	*	0.6	0.8	1.6
2005 4/.....	*	*	*	*	*	*	*	0.5	1.2	1.9
2004 4/.....	*	*	*	*	*	*	*	0.6	0.8	1.6
2003 4/.....	*	*	*	*	*	*	*	*	0.7	1.9
2002.....	*	*	*	*	*	*	*	0.5	*	1.9
2001.....	*	*	*	*	*	*	*	0.6	1.0	1.9
2000.....	*	*	*	*	*	*	*	0.6	0.9	1.8
1999.....	*	*	*	*	*	*	*	0.7	0.9	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)										
All races 3/, both sexes										
2006.....	2.8	5.8	12.7	25.8	57.4	111.8	194.8	308.5	434.8	589.1
2005.....	2.8	6.5	12.6	26.9	62.1	118.6	210.3	330.6	457.8	637.2
2004.....	2.6	5.6	11.7	25.5	60.0	113.7	200.6	315.7	434.7	601.7
2003.....	2.8	5.9	11.9	27.5	63.9	121.1	210.8	330.0	455.9	635.1
2002.....	2.9	5.8	12.0	28.2	60.8	119.0	211.5	330.1	465.7	637.6
2001.....	2.7	5.4	11.9	29.4	62.9	124.2	215.4	329.2	452.9	644.7
2000.....	2.8	5.8	11.8	29.6	62.5	124.1	218.2	327.8	473.4	648.6
1999.....	2.6	5.5	12.0	31.9	66.6	130.4	227.6	340.2	484.5	646.0
All races 3/, male										
2006.....	2.6	5.5	13.4	27.4	62.0	119.8	218.0	364.6	541.3	760.8
2005.....	2.7	6.4	13.6	28.5	67.6	132.9	240.0	393.6	564.7	819.9
2004.....	2.6	5.6	12.5	26.9	64.6	126.1	228.2	378.8	543.4	811.1
2003.....	2.5	6.3	12.3	29.4	69.2	132.6	244.0	395.8	569.1	865.9
2002.....	3.0	5.9	12.8	29.5	65.9	131.5	246.0	405.6	595.9	894.8
2001.....	2.5	5.8	12.3	31.4	67.0	137.1	252.9	406.7	583.5	916.9
2000.....	2.6	6.2	12.1	31.8	68.0	139.7	257.8	404.2	628.5	967.9
1999.....	2.4	5.7	12.1	34.3	74.6	151.0	278.3	432.4	648.3	996.7
All races 3/, female										
2006.....	2.9	6.1	12.1	24.2	53.3	104.8	175.8	267.5	368.3	508.7
2005.....	2.8	6.6	11.7	25.5	57.1	106.1	186.2	284.8	391.5	553.3
2004.....	2.7	5.5	10.9	24.2	55.8	102.9	178.4	270.4	367.7	507.4
2003.....	3.1	5.4	11.5	25.7	59.0	111.2	184.2	283.1	386.7	533.0
2002.....	2.8	5.6	11.2	26.9	56.1	108.3	183.8	276.8	386.5	526.0
2001.....	2.9	5.1	11.5	27.5	59.1	113.1	185.6	275.0	374.6	530.8
2000.....	2.9	5.5	11.5	27.5	57.5	110.7	187.0	274.6	381.9	518.5
1999.....	2.9	5.3	11.9	29.7	59.3	112.8	188.0	276.1	388.7	504.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)										
White, both sexes										
2006 4/.....	2.5	5.8	12.7	26.7	60.4	119.4	209.5	326.4	457.6	618.3
2005 4/.....	2.5	6.2	12.4	27.7	65.6	126.2	224.5	349.4	481.5	667.6
2004 4/.....	2.4	5.2	11.6	26.1	63.2	121.7	214.3	333.3	457.6	630.1
2003 4/.....	2.5	5.6	11.7	28.0	67.3	129.5	223.2	347.2	478.9	664.5
2002.....	2.5	5.5	11.7	28.9	63.8	126.8	223.3	346.9	486.5	666.2
2001.....	2.4	5.1	11.7	30.2	66.2	131.3	227.1	346.1	473.5	671.8
2000.....	2.3	5.3	11.4	30.1	65.5	130.5	229.6	343.1	495.5	675.6
1999.....	2.2	4.9	11.8	32.6	69.6	137.2	237.9	353.4	504.9	672.9
White, male										
2006 4/.....	2.3	5.3	13.1	27.9	63.3	125.0	229.8	378.5	559.5	788.6
2005 4/.....	2.4	6.0	13.2	28.5	70.2	138.6	251.2	406.6	583.5	846.1
2004 4/.....	2.4	5.3	12.3	26.6	66.5	132.4	238.5	392.8	563.4	837.7
2003 4/.....	2.2	6.1	12.0	29.4	71.5	139.3	252.4	406.9	585.6	894.4
2002.....	2.5	5.6	12.4	29.6	67.5	137.1	254.3	417.4	612.9	923.4
2001.....	2.2	5.6	11.9	31.8	69.1	142.0	260.8	418.9	600.0	943.6
2000.....	2.2	5.7	11.5	31.8	69.8	143.1	265.8	415.5	648.5	997.4
1999.....	2.0	5.0	11.6	34.5	76.3	155.7	285.6	441.3	664.4	1,029.4
White, female										
2006 4/.....	2.6	6.2	12.3	25.5	57.6	114.5	192.6	287.6	393.2	538.4
2005 4/.....	2.6	6.3	11.7	26.9	61.4	115.1	202.4	307.0	417.6	585.4
2004 4/.....	2.5	5.1	11.0	25.6	60.1	112.2	194.4	289.7	391.9	536.5
2003 4/.....	2.8	5.1	11.4	26.7	63.5	120.8	199.3	304.0	413.2	562.8
2002.....	2.5	5.4	11.0	28.3	60.4	117.6	198.0	296.4	408.9	554.5
2001.....	2.6	4.6	11.6	28.6	63.5	122.0	199.7	294.5	397.1	557.9
2000.....	2.4	4.9	11.3	28.4	61.5	119.4	200.5	292.1	404.4	544.8
1999.....	2.5	4.7	12.0	30.8	63.5	121.1	200.0	291.6	410.7	529.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic lower respiratory diseases (J40-J47)										
Black, both sexes										
2006 4/.....	5.2	7.7	17.1	27.7	51.9	83.5	122.6	199.9	262.7	331.8
2005 4/.....	5.0	10.5	17.9	30.4	52.4	92.0	142.8	220.8	271.1	366.7
2004 4/.....	4.8	9.6	16.3	27.9	52.1	81.7	128.1	204.6	249.3	333.9
2003 4/.....	5.7	9.1	16.7	31.3	52.7	86.5	146.7	211.9	271.9	353.0
2002.....	6.5	9.1	17.4	30.9	53.4	86.1	148.1	215.9	298.5	368.2
2001.....	5.1	9.1	16.6	31.6	52.2	94.3	149.6	206.9	276.5	375.8
2000.....	6.1	11.0	18.2	33.0	57.4	97.3	151.2	211.6	271.3	373.8
1999.....	6.1	11.2	17.2	36.1	57.6	100.0	163.3	242.4	295.3	376.2
Black, male										
2006 4/.....	5.0	7.8	19.9	33.6	70.2	112.0	169.2	287.7	417.6	512.0
2005 4/.....	4.9	11.0	20.1	37.5	66.0	126.0	198.9	345.5	422.8	594.9
2004 4/.....	4.9	9.9	18.9	37.7	67.6	109.0	182.3	300.7	392.4	566.3
2003 4/.....	5.1	9.8	17.7	38.0	67.7	112.0	212.5	326.8	471.8	601.6
2002.....	6.9	9.9	19.9	38.1	71.1	116.2	214.0	340.8	477.6	645.4
2001.....	5.1	8.9	18.8	37.1	66.3	127.1	222.2	329.9	468.7	671.7
2000.....	5.9	11.4	19.6	39.5	73.5	138.4	226.3	326.5	448.5	693.1
1999.....	6.2	11.9	19.7	43.2	79.3	139.4	243.5	387.0	521.4	701.9
Black, female										
2006 4/.....	5.4	7.5	14.7	22.8	37.7	62.5	90.7	147.3	180.9	258.1
2005 4/.....	5.1	10.1	16.1	24.6	41.8	66.8	104.3	146.4	191.0	274.6
2004 4/.....	4.8	9.3	14.2	19.9	40.1	61.3	91.4	147.7	173.9	241.8
2003 4/.....	6.2	8.6	15.8	25.7	41.1	67.6	102.4	143.5	167.1	256.4
2002.....	6.1	8.5	15.3	25.1	39.6	63.8	104.0	141.3	205.3	262.0
2001.....	5.1	9.2	14.7	27.1	41.1	70.0	101.3	133.4	178.7	266.1
2000.....	6.2	10.6	16.9	27.7	44.6	67.0	101.2	142.4	182.1	255.0
1999.....	6.1	10.7	15.0	30.2	40.6	70.8	110.0	155.3	183.2	254.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
All races 3/, both sexes										
2006.....	2.7	5.7	12.6	25.7	57.2	111.3	193.6	306.6	430.7	582.6
2005.....	2.7	6.4	12.6	26.8	61.9	118.2	209.2	328.1	453.8	630.8
2004.....	2.6	5.5	11.6	25.4	59.7	113.3	199.5	313.2	430.7	595.3
2003.....	2.8	5.8	11.8	27.3	63.6	120.6	209.8	327.7	451.8	629.1
2002.....	2.9	5.7	11.9	28.0	60.4	118.4	210.3	327.6	461.0	631.8
2001.....	2.7	5.4	11.8	29.3	62.5	123.4	214.2	326.9	449.0	637.7
2000.....	2.7	5.8	11.8	29.3	62.2	123.4	216.9	325.4	468.8	642.4
1999.....	2.6	5.4	11.9	31.7	66.2	129.6	226.5	337.6	479.9	639.8
All races 3/, male										
2006.....	2.6	5.4	13.2	27.3	61.8	119.5	217.2	363.0	538.3	756.9
2005.....	2.6	6.3	13.5	28.3	67.4	132.5	239.3	392.2	562.2	815.4
2004.....	2.6	5.6	12.4	26.8	64.3	125.8	227.5	376.9	541.1	806.6
2003.....	2.5	6.3	12.2	29.3	68.9	132.4	243.3	394.2	565.9	861.8
2002.....	2.9	5.9	12.7	29.4	65.6	131.1	245.1	404.0	592.8	890.1
2001.....	2.5	5.7	12.3	31.3	66.6	136.5	252.0	405.0	581.3	912.1
2000.....	2.6	6.2	12.1	31.6	67.7	139.3	256.8	402.6	625.4	963.6
1999.....	2.4	5.6	12.1	34.2	74.3	150.6	277.3	429.9	644.2	990.1
All races 3/, female										
2006.....	2.9	6.1	12.0	24.1	53.0	104.1	174.4	265.4	363.5	501.1
2005.....	2.8	6.5	11.6	25.4	56.9	105.6	184.8	281.6	386.5	546.0
2004.....	2.7	5.5	10.8	24.0	55.5	102.3	176.9	267.5	362.8	500.3
2003.....	3.1	5.3	11.4	25.4	58.7	110.4	182.9	280.4	382.0	526.2
2002.....	2.8	5.6	11.1	26.7	55.6	107.4	182.4	273.7	380.9	519.7
2001.....	2.8	5.1	11.4	27.4	58.7	112.2	184.2	272.2	369.7	522.9
2000.....	2.8	5.4	11.4	27.2	57.2	109.9	185.6	271.7	376.5	511.6
1999.....	2.8	5.2	11.7	29.4	58.9	111.6	186.8	273.5	383.7	498.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
White, both sexes										
2006 4/.....	2.4	5.7	12.6	26.6	60.1	118.8	208.3	324.5	453.3	611.5
2005 4/.....	2.5	6.1	12.4	27.5	65.4	125.7	223.3	346.8	477.3	660.6
2004 4/.....	2.4	5.2	11.5	26.0	62.9	121.2	213.1	330.6	453.3	623.4
2003 4/.....	2.5	5.6	11.7	27.9	67.0	129.0	222.1	344.7	474.6	658.1
2002.....	2.5	5.5	11.6	28.8	63.3	126.2	222.0	344.2	481.6	660.2
2001.....	2.4	5.1	11.6	30.1	65.8	130.6	225.8	343.6	469.4	664.3
2000.....	2.3	5.2	11.4	29.8	65.2	129.8	228.4	340.6	490.6	669.0
1999.....	2.2	4.8	11.7	32.3	69.2	136.4	236.8	350.8	500.1	666.2
White, male										
2006 4/.....	2.3	5.3	13.0	27.8	63.1	124.6	229.0	377.0	556.4	784.8
2005 4/.....	2.4	6.0	13.2	28.3	69.9	138.2	250.4	405.1	580.9	841.5
2004 4/.....	2.4	5.2	12.1	26.5	66.2	132.0	237.7	390.9	560.9	832.8
2003 4/.....	2.2	6.1	12.0	29.4	71.2	139.1	251.7	405.2	582.3	889.8
2002.....	2.5	5.5	12.3	29.5	67.1	136.8	253.5	415.7	609.7	918.7
2001.....	2.2	5.5	11.8	31.7	68.6	141.6	260.0	417.3	597.9	938.8
2000.....	2.2	5.6	11.5	31.6	69.6	142.8	264.8	414.0	645.1	992.8
1999.....	2.0	5.0	11.6	34.3	76.0	155.3	284.6	438.9	660.3	1,022.5
White, female										
2006 4/.....	2.6	6.2	12.2	25.4	57.3	113.7	191.2	285.3	388.1	530.2
2005 4/.....	2.6	6.3	11.6	26.7	61.2	114.7	200.9	303.7	412.3	577.4
2004 4/.....	2.5	5.1	10.9	25.6	59.9	111.6	192.9	286.5	386.5	529.0
2003 4/.....	2.7	5.1	11.4	26.5	63.2	120.0	197.9	301.0	408.2	555.5
2002.....	2.4	5.4	11.0	28.1	59.7	116.8	196.4	293.1	403.0	547.9
2001.....	2.6	4.6	11.5	28.5	63.2	121.0	198.2	291.4	391.8	549.4
2000.....	2.4	4.9	11.2	28.1	61.0	118.4	199.2	289.0	398.7	537.5
1999.....	2.5	4.7	11.8	30.4	63.0	119.9	198.8	288.9	405.5	523.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchitis, chronic and unspecified, emphysema and asthma (J40-J46)										
Black, both sexes										
2006 4/.....	5.1	7.7	17.0	27.5	51.7	83.4	121.9	199.0	261.5	330.0
2005 4/.....	5.0	10.3	17.8	30.3	52.2	91.6	142.4	220.1	270.6	365.4
2004 4/.....	4.8	9.5	16.3	27.8	52.0	81.5	127.5	204.0	248.5	332.5
2003 4/.....	5.6	9.1	16.5	30.9	52.6	86.4	146.3	211.3	270.8	352.7
2002.....	6.5	9.0	17.2	30.9	53.3	85.6	147.6	215.4	297.0	367.0
2001.....	5.1	9.0	16.6	31.5	51.9	94.0	149.2	206.0	275.2	374.6
2000.....	6.1	10.9	18.1	32.6	57.1	96.6	150.4	211.1	270.5	373.2
1999.....	6.1	11.1	17.1	35.8	57.4	99.7	162.5	241.0	293.6	375.3
Black, male										
2006 4/.....	4.9	7.8	19.8	33.3	69.6	111.8	168.2	285.9	416.2	509.4
2005 4/.....	4.9	10.7	19.9	37.3	66.0	126.0	198.3	345.1	422.1	592.1
2004 4/.....	4.9	9.8	18.9	37.5	67.5	109.0	181.9	300.7	392.4	564.3
2003 4/.....	5.1	9.8	17.5	37.7	67.7	112.0	211.5	326.4	470.3	601.6
2002.....	6.9	9.8	19.7	38.0	70.9	115.7	212.6	339.9	476.1	645.4
2001.....	5.1	8.8	18.8	36.9	66.1	126.5	221.5	328.5	465.5	669.4
2000.....	5.9	11.3	19.6	39.5	73.1	137.1	225.0	325.1	448.5	693.1
1999.....	6.1	11.8	19.6	42.9	78.9	139.4	242.9	385.1	520.6	700.7
Black, female										
2006 4/.....	5.3	7.5	14.6	22.8	37.7	62.5	90.0	147.0	179.8	256.6
2005 4/.....	5.1	9.9	16.0	24.6	41.5	66.1	104.1	145.6	190.7	273.8
2004 4/.....	4.8	9.2	14.2	19.8	39.9	61.0	90.5	146.6	172.7	240.6
2003 4/.....	6.1	8.4	15.7	25.3	40.9	67.4	102.4	143.0	166.3	256.0
2002.....	6.1	8.3	15.1	25.1	39.5	63.3	104.0	141.1	203.7	260.4
2001.....	5.1	9.2	14.7	27.1	40.8	70.0	101.1	132.9	178.2	265.2
2000.....	6.2	10.6	16.8	27.0	44.5	66.6	100.8	142.4	180.9	254.1
1999.....	6.1	10.6	14.9	30.0	40.4	70.2	109.1	154.2	181.1	253.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Bronchitis, not specified as acute or chronic (J40)

All races 3/, both sexes

2006.....	*	*	*	*	*	*	0.3	0.6	1.2	3.3
2005.....	*	*	*	*	0.2	0.3	0.4	0.5	1.3	4.2
2004.....	*	*	*	*	*	0.2	0.3	0.5	1.0	3.8
2003.....	*	*	*	*	*	*	0.3	0.7	1.1	4.6
2002.....	*	*	*	*	*	*	0.4	0.7	1.4	5.2
2001.....	*	*	*	*	*	0.2	0.4	1.0	1.3	5.2
2000.....	*	*	*	*	0.2	0.3	0.4	1.1	2.0	6.8
1999.....	*	*	*	*	0.3	0.3	0.7	1.0	2.2	6.2

All races 3/, male

2006.....	*	*	*	*	*	*	*	*	1.2	3.4
2005.....	*	*	*	*	*	*	*	0.7	1.5	3.9
2004.....	*	*	*	*	*	*	*	*	*	3.9
2003.....	*	*	*	*	*	*	*	0.8	1.1	3.9
2002.....	*	*	*	*	*	*	*	0.8	1.2	4.7
2001.....	*	*	*	*	*	*	*	1.0	1.2	5.1
2000.....	*	*	*	*	*	*	*	1.3	3.0	6.4
1999.....	*	*	*	*	*	*	0.7	1.2	2.3	5.9

All races 3/, female

2006.....	*	*	*	*	*	*	*	0.6	1.1	3.3
2005.....	*	*	*	*	*	*	*	*	1.1	4.4
2004.....	*	*	*	*	*	*	*	*	1.0	3.7
2003.....	*	*	*	*	*	*	*	0.6	1.0	4.9
2002.....	*	*	*	*	*	*	0.5	0.7	1.5	5.4
2001.....	*	*	*	*	*	*	*	0.9	1.4	5.3
2000.....	*	*	*	*	*	*	0.5	1.0	1.4	7.0
1999.....	*	*	*	*	*	*	0.6	0.8	2.1	6.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Bronchitis, not specified as acute or chronic (J40)

White, both sexes

2006 4/.....	*	*	*	*	*	*	0.3	0.6	1.2	3.5
2005 4/.....	*	*	*	*	*	0.3	0.4	0.5	1.3	4.5
2004 4/.....	*	*	*	*	*	*	0.3	0.5	1.0	3.9
2003 4/.....	*	*	*	*	*	*	0.3	0.8	1.1	4.7
2002.....	*	*	*	*	*	*	0.5	0.8	1.5	5.5
2001.....	*	*	*	*	*	*	0.4	1.0	1.4	5.5
2000.....	*	*	*	*	0.2	0.3	0.4	1.2	2.1	7.3
1999.....	*	*	*	*	0.3	0.3	0.7	0.9	2.2	6.4

White, male

2006 4/.....	*	*	*	*	*	*	*	*	1.3	3.7
2005 4/.....	*	*	*	*	*	*	*	*	1.6	3.9
2004 4/.....	*	*	*	*	*	*	*	*	*	4.1
2003 4/.....	*	*	*	*	*	*	*	0.9	1.2	4.0
2002.....	*	*	*	*	*	*	*	0.8	1.3	5.1
2001.....	*	*	*	*	*	*	*	1.0	1.2	5.3
2000.....	*	*	*	*	*	*	*	1.4	3.3	6.9
1999.....	*	*	*	*	*	*	0.7	1.1	2.2	6.3

White, female

2006 4/.....	*	*	*	*	*	*	*	0.6	1.1	3.4
2005 4/.....	*	*	*	*	*	*	*	*	1.2	4.7
2004 4/.....	*	*	*	*	*	*	*	*	1.1	3.8
2003 4/.....	*	*	*	*	*	*	*	0.7	1.0	5.0
2002.....	*	*	*	*	*	*	0.5	0.8	1.5	5.7
2001.....	*	*	*	*	*	*	*	1.0	1.5	5.7
2000.....	*	*	*	*	*	*	*	1.0	1.4	7.5
1999.....	*	*	*	*	*	*	0.6	0.7	2.2	6.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic bronchitis (J41-J42)										
All races 3/, both sexes										
2006.....	*	*	*	*	*	*	0.4	0.6	0.9	2.7
2005.....	*	*	*	0.1	*	*	0.5	0.6	0.9	3.2
2004.....	*	*	*	0.1	*	0.2	0.4	0.7	1.1	2.8
2003.....	*	*	*	*	*	0.3	0.4	0.7	1.5	2.7
2002.....	*	*	*	*	0.2	0.4	0.5	1.0	1.5	2.9
2001.....	*	*	*	*	0.2	0.3	0.4	0.8	1.2	3.9
2000.....	*	*	*	*	0.2	0.4	0.6	1.0	1.6	5.0
1999.....	*	*	*	0.2	0.3	0.3	0.6	0.9	1.9	4.5
All races 3/, male										
2006.....	*	*	*	*	*	*	*	*	1.0	2.5
2005.....	*	*	*	*	*	*	*	0.7	1.0	3.4
2004.....	*	*	*	*	*	*	*	*	1.3	3.1
2003.....	*	*	*	*	*	*	*	1.1	1.5	3.0
2002.....	*	*	*	*	*	0.5	0.7	1.5	1.6	4.0
2001.....	*	*	*	*	*	0.5	0.6	1.1	1.2	4.1
2000.....	*	*	*	*	*	0.6	0.6	1.4	1.9	6.4
1999.....	*	*	*	*	*	0.5	0.7	1.3	2.3	5.9
All races 3/, female										
2006.....	*	*	*	*	*	*	*	0.6	0.8	2.7
2005.....	*	*	*	*	*	*	0.5	0.5	0.9	3.0
2004.....	*	*	*	*	*	*	0.4	0.7	0.9	2.6
2003.....	*	*	*	*	*	*	0.5	0.5	1.5	2.6
2002.....	*	*	*	*	*	*	*	0.7	1.4	2.4
2001.....	*	*	*	*	*	*	*	0.5	1.2	3.8
2000.....	*	*	*	*	*	*	0.6	0.7	1.4	4.4
1999.....	*	*	*	*	0.4	*	0.4	0.7	1.7	4.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Chronic bronchitis (J41-J42)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	0.4	0.6	0.9	2.8
2005 4/.....	*	*	*	*	*	*	0.5	0.6	0.9	3.4
2004 4/.....	*	*	*	*	*	*	0.4	0.7	1.1	2.8
2003 4/.....	*	*	*	*	*	0.3	0.4	0.7	1.5	2.8
2002.....	*	*	*	*	*	0.4	0.6	1.1	1.5	3.1
2001.....	*	*	*	*	0.2	0.4	0.4	0.8	1.3	4.1
2000.....	*	*	*	*	0.2	0.4	0.6	1.0	1.6	5.3
1999.....	*	*	*	0.2	0.3	0.4	0.6	1.0	1.9	4.8
White, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	2.6
2005 4/.....	*	*	*	*	*	*	*	0.8	*	3.5
2004 4/.....	*	*	*	*	*	*	*	*	1.4	3.0
2003 4/.....	*	*	*	*	*	*	*	1.1	1.3	3.3
2002.....	*	*	*	*	*	*	0.7	1.5	1.6	4.2
2001.....	*	*	*	*	*	0.6	0.6	1.2	1.3	4.2
2000.....	*	*	*	*	*	0.6	0.6	1.5	1.9	6.9
1999.....	*	*	*	*	*	0.5	0.7	1.3	2.2	6.1
White, female										
2006 4/.....	*	*	*	*	*	*	*	0.6	0.8	2.8
2005 4/.....	*	*	*	*	*	*	0.5	*	0.9	3.3
2004 4/.....	*	*	*	*	*	*	*	0.7	0.9	2.6
2003 4/.....	*	*	*	*	*	*	0.5	*	1.6	2.6
2002.....	*	*	*	*	*	*	*	0.8	1.4	2.6
2001.....	*	*	*	*	*	*	*	0.5	1.3	4.0
2000.....	*	*	*	*	*	*	0.7	0.7	1.5	4.7
1999.....	*	*	*	*	*	*	*	0.7	1.7	4.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
All races 3/, both sexes										
2006.....	0.3	0.7	1.4	3.2	7.1	13.2	22.3	33.1	41.6	45.3
2005.....	0.2	0.8	1.4	3.7	8.0	15.3	25.8	37.8	46.0	51.9
2004.....	0.2	0.6	1.5	3.4	8.1	15.8	26.1	37.5	45.7	49.9
2003.....	0.3	0.7	1.5	4.0	9.3	16.6	28.5	41.9	51.6	54.6
2002.....	0.3	0.7	1.6	4.3	9.1	17.4	31.3	43.2	54.7	59.3
2001.....	0.2	0.7	1.7	5.0	10.1	19.8	33.2	46.9	55.5	62.1
2000.....	0.3	0.8	1.8	5.0	10.4	20.7	35.5	47.5	60.4	63.3
1999.....	0.4	0.7	2.0	5.5	12.1	23.5	38.4	51.3	65.3	62.1
All races 3/, male										
2006.....	0.4	0.7	1.9	3.5	7.9	15.1	25.2	40.3	52.8	63.7
2005.....	0.3	1.1	1.6	4.4	9.7	18.4	30.7	47.5	59.5	71.5
2004.....	0.3	0.8	1.6	3.8	9.5	18.4	31.6	46.5	60.5	74.4
2003.....	0.3	1.0	1.9	4.4	10.3	18.3	34.4	51.9	67.5	79.2
2002.....	0.5	0.9	1.9	4.6	10.7	19.7	37.8	56.1	74.9	90.0
2001.....	0.3	0.9	2.0	5.8	10.8	22.4	40.3	59.2	74.9	101.2
2000.....	0.3	1.0	2.1	5.6	11.2	24.5	42.6	60.4	83.6	104.2
1999.....	0.4	0.9	2.2	6.2	13.6	28.0	48.9	67.3	91.5	105.4
All races 3/, female										
2006.....	0.3	0.7	1.0	2.8	6.4	11.5	19.9	27.8	34.6	36.7
2005.....	*	0.5	1.1	3.0	6.5	12.6	21.9	30.8	37.6	42.8
2004.....	0.2	0.4	1.3	3.1	6.8	13.5	21.7	31.1	36.6	38.9
2003.....	0.3	0.5	1.2	3.6	8.4	15.2	23.7	34.8	41.9	43.7
2002.....	0.2	0.6	1.3	4.0	7.7	15.4	26.1	34.0	42.4	46.0
2001.....	*	0.5	1.4	4.2	9.5	17.6	27.5	38.3	43.8	45.8
2000.....	0.2	0.6	1.5	4.5	9.7	17.5	29.9	38.5	46.8	46.7
1999.....	0.3	0.5	1.7	4.9	10.8	19.8	30.2	40.2	49.9	44.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
White, both sexes										
2006 4/.....	0.3	0.7	1.5	3.4	7.5	14.2	24.3	35.2	43.9	47.8
2005 4/.....	0.2	0.8	1.3	3.9	8.6	16.4	27.9	39.9	48.2	54.5
2004 4/.....	0.2	0.6	1.4	3.6	8.6	17.2	28.2	40.1	48.4	52.5
2003 4/.....	0.3	0.7	1.6	4.2	9.9	18.0	30.5	44.3	54.2	57.0
2002.....	0.3	0.7	1.7	4.6	9.8	18.6	33.3	45.8	57.4	62.3
2001.....	0.2	0.7	1.7	5.1	10.9	21.5	35.2	49.8	58.7	65.1
2000.....	0.2	0.7	1.8	5.3	11.2	22.0	37.9	50.2	63.6	65.8
1999.....	0.3	0.6	2.0	5.8	13.0	25.4	40.9	53.8	68.8	64.7
White, male										
2006 4/.....	0.4	0.7	1.9	3.6	7.9	15.8	27.1	41.7	54.6	66.5
2005 4/.....	0.3	1.1	1.5	4.5	10.1	19.2	32.4	49.3	60.6	73.6
2004 4/.....	0.3	0.8	1.6	3.8	9.8	19.4	33.4	48.5	63.2	77.1
2003 4/.....	0.3	1.0	1.9	4.5	10.6	19.5	35.9	53.2	69.6	80.4
2002.....	0.4	0.9	1.9	4.8	11.1	20.2	39.5	58.2	77.4	93.0
2001.....	0.3	1.0	1.9	5.8	11.3	23.9	41.6	61.7	78.0	105.1
2000.....	0.2	0.9	2.0	5.8	11.8	25.0	44.6	63.1	87.0	105.5
1999.....	0.3	0.8	2.1	6.4	14.2	29.4	50.6	69.6	94.4	109.2
White, female										
2006 4/.....	0.3	0.7	1.2	3.2	7.1	12.7	21.9	30.3	37.1	39.1
2005 4/.....	*	0.5	1.2	3.3	7.2	13.8	24.2	33.0	40.5	45.7
2004 4/.....	*	0.4	1.3	3.4	7.5	15.2	24.0	33.9	39.2	41.4
2003 4/.....	0.3	0.5	1.4	3.9	9.2	16.7	26.0	37.8	44.7	46.6
2002.....	*	0.6	1.5	4.4	8.6	17.3	28.3	37.0	45.1	48.9
2001.....	*	0.5	1.4	4.5	10.6	19.4	30.0	41.3	47.0	48.4
2000.....	0.2	0.6	1.5	4.8	10.7	19.3	32.5	41.1	49.8	49.7
1999.....	0.3	0.5	1.9	5.3	11.9	21.9	33.1	42.6	53.7	46.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Emphysema (J43)										
Black, both sexes										
2006 4/.....	*	0.7	1.3	2.6	6.7	8.9	11.5	20.4	24.4	23.9
2005 4/.....	*	0.9	1.9	3.5	6.0	11.6	13.6	26.2	26.2	27.3
2004 4/.....	*	*	2.1	3.1	6.5	8.4	13.7	20.7	24.3	27.6
2003 4/.....	*	1.0	1.3	3.6	7.2	10.2	16.4	25.2	30.5	30.9
2002.....	0.8	1.0	1.5	3.3	5.9	11.9	20.9	23.5	34.8	32.1
2001.....	*	*	2.3	4.9	6.4	11.6	20.1	21.9	26.3	35.3
2000.....	*	1.6	2.5	4.7	7.7	14.9	19.5	24.5	30.8	37.2
1999.....	0.8	1.4	2.4	4.6	8.0	14.0	20.8	31.4	31.2	37.6
Black, male										
2006 4/.....	*	*	2.5	3.9	11.0	13.4	14.5	32.2	41.1	40.2
2005 4/.....	*	*	2.8	5.6	9.8	17.8	20.2	39.8	45.6	51.1
2004 4/.....	*	*	3.0	4.9	9.7	13.8	21.4	34.6	39.5	56.6
2003 4/.....	*	*	2.0	5.2	10.2	13.3	24.6	43.4	50.2	70.9
2002.....	*	*	2.4	4.5	9.3	20.4	30.7	41.3	61.1	64.3
2001.....	*	*	2.9	6.9	9.6	15.8	31.6	34.4	49.0	68.1
2000.....	*	2.4	3.5	5.9	10.8	25.2	30.1	35.0	53.1	87.6
1999.....	*	2.0	4.1	6.4	11.9	22.8	37.0	48.9	65.4	72.6
Black, female										
2006 4/.....	*	*	*	*	3.3	5.5	9.5	13.4	15.6	17.2
2005 4/.....	*	*	*	*	3.0	7.1	9.1	18.1	16.0	17.6
2004 4/.....	*	*	*	*	3.9	4.3	8.6	12.4	16.3	16.1
2003 4/.....	*	*	*	2.3	4.8	7.9	10.9	14.4	20.1	15.4
2002.....	*	*	*	*	3.3	5.6	14.3	12.9	21.1	19.7
2001.....	*	*	*	3.3	4.0	8.4	12.3	14.4	14.7	23.2
2000.....	*	*	*	3.6	5.2	7.3	12.4	18.2	19.5	18.5
1999.....	*	*	*	3.0	5.0	7.5	10.1	20.9	14.3	24.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic obstructive pulmonary disease (J44)										
All races 3/, both sexes										
2006.....	1.3	3.8	9.7	21.0	48.1	95.7	167.9	268.8	380.7	519.0
2005.....	1.4	4.2	9.7	21.4	51.6	100.1	179.6	285.1	398.5	557.0
2004.....	1.4	3.6	8.6	20.1	49.4	94.6	169.7	270.4	376.4	524.8
2003.....	1.3	3.5	8.5	21.3	51.9	100.9	177.4	279.4	390.4	552.9
2002.....	1.4	3.5	8.6	21.6	48.8	97.8	174.2	277.3	395.7	549.7
2001.....	1.3	3.3	8.3	22.3	49.4	100.0	176.2	273.1	382.9	550.3
2000.....	1.2	3.3	8.2	22.1	48.8	98.6	175.8	270.2	396.1	550.6
1999.....	1.1	3.1	8.0	23.3	51.2	101.7	182.1	278.2	400.8	549.6
All races 3/, male										
2006.....	1.3	3.8	10.1	22.7	52.5	102.7	189.3	318.7	478.8	678.3
2005.....	1.4	4.2	10.6	22.6	56.1	112.1	205.8	339.6	495.4	727.8
2004.....	1.4	3.6	9.7	21.4	53.3	104.9	192.8	326.3	474.0	714.1
2003.....	1.2	4.0	9.1	23.2	56.7	111.9	205.7	336.6	490.1	764.9
2002.....	1.5	3.7	9.4	23.2	52.9	108.7	203.4	341.3	508.8	780.4
2001.....	1.3	3.7	8.9	24.1	53.5	111.0	207.7	339.9	498.3	789.6
2000.....	1.2	3.8	8.7	24.2	54.2	111.1	209.8	335.5	530.2	833.7
1999.....	1.2	3.6	8.5	25.7	58.4	119.2	222.9	355.4	540.9	860.0
All races 3/, female										
2006.....	1.4	3.8	9.2	19.4	44.0	89.6	150.6	232.2	319.4	444.4
2005.....	1.4	4.2	8.8	20.2	47.5	89.7	158.4	245.5	338.3	478.5
2004.....	1.3	3.5	7.7	18.8	46.0	85.6	151.1	230.1	316.4	439.6
2003.....	1.4	3.1	8.0	19.5	47.5	91.3	154.7	238.7	329.5	459.2
2002.....	1.2	3.2	7.8	20.2	45.1	88.4	150.9	232.1	326.9	449.5
2001.....	1.4	2.8	7.7	20.6	45.7	90.5	151.2	226.3	313.7	450.2
2000.....	1.2	2.9	7.7	20.2	43.9	87.8	149.0	224.6	317.1	435.3
1999.....	1.1	2.5	7.6	21.1	44.7	86.7	150.2	224.6	318.9	424.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic obstructive pulmonary disease (J44)

White, both sexes

2006 4/.....	1.3	4.1	10.0	22.0	51.0	102.6	181.1	284.8	401.3	545.3
2005 4/.....	1.4	4.3	9.8	22.4	54.9	107.1	192.2	301.9	420.3	583.8
2004 4/.....	1.4	3.7	8.9	20.9	52.6	101.4	181.4	285.7	396.6	550.5
2003 4/.....	1.3	3.7	8.8	22.1	55.2	108.2	188.3	294.5	411.1	579.4
2002.....	1.4	3.6	8.8	22.7	51.4	104.8	184.5	291.4	413.7	574.7
2001.....	1.3	3.3	8.5	23.3	52.7	105.9	186.4	287.4	400.3	573.9
2000.....	1.1	3.3	8.3	23.0	51.5	104.3	185.5	283.0	414.8	574.7
1999.....	1.1	3.0	8.3	24.2	53.8	107.3	190.7	289.4	417.8	573.7

White, male

2006 4/.....	1.3	4.0	10.3	23.4	54.1	107.5	199.6	331.6	495.6	702.9
2005 4/.....	1.4	4.2	10.5	22.9	58.5	117.4	215.6	351.3	513.7	751.8
2004 4/.....	1.5	3.7	9.8	21.5	55.3	110.4	201.6	339.0	491.5	738.3
2003 4/.....	1.2	4.2	9.2	23.5	59.1	117.6	213.1	347.1	504.7	791.5
2002.....	1.5	3.8	9.4	23.6	54.4	114.4	210.6	351.3	523.6	805.8
2001.....	1.3	3.8	8.9	24.7	55.8	114.9	214.9	350.1	512.4	812.9
2000.....	1.2	3.7	8.5	24.7	56.0	114.7	216.3	344.3	546.6	862.1
1999.....	1.1	3.5	8.6	26.1	60.0	123.0	229.5	362.6	554.3	890.0

White, female

2006 4/.....	1.3	4.1	9.6	20.7	48.0	98.3	165.7	249.8	341.7	471.4
2005 4/.....	1.4	4.5	9.1	21.8	51.6	98.0	172.9	265.4	361.9	506.4
2004 4/.....	1.4	3.6	8.1	20.4	50.1	93.5	164.8	246.7	337.6	465.8
2003 4/.....	1.4	3.3	8.5	20.7	51.5	99.9	168.0	256.6	353.4	485.4
2002.....	1.3	3.5	8.1	21.9	48.7	96.3	163.2	248.5	346.4	474.3
2001.....	1.4	2.8	8.1	22.0	49.8	98.0	163.3	243.0	332.6	473.8
2000.....	1.1	3.0	8.1	21.4	47.3	95.2	160.8	239.8	336.4	457.9
1999.....	1.1	2.5	8.1	22.4	48.2	93.5	159.9	237.9	337.2	446.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other chronic obstructive pulmonary disease (J44)										
Black, both sexes										
2006 4/.....	1.8	3.3	11.3	20.2	38.9	69.4	103.3	172.2	227.9	290.0
2005 4/.....	1.8	4.9	11.9	21.1	40.6	73.5	119.5	185.4	230.4	324.7
2004 4/.....	1.5	4.0	9.8	19.3	38.8	67.3	107.6	173.5	214.1	284.5
2003 4/.....	1.7	3.7	9.3	21.7	40.3	69.4	121.9	177.5	227.7	304.3
2002.....	1.7	3.8	10.0	20.3	41.6	67.2	116.4	183.8	250.9	318.9
2001.....	1.6	4.0	9.3	20.6	37.3	75.5	118.7	172.5	237.0	319.7
2000.....	1.8	3.8	10.1	21.3	41.9	72.0	119.9	174.1	227.0	310.0
1999.....	1.9	4.7	8.4	23.6	42.8	76.8	129.1	197.6	247.0	312.8
Black, male										
2006 4/.....	1.5	3.8	13.0	24.8	52.8	94.2	147.3	247.9	367.6	461.1
2005 4/.....	1.7	6.0	14.5	27.2	52.4	102.7	170.8	293.7	362.4	529.9
2004 4/.....	1.5	3.8	12.3	27.8	51.4	90.1	153.4	257.5	342.9	490.8
2003 4/.....	1.8	4.5	10.8	28.5	52.4	93.8	180.3	272.5	409.0	520.3
2002.....	2.0	4.7	12.5	27.4	56.0	88.3	172.9	290.7	405.8	565.0
2001.....	1.6	4.3	11.3	26.4	50.2	105.2	178.7	283.9	405.3	580.9
2000.....	1.7	4.9	12.5	27.3	55.7	102.8	186.5	283.5	383.6	576.6
1999.....	2.4	5.9	10.8	30.9	60.9	109.6	191.6	323.9	440.3	605.8
Black, female										
2006 4/.....	2.1	2.8	9.8	16.5	28.0	51.1	73.1	127.0	154.1	220.1
2005 4/.....	1.8	4.0	9.7	16.1	31.5	51.8	84.4	120.8	160.6	241.9
2004 4/.....	1.4	4.2	7.6	12.3	29.0	50.4	76.4	123.7	146.3	202.8
2003 4/.....	1.5	3.0	7.9	16.2	30.8	51.4	82.4	121.0	132.7	220.4
2002.....	1.4	2.9	7.8	14.4	30.4	51.6	78.4	119.9	170.3	224.7
2001.....	1.5	3.7	7.7	15.8	27.2	53.5	78.7	105.9	151.3	222.7
2000.....	1.9	2.8	7.9	16.3	31.1	49.3	75.7	108.2	148.2	210.8
1999.....	1.5	3.7	6.4	17.7	28.6	52.5	87.5	121.6	151.2	203.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
All races 3/, both sexes										
2006.....	1.0	1.2	1.4	1.4	1.8	2.0	2.7	3.6	6.4	12.3
2005.....	1.0	1.4	1.4	1.5	1.9	2.3	2.9	4.1	7.1	14.5
2004.....	0.9	1.3	1.4	1.7	1.9	2.5	3.0	4.2	6.6	14.1
2003.....	1.1	1.5	1.7	1.9	2.2	2.7	3.1	4.9	7.2	14.2
2002.....	1.2	1.4	1.6	1.9	2.1	2.7	3.8	5.4	7.7	14.7
2001.....	1.1	1.3	1.8	1.9	2.6	3.0	3.9	5.2	8.1	16.1
2000.....	1.2	1.6	1.7	2.0	2.5	3.4	4.6	5.7	8.7	16.7
1999.....	1.1	1.6	1.7	2.6	2.2	3.7	4.7	6.2	9.6	17.3
All races 3/, male										
2006.....	0.8	0.9	1.1	1.0	1.2	1.4	2.1	2.9	4.4	9.1
2005.....	0.8	0.9	1.2	1.1	1.2	1.7	1.9	3.7	4.8	8.7
2004.....	0.8	1.1	1.0	1.4	1.3	2.0	2.3	2.8	4.4	11.1
2003.....	0.9	1.2	1.2	1.5	1.7	1.8	2.4	3.7	5.7	10.7
2002.....	0.9	1.1	1.3	1.5	1.7	2.0	2.8	4.3	6.2	10.9
2001.....	0.9	1.0	1.3	1.2	1.9	2.4	3.0	3.8	5.6	12.1
2000.....	1.0	1.3	1.3	1.6	1.8	2.7	3.4	3.9	6.7	12.9
1999.....	0.8	1.0	1.2	1.9	1.7	2.6	3.9	4.8	7.2	12.7
All races 3/, female										
2006.....	1.2	1.6	1.7	1.7	2.3	2.6	3.2	4.2	7.6	13.9
2005.....	1.2	1.8	1.7	1.9	2.6	2.8	3.7	4.4	8.6	17.2
2004.....	1.1	1.5	1.8	2.0	2.5	2.9	3.6	5.1	7.9	15.4
2003.....	1.4	1.7	2.1	2.2	2.6	3.5	3.7	5.7	8.0	15.8
2002.....	1.4	1.8	1.9	2.3	2.5	3.3	4.5	6.2	8.7	16.3
2001.....	1.3	1.7	2.2	2.5	3.3	3.6	4.7	6.1	9.6	17.8
2000.....	1.4	1.9	2.1	2.4	3.1	4.0	5.5	6.9	9.8	18.2
1999.....	1.3	2.2	2.2	3.3	2.7	4.6	5.4	7.2	11.1	19.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
White, both sexes										
2006 4/.....	0.8	0.9	1.0	1.0	1.4	1.7	2.3	3.4	6.0	12.1
2005 4/.....	0.8	1.0	1.1	1.1	1.6	1.8	2.3	3.8	6.4	14.5
2004 4/.....	0.7	0.8	1.0	1.3	1.4	2.1	2.8	3.6	6.2	13.7
2003 4/.....	0.8	1.1	1.1	1.5	1.8	2.3	2.6	4.4	6.7	14.3
2002.....	0.8	1.1	1.1	1.4	1.8	2.2	3.2	5.1	7.5	14.7
2001.....	0.8	1.0	1.4	1.4	1.9	2.6	3.4	4.7	7.8	15.7
2000.....	0.9	1.1	1.2	1.4	2.0	2.8	4.0	5.2	8.4	15.9
1999.....	0.8	1.2	1.3	2.0	1.8	3.1	4.0	5.8	9.4	16.6
White, male										
2006 4/.....	0.6	0.6	0.7	0.6	0.8	1.1	1.6	2.7	3.9	9.1
2005 4/.....	0.6	0.6	1.0	0.8	0.9	1.3	1.4	3.1	4.1	8.6
2004 4/.....	0.6	0.6	0.7	0.9	0.8	1.6	1.9	2.2	3.8	10.3
2003 4/.....	0.6	0.9	0.8	1.2	1.2	1.5	2.0	3.0	5.4	10.6
2002.....	0.6	0.8	0.9	1.0	1.4	1.5	2.3	3.8	5.8	10.7
2001.....	0.6	0.7	0.9	1.0	1.2	2.0	2.4	3.3	5.1	11.3
2000.....	0.7	1.0	1.0	1.0	1.4	2.0	3.0	3.7	6.3	11.4
1999.....	0.5	0.7	0.9	1.5	1.2	2.0	3.0	4.2	7.0	10.8
White, female										
2006 4/.....	1.0	1.2	1.3	1.4	1.8	2.3	2.9	3.9	7.4	13.6
2005 4/.....	0.9	1.3	1.2	1.4	2.2	2.2	3.0	4.3	7.9	17.2
2004 4/.....	0.8	1.0	1.4	1.6	2.1	2.6	3.5	4.7	7.7	15.2
2003 4/.....	1.0	1.3	1.5	1.7	2.3	3.0	3.1	5.4	7.5	15.9
2002.....	1.0	1.3	1.3	1.7	2.2	2.8	3.9	6.1	8.6	16.4
2001.....	1.0	1.2	1.8	1.9	2.5	3.2	4.2	5.7	9.4	17.6
2000.....	1.0	1.2	1.5	1.8	2.6	3.4	4.8	6.3	9.7	17.7
1999.....	1.1	1.7	1.7	2.6	2.2	4.0	4.8	6.9	10.8	18.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Asthma (J45-J46)										
Black, both sexes										
2006 4/.....	2.7	3.6	4.3	4.4	6.1	4.9	6.3	5.0	7.3	12.5
2005 4/.....	2.8	4.4	3.9	5.4	5.3	6.0	8.7	7.9	12.6	11.8
2004 4/.....	2.9	4.8	4.1	5.1	6.3	5.7	5.6	9.0	9.4	15.5
2003 4/.....	3.4	4.1	5.7	5.2	5.0	6.6	7.5	8.2	10.4	12.0
2002.....	4.0	4.1	5.7	7.1	5.6	6.1	10.1	7.4	8.9	11.9
2001.....	3.1	4.3	4.9	5.9	7.9	6.3	9.4	10.2	11.1	16.3
2000.....	3.6	5.4	5.3	6.6	7.2	8.8	9.7	10.7	9.9	22.2
1999.....	3.3	4.8	5.8	7.3	6.2	8.4	11.1	9.8	11.3	20.4
Black, male										
2006 4/.....	2.6	2.9	4.2	4.1	5.7	*	*	*	*	*
2005 4/.....	2.5	3.3	2.5	4.3	*	4.8	6.3	9.8	*	*
2004 4/.....	2.7	4.7	2.9	4.8	5.8	4.8	6.5	*	*	*
2003 4/.....	2.7	3.5	4.4	3.8	4.8	4.9	*	9.1	*	*
2002.....	3.7	3.4	4.6	6.0	5.3	6.2	8.7	*	*	*
2001.....	2.9	3.8	4.5	3.6	5.9	*	9.8	*	*	*
2000.....	3.1	4.0	3.5	6.3	6.0	8.6	7.0	*	*	25.4
1999.....	2.9	3.6	4.2	5.1	5.7	6.8	12.2	9.5	*	*
Black, female										
2006 4/.....	2.7	4.2	4.4	4.6	6.5	5.5	6.8	5.9	8.3	15.0
2005 4/.....	3.1	5.3	5.1	6.4	6.7	6.9	10.4	6.7	13.0	13.5
2004 4/.....	3.1	4.9	5.1	5.4	6.7	6.3	5.1	9.7	9.1	16.9
2003 4/.....	4.1	4.7	6.7	6.3	5.1	7.9	8.6	7.6	11.6	13.8
2002.....	4.2	4.7	6.6	8.1	5.7	5.9	11.0	8.0	9.5	12.3
2001.....	3.3	4.8	5.3	7.8	9.4	7.5	9.2	11.4	11.9	16.8
2000.....	4.0	6.6	6.9	6.9	8.2	9.0	11.5	14.2	10.2	21.0
1999.....	3.6	5.9	7.1	9.1	6.5	9.6	10.4	10.0	13.8	21.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchiectasis (J47)										
All races 3/, both sexes										
2006.....	*	*	0.1	0.1	0.3	0.5	1.1	1.9	4.1	6.5
2005.....	*	*	*	0.1	0.2	0.4	1.1	2.4	4.0	6.5
2004.....	*	*	0.1	0.1	0.2	0.5	1.2	2.5	4.0	6.3
2003.....	*	*	*	0.2	0.3	0.5	1.1	2.3	4.1	6.0
2002.....	*	*	*	0.1	0.4	0.6	1.2	2.5	4.6	5.8
2001.....	*	*	*	0.2	0.4	0.7	1.2	2.3	3.9	7.0
2000.....	*	*	*	0.3	0.3	0.7	1.2	2.4	4.6	6.1
1999.....	*	*	*	0.2	0.4	0.8	1.1	2.6	4.7	6.2
All races 3/, male										
2006.....	*	*	*	*	*	*	0.8	1.6	3.0	3.9
2005.....	*	*	*	*	*	*	0.8	1.4	2.5	4.6
2004.....	*	*	*	*	*	*	0.8	1.9	2.3	4.5
2003.....	*	*	*	*	*	*	0.8	1.6	3.2	4.2
2002.....	*	*	*	*	*	*	0.9	1.6	3.1	4.7
2001.....	*	*	*	*	0.4	0.6	0.8	1.6	2.2	4.8
2000.....	*	*	*	*	*	0.5	1.0	1.6	3.2	4.3
1999.....	*	*	*	*	*	*	1.0	2.5	4.1	6.6
All races 3/, female										
2006.....	*	*	*	*	0.3	0.7	1.4	2.1	4.8	7.7
2005.....	*	*	*	*	*	0.5	1.4	3.2	4.9	7.4
2004.....	*	*	*	*	*	0.6	1.5	2.9	5.0	7.2
2003.....	*	*	*	0.3	*	0.8	1.3	2.8	4.7	6.8
2002.....	*	*	*	*	0.5	0.9	1.4	3.1	5.5	6.3
2001.....	*	*	*	*	0.4	0.9	1.5	2.8	4.9	7.9
2000.....	*	*	*	0.3	0.4	0.9	1.4	3.0	5.5	6.9
1999.....	*	*	*	0.3	0.4	1.1	1.2	2.6	5.1	6.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Bronchiectasis (J47)										
White, both sexes										
2006 4/.....	*	*	*	*	0.3	0.6	1.2	2.0	4.3	6.8
2005 4/.....	*	*	*	0.2	0.2	0.5	1.2	2.6	4.3	6.9
2004 4/.....	*	*	0.1	*	0.3	0.5	1.2	2.6	4.3	6.7
2003 4/.....	*	*	*	0.1	0.3	0.5	1.1	2.4	4.3	6.4
2002.....	*	*	*	0.2	0.5	0.6	1.3	2.7	4.9	6.1
2001.....	*	*	*	*	0.4	0.8	1.2	2.4	4.1	7.4
2000.....	*	*	*	0.2	0.3	0.7	1.2	2.5	4.9	6.6
1999.....	*	*	*	0.2	0.4	0.8	1.1	2.6	4.8	6.7
White, male										
2006 4/.....	*	*	*	*	*	*	0.8	1.5	3.1	3.7
2005 4/.....	*	*	*	*	*	*	0.8	1.5	2.6	4.6
2004 4/.....	*	*	*	*	*	*	0.8	1.9	2.5	4.8
2003 4/.....	*	*	*	*	*	*	0.7	1.7	3.3	4.5
2002.....	*	*	*	*	*	*	0.8	1.7	3.2	4.7
2001.....	*	*	*	*	0.5	*	0.8	1.5	2.1	4.8
2000.....	*	*	*	*	*	*	1.0	1.5	3.5	4.6
1999.....	*	*	*	*	*	*	1.0	2.4	4.1	6.9
White, female										
2006 4/.....	*	*	*	*	0.3	0.8	1.5	2.3	5.1	8.2
2005 4/.....	*	*	*	*	*	0.5	1.5	3.3	5.3	8.0
2004 4/.....	*	*	*	*	*	0.7	1.6	3.1	5.4	7.6
2003 4/.....	*	*	*	*	*	0.8	1.4	3.0	5.0	7.3
2002.....	*	*	*	*	0.6	0.9	1.6	3.3	5.9	6.7
2001.....	*	*	*	*	*	1.0	1.6	3.0	5.3	8.5
2000.....	*	*	*	*	0.4	1.0	1.3	3.2	5.7	7.4
1999.....	*	*	*	*	0.5	1.2	1.2	2.7	5.2	6.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lung diseases due to external agents (J60-J70)										
All races 3/, both sexes										
2006.....	0.5	0.9	1.3	2.1	3.9	7.0	14.4	28.9	63.1	161.1
2005.....	0.4	0.9	1.6	2.2	4.1	7.0	14.4	31.2	65.9	169.8
2004.....	0.5	0.8	1.3	2.1	3.8	7.0	15.4	31.6	66.5	171.1
2003.....	0.5	0.9	1.2	2.2	4.2	7.3	15.1	33.3	70.6	182.8
2002.....	0.6	0.7	1.2	2.0	3.9	7.5	15.2	34.9	71.7	192.9
2001.....	0.5	0.7	1.3	2.1	4.0	7.3	16.2	34.0	73.3	195.9
2000.....	0.5	0.8	1.2	2.1	3.5	7.7	16.5	34.0	70.6	195.3
1999.....	0.5	0.7	1.0	2.1	3.5	7.3	15.3	31.9	66.1	181.8
All races 3/, male										
2006.....	0.6	1.0	1.8	2.8	5.4	10.0	20.9	42.6	96.3	240.5
2005.....	0.5	1.0	1.9	2.8	5.5	9.7	20.8	44.4	97.9	246.9
2004.....	0.6	1.0	1.6	2.6	5.3	10.0	22.2	46.0	98.0	256.1
2003.....	0.5	1.1	1.5	2.9	5.6	10.7	21.5	48.8	105.0	266.2
2002.....	0.7	0.9	1.6	2.6	5.1	10.5	22.0	51.6	105.9	290.1
2001.....	0.7	0.8	1.8	2.8	5.0	10.4	23.7	49.1	111.6	291.3
2000.....	0.7	1.1	1.6	2.7	4.6	11.2	25.2	50.6	107.8	301.5
1999.....	0.6	0.9	1.2	2.7	5.1	10.7	22.9	50.0	103.9	279.1
All races 3/, female										
2006.....	0.4	0.7	0.9	1.5	2.6	4.4	9.2	18.8	42.3	124.0
2005.....	0.4	0.7	1.2	1.7	2.7	4.7	9.2	21.6	46.1	134.4
2004.....	0.3	0.6	0.9	1.5	2.5	4.3	10.0	21.2	47.1	132.9
2003.....	0.5	0.7	0.8	1.5	3.0	4.3	10.0	22.3	49.6	145.9
2002.....	0.4	0.5	0.8	1.5	2.9	5.0	9.9	23.2	50.9	150.7
2001.....	0.4	0.6	0.9	1.4	3.0	4.6	10.3	23.5	50.3	156.0
2000.....	0.4	0.6	0.9	1.6	2.5	4.8	9.7	22.4	48.6	152.1
1999.....	0.4	0.5	0.9	1.6	2.1	4.3	9.3	19.3	44.0	142.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lung diseases due to external agents (J60-J70)										
White, both sexes										
2006 4/.....	0.5	0.8	1.3	2.1	4.0	6.9	14.2	28.9	64.6	165.7
2005 4/.....	0.5	0.9	1.5	2.2	3.9	6.8	14.4	30.8	66.5	172.7
2004 4/.....	0.4	0.8	1.2	2.0	3.5	6.9	15.2	31.7	67.4	174.3
2003 4/.....	0.4	0.9	1.2	2.2	3.8	7.0	15.1	33.1	70.4	185.7
2002.....	0.5	0.7	1.1	2.0	3.8	7.3	15.1	34.7	72.8	196.8
2001.....	0.5	0.7	1.2	2.1	3.8	7.2	15.8	34.1	73.7	197.6
2000.....	0.6	0.7	1.2	2.1	3.3	7.5	15.9	33.8	70.8	199.2
1999.....	0.4	0.6	0.9	2.0	3.3	7.0	14.7	31.5	66.1	184.4
White, male										
2006 4/.....	0.6	1.0	1.7	2.7	5.4	9.7	20.6	42.1	98.7	248.5
2005 4/.....	0.5	1.0	1.8	2.8	5.2	9.2	20.8	43.6	98.5	252.7
2004 4/.....	0.5	1.0	1.6	2.5	4.7	9.9	21.6	46.1	99.6	261.3
2003 4/.....	0.5	1.1	1.6	2.9	5.1	10.5	21.2	48.2	105.0	272.3
2002.....	0.7	0.9	1.5	2.6	4.9	10.1	21.6	51.1	107.8	297.5
2001.....	0.7	0.8	1.6	2.8	4.7	10.4	23.2	49.4	112.8	297.5
2000.....	0.7	0.9	1.4	2.7	4.4	10.6	23.9	50.0	109.3	311.0
1999.....	0.5	0.8	1.0	2.5	4.7	10.2	21.9	49.2	103.7	285.0
White, female										
2006 4/.....	0.4	0.7	0.9	1.5	2.7	4.4	8.9	19.1	43.1	126.8
2005 4/.....	0.4	0.7	1.2	1.6	2.7	4.6	9.1	21.4	46.4	135.9
2004 4/.....	0.3	0.6	0.8	1.6	2.4	4.3	10.0	21.3	47.4	135.0
2003 4/.....	0.4	0.7	0.8	1.5	2.7	3.9	10.0	22.1	49.1	147.3
2002.....	0.4	0.5	0.8	1.5	2.7	4.7	9.8	23.0	51.3	153.0
2001.....	0.4	0.5	0.9	1.4	3.0	4.3	9.8	23.3	50.0	155.8
2000.....	0.4	0.5	0.9	1.5	2.3	4.8	9.4	22.5	47.9	153.8
1999.....	0.4	0.4	0.9	1.5	2.1	4.2	9.0	19.0	43.9	144.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Lung diseases due to external agents (J60-J70)										
Black, both sexes										
2006 4/.....	0.9	1.2	1.9	3.2	5.0	9.6	20.5	34.2	58.0	131.0
2005 4/.....	*	1.2	2.5	3.2	6.8	10.7	17.2	42.3	71.8	156.9
2004 4/.....	0.9	1.0	1.7	3.1	7.6	9.5	21.5	35.4	68.1	149.7
2003 4/.....	0.9	1.4	1.3	2.8	8.6	11.6	17.9	42.2	78.9	166.0
2002.....	1.0	0.9	2.1	2.6	5.9	12.0	19.8	44.0	70.6	169.3
2001.....	0.8	1.1	2.5	2.6	6.4	9.8	23.4	39.9	79.6	189.2
2000.....	*	1.7	2.2	3.0	5.9	11.8	26.8	41.6	74.3	163.6
1999.....	1.0	1.2	2.0	4.0	5.9	10.9	24.8	42.7	73.4	172.2
Black, male										
2006 4/.....	*	*	2.9	4.2	7.8	15.7	29.8	58.6	92.4	181.4
2005 4/.....	*	1.6	3.5	3.6	10.5	16.2	26.2	65.7	112.4	214.7
2004 4/.....	*	*	2.3	4.5	11.8	15.3	33.7	53.2	101.9	220.5
2003 4/.....	*	1.8	*	3.9	11.6	16.0	27.9	65.7	115.2	221.0
2002.....	*	*	3.2	3.7	7.3	17.5	30.7	67.5	99.3	247.7
2001.....	*	*	3.5	3.6	9.2	13.5	33.7	59.9	118.8	257.6
2000.....	*	2.4	3.3	3.3	7.7	18.7	45.1	69.0	103.7	223.7
1999.....	*	*	2.7	5.3	10.0	17.4	40.0	69.8	116.0	246.1
Black, female										
2006 4/.....	*	*	*	2.3	2.8	5.0	14.2	19.6	39.8	110.4
2005 4/.....	*	*	1.6	2.9	3.8	6.7	11.0	28.3	50.4	133.5
2004 4/.....	*	*	*	*	4.4	5.2	13.2	24.8	50.3	121.7
2003 4/.....	*	*	*	*	6.3	8.4	11.1	28.2	60.0	144.6
2002.....	*	*	*	*	4.8	8.0	12.5	29.9	55.7	139.2
2001.....	*	*	*	*	4.3	7.1	16.6	28.0	59.7	163.8
2000.....	*	*	*	2.7	4.6	6.7	14.7	25.1	59.4	141.2
1999.....	*	*	*	2.9	*	6.2	14.6	26.3	52.3	144.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)										
All races 3/, both sexes										
2006.....	0.5	0.8	1.2	2.0	3.7	6.3	12.8	26.5	58.7	155.8
2005.....	0.4	0.8	1.5	2.0	3.7	6.4	12.9	28.5	61.1	163.9
2004.....	0.5	0.8	1.2	1.9	3.4	6.0	13.7	28.4	62.2	164.9
2003.....	0.4	0.9	1.1	2.0	3.8	6.3	13.3	30.1	65.4	176.2
2002.....	0.5	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.5	0.7	1.3	1.8	3.6	6.2	14.1	30.4	68.0	189.4
2000.....	0.5	0.8	1.2	1.9	3.2	6.7	14.2	30.4	65.7	187.6
1999.....	0.5	0.7	0.9	2.0	3.1	6.1	13.2	28.2	60.7	175.6
All races 3/, male										
2006.....	0.6	1.0	1.6	2.5	4.9	8.7	17.5	37.3	85.7	224.8
2005.....	0.4	0.9	1.8	2.4	4.9	8.4	17.5	38.3	86.2	229.3
2004.....	0.6	1.0	1.5	2.3	4.5	8.1	18.7	39.0	87.4	237.2
2003.....	0.4	1.0	1.4	2.6	4.8	8.6	17.8	41.5	92.3	246.2
2002.....	0.7	0.9	1.5	2.2	4.2	8.6	18.2	43.4	95.1	268.5
2001.....	0.7	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.1	270.3
2000.....	0.6	1.0	1.5	2.4	3.9	9.2	20.2	42.4	95.2	276.0
1999.....	0.6	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4
All races 3/, female										
2006.....	0.4	0.7	0.9	1.5	2.6	4.3	8.9	18.6	41.9	123.5
2005.....	0.4	0.7	1.1	1.6	2.7	4.6	9.1	21.3	45.5	133.9
2004.....	0.3	0.5	0.8	1.5	2.5	4.2	9.8	20.9	46.6	132.4
2003.....	0.4	0.7	0.8	1.5	2.9	4.3	9.7	22.1	49.0	145.3
2002.....	0.4	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.4	150.2
2001.....	0.4	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.6
2000.....	0.4	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999.....	0.4	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.5	142.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)										
White, both sexes										
2006 4/.....	0.5	0.8	1.2	1.9	3.7	6.2	12.4	26.3	60.1	160.0
2005 4/.....	0.5	0.8	1.4	2.0	3.5	6.0	12.7	27.9	61.4	166.5
2004 4/.....	0.4	0.7	1.1	1.8	3.1	5.9	13.4	28.2	62.9	167.8
2003 4/.....	0.4	0.8	1.1	2.0	3.3	5.9	13.1	29.7	65.0	178.7
2002.....	0.5	0.7	1.0	1.8	3.3	6.1	13.1	30.9	68.1	189.5
2001.....	0.5	0.7	1.2	1.8	3.5	6.0	13.6	30.2	68.1	190.8
2000.....	0.5	0.7	1.1	1.9	2.9	6.4	13.4	30.1	65.7	191.0
1999.....	0.4	0.6	0.8	1.8	2.9	5.8	12.5	27.6	60.3	178.0
White, male										
2006 4/.....	0.6	1.0	1.6	2.4	4.9	8.3	17.0	36.4	87.5	231.8
2005 4/.....	0.5	0.9	1.7	2.4	4.5	7.8	17.1	37.1	86.1	234.2
2004 4/.....	0.5	1.0	1.4	2.2	3.9	7.8	17.9	38.3	88.4	241.7
2003 4/.....	0.4	1.0	1.4	2.5	4.2	8.2	17.2	40.6	91.7	250.8
2002.....	0.6	0.8	1.3	2.1	4.1	8.0	17.6	42.5	96.5	274.6
2001.....	0.7	0.8	1.5	2.3	4.0	8.0	18.4	40.6	98.7	275.4
2000.....	0.7	0.9	1.3	2.3	3.6	8.5	18.7	41.4	96.2	284.0
1999.....	0.5	0.8	0.9	2.2	3.9	7.7	17.1	40.4	88.9	263.6
White, female										
2006 4/.....	0.3	0.6	0.9	1.4	2.7	4.3	8.6	18.8	42.7	126.3
2005 4/.....	0.4	0.7	1.1	1.6	2.6	4.5	9.0	21.1	45.9	135.3
2004 4/.....	0.3	0.5	0.8	1.5	2.3	4.2	9.7	20.8	47.0	134.5
2003 4/.....	0.4	0.7	0.8	1.5	2.5	3.9	9.7	21.8	48.6	146.7
2002.....	0.4	0.5	0.7	1.5	2.6	4.5	9.5	22.6	50.7	152.5
2001.....	0.4	0.5	0.8	1.3	3.0	4.2	9.7	22.8	49.6	155.3
2000.....	0.4	0.5	0.9	1.4	2.3	4.6	9.1	22.2	47.5	153.2
1999.....	0.4	0.4	0.8	1.5	2.0	4.1	8.8	18.6	43.4	143.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to solids and liquids (J69)										
Black, both sexes										
2006 4/.....	0.9	1.2	1.7	2.9	4.9	9.2	19.8	32.8	55.1	128.4
2005 4/.....	*	1.2	2.4	3.1	6.4	10.4	17.0	40.9	69.2	153.1
2004 4/.....	0.9	1.0	1.7	3.0	7.3	8.8	20.4	34.7	64.9	144.7
2003 4/.....	0.9	1.3	1.2	2.7	8.5	11.0	17.2	39.8	76.2	163.4
2002.....	0.9	0.8	2.1	2.4	5.5	11.7	18.8	41.9	67.8	165.1
2001.....	0.7	1.0	2.4	2.4	6.0	9.3	22.3	38.6	76.1	185.5
2000.....	*	1.7	2.2	2.9	5.8	11.3	25.1	38.9	71.4	159.8
1999.....	1.0	1.2	1.8	3.9	5.0	10.1	23.7	40.5	71.1	167.1
Black, male										
2006 4/.....	*	*	2.4	3.7	7.6	15.0	28.0	55.0	84.9	173.4
2005 4/.....	*	1.5	3.4	3.6	9.8	15.4	25.6	62.1	106.0	201.7
2004 4/.....	*	*	2.2	4.4	11.0	13.6	31.4	51.4	93.3	202.7
2003 4/.....	*	1.7	*	3.8	11.6	14.5	26.3	59.3	107.8	212.7
2002.....	*	*	3.1	3.7	6.3	16.7	28.4	62.5	91.7	233.7
2001.....	*	*	3.4	3.1	8.1	12.5	31.0	56.2	108.3	243.9
2000.....	*	2.4	3.1	3.3	7.2	17.4	41.1	62.4	96.1	209.9
1999.....	*	*	*	5.1	8.1	15.6	37.3	65.1	109.9	228.5
Black, female										
2006 4/.....	*	*	*	2.3	2.8	4.8	14.2	19.6	39.4	110.0
2005 4/.....	*	*	*	2.7	3.7	6.7	11.0	28.3	49.7	133.5
2004 4/.....	*	*	*	*	4.4	5.2	13.0	24.8	49.9	121.7
2003 4/.....	*	*	*	*	6.0	8.4	11.1	28.2	59.6	144.2
2002.....	*	*	*	*	4.8	8.0	12.3	29.6	55.3	138.8
2001.....	*	*	*	*	4.3	6.9	16.6	28.0	59.7	163.8
2000.....	*	*	*	*	4.6	6.7	14.4	24.8	59.0	141.2
1999.....	*	*	*	2.9	*	6.0	14.6	25.7	51.9	144.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)										
All races 3/, both sexes										
2006.....	0.5	0.8	1.2	2.0	3.7	6.3	12.8	26.5	58.7	155.8
2005.....	0.4	0.8	1.5	2.0	3.7	6.4	12.9	28.5	61.1	163.9
2004.....	0.5	0.8	1.2	1.9	3.4	6.0	13.7	28.4	62.2	164.9
2003.....	0.4	0.9	1.1	2.0	3.8	6.3	13.3	30.1	65.4	176.2
2002.....	0.5	0.7	1.1	1.8	3.4	6.5	13.4	31.3	67.3	186.0
2001.....	0.5	0.7	1.3	1.8	3.6	6.2	14.1	30.4	67.9	189.4
2000.....	0.5	0.8	1.2	1.9	3.1	6.7	14.2	30.4	65.7	187.6
1999.....	0.5	0.7	0.9	2.0	3.0	6.1	13.2	28.1	60.6	175.6
All races 3/, male										
2006.....	0.6	1.0	1.6	2.5	4.9	8.7	17.5	37.2	85.7	224.8
2005.....	0.4	0.9	1.8	2.4	4.9	8.4	17.5	38.3	86.2	229.3
2004.....	0.6	1.0	1.5	2.3	4.5	8.1	18.7	38.9	87.3	237.2
2003.....	0.4	1.0	1.4	2.6	4.8	8.6	17.8	41.5	92.2	246.2
2002.....	0.7	0.9	1.5	2.2	4.2	8.5	18.2	43.4	95.1	268.5
2001.....	0.7	0.8	1.7	2.3	4.2	8.2	19.1	41.0	98.0	270.2
2000.....	0.6	1.0	1.5	2.3	3.9	9.2	20.2	42.4	95.2	276.0
1999.....	0.6	0.9	1.0	2.4	4.2	8.3	18.4	41.6	90.1	258.4
All races 3/, female										
2006.....	0.4	0.7	0.9	1.5	2.6	4.3	8.9	18.6	41.9	123.5
2005.....	0.4	0.7	1.1	1.6	2.7	4.6	9.1	21.3	45.5	133.9
2004.....	0.3	0.5	0.8	1.5	2.5	4.2	9.8	20.9	46.6	132.4
2003.....	0.4	0.7	0.8	1.5	2.9	4.3	9.7	22.0	49.0	145.3
2002.....	0.4	0.5	0.8	1.4	2.8	4.8	9.6	22.8	50.3	150.2
2001.....	0.4	0.6	0.9	1.4	3.0	4.4	10.2	23.0	49.9	155.5
2000.....	0.4	0.6	0.9	1.5	2.5	4.6	9.4	22.1	48.2	151.6
1999.....	0.4	0.5	0.8	1.6	2.0	4.2	9.2	18.8	43.4	142.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)										
White, both sexes										
2006 4/.....	0.5	0.8	1.2	1.9	3.7	6.2	12.4	26.3	60.1	160.0
2005 4/.....	0.5	0.8	1.4	2.0	3.5	6.0	12.7	27.9	61.4	166.5
2004 4/.....	0.4	0.7	1.1	1.8	3.1	5.9	13.4	28.2	62.9	167.8
2003 4/.....	0.4	0.8	1.1	2.0	3.3	5.9	13.1	29.7	65.0	178.7
2002.....	0.5	0.7	1.0	1.8	3.3	6.1	13.1	30.9	68.1	189.5
2001.....	0.5	0.7	1.2	1.8	3.5	5.9	13.6	30.2	68.1	190.7
2000.....	0.5	0.7	1.1	1.8	2.9	6.4	13.4	30.1	65.7	191.0
1999.....	0.4	0.6	0.8	1.8	2.9	5.7	12.5	27.6	60.2	178.0
White, male										
2006 4/.....	0.6	1.0	1.6	2.4	4.9	8.3	17.0	36.4	87.5	231.8
2005 4/.....	0.5	0.9	1.7	2.4	4.5	7.8	17.1	37.1	86.1	234.2
2004 4/.....	0.5	1.0	1.4	2.2	3.9	7.8	17.9	38.3	88.4	241.7
2003 4/.....	0.4	1.0	1.4	2.5	4.2	8.2	17.2	40.6	91.6	250.8
2002.....	0.6	0.8	1.3	2.1	4.1	8.0	17.6	42.5	96.5	274.6
2001.....	0.7	0.8	1.5	2.3	4.0	8.0	18.4	40.6	98.6	275.3
2000.....	0.7	0.9	1.3	2.3	3.6	8.5	18.7	41.4	96.2	284.0
1999.....	0.5	0.8	0.9	2.2	3.9	7.7	17.1	40.4	88.9	263.6
White, female										
2006 4/.....	0.3	0.6	0.9	1.4	2.7	4.3	8.6	18.8	42.7	126.3
2005 4/.....	0.4	0.7	1.1	1.6	2.6	4.5	9.0	21.1	45.9	135.3
2004 4/.....	0.3	0.5	0.8	1.5	2.3	4.2	9.7	20.8	47.0	134.5
2003 4/.....	0.4	0.7	0.8	1.5	2.5	3.9	9.7	21.8	48.5	146.7
2002.....	0.4	0.5	0.7	1.5	2.6	4.4	9.5	22.5	50.7	152.5
2001.....	0.4	0.5	0.8	1.3	3.0	4.2	9.7	22.8	49.6	155.3
2000.....	0.4	0.5	0.9	1.4	2.3	4.6	9.1	22.2	47.5	153.2
1999.....	0.4	0.4	0.8	1.5	2.0	4.1	8.8	18.6	43.3	143.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pneumonitis due to food and vomit (J69.0)										
Black, both sexes										
2006 4/.....	0.9	1.2	1.7	2.9	4.9	9.2	19.8	32.8	55.1	128.4
2005 4/.....	*	1.2	2.4	3.1	6.4	10.4	17.0	40.9	69.2	153.1
2004 4/.....	0.9	1.0	1.7	3.0	7.3	8.8	20.4	34.7	64.6	144.7
2003 4/.....	0.9	1.3	1.2	2.7	8.5	11.0	17.2	39.8	76.2	163.1
2002.....	0.9	0.8	2.1	2.4	5.5	11.7	18.8	41.9	67.5	165.1
2001.....	0.7	1.0	2.4	2.4	6.0	9.3	22.3	38.6	76.1	185.5
2000.....	*	1.7	2.2	2.9	5.8	11.3	25.1	38.9	71.4	159.8
1999.....	1.0	1.2	1.8	3.9	5.0	9.9	23.7	40.3	71.1	167.1
Black, male										
2006 4/.....	*	*	2.4	3.6	7.6	15.0	28.0	55.0	84.9	173.4
2005 4/.....	*	1.5	3.4	3.6	9.8	15.4	25.6	62.1	106.0	201.7
2004 4/.....	*	*	2.2	4.4	11.0	13.6	31.4	51.4	92.5	202.7
2003 4/.....	*	1.7	*	3.8	11.6	14.5	26.3	59.3	107.8	212.7
2002.....	*	*	3.1	3.7	6.3	16.7	28.4	62.5	91.7	233.7
2001.....	*	*	3.4	3.1	8.1	12.5	31.0	56.2	108.3	243.9
2000.....	*	2.4	3.1	3.3	7.2	17.4	41.1	62.4	96.1	209.9
1999.....	*	*	*	5.1	8.1	15.3	37.3	65.1	109.9	228.5
Black, female										
2006 4/.....	*	*	*	2.3	2.8	4.8	14.2	19.6	39.4	110.0
2005 4/.....	*	*	*	2.7	3.7	6.7	11.0	28.3	49.7	133.5
2004 4/.....	*	*	*	*	4.4	5.2	13.0	24.8	49.9	121.7
2003 4/.....	*	*	*	*	6.0	8.4	11.1	28.2	59.6	143.8
2002.....	*	*	*	*	4.8	8.0	12.3	29.6	54.9	138.8
2001.....	*	*	*	*	4.3	6.9	16.6	28.0	59.7	163.8
2000.....	*	*	*	*	4.6	6.7	14.4	24.8	59.0	141.2
1999.....	*	*	*	2.9	*	6.0	14.6	25.5	51.9	144.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Suppurative and necrotic conditions of lower respiratory tract (J85-J86)

All races 3/, both sexes

2006.....	0.2	0.2	0.3	0.2	0.4	0.6	0.9	1.0	1.6	2.0
2005.....	0.1	0.2	0.3	0.4	0.4	0.6	0.6	1.1	1.9	2.2
2004.....	0.1	0.2	0.3	0.3	0.4	0.5	0.8	1.0	1.6	2.2
2003.....	0.1	0.3	0.3	0.4	0.5	0.6	0.9	1.1	1.7	2.1
2002.....	0.1	0.2	0.3	0.3	0.5	0.6	1.0	1.3	1.9	2.3
2001.....	0.1	0.2	0.3	0.4	0.5	0.7	0.9	1.5	1.8	3.0
2000.....	0.2	0.2	0.3	0.3	0.5	0.6	1.1	1.5	1.6	2.9
1999.....	0.2	0.2	0.3	0.4	0.5	1.0	1.0	1.3	1.7	2.8

All races 3/, male

2006.....	0.2	0.2	0.4	0.3	0.5	0.9	1.1	1.4	2.4	3.4
2005.....	0.2	0.3	0.4	0.5	0.5	1.0	1.0	1.6	2.7	3.2
2004.....	0.2	0.2	0.4	0.4	0.5	0.8	1.2	1.1	2.7	3.8
2003.....	0.2	0.3	0.4	0.5	0.6	0.9	1.3	1.2	2.2	3.3
2002.....	0.2	0.4	0.4	0.4	0.8	0.9	1.3	1.9	3.0	3.4
2001.....	0.2	0.2	0.5	0.5	0.7	0.9	1.3	2.2	2.5	4.7
2000.....	0.2	0.3	0.4	0.5	0.6	1.0	1.4	2.2	2.1	5.8
1999.....	0.3	0.3	0.5	0.5	0.7	1.6	1.5	1.7	2.8	4.0

All races 3/, female

2006.....	*	0.2	*	*	*	0.4	0.7	0.8	1.1	1.4
2005.....	*	*	*	0.3	*	*	*	0.8	1.4	1.7
2004.....	*	0.2	*	*	*	*	0.5	0.9	0.9	1.5
2003.....	*	0.2	*	0.2	0.4	*	0.7	1.0	1.3	1.5
2002.....	*	*	*	*	*	0.4	0.6	0.9	1.1	1.8
2001.....	*	*	*	0.3	*	0.5	0.7	1.1	1.4	2.3
2000.....	*	*	*	*	0.4	*	0.8	0.9	1.4	1.7
1999.....	*	*	*	*	*	0.4	0.6	1.1	1.0	2.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Suppurative and necrotic conditions of lower respiratory tract (J85-J86)										
White, both sexes										
2006 4/.....	0.1	0.2	0.3	0.2	0.4	0.7	0.8	1.1	1.5	2.1
2005 4/.....	0.1	0.2	0.2	0.4	0.4	0.7	0.6	1.1	2.0	2.2
2004 4/.....	0.1	0.2	0.2	0.3	0.4	0.5	0.8	0.9	1.5	2.2
2003 4/.....	0.1	0.2	0.3	0.3	0.4	0.6	0.9	1.0	1.5	2.1
2002.....	0.1	0.2	0.2	0.3	0.4	0.6	0.9	1.1	1.8	2.2
2001.....	0.1	0.2	0.3	0.4	0.5	0.7	0.9	1.5	1.8	3.0
2000.....	0.1	0.1	0.3	0.3	0.4	0.6	1.0	1.5	1.6	2.7
1999.....	0.2	0.2	0.3	0.3	0.5	0.9	1.0	1.3	1.7	2.5
White, male										
2006 4/.....	*	*	0.4	0.3	0.5	1.0	1.1	1.4	2.2	3.6
2005 4/.....	*	0.2	0.3	0.4	0.5	1.0	0.9	1.5	2.7	3.1
2004 4/.....	*	0.2	0.4	0.4	0.5	0.8	1.2	1.0	2.6	3.8
2003 4/.....	*	0.3	0.4	0.5	0.5	0.9	1.2	1.3	1.8	3.2
2002.....	*	0.3	0.4	0.4	0.7	0.9	1.3	1.6	3.0	3.2
2001.....	*	0.3	0.4	0.5	0.7	0.7	1.1	2.1	2.4	4.7
2000.....	*	*	0.4	0.4	0.5	0.8	1.4	2.2	2.0	5.5
1999.....	0.2	0.3	0.5	*	0.7	1.4	1.5	1.7	2.9	3.3
White, female										
2006 4/.....	*	*	*	*	*	*	0.6	0.8	1.0	1.4
2005 4/.....	*	*	*	0.3	*	*	*	0.8	1.5	1.8
2004 4/.....	*	0.3	*	*	*	*	*	0.8	0.9	1.4
2003 4/.....	*	*	*	*	*	*	0.7	0.9	1.4	1.6
2002.....	*	*	*	*	*	*	0.6	0.8	1.1	1.7
2001.....	*	*	*	*	*	0.6	0.7	1.1	1.4	2.3
2000.....	*	*	*	*	*	*	0.7	0.9	1.4	1.5
1999.....	*	*	*	*	*	*	0.6	1.1	0.9	2.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pleural effusion and plaque (J90,J92)

All races 3/, both sexes

2006.....	*	*	*	0.1	0.3	0.5	1.0	1.6	3.7	8.6
2005.....	*	*	*	0.1	0.2	0.3	0.9	1.6	4.2	9.3
2004.....	*	*	*	0.2	0.2	0.5	0.9	1.7	3.6	8.8
2003.....	*	*	*	0.1	0.2	0.5	0.9	1.6	3.5	8.8
2002.....	*	*	*	*	0.2	0.5	0.9	1.6	3.7	9.9
2001.....	*	*	*	*	0.3	0.4	0.9	1.6	3.8	10.0
2000.....	*	*	*	*	0.2	0.5	0.8	1.6	3.6	8.9
1999.....	*	*	*	0.2	0.2	0.5	0.8	1.7	3.4	9.0

All races 3/, male

2006.....	*	*	*	*	0.4	0.4	1.1	1.6	4.2	10.5
2005.....	*	*	*	*	*	*	1.1	1.7	5.2	10.8
2004.....	*	*	*	0.3	*	0.7	0.9	2.2	4.3	11.1
2003.....	*	*	*	*	*	0.5	0.9	1.8	4.3	11.3
2002.....	*	*	*	*	*	*	0.9	1.8	4.7	11.0
2001.....	*	*	*	*	*	0.5	1.2	1.6	4.8	12.2
2000.....	*	*	*	*	*	0.5	1.0	2.0	4.2	10.8
1999.....	*	*	*	*	*	0.6	1.1	2.2	4.2	11.5

All races 3/, female

2006.....	*	*	*	*	*	0.5	0.8	1.7	3.4	7.7
2005.....	*	*	*	*	*	*	0.8	1.5	3.5	8.6
2004.....	*	*	*	*	0.3	*	0.9	1.3	3.1	7.8
2003.....	*	*	*	*	*	0.4	0.9	1.5	3.0	7.7
2002.....	*	*	*	*	*	0.6	0.8	1.4	3.1	9.4
2001.....	*	*	*	*	*	*	0.7	1.7	3.3	9.1
2000.....	*	*	*	*	*	0.4	0.6	1.3	3.2	8.1
1999.....	*	*	*	*	*	*	0.7	1.3	2.9	8.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pleural effusion and plaque (J90,J92)										
White, both sexes										
2006 4/.....	*	*	*	*	0.3	0.5	0.9	1.7	3.6	9.0
2005 4/.....	*	*	*	*	0.2	0.3	0.9	1.6	4.3	9.6
2004 4/.....	*	*	*	0.2	0.3	0.5	0.9	1.7	3.6	9.0
2003 4/.....	*	*	*	*	0.2	0.5	0.9	1.5	3.6	9.2
2002.....	*	*	*	*	0.2	0.5	0.8	1.6	3.8	10.2
2001.....	*	*	*	*	0.3	0.4	0.9	1.6	3.9	10.0
2000.....	*	*	*	*	*	0.4	0.8	1.6	3.5	9.0
1999.....	*	*	*	*	0.3	0.5	0.8	1.7	3.5	9.0
White, male										
2006 4/.....	*	*	*	*	*	*	1.1	1.7	4.1	10.9
2005 4/.....	*	*	*	*	*	*	1.1	1.6	5.3	11.3
2004 4/.....	*	*	*	*	*	0.7	1.0	2.2	4.4	11.5
2003 4/.....	*	*	*	*	*	0.5	0.9	1.5	4.5	11.9
2002.....	*	*	*	*	*	*	0.9	1.8	4.8	11.6
2001.....	*	*	*	*	*	0.6	1.2	1.5	4.8	12.5
2000.....	*	*	*	*	*	*	0.9	2.0	4.2	10.9
1999.....	*	*	*	*	*	0.6	1.1	2.2	4.3	11.6
White, female										
2006 4/.....	*	*	*	*	*	0.5	0.8	1.7	3.3	8.2
2005 4/.....	*	*	*	*	*	*	0.8	1.6	3.6	8.8
2004 4/.....	*	*	*	*	*	*	0.9	1.3	3.1	7.8
2003 4/.....	*	*	*	*	*	*	1.0	1.5	3.1	7.9
2002.....	*	*	*	*	*	0.6	0.7	1.4	3.1	9.6
2001.....	*	*	*	*	*	*	0.7	1.7	3.3	8.9
2000.....	*	*	*	*	*	*	0.6	1.3	3.1	8.3
1999.....	*	*	*	*	*	*	0.6	1.3	3.0	7.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of respiratory system (J80-J84,J93-J98)										
All races 3/, both sexes										
2006.....	1.2	2.0	3.4	6.0	10.9	19.8	34.1	55.2	85.3	122.7
2005.....	1.2	2.0	3.5	5.7	11.0	20.2	35.3	55.8	81.9	121.0
2004.....	1.3	2.2	3.5	6.6	11.5	20.3	37.3	57.6	84.7	125.2
2003.....	1.1	2.2	3.5	5.8	10.4	19.5	33.7	53.6	77.0	123.3
2002.....	1.2	2.0	3.5	6.3	10.9	19.0	35.0	53.0	77.0	118.1
2001.....	1.3	2.2	3.8	6.6	11.5	21.2	35.6	57.6	82.1	125.7
2000.....	1.4	2.2	4.0	7.1	11.7	21.3	34.2	56.4	82.3	125.8
1999.....	1.3	2.2	3.5	6.1	11.5	20.5	34.2	55.1	80.6	131.6
All races 3/, male										
2006.....	1.2	2.0	3.5	7.0	12.8	24.2	43.3	71.1	109.7	145.7
2005.....	1.1	2.3	3.8	6.4	13.1	24.2	43.0	70.1	104.1	139.4
2004.....	1.2	2.2	3.9	7.1	13.2	24.7	46.6	72.7	109.1	149.7
2003.....	1.1	2.3	3.7	6.6	12.4	23.9	40.9	67.3	96.9	145.5
2002.....	1.3	2.0	3.6	7.0	12.5	22.7	44.7	65.9	95.2	138.9
2001.....	1.4	2.4	4.2	7.1	13.5	25.3	44.1	71.3	100.2	147.4
2000.....	1.4	2.3	4.3	8.0	13.5	24.6	42.3	71.9	103.5	158.3
1999.....	1.4	2.2	3.6	6.5	13.2	24.3	43.0	69.3	105.7	161.8
All races 3/, female										
2006.....	1.3	1.9	3.3	5.1	9.2	16.0	26.6	43.6	70.0	111.9
2005.....	1.3	1.8	3.3	5.2	9.1	16.7	29.0	45.4	68.1	112.6
2004.....	1.3	2.2	3.0	6.2	9.9	16.5	29.8	46.8	69.7	114.2
2003.....	1.1	2.1	3.3	5.1	8.6	15.8	28.0	43.8	64.9	113.5
2002.....	1.2	2.1	3.5	5.7	9.5	15.8	27.3	43.9	65.9	109.0
2001.....	1.2	2.0	3.5	6.2	9.6	17.7	28.9	48.0	71.2	116.6
2000.....	1.4	2.1	3.7	6.3	10.0	18.4	27.8	45.6	69.8	112.6
1999.....	1.3	2.2	3.4	5.8	10.0	17.3	27.4	45.3	65.9	119.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of respiratory system (J80-J84,J93-J98)										
White, both sexes										
2006 4/.....	1.1	1.7	3.1	5.7	10.7	20.2	35.5	57.2	88.6	126.8
2005 4/.....	1.1	1.8	3.2	5.5	10.8	20.3	36.2	57.1	84.7	124.2
2004 4/.....	1.1	1.8	3.2	6.4	11.3	20.6	38.7	59.8	87.5	129.8
2003 4/.....	0.9	1.9	3.2	5.6	10.5	20.0	34.9	55.1	80.6	128.0
2002.....	1.1	1.7	3.3	6.0	10.7	19.4	35.8	55.0	78.9	122.1
2001.....	1.1	1.9	3.4	6.3	11.3	21.4	36.5	59.3	84.7	129.0
2000.....	1.3	1.9	3.7	6.9	11.5	21.4	35.1	57.8	84.3	130.0
1999.....	1.1	1.9	3.2	5.9	11.5	20.8	34.5	56.2	82.4	134.9
White, male										
2006 4/.....	1.0	1.8	3.3	6.9	12.5	24.7	44.6	73.7	113.9	150.0
2005 4/.....	1.0	2.0	3.5	6.0	12.7	24.7	44.3	71.2	107.0	141.1
2004 4/.....	1.1	1.8	3.7	6.9	13.1	25.0	48.0	75.1	112.5	154.6
2003 4/.....	0.9	2.1	3.5	6.3	12.4	24.4	42.3	68.6	101.3	150.4
2002.....	1.2	1.7	3.3	6.8	12.2	23.6	45.3	68.1	97.9	143.8
2001.....	1.4	2.1	3.8	7.0	13.1	25.8	45.5	73.7	103.5	151.5
2000.....	1.3	2.0	3.9	7.8	13.4	25.0	43.1	73.5	106.2	163.4
1999.....	1.2	1.9	3.3	6.1	13.3	24.9	43.5	71.0	108.0	166.3
White, female										
2006 4/.....	1.2	1.6	2.9	4.5	9.1	16.1	28.0	45.0	72.6	116.0
2005 4/.....	1.1	1.6	2.9	5.0	9.0	16.3	29.5	46.6	70.8	116.4
2004 4/.....	1.2	1.8	2.7	6.0	9.6	16.6	31.0	48.6	71.9	118.7
2003 4/.....	0.9	1.7	3.0	4.9	8.7	16.2	28.8	45.3	67.8	118.0
2002.....	1.1	1.8	3.3	5.3	9.3	15.7	28.0	45.6	67.3	112.7
2001.....	0.9	1.7	3.0	5.7	9.6	17.6	29.2	49.1	73.3	119.6
2000.....	1.3	1.9	3.5	6.1	9.8	18.2	28.7	46.8	71.3	116.4
1999.....	1.0	1.9	3.1	5.6	9.9	17.1	27.3	45.9	67.4	122.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of respiratory system (J80-J84,J93-J98)										
Black, both sexes										
2006 4/.....	2.2	4.0	5.9	9.9	14.3	18.2	28.4	41.2	54.0	84.8
2005 4/.....	2.2	4.1	6.4	8.7	14.3	22.4	31.5	48.9	54.4	93.3
2004 4/.....	2.6	5.1	6.1	9.9	15.4	21.3	30.1	41.5	57.4	79.0
2003 4/.....	2.6	4.7	6.2	9.2	12.3	18.5	26.5	44.9	43.7	77.3
2002.....	2.4	4.5	6.1	9.7	15.1	18.8	30.3	35.2	59.9	76.9
2001.....	2.6	5.0	7.9	10.5	15.2	22.2	32.0	44.6	59.0	90.0
2000.....	2.4	5.2	7.2	10.5	14.2	24.5	29.2	45.3	64.1	82.3
1999.....	3.0	5.0	6.4	9.1	13.3	22.0	34.8	48.2	58.9	94.1
Black, male										
2006 4/.....	2.2	3.8	5.4	9.3	17.5	21.8	40.3	53.3	70.5	101.9
2005 4/.....	2.1	4.4	6.5	10.0	18.5	25.0	36.1	64.4	73.7	119.0
2004 4/.....	2.7	5.7	6.2	11.1	17.0	26.9	38.5	55.5	71.0	100.3
2003 4/.....	2.1	4.2	5.9	10.9	16.0	22.9	33.8	61.6	56.1	98.0
2002.....	2.8	4.2	6.3	10.4	18.6	20.2	40.4	45.0	73.4	86.8
2001.....	2.3	5.3	7.7	9.9	18.9	25.2	35.3	51.1	71.4	106.7
2000.....	2.5	5.9	8.0	11.7	15.7	27.1	35.4	61.4	74.2	104.9
1999.....	2.8	5.4	6.8	9.7	14.9	23.6	41.0	59.4	78.5	114.8
Black, female										
2006 4/.....	2.2	4.2	6.4	10.4	11.8	15.6	20.3	34.0	45.2	77.9
2005 4/.....	2.4	3.8	6.2	7.6	10.9	20.4	28.4	39.7	44.1	82.9
2004 4/.....	2.5	4.6	6.1	9.0	14.1	17.1	24.4	33.2	50.3	70.5
2003 4/.....	3.1	5.3	6.5	7.7	9.3	15.2	21.5	35.0	37.1	69.3
2002.....	2.0	4.8	5.9	9.2	12.4	17.8	23.5	29.4	52.9	73.1
2001.....	3.0	4.6	8.0	11.0	12.3	20.1	29.8	40.8	52.7	83.8
2000.....	2.3	4.5	6.4	9.6	13.1	22.7	25.1	35.6	59.0	73.8
1999.....	3.3	4.6	6.1	8.7	12.1	20.8	30.6	41.5	49.3	86.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XI. Diseases of the digestive system (K00-K92)

All races 3/, both sexes

2006.....	13.6	23.8	35.8	42.0	50.9	63.6	89.9	128.9	194.5	363.3
2005.....	13.9	24.1	35.1	41.7	50.8	65.7	92.4	130.7	199.6	376.8
2004.....	14.4	24.6	35.1	40.1	50.3	65.7	93.3	134.1	200.3	387.5
2003.....	15.4	25.5	35.5	40.5	51.9	70.5	96.1	137.2	208.3	403.8
2002.....	15.9	26.3	33.8	40.2	50.5	68.6	96.0	139.2	212.9	415.3
2001.....	15.2	26.8	33.0	39.0	53.2	71.4	97.3	138.2	212.8	431.4
2000.....	15.9	25.8	31.7	39.8	52.9	68.9	96.3	140.7	214.9	440.7
1999.....	15.8	25.3	31.0	39.6	53.5	72.3	99.0	138.5	214.9	436.4

All races 3/, male

2006.....	17.2	31.9	50.2	57.7	65.8	77.8	103.1	143.4	209.3	347.1
2005.....	17.9	32.9	50.1	57.8	66.4	78.5	106.6	146.4	214.4	364.8
2004.....	18.8	33.4	50.3	54.7	63.8	79.2	106.7	151.3	212.4	366.9
2003.....	19.9	35.4	50.9	54.8	68.1	85.5	111.9	151.8	226.6	388.9
2002.....	20.7	37.0	48.7	54.2	65.3	83.9	111.8	155.6	227.7	404.4
2001.....	20.1	37.5	46.8	53.1	68.5	86.9	113.9	153.0	227.3	428.2
2000.....	21.5	37.2	44.7	53.3	68.7	84.7	112.4	160.6	229.3	441.9
1999.....	22.0	37.0	44.4	54.2	68.6	89.0	117.3	154.4	238.2	438.5

All races 3/, female

2006.....	10.2	15.8	21.9	27.2	37.2	51.1	79.2	118.3	185.2	370.8
2005.....	10.0	15.5	20.6	26.5	36.6	54.5	80.8	119.3	190.4	382.3
2004.....	10.1	16.1	20.5	26.2	38.0	53.9	82.5	121.7	192.8	396.8
2003.....	10.9	15.9	20.7	27.0	37.1	57.4	83.5	126.8	197.1	410.4
2002.....	11.2	16.0	19.5	27.1	37.0	55.5	83.4	127.6	203.9	420.0
2001.....	10.4	16.4	19.8	25.9	39.3	58.1	84.1	127.8	204.1	432.7
2000.....	10.4	14.9	19.3	27.1	38.7	55.5	83.7	126.8	206.5	440.2
1999.....	9.7	14.0	18.2	25.9	39.8	57.9	84.7	127.4	201.3	435.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XI. Diseases of the digestive system (K00-K92)

White, both sexes

2006 4/.....	13.9	24.1	35.6	41.5	50.9	63.7	91.0	130.5	198.3	372.9
2005 4/.....	14.1	23.9	34.4	41.5	50.0	65.5	93.4	131.6	201.8	385.4
2004 4/.....	14.3	24.4	34.6	38.9	49.4	65.9	93.8	136.1	202.7	396.3
2003 4/.....	15.3	25.2	34.2	39.3	50.6	70.4	97.0	139.0	211.9	412.6
2002.....	15.9	25.7	32.8	39.3	49.5	69.0	96.4	141.0	215.3	426.3
2001.....	14.7	26.0	31.4	37.0	52.3	71.1	97.7	138.3	216.0	441.6
2000.....	15.4	24.9	29.9	38.2	51.5	68.0	96.5	141.2	217.1	449.3
1999.....	15.0	23.8	29.7	37.8	52.0	71.2	98.7	138.4	217.3	445.2

White, male

2006 4/.....	17.5	32.5	50.0	56.8	65.5	77.7	103.6	143.2	213.0	356.2
2005 4/.....	18.2	32.5	49.3	57.3	65.0	77.4	107.1	145.9	216.3	372.5
2004 4/.....	19.1	33.0	49.9	52.8	62.3	78.4	106.4	152.1	213.2	375.1
2003 4/.....	20.0	35.4	49.1	52.8	66.4	84.9	111.5	152.8	228.0	395.9
2002.....	21.0	36.4	47.4	52.9	63.7	83.9	111.4	157.0	228.7	412.5
2001.....	19.4	36.7	44.6	50.1	67.0	85.5	113.6	151.9	230.6	434.7
2000.....	20.9	36.5	42.1	51.4	66.8	83.5	111.8	160.1	230.7	448.1
1999.....	21.1	35.0	42.5	51.8	67.1	87.8	117.1	154.3	240.5	443.1

White, female

2006 4/.....	10.2	15.8	21.4	26.7	37.4	51.3	80.5	121.0	189.1	380.8
2005 4/.....	9.9	15.2	19.7	26.2	36.0	54.8	82.1	121.0	192.7	391.3
2004 4/.....	9.5	15.7	19.6	25.6	37.4	54.7	83.5	124.4	196.2	405.9
2003 4/.....	10.6	15.1	19.7	26.5	36.0	57.5	85.1	129.1	202.0	420.0
2002.....	10.9	15.1	18.5	26.4	36.4	55.9	84.2	129.6	207.2	432.3
2001.....	10.0	15.5	18.5	24.5	38.7	58.4	84.9	128.7	207.2	444.5
2000.....	9.8	13.4	18.0	25.6	37.4	54.4	84.3	127.9	208.9	449.8
1999.....	8.9	12.7	17.2	24.6	38.1	56.7	84.1	127.3	203.5	446.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XI. Diseases of the digestive system (K00-K92)										
Black, both sexes										
2006 4/.....	13.6	24.2	44.7	55.0	58.7	72.1	92.6	130.7	171.1	306.4
2005 4/.....	13.8	29.3	45.4	53.3	65.2	77.5	98.2	137.1	198.8	320.7
2004 4/.....	16.4	30.8	46.1	56.7	66.8	75.3	105.2	130.5	199.5	329.7
2003 4/.....	17.3	30.9	51.6	58.1	70.0	82.9	102.7	137.5	187.3	346.0
2002.....	17.6	35.1	48.3	54.2	65.2	76.8	105.5	137.1	208.0	333.2
2001.....	20.1	36.6	51.1	62.2	68.6	87.7	108.1	151.8	201.0	351.0
2000.....	21.3	37.2	51.0	58.7	72.2	86.3	109.0	147.5	207.0	382.0
1999.....	22.6	40.1	46.6	59.1	74.3	90.3	112.7	150.0	207.5	364.4
Black, male										
2006 4/.....	17.0	31.9	62.5	78.2	80.2	95.8	113.2	168.3	191.0	287.7
2005 4/.....	17.4	40.8	64.4	75.2	90.0	104.8	123.7	171.2	217.0	304.0
2004 4/.....	19.4	42.6	65.0	83.4	91.8	102.2	134.3	162.9	229.6	322.9
2003 4/.....	21.3	40.8	76.5	85.4	96.4	107.4	136.0	162.5	237.7	345.1
2002.....	20.9	47.7	69.4	76.8	89.3	99.5	133.2	158.0	241.5	355.9
2001.....	26.6	51.3	75.3	90.1	91.9	118.7	138.0	184.0	217.5	386.9
2000.....	28.0	49.6	73.8	81.0	97.3	109.0	138.7	181.4	241.1	392.1
1999.....	30.0	59.2	68.5	84.6	95.0	112.4	135.0	171.0	239.8	400.7
Black, female										
2006 4/.....	10.5	17.4	29.7	35.8	42.0	54.5	78.4	108.2	160.6	314.0
2005 4/.....	10.7	19.2	29.3	35.2	45.9	57.3	80.7	116.8	189.2	327.5
2004 4/.....	13.7	20.5	30.1	34.6	47.3	55.2	85.4	111.3	183.7	332.3
2003 4/.....	13.8	22.3	30.5	35.5	49.3	64.7	80.2	122.6	160.9	346.4
2002.....	14.6	24.1	30.4	35.6	46.3	59.9	86.9	124.6	190.6	324.5
2001.....	14.3	23.9	30.5	39.4	50.2	64.8	88.1	132.6	192.6	337.6
2000.....	15.4	26.3	31.5	40.5	52.5	69.5	89.2	127.0	189.8	378.2
1999.....	16.1	23.6	28.0	38.2	58.1	73.9	97.9	137.3	191.4	350.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of other parts of digestive system (K20-K92)										
All races 3/, both sexes										
2006.....	13.6	23.8	35.7	41.9	50.8	63.5	89.7	128.7	193.9	361.6
2005.....	13.9	24.0	35.0	41.7	50.7	65.5	92.1	130.3	199.0	375.2
2004.....	14.4	24.6	35.1	40.0	50.2	65.5	93.1	133.8	199.6	385.7
2003.....	15.3	25.5	35.5	40.5	51.8	70.4	96.0	137.0	208.0	402.6
2002.....	15.9	26.3	33.8	40.2	50.4	68.5	95.9	139.0	212.5	413.8
2001.....	15.2	26.8	33.0	39.0	53.1	71.4	97.2	137.9	212.2	430.2
2000.....	15.9	25.8	31.7	39.8	52.9	68.9	96.2	140.6	214.5	439.5
1999.....	15.8	25.3	31.0	39.6	53.5	72.2	98.9	138.3	214.6	435.4
All races 3/, male										
2006.....	17.1	31.9	50.1	57.6	65.8	77.7	102.9	143.1	208.8	346.0
2005.....	17.9	32.9	50.0	57.7	66.3	78.4	106.2	145.8	213.6	363.2
2004.....	18.8	33.4	50.3	54.7	63.7	79.1	106.6	150.9	211.7	365.5
2003.....	19.8	35.4	50.9	54.8	68.0	85.5	111.8	151.6	226.3	388.0
2002.....	20.7	37.0	48.7	54.1	65.2	83.8	111.6	155.3	227.4	403.1
2001.....	20.1	37.5	46.8	53.0	68.4	86.9	113.8	152.8	226.7	427.0
2000.....	21.4	37.1	44.7	53.3	68.6	84.6	112.2	160.4	229.0	440.8
1999.....	21.9	37.0	44.4	54.2	68.6	88.9	117.1	154.2	238.0	437.4
All races 3/, female										
2006.....	10.2	15.8	21.9	27.2	37.1	51.1	79.0	118.1	184.5	368.9
2005.....	10.0	15.4	20.6	26.5	36.5	54.3	80.7	119.0	189.8	380.7
2004.....	10.1	16.0	20.5	26.2	37.9	53.8	82.3	121.5	192.2	394.7
2003.....	10.9	15.9	20.7	26.9	37.0	57.3	83.4	126.7	196.7	409.1
2002.....	11.2	16.0	19.5	27.0	37.0	55.4	83.3	127.5	203.4	418.4
2001.....	10.4	16.4	19.8	25.9	39.3	58.1	84.1	127.5	203.6	431.5
2000.....	10.4	14.9	19.2	27.1	38.7	55.4	83.6	126.7	206.0	439.0
1999.....	9.7	13.9	18.2	25.9	39.7	57.8	84.6	127.1	200.9	434.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of other parts of digestive system (K20-K92)										
White, both sexes										
2006 4/.....	13.9	24.1	35.5	41.4	50.9	63.6	90.9	130.3	197.7	371.1
2005 4/.....	14.1	23.8	34.3	41.4	49.9	65.3	93.1	131.2	201.2	383.6
2004 4/.....	14.3	24.3	34.6	38.8	49.2	65.7	93.7	135.8	202.0	394.3
2003 4/.....	15.3	25.2	34.2	39.3	50.6	70.3	96.9	138.8	211.5	411.3
2002.....	15.9	25.7	32.8	39.3	49.5	68.9	96.3	140.8	214.9	424.8
2001.....	14.7	26.0	31.4	37.0	52.2	71.0	97.7	138.1	215.4	440.4
2000.....	15.3	24.9	29.9	38.2	51.5	67.9	96.4	141.1	216.7	448.1
1999.....	15.0	23.7	29.7	37.8	52.0	71.1	98.6	138.2	217.0	444.1
White, male										
2006 4/.....	17.5	32.4	49.9	56.7	65.5	77.5	103.4	143.0	212.5	355.0
2005 4/.....	18.2	32.5	49.2	57.2	64.9	77.2	106.7	145.4	215.7	370.8
2004 4/.....	19.1	33.0	49.9	52.7	62.2	78.3	106.3	151.6	212.6	373.5
2003 4/.....	20.0	35.4	49.1	52.7	66.4	84.9	111.5	152.6	227.7	394.9
2002.....	21.0	36.3	47.4	52.8	63.6	83.8	111.2	156.7	228.3	411.2
2001.....	19.4	36.6	44.6	50.0	67.0	85.5	113.6	151.7	230.0	433.4
2000.....	20.9	36.4	42.0	51.4	66.7	83.4	111.6	160.0	230.4	446.9
1999.....	21.1	34.9	42.5	51.7	67.1	87.7	116.9	154.1	240.4	441.9
White, female										
2006 4/.....	10.2	15.8	21.4	26.7	37.3	51.3	80.4	120.8	188.4	378.7
2005 4/.....	9.9	15.2	19.7	26.2	35.9	54.7	81.9	120.7	192.2	389.5
2004 4/.....	9.5	15.7	19.6	25.5	37.3	54.6	83.3	124.2	195.5	403.7
2003 4/.....	10.5	15.1	19.7	26.5	36.0	57.4	85.0	128.9	201.6	418.6
2002.....	10.9	15.1	18.5	26.4	36.4	55.8	84.1	129.5	206.7	430.6
2001.....	10.0	15.5	18.5	24.5	38.7	58.3	84.8	128.5	206.6	443.3
2000.....	9.8	13.4	18.0	25.6	37.4	54.4	84.2	127.8	208.5	448.6
1999.....	8.8	12.7	17.2	24.6	38.0	56.6	84.1	127.1	203.2	445.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of other parts of digestive system (K20-K92)										
Black, both sexes										
2006 4/.....	13.5	24.2	44.6	55.0	58.7	72.0	92.2	130.5	170.6	305.4
2005 4/.....	13.7	29.2	45.1	53.2	65.1	77.4	98.2	136.8	198.1	320.5
2004 4/.....	16.4	30.7	46.0	56.7	66.7	75.0	105.0	130.3	199.3	329.4
2003 4/.....	17.3	30.9	51.6	58.0	69.8	82.9	102.4	137.5	187.3	346.0
2002.....	17.6	35.0	48.3	54.1	65.2	76.7	105.5	136.9	207.7	332.3
2001.....	20.1	36.6	51.1	62.2	68.6	87.7	108.1	150.9	200.2	350.1
2000.....	21.3	37.2	51.0	58.7	72.2	86.2	108.9	147.5	206.1	381.6
1999.....	22.5	40.0	46.6	59.1	74.2	90.3	112.6	149.6	206.9	363.8
Black, male										
2006 4/.....	17.0	31.9	62.3	78.2	80.2	95.6	112.9	167.8	191.0	286.0
2005 4/.....	17.2	40.7	64.2	75.1	90.0	104.8	123.7	170.8	215.6	304.0
2004 4/.....	19.4	42.5	64.8	83.4	91.8	102.0	134.0	162.4	229.6	322.9
2003 4/.....	21.2	40.8	76.5	85.3	96.4	107.4	135.7	162.5	237.7	345.1
2002.....	20.9	47.5	69.3	76.8	89.3	99.2	133.2	158.0	240.7	355.9
2001.....	26.6	51.3	75.3	90.1	91.9	118.7	138.0	183.1	215.9	385.8
2000.....	28.0	49.6	73.8	81.0	97.3	109.0	138.4	181.4	241.1	392.1
1999.....	29.9	59.2	68.5	84.6	94.8	112.4	135.0	171.0	238.9	400.7
Black, female										
2006 4/.....	10.4	17.4	29.7	35.8	42.0	54.5	78.0	108.2	159.9	313.3
2005 4/.....	10.7	19.2	28.9	35.2	45.8	57.1	80.7	116.5	188.8	327.1
2004 4/.....	13.7	20.5	30.0	34.6	47.2	54.9	85.2	111.3	183.3	332.0
2003 4/.....	13.8	22.3	30.5	35.5	49.0	64.7	80.0	122.6	160.9	346.4
2002.....	14.6	24.1	30.4	35.4	46.3	59.9	86.9	124.3	190.6	323.2
2001.....	14.3	23.9	30.5	39.4	50.2	64.8	88.1	131.7	192.1	336.8
2000.....	15.4	26.3	31.5	40.5	52.5	69.3	89.2	127.0	188.5	377.7
1999.....	16.1	23.5	28.0	38.2	58.1	73.9	97.7	136.7	191.0	350.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of esophagus (K20-K22)										
All races 3/, both sexes										
2006.....	0.1	0.2	0.3	0.4	0.6	0.9	1.4	2.3	5.0	15.8
2005.....	0.1	0.2	0.3	0.3	0.6	0.7	1.3	2.9	6.8	17.2
2004.....	0.1	0.2	0.2	0.3	0.6	1.0	1.5	2.8	5.8	17.9
2003.....	0.1	0.1	0.3	0.4	0.6	0.9	1.6	3.1	6.1	19.3
2002.....	0.2	0.3	0.3	0.3	0.5	0.9	1.7	3.5	6.8	20.7
2001.....	*	0.1	0.2	0.4	0.6	1.0	1.8	3.5	7.4	21.2
2000.....	0.1	0.2	0.3	0.4	0.6	1.1	1.8	3.6	7.7	23.3
1999.....	0.1	0.2	0.3	0.4	0.6	1.1	1.9	3.3	7.7	23.5
All races 3/, male										
2006.....	*	0.2	0.4	0.6	0.8	1.1	1.7	2.9	6.3	18.1
2005.....	*	0.3	0.4	0.4	0.7	1.0	1.3	3.9	8.5	20.5
2004.....	0.2	0.2	0.4	0.4	0.8	1.1	1.6	3.8	6.7	19.1
2003.....	*	0.2	0.4	0.5	0.8	1.3	2.1	3.6	7.0	23.6
2002.....	0.2	0.4	0.4	0.3	0.6	1.4	1.9	4.2	8.3	25.4
2001.....	*	*	0.3	0.5	0.9	1.3	2.1	4.1	8.8	23.6
2000.....	0.2	0.3	0.4	0.5	1.0	1.5	2.1	4.6	9.3	28.6
1999.....	*	0.3	0.3	0.4	0.8	1.5	2.7	4.4	9.9	28.3
All races 3/, female										
2006.....	*	0.2	*	*	0.4	0.8	1.1	1.8	4.2	14.7
2005.....	*	*	*	0.3	0.5	0.5	1.3	2.1	5.7	15.7
2004.....	*	*	*	*	0.4	0.8	1.4	2.1	5.3	17.3
2003.....	*	*	0.2	0.2	0.3	0.7	1.2	2.8	5.7	17.4
2002.....	*	*	*	0.3	0.4	0.6	1.6	3.0	5.9	18.7
2001.....	*	*	*	*	0.4	0.8	1.5	3.0	6.6	20.2
2000.....	*	*	*	*	*	0.7	1.6	2.8	6.8	21.1
1999.....	*	*	0.2	0.4	0.4	0.8	1.3	2.5	6.4	21.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of esophagus (K20-K22)										
White, both sexes										
2006 4/.....	0.1	0.2	0.3	0.4	0.6	0.9	1.4	2.5	5.2	16.8
2005 4/.....	0.1	0.2	0.3	0.3	0.6	0.6	1.2	3.0	7.0	18.1
2004 4/.....	0.1	0.1	0.3	0.3	0.5	0.9	1.5	2.8	6.1	18.8
2003 4/.....	0.1	0.2	0.3	0.4	0.6	1.0	1.7	3.2	6.4	20.2
2002.....	0.2	0.3	0.3	0.3	0.5	0.9	1.7	3.4	7.0	21.7
2001.....	*	0.1	0.2	0.4	0.6	1.0	1.7	3.6	7.6	22.2
2000.....	0.1	0.2	0.3	0.4	0.6	1.1	1.8	3.7	8.1	24.4
1999.....	*	0.2	0.2	0.4	0.6	1.1	1.9	3.4	7.9	24.5
White, male										
2006 4/.....	*	0.3	0.4	0.6	0.8	1.0	1.7	3.1	6.8	19.1
2005 4/.....	*	0.3	0.4	0.4	0.7	0.8	1.3	4.0	8.9	21.4
2004 4/.....	*	*	0.4	0.4	0.7	1.0	1.6	3.8	7.0	20.1
2003 4/.....	*	0.2	0.3	0.5	0.7	1.2	2.3	3.7	7.3	24.9
2002.....	0.3	0.4	0.4	*	0.6	1.2	2.0	4.1	8.5	26.7
2001.....	*	*	0.3	0.5	0.8	1.2	2.0	4.0	9.0	24.7
2000.....	*	0.3	0.4	0.5	0.9	1.5	2.0	4.6	9.6	30.0
1999.....	*	*	*	0.5	0.7	1.5	2.6	4.5	10.0	29.8
White, female										
2006 4/.....	*	*	*	*	0.4	0.9	1.1	2.0	4.3	15.7
2005 4/.....	*	*	*	0.3	0.5	0.5	1.2	2.2	5.8	16.6
2004 4/.....	*	*	*	*	0.4	0.9	1.4	2.2	5.5	18.2
2003 4/.....	*	*	*	*	0.4	0.7	1.2	2.8	5.9	18.1
2002.....	*	*	*	*	0.4	0.6	1.5	3.0	6.1	19.5
2001.....	*	*	*	*	0.4	0.9	1.5	3.2	6.8	21.2
2000.....	*	*	*	*	*	0.7	1.6	3.0	7.2	22.1
1999.....	*	*	*	0.4	0.5	0.8	1.4	2.6	6.7	22.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
All races 3/, both sexes										
2006.....	0.2	0.5	0.8	1.0	1.5	2.1	3.5	5.3	9.6	20.7
2005.....	0.3	0.5	0.8	0.9	1.6	2.2	3.8	5.8	10.8	21.4
2004.....	0.3	0.5	0.8	1.0	1.7	2.5	3.9	6.8	11.6	24.4
2003.....	0.3	0.6	0.8	1.2	1.7	2.3	4.2	7.3	13.6	26.4
2002.....	0.3	0.6	0.8	1.4	1.7	2.7	4.7	8.0	13.2	28.3
2001.....	0.4	0.7	0.8	1.3	2.1	3.1	4.8	8.3	16.0	33.4
2000.....	0.3	0.6	1.0	1.5	2.2	3.1	5.1	8.7	15.6	35.6
1999.....	0.4	0.6	0.8	1.4	2.2	3.3	5.9	9.1	17.1	34.6
All races 3/, male										
2006.....	*	0.8	1.0	1.4	1.9	2.6	4.0	6.3	11.0	22.0
2005.....	0.5	0.8	1.2	1.3	2.2	2.6	4.8	7.3	13.1	22.4
2004.....	0.3	0.7	1.0	1.2	2.2	2.9	4.5	8.3	12.9	24.6
2003.....	0.4	0.8	1.2	1.4	2.3	3.1	5.4	8.8	16.2	28.9
2002.....	0.4	0.7	1.2	1.8	2.1	3.2	6.1	9.3	16.7	31.6
2001.....	0.5	0.9	1.1	1.8	2.8	4.1	5.6	10.3	17.3	34.9
2000.....	0.4	0.9	1.5	2.1	2.9	3.9	6.9	11.2	17.8	40.7
1999.....	0.4	0.7	1.2	1.8	3.0	4.5	7.9	11.0	21.1	37.5
All races 3/, female										
2006.....	0.2	0.3	0.5	0.6	1.1	1.7	3.1	4.6	8.8	20.1
2005.....	0.2	0.3	0.5	0.6	1.1	1.8	3.0	4.7	9.4	21.0
2004.....	0.2	0.4	0.5	0.7	1.3	2.2	3.4	5.8	10.8	24.3
2003.....	0.2	0.4	0.5	1.0	1.1	1.7	3.2	6.2	11.9	25.3
2002.....	0.3	0.4	0.5	1.0	1.4	2.2	3.6	7.0	11.0	26.9
2001.....	0.2	0.4	0.6	0.8	1.5	2.2	4.2	6.9	15.2	32.7
2000.....	0.2	0.4	0.5	0.9	1.6	2.5	3.6	6.9	14.4	33.5
1999.....	0.3	0.5	0.5	1.1	1.5	2.3	4.3	7.7	14.7	33.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
White, both sexes										
2006 4/.....	0.2	0.5	0.7	0.9	1.4	2.0	3.4	5.2	9.9	21.2
2005 4/.....	0.3	0.4	0.8	0.9	1.6	2.2	3.7	5.7	10.9	21.6
2004 4/.....	0.2	0.4	0.7	0.9	1.7	2.5	3.8	6.8	11.7	25.1
2003 4/.....	0.3	0.5	0.7	1.1	1.6	2.4	4.2	7.2	13.4	26.9
2002.....	0.2	0.5	0.8	1.3	1.6	2.6	4.8	7.9	13.6	29.1
2001.....	0.3	0.6	0.7	1.1	2.0	3.0	4.8	8.3	16.2	34.4
2000.....	0.3	0.6	0.9	1.4	2.2	3.0	5.1	8.5	16.0	36.3
1999.....	0.3	0.5	0.7	1.4	2.2	3.1	5.8	9.1	17.1	35.3
White, male										
2006 4/.....	*	0.7	0.9	1.4	1.8	2.5	3.6	6.0	11.3	22.0
2005 4/.....	0.4	0.6	1.1	1.2	2.1	2.5	4.5	7.0	13.5	22.5
2004 4/.....	0.3	0.6	1.0	1.2	2.1	2.9	4.3	8.1	12.6	25.5
2003 4/.....	0.3	0.7	1.0	1.3	2.2	3.1	5.3	8.6	15.6	29.0
2002.....	0.3	0.6	1.0	1.6	2.0	3.1	6.0	9.2	17.1	32.5
2001.....	0.5	0.8	0.9	1.5	2.7	3.8	5.3	10.1	17.3	35.7
2000.....	0.4	0.8	1.3	2.0	2.9	3.8	6.7	10.9	17.8	40.7
1999.....	0.3	0.6	1.1	1.6	2.9	4.1	7.8	11.1	21.1	38.4
White, female										
2006 4/.....	*	0.3	0.5	0.5	1.1	1.6	3.2	4.6	9.0	20.8
2005 4/.....	*	0.2	0.5	0.7	1.1	1.9	3.0	4.7	9.3	21.2
2004 4/.....	*	0.3	0.5	0.7	1.3	2.1	3.3	5.8	11.1	25.0
2003 4/.....	0.2	0.3	0.5	0.9	1.0	1.7	3.3	6.2	12.1	26.0
2002.....	*	0.4	0.5	1.0	1.3	2.1	3.8	7.0	11.5	27.6
2001.....	*	0.4	0.6	0.8	1.4	2.2	4.4	7.0	15.6	33.9
2000.....	*	0.3	0.5	0.9	1.6	2.4	3.8	6.9	14.9	34.5
1999.....	0.3	0.4	0.4	1.1	1.5	2.2	4.2	7.7	14.8	34.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer (K25-K28)										
Black, both sexes										
2006 4/.....	*	1.0	1.6	1.7	2.6	2.9	4.8	7.0	7.3	14.8
2005 4/.....	0.7	1.1	1.4	1.2	2.2	2.7	5.4	5.8	9.7	17.4
2004 4/.....	*	1.3	1.3	1.6	2.6	3.5	5.1	6.6	9.4	16.1
2003 4/.....	*	1.2	1.8	2.1	2.4	2.1	4.5	8.7	14.5	16.9
2002.....	0.9	0.9	1.6	2.3	2.9	3.2	5.2	8.8	6.8	19.0
2001.....	*	1.1	1.6	2.5	3.0	3.6	5.5	7.8	11.1	20.9
2000.....	*	1.4	2.0	2.6	2.7	4.1	5.7	9.4	11.0	26.6
1999.....	1.0	1.2	1.8	2.4	3.2	5.0	6.5	7.7	16.2	25.5
Black, male										
2006 4/.....	*	1.8	2.8	2.5	*	*	8.3	9.8	*	*
2005 4/.....	*	1.9	2.5	*	*	*	7.9	9.4	*	20.5
2004 4/.....	*	2.1	*	*	3.9	*	*	10.0	*	*
2003 4/.....	*	1.6	3.1	3.1	3.9	*	7.2	11.4	23.6	*
2002.....	*	*	2.8	3.6	4.2	*	8.0	11.9	*	*
2001.....	*	1.8	2.7	4.1	4.5	5.3	9.1	10.7	*	23.8
2000.....	*	*	3.3	3.5	*	*	10.0	12.8	*	31.1
1999.....	1.5	1.9	2.8	3.7	5.3	7.5	8.8	*	18.3	28.1
Black, female										
2006 4/.....	*	*	*	*	*	*	*	5.4	*	13.9
2005 4/.....	*	*	*	*	*	*	*	*	10.4	16.1
2004 4/.....	*	*	*	*	*	3.6	4.4	*	7.6	16.5
2003 4/.....	*	*	*	*	*	*	*	7.1	9.7	15.8
2002.....	*	*	*	*	*	*	*	6.9	*	18.5
2001.....	*	*	*	*	*	*	*	6.1	9.0	19.8
2000.....	*	*	*	*	*	*	*	7.4	8.5	24.9
1999.....	*	*	*	*	*	*	5.0	6.9	15.1	24.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Gastric ulcer (K25)										
All races 3/, both sexes										
2006.....	*	0.2	0.3	0.3	0.5	0.6	1.2	1.6	2.6	5.9
2005.....	0.1	0.2	0.3	0.3	0.5	0.7	0.9	1.6	3.1	6.1
2004.....	0.1	0.2	0.3	0.3	0.6	0.8	1.1	2.1	3.4	6.7
2003.....	0.1	0.2	0.3	0.5	0.6	0.7	1.1	2.1	3.8	7.0
2002.....	0.1	0.2	0.3	0.5	0.5	0.8	1.4	2.1	3.8	7.7
2001.....	0.1	0.3	0.3	0.4	0.6	1.0	1.5	2.4	4.6	8.0
2000.....	0.1	0.3	0.4	0.5	0.7	1.0	1.5	2.4	4.0	9.5
1999.....	0.2	0.2	0.3	0.5	0.6	0.9	1.8	2.7	4.8	8.8
All races 3/, male										
2006.....	*	0.2	0.4	0.4	0.6	0.7	1.3	1.5	2.8	5.8
2005.....	0.2	0.3	0.5	0.4	0.6	0.6	1.0	1.9	3.5	5.4
2004.....	*	0.3	0.4	0.4	0.7	0.8	1.2	2.1	4.0	6.3
2003.....	*	0.3	0.5	0.5	0.7	0.9	1.1	2.1	3.9	6.4
2002.....	*	0.2	0.3	0.6	0.8	1.0	1.8	2.2	3.9	8.1
2001.....	0.2	0.3	0.3	0.5	0.7	1.2	1.8	2.7	4.6	6.0
2000.....	*	0.4	0.5	0.5	0.8	1.2	2.0	3.0	3.8	9.9
1999.....	0.2	0.3	0.5	0.6	0.7	1.0	2.2	2.7	5.4	8.7
All races 3/, female										
2006.....	*	*	0.2	*	0.4	0.6	1.2	1.7	2.6	5.9
2005.....	*	*	0.2	*	0.5	0.7	0.9	1.4	2.8	6.5
2004.....	*	*	0.3	0.2	0.5	0.8	1.1	2.1	3.1	6.9
2003.....	*	*	*	0.4	0.4	0.6	1.1	2.1	3.7	7.3
2002.....	*	*	*	0.4	*	0.7	1.1	2.0	3.8	7.5
2001.....	*	*	0.2	*	0.6	0.9	1.4	2.2	4.6	8.8
2000.....	*	*	0.2	0.4	0.7	0.9	1.2	2.0	4.1	9.4
1999.....	*	*	*	0.4	0.5	0.8	1.5	2.7	4.4	8.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Gastric ulcer (K25)										
White, both sexes										
2006 4/.....	*	0.1	0.2	0.3	0.5	0.6	1.2	1.6	2.7	6.1
2005 4/.....	0.1	0.2	0.3	0.3	0.5	0.6	0.9	1.6	3.0	6.2
2004 4/.....	*	0.2	0.3	0.3	0.5	0.7	1.1	2.1	3.4	6.8
2003 4/.....	*	0.2	0.2	0.5	0.5	0.7	1.1	2.0	3.7	7.0
2002.....	*	0.2	0.2	0.5	0.5	0.8	1.5	2.0	3.8	7.8
2001.....	*	0.2	0.2	0.3	0.6	0.9	1.5	2.4	4.6	8.0
2000.....	*	0.2	0.3	0.4	0.7	1.0	1.5	2.3	4.0	9.4
1999.....	0.1	0.2	0.3	0.4	0.6	0.7	1.8	2.7	4.8	8.6
White, male										
2006 4/.....	*	*	0.3	0.4	0.6	0.6	1.0	1.4	2.9	6.1
2005 4/.....	*	0.2	0.4	0.4	0.5	0.6	1.0	1.7	3.5	5.7
2004 4/.....	*	*	0.4	0.3	0.6	0.8	1.1	2.0	3.8	6.7
2003 4/.....	*	0.2	0.3	0.5	0.7	0.8	1.1	1.9	3.8	6.1
2002.....	*	*	0.3	0.5	0.6	0.9	1.8	2.1	3.8	8.3
2001.....	*	0.3	0.3	0.4	0.6	0.9	1.6	2.7	4.5	5.9
2000.....	*	0.3	0.5	0.4	0.7	1.1	1.8	2.8	3.6	9.3
1999.....	*	0.2	0.4	0.5	0.7	0.8	2.1	2.7	5.3	8.1
White, female										
2006 4/.....	*	*	*	*	0.4	0.5	1.3	1.7	2.6	6.1
2005 4/.....	*	*	*	*	0.5	0.7	0.9	1.4	2.7	6.3
2004 4/.....	*	*	*	*	0.5	0.6	1.1	2.1	3.1	6.9
2003 4/.....	*	*	*	0.4	0.4	0.6	1.2	2.0	3.6	7.4
2002.....	*	*	*	0.5	*	0.8	1.2	1.9	3.7	7.5
2001.....	*	*	*	*	0.6	0.8	1.4	2.2	4.7	8.8
2000.....	*	*	*	0.4	0.7	0.9	1.2	2.0	4.3	9.5
1999.....	*	*	*	0.4	0.5	0.7	1.5	2.7	4.4	8.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Duodenal ulcer (K26)										
All races 3/, both sexes										
2006.....	*	0.2	0.3	0.4	0.6	0.7	1.2	1.8	3.6	6.8
2005.....	0.1	0.2	0.3	0.4	0.7	0.8	1.4	2.2	3.7	6.9
2004.....	0.1	0.2	0.2	0.4	0.6	0.9	1.5	2.5	3.9	7.7
2003.....	0.1	0.3	0.3	0.4	0.5	0.8	1.5	2.5	4.6	8.5
2002.....	0.1	0.2	0.3	0.5	0.7	0.9	1.7	3.1	4.3	9.0
2001.....	0.2	0.3	0.3	0.4	0.8	1.1	1.6	2.9	5.4	11.0
2000.....	0.1	0.2	0.4	0.6	0.9	1.0	1.7	3.0	5.6	10.1
1999.....	0.1	0.2	0.3	0.4	1.0	1.2	1.9	2.9	4.9	10.0
All races 3/, male										
2006.....	*	0.4	0.5	0.6	0.7	1.0	1.5	2.2	4.6	7.6
2005.....	0.2	0.4	0.4	0.6	1.1	1.1	2.1	2.9	5.0	7.7
2004.....	*	0.3	0.3	0.5	0.9	1.1	2.0	3.3	4.2	9.2
2003.....	*	0.4	0.4	0.4	0.8	1.1	2.2	3.4	6.1	10.2
2002.....	*	0.3	0.4	0.8	0.9	1.2	2.2	3.6	6.0	10.1
2001.....	0.3	0.4	0.5	0.6	1.1	1.4	2.0	3.6	6.4	13.8
2000.....	0.2	0.3	0.7	0.8	1.1	1.4	2.3	3.8	6.4	12.3
1999.....	*	0.3	0.5	0.5	1.4	1.9	2.9	4.1	6.7	11.9
All races 3/, female										
2006.....	*	*	*	*	0.4	0.5	1.0	1.5	3.0	6.4
2005.....	*	*	*	*	0.3	0.5	0.8	1.7	3.0	6.6
2004.....	*	*	*	0.3	0.4	0.7	1.1	1.9	3.7	7.1
2003.....	*	*	*	0.3	*	0.5	1.0	1.8	3.6	7.8
2002.....	*	0.2	*	0.3	0.5	0.6	1.3	2.8	3.3	8.5
2001.....	*	*	*	0.3	0.4	0.8	1.3	2.4	4.8	9.8
2000.....	*	*	*	0.3	0.7	0.7	1.3	2.5	5.1	9.2
1999.....	*	*	*	0.3	0.6	0.6	1.2	2.1	3.9	9.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Duodenal ulcer (K26)										
White, both sexes										
2006 4/.....	*	0.2	0.3	0.4	0.5	0.7	1.2	1.7	3.8	7.1
2005 4/.....	0.1	0.2	0.3	0.4	0.7	0.8	1.4	2.1	3.9	7.0
2004 4/.....	*	0.2	0.3	0.4	0.6	0.9	1.5	2.6	4.0	8.1
2003 4/.....	*	0.2	0.3	0.3	0.5	0.8	1.6	2.4	4.6	8.8
2002.....	0.1	0.2	0.2	0.5	0.7	0.9	1.8	3.2	4.7	9.5
2001.....	0.2	0.2	0.3	0.4	0.8	1.1	1.6	2.9	5.6	11.6
2000.....	*	0.2	0.4	0.6	0.9	1.0	1.7	3.1	5.8	10.5
1999.....	0.1	0.2	0.3	0.4	1.0	1.2	1.9	3.0	5.1	10.2
White, male										
2006 4/.....	*	0.3	0.4	0.6	0.7	0.9	1.4	2.2	5.0	7.5
2005 4/.....	*	0.3	0.4	0.6	1.0	1.1	2.0	2.8	5.3	7.4
2004 4/.....	*	0.3	0.3	0.5	0.9	1.1	1.9	3.4	4.2	9.4
2003 4/.....	*	0.3	0.4	0.3	0.7	1.1	2.2	3.3	6.1	10.2
2002.....	*	0.3	0.3	0.7	0.9	1.2	2.2	3.6	6.5	10.9
2001.....	0.3	0.3	0.4	0.5	1.1	1.5	1.9	3.5	6.6	14.5
2000.....	*	0.3	0.6	0.8	1.2	1.4	2.3	3.8	6.6	12.7
1999.....	*	*	0.4	0.5	1.4	1.8	3.0	4.2	6.9	12.5
White, female										
2006 4/.....	*	*	*	*	0.4	0.4	1.0	1.4	3.1	6.9
2005 4/.....	*	*	*	*	*	0.6	0.8	1.6	3.0	6.9
2004 4/.....	*	*	*	*	0.4	0.8	1.2	2.0	3.8	7.5
2003 4/.....	*	*	*	0.3	*	0.5	1.0	1.7	3.6	8.2
2002.....	*	*	*	*	0.5	0.6	1.4	2.9	3.5	8.9
2001.....	*	*	*	*	0.4	0.8	1.4	2.4	5.1	10.4
2000.....	*	*	*	0.3	0.6	0.7	1.3	2.5	5.3	9.6
1999.....	*	*	*	*	0.6	0.6	1.1	2.1	3.9	9.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer, site unspecified (K27)										
All races 3/, both sexes										
2006.....	*	0.2	0.2	0.3	0.4	0.8	1.0	1.8	3.3	7.9
2005.....	*	0.1	0.2	0.3	0.4	0.7	1.4	2.0	3.9	8.2
2004.....	*	0.1	0.2	0.3	0.5	0.8	1.2	2.1	4.1	9.7
2003.....	*	0.1	0.2	0.3	0.6	0.8	1.4	2.7	5.0	10.5
2002.....	*	0.1	0.3	0.3	0.5	0.8	1.5	2.7	4.9	11.3
2001.....	*	0.1	0.2	0.5	0.7	1.0	1.7	2.9	5.8	14.2
2000.....	*	0.2	0.2	0.5	0.6	1.0	1.7	3.1	5.9	15.5
1999.....	*	0.1	0.2	0.5	0.6	1.2	2.1	3.4	7.2	15.6
All races 3/, male										
2006.....	*	0.2	*	0.3	0.5	0.9	1.3	2.5	3.6	8.6
2005.....	*	*	0.3	0.3	0.5	0.9	1.5	2.6	4.4	9.2
2004.....	*	*	0.3	0.4	0.6	0.9	1.3	2.8	4.5	9.0
2003.....	*	*	0.3	0.4	0.8	1.0	1.8	3.3	5.8	11.9
2002.....	*	*	0.4	0.4	0.5	1.0	2.0	3.4	6.5	12.8
2001.....	*	*	0.3	0.7	0.9	1.4	1.8	3.9	6.2	14.9
2000.....	*	0.2	0.3	0.7	1.0	1.3	2.5	4.3	7.2	18.2
1999.....	*	*	0.3	0.6	0.9	1.6	2.7	4.1	8.8	16.4
All races 3/, female										
2006.....	*	*	*	*	0.3	0.6	0.9	1.3	3.2	7.6
2005.....	*	*	*	*	0.3	0.5	1.3	1.6	3.6	7.8
2004.....	*	*	*	*	0.4	0.7	1.2	1.7	3.8	10.0
2003.....	*	*	*	*	0.3	0.6	1.1	2.2	4.5	9.9
2002.....	*	*	*	0.3	0.5	0.7	1.1	2.2	3.8	10.7
2001.....	*	*	*	0.3	0.4	0.6	1.5	2.2	5.6	13.9
2000.....	*	*	*	*	*	0.8	1.2	2.3	5.1	14.5
1999.....	*	*	*	0.3	0.4	0.9	1.6	2.9	6.3	15.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Peptic ulcer, site unspecified (K27)										
White, both sexes										
2006 4/.....	*	0.1	0.2	0.3	0.4	0.8	1.0	1.8	3.3	7.9
2005 4/.....	*	*	0.2	0.3	0.4	0.7	1.3	2.0	3.9	8.3
2004 4/.....	*	*	0.2	0.3	0.5	0.8	1.2	2.1	4.1	9.9
2003 4/.....	*	*	0.2	0.3	0.6	0.8	1.4	2.7	4.9	10.7
2002.....	*	0.1	0.3	0.3	0.5	0.8	1.5	2.6	5.0	11.5
2001.....	*	0.2	0.2	0.4	0.6	1.0	1.7	2.9	5.8	14.6
2000.....	*	0.2	0.2	0.5	0.6	1.0	1.8	3.1	6.0	15.9
1999.....	*	*	0.2	0.5	0.6	1.2	2.0	3.4	7.2	16.2
White, male										
2006 4/.....	*	*	*	0.4	0.5	0.9	1.2	2.3	3.4	8.4
2005 4/.....	*	*	0.3	0.3	0.5	0.8	1.4	2.5	4.5	9.2
2004 4/.....	*	*	0.3	0.3	0.5	1.0	1.3	2.7	4.4	9.2
2003 4/.....	*	*	0.3	0.4	0.8	1.1	1.8	3.2	5.3	12.2
2002.....	*	*	0.4	*	0.5	1.0	1.9	3.3	6.5	12.9
2001.....	*	*	*	0.5	0.9	1.4	1.8	3.8	6.1	15.0
2000.....	*	*	*	0.7	1.0	1.3	2.5	4.1	7.4	18.4
1999.....	*	*	*	0.6	0.8	1.5	2.7	4.1	8.7	17.4
White, female										
2006 4/.....	*	*	*	*	*	0.6	0.9	1.4	3.3	7.6
2005 4/.....	*	*	*	*	*	0.6	1.3	1.6	3.6	7.9
2004 4/.....	*	*	*	*	0.4	0.7	1.1	1.6	3.9	10.3
2003 4/.....	*	*	*	*	*	0.6	1.1	2.3	4.7	10.1
2002.....	*	*	*	*	0.5	0.7	1.1	2.2	4.1	10.9
2001.....	*	*	*	*	0.4	0.6	1.6	2.3	5.7	14.4
2000.....	*	*	*	*	*	0.7	1.2	2.3	5.2	15.0
1999.....	*	*	*	0.4	*	0.9	1.5	2.8	6.3	15.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Gastritis and duodenitis (K29)										
All races 3/, both sexes										
2006.....	*	*	*	0.1	*	*	0.2	0.4	0.7	2.1
2005.....	*	0.1	*	0.1	*	*	0.3	0.5	0.9	2.2
2004.....	*	*	*	0.2	0.2	*	0.3	0.7	1.0	2.6
2003.....	*	*	0.1	*	0.2	*	0.4	0.5	1.0	2.9
2002.....	*	*	*	*	*	*	0.3	0.6	1.1	2.9
2001.....	*	*	*	*	*	0.2	0.3	0.6	1.0	3.8
2000.....	*	*	*	0.2	0.2	0.2	0.3	0.6	1.2	3.5
1999.....	*	*	*	*	*	0.2	0.3	0.6	1.5	3.8
All races 3/, male										
2006.....	*	*	*	0.2	*	*	*	*	1.1	2.4
2005.....	*	*	*	*	*	*	*	0.8	1.0	1.8
2004.....	*	*	*	*	0.3	*	*	0.7	1.1	3.0
2003.....	*	*	0.2	*	*	*	*	0.7	1.2	2.8
2002.....	*	*	*	*	*	*	*	0.8	1.0	2.9
2001.....	*	*	*	*	*	*	*	*	*	4.8
2000.....	*	*	*	*	*	*	*	*	1.2	3.3
1999.....	*	*	*	*	*	*	*	0.8	1.7	3.7
All races 3/, female										
2006.....	*	*	*	*	*	*	*	*	*	1.9
2005.....	*	*	*	*	*	*	*	*	0.8	2.3
2004.....	*	*	*	*	*	*	*	0.6	0.9	2.4
2003.....	*	*	*	*	*	*	*	*	0.9	3.0
2002.....	*	*	*	*	*	*	*	0.5	1.1	2.9
2001.....	*	*	*	*	*	*	*	0.5	1.0	3.3
2000.....	*	*	*	*	*	*	*	0.6	1.2	3.6
1999.....	*	*	*	*	*	*	*	*	1.4	3.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Gastritis and duodenitis (K29)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*	0.4	0.7	2.0
2005 4/.....	*	*	*	0.1	*	*	0.3	0.5	0.9	2.2
2004 4/.....	*	*	*	0.2	*	*	0.3	0.6	0.8	2.6
2003 4/.....	*	*	0.1	*	*	*	0.4	0.4	0.9	2.9
2002.....	*	*	*	*	*	*	0.3	0.7	1.0	2.9
2001.....	*	*	*	*	*	*	0.3	0.6	0.9	4.0
2000.....	*	*	*	0.2	*	*	0.3	0.6	1.3	3.5
1999.....	*	*	*	*	*	*	0.3	0.5	1.3	3.9

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	2.2
2005 4/.....	*	*	*	*	*	*	*	0.8	*	1.9
2004 4/.....	*	*	*	*	*	*	*	0.8	*	3.1
2003 4/.....	*	*	*	*	*	*	*	*	1.1	2.8
2002.....	*	*	*	*	*	*	*	0.9	1.2	2.9
2001.....	*	*	*	*	*	*	*	*	*	5.0
2000.....	*	*	*	*	*	*	*	*	1.3	3.3
1999.....	*	*	*	*	*	*	*	0.7	1.3	3.8

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	1.9
2005 4/.....	*	*	*	*	*	*	*	*	0.8	2.4
2004 4/.....	*	*	*	*	*	*	*	*	0.8	2.4
2003 4/.....	*	*	*	*	*	*	*	*	0.8	2.9
2002.....	*	*	*	*	*	*	*	*	1.0	2.9
2001.....	*	*	*	*	*	*	*	0.5	0.9	3.6
2000.....	*	*	*	*	*	*	*	0.6	1.3	3.6
1999.....	*	*	*	*	*	*	*	*	1.3	3.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Dyspepsia and other diseases of stomach and duodenum (K30-K31)

All races 3/, both sexes

2006.....	*	*	0.1	0.2	0.2	0.4	0.5	1.0	1.7	3.1
2005.....	*	0.1	0.2	0.2	0.3	0.4	0.5	0.8	1.3	3.3
2004.....	*	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.7	3.0
2003.....	0.1	*	0.1	0.2	0.2	0.4	0.7	1.0	1.9	3.5
2002.....	0.1	*	0.1	0.2	0.3	0.4	0.6	1.1	1.7	3.2
2001.....	*	*	0.2	*	0.2	0.4	0.6	1.0	1.7	3.6
2000.....	*	*	0.2	0.2	0.2	0.5	0.7	1.2	1.6	4.0
1999.....	*	0.1	*	*	0.3	0.3	0.6	1.0	1.9	4.2

All races 3/, male

2006.....	*	*	*	*	*	*	0.6	1.1	1.3	3.1
2005.....	*	*	*	0.3	*	*	0.6	0.8	1.7	3.2
2004.....	*	*	*	*	*	0.5	0.6	1.2	2.0	3.6
2003.....	*	*	*	*	*	*	0.9	1.1	2.0	2.7
2002.....	*	*	*	*	*	0.6	0.7	1.3	1.6	2.9
2001.....	*	*	*	*	*	0.5	0.6	0.9	2.2	3.3
2000.....	*	*	*	*	*	0.7	0.9	1.2	1.7	4.3
1999.....	*	*	*	*	*	*	0.6	1.2	1.6	4.8

All races 3/, female

2006.....	*	*	*	*	*	0.4	0.4	1.0	2.0	3.0
2005.....	*	*	*	*	0.3	0.5	0.5	0.9	1.0	3.3
2004.....	*	*	0.2	*	0.3	*	0.5	1.1	1.5	2.8
2003.....	*	*	*	*	*	0.5	0.6	1.0	1.8	3.9
2002.....	*	*	*	0.3	*	*	0.5	1.1	1.8	3.4
2001.....	*	*	*	*	*	*	0.6	1.0	1.5	3.7
2000.....	*	*	*	*	*	*	0.6	1.1	1.5	3.9
1999.....	*	*	*	*	0.4	*	0.6	0.9	2.1	4.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Dyspepsia and other diseases of stomach and duodenum (K30-K31)

White, both sexes

2006 4/.....	*	*	0.1	0.2	0.3	0.3	0.6	1.1	1.7	3.1
2005 4/.....	*	*	0.2	0.2	0.3	0.4	0.5	0.8	1.3	3.3
2004 4/.....	*	*	0.2	0.2	0.3	0.4	0.5	1.2	1.8	3.0
2003 4/.....	*	*	0.1	0.2	0.2	0.4	0.8	0.9	1.8	3.6
2002.....	*	*	0.1	0.2	0.3	0.4	0.6	1.2	1.6	3.3
2001.....	*	*	0.2	*	*	0.4	0.6	0.9	1.8	3.7
2000.....	*	*	0.2	0.2	0.2	0.4	0.7	1.1	1.6	4.1
1999.....	*	*	*	*	0.3	0.3	0.6	1.1	1.9	4.2

White, male

2006 4/.....	*	*	*	*	*	*	0.7	1.2	1.2	3.2
2005 4/.....	*	*	*	0.3	*	*	*	0.7	1.5	3.3
2004 4/.....	*	*	*	*	*	*	*	1.2	2.0	3.5
2003 4/.....	*	*	*	*	*	*	0.9	1.0	1.9	2.9
2002.....	*	*	*	*	*	0.5	0.6	1.3	1.5	2.8
2001.....	*	*	*	*	*	*	*	0.8	2.3	3.5
2000.....	*	*	*	*	*	0.6	0.9	1.2	1.9	4.4
1999.....	*	*	*	*	*	*	*	1.4	1.6	4.9

White, female

2006 4/.....	*	*	*	*	*	*	0.5	1.0	2.0	3.1
2005 4/.....	*	*	*	*	*	0.5	0.5	0.9	1.1	3.3
2004 4/.....	*	*	*	*	0.4	*	0.5	1.1	1.6	2.8
2003 4/.....	*	*	*	*	*	0.5	0.6	0.9	1.8	3.9
2002.....	*	*	*	*	*	*	0.6	1.1	1.7	3.5
2001.....	*	*	*	*	*	*	0.6	1.0	1.5	3.8
2000.....	*	*	*	*	*	*	0.6	1.1	1.5	4.0
1999.....	*	*	*	*	*	*	0.6	0.9	2.1	4.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of appendix (K35-K38)										
All races 3/, both sexes										
2006.....	*	*	*	0.2	0.2	0.4	0.5	0.6	1.0	1.7
2005.....	*	*	0.1	0.2	0.2	0.4	0.4	0.8	0.9	2.1
2004.....	*	*	0.2	0.1	0.2	0.3	0.6	0.8	1.4	1.7
2003.....	*	0.1	0.1	*	0.2	0.3	0.4	0.8	1.4	1.7
2002.....	*	0.1	0.2	0.2	0.3	0.3	0.7	0.8	1.1	2.0
2001.....	*	0.1	*	0.2	0.2	0.4	0.6	0.5	1.0	2.1
2000.....	*	*	0.1	0.2	0.3	0.3	0.5	0.7	1.4	1.6
1999.....	*	0.1	0.1	0.2	0.2	0.3	0.4	0.6	1.1	1.5
All races 3/, male										
2006.....	*	*	*	0.2	*	0.5	0.8	1.1	1.5	2.3
2005.....	*	*	*	0.3	*	0.6	0.7	0.9	1.3	2.8
2004.....	*	*	0.2	*	*	*	0.7	1.1	1.7	2.5
2003.....	*	*	*	*	*	0.4	0.7	0.9	2.3	2.4
2002.....	*	*	0.2	*	0.5	0.5	0.8	1.1	1.6	2.2
2001.....	*	*	*	*	*	0.5	0.9	0.7	1.6	2.8
2000.....	*	*	*	0.3	0.4	*	0.6	1.0	1.6	2.0
1999.....	*	*	*	*	*	*	0.5	0.9	*	2.9
All races 3/, female										
2006.....	*	*	*	*	*	*	*	*	0.7	1.5
2005.....	*	*	*	*	*	*	*	0.7	0.7	1.8
2004.....	*	*	*	*	*	*	0.5	0.6	1.2	1.4
2003.....	*	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	*	0.5	0.6	0.8	2.0
2001.....	*	*	*	*	*	*	0.4	*	0.7	1.8
2000.....	*	*	*	*	*	*	0.4	0.5	1.2	1.5
1999.....	*	*	*	*	*	*	*	*	1.1	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of appendix (K35-K38)										
White, both sexes										
2006 4/.....	*	*	*	0.2	*	0.4	0.5	0.6	1.0	1.8
2005 4/.....	*	*	*	0.2	0.2	0.4	0.4	0.7	1.0	2.2
2004 4/.....	*	*	0.1	0.2	0.2	0.3	0.6	0.8	1.3	1.8
2003 4/.....	*	*	*	*	*	0.3	0.4	0.7	1.5	1.7
2002.....	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.2	2.1
2001.....	*	*	*	*	*	0.3	0.7	0.5	0.9	2.1
2000.....	*	*	*	0.2	0.3	0.2	0.4	0.6	1.3	1.7
1999.....	*	*	0.2	*	*	0.3	0.4	0.6	1.0	1.5
White, male										
2006 4/.....	*	*	*	*	*	0.6	0.7	1.0	1.4	2.3
2005 4/.....	*	*	*	*	*	0.6	0.7	0.9	1.4	2.9
2004 4/.....	*	*	*	*	*	*	0.7	1.1	1.7	2.3
2003 4/.....	*	*	*	*	*	*	0.7	0.8	2.4	2.5
2002.....	*	*	*	*	0.5	*	0.7	1.2	1.6	2.0
2001.....	*	*	*	*	*	*	1.0	0.7	1.5	2.5
2000.....	*	*	*	*	0.5	*	*	0.9	1.5	2.0
1999.....	*	*	*	*	*	*	*	0.9	*	3.1
White, female										
2006 4/.....	*	*	*	*	*	*	*	*	0.7	1.5
2005 4/.....	*	*	*	*	*	*	*	0.6	0.7	1.9
2004 4/.....	*	*	*	*	*	*	0.5	0.6	1.0	1.6
2003 4/.....	*	*	*	*	*	*	*	0.6	0.9	1.3
2002.....	*	*	*	*	*	*	0.5	0.6	0.9	2.1
2001.....	*	*	*	*	*	*	*	*	*	1.9
2000.....	*	*	*	*	*	*	*	*	1.2	1.6
1999.....	*	*	*	*	*	*	*	*	1.1	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hernia (K40-K46)										
All races 3/, both sexes										
2006.....	0.1	0.1	0.3	0.5	0.7	0.9	1.7	2.8	5.4	12.1
2005.....	0.1	0.1	0.3	0.4	0.9	0.9	1.4	2.5	4.8	12.1
2004.....	0.1	0.2	0.2	0.4	0.5	0.9	1.8	2.9	5.3	11.0
2003.....	0.1	0.2	0.3	0.5	0.7	1.0	1.6	2.7	4.8	11.9
2002.....	0.1	0.2	0.2	0.4	0.6	0.9	1.6	2.8	5.3	12.8
2001.....	0.1	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.6
2000.....	*	0.1	0.3	0.4	0.6	0.9	1.4	3.0	4.9	13.1
1999.....	0.1	*	0.2	0.3	0.7	0.8	1.6	2.5	5.4	12.7
All races 3/, male										
2006.....	*	*	0.3	0.4	0.8	0.8	1.9	2.9	5.7	13.5
2005.....	*	*	0.3	0.4	0.9	0.7	1.3	2.4	4.6	12.7
2004.....	*	0.2	0.3	0.4	0.5	0.8	2.2	3.2	4.7	11.3
2003.....	*	0.2	0.4	0.4	0.7	1.1	1.6	3.0	5.2	12.3
2002.....	*	*	*	0.4	0.5	0.9	1.6	2.7	5.2	11.7
2001.....	*	*	0.3	0.3	0.8	0.9	1.4	2.3	5.3	12.0
2000.....	*	*	0.4	0.3	0.6	0.9	1.4	3.0	5.7	12.6
1999.....	*	*	0.2	0.3	0.8	0.8	1.5	2.5	6.5	13.2
All races 3/, female										
2006.....	*	*	0.3	0.5	0.7	1.0	1.5	2.7	5.1	11.4
2005.....	*	*	0.2	0.4	0.8	1.1	1.4	2.5	5.0	11.8
2004.....	*	0.2	0.2	0.4	0.5	1.0	1.4	2.7	5.6	10.8
2003.....	*	0.2	0.3	0.6	0.7	1.0	1.5	2.5	4.6	11.8
2002.....	*	*	0.3	0.4	0.7	1.0	1.5	2.9	5.3	13.3
2001.....	*	*	0.3	0.5	0.6	0.8	1.6	2.6	4.8	12.8
2000.....	*	*	0.3	0.4	0.6	0.9	1.5	3.0	4.4	13.3
1999.....	*	*	*	*	0.6	0.8	1.6	2.6	4.8	12.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Hernia (K40-K46)										
White, both sexes										
2006 4/.....	0.1	0.1	0.3	0.4	0.8	0.9	1.6	2.8	5.3	12.6
2005 4/.....	*	0.1	0.3	0.4	0.9	0.9	1.4	2.5	5.0	12.5
2004 4/.....	*	0.2	0.2	0.4	0.5	0.9	1.8	3.1	5.4	11.2
2003 4/.....	*	0.2	0.3	0.4	0.7	1.1	1.6	2.8	4.7	12.3
2002.....	*	0.1	0.2	0.4	0.6	0.9	1.6	3.0	5.4	13.4
2001.....	*	0.1	0.3	0.4	0.7	0.8	1.5	2.5	5.0	12.8
2000.....	*	0.1	0.3	0.4	0.6	0.8	1.5	3.0	5.0	13.3
1999.....	*	*	0.2	0.3	0.7	0.7	1.6	2.5	5.5	13.0
White, male										
2006 4/.....	*	*	0.3	0.3	0.8	0.8	1.9	2.9	5.6	14.0
2005 4/.....	*	*	0.3	0.4	0.9	0.7	1.4	2.4	4.6	12.8
2004 4/.....	*	*	0.3	0.4	0.5	0.7	2.2	3.3	4.7	11.4
2003 4/.....	*	*	0.4	0.4	0.7	1.1	1.6	3.1	5.1	12.5
2002.....	*	*	*	0.4	0.5	0.8	1.6	2.9	5.4	11.9
2001.....	*	*	0.3	*	0.7	0.9	1.4	2.3	5.2	11.8
2000.....	*	*	0.3	*	0.6	0.8	1.5	3.1	5.7	12.7
1999.....	*	*	*	0.4	0.8	0.8	1.6	2.5	6.6	12.9
White, female										
2006 4/.....	*	*	0.2	0.5	0.7	1.0	1.4	2.7	5.2	11.9
2005 4/.....	*	*	*	0.4	0.8	1.1	1.4	2.6	5.2	12.3
2004 4/.....	*	*	*	0.4	0.5	1.1	1.4	2.9	5.8	11.2
2003 4/.....	*	*	*	0.5	0.7	1.0	1.6	2.6	4.5	12.2
2002.....	*	*	*	0.4	0.6	0.9	1.6	3.0	5.4	14.0
2001.....	*	*	0.3	0.5	0.6	0.8	1.6	2.6	5.0	13.2
2000.....	*	*	0.3	0.5	0.6	0.8	1.5	2.9	4.6	13.6
1999.....	*	*	*	*	0.5	0.7	1.5	2.5	4.9	13.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Crohn's disease and ulcerative colitis (K50-K51)

All races 3/, both sexes

2006.....	0.1	0.1	0.3	0.4	0.6	0.8	1.2	1.7	2.3	3.0
2005.....	0.1	0.2	0.3	0.3	0.5	0.7	1.1	1.7	2.5	3.5
2004.....	0.1	0.2	0.3	0.4	0.6	0.7	1.1	1.8	2.4	3.3
2003.....	0.1	0.2	0.3	0.4	0.5	0.8	1.1	1.6	2.3	3.1
2002.....	0.2	0.2	0.3	0.3	0.4	0.7	1.0	1.7	2.3	3.1
2001.....	*	0.2	0.2	0.5	0.5	0.7	1.1	1.6	2.4	3.8
2000.....	*	0.1	0.2	0.4	0.4	0.6	1.1	1.6	2.2	3.3
1999.....	0.1	0.2	0.3	0.3	0.4	0.6	1.1	1.6	2.5	3.3

All races 3/, male

2006.....	*	*	0.2	0.4	0.5	0.8	1.1	1.8	2.5	3.0
2005.....	*	0.2	0.2	0.3	0.5	0.8	1.0	1.6	2.2	2.4
2004.....	*	0.2	0.3	0.4	0.7	0.7	1.1	1.8	2.2	2.7
2003.....	*	0.2	0.2	0.4	0.5	0.7	1.1	1.7	2.3	3.5
2002.....	0.2	0.2	*	*	*	0.8	1.0	1.7	2.3	2.7
2001.....	*	*	0.2	0.5	0.5	0.7	1.3	1.6	2.3	3.5
2000.....	*	*	*	0.3	*	0.6	1.1	1.6	1.6	2.9
1999.....	*	*	0.3	0.3	*	0.7	1.1	1.6	2.5	2.7

All races 3/, female

2006.....	*	*	0.3	0.4	0.6	0.9	1.3	1.7	2.2	2.9
2005.....	*	0.2	0.3	0.3	0.4	0.7	1.1	1.7	2.7	4.0
2004.....	*	0.2	0.3	0.4	0.5	0.7	1.1	1.8	2.6	3.6
2003.....	*	0.3	0.3	0.3	0.5	0.9	1.0	1.5	2.3	2.8
2002.....	0.2	0.2	0.3	0.3	0.5	0.7	1.0	1.7	2.3	3.2
2001.....	*	*	0.2	0.5	0.4	0.6	0.9	1.5	2.5	3.9
2000.....	*	*	0.3	0.4	0.4	0.7	1.2	1.6	2.6	3.4
1999.....	*	0.3	0.3	0.3	0.5	0.5	1.1	1.6	2.6	3.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Crohn's disease and ulcerative colitis (K50-K51)										
White, both sexes										
2006 4/.....	0.1	0.1	0.3	0.4	0.6	0.9	1.3	1.9	2.5	3.2
2005 4/.....	*	0.2	0.3	0.3	0.5	0.8	1.2	1.9	2.7	3.7
2004 4/.....	0.1	0.2	0.3	0.4	0.6	0.7	1.1	2.0	2.6	3.6
2003 4/.....	0.1	0.2	0.3	0.4	0.5	0.8	1.1	1.8	2.4	3.3
2002.....	0.2	0.2	0.3	0.3	0.5	0.8	1.0	1.8	2.5	3.4
2001.....	*	0.2	0.2	0.5	0.5	0.7	1.2	1.7	2.5	3.9
2000.....	*	0.2	0.2	0.4	0.4	0.7	1.2	1.7	2.4	3.5
1999.....	0.2	0.2	0.3	0.3	0.4	0.7	1.2	1.6	2.8	3.5
White, male										
2006 4/.....	*	*	*	0.4	0.5	0.9	1.1	1.9	2.7	3.4
2005 4/.....	*	0.2	*	0.3	0.5	0.9	1.1	1.8	2.5	2.6
2004 4/.....	*	0.2	0.3	0.4	0.7	0.7	1.2	1.9	2.3	2.9
2003 4/.....	*	*	0.3	0.5	0.5	0.7	1.1	1.8	2.5	3.7
2002.....	*	0.2	*	*	*	0.9	1.0	1.7	2.5	2.9
2001.....	*	*	*	0.5	0.5	0.8	1.4	1.7	2.5	3.5
2000.....	*	*	*	0.4	*	0.6	1.1	1.7	1.8	3.2
1999.....	*	*	*	0.4	*	0.8	1.2	1.6	2.7	3.0
White, female										
2006 4/.....	*	*	0.4	0.4	0.7	0.9	1.4	1.9	2.4	3.1
2005 4/.....	*	0.2	0.3	0.3	0.5	0.7	1.2	1.9	2.9	4.1
2004 4/.....	*	0.2	0.4	0.5	0.6	0.8	1.1	2.0	2.9	3.9
2003 4/.....	*	0.3	0.3	0.4	0.6	1.0	1.1	1.7	2.4	3.1
2002.....	*	0.2	0.3	0.3	0.5	0.8	1.0	1.9	2.5	3.6
2001.....	*	*	*	0.5	0.5	0.7	1.0	1.6	2.6	4.0
2000.....	*	*	0.3	0.4	0.5	0.8	1.3	1.7	2.8	3.6
1999.....	*	0.3	0.4	*	0.5	0.6	1.2	1.7	2.8	3.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

All races 3/, both sexes

2006.....	0.5	0.8	1.6	2.7	4.9	8.3	16.3	28.4	50.6	99.2
2005.....	0.4	0.9	1.5	2.6	4.6	8.5	17.0	28.5	49.6	103.1
2004.....	0.6	0.8	1.5	2.6	4.8	8.7	16.1	29.5	49.9	106.9
2003.....	0.5	0.9	1.6	2.9	4.8	9.7	17.4	29.7	52.3	111.5
2002.....	0.5	0.9	1.4	2.8	4.8	9.0	17.0	29.6	53.0	113.3
2001.....	0.4	0.8	1.6	2.7	5.4	10.1	17.8	29.7	52.1	113.5
2000.....	0.5	0.9	1.7	2.8	5.3	9.5	17.5	31.5	52.9	116.2
1999.....	0.5	0.8	1.5	2.9	5.3	10.1	17.1	29.9	52.5	116.2

All races 3/, male

2006.....	0.6	0.8	1.6	2.5	4.8	7.8	15.1	26.6	47.4	82.9
2005.....	0.4	0.9	1.6	2.6	4.6	8.0	15.9	26.8	44.0	88.0
2004.....	0.6	0.8	1.5	2.4	4.5	7.9	14.6	28.0	45.1	88.8
2003.....	0.5	0.8	1.6	3.1	4.5	8.8	16.2	26.8	48.8	92.6
2002.....	0.6	0.8	1.5	2.5	4.7	8.1	15.8	26.8	49.7	96.7
2001.....	0.5	0.8	1.6	2.3	5.1	9.1	17.6	27.2	47.9	98.5
2000.....	0.4	0.9	1.6	2.7	5.2	8.9	16.6	30.0	48.2	98.5
1999.....	0.6	0.8	1.4	2.9	4.9	8.7	16.7	27.9	48.7	95.6

All races 3/, female

2006.....	0.5	0.7	1.6	2.8	5.0	8.7	17.3	29.8	52.7	106.8
2005.....	0.4	0.8	1.5	2.7	4.6	9.0	17.9	29.7	53.0	110.1
2004.....	0.5	0.9	1.4	2.7	5.1	9.5	17.3	30.5	53.0	115.1
2003.....	0.5	1.0	1.6	2.7	5.1	10.5	18.3	31.8	54.5	119.8
2002.....	0.4	0.9	1.4	3.0	4.9	9.8	18.0	31.6	55.0	120.5
2001.....	0.4	0.9	1.6	3.0	5.6	11.0	18.0	31.5	54.6	119.8
2000.....	0.5	0.9	1.8	2.8	5.4	10.0	18.2	32.5	55.7	123.4
1999.....	0.5	0.7	1.6	3.0	5.7	11.3	17.5	31.3	54.8	124.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

White, both sexes

2006 4/.....	0.5	0.7	1.4	2.4	4.8	8.1	16.5	28.5	51.0	101.7
2005 4/.....	0.4	0.8	1.4	2.5	4.3	8.1	17.0	28.3	49.9	104.7
2004 4/.....	0.5	0.7	1.4	2.3	4.6	8.7	16.0	29.2	50.2	108.5
2003 4/.....	0.4	0.8	1.4	2.6	4.5	9.5	17.2	29.4	52.8	113.7
2002.....	0.5	0.8	1.3	2.6	4.6	8.7	16.4	29.8	53.0	115.9
2001.....	0.4	0.8	1.3	2.4	5.0	9.7	17.7	29.1	52.9	116.2
2000.....	0.4	0.7	1.5	2.6	4.8	9.2	17.3	31.1	52.9	118.0
1999.....	0.4	0.6	1.4	2.9	5.3	9.7	16.5	29.2	52.9	118.0

White, male

2006 4/.....	0.5	0.7	1.5	2.2	4.5	7.4	14.8	25.8	46.9	84.5
2005 4/.....	0.4	0.8	1.4	2.4	4.2	7.2	15.4	26.2	43.6	88.4
2004 4/.....	0.6	0.7	1.4	2.2	4.2	7.6	13.8	27.0	44.1	88.6
2003 4/.....	0.4	0.7	1.4	2.6	4.0	8.5	15.2	26.0	48.0	92.9
2002.....	0.5	0.7	1.3	2.3	4.2	7.8	14.9	26.7	48.7	97.5
2001.....	0.4	0.7	1.3	2.1	4.8	8.4	17.3	26.2	48.3	98.9
2000.....	0.3	0.6	1.4	2.4	4.5	8.2	16.2	29.0	47.0	98.9
1999.....	0.5	0.7	1.2	2.8	4.7	8.2	16.0	27.3	47.9	94.8

White, female

2006 4/.....	0.5	0.7	1.4	2.6	5.1	8.7	17.8	30.5	53.6	109.8
2005 4/.....	0.4	0.8	1.4	2.5	4.5	8.9	18.3	29.9	53.9	112.1
2004 4/.....	0.4	0.8	1.3	2.5	5.1	9.7	17.7	30.8	54.0	117.5
2003 4/.....	0.5	1.0	1.4	2.6	4.9	10.5	18.9	31.9	55.8	122.9
2002.....	0.4	0.8	1.2	2.9	4.9	9.5	17.7	32.0	55.6	123.9
2001.....	0.4	0.8	1.4	2.7	5.2	10.9	18.0	31.1	55.7	123.4
2000.....	0.5	0.7	1.5	2.8	5.1	9.9	18.2	32.5	56.4	125.8
1999.....	0.4	0.6	1.5	2.9	5.8	11.0	16.8	30.6	55.9	127.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Vascular disorders and obstruction of intestine without hernia (K55-K56)

Black, both sexes

2006 4/.....	1.0	1.4	3.2	5.3	6.7	11.5	18.8	34.5	55.9	91.6
2005 4/.....	0.8	1.6	2.8	4.5	8.0	12.8	22.0	36.2	52.2	102.4
2004 4/.....	1.1	1.6	2.5	5.0	7.4	10.5	20.8	37.7	56.7	106.0
2003 4/.....	1.0	1.5	3.5	5.4	8.2	12.9	22.4	37.6	54.3	102.1
2002.....	1.0	1.8	3.0	4.6	7.2	13.3	25.3	32.3	63.3	99.8
2001.....	0.8	1.8	3.9	5.1	10.2	15.4	24.0	40.1	53.6	96.7
2000.....	0.9	2.4	4.0	5.4	10.4	14.3	21.7	40.7	59.0	112.0
1999.....	1.4	1.8	3.2	4.3	7.4	15.9	26.1	42.3	54.9	108.7

Black, male

2006 4/.....	*	1.9	3.4	5.6	8.1	13.6	21.5	42.5	61.6	85.8
2005 4/.....	*	1.9	3.2	4.7	10.0	16.9	25.3	38.0	55.5	97.6
2004 4/.....	*	1.7	2.4	5.0	9.3	12.4	24.9	44.6	66.0	107.3
2003 4/.....	*	1.9	4.1	7.3	9.4	14.3	29.9	41.1	67.9	101.1
2002.....	*	1.9	3.5	4.8	9.5	12.5	27.4	30.3	71.8	105.1
2001.....	*	1.8	4.0	5.2	9.6	18.8	26.9	42.7	57.0	120.3
2000.....	*	3.0	4.1	6.9	13.0	16.4	23.4	45.8	70.8	110.7
1999.....	1.8	2.3	3.3	4.6	8.3	15.3	25.1	39.9	68.0	110.1

Black, female

2006 4/.....	*	*	3.1	5.0	5.6	10.0	16.9	29.7	52.8	94.0
2005 4/.....	*	*	2.4	4.4	6.5	9.7	19.7	35.2	50.4	104.3
2004 4/.....	1.3	1.5	2.6	5.0	6.0	9.2	18.0	33.7	51.8	105.5
2003 4/.....	*	*	3.0	3.8	7.2	11.9	17.3	35.5	47.2	102.5
2002.....	*	1.8	2.6	4.5	5.4	13.9	24.0	33.5	58.9	97.8
2001.....	*	1.8	3.8	5.1	10.7	13.0	22.0	38.6	51.9	88.0
2000.....	*	1.8	4.0	4.2	8.3	12.7	20.6	37.6	53.1	112.5
1999.....	*	*	3.0	4.2	6.7	16.4	26.8	43.8	48.4	108.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diverticular disease of intestine (K57)										
All races 3/, both sexes										
2006.....	*	0.2	0.3	0.4	0.9	1.9	3.2	6.4	11.7	24.1
2005.....	*	0.2	0.3	0.6	0.9	1.6	3.1	6.2	11.0	25.8
2004.....	0.1	*	0.3	0.5	1.1	1.8	3.8	6.5	13.0	27.0
2003.....	0.1	0.1	0.3	0.6	1.1	1.9	3.6	7.3	13.3	28.9
2002.....	*	0.2	0.3	0.5	0.9	1.9	3.5	7.3	13.4	27.7
2001.....	*	0.2	0.3	0.6	1.3	2.2	3.6	7.0	12.8	31.4
2000.....	0.1	0.2	0.3	0.4	1.1	2.2	3.7	6.5	13.3	30.9
1999.....	*	0.2	0.3	0.7	1.1	2.2	4.1	6.9	12.8	33.4
All races 3/, male										
2006.....	*	*	0.2	0.4	1.0	1.6	3.0	5.1	9.9	17.6
2005.....	*	0.2	0.3	0.6	0.7	1.4	2.8	4.9	9.2	19.1
2004.....	*	*	0.4	0.5	1.1	1.6	3.3	5.8	10.7	18.8
2003.....	*	*	0.3	0.5	1.1	1.7	3.2	5.7	10.3	22.3
2002.....	*	0.3	0.4	0.6	0.9	1.7	3.1	6.5	10.4	19.9
2001.....	*	0.3	0.3	0.6	1.1	1.6	3.2	6.6	10.7	23.4
2000.....	*	0.2	0.3	0.6	1.0	2.1	3.4	5.2	10.8	22.3
1999.....	*	*	0.2	0.6	1.2	1.9	3.7	5.5	10.4	25.8
All races 3/, female										
2006.....	*	*	0.3	0.4	0.9	2.1	3.4	7.4	12.9	27.2
2005.....	*	*	0.2	0.6	1.0	1.8	3.3	7.0	12.1	28.9
2004.....	*	*	0.2	0.5	1.1	2.0	4.3	7.0	14.4	30.6
2003.....	*	*	0.3	0.7	1.2	2.1	4.0	8.4	15.1	31.9
2002.....	*	*	*	0.5	1.0	2.0	3.8	7.9	15.2	31.0
2001.....	*	*	0.2	0.6	1.4	2.7	3.9	7.3	14.1	34.7
2000.....	*	*	0.3	0.3	1.2	2.4	4.0	7.4	14.8	34.4
1999.....	*	*	0.3	0.8	1.0	2.6	4.3	7.9	14.1	36.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diverticular disease of intestine (K57)										
White, both sexes										
2006 4/.....	*	0.2	0.3	0.4	1.0	1.8	3.3	6.5	12.2	25.2
2005 4/.....	*	0.2	0.3	0.5	0.9	1.6	3.2	6.3	11.3	27.1
2004 4/.....	*	*	0.3	0.6	1.1	1.8	4.0	6.8	13.4	27.9
2003 4/.....	*	0.1	0.3	0.5	1.1	2.0	3.7	7.5	13.7	30.3
2002.....	*	0.2	0.3	0.5	0.9	1.9	3.6	7.4	13.7	28.5
2001.....	*	0.2	0.3	0.6	1.3	2.0	3.7	7.2	13.5	32.4
2000.....	*	0.2	0.3	0.4	1.0	2.2	3.8	6.7	13.7	32.3
1999.....	*	0.2	0.3	0.7	1.1	2.2	4.2	7.2	13.1	34.6
White, male										
2006 4/.....	*	*	0.3	0.4	1.0	1.6	3.1	4.8	10.3	18.1
2005 4/.....	*	*	0.3	0.6	0.7	1.3	2.8	5.1	9.3	20.5
2004 4/.....	*	*	0.4	0.6	1.1	1.5	3.4	6.0	10.7	19.3
2003 4/.....	*	*	0.4	0.5	1.1	1.8	3.2	6.1	10.5	23.4
2002.....	*	0.3	0.5	0.5	0.9	1.7	3.3	6.3	10.3	20.5
2001.....	*	0.3	0.3	0.7	1.1	1.5	3.3	6.7	11.1	24.3
2000.....	*	*	0.3	0.6	0.8	2.1	3.4	5.4	11.3	23.7
1999.....	*	*	*	0.5	1.1	2.0	3.8	5.7	10.7	27.0
White, female										
2006 4/.....	*	*	0.3	0.4	0.9	2.1	3.5	7.8	13.4	28.5
2005 4/.....	*	*	*	0.5	1.0	1.9	3.5	7.2	12.6	30.1
2004 4/.....	*	*	*	0.6	1.1	2.0	4.5	7.4	15.1	31.8
2003 4/.....	*	*	0.3	0.6	1.1	2.2	4.1	8.6	15.7	33.3
2002.....	*	*	*	0.5	0.9	2.1	3.9	8.3	15.7	32.0
2001.....	*	*	*	0.6	1.4	2.5	4.0	7.6	15.0	35.8
2000.....	*	*	0.3	*	1.1	2.4	4.1	7.6	15.1	35.7
1999.....	*	*	0.3	0.8	1.0	2.5	4.4	8.3	14.6	37.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diverticular disease of intestine (K57)

Black, both sexes

2006 4/.....	*	*	*	*	*	2.5	3.4	6.5	9.7	17.1
2005 4/.....	*	*	*	1.2	*	*	2.6	6.3	10.0	17.9
2004 4/.....	*	*	*	*	*	2.5	3.7	4.6	11.1	21.7
2003 4/.....	*	*	*	1.3	1.7	*	3.7	6.0	10.9	18.4
2002.....	*	*	*	*	*	2.1	3.1	7.4	13.3	21.7
2001.....	*	*	*	*	*	3.8	4.2	6.6	7.0	24.3
2000.....	*	*	*	*	2.1	2.9	4.7	5.0	10.7	20.0
1999.....	*	*	*	*	*	3.0	4.1	5.2	10.7	23.6

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	9.6	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	*	*	*	*	*	*	*	6.2	10.9	19.4
2005 4/.....	*	*	*	*	*	*	*	7.5	10.4	21.8
2004 4/.....	*	*	*	*	*	*	4.4	*	10.2	23.2
2003 4/.....	*	*	*	*	*	*	*	7.6	10.8	20.3
2002.....	*	*	*	*	*	*	*	6.0	13.1	24.6
2001.....	*	*	*	*	*	4.7	4.7	6.1	*	27.8
2000.....	*	*	*	*	*	*	*	6.0	12.3	24.5
1999.....	*	*	*	*	*	4.2	*	*	11.7	27.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of intestines and peritoneum (K52,K58-K63)										
All races 3/, both sexes										
2006.....	0.3	0.4	0.9	1.1	2.2	3.3	6.0	10.4	17.5	36.3
2005.....	0.3	0.4	0.7	1.1	2.2	3.6	6.2	11.0	17.8	34.7
2004.....	0.2	0.4	0.7	1.0	1.9	3.2	5.7	9.4	17.3	33.3
2003.....	0.2	0.4	0.7	1.2	2.1	3.3	5.5	9.8	16.5	32.8
2002.....	0.3	0.4	0.7	1.2	2.1	3.5	5.2	9.9	16.2	32.8
2001.....	0.3	0.4	0.7	1.1	1.9	3.5	5.6	10.1	16.8	34.1
2000.....	0.3	0.5	0.6	1.2	2.0	3.0	5.3	9.5	16.9	33.3
1999.....	0.3	0.4	0.7	1.2	1.9	3.4	5.6	9.7	15.7	32.0
All races 3/, male										
2006.....	0.2	0.5	0.9	1.3	2.0	3.6	6.0	9.9	16.6	28.7
2005.....	0.3	0.4	0.8	1.1	2.0	3.5	6.4	10.3	17.0	28.4
2004.....	*	0.4	0.7	1.0	1.5	2.9	5.5	9.8	16.5	26.4
2003.....	*	0.4	0.8	1.2	2.2	3.0	5.5	10.1	15.7	27.1
2002.....	0.3	0.5	0.7	1.2	2.0	3.4	5.5	10.5	13.8	29.4
2001.....	0.2	0.4	0.7	1.0	1.9	3.2	5.4	9.9	15.6	29.9
2000.....	0.3	0.4	0.5	1.3	2.1	2.7	5.6	9.1	15.1	28.7
1999.....	0.3	0.6	0.7	1.2	1.8	3.1	5.5	9.8	15.5	28.9
All races 3/, female										
2006.....	0.3	0.4	0.9	1.0	2.4	3.1	6.1	10.7	18.1	39.8
2005.....	0.2	0.5	0.7	1.1	2.4	3.6	6.0	11.6	18.3	37.6
2004.....	0.3	0.3	0.7	1.1	2.2	3.4	6.0	9.1	17.8	36.4
2003.....	0.3	0.3	0.7	1.1	2.0	3.6	5.4	9.6	17.0	35.2
2002.....	0.3	0.4	0.7	1.2	2.3	3.5	5.1	9.4	17.7	34.3
2001.....	0.3	0.5	0.8	1.1	2.0	3.7	5.7	10.2	17.5	35.8
2000.....	0.2	0.5	0.8	1.1	1.8	3.3	5.1	9.7	17.9	35.2
1999.....	0.3	0.3	0.6	1.2	2.1	3.6	5.7	9.7	15.8	33.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of intestines and peritoneum (K52,K58-K63)

White, both sexes

2006 4/.....	0.3	0.4	0.8	1.1	2.3	3.4	6.2	10.5	18.2	37.5
2005 4/.....	0.3	0.4	0.7	1.0	2.1	3.5	6.3	11.2	18.1	35.7
2004 4/.....	0.2	0.3	0.7	1.0	1.8	3.1	5.8	9.6	17.8	34.4
2003 4/.....	0.2	0.4	0.7	1.1	2.0	3.2	5.4	10.0	17.0	33.8
2002.....	0.3	0.4	0.7	1.2	2.1	3.6	5.1	10.1	16.7	34.1
2001.....	0.3	0.4	0.7	0.9	1.8	3.5	5.7	10.0	16.8	35.2
2000.....	0.2	0.5	0.6	1.2	1.8	3.0	5.4	9.6	17.2	33.7
1999.....	0.3	0.4	0.6	1.2	1.9	3.3	5.6	9.8	15.9	33.0

White, male

2006 4/.....	*	0.5	0.8	1.1	1.9	3.7	6.2	9.8	17.2	29.5
2005 4/.....	0.3	0.4	0.7	1.0	1.8	3.5	6.4	10.2	17.2	28.2
2004 4/.....	*	0.4	0.7	0.9	1.5	2.6	5.5	9.8	16.8	26.9
2003 4/.....	*	0.4	0.7	1.1	2.1	3.0	5.4	10.1	15.9	27.5
2002.....	0.3	0.5	0.7	1.2	1.9	3.4	5.1	10.7	13.7	30.3
2001.....	0.2	0.4	0.7	0.9	1.7	3.0	5.1	9.8	15.7	31.2
2000.....	0.3	0.4	0.4	1.2	1.8	2.7	5.7	9.1	15.2	28.1
1999.....	0.3	0.5	0.7	1.2	1.7	3.1	5.6	9.6	15.7	28.3

White, female

2006 4/.....	0.3	0.3	0.8	1.1	2.6	3.1	6.2	11.1	18.8	41.2
2005 4/.....	*	0.4	0.7	1.1	2.4	3.5	6.2	12.0	18.6	39.1
2004 4/.....	0.2	0.3	0.7	1.1	2.1	3.5	6.1	9.4	18.4	37.8
2003 4/.....	0.3	0.3	0.6	1.2	1.8	3.4	5.5	9.8	17.7	36.5
2002.....	0.3	0.4	0.7	1.2	2.2	3.8	5.1	9.7	18.5	35.8
2001.....	0.3	0.4	0.8	1.0	1.9	3.9	6.1	10.2	17.4	36.8
2000.....	*	0.5	0.7	1.1	1.8	3.2	5.1	10.0	18.4	36.1
1999.....	0.2	0.3	0.6	1.2	2.1	3.6	5.7	9.9	16.0	34.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of intestines and peritoneum (K52,K58-K63)

Black, both sexes

2006 4/.....	*	0.9	1.7	2.0	2.9	3.6	5.8	8.9	12.5	28.0
2005 4/.....	*	0.8	1.0	1.4	3.5	4.6	5.9	10.7	18.4	29.4
2004 4/.....	*	*	*	1.5	3.1	4.3	6.5	9.1	14.9	24.0
2003 4/.....	*	*	1.5	1.8	3.5	5.5	6.6	10.4	14.7	25.7
2002.....	*	*	1.0	1.7	3.3	3.3	7.8	8.9	13.9	23.2
2001.....	*	0.8	1.1	2.3	2.7	4.1	6.1	13.0	19.8	27.0
2000.....	*	*	1.1	1.8	3.5	3.4	6.1	8.0	15.0	29.7
1999.....	*	0.9	1.2	1.5	2.3	4.3	6.4	8.9	16.2	23.6

Black, male

2006 4/.....	*	*	*	2.9	3.7	*	*	9.8	*	24.1
2005 4/.....	*	*	*	*	4.0	*	7.3	13.4	19.0	34.4
2004 4/.....	*	*	*	*	*	6.1	7.4	11.8	15.8	24.8
2003 4/.....	*	*	*	*	*	4.9	7.9	11.9	16.2	25.0
2002.....	*	*	*	*	*	5.0	10.7	10.6	16.0	24.7
2001.....	*	*	*	*	*	5.8	8.8	15.8	16.9	*
2000.....	*	*	*	*	4.1	*	*	10.9	17.7	30.0
1999.....	*	*	*	*	*	*	*	10.4	17.4	39.8

Black, female

2006 4/.....	*	*	1.7	*	*	*	6.6	8.3	12.7	29.6
2005 4/.....	*	*	*	*	3.1	4.6	5.0	9.1	18.2	27.4
2004 4/.....	*	*	*	*	3.5	*	5.9	7.5	14.4	23.6
2003 4/.....	*	*	*	*	3.5	5.8	5.8	9.5	13.9	25.9
2002.....	*	*	*	*	3.4	*	5.8	8.0	12.7	22.6
2001.....	*	*	*	*	*	*	*	11.4	21.3	29.5
2000.....	*	*	*	*	*	4.0	6.4	6.3	13.6	29.6
1999.....	*	*	*	*	*	4.2	7.2	8.0	15.6	17.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of peritoneum (K65-K66)										
All races 3/, both sexes										
2006.....	0.2	0.3	0.5	0.6	0.8	1.6	2.1	2.9	4.8	7.9
2005.....	0.1	0.4	0.5	0.7	1.1	1.5	2.1	3.3	5.0	8.3
2004.....	0.2	0.4	0.4	0.6	0.9	1.4	2.1	3.4	5.1	8.6
2003.....	0.2	0.4	0.6	0.6	1.1	1.7	2.2	3.4	5.1	9.2
2002.....	0.2	0.4	0.6	0.7	1.0	1.3	2.3	3.7	6.2	10.3
2001.....	0.2	0.4	0.5	0.8	0.9	1.4	2.3	3.6	5.7	10.4
2000.....	0.3	0.4	0.5	0.9	1.2	1.4	2.7	3.9	5.3	10.6
1999.....	0.3	0.4	0.5	0.8	1.1	1.6	2.8	3.7	5.5	9.7
All races 3/, male										
2006.....	0.2	0.3	0.5	0.7	0.9	1.8	1.9	2.5	5.0	7.2
2005.....	*	0.4	0.7	0.7	1.2	1.6	2.0	3.6	5.7	6.1
2004.....	0.2	0.4	0.6	0.6	0.9	1.7	2.3	3.4	5.9	7.7
2003.....	0.3	0.3	0.6	0.7	1.2	1.7	2.3	3.7	5.7	8.7
2002.....	0.2	0.4	0.7	0.8	1.1	1.3	2.0	3.7	5.1	9.1
2001.....	0.3	0.5	0.6	0.9	1.0	1.7	2.2	3.6	6.2	10.1
2000.....	0.3	0.4	0.5	1.0	1.4	1.5	2.7	4.3	5.2	11.0
1999.....	0.3	0.5	0.7	0.8	1.3	1.7	3.1	4.1	6.3	10.1
All races 3/, female										
2006.....	0.2	0.3	0.5	0.5	0.7	1.3	2.2	3.2	4.7	8.2
2005.....	*	0.3	0.4	0.7	1.1	1.4	2.1	3.1	4.6	9.3
2004.....	0.2	0.3	0.3	0.6	0.8	1.2	1.9	3.4	4.7	9.0
2003.....	*	0.4	0.5	0.5	0.9	1.6	2.2	3.3	4.6	9.4
2002.....	0.2	0.3	0.5	0.6	1.0	1.3	2.5	3.7	6.8	10.8
2001.....	0.2	0.4	0.5	0.6	0.8	1.1	2.3	3.5	5.4	10.5
2000.....	0.3	0.3	0.5	0.9	1.0	1.4	2.6	3.6	5.3	10.5
1999.....	0.2	0.2	0.4	0.8	0.9	1.5	2.5	3.5	5.1	9.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of peritoneum (K65-K66)										
White, both sexes										
2006 4/.....	0.2	0.3	0.4	0.5	0.8	1.6	2.0	3.0	5.0	8.1
2005 4/.....	0.1	0.3	0.5	0.6	1.0	1.5	1.9	3.3	5.0	8.5
2004 4/.....	0.2	0.3	0.4	0.6	0.8	1.4	2.0	3.5	5.0	8.8
2003 4/.....	0.2	0.3	0.5	0.5	0.9	1.6	2.1	3.5	5.1	9.1
2002.....	0.2	0.3	0.5	0.7	0.9	1.2	2.2	3.8	6.1	10.5
2001.....	0.2	0.4	0.5	0.6	0.7	1.3	2.2	3.5	5.6	10.4
2000.....	0.2	0.3	0.5	0.8	1.1	1.3	2.6	3.8	5.2	10.6
1999.....	0.2	0.4	0.5	0.7	1.0	1.4	2.7	3.5	5.6	9.6
White, male										
2006 4/.....	*	0.3	0.5	0.6	0.9	1.7	1.7	2.6	5.4	7.4
2005 4/.....	*	0.3	0.6	0.6	1.0	1.6	2.0	3.6	5.9	6.0
2004 4/.....	*	0.4	0.5	0.6	0.9	1.6	2.2	3.4	5.9	7.9
2003 4/.....	0.3	0.3	0.5	0.6	1.0	1.6	2.3	3.8	5.7	8.8
2002.....	*	0.3	0.6	0.8	0.9	1.3	1.9	3.6	5.0	9.3
2001.....	*	0.5	0.5	0.7	0.9	1.7	2.1	3.5	6.2	9.7
2000.....	0.2	0.3	0.4	0.9	1.3	1.3	2.5	4.2	5.0	10.8
1999.....	0.3	0.5	0.5	0.7	1.1	1.5	3.0	3.9	6.3	10.2
White, female										
2006 4/.....	*	0.3	0.4	0.4	0.7	1.4	2.2	3.3	4.8	8.5
2005 4/.....	*	0.3	0.3	0.6	1.0	1.5	1.8	3.0	4.5	9.6
2004 4/.....	*	0.2	0.3	0.6	0.8	1.2	1.8	3.5	4.5	9.2
2003 4/.....	*	0.2	0.4	0.4	0.8	1.6	2.0	3.3	4.7	9.3
2002.....	0.2	0.3	0.4	0.6	0.9	1.1	2.4	3.9	6.8	11.0
2001.....	*	0.3	0.4	0.5	0.6	1.0	2.2	3.4	5.2	10.7
2000.....	0.2	*	0.5	0.7	1.0	1.3	2.6	3.5	5.3	10.5
1999.....	*	*	0.4	0.6	0.9	1.4	2.3	3.3	5.2	9.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of peritoneum (K65-K66)

Black, both sexes

2006 4/.....	*	*	*	1.3	*	2.0	3.0	3.4	*	6.0
2005 4/.....	*	*	1.0	1.1	2.4	2.0	4.1	3.8	6.1	7.5
2004 4/.....	*	*	*	*	*	2.1	3.5	*	7.9	7.9
2003 4/.....	*	0.9	1.2	1.5	2.5	*	3.6	3.6	5.6	12.0
2002.....	*	1.0	1.1	*	2.2	2.1	3.6	4.1	7.3	10.4
2001.....	*	1.1	1.1	1.8	2.3	2.6	3.2	4.9	7.9	11.7
2000.....	*	1.0	1.2	2.2	2.1	2.9	3.3	5.3	7.9	11.6
1999.....	*	*	1.5	2.3	*	3.1	4.7	5.9	6.1	12.1

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	4.8	*	*	7.5
2004 4/.....	*	*	*	*	*	*	*	*	7.6	7.9
2003 4/.....	*	*	*	*	*	*	*	*	*	12.6
2002.....	*	*	*	*	*	*	*	*	8.0	11.1
2001.....	*	*	*	*	*	*	*	*	8.2	10.5
2000.....	*	*	*	*	*	*	*	*	*	10.7
1999.....	*	*	*	*	*	*	4.5	*	*	12.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of liver (K70-K76)										
All races 3/, both sexes										
2006.....	10.4	18.7	27.4	30.2	31.2	33.3	37.6	39.8	40.6	30.6
2005.....	10.8	19.0	26.7	30.0	31.6	34.3	39.4	40.4	40.5	32.3
2004.....	11.1	19.3	27.1	28.2	31.1	34.5	39.6	41.4	38.2	32.8
2003.....	11.9	20.2	27.1	28.5	32.1	37.3	40.4	41.5	40.2	33.6
2002.....	12.3	20.7	25.7	27.5	31.2	36.0	40.3	41.9	41.9	35.7
2001.....	11.9	21.1	25.0	26.4	31.7	36.6	39.7	42.2	40.2	36.1
2000.....	12.4	20.4	23.3	26.9	32.3	36.3	39.7	41.5	40.7	36.7
1999.....	12.1	19.9	23.0	26.6	32.9	38.0	40.9	42.4	41.9	37.2
All races 3/, male										
2006.....	13.3	25.5	40.3	43.9	43.9	43.9	46.8	49.2	48.2	39.2
2005.....	14.0	26.7	39.8	43.9	45.0	44.3	49.8	49.5	50.3	43.1
2004.....	15.0	27.0	40.5	41.0	42.6	45.3	50.3	51.3	46.8	40.7
2003.....	15.8	29.2	40.8	40.8	45.7	49.6	51.2	50.0	49.1	44.4
2002.....	16.3	29.8	38.7	39.7	43.1	47.8	51.6	51.0	52.8	47.3
2001.....	16.0	30.6	37.2	38.3	44.4	49.4	51.0	51.2	50.4	49.4
2000.....	17.2	30.5	34.8	38.2	45.2	48.4	51.1	52.8	48.1	52.2
1999.....	17.4	30.2	34.9	39.4	45.5	52.4	52.6	51.1	53.5	52.3
All races 3/, female										
2006.....	7.4	12.0	15.1	17.3	19.6	24.0	30.1	32.8	35.9	26.7
2005.....	7.6	11.4	14.2	16.8	19.3	25.5	31.0	33.8	34.4	27.4
2004.....	7.3	11.9	14.3	16.1	20.6	25.1	31.0	34.2	33.0	29.3
2003.....	8.1	11.4	14.1	16.8	19.8	26.6	31.7	35.5	34.8	28.9
2002.....	8.4	11.8	13.3	15.9	20.4	25.8	31.3	35.5	35.2	30.7
2001.....	7.9	11.8	13.3	15.3	20.3	25.5	30.7	35.9	34.1	30.5
2000.....	7.7	10.6	12.4	16.3	20.7	25.9	30.8	33.6	36.4	30.4
1999.....	6.9	10.0	11.6	14.6	21.5	25.8	31.9	36.3	35.0	31.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of liver (K70-K76)										
White, both sexes										
2006 4/.....	10.7	19.3	28.0	30.5	31.6	34.3	39.0	41.1	42.1	31.6
2005 4/.....	11.2	19.2	26.8	30.3	31.7	35.2	40.9	41.6	41.0	32.8
2004 4/.....	11.3	19.6	27.2	27.7	31.1	35.3	41.0	43.2	39.3	33.4
2003 4/.....	12.2	20.4	26.7	28.2	32.1	38.1	41.8	43.4	41.7	34.3
2002.....	12.6	20.5	25.3	27.2	31.4	37.0	41.9	43.6	42.9	36.6
2001.....	11.8	20.9	24.2	25.6	32.1	37.4	40.6	43.4	41.4	37.0
2000.....	12.3	20.2	22.3	26.2	32.3	36.8	40.7	42.9	41.6	37.5
1999.....	11.7	19.0	22.5	25.6	32.3	38.4	41.6	43.4	43.2	37.7
White, male										
2006 4/.....	13.9	26.5	41.0	44.0	44.5	44.8	48.4	50.6	50.3	40.7
2005 4/.....	14.6	27.0	40.0	44.2	45.0	45.1	51.8	50.6	50.7	44.1
2004 4/.....	15.3	27.2	40.8	40.1	42.5	46.0	51.7	53.2	48.3	42.6
2003 4/.....	16.5	29.7	40.0	40.0	45.6	50.4	52.7	52.1	50.6	45.1
2002.....	16.9	30.0	38.2	39.3	43.2	48.8	53.3	53.0	53.9	48.5
2001.....	15.8	30.6	36.1	36.9	44.5	50.2	52.1	52.3	51.8	50.8
2000.....	17.2	30.7	33.3	37.4	45.3	49.2	52.0	54.3	49.8	53.3
1999.....	17.1	29.1	34.1	38.0	45.2	52.7	53.4	52.3	55.7	53.8
White, female										
2006 4/.....	7.5	12.2	15.2	17.4	19.6	24.9	31.1	34.0	37.0	27.4
2005 4/.....	7.7	11.4	13.8	16.9	19.3	26.3	32.0	34.9	35.0	27.6
2004 4/.....	7.2	12.0	13.8	15.8	20.5	25.8	32.2	35.9	33.7	29.3
2003 4/.....	7.9	11.1	13.7	16.8	19.7	27.3	32.9	37.1	36.3	29.5
2002.....	8.3	11.1	12.8	15.6	20.4	26.6	32.5	36.9	36.1	31.4
2001.....	7.8	11.3	12.6	14.9	20.7	26.2	31.4	37.2	35.1	31.2
2000.....	7.3	9.8	11.6	15.6	20.3	26.0	31.5	34.8	36.8	31.1
1999.....	6.4	9.1	11.1	13.8	20.5	26.0	32.3	37.1	35.8	31.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of liver (K70-K76)										
Black, both sexes										
2006 4/.....	8.7	15.8	29.0	33.7	31.4	30.0	28.8	28.3	23.0	21.3
2005 4/.....	8.9	20.0	30.3	34.2	33.8	31.4	30.1	28.9	34.5	24.3
2004 4/.....	10.7	21.0	31.6	35.8	35.0	32.9	31.7	23.9	27.2	24.3
2003 4/.....	10.7	20.9	35.3	35.6	35.8	36.7	32.4	24.5	23.6	27.1
2002.....	11.2	24.4	33.4	34.0	33.9	32.6	30.4	30.1	28.8	27.6
2001.....	13.6	25.5	34.4	37.6	33.0	36.0	34.0	33.0	29.0	21.8
2000.....	14.2	25.2	34.8	34.5	36.0	36.7	35.6	29.1	28.2	26.0
1999.....	15.6	29.0	30.9	36.9	41.6	38.1	36.5	34.5	28.0	31.2
Black, male										
2006 4/.....	11.4	21.1	42.2	52.1	44.7	46.2	37.8	42.1	26.0	24.1
2005 4/.....	11.0	28.8	44.5	51.6	51.6	45.2	42.4	41.6	43.5	26.0
2004 4/.....	14.0	29.9	46.4	56.5	50.8	47.5	46.6	33.2	33.7	19.9
2003 4/.....	12.4	29.1	55.5	56.0	54.6	53.7	44.7	31.0	34.7	40.7
2002.....	13.4	32.7	50.1	51.5	49.9	47.1	41.7	36.7	39.0	36.5
2001.....	18.6	36.9	53.4	56.8	50.6	53.4	48.1	44.1	37.7	27.2
2000.....	18.5	34.6	53.4	50.7	50.5	51.0	50.5	41.1	29.5	35.8
1999.....	21.2	43.5	47.7	55.8	56.4	55.8	50.5	42.3	33.1	37.5
Black, female										
2006 4/.....	6.3	11.1	17.9	18.6	21.1	18.0	22.6	20.1	21.3	20.1
2005 4/.....	7.0	12.4	18.2	19.9	20.0	21.2	21.6	21.3	29.7	23.6
2004 4/.....	7.9	13.3	19.0	18.8	22.6	22.1	21.5	18.3	23.8	26.0
2003 4/.....	9.2	13.7	18.3	18.9	21.2	24.1	24.1	20.6	17.8	21.9
2002.....	9.4	17.1	19.3	19.6	21.4	21.8	22.9	26.1	23.5	24.2
2001.....	9.2	15.7	18.3	21.8	19.0	23.1	24.7	26.3	24.5	19.8
2000.....	10.4	17.0	18.8	21.2	24.5	26.1	25.8	21.9	27.6	22.3
1999.....	10.6	16.4	16.7	21.5	30.0	25.0	27.2	29.8	25.5	28.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
All races 3/, both sexes										
2006.....	5.5	9.2	12.0	12.5	11.6	10.1	8.6	7.0	4.5	2.2
2005.....	5.7	9.3	12.0	12.4	12.0	10.3	9.4	6.4	4.8	1.8
2004.....	5.8	9.1	12.2	11.8	11.5	10.8	9.0	6.5	4.1	2.1
2003.....	5.9	9.4	11.8	11.6	11.6	11.0	8.3	6.6	4.4	1.8
2002.....	5.9	9.4	11.4	11.4	11.8	10.7	9.5	6.7	4.4	2.1
2001.....	6.1	10.2	11.3	11.1	12.0	11.1	9.0	6.9	3.9	2.4
2000.....	6.6	10.0	10.7	11.7	12.9	11.4	9.1	7.2	4.4	2.2
1999.....	6.4	10.0	10.9	11.5	12.6	11.6	9.4	7.3	5.2	2.8
All races 3/, male										
2006.....	7.1	12.9	18.4	19.4	18.1	16.9	14.1	12.0	7.5	4.7
2005.....	7.5	13.2	18.5	19.5	19.4	16.5	15.0	11.2	8.7	4.2
2004.....	8.0	12.9	18.6	18.2	18.3	17.8	15.0	11.5	8.0	4.6
2003.....	8.0	14.2	18.3	18.1	18.6	17.7	14.2	11.3	8.2	3.6
2002.....	8.0	13.8	17.7	17.6	19.2	17.4	15.4	10.9	8.0	4.7
2001.....	8.4	15.3	17.0	18.0	19.0	17.9	15.2	12.1	7.0	5.5
2000.....	9.4	15.3	16.9	18.0	20.3	18.7	15.7	13.8	7.7	5.3
1999.....	9.4	15.5	17.2	18.7	20.0	19.7	16.2	12.1	9.7	6.1
All races 3/, female										
2006.....	3.8	5.6	5.9	6.0	5.6	4.2	4.1	3.4	2.6	1.1
2005.....	3.9	5.4	5.8	5.7	5.3	4.9	4.8	3.0	2.4	0.7
2004.....	3.6	5.5	6.0	5.7	5.4	4.8	4.1	2.9	1.7	1.0
2003.....	3.8	4.8	5.5	5.5	5.3	5.3	3.6	3.3	2.1	1.0
2002.....	3.9	5.1	5.4	5.6	5.1	4.9	4.8	3.7	2.2	1.0
2001.....	3.9	5.3	5.8	4.7	5.6	5.3	4.0	3.3	2.1	1.0
2000.....	3.9	4.8	4.8	5.8	6.2	5.1	3.9	2.6	2.4	1.0
1999.....	3.5	4.6	4.8	4.9	5.9	4.6	4.0	3.9	2.7	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
White, both sexes										
2006 4/.....	5.8	9.7	12.6	13.0	12.1	10.4	8.9	7.3	4.7	2.4
2005 4/.....	5.9	9.6	12.5	12.9	12.1	10.7	9.7	6.6	4.9	1.9
2004 4/.....	6.1	9.5	12.6	11.8	11.7	11.1	9.3	6.8	4.1	2.2
2003 4/.....	6.2	9.7	11.8	11.8	11.6	11.2	8.6	7.0	4.5	1.9
2002.....	6.2	9.6	11.5	11.4	12.0	11.0	9.8	6.8	4.5	2.1
2001.....	6.2	10.5	11.0	11.0	12.2	11.2	9.2	7.1	4.0	2.3
2000.....	6.6	10.0	10.6	11.5	13.0	11.6	9.3	7.3	4.3	2.3
1999.....	6.2	9.8	10.8	11.2	12.4	11.5	9.4	7.2	5.3	2.7
White, male										
2006 4/.....	7.6	13.5	19.3	20.0	19.0	17.1	14.7	12.4	7.9	4.9
2005 4/.....	7.8	13.7	19.2	20.3	19.5	17.0	15.4	11.3	8.9	4.4
2004 4/.....	8.5	13.3	19.4	18.1	18.4	18.1	15.3	12.0	8.0	4.8
2003 4/.....	8.4	14.6	18.4	18.3	18.7	17.9	14.6	11.8	8.2	3.7
2002.....	8.3	14.3	17.9	17.9	19.4	17.7	15.8	11.1	8.2	4.5
2001.....	8.5	15.9	16.8	17.6	19.4	17.8	15.4	12.5	7.2	5.4
2000.....	9.4	15.6	16.8	18.0	20.6	19.0	15.9	14.1	7.8	5.4
1999.....	9.2	15.4	17.1	18.2	20.0	19.5	16.2	11.9	9.8	5.8
White, female										
2006 4/.....	3.9	5.9	6.1	6.2	5.7	4.4	4.1	3.5	2.6	1.1
2005 4/.....	3.9	5.6	6.0	5.8	5.2	5.1	5.0	3.1	2.4	0.8
2004 4/.....	3.7	5.7	5.9	5.7	5.4	4.9	4.4	3.0	1.7	1.0
2003 4/.....	3.9	4.9	5.4	5.5	5.1	5.3	3.7	3.5	2.2	1.1
2002.....	4.0	5.0	5.2	5.3	5.1	5.0	4.9	3.7	2.2	1.1
2001.....	3.9	5.2	5.4	4.6	5.7	5.4	4.2	3.3	2.1	1.0
2000.....	3.8	4.4	4.5	5.3	6.0	5.1	4.0	2.6	2.3	1.0
1999.....	3.3	4.3	4.5	4.7	5.5	4.6	4.0	3.9	2.6	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Alcoholic liver disease (K70)										
Black, both sexes										
2006 4/.....	3.5	6.0	9.4	10.0	8.6	8.9	6.5	5.2	*	*
2005 4/.....	3.9	7.8	10.1	11.0	11.9	9.0	8.4	5.3	*	*
2004 4/.....	4.1	7.5	10.8	12.9	12.0	10.8	6.5	*	*	*
2003 4/.....	4.2	7.6	12.7	12.2	12.9	11.2	6.9	*	*	*
2002.....	4.2	8.2	12.1	12.5	12.0	9.6	7.4	6.2	*	*
2001.....	5.9	9.0	14.7	14.1	11.5	13.0	7.8	5.7	*	*
2000.....	6.9	11.0	13.3	13.9	12.6	11.8	8.4	6.2	*	*
1999.....	7.2	12.8	12.8	14.8	15.6	12.9	10.4	8.0	*	*
Black, male										
2006 4/.....	4.5	8.8	14.1	16.3	12.7	16.9	10.2	9.8	*	*
2005 4/.....	4.6	11.7	15.0	16.8	20.5	15.4	15.2	12.1	*	*
2004 4/.....	5.6	10.7	15.6	20.8	19.4	19.4	13.6	*	*	*
2003 4/.....	5.4	12.0	20.6	19.5	20.8	18.7	12.5	*	*	*
2002.....	5.6	11.5	18.7	17.4	20.4	16.5	13.0	10.6	*	*
2001.....	8.2	12.9	22.3	23.5	18.7	22.9	15.1	*	*	*
2000.....	9.6	15.0	20.6	20.4	18.0	19.3	16.7	11.3	*	*
1999.....	10.1	19.2	20.2	24.9	23.3	24.1	18.7	14.2	*	*
Black, female										
2006 4/.....	2.6	3.5	5.5	4.7	5.4	*	*	*	*	*
2005 4/.....	3.3	4.3	5.9	6.2	5.1	4.2	*	*	*	*
2004 4/.....	2.8	4.6	6.8	6.3	6.1	4.5	*	*	*	*
2003 4/.....	3.0	3.7	6.0	6.1	6.8	5.7	*	*	*	*
2002.....	3.0	5.3	6.6	8.4	5.4	4.4	*	*	*	*
2001.....	3.9	5.5	8.2	6.4	5.8	5.6	*	*	*	*
2000.....	4.5	7.5	7.1	8.5	8.3	6.3	*	*	*	*
1999.....	4.7	7.2	6.6	6.5	9.5	4.6	5.0	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic hepatitis, not elsewhere classified (K73)

All races 3/, both sexes

2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	*
2004.....	*	0.1	*	*	*	*	*	*	*	*
2003.....	*	*	0.1	*	*	0.2	*	*	*	*
2002.....	0.1	0.2	0.2	0.2	0.4	0.7	0.8	1.0	1.0	0.8
2001.....	*	0.2	0.2	0.3	0.3	0.6	0.8	0.8	0.8	0.7
2000.....	0.1	0.1	0.2	0.3	0.3	0.7	0.6	0.9	0.8	0.8
1999.....	*	0.2	0.2	0.3	0.4	0.4	0.7	0.8	0.9	0.6

All races 3/, male

2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	0.2	*	0.4	0.6	1.0	1.1	1.3	1.3	0.9
2001.....	*	0.2	0.2	0.4	0.5	0.9	1.0	1.1	1.0	0.7
2000.....	*	*	*	0.4	0.6	1.1	0.9	1.2	1.0	0.9
1999.....	*	*	*	0.4	0.6	0.6	0.9	1.0	1.2	0.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic hepatitis, not elsewhere classified (K73)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	0.2	0.2	0.2	0.4	0.7	0.9	1.1	0.9	0.8
2001.....	*	0.2	0.2	0.3	0.3	0.6	0.7	0.8	0.9	0.8
2000.....	*	*	0.2	0.3	0.4	0.7	0.6	0.9	0.8	0.9
1999.....	*	0.2	0.2	0.3	0.4	0.4	0.7	0.8	0.9	0.6

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	0.4	0.6	1.0	1.1	1.5	1.2	0.9
2001.....	*	*	*	0.5	0.6	0.9	0.9	1.1	1.1	0.8
2000.....	*	*	*	0.4	0.6	1.0	1.0	1.3	1.0	0.9
1999.....	*	*	*	*	0.5	0.6	0.9	1.0	1.2	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fibrosis and cirrhosis of liver (K74)										
All races 3/, both sexes										
2006.....	2.7	5.5	9.1	10.2	11.3	14.6	18.8	22.4	23.6	17.0
2005.....	2.9	5.7	8.6	10.6	12.0	14.9	19.8	23.2	23.0	17.7
2004.....	2.9	5.9	9.0	9.9	12.1	15.3	20.4	23.5	22.7	17.4
2003.....	3.3	6.4	9.2	10.3	12.6	16.8	22.7	24.3	23.9	18.0
2002.....	3.6	6.4	8.7	10.1	12.2	16.5	20.9	24.8	24.5	18.5
2001.....	3.6	6.7	8.6	9.7	12.4	17.2	21.4	23.5	23.9	19.1
2000.....	3.5	6.7	7.7	9.9	12.9	16.8	21.1	23.9	24.4	20.0
1999.....	3.4	6.3	7.5	9.6	13.5	17.7	21.6	24.6	24.6	19.8
All races 3/, male										
2006.....	3.6	7.7	13.3	14.2	14.9	17.1	21.3	24.7	26.9	21.2
2005.....	4.0	8.0	12.7	14.9	15.8	17.3	23.0	25.9	26.4	23.2
2004.....	4.1	8.5	13.2	14.0	15.1	17.8	23.5	27.0	25.9	20.9
2003.....	4.7	9.1	13.8	14.3	16.9	20.5	25.6	26.7	26.7	23.7
2002.....	4.8	9.3	13.1	14.3	15.1	19.6	24.1	27.9	28.5	23.2
2001.....	4.7	9.8	12.9	13.3	16.2	21.5	25.2	25.2	27.8	25.7
2000.....	4.9	10.1	11.5	13.6	17.3	20.4	24.8	26.6	26.4	28.4
1999.....	5.1	9.6	11.3	13.6	17.0	21.9	24.9	27.5	28.5	26.6
All races 3/, female										
2006.....	1.8	3.3	5.1	6.4	8.0	12.4	16.7	20.6	21.5	15.0
2005.....	1.8	3.5	4.7	6.5	8.5	12.8	17.3	21.2	20.9	15.2
2004.....	1.8	3.4	5.0	6.0	9.4	13.1	17.9	20.9	20.8	15.8
2003.....	2.0	3.8	4.8	6.6	8.8	13.5	20.4	22.6	22.2	15.5
2002.....	2.5	3.6	4.5	6.1	9.6	13.9	18.3	22.6	22.1	16.5
2001.....	2.4	3.7	4.5	6.3	9.1	13.4	18.5	22.4	21.6	16.4
2000.....	2.1	3.5	4.1	6.5	8.9	13.7	18.2	21.9	23.2	16.6
1999.....	1.8	3.0	3.8	5.8	10.2	14.1	19.0	22.5	22.4	17.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fibrosis and cirrhosis of liver (K74)										
White, both sexes										
2006 4/.....	2.7	5.7	9.2	10.2	11.5	15.2	20.0	23.3	24.8	17.7
2005 4/.....	3.1	5.7	8.5	10.5	12.0	15.7	21.0	24.1	23.8	18.2
2004 4/.....	2.9	6.0	8.9	9.8	12.3	15.9	21.6	24.8	23.8	17.9
2003 4/.....	3.4	6.4	9.1	10.1	12.9	17.5	24.0	25.8	25.1	18.8
2002.....	3.8	6.3	8.5	10.1	12.4	17.3	22.0	26.2	25.5	19.4
2001.....	3.5	6.5	8.3	9.4	12.7	17.9	22.4	24.7	25.1	20.1
2000.....	3.5	6.8	7.4	9.9	13.1	17.2	21.9	25.2	25.4	20.8
1999.....	3.4	5.9	7.4	9.3	13.5	18.3	22.5	25.7	25.8	20.4
White, male										
2006 4/.....	3.6	8.1	13.2	14.2	14.9	17.7	22.6	25.4	28.4	22.0
2005 4/.....	4.2	8.0	12.6	14.6	15.6	18.0	24.3	26.6	26.9	24.0
2004 4/.....	4.1	8.5	13.0	13.8	15.2	18.2	24.7	28.4	27.2	21.9
2003 4/.....	4.9	9.4	13.5	13.7	17.1	21.3	27.0	28.1	27.8	24.5
2002.....	5.1	9.3	12.8	14.2	15.1	20.2	25.3	29.8	29.5	24.4
2001.....	4.6	9.5	12.3	12.8	16.2	22.2	26.2	26.2	29.5	27.3
2000.....	5.0	10.2	10.9	13.4	17.5	21.0	25.7	27.8	27.7	29.4
1999.....	5.0	9.1	11.1	13.2	17.1	22.6	25.9	28.7	30.0	27.9
White, female										
2006 4/.....	1.8	3.3	5.2	6.5	8.2	13.0	17.9	21.8	22.5	15.6
2005 4/.....	2.0	3.5	4.3	6.6	8.6	13.8	18.3	22.2	21.8	15.6
2004 4/.....	1.7	3.5	4.9	5.9	9.5	13.8	19.0	22.2	21.7	16.1
2003 4/.....	1.9	3.5	4.7	6.6	9.1	14.1	21.6	24.1	23.4	16.3
2002.....	2.5	3.3	4.4	6.2	9.9	14.7	19.3	23.7	23.0	17.2
2001.....	2.4	3.5	4.4	6.1	9.6	14.1	19.3	23.6	22.4	17.1
2000.....	2.0	3.4	4.0	6.5	9.0	13.8	18.9	23.3	24.1	17.3
1999.....	1.7	2.8	3.7	5.7	10.2	14.6	19.7	23.6	23.4	17.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fibrosis and cirrhosis of liver (K74)										
Black, both sexes										
2006 4/.....	2.9	5.0	11.2	13.1	11.8	12.0	9.6	14.1	9.5	9.6
2005 4/.....	2.2	6.5	11.4	13.6	13.4	11.0	12.1	14.4	14.1	10.7
2004 4/.....	3.6	6.9	12.1	12.7	12.8	11.8	13.6	11.2	11.4	11.3
2003 4/.....	3.5	7.4	12.8	14.2	12.1	13.8	14.6	11.1	10.9	10.8
2002.....	3.4	8.8	12.4	12.0	13.0	12.6	12.3	13.9	11.0	9.2
2001.....	4.5	9.9	12.0	13.5	12.0	13.5	14.1	14.1	11.4	7.4
2000.....	3.9	8.0	12.2	11.8	13.2	14.8	15.9	11.9	12.7	11.6
1999.....	4.6	9.5	9.8	13.1	14.5	14.1	14.1	14.6	12.4	12.1
Black, male										
2006 4/.....	4.4	6.4	17.1	19.5	17.0	17.6	12.3	21.9	*	*
2005 4/.....	3.5	9.1	16.5	21.4	20.0	15.4	16.4	20.6	18.3	*
2004 4/.....	4.9	10.4	18.5	19.9	17.0	15.7	18.5	15.9	14.3	*
2003 4/.....	4.2	9.4	20.9	22.6	18.1	17.7	16.1	14.6	*	*
2002.....	4.0	11.5	19.2	19.5	18.6	18.2	16.0	12.4	*	*
2001.....	6.2	15.1	19.8	20.3	19.5	19.8	18.5	18.1	*	*
2000.....	4.8	11.3	19.8	17.9	19.3	18.5	20.7	16.1	*	*
1999.....	7.0	15.0	16.1	19.2	19.1	19.2	18.0	17.6	*	*
Black, female										
2006 4/.....	1.7	3.8	6.2	7.8	7.8	8.0	7.8	9.4	9.4	8.4
2005 4/.....	*	4.2	7.1	7.1	8.2	7.8	9.1	10.7	11.9	10.1
2004 4/.....	2.4	3.8	6.7	6.8	9.5	8.8	10.3	8.4	9.8	11.0
2003 4/.....	2.9	5.6	5.9	7.3	7.4	11.0	13.5	9.0	10.1	8.1
2002.....	2.8	6.4	6.7	5.9	8.7	8.5	9.9	14.8	9.5	9.0
2001.....	2.9	5.4	5.3	7.8	6.1	8.8	11.2	11.6	12.7	*
2000.....	3.1	5.1	5.8	6.7	8.3	12.1	12.7	9.4	13.2	10.3
1999.....	2.5	4.8	4.5	8.2	10.8	10.4	11.5	12.9	11.7	12.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of liver (K71-K72,K75-K76)										
All races 3/, both sexes										
2006.....	2.2	3.9	6.2	7.4	8.3	8.5	10.1	10.2	12.4	11.3
2005.....	2.2	3.9	6.0	7.0	7.5	8.9	10.1	10.6	12.6	12.7
2004.....	2.4	4.2	5.8	6.5	7.3	8.3	10.1	11.3	11.2	13.1
2003.....	2.6	4.3	6.1	6.4	7.7	9.3	9.2	10.5	11.7	13.5
2002.....	2.7	4.7	5.4	5.7	6.8	8.1	9.2	9.4	12.0	14.3
2001.....	2.1	3.9	4.9	5.3	7.0	7.7	8.6	10.9	11.5	13.8
2000.....	2.2	3.6	4.7	5.0	6.2	7.4	8.9	9.5	11.2	13.6
1999.....	2.2	3.5	4.5	5.1	6.4	8.4	9.3	9.8	11.0	14.0
All races 3/, male										
2006.....	2.6	4.8	8.5	10.2	10.8	9.8	11.3	12.3	13.8	13.0
2005.....	2.5	5.5	8.5	9.4	9.7	10.4	11.7	12.3	15.2	15.6
2004.....	2.8	5.5	8.6	8.8	9.2	9.7	11.5	12.7	12.8	15.1
2003.....	3.0	5.8	8.5	8.4	10.1	11.2	11.2	11.9	14.1	16.9
2002.....	3.4	6.5	7.7	7.7	8.7	10.5	11.6	11.6	15.9	18.9
2001.....	2.8	5.4	7.1	6.8	9.1	9.8	10.2	13.5	15.0	17.4
2000.....	2.8	5.1	6.2	6.4	7.4	9.1	10.4	12.1	13.6	17.8
1999.....	2.9	4.9	6.1	6.9	8.2	10.6	11.0	11.0	14.9	19.2
All races 3/, female										
2006.....	1.8	3.0	4.0	4.8	6.0	7.4	9.2	8.7	11.6	10.4
2005.....	1.8	2.5	3.6	4.6	5.5	7.6	8.7	9.4	11.0	11.3
2004.....	1.9	2.9	3.2	4.3	5.7	7.2	8.9	10.2	10.3	12.3
2003.....	2.2	2.8	3.8	4.6	5.6	7.6	7.6	9.5	10.3	12.1
2002.....	2.0	2.9	3.2	3.8	5.0	6.0	7.2	7.9	9.6	12.3
2001.....	1.5	2.6	2.8	3.9	5.1	6.0	7.2	9.0	9.4	12.3
2000.....	1.6	2.2	3.2	3.7	5.1	6.0	7.7	7.8	9.7	11.9
1999.....	1.6	2.1	2.9	3.5	4.7	6.4	7.9	8.9	8.8	11.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of liver (K71-K72,K75-K76)										
White, both sexes										
2006 4/.....	2.2	3.9	6.1	7.1	8.0	8.6	9.9	10.3	12.5	11.4
2005 4/.....	2.1	3.7	5.8	6.8	7.6	8.7	10.0	10.8	12.3	12.6
2004 4/.....	2.3	4.0	5.5	6.1	7.0	8.2	9.9	11.4	11.2	13.1
2003 4/.....	2.6	4.2	5.8	6.2	7.5	9.2	9.1	10.5	11.9	13.4
2002.....	2.6	4.4	5.1	5.4	6.6	8.1	9.2	9.4	11.9	14.3
2001.....	2.0	3.7	4.7	4.9	6.8	7.8	8.4	10.8	11.4	13.8
2000.....	2.1	3.4	4.2	4.6	5.8	7.4	8.8	9.5	11.1	13.6
1999.....	2.1	3.2	4.1	4.8	6.0	8.2	9.1	9.7	11.2	13.9
White, male										
2006 4/.....	2.6	4.8	8.3	9.7	10.6	9.9	11.2	12.6	14.0	13.5
2005 4/.....	2.5	5.2	8.2	9.2	9.8	10.1	11.9	12.5	14.8	15.6
2004 4/.....	2.7	5.2	8.3	8.2	8.7	9.6	11.4	12.6	13.0	15.9
2003 4/.....	3.1	5.7	8.0	7.9	9.7	10.9	10.9	12.0	14.4	16.7
2002.....	3.3	6.2	7.3	7.1	8.5	10.6	11.7	11.3	15.8	18.9
2001.....	2.6	5.1	6.8	6.3	8.9	10.1	10.0	13.2	14.6	17.3
2000.....	2.8	4.8	5.5	5.9	7.0	8.9	10.3	12.0	13.9	17.8
1999.....	2.8	4.5	5.7	6.5	7.8	10.5	10.9	11.2	15.4	19.6
White, female										
2006 4/.....	1.8	2.9	3.9	4.6	5.7	7.4	8.9	8.6	11.6	10.5
2005 4/.....	1.7	2.3	3.4	4.5	5.4	7.4	8.5	9.5	10.7	11.2
2004 4/.....	1.8	2.7	2.9	4.2	5.5	7.0	8.7	10.4	10.2	11.9
2003 4/.....	2.1	2.7	3.5	4.6	5.4	7.7	7.5	9.4	10.4	11.9
2002.....	1.8	2.6	3.0	3.7	4.9	5.9	7.2	8.0	9.6	12.3
2001.....	1.4	2.4	2.6	3.6	4.9	5.8	7.0	9.1	9.4	12.4
2000.....	1.4	2.0	2.9	3.3	4.7	6.0	7.6	7.6	9.4	11.9
1999.....	1.3	1.8	2.6	3.1	4.3	6.2	7.6	8.7	8.7	11.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other diseases of liver (K71-K72,K75-K76)										
Black, both sexes										
2006 4/.....	2.2	4.8	8.2	10.6	11.0	9.0	12.7	8.9	10.9	10.6
2005 4/.....	2.7	5.8	8.7	9.6	8.4	11.3	9.6	9.0	16.5	12.0
2004 4/.....	3.0	6.6	8.6	10.1	10.2	10.3	11.4	10.0	12.1	12.1
2003 4/.....	3.0	5.9	9.7	9.2	10.8	11.3	10.7	10.4	8.9	14.9
2002.....	3.3	7.0	8.5	9.3	8.4	9.7	10.5	10.0	12.8	15.4
2001.....	3.0	6.4	7.6	9.9	9.3	9.1	11.4	12.0	14.4	11.4
2000.....	3.3	6.0	8.9	8.6	10.0	9.2	10.9	10.8	10.7	12.2
1999.....	3.7	6.4	8.0	8.7	10.9	10.4	11.2	11.2	9.2	14.0
Black, male										
2006 4/.....	2.5	5.9	10.7	16.2	15.0	11.7	15.4	10.3	*	*
2005 4/.....	2.9	8.0	12.8	13.2	11.1	14.3	10.8	8.9	20.4	*
2004 4/.....	3.5	8.6	12.1	15.6	14.4	12.4	14.6	13.2	*	*
2003 4/.....	2.6	7.7	14.0	13.7	15.6	17.0	15.8	11.4	*	*
2002.....	3.6	9.4	11.9	14.6	10.9	12.2	12.7	13.8	17.6	*
2001.....	4.1	8.7	11.1	13.0	12.2	10.2	14.5	17.2	23.3	*
2000.....	3.9	8.2	12.8	12.2	13.3	13.0	13.0	13.7	*	*
1999.....	4.1	9.0	11.1	11.7	14.0	12.5	13.2	10.4	*	*
Black, female										
2006 4/.....	2.0	3.8	6.2	6.1	7.8	6.9	10.8	8.0	10.1	11.3
2005 4/.....	2.4	3.9	5.1	6.7	6.4	9.2	8.9	9.1	14.5	12.8
2004 4/.....	2.6	4.9	5.5	5.6	7.0	8.8	9.2	8.1	12.1	14.6
2003 4/.....	3.3	4.4	6.1	5.5	7.1	7.1	7.3	9.8	*	13.4
2002.....	3.0	4.9	5.5	5.0	6.5	7.8	9.0	7.7	10.3	13.6
2001.....	2.1	4.4	4.5	7.3	7.0	8.3	9.4	8.9	9.8	10.1
2000.....	2.7	4.0	5.5	5.7	7.5	6.3	9.5	9.1	11.0	10.7
1999.....	3.3	4.2	5.3	6.2	8.5	8.9	9.9	11.7	9.9	13.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)										
All races 3/, both sexes										
2006.....	0.1	0.2	0.4	0.6	1.2	1.8	3.3	6.2	9.9	21.8
2005.....	0.1	0.2	0.4	0.5	0.9	1.7	3.3	5.8	11.1	22.9
2004.....	0.2	0.3	0.4	0.6	1.2	1.8	3.2	5.8	10.4	23.8
2003.....	0.1	0.2	0.4	0.5	0.9	2.0	3.4	5.9	11.3	21.8
2002.....	0.2	0.2	0.4	0.6	1.1	2.0	3.6	5.7	10.8	22.9
2001.....	0.1	0.3	0.4	0.7	1.3	2.0	3.8	5.8	10.0	23.7
2000.....	0.1	0.2	0.4	0.6	1.0	1.9	3.0	5.7	10.3	24.8
1999.....	0.2	0.3	0.4	0.7	1.0	1.8	3.4	5.7	10.9	23.9
All races 3/, male										
2006.....	*	0.2	0.5	0.6	1.2	2.3	4.0	7.6	12.5	21.7
2005.....	*	*	0.5	0.5	1.2	2.3	4.3	7.1	12.7	24.3
2004.....	*	0.3	0.4	0.7	1.4	2.2	3.8	7.0	12.5	25.5
2003.....	*	0.2	0.5	0.6	1.1	2.2	4.3	7.5	13.9	22.7
2002.....	0.2	0.3	0.5	0.8	1.1	2.3	4.0	7.5	13.5	24.1
2001.....	*	0.4	0.4	0.9	1.5	2.2	4.4	7.6	12.1	26.9
2000.....	*	0.2	0.3	0.7	1.2	2.3	3.6	7.4	12.9	27.6
1999.....	*	0.3	0.4	0.9	1.3	2.0	4.2	6.9	13.3	27.7
All races 3/, female										
2006.....	*	0.2	0.3	0.5	1.2	1.3	2.7	5.2	8.3	21.9
2005.....	*	0.2	0.3	0.6	0.7	1.2	2.4	4.8	10.1	22.3
2004.....	*	0.3	0.4	0.5	1.0	1.5	2.8	5.0	9.0	23.0
2003.....	*	0.2	0.4	0.4	0.7	1.8	2.8	4.8	9.8	21.4
2002.....	*	*	0.3	0.4	1.0	1.8	3.2	4.4	9.1	22.4
2001.....	*	0.3	0.3	0.5	1.1	1.9	3.3	4.6	8.7	22.4
2000.....	*	0.2	0.4	0.5	0.8	1.6	2.5	4.5	8.7	23.7
1999.....	*	0.3	0.5	0.6	0.7	1.6	2.8	4.8	9.5	22.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)										
White, both sexes										
2006 4/.....	*	0.2	0.3	0.5	1.1	1.6	3.0	6.1	10.0	22.3
2005 4/.....	*	0.1	0.3	0.5	0.9	1.5	3.1	5.7	11.1	23.1
2004 4/.....	0.2	0.2	0.3	0.5	1.1	1.8	3.0	5.7	10.3	24.2
2003 4/.....	*	0.2	0.3	0.4	0.8	1.8	3.6	5.9	11.3	21.9
2002.....	0.2	0.2	0.3	0.6	1.0	2.0	3.5	5.5	10.6	23.5
2001.....	*	0.3	0.3	0.6	1.3	1.8	3.7	5.7	9.8	23.9
2000.....	0.1	0.2	0.3	0.6	0.9	1.8	2.7	5.6	10.1	25.6
1999.....	0.1	0.3	0.4	0.6	0.9	1.6	3.4	5.7	11.0	24.4
White, male										
2006 4/.....	*	*	0.4	0.5	1.1	2.1	3.8	7.5	12.7	22.8
2005 4/.....	*	*	0.4	0.5	1.1	2.1	4.1	7.1	13.1	24.8
2004 4/.....	*	0.2	0.3	0.6	1.4	2.2	3.7	6.9	12.7	26.7
2003 4/.....	*	*	0.4	0.5	0.9	2.1	4.4	7.5	13.9	23.0
2002.....	*	0.3	0.5	0.8	1.1	2.3	4.0	7.4	13.3	24.7
2001.....	*	0.3	0.4	0.8	1.4	1.9	4.4	7.6	12.0	26.7
2000.....	*	*	0.3	0.7	1.1	2.3	3.4	7.4	13.2	28.9
1999.....	*	0.3	0.3	0.7	1.2	1.9	4.2	7.0	13.8	28.1
White, female										
2006 4/.....	*	*	0.3	0.5	1.2	1.1	2.4	5.0	8.3	22.0
2005 4/.....	*	*	0.2	0.5	0.7	0.9	2.3	4.7	9.8	22.4
2004 4/.....	*	0.2	0.3	0.5	0.8	1.4	2.4	4.8	8.8	23.1
2003 4/.....	*	*	0.3	0.3	0.7	1.6	2.8	4.7	9.8	21.4
2002.....	*	*	0.3	0.4	0.9	1.7	3.1	4.1	9.0	23.0
2001.....	*	0.3	*	0.4	1.1	1.8	3.1	4.4	8.4	22.8
2000.....	*	*	0.3	0.4	0.7	1.4	2.2	4.4	8.3	24.3
1999.....	*	0.3	0.4	0.6	0.6	1.4	2.7	4.8	9.3	22.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Cholelithiasis and other disorders of gallbladder (K80-K82)										
Black, both sexes										
2006 4/.....	*	*	0.9	1.1	1.9	3.4	4.6	9.0	9.0	17.1
2005 4/.....	*	*	*	*	*	3.6	4.6	7.0	12.4	20.6
2004 4/.....	*	*	1.0	1.3	2.0	2.3	5.1	7.1	12.1	19.5
2003 4/.....	*	*	1.0	*	*	3.0	2.8	6.0	10.9	20.7
2002.....	*	*	*	*	*	2.3	4.6	7.7	12.8	14.8
2001.....	*	*	*	1.4	2.0	3.5	6.2	6.1	11.6	19.7
2000.....	*	*	*	*	*	3.1	4.9	6.6	11.0	17.8
1999.....	*	*	*	*	2.0	3.2	4.1	4.6	11.3	15.3
Black, male										
2006 4/.....	*	*	*	*	*	*	*	10.3	*	*
2005 4/.....	*	*	*	*	*	*	6.3	9.8	*	19.5
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	20.9
2002.....	*	*	*	*	*	*	*	*	16.0	*
2001.....	*	*	*	*	*	*	*	*	*	27.2
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	*	*	*	*	*	*	5.3	8.3	9.4	19.7
2005 4/.....	*	*	*	*	*	*	*	5.3	13.4	21.0
2004 4/.....	*	*	*	*	*	*	5.5	6.7	12.9	21.7
2003 4/.....	*	*	*	*	*	*	*	*	9.3	20.7
2002.....	*	*	*	*	*	*	5.4	7.7	11.1	16.0
2001.....	*	*	*	*	*	*	6.3	6.9	13.1	16.8
2000.....	*	*	*	*	*	*	4.9	6.3	11.5	19.3
1999.....	*	*	*	*	*	*	*	*	12.1	15.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of biliary tract and pancreas (K83-K86)										
All races 3/, both sexes										
2006.....	1.0	1.3	1.5	1.8	2.9	3.7	5.2	7.7	11.0	21.2
2005.....	0.9	1.1	1.7	1.8	2.6	4.1	4.9	7.5	12.0	21.0
2004.....	0.9	1.3	1.6	2.0	2.5	3.9	5.6	7.9	11.5	20.8
2003.....	1.0	1.3	1.5	1.9	2.9	3.9	5.9	8.3	12.3	21.9
2002.....	1.0	1.3	1.5	2.3	2.7	4.0	5.8	8.1	12.0	21.7
2001.....	1.0	1.4	1.7	2.1	3.1	4.2	6.1	7.6	11.9	22.9
2000.....	1.0	1.3	1.5	2.3	2.7	3.7	5.5	7.9	11.5	22.0
1999.....	1.0	1.3	1.6	2.0	2.8	3.6	5.4	7.2	10.3	22.2
All races 3/, male										
2006.....	1.2	1.8	2.2	2.5	3.7	5.3	7.2	9.0	12.5	18.5
2005.....	1.3	1.5	2.3	2.7	3.4	5.0	5.9	9.6	14.2	21.5
2004.....	1.2	1.8	2.2	2.7	3.4	5.1	6.5	9.3	12.6	21.1
2003.....	1.2	1.7	2.1	2.6	3.7	5.1	7.3	10.1	14.4	20.7
2002.....	1.3	1.9	2.2	3.0	4.0	5.5	7.4	10.3	14.3	21.2
2001.....	1.3	1.9	2.2	2.8	3.9	5.1	8.0	8.8	12.5	22.4
2000.....	1.5	1.8	2.1	2.9	3.1	5.2	6.7	10.2	13.3	22.9
1999.....	1.4	1.8	2.1	2.8	3.6	4.6	6.7	9.1	11.7	23.1
All races 3/, female										
2006.....	0.7	0.7	0.9	1.2	2.1	2.3	3.6	6.7	10.0	22.5
2005.....	0.5	0.7	1.1	1.1	1.8	3.4	4.0	6.0	10.6	20.7
2004.....	0.6	0.9	1.0	1.2	1.7	2.8	4.9	6.8	10.8	20.7
2003.....	0.7	0.8	0.9	1.1	2.1	2.8	4.8	7.1	10.9	22.4
2002.....	0.7	0.7	0.9	1.7	1.6	2.7	4.4	6.6	10.5	21.9
2001.....	0.7	1.0	1.2	1.4	2.4	3.3	4.5	6.8	11.6	23.2
2000.....	0.6	0.9	1.0	1.7	2.3	2.5	4.6	6.3	10.5	21.6
1999.....	0.7	0.8	1.1	1.4	2.2	2.8	4.4	5.9	9.5	21.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of biliary tract and pancreas (K83-K86)										
White, both sexes										
2006 4/.....	0.9	1.2	1.4	1.7	2.9	3.7	5.2	7.8	11.1	21.2
2005 4/.....	0.9	1.0	1.5	1.7	2.4	4.1	4.8	7.4	11.9	21.2
2004 4/.....	0.8	1.2	1.4	1.7	2.3	3.9	5.4	7.8	11.5	21.0
2003 4/.....	0.8	1.1	1.3	1.7	2.7	3.7	5.6	8.5	12.4	22.0
2002.....	0.9	1.1	1.3	2.2	2.5	3.9	5.7	8.2	12.1	22.0
2001.....	0.9	1.3	1.4	1.9	3.0	4.1	6.0	7.6	12.0	23.0
2000.....	0.9	1.1	1.4	2.0	2.5	3.4	5.4	7.8	11.5	21.9
1999.....	1.0	1.0	1.4	1.8	2.6	3.5	5.5	7.1	10.2	22.0
White, male										
2006 4/.....	1.1	1.7	2.0	2.3	3.6	5.4	7.1	9.1	12.6	18.9
2005 4/.....	1.3	1.3	2.0	2.5	3.2	5.0	5.7	9.4	14.3	22.2
2004 4/.....	1.2	1.6	1.9	2.4	3.1	5.1	6.3	9.2	12.8	21.4
2003 4/.....	0.9	1.5	1.8	2.3	3.4	4.9	7.0	10.1	14.5	20.5
2002.....	1.2	1.7	1.9	2.8	3.6	5.5	7.2	10.5	14.5	21.2
2001.....	1.2	1.6	1.9	2.6	3.7	5.0	8.0	8.7	12.9	22.2
2000.....	1.3	1.5	1.8	2.7	2.9	4.7	6.6	10.2	13.2	22.0
1999.....	1.3	1.4	1.8	2.4	3.3	4.6	6.8	9.0	11.7	22.0
White, female										
2006 4/.....	0.6	0.7	0.8	1.1	2.2	2.2	3.6	6.9	10.2	22.3
2005 4/.....	0.5	0.6	1.0	1.0	1.7	3.2	4.1	6.0	10.5	20.8
2004 4/.....	0.5	0.8	0.9	1.1	1.6	2.8	4.7	6.8	10.6	20.8
2003 4/.....	0.6	0.6	0.8	1.0	2.0	2.6	4.4	7.3	11.1	22.6
2002.....	0.5	0.6	0.8	1.7	1.4	2.6	4.5	6.6	10.6	22.3
2001.....	0.6	0.9	1.0	1.3	2.3	3.3	4.3	6.8	11.5	23.4
2000.....	0.6	0.8	1.0	1.5	2.2	2.2	4.5	6.2	10.4	21.9
1999.....	0.6	0.6	1.1	1.3	1.9	2.6	4.4	5.7	9.3	22.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of biliary tract and pancreas (K83-K86)										
Black, both sexes										
2006 4/.....	1.6	2.1	2.6	3.8	3.9	4.6	6.1	6.5	9.7	21.8
2005 4/.....	1.2	2.2	3.3	3.4	4.1	5.4	5.4	9.2	14.1	19.8
2004 4/.....	1.8	2.4	3.5	4.5	5.3	4.3	8.8	10.0	11.4	20.3
2003 4/.....	2.3	2.8	3.4	3.9	5.3	5.9	9.3	7.8	10.4	23.3
2002.....	1.8	2.9	3.2	3.8	5.3	5.0	7.1	6.7	11.5	21.4
2001.....	2.2	3.2	4.1	3.9	4.4	6.0	7.5	8.0	11.6	22.1
2000.....	2.0	3.1	2.8	4.5	4.6	6.7	7.7	8.7	12.1	23.5
1999.....	1.7	3.3	3.3	4.1	5.5	5.6	5.3	8.4	12.4	23.0
Black, male										
2006 4/.....	2.0	3.1	4.3	5.4	6.9	6.1	9.2	*	*	*
2005 4/.....	1.8	3.3	4.7	5.0	5.5	6.4	7.6	12.5	15.5	*
2004 4/.....	1.8	3.5	5.2	6.3	7.7	6.1	11.7	13.6	*	*
2003 4/.....	3.6	3.5	4.8	6.2	7.7	7.1	11.8	10.0	14.8	26.1
2002.....	2.1	4.5	4.7	5.7	7.3	6.0	11.7	*	*	25.7
2001.....	2.7	4.7	5.5	5.9	5.9	8.4	9.1	11.2	*	*
2000.....	3.0	3.9	4.4	6.3	5.6	10.4	9.7	10.9	*	32.3
1999.....	2.5	5.0	5.4	6.6	6.4	6.8	*	10.4	*	32.8
Black, female										
2006 4/.....	*	*	*	2.5	*	3.5	*	5.9	8.3	25.6
2005 4/.....	*	*	2.1	2.1	3.1	4.6	*	7.2	13.4	20.6
2004 4/.....	1.7	1.5	2.1	2.9	3.5	*	6.8	7.8	12.5	20.9
2003 4/.....	*	2.3	2.1	*	3.5	4.9	7.5	6.5	8.1	22.3
2002.....	1.5	1.5	1.9	*	3.7	4.3	*	5.5	11.5	19.7
2001.....	1.8	2.0	2.9	*	3.2	4.3	6.5	6.1	13.5	22.3
2000.....	*	2.4	*	3.1	3.8	4.0	6.4	7.4	11.0	20.2
1999.....	*	1.8	*	*	4.8	4.8	5.2	7.2	13.0	19.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pancreatitis (K85,K86.0-K86.1)										
All races 3/, both sexes										
2006.....	0.8	1.1	1.3	1.5	2.2	2.8	3.8	5.0	6.5	11.7
2005.....	0.8	1.0	1.5	1.5	1.8	3.0	3.5	4.9	7.5	10.9
2004.....	0.8	1.2	1.3	1.6	2.0	2.9	4.0	5.0	7.0	11.4
2003.....	0.9	1.1	1.3	1.5	2.2	2.8	4.1	5.7	7.9	11.8
2002.....	0.9	1.2	1.3	1.9	2.2	2.9	4.0	5.7	7.8	11.9
2001.....	0.9	1.2	1.4	1.8	2.5	3.2	4.5	5.4	7.7	13.2
2000.....	0.9	1.1	1.3	1.8	2.3	2.8	4.0	5.4	7.6	12.1
1999.....	0.9	1.1	1.4	1.6	2.2	2.9	4.0	5.2	6.8	12.8
All races 3/, male										
2006.....	1.1	1.6	1.9	2.0	3.0	4.0	5.3	6.1	7.8	9.6
2005.....	1.2	1.4	2.0	2.1	2.4	3.6	4.4	6.2	8.8	11.3
2004.....	1.2	1.7	1.9	2.4	2.7	3.9	4.8	6.1	7.2	10.3
2003.....	1.1	1.5	1.8	2.1	3.1	4.0	5.1	6.7	9.1	10.6
2002.....	1.1	1.8	1.9	2.5	3.2	4.2	5.1	7.1	8.8	11.2
2001.....	1.2	1.7	1.8	2.5	3.1	3.9	6.1	6.1	7.8	12.9
2000.....	1.4	1.4	1.7	2.4	2.6	4.1	5.0	6.8	8.3	11.9
1999.....	1.3	1.6	1.8	2.2	2.9	3.8	5.1	6.4	7.5	12.7
All races 3/, female										
2006.....	0.6	0.6	0.7	1.0	1.6	1.6	2.5	4.2	5.7	12.7
2005.....	0.5	0.6	0.9	0.9	1.2	2.5	2.7	3.9	6.6	10.8
2004.....	0.5	0.7	0.8	1.0	1.3	2.0	3.4	4.3	6.9	11.8
2003.....	0.6	0.7	0.8	0.8	1.5	1.9	3.4	5.0	7.2	12.3
2002.....	0.6	0.6	0.8	1.4	1.2	1.8	3.1	4.7	7.1	12.2
2001.....	0.6	0.8	1.0	1.1	1.9	2.7	3.3	4.9	7.7	13.4
2000.....	0.5	0.8	0.8	1.2	1.9	1.8	3.2	4.4	7.2	12.2
1999.....	0.6	0.7	0.9	1.0	1.6	2.2	3.2	4.3	6.5	12.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pancreatitis (K85,K86.0-K86.1)										
White, both sexes										
2006 4/.....	0.8	1.0	1.2	1.4	2.2	2.7	3.8	5.1	6.8	11.8
2005 4/.....	0.8	0.8	1.3	1.4	1.7	3.0	3.4	4.8	7.5	11.0
2004 4/.....	0.8	1.1	1.1	1.4	1.8	2.9	3.9	5.0	6.9	11.4
2003 4/.....	0.7	0.9	1.1	1.3	2.1	2.8	3.8	5.9	8.0	11.6
2002.....	0.8	1.0	1.1	1.9	1.9	2.9	4.0	5.8	7.9	12.1
2001.....	0.8	1.1	1.1	1.7	2.4	3.2	4.5	5.5	7.7	13.3
2000.....	0.8	0.9	1.2	1.6	2.1	2.6	3.9	5.3	7.7	12.2
1999.....	0.9	0.9	1.2	1.4	2.0	2.9	4.0	5.0	6.8	12.8
White, male										
2006 4/.....	1.0	1.5	1.6	1.9	2.8	4.1	5.4	6.1	8.1	9.9
2005 4/.....	1.2	1.1	1.8	2.0	2.3	3.6	4.2	6.1	9.0	11.9
2004 4/.....	1.1	1.5	1.6	2.0	2.4	3.8	4.6	6.1	7.1	10.4
2003 4/.....	0.9	1.3	1.5	1.9	2.7	3.8	4.7	6.7	9.1	10.3
2002.....	1.1	1.5	1.6	2.4	2.8	4.3	5.1	7.2	8.9	11.0
2001.....	1.1	1.5	1.5	2.3	2.9	3.8	6.2	6.1	7.8	12.9
2000.....	1.2	1.2	1.5	2.1	2.4	3.7	4.9	6.7	8.3	11.5
1999.....	1.2	1.2	1.5	1.9	2.7	3.7	5.2	6.5	7.4	12.0
White, female										
2006 4/.....	0.5	0.6	0.7	0.9	1.6	1.5	2.5	4.3	5.9	12.7
2005 4/.....	0.5	0.5	0.9	0.8	1.2	2.4	2.7	3.9	6.6	10.6
2004 4/.....	0.4	0.6	0.6	0.8	1.2	2.1	3.2	4.3	6.9	11.8
2003 4/.....	0.5	0.5	0.7	0.8	1.4	1.9	3.1	5.2	7.3	12.2
2002.....	0.5	0.5	0.6	1.4	1.1	1.8	3.2	4.8	7.2	12.6
2001.....	0.5	0.7	0.8	1.1	1.9	2.6	3.1	5.0	7.7	13.4
2000.....	0.5	0.7	0.8	1.0	1.8	1.6	3.2	4.4	7.3	12.5
1999.....	0.5	0.6	0.9	1.0	1.3	2.1	3.1	4.1	6.4	13.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pancreatitis (K85,K86.0-K86.1)										
Black, both sexes										
2006 4/.....	1.4	1.9	2.3	3.1	3.3	3.6	4.5	4.9	*	10.9
2005 4/.....	1.0	1.9	2.8	2.7	2.9	3.8	4.4	6.3	8.3	10.4
2004 4/.....	1.6	2.3	3.1	4.0	4.1	3.3	6.4	6.4	8.2	13.3
2003 4/.....	2.2	2.5	2.9	3.2	4.3	4.0	7.0	5.1	6.9	14.0
2002.....	1.4	2.5	2.8	3.0	4.7	3.5	3.9	4.3	6.5	11.3
2001.....	2.1	2.8	3.8	3.1	3.8	4.2	5.1	5.0	8.7	13.2
2000.....	1.9	2.7	2.4	3.9	3.8	5.2	5.6	5.3	7.6	11.9
1999.....	1.6	3.0	3.1	3.3	4.3	4.5	4.2	6.4	8.1	12.8
Black, male										
2006 4/.....	1.8	3.0	4.1	4.2	6.0	4.7	6.5	*	*	*
2005 4/.....	1.6	2.9	4.2	4.0	3.6	4.8	6.3	*	*	*
2004 4/.....	1.7	3.3	4.6	5.8	5.8	5.3	8.7	9.6	*	*
2003 4/.....	3.3	3.3	4.2	5.5	6.7	5.9	9.2	*	*	*
2002.....	1.8	3.9	4.2	4.5	6.9	*	*	*	*	*
2001.....	2.6	4.2	4.9	5.0	5.5	5.6	*	*	*	*
2000.....	3.0	3.2	3.8	5.4	4.8	8.3	7.7	*	*	*
1999.....	2.4	4.5	5.1	5.4	4.9	6.0	*	*	*	*
Black, female										
2006 4/.....	*	*	*	2.1	*	*	*	*	*	12.4
2005 4/.....	*	*	1.6	*	*	*	*	*	8.2	11.6
2004 4/.....	1.4	1.4	1.9	2.5	*	*	4.8	*	8.7	13.4
2003 4/.....	*	1.8	1.7	*	*	*	5.5	*	*	13.4
2002.....	*	*	*	*	*	*	*	*	*	9.4
2001.....	1.6	1.7	2.9	*	*	*	4.7	*	9.4	13.9
2000.....	*	2.2	*	2.7	*	*	*	*	*	9.9
1999.....	*	1.6	*	*	3.8	*	*	6.0	*	10.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)

All races 3/, both sexes

2006.....	0.1	0.2	0.2	0.3	0.6	1.0	1.4	2.7	4.5	9.5
2005.....	0.1	0.2	0.2	0.3	0.7	1.1	1.4	2.6	4.6	10.0
2004.....	0.1	0.1	0.2	0.3	0.6	1.0	1.6	2.8	4.4	9.4
2003.....	0.1	0.2	0.2	0.4	0.6	1.0	1.8	2.6	4.3	10.1
2002.....	0.1	0.1	0.2	0.4	0.6	1.1	1.8	2.5	4.2	9.8
2001.....	*	0.2	0.3	0.3	0.6	1.0	1.5	2.2	4.2	9.7
2000.....	0.1	0.2	0.3	0.5	0.4	0.9	1.5	2.5	3.9	9.9
1999.....	0.1	0.1	0.2	0.5	0.6	0.7	1.4	2.0	3.5	9.4

All races 3/, male

2006.....	*	0.2	0.3	0.5	0.7	1.3	1.8	2.9	4.7	8.9
2005.....	*	0.2	0.2	0.5	1.0	1.4	1.5	3.3	5.4	10.2
2004.....	*	*	0.2	0.4	0.8	1.2	1.7	3.2	5.4	10.7
2003.....	*	*	0.4	0.5	0.7	1.1	2.2	3.4	5.3	10.1
2002.....	*	*	0.3	0.5	0.8	1.3	2.3	3.3	5.5	10.0
2001.....	*	0.2	0.4	0.3	0.7	1.3	1.9	2.6	4.7	9.5
2000.....	*	0.3	0.4	0.6	0.5	1.1	1.6	3.5	5.0	11.0
1999.....	*	0.2	0.3	0.6	0.7	0.8	1.6	2.7	4.2	10.5

All races 3/, female

2006.....	*	*	*	0.2	0.5	0.7	1.0	2.5	4.3	9.8
2005.....	*	*	0.2	*	0.5	0.9	1.3	2.1	4.0	10.0
2004.....	*	*	0.2	0.3	0.4	0.8	1.5	2.6	3.9	8.9
2003.....	*	*	*	0.3	0.6	0.9	1.4	2.1	3.7	10.2
2002.....	*	*	*	0.3	0.3	0.9	1.4	1.9	3.4	9.8
2001.....	*	0.2	*	0.3	0.5	0.7	1.2	1.9	3.9	9.8
2000.....	*	*	*	0.4	0.4	0.7	1.4	1.9	3.2	9.4
1999.....	*	*	*	0.4	0.6	0.6	1.2	1.6	3.1	9.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of biliary tract and pancreas (K83,K86.2-K86.9)

White, both sexes

2006 4/.....	0.1	0.2	0.2	0.3	0.6	1.0	1.4	2.8	4.3	9.4
2005 4/.....	*	0.1	0.2	0.3	0.7	1.1	1.4	2.6	4.4	10.2
2004 4/.....	*	0.1	0.2	0.3	0.5	1.0	1.5	2.8	4.5	9.6
2003 4/.....	*	0.1	0.2	0.3	0.6	0.9	1.8	2.7	4.4	10.3
2002.....	*	0.1	0.2	0.4	0.5	1.0	1.7	2.4	4.2	9.9
2001.....	*	0.2	0.3	0.2	0.6	0.9	1.5	2.1	4.3	9.7
2000.....	*	0.2	0.2	0.5	0.4	0.8	1.5	2.5	3.8	9.7
1999.....	*	0.1	0.3	0.4	0.6	0.7	1.4	2.0	3.4	9.2

White, male

2006 4/.....	*	*	0.3	0.4	0.7	1.3	1.8	3.0	4.5	9.1
2005 4/.....	*	*	*	0.5	0.9	1.4	1.5	3.3	5.2	10.3
2004 4/.....	*	*	*	0.3	0.7	1.3	1.6	3.1	5.7	10.9
2003 4/.....	*	*	0.3	0.5	0.6	1.1	2.2	3.4	5.4	10.2
2002.....	*	*	0.3	0.4	0.8	1.2	2.0	3.3	5.6	10.3
2001.....	*	*	0.4	*	0.8	1.1	1.8	2.6	5.1	9.3
2000.....	*	0.3	0.3	0.6	0.5	1.0	1.6	3.5	5.0	10.6
1999.....	*	*	0.3	0.5	0.6	0.9	1.7	2.6	4.3	10.0

White, female

2006 4/.....	*	*	*	*	0.6	0.6	1.1	2.6	4.2	9.6
2005 4/.....	*	*	*	*	0.5	0.8	1.4	2.1	3.9	10.1
2004 4/.....	*	*	0.2	*	0.4	0.7	1.5	2.5	3.8	9.0
2003 4/.....	*	*	*	*	0.6	0.7	1.4	2.1	3.8	10.4
2002.....	*	*	*	*	*	0.8	1.4	1.8	3.3	9.7
2001.....	*	*	*	*	0.4	0.7	1.2	1.8	3.8	9.9
2000.....	*	*	*	0.4	*	0.6	1.3	1.8	3.1	9.4
1999.....	*	*	*	0.3	0.6	0.5	1.3	1.7	2.9	8.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of digestive system (K90,K92)										
All races 3/, both sexes										
2006.....	0.5	0.8	1.2	1.8	2.6	4.1	7.0	12.7	21.9	62.0
2005.....	0.4	0.7	1.2	1.9	2.6	4.7	7.5	12.8	23.9	65.2
2004.....	0.3	0.7	1.2	1.8	2.7	4.2	7.3	13.0	25.0	68.5
2003.....	0.4	0.6	1.1	1.7	2.9	4.7	7.7	14.0	26.0	74.2
2002.....	0.4	0.8	1.2	1.7	2.6	4.7	7.6	14.1	27.6	76.2
2001.....	0.4	0.7	1.0	1.8	3.1	4.8	7.7	14.1	28.2	77.8
2000.....	0.4	0.7	1.1	1.5	2.8	4.1	7.9	15.0	29.0	80.6
1999.....	0.4	0.7	1.0	1.7	2.8	4.7	7.7	14.1	27.8	77.2
All races 3/, male										
2006.....	0.7	1.1	1.7	2.3	3.6	5.0	8.6	16.7	27.3	65.8
2005.....	0.5	0.9	1.6	2.5	3.3	6.0	9.1	16.3	28.1	66.8
2004.....	0.4	1.0	1.7	2.6	3.3	5.7	9.4	16.1	30.2	69.8
2003.....	0.6	0.9	1.5	2.3	3.6	6.1	9.7	17.8	32.3	73.3
2002.....	0.6	1.2	1.6	2.3	3.8	6.1	9.6	17.8	31.0	76.2
2001.....	0.6	1.0	1.6	2.5	3.9	6.3	9.7	17.5	33.0	81.4
2000.....	0.6	1.0	1.6	2.0	3.8	5.2	9.2	18.3	36.5	83.1
1999.....	0.5	1.0	1.4	2.3	3.6	6.1	10.0	17.3	34.3	80.8
All races 3/, female										
2006.....	0.3	0.6	0.8	1.3	1.7	3.4	5.7	9.8	18.4	60.2
2005.....	0.3	0.4	0.7	1.2	2.1	3.6	6.2	10.2	21.3	64.5
2004.....	0.3	0.4	0.7	1.0	2.1	2.9	5.7	10.9	21.8	67.9
2003.....	0.2	0.4	0.6	1.1	2.2	3.5	6.1	11.3	22.1	74.6
2002.....	0.2	0.4	0.7	1.2	1.6	3.4	6.0	11.5	25.5	76.2
2001.....	0.2	0.4	0.5	1.1	2.4	3.6	6.1	11.8	25.3	76.3
2000.....	0.2	0.4	0.6	1.0	1.9	3.0	6.8	12.7	24.6	79.6
1999.....	0.2	0.4	0.6	1.2	2.1	3.6	6.0	11.8	24.0	75.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other diseases of digestive system (K90,K92)										
White, both sexes										
2006 4/.....	0.5	0.8	1.1	1.7	2.5	3.7	6.7	12.4	21.8	62.9
2005 4/.....	0.3	0.6	1.0	1.8	2.4	4.4	7.1	12.2	24.1	66.9
2004 4/.....	0.3	0.7	1.1	1.6	2.5	3.9	7.0	12.8	24.8	69.8
2003 4/.....	0.4	0.6	1.0	1.6	2.6	4.3	7.3	13.6	26.2	75.5
2002.....	0.3	0.7	1.1	1.5	2.4	4.5	7.3	13.6	27.6	77.8
2001.....	0.3	0.6	0.9	1.6	2.8	4.5	7.5	13.7	28.4	79.1
2000.....	0.3	0.6	0.9	1.3	2.4	3.7	7.6	14.4	28.8	81.7
1999.....	0.3	0.6	0.8	1.6	2.5	4.4	7.5	13.6	27.6	78.9
White, male										
2006 4/.....	0.7	1.1	1.5	2.2	3.4	4.7	8.1	16.4	27.1	67.0
2005 4/.....	0.4	0.9	1.5	2.4	3.0	5.4	8.7	15.7	28.1	69.1
2004 4/.....	0.4	1.0	1.6	2.3	2.9	5.3	8.9	15.9	29.9	71.6
2003 4/.....	0.6	0.8	1.5	2.1	3.4	5.7	9.1	17.2	32.6	75.2
2002.....	0.5	1.0	1.6	2.1	3.5	5.8	9.1	17.3	31.2	77.3
2001.....	0.5	0.9	1.3	2.1	3.5	5.8	9.3	16.7	33.5	82.8
2000.....	0.5	0.8	1.4	1.7	3.4	4.9	8.8	17.5	36.2	84.9
1999.....	0.5	0.9	1.2	2.0	3.4	5.7	9.6	16.6	34.4	82.1
White, female										
2006 4/.....	0.2	0.5	0.7	1.2	1.6	2.8	5.4	9.3	18.4	61.0
2005 4/.....	0.2	0.4	0.6	1.2	1.9	3.4	5.8	9.7	21.6	65.9
2004 4/.....	0.2	0.3	0.6	1.0	2.1	2.6	5.5	10.6	21.7	69.0
2003 4/.....	0.3	0.4	0.6	1.0	1.9	3.0	5.9	11.0	22.2	75.6
2002.....	*	0.4	0.6	1.0	1.4	3.3	5.8	11.0	25.4	78.1
2001.....	*	0.4	0.5	1.0	2.1	3.4	6.0	11.5	25.3	77.5
2000.....	*	0.3	0.5	0.8	1.6	2.7	6.6	12.2	24.4	80.5
1999.....	*	0.3	0.5	1.1	1.8	3.3	5.8	11.4	23.6	77.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases of digestive system (K90,K92)

Black, both sexes

2006 4/.....	*	1.2	2.7	2.9	4.3	8.1	11.1	19.6	25.1	62.0
2005 4/.....	*	1.1	2.5	3.1	5.1	8.8	12.5	20.7	27.4	56.1
2004 4/.....	*	1.0	2.4	3.5	4.8	7.7	12.3	19.0	32.4	62.0
2003 4/.....	0.7	1.1	1.9	3.2	5.5	9.0	12.3	22.5	27.4	70.3
2002.....	0.8	1.6	2.0	3.6	5.3	7.8	12.2	21.6	33.5	65.3
2001.....	0.9	1.3	2.2	4.2	6.4	8.5	11.2	22.1	31.4	73.1
2000.....	0.9	1.6	2.5	3.9	6.3	7.2	12.9	22.6	36.7	77.9
1999.....	*	1.2	2.2	3.1	5.5	7.8	12.2	22.0	33.2	66.0

Black, male

2006 4/.....	*	*	4.2	3.6	6.4	8.2	14.1	26.0	35.6	64.3
2005 4/.....	*	1.6	3.2	4.5	6.5	13.3	15.2	26.4	35.1	47.4
2004 4/.....	*	*	3.2	5.7	7.5	10.9	17.5	22.7	39.5	58.6
2003 4/.....	*	*	2.6	4.8	6.7	10.6	18.1	29.7	34.0	59.4
2002.....	*	2.7	2.5	4.5	7.3	11.2	17.0	26.2	36.7	70.8
2001.....	*	2.2	3.7	6.3	7.9	11.4	16.5	30.2	36.1	71.5
2000.....	*	2.4	3.3	4.9	8.7	7.8	17.4	29.3	47.2	78.4
1999.....	*	*	3.3	5.3	6.2	9.9	16.6	29.0	38.4	71.5

Black, female

2006 4/.....	*	*	*	2.4	2.8	8.0	9.1	15.8	19.5	61.0
2005 4/.....	*	*	1.8	*	4.0	5.5	10.6	17.3	23.4	59.6
2004 4/.....	*	*	1.7	*	*	5.4	8.8	16.7	28.7	63.4
2003 4/.....	*	*	*	*	4.5	7.9	8.4	18.2	24.0	74.5
2002.....	*	*	*	2.8	3.7	5.2	9.0	18.9	31.8	63.3
2001.....	*	*	*	*	5.3	6.4	7.6	17.2	29.0	73.7
2000.....	*	*	*	3.0	4.4	6.7	10.0	18.5	31.4	77.7
1999.....	*	*	*	*	5.0	6.3	9.2	17.7	30.7	64.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XII. Diseases of the skin and subcutaneous tissue (L00-L98)

All races 3/, both sexes

2006.....	0.3	0.4	0.8	1.3	1.9	2.7	3.7	6.1	11.3	23.4
2005.....	0.3	0.6	0.8	1.2	1.6	2.5	4.2	7.2	12.7	29.2
2004.....	0.4	0.6	0.7	1.2	1.7	2.7	4.3	7.1	12.7	28.8
2003.....	0.4	0.5	0.8	1.2	1.7	2.4	4.3	6.7	12.5	30.5
2002.....	0.4	0.5	0.7	1.1	1.4	2.0	4.2	6.5	11.8	32.8
2001.....	0.3	0.5	0.7	1.0	1.4	2.2	3.5	6.5	12.0	32.2
2000.....	0.3	0.4	0.5	0.9	1.5	2.1	4.0	6.4	12.7	34.7
1999.....	0.3	0.5	0.6	0.8	1.3	2.2	3.5	6.6	12.7	35.0

All races 3/, male

2006.....	0.4	0.4	0.9	1.2	1.8	2.9	3.7	6.1	10.6	21.3
2005.....	0.4	0.5	0.9	1.3	1.6	2.6	4.1	7.1	12.2	26.3
2004.....	0.3	0.6	0.7	1.1	1.7	2.2	4.1	7.0	11.9	25.3
2003.....	0.4	0.5	0.8	1.1	1.8	2.3	4.4	6.3	11.7	26.6
2002.....	0.3	0.5	0.7	1.1	1.2	2.0	4.5	6.5	11.9	28.6
2001.....	0.2	0.6	0.7	1.0	1.4	2.5	3.6	6.7	10.8	24.9
2000.....	0.3	0.4	0.4	1.0	1.6	1.6	4.0	6.1	12.3	29.7
1999.....	0.2	0.5	0.6	0.8	1.1	2.3	3.5	6.1	12.2	31.7

All races 3/, female

2006.....	0.3	0.4	0.7	1.3	2.0	2.5	3.7	6.1	11.7	24.3
2005.....	0.3	0.6	0.7	1.1	1.6	2.5	4.3	7.2	12.9	30.6
2004.....	0.4	0.5	0.7	1.3	1.8	3.1	4.4	7.1	13.2	30.4
2003.....	0.4	0.5	0.8	1.2	1.6	2.5	4.3	7.1	13.0	32.2
2002.....	0.4	0.4	0.7	1.1	1.5	2.1	3.9	6.6	11.7	34.7
2001.....	0.3	0.5	0.6	1.0	1.5	1.9	3.4	6.4	12.7	35.2
2000.....	0.3	0.5	0.5	0.9	1.3	2.4	3.9	6.6	13.0	36.7
1999.....	0.4	0.5	0.6	0.8	1.4	2.2	3.6	7.0	12.9	36.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XII. Diseases of the skin and subcutaneous tissue (L00-L98)

White, both sexes

2006 4/.....	0.3	0.4	0.7	1.1	1.7	2.4	3.1	5.3	10.1	21.6
2005 4/.....	0.3	0.5	0.6	1.0	1.3	2.2	3.6	6.1	11.3	26.5
2004 4/.....	0.3	0.5	0.7	1.0	1.5	2.3	3.4	6.4	11.3	25.7
2003 4/.....	0.3	0.4	0.6	1.0	1.4	2.1	3.4	5.9	11.1	27.2
2002.....	0.3	0.4	0.5	0.9	1.0	1.7	3.3	5.6	10.0	28.3
2001.....	0.2	0.4	0.6	0.8	1.2	1.8	3.0	5.6	10.4	28.1
2000.....	0.2	0.3	0.4	0.8	1.2	1.7	3.1	5.2	11.0	29.7
1999.....	0.2	0.4	0.4	0.6	1.0	1.8	2.9	5.3	10.8	29.3

White, male

2006 4/.....	0.3	0.4	0.9	1.0	1.6	2.5	3.2	5.5	10.0	19.9
2005 4/.....	0.3	0.5	0.7	1.1	1.3	2.3	3.4	6.2	11.0	25.1
2004 4/.....	0.3	0.5	0.7	0.9	1.5	1.9	3.3	6.5	10.8	23.1
2003 4/.....	0.3	0.4	0.7	1.0	1.5	1.8	3.3	5.7	10.1	24.0
2002.....	0.3	0.5	0.5	1.0	0.9	1.6	3.6	5.7	10.7	26.1
2001.....	*	0.5	0.5	0.9	1.2	2.2	3.3	5.5	9.4	21.8
2000.....	*	0.3	0.4	0.8	1.4	1.2	3.1	4.9	10.4	25.8
1999.....	*	0.3	0.4	0.6	0.9	1.9	2.8	4.7	10.4	27.8

White, female

2006 4/.....	*	0.3	0.6	1.1	1.8	2.3	3.1	5.1	10.2	22.4
2005 4/.....	0.2	0.5	0.5	0.8	1.2	2.1	3.7	6.0	11.5	27.2
2004 4/.....	0.3	0.5	0.6	1.0	1.5	2.6	3.6	6.3	11.7	27.0
2003 4/.....	0.3	0.4	0.6	1.1	1.4	2.4	3.6	6.1	11.7	28.7
2002.....	0.3	0.3	0.5	0.9	1.1	1.8	3.1	5.5	9.6	29.3
2001.....	0.2	0.4	0.6	0.8	1.1	1.5	2.8	5.6	11.0	30.8
2000.....	0.2	0.2	0.4	0.7	1.1	2.1	3.2	5.5	11.4	31.2
1999.....	0.3	0.4	0.4	0.6	1.0	1.7	3.0	5.6	11.0	30.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XII. Diseases of the skin and subcutaneous tissue (L00-L98)

Black, both sexes

2006 4/.....	*	0.9	1.5	3.2	3.9	6.2	9.3	16.9	27.5	48.3
2005 4/.....	0.7	1.1	2.0	3.2	5.0	5.9	10.9	19.9	30.3	66.8
2004 4/.....	0.7	1.3	1.7	3.3	3.9	6.6	13.3	16.8	33.2	70.8
2003 4/.....	0.8	1.4	2.2	2.8	4.4	5.7	14.3	18.0	32.2	73.2
2002.....	1.1	1.0	2.0	3.2	4.4	5.1	12.5	19.1	35.3	92.9
2001.....	0.8	1.3	1.5	2.5	3.8	5.9	9.0	19.1	33.9	88.4
2000.....	0.8	1.4	1.1	2.5	3.7	6.0	12.5	19.9	36.1	97.3
1999.....	1.0	1.7	2.0	2.9	4.2	6.7	11.5	23.2	38.7	105.2

Black, male

2006 4/.....	*	*	*	3.0	3.7	7.3	9.8	15.2	24.6	43.8
2005 4/.....	*	*	2.2	3.6	4.9	5.5	12.0	20.6	29.5	47.4
2004 4/.....	*	*	*	2.6	*	6.1	12.9	16.4	30.8	57.6
2003 4/.....	*	*	*	*	5.6	6.6	17.7	16.4	34.7	65.7
2002.....	*	*	2.2	*	*	6.0	15.4	18.4	32.9	63.3
2001.....	*	*	*	*	*	6.6	8.1	24.6	34.5	71.5
2000.....	*	*	*	*	*	6.2	15.0	20.3	41.3	81.9
1999.....	*	2.2	*	*	*	6.8	12.2	24.2	39.2	85.5

Black, female

2006 4/.....	*	*	1.6	3.3	4.1	5.4	8.9	17.9	28.9	50.1
2005 4/.....	*	1.4	1.8	2.8	5.1	6.2	10.2	19.5	30.8	74.6
2004 4/.....	*	*	1.7	3.9	4.4	7.0	13.6	17.0	34.4	76.0
2003 4/.....	*	1.5	2.5	3.1	3.5	4.9	12.0	19.0	30.9	76.2
2002.....	1.4	*	1.8	3.7	5.1	4.4	10.5	19.5	36.6	104.3
2001.....	*	*	*	3.0	4.2	5.4	9.6	15.8	33.5	94.7
2000.....	*	2.0	*	2.7	3.6	5.8	10.9	19.7	33.5	103.0
1999.....	*	*	*	2.9	4.7	6.7	11.0	22.6	38.5	112.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Infections of skin and subcutaneous tissue (I00-L08)

All races 3/, both sexes

2006.....	0.2	0.2	0.5	0.8	0.9	1.4	1.5	3.0	5.1	9.7
2005.....	0.2	0.3	0.5	0.6	0.7	1.1	1.9	3.0	4.6	10.1
2004.....	0.2	0.3	0.4	0.6	0.8	1.2	1.7	3.0	4.9	10.4
2003.....	0.1	0.2	0.4	0.6	0.8	1.0	1.7	2.7	4.5	10.6
2002.....	0.1	0.2	0.3	0.5	0.6	0.8	1.3	2.2	3.6	9.3
2001.....	0.1	0.3	0.3	0.4	0.7	0.9	1.2	2.0	3.4	9.4
2000.....	0.1	0.1	0.2	0.4	0.6	0.7	1.4	2.0	3.4	9.3
1999.....	*	0.2	0.2	0.3	0.5	0.7	1.1	1.5	2.9	8.3

All races 3/, male

2006.....	0.3	0.3	0.6	0.7	0.8	1.7	1.6	3.1	4.8	9.0
2005.....	0.2	0.3	0.6	0.7	0.8	1.1	2.0	3.2	4.8	10.4
2004.....	0.2	0.4	0.5	0.6	0.8	1.1	1.8	3.3	5.0	10.3
2003.....	*	0.3	0.5	0.7	0.9	1.1	1.8	2.6	4.8	10.4
2002.....	0.2	0.3	0.4	0.6	0.6	0.9	1.5	2.2	4.1	9.1
2001.....	*	0.4	0.4	0.5	0.8	1.1	1.4	2.2	2.9	8.1
2000.....	*	*	*	0.6	0.7	0.7	1.6	2.1	4.0	8.9
1999.....	*	0.3	0.3	0.4	0.4	0.7	1.0	1.3	3.3	9.1

All races 3/, female

2006.....	*	*	0.3	0.8	0.9	1.2	1.5	2.9	5.3	10.1
2005.....	*	0.3	0.3	0.5	0.5	1.1	1.8	2.8	4.4	10.0
2004.....	*	0.2	0.3	0.5	0.9	1.3	1.6	2.8	4.9	10.4
2003.....	*	0.2	0.3	0.5	0.7	1.0	1.6	2.7	4.4	10.7
2002.....	*	*	0.2	0.4	0.5	0.8	1.2	2.2	3.2	9.4
2001.....	*	*	0.3	0.4	0.6	0.7	1.0	1.9	3.8	10.0
2000.....	*	*	*	0.3	0.5	0.7	1.2	1.9	3.1	9.5
1999.....	*	*	*	*	0.5	0.7	1.1	1.6	2.7	8.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Infections of skin and subcutaneous tissue (I00-L08)

White, both sexes

2006 4/.....	0.2	0.2	0.4	0.7	0.9	1.4	1.5	2.8	5.0	9.8
2005 4/.....	0.2	0.3	0.4	0.6	0.6	1.1	1.9	2.9	4.6	10.3
2004 4/.....	0.2	0.3	0.4	0.5	0.8	1.2	1.6	3.1	4.9	10.4
2003 4/.....	0.1	0.2	0.4	0.6	0.8	1.0	1.6	2.7	4.5	10.8
2002.....	0.1	0.2	0.3	0.5	0.6	0.7	1.2	2.1	3.6	9.7
2001.....	*	0.2	0.3	0.4	0.6	0.9	1.2	2.0	3.4	9.6
2000.....	*	0.1	0.2	0.4	0.6	0.7	1.3	1.9	3.4	9.3
1999.....	*	0.2	0.2	0.3	0.4	0.7	1.1	1.5	3.0	8.3

White, male

2006 4/.....	0.3	0.3	0.7	0.7	0.9	1.6	1.4	3.0	4.8	9.3
2005 4/.....	*	0.3	0.5	0.7	0.8	1.1	1.9	3.2	4.7	10.8
2004 4/.....	*	0.4	0.4	0.5	0.8	1.0	1.7	3.4	5.1	10.8
2003 4/.....	*	0.2	0.5	0.7	0.9	1.0	1.7	2.6	4.8	10.6
2002.....	*	0.3	0.4	0.6	0.6	0.8	1.4	2.2	4.3	9.4
2001.....	*	0.3	0.3	0.5	0.8	1.1	1.5	2.1	3.0	8.4
2000.....	*	*	*	0.6	0.8	1.5	2.0	4.0	8.7	
1999.....	*	*	*	*	*	0.8	1.0	1.2	3.3	8.9

White, female

2006 4/.....	*	*	*	0.7	0.9	1.2	1.5	2.6	5.1	10.1
2005 4/.....	*	0.3	0.3	0.5	0.4	1.1	1.9	2.7	4.5	10.0
2004 4/.....	*	0.2	0.3	0.5	0.8	1.3	1.6	2.9	4.8	10.2
2003 4/.....	*	*	0.3	0.5	0.6	1.0	1.4	2.8	4.4	10.9
2002.....	*	*	*	0.4	0.5	0.7	1.1	2.1	3.1	9.8
2001.....	*	*	0.3	*	0.4	0.7	1.0	2.0	3.6	10.1
2000.....	*	*	*	*	0.5	0.7	1.2	1.8	3.1	9.5
1999.....	*	*	*	*	0.4	0.6	1.1	1.7	2.8	8.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)

All races 3/, both sexes

2006.....	0.1	0.2	0.4	0.5	1.0	1.3	2.1	3.2	6.2	13.6
2005.....	0.2	0.3	0.3	0.5	0.9	1.4	2.4	4.2	8.1	19.1
2004.....	0.2	0.3	0.4	0.6	0.9	1.5	2.6	4.1	7.8	18.4
2003.....	0.2	0.3	0.4	0.6	0.9	1.4	2.7	4.1	8.0	19.9
2002.....	0.2	0.3	0.4	0.6	0.8	1.2	2.8	4.3	8.2	23.5
2001.....	0.2	0.3	0.3	0.6	0.8	1.3	2.3	4.5	8.5	22.7
2000.....	0.2	0.3	0.3	0.5	0.8	1.4	2.6	4.4	9.3	25.4
1999.....	0.2	0.3	0.4	0.5	0.8	1.5	2.5	5.1	9.7	26.6

All races 3/, male

2006.....	*	*	0.3	0.5	0.9	1.2	2.1	3.0	5.9	12.3
2005.....	*	0.2	0.3	0.6	0.8	1.5	2.2	3.9	7.4	15.9
2004.....	*	0.2	0.3	0.5	0.9	1.1	2.3	3.7	6.9	14.9
2003.....	0.2	0.2	0.3	0.4	1.0	1.2	2.6	3.6	6.9	16.2
2002.....	*	0.2	0.3	0.5	0.5	1.1	3.0	4.3	7.8	19.4
2001.....	*	0.2	0.3	0.5	0.6	1.4	2.2	4.5	7.9	16.8
2000.....	0.2	*	*	0.4	0.9	0.9	2.4	4.0	8.3	20.8
1999.....	*	0.2	0.3	0.4	0.7	1.6	2.4	4.8	8.9	22.6

All races 3/, female

2006.....	*	0.3	0.4	0.6	1.0	1.3	2.2	3.2	6.5	14.2
2005.....	0.2	0.3	0.3	0.5	1.1	1.4	2.5	4.4	8.5	20.6
2004.....	0.2	0.3	0.4	0.8	0.9	1.7	2.8	4.3	8.3	20.0
2003.....	0.3	0.4	0.5	0.7	0.9	1.6	2.7	4.4	8.6	21.5
2002.....	0.3	0.3	0.5	0.7	1.0	1.3	2.7	4.3	8.5	25.3
2001.....	0.2	0.3	0.4	0.6	0.9	1.2	2.4	4.5	8.9	25.2
2000.....	0.2	0.4	0.4	0.6	0.8	1.7	2.7	4.7	9.9	27.3
1999.....	0.3	0.3	0.4	0.6	0.8	1.5	2.5	5.4	10.2	28.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)

White, both sexes

2006 4/.....	*	0.1	0.3	0.4	0.8	1.0	1.6	2.5	5.1	11.8
2005 4/.....	*	0.2	0.2	0.4	0.7	1.1	1.7	3.2	6.7	16.3
2004 4/.....	0.1	0.2	0.3	0.5	0.7	1.1	1.8	3.3	6.4	15.3
2003 4/.....	0.2	0.2	0.3	0.4	0.7	1.1	1.9	3.2	6.5	16.4
2002.....	0.2	0.2	0.2	0.4	0.5	1.0	2.1	3.5	6.4	18.7
2001.....	0.1	0.2	0.3	0.4	0.6	1.0	1.8	3.5	7.0	18.5
2000.....	0.1	0.2	0.2	0.3	0.6	1.0	1.8	3.3	7.6	20.4
1999.....	0.1	0.2	0.2	0.3	0.5	1.1	1.8	3.8	7.7	21.0

White, male

2006 4/.....	*	*	0.2	0.3	0.7	0.9	1.7	2.5	5.2	10.6
2005 4/.....	*	*	*	0.4	0.5	1.2	1.4	3.1	6.3	14.3
2004 4/.....	*	*	0.2	0.4	0.8	0.9	1.6	3.1	5.7	12.3
2003 4/.....	*	*	*	*	0.7	0.9	1.6	3.1	5.3	13.5
2002.....	*	*	*	0.4	*	0.8	2.2	3.6	6.4	16.7
2001.....	*	*	*	0.4	0.4	1.1	1.8	3.3	6.4	13.3
2000.....	*	*	*	*	0.7	0.6	1.5	2.9	6.5	17.0
1999.....	*	*	*	*	0.5	1.1	1.7	3.6	7.0	18.9

White, female

2006 4/.....	*	*	0.4	0.4	0.8	1.1	1.5	2.5	5.0	12.3
2005 4/.....	*	0.2	*	0.3	0.9	1.0	1.9	3.3	7.0	17.2
2004 4/.....	*	0.2	0.3	0.5	0.7	1.3	2.0	3.4	6.9	16.7
2003 4/.....	*	0.2	0.3	0.5	0.7	1.3	2.2	3.3	7.3	17.8
2002.....	0.2	*	0.3	0.5	0.6	1.1	2.0	3.4	6.5	19.6
2001.....	*	0.2	0.3	0.5	0.7	0.9	1.8	3.6	7.4	20.6
2000.....	*	*	0.3	0.4	0.6	1.4	2.0	3.6	8.3	21.7
1999.....	0.2	0.3	0.3	0.4	0.6	1.1	1.9	3.9	8.1	21.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified diseases of skin and subcutaneous tissue (L10-L98)

Black, both sexes

2006 4/.....	*	*	*	1.6	3.3	4.1	6.9	11.4	20.6	37.9
2005 4/.....	*	*	1.2	2.1	3.2	4.5	8.7	15.5	25.5	56.7
2004 4/.....	*	0.8	1.1	2.4	2.7	4.9	10.6	13.9	26.7	58.1
2003 4/.....	*	0.9	1.5	2.2	3.1	4.1	11.1	14.8	27.4	63.6
2002.....	0.7	*	1.4	2.1	3.6	3.3	10.3	15.3	31.7	85.8
2001.....	*	*	1.0	1.7	2.6	4.5	8.2	16.8	29.3	79.2
2000.....	*	1.1	*	1.8	2.9	4.6	10.9	17.2	32.2	87.3
1999.....	0.8	0.9	1.5	1.9	3.1	5.9	10.1	22.0	36.4	96.3

Black, male

2006 4/.....	*	*	*	*	*	*	6.5	10.7	17.8	36.6
2005 4/.....	*	*	*	*	*	*	9.2	16.1	23.2	40.9
2004 4/.....	*	*	*	*	*	*	10.7	12.7	25.1	48.7
2003 4/.....	*	*	*	*	4.0	*	15.1	12.8	30.3	55.3
2002.....	*	*	*	*	*	*	12.7	15.2	29.8	56.8
2001.....	*	*	*	*	*	5.3	7.4	21.4	32.1	65.8
2000.....	*	*	*	*	*	*	12.7	18.0	36.2	72.7
1999.....	*	*	*	*	*	6.8	11.2	21.4	35.8	73.8

Black, female

2006 4/.....	*	*	*	*	3.2	3.8	7.2	11.8	22.1	38.4
2005 4/.....	*	*	*	*	3.0	4.6	8.4	15.2	26.7	63.0
2004 4/.....	*	*	*	3.0	2.9	5.6	10.5	14.6	27.6	61.8
2003 4/.....	*	*	2.0	2.7	*	3.7	8.4	16.0	25.9	66.8
2002.....	*	*	*	2.7	4.0	*	8.7	15.4	32.6	96.9
2001.....	*	*	*	*	*	3.9	8.7	14.1	27.8	84.2
2000.....	*	1.7	*	*	*	4.8	9.8	16.8	30.1	92.7
1999.....	*	*	*	*	*	5.2	9.5	22.3	36.7	104.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

All races 3/, both sexes

2006.....	1.6	2.3	2.9	4.5	6.3	9.2	14.9	23.2	36.4	75.8
2005.....	1.6	2.2	3.2	4.6	6.4	9.5	14.8	23.5	40.3	84.4
2004.....	1.5	2.2	3.3	4.6	6.6	9.5	15.0	24.3	40.9	85.9
2003.....	1.6	2.2	3.3	4.4	7.0	9.8	15.9	25.3	40.5	90.4
2002.....	1.5	2.3	3.0	4.3	7.0	10.1	15.9	25.6	40.2	94.9
2001.....	1.7	2.1	3.1	4.2	6.6	10.1	15.8	25.4	42.0	93.4
2000.....	1.7	2.3	3.0	4.3	6.7	10.3	15.2	25.9	41.8	94.0
1999.....	1.6	2.4	3.1	4.3	6.7	10.1	15.5	25.5	39.3	91.2

All races 3/, male

2006.....	1.0	1.8	2.3	3.5	4.9	7.9	11.6	17.7	26.1	54.6
2005.....	1.1	1.6	2.5	3.4	5.0	7.2	11.6	19.2	31.5	58.5
2004.....	0.9	1.5	2.7	3.2	5.4	7.2	11.3	19.5	31.0	59.1
2003.....	1.2	1.5	2.4	3.3	5.1	8.4	12.6	20.8	30.1	59.9
2002.....	1.0	1.7	2.3	3.0	5.3	7.8	13.0	19.4	31.3	63.5
2001.....	1.1	1.5	2.4	3.2	4.7	7.1	12.3	20.1	32.4	63.2
2000.....	1.2	1.6	1.8	3.4	4.8	7.7	12.1	20.6	33.0	67.5
1999.....	1.1	1.5	2.1	3.1	4.7	6.9	12.5	20.2	30.9	63.7

All races 3/, female

2006.....	2.2	2.8	3.6	5.5	7.6	10.4	17.5	27.3	42.9	85.8
2005.....	2.2	2.9	3.8	5.7	7.6	11.6	17.4	26.6	45.8	96.2
2004.....	2.1	2.9	3.9	5.9	7.6	11.6	18.0	27.8	47.0	98.0
2003.....	2.1	3.0	4.2	5.5	8.8	11.0	18.5	28.5	46.8	103.8
2002.....	2.1	3.0	3.8	5.6	8.6	12.2	18.2	30.1	45.6	108.5
2001.....	2.2	2.8	3.7	5.2	8.3	12.7	18.5	29.1	47.8	106.0
2000.....	2.2	3.0	4.1	5.2	8.4	12.4	17.6	29.6	46.9	104.8
1999.....	2.1	3.3	4.1	5.5	8.5	12.8	18.0	29.3	44.3	102.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

White, both sexes

2006 4/.....	1.2	1.9	2.5	4.0	6.1	8.9	15.0	23.8	37.4	79.0
2005 4/.....	1.2	1.7	2.8	4.0	5.8	9.4	14.9	23.9	41.4	87.8
2004 4/.....	1.1	1.6	2.7	4.0	6.2	9.2	15.0	24.4	41.8	88.8
2003 4/.....	1.4	1.7	2.8	4.0	6.6	9.5	15.7	25.5	40.9	92.9
2002.....	1.1	1.8	2.7	3.9	6.6	9.9	15.8	25.9	40.7	98.2
2001.....	1.2	1.7	2.6	3.9	6.1	9.9	15.5	25.8	43.0	95.8
2000.....	1.2	1.7	2.5	3.8	6.3	10.2	15.0	26.0	42.5	96.9
1999.....	1.1	1.8	2.6	3.8	6.4	9.7	15.6	25.8	40.6	94.4

White, male

2006 4/.....	0.8	1.6	2.1	3.1	4.8	7.8	11.4	17.5	26.5	56.7
2005 4/.....	0.9	1.4	2.3	3.0	4.5	7.0	11.5	19.2	32.2	60.9
2004 4/.....	0.7	1.1	2.2	2.9	5.2	6.9	10.9	19.3	31.4	59.8
2003 4/.....	1.1	1.2	2.1	2.9	4.7	7.8	12.0	20.2	29.8	60.1
2002.....	0.8	1.4	2.1	2.7	5.0	7.3	12.7	19.4	31.0	64.3
2001.....	0.8	1.2	2.1	2.8	4.4	6.8	11.9	20.3	32.7	64.3
2000.....	0.9	1.3	1.6	3.2	4.5	7.6	11.3	20.2	32.9	67.3
1999.....	0.8	1.2	1.9	2.7	4.6	6.6	12.6	19.4	31.9	64.4

White, female

2006 4/.....	1.5	2.1	2.8	4.9	7.3	9.9	18.1	28.4	44.3	89.4
2005 4/.....	1.6	2.1	3.2	5.0	7.1	11.5	17.8	27.4	47.2	100.2
2004 4/.....	1.5	2.2	3.1	5.2	7.1	11.2	18.5	28.2	48.2	101.8
2003 4/.....	1.8	2.1	3.5	5.1	8.3	10.9	18.7	29.2	47.8	107.4
2002.....	1.4	2.3	3.2	5.0	8.1	12.2	18.4	30.6	46.6	112.9
2001.....	1.6	2.2	3.1	4.8	7.6	12.6	18.4	29.6	49.3	109.0
2000.....	1.5	2.0	3.4	4.4	7.9	12.6	17.9	30.1	48.2	108.9
1999.....	1.4	2.5	3.3	4.9	8.0	12.4	18.1	30.2	45.7	106.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XIII. Diseases of the musculoskeletal system and connective tissue (M00-M99)

Black, both sexes

2006 4/.....	4.6	5.6	7.1	9.3	9.8	12.6	16.0	21.8	33.1	52.9
2005 4/.....	4.6	5.9	6.5	9.7	11.9	12.4	16.1	24.7	35.4	58.5
2004 4/.....	4.0	6.6	8.2	9.4	11.2	14.5	17.7	27.1	36.4	64.9
2003 4/.....	3.4	6.6	7.7	8.5	11.8	14.2	20.2	27.2	38.8	68.0
2002.....	4.5	6.0	6.5	8.9	11.4	13.5	18.8	25.6	39.0	67.4
2001.....	4.9	5.5	7.5	8.0	11.2	14.1	19.6	25.0	36.0	75.5
2000.....	5.4	7.2	6.9	8.9	11.1	12.4	18.7	26.8	37.6	68.2
1999.....	5.4	6.8	7.3	8.9	10.5	15.3	17.5	27.0	27.2	60.3

Black, male

2006 4/.....	2.6	3.6	4.9	7.2	8.0	11.0	16.0	24.2	28.8	44.7
2005 4/.....	2.8	3.5	4.4	7.1	10.5	10.0	13.6	24.1	28.8	39.0
2004 4/.....	2.1	4.5	6.5	5.8	9.2	11.6	18.5	27.8	30.8	60.6
2003 4/.....	2.0	3.5	4.9	6.7	10.0	15.8	21.0	30.6	38.4	58.4
2002.....	2.5	3.9	4.3	6.7	9.1	12.5	18.7	21.6	39.7	61.1
2001.....	3.4	4.2	5.7	6.6	9.2	11.7	17.2	22.8	36.9	59.0
2000.....	3.6	4.2	3.7	6.3	8.3	12.0	21.7	25.5	39.6	78.4
1999.....	3.4	4.0	4.2	6.4	6.8	12.2	14.2	33.7	20.9	63.3

Black, female

2006 4/.....	6.3	7.3	9.0	11.0	11.3	13.8	16.1	20.4	35.4	56.3
2005 4/.....	6.1	8.1	8.4	11.9	13.0	14.3	17.7	25.1	38.9	66.4
2004 4/.....	5.8	8.3	9.7	12.4	12.8	16.5	17.1	26.7	39.3	66.5
2003 4/.....	4.7	9.4	10.0	9.9	13.2	13.0	19.7	25.2	39.1	71.7
2002.....	6.3	7.8	8.4	10.6	13.2	14.2	18.8	28.0	38.6	69.8
2001.....	6.3	6.6	8.9	9.1	12.8	16.0	21.3	26.3	35.6	81.7
2000.....	7.1	9.8	9.6	11.1	13.2	12.7	16.6	27.6	36.5	64.4
1999.....	7.2	9.3	10.0	10.9	13.5	17.5	19.6	22.9	30.2	59.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)

All races 3/, both sexes

2006.....	*	0.2	0.3	0.5	1.1	2.1	3.8	6.7	9.2	10.3
2005.....	*	0.1	0.3	0.6	1.0	2.1	3.7	6.5	10.0	11.5
2004.....	*	0.2	0.3	0.7	1.1	2.3	4.1	6.8	9.6	11.9
2003.....	0.1	0.2	0.3	0.6	1.3	2.2	4.2	7.0	10.9	12.8
2002.....	*	0.2	0.3	0.7	1.4	2.5	4.2	8.0	10.6	14.2
2001.....	*	0.1	0.3	0.7	1.4	2.5	4.3	7.6	10.5	12.7
2000.....	*	0.2	0.3	0.6	1.4	2.7	4.4	7.5	11.6	12.9
1999.....	*	0.2	0.3	0.8	1.6	2.5	4.5	8.7	10.6	13.9

All races 3/, male

2006.....	*	*	0.2	0.4	0.7	1.4	2.3	3.3	5.2	5.8
2005.....	*	*	0.2	0.4	0.6	1.2	2.5	4.2	5.4	5.4
2004.....	*	*	*	0.4	0.9	1.6	2.2	4.1	4.6	6.5
2003.....	*	*	*	0.4	1.0	1.3	2.6	4.5	5.6	6.5
2002.....	*	*	*	0.5	1.0	1.7	2.6	4.5	6.6	7.6
2001.....	*	*	*	0.6	0.9	1.5	2.6	4.6	5.8	5.5
2000.....	*	*	*	0.4	0.9	1.7	3.1	4.4	6.6	6.9
1999.....	*	*	*	0.5	1.1	1.6	3.1	5.0	5.5	7.2

All races 3/, female

2006.....	*	0.2	0.4	0.7	1.5	2.7	5.1	9.1	11.7	12.5
2005.....	*	0.2	0.4	0.8	1.5	2.8	4.7	8.2	12.8	14.3
2004.....	*	0.2	0.4	0.9	1.4	2.9	5.6	8.7	12.7	14.3
2003.....	*	0.2	0.4	0.7	1.6	2.9	5.5	8.8	14.1	15.6
2002.....	*	0.2	0.4	0.9	1.8	3.2	5.4	10.4	13.1	17.0
2001.....	*	0.2	0.5	0.9	1.8	3.3	5.5	9.7	13.3	15.7
2000.....	*	0.3	0.5	0.8	1.9	3.5	5.4	9.7	14.6	15.3
1999.....	*	0.3	0.3	1.0	2.0	3.3	5.6	11.3	13.5	16.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)										
White, both sexes										
2006 4/.....	*	0.2	0.4	0.5	1.1	2.1	4.0	7.1	9.5	10.7
2005 4/.....	*	0.2	0.3	0.7	1.0	2.2	3.9	6.8	10.5	12.0
2004 4/.....	*	0.2	0.3	0.7	1.2	2.3	4.3	7.0	9.8	12.1
2003 4/.....	*	0.2	0.3	0.6	1.4	2.2	4.5	7.3	11.1	13.2
2002.....	*	0.2	0.3	0.7	1.4	2.6	4.3	8.2	10.9	14.8
2001.....	*	0.1	0.3	0.7	1.4	2.7	4.4	8.0	10.9	12.8
2000.....	*	0.2	0.3	0.6	1.5	2.8	4.7	7.8	12.2	13.4
1999.....	*	0.2	0.3	0.8	1.7	2.6	4.8	9.0	11.0	14.9
White, male										
2006 4/.....	*	*	0.3	0.4	0.7	1.4	2.3	3.5	5.3	6.4
2005 4/.....	*	*	0.3	0.4	0.6	1.3	2.7	4.4	5.4	5.7
2004 4/.....	*	*	*	0.4	0.9	1.6	2.3	4.3	4.8	6.6
2003 4/.....	*	*	*	0.4	1.1	1.5	2.7	4.6	5.7	6.7
2002.....	*	*	*	0.5	1.0	1.7	2.7	4.7	6.6	8.0
2001.....	*	*	*	0.6	0.9	1.6	2.7	4.9	5.8	5.5
2000.....	*	*	*	0.5	0.9	1.8	3.2	4.5	6.9	7.3
1999.....	*	*	*	0.5	1.2	1.6	3.3	5.1	5.8	7.8
White, female										
2006 4/.....	*	0.2	0.4	0.7	1.4	2.8	5.4	9.8	12.1	12.7
2005 4/.....	*	*	0.4	0.9	1.4	3.1	4.9	8.7	13.7	14.9
2004 4/.....	*	0.2	0.4	1.0	1.4	3.0	5.9	8.9	12.9	14.6
2003 4/.....	*	*	0.5	0.8	1.7	2.9	5.9	9.3	14.4	16.0
2002.....	*	0.3	0.4	0.9	1.7	3.5	5.6	10.8	13.6	17.7
2001.....	*	*	0.5	0.9	1.8	3.6	5.8	10.2	14.0	15.9
2000.....	*	0.2	0.5	0.8	2.0	3.8	5.8	10.1	15.3	15.8
1999.....	*	0.3	0.4	1.0	2.2	3.5	6.0	11.8	14.1	17.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Rheumatoid arthritis and related inflammatory polyarthropathies (M05-M08)

Black, both sexes

2006 4/.....	*	*	*	*	*	*	2.9	3.4	7.1	7.3
2005 4/.....	*	*	*	*	1.8	*	3.0	4.2	*	8.3
2004 4/.....	*	*	*	*	*	2.1	3.0	5.6	7.4	9.3
2003 4/.....	*	*	*	*	*	*	2.9	4.9	9.6	9.6
2002.....	*	*	*	*	*	*	3.4	5.5	8.1	8.0
2001.....	*	*	*	*	*	*	3.2	4.0	6.5	12.0
2000.....	*	*	*	*	*	*	*	4.6	5.9	7.8
1999.....	*	*	*	*	*	*	*	6.4	*	*

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	*	*	*	*	*	*	*	*	8.7	10.2
2005 4/.....	*	*	*	*	*	*	*	5.3	*	10.9
2004 4/.....	*	*	*	*	*	*	*	6.7	10.2	11.0
2003 4/.....	*	*	*	*	*	*	*	5.4	11.6	11.3
2002.....	*	*	*	*	*	*	*	7.1	9.2	9.9
2001.....	*	*	*	*	*	*	*	*	*	14.7
2000.....	*	*	*	*	*	*	*	6.0	*	9.4
1999.....	*	*	*	*	*	*	*	6.3	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Systemic lupus erythematosus (M32)										
All races 3/, both sexes										
2006.....	0.5	0.5	0.6	0.8	0.7	0.7	0.9	1.1	1.3	1.2
2005.....	0.6	0.6	0.7	0.8	0.9	0.9	1.1	1.2	1.4	1.3
2004.....	0.5	0.5	0.8	0.8	0.8	0.7	1.0	1.2	1.5	1.1
2003.....	0.5	0.6	0.7	0.7	0.9	0.8	0.9	1.4	1.5	1.1
2002.....	0.5	0.6	0.6	0.7	1.0	0.9	1.0	1.3	1.4	1.3
2001.....	0.5	0.6	0.7	0.7	0.9	0.9	1.2	1.4	1.6	1.2
2000.....	0.6	0.6	0.7	0.8	1.0	1.0	0.9	1.4	1.5	1.2
1999.....	0.6	0.7	0.7	0.7	0.8	1.0	1.2	1.3	1.3	1.4
All races 3/, male										
2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	0.2	0.2	*	*	*	*	*	*
2004.....	*	*	0.3	*	*	*	*	*	*	*
2003.....	0.2	*	0.2	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	0.2	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2006.....	0.9	0.9	1.0	1.3	1.2	1.0	1.3	1.7	1.8	1.7
2005.....	1.0	1.0	1.2	1.4	1.4	1.5	1.6	1.8	2.0	1.8
2004.....	0.8	0.8	1.2	1.3	1.3	1.2	1.4	1.7	2.0	1.3
2003.....	0.8	0.9	1.2	1.3	1.5	1.2	1.4	2.2	2.1	1.3
2002.....	0.8	1.1	0.9	1.2	1.7	1.4	1.5	1.7	1.9	1.7
2001.....	0.8	1.1	1.1	1.1	1.5	1.5	1.8	2.2	2.0	1.3
2000.....	1.1	1.1	1.2	1.3	1.5	1.5	1.2	2.0	2.0	1.4
1999.....	1.0	1.2	1.2	1.2	1.4	1.7	1.8	1.8	1.7	1.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Systemic lupus erythematosus (M32)										
White, both sexes										
2006 4/.....	0.3	0.3	0.4	0.6	0.6	0.6	0.8	1.1	1.3	1.3
2005 4/.....	0.3	0.3	0.5	0.6	0.7	0.8	1.0	1.1	1.4	1.2
2004 4/.....	0.3	0.3	0.5	0.6	0.6	0.6	0.9	1.1	1.4	1.2
2003 4/.....	0.4	0.3	0.5	0.5	0.8	0.7	0.9	1.4	1.5	1.1
2002.....	0.3	0.4	0.4	0.5	0.7	0.7	1.0	1.3	1.4	1.4
2001.....	0.2	0.4	0.5	0.6	0.7	0.8	1.1	1.4	1.7	1.2
2000.....	0.4	0.3	0.5	0.6	0.8	0.9	0.8	1.4	1.6	1.3
1999.....	0.3	0.5	0.5	0.6	0.7	0.8	1.1	1.3	1.3	1.5
White, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
White, female										
2006 4/.....	0.5	0.5	0.7	1.0	1.0	0.8	1.2	1.6	1.9	1.8
2005 4/.....	0.6	0.5	0.9	1.0	1.2	1.3	1.5	1.7	2.0	1.7
2004 4/.....	0.5	0.5	0.8	1.0	0.9	1.0	1.2	1.6	1.9	1.4
2003 4/.....	0.6	0.5	0.8	1.0	1.3	0.9	1.2	2.2	2.0	1.3
2002.....	0.5	0.7	0.7	0.9	1.3	1.0	1.4	1.7	1.9	1.8
2001.....	0.4	0.8	0.8	1.0	1.2	1.3	1.7	2.1	2.0	1.1
2000.....	0.6	0.6	0.9	1.0	1.3	1.3	1.0	1.9	2.1	1.5
1999.....	0.6	0.8	0.8	0.9	1.2	1.3	1.6	1.9	1.7	1.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Systemic lupus erythematosus (M32)										
Black, both sexes										
2006 4/.....	1.7	1.8	1.9	2.3	1.5	*	*	*	*	*
2005 4/.....	2.0	2.4	2.0	2.4	2.3	*	*	*	*	*
2004 4/.....	1.7	2.0	2.6	2.6	2.9	*	*	*	*	*
2003 4/.....	1.4	2.4	2.3	2.2	1.9	2.1	*	*	*	*
2002.....	1.8	2.2	1.7	1.9	3.0	2.9	*	*	*	*
2001.....	2.1	2.0	2.0	1.8	2.4	*	*	*	*	*
2000.....	2.2	2.6	2.0	2.5	2.3	*	*	*	*	*
1999.....	2.4	2.4	1.9	2.1	*	3.1	*	*	*	*
Black, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
Black, female										
2006 4/.....	2.9	3.0	3.1	3.5	*	*	*	*	*	*
2005 4/.....	3.6	3.8	3.3	4.0	3.8	*	*	*	*	*
2004 4/.....	2.6	3.3	4.0	4.2	4.5	*	*	*	*	*
2003 4/.....	2.3	3.8	3.8	3.6	*	*	*	*	*	*
2002.....	3.1	4.0	2.7	3.5	5.1	4.4	*	*	*	*
2001.....	3.4	3.3	3.2	2.7	3.5	*	*	*	*	*
2000.....	3.6	4.4	3.4	3.6	3.8	*	*	*	*	*
1999.....	3.9	3.9	3.3	3.2	*	4.8	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)										
All races 3/, both sexes										
2006.....	0.6	1.0	1.1	1.8	2.4	3.7	5.3	7.0	10.3	21.7
2005.....	0.6	0.9	1.2	1.9	2.5	3.9	5.0	7.4	11.6	24.9
2004.....	0.6	0.9	1.3	1.9	2.6	3.8	5.0	7.7	12.1	24.5
2003.....	0.7	0.9	1.3	1.8	2.8	3.8	5.6	8.3	10.9	25.9
2002.....	0.7	0.9	1.4	1.7	2.9	3.9	5.6	8.1	11.9	27.2
2001.....	0.7	0.9	1.3	1.7	2.7	4.1	5.9	8.2	13.0	27.7
2000.....	0.7	1.0	1.3	1.8	2.8	4.0	6.0	8.7	13.1	28.0
1999.....	0.7	1.1	1.5	1.8	2.9	4.2	6.0	7.9	11.9	27.6
All races 3/, male										
2006.....	0.4	0.8	0.9	1.3	1.6	2.9	3.5	5.0	6.8	15.8
2005.....	0.5	0.6	0.9	1.3	2.0	2.9	3.4	5.5	8.9	19.0
2004.....	0.4	0.6	1.0	1.2	2.0	2.5	3.4	5.7	9.4	17.5
2003.....	0.5	0.6	0.9	1.3	1.8	2.9	4.2	7.2	8.4	19.4
2002.....	0.4	0.7	0.9	1.3	2.0	2.5	4.2	6.3	9.6	18.6
2001.....	0.5	0.8	1.0	1.1	2.0	2.3	4.6	6.5	10.6	20.5
2000.....	0.6	0.7	0.8	1.4	1.9	2.8	4.0	7.3	11.2	21.1
1999.....	0.5	0.7	1.1	1.3	1.9	2.8	4.4	6.5	10.0	22.5
All races 3/, female										
2006.....	0.9	1.1	1.4	2.3	3.1	4.4	6.7	8.5	12.4	24.5
2005.....	0.8	1.2	1.6	2.5	2.9	4.7	6.3	8.8	13.3	27.7
2004.....	0.9	1.2	1.7	2.5	3.2	4.9	6.4	9.0	13.7	27.7
2003.....	0.9	1.2	1.7	2.3	3.8	4.5	6.7	9.0	12.4	28.8
2002.....	0.9	1.2	1.9	2.2	3.6	5.1	6.7	9.4	13.3	31.0
2001.....	1.0	1.1	1.5	2.2	3.3	5.6	7.0	9.4	14.4	30.7
2000.....	0.9	1.2	1.7	2.2	3.5	5.1	7.5	9.6	14.3	30.8
1999.....	0.8	1.4	1.9	2.3	3.8	5.4	7.2	8.8	13.0	29.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)										
White, both sexes										
2006 4/.....	0.4	0.7	0.9	1.6	2.3	3.5	5.5	7.3	10.9	22.8
2005 4/.....	0.5	0.7	1.0	1.6	2.4	3.8	5.2	7.7	12.0	26.0
2004 4/.....	0.5	0.6	1.1	1.6	2.5	3.6	5.1	7.7	12.5	25.5
2003 4/.....	0.6	0.6	1.0	1.6	2.6	3.7	5.5	8.3	11.1	26.4
2002.....	0.5	0.7	1.2	1.6	2.8	4.0	5.5	8.3	12.1	28.0
2001.....	0.5	0.7	1.0	1.6	2.5	3.9	5.9	8.5	13.4	28.6
2000.....	0.5	0.7	1.0	1.6	2.6	4.1	5.9	8.8	13.4	28.8
1999.....	0.5	0.7	1.1	1.6	2.6	4.1	5.9	8.0	12.2	28.4
White, male										
2006 4/.....	0.3	0.6	0.8	1.1	1.5	2.7	3.7	5.0	7.1	16.8
2005 4/.....	0.3	0.5	0.8	1.1	1.8	2.9	3.5	5.9	9.0	19.6
2004 4/.....	0.3	0.3	0.9	1.0	1.9	2.4	3.4	5.8	9.8	18.0
2003 4/.....	0.5	0.5	0.7	1.1	1.6	2.7	4.0	7.1	8.2	19.9
2002.....	0.3	0.5	0.8	1.1	1.9	2.5	4.0	6.5	9.7	19.1
2001.....	0.3	0.5	0.8	1.0	1.9	2.2	4.5	6.7	10.8	21.1
2000.....	0.4	0.6	0.7	1.3	1.7	2.7	3.8	7.2	11.1	20.6
1999.....	0.3	0.5	0.9	1.1	1.8	2.7	4.4	6.4	10.4	23.3
White, female										
2006 4/.....	0.6	0.8	1.1	2.1	3.1	4.1	7.0	9.0	13.2	25.6
2005 4/.....	0.6	0.8	1.3	2.1	2.8	4.5	6.6	9.0	13.8	29.0
2004 4/.....	0.6	0.8	1.4	2.1	3.1	4.7	6.5	9.1	14.3	28.9
2003 4/.....	0.7	0.8	1.4	2.1	3.5	4.5	6.7	9.3	12.8	29.2
2002.....	0.6	0.9	1.6	2.0	3.7	5.3	6.8	9.6	13.6	31.9
2001.....	0.8	0.9	1.2	2.1	3.1	5.5	7.0	9.8	14.9	31.7
2000.....	0.6	0.8	1.4	1.9	3.4	5.3	7.5	10.0	14.7	32.1
1999.....	0.6	1.0	1.4	2.1	3.4	5.3	7.2	9.2	13.3	30.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other arthropathies and related disorders (M00,M02,M10-M31,M33-M35)

Black, both sexes

2006 4/.....	2.0	2.7	2.7	3.3	3.6	5.6	4.3	4.5	6.4	14.8
2005 4/.....	1.8	2.4	2.7	4.5	4.0	4.8	4.1	6.5	10.0	16.8
2004 4/.....	1.7	3.1	3.3	3.8	4.2	5.3	5.4	6.9	8.9	16.6
2003 4/.....	1.5	2.8	3.6	3.6	5.1	5.2	6.9	8.7	9.9	20.4
2002.....	2.1	2.5	3.0	3.6	4.1	4.4	7.0	7.4	10.7	21.7
2001.....	2.2	2.7	3.6	2.7	3.7	6.0	6.1	6.4	11.6	22.4
2000.....	2.3	3.0	3.3	3.6	4.4	4.0	7.9	7.5	12.1	21.9
1999.....	2.4	3.3	4.1	3.7	5.9	5.3	7.0	7.3	8.7	20.4

Black, male

2006 4/.....	*	2.0	*	2.4	3.7	5.2	*	*	*	*
2005 4/.....	1.7	1.5	1.9	2.8	3.8	*	*	*	*	*
2004 4/.....	*	2.3	2.4	*	*	*	*	*	*	*
2003 4/.....	*	*	2.4	3.1	*	5.7	*	10.5	*	*
2002.....	*	2.0	2.2	3.6	4.2	*	*	*	*	*
2001.....	1.6	2.6	2.9	*	*	*	*	*	*	*
2000.....	1.8	*	*	*	4.1	*	*	*	*	27.7
1999.....	2.0	2.3	2.5	*	*	*	*	9.5	*	*

Black, female

2006 4/.....	2.8	3.2	3.6	4.1	3.6	5.9	5.3	*	7.6	17.2
2005 4/.....	1.9	3.2	3.4	5.9	4.1	5.6	5.2	9.3	11.1	18.8
2004 4/.....	2.5	3.7	4.1	4.9	4.7	6.7	5.9	8.1	9.8	17.7
2003 4/.....	2.0	4.2	4.5	4.0	6.2	4.9	7.3	7.6	9.7	23.5
2002.....	2.8	3.0	3.8	3.5	4.0	4.8	7.4	8.8	11.9	24.6
2001.....	2.7	2.8	4.2	3.3	4.5	7.3	7.2	6.7	12.7	24.4
2000.....	2.8	4.2	4.5	4.3	4.6	4.0	8.9	7.1	10.6	19.7
1999.....	2.7	4.2	5.5	4.3	7.3	6.3	8.1	6.0	10.4	22.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Dorsopathies (M40-M54)										
All races 3/, both sexes										
2006.....	*	0.1	0.2	0.3	0.5	0.6	1.3	2.0	3.4	6.9
2005.....	*	0.1	0.2	0.4	0.4	0.6	1.3	2.0	3.6	6.7
2004.....	*	0.1	0.2	0.3	0.5	0.6	1.2	2.0	3.4	7.1
2003.....	*	0.1	0.2	0.3	0.5	0.7	1.1	2.2	3.4	5.7
2002.....	*	0.1	0.2	0.3	0.4	0.6	1.2	1.7	2.9	6.0
2001.....	*	*	0.2	0.3	0.3	0.5	1.0	1.7	3.3	7.4
2000.....	*	0.1	0.1	0.2	0.4	0.7	0.9	1.7	3.3	6.1
1999.....	*	0.1	0.2	0.2	0.3	0.6	0.9	1.5	2.5	5.4
All races 3/, male										
2006.....	*	0.2	*	0.3	0.5	0.6	1.3	1.9	3.5	6.0
2005.....	*	*	0.3	0.5	0.4	0.7	1.5	2.5	3.5	6.2
2004.....	*	*	0.2	0.3	0.6	0.6	1.3	2.3	3.3	6.6
2003.....	*	*	0.3	0.4	0.4	0.9	1.1	2.4	3.6	4.8
2002.....	*	*	0.3	0.3	0.4	0.5	1.3	2.0	3.1	5.5
2001.....	*	*	*	0.3	0.4	0.6	0.9	1.9	3.5	6.4
2000.....	*	*	*	*	0.4	0.8	1.2	1.5	3.8	6.5
1999.....	*	*	*	*	0.5	0.7	1.0	1.8	2.8	5.9
All races 3/, female										
2006.....	*	*	0.2	0.3	0.4	0.6	1.3	2.0	3.4	7.3
2005.....	*	*	*	0.2	0.5	0.6	1.2	1.6	3.6	6.8
2004.....	*	*	*	0.3	0.4	0.5	1.1	1.7	3.4	7.3
2003.....	*	*	*	0.3	0.5	0.5	1.2	2.0	3.3	6.1
2002.....	*	*	*	0.3	0.4	0.7	1.1	1.5	2.8	6.2
2001.....	*	*	*	*	*	0.5	1.1	1.6	3.2	7.8
2000.....	*	*	*	*	*	0.5	0.7	1.9	3.1	6.0
1999.....	*	*	0.2	*	*	0.5	0.9	1.3	2.4	5.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Dorsopathies (M40-M54)										
White, both sexes										
2006 4/.....	*	0.2	0.2	0.3	0.5	0.6	1.3	2.0	3.5	7.2
2005 4/.....	*	0.1	0.2	0.4	0.4	0.6	1.2	2.0	3.7	7.0
2004 4/.....	*	*	0.2	0.3	0.5	0.5	1.2	2.1	3.5	7.4
2003 4/.....	*	0.1	0.2	0.3	0.4	0.7	1.0	2.1	3.5	5.9
2002.....	*	0.1	0.2	0.3	0.4	0.7	1.2	1.7	3.0	6.2
2001.....	*	*	0.2	0.2	0.3	0.5	1.0	1.7	3.3	7.8
2000.....	*	0.1	*	0.2	0.4	0.7	0.8	1.7	3.3	6.4
1999.....	*	*	0.2	0.2	0.3	0.5	0.9	1.5	2.7	5.5
White, male										
2006 4/.....	*	*	*	0.3	0.5	0.6	1.2	2.0	3.5	6.4
2005 4/.....	*	*	*	0.4	*	0.6	1.4	2.4	3.8	6.7
2004 4/.....	*	*	*	*	0.6	0.5	1.2	2.2	3.3	6.8
2003 4/.....	*	*	0.3	*	*	0.8	0.9	2.3	3.6	4.8
2002.....	*	*	0.3	*	*	*	1.3	1.8	3.2	5.3
2001.....	*	*	*	*	*	0.5	0.9	1.8	3.7	6.6
2000.....	*	*	*	*	0.5	0.7	1.1	1.4	3.8	6.9
1999.....	*	*	*	*	*	0.7	1.0	1.7	3.0	5.8
White, female										
2006 4/.....	*	*	*	0.3	0.4	0.6	1.4	2.0	3.5	7.6
2005 4/.....	*	*	*	0.3	0.5	0.6	1.1	1.8	3.7	7.1
2004 4/.....	*	*	*	0.3	0.4	0.5	1.2	1.9	3.7	7.7
2003 4/.....	*	*	*	0.3	0.4	0.6	1.0	2.0	3.4	6.3
2002.....	*	*	*	0.4	*	0.8	1.1	1.6	2.9	6.6
2001.....	*	*	*	*	*	0.5	1.1	1.6	3.1	8.3
2000.....	*	*	*	*	*	0.6	0.7	1.9	3.0	6.2
1999.....	*	*	*	*	*	*	0.9	1.3	2.5	5.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Soft tissue disorders (M60-M79)										
All races 3/, both sexes										
2006.....	0.2	0.4	0.5	0.6	0.9	1.0	1.3	2.0	2.9	4.8
2005.....	0.2	0.3	0.5	0.4	0.8	0.9	1.4	2.2	3.2	5.4
2004.....	0.2	0.4	0.5	0.6	0.7	0.9	1.5	2.1	3.4	5.1
2003.....	0.2	0.3	0.5	0.5	0.8	1.1	1.5	2.1	3.3	5.4
2002.....	0.2	0.3	0.4	0.5	0.6	0.9	1.3	1.9	3.1	5.3
2001.....	0.2	0.2	0.4	0.5	0.7	0.9	1.2	2.0	2.8	4.3
2000.....	0.2	0.4	0.4	0.5	0.6	0.8	1.2	2.0	2.3	4.2
1999.....	0.2	0.2	0.3	0.5	0.5	0.8	1.2	1.7	2.7	4.3
All races 3/, male										
2006.....	0.3	0.4	0.5	0.7	1.0	1.3	1.6	2.4	3.8	6.8
2005.....	0.3	0.3	0.6	0.4	0.9	1.0	1.7	2.8	3.8	7.9
2004.....	0.2	0.4	0.6	0.7	0.8	0.9	1.6	2.6	4.5	7.8
2003.....	0.2	0.3	0.5	0.4	0.7	1.1	1.9	2.6	4.5	7.1
2002.....	0.3	0.4	0.4	0.4	0.8	1.3	1.6	2.0	4.3	7.8
2001.....	0.3	0.3	0.6	0.6	0.6	1.1	1.6	2.7	3.9	6.2
2000.....	0.2	0.4	0.3	0.6	0.5	0.9	1.6	2.6	3.5	6.7
1999.....	0.3	0.3	0.3	0.6	0.5	0.8	1.7	2.5	4.0	6.8
All races 3/, female										
2006.....	0.2	0.3	0.4	0.5	0.7	0.7	1.0	1.8	2.4	3.9
2005.....	*	0.3	0.3	0.4	0.6	0.9	1.3	1.8	2.8	4.2
2004.....	*	0.4	0.3	0.5	0.6	0.9	1.4	1.7	2.7	3.9
2003.....	*	0.3	0.5	0.6	0.8	1.0	1.2	1.7	2.5	4.6
2002.....	*	*	0.3	0.6	0.5	0.6	1.1	1.8	2.4	4.2
2001.....	*	*	0.3	0.5	0.7	0.6	0.9	1.4	2.1	3.5
2000.....	*	0.3	0.4	0.4	0.6	0.8	0.9	1.6	1.7	3.2
1999.....	*	0.2	0.3	0.4	0.5	0.8	0.7	1.2	2.0	3.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Soft tissue disorders (M60-M79)										
White, both sexes										
2006 4/.....	0.2	0.3	0.4	0.5	0.8	0.9	1.2	2.0	2.8	4.8
2005 4/.....	0.2	0.3	0.4	0.3	0.7	0.9	1.3	2.1	3.2	5.6
2004 4/.....	0.2	0.4	0.4	0.5	0.6	0.8	1.5	2.0	3.3	4.8
2003 4/.....	0.2	0.2	0.5	0.5	0.6	1.0	1.5	1.9	3.2	5.4
2002.....	0.2	0.2	0.4	0.4	0.6	0.9	1.2	1.8	3.1	5.0
2001.....	0.2	0.2	0.4	0.5	0.5	0.8	1.1	1.8	2.7	4.2
2000.....	0.1	0.3	0.3	0.5	0.5	0.8	1.1	1.7	2.3	4.0
1999.....	0.2	0.2	0.3	0.4	0.4	0.7	1.1	1.7	2.6	4.0
White, male										
2006 4/.....	0.2	0.4	0.5	0.7	1.0	1.3	1.5	2.4	3.9	6.8
2005 4/.....	0.3	0.3	0.5	0.4	0.8	0.9	1.5	2.7	4.0	8.0
2004 4/.....	*	0.4	0.4	0.6	0.8	0.9	1.4	2.6	4.5	7.2
2003 4/.....	*	0.2	0.5	0.4	0.7	0.9	1.9	2.5	4.6	7.1
2002.....	0.2	0.3	0.4	0.4	0.7	1.2	1.4	1.9	4.2	7.6
2001.....	*	0.2	0.5	0.5	*	1.0	1.5	2.7	3.8	6.0
2000.....	*	0.3	0.3	0.6	0.5	0.9	1.3	2.4	3.5	6.3
1999.....	0.2	*	0.3	0.5	*	0.8	1.7	2.3	3.7	6.3
White, female										
2006 4/.....	*	0.3	0.3	0.4	0.6	0.6	1.0	1.7	2.1	3.9
2005 4/.....	*	0.3	0.3	0.3	0.7	0.8	1.2	1.7	2.8	4.4
2004 4/.....	*	0.4	0.3	0.4	0.5	0.8	1.5	1.7	2.6	3.7
2003 4/.....	*	0.3	0.4	0.6	0.6	1.1	1.1	1.5	2.4	4.6
2002.....	*	*	0.3	0.5	0.4	0.6	1.0	1.8	2.4	3.9
2001.....	*	*	0.3	0.4	0.6	0.5	0.8	1.1	2.1	3.4
2000.....	*	0.2	0.4	*	0.4	0.7	0.9	1.3	1.5	3.0
1999.....	*	*	0.3	*	*	0.7	0.7	1.2	2.0	3.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)

All races 3/, both sexes

2006.....	*	0.1	0.3	0.5	0.7	1.2	2.3	4.4	9.3	30.8
2005.....	0.1	0.2	0.3	0.4	0.8	1.1	2.2	4.2	10.5	34.6
2004.....	0.1	0.2	0.2	0.4	0.8	1.3	2.2	4.7	11.0	36.2
2003.....	0.1	0.2	0.3	0.5	0.7	1.3	2.5	4.3	10.5	39.5
2002.....	*	0.1	0.2	0.4	0.8	1.2	2.6	4.7	10.2	40.8
2001.....	*	0.2	0.2	0.3	0.7	1.3	2.2	4.4	10.8	40.1
2000.....	*	*	0.2	0.3	0.6	1.1	1.8	4.6	9.8	41.6
1999.....	*	0.1	0.2	0.4	0.5	1.0	1.8	4.4	10.3	38.6

All races 3/, male

2006.....	*	0.2	0.3	0.6	0.8	1.4	2.4	4.6	6.4	19.8
2005.....	*	0.3	0.4	0.5	0.8	1.1	2.2	3.8	9.4	19.9
2004.....	*	0.3	0.4	0.4	0.9	1.4	2.4	4.3	8.5	20.0
2003.....	*	0.2	0.3	0.7	0.8	1.8	2.5	3.7	7.5	21.5
2002.....	*	*	0.3	0.4	0.9	1.4	2.9	4.1	7.2	23.6
2001.....	*	*	0.2	0.5	0.7	1.3	2.1	4.0	7.7	23.6
2000.....	*	*	*	0.4	0.6	1.2	1.8	4.2	7.1	25.6
1999.....	*	*	*	0.3	0.5	0.8	1.8	3.7	7.8	20.4

All races 3/, female

2006.....	*	*	0.2	0.4	0.7	1.0	2.1	4.3	11.2	36.0
2005.....	*	*	0.2	0.4	0.7	1.1	2.3	4.5	11.2	41.4
2004.....	*	0.2	*	0.4	0.8	1.2	2.1	4.9	12.5	43.5
2003.....	*	0.2	0.2	0.3	0.7	0.9	2.6	4.8	12.4	47.5
2002.....	*	*	*	0.4	0.7	1.1	2.4	5.2	12.0	48.3
2001.....	*	*	*	*	0.7	1.3	2.2	4.8	12.7	47.0
2000.....	*	*	*	*	0.5	0.9	1.9	4.8	11.3	48.1
1999.....	*	*	*	0.4	0.6	1.2	1.7	4.8	11.8	45.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)

White, both sexes

2006 4/.....	*	0.1	0.2	0.5	0.8	1.2	2.2	4.3	9.4	32.2
2005 4/.....	*	0.2	0.3	0.4	0.7	1.1	2.3	4.2	10.5	36.0
2004 4/.....	0.1	0.2	0.2	0.4	0.8	1.3	2.1	4.5	11.1	37.7
2003 4/.....	0.1	0.2	0.3	0.4	0.7	1.2	2.5	4.4	10.6	41.0
2002.....	*	0.1	0.2	0.4	0.8	1.1	2.7	4.6	10.1	42.8
2001.....	*	0.1	0.2	0.3	0.7	1.2	2.0	4.5	11.0	41.3
2000.....	*	*	0.2	0.3	0.5	1.0	1.8	4.6	9.8	43.0
1999.....	*	0.1	0.2	0.3	0.5	0.9	1.7	4.2	10.7	40.2

White, male

2006 4/.....	*	*	0.3	0.5	0.8	1.5	2.3	4.3	6.3	20.0
2005 4/.....	*	0.2	0.4	0.4	0.7	1.0	2.2	3.5	9.4	20.7
2004 4/.....	*	*	0.3	0.4	0.8	1.4	2.2	3.9	8.4	20.4
2003 4/.....	*	*	0.3	0.6	0.6	1.6	2.2	3.5	7.2	20.9
2002.....	*	*	0.3	0.4	0.9	1.2	2.8	3.8	6.7	23.8
2001.....	*	*	*	0.4	0.6	1.2	1.8	3.8	7.6	23.8
2000.....	*	*	*	0.4	0.5	1.1	1.5	4.1	6.8	25.4
1999.....	*	*	*	*	0.5	0.6	1.8	3.3	8.2	20.2

White, female

2006 4/.....	*	*	*	0.4	0.8	0.9	2.1	4.3	11.3	38.0
2005 4/.....	*	*	*	0.4	0.6	1.1	2.4	4.7	11.3	43.1
2004 4/.....	*	*	*	0.4	0.8	1.2	2.1	5.0	12.8	45.5
2003 4/.....	*	*	*	*	0.8	0.9	2.7	5.0	12.7	49.9
2002.....	*	*	*	0.3	0.7	0.9	2.6	5.1	12.2	51.0
2001.....	*	*	*	*	0.7	1.3	2.0	4.9	13.1	48.6
2000.....	*	*	*	*	0.4	0.9	2.0	4.9	11.5	50.2
1999.....	*	*	*	0.4	0.5	1.2	1.7	4.9	12.2	48.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Osteopathies, chondropathies and other disorders of musculoskeletal system and connective tissue (M80-M99)

Black, both sexes

2006 4/.....	*	*	*	*	*	*	3.5	6.9	11.1	19.7
2005 4/.....	*	*	*	*	1.8	*	*	5.5	12.6	24.1
2004 4/.....	*	*	*	*	*	*	3.5	7.4	11.6	24.8
2003 4/.....	*	*	*	*	*	2.7	3.3	4.9	10.9	26.5
2002.....	*	*	*	*	*	2.7	2.9	6.4	12.3	24.1
2001.....	*	*	*	*	*	2.2	4.7	5.0	9.5	29.8
2000.....	*	*	*	*	*	2.2	2.8	5.3	11.0	27.5
1999.....	*	*	*	*	*	*	*	6.8	7.2	21.4

Black, male

2006 4/.....	*	*	*	*	*	*	*	8.9	*	22.3
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	10.5	*	21.9
2003 4/.....	*	*	*	*	*	4.9	*	*	*	27.1
2002.....	*	*	*	*	*	*	*	*	16.0	25.7
2001.....	*	*	*	*	*	*	*	*	*	26.1
2000.....	*	*	*	*	*	*	*	*	*	32.3
1999.....	*	*	*	*	*	*	*	*	*	24.6

Black, female

2006 4/.....	*	*	*	*	*	*	*	5.6	11.6	18.6
2005 4/.....	*	*	*	*	*	*	*	*	12.6	28.1
2004 4/.....	*	*	*	*	*	*	*	5.7	11.3	26.0
2003 4/.....	*	*	*	*	*	*	*	*	9.3	26.3
2002.....	*	*	*	*	*	*	*	6.3	10.3	23.4
2001.....	*	*	*	*	*	*	4.7	*	9.4	31.2
2000.....	*	*	*	*	*	*	*	*	10.2	25.8
1999.....	*	*	*	*	*	*	*	5.7	8.6	20.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Osteoporosis (M80-M81)										
All races 3/, both sexes										
2006.....	*	*	*	*	*	*	0.5	1.3	4.7	20.3
2005.....	*	*	*	*	*	0.2	0.5	1.6	5.1	23.3
2004.....	*	*	*	*	*	*	0.6	1.7	5.3	24.4
2003.....	*	*	*	*	*	*	0.7	1.7	5.4	26.9
2002.....	*	*	*	*	*	0.2	0.6	1.9	5.4	29.0
2001.....	*	*	*	*	*	0.3	0.5	1.7	5.6	26.8
2000.....	*	*	*	*	*	0.3	0.4	2.1	5.5	28.5
1999.....	*	*	*	*	*	0.3	0.5	2.0	5.8	27.1
All races 3/, male										
2006.....	*	*	*	*	*	*	*	*	1.3	7.2
2005.....	*	*	*	*	*	*	*	*	2.4	7.7
2004.....	*	*	*	*	*	*	*	0.7	1.9	7.2
2003.....	*	*	*	*	*	*	*	*	1.8	7.9
2002.....	*	*	*	*	*	*	*	0.7	1.7	9.9
2001.....	*	*	*	*	*	*	*	*	1.6	9.3
2000.....	*	*	*	*	*	*	*	0.9	1.9	9.9
1999.....	*	*	*	*	*	*	*	0.7	2.2	7.0
All races 3/, female										
2006.....	*	*	*	*	*	*	0.8	1.7	6.9	26.4
2005.....	*	*	*	*	*	*	0.7	2.4	6.9	30.5
2004.....	*	*	*	*	*	*	0.8	2.4	7.4	32.2
2003.....	*	*	*	*	*	*	1.1	2.5	7.6	35.2
2002.....	*	*	*	*	*	*	1.0	2.9	7.6	37.2
2001.....	*	*	*	*	*	0.4	0.6	2.4	8.0	34.1
2000.....	*	*	*	*	*	0.4	0.6	2.9	7.7	36.0
1999.....	*	*	*	*	*	0.4	0.7	2.9	7.9	35.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Osteoporosis (M80-M81)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	0.6	1.4	5.2	21.9
2005 4/.....	*	*	*	*	*	*	0.6	1.7	5.5	25.2
2004 4/.....	*	*	*	*	*	*	0.6	1.9	5.7	26.3
2003 4/.....	*	*	*	*	*	*	0.8	1.8	5.8	29.1
2002.....	*	*	*	*	*	*	0.7	2.0	5.8	31.3
2001.....	*	*	*	*	*	0.3	0.5	1.8	6.0	28.8
2000.....	*	*	*	*	*	0.3	0.4	2.3	5.9	30.7
1999.....	*	*	*	*	*	0.3	0.5	2.2	6.2	29.1
White, male										
2006 4/.....	*	*	*	*	*	*	*	*	1.4	7.8
2005 4/.....	*	*	*	*	*	*	*	*	2.6	8.3
2004 4/.....	*	*	*	*	*	*	*	0.8	1.9	7.8
2003 4/.....	*	*	*	*	*	*	*	*	1.9	8.2
2002.....	*	*	*	*	*	*	*	*	1.9	10.6
2001.....	*	*	*	*	*	*	*	*	1.7	10.0
2000.....	*	*	*	*	*	*	*	0.9	2.1	10.4
1999.....	*	*	*	*	*	*	*	0.8	2.3	7.4
White, female										
2006 4/.....	*	*	*	*	*	*	0.9	1.9	7.5	28.5
2005 4/.....	*	*	*	*	*	*	0.8	2.6	7.3	33.0
2004 4/.....	*	*	*	*	*	*	0.9	2.7	8.0	34.7
2003 4/.....	*	*	*	*	*	*	1.2	2.7	8.2	38.4
2002.....	*	*	*	*	*	*	1.0	3.0	8.2	40.3
2001.....	*	*	*	*	*	0.5	0.6	2.6	8.6	36.6
2000.....	*	*	*	*	*	0.5	0.6	3.2	8.2	38.9
1999.....	*	*	*	*	*	0.5	0.8	3.1	8.5	37.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)

All races 3/, both sexes

2006.....	*	0.1	0.2	0.5	0.7	1.0	1.7	3.2	4.6	10.6
2005.....	0.1	0.2	0.3	0.4	0.7	0.9	1.7	2.6	5.4	11.3
2004.....	0.1	0.2	0.2	0.4	0.7	1.2	1.7	3.0	5.7	11.8
2003.....	0.1	0.2	0.2	0.5	0.7	1.2	1.8	2.7	5.1	12.6
2002.....	*	0.1	0.2	0.4	0.7	1.0	2.0	2.8	4.8	11.8
2001.....	*	0.2	0.2	0.3	0.6	1.0	1.7	2.8	5.3	13.3
2000.....	*	*	0.2	0.3	0.5	0.8	1.4	2.5	4.3	13.1
1999.....	*	0.1	0.2	0.3	0.4	0.7	1.3	2.4	4.6	11.5

All races 3/, male

2006.....	*	*	0.3	0.6	0.7	1.3	2.2	3.9	5.1	12.6
2005.....	*	0.2	0.4	0.4	0.8	1.1	1.9	3.2	7.1	12.2
2004.....	*	0.2	0.3	0.3	0.8	1.3	2.1	3.6	6.6	12.9
2003.....	*	0.2	0.3	0.6	0.7	1.7	2.1	3.3	5.6	13.6
2002.....	*	*	0.3	0.4	0.9	1.3	2.6	3.4	5.5	13.7
2001.....	*	*	0.2	0.4	0.7	1.2	1.8	3.3	6.1	14.2
2000.....	*	*	*	0.4	0.6	1.2	1.6	3.4	5.2	15.6
1999.....	*	*	*	0.3	0.5	0.7	1.7	3.0	5.7	13.3

All races 3/, female

2006.....	*	*	0.2	0.4	0.7	0.7	1.3	2.6	4.3	9.6
2005.....	*	*	*	0.3	0.6	0.8	1.6	2.1	4.4	10.9
2004.....	*	0.2	*	0.4	0.7	1.0	1.4	2.5	5.1	11.3
2003.....	*	0.2	0.2	0.3	0.6	0.7	1.5	2.3	4.8	12.2
2002.....	*	*	*	0.4	0.6	0.7	1.4	2.3	4.4	11.1
2001.....	*	*	*	*	0.5	0.8	1.7	2.4	4.8	12.9
2000.....	*	*	*	*	0.4	0.5	1.3	1.9	3.7	12.0
1999.....	*	*	*	0.3	0.4	0.7	1.0	1.9	3.9	10.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other osteopathies, chondropathies and disorders of musculoskeletal system and connective tissue (M83-M99)

White, both sexes

2006 4/.....	*	0.1	0.2	0.4	0.7	1.0	1.6	2.9	4.2	10.3
2005 4/.....	*	0.2	0.3	0.3	0.6	0.9	1.7	2.4	5.1	10.8
2004 4/.....	*	0.2	0.2	0.3	0.7	1.1	1.5	2.7	5.4	11.4
2003 4/.....	*	0.2	0.2	0.4	0.6	1.0	1.7	2.6	4.8	11.9
2002.....	*	0.1	0.2	0.3	0.7	0.8	2.0	2.5	4.4	11.5
2001.....	*	0.1	0.2	0.3	0.5	0.9	1.5	2.6	5.0	12.5
2000.....	*	*	0.1	0.3	0.4	0.7	1.3	2.3	3.8	12.4
1999.....	*	0.1	0.2	0.3	0.5	0.6	1.2	2.0	4.5	11.1

White, male

2006 4/.....	*	*	0.3	0.5	0.7	1.3	2.1	3.7	4.9	12.2
2005 4/.....	*	0.2	0.4	0.4	0.7	1.0	1.9	2.9	6.8	12.4
2004 4/.....	*	*	0.3	0.3	0.8	1.3	1.8	3.1	6.5	12.6
2003 4/.....	*	*	0.3	0.6	0.6	1.5	1.8	3.0	5.3	12.7
2002.....	*	*	0.3	0.4	0.8	1.1	2.5	3.2	4.8	13.2
2001.....	*	*	*	0.3	0.5	1.0	1.5	3.1	5.9	13.8
2000.....	*	*	*	0.4	0.5	1.0	1.3	3.3	4.7	15.1
1999.....	*	*	*	*	0.5	*	1.7	2.5	5.9	12.8

White, female

2006 4/.....	*	*	*	0.4	0.7	0.7	1.2	2.4	3.8	9.4
2005 4/.....	*	*	*	0.3	0.5	0.8	1.5	2.0	4.0	10.1
2004 4/.....	*	*	*	0.3	0.7	1.0	1.3	2.3	4.8	10.8
2003 4/.....	*	*	*	*	0.7	0.7	1.5	2.3	4.5	11.5
2002.....	*	*	*	*	0.5	0.6	1.5	2.1	4.1	10.7
2001.....	*	*	*	*	0.5	0.8	1.4	2.3	4.5	12.0
2000.....	*	*	*	*	*	*	1.3	1.7	3.3	11.2
1999.....	*	*	*	*	*	0.6	0.8	1.7	3.7	10.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XIV. Diseases of the genitourinary system (N00-N98)										
All races 3/, both sexes										
2006.....	2.9	4.7	8.2	13.1	22.3	37.5	64.9	115.5	207.3	454.9
2005.....	2.6	4.8	7.3	12.9	21.8	36.7	66.5	117.9	207.8	464.4
2004.....	2.7	4.6	7.9	13.0	21.2	37.3	63.2	115.4	204.1	458.5
2003.....	2.8	4.6	7.6	12.6	22.2	37.9	65.1	114.1	208.5	469.5
2002.....	2.6	4.5	7.1	12.2	20.5	36.5	64.2	115.2	206.9	469.4
2001.....	2.6	4.5	7.1	12.1	20.6	39.2	64.4	113.7	202.0	476.4
2000.....	2.6	4.3	6.8	11.5	20.7	36.4	60.5	111.8	202.9	471.4
1999.....	2.5	3.9	6.4	11.2	19.8	35.5	61.7	109.5	197.7	471.1
All races 3/, male										
2006.....	3.2	5.4	9.5	14.7	24.7	40.8	73.7	133.4	242.4	531.8
2005.....	3.0	5.5	8.6	14.7	25.3	38.7	74.2	133.4	247.5	538.8
2004.....	3.0	5.3	9.0	14.5	23.2	40.8	69.7	132.9	246.3	533.9
2003.....	3.1	5.1	8.9	14.2	23.8	41.3	71.9	132.9	253.9	547.9
2002.....	2.7	5.1	8.1	12.9	22.3	40.4	73.7	134.0	246.8	553.0
2001.....	2.7	5.0	7.8	12.9	21.7	44.0	71.6	129.8	242.7	564.2
2000.....	2.9	5.1	7.6	12.6	22.1	39.6	67.3	131.0	243.1	567.5
1999.....	2.7	4.5	6.9	12.1	22.0	38.3	70.3	130.4	241.8	558.3
All races 3/, female										
2006.....	2.5	3.9	7.0	11.7	20.0	34.7	57.8	102.4	185.4	418.9
2005.....	2.3	4.0	6.1	11.3	18.5	35.0	60.2	106.7	183.1	430.3
2004.....	2.4	4.0	6.8	11.6	19.4	34.2	58.0	102.9	178.1	424.5
2003.....	2.6	4.2	6.3	11.1	20.6	35.0	59.6	100.7	180.8	434.9
2002.....	2.4	4.0	6.2	11.6	18.9	33.1	56.7	102.0	182.6	433.2
2001.....	2.4	3.9	6.3	11.4	19.6	35.0	58.6	102.4	177.6	439.6
2000.....	2.2	3.4	6.0	10.4	19.4	33.7	55.2	98.5	179.1	432.3
1999.....	2.3	3.3	5.9	10.4	17.9	33.1	54.9	95.0	171.9	436.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XIV. Diseases of the genitourinary system (N00-N98)										
White, both sexes										
2006 4/.....	2.0	3.7	6.0	10.4	18.1	31.3	56.8	107.7	200.1	446.7
2005 4/.....	1.8	3.5	5.6	9.9	17.5	31.0	58.7	109.8	199.9	456.5
2004 4/.....	2.1	3.3	5.8	9.8	16.9	31.5	56.3	107.0	196.2	449.9
2003 4/.....	2.0	3.3	5.4	9.7	17.7	31.9	57.1	106.2	199.8	461.3
2002.....	1.9	3.3	5.0	9.4	16.4	30.9	56.8	106.8	198.9	458.5
2001.....	1.9	3.1	5.1	9.2	16.2	32.5	57.2	105.3	192.3	465.1
2000.....	1.9	3.0	4.9	8.7	16.5	30.2	52.8	103.9	193.4	460.5
1999.....	1.7	2.7	4.6	8.8	15.8	29.3	54.0	101.0	187.8	456.0
White, male										
2006 4/.....	2.3	4.3	7.0	11.4	20.4	34.5	65.4	125.6	235.6	529.7
2005 4/.....	2.2	4.1	6.7	10.9	20.5	33.7	66.9	125.0	241.2	537.4
2004 4/.....	2.4	3.7	6.4	11.2	18.6	34.3	63.1	124.6	240.8	527.3
2003 4/.....	2.3	3.7	6.1	10.9	19.2	35.1	63.6	125.4	244.3	547.4
2002.....	2.0	3.6	5.6	9.8	17.5	34.7	65.9	125.5	240.2	543.2
2001.....	2.0	3.4	5.8	9.8	17.5	37.1	64.8	122.6	233.9	556.8
2000.....	2.1	3.5	5.4	9.5	17.9	33.7	60.4	123.0	234.8	561.2
1999.....	1.8	3.0	4.9	9.9	17.5	32.0	62.2	120.6	230.8	542.8
White, female										
2006 4/.....	1.7	3.0	5.1	9.4	16.0	28.3	49.7	94.3	177.6	407.8
2005 4/.....	1.5	2.9	4.6	8.9	14.6	28.6	51.9	98.5	174.0	419.3
2004 4/.....	1.7	2.9	5.1	8.4	15.4	29.0	50.7	94.1	168.5	415.0
2003 4/.....	1.6	3.0	4.8	8.7	16.4	29.2	51.8	92.3	172.3	423.2
2002.....	1.8	2.9	4.4	9.0	15.4	27.5	49.3	93.5	173.5	421.7
2001.....	1.7	2.8	4.5	8.6	14.9	28.5	51.0	93.0	167.1	426.7
2000.....	1.6	2.6	4.4	7.9	15.3	27.0	46.7	90.5	168.8	419.6
1999.....	1.6	2.3	4.2	7.7	14.1	27.0	47.5	87.2	162.4	421.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XIV. Diseases of the genitourinary system (N00-N98)										
Black, both sexes										
2006 4/.....	8.8	12.3	25.9	37.2	62.3	98.6	147.6	220.7	329.9	625.5
2005 4/.....	8.2	14.1	21.4	40.1	63.1	91.9	149.7	226.2	339.3	623.5
2004 4/.....	7.7	14.6	25.0	41.6	63.0	92.4	140.9	222.4	331.0	634.5
2003 4/.....	9.0	14.0	24.7	39.6	64.5	96.3	153.2	219.0	346.7	645.3
2002.....	7.6	13.6	24.1	37.9	58.7	91.9	150.1	225.4	332.3	662.5
2001.....	7.6	14.5	23.0	36.9	60.3	102.5	146.3	223.5	347.5	664.5
2000.....	7.5	14.0	22.6	36.6	58.1	96.6	145.9	219.5	343.1	637.5
1999.....	7.9	13.5	22.7	32.7	56.2	94.5	147.3	219.9	337.2	679.1
Black, male										
2006 4/.....	10.2	14.3	31.2	45.2	73.5	111.8	165.2	255.1	390.9	666.6
2005 4/.....	9.2	17.3	25.9	51.1	77.6	94.4	162.6	262.4	392.6	658.1
2004 4/.....	8.1	17.0	31.2	47.1	70.6	107.1	156.0	260.2	366.6	718.3
2003 4/.....	9.2	15.5	33.3	47.7	73.7	108.3	181.0	249.7	437.1	681.9
2002.....	8.3	16.4	29.0	44.4	70.1	101.7	177.3	262.7	382.1	794.4
2001.....	7.8	18.3	25.9	42.9	63.3	118.4	158.5	241.6	404.5	755.7
2000.....	8.9	18.8	27.3	42.8	63.2	102.3	155.8	259.4	398.7	716.2
1999.....	9.2	17.3	26.4	32.9	65.8	102.0	172.3	270.7	433.4	802.7
Black, female										
2006 4/.....	7.6	10.6	21.3	30.5	53.6	88.8	135.5	200.1	297.7	608.7
2005 4/.....	7.4	11.3	17.5	31.0	51.7	90.1	140.9	204.5	311.2	609.6
2004 4/.....	7.5	12.4	19.7	37.1	57.1	81.5	130.7	200.0	312.2	601.3
2003 4/.....	8.7	12.7	17.3	32.9	57.3	87.4	134.5	200.8	299.4	631.1
2002.....	7.0	11.1	20.0	32.6	49.9	84.5	132.0	203.1	306.4	612.0
2001.....	7.4	11.3	20.5	32.1	57.9	90.7	138.1	212.7	318.5	630.6
2000.....	6.3	9.9	18.6	31.6	54.1	92.4	139.4	195.4	315.0	608.3
1999.....	6.7	10.2	19.5	32.5	48.8	88.9	130.7	189.4	289.5	633.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of urinary system (N00-N39)										
All races 3/, both sexes										
2006.....	2.8	4.5	8.1	12.9	21.9	37.0	64.1	113.9	203.2	445.7
2005.....	2.6	4.6	7.2	12.7	21.5	36.0	65.5	116.1	204.2	455.1
2004.....	2.7	4.5	7.7	12.7	20.9	36.7	62.2	113.8	200.2	448.9
2003.....	2.7	4.5	7.5	12.4	21.9	37.3	64.3	112.4	205.2	460.5
2002.....	2.5	4.4	7.0	12.1	20.3	36.1	63.5	113.9	203.4	460.8
2001.....	2.5	4.3	6.9	11.9	20.4	38.7	63.3	112.0	198.9	467.4
2000.....	2.5	4.2	6.6	11.3	20.4	35.9	59.7	110.2	199.3	462.5
1999.....	2.4	3.8	6.3	11.0	19.5	35.0	60.8	107.8	194.1	462.6
All races 3/, male										
2006.....	3.2	5.4	9.4	14.4	24.3	39.9	72.5	131.0	235.2	511.6
2005.....	2.9	5.5	8.5	14.4	25.0	37.8	72.9	130.7	240.8	517.6
2004.....	3.0	5.1	8.8	14.2	22.7	40.0	68.3	130.1	238.8	513.6
2003.....	3.1	5.0	8.8	13.9	23.5	40.5	70.8	130.3	247.9	528.3
2002.....	2.7	5.0	8.0	12.8	21.9	39.8	72.4	132.0	239.9	533.8
2001.....	2.7	5.0	7.7	12.7	21.5	43.5	70.2	127.2	237.2	542.9
2000.....	2.9	5.1	7.5	12.4	21.7	39.0	66.0	128.5	236.8	546.9
1999.....	2.6	4.5	6.8	11.9	21.6	37.6	69.0	127.7	234.6	537.0
All races 3/, female										
2006.....	2.4	3.7	6.8	11.4	19.8	34.4	57.2	101.4	183.3	414.9
2005.....	2.2	3.8	5.9	11.1	18.2	34.5	59.6	105.6	181.5	426.3
2004.....	2.3	3.9	6.7	11.4	19.2	33.8	57.3	102.1	176.4	419.9
2003.....	2.4	4.0	6.1	11.0	20.3	34.5	59.1	99.6	179.2	430.5
2002.....	2.3	3.8	6.1	11.4	18.8	32.9	56.3	101.2	181.1	429.1
2001.....	2.3	3.7	6.1	11.2	19.4	34.6	57.8	101.3	175.9	435.8
2000.....	2.1	3.3	5.8	10.3	19.3	33.3	54.7	97.5	177.2	428.1
1999.....	2.1	3.1	5.8	10.2	17.6	32.8	54.4	93.9	170.4	432.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of urinary system (N00-N39)										
White, both sexes										
2006 4/.....	1.9	3.6	5.9	10.2	17.8	30.7	56.0	106.1	195.9	437.5
2005 4/.....	1.8	3.4	5.5	9.7	17.2	30.3	57.8	108.1	196.2	447.1
2004 4/.....	2.0	3.2	5.6	9.6	16.6	31.0	55.4	105.4	192.4	440.3
2003 4/.....	1.9	3.2	5.4	9.6	17.5	31.4	56.3	104.4	196.4	451.9
2002.....	1.8	3.2	4.9	9.2	16.2	30.5	56.1	105.6	195.3	449.9
2001.....	1.8	3.0	5.0	9.0	16.0	32.1	56.1	103.6	189.1	456.0
2000.....	1.8	2.9	4.8	8.6	16.3	29.7	51.9	102.4	189.9	451.6
1999.....	1.6	2.6	4.5	8.6	15.5	28.9	53.3	99.4	184.2	447.4
White, male										
2006 4/.....	2.2	4.3	6.9	11.2	20.0	33.8	64.2	123.3	228.2	509.3
2005 4/.....	2.1	4.1	6.6	10.7	20.3	32.7	65.7	122.4	234.3	515.7
2004 4/.....	2.4	3.6	6.3	11.0	18.1	33.6	61.7	122.1	233.4	506.5
2003 4/.....	2.3	3.7	6.1	10.7	18.9	34.4	62.4	122.9	238.2	527.0
2002.....	2.0	3.6	5.6	9.7	17.2	34.2	64.7	123.7	233.4	523.8
2001.....	2.0	3.3	5.7	9.7	17.4	36.5	63.4	119.9	228.5	535.0
2000.....	2.1	3.5	5.4	9.4	17.5	33.2	59.3	120.6	228.4	540.8
1999.....	1.8	3.0	4.8	9.8	17.2	31.4	61.1	117.9	223.8	521.3
White, female										
2006 4/.....	1.5	2.8	4.9	9.2	15.8	28.1	49.1	93.3	175.5	403.7
2005 4/.....	1.4	2.8	4.5	8.7	14.3	28.1	51.3	97.6	172.3	415.4
2004 4/.....	1.6	2.8	5.0	8.2	15.1	28.7	50.2	93.2	166.9	410.4
2003 4/.....	1.5	2.8	4.7	8.5	16.2	28.7	51.4	91.1	170.6	418.6
2002.....	1.7	2.8	4.3	8.8	15.2	27.3	49.0	92.6	172.0	417.8
2001.....	1.6	2.7	4.4	8.4	14.8	28.2	50.2	92.0	165.3	423.0
2000.....	1.5	2.4	4.2	7.8	15.2	26.7	46.1	89.5	167.0	415.4
1999.....	1.5	2.2	4.1	7.5	13.9	26.7	47.1	86.3	160.9	417.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Diseases of urinary system (N00-N39)										
Black, both sexes										
2006 4/.....	8.6	12.0	25.6	36.6	61.8	97.5	146.2	218.5	325.9	616.7
2005 4/.....	8.1	13.6	21.0	39.7	62.3	91.2	148.4	222.3	336.9	615.0
2004 4/.....	7.6	14.1	24.5	41.2	62.6	91.3	138.6	219.7	327.0	625.5
2003 4/.....	8.7	13.8	24.2	39.2	63.5	95.1	152.3	216.8	344.7	638.9
2002.....	7.5	13.3	23.8	37.8	58.0	90.9	148.1	222.8	328.9	652.1
2001.....	7.3	14.2	22.5	36.7	59.1	101.9	144.9	220.6	343.2	655.3
2000.....	7.2	13.9	22.5	36.1	57.6	95.6	144.5	217.0	339.1	628.5
1999.....	7.7	13.1	22.3	32.2	55.4	93.3	145.0	216.7	332.3	669.3
Black, male										
2006 4/.....	10.0	14.3	31.1	44.6	72.5	109.9	163.0	251.5	384.7	648.8
2005 4/.....	9.2	17.1	25.6	50.7	76.3	93.6	160.0	257.5	389.1	640.5
2004 4/.....	8.0	16.9	30.7	46.4	69.9	105.6	153.4	253.9	359.4	701.4
2003 4/.....	9.2	15.4	32.9	47.1	72.7	106.9	180.3	244.7	432.6	666.2
2002.....	8.3	16.4	28.8	44.4	68.9	100.5	173.6	256.7	373.7	776.2
2001.....	7.6	18.1	25.4	42.5	61.8	117.9	156.5	237.9	395.7	736.4
2000.....	8.8	18.8	27.0	42.0	62.1	101.2	153.5	254.2	392.0	695.4
1999.....	9.2	17.2	25.7	32.2	64.5	100.5	168.9	268.3	421.2	779.2
Black, female										
2006 4/.....	7.4	9.9	20.9	30.0	53.4	88.3	134.6	198.7	294.8	603.5
2005 4/.....	7.1	10.6	17.1	30.6	51.3	89.4	140.5	201.3	309.4	604.7
2004 4/.....	7.4	11.8	19.4	36.9	57.0	80.6	128.5	199.5	310.0	595.4
2003 4/.....	8.3	12.4	16.9	32.8	56.4	86.4	133.4	200.2	298.6	628.3
2002.....	6.7	10.7	19.6	32.4	49.4	83.8	131.1	202.5	305.6	604.6
2001.....	7.0	10.8	19.9	31.8	57.0	90.1	137.2	210.2	316.4	625.2
2000.....	5.8	9.6	18.6	31.3	54.1	91.4	138.5	194.5	312.5	603.5
1999.....	6.4	9.7	19.3	32.3	48.3	87.9	129.2	185.7	288.2	628.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Glomerular and renal tubulo-interstitial diseases (N00-N15)

All races 3/, both sexes

2006.....	0.2	0.3	0.5	0.7	1.0	1.7	3.0	5.1	9.5	21.7
2005.....	0.2	0.2	0.4	0.5	0.8	1.2	2.0	4.0	6.7	14.8
2004.....	0.2	0.2	0.3	0.5	0.6	1.1	1.9	3.5	6.4	12.6
2003.....	0.2	0.3	0.4	0.5	0.6	1.3	1.9	3.2	6.9	13.3
2002.....	0.1	0.2	0.3	0.5	0.6	1.1	2.0	3.9	6.2	13.9
2001.....	0.1	0.2	0.3	0.4	0.6	1.2	2.0	3.8	5.8	13.7
2000.....	0.2	0.2	0.3	0.5	0.7	1.2	1.9	3.7	6.5	15.7
1999.....	0.2	0.2	0.5	0.6	0.8	1.3	2.2	4.2	7.2	16.4

All races 3/, male

2006.....	0.2	0.2	0.5	0.7	1.1	1.8	3.3	6.2	11.7	27.1
2005.....	0.2	*	0.4	0.5	0.9	1.2	2.1	4.1	8.0	18.4
2004.....	*	0.2	0.3	0.6	0.7	1.1	2.0	3.8	7.1	16.9
2003.....	*	0.3	0.4	0.4	0.6	1.1	1.8	3.5	7.7	17.8
2002.....	*	*	0.3	0.5	0.6	1.2	2.3	4.7	6.7	18.1
2001.....	*	*	0.3	0.4	0.5	1.3	2.4	3.9	7.6	17.2
2000.....	*	0.2	0.4	0.5	0.7	1.2	1.6	4.3	7.2	20.0
1999.....	0.2	0.2	0.4	0.6	0.8	1.2	2.0	4.8	9.8	22.6

All races 3/, female

2006.....	0.2	0.3	0.4	0.7	0.8	1.7	2.8	4.3	8.1	19.2
2005.....	*	0.3	0.3	0.4	0.7	1.2	2.0	3.9	5.9	13.1
2004.....	0.2	0.3	0.4	0.4	0.5	1.0	1.8	3.3	5.9	10.7
2003.....	0.2	0.2	0.3	0.5	0.7	1.4	1.9	3.0	6.4	11.3
2002.....	*	0.3	0.3	0.4	0.7	1.1	1.7	3.3	6.0	12.0
2001.....	*	0.2	0.4	0.4	0.7	1.2	1.6	3.7	4.7	12.3
2000.....	*	0.3	0.3	0.5	0.7	1.2	2.2	3.3	6.1	13.9
1999.....	*	0.3	0.5	0.6	0.7	1.3	2.5	3.8	5.6	13.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Glomerular and renal tubulo-interstitial diseases (N00-N15)

White, both sexes

2006 4/.....	0.1	0.2	0.4	0.6	0.8	1.5	2.7	4.8	9.1	21.1
2005 4/.....	0.1	0.2	0.3	0.4	0.7	1.1	1.9	3.8	6.7	14.5
2004 4/.....	0.2	0.2	0.3	0.4	0.5	1.0	1.9	3.4	6.4	12.4
2003 4/.....	0.2	0.2	0.3	0.4	0.5	1.3	1.8	3.1	6.6	13.2
2002.....	0.2	0.2	0.3	0.4	0.6	1.2	2.1	3.8	6.2	13.9
2001.....	0.1	0.1	0.3	0.4	0.5	1.1	2.0	3.8	5.5	13.9
2000.....	0.1	0.2	0.2	0.5	0.6	1.1	1.8	3.5	6.3	15.2
1999.....	0.1	0.3	0.4	0.5	0.6	1.1	2.1	4.2	7.1	16.3

White, male

2006 4/.....	*	*	0.3	0.5	0.9	1.4	3.2	5.9	11.1	26.1
2005 4/.....	*	*	0.3	0.5	0.7	1.1	2.0	3.8	8.1	18.1
2004 4/.....	*	*	0.3	0.5	0.6	1.1	1.9	3.6	7.3	16.5
2003 4/.....	*	0.3	0.4	0.4	0.5	1.1	1.7	3.3	7.2	18.0
2002.....	*	*	0.3	0.5	0.6	1.1	2.4	4.3	6.7	18.4
2001.....	*	*	*	0.4	*	1.0	2.4	3.7	7.4	17.5
2000.....	*	0.2	*	0.5	0.6	1.0	1.6	4.1	7.1	19.5
1999.....	*	*	0.4	0.5	0.7	1.1	1.6	4.8	9.6	22.1

White, female

2006 4/.....	*	0.3	0.4	0.7	0.7	1.5	2.3	4.0	7.7	18.8
2005 4/.....	*	0.3	0.2	0.4	0.7	1.1	1.8	3.8	5.8	12.9
2004 4/.....	0.2	*	0.3	0.4	0.5	1.0	1.8	3.3	5.8	10.6
2003 4/.....	*	*	0.3	0.5	0.6	1.4	1.8	3.0	6.2	11.0
2002.....	*	*	0.3	0.3	0.6	1.2	1.8	3.4	5.9	11.9
2001.....	*	*	0.3	0.4	0.6	1.1	1.6	3.8	4.4	12.4
2000.....	*	*	*	0.5	0.6	1.1	2.0	3.1	5.8	13.5
1999.....	*	0.3	0.4	0.6	0.6	1.1	2.4	3.7	5.6	14.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Glomerular diseases (N00-N07)										
All races 3/, both sexes										
2006.....	0.1	0.1	0.3	0.4	0.6	1.1	1.8	3.4	6.2	13.9
2005.....	*	*	0.2	0.3	0.4	0.6	1.0	1.8	2.6	5.9
2004.....	*	*	0.2	0.2	0.3	0.5	0.7	1.3	2.0	3.3
2003.....	*	*	0.2	0.2	0.3	0.5	0.8	1.1	2.1	3.0
2002.....	*	0.1	0.1	0.2	0.2	0.4	0.7	1.3	2.2	3.3
2001.....	*	*	0.2	0.2	0.2	0.6	0.9	1.2	1.8	3.3
2000.....	0.1	0.1	0.2	0.2	0.3	0.6	0.8	1.2	2.4	3.6
1999.....	*	0.1	0.2	0.2	0.4	0.5	0.9	1.7	2.3	3.7
All races 3/, male										
2006.....	*	0.2	0.3	0.5	0.8	1.3	1.9	4.6	7.7	17.9
2005.....	*	*	0.3	0.3	0.5	0.6	1.1	2.2	3.3	8.2
2004.....	*	*	*	*	0.4	0.5	0.8	1.5	2.5	4.2
2003.....	*	*	0.2	*	*	0.7	0.8	1.3	2.5	3.9
2002.....	*	*	*	0.3	*	0.5	0.9	1.6	2.4	4.3
2001.....	*	*	*	*	*	0.7	1.1	1.5	2.6	4.6
2000.....	*	*	0.3	*	0.4	0.7	0.8	1.7	2.6	5.6
1999.....	*	*	*	*	0.5	0.5	0.8	2.2	3.4	5.6
All races 3/, female										
2006.....	*	*	0.3	0.4	0.4	0.9	1.8	2.5	5.2	12.1
2005.....	*	*	*	*	0.4	0.5	0.8	1.6	2.1	4.8
2004.....	*	*	*	*	*	0.5	0.6	1.1	1.6	2.9
2003.....	*	*	*	*	*	0.4	0.9	1.0	1.9	2.6
2002.....	*	*	*	*	*	*	0.6	1.1	2.0	2.8
2001.....	*	*	*	*	*	0.6	0.7	1.0	1.3	2.8
2000.....	*	*	*	*	*	0.5	0.8	0.8	2.3	2.8
1999.....	*	*	*	*	0.4	0.5	1.0	1.4	1.7	2.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Glomerular diseases (N00-N07)										
White, both sexes										
2006 4/.....	*	*	0.2	0.3	0.5	0.8	1.6	3.1	5.7	13.0
2005 4/.....	*	*	*	0.2	0.4	0.5	0.8	1.8	2.4	5.6
2004 4/.....	*	*	0.2	0.2	0.3	0.4	0.7	1.2	1.9	3.2
2003 4/.....	*	*	0.1	0.2	0.3	0.5	0.7	1.0	2.0	2.8
2002.....	*	*	*	0.2	0.2	0.4	0.7	1.2	2.0	3.0
2001.....	*	*	*	*	*	0.5	0.9	1.2	1.7	3.3
2000.....	*	*	*	0.2	0.3	0.5	0.8	1.1	2.4	3.3
1999.....	*	*	0.1	0.2	0.3	0.4	0.9	1.7	2.1	3.4
White, male										
2006 4/.....	*	*	*	0.3	0.7	1.0	1.8	4.2	7.0	16.6
2005 4/.....	*	*	*	*	0.4	0.5	1.1	2.1	3.2	8.0
2004 4/.....	*	*	*	*	*	*	0.8	1.4	2.5	3.9
2003 4/.....	*	*	*	*	*	0.6	0.7	1.0	2.3	3.7
2002.....	*	*	*	*	*	*	0.9	1.5	2.4	4.1
2001.....	*	*	*	*	*	0.5	1.2	1.6	2.7	4.8
2000.....	*	*	*	*	*	0.6	0.9	1.7	2.6	4.9
1999.....	*	*	*	*	*	*	0.8	2.2	3.0	5.1
White, female										
2006 4/.....	*	*	*	0.3	*	0.7	1.4	2.4	4.8	11.4
2005 4/.....	*	*	*	*	0.4	*	0.6	1.6	1.9	4.5
2004 4/.....	*	*	*	*	*	*	0.6	1.1	1.6	2.8
2003 4/.....	*	*	*	*	*	*	0.8	1.0	1.8	2.4
2002.....	*	*	*	*	*	*	0.6	1.1	1.9	2.5
2001.....	*	*	*	*	*	0.5	0.7	1.0	1.2	2.7
2000.....	*	*	*	*	*	*	0.8	0.7	2.2	2.6
1999.....	*	*	*	*	*	*	1.0	1.3	1.6	2.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Nephrotic syndrome (N04)										
All races 3/, both sexes										
2006.....	*	*	*	*	*	*	*	*	0.4	0.5
2005.....	*	*	*	*	*	*	*	0.3	*	0.6
2004.....	*	*	*	*	*	*	*	*	0.4	0.6
2003.....	*	*	*	*	*	*	*	*	0.5	0.5
2002.....	*	*	*	*	*	*	*	*	0.6	0.7
2001.....	*	*	*	*	*	*	*	0.3	*	0.7
2000.....	*	*	*	*	*	*	*	*	0.6	1.0
1999.....	*	*	*	*	*	*	0.3	0.4	0.5	0.7
All races 3/, male										
2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	*	*
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2006.....	*	*	*	*	*	*	*	*	*	*
2005.....	*	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	0.6
2003.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	0.7
2001.....	*	*	*	*	*	*	*	*	*	*
2000.....	*	*	*	*	*	*	*	*	0.6	0.8
1999.....	*	*	*	*	*	*	*	0.5	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified glomerular diseases (N02-N03,N05-N07)

All races 3/, both sexes

2006.....	*	0.1	0.2	0.4	0.6	1.0	1.7	3.2	5.7	13.3
2005.....	*	*	0.2	0.2	0.4	0.5	0.8	1.5	2.2	5.1
2004.....	*	*	0.1	0.2	0.3	0.4	0.5	0.9	1.5	2.5
2003.....	*	*	0.1	*	0.2	0.4	0.6	0.9	1.5	2.5
2002.....	*	0.1	0.1	0.2	*	0.3	0.6	1.1	1.6	2.4
2001.....	*	*	0.1	*	0.2	0.5	0.7	0.9	1.4	2.5
2000.....	*	*	0.1	0.2	0.2	0.4	0.6	1.0	1.7	2.5
1999.....	*	*	0.2	0.2	0.3	0.4	0.6	1.2	1.8	2.9

All races 3/, male

2006.....	*	*	0.2	0.4	0.8	1.3	1.8	4.3	7.3	16.9
2005.....	*	*	0.2	0.3	0.4	0.6	0.9	1.8	3.0	7.3
2004.....	*	*	*	*	0.3	*	0.6	1.1	1.9	3.5
2003.....	*	*	0.2	*	*	0.5	0.5	1.0	1.9	3.6
2002.....	*	*	*	*	*	*	0.8	1.5	1.8	3.3
2001.....	*	*	*	*	*	0.5	1.0	1.2	2.2	3.5
2000.....	*	*	*	*	*	0.5	0.7	1.4	1.7	3.8
1999.....	*	*	*	*	*	*	0.5	1.7	2.4	4.4

All races 3/, female

2006.....	*	*	0.2	0.4	0.4	0.8	1.7	2.3	4.8	11.6
2005.....	*	*	*	*	0.3	0.5	0.7	1.3	1.8	4.2
2004.....	*	*	*	*	*	*	0.5	0.8	1.3	2.1
2003.....	*	*	*	*	*	*	0.7	0.8	1.3	2.0
2002.....	*	*	*	*	*	*	0.4	0.8	1.4	1.9
2001.....	*	*	*	*	*	0.5	0.4	0.6	1.0	2.1
2000.....	*	*	*	*	*	*	0.6	0.7	1.7	1.9
1999.....	*	*	*	*	*	0.4	0.7	0.9	1.4	2.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified glomerular diseases (N02-N03,N05-N07)

White, both sexes

2006 4/.....	*	*	0.2	0.3	0.4	0.7	1.5	2.9	5.2	12.4
2005 4/.....	*	*	*	0.2	0.3	0.4	0.7	1.4	2.2	4.9
2004 4/.....	*	*	*	0.1	0.2	0.3	0.5	0.9	1.5	2.4
2003 4/.....	*	*	*	*	0.2	0.4	0.5	0.8	1.4	2.3
2002.....	*	*	*	0.2	*	0.3	0.6	1.0	1.5	2.2
2001.....	*	*	*	*	*	0.4	0.7	0.9	1.4	2.5
2000.....	*	*	*	0.2	*	0.4	0.6	0.9	1.6	2.1
1999.....	*	*	*	*	0.3	0.3	0.6	1.2	1.7	2.7

White, male

2006 4/.....	*	*	*	*	0.6	0.9	1.7	3.8	6.7	15.7
2005 4/.....	*	*	*	*	*	0.5	0.8	1.7	2.9	7.1
2004 4/.....	*	*	*	*	*	*	*	1.0	1.9	3.2
2003 4/.....	*	*	*	*	*	0.5	*	0.9	1.8	3.3
2002.....	*	*	*	*	*	*	0.7	1.3	1.8	3.3
2001.....	*	*	*	*	*	*	1.1	1.2	2.2	3.7
2000.....	*	*	*	*	*	*	0.7	1.3	1.8	3.2
1999.....	*	*	*	*	*	*	*	1.7	2.2	4.2

White, female

2006 4/.....	*	*	*	0.3	*	0.6	1.4	2.2	4.3	10.9
2005 4/.....	*	*	*	*	*	*	0.5	1.2	1.7	3.9
2004 4/.....	*	*	*	*	*	*	0.5	0.7	1.2	2.1
2003 4/.....	*	*	*	*	*	*	0.6	0.7	1.2	1.8
2002.....	*	*	*	*	*	*	0.5	0.7	1.3	1.7
2001.....	*	*	*	*	*	*	*	0.6	0.9	2.0
2000.....	*	*	*	*	*	*	0.6	0.6	1.5	1.6
1999.....	*	*	*	*	*	*	0.7	0.8	1.3	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Renal tubulo-interstitial diseases (N10-N15)

All races 3/, both sexes

2006.....	0.1	0.1	0.2	0.3	0.4	0.6	1.2	1.7	3.3	7.8
2005.....	0.1	0.1	0.2	0.2	0.3	0.6	1.1	2.2	4.1	8.9
2004.....	0.1	0.1	0.2	0.3	0.3	0.6	1.2	2.3	4.4	9.3
2003.....	0.1	0.2	0.2	0.3	0.4	0.7	1.0	2.1	4.8	10.3
2002.....	*	0.1	0.2	0.2	0.4	0.7	1.2	2.6	4.1	10.6
2001.....	*	0.1	0.2	0.2	0.4	0.6	1.1	2.5	4.0	10.4
2000.....	*	0.1	0.2	0.3	0.4	0.6	1.1	2.5	4.1	12.0
1999.....	0.1	0.1	0.3	0.3	0.4	0.8	1.3	2.5	4.9	12.7

All races 3/, male

2006.....	*	*	*	0.2	*	0.5	1.4	1.6	4.0	9.2
2005.....	*	*	*	*	0.4	0.6	1.0	1.9	4.7	10.2
2004.....	*	*	*	0.4	*	0.6	1.1	2.3	4.6	12.7
2003.....	*	*	*	*	0.3	0.5	1.0	2.3	5.2	13.8
2002.....	*	*	*	*	0.4	0.7	1.5	3.1	4.3	13.8
2001.....	*	*	*	*	*	0.6	1.3	2.3	5.0	12.5
2000.....	*	*	*	*	*	0.5	0.8	2.6	4.6	14.4
1999.....	*	*	*	0.3	0.4	0.8	1.1	2.6	6.4	17.0

All races 3/, female

2006.....	*	0.2	0.2	0.3	0.4	0.7	1.0	1.8	2.9	7.1
2005.....	*	0.2	0.2	0.2	0.3	0.7	1.2	2.4	3.8	8.3
2004.....	*	0.2	0.2	0.2	*	0.6	1.2	2.2	4.3	7.8
2003.....	*	0.2	0.2	0.4	0.4	1.0	1.0	2.0	4.5	8.7
2002.....	*	*	*	*	0.5	0.7	1.1	2.2	3.9	9.2
2001.....	*	*	0.3	0.3	0.5	0.5	0.9	2.7	3.4	9.5
2000.....	*	*	*	0.3	0.5	0.7	1.4	2.5	3.8	11.1
1999.....	*	*	0.3	0.4	*	0.8	1.5	2.4	4.0	11.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Renal tubulo-interstitial diseases (N10-N15)

White, both sexes

2006 4/.....	*	0.1	0.2	0.3	0.3	0.6	1.1	1.7	3.4	8.1
2005 4/.....	*	0.1	0.2	0.2	0.3	0.6	1.1	2.0	4.3	8.9
2004 4/.....	0.1	0.1	0.1	0.3	0.3	0.6	1.2	2.2	4.5	9.3
2003 4/.....	0.1	0.2	0.2	0.3	0.3	0.7	1.0	2.1	4.6	10.4
2002.....	*	*	0.2	0.2	0.4	0.8	1.3	2.5	4.2	10.9
2001.....	*	*	0.2	0.2	0.3	0.5	1.1	2.6	3.8	10.6
2000.....	*	0.1	*	0.3	0.4	0.6	1.0	2.4	3.9	12.0
1999.....	*	0.2	0.3	0.4	0.3	0.7	1.2	2.5	5.0	12.9

White, male

2006 4/.....	*	*	*	*	*	0.5	1.4	1.7	4.1	9.5
2005 4/.....	*	*	*	*	*	0.5	1.0	1.8	4.9	10.1
2004 4/.....	*	*	*	0.3	*	0.6	1.1	2.3	4.8	12.6
2003 4/.....	*	*	*	*	*	*	1.1	2.3	4.9	14.3
2002.....	*	*	*	*	*	0.6	1.5	2.9	4.3	14.3
2001.....	*	*	*	*	*	*	1.2	2.2	4.7	12.8
2000.....	*	*	*	*	*	*	0.7	2.5	4.5	14.6
1999.....	*	*	*	*	*	0.6	0.9	2.6	6.6	17.0

White, female

2006 4/.....	*	*	0.2	0.3	0.4	0.8	0.9	1.7	3.0	7.4
2005 4/.....	*	*	*	*	*	0.7	1.2	2.2	3.9	8.4
2004 4/.....	*	*	*	*	*	0.6	1.3	2.2	4.3	7.8
2003 4/.....	*	*	*	0.3	*	1.0	1.0	2.0	4.4	8.6
2002.....	*	*	*	*	0.4	0.8	1.2	2.3	4.1	9.4
2001.....	*	*	*	*	0.5	0.6	1.0	2.8	3.2	9.6
2000.....	*	*	*	*	0.4	0.7	1.3	2.4	3.5	10.9
1999.....	*	*	0.3	0.4	*	0.7	1.4	2.4	4.1	11.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
All races 3/, both sexes										
2006.....	2.2	3.7	6.4	10.2	17.5	29.3	48.4	83.0	137.5	276.4
2005.....	2.0	3.8	5.6	10.3	17.3	28.6	50.2	83.7	140.0	282.1
2004.....	2.1	3.6	6.2	10.3	17.3	29.1	48.2	84.0	136.8	282.9
2003.....	2.2	3.7	5.9	10.2	17.6	30.2	49.7	83.2	141.7	289.7
2002.....	2.0	3.6	5.7	9.8	16.5	29.0	49.1	83.9	139.8	284.8
2001.....	2.0	3.5	5.7	9.7	16.6	31.1	48.4	80.9	133.8	284.0
2000.....	2.0	3.3	5.3	9.2	16.6	28.9	46.2	78.1	130.0	273.7
1999.....	1.9	3.0	4.9	8.7	15.3	27.7	45.7	75.7	124.8	264.8
All races 3/, male										
2006.....	2.6	4.5	7.7	12.0	20.2	32.4	57.4	100.4	174.0	358.8
2005.....	2.3	4.7	6.9	12.0	20.5	30.7	57.8	100.4	180.3	368.0
2004.....	2.5	4.3	7.4	11.6	19.1	32.8	55.3	100.5	178.4	367.4
2003.....	2.6	4.3	7.4	12.0	19.3	33.9	57.2	101.3	185.2	383.1
2002.....	2.2	4.2	6.7	10.6	18.4	33.1	58.8	103.9	180.0	380.7
2001.....	2.2	4.2	6.5	10.7	17.9	35.9	55.8	98.9	173.9	388.7
2000.....	2.4	4.2	6.1	10.4	18.1	32.1	53.3	97.9	170.1	374.2
1999.....	2.2	3.6	5.6	9.7	17.3	30.5	53.9	96.7	165.1	364.6
All races 3/, female										
2006.....	1.8	2.8	5.2	8.6	15.1	26.6	41.0	70.2	114.8	237.8
2005.....	1.7	2.8	4.4	8.6	14.3	26.7	44.0	71.6	115.0	242.6
2004.....	1.8	3.0	5.0	9.2	15.6	25.8	42.5	72.1	111.2	244.9
2003.....	1.8	3.2	4.6	8.5	16.0	27.0	43.6	70.2	115.0	248.4
2002.....	1.8	2.9	4.8	9.0	14.9	25.4	41.4	69.9	115.3	243.2
2001.....	1.8	2.8	4.8	8.8	15.5	27.0	42.5	68.3	109.8	240.1
2000.....	1.6	2.4	4.5	8.0	15.2	26.1	40.6	64.4	106.4	232.8
1999.....	1.6	2.3	4.3	7.9	13.5	25.3	39.3	61.1	101.2	224.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
White, both sexes										
2006 4/.....	1.4	2.8	4.5	7.8	13.7	23.7	41.4	76.0	130.6	268.2
2005 4/.....	1.3	2.6	4.2	7.4	13.3	23.3	43.3	76.8	132.7	274.3
2004 4/.....	1.5	2.5	4.3	7.6	13.3	23.8	42.1	76.8	130.1	274.7
2003 4/.....	1.4	2.5	4.1	7.6	13.8	24.7	42.7	76.4	133.9	281.5
2002.....	1.4	2.5	3.8	7.1	12.9	23.9	42.3	76.8	133.2	275.3
2001.....	1.4	2.3	4.0	7.1	12.6	25.2	42.1	73.8	126.0	274.6
2000.....	1.4	2.1	3.7	6.6	12.8	23.3	39.6	71.7	122.6	265.0
1999.....	1.2	1.8	3.4	6.4	11.9	22.5	39.4	69.0	116.2	251.9
White, male										
2006 4/.....	1.8	3.5	5.5	9.1	16.5	26.8	50.6	93.7	167.8	356.8
2005 4/.....	1.6	3.4	5.3	8.6	16.3	26.5	51.4	93.7	174.6	365.4
2004 4/.....	1.9	2.9	5.1	8.8	14.9	27.1	49.5	94.0	173.7	362.3
2003 4/.....	1.8	3.0	4.9	9.0	15.3	28.4	49.7	95.6	177.3	379.3
2002.....	1.5	2.9	4.5	7.7	14.0	27.9	52.0	97.3	175.3	372.7
2001.....	1.6	2.7	4.7	8.0	14.1	30.0	50.1	93.0	166.1	383.9
2000.....	1.7	2.7	4.2	7.6	14.5	27.2	47.7	91.6	163.8	370.4
1999.....	1.4	2.2	3.8	7.6	13.6	25.5	48.0	89.1	156.2	354.9
White, female										
2006 4/.....	1.0	2.0	3.5	6.5	11.1	20.9	33.8	62.8	107.1	226.6
2005 4/.....	1.0	1.9	3.2	6.3	10.5	20.6	36.7	64.3	106.4	232.4
2004 4/.....	1.1	2.1	3.5	6.4	11.8	21.0	36.0	64.2	103.1	235.2
2003 4/.....	1.0	2.1	3.3	6.4	12.4	21.4	36.9	62.4	107.1	238.2
2002.....	1.2	2.1	3.2	6.6	11.8	20.3	34.4	62.1	107.4	232.9
2001.....	1.1	1.9	3.3	6.2	11.2	21.1	35.6	60.2	101.8	228.8
2000.....	1.1	1.6	3.1	5.6	11.3	19.9	33.2	57.8	98.1	222.1
1999.....	1.1	1.5	2.9	5.3	10.3	19.9	32.5	54.8	92.5	210.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Renal failure (N17-N19)										
Black, both sexes										
2006 4/.....	7.4	10.6	21.3	31.3	53.6	82.6	118.0	172.2	242.1	418.7
2005 4/.....	7.0	11.9	17.6	34.9	54.9	78.4	122.1	172.7	250.0	413.7
2004 4/.....	6.6	12.2	21.3	35.2	56.1	78.0	114.8	173.3	239.6	426.4
2003 4/.....	7.8	12.4	20.5	34.2	53.6	82.9	126.1	168.8	257.4	436.4
2002.....	6.6	11.2	21.1	33.3	50.1	78.8	125.7	175.4	238.1	434.2
2001.....	6.5	12.3	19.3	32.2	52.5	87.1	119.0	170.6	245.9	425.6
2000.....	6.1	12.3	18.9	32.0	50.1	82.1	118.6	163.1	238.9	400.4
1999.....	6.7	11.7	18.9	28.5	46.1	76.5	115.7	161.0	241.6	425.3
Black, male										
2006 4/.....	8.9	13.1	26.4	38.3	61.8	93.7	132.3	202.3	297.1	454.9
2005 4/.....	8.0	15.6	21.9	44.7	67.1	78.7	131.2	203.4	297.1	469.4
2004 4/.....	6.8	15.0	27.6	39.1	63.0	91.1	127.9	196.5	282.7	506.7
2003 4/.....	8.6	14.4	28.7	43.3	61.9	92.6	153.7	188.1	328.5	513.0
2002.....	7.4	14.4	26.1	39.4	61.2	88.8	148.2	203.5	278.2	555.3
2001.....	6.8	16.1	22.1	37.2	54.5	99.1	129.9	187.7	307.4	508.3
2000.....	7.7	17.1	22.7	37.0	53.2	84.8	127.0	196.6	287.5	480.9
1999.....	8.0	15.3	22.7	28.7	53.5	80.5	128.9	204.7	320.9	513.2
Black, female										
2006 4/.....	6.1	8.4	17.0	25.4	47.2	74.4	108.2	154.3	213.0	403.9
2005 4/.....	6.1	8.6	13.9	26.8	45.3	78.3	115.8	154.4	225.2	391.2
2004 4/.....	6.3	9.8	16.0	31.9	50.7	68.4	105.9	159.6	217.0	394.6
2003 4/.....	7.0	10.7	13.7	26.7	47.1	75.7	107.4	157.4	220.1	406.7
2002.....	5.9	8.5	16.8	28.2	41.5	71.4	110.7	158.6	217.2	387.7
2001.....	6.4	9.0	16.9	28.1	50.9	78.3	111.7	160.3	214.6	394.9
2000.....	4.7	8.2	15.7	27.8	47.6	80.1	113.0	143.0	214.4	370.4
1999.....	5.6	8.5	15.8	28.4	40.3	73.5	106.9	134.7	202.2	392.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Urolithiasis (N20-N23)										
All races 3/, both sexes										
2006.....	*	*	*	0.1	0.2	*	0.3	0.7	1.1	1.5
2005.....	*	*	*	*	0.2	0.3	0.3	0.5	1.1	2.1
2004.....	*	*	*	*	*	0.3	0.3	0.6	1.0	1.8
2003.....	*	*	*	*	*	0.2	0.3	0.6	0.9	2.2
2002.....	*	*	*	*	*	0.2	0.3	0.5	0.9	1.8
2001.....	*	*	*	*	*	*	0.3	0.6	0.7	1.8
2000.....	*	*	*	*	*	*	0.2	0.7	0.9	1.8
1999.....	*	*	*	*	0.2	*	0.3	0.3	0.9	1.9
All races 3/, male										
2006.....	*	*	*	*	*	*	*	0.8	1.1	1.7
2005.....	*	*	*	*	*	*	*	*	1.4	2.3
2004.....	*	*	*	*	*	*	*	0.7	*	1.9
2003.....	*	*	*	*	*	*	*	*	*	2.6
2002.....	*	*	*	*	*	*	*	*	*	2.0
2001.....	*	*	*	*	*	*	*	*	*	2.0
2000.....	*	*	*	*	*	*	*	*	1.1	1.7
1999.....	*	*	*	*	*	*	*	*	*	*
All races 3/, female										
2006.....	*	*	*	*	*	*	*	0.7	1.0	1.4
2005.....	*	*	*	*	*	*	*	0.6	0.9	2.0
2004.....	*	*	*	*	*	0.4	*	0.5	1.0	1.8
2003.....	*	*	*	*	*	*	*	0.6	0.8	2.1
2002.....	*	*	*	*	*	*	*	0.5	0.9	1.7
2001.....	*	*	*	*	*	*	*	0.7	0.6	1.7
2000.....	*	*	*	*	*	*	*	0.8	0.8	1.8
1999.....	*	*	*	*	*	*	*	*	0.8	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Urolithiasis (N20-N23)										
White, both sexes										
2006 4/.....	*	*	*	0.1	*	*	0.4	0.7	1.1	1.5
2005 4/.....	*	*	*	*	*	0.3	0.3	0.5	1.1	2.2
2004 4/.....	*	*	*	*	*	0.3	0.3	0.6	1.0	1.8
2003 4/.....	*	*	*	*	*	*	0.3	0.6	0.9	2.3
2002.....	*	*	*	*	*	0.2	0.3	0.5	0.9	1.9
2001.....	*	*	*	*	*	*	0.3	0.6	0.8	1.9
2000.....	*	*	*	*	*	*	*	0.7	1.0	1.8
1999.....	*	*	*	*	*	*	0.3	0.3	1.0	1.9
White, male										
2006 4/.....	*	*	*	*	*	*	*	0.8	1.2	1.8
2005 4/.....	*	*	*	*	*	*	*	*	1.4	2.4
2004 4/.....	*	*	*	*	*	*	*	0.7	*	2.0
2003 4/.....	*	*	*	*	*	*	*	*	*	2.8
2002.....	*	*	*	*	*	*	*	*	*	2.1
2001.....	*	*	*	*	*	*	*	*	*	2.1
2000.....	*	*	*	*	*	*	*	*	1.3	1.9
1999.....	*	*	*	*	*	*	*	*	*	*
White, female										
2006 4/.....	*	*	*	*	*	*	*	0.7	1.0	1.4
2005 4/.....	*	*	*	*	*	*	*	0.6	1.0	2.0
2004 4/.....	*	*	*	*	*	0.4	*	0.5	1.1	1.8
2003 4/.....	*	*	*	*	*	*	*	0.7	0.9	2.1
2002.....	*	*	*	*	*	*	*	*	1.0	1.8
2001.....	*	*	*	*	*	*	*	0.7	*	1.8
2000.....	*	*	*	*	*	*	*	0.8	0.8	1.7
1999.....	*	*	*	*	*	*	*	*	0.9	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Urinary tract infection, site not specified (N39.0)

All races 3/, both sexes

2006.....	0.3	0.4	0.9	1.4	2.8	4.9	10.8	22.9	51.5	139.1
2005.....	0.2	0.4	0.8	1.4	2.7	5.0	11.6	25.8	52.8	149.0
2004.....	0.2	0.4	0.9	1.4	2.2	5.3	10.4	23.4	51.9	143.5
2003.....	0.2	0.3	0.8	1.3	2.7	4.9	11.2	23.3	52.2	147.7
2002.....	0.2	0.4	0.7	1.4	2.4	4.8	10.8	23.7	52.2	152.2
2001.....	0.2	0.4	0.6	1.4	2.3	5.3	11.4	24.5	54.9	160.4
2000.....	0.2	0.4	0.7	1.2	2.5	4.9	9.9	25.6	57.7	163.4
1999.....	0.2	0.4	0.6	1.2	2.6	5.1	11.2	25.4	57.2	172.0

All races 3/, male

2006.....	0.2	0.4	0.8	1.2	2.3	4.5	9.7	20.6	43.2	115.3
2005.....	0.2	0.4	0.8	1.2	2.6	4.3	10.7	23.1	46.2	119.3
2004.....	0.2	0.4	0.7	1.3	2.0	4.9	9.0	22.2	46.7	117.7
2003.....	0.2	0.3	0.6	1.0	2.6	4.5	9.8	22.4	49.3	114.6
2002.....	0.2	0.4	0.5	1.2	2.2	4.3	9.5	20.5	47.1	123.5
2001.....	*	0.3	0.5	1.2	2.1	5.0	10.1	21.6	50.1	125.8
2000.....	*	0.4	0.6	1.0	2.1	4.5	9.2	23.4	52.4	139.8
1999.....	0.2	0.4	0.4	1.1	2.6	4.4	10.9	23.3	53.5	137.5

All races 3/, female

2006.....	0.4	0.5	0.9	1.6	3.3	5.3	11.6	24.5	56.6	150.2
2005.....	0.2	0.5	0.9	1.6	2.7	5.5	12.3	27.7	56.8	162.7
2004.....	0.2	0.4	1.0	1.4	2.4	5.7	11.5	24.3	55.2	155.1
2003.....	0.2	0.4	0.9	1.6	2.9	5.3	12.3	24.0	53.9	162.3
2002.....	0.2	0.4	0.8	1.6	2.6	5.3	11.9	26.0	55.3	164.7
2001.....	0.2	0.5	0.7	1.5	2.5	5.6	12.4	26.6	57.8	174.9
2000.....	*	0.4	0.7	1.3	2.8	5.1	10.5	27.1	60.8	173.0
1999.....	0.3	0.5	0.8	1.3	2.5	5.6	11.4	26.9	59.3	185.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Urinary tract infection, site not specified (N39.0)										
White, both sexes										
2006 4/.....	0.2	0.4	0.8	1.3	2.7	4.6	10.0	22.4	51.4	139.6
2005 4/.....	0.2	0.4	0.7	1.3	2.5	4.6	10.9	25.0	52.1	148.8
2004 4/.....	0.2	0.4	0.8	1.2	2.1	4.9	9.9	22.4	50.9	143.2
2003 4/.....	0.2	0.3	0.7	1.1	2.4	4.6	10.4	22.2	51.2	147.2
2002.....	0.2	0.4	0.6	1.3	2.2	4.4	10.2	22.8	50.9	150.8
2001.....	0.2	0.4	0.5	1.2	2.1	4.8	10.6	23.3	53.2	158.2
2000.....	0.1	0.4	0.5	1.1	2.2	4.5	9.1	24.4	56.0	161.9
1999.....	0.2	0.4	0.5	1.2	2.2	4.4	10.3	24.0	56.3	169.8
White, male										
2006 4/.....	*	0.4	0.7	1.1	2.0	4.4	8.7	20.2	42.8	115.7
2005 4/.....	0.2	0.4	0.7	1.1	2.3	3.6	9.9	22.0	45.4	119.7
2004 4/.....	*	0.4	0.6	1.1	1.7	4.3	8.5	20.9	45.7	115.9
2003 4/.....	*	0.3	0.5	0.9	2.3	3.9	9.0	21.0	48.2	116.5
2002.....	*	0.4	0.5	1.1	2.0	3.9	8.6	19.3	45.6	121.1
2001.....	*	0.3	0.4	0.9	1.9	4.4	9.2	20.4	49.3	122.1
2000.....	*	0.3	0.5	0.9	1.8	4.0	8.3	22.1	50.2	137.8
1999.....	*	0.3	0.4	1.1	2.2	3.6	9.4	21.3	52.1	132.1
White, female										
2006 4/.....	0.3	0.5	0.8	1.5	3.3	4.7	11.2	24.1	56.9	150.8
2005 4/.....	*	0.5	0.7	1.6	2.6	5.5	11.7	27.2	56.2	162.2
2004 4/.....	0.2	0.4	0.9	1.2	2.4	5.5	11.1	23.5	54.1	155.5
2003 4/.....	*	0.3	0.8	1.4	2.6	5.1	11.6	23.2	53.1	160.8
2002.....	*	0.4	0.7	1.6	2.3	4.7	11.5	25.3	54.2	163.8
2001.....	0.2	0.5	0.5	1.5	2.3	5.1	11.8	25.3	55.6	173.3
2000.....	*	0.4	0.6	1.3	2.5	4.9	9.7	26.0	59.4	171.7
1999.....	0.2	0.4	0.6	1.3	2.3	5.1	10.9	25.9	58.9	185.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Urinary tract infection, site not specified (N39.0)

Black, both sexes

2006 4/.....	0.8	*	1.9	2.8	4.2	9.1	18.9	33.0	63.7	159.0
2005 4/.....	*	*	1.9	2.4	4.3	9.2	19.7	38.1	73.3	174.8
2004 4/.....	*	1.0	1.8	3.7	4.2	9.4	17.5	37.2	73.3	172.9
2003 4/.....	*	*	1.8	3.0	6.0	9.2	21.2	40.3	73.4	178.5
2002.....	*	*	1.4	2.8	5.0	9.0	18.6	38.7	76.9	191.5
2001.....	*	1.0	1.4	2.5	4.4	9.6	20.3	41.7	83.7	208.5
2000.....	*	*	1.7	2.2	5.6	9.1	19.3	43.4	85.8	196.8
1999.....	*	0.9	1.4	1.8	6.0	11.9	21.6	45.7	74.3	213.0

Black, male

2006 4/.....	*	*	*	2.8	5.3	7.3	21.5	32.2	61.6	144.8
2005 4/.....	*	*	1.9	2.9	4.5	11.2	22.1	38.9	72.3	139.4
2004 4/.....	*	*	*	3.8	4.5	10.2	17.8	44.6	66.0	162.9
2003 4/.....	*	*	*	*	6.2	10.6	21.0	45.2	79.0	125.1
2002.....	*	*	*	3.0	4.2	8.2	20.4	40.9	77.9	188.7
2001.....	*	*	*	3.1	4.5	10.7	20.2	41.4	72.2	204.2
2000.....	*	*	*	*	6.0	9.9	20.1	45.4	91.0	175.3
1999.....	*	*	*	*	7.0	13.0	29.8	51.8	80.2	215.6

Black, female

2006 4/.....	*	*	2.1	2.8	3.3	10.4	17.1	33.5	64.7	164.9
2005 4/.....	*	*	1.9	*	4.1	7.8	18.0	37.6	73.8	189.1
2004 4/.....	*	*	2.0	3.6	3.9	8.8	17.4	32.9	77.1	176.8
2003 4/.....	*	*	2.0	3.7	5.9	8.2	21.3	37.4	70.4	199.3
2002.....	*	*	*	2.6	5.6	9.6	17.5	37.3	76.4	192.6
2001.....	*	*	*	*	4.3	8.8	20.4	41.9	89.5	210.1
2000.....	*	*	*	*	5.2	8.4	18.9	42.2	83.2	204.8
1999.....	*	*	*	*	5.2	11.2	16.2	42.1	71.3	212.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of urinary system (N28-N36,N39.1-N39.9)

All races 3/, both sexes

2006.....	0.1	0.1	0.3	0.4	0.5	0.9	1.5	2.2	3.4	6.9
2005.....	0.1	0.2	0.3	0.5	0.6	1.0	1.3	2.1	3.5	6.8
2004.....	*	0.2	0.2	0.4	0.6	0.9	1.3	2.1	3.8	7.8
2003.....	0.1	0.1	0.3	0.4	0.7	0.7	1.2	1.9	3.5	7.1
2002.....	0.1	0.2	0.3	0.3	0.6	0.9	1.2	1.7	3.9	7.5
2001.....	0.1	0.2	0.3	0.4	0.6	0.9	1.2	2.1	3.5	7.1
2000.....	0.2	0.2	0.3	0.4	0.6	0.8	1.2	2.0	3.9	7.5
1999.....	*	0.1	0.2	0.4	0.6	0.8	1.3	1.8	3.8	7.1

All races 3/, male

2006.....	*	*	0.3	0.5	0.6	1.1	1.7	2.9	4.7	8.4
2005.....	*	*	0.3	0.6	0.8	1.3	1.9	2.6	4.7	9.4
2004.....	*	0.2	0.3	0.5	0.8	1.0	1.7	2.7	5.4	9.2
2003.....	*	*	0.3	0.4	0.9	0.8	1.5	2.3	4.6	9.8
2002.....	*	0.2	0.4	0.4	0.7	1.0	1.4	2.1	4.9	9.0
2001.....	*	*	0.3	0.4	0.7	1.1	1.4	2.4	4.4	8.9
2000.....	0.2	*	0.3	0.5	0.6	1.1	1.5	2.3	5.6	10.4
1999.....	*	0.2	0.2	0.5	0.6	1.2	1.8	2.2	4.7	9.9

All races 3/, female

2006.....	*	*	0.2	0.4	0.5	0.7	1.4	1.7	2.6	6.2
2005.....	*	0.2	0.2	0.4	0.4	0.7	0.9	1.7	2.7	5.6
2004.....	*	*	*	0.2	0.5	0.8	1.1	1.7	2.9	7.1
2003.....	*	*	0.2	0.3	0.6	0.6	0.9	1.6	2.9	6.0
2002.....	*	*	*	*	0.4	0.8	1.0	1.4	3.4	6.9
2001.....	*	*	*	0.4	0.5	0.7	1.0	1.9	2.9	6.3
2000.....	*	*	0.2	0.4	0.5	0.6	1.0	1.8	2.9	6.3
1999.....	*	*	*	0.3	0.5	0.4	0.9	1.5	3.2	6.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of urinary system (N28-N36,N39.1-N39.9)

White, both sexes

2006 4/.....	*	0.1	0.2	0.4	0.4	0.9	1.4	2.1	3.5	6.8
2005 4/.....	*	0.1	0.3	0.4	0.5	0.9	1.2	2.0	3.4	6.9
2004 4/.....	*	0.2	0.2	0.3	0.5	0.9	1.2	2.0	3.7	7.8
2003 4/.....	*	0.1	0.2	0.3	0.5	0.7	1.1	1.9	3.6	7.2
2002.....	*	*	0.2	0.2	0.4	0.8	1.1	1.6	3.8	7.5
2001.....	*	0.1	0.2	0.3	0.5	0.8	1.0	2.1	3.4	7.2
2000.....	0.2	0.1	0.3	0.3	0.5	0.7	1.0	1.9	3.8	7.3
1999.....	*	*	0.2	0.3	0.5	0.7	1.2	1.7	3.4	7.1

White, male

2006 4/.....	*	*	*	0.4	0.5	1.0	1.5	2.7	4.9	8.6
2005 4/.....	*	*	0.3	0.5	0.7	1.2	1.9	2.4	4.6	9.8
2004 4/.....	*	*	*	0.4	0.8	1.0	1.5	2.6	5.6	9.3
2003 4/.....	*	*	*	0.4	0.7	0.8	1.5	2.3	4.5	9.7
2002.....	*	*	0.3	0.3	0.5	1.0	1.3	2.0	4.8	8.8
2001.....	*	*	*	*	0.6	0.9	1.3	2.4	4.5	9.1
2000.....	*	*	0.3	0.4	0.5	0.9	1.3	2.1	5.7	10.2
1999.....	*	*	*	0.4	0.5	0.9	1.6	2.1	4.5	10.1

White, female

2006 4/.....	*	*	*	0.3	0.4	0.7	1.3	1.6	2.7	6.0
2005 4/.....	*	*	0.3	0.4	*	0.6	0.7	1.6	2.7	5.6
2004 4/.....	*	*	*	*	*	0.8	0.9	1.6	2.5	7.1
2003 4/.....	*	*	*	*	*	0.6	0.8	1.6	3.1	6.1
2002.....	*	*	*	*	*	0.7	0.9	1.3	3.2	6.9
2001.....	*	*	*	*	0.5	0.6	0.8	1.8	2.8	6.3
2000.....	*	*	*	*	0.5	0.5	0.8	1.8	2.6	6.1
1999.....	*	*	*	*	0.5	0.5	0.9	1.4	2.8	6.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Disorders of female genital organs (N70-N98)

All races 3/, both sexes

2006.....	*	0.1	*	*	*	*	0.3	0.5	1.2	2.2
2005.....	*	*	*	*	*	0.2	0.3	0.6	0.9	2.3
2004.....	*	*	*	*	*	*	0.4	0.4	0.9	2.5
2003.....	*	*	*	*	*	0.2	0.2	0.6	1.0	2.6
2002.....	*	0.1	*	*	*	*	*	0.4	0.9	2.4
2001.....	*	*	*	*	*	*	0.4	0.5	0.9	2.1
2000.....	*	*	*	*	*	0.2	0.3	0.5	1.0	2.5
1999.....	*	*	*	*	*	*	0.2	0.6	0.8	2.0

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	*	0.2	*	*	*	*	0.5	0.9	1.9	3.2
2005.....	*	*	*	*	*	0.4	0.6	1.0	1.5	3.3
2004.....	*	*	*	*	*	*	0.6	0.7	1.5	3.7
2003.....	*	*	*	*	*	0.4	0.4	1.1	1.5	3.7
2002.....	*	0.2	*	*	*	*	*	0.8	1.4	3.4
2001.....	*	*	*	*	*	*	0.6	0.9	1.5	3.0
2000.....	*	*	*	*	*	0.4	0.5	0.9	1.6	3.6
1999.....	*	*	*	*	*	*	0.4	1.0	1.3	2.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Disorders of female genital organs (N70-N98)										
White, both sexes										
2006 4/.....	*	*	*	*	*	*	0.3	0.5	1.2	2.3
2005 4/.....	*	*	*	*	*	*	0.3	0.5	0.9	2.2
2004 4/.....	*	*	*	*	*	*	0.3	0.5	0.9	2.5
2003 4/.....	*	*	*	*	*	*	*	0.6	1.0	2.8
2002.....	*	*	*	*	*	*	*	0.5	0.9	2.3
2001.....	*	*	*	*	*	*	0.3	0.5	0.9	2.1
2000.....	*	*	*	*	*	*	0.3	0.5	1.0	2.6
1999.....	*	*	*	*	*	*	*	0.5	0.9	2.0
White, male										
2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....
White, female										
2006 4/.....	*	*	*	*	*	*	0.5	0.9	1.9	3.3
2005 4/.....	*	*	*	*	*	*	0.5	0.8	1.5	3.3
2004 4/.....	*	*	*	*	*	*	0.5	0.8	1.5	3.6
2003 4/.....	*	*	*	*	*	*	*	1.1	1.6	4.0
2002.....	*	*	*	*	*	*	*	0.8	1.5	3.2
2001.....	*	*	*	*	*	*	0.6	0.8	1.4	2.9
2000.....	*	*	*	*	*	*	0.5	0.9	1.5	3.6
1999.....	*	*	*	*	*	*	*	0.8	1.4	2.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Noninflammatory disorders of female genital tract (N80-N98)

All races 3/, both sexes

2006.....	*	*	*	*	*	*	*	0.3	0.8	1.7
2005.....	*	*	*	*	*	*	*	0.4	0.5	1.7
2004.....	*	*	*	*	*	*	0.2	0.3	0.7	1.8
2003.....	*	*	*	*	*	*	*	0.4	0.7	1.9
2002.....	*	*	*	*	*	*	*	*	0.6	1.6
2001.....	*	*	*	*	*	*	0.2	0.3	0.6	1.2
2000.....	*	*	*	*	*	*	*	0.3	0.8	1.9
1999.....	*	*	*	*	*	*	*	0.4	0.5	1.6

All races 3/, male

2006.....
2005.....
2004.....
2003.....
2002.....
2001.....
2000.....
1999.....

All races 3/, female

2006.....	*	*	*	*	*	*	*	0.6	1.3	2.5
2005.....	*	*	*	*	*	*	*	0.7	0.9	2.5
2004.....	*	*	*	*	*	*	0.4	0.5	1.2	2.7
2003.....	*	*	*	*	*	*	*	0.7	1.2	2.8
2002.....	*	*	*	*	*	*	*	*	1.0	2.3
2001.....	*	*	*	*	*	*	0.4	0.6	0.9	1.8
2000.....	*	*	*	*	*	*	*	0.5	1.2	2.7
1999.....	*	*	*	*	*	*	*	0.6	0.7	2.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Noninflammatory disorders of female genital tract (N80-N98)

White, both sexes

2006 4/.....	*	*	*	*	*	*	*	*	0.9	1.7
2005 4/.....	*	*	*	*	*	*	*	0.3	0.5	1.7
2004 4/.....	*	*	*	*	*	*	*	0.3	0.8	1.8
2003 4/.....	*	*	*	*	*	*	*	0.4	0.8	2.1
2002.....	*	*	*	*	*	*	*	*	0.7	1.5
2001.....	*	*	*	*	*	*	*	0.3	0.6	1.2
2000.....	*	*	*	*	*	*	*	0.3	0.7	2.0
1999.....	*	*	*	*	*	*	*	0.3	0.5	1.6

White, male

2006 4/.....
2005 4/.....
2004 4/.....
2003 4/.....
2002.....
2001.....
2000.....
1999.....

White, female

2006 4/.....	*	*	*	*	*	*	*	*	1.4	2.6
2005 4/.....	*	*	*	*	*	*	*	0.5	0.8	2.5
2004 4/.....	*	*	*	*	*	*	*	0.6	1.2	2.6
2003 4/.....	*	*	*	*	*	*	*	0.7	1.3	3.0
2002.....	*	*	*	*	*	*	*	*	1.1	2.1
2001.....	*	*	*	*	*	*	*	0.5	0.9	1.7
2000.....	*	*	*	*	*	*	*	0.5	1.1	2.7
1999.....	*	*	*	*	*	*	*	0.6	0.8	2.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2006.....	1.1	1.5	1.6	2.2	2.4	1.9	2.0	2.7	3.5	5.6
2005.....	1.2	1.5	1.8	2.3	2.4	1.7	2.0	2.4	4.0	6.4
2004.....	1.2	1.5	1.8	2.2	2.2	2.3	2.3	3.1	4.3	6.1
2003.....	1.4	1.5	1.8	2.4	2.4	2.3	2.2	3.0	4.4	6.8
2002.....	1.3	1.6	1.9	2.3	2.4	2.3	2.5	3.1	4.8	6.6
2001.....	1.4	1.5	1.8	2.1	2.5	2.2	2.7	3.3	4.3	7.2
2000.....	1.3	1.4	1.8	2.2	2.5	2.6	2.4	3.3	5.0	7.5
1999.....	1.2	1.7	1.8	2.3	2.4	2.6	2.5	3.4	5.0	7.8

All races 3/, male

2006.....	1.2	1.7	1.9	2.5	2.7	1.9	2.3	3.0	4.5	6.8
2005.....	1.3	1.5	1.9	2.5	2.6	1.6	2.0	2.8	4.9	6.9
2004.....	1.3	1.7	2.0	2.3	2.2	2.3	2.7	3.7	3.8	6.8
2003.....	1.5	1.6	1.8	2.5	2.6	2.4	2.4	2.8	4.5	6.7
2002.....	1.4	1.9	2.0	2.3	2.7	2.4	2.7	3.4	4.8	7.0
2001.....	1.6	1.5	1.9	2.1	2.5	2.5	3.1	3.6	4.8	7.7
2000.....	1.5	1.6	2.2	2.0	2.6	2.7	2.6	3.3	5.3	10.6
1999.....	1.3	1.9	2.0	2.5	2.5	2.4	2.6	3.4	5.7	8.5

All races 3/, female

2006.....	1.0	1.3	1.3	1.9	2.1	1.8	1.8	2.4	2.8	5.1
2005.....	1.2	1.4	1.8	2.1	2.3	1.7	2.0	2.2	3.5	6.2
2004.....	1.2	1.3	1.7	2.0	2.1	2.3	2.0	2.7	4.6	5.8
2003.....	1.3	1.3	1.8	2.3	2.3	2.2	2.1	3.2	4.4	6.8
2002.....	1.2	1.4	1.8	2.3	2.2	2.3	2.3	2.9	4.9	6.5
2001.....	1.3	1.4	1.8	2.1	2.6	2.0	2.4	3.1	4.0	7.0
2000.....	1.1	1.3	1.5	2.4	2.5	2.5	2.3	3.3	4.8	6.3
1999.....	1.2	1.6	1.7	2.2	2.3	2.8	2.3	3.4	4.5	7.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XVII. Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2006 4/.....	1.1	1.5	1.6	2.3	2.5	1.9	2.1	2.8	3.5	5.7
2005 4/.....	1.3	1.5	2.0	2.4	2.6	1.8	2.2	2.5	4.3	6.5
2004 4/.....	1.3	1.6	1.9	2.3	2.3	2.4	2.5	3.1	4.4	6.3
2003 4/.....	1.5	1.5	1.9	2.5	2.5	2.3	2.3	3.1	4.5	6.9
2002.....	1.3	1.7	2.0	2.4	2.5	2.5	2.5	3.2	4.8	6.9
2001.....	1.5	1.4	1.9	2.2	2.7	2.3	2.7	3.4	4.4	7.4
2000.....	1.3	1.5	2.0	2.2	2.6	2.6	2.4	3.4	5.2	7.8
1999.....	1.3	1.8	1.8	2.4	2.4	2.7	2.4	3.3	5.1	8.0

White, male

2006 4/.....	1.2	1.7	1.9	2.6	2.8	2.0	2.3	3.0	4.6	6.8
2005 4/.....	1.3	1.6	2.0	2.6	2.7	1.8	2.2	2.8	5.2	6.9
2004 4/.....	1.4	1.8	2.0	2.4	2.3	2.4	2.9	3.6	3.9	7.2
2003 4/.....	1.6	1.5	2.0	2.6	2.6	2.3	2.3	2.8	4.3	6.9
2002.....	1.4	1.8	2.0	2.3	2.8	2.6	2.7	3.4	4.6	7.3
2001.....	1.6	1.5	1.9	2.2	2.6	2.7	3.2	3.9	4.6	7.5
2000.....	1.4	1.6	2.3	2.0	2.8	2.6	2.6	3.4	5.6	11.1
1999.....	1.3	1.9	2.0	2.6	2.6	2.5	2.6	3.0	5.8	8.6

White, female

2006 4/.....	1.0	1.4	1.3	2.0	2.2	1.9	1.9	2.6	2.7	5.2
2005 4/.....	1.3	1.4	1.9	2.2	2.5	1.8	2.2	2.2	3.7	6.3
2004 4/.....	1.2	1.3	1.9	2.2	2.3	2.5	2.2	2.8	4.7	6.0
2003 4/.....	1.3	1.4	1.9	2.4	2.4	2.2	2.2	3.3	4.7	6.9
2002.....	1.2	1.5	2.0	2.4	2.2	2.4	2.4	3.0	5.0	6.7
2001.....	1.3	1.4	1.9	2.2	2.8	2.1	2.3	3.1	4.2	7.3
2000.....	1.1	1.3	1.7	2.4	2.4	2.7	2.3	3.4	4.9	6.5
1999.....	1.2	1.7	1.7	2.3	2.3	2.9	2.3	3.6	4.6	7.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congenital malformations of heart (Q20-Q24)										
All races 3/, both sexes										
2006.....	0.4	0.5	0.4	0.5	0.4	0.5	0.6	0.9	0.8	1.2
2005.....	0.5	0.5	0.5	0.4	0.4	0.4	0.6	0.7	1.0	1.6
2004.....	0.5	0.4	0.5	0.5	0.4	0.5	0.6	0.9	1.3	1.5
2003.....	0.6	0.5	0.5	0.5	0.5	0.7	0.6	0.8	1.5	2.1
2002.....	0.5	0.6	0.5	0.6	0.5	0.6	0.7	1.0	1.6	2.3
2001.....	0.6	0.5	0.6	0.5	0.5	0.6	0.7	1.1	1.4	2.6
2000.....	0.6	0.5	0.5	0.6	0.6	0.7	0.8	1.1	2.0	2.5
1999.....	0.6	0.6	0.6	0.6	0.7	0.9	0.9	1.2	1.7	3.3
All races 3/, male										
2006.....	0.5	0.5	0.5	0.5	0.5	0.6	0.5	0.7	1.0	*
2005.....	0.5	0.5	0.6	0.4	0.4	*	*	*	1.0	1.4
2004.....	0.5	0.4	0.6	0.6	0.5	*	0.8	0.9	1.0	1.5
2003.....	0.6	0.5	0.4	0.6	0.5	0.8	*	*	1.3	*
2002.....	0.5	0.6	0.5	0.6	0.5	0.5	0.7	1.0	1.3	1.4
2001.....	0.6	0.6	0.6	0.5	0.5	0.7	1.0	1.0	1.6	1.7
2000.....	0.6	0.5	0.6	0.6	0.6	0.8	0.8	1.1	2.0	2.7
1999.....	0.6	0.7	0.7	0.6	0.7	0.7	0.9	1.0	1.8	3.1
All races 3/, female										
2006.....	0.4	0.5	0.4	0.4	0.4	0.4	0.6	1.0	0.7	1.2
2005.....	0.5	0.5	0.5	0.3	0.4	0.4	0.7	0.8	0.9	1.7
2004.....	0.4	0.3	0.5	0.4	0.4	0.6	0.4	0.9	1.5	1.6
2003.....	0.6	0.5	0.5	0.5	0.5	0.7	0.7	1.0	1.6	2.4
2002.....	0.5	0.6	0.6	0.6	0.4	0.6	0.8	1.0	1.8	2.6
2001.....	0.6	0.5	0.6	0.4	0.5	0.6	0.5	1.2	1.3	3.0
2000.....	0.5	0.5	0.5	0.7	0.5	0.5	0.8	1.0	1.9	2.4
1999.....	0.6	0.6	0.5	0.7	0.6	1.0	0.8	1.3	1.7	3.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Congenital malformations of heart (Q20-Q24)										
White, both sexes										
2006 4/.....	0.5	0.6	0.4	0.5	0.5	0.5	0.6	0.8	0.9	1.2
2005 4/.....	0.5	0.5	0.6	0.4	0.5	0.4	0.6	0.7	1.0	1.6
2004 4/.....	0.5	0.4	0.6	0.5	0.5	0.5	0.6	0.9	1.4	1.6
2003 4/.....	0.7	0.5	0.5	0.5	0.5	0.7	0.7	0.8	1.6	2.2
2002.....	0.5	0.6	0.5	0.6	0.4	0.6	0.8	1.0	1.6	2.4
2001.....	0.6	0.6	0.6	0.4	0.5	0.6	0.7	1.2	1.5	2.6
2000.....	0.6	0.5	0.6	0.6	0.6	0.7	0.7	1.1	2.1	2.7
1999.....	0.6	0.7	0.6	0.7	0.6	0.9	0.9	1.3	1.8	3.3
White, male										
2006 4/.....	0.5	0.5	0.5	0.5	0.5	0.6	*	*	1.0	*
2005 4/.....	0.4	0.5	0.6	0.4	0.4	*	*	*	1.1	*
2004 4/.....	0.5	0.5	0.6	0.7	0.5	*	0.7	1.0	1.1	1.6
2003 4/.....	0.7	0.5	0.4	0.6	0.4	0.8	*	*	1.3	*
2002.....	0.5	0.6	0.5	0.6	0.4	0.6	0.7	1.0	1.3	1.6
2001.....	0.6	0.6	0.6	0.5	0.5	0.7	1.0	1.1	1.6	*
2000.....	0.6	0.6	0.6	0.6	0.7	0.8	0.7	1.2	2.2	2.9
1999.....	0.6	0.7	0.6	0.7	0.7	0.7	0.9	0.9	1.8	3.0
White, female										
2006 4/.....	0.4	0.6	0.3	0.4	0.4	*	0.6	1.0	0.7	1.2
2005 4/.....	0.5	0.5	0.5	0.3	0.5	0.5	0.7	0.8	1.0	1.8
2004 4/.....	0.4	0.4	0.5	0.4	0.5	0.6	*	0.9	1.5	1.6
2003 4/.....	0.6	0.5	0.5	0.5	0.6	0.6	0.8	1.0	1.8	2.6
2002.....	0.5	0.6	0.6	0.6	0.4	0.6	0.8	1.0	1.8	2.7
2001.....	0.6	0.5	0.7	0.4	0.5	0.6	0.5	1.2	1.4	3.1
2000.....	0.5	0.5	0.5	0.7	0.5	0.6	0.8	1.0	2.0	2.6
1999.....	0.7	0.6	0.5	0.7	0.6	1.0	0.8	1.5	1.8	3.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other congenital malformations of circulatory system (Q25-Q28)

All races 3/, both sexes

2006.....	*	0.1	0.1	0.1	0.2	0.2	0.3	0.5	0.6	1.4
2005.....	0.1	0.1	0.1	0.1	0.2	*	0.3	0.6	0.9	1.5
2004.....	0.2	0.2	0.2	0.3	0.2	0.3	0.4	0.9	1.0	1.6
2003.....	0.2	0.2	0.3	0.2	0.4	0.4	0.5	0.7	0.9	1.5
2002.....	0.1	0.2	0.2	0.3	0.3	0.4	0.6	0.8	1.3	1.5
2001.....	0.2	0.2	0.2	0.3	0.3	0.4	0.8	0.6	0.9	1.7
2000.....	0.2	0.2	0.2	0.3	0.4	0.3	0.6	0.7	1.2	2.1
1999.....	0.1	0.3	0.3	0.3	0.3	0.6	0.5	0.7	1.2	1.7

All races 3/, male

2006.....	*	*	*	*	*	*	*	*	*	1.7
2005.....	*	*	*	*	*	*	*	0.7	*	2.0
2004.....	0.2	0.3	0.3	0.2	*	*	*	1.1	1.0	1.9
2003.....	0.2	*	0.3	*	0.5	*	0.6	0.8	1.3	1.7
2002.....	*	0.3	0.2	0.3	0.4	0.5	0.8	0.7	1.3	1.8
2001.....	0.2	0.3	*	0.4	*	*	0.8	0.9	1.2	2.1
2000.....	0.2	0.3	0.4	*	0.5	*	0.8	0.8	1.4	3.0
1999.....	*	0.3	0.4	0.4	*	0.6	*	0.9	1.4	2.2

All races 3/, female

2006.....	*	*	*	*	*	*	*	*	*	1.3
2005.....	*	*	*	*	*	*	*	0.5	0.9	1.3
2004.....	*	0.2	*	0.3	*	*	*	0.7	1.0	1.5
2003.....	0.2	*	0.2	*	*	*	0.5	0.5	0.7	1.4
2002.....	*	0.2	0.2	0.3	*	0.4	0.4	0.9	1.3	1.4
2001.....	*	*	*	*	*	0.4	0.7	*	0.7	1.6
2000.....	*	0.2	*	0.3	0.4	*	0.4	0.7	1.1	1.7
1999.....	*	0.3	0.2	*	0.4	0.6	0.5	0.5	1.1	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other congenital malformations of circulatory system (Q25-Q28)

White, both sexes

2006 4/.....	*	0.1	*	*	0.2	*	0.4	0.5	0.6	1.4
2005 4/.....	*	0.1	*	*	*	*	0.4	0.6	0.9	1.5
2004 4/.....	0.2	0.3	0.2	0.2	0.2	0.3	0.5	0.8	1.1	1.7
2003 4/.....	0.2	0.2	0.2	0.2	0.4	0.4	0.5	0.7	0.9	1.5
2002.....	*	0.2	0.2	0.2	0.3	0.5	0.6	0.8	1.3	1.6
2001.....	0.1	0.2	0.2	0.3	0.3	0.4	0.7	0.7	1.0	1.7
2000.....	0.2	0.2	0.2	0.2	0.4	0.3	0.6	0.8	1.2	2.2
1999.....	0.1	0.3	0.3	0.3	0.3	0.6	0.4	0.6	1.2	1.7

White, male

2006 4/.....	*	*	*	*	*	*	*	*	*	1.7
2005 4/.....	*	*	*	*	*	*	*	0.7	*	2.1
2004 4/.....	*	0.3	0.3	*	*	*	*	1.0	1.0	2.0
2003 4/.....	0.2	*	0.3	*	0.5	*	*	0.8	1.2	1.7
2002.....	*	0.3	*	*	*	*	0.7	*	1.2	1.9
2001.....	*	0.2	*	0.4	*	*	0.8	1.0	1.3	2.0
2000.....	*	0.3	0.3	*	0.5	*	0.8	0.8	1.5	3.2
1999.....	*	0.3	0.4	0.4	*	0.5	*	0.8	1.4	2.2

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	1.3
2005 4/.....	*	*	*	*	*	*	*	*	0.8	1.2
2004 4/.....	*	*	*	*	*	*	*	0.7	1.1	1.6
2003 4/.....	*	*	*	*	*	*	0.5	0.5	0.7	1.4
2002.....	*	*	*	*	*	0.5	0.5	0.9	1.3	1.5
2001.....	*	*	*	*	*	*	0.7	*	0.8	1.6
2000.....	*	*	*	*	*	*	*	0.7	1.1	1.7
1999.....	*	0.3	*	*	*	0.6	0.5	*	1.1	1.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other congenital malformations and deformations (Q10-Q18,Q35-Q89)

All races 3/, both sexes

2006.....	0.2	0.4	0.3	0.5	0.6	0.5	0.8	1.1	1.8	2.6
2005.....	0.2	0.3	0.4	0.4	0.6	0.6	0.8	1.0	1.9	3.0
2004.....	0.2	0.3	0.3	0.4	0.6	0.8	0.9	1.2	1.7	2.6
2003.....	0.2	0.3	0.3	0.5	0.5	0.6	0.8	1.3	1.7	2.8
2002.....	0.3	0.4	0.4	0.5	0.6	0.6	1.0	1.1	1.7	2.7
2001.....	0.3	0.3	0.3	0.4	0.6	0.7	0.9	1.3	1.8	2.5
2000.....	0.3	0.3	0.3	0.5	0.6	0.8	0.8	1.3	1.6	2.8
1999.....	0.2	0.3	0.3	0.4	0.6	0.7	0.9	1.3	1.8	2.6

All races 3/, male

2006.....	0.2	0.4	0.4	0.6	0.7	0.5	0.9	1.5	2.3	3.6
2005.....	0.3	0.3	0.4	0.4	0.8	0.7	0.8	1.2	2.5	3.2
2004.....	0.3	0.4	0.4	0.4	0.6	0.9	0.9	1.2	1.6	3.2
2003.....	0.2	0.4	0.4	0.5	0.4	0.7	0.9	1.3	1.6	3.3
2002.....	0.4	0.5	0.5	0.4	0.7	0.7	1.2	1.4	1.8	3.6
2001.....	0.4	0.3	0.3	0.4	0.6	0.8	1.0	1.4	1.8	3.7
2000.....	0.3	0.4	0.4	0.4	0.6	0.9	0.8	1.1	1.8	4.6
1999.....	0.3	0.4	0.4	0.5	0.5	0.7	1.1	1.4	2.1	3.0

All races 3/, female

2006.....	0.2	0.3	0.3	0.4	0.6	0.6	0.7	0.8	1.4	2.2
2005.....	*	0.2	0.4	0.4	0.5	0.5	0.8	0.8	1.5	2.9
2004.....	0.2	0.2	0.3	0.3	0.6	0.7	0.9	1.1	1.8	2.4
2003.....	0.2	0.3	0.2	0.5	0.5	0.6	0.7	1.4	1.8	2.6
2002.....	0.2	0.2	0.3	0.5	0.5	0.5	0.8	0.9	1.7	2.3
2001.....	0.2	0.3	0.2	0.4	0.6	0.6	0.8	1.2	1.8	2.0
2000.....	0.2	0.2	*	0.6	0.6	0.8	0.9	1.3	1.5	2.1
1999.....	0.2	0.3	0.2	0.4	0.6	0.8	0.7	1.2	1.7	2.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other congenital malformations and deformations (Q10-Q18,Q35-Q89)

White, both sexes

2006 4/.....	0.2	0.3	0.4	0.6	0.7	0.6	0.8	1.2	1.8	2.6
2005 4/.....	0.2	0.3	0.4	0.4	0.6	0.6	0.9	1.0	2.0	3.1
2004 4/.....	0.3	0.3	0.3	0.3	0.6	0.8	1.0	1.1	1.7	2.7
2003 4/.....	0.2	0.3	0.3	0.5	0.5	0.6	0.8	1.4	1.7	2.8
2002.....	0.3	0.4	0.4	0.5	0.6	0.6	1.0	1.1	1.7	2.7
2001.....	0.3	0.3	0.3	0.4	0.6	0.7	0.9	1.3	1.7	2.6
2000.....	0.3	0.3	0.3	0.5	0.6	0.8	0.9	1.3	1.7	2.8
1999.....	0.2	0.3	0.3	0.5	0.6	0.8	0.9	1.2	1.8	2.7

White, male

2006 4/.....	*	0.4	0.5	0.6	0.7	0.6	0.9	1.6	2.5	3.5
2005 4/.....	0.3	0.3	0.4	0.4	0.8	0.7	0.9	1.2	2.6	3.2
2004 4/.....	0.3	0.4	0.3	0.3	0.6	0.9	1.0	1.2	1.5	3.3
2003 4/.....	0.3	0.3	0.4	0.5	0.4	0.7	0.9	1.3	1.4	3.4
2002.....	0.4	0.5	0.5	0.4	0.7	0.7	1.2	1.5	1.6	3.7
2001.....	0.4	0.3	0.3	0.4	0.6	0.8	1.0	1.5	1.4	3.8
2000.....	0.3	0.4	0.4	0.4	0.7	0.8	0.8	1.2	1.9	4.6
1999.....	0.3	0.3	0.4	0.5	0.5	0.8	1.2	1.2	2.2	3.2

White, female

2006 4/.....	0.2	0.3	0.3	0.5	0.6	0.6	0.7	0.9	1.4	2.3
2005 4/.....	*	0.2	0.4	0.4	0.5	0.5	0.9	0.8	1.6	3.0
2004 4/.....	0.2	0.2	0.3	0.3	0.6	0.8	0.9	1.1	1.8	2.4
2003 4/.....	*	0.3	*	0.5	0.6	0.6	0.6	1.4	1.9	2.6
2002.....	*	0.2	0.4	0.5	0.5	0.5	0.8	0.9	1.7	2.3
2001.....	0.3	0.3	*	0.4	0.7	0.6	0.8	1.2	1.8	2.1
2000.....	0.2	0.3	*	0.6	0.6	0.8	0.9	1.3	1.6	2.1
1999.....	*	0.3	0.3	0.4	0.6	0.8	0.8	1.2	1.6	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Down's syndrome (Q90)										
All races 3/, both sexes										
2006.....	0.2	0.4	0.5	1.0	1.0	0.5	*	*	*	*
2005.....	0.3	0.4	0.7	1.3	1.1	0.5	*	*	*	*
2004.....	0.2	0.4	0.7	1.0	0.8	0.6	0.3	*	*	*
2003.....	0.3	0.4	0.6	1.1	0.9	0.4	*	*	*	*
2002.....	0.2	0.3	0.7	0.9	1.0	0.6	*	*	*	*
2001.....	0.3	0.3	0.6	0.9	1.0	0.4	*	*	*	*
2000.....	0.2	0.3	0.7	0.8	0.8	0.6	*	*	*	*
1999.....	0.2	0.4	0.5	0.8	0.7	0.3	*	*	*	*
All races 3/, male										
2006.....	*	0.4	0.6	1.1	1.1	0.4	*	*	*	*
2005.....	0.2	0.4	0.7	1.3	1.0	*	*	*	*	*
2004.....	0.2	0.5	0.6	1.0	0.9	0.6	*	*	*	*
2003.....	0.3	0.4	0.6	1.1	1.1	0.4	*	*	*	*
2002.....	0.2	0.4	0.7	0.9	1.1	0.6	*	*	*	*
2001.....	0.3	0.3	0.6	0.8	1.0	0.5	*	*	*	*
2000.....	*	0.2	0.8	0.8	0.7	0.5	*	*	*	*
1999.....	*	0.4	0.4	0.8	0.8	*	*	*	*	*
All races 3/, female										
2006.....	0.2	0.3	0.5	0.9	0.8	0.6	*	*	*	*
2005.....	0.3	0.4	0.6	1.2	1.1	0.5	*	*	*	*
2004.....	0.2	0.4	0.7	0.9	0.8	0.5	*	*	*	*
2003.....	0.2	0.4	0.7	1.1	0.8	0.4	*	*	*	*
2002.....	0.2	0.3	0.6	0.9	0.9	0.6	*	*	*	*
2001.....	0.3	0.3	0.7	1.0	1.0	*	*	*	*	*
2000.....	*	0.3	0.6	0.7	0.9	0.7	*	*	*	*
1999.....	*	0.3	0.6	0.8	0.6	0.4	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Down's syndrome (Q90)										
White, both sexes										
2006 4/.....	0.2	0.4	0.6	1.0	1.1	0.6	*	*	*	*
2005 4/.....	0.3	0.4	0.7	1.4	1.2	0.5	*	*	*	*
2004 4/.....	0.2	0.5	0.7	1.1	0.9	0.6	0.3	*	*	*
2003 4/.....	0.3	0.5	0.7	1.2	1.0	0.5	*	*	*	*
2002.....	0.2	0.3	0.7	1.0	1.1	0.6	*	*	*	*
2001.....	0.3	0.3	0.7	1.0	1.1	0.5	*	*	*	*
2000.....	0.2	0.3	0.8	0.8	0.9	0.7	*	*	*	*
1999.....	0.2	0.4	0.5	0.9	0.8	0.3	*	*	*	*
White, male										
2006 4/.....	*	0.4	0.7	1.2	1.3	0.5	*	*	*	*
2005 4/.....	0.2	0.4	0.8	1.5	1.1	*	*	*	*	*
2004 4/.....	0.2	0.5	0.7	1.1	0.9	0.6	*	*	*	*
2003 4/.....	0.3	0.5	0.7	1.1	1.2	*	*	*	*	*
2002.....	0.3	0.4	0.7	0.9	1.2	0.6	*	*	*	*
2001.....	0.3	0.3	0.7	0.8	1.1	0.6	*	*	*	*
2000.....	*	0.2	0.9	0.9	0.8	*	*	*	*	*
1999.....	*	0.4	0.5	0.9	0.9	*	*	*	*	*
White, female										
2006 4/.....	*	0.4	0.5	0.9	0.9	0.6	*	*	*	*
2005 4/.....	0.4	0.4	0.7	1.4	1.3	0.6	*	*	*	*
2004 4/.....	*	0.4	0.8	1.1	0.9	0.6	*	*	*	*
2003 4/.....	0.2	0.4	0.8	1.2	0.9	0.5	*	*	*	*
2002.....	0.2	0.3	0.7	1.0	1.0	0.6	*	*	*	*
2001.....	0.3	0.3	0.7	1.2	1.1	*	*	*	*	*
2000.....	*	0.4	0.7	0.8	1.0	0.8	*	*	*	*
1999.....	*	0.3	0.6	0.9	0.7	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2006.....	4.2	4.9	5.6	5.9	7.2	9.5	15.5	28.7	58.6	236.1
2005.....	5.2	6.4	7.0	6.9	7.7	10.4	15.7	27.6	54.0	224.4
2004.....	4.9	5.7	5.8	6.3	7.9	10.2	15.4	28.0	55.5	221.5
2003.....	5.9	7.3	6.8	6.8	8.2	10.6	17.3	26.7	55.7	224.1
2002.....	6.0	6.4	6.9	7.0	8.3	10.4	16.1	28.3	53.3	208.0
2001.....	7.7	9.1	8.8	8.4	9.9	12.7	17.8	29.8	54.3	206.4
2000.....	7.6	8.6	8.0	8.6	10.2	14.2	19.0	30.1	56.5	208.1
1999.....	5.1	5.9	5.8	6.9	8.9	11.9	17.6	26.5	50.5	187.7

All races 3/, male

2006.....	5.5	6.3	7.2	7.7	9.0	11.9	18.5	32.3	61.3	191.1
2005.....	6.4	8.2	9.3	9.2	9.6	12.8	18.8	31.0	54.8	183.9
2004.....	6.5	7.4	7.7	8.3	10.1	12.6	17.9	30.6	58.1	183.6
2003.....	7.4	9.4	8.9	8.9	10.6	13.7	21.3	31.2	58.0	177.9
2002.....	7.6	8.5	9.0	9.0	11.2	12.7	19.7	32.9	60.6	175.5
2001.....	9.9	11.9	11.8	11.0	13.2	15.5	20.9	33.2	60.6	175.0
2000.....	10.2	11.7	11.2	11.4	13.4	18.9	22.4	36.1	62.3	182.7
1999.....	6.7	8.2	7.9	9.1	11.9	15.1	21.8	31.5	54.4	166.3

All races 3/, female

2006.....	3.0	3.5	4.1	4.2	5.6	7.3	13.1	26.0	56.9	257.2
2005.....	4.1	4.6	4.8	4.7	6.0	8.3	13.2	25.1	53.5	243.0
2004.....	3.3	4.1	4.0	4.5	5.9	8.1	13.4	26.1	53.8	238.5
2003.....	4.4	5.1	4.8	4.9	6.0	7.9	14.2	23.5	54.3	244.6
2002.....	4.5	4.4	5.0	5.0	5.6	8.4	13.3	25.1	48.9	222.1
2001.....	5.5	6.3	5.9	5.9	6.9	10.2	15.4	27.4	50.4	219.5
2000.....	5.0	5.6	5.0	5.9	7.3	10.1	16.4	26.0	53.0	218.5
1999.....	3.4	3.7	3.9	4.9	6.2	9.2	14.2	23.0	48.3	196.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes

2006 4/.....	4.0	4.7	5.2	5.6	7.0	9.2	15.0	28.3	58.3	241.1
2005 4/.....	5.0	5.9	6.6	6.3	7.3	9.7	15.0	27.2	53.7	228.9
2004 4/.....	4.7	5.6	5.4	5.8	7.4	9.8	14.7	27.6	55.1	224.4
2003 4/.....	5.6	6.6	6.2	6.3	7.6	10.1	16.6	26.5	56.1	228.3
2002.....	5.7	5.8	6.1	6.4	7.9	9.8	15.9	27.9	53.3	211.6
2001.....	7.6	8.8	8.3	7.9	9.4	11.8	17.2	29.5	53.9	210.0
2000.....	7.4	8.2	7.3	7.9	9.5	13.0	18.5	29.5	56.0	212.2
1999.....	4.4	5.2	5.0	6.2	8.2	11.1	17.0	25.5	49.5	190.2

White, male

2006 4/.....	5.1	5.9	6.6	7.3	8.8	11.5	17.2	31.5	60.5	193.8
2005 4/.....	6.3	7.8	8.7	8.3	8.9	11.8	17.9	30.4	53.6	185.6
2004 4/.....	6.3	7.2	6.9	7.5	9.3	11.9	17.4	30.1	57.8	186.0
2003 4/.....	7.0	8.6	8.2	8.1	9.7	12.8	20.4	31.0	57.8	180.0
2002.....	7.2	7.6	7.7	8.1	10.7	11.9	19.2	31.6	60.0	178.2
2001.....	10.0	11.6	11.2	10.1	12.2	14.3	20.3	33.2	60.6	177.0
2000.....	10.1	11.2	10.1	10.2	12.4	17.1	21.5	35.4	62.0	186.8
1999.....	5.9	7.0	6.7	8.0	10.9	13.9	21.0	30.3	52.6	167.3

White, female

2006 4/.....	2.8	3.5	3.8	3.9	5.3	7.1	13.2	25.8	56.9	263.3
2005 4/.....	3.8	4.1	4.5	4.3	5.8	7.9	12.6	24.9	53.7	248.8
2004 4/.....	3.1	4.0	3.8	4.2	5.6	7.9	12.5	25.8	53.5	241.7
2003 4/.....	4.2	4.6	4.2	4.7	5.6	7.7	13.5	23.3	55.0	249.7
2002.....	4.2	4.0	4.4	4.7	5.3	7.9	13.1	25.3	49.2	226.1
2001.....	5.3	6.0	5.5	5.7	6.7	9.7	14.6	26.8	49.9	223.8
2000.....	4.6	5.2	4.6	5.6	6.7	9.4	16.0	25.4	52.4	222.5
1999.....	2.9	3.4	3.4	4.5	5.7	8.6	13.8	22.1	47.6	199.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XVIII. Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2006 4/.....	6.9	7.6	10.2	10.3	11.1	14.9	24.4	41.0	75.7	223.9
2005 4/.....	8.0	11.2	12.4	13.9	14.2	19.2	26.2	38.8	70.9	215.1
2004 4/.....	7.6	8.6	11.0	12.5	14.6	16.1	25.9	39.9	72.0	220.8
2003 4/.....	9.7	13.4	13.4	13.0	15.7	17.9	30.3	34.9	60.9	210.9
2002.....	9.5	12.1	15.7	13.8	13.6	17.8	23.1	39.5	64.9	196.0
2001.....	9.8	13.2	14.6	14.3	15.4	21.7	27.9	38.6	66.6	191.6
2000.....	11.2	13.8	15.9	16.0	19.0	28.3	29.7	44.2	73.4	186.4
1999.....	10.3	13.1	13.3	14.9	17.0	22.3	27.2	42.7	72.5	182.4

Black, male

2006 4/.....	9.0	10.8	14.2	14.0	12.9	20.6	36.3	52.8	92.4	193.9
2005 4/.....	9.2	13.8	17.2	19.5	20.0	26.4	33.5	49.2	85.7	199.9
2004 4/.....	9.7	11.8	16.3	18.2	20.2	22.3	28.2	45.0	81.1	181.8
2003 4/.....	12.5	17.5	17.8	18.5	22.8	25.9	38.8	41.5	70.1	193.9
2002.....	12.4	17.1	22.1	19.5	20.0	22.9	31.0	57.0	87.1	181.2
2001.....	11.7	17.3	18.9	21.7	25.4	29.5	31.6	40.9	70.6	184.9
2000.....	13.5	18.2	23.6	25.0	25.5	41.4	38.1	54.8	83.5	161.5
1999.....	13.6	19.7	19.1	22.4	24.0	30.1	35.6	54.1	88.1	184.0

Black, female

2006 4/.....	5.0	4.8	6.9	7.3	9.6	10.7	16.3	34.0	66.9	236.2
2005 4/.....	6.9	8.9	8.4	9.3	9.6	13.9	21.2	32.5	63.1	221.3
2004 4/.....	5.9	5.9	6.6	7.7	10.2	11.5	24.4	36.9	67.3	236.3
2003 4/.....	7.3	9.9	9.7	8.4	10.2	12.1	24.6	30.9	56.1	217.5
2002.....	7.0	7.7	10.2	9.0	8.7	14.1	17.7	29.1	53.3	201.7
2001.....	8.1	9.7	11.0	8.2	7.5	16.0	25.3	37.2	64.6	194.1
2000.....	9.1	9.9	9.4	8.6	13.9	18.6	24.2	37.9	68.4	195.7
1999.....	7.3	7.3	8.5	8.9	11.5	16.5	21.6	35.8	64.8	181.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Senility (R54)										
All races 3/, both sexes										
2006.....	*	*	*	*	0.1	0.3	1.3	3.3	10.6	74.6
2005.....	*	*	*	*	*	0.4	1.2	3.6	9.9	73.2
2004.....	*	*	*	*	0.2	0.4	1.2	4.1	11.2	76.3
2003.....	*	*	*	*	0.2	0.4	1.4	3.5	11.2	76.7
2002.....	*	*	*	*	*	0.4	1.2	3.7	9.8	72.0
2001.....	*	*	*	*	*	0.4	1.3	3.3	10.8	73.0
2000.....	*	*	*	*	*	0.4	1.3	3.6	10.3	67.2
1999.....	*	*	*	*	*	0.3	1.1	2.9	9.0	56.5
All races 3/, male										
2006.....	*	*	*	*	*	*	1.4	3.5	11.2	58.9
2005.....	*	*	*	*	*	*	1.4	3.6	9.3	56.7
2004.....	*	*	*	*	*	0.6	1.3	4.3	11.6	57.2
2003.....	*	*	*	*	*	0.5	1.6	3.7	10.9	55.6
2002.....	*	*	*	*	*	0.5	1.2	3.6	10.3	56.5
2001.....	*	*	*	*	*	0.5	1.5	3.2	11.4	57.2
2000.....	*	*	*	*	*	*	1.3	3.8	10.5	51.7
1999.....	*	*	*	*	*	*	1.5	3.1	8.9	45.8
All races 3/, female										
2006.....	*	*	*	*	*	0.4	1.3	3.1	10.2	81.9
2005.....	*	*	*	*	*	0.4	1.0	3.6	10.3	80.8
2004.....	*	*	*	*	*	*	1.1	4.0	10.9	84.9
2003.....	*	*	*	*	*	*	1.3	3.4	11.4	86.0
2002.....	*	*	*	*	*	*	1.2	3.8	9.5	78.8
2001.....	*	*	*	*	*	*	1.2	3.3	10.4	79.6
2000.....	*	*	*	*	*	*	1.2	3.4	10.2	73.6
1999.....	*	*	*	*	*	*	0.8	2.8	9.0	60.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Senility (R54)										
White, both sexes										
2006 4/.....	*	*	*	*	*	0.4	1.4	3.5	10.8	76.5
2005 4/.....	*	*	*	*	*	0.4	1.2	3.5	10.1	75.1
2004 4/.....	*	*	*	*	*	0.4	1.2	4.2	11.2	79.0
2003 4/.....	*	*	*	*	*	0.4	1.5	3.7	11.8	79.3
2002.....	*	*	*	*	*	0.4	1.2	3.8	10.1	74.2
2001.....	*	*	*	*	*	0.3	1.4	3.3	11.0	75.1
2000.....	*	*	*	*	*	0.3	1.2	3.5	10.2	69.0
1999.....	*	*	*	*	*	0.4	1.1	2.8	8.8	58.8
White, male										
2006 4/.....	*	*	*	*	*	*	1.5	3.7	11.5	60.8
2005 4/.....	*	*	*	*	*	*	1.5	3.5	9.3	58.5
2004 4/.....	*	*	*	*	*	0.6	1.3	4.4	11.4	59.9
2003 4/.....	*	*	*	*	*	*	1.7	3.8	11.2	57.6
2002.....	*	*	*	*	*	*	1.1	3.6	10.9	58.0
2001.....	*	*	*	*	*	*	1.6	3.3	11.7	58.3
2000.....	*	*	*	*	*	*	1.4	3.7	10.5	53.9
1999.....	*	*	*	*	*	*	1.5	2.9	8.6	47.4
White, female										
2006 4/.....	*	*	*	*	*	0.4	1.4	3.3	10.4	83.9
2005 4/.....	*	*	*	*	*	*	1.0	3.5	10.6	82.7
2004 4/.....	*	*	*	*	*	*	1.2	4.1	11.1	87.6
2003 4/.....	*	*	*	*	*	*	1.4	3.6	12.1	88.9
2002.....	*	*	*	*	*	*	1.3	3.9	9.7	81.2
2001.....	*	*	*	*	*	*	1.3	3.3	10.6	82.2
2000.....	*	*	*	*	*	*	1.1	3.4	10.0	75.1
1999.....	*	*	*	*	*	*	0.8	2.8	9.0	63.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Senility (R54)										
Black, both sexes										
2006 4/.....	*	*	*	*	*	*	*	*	10.6	65.9
2005 4/.....	*	*	*	*	*	*	*	5.3	10.7	64.9
2004 4/.....	*	*	*	*	*	*	*	4.6	12.9	56.1
2003 4/.....	*	*	*	*	*	*	*	3.4	6.3	55.7
2002.....	*	*	*	*	*	*	*	3.6	7.6	55.5
2001.....	*	*	*	*	*	*	*	4.0	10.8	56.8
2000.....	*	*	*	*	*	*	*	5.0	14.1	55.7
1999.....	*	*	*	*	*	*	*	4.5	13.3	39.9
Black, male										
2006 4/.....	*	*	*	*	*	*	*	*	*	47.4
2005 4/.....	*	*	*	*	*	*	*	*	*	49.3
2004 4/.....	*	*	*	*	*	*	*	*	17.9	32.8
2003 4/.....	*	*	*	*	*	*	*	*	*	37.5
2002.....	*	*	*	*	*	*	*	*	*	49.3
2001.....	*	*	*	*	*	*	*	*	*	57.9
2000.....	*	*	*	*	*	*	*	*	*	34.6
1999.....	*	*	*	*	*	*	*	*	*	36.3
Black, female										
2006 4/.....	*	*	*	*	*	*	*	*	10.1	73.5
2005 4/.....	*	*	*	*	*	*	*	*	10.0	71.3
2004 4/.....	*	*	*	*	*	*	*	*	10.2	65.4
2003 4/.....	*	*	*	*	*	*	*	*	*	62.8
2002.....	*	*	*	*	*	*	*	*	8.8	57.9
2001.....	*	*	*	*	*	*	*	*	10.6	56.4
2000.....	*	*	*	*	*	*	*	*	14.4	63.5
1999.....	*	*	*	*	*	*	*	*	11.7	41.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ill-defined and unknown causes of mortality (R95-R99)										
All races 3/, both sexes										
2006.....	3.2	3.8	4.1	4.0	4.1	4.7	6.1	9.2	16.2	46.4
2005.....	4.4	5.4	5.7	5.3	5.3	6.3	7.5	11.3	18.3	53.9
2004.....	4.1	4.6	4.5	4.7	5.1	6.1	7.6	11.5	18.1	52.4
2003.....	4.9	5.9	5.5	5.3	5.6	6.7	9.1	11.8	19.7	59.5
2002.....	5.0	5.3	5.5	5.5	5.8	6.6	8.7	13.4	20.8	58.3
2001.....	6.6	8.0	7.6	6.8	7.6	8.8	10.4	14.9	22.9	62.1
2000.....	6.5	7.4	6.8	6.9	7.6	10.0	11.9	16.2	26.9	73.0
1999.....	4.2	4.9	4.6	5.2	6.3	7.9	10.5	13.5	22.9	66.7
All races 3/, male										
2006.....	4.2	5.0	5.4	5.4	5.5	6.4	8.1	12.0	18.7	41.9
2005.....	5.4	7.0	7.5	7.2	7.0	8.1	9.9	14.0	21.3	48.4
2004.....	5.5	6.2	6.0	6.3	7.0	7.7	9.8	14.2	20.3	50.7
2003.....	6.3	7.8	7.3	7.1	7.6	9.4	12.2	15.0	23.4	53.6
2002.....	6.5	7.2	7.4	7.3	8.1	8.4	11.4	16.4	25.5	55.0
2001.....	8.6	10.5	10.2	9.1	10.2	11.2	13.2	18.1	27.6	58.1
2000.....	8.8	10.3	9.6	9.3	10.4	13.8	14.3	20.2	31.2	72.1
1999.....	5.7	6.9	6.3	6.9	8.8	10.5	13.8	17.0	26.4	64.4
All races 3/, female										
2006.....	2.2	2.6	2.9	2.6	2.8	3.1	4.5	7.1	14.7	48.6
2005.....	3.3	3.8	3.9	3.4	3.9	4.7	5.6	9.4	16.5	56.4
2004.....	2.6	3.1	3.0	3.1	3.5	4.6	5.8	9.6	16.8	53.2
2003.....	3.5	4.1	3.8	3.6	3.7	4.3	6.5	9.4	17.4	62.1
2002.....	3.6	3.4	3.6	3.8	3.8	5.0	6.6	11.3	17.9	59.8
2001.....	4.7	5.5	5.0	4.8	5.2	6.8	8.1	12.6	20.0	63.8
2000.....	4.3	4.7	4.1	4.7	5.1	6.8	10.1	13.4	24.3	73.3
1999.....	2.8	3.1	3.0	3.6	4.0	5.7	7.9	11.2	20.9	67.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Ill-defined and unknown causes of mortality (R95-R99)

White, both sexes

2006 4/.....	3.2	3.7	4.0	3.9	4.2	4.7	6.2	9.2	16.7	48.2
2005 4/.....	4.3	5.1	5.4	4.9	5.1	6.0	7.3	11.1	18.3	55.3
2004 4/.....	4.0	4.5	4.2	4.4	5.0	5.9	7.2	11.6	18.1	53.1
2003 4/.....	4.7	5.6	5.2	5.0	5.2	6.6	8.8	11.9	19.9	60.9
2002.....	4.9	4.8	4.8	5.1	5.6	6.3	8.7	13.3	20.8	59.3
2001.....	6.8	7.8	7.3	6.6	7.4	8.6	10.1	14.8	22.6	63.2
2000.....	6.4	7.2	6.3	6.5	7.3	9.3	11.9	16.0	27.1	74.5
1999.....	3.7	4.4	4.1	4.7	5.9	7.4	10.3	13.2	22.4	67.0

White, male

2006 4/.....	4.2	4.8	5.2	5.2	5.6	6.5	7.8	11.9	19.1	43.9
2005 4/.....	5.4	6.8	7.2	6.6	6.4	7.7	9.6	13.5	21.3	49.2
2004 4/.....	5.5	6.0	5.5	5.9	6.6	7.5	9.6	14.3	20.2	51.4
2003 4/.....	6.1	7.4	6.9	6.5	7.0	9.0	11.7	15.0	23.6	54.7
2002.....	6.3	6.5	6.4	6.7	7.8	7.9	11.3	16.0	24.9	56.2
2001.....	8.9	10.3	9.9	8.5	9.9	10.8	13.0	18.1	27.2	59.3
2000.....	8.8	10.0	8.8	8.7	9.9	12.5	13.9	19.9	31.4	73.2
1999.....	5.1	6.0	5.4	6.1	8.2	9.7	13.4	16.5	25.3	63.1

White, female

2006 4/.....	2.2	2.6	2.8	2.6	2.8	3.2	4.8	7.1	15.2	50.2
2005 4/.....	3.1	3.5	3.6	3.3	3.8	4.5	5.4	9.4	16.5	58.2
2004 4/.....	2.4	3.1	3.0	3.0	3.5	4.5	5.2	9.6	16.8	53.8
2003 4/.....	3.4	3.8	3.5	3.6	3.6	4.5	6.4	9.6	17.6	63.7
2002.....	3.5	3.2	3.3	3.6	3.7	4.8	6.5	11.4	18.3	60.6
2001.....	4.6	5.3	4.7	4.7	5.1	6.7	7.9	12.4	19.8	64.9
2000.....	4.0	4.4	3.8	4.4	4.9	6.5	10.2	13.2	24.5	75.0
1999.....	2.4	2.8	2.7	3.3	3.7	5.4	7.8	10.9	20.7	68.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Ill-defined and unknown causes of mortality (R95-R99)										
Black, both sexes										
2006 4/.....	4.0	5.4	6.6	5.8	4.3	5.8	6.9	11.4	14.9	34.2
2005 4/.....	6.4	8.7	9.7	10.1	9.2	10.5	12.1	16.9	23.1	50.5
2004 4/.....	5.6	6.7	7.8	8.2	7.9	8.9	12.7	13.9	21.3	55.0
2003 4/.....	7.8	10.0	9.6	9.2	10.1	9.2	15.0	13.4	21.3	54.6
2002.....	7.4	9.5	11.9	9.9	9.1	10.9	11.0	17.9	24.3	56.1
2001.....	7.4	10.9	11.2	10.4	10.2	11.9	14.0	18.6	28.2	55.6
2000.....	9.2	11.0	12.6	11.6	12.6	18.6	15.2	21.5	30.5	66.0
1999.....	8.2	10.2	10.1	11.0	11.5	13.7	13.8	19.3	32.4	71.4
Black, male										
2006 4/.....	5.4	8.3	9.1	8.7	5.8	9.1	13.2	17.0	19.2	26.8
2005 4/.....	7.6	10.7	13.1	15.4	14.5	14.5	17.1	23.7	29.5	54.8
2004 4/.....	6.8	9.4	11.9	12.0	12.1	11.4	14.6	16.4	26.5	52.7
2003 4/.....	10.1	13.1	12.6	14.0	16.2	15.8	23.0	19.6	25.8	59.4
2002.....	9.9	13.8	17.3	14.6	14.1	15.5	16.0	25.7	40.5	53.6
2001.....	8.4	14.0	14.4	15.8	16.1	17.0	18.5	22.8	35.3	49.9
2000.....	11.2	14.8	18.7	17.0	18.2	28.6	21.4	29.3	38.8	70.3
1999.....	10.8	15.4	15.0	16.4	17.4	20.0	20.4	25.6	41.9	89.1
Black, female										
2006 4/.....	2.8	2.9	4.4	3.4	3.2	*	*	8.0	12.7	37.3
2005 4/.....	5.3	6.9	6.7	5.8	5.1	7.6	8.7	12.8	19.7	48.8
2004 4/.....	4.5	4.3	4.3	5.1	4.7	7.0	11.4	12.4	18.5	55.9
2003 4/.....	5.8	7.3	7.0	5.3	5.4	4.4	9.5	9.8	19.0	52.7
2002.....	5.1	5.7	7.2	6.1	5.3	7.6	7.6	13.2	15.9	57.1
2001.....	6.6	8.3	8.5	5.9	5.6	8.1	11.0	16.1	24.5	57.7
2000.....	7.4	7.8	7.4	7.2	8.2	11.1	11.1	16.8	26.3	64.4
1999.....	6.0	5.7	6.0	6.5	6.8	9.0	9.5	15.4	27.7	64.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other ill-defined and unknown causes of mortality (R96-R99)

All races 3/, both sexes

2006.....	3.2	3.8	4.1	4.0	4.1	4.7	6.1	9.2	16.2	46.4
2005.....	4.4	5.4	5.7	5.3	5.3	6.3	7.5	11.3	18.3	53.9
2004.....	4.1	4.6	4.5	4.7	5.1	6.1	7.6	11.5	18.1	52.4
2003.....	4.9	5.9	5.5	5.3	5.6	6.7	9.1	11.8	19.7	59.5
2002.....	5.0	5.3	5.5	5.5	5.8	6.6	8.7	13.4	20.8	58.3
2001.....	6.6	8.0	7.6	6.8	7.6	8.8	10.4	14.9	22.9	62.1
2000.....	6.5	7.4	6.8	6.9	7.6	10.0	11.9	16.2	26.9	73.0
1999.....	4.2	4.9	4.6	5.2	6.3	7.9	10.5	13.5	22.9	66.7

All races 3/, male

2006.....	4.2	5.0	5.4	5.4	5.5	6.4	8.1	12.0	18.7	41.9
2005.....	5.4	7.0	7.5	7.2	7.0	8.1	9.9	14.0	21.3	48.4
2004.....	5.5	6.2	6.0	6.3	7.0	7.7	9.8	14.2	20.3	50.7
2003.....	6.3	7.8	7.3	7.1	7.6	9.4	12.2	15.0	23.4	53.6
2002.....	6.5	7.2	7.4	7.3	8.1	8.4	11.4	16.4	25.5	55.0
2001.....	8.6	10.5	10.2	9.1	10.2	11.2	13.2	18.1	27.6	58.1
2000.....	8.8	10.3	9.6	9.3	10.4	13.8	14.3	20.2	31.2	72.1
1999.....	5.7	6.9	6.3	6.9	8.8	10.5	13.8	17.0	26.4	64.4

All races 3/, female

2006.....	2.2	2.6	2.9	2.6	2.8	3.1	4.5	7.1	14.7	48.6
2005.....	3.3	3.8	3.9	3.4	3.9	4.7	5.6	9.4	16.5	56.4
2004.....	2.6	3.1	3.0	3.1	3.5	4.6	5.8	9.6	16.8	53.2
2003.....	3.5	4.1	3.8	3.6	3.7	4.3	6.5	9.4	17.4	62.1
2002.....	3.6	3.4	3.6	3.8	3.8	5.0	6.6	11.3	17.9	59.8
2001.....	4.7	5.5	5.0	4.8	5.2	6.8	8.1	12.6	20.0	63.8
2000.....	4.3	4.7	4.1	4.7	5.1	6.8	10.1	13.4	24.3	73.3
1999.....	2.8	3.1	3.0	3.6	4.0	5.7	7.9	11.2	20.9	67.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other ill-defined and unknown causes of mortality (R96-R99)

White, both sexes

2006 4/.....	3.2	3.7	4.0	3.9	4.2	4.7	6.2	9.2	16.7	48.2
2005 4/.....	4.3	5.1	5.4	4.9	5.1	6.0	7.3	11.1	18.3	55.3
2004 4/.....	4.0	4.5	4.2	4.4	5.0	5.9	7.2	11.6	18.1	53.1
2003 4/.....	4.7	5.6	5.2	5.0	5.2	6.6	8.8	11.9	19.9	60.9
2002.....	4.9	4.8	4.8	5.1	5.6	6.3	8.7	13.3	20.8	59.3
2001.....	6.8	7.8	7.3	6.6	7.4	8.6	10.1	14.8	22.6	63.2
2000.....	6.4	7.2	6.3	6.5	7.3	9.3	11.9	16.0	27.1	74.5
1999.....	3.7	4.4	4.1	4.7	5.9	7.4	10.3	13.2	22.4	67.0

White, male

2006 4/.....	4.2	4.8	5.2	5.2	5.6	6.5	7.8	11.9	19.1	43.9
2005 4/.....	5.4	6.8	7.2	6.6	6.4	7.7	9.6	13.5	21.3	49.2
2004 4/.....	5.5	6.0	5.5	5.9	6.6	7.5	9.6	14.3	20.2	51.4
2003 4/.....	6.1	7.4	6.9	6.5	7.0	9.0	11.7	15.0	23.6	54.7
2002.....	6.3	6.5	6.4	6.7	7.8	7.9	11.3	16.0	24.9	56.2
2001.....	8.9	10.3	9.9	8.5	9.9	10.8	13.0	18.1	27.2	59.3
2000.....	8.8	10.0	8.8	8.7	9.9	12.5	13.9	19.9	31.4	73.2
1999.....	5.1	6.0	5.4	6.1	8.2	9.7	13.4	16.5	25.3	63.1

White, female

2006 4/.....	2.2	2.6	2.8	2.6	2.8	3.2	4.8	7.1	15.2	50.2
2005 4/.....	3.1	3.5	3.6	3.3	3.8	4.5	5.4	9.4	16.5	58.2
2004 4/.....	2.4	3.1	3.0	3.0	3.5	4.5	5.2	9.6	16.8	53.8
2003 4/.....	3.4	3.8	3.5	3.6	3.6	4.5	6.4	9.6	17.6	63.7
2002.....	3.5	3.2	3.3	3.6	3.7	4.8	6.5	11.4	18.3	60.6
2001.....	4.6	5.3	4.7	4.7	5.1	6.7	7.9	12.4	19.8	64.9
2000.....	4.0	4.4	3.8	4.4	4.9	6.5	10.2	13.2	24.5	75.0
1999.....	2.4	2.8	2.7	3.3	3.7	5.4	7.8	10.9	20.7	68.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other ill-defined and unknown causes of mortality (R96-R99)

Black, both sexes

2006 4/.....	4.0	5.4	6.6	5.8	4.3	5.8	6.9	11.4	14.9	34.2
2005 4/.....	6.4	8.7	9.7	10.1	9.2	10.5	12.1	16.9	23.1	50.5
2004 4/.....	5.6	6.7	7.8	8.2	7.9	8.9	12.7	13.9	21.3	55.0
2003 4/.....	7.8	10.0	9.6	9.2	10.1	9.2	15.0	13.4	21.3	54.6
2002.....	7.4	9.5	11.9	9.9	9.1	10.9	11.0	17.9	24.3	56.1
2001.....	7.4	10.9	11.2	10.4	10.2	11.9	14.0	18.6	28.2	55.6
2000.....	9.2	11.0	12.6	11.6	12.6	18.6	15.2	21.5	30.5	66.0
1999.....	8.2	10.2	10.1	11.0	11.5	13.7	13.8	19.3	32.4	71.4

Black, male

2006 4/.....	5.4	8.3	9.1	8.7	5.8	9.1	13.2	17.0	19.2	26.8
2005 4/.....	7.6	10.7	13.1	15.4	14.5	14.5	17.1	23.7	29.5	54.8
2004 4/.....	6.8	9.4	11.9	12.0	12.1	11.4	14.6	16.4	26.5	52.7
2003 4/.....	10.1	13.1	12.6	14.0	16.2	15.8	23.0	19.6	25.8	59.4
2002.....	9.9	13.8	17.3	14.6	14.1	15.5	16.0	25.7	40.5	53.6
2001.....	8.4	14.0	14.4	15.8	16.1	17.0	18.5	22.8	35.3	49.9
2000.....	11.2	14.8	18.7	17.0	18.2	28.6	21.4	29.3	38.8	70.3
1999.....	10.8	15.4	15.0	16.4	17.4	20.0	20.4	25.6	41.9	89.1

Black, female

2006 4/.....	2.8	2.9	4.4	3.4	3.2	*	*	8.0	12.7	37.3
2005 4/.....	5.3	6.9	6.7	5.8	5.1	7.6	8.7	12.8	19.7	48.8
2004 4/.....	4.5	4.3	4.3	5.1	4.7	7.0	11.4	12.4	18.5	55.9
2003 4/.....	5.8	7.3	7.0	5.3	5.4	4.4	9.5	9.8	19.0	52.7
2002.....	5.1	5.7	7.2	6.1	5.3	7.6	7.6	13.2	15.9	57.1
2001.....	6.6	8.3	8.5	5.9	5.6	8.1	11.0	16.1	24.5	57.7
2000.....	7.4	7.8	7.4	7.2	8.2	11.1	11.1	16.8	26.3	64.4
1999.....	6.0	5.7	6.0	6.5	6.8	9.0	9.5	15.4	27.7	64.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

All races 3/, both sexes

2006.....	1.0	1.1	1.5	1.9	3.0	4.4	8.1	16.2	31.7	115.1
2005.....	0.9	1.0	1.3	1.6	2.3	3.8	7.0	12.7	25.7	97.3
2004.....	0.9	1.1	1.3	1.7	2.6	3.7	6.6	12.4	26.2	92.7
2003.....	0.9	1.3	1.3	1.5	2.4	3.5	6.9	11.4	24.8	87.9
2002.....	1.0	1.1	1.5	1.4	2.3	3.4	6.2	11.2	22.8	77.7
2001.....	1.0	1.1	1.2	1.5	2.2	3.4	6.1	11.6	20.6	71.3
2000.....	1.1	1.1	1.3	1.6	2.5	3.8	5.9	10.4	19.3	67.9
1999.....	0.8	1.0	1.2	1.7	2.5	3.7	6.0	10.0	18.7	64.5

All races 3/, male

2006.....	1.3	1.2	1.8	2.3	3.3	5.2	9.0	16.8	31.4	90.3
2005.....	1.0	1.2	1.7	1.9	2.5	4.4	7.4	13.4	24.2	78.8
2004.....	1.0	1.2	1.7	1.9	2.9	4.4	6.8	12.2	26.2	75.7
2003.....	1.0	1.6	1.6	1.8	2.9	3.7	7.5	12.5	23.7	68.6
2002.....	1.1	1.3	1.6	1.7	3.0	3.8	7.1	12.9	24.8	64.0
2001.....	1.3	1.4	1.6	1.9	2.9	3.8	6.2	11.8	21.6	59.7
2000.....	1.4	1.4	1.7	2.1	2.9	4.7	6.8	12.1	20.7	58.9
1999.....	1.1	1.3	1.5	2.2	2.9	4.2	6.4	11.5	19.2	56.1

All races 3/, female

2006.....	0.8	0.9	1.2	1.5	2.6	3.8	7.3	15.9	32.0	126.8
2005.....	0.7	0.7	0.9	1.3	2.0	3.2	6.6	12.1	26.7	105.8
2004.....	0.7	1.0	1.0	1.4	2.3	3.1	6.5	12.6	26.1	100.4
2003.....	0.8	1.0	1.0	1.2	2.0	3.3	6.4	10.6	25.4	96.4
2002.....	0.9	1.0	1.3	1.2	1.7	3.1	5.5	10.1	21.5	83.6
2001.....	0.7	0.8	0.9	1.1	1.6	3.1	6.0	11.4	20.1	76.2
2000.....	0.7	0.9	0.9	1.2	2.1	3.0	5.1	9.2	18.5	71.6
1999.....	0.6	0.7	0.8	1.3	2.1	3.3	5.6	9.0	18.3	67.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

White, both sexes

2006 4/.....	0.8	1.0	1.2	1.6	2.6	4.0	7.4	15.6	30.8	116.4
2005 4/.....	0.7	0.8	1.2	1.4	2.1	3.4	6.5	12.6	25.2	98.5
2004 4/.....	0.7	1.0	1.1	1.4	2.3	3.5	6.3	11.8	25.8	92.3
2003 4/.....	0.8	1.0	1.0	1.3	2.2	3.1	6.4	11.0	24.4	88.1
2002.....	0.8	1.0	1.2	1.2	2.1	3.1	6.0	10.8	22.4	78.2
2001.....	0.9	1.0	1.0	1.3	1.9	3.0	5.6	11.4	20.3	71.6
2000.....	1.0	1.0	1.0	1.3	2.1	3.3	5.4	10.0	18.8	68.8
1999.....	0.7	0.8	1.0	1.5	2.2	3.3	5.6	9.4	18.2	64.5

White, male

2006 4/.....	1.0	1.1	1.4	2.0	3.1	4.7	7.9	15.9	29.9	89.1
2005 4/.....	0.9	1.0	1.5	1.7	2.4	3.8	6.8	13.4	23.0	77.9
2004 4/.....	0.8	1.1	1.4	1.6	2.6	3.9	6.4	11.4	26.1	74.7
2003 4/.....	0.9	1.3	1.2	1.6	2.6	3.4	7.1	12.2	23.0	67.6
2002.....	0.9	1.1	1.3	1.3	2.7	3.5	6.8	12.0	24.3	64.0
2001.....	1.1	1.3	1.3	1.5	2.3	3.1	5.8	11.8	21.8	59.4
2000.....	1.3	1.1	1.3	1.5	2.5	4.2	6.2	11.8	20.1	59.8
1999.....	0.9	1.0	1.3	1.9	2.6	3.8	6.1	10.8	18.7	56.8

White, female

2006 4/.....	0.6	0.8	1.0	1.3	2.2	3.4	6.9	15.5	31.4	129.2
2005 4/.....	0.6	0.6	0.9	1.0	1.8	3.0	6.2	12.0	26.6	108.0
2004 4/.....	0.7	0.9	0.8	1.2	2.0	3.1	6.1	12.2	25.6	100.3
2003 4/.....	0.8	0.8	0.8	1.1	1.8	2.9	5.8	10.1	25.3	97.1
2002.....	0.7	0.8	1.1	1.1	1.5	2.8	5.3	10.0	21.2	84.3
2001.....	0.6	0.7	0.7	1.0	1.6	2.8	5.5	11.1	19.4	76.7
2000.....	0.6	0.8	0.8	1.1	1.8	2.6	4.7	8.7	18.0	72.4
1999.....	0.5	0.5	0.7	1.2	1.9	2.9	5.2	8.4	17.9	67.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R53,R55-R94)

Black, both sexes

2006 4/.....	2.8	2.2	3.6	4.5	6.5	9.1	16.5	27.6	50.2	123.8
2005 4/.....	1.6	2.5	2.8	3.7	4.8	8.1	12.7	16.5	37.1	99.7
2004 4/.....	2.0	2.0	3.2	4.2	6.1	6.5	11.9	21.5	37.9	109.7
2003 4/.....	1.9	3.4	3.8	3.7	5.3	7.9	14.3	18.0	33.3	100.7
2002.....	2.2	2.5	3.8	3.8	4.5	6.4	10.3	18.0	33.0	84.3
2001.....	2.4	2.3	3.3	3.8	4.9	8.5	12.9	16.0	27.6	79.2
2000.....	2.0	2.8	3.3	4.5	6.2	8.8	12.3	17.8	28.8	64.8
1999.....	2.0	2.9	3.2	4.0	5.1	8.5	11.9	18.9	26.9	71.1

Black, male

2006 4/.....	3.6	2.5	5.0	5.3	6.5	11.5	21.8	34.5	61.6	119.7
2005 4/.....	1.7	3.1	4.1	3.9	5.3	11.4	15.2	19.2	44.2	95.7
2004 4/.....	2.8	2.4	4.3	6.1	7.3	9.9	12.0	23.7	36.6	96.4
2003 4/.....	2.4	4.4	5.1	4.5	6.4	8.4	14.5	18.3	36.2	97.0
2002.....	2.5	3.2	4.8	4.9	5.9	7.0	12.7	26.6	41.3	78.3
2001.....	3.3	3.3	4.2	5.8	8.9	11.4	12.8	14.9	24.1	77.2
2000.....	2.3	3.5	4.9	8.1	7.0	11.7	15.4	19.4	31.2	56.5
1999.....	2.9	4.3	3.9	5.9	5.9	10.1	12.9	23.3	29.6	58.6

Black, female

2006 4/.....	2.1	2.0	2.4	3.9	6.5	7.3	12.9	23.6	44.1	125.4
2005 4/.....	1.5	2.0	1.7	3.5	4.4	5.6	11.0	14.9	33.4	101.3
2004 4/.....	1.3	1.6	2.3	2.6	5.2	4.0	11.9	20.2	38.6	115.0
2003 4/.....	1.5	2.6	2.6	3.0	4.5	7.5	14.2	17.9	31.7	102.1
2002.....	1.9	1.9	3.0	2.9	3.4	5.9	8.7	12.9	28.6	86.7
2001.....	1.6	*	2.5	*	*	6.4	13.0	16.6	29.4	80.0
2000.....	1.7	2.1	2.0	*	5.6	6.7	10.2	16.8	27.6	67.8
1999.....	*	1.6	2.5	*	4.5	7.3	11.3	16.3	25.5	75.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XX. External causes of mortality (*U01-*U03,V01-Y89)

All races 3/, both sexes

2006.....	68.9	73.9	69.2	58.7	53.9	57.2	69.3	104.6	161.9	302.4
2005.....	67.9	70.7	65.8	55.5	54.7	57.2	73.7	105.9	166.0	309.1
2004.....	66.1	69.2	62.6	53.2	52.3	55.9	70.5	103.6	163.9	307.5
2003.....	67.2	66.7	59.9	53.4	50.7	55.4	71.3	102.9	161.9	310.4
2002.....	66.2	64.5	56.6	51.0	50.3	56.5	72.1	104.7	161.6	308.3
2001 5/.....	64.3	62.5	54.8	50.2	49.5	56.0	69.9	106.1	161.2	310.1
2000.....	61.0	57.9	51.4	48.3	49.1	52.5	69.9	100.7	155.4	310.5
1999.....	59.9	56.7	49.0	48.3	48.0	57.0	72.3	104.8	164.4	318.2

All races 3/, male

2006.....	98.6	106.4	100.7	85.3	77.9	83.3	97.0	148.4	223.5	393.6
2005.....	97.7	101.9	96.9	81.6	80.4	81.9	106.3	150.4	234.6	404.4
2004.....	94.1	98.8	92.1	77.7	75.6	80.7	101.1	145.9	232.1	407.1
2003.....	97.8	97.2	87.4	78.4	74.8	80.0	101.3	146.8	228.3	417.7
2002.....	96.2	94.6	83.1	76.8	73.3	82.4	104.9	150.3	226.1	414.8
2001 5/.....	95.4	94.2	81.1	74.0	71.9	80.8	100.2	152.7	231.8	429.1
2000.....	91.6	87.5	76.8	71.2	70.9	74.9	100.4	145.6	225.1	438.2
1999.....	89.5	85.9	73.2	71.9	71.0	82.8	104.4	149.4	240.9	454.8

All races 3/, female

2006.....	39.5	42.2	38.9	33.6	32.0	34.3	46.7	72.5	123.3	259.7
2005.....	38.4	40.2	36.0	31.0	31.2	35.7	47.3	73.5	123.4	265.3
2004.....	38.3	40.3	34.4	30.1	31.2	34.3	45.9	73.2	121.9	262.7
2003.....	37.0	37.0	33.6	29.8	28.8	34.0	47.2	71.6	121.3	262.9
2002.....	36.6	35.3	31.1	26.9	29.5	34.1	45.9	72.5	122.4	262.0
2001 5/.....	33.7	31.8	29.6	27.8	29.3	34.6	45.8	73.4	118.9	260.3
2000.....	30.9	29.1	27.0	26.8	29.3	33.2	45.9	69.5	114.3	258.5
1999.....	30.7	28.5	25.9	26.4	27.3	35.0	47.3	73.8	119.6	263.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

XX. External causes of mortality (*U01-*U03,V01-Y89)

White, both sexes

2006 4/.....	70.5	75.0	69.5	58.6	54.0	57.7	70.5	106.9	167.8	316.7
2005 4/.....	69.2	72.0	65.3	55.7	54.6	57.4	74.2	106.9	170.2	321.4
2004 4/.....	67.8	70.3	62.7	53.0	52.4	56.2	70.9	105.2	168.8	321.4
2003 4/.....	68.4	67.3	59.7	53.3	50.9	55.1	71.8	104.8	165.9	323.4
2002.....	67.2	64.6	56.2	51.0	50.4	56.0	73.1	106.1	166.2	319.7
2001 5/.....	64.3	61.7	54.4	49.9	48.9	55.9	69.6	107.6	164.7	319.5
2000.....	60.0	56.8	50.0	47.6	48.9	51.8	69.4	101.5	158.6	320.4
1999.....	59.0	55.2	47.8	47.4	46.6	56.5	71.4	104.5	166.6	326.8

White, male

2006 4/.....	99.7	106.3	99.5	83.7	77.1	83.7	98.0	150.3	232.7	411.3
2005 4/.....	98.3	102.4	94.9	80.5	79.1	81.8	106.5	151.6	239.3	419.8
2004 4/.....	95.6	99.3	90.8	76.0	75.0	80.8	101.4	148.3	238.1	423.8
2003 4/.....	98.6	96.8	86.3	77.4	74.7	79.5	102.4	149.2	233.1	434.4
2002.....	97.4	93.9	81.5	75.5	73.0	80.7	105.8	151.4	231.7	427.4
2001 5/.....	95.0	92.0	79.4	72.5	70.4	80.4	99.5	154.4	236.3	441.2
2000.....	89.7	84.9	73.6	69.1	69.8	72.8	99.3	146.2	228.6	451.7
1999.....	87.8	82.5	70.6	69.2	68.6	81.4	102.7	148.6	244.1	465.8

White, female

2006 4/.....	41.0	43.7	40.1	34.3	32.4	34.5	47.6	74.5	126.8	272.4
2005 4/.....	39.7	41.8	36.3	31.8	31.8	35.6	47.5	73.9	126.9	276.1
2004 4/.....	39.8	41.5	35.2	30.9	31.5	34.4	45.9	73.7	125.8	275.2
2003 4/.....	37.9	38.0	33.7	30.2	28.8	33.6	46.7	72.6	124.4	274.2
2002.....	36.9	35.5	31.5	27.5	29.5	34.2	46.4	73.7	126.0	272.8
2001 5/.....	33.4	31.7	29.9	28.4	29.2	34.3	45.4	74.5	121.4	268.6
2000.....	30.2	28.9	26.9	27.0	29.7	33.4	45.3	70.1	116.9	267.0
1999.....	30.1	28.2	25.6	26.6	26.3	34.7	46.6	73.6	120.7	270.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
XX. External causes of mortality (*U01-*U03,V01-Y89)										
Black, both sexes										
2006 4/.....	76.0	82.2	82.7	72.3	61.1	60.7	67.1	91.5	111.7	173.8
2005 4/.....	75.2	77.8	83.6	66.8	62.7	64.4	78.1	109.6	132.0	195.1
2004 4/.....	69.7	77.6	74.7	65.8	58.4	59.7	71.6	99.4	125.0	176.0
2003 4/.....	75.2	76.8	72.8	62.2	57.1	62.4	72.6	89.0	124.9	184.4
2002.....	73.4	78.0	69.9	59.7	57.4	61.7	70.5	98.6	118.5	199.6
2001 5/.....	78.0	80.6	69.3	60.7	60.8	63.5	79.8	98.6	129.2	216.8
2000.....	79.5	76.2	70.5	60.6	56.3	64.2	82.0	97.2	127.1	212.4
1999.....	77.5	78.8	65.8	61.7	62.0	68.3	85.8	115.0	147.9	239.5
Black, male										
2006 4/.....	117.4	129.5	133.6	117.2	99.7	95.8	104.3	145.0	152.7	235.0
2005 4/.....	116.2	121.9	134.2	109.6	107.1	99.1	120.5	167.6	200.2	262.1
2004 4/.....	106.3	118.0	121.9	109.0	93.4	93.8	112.7	149.7	198.7	247.4
2003 4/.....	116.4	121.6	114.5	102.0	91.6	98.2	100.8	136.9	183.1	256.5
2002.....	109.0	122.1	112.6	102.1	92.1	100.7	110.8	157.1	182.6	307.7
2001 5/.....	119.6	131.2	111.9	100.3	96.8	97.8	120.5	151.0	199.8	306.4
2000.....	124.9	123.6	116.0	101.8	91.3	106.9	123.7	150.3	199.8	318.3
1999.....	122.3	130.6	108.6	104.1	97.6	108.0	134.0	175.2	215.4	367.9
Black, female										
2006 4/.....	39.5	41.0	39.6	35.3	31.1	34.8	41.6	59.5	90.1	148.8
2005 4/.....	39.1	39.4	40.9	31.5	28.1	38.6	49.1	74.9	96.1	168.1
2004 4/.....	37.5	42.3	34.8	30.2	31.0	34.4	43.7	69.5	86.2	147.7
2003 4/.....	39.0	37.8	37.5	29.4	30.2	35.8	53.6	60.5	94.4	156.4
2002.....	42.1	39.7	33.7	24.9	30.2	32.7	43.5	63.7	85.1	158.1
2001 5/.....	41.3	36.8	32.9	28.2	32.5	38.1	52.7	67.4	93.2	183.6
2000.....	39.4	35.0	31.8	26.9	28.6	32.6	54.4	65.2	90.4	173.0
1999.....	38.1	34.1	29.4	27.0	34.1	38.9	53.8	78.7	114.5	191.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
All races 3/, both sexes										
2006.....	43.3	47.0	43.7	37.1	35.0	39.6	50.5	81.3	136.0	274.9
2005.....	41.6	44.6	41.6	35.5	36.0	40.1	53.7	82.7	137.0	279.5
2004.....	39.7	42.6	38.6	32.8	33.7	38.4	50.5	79.6	135.9	276.7
2003.....	40.4	40.7	36.6	32.8	32.9	38.1	50.9	78.8	133.5	278.9
2002.....	39.4	38.7	34.3	30.4	32.6	38.3	50.8	80.0	131.1	275.4
2001.....	36.7	36.0	31.9	29.3	31.5	36.8	49.3	80.4	130.4	276.4
2000.....	35.4	34.2	30.8	29.9	32.2	35.4	48.8	75.5	124.6	273.5
1999.....	35.4	34.1	29.2	30.1	31.2	38.7	50.9	78.6	133.3	282.4
All races 3/, male										
2006.....	60.8	66.8	63.0	52.9	49.6	55.1	66.4	107.3	176.5	337.1
2005.....	58.7	63.5	60.7	51.3	51.4	54.6	72.6	110.0	180.8	344.2
2004.....	56.0	60.5	56.5	46.9	46.8	52.8	67.0	104.0	179.7	345.5
2003.....	58.0	58.9	52.3	47.2	47.0	52.3	66.8	104.9	175.2	352.5
2002.....	56.8	56.6	49.6	44.8	45.7	53.4	68.9	106.4	170.1	346.6
2001.....	54.1	54.0	46.9	41.8	44.5	49.9	65.6	108.3	173.2	358.5
2000.....	53.1	51.6	46.0	43.1	45.3	48.1	64.6	99.8	164.8	359.7
1999.....	53.1	51.7	43.6	43.6	44.5	53.1	68.2	102.8	181.8	375.4
All races 3/, female										
2006.....	25.8	27.8	25.2	22.2	21.7	26.0	37.5	62.3	110.7	245.8
2005.....	24.6	26.0	23.3	20.7	22.0	27.4	38.5	62.8	109.7	249.8
2004.....	23.6	25.2	21.4	19.5	21.7	25.8	37.2	62.1	109.0	245.8
2003.....	23.0	23.0	21.6	19.3	20.1	25.8	38.1	60.3	108.1	246.4
2002.....	22.4	21.3	19.6	16.8	20.7	25.3	36.3	61.4	107.3	244.6
2001.....	19.6	18.5	17.5	17.6	19.6	25.5	36.3	60.9	104.8	242.0
2000.....	18.0	17.3	16.1	17.5	20.4	24.5	36.4	58.5	100.9	238.4
1999.....	18.1	17.0	15.5	17.6	19.2	26.4	37.3	61.8	105.0	244.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
White, both sexes										
2006 4/.....	44.9	47.5	43.3	36.3	34.4	39.5	50.9	83.0	140.9	288.6
2005 4/.....	43.2	45.4	40.7	34.7	35.3	39.8	53.8	83.1	140.0	290.9
2004 4/.....	41.3	43.2	37.9	32.0	33.3	38.2	50.5	80.6	140.2	289.9
2003 4/.....	41.5	41.0	35.7	32.1	32.5	37.3	50.9	80.2	136.5	291.1
2002.....	40.5	38.5	33.6	29.6	32.1	37.3	51.0	81.1	134.6	286.1
2001.....	36.8	34.9	31.1	28.5	30.5	36.4	48.9	81.2	133.3	285.5
2000.....	35.2	33.2	29.2	28.9	31.8	34.5	48.1	75.7	127.3	282.8
1999.....	35.2	32.8	27.8	28.9	29.5	37.7	49.5	78.0	135.0	290.4
White, male										
2006 4/.....	62.6	66.7	61.2	50.9	48.0	54.7	66.2	107.9	182.9	352.4
2005 4/.....	60.6	63.8	58.6	49.2	49.2	53.7	71.7	110.0	182.9	356.6
2004 4/.....	57.9	61.0	54.7	44.8	45.4	52.4	66.5	104.6	183.9	359.4
2003 4/.....	59.6	58.6	50.6	45.6	46.1	50.8	66.9	105.8	177.3	365.7
2002.....	58.2	56.4	47.8	42.8	44.5	50.9	68.6	106.5	173.0	356.6
2001.....	54.2	51.8	45.1	40.0	42.6	48.9	64.6	108.2	175.2	368.1
2000.....	52.5	49.8	43.2	40.8	43.8	45.9	63.3	98.7	166.4	369.8
1999.....	52.5	49.1	41.1	41.1	42.0	51.1	65.7	100.7	183.4	383.6
White, female										
2006 4/.....	26.9	28.4	25.8	22.3	21.7	26.0	38.3	64.3	114.4	258.7
2005 4/.....	25.6	26.9	23.0	20.8	22.4	27.5	39.0	63.2	113.1	260.7
2004 4/.....	24.5	25.5	21.5	19.7	22.0	25.6	37.3	63.0	113.0	258.6
2003 4/.....	23.4	23.4	21.1	19.2	19.9	25.4	37.9	61.7	111.3	258.0
2002.....	22.8	20.7	19.7	16.9	20.7	25.3	36.7	62.9	111.2	255.4
2001.....	19.3	18.1	17.4	17.6	19.3	25.4	36.1	62.0	107.9	251.0
2000.....	17.8	16.8	15.5	17.5	20.7	24.6	35.8	59.6	104.0	247.4
1999.....	17.9	16.6	14.9	17.4	18.0	26.1	36.6	62.0	106.5	253.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidents (unintentional injuries) (V01-X59,Y85-Y86)										
Black, both sexes										
2006 4/.....	44.2	53.6	56.7	51.2	44.0	45.3	51.4	71.7	93.7	150.5
2005 4/.....	41.1	49.5	57.2	49.7	45.6	47.5	58.9	87.4	110.9	174.8
2004 4/.....	38.4	49.0	50.7	45.2	41.7	43.8	53.8	77.9	100.5	152.3
2003 4/.....	41.7	47.9	49.1	43.7	40.4	47.7	53.8	67.7	105.1	160.2
2002.....	40.0	48.3	45.1	42.0	40.6	47.3	51.9	74.9	96.8	173.4
2001.....	44.3	50.1	44.8	40.6	43.7	45.5	58.0	78.2	104.5	183.6
2000.....	43.8	47.0	47.0	42.0	39.4	46.1	60.2	75.3	100.0	181.1
1999.....	43.6	50.4	44.4	43.4	46.3	50.7	66.6	90.3	124.2	207.6
Black, male										
2006 4/.....	65.5	81.1	91.2	81.9	72.1	69.4	77.5	109.6	125.3	199.3
2005 4/.....	60.1	75.7	90.9	80.3	79.6	73.2	90.5	130.5	167.1	225.0
2004 4/.....	56.5	72.2	82.2	73.2	66.7	67.4	82.2	116.0	157.1	213.6
2003 4/.....	60.8	73.9	75.2	70.1	64.8	74.4	74.2	105.0	152.8	225.2
2002.....	57.9	72.0	72.6	71.2	65.3	75.8	78.4	117.6	149.8	260.5
2001.....	66.6	81.7	71.8	65.9	69.8	69.1	82.5	119.4	164.5	255.3
2000.....	68.8	74.5	76.5	71.5	65.4	74.2	86.3	119.5	156.0	275.6
1999.....	69.0	82.7	73.2	71.5	70.6	77.9	101.8	135.3	177.0	316.4
Black, female										
2006 4/.....	25.5	29.6	27.6	25.9	22.2	27.5	33.4	49.0	77.0	130.5
2005 4/.....	24.3	26.7	28.7	24.5	19.1	28.4	37.2	61.6	81.2	154.5
2004 4/.....	22.6	28.8	24.2	22.0	22.3	26.3	34.5	55.3	70.7	128.0
2003 4/.....	24.8	25.3	27.0	22.0	21.4	28.0	40.1	45.6	80.1	134.9
2002.....	24.2	27.7	21.8	18.1	21.2	26.1	34.1	49.4	69.2	140.1
2001.....	24.5	22.7	21.8	19.9	23.2	28.0	41.7	53.5	74.0	157.0
2000.....	21.7	23.1	21.9	17.8	18.8	25.3	42.8	48.7	71.8	145.9
1999.....	21.3	22.5	20.1	20.4	27.3	30.6	43.2	63.2	98.1	166.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
All races 3/, both sexes										
2006.....	16.8	17.2	16.4	15.6	16.1	16.2	17.5	21.6	26.3	24.4
2005.....	17.0	16.8	16.6	16.2	16.3	17.2	18.9	23.0	26.0	26.1
2004.....	16.4	17.1	16.3	15.2	15.9	17.0	18.1	23.6	28.1	27.2
2003.....	17.5	16.8	16.3	15.9	16.0	16.2	19.0	23.9	29.4	29.7
2002.....	17.1	16.9	15.8	14.9	16.7	16.7	20.2	25.0	29.7	28.8
2001.....	17.3	16.3	15.6	15.2	15.4	16.4	19.5	26.2	30.4	28.9
2000.....	16.8	16.2	15.5	15.7	16.4	16.8	19.4	25.1	30.0	31.5
1999.....	16.6	15.9	14.9	16.3	15.7	18.1	20.9	25.9	32.1	32.0
All races 3/, male										
2006.....	24.2	25.7	24.5	22.9	22.9	22.9	23.9	30.1	37.5	42.9
2005.....	24.8	24.9	25.0	23.7	23.2	23.9	27.0	32.5	38.0	46.0
2004.....	24.1	24.9	24.4	22.2	22.4	23.2	24.3	32.3	42.6	46.5
2003.....	25.5	24.9	23.6	23.2	22.7	22.5	25.0	32.9	42.1	52.0
2002.....	24.5	25.1	22.7	22.2	22.8	23.4	27.5	34.5	42.6	53.0
2001.....	25.4	24.0	22.5	21.4	22.0	22.1	26.8	36.2	45.5	52.7
2000.....	24.8	23.6	22.3	22.4	22.6	22.6	26.3	34.5	44.0	59.8
1999.....	24.0	23.1	21.5	23.0	21.8	24.8	27.5	36.2	48.3	63.5
All races 3/, female										
2006.....	9.4	8.9	8.6	8.6	9.9	10.2	12.3	15.3	19.3	15.7
2005.....	9.2	8.9	8.6	9.0	10.0	11.3	12.4	16.1	18.5	17.0
2004.....	8.9	9.4	8.6	8.7	10.0	11.5	13.1	17.3	19.1	18.5
2003.....	9.5	9.0	9.2	9.0	9.9	10.7	14.2	17.5	21.7	19.9
2002.....	9.9	8.8	9.2	7.9	11.1	10.9	14.3	18.3	21.8	18.3
2001.....	9.4	8.8	8.9	9.4	9.3	11.6	13.8	19.2	21.3	18.9
2000.....	9.0	8.9	8.9	9.5	10.7	11.7	13.9	18.5	21.7	19.9
1999.....	9.4	8.9	8.6	10.0	10.2	12.4	15.7	18.8	22.6	19.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
White, both sexes										
2006 4/.....	17.3	17.2	16.7	15.4	15.9	16.4	17.7	21.6	26.8	25.3
2005 4/.....	17.5	17.2	16.6	16.4	16.2	17.4	19.0	23.0	26.8	27.1
2004 4/.....	16.8	17.2	16.0	15.2	15.9	17.1	18.2	23.6	29.0	28.4
2003 4/.....	17.7	16.9	16.1	15.7	15.8	16.0	19.2	23.9	30.0	30.5
2002.....	17.5	16.9	15.9	14.8	16.7	16.5	20.5	25.5	30.3	29.6
2001.....	17.5	15.9	15.7	15.1	15.2	16.4	19.5	26.6	31.2	29.9
2000.....	16.8	16.0	15.1	15.6	16.4	16.6	19.2	25.3	30.7	32.2
1999.....	16.7	15.6	14.6	16.1	15.4	18.0	20.8	26.0	32.4	32.3
White, male										
2006 4/.....	24.8	25.6	24.7	22.7	22.4	23.0	23.9	29.8	38.4	44.0
2005 4/.....	25.3	25.3	24.8	23.8	23.0	23.7	26.9	32.1	39.0	47.5
2004 4/.....	24.4	25.0	23.7	21.9	22.0	23.2	24.1	32.2	43.2	48.6
2003 4/.....	25.8	24.9	23.3	22.8	22.6	22.2	25.4	32.7	42.6	52.8
2002.....	24.9	25.3	22.5	21.9	22.8	22.7	27.5	34.4	42.8	53.8
2001.....	25.4	23.2	22.6	21.0	21.5	21.7	26.6	36.2	46.6	54.1
2000.....	24.6	23.4	21.7	21.8	22.2	21.9	25.8	34.3	44.5	61.6
1999.....	24.0	22.5	21.0	22.5	21.1	24.6	27.1	35.9	48.4	63.7
White, female										
2006 4/.....	9.6	8.9	8.8	8.5	9.9	10.4	12.6	15.5	19.5	16.5
2005 4/.....	9.5	9.1	8.5	9.2	10.0	11.7	12.5	16.2	19.2	17.7
2004 4/.....	9.0	9.5	8.5	8.8	10.3	11.6	13.4	17.3	20.1	19.3
2003 4/.....	9.6	9.1	9.1	8.8	9.6	10.5	14.0	17.6	22.2	20.6
2002.....	10.0	8.6	9.3	8.1	11.0	11.0	14.8	19.1	22.6	19.1
2001.....	9.5	8.7	8.9	9.5	9.3	11.6	13.8	19.8	21.8	19.7
2000.....	9.0	8.7	8.6	9.6	11.1	12.0	13.9	19.0	22.6	20.3
1999.....	9.4	8.8	8.4	9.9	10.1	12.3	15.8	19.1	23.0	19.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Transport accidents (V01-V99,Y85)										
Black, both sexes										
2006 4/.....	17.1	18.7	16.9	18.9	18.3	15.5	15.4	20.1	19.9	15.6
2005 4/.....	16.8	16.9	18.9	16.9	17.3	17.2	18.8	22.4	16.3	16.6
2004 4/.....	16.7	19.1	19.9	17.2	17.4	16.6	16.2	23.0	19.6	14.7
2003 4/.....	18.7	18.1	18.2	18.9	17.8	18.5	17.9	22.8	21.8	21.0
2002.....	17.1	18.9	16.5	16.8	17.4	17.3	16.9	20.1	24.1	18.4
2001.....	19.4	20.5	16.8	17.8	18.6	18.6	19.0	22.1	19.8	17.5
2000.....	19.3	19.4	19.3	18.1	16.4	18.2	21.3	19.4	20.9	21.6
1999.....	18.0	19.6	18.3	18.7	17.7	18.0	21.0	24.6	28.3	27.7
Black, male										
2006 4/.....	26.0	29.5	27.4	29.2	29.5	25.3	25.2	33.6	28.1	36.6
2005 4/.....	26.2	25.5	30.8	27.4	27.8	28.8	30.4	38.9	27.4	33.5
2004 4/.....	25.8	29.6	32.6	27.1	28.6	27.1	26.2	36.9	40.9	28.8
2003 4/.....	28.3	29.1	28.9	30.3	27.6	28.6	24.0	37.4	37.7	42.7
2002.....	25.2	29.1	26.3	29.0	27.3	28.7	28.4	38.1	44.3	46.1
2001.....	30.6	33.8	25.7	28.6	30.9	28.2	26.3	39.0	31.3	35.2
2000.....	31.0	29.3	30.0	31.0	27.5	31.2	33.4	34.5	39.6	45.0
1999.....	28.3	31.2	29.1	29.3	28.4	27.3	31.5	40.4	48.8	68.0
Black, female										
2006 4/.....	9.2	9.3	8.0	10.3	9.6	8.3	8.7	12.1	15.6	*
2005 4/.....	8.5	9.4	8.8	8.3	9.1	8.6	10.8	12.5	10.4	9.8
2004 4/.....	8.6	10.0	9.1	9.1	8.7	8.8	9.4	14.8	8.3	9.1
2003 4/.....	10.3	8.4	9.2	9.6	10.2	11.0	13.7	14.1	13.5	12.6
2002.....	10.0	10.0	8.2	6.8	9.6	8.9	9.2	9.3	13.5	*
2001.....	9.4	8.9	9.3	8.9	9.0	11.5	14.1	11.9	13.9	10.9
2000.....	9.0	10.8	10.2	7.6	7.7	8.6	13.3	10.3	11.5	12.9
1999.....	9.0	9.6	9.1	10.1	9.3	11.2	14.0	15.2	18.1	12.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2006.....	15.4	15.7	14.8	14.0	14.7	14.9	16.1	20.2	25.0	23.4
2005.....	15.6	15.3	14.8	14.6	14.8	15.8	17.8	21.6	24.6	25.1
2004.....	15.1	15.5	14.6	13.9	14.3	15.5	16.8	22.7	26.9	26.1
2003.....	16.0	15.1	14.6	14.1	14.3	14.8	17.7	22.5	28.2	28.8
2002.....	15.7	15.2	14.4	13.4	15.2	15.3	18.9	23.8	28.4	28.0
2001.....	15.5	14.5	13.9	13.7	13.7	14.8	18.0	24.6	28.9	27.8
2000.....	15.1	14.6	13.8	14.2	14.7	15.1	18.1	23.8	28.5	30.4
1999.....	14.8	14.0	13.1	14.4	13.9	16.4	19.3	24.1	30.1	30.4

All races 3/, male

2006.....	21.9	23.2	22.0	20.5	20.4	20.8	21.5	27.8	35.0	41.0
2005.....	22.6	22.2	22.1	21.0	20.7	21.5	25.2	30.1	35.7	44.1
2004.....	21.9	22.3	21.6	19.9	19.6	20.8	22.0	30.8	40.4	44.2
2003.....	23.1	22.0	20.8	20.3	19.6	20.1	22.8	30.5	40.0	50.0
2002.....	22.1	22.2	20.3	19.6	20.2	20.8	25.0	32.3	39.9	51.7
2001.....	22.5	21.2	19.7	18.9	19.2	19.2	24.2	33.6	43.1	51.1
2000.....	21.9	21.1	19.6	19.9	19.7	19.8	23.9	32.2	41.4	57.5
1999.....	21.0	20.0	18.6	19.8	18.7	22.0	24.7	33.6	45.2	59.8

All races 3/, female

2006.....	8.9	8.4	8.0	7.9	9.4	9.6	11.7	14.6	18.7	15.2
2005.....	8.7	8.5	7.9	8.5	9.4	10.7	11.8	15.5	17.8	16.4
2004.....	8.3	8.8	8.0	8.2	9.5	10.9	12.6	16.9	18.6	17.9
2003.....	9.0	8.3	8.7	8.3	9.4	10.3	13.7	16.9	21.0	19.5
2002.....	9.4	8.4	8.7	7.5	10.6	10.6	14.1	17.7	21.4	17.7
2001.....	8.7	8.0	8.3	8.8	8.8	11.0	13.1	18.2	20.4	18.0
2000.....	8.4	8.3	8.2	9.0	10.2	11.0	13.5	17.9	20.9	19.3
1999.....	8.6	8.1	8.0	9.4	9.5	11.6	15.0	17.5	21.3	18.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)										
White, both sexes										
2006 4/.....	15.8	15.7	15.1	13.9	14.4	15.0	16.3	20.3	25.4	24.4
2005 4/.....	16.1	15.7	14.9	14.7	14.7	15.9	17.8	21.6	25.5	26.1
2004 4/.....	15.4	15.6	14.3	13.9	14.3	15.6	16.8	22.8	27.7	27.3
2003 4/.....	16.2	15.2	14.4	13.8	14.0	14.6	17.8	22.6	28.8	29.5
2002.....	16.0	15.2	14.3	13.3	15.1	15.1	19.2	24.3	29.0	28.8
2001.....	15.6	14.2	13.9	13.6	13.5	14.7	17.9	25.0	29.7	28.8
2000.....	15.0	14.5	13.4	14.1	14.7	14.9	17.9	24.0	29.2	31.1
1999.....	14.8	13.7	12.9	14.1	13.5	16.3	19.2	24.3	30.6	30.7
White, male										
2006 4/.....	22.4	23.1	22.2	20.2	19.9	20.8	21.4	27.4	35.6	42.2
2005 4/.....	23.1	22.7	22.1	21.0	20.5	21.4	25.0	29.7	36.7	45.6
2004 4/.....	22.2	22.3	20.9	19.6	19.2	20.7	21.7	30.8	40.8	46.3
2003 4/.....	23.2	22.1	20.5	19.7	19.4	19.9	23.0	30.3	40.6	50.7
2002.....	22.5	22.3	20.0	19.3	20.0	20.1	25.0	32.4	40.1	52.4
2001.....	22.5	20.5	19.7	18.7	18.5	19.0	23.9	33.6	44.3	52.6
2000.....	21.6	20.9	18.9	19.3	19.2	19.1	23.4	32.0	41.7	59.2
1999.....	20.9	19.4	18.1	19.2	18.1	21.8	24.4	33.3	45.5	60.3
White, female										
2006 4/.....	9.1	8.3	8.2	7.7	9.2	9.8	12.0	14.9	19.0	16.0
2005 4/.....	9.0	8.7	7.8	8.7	9.3	11.0	11.9	15.5	18.5	17.1
2004 4/.....	8.4	8.9	7.9	8.3	9.8	11.1	12.8	16.9	19.6	18.7
2003 4/.....	9.0	8.4	8.6	8.1	9.1	10.0	13.5	17.1	21.5	20.1
2002.....	9.5	8.2	8.8	7.6	10.5	10.7	14.5	18.5	22.2	18.5
2001.....	8.8	8.0	8.2	8.8	8.8	11.0	13.0	19.0	20.9	18.9
2000.....	8.4	8.1	7.9	9.1	10.6	11.3	13.5	18.4	21.8	19.7
1999.....	8.6	8.0	7.7	9.3	9.3	11.6	15.1	18.0	21.8	18.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2006 4/.....	15.8	17.2	15.5	17.3	17.6	14.6	14.8	18.8	18.9	14.0
2005 4/.....	15.6	15.0	16.7	15.9	16.2	15.9	18.0	21.4	15.0	15.8
2004 4/.....	15.4	17.3	18.3	15.9	15.5	15.6	15.3	20.8	19.3	14.7
2003 4/.....	17.5	16.2	16.8	17.5	16.4	16.9	17.2	21.3	20.6	20.7
2002.....	15.9	16.9	15.1	15.0	16.2	15.9	15.8	18.4	22.5	18.1
2001.....	17.6	18.1	15.3	15.9	16.9	16.5	18.3	20.1	19.0	16.6
2000.....	17.9	17.6	18.3	16.7	15.4	16.6	20.1	18.1	20.0	20.3
1999.....	16.6	17.8	16.3	17.1	15.8	16.6	19.6	22.9	26.0	25.5

Black, male

2006 4/.....	24.1	26.6	24.7	26.5	27.7	23.7	23.7	31.3	27.4	33.1
2005 4/.....	24.3	22.4	26.8	25.5	26.0	25.7	29.4	36.7	26.0	32.5
2004 4/.....	23.5	26.7	29.7	24.5	25.0	25.4	24.3	33.2	40.2	28.8
2003 4/.....	26.4	25.8	26.0	27.6	24.5	25.1	23.0	35.1	34.7	41.7
2002.....	23.0	25.6	23.6	25.3	24.9	25.4	25.7	33.5	40.5	45.0
2001.....	27.5	29.9	23.2	24.7	28.5	23.9	24.6	36.2	30.5	35.2
2000.....	28.4	26.3	28.1	28.2	25.7	27.8	31.4	32.1	38.8	43.8
1999.....	25.6	28.2	25.5	26.3	24.6	24.9	28.5	38.9	45.3	60.9

Black, female

2006 4/.....	8.6	9.0	7.7	9.7	9.6	8.0	8.7	11.2	14.5	*
2005 4/.....	7.9	8.5	8.1	8.0	8.6	8.6	10.2	12.3	9.3	9.0
2004 4/.....	8.3	9.1	8.8	8.8	8.0	8.3	9.2	13.5	8.3	9.1
2003 4/.....	9.7	7.9	9.1	9.2	10.1	10.8	13.3	13.0	13.2	12.6
2002.....	9.6	9.4	8.0	6.6	9.4	8.9	9.2	9.3	13.1	*
2001.....	8.9	8.0	8.7	8.7	7.8	11.1	14.1	10.5	13.1	9.7
2000.....	8.6	10.0	9.9	7.3	7.4	8.3	12.7	9.7	10.6	11.6
1999.....	8.7	8.9	8.5	9.6	8.8	10.4	13.7	13.2	16.4	12.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pedestrian involved in collision with motor vehicle (V02-V04)										
All races 3/, both sexes										
2006.....	1.2	1.6	1.3	1.2	1.5	1.5	1.5	2.5	2.6	2.2
2005.....	1.3	1.4	1.4	1.2	1.2	1.6	1.8	2.2	2.7	2.4
2004.....	1.2	1.4	1.4	1.4	1.5	1.3	1.8	2.2	2.8	2.7
2003.....	1.4	1.5	1.3	1.4	1.5	1.6	1.6	2.3	3.1	2.9
2002.....	1.4	1.4	1.3	1.4	1.4	1.9	2.1	2.4	2.8	3.3
2001.....	1.5	1.4	1.4	1.4	1.3	1.2	2.0	2.4	3.8	3.0
2000.....	1.4	1.4	1.3	1.5	1.5	1.5	1.7	2.5	3.3	3.6
1999.....	1.3	1.3	1.2	1.6	1.5	1.4	1.9	2.3	2.8	3.7
All races 3/, male										
2006.....	1.7	2.4	2.1	1.8	2.3	2.0	1.9	3.2	3.3	3.7
2005.....	2.0	2.1	2.2	1.8	1.7	2.4	2.7	3.3	3.8	3.7
2004.....	1.9	2.0	2.3	2.0	2.2	1.8	2.1	2.9	4.2	4.9
2003.....	2.1	2.1	2.0	2.3	2.2	2.3	2.3	2.7	4.1	5.7
2002.....	2.0	2.2	2.0	2.0	1.9	2.6	2.8	3.0	3.9	5.8
2001.....	2.2	2.0	1.9	2.0	2.0	1.4	2.9	3.2	5.6	5.9
2000.....	2.0	2.2	1.9	2.3	2.1	2.1	2.3	3.4	4.5	7.0
1999.....	1.9	2.0	1.9	2.3	2.1	2.0	2.5	3.2	4.7	6.5
All races 3/, female										
2006.....	0.8	0.8	0.6	0.6	0.8	1.0	1.2	1.9	2.1	1.4
2005.....	0.6	0.7	0.6	0.7	0.7	0.9	1.0	1.4	2.0	1.7
2004.....	0.6	0.8	0.5	0.8	0.9	0.9	1.5	1.7	2.0	1.7
2003.....	0.7	0.9	0.6	0.7	0.9	1.1	1.0	2.1	2.5	1.7
2002.....	0.9	0.7	0.7	0.8	1.0	1.3	1.5	2.0	2.1	2.2
2001.....	0.7	0.8	0.8	0.8	0.7	1.0	1.3	1.8	2.7	1.8
2000.....	0.8	0.6	0.8	0.8	0.9	1.0	1.3	1.9	2.5	2.3
1999.....	0.7	0.6	0.5	0.9	0.9	0.9	1.4	1.7	1.7	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Pedestrian involved in collision with motor vehicle (V02-V04)										
White, both sexes										
2006 4/.....	1.1	1.4	1.3	1.0	1.3	1.4	1.4	2.3	2.4	2.0
2005 4/.....	1.2	1.3	1.2	1.2	1.0	1.5	1.6	2.0	2.6	2.3
2004 4/.....	1.2	1.2	1.2	1.3	1.4	1.2	1.7	2.1	2.8	2.6
2003 4/.....	1.2	1.4	1.1	1.4	1.5	1.4	1.5	2.1	2.9	2.9
2002.....	1.3	1.3	1.2	1.3	1.3	1.6	2.0	2.2	2.6	3.2
2001.....	1.2	1.3	1.2	1.3	1.2	1.1	1.8	2.2	3.5	2.9
2000.....	1.2	1.2	1.2	1.3	1.4	1.4	1.6	2.3	3.2	3.4
1999.....	1.0	1.2	1.1	1.4	1.4	1.3	1.7	2.1	2.7	3.6
White, male										
2006 4/.....	1.7	2.1	1.9	1.6	2.0	2.0	1.9	2.9	3.0	3.3
2005 4/.....	1.9	2.0	1.9	1.7	1.5	2.2	2.5	2.9	3.5	3.6
2004 4/.....	1.8	1.8	2.0	1.8	2.0	1.7	2.0	2.7	4.0	4.6
2003 4/.....	1.9	2.0	1.8	2.2	2.1	2.0	2.1	2.4	3.9	5.4
2002.....	1.8	2.0	1.8	1.9	1.8	2.3	2.7	2.7	3.6	5.4
2001.....	1.9	1.9	1.7	1.9	1.7	1.3	2.6	2.9	5.5	5.5
2000.....	1.7	1.9	1.7	2.0	1.9	1.8	2.1	3.0	4.1	6.2
1999.....	1.5	1.7	1.7	2.0	2.0	1.9	2.1	2.8	4.4	6.0
White, female										
2006 4/.....	0.6	0.7	0.6	0.5	0.7	0.9	1.1	1.9	2.1	1.4
2005 4/.....	0.6	0.6	0.5	0.7	0.6	0.9	0.9	1.2	2.1	1.8
2004 4/.....	0.5	0.7	0.5	0.9	0.8	0.8	1.5	1.6	2.0	1.7
2003 4/.....	0.6	0.8	0.5	0.6	0.8	1.0	0.9	2.0	2.4	1.8
2002.....	0.8	0.6	0.7	0.8	0.9	1.1	1.5	1.8	2.0	2.2
2001.....	0.6	0.7	0.7	0.8	0.7	0.9	1.1	1.7	2.4	1.9
2000.....	0.6	0.5	0.7	0.7	0.9	1.0	1.3	1.8	2.6	2.2
1999.....	0.6	0.6	0.5	0.8	0.8	0.7	1.4	1.6	1.7	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

All races 3/, both sexes

2006.....	1.3	1.3	1.2	0.9	1.0	1.0	1.1	1.3	1.4	1.4
2005.....	1.3	1.3	1.1	1.0	1.0	1.2	1.1	1.4	1.2	1.3
2004.....	1.4	1.3	1.1	1.1	1.2	1.2	1.3	1.4	1.8	1.2
2003.....	1.6	1.5	1.4	1.1	1.1	1.1	1.6	1.5	1.7	1.9
2002.....	1.6	1.5	1.3	1.2	1.4	1.2	1.2	1.5	1.7	1.5
2001.....	1.4	1.3	1.1	0.9	1.0	1.1	1.1	1.7	1.6	1.7
2000.....	1.2	1.1	1.0	0.9	1.0	1.0	1.3	1.3	1.5	1.7
1999.....	1.2	1.1	1.0	1.0	1.0	1.3	1.1	1.4	1.6	1.2

All races 3/, male

2006.....	1.8	1.9	1.8	1.3	1.4	1.5	1.6	1.7	2.0	2.4
2005.....	1.7	1.9	1.6	1.5	1.5	1.7	1.8	2.2	1.7	2.3
2004.....	2.0	2.0	1.6	1.7	1.7	1.7	1.9	2.4	2.6	1.9
2003.....	2.4	2.2	1.9	1.7	1.6	1.6	2.2	2.2	2.5	3.7
2002.....	2.4	2.1	1.8	1.8	2.3	1.7	1.8	2.3	2.1	3.2
2001.....	2.2	2.0	1.6	1.6	1.5	1.6	1.5	2.5	2.6	3.1
2000.....	1.8	1.5	1.5	1.3	1.5	1.5	1.8	2.0	2.2	3.0
1999.....	1.8	1.6	1.6	1.4	1.5	1.9	1.6	2.1	2.8	2.5

All races 3/, female

2006.....	0.8	0.6	0.5	0.5	0.6	0.6	0.8	1.1	1.0	0.9
2005.....	0.8	0.7	0.5	0.5	0.5	0.7	0.6	0.8	0.9	0.8
2004.....	0.7	0.6	0.6	0.5	0.7	0.8	0.9	0.7	1.3	0.9
2003.....	0.8	0.8	0.8	0.5	0.6	0.7	1.1	1.1	1.1	1.1
2002.....	0.9	0.9	0.7	0.6	0.6	0.8	0.8	0.9	1.5	0.8
2001.....	0.6	0.6	0.6	0.3	0.5	0.7	0.8	1.1	0.9	1.2
2000.....	0.7	0.7	0.5	0.5	0.6	0.6	1.0	0.8	1.0	1.2
1999.....	0.6	0.7	0.4	0.7	0.6	0.8	0.7	1.0	0.9	0.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of motor vehicle involved in collision with other (non-motorized) road vehicle, streetcar, animal or pedestrian (V40-V41,V46-V47,V50-V51,V56-V57, V60-V61,V66-V67,V70-V71,V76-V77)

White, both sexes

2006 4/.....	1.3	1.3	1.1	0.9	0.9	1.0	1.2	1.3	1.4	1.4
2005 4/.....	1.3	1.3	1.1	1.0	1.0	1.1	1.1	1.4	1.2	1.3
2004 4/.....	1.4	1.3	1.1	1.0	1.1	1.2	1.3	1.5	1.9	1.2
2003 4/.....	1.7	1.5	1.3	1.1	1.0	1.0	1.6	1.6	1.8	2.0
2002.....	1.8	1.5	1.2	1.1	1.4	1.1	1.2	1.6	1.8	1.6
2001.....	1.4	1.2	1.1	0.9	0.9	1.1	1.1	1.7	1.5	1.9
2000.....	1.2	1.0	0.9	0.8	1.0	1.0	1.3	1.2	1.5	1.8
1999.....	1.2	1.1	0.9	0.9	1.0	1.3	1.1	1.3	1.6	1.2

White, male

2006 4/.....	1.8	1.9	1.8	1.3	1.3	1.5	1.6	1.6	2.1	2.3
2005 4/.....	1.8	1.9	1.6	1.5	1.6	1.6	1.7	2.3	1.8	2.5
2004 4/.....	2.0	2.0	1.6	1.5	1.7	1.6	1.8	2.5	2.7	2.1
2003 4/.....	2.5	2.2	1.9	1.7	1.5	1.5	2.2	2.2	2.7	3.9
2002.....	2.6	2.0	1.7	1.6	2.2	1.5	1.7	2.4	2.1	3.3
2001.....	2.1	1.9	1.6	1.6	1.4	1.5	1.4	2.4	2.6	3.2
2000.....	1.7	1.5	1.3	1.2	1.5	1.4	1.6	1.8	2.2	3.2
1999.....	1.8	1.5	1.5	1.3	1.5	1.9	1.5	2.0	2.9	2.5

White, female

2006 4/.....	0.8	0.6	0.5	0.5	0.6	0.7	0.8	1.1	1.0	1.0
2005 4/.....	0.8	0.7	0.6	0.5	0.5	0.7	0.6	0.8	0.9	0.8
2004 4/.....	0.8	0.6	0.6	0.6	0.6	0.8	0.9	0.8	1.4	0.9
2003 4/.....	0.9	0.8	0.7	0.5	0.6	0.6	1.0	1.1	1.2	1.2
2002.....	0.9	0.9	0.7	0.6	0.6	0.9	0.8	1.0	1.6	0.9
2001.....	0.6	0.6	0.6	*	0.4	0.8	0.8	1.2	0.9	1.3
2000.....	0.7	0.6	0.5	0.6	0.6	0.6	1.0	0.8	1.1	1.2
1999.....	0.6	0.7	0.3	0.6	0.5	0.7	0.8	0.9	0.9	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

All races 3/, both sexes

2006.....	2.7	2.8	2.8	2.8	3.4	3.9	4.5	5.9	7.4	7.3
2005.....	3.0	2.9	3.2	3.3	3.6	4.3	4.8	6.8	7.6	8.3
2004.....	2.9	3.1	3.2	3.4	3.6	4.8	5.1	7.6	9.2	9.6
2003.....	3.3	3.1	3.4	3.3	3.7	4.1	5.8	7.7	10.0	10.2
2002.....	3.4	3.2	3.4	3.4	4.4	4.2	6.2	8.3	10.4	9.9
2001.....	3.3	3.1	3.1	3.5	3.6	4.6	5.7	7.8	9.1	10.1
2000.....	3.0	3.2	3.1	3.6	4.0	4.2	5.9	7.9	10.5	10.1
1999.....	3.3	3.2	3.2	3.8	3.7	4.5	6.1	7.7	10.4	9.2

All races 3/, male

2006.....	3.3	3.6	3.5	3.3	4.0	4.4	5.5	7.9	9.5	12.4
2005.....	3.8	3.5	4.0	4.0	4.3	5.0	5.3	8.7	11.1	13.7
2004.....	3.6	3.7	3.8	4.0	4.0	5.6	6.0	9.2	13.4	15.9
2003.....	4.2	3.9	4.2	4.0	4.4	4.7	6.3	9.7	13.4	16.6
2002.....	4.2	4.0	4.2	4.5	4.9	4.9	7.3	10.7	14.5	16.9
2001.....	4.1	3.9	3.7	4.1	4.2	5.5	6.8	9.6	13.0	18.1
2000.....	3.7	4.2	3.9	4.4	4.5	4.6	7.2	9.5	15.0	18.2
1999.....	4.1	4.0	3.9	4.2	4.4	5.2	7.3	9.7	14.9	17.8

All races 3/, female

2006.....	2.1	2.1	2.2	2.3	2.8	3.4	3.8	4.4	6.1	5.0
2005.....	2.3	2.2	2.4	2.6	2.9	3.6	4.4	5.4	5.5	5.9
2004.....	2.2	2.5	2.5	2.8	3.2	4.2	4.5	6.4	6.7	6.7
2003.....	2.4	2.3	2.6	2.7	3.1	3.6	5.3	6.3	7.9	7.4
2002.....	2.6	2.4	2.7	2.3	3.9	3.6	5.3	6.7	7.9	6.8
2001.....	2.5	2.4	2.5	2.9	3.1	3.8	4.8	6.5	6.9	6.8
2000.....	2.3	2.3	2.4	2.8	3.6	3.8	4.9	6.9	7.8	6.8
1999.....	2.4	2.4	2.5	3.4	3.1	4.0	5.2	6.4	7.8	5.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)

White, both sexes

2006 4/.....	2.6	2.8	2.8	2.7	3.4	4.0	4.6	6.1	7.9	7.8
2005 4/.....	3.1	2.9	3.1	3.3	3.6	4.4	4.9	6.9	8.1	8.8
2004 4/.....	3.0	3.1	3.1	3.4	3.7	5.0	5.3	7.7	9.6	10.1
2003 4/.....	3.3	3.2	3.3	3.1	3.6	4.0	5.9	7.9	10.5	10.6
2002.....	3.5	3.1	3.4	3.4	4.4	4.3	6.4	8.7	10.9	10.4
2001.....	3.4	3.0	3.1	3.6	3.5	4.6	5.7	8.2	9.5	10.6
2000.....	3.0	3.2	3.0	3.6	4.0	4.3	6.0	8.2	10.9	10.6
1999.....	3.2	3.2	3.1	3.8	3.7	4.6	6.2	8.0	10.7	9.4

White, male

2006 4/.....	3.1	3.5	3.4	3.1	3.9	4.5	5.4	8.0	10.3	12.9
2005 4/.....	3.8	3.5	3.9	4.0	4.3	5.0	5.2	8.7	11.8	14.3
2004 4/.....	3.7	3.6	3.8	3.9	4.0	5.8	6.0	9.3	13.5	16.6
2003 4/.....	4.1	4.0	4.1	3.7	4.3	4.6	6.4	9.8	13.8	17.3
2002.....	4.2	3.9	4.2	4.4	4.9	4.9	7.3	10.8	15.0	17.9
2001.....	4.0	3.6	3.6	4.3	4.0	5.4	6.9	9.9	13.4	18.9
2000.....	3.6	4.1	3.7	4.4	4.1	4.7	7.3	9.6	15.6	19.3
1999.....	4.0	4.0	3.7	4.1	4.3	5.0	7.3	9.8	15.2	18.1

White, female

2006 4/.....	2.1	2.1	2.2	2.3	2.9	3.5	3.9	4.7	6.4	5.3
2005 4/.....	2.3	2.3	2.4	2.6	2.9	3.7	4.7	5.6	5.8	6.2
2004 4/.....	2.2	2.6	2.5	2.8	3.4	4.3	4.8	6.6	7.1	7.2
2003 4/.....	2.4	2.3	2.7	2.6	3.0	3.5	5.4	6.5	8.5	7.7
2002.....	2.7	2.3	2.7	2.3	4.0	3.7	5.7	7.1	8.4	7.1
2001.....	2.7	2.4	2.6	3.0	3.1	3.9	4.8	6.9	7.2	7.2
2000.....	2.3	2.3	2.3	2.9	3.8	4.0	5.0	7.2	8.2	7.1
1999.....	2.5	2.4	2.6	3.5	3.2	4.2	5.3	6.7	8.1	5.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Occupant of car, pickup truck or van involved in collision with other motor vehicle (V42-V44,V49,V52-V54,V59)										
Black, both sexes										
2006 4/.....	3.5	3.2	3.3	4.0	3.4	3.5	4.8	4.5	*	*
2005 4/.....	3.2	2.7	3.7	3.4	3.7	4.6	4.4	7.2	*	5.3
2004 4/.....	2.8	3.6	3.3	4.3	2.9	4.5	4.2	6.3	7.4	5.6
2003 4/.....	3.8	3.0	3.9	4.9	4.4	5.1	4.9	6.5	6.6	6.4
2002.....	3.7	3.5	3.5	3.3	4.3	3.4	4.8	5.5	6.0	*
2001.....	3.7	4.1	3.6	3.1	4.7	5.4	6.5	4.3	6.2	*
2000.....	3.6	4.0	4.4	3.5	4.7	3.4	5.5	4.8	6.8	*
1999.....	3.7	3.5	3.7	4.1	3.9	4.5	5.7	5.2	7.8	7.7
Black, male										
2006 4/.....	5.0	4.2	4.6	5.7	4.9	*	7.4	*	*	*
2005 4/.....	4.2	3.4	5.1	4.2	5.1	5.7	7.3	10.7	*	*
2004 4/.....	3.3	5.1	3.8	5.3	4.7	5.6	*	9.1	16.5	*
2003 4/.....	5.4	3.7	5.5	6.7	5.6	6.6	*	9.6	*	*
2002.....	4.8	4.6	4.9	5.1	5.5	*	7.0	9.2	*	*
2001.....	5.6	6.6	4.9	3.8	6.1	7.6	*	*	*	*
2000.....	5.4	5.7	5.8	5.6	8.1	*	*	*	*	*
1999.....	5.5	4.8	5.1	5.4	5.3	6.8	8.8	9.5	*	*
Black, female										
2006 4/.....	2.1	2.3	2.3	2.6	*	*	*	*	*	*
2005 4/.....	2.2	2.2	2.4	2.7	*	3.7	*	*	*	*
2004 4/.....	2.3	2.4	2.8	3.6	*	3.8	*	*	*	*
2003 4/.....	2.4	2.3	2.6	3.5	3.5	4.0	4.7	*	*	*
2002.....	2.8	2.6	2.4	*	3.3	*	*	*	*	*
2001.....	2.0	1.9	2.6	2.6	3.7	3.8	6.5	*	*	*
2000.....	2.1	2.5	3.3	*	*	*	4.9	*	*	*
1999.....	2.1	2.4	2.5	3.0	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

All races 3/, both sexes

2006.....	1.4	1.2	1.1	1.0	1.0	0.8	0.9	0.9	1.2	1.0
2005.....	1.4	1.2	1.3	0.9	1.1	1.0	0.9	0.8	1.2	0.9
2004.....	1.5	1.3	1.2	1.1	1.1	1.1	0.9	0.9	0.8	1.0
2003.....	1.6	1.4	1.3	1.2	1.1	1.0	1.0	1.0	1.0	1.3
2002.....	1.5	1.4	1.2	1.1	1.3	0.9	1.0	1.0	1.0	0.9
2001.....	1.4	1.3	1.2	1.0	1.2	0.9	1.0	1.1	1.0	0.7
2000.....	1.3	1.2	1.1	0.9	1.0	0.8	0.8	0.9	0.9	1.0
1999.....	1.4	1.2	1.0	1.1	1.0	1.1	0.8	0.8	0.8	1.0

All races 3/, male

2006.....	1.9	1.9	1.5	1.5	1.3	1.2	1.5	1.3	2.1	1.7
2005.....	1.9	1.7	1.9	1.3	1.7	1.5	1.3	1.0	1.5	1.4
2004.....	2.1	1.8	1.8	1.7	1.5	1.5	1.3	1.4	1.1	1.7
2003.....	2.2	2.0	1.8	1.6	1.5	1.5	1.5	1.5	1.7	2.0
2002.....	2.2	2.2	1.7	1.6	1.7	1.4	1.5	1.5	1.6	1.8
2001.....	2.0	2.0	1.8	1.5	1.7	1.3	1.5	1.5	1.4	*
2000.....	1.9	1.9	1.6	1.3	1.4	1.1	1.3	1.4	1.3	2.2
1999.....	2.0	1.7	1.5	1.6	1.5	1.6	0.9	1.3	*	2.0

All races 3/, female

2006.....	0.9	0.6	0.8	0.6	0.8	0.5	0.4	0.7	0.7	0.7
2005.....	0.9	0.8	0.7	0.6	0.6	0.6	0.5	0.6	0.9	0.6
2004.....	0.9	0.8	0.6	0.5	0.7	0.6	0.5	0.6	*	0.7
2003.....	0.9	0.8	0.8	0.8	0.7	0.7	0.6	0.7	*	1.0
2002.....	0.8	0.7	0.7	0.6	0.9	0.6	0.6	0.7	0.6	*
2001.....	0.8	0.6	0.7	0.6	0.7	0.6	0.6	0.8	0.7	*
2000.....	0.7	0.6	0.5	0.6	0.6	0.6	*	0.5	0.7	*
1999.....	0.8	0.7	0.5	0.6	0.7	0.6	0.7	0.5	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Occupant of motor vehicle involved in noncollision accident (V48,V58,V68,V78)

White, both sexes

2006 4/.....	1.4	1.2	1.2	1.0	1.0	0.8	0.9	1.0	1.2	1.1
2005 4/.....	1.5	1.3	1.3	0.9	1.2	1.1	0.9	0.8	1.2	0.8
2004 4/.....	1.5	1.3	1.2	1.1	1.1	1.1	0.8	0.9	0.8	1.1
2003 4/.....	1.7	1.5	1.3	1.2	1.0	1.0	1.0	1.0	1.0	1.3
2002.....	1.5	1.5	1.2	1.1	1.3	1.0	1.0	1.0	1.0	0.8
2001.....	1.5	1.2	1.3	0.9	1.1	1.0	1.0	1.1	1.0	0.7
2000.....	1.3	1.3	1.0	0.8	0.9	0.9	0.8	0.9	0.9	1.0
1999.....	1.4	1.2	1.0	1.1	1.0	1.1	0.8	0.8	0.8	1.0

White, male

2006 4/.....	1.9	1.9	1.5	1.4	1.4	1.1	1.4	1.3	2.1	1.7
2005 4/.....	2.0	1.8	1.9	1.3	1.7	1.6	1.4	1.0	1.6	*
2004 4/.....	2.1	1.7	1.7	1.8	1.5	1.5	1.1	1.2	1.1	1.9
2003 4/.....	2.3	2.1	1.7	1.5	1.4	1.4	1.6	1.5	1.7	2.1
2002.....	2.1	2.3	1.6	1.5	1.7	1.4	1.5	1.6	1.5	1.7
2001.....	2.1	1.9	1.9	1.3	1.5	1.4	1.6	1.6	1.5	*
2000.....	2.0	1.9	1.6	1.1	1.4	1.1	1.3	1.4	1.3	2.2
1999.....	2.1	1.7	1.5	1.6	1.4	1.6	0.8	1.3	*	1.9

White, female

2006 4/.....	0.9	0.6	0.8	0.6	0.8	0.5	*	0.7	*	0.7
2005 4/.....	0.9	0.8	0.7	0.6	0.7	0.6	0.5	0.6	0.9	*
2004 4/.....	0.9	0.8	0.6	0.5	0.7	0.6	0.5	0.6	*	0.7
2003 4/.....	1.0	0.9	0.8	0.9	0.7	0.7	0.6	0.6	*	0.9
2002.....	0.8	0.7	0.8	0.6	0.9	0.6	0.7	0.6	0.7	*
2001.....	0.9	0.6	0.7	0.5	0.8	0.6	0.6	0.8	0.8	*
2000.....	0.7	0.7	0.5	0.5	0.5	0.7	*	0.5	0.7	*
1999.....	0.7	0.6	0.5	0.5	0.6	0.6	0.7	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

All races 3/, both sexes

2006.....	5.6	5.4	5.2	5.2	5.6	5.6	6.7	8.4	11.2	10.8
2005.....	5.4	5.4	5.1	5.5	5.8	5.7	7.7	9.1	10.8	11.1
2004.....	5.1	5.4	4.9	4.7	5.2	5.7	6.6	9.5	11.6	10.8
2003.....	5.2	4.8	4.7	4.7	5.4	5.6	6.5	8.8	11.5	11.9
2002.....	5.3	5.1	4.9	4.6	5.2	5.9	7.3	9.6	11.4	11.4
2001.....	5.4	5.1	5.1	5.3	5.2	5.8	7.2	10.4	12.4	11.7
2000.....	5.7	5.4	5.5	5.6	5.8	6.2	7.3	10.2	11.5	13.1
1999.....	5.4	5.2	5.1	5.3	5.2	6.8	8.1	10.8	13.5	13.9

All races 3/, male

2006.....	7.7	7.4	7.2	7.0	7.2	7.6	8.6	11.4	15.3	18.7
2005.....	7.6	7.7	7.0	7.6	7.6	7.2	11.1	12.1	15.4	20.1
2004.....	7.2	7.5	6.8	6.5	6.9	7.4	8.5	12.6	17.3	17.9
2003.....	7.1	6.8	6.3	6.3	7.0	7.5	8.0	12.1	16.1	20.3
2002.....	6.9	7.0	6.5	6.5	6.5	8.0	9.5	13.0	15.3	21.7
2001.....	7.5	7.2	7.1	6.8	6.9	7.0	9.4	14.3	18.2	21.2
2000.....	8.0	7.4	7.5	7.4	7.6	7.9	9.1	13.6	16.6	24.7
1999.....	7.2	7.2	6.7	7.3	6.7	8.8	9.9	15.0	19.6	27.5

All races 3/, female

2006.....	3.5	3.5	3.3	3.5	4.0	3.8	5.2	6.2	8.7	7.1
2005.....	3.2	3.3	3.2	3.5	4.2	4.4	5.1	7.0	8.0	7.0
2004.....	3.1	3.4	3.1	3.1	3.7	4.3	5.1	7.2	8.1	7.7
2003.....	3.4	2.8	3.2	3.2	3.9	4.1	5.4	6.5	8.7	8.1
2002.....	3.7	3.3	3.4	2.8	3.9	4.1	5.5	7.2	9.0	6.9
2001.....	3.4	3.1	3.2	3.9	3.7	4.8	5.4	7.7	9.0	7.7
2000.....	3.4	3.5	3.6	3.9	4.2	4.8	5.8	7.8	8.6	8.4
1999.....	3.6	3.3	3.6	3.4	3.9	5.2	6.7	7.8	9.9	8.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

White, both sexes

2006 4/.....	5.8	5.2	5.1	5.1	5.3	5.5	6.7	8.4	11.2	11.4
2005 4/.....	5.4	5.4	5.0	5.4	5.7	5.7	7.7	9.1	11.2	11.6
2004 4/.....	5.1	5.3	4.6	4.6	5.2	5.6	6.5	9.5	12.0	11.5
2003 4/.....	5.1	4.5	4.5	4.5	5.2	5.7	6.6	8.9	11.6	12.1
2002.....	5.3	5.0	4.8	4.5	5.1	5.9	7.4	9.8	11.6	11.8
2001.....	5.4	5.0	5.1	5.1	5.1	5.7	7.2	10.6	13.0	12.2
2000.....	5.7	5.2	5.3	5.5	5.9	6.1	7.1	10.3	11.9	13.5
1999.....	5.4	5.0	4.9	5.2	5.0	6.8	8.2	10.8	13.6	14.3

White, male

2006 4/.....	7.9	7.0	7.0	6.8	6.8	7.3	8.4	11.2	15.3	19.9
2005 4/.....	7.5	7.5	6.9	7.3	7.2	7.0	11.0	11.9	15.8	20.8
2004 4/.....	7.1	7.4	6.2	6.3	6.6	7.0	8.4	12.8	17.6	19.0
2003 4/.....	7.0	6.4	6.0	5.9	6.8	7.5	8.1	12.2	16.3	20.4
2002.....	6.9	6.8	6.3	6.2	6.4	7.8	9.7	12.9	15.4	21.9
2001.....	7.4	6.9	7.1	6.4	6.7	6.9	9.3	14.3	18.8	22.3
2000.....	7.8	7.1	7.2	7.1	7.5	7.5	8.7	13.8	16.8	25.8
1999.....	7.2	6.8	6.5	7.2	6.3	8.7	10.1	14.7	19.6	28.3

White, female

2006 4/.....	3.7	3.5	3.3	3.4	3.9	3.9	5.4	6.3	8.7	7.5
2005 4/.....	3.2	3.3	3.2	3.5	4.2	4.6	5.0	7.0	8.4	7.3
2004 4/.....	3.1	3.3	3.0	3.1	3.8	4.4	5.0	7.1	8.4	8.0
2003 4/.....	3.2	2.7	3.1	3.0	3.7	4.0	5.3	6.6	8.7	8.4
2002.....	3.7	3.1	3.4	2.8	3.9	4.2	5.5	7.6	9.3	7.4
2001.....	3.4	3.2	3.2	3.9	3.7	4.7	5.6	8.0	9.4	7.9
2000.....	3.5	3.4	3.6	3.9	4.4	4.8	5.8	7.9	8.9	8.5
1999.....	3.6	3.2	3.5	3.4	3.8	5.1	6.7	8.1	10.1	8.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified motor vehicle accidents (V09.0,V09.2,V80.3-V80.5,V82.0-V82.1,V87.0-V87.5,V87.7-V87.8,V88.0-V88.5,V88.7-V88.8,V89.0,V89.2)

Black, both sexes

2006 4/.....	5.5	6.8	6.4	7.0	8.2	6.5	6.5	8.5	9.9	5.7
2005 4/.....	6.3	6.2	6.1	7.0	7.5	6.2	8.9	9.9	6.6	7.2
2004 4/.....	5.7	6.8	7.6	5.5	6.1	6.4	6.0	8.5	7.7	*
2003 4/.....	6.7	6.5	6.2	6.1	7.0	6.0	6.7	6.8	9.9	10.8
2002.....	5.4	6.2	5.8	5.7	5.6	6.4	6.0	8.4	9.7	7.1
2001.....	6.4	6.7	5.6	7.2	6.1	6.9	6.9	10.4	7.0	7.1
2000.....	6.7	7.5	7.3	5.9	5.4	7.4	8.7	7.3	8.5	8.8
1999.....	5.4	7.9	6.7	5.9	6.9	7.2	8.3	11.2	13.3	10.2

Black, male

2006 4/.....	8.3	10.4	10.2	10.6	12.5	10.8	10.2	13.4	15.7	*
2005 4/.....	9.7	9.4	9.3	11.1	11.6	10.5	13.6	17.0	*	*
2004 4/.....	8.5	10.3	12.0	8.0	9.5	11.1	9.7	11.8	15.1	*
2003 4/.....	9.2	10.4	9.3	9.5	9.8	8.9	8.5	11.9	*	20.9
2002.....	7.5	8.5	8.2	9.5	8.1	10.2	9.0	16.1	*	21.4
2001.....	9.0	10.8	8.2	11.1	10.0	8.4	10.1	18.1	*	*
2000.....	10.9	10.7	11.2	10.0	8.3	11.7	14.4	11.3	16.9	*
1999.....	7.7	12.0	10.1	9.0	10.6	10.1	8.8	19.9	24.4	23.4

Black, female

2006 4/.....	3.0	3.7	3.2	4.2	4.8	*	*	5.6	*	*
2005 4/.....	3.3	3.4	3.5	3.5	4.3	*	5.6	5.6	*	*
2004 4/.....	3.3	3.7	3.9	3.4	3.4	*	*	6.5	*	*
2003 4/.....	4.4	3.1	3.5	3.3	4.8	3.8	5.5	*	8.5	*
2002.....	3.6	4.2	3.7	2.6	3.6	*	*	*	*	*
2001.....	4.1	3.0	3.4	4.0	*	5.8	4.7	5.8	*	*
2000.....	3.0	4.8	4.0	*	*	4.2	4.9	*	*	*
1999.....	3.4	4.3	3.9	3.5	4.0	5.0	7.9	6.0	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

40-44 years 45-49 years 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85 years and over

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2006.....	0.3	0.3	0.4	0.4	0.3	0.4	0.5	0.5	0.5	0.5
2005.....	0.3	0.3	0.5	0.3	0.3	0.3	0.3	0.5	0.6	*
2004.....	0.3	0.3	0.3	0.3	0.3	0.3	0.4	*	0.5	*
2003.....	0.4	0.4	0.3	0.4	0.4	0.3	0.4	0.5	0.6	0.5
2002.....	0.2	0.3	0.3	0.2	0.3	0.2	0.4	0.4	0.5	*
2001.....	0.4	0.4	0.4	0.3	0.4	0.6	0.5	0.6	0.6	0.6
2000.....	0.5	0.4	0.4	0.4	0.4	0.5	0.5	0.6	0.7	0.7
1999.....	0.6	0.7	0.6	0.6	0.6	0.8	0.7	1.2	1.2	1.2

All races 3/, male

2006.....	0.4	0.4	0.6	0.5	0.5	0.6	0.9	0.8	*	*
2005.....	0.4	0.5	0.7	0.6	0.5	0.5	*	1.0	0.9	*
2004.....	0.4	0.4	0.5	0.5	0.5	0.5	0.6	*	0.9	*
2003.....	0.5	0.6	0.5	0.6	0.6	0.6	0.7	0.7	1.1	*
2002.....	0.3	0.6	0.5	0.4	0.5	0.5	0.6	0.7	1.0	*
2001.....	0.6	0.6	0.6	0.5	0.7	1.0	0.7	0.9	*	*
2000.....	0.7	0.5	0.5	0.5	0.7	0.7	0.8	0.9	1.1	*
1999.....	0.9	1.0	0.9	1.0	0.9	1.1	1.0	1.6	1.8	2.7

All races 3/, female

2006.....	*	*	0.2	0.2	*	*	*	*	*	*
2005.....	0.2	*	0.2	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	*
2003.....	0.2	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	0.3	*	*	*	*	*	0.5	*	*
2000.....	0.2	0.3	0.3	*	*	*	*	*	*	*
1999.....	0.3	0.4	0.2	0.3	*	0.4	0.4	0.9	0.9	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified land transport accidents (V01,V06,V09.1,V09.3,V09.9,V10-V11,V16-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.7-V80.9,V87.9,V88.9,V89.1,V89.3,V89.9)

White, both sexes

2006 4/.....	0.3	0.3	0.4	0.4	0.3	0.4	0.5	0.5	0.5	0.5
2005 4/.....	0.3	0.3	0.4	0.4	0.3	0.3	0.3	0.5	0.6	*
2004 4/.....	0.3	0.3	0.3	0.3	0.4	0.4	0.4	*	0.5	*
2003 4/.....	0.4	0.3	0.3	0.4	0.4	0.3	0.4	0.5	0.6	0.5
2002.....	0.2	0.3	0.3	0.2	0.3	*	0.4	0.3	0.5	*
2001.....	0.3	0.4	0.3	0.3	0.4	0.5	0.5	0.6	0.6	0.5
2000.....	0.4	0.4	0.5	0.3	0.4	0.4	0.4	0.6	0.7	0.7
1999.....	0.6	0.7	0.6	0.7	0.6	0.7	0.6	1.1	1.2	1.2

White, male

2006 4/.....	0.4	0.4	0.6	0.5	0.5	0.6	1.0	0.9	*	*
2005 4/.....	0.3	0.5	0.6	0.6	0.4	*	*	0.9	*	*
2004 4/.....	0.4	0.4	0.4	0.5	0.5	0.5	*	*	1.0	*
2003 4/.....	0.5	0.5	0.5	0.6	0.6	0.6	0.6	*	1.1	*
2002.....	0.3	0.5	0.5	0.4	0.5	*	0.6	*	1.2	*
2001.....	0.5	0.6	0.6	0.5	0.7	0.8	0.7	0.9	*	*
2000.....	0.7	0.5	0.6	0.5	0.8	0.6	0.7	0.9	*	*
1999.....	0.9	1.0	0.9	1.1	0.9	1.2	0.9	1.5	1.8	2.6

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	0.2	*	*	*	*	*	*	*	*
2000.....	0.2	0.2	0.4	*	*	*	*	*	*	*
1999.....	0.3	0.4	*	0.3	*	*	*	0.8	0.9	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
All races 3/, both sexes										
2006.....	1.8	2.5	3.5	4.4	6.0	9.5	16.5	33.5	66.2	151.6
2005.....	1.7	2.4	3.2	3.9	5.9	8.8	16.7	31.9	63.6	147.7
2004.....	1.8	2.6	3.2	4.1	5.8	8.9	16.1	30.3	61.8	143.3
2003.....	1.7	2.3	2.8	3.7	5.3	8.1	14.7	28.7	57.5	135.9
2002.....	1.8	2.3	2.8	3.4	4.9	7.9	13.9	27.1	54.0	130.4
2001.....	1.6	2.5	2.7	3.3	4.9	7.8	12.4	25.2	49.9	121.5
2000.....	1.6	2.0	2.6	3.3	4.6	6.2	12.0	22.1	44.5	112.6
1999.....	1.7	2.3	2.2	3.0	4.6	6.9	11.3	22.3	45.5	110.2
All races 3/, male										
2006.....	2.8	4.0	5.5	6.4	8.5	13.3	20.6	42.4	83.7	181.1
2005.....	2.6	3.7	5.0	5.9	8.5	12.1	21.4	41.0	82.1	174.8
2004.....	2.9	4.1	4.9	6.2	8.1	12.4	21.6	38.4	78.7	174.4
2003.....	2.7	3.6	4.4	5.3	7.3	10.7	18.9	36.9	73.8	165.1
2002.....	2.9	3.6	4.3	5.2	7.1	10.7	18.9	35.1	67.9	158.9
2001.....	2.8	4.2	4.4	5.0	7.1	10.6	16.0	34.0	64.1	155.7
2000.....	2.7	3.4	4.2	5.0	6.9	8.6	15.7	29.4	57.3	145.4
1999.....	2.8	3.8	3.6	4.7	7.0	9.8	15.3	27.9	61.6	142.5
All races 3/, female										
2006.....	0.7	1.1	1.5	2.4	3.7	6.2	13.1	27.0	55.3	137.8
2005.....	0.7	1.2	1.4	2.1	3.6	6.0	13.0	25.4	52.2	135.2
2004.....	0.7	1.1	1.5	2.0	3.6	5.8	11.7	24.5	51.4	129.2
2003.....	0.6	1.1	1.3	2.1	3.5	5.8	11.3	22.8	47.5	122.9
2002.....	0.7	1.0	1.4	1.8	3.0	5.5	9.9	21.4	45.5	118.1
2001.....	0.6	0.9	1.2	1.6	2.9	5.4	9.5	19.0	41.3	107.2
2000.....	0.6	0.8	1.2	1.8	2.6	4.2	9.1	17.1	36.9	99.2
1999.....	0.6	0.8	0.9	1.5	2.4	4.5	8.1	18.4	36.0	97.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
White, both sexes										
2006 4/.....	1.9	2.6	3.5	4.4	6.1	9.7	17.3	35.4	69.9	161.1
2005 4/.....	1.8	2.5	3.2	4.0	6.0	9.1	17.6	33.4	66.6	156.7
2004 4/.....	1.9	2.6	3.2	4.0	5.8	9.2	16.7	32.0	65.3	152.6
2003 4/.....	1.7	2.3	2.7	3.7	5.4	8.2	15.2	30.1	60.0	144.4
2002.....	1.8	2.3	2.8	3.4	5.1	8.1	14.4	28.3	57.1	138.0
2001.....	1.7	2.5	2.8	3.2	4.9	8.0	12.8	26.3	52.4	128.0
2000.....	1.7	2.1	2.5	3.2	4.6	6.2	12.3	23.1	46.6	119.1
1999.....	1.7	2.2	2.1	2.9	4.3	6.9	11.4	23.0	47.7	115.9
White, male										
2006 4/.....	3.0	4.1	5.4	6.4	8.5	13.5	21.4	44.0	88.0	191.2
2005 4/.....	2.8	3.7	5.0	5.9	8.4	12.4	22.4	43.1	84.6	183.8
2004 4/.....	3.1	4.2	4.8	6.1	7.9	13.0	22.0	40.4	82.7	183.7
2003 4/.....	2.8	3.6	4.1	5.3	7.3	10.6	19.5	38.6	76.4	175.1
2002.....	3.0	3.7	4.2	5.1	7.2	10.8	19.3	36.4	71.2	166.4
2001.....	2.8	4.1	4.4	4.9	7.0	10.8	16.5	35.1	66.6	162.3
2000.....	2.7	3.5	3.9	4.8	6.9	8.4	16.2	30.3	59.7	152.8
1999.....	2.7	3.7	3.3	4.4	6.6	9.6	15.3	28.3	64.2	149.1
White, female										
2006 4/.....	0.7	1.1	1.6	2.5	3.8	6.3	14.0	29.1	58.4	147.0
2005 4/.....	0.8	1.3	1.4	2.1	3.7	6.2	13.6	26.3	55.3	144.2
2004 4/.....	0.7	1.1	1.6	2.1	3.8	5.8	12.3	25.9	54.6	138.6
2003 4/.....	0.6	1.0	1.3	2.1	3.6	6.1	11.6	24.0	49.9	130.8
2002.....	0.7	1.0	1.5	1.8	3.1	5.8	10.3	22.6	48.5	125.6
2001.....	0.5	0.9	1.2	1.6	2.9	5.6	9.9	20.0	43.9	113.6
2000.....	0.6	0.7	1.2	1.8	2.6	4.3	9.3	18.1	38.8	105.4
1999.....	0.6	0.8	0.9	1.5	2.2	4.6	8.3	19.3	37.9	102.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Falls (W00-W19)										
Black, both sexes										
2006 4/.....	1.2	2.6	3.8	4.5	5.0	8.9	10.5	16.4	32.9	54.0
2005 4/.....	1.0	2.4	3.9	4.1	5.7	7.2	10.9	20.5	32.5	52.7
2004 4/.....	1.7	2.6	3.5	5.1	6.3	7.9	12.8	16.8	25.5	46.5
2003 4/.....	1.5	3.0	3.6	4.3	5.5	7.6	10.6	13.4	28.2	44.1
2002.....	1.6	2.5	3.2	3.8	4.7	6.1	10.9	14.8	18.1	49.3
2001.....	2.0	2.6	3.3	3.9	4.9	7.9	9.7	16.3	21.9	44.5
2000.....	1.6	2.0	3.6	4.5	5.1	7.1	9.7	11.6	20.3	38.8
1999.....	1.7	3.2	3.6	4.5	6.7	9.0	9.5	13.4	23.1	45.3
Black, male										
2006 4/.....	2.2	4.3	6.4	7.6	8.1	12.9	16.3	25.5	45.9	67.9
2005 4/.....	1.7	4.4	6.4	6.5	9.8	10.9	14.5	23.7	52.7	63.2
2004 4/.....	2.6	4.1	6.0	8.8	11.8	9.7	20.7	23.7	38.0	71.5
2003 4/.....	2.6	4.8	5.8	6.5	8.1	12.8	12.2	18.3	35.4	56.3
2002.....	2.6	4.1	5.6	6.3	7.3	9.2	15.4	22.0	28.3	77.2
2001.....	3.3	4.9	6.0	6.4	7.7	11.9	13.5	24.2	36.1	68.1
2000.....	2.6	3.4	6.3	7.7	8.7	10.7	12.4	18.4	26.1	61.1
1999.....	3.3	5.6	6.0	7.6	10.2	14.8	14.6	20.4	31.4	58.6
Black, female										
2006 4/.....	*	*	*	2.0	*	5.9	6.6	11.0	26.0	48.3
2005 4/.....	*	*	1.8	2.1	*	4.4	8.4	18.7	21.9	48.4
2004 4/.....	*	*	*	*	*	6.7	7.5	12.7	18.9	36.6
2003 4/.....	*	1.4	1.7	2.5	3.5	3.7	9.5	10.6	24.4	39.3
2002.....	*	*	*	*	*	3.7	7.8	10.4	12.7	38.6
2001.....	*	*	*	*	*	4.9	7.2	11.6	14.7	35.8
2000.....	*	*	*	*	*	4.4	8.0	7.4	17.4	30.5
1999.....	*	*	*	*	4.0	4.6	6.1	9.2	19.0	40.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fall on same level (W00-W09,W18)										
All races 3/, both sexes										
2006.....	0.3	0.6	1.0	1.2	1.9	3.0	5.8	12.6	25.3	61.8
2005.....	0.4	0.6	0.9	1.1	1.7	3.0	5.9	12.5	24.4	60.5
2004.....	0.3	0.6	0.7	1.0	1.8	2.7	5.1	10.2	21.7	53.6
2003.....	0.4	0.4	0.8	1.0	1.5	2.3	4.4	9.5	20.5	50.3
2002.....	0.3	0.5	0.7	0.9	1.3	2.2	4.2	8.9	17.8	50.8
2001.....	0.2	0.5	0.6	0.8	1.2	1.9	3.3	6.8	14.4	37.3
2000.....	0.2	0.3	0.5	0.6	1.0	1.4	2.7	5.5	11.9	33.0
1999.....	0.1	0.3	0.3	0.4	0.5	1.1	1.9	4.2	8.2	21.4
All races 3/, male										
2006.....	0.5	0.9	1.5	1.5	2.4	4.0	6.6	16.2	31.9	71.3
2005.....	0.6	0.8	1.4	1.5	2.4	3.8	7.3	15.5	31.5	72.6
2004.....	0.5	0.9	1.2	1.3	2.4	3.6	6.3	12.9	26.8	64.1
2003.....	0.5	0.6	1.1	1.4	1.9	2.7	5.5	12.2	26.8	60.1
2002.....	0.5	0.8	1.0	1.2	1.7	2.9	5.7	11.0	21.6	60.8
2001.....	0.3	0.8	0.8	1.0	1.7	2.4	3.7	9.0	18.5	45.3
2000.....	0.4	0.4	0.8	0.8	1.2	1.7	3.4	6.6	15.2	39.6
1999.....	0.2	0.5	0.5	0.5	0.6	1.5	2.5	5.3	10.5	26.6
All races 3/, female										
2006.....	*	0.3	0.5	1.0	1.4	2.2	5.2	10.0	21.1	57.4
2005.....	0.2	0.4	0.4	0.7	1.2	2.3	4.9	10.4	20.0	54.9
2004.....	0.2	0.3	0.3	0.6	1.2	1.8	4.1	8.2	18.5	48.9
2003.....	0.2	0.3	0.4	0.7	1.1	2.0	3.6	7.6	16.7	45.9
2002.....	*	0.2	0.4	0.7	0.9	1.6	2.9	7.3	15.5	46.4
2001.....	*	0.3	0.4	0.5	0.8	1.4	2.9	5.2	11.9	33.9
2000.....	*	*	0.3	0.5	0.8	1.1	2.2	4.7	10.0	30.4
1999.....	*	*	*	*	0.4	0.8	1.4	3.5	6.8	19.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fall on same level (W00-W09,W18)										
White, both sexes										
2006 4/.....	0.4	0.6	1.0	1.3	1.8	3.2	6.2	13.3	26.6	65.4
2005 4/.....	0.4	0.6	0.9	1.1	1.7	3.0	6.2	13.1	25.5	64.0
2004 4/.....	0.3	0.6	0.8	0.9	1.7	2.8	5.2	10.5	22.9	57.5
2003 4/.....	0.4	0.4	0.7	1.0	1.4	2.4	4.6	10.2	21.6	53.5
2002.....	0.3	0.5	0.7	0.9	1.4	2.3	4.3	9.2	18.8	53.7
2001.....	0.2	0.6	0.6	0.7	1.3	2.0	3.4	6.9	15.1	39.3
2000.....	0.3	0.3	0.5	0.6	1.0	1.4	2.8	5.8	12.5	35.0
1999.....	0.2	0.3	0.3	0.3	0.4	1.1	2.0	4.3	8.4	22.7
White, male										
2006 4/.....	0.6	0.9	1.5	1.6	2.3	4.2	6.9	16.7	33.2	74.4
2005 4/.....	0.6	0.8	1.4	1.4	2.2	3.7	7.5	16.1	32.4	75.5
2004 4/.....	0.5	0.9	1.2	1.3	2.2	4.0	6.3	13.2	28.1	68.2
2003 4/.....	0.5	0.6	1.0	1.3	1.8	2.7	5.8	12.8	28.1	63.5
2002.....	0.5	0.8	1.0	1.1	1.9	3.0	5.9	11.3	22.3	63.3
2001.....	0.3	0.8	0.8	1.0	1.7	2.6	3.9	9.3	19.2	47.7
2000.....	0.4	0.5	0.7	0.8	1.2	1.6	3.6	6.8	15.7	41.9
1999.....	*	0.4	0.4	0.4	0.5	1.4	2.6	5.4	10.8	28.0
White, female										
2006 4/.....	*	0.3	0.5	1.0	1.3	2.3	5.7	10.7	22.4	61.2
2005 4/.....	0.2	0.4	0.4	0.7	1.2	2.4	5.1	10.8	21.2	58.7
2004 4/.....	*	0.3	0.3	0.6	1.2	1.8	4.2	8.5	19.7	52.7
2003 4/.....	0.2	0.3	0.5	0.6	1.1	2.2	3.7	8.3	17.6	49.1
2002.....	*	*	0.5	0.7	1.0	1.7	3.1	7.8	16.7	49.5
2001.....	*	0.3	0.4	0.5	0.8	1.6	2.9	5.3	12.7	35.7
2000.....	*	*	0.3	0.4	0.8	1.1	2.2	5.1	10.5	32.2
1999.....	*	*	*	*	*	0.8	1.4	3.6	7.0	20.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fall from one level to another (W10-W17)										
All races 3/, both sexes										
2006.....	0.8	1.1	1.3	1.5	1.8	2.2	3.3	5.0	8.2	10.1
2005.....	0.8	1.0	1.2	1.3	1.8	2.2	3.2	5.0	7.9	9.4
2004.....	0.8	1.0	1.1	1.4	1.8	2.2	3.4	4.7	7.3	10.6
2003.....	0.7	1.1	1.0	1.1	1.7	2.3	3.2	4.6	7.6	11.4
2002.....	0.8	1.0	0.9	1.3	1.7	2.3	3.4	5.1	7.5	10.9
2001.....	0.8	1.1	1.1	1.3	1.5	2.3	3.0	5.1	7.6	11.3
2000.....	0.9	1.0	1.0	1.2	1.5	1.9	3.0	4.2	6.5	8.8
1999.....	0.9	1.0	0.9	1.3	1.9	2.3	2.9	4.8	7.3	10.4
All races 3/, male										
2006.....	1.5	1.8	2.2	2.4	2.9	3.7	5.2	7.4	13.2	17.5
2005.....	1.5	1.8	2.0	2.1	3.1	3.5	5.0	8.1	12.3	14.7
2004.....	1.5	1.7	2.0	2.4	3.0	3.6	5.5	6.7	11.2	16.4
2003.....	1.3	1.8	1.7	1.9	2.7	3.6	5.1	7.3	11.3	18.1
2002.....	1.5	1.7	1.5	2.2	2.8	3.5	5.3	8.2	12.0	17.4
2001.....	1.4	2.0	1.9	2.2	2.6	3.9	4.6	8.4	11.4	18.2
2000.....	1.5	1.7	1.8	2.1	2.4	2.9	4.9	7.0	10.8	16.4
1999.....	1.5	1.7	1.5	2.1	3.1	4.0	4.2	6.9	12.7	17.8
All races 3/, female										
2006.....	0.2	0.4	0.5	0.5	0.8	0.8	1.8	3.2	5.1	6.7
2005.....	0.2	0.3	0.4	0.5	0.7	1.0	1.7	2.6	5.2	7.0
2004.....	0.2	0.3	0.3	0.5	0.7	1.0	1.7	3.2	5.0	7.9
2003.....	*	0.3	0.3	0.4	0.8	1.2	1.6	2.8	5.2	8.5
2002.....	0.2	0.2	0.4	0.4	0.7	1.2	1.9	3.0	4.8	8.1
2001.....	*	0.2	0.3	0.4	0.5	1.0	1.8	2.8	5.3	8.3
2000.....	0.2	0.3	0.3	0.5	0.6	1.0	1.5	2.2	3.9	5.7
1999.....	0.2	0.2	0.2	0.5	0.7	0.9	1.8	3.4	4.2	7.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Fall from one level to another (W10-W17)										
White, both sexes										
2006 4/.....	0.9	1.2	1.3	1.5	1.9	2.2	3.5	5.3	8.8	10.6
2005 4/.....	0.9	1.1	1.2	1.4	1.9	2.2	3.5	5.2	8.3	10.0
2004 4/.....	0.9	1.0	1.2	1.5	1.9	2.4	3.6	5.0	7.7	11.2
2003 4/.....	0.7	1.0	1.0	1.2	1.8	2.3	3.3	4.9	7.9	12.1
2002.....	0.9	1.0	1.0	1.3	1.8	2.4	3.6	5.4	7.9	11.2
2001.....	0.8	1.1	1.1	1.3	1.5	2.4	3.2	5.3	8.1	11.7
2000.....	0.9	1.0	1.0	1.3	1.5	1.8	3.1	4.4	6.8	9.2
1999.....	0.9	1.0	0.8	1.3	2.0	2.4	3.0	5.0	7.7	10.9
White, male										
2006 4/.....	1.5	1.9	2.2	2.5	3.0	3.8	5.5	7.9	14.1	18.4
2005 4/.....	1.6	1.8	2.0	2.2	3.2	3.7	5.4	8.6	12.9	15.5
2004 4/.....	1.5	1.8	2.0	2.4	3.0	3.9	5.7	7.2	11.7	17.5
2003 4/.....	1.3	1.8	1.7	2.0	2.8	3.4	5.2	7.5	11.9	19.4
2002.....	1.6	1.8	1.6	2.2	3.0	3.7	5.6	8.5	12.8	18.1
2001.....	1.5	1.9	1.9	2.2	2.5	4.0	4.8	8.6	12.0	19.0
2000.....	1.6	1.8	1.7	2.1	2.6	2.9	5.0	7.3	11.4	17.0
1999.....	1.5	1.8	1.5	2.1	3.3	4.1	4.2	7.0	13.1	19.2
White, female										
2006 4/.....	*	0.4	0.5	0.6	0.8	0.9	1.9	3.4	5.5	6.9
2005 4/.....	0.2	0.4	0.4	0.6	0.8	1.0	1.9	2.7	5.4	7.5
2004 4/.....	*	0.3	0.3	0.5	0.8	1.0	1.9	3.3	5.2	8.3
2003 4/.....	*	0.3	0.3	0.4	0.8	1.3	1.7	2.9	5.4	8.8
2002.....	*	0.2	0.4	0.4	0.7	1.2	1.9	3.1	4.9	8.2
2001.....	*	0.3	0.3	0.4	0.6	1.1	1.9	3.0	5.7	8.7
2000.....	0.2	0.3	0.3	0.5	0.6	0.9	1.6	2.4	4.1	6.0
1999.....	0.2	*	*	0.5	0.8	1.0	2.0	3.6	4.5	7.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified fall (W19)										
All races 3/, both sexes										
2006.....	0.6	0.8	1.2	1.7	2.3	4.3	7.3	15.9	32.7	79.7
2005.....	0.5	0.8	1.1	1.5	2.4	3.7	7.6	14.4	31.3	77.8
2004.....	0.7	1.0	1.3	1.7	2.2	4.0	7.6	15.5	32.8	79.1
2003.....	0.6	0.8	1.1	1.5	2.1	3.5	7.1	14.5	29.4	74.2
2002.....	0.6	0.8	1.2	1.3	2.0	3.5	6.3	13.1	28.6	68.8
2001.....	0.6	0.9	1.1	1.3	2.1	3.6	6.1	13.4	27.9	73.0
2000.....	0.5	0.8	1.1	1.4	2.2	3.0	6.3	12.5	26.1	70.7
1999.....	0.7	1.0	1.0	1.4	2.2	3.5	6.5	13.2	30.0	78.4
All races 3/, male										
2006.....	0.8	1.2	1.8	2.5	3.1	5.6	8.8	18.8	38.6	92.3
2005.....	0.6	1.1	1.6	2.3	3.1	4.7	9.1	17.4	38.3	87.5
2004.....	1.0	1.5	1.7	2.5	2.8	5.2	9.8	18.8	40.7	93.9
2003.....	0.9	1.2	1.6	2.0	2.7	4.5	8.3	17.5	35.6	86.9
2002.....	0.9	1.1	1.8	1.9	2.6	4.3	7.8	15.9	34.3	80.7
2001.....	1.0	1.3	1.7	1.8	2.8	4.3	7.6	16.6	34.2	92.1
2000.....	0.8	1.2	1.6	2.1	3.3	4.0	7.5	15.9	31.3	89.4
1999.....	1.0	1.6	1.5	2.1	3.2	4.3	8.5	15.7	38.4	98.1
All races 3/, female										
2006.....	0.3	0.4	0.6	0.9	1.6	3.2	6.1	13.8	29.1	73.7
2005.....	0.3	0.5	0.6	0.8	1.7	2.7	6.4	12.3	27.0	73.3
2004.....	0.3	0.5	0.9	0.9	1.6	3.0	5.9	13.1	27.9	72.4
2003.....	0.3	0.4	0.6	1.0	1.6	2.6	6.1	12.3	25.6	68.5
2002.....	0.3	0.6	0.6	0.7	1.4	2.8	5.1	11.2	25.2	63.6
2001.....	0.3	0.4	0.5	0.7	1.5	3.0	4.8	11.1	24.2	64.9
2000.....	0.3	0.3	0.6	0.8	1.2	2.2	5.5	10.1	23.1	63.1
1999.....	0.3	0.4	0.5	0.8	1.3	2.7	4.9	11.5	25.0	70.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified fall (W19)										
White, both sexes										
2006 4/.....	0.6	0.9	1.2	1.6	2.4	4.3	7.6	16.9	34.5	85.1
2005 4/.....	0.5	0.8	1.1	1.5	2.4	3.8	7.9	15.1	32.8	82.7
2004 4/.....	0.7	1.0	1.3	1.6	2.2	4.0	7.9	16.5	34.7	83.9
2003 4/.....	0.6	0.8	1.0	1.5	2.2	3.5	7.3	15.1	30.5	78.8
2002.....	0.6	0.8	1.1	1.2	1.9	3.4	6.4	13.7	30.4	73.0
2001.....	0.6	0.9	1.1	1.2	2.1	3.6	6.3	14.0	29.2	77.0
2000.....	0.5	0.8	1.0	1.4	2.1	3.0	6.4	12.9	27.3	74.9
1999.....	0.6	0.9	1.0	1.3	1.9	3.4	6.4	13.7	31.6	82.3
White, male										
2006 4/.....	0.9	1.3	1.8	2.3	3.2	5.5	9.0	19.4	40.7	98.4
2005 4/.....	0.7	1.0	1.6	2.3	3.0	4.9	9.5	18.3	39.3	92.7
2004 4/.....	1.0	1.5	1.6	2.4	2.7	5.2	10.0	19.9	42.9	98.0
2003 4/.....	0.9	1.2	1.5	2.0	2.7	4.5	8.5	18.3	36.4	92.2
2002.....	0.9	1.1	1.6	1.8	2.3	4.1	7.9	16.6	36.1	85.1
2001.....	1.0	1.3	1.7	1.7	2.8	4.2	7.9	17.3	35.3	95.7
2000.....	0.7	1.3	1.5	1.9	3.1	3.9	7.6	16.2	32.6	94.0
1999.....	1.0	1.5	1.4	1.8	2.8	4.1	8.4	15.9	40.3	101.9
White, female										
2006 4/.....	0.4	0.5	0.5	0.9	1.6	3.1	6.4	14.9	30.5	78.9
2005 4/.....	0.3	0.5	0.6	0.8	1.7	2.9	6.6	12.7	28.7	78.1
2004 4/.....	0.3	0.5	0.9	0.9	1.7	3.0	6.2	14.0	29.7	77.6
2003 4/.....	0.3	0.5	0.5	1.0	1.7	2.6	6.3	12.7	26.9	72.8
2002.....	0.3	0.6	0.6	0.7	1.4	2.9	5.3	11.7	27.0	67.8
2001.....	0.2	0.4	0.5	0.7	1.5	3.0	5.0	11.7	25.5	69.2
2000.....	0.3	0.3	0.6	0.8	1.2	2.2	5.5	10.5	24.2	67.2
1999.....	0.3	0.3	0.5	0.7	1.1	2.8	4.8	12.1	26.4	74.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Unspecified fall (W19)										
Black, both sexes										
2006 4/.....	*	0.7	1.6	2.3	2.2	5.0	5.9	9.4	17.5	27.8
2005 4/.....	*	1.0	1.5	1.9	2.9	2.6	5.3	10.7	19.2	31.5
2004 4/.....	*	1.0	1.6	2.4	2.5	5.0	6.3	6.6	13.9	30.2
2003 4/.....	*	1.2	1.7	2.0	2.5	3.9	6.5	8.3	17.5	28.6
2002.....	*	1.1	2.0	1.8	3.0	3.6	5.6	7.6	10.5	25.2
2001.....	0.8	1.0	1.2	1.8	2.5	4.8	4.7	8.9	13.8	26.4
2000.....	0.9	0.9	1.5	2.1	2.8	3.8	6.3	8.0	13.3	25.7
1999.....	0.8	1.9	1.6	2.6	4.5	4.9	6.4	8.6	15.0	33.5
Black, male										
2006 4/.....	*	*	2.4	4.1	*	6.1	8.3	13.9	19.2	33.1
2005 4/.....	*	1.8	2.2	2.8	4.7	*	6.3	10.3	31.6	36.3
2004 4/.....	*	*	2.5	4.4	4.7	6.5	10.0	9.6	20.8	48.7
2003 4/.....	*	2.0	2.5	2.9	*	5.2	7.9	*	23.6	36.5
2002.....	*	1.8	3.7	3.0	5.1	5.2	8.0	*	*	38.6
2001.....	*	1.9	*	*	*	6.1	*	12.5	20.9	46.5
2000.....	*	*	2.7	3.6	5.2	5.2	7.4	11.8	18.5	38.1
1999.....	*	3.2	3.0	4.4	6.8	7.5	9.2	12.8	19.2	46.9
Black, female										
2006 4/.....	*	*	*	*	*	4.2	4.2	6.7	16.6	25.6
2005 4/.....	*	*	*	*	*	*	4.5	10.9	12.6	29.6
2004 4/.....	*	*	*	*	*	3.8	*	*	10.2	22.8
2003 4/.....	*	*	*	*	*	*	5.5	8.7	14.3	25.5
2002.....	*	*	*	*	*	*	*	7.1	8.4	20.1
2001.....	*	*	*	*	*	3.8	*	6.7	10.2	18.9
2000.....	*	*	*	*	*	*	5.5	5.7	10.6	21.0
1999.....	*	*	*	*	*	*	4.5	6.0	13.0	28.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental drowning and submersion (W65-W74)

All races 3/, both sexes

2006.....	1.1	1.2	1.0	1.0	0.9	1.1	1.2	1.6	1.4	1.4
2005.....	1.3	1.2	1.1	0.9	0.9	0.9	1.2	1.4	1.4	1.8
2004.....	1.0	1.1	1.0	0.9	0.9	1.1	1.0	1.0	1.5	1.6
2003.....	1.1	1.0	1.0	1.0	0.9	1.0	1.2	1.3	1.3	1.8
2002.....	1.1	1.0	0.9	0.8	0.8	0.8	1.0	1.2	1.5	1.7
2001.....	1.1	1.0	0.8	0.8	0.9	0.8	0.9	1.1	1.4	1.7
2000.....	1.0	0.9	1.0	0.9	0.9	0.8	1.1	1.0	1.7	1.5
1999.....	1.0	1.0	1.0	0.9	0.9	0.9	1.1	1.2	1.4	1.6

All races 3/, male

2006.....	1.8	2.0	1.6	1.6	1.4	1.6	1.9	2.5	2.3	2.4
2005.....	2.0	2.0	1.7	1.5	1.5	1.4	1.8	2.1	2.5	3.9
2004.....	1.7	1.8	1.7	1.5	1.5	1.7	1.6	1.7	2.8	2.9
2003.....	1.8	1.6	1.6	1.6	1.5	1.6	1.9	1.8	2.3	3.3
2002.....	1.9	1.6	1.5	1.3	1.2	1.4	1.8	1.8	2.5	3.6
2001.....	1.7	1.8	1.3	1.2	1.4	1.4	1.6	1.8	2.4	3.5
2000.....	1.7	1.6	1.7	1.6	1.5	1.4	1.8	1.2	2.7	2.8
1999.....	1.7	1.7	1.8	1.5	1.5	1.5	1.7	2.1	2.7	2.3

All races 3/, female

2006.....	0.5	0.5	0.4	0.5	0.4	0.7	0.6	0.9	0.8	0.9
2005.....	0.5	0.5	0.4	0.3	0.3	0.5	0.7	0.8	0.7	0.9
2004.....	0.4	0.4	0.5	0.4	0.5	0.5	0.5	0.5	0.7	1.0
2003.....	0.3	0.3	0.4	0.3	0.4	0.6	0.6	0.8	0.6	1.1
2002.....	0.4	0.4	0.3	0.3	0.4	*	0.4	0.7	0.9	0.9
2001.....	0.4	0.3	0.3	0.4	0.4	*	*	0.7	0.8	1.0
2000.....	0.3	0.3	0.3	*	0.4	*	0.6	0.8	1.1	1.0
1999.....	0.3	0.3	0.3	0.4	0.4	0.4	0.7	0.6	0.7	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental drowning and submersion (W65-W74)										
White, both sexes										
2006 4/.....	1.2	1.3	1.0	1.0	0.9	1.1	1.1	1.5	1.5	1.4
2005 4/.....	1.3	1.2	1.0	0.9	0.9	0.9	1.2	1.3	1.2	1.7
2004 4/.....	1.0	1.1	1.0	0.9	0.9	1.1	0.9	1.0	1.4	1.5
2003 4/.....	1.0	1.0	1.0	1.0	0.9	1.1	1.1	1.2	1.3	1.6
2002.....	1.2	0.9	0.8	0.8	0.8	0.8	0.9	1.1	1.4	1.6
2001.....	1.0	1.0	0.8	0.7	0.9	0.8	0.9	1.1	1.3	1.7
2000.....	1.0	0.9	0.9	0.9	0.9	0.8	1.1	0.9	1.8	1.5
1999.....	1.0	1.1	1.0	0.9	0.9	0.9	1.0	1.1	1.4	1.6
White, male										
2006 4/.....	1.8	2.0	1.6	1.6	1.3	1.7	1.9	2.5	2.5	2.6
2005 4/.....	2.0	2.0	1.6	1.5	1.4	1.3	1.7	2.1	2.3	3.9
2004 4/.....	1.7	1.8	1.6	1.4	1.4	1.8	1.4	1.6	2.8	2.8
2003 4/.....	1.8	1.5	1.5	1.6	1.4	1.7	1.9	1.8	2.4	3.3
2002.....	1.9	1.4	1.4	1.3	1.2	1.4	1.6	1.8	2.4	3.7
2001.....	1.6	1.7	1.3	1.1	1.3	1.4	1.6	1.8	2.3	3.5
2000.....	1.6	1.5	1.5	1.6	1.6	1.4	1.7	1.1	3.0	2.7
1999.....	1.7	1.8	1.7	1.4	1.4	1.4	1.5	1.9	2.7	2.4
White, female										
2006 4/.....	0.5	0.5	0.4	0.5	0.5	0.7	0.5	0.8	0.8	0.8
2005 4/.....	0.6	0.5	0.5	0.3	0.4	0.4	0.7	0.7	*	0.8
2004 4/.....	0.3	0.4	0.4	0.4	0.5	*	*	*	*	0.9
2003 4/.....	0.3	0.4	0.4	0.4	0.4	0.6	0.6	0.8	0.7	0.9
2002.....	0.4	0.4	0.3	0.3	0.4	*	*	0.6	0.8	0.7
2001.....	0.4	0.3	0.3	0.4	0.5	*	*	0.7	0.7	1.0
2000.....	0.3	0.3	0.3	*	*	*	0.6	0.7	1.0	1.0
1999.....	0.4	0.4	0.3	0.4	*	*	0.7	0.5	0.7	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

All races 3/, both sexes

2006.....	0.5	0.7	0.8	1.0	1.6	2.6	3.5	6.6	10.9	26.8
2005.....	0.5	0.7	0.9	1.1	1.7	2.4	3.7	6.5	12.2	27.6
2004.....	0.5	0.7	0.9	1.1	1.6	2.1	3.9	7.1	13.5	29.9
2003.....	0.5	0.7	0.9	1.2	1.7	2.4	3.9	6.6	13.0	27.4
2002.....	0.4	0.5	0.8	1.1	1.3	2.6	3.6	7.3	12.6	28.4
2001.....	0.5	0.8	1.0	1.2	1.5	2.3	3.6	7.2	12.8	30.0
2000.....	0.4	0.7	0.9	1.2	1.6	2.3	4.0	7.7	13.4	32.6
1999.....	0.5	0.5	0.8	1.0	1.4	2.2	4.0	6.6	12.1	30.3

All races 3/, male

2006.....	0.6	0.9	1.1	1.2	2.1	3.2	4.3	8.0	14.7	36.5
2005.....	0.5	0.9	1.0	1.4	2.1	2.9	4.8	7.6	15.4	35.8
2004.....	0.6	0.9	1.0	1.4	1.9	2.6	4.7	9.5	17.0	39.0
2003.....	0.6	0.8	0.9	1.4	2.3	2.8	4.6	9.0	17.9	37.5
2002.....	0.6	0.7	0.9	1.5	1.7	3.4	4.4	9.6	16.0	37.0
2001.....	0.6	1.0	1.2	1.6	1.6	2.9	4.3	9.2	17.5	39.8
2000.....	0.5	1.0	1.2	1.4	1.9	2.8	4.5	9.7	18.9	41.0
1999.....	0.6	0.6	1.2	1.1	1.6	2.5	5.2	7.9	15.9	39.2

All races 3/, female

2006.....	0.4	0.5	0.6	0.8	1.1	2.0	3.0	5.6	8.6	22.2
2005.....	0.5	0.5	0.8	0.9	1.3	2.0	2.9	5.7	10.3	23.9
2004.....	0.4	0.5	0.8	0.8	1.3	1.6	3.2	5.3	11.3	25.8
2003.....	0.4	0.6	0.8	0.9	1.1	2.0	3.3	5.0	10.1	23.0
2002.....	0.3	0.4	0.6	0.8	1.0	2.0	2.9	5.6	10.6	24.6
2001.....	0.3	0.6	0.8	0.8	1.3	1.8	3.1	5.8	9.9	26.0
2000.....	0.4	0.4	0.6	0.9	1.4	1.8	3.5	6.3	10.1	29.1
1999.....	0.3	0.5	0.4	0.8	1.2	2.0	3.0	5.6	9.9	26.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

White, both sexes

2006 4/.....	0.5	0.7	0.8	1.0	1.6	2.5	3.3	6.4	11.1	27.1
2005 4/.....	0.4	0.7	0.9	1.1	1.7	2.3	3.6	6.5	11.9	27.7
2004 4/.....	0.5	0.7	0.9	1.1	1.5	1.9	3.6	6.9	13.4	30.1
2003 4/.....	0.5	0.7	0.8	1.1	1.6	2.3	3.7	6.6	12.9	27.9
2002.....	0.4	0.5	0.7	1.1	1.2	2.3	3.4	7.1	12.5	28.3
2001.....	0.4	0.7	1.0	1.1	1.4	2.3	3.3	6.8	12.4	29.7
2000.....	0.5	0.7	0.8	1.1	1.5	2.1	3.7	7.6	13.3	32.5
1999.....	0.5	0.5	0.8	0.9	1.2	2.1	3.7	6.2	11.7	30.4

White, male

2006 4/.....	0.6	0.9	1.1	1.2	2.1	3.0	4.0	7.9	15.1	37.4
2005 4/.....	0.5	0.9	1.0	1.3	2.0	2.8	4.5	7.5	15.0	36.4
2004 4/.....	0.6	0.9	1.0	1.4	1.8	2.4	4.4	9.5	16.8	39.2
2003 4/.....	0.5	0.8	0.9	1.3	2.1	2.8	4.4	8.9	17.5	38.2
2002.....	0.6	0.7	0.8	1.4	1.6	2.9	4.3	9.4	15.6	37.2
2001.....	0.6	0.9	1.1	1.3	1.5	2.8	3.9	8.6	16.4	39.3
2000.....	0.6	0.9	1.1	1.4	1.8	2.5	4.2	9.5	18.4	40.2
1999.....	0.6	0.6	1.1	1.0	1.3	2.4	4.9	7.3	15.4	39.7

White, female

2006 4/.....	0.4	0.5	0.5	0.8	1.1	2.0	2.7	5.3	8.5	22.2
2005 4/.....	0.4	0.5	0.7	0.8	1.4	1.9	2.8	5.8	9.9	23.6
2004 4/.....	0.5	0.6	0.9	0.8	1.2	1.5	3.0	5.0	11.3	26.0
2003 4/.....	0.4	0.6	0.7	0.9	1.1	1.8	3.1	5.0	10.0	23.3
2002.....	0.3	0.4	0.6	0.8	0.9	1.8	2.7	5.6	10.6	24.5
2001.....	0.3	0.6	0.9	0.8	1.2	1.8	2.9	5.6	10.0	25.7
2000.....	0.4	0.5	0.6	0.9	1.3	1.7	3.3	6.3	10.2	29.4
1999.....	0.3	0.5	0.4	0.7	1.0	1.9	2.7	5.3	9.5	26.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental inhalation and ingestion of food or other objects causing obstruction of respiratory tract (W78-W80)

Black, both sexes

2006 4/.....	*	0.9	1.3	1.3	2.3	3.7	6.3	10.4	10.9	27.0
2005 4/.....	0.8	0.9	1.4	2.0	2.1	3.9	6.0	9.2	18.0	30.2
2004 4/.....	*	*	1.3	1.5	2.4	4.0	6.7	10.3	17.3	30.5
2003 4/.....	0.8	*	1.4	1.5	2.9	3.7	6.5	6.6	16.5	24.5
2002.....	*	*	1.4	1.9	2.2	5.8	7.1	11.0	16.0	33.0
2001.....	*	1.3	1.0	2.3	2.8	3.6	7.5	11.5	18.7	36.8
2000.....	*	0.8	1.5	1.9	2.6	5.0	7.2	10.8	16.7	35.3
1999.....	*	1.0	1.2	2.1	3.0	3.6	7.7	12.3	18.2	33.2

Black, male

2006 4/.....	*	*	*	*	*	4.9	7.4	11.6	*	33.1
2005 4/.....	*	*	*	2.6	3.6	*	8.2	13.0	23.2	32.5
2004 4/.....	*	*	*	*	*	5.6	8.7	11.4	25.8	42.7
2003 4/.....	*	*	*	*	4.4	*	7.9	10.5	25.8	30.2
2002.....	*	*	2.2	*	*	8.7	8.0	16.5	23.7	41.8
2001.....	*	*	*	3.6	*	*	9.8	15.8	36.1	51.1
2000.....	*	*	2.4	*	*	6.5	8.4	15.6	27.8	51.9
1999.....	*	*	*	*	*	*	10.5	17.6	25.3	43.4

Black, female

2006 4/.....	*	*	*	*	*	*	5.5	9.6	9.8	24.5
2005 4/.....	*	*	*	*	*	3.7	4.5	6.9	15.2	29.3
2004 4/.....	*	*	*	*	*	*	5.3	9.7	12.9	25.6
2003 4/.....	*	*	*	*	*	3.7	5.5	*	11.6	22.3
2002.....	*	*	*	*	*	3.7	6.5	7.7	11.9	29.6
2001.....	*	*	*	*	*	*	6.1	8.9	9.8	31.6
2000.....	*	*	*	*	*	3.8	6.4	8.0	11.0	29.2
1999.....	*	*	*	*	*	*	5.9	9.2	14.7	29.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other accidental and unspecified threats to breathing (W75-W77,W81-W84)										
All races 3/, both sexes										
2006.....	0.4	0.4	0.3	0.2	0.2	0.2	*	0.4	*	0.7
2005.....	0.3	0.3	0.4	0.3	0.3	0.2	0.3	0.4	0.4	0.5
2004.....	0.3	0.3	0.3	0.2	0.2	0.2	*	0.3	*	0.6
2003.....	0.3	0.3	0.3	0.2	0.2	*	0.3	*	0.4	0.8
2002.....	0.3	0.3	0.3	0.3	0.2	0.3	0.2	0.4	*	1.0
2001.....	0.4	0.4	0.2	0.2	0.2	0.2	0.4	0.4	0.4	0.9
2000.....	0.3	0.2	0.3	0.4	0.3	0.3	0.4	0.5	0.5	1.2
1999.....	0.4	0.4	0.3	0.3	0.3	0.5	0.8	1.4	1.8	4.7
All races 3/, male										
2006.....	0.6	0.6	0.5	0.3	0.3	*	*	*	*	*
2005.....	0.6	0.5	0.6	0.5	0.4	*	0.5	*	*	*
2004.....	0.6	0.5	0.5	0.3	0.3	*	*	*	*	*
2003.....	0.5	0.4	0.5	0.4	0.3	*	*	*	*	*
2002.....	0.5	0.5	0.5	0.4	*	*	*	*	*	*
2001.....	0.6	0.6	0.4	0.3	*	*	*	*	*	*
2000.....	0.6	0.4	0.6	0.6	0.5	*	*	*	*	*
1999.....	0.6	0.6	0.6	0.6	*	0.7	1.1	1.9	1.8	5.4
All races 3/, female										
2006.....	*	*	*	*	*	*	*	*	*	0.7
2005.....	*	*	*	*	*	*	*	*	*	*
2004.....	*	*	*	*	*	*	*	*	*	0.6
2003.....	*	*	*	*	*	*	*	*	*	0.7
2002.....	*	*	*	*	*	*	*	*	*	1.1
2001.....	*	*	*	*	*	*	*	*	*	0.9
2000.....	*	*	*	*	*	*	*	*	*	1.1
1999.....	*	*	*	*	*	*	0.6	1.1	1.7	4.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other accidental and unspecified threats to breathing (W75-W77,W81-W84)

White, both sexes

2006 4/.....	0.4	0.4	0.3	0.2	0.2	*	*	0.4	*	0.7
2005 4/.....	0.4	0.3	0.4	0.3	0.3	0.2	0.3	0.4	0.5	0.5
2004 4/.....	0.4	0.4	0.4	0.2	0.2	0.2	*	*	*	0.6
2003 4/.....	0.4	0.3	0.3	0.2	0.2	*	0.3	*	0.4	0.8
2002.....	0.3	0.3	0.3	0.3	0.3	0.3	*	0.4	*	1.0
2001.....	0.4	0.4	0.3	0.2	*	0.3	0.4	0.5	0.4	0.9
2000.....	0.4	0.3	0.4	0.4	0.3	0.3	0.3	0.5	0.6	1.2
1999.....	0.4	0.4	0.4	0.3	0.3	0.5	0.8	1.3	1.8	4.9

White, male

2006 4/.....	0.6	0.7	0.5	0.3	*	*	*	*	*	*
2005 4/.....	0.7	0.5	0.6	0.6	0.5	*	*	*	*	*
2004 4/.....	0.6	0.6	0.6	0.3	*	*	*	*	*	*
2003 4/.....	0.6	0.5	0.5	0.4	*	*	*	*	*	*
2002.....	0.5	0.6	0.6	0.4	*	*	*	*	*	*
2001.....	0.7	0.7	0.5	0.3	*	*	*	*	*	*
2000.....	0.6	0.4	0.6	0.6	0.6	*	*	*	*	*
1999.....	0.7	0.7	0.6	0.5	*	0.7	0.9	1.8	1.8	5.7

White, female

2006 4/.....	*	*	*	*	*	*	*	*	*	0.7
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	*	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	0.8
2002.....	*	*	*	*	*	*	*	*	*	1.1
2001.....	*	*	*	*	*	*	*	*	*	0.9
2000.....	*	*	*	*	*	*	*	*	*	1.1
1999.....	*	*	*	*	*	*	0.7	1.0	1.7	4.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes										
2006.....	0.7	1.1	1.1	1.4	1.5	2.1	2.2	3.2	3.8	4.3
2005.....	0.9	1.2	1.1	1.1	1.6	1.9	2.3	3.1	4.2	5.0
2004.....	0.7	1.1	1.2	1.2	1.6	1.8	2.0	2.9	4.4	5.6
2003.....	1.0	1.2	1.3	1.3	1.6	2.0	2.6	3.0	4.3	5.4
2002.....	0.9	1.0	1.3	1.2	1.4	1.8	2.4	2.8	4.2	4.8
2001.....	1.1	1.0	1.1	1.5	1.5	2.0	2.5	3.0	3.8	5.7
2000.....	1.0	1.1	1.3	1.4	1.7	2.0	2.4	3.5	4.3	6.2
1999.....	1.0	1.0	1.0	1.2	1.5	2.0	2.5	3.5	4.7	5.9
All races 3/, male										
2006.....	0.9	1.4	1.5	1.9	1.9	2.7	2.5	4.0	5.1	6.3
2005.....	1.2	1.5	1.6	1.5	2.0	2.5	2.8	3.9	5.6	7.0
2004.....	1.0	1.5	1.8	1.6	2.1	2.6	2.3	3.5	5.9	7.2
2003.....	1.2	1.7	1.6	1.8	2.0	2.5	3.4	4.0	5.1	7.4
2002.....	1.3	1.4	1.6	1.5	2.0	2.6	3.2	3.5	6.1	7.0
2001.....	1.6	1.5	1.5	2.1	1.8	2.5	3.0	3.5	5.1	8.3
2000.....	1.2	1.5	1.8	1.9	2.2	2.4	3.2	4.5	5.9	9.4
1999.....	1.5	1.5	1.3	1.6	1.9	2.5	3.4	4.3	6.6	9.3
All races 3/, female										
2006.....	0.5	0.8	0.8	1.0	1.1	1.5	2.0	2.6	2.9	3.4
2005.....	0.5	0.9	0.7	0.8	1.2	1.4	1.9	2.5	3.4	4.0
2004.....	0.5	0.8	0.7	0.9	1.3	1.2	1.6	2.5	3.5	4.9
2003.....	0.7	0.8	1.1	0.8	1.1	1.6	1.9	2.3	3.9	4.5
2002.....	0.6	0.6	0.9	0.9	0.8	1.1	1.8	2.4	3.0	3.9
2001.....	0.7	0.6	0.7	1.0	1.2	1.5	2.1	2.7	3.1	4.6
2000.....	0.7	0.7	0.7	0.9	1.2	1.6	1.8	2.8	3.4	4.9
1999.....	0.5	0.5	0.7	0.9	1.2	1.6	1.8	3.0	3.6	4.5

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental exposure to smoke, fire and flames (X00-X09)										
White, both sexes										
2006 4/.....	0.7	1.0	1.1	1.3	1.4	1.9	2.1	2.8	3.3	3.7
2005 4/.....	0.8	1.2	1.0	0.8	1.4	1.7	2.1	2.7	3.8	4.2
2004 4/.....	0.7	1.0	1.1	1.1	1.4	1.6	1.6	2.5	3.9	4.9
2003 4/.....	1.0	1.2	1.2	1.2	1.4	1.8	2.2	2.6	3.7	4.7
2002.....	0.9	0.9	1.1	1.1	1.1	1.6	2.1	2.5	3.7	4.0
2001.....	1.0	0.9	1.0	1.3	1.3	1.7	2.1	2.6	3.2	4.8
2000.....	0.9	0.9	1.2	1.1	1.4	1.7	2.0	2.8	3.5	5.1
1999.....	0.9	0.9	0.9	1.1	1.2	1.6	2.0	2.8	3.9	4.7
White, male										
2006 4/.....	0.9	1.2	1.4	1.8	1.8	2.4	2.4	3.3	4.7	6.0
2005 4/.....	1.1	1.5	1.4	1.1	1.7	2.1	2.6	3.4	5.1	6.0
2004 4/.....	0.9	1.2	1.6	1.3	1.9	2.2	1.9	2.9	5.4	6.5
2003 4/.....	1.3	1.5	1.5	1.6	1.9	2.2	3.0	3.2	4.2	6.7
2002.....	1.3	1.3	1.4	1.4	1.6	2.2	2.7	3.1	5.8	5.8
2001.....	1.4	1.2	1.2	1.8	1.5	2.1	2.4	2.8	4.4	7.0
2000.....	1.1	1.3	1.7	1.6	1.8	2.0	2.7	3.6	4.8	8.2
1999.....	1.4	1.3	1.2	1.4	1.7	2.0	2.6	3.3	5.8	8.0
White, female										
2006 4/.....	0.5	0.8	0.7	1.0	1.0	1.4	1.8	2.3	2.4	2.7
2005 4/.....	0.6	0.9	0.7	0.6	1.2	1.3	1.8	2.2	2.9	3.4
2004 4/.....	0.4	0.7	0.7	0.9	1.0	1.1	1.3	2.2	2.9	4.2
2003 4/.....	0.7	0.8	0.9	0.8	1.0	1.5	1.6	2.2	3.4	3.8
2002.....	0.6	0.6	0.9	0.8	0.6	1.0	1.6	2.1	2.5	3.2
2001.....	0.6	0.5	0.7	0.8	1.1	1.3	1.9	2.4	2.5	3.8
2000.....	0.6	0.6	0.7	0.7	1.1	1.4	1.5	2.3	2.7	3.9
1999.....	0.5	0.5	0.6	0.9	0.8	1.3	1.5	2.5	2.8	3.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Black, both sexes

2006 4/.....	1.0	1.9	1.9	2.7	2.6	4.3	4.1	9.0	9.9	12.2
2005 4/.....	1.2	1.7	2.2	3.7	3.2	4.4	4.2	8.0	11.2	14.4
2004 4/.....	1.4	2.6	2.2	2.8	4.2	4.0	5.9	8.0	12.1	14.9
2003 4/.....	1.4	2.3	2.9	2.8	2.9	4.4	6.4	8.7	12.4	14.9
2002.....	1.1	1.6	2.8	2.6	4.2	4.0	6.3	7.2	10.7	15.7
2001.....	2.1	2.3	2.5	3.7	3.3	5.2	7.3	8.5	11.6	17.8
2000.....	2.0	2.2	2.6	3.9	4.5	5.0	7.1	11.4	16.1	20.3
1999.....	1.7	2.0	2.3	2.6	4.3	5.7	8.3	12.0	15.3	21.7

Black, male

2006 4/.....	*	2.5	2.6	3.8	4.1	6.6	*	14.8	*	*
2005 4/.....	1.9	2.3	3.8	5.4	5.3	6.7	*	12.1	15.5	22.3
2004 4/.....	2.0	3.9	3.7	4.2	4.7	7.3	7.8	12.3	15.1	*
2003 4/.....	1.5	3.3	3.2	4.6	3.9	6.4	8.9	15.5	17.0	*
2002.....	1.8	2.1	4.0	3.4	6.3	6.7	9.3	9.2	*	25.7
2001.....	2.9	3.7	4.2	5.0	4.9	7.4	11.1	12.1	*	27.2
2000.....	2.7	3.2	4.0	5.8	7.0	7.5	10.4	16.5	21.9	27.7
1999.....	2.5	3.0	3.3	4.1	4.2	7.8	13.2	17.1	20.1	29.3

Black, female

2006 4/.....	*	1.4	*	*	*	*	*	5.6	8.7	11.7
2005 4/.....	*	*	*	2.3	*	*	*	5.6	8.9	11.3
2004 4/.....	*	1.5	*	*	3.8	*	4.6	5.4	10.6	13.4
2003 4/.....	*	1.4	2.6	*	*	*	4.7	*	10.1	13.4
2002.....	*	*	*	*	*	*	*	6.0	9.2	11.9
2001.....	1.3	*	*	2.7	*	*	4.7	6.4	10.2	14.3
2000.....	1.3	*	*	*	*	*	4.9	8.3	13.2	17.6
1999.....	*	*	*	*	4.3	4.2	5.0	8.9	13.0	18.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes

2006.....	19.3	20.7	17.1	9.8	4.6	2.8	2.2	2.4	2.9	3.8
2005.....	17.1	18.2	14.4	8.3	4.4	2.6	2.0	2.3	2.6	3.5
2004.....	16.3	16.7	12.0	6.7	3.7	2.4	2.1	2.4	2.5	3.4
2003.....	15.5	15.4	10.9	6.1	3.4	2.3	2.0	2.0	2.5	3.5
2002.....	14.8	13.8	9.2	5.0	3.0	2.3	1.9	2.1	2.7	3.9
2001.....	12.0	10.9	7.0	3.7	2.4	1.7	1.5	2.1	2.4	3.5
2000.....	11.2	9.9	6.1	3.3	2.2	1.5	1.5	1.6	2.6	4.3
1999.....	11.1	9.8	5.4	3.4	2.0	1.7	1.6	1.9	2.6	3.4

All races 3/, male

2006.....	25.4	26.9	22.5	12.7	6.1	3.0	2.7	3.0	2.7	4.6
2005.....	22.1	24.0	19.1	10.9	5.7	3.0	2.3	3.2	2.8	4.6
2004.....	20.9	21.6	16.3	8.4	4.2	2.7	2.5	2.6	2.9	4.2
2003.....	20.7	20.8	14.7	7.6	4.5	3.0	2.2	2.3	3.5	4.0
2002.....	20.3	18.7	12.8	6.8	3.6	2.8	2.1	2.3	3.2	5.1
2001.....	16.8	15.7	9.9	5.0	3.0	2.0	2.1	2.7	2.8	3.8
2000.....	16.4	14.7	8.9	4.0	2.7	1.8	1.7	1.6	2.9	6.2
1999.....	16.6	15.1	7.9	4.7	3.0	2.2	1.9	2.1	2.9	4.4

All races 3/, female

2006.....	13.4	14.7	11.9	7.1	3.4	2.6	1.9	1.9	3.0	3.4
2005.....	12.0	12.6	10.0	5.8	3.2	2.3	1.7	1.7	2.4	3.0
2004.....	11.7	11.9	7.9	5.1	3.2	2.1	1.8	2.2	2.3	3.0
2003.....	10.3	10.2	7.3	4.6	2.5	1.8	1.8	1.7	2.0	3.3
2002.....	9.4	9.1	5.8	3.4	2.4	1.8	1.7	1.9	2.3	3.4
2001.....	7.3	6.2	4.1	2.6	1.9	1.4	1.0	1.6	2.2	3.3
2000.....	6.0	5.3	3.4	2.7	1.8	1.3	1.3	1.6	2.4	3.6
1999.....	5.7	4.7	3.1	2.2	1.2	1.2	1.4	1.7	2.4	3.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)										
White, both sexes										
2006 4/.....	20.3	21.1	16.6	9.3	4.2	2.5	2.1	2.4	2.9	3.8
2005 4/.....	18.0	18.7	13.8	7.6	4.1	2.4	1.9	2.3	2.5	3.6
2004 4/.....	17.3	17.1	11.7	6.2	3.5	2.3	2.1	2.3	2.4	3.4
2003 4/.....	16.3	15.7	10.6	5.8	3.4	2.0	2.0	2.0	2.7	3.5
2002.....	15.4	13.7	8.7	4.6	2.7	2.2	1.9	2.1	2.6	3.9
2001.....	12.1	10.5	6.1	3.4	2.1	1.6	1.4	2.0	2.3	3.3
2000.....	11.0	9.4	5.3	3.0	2.1	1.5	1.4	1.7	2.6	4.1
1999.....	10.8	9.1	4.8	3.1	1.7	1.5	1.4	1.9	2.6	3.3
White, male										
2006 4/.....	26.3	27.0	21.0	11.3	5.1	2.5	2.3	2.9	2.8	4.7
2005 4/.....	23.3	24.2	17.9	9.4	4.9	2.3	2.0	3.1	2.6	4.8
2004 4/.....	22.2	22.0	15.4	7.3	3.6	2.3	2.4	2.4	2.7	4.3
2003 4/.....	21.8	20.9	13.8	7.0	4.2	2.3	2.2	2.3	3.6	3.7
2002.....	21.0	18.3	11.7	5.8	2.9	2.6	2.0	2.3	3.0	5.2
2001.....	17.1	14.9	8.5	4.3	2.3	1.9	2.0	2.4	2.7	3.7
2000.....	16.2	13.7	7.6	3.3	2.4	1.6	1.6	1.7	2.9	5.9
1999.....	16.2	13.8	6.6	4.0	2.3	2.0	1.7	2.1	3.0	4.5
White, female										
2006 4/.....	14.1	15.3	12.2	7.3	3.5	2.6	2.0	2.0	3.0	3.3
2005 4/.....	12.6	13.3	9.9	5.9	3.4	2.4	1.8	1.7	2.5	3.0
2004 4/.....	12.5	12.2	8.2	5.2	3.4	2.2	1.8	2.3	2.2	3.0
2003 4/.....	10.8	10.5	7.3	4.7	2.6	1.7	1.8	1.8	2.1	3.4
2002.....	9.8	9.0	5.8	3.4	2.5	1.9	1.8	1.9	2.5	3.4
2001.....	7.1	6.1	3.8	2.6	1.9	1.4	1.0	1.7	2.0	3.1
2000.....	5.8	5.1	3.1	2.7	1.8	1.4	1.3	1.7	2.5	3.4
1999.....	5.4	4.5	3.0	2.3	1.2	1.1	1.2	1.8	2.4	2.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental poisoning and exposure to noxious substances (X40-X49)										
Black, both sexes										
2006 4/.....	20.3	24.3	26.9	17.7	9.3	5.4	3.5	*	*	5.4
2005 4/.....	17.0	20.7	23.1	16.3	8.1	5.0	3.7	*	*	*
2004 4/.....	14.6	19.6	17.9	12.6	6.0	4.1	3.1	*	*	*
2003 4/.....	15.2	18.8	16.9	9.6	4.8	5.5	*	*	*	*
2002.....	15.7	19.4	16.1	10.3	6.0	3.0	*	*	*	*
2001.....	15.2	17.7	15.4	7.4	5.4	2.6	*	*	*	6.4
2000.....	15.3	16.5	14.2	6.8	3.9	2.7	*	*	*	8.1
1999.....	16.1	17.9	12.3	6.5	5.5	2.9	3.2	*	*	*
Black, male										
2006 4/.....	27.8	34.5	41.7	29.5	17.0	8.4	6.5	*	*	*
2005 4/.....	22.0	30.3	34.6	26.9	15.1	9.3	6.6	*	*	*
2004 4/.....	19.1	25.8	28.7	20.5	10.1	7.3	*	*	*	*
2003 4/.....	20.5	26.7	25.7	15.8	8.7	10.3	*	*	*	*
2002.....	21.6	27.2	25.6	17.7	11.3	5.5	*	*	*	*
2001.....	20.3	28.0	24.8	12.7	9.6	*	*	*	*	*
2000.....	22.8	26.3	23.1	11.5	7.0	*	*	*	*	*
1999.....	24.2	29.7	20.9	11.9	10.0	*	*	*	*	*
Black, female										
2006 4/.....	13.6	15.3	14.4	8.0	3.3	*	*	*	*	*
2005 4/.....	12.6	12.4	13.4	7.6	*	*	*	*	*	*
2004 4/.....	10.6	14.2	8.9	6.1	*	*	*	*	*	*
2003 4/.....	10.5	11.9	9.5	4.5	*	*	*	*	*	*
2002.....	10.5	12.5	8.1	4.3	*	*	*	*	*	*
2001.....	10.7	8.9	7.5	3.1	*	*	*	*	*	*
2000.....	8.6	8.0	6.6	3.0	*	*	*	*	*	*
1999.....	9.1	7.8	4.9	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

All races 3/, both sexes

2006.....	18.7	20.1	16.6	9.4	4.2	2.5	1.9	1.7	2.4	2.9
2005.....	16.5	17.5	13.8	7.7	3.9	2.2	1.6	1.8	1.9	2.6
2004.....	15.7	16.0	11.4	6.3	3.3	2.0	1.6	1.8	1.8	2.5
2003.....	14.8	14.7	10.4	5.5	2.9	1.9	1.4	1.4	2.1	2.8
2002.....	14.2	13.2	8.7	4.4	2.5	1.8	1.3	1.5	1.9	3.1
2001.....	11.4	10.4	6.4	3.3	2.0	1.4	1.2	1.5	1.7	2.5
2000.....	10.6	9.3	5.6	2.8	1.7	1.2	1.0	1.2	1.7	3.3
1999.....	10.5	9.2	4.9	2.9	1.6	1.3	1.1	1.4	1.9	2.8

All races 3/, male

2006.....	24.4	25.9	21.7	12.1	5.4	2.5	2.2	2.1	2.1	3.0
2005.....	21.3	22.9	18.0	10.0	5.0	2.4	1.8	2.4	1.9	3.2
2004.....	20.0	20.6	15.3	7.8	3.7	2.1	1.7	1.9	1.7	2.6
2003.....	19.6	19.6	13.9	6.8	3.6	2.4	1.4	1.7	2.7	2.7
2002.....	19.2	17.7	12.0	5.8	2.9	2.0	1.4	1.6	2.1	3.7
2001.....	15.8	14.9	9.1	4.4	2.4	1.5	1.5	1.8	1.7	1.8
2000.....	15.6	13.9	8.2	3.2	2.0	1.5	1.2	1.1	1.5	4.4
1999.....	15.5	14.2	7.1	4.0	2.2	1.6	1.3	1.3	1.7	3.3

All races 3/, female

2006.....	13.0	14.4	11.7	6.8	3.2	2.5	1.7	1.5	2.6	2.9
2005.....	11.7	12.3	9.7	5.6	3.0	2.1	1.4	1.4	2.0	2.3
2004.....	11.5	11.6	7.6	4.8	3.0	1.9	1.4	1.8	1.8	2.5
2003.....	10.1	9.9	7.1	4.3	2.3	1.5	1.4	1.3	1.7	2.9
2002.....	9.2	8.9	5.6	3.2	2.1	1.5	1.3	1.4	1.8	2.8
2001.....	7.0	6.0	3.9	2.3	1.7	1.2	0.9	1.2	1.7	2.7
2000.....	5.8	4.9	3.2	2.4	1.4	1.0	0.9	1.3	1.9	2.8
1999.....	5.5	4.5	2.9	1.9	1.1	1.0	1.0	1.4	2.0	2.6

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

White, both sexes

2006 4/.....	19.6	20.5	16.0	8.8	3.8	2.3	1.8	1.8	2.4	2.9
2005 4/.....	17.4	18.0	13.2	7.1	3.7	2.0	1.5	1.8	2.0	2.7
2004 4/.....	16.8	16.4	11.1	5.7	3.2	1.9	1.6	1.8	1.7	2.6
2003 4/.....	15.6	15.0	10.0	5.3	2.9	1.7	1.4	1.5	2.1	2.8
2002.....	14.8	13.1	8.2	4.1	2.2	1.7	1.4	1.5	1.9	3.2
2001.....	11.5	9.9	5.7	3.0	1.8	1.3	1.1	1.4	1.7	2.3
2000.....	10.5	8.8	4.9	2.5	1.5	1.2	1.0	1.3	1.7	3.1
1999.....	10.2	8.6	4.3	2.7	1.3	1.2	1.0	1.4	1.9	2.6

White, male

2006 4/.....	25.4	26.0	20.2	10.7	4.4	2.0	1.8	2.0	2.1	3.2
2005 4/.....	22.5	23.0	16.7	8.7	4.3	1.8	1.5	2.4	1.8	3.5
2004 4/.....	21.2	20.9	14.5	6.6	3.2	1.7	1.7	1.8	1.6	2.7
2003 4/.....	20.6	19.8	13.0	6.2	3.4	1.9	1.4	1.7	2.7	2.4
2002.....	19.9	17.4	10.9	5.0	2.2	1.8	1.3	1.5	1.9	3.8
2001.....	16.1	14.0	7.8	3.7	1.8	1.4	1.4	1.6	1.8	1.7
2000.....	15.3	12.9	6.9	2.6	1.6	1.2	1.0	1.2	1.4	4.1
1999.....	15.1	12.9	5.9	3.4	1.6	1.4	1.2	1.4	1.7	3.2

White, female

2006 4/.....	13.7	15.1	11.9	7.1	3.3	2.5	1.7	1.6	2.5	2.8
2005 4/.....	12.3	13.0	9.7	5.7	3.1	2.1	1.4	1.4	2.1	2.3
2004 4/.....	12.3	11.9	7.8	4.9	3.2	2.1	1.5	1.9	1.8	2.5
2003 4/.....	10.5	10.2	7.1	4.5	2.4	1.5	1.4	1.4	1.8	3.0
2002.....	9.5	8.8	5.6	3.2	2.3	1.6	1.4	1.5	2.0	2.9
2001.....	6.8	5.9	3.6	2.3	1.7	1.2	0.9	1.3	1.7	2.6
2000.....	5.6	4.7	2.9	2.5	1.4	1.1	0.9	1.3	1.9	2.6
1999.....	5.2	4.3	2.8	2.0	1.1	1.0	0.8	1.4	2.0	2.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning by and exposure to drugs and other biological substances (X40-X44)

Black, both sexes

2006 4/.....	19.6	23.5	26.3	17.1	8.7	4.9	3.0	*	*	*
2005 4/.....	16.3	20.2	22.2	15.6	7.3	4.4	3.0	*	*	*
2004 4/.....	14.2	19.0	17.4	12.0	5.4	3.5	*	*	*	*
2003 4/.....	14.6	18.1	16.3	9.0	4.2	4.5	*	*	*	*
2002.....	15.0	18.5	15.4	9.1	5.5	2.4	*	*	*	*
2001.....	14.6	17.2	14.3	7.0	4.7	2.2	*	*	*	*
2000.....	14.6	15.8	13.5	5.9	3.4	2.3	*	*	*	6.3
1999.....	15.7	17.1	11.4	5.6	4.7	2.4	*	*	*	*

Black, male

2006 4/.....	26.8	33.3	40.6	28.4	16.1	8.2	*	*	*	*
2005 4/.....	20.9	29.2	33.3	25.3	13.5	8.3	*	*	*	*
2004 4/.....	18.4	24.8	27.6	19.7	9.0	6.5	*	*	*	*
2003 4/.....	19.6	25.3	24.6	14.7	7.3	8.4	*	*	*	*
2002.....	20.5	25.8	24.4	15.3	10.5	*	*	*	*	*
2001.....	19.2	27.2	23.1	12.2	8.1	*	*	*	*	*
2000.....	21.8	25.2	22.3	9.9	6.2	*	*	*	*	*
1999.....	23.5	28.3	19.5	10.3	8.5	*	*	*	*	*

Black, female

2006 4/.....	13.3	15.1	14.2	7.8	3.0	*	*	*	*	*
2005 4/.....	12.2	12.4	12.8	7.5	*	*	*	*	*	*
2004 4/.....	10.5	14.0	8.7	5.6	*	*	*	*	*	*
2003 4/.....	10.3	11.9	9.3	4.3	*	*	*	*	*	*
2002.....	10.1	12.1	7.7	3.9	*	*	*	*	*	*
2001.....	10.4	8.5	6.8	2.8	*	*	*	*	*	*
2000.....	8.3	7.8	6.0	*	*	*	*	*	*	*
1999.....	8.8	7.4	4.6	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

All races 3/, both sexes

2006.....	2.1	2.6	3.0	3.2	3.8	4.8	6.8	11.4	23.7	61.2
2005.....	2.3	2.8	3.0	3.0	3.9	5.0	7.2	12.0	23.6	63.6
2004.....	2.0	2.4	2.9	2.8	3.6	4.4	6.6	11.7	23.2	64.5
2003.....	2.2	2.5	2.6	3.1	3.4	5.3	6.7	12.7	24.3	73.5
2002.....	2.2	2.2	2.7	3.2	3.9	5.5	7.0	13.5	25.6	75.8
2001.....	2.0	2.5	2.9	2.9	4.3	4.9	8.0	14.4	28.4	83.4
2000.....	2.3	2.6	2.7	3.1	4.1	5.0	7.6	13.6	26.9	82.6
1999.....	2.4	2.5	2.9	3.2	4.3	5.9	8.1	15.3	32.3	92.8

All races 3/, male

2006.....	3.5	4.2	5.0	5.1	5.8	7.5	9.7	15.5	29.1	61.2
2005.....	3.8	4.6	5.2	4.6	6.3	7.3	10.1	16.4	30.3	66.7
2004.....	3.2	4.1	4.8	4.4	5.7	6.5	9.0	14.9	27.9	69.6
2003.....	3.8	4.1	4.2	4.9	5.6	7.9	9.5	16.9	28.7	80.4
2002.....	3.5	3.8	4.2	5.1	6.1	7.9	9.8	17.8	30.3	79.9
2001.....	3.3	4.2	4.7	4.4	6.6	7.1	10.6	19.0	33.9	92.4
2000.....	3.9	4.4	4.5	5.2	6.1	7.1	10.0	17.5	31.1	91.7
1999.....	3.9	4.1	4.9	5.2	6.6	8.2	11.1	19.5	40.3	107.0

All races 3/, female

2006.....	0.7	1.0	1.1	1.5	1.9	2.5	4.5	8.4	20.2	61.3
2005.....	0.8	1.0	0.9	1.4	1.7	2.9	4.9	8.8	19.5	62.2
2004.....	0.8	0.8	1.2	1.3	1.7	2.7	4.8	9.3	20.3	62.2
2003.....	0.7	0.9	1.1	1.4	1.4	3.1	4.5	9.7	21.6	70.5
2002.....	0.8	0.7	1.2	1.4	1.8	3.3	4.7	10.4	22.7	74.0
2001.....	0.7	0.9	1.3	1.4	2.2	3.1	5.8	11.2	25.1	79.6
2000.....	0.7	0.8	0.9	1.2	2.2	3.2	5.6	10.9	24.3	78.9
1999.....	0.9	0.9	1.0	1.4	2.2	3.9	5.8	12.4	27.6	87.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)

White, both sexes

2006 4/.....	2.1	2.5	2.9	3.2	3.7	4.8	6.8	11.7	24.7	64.9
2005 4/.....	2.3	2.7	2.9	2.9	3.8	5.0	7.3	12.1	24.3	66.8
2004 4/.....	2.1	2.5	2.9	2.7	3.6	4.4	6.7	11.6	23.8	67.6
2003 4/.....	2.2	2.3	2.4	2.9	3.3	5.3	6.7	12.9	24.8	77.0
2002.....	2.2	2.2	2.6	3.1	3.8	5.1	7.1	13.4	25.9	79.1
2001.....	1.9	2.4	2.8	2.9	4.1	4.8	7.9	14.6	29.2	86.4
2000.....	2.2	2.4	2.6	3.0	4.0	4.8	7.5	13.4	27.5	86.2
1999.....	2.3	2.2	2.8	3.0	4.0	5.6	7.8	15.0	32.6	96.1

White, male

2006 4/.....	3.5	4.1	4.7	4.9	5.8	7.5	9.5	15.8	30.2	64.5
2005 4/.....	3.9	4.4	5.0	4.5	6.1	7.4	9.9	16.4	31.1	70.1
2004 4/.....	3.4	4.2	4.7	4.2	5.7	6.4	9.1	14.7	28.5	72.6
2003 4/.....	3.8	3.9	3.9	4.6	5.4	7.8	9.2	17.1	28.8	83.8
2002.....	3.6	3.8	4.1	4.8	6.0	7.3	9.9	17.6	30.7	82.5
2001.....	3.1	4.1	4.4	4.4	6.3	6.9	10.5	19.3	34.4	95.8
2000.....	3.7	4.0	4.3	4.7	5.8	6.7	10.0	16.9	31.4	95.0
1999.....	3.7	3.6	4.6	4.8	6.3	7.7	10.5	19.0	40.5	108.9

White, female

2006 4/.....	0.6	1.0	1.2	1.5	1.9	2.3	4.4	8.6	21.2	65.1
2005 4/.....	0.7	1.0	0.8	1.4	1.7	2.9	5.1	8.9	20.1	65.3
2004 4/.....	0.9	0.8	1.1	1.3	1.6	2.6	4.7	9.4	20.9	65.4
2003 4/.....	0.6	0.8	1.0	1.4	1.5	3.0	4.6	9.9	22.2	74.0
2002.....	0.7	0.6	1.2	1.5	1.8	3.1	4.8	10.4	23.0	77.6
2001.....	0.6	0.8	1.3	1.5	2.0	3.1	5.8	11.3	26.0	82.4
2000.....	0.6	0.7	0.9	1.3	2.3	3.2	5.4	11.0	25.3	82.6
1999.....	0.9	0.9	1.0	1.4	1.9	3.8	5.7	12.2	27.9	91.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
All other and unspecified accidents and their sequelae (W20-W31,W35-W38,W41-W64,W92-W99,X20-X32,X39,X50-X59,Y86)										
Black, both sexes										
2006 4/.....	2.5	3.5	4.0	4.5	4.9	5.8	9.3	10.9	16.8	32.7
2005 4/.....	2.3	3.8	4.0	3.5	4.9	5.7	8.1	14.4	18.4	38.8
2004 4/.....	1.9	2.7	3.8	4.3	4.3	5.7	6.7	14.6	19.6	38.4
2003 4/.....	2.5	3.8	4.3	4.6	4.6	7.0	8.1	11.9	22.3	45.8
2002.....	2.4	3.0	3.3	4.7	4.7	9.3	6.3	15.6	22.8	46.9
2001.....	2.8	3.5	4.3	2.8	6.8	6.4	9.6	15.1	23.0	58.3
2000.....	3.1	4.2	3.6	5.3	5.4	6.8	9.3	17.8	21.2	51.0
1999.....	3.4	4.7	4.6	5.7	7.1	9.4	13.1	20.7	32.6	67.0
Black, male										
2006 4/.....	4.2	5.9	7.6	7.0	7.2	8.9	14.5	16.6	22.6	37.5
2005 4/.....	3.7	6.7	6.9	5.9	9.5	8.6	12.3	21.0	24.6	42.8
2004 4/.....	3.0	4.6	6.3	7.5	7.1	8.5	9.7	21.4	28.0	42.7
2003 4/.....	4.0	6.3	6.9	7.7	8.5	10.8	14.1	16.9	29.5	54.2
2002.....	3.2	4.6	5.5	8.6	7.9	13.5	9.0	22.5	29.8	57.9
2001.....	4.7	5.4	7.3	5.0	10.0	10.4	13.1	18.6	32.9	62.4
2000.....	5.4	7.8	7.1	10.4	9.3	10.9	11.7	27.9	32.9	69.2
1999.....	5.9	8.4	7.8	9.8	10.8	15.1	20.4	28.0	41.0	106.6
Black, female										
2006 4/.....	*	1.5	*	2.4	3.0	3.5	5.7	7.5	13.7	30.7
2005 4/.....	*	*	*	*	*	3.5	5.2	10.4	15.2	37.1
2004 4/.....	*	*	1.7	*	*	3.6	4.6	10.5	15.1	36.6
2003 4/.....	1.3	1.6	2.2	*	*	4.2	*	9.0	18.6	42.5
2002.....	1.8	1.6	*	*	*	6.3	4.5	11.5	19.1	42.7
2001.....	*	1.7	*	*	4.3	*	7.2	13.0	18.0	56.8
2000.....	*	*	*	*	*	3.8	7.8	11.7	15.3	44.2
1999.....	*	*	*	*	4.2	5.2	8.3	16.3	28.5	52.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All races 3/, both sexes

2006.....	15.5	17.1	17.3	15.3	13.4	12.6	12.6	15.6	16.3	15.9
2005.....	16.0	16.7	16.1	14.3	13.3	11.8	13.5	15.4	18.7	16.9
2004.....	16.0	16.9	16.2	14.2	13.2	11.4	13.5	16.0	16.7	16.4
2003.....	15.9	16.0	15.7	14.7	12.6	12.0	13.6	15.6	17.7	16.9
2002.....	16.1	16.3	15.1	14.6	12.3	12.5	14.6	16.6	19.3	18.0
2001 5/.....	15.2	15.7	14.6	14.0	12.0	12.7	13.9	16.4	18.9	17.5
2000.....	15.4	14.9	13.9	12.7	11.4	11.3	13.8	16.5	19.3	19.6
1999.....	14.3	14.3	13.4	12.8	11.4	12.4	14.5	17.4	19.2	19.3

All races 3/, male

2006.....	23.6	26.0	26.4	23.9	21.1	22.0	23.6	31.6	35.7	43.2
2005.....	24.6	25.4	25.0	22.4	22.0	20.6	25.4	31.9	41.4	45.0
2004.....	24.2	25.2	24.4	22.3	21.7	20.2	25.4	32.4	38.3	45.0
2003.....	24.5	24.4	24.4	23.1	21.1	21.2	26.0	31.8	40.1	47.8
2002.....	24.7	24.9	23.8	23.5	20.4	22.0	27.9	34.3	44.0	50.7
2001 5/.....	23.6	24.3	22.4	22.3	19.5	22.7	26.7	33.8	44.2	51.1
2000.....	23.9	23.3	21.4	20.1	18.7	19.8	26.1	34.9	44.8	57.5
1999.....	22.1	22.3	20.6	20.3	19.1	22.1	27.7	35.6	44.1	57.1

All races 3/, female

2006.....	7.4	8.4	8.4	7.2	6.3	4.4	3.7	3.8	4.2	3.1
2005.....	7.4	8.2	7.7	6.6	5.5	4.2	3.8	3.5	4.7	4.0
2004.....	7.9	8.9	8.4	6.6	5.5	3.7	3.9	4.3	3.5	3.6
2003.....	7.4	7.9	7.5	6.7	4.9	4.0	3.6	4.0	3.9	3.3
2002.....	7.7	8.0	6.7	6.2	5.0	4.3	3.9	4.1	4.4	3.8
2001 5/.....	6.9	7.4	7.0	6.2	5.2	4.1	3.7	4.3	3.8	3.4
2000.....	7.0	6.7	6.8	5.8	4.9	4.0	4.1	3.8	4.2	4.2
1999.....	6.7	6.7	6.5	5.8	4.4	4.2	4.1	4.8	4.6	4.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)										
White, both sexes										
2006 4/.....	17.5	19.3	19.5	17.1	14.8	13.8	13.8	16.9	17.6	16.9
2005 4/.....	18.0	18.9	18.1	15.9	14.7	13.0	14.7	16.7	20.0	17.9
2004 4/.....	18.1	19.0	18.3	15.7	14.4	12.4	14.5	17.3	17.7	17.5
2003 4/.....	17.9	18.1	17.7	16.2	13.8	13.0	14.7	16.7	18.9	18.1
2002.....	18.2	18.4	16.8	16.1	13.6	13.7	15.7	17.6	20.9	19.1
2001 5/.....	16.9	17.6	16.2	15.3	13.1	13.8	14.9	17.6	20.0	18.6
2000.....	17.0	16.5	15.3	13.9	12.6	12.3	14.7	17.6	20.4	21.0
1999.....	16.0	15.9	14.8	14.0	12.3	13.3	15.6	18.6	20.4	20.7
White, male										
2006 4/.....	26.3	29.0	29.5	26.4	23.0	23.8	25.9	34.1	38.7	46.1
2005 4/.....	27.3	28.5	27.7	24.8	24.1	22.6	27.6	34.2	44.3	48.2
2004 4/.....	27.0	27.8	27.3	24.1	23.6	21.9	27.1	34.8	40.4	48.4
2003 4/.....	27.3	27.3	27.2	25.4	22.9	22.9	27.9	34.0	42.9	51.4
2002.....	27.6	27.6	26.4	25.7	22.5	23.8	30.1	36.3	47.2	53.9
2001 5/.....	26.1	27.1	24.7	24.3	21.0	24.5	28.3	36.0	46.8	55.0
2000.....	26.2	25.6	23.2	21.8	20.4	21.3	27.6	37.1	47.7	61.6
1999.....	24.5	24.4	22.3	21.9	20.5	23.6	29.5	37.9	46.8	61.5
White, female										
2006 4/.....	8.5	9.6	9.7	8.3	7.0	4.8	3.8	4.1	4.3	3.2
2005 4/.....	8.5	9.4	8.7	7.4	6.0	4.4	4.1	3.7	4.8	3.9
2004 4/.....	9.1	10.2	9.5	7.6	5.8	4.1	4.1	4.5	3.6	3.6
2003 4/.....	8.4	9.0	8.3	7.4	5.5	4.3	3.9	4.2	4.1	3.3
2002.....	8.7	9.3	7.4	6.9	5.3	4.8	3.9	4.2	4.7	4.0
2001 5/.....	7.8	8.1	7.9	6.8	5.8	4.3	4.0	4.6	3.8	3.4
2000.....	7.9	7.5	7.5	6.4	5.5	4.4	4.4	3.9	4.2	4.5
1999.....	7.5	7.4	7.4	6.4	4.8	4.4	4.5	5.0	4.8	4.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

All races 3/, both sexes

2006.....	2.8	3.1	3.5	2.8	1.9	1.2	0.8	1.0	1.0	0.9
2005.....	2.8	3.2	2.9	2.3	1.9	1.1	1.0	0.9	1.3	1.0
2004.....	2.8	3.1	3.1	2.3	1.6	0.9	0.9	1.0	0.8	1.1
2003.....	2.8	3.0	2.8	2.1	1.4	1.1	0.8	0.9	0.9	0.9
2002.....	2.9	3.1	2.6	2.0	1.4	1.0	1.0	1.0	1.1	0.8
2001.....	2.7	2.9	2.4	1.7	1.2	1.0	0.9	0.9	0.6	0.9
2000.....	2.6	2.5	2.3	1.8	1.0	0.7	0.9	0.8	0.9	1.0
1999.....	2.5	2.4	2.2	1.6	1.1	1.0	0.9	1.1	1.1	0.8

All races 3/, male

2006.....	2.7	2.9	3.3	2.7	1.5	0.9	0.7	0.9	1.0	*
2005.....	2.8	2.7	2.8	2.2	1.6	1.0	0.9	1.0	1.2	1.3
2004.....	2.5	2.8	2.9	2.1	1.6	0.6	0.9	1.0	*	1.6
2003.....	2.9	2.7	2.6	1.9	1.2	0.9	0.8	0.7	*	1.7
2002.....	2.8	2.7	2.6	1.7	1.2	0.9	0.9	1.1	1.3	*
2001.....	2.8	2.8	2.0	1.4	0.8	1.2	0.8	0.8	*	*
2000.....	2.5	2.5	2.0	1.6	0.7	0.5	0.8	0.9	1.3	*
1999.....	2.5	2.5	2.0	1.4	1.0	1.1	0.9	0.7	1.2	*

All races 3/, female

2006.....	3.0	3.4	3.7	3.0	2.2	1.4	0.9	1.1	1.0	0.9
2005.....	2.8	3.6	3.1	2.4	2.1	1.2	1.1	0.9	1.4	0.9
2004.....	3.2	3.4	3.3	2.6	1.6	1.0	0.9	1.0	0.8	0.8
2003.....	2.7	3.3	2.9	2.3	1.6	1.3	0.9	1.1	1.0	*
2002.....	3.1	3.5	2.5	2.3	1.6	1.1	1.1	0.9	0.9	0.7
2001.....	2.6	3.1	2.7	1.9	1.5	0.9	1.0	0.9	0.6	0.6
2000.....	2.8	2.5	2.5	1.9	1.3	0.7	1.1	0.8	0.7	0.9
1999.....	2.5	2.4	2.4	1.8	1.2	0.9	0.9	1.3	1.0	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-poisoning (suicide) by and exposure to drugs and other biological substances (X60-X64)

White, both sexes

2006 4/.....	3.3	3.6	4.0	3.2	2.0	1.4	0.8	1.1	1.1	1.0
2005 4/.....	3.3	3.6	3.3	2.6	2.1	1.3	1.2	0.9	1.4	1.1
2004 4/.....	3.3	3.6	3.5	2.6	1.8	0.9	1.0	1.1	0.9	1.1
2003 4/.....	3.1	3.4	3.2	2.3	1.6	1.2	0.9	1.0	1.0	1.0
2002.....	3.4	3.6	2.9	2.3	1.5	1.1	1.0	1.1	1.1	0.9
2001.....	3.1	3.3	2.7	1.9	1.3	1.1	1.0	1.0	0.7	0.9
2000.....	3.0	2.8	2.5	2.0	1.2	0.7	1.1	0.9	1.0	1.1
1999.....	2.8	2.8	2.5	1.8	1.2	1.1	1.0	1.1	1.2	0.8

White, male

2006 4/.....	3.1	3.3	3.7	2.9	1.6	1.1	0.6	1.0	1.1	*
2005 4/.....	3.2	3.1	3.2	2.4	1.8	1.1	1.1	1.0	1.3	1.5
2004 4/.....	2.9	3.2	3.2	2.2	1.8	0.6	0.9	1.1	*	1.7
2003 4/.....	3.2	3.1	3.1	2.1	1.3	1.0	0.8	0.8	*	1.8
2002.....	3.2	3.1	3.0	2.0	1.3	0.9	0.9	1.1	1.3	*
2001.....	3.2	3.1	2.3	1.6	0.9	1.3	0.8	0.9	*	*
2000.....	2.8	2.7	2.2	1.8	0.8	0.6	0.8	0.9	1.4	*
1999.....	2.9	2.8	2.2	1.6	1.0	1.2	1.0	0.7	1.3	*

White, female

2006 4/.....	3.5	4.0	4.2	3.5	2.4	1.6	1.0	1.2	1.1	1.0
2005 4/.....	3.3	4.2	3.5	2.7	2.4	1.4	1.2	0.9	1.5	1.0
2004 4/.....	3.7	4.0	3.8	3.0	1.8	1.2	1.1	1.1	0.9	0.8
2003 4/.....	3.1	3.6	3.3	2.5	1.8	1.4	1.0	1.2	1.1	*
2002.....	3.5	4.0	2.8	2.6	1.7	1.3	1.1	1.0	1.0	0.8
2001.....	3.0	3.5	3.2	2.1	1.7	0.9	1.1	1.0	0.7	*
2000.....	3.1	2.9	2.8	2.2	1.6	0.8	1.2	0.9	0.7	1.0
1999.....	2.8	2.8	2.7	2.0	1.4	1.0	1.0	1.4	1.1	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

All races 3/, both sexes

2006.....	0.7	0.8	0.8	0.8	0.5	0.5	0.4	0.4	0.5	0.6
2005.....	0.8	0.8	0.8	0.6	0.5	0.4	0.5	0.4	0.6	0.6
2004.....	0.9	0.9	0.9	0.6	0.5	0.3	0.4	0.5	0.5	0.6
2003.....	0.9	0.8	0.8	0.8	0.5	0.3	0.5	0.4	0.6	0.7
2002.....	0.9	1.0	0.9	0.8	0.4	0.4	0.4	0.6	0.7	0.8
2001.....	0.9	1.0	0.8	0.9	0.5	0.5	0.4	0.6	0.7	0.8
2000.....	0.9	1.1	0.8	0.8	0.5	0.4	0.4	0.5	0.7	0.8
1999.....	1.0	1.0	0.9	0.7	0.6	0.3	0.4	0.8	0.9	1.2

All races 3/, male

2006.....	1.1	1.2	1.2	1.3	0.8	0.9	*	0.7	1.0	1.5
2005.....	1.3	1.3	1.3	0.9	0.8	0.6	0.7	0.8	1.1	1.2
2004.....	1.4	1.4	1.5	1.0	0.9	*	0.6	0.8	1.0	*
2003.....	1.4	1.3	1.2	1.2	0.9	0.6	0.9	0.6	1.0	1.6
2002.....	1.5	1.7	1.3	1.3	0.7	0.7	0.6	1.1	1.5	2.2
2001.....	1.4	1.6	1.2	1.3	0.7	0.8	0.7	1.0	1.5	1.9
2000.....	1.4	1.7	1.2	1.3	0.7	0.6	0.6	1.0	1.6	2.3
1999.....	1.5	1.5	1.2	1.0	0.8	0.6	0.6	1.3	1.7	3.9

All races 3/, female

2006.....	0.3	0.4	0.4	0.4	0.3	*	*	*	*	*
2005.....	0.4	0.3	0.4	0.3	0.3	*	*	*	*	*
2004.....	0.3	0.4	0.4	0.3	*	*	*	*	*	*
2003.....	0.4	0.3	0.4	0.3	*	*	*	*	*	*
2002.....	0.4	0.4	0.4	0.3	*	*	*	*	*	*
2001.....	0.4	0.4	0.4	0.6	*	*	*	*	*	*
2000.....	0.4	0.4	0.5	0.4	*	*	*	*	*	*
1999.....	0.4	0.5	0.5	0.4	0.4	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-poisoning (suicide) by and exposure to other gases and vapors (X67)

White, both sexes

2006 4/.....	0.9	0.9	0.9	1.0	0.6	0.5	0.4	0.5	0.6	0.6
2005 4/.....	1.0	0.9	1.0	0.7	0.6	0.4	0.5	0.5	0.6	0.7
2004 4/.....	1.0	1.1	1.1	0.7	0.6	0.3	0.5	0.6	0.5	0.7
2003 4/.....	1.0	1.0	0.9	0.9	0.5	0.4	0.6	0.5	0.7	0.8
2002.....	1.1	1.2	1.0	0.9	0.4	0.5	0.5	0.7	0.8	0.9
2001.....	1.1	1.1	1.0	1.0	0.6	0.6	0.5	0.7	0.8	0.9
2000.....	1.1	1.2	0.9	0.9	0.6	0.4	0.4	0.5	0.8	0.9
1999.....	1.1	1.1	1.0	0.8	0.7	0.4	0.4	0.8	0.9	1.3

White, male

2006 4/.....	1.3	1.3	1.4	1.5	0.9	0.9	*	0.8	1.1	1.6
2005 4/.....	1.6	1.5	1.4	1.0	0.9	0.7	0.8	0.9	1.1	1.4
2004 4/.....	1.6	1.7	1.7	1.1	1.0	*	0.7	0.9	1.1	*
2003 4/.....	1.6	1.6	1.4	1.4	0.8	0.7	1.0	0.7	1.1	1.8
2002.....	1.7	1.9	1.5	1.4	0.8	0.7	0.7	1.1	1.6	2.3
2001.....	1.6	1.8	1.4	1.5	0.8	0.9	0.7	1.1	1.7	2.1
2000.....	1.7	1.9	1.3	1.5	0.9	0.6	0.7	1.1	1.7	2.5
1999.....	1.8	1.7	1.4	1.2	0.9	0.7	0.7	1.5	1.8	4.4

White, female

2006 4/.....	0.4	0.4	0.5	0.4	0.4	*	*	*	*	*
2005 4/.....	0.4	0.3	0.5	0.4	0.3	*	*	*	*	*
2004 4/.....	0.4	0.5	0.5	0.3	*	*	*	*	*	*
2003 4/.....	0.4	0.4	0.5	0.4	*	*	*	*	*	*
2002.....	0.4	0.5	0.5	0.4	*	*	*	*	*	*
2001.....	0.5	0.5	0.5	0.6	*	*	*	*	*	*
2000.....	0.5	0.5	0.6	0.4	*	*	*	*	*	*
1999.....	0.4	0.6	0.6	0.4	0.4	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

All races 3/, both sexes

2006.....	3.7	3.6	2.8	2.1	1.6	1.3	1.2	1.4	1.6	2.1
2005.....	3.8	3.1	2.5	2.1	1.6	1.2	1.3	1.2	1.8	2.2
2004.....	3.7	3.3	2.5	2.2	1.6	1.3	1.2	1.5	1.7	2.2
2003.....	3.4	2.9	2.4	2.1	1.4	1.2	1.2	1.5	1.9	2.1
2002.....	3.4	3.0	2.4	1.9	1.4	1.4	1.3	1.5	1.8	2.3
2001.....	3.2	2.7	2.1	1.7	1.4	1.3	1.3	1.4	2.1	2.1
2000.....	3.1	2.5	1.9	1.5	1.3	1.1	1.2	1.7	2.0	2.9
1999.....	2.8	2.1	1.9	1.4	1.3	1.4	1.3	1.8	2.0	2.8

All races 3/, male

2006.....	6.4	6.2	4.8	3.5	2.6	2.1	2.0	2.5	2.6	4.7
2005.....	6.4	5.1	4.1	3.5	2.6	2.1	2.2	2.0	2.9	4.7
2004.....	6.4	5.4	4.1	3.6	2.7	2.2	2.0	2.5	3.4	4.8
2003.....	5.8	4.9	4.0	3.5	2.2	2.1	2.3	2.6	3.7	5.0
2002.....	5.6	5.0	3.8	3.3	2.4	2.4	2.2	2.6	3.0	5.2
2001.....	5.5	4.7	3.6	2.9	2.1	2.2	2.4	2.3	4.3	5.2
2000.....	5.4	4.3	3.2	2.4	2.0	1.5	2.1	3.0	3.8	7.2
1999.....	4.7	3.6	3.0	2.5	2.1	2.1	2.2	2.9	3.6	6.6

All races 3/, female

2006.....	1.0	1.0	0.9	0.8	0.7	0.6	0.6	0.7	1.0	0.9
2005.....	1.2	1.1	1.0	0.9	0.7	0.5	0.5	0.6	1.1	1.0
2004.....	1.1	1.3	1.1	0.9	0.7	0.5	0.5	0.7	0.7	1.1
2003.....	1.1	0.9	0.9	0.7	0.6	0.5	*	0.6	0.8	0.9
2002.....	1.1	1.0	1.0	0.6	0.6	0.6	0.6	0.7	1.0	1.0
2001.....	0.9	0.8	0.7	0.6	0.7	0.6	*	0.7	0.8	0.8
2000.....	0.9	0.8	0.7	0.7	0.7	0.8	0.5	0.8	1.0	1.2
1999.....	0.8	0.7	0.7	0.4	0.6	0.8	0.5	1.0	1.1	1.3

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by hanging, strangulation and suffocation (X70)

White, both sexes

2006 4/.....	4.0	3.9	3.1	2.2	1.7	1.3	1.2	1.4	1.5	2.2
2005 4/.....	4.1	3.4	2.8	2.2	1.7	1.2	1.3	1.3	1.7	2.1
2004 4/.....	4.1	3.6	2.7	2.4	1.6	1.3	1.1	1.4	1.8	2.3
2003 4/.....	3.7	3.1	2.6	2.1	1.4	1.1	1.2	1.5	2.0	2.1
2002.....	3.7	3.2	2.6	2.0	1.6	1.3	1.3	1.5	1.8	2.3
2001.....	3.5	3.0	2.2	1.8	1.3	1.3	1.2	1.4	2.1	2.1
2000.....	3.3	2.7	2.0	1.6	1.4	1.1	1.2	1.6	2.0	2.9
1999.....	3.0	2.3	1.9	1.5	1.3	1.4	1.3	1.7	2.1	2.9

White, male

2006 4/.....	6.9	6.7	5.2	3.6	2.7	2.2	2.2	2.4	2.6	4.9
2005 4/.....	7.0	5.5	4.5	3.6	2.8	2.2	2.4	2.1	2.9	4.8
2004 4/.....	7.0	5.7	4.4	3.8	2.7	2.2	1.9	2.6	3.4	4.9
2003 4/.....	6.2	5.3	4.3	3.5	2.2	1.9	2.2	2.7	3.8	5.1
2002.....	6.1	5.3	4.1	3.4	2.6	2.3	2.4	2.6	3.1	5.2
2001.....	6.0	5.2	3.8	3.0	2.0	2.1	2.3	2.3	4.4	5.3
2000.....	5.7	4.6	3.3	2.5	2.1	1.4	2.0	2.9	3.8	7.1
1999.....	5.1	3.8	3.1	2.6	2.1	2.1	2.2	2.8	3.7	6.9

White, female

2006 4/.....	1.1	1.0	1.0	0.9	0.7	0.6	*	0.7	0.8	0.9
2005 4/.....	1.3	1.2	1.1	0.9	0.7	*	*	0.6	1.0	0.8
2004 4/.....	1.2	1.5	1.1	1.0	0.6	0.5	*	0.6	0.7	1.1
2003 4/.....	1.2	1.0	1.0	0.7	0.6	0.4	*	0.6	0.8	0.8
2002.....	1.2	1.2	1.1	0.5	0.6	0.5	0.5	0.6	1.0	1.0
2001.....	1.0	0.8	0.7	0.6	0.7	0.6	*	0.7	0.8	0.8
2000.....	0.9	0.8	0.7	0.7	0.6	0.8	0.5	0.8	0.9	1.2
1999.....	0.9	0.8	0.8	0.5	0.5	0.7	0.5	1.0	1.1	1.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes

2006.....	6.7	7.8	8.3	8.1	8.1	8.5	9.4	11.8	12.2	10.9
2005.....	7.0	8.0	8.3	8.1	8.2	8.2	9.9	11.9	13.6	11.2
2004.....	7.1	8.1	8.0	7.8	8.3	8.0	9.8	12.2	12.3	11.1
2003.....	7.2	7.8	8.3	8.4	8.2	8.5	10.1	11.9	13.1	11.9
2002.....	7.4	7.8	7.9	8.6	8.1	8.8	10.7	12.4	14.3	12.0
2001.....	7.0	7.5	8.0	8.4	8.0	9.0	10.3	12.4	14.1	12.2
2000.....	7.4	7.4	7.8	7.4	7.7	8.3	10.3	12.4	14.0	13.2
1999.....	6.7	7.4	7.2	8.1	7.5	8.7	10.9	12.9	13.5	12.6

All races 3/, male

2006.....	11.2	13.1	14.3	14.3	14.4	16.3	19.2	26.4	29.7	32.7
2005.....	11.8	13.7	14.6	14.1	15.3	15.5	20.2	26.6	33.3	33.8
2004.....	11.7	13.4	13.6	13.8	14.9	15.5	20.0	26.7	30.7	34.4
2003.....	12.1	13.2	14.5	14.6	15.3	16.5	20.7	26.3	32.4	36.6
2002.....	12.6	13.3	14.0	15.4	14.8	17.0	22.5	27.8	35.9	37.8
2001.....	12.0	12.8	13.8	14.9	14.2	17.3	21.4	27.7	35.5	38.9
2000.....	12.6	12.8	13.3	13.1	13.9	15.9	21.3	28.1	35.4	42.9
1999.....	11.2	12.6	12.4	14.0	14.0	16.9	22.4	29.1	34.6	41.6

All races 3/, female

2006.....	2.3	2.7	2.6	2.3	2.4	1.8	1.4	1.1	1.3	0.7
2005.....	2.3	2.5	2.4	2.3	1.8	1.7	1.6	1.3	1.3	0.8
2004.....	2.6	2.8	2.7	2.1	2.3	1.4	1.7	1.7	1.0	*
2003.....	2.3	2.6	2.3	2.5	1.7	1.6	1.6	1.6	1.3	0.9
2002.....	2.3	2.4	2.1	2.3	2.0	1.8	1.3	1.5	1.2	0.8
2001.....	2.2	2.3	2.4	2.4	2.3	1.9	1.4	1.7	1.3	1.0
2000.....	2.4	2.2	2.5	2.1	2.1	1.8	1.6	1.4	1.4	1.1
1999.....	2.2	2.4	2.1	2.6	1.6	1.7	1.8	1.6	1.2	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

White, both sexes

2006 4/.....	7.7	9.0	9.5	9.3	9.0	9.5	10.5	13.0	13.4	11.7
2005 4/.....	8.0	9.2	9.5	9.2	9.2	9.2	10.9	13.1	14.7	12.2
2004 4/.....	8.1	9.2	9.2	8.7	9.3	8.8	10.8	13.3	13.3	12.1
2003 4/.....	8.3	9.1	9.4	9.5	9.1	9.5	11.1	12.9	14.1	12.9
2002.....	8.5	8.9	8.9	9.7	9.0	9.9	11.7	13.3	15.6	12.9
2001.....	7.9	8.5	9.0	9.4	8.9	10.0	11.1	13.4	15.0	13.2
2000.....	8.4	8.3	8.7	8.2	8.5	9.2	11.1	13.4	15.2	14.3
1999.....	7.5	8.3	8.0	8.9	8.2	9.5	11.8	13.9	14.4	13.6

White, male

2006 4/.....	12.6	15.0	16.2	16.1	15.8	17.9	21.2	28.6	32.3	35.1
2005 4/.....	13.3	15.5	16.3	15.9	17.0	17.4	22.0	28.7	36.0	36.8
2004 4/.....	13.1	15.1	15.5	15.1	16.6	16.9	21.6	28.7	32.9	37.5
2003 4/.....	13.8	15.1	16.2	16.4	16.8	18.3	22.5	28.3	34.8	39.9
2002.....	14.3	15.0	15.5	16.9	16.3	18.8	24.4	29.6	38.8	40.5
2001.....	13.3	14.5	15.4	16.3	15.7	19.0	22.9	29.7	37.8	42.1
2000.....	14.0	14.2	14.7	14.2	15.3	17.3	22.7	30.4	38.1	46.5
1999.....	12.5	14.0	13.7	15.1	15.2	18.3	24.0	31.2	36.8	44.9

White, female

2006 4/.....	2.7	3.1	3.0	2.7	2.7	2.0	1.6	1.3	1.4	0.7
2005 4/.....	2.7	2.9	2.8	2.7	2.0	1.9	1.7	1.5	1.4	0.9
2004 4/.....	3.0	3.3	3.1	2.5	2.6	1.7	1.9	2.0	1.1	*
2003 4/.....	2.7	3.1	2.6	2.9	2.0	1.8	1.8	1.8	1.4	0.9
2002.....	2.7	2.8	2.4	2.7	2.3	2.1	1.4	1.6	1.3	0.9
2001.....	2.4	2.6	2.8	2.7	2.7	2.0	1.5	1.8	1.3	1.1
2000.....	2.7	2.5	2.8	2.4	2.3	2.1	1.8	1.5	1.5	1.2
1999.....	2.5	2.7	2.5	3.0	1.8	1.8	2.1	1.8	1.3	1.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,Y87.0)

All races 3/, both sexes

2006.....	1.0	1.2	1.3	1.0	1.0	0.8	0.6	0.6	0.6	0.8
2005.....	1.1	1.2	1.0	0.8	0.9	0.6	0.6	0.7	0.9	1.1
2004.....	1.0	1.1	1.1	0.8	0.8	0.7	0.8	0.5	0.8	0.9
2003.....	1.1	1.0	1.0	0.9	0.8	0.5	0.6	0.6	0.8	0.7
2002.....	1.0	1.0	1.0	0.9	0.7	0.6	0.8	0.8	0.9	1.4
2001 5/.....	0.8	1.1	0.9	0.8	0.6	0.6	0.7	0.8	1.0	1.0
2000.....	0.9	0.9	0.8	0.9	0.7	0.6	0.7	0.8	1.0	1.2
1999.....	0.9	0.9	0.9	0.6	0.6	0.7	0.7	0.6	1.1	1.3

All races 3/, male

2006.....	1.6	1.8	1.9	1.4	1.4	1.4	1.0	0.7	1.0	2.1
2005.....	1.6	2.0	1.5	1.2	1.3	1.0	0.9	1.2	1.9	2.2
2004.....	1.5	1.6	1.8	1.2	1.1	1.0	1.3	0.9	1.4	2.2
2003.....	1.6	1.6	1.5	1.3	1.1	0.7	1.0	1.1	1.5	1.7
2002.....	1.6	1.6	1.6	1.3	1.0	0.9	1.2	1.3	1.5	3.1
2001 5/.....	1.2	1.8	1.3	1.2	1.1	0.9	1.0	1.2	1.8	2.5
2000.....	1.4	1.4	1.1	1.4	0.9	1.0	1.0	1.4	1.8	2.4
1999.....	1.3	1.3	1.3	1.0	0.9	1.0	1.0	1.1	1.9	2.5

All races 3/, female

2006.....	0.5	0.6	0.7	0.6	0.6	*	*	0.5	*	*
2005.....	0.6	0.5	0.5	0.5	0.4	*	*	*	*	0.6
2004.....	0.5	0.7	0.6	0.5	0.4	0.4	0.4	*	*	*
2003.....	0.6	0.5	0.5	0.5	0.5	*	*	*	*	*
2002.....	0.5	0.4	0.4	0.5	0.5	*	0.5	*	*	0.7
2001 5/.....	0.5	0.5	0.5	0.5	*	*	0.4	0.5	*	*
2000.....	0.4	0.5	0.5	0.4	0.4	*	0.4	*	*	0.7
1999.....	0.6	0.4	0.5	0.3	0.4	0.5	0.5	*	*	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by all other and unspecified means and their sequelae (*U03,X71,X75-X79,X81-X84,Y87.0)

White, both sexes

2006 4/.....	1.1	1.3	1.4	1.0	1.0	0.8	0.7	0.6	0.7	0.9
2005 4/.....	1.1	1.3	1.0	0.9	0.9	0.6	0.6	0.7	0.9	1.0
2004 4/.....	1.0	1.1	1.2	0.8	0.8	0.7	0.8	0.5	0.7	0.9
2003 4/.....	1.2	1.1	1.1	1.0	0.8	0.5	0.6	0.5	0.8	0.7
2002.....	1.1	1.1	1.0	0.9	0.8	0.6	0.8	0.8	0.9	1.5
2001 5/.....	0.9	1.2	0.9	0.9	0.7	0.5	0.7	0.8	1.0	1.1
2000.....	0.9	1.0	0.8	0.9	0.7	0.6	0.7	0.9	1.0	1.2
1999.....	1.0	0.9	0.9	0.6	0.6	0.7	0.7	0.7	1.1	1.4

White, male

2006 4/.....	1.7	1.9	2.0	1.5	1.5	1.3	1.1	0.8	1.0	2.2
2005 4/.....	1.6	2.1	1.6	1.3	1.4	1.0	1.0	1.3	1.9	2.1
2004 4/.....	1.5	1.5	1.9	1.2	1.1	1.1	1.3	0.9	1.2	2.1
2003 4/.....	1.7	1.6	1.6	1.5	1.2	0.8	1.1	1.0	1.6	1.8
2002.....	1.6	1.7	1.7	1.4	1.1	0.9	1.2	1.3	1.4	3.3
2001 5/.....	1.3	1.8	1.3	1.3	1.1	0.8	1.0	1.2	1.9	2.8
2000.....	1.4	1.4	1.0	1.4	1.0	1.0	1.0	1.5	1.8	2.7
1999.....	1.3	1.4	1.4	1.0	0.9	0.9	1.0	1.2	2.0	2.8

White, female

2006 4/.....	0.5	0.7	0.8	0.5	0.6	*	*	*	*	*
2005 4/.....	0.6	0.5	0.5	0.5	0.4	*	*	*	*	*
2004 4/.....	0.5	0.7	0.6	0.5	0.4	0.4	*	*	*	*
2003 4/.....	0.7	0.6	0.6	0.5	0.5	*	*	*	*	*
2002.....	0.5	0.4	0.4	0.5	0.5	*	*	*	*	0.7
2001 5/.....	0.5	0.6	0.5	0.5	*	*	*	*	*	*
2000.....	0.4	0.5	0.5	0.4	0.4	*	0.5	*	*	*
1999.....	0.6	0.4	0.5	*	*	0.5	*	*	*	0.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
All races 3/, both sexes										
2006.....	6.3	5.6	4.5	3.4	2.9	2.0	2.2	2.0	2.3	1.9
2005.....	6.7	5.4	4.2	2.9	2.7	2.3	2.4	2.2	2.1	2.1
2004.....	6.3	5.3	4.2	3.2	2.8	2.6	2.2	2.2	2.2	2.1
2003.....	6.4	5.6	4.2	3.1	2.4	2.3	2.4	2.5	2.4	2.2
2002.....	6.5	5.4	4.0	3.5	2.7	2.4	2.2	2.3	2.4	2.1
2001 5/.....	8.6	7.0	5.5	4.4	3.5	3.3	2.5	2.5	2.5	2.4
2000.....	6.6	5.3	4.0	3.4	2.6	2.4	2.4	2.3	2.6	2.4
1999.....	6.6	5.1	4.0	3.2	2.8	2.7	2.4	2.5	2.4	2.4
All races 3/, male										
2006.....	9.4	8.5	6.7	5.2	4.4	2.9	3.2	2.9	2.3	2.3
2005.....	10.0	8.3	6.7	4.5	4.2	3.1	3.5	2.8	2.4	2.7
2004.....	9.0	7.8	6.7	4.9	4.1	3.8	3.3	2.6	2.7	2.7
2003.....	9.5	8.3	6.4	4.6	3.7	3.4	3.4	3.5	3.0	3.2
2002.....	9.6	8.0	6.0	5.3	4.2	3.4	3.2	3.3	2.9	3.0
2001 5/.....	12.6	10.6	8.3	6.7	5.0	4.6	3.3	3.2	3.0	3.2
2000.....	9.7	7.8	5.9	5.2	3.9	3.3	3.4	2.8	3.8	3.3
1999.....	9.5	7.7	5.8	5.1	4.3	4.2	3.1	3.4	3.4	3.4
All races 3/, female										
2006.....	3.3	2.8	2.5	1.7	1.6	1.3	1.4	1.4	2.3	1.8
2005.....	3.4	2.6	1.8	1.5	1.4	1.6	1.5	1.8	2.0	1.9
2004.....	3.6	3.0	1.9	1.6	1.6	1.6	1.3	1.9	2.0	1.9
2003.....	3.4	3.0	2.1	1.7	1.3	1.4	1.6	1.8	2.1	1.8
2002.....	3.5	2.9	2.2	1.8	1.4	1.5	1.4	1.6	2.1	1.7
2001 5/.....	4.7	3.6	2.7	2.3	2.2	2.1	1.9	1.9	2.2	2.0
2000.....	3.6	2.8	2.1	1.7	1.5	1.7	1.6	2.0	1.9	2.0
1999.....	3.6	2.5	2.2	1.4	1.5	1.5	1.9	1.9	1.9	2.0

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
White, both sexes										
2006 4/.....	4.3	3.9	3.2	2.5	2.4	1.5	1.9	1.7	2.2	1.7
2005 4/.....	4.4	3.8	2.9	2.3	2.1	1.8	1.9	2.0	1.9	1.9
2004 4/.....	4.3	3.8	3.0	2.4	2.3	2.2	1.8	1.8	2.0	1.9
2003 4/.....	4.2	3.9	3.1	2.4	2.0	1.9	1.9	2.2	2.1	2.0
2002.....	4.5	3.8	2.9	2.9	2.2	1.9	1.9	1.8	2.1	1.8
2001 5/.....	6.8	5.6	4.4	3.8	3.0	2.7	2.2	2.1	2.2	2.2
2000.....	4.4	3.7	2.9	2.7	2.0	1.8	2.0	2.1	2.0	2.2
1999.....	4.5	3.6	3.0	2.6	2.3	2.3	2.0	2.0	2.1	2.1
White, male										
2006 4/.....	6.1	5.7	4.5	3.5	3.5	1.9	2.4	2.3	2.1	2.0
2005 4/.....	6.2	5.6	4.5	3.3	3.3	2.2	2.7	2.2	2.0	1.9
2004 4/.....	5.8	5.2	4.6	3.6	3.2	3.0	2.6	2.0	2.2	2.3
2003 4/.....	5.8	5.4	4.5	3.3	2.9	2.7	2.6	2.9	2.5	3.0
2002.....	6.5	5.1	4.0	4.1	3.2	2.7	2.6	2.4	2.3	2.7
2001 5/.....	9.8	8.2	6.5	5.5	4.2	3.7	2.6	2.7	2.6	2.8
2000.....	6.4	5.1	4.0	4.0	2.8	2.2	2.6	2.4	2.5	2.8
1999.....	6.3	5.3	4.3	3.8	3.3	3.4	2.5	2.6	2.9	2.7
White, female										
2006 4/.....	2.4	2.2	2.0	1.5	1.3	1.2	1.4	1.3	2.3	1.6
2005 4/.....	2.6	2.0	1.4	1.3	1.1	1.4	1.3	1.8	1.8	1.9
2004 4/.....	2.9	2.3	1.4	1.3	1.4	1.6	1.1	1.6	1.9	1.8
2003 4/.....	2.7	2.4	1.7	1.6	1.0	1.2	1.3	1.6	1.8	1.6
2002.....	2.5	2.5	1.8	1.7	1.3	1.3	1.4	1.4	1.9	1.5
2001 5/.....	3.8	3.0	2.4	2.1	1.9	1.9	1.8	1.7	1.9	2.0
2000.....	2.5	2.3	1.8	1.4	1.3	1.4	1.5	1.8	1.7	1.9
1999.....	2.6	1.9	1.7	1.3	1.3	1.4	1.7	1.6	1.6	1.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)										
Black, both sexes										
2006 4/.....	20.5	17.1	14.5	10.7	8.0	6.6	5.4	5.0	*	*
2005 4/.....	21.6	17.2	13.9	8.3	7.8	7.1	7.1	5.8	5.6	*
2004 4/.....	20.2	16.8	13.4	9.4	7.9	5.7	6.4	7.1	5.2	*
2003 4/.....	21.4	17.7	12.7	9.1	6.8	5.7	6.9	6.6	6.6	*
2002.....	20.6	17.6	13.0	8.9	7.5	5.8	5.6	8.2	6.0	*
2001 5/.....	21.7	18.1	13.6	9.9	7.5	7.7	6.5	6.9	6.2	*
2000.....	21.8	16.8	12.4	9.1	7.4	8.2	6.8	5.3	9.6	*
1999.....	21.2	16.2	11.2	8.8	7.4	6.6	6.0	8.2	6.1	*
Black, male										
2006 4/.....	33.6	29.3	24.5	19.3	13.6	12.4	10.2	10.3	*	*
2005 4/.....	36.7	29.3	24.1	15.3	12.9	12.6	12.7	11.2	*	*
2004 4/.....	32.9	27.8	23.8	16.5	13.1	11.6	11.3	10.5	*	*
2003 4/.....	36.8	30.0	21.8	17.3	11.4	10.8	11.5	12.3	*	*
2002.....	32.4	30.8	22.5	16.8	13.5	10.0	11.3	15.2	*	*
2001 5/.....	34.2	30.0	23.1	17.4	12.0	13.2	11.8	11.6	*	*
2000.....	34.1	28.7	22.4	16.0	13.0	15.1	12.4	*	19.4	*
1999.....	34.1	27.3	18.5	17.0	12.7	13.0	10.2	13.8	*	*
Black, female										
2006 4/.....	8.9	6.4	6.1	3.6	3.6	*	*	*	*	*
2005 4/.....	8.3	6.7	5.2	2.5	3.8	*	*	*	*	*
2004 4/.....	8.9	7.2	4.6	3.6	3.8	*	*	*	*	*
2003 4/.....	7.9	7.0	5.0	2.3	3.2	*	*	*	*	*
2002.....	10.1	6.1	4.9	*	*	*	*	*	*	*
2001 5/.....	10.6	7.7	5.6	3.8	4.0	*	*	*	*	*
2000.....	10.9	6.5	3.8	3.5	*	*	*	*	*	*
1999.....	9.8	6.7	5.0	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All races 3/, both sexes

2006.....	3.9	3.1	2.4	1.6	1.4	0.9	0.8	0.7	0.6	0.4
2005.....	4.1	3.0	2.1	1.4	1.3	1.1	1.1	0.8	0.6	0.5
2004.....	3.8	2.9	2.2	1.7	1.5	1.2	0.8	0.8	0.7	0.6
2003.....	3.8	3.1	2.2	1.6	1.2	0.9	0.9	0.7	0.6	0.5
2002.....	3.9	2.8	2.2	1.8	1.3	1.0	0.8	1.0	0.8	*
2001.....	3.7	2.7	2.1	1.5	1.4	1.2	0.9	0.8	0.7	*
2000.....	3.8	2.9	2.1	1.7	1.2	1.0	1.0	0.7	0.5	0.7
1999.....	3.7	2.8	2.0	1.6	1.6	1.1	1.0	0.8	0.7	0.6

All races 3/, male

2006.....	6.2	4.9	3.6	2.3	2.1	1.2	1.3	1.0	*	*
2005.....	6.6	4.7	3.4	2.2	2.0	1.4	1.7	1.0	*	*
2004.....	5.9	4.4	3.5	2.7	2.2	1.8	1.4	1.0	*	*
2003.....	5.9	5.0	3.5	2.4	1.8	1.4	1.4	0.9	*	*
2002.....	6.2	4.2	3.4	2.7	1.9	1.4	1.0	1.4	1.0	*
2001.....	5.6	4.2	3.2	2.3	2.0	1.8	1.1	1.2	*	*
2000.....	5.8	4.5	3.3	2.6	1.8	1.3	1.3	0.8	*	*
1999.....	5.5	4.4	3.0	2.6	2.5	1.5	1.4	1.1	*	*

All races 3/, female

2006.....	1.6	1.3	1.2	0.9	0.7	0.6	0.4	*	0.7	*
2005.....	1.7	1.4	0.8	0.6	0.6	0.8	0.6	0.7	0.6	*
2004.....	1.6	1.4	0.9	0.7	0.9	0.7	*	0.7	0.8	*
2003.....	1.7	1.4	1.0	0.8	0.6	0.4	0.5	0.6	0.7	*
2002.....	1.7	1.5	1.1	0.9	0.7	0.6	0.6	0.6	0.7	*
2001.....	1.7	1.1	1.0	0.7	0.8	0.7	0.7	0.5	0.7	*
2000.....	1.7	1.3	1.0	0.9	0.7	0.8	0.7	0.7	*	*
1999.....	1.8	1.3	1.1	0.6	0.7	0.7	0.7	0.6	0.6	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

White, both sexes

2006 4/.....	2.5	2.0	1.5	1.2	1.1	0.6	0.6	0.5	0.7	*
2005 4/.....	2.6	2.1	1.5	1.1	1.0	0.9	0.9	0.8	0.5	*
2004 4/.....	2.5	2.0	1.5	1.3	1.2	1.1	0.7	0.7	0.7	0.5
2003 4/.....	2.4	2.2	1.7	1.2	1.0	0.7	0.8	0.6	0.6	0.5
2002.....	2.6	1.9	1.6	1.5	0.9	0.8	0.7	0.8	0.8	*
2001.....	2.6	1.9	1.5	1.2	1.2	1.0	0.7	0.7	0.7	*
2000.....	2.5	2.0	1.5	1.4	1.0	0.8	0.8	0.7	*	0.6
1999.....	2.5	1.9	1.5	1.2	1.2	1.0	0.9	0.7	0.7	0.6

White, male

2006 4/.....	3.9	2.8	2.2	1.6	1.7	0.6	0.9	*	*	*
2005 4/.....	3.9	3.0	2.3	1.6	1.6	1.1	1.2	0.9	*	*
2004 4/.....	3.5	2.9	2.3	1.9	1.7	1.4	1.0	0.9	*	*
2003 4/.....	3.4	3.1	2.5	1.8	1.4	1.1	1.1	0.7	*	*
2002.....	4.0	2.6	2.4	2.1	1.4	1.0	0.8	1.2	*	*
2001.....	3.7	2.8	2.3	1.6	1.7	1.3	0.8	1.0	*	*
2000.....	3.7	3.0	2.2	2.1	1.4	1.0	0.9	0.7	*	*
1999.....	3.6	2.8	2.3	1.9	1.8	1.3	1.1	0.7	*	*

White, female

2006 4/.....	1.2	1.2	0.9	0.8	0.6	0.6	*	*	0.8	*
2005 4/.....	1.4	1.1	0.7	0.6	0.5	0.7	0.6	0.8	*	*
2004 4/.....	1.5	1.1	0.8	0.7	0.8	0.8	*	0.6	0.9	*
2003 4/.....	1.3	1.2	0.9	0.8	0.5	*	0.5	0.5	0.7	*
2002.....	1.3	1.3	0.9	0.9	0.5	0.5	0.6	0.6	0.7	*
2001.....	1.4	0.9	0.8	0.8	0.7	0.8	0.7	0.5	0.7	*
2000.....	1.3	1.1	0.9	0.8	0.7	0.7	0.7	0.7	*	*
1999.....	1.4	1.0	0.7	0.6	0.7	0.7	0.7	0.6	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

Black, both sexes

2006 4/.....	13.7	10.3	8.4	4.5	3.7	3.3	*	*	*	*
2005 4/.....	14.4	9.9	6.9	3.7	3.3	2.8	3.3	*	*	*
2004 4/.....	12.8	9.4	7.2	5.2	4.2	2.6	*	*	*	*
2003 4/.....	13.7	10.2	6.8	4.6	3.1	2.4	*	*	*	*
2002.....	13.1	9.6	7.1	4.2	4.0	2.8	*	*	*	*
2001.....	11.6	8.4	6.5	4.3	3.0	2.9	*	*	*	*
2000.....	12.6	9.4	7.0	4.2	2.6	2.8	2.7	*	*	*
1999.....	12.2	9.7	5.9	4.5	4.2	2.2	*	*	*	*

Black, male

2006 4/.....	23.9	19.6	15.0	8.4	7.1	6.3	*	*	*	*
2005 4/.....	26.1	17.9	12.6	7.3	5.6	5.2	7.0	*	*	*
2004 4/.....	23.4	16.5	13.7	9.6	7.1	5.6	*	*	*	*
2003 4/.....	24.3	19.1	11.9	9.1	5.2	*	*	*	*	*
2002.....	22.7	17.4	12.3	8.3	6.5	5.2	*	*	*	*
2001.....	20.3	15.3	11.8	8.6	5.1	6.4	*	*	*	*
2000.....	21.6	16.7	13.1	7.1	5.4	5.2	*	*	*	*
1999.....	20.5	17.3	9.5	9.3	8.3	*	*	*	*	*

Black, female

2006 4/.....	4.6	2.2	2.9	*	*	*	*	*	*	*
2005 4/.....	4.2	3.0	2.1	*	*	*	*	*	*	*
2004 4/.....	3.5	3.2	1.7	*	*	*	*	*	*	*
2003 4/.....	4.4	2.5	2.5	*	*	*	*	*	*	*
2002.....	4.6	2.8	2.8	*	*	*	*	*	*	*
2001.....	3.9	2.4	1.9	*	*	*	*	*	*	*
2000.....	4.7	3.2	*	*	*	*	*	*	*	*
1999.....	4.8	3.2	2.8	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Assault (homicide) by sharp object (X99)										
All races 3/, both sexes										
2006.....	0.9	1.0	0.7	0.6	0.4	0.4	0.3	0.4	0.4	*
2005.....	1.0	1.0	0.7	0.5	0.4	0.4	0.5	0.3	*	*
2004.....	1.0	0.9	0.7	0.5	0.4	0.5	0.3	0.4	0.4	*
2003.....	1.0	0.9	0.7	0.5	0.4	0.4	0.5	0.5	0.4	*
2002.....	1.1	0.9	0.6	0.6	0.4	0.4	0.5	0.3	0.4	*
2001.....	0.9	0.8	0.6	0.5	0.3	0.6	0.3	0.5	*	*
2000.....	1.0	0.7	0.5	0.5	0.4	0.4	0.3	0.4	0.5	*
1999.....	1.0	0.8	0.6	0.5	0.4	0.3	0.3	0.4	*	*
All races 3/, male										
2006.....	1.2	1.4	1.0	0.9	0.7	0.5	*	*	*	*
2005.....	1.4	1.5	1.1	0.6	0.7	0.6	0.6	*	*	*
2004.....	1.4	1.2	1.0	0.7	0.6	0.8	*	*	*	*
2003.....	1.5	1.3	1.0	0.6	0.5	*	0.7	0.7	*	*
2002.....	1.5	1.3	1.0	1.0	0.5	0.5	0.8	*	*	*
2001.....	1.3	1.2	0.9	0.8	*	0.7	*	*	*	*
2000.....	1.4	1.0	0.6	0.8	0.5	0.5	0.6	0.7	*	*
1999.....	1.6	1.1	0.9	0.7	0.6	0.5	*	*	*	*
All races 3/, female										
2006.....	0.6	0.5	0.4	0.4	*	*	*	*	*	*
2005.....	0.5	0.5	0.3	0.3	*	*	*	*	*	*
2004.....	0.6	0.6	0.3	0.2	*	*	*	*	*	*
2003.....	0.6	0.5	0.3	0.3	*	*	*	*	*	*
2002.....	0.7	0.4	0.3	0.3	*	*	*	*	*	*
2001.....	0.6	0.4	0.3	0.3	0.3	0.5	*	0.5	*	*
2000.....	0.6	0.4	0.4	*	*	*	*	*	*	*
1999.....	0.4	0.4	0.3	*	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

All races 3/, both sexes

2006.....	1.1	1.1	1.1	0.9	0.9	0.6	0.8	0.7	0.8	1.0
2005.....	1.1	1.1	1.1	0.8	0.8	0.5	0.6	0.7	1.0	0.8
2004.....	1.0	1.2	1.1	0.8	0.7	0.7	0.8	0.7	0.7	0.9
2003.....	1.1	1.2	0.9	0.8	0.7	0.8	0.7	0.9	0.9	1.1
2002.....	1.1	1.2	0.8	0.8	0.8	0.8	0.7	0.8	0.7	0.9
2001 5/.....	3.4	3.1	2.5	2.1	1.4	1.1	0.9	0.8	1.0	1.1
2000.....	1.3	1.2	1.0	0.9	0.7	0.7	0.8	0.9	1.3	0.9
1999.....	1.3	1.1	1.0	0.8	0.7	0.8	0.7	0.9	1.0	1.1

All races 3/, male

2006.....	1.6	1.7	1.7	1.6	1.4	1.0	1.2	1.2	1.0	1.4
2005.....	1.6	1.8	1.8	1.4	1.3	0.8	0.9	1.2	1.3	*
2004.....	1.3	1.8	1.8	1.2	1.1	1.1	1.1	0.8	1.3	*
2003.....	1.7	1.7	1.5	1.3	1.2	1.3	1.0	1.5	1.5	2.0
2002.....	1.6	1.8	1.2	1.3	1.4	1.2	1.1	1.2	1.1	*
2001 5/.....	5.1	4.5	3.8	3.2	2.2	1.7	1.4	1.1	1.4	1.6
2000.....	1.9	1.8	1.6	1.3	1.3	1.2	1.3	1.1	2.0	*
1999.....	1.9	1.7	1.5	1.3	1.0	1.4	0.9	1.4	1.8	*

All races 3/, female

2006.....	0.5	0.5	0.5	0.3	0.5	*	0.5	*	0.6	0.8
2005.....	0.7	0.4	0.5	0.3	0.3	*	*	*	0.7	0.7
2004.....	0.7	0.6	0.5	0.4	0.3	*	0.5	0.6	*	0.8
2003.....	0.6	0.7	0.4	0.3	*	*	0.5	0.6	0.6	0.7
2002.....	0.6	0.6	0.4	0.4	*	0.4	*	0.5	*	0.8
2001 5/.....	1.7	1.6	1.2	1.1	0.8	0.6	0.4	0.6	0.8	0.9
2000.....	0.7	0.7	0.4	0.5	*	*	*	0.7	0.8	0.7
1999.....	0.8	0.6	0.5	0.4	*	*	0.5	0.6	*	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by all other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5,*U01.8-*U01.9,*U02,X92,X96-X98,Y01-Y03,Y08-Y09,Y87.1)

White, both sexes

2006 4/.....	0.8	0.9	0.9	0.7	0.8	0.5	0.7	0.7	0.7	0.9
2005 4/.....	0.8	0.9	0.8	0.7	0.6	0.3	0.4	0.6	0.9	0.7
2004 4/.....	0.8	0.9	0.9	0.7	0.5	0.6	0.7	0.6	0.7	0.9
2003 4/.....	0.9	0.9	0.7	0.6	0.6	0.6	0.6	0.9	0.8	1.1
2002.....	0.9	0.9	0.6	0.7	0.7	0.6	0.6	0.6	0.6	0.7
2001 5/.....	3.2	2.8	2.2	2.0	1.3	0.9	0.7	0.7	0.9	1.0
2000.....	0.9	0.9	0.7	0.7	0.5	0.5	0.6	0.7	1.0	0.8
1999.....	0.9	0.9	0.7	0.7	0.6	0.7	0.6	0.8	0.9	0.9

White, male

2006 4/.....	1.2	1.4	1.3	1.1	1.1	0.8	0.9	1.1	*	1.3
2005 4/.....	1.2	1.3	1.3	1.1	0.9	*	0.6	0.9	1.2	*
2004 4/.....	1.1	1.4	1.4	1.0	0.8	0.9	0.9	*	1.0	*
2003 4/.....	1.3	1.2	1.1	1.0	1.0	1.0	0.8	1.3	1.3	1.9
2002.....	1.3	1.3	0.8	1.1	1.2	0.9	1.0	0.8	*	*
2001 5/.....	4.9	4.2	3.3	3.1	1.8	1.5	1.1	0.9	1.2	*
2000.....	1.4	1.1	1.1	1.0	0.9	0.8	1.0	0.9	1.3	*
1999.....	1.4	1.3	1.0	1.1	0.9	1.1	0.7	1.3	1.5	*

White, female

2006 4/.....	0.5	0.4	0.4	0.3	0.4	*	*	*	0.6	0.7
2005 4/.....	0.5	0.4	0.3	*	*	*	*	*	0.7	0.6
2004 4/.....	0.6	0.5	0.4	0.3	*	*	*	*	*	0.8
2003 4/.....	0.5	0.6	0.4	0.3	*	*	*	0.5	*	0.7
2002.....	0.4	0.5	0.4	0.4	*	*	*	*	*	0.7
2001 5/.....	1.5	1.4	1.0	1.0	0.7	*	*	0.5	0.7	0.9
2000.....	0.5	0.6	0.4	0.4	*	*	*	0.6	0.7	*
1999.....	0.5	0.4	0.5	*	*	*	0.5	*	*	0.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
All races 3/, both sexes										
2006.....	3.2	3.5	2.8	1.8	1.2	0.9	0.8	1.0	1.1	1.4
2005.....	3.1	3.2	2.9	1.6	1.1	0.9	0.7	0.8	0.9	1.5
2004.....	3.4	3.6	2.6	1.7	1.0	0.9	0.8	0.7	1.3	1.6
2003.....	3.9	3.6	2.6	1.6	1.0	0.8	0.7	0.6	0.9	1.1
2002.....	3.4	3.3	2.3	1.4	1.0	0.8	0.7	0.6	1.0	1.4
2001.....	3.2	3.0	2.0	1.1	0.7	0.6	0.5	0.7	1.1	1.0
2000.....	3.1	2.7	1.7	1.1	0.7	0.6	0.6	0.6	0.7	1.3
1999.....	2.9	2.7	1.6	1.0	0.8	0.8	0.5	0.7	0.9	1.4
All races 3/, male										
2006.....	3.8	4.2	3.5	2.3	1.4	1.2	0.9	1.2	1.7	2.2
2005.....	3.6	3.8	3.5	2.1	1.1	1.1	1.0	1.0	*	1.6
2004.....	4.1	4.5	3.3	2.0	1.1	1.2	0.9	0.8	1.3	1.9
2003.....	4.9	4.6	3.4	2.0	1.2	0.8	0.7	0.8	1.1	1.5
2002.....	4.3	4.1	2.9	1.8	1.3	1.0	1.0	0.8	1.1	2.2
2001.....	4.4	4.2	2.5	1.5	0.8	0.9	0.6	0.9	1.5	1.5
2000.....	4.2	3.8	2.3	1.6	0.9	0.9	0.8	1.0	1.1	1.7
1999.....	4.0	3.7	2.1	1.5	1.1	1.1	0.6	1.0	1.3	2.1
All races 3/, female										
2006.....	2.6	2.7	2.0	1.5	0.9	0.7	0.7	0.8	0.8	1.0
2005.....	2.6	2.7	2.4	1.2	1.1	0.8	0.5	0.6	0.9	1.4
2004.....	2.7	2.6	2.0	1.3	0.8	0.7	0.6	0.6	1.2	1.6
2003.....	2.9	2.6	1.8	1.1	0.9	0.7	0.6	*	0.8	0.9
2002.....	2.6	2.6	1.7	1.0	0.7	0.7	0.5	0.5	0.9	1.1
2001.....	2.0	1.8	1.5	0.6	0.5	*	*	0.6	0.8	0.8
2000.....	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.2
1999.....	1.9	1.7	1.0	0.6	0.6	0.5	*	0.5	0.7	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Event of undetermined intent (Y10-Y34,Y87.2,Y89.9)										
White, both sexes										
2006 4/.....	3.3	3.5	2.7	1.7	1.2	0.9	0.7	0.8	1.1	1.1
2005 4/.....	3.1	3.3	2.9	1.6	1.1	0.9	0.7	0.7	0.9	1.5
2004 4/.....	3.5	3.7	2.6	1.7	1.0	0.8	0.8	0.6	1.3	1.6
2003 4/.....	4.0	3.6	2.5	1.5	0.9	0.7	0.6	0.6	0.9	1.2
2002.....	3.5	3.3	2.2	1.4	0.9	0.8	0.7	0.6	0.9	1.4
2001.....	3.2	2.9	1.9	1.1	0.6	0.6	0.5	0.7	1.0	1.0
2000.....	2.9	2.6	1.6	1.0	0.7	0.5	0.5	0.6	0.8	1.3
1999.....	2.8	2.5	1.4	1.0	0.8	0.8	0.4	0.7	0.9	1.4
White, male										
2006 4/.....	3.8	4.1	3.3	2.0	1.4	1.1	0.7	0.8	1.6	2.0
2005 4/.....	3.5	3.7	3.3	2.0	0.9	1.1	1.0	0.8	*	1.7
2004 4/.....	4.1	4.5	3.1	2.0	1.0	1.0	0.9	*	1.4	1.9
2003 4/.....	5.0	4.5	3.2	1.8	1.0	0.7	0.7	0.9	1.1	1.5
2002.....	4.3	4.0	2.5	1.8	1.2	0.9	0.8	*	1.1	2.3
2001.....	4.2	4.0	2.3	1.4	0.7	0.8	0.6	0.9	1.5	*
2000.....	4.0	3.5	2.1	1.3	0.9	0.8	0.7	1.0	*	*
1999.....	3.8	3.2	1.8	1.3	1.0	1.1	*	1.0	1.3	2.1
White, female										
2006 4/.....	2.9	2.9	2.1	1.5	1.0	0.7	0.7	0.8	0.7	0.7
2005 4/.....	2.6	2.9	2.5	1.3	1.2	0.8	0.5	0.6	0.8	1.3
2004 4/.....	2.9	2.9	2.0	1.4	0.9	0.6	0.6	0.6	1.2	1.5
2003 4/.....	3.1	2.8	1.9	1.2	0.8	0.7	0.5	*	0.8	1.0
2002.....	2.6	2.6	1.9	1.1	0.7	0.7	0.5	*	0.7	1.1
2001.....	2.1	1.9	1.6	0.7	0.6	*	*	0.6	0.7	0.8
2000.....	1.9	1.7	1.1	0.7	0.5	*	*	*	*	1.1
1999.....	1.7	1.8	1.0	0.7	0.7	0.5	*	*	0.8	1.1

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

All races 3/, both sexes

2006.....	2.5	2.6	2.0	1.3	0.6	0.4	*	*	*	*
2005.....	2.4	2.5	2.0	1.0	0.6	0.3	*	*	*	*
2004.....	2.6	2.8	1.9	1.1	0.5	0.3	*	*	*	*
2003.....	3.2	2.8	1.9	1.0	0.5	0.3	*	*	*	*
2002.....	2.7	2.6	1.7	0.8	0.5	0.3	*	*	*	*
2001.....	2.5	2.2	1.4	0.6	0.3	*	*	*	*	*
2000.....	2.4	2.1	1.1	0.5	0.3	*	*	*	*	*
1999.....	2.3	2.0	1.0	0.6	0.3	0.3	*	*	*	*

All races 3/, male

2006.....	2.7	2.9	2.4	1.3	0.5	0.5	*	*	*	*
2005.....	2.7	2.7	2.1	1.2	0.4	*	*	*	*	*
2004.....	3.1	3.4	2.0	1.1	0.5	*	*	*	*	*
2003.....	3.9	3.5	2.2	1.2	0.5	*	*	*	*	*
2002.....	3.2	3.0	1.9	0.9	0.6	*	*	*	*	*
2001.....	3.3	3.2	1.7	0.7	*	*	*	*	*	*
2000.....	3.3	2.8	1.5	0.7	*	*	*	*	*	*
1999.....	3.1	2.6	1.3	0.7	*	*	*	*	*	*

All races 3/, female

2006.....	2.2	2.3	1.7	1.2	0.7	0.4	*	*	*	*
2005.....	2.1	2.3	1.9	0.9	0.7	*	*	*	*	*
2004.....	2.2	2.2	1.7	1.1	0.5	*	*	*	*	*
2003.....	2.5	2.1	1.5	0.9	0.5	*	*	*	*	*
2002.....	2.2	2.2	1.5	0.7	0.4	*	*	*	*	*
2001.....	1.7	1.3	1.1	0.5	0.3	*	*	*	*	*
2000.....	1.6	1.4	0.8	0.4	*	*	*	*	*	*
1999.....	1.6	1.4	0.7	0.4	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Poisoning by and exposure to drugs and biological substances, undetermined intent (Y10-Y14)

White, both sexes

2006 4/.....	2.6	2.7	2.1	1.2	0.7	0.5	*	*	*	*
2005 4/.....	2.5	2.6	2.1	1.1	0.6	0.3	*	*	*	*
2004 4/.....	2.8	2.9	1.9	1.1	0.5	0.3	*	*	*	*
2003 4/.....	3.4	2.9	1.8	1.0	0.5	0.3	*	*	*	*
2002.....	2.7	2.6	1.6	0.8	0.5	0.3	*	*	*	*
2001.....	2.5	2.2	1.4	0.6	0.3	*	*	*	*	*
2000.....	2.3	2.0	1.1	0.5	0.3	*	*	*	*	*
1999.....	2.2	1.9	0.9	0.5	0.3	0.3	*	*	*	*

White, male

2006 4/.....	2.8	2.9	2.4	1.2	0.5	*	*	*	*	*
2005 4/.....	2.8	2.7	2.1	1.3	0.4	*	*	*	*	*
2004 4/.....	3.1	3.4	2.0	1.0	0.4	*	*	*	*	*
2003 4/.....	4.0	3.5	2.1	1.1	0.4	*	*	*	*	*
2002.....	3.3	3.0	1.7	0.8	0.6	*	*	*	*	*
2001.....	3.2	3.0	1.6	0.7	*	*	*	*	*	*
2000.....	3.1	2.7	1.4	0.6	*	*	*	*	*	*
1999.....	2.9	2.2	1.1	0.6	*	*	*	*	*	*

White, female

2006 4/.....	2.4	2.5	1.8	1.2	0.8	0.5	*	*	*	*
2005 4/.....	2.2	2.4	2.1	1.0	0.8	*	*	*	*	*
2004 4/.....	2.4	2.4	1.8	1.2	0.6	*	*	*	*	*
2003 4/.....	2.7	2.2	1.6	1.0	0.5	*	*	*	*	*
2002.....	2.2	2.2	1.6	0.7	0.5	*	*	*	*	*
2001.....	1.9	1.5	1.2	0.6	0.4	*	*	*	*	*
2000.....	1.5	1.3	0.8	0.4	*	*	*	*	*	*
1999.....	1.5	1.5	0.7	0.5	*	*	*	*	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2006.....	0.4	0.5	0.8	0.9	1.3	2.0	3.1	4.7	6.1	8.3
2005.....	0.4	0.5	0.8	1.0	1.4	2.0	3.3	4.7	7.2	9.1
2004.....	0.4	0.6	0.8	1.2	1.6	2.6	3.5	5.0	7.6	10.6
2003.....	0.5	0.7	0.7	1.1	1.7	2.2	3.7	5.3	7.3	11.2
2002.....	0.4	0.5	0.8	1.1	1.6	2.3	3.9	5.1	7.7	11.3
2001.....	0.4	0.7	0.8	1.3	1.9	2.6	3.6	5.9	8.3	12.8
2000.....	0.4	0.6	0.9	1.2	2.0	2.8	4.3	5.8	8.0	13.6
1999.....	0.4	0.5	0.7	1.1	1.7	2.4	4.0	5.5	8.4	12.7

All races 3/, male

2006.....	0.3	0.5	0.9	1.0	1.2	2.0	2.9	5.3	7.1	8.8
2005.....	0.4	0.5	0.8	1.1	1.7	2.3	3.7	4.6	8.9	10.8
2004.....	0.4	0.6	0.8	1.5	1.7	2.5	4.3	5.8	9.6	11.9
2003.....	0.5	0.7	0.7	1.3	1.6	2.3	4.2	5.6	8.8	12.9
2002.....	0.4	0.6	0.7	1.2	1.5	2.5	3.8	5.4	7.7	12.3
2001.....	0.4	0.7	0.8	1.3	2.0	2.6	3.8	6.3	9.9	14.5
2000.....	0.4	0.6	1.0	1.2	2.0	2.8	5.3	7.0	10.4	16.0
1999.....	0.4	0.4	0.7	1.2	1.7	2.3	4.5	6.4	10.1	16.5

All races 3/, female

2006.....	0.4	0.5	0.7	0.9	1.4	2.0	3.3	4.2	5.4	8.1
2005.....	0.4	0.6	0.9	1.0	1.2	1.8	3.0	4.8	6.2	8.3
2004.....	0.5	0.6	0.8	1.0	1.5	2.7	2.9	4.3	6.3	9.9
2003.....	0.4	0.6	0.6	0.9	1.7	2.1	3.3	5.1	6.4	10.5
2002.....	0.4	0.5	0.9	1.0	1.7	2.2	3.9	5.0	7.7	10.9
2001.....	0.5	0.6	0.8	1.2	1.7	2.6	3.5	5.7	7.3	12.1
2000.....	0.4	0.7	0.9	1.1	2.0	2.7	3.6	4.9	6.7	12.7
1999.....	0.4	0.5	0.7	0.9	1.6	2.5	3.6	4.8	7.4	11.2

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Complications of medical and surgical care (Y40-Y84,Y88)										
White, both sexes										
2006 4/.....	0.3	0.5	0.7	0.8	1.2	1.9	3.1	4.5	5.9	8.3
2005 4/.....	0.4	0.5	0.7	0.9	1.3	1.8	3.0	4.4	7.3	9.2
2004 4/.....	0.4	0.6	0.8	1.1	1.5	2.4	3.4	4.8	7.3	10.4
2003 4/.....	0.4	0.6	0.6	1.0	1.6	2.1	3.6	5.1	7.4	11.0
2002.....	0.4	0.5	0.7	1.0	1.5	2.2	3.8	5.0	7.6	11.2
2001.....	0.4	0.6	0.7	1.1	1.6	2.4	3.2	5.8	8.2	12.1
2000.....	0.3	0.6	0.9	1.0	1.8	2.6	4.1	5.4	8.0	13.1
1999.....	0.3	0.4	0.6	0.9	1.5	2.3	3.8	5.2	8.0	12.1
White, male										
2006 4/.....	0.3	0.4	0.8	0.8	1.1	2.0	2.8	5.1	7.2	8.8
2005 4/.....	0.4	0.4	0.7	0.9	1.5	2.3	3.4	4.3	8.8	11.2
2004 4/.....	0.3	0.6	0.8	1.3	1.7	2.5	4.2	5.8	9.5	11.8
2003 4/.....	0.6	0.6	0.5	1.2	1.6	2.3	4.1	5.5	9.1	12.8
2002.....	0.4	0.5	0.6	1.0	1.5	2.2	3.7	5.3	7.8	11.8
2001.....	0.3	0.6	0.6	1.1	1.7	2.5	3.2	6.2	10.0	13.6
2000.....	0.4	0.5	1.0	1.1	1.8	2.5	5.0	6.8	10.6	15.8
1999.....	0.3	0.3	0.6	1.0	1.6	2.3	4.2	6.2	9.7	15.6
White, female										
2006 4/.....	0.3	0.5	0.5	0.7	1.3	1.8	3.3	4.0	5.0	8.0
2005 4/.....	0.4	0.5	0.7	1.0	1.1	1.5	2.6	4.5	6.3	8.3
2004 4/.....	0.4	0.5	0.7	0.9	1.4	2.4	2.7	4.0	6.0	9.8
2003 4/.....	0.3	0.5	0.6	0.8	1.6	2.0	3.1	4.7	6.3	10.2
2002.....	0.3	0.4	0.7	0.9	1.5	2.1	3.9	4.8	7.5	10.9
2001.....	0.4	0.5	0.7	1.1	1.6	2.4	3.2	5.6	7.0	11.5
2000.....	0.3	0.6	0.9	0.9	1.8	2.7	3.3	4.5	6.4	12.0
1999.....	0.4	0.5	0.6	0.7	1.5	2.4	3.4	4.5	7.1	10.7

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Black, both sexes

2006 4/.....	0.7	1.0	1.7	2.3	3.1	3.9	4.3	8.2	9.5	10.4
2005 4/.....	*	1.0	2.1	2.2	2.9	4.5	6.4	9.4	8.5	9.1
2004 4/.....	0.9	1.0	1.4	2.7	2.5	4.5	5.5	7.6	11.9	13.8
2003 4/.....	*	1.3	1.6	2.5	3.0	3.6	5.8	9.5	7.6	15.2
2002.....	1.0	1.2	2.1	2.6	2.8	4.0	5.2	7.6	9.4	14.0
2001.....	1.0	1.6	1.5	2.7	4.2	4.6	8.5	8.0	11.1	23.3
2000.....	*	1.1	1.7	2.6	4.7	4.6	7.9	10.5	10.4	21.0
1999.....	0.8	0.9	1.8	2.6	2.6	3.5	7.3	9.5	12.4	20.1

Black, male

2006 4/.....	*	*	*	*	*	*	*	9.4	*	*
2005 4/.....	*	*	2.2	2.6	3.8	*	*	9.4	*	*
2004 4/.....	*	*	*	3.4	*	*	*	*	*	*
2003 4/.....	*	*	2.1	2.8	*	*	*	*	*	*
2002.....	*	*	*	*	*	5.0	*	*	*	*
2001.....	*	1.7	*	3.4	5.1	5.3	11.1	*	*	29.5
2000.....	*	*	*	*	4.6	6.2	9.4	10.4	*	*
1999.....	*	*	*	*	*	*	9.2	10.4	*	23.4

Black, female

2006 4/.....	*	*	1.9	2.6	2.9	4.2	*	7.5	9.4	10.2
2005 4/.....	*	*	2.1	*	*	4.8	6.7	9.3	7.4	9.0
2004 4/.....	*	*	1.7	*	*	5.4	5.1	7.5	11.0	13.4
2003 4/.....	*	*	*	*	3.0	3.8	6.4	10.6	7.7	15.4
2002.....	*	*	2.5	2.4	3.1	*	4.9	8.2	9.9	11.5
2001.....	*	*	*	*	3.5	4.1	6.7	8.0	11.0	21.0
2000.....	*	*	*	3.1	4.7	*	6.9	10.5	10.6	21.9
1999.....	*	*	*	*	*	*	6.1	8.9	10.8	18.8

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

All races 3/, both sexes

2006.....	0.2	0.4	0.6	0.7	1.0	1.6	2.5	4.0	5.2	7.4
2005.....	0.3	0.4	0.6	0.8	1.1	1.7	2.9	4.1	6.0	8.2
2004.....	0.3	0.4	0.7	1.0	1.2	2.2	3.1	4.1	6.2	9.3
2003.....	0.3	0.5	0.5	0.9	1.3	1.8	3.3	4.7	6.5	9.9
2002.....	0.3	0.4	0.7	0.8	1.4	2.0	3.3	4.6	6.8	10.4
2001.....	0.3	0.5	0.6	1.1	1.6	2.2	3.2	5.2	7.4	11.4
2000.....	0.3	0.5	0.8	0.9	1.8	2.4	3.8	5.0	7.0	12.2
1999.....	0.3	0.3	0.6	0.8	1.4	2.0	3.5	4.9	7.5	11.8

All races 3/, male

2006.....	0.2	0.4	0.7	0.7	1.0	1.6	2.3	4.6	6.1	7.9
2005.....	0.2	0.4	0.6	0.8	1.3	1.9	3.3	4.1	7.3	9.7
2004.....	0.3	0.4	0.7	1.2	1.4	2.2	3.8	4.7	8.4	10.7
2003.....	0.4	0.6	0.6	1.1	1.2	1.9	3.7	4.9	8.1	11.5
2002.....	0.3	0.4	0.6	0.9	1.3	2.1	3.3	4.9	7.0	11.7
2001.....	0.2	0.6	0.6	1.2	1.6	2.3	3.4	5.5	8.8	13.2
2000.....	0.3	0.5	0.8	1.0	1.8	2.5	4.7	6.0	8.9	14.3
1999.....	0.3	0.3	0.6	1.0	1.5	2.0	4.0	5.8	9.1	15.4

All races 3/, female

2006.....	0.2	0.3	0.5	0.7	1.0	1.6	2.7	3.6	4.6	7.1
2005.....	0.3	0.4	0.6	0.8	0.9	1.5	2.5	4.1	5.2	7.4
2004.....	0.3	0.4	0.6	0.8	1.0	2.2	2.5	3.7	4.9	8.7
2003.....	0.3	0.4	0.5	0.8	1.4	1.6	3.0	4.5	5.5	9.2
2002.....	0.4	0.4	0.7	0.8	1.5	1.9	3.3	4.4	6.7	9.9
2001.....	0.4	0.5	0.6	1.1	1.6	2.1	3.0	4.9	6.5	10.6
2000.....	0.3	0.5	0.7	0.9	1.8	2.3	3.1	4.4	5.9	11.4
1999.....	0.3	0.4	0.5	0.7	1.3	2.1	3.1	4.3	6.6	10.4

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

White, both sexes

2006 4/.....	0.2	0.3	0.5	0.6	0.9	1.5	2.5	3.8	5.0	7.3
2005 4/.....	0.2	0.4	0.5	0.7	1.0	1.5	2.5	3.8	6.0	8.2
2004 4/.....	0.3	0.4	0.6	0.9	1.2	2.1	3.0	4.0	6.1	9.1
2003 4/.....	0.3	0.4	0.4	0.8	1.3	1.7	3.2	4.4	6.6	9.6
2002.....	0.3	0.4	0.6	0.7	1.3	1.8	3.3	4.5	6.7	10.2
2001.....	0.3	0.5	0.5	1.0	1.4	2.0	2.8	5.1	7.2	10.7
2000.....	0.3	0.5	0.8	0.8	1.7	2.3	3.6	4.7	7.0	11.8
1999.....	0.3	0.3	0.5	0.7	1.3	2.0	3.4	4.7	7.2	11.2

White, male

2006 4/.....	*	0.3	0.6	0.7	0.9	1.6	2.3	4.5	6.1	7.9
2005 4/.....	0.2	0.4	0.5	0.7	1.1	1.8	3.1	3.9	7.2	10.0
2004 4/.....	0.2	0.4	0.7	1.1	1.4	2.2	3.7	4.7	8.4	10.4
2003 4/.....	0.4	0.5	0.4	1.0	1.3	2.0	3.6	4.8	8.4	11.5
2002.....	0.3	0.4	0.5	0.8	1.3	1.8	3.2	4.9	7.1	11.2
2001.....	*	0.6	0.5	1.0	1.4	2.2	2.9	5.5	8.9	12.3
2000.....	0.3	0.4	0.8	0.9	1.6	2.3	4.5	5.8	9.1	14.0
1999.....	0.2	*	0.5	0.7	1.5	1.9	3.8	5.6	8.7	14.5

White, female

2006 4/.....	*	0.3	0.4	0.6	0.9	1.4	2.8	3.4	4.3	7.0
2005 4/.....	0.2	0.4	0.5	0.7	0.9	1.2	2.1	3.8	5.3	7.4
2004 4/.....	0.3	0.3	0.5	0.7	1.0	1.9	2.3	3.4	4.7	8.4
2003 4/.....	0.2	0.4	0.4	0.6	1.3	1.5	2.8	4.2	5.5	8.8
2002.....	0.2	0.4	0.6	0.7	1.4	1.8	3.4	4.3	6.4	9.8
2001.....	0.4	0.4	0.6	1.0	1.5	1.9	2.8	4.9	6.2	10.0
2000.....	*	0.5	0.7	0.7	1.7	2.2	2.9	3.9	5.7	10.8
1999.....	0.3	0.4	0.5	0.6	1.2	2.0	3.0	4.0	6.2	9.9

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)

Black, both sexes

2006 4/.....	*	*	1.4	1.5	2.0	3.3	3.0	7.2	8.5	9.1
2005 4/.....	*	*	1.7	1.9	2.4	3.7	5.9	7.7	7.0	8.6
2004 4/.....	*	*	1.2	2.2	1.7	3.7	5.1	6.9	9.4	13.0
2003 4/.....	*	1.0	1.3	2.1	2.1	2.9	5.2	8.7	6.6	14.0
2002.....	0.9	0.9	1.7	1.8	2.3	3.8	4.2	6.7	9.2	13.4
2001.....	0.8	1.0	1.3	2.5	3.6	3.8	7.3	6.4	10.3	22.1
2000.....	*	0.8	1.2	2.4	3.6	4.0	6.7	9.8	9.0	19.1
1999.....	*	*	1.5	2.2	2.5	2.9	6.0	8.4	11.0	19.1

Black, male

2006 4/.....	*	*	*	*	*	*	*	*	*	*
2005 4/.....	*	*	*	*	*	*	*	*	*	*
2004 4/.....	*	*	*	2.6	*	*	*	*	*	*
2003 4/.....	*	*	*	*	*	*	*	*	*	*
2002.....	*	*	*	*	*	*	*	*	*	*
2001.....	*	*	*	3.3	4.7	*	9.4	*	*	28.4
2000.....	*	*	*	*	*	5.2	7.7	9.9	*	*
1999.....	*	*	*	*	*	*	8.1	9.5	*	*

Worktable 292R. Death rates for 358 selected causes, by 5-year age groups, race, and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other complications of medical and surgical care and their sequelae (Y70-Y84,Y88.2-Y88.3)										
Black, female										
2006 4/.....	*	*	*	*	*	*	*	6.4	8.7	8.8
2005 4/.....	*	*	*	*	*	3.9	6.1	7.5	*	8.3
2004 4/.....	*	*	*	*	*	4.5	4.8	7.0	8.7	12.6
2003 4/.....	*	*	*	*	*	*	6.0	9.5	*	15.0
2002.....	*	*	1.8	*	*	*	*	7.1	9.5	11.1
2001.....	*	*	*	*	*	*	5.8	6.4	10.2	19.8
2000.....	*	*	*	2.8	3.4	*	6.0	9.7	8.9	19.3
1999.....	*	*	*	*	*	*	4.5	7.7	9.5	18.0

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Includes races other than white and black.

4/ Multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2006, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data-mortality."

5/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02;

see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

Source: CDC/NCHS, National Vital Statistics System, Mortality.