

Person with disabilities experiencing problems accessing
NCHS birth and death tables should contact
nchsed@cdc.gov, or call 301-458-4688

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All causes

All origins 3/, both sexes.....	810.4	690.7	28.4	13.9	16.6	64.4	100.2	100.9	111.9	148.7
Male.....	814.8	756.3	30.5	15.4	19.6	90.7	148.0	143.4	150.4	189.0
Female.....	806.1	622.0	26.3	12.3	13.3	36.8	49.0	56.5	72.5	107.9
Hispanic, both sexes.....	300.1	590.6	26.4	11.8	16.6	65.3	95.0	79.4	83.7	108.2
Male.....	323.9	640.7	28.8	13.5	19.3	98.0	141.4	111.7	113.9	142.4
Female.....	274.6	538.3	24.0	10.1	13.7	30.5	40.0	38.7	47.7	69.0
Mexican, both sexes.....	260.5	572.6	27.1	12.1	17.9	70.5	97.3	79.5	82.5	104.4
Male.....	286.3	629.6	30.2	13.3	21.1	105.2	144.3	111.5	114.0	137.1
Female.....	232.1	515.6	24.1	10.9	14.6	32.7	39.4	38.4	44.3	64.7
Puerto Rican, both sexes.....	442.3	718.8	21.6	12.7	11.4	54.2	92.7	83.7	111.6	192.2
Male.....	502.9	*	23.8	14.6	*	78.4	142.7	132.8	158.3	249.5
Female.....	385.1	*	*	*	*	30.3	42.9	36.0	64.6	134.9
Cuban, both sexes.....	785.5	*	*	*	*	59.5	84.3	55.9	54.7	80.1
Male.....	800.9	*	*	*	*	*	*	*	*	*
Female.....	769.9	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	281.1	636.4	27.2	10.3	14.6	55.1	90.3	79.9	81.8	96.2
Male.....	293.6	723.6	26.4	13.9	16.8	84.2	131.3	107.3	105.1	130.7
Female.....	268.0	549.3	28.0	6.4	12.2	24.8	42.0	42.2	53.8	61.0
Non-Hispanic 5/, both sexes.....	897.1	714.6	28.9	14.4	16.5	64.1	100.9	106.0	118.5	156.3
Male.....	902.8	784.5	30.8	15.9	19.6	88.9	149.0	151.8	159.7	198.4
Female.....	891.7	641.4	26.9	12.8	13.2	38.0	50.6	60.3	77.8	114.9
Non-Hispanic white 6/, both sexes....	968.5	564.2	25.0	12.9	15.0	59.1	93.2	96.5	107.9	143.5
Male.....	962.0	621.9	26.7	14.2	18.0	79.0	136.5	137.2	145.2	183.0
Female.....	974.7	503.7	23.2	11.7	11.9	38.1	47.6	55.3	70.2	103.7
Non-Hispanic black 6/, both sexes....	759.1	1,339.2	44.3	20.6	23.1	87.1	146.7	171.3	201.6	262.6
Male.....	815.3	1,453.3	48.2	23.7	27.6	134.7	223.0	253.7	279.9	336.8
Female.....	708.0	1,220.1	40.3	17.3	18.4	38.2	68.6	93.5	130.9	196.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Septicemia (A40-A41)

All origins 3/, both sexes.....	11.4	6.5	0.5	0.2	0.2	0.2	0.4	0.6	0.8	1.5
Male.....	10.5	6.8	0.5	0.2	0.3	0.2	0.4	0.6	0.9	1.5
Female.....	12.3	6.2	0.6	*	*	0.3	0.4	0.6	0.8	1.6
Hispanic, both sexes.....	4.1	6.9	*	*	*	*	0.5	*	0.5	1.0
Male.....	4.1	7.5	*	*	*	*	*	*	*	*
Female.....	4.2	6.3	*	*	*	*	*	*	*	*
Mexican, both sexes.....	3.6	6.5	*	*	*	*	*	*	*	0.9
Male.....	3.5	7.0	*	*	*	*	*	*	*	*
Female.....	3.7	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	7.7	*	*	*	*	*	*	*	*	*
Male.....	8.7	*	*	*	*	*	*	*	*	*
Female.....	6.8	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	10.3	*	*	*	*	*	*	*	*	*
Male.....	9.3	*	*	*	*	*	*	*	*	*
Female.....	11.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.4	*	*	*	*	*	*	*	*	*
Male.....	3.3	*	*	*	*	*	*	*	*	*
Female.....	3.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	12.7	6.3	0.6	0.2	0.2	0.2	0.4	0.6	0.9	1.6
Male.....	11.7	6.5	0.6	*	0.3	0.2	0.4	0.7	0.9	1.6
Female.....	13.6	6.2	0.5	*	0.3	0.4	0.6	0.9	1.7	1.7
Non-Hispanic white 6/, both sexes....	12.7	4.0	0.5	0.2	0.2	0.2	0.3	0.5	0.7	1.3
Male.....	11.7	4.1	0.5	*	*	*	*	0.5	0.7	1.3
Female.....	13.7	3.9	0.4	*	*	*	*	0.4	0.7	1.2
Non-Hispanic black 6/, both sexes....	16.0	15.6	*	*	*	*	1.0	1.5	1.9	4.0
Male.....	14.9	16.3	*	*	*	*	*	*	2.2	3.7
Female.....	17.0	14.8	*	*	*	*	*	1.6	1.6	4.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins 3/, both sexes.....	4.0	*	*	*	*	0.2	0.8	2.0	3.9	7.5
Male.....	5.9	*	*	*	*	0.2	0.9	2.4	4.7	10.1
Female.....	2.2	*	*	*	*	*	0.7	1.6	3.0	4.9
Hispanic, both sexes.....	3.6	*	*	*	*	*	0.6	1.5	3.4	7.2
Male.....	5.6	*	*	*	*	*	*	2.0	4.9	10.5
Female.....	1.6	*	*	*	*	*	*	*	1.6	3.3
Mexican, both sexes.....	2.1	*	*	*	*	*	*	1.3	2.6	5.0
Male.....	3.5	*	*	*	*	*	*	1.8	4.1	8.5
Female.....	0.6	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	14.7	*	*	*	*	*	*	*	*	27.9
Male.....	20.7	*	*	*	*	*	*	*	*	34.3
Female.....	9.0	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	6.7	*	*	*	*	*	*	*	*	*
Male.....	12.1	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.4	*	*	*	*	*	*	2.3	4.5	6.6
Male.....	5.1	*	*	*	*	*	*	*	6.8	9.2
Female.....	1.6	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.1	*	*	*	*	0.2	0.8	2.1	4.0	7.5
Male.....	6.0	*	*	*	*	*	0.9	2.5	4.7	10.0
Female.....	2.3	*	*	*	*	*	0.7	1.8	3.3	5.1
Non-Hispanic white 6/, both sexes....	1.8	*	*	*	*	*	0.2	0.5	1.3	3.2
Male.....	3.0	*	*	*	*	*	*	0.7	1.8	4.9
Female.....	0.6	*	*	*	*	*	*	*	0.7	1.4
Non-Hispanic black 6/, both sexes....	17.9	*	*	*	*	0.9	3.7	10.4	18.6	33.1
Male.....	24.4	*	*	*	*	*	4.3	11.9	21.0	41.8
Female.....	12.1	*	*	*	*	*	3.1	9.0	16.5	25.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms (C00-C97)

All origins 3/, both sexes.....	187.0	1.8	2.3	2.3	2.2	3.2	4.6	6.7	11.5	20.8
Male.....	196.6	1.8	2.5	2.6	2.4	3.8	5.4	7.0	10.3	17.2
Female.....	177.6	1.8	2.1	2.1	2.0	2.5	3.7	6.4	12.7	24.4
Hispanic, both sexes.....	60.1	*	2.2	2.2	2.4	3.4	5.3	5.8	8.6	15.5
Male.....	60.4	*	2.6	2.4	2.2	3.8	6.4	5.6	6.5	11.0
Female.....	59.7	*	1.8	2.0	2.5	2.9	4.0	6.1	11.0	20.6
Mexican, both sexes.....	50.1	*	2.2	2.1	2.8	3.8	5.6	5.8	8.4	15.0
Male.....	49.9	*	2.5	2.0	2.6	4.2	6.4	5.6	6.4	10.9
Female.....	50.3	*	1.8	2.2	2.9	3.4	4.7	6.0	10.7	20.1
Puerto Rican, both sexes.....	84.9	*	*	*	*	*	*	7.0	9.8	23.1
Male.....	94.0	*	*	*	*	*	*	*	*	*
Female.....	76.2	*	*	*	*	*	*	*	*	32.8
Cuban, both sexes.....	172.3	*	*	*	*	*	*	*	*	*
Male.....	193.8	*	*	*	*	*	*	*	*	*
Female.....	150.4	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	61.3	*	*	*	*	2.6	5.5	5.4	8.8	14.9
Male.....	58.2	*	*	*	*	*	7.4	5.4	6.0	11.1
Female.....	64.7	*	*	*	*	*	*	*	12.3	18.8
Non-Hispanic 5/, both sexes.....	208.7	1.8	2.3	2.4	2.1	3.1	4.4	6.9	12.2	21.8
Male.....	221.3	1.7	2.5	2.6	2.4	3.8	5.2	7.4	11.3	18.5
Female.....	196.7	1.9	2.2	2.1	1.8	2.4	3.6	6.5	13.1	25.1
Non-Hispanic white 6/, both sexes....	227.1	2.1	2.3	2.4	2.1	3.1	4.3	6.7	11.9	21.1
Male.....	239.9	1.9	2.5	2.7	2.3	3.8	5.2	7.4	11.3	18.4
Female.....	214.7	2.2	2.1	2.2	1.8	2.4	3.4	5.9	12.6	23.8
Non-Hispanic black 6/, both sexes....	165.5	*	2.5	2.3	2.2	3.1	5.1	8.4	15.8	29.3
Male.....	179.2	*	2.7	2.6	2.5	3.9	5.5	8.0	13.5	23.0
Female.....	153.0	*	2.4	2.1	1.8	2.3	4.7	8.9	17.8	34.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All origins 3/, both sexes.....	17.9	*	*	*	*	*	0.2	0.5	1.1	2.0
Male.....	18.3	*	*	*	*	*	*	0.5	1.1	2.1
Female.....	17.5	*	*	*	*	*	0.2	0.5	1.1	1.9
Hispanic, both sexes.....	6.2	*	*	*	*	*	*	*	0.6	1.5
Male.....	6.3	*	*	*	*	*	*	*	*	1.3
Female.....	6.1	*	*	*	*	*	*	*	*	1.6
Mexican, both sexes.....	4.9	*	*	*	*	*	*	*	*	1.4
Male.....	5.2	*	*	*	*	*	*	*	*	*
Female.....	4.6	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	9.8	*	*	*	*	*	*	*	*	*
Male.....	10.0	*	*	*	*	*	*	*	*	*
Female.....	9.6	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	21.5	*	*	*	*	*	*	*	*	*
Male.....	21.8	*	*	*	*	*	*	*	*	*
Female.....	21.1	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.1	*	*	*	*	*	*	*	*	*
Male.....	5.7	*	*	*	*	*	*	*	*	*
Female.....	6.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	19.9	*	*	*	*	*	0.1	0.5	1.2	2.1
Male.....	20.4	*	*	*	*	*	*	0.5	1.2	2.3
Female.....	19.4	*	*	*	*	*	*	0.5	1.2	1.9
Non-Hispanic white 6/, both sexes....	21.2	*	*	*	*	*	*	0.5	1.1	2.0
Male.....	21.6	*	*	*	*	*	*	0.4	1.2	2.1
Female.....	20.7	*	*	*	*	*	*	0.5	1.0	1.9
Non-Hispanic black 6/, both sexes....	18.1	*	*	*	*	*	*	0.7	2.0	3.2
Male.....	18.9	*	*	*	*	*	*	*	1.8	3.7
Female.....	17.3	*	*	*	*	*	*	*	2.1	2.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of breast (C50)

All origins 3/, both sexes.....	13.8	*	*	*	*	*	*	0.4	1.5	3.8
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	26.9	*	*	*	*	*	*	0.7	3.0	7.5
Hispanic, both sexes.....	4.7	*	*	*	*	*	*	*	1.3	2.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	9.6	*	*	*	*	*	*	*	2.8	5.5
Mexican, both sexes.....	3.8	*	*	*	*	*	*	*	1.3	2.3
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	7.9	*	*	*	*	*	*	*	2.9	5.1
Puerto Rican, both sexes.....	7.1	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	13.7	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	12.0	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	24.2	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.0	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	10.2	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	15.3	*	*	*	*	*	*	0.4	1.5	4.0
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	29.7	*	*	*	*	*	*	0.7	3.0	7.9
Non-Hispanic white 6/, both sexes....	16.2	*	*	*	*	*	*	0.3	1.3	3.4
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	31.4	*	*	*	*	*	*	0.5	2.5	6.8
Non-Hispanic black 6/, both sexes....	15.1	*	*	*	*	*	*	0.8	3.4	8.0
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	28.5	*	*	*	*	*	*	1.6	6.4	15.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasm of cervix uteri (C53)

All origins 3/, both sexes.....	1.3	*	*	*	*	*	*	0.2	0.7	1.2
Male.....
Female.....	2.6	*	*	*	*	*	*	0.5	1.4	2.4
Hispanic, both sexes.....	1.0	*	*	*	*	*	*	*	0.6	1.3
Male.....
Female.....	2.2	*	*	*	*	*	*	*	1.2	2.7
Mexican, both sexes.....	1.0	*	*	*	*	*	*	*	*	1.3
Male.....
Female.....	2.0	*	*	*	*	*	*	*	*	2.8
Puerto Rican, both sexes.....	1.8	*	*	*	*	*	*	*	*	*
Male.....
Female.....	3.5	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	1.0	*	*	*	*	*	*	*	*	*
Male.....
Female.....	2.1	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.4	*	*	*	*	*	*	0.2	0.7	1.1
Male.....
Female.....	2.7	*	*	*	*	*	*	0.4	1.4	2.3
Non-Hispanic white 6/, both sexes....	1.3	*	*	*	*	*	*	0.2	0.6	1.1
Male.....
Female.....	2.5	*	*	*	*	*	*	0.4	1.2	2.3
Non-Hispanic black 6/, both sexes....	2.1	*	*	*	*	*	*	*	1.2	1.6
Male.....
Female.....	3.9	*	*	*	*	*	*	*	2.3	3.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All origins 3/, both sexes.....	18.4	0.8	0.8	0.6	0.9	1.3	1.9	1.9	2.2	2.8
Male.....	20.3	*	0.9	0.7	1.0	1.6	2.3	2.3	2.6	3.1
Female.....	16.5	*	0.8	0.5	0.7	0.9	1.3	1.5	1.8	2.4
Hispanic, both sexes.....	7.0	*	0.7	0.6	1.0	1.8	2.2	1.9	2.0	2.3
Male.....	7.4	*	*	*	*	2.1	2.9	2.2	2.0	2.3
Female.....	6.6	*	*	*	*	1.4	1.4	1.6	2.0	2.3
Mexican, both sexes.....	5.9	*	*	*	1.1	2.1	2.3	1.9	1.9	2.4
Male.....	6.2	*	*	*	*	2.5	2.9	2.1	1.8	2.2
Female.....	5.7	*	*	*	*	1.8	*	1.7	1.9	2.5
Puerto Rican, both sexes.....	8.9	*	*	*	*	*	*	*	*	*
Male.....	10.7	*	*	*	*	*	*	*	*	*
Female.....	7.2	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	17.7	*	*	*	*	*	*	*	*	*
Male.....	18.2	*	*	*	*	*	*	*	*	*
Female.....	17.1	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.5	*	*	*	*	*	2.5	*	*	*
Male.....	7.8	*	*	*	*	*	*	*	*	*
Female.....	7.2	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	20.3	0.8	0.8	0.6	0.9	1.2	1.8	1.9	2.2	2.8
Male.....	22.7	*	0.9	0.7	1.0	1.5	2.2	2.3	2.7	3.3
Female.....	18.1	*	0.8	0.5	0.7	0.8	1.3	1.5	1.7	2.4
Non-Hispanic white 6/, both sexes....	22.6	*	0.9	0.6	0.8	1.2	1.7	2.0	2.1	2.7
Male.....	25.2	*	0.9	0.6	1.0	1.5	2.2	2.4	2.6	3.1
Female.....	20.2	*	0.9	0.5	0.6	0.7	1.2	1.5	1.6	2.3
Non-Hispanic black 6/, both sexes....	13.8	*	*	*	0.9	1.3	2.0	2.0	3.0	4.3
Male.....	15.5	*	*	*	*	1.6	2.3	2.2	3.7	5.1
Female.....	12.3	*	*	*	*	*	1.7	1.9	2.4	3.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All origins 3/, both sexes.....	21.1	0.5	0.7	0.7	0.5	1.1	1.3	1.6	1.9	2.7
Male.....	21.9	*	0.7	0.8	0.5	1.3	1.8	1.9	2.4	2.9
Female.....	20.2	*	0.6	0.6	0.5	0.9	0.8	1.3	1.4	2.5
Hispanic, both sexes.....	7.7	*	0.6	0.5	0.7	1.2	1.7	1.1	1.5	2.1
Male.....	7.6	*	*	*	*	1.3	2.5	1.4	2.0	1.9
Female.....	7.8	*	*	*	*	*	*	*	*	2.4
Mexican, both sexes.....	6.8	*	*	*	0.9	1.3	1.9	1.1	1.7	2.1
Male.....	6.7	*	*	*	*	*	2.5	1.4	2.2	2.0
Female.....	6.9	*	*	*	*	*	*	*	*	2.1
Puerto Rican, both sexes.....	9.9	*	*	*	*	*	*	*	*	*
Male.....	10.6	*	*	*	*	*	*	*	*	*
Female.....	9.2	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	17.1	*	*	*	*	*	*	*	*	*
Male.....	18.8	*	*	*	*	*	*	*	*	*
Female.....	15.4	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.9	*	*	*	*	*	*	*	*	*
Male.....	7.5	*	*	*	*	*	*	*	*	*
Female.....	8.4	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	23.4	*	0.7	0.7	0.5	1.1	1.3	1.7	2.0	2.8
Male.....	24.5	*	0.8	0.8	0.5	1.3	1.7	2.1	2.5	3.1
Female.....	22.2	*	0.6	0.6	0.4	0.8	0.8	1.4	1.6	2.5
Non-Hispanic white 6/, both sexes....	25.6	*	0.6	0.8	0.5	1.1	1.3	1.6	2.1	2.8
Male.....	27.0	*	0.7	0.9	0.5	1.3	1.8	2.1	2.6	3.1
Female.....	24.3	*	0.5	0.6	0.5	0.8	0.8	1.2	1.6	2.4
Non-Hispanic black 6/, both sexes....	17.5	*	1.0	0.7	*	1.1	1.4	2.2	2.2	3.3
Male.....	18.1	*	*	*	*	1.4	1.4	2.2	2.8	3.4
Female.....	16.9	*	*	*	*	*	1.3	2.3	1.7	3.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Anemias (D50-D64)

All origins 3/, both sexes.....	1.3	*	0.2	*	*	0.2	0.3	0.4	0.3	0.4
Male.....	1.1	*	*	*	*	0.2	0.3	0.4	0.3	0.3
Female.....	1.6	*	*	*	*	*	0.2	0.3	0.4	0.5
Hispanic, both sexes.....	0.4	*	*	*	*	*	*	*	*	*
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	0.5	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	0.3	*	*	*	*	*	*	*	*	*
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	0.4	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	1.8	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.5	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	0.6	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.5	*	0.2	*	*	0.2	0.3	0.4	0.4	0.4
Male.....	1.2	*	*	*	*	*	0.4	0.5	0.3	0.4
Female.....	1.7	*	*	*	*	*	0.3	0.4	0.4	0.5
Non-Hispanic white 6/, both sexes....	1.4	*	*	*	*	*	*	*	*	*
Male.....	1.1	*	*	*	*	*	*	*	*	*
Female.....	1.7	*	*	*	*	*	*	*	*	*
Non-Hispanic black 6/, both sexes....	2.4	*	*	*	*	0.7	1.4	2.0	2.1	2.3
Male.....	2.1	*	*	*	*	*	1.6	2.2	2.1	2.0
Female.....	2.6	*	*	*	*	*	*	1.9	2.2	2.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diabetes mellitus (E10-E14)

All origins 3/, both sexes.....	24.2	*	*	*	0.1	0.2	0.6	1.2	2.1	3.3
Male.....	24.4	*	*	*	*	0.2	0.7	1.3	2.4	4.0
Female.....	24.0	*	*	*	*	*	0.5	1.1	1.9	2.7
Hispanic, both sexes.....	14.2	*	*	*	*	*	*	0.7	1.1	1.9
Male.....	13.7	*	*	*	*	*	*	*	1.1	2.2
Female.....	14.7	*	*	*	*	*	*	*	*	1.5
Mexican, both sexes.....	13.7	*	*	*	*	*	*	0.9	1.1	1.8
Male.....	13.1	*	*	*	*	*	*	*	*	1.9
Female.....	14.3	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	22.4	*	*	*	*	*	*	*	*	*
Male.....	22.4	*	*	*	*	*	*	*	*	*
Female.....	22.4	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	28.7	*	*	*	*	*	*	*	*	*
Male.....	29.1	*	*	*	*	*	*	*	*	*
Female.....	28.2	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	10.2	*	*	*	*	*	*	*	*	*
Male.....	9.8	*	*	*	*	*	*	*	*	*
Female.....	10.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	25.9	*	*	*	*	0.2	0.6	1.3	2.4	3.6
Male.....	26.3	*	*	*	*	0.2	0.7	1.5	2.8	4.4
Female.....	25.5	*	*	*	*	*	0.6	1.1	2.1	2.9
Non-Hispanic white 6/, both sexes....	25.4	*	*	*	*	0.2	0.5	0.9	1.9	3.0
Male.....	26.3	*	*	*	*	*	0.5	1.1	2.1	3.6
Female.....	24.5	*	*	*	*	*	0.5	0.8	1.7	2.5
Non-Hispanic black 6/, both sexes....	33.6	*	*	*	*	*	1.3	3.5	5.5	8.2
Male.....	31.7	*	*	*	*	*	1.5	3.7	7.1	10.3
Female.....	35.3	*	*	*	*	*	*	3.3	4.1	6.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Major cardiovascular diseases (I00-I78)

All origins 3/, both sexes.....	275.1	12.3	1.3	0.7	1.1	2.2	4.3	7.2	13.2	23.4
Male.....	267.7	13.4	1.4	0.8	1.2	2.6	5.4	9.5	17.6	31.5
Female.....	282.4	11.2	1.3	0.6	0.9	1.7	3.1	4.8	8.8	15.3
Hispanic, both sexes.....	86.5	10.2	1.1	0.6	1.0	2.1	3.4	4.8	7.8	13.0
Male.....	87.1	11.1	1.3	*	1.2	2.7	4.4	6.5	10.7	18.0
Female.....	85.9	9.3	*	*	*	1.4	2.2	2.7	4.4	7.3
Mexican, both sexes.....	70.5	9.4	1.1	*	0.8	2.0	3.2	4.9	7.8	13.0
Male.....	72.5	11.0	*	*	*	2.4	4.4	6.6	10.7	16.7
Female.....	68.3	7.9	*	*	*	*	1.8	2.6	4.2	8.4
Puerto Rican, both sexes.....	134.7	*	*	*	*	*	*	*	11.2	21.2
Male.....	141.1	*	*	*	*	*	*	*	*	35.0
Female.....	128.7	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	302.9	*	*	*	*	*	*	*	*	*
Male.....	295.7	*	*	*	*	*	*	*	*	*
Female.....	310.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	79.1	*	*	*	*	2.6	3.4	4.6	6.9	10.4
Male.....	77.5	*	*	*	*	*	*	6.0	8.4	15.7
Female.....	80.9	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	307.3	12.9	1.4	0.7	1.1	2.2	4.5	7.8	14.5	25.4
Male.....	300.1	14.1	1.4	0.8	1.2	2.6	5.7	10.4	19.4	34.2
Female.....	314.1	11.7	1.4	0.7	1.0	1.7	3.3	5.2	9.7	16.7
Non-Hispanic white 6/, both sexes....	333.1	9.5	1.2	0.6	1.0	1.9	3.9	6.2	11.7	21.0
Male.....	322.3	10.2	1.1	0.7	1.2	2.3	5.1	8.4	16.0	28.8
Female.....	343.6	8.8	1.3	0.6	0.7	1.4	2.7	4.0	7.3	13.2
Non-Hispanic black 6/, both sexes....	255.2	26.3	2.2	1.0	1.7	3.8	7.9	16.3	32.1	55.0
Male.....	259.6	29.6	2.7	*	1.6	4.4	9.0	21.5	41.5	72.7
Female.....	251.3	22.8	1.8	*	1.8	3.2	6.7	11.4	23.7	39.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All origins 3/, both sexes.....	211.0	8.4	1.0	0.5	0.8	1.7	3.4	5.8	10.6	18.6
Male.....	214.0	8.8	1.1	0.5	0.9	2.2	4.4	7.9	14.7	25.9
Female.....	208.0	7.9	0.9	0.4	0.7	1.2	2.4	3.7	6.5	11.3
Hispanic, both sexes.....	65.3	6.5	0.8	*	0.7	1.3	2.2	3.4	5.9	9.4
Male.....	67.7	7.1	*	*	*	1.9	3.1	4.8	8.4	13.3
Female.....	62.6	5.9	*	*	*	*	*	1.7	2.8	5.0
Mexican, both sexes.....	51.4	5.8	0.9	*	*	1.1	2.1	3.3	6.1	9.0
Male.....	54.8	7.0	*	*	*	1.6	3.2	4.6	8.5	11.8
Female.....	47.5	*	*	*	*	*	*	1.7	3.1	5.6
Puerto Rican, both sexes.....	107.7	*	*	*	*	*	*	*	*	17.1
Male.....	113.5	*	*	*	*	*	*	*	*	28.3
Female.....	102.1	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	248.8	*	*	*	*	*	*	*	*	*
Male.....	245.7	*	*	*	*	*	*	*	*	*
Female.....	252.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	59.3	*	*	*	*	*	*	3.3	4.7	7.6
Male.....	60.6	*	*	*	*	*	*	4.8	6.4	11.5
Female.....	58.0	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	235.8	8.9	1.0	0.5	0.8	1.8	3.7	6.4	11.8	20.4
Male.....	240.3	9.4	1.1	0.5	0.9	2.2	4.6	8.7	16.3	28.5
Female.....	231.4	8.4	1.0	0.4	0.7	1.3	2.6	4.2	7.3	12.5
Non-Hispanic white 6/, both sexes....	257.4	6.5	0.9	0.4	0.7	1.5	3.2	5.2	9.7	17.4
Male.....	260.3	6.8	0.9	0.4	0.8	1.9	4.2	7.1	13.7	24.8
Female.....	254.7	6.1	0.9	0.4	0.5	1.0	2.1	3.2	5.6	9.9
Non-Hispanic black 6/, both sexes....	189.3	17.9	1.8	*	1.4	3.3	6.4	13.3	25.5	42.4
Male.....	198.8	18.8	2.1	*	1.5	3.9	7.5	17.7	34.2	57.0
Female.....	180.6	17.0	*	*	*	2.6	5.3	9.2	17.6	29.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),
Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Hypertensive heart disease (I11)										
All origins 3/, both sexes.....	9.9	*	*	*	*	*	0.2	0.5	0.9	2.2
Male.....	9.3	*	*	*	*	*	0.3	0.7	1.3	3.2
Female.....	10.6	*	*	*	*	*	*	0.3	0.5	1.1
Hispanic, both sexes.....	3.6	*	*	*	*	*	*	*	0.5	1.2
Male.....	3.7	*	*	*	*	*	*	*	*	1.8
Female.....	3.6	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	2.7	*	*	*	*	*	*	*	*	1.3
Male.....	2.7	*	*	*	*	*	*	*	*	1.6
Female.....	2.6	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	7.3	*	*	*	*	*	*	*	*	*
Male.....	7.4	*	*	*	*	*	*	*	*	*
Female.....	7.2	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	11.8	*	*	*	*	*	*	*	*	*
Male.....	12.1	*	*	*	*	*	*	*	*	*
Female.....	11.5	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.7	*	*	*	*	*	*	*	*	*
Male.....	3.8	*	*	*	*	*	*	*	*	*
Female.....	3.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	11.0	*	*	*	*	*	0.2	0.6	1.0	2.3
Male.....	10.3	*	*	*	*	*	0.4	0.8	1.5	3.4
Female.....	11.7	*	*	*	*	*	*	0.3	0.6	1.2
Non-Hispanic white 6/, both sexes....	10.3	*	*	*	*	*	0.2	0.3	0.6	1.7
Male.....	9.0	*	*	*	*	*	0.3	0.4	0.9	2.6
Female.....	11.4	*	*	*	*	*	*	*	*	0.7
Non-Hispanic black 6/, both sexes....	17.9	*	*	*	*	*	*	1.8	3.4	6.6
Male.....	19.5	*	*	*	*	*	*	2.8	4.7	9.3
Female.....	16.5	*	*	*	*	*	*	*	2.3	4.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Ischemic heart diseases (I20-I25)										
All origins 3/, both sexes.....	142.1	0.5	*	*	*	0.2	0.5	1.4	3.9	8.8
Male.....	152.2	*	*	*	*	0.3	0.7	2.1	6.1	13.2
Female.....	132.3	*	*	*	*	*	0.2	0.6	1.7	4.4
Hispanic, both sexes.....	47.2	*	*	*	*	*	*	0.7	1.9	4.4
Male.....	49.9	*	*	*	*	*	*	1.2	2.7	6.9
Female.....	44.4	*	*	*	*	*	*	*	*	1.6
Mexican, both sexes.....	36.4	*	*	*	*	*	*	*	2.0	4.1
Male.....	39.9	*	*	*	*	*	*	*	2.7	6.1
Female.....	32.6	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	80.3	*	*	*	*	*	*	*	*	*
Male.....	85.8	*	*	*	*	*	*	*	*	*
Female.....	75.1	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	192.1	*	*	*	*	*	*	*	*	*
Male.....	189.6	*	*	*	*	*	*	*	*	*
Female.....	194.7	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	42.4	*	*	*	*	*	*	*	*	4.1
Male.....	44.4	*	*	*	*	*	*	*	*	6.9
Female.....	40.2	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	158.2	0.6	*	*	*	0.2	0.5	1.5	4.4	9.7
Male.....	170.6	*	*	*	*	0.3	0.7	2.4	6.9	14.6
Female.....	146.4	*	*	*	*	*	0.2	0.7	1.8	4.9
Non-Hispanic white 6/, both sexes....	174.5	*	*	*	*	0.2	0.5	1.4	4.1	9.2
Male.....	187.5	*	*	*	*	0.3	0.7	2.3	6.7	14.0
Female.....	161.9	*	*	*	*	*	*	0.6	1.6	4.3
Non-Hispanic black 6/, both sexes....	116.5	*	*	*	*	*	0.8	2.4	6.5	15.3
Male.....	124.5	*	*	*	*	*	*	3.4	9.7	21.5
Female.....	109.2	*	*	*	*	*	*	1.5	3.5	9.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Acute myocardial infarction (I21-I22)

All origins 3/, both sexes.....	47.2	*	*	*	*	0.1	0.2	0.6	1.5	3.6
Male.....	51.6	*	*	*	*	*	0.2	0.9	2.3	5.2
Female.....	43.0	*	*	*	*	*	*	0.3	0.8	1.9
Hispanic, both sexes.....	15.7	*	*	*	*	*	*	*	0.8	1.6
Male.....	16.5	*	*	*	*	*	*	*	1.0	2.3
Female.....	14.9	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	13.1	*	*	*	*	*	*	*	0.9	1.7
Male.....	14.1	*	*	*	*	*	*	*	*	2.2
Female.....	12.0	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	22.1	*	*	*	*	*	*	*	*	*
Male.....	25.0	*	*	*	*	*	*	*	*	*
Female.....	19.5	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	65.2	*	*	*	*	*	*	*	*	*
Male.....	62.5	*	*	*	*	*	*	*	*	*
Female.....	68.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	12.7	*	*	*	*	*	*	*	*	*
Male.....	13.0	*	*	*	*	*	*	*	*	*
Female.....	12.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	52.6	*	*	*	*	*	0.2	0.7	1.7	4.0
Male.....	57.9	*	*	*	*	*	0.2	1.0	2.6	5.9
Female.....	47.6	*	*	*	*	*	*	0.4	0.8	2.1
Non-Hispanic white 6/, both sexes....	58.0	*	*	*	*	*	0.2	0.7	1.6	4.0
Male.....	64.0	*	*	*	*	*	*	0.9	2.6	6.0
Female.....	52.2	*	*	*	*	*	*	0.4	0.7	1.9
Non-Hispanic black 6/, both sexes....	38.7	*	*	*	*	*	*	1.0	2.4	5.4
Male.....	40.6	*	*	*	*	*	*	1.5	3.5	7.1
Female.....	37.0	*	*	*	*	*	*	*	1.5	3.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other forms of chronic ischemic heart disease (I20,I25)

All origins 3/, both sexes.....	93.5	*	*	*	*	*	0.3	0.8	2.3	5.1
Male.....	99.2	*	*	*	*	*	0.4	1.2	3.7	7.9
Female.....	88.0	*	*	*	*	*	*	0.3	0.9	2.3
Hispanic, both sexes.....	31.3	*	*	*	*	*	*	*	1.1	2.8
Male.....	33.3	*	*	*	*	*	*	*	1.7	4.6
Female.....	29.2	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	23.1	*	*	*	*	*	*	*	1.0	2.5
Male.....	25.5	*	*	*	*	*	*	*	1.5	3.8
Female.....	20.4	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	57.8	*	*	*	*	*	*	*	*	*
Male.....	60.7	*	*	*	*	*	*	*	*	*
Female.....	55.1	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	126.5	*	*	*	*	*	*	*	*	*
Male.....	126.8	*	*	*	*	*	*	*	*	*
Female.....	126.1	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	29.4	*	*	*	*	*	*	*	*	2.8
Male.....	31.1	*	*	*	*	*	*	*	*	*
Female.....	27.6	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	104.1	*	*	*	*	*	0.3	0.8	2.6	5.6
Male.....	111.1	*	*	*	*	*	0.5	1.3	4.2	8.6
Female.....	97.4	*	*	*	*	*	*	0.3	1.0	2.6
Non-Hispanic white 6/, both sexes....	114.9	*	*	*	*	*	0.3	0.7	2.4	5.1
Male.....	121.9	*	*	*	*	*	0.4	1.3	4.0	7.9
Female.....	108.1	*	*	*	*	*	*	*	0.9	2.3
Non-Hispanic black 6/, both sexes....	76.4	*	*	*	*	*	*	1.4	3.8	9.4
Male.....	82.3	*	*	*	*	*	*	1.8	5.9	13.8
Female.....	71.0	*	*	*	*	*	*	*	1.9	5.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other heart diseases (I26-I51)

All origins 3/, both sexes.....	56.9	7.7	0.9	0.4	0.7	1.5	2.7	3.9	5.6	7.4
Male.....	51.0	8.0	1.0	0.5	0.8	1.8	3.3	4.9	7.0	9.2
Female.....	62.6	7.5	0.9	0.4	0.6	1.1	2.0	2.8	4.2	5.6
Hispanic, both sexes.....	13.7	6.3	0.8	*	0.6	1.1	1.7	2.4	3.3	3.6
Male.....	13.5	6.9	*	*	*	1.4	2.4	3.2	4.6	4.4
Female.....	13.9	5.7	*	*	*	*	*	1.3	1.7	2.6
Mexican, both sexes.....	11.6	5.6	0.8	*	*	0.9	1.7	2.5	3.4	3.5
Male.....	11.7	6.8	*	*	*	*	2.6	3.4	4.8	4.1
Female.....	11.5	*	*	*	*	*	*	*	1.8	2.8
Puerto Rican, both sexes.....	19.0	*	*	*	*	*	*	*	*	*
Male.....	19.2	*	*	*	*	*	*	*	*	*
Female.....	18.8	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	43.9	*	*	*	*	*	*	*	*	*
Male.....	43.3	*	*	*	*	*	*	*	*	*
Female.....	44.5	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	12.6	*	*	*	*	*	*	*	2.8	2.9
Male.....	11.8	*	*	*	*	*	*	*	*	*
Female.....	13.4	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	64.3	8.2	1.0	0.4	0.7	1.5	2.9	4.2	6.1	8.1
Male.....	57.8	8.3	1.0	0.4	0.8	1.9	3.5	5.4	7.6	10.2
Female.....	70.4	8.0	0.9	0.4	0.7	1.1	2.3	3.1	4.7	6.1
Non-Hispanic white 6/, both sexes....	70.4	5.8	0.8	0.4	0.6	1.3	2.5	3.4	4.8	6.4
Male.....	62.1	5.7	0.8	0.4	0.7	1.6	3.1	4.3	5.9	8.0
Female.....	78.4	5.8	0.8	0.4	0.5	0.9	1.8	2.5	3.6	4.8
Non-Hispanic black 6/, both sexes....	52.0	16.8	1.7	*	1.2	2.9	5.1	9.0	14.8	19.5
Male.....	52.1	17.9	2.0	*	1.4	3.6	5.5	11.4	18.7	25.0
Female.....	51.9	15.8	*	*	*	2.1	4.6	6.7	11.4	14.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All origins 3/, both sexes.....	36.0	6.9	0.7	0.3	0.6	1.3	2.4	3.5	5.1	6.6
Male.....	34.0	6.9	0.8	0.4	0.7	1.7	3.0	4.6	6.3	8.2
Female.....	38.0	6.8	0.7	0.3	0.5	1.0	1.8	2.5	3.8	4.9
Hispanic, both sexes.....	9.2	5.3	0.6	*	*	0.9	1.5	2.2	2.9	3.3
Male.....	9.6	5.8	*	*	*	1.2	2.1	3.1	4.2	4.1
Female.....	8.8	4.6	*	*	*	*	*	1.1	1.4	2.3
Mexican, both sexes.....	7.5	4.6	*	*	*	*	1.4	2.3	3.2	3.3
Male.....	8.1	*	*	*	*	*	2.2	3.1	4.5	3.9
Female.....	6.9	*	*	*	*	*	*	*	*	2.5
Puerto Rican, both sexes.....	13.2	*	*	*	*	*	*	*	*	*
Male.....	13.1	*	*	*	*	*	*	*	*	*
Female.....	13.3	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	34.2	*	*	*	*	*	*	*	*	*
Male.....	34.9	*	*	*	*	*	*	*	*	*
Female.....	33.5	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	8.4	*	*	*	*	*	*	*	*	2.6
Male.....	8.5	*	*	*	*	*	*	*	*	*
Female.....	8.3	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	40.6	7.3	0.8	0.3	0.7	1.4	2.6	3.9	5.6	7.2
Male.....	38.5	7.2	0.8	0.4	0.8	1.8	3.2	5.0	6.9	9.0
Female.....	42.7	7.4	0.8	0.3	0.6	1.0	2.0	2.8	4.4	5.4
Non-Hispanic white 6/, both sexes....	43.7	5.4	0.6	0.3	0.6	1.2	2.2	3.1	4.4	5.7
Male.....	40.5	5.2	0.6	*	0.7	1.5	2.9	4.0	5.4	7.2
Female.....	46.7	5.6	0.6	*	0.5	0.9	1.5	2.2	3.4	4.2
Non-Hispanic black 6/, both sexes....	36.5	14.3	1.5	*	1.1	2.6	4.6	8.2	13.2	17.0
Male.....	38.2	14.8	1.7	*	1.3	3.2	4.9	10.7	16.5	21.6
Female.....	34.9	13.8	*	*	*	2.0	4.2	5.8	10.3	12.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

All origins 3/, both sexes.....	8.0	*	*	*	*	*	*	0.1	0.4	0.5
Male.....	6.4	*	*	*	*	*	*	*	0.5	0.7
Female.....	9.5	*	*	*	*	*	*	*	0.4	0.4
Hispanic, both sexes.....	2.8	*	*	*	*	*	*	*	*	*
Male.....	2.4	*	*	*	*	*	*	*	*	*
Female.....	3.3	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	2.6	*	*	*	*	*	*	*	*	*
Male.....	2.1	*	*	*	*	*	*	*	*	*
Female.....	3.1	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	4.3	*	*	*	*	*	*	*	*	*
Male.....	4.7	*	*	*	*	*	*	*	*	*
Female.....	3.9	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	5.9	*	*	*	*	*	*	*	*	*
Male.....	5.0	*	*	*	*	*	*	*	*	*
Female.....	6.8	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.4	*	*	*	*	*	*	*	*	*
Male.....	2.0	*	*	*	*	*	*	*	*	*
Female.....	2.8	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	8.8	*	*	*	*	*	*	0.1	0.5	0.6
Male.....	7.1	*	*	*	*	*	*	*	0.6	0.8
Female.....	10.5	*	*	*	*	*	*	*	0.4	0.4
Non-Hispanic white 6/, both sexes....	8.5	*	*	*	*	*	*	*	0.2	0.3
Male.....	6.5	*	*	*	*	*	*	*	*	0.4
Female.....	10.3	*	*	*	*	*	*	*	*	*
Non-Hispanic black 6/, both sexes....	12.8	*	*	*	*	*	*	*	2.0	2.3
Male.....	11.7	*	*	*	*	*	*	*	2.2	3.1
Female.....	13.9	*	*	*	*	*	*	*	1.7	1.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Cerebrovascular diseases (I60-I69)

All origins 3/, both sexes.....	45.8	3.4	0.3	0.2	0.2	0.3	0.6	0.9	1.7	3.4
Male.....	37.0	3.9	0.3	0.3	0.3	0.3	0.7	1.0	1.8	3.7
Female.....	54.4	2.9	0.4	*	0.2	0.4	0.6	0.8	1.6	3.1
Hispanic, both sexes.....	15.8	3.4	*	*	*	0.6	0.9	0.9	1.4	2.8
Male.....	14.3	*	*	*	*	*	*	1.2	1.6	3.6
Female.....	17.5	*	*	*	*	*	*	*	1.2	1.9
Mexican, both sexes.....	14.3	3.4	*	*	*	*	0.8	1.1	1.2	3.2
Male.....	13.2	*	*	*	*	*	*	1.4	1.7	4.0
Female.....	15.5	*	*	*	*	*	*	*	*	2.3
Puerto Rican, both sexes.....	19.0	*	*	*	*	*	*	*	*	*
Male.....	18.6	*	*	*	*	*	*	*	*	*
Female.....	19.4	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	41.0	*	*	*	*	*	*	*	*	*
Male.....	36.0	*	*	*	*	*	*	*	*	*
Female.....	46.0	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	14.8	*	*	*	*	*	*	*	*	*
Male.....	12.2	*	*	*	*	*	*	*	*	*
Female.....	17.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	50.9	3.5	0.4	0.2	0.2	0.3	0.6	0.9	1.8	3.5
Male.....	41.1	4.1	0.3	0.3	0.3	0.3	0.7	1.0	1.9	3.7
Female.....	60.3	2.8	0.4	*	*	0.3	0.5	0.8	1.7	3.3
Non-Hispanic white 6/, both sexes....	54.2	2.4	0.4	0.2	0.2	0.3	0.5	0.7	1.4	2.5
Male.....	42.6	2.6	*	*	*	*	0.6	0.7	1.5	2.5
Female.....	65.5	2.2	0.4	*	*	*	0.4	0.7	1.3	2.4
Non-Hispanic black 6/, both sexes....	44.7	7.7	*	*	*	*	1.0	1.8	3.9	8.8
Male.....	40.8	10.2	*	*	*	*	*	2.1	3.9	10.3
Female.....	48.3	*	*	*	*	*	*	1.6	3.8	7.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Influenza and pneumonia (J10-J18)

All origins 3/, both sexes.....	18.8	6.4	0.8	0.2	0.1	0.3	0.5	0.7	1.0	1.4
Male.....	17.4	7.7	0.8	*	*	0.3	0.6	0.8	0.9	1.5
Female.....	20.2	5.0	0.8	0.2	*	0.4	0.5	0.6	1.0	1.3
Hispanic, both sexes.....	6.7	6.2	0.7	*	*	*	*	0.6	0.7	0.8
Male.....	6.3	7.3	*	*	*	*	*	*	*	1.1
Female.....	7.1	5.1	*	*	*	*	*	*	*	*
Mexican, both sexes.....	5.6	6.9	*	*	*	*	*	*	*	*
Male.....	5.3	8.2	*	*	*	*	*	*	*	*
Female.....	6.0	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	11.8	*	*	*	*	*	*	*	*	*
Male.....	12.9	*	*	*	*	*	*	*	*	*
Female.....	10.7	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	15.0	*	*	*	*	*	*	*	*	*
Male.....	13.6	*	*	*	*	*	*	*	*	*
Female.....	16.5	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.6	*	*	*	*	*	*	*	*	*
Male.....	5.7	*	*	*	*	*	*	*	*	*
Female.....	7.4	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	20.9	6.3	0.8	0.2	0.1	0.4	0.5	0.7	1.1	1.5
Male.....	19.4	7.8	0.8	*	*	0.3	0.6	0.8	1.0	1.6
Female.....	22.3	4.8	0.8	*	*	0.4	0.4	0.6	1.2	1.4
Non-Hispanic white 6/, both sexes....	23.1	4.5	0.6	0.2	*	0.4	0.5	0.7	0.9	1.4
Male.....	21.2	5.8	0.6	*	*	0.3	0.6	0.7	0.9	1.4
Female.....	25.0	3.0	0.6	*	*	0.4	0.4	0.6	1.0	1.4
Non-Hispanic black 6/, both sexes....	13.9	13.7	1.4	*	*	*	1.0	1.1	1.8	2.5
Male.....	13.6	16.3	*	*	*	*	*	1.5	*	3.0
Female.....	14.1	10.9	*	*	*	*	*	*	2.3	2.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pneumonia (J12-J18)

All origins 3/, both sexes.....	18.5	5.9	0.7	0.1	*	0.3	0.5	0.7	1.0	1.4
Male.....	17.1	7.2	0.6	*	*	0.3	0.6	0.8	0.9	1.5
Female.....	19.9	4.6	0.7	*	*	0.3	0.4	0.6	1.0	1.3
Hispanic, both sexes.....	6.6	5.7	0.5	*	*	*	*	0.5	0.7	0.8
Male.....	6.2	6.4	*	*	*	*	*	*	*	1.1
Female.....	7.0	4.8	*	*	*	*	*	*	*	*
Mexican, both sexes.....	5.5	6.3	*	*	*	*	*	*	*	*
Male.....	5.2	7.0	*	*	*	*	*	*	*	*
Female.....	5.9	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	11.7	*	*	*	*	*	*	*	*	*
Male.....	12.9	*	*	*	*	*	*	*	*	*
Female.....	10.6	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	14.9	*	*	*	*	*	*	*	*	*
Male.....	13.5	*	*	*	*	*	*	*	*	*
Female.....	16.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.5	*	*	*	*	*	*	*	*	*
Male.....	5.6	*	*	*	*	*	*	*	*	*
Female.....	7.4	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	20.6	5.9	0.7	*	*	0.3	0.5	0.7	1.0	1.5
Male.....	19.1	7.4	0.7	*	*	0.3	0.6	0.8	0.9	1.5
Female.....	21.9	4.4	0.7	*	*	0.3	0.4	0.6	1.2	1.4
Non-Hispanic white 6/, both sexes....	22.7	4.1	0.6	*	*	0.4	0.4	0.6	0.9	1.3
Male.....	20.9	5.4	0.6	*	*	0.3	0.6	0.7	0.9	1.4
Female.....	24.6	2.8	0.6	*	*	0.4	0.3	0.6	1.0	1.3
Non-Hispanic black 6/, both sexes....	13.8	12.9	1.2	*	*	*	0.9	1.1	1.7	2.5
Male.....	13.5	15.7	*	*	*	*	*	1.5	*	3.0
Female.....	14.0	10.0	*	*	*	*	*	*	2.2	2.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Chronic lower respiratory diseases (J40-J47)

All origins 3/, both sexes.....	41.6	0.7	0.3	0.3	0.3	0.3	0.4	0.5	0.7	1.0
Male.....	40.2	0.9	0.3	0.3	0.4	0.4	0.5	0.6	0.7	1.0
Female.....	43.0	*	*	*	*	0.2	0.4	0.5	0.7	1.1
Hispanic, both sexes.....	7.5	*	*	*	*	*	*	*	*	0.6
Male.....	7.5	*	*	*	*	*	*	*	*	*
Female.....	7.5	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	5.4	*	*	*	*	*	*	*	*	*
Male.....	5.8	*	*	*	*	*	*	*	*	*
Female.....	5.1	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	13.4	*	*	*	*	*	*	*	*	*
Male.....	13.3	*	*	*	*	*	*	*	*	*
Female.....	13.5	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	30.0	*	*	*	*	*	*	*	*	*
Male.....	27.8	*	*	*	*	*	*	*	*	*
Female.....	32.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.4	*	*	*	*	*	*	*	*	*
Male.....	7.0	*	*	*	*	*	*	*	*	*
Female.....	7.8	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	47.5	0.8	0.3	0.3	0.3	0.3	0.5	0.6	0.8	1.1
Male.....	46.1	*	0.3	0.3	0.5	0.4	0.6	0.7	0.9	1.0
Female.....	48.8	*	*	*	*	0.2	0.4	0.5	0.8	1.2
Non-Hispanic white 6/, both sexes....	55.6	*	*	*	*	*	0.3	0.4	0.6	0.9
Male.....	53.0	*	*	*	*	*	0.4	0.4	0.5	0.8
Female.....	58.0	*	*	*	*	*	*	0.3	0.7	1.0
Non-Hispanic black 6/, both sexes....	20.3	*	*	0.9	1.2	1.1	1.3	1.6	2.4	2.7
Male.....	22.7	*	*	*	1.6	1.4	1.4	1.8	3.0	2.6
Female.....	18.1	*	*	*	*	*	*	1.4	1.9	2.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Asthma (J45-J46)

All origins 3/, both sexes.....	1.2	*	0.2	0.2	0.3	0.3	0.4	0.4	0.5	0.7
Male.....	0.9	*	*	0.3	0.4	0.3	0.4	0.5	0.5	0.6
Female.....	1.5	*	*	*	*	0.2	0.3	0.4	0.5	0.7
Hispanic, both sexes.....	0.6	*	*	*	*	*	*	*	*	*
Male.....	0.4	*	*	*	*	*	*	*	*	*
Female.....	0.8	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	0.4	*	*	*	*	*	*	*	*	*
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	0.5	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	1.9	*	*	*	*	*	*	*	*	*
Male.....	1.5	*	*	*	*	*	*	*	*	*
Female.....	2.3	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.6	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	0.8	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.3	*	0.2	0.2	0.3	0.3	0.4	0.5	0.6	0.7
Male.....	1.0	*	*	0.3	0.4	0.3	0.5	0.6	0.6	0.6
Female.....	1.6	*	*	*	*	*	0.4	0.4	0.6	0.8
Non-Hispanic white 6/, both sexes....	1.1	*	*	*	*	*	0.3	0.3	0.3	0.5
Male.....	0.7	*	*	*	*	*	*	0.4	*	0.4
Female.....	1.5	*	*	*	*	*	*	*	0.4	0.6
Non-Hispanic black 6/, both sexes....	2.5	*	*	0.8	1.1	1.1	1.2	1.4	2.0	2.2
Male.....	2.3	*	*	*	1.5	1.4	1.4	1.6	2.5	2.0
Female.....	2.7	*	*	*	*	*	*	*	1.6	2.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All origins 3/, both sexes.....	15.1	3.9	*	*	*	0.1	0.3	0.4	1.0	1.3
Male.....	15.0	4.8	*	*	*	*	0.4	0.4	1.2	1.4
Female.....	15.3	3.0	*	*	*	*	0.3	0.4	0.8	1.1
Hispanic, both sexes.....	5.8	3.7	*	*	*	*	*	*	0.7	0.7
Male.....	5.6	*	*	*	*	*	*	*	*	*
Female.....	6.1	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	5.4	3.7	*	*	*	*	*	*	*	0.9
Male.....	5.1	*	*	*	*	*	*	*	*	*
Female.....	5.7	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	8.6	*	*	*	*	*	*	*	*	*
Male.....	8.6	*	*	*	*	*	*	*	*	*
Female.....	8.5	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	15.2	*	*	*	*	*	*	*	*	*
Male.....	17.1	*	*	*	*	*	*	*	*	*
Female.....	13.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.6	*	*	*	*	*	*	*	*	*
Male.....	4.3	*	*	*	*	*	*	*	*	*
Female.....	4.9	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	16.7	4.0	*	*	*	0.1	0.4	0.5	1.0	1.4
Male.....	16.7	5.1	*	*	*	*	0.4	0.4	1.3	1.6
Female.....	16.8	2.9	*	*	*	*	0.3	0.5	0.8	1.2
Non-Hispanic white 6/, both sexes....	16.5	3.4	*	*	*	*	0.3	0.3	0.7	0.9
Male.....	16.7	4.6	*	*	*	*	0.4	*	1.0	1.0
Female.....	16.4	2.1	*	*	*	*	*	*	0.5	0.7
Non-Hispanic black 6/, both sexes....	22.1	6.8	*	*	*	*	0.8	1.4	2.6	4.2
Male.....	21.0	7.4	*	*	*	*	*	*	3.2	4.6
Female.....	23.0	*	*	*	*	*	*	1.5	2.2	3.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Renal failure (N17-N19)

All origins 3/, both sexes.....	14.5	3.7	*	*	*	0.1	0.3	0.4	0.9	1.2
Male.....	14.3	4.4	*	*	*	*	0.3	0.3	1.0	1.3
Female.....	14.6	3.0	*	*	*	*	0.3	0.4	0.7	1.1
Hispanic, both sexes.....	5.6	3.7	*	*	*	*	*	*	0.6	0.7
Male.....	5.4	*	*	*	*	*	*	*	*	*
Female.....	5.8	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	5.1	3.7	*	*	*	*	*	*	*	0.9
Male.....	4.9	*	*	*	*	*	*	*	*	*
Female.....	5.4	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	8.4	*	*	*	*	*	*	*	*	*
Male.....	8.6	*	*	*	*	*	*	*	*	*
Female.....	8.2	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	14.7	*	*	*	*	*	*	*	*	*
Male.....	16.3	*	*	*	*	*	*	*	*	*
Female.....	13.1	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.3	*	*	*	*	*	*	*	*	*
Male.....	4.0	*	*	*	*	*	*	*	*	*
Female.....	4.7	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	16.0	3.7	*	*	*	0.1	0.3	0.4	0.9	1.3
Male.....	15.9	4.6	*	*	*	*	0.4	0.4	1.2	1.4
Female.....	16.1	2.8	*	*	*	*	0.3	0.5	0.7	1.1
Non-Hispanic white 6/, both sexes....	15.8	3.1	*	*	*	*	0.3	0.3	0.7	0.8
Male.....	15.9	4.1	*	*	*	*	0.4	*	0.9	0.9
Female.....	15.7	2.1	*	*	*	*	*	*	0.4	0.7
Non-Hispanic black 6/, both sexes....	21.1	6.6	*	*	*	*	0.7	1.2	2.4	3.9
Male.....	20.0	7.1	*	*	*	*	*	*	2.7	4.4
Female.....	22.1	*	*	*	*	*	*	1.4	2.1	3.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy, childbirth and the puerperium (000-099)

All origins 3/, both sexes.....	0.3	*	0.2	0.7	0.8	0.8	0.7
Male.....
Female.....	0.5	*	0.3	1.4	1.7	1.6	1.3
Hispanic, both sexes.....	0.3	*	*	0.8	0.8	0.9	0.6
Male.....
Female.....	0.7	*	*	1.8	1.8	2.0	1.3
Mexican, both sexes.....	0.4	*	*	1.0	0.9	1.0	*
Male.....
Female.....	0.8	*	*	2.2	2.1	2.3	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.3	*	*	*	*	*	*
Male.....
Female.....	0.6	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.2	*	0.1	0.7	0.8	0.7	0.7
Male.....
Female.....	0.5	*	0.3	1.4	1.7	1.4	1.3
Non-Hispanic white 6/, both sexes....	0.1	*	*	0.5	0.5	0.5	0.4
Male.....
Female.....	0.3	*	*	1.0	1.0	0.9	0.8
Non-Hispanic black 6/, both sexes....	0.7	*	*	1.6	2.3	1.9	1.9
Male.....
Female.....	1.3	*	*	3.2	4.6	3.7	3.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

All origins 3/, both sexes.....	0.0	*	*	*	*	*	*
Male.....
Female.....	0.0	*	*	*	*	*	*
Hispanic, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.0	*	*	*	*	*	*
Male.....
Female.....	0.0	*	*	*	*	*	*
Non-Hispanic white 6/, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Non-Hispanic black 6/, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

All origins 3/, both sexes.....	0.2	*	0.2	0.7	0.8	0.7	0.6
Male.....
Female.....	0.5	*	0.3	1.4	1.6	1.5	1.3
Hispanic, both sexes.....	0.3	*	*	0.8	0.8	0.9	0.6
Male.....
Female.....	0.7	*	*	1.7	1.8	1.9	1.3
Mexican, both sexes.....	0.4	*	*	0.9	0.9	1.0	*
Male.....
Female.....	0.8	*	*	2.0	2.1	2.2	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.3	*	*	*	*	*	*
Male.....
Female.....	0.6	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.2	*	0.1	0.6	0.8	0.7	0.7
Male.....
Female.....	0.4	*	0.3	1.3	1.6	1.4	1.3
Non-Hispanic white 6/, both sexes....	0.1	*	*	0.5	0.5	0.4	0.4
Male.....
Female.....	0.3	*	*	1.0	0.9	0.9	0.8
Non-Hispanic black 6/, both sexes....	0.6	*	*	1.5	2.2	1.8	1.9
Male.....
Female.....	1.2	*	*	3.1	4.4	3.4	3.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Congenital malformations, deformations and chromosomal abnormalities (000-Q99)

All origins 3/, both sexes.....	3.5	140.9	3.2	0.9	0.8	1.1	1.1	1.1	1.0	1.1
Male.....	3.8	146.6	2.9	0.9	0.9	1.3	1.3	1.4	1.2	1.3
Female.....	3.2	134.9	3.4	1.0	0.7	0.8	0.9	0.9	0.9	0.9
Hispanic, both sexes.....	4.4	151.0	3.3	0.7	0.8	1.1	0.8	0.8	0.6	0.7
Male.....	4.6	153.4	3.6	*	*	1.5	1.0	0.9	*	*
Female.....	4.2	148.5	3.1	*	*	*	*	*	*	*
Mexican, both sexes.....	5.0	158.2	3.5	0.8	0.8	1.3	*	0.8	*	*
Male.....	5.1	161.5	4.1	*	*	1.6	*	*	*	*
Female.....	4.9	154.8	3.0	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	3.4	114.4	*	*	*	*	*	*	*	*
Male.....	3.7	*	*	*	*	*	*	*	*	*
Female.....	3.1	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	1.8	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.4	147.4	3.5	*	*	*	*	*	*	*
Male.....	3.8	163.8	*	*	*	*	*	*	*	*
Female.....	3.1	131.0	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.3	136.8	3.1	1.0	0.8	1.0	1.2	1.2	1.1	1.2
Male.....	3.6	143.6	2.7	0.9	0.8	1.2	1.3	1.5	1.2	1.4
Female.....	3.0	129.7	3.5	1.0	0.7	0.9	1.0	0.9	1.0	1.0
Non-Hispanic white 6/, both sexes....	3.2	129.5	2.8	1.0	0.7	1.0	1.1	1.2	1.2	1.2
Male.....	3.4	135.2	2.6	1.0	0.8	1.1	1.2	1.5	1.3	1.4
Female.....	2.9	123.5	2.9	0.9	0.5	0.8	1.0	0.8	1.1	0.9
Non-Hispanic black 6/, both sexes....	4.6	174.3	4.2	1.2	1.0	1.4	1.3	1.6	1.1	1.7
Male.....	5.1	182.2	3.2	*	*	1.7	1.6	1.9	*	1.8
Female.....	4.2	166.0	5.3	1.4	*	*	*	*	*	1.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins 3/, both sexes.....	10.6	83.8	1.5	0.3	0.3	0.9	2.2	2.4	2.6	3.2
Male.....	9.5	95.2	1.6	0.4	0.4	1.1	2.9	3.2	3.7	4.0
Female.....	11.7	71.9	1.3	0.3	0.2	0.7	1.5	1.5	1.5	2.4
Hispanic, both sexes.....	4.0	52.5	1.2	*	*	0.8	1.8	1.8	2.1	2.5
Male.....	4.5	62.1	1.1	*	*	1.3	2.5	2.7	3.1	3.8
Female.....	3.4	42.5	1.3	*	*	*	*	*	*	*
Mexican, both sexes.....	3.6	47.4	1.1	*	*	0.9	1.9	1.7	1.9	2.1
Male.....	4.2	57.1	*	*	*	*	2.7	2.4	2.8	3.2
Female.....	2.9	37.7	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	7.2	101.9	*	*	*	*	*	*	*	*
Male.....	8.5	*	*	*	*	*	*	*	*	*
Female.....	6.0	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	5.4	*	*	*	*	*	*	*	*	*
Male.....	5.9	*	*	*	*	*	*	*	*	*
Female.....	4.8	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.7	55.9	*	*	*	*	*	*	*	*
Male.....	4.0	66.2	*	*	*	*	*	*	*	*
Female.....	3.3	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	11.7	92.7	1.5	0.4	0.3	0.9	2.3	2.5	2.8	3.3
Male.....	10.3	104.3	1.7	0.4	0.4	1.1	2.9	3.3	3.9	4.1
Female.....	13.0	80.5	1.3	0.3	0.3	0.7	1.6	1.7	1.7	2.6
Non-Hispanic white 6/, both sexes....	12.4	77.1	1.2	0.3	0.3	0.9	2.3	2.6	2.7	3.3
Male.....	10.5	89.1	1.4	0.4	0.4	1.0	3.0	3.4	3.7	4.1
Female.....	14.2	64.5	1.1	*	*	0.7	1.5	1.7	1.6	2.6
Non-Hispanic black 6/, both sexes....	11.4	161.4	2.8	*	*	1.2	2.6	3.0	4.0	4.4
Male.....	12.0	172.0	3.1	*	*	1.5	3.2	3.7	5.8	5.3
Female.....	10.8	150.2	2.4	*	*	*	1.9	2.2	2.4	3.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

All other diseases (Residual) (D65-E07, E15-E34, E65-F99, G04-G12, G23-G25, G31-H93, K00-K22, K29-K31, K50-K66, K71-K72, K75-K76, K83-M99, N13.0-N13.5, N13.7-N13.9, N14, N15.0, N15.8-N15.9, N20-N23, N28-N39, N41-N64, N80-N98)

All origins 3/, both sexes.....	79.3	28.4	3.4	1.8	2.2	3.9	6.1	7.4	9.2	13.4
Male.....	65.2	32.5	3.6	1.9	2.3	4.6	7.0	8.6	10.5	15.4
Female.....	93.0	24.1	3.1	1.7	2.0	3.1	5.2	6.3	8.0	11.5
Hispanic, both sexes.....	26.4	24.6	3.3	1.3	2.3	3.7	5.2	5.8	6.5	10.0
Male.....	24.9	28.0	3.3	1.5	2.3	4.8	5.8	6.8	8.0	11.7
Female.....	28.1	21.1	3.4	1.2	2.3	2.5	4.5	4.5	4.7	8.0
Mexican, both sexes.....	22.1	23.5	3.4	1.5	2.5	3.9	5.0	5.6	6.8	9.3
Male.....	21.5	28.1	3.3	1.7	2.4	4.9	5.6	6.5	8.9	11.4
Female.....	22.6	18.8	3.5	*	2.6	2.8	4.3	4.4	4.3	6.8
Puerto Rican, both sexes.....	40.0	*	*	*	*	*	*	*	*	18.3
Male.....	40.7	*	*	*	*	*	*	*	*	*
Female.....	39.2	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	74.3	*	*	*	*	*	*	*	*	*
Male.....	59.6	*	*	*	*	*	*	*	*	*
Female.....	89.3	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	26.1	33.1	4.1	*	*	2.7	6.1	6.8	5.7	9.4
Male.....	23.7	*	*	*	*	*	6.2	8.3	6.2	12.0
Female.....	28.7	*	*	*	*	*	6.0	*	*	6.8
Non-Hispanic 5/, both sexes.....	88.3	29.5	3.4	1.9	2.1	3.9	6.3	7.8	9.9	14.1
Male.....	72.4	33.8	3.7	2.0	2.3	4.6	7.2	9.0	11.1	16.1
Female.....	103.5	25.0	3.0	1.8	1.9	3.3	5.3	6.7	8.7	12.1
Non-Hispanic white 6/, both sexes....	96.9	23.3	3.0	1.8	2.0	3.6	6.1	7.4	9.0	12.7
Male.....	78.2	26.8	3.3	1.8	2.2	4.3	7.5	9.0	10.8	15.0
Female.....	114.9	19.5	2.7	1.7	1.7	2.8	4.6	5.8	7.2	10.4
Non-Hispanic black 6/, both sexes....	68.7	53.8	4.6	2.3	2.9	5.5	8.1	11.7	16.7	23.2
Male.....	61.5	61.0	5.0	2.8	3.2	6.2	7.5	11.6	15.5	24.6
Female.....	75.2	46.3	4.1	1.8	2.6	4.8	8.7	11.9	17.8	22.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/, both sexes.....	40.6	27.8	9.9	5.3	5.9	31.2	45.3	38.8	35.1	36.9
Male.....	53.5	31.6	11.1	6.1	7.3	43.0	69.1	59.2	52.1	52.7
Female.....	28.1	23.8	8.6	4.4	4.4	18.9	19.9	17.6	17.6	20.8
Hispanic, both sexes.....	27.2	19.5	9.7	4.3	6.0	29.2	42.8	30.9	28.4	28.9
Male.....	39.7	20.8	11.3	5.4	7.3	43.8	66.3	46.5	43.1	43.8
Female.....	13.8	18.1	8.0	3.3	4.6	13.7	14.8	11.2	10.8	11.9
Mexican, both sexes.....	26.9	18.3	10.7	4.6	6.7	31.6	44.4	31.3	27.2	28.3
Male.....	39.4	19.1	12.5	5.4	8.2	47.2	68.5	46.8	42.1	43.0
Female.....	13.1	17.4	8.8	3.7	5.2	14.6	14.8	11.3	9.1	10.5
Puerto Rican, both sexes.....	30.6	*	*	*	*	20.0	38.7	32.0	41.6	41.7
Male.....	47.6	*	*	*	*	27.7	58.4	58.1	66.3	64.8
Female.....	14.5	*	*	*	*	*	19.2	*	*	*
Cuban, both sexes.....	30.3	*	*	*	*	*	48.9	*	*	*
Male.....	42.8	*	*	*	*	*	*	*	*	*
Female.....	17.5	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	26.3	24.1	7.5	3.2	4.2	24.8	38.3	29.9	29.0	28.2
Male.....	37.2	*	7.9	*	*	37.6	59.5	41.6	41.4	42.4
Female.....	15.0	*	*	*	*	11.4	13.4	13.8	14.2	13.7
Non-Hispanic 5/, both sexes.....	42.8	30.2	9.8	5.5	5.9	31.6	45.8	40.7	36.6	38.3
Male.....	55.8	34.7	10.9	6.3	7.2	42.6	69.5	62.5	54.4	54.5
Female.....	30.4	25.5	8.7	4.7	4.4	19.9	20.9	18.9	19.0	22.4
Non-Hispanic white 6/, both sexes....	45.7	24.3	8.9	4.8	5.8	34.5	50.5	44.3	39.3	40.5
Male.....	58.6	27.6	9.8	5.4	7.0	45.8	76.9	67.4	57.3	56.5
Female.....	33.3	20.8	8.0	4.2	4.4	22.6	22.7	20.8	21.0	24.3
Non-Hispanic black 6/, both sexes....	36.2	54.8	14.2	8.3	6.5	22.1	31.9	34.9	36.4	39.2
Male.....	52.5	64.1	16.3	10.0	8.3	33.5	47.4	56.0	59.6	60.8
Female.....	21.5	45.0	12.0	6.6	4.7	10.4	16.0	15.0	15.5	20.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Transport accidents (V01-V99,Y85)

All origins 3/, both sexes.....	16.2	3.4	3.8	3.1	4.1	23.9	29.8	20.9	17.4	16.4
Male.....	23.1	3.3	4.0	3.5	4.7	31.2	44.6	31.8	25.9	23.7
Female.....	9.4	3.6	3.6	2.8	3.3	16.2	14.0	9.5	8.6	9.1
Hispanic, both sexes.....	14.8	4.6	4.7	2.8	4.5	23.4	32.1	20.0	15.5	14.9
Male.....	21.7	4.6	5.3	3.4	5.2	34.2	49.1	29.7	23.0	21.6
Female.....	7.4	4.6	4.0	2.2	3.7	12.0	11.8	7.7	6.5	7.2
Mexican, both sexes.....	15.6	5.1	5.5	3.0	5.1	25.9	33.4	20.6	15.4	16.0
Male.....	22.8	*	6.1	3.5	5.9	37.8	50.6	30.5	23.3	23.3
Female.....	7.6	*	4.8	2.5	4.2	12.9	12.1	8.1	5.9	7.3
Puerto Rican, both sexes.....	11.4	*	*	*	*	15.6	25.8	16.1	14.0	11.9
Male.....	17.5	*	*	*	*	20.5	39.2	29.0	20.9	*
Female.....	5.6	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	16.0	*	*	*	*	*	41.6	*	*	*
Male.....	23.8	*	*	*	*	*	*	*	*	*
Female.....	8.1	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	13.6	*	*	2.8	3.1	19.1	29.2	19.3	17.0	13.4
Male.....	19.7	*	*	*	*	27.4	44.8	27.2	24.2	19.4
Female.....	7.3	*	*	*	*	10.4	10.8	8.5	8.4	7.3
Non-Hispanic 5/, both sexes.....	16.4	3.1	3.5	3.2	3.9	23.9	29.3	21.1	17.8	16.7
Male.....	23.3	2.9	3.5	3.5	4.6	30.5	43.5	32.3	26.6	24.0
Female.....	9.8	3.2	3.5	2.9	3.3	17.0	14.4	9.9	9.1	9.4
Non-Hispanic white 6/, both sexes....	16.9	2.6	3.0	3.0	4.0	26.3	31.3	21.4	18.1	17.1
Male.....	23.8	2.6	2.9	3.1	4.6	33.0	46.5	32.6	26.4	24.2
Female.....	10.2	2.6	3.2	2.8	3.4	19.2	15.3	10.0	9.7	10.0
Non-Hispanic black 6/, both sexes....	15.6	4.9	5.2	4.1	3.9	16.0	23.5	22.8	21.9	18.4
Male.....	23.6	*	5.8	4.9	4.8	22.8	34.6	36.3	36.5	30.0
Female.....	8.2	*	4.6	3.3	3.0	9.0	12.1	10.1	8.6	8.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8, V88.0-V88.8,V89.0,V89.2)

All origins 3/, both sexes.....	15.1	3.4	3.6	2.9	3.7	23.2	28.8	19.9	16.4	15.2
Male.....	21.4	3.3	3.8	3.2	4.2	30.2	42.9	30.1	24.4	21.7
Female.....	9.0	3.5	3.4	2.6	3.2	15.7	13.7	9.2	8.1	8.7
Hispanic, both sexes.....	14.2	4.6	4.6	2.7	4.2	22.9	31.4	19.3	14.9	14.0
Male.....	20.7	4.6	5.3	3.3	4.8	33.5	48.0	28.7	22.1	20.2
Female.....	7.2	4.6	3.9	2.1	3.6	11.7	11.5	7.3	6.4	7.0
Mexican, both sexes.....	15.0	5.1	5.4	2.9	4.8	25.3	32.8	19.8	14.8	15.1
Male.....	21.8	*	6.1	3.4	5.4	37.0	49.6	29.4	22.3	21.7
Female.....	7.4	*	4.6	2.4	4.2	12.6	11.9	7.6	5.8	7.1
Puerto Rican, both sexes.....	11.0	*	*	*	*	15.6	25.2	16.1	12.9	11.9
Male.....	17.1	*	*	*	*	20.5	37.8	29.0	*	*
Female.....	5.3	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	15.0	*	*	*	*	*	40.6	*	*	*
Male.....	22.1	*	*	*	*	*	*	*	*	*
Female.....	7.9	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	13.0	*	*	*	2.8	18.4	28.3	18.8	16.6	12.7
Male.....	18.6	*	*	*	*	26.4	43.5	26.2	23.4	18.2
Female.....	7.1	*	*	*	*	10.1	10.5	8.5	8.4	7.1
Non-Hispanic 5/, both sexes.....	15.3	3.0	3.3	3.0	3.6	23.1	28.1	20.0	16.7	15.4
Male.....	21.5	2.9	3.3	3.2	4.0	29.4	41.6	30.4	25.0	21.9
Female.....	9.3	3.1	3.3	2.8	3.1	16.5	14.0	9.6	8.5	9.0
Non-Hispanic white 6/, both sexes....	15.7	2.5	2.8	2.7	3.6	25.4	30.0	20.3	16.9	15.8
Male.....	22.0	2.6	2.7	2.8	3.9	31.8	44.4	30.7	24.7	22.0
Female.....	9.7	2.5	3.0	2.6	3.2	18.7	14.8	9.8	8.9	9.5
Non-Hispanic black 6/, both sexes....	14.7	4.7	5.0	3.9	3.8	15.5	23.0	21.7	20.9	17.1
Male.....	22.2	*	5.7	4.7	4.5	22.1	33.8	34.2	34.9	27.7
Female.....	7.9	*	4.3	3.2	3.0	8.8	11.9	9.8	8.3	7.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Nontransport accidents (W00-X59,Y86)

All origins 3/, both sexes.....	24.4	24.3	6.1	2.2	1.8	7.3	15.5	17.9	17.7	20.4
Male.....	30.4	28.3	7.2	2.7	2.5	11.7	24.5	27.3	26.3	29.1
Female.....	18.6	20.2	5.0	1.7	1.1	2.7	5.9	8.1	8.9	11.7
Hispanic, both sexes.....	12.4	14.8	5.0	1.5	1.5	5.8	10.7	10.9	12.9	14.0
Male.....	18.0	16.1	6.0	2.0	2.1	9.6	17.2	16.8	20.1	22.1
Female.....	6.4	13.5	4.0	1.1	*	1.7	3.0	3.5	4.3	4.7
Mexican, both sexes.....	11.3	13.2	5.2	1.6	1.7	5.8	11.1	10.6	11.8	12.3
Male.....	16.6	14.6	6.3	2.0	2.3	9.5	17.9	16.4	18.9	19.8
Female.....	5.5	11.8	4.0	*	*	1.8	2.7	3.2	3.2	3.2
Puerto Rican, both sexes.....	19.2	*	*	*	*	*	12.9	15.8	27.6	29.8
Male.....	30.1	*	*	*	*	*	19.2	29.0	45.3	47.7
Female.....	8.9	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	14.2	*	*	*	*	*	*	*	*	*
Male.....	19.0	*	*	*	*	*	*	*	*	*
Female.....	9.4	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	12.7	19.8	5.0	*	*	5.7	9.2	10.6	12.0	14.8
Male.....	17.5	*	*	*	*	10.2	14.7	14.5	17.2	23.0
Female.....	7.7	*	*	*	*	*	*	*	5.8	6.4
Non-Hispanic 5/, both sexes.....	26.4	27.1	6.3	2.3	1.9	7.7	16.5	19.6	18.8	21.7
Male.....	32.6	31.8	7.4	2.8	2.6	12.2	26.1	30.2	27.8	30.4
Female.....	20.6	22.2	5.2	1.8	1.1	2.9	6.5	9.0	9.9	13.0
Non-Hispanic white 6/, both sexes....	28.8	21.7	5.9	1.8	1.8	8.3	19.2	22.9	21.2	23.4
Male.....	34.8	25.0	6.9	2.3	2.4	12.8	30.4	34.8	30.9	32.4
Female.....	23.1	18.2	4.8	1.3	1.0	3.5	7.5	10.8	11.4	14.3
Non-Hispanic black 6/, both sexes....	20.7	49.9	9.0	4.2	2.6	6.1	8.4	12.1	14.5	20.9
Male.....	28.8	59.5	10.5	5.1	3.5	10.8	12.8	19.7	23.0	30.8
Female.....	13.3	39.9	7.5	3.4	1.6	1.4	3.9	4.9	6.8	12.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental exposure to smoke, fire and flames (X00-X09)

All origins 3/, both sexes.....	1.0	0.7	1.2	0.6	0.3	0.4	0.5	0.5	0.5	0.6
Male.....	1.2	*	1.4	0.6	0.3	0.4	0.6	0.7	0.6	0.8
Female.....	0.8	*	1.0	0.6	0.3	0.3	0.3	0.3	0.4	0.5
Hispanic, both sexes.....	0.5	*	0.8	*	*	*	*	*	*	*
Male.....	0.7	*	*	*	*	*	*	*	*	*
Female.....	0.3	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	0.5	*	0.8	*	*	*	*	*	*	*
Male.....	0.7	*	*	*	*	*	*	*	*	*
Female.....	0.3	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	0.7	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	0.5	*	*	*	*	*	*	*	*	*
Male.....	0.7	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.1	0.8	1.3	0.6	0.3	0.4	0.5	0.5	0.5	0.7
Male.....	1.3	*	1.5	0.6	0.3	0.4	0.6	0.7	0.6	0.8
Female.....	0.9	*	1.2	0.7	0.4	0.3	0.4	0.4	0.4	0.5
Non-Hispanic white 6/, both sexes....	1.1	*	0.9	0.5	0.2	0.4	0.5	0.5	0.5	0.6
Male.....	1.3	*	1.2	0.4	*	0.5	0.6	0.6	0.6	0.7
Female.....	0.9	*	0.7	0.5	*	0.3	*	0.3	0.4	0.5
Non-Hispanic black 6/, both sexes....	1.8	*	3.1	1.6	*	*	0.7	0.9	0.9	1.1
Male.....	2.1	*	3.1	1.5	*	*	*	*	*	*
Female.....	1.6	*	3.1	1.6	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All origins 3/, both sexes.....	9.2	*	0.2	*	0.2	3.5	10.4	13.0	13.0	15.1
Male.....	12.6	*	*	*	0.2	5.2	16.0	19.2	18.6	20.5
Female.....	5.9	*	*	*	*	1.6	4.4	6.5	7.3	9.6
Hispanic, both sexes.....	5.2	*	*	*	*	2.0	4.6	5.8	8.1	9.0
Male.....	7.8	*	*	*	*	2.8	6.7	8.5	12.2	13.5
Female.....	2.4	*	*	*	*	*	2.0	2.4	3.3	3.9
Mexican, both sexes.....	4.3	*	*	*	*	1.9	4.3	5.1	6.7	6.9
Male.....	6.5	*	*	*	*	2.8	6.5	7.3	10.5	10.6
Female.....	1.9	*	*	*	*	*	*	2.3	2.2	2.5
Puerto Rican, both sexes.....	12.3	*	*	*	*	*	7.6	13.4	23.8	26.8
Male.....	20.3	*	*	*	*	*	*	24.7	39.1	43.2
Female.....	4.8	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	4.0	*	*	*	*	*	*	*	*	*
Male.....	5.6	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.4	*	*	*	*	*	4.1	5.7	7.5	9.5
Male.....	7.7	*	*	*	*	*	6.0	7.5	9.8	13.8
Female.....	3.0	*	*	*	*	*	*	*	*	5.2
Non-Hispanic 5/, both sexes.....	9.8	*	0.2	*	0.2	3.8	11.7	14.8	14.2	16.2
Male.....	13.4	*	*	*	0.3	5.7	18.1	22.2	20.3	21.9
Female.....	6.4	*	*	*	*	1.7	4.9	7.4	8.1	10.7
Non-Hispanic white 6/, both sexes....	10.5	*	*	*	0.2	4.7	14.3	17.9	16.5	17.8
Male.....	14.2	*	*	*	*	7.2	22.3	26.6	23.2	23.7
Female.....	7.0	*	*	*	*	2.1	5.9	9.2	9.6	11.9
Non-Hispanic black 6/, both sexes....	9.2	*	*	*	*	0.8	3.7	7.0	9.6	15.0
Male.....	13.3	*	*	*	*	1.2	5.1	11.4	14.9	21.2
Female.....	5.4	*	*	*	*	*	2.2	2.8	4.9	9.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All origins 3/, both sexes.....	5.8	21.5	1.6	0.6	0.6	1.2	1.9	2.1	2.2	2.5
Male.....	7.0	25.2	1.7	0.8	0.8	2.0	3.2	3.5	3.5	4.2
Female.....	4.7	17.6	1.5	0.4	0.3	0.4	0.6	0.7	0.8	0.8
Hispanic, both sexes.....	2.7	12.4	1.4	*	*	0.8	2.1	1.9	2.1	2.4
Male.....	3.9	13.7	1.4	*	*	1.3	3.6	3.1	3.4	4.3
Female.....	1.5	11.0	1.5	*	*	*	*	*	*	*
Mexican, both sexes.....	2.8	10.7	1.7	*	*	*	2.3	2.0	2.4	2.7
Male.....	4.0	12.1	1.8	*	*	*	3.9	3.5	3.9	4.9
Female.....	1.4	9.3	1.6	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	2.3	*	*	*	*	*	*	*	*	*
Male.....	3.2	*	*	*	*	*	*	*	*	*
Female.....	1.4	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	3.3	*	*	*	*	*	*	*	*	*
Male.....	4.7	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.5	17.4	*	*	*	*	*	*	*	*
Male.....	3.5	*	*	*	*	*	*	*	*	*
Female.....	1.5	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	6.4	24.2	1.6	0.6	0.6	1.3	1.9	2.2	2.1	2.5
Male.....	7.5	28.6	1.7	0.8	0.9	2.1	3.1	3.6	3.5	4.2
Female.....	5.2	19.6	1.5	0.4	0.3	0.4	0.6	0.7	0.8	0.9
Non-Hispanic white 6/, both sexes....	6.9	18.9	1.5	0.6	0.6	1.3	2.1	2.4	2.3	2.7
Male.....	8.0	22.0	1.7	0.8	1.0	2.1	3.6	4.0	3.7	4.3
Female.....	5.8	15.6	1.4	0.4	*	0.4	0.6	0.7	0.8	1.0
Non-Hispanic black 6/, both sexes....	5.1	46.4	2.2	0.7	0.7	1.1	1.1	1.9	1.9	2.5
Male.....	6.6	55.5	2.1	*	*	1.9	1.4	2.5	3.2	4.2
Female.....	3.7	37.0	2.3	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All origins 3/, both sexes.....	11.1	*	1.0	7.3	12.5	12.4	12.3	14.7
Male.....	17.8	*	1.4	11.5	20.8	19.7	19.7	22.7
Female.....	4.6	*	0.6	2.8	3.6	4.7	4.7	6.6
Hispanic, both sexes.....	4.9	*	0.6	5.5	9.2	7.6	6.6	6.4
Male.....	7.9	*	*	7.9	15.0	11.5	10.3	10.2
Female.....	1.7	*	*	2.9	2.3	2.7	2.2	2.1
Mexican, both sexes.....	4.4	*	*	5.8	9.1	7.7	6.5	6.1
Male.....	7.1	*	*	8.3	14.4	11.6	10.3	9.1
Female.....	1.5	*	*	3.0	2.6	2.7	1.9	2.5
Puerto Rican, both sexes.....	5.4	*	*	*	8.3	7.0	*	11.5
Male.....	8.8	*	*	*	*	*	*	*
Female.....	2.1	*	*	*	*	*	*	*
Cuban, both sexes.....	9.8	*	*	*	*	*	*	*
Male.....	17.0	*	*	*	*	*	*	*
Female.....	2.6	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.3	*	*	5.4	9.3	7.6	6.7	5.6
Male.....	8.5	*	*	7.8	15.2	10.8	9.8	10.1
Female.....	1.9	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	12.2	*	1.1	7.6	13.1	13.6	13.7	16.3
Male.....	19.6	*	1.6	12.3	22.0	22.0	22.2	25.4
Female.....	5.1	*	0.7	2.8	3.8	5.1	5.3	7.4
Non-Hispanic white 6/, both sexes....	13.9	*	1.1	8.3	14.1	15.1	15.5	18.8
Male.....	22.3	*	1.6	13.3	23.7	24.2	24.7	28.7
Female.....	5.9	*	0.7	3.0	4.0	6.0	6.2	8.9
Non-Hispanic black 6/, both sexes....	5.1	*	1.1	4.3	8.7	8.9	8.5	8.5
Male.....	9.1	*	1.7	7.2	15.0	16.3	16.0	15.4
Female.....	1.4	*	*	1.3	2.3	2.0	1.7	2.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All origins 3/, both sexes.....	5.6	*	0.3	3.3	6.0	5.4	5.2	6.2
Male.....	10.0	*	0.5	5.8	10.6	9.2	9.2	10.4
Female.....	1.4	*	*	0.6	1.2	1.4	1.2	1.9
Hispanic, both sexes.....	1.8	*	*	2.2	3.9	2.8	2.3	2.2
Male.....	3.3	*	*	4.1	6.8	4.4	4.1	3.8
Female.....	0.3	*	*	*	*	*	*	*
Mexican, both sexes.....	1.7	*	*	2.3	3.6	2.9	2.5	2.2
Male.....	3.1	*	*	4.2	6.2	4.5	4.4	3.8
Female.....	0.3	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	1.6	*	*	*	*	*	*	*
Male.....	3.0	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*
Cuban, both sexes.....	3.8	*	*	*	*	*	*	*
Male.....	7.3	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	1.9	*	*	*	4.5	2.5	2.3	*
Male.....	3.4	*	*	*	7.6	*	*	*
Female.....	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	6.3	*	0.3	3.5	6.5	6.1	5.9	7.0
Male.....	11.2	*	0.5	6.2	11.5	10.6	10.5	11.8
Female.....	1.6	*	*	0.7	1.3	1.5	1.4	2.2
Non-Hispanic white 6/, both sexes....	7.3	*	0.4	3.8	6.9	6.6	6.7	8.2
Male.....	13.0	*	0.7	6.7	12.2	11.4	11.8	13.5
Female.....	1.9	*	*	0.7	1.4	1.8	1.7	2.8
Non-Hispanic black 6/, both sexes....	2.6	*	*	2.4	5.4	5.3	4.3	3.5
Male.....	5.0	*	*	4.3	9.8	10.0	8.2	6.9
Female.....	0.5	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All origins 3/, both sexes.....	5.5	*	0.7	4.0	6.4	6.9	7.1	8.5
Male.....	7.8	*	0.9	5.7	10.2	10.5	10.5	12.3
Female.....	3.2	*	0.5	2.2	2.4	3.3	3.5	4.7
Hispanic, both sexes.....	3.1	*	*	3.2	5.3	4.8	4.3	4.2
Male.....	4.6	*	*	3.7	8.1	7.1	6.2	6.4
Female.....	1.4	*	*	2.7	1.8	1.8	1.9	1.8
Mexican, both sexes.....	2.7	*	*	3.4	5.5	4.8	4.0	3.9
Male.....	4.0	*	*	4.1	8.2	7.1	5.9	5.3
Female.....	1.2	*	*	2.7	2.1	1.8	1.7	2.2
Puerto Rican, both sexes.....	3.8	*	*	*	*	*	*	*
Male.....	5.8	*	*	*	*	*	*	*
Female.....	1.9	*	*	*	*	*	*	*
Cuban, both sexes.....	6.1	*	*	*	*	*	*	*
Male.....	9.7	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.4	*	*	3.1	4.8	5.1	4.4	4.0
Male.....	5.2	*	*	*	7.6	6.9	6.0	7.1
Female.....	1.5	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	5.9	*	0.8	4.2	6.6	7.5	7.8	9.4
Male.....	8.4	*	1.0	6.1	10.6	11.4	11.7	13.6
Female.....	3.5	*	0.6	2.1	2.5	3.6	3.9	5.2
Non-Hispanic white 6/, both sexes....	6.6	*	0.8	4.5	7.1	8.5	8.8	10.7
Male.....	9.3	*	0.9	6.6	11.4	12.8	12.9	15.2
Female.....	4.0	*	0.6	2.3	2.6	4.1	4.6	6.1
Non-Hispanic black 6/, both sexes....	2.4	*	1.0	1.9	3.3	3.6	4.1	4.9
Male.....	4.0	*	1.6	2.9	5.2	6.2	7.7	8.5
Female.....	1.0	*	*	*	1.4	*	*	1.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All origins 3/, both sexes.....	6.2	8.1	2.2	0.8	1.2	10.7	16.2	13.5	9.8	7.6
Male.....	10.0	9.4	2.5	0.8	1.6	18.2	27.5	22.4	16.2	11.4
Female.....	2.5	6.8	2.0	0.7	0.7	2.9	4.2	4.2	3.3	3.6
Hispanic, both sexes.....	8.0	8.3	1.7	0.7	1.7	15.8	20.5	14.5	9.9	9.2
Male.....	13.1	10.1	1.8	*	2.5	28.0	33.8	23.5	15.7	14.3
Female.....	2.4	6.5	1.5	*	*	3.0	4.7	3.3	3.1	3.4
Mexican, both sexes.....	8.5	8.2	1.7	0.7	1.9	17.7	21.8	15.0	10.4	9.7
Male.....	14.1	10.1	1.9	*	2.9	30.9	36.0	24.1	16.6	15.1
Female.....	2.4	6.2	*	*	*	3.3	4.3	3.3	3.0	3.2
Puerto Rican, both sexes.....	8.4	*	*	*	*	14.7	23.5	14.6	10.8	12.7
Male.....	13.8	*	*	*	*	26.5	40.5	27.2	*	*
Female.....	3.2	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	4.5	*	*	*	*	*	*	*	*	*
Male.....	6.9	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.7	*	*	*	*	11.5	17.0	14.3	8.8	7.2
Male.....	10.8	*	*	*	*	21.1	26.8	22.0	13.8	11.1
Female.....	2.4	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	5.9	8.0	2.4	0.8	1.0	9.7	15.2	13.2	9.7	7.2
Male.....	9.3	9.1	2.7	0.8	1.4	16.2	25.9	22.0	16.2	10.8
Female.....	2.5	6.8	2.1	0.7	0.7	2.9	4.0	4.4	3.3	3.7
Non-Hispanic white 6/, both sexes....	2.7	6.3	1.5	0.6	0.6	2.6	4.3	4.1	3.6	3.5
Male.....	3.6	6.5	1.6	0.5	0.6	3.4	6.1	5.8	5.1	4.6
Female.....	1.8	6.2	1.3	0.7	0.5	1.7	2.4	2.5	2.0	2.4
Non-Hispanic black 6/, both sexes....	23.6	15.9	6.0	1.7	3.0	38.8	64.7	56.4	41.4	27.9
Male.....	42.0	20.3	6.4	2.2	4.3	68.8	116.4	101.9	76.9	47.9
Female.....	6.8	11.3	5.6	*	1.7	7.9	11.8	13.5	9.4	10.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All origins 3/, both sexes.....	4.3	*	0.3	0.3	0.8	9.1	13.7	10.8	7.7	5.2
Male.....	7.4	*	0.3	0.3	1.3	16.0	24.1	18.7	13.4	8.4
Female.....	1.3	*	*	0.3	0.4	1.8	2.6	2.5	1.9	2.0
Hispanic, both sexes.....	5.6	*	*	*	1.3	13.4	16.7	11.1	7.5	5.8
Male.....	9.7	*	*	*	2.1	24.0	28.7	18.3	12.4	9.3
Female.....	1.2	*	*	*	*	2.1	2.4	2.0	1.6	1.8
Mexican, both sexes.....	6.3	*	*	*	1.6	15.4	18.3	11.9	8.2	6.5
Male.....	10.9	*	*	*	2.6	27.2	31.2	19.7	13.6	10.0
Female.....	1.2	*	*	*	*	2.5	2.3	2.0	1.7	2.2
Puerto Rican, both sexes.....	5.0	*	*	*	*	10.8	18.5	11.3	8.4	*
Male.....	9.2	*	*	*	*	21.1	33.8	20.4	*	*
Female.....	1.1	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	2.6	*	*	*	*	*	*	*	*	*
Male.....	4.1	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.2	*	*	*	*	8.9	12.7	9.4	5.6	4.5
Male.....	7.3	*	*	*	*	16.7	21.2	14.6	8.8	8.1
Female.....	1.0	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.0	*	0.3	0.3	0.7	8.2	12.9	10.7	7.8	5.1
Male.....	6.9	*	*	0.3	1.1	14.3	22.8	18.7	13.7	8.1
Female.....	1.3	*	*	0.3	0.4	1.8	2.6	2.6	2.0	2.1
Non-Hispanic white 6/, both sexes....	1.4	*	*	0.2	0.3	1.7	2.7	2.6	2.4	2.2
Male.....	2.0	*	*	*	0.4	2.6	4.2	4.0	3.6	3.1
Female.....	0.8	*	*	*	*	0.9	1.2	1.3	1.1	1.4
Non-Hispanic black 6/, both sexes....	18.6	*	*	*	2.5	34.7	59.2	49.5	35.9	21.4
Male.....	35.0	*	*	*	3.8	63.2	108.4	92.1	68.9	39.2
Female.....	3.7	*	*	*	*	5.6	8.9	9.3	6.1	5.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All origins 3/, both sexes.....	1.9	8.0	2.0	0.4	0.3	1.6	2.6	2.7	2.0	2.3
Male.....	2.6	9.3	2.2	0.5	0.3	2.2	3.5	3.7	2.7	3.1
Female.....	1.3	6.6	1.8	0.4	0.3	1.1	1.6	1.7	1.3	1.6
Hispanic, both sexes.....	2.4	8.2	1.4	*	*	2.5	3.8	3.4	2.5	3.4
Male.....	3.4	10.1	1.5	*	*	4.0	5.1	5.1	3.3	4.9
Female.....	1.3	6.3	1.3	*	*	*	2.3	1.2	1.4	1.6
Mexican, both sexes.....	2.2	8.0	1.4	*	*	2.3	3.6	3.0	2.2	3.2
Male.....	3.3	10.1	1.6	*	*	3.8	4.8	4.4	2.9	5.1
Female.....	1.1	*	*	*	*	*	2.0	*	*	*
Puerto Rican, both sexes.....	3.3	*	*	*	*	*	*	*	*	*
Male.....	4.6	*	*	*	*	*	*	*	*	*
Female.....	2.1	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	1.9	*	*	*	*	*	*	*	*	*
Male.....	2.8	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.5	*	*	*	*	2.6	4.3	4.9	3.3	2.7
Male.....	3.5	*	*	*	*	*	5.6	7.3	5.0	*
Female.....	1.4	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.8	7.8	2.2	0.4	0.3	1.5	2.3	2.5	1.9	2.1
Male.....	2.4	9.0	2.4	0.5	0.3	1.8	3.1	3.2	2.6	2.6
Female.....	1.3	6.6	1.9	0.4	0.3	1.1	1.4	1.8	1.3	1.6
Non-Hispanic white 6/, both sexes....	1.3	6.3	1.3	0.3	0.2	0.8	1.6	1.5	1.2	1.3
Male.....	1.6	6.5	1.5	*	*	0.8	1.9	1.8	1.5	1.6
Female.....	1.0	6.1	1.1	0.4	*	0.8	1.2	1.2	0.9	1.0
Non-Hispanic black 6/, both sexes....	5.0	15.3	5.2	1.0	*	4.0	5.5	6.9	5.5	6.5
Male.....	7.1	19.7	5.6	1.5	*	5.7	8.1	9.8	8.0	8.7
Female.....	3.1	10.6	4.9	*	*	2.4	2.9	4.2	3.3	4.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

All origins 3/, both sexes.....	12.8	1.1	0.2	*	0.2	4.1	13.1	16.5	17.1	20.2
Male.....	16.6	1.1	0.2	*	0.2	6.0	19.5	23.3	23.5	25.8
Female.....	9.1	1.0	*	*	*	2.2	6.3	9.3	10.7	14.5
Hispanic, both sexes.....	6.5	*	*	*	*	2.1	5.2	6.6	9.6	11.0
Male.....	9.3	*	*	*	*	2.9	7.2	9.4	13.7	15.2
Female.....	3.4	*	*	*	*	1.2	2.8	3.1	4.8	6.1
Mexican, both sexes.....	5.4	*	*	*	*	2.2	4.9	5.7	8.3	8.5
Male.....	7.8	*	*	*	*	3.1	7.0	7.9	12.2	11.8
Female.....	2.7	*	*	*	*	*	2.2	2.8	3.6	4.5
Puerto Rican, both sexes.....	15.0	*	*	*	*	*	7.9	14.3	25.9	32.0
Male.....	23.9	*	*	*	*	*	*	24.7	41.2	48.4
Female.....	6.6	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	5.7	*	*	*	*	*	*	*	*	*
Male.....	7.4	*	*	*	*	*	*	*	*	*
Female.....	3.9	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.6	*	*	*	*	*	4.8	7.3	8.5	11.6
Male.....	9.0	*	*	*	*	*	6.0	9.6	10.4	15.9
Female.....	4.1	*	*	*	*	*	*	*	6.2	7.3
Non-Hispanic 5/, both sexes.....	13.9	1.1	0.3	*	0.2	4.5	14.8	18.9	18.9	22.0
Male.....	17.9	*	*	*	0.3	6.6	22.3	27.2	26.0	28.0
Female.....	10.1	*	*	*	*	2.4	7.0	10.7	11.9	16.1
Non-Hispanic white 6/, both sexes....	14.9	Non-	0.2	*	0.3	5.7	18.2	23.0	22.0	24.2
Male.....	18.9	*	*	*	*	8.3	27.6	32.7	29.6	30.3
Female.....	11.1	*	*	*	*	2.9	8.4	13.2	14.2	18.0
Non-Hispanic black 6/, both sexes....	12.5	*	*	*	*	0.9	4.5	8.5	12.6	19.5
Male.....	17.8	*	*	*	*	*	6.1	13.3	19.4	26.9
Female.....	7.6	*	*	*	*	*	2.9	4.0	6.5	12.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	All ages 1/	Under 1 year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

All origins 3/, both sexes.....	10.3	*	0.3	0.4	1.3	13.2	20.6	17.0	13.5	11.8
Male.....	18.1	*	0.5	0.4	2.0	23.3	36.2	29.4	23.6	19.5
Female.....	2.8	*	*	0.4	0.5	2.5	3.8	4.1	3.2	4.1
Hispanic, both sexes.....	7.8	*	*	*	1.5	16.5	21.5	14.6	10.5	8.4
Male.....	13.7	*	*	*	2.5	29.8	37.1	24.0	17.5	13.8
Female.....	1.5	*	*	*	*	2.3	3.0	2.8	2.1	2.2
Mexican, both sexes.....	8.5	*	*	*	1.9	18.8	22.9	15.7	11.5	9.2
Male.....	14.7	*	*	*	3.1	33.4	39.1	25.7	19.1	14.6
Female.....	1.6	*	*	*	*	2.9	2.8	2.9	2.2	2.6
Puerto Rican, both sexes.....	7.1	*	*	*	*	12.6	24.2	15.2	11.5	9.7
Male.....	13.1	*	*	*	*	24.7	44.5	27.2	21.6	*
Female.....	1.4	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	6.6	*	*	*	*	*	*	*	*	*
Male.....	11.9	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	6.3	*	*	*	*	12.0	17.8	12.2	8.2	6.4
Male.....	11.2	*	*	*	*	22.6	30.1	19.1	13.2	11.5
Female.....	1.3	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	10.7	*	0.4	0.4	1.2	12.5	20.3	17.5	14.2	12.5
Male.....	18.8	*	0.5	0.5	1.9	21.9	35.8	30.8	25.1	20.7
Female.....	2.9	*	*	0.4	0.5	2.5	4.0	4.3	3.5	4.4
Non-Hispanic white 6/, both sexes....	9.1	*	*	0.4	0.9	6.1	10.3	9.8	9.6	10.9
Male.....	15.6	*	*	0.4	1.3	10.3	17.5	16.4	16.2	17.3
Female.....	2.8	*	*	*	0.3	1.7	2.7	3.2	2.9	4.4
Non-Hispanic black 6/, both sexes....	22.0	*	1.1	0.7	2.8	38.9	66.5	56.7	41.3	25.6
Male.....	41.4	*	*	*	4.5	70.8	122.0	105.9	79.0	47.5
Female.....	4.2	*	*	*	*	6.1	9.9	10.2	7.1	6.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All causes

All origins 3/, both sexes.....	229.3	347.7	516.4	730.1	1,110.2	1,656.6	2,554.8	4,033.4	6,524.0	13,253.1
Male.....	285.9	435.3	659.7	920.0	1,373.6	2,040.2	3,117.5	4,944.6	7,942.7	14,309.1
Female.....	173.1	262.2	378.9	551.0	869.7	1,321.4	2,097.0	3,366.6	5,637.2	12,759.0
Hispanic, both sexes.....	164.0	253.6	385.2	549.3	813.9	1,229.6	1,922.0	3,052.6	4,794.7	9,022.8
Male.....	215.6	326.7	507.1	718.7	1,030.0	1,553.4	2,393.6	3,720.7	5,719.9	9,435.5
Female.....	107.0	177.0	263.9	389.3	622.4	957.8	1,554.8	2,569.3	4,178.9	8,803.5
Mexican, both sexes.....	163.9	247.3	368.8	556.8	803.2	1,332.5	1,839.5	3,052.6	4,862.0	9,619.7
Male.....	211.0	314.7	469.7	696.3	981.5	1,525.8	2,242.9	3,774.3	6,348.8	*
Female.....	108.6	174.1	257.3	411.9	637.2	1,141.6	1,510.8	2,543.5	3,968.8	9,423.5
Puerto Rican, both sexes.....	265.7	411.5	565.5	745.7	1,175.2	1,318.7	3,227.8	*	*	*
Male.....	375.4	606.6	827.6	969.3	*	*	*	*	*	*
Female.....	170.0	256.1	356.5	513.3	782.1	928.1	*	*	*	*
Cuban, both sexes.....	125.0	276.1	461.6	568.7	844.0	1,486.3	1,945.8	2,601.2	*	*
Male.....	166.8	330.6	*	*	*	*	*	*	*	*
Female.....	78.4	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	140.0	208.3	344.3	445.9	666.5	913.7	1,613.4	2,908.6	3,952.7	7,592.7
Male.....	190.5	266.3	465.8	590.2	848.9	1,298.0	1,891.2	*	*	*
Female.....	87.7	149.8	241.9	330.2	519.1	663.0	1,397.9	2,668.1	*	6,967.7
Non-Hispanic 5/, both sexes.....	238.9	358.3	528.6	744.2	1,132.1	1,686.9	2,597.2	4,092.4	6,612.2	13,450.1
Male.....	296.7	447.7	673.6	934.9	1,397.6	2,072.6	3,162.7	5,017.0	8,061.4	14,558.8
Female.....	182.4	271.9	389.9	564.3	889.0	1,348.6	2,135.6	3,415.2	5,709.7	12,934.8
Non-Hispanic white 6/, both sexes....	221.5	331.4	487.4	696.1	1,081.7	1,634.0	2,559.2	4,077.7	6,640.0	13,688.2
Male.....	278.0	417.1	621.0	869.7	1,327.5	2,000.1	3,100.3	4,983.5	8,103.3	14,841.1
Female.....	165.1	246.7	356.7	528.4	851.6	1,306.2	2,107.8	3,400.4	5,718.8	13,150.7
Non-Hispanic black 6/, both sexes....	384.5	590.6	914.1	1,260.3	1,789.0	2,454.3	3,432.1	4,987.2	7,380.4	12,655.4
Male.....	471.1	732.9	1,199.1	1,662.0	2,355.8	3,190.8	4,451.9	6,441.8	9,165.9	13,403.1
Female.....	308.4	467.0	673.7	929.8	1,348.2	1,911.3	2,732.5	4,118.7	6,438.9	12,350.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain other intestinal infections (A04,A07-A09)

All origins 3/, both sexes.....	0.1	0.2	0.4	0.9	1.5	3.3	6.7	13.5	26.2	50.5
Male.....	*	0.2	0.4	0.9	1.8	3.7	6.3	13.9	25.3	50.0
Female.....	*	*	0.4	0.8	1.3	3.0	6.9	13.3	26.8	50.8
Hispanic, both sexes.....	*	*	*	*	1.9	*	5.3	13.4	19.0	24.6
Male.....	*	*	*	*	*	*	*	*	18.0	27.1
Female.....	*	*	*	*	*	*	*	16.0	19.6	23.4
Mexican, both sexes.....	*	*	*	*	*	*	*	12.0	22.8	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	*	0.2	0.4	0.9	1.5	3.4	6.7	13.5	26.6	51.8
Male.....	*	0.3	0.4	0.9	1.7	3.8	6.3	14.1	25.7	51.2
Female.....	*	*	0.5	0.8	1.4	3.1	7.1	13.1	27.1	52.1
Non-Hispanic white 6/, both sexes.....	*	0.2	0.4	0.8	1.5	3.6	7.0	14.4	28.4	55.1
Male.....	*	*	0.3	0.9	1.6	4.0	6.5	14.8	27.1	54.4
Female.....	*	*	0.4	0.7	1.3	3.2	7.5	14.1	29.2	55.4
Non-Hispanic black 6/, both sexes.....	*	*	1.0	1.3	2.3	2.9	6.4	8.3	13.6	24.2
Male.....	*	*	*	*	*	*	*	9.2	14.0	20.2
Female.....	*	*	*	*	*	*	6.7	7.7	13.3	25.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Septicemia (A40-A41)

All origins 3/, both sexes.....	2.4	3.9	6.6	10.3	16.2	25.3	40.3	66.0	103.7	177.3
Male.....	2.6	4.2	7.7	11.7	17.4	27.6	43.9	75.3	117.7	194.8
Female.....	2.2	3.7	5.6	8.9	15.1	23.2	37.4	59.2	94.9	169.2
Hispanic, both sexes.....	1.4	3.0	5.3	7.3	13.6	19.1	32.4	54.3	81.5	122.9
Male.....	1.5	3.5	6.8	9.1	14.7	24.5	38.5	64.4	95.8	138.7
Female.....	1.3	2.4	3.8	5.7	12.5	14.4	27.6	47.0	71.9	114.4
Mexican, both sexes.....	1.7	3.1	4.5	8.6	14.2	20.6	28.3	55.0	83.3	141.7
Male.....	*	3.7	5.4	9.7	15.1	23.8	35.8	59.7	*	*
Female.....	*	*	*	7.4	13.4	17.5	22.1	51.6	65.2	137.7
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	4.5	*	9.8	13.3	29.1	42.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.6	4.1	6.8	10.5	16.4	25.7	40.8	66.7	104.7	179.7
Male.....	2.8	4.3	7.7	11.9	17.5	27.8	44.2	76.0	118.8	197.5
Female.....	2.3	3.8	5.8	9.1	15.3	23.9	38.1	59.9	95.9	171.4
Non-Hispanic white 6/, both sexes....	2.1	3.2	5.4	8.7	14.0	22.7	36.5	62.2	98.9	173.2
Male.....	2.3	3.5	6.2	9.7	14.7	24.8	39.9	71.1	113.1	193.3
Female.....	1.9	2.9	4.7	7.7	13.4	20.9	33.5	55.6	90.0	163.9
Non-Hispanic black 6/, both sexes....	5.9	10.3	17.2	26.4	39.3	56.7	87.2	127.7	189.4	290.9
Male.....	6.6	10.4	20.9	32.2	46.5	62.6	96.2	157.0	214.2	300.5
Female.....	5.3	10.2	14.0	21.7	33.6	52.4	81.0	110.2	176.3	287.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Viral hepatitis (B15-B19)

All origins 3/, both sexes.....	1.8	4.8	8.9	8.0	5.0	4.5	4.5	5.0	4.0	3.1
Male.....	2.5	6.8	13.4	11.9	7.0	5.2	4.7	5.1	3.8	3.0
Female.....	1.1	2.8	4.6	4.2	3.3	3.8	4.3	5.0	4.1	3.1
Hispanic, both sexes.....	2.6	7.0	12.5	14.5	10.7	10.7	10.9	11.2	8.5	8.6
Male.....	4.1	10.0	19.2	20.2	13.7	12.1	11.8	10.3	*	*
Female.....	*	3.8	5.8	9.2	8.0	9.5	10.1	11.9	10.9	*
Mexican, both sexes.....	3.0	7.3	11.4	13.5	11.7	14.1	9.8	11.6	*	*
Male.....	4.7	9.7	17.0	16.3	14.7	*	*	*	*	*
Female.....	*	4.7	5.3	10.5	8.8	15.5	*	*	*	*
Puerto Rican, both sexes.....	*	13.8	27.5	29.0	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	4.7	10.2	12.1	*	*	*	*	*	*
Male.....	*	7.8	16.1	20.4	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.7	4.5	8.5	7.4	4.5	3.9	3.9	4.6	3.7	2.8
Male.....	2.2	6.3	12.7	11.2	6.4	4.7	4.1	4.8	3.7	3.0
Female.....	1.1	2.7	4.5	3.8	2.8	3.3	3.8	4.5	3.8	2.7
Non-Hispanic white 6/, both sexes....	1.7	4.5	8.3	6.4	3.9	3.1	3.4	4.0	3.2	2.5
Male.....	2.2	6.4	12.4	9.8	5.5	3.7	3.6	4.2	3.5	3.1
Female.....	1.1	2.6	4.2	3.2	2.4	2.6	3.3	3.8	3.0	2.2
Non-Hispanic black 6/, both sexes....	1.9	4.9	11.3	14.8	9.2	9.1	5.0	6.7	6.8	*
Male.....	2.4	6.4	16.7	23.8	14.2	11.6	6.3	9.2	*	*
Female.....	1.4	3.5	6.8	7.3	5.3	7.3	*	*	7.8	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All origins 3/, both sexes.....	10.8	10.9	9.3	6.5	4.1	3.1	1.8	1.0	0.6	*
Male.....	15.6	16.1	14.4	10.3	6.9	5.0	3.1	1.9	1.1	*
Female.....	5.9	5.7	4.3	2.8	1.5	1.4	0.8	*	*	*
Hispanic, both sexes.....	10.6	11.8	11.5	7.8	7.4	5.2	4.0	*	*	*
Male.....	15.7	18.9	18.2	12.8	13.1	8.2	7.6	*	*	*
Female.....	4.9	4.3	4.8	3.1	*	*	*	*	*	*
Mexican, both sexes.....	6.0	5.9	6.0	5.4	4.2	*	*	*	*	*
Male.....	9.7	9.9	10.5	8.5	8.4	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	47.1	56.2	53.9	26.1	*	*	*	*	*	*
Male.....	68.0	93.7	82.6	*	*	*	*	*	*	*
Female.....	28.9	26.4	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	8.5	7.8	7.8	*	*	*	*	*	*	*
Male.....	13.4	13.5	13.4	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	10.7	10.6	9.0	6.3	3.7	2.9	1.7	0.9	0.5	*
Male.....	15.5	15.6	13.9	10.1	6.3	4.7	2.8	1.7	*	*
Female.....	6.1	5.8	4.3	2.8	1.4	1.3	0.7	*	*	*
Non-Hispanic white 6/, both sexes....	5.0	4.6	3.9	2.6	1.7	1.3	0.6	0.4	*	*
Male.....	8.5	8.1	6.6	4.5	3.2	2.3	1.2	0.8	*	*
Female.....	1.5	1.2	1.3	0.7	*	*	*	*	*	*
Non-Hispanic black 6/, both sexes....	45.9	50.4	45.7	36.7	22.0	17.0	10.8	5.9	*	*
Male.....	62.3	69.6	71.1	58.8	37.2	27.8	19.3	12.4	*	*
Female.....	31.4	33.8	24.2	18.5	10.2	9.1	5.0	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All origins 3/, both sexes.....	0.8	1.0	1.6	2.3	3.3	4.8	7.1	9.8	13.8	20.4
Male.....	1.0	1.2	2.1	2.8	3.8	5.6	8.1	11.6	17.2	25.1
Female.....	0.6	0.8	1.2	1.8	2.9	4.2	6.3	8.4	11.7	18.2
Hispanic, both sexes.....	1.0	1.1	1.3	1.7	2.7	3.7	7.2	7.1	9.2	10.2
Male.....	*	1.6	2.0	*	*	5.6	*	*	*	*
Female.....	*	*	*	*	*	*	7.4	*	*	*
Mexican, both sexes.....	1.2	1.4	*	*	*	*	7.1	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.7	1.0	1.6	2.3	3.4	4.9	7.1	9.9	14.0	20.9
Male.....	0.9	1.2	2.0	2.9	3.9	5.6	8.2	11.8	17.3	25.8
Female.....	0.5	0.9	1.2	1.8	2.9	4.3	6.2	8.6	12.0	18.6
Non-Hispanic white 6/, both sexes....	0.6	0.8	1.5	2.2	3.2	4.8	7.1	10.0	14.0	21.1
Male.....	0.7	1.0	1.8	2.6	3.5	5.6	8.1	11.6	17.3	26.5
Female.....	0.5	0.7	1.2	1.8	2.9	4.1	6.2	8.8	11.9	18.6
Non-Hispanic black 6/, both sexes....	1.5	2.1	3.3	4.2	4.8	6.7	8.4	11.0	14.3	22.6
Male.....	2.2	2.5	4.8	5.8	6.6	6.3	10.1	15.7	15.4	21.1
Female.....	*	1.8	2.0	2.9	3.4	6.9	7.2	8.2	13.7	23.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms (C00-C97)

All origins 3/, both sexes.....	42.3	84.1	152.1	252.7	414.6	615.1	863.3	1,144.9	1,418.8	1,606.1
Male.....	37.1	80.7	162.1	282.3	476.3	731.6	1,045.7	1,442.3	1,901.9	2,248.7
Female.....	47.6	87.5	142.6	224.8	358.2	513.3	714.8	927.2	1,116.9	1,305.5
Hispanic, both sexes.....	28.6	54.3	101.3	159.3	255.2	386.1	560.2	773.4	950.9	1,077.5
Male.....	21.8	49.1	101.3	183.0	287.2	476.3	708.0	980.6	1,285.7	1,440.1
Female.....	36.1	59.7	101.4	136.9	226.9	310.4	445.1	623.4	728.1	884.9
Mexican, both sexes.....	27.6	52.5	93.6	151.3	242.3	400.8	517.5	770.4	965.4	1,161.8
Male.....	20.1	45.3	91.0	165.3	263.5	457.8	645.2	983.3	1,422.7	*
Female.....	36.4	60.4	96.4	136.7	222.5	344.4	413.5	620.2	690.6	970.9
Puerto Rican, both sexes.....	35.0	67.1	128.8	206.1	349.6	380.6	859.5	*	*	*
Male.....	29.1	71.0	148.3	258.0	*	*	*	*	*	*
Female.....	40.2	64.0	113.3	152.2	256.8	284.1	*	*	*	*
Cuban, both sexes.....	30.1	67.6	136.5	206.8	324.6	552.7	628.8	684.5	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	28.4	51.6	104.1	147.0	221.5	312.7	516.8	805.3	840.3	932.7
Male.....	23.3	48.5	101.0	160.0	233.8	423.8	620.9	*	*	*
Female.....	33.7	54.8	106.6	136.5	211.6	240.2	436.1	723.5	*	*
Non-Hispanic 5/, both sexes.....	44.5	87.7	157.2	260.6	427.2	632.4	884.8	1,168.3	1,443.4	1,630.6
Male.....	39.6	84.6	168.2	290.6	490.8	750.2	1,068.5	1,471.1	1,935.4	2,289.8
Female.....	49.2	90.7	146.7	232.4	369.0	529.1	734.9	946.6	1,136.9	1,324.2
Non-Hispanic white 6/, both sexes....	42.5	83.6	150.5	253.4	420.7	630.5	889.2	1,176.2	1,452.4	1,643.1
Male.....	38.3	80.9	159.5	278.9	478.0	739.5	1,061.5	1,468.4	1,937.1	2,300.2
Female.....	46.6	86.2	141.7	228.7	367.2	532.8	745.4	957.8	1,147.3	1,336.7
Non-Hispanic black 6/, both sexes....	61.5	126.9	229.8	367.1	573.9	786.1	1,021.8	1,306.2	1,581.1	1,700.4
Male.....	52.5	123.6	263.4	450.9	720.7	1,027.7	1,377.1	1,816.6	2,319.6	2,577.4
Female.....	69.5	129.8	201.4	298.1	459.7	607.9	778.1	1,001.4	1,191.6	1,342.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All origins 3/, both sexes.....	0.9	1.8	3.5	5.2	6.6	8.7	10.6	13.0	14.1	16.3
Male.....	1.2	2.9	5.6	8.4	10.6	13.9	16.0	19.7	21.4	21.9
Female.....	0.5	0.8	1.4	2.1	2.9	4.2	6.3	8.1	9.6	13.7
Hispanic, both sexes.....	*	*	1.9	2.5	2.6	6.6	7.3	8.2	9.8	12.9
Male.....	*	*	2.8	4.8	4.5	11.0	12.6	16.5	*	*
Female.....	*	*	*	*	*	*	*	*	*	12.6
Mexican, both sexes.....	*	*	1.7	*	*	7.3	6.8	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.9	2.0	3.6	5.4	6.9	8.8	10.8	13.3	14.3	16.5
Male.....	1.4	3.1	5.9	8.7	11.1	14.0	16.2	19.9	21.6	22.3
Female.....	0.5	0.9	1.5	2.2	3.1	4.3	6.5	8.5	9.7	13.8
Non-Hispanic white 6/, both sexes....	0.8	1.8	3.4	5.0	6.5	8.4	10.6	13.3	14.5	17.1
Male.....	1.2	2.8	5.5	8.0	10.3	13.0	15.5	19.6	21.5	23.0
Female.....	0.5	0.8	1.3	2.1	3.0	4.3	6.5	8.7	10.1	14.4
Non-Hispanic black 6/, both sexes....	1.2	3.3	5.7	8.7	11.3	13.5	14.4	14.5	12.9	10.1
Male.....	1.7	5.2	9.4	15.5	20.3	25.6	25.3	25.8	26.0	*
Female.....	*	1.6	2.6	3.1	4.3	4.6	6.9	7.7	*	8.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of esophagus (C15)

All origins 3/, both sexes.....	0.9	2.3	4.8	8.4	13.2	17.4	22.0	26.9	27.0	25.6
Male.....	1.5	4.1	8.1	15.0	23.2	30.8	38.7	47.0	51.3	48.5
Female.....	0.2	0.7	1.7	2.2	4.1	5.8	8.4	12.2	11.8	14.9
Hispanic, both sexes.....	*	0.9	2.4	4.1	6.1	7.5	13.0	15.4	16.7	14.9
Male.....	*	*	4.4	7.4	10.4	14.4	21.7	27.3	31.1	25.9
Female.....	*	*	*	*	*	*	6.2	*	*	*
Mexican, both sexes.....	*	*	2.2	3.5	5.8	5.5	11.6	15.1	19.1	*
Male.....	*	*	3.8	5.5	10.2	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.0	2.5	5.1	8.7	13.7	18.2	22.6	27.6	27.6	26.1
Male.....	1.7	4.4	8.5	15.6	24.1	32.0	39.8	48.3	52.6	49.8
Female.....	0.2	0.7	1.8	2.3	4.2	6.1	8.6	12.5	12.0	15.1
Non-Hispanic white 6/, both sexes....	1.0	2.5	4.9	8.7	13.9	18.4	23.2	28.2	28.5	27.0
Male.....	1.9	4.5	8.4	15.7	24.4	32.4	40.6	49.3	55.0	51.3
Female.....	*	0.6	1.6	1.8	4.0	5.9	8.6	12.5	11.8	15.7
Non-Hispanic black 6/, both sexes....	0.9	3.1	7.6	11.9	16.9	21.9	23.2	25.5	24.0	18.3
Male.....	*	5.0	12.5	19.3	29.2	38.9	42.7	45.1	40.0	35.7
Female.....	*	1.4	3.5	5.8	7.4	9.4	9.8	13.7	15.6	11.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of stomach (C16)

All origins 3/, both sexes.....	1.3	2.1	3.3	5.0	7.2	11.0	15.5	22.3	29.9	35.8
Male.....	1.4	2.6	4.4	7.0	10.1	15.4	21.4	31.3	44.7	50.2
Female.....	1.2	1.7	2.3	3.1	4.5	7.2	10.6	15.8	20.7	29.1
Hispanic, both sexes.....	2.0	3.1	4.9	9.1	11.7	14.8	27.5	35.5	39.9	47.0
Male.....	1.8	3.6	5.6	11.1	16.0	20.3	36.2	43.8	58.1	55.3
Female.....	2.3	2.6	4.2	7.1	8.0	10.2	20.8	29.4	27.8	42.5
Mexican, both sexes.....	1.9	3.0	4.8	9.0	12.2	16.6	26.8	33.9	46.3	55.7
Male.....	*	3.5	5.9	11.5	15.8	19.7	32.5	41.4	*	*
Female.....	*	*	3.7	6.5	8.8	13.5	22.1	28.5	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.1	3.8	6.1	12.1	11.7	12.4	28.4	45.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	11.8	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.2	2.0	3.2	4.6	6.8	10.7	14.5	21.4	29.3	35.2
Male.....	1.3	2.4	4.3	6.6	9.6	15.0	20.3	30.4	43.9	49.9
Female.....	1.1	1.5	2.1	2.7	4.2	6.9	9.9	14.8	20.3	28.4
Non-Hispanic white 6/, both sexes....	0.8	1.5	2.7	3.8	5.5	8.9	12.2	18.3	25.2	31.5
Male.....	0.9	2.0	3.7	5.6	7.8	12.7	17.2	26.5	37.9	45.2
Female.....	0.8	1.1	1.7	2.1	3.4	5.5	7.9	12.2	17.2	25.2
Non-Hispanic black 6/, both sexes....	2.4	3.6	5.5	8.5	15.2	21.3	29.1	42.7	65.2	68.2
Male.....	3.1	4.8	7.9	12.8	22.6	31.7	38.9	68.6	102.5	85.2
Female.....	1.8	2.5	3.4	4.9	9.4	13.7	22.4	27.2	45.5	61.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All origins 3/, both sexes.....	4.0	7.6	13.5	21.8	34.8	52.0	73.9	104.4	146.8	207.5
Male.....	4.2	8.5	15.1	25.6	43.1	64.9	91.9	127.3	181.3	234.6
Female.....	3.8	6.7	11.9	18.2	27.2	40.8	59.2	87.6	125.3	194.9
Hispanic, both sexes.....	2.6	5.2	10.5	17.1	25.2	38.4	55.1	82.4	110.9	142.4
Male.....	2.2	5.7	10.6	22.4	30.1	51.9	71.7	101.5	131.0	157.9
Female.....	3.1	4.8	10.3	12.2	20.9	27.0	42.2	68.6	97.6	134.2
Mexican, both sexes.....	2.3	5.4	9.9	17.5	24.4	38.2	45.5	82.5	88.3	146.1
Male.....	*	5.3	10.3	22.3	29.5	51.6	62.9	104.0	*	*
Female.....	2.8	5.4	9.5	12.4	19.6	25.0	31.3	67.3	67.2	133.5
Puerto Rican, both sexes.....	*	*	*	16.6	32.0	43.5	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.5	4.1	10.2	15.6	22.2	31.9	46.4	76.1	109.3	107.6
Male.....	*	*	*	19.2	*	*	*	*	*	*
Female.....	*	*	9.8	12.7	20.5	26.0	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.2	7.9	13.7	22.2	35.6	53.1	75.1	105.8	148.7	210.6
Male.....	4.6	8.9	15.5	25.8	44.1	65.9	93.0	128.9	184.1	238.4
Female.....	3.9	7.0	12.1	18.7	27.8	41.9	60.5	88.8	126.6	197.7
Non-Hispanic white 6/, both sexes....	3.8	7.3	12.7	21.0	33.4	50.9	73.7	103.0	146.2	210.5
Male.....	4.2	8.1	14.3	24.6	41.6	63.4	90.5	125.8	179.6	236.7
Female.....	3.5	6.5	11.2	17.4	25.8	39.7	59.7	86.0	125.2	198.2
Non-Hispanic black 6/, both sexes....	7.1	12.9	22.9	34.7	58.5	80.6	102.9	150.4	200.5	244.7
Male.....	7.6	15.1	26.9	40.7	72.1	102.2	139.9	192.0	271.8	314.3
Female.....	6.7	11.0	19.6	29.7	48.0	64.7	77.5	125.6	162.9	216.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All origins 3/, both sexes.....	1.1	3.2	8.2	11.5	13.1	17.2	23.9	30.2	34.1	33.5
Male.....	1.6	5.0	13.3	18.5	19.6	25.7	33.7	42.2	48.0	43.8
Female.....	0.7	1.5	3.3	4.9	7.1	9.8	15.9	21.4	25.4	28.7
Hispanic, both sexes.....	1.2	4.1	11.5	15.4	19.8	27.2	37.2	50.6	53.0	55.2
Male.....	1.8	6.1	18.6	26.1	29.7	40.1	53.4	64.9	76.2	62.0
Female.....	*	2.0	4.5	5.3	11.1	16.3	24.6	40.2	37.6	51.5
Mexican, both sexes.....	1.5	4.8	13.0	14.7	21.5	31.9	38.1	60.2	60.5	67.8
Male.....	2.4	6.9	19.3	23.0	30.5	42.5	52.3	75.1	*	*
Female.....	*	*	6.0	6.0	13.1	21.5	26.4	49.6	40.5	*
Puerto Rican, both sexes.....	*	*	14.0	32.0	36.3	*	*	*	*	*
Male.....	*	*	*	59.3	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	8.6	11.3	9.4	20.0	32.5	43.1	*	*
Male.....	*	*	15.6	18.7	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.1	3.1	7.8	11.1	12.5	16.4	22.9	28.8	33.0	32.3
Male.....	1.6	4.8	12.6	17.8	18.8	24.5	32.2	40.6	46.2	42.6
Female.....	0.7	1.4	3.1	4.8	6.7	9.3	15.2	20.1	24.8	27.6
Non-Hispanic white 6/, both sexes....	0.9	2.5	6.6	9.3	10.8	14.6	21.4	27.2	31.5	31.7
Male.....	1.1	3.8	10.6	14.4	15.7	21.4	30.1	38.7	44.8	42.3
Female.....	0.7	1.1	2.7	4.3	6.2	8.5	14.1	18.6	23.1	26.8
Non-Hispanic black 6/, both sexes....	1.8	5.4	14.7	23.1	21.0	22.6	27.0	31.2	36.6	31.6
Male.....	3.1	8.0	25.1	43.4	36.3	37.7	43.1	46.0	45.6	34.8
Female.....	*	3.1	5.9	6.4	9.1	11.4	16.1	22.3	31.8	30.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of pancreas (C25)

All origins 3/, both sexes.....	1.7	4.4	8.5	15.3	26.2	37.7	53.4	70.2	89.7	96.9
Male.....	2.0	5.2	10.7	19.2	31.9	45.3	61.2	77.3	99.0	102.1
Female.....	1.4	3.5	6.3	11.7	21.0	31.1	47.0	64.9	83.9	94.4
Hispanic, both sexes.....	0.8	3.2	5.8	9.5	19.9	28.8	40.1	57.9	62.8	77.1
Male.....	*	3.4	7.1	11.1	21.7	35.6	45.4	65.4	74.5	65.4
Female.....	*	3.0	4.6	8.0	18.3	23.2	35.9	52.5	55.0	83.3
Mexican, both sexes.....	*	3.0	4.7	10.3	19.0	29.4	36.9	58.6	65.4	76.5
Male.....	*	2.7	5.5	12.0	20.4	34.4	39.1	62.6	*	*
Female.....	*	3.2	3.7	8.6	17.6	24.5	35.1	55.7	53.4	91.8
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	3.6	6.3	7.5	17.9	30.0	38.7	70.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	20.5	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.8	4.5	8.7	15.8	26.7	38.4	54.3	70.8	91.1	97.7
Male.....	2.2	5.4	11.1	19.9	32.6	46.0	62.2	77.9	100.2	104.0
Female.....	1.5	3.6	6.5	12.0	21.3	31.7	47.8	65.7	85.4	94.8
Non-Hispanic white 6/, both sexes....	1.8	4.3	8.4	15.4	26.0	37.9	53.4	70.6	90.8	96.9
Male.....	2.1	5.3	10.7	19.4	32.1	45.8	61.9	78.4	101.5	105.3
Female.....	1.4	3.3	6.2	11.5	20.4	30.8	46.4	64.7	84.1	93.0
Non-Hispanic black 6/, both sexes....	2.4	6.4	13.0	22.2	38.1	50.4	68.5	86.7	102.1	114.7
Male.....	2.8	6.8	17.3	29.3	46.9	59.9	76.9	88.8	100.4	92.5
Female.....	2.0	6.0	9.3	16.3	31.2	43.3	62.8	85.5	102.9	123.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of larynx (C32)

All origins 3/, both sexes.....	0.3	0.7	1.4	2.6	3.8	5.2	6.2	6.9	7.4	6.3
Male.....	0.3	1.2	2.3	4.4	6.2	9.2	10.5	12.5	14.7	14.8
Female.....	*	0.3	0.5	1.0	1.5	1.7	2.7	2.8	2.8	2.3
Hispanic, both sexes.....	*	*	*	1.5	*	3.0	3.7	5.6	8.5	*
Male.....	*	*	*	*	*	5.9	7.6	11.8	18.8	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.3	0.8	1.4	2.7	3.9	5.4	6.4	7.0	7.3	6.3
Male.....	0.4	1.2	2.4	4.5	6.4	9.4	10.7	12.5	14.4	14.7
Female.....	*	0.3	0.5	1.0	1.6	1.8	2.9	3.0	2.9	2.4
Non-Hispanic white 6/, both sexes....	0.2	0.7	1.3	2.5	3.7	4.9	6.2	6.9	7.1	6.2
Male.....	0.3	1.0	2.1	4.1	6.0	8.4	10.1	12.0	13.9	14.5
Female.....	*	0.3	0.5	0.9	1.6	1.8	3.0	3.1	2.8	2.3
Non-Hispanic black 6/, both sexes.....	*	1.6	2.9	5.5	6.8	11.2	9.9	10.5	9.7	8.2
Male.....	*	2.9	5.2	9.6	12.6	22.2	19.9	23.0	21.8	19.2
Female.....	*	*	*	2.0	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All origins 3/, both sexes.....	7.1	19.9	39.4	71.8	136.5	216.9	297.1	366.8	382.3	300.5
Male.....	7.3	21.6	47.5	88.3	167.8	272.4	378.8	486.0	544.1	457.7
Female.....	6.9	18.3	31.6	56.2	107.8	168.4	230.7	279.5	281.3	226.9
Hispanic, both sexes.....	2.5	5.6	12.1	23.0	43.4	87.0	114.0	159.6	180.6	160.8
Male.....	2.5	6.3	16.1	31.6	64.5	122.7	163.6	236.9	281.7	263.9
Female.....	2.4	4.8	8.2	14.9	24.7	57.1	75.4	103.6	113.4	106.0
Mexican, both sexes.....	1.9	4.1	8.5	18.9	35.2	83.5	104.7	156.6	183.3	189.6
Male.....	*	5.0	10.6	23.7	51.2	109.8	147.7	229.2	312.5	*
Female.....	*	3.2	6.1	13.9	20.3	57.5	69.6	105.3	105.7	132.1
Puerto Rican, both sexes.....	*	10.5	22.1	36.7	60.6	83.3	*	*	*	*
Male.....	*	*	*	53.5	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	93.5	166.8	161.1	156.5	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	5.6	12.9	19.4	39.4	72.4	98.2	170.3	153.0	106.1
Male.....	*	*	18.8	23.2	51.5	113.5	*	*	*	*
Female.....	*	*	*	16.3	29.6	45.7	78.6	*	*	*
Non-Hispanic 5/, both sexes.....	7.8	21.7	42.2	76.0	144.0	227.0	310.5	380.2	393.3	306.9
Male.....	8.1	23.6	50.8	93.2	175.9	283.6	393.9	502.0	558.8	467.4
Female.....	7.6	19.9	34.0	59.8	114.8	177.4	242.5	290.9	290.2	232.3
Non-Hispanic white 6/, both sexes....	7.8	21.1	40.4	75.4	144.8	232.0	318.8	390.7	400.4	313.0
Male.....	8.0	22.4	47.6	89.8	172.3	283.6	396.7	507.6	564.4	474.2
Female.....	7.6	19.7	33.4	61.5	119.1	185.8	253.8	303.2	297.2	237.8
Non-Hispanic black 6/, both sexes....	9.7	31.2	64.1	100.2	177.7	250.8	311.7	361.4	384.1	268.4
Male.....	10.6	36.9	84.5	144.3	259.1	371.5	461.2	553.8	606.1	448.1
Female.....	9.0	26.2	46.9	63.9	114.4	161.8	209.1	246.6	267.0	195.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of breast (C50)

All origins 3/, both sexes.....	6.9	11.6	17.0	24.7	32.7	38.0	47.1	63.1	82.7	116.7
Male.....	*	*	0.3	0.4	0.8	0.8	1.2	1.8	2.6	3.0
Female.....	13.8	22.7	33.0	47.5	61.9	70.4	84.4	108.0	132.9	169.9
Hispanic, both sexes.....	5.1	7.9	12.1	16.0	23.4	23.4	30.1	38.5	44.8	58.7
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	10.6	16.2	24.0	31.1	43.9	42.6	53.2	65.2	74.1	89.9
Mexican, both sexes.....	4.9	8.0	10.9	15.2	22.3	22.6	23.5	36.3	44.4	55.7
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	10.5	16.5	22.8	31.1	42.8	45.0	42.1	61.8	70.1	89.0
Puerto Rican, both sexes.....	*	10.1	14.6	16.6	33.7	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.3	6.8	13.7	17.4	19.9	20.0	33.9	37.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	8.8	13.6	25.2	31.3	36.0	33.1	60.2	*	*	*
Non-Hispanic 5/, both sexes.....	7.2	12.0	17.5	25.4	33.5	39.1	48.3	64.7	84.8	119.4
Male.....	*	*	0.3	0.5	0.8	0.8	1.3	1.8	2.7	3.2
Female.....	14.2	23.5	33.9	48.9	63.4	72.6	86.7	110.8	135.9	173.5
Non-Hispanic white 6/, both sexes....	6.5	10.7	16.1	23.9	32.6	38.9	48.1	65.0	85.1	119.8
Male.....	*	*	*	0.4	0.8	0.8	1.2	1.8	2.6	3.3
Female.....	12.8	21.2	31.7	46.6	62.3	72.9	87.3	112.2	137.0	174.1
Non-Hispanic black 6/, both sexes....	13.0	22.0	28.3	40.1	48.1	50.5	59.8	79.5	106.7	144.7
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	24.3	40.8	51.4	72.4	84.6	86.6	99.2	125.9	160.7	202.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of cervix uteri (C53)

All origins 3/, both sexes.....	1.7	2.1	2.4	2.5	2.7	3.1	3.4	3.6	4.1	5.2
Male.....
Female.....	3.4	4.1	4.7	4.9	5.2	5.7	6.1	6.2	6.6	7.7
Hispanic, both sexes.....	1.5	2.3	3.6	3.3	3.2	3.9	5.0	4.8	9.5	*
Male.....
Female.....	3.1	4.6	7.1	6.5	6.0	7.1	8.9	8.2	15.8	*
Mexican, both sexes.....	1.7	2.5	3.9	2.9	*	*	*	*	*	*
Male.....
Female.....	3.7	5.1	8.2	6.0	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.7	2.0	2.3	2.5	2.7	3.0	3.2	3.5	3.7	5.2
Male.....
Female.....	3.4	4.0	4.5	4.8	5.1	5.6	5.9	6.0	6.1	7.6
Non-Hispanic white 6/, both sexes....	1.5	1.9	2.0	2.3	2.4	2.5	2.6	2.9	3.3	4.1
Male.....
Female.....	3.0	3.7	4.0	4.6	4.6	4.8	4.8	5.1	5.3	6.1
Non-Hispanic black 6/, both sexes....	3.3	3.2	4.1	3.4	5.0	6.9	8.8	8.4	9.2	17.5
Male.....
Female.....	6.2	6.0	7.6	6.2	8.9	11.9	14.8	13.5	14.1	24.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All origins 3/, both sexes.....	0.4	1.0	1.9	3.8	6.2	8.8	11.6	14.3	17.6	21.6
Male.....
Female.....	0.9	1.9	3.6	7.4	11.9	16.4	21.0	24.7	28.6	31.6
Hispanic, both sexes.....	*	1.0	1.4	2.9	4.4	5.7	7.7	11.5	11.8	15.3
Male.....
Female.....	*	2.1	2.9	5.6	8.3	10.4	13.7	19.7	19.6	23.4
Mexican, both sexes.....	*	*	*	*	*	6.0	6.8	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.5	1.0	1.9	3.9	6.4	9.0	11.9	14.4	17.9	21.9
Male.....
Female.....	0.9	1.9	3.7	7.5	12.2	16.8	21.6	25.0	29.0	32.0
Non-Hispanic white 6/, both sexes....	0.5	0.9	1.8	3.5	5.6	7.7	10.5	13.4	17.2	22.0
Male.....
Female.....	0.9	1.7	3.7	6.9	10.8	14.6	19.3	23.5	28.0	32.2
Non-Hispanic black 6/, both sexes.....	*	1.5	2.1	7.1	14.3	21.0	25.9	27.9	31.0	26.3
Male.....
Female.....	*	2.7	3.9	13.0	25.4	36.5	43.7	44.5	47.4	37.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of ovary (C56)

All origins 3/, both sexes.....	1.3	2.8	5.1	7.7	12.4	16.3	22.0	28.3	34.8	37.8
Male.....
Female.....	2.5	5.5	9.9	14.9	23.7	30.6	39.9	49.1	56.5	55.5
Hispanic, both sexes.....	0.9	1.9	3.5	4.1	8.8	10.2	16.2	24.9	22.6	27.8
Male.....
Female.....	1.8	3.8	7.0	8.0	16.7	18.7	28.8	42.9	37.6	42.5
Mexican, both sexes.....	*	1.9	3.6	4.1	8.0	11.1	13.4	25.1	19.1	*
Male.....
Female.....	*	3.9	7.5	8.4	15.4	22.0	24.3	42.8	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	9.4	*	20.1	*	*	*
Male.....
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.3	2.9	5.2	8.0	12.7	16.8	22.4	28.5	35.4	38.3
Male.....
Female.....	2.6	5.6	10.2	15.5	24.3	31.5	40.7	49.5	57.4	56.0
Non-Hispanic white 6/, both sexes....	1.4	3.0	5.4	8.3	13.1	17.5	23.7	29.3	36.7	39.3
Male.....
Female.....	2.7	5.9	10.6	16.3	25.5	33.1	43.4	51.1	59.8	57.6
Non-Hispanic black 6/, both sexes....	1.4	2.7	4.5	6.6	11.5	15.8	16.5	25.8	25.2	33.7
Male.....
Female.....	2.6	5.1	8.3	12.1	20.4	27.4	27.8	41.2	38.5	47.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All origins 3/, both sexes.....	0.8	2.0	3.7	6.5	9.6	14.8	18.2	24.0	28.9	34.3
Male.....	1.2	2.8	5.4	9.5	14.0	21.6	25.0	34.5	40.8	52.1
Female.....	0.4	1.2	2.1	3.6	5.6	8.8	12.6	16.4	21.5	25.9
Hispanic, both sexes.....	*	1.5	3.4	6.5	7.4	10.6	16.5	20.5	24.5	23.5
Male.....	*	2.1	4.9	8.9	10.8	14.7	26.7	30.9	34.4	32.7
Female.....	*	*	*	4.3	4.4	7.1	8.6	13.0	18.0	18.6
Mexican, both sexes.....	*	1.7	3.8	7.7	8.6	14.3	17.5	25.5	29.6	29.6
Male.....	*	*	5.2	9.7	11.9	17.7	27.2	37.6	*	*
Female.....	*	*	*	5.7	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.9	2.1	3.7	6.4	9.8	15.1	18.3	24.2	29.2	34.8
Male.....	1.4	2.9	5.4	9.5	14.3	22.1	24.8	34.6	41.1	53.2
Female.....	0.4	1.2	2.1	3.5	5.7	8.9	12.9	16.6	21.7	26.2
Non-Hispanic white 6/, both sexes....	0.8	2.1	3.7	6.7	10.2	15.6	18.8	25.1	29.9	35.5
Male.....	1.3	2.9	5.5	9.6	14.7	22.6	25.5	35.8	42.2	54.6
Female.....	0.4	1.3	2.0	3.8	6.0	9.4	13.2	17.1	22.1	26.6
Non-Hispanic black 6/, both sexes....	1.4	2.5	4.9	6.7	8.6	13.6	16.9	19.8	26.4	33.7
Male.....	2.2	4.0	6.9	11.3	13.9	22.7	23.1	28.1	40.0	49.5
Female.....	*	*	3.3	3.0	4.5	6.9	12.6	14.8	19.3	27.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of bladder (C67)

All origins 3/, both sexes.....	0.3	0.8	1.7	3.5	7.0	11.2	19.2	30.0	46.9	67.4
Male.....	0.5	1.1	2.6	5.5	11.1	17.7	31.4	51.4	85.8	131.6
Female.....	*	0.5	0.8	1.6	3.3	5.6	9.4	14.4	22.6	37.4
Hispanic, both sexes.....	*	*	*	1.5	3.4	7.2	9.4	16.2	24.9	33.3
Male.....	*	*	*	*	5.5	9.3	13.4	27.8	46.7	54.1
Female.....	*	*	*	*	*	5.4	6.2	7.8	*	22.2
Mexican, both sexes.....	*	*	*	*	*	*	*	12.7	26.5	29.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.3	0.8	1.8	3.6	7.3	11.6	20.0	30.9	48.1	69.0
Male.....	0.5	1.2	2.8	5.8	11.5	18.4	32.7	52.9	88.0	135.5
Female.....	*	0.5	0.9	1.6	3.4	5.6	9.6	14.8	23.2	38.1
Non-Hispanic white 6/, both sexes....	0.3	0.9	1.9	3.8	7.7	12.2	20.8	32.3	50.0	71.5
Male.....	0.5	1.3	3.0	6.0	12.1	19.6	34.7	56.0	92.5	142.3
Female.....	*	0.5	0.9	1.6	3.6	5.6	9.2	14.5	23.2	38.5
Non-Hispanic black 6/, both sexes.....	*	0.8	1.7	3.9	6.4	9.8	18.4	26.2	39.5	55.7
Male.....	*	*	2.9	5.7	10.8	12.8	25.3	33.1	57.6	95.3
Female.....	*	*	*	2.4	3.0	7.7	13.7	22.0	30.0	39.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All origins 3/, both sexes.....	2.3	3.5	5.7	8.1	10.8	13.0	16.8	18.5	19.6	16.7
Male.....	2.9	4.3	6.8	9.9	13.5	16.4	20.4	22.9	25.6	21.2
Female.....	1.8	2.7	4.6	6.4	8.4	10.0	13.9	15.3	15.8	14.6
Hispanic, both sexes.....	1.6	2.3	3.5	4.5	6.6	6.6	8.7	14.1	11.5	12.1
Male.....	2.1	2.9	3.9	6.2	5.7	7.6	11.4	17.0	*	*
Female.....	*	1.8	3.2	3.0	7.4	5.7	6.5	11.9	*	14.4
Mexican, both sexes.....	1.3	2.1	2.9	4.3	7.6	6.0	8.0	12.3	*	*
Male.....	*	2.6	*	5.3	*	*	*	*	*	*
Female.....	*	*	*	*	8.2	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	5.3	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.5	3.6	5.9	8.4	11.2	13.5	17.4	18.8	20.0	16.9
Male.....	3.1	4.4	7.1	10.2	14.2	17.1	21.0	23.2	26.2	21.9
Female.....	1.9	2.8	4.7	6.7	8.5	10.3	14.4	15.5	16.2	14.6
Non-Hispanic white 6/, both sexes....	2.8	4.1	6.6	9.1	12.4	14.6	18.8	20.4	21.1	17.7
Male.....	3.5	4.9	7.8	10.9	15.6	18.5	22.6	25.1	27.2	22.9
Female.....	2.1	3.2	5.3	7.4	9.4	11.2	15.7	16.9	17.3	15.3
Non-Hispanic black 6/, both sexes....	1.3	1.9	3.3	5.6	5.1	7.2	9.4	7.4	12.6	9.6
Male.....	1.5	2.5	4.1	7.6	6.6	8.7	12.7	*	19.7	*
Female.....	*	1.5	2.6	4.0	4.0	6.1	7.2	7.1	8.9	9.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All origins 3/, both sexes.....	4.2	6.3	11.0	18.5	32.4	51.7	81.7	119.0	162.2	186.1
Male.....	5.1	7.4	14.0	23.4	41.2	65.7	103.4	157.6	219.4	244.7
Female.....	3.3	5.2	8.2	13.9	24.3	39.4	64.0	90.8	126.3	158.7
Hispanic, both sexes.....	4.1	5.3	9.5	14.9	24.0	41.8	63.6	82.2	110.9	100.9
Male.....	4.8	6.6	10.5	18.4	28.5	48.0	77.4	103.0	135.9	129.7
Female.....	3.3	3.9	8.6	11.5	20.1	36.7	52.9	67.1	94.3	85.7
Mexican, both sexes.....	3.8	5.2	8.2	13.0	23.7	45.0	65.4	70.9	113.6	106.1
Male.....	4.1	6.1	9.2	15.7	27.7	45.5	77.5	90.5	*	*
Female.....	3.5	4.2	7.0	10.3	19.9	44.5	55.6	57.1	91.9	89.0
Puerto Rican, both sexes.....	*	*	*	20.1	31.2	34.3	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	92.5	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.6	5.4	10.8	16.1	22.2	32.9	50.5	97.2	112.0	98.7
Male.....	5.4	7.2	*	21.5	*	*	*	*	*	*
Female.....	*	*	11.3	11.8	20.5	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.3	6.4	11.2	18.8	33.0	52.4	82.9	121.4	164.8	190.1
Male.....	5.2	7.5	14.3	23.8	42.2	67.0	105.2	161.1	224.0	250.7
Female.....	3.3	5.4	8.2	14.1	24.6	39.5	64.8	92.3	128.0	162.0
Non-Hispanic white 6/, both sexes....	4.1	6.0	11.0	18.4	32.6	53.1	84.9	125.0	171.1	196.6
Male.....	4.9	6.9	14.0	23.2	41.6	67.8	107.5	165.4	232.9	258.6
Female.....	3.3	5.0	8.0	13.7	24.1	40.0	66.0	94.7	132.2	167.7
Non-Hispanic black 6/, both sexes....	6.2	10.3	14.9	26.6	43.8	57.8	85.0	111.2	129.2	145.5
Male.....	8.3	12.7	19.6	35.3	57.3	77.3	112.7	146.9	169.9	198.8
Female.....	4.4	8.2	11.0	19.5	33.4	43.5	66.0	89.9	107.8	123.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Non-Hodgkin's lymphoma (C82-C85)

All origins 3/, both sexes.....	1.5	2.5	4.2	6.7	12.5	19.0	30.1	46.2	63.4	72.3
Male.....	1.9	3.1	5.6	8.8	16.0	23.8	37.2	57.3	80.7	89.6
Female.....	1.1	1.8	2.8	4.7	9.3	14.8	24.3	38.2	52.6	64.2
Hispanic, both sexes.....	1.4	1.9	3.8	6.5	10.6	15.6	23.2	34.8	48.1	40.3
Male.....	1.7	2.4	4.2	8.6	12.5	17.5	28.2	40.7	59.0	54.1
Female.....	*	*	3.4	4.5	8.9	14.0	19.3	30.6	40.9	33.0
Mexican, both sexes.....	1.5	1.5	3.2	5.5	11.7	18.1	23.8	32.7	51.2	37.4
Male.....	*	*	3.6	7.6	13.0	18.2	25.8	39.5	*	*
Female.....	*	*	*	*	10.5	18.0	22.1	27.9	41.5	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	6.8	*	12.4	*	39.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.5	2.5	4.2	6.7	12.6	19.2	30.6	47.0	64.3	73.9
Male.....	2.0	3.2	5.8	8.8	16.2	24.3	37.8	58.3	81.9	91.5
Female.....	1.1	1.9	2.7	4.7	9.3	14.8	24.7	38.6	53.3	65.7
Non-Hispanic white 6/, both sexes....	1.3	2.3	4.3	6.6	12.9	20.1	32.6	50.0	68.0	78.2
Male.....	1.8	2.9	5.8	8.7	16.6	25.1	40.1	61.8	87.0	96.2
Female.....	0.9	1.8	2.8	4.7	9.5	15.6	26.3	41.2	56.1	69.8
Non-Hispanic black 6/, both sexes....	2.5	4.3	4.8	8.7	12.5	15.4	19.7	25.6	32.0	34.5
Male.....	3.5	6.1	7.2	12.2	16.6	20.5	24.1	30.8	35.8	43.1
Female.....	1.6	2.7	2.9	5.8	9.4	11.6	16.7	22.5	30.0	31.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Leukemia (C91-C95)

All origins 3/, both sexes.....	1.8	2.5	4.1	7.0	11.6	19.9	31.3	44.7	62.9	76.9
Male.....	2.0	2.7	4.9	8.7	15.1	26.3	41.6	64.8	91.8	105.1
Female.....	1.7	2.3	3.3	5.3	8.5	14.3	22.8	30.0	44.8	63.7
Hispanic, both sexes.....	1.9	2.3	3.3	4.8	7.5	13.3	21.7	24.9	36.3	35.2
Male.....	2.0	2.9	3.2	5.5	9.4	16.6	26.7	36.1	47.5	42.9
Female.....	1.7	1.8	3.4	4.1	5.8	10.4	17.8	16.8	28.9	31.2
Mexican, both sexes.....	1.8	2.6	2.8	4.2	6.8	13.6	22.9	21.1	35.8	44.4
Male.....	*	3.0	*	*	8.4	17.2	31.1	*	*	*
Female.....	*	*	*	*	*	*	16.2	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	5.8	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.8	2.5	4.1	7.1	11.9	20.4	32.0	46.0	64.3	78.9
Male.....	2.0	2.7	5.1	9.0	15.5	27.1	42.7	66.7	94.2	108.3
Female.....	1.6	2.3	3.3	5.4	8.7	14.6	23.2	30.8	45.7	65.2
Non-Hispanic white 6/, both sexes....	1.8	2.5	4.3	7.3	12.1	21.3	33.3	47.4	67.6	82.0
Male.....	1.9	2.7	5.2	9.2	15.8	28.0	44.4	68.4	98.8	112.1
Female.....	1.7	2.3	3.4	5.6	8.7	15.4	24.0	31.7	47.9	68.0
Non-Hispanic black 6/, both sexes....	2.2	2.7	4.2	7.5	13.1	18.2	29.2	39.9	42.4	56.8
Male.....	3.1	2.7	5.3	10.1	17.0	27.3	39.3	59.4	62.5	87.0
Female.....	1.5	2.7	3.2	5.4	10.1	11.4	22.4	28.3	31.8	44.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All origins 3/, both sexes.....	0.5	1.1	2.4	4.3	7.5	11.8	19.0	25.8	33.3	33.8
Male.....	0.6	1.2	2.9	5.1	9.2	14.3	22.9	33.1	43.8	45.8
Female.....	0.4	0.9	1.9	3.6	6.0	9.6	15.8	20.5	26.8	28.2
Hispanic, both sexes.....	*	0.9	2.2	3.1	5.2	12.0	17.0	20.1	23.9	22.7
Male.....	*	*	2.8	3.7	5.9	12.4	20.6	24.2	25.4	30.4
Female.....	*	*	*	2.6	4.5	11.6	14.3	17.1	22.9	18.6
Mexican, both sexes.....	*	*	1.8	2.9	4.7	11.8	17.2	13.5	22.8	*
Male.....	*	*	*	*	*	*	19.2	*	*	*
Female.....	*	*	*	*	*	15.5	15.7	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.5	1.1	2.4	4.4	7.7	11.7	19.1	26.2	33.8	34.4
Male.....	0.6	1.2	2.9	5.2	9.4	14.4	23.1	33.6	44.7	46.6
Female.....	0.4	1.0	2.0	3.6	6.2	9.4	15.9	20.7	26.9	28.6
Non-Hispanic white 6/, both sexes....	0.5	0.8	2.0	3.8	6.8	10.7	17.7	25.1	32.8	33.3
Male.....	0.6	1.0	2.5	4.6	8.2	13.4	21.3	32.5	43.9	45.9
Female.....	0.4	0.7	1.5	3.1	5.4	8.3	14.7	19.6	25.9	27.5
Non-Hispanic black 6/, both sexes....	1.0	2.7	5.5	9.8	17.6	23.4	34.8	44.4	53.1	52.0
Male.....	*	3.2	6.4	12.0	23.0	27.8	47.8	55.2	68.8	66.0
Female.....	*	2.3	4.8	8.0	13.3	20.1	25.8	37.9	44.8	46.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All origins 3/, both sexes.....	5.4	9.4	16.5	26.9	42.5	62.5	90.2	122.7	162.9	210.1
Male.....	5.6	10.5	18.5	31.5	51.7	74.1	107.7	150.6	207.3	259.6
Female.....	5.2	8.3	14.5	22.5	34.2	52.4	76.0	102.3	135.1	186.9
Hispanic, both sexes.....	4.2	7.5	11.9	18.6	32.7	42.5	69.8	88.9	110.9	145.5
Male.....	3.5	7.6	11.5	21.5	35.2	50.5	89.6	103.5	130.2	163.5
Female.....	4.8	7.5	12.3	15.8	30.5	35.8	54.3	78.3	98.1	136.0
Mexican, both sexes.....	4.2	7.0	11.1	18.5	32.1	46.3	68.7	97.2	115.4	173.1
Male.....	3.3	6.8	11.1	20.9	30.9	50.6	91.4	115.6	*	*
Female.....	5.3	7.1	11.2	16.0	33.3	42.0	50.2	84.2	99.8	172.5
Puerto Rican, both sexes.....	*	*	14.0	19.5	43.3	47.2	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.4	8.3	12.7	17.4	30.4	31.9	68.5	78.1	99.7	131.1
Male.....	*	8.7	*	21.5	34.9	*	*	*	*	*
Female.....	*	7.9	14.7	14.1	26.8	28.3	61.4	*	*	*
Non-Hispanic 5/, both sexes.....	5.6	9.6	16.9	27.6	43.2	64.0	91.6	124.8	165.7	213.1
Male.....	6.0	10.8	19.3	32.3	52.9	75.8	108.8	153.4	211.6	264.6
Female.....	5.2	8.4	14.7	23.1	34.4	53.7	77.6	103.8	137.1	189.1
Non-Hispanic white 6/, both sexes....	5.5	9.2	16.2	26.8	42.5	63.7	93.1	126.4	169.4	217.1
Male.....	5.9	10.3	18.2	31.4	52.5	76.0	110.8	155.3	216.7	271.8
Female.....	5.0	8.1	14.1	22.4	33.1	52.8	78.4	104.8	139.7	191.5
Non-Hispanic black 6/, both sexes....	7.4	13.4	25.2	39.8	59.0	79.0	98.0	130.3	152.5	188.7
Male.....	7.6	16.2	31.1	48.6	71.5	92.8	118.4	165.3	203.7	216.2
Female.....	7.3	10.9	20.2	32.6	49.3	68.8	84.0	109.4	125.5	177.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All origins 3/, both sexes.....	0.8	1.5	2.2	3.1	5.6	9.0	16.8	28.8	46.3	78.0
Male.....	0.9	1.6	2.5	3.8	6.3	11.1	21.5	38.2	60.8	92.3
Female.....	0.8	1.4	1.9	2.5	4.9	7.2	13.0	21.8	37.3	71.4
Hispanic, both sexes.....	*	1.1	1.7	2.3	5.3	6.8	11.5	19.2	33.7	48.5
Male.....	*	*	2.5	2.7	6.1	7.6	16.0	24.7	39.3	53.0
Female.....	*	*	*	*	4.5	6.2	8.0	15.3	30.0	46.1
Mexican, both sexes.....	*	*	*	*	6.3	7.5	8.3	19.1	33.9	52.2
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.9	1.5	2.3	3.2	5.6	9.2	17.1	29.4	46.9	79.5
Male.....	0.9	1.6	2.5	3.8	6.3	11.4	21.8	39.1	61.9	94.5
Female.....	0.9	1.5	2.0	2.6	5.0	7.3	13.3	22.3	37.6	72.5
Non-Hispanic white 6/, both sexes....	0.8	1.4	2.3	3.2	5.4	9.4	17.7	30.1	48.4	81.6
Male.....	0.9	1.5	2.6	3.9	6.2	11.6	22.7	40.4	63.7	96.0
Female.....	0.7	1.3	1.9	2.6	4.7	7.5	13.5	22.4	38.9	74.9
Non-Hispanic black 6/, both sexes....	1.4	2.4	2.8	3.9	8.1	9.3	15.1	26.7	40.0	63.4
Male.....	*	2.4	2.5	4.9	8.9	12.3	18.7	33.6	55.5	82.5
Female.....	1.7	2.3	3.1	3.1	7.4	7.1	12.6	22.5	31.8	55.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Anemias (D50-D64)

All origins 3/, both sexes.....	0.5	0.5	0.5	0.8	1.2	1.4	3.0	5.1	10.5	30.0
Male.....	0.5	0.5	0.5	0.8	1.1	1.5	2.8	5.4	11.6	29.6
Female.....	0.5	0.6	0.6	0.8	1.2	1.3	3.2	5.0	9.8	30.2
Hispanic, both sexes.....	*	*	*	*	*	*	*	5.8	6.5	21.1
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	23.4
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.5	0.6	0.6	0.8	1.2	1.4	3.1	5.1	10.7	30.3
Male.....	0.5	0.5	0.5	0.8	1.2	1.6	2.8	5.3	12.0	30.2
Female.....	0.5	0.7	0.6	0.8	1.2	1.3	3.3	5.0	9.9	30.4
Non-Hispanic white 6/, both sexes.....	*	0.2	0.3	0.6	0.9	1.3	2.9	4.9	10.6	30.0
Male.....	*	*	*	0.5	0.9	1.5	2.6	5.0	11.9	30.6
Female.....	*	0.3	0.3	0.6	1.0	1.1	3.1	4.8	9.8	29.7
Non-Hispanic black 6/, both sexes....	3.0	2.7	2.5	3.0	3.5	3.1	5.7	8.3	13.6	39.3
Male.....	3.0	2.5	2.2	3.4	4.0	*	*	9.7	15.4	30.2
Female.....	3.0	2.8	2.8	2.6	3.1	*	5.9	7.4	12.6	43.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diabetes mellitus (E10-E14)

All origins 3/, both sexes.....	6.2	9.6	17.1	28.7	46.4	68.3	98.3	141.9	199.3	285.2
Male.....	7.6	12.3	21.3	34.5	56.8	80.7	115.9	167.5	232.2	321.0
Female.....	4.7	6.9	13.2	23.3	36.9	57.5	84.0	123.1	178.7	268.4
Hispanic, both sexes.....	3.8	9.4	18.1	34.5	54.1	96.3	146.1	205.6	264.1	369.7
Male.....	4.7	11.8	22.0	41.8	62.0	110.3	169.7	225.1	309.5	381.2
Female.....	2.8	6.8	14.3	27.6	47.1	84.5	127.7	191.5	233.8	363.7
Mexican, both sexes.....	4.6	10.1	19.5	40.7	58.9	122.7	165.7	244.6	311.7	446.1
Male.....	5.6	12.3	22.0	47.6	64.2	128.0	186.1	272.6	389.8	*
Female.....	3.5	7.7	16.6	33.5	53.9	117.5	149.0	224.9	264.8	449.3
Puerto Rican, both sexes.....	*	13.8	23.2	46.8	81.3	88.0	*	*	*	*
Male.....	*	*	*	54.6	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	109.6	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	5.7	14.3	17.6	37.1	53.4	92.7	150.2	154.4	278.5
Male.....	*	7.8	20.6	19.8	46.2	71.2	*	*	*	*
Female.....	*	*	9.0	15.9	29.6	41.7	79.8	*	*	*
Non-Hispanic 5/, both sexes.....	6.5	9.6	17.0	28.1	45.6	65.9	94.5	137.4	195.2	280.5
Male.....	8.1	12.3	21.1	33.7	56.1	78.1	111.8	163.5	227.4	317.0
Female.....	5.0	6.9	13.0	22.8	35.9	55.2	80.5	118.2	175.1	263.5
Non-Hispanic white 6/, both sexes....	5.6	8.1	14.1	23.8	39.0	56.6	83.6	124.4	180.3	264.2
Male.....	7.0	10.7	17.9	28.9	49.6	69.7	101.6	152.9	217.1	306.8
Female.....	4.1	5.6	10.3	18.8	29.1	44.9	68.6	103.1	157.1	244.3
Non-Hispanic black 6/, both sexes....	13.0	19.6	38.0	63.0	103.6	148.4	197.1	277.5	371.2	475.9
Male.....	16.4	24.3	46.3	73.6	121.9	165.6	225.1	303.8	380.6	468.2
Female.....	9.9	15.5	31.0	54.3	89.3	135.8	177.9	261.7	366.2	479.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nutritional deficiencies (E40-E64)

All origins 3/, both sexes.....	0.1	0.2	0.3	0.5	0.8	1.1	1.9	4.0	7.5	22.7
Male.....	*	*	0.3	0.4	0.7	1.1	2.1	3.9	8.1	18.7
Female.....	*	0.2	0.3	0.5	0.8	1.0	1.8	4.0	7.2	24.5
Hispanic, both sexes.....	*	*	*	*	*	*	*	4.5	7.5	19.2
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	22.8
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	31.3
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.2	0.2	0.3	0.5	0.8	1.1	2.0	3.9	7.5	22.8
Male.....	*	*	0.3	0.5	0.7	1.2	2.1	3.8	8.2	18.9
Female.....	*	0.2	0.3	0.5	0.9	1.1	1.9	4.0	7.1	24.6
Non-Hispanic white 6/, both sexes....	0.2	0.2	0.3	0.4	0.7	1.0	1.9	3.9	7.0	22.2
Male.....	*	*	*	0.3	0.6	1.0	2.0	3.5	7.3	18.4
Female.....	*	*	0.3	0.5	0.9	1.0	1.8	4.1	6.9	24.0
Non-Hispanic black 6/, both sexes.....	*	*	*	*	*	2.7	3.5	5.2	13.8	31.3
Male.....	*	*	*	*	*	*	*	*	19.0	29.3
Female.....	*	*	*	*	*	*	*	*	11.1	32.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malnutrition (E40-E46)

All origins 3/, both sexes.....	0.1	0.2	0.3	0.4	0.7	1.0	1.8	3.7	7.0	20.9
Male.....	*	*	0.2	0.4	0.6	1.0	1.9	3.7	7.4	17.5
Female.....	*	0.2	0.3	0.5	0.7	1.0	1.8	3.8	6.8	22.4
Hispanic, both sexes.....	*	*	*	*	*	*	*	4.3	7.5	17.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	21.0
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	28.7
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.1	0.2	0.3	0.5	0.7	1.1	1.9	3.7	7.0	21.0
Male.....	*	*	0.2	0.4	0.6	1.1	2.0	3.6	7.5	17.9
Female.....	*	0.2	0.3	0.5	0.8	1.1	1.8	3.8	6.7	22.5
Non-Hispanic white 6/, both sexes....	0.2	0.2	0.2	0.4	0.7	0.9	1.8	3.6	6.5	20.3
Male.....	*	*	*	0.3	0.6	0.9	1.8	3.2	6.7	17.2
Female.....	*	*	0.3	0.5	0.8	1.0	1.8	3.9	6.4	21.8
Non-Hispanic black 6/, both sexes.....	*	*	*	*	*	2.7	3.5	4.5	13.3	30.0
Male.....	*	*	*	*	*	*	*	*	17.6	29.3
Female.....	*	*	*	*	*	*	*	*	11.1	30.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Parkinson's disease (G20-G21)

All origins 3/, both sexes.....	*	*	0.3	0.7	2.1	6.0	19.8	50.1	95.4	144.8
Male.....	*	*	0.4	0.9	2.8	8.5	29.0	76.0	145.6	231.7
Female.....	*	*	*	0.4	1.3	3.8	12.2	31.0	64.0	104.1
Hispanic, both sexes.....	*	*	*	*	*	3.9	15.4	30.3	58.6	93.9
Male.....	*	*	*	*	*	*	21.4	47.4	90.1	151.1
Female.....	*	*	*	*	*	*	10.7	17.9	37.6	63.5
Mexican, both sexes.....	*	*	*	*	*	*	12.2	26.7	57.4	93.9
Male.....	*	*	*	*	*	*	*	43.3	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	*	*	0.3	0.7	2.1	6.2	20.1	51.4	97.4	147.2
Male.....	*	*	0.4	0.9	2.9	8.8	29.6	77.9	148.8	235.9
Female.....	*	*	*	0.4	1.4	3.9	12.3	31.9	65.4	106.0
Non-Hispanic white 6/, both sexes.....	*	*	0.3	0.7	2.2	6.5	21.6	55.1	104.3	157.0
Male.....	*	*	0.4	1.0	3.0	9.1	31.4	82.8	158.0	252.4
Female.....	*	*	*	0.5	1.4	4.1	13.4	34.5	70.6	112.5
Non-Hispanic black 6/, both sexes.....	*	*	*	*	1.6	4.2	11.7	25.5	37.8	51.8
Male.....	*	*	*	*	*	6.8	19.0	43.3	63.9	71.5
Female.....	*	*	*	*	*	*	6.7	14.8	24.1	43.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alzheimer's disease (G30)

All origins 3/, both sexes.....	*	0.1	0.4	1.1	3.4	9.6	32.9	101.2	272.6	848.3
Male.....	*	*	0.4	1.0	3.4	10.3	31.6	95.1	244.5	638.6
Female.....	*	*	0.3	1.3	3.3	9.0	34.0	105.7	290.1	946.4
Hispanic, both sexes.....	*	*	*	*	3.4	6.8	19.9	63.1	181.9	514.5
Male.....	*	*	*	*	4.1	8.7	19.1	56.1	188.3	404.8
Female.....	*	*	*	*	*	5.2	20.5	68.2	177.7	572.8
Mexican, both sexes.....	*	*	*	*	*	6.0	16.1	65.7	182.7	519.2
Male.....	*	*	*	*	*	*	*	60.7	199.0	*
Female.....	*	*	*	*	*	*	17.3	69.3	172.9	582.8
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	47.1	125.7	427.3
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	*	0.1	0.4	1.2	3.4	9.9	33.8	103.6	277.5	864.3
Male.....	*	*	0.4	1.0	3.4	10.4	32.4	97.5	247.6	651.1
Female.....	*	*	0.3	1.3	3.4	9.4	34.9	108.1	296.1	963.4
Non-Hispanic white 6/, both sexes.....	*	0.1	0.4	1.3	3.6	10.5	35.2	107.5	288.3	896.5
Male.....	*	*	0.4	1.1	3.5	11.0	33.5	101.1	257.2	676.4
Female.....	*	*	0.3	1.5	3.6	10.1	36.6	112.3	307.9	999.2
Non-Hispanic black 6/, both sexes.....	*	*	*	*	3.1	7.7	33.4	92.4	229.4	674.5
Male.....	*	*	*	*	*	8.5	33.6	90.7	202.3	478.3
Female.....	*	*	*	*	2.8	7.1	33.2	93.5	243.7	754.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

All origins 3/, both sexes.....	46.5	83.2	137.0	205.7	323.5	490.4	798.7	1,368.3	2,468.6	5,928.2
Male.....	62.7	115.5	195.7	291.1	443.8	657.9	1,027.7	1,713.4	2,982.2	6,151.8
Female.....	30.5	51.6	80.8	125.2	213.5	344.0	612.4	1,115.8	2,147.6	5,823.6
Hispanic, both sexes.....	27.7	52.9	93.3	151.9	241.0	383.6	643.8	1,092.3	1,891.9	4,096.9
Male.....	38.1	70.0	134.7	210.2	330.8	511.9	825.5	1,379.3	2,203.6	4,103.7
Female.....	16.3	34.9	52.0	96.9	161.4	275.9	502.4	884.6	1,684.5	4,093.2
Mexican, both sexes.....	29.6	51.8	91.5	157.7	238.7	411.1	609.4	1,066.0	1,870.8	4,287.2
Male.....	39.3	69.4	125.7	208.3	317.6	496.8	757.1	1,380.1	2,370.1	*
Female.....	18.3	32.8	53.7	105.2	165.4	326.4	489.0	844.4	1,570.9	4,328.5
Puerto Rican, both sexes.....	39.6	76.3	127.2	202.6	357.4	442.7	1,130.6	*	*	*
Male.....	59.1	99.4	187.2	255.7	*	*	*	*	*	*
Female.....	22.6	58.0	79.4	147.4	221.6	293.6	*	*	*	*
Cuban, both sexes.....	21.4	72.4	122.0	156.3	254.5	480.7	675.7	997.5	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	21.0	43.8	79.9	117.0	189.5	272.6	527.2	1,001.6	1,489.3	3,408.1
Male.....	30.1	57.2	128.7	171.3	268.7	408.1	647.8	*	*	*
Female.....	11.5	30.3	38.8	73.5	125.5	184.3	433.6	869.2	*	3,167.0
Non-Hispanic 5/, both sexes.....	49.4	86.8	141.2	209.8	329.4	497.4	808.3	1,384.1	2,497.2	6,013.2
Male.....	66.7	121.1	201.5	297.1	451.7	666.9	1,039.3	1,731.8	3,023.0	6,255.9
Female.....	32.5	53.6	83.6	127.5	217.3	348.7	619.7	1,129.4	2,169.8	5,900.4
Non-Hispanic white 6/, both sexes....	43.4	75.9	123.3	187.3	301.9	461.8	771.3	1,354.1	2,484.7	6,119.6
Male.....	60.7	109.0	180.0	269.0	417.5	624.6	996.4	1,701.5	3,029.3	6,398.5
Female.....	26.1	43.2	67.7	108.5	193.6	316.0	583.4	1,094.3	2,141.9	5,989.5
Non-Hispanic black 6/, both sexes....	93.2	170.1	291.6	420.6	625.5	873.1	1,274.7	1,921.7	3,021.0	5,656.4
Male.....	115.5	222.0	400.0	582.9	867.2	1,177.5	1,669.0	2,419.2	3,552.0	5,493.9
Female.....	73.6	125.0	200.1	287.1	437.6	648.7	1,004.2	1,624.6	2,741.0	5,722.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All origins 3/, both sexes.....	37.3	67.4	111.0	167.7	261.3	386.9	615.9	1,031.7	1,841.0	4,480.8
Male.....	52.4	97.4	164.1	244.4	369.3	534.4	819.7	1,341.1	2,319.5	4,819.9
Female.....	22.4	38.1	60.0	95.4	162.6	258.0	450.1	805.2	1,541.9	4,322.1
Hispanic, both sexes.....	19.1	38.4	69.1	113.0	184.3	290.2	490.1	832.8	1,416.2	3,140.6
Male.....	28.0	53.8	104.7	163.8	259.7	401.8	645.8	1,056.9	1,708.2	3,257.9
Female.....	9.4	22.2	33.7	65.0	117.5	196.5	368.8	670.7	1,221.8	3,078.3
Mexican, both sexes.....	19.6	36.6	66.3	114.3	175.5	299.7	451.5	789.5	1,357.9	3,148.9
Male.....	28.0	51.9	96.0	157.0	243.5	375.4	582.9	1,029.5	1,776.3	*
Female.....	9.7	20.0	33.4	70.0	112.1	224.9	344.4	620.2	1,106.6	3,065.6
Puerto Rican, both sexes.....	30.1	60.4	96.5	155.8	299.4	355.6	885.3	*	*	*
Male.....	44.5	79.5	141.0	206.9	*	*	*	*	*	*
Female.....	*	45.2	61.0	102.7	182.0	230.5	*	*	*	*
Cuban, both sexes.....	19.7	60.8	94.3	126.8	198.7	397.3	563.1	813.9	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	14.5	31.2	61.5	89.1	148.6	207.3	399.2	771.2	1,060.2	2,571.2
Male.....	22.4	44.3	105.9	140.2	212.9	333.3	500.5	*	*	*
Female.....	6.4	17.9	24.1	48.1	96.6	125.2	320.6	668.3	*	2,340.8
Non-Hispanic 5/, both sexes.....	40.1	70.9	115.0	172.0	266.9	393.4	623.6	1,042.7	1,862.1	4,542.5
Male.....	56.4	102.8	170.0	250.6	377.2	542.8	829.7	1,357.0	2,351.5	4,899.0
Female.....	24.2	40.0	62.5	97.9	166.0	262.5	455.4	812.6	1,557.3	4,376.7
Non-Hispanic white 6/, both sexes....	36.8	63.8	103.7	157.5	249.2	370.9	600.8	1,027.0	1,862.2	4,642.7
Male.....	53.2	94.8	156.5	232.8	354.7	515.9	804.1	1,341.0	2,368.5	5,032.8
Female.....	20.4	33.2	52.0	84.8	150.5	241.0	431.3	792.2	1,543.5	4,460.8
Non-Hispanic black 6/, both sexes....	68.5	129.3	218.6	320.2	475.6	655.4	950.3	1,403.3	2,189.6	4,103.9
Male.....	88.2	175.8	306.5	448.6	675.4	903.7	1,271.0	1,836.4	2,664.4	4,129.6
Female.....	51.1	88.9	144.3	214.7	320.2	472.3	730.4	1,144.6	1,939.3	4,093.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive heart disease (I11)

All origins 3/, both sexes.....	4.0	6.8	10.0	11.8	14.9	18.2	23.1	35.2	64.2	194.0
Male.....	5.2	9.8	13.8	16.2	20.4	24.0	27.8	40.2	63.1	143.0
Female.....	2.8	3.8	6.3	7.5	9.8	13.2	19.2	31.4	64.9	217.9
Hispanic, both sexes.....	2.5	3.8	6.4	8.1	11.6	17.3	19.9	35.0	55.3	157.7
Male.....	3.8	5.7	8.9	11.4	15.6	23.1	24.8	37.6	55.7	125.2
Female.....	*	1.7	3.9	4.9	8.2	12.3	16.0	33.2	55.0	174.9
Mexican, both sexes.....	2.4	2.9	5.9	7.4	8.1	14.8	15.5	30.7	48.8	159.1
Male.....	3.6	4.3	7.9	9.9	11.2	16.2	20.5	*	*	*
Female.....	*	*	3.7	*	*	13.5	*	28.5	47.4	180.8
Puerto Rican, both sexes.....	*	*	*	*	28.6	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	3.5	6.5	7.0	9.8	*	20.8	38.1	*	142.9
Male.....	*	*	11.2	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.2	7.1	10.3	12.0	15.1	18.2	23.2	35.1	64.5	195.6
Male.....	5.5	10.3	14.3	16.5	20.7	23.9	28.0	40.3	63.2	143.8
Female.....	3.0	4.1	6.5	7.7	9.9	13.2	19.4	31.2	65.3	219.6
Non-Hispanic white 6/, both sexes....	3.1	5.2	7.4	9.0	11.4	13.8	17.6	29.3	57.9	188.0
Male.....	4.3	7.9	11.0	12.8	15.7	17.7	21.3	32.9	55.7	134.7
Female.....	1.9	2.5	3.9	5.3	7.3	10.3	14.5	26.7	59.4	212.9
Non-Hispanic black 6/, both sexes....	11.6	20.7	32.1	37.9	49.8	57.9	76.5	98.3	148.1	304.2
Male.....	13.2	28.2	41.5	51.0	72.8	85.3	96.6	133.0	168.5	278.5
Female.....	10.2	14.2	24.1	27.1	31.9	37.8	62.8	77.5	137.4	314.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive heart and renal disease (I13)

All origins 3/, both sexes.....	0.2	0.3	0.5	0.7	1.0	1.4	2.4	4.3	8.6	22.6
Male.....	0.3	0.3	0.6	0.9	1.3	1.6	2.6	4.7	10.0	23.8
Female.....	*	0.3	0.4	0.4	0.8	1.3	2.3	4.0	7.8	22.1
Hispanic, both sexes.....	*	*	*	*	*	*	*	4.5	*	12.9
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	12.6
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.2	0.3	0.5	0.6	1.0	1.4	2.4	4.3	8.8	23.0
Male.....	0.3	0.4	0.7	0.9	1.3	1.5	2.6	4.7	10.2	24.3
Female.....	*	0.3	0.4	0.4	0.8	1.3	2.3	4.0	8.0	22.5
Non-Hispanic white 6/, both sexes.....	*	0.1	0.2	0.3	0.5	0.7	1.5	3.0	7.8	21.3
Male.....	*	*	*	0.4	0.6	0.7	1.5	3.1	9.2	22.9
Female.....	*	*	*	*	0.4	0.8	1.4	2.9	6.9	20.5
Non-Hispanic black 6/, both sexes....	1.0	1.7	2.8	3.5	5.9	6.7	11.2	16.9	21.6	42.5
Male.....	1.5	2.0	3.6	5.2	7.8	8.5	13.6	23.0	23.9	40.3
Female.....	*	1.4	2.0	2.1	4.4	5.3	9.6	13.2	20.4	43.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Ischemic heart diseases (I20-I25)

All origins 3/, both sexes.....	21.5	42.2	75.0	118.1	188.5	279.1	441.7	726.4	1,263.9	2,904.7
Male.....	32.7	64.4	116.9	180.0	277.1	400.1	613.0	988.2	1,672.7	3,303.0
Female.....	10.3	20.6	34.8	59.7	107.5	173.3	302.4	534.8	1,008.4	2,718.3
Hispanic, both sexes.....	10.3	24.0	47.8	84.0	139.6	220.8	371.8	642.6	1,071.6	2,296.7
Male.....	15.6	35.2	76.1	125.7	202.6	315.2	500.9	849.8	1,297.9	2,453.9
Female.....	4.4	12.2	19.6	44.6	83.8	141.6	271.4	492.6	921.0	2,213.2
Mexican, both sexes.....	10.2	22.2	45.7	83.6	135.9	228.8	337.9	601.1	1,019.1	2,230.6
Male.....	15.5	33.1	69.2	118.1	195.1	299.5	437.2	827.3	1,347.1	*
Female.....	4.1	10.5	19.6	47.8	80.7	158.9	256.9	441.6	822.0	2,096.1
Puerto Rican, both sexes.....	15.8	39.8	66.3	119.6	215.5	277.8	684.3	*	*	*
Male.....	*	55.8	104.5	167.4	*	*	*	*	*	*
Female.....	*	27.1	35.8	70.1	120.3	186.3	*	*	*	*
Cuban, both sexes.....	*	42.5	72.5	97.3	149.3	291.0	442.7	640.3	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.8	19.4	41.9	66.9	111.2	155.9	302.3	576.9	788.3	1,856.5
Male.....	11.5	29.1	76.4	108.5	161.4	260.8	386.5	*	*	*
Female.....	*	9.7	12.8	33.6	70.5	87.4	237.1	478.2	*	*
Non-Hispanic 5/, both sexes.....	23.2	44.4	77.7	120.7	191.9	282.8	445.6	730.5	1,272.7	2,931.2
Male.....	35.6	68.0	120.9	184.1	282.2	405.2	619.0	995.2	1,692.0	3,344.5
Female.....	11.1	21.6	36.3	60.9	109.3	175.5	304.0	536.6	1,011.5	2,739.1
Non-Hispanic white 6/, both sexes....	22.9	42.3	73.6	114.3	183.6	271.4	434.8	723.3	1,275.7	2,992.4
Male.....	35.9	65.9	116.4	176.2	271.8	392.4	608.2	990.1	1,709.8	3,441.2
Female.....	10.0	18.8	31.7	54.5	101.0	163.0	290.1	523.8	1,002.4	2,783.1
Non-Hispanic black 6/, both sexes....	30.6	66.4	124.2	195.6	303.0	429.1	626.5	935.1	1,436.6	2,640.8
Male.....	42.4	93.6	182.4	286.2	441.3	604.5	864.9	1,257.7	1,811.1	2,685.6
Female.....	20.2	42.8	75.1	121.1	195.4	299.9	463.0	742.5	1,239.0	2,622.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute myocardial infarction (I21-I22)

All origins 3/, both sexes.....	8.5	16.5	29.2	46.7	73.9	107.0	160.4	252.0	406.4	825.8
Male.....	13.0	24.7	44.8	70.0	107.3	151.8	219.9	331.3	527.1	936.3
Female.....	4.1	8.4	14.3	24.7	43.4	67.8	112.0	194.0	330.9	774.1
Hispanic, both sexes.....	4.0	8.4	15.6	30.3	48.4	80.1	127.1	225.5	346.5	692.9
Male.....	5.7	11.7	23.5	42.9	67.3	112.0	172.4	293.1	411.1	742.0
Female.....	2.1	5.0	7.7	18.4	31.6	53.3	91.8	176.6	303.6	666.8
Mexican, both sexes.....	4.1	8.1	16.6	31.5	51.1	87.2	126.1	225.9	357.4	724.4
Male.....	5.9	11.5	23.3	42.6	74.4	112.3	162.3	306.3	458.9	*
Female.....	*	4.4	9.3	20.1	29.4	62.5	96.6	169.2	296.4	698.2
Puerto Rican, both sexes.....	*	13.0	14.0	36.7	64.0	81.5	*	*	*	*
Male.....	*	*	*	54.6	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	156.4	219.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.6	6.9	12.9	22.4	32.8	53.4	92.7	187.3	228.2	495.1
Male.....	*	9.9	23.2	30.5	37.5	79.7	*	*	*	*
Female.....	*	*	*	15.9	28.9	36.2	78.6	*	*	*
Non-Hispanic 5/, both sexes.....	9.2	17.5	30.6	48.0	75.8	108.8	162.5	253.3	409.1	831.0
Male.....	14.2	26.4	46.9	72.2	110.2	154.3	222.7	333.3	533.2	945.2
Female.....	4.4	8.9	15.0	25.2	44.3	68.9	113.4	194.6	331.9	778.0
Non-Hispanic white 6/, both sexes....	9.3	17.2	30.1	46.9	74.8	107.0	160.7	251.4	410.4	841.4
Male.....	14.5	26.4	46.9	71.4	109.3	153.8	222.5	333.6	538.4	964.6
Female.....	4.2	8.0	13.6	23.3	42.5	65.0	109.1	190.0	329.8	783.9
Non-Hispanic black 6/, both sexes....	11.5	23.8	42.7	68.5	103.3	147.3	211.4	316.7	455.1	811.7
Male.....	16.4	32.4	60.4	97.2	148.2	198.9	277.6	397.7	563.9	824.6
Female.....	7.1	16.3	27.8	45.0	68.4	109.2	165.9	268.3	397.7	806.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute ischemic heart diseases (I24)

All origins 3/, both sexes.....	0.3	0.6	1.1	1.5	2.5	2.9	4.4	6.4	10.0	21.5
Male.....	0.4	0.8	1.6	1.9	3.6	3.6	5.9	8.0	12.1	21.8
Female.....	0.2	0.3	0.6	1.0	1.5	2.2	3.3	5.3	8.8	21.4
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	8.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.3	0.6	1.2	1.5	2.6	3.0	4.6	6.6	10.3	22.1
Male.....	0.4	0.9	1.8	2.1	3.8	3.8	6.1	8.3	12.4	22.6
Female.....	*	0.4	0.6	1.0	1.6	2.3	3.4	5.4	9.1	21.9
Non-Hispanic white 6/, both sexes....	0.3	0.5	1.0	1.4	2.5	2.9	4.3	6.5	10.1	22.3
Male.....	0.4	0.8	1.6	1.9	3.6	3.7	5.7	8.0	11.9	22.7
Female.....	*	0.3	0.5	1.0	1.4	2.1	3.2	5.4	9.0	22.2
Non-Hispanic black 6/, both sexes.....	*	1.2	2.4	2.7	4.6	4.8	7.7	9.0	14.5	21.0
Male.....	*	*	3.3	4.0	7.1	5.8	11.1	12.9	19.7	23.8
Female.....	*	*	*	*	*	4.1	5.4	6.6	11.8	19.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other forms of chronic ischemic heart disease (I20,I25)

All origins 3/, both sexes.....	12.7	25.2	44.7	70.0	112.1	169.3	276.9	468.0	847.5	2,057.4
Male.....	19.4	38.8	70.5	108.1	166.2	244.7	387.2	649.0	1,133.5	2,344.9
Female.....	6.0	11.8	19.9	34.1	62.7	103.3	187.1	335.6	668.6	1,922.9
Hispanic, both sexes.....	6.2	15.5	32.0	53.2	90.7	139.6	242.9	414.2	720.2	1,595.2
Male.....	9.9	23.4	52.3	82.3	134.5	201.8	325.8	555.2	880.3	1,705.1
Female.....	2.2	7.2	11.8	25.7	51.9	87.4	178.5	312.3	613.6	1,536.8
Mexican, both sexes.....	6.0	14.0	28.8	51.6	84.1	140.5	209.7	371.3	657.4	1,494.9
Male.....	9.5	21.3	45.6	75.0	119.7	186.2	271.6	518.1	883.2	*
Female.....	*	6.0	10.2	27.3	51.0	95.5	159.2	267.7	521.7	1,384.0
Puerto Rican, both sexes.....	10.2	26.8	51.7	82.3	150.6	194.5	492.5	*	*	*
Male.....	*	39.7	82.6	112.7	*	*	*	*	*	*
Female.....	*	*	*	*	*	137.3	*	*	*	*
Cuban, both sexes.....	*	*	*	*	92.2	166.8	284.7	419.8	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.2	12.5	29.0	44.0	78.4	101.5	208.2	389.6	552.0	1,354.1
Male.....	8.2	19.2	53.2	77.4	123.9	178.7	274.0	*	*	*
Female.....	*	*	*	17.2	41.6	51.2	157.2	332.4	*	*
Non-Hispanic 5/, both sexes.....	13.7	26.3	45.9	71.2	113.5	171.0	278.5	470.6	853.2	2,078.0
Male.....	21.0	40.7	72.2	109.8	168.2	247.1	390.2	653.5	1,146.4	2,376.7
Female.....	6.5	12.4	20.7	34.7	63.4	104.3	187.3	336.6	670.6	1,939.2
Non-Hispanic white 6/, both sexes....	13.3	24.6	42.5	65.9	106.3	161.6	269.7	465.4	855.2	2,128.7
Male.....	21.0	38.7	67.9	102.9	158.9	235.0	379.9	648.5	1,159.5	2,453.9
Female.....	5.7	10.6	17.6	30.3	57.1	95.9	177.8	328.4	663.6	1,977.0
Non-Hispanic black 6/, both sexes....	18.6	41.5	79.1	124.4	195.0	277.0	407.4	609.5	966.9	1,808.1
Male.....	25.3	59.7	118.6	184.9	286.0	399.8	576.2	847.1	1,227.6	1,837.1
Female.....	12.8	25.6	45.8	74.5	124.3	186.5	291.7	467.6	829.5	1,796.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All origins 3/, both sexes.....	4.6	9.4	17.9	26.1	38.5	47.4	64.4	89.1	147.1	345.5
Male.....	7.0	14.9	29.3	41.5	58.9	69.9	90.6	117.9	188.1	365.3
Female.....	2.2	4.1	7.1	11.5	20.0	27.7	43.1	67.9	121.4	336.1
Hispanic, both sexes.....	2.5	6.1	13.0	23.0	35.8	45.2	65.3	81.7	124.7	260.2
Male.....	3.8	9.2	23.5	39.2	57.1	70.3	95.7	105.1	161.3	277.4
Female.....	*	2.9	2.5	7.7	16.9	24.2	41.6	64.8	100.3	251.0
Mexican, both sexes.....	2.3	5.1	12.1	22.6	31.1	46.0	60.4	76.1	140.7	300.0
Male.....	3.8	7.9	20.6	37.8	48.8	65.8	88.8	100.2	207.2	*
Female.....	*	*	*	6.9	14.7	26.5	37.2	59.1	100.8	278.2
Puerto Rican, both sexes.....	*	13.0	19.4	32.0	56.2	64.8	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	5.3	12.9	21.1	38.6	34.8	67.8	94.1	97.0	247.5
Male.....	*	7.8	27.3	39.0	65.4	62.8	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.9	9.8	18.4	26.1	38.5	47.2	63.9	89.1	147.9	349.3
Male.....	7.6	15.6	29.7	41.4	58.6	69.3	89.4	118.0	188.9	369.6
Female.....	2.3	4.2	7.5	11.8	20.2	27.8	43.0	67.9	122.3	339.9
Non-Hispanic white 6/, both sexes....	4.6	8.9	16.4	23.2	34.4	41.9	57.6	82.7	141.3	346.8
Male.....	7.2	14.4	26.9	37.0	52.2	61.4	80.3	109.5	180.6	366.7
Female.....	1.9	3.5	6.1	9.8	17.8	24.5	38.7	62.6	116.5	337.5
Non-Hispanic black 6/, both sexes....	7.9	17.0	35.6	53.0	79.7	98.6	128.0	168.3	247.5	433.7
Male.....	10.9	25.2	55.4	83.0	128.3	154.0	193.1	238.9	332.2	471.0
Female.....	5.3	9.8	18.8	28.3	41.9	57.7	83.4	126.2	202.9	418.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All origins 3/, both sexes.....	8.1	15.7	26.8	43.9	73.6	121.9	212.5	379.0	700.4	1,712.0
Male.....	12.3	23.9	41.2	66.6	107.4	174.8	296.7	531.1	945.4	1,979.5
Female.....	3.8	7.7	12.9	22.6	42.7	75.6	144.0	267.7	547.2	1,586.8
Hispanic, both sexes.....	3.8	9.4	19.0	30.2	54.9	94.4	177.7	332.5	595.5	1,335.0
Male.....	6.0	14.2	28.7	43.1	77.4	131.5	230.0	450.1	719.0	1,427.7
Female.....	*	4.3	9.3	18.0	35.0	63.2	136.9	247.4	513.4	1,285.7
Mexican, both sexes.....	3.7	8.9	16.6	29.0	53.0	94.5	149.3	295.2	516.6	1,194.9
Male.....	5.7	13.4	25.0	37.3	70.9	120.4	182.8	418.0	676.0	*
Female.....	*	4.0	7.4	20.3	36.3	69.0	122.0	208.6	420.9	1,105.8
Puerto Rican, both sexes.....	*	13.8	32.3	50.3	94.3	129.6	385.5	*	*	*
Male.....	*	*	*	66.2	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	234.6	360.0	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.8	7.2	16.2	22.9	39.8	66.7	140.4	295.5	455.0	1,106.6
Male.....	*	11.4	25.9	38.4	58.5	115.9	183.7	*	*	*
Female.....	*	*	*	*	24.7	34.6	106.9	252.4	*	*
Non-Hispanic 5/, both sexes.....	8.7	16.5	27.5	45.0	75.0	123.8	214.6	381.5	705.4	1,728.7
Male.....	13.4	25.1	42.5	68.5	109.6	177.8	300.8	535.6	957.6	2,007.1
Female.....	4.2	8.2	13.2	22.9	43.2	76.5	144.2	268.7	548.3	1,599.3
Non-Hispanic white 6/, both sexes....	8.7	15.7	26.2	42.7	71.9	119.7	212.2	382.7	713.9	1,781.9
Male.....	13.7	24.3	41.0	65.9	106.7	173.5	299.7	539.0	978.9	2,087.2
Female.....	3.8	7.1	11.6	20.4	39.3	71.4	139.1	265.8	547.1	1,639.5
Non-Hispanic black 6/, both sexes....	10.7	24.5	43.5	71.4	115.4	178.5	279.4	441.1	719.4	1,374.4
Male.....	14.5	34.5	63.2	101.9	157.7	245.8	383.0	608.1	895.4	1,366.1
Female.....	7.5	15.8	27.0	46.2	82.5	128.8	208.3	341.4	626.6	1,377.8

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other heart diseases (I26-I51)

All origins 3/, both sexes.....	11.4	17.7	25.0	36.3	55.5	85.7	144.6	259.0	493.3	1,342.8
Male.....	14.0	22.6	32.4	46.6	69.5	106.9	173.2	302.8	566.0	1,338.1
Female.....	8.9	12.9	18.0	26.6	42.7	67.2	121.3	226.9	447.9	1,345.0
Hispanic, both sexes.....	6.0	10.0	14.3	19.3	30.5	48.5	92.0	144.6	276.2	663.2
Male.....	8.0	12.2	19.0	25.0	38.9	59.3	114.8	162.2	341.5	658.6
Female.....	3.8	7.7	9.6	14.0	23.0	39.5	74.2	131.9	232.7	665.6
Mexican, both sexes.....	6.7	10.9	14.1	21.5	29.3	50.8	91.3	143.4	274.7	734.0
Male.....	8.6	13.9	18.2	26.9	35.1	53.6	120.6	158.9	361.8	*
Female.....	4.5	7.7	9.5	15.8	23.9	48.0	67.5	132.5	222.3	760.8
Puerto Rican, both sexes.....	9.4	*	21.0	23.7	50.2	45.4	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	139.5	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.2	7.5	12.5	13.6	25.0	39.6	68.5	147.2	209.0	551.1
Male.....	6.4	8.4	17.4	18.7	33.2	55.5	*	*	*	*
Female.....	*	*	*	*	*	29.1	60.2	*	*	*
Non-Hispanic 5/, both sexes.....	12.2	18.7	26.1	37.7	57.5	88.4	148.2	266.1	505.0	1,375.6
Male.....	14.9	23.9	33.7	48.3	71.9	110.2	177.0	311.5	578.4	1,374.1
Female.....	9.6	13.6	18.8	27.7	44.3	69.4	124.7	232.9	459.2	1,376.4
Non-Hispanic white 6/, both sexes....	10.5	15.9	22.1	33.1	52.6	82.4	142.7	264.3	509.0	1,422.6
Male.....	12.8	20.6	28.6	42.8	65.6	103.2	169.9	309.4	585.4	1,420.8
Female.....	8.3	11.4	15.8	23.7	40.3	63.8	120.1	230.6	460.8	1,423.5
Non-Hispanic black 6/, both sexes....	24.8	39.6	58.8	82.0	114.3	158.8	232.9	348.4	577.3	1,111.9
Male.....	30.9	51.3	78.4	105.0	151.1	203.0	293.1	418.0	658.7	1,122.4
Female.....	19.4	29.6	42.3	63.1	85.7	126.1	191.5	306.8	534.4	1,107.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Heart failure (I50)

All origins 3/, both sexes.....	1.0	2.0	3.7	6.6	12.0	21.9	43.1	88.7	190.0	614.4
Male.....	1.2	2.6	4.9	8.2	14.5	26.6	50.0	103.3	213.5	599.0
Female.....	0.8	1.5	2.6	5.1	9.7	17.9	37.6	78.0	175.2	621.6
Hispanic, both sexes.....	*	1.0	2.1	3.2	6.3	12.6	26.9	48.6	113.5	304.8
Male.....	*	*	3.0	3.7	6.8	14.4	34.3	56.7	142.5	276.3
Female.....	*	*	*	2.7	5.8	11.1	21.1	42.9	94.3	319.9
Mexican, both sexes.....	*	1.4	2.3	4.0	6.6	15.6	31.2	54.6	133.9	382.6
Male.....	*	*	3.3	5.1	*	16.2	41.7	64.5	184.2	*
Female.....	*	*	*	*	*	15.0	22.7	47.6	103.7	429.8
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	50.1	80.6	274.1
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.1	2.2	3.9	6.9	12.4	22.6	44.3	91.2	194.0	629.3
Male.....	1.3	2.8	5.0	8.6	15.1	27.4	51.0	106.2	217.3	616.3
Female.....	0.8	1.6	2.8	5.3	10.0	18.4	38.7	80.2	179.6	635.3
Non-Hispanic white 6/, both sexes....	0.8	1.5	2.9	5.8	11.4	21.1	43.3	91.6	197.7	653.7
Male.....	0.9	2.0	3.7	7.6	13.9	26.0	49.3	107.3	222.0	641.8
Female.....	0.7	1.0	2.1	4.2	9.1	16.7	38.4	79.9	182.4	659.3
Non-Hispanic black 6/, both sexes....	2.9	6.6	11.1	16.6	25.1	41.9	65.7	111.4	204.1	492.6
Male.....	3.9	8.3	15.5	18.8	32.3	50.3	83.3	128.4	229.6	473.7
Female.....	2.0	5.2	7.4	14.7	19.4	35.8	53.6	101.2	190.7	500.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All origins 3/, both sexes.....	10.0	15.0	20.5	28.7	42.2	62.4	99.2	167.0	299.4	724.4
Male.....	12.3	19.1	26.5	37.0	53.5	78.5	120.7	196.1	347.5	735.0
Female.....	7.8	11.1	14.8	20.9	31.9	48.3	81.7	145.7	269.3	719.5
Hispanic, both sexes.....	5.3	8.5	11.5	15.4	23.3	35.0	63.4	93.2	161.0	354.5
Male.....	7.1	10.1	15.0	20.3	30.9	43.2	78.2	103.0	198.2	378.9
Female.....	3.3	6.9	7.9	10.8	16.5	28.2	52.0	86.1	136.2	341.5
Mexican, both sexes.....	5.8	9.0	11.0	16.9	22.3	34.9	59.5	85.2	138.3	347.8
Male.....	7.6	11.2	13.8	20.9	28.8	36.9	78.2	91.5	176.0	*
Female.....	3.7	6.6	7.9	12.7	16.3	33.0	44.3	80.8	115.6	328.3
Puerto Rican, both sexes.....	*	*	18.9	16.6	35.5	38.9	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	115.3	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.6	6.8	10.0	11.6	19.1	28.1	48.4	96.2	128.4	275.5
Male.....	5.7	7.5	13.9	15.8	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	10.7	15.8	21.4	29.8	43.8	64.4	101.7	171.6	306.8	742.4
Male.....	13.1	20.2	27.6	38.4	55.3	81.0	123.5	201.8	355.8	753.7
Female.....	8.4	11.6	15.5	21.7	33.2	49.8	83.9	149.4	276.3	737.2
Non-Hispanic white 6/, both sexes....	9.3	13.8	18.6	26.4	40.0	59.8	97.2	169.5	307.0	765.0
Male.....	11.4	17.7	24.0	34.1	50.3	75.4	118.0	198.6	358.0	774.7
Female.....	7.2	9.9	13.2	19.0	30.3	45.9	79.8	147.7	274.8	760.4
Non-Hispanic black 6/, both sexes....	21.3	31.8	45.9	63.0	86.8	115.0	164.4	231.5	369.7	614.7
Male.....	26.5	41.3	60.2	82.7	116.2	149.4	208.0	285.4	423.5	647.8
Female.....	16.8	23.5	33.8	46.7	63.9	89.6	134.4	199.3	341.4	601.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

All origins 3/, both sexes.....	1.2	2.1	4.0	5.5	8.8	13.3	21.0	36.6	69.7	189.4
Male.....	1.5	2.6	5.1	6.7	10.6	15.1	22.8	36.6	72.2	159.2
Female.....	0.9	1.6	2.8	4.4	7.1	11.6	19.6	36.7	68.1	203.5
Hispanic, both sexes.....	0.7	1.6	3.4	6.1	8.6	13.0	22.0	27.7	58.9	147.1
Male.....	*	1.7	4.2	6.6	11.1	16.1	22.9	30.4	52.4	131.9
Female.....	*	*	2.5	5.7	6.3	10.4	21.4	25.7	63.2	155.2
Mexican, both sexes.....	*	1.6	3.7	6.1	8.5	16.8	22.0	29.5	73.5	197.4
Male.....	*	*	3.6	6.9	10.2	17.7	22.5	*	*	*
Female.....	*	*	3.7	5.3	*	16.0	21.6	26.5	76.1	219.8
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	107.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.3	2.2	4.0	5.4	8.8	13.3	20.9	37.2	70.2	191.2
Male.....	1.6	2.8	5.2	6.7	10.6	15.1	22.7	36.9	73.2	160.5
Female.....	1.0	1.7	2.9	4.2	7.1	11.7	19.5	37.4	68.3	205.5
Non-Hispanic white 6/, both sexes....	0.6	1.3	2.5	3.7	6.4	9.8	16.9	31.8	63.6	180.8
Male.....	0.8	1.8	3.3	4.5	8.3	11.3	17.8	31.9	65.7	152.0
Female.....	0.4	0.9	1.6	2.8	4.7	8.5	16.2	31.7	62.2	194.2
Non-Hispanic black 6/, both sexes....	5.3	7.9	15.3	19.5	29.2	43.6	58.4	95.7	151.5	322.8
Male.....	6.7	9.5	19.9	25.4	33.8	50.5	73.8	99.4	174.2	274.9
Female.....	4.0	6.5	11.4	14.6	25.7	38.5	47.8	93.5	139.6	342.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cerebrovascular diseases (I60-I69)

All origins 3/, both sexes.....	6.7	11.2	18.5	26.4	42.7	70.7	127.5	240.3	458.6	1,039.6
Male.....	6.9	12.0	21.2	31.2	49.2	81.9	140.9	259.6	468.5	932.4
Female.....	6.5	10.4	15.8	21.9	36.8	60.9	116.5	226.2	452.3	1,089.8
Hispanic, both sexes.....	6.7	11.5	18.2	29.2	41.4	67.2	113.4	196.5	361.2	687.0
Male.....	7.6	12.8	22.6	34.6	51.0	76.2	132.4	244.6	369.3	581.9
Female.....	5.6	10.2	13.8	24.1	33.0	59.7	98.6	161.7	355.9	742.9
Mexican, both sexes.....	7.7	12.1	19.3	33.5	48.4	81.5	120.7	214.7	380.8	795.7
Male.....	8.7	14.0	23.0	39.8	57.5	88.5	131.2	272.6	442.4	*
Female.....	6.6	10.1	15.2	27.0	39.9	74.5	112.3	173.9	343.8	899.9
Puerto Rican, both sexes.....	*	12.6	18.9	31.4	37.2	69.5	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	133.8	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.2	10.2	14.9	19.6	29.3	43.8	87.9	174.3	348.4	613.0
Male.....	5.9	10.2	17.9	20.9	35.8	*	*	*	*	*
Female.....	*	10.3	12.4	18.6	24.0	43.3	72.5	*	*	*
Non-Hispanic 5/, both sexes.....	6.7	11.1	18.4	26.1	42.7	70.7	128.3	242.9	463.3	1,056.5
Male.....	6.7	11.9	21.0	30.8	48.9	82.1	141.3	260.0	473.7	950.4
Female.....	6.6	10.4	16.0	21.7	36.9	60.7	117.8	230.3	456.8	1,105.8
Non-Hispanic white 6/, both sexes....	4.9	8.5	14.0	20.4	35.5	61.6	117.7	233.4	457.3	1,068.3
Male.....	5.1	9.0	15.4	23.3	39.8	71.0	128.2	249.2	469.2	960.2
Female.....	4.8	7.9	12.5	17.5	31.5	53.2	109.0	221.5	449.8	1,118.7
Non-Hispanic black 6/, both sexes....	16.4	28.0	49.7	69.2	103.7	145.9	225.8	355.9	564.9	1,023.0
Male.....	16.1	29.8	62.5	91.0	135.6	182.7	275.1	407.0	589.9	894.3
Female.....	16.7	26.4	38.8	51.2	78.9	118.8	192.0	325.5	551.8	1,075.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerosis (I70)

All origins 3/, both sexes.....	0.1	0.2	0.4	0.9	1.7	3.2	6.1	12.6	27.2	89.9
Male.....	*	0.3	0.6	1.2	2.2	3.8	7.3	14.2	29.5	85.3
Female.....	*	*	0.2	0.5	1.3	2.6	5.1	11.4	25.7	92.1
Hispanic, both sexes.....	*	*	*	*	*	*	*	8.0	16.7	50.5
Male.....	*	*	*	*	*	*	*	*	18.8	45.1
Female.....	*	*	*	*	*	*	*	7.8	15.3	53.3
Mexican, both sexes.....	*	*	*	*	*	*	*	*	19.8	64.4
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.1	0.2	0.4	0.9	1.8	3.2	6.3	12.9	27.7	91.8
Male.....	*	0.4	0.6	1.2	2.3	3.9	7.5	14.5	30.0	87.5
Female.....	*	*	0.3	0.6	1.4	2.7	5.4	11.6	26.3	93.9
Non-Hispanic white 6/, both sexes.....	*	0.2	0.4	0.8	1.7	3.2	6.3	13.0	28.0	95.2
Male.....	*	0.3	0.6	1.1	2.2	3.7	7.3	14.6	29.8	90.3
Female.....	*	*	*	0.6	1.2	2.7	5.4	11.8	26.8	97.6
Non-Hispanic black 6/, both sexes.....	*	*	*	1.4	3.2	4.6	9.0	14.6	29.8	72.7
Male.....	*	*	*	2.6	3.8	6.8	11.7	17.0	40.0	76.0
Female.....	*	*	*	*	*	*	7.2	13.2	24.4	71.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

All origins 3/, both sexes.....	1.2	2.2	3.2	5.3	9.0	16.4	28.2	47.1	72.2	128.5
Male.....	1.8	3.1	4.5	7.6	12.4	22.7	37.0	61.9	92.5	155.0
Female.....	0.7	1.4	1.9	3.0	5.8	10.8	21.0	36.3	59.5	116.1
Hispanic, both sexes.....	1.2	1.3	2.2	3.2	5.7	10.8	15.4	27.2	38.9	71.6
Male.....	1.5	1.6	2.7	4.7	8.0	15.0	20.2	39.1	54.9	86.8
Female.....	*	*	*	*	3.6	7.3	11.6	18.6	28.3	63.5
Mexican, both sexes.....	1.4	1.4	1.7	3.3	5.2	10.3	11.9	25.5	38.9	80.9
Male.....	*	*	*	*	*	*	*	34.7	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	20.1	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.2	2.3	3.3	5.4	9.2	16.7	29.1	48.4	73.9	131.2
Male.....	1.9	3.3	4.7	7.8	12.7	23.1	38.1	63.3	94.6	158.5
Female.....	0.6	1.4	1.9	3.1	6.0	11.1	21.7	37.5	61.1	118.5
Non-Hispanic white 6/, both sexes....	1.0	2.1	2.8	5.0	9.0	16.4	29.6	48.9	73.7	132.6
Male.....	1.6	3.0	4.1	7.2	12.5	22.8	39.1	64.8	96.2	163.3
Female.....	0.5	1.1	1.5	2.8	5.6	10.6	21.6	37.0	59.6	118.3
Non-Hispanic black 6/, both sexes....	2.8	4.5	7.3	10.3	13.8	23.6	31.2	52.2	85.1	134.0
Male.....	4.1	6.2	10.2	15.3	18.6	33.8	37.4	59.4	83.6	119.1
Female.....	1.7	3.0	4.9	6.1	10.1	16.0	26.9	47.8	85.9	140.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Aortic aneurysm and dissection (I71)

All origins 3/, both sexes.....	1.0	1.6	2.1	3.4	5.6	10.6	18.0	29.5	43.2	62.7
Male.....	1.5	2.4	3.3	5.3	8.4	15.2	24.8	41.1	59.0	87.6
Female.....	0.4	0.8	1.0	1.6	3.1	6.5	12.5	21.0	33.3	51.0
Hispanic, both sexes.....	0.8	0.8	1.4	1.8	3.4	5.9	8.7	14.3	19.3	31.7
Male.....	1.3	*	*	3.0	5.1	9.3	13.0	21.1	28.7	47.4
Female.....	*	*	*	*	*	*	*	9.3	13.1	23.4
Mexican, both sexes.....	*	*	*	*	*	*	*	13.1	*	29.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.0	1.6	2.2	3.5	5.8	10.9	18.7	30.5	44.4	64.2
Male.....	1.6	2.5	3.4	5.5	8.6	15.6	25.7	42.3	60.7	89.8
Female.....	0.4	0.8	1.1	1.7	3.2	6.8	13.0	21.8	34.3	52.3
Non-Hispanic white 6/, both sexes....	0.8	1.5	1.8	3.3	5.8	10.9	19.3	31.5	45.6	66.0
Male.....	1.3	2.3	3.0	5.2	8.7	15.7	26.7	44.2	63.0	92.8
Female.....	0.3	0.6	0.7	1.4	3.0	6.6	13.1	22.1	34.6	53.4
Non-Hispanic black 6/, both sexes....	2.3	3.1	5.2	6.2	7.6	13.2	16.2	24.1	34.2	47.8
Male.....	3.3	4.6	7.5	9.4	10.4	18.6	20.3	27.2	35.1	52.2
Female.....	1.3	1.9	3.1	3.5	5.4	9.3	13.5	22.3	33.7	46.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All origins 3/, both sexes.....	0.3	0.7	1.0	1.9	3.3	5.8	10.2	17.6	29.1	65.8
Male.....	0.3	0.7	1.2	2.3	4.0	7.5	12.1	20.8	33.5	67.3
Female.....	0.2	0.6	0.9	1.5	2.7	4.3	8.6	15.3	26.2	65.0
Hispanic, both sexes.....	*	*	*	1.4	2.3	4.9	6.7	13.0	19.6	39.9
Male.....	*	*	*	*	*	5.6	*	18.0	26.2	39.5
Female.....	*	*	*	*	*	*	6.2	9.3	15.3	40.1
Mexican, both sexes.....	*	*	*	*	*	6.3	*	12.3	24.1	51.3
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.2	0.7	1.1	1.9	3.4	5.8	10.4	17.9	29.5	67.0
Male.....	0.3	0.8	1.3	2.4	4.1	7.6	12.4	21.0	33.9	68.7
Female.....	*	0.6	0.9	1.5	2.8	4.3	8.7	15.7	26.8	66.2
Non-Hispanic white 6/, both sexes....	0.2	0.6	0.9	1.7	3.2	5.5	10.3	17.4	28.2	66.6
Male.....	*	0.7	1.1	2.0	3.8	7.1	12.4	20.6	33.2	70.5
Female.....	*	0.5	0.8	1.4	2.6	4.0	8.5	14.9	25.0	64.9
Non-Hispanic black 6/, both sexes.....	*	1.4	2.2	4.1	6.2	10.4	14.9	28.1	50.9	86.3
Male.....	*	1.7	2.7	5.9	8.2	15.2	17.1	32.2	48.5	66.9
Female.....	*	*	1.8	2.6	4.7	6.8	13.5	25.6	52.2	94.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

All origins 3/, both sexes.....	0.7	1.1	1.4	1.6	1.9	2.4	3.5	5.6	8.7	18.4
Male.....	0.8	1.3	1.8	1.9	2.3	2.4	3.5	6.1	8.3	16.2
Female.....	0.7	1.0	0.9	1.2	1.6	2.5	3.6	5.3	9.0	19.5
Hispanic, both sexes.....	*	*	1.3	*	*	*	*	*	6.9	11.3
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.8	1.2	1.3	1.6	2.0	2.4	3.6	5.8	8.8	18.8
Male.....	0.8	1.3	1.8	2.0	2.3	2.4	3.6	6.2	8.5	16.5
Female.....	0.7	1.1	0.9	1.3	1.7	2.4	3.7	5.4	9.1	19.8
Non-Hispanic white 6/, both sexes....	0.6	1.0	1.2	1.5	1.7	2.3	3.5	5.7	8.5	18.5
Male.....	0.7	1.1	1.5	1.9	1.9	2.3	3.5	6.1	7.7	16.3
Female.....	0.5	0.9	0.8	1.1	1.5	2.2	3.5	5.4	9.0	19.5
Non-Hispanic black 6/, both sexes....	1.9	2.8	3.0	2.8	4.5	4.1	5.9	8.3	15.0	26.0
Male.....	1.8	2.7	4.6	3.0	6.2	*	*	*	17.6	22.9
Female.....	2.0	2.9	1.6	2.7	3.1	3.7	6.1	8.2	13.7	27.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Influenza and pneumonia (J10-J18)

All origins 3/, both sexes.....	2.4	3.6	5.7	7.6	13.2	23.1	42.9	86.2	181.8	502.5
Male.....	2.7	4.2	7.1	9.4	16.4	28.1	52.5	108.1	226.8	586.4
Female.....	2.2	3.1	4.4	6.0	10.3	18.8	35.0	70.2	153.7	463.3
Hispanic, both sexes.....	1.5	3.3	4.1	6.9	14.0	19.8	44.9	87.6	154.4	390.9
Male.....	1.8	3.7	5.5	9.3	17.4	24.8	52.6	104.6	174.4	440.9
Female.....	*	3.0	2.6	4.5	11.1	15.6	38.9	75.3	141.1	364.3
Mexican, both sexes.....	1.4	3.4	3.7	6.3	12.9	22.6	42.8	85.6	167.3	440.9
Male.....	*	3.5	4.6	7.8	15.1	23.8	48.4	103.1	199.0	*
Female.....	*	3.4	*	*	10.8	21.5	38.3	73.4	148.2	404.8
Puerto Rican, both sexes.....	*	*	*	*	31.2	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	6.0	10.9	12.4	39.4	89.1	128.4	368.4
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.6	3.7	5.9	7.7	13.1	23.3	42.5	86.0	183.1	507.5
Male.....	2.8	4.2	7.3	9.4	16.3	28.2	52.2	108.2	229.4	593.8
Female.....	2.3	3.1	4.6	6.1	10.2	19.0	34.6	69.7	154.2	467.4
Non-Hispanic white 6/, both sexes....	2.4	3.2	5.3	7.2	12.5	21.7	42.0	85.0	182.7	514.4
Male.....	2.7	3.7	6.3	8.8	15.2	26.1	50.7	106.2	228.4	598.8
Female.....	2.1	2.7	4.4	5.6	10.0	17.8	34.7	69.1	153.9	475.0
Non-Hispanic black 6/, both sexes....	4.2	7.3	11.0	13.2	20.8	40.5	51.7	101.4	197.4	439.6
Male.....	4.4	8.4	15.7	16.4	30.3	51.5	70.0	141.8	243.7	512.2
Female.....	4.0	6.3	7.0	10.5	13.5	32.4	39.1	77.2	172.9	409.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonia (J12-J18)

All origins 3/, both sexes.....	2.4	3.6	5.6	7.5	13.0	22.8	42.3	85.0	179.3	495.5
Male.....	2.7	4.2	7.0	9.3	16.3	27.8	51.8	106.9	223.8	578.3
Female.....	2.1	3.1	4.3	5.9	10.1	18.5	34.5	69.0	151.5	456.7
Hispanic, both sexes.....	1.5	3.3	4.0	6.9	13.8	19.7	44.9	87.3	153.8	388.9
Male.....	1.8	3.6	5.4	9.3	17.2	24.5	52.6	104.6	174.4	438.7
Female.....	*	3.0	2.5	4.5	10.7	15.6	38.9	74.9	140.1	362.5
Mexican, both sexes.....	1.4	3.4	3.6	6.3	12.7	22.6	42.8	85.6	166.7	437.4
Male.....	*	3.3	4.6	7.8	14.7	23.8	48.4	103.1	199.0	*
Female.....	*	3.4	*	*	10.8	21.5	38.3	73.4	147.2	400.6
Puerto Rican, both sexes.....	*	*	*	*	30.3	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	6.0	10.9	11.9	39.4	89.1	127.1	366.9
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.5	3.6	5.8	7.6	13.0	23.0	41.9	84.6	180.4	500.2
Male.....	2.8	4.2	7.2	9.3	16.2	27.9	51.5	106.8	226.2	585.5
Female.....	2.3	3.1	4.5	5.9	10.0	18.7	34.1	68.4	151.9	460.6
Non-Hispanic white 6/, both sexes....	2.4	3.2	5.2	7.0	12.3	21.4	41.3	83.5	179.9	506.5
Male.....	2.6	3.6	6.2	8.7	15.0	25.7	49.9	104.8	225.2	589.9
Female.....	2.1	2.7	4.2	5.5	9.8	17.5	34.1	67.7	151.4	467.7
Non-Hispanic black 6/, both sexes....	4.2	7.2	10.9	13.2	20.7	40.3	51.5	101.0	195.4	437.7
Male.....	4.4	8.4	15.6	16.4	30.1	51.2	70.0	141.3	240.2	509.4
Female.....	3.9	6.3	7.0	10.5	13.3	32.2	38.9	77.0	171.8	408.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic lower respiratory diseases (J40-J47)

All origins 3/, both sexes.....	2.8	5.8	12.7	25.8	57.4	111.8	194.8	308.5	434.8	589.1
Male.....	2.6	5.5	13.4	27.4	62.0	119.8	218.0	364.6	541.3	760.8
Female.....	2.9	6.1	12.1	24.2	53.3	104.8	175.8	267.5	368.3	508.7
Hispanic, both sexes.....	0.9	1.8	3.7	8.1	17.7	32.1	62.9	120.6	203.2	374.4
Male.....	*	1.8	4.0	8.9	21.7	39.5	79.7	155.0	262.0	484.9
Female.....	*	1.8	3.4	7.3	14.1	25.8	49.9	95.8	164.0	315.7
Mexican, both sexes.....	*	1.4	2.7	6.8	14.4	32.9	51.5	114.7	185.8	377.4
Male.....	*	*	*	8.1	16.8	41.0	63.6	154.1	271.4	*
Female.....	*	*	*	5.5	12.1	25.0	41.6	87.0	134.4	282.4
Puerto Rican, both sexes.....	*	*	*	19.0	34.6	37.0	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	108.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	15.6	25.3	54.7	124.2	199.5	315.3
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.0	6.3	13.6	27.3	60.5	118.0	204.3	320.5	447.3	599.3
Male.....	2.9	5.9	14.3	29.0	65.0	125.9	227.6	377.8	557.0	775.2
Female.....	3.2	6.6	13.0	25.7	56.5	111.0	185.4	278.6	379.0	517.5
Non-Hispanic white 6/, both sexes....	2.7	6.3	13.8	28.4	64.1	126.9	221.1	340.5	472.0	629.8
Male.....	2.6	5.8	14.2	29.6	66.7	132.0	241.0	393.5	577.2	804.2
Female.....	2.8	6.8	13.3	27.2	61.6	122.3	204.5	300.9	405.8	548.5
Non-Hispanic black 6/, both sexes....	5.4	7.8	17.5	28.2	52.9	85.2	125.1	203.8	266.7	337.4
Male.....	5.3	8.0	20.3	34.0	71.5	114.6	172.5	294.2	422.8	522.3
Female.....	5.6	7.7	15.1	23.4	38.5	63.6	92.5	149.8	184.4	262.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Emphysema (J43)

All origins 3/, both sexes.....	0.3	0.7	1.4	3.2	7.1	13.2	22.3	33.1	41.6	45.3
Male.....	0.4	0.7	1.9	3.5	7.9	15.1	25.2	40.3	52.8	63.7
Female.....	0.3	0.7	1.0	2.8	6.4	11.5	19.9	27.8	34.6	36.7
Hispanic, both sexes.....	*	*	*	*	2.2	3.5	4.7	13.2	19.3	20.7
Male.....	*	*	*	*	*	*	*	21.1	22.9	31.6
Female.....	*	*	*	*	*	*	*	7.5	16.9	15.0
Mexican, both sexes.....	*	*	*	*	*	*	*	10.4	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.4	0.8	1.5	3.3	7.5	13.9	23.6	34.4	42.8	46.5
Male.....	0.4	0.8	2.0	3.6	8.2	15.9	26.5	41.6	54.5	65.3
Female.....	0.3	0.7	1.1	3.0	6.8	12.3	21.2	29.2	35.5	37.8
Non-Hispanic white 6/, both sexes....	0.4	0.8	1.7	3.6	8.0	15.1	25.9	36.7	45.3	49.2
Male.....	0.4	0.8	2.0	3.8	8.3	16.6	28.6	43.1	56.5	68.4
Female.....	0.3	0.8	1.3	3.4	7.6	13.6	23.6	31.9	38.2	40.3
Non-Hispanic black 6/, both sexes.....	*	*	1.3	2.6	6.7	9.1	11.9	21.0	25.0	24.2
Male.....	*	*	2.4	3.9	11.1	13.8	14.9	33.1	42.1	40.3
Female.....	*	*	*	*	3.3	5.7	9.8	13.7	15.9	17.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Asthma (J45-J46)

All origins 3/, both sexes.....	1.0	1.2	1.4	1.4	1.8	2.0	2.7	3.6	6.4	12.3
Male.....	0.8	0.9	1.1	1.0	1.2	1.4	2.1	2.9	4.4	9.1
Female.....	1.2	1.6	1.7	1.7	2.3	2.6	3.2	4.2	7.6	13.9
Hispanic, both sexes.....	*	*	*	*	2.2	*	*	*	*	16.4
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	19.8
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.1	1.3	1.5	1.4	1.8	2.0	2.7	3.6	6.4	12.1
Male.....	0.9	0.9	1.2	1.0	1.2	1.4	2.1	2.9	4.4	9.0
Female.....	1.3	1.7	1.7	1.8	2.3	2.5	3.2	4.1	7.6	13.6
Non-Hispanic white 6/, both sexes....	0.8	0.9	1.1	1.0	1.3	1.6	2.3	3.4	6.0	11.8
Male.....	0.7	0.6	0.7	0.6	0.8	1.0	1.6	2.8	3.9	9.0
Female.....	1.0	1.3	1.4	1.4	1.8	2.1	2.8	3.8	7.4	13.2
Non-Hispanic black 6/, both sexes....	2.8	3.7	4.4	4.5	6.2	4.9	6.3	5.2	7.5	12.7
Male.....	2.8	2.9	4.3	4.1	5.7	*	*	*	*	*
Female.....	2.8	4.4	4.5	4.8	6.7	5.5	6.7	6.0	8.5	15.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic lower respiratory diseases (J44,J47)

All origins 3/, both sexes.....	1.4	3.8	9.8	21.1	48.3	96.3	169.1	270.7	384.8	525.4
Male.....	1.3	3.9	10.3	22.8	52.7	103.1	190.1	320.3	481.9	682.2
Female.....	1.4	3.8	9.3	19.6	44.3	90.3	152.0	234.3	324.2	452.1
Hispanic, both sexes.....	*	1.1	2.4	5.9	13.3	26.0	55.8	103.6	175.4	331.8
Male.....	*	*	2.7	6.9	16.4	32.7	71.0	131.9	230.9	439.8
Female.....	*	*	2.1	4.9	10.5	20.4	43.9	83.1	138.4	274.4
Mexican, both sexes.....	*	*	2.1	4.9	10.8	28.4	43.4	100.4	160.5	344.4
Male.....	*	*	*	6.0	11.6	35.4	54.3	133.9	233.6	*
Female.....	*	*	*	*	10.1	21.5	34.5	76.8	116.6	251.8
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	92.5	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	13.3	19.1	47.7	110.2	172.2	268.2
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.5	4.2	10.5	22.4	51.1	101.7	177.3	281.4	396.1	534.6
Male.....	1.5	4.2	11.1	24.1	55.4	108.3	198.3	332.2	495.9	694.8
Female.....	1.5	4.2	10.0	20.8	47.1	95.9	160.2	244.1	333.9	460.1
Non-Hispanic white 6/, both sexes....	1.5	4.5	10.9	23.7	54.6	109.8	192.2	299.3	418.6	562.5
Male.....	1.5	4.4	11.4	25.1	57.5	113.9	210.0	346.5	514.6	720.3
Female.....	1.5	4.6	10.5	22.4	51.8	106.2	177.4	263.9	358.2	488.9
Non-Hispanic black 6/, both sexes....	2.0	3.3	11.7	20.8	39.9	70.9	106.1	176.2	232.3	296.8
Male.....	1.8	3.9	13.6	25.5	54.6	96.4	151.0	255.0	372.9	473.7
Female.....	2.2	2.9	10.1	16.9	28.5	52.0	75.4	129.2	158.1	224.6

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonitis due to solids and liquids (J69)

All origins 3/, both sexes.....	0.5	0.8	1.2	2.0	3.7	6.3	12.8	26.5	58.7	155.8
Male.....	0.6	1.0	1.6	2.5	4.9	8.7	17.5	37.3	85.7	224.8
Female.....	0.4	0.7	0.9	1.5	2.6	4.3	8.9	18.6	41.9	123.5
Hispanic, both sexes.....	*	*	*	*	*	3.5	8.5	18.6	37.0	77.5
Male.....	*	*	*	*	*	*	11.1	24.2	44.2	86.8
Female.....	*	*	*	*	*	*	6.5	14.5	32.2	72.5
Mexican, both sexes.....	*	*	*	*	*	*	8.3	19.5	34.6	84.4
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.5	0.9	1.3	2.1	3.8	6.5	13.1	27.0	60.0	159.6
Male.....	0.6	1.0	1.7	2.6	5.1	9.0	18.0	38.1	88.1	232.3
Female.....	0.4	0.7	0.9	1.6	2.7	4.4	9.1	18.9	42.4	125.8
Non-Hispanic white 6/, both sexes....	0.5	0.8	1.3	2.0	3.9	6.4	12.7	26.8	61.4	164.1
Male.....	0.6	1.0	1.7	2.5	5.1	8.6	17.4	37.2	90.1	239.9
Female.....	0.4	0.6	0.9	1.5	2.8	4.4	8.8	19.0	43.3	128.7
Non-Hispanic black 6/, both sexes....	1.0	1.2	1.7	3.0	5.0	9.3	20.1	33.6	56.0	130.9
Male.....	*	*	2.4	3.8	7.7	15.5	28.8	56.2	87.1	177.8
Female.....	*	*	*	2.3	2.8	4.8	14.1	20.1	39.6	111.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All origins 3/, both sexes.....	1.5	2.3	4.0	6.7	11.9	21.4	36.6	59.1	91.9	136.6
Male.....	1.5	2.4	4.3	7.7	14.0	26.0	46.3	75.7	118.0	163.7
Female.....	1.5	2.3	3.8	5.7	10.0	17.3	28.7	47.0	75.6	124.0
Hispanic, both sexes.....	1.0	2.0	3.4	5.6	10.5	20.3	32.1	46.9	78.5	106.8
Male.....	*	2.1	3.6	7.7	13.7	23.4	36.2	55.6	86.0	112.8
Female.....	*	2.0	3.3	3.6	7.6	17.8	28.8	40.6	73.6	103.6
Mexican, both sexes.....	*	2.2	3.6	6.3	11.3	20.6	33.0	47.8	73.5	134.8
Male.....	*	*	*	8.7	15.1	21.8	37.1	58.7	*	*
Female.....	*	*	4.2	*	7.8	19.5	29.7	40.1	67.2	126.6
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	10.1	20.5	31.8	50.1	94.3	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.6	2.4	4.1	6.7	12.0	21.4	36.9	59.9	92.6	138.1
Male.....	1.5	2.5	4.3	7.7	14.0	26.2	47.0	77.0	119.9	166.4
Female.....	1.6	2.3	3.8	5.8	10.2	17.2	28.7	47.3	75.5	124.9
Non-Hispanic white 6/, both sexes....	1.4	2.0	3.7	6.3	11.7	21.7	38.3	62.1	96.0	142.9
Male.....	1.3	2.2	4.0	7.5	13.6	26.7	48.4	79.8	124.1	171.5
Female.....	1.4	1.9	3.3	5.1	10.0	17.2	29.9	48.8	78.3	129.6
Non-Hispanic black 6/, both sexes....	2.9	5.0	7.2	11.3	16.4	20.5	32.8	44.8	62.1	96.6
Male.....	3.0	5.0	6.8	10.6	20.3	24.9	44.6	58.5	81.5	119.1
Female.....	2.9	5.1	7.5	11.9	13.5	17.3	24.8	36.6	51.8	87.4

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Peptic ulcer (K25-K28)

All origins 3/, both sexes.....	0.2	0.5	0.8	1.0	1.5	2.1	3.5	5.3	9.6	20.7
Male.....	*	0.8	1.0	1.4	1.9	2.6	4.0	6.3	11.0	22.0
Female.....	0.2	0.3	0.5	0.6	1.1	1.7	3.1	4.6	8.8	20.1
Hispanic, both sexes.....	*	*	*	*	*	*	*	5.2	*	19.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	21.0
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.2	0.6	0.8	1.0	1.6	2.1	3.6	5.3	9.8	20.7
Male.....	*	0.8	1.1	1.5	1.9	2.5	4.0	6.2	11.3	22.2
Female.....	0.3	0.3	0.6	0.6	1.2	1.7	3.2	4.6	8.9	20.0
Non-Hispanic white 6/, both sexes....	0.2	0.5	0.7	1.0	1.4	2.0	3.4	5.2	10.1	21.2
Male.....	*	0.7	0.8	1.4	1.8	2.3	3.6	5.9	11.6	22.2
Female.....	*	0.3	0.5	0.5	1.1	1.6	3.3	4.6	9.1	20.7
Non-Hispanic black 6/, both sexes.....	*	1.1	1.7	1.8	2.6	3.0	4.8	7.2	7.5	14.9
Male.....	*	1.8	2.9	2.6	*	*	8.2	10.1	*	*
Female.....	*	*	*	*	*	*	*	5.5	*	14.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All origins 3/, both sexes.....	8.2	14.8	21.2	22.8	22.9	24.8	27.4	29.6	28.2	19.4
Male.....	10.8	20.7	31.8	33.7	33.1	34.1	35.5	36.9	34.4	26.1
Female.....	5.6	9.0	11.1	12.5	13.6	16.7	20.9	24.2	24.3	16.2
Hispanic, both sexes.....	9.4	19.4	29.4	33.0	36.1	41.2	45.2	52.3	57.6	40.3
Male.....	14.1	30.7	46.3	52.7	56.3	55.6	58.7	62.3	59.0	51.9
Female.....	4.3	7.5	12.6	14.4	18.1	29.1	34.7	45.1	56.7	34.2
Mexican, both sexes.....	11.4	22.8	32.7	37.2	40.1	52.5	54.7	60.5	65.4	51.3
Male.....	17.1	35.4	50.7	56.6	60.7	64.2	70.2	78.0	*	*
Female.....	4.8	9.1	12.8	17.0	20.9	41.0	42.1	48.2	60.3	*
Puerto Rican, both sexes.....	*	21.0	30.7	42.6	51.9	38.0	*	*	*	*
Male.....	*	40.7	*	67.4	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.2	13.1	23.5	23.4	25.7	26.7	30.4	54.1	*	*
Male.....	9.9	21.0	35.8	39.0	40.1	*	*	*	*	*
Female.....	*	*	13.2	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	8.0	14.1	20.3	21.8	21.7	23.4	26.1	28.0	26.4	18.3
Male.....	10.2	19.3	30.2	31.9	31.1	32.2	33.7	35.1	32.8	24.6
Female.....	5.8	9.1	10.9	12.3	13.2	15.7	19.8	22.8	22.5	15.3
Non-Hispanic white 6/, both sexes....	8.2	14.7	20.7	22.1	22.2	24.1	27.6	29.0	27.7	19.0
Male.....	10.5	20.1	30.6	32.1	31.6	32.8	35.4	36.0	34.7	25.6
Female.....	6.0	9.4	11.0	12.5	13.4	16.3	21.0	23.8	23.4	15.9
Non-Hispanic black 6/, both sexes....	6.6	11.2	21.3	23.5	20.8	21.1	16.4	19.6	11.6	10.9
Male.....	9.1	15.3	32.2	36.4	30.1	34.6	22.5	32.2	*	*
Female.....	4.4	7.6	12.2	12.9	13.6	11.2	12.2	12.1	11.1	9.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcoholic liver disease (K70)

All origins 3/, both sexes.....	5.5	9.2	12.0	12.5	11.6	10.1	8.6	7.0	4.5	2.2
Male.....	7.1	12.9	18.4	19.4	18.1	16.9	14.1	12.0	7.5	4.7
Female.....	3.8	5.6	5.9	6.0	5.6	4.2	4.1	3.4	2.6	1.1
Hispanic, both sexes.....	6.5	12.7	18.3	18.9	20.0	17.5	16.2	16.2	11.8	*
Male.....	9.6	20.5	30.4	32.5	35.8	31.9	32.4	30.9	21.3	*
Female.....	3.0	4.5	6.3	6.0	6.0	5.4	*	*	*	*
Mexican, both sexes.....	7.8	15.0	21.1	22.6	22.7	22.9	21.4	20.3	17.9	*
Male.....	11.4	24.3	34.4	37.1	38.9	37.9	41.1	40.4	*	*
Female.....	3.5	4.8	6.5	7.6	7.5	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	20.7	32.0	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.2	9.9	15.3	12.8	12.1	13.8	*	*	*	*
Male.....	7.3	15.3	25.9	24.3	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	5.3	8.7	11.4	11.9	10.8	9.5	8.0	6.4	4.0	1.9
Male.....	6.6	11.8	17.1	18.1	16.6	15.6	12.8	10.7	6.7	3.9
Female.....	4.0	5.7	5.9	6.0	5.5	4.1	4.1	3.3	2.4	1.0
Non-Hispanic white 6/, both sexes....	5.6	9.2	11.8	12.3	11.2	9.6	8.3	6.6	4.2	2.0
Male.....	7.1	12.3	17.7	18.6	17.2	15.6	13.2	11.0	7.1	4.0
Female.....	4.1	6.1	6.0	6.2	5.6	4.3	4.2	3.4	2.5	1.1
Non-Hispanic black 6/, both sexes....	3.5	6.0	9.7	10.0	8.8	8.8	6.4	5.3	*	*
Male.....	4.4	8.8	14.3	16.4	13.0	16.7	9.8	10.1	*	*
Female.....	2.6	3.6	5.7	4.8	5.5	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic liver disease and cirrhosis (K73-K74)

All origins 3/, both sexes.....	2.7	5.6	9.2	10.3	11.4	14.7	18.9	22.5	23.7	17.2
Male.....	3.7	7.8	13.4	14.4	15.0	17.2	21.3	24.9	26.9	21.4
Female.....	1.8	3.4	5.1	6.5	8.1	12.5	16.8	20.7	21.7	15.2
Hispanic, both sexes.....	2.9	6.7	11.1	14.1	16.1	23.7	29.1	36.1	45.8	32.9
Male.....	4.4	10.2	16.0	20.2	20.5	23.7	26.3	31.4	37.7	33.8
Female.....	*	3.0	6.3	8.4	12.2	23.7	31.2	39.5	51.2	32.4
Mexican, both sexes.....	3.7	7.9	11.6	14.5	17.4	29.7	33.3	40.2	47.5	41.7
Male.....	5.7	11.2	16.3	19.6	21.8	26.3	29.1	37.6	*	*
Female.....	*	4.3	6.3	9.3	13.4	33.0	36.7	42.1	51.4	*
Puerto Rican, both sexes.....	*	11.3	17.2	21.9	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	3.2	8.2	10.6	13.7	12.9	21.4	39.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.7	5.4	8.9	9.9	10.9	13.9	18.1	21.6	22.4	16.4
Male.....	3.5	7.5	13.1	13.8	14.5	16.7	21.0	24.4	26.1	20.8
Female.....	1.9	3.4	5.0	6.3	7.7	11.5	15.7	19.5	20.1	14.3
Non-Hispanic white 6/, both sexes....	2.7	5.6	8.9	9.8	11.0	14.4	19.3	22.4	23.5	16.9
Male.....	3.4	7.8	12.9	13.5	14.3	17.2	22.2	25.0	27.6	21.5
Female.....	2.0	3.3	5.1	6.2	7.8	12.0	16.8	20.4	20.9	14.8
Non-Hispanic black 6/, both sexes....	3.1	5.1	11.7	13.5	12.1	12.3	9.9	14.3	9.2	10.1
Male.....	4.7	6.5	17.8	20.0	17.2	17.9	12.7	22.1	*	*
Female.....	1.8	4.0	6.4	8.1	8.1	8.2	8.0	9.6	9.3	9.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All origins 3/, both sexes.....	0.1	0.2	0.4	0.6	1.2	1.8	3.3	6.2	9.9	21.8
Male.....	*	0.2	0.5	0.6	1.2	2.3	4.0	7.6	12.5	21.7
Female.....	*	0.2	0.3	0.5	1.2	1.3	2.7	5.2	8.3	21.9
Hispanic, both sexes.....	*	*	*	*	*	*	*	7.6	10.1	20.0
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	22.8
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.1	0.2	0.4	0.6	1.2	1.8	3.3	6.1	9.9	21.9
Male.....	*	0.2	0.5	0.6	1.2	2.4	4.0	7.5	12.6	22.0
Female.....	*	0.2	0.3	0.5	1.2	1.3	2.7	5.1	8.2	21.9
Non-Hispanic white 6/, both sexes.....	*	0.2	0.3	0.5	1.1	1.5	3.0	5.9	9.9	22.4
Male.....	*	*	0.4	0.5	1.1	2.1	3.7	7.4	12.8	23.2
Female.....	*	*	*	0.5	1.1	1.0	2.3	4.9	8.1	22.0
Non-Hispanic black 6/, both sexes.....	*	*	1.0	1.1	1.9	3.5	4.8	9.3	9.2	17.5
Male.....	*	*	*	*	*	*	*	10.6	*	*
Female.....	*	*	*	*	*	*	5.4	8.5	9.6	20.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All origins 3/, both sexes.....	2.3	3.8	6.7	10.7	18.1	30.5	50.3	86.4	144.0	290.5
Male.....	2.7	4.7	8.0	12.4	21.0	33.8	59.4	105.1	182.1	377.1
Female.....	1.9	2.9	5.5	9.0	15.5	27.6	42.8	72.7	120.2	250.0
Hispanic, both sexes.....	1.4	3.5	6.7	11.3	18.9	34.6	47.6	78.9	130.6	206.2
Male.....	1.9	4.7	8.2	12.2	21.1	41.5	57.2	89.1	153.1	241.3
Female.....	*	2.3	5.3	10.4	17.0	28.9	40.1	71.5	115.5	187.5
Mexican, both sexes.....	1.4	3.6	7.2	14.5	20.3	41.2	51.5	84.1	148.8	241.8
Male.....	*	4.6	8.4	15.0	20.0	45.0	60.3	92.5	184.2	*
Female.....	*	*	5.8	14.1	20.6	37.5	44.3	78.1	127.5	228.1
Puerto Rican, both sexes.....	*	*	*	*	*	39.8	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	6.0	16.8	23.4	25.6	62.1	88.8	159.1
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.4	3.8	6.7	10.6	18.0	30.1	50.3	86.7	144.6	294.4
Male.....	2.9	4.7	8.0	12.4	21.0	33.1	59.5	105.9	183.5	384.0
Female.....	2.0	3.0	5.5	8.9	15.3	27.4	42.9	72.6	120.4	252.7
Non-Hispanic white 6/, both sexes....	1.5	2.7	4.4	7.7	13.7	23.5	42.4	78.9	136.4	284.8
Male.....	1.8	3.4	5.3	9.1	16.7	26.4	51.8	98.2	176.0	380.7
Female.....	1.1	2.0	3.5	6.5	10.9	20.9	34.6	64.5	111.5	240.1
Non-Hispanic black 6/, both sexes....	7.9	11.3	23.1	33.4	56.7	88.2	125.1	181.2	259.7	449.4
Male.....	9.4	14.1	28.5	41.3	66.1	101.3	138.7	215.0	318.8	496.6
Female.....	6.7	8.9	18.4	26.9	49.4	78.6	115.8	161.1	228.5	430.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All origins 3/, both sexes.....	*	0.1	0.3	0.4	0.6	1.0	1.8	3.2	6.0	13.4
Male.....	*	*	0.3	0.4	0.8	1.3	1.8	4.3	7.7	17.2
Female.....	*	*	0.3	0.4	0.4	0.8	1.7	2.3	5.0	11.6
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	8.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	*	0.1	0.3	0.4	0.6	1.0	1.8	3.2	6.0	13.6
Male.....	*	*	0.3	0.4	0.8	1.3	1.8	4.3	7.7	17.4
Female.....	*	*	0.3	0.3	0.4	0.8	1.8	2.4	4.9	11.9
Non-Hispanic white 6/, both sexes.....	*	*	0.2	0.3	0.4	0.7	1.6	2.9	5.4	12.7
Male.....	*	*	*	*	0.6	0.9	1.7	3.9	7.2	16.2
Female.....	*	*	*	0.3	*	0.5	1.5	2.2	4.2	11.1
Non-Hispanic black 6/, both sexes.....	*	*	0.9	1.4	1.8	3.7	4.3	5.7	12.1	23.9
Male.....	*	*	*	*	*	5.1	*	*	14.7	33.0
Female.....	*	*	*	*	*	*	4.8	*	10.7	20.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Renal failure (N17-N19)

All origins 3/, both sexes.....	2.2	3.7	6.4	10.2	17.5	29.3	48.4	83.0	137.5	276.4
Male.....	2.6	4.5	7.7	12.0	20.2	32.4	57.4	100.4	174.0	358.8
Female.....	1.8	2.8	5.2	8.6	15.1	26.6	41.0	70.2	114.8	237.8
Hispanic, both sexes.....	1.2	3.4	6.4	10.9	18.1	33.2	45.9	76.1	123.7	197.2
Male.....	1.7	4.5	8.0	11.9	19.9	40.4	54.9	85.0	146.6	227.8
Female.....	*	2.3	4.9	9.9	16.5	27.2	38.9	69.7	108.5	180.9
Mexican, both sexes.....	1.2	3.5	6.7	14.0	19.5	40.0	50.0	81.3	138.9	228.7
Male.....	*	4.3	8.1	14.5	18.9	44.0	57.6	88.6	176.0	*
Female.....	*	*	5.3	13.4	19.9	36.0	43.7	76.1	116.6	215.6
Puerto Rican, both sexes.....	*	*	*	*	*	37.0	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	6.0	15.2	22.4	25.6	59.1	86.1	148.8
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.4	3.7	6.4	10.1	17.4	28.9	48.4	83.2	138.2	280.0
Male.....	2.8	4.5	7.6	11.9	20.2	31.6	57.5	101.3	175.2	365.5
Female.....	1.9	2.9	5.2	8.5	14.9	26.5	41.0	70.0	115.0	240.2
Non-Hispanic white 6/, both sexes....	1.4	2.6	4.2	7.4	13.2	22.7	40.8	75.7	130.6	271.3
Male.....	1.8	3.3	5.1	8.7	16.1	25.4	50.0	93.9	168.4	363.4
Female.....	1.1	1.9	3.3	6.1	10.6	20.3	33.1	62.1	106.8	228.4
Non-Hispanic black 6/, both sexes....	7.7	10.9	21.9	32.0	54.8	84.2	120.7	175.6	246.8	425.2
Male.....	9.3	13.5	27.2	39.1	63.3	96.0	135.2	207.2	302.7	462.7
Female.....	6.3	8.6	17.5	26.1	48.1	75.5	110.8	156.7	217.4	409.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All origins 3/, both sexes.....	1.1	1.5	1.6	2.2	2.4	1.9	2.0	2.7	3.5	5.6
Male.....	1.2	1.7	1.9	2.5	2.7	1.9	2.3	3.0	4.5	6.8
Female.....	1.0	1.3	1.3	1.9	2.1	1.8	1.8	2.4	2.8	5.1
Hispanic, both sexes.....	*	1.2	1.1	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	1.4	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.1	1.5	1.6	2.3	2.4	1.9	2.0	2.7	3.5	5.8
Male.....	1.3	1.8	2.0	2.6	2.8	1.9	2.3	3.0	4.5	6.8
Female.....	1.0	1.3	1.3	2.0	2.1	1.9	1.8	2.5	2.9	5.3
Non-Hispanic white 6/, both sexes....	1.2	1.6	1.6	2.4	2.6	1.9	2.1	2.9	3.5	5.9
Male.....	1.3	1.7	2.0	2.7	2.9	1.9	2.3	3.0	4.7	6.8
Female.....	1.1	1.4	1.3	2.2	2.3	1.9	1.9	2.7	2.8	5.5
Non-Hispanic black 6/, both sexes....	1.1	1.7	2.1	2.1	1.8	*	*	*	*	5.6
Male.....	*	2.2	2.6	2.6	*	*	*	*	*	*
Female.....	*	*	1.7	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All origins 3/, both sexes.....	4.2	4.9	5.6	5.9	7.2	9.5	15.5	28.7	58.6	236.1
Male.....	5.5	6.3	7.2	7.7	9.0	11.9	18.5	32.3	61.3	191.1
Female.....	3.0	3.5	4.1	4.2	5.6	7.3	13.1	26.0	56.9	257.2
Hispanic, both sexes.....	2.5	3.1	2.9	3.4	4.4	5.5	8.7	13.2	27.2	117.0
Male.....	4.0	4.3	4.1	4.5	6.8	7.9	12.2	12.4	25.4	90.2
Female.....	*	1.8	*	*	*	*	5.9	13.8	28.3	131.2
Mexican, both sexes.....	2.5	2.6	2.4	3.4	3.9	6.0	8.9	13.5	28.4	126.1
Male.....	3.9	3.3	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	141.9
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	101.7
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	4.4	5.0	5.9	6.1	7.4	9.7	16.0	29.7	60.3	241.9
Male.....	5.7	6.5	7.5	7.9	9.1	12.2	18.8	33.5	63.4	196.3
Female.....	3.3	3.6	4.4	4.3	5.9	7.6	13.6	26.8	58.4	263.1
Non-Hispanic white 6/, both sexes....	4.2	4.9	5.5	5.7	7.2	9.4	15.4	29.2	60.1	247.4
Male.....	5.3	6.1	6.9	7.4	8.9	11.7	17.4	32.7	62.6	199.4
Female.....	3.2	3.6	4.1	4.0	5.5	7.3	13.7	26.7	58.5	269.7
Non-Hispanic black 6/, both sexes....	7.0	7.7	10.4	10.6	11.3	15.3	25.0	42.2	77.6	226.9
Male.....	9.3	11.0	14.4	14.5	13.3	21.0	37.0	54.3	94.8	196.1
Female.....	5.0	4.7	7.0	7.4	9.8	11.0	16.7	34.9	68.5	239.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All origins 3/, both sexes.....	23.0	33.3	47.6	61.8	85.0	118.6	191.7	335.2	646.8	1,724.2
Male.....	27.2	39.7	57.9	73.0	96.3	131.0	207.6	349.9	652.5	1,455.0
Female.....	19.0	27.0	37.8	51.3	74.7	107.8	178.8	324.4	643.2	1,850.2
Hispanic, both sexes.....	18.2	23.9	35.5	47.1	61.2	89.9	146.3	249.1	443.0	1,067.7
Male.....	24.4	30.2	46.7	59.4	71.4	104.1	159.4	254.4	461.0	897.6
Female.....	11.2	17.3	24.2	35.5	52.0	77.9	136.0	245.2	431.1	1,158.1
Mexican, both sexes.....	17.4	22.4	34.0	47.4	61.3	93.3	138.0	238.6	429.6	1,107.0
Male.....	23.1	27.9	42.9	58.2	69.1	89.5	149.7	242.7	514.8	*
Female.....	10.8	16.5	24.2	36.1	53.9	97.0	128.5	235.7	378.4	1,201.8
Puerto Rican, both sexes.....	33.9	45.3	59.8	61.6	84.8	96.3	*	*	*	*
Male.....	51.0	69.1	92.4	72.1	*	*	*	*	*	*
Female.....	*	26.4	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	125.1	219.1	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	16.2	20.0	29.6	41.0	55.0	78.2	132.8	242.4	400.3	894.4
Male.....	21.4	24.3	40.7	54.8	66.3	106.3	*	*	*	*
Female.....	10.7	15.7	20.3	29.9	45.8	59.8	137.6	263.1	*	*
Non-Hispanic 5/, both sexes.....	23.8	34.4	48.7	63.0	86.8	120.7	194.8	340.4	657.7	1,755.2
Male.....	27.5	40.8	58.8	73.9	98.1	132.7	210.5	355.5	663.2	1,484.0
Female.....	20.1	28.1	39.1	52.6	76.5	110.2	181.9	329.4	654.3	1,881.3
Non-Hispanic white 6/, both sexes....	21.9	32.0	45.5	59.0	84.3	117.9	192.2	341.4	661.8	1,790.9
Male.....	26.0	38.9	55.6	69.2	94.6	130.1	205.7	353.3	663.9	1,512.3
Female.....	17.9	25.2	35.6	49.1	74.7	107.0	181.0	332.5	660.5	1,920.7
Non-Hispanic black 6/, both sexes....	38.5	56.3	82.8	108.4	129.4	169.1	261.0	406.5	743.9	1,660.0
Male.....	41.5	63.0	97.5	129.9	157.9	192.9	309.3	464.0	816.7	1,437.6
Female.....	35.9	50.4	70.4	90.6	107.3	151.6	227.8	372.2	705.4	1,750.7

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All origins 3/, both sexes.....	43.3	47.0	43.7	37.1	35.0	39.6	50.5	81.3	136.0	274.9
Male.....	60.8	66.8	63.0	52.9	49.6	55.1	66.4	107.3	176.5	337.1
Female.....	25.8	27.8	25.2	22.2	21.7	26.0	37.5	62.3	110.7	245.8
Hispanic, both sexes.....	34.6	36.2	34.9	34.1	31.9	35.0	40.9	63.8	91.6	142.8
Male.....	52.8	53.6	54.6	52.4	49.3	51.9	57.2	84.5	113.8	174.8
Female.....	14.5	17.9	15.2	16.9	16.5	20.8	28.2	48.8	76.8	125.8
Mexican, both sexes.....	34.6	36.2	35.0	35.9	33.7	40.7	39.9	60.5	104.3	148.7
Male.....	51.1	53.8	53.9	53.4	49.8	53.6	58.3	85.7	*	*
Female.....	15.2	17.2	14.2	17.7	18.6	28.0	24.8	42.8	80.0	122.4
Puerto Rican, both sexes.....	50.9	60.0	38.3	31.4	*	*	*	*	*	*
Male.....	88.2	99.4	71.7	*	*	*	*	*	*	*
Female.....	*	28.6	*	*	*	*	*	*	*	*
Cuban, both sexes.....	22.6	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	31.8	28.6	32.9	32.0	29.6	25.3	33.2	73.1	88.8	154.7
Male.....	51.1	42.5	49.6	49.7	46.2	*	*	*	*	*
Female.....	11.7	14.5	18.8	17.7	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	44.5	48.2	44.4	37.3	35.1	39.9	51.1	82.3	138.1	281.3
Male.....	61.9	68.2	63.6	52.8	49.4	55.2	66.9	108.5	179.5	345.9
Female.....	27.4	28.9	26.1	22.6	22.1	26.4	38.2	63.1	112.4	251.3
Non-Hispanic white 6/, both sexes....	46.4	48.7	44.0	36.2	34.3	39.7	51.5	84.1	143.4	295.9
Male.....	63.9	67.9	61.5	50.3	47.4	54.6	66.6	109.3	186.3	362.3
Female.....	29.0	29.7	26.9	22.7	22.1	26.4	39.0	65.3	116.4	264.9
Non-Hispanic black 6/, both sexes....	45.4	55.1	57.6	52.4	45.0	46.1	52.0	72.5	95.5	153.2
Male.....	67.2	83.8	93.3	83.8	73.5	70.6	78.2	110.9	127.1	202.5
Female.....	26.2	30.1	27.5	26.6	22.9	28.0	34.1	49.5	78.9	133.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Transport accidents (V01-V99,Y85)

All origins 3/, both sexes.....	16.8	17.2	16.4	15.6	16.1	16.2	17.5	21.6	26.3	24.4
Male.....	24.2	25.7	24.5	22.9	22.9	22.9	23.9	30.1	37.5	42.9
Female.....	9.4	8.9	8.6	8.6	9.9	10.2	12.3	15.3	19.3	15.7
Hispanic, both sexes.....	15.9	15.7	15.6	15.2	15.8	17.6	17.0	22.9	17.3	11.0
Male.....	23.3	22.7	23.4	22.1	23.5	24.0	24.4	35.0	20.5	*
Female.....	7.6	8.4	7.7	8.7	8.9	12.3	11.3	14.2	15.3	*
Mexican, both sexes.....	17.5	17.9	16.7	16.8	16.4	20.9	16.4	21.1	21.0	*
Male.....	24.6	25.9	24.9	23.9	24.6	25.3	26.5	35.6	*	*
Female.....	9.3	9.1	7.7	9.3	8.8	16.5	*	*	*	*
Puerto Rican, both sexes.....	12.1	12.6	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	14.0	11.6	14.7	12.3	14.8	13.8	*	*	*	*
Male.....	21.9	16.2	20.6	17.5	*	*	*	*	*	*
Female.....	5.9	7.0	9.8	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	16.8	17.3	16.4	15.5	16.1	16.0	17.5	21.4	26.7	25.0
Male.....	24.2	25.9	24.5	22.9	22.7	22.8	23.8	29.7	38.3	44.2
Female.....	9.6	8.9	8.6	8.6	10.0	10.0	12.3	15.3	19.5	16.1
Non-Hispanic white 6/, both sexes....	17.3	17.3	16.7	15.3	15.8	16.2	17.7	21.5	27.2	26.0
Male.....	24.8	25.7	24.7	22.5	22.2	22.8	23.7	29.4	39.2	45.4
Female.....	9.9	8.9	8.9	8.4	9.9	10.2	12.7	15.6	19.7	17.0
Non-Hispanic black 6/, both sexes....	17.5	19.2	17.1	19.4	18.8	15.7	15.6	20.0	20.4	15.9
Male.....	26.8	30.4	28.1	30.0	30.1	25.6	25.6	33.6	28.8	37.6
Female.....	9.4	9.5	7.8	10.6	9.9	8.4	8.7	11.8	15.9	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8, V88.0-V88.8,V89.0,V89.2)

All origins 3/, both sexes.....	15.4	15.7	14.8	14.0	14.7	14.9	16.1	20.2	25.0	23.4
Male.....	21.9	23.2	22.0	20.5	20.4	20.8	21.5	27.8	35.0	41.0
Female.....	8.9	8.4	8.0	7.9	9.4	9.6	11.7	14.6	18.7	15.2
Hispanic, both sexes.....	14.6	14.7	14.6	14.5	15.0	16.6	16.4	21.2	16.7	10.6
Male.....	21.1	20.9	21.8	21.1	22.1	22.3	23.7	31.9	18.8	*
Female.....	7.5	8.1	7.4	8.3	8.7	11.8	10.7	13.4	15.3	*
Mexican, both sexes.....	16.2	17.0	15.8	16.1	15.7	19.4	15.8	19.9	19.8	*
Male.....	22.2	24.3	23.3	23.0	23.2	23.3	25.8	*	*	*
Female.....	9.1	9.0	7.5	8.8	8.8	15.5	*	*	*	*
Puerto Rican, both sexes.....	10.9	12.2	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	12.8	10.1	13.7	11.6	14.4	13.3	*	*	*	*
Male.....	19.8	13.8	18.8	16.4	*	*	*	*	*	*
Female.....	5.6	*	9.4	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	15.4	15.8	14.8	14.0	14.6	14.7	16.0	20.1	25.3	24.1
Male.....	21.9	23.4	21.9	20.4	20.2	20.6	21.3	27.5	35.7	42.3
Female.....	9.0	8.4	8.0	7.9	9.4	9.4	11.8	14.7	18.9	15.6
Non-Hispanic white 6/, both sexes....	15.8	15.7	15.1	13.7	14.2	14.7	16.2	20.2	25.8	25.1
Male.....	22.3	23.1	22.1	20.0	19.6	20.6	21.1	27.0	36.4	43.6
Female.....	9.2	8.3	8.2	7.6	9.2	9.5	12.0	15.0	19.1	16.5
Non-Hispanic black 6/, both sexes....	16.3	17.6	15.6	17.8	18.0	14.8	14.9	18.8	19.4	14.3
Male.....	24.9	27.4	25.2	27.1	28.3	23.9	24.1	31.8	28.1	33.9
Female.....	8.8	9.1	7.6	10.0	9.9	8.0	8.7	11.0	14.8	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nontransport accidents (W00-X59,Y86)

All origins 3/, both sexes.....	26.5	29.9	27.4	21.6	18.9	23.4	33.0	59.8	109.7	250.5
Male.....	36.6	41.1	38.5	30.0	26.8	32.2	42.5	77.2	139.0	294.3
Female.....	16.4	18.9	16.7	13.6	11.8	15.8	25.3	47.0	91.4	230.1
Hispanic, both sexes.....	18.7	20.5	19.3	18.9	16.2	17.4	23.9	40.9	74.3	131.9
Male.....	29.4	30.9	31.2	30.3	25.8	27.9	32.8	49.4	93.4	155.6
Female.....	6.9	9.5	7.4	8.2	7.6	8.5	16.9	34.7	61.6	119.2
Mexican, both sexes.....	17.1	18.4	18.3	19.1	17.3	19.9	23.5	39.4	83.3	137.4
Male.....	26.6	27.9	29.0	29.5	25.3	28.3	31.8	50.1	*	*
Female.....	6.0	8.1	6.5	8.4	9.8	*	16.7	31.9	66.2	116.8
Puerto Rican, both sexes.....	38.8	47.4	31.3	20.7	*	*	*	*	*	*
Male.....	65.5	80.4	58.3	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	17.7	17.0	18.2	19.6	14.8	*	20.8	53.1	*	144.4
Male.....	29.2	26.4	29.1	32.2	*	*	*	*	*	*
Female.....	5.9	7.6	9.0	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	27.6	30.9	28.0	21.7	19.0	23.9	33.6	60.9	111.4	256.3
Male.....	37.7	42.2	39.1	29.9	26.6	32.4	43.1	78.8	141.2	301.7
Female.....	17.8	20.0	17.5	14.0	12.1	16.4	25.9	47.8	92.9	235.2
Non-Hispanic white 6/, both sexes....	29.1	31.5	27.3	20.9	18.5	23.5	33.8	62.6	116.2	269.8
Male.....	39.2	42.2	36.8	27.7	25.3	31.8	42.8	79.9	147.1	316.9
Female.....	19.1	20.9	18.0	14.3	12.2	16.2	26.3	49.7	96.7	247.9
Non-Hispanic black 6/, both sexes....	27.8	35.9	40.6	33.1	26.3	30.4	36.5	52.5	75.2	137.2
Male.....	40.5	53.4	65.3	53.8	43.4	45.0	52.6	77.3	98.3	164.9
Female.....	16.7	20.6	19.7	16.0	12.9	19.6	25.4	37.7	63.0	125.9

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Falls (W00-W19)

All origins 3/, both sexes.....	1.8	2.5	3.5	4.4	6.0	9.5	16.5	33.5	66.2	151.6
Male.....	2.8	4.0	5.5	6.4	8.5	13.3	20.6	42.4	83.7	181.1
Female.....	0.7	1.1	1.5	2.4	3.7	6.2	13.1	27.0	55.3	137.8
Hispanic, both sexes.....	2.1	2.5	4.0	4.3	5.0	8.9	13.0	25.9	47.4	90.4
Male.....	3.6	4.4	7.0	7.1	8.6	14.1	16.4	30.4	59.0	102.6
Female.....	*	*	*	*	*	*	10.4	22.7	39.8	83.9
Mexican, both sexes.....	2.4	2.7	4.2	3.8	5.8	9.1	13.1	25.5	54.3	86.1
Male.....	4.0	4.6	7.0	6.0	9.8	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	19.7	45.4	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	107.6
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	1.7	2.5	3.4	4.4	6.0	9.6	16.7	33.9	67.2	154.6
Male.....	2.7	3.9	5.3	6.3	8.4	13.2	20.9	43.1	85.0	185.4
Female.....	0.7	1.2	1.6	2.5	3.8	6.4	13.3	27.2	56.1	140.2
Non-Hispanic white 6/, both sexes....	1.8	2.6	3.4	4.4	6.1	9.8	17.6	36.0	71.0	164.5
Male.....	2.8	4.0	5.2	6.2	8.4	13.4	21.7	44.7	89.5	196.1
Female.....	0.8	1.2	1.7	2.6	3.9	6.5	14.2	29.4	59.3	149.8
Non-Hispanic black 6/, both sexes....	1.3	2.6	3.7	4.6	5.2	8.9	10.8	16.9	33.7	54.9
Male.....	2.2	4.4	6.5	7.7	8.4	12.8	16.8	26.2	47.1	68.7
Female.....	*	*	*	2.0	*	6.1	6.7	11.3	26.7	49.3

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

All origins 3/, both sexes.....	0.7	1.1	1.1	1.4	1.5	2.1	2.2	3.2	3.8	4.3
Male.....	0.9	1.4	1.5	1.9	1.9	2.7	2.5	4.0	5.1	6.3
Female.....	0.5	0.8	0.8	1.0	1.1	1.5	2.0	2.6	2.9	3.4
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.8	1.1	1.2	1.5	1.5	2.1	2.3	3.3	3.9	4.5
Male.....	1.0	1.4	1.5	2.0	2.0	2.7	2.6	4.1	5.3	6.5
Female.....	0.6	0.9	0.8	1.1	1.0	1.6	2.1	2.7	3.0	3.5
Non-Hispanic white 6/, both sexes....	0.8	1.1	1.1	1.4	1.4	2.0	2.2	2.9	3.4	3.9
Male.....	0.9	1.3	1.5	1.8	1.8	2.4	2.5	3.4	4.9	6.2
Female.....	0.6	0.9	0.8	1.0	1.0	1.6	1.9	2.4	2.5	2.8
Non-Hispanic black 6/, both sexes....	1.0	2.0	1.9	2.8	2.7	4.3	4.0	9.1	10.2	12.5
Male.....	*	2.6	2.7	3.9	4.2	6.8	*	14.7	*	*
Female.....	*	1.4	*	*	*	*	*	5.8	8.9	12.0

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All origins 3/, both sexes.....	19.3	20.7	17.1	9.8	4.6	2.8	2.2	2.4	2.9	3.8
Male.....	25.4	26.9	22.5	12.7	6.1	3.0	2.7	3.0	2.7	4.6
Female.....	13.4	14.7	11.9	7.1	3.4	2.6	1.9	1.9	3.0	3.4
Hispanic, both sexes.....	12.5	13.4	11.0	8.7	4.7	*	*	*	*	*
Male.....	18.6	19.1	16.8	13.9	8.4	*	*	*	*	*
Female.....	5.8	7.4	5.2	3.8	*	*	*	*	*	*
Mexican, both sexes.....	10.7	10.5	9.4	8.1	4.2	*	*	*	*	*
Male.....	15.6	15.0	14.6	12.9	*	*	*	*	*	*
Female.....	4.9	5.6	3.7	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	33.9	39.0	24.3	*	*	*	*	*	*	*
Male.....	55.8	67.2	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	11.0	11.7	10.0	10.8	*	*	*	*	*	*
Male.....	17.0	16.5	13.0	15.8	*	*	*	*	*	*
Female.....	*	7.0	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	20.4	21.6	17.7	9.9	4.6	2.9	2.3	2.4	2.9	3.9
Male.....	26.4	27.8	23.0	12.6	5.7	3.1	2.7	3.0	2.7	4.7
Female.....	14.5	15.6	12.6	7.4	3.5	2.8	2.0	2.0	3.0	3.5
Non-Hispanic white 6/, both sexes....	21.6	22.1	17.1	9.2	4.1	2.6	2.2	2.4	2.9	3.8
Male.....	27.7	27.9	21.4	10.9	4.6	2.5	2.3	2.9	2.8	4.8
Female.....	15.5	16.4	12.9	7.7	3.6	2.7	2.1	2.0	3.0	3.3
Non-Hispanic black 6/, both sexes....	20.7	24.9	27.5	18.1	9.4	5.5	3.6	*	*	5.6
Male.....	28.3	35.6	42.7	30.1	17.2	8.7	6.6	*	*	*
Female.....	14.1	15.7	14.7	8.2	3.4	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All origins 3/, both sexes.....	3.3	4.0	4.5	4.7	5.8	7.7	10.7	18.8	35.2	89.3
Male.....	5.4	6.5	7.1	7.1	8.6	11.2	14.5	24.5	44.8	99.4
Female.....	1.3	1.6	2.0	2.5	3.2	4.7	7.7	14.6	29.2	84.5
Hispanic, both sexes.....	2.7	3.1	3.2	4.3	4.5	5.4	7.7	10.2	21.6	37.2
Male.....	4.8	5.3	5.5	6.6	6.5	8.7	11.1	12.4	27.8	45.1
Female.....	*	*	*	*	*	*	*	8.6	17.4	33.0
Mexican, both sexes.....	2.9	3.6	3.2	6.1	4.9	6.8	7.4	*	25.3	44.4
Male.....	5.0	6.1	5.2	8.7	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	3.0	*	*	*	*	*	*	*	*	*
Male.....	5.7	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.4	4.1	4.6	4.7	5.9	7.9	10.9	19.3	35.9	91.8
Male.....	5.5	6.6	7.2	7.1	8.8	11.4	14.7	25.3	45.5	102.2
Female.....	1.4	1.7	2.1	2.6	3.2	4.9	7.9	15.0	29.9	87.0
Non-Hispanic white 6/, both sexes....	3.5	4.1	4.5	4.7	5.8	7.8	10.6	19.5	37.2	96.0
Male.....	5.5	6.5	6.8	6.9	8.7	11.2	14.3	25.5	47.1	106.7
Female.....	1.4	1.7	2.1	2.5	3.1	4.7	7.6	14.9	31.0	91.0
Non-Hispanic black 6/, both sexes....	3.8	5.1	6.1	6.3	7.7	10.3	16.2	22.4	28.4	62.6
Male.....	6.5	8.5	10.8	9.6	11.3	14.7	22.8	29.5	35.8	76.0
Female.....	1.4	2.1	2.1	3.5	4.8	6.9	11.7	18.1	24.4	57.2

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)

All origins 3/, both sexes.....	15.5	17.1	17.3	15.3	13.4	12.6	12.6	15.6	16.3	15.9
Male.....	23.6	26.0	26.4	23.9	21.1	22.0	23.6	31.6	35.7	43.2
Female.....	7.4	8.4	8.4	7.2	6.3	4.4	3.7	3.8	4.2	3.1
Hispanic, both sexes.....	6.9	7.2	5.7	5.4	6.9	4.8	5.7	6.7	8.8	*
Male.....	11.0	11.1	8.9	9.5	12.1	8.2	10.3	14.4	18.8	*
Female.....	2.3	3.0	2.4	*	*	*	*	*	*	*
Mexican, both sexes.....	6.4	6.0	3.7	4.1	5.4	*	*	*	*	*
Male.....	10.2	9.5	5.9	6.9	8.1	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	10.1	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	7.3	7.5	7.4	5.8	9.8	*	*	*	*	*
Male.....	12.7	11.1	11.6	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	16.8	18.3	18.4	16.2	13.9	13.1	13.1	16.1	16.7	16.3
Male.....	25.7	27.8	28.2	25.1	21.8	22.9	24.6	32.6	36.7	44.4
Female.....	8.1	9.0	9.0	7.7	6.7	4.5	3.8	4.0	4.3	3.2
Non-Hispanic white 6/, both sexes....	19.4	21.0	21.0	18.3	15.5	14.5	14.5	17.6	18.1	17.4
Male.....	29.2	31.6	31.8	27.9	24.0	25.1	27.0	35.3	39.8	47.6
Female.....	9.6	10.5	10.5	8.9	7.5	5.0	4.0	4.3	4.4	3.4
Non-Hispanic black 6/, both sexes....	6.5	6.2	5.9	5.0	4.7	4.0	4.4	4.5	*	5.8
Male.....	11.5	10.9	10.1	9.5	8.4	8.5	9.2	11.5	*	*
Female.....	2.2	2.1	2.3	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (*U03,X60-X71,X75-X84,Y87.0)

All origins 3/, both sexes.....	8.7	9.3	9.0	7.2	5.3	4.0	3.2	3.8	4.1	5.0
Male.....	12.4	12.9	12.2	9.6	6.8	5.7	4.4	5.3	6.0	10.5
Female.....	5.1	5.7	5.9	4.9	4.0	2.6	2.3	2.7	2.9	2.4
Hispanic, both sexes.....	4.6	5.1	3.5	3.6	4.1	*	4.0	*	*	*
Male.....	6.9	7.7	5.1	6.2	6.3	*	*	*	*	*
Female.....	2.0	2.3	*	*	*	*	*	*	*	*
Mexican, both sexes.....	4.0	3.9	1.9	*	*	*	*	*	*	*
Male.....	6.3	6.2	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	4.8	5.6	4.7	*	*	*	*	*	*	*
Male.....	7.8	8.1	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	9.4	9.7	9.5	7.5	5.4	4.1	3.2	3.8	4.2	5.1
Male.....	13.3	13.5	12.9	9.9	6.8	5.9	4.3	5.1	6.1	10.6
Female.....	5.5	6.1	6.3	5.3	4.1	2.6	2.3	2.8	3.0	2.5
Non-Hispanic white 6/, both sexes....	10.7	11.0	10.7	8.3	5.9	4.4	3.2	4.0	4.3	5.4
Male.....	15.0	14.9	14.2	10.7	7.3	6.0	4.5	5.3	6.5	11.2
Female.....	6.4	7.1	7.2	6.0	4.6	2.9	2.2	3.0	3.0	2.7
Non-Hispanic black 6/, both sexes....	3.5	3.6	3.4	2.9	1.6	*	*	*	*	*
Male.....	5.7	6.1	5.4	5.1	*	*	*	*	*	*
Female.....	1.7	1.4	1.7	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)

All origins 3/, both sexes.....	6.3	5.6	4.5	3.4	2.9	2.0	2.2	2.0	2.3	1.9
Male.....	9.4	8.5	6.7	5.2	4.4	2.9	3.2	2.9	2.3	2.3
Female.....	3.3	2.8	2.5	1.7	1.6	1.3	1.4	1.4	2.3	1.8
Hispanic, both sexes.....	6.8	6.4	5.1	3.5	3.9	2.6	*	*	*	*
Male.....	10.4	10.3	7.6	6.3	7.4	*	*	*	*	*
Female.....	2.8	2.3	2.5	*	*	*	*	*	*	*
Mexican, both sexes.....	7.8	7.1	6.0	3.2	3.7	*	*	*	*	*
Male.....	12.3	11.5	8.7	5.5	*	*	*	*	*	*
Female.....	2.5	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.0	5.3	*	*	*	*	*	*	*	*
Male.....	7.5	8.1	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	6.2	5.5	4.4	3.4	2.8	2.0	2.2	2.0	2.3	1.9
Male.....	9.1	8.2	6.5	5.0	4.1	2.9	3.1	2.9	2.3	2.1
Female.....	3.3	2.8	2.5	1.8	1.7	1.2	1.5	1.4	2.4	1.8
Non-Hispanic white 6/, both sexes....	3.7	3.5	3.0	2.4	2.2	1.4	1.9	1.7	2.3	1.7
Male.....	5.1	4.9	4.1	3.2	3.1	1.9	2.3	2.3	2.1	1.8
Female.....	2.3	2.1	1.9	1.6	1.4	1.1	1.5	1.3	2.4	1.7
Non-Hispanic black 6/, both sexes....	21.2	17.5	14.7	10.9	8.0	6.6	5.4	5.2	*	*
Male.....	34.7	30.1	24.8	19.7	13.5	12.3	10.4	10.6	*	*
Female.....	9.3	6.5	6.2	3.7	3.7	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (*U01.4,X93-X95)

All origins 3/, both sexes.....	3.9	3.1	2.4	1.6	1.4	0.9	0.8	0.7	0.6	0.4
Male.....	6.2	4.9	3.6	2.3	2.1	1.2	1.3	1.0	*	*
Female.....	1.6	1.3	1.2	0.9	0.7	0.6	0.4	*	0.7	*
Hispanic, both sexes.....	4.4	3.5	2.4	*	2.1	*	*	*	*	*
Male.....	6.9	5.7	4.0	*	4.1	*	*	*	*	*
Female.....	1.5	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	5.3	4.1	2.8	*	*	*	*	*	*	*
Male.....	8.5	6.8	4.4	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	2.9	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.8	3.0	2.3	1.6	1.3	0.9	0.8	0.7	0.7	0.4
Male.....	6.0	4.8	3.5	2.3	2.0	1.2	1.3	1.0	*	*
Female.....	1.7	1.3	1.2	0.9	0.7	0.6	0.5	*	0.8	*
Non-Hispanic white 6/, both sexes....	2.1	1.8	1.4	1.2	1.0	0.6	0.7	0.5	0.7	*
Male.....	3.1	2.4	2.0	1.5	1.4	0.6	0.9	*	*	*
Female.....	1.1	1.2	0.9	0.8	0.7	0.6	*	*	0.8	*
Non-Hispanic black 6/, both sexes....	14.2	10.5	8.5	4.7	3.7	3.3	*	*	*	*
Male.....	24.8	20.1	15.2	8.7	6.9	6.3	*	*	*	*
Female.....	4.8	2.2	2.9	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (*U01.0-*U01.3,*U01.5-*U01.9,*U02,X85-X92,X96-Y09,Y87.1)

All origins 3/, both sexes.....	2.4	2.5	2.2	1.8	1.6	1.2	1.4	1.4	1.7	1.5
Male.....	3.2	3.6	3.1	2.9	2.3	1.7	1.9	2.0	1.8	1.8
Female.....	1.6	1.5	1.3	0.9	0.9	0.7	1.0	0.9	1.6	1.4
Hispanic, both sexes.....	2.4	2.9	2.6	2.3	*	*	*	*	*	*
Male.....	3.5	4.7	3.6	4.4	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	2.4	3.0	3.2	*	*	*	*	*	*	*
Male.....	3.8	4.7	4.3	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	3.3	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	2.4	2.5	2.1	1.8	1.5	1.1	1.4	1.4	1.7	1.5
Male.....	3.1	3.5	3.0	2.7	2.2	1.7	1.8	2.0	1.8	1.7
Female.....	1.7	1.5	1.2	0.9	0.9	0.6	1.0	0.9	1.6	1.4
Non-Hispanic white 6/, both sexes....	1.6	1.7	1.5	1.2	1.2	0.8	1.2	1.2	1.6	1.4
Male.....	1.9	2.5	2.1	1.6	1.7	1.2	1.4	1.7	1.6	1.6
Female.....	1.2	1.0	0.9	0.7	0.7	0.5	1.1	0.8	1.5	1.3
Non-Hispanic black 6/, both sexes....	7.0	7.0	6.2	6.3	4.3	3.3	3.1	*	*	*
Male.....	9.9	10.0	9.6	11.0	6.6	6.0	*	*	*	*
Female.....	4.4	4.3	3.3	2.3	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All origins 3/, both sexes.....	3.2	3.5	2.8	1.8	1.2	0.9	0.8	1.0	1.1	1.4
Male.....	3.8	4.2	3.5	2.3	1.4	1.2	0.9	1.2	1.7	2.2
Female.....	2.6	2.7	2.0	1.5	0.9	0.7	0.7	0.8	0.8	1.0
Hispanic, both sexes.....	1.3	1.6	1.3	*	*	*	*	*	*	*
Male.....	1.9	2.1	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	1.2	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.5	3.7	2.9	1.9	1.2	0.9	0.9	1.0	1.2	1.3
Male.....	4.2	4.5	3.7	2.3	1.4	1.2	0.9	1.3	1.8	2.2
Female.....	2.9	3.0	2.1	1.5	1.0	0.7	0.8	0.8	0.8	1.0
Non-Hispanic white 6/, both sexes....	3.7	3.8	2.8	1.8	1.2	0.9	0.7	0.8	1.1	1.1
Male.....	4.2	4.3	3.5	2.0	1.3	1.1	0.7	0.8	1.7	1.9
Female.....	3.2	3.2	2.2	1.5	1.0	0.8	0.8	0.8	0.8	0.7
Non-Hispanic black 6/, both sexes....	3.7	4.3	3.8	3.1	*	*	*	*	*	*
Male.....	5.6	6.6	6.2	4.7	*	*	*	*	*	*
Female.....	2.1	2.3	1.8	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All origins 3/, both sexes.....	3.2	3.4	2.7	1.8	1.1	0.9	0.8	0.9	1.1	1.4
Male.....	3.8	4.2	3.5	2.2	1.3	1.1	0.8	1.2	1.7	2.2
Female.....	2.6	2.7	2.0	1.4	0.9	0.6	0.7	0.8	0.8	1.0
Hispanic, both sexes.....	1.2	1.6	1.3	*	*	*	*	*	*	*
Male.....	1.8	2.1	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	1.1	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	3.5	3.7	2.9	1.8	1.1	0.9	0.8	1.0	1.2	1.3
Male.....	4.1	4.4	3.7	2.2	1.3	1.1	0.9	1.2	1.8	2.2
Female.....	2.9	2.9	2.1	1.5	0.9	0.7	0.8	0.8	0.8	1.0
Non-Hispanic white 6/, both sexes....	3.6	3.7	2.8	1.7	1.1	0.9	0.7	0.8	1.1	1.1
Male.....	4.1	4.2	3.4	1.9	1.2	1.0	0.7	0.8	1.6	1.9
Female.....	3.2	3.2	2.2	1.5	1.0	0.8	0.7	0.8	0.8	0.7
Non-Hispanic black 6/, both sexes....	3.6	4.3	3.7	3.1	*	*	*	*	*	*
Male.....	5.5	6.6	6.0	4.7	*	*	*	*	*	*
Female.....	2.0	2.3	1.8	*	*	*	*	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

All origins 3/, both sexes.....	0.4	0.5	0.8	0.9	1.3	2.0	3.1	4.7	6.1	8.3
Male.....	0.3	0.5	0.9	1.0	1.2	2.0	2.9	5.3	7.1	8.8
Female.....	0.4	0.5	0.7	0.9	1.4	2.0	3.3	4.2	5.4	8.1
Hispanic, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Mexican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	0.4	0.5	0.8	1.0	1.4	2.0	3.1	4.9	6.1	8.6
Male.....	0.3	0.5	0.9	1.0	1.3	2.0	2.9	5.4	7.2	9.1
Female.....	0.4	0.6	0.7	0.9	1.5	2.0	3.3	4.4	5.4	8.3
Non-Hispanic white 6/, both sexes....	0.3	0.5	0.7	0.8	1.3	1.9	3.1	4.6	5.9	8.5
Male.....	0.4	0.4	0.9	0.9	1.2	1.9	2.8	5.3	7.3	9.1
Female.....	0.3	0.5	0.5	0.7	1.3	1.8	3.3	4.2	5.0	8.2
Non-Hispanic black 6/, both sexes....	0.7	1.0	1.7	2.4	3.2	4.0	4.3	8.4	9.7	10.4
Male.....	*	*	*	*	*	*	*	9.7	*	*
Female.....	*	*	1.9	2.6	3.0	4.3	*	7.7	9.6	10.1

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 7/

All origins 3/, both sexes.....	10.1	15.7	19.7	19.9	18.6	16.0	13.2	11.0	8.1	4.9
Male.....	14.1	22.8	30.6	31.8	30.1	27.1	22.2	19.1	14.1	9.5
Female.....	6.2	8.7	9.1	8.7	8.2	6.4	5.9	5.0	4.3	2.7
Hispanic, both sexes.....	11.2	17.6	23.6	26.5	25.9	23.0	19.9	20.5	16.7	9.4
Male.....	17.2	28.6	40.0	46.2	45.8	42.3	40.1	39.1	27.0	*
Female.....	4.5	6.1	7.3	7.9	8.2	6.9	*	*	*	*
Mexican, both sexes.....	12.0	19.8	26.0	29.6	27.4	28.4	24.4	23.1	21.6	*
Male.....	18.1	32.2	43.3	49.5	47.7	48.6	47.7	45.3	*	*
Female.....	4.8	6.3	7.0	8.8	8.5	*	*	*	*	*
Puerto Rican, both sexes.....	9.8	16.8	17.2	28.4	41.5	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	11.5	14.9	21.5	22.4	20.3	20.0	*	*	*	*
Male.....	18.1	23.4	37.5	43.0	35.8	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	9.9	15.4	19.2	19.3	18.0	15.4	12.6	10.3	7.6	4.6
Male.....	13.5	21.9	29.5	30.4	28.6	25.6	20.8	17.7	13.4	8.8
Female.....	6.4	9.0	9.3	8.8	8.2	6.3	6.0	4.9	4.0	2.7
Non-Hispanic white 6/, both sexes....	10.3	16.0	19.7	19.3	18.2	15.3	12.7	10.6	7.8	4.7
Male.....	14.0	22.7	29.9	29.9	28.7	25.2	20.5	18.2	13.7	8.9
Female.....	6.6	9.4	9.6	9.0	8.4	6.5	6.2	5.0	4.1	2.7
Non-Hispanic black 6/, both sexes....	7.4	12.1	18.3	21.5	19.6	17.5	13.5	9.3	6.5	*
Male.....	9.9	18.2	29.6	37.7	34.5	33.6	25.6	18.0	*	*
Female.....	5.1	6.8	8.8	8.1	8.1	5.7	5.2	*	*	*

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,K85.3,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 7/

All origins 3/, both sexes.....	26.2	28.4	24.8	15.2	8.1	5.7	4.9	5.5	6.8	8.1
Male.....	32.9	35.0	31.3	18.7	9.5	6.0	5.5	6.2	8.0	10.7
Female.....	19.6	21.9	18.6	12.0	6.9	5.4	4.3	5.0	6.0	6.8
Hispanic, both sexes.....	15.7	17.2	15.2	11.1	6.0	3.2	*	*	*	*
Male.....	22.7	23.8	22.3	17.7	9.6	*	*	*	*	*
Female.....	7.9	10.2	8.0	4.9	*	*	*	*	*	*
Mexican, both sexes.....	13.6	13.7	13.0	10.4	5.4	*	*	*	*	*
Male.....	19.1	18.3	19.6	16.3	*	*	*	*	*	*
Female.....	7.1	8.6	5.6	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	43.3	47.8	32.9	*	*	*	*	*	*	*
Male.....	71.2	80.4	58.3	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	13.2	14.6	14.1	12.6	*	*	*	*	*	*
Male.....	20.0	20.7	17.4	19.2	*	*	*	*	*	*
Female.....	6.1	8.5	11.3	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	27.7	29.6	25.7	15.6	8.2	5.9	5.0	5.6	6.9	8.2
Male.....	34.4	36.3	32.1	18.7	9.3	6.1	5.6	6.3	8.2	10.8
Female.....	21.3	23.2	19.6	12.6	7.3	5.7	4.5	5.1	6.0	7.0
Non-Hispanic white 6/, both sexes....	29.5	30.4	25.2	14.9	7.7	5.8	4.8	5.8	7.1	8.4
Male.....	36.1	36.2	30.1	16.6	7.8	5.5	4.9	6.4	8.7	11.3
Female.....	23.0	24.7	20.4	13.3	7.7	6.1	4.8	5.4	6.1	7.1
Non-Hispanic black 6/, both sexes....	27.9	34.4	37.6	25.8	15.1	7.6	6.4	4.5	*	6.6
Male.....	37.9	48.4	57.9	42.1	26.1	13.5	12.0	*	*	*
Female.....	19.2	22.3	20.4	12.4	6.5	*	*	*	*	7.5

Worktable Orig210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, specified Hispanic origin, race for non-Hispanic population, and sex: United States, 2006

[Rates per 100,000 population in specified group. Rates based on populations estimated as of July 1, 2006. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated separately by race; data for non-Hispanic persons are tabulated by race. Data for Hispanic origin should be interpreted with caution because of inconsistencies between reporting Hispanic origin on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004),

Hispanic origin, race for non-Hispanic population, and sex	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 7/

All origins 3/, both sexes.....	11.1	11.3	11.0	9.9	9.7	9.7	10.4	12.8	13.1	11.5
Male.....	18.3	18.7	18.4	17.0	17.0	18.0	20.8	28.0	30.7	33.6
Female.....	4.1	4.1	3.9	3.2	3.1	2.4	1.9	1.6	2.1	1.1
Hispanic, both sexes.....	7.2	5.8	4.9	3.2	5.1	3.1	*	*	6.5	*
Male.....	11.9	9.5	8.5	5.6	10.2	5.6	*	*	*	*
Female.....	2.0	1.9	*	*	*	*	*	*	*	*
Mexican, both sexes.....	8.3	6.4	4.8	3.8	4.7	*	*	*	*	*
Male.....	13.5	10.5	7.9	6.4	9.1	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Puerto Rican, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Cuban, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Other Hispanic 4/, both sexes.....	5.5	4.1	5.1	*	*	*	*	*	*	*
Male.....	9.4	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Non-Hispanic 5/, both sexes.....	11.7	12.0	11.6	10.5	10.1	10.2	11.0	13.3	13.5	11.8
Male.....	19.3	19.8	19.4	18.0	17.5	18.9	22.0	29.2	31.6	34.7
Female.....	4.3	4.4	4.1	3.4	3.3	2.5	2.1	1.7	2.1	1.2
Non-Hispanic white 6/, both sexes....	11.3	12.2	12.1	11.5	10.9	11.1	12.1	14.5	14.7	12.6
Male.....	18.2	19.8	20.1	19.3	18.7	20.4	23.9	31.3	34.4	37.0
Female.....	4.4	4.8	4.3	3.9	3.6	2.8	2.3	1.9	2.4	1.1
Non-Hispanic black 6/, both sexes....	17.7	13.5	11.5	6.8	6.9	6.4	5.4	6.0	*	6.4
Male.....	31.7	25.6	20.8	13.3	13.5	13.1	11.7	15.7	*	*
Female.....	5.4	3.0	3.7	*	*	*	*	*	*	*

* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Figures for origin not stated are included in "All origins" but not distributed among specified origins.

4/ Includes Central and South American and Other and unknown Hispanic.

5/ Includes races other than white and black.

6/ Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2006, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality."

7/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.