

Person with disabilities experiencing problems accessing  
NCHS birth and death tables should contact  
[nchsed@cdc.gov](mailto:nchsed@cdc.gov), or call 301-458-4688

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All causes

All races 3/, both sexes

2006.....	810.4	690.7	28.4	15.2	82.2	106.3	190.2	427.5	890.9	2,062.1	5,115.0	13,253.1
2005.....	825.9	692.5	29.4	16.3	81.4	104.4	193.3	432.0	906.9	2,137.1	5,260.0	13,798.6
2004.....	816.5	685.2	29.9	16.8	80.1	102.1	193.5	427.0	910.3	2,164.6	5,275.1	13,823.5
2003.....	841.9	700.0	31.5	17.0	81.5	103.6	201.6	433.2	940.9	2,255.0	5,463.1	14,593.3
2002.....	847.3	695.0	31.2	17.4	81.4	103.6	202.9	430.1	952.4	2,314.7	5,556.9	14,828.3
2001.....	848.5	683.4	33.3	17.3	80.7	105.2	203.6	428.9	964.6	2,353.3	5,582.4	15,112.8
2000.....	854.0	736.7	32.4	18.0	79.9	101.4	198.9	425.6	992.2	2,399.1	5,666.5	15,524.4
1999.....	857.0	736.0	34.2	18.6	79.3	102.2	198.0	418.2	1,005.0	2,457.3	5,714.5	15,554.6

All races 3/, male

2006.....	814.8	756.3	30.5	17.6	119.3	146.8	238.7	541.0	1,110.0	2,516.2	6,177.7	14,309.1
2005.....	827.2	762.3	33.4	18.6	117.8	143.4	243.0	547.8	1,131.0	2,612.2	6,349.8	14,889.4
2004.....	817.6	753.7	32.4	19.2	114.7	139.5	243.6	543.5	1,128.8	2,644.8	6,394.3	15,031.1
2003.....	840.3	777.4	35.1	19.8	116.5	141.4	255.0	552.2	1,165.5	2,771.7	6,641.8	15,794.0
2002.....	846.6	761.5	35.2	20.0	117.3	142.2	257.5	547.5	1,184.0	2,855.3	6,760.5	16,254.5
2001.....	846.4	749.8	37.0	19.8	117.0	143.7	259.6	545.1	1,192.7	2,911.5	6,833.0	16,744.8
2000.....	853.0	806.5	35.9	20.9	114.9	138.6	255.2	542.8	1,230.7	2,979.6	6,972.6	17,501.4
1999.....	859.2	805.0	37.9	21.5	113.1	139.7	254.9	533.1	1,252.0	3,073.7	7,083.3	17,597.2

All races 3/, female

2006.....	806.1	622.0	26.3	12.8	42.8	64.3	141.6	317.7	687.0	1,677.9	4,388.3	12,759.0
2005.....	824.6	619.4	25.1	13.9	42.7	64.1	143.6	319.9	698.5	1,736.3	4,520.0	13,297.7
2004.....	815.4	613.4	27.3	14.3	43.6	63.5	143.5	314.3	707.4	1,761.4	4,521.8	13,280.3
2003.....	843.4	619.1	27.8	14.0	44.4	64.6	148.5	318.4	732.7	1,823.0	4,675.5	14,062.5
2002.....	848.0	625.3	27.0	14.7	43.7	64.0	148.8	316.9	738.0	1,864.7	4,757.9	14,209.6
2001.....	850.4	613.9	29.5	14.6	42.6	66.0	148.2	316.8	754.0	1,890.8	4,760.5	14,429.9
2000.....	855.0	663.4	28.7	15.0	43.1	63.5	143.2	312.5	772.2	1,921.2	4,814.7	14,719.2
1999.....	854.9	663.6	30.3	15.6	43.7	64.1	141.8	307.6	777.6	1,952.3	4,825.4	14,731.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

All causes

White, both sexes

2006 4/.....	858.1	576.0	25.5	14.2	78.0	98.3	177.3	397.9	847.5	2,021.1	5,127.4	13,480.4
2005 4/.....	873.7	579.1	27.0	15.0	77.1	95.5	180.1	399.6	860.7	2,094.5	5,263.2	14,021.1
2004 4/.....	863.2	574.0	27.0	15.5	76.4	92.9	179.8	394.0	863.5	2,120.4	5,272.8	14,010.1
2003 4/.....	890.1	591.3	28.5	15.8	77.1	94.3	185.7	397.5	892.2	2,207.0	5,457.8	14,792.3
2002.....	895.7	586.7	28.1	16.1	77.1	93.5	186.6	395.3	902.5	2,267.6	5,544.2	15,015.9
2001.....	895.1	570.9	30.7	16.2	75.5	95.3	185.6	393.2	913.5	2,298.9	5,560.9	15,287.5
2000.....	900.2	610.6	29.2	17.0	74.5	90.3	179.9	388.3	940.2	2,341.5	5,637.7	15,707.0
1999.....	901.4	606.1	30.8	17.2	73.2	91.4	177.3	380.4	948.1	2,396.2	5,679.5	15,718.0

White, male

2006 4/.....	852.3	632.7	27.5	16.4	111.8	135.4	224.4	505.2	1,050.6	2,455.8	6,182.2	14,576.8
2005 4/.....	864.5	640.0	30.9	17.1	110.4	130.8	228.5	509.3	1,068.1	2,552.7	6,343.2	15,156.5
2004 4/.....	854.2	631.6	29.4	17.9	108.3	127.0	229.1	504.0	1,065.9	2,584.0	6,384.8	15,250.7
2003 4/.....	877.6	658.7	31.5	18.4	108.9	128.8	237.6	509.2	1,102.3	2,707.3	6,620.6	16,037.9
2002.....	884.0	650.9	31.5	18.4	109.7	128.3	239.3	505.4	1,118.6	2,795.4	6,738.8	16,473.2
2001.....	881.9	627.6	34.2	18.4	108.0	130.3	239.7	501.3	1,127.5	2,842.3	6,799.7	16,935.4
2000.....	887.8	667.6	32.6	19.8	105.8	124.1	233.6	496.9	1,163.3	2,905.7	6,933.1	17,716.4
1999.....	892.1	667.0	33.9	19.7	102.8	125.4	230.8	484.6	1,179.7	2,998.7	7,040.1	17,752.9

White, female

2006 4/.....	863.9	516.5	23.5	11.9	41.7	58.9	129.0	291.6	654.6	1,646.0	4,395.1	12,965.7
2005 4/.....	882.8	515.3	22.9	12.8	41.5	58.0	130.4	291.1	663.9	1,700.4	4,519.4	13,498.3
2004 4/.....	871.9	513.6	24.4	13.0	42.4	56.8	129.5	285.4	671.8	1,723.6	4,514.4	13,450.9
2003 4/.....	902.3	520.7	25.5	13.1	43.2	58.0	132.9	287.2	693.7	1,780.6	4,671.6	14,240.6
2002.....	907.0	519.4	24.5	13.7	42.4	56.9	133.2	286.8	698.7	1,819.7	4,742.5	14,382.8
2001.....	907.9	511.5	27.1	13.9	40.8	58.6	131.0	286.8	712.2	1,840.2	4,738.5	14,597.6
2000.....	912.3	550.5	25.5	14.1	41.1	55.1	125.7	281.4	730.9	1,868.3	4,785.3	14,890.7
1999.....	910.4	542.0	27.5	14.6	41.5	56.2	123.2	277.9	731.0	1,893.9	4,787.8	14,900.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

All causes

Black, both sexes

2006 4/.....	733.0	1,303.1	43.3	21.2	112.1	177.7	314.2	720.8	1,440.3	2,835.9	5,886.0	12,489.7
2005 4/.....	749.4	1,311.2	41.8	23.3	112.3	179.2	318.4	744.2	1,483.3	2,928.1	6,134.3	13,083.1
2004 4/.....	744.3	1,284.7	44.8	23.6	109.4	179.3	322.3	744.8	1,499.6	2,982.6	6,170.4	13,338.5
2003 4/.....	763.6	1,273.8	46.8	22.9	113.1	181.6	343.6	771.4	1,549.9	3,108.4	6,353.9	13,976.7
2002.....	768.4	1,263.6	47.1	24.5	113.7	186.8	348.6	765.8	1,565.2	3,159.3	6,518.9	14,260.8
2001.....	773.5	1,304.5	47.5	23.3	117.7	185.0	356.5	771.9	1,576.8	3,257.6	6,585.6	14,550.8
2000.....	781.1	1,426.1	49.9	24.2	119.4	187.8	356.9	786.9	1,607.8	3,335.6	6,701.3	14,714.3
1999.....	788.1	1,457.3	52.6	26.5	121.2	184.4	368.4	782.4	1,655.6	3,413.6	6,753.3	14,757.9

Black, male

2006 4/.....	786.7	1,407.1	47.1	24.8	171.3	254.2	392.3	921.9	1,891.8	3,669.2	7,393.2	13,206.0
2005 4/.....	799.2	1,437.2	46.7	27.0	172.1	254.3	395.5	948.6	1,954.3	3,747.3	7,667.1	13,809.8
2004 4/.....	792.6	1,414.2	48.6	26.0	164.3	252.3	397.0	954.9	1,960.8	3,818.3	7,710.3	14,452.5
2003 4/.....	813.7	1,410.0	53.7	26.8	171.3	256.2	426.7	991.4	2,011.2	3,981.3	8,067.0	14,903.4
2002.....	816.7	1,351.5	54.4	28.9	172.6	264.5	434.7	983.0	2,039.2	4,024.5	8,169.6	15,635.5
2001.....	823.9	1,426.4	53.2	27.7	180.7	259.2	445.0	996.1	2,025.3	4,166.6	8,355.9	16,439.9
2000.....	834.1	1,567.6	54.5	28.2	181.4	261.0	453.0	1,017.7	2,080.1	4,253.5	8,486.0	16,791.0
1999.....	847.4	1,592.8	59.0	32.0	184.6	258.6	469.2	1,030.7	2,145.6	4,352.3	8,559.1	17,304.5

Black, female

2006 4/.....	684.0	1,194.6	39.4	17.4	51.3	106.6	245.0	548.1	1,076.3	2,239.7	5,028.9	12,196.7
2005 4/.....	703.9	1,179.7	36.7	19.4	51.2	109.8	250.0	568.4	1,103.6	2,341.5	5,263.7	12,789.9
2004 4/.....	700.3	1,149.9	40.9	21.1	53.7	112.3	256.0	564.1	1,128.6	2,386.1	5,300.0	12,896.9
2003 4/.....	717.9	1,132.2	39.7	18.9	54.0	113.5	270.0	582.3	1,178.8	2,487.2	5,385.5	13,616.7
2002.....	724.4	1,172.0	39.5	19.9	54.4	116.4	272.3	579.4	1,184.2	2,545.0	5,584.4	13,734.2
2001.....	727.7	1,176.3	41.7	18.7	54.9	117.9	278.0	579.1	1,215.5	2,616.1	5,591.6	13,849.9
2000.....	733.0	1,279.8	45.3	20.0	58.3	121.8	271.9	588.3	1,227.2	2,689.6	5,696.5	13,941.3
1999.....	734.3	1,317.4	46.1	20.9	58.6	117.6	279.4	569.8	1,262.7	2,751.5	5,742.4	13,805.9















Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Certain other intestinal infections (A04,A07-A09)

All races 3/, both sexes

2006.....	2.2	*	*	*	*	*	0.1	0.3	1.1	4.8	19.0	50.5
2005.....	1.9	*	*	*	*	*	0.1	0.2	0.8	4.4	15.9	45.8
2004.....	1.5	*	*	*	*	*	0.1	0.2	0.9	3.3	12.8	34.3
2003.....	1.1	*	*	*	*	*	*	0.2	0.6	2.4	9.3	24.8
2002.....	0.9	*	*	*	*	*	*	0.1	0.5	2.0	7.0	21.0
2001.....	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.5	12.8
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	1.2	3.7	11.8
1999.....	0.4	*	*	*	*	*	*	0.1	0.2	1.0	3.2	10.3

All races 3/, male

2006.....	1.7	*	*	*	*	*	0.1	0.3	1.3	4.8	18.6	50.0
2005.....	1.5	*	*	*	*	*	*	0.3	0.9	4.4	16.4	46.1
2004.....	1.1	*	*	*	*	*	*	0.2	0.9	3.1	12.7	32.0
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	2.2	9.3	21.4
2002.....	0.7	*	*	*	*	*	*	0.1	0.6	2.2	7.0	20.4
2001.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.6	12.1
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.0	11.4
1999.....	0.3	*	*	*	*	*	*	*	0.2	0.9	3.5	9.5

All races 3/, female

2006.....	2.7	*	*	*	*	*	*	0.3	1.0	4.8	19.4	50.8
2005.....	2.3	*	*	*	*	*	*	0.2	0.8	4.5	15.5	45.6
2004.....	1.8	*	*	*	*	*	*	0.2	0.8	3.4	12.9	35.4
2003.....	1.3	*	*	*	*	*	*	0.2	0.6	2.6	9.3	26.3
2002.....	1.0	*	*	*	*	*	*	0.1	0.5	1.8	6.9	21.3
2001.....	0.7	*	*	*	*	*	*	0.1	0.3	1.3	4.5	13.1
2000.....	0.6	*	*	*	*	*	*	*	0.2	1.2	3.6	11.9
1999.....	0.5	*	*	*	*	*	*	*	0.2	1.0	3.0	10.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Certain other intestinal infections (A04,A07-A09)

White, both sexes

2006 4/.....	2.5	*	*	*	*	*	0.1	0.2	1.1	5.0	20.3	53.6
2005 4/.....	2.2	*	*	*	*	*	0.1	0.2	0.8	4.7	16.8	48.6
2004 4/.....	1.7	*	*	*	*	*	0.1	0.2	0.9	3.5	13.5	36.6
2003 4/.....	1.2	*	*	*	*	*	*	0.1	0.6	2.6	9.7	26.1
2002.....	1.0	*	*	*	*	*	*	0.1	0.5	2.1	7.4	22.0
2001.....	0.6	*	*	*	*	*	*	0.1	0.3	1.3	4.7	13.5
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	1.3	4.0	12.2
1999.....	0.4	*	*	*	*	*	*	*	0.2	1.0	3.3	10.9

White, male

2006 4/.....	2.0	*	*	*	*	*	*	0.3	1.2	5.0	19.5	53.1
2005 4/.....	1.7	*	*	*	*	*	*	0.2	0.9	4.7	17.0	49.3
2004 4/.....	1.3	*	*	*	*	*	*	0.2	0.9	3.2	13.4	33.9
2003 4/.....	0.9	*	*	*	*	*	*	0.2	0.6	2.4	9.8	22.7
2002.....	0.8	*	*	*	*	*	*	0.2	0.5	2.3	7.4	21.4
2001.....	0.5	*	*	*	*	*	*	*	0.3	1.3	4.6	12.7
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.3	4.2	11.9
1999.....	0.3	*	*	*	*	*	*	*	0.2	1.0	3.7	9.9

White, female

2006 4/.....	3.1	*	*	*	*	*	*	0.2	1.0	5.0	20.8	53.9
2005 4/.....	2.7	*	*	*	*	*	*	0.2	0.7	4.7	16.6	48.3
2004 4/.....	2.1	*	*	*	*	*	*	0.2	0.9	3.7	13.6	37.8
2003 4/.....	1.5	*	*	*	*	*	*	0.1	0.7	2.8	9.7	27.6
2002.....	1.2	*	*	*	*	*	*	0.1	0.5	1.9	7.4	22.3
2001.....	0.8	*	*	*	*	*	*	0.2	0.3	1.3	4.7	13.9
2000.....	0.6	*	*	*	*	*	*	*	0.3	1.2	3.8	12.3
1999.....	0.5	*	*	*	*	*	*	*	*	1.0	3.1	11.3



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Tuberculosis (A16-A19)

All races 3/, both sexes

2006.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.6	1.3	2.1
2005.....	0.2	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.3	2.2
2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.4
2003.....	0.2	*	*	*	*	*	0.1	0.3	0.4	0.6	1.7	2.3
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.9	1.8	2.6
2001.....	0.3	*	*	*	*	0.1	0.1	0.2	0.5	0.8	1.6	2.8
2000.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.8	2.0	2.7
1999.....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.9	2.1	4.2

All races 3/, male

2006.....	0.3	*	*	*	*	*	0.1	0.4	0.5	0.9	1.8	3.1
2005.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	1.7	3.3
2004.....	0.3	*	*	*	*	*	0.2	0.3	0.4	1.1	1.6	3.8
2003.....	0.3	*	*	*	*	*	0.1	0.4	0.5	0.9	2.3	3.1
2002.....	0.3	*	*	*	*	*	0.2	0.3	0.6	1.2	2.4	4.0
2001.....	0.3	*	*	*	*	*	0.1	0.3	0.7	1.2	2.0	4.4
2000.....	0.3	*	*	*	*	*	0.1	0.4	0.5	1.0	2.8	4.6
1999.....	0.4	*	*	*	*	*	0.2	0.5	0.8	1.4	2.8	7.1

All races 3/, female

2006.....	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.9	1.6
2005.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.7
2004.....	0.2	*	*	*	*	*	*	0.1	*	0.4	1.0	1.7
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.9
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.1
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.7	1.5	2.0
1999.....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.6	3.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Tuberculosis (A16-A19)

White, both sexes

2006 4/.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.6
2005 4/.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	0.8	1.7
2004 4/.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.0	1.7
2003 4/.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.3	1.8
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.4	2.1
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.6	1.2	2.2
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.2
1999.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.7	1.6	3.2

White, male

2006 4/.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.3	2.1
2005 4/.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.4
2004 4/.....	0.2	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.8
2003 4/.....	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.7	2.4
2002.....	0.2	*	*	*	*	*	*	0.2	0.3	0.9	1.8	3.2
2001.....	0.2	*	*	*	*	*	*	0.2	0.5	0.7	1.5	3.1
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.7	1.8	3.8
1999.....	0.3	*	*	*	*	*	*	0.3	0.5	1.0	2.1	5.4

White, female

2006 4/.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.4
2005 4/.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.4
2004 4/.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.3
2003 4/.....	0.2	*	*	*	*	*	*	*	*	0.3	1.1	1.6
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.3	1.2	1.6
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.0	1.8
2000.....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.6
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.2	2.4





Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Respiratory tuberculosis (A16)

All races 3/, both sexes

2006.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.4	1.0	1.6
2005.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.9
2004.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	0.9	2.2
2003.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.5	1.4	1.8
2002.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.5	2.3
2001.....	0.2	*	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4
2000.....	0.2	*	*	*	*	*	0.1	0.2	0.2	0.6	1.7	2.3
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.5	0.8	1.8	3.7

All races 3/, male

2006.....	0.2	*	*	*	*	*	0.1	0.3	0.5	0.7	1.5	2.4
2005.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.7	1.3	2.7
2004.....	0.2	*	*	*	*	*	0.1	0.2	0.3	0.9	1.2	3.7
2003.....	0.2	*	*	*	*	*	*	0.3	0.4	0.7	1.9	2.5
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	1.9	3.5
2001.....	0.3	*	*	*	*	*	*	0.3	0.6	0.9	1.7	3.8
2000.....	0.3	*	*	*	*	*	0.1	0.3	0.4	0.9	2.4	4.1
1999.....	0.4	*	*	*	*	*	0.1	0.4	0.7	1.2	2.5	6.4

All races 3/, female

2006.....	0.1	*	*	*	*	*	*	*	*	0.2	0.6	1.2
2005.....	0.1	*	*	*	*	*	*	*	*	0.2	0.8	1.5
2004.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.5
2003.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	1.0	1.5
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.3	1.8
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.1	1.8
2000.....	0.2	*	*	*	*	*	*	*	*	0.5	1.2	1.6
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.3	2.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Respiratory tuberculosis (A16)

White, both sexes

2006 4/.....	0.1	*	*	*	*	*	*	0.1	0.2	0.3	0.8	1.2
2005 4/.....	0.1	*	*	*	*	*	*	0.1	0.1	0.3	0.6	1.3
2004 4/.....	0.1	*	*	*	*	*	0.1	0.1	0.1	0.4	0.7	1.6
2003 4/.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.4
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.1	1.8
2001.....	0.2	*	*	*	*	*	*	0.1	0.3	0.4	1.0	1.8
2000.....	0.2	*	*	*	*	*	*	0.1	0.1	0.4	1.2	1.9
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.3	3.0

White, male

2006 4/.....	0.2	*	*	*	*	*	*	0.1	0.3	0.4	1.1	1.5
2005 4/.....	0.1	*	*	*	*	*	*	0.2	0.2	0.4	0.7	1.9
2004 4/.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.0	2.6
2003 4/.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.4	1.8
2002.....	0.2	*	*	*	*	*	*	0.1	0.3	0.7	1.4	2.9
2001.....	0.2	*	*	*	*	*	*	0.1	0.5	0.6	1.3	2.6
2000.....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.5	3.4
1999.....	0.3	*	*	*	*	*	*	0.2	0.5	0.8	1.8	5.2

White, female

2006 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.0
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.6	1.1
2004 4/.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	1.1
2003 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.9	1.2
2002.....	0.1	*	*	*	*	*	*	*	*	*	1.0	1.3
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.4
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	1.0	1.2
1999.....	0.2	*	*	*	*	*	*	*	*	0.3	1.0	2.1























Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Meningococcal infection (A39)

All races 3/, both sexes

2006.....	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2004.....	0.0	*	*	*	0.1	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002.....	0.1	*	*	0.0	0.1	*	*	*	*	*	*	*
2001.....	0.1	*	*	0.1	0.1	*	*	*	*	*	*	*
2000.....	0.1	0.6	*	*	0.1	*	*	0.1	*	*	*	*
1999.....	0.1	0.8	0.2	0.1	0.1	0.1	0.1	*	*	*	*	*

All races 3/, male

2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.2	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*
1999.....	0.1	*	*	*	0.1	*	*	*	*	*	*	*





Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Septicemia (A40-A41)

All races 3/, both sexes

2006.....	11.4	6.5	0.5	0.2	0.3	0.7	2.0	5.2	12.8	32.1	82.4	177.3
2005.....	11.5	7.4	0.5	0.2	0.4	0.8	1.9	5.2	12.9	32.6	81.4	187.3
2004.....	11.4	6.6	0.5	0.2	0.3	0.8	1.9	5.4	12.9	32.4	81.6	186.7
2003.....	11.7	6.9	0.5	0.2	0.4	0.8	2.1	5.3	13.1	32.6	85.0	202.5
2002.....	11.7	7.3	0.5	0.2	0.3	0.8	1.9	5.2	12.6	34.7	86.5	203.0
2001.....	11.3	7.7	0.7	0.2	0.3	0.7	1.8	5.0	12.3	32.8	82.3	205.9
2000.....	11.1	7.2	0.6	0.2	0.3	0.7	1.9	4.9	11.9	31.0	80.4	215.7
1999.....	11.0	7.5	0.6	0.2	0.3	0.7	1.8	4.6	11.4	31.2	79.4	220.7

All races 3/, male

2006.....	10.5	6.8	0.5	0.3	0.3	0.7	2.1	5.9	14.1	34.8	92.7	194.8
2005.....	10.5	8.1	0.6	0.2	0.4	0.7	1.9	5.6	14.2	36.3	91.4	201.9
2004.....	10.4	8.0	0.5	0.1	0.3	0.8	2.0	5.9	13.8	35.8	93.3	203.9
2003.....	10.5	7.4	0.7	0.2	0.4	0.8	2.1	5.8	14.1	36.2	95.0	213.8
2002.....	10.6	7.7	0.5	0.2	0.3	0.8	2.0	5.7	13.7	38.0	99.0	213.9
2001.....	10.2	8.4	0.8	0.2	0.4	0.7	2.0	5.4	13.2	36.7	95.5	224.3
2000.....	9.8	7.1	0.7	0.2	0.3	0.8	2.0	5.6	13.0	34.5	91.9	223.0
1999.....	9.8	9.0	0.6	0.2	0.3	0.7	2.1	5.1	12.4	35.2	91.9	233.5

All races 3/, female

2006.....	12.3	6.2	0.6	0.2	0.3	0.7	1.9	4.6	11.5	29.7	75.3	169.2
2005.....	12.5	6.5	0.5	0.2	0.3	0.8	1.9	4.8	11.7	29.5	74.6	180.6
2004.....	12.3	5.2	0.6	0.2	0.3	0.8	1.9	4.9	12.0	29.6	73.7	179.0
2003.....	12.9	6.5	0.4	0.2	0.4	0.8	2.0	4.8	12.2	29.5	78.3	197.5
2002.....	12.9	7.0	0.5	0.2	0.3	0.8	1.8	4.7	11.7	31.9	78.2	198.3
2001.....	12.4	7.0	0.6	0.1	0.3	0.7	1.7	4.7	11.4	29.5	73.7	198.3
2000.....	12.3	7.3	0.6	0.1	0.2	0.6	1.9	4.2	11.0	28.1	72.9	212.7
1999.....	12.2	5.8	0.5	0.2	0.3	0.6	1.6	4.1	10.5	28.0	71.3	215.5



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Septicemia (A40-A41)

White, both sexes

2006 4/.....	11.3	4.9	0.5	0.2	0.3	0.6	1.6	4.2	10.9	28.7	77.8	171.0
2005 4/.....	11.3	5.0	0.5	0.2	0.3	0.6	1.6	4.3	11.1	29.6	76.1	177.3
2004 4/.....	11.2	5.2	0.5	0.2	0.3	0.7	1.6	4.4	11.0	29.1	76.6	178.1
2003 4/.....	11.5	4.6	0.4	0.2	0.3	0.6	1.7	4.3	11.2	28.9	80.1	191.1
2002.....	11.5	6.1	0.5	0.2	0.3	0.6	1.6	4.2	10.6	31.1	80.7	191.9
2001.....	11.1	5.4	0.7	0.2	0.3	0.5	1.4	4.0	10.3	29.3	77.1	195.3
2000.....	10.8	5.7	0.6	0.2	0.2	0.5	1.6	3.9	9.9	27.4	73.9	203.0
1999.....	10.6	5.5	0.5	0.2	0.3	0.6	1.4	3.6	9.4	27.3	72.8	205.9

White, male

2006 4/.....	10.4	5.1	0.4	0.2	0.3	0.6	1.7	4.8	11.9	31.5	88.1	190.7
2005 4/.....	10.2	5.9	0.5	0.2	0.3	0.6	1.5	4.6	12.3	32.6	85.7	190.7
2004 4/.....	10.2	6.3	0.5	0.2	0.3	0.6	1.6	4.8	11.7	32.2	87.6	196.5
2003 4/.....	10.2	4.6	0.6	0.2	0.3	0.6	1.7	4.7	11.9	32.1	89.5	203.3
2002.....	10.3	6.2	0.5	0.3	0.3	0.7	1.6	4.7	11.7	34.1	91.5	202.9
2001.....	9.9	6.2	0.8	0.2	0.3	0.5	1.7	4.3	11.1	32.5	89.4	211.8
2000.....	9.4	5.8	0.6	0.2	0.2	0.6	1.7	4.4	10.9	30.8	84.6	209.8
1999.....	9.3	6.9	0.5	0.2	0.3	0.6	1.6	4.0	10.0	30.6	84.6	217.4

White, female

2006 4/.....	12.2	4.6	0.5	0.2	0.3	0.6	1.5	3.6	10.0	26.3	70.7	161.8
2005 4/.....	12.4	4.1	0.4	0.2	0.2	0.6	1.6	4.0	10.1	27.0	69.4	171.0
2004 4/.....	12.2	4.0	0.5	0.2	0.2	0.7	1.5	4.0	10.3	26.5	69.2	169.9
2003 4/.....	12.8	4.5	*	0.2	0.3	0.6	1.7	3.8	10.5	26.2	73.8	185.8
2002.....	12.8	6.0	0.4	0.2	0.3	0.6	1.5	3.8	9.6	28.6	73.4	187.1
2001.....	12.3	4.6	0.6	0.1	0.2	0.5	1.2	3.7	9.7	26.6	69.0	188.3
2000.....	12.1	5.6	0.5	*	0.1	0.4	1.5	3.3	9.0	24.6	66.8	200.2
1999.....	11.9	4.0	0.5	0.2	0.2	0.5	1.2	3.2	9.0	24.5	65.1	201.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Septicemia (A40-A41)

Black, both sexes

2006 4/.....	15.4	15.0	0.8	0.3	0.6	1.6	4.8	13.1	30.9	69.1	150.3	287.5
2005 4/.....	15.9	19.8	0.8	0.3	0.7	1.9	4.6	13.0	30.8	67.0	157.3	336.8
2004 4/.....	15.6	14.5	*	*	0.7	1.9	4.5	13.5	31.6	69.9	155.6	320.3
2003 4/.....	16.3	18.6	0.9	*	0.6	1.9	4.9	13.7	32.2	73.0	157.5	372.0
2002.....	16.3	13.5	*	0.3	0.4	1.9	4.4	12.9	33.2	74.7	171.2	367.0
2001.....	15.8	19.3	0.9	*	0.7	1.8	4.6	13.4	32.1	71.8	160.2	364.8
2000.....	16.2	16.4	1.0	*	0.6	2.0	4.7	13.6	31.4	71.9	173.4	392.6
1999.....	16.1	17.7	*	*	0.6	1.5	4.9	12.9	30.3	74.4	168.3	412.3

Black, male

2006 4/.....	14.4	15.6	*	*	0.6	1.7	5.0	14.8	37.1	75.8	175.3	296.7
2005 4/.....	14.9	20.4	*	*	0.7	1.9	4.5	14.8	35.7	80.6	180.6	374.6
2004 4/.....	14.6	16.3	*	*	0.7	2.0	4.7	15.1	36.2	81.1	189.0	341.7
2003 4/.....	15.3	21.1	*	*	*	2.1	5.0	15.3	37.8	85.7	189.0	404.5
2002.....	15.0	14.8	*	*	*	1.9	4.9	14.7	35.9	86.1	212.3	393.5
2001.....	14.9	20.1	*	*	0.8	1.8	4.7	14.9	36.9	85.6	195.1	424.4
2000.....	14.9	16.0	*	*	*	2.3	4.6	16.3	35.5	81.2	207.1	427.8
1999.....	15.3	19.8	*	*	*	1.6	5.8	15.1	37.1	90.3	201.7	453.5

Black, female

2006 4/.....	16.4	14.4	*	*	0.6	1.5	4.6	11.7	26.0	64.2	136.0	283.7
2005 4/.....	16.8	19.1	*	*	0.6	2.0	4.6	11.4	26.8	57.2	144.1	321.5
2004 4/.....	16.4	12.7	*	*	0.8	1.8	4.3	12.0	27.9	61.9	136.7	311.9
2003 4/.....	17.2	16.1	*	*	0.7	1.6	4.9	12.3	27.8	63.9	139.7	359.3
2002.....	17.4	12.1	*	*	*	1.9	4.1	11.4	30.9	66.6	148.0	356.9
2001.....	16.7	18.6	*	*	*	1.8	4.6	12.1	28.3	62.0	140.6	342.7
2000.....	17.4	16.9	*	*	*	1.7	4.8	11.3	28.2	65.3	154.4	379.5
1999.....	16.9	15.4	*	*	*	1.5	4.2	11.1	24.8	63.2	149.6	396.9



























Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Viral hepatitis (B15-B19)

All races 3/, both sexes

2006.....	2.4	*	*	*	*	0.2	1.2	6.7	6.7	4.5	4.6	3.1
2005.....	1.9	*	*	*	*	0.2	1.2	5.4	4.5	3.5	3.7	2.4
2004.....	1.8	*	*	*	*	0.2	1.4	5.3	4.3	3.4	3.6	2.7
2003.....	1.9	*	*	*	*	0.2	1.5	5.5	3.9	3.8	4.2	2.4
2002.....	2.0	*	*	*	0.1	0.2	1.6	5.8	4.0	4.2	4.6	3.9
2001.....	2.0	*	*	*	0.1	0.2	1.8	5.3	4.0	4.6	4.9	3.8
2000.....	1.9	*	*	*	*	0.2	1.7	5.0	3.7	4.6	5.3	4.0
1999.....	1.7	*	*	*	*	0.2	1.6	4.1	3.3	4.8	5.3	4.7

All races 3/, male

2006.....	3.2	*	*	*	*	0.2	1.7	9.9	9.9	5.0	4.6	3.0
2005.....	2.5	*	*	*	*	0.2	1.6	8.1	6.7	3.7	3.7	2.6
2004.....	2.4	*	*	*	*	0.2	1.8	8.0	6.1	4.0	3.6	3.0
2003.....	2.5	*	*	*	*	0.2	2.1	8.3	5.4	4.2	4.8	2.8
2002.....	2.7	*	*	*	*	0.3	2.3	8.9	5.5	4.5	5.3	5.0
2001.....	2.5	*	*	*	*	0.2	2.5	8.2	5.3	4.8	5.3	4.4
2000.....	2.4	*	*	*	*	0.2	2.4	7.7	5.0	4.8	6.4	5.7
1999.....	2.2	*	*	*	*	0.3	2.3	6.3	4.3	5.1	6.6	6.9

All races 3/, female

2006.....	1.6	*	*	*	*	0.1	0.8	3.7	3.8	4.0	4.6	3.1
2005.....	1.3	*	*	*	*	0.1	0.7	2.9	2.5	3.3	3.7	2.3
2004.....	1.3	*	*	*	*	0.1	1.0	2.7	2.5	3.0	3.5	2.5
2003.....	1.3	*	*	*	*	0.1	0.9	2.9	2.5	3.5	3.8	2.3
2002.....	1.4	*	*	*	*	0.2	1.0	2.8	2.7	3.9	4.1	3.3
2001.....	1.4	*	*	*	*	0.2	1.1	2.5	2.8	4.4	4.6	3.5
2000.....	1.4	*	*	*	*	0.2	1.1	2.4	2.6	4.5	4.7	3.3
1999.....	1.3	*	*	*	*	0.2	0.9	2.0	2.4	4.5	4.5	3.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Viral hepatitis (B15-B19)

White, both sexes

2006 4/.....	2.4	*	*	*	*	0.1	1.2	6.8	6.1	3.9	4.0	2.8
2005 4/.....	1.8	*	*	*	*	0.1	1.2	5.4	4.1	2.9	3.4	2.2
2004 4/.....	1.8	*	*	*	*	0.1	1.4	5.2	3.8	3.1	3.2	2.5
2003 4/.....	1.8	*	*	*	*	0.2	1.5	5.4	3.3	3.4	3.9	2.3
2002.....	2.0	*	*	*	0.1	0.2	1.6	5.6	3.5	3.7	4.4	3.7
2001.....	1.9	*	*	*	*	0.2	1.8	5.0	3.6	4.1	4.7	3.6
2000.....	1.8	*	*	*	*	0.2	1.7	4.6	3.2	4.2	5.1	3.9
1999.....	1.7	*	*	*	*	0.2	1.5	3.8	2.9	4.3	5.1	4.5

White, male

2006 4/.....	3.2	*	*	*	*	0.2	1.6	10.0	8.8	4.4	4.2	3.1
2005 4/.....	2.5	*	*	*	*	0.1	1.7	8.1	6.0	3.2	3.6	2.4
2004 4/.....	2.4	*	*	*	*	0.1	1.9	7.8	5.3	3.6	3.3	2.7
2003 4/.....	2.4	*	*	*	*	0.2	2.1	8.2	4.7	3.8	4.4	2.8
2002.....	2.6	*	*	*	*	0.3	2.3	8.7	4.8	4.1	5.2	4.9
2001.....	2.5	*	*	*	*	0.2	2.5	7.8	4.8	4.5	5.4	4.5
2000.....	2.4	*	*	*	*	0.2	2.4	7.2	4.3	4.3	6.2	5.7
1999.....	2.2	*	*	*	*	0.3	2.3	5.7	3.9	4.8	6.5	6.5

White, female

2006 4/.....	1.6	*	*	*	*	*	0.8	3.6	3.4	3.5	4.0	2.7
2005 4/.....	1.2	*	*	*	*	*	0.7	2.8	2.3	2.7	3.3	2.2
2004 4/.....	1.2	*	*	*	*	*	0.9	2.6	2.3	2.7	3.2	2.4
2003 4/.....	1.2	*	*	*	*	*	0.9	2.7	2.0	3.0	3.5	2.1
2002.....	1.3	*	*	*	*	*	0.9	2.7	2.4	3.3	3.9	3.2
2001.....	1.3	*	*	*	*	0.1	1.0	2.2	2.4	3.7	4.2	3.2
2000.....	1.3	*	*	*	*	0.1	1.0	2.1	2.2	4.1	4.5	3.2
1999.....	1.2	*	*	*	*	0.2	0.8	1.9	2.0	3.9	4.1	3.7



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Viral hepatitis (B15-B19)

Black, both sexes

2006 4/.....	2.8	*	*	*	*	*	1.3	7.6	12.3	7.2	7.0	*
2005 4/.....	2.2	*	*	*	*	*	1.1	7.0	8.4	6.1	5.2	*
2004 4/.....	2.2	*	*	*	*	*	1.4	7.5	8.7	4.6	5.0	*
2003 4/.....	2.2	*	*	*	*	*	1.7	7.4	8.0	5.4	6.0	*
2002.....	2.3	*	*	*	*	0.4	1.9	8.0	7.1	6.1	5.2	*
2001.....	2.2	*	*	*	*	*	2.1	7.9	6.7	5.6	5.7	*
2000.....	2.2	*	*	*	*	0.4	2.0	7.9	5.9	5.7	6.2	*
1999.....	2.0	*	*	*	*	*	2.1	6.9	5.0	6.9	6.1	6.4

Black, male

2006 4/.....	3.7	*	*	*	*	*	1.6	10.8	19.7	9.2	7.8	*
2005 4/.....	2.8	*	*	*	*	*	1.5	10.2	12.8	7.2	*	*
2004 4/.....	3.1	*	*	*	*	*	1.5	11.7	13.8	6.7	*	*
2003 4/.....	3.0	*	*	*	*	*	2.4	11.5	11.8	6.1	6.2	*
2002.....	3.0	*	*	*	*	*	2.5	12.4	10.5	7.1	*	*
2001.....	2.7	*	*	*	*	*	2.9	11.8	8.9	4.9	*	*
2000.....	2.9	*	*	*	*	*	2.9	12.3	8.1	7.0	7.6	*
1999.....	2.6	*	*	*	*	*	3.0	11.6	6.2	5.9	*	*

Black, female

2006 4/.....	1.9	*	*	*	*	*	1.0	4.9	6.3	5.7	6.5	*
2005 4/.....	1.7	*	*	*	*	*	0.9	4.2	4.8	5.2	6.2	*
2004 4/.....	1.5	*	*	*	*	*	1.4	3.8	4.5	3.2	5.2	*
2003 4/.....	1.6	*	*	*	*	*	1.1	4.0	5.0	4.9	5.9	*
2002.....	1.6	*	*	*	*	*	1.4	4.3	4.3	5.4	5.4	*
2001.....	1.8	*	*	*	*	*	1.4	4.7	4.9	6.0	6.8	*
2000.....	1.5	*	*	*	*	*	1.3	4.2	4.1	4.7	5.5	*
1999.....	1.5	*	*	*	*	*	1.3	2.8	4.1	7.6	6.5	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

All races 3/, both sexes

2006.....	4.0	*	*	*	0.5	2.9	9.2	10.1	5.5	2.5	0.8	*
2005.....	4.2	*	*	*	0.4	3.3	9.9	10.6	5.3	2.3	0.8	*
2004.....	4.4	*	*	0.1	0.5	3.7	10.9	10.6	5.4	2.4	0.8	*
2003.....	4.7	*	*	0.1	0.4	4.0	12.0	10.9	5.4	2.4	0.7	*
2002.....	4.9	*	*	0.1	0.4	4.6	12.7	11.2	5.1	2.2	0.8	*
2001.....	5.0	*	*	0.1	0.6	5.3	13.0	10.5	5.2	2.1	0.7	*
2000.....	5.1	*	*	0.1	0.5	6.1	13.1	11.0	5.1	2.2	0.7	*
1999.....	5.3	*	0.2	0.2	0.5	6.8	13.8	10.7	4.8	2.2	0.6	*

All races 3/, male

2006.....	5.9	*	*	*	0.6	3.5	12.9	15.3	8.9	4.2	1.6	*
2005.....	6.3	*	*	*	0.4	4.0	14.3	16.4	8.8	4.1	1.4	*
2004.....	6.6	*	*	*	0.5	4.5	15.7	16.3	9.0	4.0	1.4	*
2003.....	7.1	*	*	*	0.4	5.1	17.5	17.2	9.1	4.0	1.5	*
2002.....	7.4	*	*	*	0.4	5.9	18.8	17.7	8.5	3.9	1.4	*
2001.....	7.6	*	*	0.1	0.5	7.1	19.5	16.8	8.6	3.5	1.5	*
2000.....	7.9	*	*	0.1	0.5	8.0	19.8	17.8	8.7	3.8	1.3	*
1999.....	8.2	*	*	0.2	0.5	9.5	21.0	17.5	8.3	3.8	1.0	*

All races 3/, female

2006.....	2.2	*	*	*	0.4	2.3	5.4	5.1	2.3	1.1	0.3	*
2005.....	2.2	*	*	*	0.3	2.6	5.6	5.1	2.0	0.9	0.4	*
2004.....	2.4	*	*	*	0.4	2.8	6.2	5.2	2.0	1.0	0.5	*
2003.....	2.4	*	*	*	0.4	2.8	6.5	4.8	2.1	1.0	0.3	*
2002.....	2.5	*	*	*	0.4	3.3	6.7	4.8	1.9	0.8	0.3	*
2001.....	2.5	*	*	*	0.6	3.5	6.7	4.4	2.0	0.9	*	*
2000.....	2.5	*	*	0.1	0.4	4.2	6.5	4.4	1.8	0.8	0.3	*
1999.....	2.5	*	*	0.2	0.5	4.1	6.7	4.1	1.6	0.8	0.3	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

White, both sexes

2006 4/.....	2.1	*	*	*	0.2	1.3	5.0	5.3	2.7	1.3	0.5	*
2005 4/.....	2.2	*	*	*	0.2	1.5	5.5	5.5	2.6	1.2	0.5	*
2004 4/.....	2.4	*	*	*	0.1	1.6	6.2	5.5	2.8	1.2	0.5	*
2003 4/.....	2.5	*	*	*	0.2	2.0	6.8	5.8	2.9	1.2	0.4	*
2002.....	2.6	*	*	*	0.2	2.2	7.0	5.8	2.8	1.0	0.4	*
2001.....	2.7	*	*	*	0.2	2.6	7.3	5.5	2.8	1.0	0.3	*
2000.....	2.8	*	*	0.1	0.2	3.1	7.5	5.9	2.8	1.0	0.4	*
1999.....	3.0	*	*	0.1	0.2	3.7	8.0	5.7	2.5	1.1	0.3	*

White, male

2006 4/.....	3.5	*	*	*	0.2	1.9	8.1	8.9	4.8	2.3	0.9	*
2005 4/.....	3.7	*	*	*	0.2	2.1	9.0	9.2	4.6	2.1	0.9	*
2004 4/.....	3.9	*	*	*	0.2	2.4	9.9	9.2	5.0	2.2	0.9	*
2003 4/.....	4.2	*	*	*	0.2	3.0	11.0	10.0	5.0	2.1	0.9	*
2002.....	4.3	*	*	*	0.2	3.3	11.6	9.8	4.9	1.8	0.6	*
2001.....	4.5	*	*	*	0.2	3.9	12.1	9.5	4.8	1.9	0.7	*
2000.....	4.7	*	*	*	0.2	4.6	12.4	10.1	5.0	1.9	0.7	*
1999.....	5.0	*	*	*	0.3	5.8	13.2	9.9	4.7	2.0	0.7	*

White, female

2006 4/.....	0.7	*	*	*	0.1	0.7	1.9	1.7	0.7	0.4	*	*
2005 4/.....	0.8	*	*	*	*	0.8	2.0	1.9	0.8	0.3	*	*
2004 4/.....	0.8	*	*	*	*	0.7	2.4	1.9	0.7	0.4	*	*
2003 4/.....	0.9	*	*	*	0.1	0.9	2.5	1.7	0.9	0.4	*	*
2002.....	0.9	*	*	*	0.1	1.0	2.5	1.8	0.8	*	*	*
2001.....	0.9	*	*	*	0.2	1.2	2.4	1.5	0.8	0.3	*	*
2000.....	1.0	*	*	*	*	1.5	2.6	1.7	0.8	0.3	*	*
1999.....	1.0	*	*	*	0.2	1.6	2.7	1.6	0.6	0.3	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Human immunodeficiency virus (HIV) disease (B20-B24)

Black, both sexes

2006 4/.....	17.3	*	*	*	2.2	13.6	38.4	47.2	30.3	14.1	4.8	*
2005 4/.....	18.0	*	*	*	1.7	14.9	40.9	49.7	29.8	13.9	4.9	*
2004 4/.....	18.8	*	*	*	2.2	17.1	44.6	50.0	29.7	13.6	4.5	*
2003 4/.....	19.6	*	*	0.5	1.9	17.3	49.2	50.6	29.6	14.5	4.7	*
2002.....	20.8	*	*	*	2.0	20.6	52.8	53.9	26.1	14.4	5.6	*
2001.....	21.1	*	*	0.4	2.7	23.2	53.7	51.2	27.7	12.8	5.3	*
2000.....	21.4	*	*	0.3	2.1	26.3	53.5	53.4	26.0	13.5	4.8	*
1999.....	21.8	*	*	0.6	2.3	27.3	56.1	51.8	26.2	13.5	3.6	*

Black, male

2006 4/.....	23.5	*	*	*	2.5	15.4	50.6	68.6	49.3	23.8	10.3	*
2005 4/.....	25.1	*	*	*	1.7	17.0	55.4	75.0	50.9	25.0	8.7	*
2004 4/.....	25.9	*	*	*	2.2	19.2	60.3	74.9	50.3	23.4	7.5	*
2003 4/.....	27.7	*	*	*	1.9	20.8	68.0	77.8	50.7	25.8	9.0	*
2002.....	29.5	*	*	*	1.9	24.5	74.0	85.0	44.4	26.3	12.0	*
2001.....	30.1	*	*	*	2.3	29.8	75.9	80.7	47.1	21.7	11.5	*
2000.....	31.0	*	*	*	2.0	32.3	77.5	86.0	46.0	24.6	9.4	*
1999.....	31.9	*	*	0.7	1.9	36.0	81.9	85.0	45.5	24.7	6.5	*

Black, female

2006 4/.....	11.7	*	*	*	1.8	12.0	27.5	28.8	15.0	7.1	*	*
2005 4/.....	11.5	*	*	*	1.6	13.0	28.0	27.9	12.7	6.0	*	*
2004 4/.....	12.4	*	*	*	2.2	15.1	30.6	28.6	13.1	6.5	*	*
2003 4/.....	12.3	*	*	*	1.9	14.2	32.5	27.3	12.5	6.4	*	*
2002.....	12.8	*	*	*	2.1	17.1	34.0	27.2	11.3	6.0	*	*
2001.....	12.9	*	*	*	3.1	17.3	34.1	25.9	12.1	6.5	*	*
2000.....	12.8	*	*	*	2.2	20.8	32.2	25.3	10.0	5.8	*	*
1999.....	12.6	*	*	*	2.7	19.5	33.4	23.4	10.7	5.6	*	*







Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

All races 3/, both sexes

2006.....	2.0	4.0	0.4	0.2	0.2	0.3	0.6	1.3	2.7	5.8	11.5	20.4
2005.....	2.6	4.1	0.4	0.2	0.2	0.3	0.9	3.2	4.2	7.0	13.7	21.2
2004.....	2.6	4.1	0.4	0.1	0.2	0.4	1.0	3.0	4.3	7.1	13.2	22.5
2003.....	2.6	4.3	0.5	0.1	0.2	0.4	1.0	2.9	4.1	7.3	13.3	22.5
2002.....	2.3	5.6	0.4	0.2	0.2	0.4	1.0	2.3	3.3	6.7	12.5	21.7
2001.....	2.0	4.7	0.4	0.2	0.2	0.4	0.8	1.7	3.0	6.2	10.6	20.1
2000.....	2.0	4.4	0.5	0.2	0.2	0.4	0.8	1.6	2.9	5.8	11.1	21.2
1999.....	2.6	5.3	0.5	0.2	0.2	0.6	1.2	2.3	3.8	7.8	14.2	27.0

All races 3/, male

2006.....	2.1	4.5	0.5	0.2	0.2	0.3	0.8	1.6	3.2	6.7	13.9	25.1
2005.....	2.9	4.8	0.5	0.1	0.2	0.4	1.2	4.5	5.3	8.3	16.4	23.6
2004.....	2.8	4.8	0.4	0.1	0.3	0.4	1.2	4.1	5.4	8.4	15.1	29.6
2003.....	2.8	4.2	0.5	0.1	0.2	0.4	1.3	4.0	5.1	8.5	15.8	26.5
2002.....	2.6	6.1	0.5	0.2	0.2	0.5	1.3	3.2	4.2	8.0	15.4	26.8
2001.....	2.1	4.9	0.5	0.1	0.2	0.4	1.0	2.2	3.7	6.8	13.2	23.6
2000.....	2.1	4.8	0.5	0.2	0.3	0.4	1.0	1.9	3.3	7.2	13.6	27.6
1999.....	2.7	6.1	0.6	0.2	0.2	0.6	1.5	2.7	4.2	9.2	17.2	31.9

All races 3/, female

2006.....	1.9	3.6	0.3	0.2	0.2	0.3	0.5	1.0	2.3	5.1	9.9	18.2
2005.....	2.3	3.4	0.3	0.2	0.2	0.3	0.6	2.0	3.2	5.9	11.8	20.1
2004.....	2.3	3.4	0.4	0.1	0.2	0.3	0.8	1.9	3.2	6.0	11.9	19.3
2003.....	2.3	4.5	0.5	0.1	0.2	0.3	0.7	1.9	3.1	6.3	11.7	20.8
2002.....	2.1	5.0	0.3	0.2	0.2	0.4	0.7	1.4	2.6	5.6	10.6	19.5
2001.....	1.9	4.6	0.4	0.2	0.2	0.3	0.6	1.2	2.4	5.8	8.9	18.7
2000.....	1.9	3.9	0.4	0.2	0.2	0.3	0.6	1.2	2.6	4.7	9.5	18.6
1999.....	2.5	4.5	0.4	0.2	0.2	0.6	0.9	1.9	3.4	6.7	12.3	25.1



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

White, both sexes

2006 4/.....	2.0	2.8	0.4	0.2	0.2	0.2	0.5	1.2	2.6	5.8	11.5	20.6
2005 4/.....	2.7	3.7	0.4	0.2	0.2	0.3	0.8	3.1	4.0	6.9	13.4	21.2
2004 4/.....	2.6	3.7	0.4	0.1	0.2	0.3	0.9	3.0	4.0	6.8	13.0	22.5
2003 4/.....	2.6	3.5	0.4	0.1	0.2	0.3	0.9	2.8	3.8	7.3	13.0	23.0
2002.....	2.4	5.0	0.4	0.2	0.2	0.4	0.8	2.1	3.0	6.6	12.4	22.0
2001.....	2.0	3.8	0.4	0.2	0.2	0.3	0.7	1.6	2.8	6.0	10.5	20.1
2000.....	2.0	3.4	0.4	0.2	0.2	0.3	0.7	1.4	2.7	5.7	11.0	21.0
1999.....	2.5	4.5	0.5	0.2	0.2	0.4	1.0	1.9	3.4	7.5	13.7	26.5

White, male

2006 4/.....	2.1	3.3	0.5	0.2	0.2	0.3	0.6	1.4	2.9	6.7	13.8	25.7
2005 4/.....	3.0	4.2	0.4	0.1	0.2	0.3	1.1	4.4	5.0	8.3	16.1	23.7
2004 4/.....	2.9	4.2	0.4	*	0.3	0.3	1.1	4.1	5.1	8.1	14.9	30.1
2003 4/.....	2.8	3.2	0.4	*	0.2	0.4	1.2	3.8	4.8	8.3	15.5	27.2
2002.....	2.6	5.3	0.5	0.2	0.2	0.4	1.1	3.0	3.7	7.9	15.2	27.2
2001.....	2.1	3.9	0.4	0.1	0.2	0.4	0.9	2.0	3.4	6.6	13.1	23.8
2000.....	2.1	3.9	0.4	0.2	0.3	0.3	0.8	1.7	3.1	7.0	13.3	26.9
1999.....	2.6	5.6	0.6	0.2	0.2	0.5	1.3	2.3	3.8	8.8	16.3	30.6

White, female

2006 4/.....	2.0	2.2	*	0.2	0.2	0.2	0.4	0.9	2.2	5.0	10.0	18.1
2005 4/.....	2.4	3.1	*	0.2	0.2	0.2	0.5	1.8	3.1	5.7	11.6	20.0
2004 4/.....	2.4	3.0	0.3	*	0.2	0.3	0.7	1.9	3.0	5.7	11.7	19.0
2003 4/.....	2.4	3.9	0.5	0.1	0.1	0.3	0.6	1.8	2.9	6.4	11.3	21.2
2002.....	2.2	4.6	*	0.2	0.2	0.4	0.6	1.3	2.4	5.5	10.5	19.8
2001.....	1.9	3.8	*	0.2	0.2	0.2	0.5	1.1	2.2	5.6	8.7	18.5
2000.....	1.9	2.9	0.4	0.2	0.2	0.2	0.5	1.1	2.3	4.6	9.5	18.7
1999.....	2.5	3.4	*	0.2	0.2	0.4	0.6	1.5	3.0	6.4	12.0	24.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1,A85.8,A86-B04,B06-B09, B25-B49,B55-B99)

Black, both sexes

2006 4/.....	2.1	9.9	*	*	*	0.7	1.3	2.6	4.3	7.2	12.4	22.1
2005 4/.....	2.5	6.5	*	*	0.3	0.7	1.5	4.1	5.9	7.8	14.9	22.7
2004 4/.....	2.5	6.7	*	*	0.3	0.8	1.6	3.4	6.2	8.7	16.5	24.3
2003 4/.....	2.5	8.0	1.0	*	*	0.6	1.8	4.3	6.1	7.0	16.2	19.5
2002.....	2.5	8.7	*	*	0.4	0.9	2.2	3.9	5.8	7.2	13.8	21.7
2001.....	2.2	9.5	*	*	0.4	0.8	1.6	2.9	5.1	8.3	11.1	20.6
2000.....	2.3	9.9	*	*	0.4	0.9	1.6	3.0	4.9	7.8	13.3	24.1
1999.....	3.5	9.9	0.9	*	0.3	1.6	3.0	5.4	7.2	11.8	20.5	34.1

Black, male

2006 4/.....	2.3	10.1	*	*	*	0.7	1.8	3.4	5.9	7.7	16.0	20.6
2005 4/.....	2.8	8.0	*	*	*	0.8	2.0	5.1	8.0	9.4	19.4	23.2
2004 4/.....	2.9	7.4	*	*	*	0.9	1.8	4.9	8.9	10.5	19.8	29.8
2003 4/.....	2.9	8.9	*	*	*	0.8	2.1	5.7	8.0	10.7	18.6	22.9
2002.....	2.9	9.6	*	*	*	1.1	2.6	5.5	8.4	8.8	16.9	28.9
2001.....	2.4	10.8	*	*	*	*	2.2	3.9	6.3	10.0	12.4	*
2000.....	2.5	9.9	*	*	*	0.9	2.1	3.9	5.7	10.2	18.2	32.3
1999.....	3.6	10.1	*	*	*	1.8	3.1	6.1	7.8	14.0	28.9	45.7

Black, female

2006 4/.....	2.0	9.6	*	*	*	*	0.9	1.8	3.0	6.9	10.3	22.7
2005 4/.....	2.2	*	*	*	*	*	1.0	3.3	4.2	6.6	12.4	22.5
2004 4/.....	2.2	*	*	*	*	*	1.4	2.1	4.0	7.4	14.6	22.1
2003 4/.....	2.2	7.1	*	*	*	*	1.4	3.1	4.6	4.3	14.8	18.2
2002.....	2.1	7.9	*	*	*	*	1.8	2.5	3.6	6.1	12.0	18.9
2001.....	2.0	8.2	*	*	*	0.9	1.1	2.1	4.2	7.2	10.4	20.2
2000.....	2.1	9.9	*	*	*	1.0	1.2	2.2	4.2	6.1	10.6	21.0
1999.....	3.4	9.7	*	*	*	1.4	2.9	4.8	6.7	10.3	15.8	29.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasms (C00-C97)

All races 3/, both sexes

2006.....	187.0	1.8	2.3	2.2	3.9	9.0	31.9	116.3	321.2	727.2	1,263.8	1,606.1
2005.....	188.7	1.8	2.3	2.5	4.1	9.0	33.2	118.6	326.9	742.7	1,274.8	1,637.7
2004.....	188.6	1.8	2.5	2.5	4.1	9.1	33.4	119.0	333.4	755.1	1,280.4	1,653.3
2003.....	191.5	1.9	2.5	2.6	4.0	9.4	35.0	122.2	343.0	770.3	1,302.5	1,698.2
2002.....	193.2	1.8	2.6	2.6	4.3	9.7	35.8	123.8	351.1	792.1	1,311.9	1,723.9
2001.....	194.4	1.6	2.7	2.5	4.3	10.1	36.8	126.5	356.5	802.8	1,315.8	1,765.6
2000.....	196.5	2.4	2.7	2.5	4.4	9.8	36.6	127.5	366.7	816.3	1,335.6	1,819.4
1999.....	197.0	1.8	2.7	2.5	4.5	10.0	37.1	127.6	374.6	827.1	1,331.5	1,805.8

All races 3/, male

2006.....	196.6	1.8	2.5	2.5	4.6	8.6	27.4	119.0	363.6	870.4	1,631.3	2,248.7
2005.....	198.9	2.1	2.6	2.7	4.8	8.8	28.9	121.6	369.5	899.1	1,649.7	2,319.3
2004.....	198.4	1.8	2.6	2.7	4.8	8.6	29.1	124.3	376.7	907.6	1,662.1	2,349.5
2003.....	201.3	1.7	2.8	2.8	4.6	8.9	30.8	127.4	386.8	931.7	1,695.4	2,413.8
2002.....	203.8	2.0	2.7	2.9	4.9	9.2	31.5	128.0	399.8	964.8	1,711.3	2,491.1
2001.....	205.3	1.5	2.9	2.5	5.0	9.3	32.6	130.3	405.2	984.6	1,727.1	2,613.6
2000.....	207.2	2.6	3.0	2.7	5.1	9.2	32.7	130.9	415.8	1,001.9	1,760.6	2,710.7
1999.....	208.9	1.9	2.9	2.7	5.2	9.7	33.3	131.7	428.4	1,019.4	1,767.0	2,722.2

All races 3/, female

2006.....	177.6	1.8	2.1	2.0	3.1	9.5	36.4	113.7	281.8	605.9	1,012.5	1,305.5
2005.....	178.8	1.5	2.0	2.2	3.3	9.1	37.5	115.8	287.4	610.9	1,020.3	1,324.6
2004.....	179.1	1.9	2.4	2.2	3.4	9.6	37.7	113.8	293.2	627.1	1,023.5	1,340.1
2003.....	182.0	2.1	2.1	2.4	3.4	9.9	39.1	117.1	302.3	635.3	1,040.1	1,381.9
2002.....	183.0	1.6	2.4	2.4	3.6	10.2	40.0	119.8	306.0	648.5	1,046.7	1,391.1
2001.....	183.9	1.8	2.5	2.4	3.5	10.9	41.0	122.7	311.5	652.2	1,045.4	1,410.7
2000.....	186.2	2.3	2.5	2.2	3.6	10.4	40.4	124.2	321.3	663.6	1,058.5	1,456.4
1999.....	185.6	1.7	2.5	2.3	3.7	10.3	40.9	123.7	325.0	669.5	1,048.5	1,436.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasms (C00-C97)

White, both sexes

2006 4/.....	199.3	2.0	2.3	2.3	3.9	8.8	30.7	111.5	315.0	727.5	1,272.3	1,618.1
2005 4/.....	200.8	2.0	2.3	2.5	4.1	8.5	32.1	113.2	319.3	742.7	1,281.6	1,645.4
2004 4/.....	200.7	2.0	2.5	2.5	4.0	8.8	32.0	113.3	326.9	753.3	1,284.3	1,659.7
2003 4/.....	203.7	2.1	2.6	2.7	4.1	9.1	33.6	116.0	336.6	769.5	1,303.4	1,701.4
2002.....	205.5	1.8	2.5	2.7	4.4	9.5	34.5	118.5	343.3	790.1	1,311.0	1,725.7
2001.....	206.4	1.7	3.0	2.4	4.3	9.8	35.3	121.2	349.1	797.8	1,311.1	1,765.2
2000.....	208.6	2.5	2.8	2.6	4.3	9.7	34.5	121.8	359.4	811.0	1,330.3	1,815.7
1999.....	208.7	1.8	2.9	2.4	4.3	9.8	35.3	121.9	365.2	820.7	1,323.7	1,798.8

White, male

2006 4/.....	208.7	2.0	2.5	2.5	4.7	8.6	26.7	113.6	352.9	862.0	1,631.3	2,258.3
2005 4/.....	210.6	2.3	2.6	2.7	4.9	8.5	28.4	115.7	356.5	889.9	1,646.2	2,322.7
2004 4/.....	209.9	1.9	2.5	2.8	4.7	8.6	28.2	117.5	364.9	896.3	1,652.7	2,348.9
2003 4/.....	213.1	1.8	2.9	2.9	4.7	8.9	29.9	119.9	375.6	922.7	1,683.6	2,412.1
2002.....	215.5	2.2	2.7	2.8	5.0	9.1	30.5	121.8	386.0	954.8	1,695.3	2,486.8
2001.....	216.4	1.5	3.1	2.4	5.1	9.3	31.3	123.6	392.1	969.4	1,704.6	2,597.6
2000.....	218.1	2.8	3.1	2.8	5.0	9.2	30.9	123.5	401.9	984.3	1,736.0	2,693.7
1999.....	219.5	2.0	3.2	2.7	5.0	9.5	31.9	123.5	412.7	1,002.3	1,741.8	2,696.3

White, female

2006 4/.....	190.1	2.0	2.0	2.1	3.0	9.1	34.9	109.5	279.1	611.5	1,023.0	1,317.5
2005 4/.....	191.1	1.7	2.0	2.3	3.4	8.6	36.0	110.7	284.0	616.2	1,030.5	1,333.6
2004 4/.....	191.7	2.1	2.5	2.2	3.3	9.0	35.8	109.2	290.8	630.8	1,033.1	1,348.9
2003 4/.....	194.6	2.3	2.3	2.5	3.4	9.4	37.3	112.1	299.8	638.9	1,046.3	1,386.5
2002.....	195.8	1.4	2.3	2.5	3.7	9.9	38.5	115.3	303.1	650.4	1,053.1	1,395.1
2001.....	196.7	2.0	2.8	2.5	3.5	10.4	39.3	118.9	308.6	652.9	1,049.8	1,416.7
2000.....	199.4	2.3	2.4	2.3	3.6	10.1	38.2	120.1	319.7	665.6	1,063.4	1,459.1
1999.....	198.2	1.5	2.7	2.2	3.7	10.1	38.7	120.2	320.7	669.4	1,049.6	1,438.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms (C00-C97)

Black, both sexes

2006 4/.....	159.5	*	2.5	2.2	4.0	11.3	44.1	169.5	439.9	874.6	1,396.6	1,678.3
2005 4/.....	161.7	*	2.4	2.6	4.0	12.3	44.8	175.5	455.9	889.9	1,414.4	1,744.7
2004 4/.....	161.9	*	2.5	2.5	4.8	12.0	46.8	177.6	458.5	915.9	1,446.0	1,774.0
2003 4/.....	164.3	*	2.2	2.6	3.9	12.2	48.9	185.6	471.0	925.4	1,487.9	1,853.1
2002.....	165.9	*	2.9	2.5	4.0	12.3	50.0	181.7	485.6	962.6	1,520.7	1,885.7
2001.....	167.1	*	1.9	2.6	4.3	12.9	51.4	184.4	486.1	991.2	1,543.9	1,931.7
2000.....	169.3	*	2.8	2.2	5.1	11.9	54.0	192.6	496.0	1,005.3	1,566.3	1,998.9
1999.....	171.3	*	2.0	2.7	5.5	12.2	53.9	192.9	521.4	1,019.7	1,579.3	2,008.4

Black, male

2006 4/.....	172.3	*	2.6	2.5	4.5	10.0	36.5	182.2	542.9	1,156.5	1,979.1	2,543.3
2005 4/.....	175.4	*	2.8	3.1	4.7	11.9	36.2	186.1	568.3	1,183.8	2,017.5	2,683.7
2004 4/.....	176.2	*	2.8	2.5	5.4	10.0	38.4	197.0	569.2	1,209.7	2,087.2	2,748.8
2003 4/.....	178.3	*	2.4	2.8	4.5	10.3	41.7	207.0	583.8	1,221.5	2,144.2	2,825.5
2002.....	181.5	*	2.6	3.1	4.5	11.2	43.0	197.3	610.3	1,274.7	2,223.0	2,976.1
2001.....	184.5	*	2.4	2.9	5.0	10.5	44.6	204.8	604.2	1,335.3	2,290.0	3,209.9
2000.....	188.5	*	2.7	2.5	6.1	10.1	48.4	214.2	626.4	1,363.8	2,351.8	3,264.8
1999.....	191.0	*	2.1	2.8	6.4	11.4	47.3	221.3	659.0	1,369.8	2,359.4	3,340.7

Black, female

2006 4/.....	147.7	*	2.3	1.9	3.4	12.5	50.8	158.7	356.9	672.9	1,065.3	1,324.4
2005 4/.....	149.1	*	2.1	2.1	3.4	12.6	52.5	166.3	365.4	679.6	1,071.9	1,365.8
2004 4/.....	148.9	*	2.2	2.6	4.2	13.9	54.2	160.9	369.4	706.2	1,083.6	1,387.7
2003 4/.....	151.4	*	2.1	2.3	3.3	13.9	55.4	167.2	380.4	714.6	1,116.9	1,475.3
2002.....	151.7	*	3.2	2.0	3.6	13.3	56.2	168.2	385.4	741.1	1,123.1	1,468.0
2001.....	151.3	*	*	2.2	3.6	15.0	57.3	166.8	390.9	748.4	1,125.0	1,457.5
2000.....	151.8	*	2.9	1.9	4.1	13.5	58.9	173.9	391.0	753.1	1,124.0	1,527.7
1999.....	153.3	*	1.9	2.6	4.6	13.0	59.8	168.5	411.1	772.7	1,142.6	1,510.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/ 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes

2006.....	2.6	*	*	*	*	0.1	0.6	2.6	5.8	9.6	13.5	16.3
2005.....	2.6	*	*	*	0.1	0.1	0.6	2.6	6.4	9.8	12.8	17.0
2004.....	2.7	*	*	*	0.1	0.1	0.6	2.9	6.1	10.1	13.7	18.0
2003.....	2.7	*	*	*	0.1	0.1	0.6	2.8	6.5	10.2	13.1	18.6
2002.....	2.7	*	*	*	0.1	0.1	0.6	2.7	6.8	10.4	13.5	18.7
2001.....	2.7	*	*	*	0.1	0.1	0.6	2.7	6.9	10.4	14.2	18.5
2000.....	2.7	*	*	*	0.1	0.1	0.6	2.8	6.8	10.7	13.2	19.4
1999.....	2.7	*	*	*	0.1	0.1	0.6	2.9	6.8	10.3	14.3	19.3

All races 3/, male

2006.....	3.6	*	*	*	*	0.1	0.9	4.1	9.3	14.8	20.4	21.9
2005.....	3.6	*	*	*	*	0.2	0.9	4.2	10.3	15.3	18.9	22.0
2004.....	3.7	*	*	*	*	0.2	0.8	4.6	10.0	15.6	20.3	24.9
2003.....	3.7	*	*	*	0.1	0.2	0.9	4.6	10.6	15.6	19.4	26.9
2002.....	3.7	*	*	*	*	0.2	0.8	4.4	10.9	15.9	19.6	26.9
2001.....	3.7	*	*	*	*	0.2	0.9	4.4	11.3	16.0	20.6	25.8
2000.....	3.6	*	*	*	*	0.1	1.0	4.5	10.9	16.0	19.2	27.1
1999.....	3.6	*	*	*	0.1	0.1	0.9	4.6	10.8	15.8	20.9	27.1

All races 3/, female

2006.....	1.6	*	*	*	*	*	0.4	1.1	2.4	5.2	8.8	13.7
2005.....	1.7	*	*	*	*	0.1	0.3	1.2	2.8	5.2	8.7	14.7
2004.....	1.7	*	*	*	*	*	0.4	1.2	2.5	5.6	9.2	14.9
2003.....	1.7	*	*	*	*	0.1	0.4	1.0	2.8	5.7	9.0	15.0
2002.....	1.7	*	*	*	*	*	0.4	1.1	3.0	5.8	9.5	15.1
2001.....	1.8	*	*	*	*	0.1	0.4	1.2	2.8	5.8	10.0	15.4
2000.....	1.8	*	*	*	*	0.1	0.3	1.3	2.9	6.3	9.2	16.3
1999.....	1.8	*	*	*	*	*	0.4	1.3	3.1	5.8	9.9	16.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

White, both sexes

2006 4/.....	2.7	*	*	*	*	0.1	0.6	2.4	5.4	9.2	13.6	16.9
2005 4/.....	2.7	*	*	*	*	0.1	0.5	2.4	6.0	9.5	12.8	17.3
2004 4/.....	2.7	*	*	*	*	0.1	0.5	2.5	5.6	9.9	13.7	18.6
2003 4/.....	2.7	*	*	*	*	0.1	0.5	2.4	6.2	9.8	12.9	19.0
2002.....	2.7	*	*	*	*	0.1	0.5	2.4	6.3	10.2	13.7	18.9
2001.....	2.7	*	*	*	0.1	0.1	0.5	2.4	6.3	10.0	14.4	19.0
2000.....	2.7	*	*	*	*	0.1	0.5	2.4	6.0	10.2	13.1	19.7
1999.....	2.7	*	*	*	*	0.1	0.5	2.3	6.1	9.9	14.1	19.8

White, male

2006 4/.....	3.6	*	*	*	*	0.1	0.8	3.8	8.6	14.0	20.2	22.4
2005 4/.....	3.6	*	*	*	*	*	0.7	3.7	9.4	14.5	18.4	22.1
2004 4/.....	3.6	*	*	*	*	0.2	0.6	3.9	9.1	14.8	19.9	24.8
2003 4/.....	3.7	*	*	*	*	0.2	0.8	4.0	9.7	14.6	18.6	27.2
2002.....	3.7	*	*	*	*	0.2	0.7	3.8	9.9	15.3	19.7	27.0
2001.....	3.6	*	*	*	*	0.1	0.7	3.9	10.1	15.1	20.4	26.4
2000.....	3.5	*	*	*	*	0.1	0.7	3.7	9.6	14.9	18.7	27.6
1999.....	3.5	*	*	*	*	*	0.7	3.6	9.6	15.1	20.3	27.3

White, female

2006 4/.....	1.7	*	*	*	*	*	0.3	1.0	2.3	5.1	9.0	14.3
2005 4/.....	1.8	*	*	*	*	*	0.3	1.1	2.7	5.3	8.9	15.1
2004 4/.....	1.8	*	*	*	*	*	0.3	1.1	2.4	5.8	9.4	15.7
2003 4/.....	1.8	*	*	*	*	*	0.2	0.9	2.8	5.7	9.1	15.4
2002.....	1.8	*	*	*	*	*	0.3	1.0	2.8	5.9	9.6	15.3
2001.....	1.9	*	*	*	*	*	0.3	1.0	2.7	5.8	10.3	15.9
2000.....	1.9	*	*	*	*	*	0.3	1.2	2.7	6.3	9.4	16.5
1999.....	1.9	*	*	*	*	*	0.3	1.0	2.8	5.6	10.1	16.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

Black, both sexes

2006 4/.....	2.6	*	*	*	*	*	0.8	4.3	9.5	13.8	13.5	10.4
2005 4/.....	2.8	*	*	*	*	*	0.9	5.0	10.6	13.9	13.6	12.6
2004 4/.....	3.0	*	*	*	*	*	1.3	6.0	11.1	13.8	15.3	12.4
2003 4/.....	2.9	*	*	*	*	*	1.1	5.1	11.2	15.4	15.3	15.5
2002.....	2.8	*	*	*	*	*	1.1	5.5	12.1	12.9	11.8	16.0
2001.....	2.9	*	*	*	*	*	1.2	5.1	12.6	15.5	13.9	12.9
2000.....	3.0	*	*	*	*	*	1.1	6.0	13.5	15.5	15.1	15.3
1999.....	3.2	*	*	*	*	*	1.2	7.1	13.5	15.2	16.9	14.3

Black, male

2006 4/.....	4.0	*	*	*	*	*	1.2	6.9	16.9	25.3	25.2	*
2005 4/.....	4.5	*	*	*	*	*	1.5	8.6	19.5	26.3	27.6	21.4
2004 4/.....	4.7	*	*	*	*	*	1.9	10.4	19.8	26.6	26.4	26.8
2003 4/.....	4.6	*	*	*	*	*	1.2	9.0	21.1	28.7	27.9	25.0
2002.....	4.3	*	*	*	*	*	1.6	9.5	21.9	23.0	18.4	26.8
2001.....	4.4	*	*	*	*	*	1.7	8.7	23.3	27.1	23.5	*
2000.....	4.8	*	*	*	*	*	1.7	10.5	24.3	28.2	28.8	*
1999.....	4.8	*	*	*	*	*	1.8	12.1	23.2	25.1	31.7	24.6

Black, female

2006 4/.....	1.3	*	*	*	*	*	*	2.0	3.5	5.6	6.9	8.8
2005 4/.....	1.2	*	*	*	*	*	*	1.8	3.5	5.1	5.6	9.0
2004 4/.....	1.4	*	*	*	*	*	0.8	2.3	4.2	4.6	9.0	*
2003 4/.....	1.4	*	*	*	*	*	1.0	1.7	3.2	6.0	8.1	11.7
2002.....	1.4	*	*	*	*	*	0.7	2.0	4.2	5.8	8.1	11.9
2001.....	1.5	*	*	*	*	*	0.8	2.1	3.9	7.4	8.4	10.5
2000.....	1.4	*	*	*	*	*	*	2.0	4.8	6.5	7.3	14.2
1999.....	1.7	*	*	*	*	*	0.7	2.9	5.7	8.2	8.6	10.5



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of esophagus (C15)

All races 3/, both sexes

2006.....	4.6	*	*	*	*	0.1	0.6	3.5	10.4	19.5	26.9	25.6
2005.....	4.6	*	*	*	*	0.1	0.7	3.6	10.2	19.8	27.4	24.9
2004.....	4.4	*	*	*	*	0.1	0.7	3.4	10.3	19.2	26.7	25.8
2003.....	4.4	*	*	*	*	0.1	0.6	3.4	10.5	19.7	26.4	25.0
2002.....	4.4	*	*	*	*	*	0.7	3.5	10.4	20.2	26.6	24.6
2001.....	4.4	*	*	*	*	0.1	0.7	3.6	10.1	20.6	26.3	25.8
2000.....	4.3	*	*	*	*	0.1	0.7	3.6	10.8	20.0	25.6	25.9
1999.....	4.3	*	*	*	*	0.1	0.7	3.5	10.6	20.1	24.4	26.8

All races 3/, male

2006.....	7.3	*	*	*	*	0.1	1.1	6.0	18.4	34.3	48.8	48.5
2005.....	7.3	*	*	*	*	0.1	1.2	6.2	17.9	34.9	50.7	46.0
2004.....	7.0	*	*	*	*	0.1	1.1	5.9	17.9	33.4	48.8	48.5
2003.....	6.9	*	*	*	*	0.1	1.1	6.0	18.5	34.1	47.8	45.3
2002.....	6.9	*	*	*	*	*	1.1	6.0	18.2	35.6	48.3	43.4
2001.....	6.9	*	*	*	*	0.1	1.1	6.3	17.6	36.0	47.4	48.1
2000.....	6.7	*	*	*	*	0.2	1.2	6.3	18.6	34.1	45.8	47.8
1999.....	6.6	*	*	*	*	0.1	1.1	6.1	18.3	35.3	44.4	48.0

All races 3/, female

2006.....	2.0	*	*	*	*	*	0.2	1.1	3.0	7.0	12.0	14.9
2005.....	1.9	*	*	*	*	*	0.2	1.0	3.1	7.0	11.6	15.3
2004.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.3	11.9	15.7
2003.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.7	12.0	16.1
2002.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.4	12.2	16.5
2001.....	2.0	*	*	*	*	*	0.2	1.0	3.2	7.9	12.4	16.5
2000.....	2.1	*	*	*	*	*	0.1	1.0	3.5	8.3	12.4	17.1
1999.....	2.0	*	*	*	*	*	0.2	0.9	3.6	7.6	11.5	18.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of esophagus (C15)

White, both sexes

2006 4/.....	4.9	*	*	*	*	0.1	0.7	3.5	10.4	19.8	27.6	26.4
2005 4/.....	4.9	*	*	*	*	0.1	0.7	3.5	10.1	20.0	28.0	25.9
2004 4/.....	4.7	*	*	*	*	0.1	0.7	3.3	10.0	19.2	27.1	26.2
2003 4/.....	4.7	*	*	*	*	0.1	0.7	3.2	10.2	19.6	26.7	25.8
2002.....	4.6	*	*	*	*	*	0.7	3.4	9.9	19.8	26.7	25.1
2001.....	4.6	*	*	*	*	0.1	0.7	3.4	9.4	19.8	26.5	26.3
2000.....	4.5	*	*	*	*	0.1	0.7	3.3	10.0	19.5	25.3	26.6
1999.....	4.4	*	*	*	*	0.1	0.6	3.2	9.8	19.4	24.2	27.2

White, male

2006 4/.....	7.9	*	*	*	*	0.1	1.2	5.9	18.6	34.7	50.3	49.9
2005 4/.....	7.9	*	*	*	*	0.2	1.2	6.3	17.8	35.3	51.9	48.5
2004 4/.....	7.5	*	*	*	*	0.1	1.2	5.8	17.6	33.5	49.7	49.2
2003 4/.....	7.4	*	*	*	*	0.1	1.1	5.7	18.0	33.9	48.8	46.7
2002.....	7.3	*	*	*	*	*	1.2	6.0	17.5	35.2	48.8	44.6
2001.....	7.2	*	*	*	*	0.1	1.1	5.9	16.4	34.9	47.9	48.8
2000.....	7.0	*	*	*	*	0.2	1.2	5.9	17.5	33.6	45.6	49.3
1999.....	6.9	*	*	*	*	0.1	1.1	5.7	17.0	34.4	44.0	48.0

White, female

2006 4/.....	2.0	*	*	*	*	*	0.2	1.0	2.6	6.9	11.9	15.3
2005 4/.....	2.0	*	*	*	*	*	0.2	0.8	2.8	6.8	11.6	15.5
2004 4/.....	2.0	*	*	*	*	*	0.2	0.8	2.8	6.9	11.7	15.9
2003 4/.....	2.0	*	*	*	*	*	0.2	0.8	2.7	7.5	11.7	16.5
2002.....	2.0	*	*	*	*	*	0.2	0.8	2.8	6.8	11.9	16.7
2001.....	2.1	*	*	*	*	*	0.2	0.9	2.7	7.1	12.3	16.8
2000.....	2.1	*	*	*	*	*	0.1	0.8	2.9	7.8	11.9	17.3
1999.....	2.0	*	*	*	*	*	0.1	0.7	3.1	6.9	11.2	18.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of esophagus (C15)

Black, both sexes

2006 4/.....	3.7	*	*	*	*	*	0.5	5.0	13.7	22.1	24.5	18.2
2005 4/.....	3.8	*	*	*	*	*	0.9	4.8	14.1	22.5	27.1	17.4
2004 4/.....	4.0	*	*	*	*	*	0.7	5.2	15.6	24.7	27.2	21.4
2003 4/.....	4.2	*	*	*	*	*	0.8	6.0	16.9	25.6	27.9	21.3
2002.....	4.2	*	*	*	*	*	1.0	5.1	17.3	28.5	29.6	20.8
2001.....	4.5	*	*	*	*	*	0.9	5.9	18.5	32.3	29.5	23.6
2000.....	4.5	*	*	*	*	*	0.8	6.6	19.8	28.2	33.7	22.2
1999.....	4.6	*	*	*	*	*	0.9	6.5	20.0	29.9	31.1	23.9

Black, male

2006 4/.....	5.5	*	*	*	*	*	0.9	8.2	22.7	39.9	42.2	34.9
2005 4/.....	5.8	*	*	*	*	*	1.5	7.9	24.4	40.4	51.1	27.0
2004 4/.....	5.8	*	*	*	*	*	1.0	8.3	25.5	41.3	47.9	38.7
2003 4/.....	6.2	*	*	*	*	*	1.3	10.2	28.7	44.3	46.0	38.6
2002.....	6.3	*	*	*	*	*	1.4	8.3	30.1	49.2	51.9	37.5
2001.....	7.0	*	*	*	*	*	1.5	10.2	32.5	55.3	54.7	46.5
2000.....	6.7	*	*	*	*	*	1.2	11.2	34.0	47.0	58.4	38.1
1999.....	7.0	*	*	*	*	*	1.7	11.3	34.8	50.6	56.9	51.6

Black, female

2006 4/.....	2.0	*	*	*	*	*	*	2.3	6.4	9.3	14.5	11.3
2005 4/.....	1.9	*	*	*	*	*	*	2.2	5.7	9.6	13.5	13.5
2004 4/.....	2.3	*	*	*	*	*	*	2.5	7.5	12.9	15.6	14.6
2003 4/.....	2.3	*	*	*	*	*	*	2.5	7.4	12.2	17.7	14.6
2002.....	2.3	*	*	*	*	*	*	2.4	7.0	13.8	16.9	14.4
2001.....	2.3	*	*	*	*	*	*	2.2	7.3	16.1	15.4	15.2
2000.....	2.5	*	*	*	*	*	*	2.7	8.3	14.9	19.8	16.3
1999.....	2.3	*	*	*	*	*	*	2.4	8.2	15.2	16.7	13.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of stomach (C16)

All races 3/, both sexes

2006.....	3.8	*	*	*	*	0.3	1.1	2.7	5.9	13.0	25.6	35.8
2005.....	3.9	*	*	*	0.0	0.3	1.0	2.6	5.9	14.0	26.6	38.5
2004.....	4.0	*	*	*	*	0.3	1.1	2.7	6.5	14.9	26.3	42.2
2003.....	4.2	*	*	*	*	0.3	1.1	3.0	6.7	15.2	27.6	43.6
2002.....	4.2	*	*	*	0.1	0.3	1.0	2.8	6.9	15.4	28.7	47.6
2001.....	4.3	*	*	*	0.1	0.3	1.0	2.9	7.0	16.5	30.2	45.4
2000.....	4.5	*	*	*	*	0.3	1.1	2.9	7.4	16.6	31.0	52.7
1999.....	4.6	*	*	*	*	0.3	1.1	3.0	7.8	17.1	31.6	50.1

All races 3/, male

2006.....	4.5	*	*	*	*	0.3	1.1	3.4	8.3	18.0	36.8	50.2
2005.....	4.6	*	*	*	*	0.3	1.2	3.3	8.2	19.7	37.4	54.2
2004.....	4.8	*	*	*	*	0.3	1.2	3.5	9.1	20.6	37.2	62.9
2003.....	4.9	*	*	*	*	0.3	1.2	3.8	9.5	21.9	38.4	60.0
2002.....	5.0	*	*	*	*	0.4	1.1	3.7	9.5	22.5	40.6	67.2
2001.....	5.1	*	*	*	*	0.4	1.1	3.8	10.2	23.2	43.1	62.6
2000.....	5.3	*	*	*	*	0.3	1.3	3.8	10.5	23.7	43.9	75.6
1999.....	5.5	*	*	*	*	0.4	1.3	4.2	11.0	24.2	46.5	71.3

All races 3/, female

2006.....	3.1	*	*	*	*	0.3	1.1	2.0	3.7	8.8	18.0	29.1
2005.....	3.2	*	*	*	*	0.3	0.8	1.9	3.7	9.2	19.3	31.2
2004.....	3.3	*	*	*	*	0.3	0.9	2.0	4.1	10.1	19.0	32.9
2003.....	3.4	*	*	*	*	0.3	1.0	2.1	4.1	9.7	20.3	36.4
2002.....	3.5	*	*	*	0.1	0.3	0.9	1.9	4.5	9.5	20.8	39.1
2001.....	3.5	*	*	*	*	0.3	0.9	2.0	4.1	10.9	21.7	38.3
2000.....	3.7	*	*	*	*	0.3	0.9	2.1	4.6	10.8	22.6	43.4
1999.....	3.7	*	*	*	*	0.3	1.0	1.9	4.9	11.3	21.9	41.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of stomach (C16)

White, both sexes

2006 4/.....	3.5	*	*	*	*	0.3	0.9	2.3	5.1	11.3	22.4	32.4
2005 4/.....	3.6	*	*	*	*	0.2	0.8	2.2	5.0	12.1	23.3	34.8
2004 4/.....	3.8	*	*	*	*	0.3	0.9	2.3	5.5	12.9	23.7	38.8
2003 4/.....	3.9	*	*	*	*	0.3	0.9	2.4	5.9	13.5	24.6	38.7
2002.....	4.0	*	*	*	0.1	0.3	0.8	2.3	5.9	13.4	25.4	42.2
2001.....	4.1	*	*	*	*	0.3	0.8	2.4	6.1	14.5	26.9	41.1
2000.....	4.2	*	*	*	*	0.2	0.9	2.5	6.5	14.5	27.3	47.3
1999.....	4.3	*	*	*	*	0.3	0.9	2.5	6.6	15.2	28.1	44.8

White, male

2006 4/.....	4.3	*	*	*	*	0.2	0.9	3.0	7.2	15.8	32.5	45.8
2005 4/.....	4.3	*	*	*	*	0.3	1.0	2.8	7.0	16.8	32.9	49.0
2004 4/.....	4.5	*	*	*	*	0.3	1.1	3.0	7.8	17.5	33.2	58.4
2003 4/.....	4.6	*	*	*	*	0.3	1.0	3.1	8.3	19.2	34.3	53.5
2002.....	4.8	*	*	*	*	0.4	0.9	3.1	8.3	19.7	36.6	60.5
2001.....	4.9	*	*	*	*	0.3	1.0	3.2	8.8	20.2	38.7	57.7
2000.....	5.0	*	*	*	*	0.3	1.0	3.2	9.3	20.9	39.0	68.9
1999.....	5.1	*	*	*	*	0.3	1.0	3.5	9.3	21.7	41.5	63.4

White, female

2006 4/.....	2.8	*	*	*	*	0.3	0.9	1.7	3.1	7.4	15.5	26.1
2005 4/.....	3.0	*	*	*	*	0.2	0.7	1.6	3.2	8.1	16.7	28.2
2004 4/.....	3.1	*	*	*	*	0.2	0.8	1.6	3.4	8.8	17.1	29.9
2003 4/.....	3.2	*	*	*	*	0.3	0.8	1.7	3.6	8.7	17.9	32.1
2002.....	3.2	*	*	*	*	0.2	0.7	1.5	3.7	8.0	17.9	34.2
2001.....	3.3	*	*	*	*	0.2	0.7	1.7	3.5	9.7	19.0	34.1
2000.....	3.5	*	*	*	*	0.2	0.7	1.7	3.9	9.2	19.6	38.6
1999.....	3.4	*	*	*	*	0.2	0.8	1.5	4.0	9.8	19.3	37.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of stomach (C16)

Black, both sexes

2006 4/.....	4.8	*	*	*	*	0.4	1.7	4.4	10.9	24.1	51.3	66.9
2005 4/.....	5.0	*	*	*	*	0.4	1.8	4.6	11.2	26.4	52.1	74.3
2004 4/.....	5.3	*	*	*	*	0.5	1.6	5.3	12.8	30.2	50.4	71.3
2003 4/.....	5.4	*	*	*	*	0.4	1.8	5.7	12.7	27.1	54.3	85.8
2002.....	5.6	*	*	*	*	0.6	1.6	5.4	12.9	30.9	56.8	100.4
2001.....	5.6	*	*	*	*	0.5	2.1	5.4	12.9	30.5	58.4	88.1
2000.....	5.9	*	*	*	*	0.7	2.0	5.7	13.8	32.0	63.9	101.7
1999.....	6.1	*	*	*	*	0.7	2.2	5.9	15.9	31.1	64.6	102.4

Black, male

2006 4/.....	5.8	*	*	*	*	*	1.9	6.0	16.2	33.9	80.6	84.0
2005 4/.....	6.2	*	*	*	*	*	2.5	6.4	17.8	40.7	72.1	114.3
2004 4/.....	6.5	*	*	*	*	*	1.9	7.2	19.4	47.3	75.2	100.3
2003 4/.....	6.5	*	*	*	*	*	2.0	8.3	19.6	44.0	79.5	106.3
2002.....	6.6	*	*	*	*	0.8	1.8	7.7	18.9	46.7	79.8	137.2
2001.....	6.7	*	*	*	*	*	2.3	7.9	19.7	48.2	87.1	104.4
2000.....	7.0	*	*	*	*	0.8	2.9	7.9	20.6	46.7	91.1	136.1
1999.....	7.3	*	*	*	*	*	2.7	8.5	23.1	46.0	100.5	147.6

Black, female

2006 4/.....	4.0	*	*	*	*	*	1.6	2.9	6.6	17.1	34.6	60.0
2005 4/.....	4.0	*	*	*	*	*	1.1	2.9	5.9	16.1	40.8	58.1
2004 4/.....	4.2	*	*	*	*	*	1.4	3.6	7.5	18.1	36.3	59.9
2003 4/.....	4.3	*	*	*	*	*	1.6	3.5	7.1	15.0	40.0	77.8
2002.....	4.7	*	*	*	*	*	1.4	3.4	8.0	19.8	43.9	86.3
2001.....	4.5	*	*	*	*	*	1.9	3.2	7.5	18.0	42.3	82.1
2000.....	5.0	*	*	*	*	*	1.3	3.7	8.4	21.6	48.6	88.9
1999.....	5.0	*	*	*	*	*	1.8	3.7	10.1	20.6	44.4	85.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/ 1 year	1-4 2/ years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	-----------------	--------------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races 3/, both sexes

2006.....	17.9	*	*	*	0.1	0.8	3.0	10.4	27.3	61.9	122.8	207.5
2005.....	18.0	*	*	*	0.1	0.8	3.0	10.2	26.9	63.3	124.5	216.9
2004.....	18.3	*	*	*	0.1	0.7	3.0	10.3	28.2	64.7	128.9	225.6
2003.....	19.2	*	*	*	0.1	0.7	3.0	10.9	29.8	70.5	134.7	239.7
2002.....	19.7	*	*	*	0.1	0.7	3.2	11.4	31.7	73.3	137.4	241.4
2001.....	20.0	*	*	*	0.1	0.8	3.2	11.6	31.5	75.3	139.4	255.7
2000.....	20.4	*	*	*	0.1	0.7	3.0	11.2	32.7	76.7	146.0	270.4
1999.....	20.5	*	*	*	0.1	0.7	3.0	11.1	33.5	77.4	146.7	271.7

All races 3/, male

2006.....	18.3	*	*	*	0.1	0.8	3.2	11.6	33.0	76.8	149.5	234.6
2005.....	18.4	*	*	*	0.1	0.8	3.2	11.5	32.9	79.6	151.6	245.9
2004.....	18.6	*	*	*	0.2	0.7	3.2	11.9	34.2	80.6	156.9	254.7
2003.....	19.6	*	*	*	0.1	0.8	3.2	12.5	36.0	88.0	164.4	273.3
2002.....	20.1	*	*	*	0.2	0.8	3.6	13.0	39.4	91.3	167.4	275.1
2001.....	20.2	*	*	*	0.1	0.8	3.5	13.1	38.3	93.7	169.5	298.3
2000.....	20.6	*	*	*	0.1	0.8	3.4	12.8	39.6	95.4	179.5	315.6
1999.....	20.6	*	*	*	0.1	0.8	3.0	12.5	40.5	95.1	181.9	324.0

All races 3/, female

2006.....	17.5	*	*	*	0.1	0.8	2.9	9.2	22.1	49.3	104.6	194.9
2005.....	17.6	*	*	*	0.1	0.7	2.8	9.0	21.2	49.5	106.1	203.6
2004.....	18.0	*	*	*	0.1	0.6	2.9	8.7	22.7	51.4	110.0	212.5
2003.....	18.9	*	*	*	*	0.7	2.7	9.3	24.0	55.9	114.8	224.8
2002.....	19.2	*	*	*	*	0.7	2.8	9.8	24.5	58.3	117.5	226.8
2001.....	19.8	*	*	*	0.1	0.7	2.9	10.2	25.3	60.1	119.6	237.8
2000.....	20.2	*	*	*	*	0.6	2.7	9.8	26.4	61.2	124.2	251.9
1999.....	20.3	*	*	*	0.1	0.6	2.9	9.7	27.0	62.9	123.8	250.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasms of colon, rectum and anus (C18-C21)

White, both sexes

2006 4/.....	18.7	*	*	*	0.1	0.7	2.8	9.7	25.9	60.2	120.5	207.4
2005 4/.....	18.7	*	*	*	0.1	0.7	2.8	9.4	25.2	61.4	122.4	215.7
2004 4/.....	19.2	*	*	*	0.1	0.7	2.8	9.6	26.9	62.8	127.4	225.1
2003 4/.....	20.2	*	*	*	0.1	0.7	2.7	10.0	28.2	68.7	133.3	238.5
2002.....	20.7	*	*	*	0.1	0.7	2.9	10.7	30.1	71.8	135.2	240.7
2001.....	21.0	*	*	*	0.1	0.7	3.0	10.8	30.1	73.4	136.9	255.3
2000.....	21.5	*	*	*	0.1	0.6	2.7	10.4	31.2	75.1	144.0	269.2
1999.....	21.6	*	*	*	0.1	0.7	2.8	10.4	32.0	75.6	144.5	270.9

White, male

2006 4/.....	19.0	*	*	*	*	0.7	2.9	10.8	31.5	74.6	146.2	233.0
2005 4/.....	19.2	*	*	*	0.1	0.7	3.1	10.6	30.9	77.4	149.2	245.3
2004 4/.....	19.5	*	*	*	0.1	0.7	3.0	11.2	32.8	78.2	154.6	255.4
2003 4/.....	20.5	*	*	*	0.1	0.7	3.0	11.5	34.3	86.2	162.8	272.9
2002.....	21.1	*	*	*	0.1	0.8	3.3	12.2	37.6	89.8	164.2	273.6
2001.....	21.2	*	*	*	*	0.7	3.2	12.3	36.8	91.8	167.0	297.3
2000.....	21.6	*	*	*	*	0.7	3.1	12.0	37.6	93.4	176.9	313.3
1999.....	21.8	*	*	*	*	0.7	2.9	11.9	38.8	93.7	179.3	324.5

White, female

2006 4/.....	18.4	*	*	*	*	0.7	2.7	8.6	20.6	47.8	102.6	195.4
2005 4/.....	18.3	*	*	*	*	0.7	2.5	8.1	19.8	47.7	103.9	202.0
2004 4/.....	18.9	*	*	*	*	0.6	2.6	7.9	21.3	49.6	108.9	211.4
2003 4/.....	19.8	*	*	*	*	0.6	2.4	8.4	22.5	53.8	113.4	223.2
2002.....	20.3	*	*	*	*	0.7	2.6	9.1	23.1	56.4	115.7	226.4
2001.....	20.7	*	*	*	*	0.7	2.7	9.3	23.7	57.9	116.8	237.7
2000.....	21.4	*	*	*	*	0.5	2.4	9.0	25.2	59.7	122.3	251.3
1999.....	21.4	*	*	*	*	0.6	2.6	8.9	25.7	60.5	121.7	249.3



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of colon, rectum and anus (C18-C21)

Black, both sexes

2006 4/.....	17.4	*	*	*	*	1.3	5.0	17.1	43.2	89.1	168.5	240.8
2005 4/.....	17.6	*	*	*	*	1.1	4.3	17.7	45.1	91.0	168.4	261.7
2004 4/.....	17.2	*	*	*	*	1.0	4.4	16.6	44.8	94.6	165.1	260.8
2003 4/.....	18.2	*	*	*	*	0.9	5.0	18.7	48.1	100.4	172.0	281.5
2002.....	18.2	*	*	*	*	0.7	4.9	18.6	49.5	99.2	183.6	274.7
2001.....	18.6	*	*	*	*	1.0	4.9	19.0	49.8	104.0	187.6	286.5
2000.....	18.8	*	*	*	*	1.0	5.1	19.5	50.1	104.7	190.1	310.0
1999.....	18.7	*	*	*	*	1.0	4.8	17.7	52.7	106.2	193.1	306.1

Black, male

2006 4/.....	18.2	*	*	*	*	1.2	5.5	20.1	51.7	116.2	219.2	309.2
2005 4/.....	17.9	*	*	*	*	1.4	4.3	19.8	57.4	114.5	210.6	295.6
2004 4/.....	17.4	*	*	*	*	1.0	4.5	19.1	53.4	120.4	214.6	278.2
2003 4/.....	18.3	*	*	*	*	1.0	5.2	22.0	58.9	125.8	213.0	317.0
2002.....	18.6	*	*	*	*	1.0	5.8	21.2	61.2	121.6	237.5	328.1
2001.....	18.4	*	*	*	*	1.2	5.7	20.8	58.4	128.1	228.4	347.2
2000.....	19.1	*	*	*	*	1.0	6.0	22.0	62.5	131.5	241.0	377.1
1999.....	18.6	*	*	*	*	1.2	4.7	20.1	65.8	126.6	244.4	350.4

Black, female

2006 4/.....	16.7	*	*	*	*	1.3	4.6	14.5	36.4	69.6	139.7	212.8
2005 4/.....	17.3	*	*	*	*	0.8	4.2	15.8	35.3	74.1	144.4	248.0
2004 4/.....	17.0	*	*	*	*	0.9	4.3	14.4	37.8	76.2	137.0	254.0
2003 4/.....	18.1	*	*	*	*	0.9	4.8	16.0	39.4	82.3	148.8	267.8
2002.....	17.8	*	*	*	*	*	4.1	16.3	40.2	83.3	153.0	254.2
2001.....	18.8	*	*	*	*	0.9	4.3	17.5	42.8	87.0	164.7	264.0
2000.....	18.5	*	*	*	*	1.0	4.3	17.2	40.2	85.8	161.4	285.0
1999.....	18.9	*	*	*	*	0.9	4.9	15.6	42.2	91.8	164.4	289.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races 3/, both sexes

2006.....	5.5	*	*	*	0.1	0.2	0.8	5.6	12.2	20.2	31.9	33.5
2005.....	5.4	*	0.1	*	0.1	0.2	0.9	5.9	11.3	20.1	32.0	32.5
2004.....	5.2	*	0.1	*	0.1	0.2	0.9	5.6	10.7	19.9	31.3	32.4
2003.....	5.1	*	0.1	*	0.1	0.2	0.9	5.4	9.9	19.8	30.7	33.4
2002.....	4.9	*	0.2	*	0.1	0.3	0.9	4.9	9.6	19.6	29.3	35.0
2001.....	4.7	*	*	*	0.1	0.2	0.9	4.8	9.3	18.8	29.0	34.1
2000.....	4.6	*	0.1	0.0	0.1	0.2	1.0	4.5	9.3	18.7	28.8	32.7
1999.....	4.4	*	0.2	*	0.1	0.2	1.0	4.1	8.8	19.0	28.0	30.6

All races 3/, male

2006.....	7.3	*	*	*	*	0.3	1.1	8.9	19.0	29.3	44.6	43.8
2005.....	7.2	*	*	*	*	0.3	1.2	9.6	17.9	29.0	45.0	44.8
2004.....	6.8	*	*	*	0.1	0.3	1.3	9.2	16.6	28.3	42.7	41.5
2003.....	6.6	*	*	*	0.1	0.3	1.3	8.9	14.9	27.9	44.6	47.5
2002.....	6.4	*	*	*	*	0.3	1.4	8.2	14.5	28.5	41.6	46.2
2001.....	6.1	*	*	*	*	0.2	1.3	7.7	13.9	26.8	41.0	50.3
2000.....	5.9	*	*	*	0.1	0.3	1.3	7.2	13.8	27.0	42.1	44.7
1999.....	5.6	*	0.3	*	0.1	0.2	1.4	6.5	12.9	27.5	39.0	40.1

All races 3/, female

2006.....	3.8	*	*	*	*	0.2	0.5	2.3	5.8	12.6	23.2	28.7
2005.....	3.7	*	*	*	*	0.1	0.6	2.4	5.2	12.6	23.2	26.8
2004.....	3.7	*	*	*	*	0.1	0.4	2.2	5.3	12.8	23.6	28.3
2003.....	3.5	*	*	*	*	0.2	0.5	2.1	5.3	13.0	21.4	27.2
2002.....	3.4	*	*	*	*	0.2	0.5	1.8	5.1	12.2	21.1	30.2
2001.....	3.4	*	*	*	*	0.2	0.4	1.9	5.0	12.1	21.1	27.2
2000.....	3.3	*	*	*	*	0.1	0.6	1.9	5.1	11.8	20.1	27.8
1999.....	3.3	*	*	*	*	0.2	0.6	1.8	5.1	11.9	20.9	26.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

White, both sexes

2006 4/.....	5.4	*	*	*	0.1	0.2	0.6	4.8	10.7	18.9	30.6	33.1
2005 4/.....	5.3	*	*	*	*	0.1	0.7	5.2	9.9	18.6	30.7	31.7
2004 4/.....	5.1	*	*	*	0.1	0.1	0.7	4.8	9.4	18.3	30.0	31.2
2003 4/.....	4.9	*	*	*	0.1	0.1	0.7	4.6	8.7	18.3	29.5	32.1
2002.....	4.8	*	0.2	*	*	0.2	0.8	4.1	8.5	18.3	28.4	33.8
2001.....	4.6	*	*	*	0.1	0.2	0.6	4.0	8.2	17.6	27.9	33.3
2000.....	4.5	*	*	*	0.1	0.1	0.7	3.7	8.1	17.3	27.8	32.0
1999.....	4.4	*	0.2	*	0.1	0.1	0.8	3.4	7.6	17.8	27.2	29.9

White, male

2006 4/.....	7.0	*	*	*	*	0.1	0.8	7.7	16.2	27.0	43.2	43.7
2005 4/.....	7.0	*	*	*	*	0.2	0.9	8.3	15.4	26.6	43.2	45.0
2004 4/.....	6.5	*	*	*	*	0.2	1.0	7.9	14.2	25.9	41.4	39.9
2003 4/.....	6.4	*	*	*	*	0.1	1.0	7.4	13.1	26.0	43.0	45.6
2002.....	6.1	*	*	*	*	0.2	1.1	6.8	12.6	26.5	40.0	45.1
2001.....	5.9	*	*	*	*	0.1	0.9	6.4	12.2	25.4	39.6	49.7
2000.....	5.7	*	*	*	*	0.2	1.0	5.7	11.8	25.0	41.1	43.1
1999.....	5.5	*	*	*	*	0.1	1.1	5.2	11.1	25.8	38.0	40.6

White, female

2006 4/.....	3.8	*	*	*	*	0.2	0.5	2.0	5.4	11.9	21.9	28.1
2005 4/.....	3.7	*	*	*	*	*	0.5	2.2	4.7	11.7	22.1	25.6
2004 4/.....	3.7	*	*	*	*	*	0.4	1.9	4.9	11.8	22.3	27.3
2003 4/.....	3.5	*	*	*	*	0.1	0.5	1.8	4.6	11.7	20.3	26.2
2002.....	3.5	*	*	*	*	0.2	0.4	1.5	4.5	11.4	20.6	29.0
2001.....	3.3	*	*	*	*	0.2	0.3	1.6	4.4	10.9	20.1	26.5
2000.....	3.3	*	*	*	*	*	0.4	1.7	4.6	10.9	19.0	27.4
1999.....	3.3	*	*	*	*	*	0.5	1.6	4.4	11.0	20.1	25.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

Black, both sexes

2006 4/.....	5.5	*	*	*	*	0.4	1.3	9.4	21.8	24.2	33.4	31.1
2005 4/.....	5.4	*	*	*	*	0.5	1.5	9.6	20.2	24.9	33.2	29.1
2004 4/.....	5.2	*	*	*	*	0.4	1.4	9.6	17.4	25.6	34.4	36.1
2003 4/.....	4.9	*	*	*	*	0.5	1.4	9.8	16.3	22.9	31.3	41.7
2002.....	4.6	*	*	*	*	0.5	1.4	8.7	14.8	23.0	30.9	38.0
2001.....	4.5	*	*	*	*	*	1.7	8.7	14.8	21.7	32.4	35.0
2000.....	4.4	*	*	*	*	0.4	1.8	8.6	14.1	23.6	28.7	31.6
1999.....	4.2	*	*	*	*	0.4	1.6	7.2	13.2	24.2	29.4	34.4

Black, male

2006 4/.....	8.0	*	*	*	*	*	2.1	15.4	39.5	39.4	45.5	34.9
2005 4/.....	7.9	*	*	*	*	0.7	2.2	16.9	35.7	39.6	50.8	31.6
2004 4/.....	7.3	*	*	*	*	*	2.2	16.7	29.9	40.3	44.0	46.7
2003 4/.....	6.8	*	*	*	*	*	2.2	17.1	25.9	33.5	44.3	63.6
2002.....	6.5	*	*	*	*	*	2.2	15.1	24.2	37.7	47.0	41.8
2001.....	5.9	*	*	*	*	*	2.6	14.8	23.6	30.0	43.0	42.0
2000.....	6.1	*	*	*	*	*	2.4	14.8	23.5	37.3	39.7	48.4
1999.....	5.5	*	*	*	*	*	2.4	12.8	20.0	36.5	41.2	30.5

Black, female

2006 4/.....	3.3	*	*	*	*	*	*	4.3	7.5	13.4	26.5	29.6
2005 4/.....	3.1	*	*	*	*	*	0.9	3.3	7.7	14.4	23.3	28.1
2004 4/.....	3.3	*	*	*	*	*	*	3.5	7.3	15.0	29.0	31.9
2003 4/.....	3.2	*	*	*	*	*	*	3.5	8.6	15.3	23.9	33.2
2002.....	2.9	*	*	*	*	*	0.7	3.2	7.3	12.6	21.8	36.6
2001.....	3.2	*	*	*	*	*	0.8	3.5	7.7	15.9	26.4	32.4
2000.....	2.8	*	*	*	*	*	1.2	3.3	6.5	14.0	22.5	25.3
1999.....	3.0	*	*	*	*	*	0.9	2.4	7.8	15.6	22.7	35.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of pancreas (C25)

All races 3/, both sexes

2006.....	11.2	*	*	*	*	0.1	1.2	6.3	19.9	44.8	78.6	96.9
2005.....	11.1	*	*	*	*	0.2	1.3	6.5	19.7	44.5	77.7	97.4
2004.....	10.8	*	*	*	*	0.2	1.2	6.3	19.8	44.3	76.0	97.7
2003.....	10.6	*	*	*	*	0.2	1.2	6.2	19.1	44.2	75.4	95.2
2002.....	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.8	75.0	94.5
2001.....	10.5	*	*	*	*	0.1	1.3	6.1	19.2	44.5	74.1	100.1
2000.....	10.4	*	*	*	*	0.1	1.2	6.3	19.4	44.1	76.5	96.4
1999.....	10.4	*	*	*	*	0.1	1.3	6.4	19.8	45.0	73.1	99.7

All races 3/, male

2006.....	11.2	*	*	*	*	0.1	1.4	7.8	24.5	52.3	86.2	102.1
2005.....	11.1	*	*	*	*	0.2	1.5	7.7	24.3	51.8	86.0	102.8
2004.....	10.9	*	*	*	*	0.2	1.5	8.1	24.1	52.6	84.6	102.0
2003.....	10.5	*	*	*	*	0.2	1.5	7.7	23.1	50.9	85.4	95.6
2002.....	10.5	*	*	*	*	0.1	1.6	7.8	23.6	51.4	84.0	100.9
2001.....	10.3	*	*	*	*	0.1	1.4	7.6	23.6	50.8	84.0	105.1
2000.....	10.3	*	*	*	*	0.2	1.4	8.0	23.6	50.4	85.9	102.9
1999.....	10.4	*	*	*	*	0.2	1.6	8.3	23.9	52.4	81.8	107.5

All races 3/, female

2006.....	11.1	*	*	*	*	0.1	0.9	4.9	15.7	38.4	73.4	94.4
2005.....	11.0	*	*	*	*	0.1	1.1	5.2	15.5	38.2	72.1	94.9
2004.....	10.7	*	*	*	*	0.1	0.9	4.7	15.8	37.3	70.1	95.8
2003.....	10.6	*	*	*	*	0.1	1.0	4.8	15.3	38.6	68.8	95.0
2002.....	10.5	*	*	*	*	*	1.0	4.4	15.1	39.3	69.1	91.7
2001.....	10.6	*	*	*	*	0.1	1.1	4.7	15.2	39.4	67.6	98.1
2000.....	10.5	*	*	*	*	0.1	1.0	4.6	15.5	38.8	70.4	93.8
1999.....	10.5	*	*	*	*	0.1	1.0	4.7	16.0	39.0	67.4	96.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of pancreas (C25)

White, both sexes

2006 4/.....	11.8	*	*	*	*	0.1	1.1	6.1	19.5	44.2	78.5	96.1
2005 4/.....	11.7	*	*	*	*	0.1	1.3	6.2	19.3	44.0	77.0	96.8
2004 4/.....	11.4	*	*	*	*	0.1	1.1	6.1	19.1	43.5	75.6	97.1
2003 4/.....	11.2	*	*	*	*	0.2	1.2	5.9	18.4	43.7	74.8	94.4
2002.....	11.1	*	*	*	*	0.1	1.3	5.8	18.7	43.7	74.3	94.3
2001.....	11.1	*	*	*	*	0.1	1.3	5.9	19.0	43.8	73.3	99.0
2000.....	11.0	*	*	*	*	0.1	1.2	6.1	18.9	43.1	75.3	94.7
1999.....	11.0	*	*	*	*	0.1	1.3	6.2	19.1	44.2	71.7	98.6

White, male

2006 4/.....	12.0	*	*	*	*	*	1.4	7.6	24.1	52.1	87.1	103.4
2005 4/.....	11.7	*	*	*	*	0.2	1.5	7.4	23.8	51.8	85.6	102.9
2004 4/.....	11.5	*	*	*	*	0.2	1.4	7.7	23.4	51.6	85.1	102.9
2003 4/.....	11.2	*	*	*	*	0.2	1.4	7.2	22.5	50.8	85.4	96.0
2002.....	11.1	*	*	*	*	*	1.6	7.5	23.0	50.7	84.2	101.6
2001.....	11.0	*	*	*	*	0.2	1.4	7.4	23.5	50.0	83.8	106.5
2000.....	10.9	*	*	*	*	0.2	1.4	7.8	23.0	49.8	85.2	102.4
1999.....	10.9	*	*	*	*	0.2	1.6	7.9	22.9	51.6	80.8	107.4

White, female

2006 4/.....	11.7	*	*	*	*	0.1	0.9	4.6	15.2	37.4	72.5	92.7
2005 4/.....	11.7	*	*	*	*	*	1.0	5.0	15.0	37.4	71.2	94.0
2004 4/.....	11.3	*	*	*	*	*	0.9	4.4	15.1	36.7	69.1	94.5
2003 4/.....	11.2	*	*	*	*	0.1	1.0	4.5	14.6	37.6	67.7	93.8
2002.....	11.0	*	*	*	*	*	1.0	4.2	14.5	37.8	67.7	91.1
2001.....	11.2	*	*	*	*	*	1.1	4.5	14.7	38.5	66.3	95.9
2000.....	11.1	*	*	*	*	*	0.9	4.4	15.0	37.5	68.8	91.5
1999.....	11.0	*	*	*	*	*	0.9	4.6	15.5	37.9	65.8	95.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of pancreas (C25)

Black, both sexes

2006 4/.....	9.8	*	*	*	*	*	1.6	9.1	27.9	57.2	91.2	113.4
2005 4/.....	9.7	*	*	*	*	*	1.5	9.6	26.9	56.7	94.3	111.4
2004 4/.....	9.5	*	*	*	*	*	1.6	9.3	28.4	57.9	88.0	113.6
2003 4/.....	9.5	*	*	*	*	*	1.8	10.0	27.3	56.3	93.1	112.9
2002.....	9.3	*	*	*	*	*	1.8	8.7	27.6	60.7	89.9	106.6
2001.....	9.1	*	*	*	*	*	1.6	9.0	25.5	58.4	92.7	121.3
2000.....	9.3	*	*	*	*	*	1.7	9.0	27.5	59.1	97.8	119.8
1999.....	9.4	*	*	*	*	*	2.0	9.6	28.2	59.5	94.0	118.6

Black, male

2006 4/.....	9.4	*	*	*	*	*	1.9	11.2	35.4	66.0	91.5	91.1
2005 4/.....	9.4	*	*	*	*	*	1.6	11.7	33.9	64.5	104.1	100.4
2004 4/.....	9.4	*	*	*	*	*	2.0	11.4	36.1	72.3	89.6	100.3
2003 4/.....	9.2	*	*	*	*	*	2.4	13.3	33.2	63.2	97.9	104.3
2002.....	9.0	*	*	*	*	*	2.1	11.4	34.1	68.4	96.1	100.8
2001.....	8.6	*	*	*	*	*	1.7	11.1	30.3	68.1	100.3	99.8
2000.....	9.0	*	*	*	*	*	2.0	11.8	34.2	65.8	106.9	109.6
1999.....	9.4	*	*	*	*	*	2.4	13.1	36.5	67.9	106.7	114.8

Black, female

2006 4/.....	10.2	*	*	*	*	*	1.2	7.3	21.8	50.9	91.1	122.5
2005 4/.....	9.9	*	*	*	*	*	1.4	7.8	21.2	51.2	88.7	115.9
2004 4/.....	9.6	*	*	*	*	*	1.2	7.5	22.2	47.7	87.0	118.9
2003 4/.....	9.7	*	*	*	*	*	1.3	7.2	22.5	51.5	90.4	116.3
2002.....	9.5	*	*	*	*	*	1.4	6.4	22.5	55.3	86.4	108.8
2001.....	9.6	*	*	*	*	*	1.4	7.1	21.6	51.6	88.4	129.2
2000.....	9.7	*	*	*	*	*	1.4	6.6	22.1	54.4	92.7	123.6
1999.....	9.4	*	*	*	*	*	1.6	6.6	21.6	53.6	86.9	120.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of larynx (C32)

All races 3/, both sexes

2006.....	1.3	*	*	*	*	*	0.1	1.0	3.1	5.7	7.1	6.3
2005.....	1.3	*	*	*	*	*	0.1	1.0	3.0	6.2	7.0	6.3
2004.....	1.2	*	*	*	*	*	0.1	1.1	3.2	5.7	6.6	6.5
2003.....	1.3	*	*	*	*	*	0.2	1.0	3.4	6.2	6.9	6.5
2002.....	1.3	*	*	*	*	*	0.2	1.1	3.4	5.9	7.6	5.5
2001.....	1.3	*	*	*	*	*	0.2	1.1	3.6	6.4	7.2	6.7
2000.....	1.4	*	*	*	*	*	0.1	1.1	3.6	6.6	8.1	6.5
1999.....	1.4	*	*	*	*	*	0.2	1.1	3.8	6.7	7.3	7.0

All races 3/, male

2006.....	2.0	*	*	*	*	*	0.2	1.7	5.1	9.8	13.4	14.8
2005.....	2.0	*	*	*	*	*	0.2	1.5	5.1	10.7	13.1	15.7
2004.....	2.0	*	*	*	*	*	0.2	1.8	5.4	10.1	13.0	15.6
2003.....	2.1	*	*	*	*	*	0.3	1.8	5.8	10.8	13.7	15.3
2002.....	2.1	*	*	*	*	*	0.3	1.8	5.8	10.0	14.9	13.2
2001.....	2.2	*	*	*	*	*	0.3	1.8	6.1	11.1	14.2	18.3
2000.....	2.2	*	*	*	*	*	0.2	1.8	6.2	11.4	15.7	17.4
1999.....	2.2	*	*	*	*	*	0.3	1.9	6.4	11.4	14.1	17.8

All races 3/, female

2006.....	0.5	*	*	*	*	*	*	0.4	1.2	2.2	2.8	2.3
2005.....	0.5	*	*	*	*	*	*	0.5	1.0	2.4	2.8	1.9
2004.....	0.5	*	*	*	*	*	*	0.4	1.2	2.0	2.3	2.4
2003.....	0.5	*	*	*	*	*	*	0.3	1.2	2.4	2.4	2.7
2002.....	0.5	*	*	*	*	*	*	0.4	1.2	2.5	2.8	2.2
2001.....	0.5	*	*	*	*	*	0.1	0.4	1.3	2.5	2.6	1.9
2000.....	0.6	*	*	*	*	*	*	0.4	1.2	2.7	3.1	2.1
1999.....	0.6	*	*	*	*	*	*	0.3	1.4	2.9	2.8	2.6



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of larynx (C32)

White, both sexes

2006 4/.....	1.3	*	*	*	*	*	0.1	0.9	2.9	5.4	7.0	6.2
2005 4/.....	1.3	*	*	*	*	*	0.1	0.9	2.6	5.8	7.0	6.0
2004 4/.....	1.3	*	*	*	*	*	0.1	0.9	3.0	5.4	6.4	6.3
2003 4/.....	1.3	*	*	*	*	*	0.2	0.9	3.1	5.8	6.6	6.2
2002.....	1.3	*	*	*	*	*	0.1	0.9	3.1	5.7	7.3	5.6
2001.....	1.3	*	*	*	*	*	0.2	0.9	3.2	6.0	6.9	6.6
2000.....	1.3	*	*	*	*	*	0.1	0.9	3.1	6.1	7.7	6.6
1999.....	1.4	*	*	*	*	*	0.1	0.9	3.4	6.4	7.2	6.7

White, male

2006 4/.....	2.0	*	*	*	*	*	0.1	1.5	4.7	9.0	13.0	14.7
2005 4/.....	2.0	*	*	*	*	*	0.2	1.4	4.4	9.9	12.9	14.9
2004 4/.....	2.0	*	*	*	*	*	0.2	1.5	4.8	9.5	12.4	15.0
2003 4/.....	2.1	*	*	*	*	*	0.3	1.5	5.2	9.8	12.9	14.5
2002.....	2.0	*	*	*	*	*	0.2	1.5	5.1	9.5	14.1	13.3
2001.....	2.1	*	*	*	*	*	0.3	1.4	5.4	10.3	13.3	18.3
2000.....	2.1	*	*	*	*	*	0.2	1.5	5.3	10.4	15.0	17.6
1999.....	2.1	*	*	*	*	*	0.2	1.6	5.7	10.7	13.8	16.8

White, female

2006 4/.....	0.6	*	*	*	*	*	*	0.4	1.1	2.2	2.9	2.2
2005 4/.....	0.6	*	*	*	*	*	*	0.4	1.0	2.3	2.9	1.9
2004 4/.....	0.5	*	*	*	*	*	*	0.4	1.2	1.9	2.3	2.5
2003 4/.....	0.5	*	*	*	*	*	*	0.3	1.1	2.4	2.4	2.5
2002.....	0.6	*	*	*	*	*	*	0.3	1.2	2.5	2.6	2.3
2001.....	0.5	*	*	*	*	*	*	0.3	1.2	2.3	2.7	1.7
2000.....	0.6	*	*	*	*	*	*	0.4	1.1	2.6	2.9	2.1
1999.....	0.6	*	*	*	*	*	*	0.3	1.2	2.8	2.9	2.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of larynx (C32)

Black, both sexes

2006 4/.....	1.6	*	*	*	*	*	*	2.1	5.9	10.3	9.9	8.0
2005 4/.....	1.7	*	*	*	*	*	*	2.2	7.0	11.5	9.9	10.2
2004 4/.....	1.7	*	*	*	*	*	*	2.3	6.6	10.3	11.2	8.2
2003 4/.....	1.8	*	*	*	*	*	*	2.5	7.5	11.5	11.7	11.1
2002.....	1.8	*	*	*	*	*	0.5	2.6	7.5	10.3	13.7	*
2001.....	1.8	*	*	*	*	*	0.4	2.8	7.9	11.4	11.7	8.3
2000.....	2.0	*	*	*	*	*	*	2.9	9.0	12.8	14.1	6.9
1999.....	1.8	*	*	*	*	*	0.6	2.9	8.6	11.6	10.0	8.3

Black, male

2006 4/.....	2.7	*	*	*	*	*	*	3.9	10.6	20.6	21.9	18.8
2005 4/.....	2.9	*	*	*	*	*	*	3.4	13.6	22.1	21.3	28.8
2004 4/.....	2.9	*	*	*	*	*	*	4.2	12.6	20.4	24.8	23.8
2003 4/.....	3.1	*	*	*	*	*	*	4.5	13.6	23.8	27.9	27.1
2002.....	3.0	*	*	*	*	*	0.7	4.5	14.6	19.3	29.3	*
2001.....	3.1	*	*	*	*	*	*	5.1	14.2	21.3	27.7	*
2000.....	3.3	*	*	*	*	*	*	5.1	17.2	24.3	28.5	*
1999.....	3.1	*	*	*	*	*	0.9	5.2	15.9	21.8	22.8	24.6

Black, female

2006 4/.....	0.6	*	*	*	*	*	*	*	2.1	2.9	3.1	*
2005 4/.....	0.6	*	*	*	*	*	*	1.1	1.7	3.9	3.4	*
2004 4/.....	0.5	*	*	*	*	*	*	*	1.7	3.2	3.5	*
2003 4/.....	0.6	*	*	*	*	*	*	0.8	2.5	2.8	*	*
2002.....	0.7	*	*	*	*	*	*	1.1	1.8	4.0	4.9	*
2001.....	0.7	*	*	*	*	*	*	*	2.8	4.5	*	*
2000.....	0.7	*	*	*	*	*	*	1.0	2.4	4.6	6.0	*
1999.....	0.7	*	*	*	*	*	*	0.9	2.7	4.4	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

All races 3/, both sexes

2006.....	53.0	*	*	*	0.1	0.4	4.6	29.1	99.1	253.1	373.5	300.5
2005.....	53.7	*	*	*	*	0.3	5.3	29.7	103.3	259.6	375.6	302.3
2004.....	53.8	*	*	*	0.1	0.4	5.5	29.9	106.9	265.5	373.6	297.5
2003.....	54.4	*	*	*	0.1	0.4	5.6	30.3	111.0	269.3	377.8	298.9
2002.....	54.7	*	*	*	0.1	0.4	6.0	30.3	115.3	275.0	377.6	297.2
2001.....	54.8	*	*	*	0.1	0.4	6.2	30.7	117.7	279.7	371.4	302.7
2000.....	55.3	*	*	*	0.1	0.5	6.1	31.6	122.4	284.2	370.8	302.1
1999.....	54.5	*	*	*	0.1	0.4	6.0	31.3	123.4	281.5	362.1	297.0

All races 3/, male

2006.....	60.5	*	*	*	*	0.4	4.7	33.8	121.6	319.4	509.9	457.7
2005.....	61.8	*	*	*	*	0.4	5.5	35.1	127.6	330.7	515.1	468.0
2004.....	62.0	*	*	*	*	0.4	5.8	35.8	131.7	337.6	517.3	466.8
2003.....	62.9	*	*	*	*	0.4	6.1	36.5	136.7	346.6	525.1	475.1
2002.....	63.7	*	*	*	*	0.4	6.2	36.6	144.0	355.9	527.9	482.2
2001.....	64.7	*	*	*	*	0.4	6.7	37.4	147.1	370.0	529.5	512.4
2000.....	65.5	*	*	*	*	0.5	6.9	38.5	154.0	377.9	532.2	521.2
1999.....	65.4	*	*	*	*	0.5	6.6	38.2	158.0	377.2	527.6	529.2

All races 3/, female

2006.....	45.7	*	*	*	*	0.4	4.5	24.6	78.2	197.0	280.3	226.9
2005.....	45.9	*	*	*	*	0.3	5.1	24.5	80.7	199.6	280.9	226.2
2004.....	45.9	*	*	*	*	0.3	5.2	24.2	83.9	205.0	277.0	221.3
2003.....	46.1	*	*	*	*	0.4	5.1	24.4	87.1	204.8	279.4	221.0
2002.....	46.0	*	*	*	*	0.4	5.8	24.3	88.8	207.7	277.8	217.0
2001.....	45.3	*	*	*	*	0.4	5.7	24.3	90.5	204.9	267.5	215.0
2000.....	45.4	*	*	*	*	0.5	5.3	25.0	93.3	206.9	265.6	212.8
1999.....	44.1	*	*	*	*	0.4	5.5	24.6	91.5	203.1	254.5	203.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

White, both sexes

2006 4/.....	57.3	*	*	*	*	0.4	4.6	27.8	98.9	258.3	381.3	306.1
2005 4/.....	58.1	*	*	*	*	0.3	5.2	28.6	102.5	264.8	382.8	305.6
2004 4/.....	58.1	*	*	*	*	0.3	5.3	28.7	106.3	270.4	378.6	302.0
2003 4/.....	58.7	*	*	*	0.1	0.4	5.4	28.9	110.8	274.2	382.6	302.4
2002.....	59.1	*	*	*	*	0.4	5.8	29.1	114.5	280.0	382.9	300.9
2001.....	59.1	*	*	*	*	0.4	6.0	29.6	117.8	283.4	374.7	305.4
2000.....	59.5	*	*	*	*	0.5	5.7	30.2	122.5	287.0	374.7	304.5
1999.....	58.5	*	*	*	*	0.5	5.8	29.7	122.9	283.5	363.8	298.8

White, male

2006 4/.....	64.6	*	*	*	*	0.4	4.6	31.7	118.3	319.8	514.6	464.0
2005 4/.....	65.9	*	*	*	*	0.4	5.3	33.3	123.4	331.8	519.9	469.9
2004 4/.....	65.9	*	*	*	*	0.4	5.5	33.3	127.6	338.1	517.5	472.2
2003 4/.....	66.9	*	*	*	*	0.4	5.7	34.1	133.1	347.1	524.6	478.5
2002.....	67.7	*	*	*	*	0.4	5.8	34.2	139.3	356.4	527.8	486.6
2001.....	68.6	*	*	*	*	0.4	6.3	34.9	143.8	368.1	526.4	513.9
2000.....	69.4	*	*	*	*	0.5	6.1	35.7	150.8	374.9	529.9	522.4
1999.....	69.1	*	*	*	*	0.5	6.1	35.2	153.9	373.4	523.5	531.5

White, female

2006 4/.....	50.2	*	*	*	*	0.3	4.5	24.0	80.5	205.2	288.8	231.9
2005 4/.....	50.4	*	*	*	*	0.3	5.2	24.0	82.7	207.2	288.4	229.9
2004 4/.....	50.4	*	*	*	*	0.3	5.2	24.1	86.1	212.5	283.8	225.3
2003 4/.....	50.7	*	*	*	*	0.4	5.1	23.8	89.8	212.2	286.6	224.4
2002.....	50.7	*	*	*	*	0.4	5.8	24.2	91.0	215.1	285.8	220.2
2001.....	49.8	*	*	*	*	0.4	5.7	24.3	93.3	211.9	274.0	218.1
2000.....	49.9	*	*	*	*	0.5	5.2	24.8	96.1	213.2	272.7	215.9
1999.....	48.2	*	*	*	*	0.4	5.4	24.4	93.8	208.6	259.1	205.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of trachea, bronchus and lung (C33-C34)

Black, both sexes

2006 4/.....	41.7	*	*	*	*	0.6	6.1	45.0	128.0	272.3	364.6	264.6
2005 4/.....	42.4	*	*	*	*	*	6.8	44.4	137.5	278.1	366.4	289.2
2004 4/.....	42.7	*	*	*	*	0.5	7.7	45.4	141.6	279.6	384.4	277.5
2003 4/.....	42.9	*	*	*	*	*	7.7	47.4	144.0	288.6	383.1	280.9
2002.....	43.2	*	*	*	*	0.4	8.7	46.9	152.4	295.3	381.3	272.3
2001.....	43.1	*	*	*	*	*	8.9	46.5	147.0	306.5	385.4	280.7
2000.....	43.9	*	*	*	*	0.6	10.1	50.4	152.5	315.8	375.0	288.1
1999.....	44.1	*	*	*	*	0.4	9.3	50.6	158.1	313.1	387.9	290.5

Black, male

2006 4/.....	52.0	*	*	*	*	*	6.6	56.9	184.3	402.7	563.9	442.3
2005 4/.....	53.0	*	*	*	*	*	7.6	56.9	199.1	408.8	565.2	495.5
2004 4/.....	54.3	*	*	*	*	*	8.8	62.5	206.9	410.4	611.1	481.8
2003 4/.....	54.9	*	*	*	*	*	9.3	63.6	210.0	427.1	622.2	502.5
2002.....	56.0	*	*	*	*	*	10.6	64.6	223.6	444.6	626.2	484.6
2001.....	57.2	*	*	*	*	*	11.0	66.2	215.2	476.4	659.8	549.2
2000.....	58.3	*	*	*	*	*	13.4	70.7	223.5	488.8	642.5	562.8
1999.....	58.9	*	*	*	*	*	11.3	72.6	237.6	484.4	661.0	574.2

Black, female

2006 4/.....	32.3	*	*	*	*	*	5.7	34.7	82.6	179.1	251.3	191.9
2005 4/.....	32.7	*	*	*	*	*	6.1	33.6	87.9	184.5	253.5	205.9
2004 4/.....	32.1	*	*	*	*	*	6.8	30.7	89.0	186.2	256.3	196.5
2003 4/.....	32.0	*	*	*	*	*	6.2	33.4	91.0	190.0	248.0	194.8
2002.....	31.6	*	*	*	*	*	7.0	31.7	95.3	189.3	242.6	191.0
2001.....	30.2	*	*	*	*	*	7.1	29.5	92.0	186.7	231.3	181.0
2000.....	30.8	*	*	*	*	*	7.3	32.9	95.3	194.1	224.3	185.9
1999.....	30.7	*	*	*	*	*	7.6	31.9	94.4	192.3	234.9	184.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant melanoma of skin (C43)

All races 3/, both sexes

2006.....	2.8	*	*	*	0.1	0.5	1.2	2.7	5.2	9.4	15.6	20.1
2005.....	2.8	*	*	*	0.1	0.5	1.2	2.8	5.2	9.4	15.6	19.8
2004.....	2.7	*	*	*	0.1	0.5	1.2	2.8	5.3	9.3	14.7	17.7
2003.....	2.7	*	*	*	0.1	0.5	1.3	2.9	5.5	8.7	14.6	17.6
2002.....	2.6	*	*	*	0.1	0.5	1.3	2.8	5.1	8.8	13.7	17.9
2001.....	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.2	13.3	17.8
2000.....	2.6	*	*	*	0.1	0.5	1.4	3.2	5.4	8.9	13.1	18.9
1999.....	2.6	*	*	*	0.1	0.5	1.4	3.1	5.3	9.0	13.4	16.4

All races 3/, male

2006.....	3.7	*	*	*	0.1	0.5	1.4	3.5	7.4	13.7	25.6	36.7
2005.....	3.6	*	*	*	0.1	0.7	1.4	3.6	7.0	13.8	24.5	34.9
2004.....	3.5	*	*	*	*	0.6	1.4	3.7	7.6	13.7	23.7	31.4
2003.....	3.5	*	*	*	0.1	0.6	1.5	3.8	7.8	12.6	23.3	30.2
2002.....	3.3	*	*	*	0.1	0.6	1.6	3.8	7.2	13.0	21.2	29.2
2001.....	3.4	*	*	*	0.1	0.5	1.7	3.9	7.3	13.9	21.3	29.9
2000.....	3.3	*	*	*	0.1	0.5	1.6	4.2	7.4	12.9	20.8	31.8
1999.....	3.3	*	*	*	0.1	0.5	1.8	4.0	7.5	13.2	20.9	27.5

All races 3/, female

2006.....	2.0	*	*	*	*	0.4	0.9	1.9	3.2	5.8	8.9	12.4
2005.....	2.0	*	*	*	*	0.4	1.0	2.1	3.5	5.6	9.5	12.9
2004.....	1.9	*	*	*	0.1	0.4	1.0	2.0	3.2	5.5	8.7	11.5
2003.....	1.9	*	*	*	*	0.4	1.0	2.1	3.5	5.4	8.7	12.1
2002.....	1.9	*	*	*	0.1	0.4	1.1	1.9	3.2	5.4	8.8	12.9
2001.....	1.9	*	*	*	*	0.4	1.1	2.3	3.3	5.3	8.1	12.8
2000.....	2.0	*	*	*	0.1	0.4	1.2	2.2	3.5	5.7	8.2	13.7
1999.....	1.9	*	*	*	*	0.4	1.0	2.2	3.3	5.5	8.5	11.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant melanoma of skin (C43)

White, both sexes

2006 4/.....	3.4	*	*	*	0.1	0.6	1.4	3.2	6.0	10.8	17.2	21.7
2005 4/.....	3.4	*	*	*	0.1	0.7	1.4	3.3	5.9	10.7	17.2	21.4
2004 4/.....	3.3	*	*	*	0.1	0.6	1.4	3.3	6.1	10.5	16.2	19.2
2003 4/.....	3.2	*	*	*	0.1	0.6	1.5	3.4	6.3	9.8	16.0	19.1
2002.....	3.1	*	*	*	0.1	0.6	1.6	3.3	5.8	9.9	15.1	19.2
2001.....	3.2	*	*	*	0.1	0.6	1.7	3.6	6.1	10.3	14.6	19.3
2000.....	3.1	*	*	*	0.1	0.6	1.6	3.7	6.1	9.9	14.3	20.3
1999.....	3.1	*	*	*	0.1	0.6	1.7	3.6	6.1	10.0	14.5	17.6

White, male

2006 4/.....	4.5	*	*	*	0.2	0.7	1.7	4.1	8.4	15.6	28.1	39.7
2005 4/.....	4.4	*	*	*	0.1	0.8	1.7	4.2	8.0	15.7	27.0	38.1
2004 4/.....	4.3	*	*	*	*	0.7	1.7	4.3	8.7	15.6	26.0	34.2
2003 4/.....	4.2	*	*	*	0.1	0.7	1.8	4.4	8.8	14.2	25.5	32.9
2002.....	4.0	*	*	*	*	0.7	1.9	4.4	8.1	14.4	23.3	31.7
2001.....	4.1	*	*	*	0.1	0.6	2.0	4.6	8.4	15.5	23.3	32.7
2000.....	4.0	*	*	*	0.1	0.7	1.9	4.8	8.5	14.3	22.5	34.7
1999.....	4.0	*	*	*	0.1	0.6	2.1	4.6	8.5	14.7	22.6	29.7

White, female

2006 4/.....	2.3	*	*	*	*	0.5	1.1	2.3	3.7	6.6	9.7	13.3
2005 4/.....	2.4	*	*	*	*	0.5	1.2	2.5	4.0	6.4	10.4	13.8
2004 4/.....	2.3	*	*	*	0.1	0.5	1.2	2.3	3.8	6.2	9.5	12.4
2003 4/.....	2.3	*	*	*	*	0.5	1.2	2.4	4.0	6.1	9.5	12.9
2002.....	2.3	*	*	*	0.1	0.5	1.3	2.3	3.6	6.1	9.6	13.8
2001.....	2.3	*	*	*	*	0.5	1.3	2.7	3.8	5.9	8.8	13.7
2000.....	2.3	*	*	*	*	0.5	1.4	2.6	3.9	6.3	8.8	14.5
1999.....	2.3	*	*	*	*	0.5	1.2	2.6	3.9	6.1	9.2	12.8





Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of breast (C50)

All races 3/, both sexes

2006.....	13.8	*	*	*	*	0.9	5.4	14.1	28.1	42.1	71.6	116.7
2005.....	14.0	*	*	*	*	0.9	5.7	14.7	28.5	43.5	71.9	122.7
2004.....	14.1	*	*	*	*	1.0	5.7	15.0	29.1	44.9	72.2	124.2
2003.....	14.4	*	*	*	*	1.0	6.1	15.6	29.6	45.5	74.9	132.5
2002.....	14.5	*	*	*	*	1.1	6.1	16.1	29.5	46.5	76.5	134.6
2001.....	14.7	*	*	*	*	1.2	6.3	16.8	30.2	47.6	76.8	134.2
2000.....	15.0	*	*	*	*	1.2	6.3	16.9	31.2	49.1	78.9	147.4
1999.....	14.9	*	*	*	*	1.1	6.1	16.9	31.1	49.5	78.9	144.4

All races 3/, male

2006.....	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.1	3.0
2005.....	0.3	*	*	*	*	*	*	0.2	0.5	1.3	2.1	2.4
2004.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.1	3.3
2003.....	0.3	*	*	*	*	*	*	0.2	0.5	1.1	2.0	3.9
2002.....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.2	3.4
2001.....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.3	3.5
2000.....	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	4.2
1999.....	0.3	*	*	*	*	*	*	0.2	0.6	1.2	2.0	4.5

All races 3/, female

2006.....	26.9	*	*	*	*	1.8	10.8	27.6	53.7	76.9	119.2	169.9
2005.....	27.3	*	*	*	*	1.8	11.3	28.7	54.5	79.2	119.2	177.9
2004.....	27.5	*	*	*	*	2.0	11.3	29.3	55.8	81.6	119.5	178.6
2003.....	28.2	*	*	*	*	2.1	12.2	30.4	56.6	82.6	123.7	189.4
2002.....	28.3	*	*	*	*	2.1	12.0	31.4	56.2	84.4	125.9	191.5
2001.....	28.6	*	*	*	*	2.4	12.4	32.8	57.5	85.8	125.8	188.9
2000.....	29.2	*	*	*	*	2.3	12.4	33.0	59.3	88.3	128.9	205.7
1999.....	28.9	*	*	*	*	2.2	12.0	32.9	59.2	88.9	128.9	200.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of breast (C50)

White, both sexes

2006 4/.....	14.2	*	*	*	*	0.8	4.8	12.9	26.9	41.8	71.9	116.9
2005 4/.....	14.4	*	*	*	*	0.7	5.1	13.2	27.1	43.2	72.3	123.4
2004 4/.....	14.6	*	*	*	*	0.8	5.0	13.6	28.1	44.8	73.0	124.8
2003 4/.....	15.0	*	*	*	*	0.9	5.5	14.4	28.4	45.1	75.1	132.5
2002.....	15.1	*	*	*	*	0.9	5.3	14.9	28.5	46.2	76.6	135.3
2001.....	15.2	*	*	*	*	1.0	5.6	15.7	29.1	47.0	77.1	134.8
2000.....	15.7	*	*	*	*	1.0	5.7	15.8	30.2	49.2	79.4	147.3
1999.....	15.5	*	*	*	*	0.9	5.4	15.7	29.9	49.0	79.0	146.0

White, male

2006 4/.....	0.3	*	*	*	*	*	*	0.2	0.6	1.0	2.1	3.2
2005 4/.....	0.3	*	*	*	*	*	*	0.1	0.5	1.2	2.0	2.4
2004 4/.....	0.3	*	*	*	*	*	*	0.1	0.4	1.0	2.1	3.0
2003 4/.....	0.3	*	*	*	*	*	*	0.2	0.5	1.0	2.0	3.9
2002.....	0.3	*	*	*	*	*	*	0.2	0.5	0.9	2.1	3.3
2001.....	0.3	*	*	*	*	*	*	0.2	0.6	1.4	2.2	3.3
2000.....	0.3	*	*	*	*	*	*	0.2	0.7	1.4	2.2	3.9
1999.....	0.3	*	*	*	*	*	*	0.2	0.5	1.2	2.0	4.4

White, female

2006 4/.....	27.9	*	*	*	*	1.6	9.8	25.6	52.0	77.1	120.4	170.3
2005 4/.....	28.3	*	*	*	*	1.5	10.2	26.2	52.4	79.3	120.7	179.1
2004 4/.....	28.6	*	*	*	*	1.7	10.1	27.0	54.3	82.2	121.3	179.7
2003 4/.....	29.3	*	*	*	*	1.8	11.1	28.4	54.9	82.6	124.6	189.4
2002.....	29.5	*	*	*	*	1.8	10.7	29.4	55.0	84.6	126.5	192.6
2001.....	29.8	*	*	*	*	2.0	11.3	31.0	56.0	85.6	126.9	189.9
2000.....	30.7	*	*	*	*	2.1	11.3	31.2	57.9	89.3	130.2	205.5
1999.....	30.2	*	*	*	*	1.9	10.8	30.9	57.5	88.9	129.5	202.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of breast (C50)

Black, both sexes

2006 4/.....	14.5	*	*	*	*	1.9	10.1	24.3	42.7	53.9	89.1	142.2
2005 4/.....	15.0	*	*	*	*	1.8	10.8	27.0	45.3	57.2	82.0	144.3
2004 4/.....	14.5	*	*	*	*	2.1	10.9	26.0	43.7	56.3	82.3	143.5
2003 4/.....	15.1	*	*	*	*	2.2	11.1	26.5	45.4	59.3	90.8	161.6
2002.....	14.9	*	*	*	*	2.3	11.7	27.0	43.1	60.1	94.0	152.6
2001.....	15.0	*	*	*	*	2.6	11.1	27.3	44.6	63.5	92.3	152.0
2000.....	14.8	*	*	*	*	2.3	11.2	27.8	45.4	58.6	90.5	176.4
1999.....	15.0	*	*	*	*	2.3	11.6	28.7	46.3	63.3	94.5	147.3

Black, male

2006 4/.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2005 4/.....	0.3	*	*	*	*	*	*	*	*	2.8	*	*
2004 4/.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2003 4/.....	0.3	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.4	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.4	*	*	*	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	27.5	*	*	*	*	3.7	19.0	44.5	76.3	91.2	138.2	199.2
2005 4/.....	28.4	*	*	*	*	3.5	20.2	50.0	81.1	96.2	126.6	201.8
2004 4/.....	27.5	*	*	*	*	4.0	20.5	48.0	78.2	95.1	127.4	196.9
2003 4/.....	28.5	*	*	*	*	4.1	20.8	48.9	81.1	100.1	141.0	222.4
2002.....	28.2	*	*	*	*	4.3	22.0	49.8	76.6	101.1	145.0	209.1
2001.....	28.3	*	*	*	*	5.0	20.8	50.3	79.8	106.6	141.8	205.9
2000.....	27.9	*	*	*	*	4.2	20.9	51.5	80.9	98.6	139.8	238.7
1999.....	28.2	*	*	*	*	4.3	21.7	53.0	82.2	106.2	145.4	199.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

All races 3/, both sexes

2006.....	1.3	*	*	*	*	0.4	1.4	2.2	2.6	3.2	3.8	5.2
2005.....	1.3	*	*	*	*	0.4	1.4	2.3	2.6	3.1	4.0	5.4
2004.....	1.3	*	*	*	*	0.5	1.4	2.0	2.8	3.3	3.7	5.1
2003.....	1.3	*	*	*	*	0.5	1.4	2.2	2.8	3.1	4.3	5.7
2002.....	1.4	*	*	*	*	0.5	1.5	2.2	2.7	3.4	4.1	6.2
2001.....	1.4	*	*	*	*	0.5	1.6	2.4	2.9	3.2	4.6	7.2
2000.....	1.5	*	*	*	*	0.6	1.5	2.4	3.1	3.7	4.8	7.3
1999.....	1.5	*	*	*	*	0.5	1.6	2.5	3.2	3.8	4.6	7.9

All races 3/, male

2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All races 3/, female

2006.....	2.6	*	*	*	*	0.9	2.9	4.4	5.0	5.9	6.4	7.7
2005.....	2.6	*	*	*	*	0.9	2.8	4.4	5.1	5.7	6.7	7.8
2004.....	2.6	*	*	*	*	1.1	2.8	4.0	5.4	6.0	6.2	7.3
2003.....	2.7	*	*	*	*	1.0	2.8	4.3	5.5	5.6	7.2	8.3
2002.....	2.7	*	*	*	*	1.1	3.0	4.4	5.2	6.3	6.9	8.9
2001.....	2.8	*	*	*	*	1.1	3.1	4.8	5.5	5.8	7.7	10.1
2000.....	2.9	*	*	*	*	1.2	3.1	4.8	5.9	6.8	8.0	10.3
1999.....	3.0	*	*	*	*	1.1	3.1	4.9	6.2	6.9	7.7	11.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of cervix uteri (C53)

White, both sexes

2006 4/.....	1.2	*	*	*	*	0.4	1.3	2.1	2.4	2.7	3.3	4.2
2005 4/.....	1.2	*	*	*	*	0.4	1.3	2.0	2.3	2.8	3.4	4.8
2004 4/.....	1.2	*	*	*	*	0.5	1.3	1.8	2.6	2.7	3.3	4.1
2003 4/.....	1.2	*	*	*	*	0.4	1.3	2.0	2.6	2.6	3.6	4.9
2002.....	1.3	*	*	*	*	0.5	1.4	2.0	2.4	2.9	3.6	5.1
2001.....	1.3	*	*	*	*	0.5	1.5	2.2	2.6	2.7	4.0	6.2
2000.....	1.4	*	*	*	*	0.5	1.4	2.2	2.7	3.1	4.3	6.3
1999.....	1.4	*	*	*	*	0.5	1.4	2.4	2.8	3.1	4.0	6.8

White, male

2006 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

White, female

2006 4/.....	2.5	*	*	*	*	0.8	2.7	4.1	4.8	5.1	5.6	6.2
2005 4/.....	2.5	*	*	*	*	0.9	2.7	4.0	4.5	5.1	5.8	6.9
2004 4/.....	2.4	*	*	*	*	1.0	2.6	3.7	5.0	5.1	5.5	5.9
2003 4/.....	2.5	*	*	*	*	0.9	2.7	4.0	5.0	4.8	6.0	7.1
2002.....	2.5	*	*	*	*	1.0	2.8	4.0	4.7	5.3	6.0	7.3
2001.....	2.7	*	*	*	*	1.1	3.0	4.5	5.0	5.0	6.6	8.8
2000.....	2.7	*	*	*	*	1.1	2.8	4.3	5.2	5.7	7.1	8.8
1999.....	2.8	*	*	*	*	1.1	2.9	4.7	5.4	5.7	6.7	9.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of cervix uteri (C53)

Black, both sexes

2006 4/.....	2.0	*	*	*	*	0.7	2.4	3.5	4.0	7.5	8.5	17.6
2005 4/.....	2.0	*	*	*	*	0.5	2.3	3.7	5.2	6.3	10.6	13.1
2004 4/.....	2.0	*	*	*	*	0.8	2.2	3.7	5.3	7.6	8.8	16.1
2003 4/.....	2.1	*	*	*	*	0.9	2.2	3.6	5.5	7.4	11.3	14.6
2002.....	2.2	*	*	*	*	0.8	2.5	4.1	5.0	9.0	10.8	17.5
2001.....	2.1	*	*	*	*	0.8	2.4	3.8	5.2	7.8	11.9	19.0
2000.....	2.4	*	*	*	*	0.9	2.6	4.3	6.3	9.5	11.9	19.7
1999.....	2.3	*	*	*	*	0.7	2.5	3.4	7.1	10.7	11.3	21.7

Black, male

2006 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Black, female

2006 4/.....	3.8	*	*	*	*	1.4	4.5	6.6	7.2	12.8	13.4	24.9
2005 4/.....	3.8	*	*	*	*	1.0	4.4	6.9	9.4	10.9	16.6	18.4
2004 4/.....	3.9	*	*	*	*	1.6	4.2	6.9	9.6	13.0	13.8	22.4
2003 4/.....	4.0	*	*	*	*	1.7	4.2	6.8	10.0	12.6	17.7	20.3
2002.....	4.2	*	*	*	*	1.5	4.7	7.6	8.9	15.4	16.9	24.2
2001.....	4.0	*	*	*	*	1.5	4.5	7.0	9.3	13.3	18.5	26.1
2000.....	4.5	*	*	*	*	1.8	4.9	7.9	11.3	16.3	18.6	27.0
1999.....	4.4	*	*	*	*	1.2	4.7	6.4	12.8	18.3	17.6	29.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

All races 3/, both sexes

2006.....	2.5	*	*	*	*	0.1	0.3	1.4	4.8	10.0	15.7	21.6
2005.....	2.4	*	*	*	*	0.1	0.3	1.3	4.8	9.7	15.0	22.1
2004.....	2.4	*	*	*	*	0.0	0.4	1.3	4.4	10.1	15.5	22.3
2003.....	2.4	*	*	*	*	0.1	0.3	1.3	4.7	10.1	15.2	22.8
2002.....	2.4	*	*	*	*	0.1	0.3	1.4	4.5	10.0	15.4	24.6
2001.....	2.4	*	*	*	*	0.1	0.3	1.4	4.4	9.7	16.4	24.7
2000.....	2.3	*	*	*	*	0.1	0.3	1.3	4.6	9.7	15.8	24.7
1999.....	2.3	*	*	*	*	0.1	0.3	1.3	4.3	9.8	16.4	24.0

All races 3/, male

2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All races 3/, female

2006.....	4.9	*	*	*	*	0.1	0.7	2.7	9.3	18.5	26.4	31.6
2005.....	4.7	*	*	*	*	0.1	0.7	2.6	9.3	17.9	25.1	32.2
2004.....	4.7	*	*	*	*	0.1	0.8	2.6	8.6	18.5	25.9	32.3
2003.....	4.7	*	*	*	*	0.1	0.7	2.6	9.0	18.5	25.3	33.0
2002.....	4.7	*	*	*	*	0.1	0.6	2.8	8.7	18.4	25.6	35.2
2001.....	4.7	*	*	*	*	0.2	0.6	2.8	8.4	17.8	27.2	35.0
2000.....	4.6	*	*	*	*	0.1	0.6	2.6	8.9	17.7	26.2	34.8
1999.....	4.5	*	*	*	*	0.1	0.6	2.5	8.2	17.8	27.1	33.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

White, both sexes

2006 4/.....	2.5	*	*	*	*	0.1	0.3	1.3	4.3	8.8	14.9	21.7
2005 4/.....	2.4	*	*	*	*	*	0.3	1.3	4.3	8.8	14.1	21.8
2004 4/.....	2.4	*	*	*	*	*	0.4	1.3	4.1	8.8	14.9	22.4
2003 4/.....	2.4	*	*	*	*	*	0.3	1.3	4.3	9.1	14.4	22.7
2002.....	2.4	*	*	*	*	*	0.3	1.3	4.1	8.9	14.4	23.8
2001.....	2.4	*	*	*	*	0.1	0.3	1.4	4.0	8.8	15.7	24.5
2000.....	2.3	*	*	*	*	0.1	0.3	1.3	4.3	8.5	14.8	24.4
1999.....	2.4	*	*	*	*	*	0.3	1.3	4.0	8.9	15.8	23.4

White, male

2006 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

White, female

2006 4/.....	4.9	*	*	*	*	0.1	0.7	2.6	8.4	16.4	25.2	31.8
2005 4/.....	4.7	*	*	*	*	*	0.6	2.5	8.4	16.4	23.8	31.8
2004 4/.....	4.7	*	*	*	*	*	0.7	2.5	7.9	16.4	25.1	32.6
2003 4/.....	4.7	*	*	*	*	*	0.6	2.5	8.3	16.9	24.1	32.8
2002.....	4.7	*	*	*	*	*	0.6	2.6	8.0	16.5	24.1	34.2
2001.....	4.8	*	*	*	*	0.2	0.6	2.7	7.8	16.2	26.2	34.7
2000.....	4.6	*	*	*	*	0.1	0.6	2.6	8.3	15.7	24.6	34.3
1999.....	4.6	*	*	*	*	*	0.6	2.5	7.7	16.3	26.1	32.8



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)

Black, both sexes

2006 4/.....	3.1	*	*	*	*	*	*	1.7	9.7	23.0	28.9	25.9
2005 4/.....	3.0	*	*	*	*	*	0.4	1.9	9.8	20.1	27.5	31.3
2004 4/.....	2.9	*	*	*	*	*	0.5	1.9	8.0	23.0	26.1	26.2
2003 4/.....	2.9	*	*	*	*	*	0.5	1.8	9.0	21.0	27.8	30.0
2002.....	3.0	*	*	*	*	*	0.4	1.9	8.4	22.3	28.3	38.9
2001.....	2.8	*	*	*	*	*	0.4	2.1	7.7	21.2	28.4	32.2
2000.....	2.9	*	*	*	*	*	0.4	1.6	8.3	22.7	30.5	33.2
1999.....	2.7	*	*	*	*	*	0.5	1.4	7.9	21.3	27.3	35.4

Black, male

2006 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Black, female

2006 4/.....	6.0	*	*	*	*	*	*	3.2	17.6	39.5	45.4	36.6
2005 4/.....	5.8	*	*	*	*	*	0.8	3.5	17.8	34.5	43.1	43.9
2004 4/.....	5.5	*	*	*	*	*	1.0	3.6	14.4	39.4	40.9	36.6
2003 4/.....	5.5	*	*	*	*	*	1.0	3.3	16.2	36.0	43.5	41.7
2002.....	5.7	*	*	*	*	*	0.7	3.6	15.2	38.1	44.3	53.8
2001.....	5.4	*	*	*	*	*	0.7	3.9	13.9	36.2	44.3	44.2
2000.....	5.5	*	*	*	*	*	0.7	3.0	14.9	38.6	47.7	45.5
1999.....	5.2	*	*	*	*	*	0.9	2.6	14.3	36.3	42.5	48.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of ovary (C56)

All races 3/, both sexes

2006.....	5.0	*	*	*	0.1	0.2	0.9	3.8	9.6	18.9	31.1	37.8
2005.....	5.0	*	*	*	*	0.2	0.9	4.0	10.0	19.1	30.9	38.1
2004.....	5.0	*	*	*	0.1	0.2	1.0	4.0	9.6	20.0	32.1	37.7
2003.....	5.0	*	*	*	0.1	0.2	1.0	4.0	10.4	19.5	32.6	38.4
2002.....	5.1	*	*	*	0.1	0.2	1.0	4.3	10.3	19.7	33.4	38.1
2001.....	5.1	*	*	*	0.1	0.3	1.0	4.1	10.3	20.0	33.2	39.6
2000.....	5.0	*	*	*	0.1	0.3	1.0	4.2	10.1	19.6	33.2	40.9
1999.....	4.9	*	*	*	0.1	0.2	1.1	4.2	10.2	19.6	31.7	37.6

All races 3/, male

2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All races 3/, female

2006.....	9.8	*	*	*	0.1	0.4	1.8	7.6	18.6	34.9	52.4	55.5
2005.....	9.8	*	*	*	*	0.4	1.9	7.8	19.4	35.3	51.8	55.6
2004.....	9.9	*	*	*	0.1	0.5	2.1	7.8	18.5	36.8	53.7	54.6
2003.....	9.9	*	*	*	0.1	0.4	2.0	7.8	20.0	35.7	54.4	55.4
2002.....	10.0	*	*	*	0.2	0.5	2.1	8.4	19.9	36.1	55.6	54.6
2001.....	9.9	*	*	*	0.1	0.5	2.1	8.1	19.9	36.6	55.0	56.1
2000.....	9.8	*	*	*	0.1	0.5	2.1	8.3	19.4	35.7	54.8	57.6
1999.....	9.6	*	*	*	0.1	0.4	2.2	8.3	19.6	35.7	52.3	52.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of ovary (C56)

White, both sexes

2006 4/.....	5.5	*	*	*	0.1	0.2	1.0	4.0	10.0	19.8	32.1	38.8
2005 4/.....	5.5	*	*	*	*	0.2	0.9	4.1	10.4	19.6	31.7	39.4
2004 4/.....	5.5	*	*	*	0.1	0.2	1.1	4.0	9.9	20.5	33.3	38.6
2003 4/.....	5.6	*	*	*	0.1	0.2	1.0	4.1	10.8	20.0	33.6	39.1
2002.....	5.6	*	*	*	0.1	0.2	1.1	4.5	10.7	20.4	34.3	39.0
2001.....	5.6	*	*	*	*	0.3	1.1	4.4	10.5	20.6	34.2	40.4
2000.....	5.5	*	*	*	*	0.3	1.1	4.4	10.5	20.1	34.3	42.0
1999.....	5.4	*	*	*	0.1	0.2	1.1	4.4	10.6	20.1	32.4	38.7

White, male

2006 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

White, female

2006 4/.....	10.9	*	*	*	0.1	0.4	1.9	7.9	19.5	36.8	54.4	57.0
2005 4/.....	10.9	*	*	*	*	0.4	1.9	8.2	20.4	36.5	53.6	57.5
2004 4/.....	10.9	*	*	*	0.1	0.5	2.1	8.0	19.3	38.0	56.0	56.0
2003 4/.....	11.0	*	*	*	0.1	0.5	2.1	8.1	21.0	37.1	56.3	56.4
2002.....	11.1	*	*	*	0.2	0.5	2.2	8.9	20.8	37.7	57.4	55.9
2001.....	11.0	*	*	*	*	0.5	2.1	8.6	20.5	38.0	56.8	57.3
2000.....	10.9	*	*	*	*	0.5	2.2	8.7	20.4	37.0	56.8	59.0
1999.....	10.6	*	*	*	0.1	0.4	2.3	8.7	20.5	36.9	53.6	54.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of ovary (C56)

Black, both sexes

2006 4/.....	3.0	*	*	*	*	*	0.9	3.5	8.5	15.9	25.1	33.5
2005 4/.....	3.1	*	*	*	*	*	1.0	3.2	8.6	18.6	27.6	31.3
2004 4/.....	3.3	*	*	*	*	*	0.9	3.9	8.6	20.1	27.3	32.7
2003 4/.....	3.1	*	*	*	*	*	1.0	3.8	8.6	17.7	27.5	36.8
2002.....	3.1	*	*	*	*	*	1.0	3.4	9.1	17.1	28.8	34.7
2001.....	3.1	*	*	*	*	*	1.1	3.0	10.0	18.3	26.9	36.5
2000.....	2.9	*	*	*	*	*	1.0	3.3	8.4	18.1	26.6	34.4
1999.....	3.0	*	*	*	*	*	0.8	3.5	9.1	18.5	28.7	28.7

Black, male

2006 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Black, female

2006 4/.....	5.7	*	*	*	*	*	1.6	6.4	15.4	27.2	39.4	47.2
2005 4/.....	6.0	*	*	*	*	*	1.8	5.9	15.5	31.9	43.3	43.9
2004 4/.....	6.2	*	*	*	*	*	1.7	7.2	15.6	34.5	42.8	45.7
2003 4/.....	6.0	*	*	*	*	*	1.9	7.1	15.5	30.2	43.1	51.0
2002.....	5.9	*	*	*	*	*	1.9	6.4	16.4	29.3	45.2	48.1
2001.....	5.9	*	*	*	*	*	2.1	5.7	18.0	31.2	42.0	50.1
2000.....	5.6	*	*	*	*	*	2.0	6.2	15.2	30.8	41.6	47.2
1999.....	5.7	*	*	*	*	*	1.5	6.6	16.4	31.5	44.8	39.4







Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

All races 3/, both sexes

2006.....	4.1	*	*	0.1	0.0	0.1	0.6	2.8	7.8	16.3	26.2	34.3
2005.....	4.2	*	*	0.1	*	0.1	0.7	3.0	8.3	16.9	26.9	32.6
2004.....	4.2	*	*	0.1	*	0.1	0.6	3.0	8.7	16.7	26.8	32.7
2003.....	4.2	*	*	0.1	0.1	0.1	0.7	2.9	9.0	16.9	27.0	33.2
2002.....	4.2	*	*	0.1	0.0	0.1	0.7	3.3	8.8	17.7	26.3	32.5
2001.....	4.2	*	0.1	0.1	0.1	0.1	0.8	3.3	9.2	17.6	26.3	32.3
2000.....	4.2	*	*	0.1	0.1	0.1	0.7	3.3	9.0	17.1	26.6	33.8
1999.....	4.0	*	*	0.1	*	0.1	0.8	3.1	8.9	16.7	24.6	31.4

All races 3/, male

2006.....	5.2	*	*	*	*	0.2	0.9	4.0	11.4	23.1	37.1	52.1
2005.....	5.3	*	*	*	*	0.2	0.8	4.3	12.0	24.4	37.9	50.6
2004.....	5.3	*	*	*	*	0.2	0.8	4.3	12.8	23.9	38.2	47.0
2003.....	5.4	*	*	*	*	0.2	1.0	4.3	13.3	23.7	39.5	48.7
2002.....	5.4	*	*	*	*	0.2	1.0	4.8	13.0	25.1	37.9	50.1
2001.....	5.4	*	*	*	*	0.1	1.1	4.9	13.4	25.2	37.4	52.9
2000.....	5.3	*	*	*	*	0.1	1.0	4.8	13.1	24.2	38.2	55.1
1999.....	5.0	*	*	*	*	0.2	1.1	4.5	12.9	23.6	35.7	49.4

All races 3/, female

2006.....	3.1	*	*	*	*	0.1	0.4	1.6	4.5	10.5	18.7	25.9
2005.....	3.1	*	*	*	*	*	0.5	1.8	4.8	10.5	19.4	24.3
2004.....	3.1	*	*	*	*	*	0.4	1.7	5.0	10.6	19.1	26.3
2003.....	3.1	*	*	0.1	*	0.1	0.4	1.6	4.9	11.2	18.7	26.4
2002.....	3.1	*	*	*	*	*	0.4	1.8	4.9	11.6	18.7	24.9
2001.....	3.1	*	*	0.1	*	0.1	0.5	1.9	5.3	11.3	19.0	23.7
2000.....	3.1	*	*	*	*	*	0.5	1.9	5.3	11.3	18.9	25.1
1999.....	3.0	*	*	*	*	*	0.5	1.8	5.1	11.0	17.4	24.1



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

White, both sexes

2006 4/.....	4.5	*	*	0.1	*	0.1	0.6	2.8	8.1	16.8	27.0	34.9
2005 4/.....	4.6	*	*	0.1	*	0.1	0.7	3.1	8.5	17.5	27.6	33.1
2004 4/.....	4.6	*	*	0.1	*	0.1	0.6	3.1	9.1	17.1	27.3	33.7
2003 4/.....	4.6	*	*	0.1	*	0.1	0.7	3.0	9.1	17.5	27.7	33.8
2002.....	4.6	*	*	0.1	*	0.1	0.7	3.3	9.1	18.3	26.8	33.2
2001.....	4.6	*	*	0.1	*	0.1	0.7	3.4	9.4	18.2	26.7	32.8
2000.....	4.6	*	*	*	*	0.1	0.7	3.4	9.3	17.5	27.0	34.5
1999.....	4.3	*	*	0.1	*	0.1	0.7	3.1	9.1	17.1	24.9	31.5

White, male

2006 4/.....	5.7	*	*	*	*	*	0.8	4.0	11.6	23.7	38.2	53.4
2005 4/.....	5.8	*	*	*	*	*	0.8	4.3	12.1	25.0	38.7	52.0
2004 4/.....	5.8	*	*	*	*	*	0.8	4.5	13.2	24.4	39.0	48.6
2003 4/.....	5.9	*	*	*	*	0.1	1.0	4.4	13.6	24.3	40.4	50.0
2002.....	5.9	*	*	*	*	*	1.0	4.9	13.3	25.9	38.6	51.9
2001.....	5.8	*	*	*	*	*	1.0	5.0	13.6	26.0	37.7	54.4
2000.....	5.7	*	*	*	*	*	1.0	4.9	13.4	24.4	38.7	56.6
1999.....	5.4	*	*	*	*	0.1	1.0	4.5	13.3	23.9	36.3	49.8

White, female

2006 4/.....	3.4	*	*	*	*	*	0.3	1.6	4.7	10.9	19.2	26.2
2005 4/.....	3.5	*	*	*	*	*	0.5	1.9	5.0	11.0	20.0	24.3
2004 4/.....	3.4	*	*	*	*	*	0.4	1.8	5.1	10.9	19.4	27.0
2003 4/.....	3.4	*	*	*	*	*	0.4	1.7	4.9	11.6	19.1	26.6
2002.....	3.4	*	*	*	*	*	0.4	1.8	5.1	11.9	18.9	25.1
2001.....	3.4	*	*	*	*	*	0.4	1.8	5.4	11.6	19.4	23.8
2000.....	3.4	*	*	*	*	*	0.4	1.9	5.4	11.6	19.2	25.5
1999.....	3.2	*	*	*	*	*	0.4	1.8	5.3	11.3	17.5	24.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of kidney and renal pelvis (C64-C65)

Black, both sexes

2006 4/.....	2.9	*	*	*	*	0.4	1.0	3.6	7.4	14.8	22.1	33.5
2005 4/.....	3.0	*	*	*	*	0.4	0.9	3.3	8.9	15.2	23.8	30.5
2004 4/.....	2.8	*	*	*	*	0.4	0.8	3.1	8.0	15.8	25.8	28.2
2003 4/.....	2.9	*	*	*	*	*	1.1	3.2	9.5	14.6	23.8	33.8
2002.....	2.8	*	*	*	*	*	0.8	3.5	8.6	15.4	26.1	28.2
2001.....	3.0	*	*	*	*	*	1.2	3.7	9.5	15.0	25.6	29.8
2000.....	2.9	*	*	*	*	*	0.9	3.4	8.8	16.5	25.6	29.7
1999.....	2.9	*	*	*	*	0.4	1.1	3.8	8.6	16.1	23.5	31.2

Black, male

2006 4/.....	3.8	*	*	*	*	0.7	1.6	5.2	12.0	22.5	32.2	49.1
2005 4/.....	3.9	*	*	*	*	*	1.3	5.2	13.9	23.9	35.0	39.0
2004 4/.....	3.6	*	*	*	*	*	1.1	4.5	12.1	24.0	37.3	38.7
2003 4/.....	3.7	*	*	*	*	*	1.6	4.9	13.8	22.2	37.0	43.8
2002.....	3.5	*	*	*	*	*	1.2	4.8	13.9	21.6	36.1	39.7
2001.....	3.8	*	*	*	*	*	1.5	5.4	14.9	22.3	40.3	46.5
2000.....	3.6	*	*	*	*	*	1.0	4.9	13.1	26.3	38.2	49.6
1999.....	3.6	*	*	*	*	*	1.6	5.8	13.2	23.5	32.0	49.2

Black, female

2006 4/.....	2.1	*	*	*	*	*	*	2.2	3.6	9.2	16.3	27.1
2005 4/.....	2.1	*	*	*	*	*	*	1.7	4.8	9.0	17.5	27.0
2004 4/.....	2.2	*	*	*	*	*	*	1.8	4.7	10.0	19.4	24.0
2003 4/.....	2.2	*	*	*	*	*	0.7	1.8	6.0	9.2	16.4	30.0
2002.....	2.2	*	*	*	*	*	*	2.4	4.2	11.0	20.5	23.8
2001.....	2.2	*	*	*	*	*	1.0	2.2	5.2	9.9	17.4	23.6
2000.....	2.1	*	*	*	*	*	0.8	2.2	5.2	9.6	18.6	22.3
1999.....	2.2	*	*	*	*	*	*	2.1	5.0	10.9	18.8	24.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasm of bladder (C67)

All races 3/, both sexes

2006.....	4.5	*	*	*	*	*	0.2	1.2	5.0	14.9	37.4	67.4
2005.....	4.5	*	*	*	*	*	0.3	1.2	4.9	14.9	37.2	68.3
2004.....	4.4	*	*	*	*	*	0.3	1.3	5.0	14.9	37.4	67.9
2003.....	4.3	*	*	*	*	*	0.2	1.3	5.0	14.9	36.4	64.8
2002.....	4.4	*	*	*	*	*	0.3	1.3	4.9	15.9	37.0	66.6
2001.....	4.3	*	*	*	*	*	0.3	1.2	5.0	15.4	35.7	68.9
2000.....	4.3	*	*	*	*	*	0.2	1.2	5.0	15.5	35.7	69.6
1999.....	4.3	*	*	*	*	*	0.2	1.3	5.2	15.9	35.3	68.5

All races 3/, male

2006.....	6.4	*	*	*	*	*	0.3	1.8	7.8	23.7	65.5	131.6
2005.....	6.3	*	*	*	*	*	0.4	1.7	7.6	24.3	65.1	132.1
2004.....	6.2	*	*	*	*	*	0.3	2.0	7.9	24.0	64.7	133.0
2003.....	6.0	*	*	*	*	*	0.3	2.0	7.8	23.9	63.1	125.1
2002.....	6.0	*	*	*	*	*	0.3	1.8	7.7	25.2	63.0	128.9
2001.....	6.0	*	*	*	*	*	0.3	1.8	7.9	25.2	62.5	134.8
2000.....	5.9	*	*	*	*	*	0.3	1.7	7.9	25.2	62.6	139.4
1999.....	5.9	*	*	*	*	*	0.3	2.0	8.2	25.7	62.5	134.8

All races 3/, female

2006.....	2.7	*	*	*	*	*	0.2	0.6	2.3	7.3	18.1	37.4
2005.....	2.7	*	*	*	*	*	0.2	0.8	2.3	6.9	18.3	39.0
2004.....	2.7	*	*	*	*	*	0.2	0.6	2.3	7.3	19.0	38.6
2003.....	2.6	*	*	*	*	*	0.2	0.6	2.4	7.4	18.5	38.1
2002.....	2.8	*	*	*	*	*	0.2	0.7	2.4	8.2	19.7	39.6
2001.....	2.7	*	*	*	*	*	0.3	0.6	2.3	7.2	18.1	41.4
2000.....	2.7	*	*	*	*	*	0.2	0.6	2.4	7.5	18.1	41.2
1999.....	2.7	*	*	*	*	*	0.2	0.6	2.5	7.8	17.7	41.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of bladder (C67)

White, both sexes

2006 4/.....	5.1	*	*	*	*	*	0.2	1.3	5.2	15.5	38.9	69.8
2005 4/.....	5.0	*	*	*	*	*	0.3	1.3	5.1	15.6	38.8	70.0
2004 4/.....	5.0	*	*	*	*	*	0.2	1.3	5.3	15.8	39.1	69.6
2003 4/.....	4.8	*	*	*	*	*	0.2	1.3	5.3	15.7	37.5	66.6
2002.....	4.9	*	*	*	*	*	0.3	1.3	5.1	16.5	38.1	67.7
2001.....	4.8	*	*	*	*	*	0.3	1.2	5.2	16.1	36.7	70.8
2000.....	4.8	*	*	*	*	*	0.3	1.2	5.3	16.2	36.4	72.0
1999.....	4.8	*	*	*	*	*	0.2	1.3	5.4	16.3	36.3	69.8

White, male

2006 4/.....	7.3	*	*	*	*	*	0.3	1.9	8.2	25.2	69.0	137.9
2005 4/.....	7.2	*	*	*	*	*	0.4	1.8	8.0	25.7	68.1	137.4
2004 4/.....	7.1	*	*	*	*	*	0.3	2.1	8.4	25.7	68.3	139.3
2003 4/.....	6.8	*	*	*	*	*	0.3	2.0	8.3	25.3	65.7	131.0
2002.....	6.8	*	*	*	*	*	0.4	1.9	8.2	26.5	65.3	134.1
2001.....	6.8	*	*	*	*	*	0.3	1.9	8.3	26.6	65.0	141.1
2000.....	6.7	*	*	*	*	*	0.3	1.9	8.4	26.5	64.7	145.7
1999.....	6.7	*	*	*	*	*	0.3	1.9	8.5	26.9	64.8	140.7

White, female

2006 4/.....	2.9	*	*	*	*	*	0.2	0.7	2.4	7.2	17.9	37.9
2005 4/.....	3.0	*	*	*	*	*	0.2	0.7	2.3	7.0	18.5	39.0
2004 4/.....	2.9	*	*	*	*	*	0.2	0.6	2.4	7.3	19.2	38.1
2003 4/.....	2.9	*	*	*	*	*	0.1	0.5	2.4	7.6	18.4	38.1
2002.....	3.0	*	*	*	*	*	0.2	0.7	2.3	8.1	19.8	38.9
2001.....	2.9	*	*	*	*	*	0.3	0.6	2.2	7.2	17.8	41.3
2000.....	2.9	*	*	*	*	*	0.2	0.6	2.5	7.5	17.8	42.1
1999.....	2.9	*	*	*	*	*	0.2	0.6	2.4	7.6	17.7	41.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasm of bladder (C67)

Black, both sexes

2006 4/.....	2.5	*	*	*	*	*	*	1.2	4.8	13.4	31.0	54.5
2005 4/.....	2.5	*	*	*	*	*	*	1.5	4.6	12.3	29.6	57.5
2004 4/.....	2.4	*	*	*	*	*	0.5	1.6	4.1	11.7	27.2	60.1
2003 4/.....	2.4	*	*	*	*	*	0.4	1.6	3.9	12.0	32.3	53.4
2002.....	2.5	*	*	*	*	*	*	1.1	4.5	14.3	31.0	63.3
2001.....	2.4	*	*	*	*	*	0.4	1.3	4.9	11.9	31.2	55.0
2000.....	2.3	*	*	*	*	*	*	1.0	4.1	13.3	34.7	49.7
1999.....	2.5	*	*	*	*	*	*	1.5	5.6	15.0	29.1	60.3

Black, male

2006 4/.....	3.0	*	*	*	*	*	*	1.8	7.5	17.8	41.9	92.9
2005 4/.....	2.9	*	*	*	*	*	*	1.7	6.6	18.2	45.3	101.3
2004 4/.....	2.6	*	*	*	*	*	*	2.3	5.9	14.5	39.5	81.5
2003 4/.....	2.8	*	*	*	*	*	*	2.2	5.4	17.5	48.8	80.3
2002.....	2.7	*	*	*	*	*	*	1.2	5.6	18.3	47.6	91.1
2001.....	2.5	*	*	*	*	*	*	1.7	6.4	16.1	43.0	78.3
2000.....	2.7	*	*	*	*	*	*	1.4	5.8	17.6	52.7	88.8
1999.....	2.9	*	*	*	*	*	*	2.6	7.4	19.1	46.1	86.7

Black, female

2006 4/.....	2.2	*	*	*	*	*	*	0.7	2.6	10.3	24.8	38.7
2005 4/.....	2.0	*	*	*	*	*	*	1.3	3.0	8.2	20.6	39.8
2004 4/.....	2.2	*	*	*	*	*	*	0.9	2.6	9.7	20.3	51.6
2003 4/.....	2.1	*	*	*	*	*	*	1.2	2.8	8.1	23.0	42.9
2002.....	2.3	*	*	*	*	*	*	0.9	3.7	11.6	21.6	52.6
2001.....	2.2	*	*	*	*	*	*	0.8	3.8	8.9	24.6	46.3
2000.....	2.0	*	*	*	*	*	*	*	2.7	10.3	24.5	35.2
1999.....	2.2	*	*	*	*	*	*	*	4.2	12.1	19.6	50.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes

2006.....	4.3	*	0.6	0.8	0.5	0.9	2.0	4.5	9.3	14.7	19.0	16.7
2005.....	4.4	*	0.5	0.8	0.6	0.9	2.2	4.6	9.4	15.8	20.2	15.8
2004.....	4.4	*	0.6	0.8	0.5	0.9	2.0	4.6	9.5	15.7	19.6	16.6
2003.....	4.4	*	0.7	0.8	0.6	1.0	2.4	4.8	9.7	15.5	19.5	15.2
2002.....	4.4	*	0.7	0.8	0.5	1.0	2.2	5.1	9.6	16.2	19.1	16.4
2001.....	4.4	*	0.7	0.7	0.6	1.1	2.3	5.0	9.7	15.4	20.4	16.4
2000.....	4.5	0.5	0.7	0.8	0.6	1.0	2.4	5.1	9.6	16.5	20.0	16.9
1999.....	4.6	*	0.7	0.8	0.6	1.1	2.4	5.1	10.3	16.9	20.4	16.0

All races 3/, male

2006.....	4.9	*	0.7	0.8	0.5	1.0	2.4	5.5	11.4	18.2	24.0	21.2
2005.....	5.0	*	0.5	0.8	0.7	1.2	2.8	5.6	11.3	19.3	25.3	19.3
2004.....	4.9	*	0.7	0.8	0.6	1.1	2.5	5.8	11.8	18.4	23.7	19.4
2003.....	5.0	*	0.7	0.8	0.6	1.1	2.9	6.2	11.4	18.6	24.6	20.0
2002.....	5.0	*	0.8	0.9	0.7	1.1	2.8	6.2	11.7	20.2	23.8	20.1
2001.....	5.0	*	0.7	0.7	0.6	1.3	2.8	6.1	12.1	19.2	25.8	20.1
2000.....	5.1	*	0.7	0.8	0.7	1.3	3.1	6.2	11.8	20.2	24.3	22.4
1999.....	5.1	*	0.7	0.7	0.7	1.4	2.8	6.2	12.4	20.8	25.5	20.7

All races 3/, female

2006.....	3.7	*	0.6	0.7	0.4	0.8	1.5	3.6	7.3	11.8	15.5	14.6
2005.....	3.9	*	0.6	0.8	0.4	0.7	1.6	3.6	7.5	12.9	16.7	14.3
2004.....	3.9	*	0.6	0.8	0.4	0.7	1.6	3.4	7.3	13.4	16.8	15.4
2003.....	3.9	*	0.6	0.8	0.5	0.9	2.0	3.5	8.1	12.8	16.1	13.1
2002.....	3.9	*	0.7	0.7	0.4	0.8	1.7	4.1	7.7	12.8	16.0	14.8
2001.....	3.9	*	0.6	0.8	0.5	0.8	1.7	3.9	7.6	12.3	16.8	14.8
2000.....	3.9	*	0.6	0.7	0.5	0.7	1.7	4.0	7.6	13.4	17.3	14.7
1999.....	4.1	*	0.8	0.8	0.5	0.8	1.9	4.1	8.4	13.7	17.1	14.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

White, both sexes

2006 4/.....	4.9	*	0.6	0.8	0.5	1.0	2.2	5.0	10.1	15.9	20.3	17.4
2005 4/.....	5.0	*	0.5	0.8	0.6	1.0	2.4	5.0	10.2	17.1	21.3	16.5
2004 4/.....	4.9	*	0.7	0.8	0.6	1.0	2.3	5.0	10.3	17.0	20.8	17.3
2003 4/.....	5.0	*	0.7	0.9	0.6	1.1	2.7	5.2	10.5	16.6	20.7	15.7
2002.....	5.0	*	0.7	0.8	0.5	1.1	2.5	5.6	10.4	17.4	20.1	17.1
2001.....	5.0	*	0.7	0.8	0.6	1.2	2.6	5.4	10.5	16.5	21.6	17.0
2000.....	5.1	*	0.7	0.8	0.6	1.1	2.6	5.6	10.4	17.8	21.0	17.5
1999.....	5.2	*	0.8	0.8	0.6	1.3	2.6	5.6	11.1	18.2	21.4	16.6

White, male

2006 4/.....	5.5	*	0.7	0.9	0.5	1.2	2.7	6.0	12.3	19.5	25.4	22.1
2005 4/.....	5.6	*	0.4	0.8	0.7	1.3	3.1	6.0	12.1	20.7	26.4	20.2
2004 4/.....	5.5	*	0.7	0.8	0.7	1.1	2.7	6.2	12.8	20.0	24.8	20.6
2003 4/.....	5.6	*	0.8	0.9	0.7	1.3	3.2	6.7	12.3	19.8	25.8	20.7
2002.....	5.7	*	0.7	0.9	0.7	1.2	3.1	6.8	12.4	21.6	24.8	21.2
2001.....	5.7	*	0.8	0.7	0.7	1.5	3.2	6.7	12.9	20.5	27.1	20.5
2000.....	5.7	*	0.7	0.9	0.7	1.4	3.4	6.8	12.8	21.7	25.5	23.5
1999.....	5.8	*	0.8	0.8	0.6	1.5	3.1	6.7	13.3	22.3	26.8	21.8

White, female

2006 4/.....	4.3	*	0.5	0.7	0.4	0.9	1.6	4.0	8.0	12.8	16.7	15.3
2005 4/.....	4.4	*	0.6	0.8	0.5	0.8	1.7	4.0	8.3	14.0	17.8	14.8
2004 4/.....	4.4	*	0.6	0.7	0.4	0.8	1.8	3.7	8.0	14.4	18.1	15.8
2003 4/.....	4.4	*	0.6	0.9	0.5	1.0	2.3	3.8	8.7	13.9	17.2	13.4
2002.....	4.4	*	0.6	0.8	0.4	1.0	1.9	4.5	8.5	13.8	16.9	15.4
2001.....	4.4	*	0.7	0.8	0.5	0.9	1.9	4.3	8.2	13.0	18.0	15.6
2000.....	4.4	*	0.6	0.7	0.5	0.9	1.8	4.5	8.2	14.5	18.0	15.0
1999.....	4.6	*	0.8	0.8	0.5	1.0	2.1	4.5	9.1	14.8	17.9	14.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

Black, both sexes

2006 4/.....	1.9	*	*	0.7	0.4	0.4	1.2	2.5	5.3	8.0	9.4	9.6
2005 4/.....	2.0	*	*	0.8	0.4	0.5	1.1	2.7	5.4	8.3	12.1	9.4
2004 4/.....	2.0	*	*	0.7	0.5	0.7	1.3	2.8	4.9	8.0	10.0	10.4
2003 4/.....	2.1	*	*	0.7	0.5	0.7	1.4	3.1	5.5	8.9	10.5	10.5
2002.....	2.1	*	1.1	0.8	0.5	0.6	1.2	2.7	5.6	8.7	11.8	9.5
2001.....	2.0	*	*	0.6	0.5	0.5	1.1	3.0	5.8	9.2	9.9	11.7
2000.....	2.1	*	*	0.8	0.6	0.7	1.6	2.9	5.7	8.1	11.9	10.6
1999.....	2.1	*	*	0.8	0.6	0.7	1.3	3.2	6.0	8.2	11.8	9.9

Black, male

2006 4/.....	2.2	*	*	0.6	0.6	*	1.5	3.1	7.0	10.1	12.4	*
2005 4/.....	2.4	*	*	0.8	*	0.9	1.3	3.6	7.0	10.3	17.5	*
2004 4/.....	2.2	*	*	0.7	*	0.8	1.9	3.9	6.7	7.9	14.2	*
2003 4/.....	2.5	*	*	0.8	0.7	*	2.1	4.0	6.7	12.0	15.2	*
2002.....	2.3	*	*	1.0	*	0.8	1.3	2.7	8.1	10.4	16.4	*
2001.....	2.2	*	*	0.7	*	*	1.2	3.6	7.9	10.1	12.4	*
2000.....	2.4	*	*	0.9	0.7	1.1	2.0	3.7	7.1	9.8	13.0	*
1999.....	2.3	*	*	0.7	0.8	0.9	1.8	4.4	7.2	10.0	12.6	*

Black, female

2006 4/.....	1.7	*	*	0.8	*	*	0.9	2.0	3.9	6.5	7.7	9.1
2005 4/.....	1.7	*	*	0.7	*	*	1.0	1.9	4.1	6.9	9.0	9.8
2004 4/.....	1.8	*	*	0.8	*	*	0.7	1.9	3.4	8.1	7.6	11.4
2003 4/.....	1.8	*	*	0.6	*	*	0.9	2.2	4.6	6.7	7.8	11.7
2002.....	1.9	*	*	*	*	*	1.1	2.8	3.7	7.4	9.3	10.3
2001.....	1.8	*	*	*	*	*	1.0	2.4	4.2	8.5	8.6	9.7
2000.....	1.9	*	*	0.6	*	*	1.2	2.2	4.6	6.9	11.3	10.7
1999.....	1.9	*	*	0.9	*	*	1.0	2.1	5.0	6.9	11.4	9.2



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes

2006.....	18.4	0.8	0.8	0.7	1.6	2.0	3.5	8.5	24.4	65.2	137.8	186.1
2005.....	18.6	0.6	0.8	0.8	1.6	2.0	3.4	8.7	25.1	67.0	138.7	189.8
2004.....	18.6	0.6	0.8	0.9	1.5	2.1	3.4	9.0	25.8	68.2	139.8	189.4
2003.....	19.1	0.8	0.9	1.0	1.5	2.2	3.9	9.3	26.8	70.3	143.3	198.0
2002.....	19.5	0.7	0.8	0.9	1.7	2.3	3.9	9.9	28.2	73.9	143.6	197.5
2001.....	19.8	0.7	1.0	0.9	1.8	2.5	4.1	10.4	29.5	74.6	145.2	199.9
2000.....	20.1	1.0	0.9	0.9	1.8	2.3	4.1	10.7	30.0	76.3	149.2	204.6
1999.....	20.2	0.8	0.9	0.9	1.9	2.4	4.2	10.8	31.8	77.8	148.0	199.4

All races 3/, male

2006.....	20.3	*	0.9	0.9	2.0	2.4	4.1	10.5	30.8	82.4	183.1	244.7
2005.....	20.6	*	1.0	1.0	1.9	2.3	4.0	10.4	31.1	86.6	183.0	262.4
2004.....	20.4	*	0.9	1.0	1.9	2.4	4.2	11.0	32.1	87.4	181.3	254.9
2003.....	21.0	*	1.0	1.1	1.8	2.6	4.7	11.4	33.9	90.4	184.8	268.7
2002.....	21.4	*	0.8	1.0	2.0	2.7	4.6	11.9	35.0	95.2	188.0	269.1
2001.....	21.5	*	1.0	1.0	2.1	2.9	5.1	12.7	36.3	94.7	187.0	275.3
2000.....	21.6	1.1	1.0	1.1	2.3	2.7	4.8	13.0	36.1	96.8	192.6	281.8
1999.....	21.7	*	1.0	1.0	2.2	2.8	5.1	13.2	38.8	97.3	189.0	279.0

All races 3/, female

2006.....	16.5	*	0.8	0.6	1.1	1.7	2.9	6.6	18.3	50.7	106.8	158.7
2005.....	16.6	*	0.6	0.7	1.2	1.8	2.8	7.0	19.6	50.5	108.6	156.4
2004.....	16.8	*	0.6	0.7	1.2	1.7	2.7	7.0	19.9	52.0	111.8	160.0
2003.....	17.3	1.1	0.8	0.8	1.2	1.7	3.1	7.2	20.1	53.4	115.6	166.8
2002.....	17.7	*	0.7	0.8	1.4	1.9	3.2	7.9	21.9	56.2	114.1	166.5
2001.....	18.1	*	1.0	0.7	1.4	2.1	3.2	8.1	23.1	57.9	117.7	168.4
2000.....	18.6	*	0.9	0.7	1.4	1.9	3.4	8.5	24.4	59.5	120.9	173.1
1999.....	18.7	*	0.8	0.7	1.5	2.0	3.4	8.5	25.3	61.9	121.3	167.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

White, both sexes

2006 4/.....	20.1	0.8	0.9	0.8	1.6	2.0	3.4	8.3	24.1	66.5	142.3	192.2
2005 4/.....	20.3	0.6	0.7	0.9	1.6	1.8	3.3	8.3	25.0	68.3	143.6	197.3
2004 4/.....	20.2	0.6	0.8	0.9	1.6	2.0	3.4	8.7	25.7	68.9	143.4	194.9
2003 4/.....	20.9	0.9	0.9	1.0	1.5	2.1	3.8	9.0	27.0	71.5	146.9	204.5
2002.....	21.3	*	0.8	1.0	1.8	2.2	3.9	9.6	28.0	75.4	147.6	204.0
2001.....	21.5	0.7	1.1	0.9	1.8	2.4	4.0	10.2	29.2	75.8	148.7	205.9
2000.....	21.9	1.2	1.0	1.0	1.8	2.2	4.0	10.4	30.1	78.0	153.3	208.7
1999.....	21.9	0.8	1.0	0.9	1.8	2.3	4.2	10.7	31.5	79.3	151.3	204.1

White, male

2006 4/.....	22.1	*	0.9	0.9	2.0	2.4	4.0	10.1	30.4	83.8	188.9	252.0
2005 4/.....	22.5	*	0.9	1.0	1.9	2.1	3.9	9.9	31.0	88.4	188.8	275.3
2004 4/.....	22.2	*	0.8	1.1	1.9	2.4	4.1	10.7	32.2	88.2	186.3	264.0
2003 4/.....	23.1	*	1.1	1.1	1.8	2.6	4.6	11.1	34.2	92.8	191.0	279.5
2002.....	23.4	*	0.9	1.0	2.1	2.6	4.5	11.6	34.8	97.3	193.7	279.4
2001.....	23.3	*	1.1	1.0	2.1	2.8	4.8	12.6	35.9	96.3	191.8	284.8
2000.....	23.6	*	1.0	1.1	2.2	2.6	4.8	12.6	36.1	98.7	198.2	290.2
1999.....	23.6	*	1.0	1.0	2.2	2.6	5.0	13.0	38.6	99.1	193.4	285.8

White, female

2006 4/.....	18.0	*	0.8	0.6	1.1	1.6	2.8	6.4	18.0	51.6	110.0	164.1
2005 4/.....	18.1	*	0.5	0.7	1.2	1.6	2.7	6.7	19.3	51.0	112.4	161.4
2004 4/.....	18.3	*	0.7	0.7	1.2	1.6	2.6	6.8	19.5	52.4	114.2	163.8
2003 4/.....	18.8	1.4	0.8	0.8	1.2	1.6	3.0	7.0	20.1	53.4	117.2	171.3
2002.....	19.2	*	0.8	0.9	1.4	1.8	3.2	7.5	21.5	56.8	116.7	171.2
2001.....	19.7	*	1.1	0.8	1.4	2.0	3.2	7.9	22.8	58.5	120.1	172.8
2000.....	20.3	*	1.0	0.8	1.3	1.8	3.2	8.3	24.4	60.6	123.8	175.5
1999.....	20.4	*	0.9	0.7	1.5	2.0	3.3	8.4	24.9	62.8	123.7	171.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

Black, both sexes

2006 4/.....	13.3	*	*	0.7	1.6	2.4	5.0	12.1	32.7	68.9	116.9	143.5
2005 4/.....	12.9	*	1.0	0.7	1.6	3.3	4.6	12.3	30.6	68.5	113.0	136.8
2004 4/.....	13.6	*	*	0.7	1.6	3.0	4.5	12.6	32.4	75.1	123.4	154.0
2003 4/.....	13.4	*	*	0.8	1.4	2.8	5.2	13.0	31.1	73.1	126.1	150.5
2002.....	13.6	*	*	0.8	1.5	2.9	4.8	13.6	35.9	76.0	122.3	152.3
2001.....	14.1	*	*	1.0	1.8	3.4	5.6	13.2	36.7	78.0	126.9	155.7
2000.....	13.9	*	0.8	0.6	2.2	2.9	5.3	14.7	35.8	75.0	121.0	175.5
1999.....	14.1	*	*	0.9	2.3	2.9	5.5	13.8	38.9	76.2	126.8	162.0

Black, male

2006 4/.....	14.9	*	*	1.0	1.9	2.8	6.5	15.3	42.9	91.1	153.7	194.8
2005 4/.....	14.1	*	*	1.0	1.9	3.7	5.3	15.5	38.6	88.3	151.9	164.5
2004 4/.....	14.4	*	*	0.8	1.9	3.3	5.2	15.8	39.7	98.5	150.6	193.7
2003 4/.....	13.9	*	*	0.9	1.6	3.4	6.4	16.6	40.0	87.0	148.9	184.5
2002.....	14.6	*	*	1.0	1.4	3.6	6.2	16.1	44.4	97.8	152.6	202.6
2001.....	15.1	*	*	1.3	2.0	3.9	7.3	16.0	46.0	99.2	159.5	206.5
2000.....	14.7	*	*	0.8	2.7	3.5	5.7	17.9	44.5	96.0	155.0	224.9
1999.....	15.1	*	*	1.1	2.6	3.8	6.8	17.3	47.4	95.4	161.1	220.3

Black, female

2006 4/.....	11.9	*	*	*	1.3	2.0	3.8	9.3	24.6	53.0	96.0	122.5
2005 4/.....	11.8	*	*	*	1.2	2.8	3.9	9.5	24.1	54.3	90.9	125.7
2004 4/.....	12.8	*	*	0.7	1.3	2.7	3.9	9.8	26.6	58.3	108.1	138.2
2003 4/.....	12.9	*	*	0.8	1.1	2.4	4.2	9.9	23.8	63.3	113.2	137.3
2002.....	12.8	*	*	0.6	1.5	2.2	3.5	11.4	29.1	60.5	105.1	133.1
2001.....	13.1	*	*	0.6	1.6	2.9	4.0	10.8	29.2	63.0	108.6	136.8
2000.....	13.1	*	*	*	1.8	2.4	5.0	12.0	28.8	60.2	101.8	157.1
1999.....	13.2	*	*	0.7	2.0	2.1	4.4	10.8	32.0	62.6	107.6	140.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hodgkin's disease (C81)

All races 3/, both sexes

2006.....	0.4	*	*	*	0.2	0.3	0.4	0.3	0.6	1.1	2.2	2.6
2005.....	0.4	*	*	*	0.1	0.3	0.3	0.4	0.6	1.1	2.3	2.1
2004.....	0.4	*	*	*	0.2	0.3	0.3	0.4	0.7	1.2	1.9	2.2
2003.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.2	2.2	2.4
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.6	1.3	1.9	2.7
2001.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.2	2.2	2.2
2000.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.1	2.5
1999.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.5	2.1	2.4

All races 3/, male

2006.....	0.5	*	*	*	0.2	0.4	0.5	0.5	0.8	1.4	2.6	3.7
2005.....	0.5	*	*	*	0.1	0.3	0.3	0.4	0.8	1.4	2.5	3.1
2004.....	0.5	*	*	*	0.2	0.3	0.3	0.5	0.9	1.6	2.5	3.2
2003.....	0.5	*	*	*	0.1	0.4	0.4	0.5	0.9	1.5	2.6	2.6
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.8	1.8	2.5	3.6
2001.....	0.5	*	*	*	0.2	0.4	0.5	0.6	0.7	1.6	2.6	2.6
2000.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.7	1.6	2.5	3.7
1999.....	0.6	*	*	*	0.3	0.4	0.5	0.5	1.0	1.9	2.5	3.8

All races 3/, female

2006.....	0.4	*	*	*	0.1	0.3	0.2	0.2	0.4	0.8	2.0	2.1
2005.....	0.4	*	*	*	0.2	0.3	0.2	0.3	0.4	0.9	2.2	1.6
2004.....	0.4	*	*	*	0.1	0.3	0.2	0.3	0.5	0.9	1.5	1.7
2003.....	0.4	*	*	*	0.2	0.4	0.3	0.3	0.5	0.9	1.9	2.3
2002.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	1.6	2.3
2001.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.8	2.0	2.0
2000.....	0.4	*	*	*	0.2	0.3	0.3	0.3	0.6	1.0	1.8	2.0
1999.....	0.4	*	*	*	0.2	0.3	0.3	0.4	0.5	1.2	1.9	1.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hodgkin's disease (C81)

White, both sexes

2006 4/.....	0.5	*	*	*	0.2	0.3	0.4	0.3	0.6	1.1	2.3	2.6
2005 4/.....	0.5	*	*	*	0.1	0.2	0.2	0.4	0.6	1.2	2.5	2.1
2004 4/.....	0.5	*	*	*	0.2	0.3	0.3	0.4	0.7	1.3	2.0	2.3
2003 4/.....	0.5	*	*	*	0.2	0.4	0.4	0.4	0.7	1.2	2.2	2.5
2002.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.3	2.1	2.9
2001.....	0.5	*	*	*	0.2	0.4	0.4	0.5	0.6	1.3	2.3	2.4
2000.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.7	1.3	2.2	2.6
1999.....	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.6	2.3	2.4

White, male

2006 4/.....	0.6	*	*	*	0.2	0.4	0.5	0.5	0.8	1.4	2.7	3.7
2005 4/.....	0.5	*	*	*	0.1	0.2	0.3	0.4	0.8	1.5	2.6	3.4
2004 4/.....	0.5	*	*	*	0.2	0.3	0.3	0.5	1.0	1.7	2.6	3.5
2003 4/.....	0.5	*	*	*	0.1	0.5	0.5	0.5	0.9	1.5	2.7	2.7
2002.....	0.6	*	*	*	0.2	0.4	0.4	0.5	0.8	1.9	2.6	3.9
2001.....	0.6	*	*	*	0.2	0.4	0.5	0.6	0.7	1.7	2.7	2.7
2000.....	0.5	*	*	*	0.2	0.4	0.4	0.6	0.7	1.7	2.7	3.9
1999.....	0.6	*	*	*	0.3	0.4	0.5	0.5	1.1	2.0	2.7	3.8

White, female

2006 4/.....	0.4	*	*	*	*	0.3	0.3	0.2	0.4	0.9	2.1	2.1
2005 4/.....	0.4	*	*	*	0.2	0.2	0.2	0.3	0.5	0.9	2.4	1.5
2004 4/.....	0.4	*	*	*	0.1	0.3	0.2	0.3	0.5	0.9	1.6	1.7
2003 4/.....	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	2.0	2.4
2002.....	0.5	*	*	*	0.2	0.4	0.3	0.4	0.5	0.9	1.7	2.5
2001.....	0.4	*	*	*	0.2	0.4	0.3	0.4	0.4	0.9	2.0	2.2
2000.....	0.5	*	*	*	0.2	0.4	0.3	0.3	0.6	1.0	1.9	2.1
1999.....	0.5	*	*	*	0.2	0.3	0.3	0.4	0.5	1.3	2.1	1.8



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Non-Hodgkin's lymphoma (C82-C85)

All races 3/, both sexes

2006.....	6.9	*	*	0.1	0.3	0.5	1.2	3.3	9.1	24.0	53.7	72.3
2005.....	7.0	*	*	0.1	0.3	0.6	1.3	3.3	9.6	24.8	54.5	75.2
2004.....	7.1	*	*	0.1	0.3	0.6	1.2	3.7	9.9	26.1	54.6	75.0
2003.....	7.4	*	*	0.1	0.3	0.6	1.4	3.8	10.8	26.8	56.1	78.8
2002.....	7.6	*	*	0.1	0.3	0.7	1.5	3.9	11.5	28.8	57.2	77.3
2001.....	7.8	*	*	0.1	0.4	0.8	1.7	4.4	12.2	29.3	58.6	79.6
2000.....	8.1	*	*	0.1	0.5	0.7	1.7	4.6	13.1	30.4	61.0	80.2
1999.....	8.2	*	*	0.1	0.4	0.7	1.8	4.8	13.3	31.7	61.8	77.7

All races 3/, male

2006.....	7.4	*	*	0.1	0.4	0.7	1.5	4.3	11.8	29.7	66.9	89.6
2005.....	7.6	*	*	0.2	0.4	0.7	1.7	4.4	11.9	31.4	67.4	100.5
2004.....	7.6	*	*	0.1	0.4	0.8	1.7	4.8	12.3	32.5	67.3	95.8
2003.....	8.0	*	*	0.1	0.4	0.8	1.8	4.8	13.8	33.9	69.1	103.6
2002.....	8.1	*	*	0.2	0.4	0.9	2.0	5.0	14.4	36.0	70.7	99.7
2001.....	8.4	*	*	0.1	0.5	0.9	2.2	5.8	15.0	36.5	71.3	106.6
2000.....	8.6	*	*	0.1	0.6	0.9	2.1	6.0	15.9	37.4	74.5	105.9
1999.....	8.6	*	*	0.1	0.5	0.9	2.3	6.2	16.2	38.3	74.7	104.4

All races 3/, female

2006.....	6.4	*	*	*	0.2	0.4	0.9	2.3	6.7	19.2	44.7	64.2
2005.....	6.5	*	*	*	0.2	0.4	0.9	2.3	7.5	19.3	45.7	63.5
2004.....	6.6	*	*	*	0.3	0.5	0.8	2.6	7.6	20.7	46.0	65.7
2003.....	6.8	*	*	*	0.3	0.4	1.0	2.8	8.0	20.9	47.4	67.8
2002.....	7.1	*	*	*	0.2	0.5	1.1	3.0	8.8	22.8	48.2	67.6
2001.....	7.3	*	*	*	0.3	0.6	1.1	3.1	9.7	23.3	50.2	68.3
2000.....	7.6	*	*	*	0.3	0.6	1.2	3.3	10.5	24.7	52.1	69.8
1999.....	7.7	*	*	*	0.3	0.6	1.2	3.4	10.6	26.4	53.4	66.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Non-Hodgkin's lymphoma (C82-C85)

White, both sexes

2006 4/.....	7.7	*	*	0.1	0.3	0.5	1.1	3.2	9.3	25.3	56.9	76.4
2005 4/.....	7.9	*	*	0.1	0.3	0.5	1.2	3.2	9.9	26.2	57.8	79.7
2004 4/.....	8.0	*	*	0.1	0.3	0.6	1.2	3.6	10.2	27.4	57.6	78.8
2003 4/.....	8.3	*	*	0.1	0.3	0.6	1.4	3.8	11.2	28.2	59.1	83.3
2002.....	8.5	*	*	0.1	0.3	0.6	1.5	3.9	11.9	30.2	60.3	82.1
2001.....	8.8	*	*	0.1	0.3	0.7	1.6	4.5	12.6	30.7	61.6	83.4
2000.....	9.1	*	*	0.1	0.4	0.7	1.7	4.7	13.7	32.1	64.3	84.3
1999.....	9.2	*	*	0.1	0.4	0.7	1.7	4.9	13.8	33.5	64.9	81.2

White, male

2006 4/.....	8.2	*	*	0.1	0.4	0.6	1.3	4.2	11.9	31.1	70.9	94.1
2005 4/.....	8.5	*	*	0.2	0.4	0.5	1.6	4.2	12.2	33.0	71.4	107.0
2004 4/.....	8.5	*	*	0.1	0.4	0.8	1.7	4.7	12.6	33.9	70.8	101.2
2003 4/.....	8.9	*	*	*	0.4	0.7	1.8	4.8	14.4	35.9	73.0	109.7
2002.....	9.1	*	*	0.2	0.5	0.8	1.9	5.0	14.8	37.7	74.4	105.6
2001.....	9.3	*	*	0.1	0.4	0.9	2.1	5.8	15.3	38.0	74.7	111.5
2000.....	9.6	*	*	0.1	0.6	0.8	2.2	6.0	16.5	39.2	78.4	111.8
1999.....	9.6	*	*	0.1	0.5	0.8	2.2	6.3	16.6	40.2	78.0	108.0

White, female

2006 4/.....	7.2	*	*	*	0.1	0.4	0.8	2.3	6.8	20.3	47.2	68.1
2005 4/.....	7.3	*	*	*	0.1	0.4	0.8	2.3	7.7	20.3	48.4	67.1
2004 4/.....	7.5	*	*	*	0.3	0.4	0.8	2.6	7.9	21.9	48.6	68.7
2003 4/.....	7.7	*	*	*	0.3	0.4	1.0	2.8	8.3	21.7	49.7	71.6
2002.....	8.0	*	*	*	0.2	0.5	1.1	2.9	9.1	23.9	50.8	71.8
2001.....	8.2	*	*	*	0.3	0.5	1.1	3.1	10.1	24.5	52.9	71.6
2000.....	8.6	*	*	*	0.2	0.6	1.1	3.4	11.1	26.1	55.0	73.1
1999.....	8.8	*	*	*	0.3	0.5	1.2	3.5	11.1	27.9	56.3	70.4



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Non-Hodgkin's lymphoma (C82-C85)

Black, both sexes

2006 4/.....	3.7	*	*	*	0.4	0.8	2.1	4.4	10.0	17.0	27.8	34.0
2005 4/.....	3.6	*	*	*	0.5	1.0	2.0	4.2	9.1	16.8	26.3	33.7
2004 4/.....	3.7	*	*	*	0.5	1.1	1.5	4.8	9.0	18.5	27.9	38.6
2003 4/.....	3.6	*	*	*	0.4	1.1	1.9	4.8	8.9	18.2	28.3	31.2
2002.....	3.9	*	*	*	0.4	1.2	2.0	4.7	10.8	21.2	30.2	30.6
2001.....	4.0	*	*	*	0.5	1.2	2.2	5.0	10.9	21.0	30.5	39.3
2000.....	3.7	*	*	*	0.6	1.0	2.1	5.0	10.5	19.3	26.8	36.3
1999.....	4.0	*	*	*	0.5	1.1	2.4	4.7	11.1	20.1	31.8	40.5

Black, male

2006 4/.....	4.2	*	*	*	*	1.0	2.8	6.4	13.6	21.7	31.9	42.0
2005 4/.....	4.0	*	*	*	*	1.3	2.3	6.1	12.1	22.7	30.3	40.0
2004 4/.....	4.0	*	*	*	0.6	1.4	1.9	6.5	11.8	23.4	32.9	41.7
2003 4/.....	3.8	*	*	*	*	1.3	2.3	6.5	11.2	20.4	31.0	34.4
2002.....	4.3	*	*	*	*	1.5	2.9	5.5	13.7	27.8	35.3	42.9
2001.....	4.5	*	*	*	0.7	1.3	3.2	6.7	13.6	27.7	36.8	54.5
2000.....	4.2	*	*	*	*	1.2	2.6	6.9	14.5	24.9	31.2	43.8
1999.....	4.6	*	*	*	*	1.5	3.2	6.6	14.6	25.6	40.6	59.8

Black, female

2006 4/.....	3.3	*	*	*	*	0.7	1.5	2.7	7.1	13.7	25.4	30.7
2005 4/.....	3.1	*	*	*	*	0.8	1.7	2.6	6.7	12.6	24.0	31.1
2004 4/.....	3.3	*	*	*	*	0.8	1.2	3.3	6.8	14.9	25.2	37.4
2003 4/.....	3.4	*	*	*	*	0.8	1.5	3.3	7.1	16.6	26.8	30.0
2002.....	3.5	*	*	*	*	0.8	1.3	4.0	8.5	16.5	27.3	25.9
2001.....	3.6	*	*	*	*	1.1	1.3	3.6	8.7	16.3	26.9	33.7
2000.....	3.3	*	*	*	*	0.9	1.6	3.3	7.3	15.4	24.4	33.5
1999.....	3.5	*	*	*	*	*	1.7	3.2	8.2	16.3	26.9	33.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Leukemia (C91-C95)

All races 3/, both sexes

2006.....	7.3	0.8	0.8	0.6	1.1	1.1	1.6	3.2	8.9	25.0	52.6	76.9
2005.....	7.3	0.5	0.7	0.7	1.1	1.1	1.6	3.3	8.8	25.1	52.3	78.7
2004.....	7.3	0.6	0.8	0.8	1.0	1.0	1.6	3.3	9.2	25.4	52.4	78.8
2003.....	7.4	0.7	0.8	0.8	1.0	1.1	1.8	3.3	9.3	25.7	53.9	80.3
2002.....	7.5	0.5	0.7	0.8	1.2	1.2	1.7	3.6	9.8	26.4	53.1	80.5
2001.....	7.5	0.6	1.0	0.8	1.2	1.3	1.7	3.8	10.0	26.7	53.3	81.4
2000.....	7.6	1.0	0.9	0.8	1.2	1.2	1.8	3.7	9.9	27.2	54.5	83.6
1999.....	7.5	0.7	0.8	0.7	1.2	1.2	1.8	3.7	10.6	27.2	52.8	82.4

All races 3/, male

2006.....	8.4	*	0.8	0.7	1.4	1.3	1.8	3.7	11.4	33.1	75.9	105.1
2005.....	8.4	*	1.0	0.8	1.4	1.3	1.6	3.6	11.3	34.0	75.0	112.4
2004.....	8.3	*	0.9	0.9	1.3	1.2	1.8	3.8	11.9	34.4	73.2	110.7
2003.....	8.4	*	0.9	0.9	1.2	1.4	2.0	4.0	11.9	35.1	75.3	111.8
2002.....	8.5	*	0.7	0.8	1.4	1.4	1.9	4.2	12.2	36.3	74.9	114.5
2001.....	8.5	*	1.0	0.9	1.4	1.5	1.9	4.2	12.8	35.7	74.2	117.5
2000.....	8.5	1.1	0.9	0.9	1.5	1.4	1.9	4.3	12.5	37.2	75.7	119.8
1999.....	8.4	*	0.8	0.8	1.5	1.4	1.9	4.3	13.1	36.8	73.2	119.2

All races 3/, female

2006.....	6.3	*	0.8	0.6	0.8	0.9	1.5	2.7	6.7	18.2	36.7	63.7
2005.....	6.2	*	0.5	0.6	0.8	1.0	1.5	3.0	6.4	17.5	37.0	63.3
2004.....	6.3	*	0.6	0.7	0.8	0.9	1.4	2.8	6.8	17.8	38.4	64.5
2003.....	6.4	1.1	0.8	0.7	0.7	0.8	1.6	2.7	7.0	17.8	39.6	66.4
2002.....	6.5	*	0.6	0.7	1.0	1.0	1.5	3.1	7.6	18.1	38.7	65.7
2001.....	6.6	*	1.0	0.7	1.0	1.1	1.5	3.3	7.5	19.2	39.5	66.3
2000.....	6.7	*	0.9	0.6	0.9	1.0	1.7	3.2	7.6	19.0	40.6	68.8
1999.....	6.7	*	0.7	0.7	1.0	1.1	1.6	3.2	8.3	19.3	39.6	67.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Leukemia (C91-C95)

White, both sexes

2006 4/.....	8.1	0.8	0.8	0.6	1.1	1.2	1.7	3.3	9.1	26.1	54.7	79.9
2005 4/.....	8.1	*	0.7	0.7	1.2	1.1	1.6	3.3	9.0	26.1	54.6	82.4
2004 4/.....	8.0	*	0.7	0.8	1.1	1.0	1.6	3.3	9.4	26.0	54.2	81.6
2003 4/.....	8.2	0.9	0.9	0.9	1.0	1.1	1.8	3.4	9.6	26.6	56.0	83.4
2002.....	8.2	*	0.7	0.8	1.2	1.2	1.8	3.6	9.9	27.5	54.9	83.0
2001.....	8.3	0.6	1.1	0.8	1.2	1.3	1.7	3.7	10.2	27.5	55.1	84.7
2000.....	8.3	1.2	1.0	0.8	1.2	1.1	1.8	3.7	10.0	28.3	56.6	86.0
1999.....	8.3	0.8	0.9	0.7	1.2	1.2	1.8	3.7	10.8	28.0	54.5	85.4

White, male

2006 4/.....	9.3	*	0.8	0.7	1.4	1.4	1.8	3.8	11.6	34.3	78.6	108.7
2005 4/.....	9.3	*	0.9	0.8	1.4	1.3	1.6	3.6	11.6	35.4	77.6	118.7
2004 4/.....	9.1	*	0.8	0.9	1.3	1.2	1.8	3.8	12.2	35.0	75.8	115.3
2003 4/.....	9.4	*	1.0	1.0	1.2	1.3	1.9	4.1	12.2	36.4	78.7	116.9
2002.....	9.3	*	0.8	0.8	1.5	1.4	2.0	4.1	12.3	37.5	77.4	119.1
2001.....	9.3	*	1.1	0.8	1.5	1.5	1.9	4.3	13.0	36.7	76.8	123.1
2000.....	9.4	*	0.9	0.9	1.4	1.3	1.9	4.2	12.6	38.5	78.5	123.6
1999.....	9.2	*	0.9	0.8	1.5	1.3	1.9	4.2	13.3	37.9	75.5	124.1

White, female

2006 4/.....	7.0	*	0.8	0.6	0.8	0.9	1.5	2.8	6.8	19.0	38.1	66.4
2005 4/.....	6.9	*	0.5	0.6	0.9	1.0	1.5	3.0	6.6	18.1	38.7	65.7
2004 4/.....	6.9	*	0.7	0.7	0.8	0.9	1.4	2.8	6.8	18.4	39.5	66.4
2003 4/.....	7.0	1.4	0.8	0.7	0.7	0.9	1.6	2.7	7.1	18.2	40.6	68.6
2002.....	7.1	*	0.6	0.8	1.0	0.9	1.6	3.1	7.6	18.9	39.8	67.3
2001.....	7.3	*	1.1	0.7	1.0	1.1	1.6	3.2	7.6	19.7	40.8	68.6
2000.....	7.4	*	1.0	0.7	0.9	0.9	1.6	3.2	7.7	19.8	42.1	70.8
1999.....	7.3	*	0.8	0.7	1.0	1.2	1.6	3.2	8.4	19.8	40.6	69.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Leukemia (C91-C95)

Black, both sexes

2006 4/.....	4.6	*	*	0.6	1.0	1.1	1.8	3.3	9.5	22.6	40.2	55.5
2005 4/.....	4.4	*	1.0	0.5	1.0	1.5	1.6	4.0	8.4	21.2	38.4	52.1
2004 4/.....	4.8	*	*	0.6	1.0	1.3	2.0	4.0	10.3	25.2	41.1	60.1
2003 4/.....	4.6	*	*	0.7	0.9	1.2	2.3	3.7	9.6	23.1	41.6	59.2
2002.....	4.7	*	*	0.6	0.9	1.2	1.6	4.3	11.2	22.3	42.1	64.7
2001.....	4.7	*	*	0.9	1.1	1.4	2.1	4.2	9.8	24.9	41.6	57.1
2000.....	4.8	*	*	0.5	1.4	1.6	2.0	4.8	10.7	22.7	38.2	66.0
1999.....	4.7	*	*	0.8	1.4	1.2	1.9	4.3	10.6	24.8	41.2	55.8

Black, male

2006 4/.....	5.3	*	*	0.8	1.2	1.1	2.4	3.8	12.5	31.9	59.8	84.9
2005 4/.....	4.9	*	*	0.7	1.3	1.5	1.8	4.3	10.6	28.6	58.2	62.3
2004 4/.....	5.3	*	*	0.7	1.1	1.4	2.2	4.9	12.8	37.0	52.3	79.5
2003 4/.....	5.0	*	*	0.7	1.1	1.6	2.8	4.5	12.3	31.1	53.3	77.2
2002.....	5.1	*	*	0.7	1.0	1.4	1.8	5.2	13.7	31.5	56.8	86.8
2001.....	5.2	*	*	1.1	1.2	1.7	2.5	4.5	13.0	34.0	55.0	77.2
2000.....	5.3	*	*	0.6	1.9	1.9	2.0	5.5	13.8	31.6	54.8	98.0
1999.....	5.2	*	*	1.0	1.6	1.5	2.0	5.5	13.4	32.9	54.7	77.3

Black, female

2006 4/.....	3.9	*	*	*	0.8	1.1	1.4	2.9	7.1	16.0	29.1	43.5
2005 4/.....	4.0	*	*	*	0.7	1.5	1.4	3.6	6.6	15.8	27.1	48.0
2004 4/.....	4.4	*	*	*	0.8	1.2	1.8	3.2	8.3	16.7	34.8	52.4
2003 4/.....	4.2	*	*	0.7	0.6	0.8	1.9	3.0	7.4	17.4	34.9	52.3
2002.....	4.3	*	*	*	0.7	1.1	1.4	3.6	9.2	15.7	33.8	56.3
2001.....	4.3	*	*	0.6	1.0	1.1	1.7	3.9	7.3	18.4	34.0	49.7
2000.....	4.2	*	*	*	0.9	1.3	2.1	4.3	8.1	16.4	28.8	54.1
1999.....	4.3	*	*	0.6	1.3	1.0	1.7	3.3	8.3	19.0	33.6	47.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

All races 3/, both sexes

2006.....	3.7	*	*	*	*	*	0.3	1.7	5.7	15.0	29.1	33.8
2005.....	3.8	*	*	*	*	*	0.3	1.6	6.1	15.9	29.5	33.6
2004.....	3.7	*	*	*	*	*	0.3	1.6	5.9	15.4	30.7	33.1
2003.....	3.9	*	*	*	*	*	0.3	1.7	5.9	16.5	31.1	36.3
2002.....	4.0	*	*	*	*	*	0.3	1.8	6.3	17.4	31.2	36.7
2001.....	3.9	*	*	*	*	*	0.3	1.7	6.6	17.4	30.9	36.3
2000.....	3.9	*	*	*	*	*	0.3	1.9	6.3	17.3	31.5	37.8
1999.....	3.9	*	*	*	*	0.1	0.3	1.8	7.1	17.2	31.0	36.6

All races 3/, male

2006.....	4.0	*	*	*	*	*	0.4	2.0	6.8	18.1	37.5	45.8
2005.....	4.1	*	*	*	*	*	0.4	2.0	7.0	19.7	38.0	46.1
2004.....	3.9	*	*	*	*	*	0.4	1.9	6.9	18.8	38.1	44.6
2003.....	4.1	*	*	*	*	*	0.4	2.1	7.4	19.8	37.7	50.4
2002.....	4.2	*	*	*	*	*	0.3	2.2	7.6	21.0	39.7	50.8
2001.....	4.1	*	*	*	*	*	0.4	2.1	7.8	20.8	38.6	48.0
2000.....	4.0	*	*	*	*	*	0.4	2.1	6.9	20.4	39.6	51.8
1999.....	4.1	*	*	*	*	*	0.4	2.1	8.5	20.1	38.3	51.3

All races 3/, female

2006.....	3.4	*	*	*	*	*	0.3	1.4	4.6	12.4	23.3	28.2
2005.....	3.5	*	*	*	*	*	0.2	1.3	5.2	12.8	23.7	27.9
2004.....	3.5	*	*	*	*	*	0.3	1.3	5.0	12.5	25.7	28.0
2003.....	3.7	*	*	*	*	*	0.2	1.4	4.6	13.8	26.7	30.1
2002.....	3.7	*	*	*	*	*	0.3	1.5	5.1	14.3	25.5	30.6
2001.....	3.8	*	*	*	*	*	0.3	1.3	5.5	14.5	25.8	31.3
2000.....	3.8	*	*	*	*	*	0.3	1.6	5.7	14.7	26.3	32.2
1999.....	3.8	*	*	*	*	*	0.3	1.5	5.9	14.8	26.2	30.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

White, both sexes

2006 4/.....	3.7	*	*	*	*	*	0.3	1.4	5.0	14.0	28.1	32.8
2005 4/.....	3.8	*	*	*	*	*	0.3	1.3	5.4	14.7	28.6	32.9
2004 4/.....	3.8	*	*	*	*	*	0.3	1.4	5.3	14.1	29.4	32.0
2003 4/.....	3.9	*	*	*	*	*	0.3	1.4	5.4	15.4	29.5	35.0
2002.....	4.0	*	*	*	*	*	0.2	1.6	5.6	16.3	30.2	35.7
2001.....	3.9	*	*	*	*	*	0.3	1.5	5.8	16.3	29.5	35.0
2000.....	3.9	*	*	*	*	*	0.3	1.6	5.6	16.2	30.0	35.2
1999.....	3.9	*	*	*	*	*	0.3	1.6	6.1	16.1	29.4	34.8

White, male

2006 4/.....	4.0	*	*	*	*	*	0.4	1.7	6.0	16.9	36.5	45.0
2005 4/.....	4.2	*	*	*	*	*	0.4	1.7	6.3	18.5	37.0	45.8
2004 4/.....	4.0	*	*	*	*	*	0.3	1.6	6.4	17.6	36.8	43.5
2003 4/.....	4.2	*	*	*	*	*	0.4	1.8	6.8	18.9	36.5	49.7
2002.....	4.3	*	*	*	*	*	0.3	2.0	6.9	20.1	39.0	50.3
2001.....	4.2	*	*	*	*	*	0.4	1.9	6.8	19.7	37.4	47.1
2000.....	4.1	*	*	*	*	*	0.3	1.8	6.3	19.3	38.2	50.2
1999.....	4.1	*	*	*	*	*	0.3	1.9	7.6	19.0	36.9	49.7

White, female

2006 4/.....	3.4	*	*	*	*	*	0.2	1.1	4.1	11.4	22.3	27.1
2005 4/.....	3.5	*	*	*	*	*	0.2	1.0	4.6	11.5	22.8	26.9
2004 4/.....	3.5	*	*	*	*	*	0.2	1.1	4.3	11.1	24.3	26.8
2003 4/.....	3.6	*	*	*	*	*	0.2	1.1	4.1	12.4	24.8	28.5
2002.....	3.7	*	*	*	*	*	0.2	1.2	4.4	13.1	24.3	29.4
2001.....	3.7	*	*	*	*	*	0.2	1.1	4.8	13.4	24.3	29.9
2000.....	3.8	*	*	*	*	*	0.2	1.4	5.0	13.6	24.6	29.1
1999.....	3.8	*	*	*	*	*	0.2	1.2	4.8	13.7	24.5	28.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Multiple myeloma and immunoproliferative neoplasms (C88,C90)

Black, both sexes

2006 4/.....	4.7	*	*	*	*	*	0.7	3.9	12.6	28.2	47.6	51.6
2005 4/.....	4.6	*	*	*	*	*	0.6	3.5	12.4	29.7	47.1	49.4
2004 4/.....	4.7	*	*	*	*	*	0.7	3.4	12.4	30.4	53.0	53.9
2003 4/.....	4.8	*	*	*	*	*	0.7	4.2	11.8	30.8	55.3	59.8
2002.....	4.7	*	*	*	*	*	0.7	4.1	13.1	31.5	48.7	55.5
2001.....	4.9	*	*	*	*	*	0.9	3.6	15.1	31.3	53.1	58.3
2000.....	5.0	*	*	*	*	*	0.8	4.5	13.9	32.1	55.1	71.6
1999.....	5.0	*	*	*	*	*	0.9	4.3	16.6	30.2	53.1	63.1

Black, male

2006 4/.....	4.9	*	*	*	*	*	0.7	4.5	15.9	35.9	60.1	65.2
2005 4/.....	4.7	*	*	*	*	*	*	4.4	14.6	35.7	61.7	61.4
2004 4/.....	4.7	*	*	*	*	*	0.7	3.9	14.2	36.6	64.0	71.5
2003 4/.....	4.7	*	*	*	*	*	0.9	5.2	15.3	33.6	63.5	73.0
2002.....	4.7	*	*	*	*	*	0.8	4.6	15.7	37.1	59.1	71.8
2001.....	4.9	*	*	*	*	*	1.0	4.3	18.4	36.6	65.6	71.5
2000.....	4.8	*	*	*	*	*	*	5.0	15.0	38.5	67.8	79.6
1999.....	4.9	*	*	*	*	*	1.1	4.4	18.6	35.1	64.6	80.9

Black, female

2006 4/.....	4.4	*	*	*	*	*	0.7	3.4	9.9	22.6	40.5	46.1
2005 4/.....	4.4	*	*	*	*	*	*	2.8	10.7	25.4	38.8	44.6
2004 4/.....	4.7	*	*	*	*	*	0.7	3.1	11.0	26.0	46.7	46.9
2003 4/.....	4.9	*	*	*	*	*	*	3.3	9.0	28.7	50.7	54.7
2002.....	4.7	*	*	*	*	*	*	3.7	11.1	27.6	42.9	49.3
2001.....	4.9	*	*	*	*	*	0.8	3.0	12.5	27.6	46.1	53.5
2000.....	5.2	*	*	*	*	*	0.9	4.0	12.9	27.6	47.9	68.7
1999.....	5.1	*	*	*	*	*	0.7	4.2	15.0	26.8	46.6	56.5









Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes

2006.....	21.1	0.5	0.7	0.6	1.2	1.8	4.1	12.7	33.5	75.0	140.1	210.1
2005.....	21.2	*	0.8	0.7	1.4	1.7	4.2	13.1	34.3	75.2	141.6	212.9
2004.....	21.1	0.6	0.8	0.7	1.3	1.7	4.3	12.7	34.4	76.8	143.0	213.1
2003.....	21.5	*	0.7	0.7	1.3	1.7	4.2	13.8	35.2	77.9	146.7	224.0
2002.....	21.8	0.6	0.8	0.7	1.3	1.8	4.5	13.7	36.6	80.8	146.7	230.1
2001.....	22.0	*	0.8	0.7	1.3	1.8	4.6	14.1	37.3	82.0	148.7	234.5
2000.....	22.4	0.6	0.8	0.6	1.3	1.8	4.6	14.1	38.3	85.5	152.6	239.8
1999.....	23.7	*	0.8	0.7	1.3	1.9	5.1	15.0	41.5	91.4	160.8	250.4

All races 3/, male

2006.....	21.9	*	0.7	0.6	1.6	2.2	4.3	14.3	40.0	89.0	173.9	259.6
2005.....	22.1	*	0.8	0.7	1.7	2.2	4.6	14.9	41.0	90.1	173.1	263.5
2004.....	21.9	*	0.7	0.8	1.7	2.1	4.6	14.4	40.6	92.5	177.9	260.6
2003.....	22.3	*	0.8	0.7	1.6	2.1	4.7	15.7	41.5	93.4	180.8	275.0
2002.....	22.7	*	0.8	0.7	1.6	2.3	5.0	15.9	43.2	98.1	180.0	290.8
2001.....	22.6	*	0.9	0.7	1.7	2.2	4.9	16.4	43.8	98.5	181.1	295.2
2000.....	23.0	*	1.0	0.7	1.6	2.1	5.1	16.0	44.9	103.2	187.8	300.7
1999.....	24.5	*	0.8	0.8	1.7	2.4	5.8	17.3	49.1	110.4	196.1	318.9

All races 3/, female

2006.....	20.2	*	0.6	0.6	0.9	1.4	3.9	11.3	27.5	63.2	117.0	186.9
2005.....	20.4	*	0.7	0.6	1.1	1.3	3.8	11.3	28.1	62.6	120.2	189.6
2004.....	20.3	*	0.9	0.6	0.9	1.4	3.9	11.0	28.6	63.7	119.5	191.8
2003.....	20.8	*	0.5	0.6	0.9	1.3	3.8	11.9	29.3	64.8	123.9	201.4
2002.....	21.0	*	0.8	0.7	0.9	1.4	4.1	11.6	30.4	66.3	124.6	203.8
2001.....	21.4	*	0.6	0.6	0.9	1.5	4.2	11.7	31.3	68.4	127.4	209.1
2000.....	21.9	*	0.7	0.6	1.0	1.4	4.2	12.2	32.2	70.8	129.6	215.0
1999.....	23.1	*	0.7	0.7	1.0	1.4	4.5	12.8	34.5	75.8	137.9	222.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

White, both sexes

2006 4/.....	22.6	*	0.6	0.6	1.2	1.8	4.0	12.3	32.9	75.6	142.8	213.9
2005 4/.....	22.7	*	0.8	0.7	1.4	1.7	4.1	12.4	33.7	75.5	143.9	217.2
2004 4/.....	22.6	*	0.8	0.7	1.3	1.8	4.1	12.0	33.6	77.0	144.8	217.6
2003 4/.....	23.1	*	0.6	0.7	1.3	1.7	4.1	13.1	34.6	78.3	148.6	227.9
2002.....	23.3	0.7	0.7	0.7	1.3	1.8	4.4	13.1	35.8	80.7	147.8	233.0
2001.....	23.5	*	0.8	0.7	1.3	1.8	4.4	13.3	36.2	81.4	149.5	238.1
2000.....	23.9	*	0.8	0.7	1.3	1.8	4.3	13.3	37.5	85.3	153.3	243.7
1999.....	25.3	*	0.8	0.7	1.3	1.9	4.9	14.3	40.2	91.3	161.2	252.4

White, male

2006 4/.....	23.6	*	0.7	0.6	1.6	2.2	4.3	13.6	39.4	90.0	177.1	266.4
2005 4/.....	23.6	*	0.9	0.7	1.7	2.1	4.6	14.2	40.1	89.8	176.2	270.3
2004 4/.....	23.4	*	0.7	0.8	1.6	2.2	4.4	13.5	39.3	92.1	180.7	266.8
2003 4/.....	23.8	*	0.7	0.7	1.6	2.1	4.6	14.8	40.6	93.5	183.4	280.9
2002.....	24.2	*	0.8	0.7	1.6	2.3	4.7	15.3	42.0	98.4	181.6	296.5
2001.....	24.0	*	1.0	0.7	1.7	2.2	4.8	15.5	42.0	97.7	182.1	300.2
2000.....	24.4	*	1.0	0.7	1.6	2.3	4.7	15.1	43.2	103.1	188.2	307.0
1999.....	25.9	*	0.9	0.8	1.6	2.4	5.6	16.2	47.4	109.9	197.5	323.6

White, female

2006 4/.....	21.7	*	0.5	0.6	0.8	1.3	3.8	10.9	26.7	63.2	118.9	189.3
2005 4/.....	21.9	*	0.6	0.6	1.1	1.2	3.6	10.8	27.6	63.1	121.6	192.8
2004 4/.....	21.8	*	0.8	0.5	0.9	1.3	3.8	10.6	28.2	64.0	120.3	195.4
2003 4/.....	22.5	*	0.5	0.6	1.0	1.2	3.6	11.4	28.9	65.4	125.0	204.4
2002.....	22.5	*	0.7	0.7	1.0	1.4	4.1	11.1	29.9	65.6	125.1	205.5
2001.....	22.9	*	0.7	0.6	0.9	1.4	4.0	11.2	30.7	67.7	127.8	212.1
2000.....	23.5	*	0.6	0.6	1.0	1.3	3.9	11.6	32.1	70.3	130.3	218.0
1999.....	24.6	*	0.7	0.6	1.1	1.4	4.2	12.4	33.4	75.8	137.4	223.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

Black, both sexes

2006 4/.....	16.8	*	1.0	0.6	1.2	2.1	5.2	18.3	46.4	85.6	136.6	186.0
2005 4/.....	17.1	*	0.8	0.9	1.2	2.3	5.4	19.5	46.3	85.7	144.4	183.9
2004 4/.....	17.2	*	0.8	0.9	1.6	1.8	5.9	19.4	47.5	90.8	147.2	186.7
2003 4/.....	17.3	*	0.9	0.8	1.1	2.3	5.8	20.7	48.2	88.2	150.1	206.3
2002.....	18.1	*	1.1	0.8	1.3	2.3	6.3	20.1	51.0	96.2	157.6	222.1
2001.....	18.3	*	*	0.7	1.3	2.3	6.0	21.7	52.9	100.8	156.9	217.1
2000.....	18.3	*	1.0	0.6	1.4	1.6	7.5	22.1	51.9	100.1	162.0	210.8
1999.....	19.5	*	*	0.7	1.4	2.0	7.5	22.9	59.6	104.7	170.9	242.6

Black, male

2006 4/.....	17.4	*	*	0.6	1.4	2.4	5.4	22.3	56.3	101.7	176.4	213.6
2005 4/.....	18.0	*	*	1.1	1.4	2.8	5.7	22.7	59.0	111.0	172.6	221.2
2004 4/.....	18.4	*	*	0.8	1.9	1.9	6.5	24.1	60.9	116.4	182.6	222.5
2003 4/.....	18.3	*	*	0.8	1.4	2.4	6.5	25.0	61.1	111.3	187.6	241.9
2002.....	18.8	*	*	0.8	1.8	2.6	7.4	24.0	63.8	116.7	195.6	267.0
2001.....	19.4	*	*	0.8	1.7	2.7	6.3	26.6	67.5	127.9	193.6	283.7
2000.....	19.4	*	*	0.7	1.9	1.8	8.5	26.2	68.1	122.6	210.4	253.7
1999.....	20.6	*	*	0.6	2.0	2.3	8.3	28.9	76.1	133.2	206.3	292.9

Black, female

2006 4/.....	16.3	*	*	*	1.0	1.9	5.1	14.9	38.4	74.0	113.9	174.7
2005 4/.....	16.3	*	*	0.8	1.0	1.9	5.2	16.8	36.0	67.5	128.3	168.8
2004 4/.....	16.2	*	*	0.9	1.3	1.8	5.5	15.3	36.8	72.5	127.1	172.5
2003 4/.....	16.5	*	*	0.7	0.8	2.2	5.1	17.0	37.8	71.8	128.8	192.4
2002.....	17.4	*	*	0.7	0.8	2.0	5.3	16.7	40.7	81.7	136.1	205.0
2001.....	17.3	*	*	0.6	0.8	1.9	5.7	17.6	41.2	81.7	136.3	192.4
2000.....	17.3	*	*	*	1.0	1.5	6.6	18.6	38.8	84.3	134.7	194.9
1999.....	18.5	*	*	0.8	0.9	1.6	6.7	17.8	46.4	84.5	151.1	223.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes

2006.....	4.7	1.6	0.4	0.2	0.2	0.4	0.7	1.8	4.2	12.5	36.4	78.0
2005.....	4.6	1.4	0.3	0.2	0.2	0.4	0.8	1.7	4.3	12.6	35.2	76.7
2004.....	4.6	1.7	0.3	0.2	0.3	0.4	0.7	1.7	4.5	12.7	35.8	77.7
2003.....	4.7	1.5	0.3	0.2	0.2	0.4	0.8	1.7	4.4	12.8	36.4	80.5
2002.....	4.6	1.7	0.4	0.2	0.2	0.4	0.8	1.8	4.5	12.8	36.1	78.7
2001.....	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.8	5.0	14.1	37.0	82.7
2000.....	4.8	1.6	0.3	0.2	0.2	0.4	0.8	1.8	5.0	13.8	37.6	84.9
1999.....	4.7	1.5	0.4	0.2	0.3	0.4	0.8	1.8	4.8	14.6	36.6	82.9

All races 3/, male

2006.....	4.7	1.6	0.3	0.2	0.2	0.3	0.8	2.0	4.8	15.7	47.5	92.3
2005.....	4.7	1.0	0.4	0.2	0.2	0.4	0.8	1.9	5.2	15.7	45.8	92.7
2004.....	4.7	1.7	0.3	0.2	0.3	0.5	0.8	1.9	5.4	15.7	45.6	98.9
2003.....	4.7	1.6	0.4	0.2	0.3	0.5	0.9	1.8	5.1	16.0	46.2	101.7
2002.....	4.6	1.5	0.4	0.2	0.2	0.4	0.9	1.9	5.3	15.8	45.9	101.1
2001.....	4.8	1.3	0.4	0.2	0.3	0.4	0.9	2.0	5.5	17.3	48.2	107.6
2000.....	4.8	1.6	0.4	0.2	0.3	0.4	0.8	2.1	5.6	17.0	47.5	108.9
1999.....	4.8	1.5	0.4	0.2	0.3	0.5	0.9	2.0	5.8	18.4	46.1	109.2

All races 3/, female

2006.....	4.7	1.6	0.5	0.2	0.2	0.4	0.7	1.6	3.6	9.8	28.8	71.4
2005.....	4.6	1.8	0.3	0.2	0.2	0.4	0.8	1.5	3.6	10.0	28.0	69.4
2004.....	4.6	1.8	0.3	0.2	0.2	0.4	0.7	1.5	3.7	10.2	29.2	68.1
2003.....	4.7	1.5	0.3	0.2	0.2	0.4	0.6	1.5	3.8	10.2	29.9	71.1
2002.....	4.6	2.0	0.4	0.2	0.2	0.4	0.8	1.7	3.7	10.4	29.6	68.9
2001.....	4.8	1.3	0.4	0.3	0.3	0.4	0.8	1.6	4.4	11.3	29.7	72.3
2000.....	4.9	1.5	0.3	0.2	0.2	0.4	0.8	1.6	4.5	11.2	31.1	75.2
1999.....	4.7	1.5	0.4	0.3	0.2	0.4	0.6	1.6	3.9	11.5	30.4	72.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

White, both sexes

2006 4/.....	5.2	1.6	0.4	0.2	0.2	0.3	0.7	1.8	4.1	12.8	37.5	80.0
2005 4/.....	5.1	1.3	0.3	0.2	0.2	0.4	0.8	1.6	4.3	12.9	36.4	79.6
2004 4/.....	5.1	1.9	0.3	0.2	0.3	0.4	0.7	1.6	4.4	12.9	36.7	80.2
2003 4/.....	5.2	1.5	0.3	0.2	0.2	0.5	0.7	1.6	4.4	13.1	37.7	83.0
2002.....	5.1	1.9	0.4	0.2	0.2	0.4	0.8	1.7	4.5	13.2	37.0	80.7
2001.....	5.3	1.2	0.4	0.2	0.2	0.4	0.8	1.9	5.0	14.3	38.0	83.9
2000.....	5.3	1.6	0.4	0.2	0.2	0.4	0.8	1.8	5.1	13.9	38.6	87.3
1999.....	5.2	1.4	0.4	0.3	0.2	0.4	0.7	1.7	4.7	14.8	37.4	85.3

White, male

2006 4/.....	5.2	1.6	*	0.2	0.2	0.3	0.7	2.1	4.8	16.1	49.0	93.7
2005 4/.....	5.2	*	0.4	0.2	0.3	0.4	0.8	1.9	5.1	16.2	47.6	97.0
2004 4/.....	5.1	1.8	0.3	0.2	0.3	0.5	0.7	1.8	5.2	16.1	46.8	103.2
2003 4/.....	5.2	1.6	0.3	0.2	0.3	0.5	0.9	1.8	5.0	16.4	47.8	105.5
2002.....	5.1	1.7	0.3	0.2	0.2	0.4	0.9	1.9	5.3	16.3	47.0	104.2
2001.....	5.3	1.2	0.4	0.2	0.2	0.4	0.9	2.1	5.6	17.8	49.4	109.9
2000.....	5.2	1.6	0.4	0.2	0.3	0.4	0.8	2.0	5.6	17.0	49.1	112.6
1999.....	5.2	1.4	0.4	0.2	0.2	0.4	0.9	2.0	5.8	18.7	47.2	114.3

White, female

2006 4/.....	5.2	1.5	0.5	0.2	0.2	0.4	0.6	1.5	3.5	10.0	29.5	73.6
2005 4/.....	5.0	1.7	*	0.2	0.2	0.4	0.8	1.4	3.5	10.0	28.7	71.6
2004 4/.....	5.0	1.9	0.3	0.2	0.2	0.4	0.6	1.4	3.6	10.2	29.9	69.9
2003 4/.....	5.1	1.5	*	0.2	0.2	0.4	0.6	1.5	3.9	10.3	30.9	73.0
2002.....	5.1	2.1	0.4	0.2	0.2	0.3	0.7	1.6	3.8	10.5	30.3	70.5
2001.....	5.2	*	0.4	0.3	0.2	0.4	0.7	1.6	4.4	11.4	30.4	73.0
2000.....	5.4	1.6	*	0.2	0.2	0.4	0.8	1.5	4.6	11.3	31.8	77.1
1999.....	5.1	1.4	0.4	0.3	0.2	0.4	0.6	1.5	3.7	11.5	30.9	73.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

Black, both sexes

2006 4/.....	3.1	*	*	*	0.4	0.5	1.1	2.5	5.4	11.6	31.6	62.3
2005 4/.....	2.9	*	*	*	*	0.5	1.3	2.5	5.4	12.8	27.9	54.5
2004 4/.....	3.1	*	*	0.3	*	0.7	1.2	2.3	6.2	12.8	31.5	57.0
2003 4/.....	2.9	*	*	*	*	0.5	1.2	2.5	5.2	13.0	29.5	57.8
2002.....	3.0	*	*	*	*	0.6	1.2	2.5	5.5	12.6	31.3	63.6
2001.....	3.1	*	*	0.4	0.3	*	1.1	2.0	5.9	14.3	31.5	73.4
2000.....	3.1	*	*	*	*	0.6	1.1	2.7	5.5	15.0	31.0	64.1
1999.....	3.2	*	*	*	0.4	0.5	1.2	2.6	6.2	15.7	33.0	61.9

Black, male

2006 4/.....	3.0	*	*	*	*	*	0.9	2.3	6.3	14.8	41.4	80.4
2005 4/.....	2.7	*	*	*	*	*	1.3	2.7	6.2	14.0	33.9	54.8
2004 4/.....	3.0	*	*	*	*	*	1.2	2.5	7.7	15.4	39.3	61.6
2003 4/.....	2.9	*	*	*	*	*	1.1	2.5	6.7	16.2	36.7	66.7
2002.....	2.8	*	*	*	*	*	1.0	2.2	7.1	14.1	38.7	79.3
2001.....	3.0	*	*	*	*	*	1.1	1.8	6.5	17.2	41.5	88.5
2000.....	3.0	*	*	*	*	*	1.2	2.8	7.0	19.2	36.0	76.1
1999.....	3.0	*	*	*	*	0.8	1.1	2.5	7.1	19.0	38.4	63.3

Black, female

2006 4/.....	3.2	*	*	*	*	*	1.3	2.6	4.7	9.4	26.0	54.8
2005 4/.....	3.2	*	*	*	*	*	1.4	2.3	4.7	11.9	24.5	54.4
2004 4/.....	3.2	*	*	*	*	0.8	1.1	2.1	5.0	11.0	27.1	55.1
2003 4/.....	3.0	*	*	*	*	*	1.2	2.4	4.0	10.8	25.5	54.3
2002.....	3.3	*	*	*	*	0.8	1.4	2.7	4.2	11.5	27.1	57.5
2001.....	3.2	*	*	*	*	*	1.0	2.2	5.3	12.3	25.9	67.8
2000.....	3.2	*	*	*	*	0.8	1.0	2.6	4.4	12.0	28.1	59.7
1999.....	3.4	*	*	*	*	*	1.2	2.7	5.6	13.4	29.9	61.3



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Anemias (D50-D64)

All races 3/, both sexes

2006.....	1.3	*	0.2	0.1	0.2	0.3	0.4	0.5	1.0	2.1	7.5	30.0
2005.....	1.6	*	0.1	0.1	0.2	0.4	0.4	0.6	1.0	2.7	9.1	36.6
2004.....	1.6	*	0.2	0.1	0.3	0.4	0.5	0.6	1.1	2.6	8.6	38.1
2003.....	1.6	0.5	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	8.8	40.7
2002.....	1.6	*	0.1	0.1	0.3	0.5	0.4	0.6	1.0	2.8	9.1	40.3
2001.....	1.6	*	0.1	0.1	0.3	0.4	0.5	0.6	1.0	2.7	9.8	42.0
2000.....	1.6	*	0.2	0.1	0.2	0.4	0.4	0.6	1.1	2.8	9.5	42.4
1999.....	1.6	*	0.2	0.1	0.3	0.4	0.4	0.6	1.0	2.6	9.5	43.8

All races 3/, male

2006.....	1.1	*	*	*	0.3	0.3	0.4	0.5	0.9	2.1	7.9	29.6
2005.....	1.2	*	*	*	0.3	0.4	0.4	0.5	1.0	2.6	9.5	33.5
2004.....	1.3	*	*	0.1	0.3	0.4	0.5	0.5	1.2	2.7	8.9	36.5
2003.....	1.3	*	*	0.1	0.2	0.4	0.4	0.6	1.2	3.0	9.3	38.0
2002.....	1.3	*	*	0.1	0.3	0.5	0.4	0.6	1.0	2.9	8.9	38.3
2001.....	1.3	*	*	0.1	0.3	0.5	0.4	0.6	0.9	2.9	9.9	40.0
2000.....	1.2	*	*	0.1	0.2	0.3	0.3	0.6	1.1	2.8	9.5	40.1
1999.....	1.3	*	*	0.1	0.3	0.4	0.5	0.6	0.9	2.7	10.1	43.4

All races 3/, female

2006.....	1.6	*	*	*	0.2	0.4	0.5	0.6	1.0	2.2	7.1	30.2
2005.....	1.9	*	*	*	0.2	0.4	0.4	0.7	1.0	2.7	8.9	38.0
2004.....	1.8	*	*	0.1	0.3	0.4	0.4	0.6	1.0	2.5	8.4	38.8
2003.....	1.9	*	*	0.1	0.3	0.4	0.4	0.6	0.8	2.3	8.5	41.9
2002.....	1.9	*	*	0.1	0.2	0.4	0.4	0.7	1.0	2.6	9.2	41.2
2001.....	2.0	*	*	0.1	0.3	0.3	0.5	0.6	1.0	2.5	9.6	42.9
2000.....	2.0	*	*	0.1	0.2	0.4	0.5	0.6	1.0	2.8	9.4	43.3
1999.....	1.9	*	*	0.2	0.3	0.4	0.4	0.6	1.0	2.6	9.2	44.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Anemias (D50-D64)

White, both sexes

2006 4/.....	1.2	*	*	*	0.1	0.1	0.1	0.3	0.7	2.0	7.3	29.6
2005 4/.....	1.5	*	*	*	0.1	0.1	0.1	0.3	0.8	2.3	8.8	36.6
2004 4/.....	1.5	*	*	0.1	0.1	0.1	0.2	0.3	0.7	2.4	8.5	37.7
2003 4/.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.6	2.4	8.6	40.6
2002.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.7	2.5	8.7	39.7
2001.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.4	41.7
2000.....	1.5	*	*	0.1	0.1	0.1	0.1	0.3	0.8	2.5	9.3	42.0
1999.....	1.5	*	*	0.1	0.1	*	0.1	0.3	0.6	2.4	9.4	43.0

White, male

2006 4/.....	1.0	*	*	*	*	*	0.1	0.2	0.7	1.9	7.8	29.9
2005 4/.....	1.1	*	*	*	*	*	*	0.3	0.8	2.3	9.0	33.7
2004 4/.....	1.1	*	*	*	*	*	0.2	0.3	0.8	2.5	8.6	36.7
2003 4/.....	1.1	*	*	*	*	0.1	0.1	0.3	0.7	2.6	9.1	37.9
2002.....	1.1	*	*	*	*	0.1	0.1	0.3	0.8	2.6	8.5	37.2
2001.....	1.1	*	*	*	*	0.1	0.1	0.2	0.7	2.6	9.8	40.4
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	9.2	39.1
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.5	2.5	9.9	43.4

White, female

2006 4/.....	1.5	*	*	*	*	*	0.1	0.3	0.8	2.0	7.0	29.4
2005 4/.....	1.8	*	*	*	*	*	0.1	0.4	0.8	2.3	8.6	37.9
2004 4/.....	1.8	*	*	*	*	*	0.1	0.2	0.7	2.2	8.4	38.2
2003 4/.....	1.8	*	*	*	*	*	0.1	0.2	0.6	2.2	8.2	41.8
2002.....	1.8	*	*	*	*	*	*	0.3	0.7	2.4	8.8	40.8
2001.....	1.9	*	*	*	*	*	0.2	0.3	0.8	2.4	9.1	42.3
2000.....	1.9	*	*	*	*	*	0.1	0.3	0.8	2.4	9.3	43.2
1999.....	1.9	*	*	*	0.1	*	0.1	0.3	0.7	2.3	9.0	42.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Anemias (D50-D64)

Black, both sexes

2006 4/.....	2.3	*	*	0.3	1.0	2.0	2.6	2.5	3.1	4.1	10.3	38.9
2005 4/.....	2.6	*	*	*	1.2	2.3	2.6	2.7	3.0	6.2	14.7	42.5
2004 4/.....	2.7	*	*	0.3	1.4	2.3	2.4	2.7	4.0	4.8	11.8	48.2
2003 4/.....	2.7	*	*	0.4	1.3	2.1	2.3	3.0	4.2	5.6	14.5	49.3
2002.....	2.8	*	*	0.4	1.3	2.4	2.5	3.1	3.6	5.9	16.0	52.3
2001.....	2.7	*	*	0.4	1.2	2.2	2.7	3.1	3.0	5.5	16.0	48.5
2000.....	2.6	*	*	0.5	1.1	2.3	2.4	3.0	3.4	5.4	13.3	51.0
1999.....	2.8	*	0.8	0.5	1.4	2.6	2.5	3.1	4.2	5.0	13.4	52.9

Black, male

2006 4/.....	2.1	*	*	*	1.1	2.0	2.4	2.3	3.5	4.0	11.9	29.5
2005 4/.....	2.4	*	*	*	1.4	2.6	2.9	2.5	2.6	6.4	17.2	37.2
2004 4/.....	2.6	*	*	*	1.5	2.7	2.7	2.5	4.9	5.0	13.9	42.7
2003 4/.....	2.7	*	*	*	1.3	2.2	2.6	3.3	5.4	7.9	14.9	49.0
2002.....	2.7	*	*	*	1.6	2.8	2.5	2.6	3.9	7.3	16.9	56.8
2001.....	2.5	*	*	*	1.4	2.9	2.4	3.6	3.1	6.7	15.0	42.0
2000.....	2.3	*	*	*	1.0	2.1	2.0	3.3	3.4	4.5	13.9	55.4
1999.....	2.7	*	*	*	1.7	2.8	2.7	3.3	5.1	5.4	12.9	44.5

Black, female

2006 4/.....	2.6	*	*	*	0.8	1.9	2.7	2.7	2.7	4.2	9.4	42.8
2005 4/.....	2.8	*	*	*	1.0	2.1	2.2	2.9	3.4	6.0	13.3	44.6
2004 4/.....	2.8	*	*	*	1.3	2.0	2.2	2.9	3.2	4.6	10.5	50.4
2003 4/.....	2.8	*	*	*	1.2	2.0	2.1	2.8	3.2	3.9	14.2	49.4
2002.....	3.0	*	*	*	0.9	2.0	2.5	3.6	3.3	5.0	15.4	50.5
2001.....	2.8	*	*	*	1.1	1.6	2.9	2.6	2.8	4.6	16.5	50.9
2000.....	3.0	*	*	*	1.1	2.4	2.8	2.8	3.4	6.0	13.0	49.4
1999.....	3.0	*	*	*	1.2	2.3	2.3	3.0	3.6	4.7	13.6	56.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Diabetes mellitus (E10-E14)

All races 3/, both sexes

2006.....	24.2	*	*	0.1	0.4	1.7	4.8	13.2	36.2	81.8	166.8	285.2
2005.....	25.3	*	*	0.1	0.5	1.5	4.7	13.4	37.2	86.8	177.2	312.1
2004.....	24.9	*	*	0.1	0.4	1.5	4.6	13.4	37.1	87.2	176.9	307.0
2003.....	25.5	*	*	0.1	0.4	1.6	4.6	13.9	38.5	90.8	181.1	317.5
2002.....	25.4	*	*	0.1	0.4	1.6	4.8	13.7	37.7	91.4	182.8	320.6
2001.....	25.1	*	*	0.1	0.4	1.5	4.3	13.6	37.8	91.4	181.4	321.8
2000.....	24.6	*	*	0.1	0.4	1.6	4.3	13.1	37.8	90.7	179.5	319.7
1999.....	24.5	*	*	0.1	0.4	1.4	4.3	12.9	38.3	91.8	178.0	317.2

All races 3/, male

2006.....	24.4	*	*	*	0.4	1.8	5.9	16.5	43.8	96.2	194.1	321.0
2005.....	25.0	*	*	*	0.5	1.8	5.8	16.6	44.9	99.7	202.7	344.4
2004.....	24.4	*	*	*	0.4	1.7	5.7	16.5	44.0	100.0	202.5	333.9
2003.....	24.8	*	*	*	0.5	1.9	5.7	16.7	45.0	104.0	207.2	340.7
2002.....	24.2	*	*	0.1	0.5	1.9	5.9	16.4	44.5	101.4	202.6	348.5
2001.....	23.5	*	*	0.1	0.4	1.7	5.3	16.4	43.1	101.7	198.9	343.1
2000.....	22.9	*	*	*	0.5	1.9	5.1	15.3	42.4	99.1	199.6	345.1
1999.....	22.8	*	*	*	0.4	1.6	5.1	15.6	42.6	99.8	198.2	345.7

All races 3/, female

2006.....	24.0	*	*	*	0.3	1.5	3.7	9.9	29.1	69.7	148.1	268.4
2005.....	25.7	*	*	0.1	0.4	1.2	3.5	10.3	30.1	75.9	159.9	297.2
2004.....	25.4	*	*	0.1	0.4	1.3	3.5	10.4	30.7	76.4	159.6	294.9
2003.....	26.2	*	*	*	0.3	1.4	3.5	11.2	32.4	79.8	163.6	307.2
2002.....	26.5	*	*	*	0.4	1.3	3.7	11.1	31.4	83.1	169.7	308.4
2001.....	26.6	*	*	*	0.3	1.3	3.4	11.0	32.9	82.8	169.9	312.9
2000.....	26.3	*	*	*	0.4	1.2	3.4	11.1	33.6	83.7	166.3	309.4
1999.....	26.2	*	*	*	0.4	1.3	3.5	10.4	34.3	85.2	164.9	305.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diabetes mellitus (E10-E14)

White, both sexes

2006 4/.....	23.6	*	*	*	0.3	1.3	4.1	11.4	31.7	73.3	154.8	270.4
2005 4/.....	24.9	*	*	0.1	0.4	1.2	4.0	11.6	32.8	77.7	165.5	296.3
2004 4/.....	24.4	*	*	*	0.3	1.3	4.0	11.6	32.4	78.1	164.2	289.4
2003 4/.....	25.0	*	*	0.1	0.3	1.4	4.0	11.8	33.5	81.4	168.4	300.8
2002.....	24.9	*	*	0.1	0.3	1.4	4.3	11.7	32.7	81.5	170.4	303.7
2001.....	24.6	*	*	0.1	0.3	1.2	3.8	11.7	32.7	81.9	168.9	306.0
2000.....	24.1	*	*	*	0.3	1.3	3.8	11.1	33.1	81.2	166.0	303.5
1999.....	23.9	*	*	0.1	0.4	1.3	3.7	10.9	32.4	81.8	165.1	298.9

White, male

2006 4/.....	24.2	*	*	*	0.3	1.4	5.0	14.6	39.0	88.4	185.4	312.0
2005 4/.....	24.9	*	*	*	0.4	1.4	5.1	14.6	40.1	91.3	193.7	336.2
2004 4/.....	24.3	*	*	*	0.3	1.4	5.0	14.7	39.2	91.6	193.3	323.0
2003 4/.....	24.8	*	*	*	0.4	1.5	5.1	14.7	40.2	95.8	198.6	333.4
2002.....	24.2	*	*	*	0.4	1.6	5.3	14.3	39.6	93.3	194.7	344.3
2001.....	23.5	*	*	*	0.3	1.3	4.7	14.3	38.4	94.6	189.2	337.3
2000.....	22.9	*	*	*	0.4	1.6	4.7	13.3	38.0	91.8	190.5	337.4
1999.....	22.7	*	*	*	0.4	1.4	4.5	13.5	37.3	92.3	189.9	333.4

White, female

2006 4/.....	23.0	*	*	*	0.3	1.2	3.2	8.2	24.7	60.2	133.5	250.8
2005 4/.....	24.9	*	*	*	0.3	0.9	3.0	8.6	25.8	66.0	146.1	278.0
2004 4/.....	24.5	*	*	*	0.3	1.1	2.9	8.6	25.9	66.5	144.4	274.3
2003 4/.....	25.2	*	*	*	0.3	1.2	2.9	9.0	27.2	69.2	148.0	286.3
2002.....	25.6	*	*	*	0.3	1.1	3.2	9.2	26.2	71.5	154.0	286.1
2001.....	25.7	*	*	*	0.2	1.1	2.9	9.2	27.4	71.2	155.5	293.0
2000.....	25.3	*	*	*	0.3	1.0	3.0	9.0	28.5	72.4	149.8	289.7
1999.....	25.0	*	*	*	0.3	1.2	3.0	8.4	27.8	73.1	148.9	285.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Diabetes mellitus (E10-E14)

Black, both sexes

2006 4/.....	32.4	*	*	*	0.8	4.2	10.2	27.3	77.6	166.8	312.0	471.1
2005 4/.....	33.2	*	*	*	1.1	3.9	9.5	27.8	79.7	177.1	321.7	506.5
2004 4/.....	33.2	*	*	*	0.9	3.3	9.5	28.1	80.5	179.3	332.8	532.4
2003 4/.....	33.8	*	*	*	0.7	3.8	9.5	30.1	85.7	184.9	336.1	527.5
2002.....	33.6	*	*	*	1.0	3.6	9.6	30.1	84.2	188.1	336.9	537.8
2001.....	33.1	*	*	*	1.0	3.7	8.8	29.6	85.0	188.4	335.2	529.7
2000.....	32.8	*	*	*	0.9	3.5	8.0	30.2	82.5	188.4	350.6	524.9
1999.....	33.0	*	*	*	0.8	3.0	8.9	30.3	90.2	190.8	333.5	529.0

Black, male

2006 4/.....	30.6	*	*	*	0.9	5.0	12.9	33.5	90.8	188.2	327.7	463.8
2005 4/.....	30.7	*	*	*	1.2	4.8	11.5	33.3	94.2	194.0	342.3	464.8
2004 4/.....	30.2	*	*	*	1.0	3.9	11.8	33.1	91.6	198.4	340.8	505.7
2003 4/.....	29.8	*	*	*	0.8	4.7	11.5	33.7	96.0	197.2	337.3	453.5
2002.....	29.0	*	*	*	1.2	4.4	11.7	34.8	94.7	190.1	329.9	417.0
2001.....	28.5	*	*	*	1.0	4.8	10.2	34.5	92.2	187.7	341.9	450.5
2000.....	27.4	*	*	*	1.0	3.8	9.4	33.4	88.2	184.9	337.0	452.1
1999.....	27.7	*	*	*	*	3.3	10.3	34.7	93.8	186.5	314.8	506.2

Black, female

2006 4/.....	34.1	*	*	*	0.7	3.5	7.8	22.0	66.9	151.6	303.2	474.1
2005 4/.....	35.5	*	*	*	1.0	3.1	7.6	23.1	68.0	165.0	310.0	523.3
2004 4/.....	36.0	*	*	*	0.9	2.7	7.6	23.8	71.6	165.7	328.2	543.0
2003 4/.....	37.4	*	*	*	*	3.0	7.8	27.0	77.4	176.1	335.4	556.2
2002.....	37.8	*	*	*	0.7	2.9	7.8	26.0	75.8	186.7	340.9	584.1
2001.....	37.2	*	*	*	1.0	2.8	7.6	25.4	79.1	188.9	331.5	559.1
2000.....	37.8	*	*	*	0.9	3.1	6.8	27.5	77.8	190.9	358.3	552.0
1999.....	37.8	*	*	*	0.9	2.6	7.6	26.6	87.4	193.9	343.9	537.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Nutritional deficiencies (E40-E64)

All races 3/, both sexes

2006.....	0.9	*	*	*	*	*	0.1	0.2	0.6	1.5	5.5	22.7
2005.....	1.1	*	*	*	*	*	0.1	0.3	0.5	1.9	7.0	30.8
2004.....	1.0	*	*	*	*	*	0.1	0.3	0.6	1.6	6.9	30.3
2003.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.9	7.7	35.5
2002.....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.1	8.9	41.5
2001.....	1.3	*	*	*	*	*	0.1	0.3	0.6	2.0	8.7	42.9
2000.....	1.5	*	*	*	*	*	0.1	0.3	0.7	2.5	10.4	50.0
1999.....	1.5	*	*	*	*	*	0.1	0.3	0.7	2.3	10.2	53.9

All races 3/, male

2006.....	0.6	*	*	*	*	*	0.1	0.2	0.5	1.5	5.6	18.7
2005.....	0.8	*	*	*	*	*	*	0.3	0.6	1.8	6.5	27.9
2004.....	0.8	*	*	*	*	*	*	0.2	0.6	1.7	6.8	27.4
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	2.0	7.8	30.6
2002.....	0.9	*	*	*	*	*	0.1	0.2	0.7	2.2	8.5	37.0
2001.....	0.9	*	*	*	*	*	0.1	0.3	0.7	2.1	8.5	35.2
2000.....	1.0	*	*	*	*	*	*	0.2	0.8	2.7	9.7	43.7
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	10.3	48.4

All races 3/, female

2006.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.4	5.4	24.5
2005.....	1.4	*	*	*	*	*	0.1	0.3	0.4	1.9	7.4	32.1
2004.....	1.3	*	*	*	*	*	0.1	0.3	0.5	1.5	7.0	31.7
2003.....	1.5	*	*	*	*	*	*	0.3	0.6	1.7	7.7	37.6
2002.....	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	9.1	43.5
2001.....	1.7	*	*	*	*	*	0.1	0.3	0.5	2.0	8.9	46.1
2000.....	2.0	*	*	*	*	*	0.2	0.3	0.6	2.4	10.9	52.6
1999.....	2.0	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	56.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Nutritional deficiencies (E40-E64)

White, both sexes

2006 4/.....	0.9	*	*	*	*	*	0.1	0.2	0.6	1.3	5.3	22.1
2005 4/.....	1.1	*	*	*	*	*	0.1	0.3	0.5	1.7	6.6	30.0
2004 4/.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.5	6.6	29.2
2003 4/.....	1.2	*	*	*	*	*	0.1	0.2	0.5	1.7	7.4	34.2
2002.....	1.4	*	*	*	*	*	0.1	0.2	0.5	1.9	8.6	40.2
2001.....	1.4	*	*	*	*	*	0.1	0.2	0.5	1.9	8.2	42.2
2000.....	1.6	*	*	*	*	*	0.1	0.2	0.6	2.3	10.0	48.8
1999.....	1.6	*	*	*	*	*	0.1	0.2	0.6	2.0	9.5	52.8

White, male

2006 4/.....	0.6	*	*	*	*	*	*	0.2	0.5	1.4	5.1	18.2
2005 4/.....	0.8	*	*	*	*	*	*	0.2	0.5	1.7	6.0	27.2
2004 4/.....	0.7	*	*	*	*	*	*	0.2	0.5	1.5	6.4	26.3
2003 4/.....	0.8	*	*	*	*	*	*	0.2	0.5	1.8	7.1	29.6
2002.....	1.0	*	*	*	*	*	0.1	0.2	0.5	2.1	8.1	36.3
2001.....	0.9	*	*	*	*	*	*	0.2	0.5	1.8	8.0	33.8
2000.....	1.0	*	*	*	*	*	*	0.2	0.6	2.3	9.1	42.8
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	9.3	46.5

White, female

2006 4/.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.3	5.4	24.0
2005 4/.....	1.4	*	*	*	*	*	*	0.3	0.4	1.8	7.0	31.4
2004 4/.....	1.4	*	*	*	*	*	*	0.3	0.5	1.4	6.8	30.4
2003 4/.....	1.6	*	*	*	*	*	*	0.2	0.6	1.7	7.5	36.3
2002.....	1.8	*	*	*	*	*	*	0.3	0.5	1.8	8.9	41.9
2001.....	1.8	*	*	*	*	*	*	0.3	0.6	2.0	8.3	45.7
2000.....	2.1	*	*	*	*	*	0.2	0.3	0.6	2.3	10.5	51.3
1999.....	2.1	*	*	*	*	*	0.2	0.2	0.5	1.9	9.6	55.4



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Nutritional deficiencies (E40-E64)

Black, both sexes

2006 4/.....	0.9	*	*	*	*	*	*	0.5	1.1	2.9	8.5	30.6
2005 4/.....	1.1	*	*	*	*	*	*	0.6	1.1	3.5	12.5	42.5
2004 4/.....	1.1	*	*	*	*	*	*	0.6	1.2	3.3	10.6	46.8
2003 4/.....	1.2	*	*	*	*	*	*	0.7	1.4	3.2	13.2	55.1
2002.....	1.3	*	*	*	*	*	*	0.4	1.7	4.2	14.5	60.6
2001.....	1.3	*	*	*	*	*	*	0.7	1.2	3.7	15.9	54.7
2000.....	1.5	*	*	*	*	*	*	0.6	1.7	4.7	16.1	66.9
1999.....	1.6	*	*	*	*	*	*	0.7	2.0	5.5	19.4	70.8

Black, male

2006 4/.....	0.8	*	*	*	*	*	*	*	1.4	4.0	11.9	28.6
2005 4/.....	0.8	*	*	*	*	*	*	*	1.7	3.3	12.3	37.2
2004 4/.....	0.9	*	*	*	*	*	*	*	1.6	4.4	12.2	45.7
2003 4/.....	1.0	*	*	*	*	*	*	*	1.9	4.4	17.8	48.0
2002.....	1.0	*	*	*	*	*	*	*	2.8	4.4	16.4	47.2
2001.....	1.2	*	*	*	*	*	*	1.2	1.9	5.5	16.5	57.9
2000.....	1.2	*	*	*	*	*	*	*	2.8	6.4	17.9	63.4
1999.....	1.5	*	*	*	*	*	*	*	2.8	8.1	22.8	75.0

Black, female

2006 4/.....	0.9	*	*	*	*	*	*	*	*	2.2	6.6	31.4
2005 4/.....	1.3	*	*	*	*	*	*	*	*	3.6	12.6	44.6
2004 4/.....	1.2	*	*	*	*	*	*	*	*	2.6	9.6	47.3
2003 4/.....	1.4	*	*	*	*	*	*	*	*	2.3	10.7	57.9
2002.....	1.6	*	*	*	*	*	*	*	*	4.1	13.5	65.7
2001.....	1.4	*	*	*	*	*	*	*	*	2.4	15.5	53.5
2000.....	1.7	*	*	*	*	*	*	*	*	3.5	15.2	68.3
1999.....	1.8	*	*	*	*	*	*	*	*	3.6	17.6	69.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malnutrition (E40-E46)

All races 3/, both sexes

2006.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.4	5.2	20.9
2005.....	1.0	*	*	*	*	*	0.1	0.3	0.5	1.8	6.6	28.9
2004.....	1.0	*	*	*	*	*	0.1	0.3	0.5	1.6	6.7	28.3
2003.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.7	7.4	33.4
2002.....	1.2	*	*	*	*	*	0.1	0.2	0.6	1.9	8.4	38.4
2001.....	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	8.2	40.2
2000.....	1.4	*	*	*	*	*	0.1	0.2	0.7	2.3	9.7	46.3
1999.....	1.4	*	*	*	*	*	0.1	0.3	0.6	2.2	9.5	50.0

All races 3/, male

2006.....	0.6	*	*	*	*	*	*	0.2	0.5	1.4	5.2	17.5
2005.....	0.7	*	*	*	*	*	*	0.3	0.6	1.7	6.2	26.3
2004.....	0.7	*	*	*	*	*	*	0.2	0.6	1.7	6.6	25.3
2003.....	0.8	*	*	*	*	*	*	0.2	0.6	1.9	7.4	28.2
2002.....	0.9	*	*	*	*	*	0.1	0.2	0.7	2.1	8.0	35.1
2001.....	0.8	*	*	*	*	*	*	0.2	0.6	1.9	8.0	33.6
2000.....	0.9	*	*	*	*	*	*	0.2	0.7	2.5	8.9	40.7
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.5	9.4	44.4

All races 3/, female

2006.....	1.0	*	*	*	*	*	0.1	0.2	0.6	1.4	5.1	22.4
2005.....	1.3	*	*	*	*	*	0.1	0.3	0.4	1.8	6.9	30.2
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.5	1.5	6.7	29.6
2003.....	1.4	*	*	*	*	*	*	0.2	0.6	1.7	7.4	35.6
2002.....	1.6	*	*	*	*	*	0.1	0.2	0.5	1.8	8.6	39.8
2001.....	1.6	*	*	*	*	*	0.1	0.3	0.5	1.9	8.2	43.0
2000.....	1.8	*	*	*	*	*	0.2	0.3	0.6	2.1	10.2	48.6
1999.....	1.8	*	*	*	*	*	0.1	0.3	0.5	2.0	9.5	52.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malnutrition (E40-E46)

White, both sexes

2006 4/.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.3	5.0	20.3
2005 4/.....	1.0	*	*	*	*	*	0.1	0.2	0.4	1.6	6.2	28.2
2004 4/.....	1.0	*	*	*	*	*	0.1	0.2	0.5	1.4	6.3	26.9
2003 4/.....	1.1	*	*	*	*	*	0.1	0.2	0.5	1.6	7.0	32.1
2002.....	1.3	*	*	*	*	*	0.1	0.2	0.5	1.8	8.0	37.0
2001.....	1.3	*	*	*	*	*	0.1	0.2	0.5	1.8	7.6	39.5
2000.....	1.5	*	*	*	*	*	0.1	0.2	0.6	2.1	9.3	45.2
1999.....	1.5	*	*	*	*	*	0.1	0.2	0.5	1.9	8.8	48.9

White, male

2006 4/.....	0.6	*	*	*	*	*	*	0.2	0.4	1.2	4.7	16.9
2005 4/.....	0.7	*	*	*	*	*	*	0.2	0.5	1.6	5.7	25.7
2004 4/.....	0.7	*	*	*	*	*	*	0.2	0.5	1.5	6.1	24.1
2003 4/.....	0.8	*	*	*	*	*	*	0.2	0.4	1.7	6.7	27.2
2002.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.9	7.5	34.3
2001.....	0.8	*	*	*	*	*	*	0.1	0.5	1.7	7.4	32.2
2000.....	1.0	*	*	*	*	*	*	0.2	0.6	2.2	8.4	39.7
1999.....	1.0	*	*	*	*	*	*	0.2	0.6	2.0	8.5	42.2

White, female

2006 4/.....	1.1	*	*	*	*	*	0.1	0.2	0.6	1.3	5.1	21.8
2005 4/.....	1.4	*	*	*	*	*	*	0.3	0.4	1.7	6.5	29.4
2004 4/.....	1.3	*	*	*	*	*	*	0.3	0.5	1.4	6.5	28.2
2003 4/.....	1.5	*	*	*	*	*	*	0.2	0.6	1.6	7.3	34.2
2002.....	1.6	*	*	*	*	*	*	0.3	0.5	1.7	8.4	38.2
2001.....	1.7	*	*	*	*	*	*	0.3	0.5	1.9	7.7	42.6
2000.....	1.9	*	*	*	*	*	0.1	0.2	0.6	2.0	9.9	47.4
1999.....	2.0	*	*	*	*	*	0.2	0.2	0.4	1.8	9.0	51.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Malnutrition (E40-E46)

Black, both sexes

2006 4/.....	0.8	*	*	*	*	*	*	0.5	1.0	2.9	7.9	29.3
2005 4/.....	1.1	*	*	*	*	*	*	0.5	1.1	3.4	12.0	40.4
2004 4/.....	1.0	*	*	*	*	*	*	0.6	1.1	3.2	10.4	46.0
2003 4/.....	1.2	*	*	*	*	*	*	0.7	1.4	3.0	13.0	53.4
2002.....	1.3	*	*	*	*	*	*	*	1.7	3.6	14.0	57.6
2001.....	1.2	*	*	*	*	*	*	0.6	1.1	3.4	14.8	51.6
2000.....	1.4	*	*	*	*	*	*	0.5	1.5	4.7	14.9	63.8
1999.....	1.5	*	*	*	*	*	*	0.6	1.8	5.4	17.9	66.0

Black, male

2006 4/.....	0.8	*	*	*	*	*	*	*	*	4.0	10.8	28.6
2005 4/.....	0.8	*	*	*	*	*	*	*	1.6	3.3	11.7	34.4
2004 4/.....	0.9	*	*	*	*	*	*	*	1.5	4.3	12.0	43.7
2003 4/.....	1.0	*	*	*	*	*	*	*	1.9	4.2	17.2	45.9
2002.....	1.0	*	*	*	*	*	*	*	2.6	3.9	15.2	46.1
2001.....	1.1	*	*	*	*	*	*	1.0	*	4.9	15.9	55.6
2000.....	1.2	*	*	*	*	*	*	*	2.4	6.3	17.3	61.1
1999.....	1.4	*	*	*	*	*	*	*	2.5	7.8	20.3	73.8

Black, female

2006 4/.....	0.9	*	*	*	*	*	*	*	*	2.2	6.3	29.6
2005 4/.....	1.3	*	*	*	*	*	*	*	*	3.5	12.1	42.8
2004 4/.....	1.2	*	*	*	*	*	*	*	*	2.5	9.4	46.9
2003 4/.....	1.3	*	*	*	*	*	*	*	*	2.2	10.7	56.3
2002.....	1.5	*	*	*	*	*	*	*	*	3.4	13.3	62.0
2001.....	1.3	*	*	*	*	*	*	*	*	2.2	14.2	50.1
2000.....	1.6	*	*	*	*	*	*	*	*	3.5	13.6	64.8
1999.....	1.6	*	*	*	*	*	*	*	*	3.6	16.5	63.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other nutritional deficiencies (E50-E64)

All races 3/, both sexes

2006.....	0.1	*	*	*	*	*	*	*	*	*	0.4	1.8
2005.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	1.8
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
2003.....	0.1	*	*	*	*	*	*	*	*	0.1	0.3	2.1
2002.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	3.1
2001.....	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.7
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.7
1999.....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9

All races 3/, male

2006.....	0.0	*	*	*	*	*	*	*	*	*	0.4	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	1.6
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	2.1
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.4
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.6	1.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.6
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.8	3.0
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.9	3.9

All races 3/, female

2006.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
2005.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.9
2004.....	0.1	*	*	*	*	*	*	*	*	*	*	2.1
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	2.0
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.6
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.1
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	4.0
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other nutritional deficiencies (E50-E64)

White, both sexes

2006 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.3	1.9
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.4	1.8
2004 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2003 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.1
2001.....	0.1	*	*	*	*	*	*	*	*	0.1	0.6	2.7
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.7	3.7
1999.....	0.1	*	*	*	*	*	*	*	*	0.1	0.7	3.9

White, male

2006 4/.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*	*	*	*	*	1.5
2004 4/.....	0.0	*	*	*	*	*	*	*	*	*	*	2.2
2003 4/.....	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.0
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.7	3.2
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.8	4.3

White, female

2006 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.9
2004 4/.....	0.1	*	*	*	*	*	*	*	*	*	*	2.2
2003 4/.....	0.1	*	*	*	*	*	*	*	*	*	*	2.1
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.5	3.6
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.1
2000.....	0.2	*	*	*	*	*	*	*	*	0.3	0.6	3.9
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.6	3.8



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Meningitis (G00,G03)

All races 3/, both sexes

2006.....	0.2	1.5	0.1	*	0.1	0.1	0.2	0.2	0.2	0.4	0.7	0.9
2005.....	0.2	1.4	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.4	0.6	0.9
2004.....	0.2	1.6	0.2	0.1	0.1	0.1	0.2	0.3	0.3	0.5	0.7	0.9
2003.....	0.3	1.9	0.2	*	0.1	0.1	0.2	0.3	0.4	0.5	0.7	1.1
2002.....	0.2	1.8	*	0.1	0.1	0.1	0.1	0.3	0.3	0.6	0.7	1.2
2001.....	0.3	1.9	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.5	0.8	0.9
2000.....	0.3	2.4	0.2	0.1	*	0.1	0.2	0.3	0.4	0.6	1.0	1.2
1999.....	0.3	3.1	0.3	0.1	0.1	0.1	0.2	0.3	0.4	0.6	1.0	1.5

All races 3/, male

2006.....	0.2	1.4	*	*	*	0.1	0.2	0.3	0.3	0.4	0.7	*
2005.....	0.2	1.4	0.3	*	*	0.1	0.2	0.3	0.4	0.4	0.6	*
2004.....	0.2	1.5	*	*	*	0.1	0.2	0.4	0.3	0.4	0.7	*
2003.....	0.3	1.9	*	*	0.1	0.1	0.2	0.4	0.5	0.5	0.8	*
2002.....	0.3	2.0	*	*	*	*	0.2	0.4	0.4	0.7	0.8	*
2001.....	0.3	1.9	*	*	0.1	0.1	0.2	0.3	0.5	0.6	0.9	*
2000.....	0.3	2.5	0.3	*	*	0.1	0.2	0.3	0.5	0.7	1.3	*
1999.....	0.3	3.1	0.3	*	0.1	0.1	0.2	0.4	0.4	0.7	1.0	1.7

All races 3/, female

2006.....	0.2	1.6	*	*	*	*	0.2	0.2	0.2	0.4	0.6	0.8
2005.....	0.2	1.4	*	*	*	*	0.1	0.2	0.3	0.4	0.6	0.9
2004.....	0.2	1.7	*	*	*	*	0.1	0.2	0.3	0.6	0.6	0.7
2003.....	0.2	2.0	*	*	*	*	0.1	0.2	0.3	0.5	0.6	1.0
2002.....	0.2	1.7	*	*	*	*	*	0.2	0.3	0.5	0.6	1.1
2001.....	0.2	2.0	*	*	*	*	0.1	0.2	0.4	0.4	0.6	0.8
2000.....	0.2	2.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.0
1999.....	0.3	3.0	*	*	*	0.1	0.1	0.2	0.4	0.5	1.0	1.4



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Meningitis (G00,G03)

White, both sexes

2006 4/.....	0.2	1.4	*	*	*	0.1	0.1	0.2	0.2	0.4	0.7	0.9
2005 4/.....	0.2	1.1	0.2	*	0.1	0.1	0.1	0.3	0.3	0.4	0.5	0.9
2004 4/.....	0.2	1.4	*	*	*	*	0.1	0.2	0.3	0.4	0.6	0.8
2003 4/.....	0.2	1.7	*	*	0.1	0.1	0.1	0.2	0.3	0.5	0.7	1.0
2002.....	0.2	1.7	*	0.1	*	0.1	0.1	0.2	0.3	0.6	0.6	1.2
2001.....	0.2	1.5	*	*	0.1	0.1	0.1	0.2	0.4	0.5	0.7	0.8
2000.....	0.2	1.8	0.2	0.1	*	0.1	0.1	0.2	0.3	0.5	0.9	1.0
1999.....	0.3	2.3	0.3	*	0.1	0.1	0.1	0.2	0.3	0.6	0.9	1.3

White, male

2006 4/.....	0.2	1.2	*	*	*	*	0.1	0.3	0.2	0.4	0.8	*
2005 4/.....	0.2	*	*	*	*	*	0.2	0.3	0.3	0.4	0.6	*
2004 4/.....	0.2	1.5	*	*	*	*	0.2	0.3	0.3	0.3	0.7	*
2003 4/.....	0.3	2.0	*	*	*	*	0.1	0.3	0.4	0.5	0.8	*
2002.....	0.2	1.8	*	*	*	*	0.1	0.3	0.3	0.7	0.8	*
2001.....	0.2	1.4	*	*	*	*	0.1	0.3	0.4	0.6	0.9	*
2000.....	0.3	2.0	*	*	*	*	0.1	0.3	0.4	0.7	1.2	*
1999.....	0.3	3.0	0.3	*	*	*	0.1	0.3	0.4	0.6	0.9	*

White, female

2006 4/.....	0.2	1.5	*	*	*	*	0.1	0.2	0.2	0.4	0.6	0.8
2005 4/.....	0.2	*	*	*	*	*	*	0.2	0.3	0.3	0.5	0.9
2004 4/.....	0.2	1.4	*	*	*	*	*	0.1	0.3	0.5	0.5	*
2003 4/.....	0.2	1.3	*	*	*	*	*	0.2	0.2	0.5	0.6	0.9
2002.....	0.2	1.5	*	*	*	*	*	0.2	0.2	0.4	0.5	1.2
2001.....	0.2	1.6	*	*	*	*	*	0.2	0.3	0.4	0.6	0.8
2000.....	0.2	1.6	*	*	*	*	*	0.2	0.2	0.4	0.8	1.0
1999.....	0.3	1.5	*	*	*	0.1	0.1	0.2	0.3	0.5	0.9	1.3



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Parkinson's disease (G20-G21)

All races 3/, both sexes

2006.....	6.5	*	*	*	*	*	*	0.2	1.3	12.2	69.8	144.8
2005.....	6.6	*	*	*	*	*	*	0.2	1.4	13.0	71.2	143.7
2004.....	6.1	*	*	*	*	*	*	0.2	1.2	12.0	67.5	135.8
2003.....	6.2	*	*	*	*	*	*	0.2	1.3	12.7	67.8	138.2
2002.....	5.9	*	*	*	*	*	*	0.1	1.2	12.2	63.9	135.2
2001.....	5.8	*	*	*	*	*	*	0.1	1.2	11.7	64.6	134.2
2000.....	5.6	*	*	*	*	*	*	0.1	1.1	11.5	61.9	131.9
1999.....	5.2	*	*	*	*	*	*	0.1	1.0	11.0	58.2	124.4

All races 3/, male

2006.....	7.7	*	*	*	*	*	*	0.2	1.7	17.6	104.7	231.7
2005.....	7.7	*	*	*	*	*	*	0.2	1.9	18.7	107.2	228.6
2004.....	7.2	*	*	*	*	*	*	0.2	1.6	17.3	101.7	220.0
2003.....	7.1	*	*	*	*	*	*	0.2	1.6	18.2	100.7	222.4
2002.....	6.8	*	*	*	*	*	*	0.2	1.5	17.9	94.7	220.1
2001.....	6.7	*	*	*	*	*	*	0.2	1.8	16.8	97.0	224.4
2000.....	6.4	*	*	*	*	*	*	0.2	1.5	16.1	92.8	222.3
1999.....	6.0	*	*	*	*	*	*	0.2	1.4	15.7	89.5	205.6

All races 3/, female

2006.....	5.4	*	*	*	*	*	*	0.1	0.8	7.7	45.9	104.1
2005.....	5.5	*	*	*	*	*	*	0.1	1.0	8.1	46.8	104.7
2004.....	5.1	*	*	*	*	*	*	*	0.8	7.5	44.4	97.9
2003.....	5.3	*	*	*	*	*	*	0.1	0.9	8.1	45.8	101.0
2002.....	5.0	*	*	*	*	*	*	*	0.9	7.4	43.5	98.4
2001.....	4.9	*	*	*	*	*	*	*	0.7	7.5	43.3	96.4
2000.....	4.8	*	*	*	*	*	*	0.1	0.8	7.6	41.7	95.0
1999.....	4.5	*	*	*	*	*	*	*	0.7	7.1	37.8	91.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Parkinson's disease (G20-G21)

White, both sexes

2006 4/.....	7.6	*	*	*	*	*	*	0.1	1.3	13.0	74.7	154.1
2005 4/.....	7.7	*	*	*	*	*	*	0.2	1.5	13.8	75.8	152.9
2004 4/.....	7.2	*	*	*	*	*	*	0.2	1.3	13.0	71.9	143.1
2003 4/.....	7.2	*	*	*	*	*	*	0.2	1.3	13.6	72.2	146.0
2002.....	6.9	*	*	*	*	*	*	0.2	1.3	13.0	68.2	143.3
2001.....	6.8	*	*	*	*	*	*	0.1	1.2	12.6	68.8	141.8
2000.....	6.5	*	*	*	*	*	*	0.1	1.2	12.3	65.6	139.5
1999.....	6.1	*	*	*	*	*	*	0.1	1.1	11.7	61.5	131.2

White, male

2006 4/.....	8.9	*	*	*	*	*	*	0.2	1.8	18.6	111.0	247.3
2005 4/.....	9.0	*	*	*	*	*	*	0.2	2.0	19.8	113.5	242.7
2004 4/.....	8.3	*	*	*	*	*	*	0.2	1.7	18.6	107.9	231.3
2003 4/.....	8.2	*	*	*	*	*	*	0.2	1.6	19.3	106.6	233.3
2002.....	7.9	*	*	*	*	*	*	0.2	1.6	19.0	100.5	232.7
2001.....	7.8	*	*	*	*	*	*	0.2	1.9	17.9	103.0	237.6
2000.....	7.4	*	*	*	*	*	*	0.2	1.6	17.2	97.5	235.3
1999.....	7.0	*	*	*	*	*	*	0.2	1.5	16.8	94.4	216.7

White, female

2006 4/.....	6.4	*	*	*	*	*	*	*	0.9	8.3	49.5	110.3
2005 4/.....	6.5	*	*	*	*	*	*	0.1	1.1	8.7	49.8	111.5
2004 4/.....	6.0	*	*	*	*	*	*	*	0.9	8.2	47.4	103.3
2003 4/.....	6.2	*	*	*	*	*	*	0.2	1.0	8.7	48.8	107.4
2002.....	5.9	*	*	*	*	*	*	*	0.9	7.9	46.5	104.4
2001.....	5.8	*	*	*	*	*	*	*	0.7	8.1	46.2	101.7
2000.....	5.6	*	*	*	*	*	*	*	0.9	8.2	44.7	100.6
1999.....	5.2	*	*	*	*	*	*	*	0.6	7.5	40.0	96.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Parkinson's disease (G20-G21)

Black, both sexes

2006 4/.....	1.7	*	*	*	*	*	*	*	1.0	7.4	30.0	51.4
2005 4/.....	1.8	*	*	*	*	*	*	*	0.8	7.2	33.8	56.7
2004 4/.....	1.6	*	*	*	*	*	*	*	0.7	5.4	29.2	61.2
2003 4/.....	1.6	*	*	*	*	*	*	*	1.2	7.0	29.0	52.8
2002.....	1.5	*	*	*	*	*	*	*	*	6.8	25.4	51.7
2001.....	1.5	*	*	*	*	*	*	*	0.8	5.5	25.2	59.0
2000.....	1.5	*	*	*	*	*	*	*	*	5.7	27.4	51.6
1999.....	1.4	*	*	*	*	*	*	*	0.9	5.9	24.4	52.0

Black, male

2006 4/.....	2.0	*	*	*	*	*	*	*	*	11.7	50.6	70.6
2005 4/.....	2.2	*	*	*	*	*	*	*	*	12.2	54.1	97.6
2004 4/.....	1.9	*	*	*	*	*	*	*	*	8.5	43.4	106.3
2003 4/.....	1.9	*	*	*	*	*	*	*	1.7	10.3	45.4	97.0
2002.....	1.7	*	*	*	*	*	*	*	*	11.1	39.6	89.0
2001.....	1.7	*	*	*	*	*	*	*	*	8.5	37.4	99.8
2000.....	1.8	*	*	*	*	*	*	*	*	8.8	47.8	86.5
1999.....	1.6	*	*	*	*	*	*	*	*	8.7	37.8	86.7

Black, female

2006 4/.....	1.5	*	*	*	*	*	*	*	*	4.3	18.3	43.5
2005 4/.....	1.5	*	*	*	*	*	*	*	*	3.6	22.3	40.1
2004 4/.....	1.4	*	*	*	*	*	*	*	*	3.2	21.1	43.3
2003 4/.....	1.4	*	*	*	*	*	*	*	*	4.7	19.8	35.6
2002.....	1.2	*	*	*	*	*	*	*	*	3.6	17.4	37.4
2001.....	1.3	*	*	*	*	*	*	*	*	3.3	18.3	43.8
2000.....	1.2	*	*	*	*	*	*	*	*	3.5	15.9	38.6
1999.....	1.2	*	*	*	*	*	*	*	*	3.9	16.9	39.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Alzheimer's disease (G30)

All races 3/, both sexes

2006.....	24.2	*	*	*	*	*	*	0.2	2.1	20.2	175.6	848.3
2005.....	24.2	*	*	*	*	*	*	0.2	2.1	20.5	177.3	861.6
2004.....	22.5	*	*	*	*	*	*	0.2	1.9	19.7	168.7	818.8
2003.....	21.8	*	*	*	*	*	*	0.2	2.0	20.9	164.4	802.4
2002.....	20.4	*	*	*	*	*	*	0.1	1.9	19.7	158.1	752.3
2001.....	18.9	*	*	*	*	*	*	0.2	2.1	18.7	147.5	710.3
2000.....	17.6	*	*	*	*	*	*	0.2	2.0	18.7	139.6	667.7
1999.....	16.0	*	*	*	*	*	*	0.2	1.9	17.4	129.5	601.3

All races 3/, male

2006.....	14.3	*	*	*	*	*	*	0.2	2.0	19.7	156.6	638.6
2005.....	14.1	*	*	*	*	*	*	0.1	2.0	19.3	157.7	639.9
2004.....	13.1	*	*	*	*	*	*	0.2	1.8	18.5	149.0	619.6
2003.....	12.8	*	*	*	*	*	*	0.2	1.8	20.2	147.6	606.5
2002.....	12.0	*	*	*	*	*	*	0.1	1.8	18.6	142.2	572.0
2001.....	11.3	*	*	*	*	*	*	0.2	1.9	17.7	136.3	556.1
2000.....	10.5	*	*	*	*	*	*	0.2	1.8	18.7	127.6	522.1
1999.....	9.8	*	*	*	*	*	*	0.2	1.9	17.3	122.6	485.7

All races 3/, female

2006.....	33.8	*	*	*	*	*	*	0.2	2.1	20.5	188.7	946.4
2005.....	33.9	*	*	*	*	*	*	0.2	2.3	21.4	190.5	963.4
2004.....	31.5	*	*	*	*	*	*	0.3	1.9	20.8	181.9	908.4
2003.....	30.5	*	*	*	*	*	*	0.2	2.1	21.5	175.7	889.0
2002.....	28.5	*	*	*	*	*	*	0.1	2.0	20.6	168.7	830.5
2001.....	26.3	*	*	*	*	*	*	0.2	2.3	19.5	154.8	774.9
2000.....	24.5	*	*	*	*	*	*	0.2	2.2	18.6	147.4	727.0
1999.....	21.9	*	*	*	*	*	*	0.2	1.8	17.5	134.0	648.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Alzheimer's disease (G30)

White, both sexes

2006 4/.....	27.7	*	*	*	*	*	*	0.3	2.2	21.1	182.7	878.9
2005 4/.....	27.6	*	*	*	*	*	*	0.2	2.3	21.1	183.3	891.6
2004 4/.....	25.6	*	*	*	*	*	*	0.3	2.0	20.5	174.1	846.8
2003 4/.....	25.0	*	*	*	*	*	*	0.2	2.2	21.7	170.7	834.0
2002.....	23.5	*	*	*	*	*	*	0.1	2.1	20.7	164.8	781.8
2001.....	21.7	*	*	*	*	*	*	0.2	2.3	19.6	152.6	738.1
2000.....	20.2	*	*	*	*	*	*	0.2	2.2	19.5	144.8	695.3
1999.....	18.3	*	*	*	*	*	*	0.2	2.0	18.1	134.5	627.9

White, male

2006 4/.....	16.4	*	*	*	*	*	*	0.3	2.1	20.5	162.5	662.5
2005 4/.....	16.0	*	*	*	*	*	*	0.1	2.1	19.7	162.0	661.6
2004 4/.....	14.9	*	*	*	*	*	*	0.2	1.9	19.0	153.7	640.9
2003 4/.....	14.6	*	*	*	*	*	*	0.2	1.9	20.8	152.4	630.3
2002.....	13.7	*	*	*	*	*	*	0.1	1.9	19.3	147.3	594.8
2001.....	12.8	*	*	*	*	*	*	0.2	2.0	18.5	140.5	577.2
2000.....	11.9	*	*	*	*	*	*	0.2	1.9	19.5	132.5	542.1
1999.....	11.2	*	*	*	*	*	*	0.2	2.1	17.9	127.4	506.9

White, female

2006 4/.....	38.8	*	*	*	*	*	*	0.2	2.3	21.6	196.7	980.5
2005 4/.....	38.9	*	*	*	*	*	*	0.3	2.5	22.2	197.9	997.5
2004 4/.....	36.1	*	*	*	*	*	*	0.3	2.1	21.7	188.0	939.6
2003 4/.....	35.2	*	*	*	*	*	*	0.2	2.4	22.5	183.1	924.3
2002.....	33.0	*	*	*	*	*	*	0.1	2.2	21.8	176.5	863.0
2001.....	30.3	*	*	*	*	*	*	0.2	2.6	20.6	160.6	805.4
2000.....	28.2	*	*	*	*	*	*	0.2	2.4	19.5	152.9	757.6
1999.....	25.3	*	*	*	*	*	*	0.2	2.0	18.3	139.1	676.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alzheimer's disease (G30)

Black, both sexes

2006 4/.....	11.3	*	*	*	*	*	*	*	1.6	18.6	146.7	663.9
2005 4/.....	11.8	*	*	*	*	*	*	*	1.5	22.1	157.9	689.5
2004 4/.....	11.0	*	*	*	*	*	*	*	1.3	19.3	153.1	657.9
2003 4/.....	9.7	*	*	*	*	*	*	*	1.0	19.2	138.1	583.8
2002.....	8.9	*	*	*	*	*	*	*	1.2	16.8	123.1	543.7
2001.....	8.4	*	*	*	*	*	*	*	1.1	15.2	124.7	506.7
2000.....	7.5	*	*	*	*	*	*	*	1.2	15.8	109.9	445.1
1999.....	6.5	*	*	*	*	*	*	*	1.2	15.1	98.2	372.4

Black, male

2006 4/.....	6.3	*	*	*	*	*	*	*	1.7	19.0	132.9	470.9
2005 4/.....	6.9	*	*	*	*	*	*	*	*	22.1	150.0	518.7
2004 4/.....	6.2	*	*	*	*	*	*	*	*	19.1	136.7	497.7
2003 4/.....	6.0	*	*	*	*	*	*	*	*	20.3	136.2	459.8
2002.....	5.3	*	*	*	*	*	*	*	*	18.0	122.5	412.8
2001.....	5.1	*	*	*	*	*	*	*	*	16.1	121.5	414.1
2000.....	4.5	*	*	*	*	*	*	*	*	16.5	97.5	371.3
1999.....	4.0	*	*	*	*	*	*	*	*	16.5	89.8	317.5

Black, female

2006 4/.....	15.8	*	*	*	*	*	*	*	1.6	18.4	154.5	742.8
2005 4/.....	16.3	*	*	*	*	*	*	*	1.7	22.1	162.4	758.5
2004 4/.....	15.3	*	*	*	*	*	*	*	1.4	19.4	162.4	721.4
2003 4/.....	13.2	*	*	*	*	*	*	*	*	18.4	139.2	631.9
2002.....	12.0	*	*	*	*	*	*	*	*	15.9	123.4	593.9
2001.....	11.3	*	*	*	*	*	*	*	*	14.6	126.4	541.0
2000.....	10.2	*	*	*	*	*	*	*	*	15.2	116.9	472.6
1999.....	8.7	*	*	*	*	*	*	*	*	14.2	102.9	392.9



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

All races 3/, both sexes

2006.....	275.1	12.3	1.3	0.9	3.2	10.1	35.3	108.7	255.5	629.6	1,846.2	5,928.2
2005.....	288.8	12.2	1.3	0.9	3.3	10.1	36.1	110.5	263.2	667.2	1,963.8	6,398.5
2004.....	293.3	13.8	1.5	0.8	3.2	10.0	36.6	110.9	268.2	697.7	2,035.6	6,616.6
2003.....	310.3	13.9	1.5	0.8	3.5	10.3	38.2	113.4	284.9	748.3	2,170.8	7,138.0
2002.....	318.6	15.9	1.4	0.9	3.2	10.0	37.8	114.3	294.8	788.4	2,257.0	7,389.6
2001.....	323.9	14.9	1.9	0.9	3.2	10.2	37.1	113.4	301.2	810.4	2,322.2	7,673.7
2000.....	332.9	16.7	1.5	0.9	3.3	9.5	36.8	115.7	319.2	849.3	2,396.2	8,030.3
1999.....	340.6	17.0	1.5	0.9	3.4	9.6	37.7	116.2	327.3	887.3	2,474.9	8,197.1

All races 3/, male

2006.....	267.7	13.4	1.4	1.0	4.0	13.4	47.5	153.3	355.1	821.3	2,235.3	6,151.8
2005.....	277.9	13.4	1.5	1.0	4.2	13.0	48.6	155.8	365.7	864.5	2,368.4	6,615.9
2004.....	281.2	14.7	1.5	0.8	4.0	12.8	49.0	156.7	371.5	903.9	2,462.3	6,818.1
2003.....	296.3	15.2	1.4	1.0	4.2	12.9	51.1	160.9	393.3	976.3	2,636.1	7,329.3
2002.....	303.3	16.6	1.5	0.9	4.1	12.7	50.8	162.3	407.0	1,027.8	2,731.1	7,638.8
2001.....	306.2	15.3	1.9	0.9	3.9	12.8	49.9	160.4	414.0	1,057.6	2,823.3	7,936.0
2000.....	315.8	17.5	1.6	1.0	3.9	11.8	49.5	165.1	441.6	1,114.1	2,926.6	8,448.1
1999.....	323.0	17.9	1.6	1.0	4.0	12.1	51.2	165.8	452.0	1,168.4	3,024.0	8,603.2

All races 3/, female

2006.....	282.4	11.2	1.3	0.8	2.4	6.7	23.1	65.5	162.9	467.4	1,580.1	5,823.6
2005.....	299.3	11.0	1.1	0.8	2.3	7.0	23.7	66.6	167.9	500.8	1,689.1	6,298.7
2004.....	304.9	12.9	1.5	0.8	2.3	7.1	24.2	66.6	172.2	524.5	1,748.4	6,526.0
2003.....	323.9	12.5	1.5	0.7	2.8	7.5	25.3	67.5	184.4	557.8	1,860.0	7,053.5
2002.....	333.3	15.1	1.3	0.8	2.2	7.2	24.9	68.0	191.0	589.1	1,942.2	7,281.5
2001.....	340.9	14.5	1.8	0.9	2.4	7.6	24.3	68.1	196.9	605.6	1,992.9	7,563.9
2000.....	349.5	15.9	1.4	0.8	2.8	7.2	24.3	68.0	206.3	631.2	2,050.4	7,860.1
1999.....	357.5	16.0	1.4	0.8	2.7	7.2	24.4	68.5	212.5	657.0	2,118.2	8,033.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

White, both sexes

2006 4/.....	292.2	9.8	1.2	0.8	2.9	8.3	30.7	95.9	233.4	596.8	1,829.1	6,029.7
2005 4/.....	307.0	9.9	1.1	0.8	2.8	8.2	31.4	96.8	240.0	635.4	1,945.0	6,507.8
2004 4/.....	311.7	11.9	1.2	0.7	2.7	8.2	32.1	97.1	244.7	665.2	2,016.2	6,704.8
2003 4/.....	330.0	11.0	1.2	0.7	2.9	8.5	32.9	99.0	259.8	712.2	2,151.4	7,233.1
2002.....	339.4	13.4	1.3	0.8	2.7	8.1	33.0	100.6	269.9	755.1	2,234.8	7,482.7
2001.....	344.8	12.2	1.5	0.8	2.7	8.5	31.7	99.5	275.3	774.1	2,298.9	7,770.6
2000.....	354.7	14.6	1.3	0.8	2.8	7.9	31.6	102.0	292.7	814.2	2,370.8	8,133.9
1999.....	362.7	14.6	1.3	0.8	2.9	8.1	32.1	102.7	300.0	850.9	2,449.9	8,299.2

White, male

2006 4/.....	281.5	10.5	1.2	1.0	3.7	11.3	42.5	138.4	327.4	782.6	2,223.9	6,287.1
2005 4/.....	293.0	10.8	1.3	0.9	3.6	10.8	43.4	140.1	337.7	829.3	2,357.8	6,772.6
2004 4/.....	296.7	12.7	1.3	0.7	3.5	10.7	44.4	141.3	343.3	870.1	2,452.6	6,940.8
2003 4/.....	312.6	11.5	1.1	0.9	3.5	10.9	45.6	144.6	364.8	938.3	2,623.0	7,466.8
2002.....	320.9	14.3	1.5	0.8	3.5	10.5	45.7	147.1	378.3	994.8	2,721.2	7,766.6
2001.....	323.5	13.2	1.6	0.8	3.3	10.9	44.2	145.1	385.3	1,020.8	2,809.7	8,060.7
2000.....	334.2	15.3	1.6	0.9	3.3	10.1	44.1	150.2	411.8	1,080.7	2,912.4	8,585.0
1999.....	341.2	15.3	1.5	0.9	3.4	10.4	45.1	150.5	420.7	1,133.1	3,012.8	8,717.5

White, female

2006 4/.....	302.7	9.1	1.2	0.6	2.0	5.2	18.7	53.9	144.1	436.5	1,555.0	5,908.9
2005 4/.....	320.8	9.0	0.9	0.7	2.0	5.4	19.1	54.0	147.4	468.7	1,660.6	6,385.8
2004 4/.....	326.5	11.0	1.1	0.7	1.9	5.5	19.5	53.6	151.4	489.8	1,718.6	6,598.4
2003 4/.....	347.0	10.4	1.4	0.6	2.3	5.9	20.0	53.9	160.6	519.6	1,832.5	7,129.6
2002.....	357.5	12.5	1.1	0.7	1.8	5.6	20.0	54.7	167.7	551.7	1,908.4	7,359.3
2001.....	365.5	11.1	1.5	0.8	2.0	6.0	19.0	54.6	171.9	565.8	1,959.8	7,649.2
2000.....	374.8	13.9	1.1	0.6	2.2	5.5	19.0	54.6	180.9	590.8	2,014.4	7,950.6
1999.....	383.6	13.8	1.2	0.7	2.4	5.7	19.0	55.7	186.8	615.7	2,081.0	8,131.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Major cardiovascular diseases (I00-I78)

Black, both sexes

2006 4/.....	246.2	25.0	2.2	1.4	5.6	22.8	72.0	220.1	491.7	1,032.8	2,342.0	5,587.2
2005 4/.....	256.2	23.1	2.3	1.6	6.2	22.9	73.8	228.0	511.8	1,066.9	2,500.7	6,016.9
2004 4/.....	260.3	21.9	2.9	1.5	5.9	22.7	74.3	231.0	525.5	1,122.3	2,572.7	6,368.6
2003 4/.....	273.1	26.2	2.8	1.4	6.9	22.9	78.8	238.8	556.2	1,207.1	2,713.5	6,862.0
2002.....	276.1	27.6	2.0	1.4	5.9	23.3	76.0	237.4	568.3	1,226.2	2,824.0	7,081.7
2001.....	280.4	28.6	3.5	1.3	6.6	22.6	78.3	238.2	578.0	1,276.2	2,891.8	7,316.7
2000.....	285.0	27.8	2.5	1.4	6.8	21.5	77.4	241.3	600.3	1,316.2	2,968.8	7,518.9
1999.....	289.4	29.2	2.4	1.4	6.8	20.7	81.6	241.9	614.6	1,368.1	3,024.7	7,582.5

Black, male

2006 4/.....	250.3	28.2	2.6	1.3	6.4	29.5	91.3	295.7	679.8	1,366.5	2,820.4	5,418.9
2005 4/.....	255.0	26.3	2.2	1.6	8.0	29.0	93.9	304.3	701.6	1,375.8	2,958.2	5,707.6
2004 4/.....	257.2	23.7	2.8	1.4	7.6	27.7	92.6	305.7	717.0	1,432.4	3,033.4	6,198.0
2003 4/.....	268.6	31.2	3.3	1.5	8.5	28.0	98.1	317.1	742.4	1,542.1	3,265.7	6,547.7
2002.....	268.5	28.2	1.9	1.5	7.2	28.9	94.7	313.1	762.5	1,556.1	3,287.6	6,929.0
2001.....	272.4	28.1	3.6	1.2	8.0	27.9	97.4	312.5	761.1	1,620.3	3,435.3	7,363.9
2000.....	275.6	29.1	2.1	1.4	7.5	25.4	95.8	317.5	796.8	1,639.9	3,501.8	7,687.4
1999.....	282.5	30.6	2.5	1.5	8.4	25.4	102.7	323.2	822.6	1,723.6	3,526.5	7,921.1

Black, female

2006 4/.....	242.5	21.7	1.7	1.5	4.7	16.5	54.9	155.1	340.2	794.1	2,069.9	5,656.0
2005 4/.....	257.2	19.7	2.3	1.6	4.3	17.4	55.8	162.4	358.8	845.8	2,240.8	6,141.8
2004 4/.....	263.2	20.1	3.0	1.5	4.1	18.1	58.0	166.9	371.5	901.1	2,312.3	6,436.3
2003 4/.....	277.2	21.0	2.4	1.3	5.3	18.1	61.7	171.5	406.5	968.7	2,401.2	6,984.1
2002.....	283.0	26.9	2.1	1.3	4.5	18.1	59.5	172.5	412.2	992.0	2,561.6	7,140.2
2001.....	287.8	29.0	3.4	1.4	5.1	17.7	61.4	174.3	430.5	1,033.4	2,586.6	7,299.2
2000.....	293.4	26.5	2.9	1.4	6.1	17.9	61.1	175.6	441.9	1,088.3	2,668.7	7,456.2
1999.....	295.7	27.9	2.4	1.2	5.1	16.4	63.0	172.3	447.7	1,117.3	2,743.8	7,456.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races 3/, both sexes

2006.....	211.0	8.4	1.0	0.6	2.5	8.2	28.3	88.0	207.3	490.3	1,383.1	4,480.8
2005.....	220.0	8.7	0.9	0.6	2.7	8.1	28.9	89.7	214.8	518.9	1,460.8	4,778.4
2004.....	222.2	10.3	1.2	0.6	2.5	7.9	29.3	90.2	218.8	541.6	1,506.3	4,895.9
2003.....	235.6	11.0	1.2	0.6	2.7	8.2	30.7	92.5	233.2	585.0	1,611.1	5,278.4
2002.....	241.7	12.4	1.1	0.6	2.5	7.9	30.5	93.7	241.5	615.9	1,677.2	5,446.8
2001.....	245.8	11.9	1.5	0.7	2.5	8.0	29.6	92.9	246.9	635.1	1,725.7	5,664.2
2000.....	252.6	13.0	1.2	0.7	2.6	7.4	29.2	94.2	261.2	665.6	1,780.3	5,926.1
1999.....	259.9	13.8	1.2	0.7	2.8	7.6	30.2	95.7	269.9	701.7	1,849.9	6,063.0

All races 3/, male

2006.....	214.0	8.8	1.1	0.7	3.3	11.2	39.5	128.9	296.8	660.5	1,743.5	4,819.9
2005.....	221.1	9.4	1.0	0.6	3.6	10.8	40.7	131.5	306.9	692.3	1,829.4	5,143.4
2004.....	222.8	10.9	1.1	0.6	3.2	10.5	40.9	132.3	312.8	723.8	1,893.6	5,239.3
2003.....	235.0	12.1	1.1	0.7	3.4	10.5	42.8	136.2	331.7	785.3	2,030.3	5,621.5
2002.....	240.7	12.9	1.1	0.7	3.3	10.5	43.1	138.4	343.4	827.1	2,110.1	5,823.5
2001.....	242.5	11.8	1.5	0.7	3.2	10.3	41.7	136.6	349.8	851.3	2,177.3	6,040.5
2000.....	249.8	13.3	1.4	0.8	3.2	9.6	41.4	140.2	371.7	898.3	2,248.1	6,430.0
1999.....	257.0	13.9	1.3	0.8	3.4	9.8	43.0	142.0	383.0	950.6	2,336.4	6,561.6

All races 3/, female

2006.....	208.0	7.9	0.9	0.6	1.8	5.1	17.0	48.5	124.1	346.3	1,136.7	4,322.1
2005.....	218.9	8.0	0.9	0.6	1.7	5.3	17.1	49.2	129.1	372.7	1,210.5	4,610.8
2004.....	221.6	9.7	1.2	0.6	1.7	5.2	17.7	49.6	131.5	388.6	1,245.6	4,741.5
2003.....	236.2	9.8	1.3	0.5	2.1	5.7	18.6	50.2	141.9	417.5	1,331.1	5,126.7
2002.....	242.7	11.8	1.0	0.6	1.7	5.2	18.0	50.6	147.2	440.1	1,389.7	5,283.3
2001.....	249.0	12.0	1.4	0.7	1.8	5.6	17.6	50.7	151.8	455.9	1,428.9	5,506.8
2000.....	255.3	12.5	1.0	0.5	2.1	5.2	17.2	49.8	159.3	474.0	1,475.1	5,720.9
1999.....	262.6	13.7	1.1	0.6	2.1	5.4	17.5	51.0	165.6	497.9	1,533.9	5,862.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

White, both sexes

2006 4/.....	225.5	6.6	0.9	0.5	2.2	6.8	25.4	79.8	193.0	469.8	1,377.8	4,576.6
2005 4/.....	235.2	7.1	0.7	0.5	2.2	6.6	25.8	80.9	199.1	499.9	1,453.7	4,879.3
2004 4/.....	237.4	9.0	0.9	0.5	2.1	6.5	26.3	81.2	203.4	521.2	1,500.0	4,977.6
2003 4/.....	251.7	8.8	1.0	0.5	2.3	6.7	27.2	82.9	216.5	561.8	1,603.6	5,363.5
2002.....	258.5	10.6	0.9	0.6	2.1	6.5	27.3	84.8	224.5	595.1	1,666.9	5,529.1
2001.....	262.8	9.9	1.2	0.6	2.0	6.6	25.9	83.9	229.6	612.2	1,714.3	5,752.1
2000.....	270.2	11.5	1.1	0.6	2.1	6.1	25.8	85.4	243.2	642.5	1,767.8	6,020.8
1999.....	277.7	11.8	1.0	0.6	2.3	6.4	26.3	86.8	251.1	677.7	1,837.5	6,152.7

White, male

2006 4/.....	226.9	7.0	0.9	0.6	2.9	9.4	36.3	119.2	278.9	636.6	1,743.3	4,947.1
2005 4/.....	234.9	7.6	0.9	0.6	3.0	9.1	37.2	121.3	288.2	671.9	1,831.8	5,288.4
2004 4/.....	236.5	9.6	1.0	0.5	2.7	8.8	37.8	122.2	294.4	703.2	1,897.1	5,348.4
2003 4/.....	249.5	9.4	0.8	0.6	2.8	9.0	39.1	125.3	313.2	761.1	2,030.1	5,747.2
2002.....	256.0	11.3	1.0	0.6	2.8	8.8	39.7	128.6	324.0	807.8	2,112.0	5,939.8
2001.....	257.8	10.6	1.2	0.6	2.6	8.7	37.8	127.0	330.8	829.1	2,175.8	6,157.2
2000.....	265.8	11.9	1.3	0.7	2.6	8.1	37.8	130.7	351.8	877.8	2,247.0	6,560.8
1999.....	272.9	12.0	1.2	0.7	2.7	8.4	38.6	132.1	361.4	928.4	2,338.8	6,669.0

White, female

2006 4/.....	224.2	6.2	0.8	0.5	1.4	3.9	14.1	40.7	111.4	325.8	1,123.9	4,402.6
2005 4/.....	235.5	6.6	0.6	0.5	1.4	4.0	14.1	40.8	114.5	351.8	1,193.3	4,691.0
2004 4/.....	238.3	8.4	0.9	0.5	1.4	4.0	14.5	40.7	117.2	365.4	1,229.1	4,810.4
2003 4/.....	253.8	8.1	1.1	0.4	1.7	4.4	15.1	41.1	125.2	392.0	1,315.2	5,193.6
2002.....	261.0	9.9	0.8	0.5	1.4	4.0	14.8	41.7	130.6	414.7	1,368.2	5,350.6
2001.....	267.7	9.2	1.1	0.6	1.5	4.4	13.9	41.5	134.3	429.0	1,407.9	5,582.5
2000.....	274.5	11.1	0.8	0.4	1.6	4.0	13.7	40.9	141.3	445.2	1,452.4	5,801.4
1999.....	282.4	11.6	0.9	0.5	1.8	4.3	13.8	42.2	147.8	468.7	1,509.0	5,945.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

Black, both sexes

2006 4/.....	182.6	17.1	1.7	1.0	4.6	18.3	53.9	165.9	374.1	772.8	1,703.9	4,056.3
2005 4/.....	189.8	16.1	1.7	1.2	5.1	18.3	55.8	171.8	395.7	792.9	1,815.7	4,326.7
2004 4/.....	192.3	15.3	2.4	1.2	4.9	17.7	56.5	176.1	402.7	839.8	1,844.1	4,563.8
2003 4/.....	202.8	20.2	2.2	1.1	5.6	18.1	59.5	181.4	427.8	913.0	1,962.1	4,941.2
2002.....	205.6	21.3	1.6	1.0	4.9	18.4	57.8	180.9	442.9	924.5	2,048.8	5,115.2
2001.....	208.8	21.6	3.0	1.1	5.4	17.9	59.5	179.9	447.6	965.7	2,108.4	5,275.7
2000.....	211.8	19.8	2.0	1.1	5.6	17.2	57.4	181.5	466.0	1,002.6	2,159.7	5,389.5
1999.....	217.2	23.8	2.0	1.1	5.8	16.7	62.1	185.4	481.3	1,048.6	2,211.1	5,487.4

Black, male

2006 4/.....	191.8	17.9	2.0	1.0	5.4	24.4	70.5	229.8	526.4	1,044.6	2,129.9	4,073.1
2005 4/.....	194.8	18.3	*	1.1	6.9	23.5	74.0	237.4	549.1	1,041.6	2,204.1	4,230.5
2004 4/.....	196.7	16.3	2.4	1.1	6.4	22.5	72.8	240.0	560.2	1,096.6	2,235.5	4,637.3
2003 4/.....	206.0	24.1	2.5	1.2	7.3	22.2	77.1	248.1	580.9	1,195.5	2,426.6	4,850.3
2002.....	206.3	22.1	*	1.0	6.2	23.9	75.6	246.0	605.3	1,192.7	2,449.6	5,125.7
2001.....	209.0	19.5	3.2	1.0	6.9	22.8	76.7	242.6	602.2	1,245.8	2,569.3	5,459.9
2000.....	211.1	19.8	1.9	1.2	6.6	21.2	74.3	247.2	631.2	1,268.8	2,597.6	5,633.5
1999.....	218.5	23.4	2.1	1.3	7.4	21.0	81.9	255.0	660.0	1,346.4	2,623.5	5,858.8

Black, female

2006 4/.....	174.3	16.2	*	1.0	3.8	12.6	39.2	111.0	251.3	578.3	1,461.7	4,049.4
2005 4/.....	185.2	13.9	2.0	1.3	3.2	13.4	39.7	115.4	272.0	614.9	1,595.1	4,365.6
2004 4/.....	188.3	14.2	2.3	1.3	3.4	13.3	42.0	121.2	276.0	656.5	1,622.9	4,534.7
2003 4/.....	200.0	16.1	2.0	1.0	3.9	14.4	43.9	124.1	304.7	712.0	1,699.6	4,976.5
2002.....	205.0	20.6	1.8	1.0	3.7	13.5	42.1	124.9	312.3	734.0	1,821.9	5,111.2
2001.....	208.6	23.9	2.7	1.1	3.8	13.5	44.2	125.9	323.1	768.0	1,849.6	5,207.3
2000.....	212.6	19.8	2.0	1.0	4.7	13.6	42.4	125.0	332.8	815.2	1,913.1	5,298.7
1999.....	216.0	24.2	1.9	1.0	4.3	12.8	44.6	125.8	337.9	838.6	1,980.3	5,348.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes

2006.....	1.1	*	*	*	*	0.1	0.2	0.4	1.1	3.2	8.6	16.7
2005.....	1.1	*	*	*	*	0.1	0.2	0.4	1.2	3.3	8.5	19.0
2004.....	1.1	*	*	*	*	0.1	0.2	0.5	1.1	3.3	8.8	17.1
2003.....	1.2	*	*	*	*	0.1	0.2	0.5	1.3	3.8	9.6	20.5
2002.....	1.2	*	*	*	*	0.1	0.2	0.6	1.6	3.8	9.6	17.8
2001.....	1.2	*	*	*	*	0.1	0.3	0.6	1.4	4.1	9.4	17.7
2000.....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.5	9.5	16.8
1999.....	1.3	*	*	*	*	0.1	0.3	0.7	1.7	4.6	10.0	18.0

All races 3/, male

2006.....	0.7	*	*	*	*	*	0.1	0.4	0.9	2.4	6.2	12.0
2005.....	0.7	*	*	*	*	*	0.2	0.4	1.0	2.5	6.2	14.0
2004.....	0.7	*	*	*	*	*	0.2	0.4	0.8	2.4	7.3	11.5
2003.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.9	7.5	16.2
2002.....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.0	6.7	11.8
2001.....	0.7	*	*	*	*	*	0.2	0.5	1.1	2.8	6.6	12.8
2000.....	0.8	*	*	*	*	*	0.3	0.6	1.4	2.9	6.6	12.2
1999.....	0.8	*	*	*	*	*	0.2	0.5	1.2	3.5	6.8	10.3

All races 3/, female

2006.....	1.5	*	*	*	*	*	0.2	0.4	1.3	3.9	10.3	18.9
2005.....	1.5	*	*	*	*	*	0.2	0.5	1.4	4.0	10.1	21.3
2004.....	1.5	*	*	*	*	*	0.3	0.6	1.4	4.1	9.9	19.5
2003.....	1.7	*	*	*	*	*	0.2	0.7	1.5	4.5	11.0	22.3
2002.....	1.7	*	*	*	*	*	0.2	0.7	2.0	4.6	11.6	20.4
2001.....	1.7	*	*	*	*	*	0.3	0.7	1.7	5.1	11.3	19.8
2000.....	1.8	*	*	*	*	0.1	0.4	0.8	2.0	5.8	11.4	18.7
1999.....	1.9	*	*	*	*	0.1	0.3	0.8	2.2	5.6	12.2	21.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

White, both sexes

2006 4/.....	1.2	*	*	*	*	0.1	0.1	0.3	1.0	3.2	9.0	18.0
2005 4/.....	1.3	*	*	*	*	*	0.2	0.4	1.1	3.4	9.0	20.4
2004 4/.....	1.2	*	*	*	*	*	0.2	0.4	1.0	3.4	9.3	18.2
2003 4/.....	1.4	*	*	*	*	*	0.2	0.4	1.2	3.8	10.2	21.8
2002.....	1.4	*	*	*	*	0.1	0.2	0.5	1.5	4.0	10.1	18.9
2001.....	1.3	*	*	*	*	*	0.2	0.5	1.4	4.1	10.0	18.7
2000.....	1.4	*	*	*	*	0.1	0.3	0.6	1.7	4.6	10.1	17.6
1999.....	1.4	*	*	*	*	0.1	0.2	0.6	1.7	4.8	10.5	18.9

White, male

2006 4/.....	0.7	*	*	*	*	*	0.1	0.3	0.8	2.4	6.5	12.9
2005 4/.....	0.8	*	*	*	*	*	0.2	0.3	0.9	2.5	6.4	14.7
2004 4/.....	0.8	*	*	*	*	*	0.1	0.4	0.7	2.5	7.8	12.4
2003 4/.....	0.9	*	*	*	*	*	0.1	0.3	0.9	3.0	8.1	17.3
2002.....	0.8	*	*	*	*	*	0.2	0.4	1.1	3.0	7.1	12.6
2001.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.8	6.9	13.3
2000.....	0.9	*	*	*	*	*	0.2	0.6	1.5	2.9	7.0	12.7
1999.....	0.8	*	*	*	*	*	0.2	0.5	1.1	3.7	7.0	11.1

White, female

2006 4/.....	1.7	*	*	*	*	*	0.1	0.4	1.3	4.0	10.8	20.4
2005 4/.....	1.7	*	*	*	*	*	0.2	0.4	1.3	4.1	10.8	23.1
2004 4/.....	1.7	*	*	*	*	*	0.2	0.5	1.4	4.2	10.4	20.8
2003 4/.....	1.8	*	*	*	*	*	0.2	0.5	1.4	4.6	11.6	23.8
2002.....	1.9	*	*	*	*	*	0.2	0.6	1.9	4.8	12.1	21.6
2001.....	1.8	*	*	*	*	*	0.2	0.6	1.6	5.2	12.0	21.0
2000.....	1.9	*	*	*	*	*	0.3	0.7	1.9	6.0	12.2	19.6
1999.....	2.0	*	*	*	*	*	0.3	0.7	2.2	5.7	12.8	22.0



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

Black, both sexes

2006 4/.....	0.6	*	*	*	*	*	0.4	0.8	1.7	2.9	5.3	*
2005 4/.....	0.6	*	*	*	*	*	*	0.9	1.4	2.4	4.4	6.1
2004 4/.....	0.6	*	*	*	*	*	0.6	1.0	1.6	2.7	4.0	5.9
2003 4/.....	0.8	*	*	*	*	*	0.5	1.3	2.6	3.9	4.1	7.6
2002.....	0.7	*	*	*	*	*	0.3	0.9	2.4	2.9	4.4	*
2001.....	0.8	*	*	*	*	*	0.7	1.3	2.1	4.1	4.2	6.8
2000.....	0.7	*	*	*	*	*	0.6	1.0	2.0	3.5	3.9	8.4
1999.....	0.8	*	*	*	*	*	0.6	1.2	2.2	3.4	4.4	8.3

Black, male

2006 4/.....	0.5	*	*	*	*	*	*	*	1.6	*	*	*
2005 4/.....	0.4	*	*	*	*	*	*	*	*	*	*	*
2004 4/.....	0.5	*	*	*	*	*	*	0.9	*	*	*	*
2003 4/.....	0.6	*	*	*	*	*	*	0.9	2.7	3.0	*	*
2002.....	0.5	*	*	*	*	*	*	*	1.9	3.0	*	*
2001.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2000.....	0.5	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	*	*	*	*

Black, female

2006 4/.....	0.8	*	*	*	*	*	*	0.8	1.7	3.2	6.2	*
2005 4/.....	0.7	*	*	*	*	*	*	1.1	1.6	2.8	4.5	*
2004 4/.....	0.8	*	*	*	*	*	*	1.1	1.7	3.3	5.0	*
2003 4/.....	1.0	*	*	*	*	*	*	1.7	2.5	4.5	5.1	8.1
2002.....	0.8	*	*	*	*	*	*	1.1	2.8	2.8	5.0	*
2001.....	1.1	*	*	*	*	*	0.9	1.7	2.6	5.3	5.3	*
2000.....	1.0	*	*	*	*	*	0.8	1.3	2.7	4.3	4.8	*
1999.....	1.0	*	*	*	*	*	0.7	1.4	2.6	4.1	4.5	11.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive heart disease (I11)

All races 3/, both sexes

2006.....	9.9	*	*	*	0.1	0.7	3.1	8.3	13.1	20.4	47.8	194.0
2005.....	9.9	*	*	*	0.1	0.8	3.1	7.8	13.0	20.4	48.8	198.6
2004.....	9.7	*	*	*	0.1	0.8	3.0	7.8	13.4	21.7	49.3	192.8
2003.....	9.7	*	*	*	0.1	0.8	2.9	7.9	13.1	22.1	51.0	195.0
2002.....	9.2	*	*	*	0.1	0.7	2.7	7.2	12.5	21.3	49.6	187.0
2001.....	8.7	*	*	*	0.1	0.6	2.4	6.3	10.9	19.8	48.7	189.5
2000.....	8.4	*	*	*	0.1	0.5	2.2	5.6	11.0	19.6	48.2	193.1
1999.....	8.1	*	*	*	0.1	0.5	2.1	5.3	10.7	19.1	47.7	184.9

All races 3/, male

2006.....	9.3	*	*	*	0.2	1.0	4.2	11.7	18.0	25.7	49.7	143.0
2005.....	9.1	*	*	*	0.1	1.1	4.4	10.9	18.1	25.2	48.7	147.5
2004.....	8.9	*	*	*	0.1	1.1	4.1	10.7	18.5	27.3	48.0	145.4
2003.....	8.8	*	*	*	0.2	1.0	4.0	10.8	17.2	27.1	51.0	142.2
2002.....	8.2	*	*	*	0.1	1.1	3.8	9.8	16.9	25.2	49.4	134.6
2001.....	7.5	*	*	*	*	0.8	3.3	8.6	14.4	23.6	48.4	134.4
2000.....	7.2	*	*	*	0.1	0.7	2.9	7.8	14.5	23.7	46.5	139.1
1999.....	6.8	*	*	*	0.1	0.7	2.9	7.0	13.7	21.9	46.8	133.6

All races 3/, female

2006.....	10.6	*	*	*	*	0.4	2.0	5.0	8.5	15.9	46.5	217.9
2005.....	10.7	*	*	*	*	0.5	1.9	4.7	8.3	16.4	48.9	222.0
2004.....	10.5	*	*	*	*	0.4	1.9	5.1	8.6	16.9	50.2	214.1
2003.....	10.7	*	*	*	*	0.5	1.9	5.2	9.3	17.8	51.0	218.3
2002.....	10.2	*	*	*	*	0.4	1.6	4.6	8.4	18.1	49.8	209.7
2001.....	9.8	*	*	*	*	0.4	1.6	4.1	7.8	16.6	48.8	212.6
2000.....	9.7	*	*	*	*	0.4	1.5	3.5	7.8	16.2	49.4	215.1
1999.....	9.4	*	*	*	*	0.3	1.3	3.6	7.9	16.8	48.3	205.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive heart disease (I11)

White, both sexes

2006 4/.....	9.2	*	*	*	0.1	0.4	2.3	6.1	10.1	15.9	42.2	186.9
2005 4/.....	9.0	*	*	*	*	0.5	2.2	5.5	10.1	16.2	42.4	188.9
2004 4/.....	8.9	*	*	*	0.1	0.5	2.1	5.6	10.0	17.6	43.2	183.7
2003 4/.....	8.8	*	*	*	0.1	0.5	2.0	5.4	9.8	17.3	45.1	185.3
2002.....	8.3	*	*	*	*	0.4	1.9	5.1	9.1	16.7	42.9	177.3
2001.....	7.8	*	*	*	*	0.4	1.6	4.2	7.9	15.2	42.6	179.9
2000.....	7.6	*	*	*	*	0.3	1.5	3.7	7.6	15.0	41.5	181.8
1999.....	7.3	*	*	*	*	0.3	1.3	3.4	7.5	14.4	40.9	175.3

White, male

2006 4/.....	8.1	*	*	*	0.2	0.7	3.4	9.1	14.1	19.8	42.8	134.5
2005 4/.....	7.9	*	*	*	*	0.8	3.2	8.1	14.3	20.0	42.3	136.9
2004 4/.....	7.8	*	*	*	*	0.7	3.1	8.1	13.9	22.3	41.9	136.4
2003 4/.....	7.7	*	*	*	0.1	0.7	3.1	7.9	13.4	21.3	44.8	131.9
2002.....	7.1	*	*	*	*	0.7	2.8	7.4	12.7	19.9	42.6	124.8
2001.....	6.4	*	*	*	*	0.6	2.4	6.1	10.9	18.8	41.9	123.7
2000.....	6.1	*	*	*	*	0.4	2.1	5.5	10.4	18.6	39.6	129.3
1999.....	5.8	*	*	*	*	0.6	2.0	4.9	10.0	16.6	39.7	123.7

White, female

2006 4/.....	10.2	*	*	*	*	0.2	1.2	3.2	6.3	12.5	41.7	211.5
2005 4/.....	10.1	*	*	*	*	0.3	1.1	2.9	6.1	12.9	42.5	212.9
2004 4/.....	9.9	*	*	*	*	0.2	1.1	3.1	6.3	13.5	44.0	205.0
2003 4/.....	10.0	*	*	*	*	0.2	1.0	2.9	6.4	13.9	45.2	209.0
2002.....	9.5	*	*	*	*	0.2	0.9	2.8	5.8	14.1	43.1	200.1
2001.....	9.2	*	*	*	*	0.2	0.8	2.4	5.1	12.3	43.0	203.5
2000.....	9.0	*	*	*	*	0.2	0.9	1.9	5.0	11.9	42.9	203.1
1999.....	8.7	*	*	*	*	*	0.7	2.0	5.1	12.5	41.7	196.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Hypertensive heart disease (I11)

Black, both sexes

2006 4/.....	17.3	*	*	*	*	2.5	8.9	25.3	41.8	65.0	116.9	299.6
2005 4/.....	17.7	*	*	*	0.4	2.5	9.5	25.1	41.2	63.1	128.4	338.1
2004 4/.....	17.6	*	*	*	0.3	2.7	9.2	26.2	45.1	64.6	126.8	314.7
2003 4/.....	17.9	*	*	*	0.4	2.5	9.4	27.9	44.7	69.3	126.3	332.3
2002.....	17.2	*	*	*	*	2.8	8.6	24.3	44.4	68.0	133.0	320.4
2001.....	16.0	*	*	*	*	2.1	8.3	23.0	39.2	65.6	124.4	326.1
2000.....	16.1	*	*	*	*	2.1	7.4	21.2	42.2	66.8	131.9	340.0
1999.....	15.5	*	*	*	0.3	1.8	7.9	20.9	39.6	67.2	131.9	305.1

Black, male

2006 4/.....	18.8	*	*	*	*	3.5	11.0	33.7	58.6	88.7	143.7	272.6
2005 4/.....	18.5	*	*	*	*	2.9	12.8	33.9	58.1	81.4	138.5	301.2
2004 4/.....	18.4	*	*	*	*	3.5	11.7	33.6	64.6	86.6	131.7	267.2
2003 4/.....	18.1	*	*	*	*	3.0	11.2	35.0	57.1	92.3	137.1	287.8
2002.....	17.5	*	*	*	*	3.9	11.6	31.3	60.8	83.6	141.1	267.0
2001.....	15.9	*	*	*	*	2.6	10.3	30.2	49.8	79.1	137.1	284.8
2000.....	15.6	*	*	*	*	2.6	9.2	27.8	54.9	81.4	139.3	273.3
1999.....	15.2	*	*	*	*	2.1	10.1	26.3	50.6	80.4	142.3	263.6

Black, female

2006 4/.....	15.9	*	*	*	*	1.5	7.0	18.2	28.3	48.0	101.7	310.7
2005 4/.....	17.0	*	*	*	*	2.1	6.6	17.5	27.6	50.0	122.7	353.0
2004 4/.....	16.9	*	*	*	*	1.9	7.0	19.8	29.4	48.9	124.0	333.5
2003 4/.....	17.8	*	*	*	*	2.0	7.9	21.8	34.7	52.9	120.2	349.6
2002.....	16.9	*	*	*	*	1.7	6.0	18.3	31.2	56.8	128.5	340.9
2001.....	16.2	*	*	*	*	1.7	6.5	16.8	30.6	56.1	117.3	341.4
2000.....	16.5	*	*	*	*	1.7	5.8	15.5	31.9	56.6	127.7	364.9
1999.....	15.9	*	*	*	*	1.5	5.9	16.2	30.8	57.9	126.0	320.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Hypertensive heart and renal disease (I13)

All races 3/, both sexes

2006.....	1.0	*	*	*	*	0.1	0.2	0.4	0.8	1.9	6.2	22.6
2005.....	1.1	*	*	*	*	0.1	0.2	0.4	0.8	2.3	7.0	25.3
2004.....	1.0	*	*	*	*	0.1	0.2	0.5	0.9	2.1	6.6	24.8
2003.....	1.1	*	*	*	*	0.1	0.2	0.5	1.0	2.2	7.0	25.6
2002.....	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.2	6.8	23.5
2001.....	1.0	*	*	*	*	0.1	0.2	0.4	1.0	2.3	6.8	23.3
2000.....	1.0	*	*	*	*	0.1	0.2	0.4	0.9	2.3	7.0	24.3
1999.....	1.2	*	*	*	*	0.1	0.3	0.4	1.1	2.6	8.9	29.1

All races 3/, male

2006.....	0.9	*	*	*	*	0.1	0.3	0.5	1.1	2.0	6.9	23.8
2005.....	0.9	*	*	*	*	0.1	0.2	0.6	1.0	2.5	7.2	27.4
2004.....	0.9	*	*	*	*	*	0.3	0.6	1.1	2.2	7.2	24.3
2003.....	0.9	*	*	*	*	*	0.3	0.6	1.2	2.3	7.4	26.7
2002.....	0.9	*	*	*	*	*	0.2	0.5	1.2	2.4	7.0	23.1
2001.....	0.8	*	*	*	*	*	0.3	0.5	1.0	2.5	6.8	24.2
2000.....	0.8	*	*	*	*	*	0.2	0.5	0.9	2.5	7.2	27.8
1999.....	1.0	*	*	*	*	*	0.3	0.6	1.3	2.7	9.6	31.3

All races 3/, female

2006.....	1.1	*	*	*	*	*	0.1	0.4	0.6	1.8	5.7	22.1
2005.....	1.2	*	*	*	*	*	0.1	0.3	0.6	2.1	6.8	24.3
2004.....	1.2	*	*	*	*	*	0.1	0.4	0.7	1.9	6.2	25.0
2003.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	25.2
2002.....	1.2	*	*	*	*	*	0.1	0.3	0.8	2.0	6.7	23.7
2001.....	1.1	*	*	*	*	*	0.2	0.2	0.9	2.1	6.8	22.9
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.1	6.9	22.9
1999.....	1.4	*	*	*	*	*	0.2	0.3	0.9	2.4	8.5	28.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive heart and renal disease (I13)

White, both sexes

2006 4/.....	0.8	*	*	*	*	*	0.1	0.2	0.4	1.2	5.1	20.9
2005 4/.....	0.9	*	*	*	*	*	0.1	0.2	0.4	1.5	5.7	23.9
2004 4/.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.3	5.5	23.0
2003 4/.....	0.9	*	*	*	*	*	0.1	0.2	0.5	1.4	6.0	23.4
2002.....	0.8	*	*	*	*	*	0.1	0.1	0.5	1.4	5.7	21.0
2001.....	0.8	*	*	*	*	*	0.1	0.2	0.5	1.5	5.7	21.0
2000.....	0.8	*	*	*	*	*	0.1	0.2	0.4	1.5	5.6	22.0
1999.....	1.0	*	*	*	*	*	0.1	0.2	0.5	1.5	7.3	26.1

White, male

2006 4/.....	0.7	*	*	*	*	*	0.1	0.2	0.5	1.2	5.7	22.5
2005 4/.....	0.7	*	*	*	*	*	*	0.3	0.5	1.6	5.8	25.5
2004 4/.....	0.7	*	*	*	*	*	*	0.2	0.6	1.5	6.1	22.9
2003 4/.....	0.7	*	*	*	*	*	*	0.2	0.6	1.6	6.6	24.3
2002.....	0.7	*	*	*	*	*	*	0.1	0.6	1.6	6.0	20.6
2001.....	0.7	*	*	*	*	*	*	0.2	0.5	1.6	5.9	22.7
2000.....	0.7	*	*	*	*	*	0.1	0.2	0.5	1.7	6.2	24.4
1999.....	0.8	*	*	*	*	*	*	0.2	0.6	1.7	8.2	28.8

White, female

2006 4/.....	0.9	*	*	*	*	*	*	0.2	0.3	1.2	4.7	20.1
2005 4/.....	1.1	*	*	*	*	*	*	*	0.3	1.4	5.6	23.2
2004 4/.....	1.0	*	*	*	*	*	*	0.1	0.4	1.2	5.1	23.0
2003 4/.....	1.0	*	*	*	*	*	*	*	0.4	1.3	5.5	22.9
2002.....	1.0	*	*	*	*	*	*	*	0.4	1.2	5.6	21.1
2001.....	1.0	*	*	*	*	*	*	0.1	0.5	1.4	5.5	20.3
2000.....	1.0	*	*	*	*	*	*	0.1	0.4	1.4	5.2	21.1
1999.....	1.1	*	*	*	*	*	*	0.1	0.5	1.4	6.7	25.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Hypertensive heart and renal disease (I13)

Black, both sexes

2006 4/.....	2.1	*	*	*	*	*	0.9	2.1	4.3	8.5	18.5	41.8
2005 4/.....	2.3	*	*	*	*	0.4	0.9	2.4	4.5	9.6	21.4	42.8
2004 4/.....	2.3	*	*	*	*	*	1.2	2.8	4.9	8.5	19.6	48.2
2003 4/.....	2.4	*	*	*	*	0.4	1.2	3.0	5.4	9.0	19.6	53.1
2002.....	2.4	*	*	*	*	*	0.9	2.7	6.1	9.9	19.6	56.1
2001.....	2.2	*	*	*	*	0.4	1.2	2.1	5.0	9.9	20.1	50.1
2000.....	2.2	*	*	*	*	*	0.8	2.2	4.6	9.9	22.8	50.4
1999.....	2.8	*	*	*	*	*	1.5	2.4	6.0	12.5	26.9	65.4

Black, male

2006 4/.....	2.1	*	*	*	*	*	1.2	2.6	6.1	10.5	23.0	39.3
2005 4/.....	2.3	*	*	*	*	*	1.2	3.3	6.0	11.1	23.2	50.2
2004 4/.....	2.2	*	*	*	*	*	1.9	3.5	6.1	8.7	23.1	41.7
2003 4/.....	2.3	*	*	*	*	*	1.7	4.0	7.1	8.9	18.9	50.0
2002.....	2.3	*	*	*	*	*	1.0	3.8	7.5	10.1	19.2	61.1
2001.....	2.1	*	*	*	*	*	1.6	3.2	5.9	11.9	18.5	44.3
2000.....	2.0	*	*	*	*	*	1.0	2.8	5.3	11.3	20.9	63.4
1999.....	2.5	*	*	*	*	*	1.8	3.5	7.6	13.5	25.8	59.8

Black, female

2006 4/.....	2.0	*	*	*	*	*	*	1.7	3.0	7.0	16.0	42.8
2005 4/.....	2.2	*	*	*	*	*	0.7	1.7	3.3	8.5	20.3	39.8
2004 4/.....	2.4	*	*	*	*	*	*	2.3	4.0	8.3	17.6	50.8
2003 4/.....	2.5	*	*	*	*	*	0.8	2.1	4.0	9.1	19.9	54.3
2002.....	2.5	*	*	*	*	*	0.9	1.8	4.9	9.7	19.8	54.2
2001.....	2.3	*	*	*	*	*	0.8	1.1	4.2	8.6	21.0	52.2
2000.....	2.4	*	*	*	*	*	*	1.6	4.1	9.0	23.9	45.5
1999.....	3.0	*	*	*	*	*	1.2	1.5	4.8	11.7	27.5	67.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Ischemic heart diseases (I20-I25)

All races 3/, both sexes

2006.....	142.1	0.5	*	*	0.3	2.6	15.3	57.7	147.9	352.5	959.8	2,904.7
2005.....	150.4	*	*	*	0.4	2.5	15.6	59.6	154.2	376.2	1,029.8	3,158.1
2004.....	153.7	*	*	*	0.3	2.7	16.2	59.9	158.2	395.9	1,072.6	3,287.4
2003.....	165.1	0.5	*	0.1	0.4	2.7	17.0	62.2	169.8	430.2	1,160.7	3,610.2
2002.....	171.4	0.6	*	0.0	0.3	2.5	17.1	63.8	177.7	458.9	1,220.2	3,775.0
2001.....	176.3	0.6	*	0.0	0.4	2.8	16.7	64.0	184.0	477.6	1,271.3	3,958.2
2000.....	183.1	0.8	*	*	0.3	2.3	16.7	66.1	196.7	505.1	1,324.2	4,182.5
1999.....	189.8	0.7	*	0.1	0.4	2.5	17.2	67.7	203.5	536.6	1,384.8	4,318.1

All races 3/, male

2006.....	152.2	*	*	*	0.5	4.0	23.2	89.1	220.7	494.2	1,269.7	3,303.0
2005.....	159.0	*	*	*	0.6	3.8	23.6	92.0	229.4	521.7	1,351.5	3,576.8
2004.....	161.6	*	*	*	0.4	4.0	24.4	92.8	234.8	549.1	1,410.3	3,680.3
2003.....	172.2	*	*	*	0.5	4.0	25.7	97.2	251.5	598.8	1,524.4	4,005.2
2002.....	178.4	*	*	*	0.4	3.7	26.1	99.4	263.3	640.1	1,598.9	4,205.3
2001.....	181.7	*	*	*	0.6	4.1	25.5	99.5	271.6	663.7	1,663.5	4,395.8
2000.....	188.7	*	*	*	0.4	3.5	25.8	103.5	290.2	705.2	1,736.8	4,701.7
1999.....	195.4	*	*	*	0.5	3.7	26.6	106.2	299.3	749.8	1,813.7	4,837.5

All races 3/, female

2006.....	132.3	*	*	*	0.2	1.1	7.4	27.4	80.1	232.6	747.9	2,718.3
2005.....	142.0	*	*	*	0.1	1.2	7.6	28.2	84.2	253.4	811.4	2,965.8
2004.....	146.1	*	*	*	0.1	1.4	8.0	28.0	87.1	267.3	845.3	3,110.6
2003.....	158.1	*	*	*	0.3	1.4	8.4	28.3	94.0	289.2	917.7	3,435.5
2002.....	164.7	*	*	*	0.2	1.3	8.2	29.5	98.5	308.1	968.9	3,588.3
2001.....	171.2	*	*	*	0.2	1.5	8.1	29.7	103.1	323.4	1,013.6	3,775.1
2000.....	177.6	*	*	*	0.2	1.2	7.6	29.9	110.4	340.3	1,055.2	3,971.0
1999.....	184.5	*	*	*	0.2	1.4	7.9	30.7	115.1	362.0	1,106.3	4,108.8



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Ischemic heart diseases (I20-I25)

White, both sexes

2006 4/.....	153.4	*	*	*	0.3	2.4	14.9	55.0	141.5	342.7	959.7	2,964.3
2005 4/.....	162.6	*	*	*	0.3	2.3	15.1	56.6	146.8	367.8	1,029.7	3,227.0
2004 4/.....	166.1	*	*	*	0.2	2.5	15.7	56.8	151.3	385.6	1,073.4	3,343.9
2003 4/.....	178.3	*	*	*	0.3	2.5	16.4	58.8	162.0	418.9	1,159.0	3,671.7
2002.....	185.4	*	*	*	0.3	2.3	16.5	60.7	169.5	448.8	1,217.8	3,836.4
2001.....	190.5	*	*	*	0.3	2.6	15.9	60.7	175.1	465.8	1,268.5	4,022.7
2000.....	197.9	0.8	*	*	0.3	2.1	15.8	62.8	187.3	493.4	1,320.9	4,257.7
1999.....	205.1	*	*	*	0.3	2.3	16.2	64.4	194.3	524.8	1,383.7	4,389.2

White, male

2006 4/.....	163.7	*	*	*	0.5	3.8	22.9	86.2	212.7	483.8	1,275.5	3,396.0
2005 4/.....	171.5	*	*	*	0.5	3.5	23.2	88.7	220.9	514.6	1,360.1	3,687.1
2004 4/.....	174.3	*	*	*	0.3	3.7	24.1	89.6	227.1	539.9	1,422.1	3,768.8
2003 4/.....	185.5	*	*	*	0.5	3.8	25.1	93.5	243.5	587.9	1,529.5	4,106.1
2002.....	192.5	*	*	*	0.4	3.5	25.6	96.3	254.3	632.9	1,608.0	4,299.7
2001.....	195.8	*	*	*	0.5	3.9	24.7	96.3	262.3	653.8	1,672.4	4,494.0
2000.....	203.6	*	*	*	0.4	3.3	25.1	100.5	280.3	697.3	1,744.8	4,813.9
1999.....	210.5	*	*	*	0.4	3.4	25.5	102.8	289.0	741.3	1,827.8	4,933.9

White, female

2006 4/.....	143.3	*	*	*	0.1	1.0	6.6	24.1	73.8	221.0	740.4	2,761.6
2005 4/.....	153.9	*	*	*	*	1.1	6.8	24.8	76.5	241.6	802.1	3,015.2
2004 4/.....	158.1	*	*	*	*	1.2	7.2	24.4	79.5	253.5	835.6	3,152.3
2003 4/.....	171.4	*	*	*	0.2	1.2	7.5	24.6	85.0	274.9	908.5	3,479.3
2002.....	178.4	*	*	*	0.2	1.1	7.3	25.5	89.5	292.7	956.0	3,635.2
2001.....	185.3	*	*	*	0.2	1.3	7.0	25.6	93.1	307.2	1,000.3	3,825.5
2000.....	192.4	*	*	*	0.2	0.9	6.5	25.6	100.0	322.5	1,042.0	4,031.8
1999.....	200.0	*	*	*	0.1	1.3	6.8	26.7	105.5	344.3	1,092.6	4,170.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Ischemic heart diseases (I20-I25)

Black, both sexes

2006 4/.....	112.6	*	*	*	0.5	4.1	22.4	90.6	233.6	508.0	1,127.9	2,616.1
2005 4/.....	117.8	*	*	*	0.5	3.9	23.0	94.0	250.8	522.6	1,213.7	2,804.4
2004 4/.....	121.0	*	*	*	0.7	4.7	23.2	96.0	255.0	568.6	1,243.5	3,014.3
2003 4/.....	129.2	*	*	*	0.7	4.3	24.7	99.3	273.1	615.2	1,363.0	3,310.1
2002.....	132.9	*	*	*	0.5	3.9	25.1	101.6	288.0	636.1	1,422.7	3,462.9
2001.....	136.9	*	*	*	0.9	4.6	25.6	102.3	297.9	673.0	1,479.0	3,615.1
2000.....	140.1	*	*	*	0.4	4.3	25.6	105.6	314.6	702.2	1,523.2	3,684.7
1999.....	143.7	*	*	*	0.8	3.9	27.5	108.5	320.1	735.6	1,552.7	3,805.8

Black, male

2006 4/.....	120.2	*	*	*	0.6	6.1	31.1	130.9	340.1	704.9	1,456.9	2,656.7
2005 4/.....	122.9	*	*	*	0.8	6.0	32.3	137.0	358.7	701.8	1,526.4	2,785.0
2004 4/.....	125.2	*	*	*	1.1	6.5	31.9	138.4	362.4	761.9	1,537.6	3,100.5
2003 4/.....	133.4	*	*	*	0.9	6.2	35.0	144.5	381.8	827.3	1,744.8	3,285.3
2002.....	135.3	*	*	*	*	5.1	35.2	144.3	407.3	840.8	1,748.8	3,556.2
2001.....	138.9	*	*	*	1.2	6.6	36.0	145.7	412.8	894.0	1,825.4	3,775.0
2000.....	141.5	*	*	*	*	5.9	35.6	149.4	439.2	907.9	1,877.0	3,885.2
1999.....	146.0	*	*	*	0.9	5.7	39.5	156.9	452.9	962.3	1,870.8	4,080.1

Black, female

2006 4/.....	105.6	*	*	*	*	2.3	14.7	56.0	147.8	367.2	940.8	2,599.5
2005 4/.....	113.1	*	*	*	*	2.0	14.8	57.1	163.9	394.2	1,036.2	2,812.2
2004 4/.....	117.1	*	*	*	*	3.0	15.5	59.6	168.6	430.7	1,077.3	2,980.1
2003 4/.....	125.3	*	*	*	*	2.5	15.6	60.5	185.6	464.3	1,147.2	3,319.7
2002.....	130.8	*	*	*	*	2.8	16.3	64.9	192.2	490.8	1,238.1	3,427.2
2001.....	135.2	*	*	*	*	2.8	16.4	65.0	205.3	517.1	1,284.5	3,555.7
2000.....	138.8	*	*	*	*	2.8	16.7	68.0	214.1	557.4	1,324.0	3,610.0
1999.....	141.7	*	*	*	0.7	2.3	16.9	67.0	213.6	575.8	1,374.6	3,703.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute myocardial infarction (I21-I22)

All races 3/, both sexes

2006.....	47.2	*	*	*	0.1	1.0	6.1	22.5	58.2	131.1	319.0	825.8
2005.....	50.9	*	*	*	0.2	1.0	6.2	23.7	61.1	143.1	348.2	923.1
2004.....	53.4	*	*	*	0.1	1.1	6.4	24.2	64.5	153.4	370.3	992.9
2003.....	58.7	*	*	*	0.2	1.0	7.0	25.5	70.9	168.6	412.7	1,118.8
2002.....	62.3	*	*	*	0.2	1.0	7.0	26.5	74.9	186.1	444.4	1,191.3
2001.....	64.9	*	*	*	0.2	1.2	7.1	27.6	78.4	193.2	471.2	1,265.4
2000.....	68.5	0.5	*	*	0.1	0.9	6.9	28.9	86.6	209.1	503.3	1,338.5
1999.....	71.5	*	*	*	0.2	1.1	7.2	29.6	90.0	225.0	529.3	1,379.3

All races 3/, male

2006.....	51.6	*	*	*	0.2	1.5	9.2	34.2	85.6	181.9	411.8	936.3
2005.....	54.8	*	*	*	0.2	1.5	9.2	36.2	89.9	195.5	445.9	1,045.0
2004.....	57.4	*	*	*	0.2	1.6	9.6	37.3	95.3	209.5	477.0	1,120.5
2003.....	62.6	*	*	*	0.2	1.5	10.4	39.8	104.1	231.0	530.6	1,263.5
2002.....	66.2	*	*	*	0.2	1.4	10.5	40.6	109.2	256.2	569.2	1,366.4
2001.....	68.6	*	*	*	0.2	1.7	10.6	42.4	114.0	264.7	604.1	1,461.3
2000.....	72.7	*	*	*	0.1	1.3	10.6	44.7	125.7	286.9	648.3	1,568.3
1999.....	76.1	*	*	*	0.2	1.5	11.0	46.3	131.8	310.0	684.4	1,622.1

All races 3/, female

2006.....	43.0	*	*	*	*	0.5	3.0	11.2	32.6	88.1	255.6	774.1
2005.....	47.2	*	*	*	*	0.5	3.2	11.6	34.3	98.9	281.8	867.1
2004.....	49.6	*	*	*	*	0.6	3.3	11.6	35.9	106.3	298.5	935.5
2003.....	54.8	*	*	*	0.1	0.6	3.6	11.7	40.0	116.5	333.9	1,054.8
2002.....	58.4	*	*	*	0.1	0.5	3.6	12.9	43.2	127.7	361.5	1,115.3
2001.....	61.3	*	*	*	*	0.7	3.5	13.3	45.4	134.0	383.9	1,183.5
2000.....	64.6	*	*	*	0.1	0.5	3.3	13.6	50.5	145.1	408.7	1,244.8
1999.....	67.0	*	*	*	0.1	0.6	3.4	13.5	51.5	155.4	428.6	1,281.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute myocardial infarction (I21-I22)

White, both sexes

2006 4/.....	51.0	*	*	*	0.1	1.0	6.1	22.1	57.3	129.6	319.5	836.0
2005 4/.....	55.1	*	*	*	0.2	0.9	6.1	23.3	60.0	142.3	348.3	936.8
2004 4/.....	57.8	*	*	*	0.1	1.1	6.4	23.6	63.5	151.2	372.1	1,004.1
2003 4/.....	63.4	*	*	*	0.2	1.0	6.8	24.7	69.5	166.8	412.7	1,132.6
2002.....	67.3	*	*	*	0.1	1.0	6.9	25.9	73.5	184.2	443.3	1,201.2
2001.....	70.1	*	*	*	0.1	1.1	6.8	26.9	76.8	190.8	471.1	1,277.8
2000.....	74.1	*	*	*	0.1	0.9	6.8	28.2	84.9	206.7	502.6	1,353.5
1999.....	77.4	*	*	*	0.1	1.0	7.0	29.1	88.6	223.3	530.2	1,393.2

White, male

2006 4/.....	55.8	*	*	*	0.2	1.5	9.3	34.1	84.9	181.8	414.8	955.4
2005 4/.....	59.5	*	*	*	0.2	1.3	9.2	36.1	89.2	196.8	449.5	1,070.1
2004 4/.....	62.3	*	*	*	0.1	1.6	9.7	37.0	95.1	208.6	484.2	1,143.7
2003 4/.....	67.7	*	*	*	0.2	1.4	10.3	39.2	103.6	230.7	532.6	1,289.3
2002.....	71.8	*	*	*	0.2	1.4	10.5	40.4	108.9	257.3	572.0	1,386.1
2001.....	74.4	*	*	*	0.2	1.6	10.6	42.4	113.7	264.7	609.1	1,481.1
2000.....	78.8	*	*	*	0.2	1.3	10.6	44.7	125.0	287.2	652.5	1,592.9
1999.....	82.7	*	*	*	0.2	1.4	10.9	46.4	131.3	311.1	692.3	1,644.0

White, female

2006 4/.....	46.3	*	*	*	*	0.5	2.8	10.2	31.0	84.5	253.3	780.0
2005 4/.....	50.8	*	*	*	*	0.4	2.9	10.6	32.3	95.4	278.5	875.4
2004 4/.....	53.3	*	*	*	*	0.5	3.0	10.4	33.5	102.1	295.6	941.1
2003 4/.....	59.2	*	*	*	*	0.5	3.2	10.4	37.2	112.3	331.5	1,063.2
2002.....	62.8	*	*	*	*	0.5	3.3	11.5	40.1	122.2	357.0	1,120.8
2001.....	66.0	*	*	*	*	0.7	3.1	11.7	42.0	128.4	379.6	1,192.7
2000.....	69.6	*	*	*	*	0.4	2.9	11.9	47.3	139.3	404.1	1,256.2
1999.....	72.3	*	*	*	*	0.6	3.1	12.1	48.6	150.1	423.9	1,292.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute myocardial infarction (I21-I22)

Black, both sexes

2006 4/.....	37.4	*	*	*	*	1.6	8.3	31.6	80.6	172.5	368.2	804.0
2005 4/.....	39.8	*	*	*	*	1.7	8.7	32.9	85.4	179.6	408.0	892.1
2004 4/.....	42.0	*	*	*	0.3	1.7	8.8	35.4	91.1	204.7	420.0	979.9
2003 4/.....	45.7	*	*	*	*	1.6	10.1	37.5	98.6	220.0	479.9	1,094.9
2002.....	48.5	*	*	*	*	1.6	9.7	37.9	106.9	238.8	517.3	1,201.2
2001.....	50.0	*	*	*	0.3	1.8	10.0	39.2	109.8	252.6	537.8	1,239.9
2000.....	52.2	*	*	*	*	1.7	9.9	41.1	120.1	268.7	571.1	1,275.7
1999.....	53.3	*	*	*	*	1.6	10.3	40.9	120.9	279.5	584.6	1,318.8

Black, male

2006 4/.....	39.1	*	*	*	*	2.4	11.5	44.0	114.6	228.5	454.9	814.1
2005 4/.....	40.3	*	*	*	*	2.5	11.7	45.7	118.3	228.4	492.8	885.9
2004 4/.....	42.4	*	*	*	*	2.0	11.6	50.3	123.5	265.8	486.1	1,009.3
2003 4/.....	46.6	*	*	*	*	2.3	13.8	53.0	133.3	283.0	600.2	1,113.5
2002.....	48.3	*	*	*	*	2.1	13.3	52.0	141.6	300.1	623.4	1,280.1
2001.....	49.2	*	*	*	*	2.5	13.2	51.9	143.8	317.1	638.0	1,382.0
2000.....	52.0	*	*	*	*	2.2	13.4	54.9	160.3	335.8	687.3	1,418.5
1999.....	53.4	*	*	*	*	2.3	14.2	56.6	165.5	353.6	685.6	1,498.7

Black, female

2006 4/.....	35.7	*	*	*	*	1.0	5.4	20.9	53.1	132.4	318.8	799.9
2005 4/.....	39.2	*	*	*	*	1.0	6.1	21.9	58.9	144.8	359.9	894.6
2004 4/.....	41.6	*	*	*	*	1.4	6.2	22.5	65.0	161.1	382.7	968.3
2003 4/.....	44.9	*	*	*	*	1.0	6.7	24.1	70.7	175.2	411.9	1,087.7
2002.....	48.8	*	*	*	*	1.2	6.5	25.9	79.1	195.3	457.2	1,171.0
2001.....	50.6	*	*	*	*	1.2	7.1	28.2	82.4	207.0	481.5	1,187.2
2000.....	52.5	*	*	*	*	1.3	6.8	29.1	87.7	221.6	505.6	1,222.5
1999.....	53.2	*	*	*	*	0.9	6.9	27.4	85.1	227.4	528.1	1,251.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute ischemic heart diseases (I24)

All races 3/, both sexes

2006.....	1.3	*	*	*	*	0.1	0.2	0.8	1.9	3.6	8.0	21.5
2005.....	1.2	*	*	*	*	*	0.2	0.8	1.7	3.3	7.2	20.8
2004.....	0.8	*	*	*	*	*	0.2	0.5	1.2	2.2	5.2	14.8
2003.....	1.1	*	*	*	*	*	0.2	0.7	1.4	3.2	6.5	20.0
2002.....	1.2	*	*	*	*	*	0.2	0.9	1.7	3.5	7.1	21.1
2001.....	1.2	*	*	*	*	*	0.2	0.8	1.8	3.5	6.9	21.5
2000.....	1.2	*	*	*	*	*	0.2	0.7	2.0	3.5	7.3	21.9
1999.....	1.2	*	*	*	*	*	0.2	0.7	1.9	3.7	7.0	21.9

All races 3/, male

2006.....	1.4	*	*	*	*	*	0.2	1.2	2.6	4.6	9.6	21.8
2005.....	1.2	*	*	*	*	*	0.3	1.2	2.5	4.3	8.5	20.2
2004.....	0.9	*	*	*	*	*	0.3	0.7	1.8	3.0	6.3	16.3
2003.....	1.1	*	*	*	*	*	0.3	1.0	2.1	4.3	8.0	19.5
2002.....	1.2	*	*	*	*	*	0.2	1.3	2.5	4.7	8.7	21.4
2001.....	1.2	*	*	*	*	*	0.3	1.2	2.6	4.7	8.5	20.7
2000.....	1.3	*	*	*	*	*	0.3	1.2	2.9	4.8	9.1	20.7
1999.....	1.2	*	*	*	*	*	0.3	1.0	2.8	4.9	8.8	22.3

All races 3/, female

2006.....	1.3	*	*	*	*	*	0.2	0.5	1.2	2.7	6.8	21.4
2005.....	1.2	*	*	*	*	*	0.1	0.5	0.9	2.4	6.2	21.1
2004.....	0.8	*	*	*	*	*	0.1	0.3	0.7	1.4	4.4	14.1
2003.....	1.0	*	*	*	*	*	0.1	0.4	0.9	2.2	5.5	20.2
2002.....	1.1	*	*	*	*	*	0.1	0.5	1.0	2.4	5.9	21.0
2001.....	1.1	*	*	*	*	*	0.1	0.4	1.1	2.5	5.9	21.8
2000.....	1.1	*	*	*	*	*	0.1	0.3	1.1	2.4	6.1	22.4
1999.....	1.1	*	*	*	*	*	*	0.4	1.1	2.6	5.8	21.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute ischemic heart diseases (I24)

White, both sexes

2006 4/.....	1.4	*	*	*	*	*	0.2	0.7	1.8	3.4	7.8	21.6
2005 4/.....	1.2	*	*	*	*	*	0.2	0.7	1.5	3.1	6.9	20.5
2004 4/.....	0.8	*	*	*	*	*	0.2	0.4	1.1	2.0	4.9	15.0
2003 4/.....	1.1	*	*	*	*	*	0.2	0.6	1.3	3.0	6.3	20.2
2002.....	1.2	*	*	*	*	*	0.2	0.8	1.6	3.3	7.0	21.4
2001.....	1.2	*	*	*	*	*	0.2	0.7	1.6	3.3	6.8	21.5
2000.....	1.2	*	*	*	*	*	0.2	0.7	1.8	3.3	7.1	22.1
1999.....	1.2	*	*	*	*	*	0.1	0.6	1.7	3.5	6.8	22.4

White, male

2006 4/.....	1.4	*	*	*	*	*	0.2	1.1	2.5	4.4	9.2	21.8
2005 4/.....	1.2	*	*	*	*	*	0.3	1.0	2.2	4.0	8.1	19.7
2004 4/.....	0.9	*	*	*	*	*	0.3	0.6	1.6	2.6	6.1	16.2
2003 4/.....	1.1	*	*	*	*	*	0.3	0.9	1.8	4.1	7.6	19.7
2002.....	1.3	*	*	*	*	*	0.2	1.1	2.3	4.4	8.6	21.9
2001.....	1.2	*	*	*	*	*	0.2	1.1	2.4	4.4	8.5	20.7
2000.....	1.3	*	*	*	*	*	0.3	1.1	2.6	4.4	8.9	21.0
1999.....	1.2	*	*	*	*	*	0.2	0.8	2.5	4.7	8.6	22.9

White, female

2006 4/.....	1.3	*	*	*	*	*	0.1	0.3	1.1	2.5	6.8	21.6
2005 4/.....	1.2	*	*	*	*	*	*	0.3	0.8	2.3	6.1	20.8
2004 4/.....	0.8	*	*	*	*	*	*	0.2	0.6	1.4	4.2	14.4
2003 4/.....	1.1	*	*	*	*	*	*	0.3	0.8	2.0	5.5	20.5
2002.....	1.2	*	*	*	*	*	*	0.4	0.9	2.3	5.9	21.1
2001.....	1.2	*	*	*	*	*	*	0.3	0.9	2.4	5.7	21.7
2000.....	1.2	*	*	*	*	*	*	0.3	0.9	2.3	5.9	22.6
1999.....	1.2	*	*	*	*	*	*	0.3	1.1	2.5	5.6	22.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute ischemic heart diseases (I24)

Black, both sexes

2006 4/.....	1.3	*	*	*	*	*	0.5	1.7	3.4	6.0	11.2	20.8
2005 4/.....	1.4	*	*	*	*	*	*	2.0	3.8	5.5	11.8	27.3
2004 4/.....	1.0	*	*	*	*	*	*	1.4	3.0	4.1	8.8	15.2
2003 4/.....	1.1	*	*	*	*	*	0.5	1.3	3.0	5.5	9.9	19.8
2002.....	1.2	*	*	*	*	*	0.3	1.7	3.0	5.9	9.3	21.1
2001.....	1.2	*	*	*	*	*	0.5	1.5	3.9	5.7	8.5	24.0
2000.....	1.2	*	*	*	*	*	0.4	1.5	4.2	5.6	10.8	21.9
1999.....	1.2	*	*	*	*	*	0.5	1.8	3.6	5.7	10.7	17.2

Black, male

2006 4/.....	1.5	*	*	*	*	*	*	2.3	5.1	8.0	15.2	23.2
2005 4/.....	1.6	*	*	*	*	*	*	2.4	5.8	7.9	15.8	27.9
2004 4/.....	1.3	*	*	*	*	*	*	1.9	4.6	7.1	11.7	22.8
2003 4/.....	1.4	*	*	*	*	*	*	1.9	4.8	7.6	16.1	24.0
2002.....	1.4	*	*	*	*	*	*	2.7	4.2	9.1	11.8	21.4
2001.....	1.4	*	*	*	*	*	*	2.4	5.5	8.5	10.3	23.8
2000.....	1.5	*	*	*	*	*	*	2.4	6.1	8.6	13.9	*
1999.....	1.4	*	*	*	*	*	0.7	2.6	6.1	8.2	12.6	*

Black, female

2006 4/.....	1.2	*	*	*	*	*	*	1.1	2.0	4.6	8.9	19.7
2005 4/.....	1.3	*	*	*	*	*	*	1.7	2.1	3.9	9.5	27.0
2004 4/.....	0.8	*	*	*	*	*	*	0.9	1.7	2.0	7.2	12.2
2003 4/.....	0.9	*	*	*	*	*	*	*	1.5	4.0	6.4	18.2
2002.....	1.0	*	*	*	*	*	*	0.8	2.0	3.6	8.0	20.9
2001.....	1.1	*	*	*	*	*	*	*	2.6	3.7	7.4	24.0
2000.....	1.0	*	*	*	*	*	*	*	2.7	3.5	9.0	22.3
1999.....	1.0	*	*	*	*	*	*	1.1	1.5	3.8	9.6	16.2



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/ 1 year	1-4 2/ years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	-----------------	--------------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other forms of chronic ischemic heart disease (I20,I25)

All races 3/, both sexes

2006.....	93.5	*	*	*	0.2	1.5	9.0	34.4	87.8	217.9	632.8	2,057.4
2005.....	98.2	*	*	*	0.2	1.5	9.2	35.0	91.4	229.8	674.5	2,214.2
2004.....	99.5	*	*	*	0.2	1.6	9.6	35.1	92.5	240.3	697.1	2,279.7
2003.....	105.3	*	*	*	0.2	1.7	9.8	35.9	97.5	258.4	741.5	2,471.4
2002.....	108.0	*	*	*	0.2	1.4	9.9	36.5	101.0	269.4	768.8	2,562.5
2001.....	110.3	*	*	*	0.2	1.6	9.5	35.6	103.8	280.9	793.1	2,671.3
2000.....	113.3	*	*	*	0.2	1.4	9.5	36.4	108.1	292.4	813.6	2,822.1
1999.....	117.2	*	*	*	0.2	1.4	9.8	37.5	111.5	308.0	848.5	2,917.0

All races 3/, male

2006.....	99.2	*	*	*	0.3	2.4	13.8	53.7	132.4	307.7	848.3	2,344.9
2005.....	102.9	*	*	*	0.3	2.3	14.1	54.7	136.9	321.9	897.1	2,511.6
2004.....	103.3	*	*	*	0.2	2.4	14.5	54.8	137.7	336.5	926.9	2,543.6
2003.....	108.5	*	*	*	0.3	2.5	14.9	56.4	145.4	363.5	985.8	2,722.2
2002.....	110.9	*	*	*	0.2	2.2	15.4	57.5	151.5	379.3	1,021.0	2,817.4
2001.....	111.9	*	*	*	0.3	2.4	14.6	55.9	154.9	394.3	1,050.9	2,913.8
2000.....	114.8	*	*	*	0.2	2.1	14.8	57.7	161.6	413.5	1,079.4	3,112.7
1999.....	118.0	*	*	*	0.3	2.1	15.3	59.0	164.7	434.9	1,120.5	3,193.1

All races 3/, female

2006.....	88.0	*	*	*	*	0.6	4.2	15.7	46.3	141.9	485.5	1,922.9
2005.....	93.7	*	*	*	*	0.7	4.3	16.0	49.0	152.2	523.3	2,077.6
2004.....	95.7	*	*	*	*	0.7	4.6	16.2	50.5	159.6	542.4	2,161.0
2003.....	102.3	*	*	*	0.1	0.8	4.7	16.2	53.1	170.5	578.3	2,360.5
2002.....	105.2	*	*	*	*	0.7	4.4	16.2	54.3	178.0	601.4	2,451.9
2001.....	108.8	*	*	*	0.1	0.7	4.4	16.0	56.7	186.9	623.7	2,569.8
2000.....	111.9	*	*	*	*	0.6	4.3	16.0	58.8	192.8	640.3	2,703.8
1999.....	116.4	*	*	*	0.1	0.7	4.3	16.7	62.6	204.0	671.8	2,805.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other forms of chronic ischemic heart disease (I20,I25)

White, both sexes

2006 4/.....	101.0	*	*	*	0.2	1.4	8.6	32.2	82.5	209.8	632.4	2,106.6
2005 4/.....	106.2	*	*	*	0.2	1.4	8.8	32.6	85.3	222.4	674.5	2,269.8
2004 4/.....	107.5	*	*	*	0.1	1.4	9.2	32.8	86.7	232.4	696.4	2,324.8
2003 4/.....	113.8	*	*	*	0.2	1.5	9.4	33.5	91.3	249.1	740.0	2,518.8
2002.....	116.9	*	*	*	0.2	1.4	9.4	34.0	94.4	261.3	767.5	2,613.9
2001.....	119.1	*	*	*	0.2	1.4	8.9	33.1	96.7	271.7	790.5	2,723.5
2000.....	122.6	*	*	*	0.2	1.2	8.9	33.9	100.6	283.4	811.2	2,882.1
1999.....	126.5	*	*	*	0.1	1.3	9.0	34.7	104.0	298.0	846.7	2,973.6

White, male

2006 4/.....	106.5	*	*	*	0.3	2.3	13.4	51.0	125.4	297.7	851.4	2,418.9
2005 4/.....	110.7	*	*	*	0.3	2.1	13.7	51.5	129.5	313.7	902.5	2,597.3
2004 4/.....	111.1	*	*	*	0.2	2.2	14.2	52.1	130.5	328.7	931.8	2,608.9
2003 4/.....	116.6	*	*	*	0.2	2.3	14.4	53.4	138.1	353.1	989.3	2,797.1
2002.....	119.4	*	*	*	0.2	2.1	14.9	54.8	143.1	371.2	1,027.4	2,891.6
2001.....	120.1	*	*	*	0.3	2.2	13.9	52.9	146.2	384.6	1,054.9	2,992.2
2000.....	123.5	*	*	*	0.2	1.9	14.2	54.8	152.7	405.6	1,083.4	3,200.0
1999.....	126.5	*	*	*	0.2	2.0	14.4	55.5	155.2	425.5	1,126.9	3,267.1

White, female

2006 4/.....	95.6	*	*	*	*	0.5	3.7	13.5	41.6	134.0	480.3	1,960.0
2005 4/.....	101.9	*	*	*	*	0.7	3.8	13.9	43.5	143.9	517.5	2,119.0
2004 4/.....	103.9	*	*	*	*	0.6	4.1	13.8	45.4	149.9	535.9	2,196.8
2003 4/.....	111.0	*	*	*	*	0.7	4.2	13.9	47.0	160.6	571.4	2,395.5
2002.....	114.4	*	*	*	*	0.6	3.9	13.6	48.5	168.1	593.1	2,493.2
2001.....	118.1	*	*	*	*	0.6	3.9	13.6	50.2	176.4	615.0	2,611.0
2000.....	121.6	*	*	*	*	0.5	3.5	13.4	51.8	180.9	632.0	2,753.0
1999.....	126.5	*	*	*	*	0.6	3.6	14.3	55.9	191.7	663.1	2,855.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other forms of chronic ischemic heart disease (I20,I25)

Black, both sexes

2006 4/.....	73.9	*	*	*	*	2.4	13.7	57.4	149.7	329.6	748.6	1,791.4
2005 4/.....	76.6	*	*	*	0.3	2.2	14.0	59.1	161.6	337.4	793.9	1,885.0
2004 4/.....	78.0	*	*	*	0.4	2.9	14.2	59.2	161.0	359.8	814.7	2,019.1
2003 4/.....	82.3	*	*	*	0.4	2.6	14.2	60.6	171.5	389.7	873.2	2,195.3
2002.....	83.2	*	*	*	*	2.2	15.1	61.9	178.1	391.4	896.1	2,240.6
2001.....	85.8	*	*	*	0.5	2.7	15.1	61.6	184.2	414.8	932.7	2,351.2
2000.....	86.6	*	*	*	*	2.5	15.3	63.1	190.2	427.8	941.3	2,387.1
1999.....	89.3	*	*	*	0.5	2.3	16.8	65.8	195.7	450.4	957.4	2,469.8

Black, male

2006 4/.....	79.6	*	*	*	*	3.6	19.0	84.7	220.4	468.4	986.9	1,819.4
2005 4/.....	81.0	*	*	*	*	3.4	20.1	88.9	234.6	465.6	1,017.8	1,871.2
2004 4/.....	81.5	*	*	*	*	4.3	19.8	86.2	234.3	489.0	1,039.8	2,068.3
2003 4/.....	85.5	*	*	*	*	3.9	20.6	89.6	243.7	536.7	1,128.5	2,147.8
2002.....	85.7	*	*	*	*	2.9	21.7	89.5	261.4	531.5	1,113.6	2,254.6
2001.....	88.2	*	*	*	0.7	4.2	22.2	91.3	263.6	568.3	1,177.1	2,369.1
2000.....	88.0	*	*	*	*	3.6	21.6	92.1	272.7	563.5	1,175.7	2,446.0
1999.....	91.1	*	*	*	*	3.3	24.6	97.7	281.3	600.5	1,172.6	2,561.5

Black, female

2006 4/.....	68.7	*	*	*	*	1.3	8.9	33.9	92.6	230.3	613.1	1,779.9
2005 4/.....	72.6	*	*	*	*	1.0	8.5	33.6	102.8	245.6	666.8	1,890.6
2004 4/.....	74.7	*	*	*	*	1.5	9.1	36.1	102.0	267.6	687.4	1,999.6
2003 4/.....	79.4	*	*	*	*	1.4	8.5	35.7	113.4	285.0	728.9	2,213.8
2002.....	81.0	*	*	*	*	1.5	9.4	38.2	111.1	291.9	772.9	2,235.2
2001.....	83.5	*	*	*	*	1.5	8.9	36.1	120.3	306.4	795.5	2,344.5
2000.....	85.3	*	*	*	*	1.5	9.7	38.1	123.7	332.3	809.4	2,365.2
1999.....	87.6	*	*	*	*	1.3	9.9	38.5	127.0	344.6	836.9	2,435.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

All races 3/, both sexes

2006.....	20.4	*	*	*	0.0	0.5	3.3	13.4	31.3	55.1	114.2	345.5
2005.....	21.2	*	*	*	*	0.5	3.2	13.9	31.5	55.0	124.8	376.4
2004.....	21.7	*	*	*	*	0.5	3.3	13.6	32.0	58.7	129.0	400.4
2003.....	23.1	*	*	*	*	0.5	3.4	13.9	32.7	62.7	142.5	442.9
2002.....	23.6	*	*	*	*	0.5	3.4	13.4	33.1	64.7	146.2	474.5
2001.....	23.9	*	*	*	0.1	0.5	3.2	12.9	33.3	65.6	152.7	493.6
2000.....	24.7	*	*	*	*	0.4	3.0	13.2	34.0	68.3	158.1	534.6
1999.....	25.8	*	*	*	*	0.4	3.3	13.0	35.3	72.7	166.2	563.4

All races 3/, male

2006.....	23.2	*	*	*	*	0.8	5.1	21.7	48.8	79.1	146.8	365.3
2005.....	23.6	*	*	*	*	0.7	5.1	22.2	49.0	78.7	158.8	390.7
2004.....	23.9	*	*	*	*	0.8	5.2	22.0	49.0	83.6	164.9	414.0
2003.....	25.2	*	*	*	*	0.9	5.2	22.7	49.9	88.9	181.1	462.1
2002.....	25.4	*	*	*	*	0.7	5.4	21.5	51.2	91.8	185.3	491.8
2001.....	25.3	*	*	*	*	0.7	5.0	21.0	51.4	92.4	193.9	502.0
2000.....	26.0	*	*	*	*	0.6	4.9	21.5	52.6	97.2	199.3	550.8
1999.....	26.8	*	*	*	*	0.6	5.3	20.9	53.7	103.0	207.0	578.6

All races 3/, female

2006.....	17.7	*	*	*	*	0.2	1.5	5.5	15.1	34.8	92.0	336.1
2005.....	18.8	*	*	*	*	0.2	1.4	5.7	15.3	35.0	101.6	369.9
2004.....	19.5	*	*	*	*	0.2	1.4	5.5	16.1	37.7	104.8	394.3
2003.....	21.1	*	*	*	*	0.2	1.5	5.5	16.7	40.8	116.7	434.4
2002.....	21.9	*	*	*	*	0.2	1.3	5.5	16.2	42.1	120.2	467.1
2001.....	22.5	*	*	*	*	0.2	1.4	5.1	16.7	43.3	125.5	490.1
2000.....	23.5	*	*	*	*	0.1	1.2	5.2	16.8	44.6	131.3	528.0
1999.....	24.8	*	*	*	*	0.2	1.3	5.3	18.3	47.9	139.8	557.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

White, both sexes

2006 4/.....	20.7	*	*	*	*	0.4	3.1	12.2	28.3	49.9	108.3	343.2
2005 4/.....	21.5	*	*	*	*	0.4	3.0	12.6	27.9	50.0	118.2	373.8
2004 4/.....	22.1	*	*	*	*	0.4	3.2	12.3	28.7	53.2	122.2	395.6
2003 4/.....	23.6	*	*	*	*	0.5	3.2	12.6	29.0	56.5	135.5	439.3
2002.....	24.1	*	*	*	*	0.4	3.1	12.1	29.4	58.1	138.4	470.6
2001.....	24.3	*	*	*	*	0.4	3.0	11.6	28.8	58.3	144.5	489.7
2000.....	25.1	*	*	*	*	0.3	2.8	11.8	29.4	61.2	149.2	529.5
1999.....	26.1	*	*	*	*	0.3	3.0	11.3	30.4	64.8	157.0	557.5

White, male

2006 4/.....	23.2	*	*	*	*	0.7	4.8	19.9	44.2	71.4	138.9	362.7
2005 4/.....	23.7	*	*	*	*	0.6	4.9	20.5	43.8	71.4	150.1	392.6
2004 4/.....	24.1	*	*	*	*	0.7	5.1	20.4	44.1	76.6	156.3	408.8
2003 4/.....	25.3	*	*	*	*	0.8	5.0	20.8	44.8	80.0	171.9	458.7
2002.....	25.5	*	*	*	*	0.7	5.1	19.9	45.9	82.9	174.7	488.0
2001.....	25.3	*	*	*	*	0.6	4.7	19.2	45.1	82.7	184.2	498.6
2000.....	26.0	*	*	*	*	0.5	4.6	19.6	46.3	87.8	187.5	545.6
1999.....	26.6	*	*	*	*	0.5	4.9	18.6	46.9	92.5	195.2	571.2

White, female

2006 4/.....	18.3	*	*	*	*	*	1.2	4.5	13.2	31.4	87.1	334.1
2005 4/.....	19.4	*	*	*	*	0.2	1.1	4.7	12.8	31.5	96.3	365.1
2004 4/.....	20.1	*	*	*	*	0.2	1.3	4.3	14.1	33.1	99.0	389.7
2003 4/.....	21.9	*	*	*	*	0.2	1.4	4.5	14.1	36.5	110.8	430.7
2002.....	22.7	*	*	*	*	0.1	1.1	4.5	13.8	37.0	114.0	463.0
2001.....	23.3	*	*	*	*	0.2	1.2	4.1	13.4	37.8	118.2	486.0
2000.....	24.3	*	*	*	*	*	1.0	4.2	13.5	39.0	123.9	522.9
1999.....	25.6	*	*	*	*	0.2	1.0	4.2	15.0	41.6	131.9	552.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerotic cardiovascular disease, so described (I25.0)

Black, both sexes

2006 4/.....	23.7	*	*	*	*	1.0	5.6	25.0	62.6	110.2	199.1	428.1
2005 4/.....	24.5	*	*	*	*	0.8	5.2	25.4	67.6	109.2	216.5	469.6
2004 4/.....	25.0	*	*	*	*	0.9	4.8	25.7	66.7	120.0	226.9	505.1
2003 4/.....	26.0	*	*	*	*	0.9	4.9	25.7	69.7	128.1	241.6	543.2
2002.....	27.0	*	*	*	*	0.8	5.8	25.5	71.4	136.3	259.0	575.5
2001.....	27.7	*	*	*	*	0.7	5.6	25.3	77.5	143.4	266.5	594.5
2000.....	28.5	*	*	*	*	0.7	5.1	26.9	78.2	145.5	278.4	646.6
1999.....	29.9	*	*	*	*	0.7	6.3	27.5	81.4	158.6	291.2	668.6

Black, male

2006 4/.....	28.6	*	*	*	*	1.5	8.1	38.2	99.3	168.6	274.7	465.6
2005 4/.....	28.8	*	*	*	*	1.3	7.5	38.8	104.8	166.2	294.5	450.8
2004 4/.....	28.9	*	*	*	*	1.5	6.9	38.1	104.6	172.0	306.2	531.5
2003 4/.....	30.5	*	*	*	*	1.5	7.7	40.4	106.0	190.3	331.4	583.9
2002.....	31.1	*	*	*	*	1.1	8.6	38.7	111.4	197.2	357.2	599.3
2001.....	31.5	*	*	*	*	0.9	8.5	39.2	117.4	208.2	351.7	609.3
2000.....	32.3	*	*	*	*	1.0	7.8	42.0	120.0	207.3	369.7	701.2
1999.....	33.8	*	*	*	*	1.0	9.6	42.8	123.3	229.3	380.0	715.9

Black, female

2006 4/.....	19.1	*	*	*	*	*	3.3	13.6	33.0	68.4	156.2	412.7
2005 4/.....	20.6	*	*	*	*	*	3.1	13.8	37.6	68.3	172.2	477.1
2004 4/.....	21.5	*	*	*	*	*	3.0	15.1	36.2	83.0	182.0	494.6
2003 4/.....	21.9	*	*	*	*	*	2.6	13.1	40.5	83.9	190.9	527.4
2002.....	23.3	*	*	*	*	*	3.3	14.2	39.2	93.0	203.5	566.4
2001.....	24.3	*	*	*	*	*	3.1	13.3	45.4	97.7	218.6	589.0
2000.....	25.0	*	*	*	*	*	2.8	14.0	44.6	102.0	227.1	626.3
1999.....	26.4	*	*	*	*	*	3.4	14.4	47.8	108.8	241.5	650.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes

2006.....	73.1	*	*	*	0.1	1.0	5.7	21.0	56.5	162.8	518.6	1,712.0
2005.....	77.0	*	*	*	0.2	1.1	6.0	21.2	59.8	174.9	549.7	1,837.8
2004.....	77.8	*	*	*	0.1	1.1	6.2	21.5	60.5	181.7	568.1	1,879.3
2003.....	82.2	*	*	*	0.2	1.1	6.5	22.0	64.9	195.7	599.0	2,028.5
2002.....	84.4	*	*	*	0.1	1.0	6.5	23.1	68.0	204.7	622.6	2,088.0
2001.....	86.4	*	*	*	0.2	1.1	6.2	22.7	70.5	215.3	640.5	2,177.7
2000.....	88.6	*	*	*	0.1	1.0	6.5	23.2	74.1	224.1	655.5	2,287.5
1999.....	91.4	*	*	*	0.2	1.0	6.5	24.5	76.3	235.2	682.3	2,353.6

All races 3/, male

2006.....	76.1	*	*	*	0.2	1.6	8.7	32.1	83.7	228.7	701.5	1,979.5
2005.....	79.3	*	*	*	0.2	1.6	9.0	32.5	88.0	243.3	738.3	2,120.9
2004.....	79.4	*	*	*	0.2	1.7	9.3	32.8	88.7	252.9	762.0	2,129.6
2003.....	83.3	*	*	*	0.2	1.6	9.7	33.7	95.5	274.6	804.7	2,260.1
2002.....	85.5	*	*	*	0.2	1.5	10.0	35.9	100.3	287.5	835.7	2,325.6
2001.....	86.5	*	*	*	0.2	1.8	9.5	34.9	103.5	301.9	856.9	2,411.8
2000.....	88.8	*	*	*	0.2	1.5	9.9	36.2	109.0	316.3	880.1	2,561.9
1999.....	91.2	*	*	*	0.2	1.5	10.0	38.1	111.0	331.9	913.6	2,614.4

All races 3/, female

2006.....	70.3	*	*	*	*	0.4	2.7	10.2	31.2	107.1	393.5	1,586.8
2005.....	74.8	*	*	*	*	0.5	2.9	10.3	33.7	117.2	421.7	1,707.7
2004.....	76.2	*	*	*	*	0.5	3.2	10.6	34.3	121.9	437.6	1,766.7
2003.....	81.1	*	*	*	0.1	0.6	3.2	10.8	36.4	129.7	461.6	1,926.2
2002.....	83.3	*	*	*	*	0.5	3.1	10.7	38.0	135.9	481.1	1,984.9
2001.....	86.2	*	*	*	0.1	0.5	3.0	11.0	40.0	143.6	498.2	2,079.8
2000.....	88.4	*	*	*	*	0.5	3.0	10.7	42.0	148.2	509.0	2,175.8
1999.....	91.6	*	*	*	0.1	0.5	3.0	11.4	44.3	156.1	532.1	2,248.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

White, both sexes

2006 4/.....	80.3	*	*	*	0.1	1.0	5.5	20.0	54.2	159.9	524.0	1,763.4
2005 4/.....	84.7	*	*	*	0.1	1.0	5.8	20.0	57.5	172.4	556.3	1,896.0
2004 4/.....	85.4	*	*	*	0.1	1.0	6.0	20.5	58.1	179.1	574.2	1,929.2
2003 4/.....	90.2	*	*	*	0.1	1.0	6.2	20.9	62.2	192.6	604.6	2,079.5
2002.....	92.8	*	*	*	0.1	0.9	6.3	21.9	65.1	203.2	629.1	2,143.3
2001.....	94.8	*	*	*	0.1	1.0	5.9	21.5	67.9	213.4	646.0	2,233.8
2000.....	97.4	*	*	*	0.1	0.9	6.1	22.1	71.2	222.2	662.0	2,352.7
1999.....	100.4	*	*	*	0.1	1.0	6.1	23.4	73.5	233.2	689.7	2,416.1

White, male

2006 4/.....	83.4	*	*	*	0.2	1.5	8.6	31.1	81.3	226.2	712.6	2,056.2
2005 4/.....	87.0	*	*	*	0.2	1.5	8.8	31.0	85.7	242.3	752.4	2,204.7
2004 4/.....	87.0	*	*	*	0.2	1.5	9.1	31.6	86.4	252.0	775.6	2,200.2
2003 4/.....	91.3	*	*	*	0.2	1.5	9.4	32.5	93.3	273.1	817.4	2,338.4
2002.....	93.9	*	*	*	0.1	1.4	9.8	34.9	97.3	288.3	852.7	2,403.6
2001.....	94.8	*	*	*	0.2	1.6	9.2	33.7	101.1	301.9	870.7	2,493.6
2000.....	97.5	*	*	*	0.1	1.4	9.6	35.2	106.4	317.8	895.9	2,654.4
1999.....	99.9	*	*	*	0.2	1.4	9.5	36.9	108.3	333.0	931.7	2,695.9

White, female

2006 4/.....	77.3	*	*	*	*	0.4	2.4	9.0	28.4	102.6	393.1	1,626.0
2005 4/.....	82.4	*	*	*	*	0.5	2.7	9.2	30.7	112.4	421.2	1,753.8
2004 4/.....	83.8	*	*	*	*	0.5	2.9	9.4	31.3	116.7	436.9	1,807.0
2003 4/.....	89.1	*	*	*	*	0.5	2.9	9.4	32.9	124.0	460.7	1,964.9
2002.....	91.7	*	*	*	*	0.4	2.8	9.1	34.7	131.1	479.1	2,030.2
2001.....	94.9	*	*	*	*	0.4	2.6	9.6	36.7	138.6	496.8	2,125.0
2000.....	97.3	*	*	*	*	0.4	2.5	9.3	38.3	141.9	508.1	2,230.1
1999.....	100.8	*	*	*	*	0.5	2.6	10.1	40.9	150.1	531.1	2,303.8



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

Black, both sexes

2006 4/.....	50.2	*	*	*	*	1.4	8.1	32.4	87.0	219.4	549.4	1,363.3
2005 4/.....	52.1	*	*	*	*	1.4	8.8	33.8	94.0	228.2	577.4	1,415.4
2004 4/.....	52.9	*	*	*	0.3	2.0	9.3	33.5	94.3	239.8	587.8	1,514.0
2003 4/.....	56.3	*	*	*	0.3	1.8	9.3	34.9	101.8	261.5	631.6	1,652.1
2002.....	56.2	*	*	*	*	1.4	9.4	36.4	106.7	255.1	637.0	1,665.1
2001.....	58.0	*	*	*	0.4	2.0	9.5	36.3	106.7	271.4	666.3	1,756.7
2000.....	58.1	*	*	*	*	1.8	10.2	36.1	112.0	282.3	662.9	1,740.5
1999.....	59.3	*	*	*	0.3	1.5	10.4	38.3	114.2	291.8	666.2	1,801.2

Black, male

2006 4/.....	50.9	*	*	*	*	2.1	11.0	46.4	121.1	299.8	712.2	1,353.8
2005 4/.....	52.2	*	*	*	*	2.1	12.6	50.0	129.8	299.3	723.3	1,420.4
2004 4/.....	52.6	*	*	*	*	2.8	12.9	48.1	129.7	317.0	733.6	1,536.8
2003 4/.....	55.0	*	*	*	*	2.4	13.0	49.1	137.6	346.5	797.1	1,563.9
2002.....	54.5	*	*	*	*	1.8	13.1	50.8	150.1	334.3	756.5	1,655.3
2001.....	56.7	*	*	*	*	3.2	13.7	52.1	146.2	360.1	825.4	1,759.8
2000.....	55.8	*	*	*	*	2.6	13.9	50.2	152.7	356.1	806.0	1,744.8
1999.....	57.3	*	*	*	*	2.3	15.0	54.9	157.9	371.2	792.6	1,845.5

Black, female

2006 4/.....	49.5	*	*	*	*	0.7	5.6	20.4	59.6	161.8	456.9	1,367.2
2005 4/.....	52.0	*	*	*	*	0.8	5.4	19.8	65.2	177.3	494.6	1,413.4
2004 4/.....	53.2	*	*	*	*	1.2	6.2	21.0	65.8	184.6	505.4	1,505.0
2003 4/.....	57.5	*	*	*	*	1.2	6.0	22.6	72.9	201.1	538.0	1,686.4
2002.....	57.7	*	*	*	*	1.1	6.1	24.0	71.9	198.9	569.4	1,668.8
2001.....	59.2	*	*	*	*	1.0	5.9	22.8	75.0	208.7	576.9	1,755.5
2000.....	60.3	*	*	*	*	1.1	6.9	24.1	79.2	230.3	582.3	1,738.9
1999.....	61.2	*	*	*	*	0.8	6.4	24.1	79.1	235.8	595.4	1,784.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other heart diseases (I26-I51)

All races 3/, both sexes

2006.....	56.9	7.7	0.9	0.6	2.1	4.7	9.5	21.2	44.4	112.3	360.7	1,342.8
2005.....	57.6	8.3	0.9	0.6	2.2	4.6	9.8	21.5	45.7	116.8	366.7	1,377.4
2004.....	56.6	10.0	1.1	0.5	2.1	4.3	9.7	21.5	45.2	118.6	368.9	1,373.9
2003.....	58.5	10.4	1.1	0.6	2.2	4.5	10.3	21.4	48.0	126.8	382.8	1,427.2
2002.....	58.8	11.7	1.0	0.6	2.1	4.5	10.3	21.7	48.7	129.6	390.9	1,443.5
2001.....	58.6	11.1	1.4	0.6	2.0	4.4	9.9	21.6	49.5	131.4	389.5	1,475.5
2000.....	58.8	12.0	1.1	0.6	2.2	4.4	9.9	21.5	50.9	134.2	391.3	1,509.4
1999.....	59.4	12.9	1.2	0.6	2.3	4.4	10.3	21.6	52.9	138.8	398.5	1,512.9

All races 3/, male

2006.....	51.0	8.0	1.0	0.6	2.5	5.9	11.6	27.2	56.2	136.2	411.1	1,338.1
2005.....	51.4	8.9	0.9	0.6	2.8	5.8	12.3	27.6	57.5	140.5	415.8	1,377.8
2004.....	50.6	10.5	1.1	0.5	2.6	5.3	12.0	27.7	57.7	142.8	420.8	1,377.7
2003.....	52.3	11.4	1.0	0.7	2.6	5.3	12.7	27.2	60.7	154.2	440.0	1,431.3
2002.....	52.4	12.1	1.0	0.6	2.7	5.6	12.8	28.2	60.8	156.5	448.2	1,448.6
2001.....	51.8	10.8	1.5	0.6	2.4	5.2	12.4	27.6	61.7	158.7	452.0	1,473.4
2000.....	52.2	12.2	1.2	0.7	2.6	5.2	12.2	27.8	64.7	164.0	451.0	1,549.1
1999.....	53.1	13.1	1.3	0.7	2.7	5.3	12.9	27.7	67.5	172.7	459.7	1,548.9

All races 3/, female

2006.....	62.6	7.5	0.9	0.5	1.5	3.5	7.3	15.3	33.5	92.1	326.3	1,345.0
2005.....	63.5	7.7	0.8	0.6	1.5	3.4	7.3	15.5	34.6	96.7	333.3	1,377.3
2004.....	62.4	9.4	1.1	0.6	1.5	3.3	7.4	15.5	33.6	98.4	334.0	1,372.1
2003.....	64.5	9.3	1.2	0.5	1.7	3.7	7.9	15.7	36.3	103.8	344.5	1,425.4
2002.....	65.0	11.4	1.0	0.5	1.4	3.4	7.8	15.5	37.5	107.2	352.8	1,441.3
2001.....	65.2	11.5	1.3	0.6	1.5	3.6	7.5	15.9	38.3	108.7	348.4	1,476.4
2000.....	65.1	11.8	0.9	0.5	1.8	3.5	7.5	15.3	38.2	109.6	352.3	1,493.2
1999.....	65.5	12.7	1.0	0.6	1.8	3.6	7.7	15.6	39.4	111.1	358.7	1,498.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other heart diseases (I26-I51)

White, both sexes

2006 4/.....	60.9	6.0	0.8	0.5	1.8	3.8	8.0	18.2	40.0	106.8	361.8	1,386.4
2005 4/.....	61.4	6.9	0.7	0.4	1.9	3.7	8.3	18.2	40.7	111.0	367.0	1,419.0
2004 4/.....	60.4	8.7	0.9	0.5	1.8	3.5	8.1	18.2	40.6	113.4	368.6	1,408.8
2003 4/.....	62.2	8.2	0.9	0.5	1.8	3.6	8.6	18.1	43.1	120.3	383.4	1,461.4
2002.....	62.6	10.0	0.9	0.5	1.7	3.6	8.7	18.4	43.9	124.2	390.3	1,475.5
2001.....	62.3	9.3	1.1	0.5	1.6	3.6	8.1	18.4	44.6	125.6	387.6	1,509.6
2000.....	62.5	10.6	1.0	0.5	1.7	3.6	8.1	18.2	46.1	128.1	389.6	1,541.6
1999.....	62.9	11.0	1.0	0.5	1.9	3.6	8.4	18.1	47.2	132.2	395.1	1,543.3

White, male

2006 4/.....	53.6	6.1	0.9	0.6	2.3	4.8	9.8	23.4	50.7	129.4	412.9	1,381.2
2005 4/.....	54.0	7.2	0.8	0.5	2.4	4.7	10.5	23.9	51.6	133.3	417.1	1,424.2
2004 4/.....	53.0	9.1	0.9	0.5	2.2	4.3	10.3	23.9	52.1	137.0	419.2	1,407.9
2003 4/.....	54.7	8.8	0.8	0.6	2.2	4.4	10.8	23.3	54.7	147.2	441.1	1,467.6
2002.....	54.9	10.5	1.0	0.5	2.3	4.5	11.1	24.3	55.3	150.4	448.3	1,482.1
2001.....	54.1	9.6	1.2	0.5	2.0	4.2	10.4	24.0	55.9	152.2	448.7	1,503.4
2000.....	54.6	10.8	1.2	0.6	2.1	4.3	10.3	23.9	59.2	157.4	449.5	1,580.5
1999.....	55.1	11.4	1.1	0.6	2.2	4.3	10.9	23.7	60.6	165.1	456.1	1,571.5

White, female

2006 4/.....	68.1	5.9	0.8	0.4	1.2	2.7	6.1	13.0	29.8	87.2	326.4	1,388.9
2005 4/.....	68.7	6.5	0.6	0.4	1.3	2.6	5.9	12.6	30.3	91.8	332.5	1,416.6
2004 4/.....	67.6	8.3	0.8	0.5	1.3	2.6	5.9	12.7	29.8	93.1	334.0	1,409.2
2003 4/.....	69.6	7.6	1.0	0.4	1.5	2.8	6.3	13.0	32.1	97.3	344.4	1,458.6
2002.....	70.2	9.6	0.8	0.5	1.1	2.6	6.4	12.6	33.1	101.9	351.4	1,472.6
2001.....	70.4	8.9	1.1	0.5	1.3	2.9	5.7	12.8	34.0	103.1	347.1	1,512.2
2000.....	70.2	10.4	0.7	0.4	1.4	2.8	6.0	12.5	33.9	103.4	350.1	1,525.8
1999.....	70.5	10.7	0.8	0.5	1.6	2.8	6.0	12.6	34.5	104.7	355.1	1,531.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other heart diseases (I26-I51)

Black, both sexes

2006 4/.....	50.1	16.0	1.6	0.9	3.8	11.3	21.4	47.1	92.6	188.4	435.3	1,094.3
2005 4/.....	51.4	15.2	1.7	1.1	4.1	11.2	22.0	49.3	97.7	195.3	447.8	1,135.3
2004 4/.....	50.8	14.8	2.3	1.1	3.7	10.0	22.2	50.0	96.1	195.5	450.2	1,180.7
2003 4/.....	52.5	19.6	2.0	1.0	4.4	10.9	23.6	49.9	102.1	215.7	449.2	1,238.1
2002.....	52.5	20.2	1.5	0.8	4.1	11.3	22.8	51.4	102.0	207.6	469.1	1,270.4
2001.....	52.8	20.1	2.9	1.0	4.2	10.5	23.7	51.1	103.5	213.0	480.6	1,277.7
2000.....	52.7	19.0	1.9	1.1	4.7	10.3	23.0	51.6	102.6	220.2	477.9	1,306.0
1999.....	54.4	22.3	2.0	1.0	4.5	10.5	24.7	52.4	113.3	229.9	495.2	1,302.8

Black, male

2006 4/.....	50.1	17.0	1.9	1.0	4.3	14.2	27.1	61.9	120.1	237.9	502.5	1,101.8
2005 4/.....	50.6	17.7	*	1.1	5.5	14.0	27.4	62.5	125.1	245.5	511.9	1,088.5
2004 4/.....	50.4	15.7	2.3	1.0	4.7	12.2	26.7	63.6	125.6	237.6	540.9	1,224.9
2003 4/.....	51.5	23.2	2.2	1.1	5.7	12.4	28.6	63.8	132.3	264.0	523.5	1,220.9
2002.....	50.8	20.9	*	0.9	5.1	14.2	27.4	65.9	127.9	255.2	537.3	1,238.3
2001.....	51.7	17.7	3.2	0.9	5.3	12.8	28.3	62.5	132.1	258.5	585.9	1,349.1
2000.....	51.5	18.9	1.9	1.2	5.4	12.2	28.1	66.5	130.5	266.0	558.0	1,402.3
1999.....	54.3	21.8	2.1	1.1	5.8	12.7	30.1	67.3	147.3	287.8	580.2	1,454.1

Black, female

2006 4/.....	50.0	15.0	*	0.8	3.2	8.6	16.4	34.4	70.5	152.9	397.0	1,091.2
2005 4/.....	52.2	12.6	2.0	1.2	2.7	8.7	17.3	38.0	75.6	159.3	411.4	1,154.2
2004 4/.....	51.1	13.9	2.3	1.1	2.7	8.0	18.3	38.4	72.3	165.4	398.9	1,163.2
2003 4/.....	53.4	15.8	1.8	0.8	3.2	9.4	19.1	38.0	77.8	181.3	407.1	1,244.8
2002.....	54.0	19.4	1.8	0.8	3.1	8.7	18.7	39.0	81.1	173.8	430.6	1,282.7
2001.....	53.8	22.7	2.5	1.0	3.0	8.5	19.6	41.3	80.5	180.9	421.5	1,251.2
2000.....	53.9	19.2	2.0	1.0	4.1	8.7	18.6	38.7	80.1	187.9	432.8	1,270.2
1999.....	54.5	22.8	1.8	0.8	3.3	8.5	19.9	39.7	86.1	189.1	447.7	1,246.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute and subacute endocarditis (I33)

All races 3/, both sexes

2006.....	0.4	*	*	*	*	0.1	0.2	0.5	0.7	1.2	2.4	2.2
2005.....	0.4	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.5	3.0
2004.....	0.4	*	*	*	*	0.1	0.2	0.5	0.6	1.3	2.5	3.2
2003.....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.5	3.0
2002.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.3	2.4	2.6
2001.....	0.4	*	*	*	*	0.1	0.3	0.5	0.6	1.3	2.5	2.5
2000.....	0.4	*	*	*	*	0.1	0.3	0.5	0.7	1.2	2.4	2.8
1999.....	0.4	*	*	*	*	0.1	0.3	0.4	0.7	1.3	2.4	2.9

All races 3/, male

2006.....	0.4	*	*	*	*	*	0.2	0.7	0.9	1.4	2.7	2.3
2005.....	0.5	*	*	*	*	0.1	0.2	0.5	0.9	1.5	3.3	5.1
2004.....	0.5	*	*	*	*	0.1	0.2	0.6	0.7	1.4	3.1	4.8
2003.....	0.5	*	*	*	*	*	0.3	0.6	0.9	1.4	3.2	4.1
2002.....	0.5	*	*	*	*	0.2	0.3	0.6	0.7	1.6	3.0	3.5
2001.....	0.5	*	*	*	*	*	0.3	0.7	0.8	1.6	3.5	3.8
2000.....	0.5	*	*	*	*	0.1	0.3	0.6	0.8	1.5	3.5	4.2
1999.....	0.5	*	*	*	*	0.1	0.3	0.6	0.9	1.4	3.2	4.7

All races 3/, female

2006.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	2.2	2.1
2005.....	0.3	*	*	*	*	*	0.1	0.3	0.5	1.1	2.0	2.1
2004.....	0.4	*	*	*	*	*	0.1	0.3	0.4	1.1	2.0	2.5
2003.....	0.4	*	*	*	*	*	0.2	0.3	0.6	1.1	2.0	2.5
2002.....	0.3	*	*	*	*	*	0.1	0.3	0.6	1.1	2.0	2.2
2001.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.1	1.8	2.0
2000.....	0.4	*	*	*	*	*	0.2	0.3	0.5	1.0	1.8	2.3
1999.....	0.3	*	*	*	*	*	0.2	0.2	0.5	1.2	1.8	2.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Acute and subacute endocarditis (I33)

White, both sexes

2006 4/.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.1	2.4	2.1
2005 4/.....	0.4	*	*	*	*	0.1	0.1	0.3	0.5	1.2	2.5	3.1
2004 4/.....	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.4	3.2
2003 4/.....	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.5	3.1
2002.....	0.4	*	*	*	*	0.1	0.2	0.3	0.5	1.2	2.5	2.7
2001.....	0.4	*	*	*	*	0.1	0.2	0.4	0.5	1.2	2.4	2.6
2000.....	0.4	*	*	*	*	0.1	0.2	0.4	0.6	1.2	2.4	2.7
1999.....	0.4	*	*	*	*	0.1	0.2	0.3	0.6	1.2	2.4	3.0

White, male

2006 4/.....	0.4	*	*	*	*	*	0.2	0.5	0.8	1.4	2.7	2.5
2005 4/.....	0.5	*	*	*	*	*	0.2	0.4	0.7	1.5	3.4	5.3
2004 4/.....	0.5	*	*	*	*	*	0.3	0.4	0.7	1.4	3.1	4.9
2003 4/.....	0.5	*	*	*	*	*	0.3	0.5	0.8	1.3	3.3	4.4
2002.....	0.5	*	*	*	*	0.2	0.2	0.5	0.6	1.5	3.2	3.6
2001.....	0.5	*	*	*	*	*	0.2	0.5	0.7	1.3	3.5	3.9
2000.....	0.5	*	*	*	*	*	0.3	0.5	0.8	1.4	3.5	4.2
1999.....	0.5	*	*	*	*	*	0.2	0.4	0.7	1.4	3.4	5.1

White, female

2006 4/.....	0.4	*	*	*	*	*	0.2	0.2	0.4	0.9	2.2	1.9
2005 4/.....	0.3	*	*	*	*	*	*	0.2	0.4	1.0	1.8	2.1
2004 4/.....	0.3	*	*	*	*	*	*	0.2	0.4	1.1	1.9	2.4
2003 4/.....	0.4	*	*	*	*	*	0.2	0.2	0.5	1.0	1.9	2.6
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.5	1.0	2.0	2.3
2001.....	0.3	*	*	*	*	*	0.2	0.2	0.4	1.1	1.7	2.1
2000.....	0.3	*	*	*	*	*	0.2	0.2	0.4	1.0	1.7	2.1
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.4	1.1	1.8	2.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute and subacute endocarditis (I33)

Black, both sexes

2006 4/.....	0.6	*	*	*	*	*	0.4	1.1	1.9	1.7	3.4	*
2005 4/.....	0.6	*	*	*	*	*	0.5	1.1	1.9	1.8	3.6	*
2004 4/.....	0.5	*	*	*	*	*	*	1.4	1.2	2.1	2.7	*
2003 4/.....	0.6	*	*	*	*	*	0.6	1.3	1.9	2.2	3.0	*
2002.....	0.5	*	*	*	*	*	0.4	1.2	1.6	2.3	*	*
2001.....	0.6	*	*	*	*	*	0.7	1.3	1.6	2.6	4.1	*
2000.....	0.6	*	*	*	*	*	0.5	1.3	1.6	1.9	2.8	*
1999.....	0.6	*	*	*	*	*	0.7	1.2	2.0	2.6	*	*

Black, male

2006 4/.....	0.6	*	*	*	*	*	*	1.5	2.6	*	*	*
2005 4/.....	0.7	*	*	*	*	*	*	1.4	2.7	*	*	*
2004 4/.....	0.6	*	*	*	*	*	*	1.8	*	*	*	*
2003 4/.....	0.7	*	*	*	*	*	*	2.0	2.2	*	*	*
2002.....	0.6	*	*	*	*	*	*	1.5	1.9	2.9	*	*
2001.....	0.8	*	*	*	*	*	0.7	1.9	2.1	3.8	*	*
2000.....	0.6	*	*	*	*	*	0.8	1.8	*	*	*	*
1999.....	0.7	*	*	*	*	*	1.0	1.9	2.5	*	*	*

Black, female

2006 4/.....	0.5	*	*	*	*	*	*	*	1.4	*	3.1	*
2005 4/.....	0.5	*	*	*	*	*	*	0.8	1.2	*	3.9	*
2004 4/.....	0.5	*	*	*	*	*	*	1.0	*	2.0	*	*
2003 4/.....	0.5	*	*	*	*	*	*	*	1.7	2.0	*	*
2002.....	0.5	*	*	*	*	*	*	1.0	*	*	*	*
2001.....	0.5	*	*	*	*	*	0.7	*	*	*	3.3	*
2000.....	0.5	*	*	*	*	*	*	1.0	1.6	*	*	*
1999.....	0.5	*	*	*	*	*	*	*	1.6	2.8	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes

2006.....	0.3	*	0.1	0.1	0.1	0.1	0.2	0.2	0.4	0.6	1.2	1.8
2005.....	0.3	0.5	*	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2004.....	0.3	0.5	0.2	*	0.1	0.1	0.2	0.3	0.4	0.6	1.2	1.8
2003.....	0.3	0.8	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	1.6
2002.....	0.3	0.6	0.1	0.1	0.1	0.1	0.2	0.3	0.4	0.8	1.2	2.1
2001.....	0.3	0.8	0.2	0.1	0.1	0.1	0.2	0.3	0.4	0.7	1.2	1.9
2000.....	0.3	0.5	0.2	0.0	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.8
1999.....	0.3	0.7	0.2	0.1	0.1	0.1	0.3	0.3	0.5	0.8	1.3	2.0

All races 3/, male

2006.....	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.7	1.3	1.8
2005.....	0.3	*	*	*	0.1	0.2	0.2	0.4	0.4	0.8	1.5	1.9
2004.....	0.3	*	*	*	*	0.2	0.3	0.5	0.7	1.3	2.3	
2003.....	0.3	*	*	*	0.2	0.1	0.2	0.4	0.4	1.0	1.3	1.5
2002.....	0.3	*	*	*	0.1	0.2	0.2	0.4	0.5	0.9	1.2	2.4
2001.....	0.3	*	*	*	0.1	0.2	0.2	0.3	0.5	0.7	1.2	1.8
2000.....	0.3	*	*	*	0.1	0.1	0.3	0.2	0.4	0.8	1.5	2.1
1999.....	0.3	*	*	*	0.1	0.2	0.3	0.3	0.6	0.8	1.4	2.5

All races 3/, female

2006.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.5	1.0	1.8
2005.....	0.3	*	*	0.1	*	0.1	0.2	0.3	0.3	0.6	1.0	1.9
2004.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.5	1.2	1.6
2003.....	0.3	*	*	*	0.1	*	0.2	0.2	0.3	0.7	1.1	1.7
2002.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.7	1.3	2.0
2001.....	0.3	*	*	*	*	*	0.2	0.3	0.3	0.6	1.2	2.0
2000.....	0.3	*	*	*	*	0.1	0.1	0.2	0.3	0.6	1.0	1.7
1999.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.8	1.3	1.9



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of pericardium and acute myocarditis (I30-I31,I40)

White, both sexes

2006 4/.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.6	1.2	1.8
2005 4/.....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.3	0.7	1.1	1.9
2004 4/.....	0.2	*	*	*	*	0.1	0.1	0.2	0.4	0.6	1.2	1.8
2003 4/.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.2	1.7
2002.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.8	1.1	2.2
2001.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.4	0.7	1.2	1.9
2000.....	0.3	*	*	*	0.1	0.1	0.2	0.2	0.3	0.7	1.2	1.8
1999.....	0.3	*	*	*	0.1	0.1	0.2	0.3	0.5	0.8	1.3	2.0

White, male

2006 4/.....	0.3	*	*	*	*	0.2	0.2	0.3	0.5	0.7	1.4	1.8
2005 4/.....	0.3	*	*	*	*	*	0.2	0.4	0.3	0.8	1.4	1.9
2004 4/.....	0.3	*	*	*	*	*	0.1	0.3	0.5	0.7	1.3	2.2
2003 4/.....	0.3	*	*	*	0.1	*	0.2	0.3	0.4	1.1	1.3	1.7
2002.....	0.3	*	*	*	*	0.1	0.2	0.3	0.5	0.9	1.1	2.5
2001.....	0.3	*	*	*	*	0.2	0.2	0.3	0.5	0.7	1.2	2.0
2000.....	0.3	*	*	*	*	*	0.2	0.2	0.4	0.7	1.5	2.1
1999.....	0.3	*	*	*	*	0.1	0.3	0.3	0.6	0.8	1.4	2.5

White, female

2006 4/.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.5	1.0	1.9
2005 4/.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	0.9	1.8
2004 4/.....	0.2	*	*	*	*	*	*	0.2	0.3	0.6	1.1	1.6
2003 4/.....	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.1	1.7
2002.....	0.3	*	*	*	*	*	*	0.2	0.3	0.7	1.2	2.0
2001.....	0.3	*	*	*	*	*	0.2	0.2	0.3	0.6	1.3	1.9
2000.....	0.2	*	*	*	*	*	*	0.2	0.3	0.6	1.0	1.6
1999.....	0.3	*	*	*	*	*	0.2	0.3	0.4	0.7	1.2	1.8



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Heart failure (I50)

All races 3/, both sexes

2006.....	20.2	*	*	*	0.1	0.2	0.8	2.8	8.9	31.5	132.7	614.4
2005.....	19.9	0.7	*	*	0.1	0.2	0.7	2.6	8.6	31.6	132.8	618.6
2004.....	19.5	0.7	*	*	0.1	0.2	0.7	2.5	8.4	32.2	131.2	621.2
2003.....	19.8	0.9	*	*	0.1	0.2	0.7	2.4	8.8	33.5	133.2	641.1
2002.....	19.6	0.6	*	*	0.1	0.3	0.6	2.5	8.8	33.8	132.8	645.0
2001.....	20.0	0.6	*	*	0.1	0.2	0.7	2.4	9.4	34.3	136.3	674.6
2000.....	19.8	0.8	*	0.1	0.1	0.2	0.6	2.2	9.1	34.8	136.1	684.9
1999.....	19.7	*	*	*	0.1	0.2	0.6	2.1	8.9	35.4	137.2	681.9

All races 3/, male

2006.....	16.2	*	*	*	0.1	0.3	0.9	3.7	10.8	36.9	148.6	599.0
2005.....	15.8	*	*	*	0.1	0.3	0.9	3.1	10.3	36.0	147.8	603.1
2004.....	15.4	*	*	*	*	0.3	0.8	3.0	10.1	37.5	147.0	605.5
2003.....	15.7	1.4	*	*	0.1	0.3	0.9	2.9	10.8	39.7	148.6	629.8
2002.....	15.3	*	*	*	*	0.3	0.8	3.1	10.5	39.2	149.0	621.4
2001.....	15.5	*	*	*	*	0.3	0.8	3.1	11.1	40.3	153.8	646.4
2000.....	15.3	1.1	*	*	0.1	0.2	0.7	2.7	10.9	41.1	153.7	672.2
1999.....	15.2	*	*	*	0.1	0.2	0.7	2.7	10.8	42.5	152.0	667.0

All races 3/, female

2006.....	24.0	*	*	*	*	0.2	0.6	2.0	7.0	26.9	121.8	621.6
2005.....	23.9	*	*	*	*	0.2	0.5	2.0	7.0	27.9	122.5	625.6
2004.....	23.4	*	*	*	*	0.2	0.6	2.0	6.8	27.8	120.5	628.2
2003.....	23.7	*	*	*	*	0.2	0.5	1.9	7.0	28.3	122.9	646.1
2002.....	23.7	*	*	*	*	0.2	0.5	1.8	7.2	29.2	122.0	655.3
2001.....	24.3	*	*	*	*	0.2	0.6	1.8	7.8	29.3	124.7	686.4
2000.....	24.1	*	*	*	*	0.1	0.5	1.6	7.5	29.6	124.6	690.0
1999.....	24.0	*	*	*	*	0.2	0.5	1.6	7.1	29.6	127.6	687.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Heart failure (I50)

White, both sexes

2006 4/.....	22.3	*	*	*	*	0.1	0.5	2.1	7.9	30.4	134.6	637.2
2005 4/.....	21.9	*	*	*	*	0.1	0.5	2.0	7.5	30.2	133.3	640.6
2004 4/.....	21.5	0.6	*	*	*	0.1	0.2	1.9	7.4	31.4	132.4	641.5
2003 4/.....	21.8	0.6	*	*	*	0.1	0.2	1.9	7.8	32.2	134.3	660.4
2002.....	21.6	*	*	*	*	0.2	0.5	1.9	8.0	32.5	133.3	664.5
2001.....	22.1	*	*	*	*	0.2	0.5	2.0	8.5	33.0	137.0	695.1
2000.....	21.8	*	*	*	*	0.1	0.4	1.8	8.2	33.4	136.7	703.7
1999.....	21.6	*	*	*	*	0.1	0.5	1.7	8.0	34.1	136.7	699.2

White, male

2006 4/.....	17.7	*	*	*	*	0.2	0.6	2.8	9.8	35.5	151.2	622.6
2005 4/.....	17.2	*	*	*	*	0.2	0.6	2.4	9.0	34.3	148.9	629.3
2004 4/.....	16.8	*	*	*	*	0.2	0.7	2.3	8.8	36.8	148.2	625.5
2003 4/.....	17.1	*	*	*	*	0.2	0.6	2.3	9.6	38.6	150.4	650.8
2002.....	16.7	*	*	*	*	0.2	0.6	2.5	9.5	37.7	150.3	641.9
2001.....	16.9	*	*	*	*	0.2	0.6	2.6	10.3	39.0	154.6	666.5
2000.....	16.7	*	*	*	*	0.1	0.6	2.2	10.0	40.0	154.6	690.0
1999.....	16.4	*	*	*	*	*	0.6	2.1	9.8	41.3	152.1	680.0

White, female

2006 4/.....	26.8	*	*	*	*	*	0.5	1.5	6.1	25.9	123.1	644.0
2005 4/.....	26.5	*	*	*	*	*	0.4	1.5	6.1	26.6	122.6	645.8
2004 4/.....	26.1	*	*	*	*	0.1	0.4	1.5	6.0	26.8	121.7	648.7
2003 4/.....	26.3	*	*	*	*	0.1	0.3	1.4	6.1	26.8	123.4	664.7
2002.....	26.4	*	*	*	*	0.1	0.3	1.4	6.5	28.1	121.8	674.3
2001.....	27.1	*	*	*	*	0.2	0.4	1.4	6.9	28.0	125.3	707.1
2000.....	26.7	*	*	*	*	*	0.3	1.3	6.6	27.9	124.9	709.3
1999.....	26.6	*	*	*	*	*	0.3	1.2	6.3	28.0	126.6	707.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Heart failure (I50)

Black, both sexes

2006 4/.....	14.1	*	*	*	*	0.8	2.2	8.4	19.5	51.6	147.0	484.9
2005 4/.....	14.4	*	*	*	*	0.8	2.3	7.3	20.2	53.6	161.4	489.6
2004 4/.....	13.5	*	*	*	*	0.7	1.9	7.5	19.4	48.2	150.4	506.5
2003 4/.....	14.0	*	*	*	*	0.7	2.3	6.8	20.6	54.3	150.4	531.8
2002.....	13.7	*	*	*	*	0.8	1.8	7.2	18.6	53.2	156.9	528.0
2001.....	13.8	*	*	*	*	0.6	1.9	6.4	19.6	54.5	157.3	543.8
2000.....	13.6	*	*	*	*	0.4	1.6	5.7	18.6	55.4	154.4	558.4
1999.....	13.8	*	*	*	*	0.5	1.6	6.4	18.9	56.4	163.4	552.9

Black, male

2006 4/.....	12.4	*	*	*	*	1.0	2.9	11.1	23.4	63.3	164.8	463.8
2005 4/.....	12.2	*	*	*	*	1.1	3.3	9.2	25.2	63.1	180.0	423.0
2004 4/.....	11.8	*	*	*	*	0.9	2.2	9.3	25.0	55.7	176.2	503.7
2003 4/.....	11.8	*	*	*	*	0.8	2.7	8.3	26.3	63.2	163.9	513.0
2002.....	11.5	*	*	*	*	1.0	2.2	9.0	22.9	64.7	174.4	479.2
2001.....	11.6	*	*	*	*	0.9	2.2	7.4	22.2	65.3	183.3	541.2
2000.....	11.2	*	*	*	*	*	1.7	7.1	21.4	61.9	173.5	581.2
1999.....	11.8	*	*	*	*	*	1.9	8.4	22.9	65.2	179.2	603.5

Black, female

2006 4/.....	15.6	*	*	*	*	*	1.7	6.0	16.3	43.2	137.0	493.5
2005 4/.....	16.3	*	*	*	*	*	1.4	5.7	16.2	46.7	150.8	516.5
2004 4/.....	15.1	*	*	*	*	*	1.7	5.9	15.0	42.8	135.8	507.6
2003 4/.....	15.9	*	*	*	*	*	1.9	5.6	16.0	48.0	142.7	539.2
2002.....	15.8	*	*	*	*	*	1.5	5.6	15.1	45.1	147.0	546.7
2001.....	15.8	*	*	*	*	*	1.7	5.5	17.6	46.9	142.6	544.8
2000.....	15.7	*	*	*	*	*	1.5	4.5	16.4	50.9	143.7	549.9
1999.....	15.7	*	*	*	*	*	1.4	4.7	15.7	50.2	154.6	534.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All races 3/, both sexes

2006.....	36.0	6.9	0.7	0.5	1.9	4.3	8.3	17.6	34.4	79.0	224.5	724.4
2005.....	37.0	7.1	0.7	0.5	2.0	4.2	8.7	18.2	36.0	83.2	230.2	754.0
2004.....	36.5	8.7	0.9	0.5	1.9	3.9	8.7	18.3	35.8	84.5	234.1	747.7
2003.....	38.0	8.6	0.9	0.5	1.9	4.1	9.1	18.2	38.1	91.2	246.0	781.5
2002.....	38.5	10.5	0.8	0.5	1.9	4.0	9.3	18.6	38.8	93.7	254.4	793.7
2001.....	37.9	9.7	1.1	0.5	1.8	4.0	8.8	18.5	39.1	95.1	249.5	796.5
2000.....	38.3	10.6	0.9	0.5	2.0	4.0	8.8	18.6	40.8	97.4	251.6	820.0
1999.....	39.0	11.9	0.9	0.5	2.1	4.1	9.2	18.7	42.8	101.3	257.6	826.1

All races 3/, male

2006.....	34.0	6.9	0.8	0.6	2.3	5.4	10.3	22.6	43.9	97.1	258.4	735.0
2005.....	34.9	7.7	0.8	0.5	2.6	5.2	10.9	23.6	46.0	102.3	263.3	767.7
2004.....	34.5	9.2	0.9	0.5	2.4	4.8	10.7	23.8	46.3	103.1	269.4	765.2
2003.....	35.8	9.2	0.9	0.6	2.3	4.8	11.3	23.3	48.5	112.1	287.0	795.9
2002.....	36.3	10.9	0.8	0.5	2.5	5.0	11.5	24.1	49.0	114.8	294.9	821.3
2001.....	35.6	9.1	1.2	0.5	2.2	4.7	11.1	23.6	49.4	116.1	293.4	821.3
2000.....	36.1	10.5	1.0	0.6	2.3	4.8	10.9	24.3	52.6	120.7	292.4	870.7
1999.....	37.1	12.1	1.0	0.6	2.5	4.8	11.6	24.2	55.3	127.9	303.1	874.7

All races 3/, female

2006.....	38.0	6.8	0.7	0.4	1.4	3.1	6.4	12.8	25.6	63.7	201.3	719.5
2005.....	39.0	6.4	0.7	0.4	1.3	3.1	6.5	13.0	26.8	67.1	207.8	747.6
2004.....	38.5	8.2	0.8	0.5	1.3	2.9	6.6	13.0	26.0	68.9	210.3	739.9
2003.....	40.1	7.9	1.0	0.4	1.5	3.4	7.0	13.3	28.4	73.7	218.6	775.1
2002.....	40.6	10.0	0.8	0.4	1.3	3.1	7.1	13.2	29.3	76.2	227.6	781.7
2001.....	40.2	10.4	1.0	0.5	1.3	3.2	6.5	13.5	29.6	77.7	220.7	786.1
2000.....	40.4	10.7	0.7	0.4	1.6	3.2	6.8	13.1	29.9	78.3	225.0	799.3
1999.....	40.9	11.6	0.8	0.5	1.7	3.3	6.9	13.5	31.3	79.6	228.0	806.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

White, both sexes

2006 4/.....	37.9	5.4	0.6	0.4	1.6	3.5	7.1	15.4	31.1	74.7	223.6	745.3
2005 4/.....	38.9	6.0	0.6	0.4	1.7	3.3	7.5	15.6	32.3	78.9	230.1	773.4
2004 4/.....	38.2	7.7	0.7	0.4	1.6	3.1	7.3	15.8	32.4	80.1	232.6	762.4
2003 4/.....	39.8	7.1	0.7	0.4	1.6	3.3	7.7	15.6	34.3	86.0	245.4	796.1
2002.....	40.4	9.1	0.7	0.4	1.6	3.2	7.9	15.9	34.9	89.7	253.4	806.1
2001.....	39.6	8.2	0.9	0.4	1.5	3.2	7.2	15.7	35.2	90.7	246.9	809.9
2000.....	40.0	9.6	0.8	0.4	1.6	3.3	7.3	15.8	37.0	92.8	249.3	833.4
1999.....	40.6	10.3	0.8	0.5	1.8	3.3	7.6	15.9	38.1	96.1	254.7	839.0

White, male

2006 4/.....	35.1	5.4	0.6	0.5	2.1	4.4	8.8	19.8	39.6	91.8	257.6	754.3
2005 4/.....	36.0	6.4	0.7	0.4	2.2	4.3	9.6	20.7	41.6	96.7	263.5	787.7
2004 4/.....	35.5	8.1	0.7	0.4	2.0	3.9	9.2	20.9	42.1	98.2	266.7	775.3
2003 4/.....	36.9	7.4	0.7	0.5	1.9	4.0	9.7	20.2	43.9	106.3	286.1	810.7
2002.....	37.5	9.5	0.8	0.5	2.1	4.0	10.0	21.0	44.6	110.3	293.7	834.2
2001.....	36.5	8.2	0.9	0.5	1.8	3.8	9.4	20.6	44.6	111.2	289.4	831.0
2000.....	37.1	9.4	0.9	0.5	1.9	4.0	9.2	20.9	48.0	115.3	290.0	884.2
1999.....	37.9	10.7	1.0	0.5	2.0	4.0	9.8	20.9	49.4	121.6	299.2	883.9

White, female

2006 4/.....	40.7	5.4	0.6	0.4	1.1	2.5	5.3	11.1	23.0	59.8	200.1	741.1
2005 4/.....	41.6	5.6	0.6	0.3	1.2	2.3	5.3	10.6	23.6	63.6	207.1	766.9
2004 4/.....	40.9	7.3	0.6	0.4	1.1	2.3	5.3	10.7	23.1	64.6	209.3	756.5
2003 4/.....	42.6	6.8	0.8	0.3	1.3	2.6	5.7	11.1	25.2	68.8	217.9	789.7
2002.....	43.2	8.7	0.6	0.4	1.0	2.3	5.8	10.9	25.8	72.2	226.4	793.9
2001.....	42.6	8.2	0.8	0.4	1.1	2.6	5.0	11.0	26.4	73.3	218.8	801.1
2000.....	42.8	9.7	0.7	0.3	1.2	2.5	5.4	10.8	26.6	73.9	222.6	812.7
1999.....	43.2	10.0	0.7	0.4	1.5	2.6	5.4	11.0	27.4	74.9	225.5	821.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

Black, both sexes

2006 4/.....	35.1	13.7	1.4	0.8	3.4	10.1	18.6	37.2	70.8	134.4	283.7	605.0
2005 4/.....	36.1	12.2	1.1	0.9	3.8	10.0	18.9	40.3	74.8	139.2	280.8	640.9
2004 4/.....	36.3	12.4	1.8	0.9	3.4	9.0	19.6	40.7	74.9	144.2	294.9	668.3
2003 4/.....	37.5	15.0	1.6	0.8	3.9	9.7	20.3	41.3	79.2	158.2	294.5	703.7
2002.....	37.8	17.2	1.1	0.7	3.8	10.1	20.3	42.4	81.0	150.9	308.2	738.5
2001.....	38.0	17.2	2.3	0.9	3.7	9.6	20.7	42.8	81.5	155.1	318.1	730.2
2000.....	38.2	15.6	1.4	0.9	4.3	9.4	20.5	44.0	81.7	161.8	319.6	742.0
1999.....	39.6	20.2	1.4	0.8	4.2	9.6	21.8	44.3	91.7	169.8	328.2	746.1

Black, male

2006 4/.....	36.8	14.1	1.6	0.9	3.9	12.8	23.3	48.6	93.6	171.6	333.1	637.2
2005 4/.....	37.3	14.8	*	1.0	5.2	12.3	23.0	51.3	96.4	179.9	325.9	661.9
2004 4/.....	37.5	13.3	1.8	0.8	4.4	10.8	23.7	51.7	98.6	177.9	359.4	715.3
2003 4/.....	38.6	17.8	1.8	0.9	5.0	10.9	24.8	52.9	103.3	197.6	355.1	707.9
2002.....	38.3	18.9	*	0.8	4.7	12.7	24.3	54.5	102.0	186.7	359.2	753.7
2001.....	38.9	15.0	2.6	0.8	4.8	11.4	25.1	52.4	106.8	188.2	395.5	805.6
2000.....	39.2	14.7	1.6	1.0	4.9	11.2	25.0	57.2	106.9	200.2	379.7	816.5
1999.....	41.5	19.8	*	1.1	5.4	11.5	26.6	56.4	121.0	218.8	398.1	849.5

Black, female

2006 4/.....	33.6	13.2	*	0.7	2.9	7.6	14.4	27.4	52.4	107.8	255.6	591.9
2005 4/.....	34.9	9.6	*	0.8	2.3	7.9	15.3	30.9	57.5	110.0	255.2	632.4
2004 4/.....	35.1	11.4	1.7	1.0	2.4	7.3	15.9	31.2	55.8	120.2	258.4	649.7
2003 4/.....	36.5	12.1	*	0.7	2.7	8.5	16.2	31.4	59.7	130.2	260.2	702.0
2002.....	37.4	15.4	*	*	2.9	7.7	16.7	32.0	64.1	125.5	279.3	732.8
2001.....	37.2	19.5	2.0	0.9	2.6	7.9	16.7	34.5	61.1	131.7	274.6	702.2
2000.....	37.3	16.5	*	0.8	3.7	7.8	16.5	32.7	61.4	134.8	285.7	714.3
1999.....	37.9	20.5	*	*	3.0	7.9	17.5	34.0	68.2	135.1	289.0	707.4



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/ 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

All races 3/, both sexes

2006.....	8.0	*	*	*	0.0	0.3	0.9	3.0	6.9	16.8	51.0	189.4
2005.....	8.4	*	*	*	0.1	0.2	0.9	2.7	6.4	17.7	55.6	210.0
2004.....	7.9	*	*	*	0.1	0.3	0.8	2.7	6.3	17.1	52.6	198.5
2003.....	7.5	*	*	*	0.1	0.2	0.8	2.5	6.3	16.9	51.7	188.9
2002.....	7.0	*	*	*	0.1	0.2	0.8	2.3	5.7	16.0	48.2	180.4
2001.....	6.8	*	*	*	0.1	0.3	0.7	2.4	5.8	15.5	47.7	171.9
2000.....	6.4	*	*	*	*	0.2	0.8	2.3	5.9	15.1	45.5	162.9
1999.....	6.1	*	*	*	*	0.2	0.7	2.2	5.5	15.2	43.6	152.1

All races 3/, male

2006.....	6.4	*	*	*	*	0.3	1.1	3.8	8.4	18.5	51.2	159.2
2005.....	6.5	*	*	*	*	0.2	1.1	3.5	7.9	19.2	55.3	169.3
2004.....	6.1	*	*	*	*	0.3	1.0	3.5	7.4	18.2	53.3	166.0
2003.....	5.7	*	*	*	*	0.3	1.0	3.3	7.4	18.4	48.9	153.5
2002.....	5.4	*	*	*	*	0.2	1.0	2.8	6.6	17.1	47.3	153.4
2001.....	5.2	*	*	*	*	0.3	0.9	3.0	6.5	16.6	45.8	150.8
2000.....	5.0	*	*	*	*	0.2	1.0	3.0	7.0	15.8	45.2	142.4
1999.....	4.7	*	*	*	*	0.2	0.9	2.7	6.4	16.5	42.9	132.4

All races 3/, female

2006.....	9.5	*	*	*	*	0.2	0.7	2.2	5.5	15.3	50.8	203.5
2005.....	10.3	*	*	*	*	0.2	0.7	2.0	5.1	16.5	55.8	228.7
2004.....	9.5	*	*	*	*	0.2	0.6	2.0	5.3	16.1	52.1	213.2
2003.....	9.3	*	*	*	*	0.2	0.7	1.8	5.3	15.7	53.5	204.6
2002.....	8.6	*	*	*	*	0.2	0.6	1.8	4.9	15.1	48.8	192.1
2001.....	8.3	*	*	*	*	0.2	0.6	1.8	5.2	14.5	48.9	180.8
2000.....	7.8	*	*	*	*	0.2	0.6	1.7	4.9	14.6	45.7	171.3
1999.....	7.4	*	*	*	*	0.2	0.5	1.7	4.7	14.1	43.9	160.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

White, both sexes

2006 4/.....	7.5	*	*	*	*	0.1	0.5	1.9	5.1	13.4	45.5	179.5
2005 4/.....	8.0	*	*	*	*	0.1	0.5	1.7	4.7	13.8	50.0	200.3
2004 4/.....	7.4	*	*	*	*	0.1	0.5	1.7	4.4	13.5	46.7	188.1
2003 4/.....	7.2	*	*	*	*	0.1	0.5	1.7	4.4	13.0	46.2	181.0
2002.....	6.6	*	*	*	*	0.1	0.4	1.5	4.1	12.8	42.6	170.4
2001.....	6.3	*	*	*	*	0.1	0.4	1.4	4.0	12.1	42.3	163.3
2000.....	6.0	*	*	*	*	0.1	0.4	1.4	4.1	11.8	40.4	153.2
1999.....	5.6	*	*	*	*	0.1	0.4	1.3	3.7	12.1	37.9	142.0

White, male

2006 4/.....	5.8	*	*	*	*	0.2	0.6	2.6	6.3	14.6	45.7	151.3
2005 4/.....	5.9	*	*	*	*	*	0.7	2.3	5.8	14.8	49.1	160.2
2004 4/.....	5.6	*	*	*	*	0.2	0.6	2.3	5.3	14.6	47.1	159.4
2003 4/.....	5.2	*	*	*	*	0.1	0.7	2.2	5.2	14.5	43.8	145.3
2002.....	4.9	*	*	*	*	0.1	0.5	1.9	4.9	13.8	41.5	145.2
2001.....	4.7	*	*	*	*	0.2	0.5	1.8	4.5	13.2	40.4	142.2
2000.....	4.5	*	*	*	*	0.1	0.6	1.9	5.0	12.6	40.3	132.5
1999.....	4.2	*	*	*	*	0.2	0.5	1.7	4.5	13.4	37.3	121.0

White, female

2006 4/.....	9.2	*	*	*	*	*	0.3	1.3	3.9	12.4	45.4	192.7
2005 4/.....	10.1	*	*	*	*	*	0.4	1.2	3.7	13.0	50.6	218.8
2004 4/.....	9.2	*	*	*	*	*	0.3	1.2	3.4	12.5	46.5	201.1
2003 4/.....	9.1	*	*	*	*	0.1	0.4	1.2	3.6	11.7	47.8	196.8
2002.....	8.3	*	*	*	*	*	0.3	1.0	3.3	11.9	43.4	181.3
2001.....	8.0	*	*	*	*	*	0.3	1.0	3.5	11.3	43.6	172.2
2000.....	7.4	*	*	*	*	*	0.3	0.9	3.2	11.2	40.4	161.6
1999.....	7.0	*	*	*	*	*	0.2	1.0	3.0	11.0	38.3	150.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Essential hypertension and hypertensive renal disease (I10,I12,I15)

Black, both sexes

2006 4/.....	12.4	*	*	*	*	1.1	3.7	11.0	23.0	49.2	117.0	318.3
2005 4/.....	12.7	*	*	*	*	0.9	3.4	10.5	22.6	54.2	122.1	341.0
2004 4/.....	12.3	*	*	*	*	1.0	3.1	10.2	23.9	51.9	122.6	335.6
2003 4/.....	11.8	*	*	*	*	1.0	3.1	9.3	24.1	54.4	119.5	301.1
2002.....	11.1	*	*	*	*	0.8	3.3	9.0	21.0	47.6	118.0	308.8
2001.....	10.9	*	*	*	*	1.0	3.1	9.9	22.0	47.4	112.6	288.0
2000.....	10.6	*	*	*	*	0.6	3.1	9.7	22.9	48.7	107.1	283.4
1999.....	10.3	*	*	*	*	0.7	3.1	9.1	21.5	46.7	111.7	269.4

Black, male

2006 4/.....	11.3	*	*	*	*	1.3	4.7	13.8	28.1	59.9	126.4	271.7
2005 4/.....	11.4	*	*	*	*	1.0	3.9	13.2	29.3	64.0	135.8	289.1
2004 4/.....	10.7	*	*	*	*	1.1	3.9	12.9	28.3	56.8	137.0	263.3
2003 4/.....	10.4	*	*	*	*	1.3	3.4	12.7	29.3	60.5	119.6	274.2
2002.....	9.6	*	*	*	*	0.9	4.1	11.0	23.5	52.0	127.4	274.5
2001.....	9.7	*	*	*	*	1.1	3.8	12.9	25.5	54.4	118.3	275.7
2000.....	9.1	*	*	*	*	*	3.7	11.9	27.8	50.5	109.0	269.9
1999.....	9.1	*	*	*	*	*	3.7	11.6	24.6	51.3	118.7	266.0

Black, female

2006 4/.....	13.4	*	*	*	*	1.0	2.7	8.5	18.8	41.6	111.7	337.4
2005 4/.....	13.8	*	*	*	*	0.8	3.1	8.1	17.2	47.2	114.3	362.0
2004 4/.....	13.8	*	*	*	*	0.9	2.4	7.8	20.3	48.4	114.5	364.2
2003 4/.....	13.0	*	*	*	*	*	2.7	6.4	20.0	50.1	119.4	311.5
2002.....	12.5	*	*	*	*	0.7	2.5	7.2	19.0	44.5	112.7	322.0
2001.....	11.9	*	*	*	*	0.9	2.4	7.4	19.2	42.5	109.4	292.6
2000.....	12.0	*	*	*	*	*	2.6	7.7	18.9	47.5	106.0	288.5
1999.....	11.4	*	*	*	*	0.7	2.6	7.0	19.0	43.4	107.7	270.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

All races 3/, both sexes

2006.....	45.8	3.4	0.3	0.2	0.5	1.3	5.1	14.7	33.3	96.3	335.1	1,039.6
2005.....	48.4	3.1	0.4	0.2	0.5	1.4	5.2	15.0	33.0	101.1	359.0	1,141.8
2004.....	51.1	3.1	0.3	0.2	0.5	1.4	5.4	14.9	34.3	107.8	386.2	1,245.9
2003.....	54.2	2.5	0.3	0.2	0.5	1.5	5.5	15.0	35.6	112.9	410.7	1,370.1
2002.....	56.4	2.9	0.3	0.2	0.4	1.4	5.4	15.1	37.2	120.3	431.0	1,445.9
2001.....	57.4	2.7	0.4	0.2	0.5	1.5	5.5	15.1	38.0	123.4	443.9	1,500.2
2000.....	59.6	3.3	0.3	0.2	0.5	1.5	5.8	16.0	41.0	128.6	461.3	1,589.2
1999.....	60.0	2.7	0.3	0.2	0.5	1.4	5.7	15.2	40.6	130.8	469.8	1,614.8

All races 3/, male

2006.....	37.0	3.9	0.3	0.3	0.5	1.4	5.3	16.4	38.7	108.0	345.5	932.4
2005.....	38.8	3.5	0.5	0.3	0.4	1.5	5.2	16.5	38.5	113.6	372.9	1,023.3
2004.....	40.7	3.4	0.3	0.2	0.5	1.4	5.6	16.7	39.5	121.1	402.9	1,118.1
2003.....	42.9	2.8	0.3	0.2	0.5	1.6	5.8	16.7	40.8	127.8	431.4	1,236.0
2002.....	44.2	3.2	0.4	0.2	0.5	1.4	5.3	16.7	42.7	135.0	445.9	1,317.9
2001.....	45.2	3.1	0.3	0.2	0.5	1.6	5.7	16.7	43.4	140.4	467.3	1,380.2
2000.....	46.9	3.8	*	0.2	0.5	1.5	5.8	17.5	47.2	145.0	490.8	1,484.3
1999.....	47.1	3.3	0.3	0.2	0.5	1.5	5.8	16.7	46.5	147.3	500.4	1,512.8

All races 3/, female

2006.....	54.4	2.9	0.4	0.2	0.5	1.2	4.8	13.0	28.2	86.5	328.0	1,089.8
2005.....	57.8	2.6	0.3	0.2	0.5	1.2	5.1	13.6	27.9	90.5	349.5	1,196.1
2004.....	61.2	2.8	*	0.2	0.5	1.4	5.1	13.1	29.5	96.6	374.9	1,303.4
2003.....	65.1	2.2	0.3	0.1	0.5	1.4	5.3	13.4	30.9	100.5	396.8	1,429.4
2002.....	68.2	2.5	0.3	0.2	0.3	1.4	5.5	13.6	32.1	108.1	421.2	1,501.5
2001.....	69.2	2.3	0.4	0.2	0.5	1.5	5.4	13.6	32.9	109.3	428.6	1,550.4
2000.....	71.8	2.7	0.4	0.2	0.5	1.5	5.7	14.5	35.3	115.1	442.1	1,632.0
1999.....	72.3	2.1	0.3	0.2	0.5	1.4	5.6	13.8	35.1	117.2	449.8	1,656.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

White, both sexes

2006 4/.....	47.9	2.8	0.3	0.2	0.5	1.1	4.0	11.6	27.7	87.5	328.4	1,050.8
2005 4/.....	50.7	2.5	0.4	0.2	0.4	1.1	4.1	11.6	27.7	92.3	351.8	1,153.3
2004 4/.....	53.7	2.5	0.2	0.2	0.5	1.2	4.4	11.4	28.6	99.0	377.7	1,257.8
2003 4/.....	57.0	2.0	0.2	0.2	0.5	1.2	4.2	11.4	29.4	103.7	403.0	1,382.2
2002.....	59.5	2.3	0.3	0.2	0.4	1.2	4.2	11.6	31.4	110.7	423.5	1,459.5
2001.....	60.5	1.9	0.3	0.2	0.4	1.3	4.4	11.4	31.7	113.0	436.4	1,511.2
2000.....	62.8	2.5	0.3	0.2	0.5	1.3	4.5	12.4	34.8	119.4	452.4	1,601.6
1999.....	63.3	2.3	0.3	0.2	0.4	1.3	4.4	11.7	34.0	121.1	461.6	1,631.6

White, male

2006 4/.....	37.7	2.9	*	0.3	0.5	1.2	4.2	12.8	31.5	97.1	338.5	941.3
2005 4/.....	39.7	2.8	0.4	0.2	0.4	1.2	4.2	12.8	31.7	103.0	364.8	1,033.7
2004 4/.....	41.8	2.8	*	0.2	0.5	1.2	4.7	12.8	32.4	110.8	393.7	1,129.3
2003 4/.....	44.2	2.0	*	0.2	0.5	1.3	4.4	12.9	33.3	117.3	422.4	1,247.0
2002.....	45.7	2.6	0.4	0.2	0.5	1.2	4.2	12.9	35.6	123.8	437.5	1,327.4
2001.....	46.6	2.2	*	0.2	0.5	1.4	4.5	12.7	36.1	128.5	458.8	1,386.2
2000.....	48.4	3.0	*	0.2	0.5	1.4	4.5	13.6	39.7	133.8	480.0	1,490.7
1999.....	48.7	2.8	*	0.2	0.5	1.3	4.6	12.8	38.9	136.0	490.3	1,522.7

White, female

2006 4/.....	57.9	2.6	0.4	0.1	0.4	1.0	3.7	10.4	24.1	79.3	321.5	1,102.2
2005 4/.....	61.6	2.1	*	0.2	0.4	1.0	4.0	10.5	23.8	83.2	342.9	1,208.5
2004 4/.....	65.3	2.2	*	0.2	0.4	1.1	4.1	10.1	25.1	89.0	366.8	1,315.7
2003 4/.....	69.5	2.0	*	0.1	0.4	1.2	4.0	10.0	25.8	92.1	389.9	1,442.1
2002.....	73.0	2.0	*	0.2	0.3	1.2	4.3	10.4	27.4	99.5	414.1	1,516.9
2001.....	74.0	1.6	0.4	0.2	0.4	1.3	4.2	10.2	27.6	99.9	421.6	1,563.5
2000.....	76.9	2.0	*	0.2	0.4	1.3	4.4	11.2	30.2	107.3	434.2	1,646.7
1999.....	77.6	1.9	*	0.1	0.4	1.2	4.3	10.7	29.4	108.6	442.8	1,675.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Cerebrovascular diseases (I60-I69)

Black, both sexes

2006 4/.....	43.1	7.2	*	0.3	0.7	2.7	12.2	37.0	81.1	178.0	435.7	1,008.7
2005 4/.....	44.9	6.0	*	0.4	0.8	2.9	12.1	39.5	79.4	183.2	469.6	1,106.7
2004 4/.....	46.9	6.0	*	*	0.8	3.1	12.3	39.0	83.9	195.3	511.7	1,211.2
2003 4/.....	49.3	4.9	*	*	0.9	3.1	13.8	41.0	89.8	201.1	531.1	1,348.7
2002.....	50.0	5.5	*	0.4	0.6	3.1	12.8	40.7	88.0	215.1	552.3	1,383.9
2001.....	51.1	6.6	*	*	0.8	2.9	13.2	42.6	91.1	222.5	560.3	1,452.7
2000.....	52.5	7.5	*	*	0.7	2.9	14.5	43.4	93.8	222.9	588.8	1,530.0
1999.....	52.2	4.6	*	*	0.7	2.7	14.3	41.5	96.3	231.9	590.5	1,492.5

Black, male

2006 4/.....	39.3	9.5	*	*	0.6	2.8	12.8	43.5	105.9	218.7	471.1	882.0
2005 4/.....	40.3	6.8	*	*	0.7	3.3	12.4	44.8	103.7	224.3	503.7	983.5
2004 4/.....	41.5	6.8	*	*	1.0	2.9	12.8	44.8	107.4	235.2	551.0	1,061.0
2003 4/.....	43.2	5.9	*	*	0.8	3.3	14.3	46.9	112.1	237.4	588.9	1,180.3
2002.....	43.5	5.8	*	*	0.7	2.9	12.2	46.5	110.3	262.9	587.8	1,252.2
2001.....	44.6	8.1	*	*	0.7	3.1	13.2	48.8	111.9	269.2	613.9	1,349.1
2000.....	46.1	8.6	*	*	*	2.7	14.7	49.5	115.4	268.5	659.2	1,458.8
1999.....	45.9	*	*	*	*	2.8	14.7	48.6	117.9	274.7	650.6	1,448.3

Black, female

2006 4/.....	46.5	*	*	*	0.7	2.6	11.7	31.3	61.1	148.9	415.6	1,060.5
2005 4/.....	49.1	*	*	*	0.8	2.5	11.8	35.0	59.8	153.7	450.2	1,156.5
2004 4/.....	51.9	*	*	*	*	3.3	11.8	33.9	65.0	166.8	489.5	1,270.7
2003 4/.....	54.8	*	*	*	1.1	2.8	13.3	36.0	71.8	175.3	498.3	1,414.2
2002.....	55.8	*	*	*	*	3.3	13.4	35.7	70.1	181.2	532.2	1,434.3
2001.....	56.9	*	*	*	1.0	2.8	13.1	37.3	74.4	189.5	530.3	1,491.2
2000.....	58.3	*	*	*	0.8	3.1	14.4	38.1	76.4	190.9	549.2	1,556.5
1999.....	57.9	*	*	*	0.8	2.7	13.9	35.3	78.9	201.7	556.8	1,509.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Atherosclerosis (I70)

All races 3/, both sexes

2006.....	2.9	*	*	*	*	*	0.1	0.3	1.2	4.5	18.9	89.9
2005.....	4.0	*	*	*	*	*	0.1	0.5	1.4	5.4	25.9	132.9
2004.....	4.0	*	*	*	*	*	0.1	0.4	1.6	6.1	26.3	137.0
2003.....	4.5	*	*	*	*	*	0.1	0.5	1.7	6.6	29.8	154.6
2002.....	4.8	*	*	*	*	*	0.1	0.5	1.8	7.1	31.9	168.1
2001.....	4.9	*	*	*	*	*	0.1	0.5	1.9	7.3	33.1	179.1
2000.....	5.1	*	*	*	*	*	0.1	0.5	1.9	7.7	34.6	188.8
1999.....	5.4	*	*	*	*	*	0.1	0.5	2.0	8.0	35.9	202.0

All races 3/, male

2006.....	2.3	*	*	*	*	*	0.1	0.5	1.6	5.3	20.5	85.3
2005.....	3.1	*	*	*	*	*	0.1	0.6	1.8	6.6	29.0	121.7
2004.....	3.1	*	*	*	*	*	0.1	0.5	2.0	7.3	29.7	128.2
2003.....	3.4	*	*	*	*	*	0.2	0.7	2.2	7.8	33.6	138.6
2002.....	3.7	*	*	*	*	*	0.1	0.6	2.4	8.8	35.5	156.6
2001.....	3.7	*	*	*	*	*	0.1	0.6	2.4	9.0	35.9	167.2
2000.....	3.9	*	*	*	*	*	0.2	0.6	2.5	9.6	39.7	176.5
1999.....	4.0	*	*	*	*	*	0.1	0.7	2.7	9.6	40.0	189.0

All races 3/, female

2006.....	3.5	*	*	*	*	*	*	0.2	0.9	3.8	17.9	92.1
2005.....	4.9	*	*	*	*	*	*	0.3	1.1	4.4	23.9	138.1
2004.....	4.9	*	*	*	*	*	*	0.2	1.2	5.1	24.0	141.0
2003.....	5.5	*	*	*	*	*	*	0.3	1.2	5.6	27.3	161.7
2002.....	5.9	*	*	*	*	*	0.1	0.3	1.3	5.7	29.5	173.1
2001.....	6.2	*	*	*	*	*	*	0.3	1.4	5.9	31.2	184.1
2000.....	6.3	*	*	*	*	*	*	0.3	1.2	6.2	31.3	193.8
1999.....	6.7	*	*	*	*	*	*	0.3	1.4	6.7	33.2	207.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Atherosclerosis (I70)

White, both sexes

2006 4/.....	3.2	*	*	*	*	*	0.1	0.3	1.2	4.4	19.2	93.2
2005 4/.....	4.5	*	*	*	*	*	0.1	0.4	1.4	5.3	26.6	138.7
2004 4/.....	4.6	*	*	*	*	*	0.1	0.4	1.6	6.1	27.1	142.0
2003 4/.....	5.0	*	*	*	*	*	0.1	0.4	1.6	6.5	30.3	159.5
2002.....	5.4	*	*	*	*	*	0.1	0.4	1.7	7.1	32.1	174.7
2001.....	5.6	*	*	*	*	*	0.1	0.5	1.8	7.3	33.8	185.9
2000.....	5.7	*	*	*	*	*	0.1	0.4	1.8	7.7	35.1	195.2
1999.....	6.0	*	*	*	*	*	0.1	0.5	2.0	8.0	36.4	207.7

White, male

2006 4/.....	2.5	*	*	*	*	*	*	0.4	1.5	5.2	20.5	87.9
2005 4/.....	3.4	*	*	*	*	*	*	0.6	1.8	6.4	29.8	128.3
2004 4/.....	3.5	*	*	*	*	*	*	0.5	1.9	7.4	30.4	134.4
2003 4/.....	3.7	*	*	*	*	*	0.2	0.6	2.1	7.5	33.7	143.3
2002.....	4.0	*	*	*	*	*	0.1	0.5	2.3	8.8	35.6	163.2
2001.....	4.1	*	*	*	*	*	0.1	0.6	2.3	9.0	36.4	173.9
2000.....	4.3	*	*	*	*	*	0.2	0.5	2.5	9.7	40.3	181.4
1999.....	4.4	*	*	*	*	*	0.1	0.7	2.6	9.5	40.2	193.7

White, female

2006 4/.....	4.0	*	*	*	*	*	*	0.1	0.8	3.8	18.3	95.7
2005 4/.....	5.6	*	*	*	*	*	*	0.2	1.1	4.4	24.4	143.5
2004 4/.....	5.6	*	*	*	*	*	*	0.2	1.2	5.0	24.9	145.4
2003 4/.....	6.3	*	*	*	*	*	*	0.3	1.1	5.6	28.0	166.7
2002.....	6.7	*	*	*	*	*	*	0.3	1.3	5.7	29.8	179.7
2001.....	7.0	*	*	*	*	*	*	0.3	1.3	5.9	32.1	190.9
2000.....	7.1	*	*	*	*	*	*	0.2	1.1	6.1	31.7	200.8
1999.....	7.6	*	*	*	*	*	*	0.3	1.4	6.8	34.0	213.3



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Atherosclerosis (I70)

Black, both sexes

2006 4/.....	1.8	*	*	*	*	*	*	0.6	2.1	6.4	20.5	71.6
2005 4/.....	2.2	*	*	*	*	*	*	0.7	2.1	7.4	23.5	97.8
2004 4/.....	2.2	*	*	*	*	*	*	0.5	2.3	7.4	23.9	102.9
2003 4/.....	2.6	*	*	*	*	*	*	0.8	2.6	9.2	30.7	121.1
2002.....	2.7	*	*	*	*	*	*	1.0	2.9	8.0	36.5	117.0
2001.....	2.6	*	*	*	*	*	*	0.7	3.3	8.9	30.7	125.6
2000.....	2.9	*	*	*	*	*	*	1.1	3.2	8.9	35.2	138.3
1999.....	3.0	*	*	*	*	*	*	0.7	3.2	9.2	34.9	154.3

Black, male

2006 4/.....	1.7	*	*	*	*	*	*	*	3.0	8.6	25.4	75.1
2005 4/.....	1.7	*	*	*	*	*	*	0.9	2.8	9.6	25.9	75.3
2004 4/.....	1.7	*	*	*	*	*	*	*	3.0	8.6	30.1	78.5
2003 4/.....	2.3	*	*	*	*	*	*	1.2	3.6	12.4	40.3	104.3
2002.....	2.3	*	*	*	*	*	*	1.4	4.2	10.1	44.2	110.4
2001.....	2.1	*	*	*	*	*	*	1.0	3.8	10.9	37.1	113.5
2000.....	2.4	*	*	*	*	*	*	1.4	3.9	11.0	40.3	147.6
1999.....	2.5	*	*	*	*	*	*	*	4.4	11.8	44.0	164.0

Black, female

2006 4/.....	1.9	*	*	*	*	*	*	*	1.4	4.8	17.7	70.2
2005 4/.....	2.6	*	*	*	*	*	*	*	1.5	5.8	22.2	106.9
2004 4/.....	2.6	*	*	*	*	*	*	*	1.7	6.5	20.5	112.6
2003 4/.....	2.9	*	*	*	*	*	*	*	1.8	7.0	25.2	127.6
2002.....	3.1	*	*	*	*	*	*	*	1.9	6.5	32.2	119.5
2001.....	3.1	*	*	*	*	*	*	*	2.9	7.5	27.1	130.1
2000.....	3.3	*	*	*	*	*	*	*	2.7	7.5	32.4	134.8
1999.....	3.4	*	*	*	*	*	*	*	2.3	7.4	29.8	150.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/ 1 year	1-4 2/ years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	-----------------	--------------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes

2006.....	7.5	*	*	*	0.2	0.4	1.0	2.7	6.8	21.7	58.0	128.5
2005.....	8.0	*	*	*	0.1	0.4	1.1	2.6	7.5	24.1	62.6	135.5
2004.....	8.1	*	*	*	0.1	0.4	1.0	2.7	7.2	25.1	64.3	139.3
2003.....	8.5	*	*	*	0.2	0.4	1.0	2.9	8.1	26.9	67.6	146.0
2002.....	8.6	*	*	*	0.2	0.4	1.0	2.7	8.6	29.1	68.7	148.4
2001.....	8.9	*	*	*	0.2	0.4	1.1	2.6	8.7	29.2	71.8	158.2
2000.....	9.3	*	*	*	0.2	0.4	1.0	2.8	9.2	32.2	74.6	163.2
1999.....	9.2	*	*	*	0.1	0.4	1.1	2.7	9.3	31.5	75.8	165.2

All races 3/, male

2006.....	8.0	*	*	*	0.2	0.5	1.5	3.8	9.6	29.0	74.5	155.0
2005.....	8.5	*	*	*	0.2	0.5	1.5	3.6	10.5	32.8	81.8	158.2
2004.....	8.5	*	*	*	0.2	0.6	1.4	3.8	9.9	33.4	82.8	166.5
2003.....	9.2	*	*	*	0.2	0.5	1.4	4.1	11.3	37.0	91.8	179.7
2002.....	9.4	*	*	*	0.2	0.5	1.3	3.7	11.9	39.8	92.3	187.4
2001.....	9.6	*	*	*	0.2	0.6	1.5	3.6	12.0	40.3	97.0	197.3
2000.....	10.2	*	*	*	0.2	0.5	1.2	3.8	13.1	45.4	102.9	214.9
1999.....	10.1	*	*	*	0.2	0.5	1.3	3.7	13.3	44.4	104.2	207.5

All races 3/, female

2006.....	7.0	*	*	*	0.1	0.2	0.6	1.6	4.2	15.5	46.7	116.1
2005.....	7.4	*	*	*	*	0.3	0.7	1.6	4.7	16.7	49.5	125.0
2004.....	7.6	*	*	*	*	0.3	0.7	1.6	4.7	18.1	51.8	127.0
2003.....	7.8	*	*	*	0.1	0.2	0.7	1.8	5.2	18.4	51.4	131.1
2002.....	7.9	*	*	*	0.1	0.3	0.7	1.7	5.5	20.1	53.0	131.5
2001.....	8.2	*	*	*	0.1	0.3	0.7	1.7	5.6	19.9	55.3	141.8
2000.....	8.3	*	*	*	0.1	0.3	0.8	1.8	5.6	21.4	56.2	142.2
1999.....	8.4	*	*	*	*	0.2	0.8	1.7	5.6	21.0	57.3	148.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

White, both sexes

2006 4/.....	8.1	*	*	*	0.2	0.3	0.9	2.3	6.5	21.7	58.2	129.7
2005 4/.....	8.5	*	*	*	0.1	0.3	0.9	2.2	7.1	24.1	62.8	136.1
2004 4/.....	8.7	*	*	*	0.1	0.4	0.9	2.4	6.8	25.4	64.7	139.3
2003 4/.....	9.2	*	*	*	0.2	0.3	0.9	2.5	7.9	27.3	68.3	146.9
2002.....	9.3	*	*	*	0.2	0.3	0.9	2.3	8.2	29.4	69.7	148.9
2001.....	9.6	*	*	*	0.1	0.4	0.9	2.3	8.3	29.5	72.1	158.2
2000.....	10.0	*	*	*	0.1	0.3	0.8	2.4	8.8	32.7	75.1	163.0
1999.....	10.0	*	*	*	0.1	0.3	1.0	2.4	9.1	32.1	76.4	165.1

White, male

2006 4/.....	8.6	*	*	*	0.2	0.5	1.3	3.4	9.2	29.2	76.0	159.4
2005 4/.....	9.1	*	*	*	0.2	0.4	1.2	3.0	10.1	33.1	82.3	162.0
2004 4/.....	9.2	*	*	*	0.2	0.5	1.2	3.4	9.3	34.1	84.2	169.3
2003 4/.....	10.0	*	*	*	0.2	0.4	1.2	3.6	11.0	37.9	93.1	184.1
2002.....	10.2	*	*	*	0.2	0.4	1.2	3.2	11.4	40.6	94.6	190.9
2001.....	10.4	*	*	*	0.2	0.6	1.2	3.1	11.6	41.1	98.3	201.3
2000.....	11.1	*	*	*	0.2	0.5	1.0	3.4	12.8	46.7	104.7	219.6
1999.....	11.0	*	*	*	0.2	0.5	1.2	3.3	13.4	45.8	106.2	211.1

White, female

2006 4/.....	7.5	*	*	*	0.1	0.2	0.5	1.3	3.9	15.2	45.9	115.7
2005 4/.....	8.0	*	*	*	*	0.2	0.6	1.3	4.3	16.3	49.3	124.2
2004 4/.....	8.2	*	*	*	*	0.2	0.6	1.4	4.4	17.9	51.3	125.7
2003 4/.....	8.3	*	*	*	*	0.2	0.5	1.5	4.9	18.3	51.5	130.5
2002.....	8.5	*	*	*	*	0.3	0.6	1.4	5.2	19.9	52.9	130.7
2001.....	8.8	*	*	*	*	0.3	0.6	1.5	5.2	19.7	54.7	140.1
2000.....	8.9	*	*	*	*	0.2	0.6	1.4	5.1	21.0	55.6	140.1
1999.....	9.1	*	*	*	*	0.2	0.7	1.5	5.2	20.6	56.9	146.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of circulatory system (I71-I78)

Black, both sexes

2006 4/.....	6.3	*	*	*	*	0.7	2.0	5.7	11.5	26.5	64.7	132.3
2005 4/.....	6.7	*	*	*	*	0.8	2.3	5.4	12.1	29.2	69.8	144.6
2004 4/.....	6.6	*	*	*	*	0.8	2.2	5.3	12.7	28.0	70.4	155.1
2003 4/.....	6.6	*	*	*	*	0.7	2.3	6.2	11.8	29.3	70.1	150.0
2002.....	6.7	*	*	*	*	0.9	2.0	5.9	13.5	31.1	68.3	156.8
2001.....	7.1	*	*	*	*	0.7	2.4	5.1	13.9	31.7	79.8	174.7
2000.....	7.1	*	*	*	*	0.7	2.2	5.5	14.4	33.0	78.0	177.7
1999.....	6.7	*	*	*	*	0.5	2.0	5.2	12.3	31.7	76.6	178.9

Black, male

2006 4/.....	6.3	*	*	*	*	1.0	3.0	7.9	16.3	34.7	67.6	117.1
2005 4/.....	6.9	*	*	*	*	1.0	3.6	7.9	16.8	36.2	88.8	129.2
2004 4/.....	6.7	*	*	*	*	1.2	3.0	7.5	18.1	35.2	79.9	158.0
2003 4/.....	6.7	*	*	*	*	1.1	3.0	8.2	16.4	36.3	90.3	138.7
2002.....	6.7	*	*	*	*	1.2	2.6	8.1	19.3	38.4	78.6	166.2
2001.....	6.9	*	*	*	*	0.9	3.4	7.3	17.8	40.0	96.8	165.7
2000.....	6.9	*	*	*	*	0.9	2.8	7.5	18.4	41.1	95.7	177.6
1999.....	6.5	*	*	*	*	0.9	2.3	7.2	15.7	39.4	89.8	184.0

Black, female

2006 4/.....	6.3	*	*	*	*	*	1.2	3.8	7.6	20.6	63.1	138.5
2005 4/.....	6.5	*	*	*	*	*	1.2	3.3	8.4	24.2	59.0	150.8
2004 4/.....	6.6	*	*	*	*	*	1.6	3.4	8.4	22.8	65.0	154.0
2003 4/.....	6.5	*	*	*	*	*	1.7	4.5	8.2	24.2	58.7	154.3
2002.....	6.6	*	*	*	*	*	1.4	4.0	8.9	25.8	62.5	153.2
2001.....	7.2	*	*	*	*	*	1.6	3.3	10.8	26.0	70.2	178.1
2000.....	7.2	*	*	*	*	*	1.5	3.9	11.1	27.3	68.0	177.7
1999.....	7.0	*	*	*	*	*	1.7	3.5	9.6	26.2	69.2	177.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Aortic aneurysm and dissection (I71)

All races 3/, both sexes

2006.....	4.4	*	*	*	0.1	0.3	0.8	1.8	4.3	13.9	35.4	62.7
2005.....	4.7	*	*	*	0.1	0.3	0.8	1.8	4.8	15.4	38.1	64.9
2004.....	4.7	*	*	*	0.1	0.3	0.7	1.8	4.5	15.9	39.6	64.8
2003.....	5.1	*	*	*	0.1	0.3	0.7	2.0	5.3	17.8	42.5	70.0
2002.....	5.1	*	*	*	0.1	0.3	0.7	1.8	5.5	19.1	43.3	69.4
2001.....	5.3	*	*	*	0.1	0.3	0.7	1.7	5.5	19.2	45.7	77.5
2000.....	5.6	*	*	*	0.1	0.3	0.7	1.8	6.2	21.8	47.2	79.2
1999.....	5.7	*	*	*	0.1	0.2	0.7	1.7	6.0	21.7	48.6	80.3

All races 3/, male

2006.....	5.2	*	*	*	0.1	0.4	1.2	2.8	6.6	19.5	48.4	87.6
2005.....	5.6	*	*	*	0.1	0.4	1.2	2.8	7.3	22.4	52.6	91.0
2004.....	5.6	*	*	*	0.2	0.5	1.1	2.7	6.9	23.1	54.8	95.0
2003.....	6.3	*	*	*	0.1	0.4	1.1	3.1	8.2	26.0	61.9	105.5
2002.....	6.4	*	*	*	0.2	0.4	1.0	2.7	8.5	28.5	62.7	106.6
2001.....	6.5	*	*	*	0.2	0.4	1.2	2.6	8.5	28.7	66.4	114.8
2000.....	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.9	70.6	127.3
1999.....	7.0	*	*	*	0.1	0.4	1.0	2.7	9.5	32.7	72.7	125.3

All races 3/, female

2006.....	3.6	*	*	*	*	0.1	0.3	0.9	2.2	9.3	26.5	51.0
2005.....	3.8	*	*	*	*	0.2	0.3	0.8	2.5	9.5	28.2	52.9
2004.....	3.8	*	*	*	*	0.2	0.3	0.8	2.3	9.8	29.3	51.2
2003.....	3.9	*	*	*	*	0.1	0.3	1.0	2.6	11.0	29.5	54.2
2002.....	4.0	*	*	*	*	0.1	0.3	0.8	2.7	11.2	30.5	53.3
2001.....	4.2	*	*	*	*	0.1	0.3	0.8	2.8	11.3	32.1	61.9
2000.....	4.3	*	*	*	*	0.1	0.4	0.9	3.0	12.7	32.0	59.6
1999.....	4.4	*	*	*	*	*	0.4	0.8	2.8	12.7	32.9	62.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Aortic aneurysm and dissection (I71)

White, both sexes

2006 4/.....	4.8	*	*	*	0.1	0.2	0.7	1.6	4.2	14.2	36.4	64.3
2005 4/.....	5.1	*	*	*	0.1	0.2	0.6	1.5	4.7	15.8	39.1	66.2
2004 4/.....	5.1	*	*	*	0.1	0.3	0.6	1.5	4.3	16.5	40.6	66.8
2003 4/.....	5.6	*	*	*	0.1	0.2	0.6	1.7	5.2	18.6	44.0	72.4
2002.....	5.7	*	*	*	0.1	0.2	0.6	1.5	5.4	19.9	45.0	71.2
2001.....	5.9	*	*	*	0.1	0.3	0.6	1.5	5.4	19.9	46.9	79.7
2000.....	6.2	*	*	*	0.1	0.2	0.6	1.6	6.0	22.7	48.5	80.8
1999.....	6.3	*	*	*	0.1	0.2	0.6	1.5	6.1	22.6	50.1	82.2

White, male

2006 4/.....	5.7	*	*	*	0.2	0.4	1.1	2.5	6.5	19.9	50.3	90.4
2005 4/.....	6.1	*	*	*	0.1	0.3	0.9	2.3	7.2	23.2	54.2	94.2
2004 4/.....	6.1	*	*	*	0.2	0.4	0.9	2.4	6.5	24.1	56.6	98.7
2003 4/.....	6.9	*	*	*	0.1	0.3	1.0	2.7	8.2	27.3	64.3	110.0
2002.....	7.0	*	*	*	0.1	0.3	0.9	2.3	8.4	29.8	65.5	111.8
2001.....	7.2	*	*	*	0.1	0.5	1.0	2.3	8.4	29.9	68.5	120.3
2000.....	7.8	*	*	*	*	0.3	0.8	2.5	9.5	34.5	73.0	132.8
1999.....	7.8	*	*	*	*	0.4	0.9	2.4	9.8	34.3	75.8	129.8

White, female

2006 4/.....	3.9	*	*	*	*	*	0.3	0.7	2.0	9.2	26.8	52.0
2005 4/.....	4.1	*	*	*	*	0.2	0.3	0.6	2.3	9.5	28.7	53.3
2004 4/.....	4.1	*	*	*	*	0.2	0.3	0.7	2.1	10.1	29.8	52.4
2003 4/.....	4.3	*	*	*	*	0.2	0.2	0.8	2.4	11.1	30.3	55.7
2002.....	4.3	*	*	*	*	*	0.3	0.7	2.6	11.4	31.2	53.5
2001.....	4.6	*	*	*	*	*	0.2	0.6	2.6	11.5	32.5	62.6
2000.....	4.6	*	*	*	*	0.1	0.3	0.7	2.8	12.8	32.3	59.6
1999.....	4.8	*	*	*	*	*	0.4	0.7	2.7	12.8	33.3	63.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Aortic aneurysm and dissection (I71)

Black, both sexes

2006 4/.....	3.2	*	*	*	*	0.5	1.5	4.0	6.6	14.4	28.0	47.2
2005 4/.....	3.3	*	*	*	*	0.5	1.7	4.0	7.0	14.4	30.9	50.2
2004 4/.....	3.2	*	*	*	*	0.6	1.6	3.6	7.6	13.0	32.0	43.7
2003 4/.....	3.2	*	*	*	*	0.5	1.6	4.2	7.1	14.5	30.0	46.4
2002.....	3.2	*	*	*	*	0.6	1.4	4.2	7.5	14.6	29.9	50.5
2001.....	3.4	*	*	*	*	0.4	1.6	3.5	7.6	14.6	37.6	55.3
2000.....	3.5	*	*	*	*	0.5	1.6	3.2	8.2	17.0	36.3	62.3
1999.....	3.3	*	*	*	*	0.4	1.4	3.7	6.5	16.7	33.8	60.3

Black, male

2006 4/.....	3.6	*	*	*	*	0.8	2.3	5.8	9.6	19.0	29.8	50.9
2005 4/.....	3.9	*	*	*	*	0.8	3.0	6.1	10.1	18.0	39.1	51.1
2004 4/.....	3.8	*	*	*	*	1.0	2.5	5.5	11.8	18.2	36.7	56.6
2003 4/.....	3.8	*	*	*	*	1.0	2.4	6.4	10.8	18.0	39.2	51.1
2002.....	3.8	*	*	*	*	1.0	2.2	6.6	12.1	19.0	35.0	49.3
2001.....	3.8	*	*	*	*	*	2.6	5.3	11.2	20.0	47.4	48.8
2000.....	3.9	*	*	*	*	*	2.4	4.7	11.5	22.1	46.9	69.2
1999.....	3.6	*	*	*	*	*	1.9	5.8	9.1	21.9	38.4	76.2

Black, female

2006 4/.....	2.8	*	*	*	*	*	0.8	2.4	4.2	11.0	26.9	45.7
2005 4/.....	2.9	*	*	*	*	*	*	2.2	4.4	11.9	26.2	49.9
2004 4/.....	2.6	*	*	*	*	*	0.8	2.0	4.2	9.4	29.3	38.6
2003 4/.....	2.7	*	*	*	*	*	1.0	2.4	4.2	11.9	24.7	44.6
2002.....	2.7	*	*	*	*	*	*	2.2	3.8	11.5	27.0	50.9
2001.....	3.0	*	*	*	*	*	0.7	1.9	4.6	10.8	32.1	57.7
2000.....	3.2	*	*	*	*	*	0.9	2.0	5.5	13.4	30.3	59.7
1999.....	3.0	*	*	*	*	*	0.9	2.0	4.4	13.0	31.2	54.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races 3/, both sexes

2006.....	3.1	*	*	*	0.1	0.1	0.2	0.8	2.5	7.8	22.6	65.8
2005.....	3.3	*	*	*	*	0.1	0.3	0.8	2.7	8.7	24.5	70.6
2004.....	3.4	*	*	*	*	0.1	0.3	1.0	2.7	9.2	24.7	74.5
2003.....	3.4	*	*	*	0.1	0.1	0.3	0.9	2.8	9.1	25.0	76.0
2002.....	3.5	*	*	*	0.1	0.1	0.3	0.9	3.1	10.0	25.3	79.0
2001.....	3.5	*	*	*	*	0.1	0.4	0.9	3.2	10.0	26.1	80.7
2000.....	3.6	*	*	*	0.1	0.1	0.3	1.0	3.1	10.4	27.4	84.0
1999.....	3.6	*	*	*	*	0.1	0.3	0.9	3.3	9.8	27.2	84.9

All races 3/, male

2006.....	2.8	*	*	*	*	*	0.2	1.0	3.0	9.5	26.1	67.3
2005.....	2.9	*	*	*	*	*	0.3	0.8	3.2	10.4	29.2	67.2
2004.....	2.9	*	*	*	*	0.1	0.3	1.1	3.0	10.3	28.0	71.6
2003.....	3.0	*	*	*	*	0.1	0.3	1.0	3.1	11.0	29.9	74.2
2002.....	3.0	*	*	*	*	*	0.3	1.0	3.4	11.3	29.6	80.8
2001.....	3.1	*	*	*	*	0.1	0.3	0.9	3.5	11.6	30.6	82.5
2000.....	3.2	*	*	*	*	0.1	0.3	1.1	3.6	12.5	32.3	87.6
1999.....	3.1	*	*	*	*	0.1	0.3	1.0	3.8	11.7	31.5	82.2

All races 3/, female

2006.....	3.4	*	*	*	*	0.1	0.2	0.7	2.0	6.3	20.2	65.0
2005.....	3.7	*	*	*	*	0.1	0.4	0.7	2.2	7.2	21.3	72.2
2004.....	3.9	*	*	*	*	0.1	0.4	0.8	2.4	8.3	22.5	75.8
2003.....	3.8	*	*	*	*	*	0.3	0.9	2.6	7.5	21.8	76.9
2002.....	4.0	*	*	*	*	0.2	0.3	0.8	2.8	8.9	22.5	78.2
2001.....	4.0	*	*	*	*	0.1	0.4	0.9	2.9	8.6	23.1	79.9
2000.....	4.0	*	*	*	*	0.1	0.4	0.9	2.6	8.7	24.2	82.6
1999.....	4.1	*	*	*	*	0.1	0.4	0.9	2.8	8.3	24.5	86.0



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

White, both sexes

2006 4/.....	3.3	*	*	*	*	0.1	0.2	0.8	2.3	7.5	21.8	65.4
2005 4/.....	3.5	*	*	*	*	0.1	0.3	0.7	2.5	8.3	23.7	69.9
2004 4/.....	3.5	*	*	*	*	0.1	0.3	0.9	2.5	8.8	24.0	72.5
2003 4/.....	3.6	*	*	*	*	0.1	0.3	0.8	2.7	8.7	24.2	74.6
2002.....	3.7	*	*	*	*	0.1	0.3	0.8	2.8	9.6	24.7	77.8
2001.....	3.7	*	*	*	*	0.1	0.3	0.9	2.9	9.5	25.2	78.5
2000.....	3.8	*	*	*	*	0.1	0.3	0.8	2.8	10.0	26.6	82.3
1999.....	3.8	*	*	*	*	0.1	0.3	0.9	3.0	9.5	26.3	82.9

White, male

2006 4/.....	2.9	*	*	*	*	*	0.2	0.9	2.7	9.3	25.6	69.0
2005 4/.....	3.0	*	*	*	*	*	0.2	0.7	3.0	9.9	28.2	67.8
2004 4/.....	3.0	*	*	*	*	*	0.3	1.0	2.7	10.1	27.7	70.7
2003 4/.....	3.1	*	*	*	*	*	0.2	0.9	2.9	10.6	28.8	74.1
2002.....	3.2	*	*	*	*	*	0.3	0.9	3.1	10.8	29.1	79.1
2001.....	3.2	*	*	*	*	*	0.2	0.8	3.2	11.2	29.8	80.9
2000.....	3.4	*	*	*	*	0.1	0.2	0.9	3.3	12.2	31.6	86.9
1999.....	3.2	*	*	*	*	*	0.3	0.9	3.6	11.4	30.4	81.3

White, female

2006 4/.....	3.6	*	*	*	*	*	0.2	0.7	1.9	6.0	19.1	63.7
2005 4/.....	3.9	*	*	*	*	*	0.3	0.7	2.0	6.8	20.7	70.8
2004 4/.....	4.0	*	*	*	*	*	0.3	0.7	2.3	7.8	21.6	73.3
2003 4/.....	4.0	*	*	*	*	*	0.3	0.7	2.5	7.2	21.2	74.8
2002.....	4.2	*	*	*	*	0.1	0.3	0.7	2.6	8.5	21.7	77.2
2001.....	4.2	*	*	*	*	0.1	0.4	0.9	2.6	8.2	22.1	77.5
2000.....	4.2	*	*	*	*	*	0.3	0.7	2.3	8.2	23.3	80.4
1999.....	4.3	*	*	*	*	*	0.3	0.8	2.5	7.9	23.6	83.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of arteries, arterioles and capillaries (I72-I78)

Black, both sexes

2006 4/.....	3.1	*	*	*	*	*	0.5	1.7	4.9	12.1	36.8	85.1
2005 4/.....	3.3	*	*	*	*	*	0.6	1.4	5.2	14.8	38.9	94.3
2004 4/.....	3.4	*	*	*	*	*	0.6	1.7	5.2	14.9	38.4	111.4
2003 4/.....	3.4	*	*	*	*	*	0.7	2.0	4.7	14.8	40.1	103.6
2002.....	3.5	*	*	*	*	*	0.6	1.7	6.1	16.5	38.5	106.3
2001.....	3.7	*	*	*	*	*	0.9	1.7	6.4	17.1	42.2	119.4
2000.....	3.6	*	*	*	*	*	0.6	2.3	6.2	16.0	41.7	115.4
1999.....	3.5	*	*	*	*	*	0.6	1.5	5.8	15.0	42.8	118.6

Black, male

2006 4/.....	2.7	*	*	*	*	*	*	2.0	6.7	15.7	37.9	66.1
2005 4/.....	3.0	*	*	*	*	*	*	1.8	6.7	18.2	49.7	78.1
2004 4/.....	2.9	*	*	*	*	*	*	2.0	6.3	17.0	43.2	101.3
2003 4/.....	2.9	*	*	*	*	*	*	1.8	5.6	18.3	51.1	87.6
2002.....	3.0	*	*	*	*	*	*	1.5	7.2	19.4	43.6	116.9
2001.....	3.2	*	*	*	*	*	0.9	2.0	6.5	20.0	49.4	116.9
2000.....	3.1	*	*	*	*	*	*	2.8	7.0	19.0	48.7	108.4
1999.....	2.9	*	*	*	*	*	*	1.4	6.6	17.5	51.3	107.8

Black, female

2006 4/.....	3.4	*	*	*	*	*	*	1.4	3.4	9.5	36.2	92.9
2005 4/.....	3.6	*	*	*	*	*	*	1.0	4.0	12.3	32.7	100.9
2004 4/.....	3.9	*	*	*	*	*	0.8	1.4	4.2	13.5	35.7	115.4
2003 4/.....	3.8	*	*	*	*	*	0.7	2.1	4.0	12.3	34.0	109.8
2002.....	3.9	*	*	*	*	*	0.8	1.9	5.1	14.4	35.6	102.3
2001.....	4.2	*	*	*	*	*	0.8	1.4	6.2	15.1	38.2	120.4
2000.....	4.1	*	*	*	*	*	0.7	1.8	5.5	13.9	37.7	118.0
1999.....	4.0	*	*	*	*	*	0.8	1.6	5.2	13.3	38.0	122.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/ 1 year	1-4 2/ years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	-----------------	--------------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes

2006.....	1.3	0.8	*	*	0.1	0.3	0.6	1.2	1.7	2.9	7.0	18.4
2005.....	1.6	0.6	*	*	0.1	0.3	0.8	1.5	2.1	3.8	8.6	23.0
2004.....	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.1	3.6	9.0	23.3
2003.....	1.6	0.9	*	*	0.1	0.3	0.8	1.4	2.0	4.1	9.0	24.0
2002.....	1.6	0.7	*	*	0.1	0.3	0.7	1.3	2.2	3.8	9.8	24.4
2001.....	1.6	0.5	*	*	0.1	0.3	0.6	1.2	2.2	4.6	9.4	26.1
2000.....	1.6	0.7	*	*	0.1	0.3	0.7	1.4	2.1	4.3	9.4	25.6
1999.....	1.4	0.6	*	*	0.1	0.2	0.6	1.0	1.9	3.8	9.1	23.4

All races 3/, male

2006.....	1.2	*	*	*	*	0.3	0.7	1.5	2.1	2.9	7.0	16.2
2005.....	1.5	*	*	*	*	0.3	0.7	1.8	2.6	4.4	8.4	19.2
2004.....	1.4	*	*	*	*	0.4	0.7	1.6	2.5	3.9	9.7	19.6
2003.....	1.4	1.0	*	*	0.1	0.3	0.8	1.6	2.4	4.2	8.5	19.1
2002.....	1.4	*	*	*	0.1	0.3	0.7	1.4	2.4	3.9	9.5	20.5
2001.....	1.4	*	*	*	*	0.3	0.7	1.4	2.5	4.9	9.8	22.0
2000.....	1.4	*	*	*	*	0.3	0.8	1.6	2.3	4.7	9.7	22.2
1999.....	1.3	*	*	*	*	0.2	0.7	1.1	2.0	4.2	9.6	20.8

All races 3/, female

2006.....	1.5	*	*	*	0.1	0.3	0.6	1.0	1.4	3.0	7.0	19.5
2005.....	1.8	*	*	*	0.1	0.4	0.8	1.1	1.7	3.3	8.8	24.8
2004.....	1.7	*	*	*	*	0.3	0.7	1.1	1.8	3.4	8.5	25.0
2003.....	1.9	*	*	*	0.1	0.4	0.8	1.2	1.7	4.0	9.4	26.2
2002.....	1.9	*	*	*	*	0.4	0.7	1.2	2.1	3.8	10.1	26.2
2001.....	1.9	*	*	*	0.1	0.4	0.6	1.0	2.0	4.4	9.2	27.8
2000.....	1.8	*	*	*	0.1	0.4	0.6	1.1	2.0	4.0	9.2	26.9
1999.....	1.6	*	*	*	*	0.2	0.5	1.0	1.8	3.4	8.8	24.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

White, both sexes

2006 4/.....	1.3	0.9	*	*	0.1	0.2	0.5	1.1	1.6	2.8	6.8	18.1
2005 4/.....	1.6	0.7	*	*	0.1	0.3	0.6	1.3	1.9	3.6	8.3	23.5
2004 4/.....	1.6	0.7	*	*	0.1	0.3	0.6	1.2	2.0	3.4	8.8	23.2
2003 4/.....	1.6	0.9	*	*	0.1	0.3	0.6	1.2	1.9	3.8	8.6	23.6
2002.....	1.6	0.6	*	*	0.1	0.3	0.6	1.2	2.0	3.6	9.5	24.3
2001.....	1.7	*	*	*	0.1	0.3	0.5	1.0	2.0	4.3	9.1	26.0
2000.....	1.6	0.7	*	*	0.1	0.3	0.6	1.2	1.9	4.0	9.1	25.5
1999.....	1.5	*	*	*	*	0.2	0.5	0.9	1.8	3.5	8.9	23.2

White, male

2006 4/.....	1.2	*	*	*	*	0.2	0.6	1.3	1.9	2.8	6.6	16.0
2005 4/.....	1.4	*	*	*	*	0.3	0.6	1.6	2.3	4.1	8.1	19.1
2004 4/.....	1.4	*	*	*	*	0.3	0.6	1.4	2.4	3.7	9.4	19.7
2003 4/.....	1.3	*	*	*	*	0.2	0.7	1.4	2.2	3.9	7.9	18.5
2002.....	1.3	*	*	*	*	0.3	0.6	1.3	2.2	3.6	9.0	19.9
2001.....	1.4	*	*	*	*	0.2	0.5	1.3	2.2	4.6	9.5	21.7
2000.....	1.4	*	*	*	*	0.2	0.8	1.5	2.0	4.4	9.5	22.0
1999.....	1.2	*	*	*	*	0.2	0.6	0.9	1.9	3.9	9.2	20.7

White, female

2006 4/.....	1.5	*	*	*	*	0.3	0.4	0.8	1.2	2.8	6.9	19.1
2005 4/.....	1.8	*	*	*	*	0.3	0.6	1.0	1.5	3.2	8.5	25.5
2004 4/.....	1.8	*	*	*	*	0.3	0.5	0.9	1.6	3.2	8.4	24.8
2003 4/.....	1.9	*	*	*	*	0.3	0.5	1.1	1.5	3.8	9.1	25.8
2002.....	1.9	*	*	*	*	0.3	0.6	1.1	1.9	3.6	9.8	26.3
2001.....	1.9	*	*	*	*	0.3	0.5	0.8	1.8	4.1	8.9	27.9
2000.....	1.9	*	*	*	*	0.3	0.5	1.0	1.8	3.6	8.9	26.9
1999.....	1.7	*	*	*	*	0.2	0.4	0.9	1.7	3.2	8.7	24.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other disorders of circulatory system (I80-I99)

Black, both sexes

2006 4/.....	1.8	*	*	*	*	0.7	1.6	2.9	3.4	4.9	10.9	25.7
2005 4/.....	2.0	*	*	*	*	0.8	1.7	3.0	4.5	6.4	14.0	24.3
2004 4/.....	2.1	*	*	*	*	0.9	1.9	2.9	4.2	6.6	13.6	28.8
2003 4/.....	2.2	*	*	*	*	0.9	1.9	2.6	4.3	7.5	15.4	33.6
2002.....	2.1	*	*	*	*	1.0	1.6	2.6	4.9	6.8	16.6	30.0
2001.....	2.0	*	*	*	*	0.8	1.5	2.2	4.9	7.8	14.4	30.1
2000.....	2.1	*	*	*	*	0.9	1.5	2.8	4.5	8.9	14.4	30.0
1999.....	1.7	*	*	*	*	0.5	1.4	2.4	2.9	6.6	13.8	28.4

Black, male

2006 4/.....	1.7	*	*	*	*	0.7	1.5	3.5	4.2	5.1	11.6	22.3
2005 4/.....	2.1	*	*	*	*	*	1.6	3.8	5.6	8.8	16.1	25.1
2004 4/.....	1.9	*	*	*	*	1.1	2.0	3.4	4.6	7.1	15.0	24.8
2003 4/.....	2.0	*	*	*	*	0.9	1.3	3.4	5.2	8.7	17.5	32.3
2002.....	2.0	*	*	*	*	1.0	1.8	3.0	5.6	7.7	18.4	32.2
2001.....	1.9	*	*	*	*	*	1.7	2.4	6.4	9.0	15.9	29.5
2000.....	1.9	*	*	*	*	0.9	1.8	3.4	5.1	9.1	13.3	30.0
1999.....	1.6	*	*	*	*	*	1.8	2.8	2.6	8.2	16.9	*

Black, female

2006 4/.....	1.9	*	*	*	*	0.7	1.7	2.3	2.8	4.8	10.5	27.1
2005 4/.....	2.0	*	*	*	*	0.8	1.8	2.3	3.5	4.7	12.7	24.0
2004 4/.....	2.2	*	*	*	*	0.8	1.9	2.5	3.9	6.3	12.7	30.3
2003 4/.....	2.3	*	*	*	*	0.9	2.5	1.9	3.6	6.7	14.2	34.0
2002.....	2.2	*	*	*	*	0.9	1.5	2.3	4.4	6.1	15.6	29.2
2001.....	2.1	*	*	*	*	0.9	1.4	2.1	3.6	7.1	13.6	30.3
2000.....	2.2	*	*	*	*	0.9	1.3	2.3	3.9	8.8	15.0	30.0
1999.....	1.8	*	*	*	*	*	1.1	2.0	3.1	5.5	12.0	30.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Influenza and pneumonia (J10-J18)

All races 3/, both sexes

2006.....	18.8	6.4	0.8	0.2	0.4	0.8	1.9	4.6	10.0	32.0	127.8	502.5
2005.....	21.3	6.5	0.7	0.3	0.4	0.9	2.1	5.1	11.3	35.5	142.2	593.9
2004.....	20.3	6.7	0.7	0.2	0.4	0.8	2.0	4.6	10.8	34.6	139.3	582.6
2003.....	22.4	8.0	1.0	0.4	0.5	0.9	2.2	5.2	11.2	37.3	151.1	666.1
2002.....	22.8	6.5	0.7	0.2	0.4	0.9	2.2	4.8	11.2	37.5	156.9	696.6
2001.....	21.8	7.4	0.7	0.2	0.5	0.9	2.2	4.6	10.7	36.3	148.5	685.6
2000.....	23.2	7.6	0.7	0.2	0.5	0.9	2.4	4.7	11.9	39.1	160.3	744.1
1999.....	22.8	8.4	0.8	0.2	0.5	0.8	2.4	4.6	11.0	37.2	157.0	751.8

All races 3/, male

2006.....	17.4	7.7	0.8	0.2	0.5	0.9	2.1	5.6	12.3	38.9	156.9	586.4
2005.....	19.2	7.0	0.7	0.3	0.5	1.0	2.5	6.2	13.6	42.8	174.3	673.5
2004.....	18.6	8.0	0.6	0.2	0.6	1.0	2.2	5.6	13.0	42.8	172.7	677.2
2003.....	20.1	9.4	1.1	0.3	0.6	1.1	2.5	6.4	13.2	46.5	186.8	757.8
2002.....	20.4	7.8	0.8	0.2	0.5	0.9	2.4	5.8	13.6	46.4	194.7	805.8
2001.....	19.6	8.1	0.7	0.2	0.5	1.0	2.5	5.7	13.1	45.0	186.4	808.4
2000.....	20.8	8.4	0.7	0.2	0.5	1.0	2.8	5.8	14.4	48.3	199.8	892.0
1999.....	20.3	8.8	0.8	0.3	0.5	1.0	2.6	5.9	12.8	46.9	196.5	894.7

All races 3/, female

2006.....	20.2	5.0	0.8	0.2	0.4	0.8	1.7	3.7	7.8	26.3	107.8	463.3
2005.....	23.2	5.8	0.7	0.3	0.3	0.8	1.8	4.1	9.1	29.4	120.4	557.4
2004.....	22.0	5.4	0.9	0.2	0.3	0.6	1.8	3.6	8.8	27.7	116.8	540.0
2003.....	24.6	6.6	0.9	0.4	0.5	0.7	1.9	4.1	9.4	29.5	127.2	625.6
2002.....	25.1	5.1	0.6	0.3	0.4	0.8	1.9	3.8	9.0	30.1	131.8	649.2
2001.....	23.9	6.7	0.7	0.2	0.4	0.8	1.9	3.5	8.5	29.1	123.7	634.2
2000.....	25.6	6.8	0.7	0.2	0.5	0.8	1.9	3.7	9.5	31.5	134.6	683.9
1999.....	25.3	8.0	0.9	0.2	0.5	0.7	2.1	3.4	9.4	29.3	131.3	694.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Influenza and pneumonia (J10-J18)

White, both sexes

2006 4/.....	20.4	5.0	0.7	0.2	0.4	0.8	1.8	4.2	9.5	31.1	127.5	509.3
2005 4/.....	23.1	4.9	0.6	0.2	0.4	0.8	1.9	4.6	10.5	34.2	142.6	606.4
2004 4/.....	22.0	5.1	0.6	0.2	0.4	0.7	1.8	4.1	10.1	33.4	138.5	590.7
2003 4/.....	24.4	6.7	0.9	0.3	0.5	0.8	2.0	4.7	10.3	35.9	151.5	679.3
2002.....	24.9	5.2	0.5	0.2	0.4	0.8	1.9	4.4	10.4	36.7	156.9	709.1
2001.....	23.6	5.3	0.7	0.2	0.4	0.7	1.8	4.0	9.9	35.3	147.2	695.0
2000.....	25.2	5.8	0.6	0.2	0.4	0.8	2.1	4.1	10.9	37.1	160.0	758.3
1999.....	24.8	6.2	0.6	0.2	0.4	0.7	1.9	4.0	10.1	36.1	157.8	764.3

White, male

2006 4/.....	18.6	6.3	0.7	0.2	0.4	0.8	2.0	4.9	11.7	37.3	155.8	591.4
2005 4/.....	20.5	5.9	0.6	0.2	0.5	0.9	2.2	5.5	12.3	40.6	173.8	684.4
2004 4/.....	19.8	6.2	0.5	0.2	0.6	0.9	1.9	4.9	12.1	40.6	170.2	685.5
2003 4/.....	21.4	8.0	1.1	0.3	0.6	1.0	2.2	5.6	11.6	44.0	184.9	767.1
2002.....	21.9	6.6	0.6	0.2	0.4	0.8	2.1	5.2	12.3	45.2	193.1	815.9
2001.....	20.7	6.2	0.7	0.2	0.5	0.8	2.1	4.9	11.8	43.0	182.6	814.4
2000.....	22.0	6.4	0.5	0.2	0.5	0.9	2.5	4.8	12.9	45.1	197.8	906.1
1999.....	21.5	6.7	0.5	0.3	0.4	0.8	2.1	4.9	11.5	44.7	196.1	905.7

White, female

2006 4/.....	22.2	3.7	0.6	0.1	0.4	0.7	1.6	3.4	7.5	25.8	107.9	470.7
2005 4/.....	25.7	4.0	0.6	0.3	0.3	0.8	1.7	3.8	8.7	28.7	121.1	570.4
2004 4/.....	24.2	3.9	0.7	0.2	0.3	0.6	1.6	3.2	8.3	27.3	116.9	548.0
2003 4/.....	27.3	5.4	0.8	0.4	0.5	0.6	1.8	3.7	9.0	29.0	129.0	640.4
2002.....	27.8	3.7	0.5	0.3	0.3	0.7	1.7	3.6	8.5	29.6	132.7	662.6
2001.....	26.4	4.4	0.6	0.2	0.3	0.7	1.6	3.2	8.1	28.9	123.8	645.0
2000.....	28.2	5.1	0.7	0.2	0.4	0.8	1.7	3.4	8.9	30.3	135.0	698.3
1999.....	27.9	5.6	0.7	0.2	0.4	0.6	1.8	3.0	8.7	28.9	132.6	707.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Influenza and pneumonia (J10-J18)

Black, both sexes

2006 4/.....	13.4	13.2	1.3	*	0.6	1.4	3.3	8.8	15.9	44.7	139.8	434.6
2005 4/.....	14.8	14.8	1.1	0.3	0.5	1.4	3.8	9.8	20.0	52.0	149.8	471.2
2004 4/.....	14.8	14.1	1.3	0.3	0.6	1.2	4.0	9.1	19.0	49.8	157.2	505.1
2003 4/.....	15.4	14.1	1.1	0.4	0.7	1.9	4.4	10.4	20.6	54.2	155.9	525.4
2002.....	15.6	12.9	1.6	*	0.6	1.5	4.3	8.7	21.2	50.3	166.2	575.2
2001.....	15.5	16.3	1.0	*	0.7	1.6	4.8	9.7	18.9	48.3	169.0	581.3
2000.....	16.4	16.7	1.2	*	0.7	1.7	4.5	10.0	22.0	60.3	171.1	600.9
1999.....	16.2	19.0	1.7	*	0.8	2.0	5.5	10.6	20.9	54.1	157.4	634.8

Black, male

2006 4/.....	13.2	15.8	*	*	0.7	1.4	3.7	11.5	21.3	58.3	181.0	504.9
2005 4/.....	14.6	13.9	*	*	0.7	1.7	4.8	12.9	27.6	69.5	189.0	531.7
2004 4/.....	15.1	16.3	*	*	*	1.7	4.8	11.8	24.9	72.5	219.4	598.0
2003 4/.....	16.0	16.1	*	*	0.8	2.3	5.3	13.2	29.1	78.2	218.6	641.2
2002.....	15.4	13.9	1.7	*	0.6	1.7	5.1	11.2	29.0	65.9	228.9	689.4
2001.....	15.9	15.9	*	*	0.8	2.1	5.7	13.3	26.3	69.4	242.2	714.8
2000.....	16.7	17.9	1.7	*	0.7	2.3	5.6	13.9	30.0	83.5	234.1	755.4
1999.....	16.4	18.9	2.0	*	0.8	2.2	6.6	15.1	27.2	76.9	213.7	803.8

Black, female

2006 4/.....	13.7	10.5	*	*	*	1.4	2.9	6.5	11.5	35.0	116.3	405.8
2005 4/.....	14.9	15.7	*	*	*	1.1	2.9	7.2	13.8	39.4	127.5	446.8
2004 4/.....	14.5	11.7	1.8	*	0.7	0.8	3.3	6.8	14.3	33.6	122.1	468.2
2003 4/.....	14.9	12.1	*	*	*	1.5	3.6	7.9	13.8	37.2	120.4	480.4
2002.....	15.7	11.8	*	*	*	1.3	3.7	6.5	15.0	39.1	130.7	531.5
2001.....	15.2	16.7	*	*	0.7	1.3	3.9	6.7	13.0	33.4	127.9	531.7
2000.....	16.0	15.5	*	*	0.8	1.2	3.4	6.7	15.6	43.9	135.7	543.4
1999.....	16.1	19.1	*	*	0.9	1.7	4.4	6.8	15.8	38.0	125.8	571.6



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Influenza (J10-J11)

All races 3/, both sexes

2006.....	0.3	*	*	0.1	*	*	*	0.1	0.1	0.4	1.8	7.0
2005.....	0.6	*	*	0.0	*	*	*	0.1	0.2	0.7	3.7	19.3
2004.....	0.4	*	*	*	*	*	*	0.1	0.2	0.5	2.2	11.4
2003.....	0.6	0.8	0.4	0.1	0.1	*	0.1	0.1	0.3	1.0	3.6	17.0
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.4	8.8
2001.....	0.1	*	*	*	*	*	*	*	0.1	0.1	0.4	2.2
2000.....	0.6	*	*	*	*	*	0.0	0.1	0.4	1.1	4.0	20.2
1999.....	0.6	*	*	*	*	*	*	0.1	0.3	0.9	3.7	21.2

All races 3/, male

2006.....	0.2	*	*	*	*	*	*	*	0.1	0.5	2.0	8.1
2005.....	0.5	*	*	*	*	*	*	0.1	0.3	0.7	3.6	18.2
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	11.1
2003.....	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.7	17.0
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	8.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.2
2000.....	0.5	*	*	*	*	*	*	0.1	0.4	1.3	4.0	20.7
1999.....	0.4	*	*	*	*	*	*	*	0.3	0.9	4.2	19.9

All races 3/, female

2006.....	0.3	*	*	*	*	*	*	*	0.1	0.4	1.7	6.5
2005.....	0.8	*	*	*	*	*	*	*	0.2	0.8	3.7	19.8
2004.....	0.5	*	*	*	*	*	*	*	0.2	0.4	2.1	11.5
2003.....	0.7	*	0.3	0.1	*	*	*	0.1	0.3	0.9	3.5	17.1
2002.....	0.3	*	*	*	*	*	*	*	*	0.2	1.3	8.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.2
2000.....	0.8	*	*	*	*	*	*	0.1	0.4	0.9	4.0	19.9
1999.....	0.7	*	*	*	*	*	*	*	0.3	0.9	3.4	21.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Influenza (J10-J11)

White, both sexes

2006 4/.....	0.3	*	*	0.1	*	*	*	0.1	0.2	0.5	1.9	7.5
2005 4/.....	0.7	*	*	*	*	*	*	0.1	0.2	0.8	4.0	20.8
2004 4/.....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	2.3	12.3
2003 4/.....	0.7	0.7	0.3	0.1	0.1	*	0.1	0.1	0.3	1.1	3.8	18.2
2002.....	0.3	*	*	*	*	*	*	*	0.1	0.3	1.5	9.4
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.3
2000.....	0.7	*	*	*	*	*	*	0.1	0.4	1.1	4.2	21.4
1999.....	0.7	*	*	*	*	*	*	0.1	0.3	0.9	3.9	22.8

White, male

2006 4/.....	0.3	*	*	*	*	*	*	*	*	0.5	2.0	8.6
2005 4/.....	0.5	*	*	*	*	*	*	*	0.3	0.8	3.9	19.2
2004 4/.....	0.3	*	*	*	*	*	*	*	0.2	0.7	2.4	12.1
2003 4/.....	0.5	*	0.4	*	*	*	*	0.2	0.3	1.1	3.9	18.3
2002.....	0.2	*	*	*	*	*	*	*	*	0.4	1.7	9.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.3
2000.....	0.5	*	*	*	*	*	*	*	0.4	1.3	4.2	21.6
1999.....	0.5	*	*	*	*	*	*	*	0.4	0.9	4.4	21.7

White, female

2006 4/.....	0.4	*	*	*	*	*	*	*	0.2	0.4	1.9	7.1
2005 4/.....	0.9	*	*	*	*	*	*	*	0.2	0.8	4.0	21.5
2004 4/.....	0.5	*	*	*	*	*	*	*	0.2	0.4	2.3	12.4
2003 4/.....	0.8	*	*	*	*	*	*	0.1	0.3	1.0	3.7	18.1
2002.....	0.4	*	*	*	*	*	*	*	*	*	1.4	9.3
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.3
2000.....	0.9	*	*	*	*	*	*	0.1	0.4	0.9	4.2	21.3
1999.....	0.9	*	*	*	*	*	*	*	0.3	0.9	3.5	23.2



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonia (J12-J18)

All races 3/, both sexes

2006.....	18.5	5.9	0.7	0.1	0.4	0.8	1.9	4.6	9.8	31.6	125.9	495.5
2005.....	20.6	6.0	0.6	0.2	0.4	0.9	2.1	5.0	11.1	34.8	138.5	574.7
2004.....	19.9	6.3	0.6	0.2	0.4	0.7	2.0	4.5	10.7	34.0	137.0	571.2
2003.....	21.8	7.3	0.7	0.2	0.5	0.9	2.2	5.1	10.9	36.3	147.5	649.1
2002.....	22.5	6.3	0.7	0.2	0.4	0.9	2.1	4.7	11.1	37.2	155.5	687.8
2001.....	21.7	7.2	0.7	0.2	0.4	0.8	2.2	4.6	10.6	36.2	148.1	683.4
2000.....	22.6	7.4	0.6	0.2	0.4	0.9	2.3	4.6	11.5	38.0	156.4	723.9
1999.....	22.2	8.1	0.8	0.2	0.4	0.8	2.3	4.6	10.7	36.4	153.3	730.6

All races 3/, male

2006.....	17.1	7.2	0.6	0.1	0.4	0.8	2.1	5.5	12.2	38.4	155.0	578.3
2005.....	18.8	6.6	0.6	0.2	0.5	0.9	2.4	6.1	13.3	42.1	170.7	655.3
2004.....	18.3	7.5	0.5	0.2	0.5	0.9	2.2	5.5	12.8	42.1	170.3	666.1
2003.....	19.6	8.5	0.7	0.2	0.5	1.1	2.5	6.2	12.9	45.4	183.2	740.9
2002.....	20.2	7.6	0.8	0.2	0.4	0.9	2.4	5.7	13.4	46.0	193.1	796.9
2001.....	19.5	7.8	0.7	0.2	0.5	0.9	2.5	5.7	13.0	44.9	186.0	806.2
2000.....	20.3	8.1	0.6	0.2	0.5	1.0	2.7	5.6	14.0	47.1	195.8	871.3
1999.....	19.8	8.5	0.7	0.2	0.4	0.9	2.6	5.8	12.5	46.0	192.3	874.7

All races 3/, female

2006.....	19.9	4.6	0.7	*	0.4	0.8	1.7	3.7	7.7	25.9	106.1	456.7
2005.....	22.5	5.3	0.6	0.2	0.2	0.8	1.8	4.0	8.9	28.7	116.7	537.6
2004.....	21.5	5.0	0.8	0.2	0.3	0.5	1.8	3.5	8.6	27.3	114.6	528.5
2003.....	23.9	6.0	0.6	0.3	0.4	0.7	1.9	4.0	9.1	28.6	123.7	608.5
2002.....	24.8	5.0	0.6	0.2	0.3	0.8	1.9	3.8	9.0	29.8	130.6	640.5
2001.....	23.8	6.7	0.7	0.2	0.4	0.7	1.9	3.5	8.4	29.0	123.2	632.0
2000.....	24.8	6.6	0.6	0.2	0.4	0.8	1.9	3.6	9.1	30.6	130.6	663.9
1999.....	24.6	7.7	0.8	0.2	0.4	0.7	2.0	3.4	9.1	28.4	128.0	672.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Pneumonia (J12-J18)

White, both sexes

2006 4/.....	20.1	4.6	0.6	0.1	0.4	0.7	1.8	4.1	9.4	30.7	125.6	501.7
2005 4/.....	22.4	4.5	0.5	0.2	0.4	0.8	1.9	4.5	10.2	33.4	138.6	585.6
2004 4/.....	21.6	4.7	0.5	0.1	0.4	0.7	1.7	4.0	9.9	32.8	136.2	578.4
2003 4/.....	23.7	6.0	0.6	0.2	0.5	0.8	1.9	4.5	10.0	34.8	147.7	661.1
2002.....	24.6	5.0	0.5	0.2	0.4	0.8	1.9	4.3	10.2	36.4	155.4	699.6
2001.....	23.5	5.1	0.6	0.2	0.4	0.7	1.8	4.0	9.8	35.2	146.8	692.7
2000.....	24.4	5.6	0.5	0.2	0.4	0.8	2.0	4.0	10.4	36.0	155.8	737.0
1999.....	24.1	5.9	0.6	0.2	0.4	0.7	1.9	3.9	9.8	35.2	153.9	741.6

White, male

2006 4/.....	18.3	5.7	0.5	0.1	0.4	0.8	2.0	4.9	11.5	36.8	153.7	582.8
2005 4/.....	20.0	5.3	0.4	0.2	0.5	0.9	2.1	5.4	12.0	39.8	169.9	665.2
2004 4/.....	19.5	5.8	0.4	0.2	0.6	0.9	1.9	4.9	11.9	39.8	167.8	673.4
2003 4/.....	20.9	7.1	0.7	0.2	0.5	1.0	2.2	5.5	11.3	42.8	181.0	748.8
2002.....	21.7	6.4	0.5	0.1	0.4	0.8	2.1	5.1	12.1	44.8	191.4	806.2
2001.....	20.6	5.8	0.6	0.2	0.5	0.8	2.1	4.8	11.7	42.9	182.2	812.1
2000.....	21.5	6.1	0.4	0.2	0.5	0.8	2.4	4.7	12.5	43.8	193.6	884.4
1999.....	21.0	6.7	0.4	0.2	0.4	0.8	2.1	4.8	11.1	43.8	191.7	884.0

White, female

2006 4/.....	21.8	3.5	0.6	*	0.3	0.7	1.6	3.4	7.4	25.4	106.1	463.7
2005 4/.....	24.8	3.7	0.5	0.2	0.2	0.7	1.6	3.7	8.5	27.9	117.1	548.9
2004 4/.....	23.6	3.5	0.6	0.1	0.3	0.5	1.6	3.1	8.1	26.8	114.6	535.6
2003 4/.....	26.5	4.8	0.5	0.2	0.4	0.6	1.7	3.6	8.7	28.0	125.2	622.3
2002.....	27.4	3.6	0.5	0.2	0.3	0.7	1.7	3.5	8.5	29.3	131.3	653.3
2001.....	26.2	4.4	0.6	0.2	0.3	0.7	1.6	3.1	8.0	28.7	123.3	642.8
2000.....	27.3	5.0	0.6	0.2	0.3	0.7	1.7	3.2	8.5	29.4	130.8	677.1
1999.....	27.1	5.2	0.7	0.2	0.3	0.5	1.7	3.0	8.5	27.9	129.1	684.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Pneumonia (J12-J18)

Black, both sexes

2006 4/.....	13.3	12.5	1.1	*	0.6	1.3	3.2	8.7	15.8	44.5	138.8	432.7
2005 4/.....	14.6	14.0	1.0	*	0.5	1.4	3.7	9.7	19.7	51.7	148.4	467.2
2004 4/.....	14.6	13.5	1.2	*	0.6	1.2	4.0	9.0	18.9	49.4	155.6	502.5
2003 4/.....	15.1	13.0	0.9	*	0.6	1.8	4.4	10.1	20.3	53.7	154.5	519.6
2002.....	15.5	12.7	1.5	*	0.6	1.5	4.3	8.7	21.1	49.9	165.8	572.8
2001.....	15.5	16.3	1.0	*	0.7	1.6	4.7	9.7	18.8	48.2	168.6	579.7
2000.....	16.1	16.4	1.2	*	0.7	1.7	4.4	9.9	21.8	59.3	169.7	591.5
1999.....	16.1	18.3	1.7	*	0.8	1.9	5.4	10.5	20.7	53.3	155.7	629.1

Black, male

2006 4/.....	13.0	15.3	*	*	0.6	1.3	3.7	11.5	21.2	58.1	179.4	502.2
2005 4/.....	14.5	13.9	*	*	0.7	1.7	4.7	12.7	27.2	69.2	188.2	524.3
2004 4/.....	14.9	15.1	*	*	*	1.6	4.8	11.7	24.8	71.9	216.6	597.1
2003 4/.....	15.8	15.2	*	*	0.7	2.2	5.2	12.9	28.7	77.7	217.5	638.1
2002.....	15.3	13.7	1.6	*	*	1.7	5.1	11.2	28.7	65.5	228.4	687.2
2001.....	15.8	15.9	*	*	0.7	2.0	5.7	13.3	26.1	69.1	242.2	712.6
2000.....	16.5	17.9	1.7	*	0.7	2.2	5.5	13.6	29.7	82.8	232.5	742.7
1999.....	16.3	17.9	1.9	*	0.8	2.2	6.6	15.0	27.1	76.0	211.8	800.3

Black, female

2006 4/.....	13.6	9.6	*	*	*	1.3	2.9	6.4	11.4	34.8	115.7	404.3
2005 4/.....	14.8	14.2	*	*	*	1.1	2.8	7.2	13.7	39.2	125.8	444.1
2004 4/.....	14.4	11.7	1.7	*	0.6	0.8	3.2	6.7	14.1	33.3	121.2	465.0
2003 4/.....	14.6	10.8	*	*	*	1.4	3.6	7.7	13.6	36.6	119.0	473.5
2002.....	15.6	11.8	*	*	*	1.3	3.7	6.5	14.9	38.8	130.4	529.0
2001.....	15.1	16.7	*	*	0.7	1.3	3.9	6.7	13.0	33.4	127.2	530.5
2000.....	15.8	14.9	*	*	0.8	1.2	3.4	6.6	15.4	42.7	134.3	535.3
1999.....	15.9	18.8	*	*	0.9	1.7	4.3	6.8	15.5	37.2	124.3	565.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute lower respiratory infections (J20-J22)

All races 3/, both sexes

2006.....	0.1	1.3	*	*	*	*	*	*	*	0.1	0.5	2.0
2005.....	0.1	1.2	*	*	*	*	*	*	*	0.1	0.6	3.3
2004.....	0.1	1.5	*	*	*	*	*	*	*	0.1	0.4	2.6
2003.....	0.1	1.3	0.2	*	*	*	*	*	0.1	0.2	0.6	3.5
2002.....	0.1	1.3	0.1	*	*	*	*	*	*	0.2	0.5	3.3
2001.....	0.1	1.2	*	*	*	*	*	*	*	0.2	0.5	3.3
2000.....	0.2	1.1	*	*	*	*	*	*	*	0.2	0.7	4.5
1999.....	0.2	1.7	0.2	*	*	*	*	*	0.1	0.2	0.9	6.0

All races 3/, male

2006.....	0.1	1.4	*	*	*	*	*	*	*	*	0.4	1.8
2005.....	0.1	1.4	*	*	*	*	*	*	*	*	0.8	3.1
2004.....	0.1	1.6	*	*	*	*	*	*	*	*	*	2.5
2003.....	0.1	1.7	*	*	*	*	*	*	*	*	0.7	2.4
2002.....	0.1	1.9	*	*	*	*	*	*	*	*	0.5	2.8
2001.....	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2000.....	0.1	1.3	*	*	*	*	*	*	*	0.3	0.7	3.8
1999.....	0.1	1.9	*	*	*	*	*	*	*	*	1.1	4.0

All races 3/, female

2006.....	0.1	1.1	*	*	*	*	*	*	*	*	0.5	2.1
2005.....	0.2	1.0	*	*	*	*	*	*	*	*	0.5	3.5
2004.....	0.1	1.3	*	*	*	*	*	*	*	*	0.5	2.7
2003.....	0.2	*	*	*	*	*	*	*	*	*	0.6	3.9
2002.....	0.2	*	*	*	*	*	*	*	*	0.2	0.5	3.5
2001.....	0.2	1.2	*	*	*	*	*	*	*	*	0.5	3.7
2000.....	0.2	*	*	*	*	*	*	*	*	0.2	0.6	4.8
1999.....	0.2	1.5	*	*	*	*	*	*	*	*	0.8	6.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other acute lower respiratory infections (J20-J22)

White, both sexes

2006 4/.....	0.1	1.1	*	*	*	*	*	*	*	0.1	0.5	2.2
2005 4/.....	0.1	0.9	*	*	*	*	*	*	*	0.1	0.6	3.5
2004 4/.....	0.1	1.4	*	*	*	*	*	*	*	*	0.4	2.8
2003 4/.....	0.2	1.3	*	*	*	*	*	*	*	0.2	0.7	3.7
2002.....	0.1	1.1	0.2	*	*	*	*	*	*	0.2	0.5	3.5
2001.....	0.1	1.0	*	*	*	*	*	*	*	0.1	0.5	3.5
2000.....	0.2	0.8	*	*	*	*	*	*	*	0.3	0.7	4.7
1999.....	0.2	1.2	*	*	*	*	*	*	*	0.2	0.9	6.3

White, male

2006 4/.....	0.1	1.3	*	*	*	*	*	*	*	*	0.4	1.9
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.8	3.3
2004 4/.....	0.1	1.5	*	*	*	*	*	*	*	*	*	2.6
2003 4/.....	0.1	1.7	*	*	*	*	*	*	*	*	0.7	2.7
2002.....	0.1	1.6	*	*	*	*	*	*	*	*	0.5	2.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	2.4
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	3.8
1999.....	0.1	*	*	*	*	*	*	*	*	*	1.1	4.0

White, female

2006 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.3
2005 4/.....	0.2	*	*	*	*	*	*	*	*	*	0.5	3.6
2004 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.5	2.8
2003 4/.....	0.2	*	*	*	*	*	*	*	*	*	0.6	4.2
2002.....	0.2	*	*	*	*	*	*	*	*	*	0.5	3.7
2001.....	0.2	*	*	*	*	*	*	*	*	*	0.5	3.9
2000.....	0.2	*	*	*	*	*	*	*	*	0.2	0.7	5.1
1999.....	0.3	*	*	*	*	*	*	*	*	*	0.8	7.2





Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute bronchitis and bronchiolitis (J20-J21)

All races 3/, both sexes

2006.....	0.1	1.3	*	*	*	*	*	*	*	*	0.3	1.1
2005.....	0.1	1.2	*	*	*	*	*	*	*	*	0.4	1.9
2004.....	0.1	1.4	*	*	*	*	*	*	*	*	0.2	1.4
2003.....	0.1	1.3	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2002.....	0.1	1.2	0.1	*	*	*	*	*	*	0.1	0.3	2.0
2001.....	0.1	1.2	*	*	*	*	*	*	*	0.1	0.3	2.1
2000.....	0.1	1.1	*	*	*	*	*	*	*	0.2	0.4	2.6
1999.....	0.1	1.6	0.2	*	*	*	*	*	*	0.2	0.6	3.6

All races 3/, male

2006.....	0.1	1.4	*	*	*	*	*	*	*	*	*	1.2
2005.....	0.1	1.4	*	*	*	*	*	*	*	*	0.5	1.7
2004.....	0.1	1.6	*	*	*	*	*	*	*	*	*	1.8
2003.....	0.1	1.7	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	1.8	*	*	*	*	*	*	*	*	*	1.7
2001.....	0.1	1.2	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	1.2	*	*	*	*	*	*	*	*	0.6	2.4
1999.....	0.1	1.9	*	*	*	*	*	*	*	*	0.7	2.6

All races 3/, female

2006.....	0.1	1.1	*	*	*	*	*	*	*	*	0.3	1.1
2005.....	0.1	1.0	*	*	*	*	*	*	*	*	0.3	2.0
2004.....	0.1	1.3	*	*	*	*	*	*	*	*	0.3	1.3
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
2001.....	0.1	1.2	*	*	*	*	*	*	*	*	0.4	2.4
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.7
1999.....	0.2	1.4	*	*	*	*	*	*	*	*	0.5	4.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Acute bronchitis and bronchiolitis (J20-J21)

White, both sexes

2006 4/.....	0.1	1.1	*	*	*	*	*	*	*	*	0.3	1.2
2005 4/.....	0.1	0.9	*	*	*	*	*	*	*	*	0.4	2.0
2004 4/.....	0.1	1.3	*	*	*	*	*	*	*	*	0.2	1.5
2003 4/.....	0.1	1.3	*	*	*	*	*	*	*	*	0.4	2.3
2002.....	0.1	1.0	0.2	*	*	*	*	*	*	0.1	0.3	2.1
2001.....	0.1	1.0	*	*	*	*	*	*	*	*	0.3	2.1
2000.....	0.1	0.8	*	*	*	*	*	*	*	0.2	0.4	2.7
1999.....	0.1	1.2	*	*	*	*	*	*	*	0.1	0.6	3.8

White, male

2006 4/.....	0.1	1.3	*	*	*	*	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.7
2004 4/.....	0.1	1.4	*	*	*	*	*	*	*	*	*	1.8
2003 4/.....	0.1	1.6	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	1.5	*	*	*	*	*	*	*	*	*	1.7
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.6	2.4
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.7	2.6

White, female

2006 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.4	1.1
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.1
2004 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.3	1.3
2003 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.7
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	2.3
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	2.5
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.9
1999.....	0.2	*	*	*	*	*	*	*	*	*	0.5	4.3



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Unspecified acute lower respiratory infection (J22)

All races 3/, both sexes

2006.....	0.0	*	*	*	*	*	*	*	*	*	0.2	0.9
2005.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.4
2004.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.2
2003.....	0.0	*	*	*	*	*	*	*	*	*	0.3	1.3
2002.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2000.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.9
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.4

All races 3/, male

2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*

All races 3/, female

2006.....	0.0	*	*	*	*	*	*	*	*	*	*	1.1
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2004.....	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2003.....	0.0	*	*	*	*	*	*	*	*	*	*	1.4
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	1.3
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.1
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Unspecified acute lower respiratory infection (J22)

White, both sexes

2006 4/.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.0
2005 4/.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.5
2004 4/.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.3
2003 4/.....	0.0	*	*	*	*	*	*	*	*	*	0.3	1.4
2002.....	0.0	*	*	*	*	*	*	*	*	*	0.2	1.4
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	1.3
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.0
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.5

White, male

2006 4/.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005 4/.....	0.0	*	*	*	*	*	*	*	*	*	*	1.5
2004 4/.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2003 4/.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.0	*	*	*	*	*	*	*	*	*	*	*
1999.....	0.0	*	*	*	*	*	*	*	*	*	*	*

White, female

2006 4/.....	0.0	*	*	*	*	*	*	*	*	*	*	1.2
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*	*	1.5
2004 4/.....	0.1	*	*	*	*	*	*	*	*	*	*	1.5
2003 4/.....	0.1	*	*	*	*	*	*	*	*	*	*	1.5
2002.....	0.1	*	*	*	*	*	*	*	*	*	*	1.4
2001.....	0.0	*	*	*	*	*	*	*	*	*	*	1.5
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.2
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.3	2.9



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic lower respiratory diseases (J40-J47)

All races 3/, both sexes

2006.....	41.6	0.7	0.3	0.3	0.4	0.6	1.9	9.1	39.2	149.3	363.4	589.1
2005.....	44.2	0.8	0.3	0.3	0.4	0.6	2.0	9.4	42.0	160.5	385.6	637.2
2004.....	41.5	0.9	0.3	0.3	0.4	0.6	2.0	8.4	40.4	153.8	366.7	601.7
2003.....	43.5	0.8	0.3	0.3	0.5	0.7	2.1	8.7	43.3	163.2	383.0	635.1
2002.....	43.3	1.0	0.4	0.3	0.5	0.8	2.2	8.7	42.4	163.0	386.7	637.6
2001.....	43.2	1.0	0.3	0.3	0.4	0.7	2.2	8.5	44.1	167.9	379.8	644.7
2000.....	43.4	0.9	0.3	0.3	0.5	0.7	2.1	8.6	44.2	169.4	386.1	648.6
1999.....	44.5	0.9	0.4	0.3	0.5	0.8	2.0	8.5	47.5	177.2	397.8	646.0

All races 3/, male

2006.....	40.2	0.9	0.3	0.4	0.4	0.7	1.8	9.2	41.9	163.2	437.3	760.8
2005.....	42.8	*	0.4	0.3	0.4	0.6	1.9	9.7	45.0	180.7	463.7	819.9
2004.....	40.6	1.2	0.4	0.4	0.5	0.7	2.0	8.8	43.1	172.1	445.6	811.1
2003.....	42.4	1.1	0.5	0.4	0.5	0.8	1.9	9.1	46.5	183.6	464.9	865.9
2002.....	42.9	1.1	0.6	0.4	0.6	0.8	2.2	9.1	45.2	184.8	480.8	894.8
2001.....	42.7	1.1	0.4	0.3	0.5	0.7	2.0	8.8	46.9	191.3	475.1	916.9
2000.....	43.5	1.2	0.4	0.4	0.6	0.8	1.9	9.0	47.8	195.2	488.5	967.9
1999.....	45.6	*	0.4	0.4	0.6	0.8	1.8	8.6	52.3	210.7	513.2	996.7

All races 3/, female

2006.....	43.0	*	*	0.2	0.3	0.6	2.0	8.9	36.6	137.5	312.8	508.7
2005.....	45.5	*	0.3	0.2	0.3	0.6	2.1	9.0	39.2	143.4	332.5	553.3
2004.....	42.5	*	*	0.2	0.3	0.5	2.1	8.1	38.0	138.4	313.6	507.4
2003.....	44.4	*	*	0.2	0.4	0.6	2.3	8.2	40.3	146.0	328.3	533.0
2002.....	43.7	*	0.3	0.3	0.4	0.7	2.3	8.2	39.8	144.9	324.1	526.0
2001.....	43.7	*	*	0.2	0.4	0.7	2.3	8.1	41.5	148.5	317.3	530.8
2000.....	43.2	*	0.3	0.3	0.4	0.7	2.2	8.3	41.0	148.2	319.2	518.5
1999.....	43.4	*	0.3	0.2	0.5	0.9	2.1	8.4	43.1	149.8	322.9	504.6



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Chronic lower respiratory diseases (J40-J47)

White, both sexes

2006 4/.....	47.5	*	0.2	0.1	0.2	0.4	1.7	9.1	41.1	160.3	383.8	618.3
2005 4/.....	50.3	0.6	0.2	0.1	0.2	0.5	1.8	9.1	44.0	171.3	407.0	667.6
2004 4/.....	47.4	0.7	0.2	0.2	0.3	0.5	1.8	8.2	42.3	164.7	387.0	630.1
2003 4/.....	49.5	0.7	0.2	0.2	0.3	0.5	1.9	8.5	45.2	173.8	403.0	664.5
2002.....	49.2	0.8	0.2	0.2	0.3	0.6	1.9	8.4	44.2	173.2	405.6	666.2
2001.....	49.0	0.7	0.2	0.2	0.3	0.5	1.8	8.3	46.0	177.7	398.6	671.8
2000.....	49.0	0.8	0.2	0.2	0.3	0.5	1.7	8.2	45.8	178.7	404.4	675.6
1999.....	50.2	*	0.3	0.2	0.4	0.5	1.6	8.1	49.3	186.2	414.3	672.9

White, male

2006 4/.....	45.1	*	*	0.2	0.2	0.4	1.6	9.0	42.9	171.5	453.3	788.6
2005 4/.....	47.9	*	*	0.2	0.3	0.4	1.7	9.4	46.3	189.2	479.5	846.1
2004 4/.....	45.5	*	*	0.2	0.3	0.5	1.7	8.5	43.8	180.6	462.4	837.7
2003 4/.....	47.4	1.3	*	0.2	0.4	0.5	1.7	8.9	47.6	191.6	478.5	894.4
2002.....	47.8	*	0.3	0.2	0.4	0.6	1.8	8.8	46.0	192.3	495.2	923.4
2001.....	47.6	*	*	0.2	0.3	0.5	1.7	8.6	48.0	198.3	489.4	943.6
2000.....	48.3	*	*	0.2	0.4	0.5	1.6	8.4	48.6	201.4	503.6	997.4
1999.....	50.6	*	*	0.3	0.4	0.5	1.5	8.1	53.1	217.3	525.4	1,029.4

White, female

2006 4/.....	49.9	*	*	*	0.2	0.5	1.8	9.1	39.3	150.6	335.5	538.4
2005 4/.....	52.8	*	*	*	0.2	0.5	1.9	8.9	41.9	156.0	357.1	585.4
2004 4/.....	49.3	*	*	0.1	0.2	0.4	1.9	7.9	40.8	151.1	335.5	536.5
2003 4/.....	51.5	*	*	0.1	0.3	0.6	2.1	8.1	42.9	158.6	352.0	562.8
2002.....	50.5	*	*	0.2	0.3	0.6	2.0	8.1	42.4	157.0	345.4	554.5
2001.....	50.3	*	*	0.1	0.3	0.5	1.9	8.0	44.1	160.4	338.3	557.9
2000.....	49.7	*	*	0.2	0.3	0.5	1.8	7.9	43.2	159.6	339.1	544.8
1999.....	49.8	*	*	0.2	0.4	0.6	1.8	8.1	45.6	160.3	341.5	529.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic lower respiratory diseases (J40-J47)

Black, both sexes

2006 4/.....	19.5	*	*	1.0	1.1	1.9	4.0	12.0	37.4	100.8	225.9	331.8
2005 4/.....	21.1	*	1.2	0.8	1.1	1.7	3.8	13.9	39.4	114.4	241.3	366.7
2004 4/.....	19.2	*	0.8	1.0	1.2	1.8	3.9	12.6	38.1	102.1	222.8	333.9
2003 4/.....	20.2	*	1.0	0.9	1.3	1.9	4.3	12.5	40.5	113.2	235.9	353.0
2002.....	20.7	*	1.2	1.1	1.3	1.9	5.1	12.9	40.7	113.5	248.7	368.2
2001.....	20.4	*	0.9	0.8	1.3	2.4	4.8	12.5	40.7	118.9	234.1	375.8
2000.....	20.8	*	0.8	1.0	1.4	2.1	4.8	14.2	43.9	121.7	234.7	373.8
1999.....	21.9	*	*	1.0	1.3	2.6	4.6	13.8	45.8	128.5	262.6	376.2

Black, male

2006 4/.....	21.9	*	*	1.3	1.4	2.2	3.8	13.3	48.0	136.7	339.1	512.0
2005 4/.....	23.9	*	1.6	1.0	1.4	2.0	3.8	15.1	48.9	157.3	375.6	594.9
2004 4/.....	22.0	*	*	1.3	1.5	2.5	4.0	13.9	50.1	140.4	336.3	566.3
2003 4/.....	23.3	*	1.6	1.2	1.5	2.8	4.0	13.3	50.5	155.1	382.2	601.6
2002.....	24.1	*	1.6	1.2	1.9	2.2	5.7	14.4	52.3	158.0	392.2	645.4
2001.....	23.6	*	*	0.9	1.6	2.7	4.7	13.3	49.8	168.0	380.8	671.7
2000.....	24.3	*	*	1.2	1.7	2.8	4.8	15.0	54.6	176.9	370.3	693.1
1999.....	26.2	*	*	1.4	1.8	2.7	4.7	15.3	59.3	184.6	434.4	701.9

Black, female

2006 4/.....	17.4	*	*	0.8	0.9	1.6	4.1	10.8	28.9	75.2	161.6	258.1
2005 4/.....	18.4	*	*	*	0.8	1.5	3.8	12.8	31.7	83.6	165.1	274.6
2004 4/.....	16.6	*	*	0.6	1.0	1.2	3.9	11.5	28.5	74.9	158.6	241.8
2003 4/.....	17.3	*	*	0.6	1.1	1.1	4.6	11.9	32.4	83.3	153.2	256.4
2002.....	17.7	*	*	1.0	0.6	1.6	4.6	11.6	31.5	82.0	167.4	262.0
2001.....	17.5	*	*	0.7	1.0	2.1	4.9	11.7	33.3	84.3	151.7	266.1
2000.....	17.6	*	*	0.8	1.0	1.5	4.7	13.4	35.3	82.9	158.4	255.0
1999.....	18.0	*	*	0.6	0.9	2.6	4.6	12.6	34.9	88.9	166.4	254.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Bronchitis, chronic and unspecified (J40-J42)

All races 3/, both sexes

2006.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.0
2005.....	0.3	0.6	*	*	*	*	*	0.1	0.2	0.6	1.6	7.4
2004.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.5
2003.....	0.3	0.5	*	*	*	*	*	0.1	0.2	0.6	1.9	7.3
2002.....	0.3	0.6	*	*	*	*	*	0.1	0.2	0.7	2.2	8.1
2001.....	0.3	0.5	*	*	*	*	0.0	0.1	0.2	0.7	2.1	9.1
2000.....	0.4	*	*	*	*	*	0.0	0.1	0.3	0.9	2.7	11.8
1999.....	0.4	0.5	*	*	*	*	*	0.1	0.4	0.9	2.8	10.7

All races 3/, male

2006.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.5	5.9
2005.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.9	7.3
2004.....	0.2	*	*	*	*	*	*	0.1	0.3	0.6	1.6	7.0
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	2.2	7.0
2002.....	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.5	8.8
2001.....	0.3	*	*	*	*	*	*	0.1	0.3	0.9	2.3	9.2
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.0	3.6	12.9
1999.....	0.4	*	*	*	*	*	*	*	0.4	1.1	3.3	11.9

All races 3/, female

2006.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.5	6.0
2005.....	0.3	*	*	*	*	*	*	*	0.2	0.7	1.4	7.4
2004.....	0.3	*	*	*	*	*	*	*	0.2	0.4	1.5	6.3
2003.....	0.3	*	*	*	*	*	*	*	0.1	0.6	1.7	7.5
2002.....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.1	7.8
2001.....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	1.9	9.1
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.8	2.1	11.4
1999.....	0.5	*	*	*	*	*	*	*	0.4	0.8	2.5	10.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Bronchitis, chronic and unspecified (J40-J42)

White, both sexes

2006 4/.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.5	6.2
2005 4/.....	0.3	*	*	*	*	*	*	0.1	0.2	0.7	1.6	7.9
2004 4/.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	1.6	6.7
2003 4/.....	0.3	*	*	*	*	*	*	0.1	0.2	0.6	2.0	7.5
2002.....	0.4	*	*	*	*	*	*	0.1	0.2	0.8	2.3	8.6
2001.....	0.4	*	*	*	*	*	*	0.1	0.3	0.7	2.1	9.6
2000.....	0.5	*	*	*	*	*	*	0.1	0.3	0.9	2.8	12.7
1999.....	0.5	*	*	*	*	*	*	0.1	0.4	1.0	2.8	11.2

White, male

2006 4/.....	0.2	*	*	*	*	*	*	*	*	0.5	1.5	6.3
2005 4/.....	0.3	*	*	*	*	*	*	*	0.2	0.6	1.9	7.4
2004 4/.....	0.3	*	*	*	*	*	*	*	0.2	0.7	1.7	7.1
2003 4/.....	0.3	*	*	*	*	*	*	*	0.2	0.5	2.2	7.3
2002.....	0.3	*	*	*	*	*	*	0.1	0.2	0.9	2.6	9.3
2001.....	0.3	*	*	*	*	*	*	*	0.3	0.9	2.3	9.4
2000.....	0.4	*	*	*	*	*	*	*	0.3	1.0	3.7	13.8
1999.....	0.4	*	*	*	*	*	*	*	0.5	1.2	3.2	12.4

White, female

2006 4/.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.6	6.2
2005 4/.....	0.4	*	*	*	*	*	*	*	0.2	0.7	1.4	8.1
2004 4/.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.6	6.5
2003 4/.....	0.4	*	*	*	*	*	*	*	*	0.6	1.8	7.6
2002.....	0.4	*	*	*	*	*	*	*	0.2	0.7	2.2	8.3
2001.....	0.4	*	*	*	*	*	*	*	0.2	0.6	2.0	9.7
2000.....	0.5	*	*	*	*	*	*	*	0.3	0.8	2.2	12.2
1999.....	0.5	*	*	*	*	*	*	*	0.4	0.8	2.5	10.8



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Emphysema (J43)

All races 3/, both sexes

2006.....	4.2	*	*	*	*	*	0.2	1.0	4.8	17.3	36.8	45.3
2005.....	4.7	*	*	*	*	*	0.2	1.0	5.5	20.1	41.4	51.9
2004.....	4.6	*	*	*	*	*	0.2	1.0	5.4	20.5	41.0	49.9
2003.....	5.1	*	*	*	*	*	0.2	1.1	6.3	22.2	46.0	54.6
2002.....	5.4	*	*	*	*	*	0.2	1.1	6.4	24.0	48.0	59.3
2001.....	5.7	*	*	*	*	*	0.2	1.1	7.2	26.2	50.4	62.1
2000.....	5.9	*	*	*	*	*	0.2	1.3	7.4	27.8	52.7	63.3
1999.....	6.4	*	*	*	*	*	0.2	1.3	8.5	30.7	56.9	62.1

All races 3/, male

2006.....	4.3	*	*	*	*	*	0.2	1.3	5.3	19.6	45.4	63.7
2005.....	5.0	*	*	*	*	*	0.2	1.3	6.7	23.9	52.4	71.5
2004.....	4.9	*	*	*	*	*	0.2	1.2	6.2	24.3	52.2	74.4
2003.....	5.3	*	*	*	*	*	0.2	1.4	6.9	25.6	58.1	79.2
2002.....	5.7	*	*	*	*	*	0.3	1.4	7.2	28.1	63.5	90.0
2001.....	6.0	*	*	*	*	*	0.2	1.4	7.9	30.8	65.3	101.2
2000.....	6.3	*	*	*	*	*	0.2	1.5	8.1	33.0	69.1	104.2
1999.....	6.9	*	*	*	*	*	0.2	1.5	9.5	37.8	76.4	105.4

All races 3/, female

2006.....	4.1	*	*	*	*	*	0.2	0.9	4.4	15.4	30.9	36.7
2005.....	4.5	*	*	*	*	*	0.1	0.8	4.5	16.9	33.9	42.8
2004.....	4.4	*	*	*	*	*	0.1	0.8	4.7	17.4	33.5	38.9
2003.....	4.9	*	*	*	*	*	0.2	0.8	5.7	19.2	37.9	43.7
2002.....	5.0	*	*	*	*	*	0.1	0.9	5.6	20.6	37.6	46.0
2001.....	5.4	*	*	*	*	*	0.1	0.9	6.6	22.5	40.6	45.8
2000.....	5.6	*	*	*	*	*	0.2	1.0	6.8	23.6	41.9	46.7
1999.....	5.8	*	*	*	*	*	0.2	1.1	7.5	24.9	44.2	44.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Emphysema (J43)

White, both sexes

2006 4/.....	4.8	*	*	*	*	*	0.2	1.1	5.1	18.8	39.0	47.8
2005 4/.....	5.4	*	*	*	*	*	0.2	1.0	5.9	21.7	43.5	54.5
2004 4/.....	5.4	*	*	*	*	*	0.2	1.0	5.8	22.3	43.7	52.5
2003 4/.....	5.9	*	*	*	*	*	0.2	1.1	6.7	23.9	48.5	57.0
2002.....	6.2	*	*	*	*	*	0.2	1.2	6.9	25.7	50.7	62.3
2001.....	6.6	*	*	*	*	*	0.2	1.2	7.7	28.2	53.5	65.1
2000.....	6.8	*	*	*	*	*	0.2	1.2	7.9	29.7	55.6	65.8
1999.....	7.3	*	*	*	*	*	0.2	1.3	9.1	32.9	59.8	64.7

White, male

2006 4/.....	4.8	*	*	*	*	*	0.2	1.2	5.4	20.8	47.1	66.5
2005 4/.....	5.6	*	*	*	*	*	0.2	1.3	6.9	25.2	53.9	73.6
2004 4/.....	5.5	*	*	*	*	*	0.2	1.1	6.4	25.8	54.5	77.1
2003 4/.....	5.9	*	*	*	*	*	0.2	1.4	7.2	27.1	59.8	80.4
2002.....	6.4	*	*	*	*	*	0.3	1.3	7.5	29.3	65.8	93.0
2001.....	6.8	*	*	*	*	*	0.2	1.4	8.2	32.3	68.1	105.1
2000.....	7.0	*	*	*	*	*	0.2	1.4	8.4	34.3	72.1	105.5
1999.....	7.8	*	*	*	*	*	0.2	1.4	9.9	39.4	79.0	109.2

White, female

2006 4/.....	4.8	*	*	*	*	*	0.2	1.0	4.9	17.0	33.4	39.1
2005 4/.....	5.3	*	*	*	*	*	*	0.8	5.0	18.7	36.4	45.7
2004 4/.....	5.2	*	*	*	*	*	0.1	0.8	5.2	19.3	36.3	41.4
2003 4/.....	5.8	*	*	*	*	*	0.2	0.9	6.2	21.2	40.8	46.6
2002.....	5.9	*	*	*	*	*	*	1.0	6.2	22.7	40.5	48.9
2001.....	6.3	*	*	*	*	*	0.1	0.9	7.2	24.7	43.7	48.4
2000.....	6.5	*	*	*	*	*	0.2	1.0	7.4	25.9	44.7	49.7
1999.....	6.8	*	*	*	*	*	0.2	1.1	8.3	27.4	47.2	46.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Emphysema (J43)

Black, both sexes

2006 4/.....	1.8	*	*	*	*	*	0.4	1.0	4.2	10.0	22.1	23.9
2005 4/.....	2.1	*	*	*	*	*	*	1.4	4.5	12.5	26.2	27.3
2004 4/.....	1.9	*	*	*	*	*	*	1.3	4.5	10.7	22.1	27.6
2003 4/.....	2.1	*	*	*	*	*	*	1.2	5.1	12.9	27.3	30.9
2002.....	2.3	*	*	*	*	*	0.4	1.2	4.4	15.9	28.0	32.1
2001.....	2.2	*	*	*	*	*	*	1.3	5.6	15.3	23.6	35.3
2000.....	2.5	*	*	*	*	*	*	2.0	6.0	17.0	26.9	37.2
1999.....	2.6	*	*	*	*	*	0.4	1.8	6.1	17.1	31.3	37.6

Black, male

2006 4/.....	2.3	*	*	*	*	*	*	1.7	6.7	13.8	35.7	40.2
2005 4/.....	2.7	*	*	*	*	*	*	2.0	7.3	18.9	42.1	51.1
2004 4/.....	2.5	*	*	*	*	*	*	2.0	6.9	17.0	36.5	56.6
2003 4/.....	2.7	*	*	*	*	*	*	1.7	7.3	18.2	46.0	70.9
2002.....	3.0	*	*	*	*	*	*	1.8	6.5	24.8	48.8	64.3
2001.....	2.7	*	*	*	*	*	*	1.6	8.0	22.6	39.7	68.1
2000.....	3.2	*	*	*	*	*	*	2.9	8.1	27.4	41.5	87.6
1999.....	3.5	*	*	*	*	*	*	2.9	8.9	29.0	54.7	72.6

Black, female

2006 4/.....	1.3	*	*	*	*	*	*	*	2.2	7.3	14.3	17.2
2005 4/.....	1.5	*	*	*	*	*	*	0.8	2.2	8.0	17.2	17.6
2004 4/.....	1.3	*	*	*	*	*	*	*	2.6	6.2	14.0	16.1
2003 4/.....	1.6	*	*	*	*	*	*	*	3.4	9.2	16.7	15.4
2002.....	1.6	*	*	*	*	*	*	*	2.7	9.5	16.2	19.7
2001.....	1.7	*	*	*	*	*	*	1.1	3.6	10.2	14.5	23.2
2000.....	1.8	*	*	*	*	*	*	1.3	4.4	9.7	18.8	18.5
1999.....	1.7	*	*	*	*	*	*	1.0	3.9	8.7	18.2	24.5



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Asthma (J45-J46)

All races 3/, both sexes

2006.....	1.2	*	0.2	0.2	0.3	0.5	0.9	1.3	1.6	2.3	4.8	12.3
2005.....	1.3	*	0.2	0.2	0.3	0.5	0.8	1.4	1.7	2.5	5.4	14.5
2004.....	1.3	*	0.2	0.3	0.4	0.5	0.9	1.4	1.8	2.7	5.2	14.1
2003.....	1.4	*	0.2	0.3	0.4	0.6	0.9	1.5	2.0	2.9	5.8	14.2
2002.....	1.5	*	0.3	0.3	0.4	0.6	1.1	1.5	2.0	3.2	6.4	14.7
2001.....	1.5	*	0.2	0.2	0.4	0.6	1.0	1.5	2.2	3.5	6.4	16.1
2000.....	1.6	*	0.2	0.3	0.4	0.6	1.0	1.6	2.2	4.0	6.9	16.7
1999.....	1.7	*	0.2	0.3	0.5	0.7	1.0	1.7	2.5	4.2	7.6	17.3

All races 3/, male

2006.....	0.9	*	*	0.3	0.4	0.5	0.7	1.0	1.1	1.7	3.5	9.1
2005.....	0.9	*	0.3	0.3	0.4	0.5	0.7	1.0	1.2	1.8	4.1	8.7
2004.....	1.0	*	*	0.3	0.5	0.6	0.7	1.0	1.3	2.1	3.5	11.1
2003.....	1.0	*	0.4	0.3	0.4	0.6	0.7	1.2	1.6	2.1	4.5	10.7
2002.....	1.1	*	0.4	0.3	0.5	0.6	0.9	1.2	1.6	2.4	5.0	10.9
2001.....	1.1	*	0.3	0.3	0.4	0.6	0.9	1.1	1.5	2.7	4.5	12.1
2000.....	1.2	*	0.3	0.4	0.5	0.7	0.9	1.3	1.7	3.0	5.0	12.9
1999.....	1.2	*	*	0.4	0.5	0.6	0.8	1.1	1.8	3.2	5.7	12.7

All races 3/, female

2006.....	1.5	*	*	0.1	0.3	0.4	1.0	1.6	2.0	2.9	5.7	13.9
2005.....	1.7	*	*	0.2	0.2	0.5	1.0	1.8	2.2	3.2	6.3	17.2
2004.....	1.6	*	*	0.2	0.3	0.4	1.0	1.6	2.2	3.2	6.3	15.4
2003.....	1.8	*	*	0.2	0.3	0.5	1.1	1.9	2.4	3.6	6.7	15.8
2002.....	1.8	*	*	0.3	0.3	0.5	1.2	1.8	2.4	3.9	7.3	16.3
2001.....	1.9	*	*	0.2	0.3	0.6	1.1	1.9	2.8	4.1	7.6	17.8
2000.....	2.0	*	*	0.3	0.3	0.5	1.2	2.0	2.7	4.8	8.1	18.2
1999.....	2.1	*	*	0.2	0.4	0.7	1.2	2.2	3.0	5.0	8.8	19.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Asthma (J45-J46)

White, both sexes

2006 4/.....	1.0	*	*	0.1	0.2	0.3	0.6	1.0	1.2	2.0	4.5	12.1
2005 4/.....	1.1	*	*	0.1	0.2	0.4	0.6	1.0	1.3	2.0	4.9	14.5
2004 4/.....	1.1	*	*	0.1	0.2	0.3	0.6	0.9	1.4	2.4	4.7	13.7
2003 4/.....	1.2	*	*	0.2	0.2	0.4	0.7	1.1	1.6	2.4	5.4	14.3
2002.....	1.3	*	*	0.1	0.3	0.5	0.7	1.1	1.5	2.7	6.1	14.7
2001.....	1.3	*	*	0.1	0.2	0.4	0.7	1.1	1.6	3.0	6.0	15.7
2000.....	1.4	*	*	0.2	0.3	0.4	0.7	1.1	1.7	3.4	6.5	15.9
1999.....	1.5	*	*	0.2	0.3	0.4	0.7	1.2	1.9	3.5	7.2	16.6

White, male

2006 4/.....	0.7	*	*	0.2	0.2	0.3	0.5	0.6	0.7	1.3	3.2	9.1
2005 4/.....	0.7	*	*	0.1	0.2	0.3	0.5	0.8	0.9	1.3	3.5	8.6
2004 4/.....	0.7	*	*	0.1	0.3	0.3	0.5	0.7	0.9	1.8	2.8	10.3
2003 4/.....	0.8	*	*	0.2	0.3	0.4	0.5	0.8	1.2	1.7	4.0	10.6
2002.....	0.9	*	*	0.2	0.3	0.5	0.5	0.8	1.2	1.9	4.6	10.7
2001.....	0.8	*	*	0.2	0.2	0.3	0.6	0.8	1.1	2.2	4.0	11.3
2000.....	0.9	*	*	0.2	0.3	0.4	0.6	1.0	1.2	2.5	4.7	11.4
1999.....	0.9	*	*	0.2	0.4	0.4	0.5	0.8	1.4	2.5	5.3	10.8

White, female

2006 4/.....	1.4	*	*	*	0.1	0.3	0.8	1.3	1.6	2.5	5.4	13.6
2005 4/.....	1.5	*	*	*	0.1	0.4	0.8	1.3	1.8	2.6	5.9	17.2
2004 4/.....	1.5	*	*	0.1	0.2	0.3	0.8	1.2	1.8	3.0	6.0	15.2
2003 4/.....	1.6	*	*	0.1	0.2	0.5	0.8	1.4	2.0	3.1	6.3	15.9
2002.....	1.7	*	*	*	0.2	0.5	0.9	1.3	1.9	3.3	7.2	16.4
2001.....	1.7	*	*	*	0.2	0.4	0.8	1.5	2.1	3.7	7.3	17.6
2000.....	1.8	*	*	0.2	0.2	0.4	0.8	1.3	2.1	4.1	7.7	17.7
1999.....	2.0	*	*	0.1	0.3	0.4	0.9	1.7	2.4	4.4	8.5	18.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Asthma (J45-J46)

Black, both sexes

2006 4/.....	2.4	*	*	1.0	1.1	1.6	2.4	3.9	5.1	5.5	6.0	12.5
2005 4/.....	2.6	*	1.1	0.8	1.1	1.5	2.3	4.2	5.4	7.2	9.8	11.8
2004 4/.....	2.6	*	*	0.8	1.1	1.6	2.5	4.5	5.6	5.7	9.1	15.5
2003 4/.....	2.7	*	0.9	0.9	1.2	1.6	2.7	4.8	5.1	7.0	9.1	12.0
2002.....	2.9	*	0.9	1.0	1.2	1.6	3.4	4.8	6.4	7.8	8.0	11.9
2001.....	3.0	*	0.8	0.8	1.1	2.1	3.3	4.6	6.8	7.7	10.6	16.3
2000.....	3.2	*	*	1.0	1.3	1.9	3.1	5.4	6.9	9.2	10.4	22.2
1999.....	3.2	*	*	1.0	1.2	2.4	2.9	5.2	6.8	9.6	10.4	20.4

Black, male

2006 4/.....	2.2	*	*	1.3	1.4	1.9	2.3	3.5	4.7	4.7	*	*
2005 4/.....	2.2	*	1.6	1.0	1.4	1.7	2.1	2.9	4.0	5.4	10.7	*
2004 4/.....	2.5	*	*	1.1	1.4	2.3	2.3	3.9	5.2	5.5	8.6	*
2003 4/.....	2.4	*	1.5	1.2	1.4	2.4	2.3	3.9	4.2	5.3	8.7	*
2002.....	2.8	*	*	1.1	1.8	2.0	3.6	3.9	5.7	7.3	6.9	*
2001.....	2.6	*	*	0.9	1.4	2.4	3.0	4.1	4.6	6.8	8.8	*
2000.....	2.8	*	*	1.2	1.5	2.5	2.9	3.8	6.1	7.9	6.4	25.4
1999.....	2.8	*	*	1.4	1.6	2.6	2.8	3.8	5.4	9.1	8.3	*

Black, female

2006 4/.....	2.6	*	*	0.6	0.9	1.3	2.5	4.3	5.4	6.1	6.9	15.0
2005 4/.....	3.0	*	*	*	0.8	1.4	2.5	5.2	6.5	8.5	9.3	13.5
2004 4/.....	2.7	*	*	*	0.9	1.1	2.7	5.0	6.0	5.7	9.4	16.9
2003 4/.....	2.9	*	*	*	1.0	0.9	3.1	5.6	5.8	8.2	9.2	13.8
2002.....	3.0	*	*	1.0	*	1.2	3.3	5.6	7.0	8.2	8.6	12.3
2001.....	3.3	*	*	0.7	0.9	1.9	3.5	5.0	8.5	8.3	11.6	16.8
2000.....	3.5	*	*	0.8	1.0	1.4	3.3	6.8	7.5	10.2	12.6	21.0
1999.....	3.5	*	*	0.6	0.9	2.1	3.1	6.4	7.9	10.0	11.5	21.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic lower respiratory diseases (J44,J47)

All races 3/, both sexes

2006.....	36.0	*	*	*	*	0.1	0.9	6.6	32.6	129.1	320.2	525.4
2005.....	37.8	*	*	*	*	0.1	1.0	6.9	34.5	137.2	337.2	563.5
2004.....	35.3	*	*	*	*	0.1	0.9	6.0	33.0	130.0	319.0	531.1
2003.....	36.6	*	*	*	0.1	0.1	1.0	5.9	34.8	137.5	329.2	558.9
2002.....	36.1	*	*	*	*	0.1	0.9	5.9	33.8	135.1	330.1	555.5
2001.....	35.7	*	*	*	0.1	0.1	0.9	5.7	34.5	137.5	321.0	557.4
2000.....	35.4	*	*	*	*	0.1	0.8	5.6	34.3	136.7	323.8	556.7
1999.....	36.0	*	*	*	*	0.1	0.7	5.5	36.2	141.4	330.6	555.8

All races 3/, male

2006.....	34.8	*	*	*	*	0.1	0.8	6.9	35.3	141.5	386.8	682.2
2005.....	36.6	*	*	*	*	0.1	1.0	7.3	37.0	154.4	405.2	732.4
2004.....	34.5	*	*	*	*	0.1	1.0	6.5	35.2	145.0	388.4	718.6
2003.....	35.9	*	*	*	*	0.1	1.0	6.4	37.8	155.4	400.1	769.0
2002.....	35.7	*	*	*	*	0.1	1.0	6.4	36.2	153.4	409.8	785.1
2001.....	35.3	*	*	*	*	0.1	0.9	6.2	37.2	157.0	403.0	794.5
2000.....	35.6	*	*	*	*	*	0.8	6.1	37.7	158.3	410.9	838.0
1999.....	37.1	*	*	*	*	*	0.8	5.9	40.5	168.6	427.9	866.7

All races 3/, female

2006.....	37.1	*	*	*	*	0.1	0.9	6.4	30.1	118.7	274.8	452.1
2005.....	39.0	*	*	*	*	*	1.0	6.4	32.2	122.6	291.0	485.9
2004.....	36.2	*	*	*	*	*	0.9	5.5	30.8	117.3	272.2	446.8
2003.....	37.4	*	*	*	*	*	1.0	5.5	32.1	122.6	281.9	466.0
2002.....	36.5	*	*	*	*	0.1	0.9	5.4	31.5	119.8	277.2	455.8
2001.....	36.0	*	*	*	*	0.1	1.0	5.2	32.0	121.3	267.1	458.2
2000.....	35.2	*	*	*	*	*	0.8	5.2	31.2	119.0	267.1	442.2
1999.....	35.0	*	*	*	*	0.2	0.7	5.0	32.2	119.1	267.4	430.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic lower respiratory diseases (J44,J47)

White, both sexes

2006 4/.....	41.4	*	*	*	*	0.1	0.8	6.9	34.6	139.1	338.7	552.1
2005 4/.....	43.5	*	*	*	*	0.1	1.0	7.0	36.6	147.0	356.9	590.7
2004 4/.....	40.6	*	*	*	*	0.1	1.0	6.2	34.9	139.4	336.9	557.2
2003 4/.....	42.1	*	*	*	*	0.1	1.0	6.2	36.7	146.9	347.2	585.8
2002.....	41.3	*	*	*	*	0.1	1.0	6.1	35.6	144.0	346.4	580.7
2001.....	40.8	*	*	*	*	0.1	0.9	5.8	36.5	145.9	337.0	581.3
2000.....	40.4	*	*	*	*	*	0.8	5.7	36.0	144.7	339.5	581.2
1999.....	41.0	*	*	*	*	0.1	0.7	5.6	37.8	148.8	344.5	580.3

White, male

2006 4/.....	39.3	*	*	*	*	*	0.8	7.1	36.6	148.9	401.6	706.7
2005 4/.....	41.3	*	*	*	*	*	1.0	7.3	38.3	162.1	420.1	756.5
2004 4/.....	39.0	*	*	*	*	0.1	1.0	6.6	36.3	152.3	403.4	743.2
2003 4/.....	40.4	*	*	*	*	*	0.9	6.6	39.0	162.2	412.5	796.0
2002.....	40.2	*	*	*	*	*	1.0	6.5	37.1	160.2	422.1	810.4
2001.....	39.7	*	*	*	*	0.1	0.9	6.3	38.5	162.9	415.0	817.8
2000.....	39.9	*	*	*	*	*	0.8	6.0	38.7	163.6	423.0	866.7
1999.....	41.5	*	*	*	*	*	0.7	5.9	41.4	174.2	438.0	897.0

White, female

2006 4/.....	43.4	*	*	*	*	*	0.9	6.8	32.7	130.5	295.1	479.5
2005 4/.....	45.6	*	*	*	*	*	1.0	6.7	35.0	134.0	313.3	514.4
2004 4/.....	42.2	*	*	*	*	*	0.9	5.8	33.6	128.3	291.6	473.4
2003 4/.....	43.7	*	*	*	*	*	1.0	5.8	34.5	133.8	303.0	492.7
2002.....	42.5	*	*	*	*	*	0.9	5.7	34.1	130.3	295.6	480.9
2001.....	41.8	*	*	*	*	*	1.0	5.4	34.6	131.5	285.3	482.3
2000.....	40.9	*	*	*	*	*	0.8	5.5	33.4	128.9	284.4	465.2
1999.....	40.4	*	*	*	*	0.1	0.7	5.2	34.5	127.7	283.2	453.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic lower respiratory diseases (J44,J47)

Black, both sexes

2006 4/.....	15.2	*	*	*	*	*	1.1	7.0	27.9	84.8	196.3	291.9
2005 4/.....	16.2	*	*	*	*	*	1.2	8.3	29.2	94.2	204.3	326.1
2004 4/.....	14.5	*	*	*	*	*	1.1	6.7	27.7	85.5	190.7	285.9
2003 4/.....	15.2	*	*	*	*	*	1.2	6.3	29.9	92.9	198.3	304.6
2002.....	15.4	*	*	*	*	*	1.2	6.7	29.7	89.5	211.3	320.1
2001.....	15.1	*	*	*	*	*	1.3	6.4	28.1	95.0	198.7	320.9
2000.....	14.9	*	*	*	*	*	1.3	6.6	30.9	94.5	195.2	310.6
1999.....	15.9	*	*	*	*	*	1.2	6.4	32.5	100.8	218.0	313.7

Black, male

2006 4/.....	17.2	*	*	*	*	*	1.0	8.0	36.2	117.7	296.8	463.8
2005 4/.....	18.8	*	*	*	*	*	1.2	10.1	37.4	132.2	321.0	532.6
2004 4/.....	16.8	*	*	*	*	*	1.2	7.7	37.8	117.4	290.6	492.7
2003 4/.....	18.0	*	*	*	*	*	1.3	7.4	38.7	131.3	325.5	520.3
2002.....	18.2	*	*	*	*	*	1.4	8.3	39.9	125.3	335.1	565.0
2001.....	18.1	*	*	*	*	*	1.4	7.4	37.0	137.4	330.5	583.2
2000.....	18.2	*	*	*	*	*	1.3	8.3	40.1	140.8	320.4	576.6
1999.....	19.6	*	*	*	*	*	1.5	8.2	44.6	145.4	366.5	607.0

Black, female

2006 4/.....	13.3	*	*	*	*	*	1.2	6.1	21.1	61.3	139.1	221.5
2005 4/.....	13.9	*	*	*	*	*	1.1	6.7	22.6	66.9	138.1	242.7
2004 4/.....	12.4	*	*	*	*	*	1.0	5.8	19.5	62.7	134.2	204.0
2003 4/.....	12.6	*	*	*	*	*	1.2	5.3	22.9	65.5	126.5	220.8
2002.....	12.9	*	*	*	*	*	1.1	5.3	21.5	64.0	141.3	226.3
2001.....	12.3	*	*	*	*	*	1.2	5.6	21.0	65.1	124.8	223.5
2000.....	12.0	*	*	*	*	*	1.3	5.1	23.4	62.0	124.8	211.6
1999.....	12.5	*	*	*	*	*	1.0	4.9	22.8	69.4	134.9	204.1









Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonitis due to solids and liquids (J69)

All races 3/, both sexes

2006.....	5.6	*	*	*	0.1	0.2	0.4	1.0	2.7	9.3	40.5	155.8
2005.....	5.8	*	*	*	0.1	0.2	0.4	1.1	2.7	9.3	42.6	163.9
2004.....	5.7	*	*	*	0.1	0.2	0.4	0.9	2.6	9.6	42.9	164.9
2003.....	6.0	*	*	*	0.1	0.2	0.4	1.0	2.8	9.6	45.0	176.2
2002.....	6.1	*	*	*	0.1	0.2	0.4	0.9	2.5	9.8	46.3	186.0
2001.....	6.1	*	*	*	0.1	0.2	0.4	1.0	2.6	10.0	45.8	189.4
2000.....	5.9	*	*	*	0.1	0.2	0.4	1.0	2.5	10.3	44.5	187.6
1999.....	5.5	*	*	*	0.1	0.2	0.4	0.8	2.5	9.5	41.1	175.6

All races 3/, male

2006.....	6.0	*	*	*	*	0.3	0.5	1.3	3.5	12.6	57.2	224.8
2005.....	6.0	*	*	*	0.1	0.2	0.4	1.3	3.5	12.5	57.9	229.3
2004.....	5.9	*	*	*	0.1	0.2	0.4	1.2	3.2	12.9	58.6	237.2
2003.....	6.1	*	*	*	0.1	0.2	0.4	1.2	3.5	12.8	61.7	246.2
2002.....	6.3	*	*	*	0.1	0.2	0.5	1.1	3.1	13.0	63.8	268.5
2001.....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.2	13.3	63.1	270.3
2000.....	6.1	*	*	*	0.1	0.2	0.5	1.2	3.0	14.4	62.3	276.0
1999.....	5.7	*	*	*	0.1	0.2	0.4	0.9	3.2	13.1	59.7	258.4

All races 3/, female

2006.....	5.2	*	*	*	*	0.1	0.3	0.8	1.9	6.4	29.1	123.5
2005.....	5.7	*	*	*	0.1	0.1	0.4	0.9	2.1	6.7	32.2	133.9
2004.....	5.5	*	*	*	*	0.1	0.3	0.7	1.9	6.8	32.3	134.4
2003.....	5.8	*	*	*	*	0.1	0.3	0.7	2.1	6.9	33.8	145.3
2002.....	5.9	*	*	*	*	0.2	0.3	0.6	2.0	7.1	34.7	150.2
2001.....	6.0	*	*	*	*	0.2	0.3	0.7	2.1	7.2	34.5	155.6
2000.....	5.8	*	*	*	*	0.1	0.3	0.7	1.9	7.0	33.0	151.6
1999.....	5.3	*	*	*	*	0.2	0.3	0.6	1.8	6.6	29.1	142.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonitis due to solids and liquids (J69)

White, both sexes

2006 4/.....	6.3	*	*	*	*	0.2	0.4	1.0	2.7	9.0	41.1	160.0
2005 4/.....	6.4	*	*	*	0.1	0.2	0.4	1.1	2.6	9.1	42.5	166.5
2004 4/.....	6.3	*	*	*	0.1	0.2	0.3	0.9	2.4	9.4	43.2	167.8
2003 4/.....	6.5	*	*	*	0.1	0.2	0.3	1.0	2.6	9.3	44.7	178.7
2002.....	6.7	*	*	*	0.1	0.2	0.4	0.8	2.4	9.5	46.5	189.5
2001.....	6.6	*	*	*	0.1	0.2	0.4	0.9	2.5	9.6	45.8	190.8
2000.....	6.5	*	*	*	0.1	0.1	0.4	0.9	2.3	9.8	44.4	191.0
1999.....	5.9	*	*	*	0.1	0.2	0.3	0.7	2.3	9.0	40.7	178.0

White, male

2006 4/.....	6.7	*	*	*	*	0.2	0.4	1.3	3.5	12.2	57.5	231.8
2005 4/.....	6.5	*	*	*	*	0.2	0.4	1.3	3.3	12.0	57.3	234.2
2004 4/.....	6.5	*	*	*	*	0.2	0.4	1.2	2.9	12.4	58.8	241.7
2003 4/.....	6.6	*	*	*	0.1	0.2	0.4	1.2	3.2	12.4	61.1	250.8
2002.....	6.9	*	*	*	0.1	0.2	0.5	1.1	3.0	12.5	64.0	274.6
2001.....	6.7	*	*	*	*	0.2	0.5	1.2	3.1	12.9	63.3	275.4
2000.....	6.6	*	*	*	*	0.1	0.5	1.1	2.9	13.4	62.1	284.0
1999.....	6.1	*	*	*	*	0.2	0.4	0.8	2.9	12.2	58.7	263.6

White, female

2006 4/.....	5.9	*	*	*	*	*	0.3	0.7	2.0	6.3	29.7	126.3
2005 4/.....	6.3	*	*	*	*	0.1	0.4	0.9	2.0	6.6	32.3	135.3
2004 4/.....	6.1	*	*	*	*	0.1	0.3	0.6	1.9	6.8	32.6	134.5
2003 4/.....	6.4	*	*	*	*	0.1	0.3	0.7	2.0	6.7	33.6	146.7
2002.....	6.6	*	*	*	*	0.2	0.3	0.6	1.9	6.9	34.8	152.5
2001.....	6.6	*	*	*	*	0.2	0.3	0.7	2.1	6.9	34.2	155.3
2000.....	6.3	*	*	*	*	0.2	0.3	0.7	1.8	6.8	32.8	153.2
1999.....	5.8	*	*	*	*	0.2	0.3	0.6	1.7	6.4	29.0	143.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pneumonitis due to solids and liquids (J69)

Black, both sexes

2006 4/.....	3.6	*	*	*	*	0.4	0.6	1.4	3.7	13.9	42.1	128.4
2005 4/.....	4.1	*	*	*	*	*	0.5	1.7	4.4	13.3	52.4	153.1
2004 4/.....	3.9	*	*	*	*	*	0.6	1.3	4.8	13.9	46.9	144.7
2003 4/.....	4.2	*	*	*	*	*	0.7	1.3	5.1	13.7	54.4	163.4
2002.....	4.1	*	*	*	*	*	0.7	1.4	3.7	14.8	52.2	165.1
2001.....	4.3	*	*	*	*	*	0.6	1.6	4.0	15.1	53.2	185.5
2000.....	4.2	*	*	*	*	*	0.7	1.9	4.2	17.5	51.5	159.8
1999.....	4.2	*	*	*	*	*	0.7	1.5	4.4	16.2	52.2	167.1

Black, male

2006 4/.....	4.0	*	*	*	*	*	0.8	1.7	5.2	20.6	66.8	173.4
2005 4/.....	4.4	*	*	*	*	*	*	2.4	6.1	19.8	79.2	201.7
2004 4/.....	4.2	*	*	*	*	*	0.7	1.7	7.1	21.2	67.7	202.7
2003 4/.....	4.3	*	*	*	*	*	*	1.6	7.1	19.6	77.8	212.7
2002.....	4.2	*	*	*	*	*	0.8	2.0	4.8	21.7	73.4	233.7
2001.....	4.2	*	*	*	*	*	0.7	1.9	5.3	20.4	75.3	243.9
2000.....	4.4	*	*	*	*	*	0.8	2.7	5.0	27.8	74.5	209.9
1999.....	4.5	*	*	*	*	*	0.9	1.8	6.4	25.0	80.9	228.5

Black, female

2006 4/.....	3.3	*	*	*	*	*	*	1.2	2.5	9.0	28.0	110.0
2005 4/.....	3.9	*	*	*	*	*	*	1.1	3.1	8.6	37.2	133.5
2004 4/.....	3.6	*	*	*	*	*	*	1.0	3.0	8.7	35.2	121.7
2003 4/.....	4.1	*	*	*	*	*	0.8	1.0	3.6	9.6	41.1	144.2
2002.....	3.9	*	*	*	*	*	*	0.9	2.9	9.9	40.1	138.8
2001.....	4.4	*	*	*	*	*	*	1.4	2.9	11.3	40.8	163.8
2000.....	3.9	*	*	*	*	*	*	1.2	3.5	10.3	38.5	141.2
1999.....	3.8	*	*	*	*	*	*	1.2	2.8	10.0	36.1	144.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes

2006.....	9.2	8.1	0.6	0.2	0.2	0.5	1.2	3.1	8.9	28.2	73.4	136.6
2005.....	9.1	7.4	0.6	0.2	0.3	0.5	1.1	3.2	8.8	28.8	72.5	136.5
2004.....	9.3	6.5	0.7	0.2	0.3	0.4	1.2	3.2	9.5	29.9	74.2	139.4
2003.....	8.8	7.0	0.5	0.1	0.3	0.5	1.1	3.3	8.8	28.1	68.5	138.1
2002.....	8.7	6.9	0.5	0.2	0.3	0.5	1.2	3.2	9.2	28.7	68.0	133.5
2001.....	9.2	7.3	0.6	0.2	0.4	0.5	1.3	3.5	9.6	30.1	73.2	142.7
2000.....	9.1	7.2	0.6	0.2	0.3	0.6	1.4	3.5	9.9	29.6	72.0	141.7
1999.....	9.0	6.6	0.6	0.2	0.3	0.6	1.3	3.3	9.5	29.4	70.7	147.6

All races 3/, male

2006.....	9.3	9.2	0.7	0.2	0.3	0.5	1.2	3.3	10.3	35.0	93.1	163.7
2005.....	9.0	8.9	0.7	0.2	0.3	0.4	1.0	3.5	10.2	34.9	90.7	157.7
2004.....	9.3	6.6	0.7	0.2	0.3	0.4	1.2	3.5	10.8	36.9	93.9	168.8
2003.....	8.7	8.0	0.6	0.2	0.4	0.6	1.1	3.5	10.2	34.1	85.3	165.0
2002.....	8.5	7.9	0.7	0.2	0.3	0.4	1.3	3.3	10.5	35.3	83.9	156.6
2001.....	8.9	9.0	0.6	0.2	0.4	0.5	1.4	3.8	10.9	36.5	89.3	169.1
2000.....	9.0	7.9	0.6	0.2	0.4	0.6	1.5	3.8	11.4	35.7	90.5	179.7
1999.....	8.9	7.5	0.7	0.2	0.3	0.6	1.4	3.5	10.6	36.2	89.9	182.3

All races 3/, female

2006.....	9.2	6.9	0.5	0.2	0.2	0.5	1.2	3.0	7.5	22.6	59.9	124.0
2005.....	9.2	5.8	0.5	0.1	0.2	0.5	1.1	2.9	7.6	23.7	60.2	126.7
2004.....	9.3	6.4	0.6	0.1	0.3	0.4	1.2	3.0	8.4	24.1	61.0	126.1
2003.....	8.9	5.9	0.5	0.1	0.3	0.5	1.1	3.0	7.4	23.1	57.2	126.2
2002.....	8.8	5.8	0.4	0.2	0.3	0.5	1.1	3.1	8.0	23.1	57.5	123.4
2001.....	9.4	5.5	0.6	0.2	0.3	0.6	1.2	3.1	8.4	24.8	62.6	131.6
2000.....	9.2	6.4	0.6	0.2	0.3	0.6	1.2	3.2	8.6	24.7	59.9	126.2
1999.....	9.1	5.8	0.5	0.2	0.2	0.7	1.1	3.1	8.5	23.7	58.3	133.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

White, both sexes

2006 4/.....	10.1	5.7	0.5	0.2	0.2	0.5	1.1	2.8	8.6	29.1	76.1	141.3
2005 4/.....	9.9	6.0	0.5	0.2	0.2	0.4	1.0	2.8	8.5	29.3	74.7	140.2
2004 4/.....	10.2	4.9	0.5	0.1	0.3	0.4	1.1	2.9	9.3	30.7	76.8	144.1
2003 4/.....	9.6	5.4	0.4	0.1	0.3	0.5	0.9	2.9	8.6	29.0	70.8	143.2
2002.....	9.5	4.8	0.5	0.2	0.3	0.4	1.0	2.9	8.9	29.3	69.9	137.9
2001.....	10.0	5.0	0.5	0.2	0.3	0.4	1.1	3.0	9.3	30.6	75.4	146.0
2000.....	9.9	5.6	0.5	0.2	0.3	0.6	1.2	3.2	9.7	30.0	73.7	145.9
1999.....	9.7	4.9	0.5	0.2	0.3	0.6	1.1	3.0	9.2	29.6	72.2	150.7

White, male

2006 4/.....	10.2	6.1	0.6	0.2	0.3	0.4	1.1	3.0	10.2	35.9	96.4	168.5
2005 4/.....	9.7	7.0	0.6	0.2	0.3	0.4	1.0	3.2	9.7	35.7	92.6	160.0
2004 4/.....	10.1	4.9	0.6	0.1	0.3	0.4	1.0	3.2	10.6	37.8	96.7	174.3
2003 4/.....	9.4	5.8	0.4	0.2	0.3	0.5	1.0	3.3	9.9	35.1	87.6	170.5
2002.....	9.2	5.4	0.6	0.2	0.3	0.4	1.1	3.0	10.1	36.1	86.1	162.0
2001.....	9.7	5.9	0.5	0.2	0.3	0.4	1.2	3.4	10.6	37.4	92.0	173.5
2000.....	9.7	6.6	0.5	0.2	0.3	0.6	1.3	3.4	11.1	36.1	92.5	184.4
1999.....	9.6	5.6	0.6	0.2	0.4	0.6	1.2	3.2	10.3	36.8	92.0	185.7

White, female

2006 4/.....	10.1	5.3	0.4	0.2	0.2	0.6	1.1	2.6	7.1	23.2	62.0	128.6
2005 4/.....	10.1	4.9	0.5	0.2	0.2	0.4	1.0	2.5	7.4	23.8	62.4	131.1
2004 4/.....	10.2	4.9	0.4	*	0.3	0.4	1.1	2.6	8.1	24.7	63.2	130.5
2003 4/.....	9.8	5.0	0.4	*	0.3	0.4	0.9	2.6	7.2	23.8	59.4	131.0
2002.....	9.7	4.2	*	0.2	0.2	0.4	1.0	2.8	7.7	23.4	59.1	127.4
2001.....	10.2	4.0	0.6	0.2	0.3	0.5	0.9	2.6	8.2	24.9	64.3	134.4
2000.....	10.0	4.6	0.6	0.2	0.3	0.6	1.1	2.9	8.4	25.0	61.4	130.3
1999.....	9.8	4.3	0.5	0.2	0.2	0.5	0.9	2.8	8.2	23.6	59.3	136.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

Black, both sexes

2006 4/.....	6.2	19.4	0.8	*	0.4	0.8	2.2	5.9	13.0	25.5	51.5	95.2
2005 4/.....	6.3	15.4	0.9	*	0.4	0.9	2.1	6.2	12.1	29.0	56.0	102.6
2004 4/.....	6.2	14.8	1.2	*	0.4	0.9	2.6	6.2	13.4	27.5	53.2	92.8
2003 4/.....	5.9	15.0	1.1	*	0.6	1.0	2.4	6.4	12.2	24.2	50.8	89.6
2002.....	6.0	16.6	1.1	0.4	0.7	0.9	2.3	6.1	13.6	26.5	51.5	90.3
2001.....	6.7	18.0	1.0	*	0.6	1.3	2.9	7.3	13.6	29.5	56.7	109.6
2000.....	6.5	16.7	0.8	*	0.7	1.1	2.7	6.8	13.4	30.5	58.0	98.2
1999.....	6.5	15.4	1.0	*	0.4	1.3	2.9	6.4	13.4	30.8	57.9	111.6

Black, male

2006 4/.....	6.2	23.6	*	*	0.6	0.8	2.3	5.6	14.0	32.7	66.3	116.2
2005 4/.....	6.5	20.1	*	*	*	0.8	2.0	6.7	15.3	33.7	74.8	132.0
2004 4/.....	6.3	15.7	*	*	*	1.0	3.0	6.7	15.2	34.5	69.0	115.2
2003 4/.....	6.0	18.4	*	*	0.8	1.1	2.2	6.3	15.1	30.8	68.5	113.6
2002.....	6.0	19.5	*	*	*	0.9	2.7	6.5	16.1	31.7	65.7	100.8
2001.....	6.4	23.1	*	*	*	1.3	2.6	7.6	15.3	32.9	69.4	129.3
2000.....	6.5	16.3	*	*	0.8	1.2	3.1	7.8	15.2	36.0	73.0	132.6
1999.....	6.4	17.6	*	*	*	1.0	2.9	7.1	15.5	35.1	72.6	145.3

Black, female

2006 4/.....	6.2	15.0	*	*	*	0.8	2.1	6.1	12.2	20.3	43.1	86.6
2005 4/.....	6.2	10.5	*	*	*	1.0	2.2	5.7	9.6	25.6	45.3	90.8
2004 4/.....	6.1	13.9	*	*	*	0.8	2.2	5.8	11.9	22.4	44.2	83.9
2003 4/.....	5.7	11.4	*	*	*	1.0	2.7	6.5	9.8	19.4	40.8	80.2
2002.....	6.0	13.6	*	*	0.7	0.8	2.0	5.8	11.5	22.9	43.5	86.3
2001.....	7.0	12.6	*	*	*	1.2	3.1	7.0	12.2	27.1	49.6	102.3
2000.....	6.4	17.2	*	*	*	1.0	2.2	5.9	11.9	26.6	49.6	85.4
1999.....	6.7	13.1	*	*	*	1.6	2.8	5.8	11.6	27.8	49.7	99.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Peptic ulcer (K25-K28)

All races 3/, both sexes

2006.....	1.1	*	*	*	*	*	0.2	0.7	1.2	2.7	7.2	20.7
2005.....	1.2	*	*	*	*	0.1	0.2	0.7	1.2	2.9	8.0	21.4
2004.....	1.3	*	*	*	*	*	0.2	0.6	1.3	3.2	8.9	24.4
2003.....	1.3	*	*	*	*	0.1	0.2	0.7	1.4	3.2	9.9	26.4
2002.....	1.4	*	*	*	*	0.1	0.3	0.7	1.5	3.6	10.1	28.3
2001.....	1.6	*	*	*	*	0.1	0.3	0.7	1.7	3.9	11.4	33.4
2000.....	1.6	*	*	*	*	*	0.3	0.8	1.8	4.1	11.4	35.6
1999.....	1.6	*	*	*	*	0.1	0.3	0.7	1.8	4.5	12.3	34.6

All races 3/, male

2006.....	1.1	*	*	*	*	*	0.2	0.9	1.6	3.2	8.2	22.0
2005.....	1.2	*	*	*	*	*	0.3	1.0	1.7	3.6	9.7	22.4
2004.....	1.2	*	*	*	*	*	0.3	0.9	1.6	3.6	10.2	24.6
2003.....	1.3	*	*	*	*	*	0.3	1.0	1.8	4.1	11.8	28.9
2002.....	1.4	*	*	*	*	*	0.3	0.9	1.9	4.6	12.2	31.6
2001.....	1.5	*	*	*	*	0.1	0.4	1.0	2.2	4.8	13.0	34.9
2000.....	1.6	*	*	*	*	*	0.4	1.2	2.4	5.3	13.7	40.7
1999.....	1.6	*	*	*	*	*	0.3	0.9	2.3	6.1	14.8	37.5

All races 3/, female

2006.....	1.1	*	*	*	*	*	0.1	0.4	0.8	2.3	6.5	20.1
2005.....	1.2	*	*	*	*	*	0.2	0.4	0.8	2.4	6.8	21.0
2004.....	1.3	*	*	*	*	*	0.1	0.4	1.0	2.8	8.0	24.3
2003.....	1.4	*	*	*	*	*	0.2	0.4	1.0	2.4	8.7	25.3
2002.....	1.5	*	*	*	*	*	0.2	0.5	1.2	2.9	8.7	26.9
2001.....	1.7	*	*	*	*	*	0.1	0.5	1.1	3.2	10.4	32.7
2000.....	1.6	*	*	*	*	*	0.1	0.5	1.3	3.0	10.0	33.5
1999.....	1.7	*	*	*	*	*	0.2	0.5	1.3	3.3	10.6	33.5



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Peptic ulcer (K25-K28)

White, both sexes

2006 4/.....	1.2	*	*	*	*	*	0.1	0.6	1.1	2.6	7.2	21.2
2005 4/.....	1.2	*	*	*	*	*	0.2	0.6	1.2	2.9	8.0	21.6
2004 4/.....	1.3	*	*	*	*	*	0.2	0.6	1.2	3.1	8.9	25.1
2003 4/.....	1.4	*	*	*	*	0.1	0.2	0.6	1.3	3.2	9.8	26.9
2002.....	1.5	*	*	*	*	*	0.2	0.6	1.4	3.6	10.3	29.1
2001.....	1.7	*	*	*	*	*	0.2	0.7	1.5	3.9	11.5	34.4
2000.....	1.7	*	*	*	*	*	0.2	0.7	1.8	4.0	11.5	36.3
1999.....	1.8	*	*	*	*	*	0.2	0.6	1.7	4.4	12.3	35.3

White, male

2006 4/.....	1.1	*	*	*	*	*	0.1	0.8	1.6	3.0	8.2	22.0
2005 4/.....	1.2	*	*	*	*	*	0.3	0.8	1.6	3.4	9.7	22.5
2004 4/.....	1.2	*	*	*	*	*	0.2	0.8	1.6	3.6	10.0	25.5
2003 4/.....	1.4	*	*	*	*	*	0.2	0.8	1.6	4.1	11.4	29.0
2002.....	1.5	*	*	*	*	*	0.3	0.8	1.8	4.4	12.3	32.5
2001.....	1.5	*	*	*	*	*	0.3	0.8	2.0	4.5	12.9	35.7
2000.....	1.7	*	*	*	*	*	0.3	1.0	2.4	5.2	13.5	40.7
1999.....	1.7	*	*	*	*	*	0.2	0.8	2.2	5.9	14.9	38.4

White, female

2006 4/.....	1.3	*	*	*	*	*	0.1	0.4	0.8	2.3	6.6	20.8
2005 4/.....	1.3	*	*	*	*	*	0.2	0.4	0.8	2.4	6.8	21.2
2004 4/.....	1.5	*	*	*	*	*	0.1	0.4	1.0	2.7	8.2	25.0
2003 4/.....	1.5	*	*	*	*	*	0.2	0.4	1.0	2.5	8.8	26.0
2002.....	1.6	*	*	*	*	*	0.1	0.5	1.1	2.9	9.0	27.6
2001.....	1.9	*	*	*	*	*	0.1	0.5	1.1	3.3	10.7	33.9
2000.....	1.8	*	*	*	*	*	0.1	0.4	1.2	3.0	10.2	34.5
1999.....	1.9	*	*	*	*	*	0.2	0.4	1.3	3.2	10.7	34.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Peptic ulcer (K25-K28)

Black, both sexes

2006 4/.....	0.9	*	*	*	*	*	0.4	1.3	2.0	3.7	7.2	14.8
2005 4/.....	0.9	*	*	*	*	*	0.5	1.2	1.6	3.9	7.4	17.4
2004 4/.....	0.9	*	*	*	*	*	0.5	1.3	2.0	4.2	7.7	16.1
2003 4/.....	1.0	*	*	*	*	*	0.4	1.5	2.2	3.2	11.0	16.9
2002.....	1.0	*	*	*	*	*	0.6	1.2	2.6	4.1	8.0	19.0
2001.....	1.1	*	*	*	*	*	0.6	1.3	2.7	4.4	9.1	20.9
2000.....	1.2	*	*	*	*	*	0.5	1.7	2.7	4.8	10.0	26.6
1999.....	1.2	*	*	*	*	*	0.8	1.5	2.8	5.7	10.9	25.5

Black, male

2006 4/.....	1.1	*	*	*	*	*	*	2.2	2.8	5.7	9.2	*
2005 4/.....	1.1	*	*	*	*	*	0.7	2.1	2.6	5.7	9.0	20.5
2004 4/.....	1.0	*	*	*	*	*	*	1.8	2.9	4.6	11.1	*
2003 4/.....	1.2	*	*	*	*	*	*	2.3	3.4	5.2	16.1	*
2002.....	1.2	*	*	*	*	*	0.8	1.9	3.8	5.9	11.5	*
2001.....	1.4	*	*	*	*	*	1.0	2.2	4.2	7.0	12.4	23.8
2000.....	1.3	*	*	*	*	*	0.8	2.3	3.7	7.0	13.9	31.1
1999.....	1.4	*	*	*	*	*	1.2	2.3	4.4	8.1	12.3	28.1

Black, female

2006 4/.....	0.7	*	*	*	*	*	*	*	1.5	2.3	6.0	13.9
2005 4/.....	0.7	*	*	*	*	*	*	*	*	2.6	6.5	16.1
2004 4/.....	0.9	*	*	*	*	*	*	0.8	1.3	4.0	5.8	16.5
2003 4/.....	0.8	*	*	*	*	*	*	0.8	*	*	8.1	15.8
2002.....	0.8	*	*	*	*	*	*	*	1.6	2.8	6.0	18.5
2001.....	0.9	*	*	*	*	*	*	*	1.5	2.7	7.3	19.8
2000.....	1.0	*	*	*	*	*	*	1.1	1.8	3.3	7.8	24.9
1999.....	1.1	*	*	*	*	*	*	*	*	3.9	10.2	24.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Diseases of appendix (K35-K38)

All races 3/, both sexes

2006.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.8	1.7
2005.....	0.1	*	*	*	*	*	*	0.1	0.2	0.4	0.8	2.1
2004.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	1.7
2003.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.3	1.0	1.7
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.9	2.0
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	0.7	2.1
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.4	1.0	1.6
1999.....	0.1	*	*	0.1	*	*	0.1	0.1	0.2	0.4	0.8	1.5

All races 3/, male

2006.....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.2	2.3
2005.....	0.2	*	*	*	*	*	*	*	0.3	0.6	1.1	2.8
2004.....	0.2	*	*	*	*	*	*	0.2	0.2	0.5	1.3	2.5
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.5	2.4
2002.....	0.2	*	*	*	*	*	*	0.2	0.4	0.7	1.3	2.2
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.8
2000.....	0.2	*	*	*	*	*	*	0.2	0.4	0.5	1.2	2.0
1999.....	0.2	*	*	*	*	*	*	0.2	0.3	0.5	1.0	2.9

All races 3/, female

2006.....	0.1	*	*	*	*	*	*	*	0.2	0.3	0.4	1.5
2005.....	0.1	*	*	*	*	*	*	*	0.1	0.2	0.7	1.8
2004.....	0.1	*	*	*	*	*	*	*	0.2	0.4	0.9	1.4
2003.....	0.1	*	*	*	*	*	*	0.1	*	*	0.7	1.3
2002.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	2.0
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.8
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.8	1.5
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Diseases of appendix (K35-K38)

White, both sexes

2006 4/.....	0.1	*	*	*	*	*	0.1	0.1	0.2	0.4	0.7	1.8
2005 4/.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	0.8	2.2
2004 4/.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	1.8
2003 4/.....	0.2	*	*	*	*	*	*	0.1	0.1	0.3	1.0	1.7
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.0	2.1
2001.....	0.1	*	*	*	*	*	*	0.1	0.1	0.5	0.7	2.1
2000.....	0.2	*	*	*	*	*	*	0.1	0.2	0.3	0.9	1.7
1999.....	0.1	*	*	*	*	*	*	0.1	0.1	0.4	0.8	1.5

White, male

2006 4/.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.2	2.3
2005 4/.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.1	2.9
2004 4/.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.3	2.3
2003 4/.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.4	2.5
2002.....	0.2	*	*	*	*	*	*	0.2	0.4	0.6	1.4	2.0
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.7	1.0	2.5
2000.....	0.2	*	*	*	*	*	*	*	0.3	0.4	1.2	2.0
1999.....	0.2	*	*	*	*	*	*	0.2	0.2	0.4	0.9	3.1

White, female

2006 4/.....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.5
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.7	1.9
2004 4/.....	0.1	*	*	*	*	*	*	*	*	0.4	0.8	1.6
2003 4/.....	0.1	*	*	*	*	*	*	*	*	*	0.7	1.3
2002.....	0.2	*	*	*	*	*	*	*	*	0.3	0.7	2.1
2001.....	0.1	*	*	*	*	*	*	*	*	0.3	0.5	1.9
2000.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	1.6
1999.....	0.1	*	*	*	*	*	*	*	*	0.3	0.7	0.9



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Hernia (K40-K46)

All races 3/, both sexes

2006.....	0.6	0.8	*	*	*	*	0.1	0.2	0.6	1.2	3.9	12.1
2005.....	0.6	1.0	*	*	*	*	0.1	0.2	0.6	1.1	3.5	12.1
2004.....	0.5	1.0	*	*	*	*	0.1	0.2	0.4	1.3	3.9	11.0
2003.....	0.6	0.9	*	*	*	*	0.1	0.3	0.6	1.3	3.6	11.9
2002.....	0.6	0.7	*	*	*	*	0.1	0.2	0.5	1.2	3.8	12.8
2001.....	0.5	1.1	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.6
2000.....	0.5	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.1
1999.....	0.5	1.2	*	*	*	*	0.1	0.2	0.4	1.2	3.7	12.7

All races 3/, male

2006.....	0.5	0.9	*	*	*	*	0.1	0.2	0.5	1.3	4.1	13.5
2005.....	0.4	1.3	*	*	*	*	*	0.2	0.6	1.0	3.3	12.7
2004.....	0.5	1.0	*	*	*	*	0.1	0.2	0.4	1.4	3.8	11.3
2003.....	0.5	1.0	*	*	*	*	*	0.3	0.5	1.3	3.9	12.3
2002.....	0.4	*	*	*	*	*	*	0.2	0.4	1.2	3.7	11.7
2001.....	0.4	1.4	*	*	*	*	*	0.2	0.5	1.1	3.5	12.0
2000.....	0.4	1.4	*	*	*	*	*	0.2	0.4	1.1	4.0	12.6
1999.....	0.4	1.3	*	*	*	*	0.1	0.2	0.5	1.1	4.0	13.2

All races 3/, female

2006.....	0.7	*	*	*	*	*	*	0.2	0.6	1.2	3.8	11.4
2005.....	0.7	*	*	*	*	*	0.1	0.2	0.6	1.2	3.6	11.8
2004.....	0.6	1.0	*	*	*	*	*	0.2	0.5	1.2	4.0	10.8
2003.....	0.6	*	*	*	*	*	*	0.2	0.6	1.3	3.4	11.8
2002.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.9	13.3
2001.....	0.6	*	*	*	*	*	*	0.2	0.5	1.2	3.6	12.8
2000.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.3
1999.....	0.6	*	*	*	*	*	*	0.1	0.4	1.2	3.5	12.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Hernia (K40-K46)

White, both sexes

2006 4/.....	0.6	0.8	*	*	*	*	0.1	0.2	0.6	1.2	3.9	12.6
2005 4/.....	0.6	1.1	*	*	*	*	0.1	0.2	0.6	1.1	3.6	12.5
2004 4/.....	0.6	1.0	*	*	*	*	0.1	0.2	0.4	1.3	4.1	11.2
2003 4/.....	0.6	0.9	*	*	*	*	0.1	0.3	0.5	1.3	3.6	12.3
2002.....	0.6	0.8	*	*	*	*	0.1	0.2	0.4	1.2	4.0	13.4
2001.....	0.6	1.0	*	*	*	*	0.1	0.2	0.5	1.2	3.5	12.8
2000.....	0.6	1.2	*	*	*	*	0.1	0.2	0.5	1.1	3.8	13.3
1999.....	0.6	1.1	*	*	*	*	0.1	0.1	0.5	1.1	3.7	13.0

White, male

2006 4/.....	0.5	*	*	*	*	*	*	0.2	0.5	1.3	4.0	14.0
2005 4/.....	0.5	1.3	*	*	*	*	*	0.2	0.6	1.0	3.3	12.8
2004 4/.....	0.5	*	*	*	*	*	*	0.2	0.4	1.4	3.9	11.4
2003 4/.....	0.5	*	*	*	*	*	*	0.3	0.5	1.3	3.9	12.5
2002.....	0.5	*	*	*	*	*	*	0.2	0.4	1.2	3.9	11.9
2001.....	0.4	*	*	*	*	*	*	0.2	0.5	1.2	3.4	11.8
2000.....	0.5	1.5	*	*	*	*	*	0.2	0.4	1.1	4.1	12.7
1999.....	0.5	1.3	*	*	*	*	0.1	0.2	0.6	1.2	4.1	12.9

White, female

2006 4/.....	0.7	*	*	*	*	*	*	0.2	0.6	1.2	3.9	11.9
2005 4/.....	0.7	*	*	*	*	*	*	0.2	0.6	1.2	3.7	12.3
2004 4/.....	0.7	*	*	*	*	*	*	0.2	0.4	1.2	4.2	11.2
2003 4/.....	0.7	*	*	*	*	*	*	0.2	0.6	1.3	3.4	12.2
2002.....	0.8	*	*	*	*	*	*	0.2	0.5	1.2	4.1	14.0
2001.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.2
2000.....	0.7	*	*	*	*	*	*	0.2	0.5	1.2	3.6	13.6
1999.....	0.7	*	*	*	*	*	*	*	0.4	1.1	3.5	13.1





Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes

2006.....	9.2	*	*	*	0.1	0.8	5.8	17.8	22.8	26.0	29.0	19.4
2005.....	9.3	*	*	*	0.1	0.8	6.1	17.7	23.5	27.2	29.0	19.7
2004.....	9.2	*	*	*	*	0.8	6.3	18.0	22.6	27.7	28.8	19.7
2003.....	9.5	*	*	*	*	0.9	6.8	18.3	23.0	29.5	30.0	20.1
2002.....	9.5	*	*	*	0.1	0.9	7.0	18.0	22.9	29.4	31.4	21.4
2001.....	9.5	*	*	*	0.1	1.0	7.4	18.5	22.7	30.0	30.2	22.2
2000.....	9.4	*	*	*	0.1	1.0	7.5	17.7	23.8	29.8	31.0	23.1
1999.....	9.4	*	*	*	0.1	1.0	7.3	17.4	23.7	30.6	31.9	23.2

All races 3/, male

2006.....	12.1	*	*	*	*	1.0	7.6	25.9	33.5	34.7	35.9	26.1
2005.....	12.3	*	*	*	*	1.0	8.1	26.0	34.8	35.8	36.4	27.6
2004.....	12.1	*	*	*	*	1.1	8.7	26.3	32.8	37.0	36.7	25.5
2003.....	12.5	*	*	*	*	1.2	9.2	27.5	33.8	39.1	36.9	27.5
2002.....	12.3	*	*	*	*	1.2	9.3	26.9	33.1	38.5	38.4	28.4
2001.....	12.4	*	*	*	*	1.3	9.8	27.5	33.1	40.2	36.8	32.0
2000.....	12.5	*	*	*	*	1.4	10.3	26.9	34.4	39.9	38.4	34.3
1999.....	12.5	*	*	*	*	1.3	10.3	26.9	34.6	41.6	39.6	33.1

All races 3/, female

2006.....	6.4	*	*	*	*	0.5	4.1	10.0	13.0	18.6	24.2	16.2
2005.....	6.4	*	*	*	*	0.6	4.1	9.7	12.9	19.9	23.9	16.0
2004.....	6.3	*	*	*	*	0.4	4.0	10.0	13.1	19.9	23.4	17.1
2003.....	6.5	*	*	*	*	0.5	4.4	9.4	13.1	21.4	25.4	16.8
2002.....	6.7	*	*	*	*	0.7	4.8	9.5	13.5	21.9	26.7	18.4
2001.....	6.7	*	*	*	*	0.7	5.0	9.8	13.1	21.5	25.9	18.1
2000.....	6.5	*	*	*	*	0.7	4.7	8.8	14.0	21.5	26.1	18.5
1999.....	6.4	*	*	*	*	0.8	4.4	8.3	13.7	21.6	26.9	19.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

White, both sexes

2006 4/.....	10.0	*	*	*	*	0.8	6.1	18.5	23.5	27.2	30.3	20.2
2005 4/.....	10.0	*	*	*	*	0.7	6.5	18.1	23.8	28.5	29.9	20.2
2004 4/.....	9.9	*	*	*	*	0.8	6.5	18.4	22.7	28.9	30.2	20.3
2003 4/.....	10.2	*	*	*	*	0.9	7.1	18.4	23.1	30.7	31.6	20.9
2002.....	10.1	*	*	*	0.1	0.9	7.3	18.1	23.1	30.7	32.8	22.3
2001.....	10.1	*	*	*	*	1.0	7.5	18.3	22.7	30.9	31.5	23.2
2000.....	10.0	*	*	*	0.1	1.0	7.4	17.5	23.8	30.6	32.3	23.9
1999.....	9.9	*	*	*	*	1.0	7.2	17.0	23.3	31.4	33.0	23.8

White, male

2006 4/.....	13.1	*	*	*	*	1.1	8.0	26.9	34.2	36.0	37.3	27.2
2005 4/.....	13.2	*	*	*	*	0.9	8.6	26.5	35.1	37.2	37.2	28.5
2004 4/.....	13.0	*	*	*	*	1.1	9.0	26.9	32.7	38.2	38.4	26.7
2003 4/.....	13.4	*	*	*	*	1.3	9.6	27.8	33.8	40.5	38.5	28.4
2002.....	13.2	*	*	*	*	1.2	9.7	27.1	33.2	39.8	40.2	29.6
2001.....	13.1	*	*	*	*	1.3	9.9	27.3	32.8	41.1	38.4	33.5
2000.....	13.2	*	*	*	*	1.4	10.3	26.8	34.5	40.9	39.9	35.5
1999.....	13.1	*	*	*	*	1.3	10.3	26.4	34.1	42.4	40.8	34.2

White, female

2006 4/.....	6.9	*	*	*	*	0.5	4.1	10.3	13.3	19.7	25.4	16.9
2005 4/.....	6.9	*	*	*	*	0.5	4.3	9.7	13.1	21.1	24.9	16.4
2004 4/.....	6.8	*	*	*	*	0.5	4.0	10.1	13.1	21.0	24.6	17.4
2003 4/.....	7.0	*	*	*	*	0.5	4.4	9.2	13.1	22.4	27.0	17.5
2002.....	7.2	*	*	*	*	0.6	4.8	9.1	13.5	23.0	27.9	19.1
2001.....	7.1	*	*	*	*	0.6	5.0	9.4	13.3	22.4	27.0	18.9
2000.....	6.8	*	*	*	*	0.6	4.5	8.2	13.7	21.9	27.2	19.2
1999.....	6.8	*	*	*	*	0.7	4.2	7.8	13.2	22.2	27.9	19.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic liver disease and cirrhosis (K70,K73-K74)

Black, both sexes

2006 4/.....	5.9	*	*	*	*	0.4	4.5	15.5	22.0	18.9	16.4	10.6
2005 4/.....	6.5	*	*	*	*	0.6	4.2	17.6	24.9	20.2	19.1	12.3
2004 4/.....	6.6	*	*	*	*	0.4	5.4	18.3	25.3	21.6	14.4	12.1
2003 4/.....	6.9	*	*	*	*	0.7	5.7	19.8	25.8	23.8	14.4	12.3
2002.....	6.9	*	*	*	*	0.9	5.5	20.8	25.0	21.6	18.5	12.2
2001.....	7.5	*	*	*	*	0.9	7.3	22.6	25.9	25.0	18.5	10.4
2000.....	7.5	*	*	*	*	1.0	7.8	22.1	25.9	26.2	18.0	13.8
1999.....	7.8	*	*	*	*	1.0	8.0	22.7	29.3	26.6	21.5	17.2

Black, male

2006 4/.....	8.0	*	*	*	*	*	5.8	22.6	33.5	29.3	24.6	*
2005 4/.....	8.9	*	*	*	*	*	5.4	25.7	39.2	31.2	29.0	*
2004 4/.....	9.1	*	*	*	*	*	7.4	27.1	39.1	33.8	20.6	*
2003 4/.....	9.5	*	*	*	*	1.0	7.1	30.4	40.8	33.4	20.6	21.9
2002.....	9.0	*	*	*	*	1.0	6.7	30.0	37.8	32.4	22.4	*
2001.....	10.4	*	*	*	*	1.0	10.0	34.5	41.5	39.1	22.4	*
2000.....	9.9	*	*	*	*	1.0	10.5	32.7	38.0	37.7	24.5	*
1999.....	10.7	*	*	*	*	1.1	11.1	35.4	43.3	40.7	29.5	*

Black, female

2006 4/.....	4.0	*	*	*	*	*	3.3	9.4	12.8	11.4	11.7	8.8
2005 4/.....	4.3	*	*	*	*	*	3.1	10.6	13.4	12.3	13.5	10.9
2004 4/.....	4.3	*	*	*	*	*	3.7	10.8	14.2	12.9	10.9	11.4
2003 4/.....	4.5	*	*	*	*	*	4.4	10.6	13.7	16.9	10.8	8.5
2002.....	4.9	*	*	*	*	0.8	4.4	12.9	14.7	14.0	16.2	10.7
2001.....	4.8	*	*	*	*	0.7	4.8	12.4	13.4	15.0	16.4	9.7
2000.....	5.2	*	*	*	*	1.0	5.5	13.1	16.2	18.1	14.3	11.6
1999.....	5.2	*	*	*	*	0.9	5.3	11.8	18.1	16.7	17.0	15.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcoholic liver disease (K70)

All races 3/, both sexes

2006.....	4.4	*	*	*	*	0.6	4.0	10.5	12.1	9.4	5.9	2.2
2005.....	4.4	*	*	*	*	0.5	4.1	10.6	12.3	9.9	5.7	1.8
2004.....	4.3	*	*	*	*	0.6	4.2	10.6	11.7	10.0	5.5	2.1
2003.....	4.3	*	*	*	*	0.6	4.4	10.5	11.6	9.8	5.7	1.8
2002.....	4.2	*	*	*	*	0.6	4.4	10.3	11.6	10.1	5.7	2.1
2001.....	4.3	*	*	*	*	0.7	4.8	10.7	11.5	10.1	5.7	2.4
2000.....	4.3	*	*	*	*	0.7	4.9	10.3	12.2	10.3	6.1	2.2
1999.....	4.3	*	*	*	*	0.6	4.8	10.4	12.0	10.5	6.5	2.8

All races 3/, male

2006.....	6.4	*	*	*	*	0.9	5.1	15.5	18.9	15.7	10.2	4.7
2005.....	6.5	*	*	*	*	0.7	5.4	15.7	19.5	15.9	10.2	4.2
2004.....	6.3	*	*	*	*	0.8	5.8	15.5	18.2	16.5	10.1	4.6
2003.....	6.4	*	*	*	*	0.9	6.0	16.1	18.3	16.1	10.1	3.6
2002.....	6.2	*	*	*	*	0.8	5.8	15.6	18.3	16.5	9.7	4.7
2001.....	6.4	*	*	*	*	0.9	6.5	16.1	18.4	16.6	10.1	5.5
2000.....	6.5	*	*	*	*	1.0	6.8	16.0	19.0	17.3	11.5	5.3
1999.....	6.5	*	*	*	*	0.8	6.8	16.3	19.2	18.1	11.2	6.1

All races 3/, female

2006.....	2.4	*	*	*	*	0.4	2.8	5.8	5.8	4.2	3.0	1.1
2005.....	2.3	*	*	*	*	0.4	2.8	5.6	5.5	4.9	2.7	0.7
2004.....	2.3	*	*	*	*	0.3	2.7	5.7	5.6	4.5	2.3	1.0
2003.....	2.2	*	*	*	*	0.3	2.9	5.1	5.4	4.5	2.7	1.0
2002.....	2.3	*	*	*	*	0.4	3.0	5.2	5.4	4.8	3.1	1.0
2001.....	2.3	*	*	*	*	0.4	3.1	5.5	5.1	4.7	2.8	1.0
2000.....	2.2	*	*	*	*	0.4	3.1	4.8	6.0	4.5	2.5	1.0
1999.....	2.1	*	*	*	*	0.5	2.9	4.7	5.4	4.3	3.4	1.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcoholic liver disease (K70)

White, both sexes

2006 4/.....	4.7	*	*	*	*	0.6	4.2	11.1	12.6	9.7	6.2	2.4
2005 4/.....	4.7	*	*	*	*	0.5	4.3	11.0	12.6	10.2	5.9	1.9
2004 4/.....	4.6	*	*	*	*	0.6	4.4	11.0	11.7	10.3	5.7	2.2
2003 4/.....	4.5	*	*	*	*	0.6	4.6	10.7	11.7	10.0	5.9	1.9
2002.....	4.4	*	*	*	*	0.6	4.6	10.5	11.7	10.4	5.8	2.1
2001.....	4.5	*	*	*	*	0.7	4.8	10.8	11.5	10.2	5.8	2.3
2000.....	4.5	*	*	*	*	0.7	4.9	10.3	12.2	10.5	6.1	2.3
1999.....	4.4	*	*	*	*	0.6	4.8	10.2	11.8	10.5	6.4	2.7

White, male

2006 4/.....	7.0	*	*	*	*	0.9	5.5	16.3	19.6	16.0	10.5	4.9
2005 4/.....	6.9	*	*	*	*	0.7	5.7	16.3	19.9	16.3	10.4	4.4
2004 4/.....	6.8	*	*	*	*	0.9	6.0	16.2	18.3	16.8	10.4	4.8
2003 4/.....	6.8	*	*	*	*	0.9	6.2	16.4	18.5	16.4	10.4	3.7
2002.....	6.6	*	*	*	*	0.8	6.1	16.0	18.5	16.8	9.9	4.5
2001.....	6.7	*	*	*	*	0.9	6.5	16.3	18.4	16.6	10.4	5.4
2000.....	6.8	*	*	*	*	1.0	6.8	16.2	19.2	17.5	11.7	5.4
1999.....	6.7	*	*	*	*	0.8	6.7	16.2	19.0	17.9	11.1	5.8

White, female

2006 4/.....	2.5	*	*	*	*	0.3	2.8	6.0	6.0	4.3	3.1	1.1
2005 4/.....	2.4	*	*	*	*	0.3	2.9	5.8	5.6	5.0	2.8	0.8
2004 4/.....	2.4	*	*	*	*	0.3	2.8	5.8	5.6	4.7	2.4	1.0
2003 4/.....	2.3	*	*	*	*	0.3	3.0	5.1	5.3	4.5	2.9	1.1
2002.....	2.3	*	*	*	*	0.3	3.0	5.1	5.2	5.0	3.1	1.1
2001.....	2.3	*	*	*	*	0.4	3.1	5.3	5.1	4.8	2.8	1.0
2000.....	2.1	*	*	*	*	0.3	3.0	4.4	5.6	4.6	2.4	1.0
1999.....	2.1	*	*	*	*	0.4	2.8	4.4	5.0	4.3	3.3	1.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcoholic liver disease (K70)

Black, both sexes

2006 4/.....	2.6	*	*	*	*	*	2.5	7.6	9.4	7.8	4.1	*
2005 4/.....	3.0	*	*	*	*	*	2.7	8.8	11.3	8.7	4.6	*
2004 4/.....	3.0	*	*	*	*	*	2.9	9.0	12.5	8.9	3.1	*
2003 4/.....	3.2	*	*	*	*	*	3.3	9.9	12.5	9.3	3.4	*
2002.....	3.2	*	*	*	*	0.5	2.9	10.0	12.3	8.6	5.3	*
2001.....	3.6	*	*	*	*	0.5	4.2	11.5	12.9	10.7	4.7	*
2000.....	3.8	*	*	*	*	0.6	5.0	12.0	13.3	10.3	5.5	*
1999.....	4.1	*	*	*	*	0.5	5.0	12.8	15.1	11.8	7.1	*

Black, male

2006 4/.....	3.8	*	*	*	*	*	3.1	11.2	14.9	14.0	7.6	*
2005 4/.....	4.3	*	*	*	*	*	3.1	13.2	18.3	15.3	9.0	*
2004 4/.....	4.4	*	*	*	*	*	3.9	12.9	20.3	16.9	*	*
2003 4/.....	4.8	*	*	*	*	*	4.1	15.8	20.1	16.0	6.8	*
2002.....	4.4	*	*	*	*	*	3.8	14.7	18.7	15.0	9.2	*
2001.....	5.3	*	*	*	*	*	5.9	17.1	21.4	19.5	7.7	*
2000.....	5.3	*	*	*	*	*	6.9	17.4	19.4	18.1	10.0	*
1999.....	5.9	*	*	*	*	*	6.7	19.7	24.2	21.8	12.6	*

Black, female

2006 4/.....	1.6	*	*	*	*	*	2.0	4.4	5.0	3.4	*	*
2005 4/.....	1.8	*	*	*	*	*	2.2	5.1	5.7	4.0	*	*
2004 4/.....	1.8	*	*	*	*	*	1.9	5.6	6.2	3.3	*	*
2003 4/.....	1.8	*	*	*	*	*	2.5	4.7	6.4	4.5	*	*
2002.....	2.0	*	*	*	*	*	2.2	5.9	7.1	4.1	*	*
2001.....	2.1	*	*	*	*	*	2.7	6.7	6.1	4.4	*	*
2000.....	2.4	*	*	*	*	*	3.3	7.3	8.4	4.7	*	*
1999.....	2.4	*	*	*	*	*	3.4	6.9	7.9	4.8	4.0	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic liver disease and cirrhosis (K73-K74)

All races 3/, both sexes

2006.....	4.8	*	*	*	*	0.2	1.9	7.3	10.7	16.6	23.0	17.2
2005.....	4.9	*	*	*	*	0.2	2.0	7.1	11.2	17.3	23.2	17.8
2004.....	4.9	*	*	*	*	0.2	2.1	7.5	10.9	17.8	23.3	17.6
2003.....	5.2	*	*	*	*	0.3	2.4	7.8	11.4	19.7	24.3	18.2
2002.....	5.2	*	*	*	0.1	0.3	2.6	7.7	11.3	19.3	25.7	19.3
2001.....	5.2	*	*	*	*	0.3	2.6	7.8	11.2	19.9	24.5	19.9
2000.....	5.1	*	*	*	0.1	0.3	2.5	7.3	11.6	19.5	24.9	20.9
1999.....	5.1	*	*	*	*	0.4	2.5	7.0	11.7	20.1	25.5	20.4

All races 3/, male

2006.....	5.7	*	*	*	*	0.2	2.5	10.4	14.6	19.0	25.7	21.4
2005.....	5.8	*	*	*	*	0.2	2.7	10.3	15.3	19.9	26.2	23.3
2004.....	5.8	*	*	*	*	0.3	2.9	10.8	14.6	20.5	26.6	21.0
2003.....	6.2	*	*	*	*	0.3	3.2	11.4	15.5	23.0	26.8	23.9
2002.....	6.1	*	*	*	*	0.4	3.5	11.2	14.8	22.0	28.7	23.7
2001.....	6.1	*	*	*	*	0.4	3.4	11.4	14.7	23.6	26.6	26.5
2000.....	6.0	*	*	*	*	0.4	3.5	10.9	15.4	22.7	26.9	29.0
1999.....	6.0	*	*	*	*	0.5	3.5	10.6	15.3	23.6	28.4	27.0

All races 3/, female

2006.....	4.0	*	*	*	*	0.2	1.3	4.2	7.1	14.5	21.2	15.2
2005.....	4.0	*	*	*	*	0.2	1.3	4.1	7.4	15.1	21.2	15.3
2004.....	4.1	*	*	*	*	0.1	1.3	4.2	7.5	15.4	21.1	16.1
2003.....	4.3	*	*	*	*	0.2	1.5	4.3	7.6	17.0	22.6	15.8
2002.....	4.5	*	*	*	*	0.3	1.8	4.3	8.1	17.1	23.7	17.4
2001.....	4.4	*	*	*	*	0.2	1.9	4.3	8.0	16.8	23.1	17.1
2000.....	4.3	*	*	*	*	0.3	1.6	4.0	8.0	16.9	23.6	17.5
1999.....	4.3	*	*	*	*	0.3	1.4	3.6	8.3	17.3	23.5	17.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic liver disease and cirrhosis (K73-K74)

White, both sexes

2006 4/.....	5.3	*	*	*	*	0.2	1.9	7.4	10.8	17.5	24.1	17.8
2005 4/.....	5.3	*	*	*	*	0.2	2.2	7.1	11.2	18.3	24.0	18.3
2004 4/.....	5.3	*	*	*	*	0.2	2.1	7.5	10.9	18.6	24.6	18.1
2003 4/.....	5.7	*	*	*	*	0.3	2.5	7.7	11.4	20.8	25.7	19.0
2002.....	5.7	*	*	*	0.1	0.3	2.7	7.5	11.4	20.3	27.0	20.2
2001.....	5.6	*	*	*	*	0.3	2.7	7.5	11.2	20.7	25.7	20.9
2000.....	5.5	*	*	*	*	0.3	2.5	7.2	11.6	20.1	26.1	21.6
1999.....	5.5	*	*	*	*	0.4	2.5	6.8	11.5	20.9	26.6	21.1

White, male

2006 4/.....	6.2	*	*	*	*	0.2	2.5	10.6	14.6	19.9	26.7	22.3
2005 4/.....	6.2	*	*	*	*	0.2	2.9	10.3	15.1	20.9	26.8	24.2
2004 4/.....	6.2	*	*	*	*	0.3	3.0	10.7	14.5	21.3	28.0	22.0
2003 4/.....	6.6	*	*	*	*	0.3	3.4	11.4	15.3	24.1	28.1	24.7
2002.....	6.5	*	*	*	*	0.4	3.7	11.1	14.7	23.0	30.3	25.1
2001.....	6.4	*	*	*	*	0.4	3.4	11.0	14.4	24.4	28.0	28.1
2000.....	6.4	*	*	*	*	0.3	3.5	10.6	15.3	23.4	28.2	30.1
1999.....	6.4	*	*	*	*	0.5	3.5	10.2	15.1	24.4	29.7	28.4

White, female

2006 4/.....	4.4	*	*	*	*	0.2	1.3	4.3	7.3	15.4	22.3	15.7
2005 4/.....	4.4	*	*	*	*	0.2	1.4	3.9	7.5	16.1	22.1	15.6
2004 4/.....	4.5	*	*	*	*	0.1	1.2	4.2	7.5	16.4	22.2	16.4
2003 4/.....	4.7	*	*	*	*	0.2	1.5	4.1	7.8	17.9	24.0	16.5
2002.....	4.9	*	*	*	*	0.3	1.8	4.0	8.3	18.0	24.8	18.1
2001.....	4.8	*	*	*	*	0.2	1.9	4.2	8.2	17.6	24.2	17.9
2000.....	4.7	*	*	*	*	0.2	1.5	3.8	8.1	17.3	24.8	18.2
1999.....	4.6	*	*	*	*	0.3	1.4	3.4	8.2	17.9	24.6	18.2



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other chronic liver disease and cirrhosis (K73-K74)

Black, both sexes

2006 4/.....	3.3	*	*	*	*	*	2.0	7.9	12.6	11.0	12.3	9.9
2005 4/.....	3.5	*	*	*	*	*	1.5	8.8	13.6	11.5	14.5	10.7
2004 4/.....	3.5	*	*	*	*	*	2.5	9.3	12.8	12.6	11.3	11.3
2003 4/.....	3.7	*	*	*	*	0.4	2.4	9.9	13.3	14.5	11.0	10.8
2002.....	3.7	*	*	*	*	0.5	2.5	10.8	12.7	13.0	13.2	9.8
2001.....	3.8	*	*	*	*	0.4	3.1	11.1	13.0	14.3	13.9	7.7
2000.....	3.7	*	*	*	*	0.4	2.8	10.1	12.6	16.0	12.5	12.2
1999.....	3.8	*	*	*	*	0.5	3.0	9.9	14.2	14.8	14.5	12.8

Black, male

2006 4/.....	4.3	*	*	*	*	*	2.7	11.4	18.6	15.3	17.0	*
2005 4/.....	4.6	*	*	*	*	*	2.2	12.5	20.9	15.9	19.9	*
2004 4/.....	4.7	*	*	*	*	*	3.5	14.2	18.8	16.9	15.3	*
2003 4/.....	4.7	*	*	*	*	*	3.0	14.6	20.8	17.3	13.8	*
2002.....	4.6	*	*	*	*	*	2.9	15.3	19.1	17.4	13.2	*
2001.....	5.1	*	*	*	*	*	4.1	17.4	20.1	19.5	14.7	*
2000.....	4.7	*	*	*	*	*	3.6	15.2	18.6	19.6	14.5	*
1999.....	4.8	*	*	*	*	*	4.4	15.8	19.1	19.0	16.9	*

Black, female

2006 4/.....	2.4	*	*	*	*	*	1.4	4.9	7.8	8.0	9.5	8.8
2005 4/.....	2.5	*	*	*	*	*	0.9	5.5	7.7	8.4	11.5	10.1
2004 4/.....	2.5	*	*	*	*	*	1.7	5.1	8.0	9.6	9.0	11.0
2003 4/.....	2.7	*	*	*	*	*	1.9	5.9	7.3	12.4	9.4	8.1
2002.....	2.9	*	*	*	*	*	2.2	7.0	7.6	9.9	13.2	9.9
2001.....	2.7	*	*	*	*	*	2.1	5.7	7.3	10.6	13.4	*
2000.....	2.8	*	*	*	*	*	2.2	5.8	7.8	13.4	11.4	11.2
1999.....	2.9	*	*	*	*	*	1.9	4.9	10.2	11.9	13.1	13.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes

2006.....	1.0	*	*	*	*	0.0	0.1	0.3	0.9	2.5	7.8	21.8
2005.....	1.0	*	*	*	*	*	0.1	0.3	0.7	2.4	8.1	22.9
2004.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.5	7.8	23.8
2003.....	1.0	*	*	*	*	0.1	0.1	0.3	0.7	2.7	8.2	21.8
2002.....	1.0	*	*	*	*	0.1	0.1	0.3	0.8	2.8	7.8	22.9
2001.....	1.0	*	*	*	*	*	0.1	0.3	1.0	2.9	7.5	23.7
2000.....	1.0	*	*	*	*	*	0.1	0.3	0.8	2.4	7.5	24.8
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.6	7.8	23.9

All races 3/, male

2006.....	0.9	*	*	*	*	*	0.1	0.3	0.9	3.1	9.6	21.7
2005.....	0.9	*	*	*	*	*	*	0.3	0.8	3.2	9.4	24.3
2004.....	0.9	*	*	*	*	*	0.1	0.4	1.0	2.9	9.3	25.5
2003.....	0.9	*	*	*	*	*	0.1	0.3	0.8	3.2	10.1	22.7
2002.....	0.9	*	*	*	*	*	0.1	0.4	1.0	3.1	9.8	24.1
2001.....	1.0	*	*	*	*	*	0.1	0.4	1.2	3.2	9.3	26.9
2000.....	0.9	*	*	*	*	*	0.1	0.3	0.9	2.9	9.4	27.6
1999.....	0.9	*	*	*	*	*	0.1	0.3	1.1	3.0	9.3	27.7

All races 3/, female

2006.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.0	6.6	21.9
2005.....	1.1	*	*	*	*	*	0.1	0.3	0.6	1.8	7.2	22.3
2004.....	1.2	*	*	*	*	*	0.1	0.3	0.7	2.1	6.8	23.0
2003.....	1.1	*	*	*	*	*	*	0.3	0.5	2.2	6.9	21.4
2002.....	1.1	*	*	*	*	*	0.1	0.2	0.7	2.5	6.4	22.4
2001.....	1.1	*	*	*	*	*	*	0.3	0.8	2.6	6.3	22.4
2000.....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.1	6.2	23.7
1999.....	1.1	*	*	*	*	*	0.1	0.4	0.7	2.2	6.8	22.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

White, both sexes

2006 4/.....	1.1	*	*	*	*	*	0.1	0.2	0.8	2.2	7.8	22.3
2005 4/.....	1.1	*	*	*	*	*	0.1	0.2	0.7	2.2	8.1	23.1
2004 4/.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.3	7.7	24.2
2003 4/.....	1.1	*	*	*	*	*	0.1	0.3	0.6	2.7	8.2	21.9
2002.....	1.1	*	*	*	*	*	0.1	0.3	0.8	2.7	7.6	23.5
2001.....	1.1	*	*	*	*	*	0.1	0.3	0.9	2.7	7.4	23.9
2000.....	1.1	*	*	*	*	*	0.1	0.2	0.7	2.3	7.4	25.6
1999.....	1.1	*	*	*	*	*	0.1	0.3	0.7	2.5	7.8	24.4

White, male

2006 4/.....	1.0	*	*	*	*	*	*	0.3	0.8	2.8	9.6	22.8
2005 4/.....	1.0	*	*	*	*	*	*	0.2	0.8	3.0	9.6	24.8
2004 4/.....	1.0	*	*	*	*	*	0.1	0.3	0.9	2.9	9.3	26.7
2003 4/.....	1.0	*	*	*	*	*	*	0.3	0.7	3.2	10.0	23.0
2002.....	1.0	*	*	*	*	*	0.1	0.4	0.9	3.1	9.8	24.7
2001.....	1.0	*	*	*	*	*	0.1	0.4	1.1	3.1	9.3	26.7
2000.....	1.0	*	*	*	*	*	0.1	0.2	0.9	2.8	9.6	28.9
1999.....	1.0	*	*	*	*	*	0.1	0.3	0.9	3.0	9.6	28.1

White, female

2006 4/.....	1.2	*	*	*	*	*	*	0.2	0.8	1.7	6.5	22.0
2005 4/.....	1.2	*	*	*	*	*	*	0.2	0.6	1.6	7.0	22.4
2004 4/.....	1.2	*	*	*	*	*	0.1	0.3	0.6	1.9	6.6	23.1
2003 4/.....	1.2	*	*	*	*	*	*	0.2	0.5	2.2	6.9	21.4
2002.....	1.2	*	*	*	*	*	*	0.2	0.6	2.4	6.2	23.0
2001.....	1.2	*	*	*	*	*	*	0.3	0.7	2.4	6.1	22.8
2000.....	1.2	*	*	*	*	*	*	0.2	0.5	1.8	6.0	24.3
1999.....	1.2	*	*	*	*	*	*	0.3	0.6	2.1	6.7	22.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

Black, both sexes

2006 4/.....	0.8	*	*	*	*	*	*	0.6	1.4	3.9	9.0	17.1
2005 4/.....	0.8	*	*	*	*	*	*	0.6	1.1	4.1	9.2	20.6
2004 4/.....	0.8	*	*	*	*	*	*	0.8	1.6	3.5	9.1	19.5
2003 4/.....	0.7	*	*	*	*	*	*	0.7	1.2	2.9	7.9	20.7
2002.....	0.7	*	*	*	*	*	*	0.6	1.1	3.3	9.7	14.8
2001.....	0.8	*	*	*	*	*	*	0.7	1.7	4.7	8.3	19.7
2000.....	0.8	*	*	*	*	*	*	0.7	1.2	3.9	8.3	17.8
1999.....	0.7	*	*	*	*	*	0.4	0.5	1.5	3.6	7.2	15.3

Black, male

2006 4/.....	0.6	*	*	*	*	*	*	*	1.6	3.7	9.5	*
2005 4/.....	0.7	*	*	*	*	*	*	*	*	5.2	10.1	19.5
2004 4/.....	0.7	*	*	*	*	*	*	1.0	1.6	3.6	8.9	*
2003 4/.....	0.7	*	*	*	*	*	*	*	*	3.2	10.4	20.9
2002.....	0.6	*	*	*	*	*	*	*	*	*	10.9	*
2001.....	0.7	*	*	*	*	*	*	*	2.1	4.9	6.2	27.2
2000.....	0.6	*	*	*	*	*	*	*	*	3.7	8.2	*
1999.....	0.6	*	*	*	*	*	*	*	*	3.7	7.1	*

Black, female

2006 4/.....	1.0	*	*	*	*	*	*	*	1.3	4.1	8.8	19.7
2005 4/.....	0.9	*	*	*	*	*	*	*	*	3.3	8.7	21.0
2004 4/.....	1.0	*	*	*	*	*	*	*	1.6	3.5	9.3	21.7
2003 4/.....	0.8	*	*	*	*	*	*	0.8	*	2.7	6.5	20.7
2002.....	0.8	*	*	*	*	*	*	*	*	3.7	9.1	16.0
2001.....	0.9	*	*	*	*	*	*	*	*	4.5	9.4	16.8
2000.....	1.0	*	*	*	*	*	*	*	*	4.1	8.4	19.3
1999.....	0.8	*	*	*	*	*	*	*	*	3.5	7.2	15.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races 3/, both sexes

2006.....	15.1	3.9	*	*	0.2	0.7	1.8	5.2	13.8	39.4	111.4	290.5
2005.....	14.8	3.9	*	0.1	0.2	0.7	1.7	4.8	13.6	39.3	110.3	288.3
2004.....	14.5	4.3	*	0.1	0.2	0.6	1.8	5.0	13.6	38.6	108.4	286.6
2003.....	14.6	4.5	*	0.1	0.2	0.7	1.8	4.9	13.6	40.1	109.5	293.1
2002.....	14.2	4.3	*	0.1	0.2	0.7	1.7	4.7	13.0	39.2	109.1	288.6
2001.....	13.9	3.3	*	*	0.2	0.6	1.7	4.6	13.0	40.2	104.2	287.7
2000.....	13.2	4.3	*	0.1	0.2	0.6	1.6	4.4	12.8	38.0	100.8	277.8
1999.....	12.7	4.4	*	0.1	0.2	0.6	1.6	4.0	12.0	37.1	97.6	268.9

All races 3/, male

2006.....	15.0	4.8	*	*	0.3	0.8	2.1	6.3	16.0	45.1	136.8	377.1
2005.....	14.6	4.2	*	*	0.2	0.7	1.9	5.9	16.1	43.8	135.9	376.5
2004.....	14.1	5.3	*	*	0.2	0.6	2.0	5.9	15.1	43.6	134.3	372.2
2003.....	14.3	5.2	*	*	0.3	0.7	2.1	6.0	15.4	45.3	136.8	387.6
2002.....	13.9	5.3	*	*	0.2	0.6	1.8	5.5	14.3	45.8	136.2	385.5
2001.....	13.5	3.7	*	*	0.2	0.6	1.8	5.5	14.1	46.1	130.0	393.7
2000.....	12.9	5.3	*	*	0.2	0.6	1.9	5.3	14.2	42.9	127.3	380.7
1999.....	12.4	4.6	*	*	0.2	0.6	1.8	4.7	13.4	42.3	125.3	371.1

All races 3/, female

2006.....	15.3	3.0	*	*	0.2	0.6	1.5	4.1	11.8	34.6	94.1	250.0
2005.....	15.0	3.5	*	*	0.2	0.7	1.5	3.7	11.4	35.5	92.9	247.8
2004.....	14.8	3.2	*	*	0.2	0.7	1.6	4.1	12.2	34.3	90.9	248.1
2003.....	14.9	3.8	*	*	0.2	0.7	1.5	3.9	12.0	35.6	91.3	251.4
2002.....	14.5	3.2	*	*	0.2	0.7	1.5	4.0	11.8	33.7	91.1	246.6
2001.....	14.2	2.9	*	*	0.2	0.6	1.6	3.9	11.9	35.2	87.1	243.3
2000.....	13.6	3.2	*	*	0.2	0.6	1.4	3.5	11.5	34.0	83.5	235.9
1999.....	13.0	4.1	*	*	0.2	0.6	1.3	3.4	10.8	33.0	79.6	227.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

White, both sexes

2006 4/.....	14.8	3.5	*	*	0.2	0.5	1.2	3.7	10.7	33.0	104.3	281.5
2005 4/.....	14.5	3.0	*	*	0.2	0.5	1.1	3.5	10.3	33.3	103.4	280.3
2004 4/.....	14.1	3.1	*	*	0.2	0.4	1.3	3.4	10.3	32.9	101.5	278.2
2003 4/.....	14.3	3.8	*	*	0.2	0.5	1.2	3.4	10.6	33.9	102.4	284.8
2002.....	13.9	3.6	*	*	0.1	0.5	1.1	3.2	9.9	33.4	102.2	278.8
2001.....	13.5	2.4	*	*	0.2	0.4	1.2	3.2	9.7	34.2	96.9	278.2
2000.....	12.9	2.9	*	0.1	0.2	0.4	1.1	3.0	9.7	31.9	94.0	268.7
1999.....	12.3	3.4	*	0.1	0.1	0.4	1.0	2.7	9.1	31.5	90.1	255.7

White, male

2006 4/.....	14.8	4.4	*	*	0.3	0.6	1.4	4.6	12.7	38.8	129.9	373.8
2005 4/.....	14.4	3.1	*	*	0.2	0.5	1.3	4.4	12.3	38.6	129.7	373.7
2004 4/.....	13.9	3.8	*	*	0.2	0.3	1.5	4.0	11.7	37.9	128.6	366.7
2003 4/.....	14.0	4.5	*	*	0.2	0.5	1.4	4.1	11.9	39.0	130.1	383.6
2002.....	13.7	4.4	*	*	0.1	0.5	1.2	3.8	10.7	40.0	130.4	377.4
2001.....	13.3	2.8	*	*	0.2	0.3	1.4	3.8	10.9	40.4	123.5	388.9
2000.....	12.7	3.8	*	*	0.2	0.4	1.3	3.6	11.0	37.7	121.1	376.2
1999.....	12.1	4.1	*	*	*	0.4	1.2	3.1	10.6	36.9	117.2	360.8

White, female

2006 4/.....	14.8	2.6	*	*	0.1	0.4	0.9	2.9	8.8	27.9	86.5	238.1
2005 4/.....	14.6	2.9	*	*	0.2	0.5	0.9	2.6	8.4	28.7	85.2	237.3
2004 4/.....	14.4	2.3	*	*	0.2	0.5	1.1	2.9	9.0	28.6	83.1	238.3
2003 4/.....	14.5	3.0	*	*	0.1	0.5	0.9	2.7	9.3	29.5	83.6	241.0
2002.....	14.1	2.7	*	*	0.1	0.5	1.0	2.7	9.1	27.8	83.3	235.9
2001.....	13.7	2.1	*	*	0.2	0.5	1.0	2.6	8.6	29.0	79.2	231.9
2000.....	13.0	2.0	*	*	0.2	0.4	0.9	2.4	8.4	27.1	76.2	225.0
1999.....	12.4	2.8	*	*	0.2	0.4	0.9	2.3	7.8	26.9	72.3	213.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

Black, both sexes

2006 4/.....	21.2	6.3	*	*	0.4	1.9	5.9	16.2	41.8	102.4	209.7	442.6
2005 4/.....	20.7	8.7	*	*	0.4	2.2	5.5	14.9	44.0	99.4	208.1	424.4
2004 4/.....	20.3	10.1	*	*	0.5	1.9	5.5	16.6	44.5	95.4	202.8	430.9
2003 4/.....	20.6	9.1	*	*	0.5	2.2	6.2	16.4	42.8	103.3	207.0	440.5
2002.....	19.8	8.2	*	*	0.6	2.0	5.3	16.2	41.0	100.1	203.2	441.6
2001.....	19.6	8.1	*	*	0.4	1.8	5.3	15.8	41.6	102.5	202.1	429.6
2000.....	18.9	11.5	*	*	0.4	1.8	5.3	15.8	40.6	99.9	195.1	406.4
1999.....	18.6	9.6	*	*	0.5	1.9	5.4	15.2	37.2	95.4	195.2	433.3

Black, male

2006 4/.....	20.2	6.9	*	*	*	2.1	6.8	20.0	49.9	114.8	250.8	488.8
2005 4/.....	19.5	10.6	*	*	*	2.3	6.1	18.9	54.9	102.6	245.6	482.4
2004 4/.....	19.2	12.2	*	*	*	1.9	6.2	21.0	49.8	108.1	232.7	512.6
2003 4/.....	20.0	9.5	*	*	0.6	2.4	7.1	21.2	51.5	120.5	246.0	518.2
2002.....	19.1	10.2	*	*	*	1.9	6.0	20.1	49.1	115.0	235.2	561.8
2001.....	18.0	8.7	*	*	*	2.2	5.2	19.3	45.4	114.0	233.4	511.7
2000.....	17.7	13.4	*	*	*	1.9	6.5	20.3	44.9	104.8	231.6	490.1
1999.....	17.5	8.1	*	*	*	1.8	6.5	18.9	40.4	102.9	250.0	526.1

Black, female

2006 4/.....	22.2	*	*	*	*	1.7	5.1	12.9	35.3	93.5	186.4	423.7
2005 4/.....	21.7	6.8	*	*	*	2.1	4.9	11.4	35.2	97.1	186.8	401.0
2004 4/.....	21.3	8.0	*	*	*	1.8	4.9	12.9	40.3	86.3	185.8	398.5
2003 4/.....	21.1	8.7	*	*	*	1.9	5.4	12.2	35.8	91.0	185.0	410.4
2002.....	20.5	6.1	*	*	0.7	2.1	4.7	12.8	34.4	89.6	185.0	395.5
2001.....	21.0	7.6	*	*	*	1.4	5.5	12.7	38.5	94.4	184.6	399.1
2000.....	20.0	9.6	*	*	*	1.7	4.3	11.9	37.1	96.5	174.6	375.2
1999.....	19.5	11.1	*	*	*	1.9	4.5	12.0	34.6	90.1	164.5	398.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Acute and rapidly progressive nephritic and nephrotic syndrome (N00-N01,N04)

All races 3/, both sexes

2006.....	0.0	*	*	*	*	*	*	*	*	*	0.3	0.6
2005.....	0.0	*	*	*	*	*	*	*	*	*	0.3	0.8
2004.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.8
2003.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.6
2002.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2001.....	0.1	*	*	*	*	*	*	*	*	0.1	0.4	0.9
2000.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	1.2
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.5	0.8

All races 3/, male

2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.5	*
2003.....	0.0	*	*	*	*	*	*	*	*	*	0.4	*
2002.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.5	1.8
1999.....	0.1	*	*	*	*	*	*	*	*	*	0.7	*

All races 3/, female

2006.....	0.0	*	*	*	*	*	*	*	*	*	0.3	*
2005.....	0.0	*	*	*	*	*	*	*	*	*	0.3	0.7
2004.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.8
2003.....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.6
2002.....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.9
2001.....	0.1	*	*	*	*	*	*	*	*	*	0.4	0.7
2000.....	0.1	*	*	*	*	*	*	*	*	*	0.3	0.9
1999.....	0.1	*	*	*	*	*	*	*	*	0.2	0.4	0.7







Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

All races 3/, both sexes

2006.....	0.6	*	*	*	*	0.1	0.1	0.2	0.5	1.4	4.4	13.4
2005.....	0.3	*	*	*	*	0.1	0.1	0.1	0.3	0.7	1.9	5.4
2004.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.3	2.8
2003.....	0.2	*	*	*	*	*	0.0	0.1	0.2	0.6	1.3	2.8
2002.....	0.2	*	*	*	*	0.1	0.1	0.1	0.2	0.5	1.4	2.8
2001.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.2	2.8
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.6	1.4	2.8
1999.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.5	1.6	3.2

All races 3/, male

2006.....	0.6	*	*	*	*	*	*	0.2	0.6	1.5	5.7	17.2
2005.....	0.3	*	*	*	*	*	0.1	0.2	0.4	0.8	2.4	7.5
2004.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	1.6	4.0
2003.....	0.2	*	*	*	*	*	*	0.2	0.2	0.6	1.6	4.1
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.6	1.8	3.8
2001.....	0.2	*	*	*	*	*	*	0.1	0.2	0.8	1.7	3.7
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.3	0.7	1.7	4.5
1999.....	0.2	*	*	*	*	*	*	0.1	0.3	0.5	2.2	5.0

All races 3/, female

2006.....	0.6	*	*	*	*	*	*	0.2	0.4	1.2	3.5	11.6
2005.....	0.3	*	*	*	*	*	*	*	0.3	0.6	1.6	4.4
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.1	2.2
2003.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	2.2
2002.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.1	2.3
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.2	2.1
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.3	2.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Chronic glomerulonephritis, nephritis and nephropathy not specified as acute or chronic, and renal sclerosis unspecified (N02-N03,N05-N07,N26)

White, both sexes

2006 4/.....	0.6	*	*	*	*	*	*	0.1	0.4	1.1	4.0	12.5
2005 4/.....	0.3	*	*	*	*	*	*	0.1	0.3	0.6	1.9	5.2
2004 4/.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.2	2.7
2003 4/.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.2	2.6
2002.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.6
2001.....	0.2	*	*	*	*	*	*	0.1	0.1	0.6	1.1	2.9
2000.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	2.4
1999.....	0.2	*	*	*	*	*	*	0.1	0.2	0.5	1.6	2.9

White, male

2006 4/.....	0.6	*	*	*	*	*	*	0.1	0.4	1.3	5.3	15.9
2005 4/.....	0.3	*	*	*	*	*	*	0.1	0.3	0.7	2.3	7.3
2004 4/.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.6	3.7
2003 4/.....	0.2	*	*	*	*	*	*	0.1	*	0.5	1.4	3.8
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.7	3.8
2001.....	0.2	*	*	*	*	*	*	*	*	0.7	1.7	4.0
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.7	4.0
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	2.2	4.6

White, female

2006 4/.....	0.6	*	*	*	*	*	*	0.1	0.3	1.0	3.2	10.9
2005 4/.....	0.3	*	*	*	*	*	*	*	0.2	0.5	1.6	4.2
2004 4/.....	0.2	*	*	*	*	*	*	*	*	0.4	1.0	2.3
2003 4/.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.0	2.1
2002.....	0.2	*	*	*	*	*	*	*	*	0.4	1.0	2.0
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	2.4
2000.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.1	1.8
1999.....	0.2	*	*	*	*	*	*	*	*	0.5	1.2	2.2



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Renal failure (N17-N19)

All races 3/, both sexes

2006.....	14.5	3.7	*	*	0.2	0.6	1.7	5.0	13.3	37.9	106.7	276.4
2005.....	14.5	3.7	*	*	0.2	0.7	1.6	4.6	13.3	38.4	108.0	282.1
2004.....	14.2	4.0	*	*	0.2	0.6	1.7	4.8	13.4	37.9	106.6	282.9
2003.....	14.4	4.2	*	0.1	0.2	0.7	1.7	4.8	13.4	39.3	107.8	289.7
2002.....	13.9	4.1	*	*	0.2	0.6	1.6	4.6	12.7	38.6	107.2	284.8
2001.....	13.6	3.1	*	*	0.2	0.5	1.6	4.5	12.7	39.4	102.6	284.0
2000.....	13.0	4.0	*	*	0.2	0.6	1.6	4.2	12.5	37.2	98.9	273.7
1999.....	12.4	4.3	*	0.1	0.2	0.6	1.5	3.9	11.7	36.4	95.3	264.8

All races 3/, male

2006.....	14.3	4.4	*	*	0.2	0.7	2.0	6.0	15.4	43.4	130.7	358.8
2005.....	14.2	4.1	*	*	0.2	0.7	1.8	5.7	15.6	42.8	133.1	368.0
2004.....	13.8	5.1	*	*	0.2	0.5	2.0	5.7	14.8	42.9	132.1	367.4
2003.....	14.1	4.8	*	*	0.2	0.7	2.0	5.7	15.2	44.5	134.8	383.1
2002.....	13.7	5.1	*	*	0.2	0.6	1.8	5.4	14.0	45.0	134.0	380.7
2001.....	13.2	3.5	*	*	0.2	0.5	1.8	5.3	13.8	45.2	128.0	388.7
2000.....	12.6	5.1	*	*	0.2	0.6	1.8	5.1	13.8	42.1	125.1	374.2
1999.....	12.1	4.5	*	*	0.1	0.5	1.7	4.5	13.1	41.5	122.3	364.6

All races 3/, female

2006.....	14.6	3.0	*	*	0.2	0.6	1.4	3.9	11.4	33.2	90.2	237.8
2005.....	14.7	3.2	*	*	0.2	0.6	1.4	3.6	11.0	34.8	91.0	242.6
2004.....	14.6	3.0	*	*	0.2	0.6	1.5	3.9	12.0	33.7	89.5	244.9
2003.....	14.6	3.7	*	*	0.2	0.6	1.5	3.8	11.8	34.9	89.7	248.4
2002.....	14.2	3.0	*	*	0.2	0.7	1.4	3.8	11.6	33.2	89.5	243.2
2001.....	14.0	2.7	*	*	0.2	0.5	1.5	3.7	11.8	34.5	85.9	240.1
2000.....	13.3	2.8	*	*	0.2	0.5	1.3	3.4	11.2	33.3	81.8	232.8
1999.....	12.7	4.1	*	*	0.2	0.6	1.3	3.3	10.4	32.2	77.8	224.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Renal failure (N17-N19)

White, both sexes

2006 4/.....	14.1	3.3	*	*	0.2	0.5	1.1	3.6	10.3	31.7	99.9	268.2
2005 4/.....	14.2	2.8	*	*	0.2	0.4	1.1	3.4	10.0	32.5	101.1	274.3
2004 4/.....	13.9	2.9	*	*	0.1	0.4	1.2	3.3	10.0	32.3	99.8	274.7
2003 4/.....	14.0	3.5	*	*	0.2	0.5	1.1	3.3	10.3	33.2	100.7	281.5
2002.....	13.6	3.4	*	*	0.1	0.4	1.1	3.1	9.6	32.7	100.5	275.3
2001.....	13.2	2.2	*	*	0.2	0.4	1.1	3.1	9.5	33.4	95.3	274.6
2000.....	12.6	2.7	*	*	0.2	0.4	1.1	2.9	9.4	31.2	92.2	265.0
1999.....	12.0	3.3	*	*	0.1	0.4	1.0	2.5	8.9	30.8	87.9	251.9

White, male

2006 4/.....	14.1	4.0	*	*	0.2	0.5	1.3	4.5	12.2	37.4	124.3	356.8
2005 4/.....	14.1	3.1	*	*	0.2	0.4	1.2	4.3	11.9	37.7	127.0	365.4
2004 4/.....	13.6	3.6	*	*	0.2	0.3	1.4	3.9	11.4	37.3	126.5	362.3
2003 4/.....	13.8	4.1	*	*	0.2	0.5	1.3	3.9	11.7	38.3	128.3	379.3
2002.....	13.4	4.2	*	*	*	0.4	1.2	3.6	10.4	39.2	128.3	372.7
2001.....	13.0	2.6	*	*	0.2	0.3	1.3	3.7	10.7	39.5	121.5	383.9
2000.....	12.4	3.6	*	*	0.2	0.4	1.2	3.4	10.6	36.9	118.9	370.4
1999.....	11.8	3.9	*	*	*	0.4	1.1	3.0	10.3	36.2	114.4	354.9

White, female

2006 4/.....	14.2	2.6	*	*	0.1	0.4	0.9	2.7	8.5	26.9	82.9	226.6
2005 4/.....	14.2	2.6	*	*	0.1	0.4	0.9	2.5	8.1	28.1	83.3	232.4
2004 4/.....	14.1	2.1	*	*	0.1	0.4	1.0	2.8	8.7	28.1	81.6	235.2
2003 4/.....	14.2	2.8	*	*	*	0.5	0.9	2.7	9.0	28.8	82.1	238.2
2002.....	13.8	2.6	*	*	*	0.5	1.0	2.6	8.9	27.2	81.8	232.9
2001.....	13.4	1.8	*	*	0.2	0.4	0.9	2.5	8.4	28.3	78.0	228.8
2000.....	12.8	1.7	*	*	0.2	0.4	0.9	2.3	8.2	26.5	74.6	222.1
1999.....	12.2	2.8	*	*	0.1	0.4	0.8	2.2	7.6	26.2	70.6	210.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Renal failure (N17-N19)

Black, both sexes

2006 4/.....	20.3	6.2	*	*	0.4	1.7	5.7	15.5	40.2	98.3	201.2	418.7
2005 4/.....	20.2	8.4	*	*	0.4	2.0	5.3	14.5	43.1	97.7	204.2	413.7
2004 4/.....	20.0	9.8	*	*	0.4	1.7	5.3	16.3	44.0	94.2	200.3	426.4
2003 4/.....	20.3	8.9	*	*	0.5	2.1	6.1	16.1	42.5	102.0	204.4	436.4
2002.....	19.5	7.7	*	*	0.5	1.9	5.3	15.7	40.6	99.6	200.2	434.2
2001.....	19.3	8.1	*	*	0.4	1.7	5.2	15.4	41.1	101.3	200.0	425.6
2000.....	18.5	11.1	*	*	0.4	1.7	5.1	15.2	40.1	98.6	192.4	400.4
1999.....	18.2	9.6	*	*	0.4	1.8	5.2	14.8	36.4	94.1	191.8	425.3

Black, male

2006 4/.....	19.2	6.6	*	*	*	1.8	6.7	19.1	47.6	110.4	239.7	454.9
2005 4/.....	19.1	10.6	*	*	*	2.0	5.8	18.5	53.7	101.2	239.8	469.4
2004 4/.....	18.9	11.9	*	*	*	1.9	6.0	20.6	49.0	106.8	230.0	506.7
2003 4/.....	19.7	9.2	*	*	*	2.4	7.0	20.8	51.2	118.8	241.7	513.0
2002.....	18.8	9.6	*	*	*	1.7	5.9	19.6	48.8	114.2	231.5	555.3
2001.....	17.7	8.7	*	*	*	2.1	5.0	18.8	44.7	112.4	231.6	508.3
2000.....	17.3	13.4	*	*	*	1.8	6.3	19.5	44.2	103.3	229.2	480.9
1999.....	17.1	8.1	*	*	*	1.7	6.2	18.5	39.7	101.5	245.7	513.2

Black, female

2006 4/.....	21.3	*	*	*	*	1.6	4.8	12.3	34.3	89.6	179.3	403.9
2005 4/.....	21.2	6.2	*	*	*	2.0	4.8	11.0	34.5	95.1	184.0	391.2
2004 4/.....	21.0	7.7	*	*	*	1.6	4.7	12.6	39.9	85.3	183.5	394.6
2003 4/.....	20.9	8.7	*	*	*	1.8	5.3	12.1	35.5	90.0	183.2	406.7
2002.....	20.2	*	*	*	*	2.0	4.7	12.3	34.1	89.2	182.6	387.7
2001.....	20.7	7.6	*	*	*	1.3	5.4	12.6	38.2	93.5	182.3	394.9
2000.....	19.6	8.6	*	*	*	1.6	4.0	11.5	36.7	95.3	171.7	370.4
1999.....	19.1	11.1	*	*	*	1.9	4.3	11.7	33.8	88.9	161.6	392.5









Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Infections of kidney (N10-N12,N13.6,N15.1)

All races 3/, both sexes

2006.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	4.7
2005.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.8	5.2
2004.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	5.2
2003.....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.6	1.9	5.9
2002.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.3
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	2.0	6.0
2000.....	0.3	*	*	*	*	*	0.0	0.1	0.2	0.6	2.0	7.5
1999.....	0.3	*	*	*	*	*	0.1	0.2	0.2	0.7	2.0	7.5

All races 3/, male

2006.....	0.1	*	*	*	*	*	*	*	0.1	0.5	0.8	3.7
2005.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.6	3.6
2004.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.5	4.5
2003.....	0.2	*	*	*	*	*	*	0.1	0.2	0.4	1.4	5.4
2002.....	0.2	*	*	*	*	*	*	*	0.2	0.6	1.4	5.3
2001.....	0.2	*	*	*	*	*	*	*	*	0.5	1.5	4.3
2000.....	0.2	*	*	*	*	*	*	*	*	0.3	1.5	5.6
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.7	6.8

All races 3/, female

2006.....	0.3	*	*	*	*	*	*	0.2	0.2	0.5	1.6	5.2
2005.....	0.4	*	*	*	*	*	*	0.2	0.2	0.6	2.0	5.9
2004.....	0.4	*	*	*	*	*	0.1	0.2	0.2	0.7	2.1	5.5
2003.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.7	2.2	6.1
2002.....	0.4	*	*	*	*	*	*	0.1	0.3	0.6	2.2	6.7
2001.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.5	2.3	6.7
2000.....	0.4	*	*	*	*	*	*	0.1	0.3	0.8	2.3	8.3
1999.....	0.4	*	*	*	*	*	0.1	0.2	0.3	0.9	2.1	7.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Infections of kidney (N10-N12,N13.6,N15.1)

White, both sexes

2006 4/.....	0.2	*	*	*	*	*	0.1	0.1	0.2	0.5	1.3	4.9
2005 4/.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.9	5.2
2004 4/.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.8	5.2
2003 4/.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.0
2002.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.6	1.9	6.4
2001.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.5	1.9	6.1
2000.....	0.3	*	*	*	*	*	*	0.1	0.2	0.5	1.9	7.6
1999.....	0.3	*	*	*	*	*	0.1	0.1	0.2	0.7	2.0	7.7

White, male

2006 4/.....	0.2	*	*	*	*	*	*	*	*	0.5	0.8	3.7
2005 4/.....	0.2	*	*	*	*	*	*	*	*	0.4	1.6	3.5
2004 4/.....	0.2	*	*	*	*	*	*	*	0.2	0.4	1.5	4.7
2003 4/.....	0.2	*	*	*	*	*	*	0.1	*	0.4	1.3	5.8
2002.....	0.2	*	*	*	*	*	*	*	*	0.5	1.4	5.4
2001.....	0.2	*	*	*	*	*	*	*	*	0.4	1.4	4.3
2000.....	0.2	*	*	*	*	*	*	*	*	*	1.5	5.9
1999.....	0.2	*	*	*	*	*	*	*	0.2	0.5	1.8	7.2

White, female

2006 4/.....	0.3	*	*	*	*	*	*	0.2	0.2	0.5	1.6	5.4
2005 4/.....	0.4	*	*	*	*	*	*	0.1	0.2	0.6	2.0	6.0
2004 4/.....	0.4	*	*	*	*	*	0.1	0.1	0.2	0.8	2.0	5.5
2003 4/.....	0.4	*	*	*	*	*	*	0.2	0.3	0.8	2.2	6.0
2002.....	0.4	*	*	*	*	*	*	*	0.2	0.7	2.3	6.8
2001.....	0.4	*	*	*	*	*	*	0.2	0.3	0.5	2.3	6.9
2000.....	0.4	*	*	*	*	*	*	*	0.3	0.7	2.2	8.3
1999.....	0.4	*	*	*	*	*	*	0.2	0.2	0.9	2.2	7.9

















Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes

2006.....	0.3	...	...	*	0.4	0.8	0.4	0.1	*	*	*	*
2005.....	0.3	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2004.....	0.2	...	...	*	0.4	0.7	0.4	0.2	*	*	*	*
2003.....	0.2	...	...	*	0.3	0.6	0.4	0.1	*	*	*	*
2002.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001.....	0.1	...	...	*	0.3	0.5	0.2	*	*	*	*	*
2000.....	0.1	...	...	*	0.2	0.5	0.3	*	*	*	*	*
1999.....	0.1	...	...	*	0.3	0.5	0.3	*	*	*	*	*

All races 3/, male

2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All races 3/, female

2006.....	0.5	...	...	*	0.9	1.6	0.9	0.3	*	*	*	*
2005.....	0.5	...	...	*	0.9	1.6	0.9	0.3	*	*	*	*
2004.....	0.5	...	...	*	0.8	1.5	0.7	0.4	*	*	*	*
2003.....	0.4	...	...	*	0.6	1.2	0.7	0.1	*	*	*	*
2002.....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
2001.....	0.3	...	...	*	0.6	1.0	0.5	*	*	*	*	*
2000.....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*
1999.....	0.3	...	...	*	0.5	0.9	0.5	*	*	*	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Pregnancy, childbirth and the puerperium (000-099)

White, both sexes

2006 4/.....	0.2	...	...	*	0.3	0.6	0.3	0.1	*	*	*	*
2005 4/.....	0.2	...	...	*	0.3	0.6	0.4	0.1	*	*	*	*
2004 4/.....	0.2	...	...	*	0.3	0.5	0.3	0.2	*	*	*	*
2003 4/.....	0.1	...	...	*	0.2	0.4	0.3	0.1	*	*	*	*
2002.....	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*
2001.....	0.1	...	...	*	0.2	0.4	0.1	*	*	*	*	*
2000.....	0.1	...	...	*	0.2	0.3	0.2	*	*	*	*	*
1999.....	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*

White, male

2006 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

White, female

2006 4/.....	0.4	...	...	*	0.7	1.2	0.7	0.2	*	*	*	*
2005 4/.....	0.4	...	...	*	0.7	1.2	0.7	0.2	*	*	*	*
2004 4/.....	0.3	...	...	*	0.6	1.1	0.6	0.3	*	*	*	*
2003 4/.....	0.3	...	...	*	0.4	0.9	0.5	0.1	*	*	*	*
2002.....	0.2	...	...	*	0.3	0.6	0.3	*	*	*	*	*
2001.....	0.2	...	...	*	0.4	0.7	0.3	*	*	*	*	*
2000.....	0.2	...	...	*	0.4	0.7	0.4	*	*	*	*	*
1999.....	0.2	...	...	*	0.3	0.6	0.4	*	*	*	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Pregnancy, childbirth and the puerperium (000-099)

Black, both sexes

2006 4/.....	0.7	...	...	*	0.9	2.1	1.1	0.4	*	*	*	*
2005 4/.....	0.7	...	...	*	1.1	2.1	1.1	*	*	*	*	*
2004 4/.....	0.7	...	...	*	1.1	2.0	0.8	0.5	*	*	*	*
2003 4/.....	0.5	...	...	*	0.7	1.5	1.0	*	*	*	*	*
2002.....	0.4	...	...	*	0.7	1.2	0.7	*	*	*	*	*
2001.....	0.4	...	...	*	0.7	1.2	0.7	*	*	*	*	*
2000.....	0.4	...	...	*	0.6	1.2	0.7	*	*	*	*	*
1999.....	0.4	...	...	*	0.8	1.3	0.7	*	*	*	*	*

Black, male

2006 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Black, female

2006 4/.....	1.3	...	...	*	1.8	4.0	2.0	0.8	*	*	*	*
2005 4/.....	1.3	...	...	*	2.1	4.0	2.1	*	*	*	*	*
2004 4/.....	1.2	...	...	*	2.3	3.8	1.5	0.9	*	*	*	*
2003 4/.....	1.0	...	...	*	1.4	2.9	1.9	*	*	*	*	*
2002.....	0.8	...	...	*	1.5	2.3	1.4	*	*	*	*	*
2001.....	0.8	...	...	*	1.5	2.3	1.4	*	*	*	*	*
2000.....	0.7	...	...	*	1.2	2.3	1.2	*	*	*	*	*
1999.....	0.9	...	...	*	1.6	2.5	1.4	*	*	*	*	*









Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

All races 3/, both sexes

2006.....	0.2	...	...	*	0.4	0.8	0.4	0.1	*	*	*	*
2005.....	0.2	...	...	*	0.4	0.7	0.4	0.1	*	*	*	*
2004.....	0.2	...	...	*	0.4	0.7	0.3	0.2	*	*	*	*
2003.....	0.2	...	...	*	0.3	0.5	0.4	0.1	*	*	*	*
2002.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2001.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
2000.....	0.1	...	...	*	0.2	0.4	0.2	*	*	*	*	*
1999.....	0.1	...	...	*	0.2	0.4	0.3	*	*	*	*	*

All races 3/, male

2006.....	...	...	...	...	...	...	...	...	...	...	...	...
2005.....	...	...	...	...	...	...	...	...	...	...	...	...
2004.....	...	...	...	...	...	...	...	...	...	...	...	...
2003.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

All races 3/, female

2006.....	0.5	...	...	*	0.8	1.6	0.9	0.3	*	*	*	*
2005.....	0.5	...	...	*	0.9	1.5	0.9	0.2	*	*	*	*
2004.....	0.4	...	...	*	0.8	1.4	0.7	0.4	*	*	*	*
2003.....	0.3	...	...	*	0.5	1.1	0.7	0.1	*	*	*	*
2002.....	0.2	...	...	*	0.4	0.8	0.4	*	*	*	*	*
2001.....	0.3	...	...	*	0.5	0.9	0.4	*	*	*	*	*
2000.....	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*
1999.....	0.3	...	...	*	0.5	0.8	0.5	*	*	*	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

White, both sexes

2006 4/.....	0.2	...	...	*	0.3	0.6	0.3	0.1	*	*	*	*
2005 4/.....	0.2	...	...	*	0.3	0.6	0.3	0.1	*	*	*	*
2004 4/.....	0.2	...	...	*	0.3	0.5	0.3	0.2	*	*	*	*
2003 4/.....	0.1	...	...	*	0.2	0.4	0.3	0.1	*	*	*	*
2002.....	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*
2001.....	0.1	...	...	*	0.2	0.3	0.1	*	*	*	*	*
2000.....	0.1	...	...	*	0.2	0.3	0.2	*	*	*	*	*
1999.....	0.1	...	...	*	0.1	0.3	0.2	*	*	*	*	*

White, male

2006 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

White, female

2006 4/.....	0.4	...	...	*	0.7	1.1	0.7	0.2	*	*	*	*
2005 4/.....	0.4	...	...	*	0.7	1.1	0.7	0.2	*	*	*	*
2004 4/.....	0.3	...	...	*	0.5	1.1	0.6	0.3	*	*	*	*
2003 4/.....	0.3	...	...	*	0.4	0.8	0.5	0.1	*	*	*	*
2002.....	0.2	...	...	*	0.3	0.6	0.3	*	*	*	*	*
2001.....	0.2	...	...	*	0.4	0.7	0.3	*	*	*	*	*
2000.....	0.2	...	...	*	0.4	0.6	0.4	*	*	*	*	*
1999.....	0.2	...	...	*	0.3	0.6	0.4	*	*	*	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other complications of pregnancy, childbirth and the puerperium (010-099)

Black, both sexes

2006 4/.....	0.6	...	...	*	0.8	2.0	1.0	0.4	*	*	*	*
2005 4/.....	0.7	...	...	*	1.0	1.9	1.1	*	*	*	*	*
2004 4/.....	0.6	...	...	*	1.0	1.9	0.7	0.5	*	*	*	*
2003 4/.....	0.5	...	...	*	0.6	1.4	0.9	*	*	*	*	*
2002.....	0.4	...	...	*	0.7	1.1	0.6	*	*	*	*	*
2001.....	0.4	...	...	*	0.6	1.1	0.7	*	*	*	*	*
2000.....	0.3	...	...	*	0.5	1.0	0.6	*	*	*	*	*
1999.....	0.4	...	...	*	0.7	1.1	0.7	*	*	*	*	*

Black, male

2006 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2005 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2004 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2003 4/.....	...	...	...	...	...	...	...	...	...	...	...	...
2002.....	...	...	...	...	...	...	...	...	...	...	...	...
2001.....	...	...	...	...	...	...	...	...	...	...	...	...
2000.....	...	...	...	...	...	...	...	...	...	...	...	...
1999.....	...	...	...	...	...	...	...	...	...	...	...	...

Black, female

2006 4/.....	1.2	...	...	*	1.7	3.8	1.9	0.8	*	*	*	*
2005 4/.....	1.2	...	...	*	2.1	3.7	2.0	*	*	*	*	*
2004 4/.....	1.1	...	...	*	2.1	3.6	1.3	0.9	*	*	*	*
2003 4/.....	0.9	...	...	*	1.1	2.7	1.8	*	*	*	*	*
2002.....	0.7	...	...	*	1.4	2.2	1.2	*	*	*	*	*
2001.....	0.7	...	...	*	1.3	2.1	1.3	*	*	*	*	*
2000.....	0.6	...	...	*	1.0	2.0	1.1	*	*	*	*	*
1999.....	0.8	...	...	*	1.5	2.2	1.3	*	*	*	*	*









Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes

2006.....	3.5	140.9	3.2	0.9	1.1	1.1	1.1	1.5	2.3	1.9	3.0	5.6
2005.....	3.5	135.2	3.2	1.0	1.2	1.1	1.2	1.6	2.4	1.8	3.1	6.4
2004.....	3.6	137.9	3.6	1.0	1.2	1.0	1.1	1.7	2.2	2.3	3.6	6.1
2003.....	3.6	140.4	3.4	0.9	1.1	1.1	1.3	1.6	2.4	2.3	3.6	6.8
2002.....	3.7	139.4	3.4	1.0	1.2	1.2	1.3	1.8	2.4	2.4	3.8	6.6
2001.....	3.7	136.7	3.6	0.9	1.3	1.2	1.3	1.6	2.3	2.4	3.7	7.2
2000.....	3.8	150.9	3.2	1.0	1.1	1.2	1.3	1.6	2.3	2.5	3.9	7.5
1999.....	3.7	144.2	3.6	1.0	1.1	1.2	1.2	1.8	2.3	2.5	4.0	7.8

All races 3/, male

2006.....	3.8	146.6	2.9	0.9	1.3	1.3	1.2	1.8	2.6	2.1	3.6	6.8
2005.....	3.7	141.1	3.2	0.9	1.5	1.2	1.2	1.7	2.5	1.8	3.6	6.9
2004.....	3.8	139.0	3.9	1.0	1.3	1.2	1.3	1.8	2.3	2.5	3.7	6.8
2003.....	3.8	147.6	3.4	0.9	1.3	1.2	1.4	1.7	2.5	2.4	3.5	6.7
2002.....	3.9	142.6	3.5	1.0	1.4	1.4	1.3	1.9	2.5	2.6	4.0	7.0
2001.....	3.9	143.1	3.6	1.0	1.5	1.2	1.4	1.7	2.3	2.7	4.1	7.7
2000.....	4.0	155.4	3.3	1.0	1.3	1.4	1.4	1.9	2.3	2.7	4.0	10.6
1999.....	3.9	147.9	4.0	1.0	1.3	1.2	1.3	1.9	2.5	2.5	4.2	8.5

All races 3/, female

2006.....	3.2	134.9	3.4	0.8	0.9	0.9	0.9	1.3	2.0	1.8	2.6	5.1
2005.....	3.3	129.0	3.3	1.0	0.9	1.0	1.1	1.6	2.2	1.8	2.8	6.2
2004.....	3.4	136.8	3.2	0.9	1.0	0.9	1.0	1.5	2.1	2.2	3.6	5.8
2003.....	3.4	132.8	3.5	0.9	0.9	1.0	1.2	1.5	2.3	2.2	3.7	6.8
2002.....	3.5	136.1	3.3	1.0	1.1	1.0	1.2	1.6	2.3	2.3	3.8	6.5
2001.....	3.4	129.9	3.7	0.8	1.0	1.1	1.2	1.6	2.3	2.2	3.5	7.0
2000.....	3.5	146.2	3.1	1.0	0.9	1.0	1.1	1.4	2.4	2.4	3.9	6.3
1999.....	3.5	140.3	3.2	1.1	0.9	1.1	1.1	1.6	2.2	2.6	3.9	7.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

White, both sexes

2006 4/.....	3.4	137.2	3.0	0.8	1.0	1.1	1.1	1.6	2.4	2.0	3.1	5.7
2005 4/.....	3.4	131.0	3.1	0.9	1.2	1.1	1.2	1.7	2.5	2.0	3.3	6.5
2004 4/.....	3.5	134.5	3.2	0.9	1.1	1.1	1.1	1.7	2.3	2.5	3.7	6.3
2003 4/.....	3.5	137.6	3.2	0.9	1.1	1.1	1.3	1.7	2.5	2.3	3.7	6.9
2002.....	3.6	138.3	3.3	1.0	1.2	1.2	1.3	1.8	2.4	2.5	3.9	6.9
2001.....	3.6	135.0	3.2	0.9	1.2	1.2	1.3	1.7	2.4	2.5	3.8	7.4
2000.....	3.6	147.1	2.9	0.9	1.1	1.2	1.2	1.7	2.4	2.5	4.1	7.8
1999.....	3.6	141.3	3.2	0.9	1.1	1.2	1.2	1.8	2.4	2.6	4.0	8.0

White, male

2006 4/.....	3.7	142.3	2.9	0.9	1.2	1.3	1.2	1.8	2.7	2.1	3.7	6.8
2005 4/.....	3.6	136.9	3.1	0.9	1.5	1.2	1.2	1.8	2.6	2.0	3.8	6.9
2004 4/.....	3.6	134.3	3.5	1.0	1.2	1.2	1.3	1.9	2.4	2.6	3.7	7.2
2003 4/.....	3.7	143.2	3.1	0.9	1.2	1.2	1.4	1.7	2.6	2.3	3.4	6.9
2002.....	3.8	143.3	3.4	1.0	1.3	1.4	1.3	1.9	2.5	2.6	3.9	7.3
2001.....	3.8	143.0	3.2	1.0	1.4	1.3	1.4	1.7	2.4	2.9	4.2	7.5
2000.....	3.9	151.9	2.9	0.9	1.3	1.4	1.4	1.9	2.4	2.6	4.3	11.1
1999.....	3.8	146.5	3.5	0.9	1.3	1.2	1.3	1.9	2.6	2.5	4.1	8.6

White, female

2006 4/.....	3.1	131.8	3.0	0.7	0.9	0.9	0.9	1.4	2.1	1.9	2.7	5.2
2005 4/.....	3.2	124.8	3.0	0.9	0.8	1.0	1.2	1.7	2.4	2.0	2.9	6.3
2004 4/.....	3.4	134.7	2.9	0.8	1.0	1.0	1.0	1.6	2.2	2.3	3.6	6.0
2003 4/.....	3.4	131.8	3.2	0.9	0.9	1.0	1.2	1.6	2.4	2.2	3.9	6.9
2002.....	3.5	133.1	3.1	1.0	1.1	1.0	1.2	1.7	2.3	2.4	3.9	6.7
2001.....	3.4	126.7	3.3	0.7	1.0	1.1	1.2	1.6	2.5	2.2	3.6	7.3
2000.....	3.5	142.0	2.9	0.9	0.9	1.0	1.1	1.5	2.4	2.5	4.0	6.5
1999.....	3.5	135.7	2.9	1.0	0.9	1.2	1.2	1.7	2.3	2.6	4.0	7.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

Black, both sexes

2006 4/.....	4.6	170.8	4.1	1.0	1.3	1.3	1.3	1.8	2.0	1.6	2.9	5.4
2005 4/.....	4.4	162.9	4.0	1.3	1.5	1.4	1.3	1.4	1.9	*	2.1	5.9
2004 4/.....	4.5	162.9	5.1	1.3	1.7	1.1	1.2	1.7	1.6	1.6	3.9	*
2003 4/.....	4.5	157.5	4.7	1.1	1.5	1.4	1.2	1.6	2.2	2.3	3.1	5.8
2002.....	4.7	155.2	4.5	1.2	1.6	1.4	1.6	1.9	2.6	2.4	3.3	*
2001.....	4.6	158.4	5.3	1.1	1.9	1.3	1.6	1.7	1.9	2.2	3.0	6.1
2000.....	4.9	179.2	5.0	1.3	1.5	1.4	1.6	1.6	2.4	2.6	2.7	*
1999.....	4.8	169.3	5.5	1.6	1.4	1.1	1.3	1.9	1.9	2.8	4.2	7.7

Black, male

2006 4/.....	5.0	176.8	3.1	0.9	1.6	1.6	1.4	2.3	2.3	*	*	*
2005 4/.....	4.8	166.5	3.6	1.2	1.9	1.6	1.6	1.4	2.3	*	*	*
2004 4/.....	5.1	169.9	5.8	1.1	2.2	1.4	1.6	1.8	1.9	*	*	*
2003 4/.....	5.1	170.7	4.8	1.1	1.7	1.5	1.2	1.9	2.7	3.5	*	*
2002.....	5.1	151.7	4.6	1.4	1.9	1.7	1.7	2.3	3.4	3.0	*	*
2001.....	5.0	160.8	5.5	1.0	2.4	1.2	1.7	1.7	2.0	*	*	*
2000.....	5.4	181.4	5.5	1.5	2.0	1.8	1.9	2.1	*	3.8	*	*
1999.....	5.2	172.7	6.2	1.6	1.8	1.2	1.6	2.3	2.1	*	*	*

Black, female

2006 4/.....	4.2	164.5	5.0	1.2	1.1	1.0	1.2	1.5	1.7	*	*	*
2005 4/.....	4.1	159.0	4.5	1.4	1.1	1.3	1.0	1.4	1.7	*	*	*
2004 4/.....	4.0	155.6	4.3	1.4	1.2	0.8	0.9	1.6	1.4	*	3.3	*
2003 4/.....	3.9	143.8	4.6	1.0	1.2	1.3	1.2	1.4	1.8	*	*	*
2002.....	4.3	158.9	4.4	1.0	1.3	1.1	1.5	1.5	2.0	*	*	*
2001.....	4.3	155.9	5.1	1.2	1.4	1.3	1.4	1.7	1.8	2.5	*	*
2000.....	4.4	176.9	4.6	1.2	1.1	1.1	1.4	1.1	3.0	*	*	*
1999.....	4.3	165.8	4.8	1.6	1.1	1.0	1.2	1.5	1.8	3.1	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes

2006.....	10.6	83.8	1.5	0.3	1.6	2.5	3.7	5.2	6.5	12.2	41.7	236.1
2005.....	10.8	87.4	1.4	0.4	1.8	2.9	4.7	6.7	7.2	12.8	39.0	224.4
2004.....	10.3	82.4	1.2	0.3	1.6	2.7	4.3	5.8	7.0	12.6	39.8	221.5
2003.....	10.8	82.9	1.6	0.4	2.0	3.3	5.4	7.0	7.4	13.8	38.9	224.1
2002.....	10.4	85.7	1.6	0.3	1.8	3.3	5.3	6.7	7.5	13.1	38.8	208.0
2001.....	11.4	86.5	1.8	0.4	2.4	3.9	6.9	8.9	9.1	15.1	39.8	206.4
2000.....	11.3	95.6	1.6	0.4	2.2	3.9	6.6	8.3	9.3	16.5	40.7	208.1
1999.....	9.6	93.2	1.5	0.3	1.6	2.7	4.6	5.9	7.8	14.6	36.1	187.7

All races 3/, male

2006.....	9.5	95.2	1.6	0.4	2.0	3.5	4.8	6.7	8.3	14.8	44.2	191.1
2005.....	10.0	101.8	1.6	0.5	2.5	4.0	5.8	8.7	9.3	15.5	40.7	183.9
2004.....	9.5	96.9	1.2	0.3	2.1	3.6	5.6	7.6	9.1	15.0	41.8	183.6
2003.....	10.2	96.5	1.8	0.4	2.7	4.5	6.9	9.2	9.6	17.1	41.9	177.9
2002.....	10.1	98.5	1.8	0.4	2.4	4.5	6.9	8.8	10.0	15.9	43.8	175.5
2001.....	11.3	98.9	2.1	0.5	3.2	5.2	9.0	11.9	12.0	18.0	43.8	175.0
2000.....	11.5	110.6	1.7	0.5	2.9	5.3	8.9	11.5	12.3	20.6	46.0	182.7
1999.....	9.4	106.6	1.7	0.4	2.1	3.6	6.0	8.1	10.4	18.2	40.1	166.3

All races 3/, female

2006.....	11.7	71.9	1.3	0.3	1.1	1.5	2.7	3.8	4.8	10.0	39.9	257.2
2005.....	11.5	72.3	1.3	0.3	1.1	1.9	3.6	4.7	5.3	10.6	37.8	243.0
2004.....	11.0	67.3	1.1	0.3	1.0	1.8	3.0	4.0	5.1	10.6	38.4	238.5
2003.....	11.4	68.6	1.5	0.4	1.4	2.1	3.8	5.0	5.3	10.9	36.9	244.6
2002.....	10.7	72.3	1.4	0.3	1.2	2.1	3.8	4.7	5.3	10.8	35.4	222.1
2001.....	11.4	73.6	1.5	0.4	1.5	2.5	4.9	6.1	6.3	12.7	37.2	219.5
2000.....	11.2	79.9	1.5	0.4	1.5	2.5	4.4	5.3	6.5	13.2	37.2	218.5
1999.....	9.7	79.2	1.3	0.3	1.0	1.9	3.2	3.8	5.5	11.7	33.5	196.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

White, both sexes												
2006 4/.....	11.0	70.2	1.2	0.3	1.5	2.5	3.6	4.9	6.2	11.8	41.4	241.1
2005 4/.....	11.1	72.9	1.2	0.3	1.8	2.9	4.5	6.2	6.7	12.2	38.8	228.9
2004 4/.....	10.5	65.8	1.0	0.2	1.6	2.7	4.2	5.5	6.5	12.1	39.5	224.4
2003 4/.....	11.1	68.1	1.4	0.4	2.0	3.2	5.1	6.4	6.9	13.2	39.0	228.3
2002.....	10.5	71.0	1.3	0.3	1.8	3.2	5.1	5.9	7.0	12.7	38.6	211.6
2001.....	11.6	70.9	1.5	0.4	2.4	3.8	6.8	8.6	8.5	14.4	39.6	210.0
2000.....	11.4	77.2	1.5	0.4	2.1	3.7	6.5	7.8	8.6	15.6	40.2	212.2
1999.....	9.4	74.7	1.3	0.3	1.4	2.5	4.1	5.1	7.1	13.9	35.1	190.2
White, male												
2006 4/.....	9.5	81.8	1.3	0.3	2.0	3.4	4.6	6.2	7.9	14.0	43.5	193.8
2005 4/.....	10.0	87.7	1.4	0.4	2.5	3.9	5.6	8.2	8.5	14.6	40.0	185.6
2004 4/.....	9.4	76.9	1.1	0.3	2.1	3.6	5.5	7.1	8.3	14.4	41.4	186.0
2003 4/.....	10.1	82.4	1.4	0.4	2.7	4.4	6.6	8.4	8.8	16.3	41.7	180.0
2002.....	9.8	82.5	1.4	0.3	2.4	4.3	6.5	7.7	9.2	15.3	42.9	178.2
2001.....	11.2	82.0	1.7	0.4	3.2	5.2	8.9	11.4	11.0	17.2	43.9	177.0
2000.....	11.3	89.4	1.7	0.4	2.9	5.1	8.9	10.7	11.2	19.2	45.5	186.8
1999.....	9.0	86.4	1.6	0.3	2.0	3.2	5.4	6.9	9.3	17.3	38.7	167.3
White, female												
2006 4/.....	12.5	58.1	1.1	0.2	1.0	1.5	2.6	3.6	4.5	9.9	40.0	263.3
2005 4/.....	12.2	57.4	1.1	0.2	1.1	1.8	3.4	4.3	5.0	10.1	37.9	248.8
2004 4/.....	11.6	54.2	1.0	0.2	1.0	1.7	2.8	3.9	4.8	10.1	38.2	241.7
2003 4/.....	12.0	53.3	1.3	0.4	1.3	2.0	3.7	4.4	5.1	10.5	37.2	249.7
2002.....	11.2	59.0	1.2	0.2	1.1	2.0	3.6	4.2	5.0	10.5	35.7	226.1
2001.....	11.9	59.3	1.4	0.4	1.5	2.4	4.8	5.8	6.2	12.1	36.7	223.8
2000.....	11.5	64.3	1.3	0.3	1.3	2.3	4.1	4.9	6.1	12.7	36.7	222.5
1999.....	9.9	62.4	1.1	0.3	0.8	1.7	2.7	3.4	5.0	11.2	32.8	199.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

Black, both sexes

2006 4/.....	11.1	155.8	2.8	0.5	1.8	3.4	5.6	8.8	10.6	19.1	55.4	223.9
2005 4/.....	12.3	169.0	2.5	0.9	2.5	4.2	7.0	11.7	14.0	22.3	51.8	215.1
2004 4/.....	11.6	171.7	1.9	0.5	1.8	4.0	6.5	9.7	13.3	20.4	53.0	220.8
2003 4/.....	12.3	159.3	3.1	0.8	2.4	5.0	8.5	13.4	14.2	23.4	45.3	210.9
2002.....	12.3	162.3	2.6	0.7	2.5	5.1	8.5	13.7	13.7	20.1	49.6	196.0
2001.....	12.6	169.5	2.8	0.6	2.7	5.0	9.0	13.9	14.8	24.4	49.5	191.6
2000.....	13.7	197.1	2.3	0.7	3.2	5.8	9.2	14.7	17.4	28.9	55.5	186.4
1999.....	12.7	192.8	2.7	0.5	2.6	4.8	9.0	13.2	15.9	24.5	54.1	182.4

Black, male

2006 4/.....	11.7	165.9	3.1	0.6	2.2	4.7	7.1	12.3	13.6	27.4	68.5	193.9
2005 4/.....	13.3	186.6	2.7	0.9	3.4	5.4	8.2	15.3	19.7	29.4	63.4	199.9
2004 4/.....	12.6	204.3	2.1	*	2.3	5.2	8.0	13.8	19.0	24.8	59.0	181.8
2003 4/.....	13.5	170.7	3.4	0.8	3.0	6.7	11.1	17.6	20.3	31.4	52.5	193.9
2002.....	14.1	183.0	3.5	0.8	3.0	6.8	11.3	19.4	19.7	26.4	68.3	181.2
2001.....	13.9	187.4	3.5	0.8	3.7	6.0	11.1	18.0	23.3	30.4	51.8	184.9
2000.....	15.5	227.1	2.3	0.8	3.7	7.7	10.8	20.6	25.2	39.9	65.1	161.5
1999.....	14.6	217.6	2.8	0.6	3.5	6.8	11.5	19.4	23.1	32.5	66.1	184.0

Black, female

2006 4/.....	10.5	145.3	2.5	*	1.4	2.2	4.2	5.8	8.2	13.2	48.0	236.2
2005 4/.....	11.4	150.7	2.2	0.8	1.5	3.1	6.0	8.7	9.4	17.2	45.3	221.3
2004 4/.....	10.8	137.7	1.7	0.6	1.3	3.0	5.1	6.2	8.8	17.3	49.6	236.3
2003 4/.....	11.3	147.5	2.9	0.7	1.9	3.3	6.1	9.8	9.2	17.7	41.3	217.5
2002.....	10.6	140.8	1.8	0.7	2.0	3.6	6.0	8.8	8.9	15.7	39.0	201.7
2001.....	11.4	150.6	2.0	*	1.7	4.1	7.2	10.3	7.9	20.2	48.3	194.1
2000.....	12.0	166.0	2.3	*	2.6	4.1	7.7	9.7	11.0	21.2	50.1	195.7
1999.....	10.9	167.1	2.7	*	1.8	3.1	6.8	7.8	10.1	18.9	47.3	181.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes

2006.....	79.3	28.4	3.4	2.0	5.0	8.3	18.4	40.1	71.6	151.6	470.5	1,724.2
2005.....	73.4	29.2	3.8	2.1	4.9	7.9	18.5	40.3	69.5	146.1	434.3	1,559.1
2004.....	68.8	29.2	3.8	2.2	4.7	8.1	18.3	38.9	67.0	141.4	409.6	1,461.6
2003.....	69.3	29.7	4.0	2.3	4.9	8.4	18.9	38.9	67.1	144.4	411.5	1,508.4
2002.....	67.5	28.4	4.2	2.1	4.6	7.9	18.9	37.8	65.3	140.6	405.3	1,493.0
2001.....	65.5	28.0	4.2	2.1	4.5	7.8	18.0	36.2	63.4	140.9	395.3	1,476.7
2000.....	63.4	29.0	4.0	2.1	4.5	7.8	18.0	35.4	62.4	136.0	388.5	1,445.7
1999.....	61.5	30.2	4.2	2.3	4.6	7.6	17.7	33.9	60.5	136.6	373.0	1,410.7

All races 3/, male

2006.....	65.2	32.5	3.6	2.1	5.8	9.5	21.4	48.3	82.7	164.8	474.4	1,455.0
2005.....	60.9	34.8	4.3	2.3	5.8	8.9	22.0	49.1	80.9	156.1	437.0	1,305.3
2004.....	57.2	33.8	4.0	2.3	5.4	9.2	21.5	47.8	77.1	151.9	410.5	1,236.6
2003.....	57.3	34.8	4.2	2.6	5.6	9.3	22.3	47.2	76.5	155.6	415.1	1,258.1
2002.....	55.8	33.4	4.6	2.3	5.1	9.1	22.4	46.3	74.5	152.6	406.7	1,262.9
2001.....	53.8	31.4	4.6	2.2	5.2	8.6	21.3	43.7	71.2	153.4	397.4	1,271.6
2000.....	52.3	32.3	4.3	2.3	5.3	9.0	21.4	42.8	70.7	148.4	391.5	1,266.2
1999.....	51.0	35.0	4.5	2.3	5.3	8.8	21.6	40.5	69.0	148.7	384.2	1,225.4

All races 3/, female

2006.....	93.0	24.1	3.1	1.8	4.2	7.1	15.3	32.2	61.3	140.5	467.9	1,850.2
2005.....	85.6	23.2	3.3	1.9	4.0	6.8	15.0	31.8	59.0	137.7	432.5	1,675.6
2004.....	79.9	24.5	3.6	2.0	3.9	7.0	15.1	30.4	57.6	132.5	408.9	1,562.8
2003.....	81.0	24.5	3.8	1.9	4.2	7.4	15.6	31.0	58.4	135.0	409.1	1,619.1
2002.....	78.8	23.0	3.9	1.9	4.0	6.6	15.5	29.6	56.8	130.6	404.4	1,592.8
2001.....	76.7	24.5	3.8	2.0	3.8	6.9	14.6	29.1	56.2	130.5	394.0	1,562.6
2000.....	74.0	25.6	3.6	2.0	3.7	6.5	14.6	28.2	54.8	125.8	386.5	1,518.8
1999.....	71.6	25.3	3.9	2.2	3.8	6.4	13.9	27.4	52.7	126.7	365.7	1,485.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

White, both sexes

2006 4/.....	85.2	23.8	3.1	1.9	4.8	7.7	17.1	37.7	68.7	149.3	473.8	1,757.7
2005 4/.....	78.5	25.1	3.5	1.9	4.7	7.2	17.2	37.6	66.7	143.4	435.5	1,586.5
2004 4/.....	73.3	25.4	3.5	2.1	4.4	7.1	16.9	36.2	63.8	138.4	410.8	1,486.9
2003 4/.....	73.9	25.5	3.7	2.2	4.7	7.5	17.4	35.9	63.8	142.0	412.1	1,534.2
2002.....	71.8	25.2	3.9	2.1	4.2	7.1	17.3	34.8	62.0	137.8	405.2	1,517.6
2001.....	69.4	23.7	3.9	2.1	4.2	6.9	16.0	32.8	59.7	138.1	395.1	1,498.0
2000.....	67.0	24.9	3.5	2.0	4.3	6.9	16.0	32.0	58.6	132.7	386.9	1,464.6
1999.....	65.0	25.2	4.0	2.1	4.3	7.0	15.5	30.3	56.9	132.6	371.6	1,434.0

White, male

2006 4/.....	69.0	27.3	3.3	2.0	5.8	9.3	20.4	46.1	79.0	161.7	474.0	1,481.8
2005 4/.....	64.1	30.0	4.0	2.1	5.7	8.6	21.2	46.6	77.5	152.3	434.7	1,328.2
2004 4/.....	59.9	29.6	3.7	2.3	5.2	8.4	20.6	45.1	73.3	147.6	407.8	1,251.8
2003 4/.....	59.9	31.5	3.8	2.5	5.7	8.8	21.0	44.0	72.4	151.4	411.4	1,279.1
2002.....	58.2	30.8	4.1	2.2	5.0	8.4	21.3	43.2	70.5	148.1	403.4	1,274.7
2001.....	55.8	25.9	4.4	2.1	5.1	7.8	19.5	39.9	66.8	149.4	394.8	1,282.9
2000.....	54.1	26.8	3.9	2.2	5.2	8.3	19.5	39.3	66.0	143.4	387.1	1,274.8
1999.....	52.7	29.2	4.3	2.1	5.1	8.4	19.6	36.7	64.9	143.7	379.3	1,231.6

White, female

2006 4/.....	101.1	20.2	2.9	1.8	3.7	6.1	13.6	29.3	58.8	138.6	473.6	1,887.2
2005 4/.....	92.6	19.9	3.1	1.7	3.7	5.9	13.1	28.7	56.4	135.7	436.1	1,705.4
2004 4/.....	86.4	21.0	3.3	1.9	3.5	5.8	13.1	27.4	54.8	130.6	412.9	1,592.9
2003 4/.....	87.6	19.3	3.7	1.8	3.7	6.2	13.6	27.9	55.5	133.9	412.6	1,647.2
2002.....	85.2	19.3	3.6	1.9	3.4	5.6	13.3	26.6	54.0	129.0	406.3	1,623.1
2001.....	82.7	21.4	3.4	2.0	3.4	5.8	12.6	25.8	53.0	128.6	395.2	1,588.0
2000.....	79.6	22.9	3.2	1.8	3.4	5.4	12.5	24.7	51.6	123.6	386.7	1,541.7
1999.....	77.0	20.9	3.7	2.1	3.4	5.5	11.4	24.0	49.5	123.3	366.5	1,515.3



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

Black, both sexes

2006 4/.....	66.2	51.8	4.5	2.5	6.6	13.5	30.0	66.8	114.4	205.7	537.8	1,636.8
2005 4/.....	63.6	50.5	5.0	3.3	6.4	13.3	30.8	68.1	111.0	201.3	514.9	1,501.2
2004 4/.....	60.9	49.3	5.3	2.8	6.7	15.2	30.9	67.9	110.8	200.4	482.4	1,405.8
2003 4/.....	60.9	51.8	5.3	2.7	6.5	15.1	33.0	69.6	111.4	196.9	480.2	1,435.7
2002.....	59.7	45.2	5.9	2.6	6.8	14.5	33.7	68.4	108.1	194.2	477.7	1,425.4
2001.....	59.1	51.3	5.4	2.4	6.5	14.1	33.9	70.4	107.2	196.2	470.9	1,418.0
2000.....	58.1	49.6	5.9	2.8	6.5	15.0	34.2	69.3	107.4	191.9	467.7	1,388.9
1999.....	56.2	56.7	5.3	3.0	6.7	12.9	35.3	67.8	103.4	199.8	440.9	1,286.2

Black, male

2006 4/.....	59.3	59.0	4.9	2.8	6.7	12.8	32.1	76.8	137.9	238.5	595.0	1,411.9
2005 4/.....	56.8	62.6	5.8	3.6	6.8	13.5	32.6	78.5	132.1	228.8	563.0	1,241.0
2004 4/.....	56.1	55.7	5.7	2.5	6.9	16.0	32.2	79.7	132.4	234.3	540.1	1,291.5
2003 4/.....	55.7	55.3	6.1	2.9	5.9	15.0	35.4	82.2	132.3	233.6	544.4	1,224.1
2002.....	54.8	48.8	6.4	2.8	6.7	15.4	35.3	81.5	129.1	229.4	522.1	1,319.8
2001.....	54.6	58.4	5.1	2.4	6.6	14.9	38.2	84.2	126.4	229.8	524.1	1,313.9
2000.....	54.3	59.8	6.1	2.5	7.0	15.6	38.7	81.1	130.2	228.4	523.5	1,325.1
1999.....	52.8	64.7	5.4	3.5	7.1	13.4	39.3	80.3	121.7	226.2	505.1	1,291.3

Black, female

2006 4/.....	72.6	44.2	4.0	2.2	6.5	14.1	28.2	58.2	95.4	182.3	505.2	1,728.8
2005 4/.....	69.7	37.9	4.1	3.1	6.1	13.2	29.2	59.0	94.0	181.6	487.6	1,606.2
2004 4/.....	65.4	42.6	4.9	3.0	6.6	14.6	29.8	57.8	93.3	176.2	449.9	1,451.1
2003 4/.....	65.6	48.2	4.5	2.6	7.1	15.2	30.9	58.8	94.6	170.7	444.0	1,517.9
2002.....	64.1	41.5	5.4	2.3	6.9	13.7	32.2	57.2	91.3	169.2	452.5	1,465.9
2001.....	63.1	43.8	5.6	2.5	6.3	13.3	30.0	58.5	91.8	172.6	441.0	1,456.6
2000.....	61.6	39.0	5.7	3.0	6.0	14.4	30.2	59.1	89.1	166.3	436.2	1,412.7
1999.....	59.3	48.3	5.1	2.5	6.2	12.5	31.9	57.0	88.7	181.2	405.0	1,284.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes

2006.....	40.6	27.8	9.9	5.6	38.2	37.0	40.2	45.5	36.2	44.5	105.1	274.9
2005.....	39.7	26.4	10.3	6.0	37.4	34.9	38.6	43.2	35.8	46.3	106.1	279.5
2004.....	38.1	25.8	10.3	6.5	37.0	32.6	37.3	40.7	33.2	44.0	103.7	276.7
2003.....	37.6	23.6	10.9	6.4	37.1	31.5	37.8	38.8	32.9	44.1	101.9	278.9
2002.....	37.0	23.5	10.5	6.6	38.0	31.5	37.2	36.6	31.4	44.2	101.3	275.4
2001.....	35.7	24.2	11.2	6.9	36.1	29.9	35.4	34.1	30.3	42.8	100.9	276.4
2000.....	34.8	23.1	11.9	7.3	36.0	29.5	34.1	32.6	30.9	41.9	95.1	273.5
1999.....	35.1	22.3	12.4	7.6	35.3	29.6	33.8	31.8	30.6	44.6	100.5	282.4

All races 3/, male

2006.....	53.5	31.6	11.1	6.7	56.0	55.8	56.9	65.0	51.6	60.1	135.8	337.1
2005.....	52.3	27.5	12.4	7.2	54.6	52.2	54.8	62.2	51.3	62.6	139.0	344.2
2004.....	49.8	29.0	11.8	8.0	53.3	48.5	53.0	58.6	46.9	59.2	134.7	345.5
2003.....	49.3	26.4	13.0	7.9	53.3	47.1	54.2	55.9	47.1	58.9	132.9	352.5
2002.....	48.9	26.8	12.9	8.2	54.9	47.7	53.7	53.3	45.2	60.6	131.6	346.6
2001.....	47.2	26.2	13.3	8.6	52.9	45.4	51.8	50.6	43.0	57.3	133.4	358.5
2000.....	46.2	25.7	14.0	8.8	52.1	45.0	51.0	49.0	44.1	55.9	124.3	359.7
1999.....	46.4	25.4	14.4	9.4	50.7	45.0	50.2	47.9	44.0	60.2	132.4	375.4

All races 3/, female

2006.....	28.1	23.8	8.6	4.4	19.4	17.6	23.4	26.6	22.0	31.3	84.1	245.8
2005.....	27.5	25.2	8.0	4.7	19.2	16.9	22.3	24.7	21.3	32.5	83.8	249.8
2004.....	26.8	22.5	8.7	5.1	19.9	16.0	21.7	23.4	20.5	31.1	82.9	245.8
2003.....	26.2	20.7	8.7	4.8	19.9	15.3	21.4	22.4	19.7	31.6	81.1	246.4
2002.....	25.6	20.0	8.1	5.0	20.1	14.9	20.8	20.5	18.6	30.6	81.2	244.6
2001.....	24.5	22.1	8.9	5.2	18.4	14.0	19.2	18.0	18.5	30.8	79.6	242.0
2000.....	23.8	20.5	9.7	5.6	19.1	13.7	17.5	16.8	18.8	30.3	76.1	238.4
1999.....	24.1	19.0	10.3	5.6	19.2	14.0	17.6	16.3	18.3	31.8	79.7	244.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

White, both sexes

2006 4/.....	42.9	22.8	9.2	5.3	41.7	39.4	41.8	45.5	35.5	44.7	108.3	288.6
2005 4/.....	41.8	23.0	10.0	5.7	40.4	36.9	40.1	43.1	35.0	46.2	107.9	290.9
2004 4/.....	40.2	22.2	10.0	6.2	40.3	34.2	38.9	40.7	32.6	43.9	106.3	289.9
2003 4/.....	39.5	20.0	10.6	6.1	40.1	33.0	39.0	38.5	32.3	43.8	104.0	291.1
2002.....	38.7	18.6	9.9	6.2	40.8	32.8	38.1	36.2	30.7	43.9	103.6	286.1
2001.....	37.0	20.6	11.0	6.4	38.2	30.9	35.8	33.1	29.4	42.4	102.7	285.5
2000.....	35.9	19.1	11.1	6.9	38.3	29.9	34.3	31.3	30.2	41.1	96.5	282.8
1999.....	36.0	18.9	11.5	7.1	37.2	30.0	33.8	30.5	29.2	43.5	100.9	290.4

White, male

2006 4/.....	55.7	25.8	10.4	6.3	60.8	58.6	58.6	64.1	49.7	59.8	138.9	352.4
2005 4/.....	54.3	23.7	12.3	6.8	58.9	54.7	56.5	61.4	49.2	61.8	140.0	356.6
2004 4/.....	52.0	25.5	11.5	7.5	57.8	50.5	54.8	58.0	45.1	58.8	137.0	359.4
2003 4/.....	51.3	22.5	12.8	7.6	57.4	49.3	55.6	54.9	45.8	58.3	134.4	365.7
2002.....	50.4	22.3	11.8	7.6	58.7	49.3	54.7	52.3	43.5	59.2	133.0	356.6
2001.....	48.4	22.1	13.3	7.8	55.8	46.7	52.3	48.6	41.1	56.3	134.3	368.1
2000.....	47.0	21.7	13.2	8.4	55.0	45.4	51.0	46.7	42.1	54.2	124.3	369.8
1999.....	46.9	21.8	13.2	8.8	52.8	45.4	50.0	45.4	41.5	58.0	131.8	383.6

White, female

2006 4/.....	30.3	19.8	8.0	4.3	21.2	18.9	24.6	27.1	22.1	31.7	87.1	258.7
2005 4/.....	29.5	22.2	7.6	4.5	20.6	18.0	23.4	25.1	21.5	32.9	85.8	260.7
2004 4/.....	28.8	18.8	8.4	4.7	21.5	17.0	22.6	23.6	20.7	31.2	85.4	258.6
2003 4/.....	28.0	17.4	8.3	4.7	21.7	15.9	22.1	22.3	19.5	31.4	83.5	258.0
2002.....	27.3	14.7	7.8	4.7	21.8	15.5	21.3	20.3	18.6	30.9	83.9	255.4
2001.....	25.9	18.9	8.5	5.0	19.5	14.4	19.2	17.8	18.3	30.7	81.6	251.0
2000.....	25.1	16.4	9.0	5.4	20.4	13.8	17.4	16.2	18.9	30.1	78.1	247.4
1999.....	25.3	15.8	9.6	5.4	20.5	14.1	17.4	15.8	17.7	31.3	80.6	253.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

Black, both sexes

2006 4/.....	35.2	52.7	14.0	7.2	26.2	34.2	41.1	55.0	48.3	48.0	80.8	150.5
2005 4/.....	34.9	47.4	12.5	7.6	27.1	32.1	38.7	53.0	48.0	52.5	97.0	174.8
2004 4/.....	32.8	45.8	12.5	8.5	25.8	31.4	36.7	49.8	43.7	48.2	87.1	152.3
2003 4/.....	32.4	43.5	13.4	7.7	25.6	29.9	38.6	48.4	42.3	50.4	82.7	160.2
2002.....	33.1	48.9	14.1	9.1	27.8	31.4	39.1	46.8	41.4	49.3	83.6	173.4
2001.....	33.5	43.1	13.6	9.4	28.7	30.2	40.0	47.7	42.0	51.0	88.5	183.6
2000.....	33.5	44.5	16.2	9.3	28.7	32.5	39.7	47.0	40.8	52.5	84.9	181.1
1999.....	35.2	41.6	17.7	10.5	29.5	32.2	40.0	47.8	44.7	57.8	103.3	207.6

Black, male

2006 4/.....	50.8	61.3	15.9	8.8	39.4	55.2	62.2	85.7	78.0	72.9	115.8	199.3
2005 4/.....	50.0	50.8	13.8	9.3	40.2	50.9	57.9	82.5	80.0	80.6	144.8	225.0
2004 4/.....	46.7	48.3	14.2	10.3	37.8	50.3	54.1	76.7	70.5	73.7	132.0	213.6
2003 4/.....	46.1	47.6	15.7	9.7	38.6	46.7	57.8	74.5	67.9	74.3	123.3	225.2
2002.....	47.9	51.1	18.5	11.6	41.8	50.6	58.4	72.2	68.7	76.9	129.7	260.5
2001.....	48.2	48.5	15.4	12.8	43.2	47.4	59.9	77.2	67.6	74.9	136.0	255.3
2000.....	49.0	47.7	18.8	12.0	43.0	51.8	62.0	75.4	68.8	79.4	132.6	275.6
1999.....	51.0	45.9	21.0	13.8	44.9	51.4	62.0	78.6	71.1	88.2	150.0	316.4

Black, female

2006 4/.....	20.9	43.6	12.1	5.4	12.7	14.7	22.4	28.7	24.4	30.2	60.9	130.5
2005 4/.....	21.2	43.8	11.1	5.7	13.7	14.7	21.5	27.6	22.2	32.4	69.8	154.5
2004 4/.....	20.1	43.2	10.6	6.6	13.6	14.1	21.3	26.7	22.2	30.0	61.7	128.0
2003 4/.....	19.9	39.3	11.0	5.7	12.5	14.5	21.6	26.1	21.7	33.4	59.8	134.9
2002.....	19.7	46.6	9.6	6.6	13.8	14.1	22.0	25.1	19.5	29.7	57.5	140.1
2001.....	20.1	37.5	11.9	6.0	14.2	14.7	22.4	22.3	21.4	34.2	61.8	157.0
2000.....	19.5	41.3	13.5	6.5	14.6	15.0	19.9	22.6	18.3	33.5	58.0	145.9
1999.....	20.8	37.3	14.3	7.1	14.3	14.9	20.6	21.4	23.5	36.4	77.1	166.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Transport accidents (V01-V99,Y85)

All races 3/, both sexes

2006.....	16.2	3.4	3.8	3.6	26.8	19.2	16.6	16.8	15.8	16.8	23.6	24.4
2005.....	16.3	3.6	4.0	3.8	26.8	19.0	16.6	16.7	16.2	18.0	24.3	26.1
2004.....	16.4	3.6	4.1	4.3	27.3	18.8	16.4	16.7	15.5	17.5	25.5	27.2
2003.....	16.5	3.7	4.1	4.3	27.7	18.3	17.1	16.6	15.9	17.5	26.2	29.7
2002.....	16.8	3.1	4.1	4.2	29.1	18.7	17.1	16.4	15.7	18.3	27.0	28.8
2001.....	16.6	3.7	4.4	4.4	27.9	18.8	17.4	16.0	15.3	17.9	27.9	28.9
2000.....	16.6	4.6	4.5	4.7	28.1	18.6	17.0	15.8	16.0	18.0	27.0	31.5
1999.....	16.6	5.0	4.6	4.8	27.6	18.5	16.8	15.4	16.0	19.4	28.4	32.0

All races 3/, male

2006.....	23.1	3.3	4.0	4.1	37.9	28.9	23.9	25.1	22.9	23.4	33.2	42.9
2005.....	23.3	3.5	4.5	4.4	37.9	28.6	24.2	24.9	23.5	25.3	34.8	46.0
2004.....	23.0	3.9	4.4	5.0	37.8	27.7	24.0	24.7	22.2	23.7	36.5	46.5
2003.....	23.3	4.0	4.5	5.2	38.6	27.3	24.8	24.3	23.0	23.7	36.6	52.0
2002.....	23.7	3.5	4.8	4.9	40.7	28.1	24.6	24.0	22.5	25.3	37.7	53.0
2001.....	23.5	3.3	5.0	5.3	39.6	28.3	25.1	23.3	21.6	24.3	39.8	52.7
2000.....	23.3	4.8	5.0	5.4	39.3	27.7	24.8	23.0	22.5	24.3	38.1	59.8
1999.....	23.1	5.2	5.1	5.8	38.1	27.2	24.0	22.4	22.5	26.1	40.7	63.5

All races 3/, female

2006.....	9.4	3.6	3.6	3.1	15.1	9.1	9.2	8.7	9.2	11.2	17.1	15.7
2005.....	9.5	3.6	3.5	3.2	15.1	9.1	9.1	8.8	9.4	11.8	17.1	17.0
2004.....	9.9	3.2	3.8	3.6	16.1	9.5	8.8	9.0	9.3	12.3	18.1	18.5
2003.....	10.0	3.4	3.7	3.4	16.3	9.0	9.4	9.1	9.4	12.4	19.3	19.9
2002.....	10.1	2.8	3.4	3.5	16.9	9.2	9.7	9.0	9.3	12.5	19.8	18.3
2001.....	10.0	4.2	3.8	3.5	15.6	9.2	9.8	8.9	9.4	12.7	20.1	18.9
2000.....	10.2	4.3	4.0	3.9	16.4	9.3	9.3	8.9	10.0	12.8	19.8	19.9
1999.....	10.4	4.8	4.2	3.9	16.6	9.6	9.6	8.7	10.1	14.0	20.4	19.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Transport accidents (V01-V99,Y85)

White, both sexes

2006 4/.....	16.7	3.1	3.5	3.6	28.9	19.6	17.1	17.0	15.6	17.0	23.9	25.3
2005 4/.....	16.9	3.1	3.9	3.9	28.7	19.6	17.1	16.9	16.3	18.1	24.6	27.1
2004 4/.....	16.9	3.3	4.0	4.4	29.5	19.1	16.7	16.7	15.5	17.6	25.9	28.4
2003 4/.....	17.0	3.6	4.0	4.4	29.8	18.7	17.4	16.6	15.7	17.5	26.5	30.5
2002.....	17.3	2.7	3.9	4.2	31.3	19.1	17.3	16.4	15.6	18.4	27.5	29.6
2001.....	17.0	3.7	4.3	4.3	29.5	19.2	17.6	15.8	15.1	17.9	28.5	29.9
2000.....	17.0	4.1	4.3	4.7	30.0	18.5	17.1	15.6	16.0	17.9	27.5	32.2
1999.....	17.0	4.6	4.1	4.8	29.3	18.7	16.8	15.2	15.7	19.4	28.6	32.3

White, male

2006 4/.....	23.7	3.2	3.6	4.0	40.7	29.2	24.4	25.2	22.6	23.4	33.3	44.0
2005 4/.....	24.0	3.3	4.3	4.4	40.5	29.0	24.5	25.1	23.5	25.2	34.9	47.5
2004 4/.....	23.7	3.3	4.2	5.1	40.8	27.8	24.4	24.4	22.0	23.6	36.7	48.6
2003 4/.....	23.9	4.0	4.5	5.2	41.1	27.8	25.0	24.1	22.7	23.7	36.7	52.8
2002.....	24.2	3.1	4.5	5.0	43.4	28.3	24.7	24.0	22.3	24.9	37.7	53.8
2001.....	23.9	3.2	4.9	5.1	41.7	28.7	25.1	22.9	21.2	24.0	40.3	54.1
2000.....	23.6	4.5	4.7	5.5	41.6	27.4	24.7	22.6	22.0	23.7	38.1	61.6
1999.....	23.4	5.2	4.4	5.7	40.1	27.3	23.8	21.8	21.9	25.8	40.6	63.7

White, female

2006 4/.....	9.8	3.1	3.4	3.1	16.3	9.3	9.6	8.8	9.1	11.4	17.3	16.5
2005 4/.....	10.0	2.9	3.5	3.3	16.2	9.6	9.5	8.9	9.5	12.1	17.5	17.7
2004 4/.....	10.3	3.2	3.8	3.6	17.4	9.9	8.9	9.0	9.5	12.4	18.5	19.3
2003 4/.....	10.3	3.1	3.4	3.5	17.7	9.1	9.6	9.1	9.1	12.2	19.7	20.6
2002.....	10.5	2.2	3.4	3.4	18.3	9.4	9.8	8.9	9.4	12.8	20.6	19.1
2001.....	10.3	4.1	3.6	3.5	16.5	9.3	10.0	8.8	9.4	12.7	20.7	19.7
2000.....	10.5	3.7	3.8	4.0	17.7	9.3	9.4	8.7	10.3	12.9	20.5	20.3
1999.....	10.7	3.9	3.8	3.9	17.8	9.8	9.6	8.6	10.0	14.0	20.7	19.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Transport accidents (V01-V99,Y85)

Black, both sexes

2006 4/.....	15.1	4.6	5.2	3.9	19.2	21.4	17.4	17.9	18.7	15.5	20.0	15.6
2005 4/.....	15.0	5.7	4.5	3.9	20.2	19.6	17.3	17.8	17.1	17.9	19.9	16.6
2004 4/.....	15.2	4.8	5.1	4.4	19.6	20.8	17.1	19.5	17.3	16.4	21.6	14.7
2003 4/.....	15.1	4.2	5.0	4.1	19.7	19.3	18.3	18.1	18.5	18.2	22.4	21.0
2002.....	15.4	5.3	5.0	4.6	21.2	20.4	18.2	17.8	17.1	17.1	21.7	18.4
2001.....	15.9	*	5.2	5.3	22.2	19.9	18.7	18.8	18.1	18.8	21.2	17.5
2000.....	16.0	6.5	5.6	4.7	21.7	21.4	18.6	19.3	17.4	19.6	20.0	21.6
1999.....	16.3	7.3	7.1	5.4	22.0	19.9	19.0	19.0	18.3	19.3	26.0	27.7

Black, male

2006 4/.....	22.9	*	5.7	4.7	28.0	34.8	27.4	28.6	29.4	25.3	31.4	36.6
2005 4/.....	22.7	*	4.9	4.6	29.4	32.3	27.7	27.9	27.6	29.4	34.4	33.5
2004 4/.....	22.8	5.9	5.4	4.8	27.9	33.8	26.8	30.9	27.7	26.7	38.4	28.8
2003 4/.....	22.6	*	5.1	5.1	29.3	30.6	28.4	29.0	29.1	26.6	37.5	42.7
2002.....	23.1	*	6.4	5.3	30.5	33.1	27.5	27.8	28.3	28.5	40.4	46.1
2001.....	23.8	*	5.7	6.9	32.5	31.6	28.9	30.2	29.6	27.4	36.2	35.2
2000.....	24.1	6.7	6.2	5.7	31.6	34.4	29.7	29.6	29.4	32.2	36.3	45.0
1999.....	24.2	*	8.0	6.8	32.0	31.8	29.0	30.3	28.9	29.1	43.4	68.0

Black, female

2006 4/.....	8.0	*	4.6	3.1	10.2	9.0	8.5	8.7	10.0	8.5	13.5	*
2005 4/.....	7.9	6.8	4.2	3.3	10.9	7.9	8.1	9.2	8.6	9.6	11.6	9.8
2004 4/.....	8.2	*	4.8	3.9	11.1	8.9	8.6	9.6	9.0	9.1	12.1	9.1
2003 4/.....	8.3	*	5.0	3.1	9.8	9.0	9.4	8.8	9.9	12.2	13.9	12.6
2002.....	8.3	*	3.4	3.9	11.8	9.0	9.9	9.2	8.1	9.0	11.0	*
2001.....	8.7	*	4.7	3.7	11.9	9.4	9.7	9.1	8.9	12.7	12.7	10.9
2000.....	8.5	*	5.0	3.6	11.9	9.7	8.7	10.5	7.6	10.8	10.7	12.9
1999.....	9.2	8.7	6.2	4.0	12.2	9.0	10.1	9.4	9.7	12.4	16.4	12.7

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

All races 3/, both sexes

2006.....	15.1	3.4	3.6	3.3	26.0	18.2	15.3	15.3	14.3	15.4	22.3	23.4
2005.....	15.3	3.6	3.8	3.6	25.9	18.0	15.4	15.1	14.7	16.7	22.9	25.1
2004.....	15.3	3.5	4.0	4.1	26.3	17.6	15.1	15.1	14.1	16.1	24.5	26.1
2003.....	15.4	3.6	3.9	4.0	26.6	17.1	15.7	14.9	14.2	16.2	24.9	28.8
2002.....	15.7	3.0	3.9	3.9	28.2	17.8	15.8	14.8	14.1	17.0	25.7	28.0
2001.....	15.4	3.6	4.1	4.1	26.8	17.5	15.7	14.2	13.7	16.3	26.3	27.8
2000.....	15.4	4.4	4.2	4.3	26.9	17.3	15.3	14.2	14.4	16.5	25.7	30.4
1999.....	15.2	4.8	4.2	4.3	26.2	16.9	14.9	13.6	14.2	17.8	26.5	30.4

All races 3/, male

2006.....	21.4	3.3	3.8	3.7	36.6	27.4	21.8	22.6	20.5	21.1	30.8	41.0
2005.....	21.7	3.5	4.2	4.1	36.5	26.9	22.2	22.2	20.9	23.2	32.4	44.1
2004.....	21.3	3.8	4.2	4.7	36.4	25.8	21.9	21.9	19.8	21.3	34.7	44.2
2003.....	21.4	3.9	4.4	4.8	36.9	25.5	22.5	21.5	20.0	21.3	34.3	50.0
2002.....	21.9	3.3	4.5	4.6	39.3	26.5	22.3	21.3	19.9	22.7	35.3	51.7
2001.....	21.5	3.2	4.5	4.9	38.0	26.2	22.4	20.5	19.0	21.5	37.3	51.1
2000.....	21.3	4.6	4.7	5.0	37.4	25.5	22.0	20.4	19.8	21.7	35.6	57.5
1999.....	20.9	5.0	4.7	5.1	36.0	24.7	21.1	19.4	19.3	23.3	38.0	59.8

All races 3/, female

2006.....	9.0	3.5	3.4	2.9	14.7	8.7	8.8	8.2	8.5	10.6	16.5	15.2
2005.....	9.1	3.6	3.4	3.0	14.7	8.8	8.6	8.2	8.9	11.2	16.5	16.4
2004.....	9.5	3.2	3.7	3.4	15.7	9.1	8.3	8.5	8.8	11.7	17.7	17.9
2003.....	9.5	3.3	3.5	3.2	15.8	8.5	8.9	8.5	8.8	11.9	18.7	19.5
2002.....	9.8	2.8	3.3	3.3	16.6	8.8	9.3	8.6	8.9	12.3	19.3	17.7
2001.....	9.5	4.0	3.6	3.3	15.1	8.7	9.1	8.1	8.8	12.0	19.2	18.0
2000.....	9.7	4.2	3.8	3.6	15.9	8.8	8.8	8.2	9.5	12.3	19.2	19.3
1999.....	9.7	4.7	3.8	3.6	15.9	8.9	8.8	8.0	9.5	13.3	19.1	18.5



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

White, both sexes

2006 4/.....	15.6	3.1	3.4	3.3	27.9	18.5	15.7	15.4	14.1	15.6	22.5	24.4
2005 4/.....	15.8	3.1	3.7	3.6	27.8	18.5	15.7	15.3	14.7	16.8	23.3	26.1
2004 4/.....	15.8	3.2	3.9	4.1	28.5	17.8	15.3	15.0	14.1	16.2	24.9	27.3
2003 4/.....	15.8	3.5	3.8	4.1	28.6	17.5	15.9	14.9	13.9	16.1	25.2	29.5
2002.....	16.2	2.6	3.7	3.9	30.3	18.0	15.9	14.8	14.1	17.1	26.3	28.8
2001.....	15.7	3.5	3.9	4.0	28.4	17.8	15.9	14.1	13.6	16.3	27.0	28.8
2000.....	15.7	3.9	4.0	4.3	28.7	17.1	15.4	14.0	14.3	16.4	26.1	31.1
1999.....	15.5	4.5	3.8	4.3	27.8	17.0	14.9	13.3	13.9	17.7	26.8	30.7

White, male

2006 4/.....	22.0	3.2	3.5	3.6	39.2	27.6	22.2	22.7	20.1	21.1	30.8	42.2
2005 4/.....	22.3	3.3	4.1	4.1	39.1	27.3	22.4	22.4	20.8	23.0	32.6	45.6
2004 4/.....	21.9	3.3	4.0	4.8	39.4	25.8	22.2	21.7	19.5	21.1	34.9	46.3
2003 4/.....	22.0	3.9	4.3	4.9	39.2	25.9	22.6	21.3	19.6	21.3	34.4	50.7
2002.....	22.4	2.9	4.2	4.6	41.9	26.6	22.3	21.2	19.6	22.4	35.5	52.4
2001.....	21.9	3.1	4.4	4.7	40.0	26.4	22.4	20.1	18.6	21.3	37.8	52.6
2000.....	21.6	4.2	4.5	5.0	39.6	25.1	21.8	20.0	19.2	21.1	35.6	59.2
1999.....	21.1	5.0	4.1	5.0	37.8	24.6	20.9	18.8	18.7	23.0	37.9	60.3

White, female

2006 4/.....	9.4	3.0	3.3	2.9	15.9	8.9	9.1	8.2	8.4	10.8	16.8	16.0
2005 4/.....	9.5	2.9	3.4	3.0	15.8	9.3	8.9	8.3	9.0	11.4	16.9	17.1
2004 4/.....	9.8	3.2	3.7	3.4	16.8	9.4	8.4	8.4	9.0	11.9	18.1	18.7
2003 4/.....	9.8	3.0	3.2	3.2	17.2	8.6	9.0	8.5	8.6	11.7	19.0	20.1
2002.....	10.1	2.2	3.2	3.2	17.9	9.0	9.4	8.5	8.9	12.6	20.1	18.5
2001.....	9.8	3.8	3.4	3.2	16.0	8.8	9.3	8.1	8.8	12.0	19.8	18.9
2000.....	10.0	3.5	3.6	3.7	17.1	8.9	8.9	8.0	9.8	12.4	19.8	19.7
1999.....	10.0	3.9	3.5	3.6	17.1	9.1	8.8	7.9	9.3	13.3	19.6	18.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Motor vehicle accidents (V02-V04,V09.0,V09.2,V12-V14,V19.0-V19.2,V19.4-V19.6,V20-V79,V80.3-V80.5,V81.0-V81.1,V82.0-V82.1,V83-V86,V87.0-V87.8,V88.0-V88.8,V89.0,V89.2)

Black, both sexes

2006 4/.....	14.3	4.4	5.0	3.7	18.7	20.4	16.2	16.4	17.4	14.7	18.8	14.0
2005 4/.....	14.1	5.7	4.3	3.8	19.5	18.7	16.3	15.7	16.0	16.8	18.8	15.8
2004 4/.....	14.3	4.7	4.9	4.2	18.7	19.8	16.0	17.8	15.7	15.5	20.2	14.7
2003 4/.....	14.2	4.2	4.9	3.9	18.8	18.4	17.1	16.5	17.0	17.0	21.0	20.7
2002.....	14.5	5.2	4.8	4.2	20.6	19.7	16.9	16.1	15.6	15.9	20.0	18.1
2001.....	14.8	*	5.0	4.9	21.5	18.9	17.2	16.9	16.4	17.3	19.7	16.6
2000.....	15.0	6.5	5.4	4.4	20.9	20.4	17.1	17.9	16.1	18.2	18.9	20.3
1999.....	15.1	7.1	6.4	5.0	21.1	18.6	17.4	17.1	16.5	17.9	24.1	25.5

Black, male

2006 4/.....	21.5	*	5.7	4.5	27.2	33.0	25.3	25.7	27.0	23.7	29.8	33.1
2005 4/.....	21.2	*	4.7	4.3	28.0	30.8	25.9	24.3	25.7	27.3	32.5	32.5
2004 4/.....	21.2	*	5.2	4.6	26.4	31.8	24.7	28.0	24.7	24.9	35.9	28.8
2003 4/.....	21.1	*	5.0	4.7	27.9	29.0	26.4	25.9	26.3	24.2	35.0	41.7
2002.....	21.5	*	6.3	4.9	29.6	31.7	25.3	24.7	25.2	25.5	36.1	45.0
2001.....	22.0	*	5.3	6.3	31.4	29.9	26.1	26.9	26.3	24.2	34.1	35.2
2000.....	22.5	6.7	5.8	5.4	30.2	32.6	27.2	27.1	27.1	29.4	34.5	43.8
1999.....	22.2	*	7.0	6.0	30.5	29.7	26.3	27.0	25.5	26.5	41.2	60.9

Black, female

2006 4/.....	7.7	*	4.4	3.0	10.0	8.7	8.1	8.4	9.7	8.3	12.6	*
2005 4/.....	7.5	6.8	3.9	3.2	10.7	7.5	7.7	8.3	8.3	9.3	11.0	9.0
2004 4/.....	7.9	*	4.7	3.8	10.9	8.7	8.3	9.0	8.5	8.7	11.3	9.1
2003 4/.....	8.0	*	4.8	3.1	9.6	8.7	8.9	8.4	9.6	11.9	13.1	12.6
2002.....	8.0	*	3.4	3.5	11.6	8.8	9.4	8.8	7.9	9.0	10.9	*
2001.....	8.2	*	4.6	3.3	11.7	8.9	9.2	8.3	8.3	12.5	11.6	9.7
2000.....	8.2	*	4.9	3.5	11.7	9.4	8.2	10.0	7.3	10.3	10.1	11.6
1999.....	8.7	8.7	5.8	3.9	11.8	8.6	9.6	8.7	9.2	11.9	14.5	12.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other land transport accidents (V01,V05-V06,V09.1,V09.3-V09.9,V10-V11,V15-V18,V19.3,V19.8-V19.9,V80.0-V80.2,V80.6-V80.9,V81.2-V81.9,V82.2-V82.9,V87.9,V88.9,V89.1,V89.3,V89.9)

All races 3/, both sexes

2006.....	0.4	*	*	0.1	0.4	0.3	0.5	0.6	0.5	0.5	0.6	0.5
2005.....	0.4	*	0.2	0.1	0.5	0.4	0.5	0.6	0.5	0.4	0.6	0.5
2004.....	0.4	*	*	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.6
2003.....	0.5	*	*	0.1	0.6	0.5	0.5	0.6	0.5	0.4	0.6	0.5
2002.....	0.4	*	*	0.2	0.4	0.3	0.5	0.5	0.4	0.4	0.5	*
2001.....	0.5	*	0.2	0.2	0.5	0.4	0.6	0.6	0.5	0.6	0.8	0.7
2000.....	0.5	*	0.2	0.2	0.6	0.5	0.7	0.6	0.5	0.5	0.7	0.8
1999.....	0.8	*	0.3	0.3	0.9	0.8	0.9	0.8	0.8	0.8	1.3	1.3

All races 3/, male

2006.....	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.8	1.0	*
2005.....	0.7	*	*	0.2	0.7	0.7	0.7	1.0	0.8	0.6	1.1	*
2004.....	0.6	*	*	0.2	0.7	0.7	0.8	0.9	0.7	0.7	0.9	*
2003.....	0.7	*	*	0.2	1.0	0.8	0.8	1.0	0.8	0.8	1.0	*
2002.....	0.6	*	*	0.2	0.6	0.5	0.8	0.9	0.7	0.7	1.0	*
2001.....	0.8	*	0.3	0.2	0.8	0.8	1.0	1.0	0.9	1.0	1.1	*
2000.....	0.8	*	*	0.3	1.0	0.8	1.1	0.9	0.8	0.9	1.2	*
1999.....	1.1	*	0.3	0.4	1.4	1.2	1.4	1.3	1.2	1.2	1.8	2.9

All races 3/, female

2006.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.2	0.3	*
2005.....	0.2	*	*	*	0.2	0.1	0.2	0.3	0.2	0.3	0.3	*
2004.....	0.2	*	*	*	0.2	0.2	0.2	0.2	0.2	0.3	*	*
2003.....	0.2	*	*	0.1	0.3	0.2	0.2	0.2	0.2	*	0.4	*
2002.....	0.1	*	*	0.1	0.1	0.2	0.2	0.1	*	*	*	*
2001.....	0.2	*	*	0.2	0.2	0.1	0.2	0.3	0.2	0.3	0.6	*
2000.....	0.3	*	*	0.2	0.3	0.2	0.3	0.4	0.2	0.2	0.5	*
1999.....	0.4	*	0.3	0.2	0.4	0.4	0.4	0.4	0.4	0.5	1.0	*





Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes

2006.....	0.6	*	*	0.2	0.5	0.7	0.8	0.9	1.0	0.8	0.8	0.4
2005.....	0.6	*	*	0.1	0.4	0.6	0.8	1.0	1.1	0.9	0.7	0.5
2004.....	0.7	*	*	0.1	0.5	0.8	0.8	1.1	1.0	0.9	0.5	0.5
2003.....	0.7	*	*	0.2	0.5	0.6	0.9	1.1	1.2	0.9	0.7	*
2002.....	0.7	*	*	0.1	0.5	0.6	0.9	1.0	1.1	0.9	0.8	0.4
2001.....	0.7	*	0.2	0.1	0.5	0.9	1.0	1.1	1.0	1.0	0.8	0.5
2000.....	0.7	*	*	0.1	0.6	0.8	0.9	1.0	1.1	1.0	0.6	*
1999.....	0.7	*	*	0.2	0.5	0.8	0.9	1.0	1.1	0.9	0.5	*

All races 3/, male

2006.....	1.0	*	*	0.2	0.7	1.1	1.3	1.6	1.7	1.4	1.4	*
2005.....	1.0	*	*	0.1	0.7	1.0	1.2	1.7	1.8	1.5	1.3	*
2004.....	1.1	*	*	0.1	0.7	1.2	1.4	1.8	1.7	1.7	0.9	*
2003.....	1.1	*	*	0.2	0.8	0.9	1.5	1.8	2.2	1.6	1.3	*
2002.....	1.1	*	*	0.1	0.8	1.1	1.5	1.8	1.9	1.9	1.5	*
2001.....	1.2	*	*	0.2	0.9	1.4	1.6	1.9	1.8	1.7	1.4	*
2000.....	1.2	*	*	0.2	0.9	1.3	1.6	1.7	1.8	1.7	1.3	*
1999.....	1.1	*	*	0.3	0.8	1.4	1.4	1.7	2.0	1.6	0.9	*

All races 3/, female

2006.....	0.3	*	*	0.1	0.2	0.3	0.3	0.3	0.4	0.4	0.3	*
2005.....	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.4	0.3	0.3	*
2004.....	0.3	*	*	*	0.3	0.3	0.3	0.4	0.3	0.2	*	*
2003.....	0.2	*	*	0.1	0.2	0.3	0.3	0.4	0.3	0.3	0.3	*
2002.....	0.2	*	*	0.1	0.2	0.2	0.3	0.3	0.3	*	0.3	*
2001.....	0.3	*	*	0.1	0.2	0.3	0.4	0.4	0.3	0.3	0.4	*
2000.....	0.2	*	*	*	0.2	0.2	0.2	0.3	0.3	0.3	*	*
1999.....	0.3	*	*	0.1	0.2	0.3	0.4	0.3	0.3	0.2	0.3	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

White, both sexes

2006 4/.....	0.7	*	*	0.2	0.6	0.7	0.9	1.0	1.1	0.9	0.8	*
2005 4/.....	0.7	*	*	0.1	0.5	0.7	0.8	1.1	1.2	1.0	0.7	0.5
2004 4/.....	0.7	*	*	0.1	0.6	0.8	0.9	1.2	1.0	0.9	0.5	0.6
2003 4/.....	0.7	*	*	0.2	0.5	0.7	1.0	1.2	1.3	0.9	0.7	*
2002.....	0.7	*	*	0.1	0.5	0.7	0.9	1.1	1.2	0.9	0.7	*
2001.....	0.8	*	0.2	0.1	0.6	0.9	1.1	1.2	1.1	1.0	0.8	*
2000.....	0.7	*	*	0.1	0.6	0.9	1.0	1.0	1.1	1.0	0.7	*
1999.....	0.7	*	*	0.2	0.6	0.9	1.0	1.1	1.1	0.9	0.5	*

White, male

2006 4/.....	1.1	*	*	0.2	0.8	1.1	1.4	1.7	1.8	1.5	1.4	*
2005 4/.....	1.1	*	*	0.1	0.7	1.1	1.3	1.8	2.0	1.6	1.3	*
2004 4/.....	1.2	*	*	0.1	0.8	1.3	1.4	2.0	1.8	1.8	1.0	*
2003 4/.....	1.2	*	*	0.2	0.8	1.0	1.6	2.0	2.3	1.7	1.3	*
2002.....	1.2	*	*	0.1	0.8	1.2	1.6	1.9	2.0	1.9	1.4	*
2001.....	1.3	*	*	0.2	0.9	1.5	1.8	1.9	1.8	1.8	1.4	*
2000.....	1.3	*	*	0.2	1.0	1.4	1.8	1.8	1.9	1.8	1.3	*
1999.....	1.2	*	*	0.3	0.9	1.5	1.5	1.8	2.0	1.6	0.9	*

White, female

2006 4/.....	0.3	*	*	*	0.3	0.3	0.4	0.4	0.4	0.4	0.3	*
2005 4/.....	0.3	*	*	0.1	0.2	0.2	0.3	0.3	0.4	0.4	0.3	*
2004 4/.....	0.3	*	*	*	0.3	0.3	0.4	0.4	0.3	*	*	*
2003 4/.....	0.3	*	*	0.2	0.2	0.3	0.4	0.4	0.3	0.3	*	*
2002.....	0.2	*	*	*	0.2	0.2	0.3	0.3	0.4	*	0.3	*
2001.....	0.3	*	*	*	0.3	0.4	0.5	0.4	0.4	0.3	0.4	*
2000.....	0.2	*	*	*	0.3	0.2	0.3	0.3	0.4	0.3	*	*
1999.....	0.3	*	*	0.1	0.3	0.4	0.4	0.3	0.3	0.3	*	*





Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes

2006.....	24.4	24.3	6.1	2.0	11.4	17.8	23.6	28.7	20.5	27.8	81.4	250.5
2005.....	23.4	22.8	6.3	2.1	10.6	15.8	21.9	26.5	19.6	28.3	81.9	253.4
2004.....	21.8	22.2	6.1	2.2	9.8	13.8	20.9	24.0	17.7	26.5	78.2	249.5
2003.....	21.0	19.9	6.8	2.1	9.3	13.2	20.7	22.3	16.9	26.6	75.6	249.2
2002.....	20.2	20.3	6.4	2.4	8.9	12.7	20.1	20.2	15.7	25.9	74.4	246.7
2001.....	19.0	20.5	6.8	2.5	8.2	11.1	18.0	18.1	15.0	24.9	73.0	247.5
2000.....	18.2	18.6	7.4	2.6	7.9	11.0	17.2	16.8	14.9	23.8	68.1	242.0
1999.....	18.4	17.3	7.7	2.7	7.7	11.1	17.0	16.4	14.6	25.1	72.1	250.3

All races 3/, male

2006.....	30.4	28.3	7.2	2.6	18.1	26.8	32.9	39.9	28.7	36.7	102.6	294.3
2005.....	29.0	24.0	7.9	2.8	16.7	23.6	30.6	37.3	27.8	37.4	104.2	298.3
2004.....	26.8	25.1	7.4	3.0	15.4	20.8	28.9	34.0	24.6	35.5	98.2	299.0
2003.....	26.0	22.4	8.5	2.8	14.7	19.9	29.4	31.6	24.1	35.3	96.3	300.5
2002.....	25.2	23.3	8.1	3.2	14.2	19.6	29.1	29.3	22.7	35.3	93.9	293.6
2001.....	23.8	22.9	8.3	3.3	13.3	17.1	26.8	27.3	21.4	33.0	93.6	305.8
2000.....	22.9	20.9	9.0	3.4	12.8	17.3	26.2	26.0	21.6	31.5	86.2	299.8
1999.....	23.3	20.2	9.3	3.7	12.5	17.7	26.2	25.6	21.5	34.1	91.7	311.9

All races 3/, female

2006.....	18.6	20.2	5.0	1.4	4.3	8.5	14.1	17.8	12.8	20.1	67.0	230.1
2005.....	18.0	21.5	4.5	1.4	4.1	7.8	13.2	16.0	11.8	20.7	66.7	232.8
2004.....	16.9	19.3	4.8	1.5	3.8	6.5	12.9	14.4	11.2	18.9	64.8	227.3
2003.....	16.2	17.3	5.0	1.4	3.6	6.3	12.0	13.3	10.3	19.3	61.8	226.5
2002.....	15.4	17.2	4.7	1.5	3.2	5.7	11.1	11.5	9.2	18.1	61.4	226.3
2001.....	14.5	17.9	5.1	1.6	2.8	4.9	9.4	9.2	9.1	18.1	59.5	223.1
2000.....	13.6	16.2	5.7	1.7	2.7	4.5	8.2	7.9	8.8	17.5	56.3	218.4
1999.....	13.7	14.2	6.1	1.7	2.6	4.4	8.0	7.6	8.2	17.8	59.4	225.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

White, both sexes

2006 4/.....	26.2	19.7	5.7	1.7	12.7	19.8	24.8	28.6	19.9	27.7	84.4	263.4
2005 4/.....	24.9	19.8	6.1	1.8	11.7	17.3	23.1	26.2	18.7	28.1	83.3	263.9
2004 4/.....	23.3	19.0	6.0	1.8	10.8	15.1	22.1	24.1	17.0	26.3	80.4	261.5
2003 4/.....	22.5	16.4	6.6	1.8	10.3	14.3	21.6	21.9	16.6	26.3	77.5	260.6
2002.....	21.4	15.9	5.9	1.9	9.6	13.7	20.8	19.8	15.1	25.5	76.1	256.5
2001.....	20.0	16.9	6.7	2.1	8.7	11.7	18.2	17.3	14.3	24.6	74.2	255.7
2000.....	18.9	15.1	6.9	2.2	8.2	11.4	17.2	15.8	14.2	23.2	69.0	250.6
1999.....	19.0	14.3	7.3	2.3	7.9	11.3	17.0	15.3	13.5	24.1	72.3	258.1

White, male

2006 4/.....	32.0	22.6	6.8	2.3	20.1	29.5	34.3	38.9	27.1	36.4	105.6	308.4
2005 4/.....	30.3	20.4	8.0	2.4	18.4	25.7	32.0	36.3	25.7	36.6	105.1	309.1
2004 4/.....	28.3	22.1	7.3	2.4	17.0	22.6	30.4	33.6	23.1	35.2	100.3	310.8
2003 4/.....	27.4	18.6	8.3	2.3	16.3	21.5	30.6	30.7	23.1	34.6	97.8	312.9
2002.....	26.2	19.2	7.4	2.6	15.3	20.9	29.9	28.3	21.3	34.3	95.2	302.8
2001.....	24.5	18.9	8.4	2.7	14.1	18.0	27.2	25.7	19.9	32.3	94.0	314.0
2000.....	23.4	17.3	8.4	2.9	13.4	18.0	26.2	24.1	20.2	30.4	86.2	308.2
1999.....	23.5	16.5	8.8	3.0	12.8	18.1	26.1	23.5	19.6	32.3	91.3	319.8

White, female

2006 4/.....	20.5	16.7	4.6	1.1	4.9	9.6	15.0	18.3	13.0	20.2	69.7	242.2
2005 4/.....	19.6	19.2	4.1	1.2	4.5	8.4	13.9	16.2	11.9	20.8	68.3	243.0
2004 4/.....	18.5	15.6	4.6	1.2	4.2	7.2	13.6	14.6	11.3	18.7	66.9	239.3
2003 4/.....	17.7	14.2	4.9	1.1	3.9	6.8	12.4	13.2	10.4	19.2	63.8	237.4
2002.....	16.8	12.5	4.4	1.2	3.5	6.1	11.5	11.3	9.2	18.0	63.3	236.3
2001.....	15.6	14.8	4.8	1.5	3.0	5.0	9.2	9.0	9.0	18.0	60.9	231.2
2000.....	14.6	12.8	5.2	1.4	2.7	4.5	8.0	7.5	8.6	17.2	57.7	227.2
1999.....	14.6	11.8	5.8	1.5	2.7	4.3	7.8	7.2	7.7	17.3	59.9	233.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Nontransport accidents (W00-X59,Y86)

Black, both sexes

2006 4/.....	20.1	48.1	8.9	3.3	7.0	12.8	23.7	37.1	29.7	32.5	60.8	134.9
2005 4/.....	20.0	41.6	7.9	3.6	6.9	12.4	21.3	35.2	31.0	34.6	77.1	158.2
2004 4/.....	17.6	41.0	7.4	4.1	6.2	10.6	19.6	30.4	26.4	31.7	65.4	137.6
2003 4/.....	17.2	39.3	8.4	3.6	5.9	10.6	20.3	30.3	23.9	32.2	60.3	139.2
2002.....	17.8	43.6	9.2	4.5	6.6	11.0	20.9	29.0	24.3	32.2	61.9	155.0
2001.....	17.6	40.2	8.4	4.1	6.5	10.3	21.3	28.9	23.8	32.3	67.3	166.1
2000.....	17.6	38.0	10.6	4.6	7.0	11.0	21.1	27.7	23.5	32.8	64.9	159.5
1999.....	18.9	34.4	10.6	5.1	7.5	12.3	21.0	28.7	26.4	38.5	77.2	179.8

Black, male

2006 4/.....	27.9	57.0	10.2	4.1	11.4	20.5	34.8	57.1	48.7	47.6	84.4	162.6
2005 4/.....	27.3	46.1	8.9	4.7	10.8	18.5	30.2	54.7	52.5	51.2	110.4	191.5
2004 4/.....	23.9	42.4	8.9	5.4	9.9	16.5	27.3	45.7	42.8	47.0	93.5	184.8
2003 4/.....	23.5	44.0	10.6	4.6	9.2	16.2	29.5	45.5	38.7	47.7	85.7	182.5
2002.....	24.8	45.9	12.1	6.3	11.2	17.5	30.9	44.4	40.4	48.4	89.2	214.4
2001.....	24.4	46.4	9.7	5.8	10.7	15.8	31.0	47.1	37.9	47.5	99.8	220.1
2000.....	24.9	40.9	12.6	6.3	11.4	17.4	32.3	45.8	39.3	47.3	96.3	230.6
1999.....	26.8	40.0	13.0	7.0	12.9	19.5	32.9	48.3	42.2	59.1	106.7	248.4

Black, female

2006 4/.....	12.9	38.8	7.5	2.4	2.5	5.7	13.9	19.9	14.4	21.7	47.4	123.6
2005 4/.....	13.3	37.0	6.9	2.5	2.9	6.8	13.4	18.4	13.6	22.7	58.2	144.8
2004 4/.....	11.9	39.5	5.9	2.7	2.5	5.2	12.7	17.1	13.2	20.9	49.6	118.9
2003 4/.....	11.6	34.3	6.0	2.6	2.6	5.5	12.2	17.3	11.9	21.2	45.9	122.3
2002.....	11.4	41.2	6.2	2.7	2.0	5.1	12.0	15.9	11.4	20.7	46.4	132.3
2001.....	11.5	33.7	7.1	2.4	2.4	5.3	12.6	13.2	12.5	21.6	49.1	146.1
2000.....	11.0	35.1	8.5	2.9	2.7	5.3	11.3	12.1	10.6	22.7	47.2	133.1
1999.....	11.7	28.5	8.1	3.1	2.0	5.9	10.5	12.0	13.8	24.0	60.8	154.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Falls (W00-W19)

All races 3/, both sexes

2006.....	7.0	0.6	0.2	0.1	0.6	0.8	1.4	3.0	5.0	12.6	47.7	151.6
2005.....	6.6	*	0.2	0.1	0.6	0.7	1.4	2.8	4.8	12.4	45.6	147.7
2004.....	6.4	0.6	0.3	0.1	0.6	0.8	1.5	2.8	4.8	12.2	43.8	143.3
2003.....	5.9	*	0.3	0.1	0.6	0.7	1.4	2.6	4.4	11.2	40.8	135.9
2002.....	5.6	*	0.2	0.1	0.6	0.8	1.5	2.5	4.1	10.8	38.3	130.4
2001.....	5.3	0.6	0.2	0.2	0.6	0.9	1.4	2.6	4.0	10.0	35.3	121.5
2000.....	4.7	*	0.2	0.1	0.6	0.8	1.3	2.3	3.9	9.0	31.1	112.6
1999.....	4.7	*	0.4	0.1	0.6	0.9	1.4	2.3	3.7	9.0	31.5	110.2

All races 3/, male

2006.....	7.3	*	0.3	0.1	1.0	1.3	2.3	4.7	7.3	16.5	59.4	181.1
2005.....	7.0	*	0.2	0.1	0.9	1.3	2.2	4.3	7.0	16.2	57.8	174.8
2004.....	6.8	*	0.4	0.1	1.0	1.3	2.4	4.5	7.1	16.5	54.8	174.4
2003.....	6.2	*	0.5	0.1	0.9	1.2	2.4	4.0	6.2	14.5	51.6	165.1
2002.....	6.0	*	0.4	0.1	1.0	1.3	2.3	3.9	6.0	14.5	48.1	158.9
2001.....	5.8	*	0.3	0.2	1.1	1.5	2.4	4.3	5.9	13.1	45.7	155.7
2000.....	5.2	*	0.3	0.1	1.0	1.4	2.2	3.7	5.8	11.9	39.9	145.4
1999.....	5.2	*	0.5	0.2	1.1	1.5	2.4	3.7	5.7	12.4	40.5	142.5

All races 3/, female

2006.....	6.6	*	*	*	0.1	0.2	0.5	1.3	3.0	9.4	39.8	137.8
2005.....	6.3	*	*	*	0.2	0.2	0.5	1.3	2.7	9.2	37.4	135.2
2004.....	6.0	*	*	*	0.2	0.2	0.6	1.3	2.7	8.6	36.4	129.2
2003.....	5.6	*	*	*	0.2	0.2	0.5	1.2	2.7	8.4	33.5	122.9
2002.....	5.3	*	*	*	0.2	0.2	0.6	1.2	2.3	7.7	31.8	118.1
2001.....	4.8	*	*	0.1	0.1	0.2	0.5	1.0	2.2	7.4	28.5	107.2
2000.....	4.3	*	*	*	0.2	0.2	0.5	0.9	2.1	6.6	25.3	99.2
1999.....	4.3	*	*	*	0.2	0.2	0.5	0.9	1.9	6.3	25.7	97.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Falls (W00-W19)

White, both sexes

2006 4/.....	7.9	*	0.2	0.1	0.6	0.9	1.5	3.0	5.1	13.2	50.5	161.1
2005 4/.....	7.5	*	*	0.1	0.7	0.8	1.5	2.8	4.8	13.0	47.9	156.7
2004 4/.....	7.3	*	0.3	0.1	0.7	0.9	1.6	2.9	4.8	12.7	46.4	152.6
2003 4/.....	6.7	*	0.3	0.1	0.6	0.8	1.5	2.5	4.4	11.5	42.8	144.4
2002.....	6.4	*	0.2	0.1	0.7	0.9	1.5	2.6	4.1	11.1	40.4	138.0
2001.....	6.0	*	0.2	0.2	0.8	0.9	1.5	2.6	4.0	10.4	37.0	128.0
2000.....	5.3	*	0.2	0.1	0.7	0.8	1.4	2.3	3.9	9.2	32.6	119.1
1999.....	5.2	*	0.4	0.1	0.7	0.9	1.5	2.2	3.5	9.1	32.9	115.9

White, male

2006 4/.....	8.2	*	*	*	1.1	1.5	2.5	4.7	7.3	17.0	62.2	191.2
2005 4/.....	7.8	*	*	*	1.1	1.3	2.4	4.3	7.0	16.9	60.2	183.8
2004 4/.....	7.6	*	0.4	*	1.1	1.4	2.5	4.5	6.9	17.1	57.6	183.7
2003 4/.....	6.9	*	0.5	0.2	1.0	1.3	2.4	3.9	6.2	14.7	53.8	175.1
2002.....	6.6	*	*	0.1	1.1	1.5	2.4	3.9	6.0	14.8	50.2	166.4
2001.....	6.4	*	0.3	0.2	1.3	1.6	2.5	4.2	5.8	13.5	47.4	162.3
2000.....	5.7	*	*	0.2	1.2	1.5	2.2	3.7	5.7	12.1	41.4	152.8
1999.....	5.6	*	0.5	0.2	1.2	1.5	2.4	3.5	5.4	12.3	41.9	149.1

White, female

2006 4/.....	7.7	*	*	*	0.2	0.2	0.6	1.4	3.0	9.9	42.4	147.0
2005 4/.....	7.3	*	*	*	0.2	0.2	0.6	1.4	2.8	9.7	39.4	144.2
2004 4/.....	7.0	*	*	*	0.2	0.2	0.6	1.3	2.8	8.9	38.7	138.6
2003 4/.....	6.5	*	*	*	0.2	0.3	0.5	1.1	2.8	8.8	35.4	130.8
2002.....	6.2	*	*	*	0.2	0.2	0.7	1.2	2.4	8.0	33.9	125.6
2001.....	5.5	*	*	*	0.1	0.2	0.4	1.1	2.2	7.7	30.2	113.6
2000.....	5.0	*	*	*	0.2	0.2	0.5	0.9	2.1	6.7	26.8	105.4
1999.....	4.9	*	*	*	0.2	0.2	0.6	0.8	1.8	6.4	27.1	102.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Falls (W00-W19)

Black, both sexes

2006 4/.....	2.7	*	*	*	0.3	0.4	1.1	3.1	4.7	9.6	23.2	54.0
2005 4/.....	2.6	*	*	*	*	0.5	0.9	3.1	4.8	8.8	25.4	52.7
2004 4/.....	2.6	*	*	*	*	0.7	1.4	3.0	5.6	10.1	20.3	46.5
2003 4/.....	2.4	*	*	*	*	0.5	1.3	3.3	4.8	8.9	19.4	44.1
2002.....	2.2	*	*	*	*	0.4	1.3	2.9	4.2	8.2	16.1	49.3
2001.....	2.3	*	*	*	*	0.5	1.6	2.9	4.4	8.7	18.5	44.5
2000.....	2.1	*	*	*	*	0.4	1.3	2.7	4.7	8.3	14.9	38.8
1999.....	2.4	*	*	*	0.4	0.8	1.3	3.4	5.5	9.2	17.1	45.3

Black, male

2006 4/.....	3.4	*	*	*	0.6	*	1.8	5.3	7.8	14.4	33.6	67.9
2005 4/.....	3.2	*	*	*	*	0.8	1.6	5.3	7.8	12.5	35.0	63.2
2004 4/.....	3.4	*	*	*	*	1.1	2.1	4.9	10.1	14.4	29.2	71.5
2003 4/.....	3.0	*	*	*	*	0.8	2.2	5.2	7.1	12.5	24.8	56.3
2002.....	3.0	*	*	*	*	0.8	2.2	4.8	6.7	11.8	24.4	77.2
2001.....	3.1	*	*	*	*	0.8	2.6	5.4	7.0	12.6	28.5	68.1
2000.....	2.8	*	*	*	*	*	2.1	4.7	8.2	11.4	21.2	61.1
1999.....	3.3	*	*	*	*	1.3	2.5	5.8	8.8	14.7	24.3	58.6

Black, female

2006 4/.....	2.0	*	*	*	*	*	*	1.3	2.2	6.2	17.4	48.3
2005 4/.....	2.1	*	*	*	*	*	*	1.2	2.3	6.2	20.0	48.4
2004 4/.....	1.8	*	*	*	*	*	0.8	1.3	2.0	7.0	15.3	36.6
2003 4/.....	1.9	*	*	*	*	*	*	1.6	2.9	6.3	16.3	39.3
2002.....	1.5	*	*	*	*	*	*	1.2	2.2	5.6	11.4	38.6
2001.....	1.6	*	*	*	*	*	*	*	2.3	5.9	12.9	35.8
2000.....	1.4	*	*	*	*	*	*	1.0	2.0	6.1	11.4	30.5
1999.....	1.7	*	*	*	*	*	*	1.3	2.9	5.3	13.1	40.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental discharge of firearms (W32-W34)

All races 3/, both sexes

2006.....	0.2	*	*	0.1	0.5	0.3	0.2	0.2	0.2	0.2	0.3	*
2005.....	0.3	*	0.1	0.1	0.5	0.3	0.3	0.3	0.2	0.2	0.3	*
2004.....	0.2	*	*	0.1	0.4	0.2	0.2	0.3	0.1	0.2	0.2	*
2003.....	0.3	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002.....	0.3	*	*	0.1	0.5	0.4	0.3	0.2	0.2	0.2	0.3	*
2001.....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000.....	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999.....	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*

All races 3/, male

2006.....	0.4	*	*	0.2	0.9	0.5	0.3	0.3	0.3	0.4	0.5	*
2005.....	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2004.....	0.4	*	*	0.2	0.7	0.5	0.4	0.4	0.2	0.4	0.4	*
2003.....	0.5	*	*	0.2	0.9	0.5	0.5	0.4	0.4	0.4	0.6	*
2002.....	0.5	*	*	0.2	0.9	0.6	0.4	0.4	0.4	0.4	0.5	*
2001.....	0.5	*	*	0.2	0.9	0.5	0.5	0.5	0.5	0.3	0.7	*
2000.....	0.5	*	*	0.3	0.9	0.5	0.6	0.4	0.4	0.4	0.5	*
1999.....	0.5	*	*	0.3	1.2	0.6	0.5	0.4	0.4	0.4	0.4	*

All races 3/, female

2006.....	0.0	*	*	*	*	*	*	*	*	*	*	*
2005.....	0.1	*	*	*	*	*	0.1	0.1	*	*	*	*
2004.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.1	*	*	*	*	0.1	0.1	*	*	*	*	*
1999.....	0.1	*	*	*	0.1	0.1	0.1	*	*	*	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental discharge of firearms (W32-W34)

White, both sexes

2006 4/.....	0.2	*	*	0.1	0.3	0.3	0.2	0.2	0.2	0.2	0.3	*
2005 4/.....	0.3	*	*	0.1	0.4	0.3	0.3	0.3	0.3	0.3	0.3	*
2004 4/.....	0.2	*	*	0.1	0.4	0.2	0.2	0.3	0.1	0.2	0.2	*
2003 4/.....	0.2	*	*	0.1	0.5	0.3	0.3	0.2	0.2	0.2	0.3	*
2002.....	0.3	*	*	0.1	0.4	0.3	0.3	0.2	0.2	0.2	0.2	*
2001.....	0.3	*	*	0.1	0.5	0.3	0.3	0.3	0.3	0.2	0.3	*
2000.....	0.3	*	*	0.2	0.5	0.3	0.3	0.2	0.2	0.2	0.2	*
1999.....	0.3	*	*	0.2	0.6	0.4	0.3	0.3	0.2	0.2	0.2	*

White, male

2006 4/.....	0.4	*	*	0.2	0.6	0.4	0.3	0.3	0.3	0.4	0.6	*
2005 4/.....	0.4	*	*	0.2	0.8	0.4	0.4	0.4	0.4	0.5	0.5	*
2004 4/.....	0.4	*	*	0.2	0.6	0.4	0.4	0.5	0.2	0.4	*	*
2003 4/.....	0.4	*	*	0.2	0.8	0.5	0.5	0.4	0.5	0.4	0.6	*
2002.....	0.4	*	*	0.2	0.8	0.5	0.4	0.4	0.4	0.4	0.5	*
2001.....	0.5	*	*	0.3	0.9	0.5	0.6	0.5	0.5	0.3	0.7	*
2000.....	0.5	*	*	0.3	0.8	0.5	0.6	0.4	0.4	0.4	0.5	*
1999.....	0.5	*	*	0.3	1.0	0.6	0.5	0.4	0.4	0.4	*	*

White, female

2006 4/.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2005 4/.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2004 4/.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2003 4/.....	0.1	*	*	*	*	*	*	*	*	*	*	*
2002.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2001.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
2000.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*
1999.....	0.1	*	*	*	*	*	0.1	*	*	*	*	*





Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental drowning and submersion (W65-W74)

All races 3/, both sexes

2006.....	1.2	1.2	2.8	0.6	1.5	1.1	1.1	1.1	1.0	1.1	1.5	1.4
2005.....	1.2	1.6	3.0	0.6	1.5	1.0	1.1	1.2	0.9	1.0	1.4	1.8
2004.....	1.1	1.5	2.7	0.7	1.4	1.0	1.0	1.1	0.9	1.0	1.2	1.6
2003.....	1.1	1.4	2.9	0.7	1.4	0.9	1.0	1.0	0.9	1.1	1.3	1.8
2002.....	1.2	1.6	2.9	0.8	1.5	1.1	1.2	0.9	0.8	0.9	1.3	1.7
2001.....	1.2	1.7	3.0	0.8	1.5	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000.....	1.2	2.0	3.2	0.9	1.6	1.1	1.1	0.9	0.9	1.0	1.3	1.5
1999.....	1.3	1.8	3.2	0.9	1.7	1.1	1.1	1.0	0.9	1.0	1.3	1.6

All races 3/, male

2006.....	1.9	1.3	3.5	0.9	2.5	1.9	1.7	1.8	1.5	1.7	2.4	2.4
2005.....	1.9	1.3	4.1	0.9	2.7	1.6	1.9	1.9	1.5	1.6	2.3	3.9
2004.....	1.8	1.8	3.2	0.9	2.4	1.6	1.7	1.8	1.5	1.7	2.2	2.9
2003.....	1.8	1.7	3.7	0.9	2.5	1.6	1.8	1.6	1.6	1.7	2.0	3.3
2002.....	1.9	1.9	3.9	1.2	2.8	1.9	2.0	1.5	1.3	1.6	2.1	3.6
2001.....	1.8	1.5	3.7	1.1	2.7	1.6	1.7	1.6	1.3	1.5	2.0	3.5
2000.....	2.0	2.2	4.0	1.3	2.9	1.8	1.8	1.6	1.5	1.6	1.8	2.8
1999.....	2.0	2.0	4.0	1.3	3.0	2.0	1.8	1.7	1.5	1.6	2.3	2.3

All races 3/, female

2006.....	0.5	1.2	2.1	0.3	0.3	0.2	0.4	0.5	0.5	0.7	0.8	0.9
2005.....	0.5	1.8	2.0	0.3	0.3	0.3	0.4	0.5	0.3	0.6	0.8	0.9
2004.....	0.5	1.2	2.1	0.4	0.3	0.3	0.3	0.4	0.4	0.5	0.6	1.0
2003.....	0.5	1.2	2.0	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.8	1.1
2002.....	0.5	1.2	1.9	0.4	0.3	0.3	0.4	0.4	0.4	0.4	0.8	0.9
2001.....	0.5	1.9	2.2	0.5	0.3	0.2	0.4	0.3	0.4	0.3	0.7	1.0
2000.....	0.5	1.7	2.4	0.5	0.3	0.3	0.3	0.3	0.3	0.5	0.9	1.0
1999.....	0.5	1.6	2.4	0.5	0.3	0.2	0.3	0.3	0.4	0.6	0.6	1.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental drowning and submersion (W65-W74)

White, both sexes

2006 4/.....	1.2	1.4	2.9	0.5	1.4	1.1	1.1	1.1	1.0	1.1	1.5	1.4
2005 4/.....	1.2	1.8	3.3	0.5	1.4	1.0	1.1	1.2	0.9	1.0	1.3	1.7
2004 4/.....	1.1	1.6	3.0	0.4	1.2	0.9	1.0	1.1	0.9	1.0	1.2	1.5
2003 4/.....	1.1	1.4	3.1	0.5	1.3	0.9	1.0	1.0	0.9	1.1	1.3	1.6
2002.....	1.1	1.6	3.0	0.5	1.4	1.0	1.1	0.9	0.8	0.9	1.3	1.6
2001.....	1.1	1.7	3.3	0.5	1.3	0.9	1.0	0.9	0.8	0.9	1.2	1.7
2000.....	1.2	1.6	3.5	0.7	1.5	1.0	1.0	0.9	0.9	1.0	1.2	1.5
1999.....	1.2	1.5	3.5	0.6	1.5	1.0	1.1	1.0	0.9	1.0	1.3	1.6

White, male

2006 4/.....	1.8	1.4	3.6	0.7	2.3	1.9	1.7	1.8	1.5	1.8	2.5	2.6
2005 4/.....	1.8	1.3	4.5	0.6	2.5	1.6	1.8	1.8	1.4	1.5	2.2	3.9
2004 4/.....	1.7	1.8	3.5	0.6	2.2	1.6	1.6	1.7	1.4	1.6	2.1	2.8
2003 4/.....	1.8	1.5	4.0	0.7	2.3	1.5	1.7	1.5	1.5	1.8	2.1	3.3
2002.....	1.8	1.9	4.0	0.7	2.5	1.8	1.9	1.4	1.2	1.5	2.1	3.7
2001.....	1.7	1.6	4.2	0.7	2.4	1.4	1.6	1.5	1.2	1.5	2.0	3.5
2000.....	1.9	1.8	4.4	0.9	2.6	1.7	1.7	1.5	1.6	1.6	1.8	2.7
1999.....	1.9	1.7	4.3	0.8	2.7	1.8	1.8	1.7	1.4	1.4	2.2	2.4

White, female

2006 4/.....	0.5	1.4	2.3	0.3	0.4	0.2	0.5	0.5	0.5	0.6	0.8	0.8
2005 4/.....	0.5	2.2	2.1	0.3	0.3	0.3	0.4	0.5	0.3	0.6	0.7	0.8
2004 4/.....	0.5	1.4	2.4	0.3	0.2	0.3	0.3	0.4	0.4	0.4	0.5	0.9
2003 4/.....	0.5	1.3	2.2	0.3	0.2	0.2	0.3	0.4	0.4	0.6	0.7	0.9
2002.....	0.4	*	2.0	0.3	0.2	0.3	0.4	0.3	0.4	0.3	0.7	0.7
2001.....	0.5	1.8	2.4	0.4	0.3	0.3	0.4	0.3	0.4	0.3	0.7	1.0
2000.....	0.5	*	2.5	0.4	0.3	0.2	0.3	0.3	0.3	0.5	0.8	1.0
1999.....	0.5	1.4	2.6	0.3	0.3	0.2	0.4	0.3	0.4	0.5	0.6	1.3



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

All races 3/, both sexes

2006.....	1.0	0.7	1.2	0.4	0.4	0.5	0.7	1.1	1.5	2.1	3.4	4.3
2005.....	1.1	0.8	1.3	0.5	0.4	0.6	0.8	1.2	1.3	2.1	3.6	5.0
2004.....	1.1	0.6	1.4	0.6	0.4	0.6	0.7	1.2	1.4	1.9	3.5	5.6
2003.....	1.2	0.7	1.4	0.5	0.5	0.6	0.9	1.3	1.4	2.3	3.6	5.4
2002.....	1.1	0.9	1.4	0.6	0.5	0.7	0.8	1.1	1.3	2.1	3.4	4.8
2001.....	1.2	1.2	1.5	0.6	0.5	0.6	1.0	1.1	1.5	2.2	3.4	5.7
2000.....	1.2	1.0	1.9	0.6	0.5	0.6	0.9	1.2	1.5	2.2	3.8	6.2
1999.....	1.2	1.1	2.0	0.6	0.5	0.7	0.9	1.0	1.3	2.3	4.0	5.9

All races 3/, male

2006.....	1.2	*	1.4	0.4	0.5	0.6	0.8	1.4	1.9	2.6	4.5	6.3
2005.....	1.3	*	1.5	0.6	0.5	0.7	1.0	1.6	1.7	2.6	4.6	7.0
2004.....	1.3	*	1.5	0.7	0.5	0.8	0.9	1.6	1.8	2.5	4.5	7.2
2003.....	1.4	*	1.6	0.5	0.6	0.8	1.2	1.6	1.9	2.9	4.4	7.4
2002.....	1.4	1.0	1.6	0.7	0.6	0.8	1.1	1.5	1.7	2.9	4.5	7.0
2001.....	1.4	1.5	1.8	0.7	0.7	0.8	1.4	1.5	1.9	2.8	4.1	8.3
2000.....	1.5	1.0	2.2	0.7	0.6	0.8	1.2	1.6	2.1	2.8	5.0	9.4
1999.....	1.5	1.1	2.2	0.7	0.7	1.0	1.3	1.4	1.7	2.9	5.1	9.3

All races 3/, female

2006.....	0.8	*	1.0	0.5	0.3	0.3	0.5	0.8	1.0	1.7	2.7	3.4
2005.....	0.9	*	1.0	0.5	0.3	0.4	0.5	0.8	1.0	1.6	2.9	4.0
2004.....	0.9	*	1.3	0.5	0.4	0.4	0.5	0.8	1.1	1.4	2.9	4.9
2003.....	1.0	1.1	1.2	0.5	0.4	0.5	0.6	0.9	1.0	1.8	3.0	4.5
2002.....	0.8	*	1.2	0.5	0.3	0.5	0.5	0.8	0.9	1.4	2.6	3.9
2001.....	0.9	1.0	1.1	0.5	0.4	0.4	0.6	0.7	1.1	1.8	2.9	4.6
2000.....	0.9	*	1.5	0.6	0.4	0.4	0.6	0.7	1.0	1.7	3.0	4.9
1999.....	0.9	1.1	1.7	0.6	0.4	0.4	0.5	0.6	1.0	1.7	3.2	4.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

White, both sexes

2006 4/.....	1.0	*	0.9	0.4	0.4	0.5	0.6	1.0	1.4	2.0	3.0	3.7
2005 4/.....	1.0	0.7	1.0	0.4	0.4	0.5	0.7	1.1	1.1	1.9	3.2	4.2
2004 4/.....	1.0	*	1.1	0.4	0.4	0.6	0.7	1.0	1.2	1.6	3.1	4.9
2003 4/.....	1.1	0.7	1.2	0.4	0.5	0.6	0.9	1.2	1.3	2.0	3.1	4.7
2002.....	1.0	*	1.0	0.4	0.4	0.6	0.8	1.0	1.1	1.8	3.0	4.0
2001.....	1.0	1.1	1.2	0.5	0.5	0.6	0.9	0.9	1.3	1.9	2.8	4.8
2000.....	1.0	0.8	1.3	0.5	0.5	0.5	0.8	1.0	1.3	1.8	3.1	5.1
1999.....	1.0	0.9	1.5	0.5	0.5	0.7	0.8	0.9	1.2	1.8	3.3	4.7

White, male

2006 4/.....	1.2	*	1.2	0.3	0.5	0.6	0.8	1.3	1.8	2.4	3.9	6.0
2005 4/.....	1.2	*	1.3	0.5	0.5	0.6	1.0	1.4	1.3	2.3	4.1	6.0
2004 4/.....	1.2	*	1.2	0.5	0.5	0.8	0.9	1.4	1.6	2.1	3.9	6.5
2003 4/.....	1.3	*	1.4	0.4	0.6	0.7	1.2	1.5	1.7	2.6	3.6	6.7
2002.....	1.2	*	1.1	0.6	0.6	0.8	1.1	1.3	1.5	2.5	4.1	5.8
2001.....	1.3	*	1.6	0.6	0.7	0.7	1.2	1.2	1.6	2.2	3.4	7.0
2000.....	1.3	*	1.6	0.5	0.6	0.7	1.1	1.5	1.7	2.3	4.1	8.2
1999.....	1.3	*	1.7	0.5	0.7	1.0	1.2	1.2	1.5	2.3	4.2	8.0

White, female

2006 4/.....	0.8	*	0.7	0.4	0.3	0.3	0.5	0.8	1.0	1.6	2.4	2.7
2005 4/.....	0.8	*	0.7	0.3	0.3	0.4	0.5	0.8	0.9	1.5	2.5	3.4
2004 4/.....	0.8	*	1.0	0.4	0.4	0.4	0.5	0.7	0.9	1.2	2.5	4.2
2003 4/.....	0.9	*	1.0	0.4	0.4	0.5	0.6	0.8	0.9	1.5	2.7	3.8
2002.....	0.7	*	1.0	0.3	0.3	0.4	0.5	0.7	0.7	1.3	2.2	3.2
2001.....	0.8	*	0.8	0.4	0.3	0.4	0.5	0.6	0.9	1.6	2.5	3.8
2000.....	0.8	*	1.1	0.5	0.3	0.4	0.5	0.6	0.8	1.5	2.4	3.9
1999.....	0.8	*	1.3	0.5	0.3	0.4	0.5	0.5	0.8	1.4	2.6	3.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental exposure to smoke, fire and flames (X00-X09)

Black, both sexes

2006 4/.....	1.8	*	3.0	1.0	0.5	0.9	1.1	1.9	2.7	4.2	9.4	12.2
2005 4/.....	1.9	*	2.8	1.2	0.5	0.9	1.1	1.9	3.5	4.3	9.3	14.4
2004 4/.....	2.1	*	2.8	1.6	0.6	1.0	1.2	2.4	3.4	4.8	9.7	14.9
2003 4/.....	2.0	*	3.0	1.4	0.6	0.9	1.2	2.6	2.9	5.3	10.2	14.9
2002.....	2.1	*	3.5	1.4	0.8	1.1	1.0	2.1	3.3	5.0	8.6	15.7
2001.....	2.3	*	3.3	1.3	0.7	1.2	1.8	2.4	3.5	6.1	9.7	17.8
2000.....	2.5	*	4.8	1.5	0.7	1.2	1.7	2.4	4.2	5.9	13.2	20.3
1999.....	2.5	*	4.5	1.5	0.7	1.0	1.8	2.1	3.4	6.9	13.2	21.7

Black, male

2006 4/.....	2.1	*	3.0	0.9	*	1.2	1.4	2.5	3.9	5.7	13.8	*
2005 4/.....	2.4	*	2.8	1.3	0.6	1.0	1.7	3.0	5.3	6.2	13.4	22.3
2004 4/.....	2.6	*	2.9	1.9	0.7	1.3	1.7	3.8	4.4	7.5	13.4	*
2003 4/.....	2.4	*	3.2	1.5	*	1.1	1.4	3.3	4.3	7.5	16.1	*
2002.....	2.6	*	4.1	1.6	1.0	1.4	1.7	3.0	4.7	7.8	10.9	25.7
2001.....	2.9	*	3.6	1.6	0.8	1.4	2.7	3.9	5.0	9.0	12.9	27.2
2000.....	3.2	*	5.6	1.7	0.9	1.5	2.4	3.5	6.3	8.8	18.5	27.7
1999.....	3.0	*	5.0	1.7	0.8	1.2	2.7	3.2	4.1	10.1	18.1	29.3

Black, female

2006 4/.....	1.5	*	2.9	1.1	*	*	0.7	1.3	1.7	3.1	6.9	11.7
2005 4/.....	1.5	*	2.7	1.2	*	0.9	*	1.0	2.0	2.9	7.0	11.3
2004 4/.....	1.6	*	2.7	1.2	*	*	0.7	1.2	2.6	3.0	7.6	13.4
2003 4/.....	1.7	*	2.7	1.3	0.6	*	1.0	2.0	1.7	3.7	6.9	13.4
2002.....	1.6	*	2.8	1.3	*	0.9	*	1.4	2.2	3.0	7.3	11.9
2001.....	1.7	*	2.9	1.0	*	1.1	1.0	1.1	2.4	4.1	7.9	14.3
2000.....	2.0	*	4.1	1.3	*	0.9	1.1	1.4	2.4	3.9	10.2	17.6
1999.....	2.0	*	3.9	1.3	*	0.8	1.0	1.3	2.7	4.6	10.5	18.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/ 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes

2006.....	9.2	*	0.2	0.1	6.9	13.0	17.3	19.0	7.6	2.6	2.6	3.8
2005.....	8.0	0.5	0.1	0.1	5.9	10.9	15.3	16.4	6.6	2.3	2.4	3.5
2004.....	7.1	*	*	0.1	5.4	9.1	14.6	14.5	5.4	2.3	2.4	3.4
2003.....	6.7	0.5	0.3	0.1	4.9	8.6	14.0	13.3	4.9	2.2	2.2	3.5
2002.....	6.1	0.6	0.2	0.1	4.1	7.8	13.4	11.7	4.1	2.1	2.3	3.9
2001.....	4.9	*	0.2	0.1	3.4	6.3	11.2	9.1	3.2	1.6	2.2	3.5
2000.....	4.5	*	0.2	0.1	3.0	6.0	10.3	8.1	2.8	1.5	2.0	4.3
1999.....	4.4	*	0.2	0.1	2.5	5.9	10.1	7.8	2.8	1.6	2.2	3.4

All races 3/, male

2006.....	12.6	*	*	0.2	10.6	18.9	23.0	24.9	9.9	2.9	2.9	4.6
2005.....	10.9	*	*	0.1	9.0	15.6	20.2	21.7	8.7	2.7	3.0	4.6
2004.....	9.6	*	*	0.1	8.2	13.1	19.0	19.1	6.6	2.6	2.7	4.2
2003.....	9.2	*	0.3	0.1	7.4	12.4	18.8	18.0	6.3	2.7	2.8	4.0
2002.....	8.5	1.0	*	0.1	6.3	11.5	18.4	15.9	5.4	2.5	2.7	5.1
2001.....	7.1	*	0.3	0.1	5.3	9.3	15.7	13.0	4.1	2.0	2.7	3.8
2000.....	6.6	*	0.3	0.1	4.6	9.0	15.1	12.0	3.4	1.8	2.1	6.2
1999.....	6.5	*	*	0.1	3.8	9.0	14.9	11.7	3.9	2.1	2.4	4.4

All races 3/, female

2006.....	5.9	*	*	0.1	3.0	6.9	11.6	13.4	5.5	2.3	2.4	3.4
2005.....	5.1	*	*	0.1	2.7	6.0	10.5	11.3	4.7	2.0	2.0	3.0
2004.....	4.7	*	*	0.1	2.4	4.9	10.3	10.0	4.3	2.0	2.2	3.0
2003.....	4.3	*	0.3	0.2	2.2	4.7	9.3	8.8	3.7	1.8	1.8	3.3
2002.....	3.7	*	*	*	1.9	4.0	8.4	7.6	3.0	1.7	2.1	3.4
2001.....	2.9	*	*	0.1	1.4	3.3	6.7	5.2	2.3	1.2	1.9	3.3
2000.....	2.5	*	*	*	1.3	2.8	5.6	4.4	2.3	1.3	1.9	3.6
1999.....	2.3	*	*	*	1.1	2.6	5.3	4.0	1.8	1.3	2.0	3.0



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

White, both sexes

2006 4/.....	9.7	*	*	0.1	8.3	14.9	18.3	19.0	7.1	2.3	2.6	3.8
2005 4/.....	8.4	*	*	0.1	7.0	12.3	16.4	16.4	6.1	2.1	2.4	3.6
2004 4/.....	7.6	*	*	0.1	6.4	10.3	15.7	14.6	5.0	2.2	2.4	3.4
2003 4/.....	7.1	*	0.3	0.1	5.8	9.7	14.8	13.3	4.8	2.0	2.3	3.5
2002.....	6.3	*	*	0.1	4.9	8.7	14.0	11.3	3.8	2.1	2.3	3.9
2001.....	5.0	*	0.2	0.1	3.9	6.9	11.4	8.4	2.8	1.5	2.1	3.3
2000.....	4.6	*	*	0.1	3.4	6.4	10.4	7.5	2.6	1.4	2.1	4.1
1999.....	4.3	*	0.2	0.1	2.8	6.1	10.0	7.1	2.5	1.5	2.2	3.3

White, male

2006 4/.....	13.2	*	*	0.1	12.7	21.3	24.1	24.2	8.7	2.4	2.9	4.7
2005 4/.....	11.4	*	*	0.2	10.7	17.5	21.5	21.2	7.5	2.2	2.9	4.8
2004 4/.....	10.2	*	*	0.2	9.7	14.9	20.3	18.9	5.7	2.4	2.5	4.3
2003 4/.....	9.7	*	*	0.1	8.8	13.9	19.8	17.6	5.8	2.2	2.8	3.7
2002.....	8.8	*	*	0.2	7.4	12.7	19.2	15.2	4.5	2.3	2.6	5.2
2001.....	7.2	*	*	*	6.0	10.2	16.2	11.9	3.4	1.9	2.5	3.7
2000.....	6.7	*	*	*	5.3	9.7	15.2	10.9	2.9	1.6	2.1	5.9
1999.....	6.4	*	*	*	4.2	9.5	14.9	10.5	3.2	1.9	2.5	4.5

White, female

2006 4/.....	6.3	*	*	*	3.5	8.0	12.3	13.8	5.7	2.3	2.4	3.3
2005 4/.....	5.4	*	*	*	3.1	6.8	11.2	11.7	4.8	2.1	2.1	3.0
2004 4/.....	5.0	*	*	*	2.8	5.5	11.0	10.3	4.4	2.0	2.2	3.0
2003 4/.....	4.5	*	*	*	2.6	5.2	9.8	9.0	3.8	1.8	1.9	3.4
2002.....	3.9	*	*	*	2.2	4.5	8.7	7.5	3.0	1.8	2.1	3.4
2001.....	2.9	*	*	0.1	1.6	3.5	6.6	5.0	2.3	1.2	1.8	3.1
2000.....	2.5	*	*	*	1.3	3.0	5.5	4.1	2.3	1.3	2.0	3.4
1999.....	2.3	*	*	*	1.2	2.6	5.1	3.8	1.8	1.2	2.1	2.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Accidental poisoning and exposure to noxious substances (X40-X49)

Black, both sexes

2006 4/.....	8.9	*	*	*	2.1	8.0	17.4	25.5	14.3	4.5	2.7	5.4
2005 4/.....	7.6	*	*	*	1.8	7.2	14.4	21.8	13.0	4.4	3.0	*
2004 4/.....	6.5	*	*	*	1.8	5.3	12.8	18.8	9.8	3.7	3.5	*
2003 4/.....	6.3	*	*	*	1.4	5.7	13.5	17.9	7.6	4.1	2.0	*
2002.....	6.3	*	*	*	1.4	5.6	13.8	17.9	8.4	2.7	2.8	*
2001.....	5.8	*	*	*	1.6	5.2	13.1	16.7	6.5	2.5	3.6	6.4
2000.....	5.5	*	*	*	1.3	5.6	13.0	15.5	5.5	2.6	*	8.1
1999.....	5.7	*	*	*	1.6	6.1	13.1	15.5	6.0	3.0	*	*

Black, male

2006 4/.....	12.9	*	*	*	3.0	12.6	24.3	37.8	24.6	7.6	*	*
2005 4/.....	10.8	*	*	*	2.4	10.2	18.8	32.2	22.2	8.1	*	*
2004 4/.....	8.9	*	*	*	2.7	7.6	16.4	27.1	16.2	6.2	5.6	*
2003 4/.....	8.8	*	*	*	2.0	8.0	18.6	26.2	12.8	7.6	*	*
2002.....	9.1	*	*	*	2.2	8.5	19.3	26.5	15.0	5.0	*	*
2001.....	8.4	*	*	*	2.3	7.5	17.9	26.6	11.3	3.9	5.9	*
2000.....	8.2	*	*	*	1.7	8.4	18.7	24.9	9.5	4.5	*	*
1999.....	8.7	*	*	*	2.7	9.1	19.4	25.9	11.0	4.4	*	*

Black, female

2006 4/.....	5.3	*	*	*	1.3	3.8	11.4	14.9	6.1	2.4	*	*
2005 4/.....	4.8	*	*	*	1.2	4.3	10.4	12.9	5.5	*	*	*
2004 4/.....	4.3	*	*	*	1.0	3.3	9.7	11.8	4.7	*	*	*
2003 4/.....	3.9	*	*	*	0.7	3.6	8.9	10.8	3.3	*	*	*
2002.....	3.7	*	*	*	0.7	3.0	8.9	10.5	3.2	*	*	*
2001.....	3.5	*	*	*	0.9	3.1	9.0	8.2	2.6	*	*	*
2000.....	3.1	*	*	*	1.0	3.0	7.8	7.4	2.3	*	*	*
1999.....	3.0	*	*	*	*	3.4	7.6	6.5	2.0	2.1	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes

2006.....	5.8	21.5	1.6	0.6	1.6	2.1	2.9	4.2	5.2	9.1	25.9	89.3
2005.....	6.3	19.5	1.5	0.6	1.7	2.3	3.1	4.7	5.8	10.2	28.6	95.1
2004.....	5.8	19.2	1.6	0.6	1.5	2.1	2.9	4.2	5.0	8.8	27.0	95.5
2003.....	5.9	16.9	1.8	0.5	1.6	2.0	3.0	3.9	5.1	9.6	27.5	102.5
2002.....	6.0	16.8	1.6	0.7	1.6	2.1	3.0	3.7	5.2	9.8	28.8	105.6
2001.....	6.2	16.6	1.8	0.6	1.6	2.0	3.1	4.2	5.3	10.0	30.5	115.0
2000.....	6.2	15.0	1.7	0.6	1.7	2.2	3.2	4.0	5.6	9.9	29.8	117.1
1999.....	6.6	13.8	1.9	0.8	1.7	2.2	3.2	4.1	5.6	11.0	32.9	128.9

All races 3/, male

2006.....	7.0	25.2	1.7	0.8	2.6	3.5	4.8	6.8	7.7	12.7	32.9	99.4
2005.....	7.4	21.1	1.8	0.9	2.8	3.9	4.9	7.4	8.6	13.8	35.9	107.1
2004.....	6.8	21.7	2.0	0.9	2.6	3.4	4.6	6.6	7.5	11.8	33.6	109.8
2003.....	6.9	19.5	2.2	0.8	2.6	3.4	4.8	6.0	7.8	13.1	34.9	120.2
2002.....	6.9	18.8	2.0	0.9	2.6	3.4	4.8	6.0	7.9	13.4	36.0	118.6
2001.....	7.2	18.9	2.1	0.9	2.6	3.4	5.0	6.6	7.7	13.2	38.4	134.0
2000.....	7.2	16.8	2.1	0.9	2.8	3.8	5.4	6.6	8.3	13.0	36.9	135.2
1999.....	7.6	16.2	2.3	1.1	2.9	3.7	5.3	6.6	8.3	14.8	40.9	152.7

All races 3/, female

2006.....	4.7	17.6	1.5	0.3	0.5	0.7	1.0	1.8	2.8	6.1	21.2	84.5
2005.....	5.1	17.9	1.2	0.4	0.6	0.7	1.2	2.0	3.1	7.1	23.6	89.6
2004.....	4.8	16.5	1.1	0.3	0.4	0.7	1.2	1.8	2.7	6.4	22.6	89.1
2003.....	4.9	14.1	1.3	0.3	0.5	0.7	1.2	1.9	2.6	6.7	22.6	94.7
2002.....	5.0	14.6	1.2	0.4	0.5	0.6	1.2	1.5	2.7	6.8	24.0	100.0
2001.....	5.3	14.2	1.5	0.4	0.6	0.6	1.1	1.9	3.1	7.3	25.4	107.0
2000.....	5.2	13.2	1.4	0.4	0.5	0.6	1.1	1.5	3.0	7.4	25.1	109.8
1999.....	5.6	11.3	1.5	0.5	0.5	0.7	1.2	1.7	3.1	8.0	27.7	119.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

White, both sexes

2006 4/.....	6.2	16.9	1.5	0.6	1.7	2.3	3.0	4.2	5.1	8.9	26.5	93.2
2005 4/.....	6.5	16.6	1.5	0.7	1.8	2.5	3.1	4.4	5.5	9.8	28.3	97.4
2004 4/.....	6.2	16.0	1.5	0.6	1.7	2.2	3.0	4.3	4.9	8.7	27.2	98.8
2003 4/.....	6.3	13.6	1.7	0.5	1.7	2.2	3.1	3.8	4.9	9.4	27.8	106.3
2002.....	6.3	12.9	1.4	0.7	1.7	2.2	3.1	3.7	5.1	9.4	28.8	108.8
2001.....	6.6	13.4	1.7	0.6	1.8	2.1	3.1	4.1	5.1	9.7	30.6	117.7
2000.....	6.6	12.2	1.6	0.7	1.7	2.4	3.2	3.8	5.4	9.5	29.8	120.5
1999.....	6.9	11.2	1.8	0.8	1.9	2.3	3.3	3.9	5.2	10.6	32.5	132.4

White, male

2006 4/.....	7.4	19.6	1.6	0.8	2.8	3.8	4.9	6.6	7.6	12.4	33.6	103.5
2005 4/.....	7.7	17.8	1.9	0.9	2.9	4.2	5.0	7.1	8.1	13.3	35.2	110.0
2004 4/.....	7.2	19.0	2.0	0.9	2.8	3.6	4.7	6.7	7.3	11.6	33.7	112.9
2003 4/.....	7.3	15.7	2.2	0.8	2.8	3.6	4.9	5.8	7.4	12.9	34.9	123.9
2002.....	7.3	15.3	1.8	0.9	2.9	3.7	5.0	6.0	7.6	12.7	35.7	121.2
2001.....	7.5	15.2	2.0	0.9	2.9	3.5	5.1	6.4	7.4	12.9	38.0	137.0
2000.....	7.4	14.0	2.0	0.9	2.9	4.0	5.4	6.2	7.9	12.5	36.2	137.6
1999.....	7.8	12.9	2.1	1.1	3.1	3.8	5.4	6.2	7.7	14.0	40.1	155.2

White, female

2006 4/.....	5.1	14.1	1.4	0.3	0.5	0.7	1.1	1.8	2.8	5.9	21.6	88.3
2005 4/.....	5.4	15.3	1.1	0.4	0.6	0.7	1.1	1.8	3.1	6.8	23.6	91.6
2004 4/.....	5.2	12.8	0.9	0.3	0.5	0.7	1.3	1.9	2.7	6.2	22.8	92.5
2003 4/.....	5.3	11.4	1.2	0.3	0.5	0.7	1.2	1.8	2.5	6.5	23.0	98.5
2002.....	5.4	10.5	1.1	0.4	0.5	0.7	1.1	1.5	2.6	6.5	24.2	103.4
2001.....	5.8	11.5	1.4	0.4	0.6	0.6	1.1	1.9	3.0	7.1	25.7	109.7
2000.....	5.7	10.3	1.2	0.4	0.5	0.6	1.0	1.5	3.0	7.1	25.6	113.6
1999.....	6.1	9.5	1.4	0.5	0.6	0.7	1.2	1.6	2.8	7.7	27.4	123.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

Black, both sexes

2006 4/.....	5.0	44.7	2.2	0.7	1.1	1.8	3.0	5.4	6.6	12.6	24.6	61.7
2005 4/.....	6.0	37.4	1.8	0.6	1.6	2.0	3.3	6.9	8.7	15.5	37.9	86.1
2004 4/.....	4.8	37.1	1.8	0.6	1.0	1.9	2.9	4.6	6.5	11.7	30.6	70.2
2003 4/.....	4.9	34.9	2.0	0.5	1.3	1.8	3.2	5.4	7.3	12.8	27.6	73.2
2002.....	5.2	37.5	2.4	0.8	1.1	1.7	3.1	4.7	7.3	15.2	32.3	83.1
2001.....	5.4	34.2	2.5	0.6	1.0	1.9	3.2	5.6	7.9	14.1	33.8	96.1
2000.....	5.4	30.6	2.3	0.7	1.5	2.2	3.5	5.6	8.2	14.9	33.7	90.1
1999.....	6.1	27.9	2.5	0.7	1.3	2.4	3.4	6.6	10.0	17.9	42.9	105.5

Black, male

2006 4/.....	6.4	53.3	2.2	0.8	1.6	2.8	5.2	9.2	10.0	18.0	31.9	75.1
2005 4/.....	7.6	41.0	1.9	0.8	2.4	3.1	5.2	11.4	14.9	21.3	54.1	97.6
2004 4/.....	6.1	38.3	1.7	1.0	1.5	3.3	4.7	7.5	10.3	16.4	42.3	87.4
2003 4/.....	6.4	40.5	2.8	0.7	1.9	2.9	5.1	8.5	11.9	18.3	39.8	91.8
2002.....	6.6	38.9	3.3	1.1	1.7	2.9	4.7	7.4	12.1	21.6	45.6	106.1
2001.....	6.8	40.4	2.8	0.9	1.5	3.2	5.1	9.0	11.9	19.8	48.8	114.6
2000.....	7.3	32.0	2.5	1.0	2.5	3.7	6.0	10.1	13.8	20.3	53.0	124.5
1999.....	8.0	33.2	3.4	0.9	2.3	4.0	5.7	11.1	15.7	26.9	57.2	153.5

Black, female

2006 4/.....	3.6	35.8	2.3	*	*	1.0	1.1	2.1	3.9	8.8	20.5	56.3
2005 4/.....	4.5	33.6	1.6	*	0.7	1.1	1.7	3.1	3.7	11.3	28.7	81.4
2004 4/.....	3.6	35.8	2.0	*	*	*	1.2	2.2	3.5	8.4	23.9	63.4
2003 4/.....	3.6	29.1	*	*	0.7	0.7	1.5	2.8	3.6	8.8	20.7	66.0
2002.....	4.0	36.0	*	*	*	*	1.7	2.3	3.5	10.7	24.7	74.3
2001.....	4.1	27.7	2.2	*	*	0.7	1.6	2.7	4.6	10.0	25.3	89.3
2000.....	3.8	29.1	2.0	*	*	0.9	1.3	1.8	3.6	11.1	22.8	77.3
1999.....	4.4	22.5	1.6	*	*	1.0	1.4	2.7	5.4	11.6	34.9	87.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

All races 3/, both sexes

2006.....	11.1	...	...	0.5	9.9	12.3	15.1	17.2	14.5	12.6	15.9	15.9
2005.....	11.0	...	...	0.7	10.0	12.4	14.9	16.5	13.9	12.6	16.9	16.9
2004.....	11.0	...	...	0.7	10.3	12.7	15.0	16.6	13.8	12.3	16.3	16.4
2003.....	10.8	...	...	0.6	9.7	12.7	14.9	15.9	13.8	12.7	16.4	16.9
2002.....	11.0	...	...	0.6	9.9	12.6	15.3	15.7	13.6	13.5	17.7	18.0
2001 5/.....	10.8	...	...	0.7	9.9	12.8	14.7	15.2	13.1	13.3	17.4	17.5
2000.....	10.4	...	...	0.7	10.2	12.0	14.5	14.4	12.1	12.5	17.6	19.6
1999.....	10.5	...	...	0.6	10.1	12.7	14.3	13.9	12.2	13.4	18.1	19.3

All races 3/, male

2006.....	17.8	...	...	0.7	16.2	19.7	23.2	26.2	22.7	22.7	33.3	43.2
2005.....	17.7	...	...	1.0	16.2	19.9	23.1	25.2	22.2	22.7	35.8	45.0
2004.....	17.7	...	...	0.9	16.8	20.4	23.0	24.8	22.1	22.6	34.8	45.0
2003.....	17.6	...	...	0.9	16.0	20.6	23.2	24.4	22.3	23.4	35.1	47.8
2002.....	17.9	...	...	0.9	16.5	20.5	23.7	24.4	22.2	24.7	38.1	50.7
2001 5/.....	17.6	...	...	1.0	16.6	21.0	23.1	23.4	21.1	24.6	37.8	51.1
2000.....	17.1	...	...	1.2	17.1	19.6	22.8	22.4	19.4	22.7	38.6	57.5
1999.....	17.1	...	...	0.9	16.8	20.7	22.4	21.5	19.8	24.7	38.8	57.1

All races 3/, female

2006.....	4.6	...	...	0.3	3.2	4.7	7.0	8.4	6.8	4.1	4.0	3.1
2005.....	4.5	...	...	0.3	3.5	4.7	6.8	8.0	6.1	4.0	4.0	4.0
2004.....	4.6	...	...	0.5	3.6	4.7	7.1	8.6	6.1	3.8	3.9	3.6
2003.....	4.3	...	...	0.3	3.0	4.6	6.6	7.7	5.9	3.8	4.0	3.3
2002.....	4.3	...	...	0.3	2.9	4.6	6.9	7.4	5.7	4.1	4.2	3.8
2001 5/.....	4.1	...	...	0.3	2.9	4.4	6.4	7.2	5.7	3.9	4.0	3.4
2000.....	4.0	...	...	0.3	3.0	4.3	6.4	6.7	5.4	4.0	4.0	4.2
1999.....	4.0	...	...	0.3	3.0	4.6	6.4	6.6	5.2	4.1	4.7	4.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

White, both sexes

2006 4/.....	12.4	...	...	0.5	10.5	13.4	17.0	19.4	16.1	13.8	17.2	16.9
2005 4/.....	12.3	...	...	0.7	10.7	13.4	16.8	18.5	15.4	13.8	18.1	17.9
2004 4/.....	12.3	...	...	0.7	11.0	13.6	16.9	18.6	15.1	13.4	17.5	17.5
2003 4/.....	12.1	...	...	0.6	10.2	13.7	16.6	17.9	15.2	13.8	17.6	18.1
2002.....	12.2	...	...	0.7	10.7	13.6	17.0	17.6	15.0	14.7	19.0	19.1
2001 5/.....	11.9	...	...	0.7	10.6	13.8	16.4	16.9	14.3	14.3	18.6	18.6
2000.....	11.5	...	...	0.8	10.7	12.8	16.1	15.9	13.3	13.5	18.8	21.0
1999.....	11.5	...	...	0.6	10.5	13.5	15.8	15.3	13.2	14.4	19.3	20.7

White, male

2006 4/.....	19.8	...	...	0.7	17.1	21.1	25.7	29.2	25.0	24.7	36.0	46.1
2005 4/.....	19.7	...	...	1.0	17.3	21.2	25.6	28.1	24.5	24.9	38.4	48.2
2004 4/.....	19.6	...	...	0.9	17.9	21.8	25.6	27.6	23.9	24.2	37.1	48.4
2003 4/.....	19.5	...	...	0.9	16.9	21.8	25.7	27.3	24.3	25.2	37.5	51.4
2002.....	19.9	...	...	1.0	17.7	21.6	26.2	27.0	24.3	26.8	40.6	53.9
2001 5/.....	19.5	...	...	1.0	17.6	22.4	25.4	26.0	22.8	26.3	40.2	55.0
2000.....	18.8	...	...	1.2	17.9	20.7	24.9	24.5	21.2	24.3	41.1	61.6
1999.....	18.7	...	...	1.0	17.5	21.8	24.4	23.4	21.2	26.4	41.2	61.5

White, female

2006 4/.....	5.2	...	...	0.3	3.4	5.3	8.1	9.7	7.7	4.4	4.2	3.2
2005 4/.....	5.0	...	...	0.3	3.7	5.1	7.7	9.1	6.8	4.3	4.2	3.9
2004 4/.....	5.1	...	...	0.5	3.8	5.0	8.1	9.8	6.8	4.1	4.1	3.6
2003 4/.....	4.7	...	...	0.3	3.1	5.1	7.5	8.7	6.5	4.1	4.2	3.3
2002.....	4.8	...	...	0.4	3.1	5.2	7.8	8.4	6.2	4.4	4.4	4.0
2001 5/.....	4.6	...	...	0.3	3.1	4.9	7.3	8.0	6.4	4.2	4.3	3.4
2000.....	4.4	...	...	0.3	3.1	4.6	7.2	7.5	6.0	4.4	4.1	4.5
1999.....	4.5	...	...	0.3	3.2	4.9	7.2	7.4	5.7	4.4	4.9	4.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

Black, both sexes

2006 4/.....	4.9	...	...	0.6	6.2	8.4	7.3	6.0	4.8	4.2	4.4	5.7
2005 4/.....	5.1	...	...	0.6	6.7	8.7	7.3	6.2	4.7	4.5	5.5	*
2004 4/.....	5.2	...	...	0.7	7.2	9.4	6.6	5.8	5.7	4.4	6.0	*
2003 4/.....	5.1	...	...	0.6	7.1	9.3	7.2	5.6	5.0	4.5	4.5	*
2002.....	5.1	...	...	0.5	6.5	9.3	7.6	5.9	4.8	4.9	5.5	5.9
2001 5/.....	5.3	...	...	0.6	7.1	9.1	7.3	6.2	5.2	5.5	5.7	*
2000.....	5.4	...	...	0.6	8.1	8.7	7.6	6.5	4.2	5.1	5.8	*
1999.....	5.4	...	...	0.4	8.1	9.2	7.6	5.7	4.9	5.8	5.7	*

Black, male

2006 4/.....	8.8	...	...	0.9	10.6	15.5	13.1	10.5	8.9	8.8	11.6	*
2005 4/.....	8.7	...	...	0.8	11.5	14.9	12.5	10.2	8.0	8.4	12.8	*
2004 4/.....	9.0	...	...	0.9	12.2	16.2	11.3	9.7	10.8	9.8	15.0	*
2003 4/.....	8.8	...	...	1.0	12.1	16.5	12.2	9.1	8.7	8.3	11.3	*
2002.....	9.1	...	...	0.8	11.3	17.3	13.0	10.2	8.4	9.7	13.8	*
2001 5/.....	9.2	...	...	0.9	13.0	16.5	12.3	9.9	9.4	10.7	13.5	*
2000.....	9.4	...	...	1.0	14.2	15.5	13.2	11.2	7.5	11.1	12.1	*
1999.....	9.6	...	...	0.6	14.3	16.8	13.3	10.0	9.1	11.8	13.8	*

Black, female

2006 4/.....	1.4	...	...	*	1.8	1.8	2.2	2.1	1.6	*	*	*
2005 4/.....	1.8	...	...	*	1.7	2.9	2.7	2.7	2.1	*	*	*
2004 4/.....	1.8	...	...	*	2.2	3.2	2.5	2.5	1.6	*	*	*
2003 4/.....	1.8	...	...	*	2.0	2.7	2.9	2.6	2.0	*	*	*
2002.....	1.5	...	...	*	1.7	2.0	2.9	2.2	1.9	*	*	*
2001 5/.....	1.7	...	...	*	1.3	2.4	2.9	3.0	1.8	*	*	*
2000.....	1.7	...	...	*	2.2	2.6	2.7	2.5	1.6	*	*	*
1999.....	1.6	...	...	*	1.9	2.3	2.6	2.0	1.6	*	*	*



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes

2006.....	5.6	...	...	0.2	4.7	5.3	6.5	8.0	8.1	8.9	12.0	10.9
2005.....	5.7	...	...	0.2	4.7	5.7	6.5	8.2	8.1	9.0	12.6	11.2
2004.....	5.7	...	...	0.1	5.0	5.7	6.5	8.0	8.0	8.8	12.2	11.1
2003.....	5.8	...	...	0.2	5.0	6.0	6.6	8.0	8.3	9.3	12.4	11.9
2002.....	5.9	...	...	0.2	5.1	6.0	7.0	7.8	8.4	9.7	13.2	12.0
2001.....	5.9	...	...	0.2	5.3	6.5	6.7	7.7	8.2	9.6	13.1	12.2
2000.....	5.9	...	...	0.3	5.8	6.1	7.1	7.6	7.6	9.2	13.0	13.2
1999.....	5.9	...	...	0.3	6.0	6.5	6.9	7.3	7.8	9.7	13.1	12.6

All races 3/, male

2006.....	10.0	...	...	0.3	8.2	9.2	10.8	13.7	14.3	17.6	27.7	32.7
2005.....	10.2	...	...	0.3	8.2	9.7	11.0	14.1	14.6	17.6	29.3	33.8
2004.....	10.0	...	...	0.2	8.8	9.7	10.8	13.5	14.2	17.5	28.3	34.4
2003.....	10.4	...	...	0.3	8.9	10.4	11.2	13.8	14.9	18.4	28.7	36.6
2002.....	10.6	...	...	0.3	9.1	10.3	11.9	13.6	15.1	19.5	31.0	37.8
2001.....	10.6	...	...	0.3	9.4	11.2	11.4	13.3	14.6	19.2	30.7	38.9
2000.....	10.5	...	...	0.4	10.2	10.6	11.8	13.0	13.5	18.4	30.8	42.9
1999.....	10.6	...	...	0.4	10.5	11.2	11.5	12.5	14.0	19.5	31.1	41.6

All races 3/, female

2006.....	1.4	...	...	*	0.9	1.3	2.1	2.6	2.3	1.6	1.2	0.7
2005.....	1.4	...	...	*	0.9	1.5	2.0	2.4	2.1	1.7	1.3	0.8
2004.....	1.5	...	...	*	1.1	1.5	2.2	2.8	2.2	1.6	1.4	*
2003.....	1.4	...	...	*	1.0	1.4	2.1	2.5	2.2	1.6	1.5	0.9
2002.....	1.4	...	...	*	1.0	1.6	2.2	2.2	2.2	1.6	1.4	0.8
2001.....	1.5	...	...	0.1	1.0	1.6	2.1	2.4	2.4	1.7	1.5	1.0
2000.....	1.5	...	...	0.1	1.2	1.7	2.4	2.3	2.1	1.7	1.4	1.1
1999.....	1.5	...	...	0.1	1.2	1.8	2.3	2.3	2.1	1.7	1.4	0.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

White, both sexes

2006 4/.....	6.4	...	...	0.2	4.9	5.7	7.4	9.3	9.2	10.0	13.1	11.7
2005 4/.....	6.5	...	...	0.2	5.0	6.2	7.4	9.3	9.2	10.0	13.8	12.2
2004 4/.....	6.5	...	...	0.2	5.4	6.1	7.4	9.2	9.0	9.7	13.3	12.1
2003 4/.....	6.6	...	...	0.2	5.3	6.4	7.6	9.2	9.3	10.3	13.4	12.9
2002.....	6.7	...	...	0.2	5.5	6.5	7.9	8.9	9.4	10.8	14.2	12.9
2001.....	6.7	...	...	0.2	5.6	7.1	7.5	8.7	9.2	10.5	14.1	13.2
2000.....	6.6	...	...	0.3	6.1	6.6	7.9	8.5	8.3	10.1	14.1	14.3
1999.....	6.6	...	...	0.3	6.2	7.0	7.7	8.2	8.6	10.6	14.1	13.6

White, male

2006 4/.....	11.3	...	...	0.3	8.7	9.7	12.1	15.5	16.0	19.4	30.2	35.1
2005 4/.....	11.5	...	...	0.4	8.7	10.4	12.3	15.9	16.4	19.5	31.7	36.8
2004 4/.....	11.3	...	...	0.3	9.4	10.3	12.1	15.3	15.8	19.1	30.4	37.5
2003 4/.....	11.7	...	...	0.3	9.4	10.9	12.6	15.6	16.5	20.2	30.9	39.9
2002.....	11.9	...	...	0.4	9.6	10.8	13.3	15.2	16.7	21.4	33.3	40.5
2001.....	11.8	...	...	0.4	9.7	12.0	12.6	14.9	16.1	20.9	32.9	42.1
2000.....	11.6	...	...	0.5	10.6	11.2	13.1	14.4	14.7	19.8	33.3	46.5
1999.....	11.7	...	...	0.4	10.8	11.8	12.6	13.9	15.2	21.0	33.3	44.9

White, female

2006 4/.....	1.6	...	...	*	0.9	1.5	2.5	3.1	2.7	1.9	1.3	0.7
2005 4/.....	1.6	...	...	*	1.0	1.7	2.4	2.9	2.4	1.8	1.4	0.9
2004 4/.....	1.7	...	...	*	1.1	1.7	2.6	3.2	2.5	1.8	1.6	*
2003 4/.....	1.6	...	...	*	1.0	1.7	2.4	2.9	2.5	1.8	1.6	0.9
2002.....	1.6	...	...	*	1.0	1.9	2.5	2.6	2.5	1.7	1.5	0.9
2001.....	1.7	...	...	*	1.1	1.8	2.4	2.7	2.7	1.8	1.6	1.1
2000.....	1.7	...	...	*	1.2	1.8	2.7	2.6	2.4	1.9	1.5	1.2
1999.....	1.7	...	...	0.1	1.3	2.0	2.6	2.6	2.4	1.9	1.5	1.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

Black, both sexes

2006 4/.....	2.6	...	...	*	3.7	4.6	3.2	2.5	2.4	2.9	3.8	*
2005 4/.....	2.6	...	...	*	3.8	4.4	3.3	2.7	2.4	3.2	4.1	*
2004 4/.....	2.6	...	...	*	4.0	4.8	2.7	2.4	3.0	3.5	4.8	*
2003 4/.....	2.6	...	...	*	4.3	5.1	2.6	2.5	2.7	3.3	4.0	*
2002.....	2.8	...	...	*	4.1	5.0	3.5	2.9	2.8	3.3	4.4	*
2001.....	2.9	...	...	*	4.9	4.9	3.4	2.8	2.8	3.7	5.2	*
2000.....	3.0	...	...	*	5.2	5.0	3.4	3.3	2.9	3.7	3.5	*
1999.....	3.1	...	...	*	5.6	5.0	3.5	3.0	3.4	4.0	4.7	*

Black, male

2006 4/.....	4.9	...	...	*	6.7	8.7	6.3	4.7	4.9	6.5	10.3	*
2005 4/.....	4.9	...	...	*	6.9	8.3	6.4	5.2	4.5	6.4	10.7	*
2004 4/.....	4.9	...	...	*	7.1	9.0	5.2	4.4	5.8	8.3	12.8	*
2003 4/.....	5.0	...	...	*	7.8	10.0	4.8	4.7	5.1	6.6	9.9	*
2002.....	5.4	...	...	*	7.5	10.0	6.3	5.5	5.5	7.1	11.5	*
2001.....	5.5	...	...	*	9.2	9.4	6.3	4.8	5.7	7.7	12.4	*
2000.....	5.5	...	...	*	9.6	9.3	6.2	5.8	5.7	8.8	8.2	*
1999.....	5.9	...	...	*	10.3	9.6	6.7	5.6	7.0	8.7	12.0	*

Black, female

2006 4/.....	0.4	...	...	*	0.7	0.8	*	*	*	*	*	*
2005 4/.....	0.5	...	...	*	*	0.9	*	*	*	*	*	*
2004 4/.....	0.5	...	...	*	0.9	0.9	*	*	*	*	*	*
2003 4/.....	0.5	...	...	*	0.7	*	0.7	*	*	*	*	*
2002.....	0.5	...	...	*	0.7	*	0.9	*	*	*	*	*
2001.....	0.6	...	...	*	*	0.9	0.9	1.0	*	*	*	*
2000.....	0.7	...	...	*	0.9	1.1	0.8	1.1	*	*	*	*
1999.....	0.6	...	...	*	1.0	0.8	0.7	*	*	*	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

All races 3/, both sexes

2006.....	5.5	...	...	0.4	5.2	7.0	8.6	9.1	6.4	3.7	3.9	5.0
2005.....	5.3	...	...	0.5	5.3	6.8	8.4	8.3	5.7	3.6	4.2	5.7
2004.....	5.3	...	...	0.6	5.3	7.0	8.5	8.5	5.8	3.5	4.1	5.4
2003.....	5.0	...	...	0.4	4.6	6.7	8.3	7.8	5.5	3.5	4.0	5.1
2002.....	5.0	...	...	0.4	4.7	6.6	8.2	7.9	5.2	3.8	4.5	6.0
2001 5/.....	4.8	...	...	0.5	4.6	6.3	8.0	7.4	4.9	3.7	4.3	5.3
2000.....	4.5	...	...	0.5	4.4	5.9	7.5	6.8	4.6	3.2	4.6	6.5
1999.....	4.5	...	...	0.3	4.1	6.2	7.5	6.6	4.4	3.7	5.0	6.7

All races 3/, male

2006.....	7.8	...	...	0.5	7.9	10.5	12.4	12.5	8.4	5.1	5.6	10.5
2005.....	7.5	...	...	0.6	8.0	10.2	12.1	11.1	7.6	5.1	6.5	11.2
2004.....	7.6	...	...	0.7	8.0	10.6	12.2	11.3	7.8	5.1	6.5	10.7
2003.....	7.3	...	...	0.6	7.1	10.2	12.1	10.6	7.3	5.0	6.4	11.1
2002.....	7.3	...	...	0.6	7.4	10.2	11.8	10.8	7.1	5.2	7.1	12.9
2001 5/.....	7.1	...	...	0.7	7.2	9.8	11.7	10.2	6.5	5.4	7.1	12.2
2000.....	6.6	...	...	0.7	6.9	9.0	10.9	9.4	6.0	4.3	7.8	14.6
1999.....	6.6	...	...	0.5	6.3	9.5	10.9	8.9	5.8	5.3	7.7	15.5

All races 3/, female

2006.....	3.2	...	...	0.3	2.3	3.4	4.9	5.8	4.5	2.4	2.8	2.4
2005.....	3.1	...	...	0.3	2.6	3.2	4.8	5.5	4.0	2.3	2.7	3.1
2004.....	3.1	...	...	0.4	2.5	3.2	4.9	5.9	3.9	2.2	2.5	3.0
2003.....	2.8	...	...	0.2	2.1	3.2	4.5	5.2	3.7	2.2	2.5	2.4
2002.....	2.9	...	...	0.2	2.0	3.0	4.7	5.2	3.5	2.5	2.8	3.0
2001 5/.....	2.6	...	...	0.2	1.9	2.8	4.3	4.8	3.4	2.3	2.5	2.4
2000.....	2.5	...	...	0.2	1.8	2.7	4.0	4.4	3.3	2.3	2.6	3.2
1999.....	2.5	...	...	0.1	1.8	2.8	4.1	4.3	3.0	2.4	3.3	3.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

White, both sexes

2006 4/.....	6.0	...	...	0.4	5.6	7.7	9.6	10.1	7.0	3.8	4.1	5.3
2005 4/.....	5.8	...	...	0.4	5.8	7.3	9.4	9.2	6.2	3.8	4.4	5.6
2004 4/.....	5.8	...	...	0.5	5.6	7.5	9.5	9.4	6.2	3.6	4.2	5.5
2003 4/.....	5.5	...	...	0.4	4.9	7.3	9.1	8.7	5.9	3.6	4.2	5.2
2002.....	5.5	...	...	0.4	5.2	7.1	9.1	8.8	5.6	3.9	4.7	6.2
2001 5/.....	5.3	...	...	0.5	5.0	6.8	8.8	8.2	5.2	3.8	4.5	5.5
2000.....	4.9	...	...	0.5	4.6	6.2	8.1	7.5	5.0	3.4	4.6	6.7
1999.....	4.9	...	...	0.4	4.3	6.5	8.2	7.2	4.6	3.8	5.2	7.1

White, male

2006 4/.....	8.5	...	...	0.4	8.5	11.4	13.6	13.7	9.0	5.3	5.8	11.0
2005 4/.....	8.2	...	...	0.6	8.6	10.8	13.3	12.2	8.1	5.4	6.7	11.4
2004 4/.....	8.3	...	...	0.6	8.4	11.4	13.5	12.3	8.2	5.2	6.6	10.9
2003 4/.....	7.9	...	...	0.6	7.5	10.9	13.1	11.7	7.7	5.0	6.6	11.5
2002.....	8.0	...	...	0.6	8.1	10.7	12.9	11.8	7.6	5.4	7.4	13.4
2001 5/.....	7.7	...	...	0.7	7.9	10.4	12.8	11.0	6.8	5.4	7.3	12.9
2000.....	7.1	...	...	0.7	7.3	9.5	11.8	10.0	6.5	4.5	7.8	15.2
1999.....	7.0	...	...	0.6	6.6	9.9	11.8	9.6	6.1	5.4	7.9	16.6

White, female

2006 4/.....	3.6	...	...	0.3	2.4	3.8	5.6	6.6	5.1	2.5	2.9	2.6
2005 4/.....	3.4	...	...	0.3	2.7	3.4	5.3	6.2	4.4	2.4	2.8	3.0
2004 4/.....	3.4	...	...	0.4	2.6	3.2	5.5	6.6	4.3	2.4	2.5	3.0
2003 4/.....	3.1	...	...	0.2	2.1	3.4	5.1	5.8	4.1	2.3	2.6	2.4
2002.....	3.2	...	...	0.3	2.1	3.3	5.3	5.8	3.7	2.6	2.9	3.1
2001 5/.....	2.9	...	...	0.2	2.0	3.0	4.9	5.3	3.7	2.4	2.7	2.3
2000.....	2.8	...	...	0.2	1.9	2.8	4.5	4.9	3.6	2.5	2.6	3.3
1999.....	2.8	...	...	0.1	1.9	2.9	4.6	4.8	3.3	2.5	3.4	3.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

Black, both sexes

2006 4/.....	2.4	...	...	0.5	2.5	3.8	4.1	3.5	2.4	1.2	*	*
2005 4/.....	2.5	...	...	0.5	2.9	4.2	4.0	3.5	2.3	1.2	*	*
2004 4/.....	2.6	...	...	0.6	3.2	4.6	3.9	3.5	2.7	*	*	*
2003 4/.....	2.5	...	...	0.5	2.8	4.2	4.6	3.1	2.2	1.2	*	*
2002.....	2.3	...	...	0.4	2.4	4.2	4.2	3.0	2.0	1.5	*	*
2001 5/.....	2.3	...	...	0.4	2.3	4.2	3.9	3.4	2.4	1.7	*	*
2000.....	2.4	...	...	0.5	2.9	3.7	4.2	3.2	1.3	1.4	2.3	*
1999.....	2.3	...	...	0.4	2.5	4.2	4.1	2.7	1.5	1.9	*	*

Black, male

2006 4/.....	4.0	...	...	0.8	3.8	6.7	6.9	5.8	4.0	*	*	*
2005 4/.....	3.8	...	...	0.7	4.6	6.6	6.1	5.0	3.4	*	*	*
2004 4/.....	4.1	...	...	0.7	5.1	7.2	6.1	5.3	5.0	*	*	*
2003 4/.....	3.8	...	...	0.9	4.3	6.5	7.3	4.4	3.6	*	*	*
2002.....	3.7	...	...	0.6	3.8	7.3	6.7	4.7	2.9	*	*	*
2001 5/.....	3.7	...	...	0.6	3.8	7.1	6.1	5.0	3.7	3.0	*	*
2000.....	3.9	...	...	0.9	4.6	6.3	6.9	5.4	1.8	*	*	*
1999.....	3.7	...	...	*	4.0	7.2	6.6	4.4	2.2	3.1	*	*

Black, female

2006 4/.....	0.9	...	...	*	1.1	1.0	1.7	1.5	1.2	*	*	*
2005 4/.....	1.3	...	...	*	1.2	2.0	2.1	2.2	1.4	*	*	*
2004 4/.....	1.3	...	...	*	1.3	2.3	2.0	1.9	*	*	*	*
2003 4/.....	1.3	...	...	*	1.3	2.0	2.2	2.1	*	*	*	*
2002.....	1.1	...	...	*	1.0	1.5	1.9	1.6	1.4	*	*	*
2001 5/.....	1.1	...	...	*	0.7	1.5	2.0	2.0	*	*	*	*
2000.....	1.0	...	...	*	1.3	1.5	1.9	1.4	*	*	*	*
1999.....	1.0	...	...	*	0.9	1.6	1.9	1.2	*	*	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

All races 3/, both sexes

2006.....	6.2	8.1	2.2	1.0	13.5	11.7	6.9	5.1	3.2	2.1	2.1	1.9
2005.....	6.1	7.5	2.3	0.8	13.0	11.8	7.1	4.8	2.8	2.4	2.2	2.1
2004.....	5.9	8.0	2.4	0.8	12.2	11.2	6.8	4.8	3.0	2.4	2.2	2.1
2003.....	6.1	8.5	2.4	0.8	13.0	11.3	7.0	4.9	2.8	2.4	2.5	2.2
2002.....	6.1	7.5	2.7	0.9	12.9	11.2	7.2	4.8	3.2	2.3	2.3	2.1
2001 5/.....	7.1	8.2	2.7	0.8	13.3	13.1	9.5	6.3	4.0	2.9	2.5	2.4
2000.....	6.0	9.2	2.3	0.9	12.6	10.4	7.1	4.7	3.0	2.4	2.4	2.4
1999.....	6.1	8.7	2.5	1.1	12.9	10.5	7.1	4.6	3.0	2.6	2.5	2.4

All races 3/, male

2006.....	10.0	9.4	2.5	1.2	22.8	19.4	10.4	7.6	4.8	3.0	2.7	2.3
2005.....	9.8	8.2	2.6	1.0	22.0	19.6	10.6	7.6	4.3	3.3	2.6	2.7
2004.....	9.4	8.0	2.5	1.0	20.4	18.5	10.1	7.3	4.6	3.6	2.6	2.7
2003.....	9.7	10.0	2.5	1.0	21.8	18.5	10.4	7.4	4.2	3.4	3.3	3.2
2002.....	9.6	7.9	2.9	0.9	21.5	18.2	10.7	7.1	4.8	3.3	3.1	3.0
2001 5/.....	11.1	9.5	3.0	0.9	22.2	20.8	13.9	9.5	5.9	4.0	3.1	3.2
2000.....	9.3	10.4	2.5	1.1	20.9	16.7	10.3	6.9	4.6	3.4	3.2	3.3
1999.....	9.3	9.6	2.5	1.2	21.1	16.5	10.3	6.8	4.7	3.7	3.4	3.4

All races 3/, female

2006.....	2.5	6.8	2.0	0.7	3.5	3.7	3.4	2.6	1.7	1.3	1.8	1.8
2005.....	2.5	6.6	2.0	0.7	3.4	3.8	3.6	2.2	1.4	1.6	1.9	1.9
2004.....	2.5	7.9	2.2	0.6	3.5	3.7	3.4	2.4	1.6	1.4	1.9	1.9
2003.....	2.6	6.9	2.3	0.5	3.7	3.9	3.6	2.5	1.5	1.5	1.9	1.8
2002.....	2.7	7.1	2.5	0.8	3.8	4.2	3.8	2.6	1.6	1.4	1.8	1.7
2001 5/.....	3.3	6.9	2.4	0.7	3.9	5.3	5.1	3.2	2.2	2.0	2.1	2.0
2000.....	2.8	7.9	2.1	0.7	3.9	4.1	4.0	2.5	1.6	1.6	2.0	2.0
1999.....	2.9	7.8	2.4	0.9	4.3	4.4	4.0	2.4	1.5	1.7	1.9	2.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

White, both sexes

2006 4/.....	3.7	6.9	1.5	0.7	6.7	6.0	4.5	3.6	2.4	1.7	1.9	1.7
2005 4/.....	3.7	6.1	1.7	0.5	6.6	6.4	4.6	3.4	2.2	1.8	1.9	1.9
2004 4/.....	3.6	6.5	1.6	0.6	6.5	6.1	4.5	3.4	2.3	2.0	1.9	1.9
2003 4/.....	3.7	6.9	1.7	0.5	6.7	6.1	4.6	3.5	2.2	1.9	2.1	2.0
2002.....	3.7	5.4	2.0	0.6	6.7	5.9	4.8	3.3	2.6	1.9	1.9	1.8
2001 5/.....	4.9	6.2	1.9	0.7	7.2	8.3	7.3	5.0	3.4	2.5	2.1	2.2
2000.....	3.6	6.6	1.7	0.7	6.4	5.7	4.7	3.3	2.4	1.9	2.0	2.2
1999.....	3.8	6.7	1.7	0.8	6.8	6.0	4.8	3.3	2.4	2.2	2.0	2.1

White, male

2006 4/.....	5.4	7.4	1.7	0.8	10.6	9.3	6.4	5.1	3.5	2.2	2.2	2.0
2005 4/.....	5.4	6.7	1.8	0.6	10.6	9.8	6.3	5.0	3.3	2.4	2.1	1.9
2004 4/.....	5.3	6.9	1.8	0.7	10.2	9.3	6.3	4.9	3.4	2.8	2.1	2.3
2003 4/.....	5.4	8.1	1.5	0.6	10.6	9.2	6.3	5.0	3.1	2.7	2.8	3.0
2002.....	5.4	6.2	2.1	0.6	10.6	8.9	6.8	4.6	3.7	2.6	2.4	2.7
2001 5/.....	7.2	7.3	2.2	0.7	11.2	12.3	10.7	7.4	4.9	3.2	2.6	2.8
2000.....	5.2	8.2	1.9	0.9	9.9	8.4	6.5	4.6	3.5	2.4	2.5	2.8
1999.....	5.5	7.9	1.8	0.9	10.3	8.8	6.7	4.8	3.6	2.9	2.7	2.7

White, female

2006 4/.....	1.9	6.3	1.3	0.6	2.4	2.5	2.5	2.1	1.4	1.3	1.8	1.6
2005 4/.....	1.9	5.5	1.6	0.5	2.3	2.8	2.8	1.7	1.2	1.4	1.8	1.9
2004 4/.....	1.9	6.2	1.4	0.5	2.5	2.7	2.6	1.9	1.4	1.4	1.7	1.8
2003 4/.....	2.0	5.7	1.8	0.3	2.5	2.8	2.8	2.1	1.4	1.3	1.7	1.6
2002.....	2.0	4.6	1.8	0.6	2.5	2.8	2.8	2.1	1.5	1.3	1.6	1.5
2001 5/.....	2.6	5.1	1.6	0.6	3.0	4.1	3.8	2.7	2.0	1.8	1.8	2.0
2000.....	2.1	5.0	1.5	0.6	2.7	2.9	2.8	2.1	1.4	1.5	1.8	1.9
1999.....	2.1	5.4	1.7	0.7	3.0	3.1	2.9	1.8	1.3	1.6	1.6	1.9



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

Black, both sexes

2006 4/.....	22.8	16.0	5.8	2.3	49.4	47.2	23.6	15.9	9.6	6.0	4.4	*
2005 4/.....	22.2	14.3	5.3	2.3	46.9	46.3	24.0	15.7	8.1	7.1	5.7	*
2004 4/.....	21.1	15.6	6.3	1.9	43.5	44.2	22.6	15.3	8.8	6.0	6.3	*
2003 4/.....	22.0	16.1	6.1	2.2	47.6	45.2	23.4	15.4	8.1	6.2	6.6	*
2002.....	22.0	17.3	6.3	2.0	46.8	45.3	23.7	15.5	8.3	5.7	7.4	*
2001 5/.....	22.1	18.9	6.8	1.4	47.3	43.7	25.0	16.1	8.9	7.1	6.7	*
2000.....	21.5	22.8	5.8	1.8	47.8	40.8	23.9	14.8	8.3	7.6	7.0	*
1999.....	21.1	19.5	6.1	2.2	47.7	39.3	23.2	14.0	8.2	6.3	7.4	*

Black, male

2006 4/.....	40.6	20.7	6.3	3.1	88.2	86.2	39.8	27.1	17.0	11.4	8.1	*
2005 4/.....	39.7	15.9	6.6	2.9	84.1	86.2	40.6	27.0	14.3	12.6	10.1	*
2004 4/.....	37.1	14.2	6.3	2.4	77.6	81.6	37.9	26.0	15.1	11.5	10.3	*
2003 4/.....	38.9	17.8	7.4	2.9	84.6	82.5	40.2	26.3	14.8	11.1	11.6	*
2002.....	38.4	16.3	7.1	2.5	83.1	82.2	38.8	27.1	15.4	10.6	13.8	*
2001 5/.....	38.3	21.0	8.0	2.0	85.7	78.8	38.5	26.9	15.0	12.6	10.0	*
2000.....	37.2	23.3	5.9	2.1	85.3	73.9	38.5	26.0	14.7	13.9	11.8	*
1999.....	36.1	19.5	5.9	2.7	84.7	69.2	37.2	23.4	15.1	11.8	12.0	*

Black, female

2006 4/.....	6.6	11.1	5.4	1.4	9.5	11.1	9.2	6.3	3.6	2.2	*	*
2005 4/.....	6.2	12.6	4.0	1.6	8.8	9.3	9.2	6.0	3.1	3.1	3.3	*
2004 4/.....	6.4	17.0	6.3	1.5	8.8	9.8	9.1	6.0	3.7	2.1	4.1	*
2003 4/.....	6.6	14.2	4.8	1.5	10.1	11.1	8.5	6.0	2.7	2.7	3.8	*
2002.....	7.0	18.5	5.5	1.5	10.3	11.9	10.4	5.6	2.5	2.3	3.7	*
2001 5/.....	7.4	16.7	5.6	0.8	8.9	12.1	13.0	6.8	3.9	3.3	4.8	*
2000.....	7.2	22.2	5.7	1.6	10.7	10.9	11.0	5.3	3.3	3.1	4.3	*
1999.....	7.6	19.5	6.3	1.7	11.2	12.4	10.8	5.9	2.6	2.5	4.8	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

All races 3/, both sexes

2006.....	4.3	*	0.3	0.6	11.4	9.3	4.6	2.7	1.5	0.8	0.6	0.4
2005.....	4.2	*	0.2	0.5	10.7	9.4	4.6	2.6	1.3	1.1	0.7	0.5
2004.....	4.0	*	0.2	0.5	9.9	8.8	4.3	2.6	1.6	1.0	0.8	0.6
2003.....	4.1	*	0.3	0.5	10.7	8.9	4.4	2.7	1.4	0.9	0.7	0.5
2002.....	4.1	*	0.3	0.5	10.6	8.7	4.5	2.6	1.5	0.9	0.9	*
2001.....	4.0	*	0.4	0.4	10.5	8.4	4.4	2.4	1.4	1.0	0.8	*
2000.....	3.8	*	0.2	0.5	10.1	7.7	4.2	2.5	1.5	1.0	0.7	0.7
1999.....	3.9	*	0.3	0.5	10.4	7.6	4.1	2.4	1.6	1.0	0.8	0.6

All races 3/, male

2006.....	7.4	*	0.3	0.8	20.0	16.2	7.3	4.3	2.2	1.2	0.8	*
2005.....	7.2	*	0.3	0.6	18.9	16.4	7.3	4.1	2.1	1.5	0.8	*
2004.....	6.9	*	*	0.6	17.3	15.3	7.1	4.0	2.5	1.6	0.8	*
2003.....	7.1	*	0.2	0.6	18.8	15.5	6.9	4.3	2.2	1.4	0.7	*
2002.....	7.0	*	0.4	0.6	18.5	14.9	7.3	3.9	2.4	1.2	1.3	*
2001.....	6.8	*	0.3	0.6	18.5	14.3	6.8	3.8	2.2	1.5	1.1	*
2000.....	6.5	*	*	0.7	17.7	13.2	6.6	3.9	2.3	1.3	0.8	*
1999.....	6.5	*	0.3	0.7	18.0	12.9	6.3	3.8	2.6	1.5	1.0	*

All races 3/, female

2006.....	1.3	*	*	0.4	2.2	2.2	1.8	1.3	0.8	0.5	0.6	*
2005.....	1.2	*	*	0.3	2.0	2.2	1.9	1.1	0.6	0.7	0.7	*
2004.....	1.1	*	*	0.3	2.0	2.0	1.5	1.1	0.8	0.5	0.8	*
2003.....	1.2	*	0.3	0.3	2.2	2.1	1.8	1.2	0.7	0.5	0.7	*
2002.....	1.3	*	*	0.4	2.4	2.3	1.9	1.3	0.8	0.6	0.7	*
2001.....	1.3	*	0.4	0.3	2.1	2.3	2.0	1.0	0.8	0.7	0.6	*
2000.....	1.3	*	*	0.2	2.2	2.1	1.8	1.2	0.8	0.8	0.6	*
1999.....	1.3	*	0.3	0.3	2.5	2.2	2.0	1.2	0.6	0.7	0.6	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

White, both sexes

2006 4/.....	2.2	*	0.2	0.4	5.1	4.2	2.8	1.8	1.2	0.6	0.6	*
2005 4/.....	2.2	*	0.2	0.3	4.8	4.5	2.8	1.8	1.1	0.9	0.7	*
2004 4/.....	2.1	*	*	0.3	4.7	4.3	2.7	1.8	1.3	0.9	0.7	0.5
2003 4/.....	2.2	*	0.2	0.3	5.0	4.2	2.7	1.9	1.1	0.7	0.6	0.5
2002.....	2.2	*	0.3	0.3	5.0	4.1	2.8	1.8	1.2	0.7	0.8	*
2001.....	2.2	*	0.3	0.4	5.0	4.2	2.9	1.7	1.2	0.9	0.7	*
2000.....	2.1	*	*	0.4	4.6	3.7	2.7	1.8	1.3	0.8	0.6	0.6
1999.....	2.2	*	0.2	0.4	4.9	3.8	2.7	1.7	1.2	0.9	0.7	0.6

White, male

2006 4/.....	3.5	*	*	0.5	8.5	6.9	4.1	2.5	1.6	0.8	0.5	*
2005 4/.....	3.5	*	*	0.4	8.2	7.3	4.0	2.7	1.6	1.1	0.7	*
2004 4/.....	3.5	*	*	0.4	7.8	7.0	4.1	2.6	1.8	1.2	0.7	*
2003 4/.....	3.5	*	*	0.4	8.5	6.9	3.9	2.8	1.6	1.1	0.5	*
2002.....	3.5	*	0.3	0.4	8.4	6.6	4.2	2.5	1.8	0.9	1.0	*
2001.....	3.5	*	*	0.5	8.3	6.7	4.3	2.6	1.6	1.1	0.8	*
2000.....	3.2	*	*	0.5	7.7	5.9	3.9	2.6	1.8	1.0	0.6	*
1999.....	3.4	*	*	0.5	8.0	6.1	4.0	2.6	1.9	1.2	0.7	*

White, female

2006 4/.....	0.9	*	*	0.3	1.3	1.4	1.3	1.1	0.7	0.5	0.6	*
2005 4/.....	0.9	*	*	0.2	1.2	1.6	1.5	0.9	0.5	0.7	0.7	*
2004 4/.....	0.9	*	*	0.2	1.3	1.3	1.2	0.9	0.7	0.6	0.7	*
2003 4/.....	0.9	*	*	0.1	1.3	1.4	1.5	1.0	0.7	0.4	0.6	*
2002.....	1.0	*	*	0.3	1.4	1.5	1.4	1.1	0.7	0.6	0.6	*
2001.....	1.0	*	*	0.3	1.5	1.6	1.5	0.9	0.8	0.7	0.6	*
2000.....	1.0	*	*	0.2	1.4	1.5	1.4	1.0	0.8	0.7	0.6	*
1999.....	0.9	*	*	0.3	1.5	1.5	1.5	0.9	0.6	0.7	0.6	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

Black, both sexes

2006 4/.....	18.0	*	0.8	1.5	44.8	41.2	17.1	9.4	4.2	2.8	*	*
2005 4/.....	17.2	*	*	1.3	42.1	40.4	16.8	8.6	3.5	3.1	*	*
2004 4/.....	16.1	*	0.8	1.2	38.8	37.7	15.6	8.4	4.8	2.4	*	*
2003 4/.....	16.8	*	*	1.4	41.9	38.8	15.9	8.7	3.9	2.5	*	*
2002.....	16.7	*	*	1.2	41.8	38.2	16.3	8.5	4.1	2.4	2.2	*
2001.....	15.8	*	*	0.8	41.8	35.1	15.0	7.5	3.7	2.7	*	*
2000.....	15.6	*	*	1.0	41.2	33.2	15.0	8.4	3.5	2.7	*	*
1999.....	15.4	*	*	1.2	42.0	31.4	14.1	8.0	4.4	2.4	2.3	*

Black, male

2006 4/.....	33.7	*	*	2.2	81.6	77.5	30.7	17.5	7.9	5.6	*	*
2005 4/.....	32.5	*	*	1.9	77.4	77.3	30.8	15.5	6.7	6.0	*	*
2004 4/.....	30.3	*	*	1.8	71.3	72.2	28.9	15.2	8.6	5.4	*	*
2003 4/.....	31.6	*	*	1.9	76.8	74.1	29.1	15.9	7.5	4.6	*	*
2002.....	31.0	*	*	1.6	76.1	72.8	29.2	15.1	7.6	4.7	*	*
2001.....	29.8	*	*	1.3	78.2	66.9	26.0	13.7	7.1	5.6	*	*
2000.....	29.2	*	*	1.5	76.5	63.3	26.7	15.1	6.3	5.4	*	*
1999.....	28.4	*	*	1.8	76.9	59.1	24.0	13.9	8.9	4.6	*	*

Black, female

2006 4/.....	3.6	*	*	0.8	7.0	7.5	5.0	2.5	1.2	*	*	*
2005 4/.....	3.1	*	*	0.7	6.0	6.2	4.4	2.6	*	*	*	*
2004 4/.....	3.0	*	*	*	5.8	6.1	3.8	2.5	1.7	*	*	*
2003 4/.....	3.3	*	*	0.8	6.5	6.6	4.2	2.5	*	*	*	*
2002.....	3.6	*	*	0.8	7.3	6.9	4.8	2.8	1.4	*	*	*
2001.....	3.1	*	*	*	5.4	6.3	5.4	2.2	*	*	*	*
2000.....	3.2	*	*	*	6.4	6.1	4.6	2.6	*	*	*	*
1999.....	3.7	*	*	0.6	7.6	6.5	5.3	3.0	*	*	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes

2006.....	1.9	8.0	2.0	0.4	2.1	2.4	2.4	2.4	1.7	1.3	1.5	1.5
2005.....	1.9	7.3	2.1	0.4	2.3	2.4	2.5	2.3	1.5	1.3	1.4	1.7
2004.....	2.0	7.8	2.1	0.4	2.3	2.5	2.5	2.3	1.4	1.4	1.5	1.5
2003.....	2.0	8.3	2.1	0.3	2.3	2.4	2.6	2.2	1.4	1.5	1.8	1.7
2002.....	2.0	7.3	2.4	0.4	2.2	2.6	2.7	2.2	1.6	1.4	1.4	1.7
2001 5/.....	3.1	8.0	2.3	0.4	2.7	4.8	5.1	3.9	2.6	1.9	1.7	2.0
2000.....	2.1	8.9	2.1	0.4	2.5	2.7	2.9	2.1	1.5	1.4	1.8	1.7
1999.....	2.2	8.5	2.1	0.5	2.5	2.9	3.0	2.1	1.5	1.5	1.7	1.8

All races 3/, male

2006.....	2.6	9.3	2.2	0.4	2.8	3.2	3.1	3.4	2.6	1.8	1.9	1.8
2005.....	2.6	7.9	2.3	0.4	3.1	3.3	3.3	3.4	2.2	1.8	1.8	2.1
2004.....	2.5	7.9	2.3	0.3	3.1	3.2	3.0	3.3	2.1	1.9	1.9	1.7
2003.....	2.6	9.7	2.3	0.4	3.1	3.0	3.5	3.1	2.1	2.0	2.6	2.5
2002.....	2.6	7.8	2.6	0.3	3.0	3.2	3.4	3.2	2.4	2.1	1.9	2.3
2001 5/.....	4.3	9.2	2.7	0.3	3.6	6.5	7.1	5.7	3.8	2.5	2.0	2.7
2000.....	2.8	10.0	2.3	0.4	3.3	3.5	3.7	3.0	2.3	2.1	2.4	2.5
1999.....	2.8	9.4	2.1	0.5	3.1	3.6	4.0	3.1	2.2	2.2	2.4	2.6

All races 3/, female

2006.....	1.3	6.6	1.8	0.4	1.3	1.5	1.6	1.4	0.9	0.8	1.2	1.4
2005.....	1.3	6.6	1.8	0.4	1.5	1.6	1.7	1.1	0.8	0.9	1.2	1.5
2004.....	1.4	7.7	2.0	0.4	1.5	1.7	1.9	1.3	0.8	0.9	1.2	1.5
2003.....	1.4	6.9	2.0	0.3	1.6	1.8	1.8	1.3	0.8	1.0	1.2	1.4
2002.....	1.4	6.8	2.2	0.4	1.4	1.9	1.9	1.3	0.9	0.8	1.1	1.4
2001 5/.....	2.0	6.7	2.0	0.4	1.8	3.1	3.1	2.1	1.5	1.3	1.5	1.7
2000.....	1.5	7.7	2.0	0.5	1.7	1.9	2.1	1.3	0.8	0.9	1.4	1.4
1999.....	1.6	7.6	2.1	0.6	1.8	2.2	2.0	1.2	0.8	1.0	1.3	1.5

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

White, both sexes

2006 4/.....	1.5	6.8	1.3	0.3	1.6	1.8	1.7	1.8	1.3	1.1	1.4	1.4
2005 4/.....	1.5	6.0	1.5	0.3	1.8	1.9	1.8	1.6	1.2	1.0	1.2	1.5
2004 4/.....	1.5	6.5	1.5	0.3	1.8	1.8	1.8	1.6	1.1	1.1	1.2	1.4
2003 4/.....	1.5	6.9	1.5	0.2	1.7	1.8	1.9	1.6	1.1	1.2	1.5	1.6
2002.....	1.5	5.3	1.7	0.3	1.7	1.8	2.0	1.6	1.3	1.2	1.1	1.4
2001 5/.....	2.6	6.0	1.6	0.3	2.2	4.1	4.4	3.3	2.3	1.6	1.4	1.8
2000.....	1.5	6.5	1.5	0.4	1.8	2.0	2.0	1.5	1.1	1.0	1.5	1.5
1999.....	1.6	6.6	1.5	0.4	1.9	2.1	2.0	1.6	1.2	1.3	1.4	1.5

White, male

2006 4/.....	1.9	7.4	1.5	0.3	2.1	2.4	2.3	2.6	1.9	1.4	1.7	1.7
2005 4/.....	1.9	6.4	1.6	0.2	2.4	2.4	2.3	2.4	1.7	1.3	1.4	1.6
2004 4/.....	1.9	6.8	1.7	0.3	2.4	2.2	2.2	2.3	1.6	1.5	1.4	1.5
2003 4/.....	1.9	8.0	1.4	0.3	2.1	2.3	2.4	2.2	1.5	1.5	2.2	2.3
2002.....	1.9	6.2	1.8	0.2	2.2	2.3	2.5	2.1	1.9	1.8	1.3	2.0
2001 5/.....	3.7	7.0	1.9	0.3	2.8	5.6	6.4	4.8	3.3	2.1	1.8	2.3
2000.....	2.0	8.1	1.7	0.4	2.3	2.5	2.6	2.0	1.7	1.4	1.9	2.1
1999.....	2.1	7.7	1.5	0.4	2.3	2.7	2.7	2.3	1.7	1.7	2.0	1.9

White, female

2006 4/.....	1.0	6.2	1.2	0.3	1.1	1.1	1.2	1.0	0.7	0.8	1.1	1.2
2005 4/.....	1.0	5.5	1.4	0.3	1.2	1.3	1.2	0.8	0.7	0.7	1.1	1.5
2004 4/.....	1.1	6.1	1.3	0.3	1.2	1.4	1.4	1.0	0.6	0.8	1.0	1.3
2003 4/.....	1.1	5.6	1.6	0.2	1.2	1.4	1.4	1.0	0.7	0.9	1.1	1.2
2002.....	1.1	4.5	1.6	0.4	1.1	1.3	1.4	1.0	0.8	0.7	1.0	1.2
2001 5/.....	1.6	4.9	1.4	0.4	1.5	2.5	2.3	1.8	1.3	1.1	1.2	1.6
2000.....	1.1	4.8	1.3	0.4	1.3	1.4	1.4	1.1	0.6	0.7	1.2	1.3
1999.....	1.2	5.4	1.5	0.5	1.4	1.6	1.4	0.9	0.7	0.9	1.0	1.4

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

Black, both sexes

2006 4/.....	4.9	15.4	5.1	0.7	4.6	6.0	6.5	6.5	5.4	3.2	3.0	*
2005 4/.....	5.0	14.0	4.8	0.9	4.8	5.9	7.1	7.2	4.6	4.0	4.3	*
2004 4/.....	5.0	15.1	5.4	0.7	4.7	6.4	7.0	6.9	4.0	3.6	4.9	*
2003 4/.....	5.2	15.2	5.3	0.8	5.7	6.4	7.5	6.8	4.1	3.7	4.9	*
2002.....	5.3	16.5	5.8	0.8	5.0	7.1	7.4	7.1	4.1	3.4	5.2	*
2001 5/.....	6.3	18.6	6.1	0.5	5.5	8.7	9.9	8.5	5.2	4.4	4.8	*
2000.....	5.9	21.6	5.5	0.8	6.5	7.6	8.9	6.5	4.9	4.8	5.1	*
1999.....	5.7	19.2	5.4	1.0	5.7	7.9	9.1	6.0	3.8	4.0	5.1	*

Black, male

2006 4/.....	6.9	20.2	5.4	0.9	6.6	8.6	9.1	9.6	9.2	5.9	*	*
2005 4/.....	7.2	15.4	5.9	1.0	6.7	8.9	9.8	11.5	7.7	6.6	6.8	*
2004 4/.....	6.8	13.9	5.5	0.6	6.4	9.5	9.0	10.8	6.5	6.1	8.4	*
2003 4/.....	7.4	16.1	6.5	0.9	7.7	8.4	11.1	10.5	7.3	6.5	8.7	*
2002.....	7.3	15.7	6.4	0.9	7.0	9.4	9.6	12.0	7.8	5.9	9.2	*
2001 5/.....	8.5	20.7	7.3	0.7	7.5	11.8	12.6	13.2	8.0	7.0	*	*
2000.....	8.0	21.1	5.5	*	8.8	10.6	11.7	10.8	8.3	8.5	8.2	*
1999.....	7.7	19.2	5.4	0.9	7.8	10.1	13.2	9.6	6.2	7.2	7.1	*

Black, female

2006 4/.....	3.0	10.5	4.7	*	2.6	3.5	4.2	3.8	2.4	*	*	*
2005 4/.....	3.0	12.6	3.7	0.9	2.9	3.2	4.8	3.4	2.1	2.1	*	*
2004 4/.....	3.4	16.4	5.4	0.9	3.0	3.6	5.3	3.5	2.0	*	*	*
2003 4/.....	3.3	14.2	4.1	0.7	3.6	4.6	4.2	3.6	1.6	*	*	*
2002.....	3.4	17.3	5.1	0.7	3.0	5.0	5.6	2.8	*	*	*	*
2001 5/.....	4.3	16.4	4.8	*	3.5	5.8	7.6	4.6	2.9	2.7	4.3	*
2000.....	4.0	22.2	5.5	1.1	4.3	4.8	6.4	2.7	2.1	2.3	3.4	*
1999.....	3.9	19.1	5.4	1.1	3.6	5.9	5.5	2.9	1.9	*	4.0	*









Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

---

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
--	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

---

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2006.....	1.7	2.2	0.3	0.1	1.4	2.0	2.9	3.1	1.6	0.9	1.0	1.4
2005.....	1.6	2.5	0.4	0.2	1.2	1.8	2.7	3.1	1.4	0.8	0.8	1.5
2004.....	1.7	2.6	0.3	0.2	1.2	2.0	3.1	3.1	1.4	0.8	0.9	1.6
2003.....	1.7	1.9	0.2	0.1	1.4	2.1	3.6	3.1	1.3	0.7	0.7	1.1
2002.....	1.7	2.5	0.4	0.2	1.2	2.1	3.4	2.8	1.2	0.8	0.7	1.4
2001.....	1.5	2.1	0.3	0.1	1.0	1.8	3.1	2.5	0.9	0.5	0.9	1.0
2000.....	1.4	1.7	0.3	0.1	0.9	1.7	2.8	2.2	0.9	0.6	0.6	1.3
1999.....	1.4	1.4	0.3	0.1	1.1	1.8	2.9	2.2	0.9	0.6	0.8	1.4

All races 3/, male

2006.....	2.2	2.7	0.3	0.2	2.1	2.8	3.5	3.9	1.9	1.1	1.4	2.2
2005.....	2.0	3.2	0.5	0.2	1.6	2.3	3.2	3.6	1.7	1.1	0.9	1.6
2004.....	2.2	3.1	0.3	0.2	1.8	2.7	4.0	4.0	1.7	1.1	1.0	1.9
2003.....	2.3	2.4	*	0.2	1.9	2.9	4.6	4.0	1.7	0.8	1.0	1.5
2002.....	2.2	2.5	0.4	0.2	1.7	2.9	4.4	3.5	1.6	1.0	0.9	2.2
2001.....	2.0	2.4	*	0.2	1.5	2.6	4.2	3.4	1.2	0.8	1.1	1.5
2000.....	1.9	1.4	0.3	0.2	1.4	2.4	3.9	3.1	1.3	0.8	1.0	1.7
1999.....	2.0	1.7	0.3	0.2	1.7	2.6	4.0	3.0	1.3	0.9	1.1	2.1

All races 3/, female

2006.....	1.2	1.7	0.3	*	0.7	1.2	2.3	2.4	1.2	0.7	0.8	1.0
2005.....	1.3	1.8	0.3	0.1	0.6	1.3	2.3	2.6	1.1	0.7	0.7	1.4
2004.....	1.2	2.0	0.3	0.1	0.6	1.3	2.3	2.3	1.1	0.6	0.9	1.6
2003.....	1.2	1.3	*	*	0.8	1.3	2.6	2.2	1.0	0.7	0.6	0.9
2002.....	1.2	2.5	0.4	0.1	0.7	1.2	2.4	2.2	0.8	0.6	0.6	1.1
2001.....	0.9	1.7	0.3	*	0.5	1.1	2.1	1.6	0.6	0.3	0.7	0.8
2000.....	0.8	2.0	*	*	0.4	1.0	1.8	1.4	0.6	0.3	0.4	1.2
1999.....	0.9	1.2	0.3	*	0.5	1.1	1.8	1.4	0.6	0.4	0.6	1.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

White, both sexes

2006 4/.....	1.7	2.0	0.2	0.1	1.6	2.2	3.0	3.1	1.5	0.8	0.9	1.1
2005 4/.....	1.6	2.1	0.3	0.1	1.2	2.0	2.8	3.1	1.4	0.8	0.8	1.5
2004 4/.....	1.7	2.1	0.2	0.1	1.3	2.1	3.3	3.2	1.4	0.8	0.9	1.6
2003 4/.....	1.8	1.7	0.2	0.1	1.5	2.3	3.7	3.1	1.3	0.6	0.7	1.2
2002.....	1.7	1.8	0.4	0.2	1.3	2.1	3.4	2.8	1.2	0.7	0.7	1.4
2001.....	1.5	1.4	0.3	0.1	1.1	1.9	3.1	2.5	0.9	0.5	0.9	1.0
2000.....	1.3	1.4	0.2	0.1	1.0	1.7	2.8	2.1	0.9	0.5	0.6	1.3
1999.....	1.4	1.2	0.2	0.1	1.1	1.9	2.8	2.0	0.9	0.6	0.8	1.4

White, male

2006 4/.....	2.2	2.5	*	0.2	2.3	2.9	3.5	3.7	1.7	0.9	1.2	2.0
2005 4/.....	2.0	2.8	0.4	0.2	1.8	2.5	3.2	3.5	1.6	1.0	0.8	1.7
2004 4/.....	2.2	2.7	*	0.1	1.9	2.9	4.0	3.9	1.6	1.0	1.0	1.9
2003 4/.....	2.3	1.9	*	0.2	2.1	3.0	4.7	3.9	1.5	0.7	1.0	1.5
2002.....	2.2	1.8	0.4	0.2	1.9	2.9	4.4	3.3	1.5	0.9	0.9	2.3
2001.....	2.0	1.7	*	0.2	1.6	2.7	4.0	3.2	1.1	0.7	1.2	*
2000.....	1.8	*	*	0.2	1.4	2.4	3.8	2.8	1.1	0.7	1.0	*
1999.....	1.9	1.4	*	0.2	1.8	2.5	3.8	2.6	1.1	0.8	1.1	2.1

White, female

2006 4/.....	1.3	1.4	*	*	0.8	1.4	2.5	2.5	1.3	0.7	0.8	0.7
2005 4/.....	1.3	1.5	*	*	0.7	1.5	2.3	2.7	1.2	0.7	0.7	1.3
2004 4/.....	1.3	1.5	*	*	0.7	1.3	2.5	2.5	1.2	0.6	0.9	1.5
2003 4/.....	1.3	1.4	*	*	0.9	1.5	2.7	2.4	1.0	0.6	0.6	1.0
2002.....	1.2	1.8	0.4	0.1	0.7	1.3	2.5	2.2	0.9	0.6	0.6	1.1
2001.....	1.0	*	*	*	0.6	1.2	2.2	1.7	0.6	0.4	0.6	0.8
2000.....	0.8	1.6	*	*	0.5	1.0	1.7	1.4	0.6	0.4	0.4	1.1
1999.....	0.9	*	*	*	0.4	1.2	1.8	1.4	0.7	0.4	0.6	1.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

Black, both sexes

2006 4/.....	1.8	3.4	0.8	*	0.9	1.5	3.2	3.9	2.3	1.3	*	*
2005 4/.....	1.8	4.7	0.9	*	0.9	1.6	3.3	3.7	1.7	*	*	*
2004 4/.....	1.8	4.8	*	0.4	0.9	1.8	3.3	3.9	1.7	1.3	*	*
2003 4/.....	1.8	3.2	*	*	0.8	1.8	3.8	4.0	1.9	1.2	*	*
2002.....	2.0	5.5	*	*	0.9	2.3	4.2	4.2	1.5	*	*	*
2001.....	1.7	5.5	*	*	0.7	1.7	3.9	3.8	1.3	*	*	*
2000.....	1.8	*	*	0.3	0.9	1.9	4.1	3.7	1.8	*	*	*
1999.....	1.9	*	*	*	1.2	2.2	4.2	4.1	1.2	*	*	*

Black, male

2006 4/.....	2.7	*	*	*	1.3	2.4	4.7	6.2	3.5	*	*	*
2005 4/.....	2.4	*	*	*	1.2	2.3	4.2	5.6	2.9	*	*	*
2004 4/.....	2.6	*	*	*	1.3	2.8	4.6	5.8	2.9	*	*	*
2003 4/.....	2.7	*	*	*	1.4	2.9	5.3	6.4	3.2	*	*	*
2002.....	2.9	*	*	*	1.4	3.5	6.0	6.4	2.5	*	*	*
2001.....	2.7	6.0	*	*	1.3	2.6	6.2	6.3	2.4	*	*	*
2000.....	2.8	*	*	*	1.6	3.1	6.0	5.9	3.0	*	*	*
1999.....	3.0	*	*	*	1.9	3.6	6.4	7.2	2.5	*	*	*

Black, female

2006 4/.....	1.0	*	*	*	*	0.7	1.9	2.0	1.3	*	*	*
2005 4/.....	1.1	*	*	*	*	1.0	2.5	2.2	*	*	*	*
2004 4/.....	1.1	*	*	*	*	0.9	2.2	2.2	*	*	*	*
2003 4/.....	0.9	*	*	*	*	*	2.4	1.9	*	*	*	*
2002.....	1.2	*	*	*	*	1.2	2.6	2.4	*	*	*	*
2001.....	0.9	*	*	*	*	0.9	1.9	1.6	*	*	*	*
2000.....	0.9	*	*	*	*	0.9	2.4	1.8	*	*	*	*
1999.....	0.9	*	*	*	*	1.0	2.3	1.6	*	*	*	*









Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes

2006.....	1.6	2.2	0.3	0.1	1.2	1.9	2.8	3.1	1.5	0.8	1.0	1.4
2005.....	1.5	2.5	0.4	0.1	1.0	1.8	2.7	3.0	1.3	0.8	0.8	1.4
2004.....	1.6	2.6	0.3	0.1	1.1	1.9	3.0	3.1	1.3	0.8	0.9	1.6
2003.....	1.7	1.9	0.2	0.1	1.2	2.0	3.5	3.0	1.2	0.7	0.7	1.1
2002.....	1.6	2.5	0.4	0.2	1.0	2.0	3.3	2.8	1.2	0.7	0.7	1.4
2001.....	1.4	2.1	0.3	0.1	0.9	1.7	3.0	2.5	0.8	0.5	0.8	1.0
2000.....	1.3	1.7	0.3	0.1	0.8	1.6	2.8	2.2	0.9	0.5	0.6	1.3
1999.....	1.3	1.4	0.3	0.1	0.8	1.7	2.7	2.1	0.9	0.5	0.7	1.3

All races 3/, male

2006.....	2.1	2.7	0.3	0.2	1.8	2.6	3.4	3.8	1.8	1.0	1.4	2.2
2005.....	1.8	3.2	0.5	0.2	1.3	2.2	3.1	3.5	1.6	1.0	0.9	1.5
2004.....	2.1	3.1	0.3	0.2	1.5	2.6	3.8	3.8	1.6	0.9	0.9	1.8
2003.....	2.2	2.4	*	0.1	1.7	2.7	4.5	3.9	1.5	0.7	0.8	1.4
2002.....	2.1	2.5	0.3	0.2	1.4	2.7	4.2	3.4	1.5	0.9	0.8	2.2
2001.....	1.9	2.4	*	0.2	1.3	2.4	4.0	3.3	1.1	0.7	1.0	*
2000.....	1.8	1.4	0.3	0.2	1.1	2.2	3.8	3.0	1.2	0.7	0.9	*
1999.....	1.8	1.7	*	0.1	1.2	2.4	3.8	2.8	1.2	0.7	0.9	1.8

All races 3/, female

2006.....	1.2	1.7	0.3	*	0.7	1.2	2.3	2.4	1.2	0.7	0.8	1.0
2005.....	1.2	1.8	0.3	*	0.6	1.3	2.2	2.5	1.1	0.6	0.7	1.4
2004.....	1.2	2.0	0.3	*	0.6	1.2	2.3	2.3	1.1	0.6	0.8	1.5
2003.....	1.2	1.3	*	*	0.7	1.3	2.5	2.2	1.0	0.7	0.6	0.9
2002.....	1.1	2.5	0.4	0.1	0.6	1.2	2.3	2.1	0.8	0.5	0.6	1.1
2001.....	0.9	1.7	0.3	*	0.5	1.1	2.0	1.6	0.6	0.3	0.6	0.8
2000.....	0.8	2.0	*	*	0.4	0.9	1.7	1.4	0.6	0.3	0.4	1.2
1999.....	0.8	1.2	0.3	*	0.4	1.1	1.7	1.4	0.6	0.4	0.5	1.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

White, both sexes

2006 4/.....	1.7	2.0	0.2	0.1	1.4	2.1	2.9	3.1	1.4	0.8	0.9	1.1
2005 4/.....	1.6	2.1	0.3	0.1	1.1	1.9	2.7	3.0	1.3	0.8	0.7	1.4
2004 4/.....	1.7	2.1	0.2	0.1	1.2	2.1	3.2	3.1	1.3	0.7	0.8	1.6
2003 4/.....	1.7	1.7	0.2	0.1	1.4	2.2	3.6	3.0	1.2	0.6	0.7	1.2
2002.....	1.6	1.8	0.4	0.1	1.1	2.0	3.3	2.7	1.2	0.6	0.7	1.4
2001.....	1.4	1.4	0.3	0.1	1.0	1.8	3.0	2.4	0.8	0.5	0.8	1.0
2000.....	1.3	1.4	0.2	0.1	0.8	1.6	2.7	2.1	0.8	0.5	0.6	1.2
1999.....	1.3	1.2	0.2	0.1	0.8	1.8	2.6	1.9	0.9	0.5	0.7	1.3

White, male

2006 4/.....	2.1	2.5	*	0.2	2.0	2.8	3.3	3.6	1.6	0.9	1.1	2.0
2005 4/.....	1.8	2.8	0.4	0.1	1.5	2.3	3.1	3.4	1.5	0.9	0.8	1.6
2004 4/.....	2.1	2.7	*	0.1	1.7	2.8	3.9	3.7	1.5	0.8	0.9	1.8
2003 4/.....	2.2	1.9	*	0.1	1.9	2.9	4.6	3.8	1.3	0.6	0.8	*
2002.....	2.0	1.8	0.4	0.1	1.5	2.8	4.2	3.2	1.5	0.7	0.8	2.3
2001.....	1.9	1.7	*	0.2	1.4	2.5	3.9	3.1	1.0	0.6	1.0	*
2000.....	1.7	*	*	0.2	1.2	2.2	3.7	2.8	1.0	0.6	0.9	*
1999.....	1.7	1.4	*	0.1	1.3	2.3	3.5	2.5	1.1	0.7	0.9	*

White, female

2006 4/.....	1.3	1.4	*	*	0.8	1.4	2.5	2.5	1.2	0.7	0.8	0.7
2005 4/.....	1.3	1.5	*	*	0.6	1.4	2.3	2.7	1.2	0.7	0.7	1.3
2004 4/.....	1.2	1.5	*	*	0.6	1.3	2.4	2.4	1.1	0.6	0.8	1.5
2003 4/.....	1.3	1.4	*	*	0.8	1.5	2.6	2.3	1.0	0.6	0.6	1.0
2002.....	1.2	1.8	0.4	0.1	0.7	1.3	2.4	2.2	0.9	0.6	0.6	1.1
2001.....	1.0	*	*	*	0.6	1.1	2.1	1.7	0.6	0.3	0.6	0.8
2000.....	0.8	1.6	*	*	0.5	1.0	1.7	1.4	0.6	0.3	0.4	1.1
1999.....	0.8	*	*	*	0.4	1.2	1.7	1.4	0.7	0.4	0.6	1.1

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

Black, both sexes

2006 4/.....	1.7	3.4	0.8	*	0.5	1.3	3.1	3.9	2.3	1.3	*	*
2005 4/.....	1.7	4.7	0.8	*	0.6	1.5	3.2	3.7	1.6	*	*	*
2004 4/.....	1.7	4.8	*	0.3	0.6	1.6	3.3	3.8	1.7	1.2	*	*
2003 4/.....	1.7	3.2	*	*	0.5	1.6	3.7	3.9	1.9	1.2	*	*
2002.....	1.9	5.5	*	*	0.6	2.1	4.1	4.2	1.4	*	*	*
2001.....	1.7	5.5	*	*	0.5	1.6	3.9	3.7	1.1	*	*	*
2000.....	1.7	*	*	*	0.5	1.7	3.9	3.6	1.6	*	*	*
1999.....	1.7	*	*	*	0.6	2.0	4.1	4.0	1.1	*	*	*

Black, male

2006 4/.....	2.5	*	*	*	0.7	2.0	4.5	6.1	3.5	*	*	*
2005 4/.....	2.3	*	*	*	0.8	2.1	4.1	5.5	2.9	*	*	*
2004 4/.....	2.4	*	*	*	1.0	2.5	4.5	5.7	2.9	*	*	*
2003 4/.....	2.6	*	*	*	0.8	2.7	5.2	6.3	3.2	*	*	*
2002.....	2.7	*	*	*	0.8	3.1	5.8	6.3	2.5	*	*	*
2001.....	2.5	6.0	*	*	0.8	2.4	6.2	6.1	2.1	*	*	*
2000.....	2.5	*	*	*	1.0	2.6	5.8	5.8	2.8	*	*	*
1999.....	2.7	*	*	*	0.9	3.1	6.2	6.9	2.5	*	*	*

Black, female

2006 4/.....	1.0	*	*	*	*	0.7	1.9	2.0	1.3	*	*	*
2005 4/.....	1.1	*	*	*	*	1.0	2.4	2.1	*	*	*	*
2004 4/.....	1.1	*	*	*	*	0.9	2.2	2.2	*	*	*	*
2003 4/.....	0.9	*	*	*	*	*	2.3	1.9	*	*	*	*
2002.....	1.2	*	*	*	*	1.2	2.6	2.4	*	*	*	*
2001.....	0.8	*	*	*	*	0.9	1.8	1.6	*	*	*	*
2000.....	0.9	*	*	*	*	0.8	2.3	1.8	*	*	*	*
1999.....	0.9	*	*	*	*	0.9	2.2	1.6	*	*	*	*







Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes

2006.....	0.8	0.6	0.1	*	0.1	0.2	0.3	0.6	1.1	2.5	5.3	8.3
2005.....	0.9	*	*	*	0.1	0.2	0.3	0.7	1.2	2.6	5.8	9.1
2004.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.1	10.6
2003.....	1.0	*	0.2	*	0.1	0.2	0.4	0.7	1.4	2.9	6.2	11.2
2002.....	1.0	*	*	0.1	0.1	0.2	0.4	0.7	1.3	3.1	6.2	11.3
2001.....	1.1	0.6	0.2	*	0.1	0.2	0.4	0.7	1.5	3.1	6.9	12.8
2000.....	1.1	*	0.1	0.1	0.1	0.2	0.3	0.8	1.5	3.5	6.7	13.6
1999.....	1.0	1.4	0.2	*	0.1	0.2	0.3	0.6	1.3	3.2	6.6	12.7

All races 3/, male

2006.....	0.8	*	*	*	0.1	0.1	0.2	0.7	1.1	2.4	6.0	8.8
2005.....	0.8	*	*	*	*	0.1	0.3	0.6	1.3	2.9	6.3	10.8
2004.....	0.9	*	*	*	0.1	0.2	0.3	0.7	1.6	3.3	7.4	11.9
2003.....	0.9	*	*	*	0.1	0.2	0.4	0.7	1.5	3.2	6.9	12.9
2002.....	0.9	*	*	0.1	*	0.2	0.4	0.6	1.3	3.1	6.3	12.3
2001.....	1.0	*	*	*	*	0.2	0.4	0.7	1.6	3.2	7.7	14.5
2000.....	1.0	*	*	*	*	0.2	0.3	0.8	1.6	4.0	8.2	16.0
1999.....	1.0	1.6	*	*	0.1	0.2	0.3	0.5	1.4	3.4	7.8	16.5

All races 3/, female

2006.....	0.9	*	*	*	0.1	0.2	0.3	0.6	1.1	2.6	4.8	8.1
2005.....	1.0	*	*	*	*	0.2	0.4	0.7	1.1	2.3	5.5	8.3
2004.....	1.0	*	*	*	0.1	0.2	0.4	0.7	1.2	2.8	5.2	9.9
2003.....	1.0	*	*	*	*	0.2	0.4	0.6	1.3	2.7	5.7	10.5
2002.....	1.1	*	*	*	*	0.2	0.4	0.7	1.3	3.0	6.2	10.9
2001.....	1.1	*	*	*	0.1	0.2	0.4	0.7	1.4	3.0	6.3	12.1
2000.....	1.1	*	*	*	*	0.2	0.3	0.8	1.5	3.1	5.7	12.7
1999.....	1.1	1.2	*	*	*	0.2	0.3	0.6	1.2	3.0	5.9	11.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

White, both sexes

2006 4/.....	0.8	*	*	*	0.1	0.1	0.2	0.6	1.0	2.4	5.1	8.3
2005 4/.....	0.9	*	*	*	0.1	0.2	0.3	0.6	1.1	2.4	5.7	9.2
2004 4/.....	1.0	*	*	*	0.1	0.2	0.3	0.7	1.3	2.9	5.9	10.4
2003 4/.....	1.0	*	*	*	0.1	0.2	0.4	0.6	1.2	2.8	6.1	11.0
2002.....	1.0	*	*	0.1	0.1	0.2	0.3	0.6	1.2	3.0	6.1	11.2
2001.....	1.1	*	*	*	0.1	0.2	0.3	0.6	1.4	2.8	6.8	12.1
2000.....	1.1	*	*	0.1	0.1	0.2	0.3	0.7	1.4	3.3	6.4	13.1
1999.....	1.0	1.2	0.2	*	0.1	0.2	0.3	0.5	1.2	3.0	6.3	12.1

White, male

2006 4/.....	0.8	*	*	*	*	*	0.2	0.6	0.9	2.4	6.0	8.8
2005 4/.....	0.9	*	*	*	*	0.1	0.3	0.5	1.2	2.8	6.2	11.2
2004 4/.....	1.0	*	*	*	*	0.2	0.3	0.7	1.5	3.3	7.3	11.8
2003 4/.....	1.0	*	*	*	0.1	0.2	0.4	0.6	1.4	3.1	7.0	12.8
2002.....	0.9	*	*	*	*	0.2	0.4	0.6	1.2	2.9	6.3	11.8
2001.....	0.9	*	*	*	*	0.2	0.3	0.6	1.4	2.8	7.7	13.6
2000.....	1.1	*	*	*	*	0.2	0.3	0.7	1.4	3.7	8.2	15.8
1999.....	1.0	1.4	*	*	*	0.2	0.3	0.5	1.3	3.2	7.5	15.6

White, female

2006 4/.....	0.9	*	*	*	*	0.2	0.2	0.5	1.0	2.5	4.5	8.0
2005 4/.....	1.0	*	*	*	*	0.2	0.3	0.6	1.0	2.0	5.3	8.3
2004 4/.....	1.0	*	*	*	*	0.2	0.3	0.6	1.1	2.5	4.9	9.8
2003 4/.....	1.0	*	*	*	*	0.2	0.3	0.5	1.1	2.5	5.4	10.2
2002.....	1.1	*	*	*	*	0.1	0.3	0.5	1.2	3.0	6.0	10.9
2001.....	1.2	*	*	*	*	0.2	0.3	0.6	1.3	2.7	6.2	11.5
2000.....	1.1	*	*	*	*	0.2	0.3	0.7	1.3	3.0	5.3	12.0
1999.....	1.1	*	*	*	*	0.1	0.3	0.6	1.1	2.9	5.6	10.7



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Complications of medical and surgical care (Y40-Y84,Y88)

Black, both sexes

2006 4/.....	1.1	*	*	*	*	*	0.5	1.3	2.6	4.0	8.7	10.4
2005 4/.....	1.1	*	*	*	*	*	0.5	1.5	2.5	5.3	9.0	9.1
2004 4/.....	1.1	*	*	*	*	*	0.8	1.2	2.6	5.0	9.3	13.8
2003 4/.....	1.1	*	*	*	*	*	0.6	1.4	2.7	4.6	8.8	15.2
2002.....	1.2	*	*	*	*	*	0.8	1.6	2.7	4.6	8.3	14.0
2001.....	1.4	*	*	*	*	0.4	0.9	1.6	3.4	6.3	9.2	23.3
2000.....	1.3	*	*	*	*	0.4	0.5	1.4	3.5	6.1	10.5	21.0
1999.....	1.2	*	*	*	*	*	0.6	1.3	2.6	5.2	10.6	20.1

Black, male

2006 4/.....	0.9	*	*	*	*	*	*	1.4	2.6	4.0	9.5	*
2005 4/.....	1.0	*	*	*	*	*	*	1.5	3.1	4.9	9.8	*
2004 4/.....	0.9	*	*	*	*	*	*	1.0	2.9	4.6	10.0	*
2003 4/.....	1.0	*	*	*	*	*	*	1.8	2.8	3.9	7.6	*
2002.....	1.0	*	*	*	*	*	*	1.3	2.5	5.3	7.2	*
2001.....	1.4	*	*	*	*	*	1.0	1.8	4.2	7.8	9.1	29.5
2000.....	1.1	*	*	*	*	*	*	1.3	3.1	7.6	10.3	*
1999.....	1.1	*	*	*	*	*	*	1.4	2.8	5.9	12.3	23.4

Black, female

2006 4/.....	1.2	*	*	*	*	*	0.7	1.3	2.7	4.1	8.3	10.2
2005 4/.....	1.2	*	*	*	*	*	0.7	1.6	2.0	5.6	8.5	9.0
2004 4/.....	1.3	*	*	*	*	*	1.0	1.4	2.3	5.2	9.0	13.4
2003 4/.....	1.3	*	*	*	*	*	0.7	1.1	2.5	5.0	9.4	15.4
2002.....	1.3	*	*	*	*	*	1.1	1.8	2.7	4.1	8.9	11.5
2001.....	1.4	*	*	*	*	*	0.7	1.3	2.7	5.3	9.3	21.0
2000.....	1.5	*	*	*	*	*	*	1.5	3.8	5.0	10.6	21.9
1999.....	1.3	*	*	*	*	*	*	1.2	2.4	4.8	9.6	18.8

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 6/

All races 3/, both sexes

2006.....	7.4	*	*	*	0.4	1.6	7.4	17.6	19.4	14.7	9.7	4.9
2005.....	7.3	*	*	*	0.4	1.3	7.5	17.6	19.5	15.1	9.2	4.6
2004.....	7.2	*	*	*	0.3	1.5	7.6	17.3	18.7	15.7	9.3	4.3
2003.....	7.1	*	*	*	0.3	1.5	8.1	17.3	18.5	15.1	9.2	4.1
2002.....	7.0	*	*	*	0.3	1.5	8.1	16.8	18.4	15.5	9.3	4.4
2001.....	7.1	*	*	*	0.3	1.6	8.3	17.2	18.2	15.5	9.6	5.0
2000.....	7.0	*	*	*	0.2	1.6	8.5	16.3	18.7	15.8	9.9	5.4
1999.....	7.0	*	*	*	0.3	1.6	8.5	16.4	18.7	15.9	10.6	5.5

All races 3/, male

2006.....	11.2	*	*	*	0.6	2.4	10.3	26.5	31.1	24.9	17.1	9.5
2005.....	11.1	*	*	*	0.6	2.0	10.6	26.7	31.6	25.0	17.0	9.3
2004.....	11.0	*	*	*	0.5	2.3	11.0	26.4	30.2	26.5	17.1	9.0
2003.....	10.9	*	*	*	0.5	2.3	11.4	27.0	29.8	25.4	16.8	7.8
2002.....	10.8	*	*	*	0.4	2.3	11.5	26.3	30.0	26.0	16.6	9.7
2001.....	10.8	*	*	*	0.4	2.3	11.9	26.6	29.7	26.3	17.2	11.6
2000.....	10.9	*	*	*	0.4	2.5	12.5	25.9	30.0	27.1	18.8	11.9
1999.....	10.9	*	*	*	0.4	2.3	12.6	26.2	30.5	27.4	19.4	11.8

All races 3/, female

2006.....	3.7	*	*	*	0.1	0.7	4.5	8.9	8.5	6.1	4.7	2.7
2005.....	3.6	*	*	*	0.1	0.7	4.3	8.8	8.4	6.8	3.9	2.5
2004.....	3.5	*	*	*	0.2	0.7	4.2	8.6	8.1	6.6	3.9	2.3
2003.....	3.4	*	*	*	0.1	0.6	4.7	7.9	8.1	6.5	4.1	2.5
2002.....	3.4	*	*	*	*	0.7	4.7	7.7	7.6	6.8	4.5	2.1
2001.....	3.4	*	*	*	*	0.9	4.8	8.1	7.5	6.6	4.6	2.3
2000.....	3.2	*	*	*	*	0.7	4.5	7.0	8.3	6.6	4.1	2.7
1999.....	3.2	*	*	*	0.1	0.9	4.5	7.0	7.7	6.5	4.8	3.0

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 6/

White, both sexes

2006 4/.....	7.8	*	*	*	0.4	1.6	7.7	18.3	19.7	14.8	10.0	5.0
2005 4/.....	7.7	*	*	*	0.4	1.4	7.8	18.0	19.7	15.3	9.2	4.6
2004 4/.....	7.5	*	*	*	0.3	1.5	7.9	17.8	18.5	15.4	9.4	4.5
2003 4/.....	7.4	*	*	*	0.3	1.5	8.2	17.4	18.2	15.2	9.3	4.2
2002.....	7.2	*	*	*	0.3	1.5	8.3	16.9	17.8	15.4	9.3	4.3
2001.....	7.2	*	*	*	0.2	1.6	8.3	16.8	17.6	15.3	9.6	4.8
2000.....	7.1	*	*	*	0.2	1.6	8.4	15.8	18.0	15.6	9.8	5.3
1999.....	7.0	*	*	*	0.3	1.6	8.2	15.7	17.7	15.3	10.3	5.3

White, male

2006 4/.....	11.8	*	*	*	0.6	2.5	10.8	27.4	31.2	24.7	17.5	9.7
2005 4/.....	11.7	*	*	*	0.6	2.0	11.1	27.1	31.7	24.9	16.8	9.4
2004 4/.....	11.5	*	*	*	0.5	2.3	11.4	26.9	29.5	25.6	17.3	9.3
2003 4/.....	11.3	*	*	*	0.5	2.4	11.8	26.8	29.0	25.4	16.8	7.8
2002.....	11.1	*	*	*	0.4	2.3	11.8	26.4	29.0	25.4	16.5	9.3
2001.....	11.0	*	*	*	0.4	2.3	11.7	26.0	28.4	25.4	17.2	11.1
2000.....	11.0	*	*	*	0.4	2.5	12.3	25.3	28.8	26.4	18.6	11.6
1999.....	10.9	*	*	*	0.4	2.3	12.2	25.2	28.9	26.0	18.9	11.4

White, female

2006 4/.....	3.9	*	*	*	0.1	0.6	4.6	9.2	8.7	6.3	4.8	2.8
2005 4/.....	3.7	*	*	*	*	0.7	4.3	8.9	8.4	7.0	4.0	2.5
2004 4/.....	3.6	*	*	*	0.1	0.7	4.3	8.7	8.0	6.7	4.0	2.3
2003 4/.....	3.5	*	*	*	*	0.7	4.7	8.0	8.0	6.5	4.3	2.5
2002.....	3.4	*	*	*	*	0.7	4.7	7.6	7.3	7.0	4.5	2.2
2001.....	3.4	*	*	*	*	0.8	4.8	7.6	7.4	6.7	4.6	2.2
2000.....	3.2	*	*	*	*	0.7	4.4	6.5	7.8	6.5	4.1	2.7
1999.....	3.1	*	*	*	*	0.9	4.1	6.4	7.2	6.5	4.6	2.9

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages 1/	Under 1 year 2/	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	-------------	-----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K85.2,K86.0,R78.0,X45,X65,Y15) 6/

Black, both sexes

2006 4/.....	5.5	*	*	*	*	1.0	5.3	14.7	20.4	15.7	8.2	*
2005 4/.....	5.9	*	*	*	*	0.9	5.6	16.9	21.6	16.5	9.8	5.6
2004 4/.....	6.1	*	*	*	*	1.0	5.7	17.1	23.5	20.1	7.6	*
2003 4/.....	6.3	*	*	*	*	1.1	6.8	18.4	24.2	16.9	8.9	*
2002.....	6.4	*	*	*	*	1.3	6.7	18.5	25.3	18.3	11.2	*
2001.....	7.3	*	*	*	*	1.3	8.7	22.5	25.9	20.5	11.1	*
2000.....	7.4	*	*	*	*	1.7	9.2	22.7	26.4	21.0	11.5	6.3
1999.....	7.8	*	*	*	*	1.3	9.9	23.4	29.1	23.2	14.6	7.7

Black, male

2006 4/.....	8.5	*	*	*	*	1.2	7.0	23.0	35.9	30.2	16.0	*
2005 4/.....	9.1	*	*	*	*	1.2	7.7	26.9	36.7	30.8	20.8	*
2004 4/.....	9.7	*	*	*	*	1.6	8.8	27.0	41.5	39.5	14.5	*
2003 4/.....	10.0	*	*	*	*	1.8	9.6	31.2	42.2	30.7	18.9	*
2002.....	10.0	*	*	*	*	2.0	9.4	29.3	43.5	35.8	21.8	*
2001.....	11.6	*	*	*	*	1.9	13.2	35.6	46.3	40.0	21.8	*
2000.....	11.4	*	*	*	*	2.5	13.6	36.0	43.8	39.9	24.2	*
1999.....	12.2	*	*	*	*	1.8	14.4	37.7	51.0	45.3	28.3	*

Black, female

2006 4/.....	2.8	*	*	*	*	0.7	3.8	7.5	7.9	5.3	3.8	*
2005 4/.....	3.0	*	*	*	*	*	3.8	8.3	9.4	6.3	3.6	*
2004 4/.....	2.8	*	*	*	*	*	2.9	8.6	9.0	6.2	3.8	*
2003 4/.....	2.9	*	*	*	*	*	4.3	7.4	9.8	7.1	3.2	*
2002.....	3.2	*	*	*	*	*	4.4	9.3	10.7	5.8	5.2	*
2001.....	3.5	*	*	*	*	0.8	4.6	11.1	9.4	6.8	5.1	*
2000.....	3.7	*	*	*	*	1.0	5.2	11.3	12.3	7.7	4.3	*
1999.....	3.9	*	*	*	*	0.9	5.9	11.2	11.6	7.7	6.9	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,K85.3,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

All races 3/, both sexes

2006.....	12.8	1.1	0.2	0.1	8.6	16.8	23.3	26.7	12.2	5.3	6.0	8.1
2005.....	11.3	0.9	0.2	0.1	7.4	14.3	21.3	23.6	10.7	4.8	5.4	7.6
2004.....	10.5	0.7	0.2	0.2	7.0	12.6	20.9	21.7	9.1	4.3	4.8	6.3
2003.....	9.9	0.6	0.2	0.1	6.4	12.2	20.6	20.0	8.0	4.1	4.2	6.0
2002.....	9.0	0.7	0.2	0.1	5.5	11.1	19.7	17.9	6.9	3.6	3.8	5.7
2001.....	7.6	0.5	0.2	0.1	4.5	9.4	17.0	14.8	5.4	3.0	3.5	5.1
2000.....	7.0	*	*	0.1	4.0	8.8	16.0	13.2	4.9	2.6	3.5	5.7
1999.....	6.9	0.6	0.2	0.1	3.5	8.9	15.7	12.6	4.9	3.0	3.8	4.8

All races 3/, male

2006.....	16.6	1.1	0.2	0.2	12.7	23.4	29.4	33.3	14.9	5.8	6.9	10.7
2005.....	14.5	*	0.3	0.1	10.7	19.4	26.6	29.4	13.1	5.3	6.7	9.9
2004.....	13.4	*	*	0.1	10.2	17.5	26.0	27.0	10.7	4.7	5.3	8.6
2003.....	12.9	*	0.3	0.1	9.2	16.6	26.5	25.7	9.4	4.6	4.9	7.8
2002.....	11.8	1.0	*	0.1	7.8	15.5	25.6	23.0	8.1	3.7	4.2	7.3
2001.....	10.2	*	0.3	0.1	6.6	13.2	22.7	19.7	6.3	3.5	3.8	5.7
2000.....	9.5	*	*	0.1	5.8	12.5	21.8	18.2	5.4	2.8	3.8	7.5
1999.....	9.4	*	*	*	5.0	12.7	21.9	17.5	6.0	3.4	3.8	6.2

All races 3/, female

2006.....	9.1	1.0	*	0.1	4.2	10.0	17.1	20.3	9.8	4.9	5.4	6.8
2005.....	8.2	*	*	0.1	3.8	8.9	16.0	18.1	8.4	4.3	4.5	6.6
2004.....	7.6	*	*	0.2	3.6	7.7	15.8	16.5	7.7	3.9	4.4	5.2
2003.....	7.0	*	*	0.1	3.3	7.6	14.8	14.6	6.7	3.7	3.7	5.2
2002.....	6.3	*	0.3	*	3.0	6.6	13.9	13.1	5.7	3.4	3.5	4.9
2001.....	5.1	*	*	0.1	2.4	5.6	11.5	10.0	4.6	2.6	3.3	4.9
2000.....	4.6	*	*	*	2.1	5.1	10.2	8.5	4.4	2.5	3.2	4.9
1999.....	4.4	*	*	*	2.0	5.0	9.7	8.0	3.8	2.7	3.8	4.3

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year

All ages 1/ Under 1 year 2/ 1-4 years 5-14 years 15-24 years 25-34 years 35-44 years 45-54 years 55-64 years 65-74 years 75-84 years 85 years and over

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,K85.3,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

White, both sexes

2006 4/.....	13.6	0.9	0.2	0.1	10.3	19.2	24.7	26.7	11.7	5.2	6.2	8.3
2005 4/.....	12.0	0.8	*	0.1	8.8	16.1	22.7	23.6	10.3	4.8	5.6	8.0
2004 4/.....	11.1	*	*	0.2	8.3	14.3	22.4	21.7	8.8	4.3	4.9	6.4
2003 4/.....	10.4	*	0.2	0.1	7.6	13.6	21.7	20.0	7.8	4.1	4.5	6.3
2002.....	9.4	*	*	0.1	6.4	12.3	20.6	17.5	6.6	3.7	3.9	5.8
2001.....	7.8	*	*	0.1	5.2	10.3	17.5	14.0	5.1	3.0	3.6	5.1
2000.....	7.1	*	*	0.1	4.6	9.5	16.1	12.4	4.8	2.7	3.7	5.6
1999.....	6.9	*	*	0.1	3.9	9.4	15.6	11.7	4.6	3.0	4.0	4.8

White, male

2006 4/.....	17.3	*	*	0.1	15.3	26.3	30.8	32.3	13.1	5.2	7.1	11.2
2005 4/.....	15.3	*	*	0.1	12.8	21.8	28.3	28.6	11.8	5.1	6.8	10.6
2004 4/.....	14.1	*	*	0.1	12.1	19.7	27.7	26.5	9.5	4.4	5.3	8.7
2003 4/.....	13.5	*	*	*	11.0	18.6	27.8	25.1	8.7	4.3	5.2	7.8
2002.....	12.2	*	*	0.1	9.1	17.1	26.6	21.9	7.2	3.6	4.2	7.6
2001.....	10.4	*	*	*	7.6	14.4	23.2	18.1	5.5	3.4	3.8	5.6
2000.....	9.6	*	*	0.1	6.7	13.3	21.9	16.5	4.9	2.7	4.0	7.2
1999.....	9.3	*	*	*	5.5	13.3	21.7	15.6	5.2	3.3	3.9	6.4

White, female

2006 4/.....	9.9	*	*	0.1	5.0	11.6	18.4	21.2	10.3	5.2	5.6	6.9
2005 4/.....	8.8	*	*	*	4.4	10.0	17.0	18.7	8.9	4.6	4.7	6.8
2004 4/.....	8.2	*	*	0.2	4.2	8.6	17.0	17.0	8.1	4.3	4.7	5.3
2003 4/.....	7.4	*	*	*	3.9	8.3	15.6	15.0	7.0	3.9	4.0	5.6
2002.....	6.7	*	*	*	3.5	7.3	14.5	13.2	6.0	3.7	3.8	4.9
2001.....	5.4	*	*	*	2.7	6.0	11.7	10.1	4.8	2.7	3.5	4.9
2000.....	4.7	*	*	*	2.3	5.5	10.2	8.4	4.7	2.7	3.4	4.9
1999.....	4.5	*	*	*	2.2	5.2	9.5	7.9	4.1	2.8	4.1	4.2

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,K85.3,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R50.2,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 6/

Black, both sexes

2006 4/.....	12.1	*	*	*	2.5	10.1	23.1	35.0	21.1	6.9	4.5	6.5
2005 4/.....	10.5	*	*	*	2.1	9.4	20.0	30.9	17.5	5.2	4.3	*
2004 4/.....	9.4	*	*	*	2.2	7.9	18.7	28.1	14.4	4.6	3.9	*
2003 4/.....	9.2	*	*	*	1.8	8.1	20.8	26.9	11.7	5.0	2.2	*
2002.....	9.2	*	*	*	2.1	8.5	20.7	26.8	11.5	3.1	2.7	*
2001.....	8.5	*	*	*	1.9	7.7	20.0	25.2	8.8	3.2	2.6	6.1
2000.....	8.3	*	*	*	2.0	8.3	20.1	23.8	7.5	3.0	2.4	7.5
1999.....	8.6	*	*	*	2.1	9.1	20.9	23.7	8.3	3.5	2.2	*

Black, male

2006 4/.....	17.3	*	*	*	3.3	15.5	31.7	51.3	35.1	12.5	5.7	*
2005 4/.....	14.3	*	*	*	2.9	12.9	25.3	44.5	29.2	9.0	6.6	*
2004 4/.....	12.8	*	*	*	3.2	11.1	23.8	39.1	24.1	8.5	7.0	*
2003 4/.....	12.7	*	*	*	2.5	10.7	27.7	38.9	19.0	8.6	*	*
2002.....	12.8	*	*	*	2.9	12.0	27.7	39.2	19.9	5.4	*	*
2001.....	12.2	*	*	*	2.9	10.8	27.0	39.3	15.1	5.4	*	*
2000.....	12.0	*	*	*	2.7	12.0	28.2	37.6	12.4	5.3	*	*
1999.....	12.7	*	*	*	3.2	13.3	30.1	38.2	15.2	5.4	*	*

Black, female

2006 4/.....	7.4	*	*	*	1.7	5.1	15.5	20.9	9.9	2.9	3.8	7.3
2005 4/.....	7.0	*	*	*	1.4	6.3	15.4	19.2	8.0	2.4	*	*
2004 4/.....	6.3	*	*	*	1.3	5.0	14.2	18.6	6.6	*	*	*
2003 4/.....	6.1	*	*	*	1.1	5.8	14.6	16.6	5.8	2.4	*	*
2002.....	5.8	*	*	*	1.3	5.3	14.5	16.1	4.8	*	*	*
2001.....	5.1	*	*	*	0.9	4.8	13.8	13.0	3.8	*	*	*
2000.....	4.9	*	*	*	1.3	4.9	13.0	11.9	3.5	*	*	*
1999.....	4.8	*	*	*	1.0	5.2	12.9	11.3	2.8	2.1	*	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

All races 3/, both sexes

2006.....	10.3	*	0.3	0.9	16.9	15.3	11.5	11.2	9.8	10.0	12.9	11.5
2005.....	10.4	*	0.4	0.8	16.2	15.7	11.6	11.2	9.8	10.3	13.7	12.0
2004.....	10.1	*	0.3	0.7	15.7	15.0	11.3	11.0	9.8	10.2	13.3	11.9
2003.....	10.4	*	0.3	0.8	16.6	15.5	11.5	11.2	10.1	10.4	13.5	12.5
2002.....	10.5	*	0.4	0.8	16.7	15.4	12.1	10.8	10.2	10.9	14.4	12.5
2001.....	10.4	*	0.5	0.8	16.7	15.5	11.7	10.5	10.1	10.9	14.3	12.8
2000.....	10.2	*	0.3	0.9	16.8	14.5	11.9	10.5	9.4	10.6	13.9	14.2
1999.....	10.3	*	0.4	1.0	17.6	14.9	11.6	10.2	9.7	11.0	14.2	13.5

All races 3/, male

2006.....	18.1	*	0.5	1.2	29.8	26.6	18.9	18.5	17.0	19.2	29.1	33.6
2005.....	18.3	*	0.5	1.2	28.7	27.2	19.2	18.9	17.4	19.7	30.8	35.4
2004.....	17.6	*	0.4	1.1	27.5	26.1	18.7	18.3	17.1	19.7	29.8	35.9
2003.....	18.3	*	0.3	1.2	29.2	27.1	19.1	18.8	17.7	20.3	30.2	37.8
2002.....	18.4	*	0.5	1.2	29.3	26.5	20.1	18.2	18.0	21.3	32.9	38.9
2001.....	18.2	*	0.5	1.2	29.6	26.7	19.2	17.8	17.4	21.1	32.8	40.2
2000.....	17.8	*	0.4	1.4	29.4	24.9	19.4	17.6	16.3	20.3	32.2	44.7
1999.....	18.1	*	0.5	1.5	30.6	25.5	18.9	17.0	17.1	21.5	32.8	43.3

All races 3/, female

2006.....	2.8	*	*	0.4	3.2	3.7	4.1	4.0	3.2	2.2	1.8	1.1
2005.....	2.7	*	0.3	0.4	3.0	3.8	4.0	3.7	2.8	2.5	2.1	1.3
2004.....	2.7	*	0.3	0.3	3.2	3.7	3.9	4.0	3.1	2.2	2.3	1.0
2003.....	2.7	*	0.3	0.3	3.3	3.6	4.0	3.8	2.9	2.2	2.2	1.3
2002.....	2.8	*	0.3	0.5	3.5	4.0	4.2	3.6	3.1	2.3	2.1	1.1
2001.....	2.8	*	0.4	0.4	3.2	4.0	4.3	3.5	3.3	2.4	2.2	1.3
2000.....	2.8	*	*	0.4	3.5	4.0	4.4	3.6	3.0	2.5	2.0	1.7
1999.....	2.9	*	0.4	0.5	3.9	4.1	4.4	3.6	2.9	2.4	2.1	1.5



Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1/1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	----------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

White, both sexes												
2006 4/.....	9.0	*	0.2	0.7	10.6	10.5	10.6	11.4	10.6	10.9	14.0	12.2
2005 4/.....	9.1	*	0.3	0.7	10.5	11.2	10.7	11.6	10.7	11.2	14.8	13.0
2004 4/.....	9.0	*	*	0.6	10.7	10.9	10.6	11.4	10.4	11.0	14.3	12.8
2003 4/.....	9.2	*	0.2	0.6	11.1	11.2	10.8	11.5	10.8	11.3	14.4	13.5
2002.....	9.3	*	0.3	0.7	11.3	11.2	11.3	11.1	10.9	11.8	15.3	13.4
2001.....	9.4	*	0.4	0.8	11.4	11.9	11.1	10.9	10.7	11.7	15.2	13.8
2000.....	9.1	*	0.2	0.9	11.5	10.9	11.2	10.6	9.9	11.2	15.0	15.2
1999.....	9.2	*	0.3	0.9	12.0	11.5	11.0	10.3	10.1	11.8	15.1	14.5

White, male												
2006 4/.....	15.4	*	*	1.0	18.4	17.7	17.0	18.7	18.1	20.7	31.3	35.8
2005 4/.....	15.8	*	0.3	1.0	18.2	18.6	17.2	19.3	18.7	21.2	33.0	38.0
2004 4/.....	15.5	*	*	0.9	18.4	18.2	17.1	18.7	17.9	20.9	31.8	38.9
2003 4/.....	16.0	*	*	1.0	19.2	18.8	17.5	19.2	18.9	21.9	32.3	41.0
2002.....	16.1	*	0.4	0.9	19.4	18.5	18.5	18.4	19.0	22.9	34.8	41.7
2001.....	16.2	*	0.4	1.1	19.5	19.9	18.0	18.3	18.3	22.4	34.7	43.3
2000.....	15.6	*	*	1.3	19.6	18.1	17.9	17.7	17.0	21.4	34.5	48.4
1999.....	15.9	*	*	1.3	20.5	19.2	17.6	17.2	17.6	22.8	34.7	46.8

White, female												
2006 4/.....	2.6	*	*	0.4	2.3	3.0	4.0	4.3	3.5	2.4	2.0	1.1
2005 4/.....	2.6	*	*	0.3	2.3	3.4	4.1	3.9	3.1	2.6	2.3	1.4
2004 4/.....	2.7	*	*	0.3	2.6	3.2	4.0	4.3	3.3	2.5	2.4	1.1
2003 4/.....	2.6	*	*	0.2	2.5	3.1	4.0	4.0	3.2	2.3	2.3	1.3
2002.....	2.7	*	*	0.4	2.6	3.5	4.1	3.8	3.3	2.4	2.2	1.2
2001.....	2.7	*	*	0.4	2.7	3.6	4.1	3.7	3.6	2.6	2.3	1.4
2000.....	2.7	*	*	0.3	2.8	3.5	4.3	3.7	3.2	2.8	2.1	1.8
1999.....	2.8	*	*	0.5	3.0	3.6	4.3	3.6	3.1	2.6	2.2	1.6

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Black, both sexes

2006 4/.....	21.3	*	1.1	1.7	50.3	47.2	20.9	12.4	6.7	5.9	5.3	6.2
2005 4/.....	20.4	*	1.0	1.7	47.2	46.2	20.9	11.7	6.1	6.5	5.7	*
2004 4/.....	19.3	*	1.2	1.5	44.1	43.8	18.8	11.1	8.0	6.2	6.6	*
2003 4/.....	20.1	*	0.9	1.7	47.7	45.4	19.0	11.4	6.8	6.0	6.1	*
2002.....	20.2	*	*	1.5	47.7	44.7	20.5	11.8	7.2	5.8	7.1	*
2001.....	19.3	*	0.9	1.2	48.1	40.7	19.0	10.6	6.9	6.6	7.5	*
2000.....	19.3	*	*	1.4	48.1	39.4	19.2	12.1	6.7	6.9	5.5	*
1999.....	19.4	*	1.1	1.6	49.9	37.7	18.3	11.5	8.2	6.6	7.5	*

Black, male

2006 4/.....	40.0	*	*	2.6	91.8	89.0	38.1	23.0	13.1	12.5	14.1	*
2005 4/.....	38.7	*	*	2.4	86.8	88.4	38.7	21.4	11.5	12.8	14.5	*
2004 4/.....	36.4	*	*	2.3	80.7	83.6	35.1	20.2	14.8	14.4	15.9	*
2003 4/.....	37.8	*	*	2.5	87.6	87.2	34.8	21.1	12.9	11.7	13.8	*
2002.....	37.8	*	*	2.1	87.1	85.6	36.9	21.4	13.5	12.1	17.5	*
2001.....	36.4	*	*	1.8	90.3	77.7	33.2	19.1	13.7	13.8	18.0	*
2000.....	36.1	*	*	2.2	89.3	74.8	34.3	21.7	12.6	15.1	12.1	*
1999.....	36.0	*	*	2.3	91.4	71.3	32.2	20.4	16.5	14.0	18.1	*

Worktable 290R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 10-year age groups, race and sex: United States, 1999-2006

[Rates per 100,000 population in specified group. Rates are based on populations enumerated as of April 1 for 2000 and estimated as of July 1 for all other years. Race categories are consistent with the 1977 Office of Management and Budget standards. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), race, sex, and year	All ages	Under 1 year	1-4 years	5-14 years	15-24 years	25-34 years	35-44 years	45-54 years	55-64 years	65-74 years	75-84 years	85 years and over
---	----------	--------------	-----------	------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 6/

Black, female

2006 4/.....	4.1	*	*	0.8	7.7	8.4	5.6	3.3	1.6	*	*	*
2005 4/.....	3.7	*	*	0.9	6.7	7.1	5.0	3.3	1.8	2.0	*	*
2004 4/.....	3.7	*	*	0.7	6.9	7.1	4.3	3.3	2.6	*	*	*
2003 4/.....	3.9	*	*	0.9	7.4	7.3	5.0	3.1	2.0	2.0	*	*
2002.....	4.2	*	*	0.9	8.1	7.5	5.9	3.5	2.1	*	*	*
2001.....	3.8	*	*	*	6.1	7.3	6.5	3.2	1.5	*	*	*
2000.....	4.0	*	*	*	7.6	7.4	5.7	3.8	1.9	*	*	*
1999.....	4.4	*	*	0.8	9.0	7.4	6.1	3.8	1.6	*	*	*

\* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Includes races other than white and black.

4/ Multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming in 2006, by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming in 2005, by California, Hawaii, Idaho, Maine, Michigan, Minnesota, Montana, New Hampshire, New Jersey, New York, Oklahoma, South Dakota, Washington, Wisconsin, and Wyoming in 2004, and by California, Hawaii, Idaho, Maine, Montana, New York, and Wisconsin in 2003. The multiple-race data for these areas were bridged to the single-race categories of the 1977 Office of Management and Budget standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2006 data - mortality."

5/ Figures include September 11, 2001 related deaths for which death certificates were filed as of 10/24/02; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2001."

6/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.