

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, all ages				
...	All causes	2,744,248	100.0	849.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	635,260	23.1	196.6
2	Malignant neoplasms (C00-C97)	598,038	21.8	185.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161,374	5.9	49.9
4	Chronic lower respiratory diseases (J40-J47)	154,596	5.6	47.8
5	Cerebrovascular diseases (I60-I69)	142,142	5.2	44.0
6	Alzheimer's disease (G30)	116,103	4.2	35.9
7	Diabetes mellitus (E10-E14)	80,058	2.9	24.8
8	Influenza and pneumonia (J09-J18)	51,537	1.9	15.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,046	1.8	15.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	44,965	1.6	13.9
11	Septicemia (A40-A41)	40,613	1.5	12.6
12	Chronic liver disease and cirrhosis (K70,K73-K74)	40,545	1.5	12.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	33,246	1.2	10.3
14	Parkinson's disease (G20-G21)	29,697	1.1	9.2
15	Pneumonitis due to solids and liquids (J69)	19,715	0.7	6.1
...	All other causes (residual)	546,313	19.9	169.1
All races and origins, both sexes, 1-4 years				
...	All causes	4,045	100.0	25.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,261	31.2	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	433	10.7	2.7
3	Malignant neoplasms (C00-C97)	377	9.3	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	8.4	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	118	2.9	0.7
6	Influenza and pneumonia (J09-J18)	103	2.5	0.6
7	Septicemia (A40-A41)	70	1.7	0.4
8	Certain conditions originating in the perinatal period (P00-P96)	60	1.5	0.4
9	Cerebrovascular diseases (I60-I69)	55	1.4	0.3
10	Chronic lower respiratory diseases (J40-J47)	51	1.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	49	1.2	0.3
12	Anemias (D50-D64)	20	0.5	0.1
13	Acute bronchitis and bronchiolitis (J20-J21)	18	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
15	Meningitis (G00,G03)	14	0.3	*
...	All other causes (residual)	1,062	26.3	6.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 5-9 years				
...	All causes	2,490	100.0	12.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	787	31.6	3.9
2	Malignant neoplasms (C00-C97)	449	18.0	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	203	8.2	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	5.6	0.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	3.1	0.4
6	Chronic lower respiratory diseases (J40-J47)	68	2.7	0.3
7	Influenza and pneumonia (J09-J18)	48	1.9	0.2
8	Septicemia (A40-A41)	40	1.6	0.2
9	Cerebrovascular diseases (I60-I69)	38	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	1.2	0.2
11	Anemias (D50-D64)	14	0.6	*
11	Certain conditions originating in the perinatal period (P00-P96)	14	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	11	0.4	*
14	Diabetes mellitus (E10-E14)	10	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.4	*
...	All other causes (residual)	551	22.1	2.7
All races and origins, both sexes, 10-14 years				
...	All causes	3,013	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	847	28.1	4.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	436	14.5	2.1
3	Malignant neoplasms (C00-C97)	431	14.3	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	4.9	0.7
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	146	4.8	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	111	3.7	0.5
7	Chronic lower respiratory diseases (J40-J47)	75	2.5	0.4
8	Cerebrovascular diseases (I60-I69)	50	1.7	0.2
9	Influenza and pneumonia (J09-J18)	39	1.3	0.2
10	Septicemia (A40-A41)	31	1.0	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	0.9	0.1
12	Diabetes mellitus (E10-E14)	16	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	12	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
15	Anemias (D50-D64)	10	0.3	*
...	All other causes (residual)	622	20.6	3.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 15-19 years				
...	All causes	10,812	100.0	51.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,152	38.4	19.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,117	19.6	10.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,816	16.8	8.6
4	Malignant neoplasms (C00-C97)	596	5.5	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	293	2.7	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	197	1.8	0.9
7	Chronic lower respiratory diseases (J40-J47)	80	0.7	0.4
8	Cerebrovascular diseases (I60-I69)	62	0.6	0.3
9	Diabetes mellitus (E10-E14)	54	0.5	0.3
9	Influenza and pneumonia (J09-J18)	54	0.5	0.3
11	Septicemia (A40-A41)	42	0.4	0.2
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.3	0.2
12	Anemias (D50-D64)	37	0.3	0.2
14	Legal intervention (Y35,Y89.0)	26	0.2	0.1
15	Pregnancy, childbirth and the puerperium (O00-O99)	24	0.2	0.1
...	All other causes (residual)	1,225	11.3	5.8
All races and origins, both sexes, 20-24 years				
...	All causes	21,763	100.0	97.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,743	44.8	43.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,606	16.6	16.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,356	15.4	15.0
4	Malignant neoplasms (C00-C97)	835	3.8	3.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	656	3.0	2.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	0.9	0.9
7	Pregnancy, childbirth and the puerperium (O00-O99)	160	0.7	0.7
8	Diabetes mellitus (E10-E14)	157	0.7	0.7
9	Influenza and pneumonia (J09-J18)	135	0.6	0.6
10	Chronic lower respiratory diseases (J40-J47)	126	0.6	0.6
11	Septicemia (A40-A41)	101	0.5	0.5
12	Cerebrovascular diseases (I60-I69)	86	0.4	0.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	63	0.3	0.3
14	Legal intervention (Y35,Y89.0)	51	0.2	0.2
15	Anemias (D50-D64)	41	0.2	0.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	41	0.2	0.2
...	All other causes (residual)	2,415	11.1	10.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 25-29 years				
...	All causes	27,053	100.0	118.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,941	44.1	52.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,790	14.0	16.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,997	11.1	13.1
4	Malignant neoplasms (C00-C97)	1,382	5.1	6.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,226	4.5	5.4
6	Diabetes mellitus (E10-E14)	300	1.1	1.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	236	0.9	1.0
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	235	0.9	1.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	233	0.9	1.0
10	Pregnancy, childbirth and the puerperium (O00-O99)	229	0.8	1.0
11	Cerebrovascular diseases (I60-I69)	198	0.7	0.9
12	Septicemia (A40-A41)	174	0.6	0.8
13	Influenza and pneumonia (J09-J18)	168	0.6	0.7
14	Chronic lower respiratory diseases (J40-J47)	140	0.5	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	99	0.4	0.4
...	All other causes (residual)	3,705	13.7	16.2
All races and origins, both sexes, 30-34 years				
...	All causes	30,563	100.0	140.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,043	39.4	55.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,576	11.7	16.4
3	Malignant neoplasms (C00-C97)	2,409	7.9	11.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,379	7.8	10.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,219	7.3	10.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	689	2.3	3.2
7	Diabetes mellitus (E10-E14)	492	1.6	2.3
8	Cerebrovascular diseases (I60-I69)	377	1.2	1.7
9	Human immunodeficiency virus (HIV) disease (B20-B24)	313	1.0	1.4
10	Influenza and pneumonia (J09-J18)	284	0.9	1.3
11	Septicemia (A40-A41)	245	0.8	1.1
12	Pregnancy, childbirth and the puerperium (O00-O99)	243	0.8	1.1
13	Chronic lower respiratory diseases (J40-J47)	213	0.7	1.0
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	201	0.7	0.9
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	0.5	0.8
...	All other causes (residual)	4,714	15.4	21.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 35-39 years				
...	All causes	35,277	100.0	169.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,154	31.6	53.7
2	Malignant neoplasms (C00-C97)	4,071	11.5	19.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,882	11.0	18.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,597	10.2	17.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,920	5.4	9.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,107	3.1	5.3
7	Diabetes mellitus (E10-E14)	802	2.3	3.9
8	Cerebrovascular diseases (I60-I69)	699	2.0	3.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	418	1.2	2.0
10	Influenza and pneumonia (J09-J18)	354	1.0	1.7
11	Septicemia (A40-A41)	335	0.9	1.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	310	0.9	1.5
13	Chronic lower respiratory diseases (J40-J47)	243	0.7	1.2
14	Pregnancy, childbirth and the puerperium (O00-O99)	232	0.7	1.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	198	0.6	1.0
...	All other causes (residual)	5,955	16.9	28.7
All races and origins, both sexes, 40-44 years				
...	All causes	42,515	100.0	215.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,821	23.1	49.9
2	Malignant neoplasms (C00-C97)	6,832	16.1	34.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,595	15.5	33.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,433	8.1	17.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,744	4.1	8.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,449	3.4	7.4
7	Diabetes mellitus (E10-E14)	1,247	2.9	6.3
8	Cerebrovascular diseases (I60-I69)	1,152	2.7	5.8
9	Septicemia (A40-A41)	562	1.3	2.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	553	1.3	2.8
11	Influenza and pneumonia (J09-J18)	540	1.3	2.7
12	Chronic lower respiratory diseases (J40-J47)	449	1.1	2.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	433	1.0	2.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	293	0.7	1.5
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	223	0.5	1.1
...	All other causes (residual)	7,189	16.9	36.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 45-49 years				
...	All causes	65,585	100.0	313.1
1	Malignant neoplasms (C00-C97)	13,715	20.9	65.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,176	18.6	58.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,972	16.7	52.4
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,868	5.9	18.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,044	4.6	14.5
6	Diabetes mellitus (E10-E14)	2,278	3.5	10.9
7	Cerebrovascular diseases (I60-I69)	1,989	3.0	9.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,152	1.8	5.5
9	Chronic lower respiratory diseases (J40-J47)	1,083	1.7	5.2
10	Septicemia (A40-A41)	869	1.3	4.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	820	1.3	3.9
12	Influenza and pneumonia (J09-J18)	812	1.2	3.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	779	1.2	3.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	558	0.9	2.7
15	Viral hepatitis (B15-B19)	336	0.5	1.6
...	All other causes (residual)	11,134	17.0	53.2
All races and origins, both sexes, 50-54 years				
...	All causes	107,931	100.0	494.2
1	Malignant neoplasms (C00-C97)	27,576	25.5	126.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,851	20.2	100.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,405	11.5	56.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	5,320	4.9	24.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,569	4.2	20.9
6	Diabetes mellitus (E10-E14)	3,989	3.7	18.3
7	Cerebrovascular diseases (I60-I69)	3,364	3.1	15.4
8	Chronic lower respiratory diseases (J40-J47)	3,224	3.0	14.8
9	Septicemia (A40-A41)	1,603	1.5	7.3
10	Influenza and pneumonia (J09-J18)	1,321	1.2	6.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,320	1.2	6.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1,052	1.0	4.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,048	1.0	4.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,000	0.9	4.6
15	Viral hepatitis (B15-B19)	826	0.8	3.8
...	All other causes (residual)	17,463	16.2	80.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 55-59 years				
...	All causes	162,146	100.0	737.7
1	Malignant neoplasms (C00-C97)	49,179	30.3	223.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,175	21.1	155.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,386	7.6	56.4
4	Chronic lower respiratory diseases (J40-J47)	6,927	4.3	31.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	6,872	4.2	31.3
6	Diabetes mellitus (E10-E14)	6,070	3.7	27.6
7	Cerebrovascular diseases (I60-I69)	5,191	3.2	23.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,375	2.7	19.9
9	Septicemia (A40-A41)	2,564	1.6	11.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,278	1.4	10.4
11	Influenza and pneumonia (J09-J18)	2,089	1.3	9.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,617	1.0	7.4
13	Viral hepatitis (B15-B19)	1,384	0.9	6.3
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1,025	0.6	4.7
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	803	0.5	3.7
...	All other causes (residual)	25,211	15.5	114.7
All races and origins, both sexes, 60-64 years				
...	All causes	204,299	100.0	1,048.6
1	Malignant neoplasms (C00-C97)	67,185	32.9	344.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,435	21.7	228.1
3	Chronic lower respiratory diseases (J40-J47)	10,883	5.3	55.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,474	4.6	48.6
5	Diabetes mellitus (E10-E14)	8,181	4.0	42.0
6	Cerebrovascular diseases (I60-I69)	7,119	3.5	36.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,576	3.2	33.8
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,384	1.7	17.4
9	Septicemia (A40-A41)	3,377	1.7	17.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,372	1.7	17.3
11	Influenza and pneumonia (J09-J18)	2,945	1.4	15.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,124	1.0	10.9
13	Viral hepatitis (B15-B19)	1,612	0.8	8.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	944	0.5	4.8
15	Pneumonitis due to solids and liquids (J69)	926	0.5	4.8
...	All other causes (residual)	31,762	15.5	163.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 65-69 years				
...	All causes	246,655	100.0	1,466.4
1	Malignant neoplasms (C00-C97)	82,745	33.5	491.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54,155	22.0	322.0
3	Chronic lower respiratory diseases (J40-J47)	16,508	6.7	98.1
4	Diabetes mellitus (E10-E14)	9,892	4.0	58.8
5	Cerebrovascular diseases (I60-I69)	9,637	3.9	57.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,579	3.1	45.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	5,204	2.1	30.9
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,525	1.8	26.9
9	Septicemia (A40-A41)	4,246	1.7	25.2
10	Influenza and pneumonia (J09-J18)	3,791	1.5	22.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,584	1.0	15.4
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,537	1.0	15.1
13	Alzheimer's disease (G30)	2,012	0.8	12.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,380	0.6	8.2
15	Parkinson's disease (G20-G21)	1,353	0.5	8.0
...	All other causes (residual)	38,507	15.6	228.9
All races and origins, both sexes, 70-74 years				
...	All causes	265,425	100.0	2,247.4
1	Malignant neoplasms (C00-C97)	82,838	31.2	701.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	58,206	21.9	492.8
3	Chronic lower respiratory diseases (J40-J47)	21,895	8.2	185.4
4	Cerebrovascular diseases (I60-I69)	12,131	4.6	102.7
5	Diabetes mellitus (E10-E14)	10,110	3.8	85.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,471	2.4	54.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,379	2.0	45.5
8	Alzheimer's disease (G30)	4,749	1.8	40.2
9	Septicemia (A40-A41)	4,521	1.7	38.3
10	Influenza and pneumonia (J09-J18)	4,378	1.6	37.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	3,593	1.4	30.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,870	1.1	24.3
13	Parkinson's disease (G20-G21)	2,741	1.0	23.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,866	0.7	15.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,742	0.7	14.7
...	All other causes (residual)	41,935	15.8	355.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 75-79 years				
...	All causes	294,895	100.0	3,524.1
1	Malignant neoplasms (C00-C97)	80,446	27.3	961.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	65,592	22.2	783.9
3	Chronic lower respiratory diseases (J40-J47)	24,390	8.3	291.5
4	Cerebrovascular diseases (I60-I69)	15,966	5.4	190.8
5	Alzheimer's disease (G30)	10,455	3.5	124.9
6	Diabetes mellitus (E10-E14)	9,847	3.3	117.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,111	2.4	85.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,404	2.2	76.5
9	Influenza and pneumonia (J09-J18)	5,543	1.9	66.2
10	Septicemia (A40-A41)	5,025	1.7	60.1
11	Parkinson's disease (G20-G21)	5,009	1.7	59.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,335	1.1	39.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,644	0.9	31.6
14	Pneumonitis due to solids and liquids (J69)	2,157	0.7	25.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,147	0.7	25.7
...	All other causes (residual)	48,824	16.6	583.5
All races and origins, both sexes, 80-84 years				
...	All causes	342,021	100.0	5,830.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	82,027	24.0	1,398.4
2	Malignant neoplasms (C00-C97)	73,515	21.5	1,253.3
3	Chronic lower respiratory diseases (J40-J47)	25,023	7.3	426.6
4	Cerebrovascular diseases (I60-I69)	21,823	6.4	372.0
5	Alzheimer's disease (G30)	20,024	5.9	341.4
6	Diabetes mellitus (E10-E14)	9,786	2.9	166.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,648	2.5	147.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,555	2.2	128.8
9	Parkinson's disease (G20-G21)	7,156	2.1	122.0
10	Influenza and pneumonia (J09-J18)	7,053	2.1	120.2
11	Septicemia (A40-A41)	5,497	1.6	93.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,309	1.3	73.5
13	Pneumonitis due to solids and liquids (J69)	2,983	0.9	50.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,552	0.7	43.5
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,900	0.6	32.4
...	All other causes (residual)	62,170	18.2	1,059.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 85-89 years				
...	All causes	384,479	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	102,758	26.7	---
2	Malignant neoplasms (C00-C97)	60,572	15.8	---
3	Alzheimer's disease (G30)	30,730	8.0	---
4	Cerebrovascular diseases (I60-I69)	27,012	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	23,042	6.0	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10,522	2.7	---
7	Diabetes mellitus (E10-E14)	8,984	2.3	---
8	Influenza and pneumonia (J09-J18)	8,978	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8,139	2.1	---
10	Parkinson's disease (G20-G21)	7,406	1.9	---
11	Septicemia (A40-A41)	5,626	1.5	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,546	1.4	---
13	Pneumonitis due to solids and liquids (J69)	3,897	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,742	0.7	---
15	Aortic aneurysm and dissection (I71)	1,371	0.4	---
...	All other causes (residual)	77,154	20.1	---
All races and origins, both sexes, 90-94 years				
...	All causes	310,771	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	92,612	29.8	---
2	Malignant neoplasms (C00-C97)	32,597	10.5	---
3	Alzheimer's disease (G30)	30,039	9.7	---
4	Cerebrovascular diseases (I60-I69)	23,050	7.4	---
5	Chronic lower respiratory diseases (J40-J47)	14,571	4.7	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,629	2.8	---
7	Influenza and pneumonia (J09-J18)	8,085	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,227	2.0	---
9	Diabetes mellitus (E10-E14)	5,728	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,220	1.7	---
11	Parkinson's disease (G20-G21)	4,159	1.3	---
12	Septicemia (A40-A41)	3,879	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,590	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,788	0.6	---
15	Nutritional deficiencies (E40-E64)	1,264	0.4	---
...	All other causes (residual)	69,333	22.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 95-99 years				
...	All causes	131,376	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	42,428	32.3	---
2	Alzheimer's disease (G30)	13,978	10.6	---
3	Cerebrovascular diseases (I60-I69)	10,030	7.6	---
4	Malignant neoplasms (C00-C97)	9,015	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	4,771	3.6	---
6	Influenza and pneumonia (J09-J18)	3,726	2.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,529	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,636	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,413	1.8	---
10	Diabetes mellitus (E10-E14)	1,814	1.4	---
11	Pneumonitis due to solids and liquids (J69)	1,561	1.2	---
12	Septicemia (A40-A41)	1,362	1.0	---
13	Parkinson's disease (G20-G21)	1,001	0.8	---
14	Nutritional deficiencies (E40-E64)	706	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	632	0.5	---
...	All other causes (residual)	31,774	24.2	---
All races and origins, both sexes, 100 years and over				
...	All causes	27,836	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,340	33.6	---
2	Alzheimer's disease (G30)	2,896	10.4	---
3	Cerebrovascular diseases (I60-I69)	1,981	7.1	---
4	Malignant neoplasms (C00-C97)	1,199	4.3	---
5	Influenza and pneumonia (J09-J18)	925	3.3	---
6	Chronic lower respiratory diseases (J40-J47)	802	2.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	783	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	652	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	453	1.6	---
10	Pneumonitis due to solids and liquids (J69)	304	1.1	---
11	Diabetes mellitus (E10-E14)	291	1.0	---
12	Septicemia (A40-A41)	249	0.9	---
13	Nutritional deficiencies (E40-E64)	210	0.8	---
14	Atherosclerosis (I70)	185	0.7	---
15	Parkinson's disease (G20-G21)	104	0.4	---
...	All other causes (residual)	7,462	26.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, all ages				
...	All causes	1,400,232	100.0	880.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	339,265	24.2	213.3
2	Malignant neoplasms (C00-C97)	314,571	22.5	197.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	103,864	7.4	65.3
4	Chronic lower respiratory diseases (J40-J47)	73,045	5.2	45.9
5	Cerebrovascular diseases (I60-I69)	59,355	4.2	37.3
6	Diabetes mellitus (E10-E14)	43,763	3.1	27.5
7	Alzheimer's disease (G30)	35,372	2.5	22.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	34,727	2.5	21.8
9	Chronic liver disease and cirrhosis (K70,K73-K74)	25,818	1.8	16.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,399	1.8	16.0
11	Influenza and pneumonia (J09-J18)	25,011	1.8	15.7
12	Septicemia (A40-A41)	19,678	1.4	12.4
13	Parkinson's disease (G20-G21)	17,877	1.3	11.2
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15,467	1.1	9.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14,598	1.0	9.2
...	All other causes (residual)	252,422	18.0	158.7
All races and origins, male, 1-4 years				
...	All causes	2,256	100.0	27.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	776	34.4	9.5
2	Malignant neoplasms (C00-C97)	206	9.1	2.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	206	9.1	2.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	176	7.8	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	2.7	0.8
6	Influenza and pneumonia (J09-J18)	53	2.3	0.6
7	Septicemia (A40-A41)	37	1.6	0.5
8	Certain conditions originating in the perinatal period (P00-P96)	35	1.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	34	1.5	0.4
10	Cerebrovascular diseases (I60-I69)	28	1.2	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.0	0.3
12	Acute bronchitis and bronchiolitis (J20-J21)	14	0.6	*
13	Anemias (D50-D64)	7	0.3	*
13	Meningitis (G00,G03)	7	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.3	*
...	All other causes (residual)	586	26.0	7.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 5-9 years				
...	All causes	1,377	100.0	13.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	469	34.1	4.5
2	Malignant neoplasms (C00-C97)	253	18.4	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	7.2	0.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	77	5.6	0.7
5	Chronic lower respiratory diseases (J40-J47)	39	2.8	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	2.7	0.4
7	Influenza and pneumonia (J09-J18)	27	2.0	0.3
8	Cerebrovascular diseases (I60-I69)	24	1.7	0.2
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	1.5	0.2
10	Septicemia (A40-A41)	20	1.5	0.2
11	Anemias (D50-D64)	11	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	7	0.5	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (residual)	276	20.0	2.6
All races and origins, male, 10-14 years				
...	All causes	1,764	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	521	29.5	5.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	265	15.0	2.5
3	Malignant neoplasms (C00-C97)	237	13.4	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	5.1	0.9
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	88	5.0	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	62	3.5	0.6
7	Chronic lower respiratory diseases (J40-J47)	47	2.7	0.4
8	Cerebrovascular diseases (I60-I69)	31	1.8	0.3
9	Influenza and pneumonia (J09-J18)	22	1.2	0.2
10	Septicemia (A40-A41)	17	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	0.9	*
12	Diabetes mellitus (E10-E14)	8	0.5	*
13	Certain conditions originating in the perinatal period (P00-P96)	7	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Anemias (D50-D64)	5	0.3	*
...	All other causes (residual)	341	19.3	3.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 15-19 years				
...	All causes	7,713	100.0	71.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,908	37.7	26.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,601	20.8	14.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,543	20.0	14.3
4	Malignant neoplasms (C00-C97)	368	4.8	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	180	2.3	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	126	1.6	1.2
7	Chronic lower respiratory diseases (J40-J47)	49	0.6	0.5
8	Cerebrovascular diseases (I60-I69)	37	0.5	0.3
9	Diabetes mellitus (E10-E14)	32	0.4	0.3
10	Influenza and pneumonia (J09-J18)	31	0.4	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	0.3	0.2
11	Legal intervention (Y35,Y89.0)	26	0.3	0.2
13	Septicemia (A40-A41)	22	0.3	0.2
14	Anemias (D50-D64)	20	0.3	0.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.1	*
...	All other causes (residual)	736	9.5	6.8
All races and origins, male, 20-24 years				
...	All causes	16,300	100.0	141.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,333	45.0	63.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,974	18.2	25.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,936	18.0	25.6
4	Malignant neoplasms (C00-C97)	534	3.3	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	449	2.8	3.9
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	105	0.6	0.9
7	Diabetes mellitus (E10-E14)	89	0.5	0.8
8	Chronic lower respiratory diseases (J40-J47)	78	0.5	0.7
9	Influenza and pneumonia (J09-J18)	74	0.5	0.6
10	Septicemia (A40-A41)	56	0.3	0.5
11	Cerebrovascular diseases (I60-I69)	55	0.3	0.5
12	Legal intervention (Y35,Y89.0)	49	0.3	0.4
13	Human immunodeficiency virus (HIV) disease (B20-B24)	37	0.2	0.3
14	Anemias (D50-D64)	25	0.2	0.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	21	0.1	0.2
...	All other causes (residual)	1,485	9.1	12.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 25-29 years				
...	All causes	19,511	100.0	167.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,909	45.7	76.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,104	15.9	26.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,557	13.1	22.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	814	4.2	7.0
5	Malignant neoplasms (C00-C97)	737	3.8	6.3
6	Human immunodeficiency virus (HIV) disease (B20-B24)	182	0.9	1.6
7	Diabetes mellitus (E10-E14)	171	0.9	1.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	146	0.7	1.3
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	128	0.7	1.1
10	Cerebrovascular diseases (I60-I69)	118	0.6	1.0
11	Influenza and pneumonia (J09-J18)	94	0.5	0.8
12	Septicemia (A40-A41)	88	0.5	0.8
13	Chronic lower respiratory diseases (J40-J47)	84	0.4	0.7
14	Legal intervention (Y35,Y89.0)	81	0.4	0.7
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	59	0.3	0.5
...	All other causes (residual)	2,239	11.5	19.2
All races and origins, male, 30-34 years				
...	All causes	20,746	100.0	189.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,828	42.6	80.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,783	13.4	25.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,983	9.6	18.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,490	7.2	13.6
5	Malignant neoplasms (C00-C97)	1,088	5.2	9.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	419	2.0	3.8
7	Diabetes mellitus (E10-E14)	285	1.4	2.6
8	Human immunodeficiency virus (HIV) disease (B20-B24)	221	1.1	2.0
9	Cerebrovascular diseases (I60-I69)	213	1.0	1.9
10	Influenza and pneumonia (J09-J18)	149	0.7	1.4
11	Septicemia (A40-A41)	121	0.6	1.1
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	117	0.6	1.1
13	Chronic lower respiratory diseases (J40-J47)	110	0.5	1.0
14	Legal intervention (Y35,Y89.0)	92	0.4	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	89	0.4	0.8
...	All other causes (residual)	2,758	13.3	25.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 35-39 years				
...	All causes	22,809	100.0	219.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,003	35.1	77.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,732	12.0	26.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,609	11.4	25.1
4	Malignant neoplasms (C00-C97)	1,685	7.4	16.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,566	6.9	15.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	669	2.9	6.4
7	Diabetes mellitus (E10-E14)	484	2.1	4.7
8	Cerebrovascular diseases (I60-I69)	398	1.7	3.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	287	1.3	2.8
10	Influenza and pneumonia (J09-J18)	202	0.9	1.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	181	0.8	1.7
12	Septicemia (A40-A41)	170	0.7	1.6
13	Chronic lower respiratory diseases (J40-J47)	140	0.6	1.3
14	Aortic aneurysm and dissection (I71)	105	0.5	1.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	104	0.5	1.0
...	All other causes (residual)	3,474	15.2	33.5
All races and origins, male, 40-44 years				
...	All causes	26,406	100.0	270.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,978	26.4	71.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,538	17.2	46.4
3	Malignant neoplasms (C00-C97)	2,940	11.1	30.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,562	9.7	26.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,142	4.3	11.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,092	4.1	11.2
7	Diabetes mellitus (E10-E14)	754	2.9	7.7
8	Cerebrovascular diseases (I60-I69)	662	2.5	6.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	357	1.4	3.7
10	Septicemia (A40-A41)	291	1.1	3.0
11	Influenza and pneumonia (J09-J18)	290	1.1	3.0
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	232	0.9	2.4
13	Chronic lower respiratory diseases (J40-J47)	224	0.8	2.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	175	0.7	1.8
15	Aortic aneurysm and dissection (I71)	146	0.6	1.5
...	All other causes (residual)	4,023	15.2	41.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 45-49 years				
...	All causes	39,769	100.0	383.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,516	21.4	82.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,510	18.9	72.4
3	Malignant neoplasms (C00-C97)	6,130	15.4	59.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,861	7.2	27.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,969	5.0	19.0
6	Diabetes mellitus (E10-E14)	1,464	3.7	14.1
7	Cerebrovascular diseases (I60-I69)	1,087	2.7	10.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	871	2.2	8.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	551	1.4	5.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	490	1.2	4.7
11	Chronic lower respiratory diseases (J40-J47)	482	1.2	4.6
12	Septicemia (A40-A41)	457	1.1	4.4
13	Influenza and pneumonia (J09-J18)	422	1.1	4.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	374	0.9	3.6
15	Viral hepatitis (B15-B19)	218	0.5	2.1
...	All other causes (residual)	6,367	16.0	61.4
All races and origins, male, 50-54 years				
...	All causes	65,317	100.0	608.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,238	23.3	142.0
2	Malignant neoplasms (C00-C97)	13,803	21.1	128.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,426	12.9	78.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,455	5.3	32.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,337	5.1	31.1
6	Diabetes mellitus (E10-E14)	2,520	3.9	23.5
7	Cerebrovascular diseases (I60-I69)	1,891	2.9	17.6
8	Chronic lower respiratory diseases (J40-J47)	1,449	2.2	13.5
9	Septicemia (A40-A41)	870	1.3	8.1
10	Human immunodeficiency virus (HIV) disease (B20-B24)	793	1.2	7.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	777	1.2	7.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	757	1.2	7.1
13	Influenza and pneumonia (J09-J18)	735	1.1	6.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	644	1.0	6.0
15	Viral hepatitis (B15-B19)	548	0.8	5.1
...	All other causes (residual)	10,074	15.4	93.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 55-59 years				
...	All causes	98,650	100.0	923.4
1	Malignant neoplasms (C00-C97)	26,354	26.7	246.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,836	24.2	223.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,579	8.7	80.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4,641	4.7	43.4
5	Diabetes mellitus (E10-E14)	3,807	3.9	35.6
6	Chronic lower respiratory diseases (J40-J47)	3,362	3.4	31.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,242	3.3	30.3
8	Cerebrovascular diseases (I60-I69)	2,981	3.0	27.9
9	Septicemia (A40-A41)	1,394	1.4	13.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,304	1.3	12.2
11	Influenza and pneumonia (J09-J18)	1,175	1.2	11.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,049	1.1	9.8
13	Viral hepatitis (B15-B19)	963	1.0	9.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	802	0.8	7.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	603	0.6	5.6
...	All other causes (residual)	14,558	14.8	136.3
All races and origins, male, 60-64 years				
...	All causes	123,734	100.0	1,328.2
1	Malignant neoplasms (C00-C97)	37,409	30.2	401.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,422	24.6	326.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,595	5.3	70.8
4	Chronic lower respiratory diseases (J40-J47)	5,702	4.6	61.2
5	Diabetes mellitus (E10-E14)	4,969	4.0	53.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,534	3.7	48.7
7	Cerebrovascular diseases (I60-I69)	4,179	3.4	44.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,503	2.0	26.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,934	1.6	20.8
10	Septicemia (A40-A41)	1,828	1.5	19.6
11	Influenza and pneumonia (J09-J18)	1,665	1.3	17.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,342	1.1	14.4
13	Viral hepatitis (B15-B19)	1,126	0.9	12.1
14	Pneumonitis due to solids and liquids (J69)	576	0.5	6.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	573	0.5	6.2
...	All other causes (residual)	18,377	14.9	197.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 65-69 years				
...	All causes	144,389	100.0	1,819.2
1	Malignant neoplasms (C00-C97)	46,076	31.9	580.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,695	24.7	449.7
3	Chronic lower respiratory diseases (J40-J47)	8,625	6.0	108.7
4	Diabetes mellitus (E10-E14)	5,856	4.1	73.8
5	Cerebrovascular diseases (I60-I69)	5,293	3.7	66.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,068	3.5	63.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,502	2.4	44.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,460	1.7	31.0
9	Septicemia (A40-A41)	2,274	1.6	28.7
10	Influenza and pneumonia (J09-J18)	2,153	1.5	27.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,961	1.4	24.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,461	1.0	18.4
13	Parkinson's disease (G20-G21)	896	0.6	11.3
14	Alzheimer's disease (G30)	892	0.6	11.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	789	0.5	9.9
...	All other causes (residual)	21,388	14.8	269.5
All races and origins, male, 70-74 years				
...	All causes	148,353	100.0	2,720.0
1	Malignant neoplasms (C00-C97)	45,536	30.7	834.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,134	24.4	662.5
3	Chronic lower respiratory diseases (J40-J47)	11,074	7.5	203.0
4	Cerebrovascular diseases (I60-I69)	6,292	4.2	115.4
5	Diabetes mellitus (E10-E14)	5,820	3.9	106.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,065	2.7	74.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,847	1.9	52.2
8	Influenza and pneumonia (J09-J18)	2,404	1.6	44.1
9	Septicemia (A40-A41)	2,282	1.5	41.8
10	Chronic liver disease and cirrhosis (K70,K73-K74)	2,156	1.5	39.5
11	Alzheimer's disease (G30)	1,941	1.3	35.6
12	Parkinson's disease (G20-G21)	1,821	1.2	33.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,560	1.1	28.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,502	1.0	27.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,033	0.7	18.9
...	All other causes (residual)	21,886	14.8	401.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 75-79 years				
...	All causes	156,851	100.0	4,212.3
1	Malignant neoplasms (C00-C97)	43,222	27.6	1,160.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,962	24.2	1,019.5
3	Chronic lower respiratory diseases (J40-J47)	11,997	7.6	322.2
4	Cerebrovascular diseases (I60-I69)	7,730	4.9	207.6
5	Diabetes mellitus (E10-E14)	5,433	3.5	145.9
6	Alzheimer's disease (G30)	4,181	2.7	112.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,136	2.6	111.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,363	2.1	90.3
9	Parkinson's disease (G20-G21)	3,314	2.1	89.0
10	Influenza and pneumonia (J09-J18)	2,975	1.9	79.9
11	Septicemia (A40-A41)	2,491	1.6	66.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,546	1.0	41.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,464	0.9	39.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,262	0.8	33.9
15	Pneumonitis due to solids and liquids (J69)	1,251	0.8	33.6
...	All other causes (residual)	24,524	15.6	658.6
All races and origins, male, 80-84 years				
...	All causes	167,968	100.0	6,846.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	43,079	25.6	1,756.0
2	Malignant neoplasms (C00-C97)	38,598	23.0	1,573.3
3	Chronic lower respiratory diseases (J40-J47)	11,662	6.9	475.4
4	Cerebrovascular diseases (I60-I69)	9,378	5.6	382.3
5	Alzheimer's disease (G30)	7,364	4.4	300.2
6	Diabetes mellitus (E10-E14)	4,958	3.0	202.1
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,526	2.7	184.5
8	Parkinson's disease (G20-G21)	4,481	2.7	182.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,909	2.3	159.3
10	Influenza and pneumonia (J09-J18)	3,523	2.1	143.6
11	Septicemia (A40-A41)	2,593	1.5	105.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,848	1.1	75.3
13	Pneumonitis due to solids and liquids (J69)	1,711	1.0	69.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,397	0.8	56.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	993	0.6	40.5
...	All other causes (residual)	27,948	16.6	1,139.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 85-89 years				
...	All causes	168,063	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	47,640	28.3	---
2	Malignant neoplasms (C00-C97)	30,353	18.1	---
3	Chronic lower respiratory diseases (J40-J47)	10,204	6.1	---
4	Alzheimer's disease (G30)	9,701	5.8	---
5	Cerebrovascular diseases (I60-I69)	9,628	5.7	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,913	2.9	---
7	Parkinson's disease (G20-G21)	4,268	2.5	---
8	Diabetes mellitus (E10-E14)	4,171	2.5	---
9	Influenza and pneumonia (J09-J18)	4,167	2.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,959	2.4	---
11	Septicemia (A40-A41)	2,546	1.5	---
12	Pneumonitis due to solids and liquids (J69)	2,149	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,072	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,393	0.8	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	707	0.4	---
...	All other causes (residual)	30,192	18.0	---
All races and origins, male, 90-94 years				
...	All causes	113,337	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,591	31.4	---
2	Malignant neoplasms (C00-C97)	15,034	13.3	---
3	Alzheimer's disease (G30)	7,887	7.0	---
4	Cerebrovascular diseases (I60-I69)	6,695	5.9	---
5	Chronic lower respiratory diseases (J40-J47)	5,882	5.2	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,380	3.0	---
7	Influenza and pneumonia (J09-J18)	3,290	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,719	2.4	---
9	Diabetes mellitus (E10-E14)	2,273	2.0	---
10	Parkinson's disease (G20-G21)	2,118	1.9	---
11	Pneumonitis due to solids and liquids (J69)	1,824	1.6	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,592	1.4	---
13	Septicemia (A40-A41)	1,519	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	769	0.7	---
15	Nutritional deficiencies (E40-E64)	439	0.4	---
...	All other causes (residual)	22,325	19.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 95-99 years				
...	All causes	36,909	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,850	34.8	---
2	Malignant neoplasms (C00-C97)	3,612	9.8	---
3	Alzheimer's disease (G30)	2,594	7.0	---
4	Cerebrovascular diseases (I60-I69)	2,268	6.1	---
5	Chronic lower respiratory diseases (J40-J47)	1,582	4.3	---
6	Influenza and pneumonia (J09-J18)	1,238	3.4	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,086	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	890	2.4	---
9	Pneumonitis due to solids and liquids (J69)	670	1.8	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	624	1.7	---
11	Diabetes mellitus (E10-E14)	598	1.6	---
12	Parkinson's disease (G20-G21)	442	1.2	---
13	Septicemia (A40-A41)	437	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	199	0.5	---
15	Nutritional deficiencies (E40-E64)	183	0.5	---
...	All other causes (residual)	7,636	20.7	---
All races and origins, male, 100 years and over				
...	All causes	5,046	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,889	37.4	---
2	Malignant neoplasms (C00-C97)	357	7.1	---
3	Alzheimer's disease (G30)	295	5.8	---
4	Cerebrovascular diseases (I60-I69)	285	5.6	---
5	Influenza and pneumonia (J09-J18)	216	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	195	3.9	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	119	2.4	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	117	2.3	---
10	Pneumonitis due to solids and liquids (J69)	103	2.0	---
11	Diabetes mellitus (E10-E14)	62	1.2	---
12	Septicemia (A40-A41)	53	1.1	---
13	Atherosclerosis (I70)	34	0.7	---
14	Parkinson's disease (G20-G21)	33	0.7	---
15	Nutritional deficiencies (E40-E64)	31	0.6	---
...	All other causes (residual)	1,130	22.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, all ages				
...	All causes	1,344,016	100.0	819.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	295,995	22.0	180.4
2	Malignant neoplasms (C00-C97)	283,467	21.1	172.8
3	Cerebrovascular diseases (I60-I69)	82,787	6.2	50.5
4	Chronic lower respiratory diseases (J40-J47)	81,551	6.1	49.7
5	Alzheimer's disease (G30)	80,731	6.0	49.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57,510	4.3	35.1
7	Diabetes mellitus (E10-E14)	36,295	2.7	22.1
8	Influenza and pneumonia (J09-J18)	26,526	2.0	16.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,647	1.8	15.0
10	Septicemia (A40-A41)	20,935	1.6	12.8
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18,648	1.4	11.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14,727	1.1	9.0
13	Parkinson's disease (G20-G21)	11,820	0.9	7.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,238	0.8	6.2
15	Pneumonitis due to solids and liquids (J69)	8,853	0.7	5.4
...	All other causes (residual)	289,286	21.5	176.3
All races and origins, female, 1-4 years				
...	All causes	1,789	100.0	22.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	485	27.1	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	227	12.7	2.9
3	Malignant neoplasms (C00-C97)	171	9.6	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	9.1	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	56	3.1	0.7
6	Influenza and pneumonia (J09-J18)	50	2.8	0.6
7	Septicemia (A40-A41)	33	1.8	0.4
8	Cerebrovascular diseases (I60-I69)	27	1.5	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.5	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	25	1.4	0.3
11	Chronic lower respiratory diseases (J40-J47)	17	1.0	*
12	Anemias (D50-D64)	13	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.6	*
14	Meningitis (G00,G03)	7	0.4	*
15	Diabetes mellitus (E10-E14)	5	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
...	All other causes (residual)	469	26.2	6.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 5-9 years				
...	All causes	1,113	100.0	11.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	318	28.6	3.2
2	Malignant neoplasms (C00-C97)	196	17.6	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	9.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.6	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	3.6	0.4
6	Chronic lower respiratory diseases (J40-J47)	29	2.6	0.3
7	Influenza and pneumonia (J09-J18)	21	1.9	0.2
8	Septicemia (A40-A41)	20	1.8	0.2
9	Cerebrovascular diseases (I60-I69)	14	1.3	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.9	*
11	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
12	Diabetes mellitus (E10-E14)	6	0.5	*
12	Meningitis (G00,G03)	6	0.5	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	6	0.5	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
...	All other causes (residual)	269	24.2	2.7
All races and origins, female, 10-14 years				
...	All causes	1,249	100.0	12.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	326	26.1	3.2
2	Malignant neoplasms (C00-C97)	194	15.5	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	171	13.7	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	58	4.6	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	57	4.6	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	3.9	0.5
7	Chronic lower respiratory diseases (J40-J47)	28	2.2	0.3
8	Cerebrovascular diseases (I60-I69)	19	1.5	*
9	Influenza and pneumonia (J09-J18)	17	1.4	*
10	Septicemia (A40-A41)	14	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	1.0	*
12	Diabetes mellitus (E10-E14)	8	0.6	*
13	Anemias (D50-D64)	5	0.4	*
13	Diseases of appendix (K35-K38)	5	0.4	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (residual)	271	21.7	2.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 15-19 years				
...	All causes	3,099	100.0	30.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,244	40.1	12.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	516	16.7	5.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	273	8.8	2.6
4	Malignant neoplasms (C00-C97)	228	7.4	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	113	3.6	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	2.3	0.7
7	Chronic lower respiratory diseases (J40-J47)	31	1.0	0.3
8	Cerebrovascular diseases (I60-I69)	25	0.8	0.2
9	Pregnancy, childbirth and the puerperium (O00-O99)	24	0.8	0.2
10	Influenza and pneumonia (J09-J18)	23	0.7	0.2
11	Diabetes mellitus (E10-E14)	22	0.7	0.2
12	Septicemia (A40-A41)	20	0.6	0.2
13	Anemias (D50-D64)	17	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (residual)	473	15.3	4.6
All races and origins, female, 20-24 years				
...	All causes	5,463	100.0	50.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,410	44.1	22.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	632	11.6	5.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	420	7.7	3.9
4	Malignant neoplasms (C00-C97)	301	5.5	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	207	3.8	1.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	160	2.9	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	86	1.6	0.8
8	Diabetes mellitus (E10-E14)	68	1.2	0.6
9	Influenza and pneumonia (J09-J18)	61	1.1	0.6
10	Chronic lower respiratory diseases (J40-J47)	48	0.9	0.4
11	Septicemia (A40-A41)	45	0.8	0.4
12	Cerebrovascular diseases (I60-I69)	31	0.6	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	26	0.5	0.2
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.4	0.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	20	0.4	0.2
...	All other causes (residual)	927	17.0	8.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 25-29 years				
...	All causes	7,542	100.0	67.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,032	40.2	26.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	686	9.1	6.1
3	Malignant neoplasms (C00-C97)	645	8.6	5.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	440	5.8	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	412	5.5	3.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	229	3.0	2.0
7	Diabetes mellitus (E10-E14)	129	1.7	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	107	1.4	1.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	90	1.2	0.8
10	Septicemia (A40-A41)	86	1.1	0.8
11	Cerebrovascular diseases (I60-I69)	80	1.1	0.7
12	Influenza and pneumonia (J09-J18)	74	1.0	0.7
13	Chronic lower respiratory diseases (J40-J47)	56	0.7	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	51	0.7	0.5
15	Anemias (D50-D64)	44	0.6	0.4
...	All other causes (residual)	1,381	18.3	12.3
All races and origins, female, 30-34 years				
...	All causes	9,817	100.0	90.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,215	32.7	29.7
2	Malignant neoplasms (C00-C97)	1,321	13.5	12.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	793	8.1	7.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	729	7.4	6.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	396	4.0	3.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	270	2.8	2.5
7	Pregnancy, childbirth and the puerperium (O00-O99)	243	2.5	2.2
8	Diabetes mellitus (E10-E14)	207	2.1	1.9
9	Cerebrovascular diseases (I60-I69)	164	1.7	1.5
10	Influenza and pneumonia (J09-J18)	135	1.4	1.2
11	Septicemia (A40-A41)	124	1.3	1.1
12	Chronic lower respiratory diseases (J40-J47)	103	1.0	1.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	92	0.9	0.9
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	84	0.9	0.8
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	77	0.8	0.7
...	All other causes (residual)	1,864	19.0	17.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 35-39 years				
...	All causes	12,468	100.0	119.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,151	25.3	30.3
2	Malignant neoplasms (C00-C97)	2,386	19.1	22.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,273	10.2	12.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	865	6.9	8.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	438	3.5	4.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	354	2.8	3.4
7	Diabetes mellitus (E10-E14)	318	2.6	3.1
8	Cerebrovascular diseases (I60-I69)	301	2.4	2.9
9	Pregnancy, childbirth and the puerperium (O00-O99)	232	1.9	2.2
10	Septicemia (A40-A41)	165	1.3	1.6
11	Influenza and pneumonia (J09-J18)	152	1.2	1.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	131	1.1	1.3
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	129	1.0	1.2
14	Chronic lower respiratory diseases (J40-J47)	103	0.8	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	102	0.8	1.0
...	All other causes (residual)	2,368	19.0	22.8
All races and origins, female, 40-44 years				
...	All causes	16,109	100.0	162.4
1	Malignant neoplasms (C00-C97)	3,892	24.2	39.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,843	17.6	28.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,057	12.8	20.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	871	5.4	8.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	652	4.0	6.6
6	Diabetes mellitus (E10-E14)	493	3.1	5.0
7	Cerebrovascular diseases (I60-I69)	490	3.0	4.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	307	1.9	3.1
9	Septicemia (A40-A41)	271	1.7	2.7
10	Influenza and pneumonia (J09-J18)	250	1.6	2.5
11	Chronic lower respiratory diseases (J40-J47)	225	1.4	2.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	1.2	2.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	196	1.2	2.0
14	Pregnancy, childbirth and the puerperium (O00-O99)	119	0.7	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	118	0.7	1.2
...	All other causes (residual)	3,124	19.4	31.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 45-49 years				
...	All causes	25,816	100.0	244.2
1	Malignant neoplasms (C00-C97)	7,585	29.4	71.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,660	14.2	34.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,462	13.4	32.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,075	4.2	10.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,007	3.9	9.5
6	Cerebrovascular diseases (I60-I69)	902	3.5	8.5
7	Diabetes mellitus (E10-E14)	814	3.2	7.7
8	Chronic lower respiratory diseases (J40-J47)	601	2.3	5.7
9	Septicemia (A40-A41)	412	1.6	3.9
10	Influenza and pneumonia (J09-J18)	390	1.5	3.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	330	1.3	3.1
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	281	1.1	2.7
13	Human immunodeficiency virus (HIV) disease (B20-B24)	228	0.9	2.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	184	0.7	1.7
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	148	0.6	1.4
...	All other causes (residual)	4,737	18.3	44.8
All races and origins, female, 50-54 years				
...	All causes	42,614	100.0	383.6
1	Malignant neoplasms (C00-C97)	13,773	32.3	124.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,613	15.5	59.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,979	9.3	35.8
4	Chronic liver disease and cirrhosis (K70,K73-K74)	1,865	4.4	16.8
5	Chronic lower respiratory diseases (J40-J47)	1,775	4.2	16.0
6	Cerebrovascular diseases (I60-I69)	1,473	3.5	13.3
7	Diabetes mellitus (E10-E14)	1,469	3.4	13.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,232	2.9	11.1
9	Septicemia (A40-A41)	733	1.7	6.6
10	Influenza and pneumonia (J09-J18)	586	1.4	5.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	543	1.3	4.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	404	0.9	3.6
13	Viral hepatitis (B15-B19)	278	0.7	2.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	259	0.6	2.3
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	243	0.6	2.2
...	All other causes (residual)	7,389	17.3	66.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 55-59 years				
...	All causes	63,496	100.0	562.1
1	Malignant neoplasms (C00-C97)	22,825	35.9	202.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,339	16.3	91.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,807	6.0	33.7
4	Chronic lower respiratory diseases (J40-J47)	3,565	5.6	31.6
5	Diabetes mellitus (E10-E14)	2,263	3.6	20.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,231	3.5	19.7
7	Cerebrovascular diseases (I60-I69)	2,210	3.5	19.6
8	Septicemia (A40-A41)	1,170	1.8	10.4
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,133	1.8	10.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	974	1.5	8.6
11	Influenza and pneumonia (J09-J18)	914	1.4	8.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	568	0.9	5.0
13	Viral hepatitis (B15-B19)	421	0.7	3.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	320	0.5	2.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	270	0.4	2.4
...	All other causes (residual)	10,486	16.5	92.8
All races and origins, female, 60-64 years				
...	All causes	80,565	100.0	792.4
1	Malignant neoplasms (C00-C97)	29,776	37.0	292.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,013	17.4	137.8
3	Chronic lower respiratory diseases (J40-J47)	5,181	6.4	51.0
4	Diabetes mellitus (E10-E14)	3,212	4.0	31.6
5	Cerebrovascular diseases (I60-I69)	2,940	3.6	28.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,879	3.6	28.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,042	2.5	20.1
8	Septicemia (A40-A41)	1,549	1.9	15.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,438	1.8	14.1
10	Influenza and pneumonia (J09-J18)	1,280	1.6	12.6
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	881	1.1	8.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	782	1.0	7.7
13	Viral hepatitis (B15-B19)	486	0.6	4.8
14	Alzheimer's disease (G30)	465	0.6	4.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	424	0.5	4.2
...	All other causes (residual)	13,217	16.4	130.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 65-69 years				
...	All causes	102,266	100.0	1,151.2
1	Malignant neoplasms (C00-C97)	36,669	35.9	412.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	18,460	18.1	207.8
3	Chronic lower respiratory diseases (J40-J47)	7,883	7.7	88.7
4	Cerebrovascular diseases (I60-I69)	4,344	4.2	48.9
5	Diabetes mellitus (E10-E14)	4,036	3.9	45.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,511	2.5	28.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,065	2.0	23.2
8	Septicemia (A40-A41)	1,972	1.9	22.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1,702	1.7	19.2
10	Influenza and pneumonia (J09-J18)	1,638	1.6	18.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,123	1.1	12.6
12	Alzheimer's disease (G30)	1,120	1.1	12.6
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	591	0.6	6.7
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	576	0.6	6.5
15	Pneumonitis due to solids and liquids (J69)	498	0.5	5.6
...	All other causes (residual)	17,078	16.7	192.3
All races and origins, female, 70-74 years				
...	All causes	117,072	100.0	1,841.9
1	Malignant neoplasms (C00-C97)	37,302	31.9	586.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,072	18.9	347.3
3	Chronic lower respiratory diseases (J40-J47)	10,821	9.2	170.2
4	Cerebrovascular diseases (I60-I69)	5,839	5.0	91.9
5	Diabetes mellitus (E10-E14)	4,290	3.7	67.5
6	Alzheimer's disease (G30)	2,808	2.4	44.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,532	2.2	39.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,406	2.1	37.9
9	Septicemia (A40-A41)	2,239	1.9	35.2
10	Influenza and pneumonia (J09-J18)	1,974	1.7	31.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,437	1.2	22.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,310	1.1	20.6
13	Parkinson's disease (G20-G21)	920	0.8	14.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	709	0.6	11.2
15	Pneumonitis due to solids and liquids (J69)	609	0.5	9.6
...	All other causes (residual)	19,804	16.9	311.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 75-79 years				
...	All causes	138,044	100.0	2,972.3
1	Malignant neoplasms (C00-C97)	37,224	27.0	801.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	27,630	20.0	594.9
3	Chronic lower respiratory diseases (J40-J47)	12,393	9.0	266.8
4	Cerebrovascular diseases (I60-I69)	8,236	6.0	177.3
5	Alzheimer's disease (G30)	6,274	4.5	135.1
6	Diabetes mellitus (E10-E14)	4,414	3.2	95.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,041	2.2	65.5
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,975	2.2	64.1
9	Influenza and pneumonia (J09-J18)	2,568	1.9	55.3
10	Septicemia (A40-A41)	2,534	1.8	54.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,789	1.3	38.5
12	Parkinson's disease (G20-G21)	1,695	1.2	36.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,180	0.9	25.4
14	Pneumonitis due to solids and liquids (J69)	906	0.7	19.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	885	0.6	19.1
...	All other causes (residual)	24,300	17.6	523.2
All races and origins, female, 80-84 years				
...	All causes	174,053	100.0	5,100.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	38,948	22.4	1,141.4
2	Malignant neoplasms (C00-C97)	34,917	20.1	1,023.2
3	Chronic lower respiratory diseases (J40-J47)	13,361	7.7	391.5
4	Alzheimer's disease (G30)	12,660	7.3	371.0
5	Cerebrovascular diseases (I60-I69)	12,445	7.2	364.7
6	Diabetes mellitus (E10-E14)	4,828	2.8	141.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,122	2.4	120.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,646	2.1	106.8
9	Influenza and pneumonia (J09-J18)	3,530	2.0	103.4
10	Septicemia (A40-A41)	2,904	1.7	85.1
11	Parkinson's disease (G20-G21)	2,675	1.5	78.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,461	1.4	72.1
13	Pneumonitis due to solids and liquids (J69)	1,272	0.7	37.3
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,155	0.7	33.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	907	0.5	26.6
...	All other causes (residual)	34,222	19.7	1,002.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 85-89 years				
...	All causes	216,416	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	55,118	25.5	---
2	Malignant neoplasms (C00-C97)	30,219	14.0	---
3	Alzheimer's disease (G30)	21,029	9.7	---
4	Cerebrovascular diseases (I60-I69)	17,384	8.0	---
5	Chronic lower respiratory diseases (J40-J47)	12,838	5.9	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,609	2.6	---
7	Diabetes mellitus (E10-E14)	4,813	2.2	---
8	Influenza and pneumonia (J09-J18)	4,811	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,180	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,474	1.6	---
11	Parkinson's disease (G20-G21)	3,138	1.4	---
12	Septicemia (A40-A41)	3,080	1.4	---
13	Pneumonitis due to solids and liquids (J69)	1,748	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,349	0.6	---
15	Nutritional deficiencies (E40-E64)	785	0.4	---
...	All other causes (residual)	46,841	21.6	---
All races and origins, female, 90-94 years				
...	All causes	197,434	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	57,021	28.9	---
2	Alzheimer's disease (G30)	22,152	11.2	---
3	Malignant neoplasms (C00-C97)	17,563	8.9	---
4	Cerebrovascular diseases (I60-I69)	16,355	8.3	---
5	Chronic lower respiratory diseases (J40-J47)	8,689	4.4	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,249	2.7	---
7	Influenza and pneumonia (J09-J18)	4,795	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,628	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,508	1.8	---
10	Diabetes mellitus (E10-E14)	3,455	1.7	---
11	Septicemia (A40-A41)	2,360	1.2	---
12	Parkinson's disease (G20-G21)	2,041	1.0	---
13	Pneumonitis due to solids and liquids (J69)	1,766	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,019	0.5	---
15	Nutritional deficiencies (E40-E64)	825	0.4	---
...	All other causes (residual)	47,008	23.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 95-99 years				
...	All causes	94,467	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	29,578	31.3	---
2	Alzheimer's disease (G30)	11,384	12.1	---
3	Cerebrovascular diseases (I60-I69)	7,762	8.2	---
4	Malignant neoplasms (C00-C97)	5,403	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	3,189	3.4	---
6	Influenza and pneumonia (J09-J18)	2,488	2.6	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,443	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,012	2.1	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,523	1.6	---
10	Diabetes mellitus (E10-E14)	1,216	1.3	---
11	Septicemia (A40-A41)	925	1.0	---
12	Pneumonitis due to solids and liquids (J69)	891	0.9	---
13	Parkinson's disease (G20-G21)	559	0.6	---
14	Nutritional deficiencies (E40-E64)	523	0.6	---
15	Atherosclerosis (I70)	462	0.5	---
...	All other causes (residual)	24,109	25.5	---
All races and origins, female, 100 years and over				
...	All causes	22,790	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,451	32.7	---
2	Alzheimer's disease (G30)	2,601	11.4	---
3	Cerebrovascular diseases (I60-I69)	1,696	7.4	---
4	Malignant neoplasms (C00-C97)	842	3.7	---
5	Influenza and pneumonia (J09-J18)	709	3.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	666	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	607	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	525	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	334	1.5	---
10	Diabetes mellitus (E10-E14)	229	1.0	---
11	Pneumonitis due to solids and liquids (J69)	201	0.9	---
12	Septicemia (A40-A41)	196	0.9	---
13	Nutritional deficiencies (E40-E64)	179	0.8	---
14	Atherosclerosis (I70)	151	0.7	---
15	Anemias (D50-D64)	74	0.3	---
...	All other causes (residual)	6,329	27.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, all ages				
...	All causes	188,254	100.0	327.6
1	Malignant neoplasms (C00-C97)	39,263	20.9	68.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,799	20.1	65.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	15,711	8.3	27.3
4	Cerebrovascular diseases (I60-I69)	10,283	5.5	17.9
5	Diabetes mellitus (E10-E14)	8,546	4.5	14.9
6	Alzheimer's disease (G30)	6,833	3.6	11.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,141	3.3	10.7
8	Chronic lower respiratory diseases (J40-J47)	5,287	2.8	9.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,775	2.0	6.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,668	1.9	6.4
11	Influenza and pneumonia (J09-J18)	3,641	1.9	6.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,187	1.7	5.5
13	Septicemia (A40-A41)	2,718	1.4	4.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,488	1.3	4.3
15	Certain conditions originating in the perinatal period (P00-P96)	2,415	1.3	4.2
...	All other causes (residual)	36,499	19.4	63.5
Hispanic, both sexes, 1-4 years				
...	All causes	799	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	233	29.2	5.7
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	99	12.4	2.4
3	Malignant neoplasms (C00-C97)	89	11.1	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	7.8	1.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	3.4	0.7
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	16	2.0	*
6	Influenza and pneumonia (J09-J18)	16	2.0	*
8	Septicemia (A40-A41)	14	1.8	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.5	*
10	Cerebrovascular diseases (I60-I69)	8	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	8	1.0	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
13	Anemias (D50-D64)	3	0.4	*
14	Meningitis (G00,G03)	2	0.3	*
14	Hernia (K40-K46)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
...	All other causes (residual)	200	25.0	4.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 5-9 years				
...	All causes	578	100.0	11.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	29.2	3.2
2	Malignant neoplasms (C00-C97)	123	21.3	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	8.8	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	4.0	0.4
5	Cerebrovascular diseases (I60-I69)	17	2.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	2.4	*
7	Chronic lower respiratory diseases (J40-J47)	12	2.1	*
8	Septicemia (A40-A41)	9	1.6	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.4	*
10	Influenza and pneumonia (J09-J18)	7	1.2	*
11	Anemias (D50-D64)	3	0.5	*
12	Diseases of appendix (K35-K38)	2	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
14	Meningococcal infection (A39)	1	0.2	*
14	Diabetes mellitus (E10-E14)	1	0.2	*
14	Nutritional deficiencies (E40-E64)	1	0.2	*
14	Meningitis (G00,G03)	1	0.2	*
14	Pneumonitis due to solids and liquids (J69)	1	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.2	*
...	All other causes (residual)	130	22.5	2.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 10-14 years				
...	All causes	586	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	161	27.5	3.2
2	Malignant neoplasms (C00-C97)	103	17.6	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	10.1	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	36	6.1	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	5.5	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.1	0.5
7	Cerebrovascular diseases (I60-I69)	12	2.0	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.7	*
9	Septicemia (A40-A41)	9	1.5	*
10	Influenza and pneumonia (J09-J18)	7	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
11	Anemias (D50-D64)	4	0.7	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
15	Viral hepatitis (B15-B19)	1	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Peptic ulcer (K25-K28)	1	0.2	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.2	*
15	Legal intervention (Y35,Y89.0)	1	0.2	*
...	All other causes (residual)	111	18.9	2.2
Hispanic, both sexes, 15-19 years				
...	All causes	2,017	100.0	42.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	807	40.0	16.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	363	18.0	7.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	327	16.2	6.8
4	Malignant neoplasms (C00-C97)	154	7.6	3.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	2.2	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	1.8	0.8
7	Cerebrovascular diseases (I60-I69)	19	0.9	*
8	Chronic lower respiratory diseases (J40-J47)	12	0.6	*
9	Septicemia (A40-A41)	8	0.4	*
9	Diabetes mellitus (E10-E14)	8	0.4	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	8	0.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.3	*
12	Influenza and pneumonia (J09-J18)	6	0.3	*
12	Legal intervention (Y35,Y89.0)	6	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.1	*
...	All other causes (residual)	208	10.3	4.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 20-24 years				
...	All causes	3,705	100.0	77.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,720	46.4	35.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	624	16.8	13.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	514	13.9	10.7
4	Malignant neoplasms (C00-C97)	216	5.8	4.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	2.2	1.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	1.1	0.8
7	Influenza and pneumonia (J09-J18)	28	0.8	0.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	21	0.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	18	0.5	*
10	Diabetes mellitus (E10-E14)	17	0.5	*
11	Cerebrovascular diseases (I60-I69)	15	0.4	*
11	Legal intervention (Y35,Y89.0)	15	0.4	*
13	Septicemia (A40-A41)	14	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.3	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.2	*
...	All other causes (residual)	362	9.8	7.5
Hispanic, both sexes, 25-29 years				
...	All causes	3,988	100.0	86.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,659	41.6	36.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	526	13.2	11.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	522	13.1	11.3
4	Malignant neoplasms (C00-C97)	283	7.1	6.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	170	4.3	3.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	49	1.2	1.1
7	Cerebrovascular diseases (I60-I69)	46	1.2	1.0
8	Human immunodeficiency virus (HIV) disease (B20-B24)	41	1.0	0.9
9	Pregnancy, childbirth and the puerperium (O00-O99)	39	1.0	0.8
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	1.0	0.8
11	Diabetes mellitus (E10-E14)	34	0.9	0.7
12	Influenza and pneumonia (J09-J18)	26	0.7	0.6
13	Septicemia (A40-A41)	25	0.6	0.5
14	Legal intervention (Y35,Y89.0)	17	0.4	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.4	*
...	All other causes (residual)	498	12.5	10.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 30-34 years				
...	All causes	4,207	100.0	94.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,553	36.9	34.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	438	10.4	9.8
3	Malignant neoplasms (C00-C97)	423	10.1	9.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	379	9.0	8.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	267	6.3	6.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	145	3.4	3.3
7	Diabetes mellitus (E10-E14)	65	1.5	1.5
8	Cerebrovascular diseases (I60-I69)	62	1.5	1.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	51	1.2	1.1
10	Influenza and pneumonia (J09-J18)	48	1.1	1.1
11	Septicemia (A40-A41)	44	1.0	1.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	41	1.0	0.9
13	Chronic lower respiratory diseases (J40-J47)	27	0.6	0.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	0.6	0.5
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.5	0.5
...	All other causes (residual)	619	14.7	13.9
Hispanic, both sexes, 35-39 years				
...	All causes	4,731	100.0	110.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,383	29.2	32.2
2	Malignant neoplasms (C00-C97)	704	14.9	16.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	395	8.3	9.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	391	8.3	9.1
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	352	7.4	8.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	244	5.2	5.7
7	Cerebrovascular diseases (I60-I69)	124	2.6	2.9
8	Diabetes mellitus (E10-E14)	91	1.9	2.1
9	Human immunodeficiency virus (HIV) disease (B20-B24)	75	1.6	1.7
10	Influenza and pneumonia (J09-J18)	47	1.0	1.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	44	0.9	1.0
12	Septicemia (A40-A41)	32	0.7	0.7
12	Pregnancy, childbirth and the puerperium (O00-O99)	32	0.7	0.7
14	Chronic lower respiratory diseases (J40-J47)	27	0.6	0.6
15	Aortic aneurysm and dissection (I71)	24	0.5	0.6
...	All other causes (residual)	766	16.2	17.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 40-44 years				
...	All causes	5,541	100.0	140.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,209	21.8	30.6
2	Malignant neoplasms (C00-C97)	1,089	19.7	27.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	676	12.2	17.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	378	6.8	9.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	308	5.6	7.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	224	4.0	5.7
7	Cerebrovascular diseases (I60-I69)	202	3.6	5.1
8	Diabetes mellitus (E10-E14)	177	3.2	4.5
9	Influenza and pneumonia (J09-J18)	88	1.6	2.2
10	Septicemia (A40-A41)	70	1.3	1.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	66	1.2	1.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	1.0	1.4
13	Chronic lower respiratory diseases (J40-J47)	41	0.7	1.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	38	0.7	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	0.6	0.8
...	All other causes (residual)	888	16.0	22.4
Hispanic, both sexes, 45-49 years				
...	All causes	7,347	100.0	205.7
1	Malignant neoplasms (C00-C97)	1,633	22.2	45.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,196	16.3	33.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,073	14.6	30.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	578	7.9	16.2
5	Cerebrovascular diseases (I60-I69)	313	4.3	8.8
6	Diabetes mellitus (E10-E14)	299	4.1	8.4
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	273	3.7	7.6
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	159	2.2	4.5
9	Human immunodeficiency virus (HIV) disease (B20-B24)	117	1.6	3.3
10	Septicemia (A40-A41)	109	1.5	3.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	1.4	2.9
12	Influenza and pneumonia (J09-J18)	101	1.4	2.8
13	Viral hepatitis (B15-B19)	68	0.9	1.9
14	Chronic lower respiratory diseases (J40-J47)	60	0.8	1.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	0.8	1.6
...	All other causes (residual)	1,207	16.4	33.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 50-54 years				
...	All causes	10,406	100.0	334.3
1	Malignant neoplasms (C00-C97)	2,648	25.4	85.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,807	17.4	58.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,175	11.3	37.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	850	8.2	27.3
5	Diabetes mellitus (E10-E14)	516	5.0	16.6
6	Cerebrovascular diseases (I60-I69)	413	4.0	13.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	277	2.7	8.9
8	Septicemia (A40-A41)	175	1.7	5.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	168	1.6	5.4
10	Viral hepatitis (B15-B19)	150	1.4	4.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	141	1.4	4.5
11	Influenza and pneumonia (J09-J18)	141	1.4	4.5
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	128	1.2	4.1
14	Chronic lower respiratory diseases (J40-J47)	127	1.2	4.1
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	111	1.1	3.6
...	All other causes (residual)	1,579	15.2	50.7
Hispanic, both sexes, 55-59 years				
...	All causes	12,974	100.0	511.3
1	Malignant neoplasms (C00-C97)	3,688	28.4	145.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,452	18.9	96.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,004	7.7	39.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	959	7.4	37.8
5	Diabetes mellitus (E10-E14)	798	6.2	31.4
6	Cerebrovascular diseases (I60-I69)	537	4.1	21.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	255	2.0	10.0
8	Chronic lower respiratory diseases (J40-J47)	215	1.7	8.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	207	1.6	8.2
10	Viral hepatitis (B15-B19)	204	1.6	8.0
11	Septicemia (A40-A41)	202	1.6	8.0
11	Influenza and pneumonia (J09-J18)	202	1.6	8.0
13	Human immunodeficiency virus (HIV) disease (B20-B24)	137	1.1	5.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	0.9	4.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	79	0.6	3.1
...	All other causes (residual)	1,914	14.8	75.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 60-64 years				
...	All causes	14,613	100.0	756.3
1	Malignant neoplasms (C00-C97)	4,530	31.0	234.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,890	19.8	149.6
3	Diabetes mellitus (E10-E14)	985	6.7	51.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	889	6.1	46.0
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	702	4.8	36.3
6	Cerebrovascular diseases (I60-I69)	663	4.5	34.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	349	2.4	18.1
8	Chronic lower respiratory diseases (J40-J47)	317	2.2	16.4
9	Influenza and pneumonia (J09-J18)	237	1.6	12.3
10	Septicemia (A40-A41)	218	1.5	11.3
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	196	1.3	10.1
12	Viral hepatitis (B15-B19)	184	1.3	9.5
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	130	0.9	6.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	78	0.5	4.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.5	3.4
...	All other causes (residual)	2,179	14.9	112.8
Hispanic, both sexes, 65-69 years				
...	All causes	15,582	100.0	1,086.9
1	Malignant neoplasms (C00-C97)	4,851	31.1	338.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,318	21.3	231.4
3	Diabetes mellitus (E10-E14)	970	6.2	67.7
4	Cerebrovascular diseases (I60-I69)	795	5.1	55.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	653	4.2	45.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	482	3.1	33.6
7	Chronic lower respiratory diseases (J40-J47)	427	2.7	29.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	407	2.6	28.4
9	Influenza and pneumonia (J09-J18)	319	2.0	22.3
10	Septicemia (A40-A41)	279	1.8	19.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	192	1.2	13.4
12	Viral hepatitis (B15-B19)	138	0.9	9.6
13	Alzheimer's disease (G30)	119	0.8	8.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	93	0.6	6.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	84	0.5	5.9
...	All other causes (residual)	2,455	15.8	171.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 70-74 years				
...	All causes	15,968	100.0	1,650.3
1	Malignant neoplasms (C00-C97)	4,559	28.6	471.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,513	22.0	363.1
3	Diabetes mellitus (E10-E14)	1,099	6.9	113.6
4	Cerebrovascular diseases (I60-I69)	910	5.7	94.1
5	Chronic lower respiratory diseases (J40-J47)	562	3.5	58.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	465	2.9	48.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	462	2.9	47.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	408	2.6	42.2
9	Influenza and pneumonia (J09-J18)	324	2.0	33.5
10	Alzheimer's disease (G30)	297	1.9	30.7
11	Septicemia (A40-A41)	290	1.8	30.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	230	1.4	23.8
13	Parkinson's disease (G20-G21)	140	0.9	14.5
14	Viral hepatitis (B15-B19)	86	0.5	8.9
14	Pneumonitis due to solids and liquids (J69)	86	0.5	8.9
...	All other causes (residual)	2,537	15.9	262.2
Hispanic, both sexes, 75-79 years				
...	All causes	17,961	100.0	2,681.5
1	Malignant neoplasms (C00-C97)	4,704	26.2	702.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,092	22.8	610.9
3	Cerebrovascular diseases (I60-I69)	1,172	6.5	175.0
4	Diabetes mellitus (E10-E14)	1,073	6.0	160.2
5	Chronic lower respiratory diseases (J40-J47)	708	3.9	105.7
6	Alzheimer's disease (G30)	641	3.6	95.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	483	2.7	72.1
8	Chronic liver disease and cirrhosis (K70,K73-K74)	414	2.3	61.8
9	Influenza and pneumonia (J09-J18)	413	2.3	61.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	372	2.1	55.5
11	Septicemia (A40-A41)	311	1.7	46.4
12	Parkinson's disease (G20-G21)	269	1.5	40.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	265	1.5	39.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	134	0.7	20.0
15	Pneumonitis due to solids and liquids (J69)	110	0.6	16.4
...	All other causes (residual)	2,800	15.6	418.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 80-84 years				
...	All causes	20,145	100.0	4,458.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,954	24.6	1,096.3
2	Malignant neoplasms (C00-C97)	4,228	21.0	935.7
3	Cerebrovascular diseases (I60-I69)	1,479	7.3	327.3
4	Alzheimer's disease (G30)	1,190	5.9	263.4
5	Diabetes mellitus (E10-E14)	958	4.8	212.0
6	Chronic lower respiratory diseases (J40-J47)	902	4.5	199.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	501	2.5	110.9
8	Influenza and pneumonia (J09-J18)	486	2.4	107.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	438	2.2	96.9
10	Parkinson's disease (G20-G21)	409	2.0	90.5
11	Septicemia (A40-A41)	327	1.6	72.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	305	1.5	67.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	295	1.5	65.3
14	Pneumonitis due to solids and liquids (J69)	167	0.8	37.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	120	0.6	26.6
...	All other causes (residual)	3,386	16.8	749.3
Hispanic, both sexes, 85-89 years				
...	All causes	20,899	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,525	26.4	---
2	Malignant neoplasms (C00-C97)	3,220	15.4	---
3	Alzheimer's disease (G30)	1,862	8.9	---
4	Cerebrovascular diseases (I60-I69)	1,669	8.0	---
5	Chronic lower respiratory diseases (J40-J47)	957	4.6	---
6	Diabetes mellitus (E10-E14)	850	4.1	---
7	Influenza and pneumonia (J09-J18)	520	2.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	473	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	467	2.2	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	427	2.0	---
11	Parkinson's disease (G20-G21)	399	1.9	---
12	Septicemia (A40-A41)	310	1.5	---
13	Pneumonitis due to solids and liquids (J69)	179	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	144	0.7	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	140	0.7	---
...	All other causes (residual)	3,757	18.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 90-94 years				
...	All causes	14,522	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,230	29.1	---
2	Alzheimer's disease (G30)	1,716	11.8	---
3	Malignant neoplasms (C00-C97)	1,546	10.6	---
4	Cerebrovascular diseases (I60-I69)	1,192	8.2	---
5	Chronic lower respiratory diseases (J40-J47)	593	4.1	---
6	Diabetes mellitus (E10-E14)	452	3.1	---
7	Influenza and pneumonia (J09-J18)	400	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	338	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	307	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	284	2.0	---
11	Parkinson's disease (G20-G21)	226	1.6	---
12	Septicemia (A40-A41)	164	1.1	---
13	Pneumonitis due to solids and liquids (J69)	123	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.5	---
15	Nutritional deficiencies (E40-E64)	65	0.4	---
...	All other causes (residual)	2,809	19.3	---
Hispanic, both sexes, 95-99 years				
...	All causes	5,562	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,742	31.3	---
2	Alzheimer's disease (G30)	741	13.3	---
3	Cerebrovascular diseases (I60-I69)	504	9.1	---
4	Malignant neoplasms (C00-C97)	403	7.2	---
5	Chronic lower respiratory diseases (J40-J47)	202	3.6	---
6	Influenza and pneumonia (J09-J18)	152	2.7	---
7	Diabetes mellitus (E10-E14)	136	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	127	2.3	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	1.6	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	1.5	---
11	Septicemia (A40-A41)	57	1.0	---
12	Pneumonitis due to solids and liquids (J69)	55	1.0	---
13	Parkinson's disease (G20-G21)	47	0.8	---
14	Nutritional deficiencies (E40-E64)	45	0.8	---
15	Atherosclerosis (I70)	22	0.4	---
...	All other causes (residual)	1,156	20.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 100 years and over				
...	All causes	1,293	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	448	34.6	---
2	Alzheimer's disease (G30)	174	13.5	---
3	Cerebrovascular diseases (I60-I69)	106	8.2	---
4	Chronic lower respiratory diseases (J40-J47)	51	3.9	---
5	Malignant neoplasms (C00-C97)	49	3.8	---
6	Influenza and pneumonia (J09-J18)	39	3.0	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	36	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	1.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	1.4	---
10	Diabetes mellitus (E10-E14)	16	1.2	---
11	Atherosclerosis (I70)	10	0.8	---
12	Pneumonitis due to solids and liquids (J69)	9	0.7	---
13	Nutritional deficiencies (E40-E64)	8	0.6	---
14	Septicemia (A40-A41)	6	0.5	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.4	---
...	All other causes (residual)	299	23.1	---
Hispanic, male, all ages				
...	All causes	103,532	100.0	356.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20,945	20.2	72.2
2	Malignant neoplasms (C00-C97)	20,504	19.8	70.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,510	11.1	39.7
4	Cerebrovascular diseases (I60-I69)	4,798	4.6	16.5
5	Diabetes mellitus (E10-E14)	4,603	4.4	15.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,225	4.1	14.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,944	2.8	10.1
8	Chronic lower respiratory diseases (J40-J47)	2,759	2.7	9.5
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,657	2.6	9.2
10	Alzheimer's disease (G30)	2,151	2.1	7.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,931	1.9	6.7
12	Influenza and pneumonia (J09-J18)	1,877	1.8	6.5
13	Septicemia (A40-A41)	1,392	1.3	4.8
14	Certain conditions originating in the perinatal period (P00-P96)	1,348	1.3	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,197	1.2	4.1
...	All other causes (residual)	18,691	18.1	64.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 1-4 years				
...	All causes	443	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	146	33.0	7.0
2	Malignant neoplasms (C00-C97)	48	10.8	2.3
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	10.2	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	7.4	1.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.2	*
6	Influenza and pneumonia (J09-J18)	10	2.3	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	1.8	*
8	Cerebrovascular diseases (I60-I69)	6	1.4	*
9	Septicemia (A40-A41)	5	1.1	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	5	1.1	*
12	Acute bronchitis and bronchiolitis (J20-J21)	3	0.7	*
13	Diabetes mellitus (E10-E14)	1	0.2	*
13	Meningitis (G00,G03)	1	0.2	*
13	Diseases of appendix (K35-K38)	1	0.2	*
13	Hernia (K40-K46)	1	0.2	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	108	24.4	5.1
Hispanic, male, 5-9 years				
...	All causes	313	100.0	11.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	30.4	3.5
2	Malignant neoplasms (C00-C97)	65	20.8	2.4
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	6.7	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	3.5	*
5	Cerebrovascular diseases (I60-I69)	10	3.2	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	2.9	*
7	Septicemia (A40-A41)	7	2.2	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	2.2	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.9	*
10	Anemias (D50-D64)	3	1.0	*
10	Influenza and pneumonia (J09-J18)	3	1.0	*
12	Meningococcal infection (A39)	1	0.3	*
12	Diabetes mellitus (E10-E14)	1	0.3	*
12	Diseases of appendix (K35-K38)	1	0.3	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1	0.3	*
...	All other causes (residual)	69	22.0	2.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 10-14 years				
...	All causes	328	100.0	12.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	94	28.7	3.7
2	Malignant neoplasms (C00-C97)	60	18.3	2.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	9.1	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	20	6.1	0.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	6.1	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	4.0	*
7	Cerebrovascular diseases (I60-I69)	8	2.4	*
8	Septicemia (A40-A41)	5	1.5	*
8	Influenza and pneumonia (J09-J18)	5	1.5	*
8	Chronic lower respiratory diseases (J40-J47)	5	1.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.2	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.6	*
14	Viral hepatitis (B15-B19)	1	0.3	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
14	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
14	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (residual)	56	17.1	2.2
Hispanic, male, 15-19 years				
...	All causes	1,447	100.0	59.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	568	39.3	23.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	310	21.4	12.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	236	16.3	9.6
4	Malignant neoplasms (C00-C97)	103	7.1	4.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	28	1.9	1.1
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	1.8	1.1
7	Cerebrovascular diseases (I60-I69)	12	0.8	*
8	Chronic lower respiratory diseases (J40-J47)	7	0.5	*
9	Diabetes mellitus (E10-E14)	6	0.4	*
9	Legal intervention (Y35,Y89.0)	6	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.3	*
12	Influenza and pneumonia (J09-J18)	4	0.3	*
13	Septicemia (A40-A41)	3	0.2	*
14	Anemias (D50-D64)	2	0.1	*
15	Tuberculosis (A16-A19)	1	0.1	*
15	Viral hepatitis (B15-B19)	1	0.1	*
15	Nutritional deficiencies (E40-E64)	1	0.1	*
15	Meningitis (G00,G03)	1	0.1	*
15	Aortic aneurysm and dissection (I71)	1	0.1	*
15	Pneumonitis due to solids and liquids (J69)	1	0.1	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.1	*
15	Certain conditions originating in the perinatal period (P00-P96)	1	0.1	*
...	All other causes (residual)	123	8.5	5.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 20-24 years				
...	All causes	2,857	100.0	115.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,329	46.5	53.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	559	19.6	22.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	429	15.0	17.3
4	Malignant neoplasms (C00-C97)	146	5.1	5.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	61	2.1	2.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	25	0.9	1.0
7	Influenza and pneumonia (J09-J18)	18	0.6	*
8	Legal intervention (Y35,Y89.0)	15	0.5	*
9	Diabetes mellitus (E10-E14)	11	0.4	*
10	Cerebrovascular diseases (I60-I69)	10	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	10	0.4	*
12	Septicemia (A40-A41)	9	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.1	*
14	Pneumonitis due to solids and liquids (J69)	4	0.1	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.1	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.1	*
...	All other causes (residual)	214	7.5	8.6
Hispanic, male, 25-29 years				
...	All causes	3,053	100.0	126.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,326	43.4	54.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	459	15.0	19.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	440	14.4	18.2
4	Malignant neoplasms (C00-C97)	166	5.4	6.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	123	4.0	5.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	35	1.1	1.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	35	1.1	1.4
8	Cerebrovascular diseases (I60-I69)	30	1.0	1.2
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	22	0.7	0.9
10	Influenza and pneumonia (J09-J18)	20	0.7	0.8
11	Legal intervention (Y35,Y89.0)	17	0.6	*
12	Septicemia (A40-A41)	16	0.5	*
12	Diabetes mellitus (E10-E14)	16	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.2	*
...	All other causes (residual)	328	10.7	13.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 30-34 years				
...	All causes	3,006	100.0	129.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,232	41.0	52.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	370	12.3	15.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	300	10.0	12.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	190	6.3	8.2
5	Malignant neoplasms (C00-C97)	178	5.9	7.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	106	3.5	4.5
7	Diabetes mellitus (E10-E14)	42	1.4	1.8
8	Human immunodeficiency virus (HIV) disease (B20-B24)	36	1.2	1.5
9	Cerebrovascular diseases (I60-I69)	35	1.2	1.5
10	Influenza and pneumonia (J09-J18)	30	1.0	1.3
11	Septicemia (A40-A41)	22	0.7	0.9
12	Legal intervention (Y35,Y89.0)	19	0.6	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	0.6	*
14	Chronic lower respiratory diseases (J40-J47)	14	0.5	*
15	Aortic aneurysm and dissection (I71)	10	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	0.3	*
...	All other causes (residual)	394	13.1	16.9
Hispanic, male, 35-39 years				
...	All causes	3,265	100.0	147.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,107	33.9	50.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	320	9.8	14.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	299	9.2	13.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	294	9.0	13.3
5	Malignant neoplasms (C00-C97)	268	8.2	12.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	173	5.3	7.8
7	Cerebrovascular diseases (I60-I69)	83	2.5	3.7
8	Human immunodeficiency virus (HIV) disease (B20-B24)	60	1.8	2.7
9	Diabetes mellitus (E10-E14)	57	1.7	2.6
10	Influenza and pneumonia (J09-J18)	31	0.9	1.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	0.6	0.9
12	Aortic aneurysm and dissection (I71)	19	0.6	*
13	Chronic lower respiratory diseases (J40-J47)	18	0.6	*
13	Legal intervention (Y35,Y89.0)	18	0.6	*
15	Septicemia (A40-A41)	15	0.5	*
...	All other causes (residual)	482	14.8	21.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 40-44 years				
...	All causes	3,682	100.0	183.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	964	26.2	48.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	501	13.6	25.0
3	Malignant neoplasms (C00-C97)	448	12.2	22.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	282	7.7	14.1
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	250	6.8	12.5
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	184	5.0	9.2
7	Cerebrovascular diseases (I60-I69)	126	3.4	6.3
8	Diabetes mellitus (E10-E14)	118	3.2	5.9
9	Influenza and pneumonia (J09-J18)	61	1.7	3.0
10	Human immunodeficiency virus (HIV) disease (B20-B24)	50	1.4	2.5
11	Septicemia (A40-A41)	44	1.2	2.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.8	1.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	0.7	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.5	*
15	Chronic lower respiratory diseases (J40-J47)	18	0.5	*
...	All other causes (residual)	563	15.3	28.1
Hispanic, male, 45-49 years				
...	All causes	4,775	100.0	263.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	937	19.6	51.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	819	17.2	45.2
3	Malignant neoplasms (C00-C97)	708	14.8	39.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	440	9.2	24.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	224	4.7	12.4
6	Cerebrovascular diseases (I60-I69)	189	4.0	10.4
7	Diabetes mellitus (E10-E14)	180	3.8	9.9
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	123	2.6	6.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	95	2.0	5.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	1.4	3.8
11	Influenza and pneumonia (J09-J18)	62	1.3	3.4
12	Septicemia (A40-A41)	60	1.3	3.3
13	Viral hepatitis (B15-B19)	53	1.1	2.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	0.7	1.9
15	Chronic lower respiratory diseases (J40-J47)	31	0.6	1.7
...	All other causes (residual)	751	15.7	41.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 50-54 years				
...	All causes	6,776	100.0	433.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,343	19.8	86.0
2	Malignant neoplasms (C00-C97)	1,341	19.8	85.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	896	13.2	57.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	645	9.5	41.3
5	Diabetes mellitus (E10-E14)	323	4.8	20.7
6	Cerebrovascular diseases (I60-I69)	258	3.8	16.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	222	3.3	14.2
8	Septicemia (A40-A41)	122	1.8	7.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	118	1.7	7.6
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	103	1.5	6.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	102	1.5	6.5
12	Viral hepatitis (B15-B19)	101	1.5	6.5
13	Influenza and pneumonia (J09-J18)	88	1.3	5.6
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	1.0	4.3
15	Chronic lower respiratory diseases (J40-J47)	60	0.9	3.8
...	All other causes (residual)	987	14.6	63.2
Hispanic, male, 55-59 years				
...	All causes	8,336	100.0	669.1
1	Malignant neoplasms (C00-C97)	1,960	23.5	157.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,798	21.6	144.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	777	9.3	62.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	721	8.6	57.9
5	Diabetes mellitus (E10-E14)	489	5.9	39.2
6	Cerebrovascular diseases (I60-I69)	337	4.0	27.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	158	1.9	12.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	153	1.8	12.3
9	Viral hepatitis (B15-B19)	148	1.8	11.9
10	Influenza and pneumonia (J09-J18)	131	1.6	10.5
11	Septicemia (A40-A41)	125	1.5	10.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	114	1.4	9.2
13	Chronic lower respiratory diseases (J40-J47)	110	1.3	8.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	0.9	6.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	0.8	5.5
...	All other causes (residual)	1,171	14.0	94.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 60-64 years				
...	All causes	9,080	100.0	987.0
1	Malignant neoplasms (C00-C97)	2,515	27.7	273.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,022	22.3	219.8
3	Chronic liver disease and cirrhosis (K70,K73-K74)	642	7.1	69.8
4	Diabetes mellitus (E10-E14)	596	6.6	64.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	502	5.5	54.6
6	Cerebrovascular diseases (I60-I69)	404	4.4	43.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	204	2.2	22.2
8	Chronic lower respiratory diseases (J40-J47)	186	2.0	20.2
9	Influenza and pneumonia (J09-J18)	150	1.7	16.3
10	Septicemia (A40-A41)	136	1.5	14.8
11	Viral hepatitis (B15-B19)	134	1.5	14.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	121	1.3	13.2
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	98	1.1	10.7
14	Human immunodeficiency virus (HIV) disease (B20-B24)	57	0.6	6.2
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	0.5	4.8
...	All other causes (residual)	1,269	14.0	137.9
Hispanic, male, 65-69 years				
...	All causes	9,244	100.0	1,394.6
1	Malignant neoplasms (C00-C97)	2,719	29.4	410.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,186	23.6	329.8
3	Diabetes mellitus (E10-E14)	572	6.2	86.3
4	Cerebrovascular diseases (I60-I69)	466	5.0	70.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	442	4.8	66.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	350	3.8	52.8
7	Chronic lower respiratory diseases (J40-J47)	253	2.7	38.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	208	2.3	31.4
9	Influenza and pneumonia (J09-J18)	175	1.9	26.4
10	Septicemia (A40-A41)	154	1.7	23.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	115	1.2	17.3
12	Viral hepatitis (B15-B19)	92	1.0	13.9
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	78	0.8	11.8
14	Alzheimer's disease (G30)	52	0.6	7.8
15	Parkinson's disease (G20-G21)	50	0.5	7.5
...	All other causes (residual)	1,332	14.4	200.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 70-74 years				
...	All causes	8,967	100.0	2,093.3
1	Malignant neoplasms (C00-C97)	2,505	27.9	584.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,150	24.0	501.9
3	Diabetes mellitus (E10-E14)	603	6.7	140.8
4	Cerebrovascular diseases (I60-I69)	480	5.4	112.1
5	Chronic lower respiratory diseases (J40-J47)	322	3.6	75.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	281	3.1	65.6
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	274	3.1	64.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	242	2.7	56.5
9	Influenza and pneumonia (J09-J18)	185	2.1	43.2
10	Septicemia (A40-A41)	140	1.6	32.7
10	Alzheimer's disease (G30)	140	1.6	32.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	139	1.6	32.4
13	Parkinson's disease (G20-G21)	79	0.9	18.4
14	Pneumonitis due to solids and liquids (J69)	56	0.6	13.1
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.6	11.7
...	All other causes (residual)	1,321	14.7	308.4
Hispanic, male, 75-79 years				
...	All causes	9,329	100.0	3,288.8
1	Malignant neoplasms (C00-C97)	2,502	26.8	882.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,269	24.3	799.9
3	Cerebrovascular diseases (I60-I69)	573	6.1	202.0
4	Diabetes mellitus (E10-E14)	546	5.9	192.5
5	Chronic lower respiratory diseases (J40-J47)	406	4.4	143.1
6	Alzheimer's disease (G30)	257	2.8	90.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	237	2.5	83.6
8	Chronic liver disease and cirrhosis (K70,K73-K74)	217	2.3	76.5
9	Influenza and pneumonia (J09-J18)	214	2.3	75.4
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	199	2.1	70.2
11	Parkinson's disease (G20-G21)	161	1.7	56.8
12	Septicemia (A40-A41)	140	1.5	49.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	124	1.3	43.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	77	0.8	27.1
15	Pneumonitis due to solids and liquids (J69)	62	0.7	21.9
...	All other causes (residual)	1,345	14.4	474.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 80-84 years				
...	All causes	9,519	100.0	5,251.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,365	24.8	1,304.7
2	Malignant neoplasms (C00-C97)	2,223	23.4	1,226.4
3	Cerebrovascular diseases (I60-I69)	622	6.5	343.1
4	Chronic lower respiratory diseases (J40-J47)	475	5.0	262.0
5	Diabetes mellitus (E10-E14)	442	4.6	243.8
6	Alzheimer's disease (G30)	429	4.5	236.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	247	2.6	136.3
8	Parkinson's disease (G20-G21)	240	2.5	132.4
9	Influenza and pneumonia (J09-J18)	226	2.4	124.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	2.3	119.7
11	Septicemia (A40-A41)	141	1.5	77.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	136	1.4	75.0
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	133	1.4	73.4
14	Pneumonitis due to solids and liquids (J69)	95	1.0	52.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	51	0.5	28.1
...	All other causes (residual)	1,477	15.5	814.8
Hispanic, male, 85-89 years				
...	All causes	8,963	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,427	27.1	---
2	Malignant neoplasms (C00-C97)	1,597	17.8	---
3	Cerebrovascular diseases (I60-I69)	617	6.9	---
4	Alzheimer's disease (G30)	596	6.6	---
5	Chronic lower respiratory diseases (J40-J47)	460	5.1	---
6	Diabetes mellitus (E10-E14)	386	4.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	238	2.7	---
8	Parkinson's disease (G20-G21)	224	2.5	---
8	Influenza and pneumonia (J09-J18)	224	2.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	213	2.4	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	159	1.8	---
12	Septicemia (A40-A41)	129	1.4	---
13	Pneumonitis due to solids and liquids (J69)	83	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	68	0.8	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	68	0.8	---
...	All other causes (residual)	1,474	16.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 90-94 years				
...	All causes	5,541	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,663	30.0	---
2	Malignant neoplasms (C00-C97)	742	13.4	---
3	Alzheimer's disease (G30)	471	8.5	---
4	Cerebrovascular diseases (I60-I69)	394	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	274	4.9	---
6	Diabetes mellitus (E10-E14)	169	3.0	---
7	Influenza and pneumonia (J09-J18)	158	2.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	129	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	2.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	122	2.2	---
11	Parkinson's disease (G20-G21)	106	1.9	---
12	Septicemia (A40-A41)	65	1.2	---
13	Pneumonitis due to solids and liquids (J69)	63	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	25	0.5	---
...	All other causes (residual)	1,002	18.1	---
Hispanic, male, 95-99 years				
...	All causes	1,643	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	534	32.5	---
2	Malignant neoplasms (C00-C97)	181	11.0	---
3	Alzheimer's disease (G30)	143	8.7	---
4	Cerebrovascular diseases (I60-I69)	104	6.3	---
5	Chronic lower respiratory diseases (J40-J47)	74	4.5	---
6	Influenza and pneumonia (J09-J18)	49	3.0	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	2.7	---
8	Diabetes mellitus (E10-E14)	42	2.6	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	2.2	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	31	1.9	---
11	Septicemia (A40-A41)	26	1.6	---
12	Pneumonitis due to solids and liquids (J69)	22	1.3	---
13	Parkinson's disease (G20-G21)	16	1.0	---
14	Nutritional deficiencies (E40-E64)	10	0.6	---
15	Atherosclerosis (I70)	6	0.4	---
...	All other causes (residual)	325	19.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 100 years and over				
...	All causes	278	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	103	37.1	---
2	Alzheimer's disease (G30)	22	7.9	---
3	Malignant neoplasms (C00-C97)	20	7.2	---
3	Cerebrovascular diseases (I60-I69)	20	7.2	---
5	Chronic lower respiratory diseases (J40-J47)	18	6.5	---
6	Influenza and pneumonia (J09-J18)	11	4.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	2.2	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	2.2	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	1.4	---
10	Diabetes mellitus (E10-E14)	3	1.1	---
10	Nutritional deficiencies (E40-E64)	3	1.1	---
10	Pneumonitis due to solids and liquids (J69)	3	1.1	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.7	---
13	Parkinson's disease (G20-G21)	2	0.7	---
13	Atherosclerosis (I70)	2	0.7	---
...	All other causes (residual)	53	19.1	---
Hispanic, female, all ages				
...	All causes	84,722	100.0	297.7
1	Malignant neoplasms (C00-C97)	18,759	22.1	65.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,854	19.9	59.2
3	Cerebrovascular diseases (I60-I69)	5,485	6.5	19.3
4	Alzheimer's disease (G30)	4,682	5.5	16.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,201	5.0	14.8
6	Diabetes mellitus (E10-E14)	3,943	4.7	13.9
7	Chronic lower respiratory diseases (J40-J47)	2,528	3.0	8.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,916	2.3	6.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,844	2.2	6.5
10	Influenza and pneumonia (J09-J18)	1,764	2.1	6.2
11	Septicemia (A40-A41)	1,326	1.6	4.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,291	1.5	4.5
13	Certain conditions originating in the perinatal period (P00-P96)	1,067	1.3	3.7
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	858	1.0	3.0
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	724	0.9	2.5
...	All other causes (residual)	17,480	20.6	61.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 1-4 years				
...	All causes	356	100.0	17.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	87	24.4	4.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	15.2	2.7
3	Malignant neoplasms (C00-C97)	41	11.5	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	8.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	3.7	*
6	Septicemia (A40-A41)	9	2.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.2	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	2.0	*
9	Influenza and pneumonia (J09-J18)	6	1.7	*
10	Anemias (D50-D64)	3	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	3	0.8	*
12	Cerebrovascular diseases (I60-I69)	2	0.6	*
13	Salmonella infections (A01-A02)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
13	Hernia (K40-K46)	1	0.3	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	87	24.4	4.3
Hispanic, female, 5-9 years				
...	All causes	265	100.0	10.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	74	27.9	2.9
2	Malignant neoplasms (C00-C97)	58	21.9	2.2
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	30	11.3	1.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	4.5	*
5	Cerebrovascular diseases (I60-I69)	7	2.6	*
6	Chronic lower respiratory diseases (J40-J47)	6	2.3	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.9	*
8	Influenza and pneumonia (J09-J18)	4	1.5	*
9	Septicemia (A40-A41)	2	0.8	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
10	Nutritional deficiencies (E40-E64)	1	0.4	*
10	Meningitis (G00,G03)	1	0.4	*
10	Pneumonitis due to solids and liquids (J69)	1	0.4	*
10	Diseases of appendix (K35-K38)	1	0.4	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
...	All other causes (residual)	61	23.0	2.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 10-14 years				
...	All causes	258	100.0	10.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	67	26.0	2.7
2	Malignant neoplasms (C00-C97)	43	16.7	1.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	11.2	1.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	6.2	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	4.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	4.3	*
7	Chronic lower respiratory diseases (J40-J47)	5	1.9	*
8	Septicemia (A40-A41)	4	1.6	*
8	Anemias (D50-D64)	4	1.6	*
8	Cerebrovascular diseases (I60-I69)	4	1.6	*
11	Influenza and pneumonia (J09-J18)	2	0.8	*
12	Peptic ulcer (K25-K28)	1	0.4	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	55	21.3	2.2
Hispanic, female, 15-19 years				
...	All causes	570	100.0	24.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	239	41.9	10.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	16.0	3.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	9.3	2.3
4	Malignant neoplasms (C00-C97)	51	8.9	2.2
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	3.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	11	1.9	*
7	Pregnancy, childbirth and the puerperium (O00-O99)	8	1.4	*
8	Cerebrovascular diseases (I60-I69)	7	1.2	*
9	Septicemia (A40-A41)	5	0.9	*
9	Chronic lower respiratory diseases (J40-J47)	5	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.4	*
11	Diabetes mellitus (E10-E14)	2	0.4	*
11	Influenza and pneumonia (J09-J18)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	73	12.8	3.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 20-24 years				
...	All causes	848	100.0	36.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	391	46.1	16.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	10.0	3.7
3	Malignant neoplasms (C00-C97)	70	8.3	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	65	7.7	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	2.6	0.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	21	2.5	0.9
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	15	1.8	*
8	Influenza and pneumonia (J09-J18)	10	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
10	Diabetes mellitus (E10-E14)	6	0.7	*
11	Septicemia (A40-A41)	5	0.6	*
11	Cerebrovascular diseases (I60-I69)	5	0.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.4	*
15	Aortic aneurysm and dissection (I71)	3	0.4	*
...	All other causes (residual)	127	15.0	5.5
Hispanic, female, 25-29 years				
...	All causes	935	100.0	42.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	333	35.6	15.2
2	Malignant neoplasms (C00-C97)	117	12.5	5.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	82	8.8	3.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	7.2	3.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	5.0	2.1
6	Pregnancy, childbirth and the puerperium (O00-O99)	39	4.2	1.8
7	Diabetes mellitus (E10-E14)	18	1.9	*
8	Cerebrovascular diseases (I60-I69)	16	1.7	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	1.7	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.5	*
11	Septicemia (A40-A41)	9	1.0	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.6	*
12	Influenza and pneumonia (J09-J18)	6	0.6	*
14	Anemias (D50-D64)	4	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	3	0.3	*
15	Inflammatory diseases of female pelvic organs (N70-N76)	3	0.3	*
...	All other causes (residual)	155	16.6	7.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 30-34 years				
...	All causes	1,201	100.0	56.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	26.7	15.1
2	Malignant neoplasms (C00-C97)	245	20.4	11.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	79	6.6	3.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	77	6.4	3.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	5.7	3.2
6	Pregnancy, childbirth and the puerperium (O00-O99)	51	4.2	2.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	39	3.2	1.8
8	Cerebrovascular diseases (I60-I69)	27	2.2	1.3
9	Diabetes mellitus (E10-E14)	23	1.9	1.1
10	Septicemia (A40-A41)	22	1.8	1.0
11	Influenza and pneumonia (J09-J18)	18	1.5	*
12	Chronic lower respiratory diseases (J40-J47)	13	1.1	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.4	*
...	All other causes (residual)	191	15.9	9.0
Hispanic, female, 35-39 years				
...	All causes	1,466	100.0	70.6
1	Malignant neoplasms (C00-C97)	436	29.7	21.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	276	18.8	13.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	97	6.6	4.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	75	5.1	3.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	71	4.8	3.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	3.6	2.6
7	Cerebrovascular diseases (I60-I69)	41	2.8	2.0
8	Diabetes mellitus (E10-E14)	34	2.3	1.6
9	Pregnancy, childbirth and the puerperium (O00-O99)	32	2.2	1.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	1.6	1.1
11	Septicemia (A40-A41)	17	1.2	*
12	Influenza and pneumonia (J09-J18)	16	1.1	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	15	1.0	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	12	0.8	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.8	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.8	*
...	All other causes (residual)	246	16.8	11.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 40-44 years				
...	All causes	1,859	100.0	95.3
1	Malignant neoplasms (C00-C97)	641	34.5	32.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	245	13.2	12.6
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	175	9.4	9.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	96	5.2	4.9
5	Cerebrovascular diseases (I60-I69)	76	4.1	3.9
6	Diabetes mellitus (E10-E14)	59	3.2	3.0
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	3.1	3.0
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	2.2	2.1
9	Influenza and pneumonia (J09-J18)	27	1.5	1.4
10	Septicemia (A40-A41)	26	1.4	1.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	1.4	1.3
12	Chronic lower respiratory diseases (J40-J47)	23	1.2	1.2
13	Pregnancy, childbirth and the puerperium (O00-O99)	21	1.1	1.1
14	Human immunodeficiency virus (HIV) disease (B20-B24)	16	0.9	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	0.8	*
...	All other causes (residual)	316	17.0	16.2
Hispanic, female, 45-49 years				
...	All causes	2,572	100.0	145.9
1	Malignant neoplasms (C00-C97)	925	36.0	52.5
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	259	10.1	14.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	254	9.9	14.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	138	5.4	7.8
5	Cerebrovascular diseases (I60-I69)	124	4.8	7.0
6	Diabetes mellitus (E10-E14)	119	4.6	6.8
7	Septicemia (A40-A41)	49	1.9	2.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	1.9	2.8
9	Influenza and pneumonia (J09-J18)	39	1.5	2.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	36	1.4	2.0
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	1.4	2.0
12	Chronic lower respiratory diseases (J40-J47)	29	1.1	1.6
13	Human immunodeficiency virus (HIV) disease (B20-B24)	22	0.9	1.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.9	1.2
15	Viral hepatitis (B15-B19)	15	0.6	*
...	All other causes (residual)	456	17.7	25.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 50-54 years				
...	All causes	3,630	100.0	234.1
1	Malignant neoplasms (C00-C97)	1,307	36.0	84.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	464	12.8	29.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	279	7.7	18.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	205	5.6	13.2
5	Diabetes mellitus (E10-E14)	193	5.3	12.4
6	Cerebrovascular diseases (I60-I69)	155	4.3	10.0
7	Chronic lower respiratory diseases (J40-J47)	67	1.8	4.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	1.8	4.3
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	1.5	3.5
10	Septicemia (A40-A41)	53	1.5	3.4
10	Influenza and pneumonia (J09-J18)	53	1.5	3.4
12	Viral hepatitis (B15-B19)	49	1.3	3.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.2	2.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	0.7	1.6
15	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.6	1.5
...	All other causes (residual)	592	16.3	38.2
Hispanic, female, 55-59 years				
...	All causes	4,638	100.0	359.1
1	Malignant neoplasms (C00-C97)	1,728	37.3	133.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	654	14.1	50.6
3	Diabetes mellitus (E10-E14)	309	6.7	23.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	238	5.1	18.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	227	4.9	17.6
6	Cerebrovascular diseases (I60-I69)	200	4.3	15.5
7	Chronic lower respiratory diseases (J40-J47)	105	2.3	8.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	102	2.2	7.9
9	Septicemia (A40-A41)	77	1.7	6.0
10	Influenza and pneumonia (J09-J18)	71	1.5	5.5
11	Viral hepatitis (B15-B19)	56	1.2	4.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	49	1.1	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	1.0	3.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	23	0.5	1.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	0.5	1.7
...	All other causes (residual)	732	15.8	56.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 60-64 years				
...	All causes	5,533	100.0	546.7
1	Malignant neoplasms (C00-C97)	2,015	36.4	199.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	868	15.7	85.8
3	Diabetes mellitus (E10-E14)	389	7.0	38.4
4	Cerebrovascular diseases (I60-I69)	259	4.7	25.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	247	4.5	24.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	200	3.6	19.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	2.6	14.3
8	Chronic lower respiratory diseases (J40-J47)	131	2.4	12.9
9	Influenza and pneumonia (J09-J18)	87	1.6	8.6
10	Septicemia (A40-A41)	82	1.5	8.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	1.4	7.4
12	Viral hepatitis (B15-B19)	50	0.9	4.9
13	Alzheimer's disease (G30)	36	0.7	3.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	32	0.6	3.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.5	2.9
...	All other causes (residual)	888	16.0	87.7
Hispanic, female, 65-69 years				
...	All causes	6,338	100.0	822.2
1	Malignant neoplasms (C00-C97)	2,132	33.6	276.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,132	17.9	146.9
3	Diabetes mellitus (E10-E14)	398	6.3	51.6
4	Cerebrovascular diseases (I60-I69)	329	5.2	42.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	211	3.3	27.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	3.1	25.8
7	Chronic lower respiratory diseases (J40-J47)	174	2.7	22.6
8	Influenza and pneumonia (J09-J18)	144	2.3	18.7
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	2.1	17.1
10	Septicemia (A40-A41)	125	2.0	16.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	77	1.2	10.0
12	Alzheimer's disease (G30)	67	1.1	8.7
13	Viral hepatitis (B15-B19)	46	0.7	6.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	37	0.6	4.8
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	33	0.5	4.3
...	All other causes (residual)	1,102	17.4	143.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 70-74 years				
...	All causes	7,001	100.0	1,298.4
1	Malignant neoplasms (C00-C97)	2,054	29.3	380.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,363	19.5	252.8
3	Diabetes mellitus (E10-E14)	496	7.1	92.0
4	Cerebrovascular diseases (I60-I69)	430	6.1	79.7
5	Chronic lower respiratory diseases (J40-J47)	240	3.4	44.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	223	3.2	41.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	181	2.6	33.6
8	Alzheimer's disease (G30)	157	2.2	29.1
9	Septicemia (A40-A41)	150	2.1	27.8
10	Influenza and pneumonia (J09-J18)	139	2.0	25.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	134	1.9	24.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	91	1.3	16.9
13	Parkinson's disease (G20-G21)	61	0.9	11.3
14	Viral hepatitis (B15-B19)	37	0.5	6.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.4	5.6
15	Pneumonitis due to solids and liquids (J69)	30	0.4	5.6
...	All other causes (residual)	1,185	16.9	219.8
Hispanic, female, 75-79 years				
...	All causes	8,632	100.0	2,235.4
1	Malignant neoplasms (C00-C97)	2,202	25.5	570.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,823	21.1	472.1
3	Cerebrovascular diseases (I60-I69)	599	6.9	155.1
4	Diabetes mellitus (E10-E14)	527	6.1	136.5
5	Alzheimer's disease (G30)	384	4.4	99.4
6	Chronic lower respiratory diseases (J40-J47)	302	3.5	78.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	246	2.8	63.7
8	Influenza and pneumonia (J09-J18)	199	2.3	51.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	197	2.3	51.0
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	173	2.0	44.8
11	Septicemia (A40-A41)	171	2.0	44.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	141	1.6	36.5
13	Parkinson's disease (G20-G21)	108	1.3	28.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.7	14.8
15	Pneumonitis due to solids and liquids (J69)	48	0.6	12.4
...	All other causes (residual)	1,455	16.9	376.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 80-84 years				
...	All causes	10,626	100.0	3,926.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,589	24.4	956.8
2	Malignant neoplasms (C00-C97)	2,005	18.9	741.0
3	Cerebrovascular diseases (I60-I69)	857	8.1	316.7
4	Alzheimer's disease (G30)	761	7.2	281.2
5	Diabetes mellitus (E10-E14)	516	4.9	190.7
6	Chronic lower respiratory diseases (J40-J47)	427	4.0	157.8
7	Influenza and pneumonia (J09-J18)	260	2.4	96.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	254	2.4	93.9
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	221	2.1	81.7
10	Septicemia (A40-A41)	186	1.8	68.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	172	1.6	63.6
12	Parkinson's disease (G20-G21)	169	1.6	62.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	159	1.5	58.8
14	Pneumonitis due to solids and liquids (J69)	72	0.7	26.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	69	0.6	25.5
...	All other causes (residual)	1,909	18.0	705.5
Hispanic, female, 85-89 years				
...	All causes	11,936	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,098	26.0	---
2	Malignant neoplasms (C00-C97)	1,623	13.6	---
3	Alzheimer's disease (G30)	1,266	10.6	---
4	Cerebrovascular diseases (I60-I69)	1,052	8.8	---
5	Chronic lower respiratory diseases (J40-J47)	497	4.2	---
6	Diabetes mellitus (E10-E14)	464	3.9	---
7	Influenza and pneumonia (J09-J18)	296	2.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	268	2.2	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	254	2.1	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	235	2.0	---
11	Septicemia (A40-A41)	181	1.5	---
12	Parkinson's disease (G20-G21)	175	1.5	---
13	Pneumonitis due to solids and liquids (J69)	96	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	76	0.6	---
15	Chronic liver disease and cirrhosis (K70,K73-K74)	72	0.6	---
...	All other causes (residual)	2,283	19.1	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 90-94 years				
...	All causes	8,981	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,567	28.6	---
2	Alzheimer's disease (G30)	1,245	13.9	---
3	Malignant neoplasms (C00-C97)	804	9.0	---
4	Cerebrovascular diseases (I60-I69)	798	8.9	---
5	Chronic lower respiratory diseases (J40-J47)	319	3.6	---
6	Diabetes mellitus (E10-E14)	283	3.2	---
7	Influenza and pneumonia (J09-J18)	242	2.7	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	209	2.3	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	180	2.0	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	162	1.8	---
11	Parkinson's disease (G20-G21)	120	1.3	---
12	Septicemia (A40-A41)	99	1.1	---
13	Pneumonitis due to solids and liquids (J69)	60	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.5	---
15	Nutritional deficiencies (E40-E64)	44	0.5	---
...	All other causes (residual)	1,803	20.1	---
Hispanic, female, 95-99 years				
...	All causes	3,919	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,208	30.8	---
2	Alzheimer's disease (G30)	598	15.3	---
3	Cerebrovascular diseases (I60-I69)	400	10.2	---
4	Malignant neoplasms (C00-C97)	222	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	128	3.3	---
6	Influenza and pneumonia (J09-J18)	103	2.6	---
7	Diabetes mellitus (E10-E14)	94	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	83	2.1	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60	1.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.2	---
11	Nutritional deficiencies (E40-E64)	35	0.9	---
12	Pneumonitis due to solids and liquids (J69)	33	0.8	---
13	Septicemia (A40-A41)	31	0.8	---
13	Parkinson's disease (G20-G21)	31	0.8	---
15	Atherosclerosis (I70)	16	0.4	---
...	All other causes (residual)	831	21.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 100 years and over				
...	All causes	1,015	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	345	34.0	---
2	Alzheimer's disease (G30)	152	15.0	---
3	Cerebrovascular diseases (I60-I69)	86	8.5	---
4	Chronic lower respiratory diseases (J40-J47)	33	3.3	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	3.2	---
6	Malignant neoplasms (C00-C97)	29	2.9	---
7	Influenza and pneumonia (J09-J18)	28	2.8	---
8	Diabetes mellitus (E10-E14)	13	1.3	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.3	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	1.2	---
11	Atherosclerosis (I70)	8	0.8	---
12	Pneumonitis due to solids and liquids (J69)	6	0.6	---
13	Septicemia (A40-A41)	5	0.5	---
13	Nutritional deficiencies (E40-E64)	5	0.5	---
15	Anemias (D50-D64)	4	0.4	---
...	All other causes (residual)	244	24.0	---
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,133,463	100.0	1,059.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	500,631	23.5	248.7
2	Malignant neoplasms (C00-C97)	466,467	21.9	231.7
3	Chronic lower respiratory diseases (J40-J47)	135,268	6.3	67.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121,641	5.7	60.4
5	Cerebrovascular diseases (I60-I69)	107,491	5.0	53.4
6	Alzheimer's disease (G30)	97,779	4.6	48.6
7	Diabetes mellitus (E10-E14)	53,399	2.5	26.5
8	Influenza and pneumonia (J09-J18)	39,748	1.9	19.7
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,531	1.7	18.1
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,248	1.7	17.5
11	Septicemia (A40-A41)	30,009	1.4	14.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29,432	1.4	14.6
13	Parkinson's disease (G20-G21)	25,826	1.2	12.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23,237	1.1	11.5
15	Pneumonitis due to solids and liquids (J69)	16,302	0.8	8.1
...	All other causes (residual)	414,454	19.4	205.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,978	100.0	23.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	680	34.4	8.2
2	Malignant neoplasms (C00-C97)	206	10.4	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	191	9.7	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	121	6.1	1.5
5	Influenza and pneumonia (J09-J18)	47	2.4	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	45	2.3	0.5
7	Septicemia (A40-A41)	34	1.7	0.4
8	Cerebrovascular diseases (I60-I69)	27	1.4	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	23	1.2	0.3
10	Chronic lower respiratory diseases (J40-J47)	22	1.1	0.3
11	Certain conditions originating in the perinatal period (P00-P96)	20	1.0	0.2
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	6	0.3	*
15	Meningitis (G00,G03)	5	0.3	*
15	Pneumonitis due to solids and liquids (J69)	5	0.3	*
...	All other causes (residual)	532	26.9	6.4
Non-Hispanic, white, both sexes, 5-9 years				
...	All causes	1,158	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	360	31.1	3.4
2	Malignant neoplasms (C00-C97)	221	19.1	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	108	9.3	1.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	53	4.6	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	39	3.4	0.4
6	Influenza and pneumonia (J09-J18)	26	2.2	0.2
7	Septicemia (A40-A41)	20	1.7	0.2
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.5	*
9	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
10	Cerebrovascular diseases (I60-I69)	11	0.9	*
11	Diabetes mellitus (E10-E14)	8	0.7	*
12	Certain conditions originating in the perinatal period (P00-P96)	7	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
15	Anemias (D50-D64)	4	0.3	*
...	All other causes (residual)	260	22.5	2.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 10-14 years				
...	All causes	1,623	100.0	14.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	484	29.8	4.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	309	19.0	2.8
3	Malignant neoplasms (C00-C97)	219	13.5	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	4.9	0.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	2.7	0.4
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	2.6	0.4
7	Cerebrovascular diseases (I60-I69)	20	1.2	0.2
7	Chronic lower respiratory diseases (J40-J47)	20	1.2	0.2
9	Influenza and pneumonia (J09-J18)	18	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	1.0	*
11	Septicemia (A40-A41)	14	0.9	*
12	Diabetes mellitus (E10-E14)	7	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.3	*
15	Anemias (D50-D64)	3	0.2	*
15	Aortic aneurysm and dissection (I71)	3	0.2	*
15	Pneumonitis due to solids and liquids (J69)	3	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
...	All other causes (residual)	326	20.1	2.9
Non-Hispanic, white, both sexes, 15-19 years				
...	All causes	5,668	100.0	48.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,545	44.9	21.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,422	25.1	12.1
3	Malignant neoplasms (C00-C97)	318	5.6	2.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	284	5.0	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	149	2.6	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	104	1.8	0.9
7	Diabetes mellitus (E10-E14)	33	0.6	0.3
7	Influenza and pneumonia (J09-J18)	33	0.6	0.3
9	Cerebrovascular diseases (I60-I69)	26	0.5	0.2
10	Chronic lower respiratory diseases (J40-J47)	23	0.4	0.2
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.4	0.2
12	Septicemia (A40-A41)	19	0.3	*
13	Pneumonitis due to solids and liquids (J69)	10	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	8	0.1	*
...	All other causes (residual)	664	11.7	5.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 20-24 years				
...	All causes	11,847	100.0	95.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,350	53.6	51.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,334	19.7	18.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	482	4.1	3.9
4	Malignant neoplasms (C00-C97)	426	3.6	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	332	2.8	2.7
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	106	0.9	0.9
7	Diabetes mellitus (E10-E14)	80	0.7	0.6
8	Pregnancy, childbirth and the puerperium (O00-O99)	71	0.6	0.6
9	Influenza and pneumonia (J09-J18)	67	0.6	0.5
10	Chronic lower respiratory diseases (J40-J47)	64	0.5	0.5
11	Septicemia (A40-A41)	55	0.5	0.4
12	Cerebrovascular diseases (I60-I69)	45	0.4	0.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	24	0.2	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.1	*
15	Pneumonitis due to solids and liquids (J69)	16	0.1	*
15	Legal intervention (Y35,Y89.0)	16	0.1	*
...	All other causes (residual)	1,362	11.5	10.9
Non-Hispanic, white, both sexes, 25-29 years				
...	All causes	16,143	100.0	124.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,404	52.1	64.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,616	16.2	20.1
3	Malignant neoplasms (C00-C97)	768	4.8	5.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	634	3.9	4.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	569	3.5	4.4
6	Diabetes mellitus (E10-E14)	150	0.9	1.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	137	0.8	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	136	0.8	1.0
9	Septicemia (A40-A41)	95	0.6	0.7
9	Cerebrovascular diseases (I60-I69)	95	0.6	0.7
9	Pregnancy, childbirth and the puerperium (O00-O99)	95	0.6	0.7
12	Influenza and pneumonia (J09-J18)	92	0.6	0.7
13	Chronic lower respiratory diseases (J40-J47)	63	0.4	0.5
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	42	0.3	0.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	41	0.3	0.3
...	All other causes (residual)	2,206	13.7	17.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 30-34 years				
...	All causes	19,079	100.0	151.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,766	45.9	69.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,664	14.0	21.2
3	Malignant neoplasms (C00-C97)	1,439	7.5	11.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,177	6.2	9.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	571	3.0	4.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	414	2.2	3.3
7	Diabetes mellitus (E10-E14)	254	1.3	2.0
8	Cerebrovascular diseases (I60-I69)	188	1.0	1.5
9	Influenza and pneumonia (J09-J18)	155	0.8	1.2
10	Septicemia (A40-A41)	138	0.7	1.1
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	123	0.6	1.0
12	Pregnancy, childbirth and the puerperium (O00-O99)	106	0.6	0.8
13	Chronic lower respiratory diseases (J40-J47)	104	0.5	0.8
14	Human immunodeficiency virus (HIV) disease (B20-B24)	70	0.4	0.6
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	66	0.3	0.5
...	All other causes (residual)	2,844	14.9	22.6
Non-Hispanic, white, both sexes, 35-39 years				
...	All causes	21,922	100.0	182.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,015	36.6	66.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,769	12.6	23.1
3	Malignant neoplasms (C00-C97)	2,335	10.7	19.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,155	9.8	17.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	698	3.2	5.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	530	2.4	4.4
7	Diabetes mellitus (E10-E14)	438	2.0	3.6
8	Cerebrovascular diseases (I60-I69)	351	1.6	2.9
9	Influenza and pneumonia (J09-J18)	221	1.0	1.8
10	Septicemia (A40-A41)	198	0.9	1.6
11	Chronic lower respiratory diseases (J40-J47)	144	0.7	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	139	0.6	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	128	0.6	1.1
14	Pregnancy, childbirth and the puerperium (O00-O99)	112	0.5	0.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	100	0.5	0.8
...	All other causes (residual)	3,589	16.4	29.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 40-44 years				
...	All causes	26,487	100.0	230.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,042	26.6	61.1
2	Malignant neoplasms (C00-C97)	4,079	15.4	35.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,836	14.5	33.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,759	10.4	24.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,093	4.1	9.5
6	Diabetes mellitus (E10-E14)	642	2.4	5.6
7	Cerebrovascular diseases (I60-I69)	518	2.0	4.5
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	495	1.9	4.3
9	Septicemia (A40-A41)	317	1.2	2.8
10	Influenza and pneumonia (J09-J18)	311	1.2	2.7
11	Chronic lower respiratory diseases (J40-J47)	285	1.1	2.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	183	0.7	1.6
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	160	0.6	1.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	159	0.6	1.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	130	0.5	1.1
...	All other causes (residual)	4,478	16.9	38.9
Non-Hispanic, white, both sexes, 45-49 years				
...	All causes	42,994	100.0	326.1
1	Malignant neoplasms (C00-C97)	8,999	20.9	68.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,960	18.5	60.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,668	17.8	58.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,288	7.6	24.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,059	4.8	15.6
6	Diabetes mellitus (E10-E14)	1,301	3.0	9.9
7	Cerebrovascular diseases (I60-I69)	993	2.3	7.5
8	Chronic lower respiratory diseases (J40-J47)	843	2.0	6.4
9	Septicemia (A40-A41)	507	1.2	3.8
10	Influenza and pneumonia (J09-J18)	500	1.2	3.8
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	478	1.1	3.6
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	380	0.9	2.9
13	Human immunodeficiency virus (HIV) disease (B20-B24)	274	0.6	2.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	267	0.6	2.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	233	0.5	1.8
...	All other causes (residual)	7,244	16.8	54.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 50-54 years				
...	All causes	73,640	100.0	505.0
1	Malignant neoplasms (C00-C97)	19,038	25.9	130.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,550	19.8	99.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,013	12.2	61.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,963	5.4	27.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,790	5.1	26.0
6	Chronic lower respiratory diseases (J40-J47)	2,605	3.5	17.9
7	Diabetes mellitus (E10-E14)	2,327	3.2	16.0
8	Cerebrovascular diseases (I60-I69)	1,835	2.5	12.6
9	Septicemia (A40-A41)	998	1.4	6.8
10	Influenza and pneumonia (J09-J18)	854	1.2	5.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	673	0.9	4.6
12	Viral hepatitis (B15-B19)	555	0.8	3.8
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	539	0.7	3.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	489	0.7	3.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	376	0.5	2.6
...	All other causes (residual)	12,035	16.3	82.5
Non-Hispanic, white, both sexes, 55-59 years				
...	All causes	114,682	100.0	738.9
1	Malignant neoplasms (C00-C97)	35,526	31.0	228.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,539	20.5	151.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,124	8.0	58.8
4	Chronic lower respiratory diseases (J40-J47)	5,701	5.0	36.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	4,985	4.3	32.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,878	3.4	25.0
7	Diabetes mellitus (E10-E14)	3,615	3.2	23.3
8	Cerebrovascular diseases (I60-I69)	3,119	2.7	20.1
9	Septicemia (A40-A41)	1,678	1.5	10.8
10	Influenza and pneumonia (J09-J18)	1,430	1.2	9.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,262	1.1	8.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	953	0.8	6.1
13	Viral hepatitis (B15-B19)	903	0.8	5.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	529	0.5	3.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	465	0.4	3.0
...	All other causes (residual)	17,975	15.7	115.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 60-64 years				
...	All causes	147,548	100.0	1,034.3
1	Malignant neoplasms (C00-C97)	49,585	33.6	347.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,490	21.3	220.7
3	Chronic lower respiratory diseases (J40-J47)	9,130	6.2	64.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,961	4.7	48.8
5	Diabetes mellitus (E10-E14)	5,084	3.4	35.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,815	3.3	33.8
7	Cerebrovascular diseases (I60-I69)	4,382	3.0	30.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,023	2.0	21.2
9	Septicemia (A40-A41)	2,289	1.6	16.0
10	Influenza and pneumonia (J09-J18)	2,064	1.4	14.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,914	1.3	13.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,297	0.9	9.1
13	Viral hepatitis (B15-B19)	1,075	0.7	7.5
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	710	0.5	5.0
15	Pneumonitis due to solids and liquids (J69)	696	0.5	4.9
...	All other causes (residual)	23,033	15.6	161.5
Non-Hispanic, white, both sexes, 65-69 years				
...	All causes	186,455	100.0	1,457.7
1	Malignant neoplasms (C00-C97)	64,053	34.4	500.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,929	21.4	312.2
3	Chronic lower respiratory diseases (J40-J47)	14,260	7.6	111.5
4	Diabetes mellitus (E10-E14)	6,556	3.5	51.3
5	Cerebrovascular diseases (I60-I69)	6,429	3.4	50.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,806	3.1	45.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	3,961	2.1	31.0
8	Septicemia (A40-A41)	3,016	1.6	23.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,820	1.5	22.0
10	Influenza and pneumonia (J09-J18)	2,702	1.4	21.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,292	1.2	17.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,638	0.9	12.8
13	Alzheimer's disease (G30)	1,637	0.9	12.8
14	Parkinson's disease (G20-G21)	1,123	0.6	8.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,081	0.6	8.5
...	All other causes (residual)	29,152	15.6	227.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 70-74 years				
...	All causes	208,699	100.0	2,278.8
1	Malignant neoplasms (C00-C97)	66,597	31.9	727.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	44,688	21.4	488.0
3	Chronic lower respiratory diseases (J40-J47)	19,434	9.3	212.2
4	Cerebrovascular diseases (I60-I69)	8,700	4.2	95.0
5	Diabetes mellitus (E10-E14)	6,805	3.3	74.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,185	2.5	56.6
7	Alzheimer's disease (G30)	3,887	1.9	42.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,618	1.7	39.5
9	Septicemia (A40-A41)	3,314	1.6	36.2
10	Influenza and pneumonia (J09-J18)	3,304	1.6	36.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,788	1.3	30.4
12	Parkinson's disease (G20-G21)	2,338	1.1	25.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,885	0.9	20.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,715	0.8	18.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,474	0.7	16.1
...	All other causes (residual)	32,967	15.8	360.0
Non-Hispanic, white, both sexes, 75-79 years				
...	All causes	235,476	100.0	3,606.7
1	Malignant neoplasms (C00-C97)	65,008	27.6	995.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	51,484	21.9	788.6
3	Chronic lower respiratory diseases (J40-J47)	21,798	9.3	333.9
4	Cerebrovascular diseases (I60-I69)	11,874	5.0	181.9
5	Alzheimer's disease (G30)	8,553	3.6	131.0
6	Diabetes mellitus (E10-E14)	6,686	2.8	102.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,915	2.5	90.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,518	1.9	69.2
9	Parkinson's disease (G20-G21)	4,304	1.8	65.9
10	Influenza and pneumonia (J09-J18)	4,225	1.8	64.7
11	Septicemia (A40-A41)	3,758	1.6	57.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,230	0.9	34.2
13	Chronic liver disease and cirrhosis (K70,K73-K74)	2,004	0.9	30.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,781	0.8	27.3
15	Pneumonitis due to solids and liquids (J69)	1,758	0.7	26.9
...	All other causes (residual)	39,580	16.8	606.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 80-84 years				
...	All causes	279,076	100.0	6,011.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	66,196	23.7	1,425.9
2	Malignant neoplasms (C00-C97)	60,246	21.6	1,297.7
3	Chronic lower respiratory diseases (J40-J47)	22,226	8.0	478.8
4	Cerebrovascular diseases (I60-I69)	17,115	6.1	368.7
5	Alzheimer's disease (G30)	16,710	6.0	359.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,450	2.7	160.5
7	Diabetes mellitus (E10-E14)	6,914	2.5	148.9
8	Parkinson's disease (G20-G21)	6,211	2.2	133.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,614	2.0	120.9
10	Influenza and pneumonia (J09-J18)	5,595	2.0	120.5
11	Septicemia (A40-A41)	4,300	1.5	92.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,090	1.1	66.6
13	Pneumonitis due to solids and liquids (J69)	2,445	0.9	52.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,187	0.8	47.1
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1,441	0.5	31.0
...	All other causes (residual)	51,336	18.4	1,105.8
Non-Hispanic, white, both sexes, 85-89 years				
...	All causes	324,585	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	87,001	26.8	---
2	Malignant neoplasms (C00-C97)	50,930	15.7	---
3	Alzheimer's disease (G30)	26,157	8.1	---
4	Cerebrovascular diseases (I60-I69)	22,186	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	20,545	6.3	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,324	2.9	---
7	Influenza and pneumonia (J09-J18)	7,371	2.3	---
8	Diabetes mellitus (E10-E14)	6,569	2.0	---
9	Parkinson's disease (G20-G21)	6,559	2.0	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,401	2.0	---
11	Septicemia (A40-A41)	4,595	1.4	---
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,186	1.3	---
13	Pneumonitis due to solids and liquids (J69)	3,342	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2,370	0.7	---
15	Aortic aneurysm and dissection (I71)	1,180	0.4	---
...	All other causes (residual)	65,869	20.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 90-94 years				
...	All causes	267,709	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	80,586	30.1	---
2	Malignant neoplasms (C00-C97)	27,738	10.4	---
3	Alzheimer's disease (G30)	25,656	9.6	---
4	Cerebrovascular diseases (I60-I69)	19,465	7.3	---
5	Chronic lower respiratory diseases (J40-J47)	13,053	4.9	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,839	2.9	---
7	Influenza and pneumonia (J09-J18)	6,768	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5,124	1.9	---
9	Diabetes mellitus (E10-E14)	4,344	1.6	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,109	1.5	---
11	Parkinson's disease (G20-G21)	3,694	1.4	---
12	Septicemia (A40-A41)	3,274	1.2	---
13	Pneumonitis due to solids and liquids (J69)	3,143	1.2	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,550	0.6	---
15	Nutritional deficiencies (E40-E64)	1,033	0.4	---
...	All other causes (residual)	60,333	22.5	---
Non-Hispanic, white, both sexes, 95-99 years				
...	All causes	113,496	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	37,091	32.7	---
2	Alzheimer's disease (G30)	11,849	10.4	---
3	Cerebrovascular diseases (I60-I69)	8,476	7.5	---
4	Malignant neoplasms (C00-C97)	7,699	6.8	---
5	Chronic lower respiratory diseases (J40-J47)	4,248	3.7	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,230	2.8	---
7	Influenza and pneumonia (J09-J18)	3,149	2.8	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,150	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,072	1.8	---
10	Pneumonitis due to solids and liquids (J69)	1,380	1.2	---
11	Diabetes mellitus (E10-E14)	1,367	1.2	---
12	Septicemia (A40-A41)	1,123	1.0	---
13	Parkinson's disease (G20-G21)	882	0.8	---
14	Nutritional deficiencies (E40-E64)	591	0.5	---
15	Atherosclerosis (I70)	552	0.5	---
...	All other causes (residual)	27,637	24.4	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 100 years and over				
...	All causes	23,096	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,849	34.0	---
2	Alzheimer's disease (G30)	2,350	10.2	---
3	Cerebrovascular diseases (I60-I69)	1,576	6.8	---
4	Malignant neoplasms (C00-C97)	1,004	4.3	---
5	Influenza and pneumonia (J09-J18)	755	3.3	---
6	Chronic lower respiratory diseases (J40-J47)	671	2.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	634	2.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	586	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	367	1.6	---
10	Pneumonitis due to solids and liquids (J69)	266	1.2	---
11	Diabetes mellitus (E10-E14)	216	0.9	---
12	Septicemia (A40-A41)	198	0.9	---
13	Nutritional deficiencies (E40-E64)	182	0.8	---
14	Atherosclerosis (I70)	155	0.7	---
15	Parkinson's disease (G20-G21)	96	0.4	---
...	All other causes (residual)	6,191	26.8	---
Non-Hispanic, white, male, all ages				
...	All causes	1,077,362	100.0	1,085.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	266,995	24.8	269.0
2	Malignant neoplasms (C00-C97)	247,202	22.9	249.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	76,033	7.1	76.6
4	Chronic lower respiratory diseases (J40-J47)	62,945	5.8	63.4
5	Cerebrovascular diseases (I60-I69)	43,713	4.1	44.0
6	Diabetes mellitus (E10-E14)	30,010	2.8	30.2
7	Alzheimer's disease (G30)	29,886	2.8	30.1
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28,110	2.6	28.3
9	Influenza and pneumonia (J09-J18)	19,033	1.8	19.2
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18,522	1.7	18.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,216	1.7	18.4
12	Parkinson's disease (G20-G21)	15,641	1.5	15.8
13	Septicemia (A40-A41)	14,620	1.4	14.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9,906	0.9	10.0
15	Pneumonitis due to solids and liquids (J69)	8,997	0.8	9.1
...	All other causes (residual)	187,533	17.4	189.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,113	100.0	26.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	419	37.6	9.9
2	Malignant neoplasms (C00-C97)	114	10.2	2.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	95	8.5	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	5.6	1.5
5	Septicemia (A40-A41)	23	2.1	0.5
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.1	0.5
7	Influenza and pneumonia (J09-J18)	21	1.9	0.5
8	Chronic lower respiratory diseases (J40-J47)	14	1.3	*
9	Cerebrovascular diseases (I60-I69)	12	1.1	*
9	Certain conditions originating in the perinatal period (P00-P96)	12	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.8	*
12	Acute bronchitis and bronchiolitis (J20-J21)	4	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
14	Meningococcal infection (A39)	2	0.2	*
14	Anemias (D50-D64)	2	0.2	*
14	Meningitis (G00,G03)	2	0.2	*
14	Pneumonitis due to solids and liquids (J69)	2	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.2	*
...	All other causes (residual)	288	25.9	6.8
Non-Hispanic, white, male, 5-9 years				
...	All causes	649	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	218	33.6	4.0
2	Malignant neoplasms (C00-C97)	135	20.8	2.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	59	9.1	1.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	34	5.2	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.1	0.4
6	Influenza and pneumonia (J09-J18)	14	2.2	*
7	Septicemia (A40-A41)	10	1.5	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	1.5	*
9	Cerebrovascular diseases (I60-I69)	8	1.2	*
10	Anemias (D50-D64)	3	0.5	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
10	Certain conditions originating in the perinatal period (P00-P96)	3	0.5	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
14	Diabetes mellitus (E10-E14)	2	0.3	*
14	Chronic lower respiratory diseases (J40-J47)	2	0.3	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2	0.3	*
...	All other causes (residual)	123	19.0	2.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 10-14 years				
...	All causes	976	100.0	17.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	31.3	5.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	192	19.7	3.4
3	Malignant neoplasms (C00-C97)	129	13.2	2.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	48	4.9	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	2.7	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	2.3	0.4
7	Chronic lower respiratory diseases (J40-J47)	15	1.5	*
8	Cerebrovascular diseases (I60-I69)	13	1.3	*
9	Influenza and pneumonia (J09-J18)	10	1.0	*
10	Septicemia (A40-A41)	9	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	3	0.3	*
14	Anemias (D50-D64)	2	0.2	*
14	Diabetes mellitus (E10-E14)	2	0.2	*
14	Meningitis (G00,G03)	2	0.2	*
14	Aortic aneurysm and dissection (I71)	2	0.2	*
14	Pneumonitis due to solids and liquids (J69)	2	0.2	*
...	All other causes (residual)	182	18.6	3.2
Non-Hispanic, white, male, 15-19 years				
...	All causes	3,924	100.0	65.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,772	45.2	29.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,097	28.0	18.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	206	5.2	3.4
4	Malignant neoplasms (C00-C97)	194	4.9	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	2.2	1.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	1.7	1.1
7	Diabetes mellitus (E10-E14)	20	0.5	0.3
7	Influenza and pneumonia (J09-J18)	20	0.5	0.3
9	Cerebrovascular diseases (I60-I69)	16	0.4	*
9	Chronic lower respiratory diseases (J40-J47)	16	0.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	0.4	*
12	Septicemia (A40-A41)	11	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.1	*
14	Pneumonitis due to solids and liquids (J69)	4	0.1	*
14	Legal intervention (Y35,Y89.0)	4	0.1	*
...	All other causes (residual)	395	10.1	6.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 20-24 years				
...	All causes	8,721	100.0	136.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,785	54.9	74.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,945	22.3	30.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	343	3.9	5.4
4	Malignant neoplasms (C00-C97)	270	3.1	4.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	231	2.6	3.6
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	0.6	0.8
7	Diabetes mellitus (E10-E14)	45	0.5	0.7
8	Influenza and pneumonia (J09-J18)	37	0.4	0.6
8	Chronic lower respiratory diseases (J40-J47)	37	0.4	0.6
10	Septicemia (A40-A41)	32	0.4	0.5
10	Cerebrovascular diseases (I60-I69)	32	0.4	0.5
12	Legal intervention (Y35,Y89.0)	16	0.2	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	14	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.1	*
15	Pneumonitis due to solids and liquids (J69)	9	0.1	*
...	All other causes (residual)	861	9.9	13.4
Non-Hispanic, white, male, 25-29 years				
...	All causes	11,449	100.0	173.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,208	54.2	94.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,123	18.5	32.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	425	3.7	6.4
4	Malignant neoplasms (C00-C97)	408	3.6	6.2
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	406	3.5	6.1
6	Diabetes mellitus (E10-E14)	85	0.7	1.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	79	0.7	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	66	0.6	1.0
9	Cerebrovascular diseases (I60-I69)	53	0.5	0.8
10	Influenza and pneumonia (J09-J18)	50	0.4	0.8
11	Septicemia (A40-A41)	43	0.4	0.7
12	Chronic lower respiratory diseases (J40-J47)	38	0.3	0.6
13	Legal intervention (Y35,Y89.0)	35	0.3	0.5
14	Human immunodeficiency virus (HIV) disease (B20-B24)	34	0.3	0.5
15	Pneumonitis due to solids and liquids (J69)	19	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	19	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	19	0.2	*
...	All other causes (residual)	1,339	11.7	20.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 30-34 years				
...	All causes	12,844	100.0	202.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,325	49.2	99.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,076	16.2	32.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	802	6.2	12.6
4	Malignant neoplasms (C00-C97)	685	5.3	10.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	406	3.2	6.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	240	1.9	3.8
7	Diabetes mellitus (E10-E14)	150	1.2	2.4
8	Cerebrovascular diseases (I60-I69)	103	0.8	1.6
9	Influenza and pneumonia (J09-J18)	79	0.6	1.2
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	69	0.5	1.1
11	Septicemia (A40-A41)	65	0.5	1.0
12	Human immunodeficiency virus (HIV) disease (B20-B24)	45	0.4	0.7
12	Legal intervention (Y35,Y89.0)	45	0.4	0.7
14	Chronic lower respiratory diseases (J40-J47)	44	0.3	0.7
15	Aortic aneurysm and dissection (I71)	41	0.3	0.6
...	All other causes (residual)	1,669	13.0	26.3
Non-Hispanic, white, male, 35-39 years				
...	All causes	14,148	100.0	233.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,662	40.0	93.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,079	14.7	34.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,437	10.2	23.8
4	Malignant neoplasms (C00-C97)	1,023	7.2	16.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	403	2.8	6.7
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	367	2.6	6.1
7	Diabetes mellitus (E10-E14)	281	2.0	4.6
8	Cerebrovascular diseases (I60-I69)	207	1.5	3.4
9	Influenza and pneumonia (J09-J18)	128	0.9	2.1
10	Septicemia (A40-A41)	99	0.7	1.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	83	0.6	1.4
12	Chronic lower respiratory diseases (J40-J47)	79	0.6	1.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	78	0.6	1.3
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	67	0.5	1.1
15	Aortic aneurysm and dissection (I71)	51	0.4	0.8
15	Legal intervention (Y35,Y89.0)	51	0.4	0.8
...	All other causes (residual)	2,053	14.5	33.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 40-44 years				
...	All causes	16,374	100.0	283.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,871	29.7	84.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,661	16.3	46.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,031	12.4	35.1
4	Malignant neoplasms (C00-C97)	1,847	11.3	31.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	656	4.0	11.3
6	Diabetes mellitus (E10-E14)	385	2.4	6.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	327	2.0	5.7
8	Cerebrovascular diseases (I60-I69)	278	1.7	4.8
9	Septicemia (A40-A41)	161	1.0	2.8
10	Influenza and pneumonia (J09-J18)	152	0.9	2.6
11	Chronic lower respiratory diseases (J40-J47)	142	0.9	2.5
12	Human immunodeficiency virus (HIV) disease (B20-B24)	123	0.8	2.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	101	0.6	1.7
14	Aortic aneurysm and dissection (I71)	80	0.5	1.4
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	0.5	1.3
...	All other causes (residual)	2,483	15.2	42.9
Non-Hispanic, white, male, 45-49 years				
...	All causes	26,151	100.0	396.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,422	20.7	82.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,322	20.4	80.7
3	Malignant neoplasms (C00-C97)	4,141	15.8	62.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,406	9.2	36.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,293	4.9	19.6
6	Diabetes mellitus (E10-E14)	883	3.4	13.4
7	Cerebrovascular diseases (I60-I69)	535	2.0	8.1
8	Chronic lower respiratory diseases (J40-J47)	368	1.4	5.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	331	1.3	5.0
10	Septicemia (A40-A41)	267	1.0	4.0
11	Influenza and pneumonia (J09-J18)	249	1.0	3.8
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	228	0.9	3.5
13	Human immunodeficiency virus (HIV) disease (B20-B24)	227	0.9	3.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	190	0.7	2.9
15	Viral hepatitis (B15-B19)	133	0.5	2.0
...	All other causes (residual)	4,156	15.9	63.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 50-54 years				
...	All causes	44,820	100.0	620.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	10,328	23.0	142.9
2	Malignant neoplasms (C00-C97)	9,770	21.8	135.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,962	13.3	82.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,869	6.4	39.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,388	5.3	33.0
6	Diabetes mellitus (E10-E14)	1,517	3.4	21.0
7	Chronic lower respiratory diseases (J40-J47)	1,172	2.6	16.2
8	Cerebrovascular diseases (I60-I69)	1,031	2.3	14.3
9	Septicemia (A40-A41)	538	1.2	7.4
10	Influenza and pneumonia (J09-J18)	453	1.0	6.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	406	0.9	5.6
12	Viral hepatitis (B15-B19)	376	0.8	5.2
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	353	0.8	4.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	339	0.8	4.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	323	0.7	4.5
...	All other causes (residual)	6,995	15.6	96.8
Non-Hispanic, white, male, 55-59 years				
...	All causes	70,093	100.0	919.3
1	Malignant neoplasms (C00-C97)	19,267	27.5	252.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,711	23.8	219.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,167	8.8	80.9
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,313	4.7	43.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,854	4.1	37.4
6	Chronic lower respiratory diseases (J40-J47)	2,771	4.0	36.3
7	Diabetes mellitus (E10-E14)	2,363	3.4	31.0
8	Cerebrovascular diseases (I60-I69)	1,771	2.5	23.2
9	Septicemia (A40-A41)	940	1.3	12.3
10	Influenza and pneumonia (J09-J18)	790	1.1	10.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	712	1.0	9.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	636	0.9	8.3
13	Viral hepatitis (B15-B19)	626	0.9	8.2
14	Aortic aneurysm and dissection (I71)	338	0.5	4.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	284	0.4	3.7
...	All other causes (residual)	10,550	15.1	138.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 60-64 years				
...	All causes	89,957	100.0	1,300.6
1	Malignant neoplasms (C00-C97)	27,713	30.8	400.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,059	24.5	318.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,781	5.3	69.1
4	Chronic lower respiratory diseases (J40-J47)	4,744	5.3	68.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	3,309	3.7	47.8
6	Diabetes mellitus (E10-E14)	3,176	3.5	45.9
7	Cerebrovascular diseases (I60-I69)	2,596	2.9	37.5
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,234	2.5	32.3
9	Septicemia (A40-A41)	1,245	1.4	18.0
10	Influenza and pneumonia (J09-J18)	1,136	1.3	16.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,136	1.3	16.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	823	0.9	11.9
13	Viral hepatitis (B15-B19)	755	0.8	10.9
14	Aortic aneurysm and dissection (I71)	439	0.5	6.3
15	Pneumonitis due to solids and liquids (J69)	430	0.5	6.2
...	All other causes (residual)	13,381	14.9	193.5
Non-Hispanic, white, male, 65-69 years				
...	All causes	109,782	100.0	1,790.2
1	Malignant neoplasms (C00-C97)	35,827	32.6	584.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,898	24.5	438.6
3	Chronic lower respiratory diseases (J40-J47)	7,325	6.7	119.4
4	Diabetes mellitus (E10-E14)	3,979	3.6	64.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,802	3.5	62.0
6	Cerebrovascular diseases (I60-I69)	3,478	3.2	56.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2,660	2.4	43.4
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,772	1.6	28.9
9	Septicemia (A40-A41)	1,624	1.5	26.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,570	1.4	25.6
11	Influenza and pneumonia (J09-J18)	1,537	1.4	25.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	944	0.9	15.4
13	Parkinson's disease (G20-G21)	745	0.7	12.1
14	Alzheimer's disease (G30)	725	0.7	11.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	640	0.6	10.4
...	All other causes (residual)	16,256	14.8	265.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 70-74 years				
...	All causes	117,232	100.0	2,725.5
1	Malignant neoplasms (C00-C97)	36,689	31.3	853.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,253	24.1	656.8
3	Chronic lower respiratory diseases (J40-J47)	9,722	8.3	226.0
4	Cerebrovascular diseases (I60-I69)	4,547	3.9	105.7
5	Diabetes mellitus (E10-E14)	4,053	3.5	94.2
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,254	2.8	75.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,953	1.7	45.4
8	Influenza and pneumonia (J09-J18)	1,801	1.5	41.9
9	Septicemia (A40-A41)	1,706	1.5	39.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,657	1.4	38.5
11	Parkinson's disease (G20-G21)	1,582	1.3	36.8
12	Alzheimer's disease (G30)	1,561	1.3	36.3
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,384	1.2	32.2
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,009	0.9	23.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	870	0.7	20.2
...	All other causes (residual)	17,191	14.7	399.7
Non-Hispanic, white, male, 75-79 years				
...	All causes	126,471	100.0	4,280.1
1	Malignant neoplasms (C00-C97)	35,143	27.8	1,189.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,479	24.1	1,031.5
3	Chronic lower respiratory diseases (J40-J47)	10,555	8.3	357.2
4	Cerebrovascular diseases (I60-I69)	5,790	4.6	195.9
5	Diabetes mellitus (E10-E14)	3,868	3.1	130.9
6	Alzheimer's disease (G30)	3,424	2.7	115.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,421	2.7	115.8
8	Parkinson's disease (G20-G21)	2,868	2.3	97.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,458	1.9	83.2
10	Influenza and pneumonia (J09-J18)	2,273	1.8	76.9
11	Septicemia (A40-A41)	1,891	1.5	64.0
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,149	0.9	38.9
13	Chronic liver disease and cirrhosis (K70,K73-K74)	1,128	0.9	38.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,054	0.8	35.7
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,038	0.8	35.1
...	All other causes (residual)	19,932	15.8	674.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 80-84 years				
...	All causes	138,949	100.0	7,030.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	35,674	25.7	1,805.1
2	Malignant neoplasms (C00-C97)	31,887	22.9	1,613.5
3	Chronic lower respiratory diseases (J40-J47)	10,205	7.3	516.4
4	Cerebrovascular diseases (I60-I69)	7,441	5.4	376.5
5	Alzheimer's disease (G30)	6,230	4.5	315.2
6	Parkinson's disease (G20-G21)	3,917	2.8	198.2
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,886	2.8	196.6
8	Diabetes mellitus (E10-E14)	3,663	2.6	185.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,056	2.2	154.6
10	Influenza and pneumonia (J09-J18)	2,837	2.0	143.6
11	Septicemia (A40-A41)	2,069	1.5	104.7
12	Pneumonitis due to solids and liquids (J69)	1,415	1.0	71.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,352	1.0	68.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,229	0.9	62.2
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	900	0.6	45.5
...	All other causes (residual)	23,188	16.7	1,173.3
Non-Hispanic, white, male, 85-89 years				
...	All causes	143,664	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,072	28.6	---
2	Malignant neoplasms (C00-C97)	25,749	17.9	---
3	Chronic lower respiratory diseases (J40-J47)	9,005	6.3	---
4	Alzheimer's disease (G30)	8,337	5.8	---
5	Cerebrovascular diseases (I60-I69)	7,944	5.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,362	3.0	---
7	Parkinson's disease (G20-G21)	3,801	2.6	---
8	Influenza and pneumonia (J09-J18)	3,426	2.4	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,239	2.3	---
10	Diabetes mellitus (E10-E14)	3,205	2.2	---
11	Septicemia (A40-A41)	2,137	1.5	---
12	Pneumonitis due to solids and liquids (J69)	1,890	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,588	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,234	0.9	---
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	659	0.5	---
...	All other causes (residual)	26,016	18.1	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 90-94 years				
...	All causes	98,304	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	31,401	31.9	---
2	Malignant neoplasms (C00-C97)	12,827	13.0	---
3	Alzheimer's disease (G30)	6,786	6.9	---
4	Cerebrovascular diseases (I60-I69)	5,658	5.8	---
5	Chronic lower respiratory diseases (J40-J47)	5,155	5.2	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,056	3.1	---
7	Influenza and pneumonia (J09-J18)	2,760	2.8	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,328	2.4	---
9	Parkinson's disease (G20-G21)	1,901	1.9	---
10	Diabetes mellitus (E10-E14)	1,809	1.8	---
11	Pneumonitis due to solids and liquids (J69)	1,590	1.6	---
12	Septicemia (A40-A41)	1,311	1.3	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,230	1.3	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	677	0.7	---
15	Nutritional deficiencies (E40-E64)	364	0.4	---
...	All other causes (residual)	19,451	19.8	---
Non-Hispanic, white, male, 95-99 years				
...	All causes	31,954	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	11,335	35.5	---
2	Malignant neoplasms (C00-C97)	3,074	9.6	---
3	Alzheimer's disease (G30)	2,188	6.8	---
4	Cerebrovascular diseases (I60-I69)	1,933	6.0	---
5	Chronic lower respiratory diseases (J40-J47)	1,379	4.3	---
6	Influenza and pneumonia (J09-J18)	1,042	3.3	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	984	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	766	2.4	---
9	Pneumonitis due to solids and liquids (J69)	604	1.9	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	498	1.6	---
11	Diabetes mellitus (E10-E14)	471	1.5	---
12	Parkinson's disease (G20-G21)	396	1.2	---
13	Septicemia (A40-A41)	354	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	176	0.6	---
15	Nutritional deficiencies (E40-E64)	151	0.5	---
...	All other causes (residual)	6,603	20.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 100 years and over				
...	All causes	4,132	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,580	38.2	---
2	Malignant neoplasms (C00-C97)	294	7.1	---
3	Cerebrovascular diseases (I60-I69)	229	5.5	---
4	Alzheimer's disease (G30)	228	5.5	---
5	Influenza and pneumonia (J09-J18)	174	4.2	---
6	Chronic lower respiratory diseases (J40-J47)	144	3.5	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	111	2.7	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	2.3	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	92	2.2	---
10	Pneumonitis due to solids and liquids (J69)	91	2.2	---
11	Diabetes mellitus (E10-E14)	52	1.3	---
12	Septicemia (A40-A41)	44	1.1	---
13	Parkinson's disease (G20-G21)	30	0.7	---
14	Atherosclerosis (I70)	26	0.6	---
15	Nutritional deficiencies (E40-E64)	25	0.6	---
...	All other causes (residual)	916	22.2	---
Non-Hispanic, white, female, all ages				
...	All causes	1,056,101	100.0	1,034.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	233,636	22.1	228.9
2	Malignant neoplasms (C00-C97)	219,265	20.8	214.8
3	Chronic lower respiratory diseases (J40-J47)	72,323	6.8	70.8
4	Alzheimer's disease (G30)	67,893	6.4	66.5
5	Cerebrovascular diseases (I60-I69)	63,778	6.0	62.5
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45,608	4.3	44.7
7	Diabetes mellitus (E10-E14)	23,389	2.2	22.9
8	Influenza and pneumonia (J09-J18)	20,715	2.0	20.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,032	1.6	16.7
10	Septicemia (A40-A41)	15,389	1.5	15.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,331	1.3	13.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	10,910	1.0	10.7
13	Parkinson's disease (G20-G21)	10,185	1.0	10.0
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,421	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	7,305	0.7	7.2
...	All other causes (residual)	226,921	21.5	222.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 1-4 years				
...	All causes	865	100.0	21.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	261	30.2	6.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	96	11.1	2.4
3	Malignant neoplasms (C00-C97)	92	10.6	2.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	6.8	1.5
5	Influenza and pneumonia (J09-J18)	26	3.0	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	2.5	0.5
7	Cerebrovascular diseases (I60-I69)	15	1.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.6	*
9	Septicemia (A40-A41)	11	1.3	*
10	Chronic lower respiratory diseases (J40-J47)	8	0.9	*
10	Certain conditions originating in the perinatal period (P00-P96)	8	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.5	*
14	Meningitis (G00,G03)	3	0.3	*
14	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (residual)	238	27.5	5.9
Non-Hispanic, white, female, 5-9 years				
...	All causes	509	100.0	9.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	27.9	2.7
2	Malignant neoplasms (C00-C97)	86	16.9	1.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	49	9.6	0.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	19	3.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	3.7	*
6	Influenza and pneumonia (J09-J18)	12	2.4	*
7	Septicemia (A40-A41)	10	2.0	*
7	Chronic lower respiratory diseases (J40-J47)	10	2.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	1.4	*
10	Diabetes mellitus (E10-E14)	6	1.2	*
11	Certain conditions originating in the perinatal period (P00-P96)	4	0.8	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.8	*
13	Cerebrovascular diseases (I60-I69)	3	0.6	*
14	Meningitis (G00,G03)	2	0.4	*
14	Hernia (K40-K46)	2	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
...	All other causes (residual)	132	25.9	2.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 10-14 years				
...	All causes	647	100.0	11.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	179	27.7	3.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	117	18.1	2.2
3	Malignant neoplasms (C00-C97)	90	13.9	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	32	4.9	0.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.2	0.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	18	2.8	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.4	*
8	Influenza and pneumonia (J09-J18)	8	1.2	*
9	Cerebrovascular diseases (I60-I69)	7	1.1	*
10	Septicemia (A40-A41)	5	0.8	*
10	Diabetes mellitus (E10-E14)	5	0.8	*
10	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	Diseases of appendix (K35-K38)	2	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (residual)	144	22.3	2.6
Non-Hispanic, white, female, 15-19 years				
...	All causes	1,744	100.0	30.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	773	44.3	13.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	325	18.6	5.7
3	Malignant neoplasms (C00-C97)	124	7.1	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	78	4.5	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	64	3.7	1.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	2.2	0.7
7	Diabetes mellitus (E10-E14)	13	0.7	*
7	Influenza and pneumonia (J09-J18)	13	0.7	*
9	Cerebrovascular diseases (I60-I69)	10	0.6	*
10	Septicemia (A40-A41)	8	0.5	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	8	0.5	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.4	*
14	Pneumonitis due to solids and liquids (J69)	6	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	0.3	*
...	All other causes (residual)	263	15.1	4.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 20-24 years				
...	All causes	3,126	100.0	51.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,565	50.1	25.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	389	12.4	6.4
3	Malignant neoplasms (C00-C97)	156	5.0	2.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	139	4.4	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	101	3.2	1.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	71	2.3	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	52	1.7	0.9
8	Diabetes mellitus (E10-E14)	35	1.1	0.6
9	Influenza and pneumonia (J09-J18)	30	1.0	0.5
10	Chronic lower respiratory diseases (J40-J47)	27	0.9	0.4
11	Septicemia (A40-A41)	23	0.7	0.4
12	Cerebrovascular diseases (I60-I69)	13	0.4	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	10	0.3	*
14	Aortic aneurysm and dissection (I71)	9	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.3	*
...	All other causes (residual)	498	15.9	8.2
Non-Hispanic, white, female, 25-29 years				
...	All causes	4,694	100.0	73.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,196	46.8	34.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	493	10.5	7.7
3	Malignant neoplasms (C00-C97)	360	7.7	5.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	209	4.5	3.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	3.5	2.5
6	Pregnancy, childbirth and the puerperium (O00-O99)	95	2.0	1.5
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	70	1.5	1.1
8	Diabetes mellitus (E10-E14)	65	1.4	1.0
9	Chronic liver disease and cirrhosis (K70,K73-K74)	58	1.2	0.9
10	Septicemia (A40-A41)	52	1.1	0.8
11	Cerebrovascular diseases (I60-I69)	42	0.9	0.7
11	Influenza and pneumonia (J09-J18)	42	0.9	0.7
13	Chronic lower respiratory diseases (J40-J47)	25	0.5	0.4
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.5	0.4
15	Pneumonitis due to solids and liquids (J69)	17	0.4	*
...	All other causes (residual)	784	16.7	12.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 30-34 years				
...	All causes	6,235	100.0	100.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,441	39.1	39.2
2	Malignant neoplasms (C00-C97)	754	12.1	12.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	588	9.4	9.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	375	6.0	6.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	174	2.8	2.8
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	165	2.6	2.6
7	Pregnancy, childbirth and the puerperium (O00-O99)	106	1.7	1.7
8	Diabetes mellitus (E10-E14)	104	1.7	1.7
9	Cerebrovascular diseases (I60-I69)	85	1.4	1.4
10	Influenza and pneumonia (J09-J18)	76	1.2	1.2
11	Septicemia (A40-A41)	73	1.2	1.2
12	Chronic lower respiratory diseases (J40-J47)	60	1.0	1.0
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	0.9	0.9
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	0.5	0.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	25	0.4	0.4
...	All other causes (residual)	1,125	18.0	18.0
Non-Hispanic, white, female, 35-39 years				
...	All causes	7,774	100.0	130.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,353	30.3	39.5
2	Malignant neoplasms (C00-C97)	1,312	16.9	22.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	718	9.2	12.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	690	8.9	11.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	295	3.8	4.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	163	2.1	2.7
7	Diabetes mellitus (E10-E14)	157	2.0	2.6
8	Cerebrovascular diseases (I60-I69)	144	1.9	2.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	112	1.4	1.9
10	Septicemia (A40-A41)	99	1.3	1.7
11	Influenza and pneumonia (J09-J18)	93	1.2	1.6
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	72	0.9	1.2
13	Chronic lower respiratory diseases (J40-J47)	65	0.8	1.1
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	0.6	0.8
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	0.4	0.5
...	All other causes (residual)	1,427	18.4	23.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 40-44 years				
...	All causes	10,113	100.0	176.4
1	Malignant neoplasms (C00-C97)	2,232	22.1	38.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,171	21.5	37.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,175	11.6	20.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	728	7.2	12.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	437	4.3	7.6
6	Diabetes mellitus (E10-E14)	257	2.5	4.5
7	Cerebrovascular diseases (I60-I69)	240	2.4	4.2
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	1.7	2.9
9	Influenza and pneumonia (J09-J18)	159	1.6	2.8
10	Septicemia (A40-A41)	156	1.5	2.7
11	Chronic lower respiratory diseases (J40-J47)	143	1.4	2.5
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	92	0.9	1.6
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	82	0.8	1.4
14	Pregnancy, childbirth and the puerperium (O00-O99)	56	0.6	1.0
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	0.5	0.9
...	All other causes (residual)	1,963	19.4	34.2
Non-Hispanic, white, female, 45-49 years				
...	All causes	16,843	100.0	255.7
1	Malignant neoplasms (C00-C97)	4,858	28.8	73.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,638	15.7	40.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,246	13.3	34.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	882	5.2	13.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	766	4.5	11.6
6	Chronic lower respiratory diseases (J40-J47)	475	2.8	7.2
7	Cerebrovascular diseases (I60-I69)	458	2.7	7.0
8	Diabetes mellitus (E10-E14)	418	2.5	6.3
9	Influenza and pneumonia (J09-J18)	251	1.5	3.8
10	Septicemia (A40-A41)	240	1.4	3.6
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	152	0.9	2.3
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	147	0.9	2.2
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	0.7	1.8
14	Viral hepatitis (B15-B19)	87	0.5	1.3
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	77	0.5	1.2
...	All other causes (residual)	3,030	18.0	46.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 50-54 years				
...	All causes	28,820	100.0	391.8
1	Malignant neoplasms (C00-C97)	9,268	32.2	126.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,222	14.6	57.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,051	10.6	41.5
4	Chronic lower respiratory diseases (J40-J47)	1,433	5.0	19.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,402	4.9	19.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,094	3.8	14.9
7	Diabetes mellitus (E10-E14)	810	2.8	11.0
8	Cerebrovascular diseases (I60-I69)	804	2.8	10.9
9	Septicemia (A40-A41)	460	1.6	6.3
10	Influenza and pneumonia (J09-J18)	401	1.4	5.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	267	0.9	3.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	186	0.6	2.5
13	Viral hepatitis (B15-B19)	179	0.6	2.4
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	152	0.5	2.1
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	0.5	2.0
...	All other causes (residual)	4,941	17.1	67.2
Non-Hispanic, white, female, 55-59 years				
...	All causes	44,589	100.0	564.7
1	Malignant neoplasms (C00-C97)	16,259	36.5	205.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,828	15.3	86.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,957	6.6	37.5
4	Chronic lower respiratory diseases (J40-J47)	2,930	6.6	37.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,672	3.7	21.2
6	Cerebrovascular diseases (I60-I69)	1,348	3.0	17.1
7	Diabetes mellitus (E10-E14)	1,252	2.8	15.9
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,024	2.3	13.0
9	Septicemia (A40-A41)	738	1.7	9.3
10	Influenza and pneumonia (J09-J18)	640	1.4	8.1
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	550	1.2	7.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	317	0.7	4.0
13	Viral hepatitis (B15-B19)	277	0.6	3.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	263	0.6	3.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	201	0.5	2.5
...	All other causes (residual)	7,333	16.4	92.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 60-64 years				
...	All causes	57,591	100.0	783.6
1	Malignant neoplasms (C00-C97)	21,872	38.0	297.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,431	16.4	128.3
3	Chronic lower respiratory diseases (J40-J47)	4,386	7.6	59.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,180	3.8	29.7
5	Diabetes mellitus (E10-E14)	1,908	3.3	26.0
6	Cerebrovascular diseases (I60-I69)	1,786	3.1	24.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	1,506	2.6	20.5
8	Septicemia (A40-A41)	1,044	1.8	14.2
9	Influenza and pneumonia (J09-J18)	928	1.6	12.6
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	789	1.4	10.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	778	1.4	10.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	474	0.8	6.4
13	Alzheimer's disease (G30)	375	0.7	5.1
14	Viral hepatitis (B15-B19)	320	0.6	4.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	313	0.5	4.3
...	All other causes (residual)	9,501	16.5	129.3
Non-Hispanic, white, female, 65-69 years				
...	All causes	76,673	100.0	1,151.5
1	Malignant neoplasms (C00-C97)	28,226	36.8	423.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	13,031	17.0	195.7
3	Chronic lower respiratory diseases (J40-J47)	6,935	9.0	104.2
4	Cerebrovascular diseases (I60-I69)	2,951	3.8	44.3
5	Diabetes mellitus (E10-E14)	2,577	3.4	38.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,004	2.6	30.1
7	Septicemia (A40-A41)	1,392	1.8	20.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,301	1.7	19.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,250	1.6	18.8
10	Influenza and pneumonia (J09-J18)	1,165	1.5	17.5
11	Alzheimer's disease (G30)	912	1.2	13.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	694	0.9	10.4
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	520	0.7	7.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	441	0.6	6.6
15	Pneumonitis due to solids and liquids (J69)	380	0.5	5.7
...	All other causes (residual)	12,894	16.8	193.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 70-74 years				
...	All causes	91,467	100.0	1,883.3
1	Malignant neoplasms (C00-C97)	29,908	32.7	615.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	16,435	18.0	338.4
3	Chronic lower respiratory diseases (J40-J47)	9,712	10.6	200.0
4	Cerebrovascular diseases (I60-I69)	4,153	4.5	85.5
5	Diabetes mellitus (E10-E14)	2,752	3.0	56.7
6	Alzheimer's disease (G30)	2,326	2.5	47.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,931	2.1	39.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,665	1.8	34.3
9	Septicemia (A40-A41)	1,608	1.8	33.1
10	Influenza and pneumonia (J09-J18)	1,503	1.6	30.9
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1,131	1.2	23.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	876	1.0	18.0
13	Parkinson's disease (G20-G21)	756	0.8	15.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	604	0.7	12.4
15	Pneumonitis due to solids and liquids (J69)	484	0.5	10.0
...	All other causes (residual)	15,623	17.1	321.7
Non-Hispanic, white, female, 75-79 years				
...	All causes	109,005	100.0	3,050.0
1	Malignant neoplasms (C00-C97)	29,865	27.4	835.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	21,005	19.3	587.7
3	Chronic lower respiratory diseases (J40-J47)	11,243	10.3	314.6
4	Cerebrovascular diseases (I60-I69)	6,084	5.6	170.2
5	Alzheimer's disease (G30)	5,129	4.7	143.5
6	Diabetes mellitus (E10-E14)	2,818	2.6	78.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,494	2.3	69.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,060	1.9	57.6
9	Influenza and pneumonia (J09-J18)	1,952	1.8	54.6
10	Septicemia (A40-A41)	1,867	1.7	52.2
11	Parkinson's disease (G20-G21)	1,436	1.3	40.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,192	1.1	33.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	876	0.8	24.5
14	Pneumonitis due to solids and liquids (J69)	730	0.7	20.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	727	0.7	20.3
...	All other causes (residual)	19,527	17.9	546.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 80-84 years				
...	All causes	140,127	100.0	5,255.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	30,522	21.8	1,144.8
2	Malignant neoplasms (C00-C97)	28,359	20.2	1,063.6
3	Chronic lower respiratory diseases (J40-J47)	12,021	8.6	450.9
4	Alzheimer's disease (G30)	10,480	7.5	393.1
5	Cerebrovascular diseases (I60-I69)	9,674	6.9	362.8
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,564	2.5	133.7
7	Diabetes mellitus (E10-E14)	3,251	2.3	121.9
8	Influenza and pneumonia (J09-J18)	2,758	2.0	103.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,558	1.8	95.9
10	Parkinson's disease (G20-G21)	2,294	1.6	86.0
11	Septicemia (A40-A41)	2,231	1.6	83.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,738	1.2	65.2
13	Pneumonitis due to solids and liquids (J69)	1,030	0.7	38.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	958	0.7	35.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	661	0.5	24.8
...	All other causes (residual)	28,028	20.0	1,051.2
Non-Hispanic, white, female, 85-89 years				
...	All causes	180,921	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	45,929	25.4	---
2	Malignant neoplasms (C00-C97)	25,181	13.9	---
3	Alzheimer's disease (G30)	17,820	9.8	---
4	Cerebrovascular diseases (I60-I69)	14,242	7.9	---
5	Chronic lower respiratory diseases (J40-J47)	11,540	6.4	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,962	2.7	---
7	Influenza and pneumonia (J09-J18)	3,945	2.2	---
8	Diabetes mellitus (E10-E14)	3,364	1.9	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,162	1.7	---
10	Parkinson's disease (G20-G21)	2,758	1.5	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,598	1.4	---
12	Septicemia (A40-A41)	2,458	1.4	---
13	Pneumonitis due to solids and liquids (J69)	1,452	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1,136	0.6	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	674	0.4	---
...	All other causes (residual)	39,700	21.9	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 90-94 years				
...	All causes	169,405	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49,185	29.0	---
2	Alzheimer's disease (G30)	18,870	11.1	---
3	Malignant neoplasms (C00-C97)	14,911	8.8	---
4	Cerebrovascular diseases (I60-I69)	13,807	8.2	---
5	Chronic lower respiratory diseases (J40-J47)	7,898	4.7	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,783	2.8	---
7	Influenza and pneumonia (J09-J18)	4,008	2.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,879	1.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,796	1.7	---
10	Diabetes mellitus (E10-E14)	2,535	1.5	---
11	Septicemia (A40-A41)	1,963	1.2	---
12	Parkinson's disease (G20-G21)	1,793	1.1	---
13	Pneumonitis due to solids and liquids (J69)	1,553	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	873	0.5	---
15	Nutritional deficiencies (E40-E64)	669	0.4	---
...	All other causes (residual)	40,882	24.1	---
Non-Hispanic, white, female, 95-99 years				
...	All causes	81,542	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	25,756	31.6	---
2	Alzheimer's disease (G30)	9,661	11.8	---
3	Cerebrovascular diseases (I60-I69)	6,543	8.0	---
4	Malignant neoplasms (C00-C97)	4,625	5.7	---
5	Chronic lower respiratory diseases (J40-J47)	2,869	3.5	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,246	2.8	---
7	Influenza and pneumonia (J09-J18)	2,107	2.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,652	2.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,306	1.6	---
10	Diabetes mellitus (E10-E14)	896	1.1	---
11	Pneumonitis due to solids and liquids (J69)	776	1.0	---
12	Septicemia (A40-A41)	769	0.9	---
13	Parkinson's disease (G20-G21)	486	0.6	---
14	Nutritional deficiencies (E40-E64)	440	0.5	---
15	Atherosclerosis (I70)	402	0.5	---
...	All other causes (residual)	21,008	25.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 100 years and over				
...	All causes	18,964	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,269	33.1	---
2	Alzheimer's disease (G30)	2,122	11.2	---
3	Cerebrovascular diseases (I60-I69)	1,347	7.1	---
4	Malignant neoplasms (C00-C97)	710	3.7	---
5	Influenza and pneumonia (J09-J18)	581	3.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	542	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	527	2.8	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	475	2.5	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	271	1.4	---
10	Pneumonitis due to solids and liquids (J69)	175	0.9	---
11	Diabetes mellitus (E10-E14)	164	0.9	---
12	Nutritional deficiencies (E40-E64)	157	0.8	---
13	Septicemia (A40-A41)	154	0.8	---
14	Atherosclerosis (I70)	129	0.7	---
15	Parkinson's disease (G20-G21)	66	0.3	---
...	All other causes (residual)	5,275	27.8	---
Non-Hispanic, black, both sexes, all ages				
...	All causes	326,810	100.0	775.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,608	23.4	181.8
2	Malignant neoplasms (C00-C97)	69,725	21.3	165.5
3	Cerebrovascular diseases (I60-I69)	18,189	5.6	43.2
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18,089	5.5	42.9
5	Diabetes mellitus (E10-E14)	14,053	4.3	33.3
6	Chronic lower respiratory diseases (J40-J47)	10,838	3.3	25.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,860	3.0	23.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,200	2.8	21.8
9	Alzheimer's disease (G30)	8,577	2.6	20.4
10	Septicemia (A40-A41)	6,684	2.0	15.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,924	1.8	14.1
12	Influenza and pneumonia (J09-J18)	5,468	1.7	13.0
13	Certain conditions originating in the perinatal period (P00-P96)	3,775	1.2	9.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3,152	1.0	7.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3,149	1.0	7.5
...	All other causes (residual)	63,519	19.4	150.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	1,016	100.0	41.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	277	27.3	11.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	141	13.9	5.8
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	111	10.9	4.5
4	Malignant neoplasms (C00-C97)	58	5.7	2.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	3.3	1.4
5	Influenza and pneumonia (J09-J18)	34	3.3	1.4
7	Certain conditions originating in the perinatal period (P00-P96)	26	2.6	1.1
8	Chronic lower respiratory diseases (J40-J47)	19	1.9	*
9	Cerebrovascular diseases (I60-I69)	15	1.5	*
10	Septicemia (A40-A41)	14	1.4	*
11	Anemias (D50-D64)	13	1.3	*
12	Acute bronchitis and bronchiolitis (J20-J21)	8	0.8	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
14	Meningitis (G00,G03)	6	0.6	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.5	*
...	All other causes (residual)	248	24.4	10.1
Non-Hispanic, black, both sexes, 5-9 years				
...	All causes	610	100.0	19.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	36.1	7.0
2	Malignant neoplasms (C00-C97)	79	13.0	2.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	54	8.9	1.7
4	Chronic lower respiratory diseases (J40-J47)	43	7.0	1.4
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	38	6.2	1.2
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	3.3	0.6
7	Septicemia (A40-A41)	9	1.5	*
7	Influenza and pneumonia (J09-J18)	9	1.5	*
9	Anemias (D50-D64)	7	1.1	*
9	Cerebrovascular diseases (I60-I69)	7	1.1	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	4	0.7	*
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.7	*
14	Meningitis (G00,G03)	3	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.5	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.5	*
...	All other causes (residual)	102	16.7	3.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 10-14 years				
...	All causes	623	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	168	27.0	5.5
2	Malignant neoplasms (C00-C97)	65	10.4	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	10.3	2.1
4	Chronic lower respiratory diseases (J40-J47)	45	7.2	1.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	6.4	1.3
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	31	5.0	1.0
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	24	3.9	0.8
8	Cerebrovascular diseases (I60-I69)	13	2.1	*
9	Influenza and pneumonia (J09-J18)	12	1.9	*
10	Diabetes mellitus (E10-E14)	8	1.3	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.0	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.6	*
13	Septicemia (A40-A41)	3	0.5	*
13	Anemias (D50-D64)	3	0.5	*
13	Diseases of appendix (K35-K38)	3	0.5	*
...	All other causes (residual)	134	21.5	4.4
Non-Hispanic, black, both sexes, 15-19 years				
...	All causes	2,570	100.0	80.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,122	43.7	35.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	619	24.1	19.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	197	7.7	6.1
4	Malignant neoplasms (C00-C97)	90	3.5	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	3.4	2.7
6	Chronic lower respiratory diseases (J40-J47)	40	1.6	1.2
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	39	1.5	1.2
8	Anemias (D50-D64)	28	1.1	0.9
9	Legal intervention (Y35,Y89.0)	14	0.5	*
10	Septicemia (A40-A41)	13	0.5	*
10	Cerebrovascular diseases (I60-I69)	13	0.5	*
10	Influenza and pneumonia (J09-J18)	13	0.5	*
13	Diabetes mellitus (E10-E14)	10	0.4	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.3	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	6	0.2	*
...	All other causes (residual)	271	10.5	8.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 20-24 years				
...	All causes	5,133	100.0	147.2
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,159	42.1	61.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,250	24.4	35.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	459	8.9	13.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	212	4.1	6.1
5	Malignant neoplasms (C00-C97)	140	2.7	4.0
6	Pregnancy, childbirth and the puerperium (O00-O99)	61	1.2	1.7
7	Diabetes mellitus (E10-E14)	54	1.1	1.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	49	1.0	1.4
9	Chronic lower respiratory diseases (J40-J47)	38	0.7	1.1
10	Influenza and pneumonia (J09-J18)	37	0.7	1.1
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	33	0.6	0.9
12	Anemias (D50-D64)	31	0.6	0.9
13	Septicemia (A40-A41)	30	0.6	0.9
14	Cerebrovascular diseases (I60-I69)	20	0.4	0.6
15	Legal intervention (Y35,Y89.0)	18	0.4	*
...	All other causes (residual)	542	10.6	15.5
Non-Hispanic, black, both sexes, 25-29 years				
...	All causes	5,683	100.0	167.6
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,822	32.1	53.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,428	25.1	42.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	412	7.2	12.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	352	6.2	10.4
5	Malignant neoplasms (C00-C97)	238	4.2	7.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	148	2.6	4.4
7	Diabetes mellitus (E10-E14)	102	1.8	3.0
8	Pregnancy, childbirth and the puerperium (O00-O99)	86	1.5	2.5
9	Anemias (D50-D64)	68	1.2	2.0
10	Chronic lower respiratory diseases (J40-J47)	62	1.1	1.8
11	Cerebrovascular diseases (I60-I69)	50	0.9	1.5
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	0.8	1.4
13	Septicemia (A40-A41)	44	0.8	1.3
14	Influenza and pneumonia (J09-J18)	41	0.7	1.2
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	0.7	1.1
...	All other causes (residual)	745	13.1	22.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 30-34 years				
...	All causes	5,829	100.0	201.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,323	22.7	45.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,279	21.9	44.2
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	665	11.4	23.0
4	Malignant neoplasms (C00-C97)	402	6.9	13.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	317	5.4	10.9
6	Human immunodeficiency virus (HIV) disease (B20-B24)	190	3.3	6.6
7	Diabetes mellitus (E10-E14)	147	2.5	5.1
8	Cerebrovascular diseases (I60-I69)	89	1.5	3.1
9	Chronic lower respiratory diseases (J40-J47)	72	1.2	2.5
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	70	1.2	2.4
11	Pregnancy, childbirth and the puerperium (O00-O99)	68	1.2	2.3
12	Influenza and pneumonia (J09-J18)	66	1.1	2.3
13	Anemias (D50-D64)	63	1.1	2.2
14	Septicemia (A40-A41)	54	0.9	1.9
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	42	0.7	1.5
...	All other causes (residual)	982	16.8	33.9
Non-Hispanic, black, both sexes, 35-39 years				
...	All causes	6,888	100.0	250.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,365	19.8	49.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,133	16.4	41.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	978	14.2	35.6
4	Malignant neoplasms (C00-C97)	712	10.3	25.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	242	3.5	8.8
6	Human immunodeficiency virus (HIV) disease (B20-B24)	235	3.4	8.6
7	Diabetes mellitus (E10-E14)	221	3.2	8.0
8	Cerebrovascular diseases (I60-I69)	173	2.5	6.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	1.8	4.6
10	Septicemia (A40-A41)	86	1.2	3.1
11	Anemias (D50-D64)	79	1.1	2.9
12	Pregnancy, childbirth and the puerperium (O00-O99)	73	1.1	2.7
13	Chronic liver disease and cirrhosis (K70,K73-K74)	67	1.0	2.4
14	Chronic lower respiratory diseases (J40-J47)	64	0.9	2.3
15	Influenza and pneumonia (J09-J18)	63	0.9	2.3
...	All other causes (residual)	1,271	18.5	46.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 40-44 years				
...	All causes	8,307	100.0	325.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,731	20.8	67.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,204	14.5	47.2
3	Malignant neoplasms (C00-C97)	1,162	14.0	45.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	665	8.0	26.1
5	Diabetes mellitus (E10-E14)	360	4.3	14.1
6	Cerebrovascular diseases (I60-I69)	329	4.0	12.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	306	3.7	12.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	210	2.5	8.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	174	2.1	6.8
10	Septicemia (A40-A41)	157	1.9	6.2
11	Chronic liver disease and cirrhosis (K70,K73-K74)	138	1.7	5.4
12	Influenza and pneumonia (J09-J18)	121	1.5	4.7
13	Chronic lower respiratory diseases (J40-J47)	112	1.3	4.4
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	105	1.3	4.1
15	Anemias (D50-D64)	59	0.7	2.3
...	All other causes (residual)	1,474	17.7	57.8
Non-Hispanic, black, both sexes, 45-49 years				
...	All causes	12,218	100.0	461.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,936	24.0	110.9
2	Malignant neoplasms (C00-C97)	2,264	18.5	85.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,424	11.7	53.8
4	Diabetes mellitus (E10-E14)	555	4.5	21.0
5	Cerebrovascular diseases (I60-I69)	544	4.5	20.6
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	463	3.8	17.5
7	Human immunodeficiency virus (HIV) disease (B20-B24)	365	3.0	13.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	297	2.4	11.2
9	Septicemia (A40-A41)	218	1.8	8.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	218	1.8	8.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	215	1.8	8.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	171	1.4	6.5
13	Influenza and pneumonia (J09-J18)	169	1.4	6.4
14	Chronic lower respiratory diseases (J40-J47)	158	1.3	6.0
15	Aortic aneurysm and dissection (I71)	81	0.7	3.1
...	All other causes (residual)	2,140	17.5	80.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 50-54 years				
...	All causes	19,420	100.0	711.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,761	24.5	174.5
2	Malignant neoplasms (C00-C97)	4,559	23.5	167.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,741	9.0	63.8
4	Diabetes mellitus (E10-E14)	946	4.9	34.7
5	Cerebrovascular diseases (I60-I69)	903	4.6	33.1
6	Human immunodeficiency virus (HIV) disease (B20-B24)	502	2.6	18.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	425	2.2	15.6
8	Chronic lower respiratory diseases (J40-J47)	424	2.2	15.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	424	2.2	15.5
10	Septicemia (A40-A41)	373	1.9	13.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	344	1.8	12.6
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	343	1.8	12.6
13	Influenza and pneumonia (J09-J18)	250	1.3	9.2
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	168	0.9	6.2
15	Aortic aneurysm and dissection (I71)	85	0.4	3.1
...	All other causes (residual)	3,172	16.3	116.3
Non-Hispanic, black, both sexes, 55-59 years				
...	All causes	28,391	100.0	1,079.9
1	Malignant neoplasms (C00-C97)	7,912	27.9	300.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,003	24.7	266.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,792	6.3	68.2
4	Diabetes mellitus (E10-E14)	1,367	4.8	52.0
5	Cerebrovascular diseases (I60-I69)	1,271	4.5	48.3
6	Chronic lower respiratory diseases (J40-J47)	873	3.1	33.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	690	2.4	26.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	660	2.3	25.1
9	Septicemia (A40-A41)	609	2.1	23.2
10	Human immunodeficiency virus (HIV) disease (B20-B24)	512	1.8	19.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	476	1.7	18.1
12	Influenza and pneumonia (J09-J18)	373	1.3	14.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	288	1.0	11.0
14	Viral hepatitis (B15-B19)	209	0.7	7.9
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	146	0.5	5.6
...	All other causes (residual)	4,210	14.8	160.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 60-64 years				
...	All causes	34,430	100.0	1,592.5
1	Malignant neoplasms (C00-C97)	10,298	29.9	476.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,531	24.8	394.6
3	Diabetes mellitus (E10-E14)	1,694	4.9	78.4
4	Cerebrovascular diseases (I60-I69)	1,685	4.9	77.9
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,437	4.2	66.5
6	Chronic lower respiratory diseases (J40-J47)	1,206	3.5	55.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	998	2.9	46.2
8	Septicemia (A40-A41)	769	2.2	35.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	641	1.9	29.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	553	1.6	25.6
11	Influenza and pneumonia (J09-J18)	510	1.5	23.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	408	1.2	18.9
13	Viral hepatitis (B15-B19)	272	0.8	12.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	182	0.5	8.4
15	Pneumonitis due to solids and liquids (J69)	148	0.4	6.8
...	All other causes (residual)	5,098	14.8	235.8
Non-Hispanic, black, both sexes, 65-69 years				
...	All causes	35,847	100.0	2,136.1
1	Malignant neoplasms (C00-C97)	10,844	30.3	646.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,042	25.2	538.8
3	Cerebrovascular diseases (I60-I69)	1,926	5.4	114.8
4	Diabetes mellitus (E10-E14)	1,889	5.3	112.6
5	Chronic lower respiratory diseases (J40-J47)	1,496	4.2	89.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,125	3.1	67.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	963	2.7	57.4
8	Septicemia (A40-A41)	818	2.3	48.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	647	1.8	38.6
10	Influenza and pneumonia (J09-J18)	589	1.6	35.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	414	1.2	24.7
12	Human immunodeficiency virus (HIV) disease (B20-B24)	233	0.6	13.9
13	Alzheimer's disease (G30)	218	0.6	13.0
14	Viral hepatitis (B15-B19)	182	0.5	10.8
15	Pneumonitis due to solids and liquids (J69)	174	0.5	10.4
...	All other causes (residual)	5,287	14.7	315.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 70-74 years				
...	All causes	31,781	100.0	2,966.1
1	Malignant neoplasms (C00-C97)	8,882	27.9	828.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,968	25.1	743.6
3	Cerebrovascular diseases (I60-I69)	1,965	6.2	183.4
4	Diabetes mellitus (E10-E14)	1,720	5.4	160.5
5	Chronic lower respiratory diseases (J40-J47)	1,487	4.7	138.8
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,092	3.4	101.9
7	Septicemia (A40-A41)	773	2.4	72.1
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	645	2.0	60.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	602	1.9	56.2
10	Influenza and pneumonia (J09-J18)	538	1.7	50.2
11	Alzheimer's disease (G30)	472	1.5	44.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	230	0.7	21.5
13	Pneumonitis due to solids and liquids (J69)	195	0.6	18.2
14	Parkinson's disease (G20-G21)	174	0.5	16.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	137	0.4	12.8
...	All other causes (residual)	4,901	15.4	457.4
Non-Hispanic, black, both sexes, 75-79 years				
...	All causes	31,486	100.0	4,256.9
1	Malignant neoplasms (C00-C97)	7,921	25.2	1,070.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,839	24.9	1,059.8
3	Cerebrovascular diseases (I60-I69)	2,194	7.0	296.6
4	Diabetes mellitus (E10-E14)	1,566	5.0	211.7
5	Chronic lower respiratory diseases (J40-J47)	1,469	4.7	198.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,144	3.6	154.7
7	Alzheimer's disease (G30)	1,020	3.2	137.9
8	Septicemia (A40-A41)	799	2.5	108.0
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	669	2.1	90.4
10	Influenza and pneumonia (J09-J18)	620	2.0	83.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	525	1.7	71.0
12	Parkinson's disease (G20-G21)	267	0.8	36.1
13	Pneumonitis due to solids and liquids (J69)	221	0.7	29.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	161	0.5	21.8
15	Chronic liver disease and cirrhosis (K70,K73-K74)	123	0.4	16.6
...	All other causes (residual)	4,948	15.7	669.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 80-84 years				
...	All causes	31,588	100.0	6,427.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,228	26.0	1,674.2
2	Malignant neoplasms (C00-C97)	6,486	20.5	1,319.7
3	Cerebrovascular diseases (I60-I69)	2,306	7.3	469.2
4	Alzheimer's disease (G30)	1,682	5.3	342.2
5	Diabetes mellitus (E10-E14)	1,408	4.5	286.5
6	Chronic lower respiratory diseases (J40-J47)	1,346	4.3	273.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,169	3.7	237.9
8	Septicemia (A40-A41)	732	2.3	148.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	662	2.1	134.7
10	Influenza and pneumonia (J09-J18)	620	2.0	126.2
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	471	1.5	95.8
12	Parkinson's disease (G20-G21)	312	1.0	63.5
13	Pneumonitis due to solids and liquids (J69)	270	0.9	54.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	169	0.5	34.4
15	Nutritional deficiencies (E40-E64)	114	0.4	23.2
...	All other causes (residual)	5,613	17.8	1,142.1
Non-Hispanic, black, both sexes, 85-89 years				
...	All causes	27,362	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,240	26.5	---
2	Malignant neoplasms (C00-C97)	4,533	16.6	---
3	Cerebrovascular diseases (I60-I69)	2,114	7.7	---
4	Alzheimer's disease (G30)	1,963	7.2	---
5	Diabetes mellitus (E10-E14)	1,102	4.0	---
6	Chronic lower respiratory diseases (J40-J47)	1,024	3.7	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	992	3.6	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	635	2.3	---
9	Septicemia (A40-A41)	583	2.1	---
10	Influenza and pneumonia (J09-J18)	578	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	413	1.5	---
12	Parkinson's disease (G20-G21)	257	0.9	---
13	Pneumonitis due to solids and liquids (J69)	247	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	142	0.5	---
15	Nutritional deficiencies (E40-E64)	113	0.4	---
...	All other causes (residual)	5,426	19.8	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 90-94 years				
...	All causes	19,708	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,433	27.6	---
2	Malignant neoplasms (C00-C97)	2,308	11.7	---
3	Alzheimer's disease (G30)	1,897	9.6	---
4	Cerebrovascular diseases (I60-I69)	1,608	8.2	---
5	Diabetes mellitus (E10-E14)	637	3.2	---
6	Chronic lower respiratory diseases (J40-J47)	608	3.1	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	595	3.0	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	508	2.6	---
9	Influenza and pneumonia (J09-J18)	468	2.4	---
10	Septicemia (A40-A41)	347	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	289	1.5	---
12	Pneumonitis due to solids and liquids (J69)	207	1.1	---
13	Parkinson's disease (G20-G21)	137	0.7	---
14	Nutritional deficiencies (E40-E64)	120	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	102	0.5	---
...	All other causes (residual)	4,444	22.5	---
Non-Hispanic, black, both sexes, 95-99 years				
...	All causes	8,495	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,496	29.4	---
2	Alzheimer's disease (G30)	925	10.9	---
3	Cerebrovascular diseases (I60-I69)	699	8.2	---
4	Malignant neoplasms (C00-C97)	657	7.7	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	259	3.0	---
6	Diabetes mellitus (E10-E14)	220	2.6	---
7	Influenza and pneumonia (J09-J18)	219	2.6	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	199	2.3	---
9	Chronic lower respiratory diseases (J40-J47)	189	2.2	---
10	Septicemia (A40-A41)	145	1.7	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	126	1.5	---
12	Pneumonitis due to solids and liquids (J69)	77	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.6	---
14	Nutritional deficiencies (E40-E64)	49	0.6	---
15	Parkinson's disease (G20-G21)	48	0.6	---
...	All other causes (residual)	2,137	25.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 100 years and over				
...	All causes	2,551	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	775	30.4	---
2	Alzheimer's disease (G30)	285	11.2	---
3	Cerebrovascular diseases (I60-I69)	227	8.9	---
4	Malignant neoplasms (C00-C97)	101	4.0	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	78	3.1	---
6	Influenza and pneumonia (J09-J18)	75	2.9	---
7	Chronic lower respiratory diseases (J40-J47)	51	2.0	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	51	2.0	---
9	Diabetes mellitus (E10-E14)	44	1.7	---
10	Septicemia (A40-A41)	39	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35	1.4	---
12	Pneumonitis due to solids and liquids (J69)	21	0.8	---
13	Nutritional deficiencies (E40-E64)	15	0.6	---
14	Atherosclerosis (I70)	14	0.5	---
15	Anemias (D50-D64)	12	0.5	---
...	All other causes (residual)	728	28.5	---
Non-Hispanic, black, male, all ages				
...	All causes	168,750	100.0	836.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	40,044	23.7	198.4
2	Malignant neoplasms (C00-C97)	35,215	20.9	174.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,453	7.4	61.7
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,532	5.1	42.3
5	Cerebrovascular diseases (I60-I69)	8,115	4.8	40.2
6	Diabetes mellitus (E10-E14)	6,976	4.1	34.6
7	Chronic lower respiratory diseases (J40-J47)	5,519	3.3	27.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,348	2.6	21.5
9	Septicemia (A40-A41)	3,074	1.8	15.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,805	1.7	13.9
11	Influenza and pneumonia (J09-J18)	2,705	1.6	13.4
12	Alzheimer's disease (G30)	2,451	1.5	12.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,137	1.3	10.6
14	Certain conditions originating in the perinatal period (P00-P96)	2,101	1.2	10.4
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2,072	1.2	10.3
...	All other causes (residual)	30,203	17.9	149.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 1-4 years				
...	All causes	575	100.0	46.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	167	29.0	13.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	12.7	5.9
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	54	9.4	4.3
4	Malignant neoplasms (C00-C97)	32	5.6	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.8	1.8
6	Influenza and pneumonia (J09-J18)	18	3.1	*
7	Certain conditions originating in the perinatal period (P00-P96)	17	3.0	*
8	Chronic lower respiratory diseases (J40-J47)	14	2.4	*
9	Cerebrovascular diseases (I60-I69)	7	1.2	*
9	Acute bronchitis and bronchiolitis (J20-J21)	7	1.2	*
11	Septicemia (A40-A41)	6	1.0	*
12	Anemias (D50-D64)	5	0.9	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.7	*
13	Meningitis (G00,G03)	4	0.7	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	2	0.3	*
...	All other causes (residual)	141	24.5	11.3
Non-Hispanic, black, male, 5-9 years				
...	All causes	333	100.0	21.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	38.7	8.1
2	Malignant neoplasms (C00-C97)	37	11.1	2.3
3	Chronic lower respiratory diseases (J40-J47)	31	9.3	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	29	8.7	1.8
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	17	5.1	*
6	Influenza and pneumonia (J09-J18)	7	2.1	*
7	Anemias (D50-D64)	5	1.5	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	5	1.5	*
7	Cerebrovascular diseases (I60-I69)	5	1.5	*
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.9	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.6	*
13	Septicemia (A40-A41)	1	0.3	*
13	Meningitis (G00,G03)	1	0.3	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	53	15.9	3.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 10-14 years				
...	All causes	368	100.0	23.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	104	28.3	6.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	42	11.4	2.7
3	Malignant neoplasms (C00-C97)	29	7.9	1.9
4	Chronic lower respiratory diseases (J40-J47)	27	7.3	1.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	6.8	1.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	4.9	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	16	4.3	*
8	Cerebrovascular diseases (I60-I69)	7	1.9	*
8	Influenza and pneumonia (J09-J18)	7	1.9	*
10	Diabetes mellitus (E10-E14)	5	1.4	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.1	*
12	Anemias (D50-D64)	3	0.8	*
13	Septicemia (A40-A41)	2	0.5	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
14	Meningitis (G00,G03)	1	0.3	*
14	Aortic aneurysm and dissection (I71)	1	0.3	*
14	Diseases of appendix (K35-K38)	1	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	73	19.8	4.7
Non-Hispanic, black, male, 15-19 years				
...	All causes	1,952	100.0	119.7
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	991	50.8	60.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	441	22.6	27.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	7.4	8.9
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	3.0	3.6
5	Malignant neoplasms (C00-C97)	49	2.5	3.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	27	1.4	1.7
7	Chronic lower respiratory diseases (J40-J47)	22	1.1	1.3
8	Anemias (D50-D64)	15	0.8	*
9	Legal intervention (Y35,Y89.0)	14	0.7	*
10	Septicemia (A40-A41)	7	0.4	*
10	Cerebrovascular diseases (I60-I69)	7	0.4	*
12	Influenza and pneumonia (J09-J18)	6	0.3	*
13	Diabetes mellitus (E10-E14)	5	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.2	*
...	All other causes (residual)	155	7.9	9.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 20-24 years				
...	All causes	3,957	100.0	223.1
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,964	49.6	110.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	898	22.7	50.6
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	379	9.6	21.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	136	3.4	7.7
5	Malignant neoplasms (C00-C97)	88	2.2	5.0
6	Diabetes mellitus (E10-E14)	28	0.7	1.6
7	Human immunodeficiency virus (HIV) disease (B20-B24)	27	0.7	1.5
8	Chronic lower respiratory diseases (J40-J47)	26	0.7	1.5
9	Anemias (D50-D64)	20	0.5	1.1
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	19	0.5	*
11	Influenza and pneumonia (J09-J18)	18	0.5	*
12	Legal intervention (Y35,Y89.0)	17	0.4	*
13	Septicemia (A40-A41)	14	0.4	*
14	Cerebrovascular diseases (I60-I69)	10	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	0.2	*
...	All other causes (residual)	305	7.7	17.2
Non-Hispanic, black, male, 25-29 years				
...	All causes	4,139	100.0	245.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,627	39.3	96.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,059	25.6	62.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	348	8.4	20.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	216	5.2	12.8
5	Malignant neoplasms (C00-C97)	113	2.7	6.7
6	Human immunodeficiency virus (HIV) disease (B20-B24)	111	2.7	6.6
7	Diabetes mellitus (E10-E14)	61	1.5	3.6
8	Chronic lower respiratory diseases (J40-J47)	37	0.9	2.2
9	Anemias (D50-D64)	32	0.8	1.9
10	Cerebrovascular diseases (I60-I69)	29	0.7	1.7
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	29	0.7	1.7
12	Septicemia (A40-A41)	24	0.6	1.4
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	0.6	1.4
14	Legal intervention (Y35,Y89.0)	21	0.5	1.2
15	Influenza and pneumonia (J09-J18)	19	0.5	*
...	All other causes (residual)	390	9.4	23.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 30-34 years				
...	All causes	3,967	100.0	283.9
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,135	28.6	81.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	988	24.9	70.7
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	417	10.5	29.8
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	255	6.4	18.2
5	Malignant neoplasms (C00-C97)	161	4.1	11.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	132	3.3	9.4
7	Diabetes mellitus (E10-E14)	77	1.9	5.5
8	Cerebrovascular diseases (I60-I69)	55	1.4	3.9
9	Chronic lower respiratory diseases (J40-J47)	45	1.1	3.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	39	1.0	2.8
11	Septicemia (A40-A41)	30	0.8	2.1
11	Influenza and pneumonia (J09-J18)	30	0.8	2.1
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	23	0.6	1.6
14	Anemias (D50-D64)	22	0.6	1.6
14	Legal intervention (Y35,Y89.0)	22	0.6	1.6
...	All other causes (residual)	536	13.5	38.4
Non-Hispanic, black, male, 35-39 years				
...	All causes	4,302	100.0	330.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	956	22.2	73.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	853	19.8	65.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	723	16.8	55.5
4	Malignant neoplasms (C00-C97)	269	6.3	20.6
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	190	4.4	14.6
6	Human immunodeficiency virus (HIV) disease (B20-B24)	143	3.3	11.0
7	Diabetes mellitus (E10-E14)	113	2.6	8.7
8	Cerebrovascular diseases (I60-I69)	85	2.0	6.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	1.6	5.3
10	Septicemia (A40-A41)	42	1.0	3.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	42	1.0	3.2
12	Chronic lower respiratory diseases (J40-J47)	38	0.9	2.9
13	Anemias (D50-D64)	34	0.8	2.6
14	Aortic aneurysm and dissection (I71)	30	0.7	2.3
14	Influenza and pneumonia (J09-J18)	30	0.7	2.3
...	All other causes (residual)	685	15.9	52.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 40-44 years				
...	All causes	5,014	100.0	418.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,125	22.4	93.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	883	17.6	73.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	592	11.8	49.4
4	Malignant neoplasms (C00-C97)	440	8.8	36.7
5	Diabetes mellitus (E10-E14)	209	4.2	17.4
6	Cerebrovascular diseases (I60-I69)	196	3.9	16.3
7	Human immunodeficiency virus (HIV) disease (B20-B24)	164	3.3	13.7
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	162	3.2	13.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	88	1.8	7.3
10	Septicemia (A40-A41)	78	1.6	6.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	74	1.5	6.2
12	Influenza and pneumonia (J09-J18)	66	1.3	5.5
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	61	1.2	5.1
14	Chronic lower respiratory diseases (J40-J47)	57	1.1	4.8
15	Aortic aneurysm and dissection (I71)	44	0.9	3.7
...	All other causes (residual)	775	15.5	64.6
Non-Hispanic, black, male, 45-49 years				
...	All causes	7,028	100.0	565.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,899	27.0	152.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	977	13.9	78.6
3	Malignant neoplasms (C00-C97)	931	13.2	74.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	378	5.4	30.4
5	Diabetes mellitus (E10-E14)	327	4.7	26.3
6	Cerebrovascular diseases (I60-I69)	295	4.2	23.7
7	Human immunodeficiency virus (HIV) disease (B20-B24)	209	3.0	16.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	170	2.4	13.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	1.9	11.0
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	132	1.9	10.6
11	Chronic liver disease and cirrhosis (K70,K73-K74)	125	1.8	10.1
12	Septicemia (A40-A41)	111	1.6	8.9
13	Influenza and pneumonia (J09-J18)	84	1.2	6.8
14	Chronic lower respiratory diseases (J40-J47)	71	1.0	5.7
15	Aortic aneurysm and dissection (I71)	64	0.9	5.1
...	All other causes (residual)	1,118	15.9	90.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 50-54 years				
...	All causes	11,005	100.0	861.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,987	27.1	233.8
2	Malignant neoplasms (C00-C97)	2,070	18.8	162.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,228	11.2	96.1
4	Diabetes mellitus (E10-E14)	547	5.0	42.8
5	Cerebrovascular diseases (I60-I69)	486	4.4	38.0
6	Human immunodeficiency virus (HIV) disease (B20-B24)	327	3.0	25.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	281	2.6	22.0
8	Chronic liver disease and cirrhosis (K70,K73-K74)	262	2.4	20.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	236	2.1	18.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	185	1.7	14.5
11	Chronic lower respiratory diseases (J40-J47)	183	1.7	14.3
12	Septicemia (A40-A41)	180	1.6	14.1
13	Influenza and pneumonia (J09-J18)	145	1.3	11.4
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	135	1.2	10.6
15	Aortic aneurysm and dissection (I71)	53	0.5	4.1
...	All other causes (residual)	1,700	15.4	133.1
Non-Hispanic, black, male, 55-59 years				
...	All causes	16,437	100.0	1,351.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,474	27.2	367.7
2	Malignant neoplasms (C00-C97)	4,048	24.6	332.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,299	7.9	106.8
4	Diabetes mellitus (E10-E14)	759	4.6	62.4
5	Cerebrovascular diseases (I60-I69)	714	4.3	58.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	449	2.7	36.9
7	Chronic lower respiratory diseases (J40-J47)	397	2.4	32.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	384	2.3	31.6
9	Human immunodeficiency virus (HIV) disease (B20-B24)	383	2.3	31.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	299	1.8	24.6
11	Septicemia (A40-A41)	291	1.8	23.9
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	227	1.4	18.7
13	Influenza and pneumonia (J09-J18)	201	1.2	16.5
14	Viral hepatitis (B15-B19)	140	0.9	11.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	115	0.7	9.5
...	All other causes (residual)	2,257	13.7	185.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 60-64 years				
...	All causes	20,064	100.0	2,064.3
1	Malignant neoplasms (C00-C97)	5,711	28.5	587.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,264	26.2	541.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,053	5.2	108.3
4	Cerebrovascular diseases (I60-I69)	961	4.8	98.9
5	Diabetes mellitus (E10-E14)	935	4.7	96.2
6	Chronic lower respiratory diseases (J40-J47)	646	3.2	66.5
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	530	2.6	54.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	440	2.2	45.3
9	Septicemia (A40-A41)	391	1.9	40.2
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	342	1.7	35.2
11	Influenza and pneumonia (J09-J18)	291	1.5	29.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	268	1.3	27.6
13	Viral hepatitis (B15-B19)	184	0.9	18.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	146	0.7	15.0
15	Pneumonitis due to solids and liquids (J69)	94	0.5	9.7
...	All other causes (residual)	2,808	14.0	288.9
Non-Hispanic, black, male, 65-69 years				
...	All causes	20,252	100.0	2,766.0
1	Malignant neoplasms (C00-C97)	5,881	29.0	803.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,392	26.6	736.4
3	Cerebrovascular diseases (I60-I69)	1,074	5.3	146.7
4	Diabetes mellitus (E10-E14)	1,023	5.1	139.7
5	Chronic lower respiratory diseases (J40-J47)	849	4.2	116.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	720	3.6	98.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	591	2.9	80.7
8	Septicemia (A40-A41)	434	2.1	59.3
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	346	1.7	47.3
10	Influenza and pneumonia (J09-J18)	327	1.6	44.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	290	1.4	39.6
12	Human immunodeficiency virus (HIV) disease (B20-B24)	167	0.8	22.8
13	Viral hepatitis (B15-B19)	140	0.7	19.1
14	Alzheimer's disease (G30)	100	0.5	13.7
15	Pneumonitis due to solids and liquids (J69)	97	0.5	13.2
...	All other causes (residual)	2,821	13.9	385.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 70-74 years				
...	All causes	17,049	100.0	3,791.5
1	Malignant neoplasms (C00-C97)	4,800	28.2	1,067.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,429	26.0	985.0
3	Cerebrovascular diseases (I60-I69)	983	5.8	218.6
4	Diabetes mellitus (E10-E14)	893	5.2	198.6
5	Chronic lower respiratory diseases (J40-J47)	800	4.7	177.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	546	3.2	121.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	382	2.2	85.0
8	Septicemia (A40-A41)	375	2.2	83.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	348	2.0	77.4
10	Influenza and pneumonia (J09-J18)	295	1.7	65.6
11	Alzheimer's disease (G30)	203	1.2	45.1
12	Chronic liver disease and cirrhosis (K70,K73-K74)	157	0.9	34.9
13	Pneumonitis due to solids and liquids (J69)	117	0.7	26.0
14	Parkinson's disease (G20-G21)	103	0.6	22.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	86	0.5	19.1
...	All other causes (residual)	2,532	14.9	563.1
Non-Hispanic, black, male, 75-79 years				
...	All causes	15,654	100.0	5,330.1
1	Malignant neoplasms (C00-C97)	4,050	25.9	1,379.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,988	25.5	1,357.9
3	Cerebrovascular diseases (I60-I69)	1,014	6.5	345.3
4	Chronic lower respiratory diseases (J40-J47)	794	5.1	270.4
5	Diabetes mellitus (E10-E14)	742	4.7	252.6
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	530	3.4	180.5
7	Alzheimer's disease (G30)	404	2.6	137.6
8	Septicemia (A40-A41)	370	2.4	126.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	329	2.1	112.0
10	Influenza and pneumonia (J09-J18)	323	2.1	110.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	312	2.0	106.2
12	Parkinson's disease (G20-G21)	169	1.1	57.5
13	Pneumonitis due to solids and liquids (J69)	118	0.8	40.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	94	0.6	32.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	76	0.5	25.9
...	All other causes (residual)	2,341	15.0	797.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 80-84 years				
...	All causes	13,999	100.0	7,835.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,685	26.3	2,062.5
2	Malignant neoplasms (C00-C97)	3,158	22.6	1,767.5
3	Cerebrovascular diseases (I60-I69)	936	6.7	523.9
4	Chronic lower respiratory diseases (J40-J47)	649	4.6	363.2
5	Diabetes mellitus (E10-E14)	610	4.4	341.4
6	Alzheimer's disease (G30)	560	4.0	313.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	475	3.4	265.9
8	Septicemia (A40-A41)	310	2.2	173.5
9	Influenza and pneumonia (J09-J18)	288	2.1	161.2
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	267	1.9	149.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	265	1.9	148.3
12	Parkinson's disease (G20-G21)	190	1.4	106.3
13	Pneumonitis due to solids and liquids (J69)	141	1.0	78.9
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	73	0.5	40.9
15	Chronic liver disease and cirrhosis (K70,K73-K74)	51	0.4	28.5
...	All other causes (residual)	2,341	16.7	1,310.3
Non-Hispanic, black, male, 85-89 years				
...	All causes	10,383	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,792	26.9	---
2	Malignant neoplasms (C00-C97)	2,073	20.0	---
3	Cerebrovascular diseases (I60-I69)	683	6.6	---
4	Alzheimer's disease (G30)	530	5.1	---
5	Chronic lower respiratory diseases (J40-J47)	454	4.4	---
6	Diabetes mellitus (E10-E14)	400	3.9	---
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	382	3.7	---
8	Influenza and pneumonia (J09-J18)	272	2.6	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	222	2.1	---
10	Septicemia (A40-A41)	213	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	1.7	---
12	Parkinson's disease (G20-G21)	136	1.3	---
13	Pneumonitis due to solids and liquids (J69)	110	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	54	0.5	---
15	Nutritional deficiencies (E40-E64)	46	0.4	---
...	All other causes (residual)	1,835	17.7	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 90-94 years				
...	All causes	6,095	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,628	26.7	---
2	Malignant neoplasms (C00-C97)	1,001	16.4	---
3	Alzheimer's disease (G30)	419	6.9	---
4	Cerebrovascular diseases (I60-I69)	382	6.3	---
5	Chronic lower respiratory diseases (J40-J47)	285	4.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	196	3.2	---
7	Diabetes mellitus (E10-E14)	178	2.9	---
8	Influenza and pneumonia (J09-J18)	165	2.7	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	146	2.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	115	1.9	---
11	Septicemia (A40-A41)	108	1.8	---
12	Pneumonitis due to solids and liquids (J69)	91	1.5	---
13	Parkinson's disease (G20-G21)	58	1.0	---
14	Nutritional deficiencies (E40-E64)	38	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.6	---
...	All other causes (residual)	1,249	20.5	---
Non-Hispanic, black, male, 95-99 years				
...	All causes	2,029	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	614	30.3	---
2	Malignant neoplasms (C00-C97)	239	11.8	---
3	Alzheimer's disease (G30)	153	7.5	---
4	Cerebrovascular diseases (I60-I69)	139	6.9	---
5	Chronic lower respiratory diseases (J40-J47)	66	3.3	---
6	Influenza and pneumonia (J09-J18)	65	3.2	---
7	Diabetes mellitus (E10-E14)	58	2.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	56	2.8	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	54	2.7	---
10	Septicemia (A40-A41)	42	2.1	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	1.9	---
12	Pneumonitis due to solids and liquids (J69)	22	1.1	---
13	Parkinson's disease (G20-G21)	17	0.8	---
14	Nutritional deficiencies (E40-E64)	14	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.4	---
...	All other causes (residual)	444	21.9	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 100 years and over				
...	All causes	394	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	126	32.0	---
2	Alzheimer's disease (G30)	27	6.9	---
3	Malignant neoplasms (C00-C97)	25	6.3	---
3	Cerebrovascular diseases (I60-I69)	25	6.3	---
5	Chronic lower respiratory diseases (J40-J47)	19	4.8	---
6	Influenza and pneumonia (J09-J18)	13	3.3	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	3.3	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	3.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	2.0	---
10	Septicemia (A40-A41)	7	1.8	---
11	Diabetes mellitus (E10-E14)	4	1.0	---
11	Pneumonitis due to solids and liquids (J69)	4	1.0	---
13	Anemias (D50-D64)	3	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.5	---
14	Atherosclerosis (I70)	2	0.5	---
14	Aortic aneurysm and dissection (I71)	2	0.5	---
...	All other causes (residual)	102	25.9	---
Non-Hispanic, black, female, all ages				
...	All causes	158,060	100.0	719.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,564	23.1	166.5
2	Malignant neoplasms (C00-C97)	34,510	21.8	157.1
3	Cerebrovascular diseases (I60-I69)	10,074	6.4	45.9
4	Diabetes mellitus (E10-E14)	7,077	4.5	32.2
5	Alzheimer's disease (G30)	6,126	3.9	27.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,636	3.6	25.7
7	Chronic lower respiratory diseases (J40-J47)	5,319	3.4	24.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,852	3.1	22.1
9	Septicemia (A40-A41)	3,610	2.3	16.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,119	2.0	14.2
11	Influenza and pneumonia (J09-J18)	2,763	1.7	12.6
12	Certain conditions originating in the perinatal period (P00-P96)	1,674	1.1	7.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,328	0.8	6.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,151	0.7	5.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,077	0.7	4.9
...	All other causes (residual)	33,180	21.0	151.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 1-4 years				
...	All causes	441	100.0	36.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	110	24.9	9.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	68	15.4	5.7
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	12.9	4.7
4	Malignant neoplasms (C00-C97)	26	5.9	2.2
5	Influenza and pneumonia (J09-J18)	16	3.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.7	*
7	Certain conditions originating in the perinatal period (P00-P96)	9	2.0	*
8	Septicemia (A40-A41)	8	1.8	*
8	Anemias (D50-D64)	8	1.8	*
8	Cerebrovascular diseases (I60-I69)	8	1.8	*
11	Chronic lower respiratory diseases (J40-J47)	5	1.1	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.7	*
14	Meningitis (G00,G03)	2	0.5	*
15	Diabetes mellitus (E10-E14)	1	0.2	*
15	Acute bronchitis and bronchiolitis (J20-J21)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	102	23.1	8.5
Non-Hispanic, black, female, 5-9 years				
...	All causes	277	100.0	18.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	32.9	5.9
2	Malignant neoplasms (C00-C97)	42	15.2	2.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	9.0	1.6
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	7.6	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	5.4	*
6	Chronic lower respiratory diseases (J40-J47)	12	4.3	*
7	Septicemia (A40-A41)	8	2.9	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.7	*
8	Anemias (D50-D64)	2	0.7	*
8	Meningitis (G00,G03)	2	0.7	*
8	Cerebrovascular diseases (I60-I69)	2	0.7	*
8	Influenza and pneumonia (J09-J18)	2	0.7	*
8	Pneumonitis due to solids and liquids (J69)	2	0.7	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
8	Certain conditions originating in the perinatal period (P00-P96)	2	0.7	*
8	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.7	*
...	All other causes (residual)	45	16.2	2.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 10-14 years				
...	All causes	255	100.0	16.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	25.1	4.3
2	Malignant neoplasms (C00-C97)	36	14.1	2.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	8.6	1.5
4	Chronic lower respiratory diseases (J40-J47)	18	7.1	*
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	5.9	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	5.1	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	3.1	*
8	Cerebrovascular diseases (I60-I69)	6	2.4	*
9	Influenza and pneumonia (J09-J18)	5	2.0	*
10	Diabetes mellitus (E10-E14)	3	1.2	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	1.2	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.8	*
12	Diseases of appendix (K35-K38)	2	0.8	*
14	Septicemia (A40-A41)	1	0.4	*
14	Nutritional deficiencies (E40-E64)	1	0.4	*
14	Meningitis (G00,G03)	1	0.4	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
14	Pneumonitis due to solids and liquids (J69)	1	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.4	*
...	All other causes (residual)	49	19.2	3.3
Non-Hispanic, black, female, 15-19 years				
...	All causes	618	100.0	39.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	178	28.8	11.3
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	131	21.2	8.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	8.4	3.3
4	Malignant neoplasms (C00-C97)	41	6.6	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	4.9	1.9
6	Chronic lower respiratory diseases (J40-J47)	18	2.9	*
7	Anemias (D50-D64)	13	2.1	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.9	*
9	Influenza and pneumonia (J09-J18)	7	1.1	*
10	Septicemia (A40-A41)	6	1.0	*
10	Cerebrovascular diseases (I60-I69)	6	1.0	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	6	1.0	*
13	Diabetes mellitus (E10-E14)	5	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.3	*
...	All other causes (residual)	106	17.2	6.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 20-24 years				
...	All causes	1,176	100.0	68.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	352	29.9	20.5
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	195	16.6	11.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	6.8	4.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	6.5	4.4
5	Pregnancy, childbirth and the puerperium (O00-O99)	61	5.2	3.6
6	Malignant neoplasms (C00-C97)	52	4.4	3.0
7	Diabetes mellitus (E10-E14)	26	2.2	1.5
8	Human immunodeficiency virus (HIV) disease (B20-B24)	22	1.9	1.3
9	Influenza and pneumonia (J09-J18)	19	1.6	*
10	Septicemia (A40-A41)	16	1.4	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	1.2	*
12	Chronic lower respiratory diseases (J40-J47)	12	1.0	*
13	Anemias (D50-D64)	11	0.9	*
14	Cerebrovascular diseases (I60-I69)	10	0.9	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.6	*
...	All other causes (residual)	223	19.0	13.0
Non-Hispanic, black, female, 25-29 years				
...	All causes	1,544	100.0	90.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	369	23.9	21.6
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	195	12.6	11.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	136	8.8	8.0
4	Malignant neoplasms (C00-C97)	125	8.1	7.3
5	Pregnancy, childbirth and the puerperium (O00-O99)	86	5.6	5.0
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	64	4.1	3.7
7	Diabetes mellitus (E10-E14)	41	2.7	2.4
8	Human immunodeficiency virus (HIV) disease (B20-B24)	37	2.4	2.2
9	Anemias (D50-D64)	36	2.3	2.1
10	Chronic lower respiratory diseases (J40-J47)	25	1.6	1.5
11	Influenza and pneumonia (J09-J18)	22	1.4	1.3
12	Cerebrovascular diseases (I60-I69)	21	1.4	1.2
13	Septicemia (A40-A41)	20	1.3	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	18	1.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.0	*
...	All other causes (residual)	334	21.6	19.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 30-34 years				
...	All causes	1,862	100.0	124.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	335	18.0	22.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	248	13.3	16.5
3	Malignant neoplasms (C00-C97)	241	12.9	16.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	144	7.7	9.6
5	Diabetes mellitus (E10-E14)	70	3.8	4.7
6	Pregnancy, childbirth and the puerperium (O00-O99)	68	3.7	4.5
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	62	3.3	4.1
8	Human immunodeficiency virus (HIV) disease (B20-B24)	58	3.1	3.9
9	Anemias (D50-D64)	41	2.2	2.7
10	Influenza and pneumonia (J09-J18)	36	1.9	2.4
11	Cerebrovascular diseases (I60-I69)	34	1.8	2.3
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.7	2.1
13	Chronic lower respiratory diseases (J40-J47)	27	1.5	1.8
14	Septicemia (A40-A41)	24	1.3	1.6
15	Chronic liver disease and cirrhosis (K70,K73-K74)	23	1.2	1.5
...	All other causes (residual)	420	22.6	28.0
Non-Hispanic, black, female, 35-39 years				
...	All causes	2,586	100.0	178.9
1	Malignant neoplasms (C00-C97)	443	17.1	30.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	410	15.9	28.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	409	15.8	28.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	4.8	8.6
5	Diabetes mellitus (E10-E14)	108	4.2	7.5
6	Human immunodeficiency virus (HIV) disease (B20-B24)	92	3.6	6.4
7	Cerebrovascular diseases (I60-I69)	88	3.4	6.1
8	Pregnancy, childbirth and the puerperium (O00-O99)	73	2.8	5.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	57	2.2	3.9
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	52	2.0	3.6
11	Anemias (D50-D64)	45	1.7	3.1
12	Septicemia (A40-A41)	44	1.7	3.0
13	Chronic liver disease and cirrhosis (K70,K73-K74)	39	1.5	2.7
14	Influenza and pneumonia (J09-J18)	33	1.3	2.3
15	Chronic lower respiratory diseases (J40-J47)	26	1.0	1.8
...	All other causes (residual)	542	21.0	37.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 40-44 years				
...	All causes	3,293	100.0	243.5
1	Malignant neoplasms (C00-C97)	722	21.9	53.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	606	18.4	44.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	321	9.7	23.7
4	Diabetes mellitus (E10-E14)	151	4.6	11.2
5	Human immunodeficiency virus (HIV) disease (B20-B24)	142	4.3	10.5
6	Cerebrovascular diseases (I60-I69)	133	4.0	9.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	86	2.6	6.4
8	Septicemia (A40-A41)	79	2.4	5.8
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	73	2.2	5.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	64	1.9	4.7
11	Influenza and pneumonia (J09-J18)	55	1.7	4.1
11	Chronic lower respiratory diseases (J40-J47)	55	1.7	4.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	1.5	3.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.3	3.3
15	Pregnancy, childbirth and the puerperium (O00-O99)	35	1.1	2.6
...	All other causes (residual)	679	20.6	50.2
Non-Hispanic, black, female, 45-49 years				
...	All causes	5,190	100.0	369.7
1	Malignant neoplasms (C00-C97)	1,333	25.7	95.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,037	20.0	73.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	447	8.6	31.8
4	Cerebrovascular diseases (I60-I69)	249	4.8	17.7
5	Diabetes mellitus (E10-E14)	228	4.4	16.2
6	Human immunodeficiency virus (HIV) disease (B20-B24)	156	3.0	11.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	127	2.4	9.0
8	Septicemia (A40-A41)	107	2.1	7.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	93	1.8	6.6
10	Chronic lower respiratory diseases (J40-J47)	87	1.7	6.2
11	Influenza and pneumonia (J09-J18)	85	1.6	6.1
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	85	1.6	6.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	78	1.5	5.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	0.8	2.8
15	Pregnancy, childbirth and the puerperium (O00-O99)	37	0.7	2.6
...	All other causes (residual)	1,002	19.3	71.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 50-54 years				
...	All causes	8,415	100.0	579.9
1	Malignant neoplasms (C00-C97)	2,489	29.6	171.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,774	21.1	122.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	513	6.1	35.4
4	Cerebrovascular diseases (I60-I69)	417	5.0	28.7
5	Diabetes mellitus (E10-E14)	399	4.7	27.5
6	Chronic lower respiratory diseases (J40-J47)	241	2.9	16.6
7	Septicemia (A40-A41)	193	2.3	13.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	188	2.2	13.0
9	Human immunodeficiency virus (HIV) disease (B20-B24)	175	2.1	12.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	163	1.9	11.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	159	1.9	11.0
12	Influenza and pneumonia (J09-J18)	105	1.2	7.2
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	0.7	4.3
14	Anemias (D50-D64)	33	0.4	2.3
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	0.4	2.3
...	All other causes (residual)	1,471	17.5	101.4
Non-Hispanic, black, female, 55-59 years				
...	All causes	11,954	100.0	846.3
1	Malignant neoplasms (C00-C97)	3,864	32.3	273.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,529	21.2	179.0
3	Diabetes mellitus (E10-E14)	608	5.1	43.0
4	Cerebrovascular diseases (I60-I69)	557	4.7	39.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	493	4.1	34.9
6	Chronic lower respiratory diseases (J40-J47)	476	4.0	33.7
7	Septicemia (A40-A41)	318	2.7	22.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	276	2.3	19.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	241	2.0	17.1
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	177	1.5	12.5
11	Influenza and pneumonia (J09-J18)	172	1.4	12.2
12	Human immunodeficiency virus (HIV) disease (B20-B24)	129	1.1	9.1
13	Viral hepatitis (B15-B19)	69	0.6	4.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	61	0.5	4.3
15	Anemias (D50-D64)	53	0.4	3.8
...	All other causes (residual)	1,931	16.2	136.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 60-64 years				
...	All causes	14,366	100.0	1,207.2
1	Malignant neoplasms (C00-C97)	4,587	31.9	385.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,267	22.7	274.5
3	Diabetes mellitus (E10-E14)	759	5.3	63.8
4	Cerebrovascular diseases (I60-I69)	724	5.0	60.8
5	Chronic lower respiratory diseases (J40-J47)	560	3.9	47.1
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	468	3.3	39.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	384	2.7	32.3
8	Septicemia (A40-A41)	378	2.6	31.8
9	Influenza and pneumonia (J09-J18)	219	1.5	18.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	211	1.5	17.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	201	1.4	16.9
12	Human immunodeficiency virus (HIV) disease (B20-B24)	140	1.0	11.8
13	Viral hepatitis (B15-B19)	88	0.6	7.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	57	0.4	4.8
15	Pneumonitis due to solids and liquids (J69)	54	0.4	4.5
...	All other causes (residual)	2,269	15.8	190.7
Non-Hispanic, black, female, 65-69 years				
...	All causes	15,595	100.0	1,648.5
1	Malignant neoplasms (C00-C97)	4,963	31.8	524.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,650	23.4	385.8
3	Diabetes mellitus (E10-E14)	866	5.6	91.5
4	Cerebrovascular diseases (I60-I69)	852	5.5	90.1
5	Chronic lower respiratory diseases (J40-J47)	647	4.1	68.4
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	534	3.4	56.4
7	Septicemia (A40-A41)	384	2.5	40.6
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	301	1.9	31.8
9	Influenza and pneumonia (J09-J18)	262	1.7	27.7
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	243	1.6	25.7
11	Chronic liver disease and cirrhosis (K70,K73-K74)	124	0.8	13.1
12	Alzheimer's disease (G30)	118	0.8	12.5
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	78	0.5	8.2
14	Pneumonitis due to solids and liquids (J69)	77	0.5	8.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	66	0.4	7.0
...	All other causes (residual)	2,430	15.6	256.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 70-74 years				
...	All causes	14,732	100.0	2,369.1
1	Malignant neoplasms (C00-C97)	4,082	27.7	656.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,539	24.0	569.1
3	Cerebrovascular diseases (I60-I69)	982	6.7	157.9
4	Diabetes mellitus (E10-E14)	827	5.6	133.0
5	Chronic lower respiratory diseases (J40-J47)	687	4.7	110.5
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	546	3.7	87.8
7	Septicemia (A40-A41)	398	2.7	64.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	297	2.0	47.8
9	Alzheimer's disease (G30)	269	1.8	43.3
10	Influenza and pneumonia (J09-J18)	243	1.6	39.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	1.5	35.4
12	Pneumonitis due to solids and liquids (J69)	78	0.5	12.5
13	Chronic liver disease and cirrhosis (K70,K73-K74)	73	0.5	11.7
14	Parkinson's disease (G20-G21)	71	0.5	11.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	55	0.4	8.8
...	All other causes (residual)	2,365	16.1	380.3
Non-Hispanic, black, female, 75-79 years				
...	All causes	15,832	100.0	3,550.2
1	Malignant neoplasms (C00-C97)	3,871	24.5	868.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,851	24.3	863.6
3	Cerebrovascular diseases (I60-I69)	1,180	7.5	264.6
4	Diabetes mellitus (E10-E14)	824	5.2	184.8
5	Chronic lower respiratory diseases (J40-J47)	675	4.3	151.4
6	Alzheimer's disease (G30)	616	3.9	138.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	614	3.9	137.7
8	Septicemia (A40-A41)	429	2.7	96.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	357	2.3	80.1
10	Influenza and pneumonia (J09-J18)	297	1.9	66.6
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	196	1.2	44.0
12	Pneumonitis due to solids and liquids (J69)	103	0.7	23.1
13	Parkinson's disease (G20-G21)	98	0.6	22.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	67	0.4	15.0
15	Anemias (D50-D64)	49	0.3	11.0
...	All other causes (residual)	2,605	16.5	584.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 80-84 years				
...	All causes	17,589	100.0	5,623.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,543	25.8	1,452.4
2	Malignant neoplasms (C00-C97)	3,328	18.9	1,064.0
3	Cerebrovascular diseases (I60-I69)	1,370	7.8	438.0
4	Alzheimer's disease (G30)	1,122	6.4	358.7
5	Diabetes mellitus (E10-E14)	798	4.5	255.1
6	Chronic lower respiratory diseases (J40-J47)	697	4.0	222.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	694	3.9	221.9
8	Septicemia (A40-A41)	422	2.4	134.9
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	397	2.3	126.9
10	Influenza and pneumonia (J09-J18)	332	1.9	106.1
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	204	1.2	65.2
12	Pneumonitis due to solids and liquids (J69)	129	0.7	41.2
13	Parkinson's disease (G20-G21)	122	0.7	39.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	96	0.5	30.7
15	Nutritional deficiencies (E40-E64)	64	0.4	20.5
...	All other causes (residual)	3,271	18.6	1,045.8
Non-Hispanic, black, female, 85-89 years				
...	All causes	16,979	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,448	26.2	---
2	Malignant neoplasms (C00-C97)	2,460	14.5	---
3	Alzheimer's disease (G30)	1,433	8.4	---
4	Cerebrovascular diseases (I60-I69)	1,431	8.4	---
5	Diabetes mellitus (E10-E14)	702	4.1	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	610	3.6	---
7	Chronic lower respiratory diseases (J40-J47)	570	3.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	413	2.4	---
9	Septicemia (A40-A41)	370	2.2	---
10	Influenza and pneumonia (J09-J18)	306	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	232	1.4	---
12	Pneumonitis due to solids and liquids (J69)	137	0.8	---
13	Parkinson's disease (G20-G21)	121	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	88	0.5	---
15	Nutritional deficiencies (E40-E64)	67	0.4	---
...	All other causes (residual)	3,591	21.1	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 90-94 years				
...	All causes	13,613	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,805	28.0	---
2	Alzheimer's disease (G30)	1,478	10.9	---
3	Malignant neoplasms (C00-C97)	1,307	9.6	---
4	Cerebrovascular diseases (I60-I69)	1,226	9.0	---
5	Diabetes mellitus (E10-E14)	459	3.4	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	399	2.9	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	362	2.7	---
8	Chronic lower respiratory diseases (J40-J47)	323	2.4	---
9	Influenza and pneumonia (J09-J18)	303	2.2	---
10	Septicemia (A40-A41)	239	1.8	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	1.3	---
12	Pneumonitis due to solids and liquids (J69)	116	0.9	---
13	Nutritional deficiencies (E40-E64)	82	0.6	---
14	Parkinson's disease (G20-G21)	79	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	66	0.5	---
...	All other causes (residual)	3,195	23.5	---
Non-Hispanic, black, female, 95-99 years				
...	All causes	6,466	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,882	29.1	---
2	Alzheimer's disease (G30)	772	11.9	---
3	Cerebrovascular diseases (I60-I69)	560	8.7	---
4	Malignant neoplasms (C00-C97)	418	6.5	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	205	3.2	---
6	Diabetes mellitus (E10-E14)	162	2.5	---
7	Influenza and pneumonia (J09-J18)	154	2.4	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	143	2.2	---
9	Chronic lower respiratory diseases (J40-J47)	123	1.9	---
10	Septicemia (A40-A41)	103	1.6	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	1.4	---
12	Pneumonitis due to solids and liquids (J69)	55	0.9	---
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	42	0.6	---
14	Atherosclerosis (I70)	36	0.6	---
15	Nutritional deficiencies (E40-E64)	35	0.5	---
...	All other causes (residual)	1,688	26.1	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 100 years and over				
...	All causes	2,157	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	649	30.1	---
2	Alzheimer's disease (G30)	258	12.0	---
3	Cerebrovascular diseases (I60-I69)	202	9.4	---
4	Malignant neoplasms (C00-C97)	76	3.5	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	66	3.1	---
6	Influenza and pneumonia (J09-J18)	62	2.9	---
7	Diabetes mellitus (E10-E14)	40	1.9	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	38	1.8	---
9	Septicemia (A40-A41)	32	1.5	---
9	Chronic lower respiratory diseases (J40-J47)	32	1.5	---
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	1.3	---
12	Pneumonitis due to solids and liquids (J69)	17	0.8	---
13	Nutritional deficiencies (E40-E64)	14	0.6	---
14	Atherosclerosis (I70)	12	0.6	---
15	Anemias (D50-D64)	9	0.4	---
...	All other causes (residual)	623	28.9	---
Non-Hispanic, American Indian or Alaska Native, both sexes, all ages				
...	All causes	18,595	100.0	685.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,332	17.9	122.9
2	Malignant neoplasms (C00-C97)	3,225	17.3	119.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,100	11.3	77.5
4	Diabetes mellitus (E10-E14)	1,078	5.8	39.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,010	5.4	37.3
6	Chronic lower respiratory diseases (J40-J47)	853	4.6	31.5
7	Cerebrovascular diseases (I60-I69)	632	3.4	23.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	580	3.1	21.4
9	Alzheimer's disease (G30)	377	2.0	13.9
10	Influenza and pneumonia (J09-J18)	360	1.9	13.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	343	1.8	12.7
12	Septicemia (A40-A41)	295	1.6	10.9
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	279	1.5	10.3
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	193	1.0	7.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	120	0.6	4.4
...	All other causes (residual)	3,818	20.5	140.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 1-4 years				
...	All causes	82	100.0	51.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	32.9	16.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	12.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	9.8	*
4	Malignant neoplasms (C00-C97)	3	3.7	*
4	Influenza and pneumonia (J09-J18)	3	3.7	*
6	Meningococcal infection (A39)	2	2.4	*
6	Septicemia (A40-A41)	2	2.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.4	*
9	Diabetes mellitus (E10-E14)	1	1.2	*
9	Cerebrovascular diseases (I60-I69)	1	1.2	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.2	*
...	All other causes (residual)	22	26.8	13.8
Non-Hispanic, American Indian or Alaska Native, both sexes, 5-9 years				
...	All causes	42	100.0	20.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	40.5	*
2	Malignant neoplasms (C00-C97)	2	4.8	*
2	Meningitis (G00,G03)	2	4.8	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	4.8	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.8	*
6	Meningococcal infection (A39)	1	2.4	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.4	*
6	Chronic lower respiratory diseases (J40-J47)	1	2.4	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	2.4	*
6	Complications of medical and surgical care (Y40-Y84,Y88)	1	2.4	*
...	All other causes (residual)	12	28.6	*
Non-Hispanic, American Indian or Alaska Native, both sexes, 10-14 years				
...	All causes	50	100.0	24.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	28.0	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	13	26.0	*
3	Malignant neoplasms (C00-C97)	6	12.0	*
4	Septicemia (A40-A41)	2	4.0	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	4.0	*
6	Diabetes mellitus (E10-E14)	1	2.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.0	*
6	Cerebrovascular diseases (I60-I69)	1	2.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.0	*
...	All other causes (residual)	9	18.0	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 15-19 years				
...	All causes	187	100.0	88.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	81	43.3	38.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	55	29.4	26.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	8.6	*
4	Malignant neoplasms (C00-C97)	4	2.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	1.6	*
6	Influenza and pneumonia (J09-J18)	2	1.1	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	2	1.1	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
8	Diabetes mellitus (E10-E14)	1	0.5	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.5	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.5	*
8	Legal intervention (Y35,Y89.0)	1	0.5	*
...	All other causes (residual)	19	10.2	*
Non-Hispanic, American Indian or Alaska Native, both sexes, 20-24 years				
...	All causes	404	100.0	181.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	189	46.8	84.7
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	25.0	45.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	9.9	17.9
4	Malignant neoplasms (C00-C97)	6	1.5	*
4	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.5	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.0	*
7	Diabetes mellitus (E10-E14)	3	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
9	Cerebrovascular diseases (I60-I69)	2	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
9	Legal intervention (Y35,Y89.0)	2	0.5	*
12	Septicemia (A40-A41)	1	0.2	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
12	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.2	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.2	*
...	All other causes (residual)	42	10.4	18.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 25-29 years				
...	All causes	533	100.0	247.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	40.3	100.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	17.3	42.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	41	7.7	19.1
4	Chronic liver disease and cirrhosis (K70,K73-K74)	31	5.8	14.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	27	5.1	12.6
6	Malignant neoplasms (C00-C97)	14	2.6	*
7	Diabetes mellitus (E10-E14)	7	1.3	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.3	*
9	Legal intervention (Y35,Y89.0)	5	0.9	*
10	Septicemia (A40-A41)	4	0.8	*
11	Cerebrovascular diseases (I60-I69)	3	0.6	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Viral hepatitis (B15-B19)	1	0.2	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.2	*
14	Anemias (D50-D64)	1	0.2	*
14	Influenza and pneumonia (J09-J18)	1	0.2	*
14	Hernia (K40-K46)	1	0.2	*
14	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.2	*
...	All other causes (residual)	75	14.1	34.9
Non-Hispanic, American Indian or Alaska Native, both sexes, 30-34 years				
...	All causes	570	100.0	306.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	185	32.5	99.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	78	13.7	42.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	68	11.9	36.6
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	6.7	20.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	34	6.0	18.3
6	Diabetes mellitus (E10-E14)	18	3.2	*
7	Malignant neoplasms (C00-C97)	17	3.0	*
8	Cerebrovascular diseases (I60-I69)	7	1.2	*
8	Influenza and pneumonia (J09-J18)	7	1.2	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.9	*
10	Legal intervention (Y35,Y89.0)	5	0.9	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.7	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.5	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.5	*
...	All other causes (residual)	94	16.5	50.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 35-39 years				
...	All causes	622	100.0	365.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	186	29.9	109.4
2	Chronic liver disease and cirrhosis (K70,K73-K74)	72	11.6	42.3
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	10.6	38.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	9.3	34.1
5	Malignant neoplasms (C00-C97)	35	5.6	20.6
6	Diabetes mellitus (E10-E14)	34	5.5	20.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	5.0	18.2
8	Influenza and pneumonia (J09-J18)	10	1.6	*
9	Cerebrovascular diseases (I60-I69)	9	1.4	*
10	Septicemia (A40-A41)	8	1.3	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
14	Pneumonitis due to solids and liquids (J69)	3	0.5	*
15	Viral hepatitis (B15-B19)	2	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
...	All other causes (residual)	93	15.0	54.7
Non-Hispanic, American Indian or Alaska Native, both sexes, 40-44 years				
...	All causes	710	100.0	452.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	24.9	112.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	102	14.4	65.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	99	13.9	63.1
4	Malignant neoplasms (C00-C97)	67	9.4	42.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36	5.1	23.0
6	Diabetes mellitus (E10-E14)	34	4.8	21.7
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	3.5	15.9
8	Cerebrovascular diseases (I60-I69)	15	2.1	*
9	Influenza and pneumonia (J09-J18)	10	1.4	*
10	Septicemia (A40-A41)	8	1.1	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.1	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	6	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.8	*
14	Chronic lower respiratory diseases (J40-J47)	5	0.7	*
14	Pneumonitis due to solids and liquids (J69)	5	0.7	*
...	All other causes (residual)	107	15.1	68.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 45-49 years				
...	All causes	919	100.0	563.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	18.9	106.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	134	14.6	82.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	14.0	79.0
4	Malignant neoplasms (C00-C97)	105	11.4	64.3
5	Diabetes mellitus (E10-E14)	48	5.2	29.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	2.7	15.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	23	2.5	14.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	2.2	12.3
9	Cerebrovascular diseases (I60-I69)	17	1.8	*
9	Influenza and pneumonia (J09-J18)	17	1.8	*
11	Septicemia (A40-A41)	14	1.5	*
12	Viral hepatitis (B15-B19)	10	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	10	1.1	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.7	*
15	Aortic aneurysm and dissection (I71)	4	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.4	*
...	All other causes (residual)	179	19.5	109.7
Non-Hispanic, American Indian or Alaska Native, both sexes, 50-54 years				
...	All causes	1,373	100.0	781.7
1	Malignant neoplasms (C00-C97)	218	15.9	124.1
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	217	15.8	123.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	215	15.7	122.4
4	Chronic liver disease and cirrhosis (K70,K73-K74)	153	11.1	87.1
5	Diabetes mellitus (E10-E14)	84	6.1	47.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38	2.8	21.6
7	Chronic lower respiratory diseases (J40-J47)	35	2.5	19.9
8	Cerebrovascular diseases (I60-I69)	32	2.3	18.2
9	Influenza and pneumonia (J09-J18)	25	1.8	14.2
10	Septicemia (A40-A41)	20	1.5	11.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18	1.3	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	1.1	*
13	Viral hepatitis (B15-B19)	14	1.0	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	0.9	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.7	*
...	All other causes (residual)	267	19.4	152.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 55-59 years				
...	All causes	1,647	100.0	953.2
1	Malignant neoplasms (C00-C97)	391	23.7	226.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	280	17.0	162.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	10.6	100.7
4	Chronic liver disease and cirrhosis (K70,K73-K74)	148	9.0	85.7
5	Diabetes mellitus (E10-E14)	94	5.7	54.4
6	Chronic lower respiratory diseases (J40-J47)	51	3.1	29.5
7	Cerebrovascular diseases (I60-I69)	35	2.1	20.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	33	2.0	19.1
9	Septicemia (A40-A41)	28	1.7	16.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	28	1.7	16.2
11	Influenza and pneumonia (J09-J18)	25	1.5	14.5
12	Viral hepatitis (B15-B19)	24	1.5	13.9
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11	0.7	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	0.6	*
15	Pneumonitis due to solids and liquids (J69)	8	0.5	*
...	All other causes (residual)	307	18.6	177.7
Non-Hispanic, American Indian or Alaska Native, both sexes, 60-64 years				
...	All causes	1,786	100.0	1,251.0
1	Malignant neoplasms (C00-C97)	435	24.4	304.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	342	19.1	239.5
3	Diabetes mellitus (E10-E14)	144	8.1	100.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	7.1	89.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	117	6.6	82.0
6	Chronic lower respiratory diseases (J40-J47)	99	5.5	69.3
7	Cerebrovascular diseases (I60-I69)	53	3.0	37.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	31	1.7	21.7
9	Influenza and pneumonia (J09-J18)	30	1.7	21.0
10	Septicemia (A40-A41)	27	1.5	18.9
11	Viral hepatitis (B15-B19)	21	1.2	14.7
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	1.1	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	1.0	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	13	0.7	*
15	Peptic ulcer (K25-K28)	8	0.4	*
...	All other causes (residual)	303	17.0	212.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 65-69 years				
...	All causes	1,824	100.0	1,621.4
1	Malignant neoplasms (C00-C97)	477	26.2	424.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	352	19.3	312.9
3	Diabetes mellitus (E10-E14)	146	8.0	129.8
4	Chronic lower respiratory diseases (J40-J47)	129	7.1	114.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	77	4.2	68.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	76	4.2	67.6
7	Cerebrovascular diseases (I60-I69)	67	3.7	59.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	45	2.5	40.0
9	Septicemia (A40-A41)	39	2.1	34.7
10	Influenza and pneumonia (J09-J18)	28	1.5	24.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.1	17.8
12	Alzheimer's disease (G30)	13	0.7	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	10	0.5	*
14	Viral hepatitis (B15-B19)	9	0.5	*
14	Pneumonitis due to solids and liquids (J69)	9	0.5	*
...	All other causes (residual)	327	17.9	290.7
Non-Hispanic, American Indian or Alaska Native, both sexes, 70-74 years				
...	All causes	1,813	100.0	2,530.5
1	Malignant neoplasms (C00-C97)	460	25.4	642.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	419	23.1	584.8
3	Chronic lower respiratory diseases (J40-J47)	146	8.1	203.8
4	Diabetes mellitus (E10-E14)	134	7.4	187.0
5	Cerebrovascular diseases (I60-I69)	52	2.9	72.6
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	52	2.9	72.6
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	2.6	65.6
8	Septicemia (A40-A41)	41	2.3	57.2
9	Influenza and pneumonia (J09-J18)	36	2.0	50.2
9	Chronic liver disease and cirrhosis (K70,K73-K74)	36	2.0	50.2
11	Alzheimer's disease (G30)	27	1.5	37.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	1.3	32.1
13	Pneumonitis due to solids and liquids (J69)	12	0.7	*
14	Parkinson's disease (G20-G21)	11	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.4	*
15	Anemias (D50-D64)	7	0.4	*
15	Aortic aneurysm and dissection (I71)	7	0.4	*
...	All other causes (residual)	296	16.3	413.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 75-79 years				
...	All causes	1,703	100.0	3,612.6
1	Malignant neoplasms (C00-C97)	403	23.7	854.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	366	21.5	776.4
3	Chronic lower respiratory diseases (J40-J47)	127	7.5	269.4
4	Diabetes mellitus (E10-E14)	115	6.8	243.9
5	Cerebrovascular diseases (I60-I69)	86	5.0	182.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	3.5	125.2
7	Alzheimer's disease (G30)	45	2.6	95.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	42	2.5	89.1
9	Influenza and pneumonia (J09-J18)	38	2.2	80.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	2.1	74.2
11	Septicemia (A40-A41)	29	1.7	61.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	20	1.2	42.4
13	Parkinson's disease (G20-G21)	17	1.0	*
14	Pneumonitis due to solids and liquids (J69)	14	0.8	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	11	0.6	*
...	All other causes (residual)	296	17.4	627.9
Non-Hispanic, American Indian or Alaska Native, both sexes, 80-84 years				
...	All causes	1,629	100.0	5,756.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	393	24.1	1,388.7
2	Malignant neoplasms (C00-C97)	301	18.5	1,063.6
3	Chronic lower respiratory diseases (J40-J47)	115	7.1	406.4
4	Diabetes mellitus (E10-E14)	107	6.6	378.1
5	Cerebrovascular diseases (I60-I69)	92	5.6	325.1
6	Alzheimer's disease (G30)	76	4.7	268.6
7	Influenza and pneumonia (J09-J18)	43	2.6	151.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	2.3	130.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35	2.1	123.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	1.9	109.5
11	Septicemia (A40-A41)	29	1.8	102.5
12	Parkinson's disease (G20-G21)	24	1.5	84.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	18	1.1	*
14	Pneumonitis due to solids and liquids (J69)	15	0.9	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	12	0.7	*
...	All other causes (residual)	301	18.5	1,063.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 85-89 years				
...	All causes	1,281	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	309	24.1	---
2	Malignant neoplasms (C00-C97)	172	13.4	---
3	Alzheimer's disease (G30)	101	7.9	---
4	Chronic lower respiratory diseases (J40-J47)	81	6.3	---
5	Cerebrovascular diseases (I60-I69)	79	6.2	---
6	Diabetes mellitus (E10-E14)	70	5.5	---
7	Influenza and pneumonia (J09-J18)	48	3.7	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	39	3.0	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	23	1.8	---
11	Septicemia (A40-A41)	21	1.6	---
12	Pneumonitis due to solids and liquids (J69)	14	1.1	---
13	Chronic liver disease and cirrhosis (K70,K73-K74)	9	0.7	---
14	Nutritional deficiencies (E40-E64)	8	0.6	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.5	---
15	Parkinson's disease (G20-G21)	7	0.5	---
...	All other causes (residual)	263	20.5	---
Non-Hispanic, American Indian or Alaska Native, both sexes, 90-94 years				
...	All causes	779	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	206	26.4	---
2	Malignant neoplasms (C00-C97)	88	11.3	---
3	Alzheimer's disease (G30)	68	8.7	---
4	Cerebrovascular diseases (I60-I69)	55	7.1	---
5	Chronic lower respiratory diseases (J40-J47)	38	4.9	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	3.7	---
7	Diabetes mellitus (E10-E14)	24	3.1	---
8	Influenza and pneumonia (J09-J18)	22	2.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.3	---
10	Septicemia (A40-A41)	14	1.8	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.7	---
12	Pneumonitis due to solids and liquids (J69)	9	1.2	---
13	Parkinson's disease (G20-G21)	8	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	---
14	Nutritional deficiencies (E40-E64)	6	0.8	---
...	All other causes (residual)	175	22.5	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 95-99 years				
...	All causes	269	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	24.5	---
2	Alzheimer's disease (G30)	35	13.0	---
3	Cerebrovascular diseases (I60-I69)	19	7.1	---
4	Malignant neoplasms (C00-C97)	17	6.3	---
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	4.8	---
6	Diabetes mellitus (E10-E14)	12	4.5	---
7	Influenza and pneumonia (J09-J18)	10	3.7	---
8	Pneumonitis due to solids and liquids (J69)	4	1.5	---
9	Septicemia (A40-A41)	3	1.1	---
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	1.1	---
9	Chronic lower respiratory diseases (J40-J47)	3	1.1	---
9	Pneumoconioses and chemical effects (J60-J66,J68)	3	1.1	---
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.7	---
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	---
15	Anemias (D50-D64)	1	0.4	---
15	Nutritional deficiencies (E40-E64)	1	0.4	---
15	Parkinson's disease (G20-G21)	1	0.4	---
15	Peptic ulcer (K25-K28)	1	0.4	---
15	Diseases of appendix (K35-K38)	1	0.4	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	---
15	Hyperplasia of prostate (N40)	1	0.4	---
...	All other causes (residual)	70	26.0	---
Non-Hispanic, American Indian or Alaska Native, both sexes, 100 years and over				
...	All causes	60	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	33.3	---
2	Cerebrovascular diseases (I60-I69)	7	11.7	---
3	Malignant neoplasms (C00-C97)	3	5.0	---
3	Influenza and pneumonia (J09-J18)	3	5.0	---
3	Chronic lower respiratory diseases (J40-J47)	3	5.0	---
3	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	5.0	---
7	Diabetes mellitus (E10-E14)	1	1.7	---
7	Nutritional deficiencies (E40-E64)	1	1.7	---
7	Alzheimer's disease (G30)	1	1.7	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.7	---
7	Pneumoconioses and chemical effects (J60-J66,J68)	1	1.7	---
7	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	1.7	---
...	All other causes (residual)	15	25.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, all ages				
...	All causes	10,280	100.0	772.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,951	19.0	146.7
2	Malignant neoplasms (C00-C97)	1,723	16.8	129.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,404	13.7	105.5
4	Diabetes mellitus (E10-E14)	574	5.6	43.1
5	Chronic liver disease and cirrhosis (K70,K73-K74)	554	5.4	41.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	441	4.3	33.2
7	Chronic lower respiratory diseases (J40-J47)	430	4.2	32.3
8	Cerebrovascular diseases (I60-I69)	288	2.8	21.6
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	209	2.0	15.7
10	Influenza and pneumonia (J09-J18)	176	1.7	13.2
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	166	1.6	12.5
12	Septicemia (A40-A41)	128	1.2	9.6
13	Alzheimer's disease (G30)	121	1.2	9.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	0.8	6.1
15	Certain conditions originating in the perinatal period (P00-P96)	72	0.7	5.4
...	All other causes (residual)	1,962	19.1	147.5
Non-Hispanic, American Indian or Alaska Native, male, 1-4 years				
...	All causes	38	100.0	46.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	36.8	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	10.5	*
3	Malignant neoplasms (C00-C97)	2	5.3	*
3	Influenza and pneumonia (J09-J18)	2	5.3	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	5.3	*
6	Meningococcal infection (A39)	1	2.6	*
6	Septicemia (A40-A41)	1	2.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.6	*
6	Cerebrovascular diseases (I60-I69)	1	2.6	*
...	All other causes (residual)	10	26.3	*
Non-Hispanic, American Indian or Alaska Native, male, 5-9 years				
...	All causes	25	100.0	24.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	52.0	*
2	Malignant neoplasms (C00-C97)	1	4.0	*
2	Meningitis (G00,G03)	1	4.0	*
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	4.0	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.0	*
2	Complications of medical and surgical care (Y40-Y84,Y88)	1	4.0	*
...	All other causes (residual)	7	28.0	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 10-14 years				
...	All causes	24	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	33.3	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	29.2	*
3	Malignant neoplasms (C00-C97)	2	8.3	*
4	Diabetes mellitus (E10-E14)	1	4.2	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	4.2	*
...	All other causes (residual)	5	20.8	*
Non-Hispanic, American Indian or Alaska Native, male, 15-19 years				
...	All causes	131	100.0	122.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	41.2	50.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	30.5	37.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	9.2	*
4	Malignant neoplasms (C00-C97)	4	3.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.5	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
6	Influenza and pneumonia (J09-J18)	1	0.8	*
6	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.8	*
6	Legal intervention (Y35,Y89.0)	1	0.8	*
...	All other causes (residual)	14	10.7	*
Non-Hispanic, American Indian or Alaska Native, male, 20-24 years				
...	All causes	273	100.0	239.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	48.4	116.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	27.1	65.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	11.0	26.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.5	*
5	Diabetes mellitus (E10-E14)	3	1.1	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.1	*
7	Malignant neoplasms (C00-C97)	2	0.7	*
7	Chronic liver disease and cirrhosis (K70,K73-K74)	2	0.7	*
9	Cerebrovascular diseases (I60-I69)	1	0.4	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
9	Legal intervention (Y35,Y89.0)	1	0.4	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	19	7.0	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 25-29 years				
...	All causes	362	100.0	332.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	140	38.7	128.4
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	74	20.4	67.9
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	9.1	30.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	20	5.5	18.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	18	5.0	*
6	Malignant neoplasms (C00-C97)	6	1.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.7	*
8	Diabetes mellitus (E10-E14)	5	1.4	*
8	Legal intervention (Y35,Y89.0)	5	1.4	*
10	Cerebrovascular diseases (I60-I69)	3	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.6	*
12	Septicemia (A40-A41)	1	0.3	*
12	Viral hepatitis (B15-B19)	1	0.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
12	Hernia (K40-K46)	1	0.3	*
...	All other causes (residual)	45	12.4	41.3
Non-Hispanic, American Indian or Alaska Native, male, 30-34 years				
...	All causes	367	100.0	398.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	124	33.8	134.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	59	16.1	64.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	39	10.6	42.3
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	31	8.4	33.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	5.4	21.7
6	Diabetes mellitus (E10-E14)	12	3.3	*
7	Malignant neoplasms (C00-C97)	7	1.9	*
8	Cerebrovascular diseases (I60-I69)	5	1.4	*
9	Influenza and pneumonia (J09-J18)	4	1.1	*
9	Legal intervention (Y35,Y89.0)	4	1.1	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
13	Tuberculosis (A16-A19)	1	0.3	*
13	Septicemia (A40-A41)	1	0.3	*
13	Viral hepatitis (B15-B19)	1	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.3	*
13	Aortic aneurysm and dissection (I71)	1	0.3	*
13	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	48	13.1	52.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 35-39 years				
...	All causes	397	100.0	474.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	123	31.0	147.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	48	12.1	57.4
3	Chronic liver disease and cirrhosis (K70,K73-K74)	46	11.6	55.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	9.6	45.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	6.5	31.1
6	Diabetes mellitus (E10-E14)	19	4.8	*
7	Malignant neoplasms (C00-C97)	18	4.5	*
8	Septicemia (A40-A41)	5	1.3	*
9	Cerebrovascular diseases (I60-I69)	4	1.0	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.0	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3	0.8	*
11	Influenza and pneumonia (J09-J18)	3	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
14	Viral hepatitis (B15-B19)	1	0.3	*
14	Atherosclerosis (I70)	1	0.3	*
14	Legal intervention (Y35,Y89.0)	1	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	54	13.6	64.6
Non-Hispanic, American Indian or Alaska Native, male, 40-44 years				
...	All causes	428	100.0	557.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	27.6	153.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	69	16.1	89.9
3	Chronic liver disease and cirrhosis (K70,K73-K74)	50	11.7	65.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	6.3	35.2
5	Malignant neoplasms (C00-C97)	26	6.1	33.9
6	Diabetes mellitus (E10-E14)	20	4.7	26.0
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15	3.5	*
8	Human immunodeficiency virus (HIV) disease (B20-B24)	6	1.4	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6	1.4	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	1.2	*
10	Cerebrovascular diseases (I60-I69)	5	1.2	*
10	Influenza and pneumonia (J09-J18)	5	1.2	*
13	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
13	Pneumonitis due to solids and liquids (J69)	3	0.7	*
15	Legal intervention (Y35,Y89.0)	2	0.5	*
...	All other causes (residual)	68	15.9	88.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 45-49 years				
...	All causes	546	100.0	686.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	119	21.8	149.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	15.0	103.2
3	Chronic liver disease and cirrhosis (K70,K73-K74)	71	13.0	89.3
4	Malignant neoplasms (C00-C97)	47	8.6	59.1
5	Diabetes mellitus (E10-E14)	29	5.3	36.5
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	3.8	26.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	3.8	26.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	2.0	*
9	Influenza and pneumonia (J09-J18)	10	1.8	*
10	Septicemia (A40-A41)	8	1.5	*
11	Viral hepatitis (B15-B19)	6	1.1	*
11	Cerebrovascular diseases (I60-I69)	6	1.1	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.7	*
13	Aortic aneurysm and dissection (I71)	4	0.7	*
13	Chronic lower respiratory diseases (J40-J47)	4	0.7	*
...	All other causes (residual)	103	18.9	129.6
Non-Hispanic, American Indian or Alaska Native, male, 50-54 years				
...	All causes	817	100.0	967.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	160	19.6	189.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	149	18.2	176.4
3	Malignant neoplasms (C00-C97)	98	12.0	116.0
4	Chronic liver disease and cirrhosis (K70,K73-K74)	85	10.4	100.6
5	Diabetes mellitus (E10-E14)	50	6.1	59.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	3.7	35.5
7	Cerebrovascular diseases (I60-I69)	17	2.1	*
8	Influenza and pneumonia (J09-J18)	15	1.8	*
9	Chronic lower respiratory diseases (J40-J47)	14	1.7	*
10	Septicemia (A40-A41)	11	1.3	*
10	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.2	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.7	*
14	Viral hepatitis (B15-B19)	5	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.6	*
14	Aortic aneurysm and dissection (I71)	5	0.6	*
...	All other causes (residual)	146	17.9	172.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 55-59 years				
...	All causes	1,028	100.0	1,262.8
1	Malignant neoplasms (C00-C97)	237	23.1	291.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	184	17.9	226.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	12.5	158.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	87	8.5	106.9
5	Diabetes mellitus (E10-E14)	63	6.1	77.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29	2.8	35.6
7	Chronic lower respiratory diseases (J40-J47)	26	2.5	31.9
8	Cerebrovascular diseases (I60-I69)	25	2.4	30.7
9	Influenza and pneumonia (J09-J18)	18	1.8	*
10	Viral hepatitis (B15-B19)	15	1.5	*
11	Septicemia (A40-A41)	12	1.2	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.2	*
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	0.7	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6	0.6	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.5	*
...	All other causes (residual)	173	16.8	212.5
Non-Hispanic, American Indian or Alaska Native, male, 60-64 years				
...	All causes	1,037	100.0	1,565.1
1	Malignant neoplasms (C00-C97)	238	23.0	359.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	209	20.2	315.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	93	9.0	140.4
4	Diabetes mellitus (E10-E14)	78	7.5	117.7
5	Chronic liver disease and cirrhosis (K70,K73-K74)	65	6.3	98.1
6	Chronic lower respiratory diseases (J40-J47)	50	4.8	75.5
7	Cerebrovascular diseases (I60-I69)	25	2.4	37.7
8	Influenza and pneumonia (J09-J18)	24	2.3	36.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.4	*
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	1.4	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	1.2	*
12	Septicemia (A40-A41)	11	1.1	*
12	Viral hepatitis (B15-B19)	11	1.1	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	0.8	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.4	*
15	Parkinson's disease (G20-G21)	4	0.4	*
15	Pneumonitis due to solids and liquids (J69)	4	0.4	*
15	Peptic ulcer (K25-K28)	4	0.4	*
...	All other causes (residual)	167	16.1	252.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 65-69 years				
...	All causes	1,052	100.0	1,991.8
1	Malignant neoplasms (C00-C97)	281	26.7	532.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	223	21.2	422.2
3	Diabetes mellitus (E10-E14)	86	8.2	162.8
4	Chronic lower respiratory diseases (J40-J47)	71	6.7	134.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	54	5.1	102.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	44	4.2	83.3
7	Cerebrovascular diseases (I60-I69)	37	3.5	70.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	22	2.1	41.7
9	Septicemia (A40-A41)	17	1.6	*
9	Influenza and pneumonia (J09-J18)	17	1.6	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.7	*
12	Parkinson's disease (G20-G21)	6	0.6	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	6	0.6	*
14	Anemias (D50-D64)	5	0.5	*
14	Nutritional deficiencies (E40-E64)	5	0.5	*
14	Pneumonitis due to solids and liquids (J69)	5	0.5	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.5	*
...	All other causes (residual)	161	15.3	304.8
Non-Hispanic, American Indian or Alaska Native, male, 70-74 years				
...	All causes	995	100.0	3,013.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	261	26.2	790.5
2	Malignant neoplasms (C00-C97)	259	26.0	784.5
3	Diabetes mellitus (E10-E14)	75	7.5	227.2
4	Chronic lower respiratory diseases (J40-J47)	64	6.4	193.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33	3.3	100.0
6	Cerebrovascular diseases (I60-I69)	28	2.8	84.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	26	2.6	78.7
8	Influenza and pneumonia (J09-J18)	20	2.0	60.6
9	Chronic liver disease and cirrhosis (K70,K73-K74)	16	1.6	*
10	Septicemia (A40-A41)	14	1.4	*
11	Alzheimer's disease (G30)	11	1.1	*
12	Parkinson's disease (G20-G21)	10	1.0	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	0.9	*
14	Pneumonitis due to solids and liquids (J69)	8	0.8	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.5	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	5	0.5	*
...	All other causes (residual)	151	15.2	457.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 75-79 years				
...	All causes	869	100.0	4,147.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	210	24.2	1,002.2
2	Malignant neoplasms (C00-C97)	209	24.1	997.5
3	Chronic lower respiratory diseases (J40-J47)	66	7.6	315.0
4	Diabetes mellitus (E10-E14)	52	6.0	248.2
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	4.6	190.9
6	Cerebrovascular diseases (I60-I69)	39	4.5	186.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	21	2.4	100.2
8	Alzheimer's disease (G30)	20	2.3	95.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	2.3	95.5
10	Septicemia (A40-A41)	15	1.7	*
10	Influenza and pneumonia (J09-J18)	15	1.7	*
12	Parkinson's disease (G20-G21)	9	1.0	*
13	Pneumonitis due to solids and liquids (J69)	7	0.8	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.6	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	0.6	*
...	All other causes (residual)	136	15.7	649.1
Non-Hispanic, American Indian or Alaska Native, male, 80-84 years				
...	All causes	758	100.0	6,484.7
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	199	26.3	1,702.5
2	Malignant neoplasms (C00-C97)	154	20.3	1,317.5
3	Chronic lower respiratory diseases (J40-J47)	65	8.6	556.1
4	Diabetes mellitus (E10-E14)	40	5.3	342.2
5	Cerebrovascular diseases (I60-I69)	38	5.0	325.1
6	Alzheimer's disease (G30)	22	2.9	188.2
7	Parkinson's disease (G20-G21)	19	2.5	*
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	2.5	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15	2.0	*
10	Influenza and pneumonia (J09-J18)	14	1.8	*
11	Septicemia (A40-A41)	13	1.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	1.5	*
13	Pneumonitis due to solids and liquids (J69)	9	1.2	*
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	5	0.7	*
15	Pneumoconioses and chemical effects (J60-J66,J68)	4	0.5	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.5	*
...	All other causes (residual)	127	16.8	1,086.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 85-89 years				
...	All causes	561	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	159	28.3	---
2	Malignant neoplasms (C00-C97)	86	15.3	---
3	Chronic lower respiratory diseases (J40-J47)	39	7.0	---
4	Alzheimer's disease (G30)	36	6.4	---
5	Cerebrovascular diseases (I60-I69)	29	5.2	---
6	Diabetes mellitus (E10-E14)	27	4.8	---
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	3.0	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.9	---
9	Septicemia (A40-A41)	15	2.7	---
9	Influenza and pneumonia (J09-J18)	15	2.7	---
11	Parkinson's disease (G20-G21)	5	0.9	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.9	---
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	3	0.5	---
14	Nutritional deficiencies (E40-E64)	3	0.5	---
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.5	---
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.5	---
...	All other causes (residual)	96	17.1	---
Non-Hispanic, American Indian or Alaska Native, male, 90-94 years				
...	All causes	291	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	85	29.2	---
2	Malignant neoplasms (C00-C97)	39	13.4	---
3	Chronic lower respiratory diseases (J40-J47)	20	6.9	---
4	Cerebrovascular diseases (I60-I69)	18	6.2	---
5	Alzheimer's disease (G30)	15	5.2	---
6	Diabetes mellitus (E10-E14)	10	3.4	---
7	Influenza and pneumonia (J09-J18)	9	3.1	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	2.7	---
9	Pneumonitis due to solids and liquids (J69)	7	2.4	---
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	2.1	---
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	1.4	---
11	Parkinson's disease (G20-G21)	4	1.4	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	1.4	---
14	Septicemia (A40-A41)	2	0.7	---
14	Nutritional deficiencies (E40-E64)	2	0.7	---
14	Aortic aneurysm and dissection (I71)	2	0.7	---
...	All other causes (residual)	56	19.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 95-99 years				
...	All causes	83	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	20.5	---
2	Malignant neoplasms (C00-C97)	7	8.4	---
3	Alzheimer's disease (G30)	6	7.2	---
4	Cerebrovascular diseases (I60-I69)	5	6.0	---
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	6.0	---
6	Diabetes mellitus (E10-E14)	3	3.6	---
6	Influenza and pneumonia (J09-J18)	3	3.6	---
6	Pneumoconioses and chemical effects (J60-J66,J68)	3	3.6	---
6	Pneumonitis due to solids and liquids (J69)	3	3.6	---
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.4	---
10	Chronic lower respiratory diseases (J40-J47)	2	2.4	---
12	Peptic ulcer (K25-K28)	1	1.2	---
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	1.2	---
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.2	---
12	Hyperplasia of prostate (N40)	1	1.2	---
...	All other causes (residual)	23	27.7	---
Non-Hispanic, American Indian or Alaska Native, male, 100 years and over				
...	All causes	15	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	46.7	---
2	Cerebrovascular diseases (I60-I69)	2	13.3	---
3	Diabetes mellitus (E10-E14)	1	6.7	---
3	Nutritional deficiencies (E40-E64)	1	6.7	---
3	Pneumoconioses and chemical effects (J60-J66,J68)	1	6.7	---
3	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	6.7	---
...	All other causes (residual)	2	13.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, all ages				
...	All causes	8,315	100.0	602.2
1	Malignant neoplasms (C00-C97)	1,502	18.1	108.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,381	16.6	100.0
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	696	8.4	50.4
4	Diabetes mellitus (E10-E14)	504	6.1	36.5
5	Chronic liver disease and cirrhosis (K70,K73-K74)	456	5.5	33.0
6	Chronic lower respiratory diseases (J40-J47)	423	5.1	30.6
7	Cerebrovascular diseases (I60-I69)	344	4.1	24.9
8	Alzheimer's disease (G30)	256	3.1	18.5
9	Influenza and pneumonia (J09-J18)	184	2.2	13.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	177	2.1	12.8
11	Septicemia (A40-A41)	167	2.0	12.1
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	139	1.7	10.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	112	1.3	8.1
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	70	0.8	5.1
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	65	0.8	4.7
...	All other causes (residual)	1,839	22.1	133.2
Non-Hispanic, American Indian or Alaska Native, female, 1-4 years				
...	All causes	44	100.0	55.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	29.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	18.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	9.1	*
4	Meningococcal infection (A39)	1	2.3	*
4	Septicemia (A40-A41)	1	2.3	*
4	Malignant neoplasms (C00-C97)	1	2.3	*
4	Diabetes mellitus (E10-E14)	1	2.3	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.3	*
4	Influenza and pneumonia (J09-J18)	1	2.3	*
4	Certain conditions originating in the perinatal period (P00-P96)	1	2.3	*
...	All other causes (residual)	12	27.3	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 5-9 years				
...	All causes	17	100.0	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4	23.5	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	11.8	*
3	Meningococcal infection (A39)	1	5.9	*
3	Malignant neoplasms (C00-C97)	1	5.9	*
3	Meningitis (G00,G03)	1	5.9	*
3	Chronic lower respiratory diseases (J40-J47)	1	5.9	*
3	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	5.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	5.9	*
...	All other causes (residual)	5	29.4	*
Non-Hispanic, American Indian or Alaska Native, female, 10-14 years				
...	All causes	26	100.0	25.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6	23.1	*
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6	23.1	*
3	Malignant neoplasms (C00-C97)	4	15.4	*
4	Septicemia (A40-A41)	2	7.7	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	7.7	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	3.8	*
6	Cerebrovascular diseases (I60-I69)	1	3.8	*
...	All other causes (residual)	4	15.4	*
Non-Hispanic, American Indian or Alaska Native, female, 15-19 years				
...	All causes	56	100.0	53.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	27	48.2	26.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15	26.8	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	7.1	*
4	Pregnancy, childbirth and the puerperium (O00-O99)	2	3.6	*
5	Diabetes mellitus (E10-E14)	1	1.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	1.8	*
5	Influenza and pneumonia (J09-J18)	1	1.8	*
...	All other causes (residual)	5	8.9	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 20-24 years				
...	All causes	131	100.0	119.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	57	43.5	52.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	27	20.6	24.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	7.6	*
4	Malignant neoplasms (C00-C97)	4	3.1	*
4	Chronic liver disease and cirrhosis (K70,K73-K74)	4	3.1	*
6	Septicemia (A40-A41)	1	0.8	*
6	Cerebrovascular diseases (I60-I69)	1	0.8	*
6	Chronic lower respiratory diseases (J40-J47)	1	0.8	*
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.8	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.8	*
6	Legal intervention (Y35,Y89.0)	1	0.8	*
...	All other causes (residual)	23	17.6	21.0
Non-Hispanic, American Indian or Alaska Native, female, 25-29 years				
...	All causes	171	100.0	161.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	75	43.9	70.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	10.5	*
3	Chronic liver disease and cirrhosis (K70,K73-K74)	11	6.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	5.3	*
5	Malignant neoplasms (C00-C97)	8	4.7	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	4.7	*
7	Septicemia (A40-A41)	3	1.8	*
8	Diabetes mellitus (E10-E14)	2	1.2	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.6	*
9	Anemias (D50-D64)	1	0.6	*
9	Influenza and pneumonia (J09-J18)	1	0.6	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.6	*
9	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.6	*
9	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.6	*
9	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.6	*
...	All other causes (residual)	30	17.5	28.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 30-34 years				
...	All causes	203	100.0	216.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	61	30.0	65.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	29	14.3	31.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	9.4	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	6.9	*
5	Malignant neoplasms (C00-C97)	10	4.9	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	3.4	*
7	Diabetes mellitus (E10-E14)	6	3.0	*
8	Influenza and pneumonia (J09-J18)	3	1.5	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.5	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.5	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	1.0	*
11	Cerebrovascular diseases (I60-I69)	2	1.0	*
11	Chronic lower respiratory diseases (J40-J47)	2	1.0	*
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.0	*
15	Septicemia (A40-A41)	1	0.5	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.5	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
15	Anemias (D50-D64)	1	0.5	*
15	Legal intervention (Y35,Y89.0)	1	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.5	*
...	All other causes (residual)	34	16.7	36.3
Non-Hispanic, American Indian or Alaska Native, female, 35-39 years				
...	All causes	225	100.0	260.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	63	28.0	72.9
2	Chronic liver disease and cirrhosis (K70,K73-K74)	26	11.6	30.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	20	8.9	23.1
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	18	8.0	*
5	Malignant neoplasms (C00-C97)	17	7.6	*
6	Diabetes mellitus (E10-E14)	15	6.7	*
7	Influenza and pneumonia (J09-J18)	7	3.1	*
8	Cerebrovascular diseases (I60-I69)	5	2.2	*
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	2.2	*
10	Septicemia (A40-A41)	3	1.3	*
10	Pneumonitis due to solids and liquids (J69)	3	1.3	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.9	*
14	Viral hepatitis (B15-B19)	1	0.4	*
14	Nutritional deficiencies (E40-E64)	1	0.4	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	*
14	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
...	All other causes (residual)	33	14.7	38.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 40-44 years				
...	All causes	282	100.0	352.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	59	20.9	73.7
2	Chronic liver disease and cirrhosis (K70,K73-K74)	49	17.4	61.2
3	Malignant neoplasms (C00-C97)	41	14.5	51.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	33	11.7	41.2
5	Diabetes mellitus (E10-E14)	14	5.0	*
6	Cerebrovascular diseases (I60-I69)	10	3.5	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	3.5	*
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	9	3.2	*
9	Septicemia (A40-A41)	7	2.5	*
10	Influenza and pneumonia (J09-J18)	5	1.8	*
11	Viral hepatitis (B15-B19)	3	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
12	Pneumonitis due to solids and liquids (J69)	2	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.4	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.4	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	1	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	0.4	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.4	*
...	All other causes (residual)	30	10.6	37.5
Non-Hispanic, American Indian or Alaska Native, female, 45-49 years				
...	All causes	373	100.0	445.4
1	Chronic liver disease and cirrhosis (K70,K73-K74)	63	16.9	75.2
2	Malignant neoplasms (C00-C97)	58	15.5	69.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55	14.7	65.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	12.6	56.1
5	Diabetes mellitus (E10-E14)	19	5.1	*
6	Cerebrovascular diseases (I60-I69)	11	2.9	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	2.4	*
8	Influenza and pneumonia (J09-J18)	7	1.9	*
9	Septicemia (A40-A41)	6	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	6	1.6	*
11	Viral hepatitis (B15-B19)	4	1.1	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	1.1	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.8	*
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	0.5	*
...	All other causes (residual)	75	20.1	89.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 50-54 years				
...	All causes	556	100.0	609.8
1	Malignant neoplasms (C00-C97)	120	21.6	131.6
2	Chronic liver disease and cirrhosis (K70,K73-K74)	68	12.2	74.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	68	12.2	74.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	55	9.9	60.3
5	Diabetes mellitus (E10-E14)	34	6.1	37.3
6	Chronic lower respiratory diseases (J40-J47)	21	3.8	23.0
7	Cerebrovascular diseases (I60-I69)	15	2.7	*
8	Influenza and pneumonia (J09-J18)	10	1.8	*
9	Septicemia (A40-A41)	9	1.6	*
9	Viral hepatitis (B15-B19)	9	1.6	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	8	1.4	*
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8	1.4	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	1.3	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	4	0.7	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	0.7	*
...	All other causes (residual)	116	20.9	127.2
Non-Hispanic, American Indian or Alaska Native, female, 55-59 years				
...	All causes	619	100.0	677.3
1	Malignant neoplasms (C00-C97)	154	24.9	168.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	15.5	105.0
3	Chronic liver disease and cirrhosis (K70,K73-K74)	61	9.9	66.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	45	7.3	49.2
5	Diabetes mellitus (E10-E14)	31	5.0	33.9
6	Chronic lower respiratory diseases (J40-J47)	25	4.0	27.4
7	Septicemia (A40-A41)	16	2.6	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.6	*
9	Cerebrovascular diseases (I60-I69)	10	1.6	*
10	Viral hepatitis (B15-B19)	9	1.5	*
11	Influenza and pneumonia (J09-J18)	7	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.8	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.8	*
14	Pneumonitis due to solids and liquids (J69)	4	0.6	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.6	*
...	All other causes (residual)	131	21.2	143.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 60-64 years				
...	All causes	749	100.0	979.0
1	Malignant neoplasms (C00-C97)	197	26.3	257.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	133	17.8	173.8
3	Diabetes mellitus (E10-E14)	66	8.8	86.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	52	6.9	68.0
5	Chronic lower respiratory diseases (J40-J47)	49	6.5	64.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34	4.5	44.4
7	Cerebrovascular diseases (I60-I69)	28	3.7	36.6
8	Septicemia (A40-A41)	16	2.1	*
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	16	2.1	*
10	Viral hepatitis (B15-B19)	10	1.3	*
11	Influenza and pneumonia (J09-J18)	6	0.8	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.7	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	0.7	*
14	Alzheimer's disease (G30)	4	0.5	*
14	Peptic ulcer (K25-K28)	4	0.5	*
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4	0.5	*
...	All other causes (residual)	120	16.0	156.8
Non-Hispanic, American Indian or Alaska Native, female, 65-69 years				
...	All causes	772	100.0	1,293.6
1	Malignant neoplasms (C00-C97)	196	25.4	328.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	129	16.7	216.2
3	Diabetes mellitus (E10-E14)	60	7.8	100.5
4	Chronic lower respiratory diseases (J40-J47)	58	7.5	97.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	32	4.1	53.6
6	Cerebrovascular diseases (I60-I69)	30	3.9	50.3
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	23	3.0	38.5
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	3.0	38.5
9	Septicemia (A40-A41)	22	2.8	36.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13	1.7	*
11	Influenza and pneumonia (J09-J18)	11	1.4	*
12	Alzheimer's disease (G30)	9	1.2	*
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.8	*
14	Viral hepatitis (B15-B19)	5	0.6	*
15	Pneumonitis due to solids and liquids (J69)	4	0.5	*
...	All other causes (residual)	151	19.6	253.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 70-74 years				
...	All causes	818	100.0	2,117.5
1	Malignant neoplasms (C00-C97)	201	24.6	520.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	158	19.3	409.0
3	Chronic lower respiratory diseases (J40-J47)	82	10.0	212.3
4	Diabetes mellitus (E10-E14)	59	7.2	152.7
5	Septicemia (A40-A41)	27	3.3	69.9
6	Cerebrovascular diseases (I60-I69)	24	2.9	62.1
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	2.6	54.4
8	Chronic liver disease and cirrhosis (K70,K73-K74)	20	2.4	51.8
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	2.3	*
10	Alzheimer's disease (G30)	16	2.0	*
10	Influenza and pneumonia (J09-J18)	16	2.0	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	14	1.7	*
13	Anemias (D50-D64)	4	0.5	*
13	Aortic aneurysm and dissection (I71)	4	0.5	*
13	Pneumonitis due to solids and liquids (J69)	4	0.5	*
...	All other causes (residual)	149	18.2	385.7
Non-Hispanic, American Indian or Alaska Native, female, 75-79 years				
...	All causes	834	100.0	3,184.7
1	Malignant neoplasms (C00-C97)	194	23.3	740.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	156	18.7	595.7
3	Diabetes mellitus (E10-E14)	63	7.6	240.6
4	Chronic lower respiratory diseases (J40-J47)	61	7.3	232.9
5	Cerebrovascular diseases (I60-I69)	47	5.6	179.5
6	Alzheimer's disease (G30)	25	3.0	95.5
7	Influenza and pneumonia (J09-J18)	23	2.8	87.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	21	2.5	80.2
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	2.3	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.9	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	1.8	*
12	Septicemia (A40-A41)	14	1.7	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	9	1.1	*
14	Parkinson's disease (G20-G21)	8	1.0	*
15	Pneumonitis due to solids and liquids (J69)	7	0.8	*
...	All other causes (residual)	156	18.7	595.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 80-84 years				
...	All causes	871	100.0	5,243.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	194	22.3	1,167.9
2	Malignant neoplasms (C00-C97)	147	16.9	885.0
3	Diabetes mellitus (E10-E14)	67	7.7	403.3
4	Alzheimer's disease (G30)	54	6.2	325.1
4	Cerebrovascular diseases (I60-I69)	54	6.2	325.1
6	Chronic lower respiratory diseases (J40-J47)	50	5.7	301.0
7	Influenza and pneumonia (J09-J18)	29	3.3	174.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24	2.8	144.5
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	2.1	*
10	Septicemia (A40-A41)	16	1.8	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	1.8	*
12	Chronic liver disease and cirrhosis (K70,K73-K74)	14	1.6	*
13	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	7	0.8	*
14	Pneumonitis due to solids and liquids (J69)	6	0.7	*
15	Parkinson's disease (G20-G21)	5	0.6	*
...	All other causes (residual)	170	19.5	1,023.4
Non-Hispanic, American Indian or Alaska Native, female, 85-89 years				
...	All causes	720	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	150	20.8	---
2	Malignant neoplasms (C00-C97)	86	11.9	---
3	Alzheimer's disease (G30)	65	9.0	---
4	Cerebrovascular diseases (I60-I69)	50	6.9	---
5	Diabetes mellitus (E10-E14)	43	6.0	---
6	Chronic lower respiratory diseases (J40-J47)	42	5.8	---
7	Influenza and pneumonia (J09-J18)	33	4.6	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	22	3.1	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	18	2.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.9	---
11	Pneumonitis due to solids and liquids (J69)	12	1.7	---
12	Septicemia (A40-A41)	6	0.8	---
12	Chronic liver disease and cirrhosis (K70,K73-K74)	6	0.8	---
14	Nutritional deficiencies (E40-E64)	5	0.7	---
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.6	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	4	0.6	---
...	All other causes (residual)	160	22.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 90-94 years				
...	All causes	488	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	121	24.8	---
2	Alzheimer's disease (G30)	53	10.9	---
3	Malignant neoplasms (C00-C97)	49	10.0	---
4	Cerebrovascular diseases (I60-I69)	37	7.6	---
5	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21	4.3	---
6	Chronic lower respiratory diseases (J40-J47)	18	3.7	---
7	Diabetes mellitus (E10-E14)	14	2.9	---
8	Influenza and pneumonia (J09-J18)	13	2.7	---
9	Septicemia (A40-A41)	12	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12	2.5	---
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	9	1.8	---
12	Nutritional deficiencies (E40-E64)	4	0.8	---
12	Parkinson's disease (G20-G21)	4	0.8	---
12	Atherosclerosis (I70)	4	0.8	---
15	Peptic ulcer (K25-K28)	3	0.6	---
...	All other causes (residual)	114	23.4	---
Non-Hispanic, American Indian or Alaska Native, female, 95-99 years				
...	All causes	186	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	49	26.3	---
2	Alzheimer's disease (G30)	29	15.6	---
3	Cerebrovascular diseases (I60-I69)	14	7.5	---
4	Malignant neoplasms (C00-C97)	10	5.4	---
5	Diabetes mellitus (E10-E14)	9	4.8	---
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	4.3	---
7	Influenza and pneumonia (J09-J18)	7	3.8	---
8	Septicemia (A40-A41)	3	1.6	---
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	1.1	---
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	---
10	Anemias (D50-D64)	1	0.5	---
10	Nutritional deficiencies (E40-E64)	1	0.5	---
10	Parkinson's disease (G20-G21)	1	0.5	---
10	Chronic lower respiratory diseases (J40-J47)	1	0.5	---
10	Pneumonitis due to solids and liquids (J69)	1	0.5	---
10	Diseases of appendix (K35-K38)	1	0.5	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.5	---
...	All other causes (residual)	47	25.3	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 100 years and over				
...	All causes	45	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	13	28.9	---
2	Cerebrovascular diseases (I60-I69)	5	11.1	---
3	Malignant neoplasms (C00-C97)	3	6.7	---
3	Influenza and pneumonia (J09-J18)	3	6.7	---
3	Chronic lower respiratory diseases (J40-J47)	3	6.7	---
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	4.4	---
7	Alzheimer's disease (G30)	1	2.2	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	2.2	---
7	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	2.2	---
...	All other causes (residual)	13	28.9	---
Non-Hispanic, Asian or Pacific Islander, both sexes, all ages				
...	All causes	68,235	100.0	350.3
1	Malignant neoplasms (C00-C97)	17,718	26.0	91.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,519	21.3	74.5
3	Cerebrovascular diseases (I60-I69)	5,217	7.6	26.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,101	4.5	15.9
5	Diabetes mellitus (E10-E14)	2,697	4.0	13.8
6	Alzheimer's disease (G30)	2,363	3.5	12.1
7	Influenza and pneumonia (J09-J18)	2,132	3.1	10.9
8	Chronic lower respiratory diseases (J40-J47)	1,914	2.8	9.8
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,363	2.0	7.0
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,348	2.0	6.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,280	1.9	6.6
12	Septicemia (A40-A41)	786	1.2	4.0
13	Parkinson's disease (G20-G21)	749	1.1	3.8
14	Chronic liver disease and cirrhosis (K70,K73-K74)	632	0.9	3.2
15	Certain conditions originating in the perinatal period (P00-P96)	527	0.8	2.7
...	All other causes (residual)	11,889	17.4	61.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	160	100.0	17.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	43	26.9	4.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	13.1	2.2
3	Malignant neoplasms (C00-C97)	20	12.5	2.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	6.3	*
5	Septicemia (A40-A41)	5	3.1	*
6	Cerebrovascular diseases (I60-I69)	4	2.5	*
7	Influenza and pneumonia (J09-J18)	3	1.9	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	1.3	*
9	Diabetes mellitus (E10-E14)	2	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	2	1.3	*
12	Meningitis (G00,G03)	1	0.6	*
12	Pneumonitis due to solids and liquids (J69)	1	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (residual)	40	25.0	4.3
Non-Hispanic, Asian or Pacific Islander, both sexes, 5-9 years				
...	All causes	97	100.0	8.3
1	Malignant neoplasms (C00-C97)	24	24.7	2.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	19.6	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	7.2	*
4	Influenza and pneumonia (J09-J18)	5	5.2	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	4.1	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	3	3.1	*
6	Cerebrovascular diseases (I60-I69)	3	3.1	*
8	Septicemia (A40-A41)	2	2.1	*
9	Scarlet fever and erysipelas (A38,A46)	1	1.0	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
9	Diabetes mellitus (E10-E14)	1	1.0	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (residual)	26	26.8	2.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 10-14 years				
...	All causes	121	100.0	10.2
1	Malignant neoplasms (C00-C97)	36	29.8	3.0
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	14.9	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14	11.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	8.3	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	4.1	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	5	4.1	*
7	Septicemia (A40-A41)	3	2.5	*
7	Cerebrovascular diseases (I60-I69)	3	2.5	*
7	Certain conditions originating in the perinatal period (P00-P96)	3	2.5	*
10	Influenza and pneumonia (J09-J18)	2	1.7	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.8	*
11	Meningitis (G00,G03)	1	0.8	*
11	Diseases of appendix (K35-K38)	1	0.8	*
...	All other causes (residual)	19	15.7	*
Non-Hispanic, Asian or Pacific Islander, both sexes, 15-19 years				
...	All causes	349	100.0	29.2
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	31.8	9.3
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	27.2	7.9
3	Malignant neoplasms (C00-C97)	28	8.0	2.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	28	8.0	2.3
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	4.6	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	8	2.3	*
7	Cerebrovascular diseases (I60-I69)	4	1.1	*
7	Chronic lower respiratory diseases (J40-J47)	4	1.1	*
9	Septicemia (A40-A41)	2	0.6	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
10	Anemias (D50-D64)	1	0.3	*
10	Diabetes mellitus (E10-E14)	1	0.3	*
10	Aortic aneurysm and dissection (I71)	1	0.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
10	Legal intervention (Y35,Y89.0)	1	0.3	*
...	All other causes (residual)	47	13.5	3.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 20-24 years				
...	All causes	622	100.0	44.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	212	34.1	15.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	194	31.2	13.8
3	Malignant neoplasms (C00-C97)	45	7.2	3.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	6.4	2.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	4.0	1.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	6	1.0	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	1.0	*
8	Cerebrovascular diseases (I60-I69)	4	0.6	*
9	Influenza and pneumonia (J09-J18)	3	0.5	*
9	Chronic lower respiratory diseases (J40-J47)	3	0.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
11	Diabetes mellitus (E10-E14)	2	0.3	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.3	*
11	Aortic aneurysm and dissection (I71)	2	0.3	*
15	Tuberculosis (A16-A19)	1	0.2	*
15	Septicemia (A40-A41)	1	0.2	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
15	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.2	*
...	All other causes (residual)	69	11.1	4.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 25-29 years				
...	All causes	626	100.0	37.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	198	31.6	11.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	141	22.5	8.5
3	Malignant neoplasms (C00-C97)	74	11.8	4.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	40	6.4	2.4
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	32	5.1	1.9
6	Pregnancy, childbirth and the puerperium (O00-O99)	9	1.4	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	1.1	*
8	Septicemia (A40-A41)	6	1.0	*
8	Influenza and pneumonia (J09-J18)	6	1.0	*
10	Diabetes mellitus (E10-E14)	4	0.6	*
10	Cerebrovascular diseases (I60-I69)	4	0.6	*
10	Chronic lower respiratory diseases (J40-J47)	4	0.6	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	4	0.6	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2	0.3	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.3	*
14	Aortic aneurysm and dissection (I71)	2	0.3	*
14	Pneumonitis due to solids and liquids (J69)	2	0.3	*
14	Peptic ulcer (K25-K28)	2	0.3	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
14	Legal intervention (Y35,Y89.0)	2	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	81	12.9	4.9
Non-Hispanic, Asian or Pacific Islander, both sexes, 30-34 years				
...	All causes	775	100.0	46.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	174	22.5	10.4
2	Malignant neoplasms (C00-C97)	125	16.1	7.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	123	15.9	7.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	66	8.5	4.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	5.8	2.7
6	Cerebrovascular diseases (I60-I69)	31	4.0	1.9
7	Chronic liver disease and cirrhosis (K70,K73-K74)	19	2.5	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	14	1.8	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	8	1.0	*
10	Septicemia (A40-A41)	7	0.9	*
10	Diabetes mellitus (E10-E14)	7	0.9	*
10	Influenza and pneumonia (J09-J18)	7	0.9	*
13	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.8	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
...	All other causes (residual)	132	17.0	7.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 35-39 years				
...	All causes	993	100.0	64.0
1	Malignant neoplasms (C00-C97)	276	27.8	17.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	153	15.4	9.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	133	13.4	8.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	117	11.8	7.5
5	Cerebrovascular diseases (I60-I69)	42	4.2	2.7
6	Chronic liver disease and cirrhosis (K70,K73-K74)	25	2.5	1.6
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	24	2.4	1.5
8	Diabetes mellitus (E10-E14)	16	1.6	*
9	Influenza and pneumonia (J09-J18)	13	1.3	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	13	1.3	*
11	Septicemia (A40-A41)	11	1.1	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.7	*
12	Aortic aneurysm and dissection (I71)	7	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.7	*
...	All other causes (residual)	142	14.3	9.2
Non-Hispanic, Asian or Pacific Islander, both sexes, 40-44 years				
...	All causes	1,310	100.0	86.6
1	Malignant neoplasms (C00-C97)	417	31.8	27.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	222	16.9	14.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	11.6	10.0
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	111	8.5	7.3
5	Cerebrovascular diseases (I60-I69)	81	6.2	5.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	35	2.7	2.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	32	2.4	2.1
8	Diabetes mellitus (E10-E14)	28	2.1	1.9
9	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.9	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.9	*
11	Influenza and pneumonia (J09-J18)	10	0.8	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.7	*
13	Viral hepatitis (B15-B19)	7	0.5	*
13	Aortic aneurysm and dissection (I71)	7	0.5	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.5	*
...	All other causes (residual)	168	12.8	11.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 45-49 years				
...	All causes	1,814	100.0	131.5
1	Malignant neoplasms (C00-C97)	669	36.9	48.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	309	17.0	22.4
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	156	8.6	11.3
4	Cerebrovascular diseases (I60-I69)	115	6.3	8.3
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	99	5.5	7.2
6	Diabetes mellitus (E10-E14)	66	3.6	4.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	43	2.4	3.1
8	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	26	1.4	1.9
9	Influenza and pneumonia (J09-J18)	22	1.2	1.6
10	Septicemia (A40-A41)	19	1.0	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	0.6	*
15	Aortic aneurysm and dissection (I71)	10	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	10	0.6	*
...	All other causes (residual)	220	12.1	15.9
Non-Hispanic, Asian or Pacific Islander, both sexes, 50-54 years				
...	All causes	2,552	100.0	206.0
1	Malignant neoplasms (C00-C97)	1,028	40.3	83.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	413	16.2	33.3
3	Cerebrovascular diseases (I60-I69)	171	6.7	13.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	160	6.3	12.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	105	4.1	8.5
6	Diabetes mellitus (E10-E14)	95	3.7	7.7
7	Chronic liver disease and cirrhosis (K70,K73-K74)	67	2.6	5.4
8	Influenza and pneumonia (J09-J18)	43	1.7	3.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	34	1.3	2.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	32	1.3	2.6
11	Septicemia (A40-A41)	30	1.2	2.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	0.8	1.7
13	Viral hepatitis (B15-B19)	20	0.8	1.6
14	Chronic lower respiratory diseases (J40-J47)	18	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.5	*
15	Aortic aneurysm and dissection (I71)	12	0.5	*
...	All other causes (residual)	291	11.4	23.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 55-59 years				
...	All causes	3,606	100.0	321.9
1	Malignant neoplasms (C00-C97)	1,494	41.4	133.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	709	19.7	63.3
3	Cerebrovascular diseases (I60-I69)	210	5.8	18.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	4.8	15.4
5	Diabetes mellitus (E10-E14)	159	4.4	14.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	86	2.4	7.7
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	64	1.8	5.7
8	Chronic liver disease and cirrhosis (K70,K73-K74)	55	1.5	4.9
9	Influenza and pneumonia (J09-J18)	48	1.3	4.3
10	Chronic lower respiratory diseases (J40-J47)	47	1.3	4.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	44	1.2	3.9
12	Septicemia (A40-A41)	38	1.1	3.4
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	33	0.9	2.9
14	Viral hepatitis (B15-B19)	31	0.9	2.8
15	Aortic aneurysm and dissection (I71)	14	0.4	*
...	All other causes (residual)	402	11.1	35.9
Non-Hispanic, Asian or Pacific Islander, both sexes, 60-64 years				
...	All causes	4,902	100.0	500.2
1	Malignant neoplasms (C00-C97)	2,082	42.5	212.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	917	18.7	93.6
3	Cerebrovascular diseases (I60-I69)	304	6.2	31.0
4	Diabetes mellitus (E10-E14)	238	4.9	24.3
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169	3.4	17.2
6	Influenza and pneumonia (J09-J18)	90	1.8	9.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	84	1.7	8.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	84	1.7	8.6
9	Chronic lower respiratory diseases (J40-J47)	81	1.7	8.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	68	1.4	6.9
11	Septicemia (A40-A41)	56	1.1	5.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	47	1.0	4.8
13	Viral hepatitis (B15-B19)	40	0.8	4.1
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	34	0.7	3.5
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	0.5	2.6
...	All other causes (residual)	583	11.9	59.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 65-69 years				
...	All causes	5,869	100.0	729.1
1	Malignant neoplasms (C00-C97)	2,257	38.5	280.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,177	20.1	146.2
3	Cerebrovascular diseases (I60-I69)	391	6.7	48.6
4	Diabetes mellitus (E10-E14)	285	4.9	35.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	195	3.3	24.2
6	Influenza and pneumonia (J09-J18)	130	2.2	16.1
6	Chronic lower respiratory diseases (J40-J47)	130	2.2	16.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	118	2.0	14.7
9	Septicemia (A40-A41)	83	1.4	10.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	1.3	9.3
11	Chronic liver disease and cirrhosis (K70,K73-K74)	72	1.2	8.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	58	1.0	7.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	46	0.8	5.7
14	Viral hepatitis (B15-B19)	40	0.7	5.0
14	Parkinson's disease (G20-G21)	40	0.7	5.0
...	All other causes (residual)	772	13.2	95.9
Non-Hispanic, Asian or Pacific Islander, both sexes, 70-74 years				
...	All causes	6,156	100.0	1,137.0
1	Malignant neoplasms (C00-C97)	2,098	34.1	387.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,324	21.5	244.5
3	Cerebrovascular diseases (I60-I69)	460	7.5	85.0
4	Diabetes mellitus (E10-E14)	318	5.2	58.7
5	Chronic lower respiratory diseases (J40-J47)	190	3.1	35.1
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	3.1	35.1
7	Influenza and pneumonia (J09-J18)	150	2.4	27.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	136	2.2	25.1
9	Septicemia (A40-A41)	92	1.5	17.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	81	1.3	15.0
11	Parkinson's disease (G20-G21)	75	1.2	13.9
12	Chronic liver disease and cirrhosis (K70,K73-K74)	63	1.0	11.6
13	Alzheimer's disease (G30)	49	0.8	9.1
14	Aortic aneurysm and dissection (I71)	40	0.6	7.4
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	0.6	7.2
...	All other causes (residual)	851	13.8	157.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 75-79 years				
...	All causes	7,444	100.0	1,946.1
1	Malignant neoplasms (C00-C97)	2,216	29.8	579.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,568	21.1	409.9
3	Cerebrovascular diseases (I60-I69)	601	8.1	157.1
4	Diabetes mellitus (E10-E14)	374	5.0	97.8
5	Chronic lower respiratory diseases (J40-J47)	238	3.2	62.2
6	Influenza and pneumonia (J09-J18)	232	3.1	60.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	220	3.0	57.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	201	2.7	52.5
9	Alzheimer's disease (G30)	184	2.5	48.1
10	Parkinson's disease (G20-G21)	138	1.9	36.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	133	1.8	34.8
12	Septicemia (A40-A41)	111	1.5	29.0
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	60	0.8	15.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	54	0.7	14.1
15	Pneumonitis due to solids and liquids (J69)	46	0.6	12.0
...	All other causes (residual)	1,068	14.3	279.2
Non-Hispanic, Asian or Pacific Islander, both sexes, 80-84 years				
...	All causes	8,830	100.0	3,510.7
1	Malignant neoplasms (C00-C97)	2,102	23.8	835.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,033	23.0	808.3
3	Cerebrovascular diseases (I60-I69)	793	9.0	315.3
4	Diabetes mellitus (E10-E14)	381	4.3	151.5
5	Chronic lower respiratory diseases (J40-J47)	372	4.2	147.9
6	Alzheimer's disease (G30)	340	3.9	135.2
7	Influenza and pneumonia (J09-J18)	290	3.3	115.3
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	241	2.7	95.8
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	225	2.5	89.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	206	2.3	81.9
11	Parkinson's disease (G20-G21)	189	2.1	75.1
12	Septicemia (A40-A41)	93	1.1	37.0
13	Pneumonitis due to solids and liquids (J69)	85	1.0	33.8
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	63	0.7	25.0
15	Chronic liver disease and cirrhosis (K70,K73-K74)	59	0.7	23.5
...	All other causes (residual)	1,358	15.4	539.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 85-89 years				
...	All causes	9,535	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,411	25.3	---
2	Malignant neoplasms (C00-C97)	1,601	16.8	---
3	Cerebrovascular diseases (I60-I69)	917	9.6	---
4	Alzheimer's disease (G30)	598	6.3	---
5	Influenza and pneumonia (J09-J18)	430	4.5	---
6	Chronic lower respiratory diseases (J40-J47)	395	4.1	---
7	Diabetes mellitus (E10-E14)	374	3.9	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	259	2.7	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	253	2.7	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	233	2.4	---
11	Parkinson's disease (G20-G21)	171	1.8	---
12	Septicemia (A40-A41)	108	1.1	---
13	Pneumonitis due to solids and liquids (J69)	103	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	75	0.8	---
15	Aortic aneurysm and dissection (I71)	43	0.5	---
...	All other causes (residual)	1,564	16.4	---
Non-Hispanic, Asian or Pacific Islander, both sexes, 90-94 years				
...	All causes	7,439	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,939	26.1	---
2	Malignant neoplasms (C00-C97)	862	11.6	---
3	Cerebrovascular diseases (I60-I69)	691	9.3	---
4	Alzheimer's disease (G30)	667	9.0	---
5	Influenza and pneumonia (J09-J18)	406	5.5	---
6	Chronic lower respiratory diseases (J40-J47)	258	3.5	---
7	Diabetes mellitus (E10-E14)	254	3.4	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	240	3.2	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	2.4	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	162	2.2	---
11	Pneumonitis due to solids and liquids (J69)	100	1.3	---
12	Parkinson's disease (G20-G21)	90	1.2	---
13	Septicemia (A40-A41)	70	0.9	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	50	0.7	---
15	Nutritional deficiencies (E40-E64)	37	0.5	---
...	All other causes (residual)	1,432	19.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 95-99 years				
...	All causes	3,310	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	953	28.8	---
2	Alzheimer's disease (G30)	406	12.3	---
3	Cerebrovascular diseases (I60-I69)	318	9.6	---
4	Malignant neoplasms (C00-C97)	221	6.7	---
5	Influenza and pneumonia (J09-J18)	187	5.6	---
6	Chronic lower respiratory diseases (J40-J47)	121	3.7	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	95	2.9	---
8	Diabetes mellitus (E10-E14)	78	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	64	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	52	1.6	---
11	Pneumonitis due to solids and liquids (J69)	40	1.2	---
12	Septicemia (A40-A41)	30	0.9	---
13	Parkinson's disease (G20-G21)	21	0.6	---
14	Nutritional deficiencies (E40-E64)	20	0.6	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	15	0.5	---
...	All other causes (residual)	689	20.8	---
Non-Hispanic, Asian or Pacific Islander, both sexes, 100 years and over				
...	All causes	793	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	228	28.8	---
2	Alzheimer's disease (G30)	81	10.2	---
3	Cerebrovascular diseases (I60-I69)	63	7.9	---
4	Influenza and pneumonia (J09-J18)	52	6.6	---
5	Malignant neoplasms (C00-C97)	41	5.2	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	4.0	---
7	Chronic lower respiratory diseases (J40-J47)	25	3.2	---
8	Diabetes mellitus (E10-E14)	14	1.8	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	1.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13	1.6	---
11	Pneumonitis due to solids and liquids (J69)	8	1.0	---
12	Septicemia (A40-A41)	6	0.8	---
12	Atherosclerosis (I70)	6	0.8	---
14	Aortic aneurysm and dissection (I71)	5	0.6	---
15	Anemias (D50-D64)	4	0.5	---
15	Nutritional deficiencies (E40-E64)	4	0.5	---
...	All other causes (residual)	198	25.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, all ages				
...	All causes	34,892	100.0	374.9
1	Malignant neoplasms (C00-C97)	8,978	25.7	96.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,840	22.5	84.2
3	Cerebrovascular diseases (I60-I69)	2,268	6.5	24.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,929	5.5	20.7
5	Diabetes mellitus (E10-E14)	1,414	4.1	15.2
6	Chronic lower respiratory diseases (J40-J47)	1,134	3.3	12.2
7	Influenza and pneumonia (J09-J18)	1,120	3.2	12.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	977	2.8	10.5
9	Alzheimer's disease (G30)	709	2.0	7.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	665	1.9	7.1
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	545	1.6	5.9
12	Parkinson's disease (G20-G21)	440	1.3	4.7
13	Septicemia (A40-A41)	393	1.1	4.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	388	1.1	4.2
15	Certain conditions originating in the perinatal period (P00-P96)	306	0.9	3.3
...	All other causes (residual)	5,786	16.6	62.2
Non-Hispanic, Asian or Pacific Islander, male, 1-4 years				
...	All causes	80	100.0	16.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29	36.3	6.0
2	Malignant neoplasms (C00-C97)	9	11.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	11.3	*
4	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	2.5	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.5	*
4	Cerebrovascular diseases (I60-I69)	2	2.5	*
4	Influenza and pneumonia (J09-J18)	2	2.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.5	*
9	Septicemia (A40-A41)	1	1.3	*
9	Diabetes mellitus (E10-E14)	1	1.3	*
9	Chronic lower respiratory diseases (J40-J47)	1	1.3	*
9	Pneumonitis due to solids and liquids (J69)	1	1.3	*
9	Certain conditions originating in the perinatal period (P00-P96)	1	1.3	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	1	1.3	*
...	All other causes (residual)	17	21.3	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 5-9 years				
...	All causes	55	100.0	9.2
1	Malignant neoplasms (C00-C97)	15	27.3	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	25.5	*
3	Influenza and pneumonia (J09-J18)	3	5.5	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.5	*
5	Septicemia (A40-A41)	2	3.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.8	*
7	Diabetes mellitus (E10-E14)	1	1.8	*
7	Cerebrovascular diseases (I60-I69)	1	1.8	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.8	*
...	All other causes (residual)	11	20.0	*
Non-Hispanic, Asian or Pacific Islander, male, 10-14 years				
...	All causes	63	100.0	10.6
1	Malignant neoplasms (C00-C97)	17	27.0	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	17.5	*
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	12.7	*
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8	12.7	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	4.8	*
6	Cerebrovascular diseases (I60-I69)	2	3.2	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.2	*
8	Septicemia (A40-A41)	1	1.6	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.6	*
...	All other causes (residual)	10	15.9	*
Non-Hispanic, Asian or Pacific Islander, male, 15-19 years				
...	All causes	242	100.0	40.0
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	80	33.1	13.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	28.5	11.4
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	8.7	3.5
4	Malignant neoplasms (C00-C97)	16	6.6	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	9	3.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	2.1	*
7	Chronic lower respiratory diseases (J40-J47)	3	1.2	*
8	Cerebrovascular diseases (I60-I69)	2	0.8	*
9	Septicemia (A40-A41)	1	0.4	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
9	Anemias (D50-D64)	1	0.4	*
9	Aortic aneurysm and dissection (I71)	1	0.4	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.4	*
9	Legal intervention (Y35,Y89.0)	1	0.4	*
...	All other causes (residual)	31	12.8	5.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 20-24 years				
...	All causes	454	100.0	63.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	172	37.9	24.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	143	31.5	20.0
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	6.6	4.2
4	Malignant neoplasms (C00-C97)	28	6.2	3.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	17	3.7	*
6	Chronic lower respiratory diseases (J40-J47)	3	0.7	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.7	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
8	Diabetes mellitus (E10-E14)	2	0.4	*
8	Cerebrovascular diseases (I60-I69)	2	0.4	*
8	Aortic aneurysm and dissection (I71)	2	0.4	*
12	Septicemia (A40-A41)	1	0.2	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.2	*
12	Influenza and pneumonia (J09-J18)	1	0.2	*
12	Pneumonitis due to solids and liquids (J69)	1	0.2	*
12	Cholelithiasis and other disorders of gallbladder (K80-K82)	1	0.2	*
...	All other causes (residual)	45	9.9	6.3
Non-Hispanic, Asian or Pacific Islander, male, 25-29 years				
...	All causes	451	100.0	55.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	147	32.6	18.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	113	25.1	13.8
3	Malignant neoplasms (C00-C97)	41	9.1	5.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	30	6.7	3.7
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	25	5.5	3.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	1.1	*
7	Septicemia (A40-A41)	4	0.9	*
7	Influenza and pneumonia (J09-J18)	4	0.9	*
9	Diabetes mellitus (E10-E14)	3	0.7	*
9	Cerebrovascular diseases (I60-I69)	3	0.7	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
12	Pneumonitis due to solids and liquids (J69)	2	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
12	Legal intervention (Y35,Y89.0)	2	0.4	*
...	All other causes (residual)	63	14.0	7.7

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 30-34 years				
...	All causes	493	100.0	61.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127	25.8	15.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	83	16.8	10.4
3	Malignant neoplasms (C00-C97)	57	11.6	7.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	11.0	6.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	36	7.3	4.5
6	Cerebrovascular diseases (I60-I69)	15	3.0	*
6	Chronic liver disease and cirrhosis (K70,K73-K74)	15	3.0	*
8	Influenza and pneumonia (J09-J18)	6	1.2	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	5	1.0	*
9	Chronic lower respiratory diseases (J40-J47)	5	1.0	*
11	Diabetes mellitus (E10-E14)	4	0.8	*
12	Septicemia (A40-A41)	3	0.6	*
12	Aortic aneurysm and dissection (I71)	3	0.6	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	0.6	*
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	0.6	*
...	All other causes (residual)	74	15.0	9.3
Non-Hispanic, Asian or Pacific Islander, male, 35-39 years				
...	All causes	617	100.0	85.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	19.6	16.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	108	17.5	14.9
3	Malignant neoplasms (C00-C97)	103	16.7	14.2
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	14.1	12.0
5	Cerebrovascular diseases (I60-I69)	19	3.1	*
5	Chronic liver disease and cirrhosis (K70,K73-K74)	19	3.1	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	17	2.8	*
8	Diabetes mellitus (E10-E14)	12	1.9	*
9	Influenza and pneumonia (J09-J18)	10	1.6	*
10	Septicemia (A40-A41)	9	1.5	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	5	0.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	0.8	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.8	*
11	Aortic aneurysm and dissection (I71)	5	0.8	*
11	Chronic lower respiratory diseases (J40-J47)	5	0.8	*
...	All other causes (residual)	87	14.1	12.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 40-44 years				
...	All causes	809	100.0	114.0
1	Malignant neoplasms (C00-C97)	175	21.6	24.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	168	20.8	23.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	116	14.3	16.3
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	85	10.5	12.0
5	Cerebrovascular diseases (I60-I69)	53	6.6	7.5
6	Chronic liver disease and cirrhosis (K70,K73-K74)	29	3.6	4.1
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	2.3	*
8	Diabetes mellitus (E10-E14)	17	2.1	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	11	1.4	*
10	Viral hepatitis (B15-B19)	7	0.9	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.9	*
10	Aortic aneurysm and dissection (I71)	7	0.9	*
13	Influenza and pneumonia (J09-J18)	6	0.7	*
14	Septicemia (A40-A41)	5	0.6	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.6	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.6	*
...	All other causes (residual)	94	11.6	13.2
Non-Hispanic, Asian or Pacific Islander, male, 45-49 years				
...	All causes	1,080	100.0	167.4
1	Malignant neoplasms (C00-C97)	280	25.9	43.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	252	23.3	39.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	10.6	17.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	6.3	10.5
5	Cerebrovascular diseases (I60-I69)	59	5.5	9.1
6	Diabetes mellitus (E10-E14)	42	3.9	6.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	33	3.1	5.1
8	Influenza and pneumonia (J09-J18)	17	1.6	*
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	1.5	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	12	1.1	*
11	Septicemia (A40-A41)	10	0.9	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.8	*
13	Viral hepatitis (B15-B19)	8	0.7	*
13	Aortic aneurysm and dissection (I71)	8	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.6	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.6	*
...	All other causes (residual)	138	12.8	21.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 50-54 years				
...	All causes	1,513	100.0	261.7
1	Malignant neoplasms (C00-C97)	478	31.6	82.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	331	21.9	57.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	118	7.8	20.4
4	Cerebrovascular diseases (I60-I69)	92	6.1	15.9
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	68	4.5	11.8
6	Diabetes mellitus (E10-E14)	67	4.4	11.6
7	Chronic liver disease and cirrhosis (K70,K73-K74)	46	3.0	8.0
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	1.8	4.7
8	Influenza and pneumonia (J09-J18)	27	1.8	4.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20	1.3	3.5
11	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	19	1.3	*
12	Septicemia (A40-A41)	16	1.1	*
12	Viral hepatitis (B15-B19)	16	1.1	*
14	Chronic lower respiratory diseases (J40-J47)	11	0.7	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.7	*
...	All other causes (residual)	167	11.0	28.9
Non-Hispanic, Asian or Pacific Islander, male, 55-59 years				
...	All causes	2,161	100.0	419.9
1	Malignant neoplasms (C00-C97)	743	34.4	144.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	524	24.2	101.8
3	Cerebrovascular diseases (I60-I69)	121	5.6	23.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	114	5.3	22.2
5	Diabetes mellitus (E10-E14)	105	4.9	20.4
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	3.1	12.8
7	Chronic liver disease and cirrhosis (K70,K73-K74)	47	2.2	9.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	37	1.7	7.2
9	Chronic lower respiratory diseases (J40-J47)	30	1.4	5.8
10	Influenza and pneumonia (J09-J18)	28	1.3	5.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	27	1.2	5.2
12	Viral hepatitis (B15-B19)	23	1.1	4.5
13	Septicemia (A40-A41)	21	1.0	4.1
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	21	1.0	4.1
15	Aortic aneurysm and dissection (I71)	11	0.5	*
...	All other causes (residual)	243	11.2	47.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 60-64 years				
...	All causes	2,838	100.0	643.9
1	Malignant neoplasms (C00-C97)	1,061	37.4	240.7
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	659	23.2	149.5
3	Cerebrovascular diseases (I60-I69)	168	5.9	38.1
4	Diabetes mellitus (E10-E14)	153	5.4	34.7
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	3.5	22.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	56	2.0	12.7
7	Influenza and pneumonia (J09-J18)	54	1.9	12.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	54	1.9	12.3
9	Chronic lower respiratory diseases (J40-J47)	46	1.6	10.4
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41	1.4	9.3
11	Septicemia (A40-A41)	32	1.1	7.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	31	1.1	7.0
13	Viral hepatitis (B15-B19)	29	1.0	6.6
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	14	0.5	*
15	Aortic aneurysm and dissection (I71)	13	0.5	*
15	Pneumonitis due to solids and liquids (J69)	13	0.5	*
...	All other causes (residual)	314	11.1	71.2
Non-Hispanic, Asian or Pacific Islander, male, 65-69 years				
...	All causes	3,317	100.0	929.7
1	Malignant neoplasms (C00-C97)	1,192	35.9	334.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	752	22.7	210.8
3	Cerebrovascular diseases (I60-I69)	215	6.5	60.3
4	Diabetes mellitus (E10-E14)	166	5.0	46.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	100	3.0	28.0
6	Influenza and pneumonia (J09-J18)	83	2.5	23.3
7	Chronic lower respiratory diseases (J40-J47)	82	2.5	23.0
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	63	1.9	17.7
9	Chronic liver disease and cirrhosis (K70,K73-K74)	46	1.4	12.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	45	1.4	12.6
11	Septicemia (A40-A41)	39	1.2	10.9
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	39	1.2	10.9
13	Viral hepatitis (B15-B19)	26	0.8	7.3
14	Aortic aneurysm and dissection (I71)	24	0.7	6.7
15	Parkinson's disease (G20-G21)	21	0.6	5.9
...	All other causes (residual)	424	12.8	118.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 70-74 years				
...	All causes	3,476	100.0	1,437.5
1	Malignant neoplasms (C00-C97)	1,150	33.1	475.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	840	24.2	347.4
3	Cerebrovascular diseases (I60-I69)	225	6.5	93.0
4	Diabetes mellitus (E10-E14)	176	5.1	72.8
5	Chronic lower respiratory diseases (J40-J47)	120	3.5	49.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	2.7	39.3
7	Influenza and pneumonia (J09-J18)	88	2.5	36.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	69	2.0	28.5
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	52	1.5	21.5
10	Parkinson's disease (G20-G21)	45	1.3	18.6
11	Septicemia (A40-A41)	42	1.2	17.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	34	1.0	14.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	0.8	11.6
14	Aortic aneurysm and dissection (I71)	25	0.7	10.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	20	0.6	8.3
...	All other causes (residual)	467	13.4	193.1
Non-Hispanic, Asian or Pacific Islander, male, 75-79 years				
...	All causes	4,029	100.0	2,363.7
1	Malignant neoplasms (C00-C97)	1,215	30.2	712.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	861	21.4	505.1
3	Cerebrovascular diseases (I60-I69)	291	7.2	170.7
4	Diabetes mellitus (E10-E14)	205	5.1	120.3
5	Chronic lower respiratory diseases (J40-J47)	146	3.6	85.7
6	Influenza and pneumonia (J09-J18)	143	3.5	83.9
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	132	3.3	77.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	104	2.6	61.0
9	Parkinson's disease (G20-G21)	95	2.4	55.7
10	Alzheimer's disease (G30)	71	1.8	41.7
11	Septicemia (A40-A41)	60	1.5	35.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	57	1.4	33.4
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	31	0.8	18.2
13	Pneumonitis due to solids and liquids (J69)	31	0.8	18.2
15	Aortic aneurysm and dissection (I71)	23	0.6	13.5
...	All other causes (residual)	564	14.0	330.9

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 80-84 years				
...	All causes	4,341	100.0	4,119.5
1	Malignant neoplasms (C00-C97)	1,091	25.1	1,035.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,029	23.7	976.5
3	Cerebrovascular diseases (I60-I69)	325	7.5	308.4
4	Chronic lower respiratory diseases (J40-J47)	235	5.4	223.0
5	Diabetes mellitus (E10-E14)	189	4.4	179.4
6	Influenza and pneumonia (J09-J18)	149	3.4	141.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	129	3.0	122.4
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	115	2.6	109.1
9	Alzheimer's disease (G30)	112	2.6	106.3
10	Parkinson's disease (G20-G21)	110	2.5	104.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	75	1.7	71.2
12	Pneumonitis due to solids and liquids (J69)	51	1.2	48.4
13	Septicemia (A40-A41)	50	1.2	47.4
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	36	0.8	34.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	20	0.5	19.0
...	All other causes (residual)	625	14.4	593.1
Non-Hispanic, Asian or Pacific Islander, male, 85-89 years				
...	All causes	4,107	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,051	25.6	---
2	Malignant neoplasms (C00-C97)	785	19.1	---
3	Cerebrovascular diseases (I60-I69)	337	8.2	---
4	Chronic lower respiratory diseases (J40-J47)	226	5.5	---
5	Influenza and pneumonia (J09-J18)	214	5.2	---
6	Alzheimer's disease (G30)	189	4.6	---
7	Diabetes mellitus (E10-E14)	145	3.5	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109	2.7	---
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	105	2.6	---
10	Parkinson's disease (G20-G21)	94	2.3	---
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	94	2.3	---
12	Pneumonitis due to solids and liquids (J69)	59	1.4	---
13	Septicemia (A40-A41)	46	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	32	0.8	---
15	Aortic aneurysm and dissection (I71)	21	0.5	---
...	All other causes (residual)	600	14.6	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 90-94 years				
...	All causes	2,885	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	738	25.6	---
2	Malignant neoplasms (C00-C97)	397	13.8	---
3	Cerebrovascular diseases (I60-I69)	235	8.1	---
4	Influenza and pneumonia (J09-J18)	189	6.6	---
5	Alzheimer's disease (G30)	188	6.5	---
6	Chronic lower respiratory diseases (J40-J47)	141	4.9	---
7	Diabetes mellitus (E10-E14)	100	3.5	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	80	2.8	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	73	2.5	---
10	Pneumonitis due to solids and liquids (J69)	67	2.3	---
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	1.9	---
12	Parkinson's disease (G20-G21)	48	1.7	---
13	Septicemia (A40-A41)	30	1.0	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	21	0.7	---
15	Aortic aneurysm and dissection (I71)	14	0.5	---
...	All other causes (residual)	509	17.6	---
Non-Hispanic, Asian or Pacific Islander, male, 95-99 years				
...	All causes	1,132	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	330	29.2	---
2	Malignant neoplasms (C00-C97)	106	9.4	---
3	Alzheimer's disease (G30)	100	8.8	---
4	Cerebrovascular diseases (I60-I69)	86	7.6	---
5	Influenza and pneumonia (J09-J18)	76	6.7	---
6	Chronic lower respiratory diseases (J40-J47)	57	5.0	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	28	2.5	---
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	2.4	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	2.3	---
10	Diabetes mellitus (E10-E14)	24	2.1	---
11	Septicemia (A40-A41)	15	1.3	---
11	Pneumonitis due to solids and liquids (J69)	15	1.3	---
13	Parkinson's disease (G20-G21)	12	1.1	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	0.7	---
14	Nutritional deficiencies (E40-E64)	8	0.7	---
...	All other causes (residual)	214	18.9	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 100 years and over				
...	All causes	215	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	31.2	---
2	Malignant neoplasms (C00-C97)	18	8.4	---
2	Influenza and pneumonia (J09-J18)	18	8.4	---
4	Alzheimer's disease (G30)	15	7.0	---
5	Chronic lower respiratory diseases (J40-J47)	14	6.5	---
6	Cerebrovascular diseases (I60-I69)	9	4.2	---
7	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8	3.7	---
8	Pneumonitis due to solids and liquids (J69)	5	2.3	---
9	Atherosclerosis (I70)	4	1.9	---
10	Aortic aneurysm and dissection (I71)	3	1.4	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3	1.4	---
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.9	---
12	Diabetes mellitus (E10-E14)	2	0.9	---
12	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2	0.9	---
15	Tuberculosis (A16-A19)	1	0.5	---
15	Septicemia (A40-A41)	1	0.5	---
15	Anemias (D50-D64)	1	0.5	---
15	Nutritional deficiencies (E40-E64)	1	0.5	---
15	Parkinson's disease (G20-G21)	1	0.5	---
15	Hyperplasia of prostate (N40)	1	0.5	---
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.5	---
...	All other causes (residual)	38	17.7	---
Non-Hispanic, Asian or Pacific Islander, female, all ages				
...	All causes	33,343	100.0	327.8
1	Malignant neoplasms (C00-C97)	8,740	26.2	85.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,679	20.0	65.7
3	Cerebrovascular diseases (I60-I69)	2,949	8.8	29.0
4	Alzheimer's disease (G30)	1,654	5.0	16.3
5	Diabetes mellitus (E10-E14)	1,283	3.8	12.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,172	3.5	11.5
7	Influenza and pneumonia (J09-J18)	1,012	3.0	9.9
8	Chronic lower respiratory diseases (J40-J47)	780	2.3	7.7
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	735	2.2	7.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	683	2.0	6.7
11	Septicemia (A40-A41)	393	1.2	3.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	386	1.2	3.8
13	Parkinson's disease (G20-G21)	309	0.9	3.0
14	Chronic liver disease and cirrhosis (K70,K73-K74)	244	0.7	2.4
15	Certain conditions originating in the perinatal period (P00-P96)	221	0.7	2.2
...	All other causes (residual)	6,103	18.3	60.0

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 1-4 years				
...	All causes	80	100.0	17.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	17.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	15.0	*
3	Malignant neoplasms (C00-C97)	11	13.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	10.0	*
5	Septicemia (A40-A41)	4	5.0	*
6	Cerebrovascular diseases (I60-I69)	2	2.5	*
7	Diabetes mellitus (E10-E14)	1	1.3	*
7	Meningitis (G00,G03)	1	1.3	*
7	Influenza and pneumonia (J09-J18)	1	1.3	*
7	Chronic lower respiratory diseases (J40-J47)	1	1.3	*
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.3	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	1.3	*
...	All other causes (residual)	23	28.8	5.0
Non-Hispanic, Asian or Pacific Islander, female, 5-9 years				
...	All causes	42	100.0	7.3
1	Malignant neoplasms (C00-C97)	9	21.4	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5	11.9	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	9.5	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	7.1	*
5	Cerebrovascular diseases (I60-I69)	2	4.8	*
5	Influenza and pneumonia (J09-J18)	2	4.8	*
7	Scarlet fever and erysipelas (A38,A46)	1	2.4	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.4	*
...	All other causes (residual)	15	35.7	*
Non-Hispanic, Asian or Pacific Islander, female, 10-14 years				
...	All causes	58	100.0	9.8
1	Malignant neoplasms (C00-C97)	19	32.8	*
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	17.2	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3	5.2	*
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	5.2	*
5	Septicemia (A40-A41)	2	3.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.4	*
5	Influenza and pneumonia (J09-J18)	2	3.4	*
5	Certain conditions originating in the perinatal period (P00-P96)	2	3.4	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	3.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.7	*
10	Meningitis (G00,G03)	1	1.7	*
10	Cerebrovascular diseases (I60-I69)	1	1.7	*
10	Diseases of appendix (K35-K38)	1	1.7	*
...	All other causes (residual)	9	15.5	*

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 15-19 years				
...	All causes	107	100.0	18.1
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	29.0	5.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	24.3	4.4
3	Malignant neoplasms (C00-C97)	12	11.2	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7	6.5	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	6.5	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	2.8	*
7	Cerebrovascular diseases (I60-I69)	2	1.9	*
8	Septicemia (A40-A41)	1	0.9	*
8	Diabetes mellitus (E10-E14)	1	0.9	*
8	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
...	All other causes (residual)	16	15.0	*
Non-Hispanic, Asian or Pacific Islander, female, 20-24 years				
...	All causes	168	100.0	24.2
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	51	30.4	7.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	23.8	5.8
3	Malignant neoplasms (C00-C97)	17	10.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	6.0	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	4.8	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	6	3.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.8	*
8	Cerebrovascular diseases (I60-I69)	2	1.2	*
8	Influenza and pneumonia (J09-J18)	2	1.2	*
10	Tuberculosis (A16-A19)	1	0.6	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.6	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.6	*
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
10	Inflammatory diseases of female pelvic organs (N70-N76)	1	0.6	*
...	All other causes (residual)	24	14.3	3.5

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 25-29 years				
...	All causes	175	100.0	20.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	51	29.1	6.0
2	Malignant neoplasms (C00-C97)	33	18.9	3.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	16.0	3.3
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	5.7	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	9	5.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	4.0	*
7	Septicemia (A40-A41)	2	1.1	*
7	Influenza and pneumonia (J09-J18)	2	1.1	*
7	Chronic lower respiratory diseases (J40-J47)	2	1.1	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.1	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.6	*
11	Diabetes mellitus (E10-E14)	1	0.6	*
11	Cerebrovascular diseases (I60-I69)	1	0.6	*
11	Aortic aneurysm and dissection (I71)	1	0.6	*
11	Peptic ulcer (K25-K28)	1	0.6	*
11	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.6	*
...	All other causes (residual)	22	12.6	2.6
Non-Hispanic, Asian or Pacific Islander, female, 30-34 years				
...	All causes	282	100.0	32.5
1	Malignant neoplasms (C00-C97)	68	24.1	7.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47	16.7	5.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	40	14.2	4.6
4	Cerebrovascular diseases (I60-I69)	16	5.7	*
5	Pregnancy, childbirth and the puerperium (O00-O99)	14	5.0	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	4.3	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	3.2	*
8	Septicemia (A40-A41)	4	1.4	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	4	1.4	*
10	Human immunodeficiency virus (HIV) disease (B20-B24)	3	1.1	*
10	Diabetes mellitus (E10-E14)	3	1.1	*
10	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.1	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	3	1.1	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
15	Meningococcal infection (A39)	1	0.4	*
15	Viral hepatitis (B15-B19)	1	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.4	*
15	Anemias (D50-D64)	1	0.4	*
15	Nutritional deficiencies (E40-E64)	1	0.4	*
15	Aortic aneurysm and dissection (I71)	1	0.4	*
15	Influenza and pneumonia (J09-J18)	1	0.4	*
15	Chronic lower respiratory diseases (J40-J47)	1	0.4	*
15	Diseases of appendix (K35-K38)	1	0.4	*
15	Infections of kidney (N10-N12,N13.6,N15.1)	1	0.4	*
...	All other causes (residual)	44	15.6	5.1

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 35-39 years				
...	All causes	376	100.0	45.5
1	Malignant neoplasms (C00-C97)	173	46.0	20.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	32	8.5	3.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	8.0	3.6
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	25	6.6	3.0
5	Cerebrovascular diseases (I60-I69)	23	6.1	2.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	13	3.5	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	1.9	*
8	Chronic liver disease and cirrhosis (K70,K73-K74)	6	1.6	*
9	Diabetes mellitus (E10-E14)	4	1.1	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	1.1	*
11	Influenza and pneumonia (J09-J18)	3	0.8	*
12	Tuberculosis (A16-A19)	2	0.5	*
12	Septicemia (A40-A41)	2	0.5	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2	0.5	*
12	Aortic aneurysm and dissection (I71)	2	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	2	0.5	*
...	All other causes (residual)	46	12.2	5.6
Non-Hispanic, Asian or Pacific Islander, female, 40-44 years				
...	All causes	501	100.0	62.4
1	Malignant neoplasms (C00-C97)	242	48.3	30.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	54	10.8	6.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	7.2	4.5
4	Cerebrovascular diseases (I60-I69)	28	5.6	3.5
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	26	5.2	3.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	16	3.2	*
7	Diabetes mellitus (E10-E14)	11	2.2	*
8	Pregnancy, childbirth and the puerperium (O00-O99)	6	1.2	*
9	Anemias (D50-D64)	5	1.0	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	1.0	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.8	*
11	Influenza and pneumonia (J09-J18)	4	0.8	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.8	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3	0.6	*
15	Chronic lower respiratory diseases (J40-J47)	2	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.4	*
...	All other causes (residual)	51	10.2	6.4

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 45-49 years				
...	All causes	734	100.0	99.9
1	Malignant neoplasms (C00-C97)	389	53.0	52.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	57	7.8	7.8
3	Cerebrovascular diseases (I60-I69)	56	7.6	7.6
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	5.7	5.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	31	4.2	4.2
6	Diabetes mellitus (E10-E14)	24	3.3	3.3
7	Chronic liver disease and cirrhosis (K70,K73-K74)	10	1.4	*
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	10	1.4	*
9	Septicemia (A40-A41)	9	1.2	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5	0.7	*
10	Influenza and pneumonia (J09-J18)	5	0.7	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.7	*
10	Pregnancy, childbirth and the puerperium (O00-O99)	5	0.7	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	4	0.5	*
...	All other causes (residual)	78	10.6	10.6
Non-Hispanic, Asian or Pacific Islander, female, 50-54 years				
...	All causes	1,039	100.0	157.3
1	Malignant neoplasms (C00-C97)	550	52.9	83.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	82	7.9	12.4
3	Cerebrovascular diseases (I60-I69)	79	7.6	12.0
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	42	4.0	6.4
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	37	3.6	5.6
6	Diabetes mellitus (E10-E14)	28	2.7	4.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	21	2.0	3.2
8	Influenza and pneumonia (J09-J18)	16	1.5	*
9	Septicemia (A40-A41)	14	1.3	*
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	12	1.2	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	7	0.7	*
11	Chronic lower respiratory diseases (J40-J47)	7	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.5	*
15	Viral hepatitis (B15-B19)	4	0.4	*
15	Aortic aneurysm and dissection (I71)	4	0.4	*
...	All other causes (residual)	124	11.9	18.8

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 55-59 years				
...	All causes	1,445	100.0	238.6
1	Malignant neoplasms (C00-C97)	751	52.0	124.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	185	12.8	30.6
3	Cerebrovascular diseases (I60-I69)	89	6.2	14.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	58	4.0	9.6
5	Diabetes mellitus (E10-E14)	54	3.7	8.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.9	4.5
7	Influenza and pneumonia (J09-J18)	20	1.4	3.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	1.4	3.3
9	Septicemia (A40-A41)	17	1.2	*
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17	1.2	*
9	Chronic lower respiratory diseases (J40-J47)	17	1.2	*
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.8	*
13	Viral hepatitis (B15-B19)	8	0.6	*
13	Chronic liver disease and cirrhosis (K70,K73-K74)	8	0.6	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	0.5	*
...	All other causes (residual)	155	10.7	25.6
Non-Hispanic, Asian or Pacific Islander, female, 60-64 years				
...	All causes	2,064	100.0	382.7
1	Malignant neoplasms (C00-C97)	1,021	49.5	189.3
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	258	12.5	47.8
3	Cerebrovascular diseases (I60-I69)	136	6.6	25.2
4	Diabetes mellitus (E10-E14)	85	4.1	15.8
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	69	3.3	12.8
6	Influenza and pneumonia (J09-J18)	36	1.7	6.7
7	Chronic lower respiratory diseases (J40-J47)	35	1.7	6.5
8	Chronic liver disease and cirrhosis (K70,K73-K74)	30	1.5	5.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	28	1.4	5.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	27	1.3	5.0
11	Septicemia (A40-A41)	24	1.2	4.5
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.1	4.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.8	*
14	Viral hepatitis (B15-B19)	11	0.5	*
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	0.5	*
...	All other causes (residual)	255	12.4	47.3

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016



LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 65-69 years				
...	All causes	2,552	100.0	569.4
1	Malignant neoplasms (C00-C97)	1,065	41.7	237.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	425	16.7	94.8
3	Cerebrovascular diseases (I60-I69)	176	6.9	39.3
4	Diabetes mellitus (E10-E14)	119	4.7	26.5
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	3.7	21.2
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	55	2.2	12.3
7	Chronic lower respiratory diseases (J40-J47)	48	1.9	10.7
8	Influenza and pneumonia (J09-J18)	47	1.8	10.5
9	Septicemia (A40-A41)	44	1.7	9.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	30	1.2	6.7
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	26	1.0	5.8
11	Chronic liver disease and cirrhosis (K70,K73-K74)	26	1.0	5.8
13	Parkinson's disease (G20-G21)	19	0.7	*
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	19	0.7	*
15	Viral hepatitis (B15-B19)	14	0.5	*
...	All other causes (residual)	344	13.5	76.7
Non-Hispanic, Asian or Pacific Islander, female, 70-74 years				
...	All causes	2,680	100.0	894.5
1	Malignant neoplasms (C00-C97)	948	35.4	316.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	484	18.1	161.5
3	Cerebrovascular diseases (I60-I69)	235	8.8	78.4
4	Diabetes mellitus (E10-E14)	142	5.3	47.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	95	3.5	31.7
6	Chronic lower respiratory diseases (J40-J47)	70	2.6	23.4
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	67	2.5	22.4
8	Influenza and pneumonia (J09-J18)	62	2.3	20.7
9	Septicemia (A40-A41)	50	1.9	16.7
10	Alzheimer's disease (G30)	31	1.2	10.3
11	Parkinson's disease (G20-G21)	30	1.1	10.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29	1.1	9.7
12	Chronic liver disease and cirrhosis (K70,K73-K74)	29	1.1	9.7
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	17	0.6	*
15	Viral hepatitis (B15-B19)	16	0.6	*
...	All other causes (residual)	375	14.0	125.2

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 75-79 years				
...	All causes	3,415	100.0	1,610.4
1	Malignant neoplasms (C00-C97)	1,001	29.3	472.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	707	20.7	333.4
3	Cerebrovascular diseases (I60-I69)	310	9.1	146.2
4	Diabetes mellitus (E10-E14)	169	4.9	79.7
5	Alzheimer's disease (G30)	113	3.3	53.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	97	2.8	45.7
7	Chronic lower respiratory diseases (J40-J47)	92	2.7	43.4
8	Influenza and pneumonia (J09-J18)	89	2.6	42.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	88	2.6	41.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	2.2	35.8
11	Septicemia (A40-A41)	51	1.5	24.0
12	Parkinson's disease (G20-G21)	43	1.3	20.3
13	Chronic liver disease and cirrhosis (K70,K73-K74)	36	1.1	17.0
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.8	13.7
15	Aortic aneurysm and dissection (I71)	18	0.5	*
...	All other causes (residual)	496	14.5	233.9
Non-Hispanic, Asian or Pacific Islander, female, 80-84 years				
...	All causes	4,489	100.0	3,071.8
1	Malignant neoplasms (C00-C97)	1,011	22.5	691.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,004	22.4	687.0
3	Cerebrovascular diseases (I60-I69)	468	10.4	320.2
4	Alzheimer's disease (G30)	228	5.1	156.0
5	Diabetes mellitus (E10-E14)	192	4.3	131.4
6	Influenza and pneumonia (J09-J18)	141	3.1	96.5
7	Chronic lower respiratory diseases (J40-J47)	137	3.1	93.7
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	131	2.9	89.6
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	112	2.5	76.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	110	2.5	75.3
11	Parkinson's disease (G20-G21)	79	1.8	54.1
12	Septicemia (A40-A41)	43	1.0	29.4
13	Chronic liver disease and cirrhosis (K70,K73-K74)	39	0.9	26.7
14	Pneumonitis due to solids and liquids (J69)	34	0.8	23.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	0.6	18.5
...	All other causes (residual)	733	16.3	501.6

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 85-89 years				
...	All causes	5,428	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,360	25.1	---
2	Malignant neoplasms (C00-C97)	816	15.0	---
3	Cerebrovascular diseases (I60-I69)	580	10.7	---
4	Alzheimer's disease (G30)	409	7.5	---
5	Diabetes mellitus (E10-E14)	229	4.2	---
6	Influenza and pneumonia (J09-J18)	216	4.0	---
7	Chronic lower respiratory diseases (J40-J47)	169	3.1	---
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	165	3.0	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	144	2.7	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	128	2.4	---
11	Parkinson's disease (G20-G21)	77	1.4	---
12	Septicemia (A40-A41)	62	1.1	---
13	Pneumonitis due to solids and liquids (J69)	44	0.8	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	43	0.8	---
15	Aortic aneurysm and dissection (I71)	22	0.4	---
...	All other causes (residual)	964	17.8	---
Non-Hispanic, Asian or Pacific Islander, female, 90-94 years				
...	All causes	4,554	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,201	26.4	---
2	Alzheimer's disease (G30)	479	10.5	---
3	Malignant neoplasms (C00-C97)	465	10.2	---
4	Cerebrovascular diseases (I60-I69)	456	10.0	---
5	Influenza and pneumonia (J09-J18)	217	4.8	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	160	3.5	---
7	Diabetes mellitus (E10-E14)	154	3.4	---
8	Chronic lower respiratory diseases (J40-J47)	117	2.6	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	108	2.4	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	107	2.3	---
11	Parkinson's disease (G20-G21)	42	0.9	---
12	Septicemia (A40-A41)	40	0.9	---
13	Pneumonitis due to solids and liquids (J69)	33	0.7	---
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	29	0.6	---
15	Nutritional deficiencies (E40-E64)	25	0.5	---
...	All other causes (residual)	921	20.2	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016

LCWK1. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 5-year age groups, by race and Hispanic origin, and sex: United States, 2016

[Rates per 100,000 population in specified group. Rates are not shown for age groups over 85 years. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank <sup>1</sup>	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i> ), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 95-99 years				
...	All causes	2,178	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	623	28.6	---
2	Alzheimer's disease (G30)	306	14.0	---
3	Cerebrovascular diseases (I60-I69)	232	10.7	---
4	Malignant neoplasms (C00-C97)	115	5.3	---
5	Influenza and pneumonia (J09-J18)	111	5.1	---
6	Essential hypertension and hypertensive renal disease (I10,I12,I15)	67	3.1	---
7	Chronic lower respiratory diseases (J40-J47)	64	2.9	---
8	Diabetes mellitus (E10-E14)	54	2.5	---
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	38	1.7	---
10	Pneumonitis due to solids and liquids (J69)	25	1.1	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25	1.1	---
12	Septicemia (A40-A41)	15	0.7	---
13	Nutritional deficiencies (E40-E64)	12	0.6	---
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	10	0.5	---
15	Parkinson's disease (G20-G21)	9	0.4	---
...	All other causes (residual)	472	21.7	---
Non-Hispanic, Asian or Pacific Islander, female, 100 years and over				
...	All causes	578	100.0	---
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	161	27.9	---
2	Alzheimer's disease (G30)	66	11.4	---
3	Cerebrovascular diseases (I60-I69)	54	9.3	---
4	Influenza and pneumonia (J09-J18)	34	5.9	---
5	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24	4.2	---
6	Malignant neoplasms (C00-C97)	23	4.0	---
7	Diabetes mellitus (E10-E14)	12	2.1	---
8	Chronic lower respiratory diseases (J40-J47)	11	1.9	---
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	1.9	---
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	10	1.7	---
11	Septicemia (A40-A41)	5	0.9	---
12	Anemias (D50-D64)	3	0.5	---
12	Nutritional deficiencies (E40-E64)	3	0.5	---
12	Pneumonitis due to solids and liquids (J69)	3	0.5	---
15	Parkinson's disease (G20-G21)	2	0.3	---
15	Atherosclerosis (I70)	2	0.3	---
15	Aortic aneurysm and dissection (I71)	2	0.3	---
15	Cholelithiasis and other disorders of gallbladder (K80-K82)	2	0.3	---
...	All other causes (residual)	150	26.0	---

... Category not applicable

\* Figure does not meet standards of reliability or precision

--- Data not available

<sup>1</sup> Rank based on number of deaths.

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2016