

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|-----------|-------------------------|-------|
| All races and origins, both sexes, all ages | | | | |
| ... | All causes | 2,712,630 | 100.0 | 844.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 633,842 | 23.4 | 197.2 |
| 2 | Malignant neoplasms (C00-C97) | 595,930 | 22.0 | 185.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 155,041 | 5.7 | 48.2 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 146,571 | 5.4 | 45.6 |
| 5 | Cerebrovascular diseases (I60-I69) | 140,323 | 5.2 | 43.7 |
| 6 | Alzheimer's disease (G30) | 110,561 | 4.1 | 34.4 |
| 7 | Diabetes mellitus (E10-E14) | 79,535 | 2.9 | 24.7 |
| 8 | Influenza and pneumonia (J09-J18) | 57,062 | 2.1 | 17.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 49,959 | 1.8 | 15.5 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 44,193 | 1.6 | 13.7 |
| 11 | Septicemia (A40-A41) | 40,773 | 1.5 | 12.7 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 40,326 | 1.5 | 12.5 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 32,200 | 1.2 | 10.0 |
| 14 | Parkinson's disease (G20-G21) | 27,972 | 1.0 | 8.7 |
| 15 | Pneumonitis due to solids and liquids (J69) | 19,803 | 0.7 | 6.2 |
| ... | All other causes (residual) | 538,539 | 19.9 | 167.6 |
| All races and origins, both sexes, 1-4 years | | | | |
| ... | All causes | 3,965 | 100.0 | 24.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,235 | 31.1 | 7.8 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 435 | 11.0 | 2.7 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 369 | 9.3 | 2.3 |
| 4 | Malignant neoplasms (C00-C97) | 354 | 8.9 | 2.2 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 147 | 3.7 | 0.9 |
| 6 | Influenza and pneumonia (J09-J18) | 88 | 2.2 | 0.6 |
| 7 | Septicemia (A40-A41) | 54 | 1.4 | 0.3 |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 50 | 1.3 | 0.3 |
| 9 | Cerebrovascular diseases (I60-I69) | 42 | 1.1 | 0.3 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 40 | 1.0 | 0.3 |
| 11 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 34 | 0.9 | 0.2 |
| 12 | Anemias (D50-D64) | 25 | 0.6 | 0.2 |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 18 | 0.5 | * |
| 14 | Acute bronchitis and bronchiolitis (J20-J21) | 17 | 0.4 | * |
| 15 | Meningitis (G00,G03) | 16 | 0.4 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 16 | 0.4 | * |
| ... | All other causes (residual) | 1,025 | 25.9 | 6.4 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| All races and origins, both sexes, 5-14 years | | | | |
| ... | All causes | 5,411 | 100.0 | 13.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,518 | 28.1 | 3.7 |
| 2 | Malignant neoplasms (C00-C97) | 865 | 16.0 | 2.1 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 413 | 7.6 | 1.0 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 337 | 6.2 | 0.8 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 298 | 5.5 | 0.7 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 210 | 3.9 | 0.5 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 173 | 3.2 | 0.4 |
| 8 | Cerebrovascular diseases (I60-I69) | 84 | 1.6 | 0.2 |
| 9 | Influenza and pneumonia (J09-J18) | 83 | 1.5 | 0.2 |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 72 | 1.3 | 0.2 |
| 11 | Septicemia (A40-A41) | 64 | 1.2 | 0.2 |
| 12 | Anemias (D50-D64) | 33 | 0.6 | 0.1 |
| 13 | Diabetes mellitus (E10-E14) | 23 | 0.4 | 0.1 |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 21 | 0.4 | 0.1 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 17 | 0.3 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 17 | 0.3 | * |
| ... | All other causes (residual) | 1,183 | 21.9 | 2.9 |
| All races and origins, both sexes, 15-24 years | | | | |
| ... | All causes | 30,494 | 100.0 | 69.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 12,514 | 41.0 | 28.5 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 5,491 | 18.0 | 12.5 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 4,733 | 15.5 | 10.8 |
| 4 | Malignant neoplasms (C00-C97) | 1,469 | 4.8 | 3.4 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 997 | 3.3 | 2.3 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 386 | 1.3 | 0.9 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 202 | 0.7 | 0.5 |
| 8 | Diabetes mellitus (E10-E14) | 196 | 0.6 | 0.4 |
| 9 | Influenza and pneumonia (J09-J18) | 184 | 0.6 | 0.4 |
| 10 | Cerebrovascular diseases (I60-I69) | 166 | 0.5 | 0.4 |
| 11 | Pregnancy, childbirth and the puerperium (O00-O99) | 161 | 0.5 | 0.4 |
| 12 | Septicemia (A40-A41) | 130 | 0.4 | 0.3 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 100 | 0.3 | 0.2 |
| 14 | Legal intervention (Y35,Y89.0) | 88 | 0.3 | 0.2 |
| 15 | Anemias (D50-D64) | 84 | 0.3 | 0.2 |
| ... | All other causes (residual) | 3,593 | 11.8 | 8.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| All races and origins, both sexes, 25-34 years | | | | |
| ... | All causes | 51,517 | 100.0 | 116.7 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 19,795 | 38.4 | 44.8 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 6,947 | 13.5 | 15.7 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 4,863 | 9.4 | 11.0 |
| 4 | Malignant neoplasms (C00-C97) | 3,704 | 7.2 | 8.4 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,522 | 6.8 | 8.0 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 844 | 1.6 | 1.9 |
| 7 | Diabetes mellitus (E10-E14) | 798 | 1.5 | 1.8 |
| 8 | Cerebrovascular diseases (I60-I69) | 567 | 1.1 | 1.3 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 529 | 1.0 | 1.2 |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 443 | 0.9 | 1.0 |
| 11 | Pregnancy, childbirth and the puerperium (O00-O99) | 424 | 0.8 | 1.0 |
| 12 | Influenza and pneumonia (J09-J18) | 397 | 0.8 | 0.9 |
| 13 | Septicemia (A40-A41) | 379 | 0.7 | 0.9 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 288 | 0.6 | 0.7 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 262 | 0.5 | 0.6 |
| ... | All other causes (residual) | 7,755 | 15.1 | 17.6 |
| All races and origins, both sexes, 35-44 years | | | | |
| ... | All causes | 73,088 | 100.0 | 180.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 17,818 | 24.4 | 43.9 |
| 2 | Malignant neoplasms (C00-C97) | 10,909 | 14.9 | 26.9 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,387 | 14.2 | 25.6 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 6,936 | 9.5 | 17.1 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,895 | 4.0 | 7.1 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,861 | 3.9 | 7.0 |
| 7 | Diabetes mellitus (E10-E14) | 1,986 | 2.7 | 4.9 |
| 8 | Cerebrovascular diseases (I60-I69) | 1,788 | 2.4 | 4.4 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,055 | 1.4 | 2.6 |
| 10 | Septicemia (A40-A41) | 829 | 1.1 | 2.0 |
| 11 | Influenza and pneumonia (J09-J18) | 708 | 1.0 | 1.7 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 702 | 1.0 | 1.7 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 686 | 0.9 | 1.7 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 483 | 0.7 | 1.2 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 470 | 0.6 | 1.2 |
| ... | All other causes (residual) | 12,575 | 17.2 | 31.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|-------|
| All races and origins, both sexes, 45-54 years | | | | |
| ... | All causes | 174,494 | 100.0 | 404.0 |
| 1 | Malignant neoplasms (C00-C97) | 43,054 | 24.7 | 99.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 34,248 | 19.6 | 79.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 21,499 | 12.3 | 49.8 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 8,874 | 5.1 | 20.5 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 8,751 | 5.0 | 20.3 |
| 6 | Diabetes mellitus (E10-E14) | 6,212 | 3.6 | 14.4 |
| 7 | Cerebrovascular diseases (I60-I69) | 5,307 | 3.0 | 12.3 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 4,345 | 2.5 | 10.1 |
| 9 | Septicemia (A40-A41) | 2,542 | 1.5 | 5.9 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,124 | 1.2 | 4.9 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,106 | 1.2 | 4.9 |
| 12 | Influenza and pneumonia (J09-J18) | 2,050 | 1.2 | 4.7 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2,027 | 1.2 | 4.7 |
| 14 | Viral hepatitis (B15-B19) | 1,496 | 0.9 | 3.5 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,477 | 0.8 | 3.4 |
| ... | All other causes (residual) | 28,382 | 16.3 | 65.7 |
| All races and origins, both sexes, 55-64 years | | | | |
| ... | All causes | 357,785 | 100.0 | 875.3 |
| 1 | Malignant neoplasms (C00-C97) | 116,122 | 32.5 | 284.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 76,872 | 21.5 | 188.1 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 19,488 | 5.4 | 47.7 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 17,457 | 4.9 | 42.7 |
| 5 | Diabetes mellitus (E10-E14) | 14,166 | 4.0 | 34.7 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 13,278 | 3.7 | 32.5 |
| 7 | Cerebrovascular diseases (I60-I69) | 12,116 | 3.4 | 29.6 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 7,739 | 2.2 | 18.9 |
| 9 | Septicemia (A40-A41) | 5,774 | 1.6 | 14.1 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 5,452 | 1.5 | 13.3 |
| 11 | Influenza and pneumonia (J09-J18) | 4,601 | 1.3 | 11.3 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 3,595 | 1.0 | 8.8 |
| 13 | Viral hepatitis (B15-B19) | 3,559 | 1.0 | 8.7 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,824 | 0.5 | 4.5 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,619 | 0.5 | 4.0 |
| ... | All other causes (residual) | 54,123 | 15.1 | 132.4 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| All races and origins, both sexes, 65-74 years | | | | |
| ... | All causes | 495,016 | 100.0 | 1,796.8 |
| 1 | Malignant neoplasms (C00-C97) | 163,728 | 33.1 | 594.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 107,303 | 21.7 | 389.5 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 37,642 | 7.6 | 136.6 |
| 4 | Cerebrovascular diseases (I60-I69) | 20,793 | 4.2 | 75.5 |
| 5 | Diabetes mellitus (E10-E14) | 19,453 | 3.9 | 70.6 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 12,961 | 2.6 | 47.0 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 9,662 | 2.0 | 35.1 |
| 8 | Septicemia (A40-A41) | 8,655 | 1.7 | 31.4 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 8,409 | 1.7 | 30.5 |
| 10 | Influenza and pneumonia (J09-J18) | 8,131 | 1.6 | 29.5 |
| 11 | Alzheimer's disease (G30) | 6,167 | 1.2 | 22.4 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 4,979 | 1.0 | 18.1 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 4,201 | 0.8 | 15.2 |
| 14 | Parkinson's disease (G20-G21) | 3,813 | 0.8 | 13.8 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 3,156 | 0.6 | 11.5 |
| ... | All other causes (residual) | 75,963 | 15.3 | 275.7 |
| All races and origins, both sexes, 75-84 years | | | | |
| ... | All causes | 637,566 | 100.0 | 4,579.2 |
| 1 | Malignant neoplasms (C00-C97) | 153,268 | 24.0 | 1,100.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 149,199 | 23.4 | 1,071.6 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 49,832 | 7.8 | 357.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 38,012 | 6.0 | 273.0 |
| 5 | Alzheimer's disease (G30) | 29,503 | 4.6 | 211.9 |
| 6 | Diabetes mellitus (E10-E14) | 19,904 | 3.1 | 143.0 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 15,518 | 2.4 | 111.5 |
| 8 | Influenza and pneumonia (J09-J18) | 14,149 | 2.2 | 101.6 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 13,881 | 2.2 | 99.7 |
| 10 | Parkinson's disease (G20-G21) | 11,473 | 1.8 | 82.4 |
| 11 | Septicemia (A40-A41) | 10,481 | 1.6 | 75.3 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 7,533 | 1.2 | 54.1 |
| 13 | Pneumonitis due to solids and liquids (J69) | 5,152 | 0.8 | 37.0 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4,837 | 0.8 | 34.7 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 4,446 | 0.7 | 31.9 |
| ... | All other causes (residual) | 110,378 | 17.3 | 792.8 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|-----------|-------------------------|----------|
| All races and origins, both sexes, 85 years and over | | | | |
| ... | All causes | 859,701 | 100.0 | 13,673.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 250,636 | 29.2 | 3,986.5 |
| 2 | Malignant neoplasms (C00-C97) | 102,393 | 11.9 | 1,628.6 |
| 3 | Alzheimer's disease (G30) | 73,825 | 8.6 | 1,174.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 61,351 | 7.1 | 975.8 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 44,330 | 5.2 | 705.1 |
| 6 | Influenza and pneumonia (J09-J18) | 26,494 | 3.1 | 421.4 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 22,916 | 2.7 | 364.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 17,715 | 2.1 | 281.8 |
| 9 | Diabetes mellitus (E10-E14) | 16,785 | 2.0 | 267.0 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 14,003 | 1.6 | 222.7 |
| 11 | Parkinson's disease (G20-G21) | 11,983 | 1.4 | 190.6 |
| 12 | Septicemia (A40-A41) | 11,681 | 1.4 | 185.8 |
| 13 | Pneumonitis due to solids and liquids (J69) | 9,736 | 1.1 | 154.9 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 5,346 | 0.6 | 85.0 |
| 15 | Atherosclerosis (I70) | 3,346 | 0.4 | 53.2 |
| ... | All other causes (residual) | 187,161 | 21.8 | 2,976.9 |
| All races and origins, male, all ages | | | | |
| ... | All causes | 1,373,404 | 100.0 | 868.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 335,002 | 24.4 | 211.7 |
| 2 | Malignant neoplasms (C00-C97) | 313,818 | 22.8 | 198.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 92,919 | 6.8 | 58.7 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 72,498 | 5.3 | 45.8 |
| 5 | Cerebrovascular diseases (I60-I69) | 58,288 | 4.2 | 36.8 |
| 6 | Diabetes mellitus (E10-E14) | 43,123 | 3.1 | 27.3 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 33,994 | 2.5 | 21.5 |
| 8 | Alzheimer's disease (G30) | 33,690 | 2.5 | 21.3 |
| 9 | Influenza and pneumonia (J09-J18) | 26,903 | 2.0 | 17.0 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 25,666 | 1.9 | 16.2 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 25,441 | 1.9 | 16.1 |
| 12 | Septicemia (A40-A41) | 19,385 | 1.4 | 12.3 |
| 13 | Parkinson's disease (G20-G21) | 16,867 | 1.2 | 10.7 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 14,274 | 1.0 | 9.0 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 13,934 | 1.0 | 8.8 |
| ... | All other causes (residual) | 247,602 | 18.0 | 156.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| All races and origins, male, 1-4 years | | | | |
| ... | All causes | 2,281 | 100.0 | 28.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 768 | 33.7 | 9.4 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 230 | 10.1 | 2.8 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 227 | 10.0 | 2.8 |
| 4 | Malignant neoplasms (C00-C97) | 196 | 8.6 | 2.4 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 78 | 3.4 | 1.0 |
| 6 | Influenza and pneumonia (J09-J18) | 49 | 2.1 | 0.6 |
| 7 | Certain conditions originating in the perinatal period (P00-P96) | 34 | 1.5 | 0.4 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 27 | 1.2 | 0.3 |
| 9 | Cerebrovascular diseases (I60-I69) | 26 | 1.1 | 0.3 |
| 10 | Septicemia (A40-A41) | 24 | 1.1 | 0.3 |
| 11 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 17 | 0.7 | * |
| 12 | Anemias (D50-D64) | 12 | 0.5 | * |
| 12 | Complications of medical and surgical care (Y40-Y84,Y88) | 12 | 0.5 | * |
| 14 | Acute bronchitis and bronchiolitis (J20-J21) | 11 | 0.5 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 9 | 0.4 | * |
| ... | All other causes (residual) | 561 | 24.6 | 6.9 |
| All races and origins, male, 5-14 years | | | | |
| ... | All causes | 3,153 | 100.0 | 15.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 939 | 29.8 | 4.5 |
| 2 | Malignant neoplasms (C00-C97) | 469 | 14.9 | 2.2 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 255 | 8.1 | 1.2 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 194 | 6.2 | 0.9 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 194 | 6.2 | 0.9 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 108 | 3.4 | 0.5 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 104 | 3.3 | 0.5 |
| 8 | Cerebrovascular diseases (I60-I69) | 45 | 1.4 | 0.2 |
| 9 | Influenza and pneumonia (J09-J18) | 41 | 1.3 | 0.2 |
| 10 | Septicemia (A40-A41) | 33 | 1.0 | 0.2 |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 33 | 1.0 | 0.2 |
| 12 | Anemias (D50-D64) | 14 | 0.4 | * |
| 13 | Diabetes mellitus (E10-E14) | 11 | 0.3 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 11 | 0.3 | * |
| 13 | Certain conditions originating in the perinatal period (P00-P96) | 11 | 0.3 | * |
| ... | All other causes (residual) | 691 | 21.9 | 3.3 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| All races and origins, male, 15-24 years | | | | |
| ... | All causes | 22,346 | 100.0 | 99.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 9,149 | 40.9 | 40.7 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 4,359 | 19.5 | 19.4 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 4,134 | 18.5 | 18.4 |
| 4 | Malignant neoplasms (C00-C97) | 880 | 3.9 | 3.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 667 | 3.0 | 3.0 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 220 | 1.0 | 1.0 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 130 | 0.6 | 0.6 |
| 8 | Diabetes mellitus (E10-E14) | 118 | 0.5 | 0.5 |
| 9 | Influenza and pneumonia (J09-J18) | 107 | 0.5 | 0.5 |
| 10 | Cerebrovascular diseases (I60-I69) | 89 | 0.4 | 0.4 |
| 11 | Legal intervention (Y35,Y89.0) | 84 | 0.4 | 0.4 |
| 12 | Septicemia (A40-A41) | 71 | 0.3 | 0.3 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 64 | 0.3 | 0.3 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 46 | 0.2 | 0.2 |
| 15 | Anemias (D50-D64) | 42 | 0.2 | 0.2 |
| ... | All other causes (residual) | 2,186 | 9.8 | 9.7 |
| All races and origins, male, 25-34 years | | | | |
| ... | All causes | 35,781 | 100.0 | 160.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 14,659 | 41.0 | 65.7 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 5,503 | 15.4 | 24.7 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 4,097 | 11.5 | 18.4 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,297 | 6.4 | 10.3 |
| 5 | Malignant neoplasms (C00-C97) | 1,818 | 5.1 | 8.2 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 520 | 1.5 | 2.3 |
| 7 | Diabetes mellitus (E10-E14) | 464 | 1.3 | 2.1 |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 380 | 1.1 | 1.7 |
| 9 | Cerebrovascular diseases (I60-I69) | 320 | 0.9 | 1.4 |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 262 | 0.7 | 1.2 |
| 11 | Influenza and pneumonia (J09-J18) | 234 | 0.7 | 1.0 |
| 12 | Septicemia (A40-A41) | 193 | 0.5 | 0.9 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 162 | 0.5 | 0.7 |
| 14 | Legal intervention (Y35,Y89.0) | 157 | 0.4 | 0.7 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 140 | 0.4 | 0.6 |
| ... | All other causes (residual) | 4,575 | 12.8 | 20.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|-------|
| All races and origins, male, 35-44 years | | | | |
| ... | All causes | 45,670 | 100.0 | 226.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 12,566 | 27.5 | 62.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,094 | 15.5 | 35.1 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 5,223 | 11.4 | 25.9 |
| 4 | Malignant neoplasms (C00-C97) | 4,656 | 10.2 | 23.0 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,333 | 5.1 | 11.5 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,811 | 4.0 | 9.0 |
| 7 | Diabetes mellitus (E10-E14) | 1,242 | 2.7 | 6.1 |
| 8 | Cerebrovascular diseases (I60-I69) | 990 | 2.2 | 4.9 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 691 | 1.5 | 3.4 |
| 10 | Influenza and pneumonia (J09-J18) | 402 | 0.9 | 2.0 |
| 11 | Septicemia (A40-A41) | 398 | 0.9 | 2.0 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 370 | 0.8 | 1.8 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 334 | 0.7 | 1.7 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 312 | 0.7 | 1.5 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 265 | 0.6 | 1.3 |
| ... | All other causes (residual) | 6,983 | 15.3 | 34.6 |
| All races and origins, male, 45-54 years | | | | |
| ... | All causes | 105,547 | 100.0 | 495.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 23,871 | 22.6 | 112.1 |
| 2 | Malignant neoplasms (C00-C97) | 21,171 | 20.1 | 99.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 14,364 | 13.6 | 67.4 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 6,411 | 6.1 | 30.1 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 5,765 | 5.5 | 27.1 |
| 6 | Diabetes mellitus (E10-E14) | 3,884 | 3.7 | 18.2 |
| 7 | Cerebrovascular diseases (I60-I69) | 2,900 | 2.7 | 13.6 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 2,004 | 1.9 | 9.4 |
| 9 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,626 | 1.5 | 7.6 |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,515 | 1.4 | 7.1 |
| 11 | Septicemia (A40-A41) | 1,325 | 1.3 | 6.2 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,265 | 1.2 | 5.9 |
| 13 | Influenza and pneumonia (J09-J18) | 1,169 | 1.1 | 5.5 |
| 14 | Viral hepatitis (B15-B19) | 1,027 | 1.0 | 4.8 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 945 | 0.9 | 4.4 |
| ... | All other causes (residual) | 16,305 | 15.4 | 76.6 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| All races and origins, male, 55-64 years | | | | |
| ... | All causes | 217,626 | 100.0 | 1,103.9 |
| 1 | Malignant neoplasms (C00-C97) | 64,180 | 29.5 | 325.5 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 53,108 | 24.4 | 269.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 13,433 | 6.2 | 68.1 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 9,076 | 4.2 | 46.0 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 9,005 | 4.1 | 45.7 |
| 6 | Diabetes mellitus (E10-E14) | 8,686 | 4.0 | 44.1 |
| 7 | Cerebrovascular diseases (I60-I69) | 6,970 | 3.2 | 35.4 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 5,696 | 2.6 | 28.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,183 | 1.5 | 16.1 |
| 10 | Septicemia (A40-A41) | 3,103 | 1.4 | 15.7 |
| 11 | Influenza and pneumonia (J09-J18) | 2,583 | 1.2 | 13.1 |
| 12 | Viral hepatitis (B15-B19) | 2,511 | 1.2 | 12.7 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,279 | 1.0 | 11.6 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,420 | 0.7 | 7.2 |
| 15 | Aortic aneurysm and dissection (I71) | 967 | 0.4 | 4.9 |
| ... | All other causes (residual) | 31,426 | 14.4 | 159.4 |
| All races and origins, male, 65-74 years | | | | |
| ... | All causes | 282,347 | 100.0 | 2,190.0 |
| 1 | Malignant neoplasms (C00-C97) | 90,468 | 32.0 | 701.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 68,285 | 24.2 | 529.7 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 19,129 | 6.8 | 148.4 |
| 4 | Diabetes mellitus (E10-E14) | 11,358 | 4.0 | 88.1 |
| 5 | Cerebrovascular diseases (I60-I69) | 11,120 | 3.9 | 86.3 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8,388 | 3.0 | 65.1 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 5,348 | 1.9 | 41.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 5,278 | 1.9 | 40.9 |
| 9 | Septicemia (A40-A41) | 4,500 | 1.6 | 34.9 |
| 10 | Influenza and pneumonia (J09-J18) | 4,413 | 1.6 | 34.2 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 3,372 | 1.2 | 26.2 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,731 | 1.0 | 21.2 |
| 13 | Alzheimer's disease (G30) | 2,592 | 0.9 | 20.1 |
| 14 | Parkinson's disease (G20-G21) | 2,570 | 0.9 | 19.9 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,863 | 0.7 | 14.5 |
| ... | All other causes (residual) | 40,932 | 14.5 | 317.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|----------|
| All races and origins, male, 75-84 years | | | | |
| ... | All causes | 323,846 | 100.0 | 5,376.3 |
| 1 | Malignant neoplasms (C00-C97) | 81,669 | 25.2 | 1,355.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 81,586 | 25.2 | 1,354.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 23,652 | 7.3 | 392.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 17,000 | 5.2 | 282.2 |
| 5 | Alzheimer's disease (G30) | 11,104 | 3.4 | 184.3 |
| 6 | Diabetes mellitus (E10-E14) | 10,337 | 3.2 | 171.6 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8,524 | 2.6 | 141.5 |
| 8 | Parkinson's disease (G20-G21) | 7,356 | 2.3 | 122.1 |
| 9 | Influenza and pneumonia (J09-J18) | 7,263 | 2.2 | 120.6 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7,181 | 2.2 | 119.2 |
| 11 | Septicemia (A40-A41) | 5,055 | 1.6 | 83.9 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 3,313 | 1.0 | 55.0 |
| 13 | Pneumonitis due to solids and liquids (J69) | 2,973 | 0.9 | 49.4 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2,707 | 0.8 | 44.9 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,378 | 0.7 | 39.5 |
| ... | All other causes (residual) | 51,748 | 16.0 | 859.1 |
| All races and origins, male, 85 years and over | | | | |
| ... | All causes | 321,704 | 100.0 | 14,795.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 97,736 | 30.4 | 4,495.1 |
| 2 | Malignant neoplasms (C00-C97) | 48,274 | 15.0 | 2,220.2 |
| 3 | Alzheimer's disease (G30) | 19,529 | 6.1 | 898.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 18,770 | 5.8 | 863.3 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 17,933 | 5.6 | 824.8 |
| 6 | Influenza and pneumonia (J09-J18) | 10,528 | 3.3 | 484.2 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 9,360 | 2.9 | 430.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7,937 | 2.5 | 365.0 |
| 9 | Diabetes mellitus (E10-E14) | 7,016 | 2.2 | 322.7 |
| 10 | Parkinson's disease (G20-G21) | 6,462 | 2.0 | 297.2 |
| 11 | Pneumonitis due to solids and liquids (J69) | 4,939 | 1.5 | 227.2 |
| 12 | Septicemia (A40-A41) | 4,584 | 1.4 | 210.8 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 4,271 | 1.3 | 196.4 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2,432 | 0.8 | 111.9 |
| 15 | Aortic aneurysm and dissection (I71) | 1,187 | 0.4 | 54.6 |
| ... | All other causes (residual) | 60,746 | 18.9 | 2,793.8 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|-----------|-------------------------|-------|
| All races and origins, female, all ages | | | | |
| ... | All causes | 1,339,226 | 100.0 | 820.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 298,840 | 22.3 | 183.1 |
| 2 | Malignant neoplasms (C00-C97) | 282,112 | 21.1 | 172.9 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 82,543 | 6.2 | 50.6 |
| 4 | Cerebrovascular diseases (I60-I69) | 82,035 | 6.1 | 50.3 |
| 5 | Alzheimer's disease (G30) | 76,871 | 5.7 | 47.1 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 53,652 | 4.0 | 32.9 |
| 7 | Diabetes mellitus (E10-E14) | 36,412 | 2.7 | 22.3 |
| 8 | Influenza and pneumonia (J09-J18) | 30,159 | 2.3 | 18.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 24,518 | 1.8 | 15.0 |
| 10 | Septicemia (A40-A41) | 21,388 | 1.6 | 13.1 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 18,266 | 1.4 | 11.2 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 14,660 | 1.1 | 9.0 |
| 13 | Parkinson's disease (G20-G21) | 11,105 | 0.8 | 6.8 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 10,199 | 0.8 | 6.2 |
| 15 | Pneumonitis due to solids and liquids (J69) | 8,892 | 0.7 | 5.4 |
| ... | All other causes (residual) | 287,574 | 21.5 | 176.2 |
| All races and origins, female, 1-4 years | | | | |
| ... | All causes | 1,684 | 100.0 | 21.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 467 | 27.7 | 6.0 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 208 | 12.4 | 2.7 |
| 3 | Malignant neoplasms (C00-C97) | 158 | 9.4 | 2.0 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 139 | 8.3 | 1.8 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 69 | 4.1 | 0.9 |
| 6 | Influenza and pneumonia (J09-J18) | 39 | 2.3 | 0.5 |
| 7 | Septicemia (A40-A41) | 30 | 1.8 | 0.4 |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 17 | 1.0 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 16 | 1.0 | * |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 16 | 1.0 | * |
| 11 | Anemias (D50-D64) | 13 | 0.8 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 13 | 0.8 | * |
| 13 | Meningitis (G00,G03) | 11 | 0.7 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7 | 0.4 | * |
| 15 | Acute bronchitis and bronchiolitis (J20-J21) | 6 | 0.4 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 6 | 0.4 | * |
| ... | All other causes (residual) | 469 | 27.9 | 6.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| All races and origins, female, 5-14 years | | | | |
| ... | All causes | 2,258 | 100.0 | 11.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 579 | 25.6 | 2.9 |
| 2 | Malignant neoplasms (C00-C97) | 396 | 17.5 | 2.0 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 158 | 7.0 | 0.8 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 143 | 6.3 | 0.7 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 104 | 4.6 | 0.5 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 102 | 4.5 | 0.5 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 69 | 3.1 | 0.3 |
| 8 | Influenza and pneumonia (J09-J18) | 42 | 1.9 | 0.2 |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 39 | 1.7 | 0.2 |
| 9 | Cerebrovascular diseases (I60-I69) | 39 | 1.7 | 0.2 |
| 11 | Septicemia (A40-A41) | 31 | 1.4 | 0.2 |
| 12 | Anemias (D50-D64) | 19 | 0.8 | * |
| 13 | Diabetes mellitus (E10-E14) | 12 | 0.5 | * |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 10 | 0.4 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 9 | 0.4 | * |
| ... | All other causes (residual) | 506 | 22.4 | 2.5 |
| All races and origins, female, 15-24 years | | | | |
| ... | All causes | 8,148 | 100.0 | 38.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,365 | 41.3 | 15.7 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,132 | 13.9 | 5.3 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 599 | 7.4 | 2.8 |
| 4 | Malignant neoplasms (C00-C97) | 589 | 7.2 | 2.8 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 330 | 4.1 | 1.5 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 166 | 2.0 | 0.8 |
| 7 | Pregnancy, childbirth and the puerperium (O00-O99) | 161 | 2.0 | 0.8 |
| 8 | Diabetes mellitus (E10-E14) | 78 | 1.0 | 0.4 |
| 9 | Cerebrovascular diseases (I60-I69) | 77 | 0.9 | 0.4 |
| 9 | Influenza and pneumonia (J09-J18) | 77 | 0.9 | 0.4 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 72 | 0.9 | 0.3 |
| 12 | Septicemia (A40-A41) | 59 | 0.7 | 0.3 |
| 13 | Anemias (D50-D64) | 42 | 0.5 | 0.2 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 37 | 0.5 | 0.2 |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 36 | 0.4 | 0.2 |
| ... | All other causes (residual) | 1,328 | 16.3 | 6.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| All races and origins, female, 25-34 years | | | | |
| ... | All causes | 15,736 | 100.0 | 72.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,136 | 32.6 | 23.5 |
| 2 | Malignant neoplasms (C00-C97) | 1,886 | 12.0 | 8.6 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,444 | 9.2 | 6.6 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,225 | 7.8 | 5.6 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 766 | 4.9 | 3.5 |
| 6 | Pregnancy, childbirth and the puerperium (O00-O99) | 424 | 2.7 | 1.9 |
| 7 | Diabetes mellitus (E10-E14) | 334 | 2.1 | 1.5 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 324 | 2.1 | 1.5 |
| 9 | Cerebrovascular diseases (I60-I69) | 247 | 1.6 | 1.1 |
| 10 | Septicemia (A40-A41) | 186 | 1.2 | 0.9 |
| 11 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 181 | 1.2 | 0.8 |
| 12 | Influenza and pneumonia (J09-J18) | 163 | 1.0 | 0.7 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 149 | 0.9 | 0.7 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 126 | 0.8 | 0.6 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 122 | 0.8 | 0.6 |
| ... | All other causes (residual) | 3,023 | 19.2 | 13.8 |
| All races and origins, female, 35-44 years | | | | |
| ... | All causes | 27,418 | 100.0 | 134.5 |
| 1 | Malignant neoplasms (C00-C97) | 6,253 | 22.8 | 30.7 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,252 | 19.2 | 25.8 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,293 | 12.0 | 16.2 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,713 | 6.2 | 8.4 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,050 | 3.8 | 5.2 |
| 6 | Cerebrovascular diseases (I60-I69) | 798 | 2.9 | 3.9 |
| 7 | Diabetes mellitus (E10-E14) | 744 | 2.7 | 3.6 |
| 8 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 562 | 2.0 | 2.8 |
| 9 | Septicemia (A40-A41) | 431 | 1.6 | 2.1 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 368 | 1.3 | 1.8 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 364 | 1.3 | 1.8 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 316 | 1.2 | 1.6 |
| 13 | Pregnancy, childbirth and the puerperium (O00-O99) | 310 | 1.1 | 1.5 |
| 14 | Influenza and pneumonia (J09-J18) | 306 | 1.1 | 1.5 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 218 | 0.8 | 1.1 |
| ... | All other causes (residual) | 5,440 | 19.8 | 26.7 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|-------|
| All races and origins, female, 45-54 years | | | | |
| ... | All causes | 68,947 | 100.0 | 315.0 |
| 1 | Malignant neoplasms (C00-C97) | 21,883 | 31.7 | 100.0 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,377 | 15.1 | 47.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 7,135 | 10.3 | 32.6 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,109 | 4.5 | 14.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 2,407 | 3.5 | 11.0 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 2,341 | 3.4 | 10.7 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,340 | 3.4 | 10.7 |
| 8 | Diabetes mellitus (E10-E14) | 2,328 | 3.4 | 10.6 |
| 9 | Septicemia (A40-A41) | 1,217 | 1.8 | 5.6 |
| 10 | Influenza and pneumonia (J09-J18) | 881 | 1.3 | 4.0 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 859 | 1.2 | 3.9 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 532 | 0.8 | 2.4 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 512 | 0.7 | 2.3 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 480 | 0.7 | 2.2 |
| 15 | Viral hepatitis (B15-B19) | 469 | 0.7 | 2.1 |
| ... | All other causes (residual) | 12,077 | 17.5 | 55.2 |
| All races and origins, female, 55-64 years | | | | |
| ... | All causes | 140,159 | 100.0 | 662.3 |
| 1 | Malignant neoplasms (C00-C97) | 51,942 | 37.1 | 245.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 23,764 | 17.0 | 112.3 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 8,452 | 6.0 | 39.9 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 6,055 | 4.3 | 28.6 |
| 5 | Diabetes mellitus (E10-E14) | 5,480 | 3.9 | 25.9 |
| 6 | Cerebrovascular diseases (I60-I69) | 5,146 | 3.7 | 24.3 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 4,202 | 3.0 | 19.9 |
| 8 | Septicemia (A40-A41) | 2,671 | 1.9 | 12.6 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,269 | 1.6 | 10.7 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,043 | 1.5 | 9.7 |
| 11 | Influenza and pneumonia (J09-J18) | 2,018 | 1.4 | 9.5 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,316 | 0.9 | 6.2 |
| 13 | Viral hepatitis (B15-B19) | 1,048 | 0.7 | 5.0 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 692 | 0.5 | 3.3 |
| 15 | Pneumonitis due to solids and liquids (J69) | 561 | 0.4 | 2.7 |
| ... | All other causes (residual) | 22,500 | 16.1 | 106.3 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| All races and origins, female, 65-74 years | | | | |
| ... | All causes | 212,669 | 100.0 | 1,450.9 |
| 1 | Malignant neoplasms (C00-C97) | 73,260 | 34.4 | 499.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 39,018 | 18.3 | 266.2 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 18,513 | 8.7 | 126.3 |
| 4 | Cerebrovascular diseases (I60-I69) | 9,673 | 4.5 | 66.0 |
| 5 | Diabetes mellitus (E10-E14) | 8,095 | 3.8 | 55.2 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,573 | 2.2 | 31.2 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4,384 | 2.1 | 29.9 |
| 8 | Septicemia (A40-A41) | 4,155 | 2.0 | 28.3 |
| 9 | Influenza and pneumonia (J09-J18) | 3,718 | 1.7 | 25.4 |
| 10 | Alzheimer's disease (G30) | 3,575 | 1.7 | 24.4 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,061 | 1.4 | 20.9 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,248 | 1.1 | 15.3 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,293 | 0.6 | 8.8 |
| 14 | Parkinson's disease (G20-G21) | 1,243 | 0.6 | 8.5 |
| 15 | Pneumonitis due to solids and liquids (J69) | 1,024 | 0.5 | 7.0 |
| ... | All other causes (residual) | 34,836 | 16.4 | 237.7 |
| All races and origins, female, 75-84 years | | | | |
| ... | All causes | 313,720 | 100.0 | 3,971.3 |
| 1 | Malignant neoplasms (C00-C97) | 71,599 | 22.8 | 906.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 67,613 | 21.6 | 855.9 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 26,180 | 8.3 | 331.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 21,012 | 6.7 | 266.0 |
| 5 | Alzheimer's disease (G30) | 18,399 | 5.9 | 232.9 |
| 6 | Diabetes mellitus (E10-E14) | 9,567 | 3.0 | 121.1 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 6,994 | 2.2 | 88.5 |
| 8 | Influenza and pneumonia (J09-J18) | 6,886 | 2.2 | 87.2 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 6,700 | 2.1 | 84.8 |
| 10 | Septicemia (A40-A41) | 5,426 | 1.7 | 68.7 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 4,220 | 1.3 | 53.4 |
| 12 | Parkinson's disease (G20-G21) | 4,117 | 1.3 | 52.1 |
| 13 | Pneumonitis due to solids and liquids (J69) | 2,179 | 0.7 | 27.6 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2,130 | 0.7 | 27.0 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,068 | 0.7 | 26.2 |
| ... | All other causes (residual) | 58,630 | 18.7 | 742.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|----------|
| All races and origins, female, 85 years and over | | | | |
| ... | All causes | 537,997 | 100.0 | 13,080.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 152,900 | 28.4 | 3,717.6 |
| 2 | Alzheimer's disease (G30) | 54,296 | 10.1 | 1,320.2 |
| 3 | Malignant neoplasms (C00-C97) | 54,119 | 10.1 | 1,315.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 42,581 | 7.9 | 1,035.3 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 26,397 | 4.9 | 641.8 |
| 6 | Influenza and pneumonia (J09-J18) | 15,966 | 3.0 | 388.2 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 13,556 | 2.5 | 329.6 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 9,778 | 1.8 | 237.7 |
| 9 | Diabetes mellitus (E10-E14) | 9,769 | 1.8 | 237.5 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 9,732 | 1.8 | 236.6 |
| 11 | Septicemia (A40-A41) | 7,097 | 1.3 | 172.6 |
| 12 | Parkinson's disease (G20-G21) | 5,521 | 1.0 | 134.2 |
| 13 | Pneumonitis due to solids and liquids (J69) | 4,797 | 0.9 | 116.6 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2,914 | 0.5 | 70.9 |
| 15 | Atherosclerosis (I70) | 2,244 | 0.4 | 54.6 |
| ... | All other causes (residual) | 126,330 | 23.5 | 3,071.6 |
| Hispanic, both sexes, all ages | | | | |
| ... | All causes | 179,457 | 100.0 | 317.1 |
| 1 | Malignant neoplasms (C00-C97) | 37,804 | 21.1 | 66.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 36,401 | 20.3 | 64.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 13,806 | 7.7 | 24.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 9,795 | 5.5 | 17.3 |
| 5 | Diabetes mellitus (E10-E14) | 8,278 | 4.6 | 14.6 |
| 6 | Alzheimer's disease (G30) | 6,444 | 3.6 | 11.4 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 6,018 | 3.4 | 10.6 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 5,159 | 2.9 | 9.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,581 | 2.0 | 6.3 |
| 10 | Influenza and pneumonia (J09-J18) | 3,497 | 1.9 | 6.2 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 3,303 | 1.8 | 5.8 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,886 | 1.6 | 5.1 |
| 13 | Septicemia (A40-A41) | 2,692 | 1.5 | 4.8 |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 2,454 | 1.4 | 4.3 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,384 | 1.3 | 4.2 |
| ... | All other causes (residual) | 34,955 | 19.5 | 61.8 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|----------------------------------|---|--------|-------------------------|------|
| Hispanic, both sexes, 1-4 years | | | | |
| ... | All causes | 815 | 100.0 | 19.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 258 | 31.7 | 6.3 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 115 | 14.1 | 2.8 |
| 3 | Malignant neoplasms (C00-C97) | 75 | 9.2 | 1.8 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 69 | 8.5 | 1.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 29 | 3.6 | 0.7 |
| 6 | Influenza and pneumonia (J09-J18) | 18 | 2.2 | * |
| 7 | Septicemia (A40-A41) | 17 | 2.1 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 7 | 0.9 | * |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 7 | 0.9 | * |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 6 | 0.7 | * |
| 11 | Anemias (D50-D64) | 4 | 0.5 | * |
| 11 | Meningitis (G00,G03) | 4 | 0.5 | * |
| 11 | Acute bronchitis and bronchiolitis (J20-J21) | 4 | 0.5 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 4 | 0.5 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 0.4 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 3 | 0.4 | * |
| ... | All other causes (residual) | 192 | 23.6 | 4.7 |
| Hispanic, both sexes, 5-14 years | | | | |
| ... | All causes | 1,063 | 100.0 | 10.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 308 | 29.0 | 3.0 |
| 2 | Malignant neoplasms (C00-C97) | 211 | 19.8 | 2.1 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 69 | 6.5 | 0.7 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 53 | 5.0 | 0.5 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 46 | 4.3 | 0.5 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 35 | 3.3 | 0.3 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 23 | 2.2 | 0.2 |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 17 | 1.6 | * |
| 9 | Septicemia (A40-A41) | 16 | 1.5 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 16 | 1.5 | * |
| 11 | Influenza and pneumonia (J09-J18) | 14 | 1.3 | * |
| 12 | Anemias (D50-D64) | 5 | 0.5 | * |
| 13 | Meningitis (G00,G03) | 4 | 0.4 | * |
| 13 | Certain conditions originating in the perinatal period (P00-P96) | 4 | 0.4 | * |
| 15 | Diabetes mellitus (E10-E14) | 3 | 0.3 | * |
| 15 | Acute bronchitis and bronchiolitis (J20-J21) | 3 | 0.3 | * |
| 15 | Pneumonitis due to solids and liquids (J69) | 3 | 0.3 | * |
| ... | All other causes (residual) | 233 | 21.9 | 2.3 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|------|
| Hispanic, both sexes, 15-24 years | | | | |
| ... | All causes | 5,353 | 100.0 | 56.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,261 | 42.2 | 23.7 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 921 | 17.2 | 9.7 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 807 | 15.1 | 8.5 |
| 4 | Malignant neoplasms (C00-C97) | 332 | 6.2 | 3.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 145 | 2.7 | 1.5 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 69 | 1.3 | 0.7 |
| 7 | Influenza and pneumonia (J09-J18) | 36 | 0.7 | 0.4 |
| 8 | Cerebrovascular diseases (I60-I69) | 32 | 0.6 | 0.3 |
| 9 | Septicemia (A40-A41) | 30 | 0.6 | 0.3 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 29 | 0.5 | 0.3 |
| 11 | Pregnancy, childbirth and the puerperium (O00-O99) | 27 | 0.5 | 0.3 |
| 12 | Diabetes mellitus (E10-E14) | 20 | 0.4 | 0.2 |
| 13 | Legal intervention (Y35,Y89.0) | 17 | 0.3 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 13 | 0.2 | * |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 12 | 0.2 | * |
| ... | All other causes (residual) | 602 | 11.2 | 6.3 |
| Hispanic, both sexes, 25-34 years | | | | |
| ... | All causes | 7,182 | 100.0 | 79.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,653 | 36.9 | 29.5 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 791 | 11.0 | 8.8 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 764 | 10.6 | 8.5 |
| 4 | Malignant neoplasms (C00-C97) | 714 | 9.9 | 7.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 439 | 6.1 | 4.9 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 169 | 2.4 | 1.9 |
| 7 | Cerebrovascular diseases (I60-I69) | 121 | 1.7 | 1.3 |
| 8 | Diabetes mellitus (E10-E14) | 106 | 1.5 | 1.2 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 96 | 1.3 | 1.1 |
| 10 | Pregnancy, childbirth and the puerperium (O00-O99) | 59 | 0.8 | 0.7 |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 59 | 0.8 | 0.7 |
| 12 | Influenza and pneumonia (J09-J18) | 56 | 0.8 | 0.6 |
| 13 | Septicemia (A40-A41) | 46 | 0.6 | 0.5 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 44 | 0.6 | 0.5 |
| 15 | Chronic lower respiratory diseases (J40-J47) | 39 | 0.5 | 0.4 |
| 15 | Legal intervention (Y35,Y89.0) | 39 | 0.5 | 0.4 |
| ... | All other causes (residual) | 987 | 13.7 | 11.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|-------|
| Hispanic, both sexes, 35-44 years | | | | |
| ... | All causes | 9,490 | 100.0 | 116.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,190 | 23.1 | 26.8 |
| 2 | Malignant neoplasms (C00-C97) | 1,718 | 18.1 | 21.0 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,082 | 11.4 | 13.2 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 594 | 6.3 | 7.3 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 587 | 6.2 | 7.2 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 506 | 5.3 | 6.2 |
| 7 | Cerebrovascular diseases (I60-I69) | 321 | 3.4 | 3.9 |
| 8 | Diabetes mellitus (E10-E14) | 274 | 2.9 | 3.4 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 191 | 2.0 | 2.3 |
| 10 | Septicemia (A40-A41) | 105 | 1.1 | 1.3 |
| 11 | Influenza and pneumonia (J09-J18) | 102 | 1.1 | 1.2 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 94 | 1.0 | 1.1 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 56 | 0.6 | 0.7 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 50 | 0.5 | 0.6 |
| 15 | Pregnancy, childbirth and the puerperium (O00-O99) | 48 | 0.5 | 0.6 |
| ... | All other causes (residual) | 1,572 | 16.6 | 19.2 |
| Hispanic, both sexes, 45-54 years | | | | |
| ... | All causes | 17,189 | 100.0 | 262.8 |
| 1 | Malignant neoplasms (C00-C97) | 4,286 | 24.9 | 65.5 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,844 | 16.5 | 43.5 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,090 | 12.2 | 32.0 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,477 | 8.6 | 22.6 |
| 5 | Diabetes mellitus (E10-E14) | 772 | 4.5 | 11.8 |
| 6 | Cerebrovascular diseases (I60-I69) | 665 | 3.9 | 10.2 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 504 | 2.9 | 7.7 |
| 8 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 298 | 1.7 | 4.6 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 296 | 1.7 | 4.5 |
| 10 | Viral hepatitis (B15-B19) | 280 | 1.6 | 4.3 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 271 | 1.6 | 4.1 |
| 12 | Septicemia (A40-A41) | 269 | 1.6 | 4.1 |
| 13 | Influenza and pneumonia (J09-J18) | 209 | 1.2 | 3.2 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 167 | 1.0 | 2.6 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 131 | 0.8 | 2.0 |
| ... | All other causes (residual) | 2,630 | 15.3 | 40.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|---------|
| Hispanic, both sexes, 55-64 years | | | | |
| ... | All causes | 25,987 | 100.0 | 605.7 |
| 1 | Malignant neoplasms (C00-C97) | 7,908 | 30.4 | 184.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,096 | 19.6 | 118.8 |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,767 | 6.8 | 41.2 |
| 4 | Diabetes mellitus (E10-E14) | 1,616 | 6.2 | 37.7 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,437 | 5.5 | 33.5 |
| 6 | Cerebrovascular diseases (I60-I69) | 1,107 | 4.3 | 25.8 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 511 | 2.0 | 11.9 |
| 8 | Viral hepatitis (B15-B19) | 493 | 1.9 | 11.5 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 492 | 1.9 | 11.5 |
| 10 | Septicemia (A40-A41) | 421 | 1.6 | 9.8 |
| 11 | Influenza and pneumonia (J09-J18) | 352 | 1.4 | 8.2 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 318 | 1.2 | 7.4 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 303 | 1.2 | 7.1 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 208 | 0.8 | 4.8 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 135 | 0.5 | 3.1 |
| ... | All other causes (residual) | 3,823 | 14.7 | 89.1 |
| Hispanic, both sexes, 65-74 years | | | | |
| ... | All causes | 30,275 | 100.0 | 1,332.4 |
| 1 | Malignant neoplasms (C00-C97) | 9,133 | 30.2 | 401.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,517 | 21.5 | 286.8 |
| 3 | Diabetes mellitus (E10-E14) | 2,056 | 6.8 | 90.5 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,561 | 5.2 | 68.7 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,153 | 3.8 | 50.7 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 957 | 3.2 | 42.1 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 833 | 2.8 | 36.7 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 800 | 2.6 | 35.2 |
| 9 | Septicemia (A40-A41) | 569 | 1.9 | 25.0 |
| 10 | Influenza and pneumonia (J09-J18) | 556 | 1.8 | 24.5 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 430 | 1.4 | 18.9 |
| 12 | Alzheimer's disease (G30) | 401 | 1.3 | 17.6 |
| 13 | Parkinson's disease (G20-G21) | 242 | 0.8 | 10.7 |
| 14 | Viral hepatitis (B15-B19) | 225 | 0.7 | 9.9 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 170 | 0.6 | 7.5 |
| ... | All other causes (residual) | 4,672 | 15.4 | 205.6 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|---------|
| Hispanic, both sexes, 75-84 years | | | | |
| ... | All causes | 37,089 | 100.0 | 3,430.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8,835 | 23.8 | 817.1 |
| 2 | Malignant neoplasms (C00-C97) | 8,490 | 22.9 | 785.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,648 | 7.1 | 244.9 |
| 4 | Diabetes mellitus (E10-E14) | 2,053 | 5.5 | 189.9 |
| 5 | Alzheimer's disease (G30) | 1,855 | 5.0 | 171.6 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 1,626 | 4.4 | 150.4 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 977 | 2.6 | 90.4 |
| 8 | Influenza and pneumonia (J09-J18) | 939 | 2.5 | 86.8 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 756 | 2.0 | 69.9 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 654 | 1.8 | 60.5 |
| 11 | Parkinson's disease (G20-G21) | 633 | 1.7 | 58.5 |
| 12 | Septicemia (A40-A41) | 618 | 1.7 | 57.2 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 605 | 1.6 | 56.0 |
| 14 | Pneumonitis due to solids and liquids (J69) | 280 | 0.8 | 25.9 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 235 | 0.6 | 21.7 |
| ... | All other causes (residual) | 5,885 | 15.9 | 544.3 |
| Hispanic, both sexes, 85 years and over | | | | |
| ... | All causes | 40,203 | 100.0 | 9,585.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 11,334 | 28.2 | 2,702.3 |
| 2 | Malignant neoplasms (C00-C97) | 4,924 | 12.2 | 1,174.0 |
| 3 | Alzheimer's disease (G30) | 4,101 | 10.2 | 977.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 3,297 | 8.2 | 786.1 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,770 | 4.4 | 422.0 |
| 6 | Diabetes mellitus (E10-E14) | 1,375 | 3.4 | 327.8 |
| 7 | Influenza and pneumonia (J09-J18) | 1,185 | 2.9 | 282.5 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 865 | 2.2 | 206.2 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 830 | 2.1 | 197.9 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 818 | 2.0 | 195.0 |
| 11 | Parkinson's disease (G20-G21) | 609 | 1.5 | 145.2 |
| 12 | Septicemia (A40-A41) | 560 | 1.4 | 133.5 |
| 13 | Pneumonitis due to solids and liquids (J69) | 410 | 1.0 | 97.8 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 196 | 0.5 | 46.7 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 195 | 0.5 | 46.5 |
| ... | All other causes (residual) | 7,734 | 19.2 | 1,843.9 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---------------------------|---|--------|-------------------------|-------|
| Hispanic, male, all ages | | | | |
| ... | All causes | 98,170 | 100.0 | 343.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 20,225 | 20.6 | 70.7 |
| 2 | Malignant neoplasms (C00-C97) | 19,847 | 20.2 | 69.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 10,067 | 10.3 | 35.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 4,544 | 4.6 | 15.9 |
| 5 | Diabetes mellitus (E10-E14) | 4,426 | 4.5 | 15.5 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 4,109 | 4.2 | 14.4 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 2,606 | 2.7 | 9.1 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,587 | 2.6 | 9.0 |
| 9 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,391 | 2.4 | 8.4 |
| 10 | Alzheimer's disease (G30) | 2,035 | 2.1 | 7.1 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,837 | 1.9 | 6.4 |
| 12 | Influenza and pneumonia (J09-J18) | 1,735 | 1.8 | 6.1 |
| 13 | Septicemia (A40-A41) | 1,370 | 1.4 | 4.8 |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 1,368 | 1.4 | 4.8 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,139 | 1.2 | 4.0 |
| ... | All other causes (residual) | 17,884 | 18.2 | 62.5 |
| Hispanic, male, 1-4 years | | | | |
| ... | All causes | 461 | 100.0 | 22.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 154 | 33.4 | 7.3 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 58 | 12.6 | 2.8 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 45 | 9.8 | 2.1 |
| 4 | Malignant neoplasms (C00-C97) | 40 | 8.7 | 1.9 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 21 | 4.6 | 1.0 |
| 6 | Influenza and pneumonia (J09-J18) | 12 | 2.6 | * |
| 7 | Septicemia (A40-A41) | 8 | 1.7 | * |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 5 | 1.1 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 3 | 0.7 | * |
| 9 | Chronic lower respiratory diseases (J40-J47) | 3 | 0.7 | * |
| 11 | Meningitis (G00,G03) | 2 | 0.4 | * |
| 11 | Acute bronchitis and bronchiolitis (J20-J21) | 2 | 0.4 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2 | 0.4 | * |
| 14 | Salmonella infections (A01-A02) | 1 | 0.2 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1 | 0.2 | * |
| 14 | Anemias (D50-D64) | 1 | 0.2 | * |
| 14 | Diabetes mellitus (E10-E14) | 1 | 0.2 | * |
| 14 | Nutritional deficiencies (E40-E64) | 1 | 0.2 | * |
| 14 | Pneumonitis due to solids and liquids (J69) | 1 | 0.2 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 1 | 0.2 | * |
| ... | All other causes (residual) | 99 | 21.5 | 4.7 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|------|
| Hispanic, male, 5-14 years | | | | |
| ... | All causes | 633 | 100.0 | 12.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 206 | 32.5 | 4.0 |
| 2 | Malignant neoplasms (C00-C97) | 121 | 19.1 | 2.3 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 43 | 6.8 | 0.8 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 42 | 6.6 | 0.8 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 21 | 3.3 | 0.4 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 14 | 2.2 | * |
| 7 | Chronic lower respiratory diseases (J40-J47) | 13 | 2.1 | * |
| 8 | Septicemia (A40-A41) | 12 | 1.9 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 8 | 1.3 | * |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 6 | 0.9 | * |
| 11 | Influenza and pneumonia (J09-J18) | 5 | 0.8 | * |
| 12 | Anemias (D50-D64) | 2 | 0.3 | * |
| 12 | Meningitis (G00,G03) | 2 | 0.3 | * |
| 12 | Aortic aneurysm and dissection (I71) | 2 | 0.3 | * |
| 12 | Peptic ulcer (K25-K28) | 2 | 0.3 | * |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2 | 0.3 | * |
| 12 | Certain conditions originating in the perinatal period (P00-P96) | 2 | 0.3 | * |
| ... | All other causes (residual) | 130 | 20.5 | 2.5 |
| Hispanic, male, 15-24 years | | | | |
| ... | All causes | 4,039 | 100.0 | 82.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,748 | 43.3 | 35.6 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 809 | 20.0 | 16.5 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 629 | 15.6 | 12.8 |
| 4 | Malignant neoplasms (C00-C97) | 208 | 5.1 | 4.2 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 97 | 2.4 | 2.0 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 36 | 0.9 | 0.7 |
| 7 | Influenza and pneumonia (J09-J18) | 21 | 0.5 | 0.4 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 21 | 0.5 | 0.4 |
| 9 | Septicemia (A40-A41) | 19 | 0.5 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 19 | 0.5 | * |
| 11 | Legal intervention (Y35,Y89.0) | 17 | 0.4 | * |
| 12 | Diabetes mellitus (E10-E14) | 15 | 0.4 | * |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 9 | 0.2 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 6 | 0.1 | * |
| 15 | Aortic aneurysm and dissection (I71) | 5 | 0.1 | * |
| 15 | Pneumonitis due to solids and liquids (J69) | 5 | 0.1 | * |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 5 | 0.1 | * |
| ... | All other causes (residual) | 370 | 9.2 | 7.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|-------|
| Hispanic, male, 25-34 years | | | | |
| ... | All causes | 5,268 | 100.0 | 111.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,129 | 40.4 | 45.2 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 674 | 12.8 | 14.3 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 612 | 11.6 | 13.0 |
| 4 | Malignant neoplasms (C00-C97) | 361 | 6.9 | 7.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 318 | 6.0 | 6.8 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 126 | 2.4 | 2.7 |
| 7 | Cerebrovascular diseases (I60-I69) | 84 | 1.6 | 1.8 |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 78 | 1.5 | 1.7 |
| 9 | Diabetes mellitus (E10-E14) | 69 | 1.3 | 1.5 |
| 10 | Influenza and pneumonia (J09-J18) | 38 | 0.7 | 0.8 |
| 10 | Legal intervention (Y35,Y89.0) | 38 | 0.7 | 0.8 |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 30 | 0.6 | 0.6 |
| 13 | Septicemia (A40-A41) | 27 | 0.5 | 0.6 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 24 | 0.5 | 0.5 |
| 15 | Chronic lower respiratory diseases (J40-J47) | 22 | 0.4 | 0.5 |
| ... | All other causes (residual) | 638 | 12.1 | 13.5 |
| Hispanic, male, 35-44 years | | | | |
| ... | All causes | 6,385 | 100.0 | 152.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,729 | 27.1 | 41.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 810 | 12.7 | 19.4 |
| 3 | Malignant neoplasms (C00-C97) | 712 | 11.2 | 17.0 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 473 | 7.4 | 11.3 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 434 | 6.8 | 10.4 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 413 | 6.5 | 9.9 |
| 7 | Cerebrovascular diseases (I60-I69) | 203 | 3.2 | 4.9 |
| 8 | Diabetes mellitus (E10-E14) | 189 | 3.0 | 4.5 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 164 | 2.6 | 3.9 |
| 10 | Septicemia (A40-A41) | 66 | 1.0 | 1.6 |
| 11 | Influenza and pneumonia (J09-J18) | 63 | 1.0 | 1.5 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 52 | 0.8 | 1.2 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 42 | 0.7 | 1.0 |
| 14 | Aortic aneurysm and dissection (I71) | 37 | 0.6 | 0.9 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 32 | 0.5 | 0.8 |
| ... | All other causes (residual) | 966 | 15.1 | 23.1 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|-------|
| Hispanic, male, 45-54 years | | | | |
| ... | All causes | 11,230 | 100.0 | 340.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,139 | 19.0 | 64.8 |
| 2 | Malignant neoplasms (C00-C97) | 2,088 | 18.6 | 63.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,602 | 14.3 | 48.5 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,148 | 10.2 | 34.8 |
| 5 | Diabetes mellitus (E10-E14) | 508 | 4.5 | 15.4 |
| 6 | Cerebrovascular diseases (I60-I69) | 379 | 3.4 | 11.5 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 377 | 3.4 | 11.4 |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 246 | 2.2 | 7.5 |
| 9 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 230 | 2.0 | 7.0 |
| 10 | Viral hepatitis (B15-B19) | 216 | 1.9 | 6.5 |
| 11 | Septicemia (A40-A41) | 167 | 1.5 | 5.1 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 166 | 1.5 | 5.0 |
| 13 | Influenza and pneumonia (J09-J18) | 120 | 1.1 | 3.6 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 85 | 0.8 | 2.6 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 82 | 0.7 | 2.5 |
| ... | All other causes (residual) | 1,677 | 14.9 | 50.8 |
| Hispanic, male, 55-64 years | | | | |
| ... | All causes | 16,366 | 100.0 | 788.2 |
| 1 | Malignant neoplasms (C00-C97) | 4,336 | 26.5 | 208.8 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,663 | 22.4 | 176.4 |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,272 | 7.8 | 61.3 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,082 | 6.6 | 52.1 |
| 5 | Diabetes mellitus (E10-E14) | 994 | 6.1 | 47.9 |
| 6 | Cerebrovascular diseases (I60-I69) | 680 | 4.2 | 32.7 |
| 7 | Viral hepatitis (B15-B19) | 352 | 2.2 | 17.0 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 313 | 1.9 | 15.1 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 250 | 1.5 | 12.0 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 238 | 1.5 | 11.5 |
| 11 | Septicemia (A40-A41) | 227 | 1.4 | 10.9 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 200 | 1.2 | 9.6 |
| 13 | Influenza and pneumonia (J09-J18) | 196 | 1.2 | 9.4 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 167 | 1.0 | 8.0 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 111 | 0.7 | 5.3 |
| ... | All other causes (residual) | 2,285 | 14.0 | 110.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------|---|--------|-------------------------|---------|
| Hispanic, male, 65-74 years | | | | |
| ... | All causes | 17,253 | 100.0 | 1,674.3 |
| 1 | Malignant neoplasms (C00-C97) | 5,028 | 29.1 | 487.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,076 | 23.6 | 395.6 |
| 3 | Diabetes mellitus (E10-E14) | 1,120 | 6.5 | 108.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 890 | 5.2 | 86.4 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 705 | 4.1 | 68.4 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 554 | 3.2 | 53.8 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 517 | 3.0 | 50.2 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 447 | 2.6 | 43.4 |
| 9 | Septicemia (A40-A41) | 325 | 1.9 | 31.5 |
| 10 | Influenza and pneumonia (J09-J18) | 306 | 1.8 | 29.7 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 244 | 1.4 | 23.7 |
| 12 | Alzheimer's disease (G30) | 179 | 1.0 | 17.4 |
| 13 | Parkinson's disease (G20-G21) | 145 | 0.8 | 14.1 |
| 14 | Viral hepatitis (B15-B19) | 143 | 0.8 | 13.9 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 126 | 0.7 | 12.2 |
| ... | All other causes (residual) | 2,448 | 14.2 | 237.6 |
| Hispanic, male, 75-84 years | | | | |
| ... | All causes | 18,483 | 100.0 | 4,127.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,687 | 25.4 | 1,046.6 |
| 2 | Malignant neoplasms (C00-C97) | 4,552 | 24.6 | 1,016.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,191 | 6.4 | 265.9 |
| 4 | Diabetes mellitus (E10-E14) | 963 | 5.2 | 215.0 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 830 | 4.5 | 185.3 |
| 6 | Alzheimer's disease (G30) | 688 | 3.7 | 153.6 |
| 7 | Influenza and pneumonia (J09-J18) | 483 | 2.6 | 107.9 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 464 | 2.5 | 103.6 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 400 | 2.2 | 89.3 |
| 10 | Parkinson's disease (G20-G21) | 370 | 2.0 | 82.6 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 321 | 1.7 | 71.7 |
| 12 | Septicemia (A40-A41) | 289 | 1.6 | 64.5 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 283 | 1.5 | 63.2 |
| 14 | Pneumonitis due to solids and liquids (J69) | 147 | 0.8 | 32.8 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 121 | 0.7 | 27.0 |
| ... | All other causes (residual) | 2,694 | 14.6 | 601.6 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-----------------------------------|---|--------|-------------------------|----------|
| Hispanic, male, 85 years and over | | | | |
| ... | All causes | 15,428 | 100.0 | 10,145.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,385 | 28.4 | 2,883.7 |
| 2 | Malignant neoplasms (C00-C97) | 2,393 | 15.5 | 1,573.7 |
| 3 | Alzheimer's disease (G30) | 1,132 | 7.3 | 744.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,075 | 7.0 | 706.9 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 804 | 5.2 | 528.7 |
| 6 | Diabetes mellitus (E10-E14) | 566 | 3.7 | 372.2 |
| 7 | Influenza and pneumonia (J09-J18) | 471 | 3.1 | 309.7 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 395 | 2.6 | 259.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 352 | 2.3 | 231.5 |
| 10 | Parkinson's disease (G20-G21) | 306 | 2.0 | 201.2 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 268 | 1.7 | 176.2 |
| 12 | Septicemia (A40-A41) | 210 | 1.4 | 138.1 |
| 13 | Pneumonitis due to solids and liquids (J69) | 199 | 1.3 | 130.9 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 98 | 0.6 | 64.4 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 79 | 0.5 | 52.0 |
| ... | All other causes (residual) | 2,695 | 17.5 | 1,772.3 |
| Hispanic, female, all ages | | | | |
| ... | All causes | 81,287 | 100.0 | 290.4 |
| 1 | Malignant neoplasms (C00-C97) | 17,957 | 22.1 | 64.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 16,176 | 19.9 | 57.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 5,251 | 6.5 | 18.8 |
| 4 | Alzheimer's disease (G30) | 4,409 | 5.4 | 15.8 |
| 5 | Diabetes mellitus (E10-E14) | 3,852 | 4.7 | 13.8 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,739 | 4.6 | 13.4 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 2,553 | 3.1 | 9.1 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,909 | 2.3 | 6.8 |
| 9 | Influenza and pneumonia (J09-J18) | 1,762 | 2.2 | 6.3 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,744 | 2.1 | 6.2 |
| 11 | Septicemia (A40-A41) | 1,322 | 1.6 | 4.7 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,245 | 1.5 | 4.4 |
| 13 | Certain conditions originating in the perinatal period (P00-P96) | 1,086 | 1.3 | 3.9 |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 837 | 1.0 | 3.0 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 716 | 0.9 | 2.6 |
| ... | All other causes (residual) | 16,729 | 20.6 | 59.8 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|------------------------------|---|--------|-------------------------|------|
| Hispanic, female, 1-4 years | | | | |
| ... | All causes | 354 | 100.0 | 17.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 104 | 29.4 | 5.1 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 57 | 16.1 | 2.8 |
| 3 | Malignant neoplasms (C00-C97) | 35 | 9.9 | 1.7 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 24 | 6.8 | 1.2 |
| 5 | Septicemia (A40-A41) | 9 | 2.5 | * |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 8 | 2.3 | * |
| 7 | Influenza and pneumonia (J09-J18) | 6 | 1.7 | * |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 5 | 1.4 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 4 | 1.1 | * |
| 10 | Anemias (D50-D64) | 3 | 0.8 | * |
| 11 | Meningitis (G00,G03) | 2 | 0.6 | * |
| 11 | Acute bronchitis and bronchiolitis (J20-J21) | 2 | 0.6 | * |
| 11 | Certain conditions originating in the perinatal period (P00-P96) | 2 | 0.6 | * |
| 11 | Complications of medical and surgical care (Y40-Y84,Y88) | 2 | 0.6 | * |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1 | 0.3 | * |
| 15 | Chronic lower respiratory diseases (J40-J47) | 1 | 0.3 | * |
| 15 | Pneumonitis due to solids and liquids (J69) | 1 | 0.3 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.3 | * |
| 15 | Enterocolitis due to <i>Clostridium difficile</i> (A04.7) | 1 | 0.3 | * |
| ... | All other causes (residual) | 86 | 24.3 | 4.3 |
| Hispanic, female, 5-14 years | | | | |
| ... | All causes | 430 | 100.0 | 8.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 102 | 23.7 | 2.0 |
| 2 | Malignant neoplasms (C00-C97) | 90 | 20.9 | 1.8 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 26 | 6.0 | 0.5 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 25 | 5.8 | 0.5 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 21 | 4.9 | 0.4 |
| 6 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 11 | 2.6 | * |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 11 | 2.6 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 10 | 2.3 | * |
| 9 | Influenza and pneumonia (J09-J18) | 9 | 2.1 | * |
| 10 | Cerebrovascular diseases (I60-I69) | 8 | 1.9 | * |
| 11 | Septicemia (A40-A41) | 4 | 0.9 | * |
| 12 | Anemias (D50-D64) | 3 | 0.7 | * |
| 12 | Diabetes mellitus (E10-E14) | 3 | 0.7 | * |
| 12 | Pneumonitis due to solids and liquids (J69) | 3 | 0.7 | * |
| 15 | Meningitis (G00,G03) | 2 | 0.5 | * |
| 15 | Acute bronchitis and bronchiolitis (J20-J21) | 2 | 0.5 | * |
| 15 | Certain conditions originating in the perinatal period (P00-P96) | 2 | 0.5 | * |
| ... | All other causes (residual) | 98 | 22.8 | 2.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|------|
| Hispanic, female, 15-24 years | | | | |
| ... | All causes | 1,314 | 100.0 | 28.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 513 | 39.0 | 11.1 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 178 | 13.5 | 3.9 |
| 3 | Malignant neoplasms (C00-C97) | 124 | 9.4 | 2.7 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 112 | 8.5 | 2.4 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 48 | 3.7 | 1.0 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 33 | 2.5 | 0.7 |
| 7 | Pregnancy, childbirth and the puerperium (O00-O99) | 27 | 2.1 | 0.6 |
| 8 | Influenza and pneumonia (J09-J18) | 15 | 1.1 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 13 | 1.0 | * |
| 10 | Septicemia (A40-A41) | 11 | 0.8 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 8 | 0.6 | * |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 0.5 | * |
| 13 | Diabetes mellitus (E10-E14) | 5 | 0.4 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 5 | 0.4 | * |
| 15 | Meningitis (G00,G03) | 4 | 0.3 | * |
| ... | All other causes (residual) | 211 | 16.1 | 4.6 |
| Hispanic, female, 25-34 years | | | | |
| ... | All causes | 1,914 | 100.0 | 44.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 524 | 27.4 | 12.3 |
| 2 | Malignant neoplasms (C00-C97) | 353 | 18.4 | 8.3 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 152 | 7.9 | 3.6 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 121 | 6.3 | 2.8 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 117 | 6.1 | 2.7 |
| 6 | Pregnancy, childbirth and the puerperium (O00-O99) | 59 | 3.1 | 1.4 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 43 | 2.2 | 1.0 |
| 8 | Diabetes mellitus (E10-E14) | 37 | 1.9 | 0.9 |
| 8 | Cerebrovascular diseases (I60-I69) | 37 | 1.9 | 0.9 |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 29 | 1.5 | 0.7 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 20 | 1.0 | 0.5 |
| 12 | Septicemia (A40-A41) | 19 | 1.0 | * |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 18 | 0.9 | * |
| 13 | Influenza and pneumonia (J09-J18) | 18 | 0.9 | * |
| 15 | Chronic lower respiratory diseases (J40-J47) | 17 | 0.9 | * |
| ... | All other causes (residual) | 350 | 18.3 | 8.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|-------|
| Hispanic, female, 35-44 years | | | | |
| ... | All causes | 3,105 | 100.0 | 77.7 |
| 1 | Malignant neoplasms (C00-C97) | 1,006 | 32.4 | 25.2 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 461 | 14.8 | 11.5 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 272 | 8.8 | 6.8 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 160 | 5.2 | 4.0 |
| 5 | Cerebrovascular diseases (I60-I69) | 118 | 3.8 | 3.0 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 114 | 3.7 | 2.9 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 93 | 3.0 | 2.3 |
| 8 | Diabetes mellitus (E10-E14) | 85 | 2.7 | 2.1 |
| 9 | Pregnancy, childbirth and the puerperium (O00-O99) | 48 | 1.5 | 1.2 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 42 | 1.4 | 1.1 |
| 11 | Septicemia (A40-A41) | 39 | 1.3 | 1.0 |
| 11 | Influenza and pneumonia (J09-J18) | 39 | 1.3 | 1.0 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 27 | 0.9 | 0.7 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 27 | 0.9 | 0.7 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 18 | 0.6 | * |
| ... | All other causes (residual) | 556 | 17.9 | 13.9 |
| Hispanic, female, 45-54 years | | | | |
| ... | All causes | 5,959 | 100.0 | 184.0 |
| 1 | Malignant neoplasms (C00-C97) | 2,198 | 36.9 | 67.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 705 | 11.8 | 21.8 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 488 | 8.2 | 15.1 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 329 | 5.5 | 10.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 286 | 4.8 | 8.8 |
| 6 | Diabetes mellitus (E10-E14) | 264 | 4.4 | 8.2 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 127 | 2.1 | 3.9 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 105 | 1.8 | 3.2 |
| 9 | Septicemia (A40-A41) | 102 | 1.7 | 3.1 |
| 10 | Influenza and pneumonia (J09-J18) | 89 | 1.5 | 2.7 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 82 | 1.4 | 2.5 |
| 12 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 68 | 1.1 | 2.1 |
| 13 | Viral hepatitis (B15-B19) | 64 | 1.1 | 2.0 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 50 | 0.8 | 1.5 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 49 | 0.8 | 1.5 |
| ... | All other causes (residual) | 953 | 16.0 | 29.4 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------|---|--------|-------------------------|---------|
| Hispanic, female, 55-64 years | | | | |
| ... | All causes | 9,621 | 100.0 | 434.6 |
| 1 | Malignant neoplasms (C00-C97) | 3,572 | 37.1 | 161.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,433 | 14.9 | 64.7 |
| 3 | Diabetes mellitus (E10-E14) | 622 | 6.5 | 28.1 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 495 | 5.1 | 22.4 |
| 5 | Cerebrovascular diseases (I60-I69) | 427 | 4.4 | 19.3 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 355 | 3.7 | 16.0 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 242 | 2.5 | 10.9 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 198 | 2.1 | 8.9 |
| 9 | Septicemia (A40-A41) | 194 | 2.0 | 8.8 |
| 10 | Influenza and pneumonia (J09-J18) | 156 | 1.6 | 7.0 |
| 11 | Viral hepatitis (B15-B19) | 141 | 1.5 | 6.4 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 103 | 1.1 | 4.7 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 80 | 0.8 | 3.6 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 48 | 0.5 | 2.2 |
| 15 | Alzheimer's disease (G30) | 43 | 0.4 | 1.9 |
| ... | All other causes (residual) | 1,512 | 15.7 | 68.3 |
| Hispanic, female, 65-74 years | | | | |
| ... | All causes | 13,022 | 100.0 | 1,048.7 |
| 1 | Malignant neoplasms (C00-C97) | 4,105 | 31.5 | 330.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,441 | 18.7 | 196.6 |
| 3 | Diabetes mellitus (E10-E14) | 936 | 7.2 | 75.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 671 | 5.2 | 54.0 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 448 | 3.4 | 36.1 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 403 | 3.1 | 32.5 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 386 | 3.0 | 31.1 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 283 | 2.2 | 22.8 |
| 9 | Influenza and pneumonia (J09-J18) | 250 | 1.9 | 20.1 |
| 10 | Septicemia (A40-A41) | 244 | 1.9 | 19.6 |
| 11 | Alzheimer's disease (G30) | 222 | 1.7 | 17.9 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 186 | 1.4 | 15.0 |
| 13 | Parkinson's disease (G20-G21) | 97 | 0.7 | 7.8 |
| 14 | Viral hepatitis (B15-B19) | 82 | 0.6 | 6.6 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 74 | 0.6 | 6.0 |
| ... | All other causes (residual) | 2,194 | 16.8 | 176.7 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|-------------------------------------|---|--------|-------------------------|---------|
| Hispanic, female, 75-84 years | | | | |
| ... | All causes | 18,606 | 100.0 | 2,937.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,148 | 22.3 | 654.8 |
| 2 | Malignant neoplasms (C00-C97) | 3,938 | 21.2 | 621.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,457 | 7.8 | 230.0 |
| 4 | Alzheimer's disease (G30) | 1,167 | 6.3 | 184.2 |
| 5 | Diabetes mellitus (E10-E14) | 1,090 | 5.9 | 172.1 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 796 | 4.3 | 125.7 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 513 | 2.8 | 81.0 |
| 8 | Influenza and pneumonia (J09-J18) | 456 | 2.5 | 72.0 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 356 | 1.9 | 56.2 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 333 | 1.8 | 52.6 |
| 11 | Septicemia (A40-A41) | 329 | 1.8 | 51.9 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 322 | 1.7 | 50.8 |
| 13 | Parkinson's disease (G20-G21) | 263 | 1.4 | 41.5 |
| 14 | Pneumonitis due to solids and liquids (J69) | 133 | 0.7 | 21.0 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 114 | 0.6 | 18.0 |
| ... | All other causes (residual) | 3,191 | 17.2 | 503.8 |
| Hispanic, female, 85 years and over | | | | |
| ... | All causes | 24,775 | 100.0 | 9,266.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,949 | 28.0 | 2,599.1 |
| 2 | Alzheimer's disease (G30) | 2,969 | 12.0 | 1,110.5 |
| 3 | Malignant neoplasms (C00-C97) | 2,531 | 10.2 | 946.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 2,222 | 9.0 | 831.1 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 966 | 3.9 | 361.3 |
| 6 | Diabetes mellitus (E10-E14) | 809 | 3.3 | 302.6 |
| 7 | Influenza and pneumonia (J09-J18) | 714 | 2.9 | 267.1 |
| 8 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 562 | 2.3 | 210.2 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 470 | 1.9 | 175.8 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 466 | 1.9 | 174.3 |
| 11 | Septicemia (A40-A41) | 350 | 1.4 | 130.9 |
| 12 | Parkinson's disease (G20-G21) | 303 | 1.2 | 113.3 |
| 13 | Pneumonitis due to solids and liquids (J69) | 211 | 0.9 | 78.9 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 116 | 0.5 | 43.4 |
| 15 | Enterocolitis due to <i>Clostridium difficile</i> (A04.7) | 106 | 0.4 | 39.6 |
| ... | All other causes (residual) | 5,031 | 20.3 | 1,881.7 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|-----------|-------------------------|---------|
| Non-Hispanic, white, both sexes, all ages | | | | |
| ... | All causes | 2,123,631 | 100.0 | 1,055.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 503,172 | 23.7 | 250.0 |
| 2 | Malignant neoplasms (C00-C97) | 467,208 | 22.0 | 232.2 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 136,228 | 6.4 | 67.7 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 111,827 | 5.3 | 55.6 |
| 5 | Cerebrovascular diseases (I60-I69) | 106,830 | 5.0 | 53.1 |
| 6 | Alzheimer's disease (G30) | 93,325 | 4.4 | 46.4 |
| 7 | Diabetes mellitus (E10-E14) | 53,590 | 2.5 | 26.6 |
| 8 | Influenza and pneumonia (J09-J18) | 45,242 | 2.1 | 22.5 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 36,465 | 1.7 | 18.1 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 35,481 | 1.7 | 17.6 |
| 11 | Septicemia (A40-A41) | 30,240 | 1.4 | 15.0 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 29,343 | 1.4 | 14.6 |
| 13 | Parkinson's disease (G20-G21) | 24,339 | 1.1 | 12.1 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 22,628 | 1.1 | 11.2 |
| 15 | Pneumonitis due to solids and liquids (J69) | 16,430 | 0.8 | 8.2 |
| ... | All other causes (residual) | 411,283 | 19.4 | 204.4 |
| Non-Hispanic, white, both sexes, 1-4 years | | | | |
| ... | All causes | 1,895 | 100.0 | 22.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 631 | 33.3 | 7.6 |
| 2 | Malignant neoplasms (C00-C97) | 206 | 10.9 | 2.5 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 196 | 10.3 | 2.4 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 145 | 7.7 | 1.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 55 | 2.9 | 0.7 |
| 6 | Influenza and pneumonia (J09-J18) | 37 | 2.0 | 0.4 |
| 7 | Septicemia (A40-A41) | 20 | 1.1 | 0.2 |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 19 | 1.0 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 17 | 0.9 | * |
| 10 | Certain conditions originating in the perinatal period (P00-P96) | 14 | 0.7 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 11 | 0.6 | * |
| 12 | Acute bronchitis and bronchiolitis (J20-J21) | 10 | 0.5 | * |
| 13 | Meningitis (G00,G03) | 8 | 0.4 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 8 | 0.4 | * |
| 15 | Anemias (D50-D64) | 7 | 0.4 | * |
| ... | All other causes (residual) | 511 | 27.0 | 6.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Non-Hispanic, white, both sexes, 5-14 years | | | | |
| ... | All causes | 2,835 | 100.0 | 12.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 798 | 28.1 | 3.6 |
| 2 | Malignant neoplasms (C00-C97) | 476 | 16.8 | 2.2 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 291 | 10.3 | 1.3 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 186 | 6.6 | 0.8 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 109 | 3.8 | 0.5 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 101 | 3.6 | 0.5 |
| 7 | Influenza and pneumonia (J09-J18) | 47 | 1.7 | 0.2 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 45 | 1.6 | 0.2 |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 41 | 1.4 | 0.2 |
| 9 | Cerebrovascular diseases (I60-I69) | 41 | 1.4 | 0.2 |
| 11 | Septicemia (A40-A41) | 28 | 1.0 | 0.1 |
| 12 | Diabetes mellitus (E10-E14) | 11 | 0.4 | * |
| 13 | Anemias (D50-D64) | 10 | 0.4 | * |
| 13 | Certain conditions originating in the perinatal period (P00-P96) | 10 | 0.4 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 8 | 0.3 | * |
| ... | All other causes (residual) | 633 | 22.3 | 2.9 |
| Non-Hispanic, white, both sexes, 15-24 years | | | | |
| ... | All causes | 16,536 | 100.0 | 67.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8,079 | 48.9 | 33.0 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 3,677 | 22.2 | 15.0 |
| 3 | Malignant neoplasms (C00-C97) | 781 | 4.7 | 3.2 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 683 | 4.1 | 2.8 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 495 | 3.0 | 2.0 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 224 | 1.4 | 0.9 |
| 7 | Influenza and pneumonia (J09-J18) | 107 | 0.6 | 0.4 |
| 8 | Diabetes mellitus (E10-E14) | 104 | 0.6 | 0.4 |
| 9 | Cerebrovascular diseases (I60-I69) | 83 | 0.5 | 0.3 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 72 | 0.4 | 0.3 |
| 11 | Pregnancy, childbirth and the puerperium (O00-O99) | 67 | 0.4 | 0.3 |
| 12 | Septicemia (A40-A41) | 61 | 0.4 | 0.2 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 49 | 0.3 | 0.2 |
| 14 | Legal intervention (Y35,Y89.0) | 32 | 0.2 | 0.1 |
| 15 | Pneumonitis due to solids and liquids (J69) | 29 | 0.2 | 0.1 |
| ... | All other causes (residual) | 1,993 | 12.1 | 8.1 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, white, both sexes, 25-34 years | | | | |
| ... | All causes | 31,492 | 100.0 | 124.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 14,143 | 44.9 | 55.7 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 5,177 | 16.4 | 20.4 |
| 3 | Malignant neoplasms (C00-C97) | 2,131 | 6.8 | 8.4 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,928 | 6.1 | 7.6 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,035 | 3.3 | 4.1 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 523 | 1.7 | 2.1 |
| 7 | Diabetes mellitus (E10-E14) | 408 | 1.3 | 1.6 |
| 8 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 278 | 0.9 | 1.1 |
| 9 | Cerebrovascular diseases (I60-I69) | 256 | 0.8 | 1.0 |
| 10 | Influenza and pneumonia (J09-J18) | 226 | 0.7 | 0.9 |
| 11 | Pregnancy, childbirth and the puerperium (O00-O99) | 202 | 0.6 | 0.8 |
| 12 | Septicemia (A40-A41) | 198 | 0.6 | 0.8 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 127 | 0.4 | 0.5 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 99 | 0.3 | 0.4 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 92 | 0.3 | 0.4 |
| ... | All other causes (residual) | 4,669 | 14.8 | 18.4 |
| Non-Hispanic, white, both sexes, 35-44 years | | | | |
| ... | All causes | 45,637 | 100.0 | 192.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 12,891 | 28.2 | 54.3 |
| 2 | Malignant neoplasms (C00-C97) | 6,536 | 14.3 | 27.5 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,983 | 13.1 | 25.2 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 5,623 | 12.3 | 23.7 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,832 | 4.0 | 7.7 |
| 6 | Diabetes mellitus (E10-E14) | 1,068 | 2.3 | 4.5 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 864 | 1.9 | 3.6 |
| 8 | Cerebrovascular diseases (I60-I69) | 796 | 1.7 | 3.4 |
| 9 | Septicemia (A40-A41) | 463 | 1.0 | 1.9 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 444 | 1.0 | 1.9 |
| 11 | Influenza and pneumonia (J09-J18) | 424 | 0.9 | 1.8 |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 340 | 0.7 | 1.4 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 282 | 0.6 | 1.2 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 229 | 0.5 | 1.0 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 182 | 0.4 | 0.8 |
| ... | All other causes (residual) | 7,680 | 16.8 | 32.3 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|-------|
| Non-Hispanic, white, both sexes, 45-54 years | | | | |
| ... | All causes | 118,578 | 100.0 | 418.4 |
| 1 | Malignant neoplasms (C00-C97) | 29,566 | 24.9 | 104.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 22,647 | 19.1 | 79.9 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 15,877 | 13.4 | 56.0 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 7,546 | 6.4 | 26.6 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 6,214 | 5.2 | 21.9 |
| 6 | Diabetes mellitus (E10-E14) | 3,635 | 3.1 | 12.8 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 3,463 | 2.9 | 12.2 |
| 8 | Cerebrovascular diseases (I60-I69) | 2,875 | 2.4 | 10.1 |
| 9 | Septicemia (A40-A41) | 1,550 | 1.3 | 5.5 |
| 10 | Influenza and pneumonia (J09-J18) | 1,329 | 1.1 | 4.7 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,080 | 0.9 | 3.8 |
| 12 | Viral hepatitis (B15-B19) | 982 | 0.8 | 3.5 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 924 | 0.8 | 3.3 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 781 | 0.7 | 2.8 |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 703 | 0.6 | 2.5 |
| ... | All other causes (residual) | 19,406 | 16.4 | 68.5 |
| Non-Hispanic, white, both sexes, 55-64 years | | | | |
| ... | All causes | 257,512 | 100.0 | 872.0 |
| 1 | Malignant neoplasms (C00-C97) | 85,694 | 33.3 | 290.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 53,906 | 20.9 | 182.5 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 14,657 | 5.7 | 49.6 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 14,575 | 5.7 | 49.4 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 9,728 | 3.8 | 32.9 |
| 6 | Diabetes mellitus (E10-E14) | 8,827 | 3.4 | 29.9 |
| 7 | Cerebrovascular diseases (I60-I69) | 7,348 | 2.9 | 24.9 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 6,929 | 2.7 | 23.5 |
| 9 | Septicemia (A40-A41) | 3,872 | 1.5 | 13.1 |
| 10 | Influenza and pneumonia (J09-J18) | 3,202 | 1.2 | 10.8 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,045 | 1.2 | 10.3 |
| 12 | Viral hepatitis (B15-B19) | 2,357 | 0.9 | 8.0 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,075 | 0.8 | 7.0 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,237 | 0.5 | 4.2 |
| 15 | Pneumonitis due to solids and liquids (J69) | 1,091 | 0.4 | 3.7 |
| ... | All other causes (residual) | 38,969 | 15.1 | 132.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| Non-Hispanic, white, both sexes, 65-74 years | | | | |
| ... | All causes | 384,094 | 100.0 | 1,811.0 |
| 1 | Malignant neoplasms (C00-C97) | 129,807 | 33.8 | 612.0 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 81,291 | 21.2 | 383.3 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 33,252 | 8.7 | 156.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 14,566 | 3.8 | 68.7 |
| 5 | Diabetes mellitus (E10-E14) | 13,113 | 3.4 | 61.8 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 10,311 | 2.7 | 48.6 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 6,328 | 1.6 | 29.8 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 6,327 | 1.6 | 29.8 |
| 9 | Septicemia (A40-A41) | 6,311 | 1.6 | 29.8 |
| 10 | Influenza and pneumonia (J09-J18) | 6,112 | 1.6 | 28.8 |
| 11 | Alzheimer's disease (G30) | 5,005 | 1.3 | 23.6 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 3,827 | 1.0 | 18.0 |
| 13 | Parkinson's disease (G20-G21) | 3,203 | 0.8 | 15.1 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 3,181 | 0.8 | 15.0 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2,560 | 0.7 | 12.1 |
| ... | All other causes (residual) | 58,900 | 15.3 | 277.7 |
| Non-Hispanic, white, both sexes, 75-84 years | | | | |
| ... | All causes | 517,686 | 100.0 | 4,717.9 |
| 1 | Malignant neoplasms (C00-C97) | 125,267 | 24.2 | 1,141.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 119,910 | 23.2 | 1,092.8 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 44,473 | 8.6 | 405.3 |
| 4 | Cerebrovascular diseases (I60-I69) | 29,162 | 5.6 | 265.8 |
| 5 | Alzheimer's disease (G30) | 24,522 | 4.7 | 223.5 |
| 6 | Diabetes mellitus (E10-E14) | 13,764 | 2.7 | 125.4 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 13,233 | 2.6 | 120.6 |
| 8 | Influenza and pneumonia (J09-J18) | 11,165 | 2.2 | 101.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 10,169 | 2.0 | 92.7 |
| 10 | Parkinson's disease (G20-G21) | 9,984 | 1.9 | 91.0 |
| 11 | Septicemia (A40-A41) | 8,076 | 1.6 | 73.6 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 5,255 | 1.0 | 47.9 |
| 13 | Pneumonitis due to solids and liquids (J69) | 4,241 | 0.8 | 38.6 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4,087 | 0.8 | 37.2 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,440 | 0.7 | 31.3 |
| ... | All other causes (residual) | 90,938 | 17.6 | 828.8 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|-----------|-------------------------|----------|
| Non-Hispanic, white, both sexes, 85 years and over | | | | |
| ... | All causes | 737,032 | 100.0 | 14,324.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 216,704 | 29.4 | 4,211.7 |
| 2 | Malignant neoplasms (C00-C97) | 86,709 | 11.8 | 1,685.2 |
| 3 | Alzheimer's disease (G30) | 62,966 | 8.5 | 1,223.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 51,648 | 7.0 | 1,003.8 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 39,672 | 5.4 | 771.0 |
| 6 | Influenza and pneumonia (J09-J18) | 22,515 | 3.1 | 437.6 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 20,648 | 2.8 | 401.3 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 14,400 | 2.0 | 279.9 |
| 9 | Diabetes mellitus (E10-E14) | 12,657 | 1.7 | 246.0 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 11,097 | 1.5 | 215.7 |
| 11 | Parkinson's disease (G20-G21) | 10,591 | 1.4 | 205.8 |
| 12 | Septicemia (A40-A41) | 9,589 | 1.3 | 186.4 |
| 13 | Pneumonitis due to solids and liquids (J69) | 8,448 | 1.1 | 164.2 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4,666 | 0.6 | 90.7 |
| 15 | Atherosclerosis (I70) | 2,919 | 0.4 | 56.7 |
| ... | All other causes (residual) | 161,803 | 22.0 | 3,144.7 |
| Non-Hispanic, white, male, all ages | | | | |
| ... | All causes | 1,063,705 | 100.0 | 1,072.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 264,784 | 24.9 | 267.0 |
| 2 | Malignant neoplasms (C00-C97) | 247,645 | 23.3 | 249.7 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 68,715 | 6.5 | 69.3 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 62,857 | 5.9 | 63.4 |
| 5 | Cerebrovascular diseases (I60-I69) | 43,100 | 4.1 | 43.5 |
| 6 | Diabetes mellitus (E10-E14) | 29,813 | 2.8 | 30.1 |
| 7 | Alzheimer's disease (G30) | 28,631 | 2.7 | 28.9 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 28,045 | 2.6 | 28.3 |
| 9 | Influenza and pneumonia (J09-J18) | 20,995 | 2.0 | 21.2 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 18,546 | 1.7 | 18.7 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 18,351 | 1.7 | 18.5 |
| 12 | Parkinson's disease (G20-G21) | 14,778 | 1.4 | 14.9 |
| 13 | Septicemia (A40-A41) | 14,309 | 1.3 | 14.4 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 9,526 | 0.9 | 9.6 |
| 15 | Pneumonitis due to solids and liquids (J69) | 9,085 | 0.9 | 9.2 |
| ... | All other causes (residual) | 184,525 | 17.3 | 186.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---------------------------------------|---|--------|-------------------------|------|
| Non-Hispanic, white, male, 1-4 years | | | | |
| ... | All causes | 1,083 | 100.0 | 25.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 401 | 37.0 | 9.4 |
| 2 | Malignant neoplasms (C00-C97) | 116 | 10.7 | 2.7 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 104 | 9.6 | 2.4 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 88 | 8.1 | 2.1 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 23 | 2.1 | 0.5 |
| 6 | Influenza and pneumonia (J09-J18) | 17 | 1.6 | * |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 12 | 1.1 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 10 | 0.9 | * |
| 9 | Septicemia (A40-A41) | 9 | 0.8 | * |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 9 | 0.8 | * |
| 11 | Acute bronchitis and bronchiolitis (J20-J21) | 7 | 0.6 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 7 | 0.6 | * |
| 13 | Anemias (D50-D64) | 4 | 0.4 | * |
| 14 | Pneumonitis due to solids and liquids (J69) | 3 | 0.3 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 0.3 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 3 | 0.3 | * |
| ... | All other causes (residual) | 267 | 24.7 | 6.3 |
| Non-Hispanic, white, male, 5-14 years | | | | |
| ... | All causes | 1,634 | 100.0 | 14.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 493 | 30.2 | 4.4 |
| 2 | Malignant neoplasms (C00-C97) | 247 | 15.1 | 2.2 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 187 | 11.4 | 1.7 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 106 | 6.5 | 0.9 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 61 | 3.7 | 0.5 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 55 | 3.4 | 0.5 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 27 | 1.7 | 0.2 |
| 8 | Influenza and pneumonia (J09-J18) | 25 | 1.5 | 0.2 |
| 9 | Cerebrovascular diseases (I60-I69) | 21 | 1.3 | 0.2 |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 20 | 1.2 | 0.2 |
| 11 | Septicemia (A40-A41) | 12 | 0.7 | * |
| 12 | Certain conditions originating in the perinatal period (P00-P96) | 6 | 0.4 | * |
| 13 | Anemias (D50-D64) | 5 | 0.3 | * |
| 13 | Diabetes mellitus (E10-E14) | 5 | 0.3 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 5 | 0.3 | * |
| 13 | Complications of medical and surgical care (Y40-Y84,Y88) | 5 | 0.3 | * |
| ... | All other causes (residual) | 354 | 21.7 | 3.1 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, white, male, 15-24 years | | | | |
| ... | All causes | 11,753 | 100.0 | 93.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,805 | 49.4 | 46.2 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,948 | 25.1 | 23.4 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 481 | 4.1 | 3.8 |
| 4 | Malignant neoplasms (C00-C97) | 468 | 4.0 | 3.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 337 | 2.9 | 2.7 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 123 | 1.0 | 1.0 |
| 7 | Influenza and pneumonia (J09-J18) | 62 | 0.5 | 0.5 |
| 8 | Diabetes mellitus (E10-E14) | 58 | 0.5 | 0.5 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 46 | 0.4 | 0.4 |
| 10 | Cerebrovascular diseases (I60-I69) | 43 | 0.4 | 0.3 |
| 11 | Septicemia (A40-A41) | 32 | 0.3 | 0.3 |
| 12 | Legal intervention (Y35,Y89.0) | 31 | 0.3 | 0.2 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 26 | 0.2 | 0.2 |
| 14 | Pneumonitis due to solids and liquids (J69) | 19 | 0.2 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 19 | 0.2 | * |
| ... | All other causes (residual) | 1,255 | 10.7 | 10.0 |
| Non-Hispanic, white, male, 25-34 years | | | | |
| ... | All causes | 21,596 | 100.0 | 168.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 10,306 | 47.7 | 80.2 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 4,107 | 19.0 | 32.0 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,256 | 5.8 | 9.8 |
| 4 | Malignant neoplasms (C00-C97) | 1,077 | 5.0 | 8.4 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 711 | 3.3 | 5.5 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 312 | 1.4 | 2.4 |
| 7 | Diabetes mellitus (E10-E14) | 245 | 1.1 | 1.9 |
| 8 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 168 | 0.8 | 1.3 |
| 9 | Cerebrovascular diseases (I60-I69) | 140 | 0.6 | 1.1 |
| 10 | Influenza and pneumonia (J09-J18) | 118 | 0.5 | 0.9 |
| 11 | Septicemia (A40-A41) | 98 | 0.5 | 0.8 |
| 12 | Legal intervention (Y35,Y89.0) | 72 | 0.3 | 0.6 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 69 | 0.3 | 0.5 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 63 | 0.3 | 0.5 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 60 | 0.3 | 0.5 |
| ... | All other causes (residual) | 2,794 | 12.9 | 21.7 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, white, male, 35-44 years | | | | |
| ... | All causes | 28,437 | 100.0 | 238.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8,873 | 31.2 | 74.3 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 4,188 | 14.7 | 35.1 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,111 | 14.5 | 34.4 |
| 4 | Malignant neoplasms (C00-C97) | 2,925 | 10.3 | 24.5 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,146 | 4.0 | 9.6 |
| 6 | Diabetes mellitus (E10-E14) | 667 | 2.3 | 5.6 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 595 | 2.1 | 5.0 |
| 8 | Cerebrovascular diseases (I60-I69) | 412 | 1.4 | 3.4 |
| 9 | Influenza and pneumonia (J09-J18) | 229 | 0.8 | 1.9 |
| 10 | Septicemia (A40-A41) | 215 | 0.8 | 1.8 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 202 | 0.7 | 1.7 |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 179 | 0.6 | 1.5 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 161 | 0.6 | 1.3 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 153 | 0.5 | 1.3 |
| 15 | Aortic aneurysm and dissection (I71) | 131 | 0.5 | 1.1 |
| ... | All other causes (residual) | 4,250 | 14.9 | 35.6 |
| Non-Hispanic, white, male, 45-54 years | | | | |
| ... | All causes | 71,911 | 100.0 | 510.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 16,052 | 22.3 | 113.8 |
| 2 | Malignant neoplasms (C00-C97) | 14,804 | 20.6 | 105.0 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 10,313 | 14.3 | 73.1 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 5,528 | 7.7 | 39.2 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,936 | 5.5 | 27.9 |
| 6 | Diabetes mellitus (E10-E14) | 2,325 | 3.2 | 16.5 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 1,578 | 2.2 | 11.2 |
| 8 | Cerebrovascular diseases (I60-I69) | 1,538 | 2.1 | 10.9 |
| 9 | Septicemia (A40-A41) | 795 | 1.1 | 5.6 |
| 10 | Influenza and pneumonia (J09-J18) | 735 | 1.0 | 5.2 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 658 | 0.9 | 4.7 |
| 12 | Viral hepatitis (B15-B19) | 655 | 0.9 | 4.6 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 638 | 0.9 | 4.5 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 600 | 0.8 | 4.3 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 523 | 0.7 | 3.7 |
| ... | All other causes (residual) | 11,233 | 15.6 | 79.7 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| Non-Hispanic, white, male, 55-64 years | | | | |
| ... | All causes | 157,623 | 100.0 | 1,093.3 |
| 1 | Malignant neoplasms (C00-C97) | 47,829 | 30.3 | 331.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 37,968 | 24.1 | 263.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 9,810 | 6.2 | 68.0 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 7,586 | 4.8 | 52.6 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 6,607 | 4.2 | 45.8 |
| 6 | Diabetes mellitus (E10-E14) | 5,559 | 3.5 | 38.6 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 5,094 | 3.2 | 35.3 |
| 8 | Cerebrovascular diseases (I60-I69) | 4,170 | 2.6 | 28.9 |
| 9 | Septicemia (A40-A41) | 2,121 | 1.3 | 14.7 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,803 | 1.1 | 12.5 |
| 11 | Influenza and pneumonia (J09-J18) | 1,768 | 1.1 | 12.3 |
| 12 | Viral hepatitis (B15-B19) | 1,665 | 1.1 | 11.5 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,368 | 0.9 | 9.5 |
| 14 | Aortic aneurysm and dissection (I71) | 750 | 0.5 | 5.2 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 724 | 0.5 | 5.0 |
| ... | All other causes (residual) | 22,801 | 14.5 | 158.2 |
| Non-Hispanic, white, male, 65-74 years | | | | |
| ... | All causes | 220,260 | 100.0 | 2,184.0 |
| 1 | Malignant neoplasms (C00-C97) | 71,914 | 32.6 | 713.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 52,665 | 23.9 | 522.2 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 16,633 | 7.6 | 164.9 |
| 4 | Diabetes mellitus (E10-E14) | 7,937 | 3.6 | 78.7 |
| 5 | Cerebrovascular diseases (I60-I69) | 7,720 | 3.5 | 76.5 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 6,663 | 3.0 | 66.1 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 4,050 | 1.8 | 40.2 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3,547 | 1.6 | 35.2 |
| 9 | Influenza and pneumonia (J09-J18) | 3,323 | 1.5 | 33.0 |
| 10 | Septicemia (A40-A41) | 3,222 | 1.5 | 31.9 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 3,071 | 1.4 | 30.5 |
| 12 | Parkinson's disease (G20-G21) | 2,187 | 1.0 | 21.7 |
| 13 | Alzheimer's disease (G30) | 2,089 | 0.9 | 20.7 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,789 | 0.8 | 17.7 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,529 | 0.7 | 15.2 |
| ... | All other causes (residual) | 31,921 | 14.5 | 316.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|----------|
| Non-Hispanic, white, male, 75-84 years | | | | |
| ... | All causes | 265,528 | 100.0 | 5,500.5 |
| 1 | Malignant neoplasms (C00-C97) | 67,155 | 25.3 | 1,391.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 66,590 | 25.1 | 1,379.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 20,890 | 7.9 | 432.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 13,130 | 4.9 | 272.0 |
| 5 | Alzheimer's disease (G30) | 9,296 | 3.5 | 192.6 |
| 6 | Diabetes mellitus (E10-E14) | 7,511 | 2.8 | 155.6 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 7,290 | 2.7 | 151.0 |
| 8 | Parkinson's disease (G20-G21) | 6,447 | 2.4 | 133.6 |
| 9 | Influenza and pneumonia (J09-J18) | 5,702 | 2.1 | 118.1 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 5,459 | 2.1 | 113.1 |
| 11 | Septicemia (A40-A41) | 3,950 | 1.5 | 81.8 |
| 12 | Pneumonitis due to solids and liquids (J69) | 2,480 | 0.9 | 51.4 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2,337 | 0.9 | 48.4 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,337 | 0.9 | 48.4 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,942 | 0.7 | 40.2 |
| ... | All other causes (residual) | 43,012 | 16.2 | 891.0 |
| Non-Hispanic, white, male, 85 years and over | | | | |
| ... | All causes | 278,085 | 100.0 | 15,526.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 85,635 | 30.8 | 4,781.2 |
| 2 | Malignant neoplasms (C00-C97) | 41,090 | 14.8 | 2,294.1 |
| 3 | Alzheimer's disease (G30) | 16,884 | 6.1 | 942.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 15,892 | 5.7 | 887.3 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 15,816 | 5.7 | 883.0 |
| 6 | Influenza and pneumonia (J09-J18) | 8,968 | 3.2 | 500.7 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 8,359 | 3.0 | 466.7 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 6,654 | 2.4 | 371.5 |
| 9 | Parkinson's disease (G20-G21) | 5,759 | 2.1 | 321.5 |
| 10 | Diabetes mellitus (E10-E14) | 5,505 | 2.0 | 307.4 |
| 11 | Pneumonitis due to solids and liquids (J69) | 4,324 | 1.6 | 241.4 |
| 12 | Septicemia (A40-A41) | 3,816 | 1.4 | 213.1 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 3,357 | 1.2 | 187.4 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2,180 | 0.8 | 121.7 |
| 15 | Aortic aneurysm and dissection (I71) | 1,056 | 0.4 | 59.0 |
| ... | All other causes (residual) | 52,790 | 19.0 | 2,947.4 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|-----------|-------------------------|---------|
| Non-Hispanic, white, female, all ages | | | | |
| ... | All causes | 1,059,926 | 100.0 | 1,038.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 238,388 | 22.5 | 233.6 |
| 2 | Malignant neoplasms (C00-C97) | 219,563 | 20.7 | 215.1 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 73,371 | 6.9 | 71.9 |
| 4 | Alzheimer's disease (G30) | 64,694 | 6.1 | 63.4 |
| 5 | Cerebrovascular diseases (I60-I69) | 63,730 | 6.0 | 62.4 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 43,112 | 4.1 | 42.2 |
| 7 | Influenza and pneumonia (J09-J18) | 24,247 | 2.3 | 23.8 |
| 8 | Diabetes mellitus (E10-E14) | 23,777 | 2.2 | 23.3 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 17,130 | 1.6 | 16.8 |
| 10 | Septicemia (A40-A41) | 15,931 | 1.5 | 15.6 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 13,102 | 1.2 | 12.8 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 10,797 | 1.0 | 10.6 |
| 13 | Parkinson's disease (G20-G21) | 9,561 | 0.9 | 9.4 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 8,420 | 0.8 | 8.3 |
| 15 | Pneumonitis due to solids and liquids (J69) | 7,345 | 0.7 | 7.2 |
| ... | All other causes (residual) | 226,758 | 21.4 | 222.2 |
| Non-Hispanic, white, female, 1-4 years | | | | |
| ... | All causes | 812 | 100.0 | 20.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 230 | 28.3 | 5.7 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 92 | 11.3 | 2.3 |
| 3 | Malignant neoplasms (C00-C97) | 90 | 11.1 | 2.2 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 57 | 7.0 | 1.4 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 32 | 3.9 | 0.8 |
| 6 | Influenza and pneumonia (J09-J18) | 20 | 2.5 | 0.5 |
| 7 | Septicemia (A40-A41) | 11 | 1.4 | * |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 0.9 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 7 | 0.9 | * |
| 10 | Meningitis (G00,G03) | 6 | 0.7 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 5 | 0.6 | * |
| 11 | Certain conditions originating in the perinatal period (P00-P96) | 5 | 0.6 | * |
| 13 | Chronic lower respiratory diseases (J40-J47) | 4 | 0.5 | * |
| 14 | Anemias (D50-D64) | 3 | 0.4 | * |
| 14 | Acute bronchitis and bronchiolitis (J20-J21) | 3 | 0.4 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 3 | 0.4 | * |
| ... | All other causes (residual) | 237 | 29.2 | 5.9 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Non-Hispanic, white, female, 5-14 years | | | | |
| ... | All causes | 1,201 | 100.0 | 11.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 305 | 25.4 | 2.8 |
| 2 | Malignant neoplasms (C00-C97) | 229 | 19.1 | 2.1 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 104 | 8.7 | 1.0 |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 80 | 6.7 | 0.7 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 48 | 4.0 | 0.4 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 46 | 3.8 | 0.4 |
| 7 | Influenza and pneumonia (J09-J18) | 22 | 1.8 | 0.2 |
| 8 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 21 | 1.7 | 0.2 |
| 9 | Cerebrovascular diseases (I60-I69) | 20 | 1.7 | 0.2 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 18 | 1.5 | * |
| 11 | Septicemia (A40-A41) | 16 | 1.3 | * |
| 12 | Diabetes mellitus (E10-E14) | 6 | 0.5 | * |
| 13 | Anemias (D50-D64) | 5 | 0.4 | * |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 4 | 0.3 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 0.2 | * |
| ... | All other causes (residual) | 274 | 22.8 | 2.6 |
| Non-Hispanic, white, female, 15-24 years | | | | |
| ... | All causes | 4,783 | 100.0 | 40.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,274 | 47.5 | 19.1 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 729 | 15.2 | 6.1 |
| 3 | Malignant neoplasms (C00-C97) | 313 | 6.5 | 2.6 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 202 | 4.2 | 1.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 158 | 3.3 | 1.3 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 101 | 2.1 | 0.8 |
| 7 | Pregnancy, childbirth and the puerperium (O00-O99) | 67 | 1.4 | 0.6 |
| 8 | Diabetes mellitus (E10-E14) | 46 | 1.0 | 0.4 |
| 9 | Influenza and pneumonia (J09-J18) | 45 | 0.9 | 0.4 |
| 10 | Cerebrovascular diseases (I60-I69) | 40 | 0.8 | 0.3 |
| 11 | Septicemia (A40-A41) | 29 | 0.6 | 0.2 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 26 | 0.5 | 0.2 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 23 | 0.5 | 0.2 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 13 | 0.3 | * |
| 15 | Pneumonitis due to solids and liquids (J69) | 10 | 0.2 | * |
| ... | All other causes (residual) | 707 | 14.8 | 5.9 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, white, female, 25-34 years | | | | |
| ... | All causes | 9,896 | 100.0 | 78.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,837 | 38.8 | 30.6 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,070 | 10.8 | 8.5 |
| 3 | Malignant neoplasms (C00-C97) | 1,054 | 10.7 | 8.4 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 672 | 6.8 | 5.4 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 324 | 3.3 | 2.6 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 211 | 2.1 | 1.7 |
| 7 | Pregnancy, childbirth and the puerperium (O00-O99) | 202 | 2.0 | 1.6 |
| 8 | Diabetes mellitus (E10-E14) | 163 | 1.6 | 1.3 |
| 9 | Cerebrovascular diseases (I60-I69) | 116 | 1.2 | 0.9 |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 110 | 1.1 | 0.9 |
| 11 | Influenza and pneumonia (J09-J18) | 108 | 1.1 | 0.9 |
| 12 | Septicemia (A40-A41) | 100 | 1.0 | 0.8 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 64 | 0.6 | 0.5 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 41 | 0.4 | 0.3 |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 39 | 0.4 | 0.3 |
| ... | All other causes (residual) | 1,785 | 18.0 | 14.2 |
| Non-Hispanic, white, female, 35-44 years | | | | |
| ... | All causes | 17,200 | 100.0 | 145.7 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,018 | 23.4 | 34.0 |
| 2 | Malignant neoplasms (C00-C97) | 3,611 | 21.0 | 30.6 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,872 | 10.9 | 15.9 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,435 | 8.3 | 12.2 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 686 | 4.0 | 5.8 |
| 6 | Diabetes mellitus (E10-E14) | 401 | 2.3 | 3.4 |
| 7 | Cerebrovascular diseases (I60-I69) | 384 | 2.2 | 3.3 |
| 8 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 269 | 1.6 | 2.3 |
| 9 | Septicemia (A40-A41) | 248 | 1.4 | 2.1 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 242 | 1.4 | 2.0 |
| 11 | Influenza and pneumonia (J09-J18) | 195 | 1.1 | 1.7 |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 161 | 0.9 | 1.4 |
| 13 | Pregnancy, childbirth and the puerperium (O00-O99) | 130 | 0.8 | 1.1 |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 129 | 0.8 | 1.1 |
| 15 | Viral hepatitis (B15-B19) | 80 | 0.5 | 0.7 |
| ... | All other causes (residual) | 3,339 | 19.4 | 28.3 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, white, female, 45-54 years | | | | |
| ... | All causes | 46,667 | 100.0 | 327.7 |
| 1 | Malignant neoplasms (C00-C97) | 14,762 | 31.6 | 103.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,595 | 14.1 | 46.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,564 | 11.9 | 39.1 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,278 | 4.9 | 16.0 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 2,018 | 4.3 | 14.2 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 1,885 | 4.0 | 13.2 |
| 7 | Cerebrovascular diseases (I60-I69) | 1,337 | 2.9 | 9.4 |
| 8 | Diabetes mellitus (E10-E14) | 1,310 | 2.8 | 9.2 |
| 9 | Septicemia (A40-A41) | 755 | 1.6 | 5.3 |
| 10 | Influenza and pneumonia (J09-J18) | 594 | 1.3 | 4.2 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 442 | 0.9 | 3.1 |
| 12 | Viral hepatitis (B15-B19) | 327 | 0.7 | 2.3 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 282 | 0.6 | 2.0 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 266 | 0.6 | 1.9 |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 258 | 0.6 | 1.8 |
| ... | All other causes (residual) | 7,994 | 17.1 | 56.1 |
| Non-Hispanic, white, female, 55-64 years | | | | |
| ... | All causes | 99,889 | 100.0 | 661.0 |
| 1 | Malignant neoplasms (C00-C97) | 37,865 | 37.9 | 250.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,938 | 16.0 | 105.5 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 7,071 | 7.1 | 46.8 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,765 | 4.8 | 31.5 |
| 5 | Diabetes mellitus (E10-E14) | 3,268 | 3.3 | 21.6 |
| 6 | Cerebrovascular diseases (I60-I69) | 3,178 | 3.2 | 21.0 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,121 | 3.1 | 20.7 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,835 | 1.8 | 12.1 |
| 9 | Septicemia (A40-A41) | 1,751 | 1.8 | 11.6 |
| 10 | Influenza and pneumonia (J09-J18) | 1,434 | 1.4 | 9.5 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,242 | 1.2 | 8.2 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 707 | 0.7 | 4.7 |
| 13 | Viral hepatitis (B15-B19) | 692 | 0.7 | 4.6 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 513 | 0.5 | 3.4 |
| 15 | Alzheimer's disease (G30) | 434 | 0.4 | 2.9 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 434 | 0.4 | 2.9 |
| ... | All other causes (residual) | 15,641 | 15.7 | 103.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|---------|
| Non-Hispanic, white, female, 65-74 years | | | | |
| ... | All causes | 163,834 | 100.0 | 1,472.8 |
| 1 | Malignant neoplasms (C00-C97) | 57,893 | 35.3 | 520.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 28,626 | 17.5 | 257.3 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 16,619 | 10.1 | 149.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 6,846 | 4.2 | 61.5 |
| 5 | Diabetes mellitus (E10-E14) | 5,176 | 3.2 | 46.5 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 3,648 | 2.2 | 32.8 |
| 7 | Septicemia (A40-A41) | 3,089 | 1.9 | 27.8 |
| 8 | Alzheimer's disease (G30) | 2,916 | 1.8 | 26.2 |
| 9 | Influenza and pneumonia (J09-J18) | 2,789 | 1.7 | 25.1 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,781 | 1.7 | 25.0 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 2,277 | 1.4 | 20.5 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,392 | 0.8 | 12.5 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,031 | 0.6 | 9.3 |
| 14 | Parkinson's disease (G20-G21) | 1,016 | 0.6 | 9.1 |
| 15 | Pneumonitis due to solids and liquids (J69) | 792 | 0.5 | 7.1 |
| ... | All other causes (residual) | 26,943 | 16.4 | 242.2 |
| Non-Hispanic, white, female, 75-84 years | | | | |
| ... | All causes | 252,158 | 100.0 | 4,103.1 |
| 1 | Malignant neoplasms (C00-C97) | 58,112 | 23.0 | 945.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 53,320 | 21.1 | 867.6 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 23,583 | 9.4 | 383.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 16,032 | 6.4 | 260.9 |
| 5 | Alzheimer's disease (G30) | 15,226 | 6.0 | 247.8 |
| 6 | Diabetes mellitus (E10-E14) | 6,253 | 2.5 | 101.7 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 5,943 | 2.4 | 96.7 |
| 8 | Influenza and pneumonia (J09-J18) | 5,463 | 2.2 | 88.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4,710 | 1.9 | 76.6 |
| 10 | Septicemia (A40-A41) | 4,126 | 1.6 | 67.1 |
| 11 | Parkinson's disease (G20-G21) | 3,537 | 1.4 | 57.6 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,918 | 1.2 | 47.5 |
| 13 | Pneumonitis due to solids and liquids (J69) | 1,761 | 0.7 | 28.7 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1,750 | 0.7 | 28.5 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,552 | 0.6 | 25.3 |
| ... | All other causes (residual) | 47,872 | 19.0 | 779.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|----------|
| Non-Hispanic, white, female, 85 years and over | | | | |
| ... | All causes | 458,947 | 100.0 | 13,682.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 131,069 | 28.6 | 3,907.6 |
| 2 | Alzheimer's disease (G30) | 46,082 | 10.0 | 1,373.8 |
| 3 | Malignant neoplasms (C00-C97) | 45,619 | 9.9 | 1,360.0 |
| 4 | Cerebrovascular diseases (I60-I69) | 35,756 | 7.8 | 1,066.0 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 23,856 | 5.2 | 711.2 |
| 6 | Influenza and pneumonia (J09-J18) | 13,547 | 3.0 | 403.9 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 12,289 | 2.7 | 366.4 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7,746 | 1.7 | 230.9 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 7,740 | 1.7 | 230.8 |
| 10 | Diabetes mellitus (E10-E14) | 7,152 | 1.6 | 213.2 |
| 11 | Septicemia (A40-A41) | 5,773 | 1.3 | 172.1 |
| 12 | Parkinson's disease (G20-G21) | 4,832 | 1.1 | 144.1 |
| 13 | Pneumonitis due to solids and liquids (J69) | 4,124 | 0.9 | 122.9 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2,486 | 0.5 | 74.1 |
| 15 | Atherosclerosis (I70) | 1,965 | 0.4 | 58.6 |
| ... | All other causes (residual) | 108,911 | 23.7 | 3,247.0 |
| Non-Hispanic, black, both sexes, all ages | | | | |
| ... | All causes | 315,254 | 100.0 | 754.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 74,093 | 23.5 | 177.4 |
| 2 | Malignant neoplasms (C00-C97) | 68,523 | 21.7 | 164.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 17,760 | 5.6 | 42.5 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 15,366 | 4.9 | 36.8 |
| 5 | Diabetes mellitus (E10-E14) | 13,693 | 4.3 | 32.8 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 10,327 | 3.3 | 24.7 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 9,083 | 2.9 | 21.7 |
| 8 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 9,038 | 2.9 | 21.6 |
| 9 | Alzheimer's disease (G30) | 8,072 | 2.6 | 19.3 |
| 10 | Septicemia (A40-A41) | 6,576 | 2.1 | 15.7 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 5,740 | 1.8 | 13.7 |
| 12 | Influenza and pneumonia (J09-J18) | 5,554 | 1.8 | 13.3 |
| 13 | Certain conditions originating in the perinatal period (P00-P96) | 3,827 | 1.2 | 9.2 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 3,379 | 1.1 | 8.1 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 3,185 | 1.0 | 7.6 |
| ... | All other causes (residual) | 61,038 | 19.4 | 146.1 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| Non-Hispanic, black, both sexes, 1-4 years | | | | |
| ... | All causes | 1,016 | 100.0 | 41.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 286 | 28.1 | 11.7 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 139 | 13.7 | 5.7 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 93 | 9.2 | 3.8 |
| 4 | Malignant neoplasms (C00-C97) | 53 | 5.2 | 2.2 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 52 | 5.1 | 2.1 |
| 6 | Influenza and pneumonia (J09-J18) | 27 | 2.7 | 1.1 |
| 6 | Certain conditions originating in the perinatal period (P00-P96) | 27 | 2.7 | 1.1 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 23 | 2.3 | 0.9 |
| 9 | Cerebrovascular diseases (I60-I69) | 17 | 1.7 | * |
| 10 | Septicemia (A40-A41) | 14 | 1.4 | * |
| 11 | Anemias (D50-D64) | 13 | 1.3 | * |
| 12 | Complications of medical and surgical care (Y40-Y84,Y88) | 9 | 0.9 | * |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 6 | 0.6 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4 | 0.4 | * |
| 15 | Meningitis (G00,G03) | 3 | 0.3 | * |
| ... | All other causes (residual) | 250 | 24.6 | 10.2 |
| Non-Hispanic, black, both sexes, 5-14 years | | | | |
| ... | All causes | 1,187 | 100.0 | 19.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 327 | 27.5 | 5.3 |
| 2 | Malignant neoplasms (C00-C97) | 126 | 10.6 | 2.0 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 121 | 10.2 | 2.0 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 102 | 8.6 | 1.7 |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 66 | 5.6 | 1.1 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 58 | 4.9 | 0.9 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 47 | 4.0 | 0.8 |
| 8 | Cerebrovascular diseases (I60-I69) | 22 | 1.9 | 0.4 |
| 9 | Influenza and pneumonia (J09-J18) | 19 | 1.6 | * |
| 10 | Septicemia (A40-A41) | 16 | 1.3 | * |
| 10 | Anemias (D50-D64) | 16 | 1.3 | * |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 10 | 0.8 | * |
| 13 | Diabetes mellitus (E10-E14) | 9 | 0.8 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 7 | 0.6 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 6 | 0.5 | * |
| ... | All other causes (residual) | 235 | 19.8 | 3.8 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, black, both sexes, 15-24 years | | | | |
| ... | All causes | 7,096 | 100.0 | 104.5 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 3,013 | 42.5 | 44.4 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,649 | 23.2 | 24.3 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 574 | 8.1 | 8.5 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 302 | 4.3 | 4.4 |
| 5 | Malignant neoplasms (C00-C97) | 252 | 3.6 | 3.7 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 86 | 1.2 | 1.3 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 73 | 1.0 | 1.1 |
| 8 | Anemias (D50-D64) | 70 | 1.0 | 1.0 |
| 9 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 67 | 0.9 | 1.0 |
| 10 | Diabetes mellitus (E10-E14) | 64 | 0.9 | 0.9 |
| 11 | Pregnancy, childbirth and the puerperium (O00-O99) | 63 | 0.9 | 0.9 |
| 12 | Cerebrovascular diseases (I60-I69) | 39 | 0.5 | 0.6 |
| 13 | Influenza and pneumonia (J09-J18) | 35 | 0.5 | 0.5 |
| 14 | Septicemia (A40-A41) | 34 | 0.5 | 0.5 |
| 14 | Legal intervention (Y35,Y89.0) | 34 | 0.5 | 0.5 |
| ... | All other causes (residual) | 741 | 10.4 | 10.9 |
| Non-Hispanic, black, both sexes, 25-34 years | | | | |
| ... | All causes | 10,316 | 100.0 | 169.1 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,882 | 27.9 | 47.2 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,154 | 20.9 | 35.3 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 978 | 9.5 | 16.0 |
| 4 | Malignant neoplasms (C00-C97) | 631 | 6.1 | 10.3 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 589 | 5.7 | 9.7 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 326 | 3.2 | 5.3 |
| 7 | Diabetes mellitus (E10-E14) | 252 | 2.4 | 4.1 |
| 8 | Cerebrovascular diseases (I60-I69) | 151 | 1.5 | 2.5 |
| 9 | Pregnancy, childbirth and the puerperium (O00-O99) | 133 | 1.3 | 2.2 |
| 10 | Anemias (D50-D64) | 119 | 1.2 | 2.0 |
| 11 | Septicemia (A40-A41) | 116 | 1.1 | 1.9 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 106 | 1.0 | 1.7 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 102 | 1.0 | 1.7 |
| 14 | Influenza and pneumonia (J09-J18) | 98 | 0.9 | 1.6 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 88 | 0.9 | 1.4 |
| ... | All other causes (residual) | 1,591 | 15.4 | 26.1 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, black, both sexes, 35-44 years | | | | |
| ... | All causes | 14,302 | 100.0 | 269.9 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,821 | 19.7 | 53.2 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,083 | 14.6 | 39.3 |
| 3 | Malignant neoplasms (C00-C97) | 1,908 | 13.3 | 36.0 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,408 | 9.8 | 26.6 |
| 5 | Human immunodeficiency virus (HIV) disease (B20-B24) | 593 | 4.1 | 11.2 |
| 6 | Diabetes mellitus (E10-E14) | 540 | 3.8 | 10.2 |
| 7 | Cerebrovascular diseases (I60-I69) | 524 | 3.7 | 9.9 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 428 | 3.0 | 8.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 283 | 2.0 | 5.3 |
| 10 | Septicemia (A40-A41) | 217 | 1.5 | 4.1 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 212 | 1.5 | 4.0 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 186 | 1.3 | 3.5 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 186 | 1.3 | 3.5 |
| 14 | Influenza and pneumonia (J09-J18) | 142 | 1.0 | 2.7 |
| 15 | Anemias (D50-D64) | 112 | 0.8 | 2.1 |
| ... | All other causes (residual) | 2,659 | 18.6 | 50.2 |
| Non-Hispanic, black, both sexes, 45-54 years | | | | |
| ... | All causes | 31,172 | 100.0 | 577.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,417 | 23.8 | 137.5 |
| 2 | Malignant neoplasms (C00-C97) | 7,037 | 22.6 | 130.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,718 | 8.7 | 50.4 |
| 4 | Diabetes mellitus (E10-E14) | 1,486 | 4.8 | 27.5 |
| 5 | Cerebrovascular diseases (I60-I69) | 1,434 | 4.6 | 26.6 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 954 | 3.1 | 17.7 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 790 | 2.5 | 14.6 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 729 | 2.3 | 13.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 685 | 2.2 | 12.7 |
| 10 | Septicemia (A40-A41) | 634 | 2.0 | 11.8 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 605 | 1.9 | 11.2 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 504 | 1.6 | 9.3 |
| 13 | Influenza and pneumonia (J09-J18) | 421 | 1.4 | 7.8 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 366 | 1.2 | 6.8 |
| 15 | Viral hepatitis (B15-B19) | 158 | 0.5 | 2.9 |
| ... | All other causes (residual) | 5,234 | 16.8 | 97.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Non-Hispanic, black, both sexes, 55-64 years | | | | |
| ... | All causes | 60,886 | 100.0 | 1,298.5 |
| 1 | Malignant neoplasms (C00-C97) | 17,970 | 29.5 | 383.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,068 | 24.7 | 321.4 |
| 3 | Diabetes mellitus (E10-E14) | 3,058 | 5.0 | 65.2 |
| 4 | Cerebrovascular diseases (I60-I69) | 2,998 | 4.9 | 63.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,707 | 4.4 | 57.7 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 1,921 | 3.2 | 41.0 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,667 | 2.7 | 35.6 |
| 8 | Septicemia (A40-A41) | 1,291 | 2.1 | 27.5 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,273 | 2.1 | 27.1 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,066 | 1.8 | 22.7 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 978 | 1.6 | 20.9 |
| 12 | Influenza and pneumonia (J09-J18) | 860 | 1.4 | 18.3 |
| 13 | Viral hepatitis (B15-B19) | 579 | 1.0 | 12.3 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 413 | 0.7 | 8.8 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 238 | 0.4 | 5.1 |
| ... | All other causes (residual) | 8,799 | 14.5 | 187.7 |
| Non-Hispanic, black, both sexes, 65-74 years | | | | |
| ... | All causes | 63,318 | 100.0 | 2,425.7 |
| 1 | Malignant neoplasms (C00-C97) | 18,928 | 29.9 | 725.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,771 | 24.9 | 604.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 3,632 | 5.7 | 139.1 |
| 4 | Diabetes mellitus (E10-E14) | 3,298 | 5.2 | 126.3 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 2,702 | 4.3 | 103.5 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,148 | 3.4 | 82.3 |
| 7 | Septicemia (A40-A41) | 1,494 | 2.4 | 57.2 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,301 | 2.1 | 49.8 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,151 | 1.8 | 44.1 |
| 10 | Influenza and pneumonia (J09-J18) | 1,069 | 1.7 | 41.0 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 675 | 1.1 | 25.9 |
| 12 | Alzheimer's disease (G30) | 635 | 1.0 | 24.3 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 360 | 0.6 | 13.8 |
| 14 | Pneumonitis due to solids and liquids (J69) | 324 | 0.5 | 12.4 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 319 | 0.5 | 12.2 |
| ... | All other causes (residual) | 9,511 | 15.0 | 364.4 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|----------|
| Non-Hispanic, black, both sexes, 75-84 years | | | | |
| ... | All causes | 61,461 | 100.0 | 5,152.8 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,562 | 25.3 | 1,304.7 |
| 2 | Malignant neoplasms (C00-C97) | 14,113 | 23.0 | 1,183.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 4,519 | 7.4 | 378.9 |
| 4 | Diabetes mellitus (E10-E14) | 3,024 | 4.9 | 253.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 2,729 | 4.4 | 228.8 |
| 6 | Alzheimer's disease (G30) | 2,435 | 4.0 | 204.1 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2,221 | 3.6 | 186.2 |
| 8 | Septicemia (A40-A41) | 1,485 | 2.4 | 124.5 |
| 9 | Influenza and pneumonia (J09-J18) | 1,317 | 2.1 | 110.4 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,302 | 2.1 | 109.2 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 950 | 1.5 | 79.6 |
| 12 | Parkinson's disease (G20-G21) | 495 | 0.8 | 41.5 |
| 13 | Pneumonitis due to solids and liquids (J69) | 457 | 0.7 | 38.3 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 372 | 0.6 | 31.2 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 205 | 0.3 | 17.2 |
| ... | All other causes (residual) | 10,275 | 16.7 | 861.4 |
| Non-Hispanic, black, both sexes, 85 years and over | | | | |
| ... | All causes | 57,576 | 100.0 | 12,364.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 15,971 | 27.7 | 3,429.8 |
| 2 | Malignant neoplasms (C00-C97) | 7,496 | 13.0 | 1,609.8 |
| 3 | Alzheimer's disease (G30) | 4,885 | 8.5 | 1,049.1 |
| 4 | Cerebrovascular diseases (I60-I69) | 4,397 | 7.6 | 944.3 |
| 5 | Diabetes mellitus (E10-E14) | 1,957 | 3.4 | 420.3 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,917 | 3.3 | 411.7 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 1,859 | 3.2 | 399.2 |
| 8 | Influenza and pneumonia (J09-J18) | 1,509 | 2.6 | 324.1 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,453 | 2.5 | 312.0 |
| 10 | Septicemia (A40-A41) | 1,216 | 2.1 | 261.1 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 779 | 1.4 | 167.3 |
| 12 | Pneumonitis due to solids and liquids (J69) | 547 | 1.0 | 117.5 |
| 13 | Parkinson's disease (G20-G21) | 401 | 0.7 | 86.1 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 324 | 0.6 | 69.6 |
| 15 | Nutritional deficiencies (E40-E64) | 260 | 0.5 | 55.8 |
| ... | All other causes (residual) | 12,605 | 21.9 | 2,706.9 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--------------------------------------|---|---------|-------------------------|-------|
| Non-Hispanic, black, male, all ages | | | | |
| ... | All causes | 161,850 | 100.0 | 809.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 38,638 | 23.9 | 193.2 |
| 2 | Malignant neoplasms (C00-C97) | 34,787 | 21.5 | 174.0 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 10,516 | 6.5 | 52.6 |
| 4 | Cerebrovascular diseases (I60-I69) | 7,962 | 4.9 | 39.8 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 7,909 | 4.9 | 39.6 |
| 6 | Diabetes mellitus (E10-E14) | 6,806 | 4.2 | 34.0 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 5,236 | 3.2 | 26.2 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4,351 | 2.7 | 21.8 |
| 9 | Septicemia (A40-A41) | 3,061 | 1.9 | 15.3 |
| 10 | Influenza and pneumonia (J09-J18) | 2,729 | 1.7 | 13.6 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 2,599 | 1.6 | 13.0 |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2,230 | 1.4 | 11.2 |
| 13 | Alzheimer's disease (G30) | 2,154 | 1.3 | 10.8 |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 2,130 | 1.3 | 10.7 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,951 | 1.2 | 9.8 |
| ... | All other causes (residual) | 28,791 | 17.8 | 144.0 |
| Non-Hispanic, black, male, 1-4 years | | | | |
| ... | All causes | 608 | 100.0 | 48.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 180 | 29.6 | 14.5 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 90 | 14.8 | 7.2 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 52 | 8.6 | 4.2 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 28 | 4.6 | 2.3 |
| 5 | Malignant neoplasms (C00-C97) | 26 | 4.3 | 2.1 |
| 6 | Certain conditions originating in the perinatal period (P00-P96) | 19 | 3.1 | * |
| 7 | Influenza and pneumonia (J09-J18) | 17 | 2.8 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 15 | 2.5 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 12 | 2.0 | * |
| 10 | Complications of medical and surgical care (Y40-Y84,Y88) | 8 | 1.3 | * |
| 11 | Septicemia (A40-A41) | 7 | 1.2 | * |
| 11 | Anemias (D50-D64) | 7 | 1.2 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 0.5 | * |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1 | 0.2 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1 | 0.2 | * |
| 14 | Diabetes mellitus (E10-E14) | 1 | 0.2 | * |
| 14 | Nutritional deficiencies (E40-E64) | 1 | 0.2 | * |
| 14 | Meningitis (G00,G03) | 1 | 0.2 | * |
| 14 | Pneumonitis due to solids and liquids (J69) | 1 | 0.2 | * |
| 14 | Diseases of appendix (K35-K38) | 1 | 0.2 | * |
| ... | All other causes (residual) | 137 | 22.5 | 11.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, black, male, 5-14 years | | | | |
| ... | All causes | 703 | 100.0 | 22.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 187 | 26.6 | 6.0 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 83 | 11.8 | 2.6 |
| 3 | Malignant neoplasms (C00-C97) | 68 | 9.7 | 2.2 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 63 | 9.0 | 2.0 |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 38 | 5.4 | 1.2 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 35 | 5.0 | 1.1 |
| 7 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 32 | 4.6 | 1.0 |
| 8 | Cerebrovascular diseases (I60-I69) | 13 | 1.8 | * |
| 9 | Influenza and pneumonia (J09-J18) | 9 | 1.3 | * |
| 10 | Septicemia (A40-A41) | 7 | 1.0 | * |
| 10 | Anemias (D50-D64) | 7 | 1.0 | * |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 6 | 0.9 | * |
| 12 | Diabetes mellitus (E10-E14) | 6 | 0.9 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 0.4 | * |
| 14 | Certain conditions originating in the perinatal period (P00-P96) | 3 | 0.4 | * |
| ... | All other causes (residual) | 143 | 20.3 | 4.6 |
| Non-Hispanic, black, male, 15-24 years | | | | |
| ... | All causes | 5,486 | 100.0 | 159.2 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,744 | 50.0 | 79.6 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,226 | 22.3 | 35.6 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 452 | 8.2 | 13.1 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 194 | 3.5 | 5.6 |
| 5 | Malignant neoplasms (C00-C97) | 138 | 2.5 | 4.0 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 55 | 1.0 | 1.6 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 45 | 0.8 | 1.3 |
| 7 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 45 | 0.8 | 1.3 |
| 9 | Diabetes mellitus (E10-E14) | 39 | 0.7 | 1.1 |
| 10 | Anemias (D50-D64) | 35 | 0.6 | 1.0 |
| 11 | Legal intervention (Y35,Y89.0) | 32 | 0.6 | 0.9 |
| 12 | Influenza and pneumonia (J09-J18) | 21 | 0.4 | 0.6 |
| 13 | Cerebrovascular diseases (I60-I69) | 20 | 0.4 | 0.6 |
| 14 | Septicemia (A40-A41) | 15 | 0.3 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 11 | 0.2 | * |
| ... | All other causes (residual) | 414 | 7.5 | 12.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, black, male, 25-34 years | | | | |
| ... | All causes | 7,229 | 100.0 | 243.1 |
| 1 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 2,592 | 35.9 | 87.2 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,602 | 22.2 | 53.9 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 596 | 8.2 | 20.0 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 491 | 6.8 | 16.5 |
| 5 | Malignant neoplasms (C00-C97) | 268 | 3.7 | 9.0 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 220 | 3.0 | 7.4 |
| 7 | Diabetes mellitus (E10-E14) | 131 | 1.8 | 4.4 |
| 8 | Cerebrovascular diseases (I60-I69) | 76 | 1.1 | 2.6 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 68 | 0.9 | 2.3 |
| 10 | Influenza and pneumonia (J09-J18) | 67 | 0.9 | 2.3 |
| 11 | Septicemia (A40-A41) | 61 | 0.8 | 2.1 |
| 12 | Anemias (D50-D64) | 59 | 0.8 | 2.0 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 52 | 0.7 | 1.7 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 52 | 0.7 | 1.7 |
| 15 | Legal intervention (Y35,Y89.0) | 39 | 0.5 | 1.3 |
| ... | All other causes (residual) | 855 | 11.8 | 28.8 |
| Non-Hispanic, black, male, 35-44 years | | | | |
| ... | All causes | 8,610 | 100.0 | 344.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,802 | 20.9 | 72.1 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,498 | 17.4 | 60.0 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,237 | 14.4 | 49.5 |
| 4 | Malignant neoplasms (C00-C97) | 697 | 8.1 | 27.9 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 357 | 4.1 | 14.3 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 333 | 3.9 | 13.3 |
| 7 | Diabetes mellitus (E10-E14) | 317 | 3.7 | 12.7 |
| 8 | Cerebrovascular diseases (I60-I69) | 299 | 3.5 | 12.0 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 150 | 1.7 | 6.0 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 133 | 1.5 | 5.3 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 100 | 1.2 | 4.0 |
| 12 | Septicemia (A40-A41) | 94 | 1.1 | 3.8 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 81 | 0.9 | 3.2 |
| 14 | Influenza and pneumonia (J09-J18) | 80 | 0.9 | 3.2 |
| 15 | Aortic aneurysm and dissection (I71) | 75 | 0.9 | 3.0 |
| ... | All other causes (residual) | 1,357 | 15.8 | 54.3 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Non-Hispanic, black, male, 45-54 years | | | | |
| ... | All causes | 17,812 | 100.0 | 703.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 4,660 | 26.2 | 184.1 |
| 2 | Malignant neoplasms (C00-C97) | 3,266 | 18.3 | 129.0 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,900 | 10.7 | 75.0 |
| 4 | Diabetes mellitus (E10-E14) | 858 | 4.8 | 33.9 |
| 5 | Cerebrovascular diseases (I60-I69) | 795 | 4.5 | 31.4 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 660 | 3.7 | 26.1 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 608 | 3.4 | 24.0 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 406 | 2.3 | 16.0 |
| 9 | Chronic liver disease and cirrhosis (K70,K73-K74) | 403 | 2.3 | 15.9 |
| 10 | Septicemia (A40-A41) | 317 | 1.8 | 12.5 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 293 | 1.6 | 11.6 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 286 | 1.6 | 11.3 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 277 | 1.6 | 10.9 |
| 14 | Influenza and pneumonia (J09-J18) | 256 | 1.4 | 10.1 |
| 15 | Aortic aneurysm and dissection (I71) | 110 | 0.6 | 4.3 |
| ... | All other causes (residual) | 2,717 | 15.3 | 107.3 |
| Non-Hispanic, black, male, 55-64 years | | | | |
| ... | All causes | 35,443 | 100.0 | 1,656.8 |
| 1 | Malignant neoplasms (C00-C97) | 9,646 | 27.2 | 450.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9,451 | 26.7 | 441.8 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,987 | 5.6 | 92.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,730 | 4.9 | 80.9 |
| 5 | Diabetes mellitus (E10-E14) | 1,724 | 4.9 | 80.6 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 956 | 2.7 | 44.7 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 932 | 2.6 | 43.6 |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 842 | 2.4 | 39.4 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 707 | 2.0 | 33.0 |
| 10 | Septicemia (A40-A41) | 659 | 1.9 | 30.8 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 608 | 1.7 | 28.4 |
| 12 | Influenza and pneumonia (J09-J18) | 504 | 1.4 | 23.6 |
| 13 | Viral hepatitis (B15-B19) | 405 | 1.1 | 18.9 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 332 | 0.9 | 15.5 |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 188 | 0.5 | 8.8 |
| ... | All other causes (residual) | 4,772 | 13.5 | 223.1 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Non-Hispanic, black, male, 65-74 years | | | | |
| ... | All causes | 34,867 | 100.0 | 3,109.3 |
| 1 | Malignant neoplasms (C00-C97) | 10,276 | 29.5 | 916.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9,160 | 26.3 | 816.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,957 | 5.6 | 174.5 |
| 4 | Diabetes mellitus (E10-E14) | 1,751 | 5.0 | 156.1 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,508 | 4.3 | 134.5 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,099 | 3.2 | 98.0 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 851 | 2.4 | 75.9 |
| 8 | Septicemia (A40-A41) | 784 | 2.2 | 69.9 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 585 | 1.7 | 52.2 |
| 10 | Influenza and pneumonia (J09-J18) | 553 | 1.6 | 49.3 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 460 | 1.3 | 41.0 |
| 12 | Alzheimer's disease (G30) | 265 | 0.8 | 23.6 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 252 | 0.7 | 22.5 |
| 14 | Viral hepatitis (B15-B19) | 196 | 0.6 | 17.5 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 176 | 0.5 | 15.7 |
| 15 | Pneumonitis due to solids and liquids (J69) | 176 | 0.5 | 15.7 |
| ... | All other causes (residual) | 4,818 | 13.8 | 429.7 |
| Non-Hispanic, black, male, 75-84 years | | | | |
| ... | All causes | 28,940 | 100.0 | 6,345.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,647 | 26.4 | 1,676.7 |
| 2 | Malignant neoplasms (C00-C97) | 7,136 | 24.7 | 1,564.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,915 | 6.6 | 419.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 1,391 | 4.8 | 305.0 |
| 5 | Diabetes mellitus (E10-E14) | 1,350 | 4.7 | 296.0 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 998 | 3.4 | 218.8 |
| 7 | Alzheimer's disease (G30) | 831 | 2.9 | 182.2 |
| 8 | Influenza and pneumonia (J09-J18) | 670 | 2.3 | 146.9 |
| 9 | Septicemia (A40-A41) | 664 | 2.3 | 145.6 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 536 | 1.9 | 117.5 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 533 | 1.8 | 116.9 |
| 12 | Parkinson's disease (G20-G21) | 301 | 1.0 | 66.0 |
| 13 | Pneumonitis due to solids and liquids (J69) | 250 | 0.9 | 54.8 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 175 | 0.6 | 38.4 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 101 | 0.3 | 22.1 |
| ... | All other causes (residual) | 4,442 | 15.3 | 974.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|---------|-------------------------|----------|
| Non-Hispanic, black, male, 85 years and over | | | | |
| ... | All causes | 18,325 | 100.0 | 13,344.0 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,020 | 27.4 | 3,655.5 |
| 2 | Malignant neoplasms (C00-C97) | 3,261 | 17.8 | 2,374.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,131 | 6.2 | 823.6 |
| 4 | Alzheimer's disease (G30) | 1,008 | 5.5 | 734.0 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 789 | 4.3 | 574.5 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 691 | 3.8 | 503.2 |
| 7 | Diabetes mellitus (E10-E14) | 627 | 3.4 | 456.6 |
| 8 | Influenza and pneumonia (J09-J18) | 515 | 2.8 | 375.0 |
| 9 | Septicemia (A40-A41) | 419 | 2.3 | 305.1 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 414 | 2.3 | 301.5 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 329 | 1.8 | 239.6 |
| 12 | Pneumonitis due to solids and liquids (J69) | 234 | 1.3 | 170.4 |
| 13 | Parkinson's disease (G20-G21) | 194 | 1.1 | 141.3 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 112 | 0.6 | 81.6 |
| 15 | Nutritional deficiencies (E40-E64) | 88 | 0.5 | 64.1 |
| ... | All other causes (residual) | 3,493 | 19.1 | 2,543.5 |
| Non-Hispanic, black, female, all ages | | | | |
| ... | All causes | 153,404 | 100.0 | 704.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 35,455 | 23.1 | 162.8 |
| 2 | Malignant neoplasms (C00-C97) | 33,736 | 22.0 | 154.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 9,798 | 6.4 | 45.0 |
| 4 | Diabetes mellitus (E10-E14) | 6,887 | 4.5 | 31.6 |
| 5 | Alzheimer's disease (G30) | 5,918 | 3.9 | 27.2 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 5,091 | 3.3 | 23.4 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 4,850 | 3.2 | 22.3 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4,732 | 3.1 | 21.7 |
| 9 | Septicemia (A40-A41) | 3,515 | 2.3 | 16.1 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 3,141 | 2.0 | 14.4 |
| 11 | Influenza and pneumonia (J09-J18) | 2,825 | 1.8 | 13.0 |
| 12 | Certain conditions originating in the perinatal period (P00-P96) | 1,697 | 1.1 | 7.8 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,252 | 0.8 | 5.7 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1,149 | 0.7 | 5.3 |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 1,129 | 0.7 | 5.2 |
| ... | All other causes (residual) | 32,229 | 21.0 | 148.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| Non-Hispanic, black, female, 1-4 years | | | | |
| ... | All causes | 408 | 100.0 | 34.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 106 | 26.0 | 8.8 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 49 | 12.0 | 4.1 |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 41 | 10.0 | 3.4 |
| 4 | Malignant neoplasms (C00-C97) | 27 | 6.6 | 2.3 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 24 | 5.9 | 2.0 |
| 6 | Influenza and pneumonia (J09-J18) | 10 | 2.5 | * |
| 7 | Chronic lower respiratory diseases (J40-J47) | 8 | 2.0 | * |
| 7 | Certain conditions originating in the perinatal period (P00-P96) | 8 | 2.0 | * |
| 9 | Septicemia (A40-A41) | 7 | 1.7 | * |
| 10 | Anemias (D50-D64) | 6 | 1.5 | * |
| 11 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 5 | 1.2 | * |
| 11 | Cerebrovascular diseases (I60-I69) | 5 | 1.2 | * |
| 13 | Meningitis (G00,G03) | 2 | 0.5 | * |
| 14 | Meningococcal infection (A39) | 1 | 0.2 | * |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1 | 0.2 | * |
| 14 | Nutritional deficiencies (E40-E64) | 1 | 0.2 | * |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1 | 0.2 | * |
| 14 | Acute bronchitis and bronchiolitis (J20-J21) | 1 | 0.2 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.2 | * |
| 14 | Infections of kidney (N10-N12,N13.6,N15.1) | 1 | 0.2 | * |
| 14 | Complications of medical and surgical care (Y40-Y84,Y88) | 1 | 0.2 | * |
| 14 | Enterocolitis due to <i>Clostridium difficile</i> (A04.7) | 1 | 0.2 | * |
| ... | All other causes (residual) | 101 | 24.8 | 8.4 |
| Non-Hispanic, black, female, 5-14 years | | | | |
| ... | All causes | 484 | 100.0 | 15.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 140 | 28.9 | 4.6 |
| 2 | Malignant neoplasms (C00-C97) | 58 | 12.0 | 1.9 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 39 | 8.1 | 1.3 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 38 | 7.9 | 1.2 |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 28 | 5.8 | 0.9 |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 26 | 5.4 | 0.9 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 12 | 2.5 | * |
| 8 | Influenza and pneumonia (J09-J18) | 10 | 2.1 | * |
| 9 | Septicemia (A40-A41) | 9 | 1.9 | * |
| 9 | Anemias (D50-D64) | 9 | 1.9 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 9 | 1.9 | * |
| 12 | Complications of medical and surgical care (Y40-Y84,Y88) | 5 | 1.0 | * |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4 | 0.8 | * |
| 14 | Diabetes mellitus (E10-E14) | 3 | 0.6 | * |
| 14 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 3 | 0.6 | * |
| ... | All other causes (residual) | 91 | 18.8 | 3.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Non-Hispanic, black, female, 15-24 years | | | | |
| ... | All causes | 1,610 | 100.0 | 48.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 423 | 26.3 | 12.7 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 269 | 16.7 | 8.0 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 122 | 7.6 | 3.7 |
| 4 | Malignant neoplasms (C00-C97) | 114 | 7.1 | 3.4 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 108 | 6.7 | 3.2 |
| 6 | Pregnancy, childbirth and the puerperium (O00-O99) | 63 | 3.9 | 1.9 |
| 7 | Anemias (D50-D64) | 35 | 2.2 | 1.0 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 31 | 1.9 | 0.9 |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 28 | 1.7 | 0.8 |
| 10 | Diabetes mellitus (E10-E14) | 25 | 1.6 | 0.7 |
| 11 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 22 | 1.4 | 0.7 |
| 12 | Septicemia (A40-A41) | 19 | 1.2 | * |
| 12 | Cerebrovascular diseases (I60-I69) | 19 | 1.2 | * |
| 14 | Influenza and pneumonia (J09-J18) | 14 | 0.9 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 13 | 0.8 | * |
| ... | All other causes (residual) | 305 | 18.9 | 9.1 |
| Non-Hispanic, black, female, 25-34 years | | | | |
| ... | All causes | 3,087 | 100.0 | 98.7 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 552 | 17.9 | 17.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 382 | 12.4 | 12.2 |
| 3 | Malignant neoplasms (C00-C97) | 363 | 11.8 | 11.6 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 290 | 9.4 | 9.3 |
| 5 | Pregnancy, childbirth and the puerperium (O00-O99) | 133 | 4.3 | 4.3 |
| 6 | Diabetes mellitus (E10-E14) | 121 | 3.9 | 3.9 |
| 7 | Human immunodeficiency virus (HIV) disease (B20-B24) | 106 | 3.4 | 3.4 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 98 | 3.2 | 3.1 |
| 9 | Cerebrovascular diseases (I60-I69) | 75 | 2.4 | 2.4 |
| 10 | Anemias (D50-D64) | 60 | 1.9 | 1.9 |
| 11 | Septicemia (A40-A41) | 55 | 1.8 | 1.8 |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 54 | 1.7 | 1.7 |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 36 | 1.2 | 1.2 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 34 | 1.1 | 1.1 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 33 | 1.1 | 1.1 |
| ... | All other causes (residual) | 695 | 22.5 | 22.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, black, female, 35-44 years | | | | |
| ... | All causes | 5,692 | 100.0 | 203.3 |
| 1 | Malignant neoplasms (C00-C97) | 1,211 | 21.3 | 43.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,019 | 17.9 | 36.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 585 | 10.3 | 20.9 |
| 4 | Human immunodeficiency virus (HIV) disease (B20-B24) | 260 | 4.6 | 9.3 |
| 5 | Cerebrovascular diseases (I60-I69) | 225 | 4.0 | 8.0 |
| 6 | Diabetes mellitus (E10-E14) | 223 | 3.9 | 8.0 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 171 | 3.0 | 6.1 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 133 | 2.3 | 4.7 |
| 9 | Septicemia (A40-A41) | 123 | 2.2 | 4.4 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 105 | 1.8 | 3.7 |
| 10 | Pregnancy, childbirth and the puerperium (O00-O99) | 105 | 1.8 | 3.7 |
| 12 | Chronic lower respiratory diseases (J40-J47) | 86 | 1.5 | 3.1 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 79 | 1.4 | 2.8 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 71 | 1.2 | 2.5 |
| 15 | Influenza and pneumonia (J09-J18) | 62 | 1.1 | 2.2 |
| ... | All other causes (residual) | 1,234 | 21.7 | 44.1 |
| Non-Hispanic, black, female, 45-54 years | | | | |
| ... | All causes | 13,360 | 100.0 | 466.5 |
| 1 | Malignant neoplasms (C00-C97) | 3,771 | 28.2 | 131.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,757 | 20.6 | 96.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 818 | 6.1 | 28.6 |
| 4 | Cerebrovascular diseases (I60-I69) | 639 | 4.8 | 22.3 |
| 5 | Diabetes mellitus (E10-E14) | 628 | 4.7 | 21.9 |
| 6 | Human immunodeficiency virus (HIV) disease (B20-B24) | 346 | 2.6 | 12.1 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 326 | 2.4 | 11.4 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 319 | 2.4 | 11.1 |
| 9 | Septicemia (A40-A41) | 317 | 2.4 | 11.1 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 279 | 2.1 | 9.7 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 211 | 1.6 | 7.4 |
| 12 | Influenza and pneumonia (J09-J18) | 165 | 1.2 | 5.8 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 130 | 1.0 | 4.5 |
| 14 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 89 | 0.7 | 3.1 |
| 15 | Anemias (D50-D64) | 76 | 0.6 | 2.7 |
| ... | All other causes (residual) | 2,489 | 18.6 | 86.9 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Non-Hispanic, black, female, 55-64 years | | | | |
| ... | All causes | 25,443 | 100.0 | 997.9 |
| 1 | Malignant neoplasms (C00-C97) | 8,324 | 32.7 | 326.5 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,617 | 22.1 | 220.3 |
| 3 | Diabetes mellitus (E10-E14) | 1,334 | 5.2 | 52.3 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,268 | 5.0 | 49.7 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 965 | 3.8 | 37.8 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 735 | 2.9 | 28.8 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 720 | 2.8 | 28.2 |
| 8 | Septicemia (A40-A41) | 632 | 2.5 | 24.8 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 458 | 1.8 | 18.0 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 431 | 1.7 | 16.9 |
| 11 | Influenza and pneumonia (J09-J18) | 356 | 1.4 | 14.0 |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 271 | 1.1 | 10.6 |
| 13 | Viral hepatitis (B15-B19) | 174 | 0.7 | 6.8 |
| 14 | Pneumonitis due to solids and liquids (J69) | 100 | 0.4 | 3.9 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 98 | 0.4 | 3.8 |
| ... | All other causes (residual) | 3,960 | 15.6 | 155.3 |
| Non-Hispanic, black, female, 65-74 years | | | | |
| ... | All causes | 28,451 | 100.0 | 1,910.9 |
| 1 | Malignant neoplasms (C00-C97) | 8,652 | 30.4 | 581.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,611 | 23.2 | 444.0 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,675 | 5.9 | 112.5 |
| 4 | Diabetes mellitus (E10-E14) | 1,547 | 5.4 | 103.9 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1,194 | 4.2 | 80.2 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,049 | 3.7 | 70.5 |
| 7 | Septicemia (A40-A41) | 710 | 2.5 | 47.7 |
| 8 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 566 | 2.0 | 38.0 |
| 9 | Influenza and pneumonia (J09-J18) | 516 | 1.8 | 34.7 |
| 10 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 450 | 1.6 | 30.2 |
| 11 | Alzheimer's disease (G30) | 370 | 1.3 | 24.9 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 215 | 0.8 | 14.4 |
| 13 | Pneumonitis due to solids and liquids (J69) | 148 | 0.5 | 9.9 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 143 | 0.5 | 9.6 |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 108 | 0.4 | 7.3 |
| ... | All other causes (residual) | 4,497 | 15.8 | 302.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|----------|
| Non-Hispanic, black, female, 75-84 years | | | | |
| ... | All causes | 32,521 | 100.0 | 4,414.5 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,915 | 24.3 | 1,074.4 |
| 2 | Malignant neoplasms (C00-C97) | 6,977 | 21.5 | 947.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,604 | 8.0 | 353.5 |
| 4 | Diabetes mellitus (E10-E14) | 1,674 | 5.1 | 227.2 |
| 5 | Alzheimer's disease (G30) | 1,604 | 4.9 | 217.7 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 1,338 | 4.1 | 181.6 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,223 | 3.8 | 166.0 |
| 8 | Septicemia (A40-A41) | 821 | 2.5 | 111.4 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 766 | 2.4 | 104.0 |
| 10 | Influenza and pneumonia (J09-J18) | 647 | 2.0 | 87.8 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 417 | 1.3 | 56.6 |
| 12 | Pneumonitis due to solids and liquids (J69) | 207 | 0.6 | 28.1 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 197 | 0.6 | 26.7 |
| 14 | Parkinson's disease (G20-G21) | 194 | 0.6 | 26.3 |
| 15 | Aortic aneurysm and dissection (I71) | 116 | 0.4 | 15.7 |
| ... | All other causes (residual) | 5,821 | 17.9 | 790.2 |
| Non-Hispanic, black, female, 85 years and over | | | | |
| ... | All causes | 39,251 | 100.0 | 11,954.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10,951 | 27.9 | 3,335.4 |
| 2 | Malignant neoplasms (C00-C97) | 4,235 | 10.8 | 1,289.9 |
| 3 | Alzheimer's disease (G30) | 3,877 | 9.9 | 1,180.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 3,266 | 8.3 | 994.7 |
| 5 | Diabetes mellitus (E10-E14) | 1,330 | 3.4 | 405.1 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,226 | 3.1 | 373.4 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 1,070 | 2.7 | 325.9 |
| 8 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,039 | 2.6 | 316.4 |
| 9 | Influenza and pneumonia (J09-J18) | 994 | 2.5 | 302.7 |
| 10 | Septicemia (A40-A41) | 797 | 2.0 | 242.7 |
| 11 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 450 | 1.1 | 137.1 |
| 12 | Pneumonitis due to solids and liquids (J69) | 313 | 0.8 | 95.3 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 212 | 0.5 | 64.6 |
| 14 | Parkinson's disease (G20-G21) | 207 | 0.5 | 63.0 |
| 15 | Nutritional deficiencies (E40-E64) | 172 | 0.4 | 52.4 |
| ... | All other causes (residual) | 9,112 | 23.2 | 2,775.3 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|-------|
| Non-Hispanic, American Indian or Alaska Native, both sexes, all ages | | | | |
| ... | All causes | 18,039 | 100.0 | 670.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,303 | 18.3 | 122.8 |
| 2 | Malignant neoplasms (C00-C97) | 3,192 | 17.7 | 118.7 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,953 | 10.8 | 72.6 |
| 4 | Diabetes mellitus (E10-E14) | 1,034 | 5.7 | 38.4 |
| 5 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1,006 | 5.6 | 37.4 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 859 | 4.8 | 31.9 |
| 7 | Cerebrovascular diseases (I60-I69) | 646 | 3.6 | 24.0 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 535 | 3.0 | 19.9 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 340 | 1.9 | 12.6 |
| 10 | Influenza and pneumonia (J09-J18) | 330 | 1.8 | 12.3 |
| 11 | Alzheimer's disease (G30) | 307 | 1.7 | 11.4 |
| 12 | Septicemia (A40-A41) | 296 | 1.6 | 11.0 |
| 13 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 262 | 1.5 | 9.7 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 192 | 1.1 | 7.1 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 109 | 0.6 | 4.1 |
| ... | All other causes (residual) | 3,675 | 20.4 | 136.6 |
| Non-Hispanic, American Indian or Alaska Native, both sexes, 1-4 years | | | | |
| ... | All causes | 82 | 100.0 | 51.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 30 | 36.6 | 18.7 |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 10 | 12.2 | * |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 8 | 9.8 | * |
| 4 | Malignant neoplasms (C00-C97) | 3 | 3.7 | * |
| 5 | Meningitis (G00,G03) | 1 | 1.2 | * |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1 | 1.2 | * |
| 5 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1 | 1.2 | * |
| 5 | Cerebrovascular diseases (I60-I69) | 1 | 1.2 | * |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1 | 1.2 | * |
| 5 | Pneumonitis due to solids and liquids (J69) | 1 | 1.2 | * |
| ... | All other causes (residual) | 25 | 30.5 | 15.6 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|-------|
| Non-Hispanic, American Indian or Alaska Native, both sexes, 5-14 years | | | | |
| ... | All causes | 99 | 100.0 | 24.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 31 | 31.3 | 7.5 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 14 | 14.1 | * |
| 3 | Malignant neoplasms (C00-C97) | 11 | 11.1 | * |
| 4 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 7 | 7.1 | * |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 7 | 7.1 | * |
| 6 | Chronic lower respiratory diseases (J40-J47) | 3 | 3.0 | * |
| 7 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2 | 2.0 | * |
| 7 | Certain conditions originating in the perinatal period (P00-P96) | 2 | 2.0 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1 | 1.0 | * |
| 9 | Influenza and pneumonia (J09-J18) | 1 | 1.0 | * |
| ... | All other causes (residual) | 20 | 20.2 | 4.9 |
| Non-Hispanic, American Indian or Alaska Native, both sexes, 15-24 years | | | | |
| ... | All causes | 525 | 100.0 | 119.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 205 | 39.0 | 46.6 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 150 | 28.6 | 34.1 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 52 | 9.9 | 11.8 |
| 4 | Malignant neoplasms (C00-C97) | 16 | 3.0 | * |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 14 | 2.7 | * |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 9 | 1.7 | * |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 6 | 1.1 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 5 | 1.0 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 4 | 0.8 | * |
| 10 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 3 | 0.6 | * |
| 10 | Pregnancy, childbirth and the puerperium (O00-O99) | 3 | 0.6 | * |
| 12 | Diabetes mellitus (E10-E14) | 2 | 0.4 | * |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1 | 0.2 | * |
| 13 | Meningitis (G00,G03) | 1 | 0.2 | * |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1 | 0.2 | * |
| 13 | Influenza and pneumonia (J09-J18) | 1 | 0.2 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.2 | * |
| 13 | Legal intervention (Y35,Y89.0) | 1 | 0.2 | * |
| ... | All other causes (residual) | 50 | 9.5 | 11.4 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|-------|
| Non-Hispanic, American Indian or Alaska Native, both sexes, 25-34 years | | | | |
| ... | All causes | 997 | 100.0 | 255.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 412 | 41.3 | 105.6 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 126 | 12.6 | 32.3 |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 83 | 8.3 | 21.3 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 68 | 6.8 | 17.4 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 55 | 5.5 | 14.1 |
| 6 | Malignant neoplasms (C00-C97) | 41 | 4.1 | 10.5 |
| 7 | Septicemia (A40-A41) | 11 | 1.1 | * |
| 8 | Diabetes mellitus (E10-E14) | 10 | 1.0 | * |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 8 | 0.8 | * |
| 10 | Cerebrovascular diseases (I60-I69) | 7 | 0.7 | * |
| 10 | Pregnancy, childbirth and the puerperium (O00-O99) | 7 | 0.7 | * |
| 12 | Chronic lower respiratory diseases (J40-J47) | 6 | 0.6 | * |
| 13 | Influenza and pneumonia (J09-J18) | 5 | 0.5 | * |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 5 | 0.5 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4 | 0.4 | * |
| ... | All other causes (residual) | 149 | 14.9 | 38.2 |
| Non-Hispanic, American Indian or Alaska Native, both sexes, 35-44 years | | | | |
| ... | All causes | 1,277 | 100.0 | 391.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 332 | 26.0 | 101.8 |
| 2 | Chronic liver disease and cirrhosis (K70,K73-K74) | 180 | 14.1 | 55.2 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 139 | 10.9 | 42.6 |
| 4 | Malignant neoplasms (C00-C97) | 90 | 7.0 | 27.6 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 89 | 7.0 | 27.3 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 53 | 4.2 | 16.2 |
| 7 | Diabetes mellitus (E10-E14) | 47 | 3.7 | 14.4 |
| 8 | Cerebrovascular diseases (I60-I69) | 25 | 2.0 | 7.7 |
| 9 | Septicemia (A40-A41) | 20 | 1.6 | 6.1 |
| 10 | Influenza and pneumonia (J09-J18) | 13 | 1.0 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 12 | 0.9 | * |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 9 | 0.7 | * |
| 13 | Chronic lower respiratory diseases (J40-J47) | 7 | 0.5 | * |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 7 | 0.5 | * |
| 15 | Viral hepatitis (B15-B19) | 6 | 0.5 | * |
| 15 | Pregnancy, childbirth and the puerperium (O00-O99) | 6 | 0.5 | * |
| ... | All other causes (residual) | 242 | 19.0 | 74.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|---------|
| Non-Hispanic, American Indian or Alaska Native, both sexes, 45-54 years | | | | |
| ... | All causes | 2,365 | 100.0 | 688.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 383 | 16.2 | 111.5 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 367 | 15.5 | 106.8 |
| 3 | Malignant neoplasms (C00-C97) | 345 | 14.6 | 100.4 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 310 | 13.1 | 90.2 |
| 5 | Diabetes mellitus (E10-E14) | 142 | 6.0 | 41.3 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 87 | 3.7 | 25.3 |
| 7 | Cerebrovascular diseases (I60-I69) | 55 | 2.3 | 16.0 |
| 8 | Septicemia (A40-A41) | 43 | 1.8 | 12.5 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 42 | 1.8 | 12.2 |
| 10 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 37 | 1.6 | 10.8 |
| 11 | Viral hepatitis (B15-B19) | 30 | 1.3 | 8.7 |
| 11 | Influenza and pneumonia (J09-J18) | 30 | 1.3 | 8.7 |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 26 | 1.1 | 7.6 |
| 14 | Human immunodeficiency virus (HIV) disease (B20-B24) | 18 | 0.8 | * |
| 15 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 15 | 0.6 | * |
| ... | All other causes (residual) | 435 | 18.4 | 126.6 |
| Non-Hispanic, American Indian or Alaska Native, both sexes, 55-64 years | | | | |
| ... | All causes | 3,328 | 100.0 | 1,081.4 |
| 1 | Malignant neoplasms (C00-C97) | 749 | 22.5 | 243.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 671 | 20.2 | 218.0 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 270 | 8.1 | 87.7 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 264 | 7.9 | 85.8 |
| 5 | Diabetes mellitus (E10-E14) | 242 | 7.3 | 78.6 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 170 | 5.1 | 55.2 |
| 7 | Cerebrovascular diseases (I60-I69) | 93 | 2.8 | 30.2 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 68 | 2.0 | 22.1 |
| 9 | Septicemia (A40-A41) | 61 | 1.8 | 19.8 |
| 10 | Influenza and pneumonia (J09-J18) | 45 | 1.4 | 14.6 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 43 | 1.3 | 14.0 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 41 | 1.2 | 13.3 |
| 13 | Viral hepatitis (B15-B19) | 39 | 1.2 | 12.7 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 22 | 0.7 | 7.1 |
| 15 | Pneumonitis due to solids and liquids (J69) | 16 | 0.5 | * |
| ... | All other causes (residual) | 534 | 16.0 | 173.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|---------|
| Non-Hispanic, American Indian or Alaska Native, both sexes, 65-74 years | | | | |
| ... | All causes | 3,498 | 100.0 | 2,000.9 |
| 1 | Malignant neoplasms (C00-C97) | 927 | 26.5 | 530.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 708 | 20.2 | 405.0 |
| 3 | Diabetes mellitus (E10-E14) | 299 | 8.5 | 171.0 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 259 | 7.4 | 148.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 148 | 4.2 | 84.7 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 113 | 3.2 | 64.6 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 112 | 3.2 | 64.1 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 88 | 2.5 | 50.3 |
| 9 | Influenza and pneumonia (J09-J18) | 76 | 2.2 | 43.5 |
| 10 | Septicemia (A40-A41) | 75 | 2.1 | 42.9 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 33 | 0.9 | 18.9 |
| 12 | Alzheimer's disease (G30) | 24 | 0.7 | 13.7 |
| 13 | Viral hepatitis (B15-B19) | 18 | 0.5 | * |
| 14 | Pneumonitis due to solids and liquids (J69) | 17 | 0.5 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 15 | 0.4 | * |
| ... | All other causes (residual) | 586 | 16.8 | 335.2 |
| Non-Hispanic, American Indian or Alaska Native, both sexes, 75-84 years | | | | |
| ... | All causes | 3,224 | 100.0 | 4,531.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 748 | 23.2 | 1,051.3 |
| 2 | Malignant neoplasms (C00-C97) | 705 | 21.9 | 990.9 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 237 | 7.4 | 333.1 |
| 4 | Diabetes mellitus (E10-E14) | 206 | 6.4 | 289.5 |
| 5 | Cerebrovascular diseases (I60-I69) | 161 | 5.0 | 226.3 |
| 6 | Alzheimer's disease (G30) | 120 | 3.7 | 168.7 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 85 | 2.6 | 119.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 82 | 2.5 | 115.3 |
| 9 | Influenza and pneumonia (J09-J18) | 77 | 2.4 | 108.2 |
| 10 | Septicemia (A40-A41) | 58 | 1.8 | 81.5 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 45 | 1.4 | 63.2 |
| 12 | Parkinson's disease (G20-G21) | 43 | 1.3 | 60.4 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 39 | 1.2 | 54.8 |
| 14 | Pneumonitis due to solids and liquids (J69) | 22 | 0.7 | 30.9 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 18 | 0.6 | * |
| ... | All other causes (residual) | 578 | 17.9 | 812.4 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|---------|
| Non-Hispanic, American Indian or Alaska Native, both sexes, 85 years and over | | | | |
| ... | All causes | 2,344 | 100.0 | 9,792.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 579 | 24.7 | 2,418.8 |
| 2 | Malignant neoplasms (C00-C97) | 304 | 13.0 | 1,270.0 |
| 3 | Alzheimer's disease (G30) | 160 | 6.8 | 668.4 |
| 4 | Cerebrovascular diseases (I60-I69) | 151 | 6.4 | 630.8 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 129 | 5.5 | 538.9 |
| 6 | Diabetes mellitus (E10-E14) | 86 | 3.7 | 359.3 |
| 7 | Influenza and pneumonia (J09-J18) | 78 | 3.3 | 325.9 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 76 | 3.2 | 317.5 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 58 | 2.5 | 242.3 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 50 | 2.1 | 208.9 |
| 11 | Parkinson's disease (G20-G21) | 29 | 1.2 | 121.2 |
| 12 | Septicemia (A40-A41) | 26 | 1.1 | 108.6 |
| 13 | Pneumonitis due to solids and liquids (J69) | 25 | 1.1 | 104.4 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 17 | 0.7 | * |
| 14 | Nutritional deficiencies (E40-E64) | 17 | 0.7 | * |
| ... | All other causes (residual) | 559 | 23.8 | 2,335.3 |
| Non-Hispanic, American Indian or Alaska Native, male, all ages | | | | |
| ... | All causes | 9,869 | 100.0 | 747.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,916 | 19.4 | 145.1 |
| 2 | Malignant neoplasms (C00-C97) | 1,699 | 17.2 | 128.7 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,282 | 13.0 | 97.1 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 563 | 5.7 | 42.6 |
| 5 | Diabetes mellitus (E10-E14) | 537 | 5.4 | 40.7 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 398 | 4.0 | 30.1 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 395 | 4.0 | 29.9 |
| 8 | Cerebrovascular diseases (I60-I69) | 291 | 2.9 | 22.0 |
| 9 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 209 | 2.1 | 15.8 |
| 10 | Influenza and pneumonia (J09-J18) | 181 | 1.8 | 13.7 |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 159 | 1.6 | 12.0 |
| 12 | Septicemia (A40-A41) | 130 | 1.3 | 9.8 |
| 13 | Alzheimer's disease (G30) | 100 | 1.0 | 7.6 |
| 14 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 87 | 0.9 | 6.6 |
| 15 | Viral hepatitis (B15-B19) | 70 | 0.7 | 5.3 |
| ... | All other causes (residual) | 1,852 | 18.8 | 140.3 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|-------|
| Non-Hispanic, American Indian or Alaska Native, male, 1-4 years | | | | |
| ... | All causes | 45 | 100.0 | 55.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 14 | 31.1 | * |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 4 | 8.9 | * |
| 3 | Malignant neoplasms (C00-C97) | 3 | 6.7 | * |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 3 | 6.7 | * |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1 | 2.2 | * |
| 5 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1 | 2.2 | * |
| 5 | Cerebrovascular diseases (I60-I69) | 1 | 2.2 | * |
| 5 | Chronic lower respiratory diseases (J40-J47) | 1 | 2.2 | * |
| ... | All other causes (residual) | 17 | 37.8 | * |
| Non-Hispanic, American Indian or Alaska Native, male, 5-14 years | | | | |
| ... | All causes | 51 | 100.0 | 24.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 18 | 35.3 | * |
| 2 | Malignant neoplasms (C00-C97) | 8 | 15.7 | * |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 4 | 7.8 | * |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 4 | 7.8 | * |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 2 | 3.9 | * |
| 6 | Influenza and pneumonia (J09-J18) | 1 | 2.0 | * |
| 6 | Chronic lower respiratory diseases (J40-J47) | 1 | 2.0 | * |
| ... | All other causes (residual) | 13 | 25.5 | * |
| Non-Hispanic, American Indian or Alaska Native, male, 15-24 years | | | | |
| ... | All causes | 362 | 100.0 | 161.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 138 | 38.1 | 61.7 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 110 | 30.4 | 49.2 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 49 | 13.5 | 21.9 |
| 4 | Malignant neoplasms (C00-C97) | 10 | 2.8 | * |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7 | 1.9 | * |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 5 | 1.4 | * |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 5 | 1.4 | * |
| 8 | Diabetes mellitus (E10-E14) | 2 | 0.6 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 2 | 0.6 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 2 | 0.6 | * |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 1 | 0.3 | * |
| 11 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1 | 0.3 | * |
| 11 | Influenza and pneumonia (J09-J18) | 1 | 0.3 | * |
| 11 | Legal intervention (Y35,Y89.0) | 1 | 0.3 | * |
| ... | All other causes (residual) | 28 | 7.7 | 12.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|-------|
| Non-Hispanic, American Indian or Alaska Native, male, 25-34 years | | | | |
| ... | All causes | 662 | 100.0 | 339.4 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 282 | 42.6 | 144.6 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 99 | 15.0 | 50.8 |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 52 | 7.9 | 26.7 |
| 3 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 52 | 7.9 | 26.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 36 | 5.4 | 18.5 |
| 6 | Malignant neoplasms (C00-C97) | 20 | 3.0 | 10.3 |
| 7 | Septicemia (A40-A41) | 5 | 0.8 | * |
| 8 | Viral hepatitis (B15-B19) | 3 | 0.5 | * |
| 8 | Human immunodeficiency virus (HIV) disease (B20-B24) | 3 | 0.5 | * |
| 8 | Diabetes mellitus (E10-E14) | 3 | 0.5 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 3 | 0.5 | * |
| 8 | Influenza and pneumonia (J09-J18) | 3 | 0.5 | * |
| 8 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 3 | 0.5 | * |
| 8 | Legal intervention (Y35,Y89.0) | 3 | 0.5 | * |
| 15 | Chronic lower respiratory diseases (J40-J47) | 2 | 0.3 | * |
| 15 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2 | 0.3 | * |
| ... | All other causes (residual) | 91 | 13.7 | 46.6 |
| Non-Hispanic, American Indian or Alaska Native, male, 35-44 years | | | | |
| ... | All causes | 809 | 100.0 | 504.3 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 239 | 29.5 | 149.0 |
| 2 | Chronic liver disease and cirrhosis (K70,K73-K74) | 99 | 12.2 | 61.7 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 85 | 10.5 | 53.0 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 65 | 8.0 | 40.5 |
| 5 | Malignant neoplasms (C00-C97) | 44 | 5.4 | 27.4 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 44 | 5.4 | 27.4 |
| 7 | Diabetes mellitus (E10-E14) | 31 | 3.8 | 19.3 |
| 8 | Cerebrovascular diseases (I60-I69) | 15 | 1.9 | * |
| 9 | Septicemia (A40-A41) | 12 | 1.5 | * |
| 10 | Influenza and pneumonia (J09-J18) | 9 | 1.1 | * |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 7 | 0.9 | * |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 5 | 0.6 | * |
| 13 | Viral hepatitis (B15-B19) | 4 | 0.5 | * |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 4 | 0.5 | * |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 4 | 0.5 | * |
| ... | All other causes (residual) | 142 | 17.6 | 88.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|---------|
| Non-Hispanic, American Indian or Alaska Native, male, 45-54 years | | | | |
| ... | All causes | 1,421 | 100.0 | 857.1 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 267 | 18.8 | 161.0 |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 228 | 16.0 | 137.5 |
| 3 | Malignant neoplasms (C00-C97) | 182 | 12.8 | 109.8 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 167 | 11.8 | 100.7 |
| 5 | Diabetes mellitus (E10-E14) | 76 | 5.3 | 45.8 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 66 | 4.6 | 39.8 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 32 | 2.3 | 19.3 |
| 8 | Cerebrovascular diseases (I60-I69) | 25 | 1.8 | 15.1 |
| 9 | Septicemia (A40-A41) | 24 | 1.7 | 14.5 |
| 9 | Viral hepatitis (B15-B19) | 24 | 1.7 | 14.5 |
| 11 | Chronic lower respiratory diseases (J40-J47) | 18 | 1.3 | * |
| 12 | Influenza and pneumonia (J09-J18) | 17 | 1.2 | * |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 12 | 0.8 | * |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 12 | 0.8 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 12 | 0.8 | * |
| ... | All other causes (residual) | 259 | 18.2 | 156.2 |
| Non-Hispanic, American Indian or Alaska Native, male, 55-64 years | | | | |
| ... | All causes | 1,982 | 100.0 | 1,371.7 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 470 | 23.7 | 325.3 |
| 2 | Malignant neoplasms (C00-C97) | 405 | 20.4 | 280.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 193 | 9.7 | 133.6 |
| 4 | Chronic liver disease and cirrhosis (K70,K73-K74) | 160 | 8.1 | 110.7 |
| 5 | Diabetes mellitus (E10-E14) | 141 | 7.1 | 97.6 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 72 | 3.6 | 49.8 |
| 7 | Cerebrovascular diseases (I60-I69) | 50 | 2.5 | 34.6 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 38 | 1.9 | 26.3 |
| 9 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 31 | 1.6 | 21.5 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 25 | 1.3 | 17.3 |
| 10 | Influenza and pneumonia (J09-J18) | 25 | 1.3 | 17.3 |
| 12 | Viral hepatitis (B15-B19) | 24 | 1.2 | 16.6 |
| 13 | Septicemia (A40-A41) | 23 | 1.2 | 15.9 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 15 | 0.8 | * |
| 15 | Pneumonitis due to solids and liquids (J69) | 11 | 0.6 | * |
| ... | All other causes (residual) | 299 | 15.1 | 206.9 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|---------|
| Non-Hispanic, American Indian or Alaska Native, male, 65-74 years | | | | |
| ... | All causes | 1,908 | 100.0 | 2,334.6 |
| 1 | Malignant neoplasms (C00-C97) | 520 | 27.3 | 636.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 423 | 22.2 | 517.6 |
| 3 | Diabetes mellitus (E10-E14) | 156 | 8.2 | 190.9 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 128 | 6.7 | 156.6 |
| 5 | Cerebrovascular diseases (I60-I69) | 79 | 4.1 | 96.7 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 77 | 4.0 | 94.2 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 57 | 3.0 | 69.7 |
| 8 | Influenza and pneumonia (J09-J18) | 46 | 2.4 | 56.3 |
| 9 | Septicemia (A40-A41) | 38 | 2.0 | 46.5 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 37 | 1.9 | 45.3 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 17 | 0.9 | * |
| 12 | Pneumonitis due to solids and liquids (J69) | 13 | 0.7 | * |
| 13 | Alzheimer's disease (G30) | 11 | 0.6 | * |
| 13 | Aortic aneurysm and dissection (I71) | 11 | 0.6 | * |
| 15 | Viral hepatitis (B15-B19) | 9 | 0.5 | * |
| ... | All other causes (residual) | 286 | 15.0 | 349.9 |
| Non-Hispanic, American Indian or Alaska Native, male, 75-84 years | | | | |
| ... | All causes | 1,593 | 100.0 | 5,211.3 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 399 | 25.0 | 1,305.3 |
| 2 | Malignant neoplasms (C00-C97) | 373 | 23.4 | 1,220.2 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 113 | 7.1 | 369.7 |
| 4 | Diabetes mellitus (E10-E14) | 98 | 6.2 | 320.6 |
| 5 | Cerebrovascular diseases (I60-I69) | 70 | 4.4 | 229.0 |
| 6 | Alzheimer's disease (G30) | 49 | 3.1 | 160.3 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 49 | 3.1 | 160.3 |
| 8 | Influenza and pneumonia (J09-J18) | 42 | 2.6 | 137.4 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 40 | 2.5 | 130.9 |
| 10 | Parkinson's disease (G20-G21) | 24 | 1.5 | 78.5 |
| 11 | Septicemia (A40-A41) | 20 | 1.3 | 65.4 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 19 | 1.2 | * |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 16 | 1.0 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 12 | 0.8 | * |
| 15 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 11 | 0.7 | * |
| ... | All other causes (residual) | 258 | 16.2 | 844.0 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|----------|
| Non-Hispanic, American Indian or Alaska Native, male, 85 years and over | | | | |
| ... | All causes | 875 | 100.0 | 10,272.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 227 | 25.9 | 2,664.9 |
| 2 | Malignant neoplasms (C00-C97) | 133 | 15.2 | 1,561.4 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 58 | 6.6 | 680.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 45 | 5.1 | 528.3 |
| 5 | Alzheimer's disease (G30) | 39 | 4.5 | 457.9 |
| 6 | Influenza and pneumonia (J09-J18) | 35 | 4.0 | 410.9 |
| 7 | Diabetes mellitus (E10-E14) | 30 | 3.4 | 352.2 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 27 | 3.1 | 317.0 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 24 | 2.7 | 281.8 |
| 10 | Parkinson's disease (G20-G21) | 19 | 2.2 | * |
| 11 | Pneumonitis due to solids and liquids (J69) | 17 | 1.9 | * |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 11 | 1.3 | * |
| 13 | Septicemia (A40-A41) | 7 | 0.8 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 6 | 0.7 | * |
| 14 | Cholelithiasis and other disorders of gallbladder (K80-K82) | 6 | 0.7 | * |
| ... | All other causes (residual) | 191 | 21.8 | 2,242.3 |
| Non-Hispanic, American Indian or Alaska Native, female, all ages | | | | |
| ... | All causes | 8,170 | 100.0 | 596.7 |
| 1 | Malignant neoplasms (C00-C97) | 1,493 | 18.3 | 109.0 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,387 | 17.0 | 101.3 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 671 | 8.2 | 49.0 |
| 4 | Diabetes mellitus (E10-E14) | 497 | 6.1 | 36.3 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 461 | 5.6 | 33.7 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 443 | 5.4 | 32.4 |
| 7 | Cerebrovascular diseases (I60-I69) | 355 | 4.3 | 25.9 |
| 8 | Alzheimer's disease (G30) | 207 | 2.5 | 15.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 181 | 2.2 | 13.2 |
| 10 | Septicemia (A40-A41) | 166 | 2.0 | 12.1 |
| 11 | Influenza and pneumonia (J09-J18) | 149 | 1.8 | 10.9 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 140 | 1.7 | 10.2 |
| 13 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 105 | 1.3 | 7.7 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 53 | 0.6 | 3.9 |
| 15 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 52 | 0.6 | 3.8 |
| ... | All other causes (residual) | 1,810 | 22.2 | 132.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| Non-Hispanic, American Indian or Alaska Native, female, 1-4 years | | | | |
| ... | All causes | 37 | 100.0 | 47.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 16 | 43.2 | * |
| 2 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 6 | 16.2 | * |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 5 | 13.5 | * |
| 4 | Meningitis (G00,G03) | 1 | 2.7 | * |
| 4 | Pneumonitis due to solids and liquids (J69) | 1 | 2.7 | * |
| ... | All other causes (residual) | 8 | 21.6 | * |
| Non-Hispanic, American Indian or Alaska Native, female, 5-14 years | | | | |
| ... | All causes | 48 | 100.0 | 23.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 13 | 27.1 | * |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 10 | 20.8 | * |
| 3 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 5 | 10.4 | * |
| 4 | Malignant neoplasms (C00-C97) | 3 | 6.3 | * |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 3 | 6.3 | * |
| 6 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2 | 4.2 | * |
| 6 | Chronic lower respiratory diseases (J40-J47) | 2 | 4.2 | * |
| 6 | Certain conditions originating in the perinatal period (P00-P96) | 2 | 4.2 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1 | 2.1 | * |
| ... | All other causes (residual) | 7 | 14.6 | * |
| Non-Hispanic, American Indian or Alaska Native, female, 15-24 years | | | | |
| ... | All causes | 163 | 100.0 | 75.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 67 | 41.1 | 31.0 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 40 | 24.5 | 18.5 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7 | 4.3 | * |
| 4 | Malignant neoplasms (C00-C97) | 6 | 3.7 | * |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 4 | 2.5 | * |
| 6 | Chronic lower respiratory diseases (J40-J47) | 3 | 1.8 | * |
| 6 | Pregnancy, childbirth and the puerperium (O00-O99) | 3 | 1.8 | * |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 3 | 1.8 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2 | 1.2 | * |
| 9 | Cerebrovascular diseases (I60-I69) | 2 | 1.2 | * |
| 11 | Meningitis (G00,G03) | 1 | 0.6 | * |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1 | 0.6 | * |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 1 | 0.6 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.6 | * |
| ... | All other causes (residual) | 22 | 13.5 | 10.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|-------|
| Non-Hispanic, American Indian or Alaska Native, female, 25-34 years | | | | |
| ... | All causes | 335 | 100.0 | 171.7 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 130 | 38.8 | 66.6 |
| 2 | Chronic liver disease and cirrhosis (K70,K73-K74) | 31 | 9.3 | 15.9 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 27 | 8.1 | 13.8 |
| 4 | Malignant neoplasms (C00-C97) | 21 | 6.3 | 10.8 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 19 | 5.7 | * |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 16 | 4.8 | * |
| 7 | Diabetes mellitus (E10-E14) | 7 | 2.1 | * |
| 7 | Pregnancy, childbirth and the puerperium (O00-O99) | 7 | 2.1 | * |
| 9 | Septicemia (A40-A41) | 6 | 1.8 | * |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 5 | 1.5 | * |
| 11 | Cerebrovascular diseases (I60-I69) | 4 | 1.2 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 4 | 1.2 | * |
| 13 | Influenza and pneumonia (J09-J18) | 2 | 0.6 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 2 | 0.6 | * |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 2 | 0.6 | * |
| ... | All other causes (residual) | 52 | 15.5 | 26.7 |
| Non-Hispanic, American Indian or Alaska Native, female, 35-44 years | | | | |
| ... | All causes | 468 | 100.0 | 282.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 93 | 19.9 | 56.1 |
| 2 | Chronic liver disease and cirrhosis (K70,K73-K74) | 81 | 17.3 | 48.8 |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 54 | 11.5 | 32.6 |
| 4 | Malignant neoplasms (C00-C97) | 46 | 9.8 | 27.7 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 24 | 5.1 | 14.5 |
| 6 | Diabetes mellitus (E10-E14) | 16 | 3.4 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 10 | 2.1 | * |
| 8 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 9 | 1.9 | * |
| 9 | Septicemia (A40-A41) | 8 | 1.7 | * |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7 | 1.5 | * |
| 11 | Pregnancy, childbirth and the puerperium (O00-O99) | 6 | 1.3 | * |
| 12 | Influenza and pneumonia (J09-J18) | 4 | 0.9 | * |
| 12 | Chronic lower respiratory diseases (J40-J47) | 4 | 0.9 | * |
| 14 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 3 | 0.6 | * |
| 15 | Viral hepatitis (B15-B19) | 2 | 0.4 | * |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 2 | 0.4 | * |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2 | 0.4 | * |
| 15 | Aortic aneurysm and dissection (I71) | 2 | 0.4 | * |
| 15 | Infections of kidney (N10-N12,N13.6,N15.1) | 2 | 0.4 | * |
| ... | All other causes (residual) | 93 | 19.9 | 56.1 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|-------|
| Non-Hispanic, American Indian or Alaska Native, female, 45-54 years | | | | |
| ... | All causes | 944 | 100.0 | 531.0 |
| 1 | Malignant neoplasms (C00-C97) | 163 | 17.3 | 91.7 |
| 2 | Chronic liver disease and cirrhosis (K70,K73-K74) | 143 | 15.1 | 80.4 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 139 | 14.7 | 78.2 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 116 | 12.3 | 65.3 |
| 5 | Diabetes mellitus (E10-E14) | 66 | 7.0 | 37.1 |
| 6 | Cerebrovascular diseases (I60-I69) | 30 | 3.2 | 16.9 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 24 | 2.5 | 13.5 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 21 | 2.2 | 11.8 |
| 9 | Septicemia (A40-A41) | 19 | 2.0 | * |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 14 | 1.5 | * |
| 11 | Influenza and pneumonia (J09-J18) | 13 | 1.4 | * |
| 12 | Viral hepatitis (B15-B19) | 6 | 0.6 | * |
| 12 | Human immunodeficiency virus (HIV) disease (B20-B24) | 6 | 0.6 | * |
| 14 | Pneumonitis due to solids and liquids (J69) | 5 | 0.5 | * |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 5 | 0.5 | * |
| ... | All other causes (residual) | 174 | 18.4 | 97.9 |
| Non-Hispanic, American Indian or Alaska Native, female, 55-64 years | | | | |
| ... | All causes | 1,346 | 100.0 | 824.5 |
| 1 | Malignant neoplasms (C00-C97) | 344 | 25.6 | 210.7 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 201 | 14.9 | 123.1 |
| 3 | Chronic liver disease and cirrhosis (K70,K73-K74) | 104 | 7.7 | 63.7 |
| 4 | Diabetes mellitus (E10-E14) | 101 | 7.5 | 61.9 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 98 | 7.3 | 60.0 |
| 6 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 77 | 5.7 | 47.2 |
| 7 | Cerebrovascular diseases (I60-I69) | 43 | 3.2 | 26.3 |
| 8 | Septicemia (A40-A41) | 38 | 2.8 | 23.3 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 30 | 2.2 | 18.4 |
| 10 | Influenza and pneumonia (J09-J18) | 20 | 1.5 | 12.3 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 16 | 1.2 | * |
| 12 | Viral hepatitis (B15-B19) | 15 | 1.1 | * |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 12 | 0.9 | * |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 7 | 0.5 | * |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 5 | 0.4 | * |
| 15 | Pneumonitis due to solids and liquids (J69) | 5 | 0.4 | * |
| ... | All other causes (residual) | 230 | 17.1 | 140.9 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|---------|
| Non-Hispanic, American Indian or Alaska Native, female, 65-74 years | | | | |
| ... | All causes | 1,590 | 100.0 | 1,708.0 |
| 1 | Malignant neoplasms (C00-C97) | 407 | 25.6 | 437.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 285 | 17.9 | 306.2 |
| 3 | Diabetes mellitus (E10-E14) | 143 | 9.0 | 153.6 |
| 4 | Chronic lower respiratory diseases (J40-J47) | 131 | 8.2 | 140.7 |
| 5 | Cerebrovascular diseases (I60-I69) | 69 | 4.3 | 74.1 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 55 | 3.5 | 59.1 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 51 | 3.2 | 54.8 |
| 8 | Septicemia (A40-A41) | 37 | 2.3 | 39.7 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 36 | 2.3 | 38.7 |
| 10 | Influenza and pneumonia (J09-J18) | 30 | 1.9 | 32.2 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 16 | 1.0 | * |
| 12 | Alzheimer's disease (G30) | 13 | 0.8 | * |
| 13 | Viral hepatitis (B15-B19) | 9 | 0.6 | * |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 0.4 | * |
| 15 | Parkinson's disease (G20-G21) | 6 | 0.4 | * |
| ... | All other causes (residual) | 295 | 18.6 | 316.9 |
| Non-Hispanic, American Indian or Alaska Native, female, 75-84 years | | | | |
| ... | All causes | 1,631 | 100.0 | 4,019.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 349 | 21.4 | 860.0 |
| 2 | Malignant neoplasms (C00-C97) | 332 | 20.4 | 818.1 |
| 3 | Chronic lower respiratory diseases (J40-J47) | 124 | 7.6 | 305.6 |
| 4 | Diabetes mellitus (E10-E14) | 108 | 6.6 | 266.1 |
| 5 | Cerebrovascular diseases (I60-I69) | 91 | 5.6 | 224.2 |
| 6 | Alzheimer's disease (G30) | 71 | 4.4 | 175.0 |
| 7 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 42 | 2.6 | 103.5 |
| 8 | Septicemia (A40-A41) | 38 | 2.3 | 93.6 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 36 | 2.2 | 88.7 |
| 10 | Influenza and pneumonia (J09-J18) | 35 | 2.1 | 86.2 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 29 | 1.8 | 71.5 |
| 12 | Chronic liver disease and cirrhosis (K70,K73-K74) | 20 | 1.2 | 49.3 |
| 13 | Parkinson's disease (G20-G21) | 19 | 1.2 | * |
| 14 | Pneumonitis due to solids and liquids (J69) | 14 | 0.9 | * |
| 15 | Aortic aneurysm and dissection (I71) | 7 | 0.4 | * |
| ... | All other causes (residual) | 316 | 19.4 | 778.7 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|---------|
| Non-Hispanic, American Indian or Alaska Native, female, 85 years and over | | | | |
| ... | All causes | 1,469 | 100.0 | 9,527.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 352 | 24.0 | 2,282.9 |
| 2 | Malignant neoplasms (C00-C97) | 171 | 11.6 | 1,109.0 |
| 3 | Alzheimer's disease (G30) | 121 | 8.2 | 784.7 |
| 4 | Cerebrovascular diseases (I60-I69) | 106 | 7.2 | 687.5 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 71 | 4.8 | 460.5 |
| 6 | Diabetes mellitus (E10-E14) | 56 | 3.8 | 363.2 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 49 | 3.3 | 317.8 |
| 8 | Influenza and pneumonia (J09-J18) | 43 | 2.9 | 278.9 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 39 | 2.7 | 252.9 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 34 | 2.3 | 220.5 |
| 11 | Septicemia (A40-A41) | 19 | 1.3 | * |
| 12 | Nutritional deficiencies (E40-E64) | 12 | 0.8 | * |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 11 | 0.7 | * |
| 14 | Parkinson's disease (G20-G21) | 10 | 0.7 | * |
| 15 | Atherosclerosis (I70) | 8 | 0.5 | * |
| 15 | Pneumonitis due to solids and liquids (J69) | 8 | 0.5 | * |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 8 | 0.5 | * |
| ... | All other causes (residual) | 351 | 23.9 | 2,276.4 |
| Non-Hispanic, Asian or Pacific Islander, both sexes, all ages | | | | |
| ... | All causes | 65,277 | 100.0 | 341.5 |
| 1 | Malignant neoplasms (C00-C97) | 17,306 | 26.5 | 90.5 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 13,974 | 21.4 | 73.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 4,798 | 7.4 | 25.1 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 2,839 | 4.3 | 14.9 |
| 5 | Diabetes mellitus (E10-E14) | 2,599 | 4.0 | 13.6 |
| 6 | Alzheimer's disease (G30) | 2,180 | 3.3 | 11.4 |
| 7 | Influenza and pneumonia (J09-J18) | 2,161 | 3.3 | 11.3 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 1,874 | 2.9 | 9.8 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1,318 | 2.0 | 6.9 |
| 10 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 1,273 | 2.0 | 6.7 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 1,136 | 1.7 | 5.9 |
| 12 | Septicemia (A40-A41) | 798 | 1.2 | 4.2 |
| 13 | Parkinson's disease (G20-G21) | 747 | 1.1 | 3.9 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 598 | 0.9 | 3.1 |
| 15 | Pneumonitis due to solids and liquids (J69) | 500 | 0.8 | 2.6 |
| ... | All other causes (residual) | 11,176 | 17.1 | 58.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| Non-Hispanic, Asian or Pacific Islander, both sexes, 1-4 years | | | | |
| ... | All causes | 141 | 100.0 | 15.3 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 26 | 18.4 | 2.8 |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 20 | 14.2 | 2.2 |
| 3 | Malignant neoplasms (C00-C97) | 13 | 9.2 | * |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 10 | 7.1 | * |
| 5 | Influenza and pneumonia (J09-J18) | 6 | 4.3 | * |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 6 | 4.3 | * |
| 7 | Septicemia (A40-A41) | 3 | 2.1 | * |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 3 | 2.1 | * |
| 9 | Certain conditions originating in the perinatal period (P00-P96) | 2 | 1.4 | * |
| 10 | Tuberculosis (A16-A19) | 1 | 0.7 | * |
| 10 | Anemias (D50-D64) | 1 | 0.7 | * |
| 10 | Acute bronchitis and bronchiolitis (J20-J21) | 1 | 0.7 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 1 | 0.7 | * |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.7 | * |
| ... | All other causes (residual) | 47 | 33.3 | 5.1 |
| Non-Hispanic, Asian or Pacific Islander, both sexes, 5-14 years | | | | |
| ... | All causes | 212 | 100.0 | 9.0 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 49 | 23.1 | 2.1 |
| 2 | Malignant neoplasms (C00-C97) | 40 | 18.9 | 1.7 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 15 | 7.1 | * |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 14 | 6.6 | * |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 7 | 3.3 | * |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 7 | 3.3 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 5 | 2.4 | * |
| 8 | Septicemia (A40-A41) | 4 | 1.9 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 3 | 1.4 | * |
| 10 | Anemias (D50-D64) | 2 | 0.9 | * |
| 10 | Influenza and pneumonia (J09-J18) | 2 | 0.9 | * |
| 10 | Complications of medical and surgical care (Y40-Y84,Y88) | 2 | 0.9 | * |
| 13 | Meningitis (G00,G03) | 1 | 0.5 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.5 | * |
| ... | All other causes (residual) | 60 | 28.3 | 2.6 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Non-Hispanic, Asian or Pacific Islander, both sexes, 15-24 years | | | | |
| ... | All causes | 873 | 100.0 | 33.9 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 277 | 31.7 | 10.7 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 265 | 30.4 | 10.3 |
| 3 | Malignant neoplasms (C00-C97) | 84 | 9.6 | 3.3 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 46 | 5.3 | 1.8 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 40 | 4.6 | 1.6 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 16 | 1.8 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 7 | 0.8 | * |
| 8 | Influenza and pneumonia (J09-J18) | 5 | 0.6 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 5 | 0.6 | * |
| 10 | Complications of medical and surgical care (Y40-Y84,Y88) | 4 | 0.5 | * |
| 11 | Septicemia (A40-A41) | 3 | 0.3 | * |
| 11 | Diabetes mellitus (E10-E14) | 3 | 0.3 | * |
| 11 | Aortic aneurysm and dissection (I71) | 3 | 0.3 | * |
| 11 | Pneumonitis due to solids and liquids (J69) | 3 | 0.3 | * |
| 11 | Legal intervention (Y35,Y89.0) | 3 | 0.3 | * |
| ... | All other causes (residual) | 109 | 12.5 | 4.2 |
| Non-Hispanic, Asian or Pacific Islander, both sexes, 25-34 years | | | | |
| ... | All causes | 1,332 | 100.0 | 40.8 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 358 | 26.9 | 11.0 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 266 | 20.0 | 8.2 |
| 3 | Malignant neoplasms (C00-C97) | 179 | 13.4 | 5.5 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 104 | 7.8 | 3.2 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 68 | 5.1 | 2.1 |
| 6 | Cerebrovascular diseases (I60-I69) | 29 | 2.2 | 0.9 |
| 7 | Pregnancy, childbirth and the puerperium (O00-O99) | 21 | 1.6 | 0.6 |
| 8 | Diabetes mellitus (E10-E14) | 17 | 1.3 | * |
| 9 | Influenza and pneumonia (J09-J18) | 12 | 0.9 | * |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 11 | 0.8 | * |
| 10 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 11 | 0.8 | * |
| 12 | Chronic lower respiratory diseases (J40-J47) | 9 | 0.7 | * |
| 12 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 9 | 0.7 | * |
| 14 | Septicemia (A40-A41) | 8 | 0.6 | * |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 7 | 0.5 | * |
| ... | All other causes (residual) | 223 | 16.7 | 6.8 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, Asian or Pacific Islander, both sexes, 35-44 years | | | | |
| ... | All causes | 2,047 | 100.0 | 67.3 |
| 1 | Malignant neoplasms (C00-C97) | 627 | 30.6 | 20.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 306 | 14.9 | 10.1 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 229 | 11.2 | 7.5 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 180 | 8.8 | 5.9 |
| 5 | Cerebrovascular diseases (I60-I69) | 114 | 5.6 | 3.7 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 56 | 2.7 | 1.8 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 56 | 2.7 | 1.8 |
| 8 | Diabetes mellitus (E10-E14) | 46 | 2.2 | 1.5 |
| 9 | Influenza and pneumonia (J09-J18) | 23 | 1.1 | 0.8 |
| 10 | Pregnancy, childbirth and the puerperium (O00-O99) | 21 | 1.0 | 0.7 |
| 11 | Septicemia (A40-A41) | 20 | 1.0 | 0.7 |
| 11 | Human immunodeficiency virus (HIV) disease (B20-B24) | 20 | 1.0 | 0.7 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 14 | 0.7 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 14 | 0.7 | * |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 14 | 0.7 | * |
| ... | All other causes (residual) | 307 | 15.0 | 10.1 |
| Non-Hispanic, Asian or Pacific Islander, both sexes, 45-54 years | | | | |
| ... | All causes | 4,212 | 100.0 | 164.0 |
| 1 | Malignant neoplasms (C00-C97) | 1,678 | 39.8 | 65.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 759 | 18.0 | 29.6 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 277 | 6.6 | 10.8 |
| 4 | Cerebrovascular diseases (I60-I69) | 244 | 5.8 | 9.5 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 204 | 4.8 | 7.9 |
| 6 | Diabetes mellitus (E10-E14) | 141 | 3.3 | 5.5 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 101 | 2.4 | 3.9 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 53 | 1.3 | 2.1 |
| 9 | Influenza and pneumonia (J09-J18) | 45 | 1.1 | 1.8 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 44 | 1.0 | 1.7 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 39 | 0.9 | 1.5 |
| 12 | Septicemia (A40-A41) | 35 | 0.8 | 1.4 |
| 13 | Viral hepatitis (B15-B19) | 34 | 0.8 | 1.3 |
| 13 | Chronic lower respiratory diseases (J40-J47) | 34 | 0.8 | 1.3 |
| 15 | Human immunodeficiency virus (HIV) disease (B20-B24) | 25 | 0.6 | 1.0 |
| ... | All other causes (residual) | 499 | 11.8 | 19.4 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, Asian or Pacific Islander, both sexes, 55-64 years | | | | |
| ... | All causes | 8,009 | 100.0 | 388.6 |
| 1 | Malignant neoplasms (C00-C97) | 3,384 | 42.3 | 164.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,582 | 19.8 | 76.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 514 | 6.4 | 24.9 |
| 4 | Diabetes mellitus (E10-E14) | 344 | 4.3 | 16.7 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 300 | 3.7 | 14.6 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 163 | 2.0 | 7.9 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 161 | 2.0 | 7.8 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 134 | 1.7 | 6.5 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 105 | 1.3 | 5.1 |
| 10 | Septicemia (A40-A41) | 104 | 1.3 | 5.0 |
| 10 | Influenza and pneumonia (J09-J18) | 104 | 1.3 | 5.0 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 86 | 1.1 | 4.2 |
| 13 | Viral hepatitis (B15-B19) | 57 | 0.7 | 2.8 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 52 | 0.6 | 2.5 |
| 15 | Aortic aneurysm and dissection (I71) | 46 | 0.6 | 2.2 |
| ... | All other causes (residual) | 873 | 10.9 | 42.4 |
| Non-Hispanic, Asian or Pacific Islander, both sexes, 65-74 years | | | | |
| ... | All causes | 11,496 | 100.0 | 895.3 |
| 1 | Malignant neoplasms (C00-C97) | 4,369 | 38.0 | 340.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,314 | 20.1 | 180.2 |
| 3 | Cerebrovascular diseases (I60-I69) | 787 | 6.8 | 61.3 |
| 4 | Diabetes mellitus (E10-E14) | 604 | 5.3 | 47.0 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 355 | 3.1 | 27.6 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 300 | 2.6 | 23.4 |
| 7 | Influenza and pneumonia (J09-J18) | 261 | 2.3 | 20.3 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 224 | 1.9 | 17.4 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 156 | 1.4 | 12.1 |
| 10 | Septicemia (A40-A41) | 155 | 1.3 | 12.1 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 120 | 1.0 | 9.3 |
| 12 | Parkinson's disease (G20-G21) | 113 | 1.0 | 8.8 |
| 13 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 91 | 0.8 | 7.1 |
| 14 | Alzheimer's disease (G30) | 86 | 0.7 | 6.7 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 83 | 0.7 | 6.5 |
| ... | All other causes (residual) | 1,478 | 12.9 | 115.1 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Non-Hispanic, Asian or Pacific Islander, both sexes, 75-84 years | | | | |
| ... | All causes | 15,915 | 100.0 | 2,630.2 |
| 1 | Malignant neoplasms (C00-C97) | 4,247 | 26.7 | 701.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,499 | 22.0 | 578.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,393 | 8.8 | 230.2 |
| 4 | Diabetes mellitus (E10-E14) | 788 | 5.0 | 130.2 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 614 | 3.9 | 101.5 |
| 6 | Influenza and pneumonia (J09-J18) | 585 | 3.7 | 96.7 |
| 7 | Alzheimer's disease (G30) | 508 | 3.2 | 84.0 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 450 | 2.8 | 74.4 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 393 | 2.5 | 64.9 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 299 | 1.9 | 49.4 |
| 11 | Parkinson's disease (G20-G21) | 292 | 1.8 | 48.3 |
| 12 | Septicemia (A40-A41) | 210 | 1.3 | 34.7 |
| 13 | Pneumonitis due to solids and liquids (J69) | 131 | 0.8 | 21.6 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 113 | 0.7 | 18.7 |
| 15 | Chronic liver disease and cirrhosis (K70,K73-K74) | 101 | 0.6 | 16.7 |
| ... | All other causes (residual) | 2,292 | 14.4 | 378.8 |
| Non-Hispanic, Asian or Pacific Islander, both sexes, 85 years and over | | | | |
| ... | All causes | 20,147 | 100.0 | 8,653.4 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5,332 | 26.5 | 2,290.2 |
| 2 | Malignant neoplasms (C00-C97) | 2,680 | 13.3 | 1,151.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 1,700 | 8.4 | 730.2 |
| 4 | Alzheimer's disease (G30) | 1,565 | 7.8 | 672.2 |
| 5 | Influenza and pneumonia (J09-J18) | 1,113 | 5.5 | 478.0 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 792 | 3.9 | 340.2 |
| 7 | Diabetes mellitus (E10-E14) | 656 | 3.3 | 281.8 |
| 8 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 538 | 2.7 | 231.1 |
| 9 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 495 | 2.5 | 212.6 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 484 | 2.4 | 207.9 |
| 11 | Parkinson's disease (G20-G21) | 327 | 1.6 | 140.5 |
| 12 | Pneumonitis due to solids and liquids (J69) | 276 | 1.4 | 118.5 |
| 13 | Septicemia (A40-A41) | 249 | 1.2 | 106.9 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 126 | 0.6 | 54.1 |
| 15 | Aortic aneurysm and dissection (I71) | 86 | 0.4 | 36.9 |
| ... | All other causes (residual) | 3,728 | 18.5 | 1,601.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, Asian or Pacific Islander, male, all ages | | | | |
| ... | All causes | 33,306 | 100.0 | 364.9 |
| 1 | Malignant neoplasms (C00-C97) | 8,774 | 26.3 | 96.1 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7,612 | 22.9 | 83.4 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,153 | 6.5 | 23.6 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,788 | 5.4 | 19.6 |
| 5 | Diabetes mellitus (E10-E14) | 1,322 | 4.0 | 14.5 |
| 6 | Influenza and pneumonia (J09-J18) | 1,109 | 3.3 | 12.2 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 1,061 | 3.2 | 11.6 |
| 8 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 855 | 2.6 | 9.4 |
| 9 | Alzheimer's disease (G30) | 675 | 2.0 | 7.4 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 651 | 2.0 | 7.1 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 515 | 1.5 | 5.6 |
| 12 | Parkinson's disease (G20-G21) | 448 | 1.3 | 4.9 |
| 13 | Septicemia (A40-A41) | 418 | 1.3 | 4.6 |
| 14 | Chronic liver disease and cirrhosis (K70,K73-K74) | 384 | 1.2 | 4.2 |
| 15 | Pneumonitis due to solids and liquids (J69) | 285 | 0.9 | 3.1 |
| ... | All other causes (residual) | 5,256 | 15.8 | 57.6 |
| Non-Hispanic, Asian or Pacific Islander, male, 1-4 years | | | | |
| ... | All causes | 72 | 100.0 | 15.3 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 16 | 22.2 | * |
| 2 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 9 | 12.5 | * |
| 3 | Malignant neoplasms (C00-C97) | 7 | 9.7 | * |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5 | 6.9 | * |
| 5 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 3 | 4.2 | * |
| 5 | Influenza and pneumonia (J09-J18) | 3 | 4.2 | * |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 3 | 4.2 | * |
| 8 | Acute bronchitis and bronchiolitis (J20-J21) | 1 | 1.4 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 1 | 1.4 | * |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 1.4 | * |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 1 | 1.4 | * |
| ... | All other causes (residual) | 22 | 30.6 | 4.7 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Non-Hispanic, Asian or Pacific Islander, male, 5-14 years | | | | |
| ... | All causes | 124 | 100.0 | 10.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 31 | 25.0 | 2.6 |
| 2 | Malignant neoplasms (C00-C97) | 25 | 20.2 | 2.1 |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 8 | 6.5 | * |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7 | 5.6 | * |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 4 | 3.2 | * |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 4 | 3.2 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 3 | 2.4 | * |
| 8 | Septicemia (A40-A41) | 2 | 1.6 | * |
| 9 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 1 | 0.8 | * |
| 9 | Meningitis (G00,G03) | 1 | 0.8 | * |
| 9 | Influenza and pneumonia (J09-J18) | 1 | 0.8 | * |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.8 | * |
| 9 | Complications of medical and surgical care (Y40-Y84,Y88) | 1 | 0.8 | * |
| ... | All other causes (residual) | 35 | 28.2 | 3.0 |
| Non-Hispanic, Asian or Pacific Islander, male, 15-24 years | | | | |
| ... | All causes | 630 | 100.0 | 48.2 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 208 | 33.0 | 15.9 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 203 | 32.2 | 15.5 |
| 3 | Malignant neoplasms (C00-C97) | 54 | 8.6 | 4.1 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 35 | 5.6 | 2.7 |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 31 | 4.9 | 2.4 |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 10 | 1.6 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 5 | 0.8 | * |
| 8 | Septicemia (A40-A41) | 3 | 0.5 | * |
| 8 | Aortic aneurysm and dissection (I71) | 3 | 0.5 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 3 | 0.5 | * |
| 8 | Complications of medical and surgical care (Y40-Y84,Y88) | 3 | 0.5 | * |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2 | 0.3 | * |
| 12 | Diabetes mellitus (E10-E14) | 2 | 0.3 | * |
| 12 | Influenza and pneumonia (J09-J18) | 2 | 0.3 | * |
| 12 | Legal intervention (Y35,Y89.0) | 2 | 0.3 | * |
| ... | All other causes (residual) | 64 | 10.2 | 4.9 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Non-Hispanic, Asian or Pacific Islander, male, 25-34 years | | | | |
| ... | All causes | 887 | 100.0 | 56.5 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 279 | 31.5 | 17.8 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 178 | 20.1 | 11.3 |
| 3 | Malignant neoplasms (C00-C97) | 85 | 9.6 | 5.4 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 81 | 9.1 | 5.2 |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 55 | 6.2 | 3.5 |
| 6 | Cerebrovascular diseases (I60-I69) | 17 | 1.9 | * |
| 7 | Diabetes mellitus (E10-E14) | 13 | 1.5 | * |
| 8 | Influenza and pneumonia (J09-J18) | 8 | 0.9 | * |
| 9 | Human immunodeficiency virus (HIV) disease (B20-B24) | 7 | 0.8 | * |
| 9 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 7 | 0.8 | * |
| 11 | Aortic aneurysm and dissection (I71) | 6 | 0.7 | * |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 6 | 0.7 | * |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 4 | 0.5 | * |
| 13 | Chronic lower respiratory diseases (J40-J47) | 4 | 0.5 | * |
| 13 | Legal intervention (Y35,Y89.0) | 4 | 0.5 | * |
| ... | All other causes (residual) | 133 | 15.0 | 8.5 |
| Non-Hispanic, Asian or Pacific Islander, male, 35-44 years | | | | |
| ... | All causes | 1,212 | 100.0 | 85.2 |
| 1 | Malignant neoplasms (C00-C97) | 265 | 21.9 | 18.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 244 | 20.1 | 17.2 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 166 | 13.7 | 11.7 |
| 4 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 116 | 9.6 | 8.2 |
| 5 | Cerebrovascular diseases (I60-I69) | 55 | 4.5 | 3.9 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 42 | 3.5 | 3.0 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 38 | 3.1 | 2.7 |
| 8 | Diabetes mellitus (E10-E14) | 33 | 2.7 | 2.3 |
| 9 | Influenza and pneumonia (J09-J18) | 19 | 1.6 | * |
| 10 | Human immunodeficiency virus (HIV) disease (B20-B24) | 17 | 1.4 | * |
| 11 | Septicemia (A40-A41) | 10 | 0.8 | * |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 10 | 0.8 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 10 | 0.8 | * |
| 11 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 10 | 0.8 | * |
| 15 | Aortic aneurysm and dissection (I71) | 8 | 0.7 | * |
| ... | All other causes (residual) | 169 | 13.9 | 11.9 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, Asian or Pacific Islander, male, 45-54 years | | | | |
| ... | All causes | 2,461 | 100.0 | 205.0 |
| 1 | Malignant neoplasms (C00-C97) | 748 | 30.4 | 62.3 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 594 | 24.1 | 49.5 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 193 | 7.8 | 16.1 |
| 4 | Cerebrovascular diseases (I60-I69) | 140 | 5.7 | 11.7 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 128 | 5.2 | 10.7 |
| 6 | Diabetes mellitus (E10-E14) | 87 | 3.5 | 7.2 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 76 | 3.1 | 6.3 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 37 | 1.5 | 3.1 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 33 | 1.3 | 2.7 |
| 10 | Influenza and pneumonia (J09-J18) | 31 | 1.3 | 2.6 |
| 11 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 30 | 1.2 | 2.5 |
| 12 | Viral hepatitis (B15-B19) | 23 | 0.9 | 1.9 |
| 13 | Human immunodeficiency virus (HIV) disease (B20-B24) | 22 | 0.9 | 1.8 |
| 14 | Chronic lower respiratory diseases (J40-J47) | 18 | 0.7 | * |
| 15 | Septicemia (A40-A41) | 16 | 0.7 | * |
| 15 | Aortic aneurysm and dissection (I71) | 16 | 0.7 | * |
| ... | All other causes (residual) | 269 | 10.9 | 22.4 |
| Non-Hispanic, Asian or Pacific Islander, male, 55-64 years | | | | |
| ... | All causes | 4,746 | 100.0 | 506.4 |
| 1 | Malignant neoplasms (C00-C97) | 1,709 | 36.0 | 182.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,129 | 23.8 | 120.5 |
| 3 | Cerebrovascular diseases (I60-I69) | 303 | 6.4 | 32.3 |
| 4 | Diabetes mellitus (E10-E14) | 215 | 4.5 | 22.9 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 207 | 4.4 | 22.1 |
| 6 | Chronic liver disease and cirrhosis (K70,K73-K74) | 137 | 2.9 | 14.6 |
| 7 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 104 | 2.2 | 11.1 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 77 | 1.6 | 8.2 |
| 9 | Chronic lower respiratory diseases (J40-J47) | 71 | 1.5 | 7.6 |
| 10 | Influenza and pneumonia (J09-J18) | 67 | 1.4 | 7.1 |
| 11 | Septicemia (A40-A41) | 62 | 1.3 | 6.6 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 61 | 1.3 | 6.5 |
| 13 | Viral hepatitis (B15-B19) | 38 | 0.8 | 4.1 |
| 14 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 29 | 0.6 | 3.1 |
| 15 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 27 | 0.6 | 2.9 |
| 15 | Aortic aneurysm and dissection (I71) | 27 | 0.6 | 2.9 |
| ... | All other causes (residual) | 483 | 10.2 | 51.5 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Non-Hispanic, Asian or Pacific Islander, male, 65-74 years | | | | |
| ... | All causes | 6,499 | 100.0 | 1,132.5 |
| 1 | Malignant neoplasms (C00-C97) | 2,379 | 36.6 | 414.6 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,455 | 22.4 | 253.6 |
| 3 | Cerebrovascular diseases (I60-I69) | 422 | 6.5 | 73.5 |
| 4 | Diabetes mellitus (E10-E14) | 340 | 5.2 | 59.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 215 | 3.3 | 37.5 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 190 | 2.9 | 33.1 |
| 7 | Influenza and pneumonia (J09-J18) | 146 | 2.2 | 25.4 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 122 | 1.9 | 21.3 |
| 9 | Septicemia (A40-A41) | 94 | 1.4 | 16.4 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 81 | 1.2 | 14.1 |
| 11 | Parkinson's disease (G20-G21) | 74 | 1.1 | 12.9 |
| 12 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 67 | 1.0 | 11.7 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 59 | 0.9 | 10.3 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 45 | 0.7 | 7.8 |
| 15 | Alzheimer's disease (G30) | 37 | 0.6 | 6.4 |
| 15 | Pneumonitis due to solids and liquids (J69) | 37 | 0.6 | 6.4 |
| ... | All other causes (residual) | 736 | 11.3 | 128.3 |
| Non-Hispanic, Asian or Pacific Islander, male, 75-84 years | | | | |
| ... | All causes | 8,105 | 100.0 | 3,096.7 |
| 1 | Malignant neoplasms (C00-C97) | 2,218 | 27.4 | 847.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,876 | 23.1 | 716.8 |
| 3 | Cerebrovascular diseases (I60-I69) | 634 | 7.8 | 242.2 |
| 4 | Diabetes mellitus (E10-E14) | 368 | 4.5 | 140.6 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 343 | 4.2 | 131.1 |
| 6 | Influenza and pneumonia (J09-J18) | 327 | 4.0 | 124.9 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 233 | 2.9 | 89.0 |
| 8 | Alzheimer's disease (G30) | 211 | 2.6 | 80.6 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 199 | 2.5 | 76.0 |
| 10 | Parkinson's disease (G20-G21) | 197 | 2.4 | 75.3 |
| 11 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 129 | 1.6 | 49.3 |
| 12 | Septicemia (A40-A41) | 113 | 1.4 | 43.2 |
| 13 | Pneumonitis due to solids and liquids (J69) | 73 | 0.9 | 27.9 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 55 | 0.7 | 21.0 |
| 15 | Aortic aneurysm and dissection (I71) | 45 | 0.6 | 17.2 |
| ... | All other causes (residual) | 1,084 | 13.4 | 414.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Non-Hispanic, Asian or Pacific Islander, male, 85 years and over | | | | |
| ... | All causes | 8,069 | 100.0 | 9,459.6 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 2,181 | 27.0 | 2,556.9 |
| 2 | Malignant neoplasms (C00-C97) | 1,282 | 15.9 | 1,502.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 571 | 7.1 | 669.4 |
| 4 | Influenza and pneumonia (J09-J18) | 501 | 6.2 | 587.3 |
| 5 | Chronic lower respiratory diseases (J40-J47) | 424 | 5.3 | 497.1 |
| 6 | Alzheimer's disease (G30) | 416 | 5.2 | 487.7 |
| 7 | Diabetes mellitus (E10-E14) | 264 | 3.3 | 309.5 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 225 | 2.8 | 263.8 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 201 | 2.5 | 235.6 |
| 10 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 198 | 2.5 | 232.1 |
| 11 | Parkinson's disease (G20-G21) | 168 | 2.1 | 197.0 |
| 12 | Pneumonitis due to solids and liquids (J69) | 151 | 1.9 | 177.0 |
| 13 | Septicemia (A40-A41) | 112 | 1.4 | 131.3 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 48 | 0.6 | 56.3 |
| 15 | Nutritional deficiencies (E40-E64) | 31 | 0.4 | 36.3 |
| ... | All other causes (residual) | 1,296 | 16.1 | 1,519.3 |
| Non-Hispanic, Asian or Pacific Islander, female, all ages | | | | |
| ... | All causes | 31,971 | 100.0 | 320.1 |
| 1 | Malignant neoplasms (C00-C97) | 8,532 | 26.7 | 85.4 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 6,362 | 19.9 | 63.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 2,645 | 8.3 | 26.5 |
| 4 | Alzheimer's disease (G30) | 1,505 | 4.7 | 15.1 |
| 5 | Diabetes mellitus (E10-E14) | 1,277 | 4.0 | 12.8 |
| 6 | Influenza and pneumonia (J09-J18) | 1,052 | 3.3 | 10.5 |
| 7 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 1,051 | 3.3 | 10.5 |
| 8 | Chronic lower respiratory diseases (J40-J47) | 813 | 2.5 | 8.1 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 667 | 2.1 | 6.7 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 621 | 1.9 | 6.2 |
| 11 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 418 | 1.3 | 4.2 |
| 12 | Septicemia (A40-A41) | 380 | 1.2 | 3.8 |
| 13 | Parkinson's disease (G20-G21) | 299 | 0.9 | 3.0 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 222 | 0.7 | 2.2 |
| 15 | Pneumonitis due to solids and liquids (J69) | 215 | 0.7 | 2.2 |
| ... | All other causes (residual) | 5,912 | 18.5 | 59.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|---|---|--------|-------------------------|------|
| Non-Hispanic, Asian or Pacific Islander, female, 1-4 years | | | | |
| ... | All causes | 69 | 100.0 | 15.4 |
| 1 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 11 | 15.9 | * |
| 2 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 10 | 14.5 | * |
| 3 | Malignant neoplasms (C00-C97) | 6 | 8.7 | * |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 5 | 7.2 | * |
| 5 | Septicemia (A40-A41) | 3 | 4.3 | * |
| 5 | Influenza and pneumonia (J09-J18) | 3 | 4.3 | * |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 3 | 4.3 | * |
| 8 | Tuberculosis (A16-A19) | 1 | 1.4 | * |
| 8 | Anemias (D50-D64) | 1 | 1.4 | * |
| 8 | Certain conditions originating in the perinatal period (P00-P96) | 1 | 1.4 | * |
| ... | All other causes (residual) | 25 | 36.2 | 5.6 |
| Non-Hispanic, Asian or Pacific Islander, female, 5-14 years | | | | |
| ... | All causes | 88 | 100.0 | 7.6 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 18 | 20.5 | * |
| 2 | Malignant neoplasms (C00-C97) | 15 | 17.0 | * |
| 3 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 7 | 8.0 | * |
| 3 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 7 | 8.0 | * |
| 5 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 3 | 3.4 | * |
| 5 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 3 | 3.4 | * |
| 7 | Septicemia (A40-A41) | 2 | 2.3 | * |
| 7 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 2 | 2.3 | * |
| 7 | Anemias (D50-D64) | 2 | 2.3 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 2 | 2.3 | * |
| 11 | Influenza and pneumonia (J09-J18) | 1 | 1.1 | * |
| 11 | Complications of medical and surgical care (Y40-Y84,Y88) | 1 | 1.1 | * |
| ... | All other causes (residual) | 25 | 28.4 | 2.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|------|
| Non-Hispanic, Asian or Pacific Islander, female, 15-24 years | | | | |
| ... | All causes | 243 | 100.0 | 19.1 |
| 1 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 69 | 28.4 | 5.4 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 62 | 25.5 | 4.9 |
| 3 | Malignant neoplasms (C00-C97) | 30 | 12.3 | 2.4 |
| 4 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 11 | 4.5 | * |
| 5 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 9 | 3.7 | * |
| 6 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 6 | 2.5 | * |
| 7 | Influenza and pneumonia (J09-J18) | 3 | 1.2 | * |
| 8 | Cerebrovascular diseases (I60-I69) | 2 | 0.8 | * |
| 8 | Chronic lower respiratory diseases (J40-J47) | 2 | 0.8 | * |
| 8 | Pneumonitis due to solids and liquids (J69) | 2 | 0.8 | * |
| 11 | Tuberculosis (A16-A19) | 1 | 0.4 | * |
| 11 | Anemias (D50-D64) | 1 | 0.4 | * |
| 11 | Diabetes mellitus (E10-E14) | 1 | 0.4 | * |
| 11 | Meningitis (G00,G03) | 1 | 0.4 | * |
| 11 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 1 | 0.4 | * |
| 11 | Pregnancy, childbirth and the puerperium (O00-O99) | 1 | 0.4 | * |
| 11 | Certain conditions originating in the perinatal period (P00-P96) | 1 | 0.4 | * |
| 11 | Legal intervention (Y35,Y89.0) | 1 | 0.4 | * |
| 11 | Complications of medical and surgical care (Y40-Y84,Y88) | 1 | 0.4 | * |
| 11 | Enterocolitis due to <i>Clostridium difficile</i> (A04.7) | 1 | 0.4 | * |
| ... | All other causes (residual) | 37 | 15.2 | 2.9 |
| Non-Hispanic, Asian or Pacific Islander, female, 25-34 years | | | | |
| ... | All causes | 445 | 100.0 | 26.3 |
| 1 | Malignant neoplasms (C00-C97) | 94 | 21.1 | 5.6 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 88 | 19.8 | 5.2 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 79 | 17.8 | 4.7 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 23 | 5.2 | 1.4 |
| 5 | Pregnancy, childbirth and the puerperium (O00-O99) | 21 | 4.7 | 1.2 |
| 6 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 13 | 2.9 | * |
| 7 | Cerebrovascular diseases (I60-I69) | 12 | 2.7 | * |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 7 | 1.6 | * |
| 9 | Septicemia (A40-A41) | 6 | 1.3 | * |
| 10 | Chronic lower respiratory diseases (J40-J47) | 5 | 1.1 | * |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 5 | 1.1 | * |
| 12 | Diabetes mellitus (E10-E14) | 4 | 0.9 | * |
| 12 | Influenza and pneumonia (J09-J18) | 4 | 0.9 | * |
| 12 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 4 | 0.9 | * |
| 15 | Complications of medical and surgical care (Y40-Y84,Y88) | 2 | 0.4 | * |
| ... | All other causes (residual) | 78 | 17.5 | 4.6 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, Asian or Pacific Islander, female, 35-44 years | | | | |
| ... | All causes | 835 | 100.0 | 51.6 |
| 1 | Malignant neoplasms (C00-C97) | 362 | 43.4 | 22.4 |
| 2 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 64 | 7.7 | 4.0 |
| 3 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 63 | 7.5 | 3.9 |
| 4 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 62 | 7.4 | 3.8 |
| 5 | Cerebrovascular diseases (I60-I69) | 59 | 7.1 | 3.6 |
| 6 | Pregnancy, childbirth and the puerperium (O00-O99) | 21 | 2.5 | 1.3 |
| 7 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 18 | 2.2 | * |
| 8 | Chronic liver disease and cirrhosis (K70,K73-K74) | 14 | 1.7 | * |
| 9 | Diabetes mellitus (E10-E14) | 13 | 1.6 | * |
| 10 | Septicemia (A40-A41) | 10 | 1.2 | * |
| 11 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 7 | 0.8 | * |
| 11 | Chronic lower respiratory diseases (J40-J47) | 7 | 0.8 | * |
| 13 | Anemias (D50-D64) | 4 | 0.5 | * |
| 13 | Influenza and pneumonia (J09-J18) | 4 | 0.5 | * |
| 13 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 4 | 0.5 | * |
| 13 | Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99) | 4 | 0.5 | * |
| ... | All other causes (residual) | 119 | 14.3 | 7.4 |
| Non-Hispanic, Asian or Pacific Islander, female, 45-54 years | | | | |
| ... | All causes | 1,751 | 100.0 | 128.1 |
| 1 | Malignant neoplasms (C00-C97) | 930 | 53.1 | 68.0 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 165 | 9.4 | 12.1 |
| 3 | Cerebrovascular diseases (I60-I69) | 104 | 5.9 | 7.6 |
| 4 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 84 | 4.8 | 6.1 |
| 5 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 76 | 4.3 | 5.6 |
| 6 | Diabetes mellitus (E10-E14) | 54 | 3.1 | 3.9 |
| 7 | Chronic liver disease and cirrhosis (K70,K73-K74) | 25 | 1.4 | 1.8 |
| 8 | Septicemia (A40-A41) | 19 | 1.1 | * |
| 9 | Chronic lower respiratory diseases (J40-J47) | 16 | 0.9 | * |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 16 | 0.9 | * |
| 11 | Influenza and pneumonia (J09-J18) | 14 | 0.8 | * |
| 12 | Viral hepatitis (B15-B19) | 11 | 0.6 | * |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 11 | 0.6 | * |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 11 | 0.6 | * |
| 15 | Assault (homicide) (*U01-*U02,X85-Y09,Y87.1) | 9 | 0.5 | * |
| ... | All other causes (residual) | 206 | 11.8 | 15.1 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|-------|
| Non-Hispanic, Asian or Pacific Islander, female, 55-64 years | | | | |
| ... | All causes | 3,263 | 100.0 | 290.3 |
| 1 | Malignant neoplasms (C00-C97) | 1,675 | 51.3 | 149.0 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 453 | 13.9 | 40.3 |
| 3 | Cerebrovascular diseases (I60-I69) | 211 | 6.5 | 18.8 |
| 4 | Diabetes mellitus (E10-E14) | 129 | 4.0 | 11.5 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 93 | 2.9 | 8.3 |
| 6 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 57 | 1.7 | 5.1 |
| 6 | Intentional self-harm (suicide) (*U03,X60-X84,Y87.0) | 57 | 1.7 | 5.1 |
| 8 | Septicemia (A40-A41) | 42 | 1.3 | 3.7 |
| 9 | Influenza and pneumonia (J09-J18) | 37 | 1.1 | 3.3 |
| 10 | Chronic lower respiratory diseases (J40-J47) | 34 | 1.0 | 3.0 |
| 11 | Chronic liver disease and cirrhosis (K70,K73-K74) | 26 | 0.8 | 2.3 |
| 12 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 25 | 0.8 | 2.2 |
| 12 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 25 | 0.8 | 2.2 |
| 14 | Viral hepatitis (B15-B19) | 19 | 0.6 | * |
| 14 | Aortic aneurysm and dissection (I71) | 19 | 0.6 | * |
| ... | All other causes (residual) | 361 | 11.1 | 32.1 |
| Non-Hispanic, Asian or Pacific Islander, female, 65-74 years | | | | |
| ... | All causes | 4,997 | 100.0 | 703.6 |
| 1 | Malignant neoplasms (C00-C97) | 1,990 | 39.8 | 280.2 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 859 | 17.2 | 120.9 |
| 3 | Cerebrovascular diseases (I60-I69) | 365 | 7.3 | 51.4 |
| 4 | Diabetes mellitus (E10-E14) | 264 | 5.3 | 37.2 |
| 5 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 140 | 2.8 | 19.7 |
| 6 | Influenza and pneumonia (J09-J18) | 115 | 2.3 | 16.2 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 110 | 2.2 | 15.5 |
| 8 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 102 | 2.0 | 14.4 |
| 9 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 75 | 1.5 | 10.6 |
| 10 | Septicemia (A40-A41) | 61 | 1.2 | 8.6 |
| 10 | Chronic liver disease and cirrhosis (K70,K73-K74) | 61 | 1.2 | 8.6 |
| 12 | Alzheimer's disease (G30) | 49 | 1.0 | 6.9 |
| 13 | Parkinson's disease (G20-G21) | 39 | 0.8 | 5.5 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 38 | 0.8 | 5.4 |
| 15 | Aortic aneurysm and dissection (I71) | 30 | 0.6 | 4.2 |
| ... | All other causes (residual) | 699 | 14.0 | 98.4 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.

LCWK2. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in 10-year age groups, by race and Hispanic origin, and sex: United States, 2015

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

| Rank ¹ | Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age | Number | Percent of total deaths | Rate |
|--|---|--------|-------------------------|---------|
| Non-Hispanic, Asian or Pacific Islander, female, 75-84 years | | | | |
| ... | All causes | 7,810 | 100.0 | 2,274.6 |
| 1 | Malignant neoplasms (C00-C97) | 2,029 | 26.0 | 590.9 |
| 2 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 1,623 | 20.8 | 472.7 |
| 3 | Cerebrovascular diseases (I60-I69) | 759 | 9.7 | 221.0 |
| 4 | Diabetes mellitus (E10-E14) | 420 | 5.4 | 122.3 |
| 5 | Alzheimer's disease (G30) | 297 | 3.8 | 86.5 |
| 6 | Chronic lower respiratory diseases (J40-J47) | 271 | 3.5 | 78.9 |
| 7 | Influenza and pneumonia (J09-J18) | 258 | 3.3 | 75.1 |
| 8 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 217 | 2.8 | 63.2 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 194 | 2.5 | 56.5 |
| 10 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 170 | 2.2 | 49.5 |
| 11 | Septicemia (A40-A41) | 97 | 1.2 | 28.2 |
| 12 | Parkinson's disease (G20-G21) | 95 | 1.2 | 27.7 |
| 13 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 58 | 0.7 | 16.9 |
| 13 | Pneumonitis due to solids and liquids (J69) | 58 | 0.7 | 16.9 |
| 13 | Chronic liver disease and cirrhosis (K70,K73-K74) | 58 | 0.7 | 16.9 |
| ... | All other causes (residual) | 1,206 | 15.4 | 351.2 |
| Non-Hispanic, Asian or Pacific Islander, female, 85 years and over | | | | |
| ... | All causes | 12,078 | 100.0 | 8,187.2 |
| 1 | Diseases of heart (I00-I09,I11,I13,I20-I51) | 3,151 | 26.1 | 2,135.9 |
| 2 | Malignant neoplasms (C00-C97) | 1,398 | 11.6 | 947.6 |
| 3 | Alzheimer's disease (G30) | 1,149 | 9.5 | 778.9 |
| 4 | Cerebrovascular diseases (I60-I69) | 1,129 | 9.3 | 765.3 |
| 5 | Influenza and pneumonia (J09-J18) | 612 | 5.1 | 414.9 |
| 6 | Diabetes mellitus (E10-E14) | 392 | 3.2 | 265.7 |
| 7 | Chronic lower respiratory diseases (J40-J47) | 368 | 3.0 | 249.5 |
| 8 | Essential hypertension and hypertensive renal disease (I10,I12,I15) | 337 | 2.8 | 228.4 |
| 9 | Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27) | 286 | 2.4 | 193.9 |
| 10 | Accidents (unintentional injuries) (V01-X59,Y85-Y86) | 270 | 2.2 | 183.0 |
| 11 | Parkinson's disease (G20-G21) | 159 | 1.3 | 107.8 |
| 12 | Septicemia (A40-A41) | 137 | 1.1 | 92.9 |
| 13 | Pneumonitis due to solids and liquids (J69) | 125 | 1.0 | 84.7 |
| 14 | In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48) | 78 | 0.6 | 52.9 |
| 15 | Aortic aneurysm and dissection (I71) | 57 | 0.5 | 38.6 |
| ... | All other causes (residual) | 2,430 | 20.1 | 1,647.2 |

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2015.