

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, all ages				
...	All causes	2,813,503	100.0	863.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	647,457	23.0	198.8
2	Malignant neoplasms (C00-C97)	599,108	21.3	183.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	169,936	6.0	52.2
4	Chronic lower respiratory diseases (J40-J47)	160,201	5.7	49.2
5	Cerebrovascular diseases (I60-I69)	146,383	5.2	44.9
6	Alzheimer disease (G30)	121,404	4.3	37.3
7	Diabetes mellitus (E10-E14)	83,564	3.0	25.7
8	Influenza and pneumonia (J09-J18)	55,672	2.0	17.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50,633	1.8	15.5
10	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	47,173	1.7	14.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	41,743	1.5	12.8
12	Septicemia (A40-A41)	40,922	1.5	12.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	35,316	1.3	10.8
14	Parkinson disease (G20-G21)	31,963	1.1	9.8
15	Pneumonitis due to solids and liquids (J69)	20,108	0.7	6.2
...	All other causes (residual)	561,920	20.0	172.5
All races and origins, both sexes, 1-4 years				
...	All causes	3,880	100.0	24.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,267	32.7	7.9
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	424	10.9	2.7
3	Malignant neoplasms (C00-C97)	325	8.4	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	303	7.8	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	127	3.3	0.8
6	Influenza and pneumonia (J09-J18)	104	2.7	0.7
7	Cerebrovascular diseases (I60-I69)	66	1.7	0.4
8	Septicemia (A40-A41)	48	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	44	1.1	0.3
10	Certain conditions originating in the perinatal period (P00-P96)	42	1.1	0.3
11	Chronic lower respiratory diseases (J40-J47)	31	0.8	0.2
12	Anemias (D50-D64)	24	0.6	0.2
12	Complications of medical and surgical care (Y40-Y84,Y88)	24	0.6	0.2
14	Acute bronchitis and bronchiolitis (J20-J21)	22	0.6	0.1
15	Meningitis (G00,G03)	19	0.5	*
...	All other causes (residual)	1,010	26.0	6.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 5-14 years				
...	All causes	5,571	100.0	13.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,578	28.3	3.8
2	Malignant neoplasms (C00-C97)	855	15.3	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	522	9.4	1.3
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	379	6.8	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	332	6.0	0.8
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	179	3.2	0.4
7	Chronic lower respiratory diseases (J40-J47)	134	2.4	0.3
8	Influenza and pneumonia (J09-J18)	113	2.0	0.3
9	Cerebrovascular diseases (I60-I69)	97	1.7	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	62	1.1	0.2
11	Septicemia (A40-A41)	57	1.0	0.1
12	Diabetes mellitus (E10-E14)	34	0.6	0.1
13	Complications of medical and surgical care (Y40-Y84,Y88)	25	0.4	0.1
14	Anemias (D50-D64)	24	0.4	0.1
15	Certain conditions originating in the perinatal period (P00-P96)	21	0.4	0.1
...	All other causes (residual)	1,159	20.8	2.8
All races and origins, both sexes, 15-24 years				
...	All causes	32,025	100.0	74.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,441	42.0	31.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,252	19.5	14.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,905	15.3	11.3
4	Malignant neoplasms (C00-C97)	1,374	4.3	3.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	913	2.9	2.1
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	355	1.1	0.8
7	Diabetes mellitus (E10-E14)	248	0.8	0.6
8	Influenza and pneumonia (J09-J18)	190	0.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	188	0.6	0.4
10	Pregnancy, childbirth and the puerperium (O00-O99)	168	0.5	0.4
11	Cerebrovascular diseases (I60-I69)	155	0.5	0.4
12	Septicemia (A40-A41)	123	0.4	0.3
13	Legal intervention (Y35,Y89.0)	97	0.3	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	80	0.2	0.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	69	0.2	0.2
15	Anemias (D50-D64)	69	0.2	0.2
...	All other causes (residual)	3,398	10.6	7.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 25-44 years				
...	All causes	140,011	100.0	162.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	48,497	34.6	56.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	15,283	10.9	17.7
3	Malignant neoplasms (C00-C97)	14,516	10.4	16.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,082	10.1	16.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,839	6.3	10.3
6	Chronic liver disease and cirrhosis (K70,K73-K74)	3,918	2.8	4.5
7	Diabetes mellitus (E10-E14)	2,941	2.1	3.4
8	Cerebrovascular diseases (I60-I69)	2,404	1.7	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1,344	1.0	1.6
10	Septicemia (A40-A41)	1,246	0.9	1.4
11	Influenza and pneumonia (J09-J18)	1,187	0.8	1.4
12	Chronic lower respiratory diseases (J40-J47)	1,045	0.7	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	947	0.7	1.1
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	856	0.6	1.0
15	Pregnancy, childbirth and the puerperium (O00-O99)	840	0.6	1.0
...	All other causes (residual)	22,066	15.8	25.6
All races and origins, both sexes, 45-64 years				
...	All causes	542,148	100.0	642.6
1	Malignant neoplasms (C00-C97)	154,076	28.4	182.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	112,760	20.8	133.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,869	8.8	56.7
4	Chronic lower respiratory diseases (J40-J47)	22,642	4.2	26.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	22,049	4.1	26.1
6	Diabetes mellitus (E10-E14)	21,313	3.9	25.3
7	Cerebrovascular diseases (I60-I69)	17,906	3.3	21.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	16,543	3.1	19.6
9	Septicemia (A40-A41)	8,279	1.5	9.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7,868	1.5	9.3
11	Influenza and pneumonia (J09-J18)	7,058	1.3	8.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	5,633	1.0	6.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,827	0.7	4.5
14	Viral hepatitis (B15-B19)	3,444	0.6	4.1
15	Human immunodeficiency virus (HIV) disease (B20-B24)	3,346	0.6	4.0
...	All other causes (residual)	87,535	16.1	103.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, both sexes, 65 years and over				
...	All causes	2,067,404	100.0	4,065.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	519,052	25.1	1,020.6
2	Malignant neoplasms (C00-C97)	427,896	20.7	841.3
3	Chronic lower respiratory diseases (J40-J47)	136,139	6.6	267.7
4	Cerebrovascular diseases (I60-I69)	125,653	6.1	247.1
5	Alzheimer disease (G30)	120,107	5.8	236.2
6	Diabetes mellitus (E10-E14)	59,020	2.9	116.0
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	55,951	2.7	110.0
8	Influenza and pneumonia (J09-J18)	46,862	2.3	92.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	41,670	2.0	81.9
10	Parkinson disease (G20-G21)	31,177	1.5	61.3
11	Septicemia (A40-A41)	31,022	1.5	61.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	29,036	1.4	57.1
13	Pneumonitis due to solids and liquids (J69)	17,589	0.9	34.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	15,746	0.8	31.0
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13,155	0.6	25.9
...	All other causes (residual)	397,329	19.2	781.2
All races and origins, male, all ages				
...	All causes	1,439,111	100.0	897.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	347,879	24.2	216.9
2	Malignant neoplasms (C00-C97)	315,147	21.9	196.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	109,722	7.6	68.4
4	Chronic lower respiratory diseases (J40-J47)	75,005	5.2	46.8
5	Cerebrovascular diseases (I60-I69)	61,645	4.3	38.4
6	Diabetes mellitus (E10-E14)	46,302	3.2	28.9
7	Alzheimer disease (G30)	37,325	2.6	23.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	36,782	2.6	22.9
9	Influenza and pneumonia (J09-J18)	26,558	1.8	16.6
10	Chronic liver disease and cirrhosis (K70,K73-K74)	26,451	1.8	16.5
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	25,744	1.8	16.0
12	Septicemia (A40-A41)	19,603	1.4	12.2
13	Parkinson disease (G20-G21)	19,388	1.3	12.1
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	15,749	1.1	9.8
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	15,524	1.1	9.7
...	All other causes (residual)	260,287	18.1	162.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 1-4 years				
...	All causes	2,232	100.0	27.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	770	34.5	9.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	225	10.1	2.8
3	Malignant neoplasms (C00-C97)	184	8.2	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	168	7.5	2.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	67	3.0	0.8
6	Influenza and pneumonia (J09-J18)	63	2.8	0.8
7	Cerebrovascular diseases (I60-I69)	42	1.9	0.5
8	Septicemia (A40-A41)	26	1.2	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	19	0.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	19	0.9	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	17	0.8	*
12	Chronic lower respiratory diseases (J40-J47)	16	0.7	*
13	Acute bronchitis and bronchiolitis (J20-J21)	15	0.7	*
14	Meningitis (G00,G03)	14	0.6	*
15	Anemias (D50-D64)	8	0.4	*
...	All other causes (residual)	579	25.9	7.1
All races and origins, male, 5-14 years				
...	All causes	3,269	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	985	30.1	4.7
2	Malignant neoplasms (C00-C97)	455	13.9	2.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	352	10.8	1.7
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	6.4	1.0
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	196	6.0	0.9
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	96	2.9	0.5
7	Chronic lower respiratory diseases (J40-J47)	84	2.6	0.4
8	Cerebrovascular diseases (I60-I69)	53	1.6	0.3
9	Influenza and pneumonia (J09-J18)	49	1.5	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	1.1	0.2
11	Septicemia (A40-A41)	31	0.9	0.1
12	Diabetes mellitus (E10-E14)	20	0.6	0.1
13	Anemias (D50-D64)	15	0.5	*
14	Certain conditions originating in the perinatal period (P00-P96)	12	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.4	*
...	All other causes (residual)	666	20.4	3.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 15-24 years				
...	All causes	23,503	100.0	106.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,746	41.5	44.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5,027	21.4	22.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,234	18.0	19.1
4	Malignant neoplasms (C00-C97)	833	3.5	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	620	2.6	2.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	208	0.9	0.9
7	Diabetes mellitus (E10-E14)	144	0.6	0.7
8	Chronic lower respiratory diseases (J40-J47)	115	0.5	0.5
9	Influenza and pneumonia (J09-J18)	109	0.5	0.5
10	Legal intervention (Y35,Y89.0)	90	0.4	0.4
11	Cerebrovascular diseases (I60-I69)	83	0.4	0.4
12	Septicemia (A40-A41)	62	0.3	0.3
13	Human immunodeficiency virus (HIV) disease (B20-B24)	52	0.2	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	45	0.2	0.2
15	Pneumonitis due to solids and liquids (J69)	30	0.1	0.1
...	All other causes (residual)	2,105	9.0	9.5
All races and origins, male, 25-44 years				
...	All causes	92,941	100.0	214.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	35,244	37.9	81.3
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,944	12.9	27.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	9,523	10.2	22.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7,283	7.8	16.8
5	Malignant neoplasms (C00-C97)	6,415	6.9	14.8
6	Chronic liver disease and cirrhosis (K70,K73-K74)	2,404	2.6	5.5
7	Diabetes mellitus (E10-E14)	1,769	1.9	4.1
8	Cerebrovascular diseases (I60-I69)	1,373	1.5	3.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	936	1.0	2.2
10	Septicemia (A40-A41)	656	0.7	1.5
11	Influenza and pneumonia (J09-J18)	652	0.7	1.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	535	0.6	1.2
13	Chronic lower respiratory diseases (J40-J47)	515	0.6	1.2
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	505	0.5	1.2
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	440	0.5	1.0
...	All other causes (residual)	12,747	13.7	29.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, male, 45-64 years				
...	All causes	329,139	100.0	799.6
1	Malignant neoplasms (C00-C97)	81,278	24.7	197.4
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,143	23.7	189.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	33,137	10.1	80.5
4	Chronic liver disease and cirrhosis (K70,K73-K74)	14,612	4.4	35.5
5	Diabetes mellitus (E10-E14)	13,302	4.0	32.3
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12,371	3.8	30.1
7	Chronic lower respiratory diseases (J40-J47)	11,256	3.4	27.3
8	Cerebrovascular diseases (I60-I69)	10,342	3.1	25.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,487	1.4	10.9
10	Septicemia (A40-A41)	4,483	1.4	10.9
11	Influenza and pneumonia (J09-J18)	3,902	1.2	9.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,534	1.1	8.6
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,880	0.9	7.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	2,481	0.8	6.0
15	Viral hepatitis (B15-B19)	2,363	0.7	5.7
...	All other causes (residual)	50,568	15.4	122.8
All races and origins, male, 65 years and over				
...	All causes	975,468	100.0	4,323.0
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	259,248	26.6	1,148.9
2	Malignant neoplasms (C00-C97)	225,950	23.2	1,001.3
3	Chronic lower respiratory diseases (J40-J47)	63,004	6.5	279.2
4	Cerebrovascular diseases (I60-I69)	49,693	5.1	220.2
5	Alzheimer disease (G30)	36,757	3.8	162.9
6	Diabetes mellitus (E10-E14)	31,064	3.2	137.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	29,065	3.0	128.8
8	Influenza and pneumonia (J09-J18)	21,684	2.2	96.1
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	20,644	2.1	91.5
10	Parkinson disease (G20-G21)	18,872	1.9	83.6
11	Septicemia (A40-A41)	14,266	1.5	63.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	11,762	1.2	52.1
13	Pneumonitis due to solids and liquids (J69)	9,817	1.0	43.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	9,416	1.0	41.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7,085	0.7	31.4
...	All other causes (residual)	167,141	17.1	740.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, all ages				
...	All causes	1,374,392	100.0	831.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	299,578	21.8	181.2
2	Malignant neoplasms (C00-C97)	283,961	20.7	171.8
3	Chronic lower respiratory diseases (J40-J47)	85,196	6.2	51.5
4	Cerebrovascular diseases (I60-I69)	84,738	6.2	51.3
5	Alzheimer disease (G30)	84,079	6.1	50.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	60,214	4.4	36.4
7	Diabetes mellitus (E10-E14)	37,262	2.7	22.5
8	Influenza and pneumonia (J09-J18)	29,114	2.1	17.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	24,889	1.8	15.1
10	Septicemia (A40-A41)	21,319	1.6	12.9
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	19,567	1.4	11.8
12	Chronic liver disease and cirrhosis (K70,K73-K74)	15,292	1.1	9.3
13	Parkinson disease (G20-G21)	12,575	0.9	7.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,391	0.8	6.3
15	Pneumonitis due to solids and liquids (J69)	8,758	0.6	5.3
...	All other causes (residual)	297,459	21.6	179.9
All races and origins, female, 1-4 years				
...	All causes	1,648	100.0	21.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	497	30.2	6.4
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	199	12.1	2.5
3	Malignant neoplasms (C00-C97)	141	8.6	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	135	8.2	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	60	3.6	0.8
6	Influenza and pneumonia (J09-J18)	41	2.5	0.5
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	25	1.5	0.3
8	Cerebrovascular diseases (I60-I69)	24	1.5	0.3
9	Certain conditions originating in the perinatal period (P00-P96)	23	1.4	0.3
10	Septicemia (A40-A41)	22	1.3	0.3
11	Anemias (D50-D64)	16	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	15	0.9	*
13	Acute bronchitis and bronchiolitis (J20-J21)	7	0.4	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.4	*
15	Meningitis (G00,G03)	5	0.3	*
...	All other causes (residual)	431	26.2	5.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 5-14 years				
...	All causes	2,302	100.0	11.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	593	25.8	2.9
2	Malignant neoplasms (C00-C97)	400	17.4	2.0
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	171	7.4	0.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	170	7.4	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	136	5.9	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	83	3.6	0.4
7	Influenza and pneumonia (J09-J18)	64	2.8	0.3
8	Chronic lower respiratory diseases (J40-J47)	50	2.2	0.2
9	Cerebrovascular diseases (I60-I69)	44	1.9	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	27	1.2	0.1
11	Septicemia (A40-A41)	26	1.1	0.1
12	Diabetes mellitus (E10-E14)	14	0.6	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.6	*
14	Anemias (D50-D64)	9	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	9	0.4	*
...	All other causes (residual)	493	21.4	2.5
All races and origins, female, 15-24 years				
...	All causes	8,522	100.0	40.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,695	43.4	17.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,225	14.4	5.8
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	671	7.9	3.2
4	Malignant neoplasms (C00-C97)	541	6.3	2.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	293	3.4	1.4
6	Pregnancy, childbirth and the puerperium (O00-O99)	168	2.0	0.8
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	147	1.7	0.7
8	Diabetes mellitus (E10-E14)	104	1.2	0.5
9	Influenza and pneumonia (J09-J18)	81	1.0	0.4
10	Chronic lower respiratory diseases (J40-J47)	73	0.9	0.3
11	Cerebrovascular diseases (I60-I69)	72	0.8	0.3
12	Septicemia (A40-A41)	61	0.7	0.3
13	Anemias (D50-D64)	40	0.5	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.4	0.2
15	Complications of medical and surgical care (Y40-Y84,Y88)	26	0.3	0.1
...	All other causes (residual)	1,290	15.1	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 25-44 years				
...	All causes	47,070	100.0	109.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,253	28.2	30.9
2	Malignant neoplasms (C00-C97)	8,101	17.2	18.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	4,559	9.7	10.6
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,339	7.1	7.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,556	3.3	3.6
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,514	3.2	3.5
7	Diabetes mellitus (E10-E14)	1,172	2.5	2.7
8	Cerebrovascular diseases (I60-I69)	1,031	2.2	2.4
9	Pregnancy, childbirth and the puerperium (O00-O99)	840	1.8	2.0
10	Septicemia (A40-A41)	590	1.3	1.4
11	Influenza and pneumonia (J09-J18)	535	1.1	1.2
12	Chronic lower respiratory diseases (J40-J47)	530	1.1	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	412	0.9	1.0
14	Human immunodeficiency virus (HIV) disease (B20-B24)	408	0.9	1.0
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	351	0.7	0.8
...	All other causes (residual)	8,879	18.9	20.7
All races and origins, female, 45-64 years				
...	All causes	213,009	100.0	493.0
1	Malignant neoplasms (C00-C97)	72,798	34.2	168.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	34,617	16.3	80.1
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14,732	6.9	34.1
4	Chronic lower respiratory diseases (J40-J47)	11,386	5.3	26.4
5	Diabetes mellitus (E10-E14)	8,011	3.8	18.5
6	Cerebrovascular diseases (I60-I69)	7,564	3.6	17.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	7,437	3.5	17.2
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,172	2.0	9.7
9	Septicemia (A40-A41)	3,796	1.8	8.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,381	1.6	7.8
11	Influenza and pneumonia (J09-J18)	3,156	1.5	7.3
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,099	1.0	4.9
13	Viral hepatitis (B15-B19)	1,081	0.5	2.5
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	947	0.4	2.2
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	935	0.4	2.2
...	All other causes (residual)	36,897	17.3	85.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
All races and origins, female, 65 years and over				
...	All causes	1,091,936	100.0	3,859.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	259,804	23.8	918.2
2	Malignant neoplasms (C00-C97)	201,946	18.5	713.7
3	Alzheimer disease (G30)	83,350	7.6	294.6
4	Cerebrovascular diseases (I60-I69)	75,960	7.0	268.5
5	Chronic lower respiratory diseases (J40-J47)	73,135	6.7	258.5
6	Diabetes mellitus (E10-E14)	27,956	2.6	98.8
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26,886	2.5	95.0
8	Influenza and pneumonia (J09-J18)	25,178	2.3	89.0
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	21,026	1.9	74.3
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	17,274	1.6	61.1
11	Septicemia (A40-A41)	16,756	1.5	59.2
12	Parkinson disease (G20-G21)	12,305	1.1	43.5
13	Pneumonitis due to solids and liquids (J69)	7,772	0.7	27.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	6,330	0.6	22.4
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6,168	0.6	21.8
...	All other causes (residual)	230,090	21.1	813.2
Hispanic, both sexes, all ages				
...	All causes	197,249	100.0	334.6
1	Malignant neoplasms (C00-C97)	40,550	20.6	68.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	39,402	20.0	66.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	16,757	8.5	28.4
4	Cerebrovascular diseases (I60-I69)	10,775	5.5	18.3
5	Diabetes mellitus (E10-E14)	9,216	4.7	15.6
6	Alzheimer disease (G30)	7,383	3.7	12.5
7	Chronic liver disease and cirrhosis (K70,K73-K74)	6,286	3.2	10.7
8	Chronic lower respiratory diseases (J40-J47)	5,599	2.8	9.5
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,933	2.0	6.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,928	2.0	6.7
11	Influenza and pneumonia (J09-J18)	3,851	2.0	6.5
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,186	1.6	5.4
13	Septicemia (A40-A41)	2,869	1.5	4.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,799	1.4	4.7
15	Certain conditions originating in the perinatal period (P00-P96)	2,409	1.2	4.1
...	All other causes (residual)	38,306	19.4	65.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 1-4 years				
...	All causes	805	100.0	19.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	234	29.1	5.6
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	112	13.9	2.7
3	Malignant neoplasms (C00-C97)	92	11.4	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	50	6.2	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.2	0.6
6	Influenza and pneumonia (J09-J18)	24	3.0	0.6
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.7	*
8	Cerebrovascular diseases (I60-I69)	12	1.5	*
9	Septicemia (A40-A41)	11	1.4	*
9	Certain conditions originating in the perinatal period (P00-P96)	11	1.4	*
11	Acute bronchitis and bronchiolitis (J20-J21)	6	0.7	*
12	Chronic lower respiratory diseases (J40-J47)	5	0.6	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.6	*
14	Meningitis (G00,G03)	4	0.5	*
15	Anemias (D50-D64)	3	0.4	*
...	All other causes (residual)	196	24.3	4.7
Hispanic, both sexes, 5-14 years				
...	All causes	1,165	100.0	11.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	302	25.9	2.9
2	Malignant neoplasms (C00-C97)	222	19.1	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	91	7.8	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	81	7.0	0.8
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	60	5.2	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	38	3.3	0.4
7	Cerebrovascular diseases (I60-I69)	24	2.1	0.2
8	Influenza and pneumonia (J09-J18)	21	1.8	0.2
9	Chronic lower respiratory diseases (J40-J47)	19	1.6	*
10	Septicemia (A40-A41)	18	1.5	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	14	1.2	*
12	Diabetes mellitus (E10-E14)	5	0.4	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	5	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
15	Diseases of appendix (K35-K38)	4	0.3	*
...	All other causes (residual)	256	22.0	2.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 15-24 years				
...	All causes	5,854	100.0	60.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,486	42.5	25.5
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	980	16.7	10.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	971	16.6	10.0
4	Malignant neoplasms (C00-C97)	371	6.3	3.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	131	2.2	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	71	1.2	0.7
7	Cerebrovascular diseases (I60-I69)	41	0.7	0.4
8	Pregnancy, childbirth and the puerperium (O00-O99)	34	0.6	0.3
9	Influenza and pneumonia (J09-J18)	32	0.5	0.3
10	Chronic lower respiratory diseases (J40-J47)	30	0.5	0.3
11	Diabetes mellitus (E10-E14)	29	0.5	0.3
12	Legal intervention (Y35,Y89.0)	27	0.5	0.3
13	Septicemia (A40-A41)	23	0.4	0.2
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.3	*
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.2	*
...	All other causes (residual)	601	10.3	6.2
Hispanic, both sexes, 25-44 years				
...	All causes	19,230	100.0	108.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,283	32.7	35.3
2	Malignant neoplasms (C00-C97)	2,505	13.0	14.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,668	8.7	9.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,660	8.6	9.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,489	7.7	8.4
6	Chronic liver disease and cirrhosis (K70,K73-K74)	800	4.2	4.5
7	Cerebrovascular diseases (I60-I69)	476	2.5	2.7
8	Diabetes mellitus (E10-E14)	393	2.0	2.2
9	Human immunodeficiency virus (HIV) disease (B20-B24)	229	1.2	1.3
10	Septicemia (A40-A41)	183	1.0	1.0
11	Influenza and pneumonia (J09-J18)	162	0.8	0.9
12	Pregnancy, childbirth and the puerperium (O00-O99)	136	0.7	0.8
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	134	0.7	0.8
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	118	0.6	0.7
15	Chronic lower respiratory diseases (J40-J47)	115	0.6	0.6
...	All other causes (residual)	2,879	15.0	16.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, both sexes, 45-64 years				
...	All causes	46,942	100.0	404.8
1	Malignant neoplasms (C00-C97)	12,662	27.0	109.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,534	18.2	73.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,427	9.4	38.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	3,362	7.2	29.0
5	Diabetes mellitus (E10-E14)	2,676	5.7	23.1
6	Cerebrovascular diseases (I60-I69)	2,044	4.4	17.6
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	910	1.9	7.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	891	1.9	7.7
9	Septicemia (A40-A41)	740	1.6	6.4
10	Chronic lower respiratory diseases (J40-J47)	714	1.5	6.2
11	Influenza and pneumonia (J09-J18)	688	1.5	5.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	562	1.2	4.8
13	Viral hepatitis (B15-B19)	555	1.2	4.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	490	1.0	4.2
15	Human immunodeficiency virus (HIV) disease (B20-B24)	443	0.9	3.8
...	All other causes (residual)	7,244	15.4	62.5
Hispanic, both sexes, 65 years and over				
...	All causes	118,434	100.0	2,817.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	28,956	24.4	688.8
2	Malignant neoplasms (C00-C97)	24,681	20.8	587.1
3	Cerebrovascular diseases (I60-I69)	8,166	6.9	194.2
4	Alzheimer disease (G30)	7,266	6.1	172.8
5	Diabetes mellitus (E10-E14)	6,112	5.2	145.4
6	Chronic lower respiratory diseases (J40-J47)	4,711	4.0	112.1
7	Influenza and pneumonia (J09-J18)	2,889	2.4	68.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,872	2.4	68.3
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,818	2.4	67.0
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,162	1.8	51.4
11	Chronic liver disease and cirrhosis (K70,K73-K74)	2,116	1.8	50.3
12	Septicemia (A40-A41)	1,863	1.6	44.3
13	Parkinson disease (G20-G21)	1,750	1.5	41.6
14	Pneumonitis due to solids and liquids (J69)	836	0.7	19.9
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	600	0.5	14.3
...	All other causes (residual)	20,636	17.4	490.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, all ages				
...	All causes	108,579	100.0	364.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,009	20.3	73.9
2	Malignant neoplasms (C00-C97)	21,034	19.4	70.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	12,455	11.5	41.8
4	Cerebrovascular diseases (I60-I69)	5,073	4.7	17.0
5	Diabetes mellitus (E10-E14)	5,054	4.7	17.0
6	Chronic liver disease and cirrhosis (K70,K73-K74)	4,369	4.0	14.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,175	2.9	10.7
8	Chronic lower respiratory diseases (J40-J47)	2,759	2.5	9.3
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,588	2.4	8.7
10	Alzheimer disease (G30)	2,297	2.1	7.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,988	1.8	6.7
12	Influenza and pneumonia (J09-J18)	1,911	1.8	6.4
13	Septicemia (A40-A41)	1,473	1.4	4.9
14	Certain conditions originating in the perinatal period (P00-P96)	1,356	1.2	4.6
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,354	1.2	4.5
...	All other causes (residual)	19,684	18.1	66.1
Hispanic, male, 1-4 years				
...	All causes	455	100.0	21.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	143	31.4	6.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	55	12.1	2.6
3	Malignant neoplasms (C00-C97)	50	11.0	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	30	6.6	1.4
5	Influenza and pneumonia (J09-J18)	15	3.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	14	3.1	*
7	Cerebrovascular diseases (I60-I69)	9	2.0	*
8	Septicemia (A40-A41)	7	1.5	*
8	Certain conditions originating in the perinatal period (P00-P96)	7	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	1.3	*
11	Acute bronchitis and bronchiolitis (J20-J21)	5	1.1	*
12	Chronic lower respiratory diseases (J40-J47)	4	0.9	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.9	*
14	Meningitis (G00,G03)	2	0.4	*
14	Diseases of appendix (K35-K38)	2	0.4	*
...	All other causes (residual)	102	22.4	4.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 5-14 years				
...	All causes	677	100.0	12.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	181	26.7	3.4
2	Malignant neoplasms (C00-C97)	113	16.7	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	61	9.0	1.1
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	6.9	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	5.6	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	26	3.8	0.5
7	Cerebrovascular diseases (I60-I69)	10	1.5	*
7	Influenza and pneumonia (J09-J18)	10	1.5	*
7	Chronic lower respiratory diseases (J40-J47)	10	1.5	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.3	*
11	Septicemia (A40-A41)	8	1.2	*
12	Diabetes mellitus (E10-E14)	5	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.6	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.4	*
15	Diseases of appendix (K35-K38)	2	0.3	*
...	All other causes (residual)	150	22.2	2.8
Hispanic, male, 15-24 years				
...	All causes	4,387	100.0	87.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,890	43.1	37.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	839	19.1	16.8
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	780	17.8	15.6
4	Malignant neoplasms (C00-C97)	232	5.3	4.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	99	2.3	2.0
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	40	0.9	0.8
7	Cerebrovascular diseases (I60-I69)	26	0.6	0.5
8	Legal intervention (Y35,Y89.0)	25	0.6	0.5
9	Diabetes mellitus (E10-E14)	19	0.4	*
10	Influenza and pneumonia (J09-J18)	16	0.4	*
10	Chronic lower respiratory diseases (J40-J47)	16	0.4	*
12	Septicemia (A40-A41)	15	0.3	*
13	Human immunodeficiency virus (HIV) disease (B20-B24)	9	0.2	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	7	0.2	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	6	0.1	*
15	Anemias (D50-D64)	6	0.1	*
...	All other causes (residual)	362	8.3	7.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 25-44 years				
...	All causes	13,667	100.0	148.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,062	37.0	54.8
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,365	10.0	14.8
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,238	9.1	13.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,222	8.9	13.2
5	Malignant neoplasms (C00-C97)	1,099	8.0	11.9
6	Chronic liver disease and cirrhosis (K70,K73-K74)	586	4.3	6.3
7	Cerebrovascular diseases (I60-I69)	316	2.3	3.4
8	Diabetes mellitus (E10-E14)	256	1.9	2.8
9	Human immunodeficiency virus (HIV) disease (B20-B24)	187	1.4	2.0
10	Septicemia (A40-A41)	110	0.8	1.2
11	Influenza and pneumonia (J09-J18)	101	0.7	1.1
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	84	0.6	0.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	74	0.5	0.8
14	Legal intervention (Y35,Y89.0)	71	0.5	0.8
15	Chronic lower respiratory diseases (J40-J47)	70	0.5	0.8
...	All other causes (residual)	1,826	13.4	19.8
Hispanic, male, 45-64 years				
...	All causes	30,119	100.0	523.0
1	Malignant neoplasms (C00-C97)	6,601	21.9	114.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,173	20.5	107.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,475	11.5	60.3
4	Chronic liver disease and cirrhosis (K70,K73-K74)	2,544	8.4	44.2
5	Diabetes mellitus (E10-E14)	1,644	5.5	28.5
6	Cerebrovascular diseases (I60-I69)	1,286	4.3	22.3
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	726	2.4	12.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	518	1.7	9.0
9	Septicemia (A40-A41)	438	1.5	7.6
10	Influenza and pneumonia (J09-J18)	394	1.3	6.8
11	Viral hepatitis (B15-B19)	391	1.3	6.8
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	379	1.3	6.6
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	371	1.2	6.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	359	1.2	6.2
15	Chronic lower respiratory diseases (J40-J47)	356	1.2	6.2
...	All other causes (residual)	4,464	14.8	77.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, male, 65 years and over				
...	All causes	56,631	100.0	3,105.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,426	25.5	791.1
2	Malignant neoplasms (C00-C97)	12,933	22.8	709.2
3	Cerebrovascular diseases (I60-I69)	3,421	6.0	187.6
4	Diabetes mellitus (E10-E14)	3,129	5.5	171.6
5	Chronic lower respiratory diseases (J40-J47)	2,300	4.1	126.1
6	Alzheimer disease (G30)	2,250	4.0	123.4
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,589	2.8	87.1
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,369	2.4	75.1
9	Influenza and pneumonia (J09-J18)	1,352	2.4	74.1
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,233	2.2	67.6
11	Parkinson disease (G20-G21)	1,022	1.8	56.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	927	1.6	50.8
13	Septicemia (A40-A41)	879	1.6	48.2
14	Pneumonitis due to solids and liquids (J69)	442	0.8	24.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	309	0.5	16.9
...	All other causes (residual)	9,050	16.0	496.3
Hispanic, female, all ages				
...	All causes	88,670	100.0	304.0
1	Malignant neoplasms (C00-C97)	19,516	22.0	66.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	17,393	19.6	59.6
3	Cerebrovascular diseases (I60-I69)	5,702	6.4	19.5
4	Alzheimer disease (G30)	5,086	5.7	17.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,302	4.9	14.7
6	Diabetes mellitus (E10-E14)	4,162	4.7	14.3
7	Chronic lower respiratory diseases (J40-J47)	2,840	3.2	9.7
8	Influenza and pneumonia (J09-J18)	1,940	2.2	6.7
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,940	2.2	6.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	1,917	2.2	6.6
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,445	1.6	5.0
12	Septicemia (A40-A41)	1,396	1.6	4.8
13	Certain conditions originating in the perinatal period (P00-P96)	1,053	1.2	3.6
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	873	1.0	3.0
15	Parkinson disease (G20-G21)	758	0.9	2.6
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	758	0.9	2.6
...	All other causes (residual)	17,589	19.8	60.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 1-4 years				
...	All causes	350	100.0	17.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	91	26.0	4.5
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	57	16.3	2.8
3	Malignant neoplasms (C00-C97)	42	12.0	2.1
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	20	5.7	1.0
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	3.4	*
6	Influenza and pneumonia (J09-J18)	9	2.6	*
7	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	8	2.3	*
8	Septicemia (A40-A41)	4	1.1	*
8	Certain conditions originating in the perinatal period (P00-P96)	4	1.1	*
10	Cerebrovascular diseases (I60-I69)	3	0.9	*
11	Anemias (D50-D64)	2	0.6	*
11	Meningitis (G00,G03)	2	0.6	*
13	Meningococcal infection (A39)	1	0.3	*
13	Acute bronchitis and bronchiolitis (J20-J21)	1	0.3	*
13	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
13	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
...	All other causes (residual)	91	26.0	4.5
Hispanic, female, 5-14 years				
...	All causes	488	100.0	9.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	121	24.8	2.4
2	Malignant neoplasms (C00-C97)	109	22.3	2.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	34	7.0	0.7
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	30	6.1	0.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	22	4.5	0.4
6	Cerebrovascular diseases (I60-I69)	14	2.9	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	12	2.5	*
8	Influenza and pneumonia (J09-J18)	11	2.3	*
9	Septicemia (A40-A41)	10	2.0	*
10	Chronic lower respiratory diseases (J40-J47)	9	1.8	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.0	*
12	Diseases of appendix (K35-K38)	2	0.4	*
12	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.4	*
15	Anemias (D50-D64)	1	0.2	*
15	Meningitis (G00,G03)	1	0.2	*
15	Pneumonitis due to solids and liquids (J69)	1	0.2	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.2	*
...	All other causes (residual)	101	20.7	2.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 15-24 years				
...	All causes	1,467	100.0	31.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	596	40.6	12.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	200	13.6	4.2
3	Malignant neoplasms (C00-C97)	139	9.5	2.9
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	132	9.0	2.8
5	Pregnancy, childbirth and the puerperium (O00-O99)	34	2.3	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	32	2.2	0.7
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	2.1	0.7
8	Influenza and pneumonia (J09-J18)	16	1.1	*
9	Cerebrovascular diseases (I60-I69)	15	1.0	*
10	Chronic lower respiratory diseases (J40-J47)	14	1.0	*
11	Diabetes mellitus (E10-E14)	10	0.7	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.6	*
13	Septicemia (A40-A41)	8	0.5	*
14	Cholelithiasis and other disorders of gallbladder (K80-K82)	5	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.3	*
...	All other causes (residual)	222	15.1	4.7
Hispanic, female, 25-44 years				
...	All causes	5,563	100.0	65.0
1	Malignant neoplasms (C00-C97)	1,406	25.3	16.4
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,221	21.9	14.3
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	422	7.6	4.9
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	303	5.4	3.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	267	4.8	3.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	214	3.8	2.5
7	Cerebrovascular diseases (I60-I69)	160	2.9	1.9
8	Diabetes mellitus (E10-E14)	137	2.5	1.6
9	Pregnancy, childbirth and the puerperium (O00-O99)	136	2.4	1.6
10	Septicemia (A40-A41)	73	1.3	0.9
11	Influenza and pneumonia (J09-J18)	61	1.1	0.7
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	50	0.9	0.6
13	Chronic lower respiratory diseases (J40-J47)	45	0.8	0.5
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	44	0.8	0.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	42	0.8	0.5
...	All other causes (residual)	982	17.7	11.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Hispanic, female, 45-64 years				
...	All causes	16,823	100.0	288.1
1	Malignant neoplasms (C00-C97)	6,061	36.0	103.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,361	14.0	40.4
3	Diabetes mellitus (E10-E14)	1,032	6.1	17.7
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	952	5.7	16.3
5	Chronic liver disease and cirrhosis (K70,K73-K74)	818	4.9	14.0
6	Cerebrovascular diseases (I60-I69)	758	4.5	13.0
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	373	2.2	6.4
8	Chronic lower respiratory diseases (J40-J47)	358	2.1	6.1
9	Septicemia (A40-A41)	302	1.8	5.2
10	Influenza and pneumonia (J09-J18)	294	1.7	5.0
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	191	1.1	3.3
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	184	1.1	3.2
13	Viral hepatitis (B15-B19)	164	1.0	2.8
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	111	0.7	1.9
15	Human immunodeficiency virus (HIV) disease (B20-B24)	84	0.5	1.4
...	All other causes (residual)	2,780	16.5	47.6
Hispanic, female, 65 years and over				
...	All causes	61,803	100.0	2,596.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,530	23.5	610.4
2	Malignant neoplasms (C00-C97)	11,748	19.0	493.5
3	Alzheimer disease (G30)	5,016	8.1	210.7
4	Cerebrovascular diseases (I60-I69)	4,745	7.7	199.3
5	Diabetes mellitus (E10-E14)	2,983	4.8	125.3
6	Chronic lower respiratory diseases (J40-J47)	2,411	3.9	101.3
7	Influenza and pneumonia (J09-J18)	1,537	2.5	64.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,503	2.4	63.1
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,235	2.0	51.9
10	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,229	2.0	51.6
11	Septicemia (A40-A41)	984	1.6	41.3
12	Chronic liver disease and cirrhosis (K70,K73-K74)	883	1.4	37.1
13	Parkinson disease (G20-G21)	728	1.2	30.6
14	Pneumonitis due to solids and liquids (J69)	394	0.6	16.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	291	0.5	12.2
...	All other causes (residual)	11,586	18.7	486.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, all ages				
...	All causes	2,179,857	100.0	1,083.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	508,485	23.3	252.7
2	Malignant neoplasms (C00-C97)	465,679	21.4	231.4
3	Chronic lower respiratory diseases (J40-J47)	139,833	6.4	69.5
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	127,029	5.8	63.1
5	Cerebrovascular diseases (I60-I69)	110,038	5.0	54.7
6	Alzheimer disease (G30)	101,876	4.7	50.6
7	Diabetes mellitus (E10-E14)	55,116	2.5	27.4
8	Influenza and pneumonia (J09-J18)	43,397	2.0	21.6
9	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	38,106	1.7	18.9
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	35,191	1.6	17.5
11	Chronic liver disease and cirrhosis (K70,K73-K74)	30,223	1.4	15.0
12	Septicemia (A40-A41)	30,198	1.4	15.0
13	Parkinson disease (G20-G21)	27,582	1.3	13.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	24,465	1.1	12.2
15	Pneumonitis due to solids and liquids (J69)	16,449	0.8	8.2
...	All other causes (residual)	426,190	19.6	211.8
Non-Hispanic, white, both sexes, 1-4 years				
...	All causes	1,858	100.0	22.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	677	36.4	8.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	183	9.8	2.2
3	Malignant neoplasms (C00-C97)	164	8.8	2.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	113	6.1	1.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	58	3.1	0.7
6	Influenza and pneumonia (J09-J18)	44	2.4	0.5
7	Cerebrovascular diseases (I60-I69)	29	1.6	0.3
8	Septicemia (A40-A41)	24	1.3	0.3
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	22	1.2	0.3
10	Anemias (D50-D64)	12	0.6	*
11	Acute bronchitis and bronchiolitis (J20-J21)	9	0.5	*
11	Certain conditions originating in the perinatal period (P00-P96)	9	0.5	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	9	0.5	*
14	Chronic lower respiratory diseases (J40-J47)	8	0.4	*
15	Meningitis (G00,G03)	7	0.4	*
...	All other causes (residual)	490	26.4	5.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 5-14 years				
...	All causes	2,782	100.0	12.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	819	29.4	3.8
2	Malignant neoplasms (C00-C97)	433	15.6	2.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	328	11.8	1.5
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	193	6.9	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	118	4.2	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	84	3.0	0.4
7	Influenza and pneumonia (J09-J18)	65	2.3	0.3
8	Cerebrovascular diseases (I60-I69)	45	1.6	0.2
9	Chronic lower respiratory diseases (J40-J47)	35	1.3	0.2
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	28	1.0	0.1
11	Septicemia (A40-A41)	19	0.7	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	15	0.5	*
13	Diabetes mellitus (E10-E14)	11	0.4	*
14	Certain conditions originating in the perinatal period (P00-P96)	9	0.3	*
15	Anemias (D50-D64)	7	0.3	*
15	Meningitis (G00,G03)	7	0.3	*
15	Pneumonitis due to solids and liquids (J69)	7	0.3	*
...	All other causes (residual)	559	20.1	2.6
Non-Hispanic, white, both sexes, 15-24 years				
...	All causes	17,207	100.0	72.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	8,555	49.7	35.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	4,074	23.7	17.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	763	4.4	3.2
4	Malignant neoplasms (C00-C97)	686	4.0	2.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	440	2.6	1.8
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	190	1.1	0.8
7	Diabetes mellitus (E10-E14)	129	0.7	0.5
8	Influenza and pneumonia (J09-J18)	100	0.6	0.4
9	Chronic lower respiratory diseases (J40-J47)	71	0.4	0.3
10	Pregnancy, childbirth and the puerperium (O00-O99)	69	0.4	0.3
11	Cerebrovascular diseases (I60-I69)	66	0.4	0.3
12	Septicemia (A40-A41)	65	0.4	0.3
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	48	0.3	0.2
14	Complications of medical and surgical care (Y40-Y84,Y88)	28	0.2	0.1
15	Legal intervention (Y35,Y89.0)	26	0.2	0.1
...	All other causes (residual)	1,897	11.0	8.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 25-44 years				
...	All causes	86,160	100.0	175.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,372	39.9	69.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11,450	13.3	23.3
3	Malignant neoplasms (C00-C97)	8,465	9.8	17.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,877	9.1	16.0
5	Chronic liver disease and cirrhosis (K70,K73-K74)	2,470	2.9	5.0
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,186	2.5	4.4
7	Diabetes mellitus (E10-E14)	1,447	1.7	2.9
8	Cerebrovascular diseases (I60-I69)	1,112	1.3	2.3
9	Septicemia (A40-A41)	689	0.8	1.4
10	Influenza and pneumonia (J09-J18)	683	0.8	1.4
11	Chronic lower respiratory diseases (J40-J47)	605	0.7	1.2
12	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	576	0.7	1.2
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	397	0.5	0.8
14	Pregnancy, childbirth and the puerperium (O00-O99)	396	0.5	0.8
15	Human immunodeficiency virus (HIV) disease (B20-B24)	286	0.3	0.6
...	All other causes (residual)	13,149	15.3	26.7
Non-Hispanic, white, both sexes, 45-64 years				
...	All causes	377,764	100.0	663.3
1	Malignant neoplasms (C00-C97)	110,485	29.2	194.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	76,455	20.2	134.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	34,413	9.1	60.4
4	Chronic lower respiratory diseases (J40-J47)	18,759	5.0	32.9
5	Chronic liver disease and cirrhosis (K70,K73-K74)	15,696	4.2	27.6
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	14,423	3.8	25.3
7	Diabetes mellitus (E10-E14)	12,836	3.4	22.5
8	Cerebrovascular diseases (I60-I69)	10,304	2.7	18.1
9	Septicemia (A40-A41)	5,356	1.4	9.4
10	Influenza and pneumonia (J09-J18)	4,757	1.3	8.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,087	1.1	7.2
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,126	0.8	5.5
13	Viral hepatitis (B15-B19)	2,255	0.6	4.0
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,794	0.5	3.2
15	Pneumonitis due to solids and liquids (J69)	1,588	0.4	2.8
...	All other causes (residual)	61,430	16.3	107.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, both sexes, 65 years and over				
...	All causes	1,684,667	100.0	4,285.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	423,429	25.1	1,077.1
2	Malignant neoplasms (C00-C97)	345,411	20.5	878.7
3	Chronic lower respiratory diseases (J40-J47)	120,347	7.1	306.1
4	Alzheimer disease (G30)	100,850	6.0	256.5
5	Cerebrovascular diseases (I60-I69)	98,433	5.8	250.4
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,587	2.8	121.1
7	Diabetes mellitus (E10-E14)	40,689	2.4	103.5
8	Influenza and pneumonia (J09-J18)	37,689	2.2	95.9
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	30,643	1.8	78.0
10	Parkinson disease (G20-G21)	26,963	1.6	68.6
11	Septicemia (A40-A41)	23,992	1.4	61.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	21,087	1.3	53.6
13	Pneumonitis due to solids and liquids (J69)	14,651	0.9	37.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	12,040	0.7	30.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11,231	0.7	28.6
...	All other causes (residual)	329,625	19.6	838.5
Non-Hispanic, white, male, all ages				
...	All causes	1,102,838	100.0	1,111.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	272,428	24.7	274.5
2	Malignant neoplasms (C00-C97)	247,022	22.4	248.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79,280	7.2	79.9
4	Chronic lower respiratory diseases (J40-J47)	64,673	5.9	65.2
5	Cerebrovascular diseases (I60-I69)	45,078	4.1	45.4
6	Alzheimer disease (G30)	31,577	2.9	31.8
7	Diabetes mellitus (E10-E14)	31,343	2.8	31.6
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	29,708	2.7	29.9
9	Influenza and pneumonia (J09-J18)	20,515	1.9	20.7
10	Chronic liver disease and cirrhosis (K70,K73-K74)	18,866	1.7	19.0
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	18,187	1.6	18.3
12	Parkinson disease (G20-G21)	16,841	1.5	17.0
13	Septicemia (A40-A41)	14,419	1.3	14.5
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	10,696	1.0	10.8
15	Pneumonitis due to solids and liquids (J69)	9,352	0.8	9.4
...	All other causes (residual)	192,853	17.5	194.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 1-4 years				
...	All causes	1,107	100.0	26.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	425	38.4	10.0
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	9.1	2.4
3	Malignant neoplasms (C00-C97)	93	8.4	2.2
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	64	5.8	1.5
5	Influenza and pneumonia (J09-J18)	30	2.7	0.7
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	2.6	0.7
7	Cerebrovascular diseases (I60-I69)	18	1.6	*
8	Septicemia (A40-A41)	12	1.1	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.0	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
11	Meningitis (G00,G03)	6	0.5	*
12	Acute bronchitis and bronchiolitis (J20-J21)	5	0.5	*
13	Anemias (D50-D64)	4	0.4	*
13	Certain conditions originating in the perinatal period (P00-P96)	4	0.4	*
15	Pneumonitis due to solids and liquids (J69)	3	0.3	*
...	All other causes (residual)	295	26.6	6.9
Non-Hispanic, white, male, 5-14 years				
...	All causes	1,628	100.0	14.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	514	31.6	4.6
2	Malignant neoplasms (C00-C97)	235	14.4	2.1
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	227	13.9	2.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	103	6.3	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	62	3.8	0.6
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	41	2.5	0.4
7	Cerebrovascular diseases (I60-I69)	27	1.7	0.2
8	Influenza and pneumonia (J09-J18)	25	1.5	0.2
9	Chronic lower respiratory diseases (J40-J47)	19	1.2	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	15	0.9	*
11	Septicemia (A40-A41)	13	0.8	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
13	Diabetes mellitus (E10-E14)	6	0.4	*
14	Anemias (D50-D64)	5	0.3	*
14	Meningitis (G00,G03)	5	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.3	*
...	All other causes (residual)	318	19.5	2.9

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 15-24 years				
...	All causes	12,285	100.0	100.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,120	49.8	50.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,328	27.1	27.2
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	538	4.4	4.4
4	Malignant neoplasms (C00-C97)	413	3.4	3.4
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	292	2.4	2.4
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	110	0.9	0.9
7	Diabetes mellitus (E10-E14)	74	0.6	0.6
8	Influenza and pneumonia (J09-J18)	56	0.5	0.5
9	Chronic lower respiratory diseases (J40-J47)	44	0.4	0.4
10	Cerebrovascular diseases (I60-I69)	32	0.3	0.3
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	30	0.2	0.2
12	Septicemia (A40-A41)	27	0.2	0.2
13	Legal intervention (Y35,Y89.0)	22	0.2	0.2
14	Aortic aneurysm and dissection (I71)	18	0.1	*
14	Pneumonitis due to solids and liquids (J69)	18	0.1	*
...	All other causes (residual)	1,163	9.5	9.5
Non-Hispanic, white, male, 25-44 years				
...	All causes	56,634	100.0	227.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24,406	43.1	98.2
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,912	15.7	35.9
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	5,349	9.4	21.5
4	Malignant neoplasms (C00-C97)	3,874	6.8	15.6
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,519	2.7	6.1
6	Chronic liver disease and cirrhosis (K70,K73-K74)	1,464	2.6	5.9
7	Diabetes mellitus (E10-E14)	883	1.6	3.6
8	Cerebrovascular diseases (I60-I69)	618	1.1	2.5
9	Influenza and pneumonia (J09-J18)	361	0.6	1.5
10	Septicemia (A40-A41)	355	0.6	1.4
11	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	344	0.6	1.4
12	Chronic lower respiratory diseases (J40-J47)	270	0.5	1.1
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	233	0.4	0.9
14	Human immunodeficiency virus (HIV) disease (B20-B24)	214	0.4	0.9
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	175	0.3	0.7
...	All other causes (residual)	7,657	13.5	30.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, male, 45-64 years				
...	All causes	230,079	100.0	819.5
1	Malignant neoplasms (C00-C97)	59,137	25.7	210.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	53,985	23.5	192.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,175	10.1	82.5
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	10,734	4.7	38.2
5	Chronic liver disease and cirrhosis (K70,K73-K74)	10,175	4.4	36.2
6	Chronic lower respiratory diseases (J40-J47)	9,342	4.1	33.3
7	Diabetes mellitus (E10-E14)	8,222	3.6	29.3
8	Cerebrovascular diseases (I60-I69)	5,873	2.6	20.9
9	Septicemia (A40-A41)	2,898	1.3	10.3
10	Influenza and pneumonia (J09-J18)	2,620	1.1	9.3
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,383	1.0	8.5
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,008	0.9	7.2
13	Viral hepatitis (B15-B19)	1,519	0.7	5.4
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,212	0.5	4.3
15	Aortic aneurysm and dissection (I71)	1,015	0.4	3.6
...	All other causes (residual)	35,781	15.6	127.4
Non-Hispanic, white, male, 65 years and over				
...	All causes	795,777	100.0	4,497.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	212,656	26.7	1,201.9
2	Malignant neoplasms (C00-C97)	183,250	23.0	1,035.7
3	Chronic lower respiratory diseases (J40-J47)	54,991	6.9	310.8
4	Cerebrovascular diseases (I60-I69)	38,481	4.8	217.5
5	Alzheimer disease (G30)	31,124	3.9	175.9
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	24,270	3.0	137.2
7	Diabetes mellitus (E10-E14)	22,156	2.8	125.2
8	Influenza and pneumonia (J09-J18)	17,387	2.2	98.3
9	Parkinson disease (G20-G21)	16,423	2.1	92.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,539	2.0	87.8
11	Septicemia (A40-A41)	11,084	1.4	62.6
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	8,511	1.1	48.1
13	Pneumonitis due to solids and liquids (J69)	8,261	1.0	46.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	7,216	0.9	40.8
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	6,506	0.8	36.8
...	All other causes (residual)	137,922	17.3	779.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent	
			of total deaths	Rate
Non-Hispanic, white, female, all ages				
...	All causes	1,077,019	100.0	1,055.8
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	236,057	21.9	231.4
2	Malignant neoplasms (C00-C97)	218,657	20.3	214.3
3	Chronic lower respiratory diseases (J40-J47)	75,160	7.0	73.7
4	Alzheimer disease (G30)	70,299	6.5	68.9
5	Cerebrovascular diseases (I60-I69)	64,960	6.0	63.7
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	47,749	4.4	46.8
7	Diabetes mellitus (E10-E14)	23,773	2.2	23.3
8	Influenza and pneumonia (J09-J18)	22,882	2.1	22.4
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	17,004	1.6	16.7
10	Septicemia (A40-A41)	15,779	1.5	15.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	13,769	1.3	13.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	11,357	1.1	11.1
13	Parkinson disease (G20-G21)	10,741	1.0	10.5
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	8,398	0.8	8.2
15	Pneumonitis due to solids and liquids (J69)	7,097	0.7	7.0
...	All other causes (residual)	233,337	21.7	228.7
Non-Hispanic, white, female, 1-4 years				
...	All causes	751	100.0	18.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	252	33.6	6.2
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	82	10.9	2.0
3	Malignant neoplasms (C00-C97)	71	9.5	1.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	49	6.5	1.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	29	3.9	0.7
6	Influenza and pneumonia (J09-J18)	14	1.9	*
7	Septicemia (A40-A41)	12	1.6	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	11	1.5	*
8	Cerebrovascular diseases (I60-I69)	11	1.5	*
10	Anemias (D50-D64)	8	1.1	*
11	Chronic lower respiratory diseases (J40-J47)	6	0.8	*
12	Certain conditions originating in the perinatal period (P00-P96)	5	0.7	*
13	Acute bronchitis and bronchiolitis (J20-J21)	4	0.5	*
14	Hernia (K40-K46)	3	0.4	*
15	Meningococcal infection (A39)	2	0.3	*
15	Diabetes mellitus (E10-E14)	2	0.3	*
15	Pneumonitis due to solids and liquids (J69)	2	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.3	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	184	24.5	4.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 5-14 years				
...	All causes	1,154	100.0	10.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	305	26.4	2.9
2	Malignant neoplasms (C00-C97)	198	17.2	1.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	101	8.8	1.0
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	90	7.8	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	4.9	0.5
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	43	3.7	0.4
7	Influenza and pneumonia (J09-J18)	40	3.5	0.4
8	Cerebrovascular diseases (I60-I69)	18	1.6	*
9	Chronic lower respiratory diseases (J40-J47)	16	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	13	1.1	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	7	0.6	*
12	Septicemia (A40-A41)	6	0.5	*
13	Diabetes mellitus (E10-E14)	5	0.4	*
14	Pneumonitis due to solids and liquids (J69)	4	0.3	*
14	Certain conditions originating in the perinatal period (P00-P96)	4	0.3	*
...	All other causes (residual)	248	21.5	2.3
Non-Hispanic, white, female, 15-24 years				
...	All causes	4,922	100.0	42.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,435	49.5	21.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	746	15.2	6.4
3	Malignant neoplasms (C00-C97)	273	5.5	2.4
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	225	4.6	1.9
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	148	3.0	1.3
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	1.6	0.7
7	Pregnancy, childbirth and the puerperium (O00-O99)	69	1.4	0.6
8	Diabetes mellitus (E10-E14)	55	1.1	0.5
9	Influenza and pneumonia (J09-J18)	44	0.9	0.4
10	Septicemia (A40-A41)	38	0.8	0.3
11	Cerebrovascular diseases (I60-I69)	34	0.7	0.3
12	Chronic lower respiratory diseases (J40-J47)	27	0.5	0.2
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	0.4	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	13	0.3	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	13	0.3	*
...	All other causes (residual)	704	14.3	6.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 25-44 years				
...	All causes	29,526	100.0	121.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	9,966	33.8	40.9
2	Malignant neoplasms (C00-C97)	4,591	15.5	18.9
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,538	8.6	10.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,528	8.6	10.4
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,006	3.4	4.1
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	667	2.3	2.7
7	Diabetes mellitus (E10-E14)	564	1.9	2.3
8	Cerebrovascular diseases (I60-I69)	494	1.7	2.0
9	Pregnancy, childbirth and the puerperium (O00-O99)	396	1.3	1.6
10	Chronic lower respiratory diseases (J40-J47)	335	1.1	1.4
11	Septicemia (A40-A41)	334	1.1	1.4
12	Influenza and pneumonia (J09-J18)	322	1.1	1.3
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	232	0.8	1.0
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	164	0.6	0.7
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	91	0.3	0.4
...	All other causes (residual)	5,298	17.9	21.8
Non-Hispanic, white, female, 45-64 years				
...	All causes	147,685	100.0	511.5
1	Malignant neoplasms (C00-C97)	51,348	34.8	177.8
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	22,470	15.2	77.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11,238	7.6	38.9
4	Chronic lower respiratory diseases (J40-J47)	9,417	6.4	32.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	5,521	3.7	19.1
6	Diabetes mellitus (E10-E14)	4,614	3.1	16.0
7	Cerebrovascular diseases (I60-I69)	4,431	3.0	15.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	3,689	2.5	12.8
9	Septicemia (A40-A41)	2,458	1.7	8.5
10	Influenza and pneumonia (J09-J18)	2,137	1.4	7.4
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,704	1.2	5.9
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,118	0.8	3.9
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	763	0.5	2.6
14	Viral hepatitis (B15-B19)	736	0.5	2.5
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	638	0.4	2.2
...	All other causes (residual)	25,403	17.2	88.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, white, female, 65 years and over				
...	All causes	888,890	100.0	4,112.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	210,773	23.7	975.0
2	Malignant neoplasms (C00-C97)	162,161	18.2	750.2
3	Alzheimer disease (G30)	69,726	7.8	322.6
4	Chronic lower respiratory diseases (J40-J47)	65,356	7.4	302.3
5	Cerebrovascular diseases (I60-I69)	59,952	6.7	277.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23,317	2.6	107.9
7	Influenza and pneumonia (J09-J18)	20,302	2.3	93.9
8	Diabetes mellitus (E10-E14)	18,533	2.1	85.7
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15,104	1.7	69.9
10	Septicemia (A40-A41)	12,908	1.5	59.7
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	12,576	1.4	58.2
12	Parkinson disease (G20-G21)	10,540	1.2	48.8
13	Pneumonitis due to solids and liquids (J69)	6,390	0.7	29.6
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5,223	0.6	24.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	4,824	0.5	22.3
...	All other causes (residual)	191,205	21.5	884.5
Non-Hispanic, black, both sexes, all ages				
...	All causes	335,667	100.0	787.5
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	78,161	23.3	183.4
2	Malignant neoplasms (C00-C97)	69,872	20.8	163.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19,869	5.9	46.6
4	Cerebrovascular diseases (I60-I69)	19,088	5.7	44.8
5	Diabetes mellitus (E10-E14)	14,798	4.4	34.7
6	Chronic lower respiratory diseases (J40-J47)	11,217	3.3	26.3
7	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9,908	3.0	23.2
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9,542	2.8	22.4
9	Alzheimer disease (G30)	8,991	2.7	21.1
10	Septicemia (A40-A41)	6,568	2.0	15.4
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	6,273	1.9	14.7
12	Influenza and pneumonia (J09-J18)	5,556	1.7	13.0
13	Certain conditions originating in the perinatal period (P00-P96)	3,637	1.1	8.5
14	Chronic liver disease and cirrhosis (K70,K73-K74)	3,290	1.0	7.7
15	Human immunodeficiency virus (HIV) disease (B20-B24)	2,965	0.9	7.0
...	All other causes (residual)	65,932	19.6	154.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 1-4 years				
...	All causes	985	100.0	40.1
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	295	29.9	12.0
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	125	12.7	5.1
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	101	10.3	4.1
4	Malignant neoplasms (C00-C97)	42	4.3	1.7
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.8	1.5
6	Influenza and pneumonia (J09-J18)	25	2.5	1.0
7	Cerebrovascular diseases (I60-I69)	21	2.1	0.9
8	Chronic lower respiratory diseases (J40-J47)	18	1.8	*
8	Certain conditions originating in the perinatal period (P00-P96)	18	1.8	*
10	Complications of medical and surgical care (Y40-Y84,Y88)	10	1.0	*
11	Septicemia (A40-A41)	9	0.9	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	7	0.7	*
13	Anemias (D50-D64)	6	0.6	*
13	Meningitis (G00,G03)	6	0.6	*
13	Acute bronchitis and bronchiolitis (J20-J21)	6	0.6	*
...	All other causes (residual)	259	26.3	10.6
Non-Hispanic, black, both sexes, 5-14 years				
...	All causes	1,296	100.0	20.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	377	29.1	6.1
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	11.0	2.3
3	Malignant neoplasms (C00-C97)	136	10.5	2.2
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	80	6.2	1.3
5	Chronic lower respiratory diseases (J40-J47)	77	5.9	1.2
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	66	5.1	1.1
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	47	3.6	0.8
8	Cerebrovascular diseases (I60-I69)	25	1.9	0.4
9	Influenza and pneumonia (J09-J18)	23	1.8	0.4
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	18	1.4	*
11	Diabetes mellitus (E10-E14)	16	1.2	*
12	Septicemia (A40-A41)	14	1.1	*
13	Anemias (D50-D64)	12	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	5	0.4	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	5	0.4	*
...	All other causes (residual)	253	19.5	4.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 15-24 years				
...	All causes	7,364	100.0	111.4
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3,055	41.5	46.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,830	24.9	27.7
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	710	9.6	10.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	294	4.0	4.4
5	Malignant neoplasms (C00-C97)	219	3.0	3.3
6	Diabetes mellitus (E10-E14)	83	1.1	1.3
7	Chronic lower respiratory diseases (J40-J47)	82	1.1	1.2
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	78	1.1	1.2
9	Pregnancy, childbirth and the puerperium (O00-O99)	58	0.8	0.9
10	Human immunodeficiency virus (HIV) disease (B20-B24)	47	0.6	0.7
11	Anemias (D50-D64)	44	0.6	0.7
12	Influenza and pneumonia (J09-J18)	41	0.6	0.6
13	Cerebrovascular diseases (I60-I69)	38	0.5	0.6
14	Legal intervention (Y35,Y89.0)	37	0.5	0.6
15	Septicemia (A40-A41)	29	0.4	0.4
...	All other causes (residual)	719	9.8	10.9
Non-Hispanic, black, both sexes, 25-44 years				
...	All causes	27,810	100.0	234.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,021	21.7	50.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,858	17.5	41.0
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,897	14.0	32.9
4	Malignant neoplasms (C00-C97)	2,579	9.3	21.8
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,308	4.7	11.0
6	Diabetes mellitus (E10-E14)	927	3.3	7.8
7	Human immunodeficiency virus (HIV) disease (B20-B24)	784	2.8	6.6
8	Cerebrovascular diseases (I60-I69)	617	2.2	5.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	368	1.3	3.1
10	Septicemia (A40-A41)	313	1.1	2.6
11	Chronic lower respiratory diseases (J40-J47)	290	1.0	2.4
12	Chronic liver disease and cirrhosis (K70,K73-K74)	277	1.0	2.3
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	276	1.0	2.3
14	Influenza and pneumonia (J09-J18)	266	1.0	2.2
15	Pregnancy, childbirth and the puerperium (O00-O99)	265	1.0	2.2
...	All other causes (residual)	4,764	17.1	40.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, both sexes, 45-64 years				
...	All causes	95,724	100.0	933.4
1	Malignant neoplasms (C00-C97)	24,010	25.1	234.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,580	24.6	229.9
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	7,276	7.6	70.9
4	Diabetes mellitus (E10-E14)	4,689	4.9	45.7
5	Cerebrovascular diseases (I60-I69)	4,564	4.8	44.5
6	Chronic lower respiratory diseases (J40-J47)	2,647	2.8	25.8
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,547	2.7	24.8
8	Chronic liver disease and cirrhosis (K70,K73-K74)	2,003	2.1	19.5
9	Septicemia (A40-A41)	1,895	2.0	18.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,689	1.8	16.5
11	Human immunodeficiency virus (HIV) disease (B20-B24)	1,684	1.8	16.4
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,380	1.4	13.5
13	Influenza and pneumonia (J09-J18)	1,292	1.3	12.6
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	654	0.7	6.4
15	Viral hepatitis (B15-B19)	461	0.5	4.5
...	All other causes (residual)	15,353	16.0	149.7
Non-Hispanic, black, both sexes, 65 years and over				
...	All causes	195,742	100.0	4,203.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	50,208	25.7	1,078.2
2	Malignant neoplasms (C00-C97)	42,877	21.9	920.8
3	Cerebrovascular diseases (I60-I69)	13,786	7.0	296.1
4	Diabetes mellitus (E10-E14)	9,080	4.6	195.0
5	Alzheimer disease (G30)	8,864	4.5	190.4
6	Chronic lower respiratory diseases (J40-J47)	8,098	4.1	173.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	6,584	3.4	141.4
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4,302	2.2	92.4
9	Septicemia (A40-A41)	4,257	2.2	91.4
10	Influenza and pneumonia (J09-J18)	3,854	2.0	82.8
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,614	1.8	77.6
12	Pneumonitis due to solids and liquids (J69)	1,480	0.8	31.8
13	Parkinson disease (G20-G21)	1,448	0.7	31.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,007	0.5	21.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	924	0.5	19.8
...	All other causes (residual)	35,359	18.1	759.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, all ages				
...	All causes	174,403	100.0	854.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	41,411	23.7	202.8
2	Malignant neoplasms (C00-C97)	35,261	20.2	172.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	13,844	7.9	67.8
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8,643	5.0	42.3
5	Cerebrovascular diseases (I60-I69)	8,566	4.9	42.0
6	Diabetes mellitus (E10-E14)	7,494	4.3	36.7
7	Chronic lower respiratory diseases (J40-J47)	5,605	3.2	27.5
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,555	2.6	22.3
9	Septicemia (A40-A41)	3,052	1.7	14.9
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,874	1.6	14.1
11	Influenza and pneumonia (J09-J18)	2,682	1.5	13.1
12	Alzheimer disease (G30)	2,476	1.4	12.1
13	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	2,324	1.3	11.4
14	Certain conditions originating in the perinatal period (P00-P96)	2,080	1.2	10.2
15	Chronic liver disease and cirrhosis (K70,K73-K74)	2,064	1.2	10.1
...	All other causes (residual)	31,472	18.0	154.2
Non-Hispanic, black, male, 1-4 years				
...	All causes	535	100.0	42.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	165	30.8	13.2
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	69	12.9	5.5
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	50	9.3	4.0
4	Malignant neoplasms (C00-C97)	23	4.3	1.8
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	21	3.9	1.7
6	Influenza and pneumonia (J09-J18)	13	2.4	*
7	Cerebrovascular diseases (I60-I69)	11	2.1	*
8	Chronic lower respiratory diseases (J40-J47)	10	1.9	*
9	Certain conditions originating in the perinatal period (P00-P96)	6	1.1	*
9	Complications of medical and surgical care (Y40-Y84,Y88)	6	1.1	*
11	Septicemia (A40-A41)	4	0.7	*
11	Meningitis (G00,G03)	4	0.7	*
11	Acute bronchitis and bronchiolitis (J20-J21)	4	0.7	*
14	Anemias (D50-D64)	3	0.6	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	2	0.4	*
...	All other causes (residual)	144	26.9	11.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 5-14 years				
...	All causes	784	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	247	31.5	7.9
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	87	11.1	2.8
3	Malignant neoplasms (C00-C97)	73	9.3	2.3
4	Chronic lower respiratory diseases (J40-J47)	54	6.9	1.7
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	46	5.9	1.5
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	45	5.7	1.4
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	23	2.9	0.7
8	Cerebrovascular diseases (I60-I69)	15	1.9	*
9	Influenza and pneumonia (J09-J18)	11	1.4	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.1	*
10	Diabetes mellitus (E10-E14)	9	1.1	*
12	Septicemia (A40-A41)	8	1.0	*
13	Anemias (D50-D64)	7	0.9	*
14	Certain conditions originating in the perinatal period (P00-P96)	3	0.4	*
15	Pneumonitis due to solids and liquids (J69)	2	0.3	*
...	All other causes (residual)	145	18.5	4.6
Non-Hispanic, black, male, 15-24 years				
...	All causes	5,711	100.0	170.0
1	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2,761	48.3	82.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,328	23.3	39.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	565	9.9	16.8
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	199	3.5	5.9
5	Malignant neoplasms (C00-C97)	127	2.2	3.8
6	Chronic lower respiratory diseases (J40-J47)	52	0.9	1.5
7	Diabetes mellitus (E10-E14)	47	0.8	1.4
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	47	0.8	1.4
9	Human immunodeficiency virus (HIV) disease (B20-B24)	37	0.6	1.1
10	Legal intervention (Y35,Y89.0)	36	0.6	1.1
11	Influenza and pneumonia (J09-J18)	25	0.4	0.7
12	Cerebrovascular diseases (I60-I69)	19	0.3	*
13	Anemias (D50-D64)	17	0.3	*
14	Septicemia (A40-A41)	16	0.3	*
15	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	9	0.2	*
...	All other causes (residual)	426	7.5	12.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 25-44 years				
...	All causes	18,273	100.0	318.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	4,459	24.4	77.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4,324	23.7	75.4
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,460	13.5	42.9
4	Malignant neoplasms (C00-C97)	1,045	5.7	18.2
5	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,043	5.7	18.2
6	Diabetes mellitus (E10-E14)	521	2.9	9.1
7	Human immunodeficiency virus (HIV) disease (B20-B24)	498	2.7	8.7
8	Cerebrovascular diseases (I60-I69)	320	1.8	5.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	192	1.1	3.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	182	1.0	3.2
11	Septicemia (A40-A41)	162	0.9	2.8
11	Chronic lower respiratory diseases (J40-J47)	162	0.9	2.8
13	Chronic liver disease and cirrhosis (K70,K73-K74)	145	0.8	2.5
14	Influenza and pneumonia (J09-J18)	142	0.8	2.5
15	Anemias (D50-D64)	139	0.8	2.4
...	All other causes (residual)	2,479	13.6	43.3
Non-Hispanic, black, male, 45-64 years				
...	All causes	55,733	100.0	1,173.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	14,953	26.8	314.9
2	Malignant neoplasms (C00-C97)	12,088	21.7	254.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	5,240	9.4	110.3
4	Diabetes mellitus (E10-E14)	2,714	4.9	57.2
5	Cerebrovascular diseases (I60-I69)	2,608	4.7	54.9
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,373	2.5	28.9
7	Chronic lower respiratory diseases (J40-J47)	1,276	2.3	26.9
8	Chronic liver disease and cirrhosis (K70,K73-K74)	1,267	2.3	26.7
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,168	2.1	24.6
10	Human immunodeficiency virus (HIV) disease (B20-B24)	1,092	2.0	23.0
11	Septicemia (A40-A41)	985	1.8	20.7
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	982	1.8	20.7
13	Influenza and pneumonia (J09-J18)	701	1.3	14.8
14	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	504	0.9	10.6
15	Viral hepatitis (B15-B19)	331	0.6	7.0
...	All other causes (residual)	8,451	15.2	178.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, male, 65 years and over				
...	All causes	89,597	100.0	4,768.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	23,705	26.5	1,261.5
2	Malignant neoplasms (C00-C97)	21,900	24.4	1,165.5
3	Cerebrovascular diseases (I60-I69)	5,569	6.2	296.4
4	Diabetes mellitus (E10-E14)	4,203	4.7	223.7
5	Chronic lower respiratory diseases (J40-J47)	4,046	4.5	215.3
6	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2,965	3.3	157.8
7	Alzheimer disease (G30)	2,420	2.7	128.8
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,158	2.4	114.8
9	Septicemia (A40-A41)	1,850	2.1	98.5
10	Influenza and pneumonia (J09-J18)	1,757	2.0	93.5
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,704	1.9	90.7
12	Parkinson disease (G20-G21)	849	0.9	45.2
13	Pneumonitis due to solids and liquids (J69)	753	0.8	40.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	650	0.7	34.6
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	450	0.5	23.9
...	All other causes (residual)	14,618	16.3	777.9
Non-Hispanic, black, female, all ages				
...	All causes	161,264	100.0	726.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	36,750	22.8	165.5
2	Malignant neoplasms (C00-C97)	34,611	21.5	155.8
3	Cerebrovascular diseases (I60-I69)	10,522	6.5	47.4
4	Diabetes mellitus (E10-E14)	7,304	4.5	32.9
5	Alzheimer disease (G30)	6,515	4.0	29.3
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	6,025	3.7	27.1
7	Chronic lower respiratory diseases (J40-J47)	5,612	3.5	25.3
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4,987	3.1	22.5
9	Septicemia (A40-A41)	3,516	2.2	15.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	3,399	2.1	15.3
11	Influenza and pneumonia (J09-J18)	2,874	1.8	12.9
12	Certain conditions originating in the perinatal period (P00-P96)	1,557	1.0	7.0
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1,265	0.8	5.7
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1,226	0.8	5.5
15	Human immunodeficiency virus (HIV) disease (B20-B24)	1,002	0.6	4.5
...	All other causes (residual)	34,099	21.1	153.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 1-4 years				
...	All causes	450	100.0	37.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	28.9	10.8
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	56	12.4	4.6
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	51	11.3	4.2
4	Malignant neoplasms (C00-C97)	19	4.2	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	16	3.6	*
6	Influenza and pneumonia (J09-J18)	12	2.7	*
6	Certain conditions originating in the perinatal period (P00-P96)	12	2.7	*
8	Cerebrovascular diseases (I60-I69)	10	2.2	*
9	Chronic lower respiratory diseases (J40-J47)	8	1.8	*
10	Septicemia (A40-A41)	5	1.1	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	5	1.1	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.9	*
13	Anemias (D50-D64)	3	0.7	*
14	Diabetes mellitus (E10-E14)	2	0.4	*
14	Meningitis (G00,G03)	2	0.4	*
14	Acute bronchitis and bronchiolitis (J20-J21)	2	0.4	*
...	All other causes (residual)	113	25.1	9.4
Non-Hispanic, black, female, 5-14 years				
...	All causes	512	100.0	16.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	130	25.4	4.3
2	Malignant neoplasms (C00-C97)	63	12.3	2.1
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	55	10.7	1.8
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	35	6.8	1.1
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	24	4.7	0.8
6	Chronic lower respiratory diseases (J40-J47)	23	4.5	0.8
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	20	3.9	0.7
8	Influenza and pneumonia (J09-J18)	12	2.3	*
9	Cerebrovascular diseases (I60-I69)	10	2.0	*
10	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	1.8	*
11	Diabetes mellitus (E10-E14)	7	1.4	*
12	Septicemia (A40-A41)	6	1.2	*
13	Anemias (D50-D64)	5	1.0	*
14	Complications of medical and surgical care (Y40-Y84,Y88)	4	0.8	*
15	Certain conditions originating in the perinatal period (P00-P96)	2	0.4	*
...	All other causes (residual)	107	20.9	3.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 15-24 years				
...	All causes	1,653	100.0	50.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	502	30.4	15.4
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	294	17.8	9.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	8.8	4.5
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	95	5.7	2.9
5	Malignant neoplasms (C00-C97)	92	5.6	2.8
6	Pregnancy, childbirth and the puerperium (O00-O99)	58	3.5	1.8
7	Diabetes mellitus (E10-E14)	36	2.2	1.1
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	31	1.9	1.0
9	Chronic lower respiratory diseases (J40-J47)	30	1.8	0.9
10	Anemias (D50-D64)	27	1.6	0.8
11	Cerebrovascular diseases (I60-I69)	19	1.1	*
12	Influenza and pneumonia (J09-J18)	16	1.0	*
13	Septicemia (A40-A41)	13	0.8	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	10	0.6	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	8	0.5	*
...	All other causes (residual)	277	16.8	8.5
Non-Hispanic, black, female, 25-44 years				
...	All causes	9,537	100.0	155.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,562	16.4	25.5
2	Malignant neoplasms (C00-C97)	1,534	16.1	25.1
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,437	15.1	23.5
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	534	5.6	8.7
5	Diabetes mellitus (E10-E14)	406	4.3	6.6
6	Cerebrovascular diseases (I60-I69)	297	3.1	4.9
7	Human immunodeficiency virus (HIV) disease (B20-B24)	286	3.0	4.7
8	Pregnancy, childbirth and the puerperium (O00-O99)	265	2.8	4.3
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	265	2.8	4.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	176	1.8	2.9
11	Septicemia (A40-A41)	151	1.6	2.5
12	Chronic liver disease and cirrhosis (K70,K73-K74)	132	1.4	2.2
13	Chronic lower respiratory diseases (J40-J47)	128	1.3	2.1
14	Influenza and pneumonia (J09-J18)	124	1.3	2.0
15	Anemias (D50-D64)	115	1.2	1.9
...	All other causes (residual)	2,125	22.3	34.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, black, female, 45-64 years				
...	All causes	39,991	100.0	726.3
1	Malignant neoplasms (C00-C97)	11,922	29.8	216.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,627	21.6	156.7
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,036	5.1	37.0
4	Diabetes mellitus (E10-E14)	1,975	4.9	35.9
5	Cerebrovascular diseases (I60-I69)	1,956	4.9	35.5
6	Chronic lower respiratory diseases (J40-J47)	1,371	3.4	24.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,174	2.9	21.3
8	Septicemia (A40-A41)	910	2.3	16.5
9	Chronic liver disease and cirrhosis (K70,K73-K74)	736	1.8	13.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	707	1.8	12.8
11	Human immunodeficiency virus (HIV) disease (B20-B24)	592	1.5	10.8
12	Influenza and pneumonia (J09-J18)	591	1.5	10.7
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	212	0.5	3.9
14	Anemias (D50-D64)	161	0.4	2.9
15	Pneumonitis due to solids and liquids (J69)	150	0.4	2.7
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	150	0.4	2.7
...	All other causes (residual)	6,721	16.8	122.1
Non-Hispanic, black, female, 65 years and over				
...	All causes	106,145	100.0	3,821.6
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	26,503	25.0	954.2
2	Malignant neoplasms (C00-C97)	20,977	19.8	755.2
3	Cerebrovascular diseases (I60-I69)	8,217	7.7	295.8
4	Alzheimer disease (G30)	6,444	6.1	232.0
5	Diabetes mellitus (E10-E14)	4,877	4.6	175.6
6	Chronic lower respiratory diseases (J40-J47)	4,052	3.8	145.9
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	3,619	3.4	130.3
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	2,598	2.4	93.5
9	Septicemia (A40-A41)	2,407	2.3	86.7
10	Influenza and pneumonia (J09-J18)	2,097	2.0	75.5
11	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,456	1.4	52.4
12	Pneumonitis due to solids and liquids (J69)	727	0.7	26.2
13	Parkinson disease (G20-G21)	599	0.6	21.6
14	Nutritional deficiencies (E40-E64)	478	0.5	17.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	474	0.4	17.1
...	All other causes (residual)	20,620	19.4	742.4

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, all ages				
...	All causes	19,198	100.0	703.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	3,472	18.1	127.2
2	Malignant neoplasms (C00-C97)	3,257	17.0	119.3
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,229	11.6	81.7
4	Diabetes mellitus (E10-E14)	1,114	5.8	40.8
5	Chronic liver disease and cirrhosis (K70,K73-K74)	1,047	5.5	38.4
6	Chronic lower respiratory diseases (J40-J47)	939	4.9	34.4
7	Cerebrovascular diseases (I60-I69)	737	3.8	27.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	602	3.1	22.1
9	Influenza and pneumonia (J09-J18)	391	2.0	14.3
10	Alzheimer disease (G30)	376	2.0	13.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	333	1.7	12.2
12	Septicemia (A40-A41)	295	1.5	10.8
13	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	266	1.4	9.7
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	190	1.0	7.0
15	Certain conditions originating in the perinatal period (P00-P96)	117	0.6	4.3
...	All other causes (residual)	3,833	20.0	140.4
Non-Hispanic, American Indian or Alaska Native, both sexes, 1-4 years				
...	All causes	72	100.0	45.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	25	34.7	15.7
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	8	11.1	*
3	Influenza and pneumonia (J09-J18)	5	6.9	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	6.9	*
5	Malignant neoplasms (C00-C97)	4	5.6	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.8	*
6	Cerebrovascular diseases (I60-I69)	2	2.8	*
8	Acute bronchitis and bronchiolitis (J20-J21)	1	1.4	*
8	Hernia (K40-K46)	1	1.4	*
8	Certain conditions originating in the perinatal period (P00-P96)	1	1.4	*
...	All other causes (residual)	18	25.0	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 5-14 years				
...	All causes	102	100.0	24.9
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	40	39.2	9.8
2	Malignant neoplasms (C00-C97)	16	15.7	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	12	11.8	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	6	5.9	*
5	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	3.9	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	3	2.9	*
7	Certain conditions originating in the perinatal period (P00-P96)	2	2.0	*
8	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.0	*
8	Diabetes mellitus (E10-E14)	1	1.0	*
8	Influenza and pneumonia (J09-J18)	1	1.0	*
8	Legal intervention (Y35,Y89.0)	1	1.0	*
...	All other causes (residual)	15	14.7	*
Non-Hispanic, American Indian or Alaska Native, both sexes, 15-24 years				
...	All causes	562	100.0	132.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	256	45.6	60.1
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	159	28.3	37.3
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	45	8.0	10.6
4	Malignant neoplasms (C00-C97)	16	2.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	10	1.8	*
6	Legal intervention (Y35,Y89.0)	5	0.9	*
7	Cerebrovascular diseases (I60-I69)	4	0.7	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	0.7	*
9	Diabetes mellitus (E10-E14)	3	0.5	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	3	0.5	*
11	Septicemia (A40-A41)	2	0.4	*
11	Influenza and pneumonia (J09-J18)	2	0.4	*
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.4	*
14	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.2	*
14	Chronic lower respiratory diseases (J40-J47)	1	0.2	*
14	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.2	*
...	All other causes (residual)	48	8.5	11.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 25-44 years				
...	All causes	2,603	100.0	351.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	880	33.8	118.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	291	11.2	39.3
3	Chronic liver disease and cirrhosis (K70,K73-K74)	274	10.5	37.0
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	195	7.5	26.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	150	5.8	20.3
6	Malignant neoplasms (C00-C97)	110	4.2	14.9
7	Diabetes mellitus (E10-E14)	89	3.4	12.0
8	Cerebrovascular diseases (I60-I69)	30	1.2	4.1
9	Septicemia (A40-A41)	28	1.1	3.8
9	Influenza and pneumonia (J09-J18)	28	1.1	3.8
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.6	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	13	0.5	*
12	Chronic lower respiratory diseases (J40-J47)	13	0.5	*
14	Legal intervention (Y35,Y89.0)	12	0.5	*
15	Pregnancy, childbirth and the puerperium (O00-O99)	11	0.4	*
15	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	11	0.4	*
...	All other causes (residual)	453	17.4	61.2
Non-Hispanic, American Indian or Alaska Native, both sexes, 45-64 years				
...	All causes	5,829	100.0	889.7
1	Malignant neoplasms (C00-C97)	1,183	20.3	180.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,017	17.4	155.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	672	11.5	102.6
4	Chronic liver disease and cirrhosis (K70,K73-K74)	589	10.1	89.9
5	Diabetes mellitus (E10-E14)	380	6.5	58.0
6	Chronic lower respiratory diseases (J40-J47)	223	3.8	34.0
7	Cerebrovascular diseases (I60-I69)	157	2.7	24.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	117	2.0	17.9
9	Influenza and pneumonia (J09-J18)	107	1.8	16.3
10	Septicemia (A40-A41)	103	1.8	15.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	93	1.6	14.2
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	52	0.9	7.9
13	Viral hepatitis (B15-B19)	51	0.9	7.8
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	48	0.8	7.3
15	Pneumonitis due to solids and liquids (J69)	27	0.5	4.1
...	All other causes (residual)	1,010	17.3	154.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent	
			of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, both sexes, 65 years and over				
...	All causes	9,746	100.0	3,247.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,235	22.9	744.8
2	Malignant neoplasms (C00-C97)	1,928	19.8	642.5
3	Chronic lower respiratory diseases (J40-J47)	702	7.2	233.9
4	Diabetes mellitus (E10-E14)	641	6.6	213.6
5	Cerebrovascular diseases (I60-I69)	544	5.6	181.3
6	Alzheimer disease (G30)	370	3.8	123.3
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	332	3.4	110.6
8	Influenza and pneumonia (J09-J18)	247	2.5	82.3
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	222	2.3	74.0
10	Chronic liver disease and cirrhosis (K70,K73-K74)	183	1.9	61.0
11	Septicemia (A40-A41)	159	1.6	53.0
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	137	1.4	45.7
13	Parkinson disease (G20-G21)	113	1.2	37.7
14	Pneumonitis due to solids and liquids (J69)	61	0.6	20.3
15	Nutritional deficiencies (E40-E64)	39	0.4	13.0
...	All other causes (residual)	1,833	18.8	610.9
Non-Hispanic, American Indian or Alaska Native, male, all ages				
...	All causes	10,502	100.0	784.4
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,033	19.4	151.8
2	Malignant neoplasms (C00-C97)	1,725	16.4	128.8
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,448	13.8	108.1
4	Diabetes mellitus (E10-E14)	624	5.9	46.6
5	Chronic liver disease and cirrhosis (K70,K73-K74)	559	5.3	41.8
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	448	4.3	33.5
7	Chronic lower respiratory diseases (J40-J47)	441	4.2	32.9
8	Cerebrovascular diseases (I60-I69)	322	3.1	24.0
9	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	199	1.9	14.9
10	Influenza and pneumonia (J09-J18)	186	1.8	13.9
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	154	1.5	11.5
12	Septicemia (A40-A41)	130	1.2	9.7
13	Alzheimer disease (G30)	120	1.1	9.0
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	97	0.9	7.2
15	Parkinson disease (G20-G21)	68	0.6	5.1
...	All other causes (residual)	1,948	18.5	145.5

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 1-4 years				
...	All causes	33	100.0	40.8
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	11	33.3	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	4	12.1	*
3	Malignant neoplasms (C00-C97)	3	9.1	*
4	Cerebrovascular diseases (I60-I69)	2	6.1	*
4	Influenza and pneumonia (J09-J18)	2	6.1	*
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	6.1	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	3.0	*
7	Acute bronchitis and bronchiolitis (J20-J21)	1	3.0	*
7	Hernia (K40-K46)	1	3.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	1	3.0	*
...	All other causes (residual)	5	15.2	*
Non-Hispanic, American Indian or Alaska Native, male, 5-14 years				
...	All causes	55	100.0	26.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	23	41.8	11.1
2	Malignant neoplasms (C00-C97)	9	16.4	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	5	9.1	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	7.3	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.6	*
6	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	1.8	*
6	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	1.8	*
6	Legal intervention (Y35,Y89.0)	1	1.8	*
...	All other causes (residual)	8	14.5	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates.]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 15-24 years				
...	All causes	392	100.0	181.5
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	177	45.2	82.0
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	116	29.6	53.7
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	38	9.7	17.6
4	Malignant neoplasms (C00-C97)	11	2.8	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	8	2.0	*
6	Legal intervention (Y35,Y89.0)	5	1.3	*
7	Diabetes mellitus (E10-E14)	2	0.5	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	0.5	*
9	Septicemia (A40-A41)	1	0.3	*
9	Influenza and pneumonia (J09-J18)	1	0.3	*
9	Chronic lower respiratory diseases (J40-J47)	1	0.3	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.3	*
...	All other causes (residual)	29	7.4	13.4
Non-Hispanic, American Indian or Alaska Native, male, 25-44 years				
...	All causes	1,650	100.0	447.7
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	582	35.3	157.9
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	214	13.0	58.1
3	Chronic liver disease and cirrhosis (K70,K73-K74)	144	8.7	39.1
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	119	7.2	32.3
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	110	6.7	29.8
6	Diabetes mellitus (E10-E14)	56	3.4	15.2
7	Malignant neoplasms (C00-C97)	46	2.8	12.5
8	Cerebrovascular diseases (I60-I69)	19	1.2	*
9	Influenza and pneumonia (J09-J18)	18	1.1	*
10	Septicemia (A40-A41)	15	0.9	*
11	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.7	*
11	Legal intervention (Y35,Y89.0)	12	0.7	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	11	0.7	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	6	0.4	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	4	0.2	*
15	Essential hypertension and hypertensive renal disease (I10,I12,I15)	4	0.2	*
15	Aortic aneurysm and dissection (I71)	4	0.2	*
15	Chronic lower respiratory diseases (J40-J47)	4	0.2	*
...	All other causes (residual)	270	16.4	73.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent	
			of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, male, 45-64 years				
...	All causes	3,435	100.0	1,101.1
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	672	19.6	215.4
2	Malignant neoplasms (C00-C97)	634	18.5	203.2
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	450	13.1	144.2
4	Chronic liver disease and cirrhosis (K70,K73-K74)	320	9.3	102.6
5	Diabetes mellitus (E10-E14)	247	7.2	79.2
6	Chronic lower respiratory diseases (J40-J47)	99	2.9	31.7
7	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	92	2.7	29.5
8	Cerebrovascular diseases (I60-I69)	82	2.4	26.3
9	Influenza and pneumonia (J09-J18)	57	1.7	18.3
10	Septicemia (A40-A41)	52	1.5	16.7
11	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	46	1.3	14.7
12	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	1.2	12.8
13	Viral hepatitis (B15-B19)	34	1.0	10.9
14	Essential hypertension and hypertensive renal disease (I10,I12,I15)	32	0.9	10.3
15	Human immunodeficiency virus (HIV) disease (B20-B24)	12	0.3	*
...	All other causes (residual)	566	16.5	181.4
Non-Hispanic, American Indian or Alaska Native, male, 65 years and over				
...	All causes	4,786	100.0	3,561.9
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,224	25.6	910.9
2	Malignant neoplasms (C00-C97)	1,022	21.4	760.6
3	Chronic lower respiratory diseases (J40-J47)	337	7.0	250.8
4	Diabetes mellitus (E10-E14)	319	6.7	237.4
5	Cerebrovascular diseases (I60-I69)	219	4.6	163.0
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	190	4.0	141.4
7	Alzheimer disease (G30)	117	2.4	87.1
8	Influenza and pneumonia (J09-J18)	107	2.2	79.6
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	96	2.0	71.4
10	Chronic liver disease and cirrhosis (K70,K73-K74)	95	2.0	70.7
11	Parkinson disease (G20-G21)	68	1.4	50.6
12	Septicemia (A40-A41)	61	1.3	45.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	61	1.3	45.4
14	Pneumonitis due to solids and liquids (J69)	37	0.8	27.5
15	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	21	0.4	15.6
...	All other causes (residual)	812	17.0	604.3

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, all ages				
...	All causes	8,696	100.0	625.5
1	Malignant neoplasms (C00-C97)	1,532	17.6	110.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,439	16.5	103.5
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	781	9.0	56.2
4	Chronic lower respiratory diseases (J40-J47)	498	5.7	35.8
5	Diabetes mellitus (E10-E14)	490	5.6	35.2
6	Chronic liver disease and cirrhosis (K70,K73-K74)	488	5.6	35.1
7	Cerebrovascular diseases (I60-I69)	415	4.8	29.9
8	Alzheimer disease (G30)	256	2.9	18.4
9	Influenza and pneumonia (J09-J18)	205	2.4	14.7
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	179	2.1	12.9
11	Septicemia (A40-A41)	165	1.9	11.9
12	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	154	1.8	11.1
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	93	1.1	6.7
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	67	0.8	4.8
15	Certain conditions originating in the perinatal period (P00-P96)	56	0.6	4.0
...	All other causes (residual)	1,878	21.6	135.1
Non-Hispanic, American Indian or Alaska Native, female, 1-4 years				
...	All causes	39	100.0	50.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	14	35.9	*
2	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	6	15.4	*
3	Influenza and pneumonia (J09-J18)	3	7.7	*
4	Malignant neoplasms (C00-C97)	1	2.6	*
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	1	2.6	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	1	2.6	*
...	All other causes (residual)	13	33.3	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 5-14 years				
...	All causes	47	100.0	23.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	17	36.2	*
2	Malignant neoplasms (C00-C97)	7	14.9	*
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	7	14.9	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	6.4	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	4.3	*
6	Diabetes mellitus (E10-E14)	1	2.1	*
6	Influenza and pneumonia (J09-J18)	1	2.1	*
6	Certain conditions originating in the perinatal period (P00-P96)	1	2.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	1	2.1	*
...	All other causes (residual)	7	14.9	*
Non-Hispanic, American Indian or Alaska Native, female, 15-24 years				
...	All causes	170	100.0	81.0
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	79	46.5	37.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	43	25.3	20.5
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	4.1	*
4	Malignant neoplasms (C00-C97)	5	2.9	*
5	Cerebrovascular diseases (I60-I69)	4	2.4	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	3	1.8	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.2	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	2	1.2	*
9	Septicemia (A40-A41)	1	0.6	*
9	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.6	*
9	Diabetes mellitus (E10-E14)	1	0.6	*
9	Influenza and pneumonia (J09-J18)	1	0.6	*
9	Chronic liver disease and cirrhosis (K70,K73-K74)	1	0.6	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.6	*
...	All other causes (residual)	19	11.2	*

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 25-44 years				
...	All causes	953	100.0	256.4
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	298	31.3	80.2
2	Chronic liver disease and cirrhosis (K70,K73-K74)	130	13.6	35.0
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	77	8.1	20.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	76	8.0	20.5
5	Malignant neoplasms (C00-C97)	64	6.7	17.2
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	40	4.2	10.8
7	Diabetes mellitus (E10-E14)	33	3.5	8.9
8	Septicemia (A40-A41)	13	1.4	*
9	Cerebrovascular diseases (I60-I69)	11	1.2	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	11	1.2	*
11	Influenza and pneumonia (J09-J18)	10	1.0	*
12	Chronic lower respiratory diseases (J40-J47)	9	0.9	*
13	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	5	0.5	*
14	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	4	0.4	*
15	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	3	0.3	*
...	All other causes (residual)	169	17.7	45.5
Non-Hispanic, American Indian or Alaska Native, female, 45-64 years				
...	All causes	2,394	100.0	697.6
1	Malignant neoplasms (C00-C97)	549	22.9	160.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	345	14.4	100.5
3	Chronic liver disease and cirrhosis (K70,K73-K74)	269	11.2	78.4
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	222	9.3	64.7
5	Diabetes mellitus (E10-E14)	133	5.6	38.8
6	Chronic lower respiratory diseases (J40-J47)	124	5.2	36.1
7	Cerebrovascular diseases (I60-I69)	75	3.1	21.9
8	Septicemia (A40-A41)	51	2.1	14.9
9	Influenza and pneumonia (J09-J18)	50	2.1	14.6
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	47	2.0	13.7
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	25	1.0	7.3
12	Viral hepatitis (B15-B19)	17	0.7	*
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	16	0.7	*
13	Pneumonitis due to solids and liquids (J69)	16	0.7	*
15	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	12	0.5	*
15	Complications of medical and surgical care (Y40-Y84,Y88)	12	0.5	*
...	All other causes (residual)	431	18.0	125.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, American Indian or Alaska Native, female, 65 years and over				
...	All causes	4,960	100.0	2,993.3
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,011	20.4	610.1
2	Malignant neoplasms (C00-C97)	906	18.3	546.8
3	Chronic lower respiratory diseases (J40-J47)	365	7.4	220.3
4	Cerebrovascular diseases (I60-I69)	325	6.6	196.1
5	Diabetes mellitus (E10-E14)	322	6.5	194.3
6	Alzheimer disease (G30)	253	5.1	152.7
7	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	142	2.9	85.7
8	Influenza and pneumonia (J09-J18)	140	2.8	84.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	126	2.5	76.0
10	Septicemia (A40-A41)	98	2.0	59.1
11	Chronic liver disease and cirrhosis (K70,K73-K74)	88	1.8	53.1
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	76	1.5	45.9
13	Parkinson disease (G20-G21)	45	0.9	27.2
14	Pneumonitis due to solids and liquids (J69)	24	0.5	14.5
14	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	24	0.5	14.5
...	All other causes (residual)	1,015	20.5	612.5
Non-Hispanic, Asian or Pacific Islander, both sexes, all ages				
...	All causes	72,598	100.0	359.8
1	Malignant neoplasms (C00-C97)	18,257	25.1	90.5
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	15,490	21.3	76.8
3	Cerebrovascular diseases (I60-I69)	5,430	7.5	26.9
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	3,256	4.5	16.1
5	Diabetes mellitus (E10-E14)	3,047	4.2	15.1
6	Alzheimer disease (G30)	2,583	3.6	12.8
7	Influenza and pneumonia (J09-J18)	2,286	3.1	11.3
8	Chronic lower respiratory diseases (J40-J47)	2,063	2.8	10.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,534	2.1	7.6
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,472	2.0	7.3
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	1,414	1.9	7.0
12	Septicemia (A40-A41)	879	1.2	4.4
13	Parkinson disease (G20-G21)	869	1.2	4.3
14	Chronic liver disease and cirrhosis (K70,K73-K74)	725	1.0	3.6
15	Pneumonitis due to solids and liquids (J69)	549	0.8	2.7
...	All other causes (residual)	12,744	17.6	63.2

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 1-4 years				
...	All causes	148	100.0	15.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	36	24.3	3.8
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	14.2	2.2
3	Malignant neoplasms (C00-C97)	20	13.5	2.1
4	Influenza and pneumonia (J09-J18)	6	4.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	2.7	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	4	2.7	*
7	Septicemia (A40-A41)	3	2.0	*
7	Anemias (D50-D64)	3	2.0	*
7	Certain conditions originating in the perinatal period (P00-P96)	3	2.0	*
10	Meningitis (G00,G03)	2	1.4	*
10	Cerebrovascular diseases (I60-I69)	2	1.4	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.7	*
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	0.7	*
12	Hernia (K40-K46)	1	0.7	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	0.7	*
...	All other causes (residual)	40	27.0	4.2
Non-Hispanic, Asian or Pacific Islander, both sexes, 5-14 years				
...	All causes	208	100.0	8.7
1	Malignant neoplasms (C00-C97)	42	20.2	1.7
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	37	17.8	1.5
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	22	10.6	0.9
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	21	10.1	0.9
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	9	4.3	*
6	Septicemia (A40-A41)	6	2.9	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	4	1.9	*
8	Anemias (D50-D64)	3	1.4	*
8	Influenza and pneumonia (J09-J18)	3	1.4	*
8	Chronic lower respiratory diseases (J40-J47)	3	1.4	*
11	Cerebrovascular diseases (I60-I69)	2	1.0	*
11	Acute bronchitis and bronchiolitis (J20-J21)	2	1.0	*
11	Certain conditions originating in the perinatal period (P00-P96)	2	1.0	*
14	Meningococcal infection (A39)	1	0.5	*
14	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.5	*
14	Diabetes mellitus (E10-E14)	1	0.5	*
14	Meningitis (G00,G03)	1	0.5	*
...	All other causes (residual)	48	23.1	2.0

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 15-24 years				
...	All causes	960	100.0	36.0
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	315	32.8	11.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	283	29.5	10.6
3	Malignant neoplasms (C00-C97)	79	8.2	3.0
4	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	59	6.1	2.2
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	37	3.9	1.4
6	Influenza and pneumonia (J09-J18)	15	1.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	1.3	*
8	Cerebrovascular diseases (I60-I69)	6	0.6	*
9	Septicemia (A40-A41)	4	0.4	*
9	Diabetes mellitus (E10-E14)	4	0.4	*
9	Pregnancy, childbirth and the puerperium (O00-O99)	4	0.4	*
12	Aortic aneurysm and dissection (I71)	3	0.3	*
12	Chronic lower respiratory diseases (J40-J47)	3	0.3	*
12	Pneumonitis due to solids and liquids (J69)	3	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	3	0.3	*
...	All other causes (residual)	130	13.5	4.9
Non-Hispanic, Asian or Pacific Islander, both sexes, 25-44 years				
...	All causes	3,727	100.0	56.1
1	Malignant neoplasms (C00-C97)	811	21.8	12.2
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	743	19.9	11.2
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	511	13.7	7.7
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	421	11.3	6.3
5	Cerebrovascular diseases (I60-I69)	162	4.3	2.4
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	142	3.8	2.1
7	Chronic liver disease and cirrhosis (K70,K73-K74)	87	2.3	1.3
8	Diabetes mellitus (E10-E14)	76	2.0	1.1
9	Influenza and pneumonia (J09-J18)	41	1.1	0.6
10	Pregnancy, childbirth and the puerperium (O00-O99)	31	0.8	0.5
11	Septicemia (A40-A41)	30	0.8	0.5
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	29	0.8	0.4
13	Essential hypertension and hypertensive renal disease (I10,I12,I15)	26	0.7	0.4
14	Human immunodeficiency virus (HIV) disease (B20-B24)	21	0.6	0.3
14	Aortic aneurysm and dissection (I71)	21	0.6	0.3
...	All other causes (residual)	575	15.4	8.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, both sexes, 45-64 years				
...	All causes	13,187	100.0	268.5
1	Malignant neoplasms (C00-C97)	5,259	39.9	107.1
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	2,470	18.7	50.3
3	Cerebrovascular diseases (I60-I69)	776	5.9	15.8
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	704	5.3	14.3
5	Diabetes mellitus (E10-E14)	624	4.7	12.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	364	2.8	7.4
7	Chronic liver disease and cirrhosis (K70,K73-K74)	284	2.2	5.8
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	223	1.7	4.5
9	Influenza and pneumonia (J09-J18)	180	1.4	3.7
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	173	1.3	3.5
11	Septicemia (A40-A41)	155	1.2	3.2
12	Chronic lower respiratory diseases (J40-J47)	153	1.2	3.1
13	Viral hepatitis (B15-B19)	92	0.7	1.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	90	0.7	1.8
15	Aortic aneurysm and dissection (I71)	83	0.6	1.7
...	All other causes (residual)	1,557	11.8	31.7
Non-Hispanic, Asian or Pacific Islander, both sexes, 65 years and over				
...	All causes	53,411	100.0	2,237.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	12,528	23.5	524.7
2	Malignant neoplasms (C00-C97)	12,042	22.5	504.4
3	Cerebrovascular diseases (I60-I69)	4,478	8.4	187.6
4	Alzheimer disease (G30)	2,567	4.8	107.5
5	Diabetes mellitus (E10-E14)	2,342	4.4	98.1
6	Influenza and pneumonia (J09-J18)	2,033	3.8	85.2
7	Chronic lower respiratory diseases (J40-J47)	1,884	3.5	78.9
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,420	2.7	59.5
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1,275	2.4	53.4
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1,271	2.4	53.2
11	Parkinson disease (G20-G21)	844	1.6	35.4
12	Septicemia (A40-A41)	674	1.3	28.2
13	Pneumonitis due to solids and liquids (J69)	506	0.9	21.2
14	Chronic liver disease and cirrhosis (K70,K73-K74)	354	0.7	14.8
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	343	0.6	14.4
...	All other causes (residual)	8,850	16.6	370.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, all ages				
...	All causes	37,236	100.0	386.2
1	Malignant neoplasms (C00-C97)	9,243	24.8	95.9
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	8,418	22.6	87.3
3	Cerebrovascular diseases (I60-I69)	2,442	6.6	25.3
4	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	2,074	5.6	21.5
5	Diabetes mellitus (E10-E14)	1,612	4.3	16.7
6	Chronic lower respiratory diseases (J40-J47)	1,198	3.2	12.4
7	Influenza and pneumonia (J09-J18)	1,155	3.1	12.0
8	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	990	2.7	10.3
9	Alzheimer disease (G30)	790	2.1	8.2
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	789	2.1	8.2
11	Essential hypertension and hypertensive renal disease (I10,I12,I15)	654	1.8	6.8
12	Parkinson disease (G20-G21)	489	1.3	5.1
13	Chronic liver disease and cirrhosis (K70,K73-K74)	468	1.3	4.9
14	Septicemia (A40-A41)	460	1.2	4.8
15	Pneumonitis due to solids and liquids (J69)	317	0.9	3.3
...	All other causes (residual)	6,137	16.5	63.6
Non-Hispanic, Asian or Pacific Islander, male, 1-4 years				
...	All causes	96	100.0	19.6
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	26	27.1	5.3
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	14	14.6	*
3	Malignant neoplasms (C00-C97)	13	13.5	*
4	Influenza and pneumonia (J09-J18)	3	3.1	*
5	Septicemia (A40-A41)	2	2.1	*
5	Meningitis (G00,G03)	2	2.1	*
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.1	*
5	Cerebrovascular diseases (I60-I69)	2	2.1	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.1	*
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	1	1.0	*
10	Certain conditions originating in the perinatal period (P00-P96)	1	1.0	*
...	All other causes (residual)	28	29.2	5.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 5-14 years				
...	All causes	114	100.0	9.4
1	Malignant neoplasms (C00-C97)	23	20.2	1.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	18	15.8	*
3	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	12	10.5	*
4	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	9.6	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	7	6.1	*
6	Influenza and pneumonia (J09-J18)	3	2.6	*
7	Septicemia (A40-A41)	2	1.8	*
7	Anemias (D50-D64)	2	1.8	*
7	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	1.8	*
7	Certain conditions originating in the perinatal period (P00-P96)	2	1.8	*
11	Meningococcal infection (A39)	1	0.9	*
11	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.9	*
11	Meningitis (G00,G03)	1	0.9	*
11	Acute bronchitis and bronchiolitis (J20-J21)	1	0.9	*
11	Chronic lower respiratory diseases (J40-J47)	1	0.9	*
...	All other causes (residual)	27	23.7	2.2
Non-Hispanic, Asian or Pacific Islander, male, 15-24 years				
...	All causes	667	100.0	49.6
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	228	34.2	16.9
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	203	30.4	15.1
3	Malignant neoplasms (C00-C97)	48	7.2	3.6
3	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	48	7.2	3.6
5	Diseases of heart (I00-I09,I11,I13,I20-I51)	22	3.3	1.6
6	Influenza and pneumonia (J09-J18)	11	1.6	*
7	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	1.3	*
8	Cerebrovascular diseases (I60-I69)	6	0.9	*
9	Septicemia (A40-A41)	3	0.4	*
9	Aortic aneurysm and dissection (I71)	3	0.4	*
11	Anemias (D50-D64)	2	0.3	*
11	Diabetes mellitus (E10-E14)	2	0.3	*
11	Pneumonitis due to solids and liquids (J69)	2	0.3	*
11	Legal intervention (Y35,Y89.0)	2	0.3	*
11	Complications of medical and surgical care (Y40-Y84,Y88)	2	0.3	*
...	All other causes (residual)	76	11.4	5.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent	
			of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 25-44 years				
...	All causes	2,387	100.0	75.2
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	591	24.8	18.6
2	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	366	15.3	11.5
3	Malignant neoplasms (C00-C97)	333	14.0	10.5
3	Diseases of heart (I00-I09,I11,I13,I20-I51)	333	14.0	10.5
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	98	4.1	3.1
6	Cerebrovascular diseases (I60-I69)	95	4.0	3.0
7	Chronic liver disease and cirrhosis (K70,K73-K74)	61	2.6	1.9
8	Diabetes mellitus (E10-E14)	46	1.9	1.4
9	Influenza and pneumonia (J09-J18)	24	1.0	0.8
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	22	0.9	0.7
11	Aortic aneurysm and dissection (I71)	19	0.8	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	18	0.8	*
13	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	15	0.6	*
14	Viral hepatitis (B15-B19)	14	0.6	*
15	Septicemia (A40-A41)	12	0.5	*
...	All other causes (residual)	340	14.2	10.7
Non-Hispanic, Asian or Pacific Islander, male, 45-64 years				
...	All causes	7,825	100.0	345.1
1	Malignant neoplasms (C00-C97)	2,530	32.3	111.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	1,827	23.3	80.6
3	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	489	6.2	21.6
4	Cerebrovascular diseases (I60-I69)	453	5.8	20.0
5	Diabetes mellitus (E10-E14)	401	5.1	17.7
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	254	3.2	11.2
7	Chronic liver disease and cirrhosis (K70,K73-K74)	218	2.8	9.6
8	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	145	1.9	6.4
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	117	1.5	5.2
10	Influenza and pneumonia (J09-J18)	107	1.4	4.7
11	Septicemia (A40-A41)	91	1.2	4.0
12	Chronic lower respiratory diseases (J40-J47)	86	1.1	3.8
13	Viral hepatitis (B15-B19)	66	0.8	2.9
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	63	0.8	2.8
15	Aortic aneurysm and dissection (I71)	58	0.7	2.6
...	All other causes (residual)	920	11.8	40.6

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, male, 65 years and over				
...	All causes	25,619	100.0	2,478.0
1	Malignant neoplasms (C00-C97)	6,296	24.6	609.0
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,222	24.3	601.8
3	Cerebrovascular diseases (I60-I69)	1,885	7.4	182.3
4	Diabetes mellitus (E10-E14)	1,163	4.5	112.5
5	Chronic lower respiratory diseases (J40-J47)	1,102	4.3	106.6
6	Influenza and pneumonia (J09-J18)	1,001	3.9	96.8
7	Alzheimer disease (G30)	783	3.1	75.7
8	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	726	2.8	70.2
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	626	2.4	60.5
10	Essential hypertension and hypertensive renal disease (I10,I12,I15)	513	2.0	49.6
11	Parkinson disease (G20-G21)	476	1.9	46.0
12	Septicemia (A40-A41)	347	1.4	33.6
13	Pneumonitis due to solids and liquids (J69)	291	1.1	28.1
14	Chronic liver disease and cirrhosis (K70,K73-K74)	189	0.7	18.3
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	185	0.7	17.9
...	All other causes (residual)	3,814	14.9	368.9
Non-Hispanic, Asian or Pacific Islander, female, all ages				
...	All causes	35,362	100.0	335.6
1	Malignant neoplasms (C00-C97)	9,014	25.5	85.6
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	7,072	20.0	67.1
3	Cerebrovascular diseases (I60-I69)	2,988	8.4	28.4
4	Alzheimer disease (G30)	1,793	5.1	17.0
5	Diabetes mellitus (E10-E14)	1,435	4.1	13.6
6	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	1,182	3.3	11.2
7	Influenza and pneumonia (J09-J18)	1,131	3.2	10.7
8	Chronic lower respiratory diseases (J40-J47)	865	2.4	8.2
9	Essential hypertension and hypertensive renal disease (I10,I12,I15)	818	2.3	7.8
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	745	2.1	7.1
11	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	424	1.2	4.0
12	Septicemia (A40-A41)	419	1.2	4.0
13	Parkinson disease (G20-G21)	380	1.1	3.6
14	Chronic liver disease and cirrhosis (K70,K73-K74)	257	0.7	2.4
15	Pneumonitis due to solids and liquids (J69)	232	0.7	2.2
...	All other causes (residual)	6,607	18.7	62.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 1-4 years				
...	All causes	52	100.0	11.3
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	10	19.2	*
2	Malignant neoplasms (C00-C97)	7	13.5	*
2	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	7	13.5	*
4	Anemias (D50-D64)	3	5.8	*
4	Influenza and pneumonia (J09-J18)	3	5.8	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	3.8	*
6	Certain conditions originating in the perinatal period (P00-P96)	2	3.8	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	3.8	*
9	Septicemia (A40-A41)	1	1.9	*
9	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	1.9	*
9	Hernia (K40-K46)	1	1.9	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	1	1.9	*
...	All other causes (residual)	12	23.1	*
Non-Hispanic, Asian or Pacific Islander, female, 5-14 years				
...	All causes	94	100.0	7.9
1	Malignant neoplasms (C00-C97)	19	20.2	*
1	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	19	20.2	*
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	11	11.7	*
4	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	9	9.6	*
5	Septicemia (A40-A41)	4	4.3	*
6	Diseases of heart (I00-I09,I11,I13,I20-I51)	2	2.1	*
6	Cerebrovascular diseases (I60-I69)	2	2.1	*
6	Chronic lower respiratory diseases (J40-J47)	2	2.1	*
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	2	2.1	*
10	Anemias (D50-D64)	1	1.1	*
10	Diabetes mellitus (E10-E14)	1	1.1	*
10	Acute bronchitis and bronchiolitis (J20-J21)	1	1.1	*
...	All other causes (residual)	21	22.3	1.8

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 15-24 years				
...	All causes	293	100.0	22.2
1	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	87	29.7	6.6
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	80	27.3	6.1
3	Malignant neoplasms (C00-C97)	31	10.6	2.4
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	15	5.1	*
5	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	11	3.8	*
6	Influenza and pneumonia (J09-J18)	4	1.4	*
6	Pregnancy, childbirth and the puerperium (O00-O99)	4	1.4	*
8	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	3	1.0	*
9	Diabetes mellitus (E10-E14)	2	0.7	*
9	Chronic lower respiratory diseases (J40-J47)	2	0.7	*
9	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	2	0.7	*
12	Septicemia (A40-A41)	1	0.3	*
12	Human immunodeficiency virus (HIV) disease (B20-B24)	1	0.3	*
12	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	1	0.3	*
12	Meningitis (G00,G03)	1	0.3	*
12	Pneumonitis due to solids and liquids (J69)	1	0.3	*
12	Certain conditions originating in the perinatal period (P00-P96)	1	0.3	*
12	Complications of medical and surgical care (Y40-Y84,Y88)	1	0.3	*
12	Enterocolitis due to <i>Clostridium difficile</i> (A04.7)	1	0.3	*
...	All other causes (residual)	44	15.0	3.3
Non-Hispanic, Asian or Pacific Islander, female, 25-44 years				
...	All causes	1,340	100.0	38.7
1	Malignant neoplasms (C00-C97)	478	35.7	13.8
2	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	152	11.3	4.4
3	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	145	10.8	4.2
4	Diseases of heart (I00-I09,I11,I13,I20-I51)	88	6.6	2.5
5	Cerebrovascular diseases (I60-I69)	67	5.0	1.9
6	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	44	3.3	1.3
7	Pregnancy, childbirth and the puerperium (O00-O99)	31	2.3	0.9
8	Diabetes mellitus (E10-E14)	30	2.2	0.9
9	Chronic liver disease and cirrhosis (K70,K73-K74)	26	1.9	0.8
10	Septicemia (A40-A41)	18	1.3	*
11	Influenza and pneumonia (J09-J18)	17	1.3	*
12	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	14	1.0	*
13	Chronic lower respiratory diseases (J40-J47)	12	0.9	*
14	Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)	10	0.7	*
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	9	0.7	*
...	All other causes (residual)	199	14.9	5.7

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017

LCWK3. Deaths, percent of total deaths, and death rates for the 15 leading causes of death in selected age groups, by race and Hispanic origin, and sex: United States, 2017

[Rates per 100,000 population in specified group. Rates for "all ages" include deaths for age "Under 1 year." Figures for "age not stated" are included in "all ages" but are not distributed among age groups. Race and Hispanic origin are reported separately on the death certificate. Persons of Hispanic origin may be of any race. Data for Hispanic persons are not tabulated by race; data for non-Hispanic persons are tabulated by race. Data for racial and ethnic groups other than non-Hispanic white and non-Hispanic black should be interpreted with caution because of misreporting of race and Hispanic origin on death certificates]

Rank ¹	Cause of death (based on the <i>International Classification of Diseases, Tenth Revision</i>), Hispanic origin, race, sex, and age	Number	Percent of total deaths	Rate
Non-Hispanic, Asian or Pacific Islander, female, 45-64 years				
...	All causes	5,362	100.0	202.8
1	Malignant neoplasms (C00-C97)	2,729	50.9	103.2
2	Diseases of heart (I00-I09,I11,I13,I20-I51)	643	12.0	24.3
3	Cerebrovascular diseases (I60-I69)	323	6.0	12.2
4	Diabetes mellitus (E10-E14)	223	4.2	8.4
5	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	215	4.0	8.1
6	Intentional self-harm (suicide) (*U03,X60-X84,Y87.0)	110	2.1	4.2
7	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	78	1.5	3.0
8	Influenza and pneumonia (J09-J18)	73	1.4	2.8
9	Chronic lower respiratory diseases (J40-J47)	67	1.2	2.5
10	Chronic liver disease and cirrhosis (K70,K73-K74)	66	1.2	2.5
11	Septicemia (A40-A41)	64	1.2	2.4
12	Essential hypertension and hypertensive renal disease (I10,I12,I15)	56	1.0	2.1
13	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	35	0.7	1.3
14	Assault (homicide) (*U01-*U02,X85-Y09,Y87.1)	27	0.5	1.0
15	Viral hepatitis (B15-B19)	26	0.5	1.0
...	All other causes (residual)	627	11.7	23.7
Non-Hispanic, Asian or Pacific Islander, female, 65 years and over				
...	All causes	27,792	100.0	2,053.2
1	Diseases of heart (I00-I09,I11,I13,I20-I51)	6,306	22.7	465.9
2	Malignant neoplasms (C00-C97)	5,746	20.7	424.5
3	Cerebrovascular diseases (I60-I69)	2,593	9.3	191.6
4	Alzheimer disease (G30)	1,784	6.4	131.8
5	Diabetes mellitus (E10-E14)	1,179	4.2	87.1
6	Influenza and pneumonia (J09-J18)	1,032	3.7	76.2
7	Chronic lower respiratory diseases (J40-J47)	782	2.8	57.8
8	Essential hypertension and hypertensive renal disease (I10,I12,I15)	758	2.7	56.0
9	Accidents (unintentional injuries) (V01-X59,Y85-Y86)	694	2.5	51.3
10	Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)	649	2.3	47.9
11	Parkinson disease (G20-G21)	368	1.3	27.2
12	Septicemia (A40-A41)	327	1.2	24.2
13	Pneumonitis due to solids and liquids (J69)	215	0.8	15.9
14	Chronic liver disease and cirrhosis (K70,K73-K74)	165	0.6	12.2
15	In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)	158	0.6	11.7
...	All other causes (residual)	5,036	18.1	372.1

... Category not applicable

* Figure does not meet standards of reliability or precision

¹ Rank based on number of deaths

SOURCE: CDC/NCHS, National Vital Statistics System, Mortality 2017