

Person with disabilities experiencing problems accessing  
NCHS birth and death tables should contact  
[nchsed@cdc.gov](mailto:nchsed@cdc.gov), or call 301-458-4688





















Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes.....	2.6	4.1	0.4	0.2	0.1	0.2	0.2	0.2	0.4	0.7
Male.....	2.9	4.8	0.5	*	*	0.2	0.2	0.3	0.5	0.9
Female.....	2.3	3.4	0.3	0.2	*	0.2	0.2	0.2	0.3	0.5
White, both sexes.....	2.7	3.7	0.4	0.2	0.2	0.2	0.2	0.2	0.3	0.6
Male.....	3.0	4.2	0.4	*	*	*	*	0.2	0.4	0.8
Female.....	2.4	3.1	*	*	*	*	*	*	*	0.4
Black, both sexes.....	2.5	6.5	*	*	*	*	*	*	1.1	1.2
Male.....	2.8	8.0	*	*	*	*	*	*	*	1.5
Female.....	2.2	*	*	*	*	*	*	*	*	*

Malignant neoplasms (C00-C97)

All races 3/, both sexes.....	188.7	1.8	2.3	2.5	2.5	3.5	4.7	6.7	11.3	20.8
Male.....	198.9	2.1	2.6	2.6	2.8	4.1	5.5	7.3	10.4	17.4
Female.....	178.8	1.5	2.0	2.3	2.2	2.8	3.9	6.0	12.2	24.3
White, both sexes.....	200.8	2.0	2.3	2.5	2.5	3.6	4.7	6.3	10.8	20.1
Male.....	210.6	2.3	2.6	2.6	2.8	4.2	5.4	7.0	10.0	17.1
Female.....	191.1	1.7	2.0	2.4	2.3	2.9	3.9	5.5	11.7	23.3
Black, both sexes.....	161.7	*	2.4	2.6	2.6	3.0	5.1	9.2	15.6	28.1
Male.....	175.4	*	2.8	2.9	3.2	3.4	6.1	9.3	14.7	21.7
Female.....	149.1	*	2.1	2.3	2.0	2.7	4.1	9.0	16.3	33.8









Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
<b>Malignant melanoma of skin (C43)</b>										
All races 3/, both sexes.....	2.8	*	*	*	*	*	0.2	0.5	0.6	0.9
Male.....	3.6	*	*	*	*	*	*	0.6	0.8	0.9
Female.....	2.0	*	*	*	*	*	*	0.3	0.5	0.8
White, both sexes.....	3.4	*	*	*	*	*	0.2	0.6	0.8	1.0
Male.....	4.4	*	*	*	*	*	*	0.7	0.9	1.1
Female.....	2.4	*	*	*	*	*	*	0.4	0.6	1.0
Black, both sexes.....	0.3	*	*	*	*	*	*	*	*	*
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	0.3	*	*	*	*	*	*	*	*	*
<b>Malignant neoplasm of breast (C50)</b>										
All races 3/, both sexes.....	14.0	*	*	*	*	*	*	0.4	1.4	3.8
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	27.3	*	*	*	*	*	*	0.8	2.8	7.7
White, both sexes.....	14.4	*	*	*	*	*	*	0.3	1.2	3.3
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	28.3	*	*	*	*	*	*	0.6	2.4	6.7
Black, both sexes.....	15.0	*	*	*	*	*	*	0.8	2.9	7.8
Male.....	0.3	*	*	*	*	*	*	*	*	*
Female.....	28.4	*	*	*	*	*	*	1.6	5.6	14.7









Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
<b>Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)</b>										
All races 3/, both sexes.....	4.4	*	0.5	1.0	0.7	0.6	0.6	0.8	1.1	1.6
Male.....	5.0	*	0.5	0.9	0.7	0.7	0.6	1.1	1.3	2.0
Female.....	3.9	*	0.6	1.0	0.6	0.4	0.5	0.5	0.8	1.2
White, both sexes.....	5.0	*	0.5	0.9	0.7	0.6	0.6	0.9	1.2	1.8
Male.....	5.6	*	0.4	0.9	0.8	0.8	0.7	1.2	1.5	2.2
Female.....	4.4	*	0.6	1.0	0.6	0.4	0.5	0.6	0.9	1.3
Black, both sexes.....	2.0	*	*	1.1	*	*	*	*	*	1.0
Male.....	2.4	*	*	*	*	*	*	*	*	*
Female.....	1.7	*	*	*	*	*	*	*	*	*
<b>Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)</b>										
All races 3/, both sexes.....	18.6	0.6	0.8	0.8	0.9	1.3	1.8	1.8	2.2	2.7
Male.....	20.6	*	1.0	0.9	1.0	1.6	2.2	2.1	2.4	3.0
Female.....	16.6	*	0.6	0.6	0.7	1.0	1.3	1.5	2.0	2.3
White, both sexes.....	20.3	0.6	0.7	0.8	0.9	1.4	1.8	1.6	2.1	2.7
Male.....	22.5	*	0.9	0.9	1.1	1.7	2.2	1.9	2.2	3.1
Female.....	18.1	*	0.5	0.7	0.7	1.0	1.4	1.3	1.9	2.3
Black, both sexes.....	12.9	*	1.0	*	0.8	1.3	1.9	2.7	3.8	3.2
Male.....	14.1	*	*	*	*	1.4	2.5	2.8	4.7	3.7
Female.....	11.8	*	*	*	*	1.3	*	2.6	3.0	2.8





Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races 3/, both sexes.....	0.0	*	*	*	*	*	*	*	*	*
Male.....	0.0	*	*	*	*	*	*	*	*	*
Female.....	0.0	*	*	*	*	*	*	*	*	*
White, both sexes.....	0.0	*	*	*	*	*	*	*	*	*
Male.....	0.0	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes.....	21.2	*	0.8	0.6	0.8	1.3	1.5	1.5	2.0	2.9
Male.....	22.1	*	0.8	0.6	0.8	1.4	1.9	1.9	2.4	3.1
Female.....	20.4	*	0.7	0.6	0.7	1.2	1.0	1.0	1.6	2.7
White, both sexes.....	22.7	*	0.8	0.6	0.7	1.4	1.5	1.4	1.9	2.9
Male.....	23.6	*	0.9	0.6	0.7	1.5	2.0	1.9	2.3	3.2
Female.....	21.9	*	0.6	0.5	0.8	1.2	1.0	0.9	1.5	2.6
Black, both sexes.....	17.1	*	0.8	0.7	1.1	0.9	1.5	2.1	2.6	3.6
Male.....	18.0	*	*	*	1.4	*	1.9	2.4	3.2	3.7
Female.....	16.3	*	*	*	*	*	1.7	2.0	3.6	

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes.....	4.6	1.4	0.3	0.2	0.2	0.2	0.2	0.3	0.5	0.7
Male.....	4.7	1.0	0.4	*	*	0.3	0.2	0.4	0.5	0.7
Female.....	4.6	1.8	0.3	0.2	*	0.2	*	0.3	0.4	0.7
White, both sexes.....	5.1	1.3	0.3	0.2	0.2	0.2	0.2	0.3	0.5	0.6
Male.....	5.2	*	0.4	*	*	0.3	0.3	0.4	0.5	0.7
Female.....	5.0	1.7	*	*	*	*	*	0.3	0.5	0.6
Black, both sexes.....	2.9	*	*	*	*	*	*	*	*	1.1
Male.....	2.7	*	*	*	*	*	*	*	*	*
Female.....	3.2	*	*	*	*	*	*	*	*	1.4

Anemias (D50-D64)

All races 3/, both sexes.....	1.6	*	0.1	*	*	0.2	0.3	0.4	0.4	0.4
Male.....	1.2	*	*	*	*	0.2	0.3	0.4	0.4	0.4
Female.....	1.9	*	*	*	*	*	0.3	0.4	0.3	0.4
White, both sexes.....	1.5	*	*	*	*	*	*	*	*	*
Male.....	1.1	*	*	*	*	*	*	*	*	*
Female.....	1.8	*	*	*	*	*	*	*	*	*
Black, both sexes.....	2.6	*	*	*	*	0.9	1.5	2.3	2.4	2.2
Male.....	2.4	*	*	*	*	*	1.7	2.4	2.7	2.7
Female.....	2.8	*	*	*	*	*	1.3	2.1	2.2	1.8









Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

All ages 1/  
Under 1 Year 2/  
1-4 years  
5-9 years  
10-14 years  
15-19 years  
20-24 years  
25-29 years  
30-34 years  
35-39 years

Alzheimer's disease (G30)

All races 3/, both sexes.....	24.2	*	*	*	*	*	*	*	*	*
Male.....	14.1	*	*	*	*	*	*	*	*	*
Female.....	33.9	*	*	*	*	*	*	*	*	*
White, both sexes.....	27.6	*	*	*	*	*	*	*	*	*
Male.....	16.0	*	*	*	*	*	*	*	*	*
Female.....	38.9	*	*	*	*	*	*	*	*	*
Black, both sexes.....	11.8	*	*	*	*	*	*	*	*	*
Male.....	6.9	*	*	*	*	*	*	*	*	*
Female.....	16.3	*	*	*	*	*	*	*	*	*

Major cardiovascular diseases (I00-I78)

All races 3/, both sexes.....	288.8	12.2	1.3	0.8	1.0	2.3	4.3	7.3	12.8	24.1
Male.....	277.9	13.4	1.5	0.8	1.1	3.0	5.5	9.5	16.6	32.0
Female.....	299.3	11.0	1.1	0.9	0.8	1.7	3.0	5.0	9.0	16.1
White, both sexes.....	307.0	9.9	1.1	0.7	0.8	2.0	3.6	5.9	10.5	20.3
Male.....	293.0	10.8	1.3	0.7	1.0	2.5	4.7	7.9	13.7	27.5
Female.....	320.8	9.0	0.9	0.7	0.7	1.5	2.5	3.7	7.1	13.0
Black, both sexes.....	256.2	23.1	2.3	1.6	1.5	4.2	8.3	16.6	29.5	52.6
Male.....	255.0	26.3	2.2	1.4	1.7	5.6	10.6	20.6	38.0	69.2
Female.....	257.2	19.7	2.3	1.7	1.4	2.6	6.1	12.9	22.0	37.8







Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Other acute ischemic heart diseases (I24)										
All races 3/, both sexes.....	1.2	*	*	*	*	*	*	*	*	0.1
Male.....	1.2	*	*	*	*	*	*	*	*	*
Female.....	1.2	*	*	*	*	*	*	*	*	*
White, both sexes.....	1.2	*	*	*	*	*	*	*	*	*
Male.....	1.2	*	*	*	*	*	*	*	*	*
Female.....	1.2	*	*	*	*	*	*	*	*	*
Black, both sexes.....	1.4	*	*	*	*	*	*	*	*	*
Male.....	1.6	*	*	*	*	*	*	*	*	*
Female.....	1.3	*	*	*	*	*	*	*	*	*
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes.....	98.2	*	*	*	*	*	0.3	0.8	2.3	5.5
Male.....	102.9	*	*	*	*	*	0.5	1.2	3.4	8.4
Female.....	93.7	*	*	*	*	*	*	0.3	1.2	2.5
White, both sexes.....	106.2	*	*	*	*	*	0.3	0.7	2.2	5.0
Male.....	110.7	*	*	*	*	*	0.4	1.1	3.1	7.6
Female.....	101.9	*	*	*	*	*	*	0.3	1.1	2.3
Black, both sexes.....	76.6	*	*	*	*	*	*	1.1	3.3	9.8
Male.....	81.0	*	*	*	*	*	*	2.0	4.9	15.3
Female.....	72.6	*	*	*	*	*	*	*	1.8	4.9



























Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
<b>Emphysema (J43)</b>										
All races 3/, both sexes.....	4.7	*	*	*	*	*	*	*	*	0.1
Male.....	5.0	*	*	*	*	*	*	*	*	*
Female.....	4.5	*	*	*	*	*	*	*	*	*
White, both sexes.....	5.4	*	*	*	*	*	*	*	*	*
Male.....	5.6	*	*	*	*	*	*	*	*	*
Female.....	5.3	*	*	*	*	*	*	*	*	*
Black, both sexes.....	2.1	*	*	*	*	*	*	*	*	*
Male.....	2.7	*	*	*	*	*	*	*	*	*
Female.....	1.5	*	*	*	*	*	*	*	*	*
<b>Asthma (J45-J46)</b>										
All races 3/, both sexes.....	1.3	*	0.2	0.2	0.3	0.2	0.4	0.5	0.6	0.7
Male.....	0.9	*	0.3	0.3	0.3	0.3	0.4	0.5	0.5	0.5
Female.....	1.7	*	*	*	*	*	0.3	0.4	0.7	0.8
White, both sexes.....	1.1	*	*	0.1	*	0.2	0.2	0.4	0.4	0.5
Male.....	0.7	*	*	*	*	*	0.2	0.4	0.3	0.4
Female.....	1.5	*	*	*	*	*	*	0.4	0.5	0.6
Black, both sexes.....	2.6	*	1.1	0.7	0.9	0.7	1.5	1.2	1.9	1.8
Male.....	2.2	*	1.6	*	1.2	*	1.6	1.5	1.9	1.7
Female.....	3.0	*	*	*	*	*	1.3	*	1.9	1.9













Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes.....	1.0	*	*	*	*	*	*	*	*	*
Male.....	0.9	*	*	*	*	*	*	*	*	*
Female.....	1.1	*	*	*	*	*	*	*	*	*
White, both sexes.....	1.1	*	*	*	*	*	*	*	*	*
Male.....	1.0	*	*	*	*	*	*	*	*	*
Female.....	1.2	*	*	*	*	*	*	*	*	*
Black, both sexes.....	0.8	*	*	*	*	*	*	*	*	*
Male.....	0.7	*	*	*	*	*	*	*	*	*
Female.....	0.9	*	*	*	*	*	*	*	*	*

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races 3/, both sexes.....	14.8	3.9	*	*	*	0.2	0.3	0.6	0.9	1.2
Male.....	14.6	4.2	*	*	*	*	0.3	0.5	1.0	1.3
Female.....	15.0	3.5	*	*	*	0.2	0.3	0.6	0.8	1.1
White, both sexes.....	14.5	3.0	*	*	*	0.1	0.2	0.4	0.6	0.9
Male.....	14.4	3.1	*	*	*	*	0.3	0.3	0.7	1.0
Female.....	14.6	2.9	*	*	*	*	*	0.4	0.5	0.8
Black, both sexes.....	20.7	8.7	*	*	*	*	*	1.6	2.7	3.5
Male.....	19.5	10.6	*	*	*	*	*	1.7	2.9	3.7
Female.....	21.7	6.8	*	*	*	*	*	1.6	2.6	3.4







Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

All ages 1/  
Under 1 Year 2/  
1-4 years  
5-9 years  
10-14 years  
15-19 years  
20-24 years  
25-29 years  
30-34 years  
35-39 years

Inflammatory diseases of female pelvic organs (N70-N76)

All races 3/, both sexes.....	0.0	*	*	*	*	*	*	*	*	*
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	0.1	*	*	*	*	*	*	*	*	*
White, both sexes.....	0.0	*	*	*	*	*	*	*	*	*
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	0.1	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	*	*	*	*	*	*	*	*	*	*

Pregnancy, childbirth and the puerperium (000-099)

All races 3/, both sexes.....	0.3	...	...	...	*	0.2	0.7	0.8	0.8	0.6
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	0.5	...	...	...	*	0.4	1.4	1.6	1.6	1.3
White, both sexes.....	0.2	...	...	...	*	0.1	0.5	0.5	0.6	0.5
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	0.4	...	...	...	*	0.3	1.1	1.0	1.3	1.0
Black, both sexes.....	0.7	...	...	...	*	0.6	1.5	2.3	1.9	1.5
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	1.3	...	...	...	*	1.2	3.1	4.5	3.6	2.8

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Pregnancy with abortive outcome (000-007)										
All races 3/, both sexes.....	0.0	...	...	...	*	*	*	*	*	*
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	0.0	...	...	...	*	*	*	*	*	*
White, both sexes.....	*	...	...	...	*	*	*	*	*	*
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	*	...	...	...	*	*	*	*	*	*
Black, both sexes.....	*	...	...	...	*	*	*	*	*	*
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	*	...	...	...	*	*	*	*	*	*
Other complications of pregnancy, childbirth and the puerperium (010-099)										
All races 3/, both sexes.....	0.2	...	...	...	*	0.2	0.7	0.7	0.8	0.6
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	0.5	...	...	...	*	0.4	1.4	1.5	1.5	1.2
White, both sexes.....	0.2	...	...	...	*	*	0.5	0.5	0.6	0.4
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	0.4	...	...	...	*	*	1.1	1.0	1.3	0.9
Black, both sexes.....	0.7	...	...	...	*	*	1.5	2.1	1.8	1.3
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	1.2	...	...	...	*	*	3.1	4.1	3.3	2.5

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Certain conditions originating in the perinatal period (P00-P96)

All races 3/, both sexes.....	4.9	351.2	0.4	*	*	*	*	*	*	*
Male.....	5.6	386.1	0.4	*	*	*	*	*	*	*
Female.....	4.2	314.6	0.3	*	*	*	*	*	*	*
White, both sexes.....	3.7	275.7	0.3	*	*	*	*	*	*	*
Male.....	4.2	303.8	0.3	*	*	*	*	*	*	*
Female.....	3.2	246.3	*	*	*	*	*	*	*	*
Black, both sexes.....	12.9	756.7	0.9	*	*	*	*	*	*	*
Male.....	15.2	829.5	*	*	*	*	*	*	*	*
Female.....	10.9	680.8	*	*	*	*	*	*	*	*

Congenital malformations, deformations and chromosomal abnormalities (Q00-Q99)

All races 3/, both sexes.....	3.5	135.2	3.2	1.0	1.0	1.2	1.2	1.2	1.0	1.1
Male.....	3.7	141.1	3.2	0.9	1.0	1.6	1.4	1.3	1.1	1.2
Female.....	3.3	129.0	3.3	1.2	0.9	0.8	1.0	1.0	0.9	1.0
White, both sexes.....	3.4	131.0	3.1	0.9	0.9	1.1	1.2	1.1	1.0	1.1
Male.....	3.6	136.9	3.1	0.8	1.0	1.5	1.4	1.2	1.1	1.2
Female.....	3.2	124.8	3.0	1.0	0.8	0.7	0.9	1.0	0.9	1.1
Black, both sexes.....	4.4	162.9	4.0	1.3	1.3	1.6	1.4	1.6	1.3	1.3
Male.....	4.8	166.5	3.6	*	1.3	2.1	1.8	2.1	*	1.8
Female.....	4.1	159.0	4.5	1.5	1.4	*	*	*	1.5	*



Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes.....	10.8	87.4	1.4	0.3	0.5	1.2	2.5	2.8	3.1	4.1
Male.....	10.0	101.8	1.6	0.3	0.6	1.6	3.4	3.9	4.0	5.1
Female.....	11.5	72.3	1.3	0.3	0.3	0.8	1.4	1.7	2.1	3.1
White, both sexes.....	11.1	72.9	1.2	0.2	0.4	1.1	2.4	2.7	3.1	4.0
Male.....	10.0	87.7	1.4	*	0.5	1.6	3.4	3.8	4.1	4.9
Female.....	12.2	57.4	1.1	*	*	0.7	1.4	1.5	2.0	3.0
Black, both sexes.....	12.3	169.0	2.5	0.8	0.9	1.9	3.1	4.2	4.2	6.0
Male.....	13.3	186.6	2.7	*	*	2.4	4.5	5.7	5.2	7.1
Female.....	11.4	150.7	2.2	*	*	1.3	1.7	2.8	3.3	5.0

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes.....	73.4	29.2	3.8	1.9	2.4	4.0	5.8	7.0	8.7	13.9
Male.....	60.9	34.8	4.3	2.0	2.5	5.0	6.5	7.9	9.9	16.1
Female.....	85.6	23.2	3.3	1.7	2.2	2.9	5.0	6.1	7.5	11.7
White, both sexes.....	78.5	25.1	3.5	1.7	2.1	3.9	5.6	6.5	7.9	12.6
Male.....	64.1	30.0	4.0	1.9	2.3	5.0	6.5	7.6	9.6	15.1
Female.....	92.6	19.9	3.1	1.5	1.9	2.7	4.6	5.5	6.2	10.0
Black, both sexes.....	63.6	50.5	5.0	2.8	3.8	4.9	8.1	11.3	15.5	24.7
Male.....	56.8	62.6	5.8	2.9	4.1	5.7	7.9	11.7	15.3	25.1
Female.....	69.7	37.9	4.1	2.7	3.4	4.2	8.2	10.9	15.6	24.3

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes.....	39.7	26.4	10.3	5.5	6.4	31.4	43.4	36.0	33.8	35.3
Male.....	52.3	27.5	12.4	6.3	8.1	43.0	66.2	54.8	49.7	50.6
Female.....	27.5	25.2	8.0	4.7	4.7	19.2	19.2	16.3	17.5	19.8
White, both sexes.....	41.8	23.0	10.0	5.1	6.3	33.9	46.9	38.5	35.3	36.8
Male.....	54.3	23.7	12.3	5.9	7.8	46.1	71.6	58.1	51.2	52.1
Female.....	29.5	22.2	7.6	4.3	4.7	21.0	20.3	17.4	18.6	20.9
Black, both sexes.....	34.9	47.4	12.5	7.5	7.6	22.8	31.8	29.4	34.9	36.1
Male.....	50.0	50.8	13.8	8.3	10.3	32.5	48.5	47.2	54.8	55.7
Female.....	21.2	43.8	11.1	6.7	4.8	12.7	14.9	12.6	16.8	18.5

Transport accidents (V01-V99,Y85)

All races 3/, both sexes.....	16.3	3.6	4.0	3.3	4.3	24.4	29.3	20.5	17.6	16.3
Male.....	23.3	3.5	4.5	3.7	5.1	31.8	44.0	31.2	26.0	23.5
Female.....	9.5	3.6	3.5	3.0	3.5	16.5	13.6	9.3	8.9	8.9
White, both sexes.....	16.9	3.1	3.9	3.3	4.3	26.4	31.1	21.2	17.9	16.6
Male.....	24.0	3.3	4.3	3.7	5.1	34.2	46.7	32.0	26.0	23.5
Female.....	10.0	2.9	3.5	3.0	3.6	18.2	14.2	9.7	9.5	9.4
Black, both sexes.....	15.0	5.7	4.5	3.6	4.3	17.1	23.6	19.2	20.0	17.9
Male.....	22.7	*	4.9	3.6	5.5	23.5	35.8	31.2	33.6	29.4
Female.....	7.9	6.8	4.2	3.5	3.0	10.6	11.2	7.9	7.8	7.7



Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes.....	0.6	*	*	*	0.2	0.3	0.6	0.5	0.7	0.7
Male.....	1.0	*	*	*	*	0.4	0.9	0.9	1.1	1.1
Female.....	0.2	*	*	*	*	0.2	*	*	0.2	0.3
White, both sexes.....	0.7	*	*	*	0.2	0.4	0.6	0.6	0.7	0.7
Male.....	1.1	*	*	*	*	0.5	1.0	1.0	1.2	1.1
Female.....	0.3	*	*	*	*	0.3	*	*	*	0.4
Black, both sexes.....	0.4	*	*	*	*	*	*	*	*	*
Male.....	0.6	*	*	*	*	*	*	*	*	*
Female.....	0.1	*	*	*	*	*	*	*	*	*

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes.....	23.4	22.8	6.3	2.2	2.1	7.1	14.2	15.5	16.2	19.1
Male.....	29.0	24.0	7.9	2.6	3.0	11.2	22.2	23.6	23.7	27.1
Female.....	18.0	21.5	4.5	1.7	1.2	2.7	5.6	7.0	8.6	10.9
White, both sexes.....	24.9	19.8	6.1	1.8	1.9	7.5	15.8	17.3	17.4	20.2
Male.....	30.3	20.4	8.0	2.2	2.7	11.9	24.8	26.1	25.3	28.6
Female.....	19.6	19.2	4.1	1.3	1.1	2.8	6.1	7.8	9.1	11.5
Black, both sexes.....	20.0	41.6	7.9	3.9	3.3	5.6	8.2	10.2	14.8	18.2
Male.....	27.3	46.1	8.9	4.6	4.8	9.1	12.7	16.0	21.3	26.3
Female.....	13.3	37.0	6.9	3.2	1.8	2.1	3.7	4.7	9.0	10.9



Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes.....	1.2	1.6	3.0	0.6	0.6	1.5	1.6	1.0	0.9	1.0
Male.....	1.9	1.3	4.1	0.8	1.0	2.6	2.8	1.7	1.5	1.7
Female.....	0.5	1.8	2.0	0.4	0.3	0.3	0.3	0.3	0.3	0.3
White, both sexes.....	1.2	1.8	3.3	0.5	0.5	1.3	1.6	1.0	0.9	0.9
Male.....	1.8	1.3	4.5	0.6	0.6	2.2	2.8	1.8	1.4	1.5
Female.....	0.5	2.2	2.1	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Black, both sexes.....	1.4	*	2.2	1.2	1.4	2.2	1.9	1.1	1.1	1.5
Male.....	2.5	*	2.7	1.6	2.6	4.2	3.3	1.9	2.1	2.7
Female.....	0.4	*	1.7	*	*	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes.....	1.1	0.8	1.3	0.7	0.4	0.3	0.5	0.6	0.5	0.7
Male.....	1.3	*	1.5	0.8	0.4	0.4	0.6	0.7	0.6	0.9
Female.....	0.9	*	1.0	0.6	0.4	0.2	0.4	0.4	0.5	0.4
White, both sexes.....	1.0	0.7	1.0	0.5	0.3	0.2	0.5	0.5	0.5	0.6
Male.....	1.2	*	1.3	0.6	0.3	0.3	0.6	0.7	0.6	0.8
Female.....	0.8	*	0.7	0.4	0.3	*	0.4	0.4	0.3	0.4
Black, both sexes.....	1.9	*	2.8	1.6	0.9	*	*	0.9	1.0	1.1
Male.....	2.4	*	2.8	1.7	*	*	*	*	*	*
Female.....	1.5	*	2.7	1.6	*	*	*	*	*	*

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
Accidental poisoning and exposure to noxious substances (X40-X49)										
All races 3/, both sexes.....	8.0	0.5	0.1	*	0.2	3.0	8.8	10.6	11.3	13.5
Male.....	10.9	*	*	*	0.2	4.5	13.4	15.5	15.7	18.0
Female.....	5.1	*	*	*	*	1.5	3.9	5.4	6.7	8.9
White, both sexes.....	8.4	*	*	*	0.2	3.6	10.4	12.1	12.5	14.6
Male.....	11.4	*	*	*	*	5.5	15.8	17.7	17.4	19.5
Female.....	5.4	*	*	*	*	1.7	4.4	6.2	7.3	9.5
Black, both sexes.....	7.6	*	*	*	*	0.7	3.0	5.6	8.8	11.6
Male.....	10.8	*	*	*	*	*	4.2	8.6	12.0	15.5
Female.....	4.8	*	*	*	*	*	1.7	2.8	5.9	8.1
Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)										
All races 3/, both sexes.....	6.3	19.5	1.5	0.6	0.7	1.4	2.0	2.3	2.4	2.6
Male.....	7.4	21.1	1.8	0.8	1.0	2.3	3.3	3.8	4.0	4.2
Female.....	5.1	17.9	1.2	0.5	0.3	0.6	0.7	0.7	0.8	0.9
White, both sexes.....	6.5	16.6	1.5	0.6	0.7	1.5	2.1	2.5	2.5	2.6
Male.....	7.7	17.8	1.9	0.8	1.0	2.4	3.4	4.1	4.2	4.4
Female.....	5.4	15.3	1.1	0.5	0.3	0.5	0.6	0.7	0.8	0.8
Black, both sexes.....	6.0	37.4	1.8	*	0.7	1.3	1.8	1.8	2.3	2.7
Male.....	7.6	41.0	1.9	*	*	2.0	2.8	2.9	3.3	4.3
Female.....	4.5	33.6	1.6	*	*	*	*	*	*	1.4

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) (\*U03,X60-X84,Y87.0)

All races 3/, both sexes.....	11.0	...	...	*	1.3	7.7	12.4	11.8	13.0	13.8
Male.....	17.7	...	...	*	1.9	12.1	20.2	19.2	20.6	21.4
Female.....	4.5	...	...	*	0.7	3.0	4.0	4.1	5.3	6.1
White, both sexes.....	12.3	...	...	*	1.3	8.3	13.1	12.5	14.3	15.4
Male.....	19.7	...	...	*	1.9	13.3	21.3	20.3	22.2	23.7
Female.....	5.0	...	...	*	0.6	3.1	4.3	4.2	6.0	6.8
Black, both sexes.....	5.1	...	...	*	1.1	4.4	9.1	8.8	8.5	7.2
Male.....	8.7	...	...	*	1.5	7.2	16.2	14.9	15.0	12.0
Female.....	1.8	...	...	*	*	1.4	2.0	3.0	2.7	2.8

Intentional self-harm (suicide) by discharge of firearms (X72-X74)

All races 3/, both sexes.....	5.7	...	...	*	0.4	3.5	5.8	5.6	5.7	5.9
Male.....	10.2	...	...	*	0.7	6.1	10.3	9.7	9.7	10.1
Female.....	1.4	...	...	*	*	0.8	1.1	1.3	1.6	1.7
White, both sexes.....	6.5	...	...	*	0.5	3.8	6.1	6.0	6.3	6.7
Male.....	11.5	...	...	*	0.8	6.6	10.7	10.3	10.5	11.2
Female.....	1.6	...	...	*	*	0.9	1.2	1.4	1.9	2.0
Black, both sexes.....	2.6	...	...	*	*	2.4	5.3	4.6	4.2	3.5
Male.....	4.9	...	...	*	*	4.3	9.8	8.4	8.3	6.7
Female.....	0.5	...	...	*	*	*	*	*	*	*



Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

All races 3/, both sexes.....	5.3	...	...	*	0.9	4.2	6.5	6.2	7.3	7.8
Male.....	7.5	...	...	*	1.2	6.0	10.0	9.5	10.9	11.3
Female.....	3.1	...	...	*	0.6	2.2	2.9	2.8	3.6	4.4
White, both sexes.....	5.8	...	...	*	0.8	4.5	7.0	6.5	8.0	8.8
Male.....	8.2	...	...	*	1.1	6.7	10.6	9.9	11.8	12.5
Female.....	3.4	...	...	*	0.5	2.2	3.2	2.8	4.0	4.8
Black, both sexes.....	2.5	...	...	*	1.0	2.0	3.9	4.1	4.3	3.7
Male.....	3.8	...	...	*	1.2	2.9	6.4	6.5	6.7	5.3
Female.....	1.3	...	...	*	*	*	1.3	1.9	2.1	2.3

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

All races 3/, both sexes.....	6.1	7.5	2.3	0.6	1.1	9.9	16.1	13.6	10.0	7.5
Male.....	9.8	8.2	2.6	0.7	1.3	16.8	27.2	22.8	16.4	11.2
Female.....	2.5	6.6	2.0	0.6	0.8	2.5	4.3	4.0	3.5	3.9
White, both sexes.....	3.7	6.1	1.7	0.4	0.6	5.3	7.9	7.2	5.6	4.7
Male.....	5.4	6.7	1.8	0.5	0.7	8.6	12.5	11.1	8.4	6.5
Female.....	1.9	5.5	1.6	0.4	0.6	1.8	2.9	3.0	2.7	2.9
Black, both sexes.....	22.2	14.3	5.3	1.4	3.0	33.5	61.2	53.0	39.2	26.5
Male.....	39.7	15.9	6.6	1.6	4.1	60.0	110.1	98.6	72.8	44.7
Female.....	6.2	12.6	4.0	1.3	1.9	6.3	11.5	9.9	8.8	10.1

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

All races 3/, both sexes.....	4.2	*	0.2	0.2	0.7	8.3	13.1	11.0	7.8	5.1
Male.....	7.2	*	0.3	0.3	1.0	14.7	23.1	19.3	13.5	8.1
Female.....	1.2	*	*	*	0.4	1.5	2.4	2.3	2.1	2.1
White, both sexes.....	2.2	*	0.2	0.2	0.4	4.1	5.5	5.1	4.0	3.0
Male.....	3.5	*	*	*	0.5	7.0	9.3	8.4	6.3	4.2
Female.....	0.9	*	*	*	*	1.0	1.4	1.6	1.5	1.7
Black, both sexes.....	17.2	*	*	*	2.1	30.0	55.1	46.9	33.5	19.4
Male.....	32.5	*	*	*	3.1	54.9	101.6	89.5	64.2	35.7
Female.....	3.1	*	*	*	*	4.4	7.6	6.6	5.8	4.7

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes.....	1.9	7.3	2.1	0.4	0.4	1.6	3.0	2.6	2.2	2.5
Male.....	2.6	7.9	2.3	0.4	0.3	2.1	4.0	3.5	3.0	3.1
Female.....	1.3	6.6	1.8	0.4	0.4	1.0	1.9	1.7	1.5	1.8
White, both sexes.....	1.5	6.0	1.5	0.3	0.2	1.2	2.4	2.1	1.7	1.7
Male.....	1.9	6.4	1.6	0.3	*	1.6	3.2	2.7	2.2	2.3
Female.....	1.0	5.5	1.4	*	0.4	0.8	1.5	1.3	1.2	1.2
Black, both sexes.....	5.0	14.0	4.8	0.9	0.9	3.5	6.2	6.1	5.7	7.1
Male.....	7.2	15.4	5.9	*	*	5.1	8.5	9.1	8.6	9.0
Female.....	3.0	12.6	3.7	*	*	1.9	3.9	3.3	3.0	5.4

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Legal intervention (Y35,Y89.0)

All races 3/, both sexes.....	0.1	*	*	*	*	0.1	0.3	0.2	0.3	0.4
Male.....	0.3	*	*	*	*	0.2	0.6	0.4	0.6	0.7
Female.....	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	0.1	*	*	*	*	*	0.3	0.2	0.2	0.3
Male.....	0.2	*	*	*	*	*	0.5	0.4	0.5	0.5
Female.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	0.3	*	*	*	*	*	*	*	1.0	0.8
Male.....	0.6	*	*	*	*	*	*	*	2.1	1.6
Female.....	*	*	*	*	*	*	*	*	*	*

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes.....	1.6	2.5	0.4	0.1	0.2	0.7	1.6	1.8	1.9	2.3
Male.....	2.0	3.2	0.5	*	0.3	1.0	2.3	2.4	2.3	2.7
Female.....	1.3	1.8	0.3	*	*	0.4	0.9	1.2	1.5	1.9
White, both sexes.....	1.6	2.1	0.3	*	0.2	0.7	1.7	2.0	2.0	2.4
Male.....	2.0	2.8	0.4	*	0.3	1.0	2.5	2.5	2.4	2.8
Female.....	1.3	1.5	*	*	*	0.4	0.9	1.4	1.6	2.0
Black, both sexes.....	1.8	4.7	0.9	*	*	0.6	1.1	1.4	1.9	2.5
Male.....	2.4	*	*	*	*	*	1.6	2.3	2.3	3.1
Female.....	1.1	*	*	*	*	*	*	*	1.5	2.0

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Discharge of firearms, undetermined intent (Y22-Y24)

All races 3/, both sexes.....	0.1	*	*	*	*	0.1	0.3	0.1	*	*
Male.....	0.1	*	*	*	*	0.2	0.4	*	*	*
Female.....	0.0	*	*	*	*	*	*	*	*	*
White, both sexes.....	0.1	*	*	*	*	*	0.3	*	*	*
Male.....	0.1	*	*	*	*	*	0.4	*	*	*
Female.....	0.0	*	*	*	*	*	*	*	*	*
Black, both sexes.....	0.1	*	*	*	*	*	*	*	*	*
Male.....	0.2	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes.....	1.5	2.5	0.4	*	0.2	0.6	1.3	1.7	1.8	2.3
Male.....	1.8	3.2	0.5	*	0.2	0.8	1.9	2.2	2.2	2.6
Female.....	1.2	1.8	0.3	*	*	0.4	0.8	1.1	1.5	1.9
White, both sexes.....	1.6	2.1	0.3	*	0.1	0.6	1.5	1.9	1.9	2.4
Male.....	1.8	2.8	0.4	*	*	0.8	2.1	2.4	2.3	2.7
Female.....	1.3	1.5	*	*	*	0.4	0.8	1.3	1.6	2.0
Black, both sexes.....	1.7	4.7	0.8	*	*	*	0.9	1.3	1.8	2.5
Male.....	2.3	*	*	*	*	*	*	1.9	2.2	3.0
Female.....	1.1	*	*	*	*	*	*	*	1.5	2.0



Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

All ages 1/ Under 1 Year 2/ 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

All races 3/, both sexes.....	7.3	*	*	*	*	0.2	0.5	1.0	1.7	4.7
Male.....	11.1	*	*	*	*	0.3	0.9	1.5	2.5	6.6
Female.....	3.6	*	*	*	*	*	*	0.4	1.0	2.7
White, both sexes.....	7.7	*	*	*	*	0.2	0.5	1.0	1.7	4.9
Male.....	11.7	*	*	*	*	0.3	0.9	1.5	2.5	6.9
Female.....	3.7	*	*	*	*	*	*	0.4	0.9	2.7
Black, both sexes.....	5.9	*	*	*	*	*	*	*	1.2	3.0
Male.....	9.1	*	*	*	*	*	*	*	1.5	4.1
Female.....	3.0	*	*	*	*	*	*	*	*	2.1

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

All races 3/, both sexes.....	11.3	0.9	0.2	*	0.2	3.7	11.0	13.5	15.1	18.3
Male.....	14.5	*	0.3	*	*	5.4	16.0	18.9	20.0	23.0
Female.....	8.2	*	*	*	0.2	2.0	5.6	7.7	10.1	13.5
White, both sexes.....	12.0	0.8	*	*	0.2	4.5	13.0	15.3	16.8	19.8
Male.....	15.3	*	*	*	*	6.5	19.0	21.5	22.2	24.8
Female.....	8.8	*	*	*	*	2.3	6.5	8.8	11.1	14.6
Black, both sexes.....	10.5	*	*	*	*	1.0	3.4	7.5	11.5	16.1
Male.....	14.3	*	*	*	*	*	4.8	11.1	14.8	20.1
Female.....	7.0	*	*	*	*	*	2.0	4.0	8.6	12.5

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race, and sex

	All ages 1/	Under 1 Year 2/	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years
--	-------------	-----------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

All races 3/, both sexes.....	10.4	*	0.4	0.3	1.3	12.5	20.0	17.2	14.2	11.6
Male.....	18.3	*	0.5	0.4	2.0	22.0	35.3	30.2	24.3	19.3
Female.....	2.7	*	0.3	0.3	0.6	2.4	3.6	3.7	3.8	3.9
White, both sexes.....	9.1	*	0.3	0.2	1.1	8.5	12.6	11.7	10.7	10.2
Male.....	15.8	*	0.3	0.3	1.7	14.6	21.8	19.7	17.6	16.3
Female.....	2.6	*	*	*	0.4	1.9	2.6	3.2	3.6	3.9
Black, both sexes.....	20.4	*	1.0	0.8	2.4	33.6	61.8	52.6	39.4	23.8
Male.....	38.7	*	*	*	3.7	61.5	114.1	100.1	75.9	44.4
Female.....	3.7	*	*	*	1.2	5.0	8.5	7.7	6.4	5.3





Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Shigellosis and amebiasis (A03,A06)

All races 3/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

Certain other intestinal infections (A04,A07-A09)

All races 3/, both sexes.....	0.1	0.1	0.4	0.5	1.2	2.8	6.5	11.6	21.5	45.8
Male.....	*	*	0.4	0.5	1.4	2.9	6.3	12.4	22.1	46.1
Female.....	*	*	0.4	0.6	1.0	2.6	6.6	11.0	21.1	45.6
White, both sexes.....	*	0.2	0.3	0.5	1.2	2.9	6.7	12.2	22.6	48.6
Male.....	*	*	0.3	0.4	1.5	3.2	6.5	13.0	22.7	49.3
Female.....	*	*	0.4	0.5	1.0	2.7	6.9	11.7	22.6	48.3
Black, both sexes.....	*	*	*	1.2	*	2.2	5.7	7.2	13.4	23.3
Male.....	*	*	*	*	*	*	*	9.4	18.3	*
Female.....	*	*	*	*	*	*	5.6	5.9	10.8	25.5











Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Measles (B05)

All races 3/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

Viral hepatitis (B15-B19)

All races 3/, both sexes.....	1.7	4.2	6.8	5.3	3.5	3.2	3.8	3.6	3.8	2.4
Male.....	2.3	6.3	10.2	8.2	4.7	3.5	4.0	3.3	4.2	2.6
Female.....	1.1	2.2	3.6	2.7	2.3	2.9	3.7	3.8	3.6	2.3
White, both sexes.....	1.7	4.3	6.7	4.9	3.0	2.7	3.2	3.3	3.6	2.2
Male.....	2.4	6.4	10.0	7.6	4.0	3.1	3.4	3.3	3.9	2.4
Female.....	1.1	2.2	3.4	2.4	2.0	2.4	3.0	3.2	3.4	2.2
Black, both sexes.....	1.7	4.7	9.7	9.2	7.2	5.8	6.4	5.5	4.9	*
Male.....	2.2	6.7	14.5	14.3	10.5	6.7	7.9	*	*	*
Female.....	*	3.0	5.7	5.0	4.7	5.1	5.4	7.5	*	*





Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Other and unspecified infectious and parasitic diseases and their sequelae (A00,A05,A20-A36,A42-A44,A48-A49,A54-A79,A81-A82,A85.0-A85.1, A85.8,A86-B04,B06-B09,B25-B49,B55-B99)

All races 3/, both sexes.....	1.1	2.4	4.2	3.9	4.5	5.9	8.3	12.0	15.8	21.2
Male.....	1.5	3.3	5.8	5.2	5.3	7.0	10.0	14.1	19.6	23.6
Female.....	0.7	1.5	2.5	2.7	3.8	5.0	7.0	10.5	13.4	20.1
White, both sexes.....	1.1	2.2	4.1	3.8	4.3	5.9	8.0	11.7	15.6	21.2
Male.....	1.4	3.1	5.9	5.0	5.0	7.0	9.8	13.8	19.4	23.7
Female.....	0.7	1.4	2.4	2.7	3.6	5.0	6.6	10.2	13.3	20.0
Black, both sexes.....	1.7	3.4	5.0	5.5	6.6	6.3	9.6	13.4	17.2	22.7
Male.....	2.5	4.3	6.1	7.8	8.4	8.3	10.8	18.8	20.4	23.2
Female.....	*	2.6	4.0	3.5	5.2	4.8	8.9	10.1	15.6	22.5

Malignant neoplasms (C00-C97)

All races 3/, both sexes.....	44.6	86.5	154.8	255.6	422.1	622.2	886.3	1,158.4	1,427.7	1,637.7
Male.....	39.5	83.4	164.9	284.8	484.5	743.6	1,091.8	1,475.1	1,901.6	2,319.3
Female.....	49.6	89.6	145.1	228.1	365.2	516.2	719.8	928.5	1,133.4	1,324.6
White, both sexes.....	43.0	82.0	147.8	247.2	414.2	618.5	888.9	1,164.9	1,432.7	1,645.4
Male.....	38.7	79.2	156.6	271.7	470.2	731.9	1,083.4	1,472.2	1,894.9	2,322.7
Female.....	47.3	84.8	139.2	223.6	362.1	517.7	728.2	937.5	1,143.0	1,333.6
Black, both sexes.....	60.6	129.1	231.4	374.3	573.3	782.6	1,026.2	1,296.6	1,585.5	1,744.7
Male.....	50.0	127.0	258.6	459.7	729.5	1,024.8	1,395.3	1,803.3	2,354.0	2,683.7
Female.....	70.0	131.0	208.4	303.9	451.6	603.0	773.5	994.4	1,179.6	1,365.8

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Malignant neoplasms of lip, oral cavity and pharynx (C00-C14)

All races 3/, both sexes.....	0.8	1.9	3.5	5.6	7.5	9.2	10.6	11.8	14.2	17.0
Male.....	1.2	2.8	5.7	9.1	11.9	14.8	16.0	17.9	20.4	22.0
Female.....	0.4	1.0	1.4	2.3	3.4	4.3	6.2	7.3	10.4	14.7
White, both sexes.....	0.7	1.7	3.1	5.1	7.1	8.7	10.5	11.7	14.2	17.3
Male.....	1.0	2.4	5.1	8.1	11.2	13.8	15.3	17.3	20.0	22.1
Female.....	0.5	0.9	1.3	2.2	3.3	4.2	6.5	7.6	10.6	15.1
Black, both sexes.....	1.1	3.6	6.6	9.2	12.6	14.1	13.7	13.0	14.3	12.6
Male.....	2.1	6.1	11.7	17.5	22.4	26.6	25.9	26.8	28.8	21.4
Female.....	*	1.5	2.2	2.4	5.0	4.8	5.4	*	*	9.0

Malignant neoplasm of esophagus (C15)

All races 3/, both sexes.....	1.1	2.3	4.9	8.4	12.6	17.5	22.4	27.6	27.3	24.9
Male.....	1.7	3.9	8.8	14.8	22.2	31.1	39.6	49.6	52.4	46.0
Female.....	0.4	0.7	1.2	2.5	3.9	5.6	8.5	11.6	11.6	15.3
White, both sexes.....	1.1	2.3	4.9	8.3	12.5	17.6	22.9	28.0	28.0	25.9
Male.....	1.8	3.9	9.0	14.7	22.0	31.2	40.4	50.6	53.7	48.5
Female.....	0.3	0.6	1.0	2.2	3.6	5.5	8.4	11.3	11.8	15.5
Black, both sexes.....	1.3	3.2	6.8	11.8	17.4	21.4	23.9	30.2	22.6	17.4
Male.....	2.1	5.1	11.4	19.8	31.3	39.7	41.4	52.7	48.5	27.0
Female.....	*	1.6	2.9	5.2	6.5	7.8	11.9	16.8	8.9	13.5

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Malignant neoplasm of stomach (C16)

All races 3/, both sexes.....	1.3	2.1	3.1	4.8	7.3	11.8	16.7	23.5	30.6	38.5
Male.....	1.6	2.5	4.3	6.6	10.4	16.4	23.8	32.5	44.3	54.2
Female.....	1.1	1.8	2.0	3.0	4.5	7.7	11.0	17.0	22.1	31.2
White, both sexes.....	1.1	1.7	2.7	4.2	6.2	10.0	14.7	20.5	27.0	34.8
Male.....	1.3	2.0	3.7	5.8	8.6	13.7	20.7	28.7	39.1	49.0
Female.....	0.9	1.5	1.8	2.6	3.9	6.7	9.7	14.5	19.5	28.2
Black, both sexes.....	2.3	4.0	5.2	9.0	14.4	23.5	30.1	48.6	57.3	74.3
Male.....	3.2	5.4	7.7	13.5	24.2	34.9	48.4	67.5	79.4	114.3
Female.....	1.5	2.8	3.1	5.3	6.8	15.0	17.5	37.3	45.6	58.1

Malignant neoplasms of colon, rectum and anus (C18-C21)

All races 3/, both sexes.....	4.0	7.4	13.4	21.4	34.2	52.7	75.8	105.7	149.2	216.9
Male.....	4.2	8.2	15.2	25.9	42.4	67.7	94.4	131.8	180.2	245.9
Female.....	3.7	6.6	11.6	17.1	26.7	39.7	60.8	86.8	130.0	203.6
White, both sexes.....	3.7	6.8	12.2	19.9	32.1	50.3	74.4	103.1	147.3	215.7
Male.....	4.1	7.6	13.9	24.1	40.0	64.6	92.9	128.0	179.5	245.3
Female.....	3.4	6.0	10.5	15.9	24.8	37.6	59.2	84.7	127.1	202.0
Black, both sexes.....	5.7	12.6	23.8	36.5	57.5	82.9	101.3	148.3	197.6	261.7
Male.....	5.7	13.7	27.3	46.0	74.2	106.7	124.9	197.1	231.8	295.6
Female.....	5.6	11.6	20.9	28.7	44.5	65.2	85.1	119.2	179.5	248.0

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Malignant neoplasms of liver and intrahepatic bile ducts (C22)

All races 3/, both sexes.....	1.2	3.7	8.5	10.2	12.9	16.6	24.3	30.2	34.4	32.5
Male.....	1.6	5.9	13.8	16.7	19.5	24.2	34.8	42.8	48.2	44.8
Female.....	0.8	1.5	3.4	4.1	6.8	9.9	15.8	21.0	25.9	26.8
White, both sexes.....	1.0	3.3	7.4	8.9	11.3	15.3	22.5	29.1	32.8	31.7
Male.....	1.3	5.2	11.6	14.2	17.1	22.0	32.2	41.5	45.8	45.0
Female.....	0.7	1.3	3.2	3.7	6.0	9.3	14.5	19.9	24.7	25.6
Black, both sexes.....	1.8	5.6	14.5	18.9	22.1	21.4	29.4	29.1	39.3	29.1
Male.....	2.5	8.9	26.8	35.0	36.7	35.9	44.6	45.1	59.7	31.6
Female.....	*	2.6	4.2	5.6	10.8	10.6	19.0	19.5	28.6	28.1

Malignant neoplasm of pancreas (C25)

All races 3/, both sexes.....	1.9	4.5	8.7	15.7	25.1	37.6	52.6	69.6	88.4	97.4
Male.....	2.2	5.2	10.6	19.6	30.6	43.9	61.6	78.7	96.5	102.8
Female.....	1.6	3.8	6.9	12.0	20.0	32.1	45.3	63.0	83.4	94.9
White, both sexes.....	1.9	4.4	8.2	15.2	24.6	37.1	52.2	68.8	87.7	96.8
Male.....	2.2	5.1	10.0	18.9	30.4	43.4	62.0	78.1	96.2	102.9
Female.....	1.5	3.6	6.4	11.7	19.3	31.5	44.0	61.9	82.4	94.0
Black, both sexes.....	2.2	6.2	13.7	22.3	33.6	49.1	66.4	86.9	105.1	111.4
Male.....	2.3	7.2	17.3	30.1	39.6	59.2	71.5	96.1	116.6	100.4
Female.....	2.2	5.3	10.7	15.8	28.9	41.6	63.0	81.3	99.0	115.9

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
<b>Malignant neoplasm of larynx (C32)</b>										
All races 3/, both sexes.....	0.2	0.7	1.3	2.4	3.9	5.8	6.7	6.8	7.3	6.3
Male.....	0.4	1.0	2.1	4.0	6.7	10.2	11.4	12.2	14.5	15.7
Female.....	*	0.4	0.6	0.8	1.3	1.9	3.0	2.8	2.8	1.9
White, both sexes.....	0.2	0.6	1.1	2.1	3.4	5.4	6.3	6.8	7.2	6.0
Male.....	0.4	0.9	1.8	3.5	5.7	9.5	10.5	12.1	14.0	14.9
Female.....	*	0.3	0.5	0.8	1.3	1.7	2.9	2.9	2.9	1.9
Black, both sexes.....	*	1.3	3.3	5.5	9.1	11.5	11.4	9.2	10.9	10.2
Male.....	*	2.1	5.0	10.3	18.5	22.1	22.1	18.3	26.0	28.8
Female.....	*	*	1.8	*	*	3.7	*	*	*	*
<b>Malignant neoplasms of trachea, bronchus and lung (C33-C34)</b>										
All races 3/, both sexes.....	8.2	20.4	40.1	73.8	142.7	218.7	308.3	372.3	380.0	302.3
Male.....	8.5	22.8	49.0	91.0	177.3	275.5	399.2	497.5	540.5	468.0
Female.....	8.0	18.1	31.7	57.6	111.2	169.1	234.7	281.3	280.3	226.2
White, both sexes.....	8.1	19.6	38.7	72.0	142.7	222.2	315.0	380.7	385.6	305.6
Male.....	8.1	21.5	46.6	86.3	173.3	275.0	401.3	503.1	543.8	469.9
Female.....	8.0	17.7	30.9	58.3	114.2	175.2	243.6	290.1	286.4	229.9
Black, both sexes.....	10.9	30.9	60.6	107.8	180.3	248.0	316.4	361.1	374.0	289.2
Male.....	12.1	38.4	79.5	156.2	262.7	363.2	469.6	544.5	597.6	495.5
Female.....	9.9	24.4	44.6	67.7	116.2	162.5	211.5	251.7	255.9	205.9

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
<b>Malignant melanoma of skin (C43)</b>										
All races 3/, both sexes.....	1.5	2.2	3.5	4.6	5.9	8.1	10.9	15.1	16.2	19.8
Male.....	1.9	2.7	4.6	6.4	7.9	11.9	16.2	23.5	26.0	34.9
Female.....	1.1	1.8	2.4	2.9	4.2	4.8	6.6	9.0	10.1	12.9
White, both sexes.....	1.8	2.6	4.1	5.3	6.8	9.3	12.3	16.8	17.6	21.4
Male.....	2.2	3.1	5.4	7.3	9.0	13.6	18.3	26.0	28.3	38.1
Female.....	1.3	2.1	2.8	3.4	4.7	5.5	7.4	10.0	10.9	13.8
Black, both sexes.....	*	*	*	*	*	*	*	*	*	5.3
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
<b>Malignant neoplasm of breast (C50)</b>										
All races 3/, both sexes.....	7.4	11.8	17.9	25.0	33.1	38.7	49.3	63.5	82.9	122.7
Male.....	*	*	0.3	0.4	0.7	1.0	1.7	1.8	2.5	2.4
Female.....	14.7	23.2	34.9	48.3	62.6	71.6	87.9	108.3	132.8	177.9
White, both sexes.....	6.6	10.4	16.4	23.4	32.0	38.1	49.2	63.6	83.6	123.4
Male.....	*	*	0.3	0.4	0.6	0.9	1.6	1.6	2.7	2.4
Female.....	13.3	20.7	32.3	45.7	61.1	71.1	88.6	109.5	134.4	179.1
Black, both sexes.....	13.6	23.1	31.8	43.6	47.7	53.8	61.5	75.5	91.5	144.3
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	25.4	43.0	58.4	79.1	83.9	91.8	101.5	117.3	139.5	201.8

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
<b>Malignant neoplasm of cervix uteri (C53)</b>										
All races 3/, both sexes.....	1.7	2.2	2.3	2.6	2.6	3.0	3.1	3.5	4.6	5.4
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	3.4	4.3	4.6	5.1	5.0	5.7	5.7	6.0	7.5	7.8
White, both sexes.....	1.6	1.9	2.2	2.3	2.3	2.7	2.8	3.0	4.0	4.8
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	3.1	3.8	4.3	4.5	4.5	5.1	5.2	5.2	6.6	6.9
Black, both sexes.....	3.0	4.0	3.3	5.1	5.3	6.5	6.2	9.4	12.4	13.1
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	5.7	7.6	6.2	9.3	9.5	11.3	10.4	14.9	18.9	18.4
<b>Malignant neoplasms of corpus uteri and uterus, part unspecified (C54-C55)</b>										
All races 3/, both sexes.....	0.4	0.9	1.8	3.7	6.4	8.7	10.8	13.1	17.4	22.1
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	0.8	1.8	3.5	7.2	12.1	16.4	19.6	22.6	28.2	32.2
White, both sexes.....	0.4	0.9	1.7	3.4	5.5	7.9	9.8	12.0	16.9	21.8
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	0.8	1.8	3.4	6.7	10.6	15.0	18.0	20.8	27.5	31.8
Black, both sexes.....	*	1.3	2.5	6.1	15.1	18.0	22.7	27.6	27.4	31.3
Male.....	...	...	...	...	...	...	...	...	...	...
Female.....	*	2.5	4.7	11.2	26.9	31.4	38.3	44.0	41.9	43.9





Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
<b>Malignant neoplasms of kidney and renal pelvis (C64-C65)</b>										
All races 3/, both sexes.....	0.9	2.0	4.1	6.4	10.8	14.4	19.9	24.3	30.3	32.6
Male.....	1.2	2.9	5.8	9.3	15.7	20.7	28.9	34.1	43.3	50.6
Female.....	0.7	1.2	2.4	3.6	6.4	8.8	12.5	17.2	22.2	24.3
White, both sexes.....	0.9	2.0	4.2	6.5	11.0	14.8	20.6	24.9	31.1	33.1
Male.....	1.2	2.8	5.9	9.5	15.7	21.3	29.6	34.8	44.2	52.0
Female.....	0.7	1.2	2.6	3.7	6.7	9.1	13.1	17.6	22.9	24.3
Black, both sexes.....	1.1	2.4	4.4	6.6	12.1	13.5	17.5	22.2	26.2	30.5
Male.....	1.9	3.6	7.1	10.0	19.6	20.0	29.1	32.2	39.3	39.0
Female.....	*	*	2.1	3.8	6.2	8.6	9.5	16.3	19.3	27.0
<b>Malignant neoplasm of bladder (C67)</b>										
All races 3/, both sexes.....	0.4	0.9	1.6	3.5	6.8	11.5	18.9	30.0	46.7	68.3
Male.....	0.5	1.2	2.2	5.3	10.8	18.7	31.2	50.4	86.3	132.1
Female.....	0.3	0.6	1.0	1.7	3.1	5.2	8.9	15.2	22.1	39.0
White, both sexes.....	0.4	0.9	1.6	3.6	7.1	12.0	19.8	31.1	48.7	70.0
Male.....	0.5	1.4	2.3	5.6	11.3	19.9	32.8	52.5	90.5	137.4
Female.....	0.3	0.5	0.9	1.6	3.2	5.1	9.1	15.3	22.4	39.0
Black, both sexes.....	*	1.0	2.0	3.7	5.9	9.7	15.7	25.1	36.2	57.5
Male.....	*	*	2.7	4.9	9.1	13.1	25.0	38.0	56.9	101.3
Female.....	*	*	*	2.7	3.4	7.2	9.3	17.3	25.2	39.8

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Malignant neoplasms of meninges, brain and other parts of central nervous system (C70-C72)

All races 3/, both sexes.....	2.7	3.7	5.6	8.3	10.7	14.0	18.0	20.0	20.4	15.8
Male.....	3.5	4.3	7.0	10.0	13.1	17.7	21.2	25.3	25.3	19.3
Female.....	1.9	3.0	4.3	6.8	8.6	10.7	15.4	16.1	17.4	14.3
White, both sexes.....	3.0	4.0	6.2	9.1	11.6	15.2	19.4	21.1	21.5	16.5
Male.....	3.9	4.6	7.6	10.7	14.0	19.3	22.5	26.3	26.5	20.2
Female.....	2.1	3.3	4.8	7.4	9.4	11.5	16.7	17.3	18.4	14.8
Black, both sexes.....	1.3	2.4	3.1	5.2	5.6	7.2	9.8	12.9	10.9	9.4
Male.....	1.5	2.9	4.5	6.6	7.5	8.1	13.3	19.2	14.7	*
Female.....	*	2.0	1.9	4.0	4.1	6.5	7.4	9.1	8.9	9.8

Malignant neoplasms of lymphoid, hematopoietic and related tissue (C81-C96)

All races 3/, both sexes.....	4.1	6.6	11.0	19.0	33.2	52.9	83.8	121.4	161.5	189.8
Male.....	4.9	7.8	13.3	23.9	40.7	68.1	109.5	161.0	214.7	262.4
Female.....	3.3	5.5	8.7	14.4	26.4	39.6	63.0	92.7	128.4	156.4
White, both sexes.....	3.9	6.3	10.5	18.7	33.3	53.6	85.6	125.4	167.1	197.3
Male.....	4.7	7.4	12.7	23.5	41.1	69.4	111.8	165.6	221.9	275.3
Female.....	3.0	5.3	8.3	14.1	26.1	39.5	64.0	95.7	132.7	161.4
Black, both sexes.....	5.8	9.2	16.0	24.9	38.7	58.5	81.2	103.1	127.4	136.8
Male.....	6.8	11.9	19.9	33.1	46.7	73.4	108.2	139.0	172.1	164.5
Female.....	5.0	6.7	12.8	18.2	32.4	47.4	62.8	81.6	103.9	125.7

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
<b>Hodgkin's disease (C81)</b>										
All races 3/, both sexes.....	0.3	0.4	0.4	0.5	0.7	1.0	1.2	2.2	2.5	2.1
Male.....	0.3	0.5	0.4	0.8	0.9	1.2	1.6	2.3	2.7	3.1
Female.....	0.2	0.4	0.3	0.3	0.5	0.8	1.0	2.1	2.3	1.6
White, both sexes.....	0.2	0.4	0.4	0.5	0.8	1.1	1.3	2.4	2.6	2.1
Male.....	0.3	0.5	0.4	0.7	1.0	1.3	1.6	2.4	2.9	3.4
Female.....	*	0.4	0.3	0.4	0.6	0.9	1.0	2.3	2.5	1.5
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
<b>Non-Hodgkin's lymphoma (C82-C85)</b>										
All races 3/, both sexes.....	1.6	2.4	4.3	7.6	12.3	19.2	31.5	46.9	64.4	75.2
Male.....	2.2	3.1	5.7	9.7	15.0	24.5	39.9	59.0	79.4	100.5
Female.....	1.1	1.7	3.0	5.7	9.9	14.5	24.7	38.1	55.2	63.5
White, both sexes.....	1.6	2.4	4.2	7.8	12.6	20.2	33.2	50.0	67.9	79.7
Male.....	2.2	3.0	5.5	9.8	15.3	25.7	41.9	62.8	83.6	107.0
Female.....	1.0	1.7	3.0	5.8	10.2	15.3	26.0	40.4	58.1	67.1
Black, both sexes.....	2.4	3.2	5.5	7.5	11.5	13.9	20.6	22.4	32.0	33.7
Male.....	2.8	4.4	8.2	10.4	14.7	17.8	29.1	25.9	37.2	40.0
Female.....	2.0	2.1	3.2	5.2	8.9	10.9	14.7	20.3	29.3	31.1

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
<b>Leukemia (C91-C95)</b>										
All races 3/, both sexes.....	1.7	2.7	4.1	6.5	11.8	19.8	31.3	45.7	61.0	78.7
Male.....	1.8	2.9	4.4	8.3	15.4	26.1	43.9	66.0	87.9	112.4
Female.....	1.6	2.5	3.7	4.9	8.5	14.4	21.1	31.0	44.3	63.3
White, both sexes.....	1.7	2.7	4.0	6.7	12.1	20.5	32.7	47.5	63.7	82.4
Male.....	1.7	2.9	4.5	8.5	15.8	27.0	45.6	67.9	91.4	118.7
Female.....	1.6	2.5	3.6	4.9	8.7	14.7	22.0	32.4	46.4	65.7
Black, both sexes.....	2.0	3.1	5.0	6.8	10.7	18.7	24.3	35.1	43.2	52.1
Male.....	2.2	3.7	5.1	8.3	14.0	24.5	34.2	52.7	66.7	62.3
Female.....	1.8	2.6	4.9	5.5	8.2	14.5	17.5	24.5	30.8	48.0
<b>Multiple myeloma and immunoproliferative neoplasms (C88,C90)</b>										
All races 3/, both sexes.....	0.4	1.1	2.2	4.3	8.4	12.8	19.6	26.5	33.4	33.6
Male.....	0.6	1.3	2.7	5.2	9.4	16.3	23.9	33.5	44.5	46.1
Female.....	0.3	0.9	1.7	3.5	7.4	9.9	16.1	21.4	26.5	27.9
White, both sexes.....	0.4	0.9	1.9	3.7	7.7	11.7	18.3	25.5	32.7	32.9
Male.....	0.5	1.0	2.3	4.4	8.9	15.2	22.5	32.2	43.9	45.8
Female.....	0.3	0.7	1.4	3.1	6.6	8.6	14.9	20.5	25.6	26.9
Black, both sexes.....	0.9	2.3	5.0	9.9	16.2	25.4	35.1	44.3	51.2	49.4
Male.....	*	3.0	6.2	12.7	17.5	30.4	42.7	58.1	67.4	61.4
Female.....	*	1.7	4.1	7.5	15.2	21.7	29.9	36.0	42.7	44.6

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Other and unspecified malignant neoplasms of lymphoid, hematopoietic and related tissue (C96)

All races 3/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

All other and unspecified malignant neoplasms (C17,C23-C24,C26-C31,C37-C41,C44-C49,C51-C52,C57-C60,C62-C63,C66,C68-C69,C73-C80,C97)

All races 3/, both sexes.....	5.3	9.9	16.6	27.4	43.5	62.7	90.1	121.9	167.4	212.9
Male.....	5.9	11.2	19.1	32.8	52.3	74.7	109.2	148.5	208.5	263.5
Female.....	4.7	8.7	14.2	22.4	35.6	52.1	74.7	102.5	141.9	189.6
White, both sexes.....	5.2	9.4	15.8	26.7	42.9	62.4	90.9	123.8	170.0	217.2
Male.....	5.8	10.4	18.3	31.7	51.4	73.8	109.4	151.6	211.5	270.3
Female.....	4.5	8.4	13.4	21.9	35.0	52.1	75.7	103.2	144.0	192.8
Black, both sexes.....	7.2	14.9	25.1	39.4	56.1	76.0	97.9	127.4	168.9	183.9
Male.....	7.7	18.1	28.3	50.5	71.6	98.6	127.5	144.4	217.0	221.2
Female.....	6.7	12.1	22.4	30.3	44.1	59.2	77.7	117.3	143.6	168.8

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

In situ neoplasms, benign neoplasms and neoplasms of uncertain or unknown behavior (D00-D48)

All races 3/, both sexes.....	1.0	1.4	2.1	3.3	5.7	9.5	16.3	27.9	44.8	76.7
Male.....	1.0	1.5	2.4	3.6	7.2	12.1	20.3	36.3	59.5	92.7
Female.....	1.0	1.3	1.8	3.1	4.3	7.3	13.0	21.8	35.6	69.4
White, both sexes.....	0.9	1.3	2.1	3.3	5.6	9.7	16.6	28.7	46.3	79.6
Male.....	0.9	1.4	2.4	3.7	7.1	12.4	21.0	37.5	61.9	97.0
Female.....	0.9	1.1	1.8	3.0	4.1	7.3	13.0	22.1	36.5	71.6
Black, both sexes.....	1.6	2.4	2.5	3.9	7.5	9.9	16.4	24.1	33.5	54.5
Male.....	1.7	2.6	2.8	4.3	9.1	10.9	18.0	28.2	42.8	54.8
Female.....	1.4	2.2	2.3	3.6	6.2	9.2	15.4	21.6	28.6	54.4

Anemias (D50-D64)

All races 3/, both sexes.....	0.5	0.6	0.6	0.8	1.2	2.0	3.5	6.3	12.8	36.6
Male.....	0.5	0.5	0.5	0.7	1.3	1.9	3.5	6.4	14.1	33.5
Female.....	0.5	0.7	0.7	0.9	1.2	2.0	3.5	6.3	12.1	38.0
White, both sexes.....	0.1	0.3	0.3	0.6	1.0	1.6	3.1	6.0	12.4	36.6
Male.....	*	0.3	0.3	0.6	1.0	1.5	3.3	5.9	13.5	33.7
Female.....	*	0.4	0.3	0.7	1.0	1.7	3.0	6.1	11.7	37.9
Black, both sexes.....	2.9	2.5	2.9	2.5	3.7	5.3	7.3	11.0	20.1	42.5
Male.....	3.1	2.4	2.6	*	*	6.7	*	13.4	23.2	37.2
Female.....	2.7	2.6	3.2	2.9	4.0	4.2	8.2	9.6	18.5	44.6

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Diabetes mellitus (E10-E14)

All races 3/, both sexes.....	5.6	9.9	17.3	29.7	47.3	71.7	104.9	151.4	211.2	312.1
Male.....	7.2	12.7	21.0	36.3	56.6	82.4	121.2	174.0	244.1	344.4
Female.....	4.1	7.2	13.8	23.4	38.9	62.3	91.6	134.9	190.8	297.2
White, both sexes.....	4.9	8.6	15.0	25.9	41.8	63.3	94.6	139.9	198.7	296.3
Male.....	6.5	11.3	18.4	32.0	51.0	74.8	111.5	164.6	235.4	336.2
Female.....	3.3	5.9	11.6	20.0	33.2	53.1	80.7	121.7	175.7	278.0
Black, both sexes.....	10.8	20.7	36.3	63.6	102.9	150.2	211.2	283.6	377.0	506.5
Male.....	12.3	25.9	42.2	78.0	118.2	161.8	236.9	310.7	391.9	464.8
Female.....	9.5	16.2	31.3	51.7	91.0	141.5	193.7	267.5	369.1	523.3

Nutritional deficiencies (E40-E64)

All races 3/, both sexes.....	0.1	0.2	0.4	0.5	0.6	1.2	2.7	4.7	10.0	30.8
Male.....	*	0.2	0.3	0.5	0.7	1.2	2.6	4.5	9.3	27.9
Female.....	*	0.2	0.4	0.4	0.5	1.2	2.8	5.0	10.4	32.1
White, both sexes.....	*	0.2	0.3	0.4	0.5	1.1	2.6	4.4	9.4	30.0
Male.....	*	*	0.3	0.5	0.6	1.1	2.5	4.1	8.6	27.2
Female.....	*	*	0.4	0.4	0.5	1.0	2.6	4.6	9.9	31.4
Black, both sexes.....	*	*	*	*	1.6	2.7	4.4	8.7	18.0	42.5
Male.....	*	*	*	*	*	*	*	*	19.0	37.2
Female.....	*	*	*	*	*	*	5.0	9.1	17.4	44.6





Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Meningitis (G00,G03)

All races 3/, both sexes.....	0.2	0.3	0.3	0.4	0.3	0.4	0.4	0.5	0.6	0.9
Male.....	0.2	0.3	0.3	0.4	0.3	*	*	*	*	*
Female.....	0.2	0.2	0.3	0.4	0.3	*	*	0.5	0.6	0.9
White, both sexes.....	0.2	0.3	0.2	0.4	0.3	0.3	0.4	0.5	0.6	0.9
Male.....	*	0.3	0.3	0.4	*	*	*	*	*	*
Female.....	*	*	*	0.3	*	*	*	*	*	0.9
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

Parkinson's disease (G20-G21)

All races 3/, both sexes.....	*	0.1	0.2	0.8	2.3	6.8	20.3	50.2	98.7	143.7
Male.....	*	*	0.3	0.9	3.2	9.6	30.0	75.0	153.6	228.6
Female.....	*	*	*	0.6	1.5	4.3	12.5	32.3	64.6	104.7
White, both sexes.....	*	0.1	0.2	0.8	2.4	7.1	21.8	53.7	104.4	152.9
Male.....	*	*	0.3	1.0	3.3	10.0	31.9	79.5	162.2	242.7
Female.....	*	*	*	0.7	1.6	4.6	13.4	34.6	68.2	111.5
Black, both sexes.....	*	*	*	*	*	4.3	10.9	22.7	50.0	56.7
Male.....	*	*	*	*	*	7.4	18.7	36.2	82.2	97.6
Female.....	*	*	*	*	*	*	5.6	14.7	33.0	40.1

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Alzheimer's disease (G30)

All races 3/, both sexes.....	*	0.1	0.3	1.1	3.5	9.7	33.3	102.5	275.5	861.6
Male.....	*	*	0.2	1.0	3.2	9.6	31.4	95.9	246.9	639.9
Female.....	*	*	0.4	1.2	3.7	9.8	34.8	107.3	293.2	963.4
White, both sexes.....	*	*	0.3	1.2	3.7	10.0	34.1	105.3	284.3	891.6
Male.....	*	*	0.2	1.1	3.5	9.8	31.9	97.5	254.3	661.6
Female.....	*	*	0.4	1.4	3.9	10.2	35.8	111.1	303.1	997.5
Black, both sexes.....	*	*	*	*	2.8	10.3	37.0	100.4	241.5	689.5
Male.....	*	*	*	*	*	11.6	36.1	103.7	222.6	518.7
Female.....	*	*	*	*	3.5	9.3	37.7	98.4	251.5	758.5

Major cardiovascular diseases (I00-I78)

All races 3/, both sexes.....	47.2	84.2	140.0	208.2	336.6	515.2	848.2	1,452.7	2,635.2	6,398.5
Male.....	64.1	116.8	199.9	294.8	462.0	679.9	1,093.3	1,820.2	3,159.0	6,615.9
Female.....	30.5	52.4	82.5	126.6	222.2	371.4	649.7	1,185.8	2,309.9	6,298.7
White, both sexes.....	41.4	73.7	122.4	187.2	309.5	483.0	814.8	1,420.7	2,623.8	6,507.8
Male.....	58.0	105.2	179.2	268.7	430.2	643.2	1,057.0	1,793.2	3,164.9	6,772.6
Female.....	24.6	42.4	66.8	108.8	197.3	340.5	614.5	1,145.0	2,284.8	6,385.8
Black, both sexes.....	93.7	172.2	295.3	422.3	640.6	882.2	1,301.4	2,041.3	3,168.4	6,016.9
Male.....	117.6	224.6	402.0	585.8	873.4	1,156.7	1,667.3	2,508.7	3,664.5	5,707.6
Female.....	72.7	126.5	205.1	287.3	459.1	678.7	1,051.0	1,762.4	2,906.3	6,141.8

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Diseases of heart (I00-I09,I11,I13,I20-I51)

All races 3/, both sexes.....	37.7	68.0	114.1	171.0	273.2	405.5	654.0	1,090.7	1,946.8	4,778.4
Male.....	53.6	98.7	168.8	249.4	385.1	549.7	869.1	1,414.1	2,428.3	5,143.4
Female.....	21.8	38.0	61.6	97.1	171.2	279.6	479.8	856.0	1,647.8	4,610.8
White, both sexes.....	34.0	61.3	102.5	156.6	255.0	385.2	634.7	1,073.2	1,946.5	4,879.3
Male.....	49.8	91.0	155.3	231.6	364.2	527.6	848.6	1,402.3	2,445.7	5,288.4
Female.....	18.0	31.8	50.8	84.4	153.5	258.6	457.9	829.6	1,633.6	4,691.0
Black, both sexes.....	69.4	128.5	224.1	326.8	494.7	662.2	958.8	1,494.7	2,282.2	4,326.7
Male.....	91.1	175.9	312.9	461.0	679.8	877.7	1,259.7	1,881.5	2,710.8	4,230.5
Female.....	50.2	87.1	149.0	216.0	350.4	502.4	753.0	1,264.0	2,055.7	4,365.6

Acute rheumatic fever and chronic rheumatic heart diseases (I00-I09)

All races 3/, both sexes.....	0.2	0.4	0.5	0.9	1.5	2.5	4.2	6.7	10.9	19.0
Male.....	0.2	0.3	0.5	0.9	1.1	1.9	3.1	4.9	8.0	14.0
Female.....	0.3	0.5	0.6	1.0	1.9	3.0	5.1	8.0	12.7	21.3
White, both sexes.....	0.2	0.3	0.4	0.8	1.4	2.5	4.4	7.0	11.5	20.4
Male.....	*	0.3	0.4	0.8	1.1	1.9	3.3	5.2	8.3	14.7
Female.....	*	0.4	0.5	0.9	1.8	3.1	5.3	8.4	13.6	23.1
Black, both sexes.....	*	*	1.3	1.2	1.8	2.5	*	4.2	*	6.1
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	5.3	*	*

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Hypertensive heart disease (I11)

All races 3/, both sexes.....	4.0	6.5	9.2	11.7	14.7	17.6	23.8	35.8	65.9	198.6
Male.....	5.4	9.1	13.0	16.5	20.3	22.5	28.5	39.3	62.2	147.5
Female.....	2.5	3.9	5.5	7.2	9.7	13.3	20.0	33.2	68.2	222.0
White, both sexes.....	2.8	4.6	6.5	8.9	11.7	13.5	19.4	30.4	58.0	188.9
Male.....	4.0	6.9	9.6	12.8	16.4	17.7	22.9	33.6	54.7	136.9
Female.....	1.5	2.3	3.6	5.2	7.3	9.7	16.5	28.0	60.1	212.9
Black, both sexes.....	12.2	20.8	30.3	37.8	46.0	57.8	69.9	100.9	168.5	338.1
Male.....	16.3	27.1	42.3	53.5	64.9	74.2	91.1	118.9	169.2	301.2
Female.....	8.6	15.3	20.2	24.9	31.3	45.7	55.4	90.1	168.0	353.0

Hypertensive heart and renal disease (I13)

All races 3/, both sexes.....	0.2	0.3	0.6	0.6	1.0	1.8	2.9	5.0	9.5	25.3
Male.....	0.3	0.5	0.7	0.8	1.2	2.1	2.9	5.3	9.8	27.4
Female.....	0.2	*	0.4	0.5	0.8	1.6	2.8	4.8	9.3	24.3
White, both sexes.....	*	0.1	0.2	0.3	0.5	1.1	2.0	3.9	7.9	23.9
Male.....	*	*	0.3	0.4	0.5	1.1	2.1	4.2	8.2	25.5
Female.....	*	*	*	*	0.4	1.0	1.9	3.7	7.8	23.2
Black, both sexes.....	1.2	1.7	3.4	3.7	5.7	8.7	10.7	16.5	28.4	42.8
Male.....	*	2.8	4.0	4.7	8.0	11.6	10.4	17.0	33.0	50.2
Female.....	*	*	2.9	2.9	3.8	6.5	10.8	16.3	26.0	39.8

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Ischemic heart diseases (I20-I25)

All races 3/, both sexes.....	21.6	42.9	78.3	120.9	198.6	295.4	472.3	778.6	1,359.8	3,158.1
Male.....	33.0	66.2	121.3	183.8	291.3	415.2	653.7	1,054.3	1,780.2	3,576.8
Female.....	10.4	20.2	37.0	61.5	114.1	190.9	325.4	578.5	1,098.7	2,965.8
White, both sexes.....	21.0	41.1	73.8	114.2	189.8	286.3	463.7	771.1	1,364.5	3,227.0
Male.....	32.7	64.1	116.2	175.7	281.5	406.5	646.8	1,053.1	1,798.9	3,687.1
Female.....	9.2	18.1	32.1	54.9	104.5	179.4	312.3	562.4	1,092.2	3,015.2
Black, both sexes.....	30.2	65.0	129.0	202.1	320.9	431.0	638.8	1,004.8	1,517.3	2,804.4
Male.....	41.2	95.4	187.9	294.0	454.6	584.9	857.4	1,296.8	1,887.0	2,785.0
Female.....	20.5	38.6	79.1	126.3	216.6	316.9	489.1	830.7	1,322.0	2,812.2

Acute myocardial infarction (I21-I22)

All races 3/, both sexes.....	8.7	17.3	30.9	47.8	78.9	115.0	176.6	272.1	448.1	923.1
Male.....	13.1	26.5	47.2	72.2	114.0	159.3	240.5	361.8	567.1	1,045.0
Female.....	4.4	8.4	15.2	24.7	47.0	76.4	124.8	206.9	374.1	867.1
White, both sexes.....	8.6	17.0	30.3	46.3	78.0	113.8	175.8	270.1	449.4	936.8
Male.....	13.1	26.3	47.2	70.7	114.2	160.0	241.8	363.3	572.8	1,070.1
Female.....	3.9	7.7	13.7	22.9	44.3	72.6	121.3	201.2	372.1	875.4
Black, both sexes.....	11.8	24.2	43.3	69.9	107.8	150.2	217.0	345.4	499.1	892.1
Male.....	15.8	34.4	59.5	100.2	145.2	190.8	278.3	425.6	598.3	885.9
Female.....	8.4	15.3	29.6	44.9	78.6	120.0	175.1	297.6	446.6	894.6

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Other acute ischemic heart diseases (I24)										
All races 3/, both sexes.....	0.3	0.6	1.1	1.4	2.1	2.4	4.2	5.4	9.5	20.8
Male.....	0.4	0.8	1.6	2.1	3.2	3.4	5.3	7.1	10.6	20.2
Female.....	*	0.4	0.6	0.7	1.2	1.6	3.3	4.2	8.8	21.1
White, both sexes.....	0.2	0.5	0.9	1.2	1.9	2.3	3.9	5.1	9.2	20.5
Male.....	0.3	0.7	1.4	1.8	2.8	3.2	4.9	6.6	10.3	19.7
Female.....	*	0.2	0.4	0.6	1.0	1.6	3.1	4.0	8.5	20.8
Black, both sexes.....	*	1.3	2.8	3.2	4.5	4.1	7.3	9.7	14.8	27.3
Male.....	*	*	3.6	5.0	6.9	6.4	9.8	14.8	17.6	27.9
Female.....	*	*	2.2	*	*	*	5.6	6.7	13.4	27.0
Other forms of chronic ischemic heart disease (I20,I25)										
All races 3/, both sexes.....	12.7	25.1	46.3	71.7	117.5	178.0	291.6	501.1	902.3	2,214.2
Male.....	19.5	39.0	72.5	109.5	174.1	252.6	407.9	685.4	1,202.5	2,511.6
Female.....	5.9	11.5	21.2	36.1	65.9	112.9	197.3	367.3	715.8	2,077.6
White, both sexes.....	12.2	23.6	42.6	66.7	109.9	170.2	283.9	495.8	905.8	2,269.8
Male.....	19.2	37.1	67.6	103.3	164.5	243.3	400.0	683.2	1,215.8	2,597.3
Female.....	5.1	10.2	18.0	31.4	59.2	105.2	187.9	357.2	711.6	2,119.0
Black, both sexes.....	17.9	39.5	82.8	129.0	208.5	276.7	414.4	649.7	1,003.4	1,885.0
Male.....	24.7	59.5	124.8	188.8	302.5	387.6	569.3	856.5	1,271.1	1,871.2
Female.....	11.9	22.1	47.3	79.7	135.3	194.4	308.4	526.4	862.1	1,890.6

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Atherosclerotic cardiovascular disease, so described (I25.0)

All races 3/, both sexes.....	4.4	9.6	18.6	26.9	37.7	46.8	64.6	96.5	161.9	376.4
Male.....	7.0	15.3	30.0	42.5	57.7	68.3	91.5	127.0	204.6	390.7
Female.....	1.9	4.0	7.7	12.2	19.4	28.1	42.9	74.3	135.4	369.9
White, both sexes.....	4.3	9.0	16.6	23.8	33.3	42.5	58.7	89.9	154.9	373.8
Male.....	7.0	14.6	27.2	38.2	51.2	62.2	82.8	118.8	194.8	392.6
Female.....	1.5	3.4	6.2	9.9	16.6	25.1	38.9	68.5	129.9	365.1
Black, both sexes.....	6.5	15.5	37.3	56.6	83.4	92.4	130.4	183.1	265.1	469.6
Male.....	8.9	23.1	58.1	87.6	130.3	140.2	200.8	251.7	361.7	450.8
Female.....	4.3	8.9	19.7	31.1	46.8	56.9	82.2	142.1	214.0	477.1

All other forms of chronic ischemic heart disease (I20,I25.1-I25.9)

All races 3/, both sexes.....	8.2	15.5	27.7	44.9	79.9	131.2	226.9	404.6	740.4	1,837.8
Male.....	12.5	23.7	42.4	67.0	116.4	184.3	316.4	558.3	997.9	2,120.9
Female.....	4.0	7.4	13.5	23.9	46.5	84.8	154.5	293.0	580.4	1,707.7
White, both sexes.....	8.0	14.6	26.0	42.9	76.7	127.6	225.2	405.9	750.9	1,896.0
Male.....	12.3	22.5	40.4	65.0	113.3	181.1	317.3	564.5	1,021.0	2,204.7
Female.....	3.6	6.8	11.8	21.5	42.6	80.1	149.1	288.6	581.7	1,753.8
Black, both sexes.....	11.4	24.0	45.5	72.4	125.2	184.3	284.0	466.7	738.4	1,415.4
Male.....	15.8	36.4	66.7	101.3	172.1	247.4	368.4	604.8	909.5	1,420.4
Female.....	7.6	13.2	27.6	48.5	88.6	137.5	226.2	384.3	648.0	1,413.4





Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Diseases of pericardium and acute myocarditis (I30-I31,I40)

All races 3/, both sexes.....	0.2	0.3	0.3	0.3	0.4	0.7	0.8	1.1	1.3	1.9
Male.....	0.3	0.4	0.4	0.3	0.5	0.7	1.0	1.4	1.5	1.9
Female.....	*	0.3	0.3	0.3	0.4	0.6	0.6	0.9	1.1	1.9
White, both sexes.....	0.2	0.3	0.3	0.3	0.4	0.7	0.8	1.1	1.2	1.9
Male.....	0.2	0.3	0.4	0.3	0.4	0.7	1.0	1.3	1.5	1.9
Female.....	*	*	0.3	*	0.4	0.6	0.6	0.9	1.1	1.8
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

Heart failure (I50)

All races 3/, both sexes.....	1.0	1.7	3.5	6.2	11.8	21.5	43.6	88.8	190.5	618.6
Male.....	1.3	2.1	4.3	7.6	14.0	24.6	50.1	102.1	213.7	603.1
Female.....	0.6	1.4	2.7	4.9	9.8	18.7	38.4	79.1	176.1	625.6
White, both sexes.....	0.7	1.2	2.7	5.3	10.5	19.8	42.3	87.6	192.4	640.6
Male.....	0.8	1.5	3.4	6.5	12.5	22.7	48.4	100.6	217.9	629.3
Female.....	0.5	1.0	2.1	4.2	8.6	17.3	37.2	78.0	176.5	645.8
Black, both sexes.....	3.0	5.2	10.0	15.6	26.8	41.3	69.1	123.8	216.0	489.6
Male.....	4.6	6.4	12.6	20.1	32.7	48.5	82.5	152.0	224.0	423.0
Female.....	1.6	4.0	7.7	11.9	22.1	36.0	60.0	106.9	211.8	516.5

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

All other forms of heart disease (I26-I28,I34-I38,I42-I49,I51)

All races 3/, both sexes.....	10.3	15.5	21.2	29.7	44.5	65.0	104.9	172.3	306.2	754.0
Male.....	12.9	19.8	27.9	38.7	55.8	81.5	128.0	203.6	349.3	767.7
Female.....	7.6	11.4	14.8	21.3	34.1	50.5	86.2	149.6	279.5	747.6
White, both sexes.....	8.9	13.4	18.2	26.4	40.1	60.3	100.7	169.7	308.2	773.4
Male.....	11.5	17.3	24.4	34.5	51.0	75.8	122.3	201.1	352.7	787.7
Female.....	6.3	9.4	12.0	18.6	30.0	46.6	82.9	146.5	280.4	766.9
Black, both sexes.....	21.7	33.7	48.4	63.6	91.0	118.1	165.9	239.4	341.0	640.9
Male.....	26.2	42.0	62.5	84.2	114.3	153.5	215.1	288.8	384.1	661.9
Female.....	17.6	26.4	36.4	46.5	72.8	91.8	132.2	209.9	318.3	632.4

Essential (primary) hypertension and hypertensive renal disease (I10,I12)

All races 3/, both sexes.....	1.3	2.2	3.3	5.0	8.3	13.9	22.2	40.5	75.4	210.0
Male.....	1.5	2.8	4.4	6.4	9.9	15.9	23.3	41.9	74.5	169.3
Female.....	1.0	1.7	2.3	3.8	6.8	12.2	21.4	39.4	76.0	228.7
White, both sexes.....	0.8	1.4	2.1	3.5	6.2	10.6	17.6	35.0	69.5	200.3
Male.....	1.0	1.8	2.8	4.5	7.5	12.2	18.1	35.3	68.9	160.2
Female.....	0.5	0.9	1.4	2.6	5.1	9.2	17.3	34.7	69.9	218.8
Black, both sexes.....	4.6	8.5	12.8	18.4	28.6	44.6	66.4	100.7	153.2	341.0
Male.....	5.2	9.8	17.4	24.4	36.5	52.8	79.1	123.8	154.5	289.1
Female.....	4.0	7.4	9.0	13.4	22.4	38.6	57.8	86.9	152.5	362.0

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

#### Cerebrovascular diseases (I60-I69)

All races 3/, both sexes.....	6.9	11.7	18.8	25.8	42.7	73.8	133.6	253.6	497.3	1,141.8
Male.....	7.0	12.1	21.5	30.1	50.0	84.6	149.5	275.8	513.0	1,023.3
Female.....	6.7	11.3	16.2	21.8	36.0	64.3	120.7	237.6	487.5	1,196.1
White, both sexes.....	5.3	9.1	14.5	21.1	36.2	65.6	123.8	244.2	491.2	1,153.3
Male.....	5.5	9.5	16.5	24.2	41.8	74.3	138.0	266.4	505.3	1,033.7
Female.....	5.2	8.6	12.5	18.2	31.1	57.8	112.1	227.7	482.4	1,208.5
Black, both sexes.....	17.1	30.5	50.3	65.4	99.5	145.1	231.4	372.3	611.0	1,106.7
Male.....	17.4	32.0	60.5	83.6	133.4	184.9	276.7	407.2	655.2	983.5
Female.....	16.8	29.3	41.8	50.3	73.0	115.6	200.4	351.5	587.6	1,156.5

#### Atherosclerosis (I70)

All races 3/, both sexes.....	0.1	0.3	0.7	1.0	2.1	3.9	7.2	16.9	37.8	132.9
Male.....	*	0.4	1.0	1.2	2.7	4.8	8.7	20.0	42.0	121.7
Female.....	*	0.2	0.4	0.8	1.5	3.1	5.9	14.7	35.2	138.1
White, both sexes.....	0.1	0.3	0.6	0.9	2.0	3.8	7.1	17.2	38.8	138.7
Male.....	*	0.4	0.9	1.2	2.6	4.8	8.5	20.6	42.9	128.3
Female.....	*	*	0.3	0.7	1.5	3.0	5.9	14.7	36.2	143.5
Black, both sexes.....	*	*	1.3	1.3	3.1	5.8	9.5	16.5	33.7	97.8
Male.....	*	*	*	*	4.4	7.6	12.3	17.0	40.0	75.3
Female.....	*	*	*	*	*	4.4	7.6	16.3	30.4	106.9

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Other diseases of circulatory system (I71-I78)

All races 3/, both sexes.....	1.3	2.0	3.2	5.4	10.2	18.0	31.2	50.9	77.9	135.5
Male.....	1.8	2.9	4.4	7.8	14.2	24.8	42.7	68.4	101.2	158.2
Female.....	0.9	1.1	2.0	3.2	6.6	12.1	21.9	38.2	63.4	125.0
White, both sexes.....	1.2	1.7	2.8	5.0	10.0	17.7	31.6	51.1	77.9	136.1
Male.....	1.5	2.4	3.8	7.2	14.1	24.3	43.9	68.5	102.1	162.0
Female.....	0.8	0.9	1.8	2.9	6.2	11.8	21.3	38.3	62.7	124.2
Black, both sexes.....	2.6	4.3	6.7	10.4	14.7	24.5	35.2	57.0	88.4	144.6
Male.....	3.7	6.7	9.5	15.1	19.3	33.7	39.5	79.1	103.9	129.2
Female.....	1.5	2.3	4.4	6.5	11.2	17.6	32.2	43.7	80.1	150.8

Aortic aneurysm and dissection (I71)

All races 3/, both sexes.....	0.9	1.4	2.2	3.3	6.8	11.7	19.8	31.6	46.6	64.9
Male.....	1.5	2.3	3.3	5.3	10.0	17.1	29.0	43.8	65.4	91.0
Female.....	0.4	0.6	1.1	1.5	3.8	7.0	12.4	22.7	35.0	52.9
White, both sexes.....	0.8	1.2	1.8	3.2	6.7	11.7	20.6	32.5	47.6	66.2
Male.....	1.2	1.9	2.8	5.1	10.0	17.2	30.6	45.1	67.1	94.2
Female.....	0.3	0.4	0.9	1.4	3.5	6.9	12.3	23.1	35.3	53.3
Black, both sexes.....	1.8	3.3	4.9	5.6	8.8	13.3	15.9	25.6	38.6	50.2
Male.....	3.1	5.3	7.1	8.8	12.0	18.8	17.1	34.0	47.1	51.1
Female.....	*	1.6	3.1	3.0	6.4	9.2	15.2	20.5	34.1	49.9

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Other diseases of arteries, arterioles and capillaries (I72-I78)

All races 3/, both sexes.....	0.4	0.5	1.0	2.1	3.5	6.3	11.4	19.3	31.3	70.6
Male.....	0.3	0.6	1.1	2.5	4.2	7.7	13.7	24.6	35.8	67.2
Female.....	0.5	0.5	0.9	1.7	2.8	5.2	9.6	15.5	28.4	72.2
White, both sexes.....	0.4	0.5	1.0	1.8	3.3	6.0	11.0	18.6	30.3	69.9
Male.....	0.3	0.5	1.0	2.1	4.1	7.1	13.3	23.4	35.0	67.8
Female.....	0.5	0.5	0.9	1.5	2.6	4.9	9.0	15.1	27.4	70.8
Black, both sexes.....	0.7	1.0	1.8	4.7	5.9	11.2	19.3	31.4	49.8	94.3
Male.....	*	*	2.4	6.2	7.3	15.0	22.5	45.1	56.9	78.1
Female.....	*	*	*	3.4	4.8	8.5	17.1	23.2	46.0	100.9

Other disorders of circulatory system (I80-I99)

All races 3/, both sexes.....	0.9	1.3	1.6	1.9	2.4	3.3	4.4	7.0	10.8	23.0
Male.....	1.0	1.6	2.1	2.3	3.0	4.0	5.0	6.6	11.1	19.2
Female.....	0.9	1.0	1.2	1.5	1.8	2.7	4.0	7.2	10.7	24.8
White, both sexes.....	0.7	1.2	1.4	1.7	2.1	3.2	4.2	6.7	10.5	23.5
Male.....	0.8	1.4	1.9	2.1	2.6	3.7	4.6	6.4	10.5	19.1
Female.....	0.7	1.0	1.0	1.4	1.6	2.7	3.9	6.9	10.5	25.5
Black, both sexes.....	2.3	2.6	3.5	4.0	5.2	5.5	7.6	11.5	17.5	24.3
Male.....	2.4	3.6	4.1	4.8	6.9	7.8	10.1	12.1	22.5	25.1
Female.....	2.2	1.8	3.0	3.3	3.8	3.7	5.8	11.2	14.8	24.0











Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
<b>Emphysema (J43)</b>										
All races 3/, both sexes.....	0.2	0.8	1.4	3.7	8.0	15.3	25.8	37.8	46.0	51.9
Male.....	0.3	1.1	1.6	4.4	9.7	18.4	30.7	47.5	59.5	71.5
Female.....	*	0.5	1.1	3.0	6.5	12.6	21.9	30.8	37.6	42.8
White, both sexes.....	0.2	0.8	1.3	3.9	8.6	16.4	27.9	39.9	48.2	54.5
Male.....	0.3	1.1	1.5	4.5	10.1	19.2	32.4	49.3	60.6	73.6
Female.....	*	0.5	1.2	3.3	7.2	13.8	24.2	33.0	40.5	45.7
Black, both sexes.....	*	0.9	1.9	3.5	6.0	11.6	13.6	26.2	26.2	27.3
Male.....	*	*	2.8	5.6	9.8	17.8	20.2	39.8	45.6	51.1
Female.....	*	*	*	*	3.0	7.1	9.1	18.1	16.0	17.6
<b>Asthma (J45-J46)</b>										
All races 3/, both sexes.....	1.0	1.4	1.4	1.5	1.9	2.3	2.9	4.1	7.1	14.5
Male.....	0.8	0.9	1.2	1.1	1.2	1.7	1.9	3.7	4.8	8.7
Female.....	1.2	1.8	1.7	1.9	2.6	2.8	3.7	4.4	8.6	17.2
White, both sexes.....	0.8	1.0	1.1	1.1	1.6	1.8	2.3	3.8	6.4	14.5
Male.....	0.6	0.6	1.0	0.8	0.9	1.3	1.4	3.1	4.1	8.6
Female.....	0.9	1.3	1.2	1.4	2.2	2.2	3.0	4.3	7.9	17.2
Black, both sexes.....	2.8	4.4	3.9	5.4	5.3	6.0	8.7	7.9	12.6	11.8
Male.....	2.5	3.3	2.5	4.3	*	4.8	6.3	9.8	*	*
Female.....	3.1	5.3	5.1	6.4	6.7	6.9	10.4	6.7	13.0	13.5



Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Pneumonitis due to solids and liquids (J69)

All races 3/, both sexes.....	0.4	0.8	1.5	2.0	3.7	6.4	12.9	28.5	61.1	163.9
Male.....	0.4	0.9	1.8	2.4	4.9	8.4	17.5	38.3	86.2	229.3
Female.....	0.4	0.7	1.1	1.6	2.7	4.6	9.1	21.3	45.5	133.9
White, both sexes.....	0.5	0.8	1.4	2.0	3.5	6.0	12.7	27.9	61.4	166.5
Male.....	0.5	0.9	1.7	2.4	4.5	7.8	17.1	37.1	86.1	234.2
Female.....	0.4	0.7	1.1	1.6	2.6	4.5	9.0	21.1	45.9	135.3
Black, both sexes.....	*	1.2	2.4	3.1	6.4	10.4	17.0	40.9	69.2	153.1
Male.....	*	1.5	3.4	3.6	9.8	15.4	25.6	62.1	106.0	201.7
Female.....	*	*	*	2.7	3.7	6.7	11.0	28.3	49.7	133.5

Other diseases of respiratory system (J00-J06,J30-J39,J67,J70-J98)

All races 3/, both sexes.....	1.4	2.4	4.1	6.6	11.8	21.5	37.5	59.4	89.7	136.5
Male.....	1.4	2.7	4.4	7.3	14.1	25.9	46.0	74.5	113.9	157.7
Female.....	1.4	2.1	3.8	5.9	9.8	17.7	30.6	48.5	74.6	126.7
White, both sexes.....	1.3	2.1	3.7	6.2	11.6	21.6	38.4	60.7	92.8	140.2
Male.....	1.2	2.4	4.0	6.8	13.6	26.3	47.2	75.5	117.0	160.0
Female.....	1.3	1.8	3.3	5.7	9.7	17.4	31.1	49.8	77.5	131.1
Black, both sexes.....	2.8	4.8	7.7	9.9	15.3	24.7	34.4	52.6	60.9	102.6
Male.....	2.8	5.3	8.4	12.1	20.0	28.3	40.8	70.2	82.2	132.0
Female.....	2.8	4.4	7.2	8.1	11.6	22.0	30.1	42.1	49.7	90.8



Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Hernia (K40-K46)

All races 3/, both sexes.....	0.1	0.1	0.3	0.4	0.9	0.9	1.4	2.5	4.8	12.1
Male.....	*	*	0.3	0.4	0.9	0.7	1.3	2.4	4.6	12.7
Female.....	*	*	0.2	0.4	0.8	1.1	1.4	2.5	5.0	11.8
White, both sexes.....	*	0.1	0.3	0.4	0.9	0.9	1.4	2.5	5.0	12.5
Male.....	*	*	0.3	0.4	0.9	0.7	1.4	2.4	4.6	12.8
Female.....	*	*	*	0.4	0.8	1.1	1.4	2.6	5.2	12.3
Black, both sexes.....	*	*	*	*	*	*	*	*	*	9.1
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	7.9

Chronic liver disease and cirrhosis (K70,K73-K74)

All races 3/, both sexes.....	8.6	15.0	20.7	23.0	24.1	25.4	29.3	29.8	27.9	19.7
Male.....	11.5	21.3	31.3	34.5	35.3	33.9	38.1	37.2	35.1	27.6
Female.....	5.7	8.9	10.6	12.2	13.9	17.9	22.2	24.4	23.4	16.0
White, both sexes.....	9.0	15.4	21.0	23.5	24.2	26.5	30.9	30.8	28.8	20.2
Male.....	12.1	21.8	31.9	35.0	35.2	35.0	39.8	38.1	35.9	28.5
Female.....	5.9	9.1	10.3	12.4	13.9	19.0	23.5	25.5	24.3	16.4
Black, both sexes.....	6.2	14.2	21.6	24.6	25.4	20.0	20.4	19.9	18.0	12.3
Male.....	8.1	20.8	31.7	38.3	40.5	30.9	31.6	32.6	23.2	*
Female.....	4.5	8.5	13.1	13.2	13.6	12.0	12.8	12.3	15.2	10.9

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
<b>Alcoholic liver disease (K70)</b>										
All races 3/, both sexes.....	5.7	9.3	12.0	12.4	12.0	10.3	9.4	6.4	4.8	1.8
Male.....	7.5	13.2	18.5	19.5	19.4	16.5	15.0	11.2	8.7	4.2
Female.....	3.9	5.4	5.8	5.7	5.3	4.9	4.8	3.0	2.4	0.7
White, both sexes.....	5.9	9.6	12.5	12.9	12.1	10.7	9.7	6.6	4.9	1.9
Male.....	7.8	13.7	19.2	20.3	19.5	17.0	15.4	11.3	8.9	4.4
Female.....	3.9	5.6	6.0	5.8	5.2	5.1	5.0	3.1	2.4	0.8
Black, both sexes.....	3.9	7.8	10.1	11.0	11.9	9.0	8.4	5.3	*	*
Male.....	4.6	11.7	15.0	16.8	20.5	15.4	15.2	12.1	*	*
Female.....	3.3	4.3	5.9	6.2	5.1	4.2	*	*	*	*
<b>Other chronic liver disease and cirrhosis (K73-K74)</b>										
All races 3/, both sexes.....	2.9	5.7	8.7	10.6	12.0	15.0	20.0	23.3	23.1	17.8
Male.....	4.0	8.1	12.8	14.9	15.9	17.4	23.1	26.0	26.5	23.3
Female.....	1.9	3.5	4.7	6.5	8.5	13.0	17.4	21.4	21.0	15.3
White, both sexes.....	3.1	5.8	8.5	10.6	12.0	15.8	21.2	24.2	23.8	18.3
Male.....	4.2	8.1	12.7	14.7	15.7	18.0	24.5	26.7	26.9	24.2
Female.....	2.0	3.5	4.4	6.6	8.7	13.9	18.5	22.3	21.9	15.6
Black, both sexes.....	2.3	6.5	11.5	13.6	13.5	11.0	12.1	14.5	14.6	10.7
Male.....	3.5	9.1	16.7	21.6	20.0	15.4	16.4	20.6	19.0	*
Female.....	*	4.2	7.1	7.1	8.5	7.8	9.1	10.9	12.2	10.1

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Cholelithiasis and other disorders of gallbladder (K80-K82)

All races 3/, both sexes.....	0.1	0.2	0.4	0.5	0.9	1.7	3.3	5.8	11.1	22.9
Male.....	*	*	0.5	0.5	1.2	2.3	4.3	7.1	12.7	24.3
Female.....	*	0.2	0.3	0.6	0.7	1.2	2.4	4.8	10.1	22.3
White, both sexes.....	*	0.1	0.3	0.5	0.9	1.5	3.1	5.7	11.1	23.1
Male.....	*	*	0.4	0.5	1.1	2.1	4.1	7.1	13.1	24.8
Female.....	*	*	0.2	0.5	0.7	0.9	2.3	4.7	9.8	22.4
Black, both sexes.....	*	*	*	*	*	3.6	4.6	7.0	12.4	20.6
Male.....	*	*	*	*	*	*	6.3	9.8	*	19.5
Female.....	*	*	*	*	*	*	*	5.3	13.4	21.0

Nephritis, nephrotic syndrome and nephrosis (N00-N07,N17-N19,N25-N27)

All races 3/, both sexes.....	2.1	3.8	5.8	10.6	17.7	29.2	51.3	85.6	142.8	288.3
Male.....	2.5	4.8	7.2	12.4	21.1	31.4	59.1	102.7	183.8	376.5
Female.....	1.7	2.9	4.5	8.8	14.7	27.3	44.9	73.2	117.2	247.8
White, both sexes.....	1.4	2.7	4.3	7.7	13.8	23.9	44.3	78.7	135.3	280.3
Male.....	1.7	3.4	5.5	8.9	16.8	27.1	52.7	95.9	178.1	373.7
Female.....	1.1	2.0	3.2	6.5	11.0	21.0	37.4	66.0	108.5	237.3
Black, both sexes.....	7.3	12.1	18.2	35.8	55.7	79.4	124.8	175.9	254.9	424.4
Male.....	8.5	15.9	22.6	45.8	68.3	79.4	133.5	208.8	303.4	482.4
Female.....	6.3	8.8	14.5	27.5	45.9	79.5	118.8	156.3	229.2	401.0















Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Symptoms, signs and abnormal clinical and laboratory findings, not elsewhere classified (R00-R99)

All races 3/, both sexes.....	5.2	6.4	7.0	6.9	7.7	10.4	15.7	27.6	54.0	224.4
Male.....	6.4	8.2	9.3	9.2	9.6	12.8	18.8	31.0	54.8	183.9
Female.....	4.1	4.6	4.8	4.7	6.0	8.3	13.2	25.1	53.5	243.0
White, both sexes.....	5.0	5.9	6.6	6.3	7.3	9.7	15.0	27.2	53.7	228.9
Male.....	6.3	7.8	8.7	8.3	8.9	11.8	17.9	30.4	53.6	185.6
Female.....	3.8	4.1	4.5	4.3	5.8	7.9	12.6	24.9	53.7	248.8
Black, both sexes.....	8.0	11.2	12.4	13.9	14.2	19.2	26.2	38.8	70.9	215.1
Male.....	9.2	13.8	17.2	19.5	20.0	26.4	33.5	49.2	85.7	199.9
Female.....	6.9	8.9	8.4	9.3	9.6	13.9	21.2	32.5	63.1	221.3

All other diseases (Residual) (D65-E07,E15-E34,E65-F99,G04-G12,G23-G25,G31-H93,K00-K22,K29-K31,K50-K66,K71-K72,K75-K76,K83-M99,N13.0-N13.5,N13.7-N13.9,N14,N15.0,N15.8-N15.9,N20-N23,N28-N39,N41-N64,N80-N98)

All races 3/, both sexes.....	22.8	33.6	47.8	60.3	81.9	115.6	182.4	313.2	593.5	1,559.1
Male.....	27.6	40.7	58.5	71.9	93.1	125.3	194.2	327.1	595.6	1,305.3
Female.....	17.9	26.7	37.6	49.4	71.7	107.2	172.9	303.1	592.2	1,675.6
White, both sexes.....	21.4	31.1	44.7	57.4	78.8	112.6	179.6	311.2	596.4	1,586.5
Male.....	26.8	38.6	55.5	68.5	89.5	120.8	190.8	322.7	594.7	1,328.2
Female.....	15.9	23.7	34.2	46.7	68.9	105.3	170.4	302.7	597.5	1,705.4
Black, both sexes.....	36.6	58.1	80.1	98.8	128.6	164.8	247.6	401.0	680.4	1,501.2
Male.....	39.8	65.7	94.2	119.4	151.1	194.4	274.5	447.9	743.7	1,241.0
Female.....	33.8	51.4	68.1	81.8	111.1	142.8	229.2	373.1	646.9	1,606.2

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Accidents (unintentional injuries) (V01-X59,Y85-Y86)

All races 3/, both sexes.....	41.6	44.6	41.6	35.5	36.0	40.1	53.7	82.7	137.0	279.5
Male.....	58.7	63.5	60.7	51.3	51.4	54.6	72.6	110.0	180.8	344.2
Female.....	24.6	26.0	23.3	20.7	22.0	27.4	38.5	62.8	109.7	249.8
White, both sexes.....	43.2	45.4	40.7	34.7	35.3	39.8	53.8	83.1	140.0	290.9
Male.....	60.6	63.8	58.6	49.2	49.2	53.7	71.7	110.0	182.9	356.6
Female.....	25.6	26.9	23.0	20.8	22.4	27.5	39.0	63.2	113.1	260.7
Black, both sexes.....	41.1	49.5	57.2	49.7	45.6	47.5	58.9	87.4	110.9	174.8
Male.....	60.1	75.7	90.9	80.3	79.6	73.2	90.5	130.5	167.1	225.0
Female.....	24.3	26.7	28.7	24.5	19.1	28.4	37.2	61.6	81.2	154.5

Transport accidents (V01-V99,Y85)

All races 3/, both sexes.....	17.0	16.8	16.6	16.2	16.3	17.2	18.9	23.0	26.0	26.1
Male.....	24.8	24.9	25.0	23.7	23.2	23.9	27.0	32.5	38.0	46.0
Female.....	9.2	8.9	8.6	9.0	10.0	11.3	12.4	16.1	18.5	17.0
White, both sexes.....	17.5	17.2	16.6	16.4	16.2	17.4	19.0	23.0	26.8	27.1
Male.....	25.3	25.3	24.8	23.8	23.0	23.7	26.9	32.1	39.0	47.5
Female.....	9.5	9.1	8.5	9.2	10.0	11.7	12.5	16.2	19.2	17.7
Black, both sexes.....	16.8	16.9	18.9	16.9	17.3	17.2	18.8	22.4	16.3	16.6
Male.....	26.2	25.5	30.8	27.4	27.8	28.8	30.4	38.9	27.4	33.5
Female.....	8.5	9.4	8.8	8.3	9.1	8.6	10.8	12.5	10.4	9.8





Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Water, air and space, and other and unspecified transport accidents and their sequelae (V90-V99,Y85)

All races 3/, both sexes.....	0.8	1.0	1.0	1.1	1.0	1.0	0.7	0.7	0.7	0.5
Male.....	1.4	1.7	1.8	1.9	1.7	1.7	1.3	1.3	1.2	*
Female.....	0.3	0.3	0.3	0.3	0.4	0.4	*	*	*	*
White, both sexes.....	0.9	1.0	1.1	1.2	1.1	1.1	0.8	0.8	0.7	0.5
Male.....	1.6	1.8	1.8	2.1	1.9	1.8	1.4	1.4	1.2	*
Female.....	0.3	0.3	0.4	0.3	0.4	0.4	*	*	*	*
Black, both sexes.....	*	1.1	*	*	*	*	*	*	*	*
Male.....	*	1.6	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

Nontransport accidents (W00-X59,Y86)

All races 3/, both sexes.....	24.6	27.8	25.0	19.4	19.8	22.9	34.8	59.7	111.0	253.4
Male.....	33.9	38.7	35.7	27.5	28.3	30.7	45.6	77.5	142.8	298.3
Female.....	15.3	17.1	14.7	11.7	12.0	16.0	26.1	46.7	91.3	232.8
White, both sexes.....	25.7	28.1	24.1	18.3	19.1	22.4	34.8	60.2	113.2	263.9
Male.....	35.2	38.5	33.8	25.4	26.2	29.9	44.8	77.9	143.9	309.1
Female.....	16.1	17.8	14.5	11.6	12.5	15.8	26.5	47.1	94.0	243.0
Black, both sexes.....	24.3	32.6	38.3	32.8	28.3	30.3	40.1	65.0	94.7	158.2
Male.....	34.0	50.2	60.1	52.9	51.8	44.4	60.1	91.6	139.8	191.5
Female.....	15.8	17.3	19.8	16.2	10.1	19.7	26.4	49.1	70.8	144.8



Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
Accidental drowning and submersion (W65-W74)										
All races 3/, both sexes.....	1.3	1.2	1.1	0.9	0.9	0.9	1.2	1.4	1.4	1.8
Male.....	2.0	2.0	1.7	1.5	1.5	1.4	1.8	2.1	2.5	3.9
Female.....	0.5	0.5	0.4	0.3	0.3	0.5	0.7	0.8	0.7	0.9
White, both sexes.....	1.3	1.2	1.0	0.9	0.9	0.9	1.2	1.3	1.2	1.7
Male.....	2.0	2.0	1.6	1.5	1.4	1.3	1.7	2.1	2.3	3.9
Female.....	0.6	0.5	0.5	0.3	0.4	0.4	0.7	0.7	*	0.8
Black, both sexes.....	1.1	1.2	1.2	*	*	*	*	*	*	*
Male.....	1.9	2.2	2.7	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Accidental exposure to smoke, fire and flames (X00-X09)										
All races 3/, both sexes.....	0.9	1.2	1.1	1.1	1.6	1.9	2.3	3.1	4.2	5.0
Male.....	1.2	1.5	1.6	1.5	2.0	2.5	2.8	3.9	5.6	7.0
Female.....	0.5	0.9	0.7	0.8	1.2	1.4	1.9	2.5	3.4	4.0
White, both sexes.....	0.8	1.2	1.0	0.8	1.4	1.7	2.1	2.7	3.8	4.2
Male.....	1.1	1.5	1.4	1.1	1.7	2.1	2.6	3.4	5.1	6.0
Female.....	0.6	0.9	0.7	0.6	1.2	1.3	1.8	2.2	2.9	3.4
Black, both sexes.....	1.2	1.7	2.2	3.7	3.2	4.4	4.2	8.0	11.2	14.4
Male.....	1.9	2.3	3.8	5.4	5.3	6.7	*	12.1	15.5	22.3
Female.....	*	*	*	2.3	*	*	*	5.6	8.9	11.3

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Accidental poisoning and exposure to noxious substances (X40-X49)

All races 3/, both sexes.....	17.1	18.2	14.4	8.3	4.4	2.6	2.0	2.3	2.6	3.5
Male.....	22.1	24.0	19.1	10.9	5.7	3.0	2.3	3.2	2.8	4.6
Female.....	12.0	12.6	10.0	5.8	3.2	2.3	1.7	1.7	2.4	3.0
White, both sexes.....	18.0	18.7	13.8	7.6	4.1	2.4	1.9	2.3	2.5	3.6
Male.....	23.3	24.2	17.9	9.4	4.9	2.3	2.0	3.1	2.6	4.8
Female.....	12.6	13.3	9.9	5.9	3.4	2.4	1.8	1.7	2.5	3.0
Black, both sexes.....	17.0	20.7	23.1	16.3	8.1	5.0	3.7	*	*	*
Male.....	22.0	30.3	34.6	26.9	15.1	9.3	6.6	*	*	*
Female.....	12.6	12.4	13.4	7.6	*	*	*	*	*	*

Other and unspecified nontransport accidents and their sequelae (W20-W31,W35-W64,W75-W99,X10-X39,X50-X59,Y86)

All races 3/, both sexes.....	3.5	4.4	4.9	5.0	6.8	8.4	12.3	20.7	38.9	95.1
Male.....	5.6	7.1	7.8	7.4	10.2	11.5	16.6	26.8	49.0	107.1
Female.....	1.5	1.8	2.1	2.7	3.6	5.7	8.7	16.3	32.6	89.6
White, both sexes.....	3.5	4.3	4.7	4.8	6.5	8.2	11.7	20.3	38.8	97.4
Male.....	5.6	6.8	7.4	7.0	9.5	11.4	15.5	25.9	48.6	110.0
Female.....	1.4	1.7	1.9	2.6	3.7	5.3	8.6	16.1	32.6	91.6
Black, both sexes.....	3.9	6.4	7.6	7.6	10.1	12.4	19.3	31.9	46.6	86.1
Male.....	6.0	10.8	12.1	12.1	18.9	15.4	29.1	48.3	63.2	97.6
Female.....	2.0	2.6	3.8	3.9	3.3	10.2	12.6	22.1	37.8	81.4



Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Intentional self-harm (suicide) by other and unspecified means and their sequelae (\*U03,X60-X71,X75-X84,Y87.0)

All races 3/, both sexes.....	9.0	8.7	7.8	6.2	5.1	3.7	3.6	3.5	5.2	5.7
Male.....	12.8	11.8	10.4	8.3	6.7	5.1	5.2	5.3	8.1	11.2
Female.....	5.1	5.7	5.3	4.3	3.6	2.4	2.2	2.2	3.4	3.1
White, both sexes.....	9.9	9.7	8.6	6.8	5.5	3.8	3.8	3.6	5.3	5.6
Male.....	14.0	13.0	11.4	8.9	7.2	5.3	5.6	5.5	8.4	11.4
Female.....	5.8	6.5	5.9	4.8	3.9	2.5	2.3	2.2	3.4	3.0
Black, both sexes.....	4.3	3.4	3.6	2.5	2.0	*	*	*	*	*
Male.....	6.9	5.1	4.9	3.8	*	*	*	*	*	*
Female.....	2.0	2.0	2.4	*	*	*	*	*	*	*

Assault (homicide) (\*U01-\*U02,X85-Y09,Y87.1)

All races 3/, both sexes.....	6.7	5.4	4.2	2.9	2.7	2.3	2.4	2.2	2.1	2.1
Male.....	10.0	8.3	6.7	4.5	4.2	3.1	3.5	2.8	2.4	2.7
Female.....	3.4	2.6	1.8	1.5	1.4	1.6	1.5	1.8	2.0	1.9
White, both sexes.....	4.4	3.8	2.9	2.3	2.1	1.8	1.9	2.0	1.9	1.9
Male.....	6.2	5.6	4.5	3.3	3.3	2.2	2.7	2.2	2.0	1.9
Female.....	2.6	2.0	1.4	1.3	1.1	1.4	1.3	1.8	1.8	1.9
Black, both sexes.....	21.6	17.2	13.9	8.3	7.8	7.1	7.1	5.8	5.6	*
Male.....	36.7	29.3	24.1	15.3	12.9	12.6	12.7	11.2	*	*
Female.....	8.3	6.7	5.2	2.5	3.8	*	*	*	*	*

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Assault (homicide) by discharge of firearms (\*U01.4,X93-X95)

All races 3/, both sexes.....	4.1	3.0	2.1	1.4	1.3	1.1	1.1	0.8	0.6	0.5
Male.....	6.6	4.7	3.4	2.2	2.0	1.4	1.7	1.0	*	*
Female.....	1.7	1.4	0.8	0.6	0.6	0.8	0.6	0.7	0.6	*
White, both sexes.....	2.6	2.1	1.5	1.1	1.0	0.9	0.9	0.8	0.5	*
Male.....	3.9	3.0	2.3	1.6	1.6	1.1	1.2	0.9	*	*
Female.....	1.4	1.1	0.7	0.6	0.5	0.7	0.6	0.8	*	*
Black, both sexes.....	14.4	9.9	6.9	3.7	3.3	2.8	3.3	*	*	*
Male.....	26.1	17.9	12.6	7.3	5.6	5.2	7.0	*	*	*
Female.....	4.2	3.0	2.1	*	*	*	*	*	*	*

Assault (homicide) by other and unspecified means and their sequelae (\*U01.0-\*U01.3,\*U01.5-\*U01.9,\*U02,X85-X92,X96-Y09,Y87.1)

All races 3/, both sexes.....	2.5	2.4	2.1	1.6	1.4	1.2	1.4	1.4	1.5	1.7
Male.....	3.4	3.6	3.2	2.3	2.1	1.7	1.9	1.8	1.8	2.1
Female.....	1.7	1.2	1.0	0.9	0.8	0.8	1.0	1.1	1.3	1.5
White, both sexes.....	1.8	1.7	1.4	1.2	1.1	0.9	1.1	1.1	1.3	1.5
Male.....	2.4	2.5	2.2	1.7	1.6	1.1	1.5	1.3	1.6	1.6
Female.....	1.2	0.9	0.7	0.8	0.6	0.7	0.8	1.0	1.2	1.5
Black, both sexes.....	7.2	7.3	7.0	4.6	4.5	4.3	3.7	4.5	*	*
Male.....	10.6	11.5	11.5	8.0	7.3	7.4	*	*	*	*
Female.....	4.2	3.7	3.1	*	*	*	*	*	*	*



Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Legal intervention (Y35,Y89.0)

All races 3/, both sexes.....	0.2	0.2	0.1	*	*	*	*	*	*	*
Male.....	0.4	0.3	0.2	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	0.2	0.2	*	*	*	*	*	*	*	*
Male.....	0.4	0.3	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

Events of undetermined intent (Y10-Y34,Y87.2,Y89.9)

All races 3/, both sexes.....	3.1	3.2	2.9	1.6	1.1	0.9	0.7	0.8	0.9	1.5
Male.....	3.6	3.8	3.5	2.1	1.1	1.1	1.0	1.0	*	1.6
Female.....	2.6	2.7	2.4	1.2	1.1	0.8	0.5	0.6	0.9	1.4
White, both sexes.....	3.1	3.3	2.9	1.6	1.1	0.9	0.7	0.7	0.9	1.5
Male.....	3.5	3.7	3.3	2.0	0.9	1.1	1.0	0.8	*	1.7
Female.....	2.6	2.9	2.5	1.3	1.2	0.8	0.5	0.6	0.8	1.3
Black, both sexes.....	4.1	3.7	3.8	1.8	1.6	*	*	*	*	*
Male.....	5.3	5.3	6.0	3.2	*	*	*	*	*	*
Female.....	2.9	2.3	2.0	*	*	*	*	*	*	*

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Discharge of firearms, undetermined intent (Y22-Y24)

All races 3/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

Other and unspecified events of undetermined intent and their sequelae (Y10-Y21,Y25-Y34,Y87.2,Y89.9)

All races 3/, both sexes.....	3.0	3.2	2.8	1.6	1.0	0.9	0.7	0.7	0.9	1.4
Male.....	3.5	3.7	3.3	2.0	1.0	1.1	0.9	0.9	*	1.5
Female.....	2.6	2.7	2.4	1.1	1.0	0.7	0.5	0.6	0.9	1.4
White, both sexes.....	3.0	3.3	2.8	1.6	1.0	0.9	0.6	0.7	0.8	1.4
Male.....	3.4	3.7	3.1	1.9	0.8	1.0	0.8	0.7	*	1.6
Female.....	2.6	2.9	2.5	1.2	1.1	0.8	0.5	0.6	0.8	1.3
Black, both sexes.....	3.9	3.6	3.8	1.7	*	*	*	*	*	*
Male.....	5.2	5.2	6.0	3.2	*	*	*	*	*	*
Female.....	2.9	2.2	1.9	*	*	*	*	*	*	*

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Operations of war and their sequelae (Y36,Y89.1)

All races 3/, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
White, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*
Black, both sexes.....	*	*	*	*	*	*	*	*	*	*
Male.....	*	*	*	*	*	*	*	*	*	*
Female.....	*	*	*	*	*	*	*	*	*	*

Complications of medical and surgical care (Y40-Y84,Y88)

All races 3/, both sexes.....	0.4	0.5	0.8	1.0	1.4	2.0	3.3	4.7	7.2	9.1
Male.....	0.4	0.5	0.8	1.1	1.7	2.3	3.7	4.6	8.9	10.8
Female.....	0.4	0.6	0.9	1.0	1.2	1.8	3.0	4.8	6.2	8.3
White, both sexes.....	0.4	0.5	0.7	0.9	1.3	1.8	3.0	4.4	7.3	9.2
Male.....	0.4	0.4	0.7	0.9	1.5	2.3	3.4	4.3	8.8	11.2
Female.....	0.4	0.5	0.7	1.0	1.1	1.5	2.6	4.5	6.3	8.3
Black, both sexes.....	*	1.0	2.1	2.2	2.9	4.5	6.4	9.4	8.5	9.1
Male.....	*	*	2.2	2.6	3.8	*	*	9.4	*	*
Female.....	*	*	2.1	*	*	4.8	6.7	9.3	7.4	9.0

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

40-44 years    45-49 years    50-54 years    55-59 years    60-64 years    65-69 years    70-74 years    75-79 years    80-84 years    85 years and over

Alcohol-induced causes (E24.4,F10,G31.2,G62.1,G72.1,I42.6,K29.2,K70,K86.0,R78.0,X45,X65,Y15) 4/

All races 3/, both sexes.....	10.1	15.7	19.7	19.7	19.3	16.3	13.7	10.1	8.1	4.6
Male.....	14.4	22.9	30.9	31.7	31.3	27.0	22.6	18.4	14.9	9.3
Female.....	5.8	8.6	8.9	8.4	8.4	7.1	6.5	4.0	3.8	2.5
White, both sexes.....	10.4	16.1	20.0	20.2	19.1	16.6	13.8	10.1	8.0	4.6
Male.....	14.9	23.4	31.2	32.3	30.8	27.0	22.4	18.2	14.8	9.4
Female.....	5.8	8.9	9.0	8.5	8.2	7.3	6.6	4.1	3.8	2.5
Black, both sexes.....	8.1	14.6	19.6	20.3	23.5	17.0	15.9	10.7	8.5	5.6
Male.....	11.1	23.1	31.7	33.6	41.3	31.1	30.4	24.1	15.5	*
Female.....	5.4	7.3	9.4	9.3	9.6	6.5	6.1	*	*	*

Drug-induced causes (D52.1,D59.0,D59.2,D61.1,D64.2,E06.4,E16.0,E23.1,E24.2,E27.3,E66.1,F11.0-F11.5,F11.7-F11.9,F12.0-F12.5,F12.7-F12.9,F13.0-F13.5,F13.7-F13.9,F14.0-F14.5,F14.7-F14.9,F15.0-F15.5,F15.7-F15.9,F16.0-F16.5,F16.7-F16.9,F17.0,F17.3-F17.5,F17.7-F17.9,F18.0-F18.5,F18.7-F18.9,F19.0-F19.5,F19.7-F19.9,G21.1,G24.0,G25.1,G25.4,G25.6,G44.4,G62.0,G72.0,I95.2,J70.2-J70.4,L10.5,L27.0-L27.1,M10.2,M32.0,M80.4,M81.4,M83.5,M87.1,R78.1-R78.5,X40-X44,X60-X64,X85,Y10-Y14) 4/

All races 3/, both sexes.....	24.0	25.8	21.2	12.8	7.8	5.0	4.4	4.9	6.1	7.6
Male.....	29.9	31.9	26.6	16.0	9.0	5.4	5.2	6.6	6.9	9.9
Female.....	18.2	19.8	16.1	9.8	6.6	4.8	3.8	3.7	5.5	6.6
White, both sexes.....	25.4	26.5	20.5	12.2	7.8	5.1	4.6	5.0	6.3	8.0
Male.....	31.4	32.0	24.8	14.4	8.4	5.0	5.2	6.7	6.9	10.6
Female.....	19.2	21.0	16.2	10.2	7.2	5.1	4.1	3.8	5.9	6.8
Black, both sexes.....	23.8	29.3	32.8	22.4	10.4	5.6	4.6	4.3	*	*
Male.....	30.2	41.3	48.4	36.0	19.1	10.0	7.6	*	*	*
Female.....	18.1	18.9	19.6	11.2	3.5	*	*	*	*	*

Worktable 210R. Death rates for 113 selected causes, alcohol-induced causes, drug-induced causes, and injury by firearms, by 5-year age groups, race, and sex: United States, 2005

[Rates per 100,000 population in specified group. Rates are based on populations estimated as of July 1, 2005. Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2005, multiple-race data were reported by California, Connecticut, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New York, Oklahoma, South Carolina, South Dakota, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005." For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Technical Appendix from Vital Statistics of the United States: Mortality, 2005"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, 1992), race and sex

	40-44 years	45-49 years	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85 years and over
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------------

Injury by firearms (\*U01.4,W32-W34,X72-X74,X93-X95,Y22-Y24,Y35.0) 4/

All races 3/, both sexes.....	11.6	11.4	10.9	9.8	9.9	9.5	11.3	13.0	14.5	12.0
Male.....	19.2	19.1	18.7	17.0	17.9	17.4	22.6	28.1	34.6	35.4
Female.....	4.1	4.0	3.3	3.0	2.6	2.7	2.3	2.1	2.0	1.3
White, both sexes.....	11.1	11.7	11.4	10.7	10.7	10.3	12.2	14.2	15.6	13.0
Male.....	18.0	19.2	19.4	18.2	19.2	18.9	24.0	30.0	37.2	38.0
Female.....	4.2	4.2	3.6	3.4	2.7	2.8	2.4	2.4	2.1	1.4
Black, both sexes.....	18.1	12.9	10.1	6.0	6.2	5.7	7.6	5.3	6.3	*
Male.....	33.3	23.5	18.8	11.8	11.1	10.2	16.1	14.3	14.7	*
Female.....	4.8	3.7	2.7	*	*	*	*	*	*	*

\* Figure does not meet standards of reliability or precision.

0.0 Quantity more than zero but less than 0.05.

... Category not applicable.

1/ Figures for age not stated included in "All ages" but not distributed among age groups.

2/ Death rates for "Under 1 year" (based on population estimates) differ from infant mortality rates (based on live births).

3/ Includes races other than white and black.

4/ Included in selected categories.

Source: CDC/NCHS, National Vital Statistics System, Mortality.