

Person with disabilities experiencing problems accessing  
NCHS birth and death tables should contact  
[nhsed@cdc.gov](mailto:nhsed@cdc.gov), or call 301-458-4688

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>United States</b>												
All races, both sexes.....	2,423,712	29,138	4,703	2,711	3,436	13,299	20,683	20,931	21,641	30,881	48,725	77,738
Male.....	1,203,968	16,293	2,634	1,519	2,066	9,558	15,758	15,107	14,685	19,755	30,350	47,904
Female.....	1,219,744	12,845	2,069	1,192	1,370	3,741	4,925	5,824	6,956	11,126	18,375	29,834
White, both sexes.....	2,074,151	18,807	3,287	2,001	2,479	9,765	15,372	15,382	15,799	22,804	36,767	59,635
Male.....	1,023,951	10,540	1,847	1,112	1,478	6,851	11,662	11,159	10,839	14,820	23,301	37,447
Female.....	1,050,200	8,267	1,440	889	1,001	2,914	3,710	4,223	4,960	7,984	13,466	22,188
Black, both sexes.....	289,585	8,944	1,137	577	786	2,956	4,424	4,741	4,921	6,831	10,159	15,609
Male.....	148,309	4,975	621	331	485	2,302	3,449	3,398	3,250	4,147	5,959	8,970
Female.....	141,276	3,969	516	246	301	654	975	1,343	1,671	2,684	4,200	6,639
American Indian 1/, both sexes.....	14,367	432	97	40	54	265	357	349	382	497	716	937
Male.....	7,885	241	57	25	36	182	259	250	264	315	453	562
Female.....	6,482	191	40	15	18	83	98	99	118	182	263	375
Asian or Pacific Islander 2/, both sexes.....	45,609	955	182	93	117	313	530	459	539	749	1,083	1,557
Male.....	23,823	537	109	51	67	223	388	300	332	473	637	925
Female.....	21,786	418	73	42	50	90	142	159	207	276	446	632
<b>Alabama</b>												
All races, both sexes.....	46,696	641	76	66	55	304	529	442	514	727	1,049	1,685
Male.....	23,466	353	44	31	37	216	416	318	332	428	636	1,039
Female.....	23,230	288	32	35	18	88	113	124	182	299	413	646
White, both sexes.....	35,502	350	38	39	37	209	362	271	327	464	701	1,122
Male.....	17,777	196	18	18	23	147	286	202	208	287	430	695
Female.....	17,725	154	20	21	14	62	76	69	119	177	271	427
Black, both sexes.....	11,040	284	38	26	18	94	164	166	181	258	342	554
Male.....	5,612	155	26	12	14	68	128	113	121	138	204	337
Female.....	5,428	129	12	14	4	26	36	53	60	120	138	217
American Indian 1/, both sexes.....	52	2	-	-	-	-	1	2	1	2	3	5
Male.....	27	-	-	-	-	-	1	1	-	1	1	5
Female.....	25	2	-	-	-	-	-	1	1	1	2	-
Asian or Pacific Islander 2/, both sexes.....	102	5	-	1	-	1	2	3	5	3	3	4
Male.....	50	2	-	1	-	1	1	2	3	2	1	2
Female.....	52	3	-	-	-	-	1	1	2	1	2	2

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Alaska												
All races, both sexes.....	3,463	72	14	13	15	53	76	55	65	88	133	201
Male.....	1,962	46	10	10	10	40	62	43	44	56	79	118
Female.....	1,501	26	4	3	5	13	14	12	21	32	54	83
White, both sexes.....	2,445	36	6	7	6	25	40	30	35	61	81	138
Male.....	1,400	21	5	5	3	19	34	25	25	41	53	86
Female.....	1,045	15	1	2	3	6	6	5	10	20	28	52
Black, both sexes.....	96	3	-	1	1	2	5	1	1	2	4	5
Male.....	60	2	-	1	-	2	5	1	1	2	2	2
Female.....	36	1	-	-	1	-	-	-	-	-	2	3
American Indian 1/, both sexes.....	808	29	7	4	7	24	29	22	29	23	44	52
Male.....	438	19	5	3	6	17	21	15	18	13	22	26
Female.....	370	10	2	1	1	7	8	7	11	10	22	26
Asian or Pacific Islander 2/, both sexes.....	114	4	1	1	1	2	2	2	-	2	4	6
Male.....	64	4	-	1	1	2	2	2	-	-	2	4
Female.....	50	-	1	-	-	-	-	-	-	2	2	2
Arizona												
All races, both sexes.....	45,554	711	115	65	91	349	535	497	545	659	1,013	1,479
Male.....	23,886	404	58	33	50	241	415	382	380	438	649	971
Female.....	21,668	307	57	32	41	108	120	115	165	221	364	508
White, both sexes.....	42,170	581	94	46	71	279	425	422	439	529	826	1,260
Male.....	22,027	325	48	22	39	199	329	330	306	354	529	826
Female.....	20,143	256	46	24	32	80	96	92	133	175	297	434
Black, both sexes.....	1,300	65	6	6	7	20	34	28	31	42	53	84
Male.....	709	43	2	2	5	13	23	21	23	30	31	56
Female.....	591	22	4	4	2	7	11	7	8	12	22	28
American Indian 1/, both sexes.....	1,700	54	14	9	10	48	67	42	65	81	123	120
Male.....	970	29	8	7	4	28	55	27	48	50	84	78
Female.....	730	25	6	2	6	20	12	15	17	31	39	42
Asian or Pacific Islander 2/, both sexes.....	384	11	1	4	3	2	9	5	10	7	11	15
Male.....	180	7	-	2	2	1	8	4	3	4	5	11
Female.....	204	4	1	2	1	1	1	1	7	3	6	4

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Arkansas												
All races, both sexes.....	28,191	317	72	31	50	184	253	273	259	391	642	988
Male.....	14,117	180	39	19	30	118	173	192	165	273	400	601
Female.....	14,074	137	33	12	20	66	80	81	94	118	242	387
White, both sexes.....	24,171	209	47	23	34	136	194	196	194	310	482	770
Male.....	12,105	121	25	15	20	84	129	136	130	216	299	484
Female.....	12,066	88	22	8	14	52	65	60	64	94	183	286
Black, both sexes.....	3,873	107	25	7	14	44	57	76	62	79	148	215
Male.....	1,940	58	14	3	9	31	43	55	34	55	95	114
Female.....	1,933	49	11	4	5	13	14	21	28	24	53	101
American Indian 1/, both sexes.....	67	1	-	-	1	1	1	1	1	1	6	3
Male.....	34	1	-	-	1	1	-	1	-	1	2	3
Female.....	33	-	-	-	-	-	1	-	1	-	4	-
Asian or Pacific Islander 2/, both sexes.....	80	-	-	1	1	3	1	-	2	1	6	-
Male.....	38	-	-	1	-	2	1	-	1	1	4	-
Female.....	42	-	-	-	1	1	-	-	1	-	2	-
California												
All races, both sexes.....	233,720	2,944	496	294	368	1,440	2,257	2,132	2,163	3,205	4,848	7,882
Male.....	118,560	1,634	264	167	223	1,092	1,759	1,558	1,494	2,102	3,114	4,945
Female.....	115,160	1,310	232	127	145	348	498	574	669	1,103	1,734	2,937
White, both sexes.....	194,859	2,223	381	233	293	1,092	1,742	1,649	1,652	2,468	3,821	6,213
Male.....	98,235	1,227	208	131	177	823	1,370	1,237	1,150	1,645	2,475	3,927
Female.....	96,624	996	173	102	116	269	372	412	502	823	1,346	2,286
Black, both sexes.....	18,701	421	50	27	41	219	300	294	330	424	588	992
Male.....	9,708	229	18	21	26	176	232	195	230	267	362	600
Female.....	8,993	192	32	6	15	43	68	99	100	157	226	392
American Indian 1/, both sexes.....	1,364	21	7	1	4	14	20	30	24	32	66	81
Male.....	775	12	3	-	2	11	16	26	18	19	43	57
Female.....	589	9	4	1	2	3	4	4	6	13	23	24
Asian or Pacific Islander 2/, both sexes.....	18,796	279	58	33	30	115	195	159	157	281	373	596
Male.....	9,842	166	35	15	18	82	141	100	96	171	234	361
Female.....	8,954	113	23	18	12	33	54	59	61	110	139	235

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Colorado</b>												
All races, both sexes.....	29,993	433	59	48	39	187	264	330	291	459	698	1,066
Male.....	15,147	251	31	23	28	135	205	229	206	300	441	669
Female.....	14,846	182	28	25	11	52	59	101	85	159	257	397
White, both sexes.....	28,492	375	49	48	38	172	237	305	265	410	628	966
Male.....	14,358	223	26	23	27	123	181	212	189	265	404	614
Female.....	14,134	152	23	25	11	49	56	93	76	145	224	352
Black, both sexes.....	1,050	45	9	-	-	9	21	19	18	37	52	75
Male.....	559	22	5	-	-	6	19	13	11	25	27	39
Female.....	491	23	4	-	-	3	2	6	7	12	25	36
American Indian 1/, both sexes.....	198	6	-	-	-	3	4	3	5	9	9	16
Male.....	112	3	-	-	-	3	4	2	3	8	6	10
Female.....	86	3	-	-	-	-	-	1	2	1	3	6
Asian or Pacific Islander 2/, both sexes.....	253	7	1	-	1	3	2	3	3	3	9	9
Male.....	118	3	-	-	1	3	1	2	3	2	4	6
Female.....	135	4	1	-	-	-	1	1	-	1	5	3
<b>Connecticut</b>												
All races, both sexes.....	28,651	276	39	14	23	111	163	159	179	264	453	818
Male.....	13,561	144	23	11	14	82	120	115	112	177	290	505
Female.....	15,090	132	16	3	9	29	43	44	67	87	163	313
White, both sexes.....	26,350	197	22	8	19	88	124	135	143	213	365	688
Male.....	12,350	103	13	5	12	65	92	94	91	149	239	431
Female.....	14,000	94	9	3	7	23	32	41	52	64	126	257
Black, both sexes.....	2,076	68	15	6	3	20	35	21	34	46	82	121
Male.....	1,083	34	9	6	2	16	25	18	19	26	46	68
Female.....	993	34	6	-	1	4	10	3	15	20	36	53
American Indian 1/, both sexes.....	27	1	-	-	-	-	-	-	1	1	-	2
Male.....	14	-	-	-	-	-	-	-	1	1	-	1
Female.....	13	1	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	198	10	2	-	1	3	4	3	1	4	6	7
Male.....	114	7	1	-	-	1	3	3	1	1	5	5
Female.....	84	3	1	-	1	2	1	-	-	3	1	2

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Delaware</b>												
All races, both sexes.....	7,327	91	8	2	6	35	57	59	49	119	169	274
Male.....	3,678	48	4	1	6	26	44	43	34	79	98	177
Female.....	3,649	43	4	1	-	9	13	16	15	40	71	97
White, both sexes.....	6,054	51	5	2	4	23	33	45	34	89	117	197
Male.....	3,016	26	2	1	4	14	26	35	26	62	71	129
Female.....	3,038	25	3	1	-	9	7	10	8	27	46	68
Black, both sexes.....	1,198	39	2	-	1	11	22	11	13	29	50	76
Male.....	616	21	1	-	1	11	16	6	7	16	26	48
Female.....	582	18	1	-	-	-	6	5	6	13	24	28
American Indian 1/, both sexes.....	27	-	-	-	-	-	-	-	-	1	1	-
Male.....	17	-	-	-	-	-	-	-	-	1	1	-
Female.....	10	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	48	1	1	-	1	1	2	3	2	-	1	1
Male.....	29	1	1	-	1	1	2	2	1	-	-	-
Female.....	19	-	-	-	-	-	-	1	1	-	1	1
<b>Dist. of Columbia</b>												
All races, both sexes.....	5,188	116	9	11	5	37	55	66	87	98	155	218
Male.....	2,592	53	5	3	4	29	44	53	62	68	90	131
Female.....	2,596	63	4	8	1	8	11	13	25	30	65	87
White, both sexes.....	1,155	29	2	3	-	4	6	10	11	12	12	22
Male.....	597	15	1	1	-	4	4	7	9	7	7	16
Female.....	558	14	1	2	-	-	2	3	2	5	5	6
Black, both sexes.....	3,983	87	7	8	5	33	49	55	75	84	142	195
Male.....	1,971	38	4	2	4	25	40	46	52	59	82	114
Female.....	2,012	49	3	6	1	8	9	9	23	25	60	81
American Indian 1/, both sexes.....	6	-	-	-	-	-	-	-	1	-	-	-
Male.....	3	-	-	-	-	-	-	-	1	-	-	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	44	-	-	-	-	-	-	1	-	2	1	1
Male.....	21	-	-	-	-	-	-	-	-	2	1	1
Female.....	23	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Florida</b>												
All races, both sexes.....	168,096	1,685	313	142	207	834	1,493	1,438	1,487	2,063	3,226	5,253
Male.....	87,273	946	175	76	117	607	1,153	1,026	1,007	1,345	2,038	3,289
Female.....	80,823	739	138	66	90	227	340	412	480	718	1,188	1,964
White, both sexes.....	148,198	950	204	94	130	588	1,108	1,036	1,083	1,499	2,465	4,052
Male.....	76,796	550	109	50	74	415	849	748	746	1,005	1,611	2,610
Female.....	71,402	400	95	44	56	173	259	288	337	494	854	1,442
Black, both sexes.....	18,666	709	101	46	74	234	368	378	377	535	707	1,140
Male.....	9,820	382	60	25	41	182	293	264	237	320	396	643
Female.....	8,846	327	41	21	33	52	75	114	140	215	311	497
American Indian 1/, both sexes.....	175	1	1	-	-	1	4	2	8	5	10	16
Male.....	107	1	1	-	-	-	2	1	8	5	5	9
Female.....	68	-	-	-	-	1	2	1	-	-	5	7
Asian or Pacific Islander 2/, both sexes.....	1,057	25	7	2	3	11	13	22	19	24	44	45
Male.....	550	13	5	1	2	10	9	13	16	15	26	27
Female.....	507	12	2	1	1	1	4	9	3	9	18	18
<b>Georgia</b>												
All races, both sexes.....	68,331	1,206	190	103	116	497	752	794	823	1,223	1,771	2,690
Male.....	34,246	662	95	63	71	351	570	547	515	776	1,090	1,607
Female.....	34,085	544	95	40	45	146	182	247	308	447	681	1,083
White, both sexes.....	49,118	516	96	61	55	294	464	459	471	697	1,039	1,657
Male.....	24,466	275	51	38	37	206	355	333	299	453	655	1,028
Female.....	24,652	241	45	23	18	88	109	126	172	244	384	629
Black, both sexes.....	18,590	664	88	39	56	195	277	326	338	502	715	1,000
Male.....	9,447	371	42	22	32	140	207	207	207	308	427	560
Female.....	9,143	293	46	17	24	55	70	119	131	194	288	440
American Indian 1/, both sexes.....	61	3	1	-	-	2	-	-	1	2	3	4
Male.....	35	3	-	-	-	1	-	-	-	2	2	2
Female.....	26	-	1	-	-	1	-	-	1	-	1	2
Asian or Pacific Islander 2/, both sexes.....	562	23	5	3	5	6	11	9	13	22	14	29
Male.....	298	13	2	3	2	4	8	7	9	13	6	17
Female.....	264	10	3	-	3	2	3	2	4	9	8	12

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Hawaii</b>												
All races, both sexes.....	9,495	124	20	6	18	32	71	60	78	111	169	301
Male.....	5,156	73	11	5	13	22	57	51	52	71	99	195
Female.....	4,339	51	9	1	5	10	14	9	26	40	70	106
White, both sexes.....	2,732	34	5	-	5	2	33	21	16	39	63	105
Male.....	1,557	18	2	-	2	2	27	17	12	29	39	74
Female.....	1,175	16	3	-	3	-	6	4	4	10	24	31
Black, both sexes.....	89	8	2	-	-	-	-	1	1	1	4	11
Male.....	62	5	-	-	-	-	-	1	1	1	4	9
Female.....	27	3	2	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes.....	22	-	-	-	-	-	1	1	2	1	-	-
Male.....	17	-	-	-	-	-	1	1	2	1	-	-
Female.....	5	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6,652	82	13	6	13	30	37	37	59	70	102	185
Male.....	3,520	50	9	5	11	20	29	32	37	40	56	112
Female.....	3,132	32	4	1	2	10	8	5	22	30	46	73
<b>Idaho</b>												
All races, both sexes.....	10,822	169	31	11	27	85	105	85	103	126	179	309
Male.....	5,379	94	14	4	20	50	78	62	65	70	100	191
Female.....	5,443	75	17	7	7	35	27	23	38	56	79	118
White, both sexes.....	10,612	158	29	11	24	80	99	81	96	113	171	298
Male.....	5,256	88	13	4	19	47	73	59	61	65	95	184
Female.....	5,356	70	16	7	5	33	26	22	35	48	76	114
Black, both sexes.....	26	3	-	-	2	-	-	-	-	-	2	1
Male.....	14	1	-	-	-	-	-	-	-	-	2	-
Female.....	12	2	-	-	2	-	-	-	-	-	-	1
American Indian 1/, both sexes.....	116	4	2	-	1	4	4	3	4	11	5	9
Male.....	67	3	1	-	1	3	3	2	2	4	3	6
Female.....	49	1	1	-	-	1	1	1	2	7	2	3
Asian or Pacific Islander 2/, both sexes.....	68	4	-	-	-	1	2	1	3	2	1	1
Male.....	42	2	-	-	-	-	2	1	2	1	-	1
Female.....	26	2	-	-	-	1	-	-	1	1	1	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Illinois</b>												
All races, both sexes.....	100,503	1,217	188	106	164	560	800	778	820	1,164	1,873	3,131
Male.....	48,453	679	116	64	93	408	636	563	572	775	1,102	1,908
Female.....	52,050	538	72	42	71	152	164	215	248	389	771	1,223
White, both sexes.....	83,335	725	116	75	110	382	538	511	536	783	1,296	2,158
Male.....	39,567	404	74	45	57	272	423	373	380	519	788	1,336
Female.....	43,768	321	42	30	53	110	115	138	156	264	508	822
Black, both sexes.....	15,824	452	63	30	51	170	239	256	272	354	549	936
Male.....	8,208	252	38	19	34	130	194	186	186	244	302	549
Female.....	7,616	200	25	11	17	40	45	70	86	110	247	387
American Indian 1/, both sexes.....	81	2	1	-	-	2	1	-	-	2	4	8
Male.....	40	2	1	-	-	2	1	-	-	-	2	4
Female.....	41	-	-	-	-	-	-	-	-	2	2	4
Asian or Pacific Islander 2/, both sexes.....	1,263	38	8	1	3	6	22	11	12	25	24	29
Male.....	638	21	3	-	2	4	18	4	6	12	10	19
Female.....	625	17	5	1	1	2	4	7	6	13	14	10
<b>Indiana</b>												
All races, both sexes.....	54,000	681	105	64	91	307	425	479	504	658	1,058	1,754
Male.....	26,396	379	55	37	52	226	332	349	341	427	659	1,071
Female.....	27,604	302	50	27	39	81	93	130	163	231	399	683
White, both sexes.....	49,586	513	76	52	70	258	343	391	417	560	910	1,532
Male.....	24,151	288	40	31	40	188	266	281	277	370	571	952
Female.....	25,435	225	36	21	30	70	77	110	140	190	339	580
Black, both sexes.....	4,314	166	28	11	18	48	81	87	84	91	145	216
Male.....	2,197	91	15	5	12	38	65	67	62	53	85	118
Female.....	2,117	75	13	6	6	10	16	20	22	38	60	98
American Indian 1/, both sexes.....	22	-	-	-	-	-	-	1	2	2	-	2
Male.....	11	-	-	-	-	-	-	1	1	2	-	1
Female.....	11	-	-	-	-	-	-	-	1	-	-	1
Asian or Pacific Islander 2/, both sexes.....	78	2	1	1	3	1	1	-	1	5	3	4
Male.....	37	-	-	1	-	-	1	-	1	2	3	-
Female.....	41	2	1	-	3	1	-	-	-	3	-	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Iowa</b>												
All races, both sexes.....	27,221	225	43	25	34	121	136	136	156	218	386	634
Male.....	12,981	124	29	14	20	79	108	91	93	143	237	365
Female.....	14,240	101	14	11	14	42	28	45	63	75	149	269
White, both sexes.....	26,660	201	40	25	34	112	125	122	144	201	362	602
Male.....	12,685	116	27	14	20	72	102	81	90	130	225	342
Female.....	13,975	85	13	11	14	40	23	41	54	71	137	260
Black, both sexes.....	434	22	3	-	-	8	8	12	8	12	12	25
Male.....	232	7	2	-	-	6	4	9	3	8	6	19
Female.....	202	15	1	-	-	2	4	3	5	4	6	6
American Indian 1/, both sexes.....	47	-	-	-	-	-	1	-	1	4	6	4
Male.....	25	-	-	-	-	-	1	-	-	4	4	2
Female.....	22	-	-	-	-	-	-	-	1	-	2	2
Asian or Pacific Islander 2/, both sexes.....	80	2	-	-	-	1	2	2	3	1	6	3
Male.....	39	1	-	-	-	1	1	1	-	1	2	2
Female.....	41	1	-	-	-	-	1	1	3	-	4	1
<b>Kansas</b>												
All races, both sexes.....	24,491	333	38	26	36	139	180	166	176	224	412	718
Male.....	11,825	187	23	14	19	97	134	128	115	147	235	424
Female.....	12,666	146	15	12	17	42	46	38	61	77	177	294
White, both sexes.....	22,903	260	34	20	32	114	161	136	145	189	354	626
Male.....	11,005	142	22	11	17	79	120	103	92	128	202	372
Female.....	11,898	118	12	9	15	35	41	33	53	61	152	254
Black, both sexes.....	1,225	62	2	5	4	17	17	26	23	27	46	68
Male.....	634	38	-	2	2	13	12	21	17	15	26	37
Female.....	591	24	2	3	2	4	5	5	6	12	20	31
American Indian 1/, both sexes.....	247	6	2	1	-	5	2	3	4	6	5	21
Male.....	128	4	1	1	-	4	2	3	3	3	3	12
Female.....	119	2	1	-	-	1	-	-	1	3	2	9
Asian or Pacific Islander 2/, both sexes.....	116	5	-	-	-	3	-	1	4	2	7	3
Male.....	58	3	-	-	-	1	-	1	3	1	4	3
Female.....	58	2	-	-	-	2	-	-	1	1	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Kentucky</b>												
All races, both sexes.....	40,090	397	64	39	64	210	315	423	406	570	950	1,310
Male.....	20,020	216	36	20	38	155	233	291	284	366	597	828
Female.....	20,070	181	28	19	26	55	82	132	122	204	353	482
White, both sexes.....	37,274	316	53	35	57	186	278	368	367	518	851	1,170
Male.....	18,598	175	31	18	34	134	203	248	260	334	546	750
Female.....	18,676	141	22	17	23	52	75	120	107	184	305	420
Black, both sexes.....	2,691	72	10	4	6	24	36	49	35	47	98	135
Male.....	1,354	35	4	2	3	21	29	39	22	29	50	75
Female.....	1,337	37	6	2	3	3	7	10	13	18	48	60
American Indian 1/, both sexes.....	20	-	-	-	-	-	-	2	2	-	-	1
Male.....	11	-	-	-	-	-	-	1	1	-	-	1
Female.....	9	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	105	9	1	-	1	-	1	4	2	5	1	4
Male.....	57	6	1	-	1	-	1	3	1	3	1	2
Female.....	48	3	-	-	-	-	-	1	1	2	-	2
<b>Louisiana</b>												
All races, both sexes.....	39,966	608	101	56	82	304	524	610	548	678	1,052	1,501
Male.....	20,225	347	59	39	52	228	409	445	385	427	695	918
Female.....	19,741	261	42	17	30	76	115	165	163	251	357	583
White, both sexes.....	27,926	238	41	33	51	167	297	349	299	375	595	875
Male.....	13,919	138	26	22	35	126	237	259	210	247	401	539
Female.....	14,007	100	15	11	16	41	60	90	89	128	194	336
Black, both sexes.....	11,789	365	54	22	30	131	223	248	243	297	445	616
Male.....	6,163	208	28	16	17	98	170	177	169	176	286	374
Female.....	5,626	157	26	6	13	33	53	71	74	121	159	242
American Indian 1/, both sexes.....	76	1	2	-	1	3	3	4	2	-	5	6
Male.....	44	1	1	-	-	2	1	3	2	-	4	1
Female.....	32	-	1	-	1	1	2	1	-	-	1	5
Asian or Pacific Islander 2/, both sexes.....	175	4	4	1	-	3	1	9	4	6	7	4
Male.....	99	-	4	1	-	2	1	6	4	4	4	4
Female.....	76	4	-	-	-	1	-	3	-	2	3	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Maine												
All races, both sexes.....	12,493	89	14	12	8	48	89	75	78	100	168	309
Male.....	6,191	51	9	5	5	30	66	55	60	67	111	191
Female.....	6,302	38	5	7	3	18	23	20	18	33	57	118
White, both sexes.....	12,374	85	13	11	7	46	86	69	75	98	162	296
Male.....	6,120	49	8	5	5	29	63	51	58	66	108	183
Female.....	6,254	36	5	6	2	17	23	18	17	32	54	113
Black, both sexes.....	49	4	1	1	1	2	2	3	1	-	2	6
Male.....	31	2	1	-	-	1	2	2	1	-	1	5
Female.....	18	2	-	1	1	1	-	1	-	-	1	1
American Indian 1/, both sexes.....	49	-	-	-	-	-	-	1	2	2	4	5
Male.....	30	-	-	-	-	-	-	1	1	1	2	2
Female.....	19	-	-	-	-	-	-	-	1	1	2	3
Asian or Pacific Islander 2/, both sexes.....	21	-	-	-	-	-	1	2	-	-	-	2
Male.....	10	-	-	-	-	-	1	1	-	-	-	1
Female.....	11	-	-	-	-	-	-	1	-	-	-	1
Maryland												
All races, both sexes.....	43,757	625	96	49	73	273	429	438	424	691	1,084	1,579
Male.....	21,597	337	59	29	39	205	332	314	289	439	666	968
Female.....	22,160	288	37	20	34	68	97	124	135	252	418	611
White, both sexes.....	31,111	219	45	29	39	128	211	226	201	334	568	868
Male.....	15,060	118	29	14	21	87	156	152	140	225	366	551
Female.....	16,051	101	16	15	18	41	55	74	61	109	202	317
Black, both sexes.....	11,881	371	44	20	33	137	207	200	212	347	486	682
Male.....	6,123	206	27	15	18	112	169	156	140	205	281	400
Female.....	5,758	165	17	5	15	25	38	44	72	142	205	282
American Indian 1/, both sexes.....	45	2	1	-	-	-	1	1	-	-	1	3
Male.....	28	-	1	-	-	-	1	1	-	-	1	3
Female.....	17	2	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	720	33	6	-	1	8	10	11	11	10	29	26
Male.....	386	13	2	-	-	6	6	5	9	9	18	14
Female.....	334	20	4	-	1	2	4	6	2	1	11	12

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Massachusetts</b>												
All races, both sexes.....	52,917	384	47	36	48	199	312	315	300	563	857	1,403
Male.....	24,971	214	24	19	31	142	230	234	215	358	541	876
Female.....	27,946	170	23	17	17	57	82	81	85	205	316	527
White, both sexes.....	50,002	283	41	29	34	164	260	265	263	483	747	1,259
Male.....	23,439	158	21	13	21	115	185	191	191	310	479	795
Female.....	26,563	125	20	16	13	49	75	74	72	173	268	464
Black, both sexes.....	2,256	83	2	5	11	32	49	40	26	66	89	118
Male.....	1,174	49	1	4	8	24	42	35	16	38	47	62
Female.....	1,082	34	1	1	3	8	7	5	10	28	42	56
American Indian 1/, both sexes.....	43	-	-	-	-	-	-	1	-	2	2	1
Male.....	25	-	-	-	-	-	-	1	-	1	2	-
Female.....	18	-	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes.....	616	18	4	2	3	3	3	9	11	12	19	25
Male.....	333	7	2	2	2	3	3	7	8	9	13	19
Female.....	283	11	2	-	1	-	-	2	3	3	6	6
<b>Michigan</b>												
All races, both sexes.....	86,721	995	121	87	120	439	630	650	713	1,045	1,574	2,777
Male.....	42,835	558	74	51	71	324	459	457	487	682	976	1,705
Female.....	43,886	437	47	36	49	115	171	193	226	363	598	1,072
White, both sexes.....	73,510	595	74	59	80	300	456	458	445	702	1,146	2,106
Male.....	35,999	341	47	38	48	210	328	329	303	459	726	1,323
Female.....	37,511	254	27	21	32	90	128	129	142	243	420	783
Black, both sexes.....	12,339	370	39	26	38	130	161	186	251	317	389	630
Male.....	6,400	202	24	12	22	107	124	124	173	208	229	362
Female.....	5,939	168	15	14	16	23	37	62	78	109	160	268
American Indian 1/, both sexes.....	424	8	3	1	-	4	6	3	9	17	22	26
Male.....	222	3	1	-	-	3	4	2	6	10	15	13
Female.....	202	5	2	1	-	1	2	1	3	7	7	13
Asian or Pacific Islander 2/, both sexes.....	448	22	5	1	2	5	7	3	8	9	17	15
Male.....	214	12	2	1	1	4	3	2	5	5	6	7
Female.....	234	10	3	-	1	1	4	1	3	4	11	8

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Minnesota</b>												
All races, both sexes.....	37,138	409	65	38	44	161	270	218	230	359	561	940
Male.....	18,179	256	30	22	22	104	197	177	156	215	343	550
Female.....	18,959	153	35	16	22	57	73	41	74	144	218	390
White, both sexes.....	35,348	280	51	28	35	137	218	175	182	289	468	821
Male.....	17,122	169	23	15	16	86	157	140	127	175	283	479
Female.....	18,226	111	28	13	19	51	61	35	55	114	185	342
Black, both sexes.....	971	82	7	6	5	9	32	29	30	44	57	72
Male.....	597	52	2	5	2	6	24	25	19	27	43	42
Female.....	374	30	5	1	3	3	8	4	11	17	14	30
American Indian 1/, both sexes.....	427	18	3	-	3	8	10	9	10	19	23	26
Male.....	245	12	2	-	3	7	7	7	8	10	10	17
Female.....	182	6	1	-	-	1	3	2	2	9	13	9
Asian or Pacific Islander 2/, both sexes.....	392	29	4	4	1	7	10	5	8	7	13	21
Male.....	215	23	3	2	1	5	9	5	2	3	7	12
Female.....	177	6	1	2	-	2	1	-	6	4	6	9
<b>Mississippi</b>												
All races, both sexes.....	28,255	467	92	48	63	216	318	311	324	460	732	1,080
Male.....	14,481	250	48	30	42	150	239	232	229	279	488	675
Female.....	13,774	217	44	18	21	66	79	79	95	181	244	405
White, both sexes.....	18,896	165	42	24	34	123	168	159	175	244	397	577
Male.....	9,555	95	21	17	23	85	125	122	125	147	267	367
Female.....	9,341	70	21	7	11	38	43	37	50	97	130	210
Black, both sexes.....	9,203	290	49	24	28	90	149	149	146	211	329	493
Male.....	4,837	148	26	13	18	63	114	108	101	130	219	302
Female.....	4,366	142	23	11	10	27	35	41	45	81	110	191
American Indian 1/, both sexes.....	96	9	1	-	1	2	-	1	3	5	3	6
Male.....	59	5	1	-	1	1	-	1	3	2	2	5
Female.....	37	4	-	-	-	1	-	-	-	3	1	1
Asian or Pacific Islander 2/, both sexes.....	60	3	-	-	-	1	1	2	-	-	3	4
Male.....	30	2	-	-	-	1	-	1	-	-	-	1
Female.....	30	1	-	-	-	-	1	1	-	-	3	3



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Nebraska</b>												
All races, both sexes.....	15,263	182	27	19	24	84	102	83	91	142	251	359
Male.....	7,271	95	16	7	10	52	80	60	57	86	158	217
Female.....	7,992	87	11	12	14	32	22	23	34	56	93	142
White, both sexes.....	14,545	144	21	18	20	77	86	73	77	121	224	317
Male.....	6,916	74	12	7	9	47	67	52	48	74	144	197
Female.....	7,629	70	9	11	11	30	19	21	29	47	80	120
Black, both sexes.....	554	28	2	1	3	5	12	6	7	12	26	32
Male.....	270	16	2	-	1	4	11	5	4	8	13	14
Female.....	284	12	-	1	2	1	1	1	3	4	13	18
American Indian 1/, both sexes.....	110	7	4	-	-	2	3	4	2	8	1	8
Male.....	58	3	2	-	-	1	2	3	1	3	1	6
Female.....	52	4	2	-	-	1	1	1	1	5	-	2
Asian or Pacific Islander 2/, both sexes.....	54	3	-	-	1	-	1	-	5	1	-	2
Male.....	27	2	-	-	-	-	-	-	4	1	-	-
Female.....	27	1	-	-	1	-	1	-	1	-	-	2
<b>Nevada</b>												
All races, both sexes.....	18,687	262	54	21	40	116	208	236	214	337	498	702
Male.....	10,295	146	31	12	23	81	154	167	147	220	317	442
Female.....	8,392	116	23	9	17	35	54	69	67	117	181	260
White, both sexes.....	16,519	201	43	18	30	92	175	178	175	275	397	592
Male.....	9,107	112	25	12	17	60	132	127	121	179	257	380
Female.....	7,412	89	18	6	13	32	43	51	54	96	140	212
Black, both sexes.....	1,355	47	10	2	5	21	28	44	28	42	68	71
Male.....	752	27	6	-	4	18	18	30	18	26	38	44
Female.....	603	20	4	2	1	3	10	14	10	16	30	27
American Indian 1/, both sexes.....	164	1	-	-	-	2	-	4	5	12	10	14
Male.....	97	-	-	-	-	2	-	3	2	7	8	11
Female.....	67	1	-	-	-	-	-	1	3	5	2	3
Asian or Pacific Islander 2/, both sexes.....	649	13	1	1	5	1	5	10	6	8	23	25
Male.....	339	7	-	-	2	1	4	7	6	8	14	7
Female.....	310	6	1	1	3	-	1	3	-	-	9	18



## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
New Hampshire												
All races, both sexes.....	10,303	76	11	10	12	40	69	53	59	94	186	285
Male.....	4,947	42	6	6	7	24	47	33	39	54	118	178
Female.....	5,356	34	5	4	5	16	22	20	20	40	68	107
White, both sexes.....	10,229	71	11	10	11	40	68	52	57	93	184	280
Male.....	4,910	39	6	6	6	24	46	32	38	53	118	175
Female.....	5,319	32	5	4	5	16	22	20	19	40	66	105
Black, both sexes.....	34	3	-	-	1	-	1	1	1	-	1	4
Male.....	20	1	-	-	1	-	1	1	1	-	-	2
Female.....	14	2	-	-	-	-	-	-	-	-	1	2
American Indian 1/, both sexes.....	5	-	-	-	-	-	-	-	-	-	1	1
Male.....	3	-	-	-	-	-	-	-	-	-	-	1
Female.....	2	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	35	2	-	-	-	-	-	-	1	1	-	-
Male.....	14	2	-	-	-	-	-	-	-	1	-	-
Female.....	21	-	-	-	-	-	-	-	1	-	-	-
New Jersey												
All races, both sexes.....	69,662	601	91	61	81	264	445	453	493	791	1,278	1,999
Male.....	33,211	344	47	34	43	199	333	339	326	481	787	1,207
Female.....	36,451	257	44	27	38	65	112	114	167	310	491	792
White, both sexes.....	59,371	342	52	43	52	162	298	318	319	527	895	1,455
Male.....	28,059	190	27	23	28	120	223	239	212	335	587	921
Female.....	31,312	152	25	20	24	42	75	79	107	192	308	534
Black, both sexes.....	8,919	234	37	13	22	95	130	122	154	244	355	486
Male.....	4,457	136	18	9	11	74	99	94	101	130	190	258
Female.....	4,462	98	19	4	11	21	31	28	53	114	165	228
American Indian 1/, both sexes.....	57	-	-	-	-	-	-	-	-	-	1	3
Male.....	24	-	-	-	-	-	-	-	-	-	1	3
Female.....	33	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1,315	25	2	5	7	7	17	13	20	20	27	55
Male.....	671	18	2	2	4	5	11	6	13	16	9	25
Female.....	644	7	-	3	3	2	6	7	7	4	18	30

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
New Mexico												
All races, both sexes.....	15,482	192	48	17	25	140	186	199	212	258	398	591
Male.....	8,277	115	22	7	18	108	143	143	138	188	274	389
Female.....	7,205	77	26	10	7	32	43	56	74	70	124	202
White, both sexes.....	14,016	153	34	13	18	108	132	148	162	201	326	506
Male.....	7,425	92	16	4	13	84	103	104	101	145	219	337
Female.....	6,591	61	18	9	5	24	29	44	61	56	107	169
Black, both sexes.....	251	8	2	-	1	6	7	2	4	2	12	7
Male.....	164	5	-	-	1	5	6	2	3	1	10	4
Female.....	87	3	2	-	-	1	1	-	1	1	2	3
American Indian 1/, both sexes.....	1,126	30	12	4	6	24	44	49	45	54	58	74
Male.....	654	17	6	3	4	18	32	37	34	41	44	44
Female.....	472	13	6	1	2	6	12	12	11	13	14	30
Asian or Pacific Islander 2/, both sexes.....	89	1	-	-	-	2	3	-	1	1	2	4
Male.....	34	1	-	-	-	1	2	-	-	1	1	4
Female.....	55	-	-	-	-	1	1	-	1	-	1	-
New York												
All races, both sexes.....	147,680	1,412	213	140	166	551	875	931	1,001	1,593	2,740	4,236
Male.....	70,589	782	116	82	99	394	652	683	668	985	1,712	2,545
Female.....	77,091	630	97	58	67	157	223	248	333	608	1,028	1,691
White, both sexes.....	122,670	861	140	91	113	359	597	639	681	1,042	1,877	2,967
Male.....	58,320	476	78	54	69	240	449	486	476	671	1,210	1,857
Female.....	64,350	385	62	37	44	119	148	153	205	371	667	1,110
Black, both sexes.....	21,215	482	61	43	45	170	233	264	269	490	758	1,124
Male.....	10,211	270	30	26	28	138	174	181	166	269	433	597
Female.....	11,004	212	31	17	17	32	59	83	103	221	325	527
American Indian 1/, both sexes.....	244	7	-	-	1	3	3	-	7	4	4	19
Male.....	120	4	-	-	1	1	2	-	5	4	3	12
Female.....	124	3	-	-	-	2	1	-	2	-	1	7
Asian or Pacific Islander 2/, both sexes.....	3,551	62	12	6	7	19	42	28	44	57	101	126
Male.....	1,938	32	8	2	1	15	27	16	21	41	66	79
Female.....	1,613	30	4	4	6	4	15	12	23	16	35	47



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Ohio												
All races, both sexes.....	106,534	1,160	190	91	98	471	731	762	836	1,152	1,892	3,186
Male.....	51,741	651	113	48	62	340	541	552	556	764	1,182	1,961
Female.....	54,793	509	77	43	36	131	190	210	280	388	710	1,225
White, both sexes.....	94,197	768	136	74	73	366	565	564	660	937	1,566	2,612
Male.....	45,383	433	78	39	45	257	406	406	437	624	982	1,649
Female.....	48,814	335	58	35	28	109	159	158	223	313	584	963
Black, both sexes.....	11,899	382	51	17	24	103	160	194	170	199	313	562
Male.....	6,128	213	34	9	16	82	132	143	116	127	194	304
Female.....	5,771	169	17	8	8	21	28	51	54	72	119	258
American Indian 1/, both sexes.....	68	-	-	-	-	-	-	-	-	4	3	1
Male.....	42	-	-	-	-	-	-	-	-	4	2	1
Female.....	26	-	-	-	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	370	10	3	-	1	2	6	4	6	12	10	11
Male.....	188	5	1	-	1	1	3	3	3	9	4	7
Female.....	182	5	2	-	-	1	3	1	3	3	6	4
Oklahoma												
All races, both sexes.....	36,032	469	84	61	53	210	309	334	369	477	707	1,289
Male.....	17,921	249	49	38	38	153	217	232	243	293	419	789
Female.....	18,111	220	35	23	15	57	92	102	126	184	288	500
White, both sexes.....	31,357	307	61	39	38	155	229	258	271	372	543	1,008
Male.....	15,481	157	37	25	27	112	156	177	172	236	322	624
Female.....	15,876	150	24	14	11	43	73	81	99	136	221	384
Black, both sexes.....	2,269	90	14	11	10	30	41	27	44	38	67	127
Male.....	1,180	56	7	9	7	24	30	23	35	17	38	82
Female.....	1,089	34	7	2	3	6	11	4	9	21	29	45
American Indian 1/, both sexes.....	2,256	68	8	10	5	24	34	48	52	66	93	143
Male.....	1,195	36	4	4	4	17	26	32	34	39	57	80
Female.....	1,061	32	4	6	1	7	8	16	18	27	36	63
Asian or Pacific Islander 2/, both sexes.....	150	4	1	1	-	1	5	1	2	1	4	11
Male.....	65	-	1	-	-	-	5	-	2	1	2	3
Female.....	85	4	-	1	-	1	-	1	-	-	2	8

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Oregon</b>												
All races, both sexes.....	31,403	284	45	29	37	120	206	206	240	322	511	901
Male.....	15,669	145	30	18	20	80	148	154	176	210	327	590
Female.....	15,734	139	15	11	17	40	58	52	64	112	184	311
White, both sexes.....	30,245	254	43	25	34	108	186	188	218	300	476	839
Male.....	15,041	128	28	14	18	72	136	143	160	197	306	550
Female.....	15,204	126	15	11	16	36	50	45	58	103	170	289
Black, both sexes.....	391	12	2	-	-	4	7	8	8	10	13	23
Male.....	232	7	2	-	-	3	5	7	6	7	11	15
Female.....	159	5	-	-	-	1	2	1	2	3	2	8
American Indian 1/, both sexes.....	309	8	-	2	2	7	7	3	8	5	13	17
Male.....	159	4	-	2	2	5	4	1	7	3	7	11
Female.....	150	4	-	-	-	2	3	2	1	2	6	6
Asian or Pacific Islander 2/, both sexes.....	458	10	-	2	1	1	6	7	6	7	9	22
Male.....	237	6	-	2	-	-	3	3	3	3	3	14
Female.....	221	4	-	-	1	1	3	4	3	4	6	8
<b>Pennsylvania</b>												
All races, both sexes.....	125,104	1,139	165	96	127	527	855	866	811	1,183	1,936	3,238
Male.....	60,189	641	108	56	74	385	665	632	556	714	1,235	1,987
Female.....	64,915	498	57	40	53	142	190	234	255	469	701	1,251
White, both sexes.....	112,633	721	128	68	100	378	652	665	612	934	1,570	2,616
Male.....	53,864	404	82	39	58	258	502	480	427	564	1,021	1,629
Female.....	58,769	317	46	29	42	120	150	185	185	370	549	987
Black, both sexes.....	11,751	394	32	26	26	145	193	187	188	240	342	601
Male.....	5,937	224	23	15	15	125	157	143	122	145	198	343
Female.....	5,814	170	9	11	11	20	36	44	66	95	144	258
American Indian 1/, both sexes.....	44	-	-	-	1	-	1	2	-	-	5	1
Male.....	27	-	-	-	1	-	1	2	-	-	3	1
Female.....	17	-	-	-	-	-	-	-	-	-	2	-
Asian or Pacific Islander 2/, both sexes.....	676	24	5	2	-	4	9	12	11	9	19	20
Male.....	361	13	3	2	-	2	5	7	7	5	13	14
Female.....	315	11	2	-	-	2	4	5	4	4	6	6

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Rhode Island</b>												
All races, both sexes.....	9,723	91	5	2	9	31	41	43	61	86	150	229
Male.....	4,493	54	2	1	4	16	27	30	37	60	100	139
Female.....	5,230	37	3	1	5	15	14	13	24	26	50	90
White, both sexes.....	9,322	68	5	2	7	29	37	38	56	76	128	207
Male.....	4,268	40	2	1	3	15	24	25	34	52	85	123
Female.....	5,054	28	3	1	4	14	13	13	22	24	43	84
Black, both sexes.....	311	20	-	-	1	2	1	5	4	8	19	19
Male.....	174	11	-	-	1	1	1	5	2	6	12	13
Female.....	137	9	-	-	-	1	-	-	2	2	7	6
American Indian 1/, both sexes.....	30	1	-	-	-	-	1	-	1	-	-	1
Male.....	12	1	-	-	-	-	-	-	1	-	-	1
Female.....	18	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	60	2	-	-	1	-	2	-	-	2	3	2
Male.....	39	2	-	-	-	-	2	-	-	2	3	2
Female.....	21	-	-	-	1	-	-	-	-	-	-	-
<b>South Carolina</b>												
All races, both sexes.....	39,439	539	94	38	72	259	393	376	444	584	992	1,393
Male.....	19,941	329	46	17	46	184	298	273	284	360	581	841
Female.....	19,498	210	48	21	26	75	95	103	160	224	411	552
White, both sexes.....	28,275	242	46	18	40	146	243	223	263	352	622	845
Male.....	14,268	149	27	10	24	97	189	155	172	224	375	525
Female.....	14,007	93	19	8	16	49	54	68	91	128	247	320
Black, both sexes.....	11,003	291	47	20	32	110	148	150	176	228	366	535
Male.....	5,587	178	18	7	22	85	107	116	108	132	204	308
Female.....	5,416	113	29	13	10	25	41	34	68	96	162	227
American Indian 1/, both sexes.....	56	1	1	-	-	1	-	-	3	-	1	3
Male.....	26	-	1	-	-	1	-	-	2	-	-	2
Female.....	30	1	-	-	-	-	-	-	1	-	1	1
Asian or Pacific Islander 2/, both sexes.....	105	5	-	-	-	2	2	3	2	4	3	10
Male.....	60	2	-	-	-	1	2	2	2	4	2	6
Female.....	45	3	-	-	-	1	-	1	-	-	1	4

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>South Dakota</b>												
All races, both sexes.....	6,826	79	18	12	11	48	44	59	38	46	100	174
Male.....	3,443	45	12	5	6	29	32	47	20	27	61	109
Female.....	3,383	34	6	7	5	19	12	12	18	19	39	65
White, both sexes.....	6,273	54	12	11	8	27	29	44	29	25	61	134
Male.....	3,167	30	9	5	5	14	23	34	15	16	37	85
Female.....	3,106	24	3	6	3	13	6	10	14	9	24	49
Black, both sexes.....	28	3	-	-	2	-	1	3	2	-	2	4
Male.....	18	1	-	-	1	-	-	3	1	-	1	4
Female.....	10	2	-	-	1	-	1	-	1	-	1	-
American Indian 1/, both sexes.....	509	22	6	1	1	21	14	11	7	21	37	36
Male.....	252	14	3	-	-	15	9	9	4	11	23	20
Female.....	257	8	3	1	1	6	5	2	3	10	14	16
Asian or Pacific Islander 2/, both sexes.....	16	-	-	-	-	-	-	1	-	-	-	-
Male.....	6	-	-	-	-	-	-	1	-	-	-	-
Female.....	10	-	-	-	-	-	-	-	-	-	-	-
<b>Tennessee</b>												
All races, both sexes.....	57,087	721	110	41	74	346	489	564	586	846	1,385	2,082
Male.....	28,578	424	59	19	48	247	355	400	400	546	836	1,296
Female.....	28,509	297	51	22	26	99	134	164	186	300	549	786
White, both sexes.....	48,475	421	78	32	51	269	362	403	420	644	1,074	1,602
Male.....	24,197	265	39	16	32	185	261	281	293	411	648	1,006
Female.....	24,278	156	39	16	19	84	101	122	127	233	426	596
Black, both sexes.....	8,370	295	32	9	23	73	120	155	158	199	303	463
Male.....	4,252	155	20	3	16	58	90	115	103	133	183	281
Female.....	4,118	140	12	6	7	15	30	40	55	66	120	182
American Indian 1/, both sexes.....	44	-	-	-	-	-	1	5	-	1	2	4
Male.....	30	-	-	-	-	-	1	4	-	-	1	3
Female.....	14	-	-	-	-	-	-	1	-	1	1	1
Asian or Pacific Islander 2/, both sexes.....	198	5	-	-	-	4	6	1	8	2	6	13
Male.....	99	4	-	-	-	4	3	-	4	2	4	6
Female.....	99	1	-	-	-	-	3	1	4	-	2	7

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>All causes</b>												
<b>Texas</b>												
All races, both sexes.....	160,548	2,564	497	265	296	1,106	1,783	1,780	1,834	2,516	3,816	6,009
Male.....	81,630	1,401	291	150	173	783	1,380	1,267	1,264	1,608	2,402	3,741
Female.....	78,918	1,163	206	115	123	323	403	513	570	908	1,414	2,268
White, both sexes.....	138,866	1,934	397	213	237	929	1,484	1,421	1,461	2,001	3,028	4,813
Male.....	70,498	1,064	225	119	136	648	1,141	1,009	1,025	1,314	1,919	3,050
Female.....	68,368	870	172	94	101	281	343	412	436	687	1,109	1,763
Black, both sexes.....	19,568	573	84	42	53	149	269	331	345	468	714	1,101
Male.....	10,050	309	54	25	33	115	218	238	224	262	435	635
Female.....	9,518	264	30	17	20	34	51	93	121	206	279	466
American Indian 1/, both sexes.....	260	3	1	1	-	4	5	4	3	4	5	20
Male.....	139	2	1	1	-	2	2	2	2	3	-	13
Female.....	121	1	-	-	-	2	3	2	1	1	5	7
Asian or Pacific Islander 2/, both sexes.....	1,854	54	15	9	6	24	25	24	25	43	69	75
Male.....	943	26	11	5	4	18	19	18	13	29	48	43
Female.....	911	28	4	4	2	6	6	6	12	14	21	32
<b>Utah</b>												
All races, both sexes.....	14,143	280	62	44	21	124	217	188	245	242	306	472
Male.....	7,192	154	33	27	14	98	153	142	173	145	187	283
Female.....	6,951	126	29	17	7	26	64	46	72	97	119	189
White, both sexes.....	13,647	259	60	41	20	113	198	180	226	226	289	442
Male.....	6,904	145	31	26	13	89	139	136	159	132	178	264
Female.....	6,743	114	29	15	7	24	59	44	67	94	111	178
Black, both sexes.....	95	4	-	1	-	1	3	1	3	4	6	7
Male.....	60	2	-	1	-	1	2	1	3	4	3	6
Female.....	35	2	-	-	-	-	1	-	-	-	3	1
American Indian 1/, both sexes.....	177	8	1	1	1	7	11	7	8	6	8	15
Male.....	104	3	1	-	1	5	8	5	7	4	5	7
Female.....	73	5	-	1	-	2	3	2	1	2	3	8
Asian or Pacific Islander 2/, both sexes.....	224	9	1	1	-	3	5	-	8	6	3	8
Male.....	124	4	1	-	-	3	4	-	4	5	1	6
Female.....	100	5	-	1	-	-	1	-	4	1	2	2



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
All causes												
Vermont												
All races, both sexes.....	5,179	33	6	-	6	16	33	30	33	44	87	128
Male.....	2,474	19	1	-	3	13	27	21	21	26	48	71
Female.....	2,705	14	5	-	3	3	6	9	12	18	39	57
White, both sexes.....	5,147	30	6	-	6	16	31	30	33	44	87	127
Male.....	2,457	18	1	-	3	13	25	21	21	26	48	70
Female.....	2,690	12	5	-	3	3	6	9	12	18	39	57
Black, both sexes.....	11	1	-	-	-	-	1	-	-	-	-	1
Male.....	6	1	-	-	-	-	1	-	-	-	-	1
Female.....	5	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	7	-	-	-	-	-	-	-	-	-	-	-
Male.....	3	-	-	-	-	-	-	-	-	-	-	-
Female.....	4	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	14	2	-	-	-	-	1	-	-	-	-	-
Male.....	8	-	-	-	-	-	1	-	-	-	-	-
Female.....	6	2	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes.....	58,225	848	104	64	91	283	504	533	493	734	1,205	1,895
Male.....	28,717	479	55	34	53	195	388	386	334	452	717	1,135
Female.....	29,508	369	49	30	38	88	116	147	159	282	488	760
White, both sexes.....	45,227	444	77	42	52	186	344	345	333	489	807	1,287
Male.....	22,079	255	43	22	28	116	260	247	230	304	487	799
Female.....	23,148	189	34	20	24	70	84	98	103	185	320	488
Black, both sexes.....	12,032	375	24	20	32	92	145	179	144	222	362	571
Male.....	6,135	209	10	11	20	76	114	133	94	136	211	316
Female.....	5,897	166	14	9	12	16	31	46	50	86	151	255
American Indian 1/, both sexes.....	47	3	-	-	-	-	3	1	-	2	2	1
Male.....	25	2	-	-	-	-	3	1	-	1	2	-
Female.....	22	1	-	-	-	-	-	-	-	1	-	1
Asian or Pacific Islander 2/, both sexes.....	919	26	3	2	7	5	12	8	16	21	34	36
Male.....	478	13	2	1	5	3	11	5	10	11	17	20
Female.....	441	13	1	1	2	2	1	3	6	10	17	16





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All causes

United States

All races, both sexes.....	106,948	132,458	154,652	174,991	214,247	289,029	363,653	359,224	240,180	94,477	19,766	201
Male.....	66,552	81,590	92,028	100,492	117,852	149,669	171,134	146,114	77,477	22,102	3,173	156
Female.....	40,396	50,868	62,624	74,499	96,395	139,360	192,519	213,110	162,703	72,375	16,593	45
White, both sexes.....	82,598	104,284	127,003	145,663	181,714	253,164	325,958	327,592	220,511	86,150	17,268	148
Male.....	52,109	64,772	76,013	84,199	100,629	132,199	154,625	134,013	71,426	20,093	2,695	122
Female.....	30,489	39,512	50,990	61,464	81,085	120,965	171,333	193,579	149,085	66,057	14,573	26
Black, both sexes.....	21,209	24,190	23,492	24,555	26,829	29,063	29,923	24,885	15,419	6,777	2,113	45
Male.....	12,525	14,468	13,659	13,587	14,085	14,001	12,814	9,060	4,421	1,420	354	28
Female.....	8,684	9,722	9,833	10,968	12,744	15,062	17,109	15,825	10,998	5,357	1,759	17
American Indian 1/, both sexes.....	1,021	1,186	1,181	1,222	1,314	1,351	1,195	958	540	211	62	-
Male.....	631	700	672	644	713	671	543	397	192	55	23	-
Female.....	390	486	509	578	601	680	652	561	348	156	39	-
Asian or Pacific Islander 2/, both sexes.....	2,120	2,798	2,976	3,551	4,390	5,451	6,577	5,789	3,710	1,339	323	8
Male.....	1,287	1,650	1,684	2,062	2,425	2,798	3,152	2,644	1,438	534	101	6
Female.....	833	1,148	1,292	1,489	1,965	2,653	3,425	3,145	2,272	805	222	2

Alabama

All races, both sexes.....	2,272	2,890	3,338	3,863	4,620	5,604	6,573	6,035	3,732	1,361	318	2
Male.....	1,413	1,808	1,996	2,260	2,607	2,801	2,957	2,323	1,101	301	47	2
Female.....	859	1,082	1,342	1,603	2,013	2,803	3,616	3,712	2,631	1,060	271	-
White, both sexes.....	1,486	1,987	2,431	2,946	3,546	4,481	5,346	5,039	3,054	1,060	204	2
Male.....	943	1,237	1,468	1,732	2,027	2,294	2,453	1,971	881	235	24	2
Female.....	543	750	963	1,214	1,519	2,187	2,893	3,068	2,173	825	180	-
Black, both sexes.....	773	892	893	905	1,062	1,109	1,216	988	668	296	113	-
Male.....	461	563	520	524	573	500	500	351	216	65	23	-
Female.....	312	329	373	381	489	609	716	637	452	231	90	-
American Indian 1/, both sexes.....	3	6	4	4	4	4	3	1	3	3	1	-
Male.....	2	4	3	2	2	3	-	-	2	-	-	-
Female.....	1	2	1	2	2	1	3	1	1	3	1	-
Asian or Pacific Islander 2/, both sexes.....	10	5	10	8	8	10	8	7	7	2	-	-
Male.....	7	4	5	2	5	4	4	1	2	1	-	-
Female.....	3	1	5	6	3	6	4	6	5	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All causes

Alaska

All races, both sexes.....	265	299	271	278	310	387	334	295	173	57	9	-
Male.....	171	186	174	150	187	208	170	129	52	15	2	-
Female.....	94	113	97	128	123	179	164	166	121	42	7	-
White, both sexes.....	180	218	205	209	206	271	266	231	142	45	7	-
Male.....	117	141	142	124	123	146	131	105	43	10	1	-
Female.....	63	77	63	85	83	125	135	126	99	35	6	-
Black, both sexes.....	13	11	9	4	7	10	5	6	4	2	-	-
Male.....	10	6	7	1	6	6	3	1	1	1	-	-
Female.....	3	5	2	3	1	4	2	5	3	1	-	-
American Indian 1/, both sexes.....	63	59	48	56	82	93	56	49	21	9	2	-
Male.....	39	35	19	22	50	48	32	19	5	3	1	-
Female.....	24	24	29	34	32	45	24	30	16	6	1	-
Asian or Pacific Islander 2/, both sexes.....	9	11	9	9	15	13	7	9	6	1	-	-
Male.....	5	4	6	3	8	8	4	4	3	1	-	-
Female.....	4	7	3	6	7	5	3	5	3	-	-	-

Arizona

All races, both sexes.....	1,976	2,423	3,000	3,396	4,122	5,482	6,666	6,467	4,139	1,508	286	30
Male.....	1,258	1,544	1,793	1,983	2,397	2,988	3,193	2,773	1,450	410	52	24
Female.....	718	879	1,207	1,413	1,725	2,494	3,473	3,694	2,689	1,098	234	6
White, both sexes.....	1,737	2,162	2,763	3,141	3,846	5,226	6,367	6,239	3,975	1,446	267	29
Male.....	1,104	1,383	1,664	1,844	2,248	2,862	3,069	2,682	1,403	391	46	24
Female.....	633	779	1,099	1,297	1,598	2,364	3,298	3,557	2,572	1,055	221	5
Black, both sexes.....	110	111	98	100	102	97	107	95	68	25	10	1
Male.....	67	69	58	56	60	44	41	37	18	8	2	-
Female.....	43	42	40	44	42	53	66	58	50	17	8	1
American Indian 1/, both sexes.....	110	119	104	120	135	124	144	100	72	32	7	-
Male.....	75	74	54	67	73	64	67	43	23	9	3	-
Female.....	35	45	50	53	62	60	77	57	49	23	4	-
Asian or Pacific Islander 2/, both sexes.....	19	31	35	35	39	35	48	33	24	5	2	-
Male.....	12	18	17	16	16	18	16	11	6	2	1	-
Female.....	7	13	18	19	23	17	32	22	18	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>Arkansas</b>												
All races, both sexes.....	1,306	1,597	1,995	2,266	2,671	3,381	4,053	3,738	2,477	1,023	223	1
Male.....	820	966	1,158	1,307	1,531	1,827	1,854	1,422	778	222	41	1
Female.....	486	631	837	959	1,140	1,554	2,199	2,316	1,699	801	182	-
White, both sexes.....	981	1,257	1,707	1,986	2,349	3,021	3,641	3,370	2,195	894	175	-
Male.....	632	763	985	1,140	1,355	1,655	1,681	1,308	695	200	32	-
Female.....	349	494	722	846	994	1,366	1,960	2,062	1,500	694	143	-
Black, both sexes.....	317	327	275	271	308	341	397	357	273	125	47	1
Male.....	181	197	167	162	170	163	167	112	79	21	9	1
Female.....	136	130	108	109	138	178	230	245	194	104	38	-
American Indian 1/, both sexes.....	5	4	9	5	4	5	7	7	4	1	-	-
Male.....	5	1	5	3	2	2	4	1	1	-	-	-
Female.....	-	3	4	2	2	3	3	6	3	1	-	-
Asian or Pacific Islander 2/, both sexes.....	3	9	4	4	10	14	8	4	5	3	1	-
Male.....	2	5	1	2	4	7	2	1	3	1	-	-
Female.....	1	4	3	2	6	7	6	3	2	2	1	-
<b>California</b>												
All races, both sexes.....	10,978	13,109	14,629	15,697	19,659	26,800	34,986	35,025	23,405	9,352	2,008	43
Male.....	6,949	8,121	8,678	9,093	10,717	13,943	16,480	15,101	8,161	2,561	370	34
Female.....	4,029	4,988	5,951	6,604	8,942	12,857	18,506	19,924	15,244	6,791	1,638	9
White, both sexes.....	8,596	10,310	11,869	12,620	15,884	22,511	30,057	30,609	20,663	8,232	1,708	33
Male.....	5,524	6,512	7,053	7,327	8,635	11,756	14,178	13,169	7,171	2,207	306	27
Female.....	3,072	3,798	4,816	5,293	7,249	10,755	15,879	17,440	13,492	6,025	1,402	6
Black, both sexes.....	1,419	1,662	1,534	1,618	1,836	1,810	1,859	1,643	1,015	474	141	4
Male.....	826	934	903	919	999	886	818	631	308	108	18	2
Female.....	593	728	631	699	837	924	1,041	1,012	707	366	123	2
American Indian 1/, both sexes.....	110	120	124	121	132	134	112	123	57	26	5	-
Male.....	71	80	72	65	87	60	51	50	25	5	2	-
Female.....	39	40	52	56	45	74	61	73	32	21	3	-
Asian or Pacific Islander 2/, both sexes.....	853	1,017	1,102	1,338	1,807	2,345	2,958	2,650	1,670	620	154	6
Male.....	528	595	650	782	996	1,241	1,433	1,251	657	241	44	5
Female.....	325	422	452	556	811	1,104	1,525	1,399	1,013	379	110	1

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## All causes

## Colorado

All races, both sexes.....	1,522	1,725	1,857	2,009	2,488	3,450	4,221	4,500	2,973	1,143	231	-
Male.....	974	1,069	1,120	1,126	1,371	1,824	1,978	1,864	985	273	45	-
Female.....	548	656	737	883	1,117	1,626	2,243	2,636	1,988	870	186	-
White, both sexes.....	1,406	1,608	1,752	1,911	2,351	3,295	4,082	4,363	2,898	1,111	222	-
Male.....	897	993	1,057	1,077	1,297	1,757	1,928	1,802	952	268	43	-
Female.....	509	615	695	834	1,054	1,538	2,154	2,561	1,946	843	179	-
Black, both sexes.....	86	88	76	61	100	110	84	79	51	23	7	-
Male.....	56	59	48	35	54	49	29	34	23	4	1	-
Female.....	30	29	28	26	46	61	55	45	28	19	6	-
American Indian 1/, both sexes.....	17	15	22	19	18	13	14	12	8	3	2	-
Male.....	11	9	12	6	10	7	7	6	4	-	1	-
Female.....	6	6	10	13	8	6	7	6	4	3	1	-
Asian or Pacific Islander 2/, both sexes.....	13	14	7	18	19	32	41	46	16	6	-	-
Male.....	10	8	3	8	10	11	14	22	6	1	-	-
Female.....	3	6	4	10	9	21	27	24	10	5	-	-

## Connecticut

All races, both sexes.....	995	1,256	1,577	1,687	2,170	3,317	4,562	5,156	3,569	1,542	319	2
Male.....	628	776	875	935	1,180	1,736	2,177	2,103	1,136	377	44	1
Female.....	367	480	702	752	990	1,581	2,385	3,053	2,433	1,165	275	1
White, both sexes.....	831	1,071	1,378	1,475	1,942	3,093	4,326	4,963	3,466	1,492	309	2
Male.....	523	653	765	816	1,058	1,616	2,067	2,038	1,112	365	42	1
Female.....	308	418	613	659	884	1,477	2,259	2,925	2,354	1,127	267	1
Black, both sexes.....	142	171	179	200	208	196	212	171	92	44	10	-
Male.....	90	113	97	111	111	105	100	58	17	10	2	-
Female.....	52	58	82	89	97	91	112	113	75	34	8	-
American Indian 1/, both sexes.....	1	5	5	-	3	3	1	1	1	2	-	-
Male.....	1	3	4	-	2	1	-	-	-	-	-	-
Female.....	-	2	1	-	1	2	1	1	1	2	-	-
Asian or Pacific Islander 2/, both sexes.....	21	9	15	12	17	25	23	21	10	4	-	-
Male.....	14	7	9	8	9	14	10	7	7	2	-	-
Female.....	7	2	6	4	8	11	13	14	3	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All causes

Delaware

All races, both sexes.....	315	385	461	547	696	944	1,105	1,026	681	240	54	4
Male.....	190	238	291	314	397	473	528	403	221	55	6	2
Female.....	125	147	170	233	299	471	577	623	460	185	48	2
White, both sexes.....	224	287	370	423	569	807	967	936	611	211	47	2
Male.....	143	180	234	240	327	406	470	367	199	49	4	1
Female.....	81	107	136	183	242	401	497	569	412	162	43	1
Black, both sexes.....	89	94	87	114	116	128	131	83	65	28	7	2
Male.....	46	54	56	67	64	64	56	30	18	5	2	1
Female.....	43	40	31	47	52	64	75	53	47	23	5	1
American Indian 1/, both sexes.....	1	1	-	4	6	3	4	2	3	1	-	-
Male.....	1	1	-	4	3	1	1	1	2	1	-	-
Female.....	-	-	-	-	3	2	3	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.	1	3	4	6	5	6	3	5	2	-	-	-
Male.....	-	3	1	3	3	2	1	5	2	-	-	-
Female.....	1	-	3	3	2	4	2	-	-	-	-	-

Dist. of Columbia

All races, both sexes.....	343	384	368	442	403	505	570	621	428	204	63	-
Male.....	205	227	224	249	216	257	218	239	147	53	15	-
Female.....	138	157	144	193	187	248	352	382	281	151	48	-
White, both sexes.....	34	53	63	75	71	117	159	207	166	73	26	-
Male.....	25	31	44	42	39	70	78	103	65	21	8	-
Female.....	9	22	19	33	32	47	81	104	101	52	18	-
Black, both sexes.....	306	330	301	363	326	381	402	410	258	129	37	-
Male.....	179	196	178	204	173	184	136	136	81	31	7	-
Female.....	127	134	123	159	153	197	266	274	177	98	30	-
American Indian 1/, both sexes.....	-	-	-	2	-	-	2	1	-	-	-	-
Male.....	-	-	-	2	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	3	1	4	2	6	7	7	3	4	2	-	-
Male.....	1	-	2	1	4	3	4	-	1	1	-	-
Female.....	2	1	2	1	2	4	3	3	3	1	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All causes

Florida

All races, both sexes.....	7,071	8,489	10,197	11,857	14,888	20,710	26,081	25,735	16,962	6,622	1,333	10
Male.....	4,447	5,356	6,175	7,008	8,483	11,273	13,058	11,387	6,211	1,824	263	9
Female.....	2,624	3,133	4,022	4,849	6,405	9,437	13,023	14,348	10,751	4,798	1,070	1
White, both sexes.....	5,580	6,961	8,661	10,164	12,982	18,762	24,152	24,220	16,027	6,252	1,220	8
Male.....	3,591	4,451	5,279	6,036	7,429	10,311	12,193	10,831	5,928	1,733	240	7
Female.....	1,989	2,510	3,382	4,128	5,553	8,451	11,959	13,389	10,099	4,519	980	1
Black, both sexes.....	1,418	1,444	1,436	1,564	1,771	1,791	1,810	1,428	877	346	110	2
Male.....	819	852	837	902	988	889	807	521	261	78	21	2
Female.....	599	592	599	662	783	902	1,003	907	616	268	89	-
American Indian 1/, both sexes.....	10	17	24	15	10	18	15	10	6	1	1	-
Male.....	6	11	18	9	4	9	7	7	2	1	1	-
Female.....	4	6	6	6	6	9	8	3	4	-	-	-
Asian or Pacific Islander 2/, both sexes.....	63	67	76	114	125	139	104	77	52	23	2	-
Male.....	31	42	41	61	62	64	51	28	20	12	1	-
Female.....	32	25	35	53	63	75	53	49	32	11	1	-

Georgia

All races, both sexes.....	3,633	4,461	5,123	5,679	6,260	7,473	8,945	8,448	5,424	2,297	421	2
Male.....	2,240	2,770	3,084	3,308	3,411	3,817	3,999	3,134	1,596	486	53	1
Female.....	1,393	1,691	2,039	2,371	2,849	3,656	4,946	5,314	3,828	1,811	368	1
White, both sexes.....	2,185	2,822	3,443	4,000	4,568	5,735	7,236	6,865	4,417	1,766	271	1
Male.....	1,397	1,775	2,096	2,366	2,529	2,979	3,305	2,578	1,321	358	31	1
Female.....	788	1,047	1,347	1,634	2,039	2,756	3,931	4,287	3,096	1,408	240	-
Black, both sexes.....	1,407	1,597	1,637	1,621	1,635	1,660	1,644	1,534	983	524	147	1
Male.....	822	971	966	909	845	803	661	536	268	123	20	-
Female.....	585	626	671	712	790	857	983	998	715	401	127	1
American Indian 1/, both sexes.....	4	7	7	3	6	8	4	6	-	-	-	-
Male.....	3	5	4	1	4	4	-	4	-	-	-	-
Female.....	1	2	3	2	2	4	4	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	37	35	36	55	51	70	61	43	24	7	3	-
Male.....	18	19	18	32	33	31	33	16	7	5	2	-
Female.....	19	16	18	23	18	39	28	27	17	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All causes

Hawaii

All races, both sexes.....	409	559	554	686	715	1,043	1,415	1,554	1,059	420	90	1
Male.....	269	370	338	413	420	570	753	749	423	167	34	1
Female.....	140	189	216	273	295	473	662	805	636	253	56	-
White, both sexes.....	164	191	191	211	208	272	347	422	259	120	23	1
Male.....	108	137	129	138	123	155	190	209	108	30	7	1
Female.....	56	54	62	73	85	117	157	213	151	90	16	-
Black, both sexes.....	8	11	7	5	4	10	4	9	1	1	1	-
Male.....	7	9	4	4	1	6	1	6	1	1	1	-
Female.....	1	2	3	1	3	4	3	3	-	-	-	-
American Indian 1/, both sexes.....	-	7	2	1	-	1	1	2	2	1	-	-
Male.....	-	5	2	1	-	-	1	2	-	1	-	-
Female.....	-	2	-	-	-	1	-	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	237	350	354	469	503	760	1,063	1,121	797	298	66	-
Male.....	154	219	203	270	296	409	561	532	314	135	26	-
Female.....	83	131	151	199	207	351	502	589	483	163	40	-

Idaho

All races, both sexes.....	416	523	638	742	945	1,305	1,602	1,725	1,167	457	72	-
Male.....	257	313	400	435	501	658	768	734	424	130	11	-
Female.....	159	210	238	307	444	647	834	991	743	327	61	-
White, both sexes.....	402	510	623	729	926	1,285	1,585	1,706	1,159	456	71	-
Male.....	246	308	393	428	490	648	754	723	419	129	10	-
Female.....	156	202	230	301	436	637	831	983	740	327	61	-
Black, both sexes.....	1	3	2	2	2	6	-	2	-	-	-	-
Male.....	-	3	1	1	1	4	-	1	-	-	-	-
Female.....	1	-	1	1	1	2	-	1	-	-	-	-
American Indian 1/, both sexes.....	11	9	10	7	11	6	5	5	4	-	1	-
Male.....	9	2	5	4	7	4	4	1	2	-	1	-
Female.....	2	7	5	3	4	2	1	4	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	1	3	4	6	8	12	12	4	1	-	-
Male.....	2	-	1	2	3	2	10	9	3	1	-	-
Female.....	-	1	2	2	3	6	2	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>Illinois</b>												
All races, both sexes.....	4,196	5,419	6,129	7,037	8,697	12,008	15,452	15,055	10,529	4,252	916	12
Male.....	2,533	3,246	3,524	4,041	4,689	6,098	7,145	5,933	3,258	927	134	9
Female.....	1,663	2,173	2,605	2,996	4,008	5,910	8,307	9,122	7,271	3,325	782	3
White, both sexes.....	3,025	3,911	4,733	5,512	6,996	10,207	13,586	13,682	9,717	3,915	815	6
Male.....	1,855	2,367	2,733	3,188	3,805	5,207	6,337	5,419	3,002	864	114	5
Female.....	1,170	1,544	2,000	2,324	3,191	5,000	7,249	8,263	6,715	3,051	701	1
Black, both sexes.....	1,114	1,406	1,309	1,401	1,564	1,643	1,674	1,224	709	310	93	5
Male.....	648	821	738	778	800	812	719	458	222	56	18	4
Female.....	466	585	571	623	764	831	955	766	487	254	75	1
American Indian 1/, both sexes.....	5	8	3	6	10	9	7	7	3	1	2	-
Male.....	2	5	3	2	6	5	2	1	1	-	1	-
Female.....	3	3	-	4	4	4	5	6	2	1	1	-
Asian or Pacific Islander 2/, both sexes.....	52	94	84	118	127	149	185	142	100	26	6	1
Male.....	28	53	50	73	78	74	87	55	33	7	1	-
Female.....	24	41	34	45	49	75	98	87	67	19	5	1
<b>Indiana</b>												
All races, both sexes.....	2,292	2,940	3,482	4,118	5,071	6,507	8,086	7,868	5,110	1,963	433	4
Male.....	1,444	1,761	2,024	2,253	2,820	3,265	3,792	3,120	1,546	388	53	2
Female.....	848	1,179	1,458	1,865	2,251	3,242	4,294	4,748	3,564	1,575	380	2
White, both sexes.....	1,987	2,590	3,134	3,782	4,662	6,052	7,582	7,487	4,898	1,875	413	2
Male.....	1,258	1,556	1,841	2,087	2,615	3,047	3,563	2,968	1,488	371	51	2
Female.....	729	1,034	1,293	1,695	2,047	3,005	4,019	4,519	3,410	1,504	362	-
Black, both sexes.....	298	343	340	324	401	444	497	374	209	87	20	2
Male.....	182	202	178	161	201	209	225	151	58	17	2	-
Female.....	116	141	162	163	200	235	272	223	151	70	18	2
American Indian 1/, both sexes.....	1	3	3	2	2	1	1	2	-	-	-	-
Male.....	1	1	2	-	-	1	1	-	-	-	-	-
Female.....	-	2	1	2	2	-	-	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	6	4	5	10	6	10	6	5	3	1	-	-
Male.....	3	2	3	5	4	8	3	1	-	-	-	-
Female.....	3	2	2	5	2	2	3	4	3	1	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>Iowa</b>												
All races, both sexes.....	901	1,182	1,437	1,661	2,256	3,259	4,266	4,649	3,606	1,537	352	1
Male.....	584	741	858	951	1,227	1,749	2,087	1,942	1,150	346	42	1
Female.....	317	441	579	710	1,029	1,510	2,179	2,707	2,456	1,191	310	-
White, both sexes.....	863	1,131	1,384	1,625	2,198	3,206	4,214	4,611	3,577	1,531	351	1
Male.....	561	714	829	930	1,198	1,723	2,061	1,927	1,136	345	41	1
Female.....	302	417	555	695	1,000	1,483	2,153	2,684	2,441	1,186	310	-
Black, both sexes.....	30	40	39	27	45	43	41	31	23	4	1	-
Male.....	19	20	22	15	26	21	22	13	9	-	1	-
Female.....	11	20	17	12	19	22	19	18	14	4	-	-
American Indian 1/, both sexes.....	4	4	6	2	6	1	3	3	1	1	-	-
Male.....	1	2	5	2	-	1	2	1	-	-	-	-
Female.....	3	2	1	-	6	-	1	2	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	4	7	8	7	7	9	8	4	5	1	-	-
Male.....	3	5	2	4	3	4	2	1	5	1	-	-
Female.....	1	2	6	3	4	5	6	3	-	-	-	-
<b>Kansas</b>												
All races, both sexes.....	976	1,145	1,308	1,564	1,942	2,848	3,862	3,895	2,955	1,267	281	-
Male.....	611	673	816	886	1,086	1,476	1,883	1,588	952	296	35	-
Female.....	365	472	492	678	856	1,372	1,979	2,307	2,003	971	246	-
White, both sexes.....	864	1,027	1,202	1,450	1,808	2,678	3,689	3,754	2,851	1,236	273	-
Male.....	546	604	758	819	1,020	1,390	1,803	1,537	918	288	34	-
Female.....	318	423	444	631	788	1,288	1,886	2,217	1,933	948	239	-
Black, both sexes.....	92	98	77	80	93	127	129	113	85	26	8	-
Male.....	54	59	44	50	44	65	60	39	27	8	1	-
Female.....	38	39	33	30	49	62	69	74	58	18	7	-
American Indian 1/, both sexes.....	15	14	21	22	25	31	29	20	11	4	-	-
Male.....	8	6	9	12	15	15	14	8	5	-	-	-
Female.....	7	8	12	10	10	16	15	12	6	4	-	-
Asian or Pacific Islander 2/, both sexes.....	5	6	8	12	16	12	15	8	8	1	-	-
Male.....	3	4	5	5	7	6	6	4	2	-	-	-
Female.....	2	2	3	7	9	6	9	4	6	1	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Kentucky												
All races, both sexes.....	1,882	2,445	2,834	3,474	3,994	4,905	5,772	5,194	3,324	1,268	250	-
Male.....	1,183	1,489	1,678	1,966	2,195	2,506	2,661	1,961	1,000	278	39	-
Female.....	699	956	1,156	1,508	1,799	2,399	3,111	3,233	2,324	990	211	-
White, both sexes.....	1,687	2,222	2,629	3,209	3,710	4,624	5,455	4,946	3,145	1,222	226	-
Male.....	1,080	1,349	1,557	1,826	2,042	2,368	2,520	1,866	949	271	37	-
Female.....	607	873	1,072	1,383	1,668	2,256	2,935	3,080	2,196	951	189	-
Black, both sexes.....	187	213	198	255	271	264	306	241	171	46	23	-
Male.....	101	132	118	135	145	130	136	92	47	7	2	-
Female.....	86	81	80	120	126	134	170	149	124	39	21	-
American Indian 1/, both sexes.....	1	5	3	1	-	-	1	2	2	-	-	-
Male.....	-	4	-	-	-	-	1	1	2	-	-	-
Female.....	1	1	3	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	7	5	4	9	13	17	10	5	6	-	1	-
Male.....	2	4	3	5	8	8	4	2	2	-	-	-
Female.....	5	1	1	4	5	9	6	3	4	-	1	-
Louisiana												
All races, both sexes.....	2,103	2,627	2,780	3,242	3,667	4,569	5,528	4,934	3,046	1,153	251	2
Male.....	1,291	1,628	1,632	1,856	1,977	2,270	2,526	1,851	902	239	48	1
Female.....	812	999	1,148	1,386	1,690	2,299	3,002	3,083	2,144	914	203	1
White, both sexes.....	1,242	1,589	1,804	2,192	2,577	3,434	4,351	3,940	2,421	891	165	-
Male.....	792	980	1,074	1,244	1,422	1,751	2,011	1,473	729	178	25	-
Female.....	450	609	730	948	1,155	1,683	2,340	2,467	1,692	713	140	-
Black, both sexes.....	841	1,020	957	1,037	1,059	1,110	1,154	977	613	260	85	2
Male.....	483	639	545	602	544	509	506	365	166	61	23	1
Female.....	358	381	412	435	515	601	648	612	447	199	62	1
American Indian 1/, both sexes.....	4	6	9	4	5	7	5	5	3	1	-	-
Male.....	3	2	7	3	1	4	2	5	2	-	-	-
Female.....	1	4	2	1	4	3	3	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	16	12	10	9	26	18	18	12	9	1	1	-
Male.....	13	7	6	7	10	6	7	8	5	-	-	-
Female.....	3	5	4	2	16	12	11	4	4	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>Maine</b>												
All races, both sexes.....	448	625	809	872	1,126	1,660	1,913	2,022	1,375	533	120	-
Male.....	286	391	478	536	622	879	894	825	469	141	20	-
Female.....	162	234	331	336	504	781	1,019	1,197	906	392	100	-
White, both sexes.....	438	618	801	864	1,117	1,649	1,903	2,020	1,366	530	120	-
Male.....	281	386	473	532	616	869	890	823	466	139	20	-
Female.....	157	232	328	332	501	780	1,013	1,197	900	391	100	-
Black, both sexes.....	5	3	2	-	4	4	5	-	1	2	-	-
Male.....	3	2	1	-	3	3	3	-	-	1	-	-
Female.....	2	1	1	-	1	1	2	-	1	1	-	-
American Indian 1/, both sexes.....	4	2	6	5	4	5	2	2	5	-	-	-
Male.....	1	2	4	3	3	5	1	2	2	-	-	-
Female.....	3	-	2	2	1	-	1	-	3	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	2	-	3	1	2	3	-	3	1	-	-
Male.....	1	1	-	1	-	2	-	-	1	1	-	-
Female.....	-	1	-	2	1	-	3	-	2	-	-	-
<b>Maryland</b>												
All races, both sexes.....	2,100	2,440	2,847	3,090	3,802	5,018	6,293	6,415	4,093	1,597	298	3
Male.....	1,286	1,451	1,707	1,751	2,033	2,537	2,879	2,573	1,283	374	44	2
Female.....	814	989	1,140	1,339	1,769	2,481	3,414	3,842	2,810	1,223	254	1
White, both sexes.....	1,198	1,427	1,816	2,029	2,592	3,784	5,006	5,359	3,458	1,339	233	2
Male.....	742	846	1,102	1,173	1,403	1,953	2,334	2,167	1,120	324	35	2
Female.....	456	581	714	856	1,189	1,831	2,672	3,192	2,338	1,015	198	-
Black, both sexes.....	871	967	975	994	1,128	1,143	1,203	964	591	245	60	1
Male.....	524	572	573	534	581	544	503	365	144	47	7	-
Female.....	347	395	402	460	547	599	700	599	447	198	53	1
American Indian 1/, both sexes.....	1	2	3	5	6	7	7	2	2	1	-	-
Male.....	1	1	1	5	5	3	4	1	-	-	-	-
Female.....	-	1	2	-	1	4	3	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes.....	30	44	53	62	76	84	77	90	42	12	5	-
Male.....	19	32	31	39	44	37	38	40	19	3	2	-
Female.....	11	12	22	23	32	47	39	50	23	9	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All causes

Massachusetts

All races, both sexes.....	1,827	2,513	2,869	3,286	4,256	6,260	8,572	9,278	6,360	2,629	599	4
Male.....	1,149	1,551	1,666	1,874	2,311	3,234	3,922	3,724	2,002	569	82	3
Female.....	678	962	1,203	1,412	1,945	3,026	4,650	5,554	4,358	2,060	517	1
White, both sexes.....	1,658	2,297	2,649	3,042	3,978	5,942	8,269	9,004	6,198	2,563	572	2
Male.....	1,050	1,414	1,544	1,735	2,163	3,065	3,783	3,627	1,947	554	76	2
Female.....	608	883	1,105	1,307	1,815	2,877	4,486	5,377	4,251	2,009	496	-
Black, both sexes.....	139	175	177	181	205	241	212	211	123	48	21	2
Male.....	79	116	97	102	104	126	97	74	40	8	4	1
Female.....	60	59	80	79	101	115	115	137	83	40	17	1
American Indian 1/, both sexes.....	6	2	4	6	4	7	3	3	-	1	1	-
Male.....	3	1	3	4	4	3	1	1	-	-	1	-
Female.....	3	1	1	2	-	4	2	2	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	24	39	39	57	69	70	88	60	39	17	5	-
Male.....	17	20	22	33	40	40	41	22	15	7	1	-
Female.....	7	19	17	24	29	30	47	38	24	10	4	-

Michigan

All races, both sexes.....	4,086	4,820	5,553	6,181	7,499	10,549	13,348	12,958	8,666	3,233	677	-
Male.....	2,507	2,969	3,298	3,514	3,980	5,428	6,337	5,269	2,800	763	126	-
Female.....	1,579	1,851	2,255	2,667	3,519	5,121	7,011	7,689	5,866	2,470	551	-
White, both sexes.....	3,002	3,633	4,450	5,125	6,381	9,224	11,865	11,828	7,998	2,995	588	-
Male.....	1,885	2,256	2,665	2,930	3,422	4,789	5,704	4,809	2,571	711	105	-
Female.....	1,117	1,377	1,785	2,195	2,959	4,435	6,161	7,019	5,427	2,284	483	-
Black, both sexes.....	1,035	1,122	1,032	980	1,038	1,223	1,380	1,060	626	224	82	-
Male.....	596	677	594	541	510	588	597	430	213	51	16	-
Female.....	439	445	438	439	528	635	783	630	413	173	66	-
American Indian 1/, both sexes.....	28	36	30	37	43	55	40	33	16	5	2	-
Male.....	16	20	18	19	26	29	14	17	5	-	1	-
Female.....	12	16	12	18	17	26	26	16	11	5	1	-
Asian or Pacific Islander 2/, both sexes.....	21	29	41	39	37	47	63	37	26	9	5	-
Male.....	10	16	21	24	22	22	22	13	11	1	4	-
Female.....	11	13	20	15	15	25	41	24	15	8	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All causes

Minnesota

All races, both sexes.....	1,357	1,726	2,008	2,338	2,870	4,112	5,741	6,382	4,813	2,057	439	-
Male.....	857	1,098	1,215	1,376	1,636	2,203	2,893	2,689	1,598	478	64	-
Female.....	500	628	793	962	1,234	1,909	2,848	3,693	3,215	1,579	375	-
White, both sexes.....	1,205	1,551	1,875	2,207	2,728	3,982	5,608	6,294	4,746	2,038	430	-
Male.....	759	985	1,133	1,291	1,552	2,138	2,829	2,654	1,577	471	63	-
Female.....	446	566	742	916	1,176	1,844	2,779	3,640	3,169	1,567	367	-
Black, both sexes.....	86	96	63	61	61	68	75	47	32	4	5	-
Male.....	59	68	39	42	39	32	43	20	8	-	-	-
Female.....	27	28	24	19	22	36	32	27	24	4	5	-
American Indian 1/, both sexes.....	49	48	41	38	40	34	23	10	8	6	1	-
Male.....	30	25	24	22	24	20	7	4	2	3	1	-
Female.....	19	23	17	16	16	14	16	6	6	3	-	-
Asian or Pacific Islander 2/, both sexes.....	17	31	29	32	41	28	35	31	27	9	3	-
Male.....	9	20	19	21	21	13	14	11	11	4	-	-
Female.....	8	11	10	11	20	15	21	20	16	5	3	-

Mississippi

All races, both sexes.....	1,508	1,813	2,072	2,308	2,770	3,293	3,757	3,325	2,157	896	236	9
Male.....	956	1,150	1,257	1,335	1,564	1,709	1,706	1,262	643	191	37	9
Female.....	552	663	815	973	1,206	1,584	2,051	2,063	1,514	705	199	-
White, both sexes.....	825	1,060	1,301	1,542	1,908	2,434	2,803	2,532	1,620	635	124	4
Male.....	523	689	807	896	1,127	1,262	1,310	961	448	122	12	4
Female.....	302	371	494	646	781	1,172	1,493	1,571	1,172	513	112	-
Black, both sexes.....	674	746	755	745	847	847	941	782	533	259	111	5
Male.....	426	457	438	427	431	441	391	294	192	69	24	5
Female.....	248	289	317	318	416	406	550	488	341	190	87	-
American Indian 1/, both sexes.....	8	4	12	16	8	4	5	5	1	2	-	-
Male.....	6	3	10	9	3	2	2	2	1	-	-	-
Female.....	2	1	2	7	5	2	3	-	2	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	3	4	5	7	8	8	6	3	-	1	-
Male.....	1	1	2	3	3	4	3	5	2	-	1	-
Female.....	-	2	2	2	4	4	5	1	1	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>Missouri</b>												
All races, both sexes.....	2,304	2,726	3,353	4,091	4,925	6,619	8,104	7,971	5,511	2,225	457	6
Male.....	1,445	1,687	2,030	2,353	2,755	3,434	3,822	3,117	1,724	484	67	5
Female.....	859	1,039	1,323	1,738	2,170	3,185	4,282	4,854	3,787	1,741	390	1
White, both sexes.....	1,924	2,251	2,918	3,626	4,374	5,979	7,428	7,471	5,164	2,090	404	4
Male.....	1,202	1,414	1,755	2,105	2,450	3,126	3,543	2,967	1,638	454	58	4
Female.....	722	837	1,163	1,521	1,924	2,853	3,885	4,504	3,526	1,636	346	-
Black, both sexes.....	362	442	419	439	510	602	638	473	326	133	53	2
Male.....	230	256	265	232	281	286	265	143	81	30	9	1
Female.....	132	186	154	207	229	316	373	330	245	103	44	1
American Indian 1/, both sexes.....	8	6	7	10	17	14	8	6	4	1	-	-
Male.....	5	2	4	6	10	9	1	1	1	-	-	-
Female.....	3	4	3	4	7	5	7	5	3	1	-	-
Asian or Pacific Islander 2/, both sexes.....	10	27	9	16	24	24	30	21	17	1	-	-
Male.....	8	15	6	10	14	13	13	6	4	-	-	-
Female.....	2	12	3	6	10	11	17	15	13	1	-	-
<b>Montana</b>												
All races, both sexes.....	371	427	542	603	749	989	1,263	1,365	965	381	62	-
Male.....	229	281	320	366	400	521	623	605	361	99	13	-
Female.....	142	146	222	237	349	468	640	760	604	282	49	-
White, both sexes.....	344	384	508	561	710	942	1,223	1,341	950	377	61	-
Male.....	214	254	305	345	379	494	606	591	357	98	13	-
Female.....	130	130	203	216	331	448	617	750	593	279	48	-
Black, both sexes.....	2	-	1	2	1	2	1	1	-	-	-	-
Male.....	2	-	1	2	1	2	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes.....	24	43	33	36	37	45	31	21	13	3	1	-
Male.....	13	27	14	19	19	25	14	14	4	1	-	-
Female.....	11	16	19	17	18	20	17	7	9	2	1	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	4	1	-	8	2	2	1	-	-
Male.....	-	-	-	-	1	-	3	-	-	-	-	-
Female.....	1	-	-	4	-	-	5	2	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

All causes

Nebraska

All races, both sexes.....	525	644	770	1,013	1,238	1,803	2,363	2,479	1,987	864	212	1
Male.....	348	393	466	579	678	949	1,182	1,017	622	169	29	1
Female.....	177	251	304	434	560	854	1,181	1,462	1,365	695	183	-
White, both sexes.....	474	587	701	949	1,166	1,736	2,306	2,430	1,958	851	208	1
Male.....	312	361	426	550	639	924	1,168	993	615	168	28	1
Female.....	162	226	275	399	527	812	1,138	1,437	1,343	683	180	-
Black, both sexes.....	42	47	46	54	55	50	44	38	27	13	4	-
Male.....	29	26	30	23	30	19	10	17	6	1	1	-
Female.....	13	21	16	31	25	31	34	21	21	12	3	-
American Indian 1/, both sexes.....	9	6	17	8	9	10	6	5	1	-	-	-
Male.....	7	4	7	6	6	2	2	2	-	-	-	-
Female.....	2	2	10	2	3	8	4	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	4	6	2	8	7	7	6	1	-	-	-
Male.....	-	2	3	-	3	4	2	5	1	-	-	-
Female.....	-	2	3	2	5	3	5	1	-	-	-	-

Nevada

All races, both sexes.....	1,017	1,280	1,497	1,709	1,950	2,330	2,506	2,086	1,203	345	74	2
Male.....	657	811	923	1,019	1,142	1,296	1,305	885	400	104	11	2
Female.....	360	469	574	690	808	1,034	1,201	1,201	803	241	63	-
White, both sexes.....	854	1,075	1,313	1,494	1,727	2,101	2,330	1,940	1,126	316	65	2
Male.....	563	683	820	897	1,013	1,183	1,214	834	371	94	11	2
Female.....	291	392	493	597	714	918	1,116	1,106	755	222	54	-
Black, both sexes.....	100	121	107	132	134	136	105	84	43	20	7	-
Male.....	53	77	61	76	84	68	58	26	14	6	-	-
Female.....	47	44	46	56	50	68	47	58	29	14	7	-
American Indian 1/, both sexes.....	16	16	9	13	12	18	16	8	7	1	-	-
Male.....	12	10	5	8	7	8	8	3	2	1	-	-
Female.....	4	6	4	5	5	10	8	5	5	-	-	-
Asian or Pacific Islander 2/, both sexes.....	47	68	68	70	77	75	55	54	27	8	2	-
Male.....	29	41	37	38	38	37	25	22	13	3	-	-
Female.....	18	27	31	32	39	38	30	32	14	5	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All causes

New Hampshire

All races, both sexes.....	391	501	663	680	959	1,233	1,581	1,660	1,175	462	103	-
Male.....	242	310	418	400	500	640	732	674	357	104	16	-
Female.....	149	191	245	280	459	593	849	986	818	358	87	-
White, both sexes.....	387	492	655	672	954	1,225	1,577	1,658	1,169	461	102	-
Male.....	240	305	412	396	498	639	729	673	357	103	15	-
Female.....	147	187	243	276	456	586	848	985	812	358	87	-
Black, both sexes.....	2	3	3	5	1	2	2	1	2	-	1	-
Male.....	1	3	3	3	-	-	2	-	-	-	1	-
Female.....	1	-	-	2	1	2	-	1	2	-	-	-
American Indian 1/, both sexes.....	-	-	2	-	-	-	-	1	-	-	-	-
Male.....	-	-	1	-	-	-	-	1	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	6	3	3	4	6	2	-	4	1	-	-
Male.....	1	2	2	1	2	1	1	-	-	1	-	-
Female.....	1	4	1	2	2	5	1	-	4	-	-	-

New Jersey

All races, both sexes.....	2,626	3,348	3,886	4,609	5,736	8,652	11,384	11,566	7,657	3,062	575	4
Male.....	1,618	1,966	2,295	2,587	3,086	4,390	5,283	4,583	2,452	711	98	2
Female.....	1,008	1,382	1,591	2,022	2,650	4,262	6,101	6,983	5,205	2,351	477	2
White, both sexes.....	1,963	2,548	3,081	3,682	4,687	7,529	10,255	10,675	7,148	2,818	519	3
Male.....	1,232	1,520	1,863	2,046	2,543	3,841	4,780	4,256	2,317	663	91	2
Female.....	731	1,028	1,218	1,636	2,144	3,688	5,475	6,419	4,831	2,155	428	1
Black, both sexes.....	588	709	677	799	896	966	953	743	426	218	51	1
Male.....	341	391	363	458	456	480	427	266	110	40	5	-
Female.....	247	318	314	341	440	486	526	477	316	178	46	1
American Indian 1/, both sexes.....	3	4	4	6	9	9	7	5	5	1	-	-
Male.....	1	1	3	2	5	3	4	1	-	-	-	-
Female.....	2	3	1	4	4	6	3	4	5	1	-	-
Asian or Pacific Islander 2/, both sexes.....	72	87	124	122	144	148	169	143	78	25	5	-
Male.....	44	54	66	81	82	66	72	60	25	8	2	-
Female.....	28	33	58	41	62	82	97	83	53	17	3	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>New Mexico</b>												
All races, both sexes.....	768	911	1,028	1,150	1,389	1,817	2,166	2,046	1,307	525	107	2
Male.....	498	585	608	676	785	975	1,052	882	497	139	34	1
Female.....	270	326	420	474	604	842	1,114	1,164	810	386	73	1
White, both sexes.....	681	810	940	1,049	1,246	1,699	2,027	1,938	1,243	489	91	2
Male.....	438	517	553	623	713	915	984	839	468	128	28	1
Female.....	243	293	387	426	533	784	1,043	1,099	775	361	63	1
Black, both sexes.....	15	22	22	13	28	24	40	16	15	4	1	-
Male.....	13	17	15	9	19	21	22	5	6	-	-	-
Female.....	2	5	7	4	9	3	18	11	9	4	1	-
American Indian 1/, both sexes.....	66	66	61	80	106	86	90	81	46	29	15	-
Male.....	46	45	39	41	52	37	43	32	22	11	6	-
Female.....	20	21	22	39	54	49	47	49	24	18	9	-
Asian or Pacific Islander 2/, both sexes.....	6	13	5	8	9	8	9	11	3	3	-	-
Male.....	1	6	1	3	1	2	3	6	1	-	-	-
Female.....	5	7	4	5	8	6	6	5	2	3	-	-
<b>New York</b>												
All races, both sexes.....	5,974	7,679	8,946	10,155	12,662	17,722	22,862	22,938	16,323	6,998	1,561	2
Male.....	3,688	4,625	5,217	5,726	6,745	9,072	10,627	9,176	5,209	1,547	238	1
Female.....	2,286	3,054	3,729	4,429	5,917	8,650	12,235	13,762	11,114	5,451	1,323	1
White, both sexes.....	4,279	5,690	6,932	7,958	10,265	14,943	20,024	20,565	14,900	6,386	1,359	2
Male.....	2,697	3,501	4,076	4,554	5,508	7,753	9,419	8,317	4,806	1,416	206	1
Female.....	1,582	2,189	2,856	3,404	4,757	7,190	10,605	12,248	10,094	4,970	1,153	1
Black, both sexes.....	1,518	1,724	1,732	1,897	2,011	2,285	2,321	1,975	1,143	502	168	-
Male.....	883	966	976	983	997	1,059	953	670	298	91	23	-
Female.....	635	758	756	914	1,014	1,226	1,368	1,305	845	411	145	-
American Indian 1/, both sexes.....	17	16	24	28	25	35	21	14	11	3	2	-
Male.....	9	8	14	11	13	17	7	6	1	1	1	-
Female.....	8	8	10	17	12	18	14	8	10	2	1	-
Asian or Pacific Islander 2/, both sexes.....	160	249	258	272	361	459	496	384	269	107	32	-
Male.....	99	150	151	178	227	243	248	183	104	39	8	-
Female.....	61	99	107	94	134	216	248	201	165	68	24	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
North Carolina												
All races, both sexes.....	3,476	4,431	5,399	6,138	7,388	9,131	10,969	10,289	6,474	2,385	534	6
Male.....	2,126	2,718	3,224	3,498	4,103	4,650	4,972	3,951	1,859	445	70	4
Female.....	1,350	1,713	2,175	2,640	3,285	4,481	5,997	6,338	4,615	1,940	464	2
White, both sexes.....	2,318	3,058	3,940	4,636	5,775	7,396	9,170	8,652	5,514	1,957	427	2
Male.....	1,428	1,884	2,358	2,697	3,237	3,825	4,223	3,399	1,594	374	54	1
Female.....	890	1,174	1,582	1,939	2,538	3,571	4,947	5,253	3,920	1,583	373	1
Black, both sexes.....	1,095	1,275	1,356	1,392	1,523	1,632	1,704	1,560	925	413	104	4
Male.....	662	779	806	744	817	774	710	527	255	69	15	3
Female.....	433	496	550	648	706	858	994	1,033	670	344	89	1
American Indian 1/, both sexes.....	42	67	75	70	65	63	68	50	25	10	2	-
Male.....	25	37	48	38	36	32	30	13	7	1	1	-
Female.....	17	30	27	32	29	31	38	37	18	9	1	-
Asian or Pacific Islander 2/, both sexes.....	21	31	28	40	25	40	27	27	10	5	1	-
Male.....	11	18	12	19	13	19	9	12	3	1	-	-
Female.....	10	13	16	21	12	21	18	15	7	4	1	-
North Dakota												
All races, both sexes.....	185	212	266	335	451	647	810	948	789	354	59	-
Male.....	129	139	159	177	275	350	435	396	282	80	7	-
Female.....	56	73	107	158	176	297	375	552	507	274	52	-
White, both sexes.....	168	192	241	315	432	618	802	937	783	352	59	-
Male.....	115	126	141	169	266	340	430	392	280	79	7	-
Female.....	53	66	100	146	166	278	372	545	503	273	52	-
Black, both sexes.....	1	-	-	2	-	-	-	-	-	-	-	-
Male.....	1	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	16	20	25	18	19	28	7	11	6	1	-	-
Male.....	13	13	18	7	9	10	5	4	2	-	-	-
Female.....	3	7	7	11	10	18	2	7	4	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	1	1	-	-	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Ohio												
All races, both sexes.....	4,622	5,723	6,661	7,779	9,747	13,157	16,963	16,251	10,529	3,734	798	1
Male.....	2,753	3,565	3,977	4,342	5,228	6,695	7,828	6,419	3,240	805	118	1
Female.....	1,869	2,158	2,684	3,437	4,519	6,462	9,135	9,832	7,289	2,929	680	-
White, both sexes.....	3,761	4,714	5,719	6,727	8,526	11,785	15,504	15,125	9,848	3,450	716	1
Male.....	2,267	2,980	3,422	3,771	4,564	6,000	7,149	5,974	3,041	751	107	1
Female.....	1,494	1,734	2,297	2,956	3,962	5,785	8,355	9,151	6,807	2,699	609	-
Black, both sexes.....	839	962	903	1,013	1,177	1,328	1,402	1,088	658	278	76	-
Male.....	473	562	533	547	645	668	652	429	188	50	11	-
Female.....	366	400	370	466	532	660	750	659	470	228	65	-
American Indian 1/, both sexes.....	7	10	5	8	3	6	10	3	6	1	1	-
Male.....	4	7	4	4	2	5	6	-	2	1	-	-
Female.....	3	3	1	4	1	1	4	3	4	-	1	-
Asian or Pacific Islander 2/, both sexes.....	15	37	34	31	41	38	47	35	17	5	5	-
Male.....	9	16	18	20	17	22	21	16	9	3	-	-
Female.....	6	21	16	11	24	16	26	19	8	2	5	-
Oklahoma												
All races, both sexes.....	1,656	2,081	2,473	2,847	3,517	4,502	5,214	4,822	3,083	1,219	256	-
Male.....	1,014	1,238	1,452	1,600	1,882	2,293	2,432	1,959	1,030	254	47	-
Female.....	642	843	1,021	1,247	1,635	2,209	2,782	2,863	2,053	965	209	-
White, both sexes.....	1,336	1,692	2,091	2,447	3,087	3,997	4,753	4,459	2,867	1,124	223	-
Male.....	833	1,016	1,223	1,374	1,664	2,053	2,231	1,825	950	225	42	-
Female.....	503	676	868	1,073	1,423	1,944	2,522	2,634	1,917	899	181	-
Black, both sexes.....	165	183	179	188	195	254	238	183	106	58	21	-
Male.....	91	110	114	100	103	112	98	60	43	17	4	-
Female.....	74	73	65	88	92	142	140	123	63	41	17	-
American Indian 1/, both sexes.....	145	192	186	204	218	235	207	169	105	33	11	-
Male.....	83	107	106	125	108	121	97	69	34	11	1	-
Female.....	62	85	80	79	110	114	110	100	71	22	10	-
Asian or Pacific Islander 2/, both sexes.....	10	14	17	8	17	16	16	11	5	4	1	-
Male.....	7	5	9	1	7	7	6	5	3	1	-	-
Female.....	3	9	8	7	10	9	10	6	2	3	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>Oregon</b>												
All races, both sexes.....	1,375	1,709	1,968	2,168	2,637	3,574	4,831	5,063	3,567	1,349	258	3
Male.....	858	1,016	1,164	1,254	1,452	1,888	2,386	2,115	1,232	356	47	3
Female.....	517	693	804	914	1,185	1,686	2,445	2,948	2,335	993	211	-
White, both sexes.....	1,298	1,603	1,881	2,073	2,531	3,467	4,688	4,947	3,505	1,321	257	3
Male.....	810	954	1,111	1,201	1,392	1,833	2,321	2,064	1,206	347	47	3
Female.....	488	649	770	872	1,139	1,634	2,367	2,883	2,299	974	210	-
Black, both sexes.....	31	49	23	30	37	36	31	36	20	11	-	-
Male.....	20	32	17	21	28	18	11	12	7	3	-	-
Female.....	11	17	6	9	9	18	20	24	13	8	-	-
American Indian 1/, both sexes.....	30	26	30	30	27	22	38	22	7	4	1	-
Male.....	17	11	17	14	11	10	18	9	6	-	-	-
Female.....	13	15	13	16	16	12	20	13	1	4	1	-
Asian or Pacific Islander 2/, both sexes.....	16	31	34	35	42	49	74	58	35	13	-	-
Male.....	11	19	19	18	21	27	36	30	13	6	-	-
Female.....	5	12	15	17	21	22	38	28	22	7	-	-
<b>Pennsylvania</b>												
All races, both sexes.....	4,555	5,896	7,172	8,033	10,456	15,767	20,944	21,301	13,972	5,050	999	16
Male.....	2,827	3,670	4,309	4,643	5,700	8,070	9,739	8,530	4,346	1,145	147	10
Female.....	1,728	2,226	2,863	3,390	4,756	7,697	11,205	12,771	9,626	3,905	852	6
White, both sexes.....	3,658	4,896	6,203	7,030	9,322	14,422	19,581	20,115	13,254	4,772	927	9
Male.....	2,285	3,064	3,767	4,078	5,118	7,449	9,151	8,095	4,160	1,091	138	4
Female.....	1,373	1,832	2,436	2,952	4,204	6,973	10,430	12,020	9,094	3,681	789	5
Black, both sexes.....	856	953	919	936	1,070	1,258	1,274	1,092	677	267	69	6
Male.....	516	578	511	520	548	578	549	393	171	49	9	5
Female.....	340	375	408	416	522	680	725	699	506	218	60	1
American Indian 1/, both sexes.....	2	5	3	4	1	6	5	4	3	-	1	-
Male.....	1	2	2	4	1	3	4	1	1	-	-	-
Female.....	1	3	1	-	-	3	1	3	2	-	1	-
Asian or Pacific Islander 2/, both sexes.....	39	42	47	63	63	81	84	90	38	11	2	1
Male.....	25	26	29	41	33	40	35	41	14	5	-	1
Female.....	14	16	18	22	30	41	49	49	24	6	2	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Rhode Island												
All races, both sexes.....	347	451	525	565	708	1,131	1,603	1,765	1,256	541	83	-
Male.....	201	290	310	315	403	578	747	689	372	113	5	-
Female.....	146	161	215	250	305	553	856	1,076	884	428	78	-
White, both sexes.....	318	414	494	537	672	1,083	1,562	1,734	1,239	535	81	-
Male.....	190	262	289	302	378	555	730	679	364	111	4	-
Female.....	128	152	205	235	294	528	832	1,055	875	424	77	-
Black, both sexes.....	22	31	29	17	27	36	31	23	13	3	-	-
Male.....	8	25	20	7	19	18	10	9	5	1	-	-
Female.....	14	6	9	10	8	18	21	14	8	2	-	-
American Indian 1/, both sexes.....	4	3	2	3	4	4	-	3	2	-	1	-
Male.....	1	1	1	1	1	1	-	1	1	-	1	-
Female.....	3	2	1	2	3	3	-	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	3	-	8	5	8	10	5	2	3	1	-
Male.....	2	2	-	5	5	4	7	-	2	1	-	-
Female.....	1	1	-	3	-	4	3	5	-	2	1	-
South Carolina												
All races, both sexes.....	2,068	2,578	3,158	3,386	3,844	4,646	5,385	4,820	3,042	1,094	229	5
Male.....	1,282	1,605	1,898	2,045	2,107	2,369	2,460	1,815	856	213	27	5
Female.....	786	973	1,260	1,341	1,737	2,277	2,925	3,005	2,186	881	202	-
White, both sexes.....	1,228	1,592	2,142	2,386	2,830	3,513	4,228	3,847	2,460	847	157	5
Male.....	769	989	1,296	1,451	1,600	1,859	1,971	1,486	706	171	18	5
Female.....	459	603	846	935	1,230	1,654	2,257	2,361	1,754	676	139	-
Black, both sexes.....	823	971	1,001	982	1,003	1,120	1,143	964	574	247	72	-
Male.....	500	611	595	586	499	502	482	328	148	42	9	-
Female.....	323	360	406	396	504	618	661	636	426	205	63	-
American Indian 1/, both sexes.....	6	11	5	6	3	3	5	2	5	-	-	-
Male.....	4	4	2	1	3	2	3	-	1	-	-	-
Female.....	2	7	3	5	-	1	2	2	4	-	-	-
Asian or Pacific Islander 2/, both sexes.....	11	4	10	12	8	10	9	7	3	-	-	-
Male.....	9	1	5	7	5	6	4	1	1	-	-	-
Female.....	2	3	5	5	3	4	5	6	2	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>South Dakota</b>												
All races, both sexes.....	231	295	376	446	569	767	1,004	1,111	871	421	106	-
Male.....	147	181	204	251	330	448	544	500	323	108	14	-
Female.....	84	114	172	195	239	319	460	611	548	313	92	-
White, both sexes.....	194	243	323	401	527	727	965	1,080	857	418	104	-
Male.....	130	151	186	233	310	426	529	488	319	108	14	-
Female.....	64	92	137	168	217	301	436	592	538	310	90	-
Black, both sexes.....	1	2	3	1	1	-	1	-	-	-	2	-
Male.....	1	2	1	1	1	-	1	-	-	-	-	-
Female.....	-	-	2	-	-	-	-	-	-	-	2	-
American Indian 1/, both sexes.....	36	45	47	41	41	40	35	30	14	3	-	-
Male.....	16	27	16	15	19	22	14	11	4	-	-	-
Female.....	20	18	31	26	22	18	21	19	10	3	-	-
Asian or Pacific Islander 2/, both sexes.....	-	5	3	3	-	-	3	1	-	-	-	-
Male.....	-	1	1	2	-	-	-	1	-	-	-	-
Female.....	-	4	2	1	-	-	3	-	-	-	-	-
<b>Tennessee</b>												
All races, both sexes.....	2,858	3,554	4,184	4,766	5,409	6,792	7,987	7,514	4,709	1,720	345	5
Male.....	1,777	2,226	2,534	2,723	3,031	3,496	3,667	2,754	1,328	356	53	3
Female.....	1,081	1,328	1,650	2,043	2,378	3,296	4,320	4,760	3,381	1,364	292	2
White, both sexes.....	2,217	2,812	3,505	4,040	4,671	5,993	7,093	6,755	4,231	1,517	281	4
Male.....	1,411	1,779	2,131	2,328	2,660	3,110	3,281	2,493	1,208	312	44	3
Female.....	806	1,033	1,374	1,712	2,011	2,883	3,812	4,262	3,023	1,205	237	1
Black, both sexes.....	627	723	663	698	713	774	867	742	472	198	62	1
Male.....	357	439	394	379	358	370	375	253	119	42	9	-
Female.....	270	284	269	319	355	404	492	489	353	156	53	1
American Indian 1/, both sexes.....	2	5	5	4	3	3	6	1	-	1	1	-
Male.....	2	2	4	3	2	3	4	1	-	-	-	-
Female.....	-	3	1	1	1	-	2	-	-	1	1	-
Asian or Pacific Islander 2/, both sexes.....	12	14	11	24	22	22	21	16	6	4	1	-
Male.....	7	6	5	13	11	13	7	7	1	2	-	-
Female.....	5	8	6	11	11	9	14	9	5	2	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>Texas</b>												
All races, both sexes.....	8,143	9,756	11,237	12,162	14,625	19,001	22,426	21,007	13,295	5,316	1,100	14
Male.....	5,090	6,022	6,676	6,918	8,005	9,732	10,546	8,670	4,144	1,172	183	12
Female.....	3,053	3,734	4,561	5,244	6,620	9,269	11,880	12,337	9,151	4,144	917	2
White, both sexes.....	6,520	7,935	9,417	10,322	12,615	16,784	20,252	19,231	12,169	4,757	933	13
Male.....	4,114	4,934	5,628	5,961	6,961	8,664	9,596	7,990	3,808	1,037	143	12
Female.....	2,406	3,001	3,789	4,361	5,654	8,120	10,656	11,241	8,361	3,720	790	1
Black, both sexes.....	1,508	1,666	1,646	1,659	1,799	1,964	1,915	1,591	1,014	517	159	1
Male.....	914	997	954	865	929	955	830	606	295	121	36	-
Female.....	594	669	692	794	870	1,009	1,085	985	719	396	123	1
American Indian 1/, both sexes.....	23	20	15	29	24	29	28	28	11	3	-	-
Male.....	12	13	12	14	20	12	9	16	3	-	-	-
Female.....	11	7	3	15	4	17	19	12	8	3	-	-
Asian or Pacific Islander 2/, both sexes.....	92	135	159	152	187	224	231	157	101	39	8	-
Male.....	50	78	82	78	95	101	111	58	38	14	4	-
Female.....	42	57	77	74	92	123	120	99	63	25	4	-
<b>Utah</b>												
All races, both sexes.....	611	727	796	911	1,170	1,513	2,093	2,138	1,382	538	62	1
Male.....	374	429	497	534	631	760	978	902	511	156	10	1
Female.....	237	298	299	377	539	753	1,115	1,236	871	382	52	-
White, both sexes.....	582	685	757	872	1,134	1,475	2,040	2,100	1,358	528	61	1
Male.....	352	400	472	513	613	737	953	885	503	153	10	1
Female.....	230	285	285	359	521	738	1,087	1,215	855	375	51	-
Black, both sexes.....	7	8	9	8	3	5	11	7	3	4	-	-
Male.....	7	5	7	4	2	4	4	3	1	-	-	-
Female.....	-	3	2	4	1	1	7	4	2	4	-	-
American Indian 1/, both sexes.....	14	15	12	15	11	11	10	8	5	3	-	-
Male.....	12	10	7	9	4	6	5	2	1	2	-	-
Female.....	2	5	5	6	7	5	5	6	4	1	-	-
Asian or Pacific Islander 2/, both sexes.....	8	19	18	16	22	22	32	23	16	3	1	-
Male.....	3	14	11	8	12	13	16	12	6	1	-	-
Female.....	5	5	7	8	10	9	16	11	10	2	1	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
All causes												
Vermont												
All races, both sexes.....	224	232	316	385	452	626	815	843	565	259	46	-
Male.....	124	135	200	218	255	307	395	347	178	57	8	-
Female.....	100	97	116	167	197	319	420	496	387	202	38	-
White, both sexes.....	221	229	315	382	449	620	810	841	565	259	46	-
Male.....	123	133	200	217	253	304	393	345	178	57	8	-
Female.....	98	96	115	165	196	316	417	496	387	202	38	-
Black, both sexes.....	3	1	-	-	-	2	1	1	-	-	-	-
Male.....	1	-	-	-	-	1	-	1	-	-	-	-
Female.....	2	1	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	1	-	3	2	-	-	-	-	-
Male.....	-	1	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	1	-	2	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	2	3	1	2	1	-	-	-	-
Male.....	-	1	-	1	2	1	1	1	-	-	-	-
Female.....	-	-	1	1	1	-	1	-	-	-	-	-
Virginia												
All races, both sexes.....	2,610	3,300	3,948	4,470	5,239	7,231	8,645	8,327	5,297	1,963	439	2
Male.....	1,593	2,001	2,347	2,520	2,914	3,735	3,953	3,321	1,617	428	58	2
Female.....	1,017	1,299	1,601	1,950	2,325	3,496	4,692	5,006	3,680	1,535	381	-
White, both sexes.....	1,771	2,294	2,906	3,341	4,018	5,737	7,145	7,095	4,502	1,672	340	-
Male.....	1,085	1,389	1,744	1,923	2,241	2,998	3,302	2,826	1,383	356	41	-
Female.....	686	905	1,162	1,418	1,777	2,739	3,843	4,269	3,119	1,316	299	-
Black, both sexes.....	786	946	971	1,040	1,095	1,383	1,387	1,148	734	277	97	2
Male.....	481	575	563	550	602	683	602	457	208	67	15	2
Female.....	305	371	408	490	493	700	785	691	526	210	82	-
American Indian 1/, both sexes.....	4	2	3	1	5	8	5	2	3	2	-	-
Male.....	1	2	1	1	3	4	1	1	1	1	-	-
Female.....	3	-	2	-	2	4	4	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes.....	49	58	68	88	121	103	108	82	58	12	2	-
Male.....	26	35	39	46	68	50	48	37	25	4	2	-
Female.....	23	23	29	42	53	53	60	45	33	8	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>All causes</b>												
<b>Washington</b>												
All races, both sexes.....	2,036	2,599	3,005	3,290	4,176	5,418	6,977	7,560	5,179	1,987	408	1
Male.....	1,236	1,546	1,727	1,921	2,311	2,809	3,246	3,239	1,804	543	79	1
Female.....	800	1,053	1,278	1,369	1,865	2,609	3,731	4,321	3,375	1,444	329	-
White, both sexes.....	1,792	2,317	2,726	3,001	3,851	5,063	6,576	7,225	5,003	1,932	392	1
Male.....	1,102	1,381	1,569	1,755	2,139	2,635	3,074	3,101	1,749	525	75	1
Female.....	690	936	1,157	1,246	1,712	2,428	3,502	4,124	3,254	1,407	317	-
Black, both sexes.....	113	120	103	116	107	133	105	92	53	14	10	-
Male.....	62	73	65	70	69	69	48	36	18	2	2	-
Female.....	51	47	38	46	38	64	57	56	35	12	8	-
American Indian 1/, both sexes.....	53	55	67	62	78	54	55	48	16	3	-	-
Male.....	28	33	39	27	35	27	26	20	7	-	-	-
Female.....	25	22	28	35	43	27	29	28	9	3	-	-
Asian or Pacific Islander 2/, both sexes.....	78	107	109	111	140	168	241	195	107	38	6	-
Male.....	44	59	54	69	68	78	98	82	30	16	2	-
Female.....	34	48	55	42	72	90	143	113	77	22	4	-
<b>West Virginia</b>												
All races, both sexes.....	941	1,205	1,468	1,689	2,098	2,698	3,304	2,842	1,779	663	123	-
Male.....	617	767	896	967	1,143	1,351	1,595	1,140	528	130	15	-
Female.....	324	438	572	722	955	1,347	1,709	1,702	1,251	533	108	-
White, both sexes.....	896	1,139	1,407	1,635	2,034	2,618	3,234	2,763	1,736	635	115	-
Male.....	579	723	854	938	1,116	1,310	1,565	1,118	517	127	14	-
Female.....	317	416	553	697	918	1,308	1,669	1,645	1,219	508	101	-
Black, both sexes.....	44	65	58	49	59	76	66	76	41	28	8	-
Male.....	38	43	40	26	26	38	28	21	10	3	1	-
Female.....	6	22	18	23	33	38	38	55	31	25	7	-
American Indian 1/, both sexes.....	1	1	1	1	3	1	1	-	-	-	-	-
Male.....	-	1	-	-	-	1	1	-	-	-	-	-
Female.....	1	-	1	1	3	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	4	2	3	3	3	2	-	-	-
Male.....	-	-	2	3	1	2	1	1	1	-	-	-
Female.....	-	-	-	1	1	1	2	2	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

All causes

Wisconsin

All races, both sexes.....	1,669	2,170	2,606	2,969	3,869	5,371	7,140	7,630	5,658	2,257	460	-
Male.....	1,060	1,346	1,558	1,745	2,219	2,859	3,428	3,094	1,796	474	64	-
Female.....	609	824	1,048	1,224	1,650	2,512	3,712	4,536	3,862	1,783	396	-
White, both sexes.....	1,430	1,920	2,396	2,743	3,629	5,126	6,924	7,463	5,556	2,207	449	-
Male.....	928	1,203	1,436	1,620	2,101	2,740	3,346	3,032	1,763	466	64	-
Female.....	502	717	960	1,123	1,528	2,386	3,578	4,431	3,793	1,741	385	-
Black, both sexes.....	198	197	163	169	188	186	154	121	74	38	9	-
Male.....	103	110	96	97	96	90	56	42	20	5	-	-
Female.....	95	87	67	72	92	96	98	79	54	33	9	-
American Indian 1/, both sexes.....	28	39	31	37	27	40	33	20	11	8	1	-
Male.....	21	23	16	18	14	18	13	11	6	2	-	-
Female.....	7	16	15	19	13	22	20	9	5	6	1	-
Asian or Pacific Islander 2/, both sexes.....	13	14	16	20	25	19	29	26	17	4	1	-
Male.....	8	10	10	10	8	11	13	9	7	1	-	-
Female.....	5	4	6	10	17	8	16	17	10	3	1	-

Wyoming

All races, both sexes.....	188	228	273	322	387	503	611	582	382	154	31	-
Male.....	111	146	160	195	214	273	299	236	121	46	8	-
Female.....	77	82	113	127	173	230	312	346	261	108	23	-
White, both sexes.....	184	219	258	315	379	494	599	578	379	151	31	-
Male.....	108	139	151	191	208	269	295	234	120	45	8	-
Female.....	76	80	107	124	171	225	304	344	259	106	23	-
Black, both sexes.....	-	-	1	1	3	3	1	1	2	2	-	-
Male.....	-	-	-	1	2	3	-	1	1	1	-	-
Female.....	-	-	1	-	1	-	1	-	1	1	-	-
American Indian 1/, both sexes.....	4	9	12	6	3	5	7	3	1	-	-	-
Male.....	3	7	8	3	2	1	2	1	-	-	-	-
Female.....	1	2	4	3	1	4	5	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	-	2	1	4	-	-	1	-	-
Male.....	-	-	1	-	2	-	2	-	-	-	-	-
Female.....	-	-	1	-	-	1	2	-	-	1	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>I. Certain infectious and parasitic diseases (A00-B99)</b>												
<b>Arkansas</b>												
All races, both sexes.....	719	6	4	1	2	2	1	9	12	20	31	48
Male.....	350	2	3	1	2	-	-	7	5	14	20	27
Female.....	369	4	1	-	-	2	1	2	7	6	11	21
White, both sexes.....	548	3	3	1	1	1	1	3	4	12	18	28
Male.....	256	1	2	1	1	-	-	2	2	10	11	16
Female.....	292	2	1	-	-	1	1	1	2	2	7	12
Black, both sexes.....	169	3	1	-	1	1	-	6	7	8	13	20
Male.....	93	1	1	-	1	-	-	5	3	4	9	11
Female.....	76	2	-	-	-	1	-	1	4	4	4	9
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	1	-	-	-
<b>California</b>												
All races, both sexes.....	4,856	41	21	8	8	14	32	67	78	158	340	448
Male.....	2,939	24	10	5	5	10	20	46	52	123	276	319
Female.....	1,917	17	11	3	3	4	12	21	26	35	64	129
White, both sexes.....	3,849	35	17	6	8	11	22	47	58	118	251	333
Male.....	2,310	22	8	4	5	7	14	33	39	97	214	237
Female.....	1,539	13	9	2	3	4	8	14	19	21	37	96
Black, both sexes.....	587	5	1	2	-	2	5	13	17	26	72	80
Male.....	382	1	-	1	-	2	3	8	11	19	51	56
Female.....	205	4	1	1	-	-	2	5	6	7	21	24
American Indian 1/, both sexes.....	56	-	-	-	-	-	-	1	1	6	6	6
Male.....	34	-	-	-	-	-	-	1	1	2	5	4
Female.....	22	-	-	-	-	-	-	-	-	4	1	2
Asian or Pacific Islander 2/, both sexes.....	364	1	3	-	-	1	5	6	2	8	11	29
Male.....	213	1	2	-	-	1	3	4	1	5	6	22
Female.....	151	-	1	-	-	-	2	2	1	3	5	7











## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>I. Certain infectious and parasitic diseases (A00-B99)</b>												
<b>Illinois</b>												
All races, both sexes.....	3,200	19	9	3	3	12	10	22	29	59	103	142
Male.....	1,529	10	4	2	-	6	8	12	19	42	66	96
Female.....	1,671	9	5	1	3	6	2	10	10	17	37	46
White, both sexes.....	2,341	11	7	3	3	7	5	8	9	17	52	77
Male.....	1,078	5	3	2	-	3	5	6	7	9	33	51
Female.....	1,263	6	4	1	3	4	-	2	2	8	19	26
Black, both sexes.....	809	8	1	-	-	5	5	14	20	42	51	62
Male.....	428	5	-	-	-	3	3	6	12	33	33	43
Female.....	381	3	1	-	-	2	2	8	8	9	18	19
American Indian 1/, both sexes.....	4	-	1	-	-	-	-	-	-	-	-	1
Male.....	2	-	1	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	46	-	-	-	-	-	-	-	-	-	-	2
Male.....	21	-	-	-	-	-	-	-	-	-	-	2
Female.....	25	-	-	-	-	-	-	-	-	-	-	-
<b>Indiana</b>												
All races, both sexes.....	1,318	12	5	2	1	2	6	13	12	24	51	67
Male.....	591	9	2	2	1	-	5	8	6	16	31	38
Female.....	727	3	3	-	-	2	1	5	6	8	20	29
White, both sexes.....	1,146	8	3	1	1	2	4	11	10	18	39	54
Male.....	514	6	-	1	1	-	4	7	6	12	25	33
Female.....	632	2	3	-	-	2	-	4	4	6	14	21
Black, both sexes.....	168	4	2	1	-	-	2	2	2	5	12	12
Male.....	76	3	2	1	-	-	1	1	-	4	6	5
Female.....	92	1	-	-	-	-	1	1	2	1	6	7
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	4	-	-	-	-	-	-	-	-	1	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	3	-	-	-	-	-	-	-	-	1	-	1





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>I. Certain infectious and parasitic diseases (A00-B99)</b>												
<b>Maine</b>												
All races, both sexes.....	285	2	-	-	1	-	2	-	-	-	5	8
Male.....	147	1	-	-	1	-	1	-	-	-	4	5
Female.....	138	1	-	-	-	-	1	-	-	-	1	3
White, both sexes.....	280	2	-	-	1	-	2	-	-	-	5	8
Male.....	144	1	-	-	1	-	1	-	-	-	4	5
Female.....	136	1	-	-	-	-	1	-	-	-	1	3
Black, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
<b>Maryland</b>												
All races, both sexes.....	1,820	11	9	4	3	3	9	19	31	65	107	143
Male.....	928	7	8	4	1	3	3	7	13	28	55	92
Female.....	892	4	1	-	2	-	6	12	18	37	52	51
White, both sexes.....	969	6	3	3	1	1	2	8	2	14	24	29
Male.....	461	4	3	3	1	1	-	2	-	7	12	21
Female.....	508	2	-	-	-	-	2	6	2	7	12	8
Black, both sexes.....	822	4	3	1	2	2	6	10	29	51	83	113
Male.....	448	3	3	1	-	2	2	5	13	21	43	70
Female.....	374	1	-	-	2	-	4	5	16	30	40	43
American Indian 1/, both sexes.....	3	-	1	-	-	-	-	-	-	-	-	-
Male.....	3	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	26	1	2	-	-	-	1	1	-	-	-	1
Male.....	16	-	1	-	-	-	1	-	-	-	-	1
Female.....	10	1	1	-	-	-	-	1	-	-	-	-



## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>I. Certain infectious and parasitic diseases (A00-B99)</b>												
<b>Massachusetts</b>												
All races, both sexes.....	1,496	2	3	2	-	3	2	3	7	24	44	75
Male.....	746	2	1	2	-	2	1	2	3	14	32	49
Female.....	750	-	2	-	-	1	1	1	4	10	12	26
White, both sexes.....	1,359	1	3	1	-	3	1	3	4	14	33	60
Male.....	677	1	1	1	-	2	-	2	2	11	25	45
Female.....	682	-	2	-	-	1	1	1	2	3	8	15
Black, both sexes.....	112	1	-	-	-	-	1	-	2	10	8	13
Male.....	51	1	-	-	-	-	1	-	-	3	5	2
Female.....	61	-	-	-	-	-	-	-	2	7	3	11
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	1	-
Male.....	1	-	-	-	-	-	-	-	-	-	1	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	24	-	-	1	-	-	-	-	1	-	2	2
Male.....	17	-	-	1	-	-	-	-	1	-	1	2
Female.....	7	-	-	-	-	-	-	-	-	-	1	-
<b>Michigan</b>												
All races, both sexes.....	1,769	12	3	-	1	4	4	8	25	32	70	95
Male.....	858	4	1	-	-	4	3	3	14	22	45	64
Female.....	911	8	2	-	1	-	1	5	11	10	25	31
White, both sexes.....	1,293	4	2	-	1	3	1	2	6	16	33	62
Male.....	611	2	1	-	-	3	1	1	3	11	24	43
Female.....	682	2	1	-	1	-	-	1	3	5	9	19
Black, both sexes.....	454	8	-	-	-	1	3	6	17	16	37	30
Male.....	235	2	-	-	-	1	2	2	10	11	21	19
Female.....	219	6	-	-	-	-	1	4	7	5	16	11
American Indian 1/, both sexes.....	13	-	-	-	-	-	-	-	1	-	-	1
Male.....	6	-	-	-	-	-	-	-	1	-	-	1
Female.....	7	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	9	-	1	-	-	-	-	-	1	-	-	2
Male.....	6	-	-	-	-	-	-	-	-	-	-	1
Female.....	3	-	1	-	-	-	-	-	1	-	-	1









## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>I. Certain infectious and parasitic diseases (A00-B99)</b>												
<b>New Mexico</b>												
All races, both sexes.....	354	5	1	-	-	1	-	2	9	12	14	24
Male.....	200	3	1	-	-	1	-	1	5	10	12	17
Female.....	154	2	-	-	-	-	-	1	4	2	2	7
White, both sexes.....	314	4	1	-	-	1	-	1	8	8	12	20
Male.....	176	2	1	-	-	1	-	1	4	6	10	15
Female.....	138	2	-	-	-	-	-	-	4	2	2	5
Black, both sexes.....	7	1	-	-	-	-	-	-	-	-	1	-
Male.....	6	1	-	-	-	-	-	-	-	-	1	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	31	-	-	-	-	-	-	1	1	4	1	4
Male.....	16	-	-	-	-	-	-	-	1	4	1	2
Female.....	15	-	-	-	-	-	-	1	-	-	-	2
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
<b>New York</b>												
All races, both sexes.....	4,958	21	11	7	2	13	22	38	61	173	301	416
Male.....	2,636	12	6	4	1	4	14	16	35	107	182	250
Female.....	2,322	9	5	3	1	9	8	22	26	66	119	166
White, both sexes.....	3,557	13	7	6	1	7	15	18	29	73	155	215
Male.....	1,835	6	4	3	-	2	11	8	17	56	98	139
Female.....	1,722	7	3	3	1	5	4	10	12	17	57	76
Black, both sexes.....	1,318	8	4	1	1	6	7	20	29	98	140	195
Male.....	747	6	2	1	1	2	3	8	17	49	80	106
Female.....	571	2	2	-	-	4	4	12	12	49	60	89
American Indian 1/, both sexes.....	10	-	-	-	-	-	-	-	-	-	-	2
Male.....	4	-	-	-	-	-	-	-	-	-	-	1
Female.....	6	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	73	-	-	-	-	-	-	-	3	2	6	4
Male.....	50	-	-	-	-	-	-	-	1	2	4	4
Female.....	23	-	-	-	-	-	-	-	2	-	2	-















## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>I. Certain infectious and parasitic diseases (A00-B99)</b>												
<b>Vermont</b>												
All races, both sexes.....	80	-	-	-	-	-	-	-	-	-	1	3
Male.....	43	-	-	-	-	-	-	-	-	-	-	2
Female.....	37	-	-	-	-	-	-	-	-	-	1	1
White, both sexes.....	77	-	-	-	-	-	-	-	-	-	1	2
Male.....	41	-	-	-	-	-	-	-	-	-	-	1
Female.....	36	-	-	-	-	-	-	-	-	-	1	1
Black, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	1
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
<b>Virginia</b>												
All races, both sexes.....	2,039	17	6	-	1	-	6	21	24	49	75	94
Male.....	1,033	7	5	-	1	-	-	15	18	32	51	61
Female.....	1,006	10	1	-	-	-	6	6	6	17	24	33
White, both sexes.....	1,414	9	6	-	1	-	2	11	6	22	26	48
Male.....	688	5	5	-	1	-	-	8	6	13	19	33
Female.....	726	4	1	-	-	-	2	3	-	9	7	15
Black, both sexes.....	590	6	-	-	-	-	4	9	15	27	49	46
Male.....	325	1	-	-	-	-	-	6	10	19	32	28
Female.....	265	5	-	-	-	-	4	3	5	8	17	18
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	34	2	-	-	-	-	-	1	3	-	-	-
Male.....	20	1	-	-	-	-	-	1	2	-	-	-
Female.....	14	1	-	-	-	-	-	-	1	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

United States

All races, both sexes.....	5,550	5,294	4,415	4,482	5,310	6,790	8,372	7,735	4,393	1,499	248	5
Male.....	3,723	3,458	2,596	2,422	2,726	3,210	3,672	2,987	1,450	353	42	4
Female.....	1,827	1,836	1,819	2,060	2,584	3,580	4,700	4,748	2,943	1,146	206	1
White, both sexes.....	3,547	3,610	3,178	3,400	4,159	5,652	7,221	6,842	3,892	1,312	203	1
Male.....	2,432	2,377	1,821	1,822	2,116	2,700	3,219	2,661	1,282	301	34	1
Female.....	1,115	1,233	1,357	1,578	2,043	2,952	4,002	4,181	2,610	1,011	169	-
Black, both sexes.....	1,870	1,520	1,104	945	949	954	973	756	437	173	40	3
Male.....	1,219	976	701	526	504	425	370	273	135	46	7	3
Female.....	651	544	403	419	445	529	603	483	302	127	33	-
American Indian 1/, both sexes.....	61	64	40	34	47	36	33	19	8	1	2	-
Male.....	30	36	18	19	22	15	16	4	4	-	-	-
Female.....	31	28	22	15	25	21	17	15	4	1	2	-
Asian or Pacific Islander 2/, both sexes.....	72	100	93	103	155	148	145	118	56	13	3	1
Male.....	42	69	56	55	84	70	67	49	29	6	1	-
Female.....	30	31	37	48	71	78	78	69	27	7	2	1

Alabama

All races, both sexes.....	119	94	91	96	117	128	144	119	70	25	7	-
Male.....	81	63	49	49	52	49	61	38	26	6	1	-
Female.....	38	31	42	47	65	79	83	81	44	19	6	-
White, both sexes.....	74	53	67	64	78	103	121	100	52	17	1	-
Male.....	49	32	34	31	30	37	56	34	17	3	-	-
Female.....	25	21	33	33	48	66	65	66	35	14	1	-
Black, both sexes.....	45	41	24	32	39	25	23	18	18	8	6	-
Male.....	32	31	15	18	22	12	5	4	9	3	1	-
Female.....	13	10	9	14	17	13	18	14	9	5	5	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Alaska

All races, both sexes.....	7	15	4	8	3	6	4	5	5	1	-	-
Male.....	5	8	1	3	1	4	2	2	2	-	-	-
Female.....	2	7	3	5	2	2	2	3	3	1	-	-
White, both sexes.....	5	8	3	6	1	2	1	3	4	1	-	-
Male.....	4	5	1	3	1	2	1	2	1	-	-	-
Female.....	1	3	2	3	-	-	-	1	3	1	-	-
Black, both sexes.....	1	1	-	-	-	1	-	-	-	-	-	-
Male.....	1	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	5	1	1	2	3	3	2	-	-	-	-
Male.....	-	2	-	-	-	2	1	-	-	-	-	-
Female.....	1	3	1	1	2	1	2	2	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	1	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	1	-	1	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	96	96	90	85	119	158	143	119	63	12	3	1
Male.....	69	65	52	48	62	74	63	52	22	6	-	1
Female.....	27	31	38	37	57	84	80	67	41	6	3	-
White, both sexes.....	79	83	77	76	101	144	132	112	60	12	3	1
Male.....	60	58	45	40	53	69	59	52	21	6	-	1
Female.....	19	25	32	36	48	75	73	60	39	6	3	-
Black, both sexes.....	7	5	7	2	6	5	2	4	1	-	-	-
Male.....	5	3	4	2	4	2	-	-	1	-	-	-
Female.....	2	2	3	-	2	3	2	4	-	-	-	-
American Indian 1/, both sexes.....	9	6	5	5	7	7	3	3	1	-	-	-
Male.....	4	4	2	5	3	2	1	-	-	-	-	-
Female.....	5	2	3	-	4	5	2	3	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	2	1	2	5	2	6	-	1	-	-	-
Male.....	-	-	1	1	2	1	3	-	-	-	-	-
Female.....	1	2	-	1	3	1	3	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Arkansas

All races, both sexes.....	42	58	55	43	63	72	99	83	46	18	3	1
Male.....	30	32	26	20	32	33	50	25	15	5	-	1
Female.....	12	26	29	23	31	39	49	58	31	13	3	-
White, both sexes.....	28	42	44	31	54	63	79	72	41	16	3	-
Male.....	20	21	20	13	27	28	41	23	13	4	-	-
Female.....	8	21	24	18	27	35	38	49	28	12	3	-
Black, both sexes.....	14	15	11	12	9	9	20	11	5	2	-	1
Male.....	10	10	6	7	5	5	9	2	2	1	-	1
Female.....	4	5	5	5	4	4	11	9	3	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	559	582	384	314	328	334	432	410	213	72	13	-
Male.....	387	411	251	194	180	168	198	157	80	20	3	-
Female.....	172	171	133	120	148	166	234	253	133	52	10	-
White, both sexes.....	434	449	313	254	244	260	366	358	190	63	12	-
Male.....	314	321	200	158	119	130	163	138	67	18	2	-
Female.....	120	128	113	96	125	130	203	220	123	45	10	-
Black, both sexes.....	87	92	42	35	33	26	22	15	7	5	-	-
Male.....	50	62	31	24	29	13	13	4	3	1	-	-
Female.....	37	30	11	11	4	13	9	11	4	4	-	-
American Indian 1/, both sexes.....	11	8	4	3	5	3	1	1	-	-	-	-
Male.....	6	4	2	1	5	2	1	-	-	-	-	-
Female.....	5	4	2	2	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	27	33	25	22	46	45	43	36	16	4	1	-
Male.....	17	24	18	11	27	23	21	15	10	1	1	-
Female.....	10	9	7	11	19	22	22	21	6	3	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Colorado

All races, both sexes.....	65	64	45	40	54	65	74	63	43	11	1	-
Male.....	51	39	22	20	29	35	32	28	16	3	1	-
Female.....	14	25	23	20	25	30	42	35	27	8	-	-
White, both sexes.....	59	57	38	36	50	61	73	60	42	11	1	-
Male.....	46	34	19	19	25	35	31	25	15	3	1	-
Female.....	13	23	19	17	25	26	42	35	27	8	-	-
Black, both sexes.....	2	5	5	2	3	3	1	1	1	-	-	-
Male.....	2	4	2	-	3	-	1	1	1	-	-	-
Female.....	-	1	3	2	-	3	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	2	1	-	1	-	-	-	-	-	-	-
Male.....	-	1	1	-	1	-	-	-	-	-	-	-
Female.....	1	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	1	2	-	1	-	2	-	-	-	-
Male.....	3	-	-	1	-	-	-	2	-	-	-	-
Female.....	-	-	1	1	-	1	-	-	-	-	-	-

Connecticut

All races, both sexes.....	85	57	53	63	87	93	179	183	92	39	6	-
Male.....	59	43	31	29	47	52	81	76	29	10	2	-
Female.....	26	14	22	34	40	41	98	107	63	29	4	-
White, both sexes.....	57	38	44	49	73	88	167	175	89	38	6	-
Male.....	39	27	26	22	41	51	75	73	27	10	2	-
Female.....	18	11	18	27	32	37	92	102	62	28	4	-
Black, both sexes.....	26	18	9	14	12	4	12	6	2	1	-	-
Male.....	18	15	5	7	4	1	6	2	1	-	-	-
Female.....	8	3	4	7	8	3	6	4	1	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	1	-	-	1	1	-	2	1	-	-	-
Male.....	2	1	-	-	1	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Florida

All races, both sexes.....	511	413	274	266	331	472	530	442	243	74	15	1
Male.....	362	289	171	155	184	242	255	185	84	23	1	1
Female.....	149	124	103	111	147	230	275	257	159	51	14	-
White, both sexes.....	312	290	183	204	266	409	478	408	230	67	14	-
Male.....	228	208	112	117	147	209	229	176	78	21	1	-
Female.....	84	82	71	87	119	200	249	232	152	46	13	-
Black, both sexes.....	198	122	91	59	60	56	50	32	12	7	1	1
Male.....	134	80	59	35	36	31	25	9	6	2	-	1
Female.....	64	42	32	24	24	25	25	23	6	5	1	-
American Indian 1/, both sexes.....	-	1	-	-	-	1	-	-	-	-	-	-
Male.....	-	1	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	3	5	6	2	2	1	-	-	-
Male.....	-	-	-	3	1	1	1	-	-	-	-	-
Female.....	1	-	-	-	4	5	1	2	1	-	-	-

Georgia

All races, both sexes.....	252	219	190	174	191	220	232	209	115	49	7	-
Male.....	164	138	117	96	96	109	97	63	30	9	-	-
Female.....	88	81	73	78	95	111	135	146	85	40	7	-
White, both sexes.....	107	97	91	110	111	151	173	155	83	32	5	-
Male.....	74	58	55	62	53	74	74	44	25	5	-	-
Female.....	33	39	36	48	58	77	99	111	58	27	5	-
Black, both sexes.....	144	116	94	61	77	66	59	52	32	17	2	-
Male.....	90	74	58	33	41	33	23	18	5	4	-	-
Female.....	54	42	36	28	36	33	36	34	27	13	2	-
American Indian 1/, both sexes.....	-	2	1	-	-	-	-	-	-	-	-	-
Male.....	-	2	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	4	4	3	3	3	-	2	-	-	-	-
Male.....	-	4	3	1	2	2	-	1	-	-	-	-
Female.....	1	-	1	2	1	1	-	1	-	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Maine

All races, both sexes.....	20	15	18	23	19	35	41	53	31	12	-	-
Male.....	14	11	9	17	10	18	21	20	13	2	-	-
Female.....	6	4	9	6	9	17	20	33	18	10	-	-
White, both sexes.....	17	15	18	23	18	35	41	53	30	12	-	-
Male.....	13	11	9	17	9	18	21	20	12	2	-	-
Female.....	4	4	9	6	9	17	20	33	18	10	-	-
Black, both sexes.....	1	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	160	124	104	135	120	190	193	228	124	34	4	-
Male.....	103	81	64	79	58	94	91	87	41	9	-	-
Female.....	57	43	40	56	62	96	102	141	83	25	4	-
White, both sexes.....	37	46	58	69	72	133	147	185	98	28	3	-
Male.....	20	28	32	37	33	68	71	75	34	9	-	-
Female.....	17	18	26	32	39	65	76	110	64	19	3	-
Black, both sexes.....	121	74	43	64	45	53	44	42	25	6	1	-
Male.....	82	51	29	40	22	24	18	12	7	-	-	-
Female.....	39	23	14	24	23	29	26	30	18	6	1	-
American Indian 1/, both sexes.....	-	-	-	1	1	-	-	-	-	-	-	-
Male.....	-	-	-	1	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	4	3	1	2	4	2	1	1	-	-	-
Male.....	1	2	3	1	2	2	2	-	-	-	-	-
Female.....	1	2	-	-	-	2	-	1	1	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Minnesota

All races, both sexes.....	46	45	46	42	64	61	85	88	57	14	3	-
Male.....	30	30	33	20	38	31	42	26	20	3	2	-
Female.....	16	15	13	22	26	30	43	62	37	11	1	-
White, both sexes.....	36	34	38	38	56	55	83	87	54	14	3	-
Male.....	24	22	26	17	35	29	41	26	19	3	2	-
Female.....	12	12	12	21	21	26	42	61	35	11	1	-
Black, both sexes.....	7	3	5	2	4	4	2	-	1	-	-	-
Male.....	5	2	4	1	2	1	1	-	-	-	-	-
Female.....	2	1	1	1	2	3	1	-	1	-	-	-
American Indian 1/, both sexes.....	3	5	1	1	1	2	-	-	-	-	-	-
Male.....	1	3	1	1	-	1	-	-	-	-	-	-
Female.....	2	2	-	-	1	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	3	2	1	3	-	-	1	2	-	-	-
Male.....	-	3	2	1	1	-	-	-	1	-	-	-
Female.....	-	-	-	-	2	-	-	1	1	-	-	-

Mississippi

All races, both sexes.....	62	63	62	59	72	80	90	81	52	17	3	-
Male.....	39	31	30	33	35	38	39	29	17	3	-	-
Female.....	23	32	32	26	37	42	51	52	35	14	3	-
White, both sexes.....	27	30	42	35	44	54	59	61	32	11	1	-
Male.....	18	14	20	21	20	28	29	20	8	-	-	-
Female.....	9	16	22	14	24	26	30	41	24	11	1	-
Black, both sexes.....	35	33	20	23	28	26	30	20	19	6	2	-
Male.....	21	17	10	12	15	10	10	9	8	3	-	-
Female.....	14	16	10	11	13	16	20	11	11	3	2	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Nebraska

All races, both sexes.....	18	16	19	21	19	31	35	39	27	10	-	-
Male.....	16	9	12	11	4	11	19	19	8	3	-	-
Female.....	2	7	7	10	15	20	16	20	19	7	-	-
White, both sexes.....	16	13	16	21	14	28	33	37	26	10	-	-
Male.....	14	7	10	11	3	10	18	18	8	3	-	-
Female.....	2	6	6	10	11	18	15	19	18	7	-	-
Black, both sexes.....	2	3	2	-	5	2	2	-	1	-	-	-
Male.....	2	2	1	-	1	1	1	-	-	-	-	-
Female.....	-	1	1	-	4	1	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	-	1	-	2	-	-	-	-
Male.....	-	-	1	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	47	47	48	57	78	77	83	43	32	4	2	-
Male.....	31	30	27	28	41	39	37	24	19	1	-	-
Female.....	16	17	21	29	37	38	46	19	13	3	2	-
White, both sexes.....	39	35	39	44	67	63	77	35	26	4	1	-
Male.....	25	21	22	22	37	34	34	21	15	1	-	-
Female.....	14	14	17	22	30	29	43	14	11	3	1	-
Black, both sexes.....	7	7	4	7	3	9	3	3	3	-	1	-
Male.....	5	5	4	3	2	2	3	1	2	-	-	-
Female.....	2	2	-	4	1	7	-	2	1	-	1	-
American Indian 1/, both sexes.....	1	2	1	2	-	-	-	-	1	-	-	-
Male.....	1	2	-	1	-	-	-	-	1	-	-	-
Female.....	-	-	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	3	4	4	8	5	3	5	2	-	-	-
Male.....	-	2	1	2	2	3	-	2	1	-	-	-
Female.....	-	1	3	2	6	2	3	3	1	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
New Hampshire												
All races, both sexes.....	11	22	18	16	18	25	35	41	27	9	2	-
Male.....	6	17	12	12	10	11	18	14	10	2	1	-
Female.....	5	5	6	4	8	14	17	27	17	7	1	-
White, both sexes.....	11	22	17	16	18	25	35	41	27	9	1	-
Male.....	6	17	11	12	10	11	18	14	10	2	-	-
Female.....	5	5	6	4	8	14	17	27	17	7	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	1	-
Male.....	-	-	-	-	-	-	-	-	-	-	1	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes.....	200	182	162	166	219	312	417	412	225	90	9	-
Male.....	119	119	104	87	128	158	171	141	61	17	2	-
Female.....	81	63	58	79	91	154	246	271	164	73	7	-
White, both sexes.....	98	102	110	115	161	253	361	365	207	80	9	-
Male.....	58	68	71	56	94	132	146	126	57	14	2	-
Female.....	40	34	39	59	67	121	215	239	150	66	7	-
Black, both sexes.....	100	78	45	48	49	51	49	38	18	9	-	-
Male.....	60	49	30	30	29	24	23	13	4	2	-	-
Female.....	40	29	15	18	20	27	26	25	14	7	-	-
American Indian 1/, both sexes.....	1	-	1	-	1	-	-	1	-	-	-	-
Male.....	1	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	2	6	3	8	8	7	8	-	1	-	-
Male.....	-	2	2	1	5	2	2	2	-	1	-	-
Female.....	1	-	4	2	3	6	5	6	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

New Mexico

All races, both sexes.....	38	37	28	20	30	33	46	30	19	4	1	-
Male.....	24	25	18	11	16	20	16	11	8	1	-	-
Female.....	14	12	10	9	14	13	30	19	11	3	1	-
White, both sexes.....	33	36	26	18	25	26	43	29	19	4	-	-
Male.....	21	24	17	10	14	17	14	10	8	1	-	-
Female.....	12	12	9	8	11	9	29	19	11	3	-	-
Black, both sexes.....	1	-	-	-	-	2	1	1	-	-	-	-
Male.....	1	-	-	-	-	2	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	4	-	2	1	5	5	2	-	-	-	1	-
Male.....	2	-	1	-	2	1	2	-	-	-	-	-
Female.....	2	-	1	1	3	4	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	1	-	-	-	-	-	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	453	426	309	293	349	443	601	556	316	123	24	-
Male.....	308	290	208	163	197	208	255	232	107	32	5	-
Female.....	145	136	101	130	152	235	346	324	209	91	19	-
White, both sexes.....	249	250	201	214	284	369	527	506	283	114	21	-
Male.....	170	172	125	121	164	174	225	211	95	29	5	-
Female.....	79	78	76	93	120	195	302	295	188	85	16	-
Black, both sexes.....	200	172	101	75	54	65	59	43	29	8	3	-
Male.....	136	115	77	41	25	28	21	17	10	2	-	-
Female.....	64	57	24	34	29	37	38	26	19	6	3	-
American Indian 1/, both sexes.....	1	-	1	2	2	1	1	-	-	-	-	-
Male.....	-	-	1	-	-	1	1	-	-	-	-	-
Female.....	1	-	-	2	2	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	4	6	2	9	8	14	7	4	1	-	-
Male.....	2	3	5	1	8	5	8	4	2	1	-	-
Female.....	1	1	1	1	1	3	6	3	2	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Ohio

All races, both sexes.....	176	164	157	174	193	315	346	346	209	63	9	-
Male.....	120	107	89	94	79	145	144	131	63	11	4	-
Female.....	56	57	68	80	114	170	202	215	146	52	5	-
White, both sexes.....	129	124	129	150	169	270	303	316	191	57	8	-
Male.....	91	80	76	82	66	121	129	117	57	10	4	-
Female.....	38	44	53	68	103	149	174	199	134	47	4	-
Black, both sexes.....	46	39	28	23	23	44	41	29	18	6	1	-
Male.....	28	26	13	12	13	23	13	13	6	1	-	-
Female.....	18	13	15	11	10	21	28	16	12	5	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	1	1	-	1	1	1	1	1	-	-	-	-
Male.....	1	1	-	-	-	1	1	1	-	-	-	-
Female.....	-	-	-	1	1	-	-	-	-	-	-	-

Oklahoma

All races, both sexes.....	82	74	57	58	83	74	108	106	54	20	1	-
Male.....	48	53	33	30	31	35	43	47	20	4	-	-
Female.....	34	21	24	28	52	39	65	59	34	16	1	-
White, both sexes.....	59	55	47	45	69	64	94	96	50	18	1	-
Male.....	36	42	28	21	25	30	39	42	19	2	-	-
Female.....	23	13	19	24	44	34	55	54	31	16	1	-
Black, both sexes.....	12	6	5	7	7	6	6	5	3	2	-	-
Male.....	8	3	4	6	2	2	2	4	1	2	-	-
Female.....	4	3	1	1	5	4	4	1	2	-	-	-
American Indian 1/, both sexes.....	11	13	5	6	7	4	6	4	-	-	-	-
Male.....	4	8	1	3	4	3	2	1	-	-	-	-
Female.....	7	5	4	3	3	1	4	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	-	2	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	2	1	1	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
Oregon												
All races, both sexes.....	71	74	46	40	45	53	69	61	30	13	1	-
Male.....	45	48	31	21	25	18	35	25	8	3	-	-
Female.....	26	26	15	19	20	35	34	36	22	10	1	-
White, both sexes.....	65	69	42	37	41	50	68	56	29	12	1	-
Male.....	41	46	28	19	24	17	34	24	8	3	-	-
Female.....	24	23	14	18	17	33	34	32	21	9	1	-
Black, both sexes.....	3	2	-	1	-	1	-	2	1	-	-	-
Male.....	2	2	-	1	-	1	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	2	1	-	-	-
American Indian 1/, both sexes.....	2	2	2	-	2	-	1	-	-	-	-	-
Male.....	2	-	2	-	1	-	1	-	-	-	-	-
Female.....	-	2	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	2	2	2	2	2	3	-	1	-	-
Male.....	-	-	1	1	-	-	-	1	-	-	-	-
Female.....	1	1	1	1	2	2	-	2	-	1	-	-
Pennsylvania												
All races, both sexes.....	241	231	239	283	331	492	577	600	318	107	9	-
Male.....	159	147	139	145	176	225	259	257	113	19	2	-
Female.....	82	84	100	138	155	267	318	343	205	88	7	-
White, both sexes.....	155	157	169	220	269	427	522	542	298	98	8	-
Male.....	103	100	96	108	145	195	236	229	106	17	2	-
Female.....	52	57	73	112	124	232	286	313	192	81	6	-
Black, both sexes.....	84	73	68	57	58	59	52	54	20	9	1	-
Male.....	55	47	41	32	28	28	22	26	7	2	-	-
Female.....	29	26	27	25	30	31	30	28	13	7	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	1	2	6	4	6	3	4	-	-	-	-
Male.....	1	-	2	5	3	2	1	2	-	-	-	-
Female.....	1	1	-	1	1	4	2	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Rhode Island

All races, both sexes.....	15	20	18	21	26	40	53	50	33	7	1	-
Male.....	10	16	13	9	14	22	24	23	6	2	-	-
Female.....	5	4	5	12	12	18	29	27	27	5	1	-
White, both sexes.....	12	19	18	20	24	39	51	47	32	7	1	-
Male.....	9	16	13	8	14	22	24	20	5	2	-	-
Female.....	3	3	5	12	10	17	27	27	27	5	1	-
Black, both sexes.....	2	1	-	-	2	-	2	2	1	-	-	-
Male.....	-	-	-	-	-	-	-	2	1	-	-	-
Female.....	2	1	-	-	2	-	2	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	1	-	1	-	-	-	-	-	-
Male.....	1	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

South Carolina

All races, both sexes.....	122	115	121	76	93	107	136	108	67	13	2	-
Male.....	76	69	65	41	49	56	45	47	27	3	-	-
Female.....	46	46	56	35	44	51	91	61	40	10	2	-
White, both sexes.....	45	61	65	44	57	72	98	74	55	10	1	-
Male.....	23	39	30	21	34	36	35	31	23	3	-	-
Female.....	22	22	35	23	23	36	63	43	32	7	1	-
Black, both sexes.....	77	54	55	32	36	31	37	34	12	3	1	-
Male.....	53	30	35	20	15	17	10	16	4	-	-	-
Female.....	24	24	20	12	21	14	27	18	8	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	4	1	-	-	-	-	-
Male.....	-	-	-	-	-	3	-	-	-	-	-	-
Female.....	-	-	1	-	-	1	1	-	-	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
I. Certain infectious and parasitic diseases (A00-B99)												
South Dakota												
All races, both sexes.....	7	7	9	6	9	7	18	19	11	10	-	-
Male.....	5	3	3	3	6	3	10	10	2	2	-	-
Female.....	2	4	6	3	3	4	8	9	9	8	-	-
White, both sexes.....	4	4	5	5	7	7	16	18	11	10	-	-
Male.....	4	1	2	3	5	3	10	10	2	2	-	-
Female.....	-	3	3	2	2	4	6	8	9	8	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	3	3	4	1	2	-	2	1	-	-	-	-
Male.....	1	2	1	-	1	-	-	-	-	-	-	-
Female.....	2	1	3	1	1	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes.....	136	115	101	103	107	119	166	148	88	23	-	-
Male.....	93	69	54	56	56	38	64	60	27	7	-	-
Female.....	43	46	47	47	51	81	102	88	61	16	-	-
White, both sexes.....	95	79	78	80	87	98	146	128	79	18	-	-
Male.....	66	49	43	42	46	33	60	52	25	3	-	-
Female.....	29	30	35	38	41	65	86	76	54	15	-	-
Black, both sexes.....	41	36	23	23	18	21	19	19	9	5	-	-
Male.....	27	20	11	14	9	5	4	8	2	4	-	-
Female.....	14	16	12	9	9	16	15	11	7	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	1	1	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	1	-	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
<b>I. Certain infectious and parasitic diseases (A00-B99)</b>												
<b>Texas</b>												
All races, both sexes.....	437	409	357	370	414	485	575	517	303	92	20	-
Male.....	314	265	203	192	217	222	254	187	85	23	3	-
Female.....	123	144	154	178	197	263	321	330	218	69	17	-
White, both sexes.....	323	324	275	291	340	417	506	455	273	78	17	-
Male.....	236	213	160	155	174	192	228	164	75	18	3	-
Female.....	87	111	115	136	166	225	278	291	198	60	14	-
Black, both sexes.....	112	82	76	71	68	58	59	57	28	13	3	-
Male.....	77	49	40	33	40	26	23	21	9	4	-	-
Female.....	35	33	36	38	28	32	36	36	19	9	3	-
American Indian 1/, both sexes.....	1	-	1	1	-	4	2	1	-	-	-	-
Male.....	-	-	1	1	-	2	-	-	-	-	-	-
Female.....	1	-	-	-	-	2	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	3	5	7	6	6	8	4	2	1	-	-
Male.....	1	3	2	3	3	2	3	2	1	1	-	-
Female.....	-	-	3	4	3	4	5	2	1	-	-	-
<b>Utah</b>												
All races, both sexes.....	25	22	19	12	22	19	27	24	14	2	-	-
Male.....	20	14	10	7	8	6	13	10	6	1	-	-
Female.....	5	8	9	5	14	13	14	14	8	1	-	-
White, both sexes.....	23	19	17	9	19	19	24	24	12	2	-	-
Male.....	18	11	8	5	7	6	12	10	5	1	-	-
Female.....	5	8	9	4	12	13	12	14	7	1	-	-
Black, both sexes.....	-	-	1	-	-	-	1	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	2	1	-	1	1	-	1	-	1	-	-	-
Male.....	2	1	-	1	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	1	2	2	-	1	-	1	-	-	-
Male.....	-	2	1	1	1	-	-	-	1	-	-	-
Female.....	-	-	-	1	1	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Vermont

All races, both sexes.....	11	5	4	5	7	6	10	18	8	1	1	-
Male.....	8	4	3	3	5	4	6	5	3	-	-	-
Female.....	3	1	1	2	2	2	4	13	5	1	1	-
White, both sexes.....	10	4	4	5	7	6	10	18	8	1	1	-
Male.....	8	3	3	3	5	4	6	5	3	-	-	-
Female.....	2	1	1	2	2	2	4	13	5	1	1	-
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	120	123	122	134	184	268	296	283	144	51	20	1
Male.....	75	81	76	76	105	131	140	105	40	11	2	1
Female.....	45	42	46	58	79	137	156	178	104	40	18	-
White, both sexes.....	69	67	76	81	135	216	241	234	111	41	12	-
Male.....	42	45	44	47	74	109	116	86	27	8	-	-
Female.....	27	22	32	34	61	107	125	148	84	33	12	-
Black, both sexes.....	50	52	45	48	41	48	50	49	32	10	8	1
Male.....	33	33	32	27	26	19	22	19	12	3	2	1
Female.....	17	19	13	21	15	29	28	30	20	7	6	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	1	4	1	5	7	4	5	-	1	-	-	-
Male.....	-	3	-	2	5	3	2	-	1	-	-	-
Female.....	1	1	1	3	2	1	3	-	-	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

I. Certain infectious and parasitic diseases (A00-B99)

Wisconsin

All races, both sexes.....	43	42	49	51	89	82	119	108	67	35	6	-
Male.....	30	20	21	23	37	35	46	43	23	6	2	-
Female.....	13	22	28	28	52	47	73	65	44	29	4	-
White, both sexes.....	25	31	40	47	80	81	112	102	66	33	6	-
Male.....	19	18	17	21	35	34	42	41	22	6	2	-
Female.....	6	13	23	26	45	47	70	61	44	27	4	-
Black, both sexes.....	14	9	7	3	6	1	4	4	1	1	-	-
Male.....	9	2	4	1	2	1	2	2	1	-	-	-
Female.....	5	7	3	2	4	-	2	2	-	1	-	-
American Indian 1/, both sexes.....	2	2	1	-	-	-	2	1	-	1	-	-
Male.....	1	-	-	-	-	-	1	-	-	-	-	-
Female.....	1	2	1	-	-	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	1	1	3	-	1	1	-	-	-	-
Male.....	1	-	-	1	-	-	1	-	-	-	-	-
Female.....	1	-	1	-	3	-	-	1	-	-	-	-

Wyoming

All races, both sexes.....	3	4	4	1	10	12	14	7	1	3	-	-
Male.....	1	2	3	1	7	7	8	3	1	-	-	-
Female.....	2	2	1	-	3	5	6	4	-	3	-	-
White, both sexes.....	3	4	3	1	10	12	13	6	1	3	-	-
Male.....	1	2	2	1	7	7	8	2	1	-	-	-
Female.....	2	2	1	-	3	5	5	4	-	3	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	-	1	-	-	-	-
Male.....	-	-	1	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Intestinal infectious diseases (A00-A09)</b>												
<b>Arkansas</b>												
All races, both sexes.....	45	-	-	-	-	-	-	-	-	-	-	-
Male.....	13	-	-	-	-	-	-	-	-	-	-	-
Female.....	32	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	41	-	-	-	-	-	-	-	-	-	-	-
Male.....	11	-	-	-	-	-	-	-	-	-	-	-
Female.....	30	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	4	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
<b>California</b>												
All races, both sexes.....	656	1	4	-	-	1	-	1	1	3	3	11
Male.....	261	1	3	-	-	1	-	-	-	1	1	1
Female.....	395	-	1	-	-	-	-	1	1	2	2	10
White, both sexes.....	588	1	4	-	-	1	-	-	1	2	3	8
Male.....	231	1	3	-	-	1	-	-	-	1	1	-
Female.....	357	-	1	-	-	-	-	-	1	1	2	8
Black, both sexes.....	46	-	-	-	-	-	-	1	-	-	-	1
Male.....	19	-	-	-	-	-	-	-	-	-	-	-
Female.....	27	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	1
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	1
Asian or Pacific Islander 2/, both sexes.....	21	-	-	-	-	-	-	-	-	1	-	1
Male.....	11	-	-	-	-	-	-	-	-	-	-	1
Female.....	10	-	-	-	-	-	-	-	-	1	-	-























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intestinal infectious diseases (A00-A09)

Alaska

All races, both sexes.....	-	-	-	2	-	1	-	1	1	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	1	1	-	-	-
White, both sexes.....	-	-	-	1	-	-	-	-	1	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	2	4	5	15	17	28	34	34	26	5	1	-
Male.....	-	4	4	11	9	12	7	14	7	3	-	-
Female.....	2	-	1	4	8	16	27	20	19	2	1	-
White, both sexes.....	2	4	5	12	14	28	31	32	26	5	1	-
Male.....	-	4	4	8	6	12	7	14	7	3	-	-
Female.....	2	-	1	4	8	16	24	18	19	2	1	-
Black, both sexes.....	-	-	-	1	2	-	1	1	-	-	-	-
Male.....	-	-	-	1	2	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	2	1	-	1	1	-	-	-	-
Male.....	-	-	-	2	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Intestinal infectious diseases (A00-A09)

Arkansas

All races, both sexes.....	-	1	2	4	7	8	6	8	4	4	1	-
Male.....	-	-	1	1	1	3	4	2	-	1	-	-
Female.....	-	1	1	3	6	5	2	6	4	3	1	-
White, both sexes.....	-	1	1	4	6	8	4	8	4	4	1	-
Male.....	-	-	-	1	1	3	3	2	-	1	-	-
Female.....	-	1	1	3	5	5	1	6	4	3	1	-
Black, both sexes.....	-	-	1	-	1	-	2	-	-	-	-	-
Male.....	-	-	1	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	12	20	18	36	60	95	127	151	78	30	4	-
Male.....	6	9	9	19	23	49	53	49	27	6	3	-
Female.....	6	11	9	17	37	46	74	102	51	24	1	-
White, both sexes.....	9	16	14	31	53	81	118	141	76	26	3	-
Male.....	5	7	8	17	17	42	47	47	26	6	2	-
Female.....	4	9	6	14	36	39	71	94	50	20	1	-
Black, both sexes.....	3	4	4	5	4	10	5	5	1	3	-	-
Male.....	1	2	1	2	3	5	4	-	1	-	-	-
Female.....	2	2	3	3	1	5	1	5	-	3	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	3	4	4	5	1	1	1	-
Male.....	-	-	-	-	3	2	2	2	-	-	1	-
Female.....	-	-	-	-	-	2	2	3	1	1	-	-





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Intestinal infectious diseases (A00-A09)												
New Hampshire												
All races, both sexes.....	-	-	2	1	6	12	11	15	13	5	1	-
Male.....	-	-	1	1	3	5	7	6	4	1	-	-
Female.....	-	-	1	-	3	7	4	9	9	4	1	-
White, both sexes.....	-	-	2	1	6	12	11	15	13	5	1	-
Male.....	-	-	1	1	3	5	7	6	4	1	-	-
Female.....	-	-	1	-	3	7	4	9	9	4	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes.....	1	3	6	21	20	38	62	77	36	13	1	-
Male.....	1	2	4	8	5	21	22	26	9	2	-	-
Female.....	-	1	2	13	15	17	40	51	27	11	1	-
White, both sexes.....	1	2	5	20	20	34	61	71	36	13	1	-
Male.....	1	2	3	8	5	19	21	25	9	2	-	-
Female.....	-	-	2	12	15	15	40	46	27	11	1	-
Black, both sexes.....	-	1	-	1	-	4	1	5	-	-	-	-
Male.....	-	-	-	-	-	2	1	1	-	-	-	-
Female.....	-	1	-	1	-	2	-	4	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	1	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intestinal infectious diseases (A00-A09)

New Mexico

All races, both sexes.....	1	-	1	4	2	4	7	5	5	1	-	-
Male.....	1	-	-	1	1	3	3	2	2	1	-	-
Female.....	-	-	1	3	1	1	4	3	3	-	-	-
White, both sexes.....	1	-	1	4	1	3	7	5	5	1	-	-
Male.....	1	-	-	1	-	2	3	2	2	1	-	-
Female.....	-	-	1	3	1	1	4	3	3	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	-	-	-	-	-	-
Male.....	-	-	-	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	5	11	4	18	40	60	99	93	58	21	4	-
Male.....	1	6	4	8	22	28	37	35	19	7	1	-
Female.....	4	5	-	10	18	32	62	58	39	14	3	-
White, both sexes.....	2	9	3	13	35	55	91	88	54	20	3	-
Male.....	-	5	3	6	20	26	33	33	18	7	1	-
Female.....	2	4	-	7	15	29	58	55	36	13	2	-
Black, both sexes.....	3	2	1	5	4	4	6	3	4	1	1	-
Male.....	1	1	1	2	1	2	3	1	1	-	-	-
Female.....	2	1	-	3	3	2	3	2	3	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	1	2	2	-	-	-	-
Male.....	-	-	-	-	1	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	1	1	1	-	-	-	-





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cholera (A00)

United States

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for United States: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Alabama

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alabama: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cholera (A00)

Alaska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cholera (A00)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cholera (A00)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cholera (A00)

Kentucky

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cholera (A00)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cholera (A00)

Massachusetts

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts (All races, White, Black, American Indian, Asian or Pacific Islander) split by sex.

Michigan

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan (All races, White, Black, American Indian, Asian or Pacific Islander) split by sex.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cholera (A00)

Missouri

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cholera (A00)

Nebraska

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cholera (A00)

New Mexico

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cholera (A00)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cholera (A00)

Ohio

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cholera (A00)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cholera (A00)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cholera (A00)

Wisconsin

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cholera (A00)

United States

Table with 13 columns (Age groups) and rows for United States: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Alabama

Table with 13 columns (Age groups) and rows for Alabama: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cholera (A00)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cholera (A00)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Cholera (A00)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cholera (A00)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cholera (A00)

Ohio

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Oklahoma

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other intestinal infectious diseases (A01-A08)

Arkansas

All races, both sexes.....	43	-	-	-	-	-	-	-	-	-	-	-
Male.....	13	-	-	-	-	-	-	-	-	-	-	-
Female.....	30	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	39	-	-	-	-	-	-	-	-	-	-	-
Male.....	11	-	-	-	-	-	-	-	-	-	-	-
Female.....	28	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	4	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	643	1	3	-	-	1	-	1	1	3	3	10
Male.....	254	1	3	-	-	1	-	-	-	1	1	1
Female.....	389	-	-	-	-	-	-	1	1	2	2	9
White, both sexes.....	578	1	3	-	-	1	-	-	1	2	3	8
Male.....	225	1	3	-	-	1	-	-	-	1	1	-
Female.....	353	-	-	-	-	-	-	-	1	1	2	8
Black, both sexes.....	45	-	-	-	-	-	-	1	-	-	-	1
Male.....	18	-	-	-	-	-	-	-	-	-	-	-
Female.....	27	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	20	-	-	-	-	-	-	-	-	1	-	1
Male.....	11	-	-	-	-	-	-	-	-	-	-	1
Female.....	9	-	-	-	-	-	-	-	-	1	-	-























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other intestinal infectious diseases (A01-A08)

Alaska

All races, both sexes.....	-	-	-	2	-	1	-	1	1	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	1	1	-	-	-
White, both sexes.....	-	-	-	1	-	-	-	-	1	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	-	3	5	14	17	26	33	34	25	5	1	-
Male.....	-	3	4	11	9	12	6	14	7	3	-	-
Female.....	-	-	1	3	8	14	27	20	18	2	1	-
White, both sexes.....	-	3	5	11	14	26	30	32	25	5	1	-
Male.....	-	3	4	8	6	12	6	14	7	3	-	-
Female.....	-	-	1	3	8	14	24	18	18	2	1	-
Black, both sexes.....	-	-	-	1	2	-	1	1	-	-	-	-
Male.....	-	-	-	1	2	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	2	1	-	1	1	-	-	-	-
Male.....	-	-	-	2	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other intestinal infectious diseases (A01-A08)

Arkansas

All races, both sexes.....	-	1	2	4	7	8	6	7	3	4	1	-
Male.....	-	-	1	1	1	3	4	2	-	1	-	-
Female.....	-	1	1	3	6	5	2	5	3	3	1	-
White, both sexes.....	-	1	1	4	6	8	4	7	3	4	1	-
Male.....	-	-	-	1	1	3	3	2	-	1	-	-
Female.....	-	1	1	3	5	5	1	5	3	3	1	-
Black, both sexes.....	-	-	1	-	1	-	2	-	-	-	-	-
Male.....	-	-	1	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	11	20	17	36	58	92	126	149	77	30	4	-
Male.....	5	9	9	19	21	47	53	48	26	6	3	-
Female.....	6	11	8	17	37	45	73	101	51	24	1	-
White, both sexes.....	8	16	13	31	51	79	117	140	75	26	3	-
Male.....	4	7	8	17	15	41	47	46	25	6	2	-
Female.....	4	9	5	14	36	38	70	94	50	20	1	-
Black, both sexes.....	3	4	4	5	4	9	5	5	1	3	-	-
Male.....	1	2	1	2	3	4	4	-	1	-	-	-
Female.....	2	2	3	3	1	5	1	5	-	3	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	3	4	4	4	1	1	1	-
Male.....	-	-	-	-	3	2	2	2	-	-	1	-
Female.....	-	-	-	-	-	2	2	2	1	1	-	-





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other intestinal infectious diseases (A01-A08)

New Hampshire

All races, both sexes.....	-	-	2	1	6	12	11	15	13	5	1	-
Male.....	-	-	1	1	3	5	7	6	4	1	-	-
Female.....	-	-	1	-	3	7	4	9	9	4	1	-
White, both sexes.....	-	-	2	1	6	12	11	15	13	5	1	-
Male.....	-	-	1	1	3	5	7	6	4	1	-	-
Female.....	-	-	1	-	3	7	4	9	9	4	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	1	3	6	20	19	38	58	74	36	12	1	-
Male.....	1	2	4	7	5	21	22	25	9	2	-	-
Female.....	-	1	2	13	14	17	36	49	27	10	1	-
White, both sexes.....	1	2	5	19	19	34	57	68	36	12	1	-
Male.....	1	2	3	7	5	19	21	24	9	2	-	-
Female.....	-	-	2	12	14	15	36	44	27	10	1	-
Black, both sexes.....	-	1	-	1	-	4	1	5	-	-	-	-
Male.....	-	-	-	-	-	2	1	1	-	-	-	-
Female.....	-	1	-	1	-	2	-	4	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	1	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other intestinal infectious diseases (A01-A08)

New Mexico

All races, both sexes.....	1	-	1	4	2	4	7	5	5	1	-	-
Male.....	1	-	-	1	1	3	3	2	2	1	-	-
Female.....	-	-	1	3	1	1	4	3	3	-	-	-
White, both sexes.....	1	-	1	4	1	3	7	5	5	1	-	-
Male.....	1	-	-	1	-	2	3	2	2	1	-	-
Female.....	-	-	1	3	1	1	4	3	3	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	-	-	-	-	-	-
Male.....	-	-	-	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	4	11	4	17	40	59	96	91	57	21	4	-
Male.....	1	6	4	7	22	27	35	34	19	7	1	-
Female.....	3	5	-	10	18	32	61	57	38	14	3	-
White, both sexes.....	1	9	3	13	35	54	88	86	53	20	3	-
Male.....	-	5	3	6	20	25	31	32	18	7	1	-
Female.....	1	4	-	7	15	29	57	54	35	13	2	-
Black, both sexes.....	3	2	1	4	4	4	6	3	4	1	1	-
Male.....	1	1	1	1	1	2	3	1	1	-	-	-
Female.....	2	1	-	3	3	2	3	2	3	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	1	2	2	-	-	-	-
Male.....	-	-	-	-	1	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	1	1	1	-	-	-	-





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Missouri

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

New Mexico

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Colorado

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

New Hampshire

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Paratyphoid fevers and other salmonella infections (A01.1-A01.4,A02)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Shigellosis (A03)

Kentucky

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Shigellosis (A03)

Nebraska

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Shigellosis (A03)

New Hampshire

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Shigellosis (A03)

North Carolina

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Shigellosis (A03)

Oregon

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Shigellosis (A03)

Rhode Island

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Shigellosis (A03)

Vermont

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Shigellosis (A03)

Washington

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Shigellosis (A03)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Shigellosis (A03)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Shigellosis (A03)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Shigellosis (A03)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Shigellosis (A03)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other bacterial food-borne intoxications (A05)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other bacterial food-borne intoxications (A05)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other bacterial food-borne intoxications (A05)

Vermont

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other bacterial food-borne intoxications (A05)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other bacterial food-borne intoxications (A05)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other bacterial food-borne intoxications (A05)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other bacterial food-borne intoxications (A05)

Kentucky

Table with 13 columns representing age groups and one column for 'Age not stated'. Rows include 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with sub-rows for 'Male' and 'Female'. All data points are dashes.

Louisiana

Table with 13 columns representing age groups and one column for 'Age not stated'. Rows include 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with sub-rows for 'Male' and 'Female'. All data points are dashes.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other bacterial food-borne intoxications (A05)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other bacterial food-borne intoxications (A05)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Other bacterial food-borne intoxications (A05)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other bacterial food-borne intoxications (A05)

Ohio

Table with 13 columns (Age groups) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Age groups) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Other bacterial food-borne intoxications (A05)

Texas

Table with 13 columns (Age groups) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Age groups) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Other bacterial food-borne intoxications (A05)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85-89	90-94	95-99	100 years	Age not
	years	years	years	years	years	years	years	years	years	years	and over	stated

Other bacterial food-borne intoxications (A05)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Intestinal infections due to other specified organisms (A04,A07-A08)

Arkansas

All races, both sexes.....	41	-	-	-	-	-	-	-	-	-	-	-
Male.....	12	-	-	-	-	-	-	-	-	-	-	-
Female.....	29	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	37	-	-	-	-	-	-	-	-	-	-	-
Male.....	10	-	-	-	-	-	-	-	-	-	-	-
Female.....	27	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	4	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	639	1	3	-	-	1	-	1	1	3	3	10
Male.....	251	1	3	-	-	1	-	-	-	1	1	1
Female.....	388	-	-	-	-	-	-	1	1	2	2	9
White, both sexes.....	575	1	3	-	-	1	-	-	1	2	3	8
Male.....	223	1	3	-	-	1	-	-	-	1	1	-
Female.....	352	-	-	-	-	-	-	-	1	1	2	8
Black, both sexes.....	44	-	-	-	-	-	-	1	-	-	-	1
Male.....	17	-	-	-	-	-	-	-	-	-	-	-
Female.....	27	-	-	-	-	-	-	1	-	-	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	20	-	-	-	-	-	-	-	-	1	-	1
Male.....	11	-	-	-	-	-	-	-	-	-	-	1
Female.....	9	-	-	-	-	-	-	-	-	1	-	-























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intestinal infections due to other specified organisms (A04,A07-A08)

Alaska

All races, both sexes.....	-	-	-	1	-	1	-	1	1	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	1	-	-	-
White, both sexes.....	-	-	-	1	-	-	-	-	1	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	-	3	5	14	17	26	33	34	24	5	1	-
Male.....	-	3	4	11	9	12	6	14	6	3	-	-
Female.....	-	-	1	3	8	14	27	20	18	2	1	-
White, both sexes.....	-	3	5	11	14	26	30	32	24	5	1	-
Male.....	-	3	4	8	6	12	6	14	6	3	-	-
Female.....	-	-	1	3	8	14	24	18	18	2	1	-
Black, both sexes.....	-	-	-	1	2	-	1	1	-	-	-	-
Male.....	-	-	-	1	2	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	2	1	-	1	1	-	-	-	-
Male.....	-	-	-	2	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intestinal infections due to other specified organisms (A04,A07-A08)

Arkansas

All races, both sexes.....	-	1	2	4	6	7	6	7	3	4	1	-
Male.....	-	-	1	1	-	3	4	2	-	1	-	-
Female.....	-	1	1	3	6	4	2	5	3	3	1	-
White, both sexes.....	-	1	1	4	5	7	4	7	3	4	1	-
Male.....	-	-	-	1	-	3	3	2	-	1	-	-
Female.....	-	1	1	3	5	4	1	5	3	3	1	-
Black, both sexes.....	-	-	1	-	1	-	2	-	-	-	-	-
Male.....	-	-	1	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	10	19	17	35	58	92	126	148	77	30	4	-
Male.....	4	8	9	19	21	47	53	47	26	6	3	-
Female.....	6	11	8	16	37	45	73	101	51	24	1	-
White, both sexes.....	8	15	13	30	51	79	117	139	75	26	3	-
Male.....	4	6	8	17	15	41	47	45	25	6	2	-
Female.....	4	9	5	13	36	38	70	94	50	20	1	-
Black, both sexes.....	2	4	4	5	4	9	5	5	1	3	-	-
Male.....	-	2	1	2	3	4	4	-	1	-	-	-
Female.....	2	2	3	3	1	5	1	5	-	3	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	3	4	4	4	1	1	1	-
Male.....	-	-	-	-	3	2	2	2	-	-	1	-
Female.....	-	-	-	-	-	2	2	2	1	1	-	-





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intestinal infections due to other specified organisms (A04,A07-A08)

New Hampshire

All races, both sexes.....	-	-	2	1	6	12	11	15	13	5	1	-
Male.....	-	-	1	1	3	5	7	6	4	1	-	-
Female.....	-	-	1	-	3	7	4	9	9	4	1	-
White, both sexes.....	-	-	2	1	6	12	11	15	13	5	1	-
Male.....	-	-	1	1	3	5	7	6	4	1	-	-
Female.....	-	-	1	-	3	7	4	9	9	4	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	-	3	6	20	19	38	58	74	36	12	1	-
Male.....	-	2	4	7	5	21	22	25	9	2	-	-
Female.....	-	1	2	13	14	17	36	49	27	10	1	-
White, both sexes.....	-	2	5	19	19	34	57	68	36	12	1	-
Male.....	-	2	3	7	5	19	21	24	9	2	-	-
Female.....	-	-	2	12	14	15	36	44	27	10	1	-
Black, both sexes.....	-	1	-	1	-	4	1	5	-	-	-	-
Male.....	-	-	-	-	-	2	1	1	-	-	-	-
Female.....	-	1	-	1	-	2	-	4	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	1	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Intestinal infections due to other specified organisms (A04,A07-A08)

New Mexico

All races, both sexes.....	-	-	1	4	2	4	7	5	5	1	-	-
Male.....	-	-	-	1	1	3	3	2	2	1	-	-
Female.....	-	-	1	3	1	1	4	3	3	-	-	-
White, both sexes.....	-	-	1	4	1	3	7	5	5	1	-	-
Male.....	-	-	-	1	-	2	3	2	2	1	-	-
Female.....	-	-	1	3	1	1	4	3	3	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	1	-	-	-	-	-	-
Male.....	-	-	-	-	1	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	4	11	4	17	40	58	96	91	57	21	4	-
Male.....	1	6	4	7	22	26	35	34	19	7	1	-
Female.....	3	5	-	10	18	32	61	57	38	14	3	-
White, both sexes.....	1	9	3	13	35	53	88	86	53	20	3	-
Male.....	-	5	3	6	20	24	31	32	18	7	1	-
Female.....	1	4	-	7	15	29	57	54	35	13	2	-
Black, both sexes.....	3	2	1	4	4	4	6	3	4	1	1	-
Male.....	1	1	1	1	1	2	3	1	1	-	-	-
Female.....	2	1	-	3	3	2	3	2	3	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	1	2	2	-	-	-	-
Male.....	-	-	-	-	1	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	1	1	1	-	-	-	-



































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diarrhea and gastroenteritis of infectious origin (A09)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diarrhea and gastroenteritis of infectious origin (A09)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diarrhea and gastroenteritis of infectious origin (A09)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diarrhea and gastroenteritis of infectious origin (A09)

Arkansas

All races, both sexes.....	-	-	-	-	-	-	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	1	-	1	-	2	3	1	2	1	-	-	-
Male.....	1	-	-	-	2	2	-	1	1	-	-	-
Female.....	-	-	1	-	-	1	1	1	-	-	-	-
White, both sexes.....	1	-	1	-	2	2	1	1	1	-	-	-
Male.....	1	-	-	-	2	1	-	1	1	-	-	-
Female.....	-	-	1	-	-	1	1	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diarrhea and gastroenteritis of infectious origin (A09)

Colorado

Table with 13 columns (Age groups) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Age groups) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diarrhea and gastroenteritis of infectious origin (A09)

Delaware

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diarrhea and gastroenteritis of infectious origin (A09)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diarrhea and gastroenteritis of infectious origin (A09)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diarrhea and gastroenteritis of infectious origin (A09)

New Hampshire

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diarrhea and gastroenteritis of infectious origin (A09)

North Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diarrhea and gastroenteritis of infectious origin (A09)

Rhode Island

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diarrhea and gastroenteritis of infectious origin (A09)

South Dakota

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diarrhea and gastroenteritis of infectious origin (A09)

Washington

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

West Virginia

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Tuberculosis (A16-A19)

Arkansas

All races, both sexes.....	8	-	-	-	-	-	-	-	-	-	-	-
Male.....	6	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	6	-	-	-	-	-	-	-	-	-	-	-
Male.....	4	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	2	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	104	-	-	-	-	-	1	2	2	4	1	5
Male.....	58	-	-	-	-	-	1	-	-	3	1	2
Female.....	46	-	-	-	-	-	-	2	2	1	-	3
White, both sexes.....	60	-	-	-	-	-	1	1	1	3	-	2
Male.....	33	-	-	-	-	-	1	-	-	2	-	-
Female.....	27	-	-	-	-	-	-	1	1	1	-	2
Black, both sexes.....	7	-	-	-	-	-	-	-	-	-	1	1
Male.....	4	-	-	-	-	-	-	-	-	-	1	-
Female.....	3	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes.....	3	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	34	-	-	-	-	-	-	1	1	1	-	2
Male.....	20	-	-	-	-	-	-	-	-	1	-	2
Female.....	14	-	-	-	-	-	-	1	1	-	-	-

























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Tuberculosis (A16-A19)

Arkansas

All races, both sexes.....	-	1	-	-	-	1	1	2	2	1	-	-
Male.....	-	1	-	-	-	1	1	1	1	1	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes.....	-	-	-	-	-	1	1	2	2	-	-	-
Male.....	-	-	-	-	-	1	1	1	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes.....	-	1	-	-	-	-	-	-	-	1	-	-
Male.....	-	1	-	-	-	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	9	6	8	5	13	13	18	13	4	-	-	-
Male.....	7	3	6	3	6	9	9	6	2	-	-	-
Female.....	2	3	2	2	7	4	9	7	2	-	-	-
White, both sexes.....	5	2	7	3	3	9	9	12	2	-	-	-
Male.....	4	1	5	2	1	6	5	6	-	-	-	-
Female.....	1	1	2	1	2	3	4	6	2	-	-	-
Black, both sexes.....	1	2	-	-	2	-	-	-	-	-	-	-
Male.....	1	-	-	-	2	-	-	-	-	-	-	-
Female.....	-	2	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	1	-	1	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	1	1	1	8	3	9	1	2	-	-	-
Male.....	2	1	1	1	3	3	4	-	2	-	-	-
Female.....	1	-	-	-	5	-	5	1	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Tuberculosis (A16-A19)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Tuberculosis (A16-A19)

New Hampshire

All races, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	1	-	1	3	3	2	3	2	1	-	-	-
Male.....	1	-	1	1	2	-	1	1	-	-	-	-
Female.....	-	-	-	2	1	2	2	1	1	-	-	-
White, both sexes.....	1	-	1	2	1	1	2	2	-	-	-	-
Male.....	1	-	1	1	1	-	1	1	-	-	-	-
Female.....	-	-	-	1	-	1	1	1	-	-	-	-
Black, both sexes.....	-	-	-	-	1	-	1	-	1	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	1	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	1	1	-	-	-	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Tuberculosis (A16-A19)												
Ohio												
All races, both sexes.....	1	2	2	1	3	3	4	1	1	1	-	-
Male.....	1	2	1	-	1	2	3	1	-	-	-	-
Female.....	-	-	1	1	2	1	1	-	1	1	-	-
White, both sexes.....	1	1	-	1	3	2	4	1	1	-	-	-
Male.....	1	1	-	-	1	2	3	1	-	-	-	-
Female.....	-	-	-	1	2	-	1	-	1	-	-	-
Black, both sexes.....	-	-	2	-	-	1	-	-	-	1	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes.....	1	2	-	-	-	1	2	2	-	-	-	-
Male.....	-	2	-	-	-	-	1	1	-	-	-	-
Female.....	1	-	-	-	-	1	1	1	-	-	-	-
White, both sexes.....	1	1	-	-	-	1	2	1	-	-	-	-
Male.....	-	1	-	-	-	-	1	1	-	-	-	-
Female.....	1	-	-	-	-	1	1	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Respiratory tuberculosis (A16)												
Arkansas												
All races, both sexes.....	6	-	-	-	-	-	-	-	-	-	-	-
Male.....	4	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	5	-	-	-	-	-	-	-	-	-	-	-
Male.....	3	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	80	-	-	-	-	-	-	2	1	4	1	3
Male.....	48	-	-	-	-	-	-	-	-	3	1	2
Female.....	32	-	-	-	-	-	-	2	1	1	-	1
White, both sexes.....	44	-	-	-	-	-	-	1	-	3	-	1
Male.....	26	-	-	-	-	-	-	-	-	2	-	-
Female.....	18	-	-	-	-	-	-	1	-	1	-	1
Black, both sexes.....	4	-	-	-	-	-	-	-	-	-	1	-
Male.....	3	-	-	-	-	-	-	-	-	-	1	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	30	-	-	-	-	-	-	1	1	1	-	2
Male.....	19	-	-	-	-	-	-	-	-	1	-	2
Female.....	11	-	-	-	-	-	-	1	1	-	-	-



























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

Arkansas

All races, both sexes.....	-	-	-	-	-	1	-	2	2	1	-	-
Male.....	-	-	-	-	-	1	-	1	1	1	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-
White, both sexes.....	-	-	-	-	-	1	-	2	2	-	-	-
Male.....	-	-	-	-	-	1	-	1	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	1	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	6	4	7	4	9	10	14	11	4	-	-	-
Male.....	5	2	6	3	5	7	7	5	2	-	-	-
Female.....	1	2	1	1	4	3	7	6	2	-	-	-
White, both sexes.....	2	2	6	2	2	6	7	10	2	-	-	-
Male.....	2	1	5	2	1	4	4	5	-	-	-	-
Female.....	-	1	1	-	1	2	3	5	2	-	-	-
Black, both sexes.....	1	1	-	-	1	-	-	-	-	-	-	-
Male.....	1	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	1	1	1	6	3	7	1	2	-	-	-
Male.....	2	1	1	1	3	3	3	-	2	-	-	-
Female.....	1	-	-	-	3	-	4	1	-	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Respiratory tuberculosis (A16)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Respiratory tuberculosis (A16)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Respiratory tuberculosis (A16)

New Hampshire

All races, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	1	-	1	2	2	1	2	2	1	-	-	-
Male.....	1	-	1	-	1	-	-	1	-	-	-	-
Female.....	-	-	-	2	1	1	2	1	1	-	-	-
White, both sexes.....	1	-	1	1	-	1	1	2	-	-	-	-
Male.....	1	-	1	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	1	-	1	1	1	-	-	-	-
Black, both sexes.....	-	-	-	-	1	-	1	-	1	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	1	-	-	-	-	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Respiratory tuberculosis (A16)

Ohio

All races, both sexes.....	-	1	2	-	2	2	2	1	-	1	-	-
Male.....	-	1	1	-	1	1	2	1	-	-	-	-
Female.....	-	-	1	-	1	1	-	-	-	1	-	-
White, both sexes.....	-	1	-	-	2	1	2	1	-	-	-	-
Male.....	-	1	-	-	1	1	2	1	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Black, both sexes.....	-	-	2	-	-	1	-	-	-	1	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	1	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Oklahoma

All races, both sexes.....	1	1	-	-	-	1	1	2	-	-	-	-
Male.....	-	1	-	-	-	-	1	1	-	-	-	-
Female.....	1	-	-	-	-	1	-	1	-	-	-	-
White, both sexes.....	1	1	-	-	-	1	1	1	-	-	-	-
Male.....	-	1	-	-	-	-	1	1	-	-	-	-
Female.....	1	-	-	-	-	1	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-















































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other tuberculosis (A17-A19)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other tuberculosis (A17-A19)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other tuberculosis (A17-A19)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other tuberculosis (A17-A19)

Arkansas

All races, both sexes.....	-	1	-	-	-	-	1	-	-	-	-	-
Male.....	-	1	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	3	2	1	1	4	3	4	2	-	-	-	-
Male.....	2	1	-	-	1	2	2	1	-	-	-	-
Female.....	1	1	1	1	3	1	2	1	-	-	-	-
White, both sexes.....	3	-	1	1	1	3	2	2	-	-	-	-
Male.....	2	-	-	-	-	2	1	1	-	-	-	-
Female.....	1	-	1	1	1	1	1	1	-	-	-	-
Black, both sexes.....	-	1	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	2	-	2	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	2	-	1	-	-	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other tuberculosis (A17-A19)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other tuberculosis (A17-A19)

New Hampshire

All races, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	-	-	-	1	1	1	1	-	-	-	-	-
Male.....	-	-	-	1	1	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
White, both sexes.....	-	-	-	1	1	-	1	-	-	-	-	-
Male.....	-	-	-	1	1	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>Zoonotic and other bacterial diseases (A20-A49)</b>												
<b>Arkansas</b>												
All races, both sexes.....	476	3	2	-	1	2	1	1	4	6	6	19
Male.....	202	1	2	-	1	-	-	-	2	3	2	10
Female.....	274	2	-	-	-	2	1	1	2	3	4	9
White, both sexes.....	385	1	1	-	1	1	1	1	4	4	6	14
Male.....	163	1	1	-	1	-	-	-	2	2	2	8
Female.....	222	-	-	-	-	1	1	1	2	2	4	6
Black, both sexes.....	91	2	1	-	-	1	-	-	-	2	-	5
Male.....	39	-	1	-	-	-	-	-	-	1	-	2
Female.....	52	2	-	-	-	1	-	-	-	1	-	3
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
<b>California</b>												
All races, both sexes.....	1,208	19	6	4	3	4	4	6	9	15	33	41
Male.....	584	9	2	2	2	2	1	1	2	9	22	16
Female.....	624	10	4	2	1	2	3	5	7	6	11	25
White, both sexes.....	991	15	4	3	3	4	2	6	6	15	23	27
Male.....	466	9	2	2	2	2	1	1	1	9	17	13
Female.....	525	6	2	1	1	2	1	5	5	6	6	14
Black, both sexes.....	109	4	1	1	-	-	1	-	3	-	5	10
Male.....	49	-	-	-	-	-	-	-	1	-	1	1
Female.....	60	4	1	1	-	-	1	-	2	-	4	9
American Indian 1/, both sexes.....	9	-	-	-	-	-	-	-	-	-	2	-
Male.....	8	-	-	-	-	-	-	-	-	-	2	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	99	-	1	-	-	-	1	-	-	-	3	4
Male.....	61	-	-	-	-	-	-	-	-	-	2	2
Female.....	38	-	1	-	-	-	1	-	-	-	1	2











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Zoonotic and other bacterial diseases (A20-A49)												
Illinois												
All races, both sexes.....	2,184	15	5	-	3	7	4	11	5	18	37	48
Male.....	989	8	3	-	-	4	4	5	1	10	19	31
Female.....	1,195	7	2	-	3	3	-	6	4	8	18	17
White, both sexes.....	1,642	8	5	-	3	5	2	5	2	8	22	29
Male.....	744	4	3	-	-	2	2	3	-	2	10	18
Female.....	898	4	2	-	3	3	-	2	2	6	12	11
Black, both sexes.....	506	7	-	-	-	2	2	6	3	10	15	18
Male.....	232	4	-	-	-	2	2	2	1	8	9	12
Female.....	274	3	-	-	-	-	-	4	2	2	6	6
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	35	-	-	-	-	-	-	-	-	-	-	1
Male.....	13	-	-	-	-	-	-	-	-	-	-	1
Female.....	22	-	-	-	-	-	-	-	-	-	-	-
Indiana												
All races, both sexes.....	899	6	5	1	1	-	3	7	4	9	18	22
Male.....	378	5	2	1	1	-	3	4	2	4	9	7
Female.....	521	1	3	-	-	-	-	3	2	5	9	15
White, both sexes.....	778	4	3	-	1	-	2	6	3	7	13	15
Male.....	329	3	-	-	1	-	2	4	2	3	7	5
Female.....	449	1	3	-	-	-	-	2	1	4	6	10
Black, both sexes.....	118	2	2	1	-	-	1	1	1	1	5	6
Male.....	48	2	2	1	-	-	1	-	-	1	2	2
Female.....	70	-	-	-	-	-	-	1	1	-	3	4
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	-	-	-	1	-	1
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	1	-	1



































## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

## Zoonotic and other bacterial diseases (A20-A49)

## Vermont

All races, both sexes.....	35	-	-	-	-	-	-	-	-	-	-	1
Male.....	13	-	-	-	-	-	-	-	-	-	-	-
Female.....	22	-	-	-	-	-	-	-	-	-	-	1
White, both sexes.....	34	-	-	-	-	-	-	-	-	-	-	1
Male.....	13	-	-	-	-	-	-	-	-	-	-	-
Female.....	21	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

## Virginia

All races, both sexes.....	1,417	13	3	-	1	-	2	13	9	12	20	31
Male.....	655	4	3	-	1	-	-	8	6	5	9	20
Female.....	762	9	-	-	-	-	2	5	3	7	11	11
White, both sexes.....	1,030	7	3	-	1	-	1	8	3	8	7	20
Male.....	475	3	3	-	1	-	-	6	3	2	4	14
Female.....	555	4	-	-	-	-	1	2	-	6	3	6
Black, both sexes.....	361	4	-	-	-	-	1	4	3	4	13	11
Male.....	165	-	-	-	-	-	-	1	1	3	5	6
Female.....	196	4	-	-	-	-	1	3	2	1	8	5
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	26	2	-	-	-	-	-	1	3	-	-	-
Male.....	15	1	-	-	-	-	-	1	2	-	-	-
Female.....	11	1	-	-	-	-	-	-	1	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

United States

All races, both sexes.....	1,497	2,070	2,405	2,935	3,711	4,845	6,052	5,572	3,187	1,074	191	2
Male.....	808	1,119	1,226	1,518	1,902	2,307	2,714	2,175	1,070	261	26	1
Female.....	689	951	1,179	1,417	1,809	2,538	3,338	3,397	2,117	813	165	1
White, both sexes.....	1,012	1,483	1,824	2,224	2,870	3,974	5,136	4,833	2,756	919	150	1
Male.....	541	804	924	1,149	1,473	1,922	2,363	1,904	925	214	19	1
Female.....	471	679	900	1,075	1,397	2,052	2,773	2,929	1,831	705	131	-
Black, both sexes.....	438	535	520	642	724	764	807	644	380	143	37	-
Male.....	242	284	269	326	359	332	303	233	118	41	7	-
Female.....	196	251	251	316	365	432	504	411	262	102	30	-
American Indian 1/, both sexes.....	27	22	24	19	33	23	22	14	8	1	2	-
Male.....	13	10	11	10	15	9	10	2	4	-	-	-
Female.....	14	12	13	9	18	14	12	12	4	1	2	-
Asian or Pacific Islander 2/, both sexes.....	20	30	37	50	84	84	87	81	43	11	2	1
Male.....	12	21	22	33	55	44	38	36	23	6	-	-
Female.....	8	9	15	17	29	40	49	45	20	5	2	1

Alabama

All races, both sexes.....	48	48	62	72	95	111	115	98	51	22	6	-
Male.....	21	29	28	40	42	45	42	33	22	5	1	-
Female.....	27	19	34	32	53	66	73	65	29	17	5	-
White, both sexes.....	33	28	47	48	58	88	93	80	36	14	1	-
Male.....	15	17	20	26	22	34	38	29	15	2	-	-
Female.....	18	11	27	22	36	54	55	51	21	12	1	-
Black, both sexes.....	15	20	15	24	37	23	22	17	15	8	5	-
Male.....	6	12	8	14	20	11	4	4	7	3	1	-
Female.....	9	8	7	10	17	12	18	13	8	5	4	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

Alaska

All races, both sexes.....	2	7	2	3	2	4	4	4	4	-	-	-
Male.....	1	3	-	1	-	3	2	2	2	-	-	-
Female.....	1	4	2	2	2	1	2	2	2	-	-	-
White, both sexes.....	1	2	1	3	-	2	1	3	3	-	-	-
Male.....	1	-	-	1	-	2	1	2	1	-	-	-
Female.....	-	2	1	2	-	-	-	1	2	-	-	-
Black, both sexes.....	-	1	-	-	-	1	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	4	1	-	2	1	3	1	-	-	-	-
Male.....	-	2	-	-	-	1	1	-	-	-	-	-
Female.....	1	2	1	-	2	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	31	36	51	49	73	102	86	70	33	6	1	1
Male.....	17	18	26	25	36	44	42	31	13	2	-	1
Female.....	14	18	25	24	37	58	44	39	20	4	1	-
White, both sexes.....	20	33	44	45	61	89	81	67	30	6	1	1
Male.....	11	17	23	21	31	40	40	31	12	2	-	1
Female.....	9	16	21	24	30	49	41	36	18	4	1	-
Black, both sexes.....	3	2	3	1	3	5	1	1	1	-	-	-
Male.....	2	1	2	1	1	2	-	-	1	-	-	-
Female.....	1	1	1	-	2	3	1	1	-	-	-	-
American Indian 1/, both sexes.....	8	-	4	2	5	7	1	2	1	-	-	-
Male.....	4	-	1	2	2	2	-	-	-	-	-	-
Female.....	4	-	3	-	3	5	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	1	-	1	4	1	3	-	1	-	-	-
Male.....	-	-	-	1	2	-	2	-	-	-	-	-
Female.....	-	1	-	-	2	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

Arkansas

All races, both sexes.....	13	31	37	35	49	57	88	68	39	12	2	-
Male.....	7	13	16	17	27	25	41	19	13	3	-	-
Female.....	6	18	21	18	22	32	47	49	26	9	2	-
White, both sexes.....	7	25	29	25	41	48	71	58	34	11	2	-
Male.....	4	10	12	12	22	20	34	18	11	3	-	-
Female.....	3	15	17	13	19	28	37	40	23	8	2	-
Black, both sexes.....	6	6	8	10	8	9	17	10	5	1	-	-
Male.....	3	3	4	5	5	5	7	1	2	-	-	-
Female.....	3	3	4	5	3	4	10	9	3	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	48	84	78	90	121	123	196	180	101	36	7	-
Male.....	26	49	35	47	76	61	95	74	39	14	-	-
Female.....	22	35	43	43	45	62	101	106	62	22	7	-
White, both sexes.....	40	69	66	77	92	98	167	148	87	32	7	-
Male.....	23	38	27	39	50	46	81	60	31	12	-	-
Female.....	17	31	39	38	42	52	86	88	56	20	7	-
Black, both sexes.....	7	10	5	10	13	11	12	9	6	1	-	-
Male.....	3	6	2	6	11	4	7	4	2	1	-	-
Female.....	4	4	3	4	2	7	5	5	4	-	-	-
American Indian 1/, both sexes.....	-	1	1	-	3	1	-	1	-	-	-	-
Male.....	-	1	1	-	3	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	4	6	3	13	13	17	22	8	3	-	-
Male.....	-	4	5	2	12	10	7	10	6	1	-	-
Female.....	1	-	1	1	1	3	10	12	2	2	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

Colorado

All races, both sexes.....	13	24	19	26	36	45	51	40	34	5	1	-
Male.....	11	11	6	12	21	24	25	17	13	1	1	-
Female.....	2	13	13	14	15	21	26	23	21	4	-	-
White, both sexes.....	12	22	17	23	33	42	50	38	33	5	1	-
Male.....	10	10	5	11	18	24	24	15	12	1	1	-
Female.....	2	12	12	12	15	18	26	23	21	4	-	-
Black, both sexes.....	1	2	2	2	2	2	1	1	1	-	-	-
Male.....	1	1	1	-	2	-	1	1	1	-	-	-
Female.....	-	1	1	2	-	2	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	1	-	1	-	-	-	-
Male.....	-	-	-	1	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Connecticut

All races, both sexes.....	23	22	37	44	51	68	133	133	65	27	4	-
Male.....	10	13	21	19	26	41	60	54	19	6	-	-
Female.....	13	9	16	25	25	27	73	79	46	21	4	-
White, both sexes.....	17	18	32	34	43	65	122	128	63	27	4	-
Male.....	6	12	18	14	23	40	55	52	17	6	-	-
Female.....	11	6	14	20	20	25	67	76	46	21	4	-
Black, both sexes.....	5	4	5	10	7	2	11	3	1	-	-	-
Male.....	3	1	3	5	2	1	5	1	1	-	-	-
Female.....	2	3	2	5	5	1	6	2	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	1	-	2	1	-	-	-
Male.....	1	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

Florida

All races, both sexes.....	86	115	97	147	198	301	319	302	173	54	10	-
Male.....	43	66	49	79	108	158	159	128	60	18	1	-
Female.....	43	49	48	68	90	143	160	174	113	36	9	-
White, both sexes.....	58	90	78	119	157	257	281	278	161	48	9	-
Male.....	28	53	43	62	88	138	141	123	54	16	1	-
Female.....	30	37	35	57	69	119	140	155	107	32	8	-
Black, both sexes.....	28	25	19	25	40	39	36	24	12	6	1	-
Male.....	15	13	6	14	20	19	17	5	6	2	-	-
Female.....	13	12	13	11	20	20	19	19	6	4	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	3	1	5	2	-	-	-	-	-
Male.....	-	-	-	3	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	1	4	1	-	-	-	-	-

Georgia

All races, both sexes.....	78	114	124	129	154	190	196	190	94	42	7	-
Male.....	45	64	71	68	74	92	85	58	23	9	-	-
Female.....	33	50	53	61	80	98	111	132	71	33	7	-
White, both sexes.....	39	59	64	80	91	130	138	140	64	25	5	-
Male.....	25	29	37	42	41	64	62	42	18	5	-	-
Female.....	14	30	27	38	50	66	76	98	46	20	5	-
Black, both sexes.....	39	52	55	48	61	57	58	49	30	17	2	-
Male.....	20	32	30	25	31	26	23	16	5	4	-	-
Female.....	19	20	25	23	30	31	35	33	25	13	2	-
American Indian 1/, both sexes.....	-	1	1	-	-	-	-	-	-	-	-	-
Male.....	-	1	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	4	1	2	3	-	1	-	-	-	-
Male.....	-	2	3	1	2	2	-	-	-	-	-	-
Female.....	-	-	1	-	-	1	-	1	-	-	-	-











## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

## Zoonotic and other bacterial diseases (A20-A49)

## Maine

All races, both sexes.....	4	9	13	13	9	17	22	31	13	7	-	-
Male.....	2	6	5	8	5	8	13	14	4	1	-	-
Female.....	2	3	8	5	4	9	9	17	9	6	-	-
White, both sexes.....	3	9	13	13	9	17	22	31	12	7	-	-
Male.....	1	6	5	8	5	8	13	14	3	1	-	-
Female.....	2	3	8	5	4	9	9	17	9	6	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

## Maryland

All races, both sexes.....	41	45	59	91	87	142	144	169	87	25	3	-
Male.....	15	28	29	53	43	72	66	64	29	6	-	-
Female.....	26	17	30	38	44	70	78	105	58	19	3	-
White, both sexes.....	17	17	42	44	48	97	108	133	64	20	2	-
Male.....	5	11	22	25	24	50	52	53	24	6	-	-
Female.....	12	6	20	19	24	47	56	80	40	14	2	-
Black, both sexes.....	23	26	16	45	36	43	35	35	22	5	1	-
Male.....	10	15	6	26	16	21	13	11	5	-	-	-
Female.....	13	11	10	19	20	22	22	24	17	5	1	-
American Indian 1/, both sexes.....	-	-	-	1	1	-	-	-	-	-	-	-
Male.....	-	-	-	1	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	2	1	1	2	2	1	1	1	-	-	-
Male.....	-	2	1	1	2	1	1	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	1	1	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

Nebraska

All races, both sexes.....	8	3	11	15	14	17	24	24	15	8	-	-
Male.....	7	2	8	7	4	6	15	11	4	3	-	-
Female.....	1	1	3	8	10	11	9	13	11	5	-	-
White, both sexes.....	7	3	10	15	11	15	22	22	14	8	-	-
Male.....	6	2	7	7	3	5	14	10	4	3	-	-
Female.....	1	1	3	8	8	10	8	12	10	5	-	-
Black, both sexes.....	1	-	-	-	3	1	2	-	1	-	-	-
Male.....	1	-	-	-	1	1	1	-	-	-	-	-
Female.....	-	-	-	-	2	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	-	1	-	2	-	-	-	-
Male.....	-	-	1	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	24	24	33	48	68	58	71	34	27	4	2	-
Male.....	14	12	16	25	34	31	28	20	16	1	-	-
Female.....	10	12	17	23	34	27	43	14	11	3	2	-
White, both sexes.....	19	19	28	38	57	48	66	27	22	4	1	-
Male.....	10	10	13	20	30	28	26	18	13	1	-	-
Female.....	9	9	15	18	27	20	40	9	9	3	1	-
Black, both sexes.....	5	3	2	6	3	9	2	3	2	-	1	-
Male.....	4	1	2	3	2	2	2	1	1	-	-	-
Female.....	1	2	-	3	1	7	-	2	1	-	1	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	3	3	8	1	3	4	2	-	-	-
Male.....	-	1	1	2	2	1	-	1	1	-	-	-
Female.....	-	1	2	1	6	-	3	3	1	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
New Hampshire												
All races, both sexes.....	6	10	12	9	10	10	17	24	13	4	1	-
Male.....	3	7	9	7	7	6	7	8	6	1	1	-
Female.....	3	3	3	2	3	4	10	16	7	3	-	-
White, both sexes.....	6	10	11	9	10	10	17	24	13	4	-	-
Male.....	3	7	8	7	7	6	7	8	6	1	-	-
Female.....	3	3	3	2	3	4	10	16	7	3	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	1	-
Male.....	-	-	-	-	-	-	-	-	-	-	1	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes.....	62	69	94	105	178	239	326	318	183	75	8	-
Male.....	30	39	50	52	109	121	139	111	52	14	2	-
Female.....	32	30	44	53	69	118	187	207	131	61	6	-
White, both sexes.....	39	44	70	71	131	196	279	280	167	65	8	-
Male.....	18	24	37	33	82	100	118	97	48	11	2	-
Female.....	21	20	33	38	49	96	161	183	119	54	6	-
Black, both sexes.....	22	24	21	32	40	38	42	32	16	9	-	-
Male.....	11	14	11	18	22	19	19	12	4	2	-	-
Female.....	11	10	10	14	18	19	23	20	12	7	-	-
American Indian 1/, both sexes.....	1	-	1	-	1	-	-	1	-	-	-	-
Male.....	1	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	2	2	6	5	5	5	-	1	-	-
Male.....	-	1	1	1	5	2	2	2	-	1	-	-
Female.....	-	-	1	1	1	3	3	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

New Mexico

All races, both sexes.....	13	13	11	9	21	18	30	21	12	3	1	-
Male.....	6	9	6	6	9	11	10	8	6	-	-	-
Female.....	7	4	5	3	12	7	20	13	6	3	1	-
White, both sexes.....	11	12	9	7	18	16	27	20	12	3	-	-
Male.....	5	8	5	5	9	11	8	7	6	-	-	-
Female.....	6	4	4	2	9	5	19	13	6	3	-	-
Black, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	2	-	2	1	3	2	2	-	-	-	1	-
Male.....	1	-	1	-	-	-	2	-	-	-	-	-
Female.....	1	-	1	1	3	2	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	1	-	-	-	-	-	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	80	104	130	173	226	322	451	428	237	97	19	-
Male.....	46	58	67	87	120	155	195	177	76	24	3	-
Female.....	34	46	63	86	106	167	256	251	161	73	16	-
White, both sexes.....	54	76	106	127	190	270	397	392	209	90	17	-
Male.....	30	40	53	68	104	132	173	164	66	21	3	-
Female.....	24	36	53	59	86	138	224	228	143	69	14	-
Black, both sexes.....	23	28	20	44	32	48	45	32	25	6	2	-
Male.....	14	18	10	19	13	19	17	10	9	2	-	-
Female.....	9	10	10	25	19	29	28	22	16	4	2	-
American Indian 1/, both sexes.....	1	-	1	2	1	1	1	-	-	-	-	-
Male.....	-	-	1	-	-	1	1	-	-	-	-	-
Female.....	1	-	-	2	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	3	-	3	3	8	4	3	1	-	-
Male.....	2	-	3	-	3	3	4	3	1	1	-	-
Female.....	-	-	-	-	-	-	4	1	2	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

Ohio

All races, both sexes.....	55	80	95	120	126	181	201	183	128	28	6	-
Male.....	33	48	47	69	52	86	101	59	40	5	2	-
Female.....	22	32	48	51	74	95	100	124	88	23	4	-
White, both sexes.....	41	60	84	106	107	151	168	163	113	24	5	-
Male.....	27	37	45	60	44	71	88	52	34	4	2	-
Female.....	14	23	39	46	63	80	80	111	79	20	3	-
Black, both sexes.....	14	20	11	14	18	29	31	20	15	4	1	-
Male.....	6	11	2	9	8	14	11	7	6	1	-	-
Female.....	8	9	9	5	10	15	20	13	9	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-

Oklahoma

All races, both sexes.....	31	24	30	41	52	45	79	66	40	9	1	-
Male.....	15	10	19	19	17	19	32	31	16	1	-	-
Female.....	16	14	11	22	35	26	47	35	24	8	1	-
White, both sexes.....	17	17	23	32	41	37	68	58	36	8	1	-
Male.....	8	10	16	13	13	15	28	26	15	-	-	-
Female.....	9	7	7	19	28	22	40	32	21	8	1	-
Black, both sexes.....	9	2	4	5	6	5	5	4	3	1	-	-
Male.....	6	-	3	4	2	2	2	4	1	1	-	-
Female.....	3	2	1	1	4	3	3	-	2	-	-	-
American Indian 1/, both sexes.....	5	5	3	4	5	3	4	4	-	-	-	-
Male.....	1	-	-	2	2	2	2	1	-	-	-	-
Female.....	4	5	3	2	3	1	2	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

Oregon

All races, both sexes.....	15	12	12	24	23	27	52	39	17	7	-	-
Male.....	8	9	5	11	14	8	25	17	5	1	-	-
Female.....	7	3	7	13	9	19	27	22	12	6	-	-
White, both sexes.....	14	11	11	24	22	26	52	35	17	7	-	-
Male.....	7	8	4	11	13	7	25	16	5	1	-	-
Female.....	7	3	7	13	9	19	27	19	12	6	-	-
Black, both sexes.....	-	1	-	-	-	1	-	2	-	-	-	-
Male.....	-	1	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes.....	1	-	1	-	1	-	-	-	-	-	-	-
Male.....	1	-	1	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	2	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Pennsylvania

All races, both sexes.....	89	120	170	210	252	379	439	448	243	80	8	-
Male.....	54	68	91	98	133	176	198	195	89	15	1	-
Female.....	35	52	79	112	119	203	241	253	154	65	7	-
White, both sexes.....	63	92	135	167	201	327	396	397	227	75	7	-
Male.....	37	53	74	76	108	152	178	169	83	13	1	-
Female.....	26	39	61	91	93	175	218	228	144	62	6	-
Black, both sexes.....	25	28	34	39	48	48	41	47	16	5	1	-
Male.....	17	15	16	19	23	22	19	24	6	2	-	-
Female.....	8	13	18	20	25	26	22	23	10	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	1	4	3	4	2	4	-	-	-	-
Male.....	-	-	1	3	2	2	1	2	-	-	-	-
Female.....	1	-	-	1	1	2	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

Rhode Island

All races, both sexes.....	7	7	10	12	15	15	34	32	17	5	-	-
Male.....	6	5	6	6	8	8	16	14	3	1	-	-
Female.....	1	2	4	6	7	7	18	18	14	4	-	-
White, both sexes.....	6	7	10	11	14	14	32	31	16	5	-	-
Male.....	5	5	6	5	8	8	16	13	2	1	-	-
Female.....	1	2	4	6	6	6	16	18	14	4	-	-
Black, both sexes.....	-	-	-	-	1	-	2	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	1	-	1	-	-	-	-	-	-
Male.....	1	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

South Carolina

All races, both sexes.....	27	53	83	61	74	89	109	89	55	11	2	-
Male.....	8	27	37	34	39	43	36	43	23	3	-	-
Female.....	19	26	46	27	35	46	73	46	32	8	2	-
White, both sexes.....	14	31	48	34	44	60	78	58	47	8	1	-
Male.....	3	17	19	16	28	28	28	28	19	3	-	-
Female.....	11	14	29	18	16	32	50	30	28	5	1	-
Black, both sexes.....	13	22	34	27	30	26	30	31	8	3	1	-
Male.....	5	10	18	18	11	13	8	15	4	-	-	-
Female.....	8	12	16	9	19	13	22	16	4	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	3	1	-	-	-	-	-
Male.....	-	-	-	-	-	2	-	-	-	-	-	-
Female.....	-	-	1	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

South Dakota

All races, both sexes.....	1	5	5	5	5	7	10	11	6	7	-	-
Male.....	-	1	2	3	4	3	6	5	-	2	-	-
Female.....	1	4	3	2	1	4	4	6	6	5	-	-
White, both sexes.....	-	3	3	4	4	7	9	10	6	7	-	-
Male.....	-	-	1	3	3	3	6	5	-	2	-	-
Female.....	-	3	2	1	1	4	3	5	6	5	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	2	2	1	1	-	1	1	-	-	-	-
Male.....	-	1	1	-	1	-	-	-	-	-	-	-
Female.....	1	1	1	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Tennessee

All races, both sexes.....	34	54	65	63	74	79	114	108	58	15	-	-
Male.....	20	30	32	32	38	26	52	48	20	4	-	-
Female.....	14	24	33	31	36	53	62	60	38	11	-	-
White, both sexes.....	26	36	54	44	61	64	99	91	51	11	-	-
Male.....	16	21	29	22	31	22	49	40	19	1	-	-
Female.....	10	15	25	22	30	42	50	51	32	10	-	-
Black, both sexes.....	8	18	11	19	13	15	14	16	7	4	-	-
Male.....	4	9	3	10	7	4	3	8	1	3	-	-
Female.....	4	9	8	9	6	11	11	8	6	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Zoonotic and other bacterial diseases (A20-A49)

Texas

All races, both sexes.....	127	195	193	250	303	375	466	417	253	77	17	-
Male.....	78	106	101	123	165	172	213	147	69	18	2	-
Female.....	49	89	92	127	138	203	253	270	184	59	15	-
White, both sexes.....	100	157	153	194	247	321	403	363	226	63	14	-
Male.....	59	86	83	99	131	150	191	129	60	13	2	-
Female.....	41	71	70	95	116	171	212	234	166	50	12	-
Black, both sexes.....	26	37	39	52	53	47	55	51	25	13	3	-
Male.....	18	19	17	22	31	20	19	17	8	4	-	-
Female.....	8	18	22	30	22	27	36	34	17	9	3	-
American Indian 1/, both sexes.....	-	-	1	1	-	3	2	-	-	-	-	-
Male.....	-	-	1	1	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	2	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	3	3	4	6	3	2	1	-	-
Male.....	1	1	-	1	3	1	3	1	1	1	-	-
Female.....	-	-	-	2	-	3	3	2	1	-	-	-

Utah

All races, both sexes.....	5	9	8	7	9	11	20	19	10	2	-	-
Male.....	2	6	3	5	3	3	9	9	5	1	-	-
Female.....	3	3	5	2	6	8	11	10	5	1	-	-
White, both sexes.....	5	7	8	5	8	11	19	19	8	2	-	-
Male.....	2	4	3	3	3	3	9	9	4	1	-	-
Female.....	3	3	5	2	5	8	10	10	4	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	1	1	-	-	-	1	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	1	-	-	1	-	1	-	-	-
Male.....	-	1	-	1	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Zoonotic and other bacterial diseases (A20-A49)												
Vermont												
All races, both sexes.....	3	2	2	3	4	3	4	9	2	1	1	-
Male.....	1	1	1	2	3	1	2	2	-	-	-	-
Female.....	2	1	1	1	1	2	2	7	2	1	1	-
White, both sexes.....	2	2	2	3	4	3	4	9	2	1	1	-
Male.....	1	1	1	2	3	1	2	2	-	-	-	-
Female.....	1	1	1	1	1	2	2	7	2	1	1	-
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes.....	42	67	85	107	147	215	239	226	123	43	19	-
Male.....	17	38	49	60	86	105	109	87	36	10	2	-
Female.....	25	29	36	47	61	110	130	139	87	33	17	-
White, both sexes.....	28	41	51	69	109	169	190	179	92	33	11	-
Male.....	11	25	25	41	62	86	88	70	24	7	-	-
Female.....	17	16	26	28	47	83	102	109	68	26	11	-
Black, both sexes.....	13	25	33	35	32	43	45	47	30	10	8	-
Male.....	6	12	24	18	20	17	19	17	11	3	2	-
Female.....	7	13	9	17	12	26	26	30	19	7	6	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	1	3	6	3	4	-	1	-	-	-
Male.....	-	1	-	1	4	2	2	-	1	-	-	-
Female.....	1	-	1	2	2	1	2	-	-	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Delaware

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Florida

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Hawaii

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Illinois

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Maine

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.

Maryland

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Massachusetts

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts (All races, White, Black, American Indian, Asian or Pacific Islander).

Michigan

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan (All races, White, Black, American Indian, Asian or Pacific Islander).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Missouri

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

New Hampshire

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

North Carolina

Table with 12 columns for age groups and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 12 columns for age groups and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Ohio

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Oregon

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Texas

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Vermont

Table with 12 columns for age groups and 12 rows for race and sex categories (All races, White, Black, American Indian, Asian or Pacific Islander) for Vermont.

Virginia

Table with 12 columns for age groups and 12 rows for race and sex categories (All races, White, Black, American Indian, Asian or Pacific Islander) for Virginia.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Plague (A20)

Washington

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Plague (A20)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Plague (A20)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Plague (A20)

Hawaii

Table with 13 columns (Age groups) and 13 rows (Race categories) for Hawaii. All cells contain dashes (-).

Idaho

Table with 13 columns (Age groups) and 13 rows (Race categories) for Idaho. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Plague (A20)

Illinois

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Illinois. All cells contain dashes.

Indiana

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Indiana. All cells contain dashes.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Plague (A20)

Iowa

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Iowa. All cells contain dashes.

Kansas

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Kansas. All cells contain dashes.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Plague (A20)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Plague (A20)

Massachusetts

Table with 13 columns (Age groups) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Age groups) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Plague (A20)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not years years years years years years years years years and over stated

Plague (A20)

Nebraska

Table with 13 columns for age groups and 1 row for Nebraska. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns for age groups and 1 row for Nevada. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Plague (A20)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Plague (A20)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Plague (A20)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Plague (A20)

Texas

Table with 13 columns (Age groups) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Age groups) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Illinois

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Iowa

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Massachusetts

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Minnesota

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

South Dakota

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Vermont

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Brucellosis (A23)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Brucellosis (A23)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Brucellosis (A23)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Brucellosis (A23)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Brucellosis (A23)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Brucellosis (A23)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Brucellosis (A23)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Delaware

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Illinois

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Iowa

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Kentucky

Table with 12 columns for age groups and rows for 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with 'Male' and 'Female' sub-rows.

Louisiana

Table with 12 columns for age groups and rows for 'All races, both sexes', 'White, both sexes', 'Black, both sexes', 'American Indian 1/, both sexes', and 'Asian or Pacific Islander 2/, both sexes', each with 'Male' and 'Female' sub-rows.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Maine

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Massachusetts

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Minnesota

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Missouri

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

New Mexico

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Ohio

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Rhode Island

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

South Dakota

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Vermont

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Tetanus (A33-A35)

Wisconsin

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Tetanus (A33-A35)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Tetanus (A33-A35)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years Age not years years years years years years years years years and over stated

Tetanus (A33-A35)

Maine

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Maryland

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Tetanus (A33-A35)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Diphtheria (A36)

United States

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for United States: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Alabama

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alabama: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Alaska

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Arkansas

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Delaware

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Florida

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Hawaii

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Illinois

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Iowa

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Kentucky

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Massachusetts

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Missouri

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

New Hampshire

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

North Carolina

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Ohio

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

South Dakota

Table with 12 columns for age groups and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 12 columns for age groups and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diphtheria (A36)

Wisconsin

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diphtheria (A36)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diphtheria (A36)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Diphtheria (A36)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diphtheria (A36)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Diphtheria (A36)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Diphtheria (A36)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Whooping cough (A37)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Whooping cough (A37)

Nebraska

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Whooping cough (A37)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Whooping cough (A37)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Whooping cough (A37)

South Dakota

Table with 12 columns for age groups and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 12 columns for age groups and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Whooping cough (A37)

Washington

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Whooping cough (A37)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Whooping cough (A37)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Whooping cough (A37)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Whooping cough (A37)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Alaska

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows for Arkansas (All races, White, Black, American Indian, Asian or Pacific Islander).

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows for California (All races, White, Black, American Indian, Asian or Pacific Islander).



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Colorado

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Florida

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Illinois

Table with 12 columns for age groups and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Indiana

Table with 12 columns for age groups and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Massachusetts

Table with 12 columns for age groups and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Michigan

Table with 12 columns for age groups and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Missouri

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Scarlet fever and erysipelas (A38,A46)

New Hampshire

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

New Mexico

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Rhode Island

Table with 12 columns for age groups and 12 rows for race/sex categories (All races, White, Black, American Indian, Asian or Pacific Islander) for Rhode Island.

South Carolina

Table with 12 columns for age groups and 12 rows for race/sex categories (All races, White, Black, American Indian, Asian or Pacific Islander) for South Carolina.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

South Dakota

Table with 12 columns for age groups and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 12 columns for age groups and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Vermont

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Scarlet fever and erysipelas (A38,A46)

Wisconsin

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Scarlet fever and erysipelas (A38,A46)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Scarlet fever and erysipelas (A38,A46)

Maine

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Maine. All cells contain dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Maryland. All cells contain dashes (-).



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Scarlet fever and erysipelas (A38,A46)

Minnesota

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Minnesota. All cells contain dashes (-).

Mississippi

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Mississippi. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54	55-59	60-64	65-69	70-74	75-79	80-84	85-89	90-94	95-99	100 years	Age not
	years	years	years	years	years	years	years	years	years	years	and over	stated

Scarlet fever and erysipelas (A38,A46)

Missouri

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Scarlet fever and erysipelas (A38,A46)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Scarlet fever and erysipelas (A38,A46)

Ohio

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Ohio. All cells contain dashes (-).

Oklahoma

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Oklahoma. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Scarlet fever and erysipelas (A38,A46)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Scarlet fever and erysipelas (A38,A46)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Meningococcal infection (A39)

Alaska

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Meningococcal infection (A39)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Meningococcal infection (A39)

Massachusetts

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Meningococcal infection (A39)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Meningococcal infection (A39)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Meningococcal infection (A39)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Meningococcal infection (A39)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Meningococcal infection (A39)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Meningococcal infection (A39)

Colorado

Table with 13 columns (Age groups) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Age groups) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Meningococcal infection (A39)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Meningococcal infection (A39)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Meningococcal infection (A39)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Meningococcal infection (A39)

Texas

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Utah

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Meningococcal infection (A39)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Meningococcal infection (A39)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Arkansas												
All races, both sexes.....	461	3	2	-	1	2	1	-	4	5	6	19
Male.....	194	1	2	-	1	-	-	-	2	3	2	10
Female.....	267	2	-	-	-	2	1	-	2	2	4	9
White, both sexes.....	374	1	1	-	1	1	1	-	4	4	6	14
Male.....	157	1	1	-	1	-	-	-	2	2	2	8
Female.....	217	-	-	-	-	1	1	-	2	2	4	6
Black, both sexes.....	87	2	1	-	-	1	-	-	-	1	-	5
Male.....	37	-	1	-	-	-	-	-	-	1	-	2
Female.....	50	2	-	-	-	1	-	-	-	-	-	3
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	1,073	14	5	2	1	3	4	5	9	11	29	33
Male.....	509	6	1	1	1	1	1	1	2	8	20	12
Female.....	564	8	4	1	-	2	3	4	7	3	9	21
White, both sexes.....	876	10	3	2	1	3	2	5	6	11	19	20
Male.....	403	6	1	1	1	1	1	1	1	8	15	9
Female.....	473	4	2	1	-	2	1	4	5	3	4	11
Black, both sexes.....	100	4	1	-	-	-	1	-	3	-	5	9
Male.....	45	-	-	-	-	-	-	-	1	-	1	1
Female.....	55	4	1	-	-	-	1	-	2	-	4	8
American Indian 1/, both sexes.....	8	-	-	-	-	-	-	-	-	-	2	-
Male.....	7	-	-	-	-	-	-	-	-	-	2	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	89	-	1	-	-	-	1	-	-	-	3	4
Male.....	54	-	-	-	-	-	-	-	-	-	2	2
Female.....	35	-	1	-	-	-	1	-	-	-	1	2















































## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Septicemia (A40-A41)												
Vermont												
All races, both sexes.....	30	-	-	-	-	-	-	-	-	-	-	1
Male.....	10	-	-	-	-	-	-	-	-	-	-	-
Female.....	20	-	-	-	-	-	-	-	-	-	-	1
White, both sexes.....	29	-	-	-	-	-	-	-	-	-	-	1
Male.....	10	-	-	-	-	-	-	-	-	-	-	-
Female.....	19	-	-	-	-	-	-	-	-	-	-	1
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes.....	1,363	13	3	-	1	-	2	9	9	11	20	30
Male.....	632	4	3	-	1	-	-	7	6	5	9	19
Female.....	731	9	-	-	-	-	2	2	3	6	11	11
White, both sexes.....	988	7	3	-	1	-	1	6	3	7	7	20
Male.....	457	3	3	-	1	-	-	5	3	2	4	14
Female.....	531	4	-	-	-	-	1	1	-	5	3	6
Black, both sexes.....	351	4	-	-	-	-	1	2	3	4	13	10
Male.....	161	-	-	-	-	-	-	1	1	3	5	5
Female.....	190	4	-	-	-	-	1	1	2	1	8	5
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	24	2	-	-	-	-	-	1	3	-	-	-
Male.....	14	1	-	-	-	-	-	1	2	-	-	-
Female.....	10	1	-	-	-	-	-	-	1	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

United States

All races, both sexes.....	1,429	1,958	2,273	2,796	3,549	4,629	5,774	5,335	3,059	1,033	187	2
Male.....	765	1,051	1,152	1,442	1,814	2,200	2,568	2,064	1,021	253	26	1
Female.....	664	907	1,121	1,354	1,735	2,429	3,206	3,271	2,038	780	161	1
White, both sexes.....	965	1,400	1,720	2,113	2,739	3,790	4,894	4,615	2,636	882	147	1
Male.....	512	748	868	1,089	1,405	1,833	2,235	1,801	880	206	19	1
Female.....	453	652	852	1,024	1,334	1,957	2,659	2,814	1,756	676	128	-
Black, both sexes.....	419	507	497	620	700	741	778	630	373	140	36	-
Male.....	228	273	255	314	345	319	290	227	115	41	7	-
Female.....	191	234	242	306	355	422	488	403	258	99	29	-
American Indian 1/, both sexes.....	26	22	22	18	31	20	21	13	8	1	2	-
Male.....	13	10	9	9	13	7	9	2	4	-	-	-
Female.....	13	12	13	9	18	13	12	11	4	1	2	-
Asian or Pacific Islander 2/, both sexes.....	19	29	34	45	79	78	81	77	42	10	2	1
Male.....	12	20	20	30	51	41	34	34	22	6	-	-
Female.....	7	9	14	15	28	37	47	43	20	4	2	1

Alabama

All races, both sexes.....	45	46	58	71	91	109	109	95	49	22	6	-
Male.....	19	27	25	39	40	43	39	31	21	5	1	-
Female.....	26	19	33	32	51	66	70	64	28	17	5	-
White, both sexes.....	31	26	43	47	56	86	87	77	34	14	1	-
Male.....	14	15	17	25	22	32	35	27	14	2	-	-
Female.....	17	11	26	22	34	54	52	50	20	12	1	-
Black, both sexes.....	14	20	15	24	35	23	22	17	15	8	5	-
Male.....	5	12	8	14	18	11	4	4	7	3	1	-
Female.....	9	8	7	10	17	12	18	13	8	5	4	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

Alaska

All races, both sexes.....	1	6	2	2	2	4	4	4	4	-	-	-
Male.....	1	2	-	1	-	3	2	2	2	-	-	-
Female.....	-	4	2	1	2	1	2	2	2	-	-	-
White, both sexes.....	1	2	1	2	-	2	1	3	3	-	-	-
Male.....	1	-	-	1	-	2	1	2	1	-	-	-
Female.....	-	2	1	1	-	-	-	1	2	-	-	-
Black, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	4	1	-	2	1	3	1	-	-	-	-
Male.....	-	2	-	-	-	1	1	-	-	-	-	-
Female.....	-	2	1	-	2	-	2	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	29	32	48	47	69	99	81	61	31	6	1	1
Male.....	16	15	25	25	34	43	37	23	12	2	-	1
Female.....	13	17	23	22	35	56	44	38	19	4	1	-
White, both sexes.....	18	29	41	43	57	88	77	58	28	6	1	1
Male.....	10	14	22	21	29	40	36	23	11	2	-	1
Female.....	8	15	19	22	28	48	41	35	17	4	1	-
Black, both sexes.....	3	2	3	1	3	5	1	1	1	-	-	-
Male.....	2	1	2	1	1	2	-	-	1	-	-	-
Female.....	1	1	1	-	2	3	1	1	-	-	-	-
American Indian 1/, both sexes.....	8	-	4	2	5	5	1	2	1	-	-	-
Male.....	4	-	1	2	2	1	-	-	-	-	-	-
Female.....	4	-	3	-	3	4	1	2	1	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	1	-	1	4	1	2	-	1	-	-	-
Male.....	-	-	-	1	2	-	1	-	-	-	-	-
Female.....	-	1	-	-	2	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

Arkansas

All races, both sexes.....	11	30	37	33	47	56	88	65	37	12	2	-
Male.....	5	12	16	16	25	24	41	18	13	3	-	-
Female.....	6	18	21	17	22	32	47	47	24	9	2	-
White, both sexes.....	6	24	29	23	39	48	71	56	32	11	2	-
Male.....	3	9	12	11	20	20	34	17	11	3	-	-
Female.....	3	15	17	12	19	28	37	39	21	8	2	-
Black, both sexes.....	5	6	8	10	8	8	17	9	5	1	-	-
Male.....	2	3	4	5	5	4	7	1	2	-	-	-
Female.....	3	3	4	5	3	4	10	8	3	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	40	77	70	84	110	115	175	159	93	28	6	-
Male.....	21	43	30	44	69	55	81	64	35	13	-	-
Female.....	19	34	40	40	41	60	94	95	58	15	6	-
White, both sexes.....	34	62	58	72	84	91	151	131	80	25	6	-
Male.....	19	32	22	36	46	41	71	52	28	11	-	-
Female.....	15	30	36	36	38	50	80	79	52	14	6	-
Black, both sexes.....	5	10	5	10	13	11	9	7	6	1	-	-
Male.....	2	6	2	6	11	4	5	3	2	1	-	-
Female.....	3	4	3	4	2	7	4	4	4	-	-	-
American Indian 1/, both sexes.....	-	1	1	-	2	1	-	1	-	-	-	-
Male.....	-	1	1	-	2	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	4	6	2	11	12	15	20	7	2	-	-
Male.....	-	4	5	2	10	9	5	9	5	1	-	-
Female.....	1	-	1	-	1	3	10	11	2	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Colorado												
All races, both sexes.....	11	21	17	24	34	41	48	39	33	4	1	-
Male.....	10	11	6	10	21	21	22	16	13	1	1	-
Female.....	1	10	11	14	13	20	26	23	20	3	-	-
White, both sexes.....	10	20	16	21	31	38	48	37	32	4	1	-
Male.....	9	10	5	9	18	21	22	14	12	1	1	-
Female.....	1	10	11	12	13	17	26	23	20	3	-	-
Black, both sexes.....	1	1	1	2	2	2	-	1	1	-	-	-
Male.....	1	1	1	-	2	-	-	1	1	-	-	-
Female.....	-	-	-	2	-	2	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	1	-	1	-	-	-	-
Male.....	-	-	-	1	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
Connecticut												
All races, both sexes.....	22	20	34	41	48	65	129	131	62	25	4	-
Male.....	9	12	20	17	25	39	59	53	19	5	-	-
Female.....	13	8	14	24	23	26	70	78	43	20	4	-
White, both sexes.....	16	16	29	31	41	62	118	126	60	25	4	-
Male.....	5	11	17	12	22	38	54	51	17	5	-	-
Female.....	11	5	12	19	19	24	64	75	43	20	4	-
Black, both sexes.....	5	4	5	10	6	2	11	3	1	-	-	-
Male.....	3	1	3	5	2	1	5	1	1	-	-	-
Female.....	2	3	2	5	4	1	6	2	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	1	-	2	1	-	-	-
Male.....	1	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

Florida

All races, both sexes.....	81	107	95	136	189	287	305	284	162	54	10	-
Male.....	38	60	48	72	102	153	150	119	57	18	1	-
Female.....	43	47	47	64	87	134	155	165	105	36	9	-
White, both sexes.....	54	83	76	109	152	243	267	260	150	48	9	-
Male.....	24	47	42	56	84	133	132	114	51	16	1	-
Female.....	30	36	34	53	68	110	135	146	99	32	8	-
Black, both sexes.....	27	24	19	25	36	39	36	24	12	6	1	-
Male.....	14	13	6	14	18	19	17	5	6	2	-	-
Female.....	13	11	13	11	18	20	19	19	6	4	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	1	5	2	-	-	-	-	-
Male.....	-	-	-	2	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	1	4	1	-	-	-	-	-

Georgia

All races, both sexes.....	77	112	120	125	150	187	189	185	93	41	7	-
Male.....	45	62	69	65	73	90	84	55	22	9	-	-
Female.....	32	50	51	60	77	97	105	130	71	32	7	-
White, both sexes.....	38	58	63	78	88	128	132	135	64	25	5	-
Male.....	25	28	37	40	41	63	61	39	18	5	-	-
Female.....	13	30	26	38	47	65	71	96	46	20	5	-
Black, both sexes.....	39	51	53	46	60	56	57	49	29	16	2	-
Male.....	20	31	29	24	30	25	23	16	4	4	-	-
Female.....	19	20	24	22	30	31	34	33	25	12	2	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	4	1	2	3	-	1	-	-	-	-
Male.....	-	2	3	1	2	2	-	-	-	-	-	-
Female.....	-	-	1	-	-	1	-	1	-	-	-	-











## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Maine												
All races, both sexes.....	4	7	13	11	8	16	21	27	12	6	-	-
Male.....	2	4	5	7	4	7	12	12	3	-	-	-
Female.....	2	3	8	4	4	9	9	15	9	6	-	-
White, both sexes.....	3	7	13	11	8	16	21	27	11	6	-	-
Male.....	1	4	5	7	4	7	12	12	2	-	-	-
Female.....	2	3	8	4	4	9	9	15	9	6	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Maryland												
All races, both sexes.....	39	39	53	88	84	137	140	166	83	25	3	-
Male.....	15	25	26	52	42	67	65	63	28	6	-	-
Female.....	24	14	27	36	42	70	75	103	55	19	3	-
White, both sexes.....	16	14	36	43	46	93	104	130	60	20	2	-
Male.....	5	9	19	25	24	46	51	52	23	6	-	-
Female.....	11	5	17	18	22	47	53	78	37	14	2	-
Black, both sexes.....	22	23	16	43	35	42	35	35	22	5	1	-
Male.....	10	14	6	25	15	20	13	11	5	-	-	-
Female.....	12	9	10	18	20	22	22	24	17	5	1	-
American Indian 1/, both sexes.....	-	-	-	1	1	-	-	-	-	-	-	-
Male.....	-	-	-	1	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	2	1	1	2	2	1	1	1	-	-	-
Male.....	-	2	1	1	2	1	1	-	-	-	-	-
Female.....	1	-	-	-	-	1	-	1	1	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

Nebraska

All races, both sexes.....	6	3	11	14	13	16	21	22	15	8	-	-
Male.....	5	2	8	6	3	6	14	10	4	3	-	-
Female.....	1	1	3	8	10	10	7	12	11	5	-	-
White, both sexes.....	5	3	10	14	10	14	19	20	14	8	-	-
Male.....	4	2	7	6	2	5	13	9	4	3	-	-
Female.....	1	1	3	8	8	9	6	11	10	5	-	-
Black, both sexes.....	1	-	-	-	3	1	2	-	1	-	-	-
Male.....	1	-	-	-	1	1	1	-	-	-	-	-
Female.....	-	-	-	-	2	-	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	1	-	-	1	-	2	-	-	-	-
Male.....	-	-	1	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	23	24	32	46	66	56	67	34	25	4	1	-
Male.....	14	12	16	25	33	30	27	20	15	1	-	-
Female.....	9	12	16	21	33	26	40	14	10	3	1	-
White, both sexes.....	18	19	28	37	56	47	62	27	21	4	1	-
Male.....	10	10	13	20	30	27	25	18	12	1	-	-
Female.....	8	9	15	17	26	20	37	9	9	3	1	-
Black, both sexes.....	5	3	2	5	2	8	2	3	1	-	-	-
Male.....	4	1	2	3	1	2	2	1	1	-	-	-
Female.....	1	2	-	2	1	6	-	2	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	1	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	2	3	8	1	3	4	2	-	-	-
Male.....	-	1	1	2	2	1	-	1	1	-	-	-
Female.....	-	1	1	1	6	-	3	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

New Hampshire

All races, both sexes.....	6	9	11	9	9	8	17	22	11	4	1	-
Male.....	3	6	9	7	6	4	7	6	5	1	1	-
Female.....	3	3	2	2	3	4	10	16	6	3	-	-
White, both sexes.....	6	9	10	9	9	8	17	22	11	4	-	-
Male.....	3	6	8	7	6	4	7	6	5	1	-	-
Female.....	3	3	2	2	3	4	10	16	6	3	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	1	-
Male.....	-	-	-	-	-	-	-	-	-	-	1	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	61	66	90	104	175	231	316	310	181	75	8	-
Male.....	29	38	47	52	108	115	133	109	50	14	2	-
Female.....	32	28	43	52	67	116	183	201	131	61	6	-
White, both sexes.....	38	42	66	70	128	190	270	273	165	65	8	-
Male.....	17	23	34	33	81	96	113	95	46	11	2	-
Female.....	21	19	32	37	47	94	157	178	119	54	6	-
Black, both sexes.....	22	23	21	32	40	36	41	32	16	9	-	-
Male.....	11	14	11	18	22	17	18	12	4	2	-	-
Female.....	11	9	10	14	18	19	23	20	12	7	-	-
American Indian 1/, both sexes.....	1	-	1	-	1	-	-	-	-	-	-	-
Male.....	1	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	2	2	6	5	5	5	-	1	-	-
Male.....	-	1	1	1	5	2	2	2	-	1	-	-
Female.....	-	-	1	1	1	3	3	3	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

New Mexico

All races, both sexes.....	13	12	10	9	20	17	29	20	12	3	1	-
Male.....	6	8	6	6	9	10	10	8	6	-	-	-
Female.....	7	4	4	3	11	7	19	12	6	3	1	-
White, both sexes.....	11	12	8	7	17	15	26	19	12	3	-	-
Male.....	5	8	5	5	9	10	8	7	6	-	-	-
Female.....	6	4	3	2	8	5	18	12	6	3	-	-
Black, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	2	-	2	1	3	2	2	-	-	-	1	-
Male.....	1	-	1	-	-	-	2	-	-	-	-	-
Female.....	1	-	1	1	3	2	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	78	98	124	161	221	314	437	420	229	94	18	-
Male.....	45	54	63	81	117	151	186	173	71	22	3	-
Female.....	33	44	61	80	104	163	251	247	158	72	15	-
White, both sexes.....	53	72	102	118	187	263	384	384	201	87	16	-
Male.....	30	38	51	62	103	129	165	160	61	19	3	-
Female.....	23	34	51	56	84	134	219	224	140	68	13	-
Black, both sexes.....	22	26	19	41	30	47	44	32	25	6	2	-
Male.....	13	16	9	19	11	18	16	10	9	2	-	-
Female.....	9	10	10	22	19	29	28	22	16	4	2	-
American Indian 1/, both sexes.....	1	-	1	2	1	1	1	-	-	-	-	-
Male.....	-	-	1	-	-	1	1	-	-	-	-	-
Female.....	1	-	-	2	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	2	-	3	3	8	4	3	1	-	-
Male.....	2	-	2	-	3	3	4	3	1	1	-	-
Female.....	-	-	-	-	-	-	4	1	2	-	-	-





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
Ohio												
All races, both sexes.....	53	77	85	107	120	171	186	169	118	26	6	-
Male.....	32	48	40	63	50	81	90	56	38	5	2	-
Female.....	21	29	45	44	70	90	96	113	80	21	4	-
White, both sexes.....	39	59	75	94	102	141	155	150	103	22	5	-
Male.....	26	37	38	54	42	66	78	49	32	4	2	-
Female.....	13	22	37	40	60	75	77	101	71	18	3	-
Black, both sexes.....	14	18	10	13	17	29	30	19	15	4	1	-
Male.....	6	11	2	9	8	14	11	7	6	1	-	-
Female.....	8	7	8	4	9	15	19	12	9	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Oklahoma												
All races, both sexes.....	28	22	30	37	50	42	77	62	38	9	1	-
Male.....	13	9	19	17	17	18	31	30	15	1	-	-
Female.....	15	13	11	20	33	24	46	32	23	8	1	-
White, both sexes.....	16	16	23	29	40	34	66	54	34	8	1	-
Male.....	8	9	16	12	13	14	27	25	14	-	-	-
Female.....	8	7	7	17	27	20	39	29	20	8	1	-
Black, both sexes.....	7	1	4	4	5	5	5	4	3	1	-	-
Male.....	4	-	3	3	2	2	2	4	1	1	-	-
Female.....	3	1	1	1	3	3	3	-	2	-	-	-
American Indian 1/, both sexes.....	5	5	3	4	5	3	4	4	-	-	-	-
Male.....	1	-	-	2	2	2	2	1	-	-	-	-
Female.....	4	5	3	2	3	1	2	3	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	2	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

Oregon

All races, both sexes.....	15	11	9	24	23	22	48	35	14	7	-	-
Male.....	8	8	2	11	14	6	23	14	5	1	-	-
Female.....	7	3	7	13	9	16	25	21	9	6	-	-
White, both sexes.....	14	10	9	24	22	22	48	32	14	7	-	-
Male.....	7	7	2	11	13	6	23	14	5	1	-	-
Female.....	7	3	7	13	9	16	25	18	9	6	-	-
Black, both sexes.....	-	1	-	-	-	-	-	2	-	-	-	-
Male.....	-	1	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	2	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	1	-	-	-	-	-	-	-
Male.....	1	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Pennsylvania

All races, both sexes.....	86	119	162	206	240	362	424	428	233	78	8	-
Male.....	51	68	87	95	127	171	192	181	84	15	1	-
Female.....	35	51	75	111	113	191	232	247	149	63	7	-
White, both sexes.....	60	91	129	165	192	310	382	380	217	73	7	-
Male.....	34	53	71	75	104	147	173	157	78	13	1	-
Female.....	26	38	58	90	88	163	209	223	139	60	6	-
Black, both sexes.....	25	28	32	38	46	48	40	44	16	5	1	-
Male.....	17	15	15	18	22	22	18	22	6	2	-	-
Female.....	8	13	17	20	24	26	22	22	10	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	1	3	2	4	2	4	-	-	-	-
Male.....	-	-	1	2	1	2	1	2	-	-	-	-
Female.....	1	-	-	1	1	2	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

Rhode Island

All races, both sexes.....	7	6	9	12	13	13	33	29	16	5	-	-
Male.....	6	4	5	6	7	6	15	14	3	1	-	-
Female.....	1	2	4	6	6	7	18	15	13	4	-	-
White, both sexes.....	6	6	9	11	12	12	31	28	15	5	-	-
Male.....	5	4	5	5	7	6	15	13	2	1	-	-
Female.....	1	2	4	6	5	6	16	15	13	4	-	-
Black, both sexes.....	-	-	-	-	1	-	2	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	1	-	-	-
Female.....	-	-	-	-	1	-	2	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	1	-	-	1	-	1	-	-	-	-	-	-
Male.....	1	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

South Carolina

All races, both sexes.....	25	48	76	61	72	85	102	87	53	10	2	-
Male.....	7	25	33	34	39	41	34	43	21	3	-	-
Female.....	18	23	43	27	33	44	68	44	32	7	2	-
White, both sexes.....	13	28	45	34	43	57	72	57	46	7	1	-
Male.....	3	16	18	16	28	27	26	28	18	3	-	-
Female.....	10	12	27	18	15	30	46	29	28	4	1	-
Black, both sexes.....	12	20	30	27	29	25	29	30	7	3	1	-
Male.....	4	9	15	18	11	12	8	15	3	-	-	-
Female.....	8	11	15	9	18	13	21	15	4	3	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	1	-	-	3	1	-	-	-	-	-
Male.....	-	-	-	-	-	2	-	-	-	-	-	-
Female.....	-	-	1	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Septicemia (A40-A41)												
South Dakota												
All races, both sexes.....	1	5	5	4	4	7	9	11	6	4	-	-
Male.....	-	1	2	2	3	3	5	5	-	2	-	-
Female.....	1	4	3	2	1	4	4	6	6	2	-	-
White, both sexes.....	-	3	3	3	4	7	8	10	6	4	-	-
Male.....	-	-	1	2	3	3	5	5	-	2	-	-
Female.....	-	3	2	1	1	4	3	5	6	2	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	2	2	1	-	-	1	1	-	-	-	-
Male.....	-	1	1	-	-	-	-	-	-	-	-	-
Female.....	1	1	1	1	-	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Tennessee												
All races, both sexes.....	34	49	64	60	71	76	105	103	56	14	-	-
Male.....	20	28	32	30	37	25	47	45	19	4	-	-
Female.....	14	21	32	30	34	51	58	58	37	10	-	-
White, both sexes.....	26	34	53	42	60	62	92	88	49	10	-	-
Male.....	16	20	29	20	31	21	44	38	18	1	-	-
Female.....	10	14	24	22	29	41	48	50	31	9	-	-
Black, both sexes.....	8	15	11	18	11	14	12	14	7	4	-	-
Male.....	4	8	3	10	6	4	3	7	1	3	-	-
Female.....	4	7	8	8	5	10	9	7	6	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

Texas

All races, both sexes.....	123	191	186	239	291	368	449	409	245	76	17	-
Male.....	76	102	97	119	162	169	202	145	66	18	2	-
Female.....	47	89	89	120	129	199	247	264	179	58	15	-
White, both sexes.....	97	153	146	183	236	318	388	356	219	62	14	-
Male.....	57	82	79	95	128	149	182	127	57	13	2	-
Female.....	40	71	67	88	108	169	206	229	162	49	12	-
Black, both sexes.....	25	37	39	52	52	44	53	50	24	13	3	-
Male.....	18	19	17	22	31	19	17	17	8	4	-	-
Female.....	7	18	22	30	21	25	36	33	16	9	3	-
American Indian 1/, both sexes.....	-	-	1	1	-	2	2	-	-	-	-	-
Male.....	-	-	1	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	2	2	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	3	3	4	6	3	2	1	-	-
Male.....	1	1	-	1	3	1	3	1	1	1	-	-
Female.....	-	-	-	2	-	3	3	2	1	-	-	-

Utah

All races, both sexes.....	5	8	8	7	9	11	19	19	10	2	-	-
Male.....	2	5	3	5	3	3	9	9	5	1	-	-
Female.....	3	3	5	2	6	8	10	10	5	1	-	-
White, both sexes.....	5	6	8	5	8	11	18	19	8	2	-	-
Male.....	2	3	3	3	3	3	9	9	4	1	-	-
Female.....	3	3	5	2	5	8	9	10	4	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	-	1	1	-	-	-	1	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	1	-	-	1	-	1	-	-	-
Male.....	-	1	-	1	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Septicemia (A40-A41)

Vermont

All races, both sexes.....	3	2	2	3	3	2	3	7	2	1	1	-
Male.....	1	1	1	2	2	1	1	1	-	-	-	-
Female.....	2	1	1	1	1	1	2	6	2	1	1	-
White, both sexes.....	2	2	2	3	3	2	3	7	2	1	1	-
Male.....	1	1	1	2	2	1	1	1	-	-	-	-
Female.....	1	1	1	1	1	1	2	6	2	1	1	-
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	42	64	81	105	142	200	232	222	118	40	19	-
Male.....	17	37	47	58	83	98	105	87	35	9	2	-
Female.....	25	27	34	47	59	102	127	135	83	31	17	-
White, both sexes.....	28	40	49	69	104	156	183	175	88	30	11	-
Male.....	11	24	24	41	59	80	84	70	23	6	-	-
Female.....	17	16	25	28	45	76	99	105	65	24	11	-
Black, both sexes.....	13	23	31	33	32	43	45	47	29	10	8	-
Male.....	6	12	23	16	20	17	19	17	11	3	2	-
Female.....	7	11	8	17	12	26	26	30	18	7	6	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	1	3	6	1	4	-	1	-	-	-
Male.....	-	1	-	1	4	1	2	-	1	-	-	-
Female.....	1	-	1	2	2	-	2	-	-	-	-	-























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

Illinois

All races, both sexes.....	82	-	1	-	1	-	-	1	-	-	-	1
Male.....	35	-	1	-	-	-	-	-	-	-	-	-
Female.....	47	-	-	-	1	-	-	1	-	-	-	1
White, both sexes.....	66	-	1	-	1	-	-	1	-	-	-	-
Male.....	27	-	1	-	-	-	-	-	-	-	-	-
Female.....	39	-	-	-	1	-	-	1	-	-	-	-
Black, both sexes.....	15	-	-	-	-	-	-	-	-	-	-	1
Male.....	8	-	-	-	-	-	-	-	-	-	-	-
Female.....	7	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-

Indiana

All races, both sexes.....	60	-	-	-	-	-	1	-	-	2	3	2
Male.....	28	-	-	-	-	-	1	-	-	-	2	-
Female.....	32	-	-	-	-	-	-	-	-	2	1	2
White, both sexes.....	52	-	-	-	-	-	1	-	-	1	3	1
Male.....	25	-	-	-	-	-	1	-	-	-	2	-
Female.....	27	-	-	-	-	-	-	-	-	1	1	1
Black, both sexes.....	7	-	-	-	-	-	-	-	-	-	-	1
Male.....	3	-	-	-	-	-	-	-	-	-	-	-
Female.....	4	-	-	-	-	-	-	-	-	-	-	1
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	1	-	-













































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

Arkansas

All races, both sexes.....	2	1	-	2	2	1	-	3	2	-	-	-
Male.....	2	1	-	1	2	1	-	1	-	-	-	-
Female.....	-	-	-	1	-	-	-	2	2	-	-	-
White, both sexes.....	1	1	-	2	2	-	-	2	2	-	-	-
Male.....	1	1	-	1	2	-	-	1	-	-	-	-
Female.....	-	-	-	1	-	-	-	1	2	-	-	-
Black, both sexes.....	1	-	-	-	-	1	-	1	-	-	-	-
Male.....	1	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	8	5	7	6	9	7	21	20	8	8	-	-
Male.....	5	5	4	3	6	6	14	10	4	1	-	-
Female.....	3	-	3	3	3	1	7	10	4	7	-	-
White, both sexes.....	6	5	7	5	6	6	16	16	7	7	-	-
Male.....	4	5	4	3	3	5	10	8	3	1	-	-
Female.....	2	-	3	2	3	1	6	8	4	6	-	-
Black, both sexes.....	2	-	-	-	-	-	3	2	-	-	-	-
Male.....	1	-	-	-	-	-	2	1	-	-	-	-
Female.....	1	-	-	-	-	-	1	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	2	1	2	2	1	1	-	-
Male.....	-	-	-	-	2	1	2	1	1	-	-	-
Female.....	-	-	-	1	-	-	-	1	-	1	-	-















































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other zoonotic and bacterial diseases (A21-A22,A24-A32,A42-A44,A48-A49)

Vermont

All races, both sexes.....	-	-	-	-	1	1	1	2	-	-	-	-
Male.....	-	-	-	-	1	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes.....	-	-	-	-	1	1	1	2	-	-	-	-
Male.....	-	-	-	-	1	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	-	3	4	2	5	15	7	4	5	3	-	-
Male.....	-	1	2	2	3	7	4	-	1	1	-	-
Female.....	-	2	2	-	2	8	3	4	4	2	-	-
White, both sexes.....	-	1	2	-	5	13	7	4	4	3	-	-
Male.....	-	1	1	-	3	6	4	-	1	1	-	-
Female.....	-	-	1	-	2	7	3	4	3	2	-	-
Black, both sexes.....	-	2	2	2	-	-	-	-	1	-	-	-
Male.....	-	-	1	2	-	-	-	-	-	-	-	-
Female.....	-	2	1	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections with a predominately sexual mode of transmission (A50-A64)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections with a predominately sexual mode of transmission (A50-A64)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections with a predominately sexual mode of transmission (A50-A64)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections with a predominately sexual mode of transmission (A50-A64)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections with a predominately sexual mode of transmission (A50-A64)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections with a predominately sexual mode of transmission (A50-A64)

Missouri

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections with a predominately sexual mode of transmission (A50-A64)

New Hampshire

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections with a predominately sexual mode of transmission (A50-A64)

New Mexico

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections with a predominately sexual mode of transmission (A50-A64)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Delaware

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Hawaii

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Hawaii. All cells contain dashes (-).

Idaho

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Idaho. All cells contain dashes (-).



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Massachusetts

Table with 13 columns (Age groups) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Age groups) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

New Hampshire

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

South Dakota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections with a predominately sexual mode of transmission (A50-A64)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Syphilis (A50-A53)

Alaska

Table with 12 columns for age groups and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 12 columns for age groups and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Syphilis (A50-A53)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Syphilis (A50-A53)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Syphilis (A50-A53)

Delaware

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Syphilis (A50-A53)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Syphilis (A50-A53)

Minnesota

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Syphilis (A50-A53)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Syphilis (A50-A53)

New Mexico

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Syphilis (A50-A53)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Syphilis (A50-A53)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Syphilis (A50-A53)

Vermont

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Syphilis (A50-A53)

Wisconsin

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.

Wyoming

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each with Male and Female sub-rows.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Syphilis (A50-A53)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Syphilis (A50-A53)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Syphilis (A50-A53)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex
All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cardiovascular syphilis (A52.0)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cardiovascular syphilis (A52.0)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cardiovascular syphilis (A52.0)

Nebraska

Table with 12 columns for age groups and 13 rows for race/sex categories (All races, White, Black, American Indian, Asian or Pacific Islander) in Nebraska.

Nevada

Table with 12 columns for age groups and 13 rows for race/sex categories (All races, White, Black, American Indian, Asian or Pacific Islander) in Nevada.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cardiovascular syphilis (A52.0)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cardiovascular syphilis (A52.0)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Cardiovascular syphilis (A52.0)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cardiovascular syphilis (A52.0)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Cardiovascular syphilis (A52.0)

Wisconsin

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cardiovascular syphilis (A52.0)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cardiovascular syphilis (A52.0)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cardiovascular syphilis (A52.0)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cardiovascular syphilis (A52.0)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cardiovascular syphilis (A52.0)

Ohio

Table with 13 columns (Age groups) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Age groups) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Cardiovascular syphilis (A52.0)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Cardiovascular syphilis (A52.0)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Cardiovascular syphilis (A52.0)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Alaska

Table with 12 columns for age groups and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 12 columns for age groups and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Colorado

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Delaware

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Neurosyphilis (A52.1-A52.3)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Iowa

Table with 12 columns for age groups and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 12 columns for age groups and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Massachusetts

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Neurosyphilis (A52.1-A52.3)

Missouri

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

New Hampshire

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

New Mexico

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Neurosyphilis (A52.1-A52.3)

Wisconsin

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Neurosyphilis (A52.1-A52.3)

Colorado

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Colorado. All cells contain dashes (-).

Connecticut

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Connecticut. All cells contain dashes (-).









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Neurosyphilis (A52.1-A52.3)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Neurosyphilis (A52.1-A52.3)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Neurosyphilis (A52.1-A52.3)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Alaska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

New Mexico

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

United States

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Alabama

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified syphilis (A50-A51,A52.7-A52.9,A53)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Delaware

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Gonococcal infection (A54)

Hawaii

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex
All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Gonococcal infection (A54)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Maine

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Massachusetts

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Michigan

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Nebraska

Table with 12 columns for age groups and 13 rows for race and sex categories in Nebraska, showing death counts for Gonococcal infection (A54).

Nevada

Table with 12 columns for age groups and 13 rows for race and sex categories in Nevada, showing death counts for Gonococcal infection (A54).







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

North Carolina

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Ohio

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Rhode Island

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

South Dakota

Table with 12 columns for age groups and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 12 columns for age groups and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Vermont

Table with 12 columns for age groups and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns for age groups and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Gonococcal infection (A54)

Washington

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Gonococcal infection (A54)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Gonococcal infection (A54)

Ohio

Table with 13 columns (Age groups) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Age groups) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Gonococcal infection (A54)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Gonococcal infection (A54)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other infections with a predominately sexual mode of transmission (A55-A64)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other infections with a predominately sexual mode of transmission (A55-A64)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other infections with a predominately sexual mode of transmission (A55-A64)

Wisconsin

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections with a predominately sexual mode of transmission (A55-A64)

Colorado

Table with 13 columns (Age groups) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Age groups) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections with a predominately sexual mode of transmission (A55-A64)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections with a predominately sexual mode of transmission (A55-A64)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections with a predominately sexual mode of transmission (A55-A64)

Massachusetts

Table with 13 columns (Age groups) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Age groups) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections with a predominately sexual mode of transmission (A55-A64)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections with a predominately sexual mode of transmission (A55-A64)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

Delaware

Table with 13 columns representing age groups and 13 rows representing race and sex categories for Delaware. All cells contain dashes (-).

Dist. of Columbia

Table with 13 columns representing age groups and 13 rows representing race and sex categories for District of Columbia. All cells contain dashes (-).

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

Arkansas

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Arkansas. All values are dashes (-).

California

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for California. Values are dashes (-) or 1.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

Colorado

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

Kentucky

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

Massachusetts

Table with 13 columns (Age groups) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Age groups) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other infections caused by spirochetes, chlamydia or rickettsia (A65-A79)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Lyme disease (A69.2)

United States

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for United States: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Alabama

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alabama: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

Alaska

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

Delaware

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

Illinois

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

Massachusetts

Table with 12 columns for age groups and rows for race and sex categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 12 columns for age groups and rows for race and sex categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

Minnesota

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

New Mexico

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

South Dakota

Table with 12 columns for age groups and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 12 columns for age groups and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

Vermont

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Lyme disease (A69.2)

Wisconsin

Table with 12 columns for age groups and rows for All races, White, Black, American Indian 1/, and Asian or Pacific Islander 2/ for both sexes, Male, and Female.

Wyoming

Table with 12 columns for age groups and rows for All races, White, Black, American Indian 1/, and Asian or Pacific Islander 2/ for both sexes, Male, and Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Lyme disease (A69.2)

United States

Table with 13 columns (Age groups) and rows for United States: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Alabama

Table with 13 columns (Age groups) and rows for Alabama: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Lyme disease (A69.2)

Colorado

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Colorado. All cells contain dashes (-).

Connecticut

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Connecticut. All cells contain dashes (-).



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Lyme disease (A69.2)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Lyme disease (A69.2)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Typhus fever (A75)

United States

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for United States: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Alabama

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alabama: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Alaska

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Colorado

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Delaware

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Illinois

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Iowa

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Kentucky

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Massachusetts

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Missouri

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Nebraska

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

New Hampshire

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Rhode Island

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Vermont

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Typhus fever (A75)

Wisconsin

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 12 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Typhus fever (A75)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Typhus fever (A75)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







































































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex
All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other infections caused by spirochetes, chlamydia or rickettsia (A65-A68,A69.0-A69.1,A69.8-A69.9,A70-A74,A77-A79)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





























































