

Person with disabilities experiencing problems accessing  
NCHS birth and death tables should contact  
[nchsed@cdc.gov](mailto:nchsed@cdc.gov), or call 301-458-4688





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XIV. Diseases of the genitourinary system (N00-N98)

Arkansas

All races, both sexes.....	869	2	-	-	1	-	2	3	6	4	16	23
Male.....	401	-	-	-	-	-	1	2	3	4	10	10
Female.....	468	2	-	-	1	-	1	1	3	-	6	13
White, both sexes.....	703	1	-	-	1	-	2	2	6	3	10	14
Male.....	322	-	-	-	-	-	1	2	3	3	6	6
Female.....	381	1	-	-	1	-	1	-	3	-	4	8
Black, both sexes.....	162	1	-	-	-	-	-	1	-	1	6	9
Male.....	78	-	-	-	-	-	-	-	-	1	4	4
Female.....	84	1	-	-	-	-	-	1	-	-	2	5
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	4,794	13	3	2	-	2	13	11	14	32	49	88
Male.....	2,205	7	-	-	-	1	8	7	6	19	27	39
Female.....	2,589	6	3	2	-	1	5	4	8	13	22	49
White, both sexes.....	3,821	10	1	2	-	-	9	8	8	24	36	58
Male.....	1,777	5	-	-	-	-	6	5	4	16	19	24
Female.....	2,044	5	1	2	-	-	3	3	4	8	17	34
Black, both sexes.....	495	3	1	-	-	-	2	2	5	8	11	23
Male.....	203	2	-	-	-	-	2	2	2	3	7	11
Female.....	292	1	1	-	-	-	-	-	3	5	4	12
American Indian 1/, both sexes.....	31	-	-	-	-	-	-	-	-	-	-	-
Male.....	10	-	-	-	-	-	-	-	-	-	-	-
Female.....	21	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	447	-	1	-	-	2	2	1	1	-	2	7
Male.....	215	-	-	-	-	1	-	-	-	-	1	4
Female.....	232	-	1	-	-	1	2	1	1	-	1	3























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

XIV. Diseases of the genitourinary system (N00-N98)

Alaska

All races, both sexes.....	1	4	5	4	5	13	6	7	6	2	-	-
Male.....	1	1	4	2	1	5	2	4	3	-	-	-
Female.....	-	3	1	2	4	8	4	3	3	2	-	-
White, both sexes.....	1	3	3	3	4	10	4	4	4	2	-	-
Male.....	1	1	3	1	1	4	2	2	2	-	-	-
Female.....	-	2	-	2	3	6	2	2	2	2	-	-
Black, both sexes.....	-	-	1	-	1	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	1	1	-	3	2	3	1	-	-	-
Male.....	-	-	-	1	-	1	-	2	-	-	-	-
Female.....	-	1	1	-	-	2	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Arizona

All races, both sexes.....	30	21	42	63	73	114	149	128	103	40	4	-
Male.....	18	15	25	30	40	58	77	46	50	9	-	-
Female.....	12	6	17	33	33	56	72	82	53	31	4	-
White, both sexes.....	23	20	36	50	67	104	134	114	91	38	3	-
Male.....	14	14	23	25	37	51	69	42	45	9	-	-
Female.....	9	6	13	25	30	53	65	72	46	29	3	-
Black, both sexes.....	4	-	-	6	-	8	1	7	5	2	-	-
Male.....	2	-	-	3	-	7	1	3	2	-	-	-
Female.....	2	-	-	3	-	1	-	4	3	2	-	-
American Indian 1/, both sexes.....	3	1	6	6	6	2	12	7	6	-	1	-
Male.....	2	1	2	2	3	-	5	1	3	-	-	-
Female.....	1	-	4	4	3	2	7	6	3	-	1	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	2	-	1	-	-	-
Male.....	-	-	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	1	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

## XIV. Diseases of the genitourinary system (N00-N98)

## Arkansas

All races, both sexes.....	22	20	30	50	91	116	147	159	114	51	12	-
Male.....	13	16	15	31	54	56	66	66	34	17	3	-
Female.....	9	4	15	19	37	60	81	93	80	34	9	-
White, both sexes.....	13	14	17	38	70	96	129	140	95	44	8	-
Male.....	5	10	9	22	46	48	56	60	29	15	1	-
Female.....	8	4	8	16	24	48	73	80	66	29	7	-
Black, both sexes.....	9	6	13	12	19	20	17	19	18	7	4	-
Male.....	8	6	6	9	8	8	10	6	4	2	2	-
Female.....	1	-	7	3	11	12	7	13	14	5	2	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-

## California

All races, both sexes.....	145	219	243	304	408	618	842	888	611	232	57	-
Male.....	81	110	128	156	196	288	404	382	245	88	13	-
Female.....	64	109	115	148	212	330	438	506	366	144	44	-
White, both sexes.....	104	153	169	233	311	504	696	740	507	203	45	-
Male.....	58	75	88	121	148	245	347	322	209	78	7	-
Female.....	46	78	81	112	163	259	349	418	298	125	38	-
Black, both sexes.....	28	42	44	36	51	60	54	66	42	14	3	-
Male.....	17	19	22	16	21	21	16	25	13	3	1	-
Female.....	11	23	22	20	30	39	38	41	29	11	2	-
American Indian 1/, both sexes.....	1	2	8	4	3	4	2	2	4	1	-	-
Male.....	-	-	4	1	1	-	1	1	2	-	-	-
Female.....	1	2	4	3	2	4	1	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes.....	12	22	22	31	43	50	90	80	58	14	9	-
Male.....	6	16	14	18	26	22	40	34	21	7	5	-
Female.....	6	6	8	13	17	28	50	46	37	7	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

XIV. Diseases of the genitourinary system (N00-N98)

Colorado

All races, both sexes.....	13	21	28	38	64	82	118	149	102	28	8	-
Male.....	9	13	17	17	30	42	41	61	42	9	3	-
Female.....	4	8	11	21	34	40	77	88	60	19	5	-
White, both sexes.....	11	21	24	35	60	78	114	143	101	28	7	-
Male.....	7	13	14	15	27	41	40	59	42	9	3	-
Female.....	4	8	10	20	33	37	74	84	59	19	4	-
Black, both sexes.....	1	-	2	1	4	2	2	5	-	-	-	-
Male.....	1	-	1	1	3	1	-	1	-	-	-	-
Female.....	-	-	1	-	1	1	2	4	-	-	-	-
American Indian 1/, both sexes.....	1	-	2	1	-	-	1	-	-	-	1	-
Male.....	1	-	2	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	2	1	1	1	-	-	-
Male.....	-	-	-	1	-	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	2	-	-	1	-	-	-

Connecticut

All races, both sexes.....	14	28	29	37	55	88	136	172	144	55	10	-
Male.....	8	18	14	22	26	47	75	77	52	18	2	-
Female.....	6	10	15	15	29	41	61	95	92	37	8	-
White, both sexes.....	9	24	27	34	38	85	120	166	137	55	10	-
Male.....	4	14	13	20	21	44	63	76	50	18	2	-
Female.....	5	10	14	14	17	41	57	90	87	37	8	-
Black, both sexes.....	4	3	2	3	17	3	14	5	7	-	-	-
Male.....	3	3	1	2	5	3	10	1	2	-	-	-
Female.....	1	-	1	1	12	-	4	4	5	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	-	-	-	2	1	-	-	-	-
Male.....	1	1	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

## XIV. Diseases of the genitourinary system (N00-N98)

## Florida

All races, both sexes.....	94	154	181	231	344	527	693	721	482	178	30	-
Male.....	55	85	102	122	209	295	359	365	196	70	7	-
Female.....	39	69	79	109	135	232	334	356	286	108	23	-
White, both sexes.....	70	103	135	167	263	453	615	652	443	167	27	-
Male.....	38	61	81	91	161	256	334	343	183	68	7	-
Female.....	32	42	54	76	102	197	281	309	260	99	20	-
Black, both sexes.....	22	48	44	59	76	69	69	65	35	10	3	-
Male.....	15	23	20	29	45	39	22	21	10	2	-	-
Female.....	7	25	24	30	31	30	47	44	25	8	3	-
American Indian 1/, both sexes.....	-	1	1	1	1	1	2	-	1	-	-	-
Male.....	-	-	1	-	1	-	-	-	-	-	-	-
Female.....	-	1	-	1	-	1	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	2	1	4	4	4	7	4	3	1	-	-
Male.....	2	1	-	2	2	-	3	1	3	-	-	-
Female.....	-	1	1	2	2	4	4	3	-	1	-	-

## Georgia

All races, both sexes.....	75	105	142	175	223	312	330	369	189	72	14	-
Male.....	32	57	77	81	104	136	143	133	70	20	3	-
Female.....	43	48	65	94	119	176	187	236	119	52	11	-
White, both sexes.....	23	54	63	92	123	204	236	267	137	49	7	-
Male.....	10	29	35	44	56	96	110	101	47	13	2	-
Female.....	13	25	28	48	67	108	126	166	90	36	5	-
Black, both sexes.....	52	51	78	81	95	103	91	99	52	23	7	-
Male.....	22	28	41	37	43	38	32	30	23	7	1	-
Female.....	30	23	37	44	52	65	59	69	29	16	6	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	5	5	3	3	-	-	-	-
Male.....	-	-	-	-	5	2	1	2	-	-	-	-
Female.....	-	-	-	2	-	3	2	1	-	-	-	-





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

## XIV. Diseases of the genitourinary system (N00-N98)

## Iowa

All races, both sexes.....	10	6	12	15	35	56	91	123	103	44	10	-
Male.....	2	4	9	5	16	34	40	52	32	14	2	-
Female.....	8	2	3	10	19	22	51	71	71	30	8	-
White, both sexes.....	10	6	10	15	33	54	86	123	99	44	10	-
Male.....	2	4	7	5	14	32	39	52	31	14	2	-
Female.....	8	2	3	10	19	22	47	71	68	30	8	-
Black, both sexes.....	-	-	2	-	1	2	5	-	3	-	-	-
Male.....	-	-	2	-	1	2	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	4	-	3	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	1	-	-	-
Male.....	-	-	-	-	1	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

## Kansas

All races, both sexes.....	13	28	26	43	54	94	132	144	108	44	5	-
Male.....	8	14	11	23	35	48	62	61	37	17	1	-
Female.....	5	14	15	20	19	46	70	83	71	27	4	-
White, both sexes.....	11	22	20	36	47	86	124	135	103	44	4	-
Male.....	6	10	10	21	31	44	58	58	36	17	1	-
Female.....	5	12	10	15	16	42	66	77	67	27	3	-
Black, both sexes.....	2	6	4	6	5	7	6	9	5	-	1	-
Male.....	2	4	1	1	3	4	3	3	1	-	-	-
Female.....	-	2	3	5	2	3	3	6	4	-	1	-
American Indian 1/, both sexes.....	-	-	2	1	2	-	2	-	-	-	-	-
Male.....	-	-	-	1	1	-	1	-	-	-	-	-
Female.....	-	-	2	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

XIV. Diseases of the genitourinary system (N00-N98)

Kentucky

All races, both sexes.....	31	50	66	80	146	182	232	233	156	45	9	-
Male.....	14	26	47	36	74	84	101	85	52	14	1	-
Female.....	17	24	19	44	72	98	131	148	104	31	8	-
White, both sexes.....	25	43	55	68	129	167	217	215	143	43	7	-
Male.....	10	23	38	26	66	78	94	79	49	14	1	-
Female.....	15	20	17	42	63	89	123	136	94	29	6	-
Black, both sexes.....	6	7	11	12	16	15	14	18	13	2	2	-
Male.....	4	3	9	10	8	6	7	6	3	-	-	-
Female.....	2	4	2	2	8	9	7	12	10	2	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-

Louisiana

All races, both sexes.....	52	70	76	110	140	190	264	236	153	53	13	-
Male.....	28	31	45	46	60	98	126	97	53	14	2	-
Female.....	24	39	31	64	80	92	138	139	100	39	11	-
White, both sexes.....	25	29	36	67	69	123	193	175	104	40	5	-
Male.....	16	14	23	29	31	69	97	72	41	8	1	-
Female.....	9	15	13	38	38	54	96	103	63	32	4	-
Black, both sexes.....	27	40	40	43	70	65	69	59	47	13	8	-
Male.....	12	16	22	17	29	28	28	23	11	6	1	-
Female.....	15	24	18	26	41	37	41	36	36	7	7	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	1	2	1	2	2	-	-	-
Male.....	-	1	-	-	-	1	-	2	1	-	-	-
Female.....	-	-	-	-	1	1	1	-	1	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

## XIV. Diseases of the genitourinary system (N00-N98)

## Maine

All races, both sexes.....	5	8	15	21	24	49	76	75	47	21	6	-
Male.....	4	8	9	9	12	19	38	30	24	9	1	-
Female.....	1	-	6	12	12	30	38	45	23	12	5	-
White, both sexes.....	5	8	15	21	24	48	75	75	45	21	6	-
Male.....	4	8	9	9	12	18	38	30	23	9	1	-
Female.....	1	-	6	12	12	30	37	45	22	12	5	-
Black, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	1	-	-	-
Male.....	-	-	-	-	-	1	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

## Maryland

All races, both sexes.....	29	48	68	80	98	132	226	230	120	49	9	-
Male.....	19	28	39	35	47	57	95	82	49	14	1	-
Female.....	10	20	29	45	51	75	131	148	71	35	8	-
White, both sexes.....	12	25	27	47	64	87	166	176	102	38	8	-
Male.....	9	13	17	21	29	38	69	67	39	11	-	-
Female.....	3	12	10	26	35	49	97	109	63	27	8	-
Black, both sexes.....	16	22	39	32	29	44	59	47	15	9	1	-
Male.....	9	14	21	14	15	19	25	14	7	2	1	-
Female.....	7	8	18	18	14	25	34	33	8	7	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	2	1	5	1	1	7	3	2	-	-
Male.....	1	1	1	-	3	-	1	1	3	1	-	-
Female.....	-	-	1	1	2	1	-	6	-	1	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

## XIV. Diseases of the genitourinary system (N00-N98)

## Massachusetts

All races, both sexes.....	24	38	61	90	138	234	358	391	291	113	20	-
Male.....	17	21	34	54	68	114	169	183	111	39	5	-
Female.....	7	17	27	36	70	120	189	208	180	74	15	-
White, both sexes.....	21	34	54	74	121	209	342	365	279	109	18	-
Male.....	14	19	30	45	61	97	163	174	106	38	5	-
Female.....	7	15	24	29	60	112	179	191	173	71	13	-
Black, both sexes.....	2	4	6	14	14	21	13	23	10	4	2	-
Male.....	2	2	3	7	6	14	6	8	5	1	-	-
Female.....	-	2	3	7	8	7	7	15	5	3	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	1	2	3	4	3	3	2	-	-	-
Male.....	1	-	1	2	1	3	-	1	-	-	-	-
Female.....	-	-	-	-	2	1	3	2	2	-	-	-

## Michigan

All races, both sexes.....	57	81	94	125	165	299	443	478	277	99	20	-
Male.....	31	41	46	63	76	128	217	204	103	26	10	-
Female.....	26	40	48	62	89	171	226	274	174	73	10	-
White, both sexes.....	33	53	64	93	137	244	385	421	251	83	16	-
Male.....	18	24	31	44	62	106	188	181	94	20	8	-
Female.....	15	29	33	49	75	138	197	240	157	63	8	-
Black, both sexes.....	23	26	29	31	28	48	54	54	25	15	3	-
Male.....	13	15	14	18	14	19	26	22	8	6	1	-
Female.....	10	11	15	13	14	29	28	32	17	9	2	-
American Indian 1/, both sexes.....	1	2	1	-	-	5	1	1	-	1	-	-
Male.....	-	2	1	-	-	2	1	1	-	-	-	-
Female.....	1	-	-	-	-	3	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	2	3	2	1	-	1	-
Male.....	-	-	-	1	-	1	2	-	1	-	1	-
Female.....	-	-	-	-	-	1	1	2	-	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

## XIV. Diseases of the genitourinary system (N00-N98)

## Minnesota

All races, both sexes.....	13	26	34	31	71	103	201	226	169	66	13	-
Male.....	9	14	17	15	50	57	101	108	64	24	3	-
Female.....	4	12	17	16	21	46	100	118	105	42	10	-
White, both sexes.....	11	20	29	28	59	91	196	225	164	65	13	-
Male.....	8	10	15	14	42	52	99	107	63	24	3	-
Female.....	3	10	14	14	17	39	97	118	101	41	10	-
Black, both sexes.....	2	4	1	1	4	7	2	1	1	1	-	-
Male.....	1	3	-	-	3	3	1	1	-	-	-	-
Female.....	1	1	1	1	1	4	1	-	1	1	-	-
American Indian 1/, both sexes.....	-	-	2	-	4	3	1	-	1	-	-	-
Male.....	-	-	1	-	3	1	-	-	-	-	-	-
Female.....	-	-	1	-	1	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	2	2	4	2	2	-	3	-	-	-
Male.....	-	1	1	1	2	1	1	-	1	-	-	-
Female.....	-	1	1	1	2	1	1	-	2	-	-	-

## Mississippi

All races, both sexes.....	38	40	58	78	98	133	161	142	98	37	10	-
Male.....	23	17	30	33	46	69	80	51	39	10	2	-
Female.....	15	23	28	45	52	64	81	91	59	27	8	-
White, both sexes.....	17	13	29	33	48	89	99	95	68	20	5	-
Male.....	10	5	14	15	26	53	51	37	27	4	-	-
Female.....	7	8	15	18	22	36	48	58	41	16	5	-
Black, both sexes.....	21	27	27	43	50	43	62	47	29	17	5	-
Male.....	13	12	14	17	20	16	29	14	12	6	2	-
Female.....	8	15	13	26	30	27	33	33	17	11	3	-
American Indian 1/, both sexes.....	-	-	2	1	-	-	-	-	-	-	-	-
Male.....	-	-	2	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	1	-	-	1	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	1	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

XIV. Diseases of the genitourinary system (N00-N98)

Nebraska

All races, both sexes.....	5	12	17	12	33	40	62	87	67	24	7	-
Male.....	2	7	12	3	21	22	28	31	23	5	1	-
Female.....	3	5	5	9	12	18	34	56	44	19	6	-
White, both sexes.....	5	9	10	9	30	37	61	87	65	24	7	-
Male.....	2	6	10	3	19	20	28	31	22	5	1	-
Female.....	3	3	-	6	11	17	33	56	43	19	6	-
Black, both sexes.....	-	3	4	3	3	2	1	-	2	-	-	-
Male.....	-	1	1	-	2	1	-	-	1	-	-	-
Female.....	-	2	3	3	1	1	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	3	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	20	23	32	44	79	81	95	84	42	13	1	-
Male.....	11	11	20	27	50	43	52	39	17	8	-	-
Female.....	9	12	12	17	29	38	43	45	25	5	1	-
White, both sexes.....	16	20	24	36	63	66	84	74	35	11	-	-
Male.....	10	9	15	23	42	36	43	34	15	7	-	-
Female.....	6	11	9	13	21	30	41	40	20	4	-	-
Black, both sexes.....	2	2	4	5	11	11	6	7	4	1	-	-
Male.....	1	1	2	2	6	4	5	4	-	-	-	-
Female.....	1	1	2	3	5	7	1	3	4	1	-	-
American Indian 1/, both sexes.....	1	-	-	2	-	1	2	-	-	-	-	-
Male.....	-	-	-	2	-	1	1	-	-	-	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	4	1	5	3	3	3	3	1	1	-
Male.....	-	1	3	-	2	2	3	1	2	1	-	-
Female.....	1	-	1	1	3	1	-	2	1	-	1	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

## XIV. Diseases of the genitourinary system (N00-N98)

## New Hampshire

All races, both sexes.....	2	2	7	8	17	27	48	56	31	12	2	-
Male.....	1	1	4	4	8	11	22	25	14	3	-	-
Female.....	1	1	3	4	9	16	26	31	17	9	2	-
White, both sexes.....	2	2	7	8	17	27	47	56	31	12	2	-
Male.....	1	1	4	4	8	11	22	25	14	3	-	-
Female.....	1	1	3	4	9	16	25	31	17	9	2	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

## New Jersey

All races, both sexes.....	52	67	101	113	159	303	415	456	341	105	11	-
Male.....	26	31	59	68	84	136	213	202	129	24	4	-
Female.....	26	36	42	45	75	167	202	254	212	81	7	-
White, both sexes.....	34	41	57	79	111	240	372	411	316	91	9	-
Male.....	16	23	34	50	60	115	194	187	120	21	3	-
Female.....	18	18	23	29	51	125	178	224	196	70	6	-
Black, both sexes.....	16	23	40	33	42	53	36	39	17	14	1	-
Male.....	8	6	23	17	19	17	14	12	4	3	-	-
Female.....	8	17	17	16	23	36	22	27	13	11	1	-
American Indian 1/, both sexes.....	-	-	1	-	1	1	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	3	3	1	5	9	7	6	8	-	1	-
Male.....	2	2	2	1	5	3	5	3	5	-	1	-
Female.....	-	1	1	-	-	6	2	3	3	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

## XIV. Diseases of the genitourinary system (N00-N98)

## New Mexico

All races, both sexes.....	8	15	16	21	33	48	77	69	34	17	6	-
Male.....	4	12	10	11	15	28	35	31	22	8	5	-
Female.....	4	3	6	10	18	20	42	38	12	9	1	-
White, both sexes.....	6	13	13	18	29	45	68	65	33	15	6	-
Male.....	2	11	8	10	14	26	32	29	21	8	5	-
Female.....	4	2	5	8	15	19	36	36	12	7	1	-
Black, both sexes.....	-	1	-	1	-	1	4	-	-	-	-	-
Male.....	-	1	-	1	-	1	2	-	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes.....	2	1	2	2	4	2	5	4	1	2	-	-
Male.....	2	-	2	-	1	1	1	2	1	-	-	-
Female.....	-	1	-	2	3	1	4	2	-	2	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-

## New York

All races, both sexes.....	67	112	141	206	263	502	694	829	522	198	55	-
Male.....	36	62	64	91	139	248	311	346	177	48	11	-
Female.....	31	50	77	115	124	254	383	483	345	150	44	-
White, both sexes.....	45	78	96	154	197	415	601	755	468	178	44	-
Male.....	23	42	43	75	109	208	273	319	161	47	6	-
Female.....	22	36	53	79	88	207	328	436	307	131	38	-
Black, both sexes.....	20	30	42	46	57	73	77	67	48	19	9	-
Male.....	11	17	21	11	25	36	30	23	14	1	4	-
Female.....	9	13	21	35	32	37	47	44	34	18	5	-
American Indian 1/, both sexes.....	2	2	-	2	2	-	1	1	-	-	-	-
Male.....	2	1	-	2	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	2	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	3	4	7	14	15	6	6	1	2	-
Male.....	-	2	-	3	5	4	8	4	2	-	1	-
Female.....	-	-	3	1	2	10	7	2	4	1	1	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

XIV. Diseases of the genitourinary system (N00-N98)

Oregon

All races, both sexes.....	14	20	23	28	50	78	129	141	104	40	8	-
Male.....	12	8	9	11	24	35	59	59	31	10	6	-
Female.....	2	12	14	17	26	43	70	82	73	30	2	-
White, both sexes.....	10	15	20	26	48	77	125	135	102	40	8	-
Male.....	9	6	8	11	24	34	57	55	31	10	6	-
Female.....	1	9	12	15	24	43	68	80	71	30	2	-
Black, both sexes.....	3	2	-	-	1	-	-	4	-	-	-	-
Male.....	2	1	-	-	-	-	-	3	-	-	-	-
Female.....	1	1	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	1	1	1	1	-	-	-	1	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	1	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	2	1	1	1	4	1	2	-	-	-
Male.....	-	1	1	-	-	1	2	1	-	-	-	-
Female.....	-	1	1	1	1	-	2	-	2	-	-	-

Pennsylvania

All races, both sexes.....	70	95	136	215	333	549	762	794	536	173	30	-
Male.....	37	60	71	116	153	273	331	329	175	43	7	-
Female.....	33	35	65	99	180	276	431	465	361	130	23	-
White, both sexes.....	52	73	106	168	284	491	690	739	497	157	28	-
Male.....	25	44	54	85	136	252	305	306	164	39	6	-
Female.....	27	29	52	83	148	239	385	433	333	118	22	-
Black, both sexes.....	18	22	28	44	49	55	68	53	39	16	2	-
Male.....	12	16	15	29	17	19	22	23	11	4	1	-
Female.....	6	6	13	15	32	36	46	30	28	12	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	3	-	2	4	2	-	-	-	-
Male.....	-	-	2	2	-	1	4	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	2	-	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

XIV. Diseases of the genitourinary system (N00-N98)

Texas

All races, both sexes.....	156	196	258	329	453	601	733	736	513	185	47	-
Male.....	84	98	138	173	223	291	331	325	187	50	8	-
Female.....	72	98	120	156	230	310	402	411	326	135	39	-
White, both sexes.....	108	145	191	246	368	510	643	631	464	166	39	-
Male.....	64	68	98	133	184	250	293	284	172	44	5	-
Female.....	44	77	93	113	184	260	350	347	292	122	34	-
Black, both sexes.....	46	43	59	76	78	78	77	93	46	19	7	-
Male.....	18	25	33	37	33	37	31	36	13	6	3	-
Female.....	28	18	26	39	45	41	46	57	33	13	4	-
American Indian 1/, both sexes.....	-	3	1	-	1	-	-	2	-	-	-	-
Male.....	-	2	1	-	1	-	-	2	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	5	7	7	6	13	13	10	3	-	1	-
Male.....	2	3	6	3	5	4	7	3	2	-	-	-
Female.....	-	2	1	4	1	9	6	7	1	-	1	-

Utah

All races, both sexes.....	4	15	13	17	41	46	60	56	38	21	3	-
Male.....	2	7	5	4	22	18	33	22	11	10	-	-
Female.....	2	8	8	13	19	28	27	34	27	11	3	-
White, both sexes.....	3	14	13	14	40	43	59	55	36	20	3	-
Male.....	1	7	5	4	22	17	32	22	11	10	-	-
Female.....	2	7	8	10	18	26	27	33	25	10	3	-
Black, both sexes.....	-	-	-	1	1	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	1	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	2	-	3	-	1	2	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	1	-	2	-	2	-	1	2	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

## XIV. Diseases of the genitourinary system (N00-N98)

## Vermont

All races, both sexes.....	-	4	2	5	8	10	12	22	14	4	-	-
Male.....	-	3	2	2	5	6	5	7	2	-	-	-
Female.....	-	1	-	3	3	4	7	15	12	4	-	-
White, both sexes.....	-	4	2	5	8	10	12	22	14	4	-	-
Male.....	-	3	2	2	5	6	5	7	2	-	-	-
Female.....	-	1	-	3	3	4	7	15	12	4	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

## Virginia

All races, both sexes.....	49	78	92	142	190	259	353	358	222	78	14	-
Male.....	27	46	40	68	92	136	154	146	86	14	1	-
Female.....	22	32	52	74	98	123	199	212	136	64	13	-
White, both sexes.....	25	38	50	90	120	182	282	277	179	64	7	-
Male.....	15	25	25	47	59	96	123	115	73	12	1	-
Female.....	10	13	25	43	61	86	159	162	106	52	6	-
Black, both sexes.....	23	40	40	49	62	73	66	75	41	14	7	-
Male.....	11	21	14	20	29	40	30	30	13	2	-	-
Female.....	12	19	26	29	33	33	36	45	28	12	7	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	2	3	8	4	5	5	2	-	-	-
Male.....	1	-	1	1	4	-	1	1	-	-	-	-
Female.....	-	-	1	2	4	4	4	4	2	-	-	-









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Diseases of urinary system (N00-N39)												
Arkansas												
All races, both sexes.....	865	2	-	-	1	-	2	3	6	4	15	23
Male.....	397	-	-	-	-	-	1	2	3	4	9	10
Female.....	468	2	-	-	1	-	1	1	3	-	6	13
White, both sexes.....	700	1	-	-	1	-	2	2	6	3	9	14
Male.....	319	-	-	-	-	-	1	2	3	3	5	6
Female.....	381	1	-	-	1	-	1	-	3	-	4	8
Black, both sexes.....	161	1	-	-	-	-	-	1	-	1	6	9
Male.....	77	-	-	-	-	-	-	-	-	1	4	4
Female.....	84	1	-	-	-	-	-	1	-	-	2	5
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	4,629	12	3	2	-	2	11	10	13	29	46	84
Male.....	2,086	7	-	-	-	1	8	7	6	18	27	37
Female.....	2,543	5	3	2	-	1	3	3	7	11	19	47
White, both sexes.....	3,677	10	1	2	-	-	9	7	8	21	34	54
Male.....	1,670	5	-	-	-	-	6	5	4	15	19	22
Female.....	2,007	5	1	2	-	-	3	2	4	6	15	32
Black, both sexes.....	484	2	1	-	-	-	2	2	5	8	10	23
Male.....	197	2	-	-	-	-	2	2	2	3	7	11
Female.....	287	-	1	-	-	-	-	-	3	5	3	12
American Indian 1/, both sexes.....	31	-	-	-	-	-	-	-	-	-	-	-
Male.....	10	-	-	-	-	-	-	-	-	-	-	-
Female.....	21	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	437	-	1	-	-	2	-	1	-	-	2	7
Male.....	209	-	-	-	-	1	-	-	-	-	1	4
Female.....	228	-	1	-	-	1	-	1	-	-	1	3

























































## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of urinary system (N00-N39)												
Alaska												
All races, both sexes.....	1	4	5	4	5	12	6	6	6	2	-	-
Male.....	1	1	4	2	1	4	2	3	3	-	-	-
Female.....	-	3	1	2	4	8	4	3	3	2	-	-
White, both sexes.....	1	3	3	3	4	9	4	4	4	2	-	-
Male.....	1	1	3	1	1	3	2	2	2	-	-	-
Female.....	-	2	-	2	3	6	2	2	2	2	-	-
Black, both sexes.....	-	-	1	-	1	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	1	1	-	3	2	2	1	-	-	-
Male.....	-	-	-	1	-	1	-	1	-	-	-	-
Female.....	-	1	1	-	-	2	2	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes.....	29	21	42	63	72	114	148	121	101	40	4	-
Male.....	18	15	25	30	39	58	76	42	48	9	-	-
Female.....	11	6	17	33	33	56	72	79	53	31	4	-
White, both sexes.....	22	20	36	50	66	104	133	109	89	38	3	-
Male.....	14	14	23	25	36	51	68	39	43	9	-	-
Female.....	8	6	13	25	30	53	65	70	46	29	3	-
Black, both sexes.....	4	-	-	6	-	8	1	7	5	2	-	-
Male.....	2	-	-	3	-	7	1	3	2	-	-	-
Female.....	2	-	-	3	-	1	-	4	3	2	-	-
American Indian 1/, both sexes.....	3	1	6	6	6	2	12	5	6	-	1	-
Male.....	2	1	2	2	3	-	5	-	3	-	-	-
Female.....	1	-	4	4	3	2	7	5	3	-	1	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	2	-	1	-	-	-
Male.....	-	-	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Diseases of urinary system (N00-N39)

Arkansas

All races, both sexes.....	22	20	30	50	91	116	146	159	114	49	12	-
Male.....	13	16	15	31	54	56	65	66	34	15	3	-
Female.....	9	4	15	19	37	60	81	93	80	34	9	-
White, both sexes.....	13	14	17	38	70	96	129	140	95	42	8	-
Male.....	5	10	9	22	46	48	56	60	29	13	1	-
Female.....	8	4	8	16	24	48	73	80	66	29	7	-
Black, both sexes.....	9	6	13	12	19	20	16	19	18	7	4	-
Male.....	8	6	6	9	8	8	9	6	4	2	2	-
Female.....	1	-	7	3	11	12	7	13	14	5	2	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-

California

All races, both sexes.....	143	209	236	288	403	606	812	854	590	224	52	-
Male.....	80	105	125	142	195	279	381	352	226	81	9	-
Female.....	63	104	111	146	208	327	431	502	364	143	43	-
White, both sexes.....	102	144	165	218	307	493	668	709	489	196	40	-
Male.....	57	71	87	107	147	237	326	295	192	72	3	-
Female.....	45	73	78	111	160	256	342	414	297	124	37	-
Black, both sexes.....	28	42	41	35	50	59	54	64	42	13	3	-
Male.....	17	19	20	16	21	20	16	23	13	2	1	-
Female.....	11	23	21	19	29	39	38	41	29	11	2	-
American Indian 1/, both sexes.....	1	2	8	4	3	4	2	2	4	1	-	-
Male.....	-	-	4	1	1	-	1	1	2	-	-	-
Female.....	1	2	4	3	2	4	1	1	2	1	-	-
Asian or Pacific Islander 2/, both sexes.....	12	21	22	31	43	50	88	79	55	14	9	-
Male.....	6	15	14	18	26	22	38	33	19	7	5	-
Female.....	6	6	8	13	17	28	50	46	36	7	4	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Diseases of urinary system (N00-N39)

Colorado

All races, both sexes.....	13	21	28	37	64	81	117	145	98	27	7	-
Male.....	9	13	17	17	30	41	41	59	38	8	2	-
Female.....	4	8	11	20	34	40	76	86	60	19	5	-
White, both sexes.....	11	21	24	34	60	77	113	139	97	27	6	-
Male.....	7	13	14	15	27	40	40	57	38	8	2	-
Female.....	4	8	10	19	33	37	73	82	59	19	4	-
Black, both sexes.....	1	-	2	1	4	2	2	5	-	-	-	-
Male.....	1	-	1	1	3	1	-	1	-	-	-	-
Female.....	-	-	1	-	1	1	2	4	-	-	-	-
American Indian 1/, both sexes.....	1	-	2	1	-	-	1	-	-	-	1	-
Male.....	1	-	2	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	1	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	2	1	1	1	-	-	-
Male.....	-	-	-	1	-	-	1	1	-	-	-	-
Female.....	-	-	-	-	-	2	-	-	1	-	-	-

Connecticut

All races, both sexes.....	12	27	28	37	54	86	134	170	141	55	10	-
Male.....	8	18	13	22	25	46	74	75	49	18	2	-
Female.....	4	9	15	15	29	40	60	95	92	37	8	-
White, both sexes.....	8	23	26	34	37	83	118	164	135	55	10	-
Male.....	4	14	12	20	20	43	62	74	48	18	2	-
Female.....	4	9	14	14	17	40	56	90	87	37	8	-
Black, both sexes.....	3	3	2	3	17	3	14	5	6	-	-	-
Male.....	3	3	1	2	5	3	10	1	1	-	-	-
Female.....	-	-	1	1	12	-	4	4	5	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	-	-	-	2	1	-	-	-	-
Male.....	1	1	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-



## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of urinary system (N00-N39)												
Florida												
All races, both sexes.....	94	153	172	231	338	516	683	709	478	174	29	-
Male.....	55	85	94	122	206	292	353	358	194	66	7	-
Female.....	39	68	78	109	132	224	330	351	284	108	22	-
White, both sexes.....	70	102	127	167	258	443	605	641	441	163	26	-
Male.....	38	61	74	91	158	253	328	336	181	64	7	-
Female.....	32	41	53	76	100	190	277	305	260	99	19	-
Black, both sexes.....	22	48	43	59	75	68	69	64	33	10	3	-
Male.....	15	23	19	29	45	39	22	21	10	2	-	-
Female.....	7	25	24	30	30	29	47	43	23	8	3	-
American Indian 1/, both sexes.....	-	1	1	1	1	1	2	-	1	-	-	-
Male.....	-	-	1	-	1	-	-	-	-	-	-	-
Female.....	-	1	-	1	-	1	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	2	1	4	4	4	7	4	3	1	-	-
Male.....	2	1	-	2	2	-	3	1	3	-	-	-
Female.....	-	1	1	2	2	4	4	3	-	1	-	-
Georgia												
All races, both sexes.....	74	103	140	174	219	307	329	360	187	70	14	-
Male.....	32	55	77	80	102	132	142	127	68	19	3	-
Female.....	42	48	63	94	117	175	187	233	119	51	11	-
White, both sexes.....	23	54	62	91	123	201	235	259	137	48	7	-
Male.....	10	29	35	43	56	94	109	96	47	13	2	-
Female.....	13	25	27	48	67	107	126	163	90	35	5	-
Black, both sexes.....	51	49	77	81	91	101	91	98	50	22	7	-
Male.....	22	26	41	37	41	36	32	29	21	6	1	-
Female.....	29	23	36	44	50	65	59	69	29	16	6	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	2	5	5	3	3	-	-	-	-
Male.....	-	-	-	-	5	2	1	2	-	-	-	-
Female.....	-	-	-	2	-	3	2	1	-	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Diseases of urinary system (N00-N39)

Iowa

All races, both sexes.....	10	5	12	14	34	55	87	114	100	44	10	-
Male.....	2	4	9	4	15	33	37	45	31	14	2	-
Female.....	8	1	3	10	19	22	50	69	69	30	8	-
White, both sexes.....	10	5	10	14	32	54	82	114	96	44	10	-
Male.....	2	4	7	4	13	32	36	45	30	14	2	-
Female.....	8	1	3	10	19	22	46	69	66	30	8	-
Black, both sexes.....	-	-	2	-	1	1	5	-	3	-	-	-
Male.....	-	-	2	-	1	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	4	-	3	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	1	-	-	-
Male.....	-	-	-	-	1	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas

All races, both sexes.....	13	27	26	43	54	92	131	142	107	43	5	-
Male.....	8	14	11	23	35	47	62	60	36	16	1	-
Female.....	5	13	15	20	19	45	69	82	71	27	4	-
White, both sexes.....	11	21	20	36	47	85	123	133	102	43	4	-
Male.....	6	10	10	21	31	44	58	57	35	16	1	-
Female.....	5	11	10	15	16	41	65	76	67	27	3	-
Black, both sexes.....	2	6	4	6	5	6	6	9	5	-	1	-
Male.....	2	4	1	1	3	3	3	3	1	-	-	-
Female.....	-	2	3	5	2	3	3	6	4	-	1	-
American Indian 1/, both sexes.....	-	-	2	1	2	-	2	-	-	-	-	-
Male.....	-	-	-	1	1	-	1	-	-	-	-	-
Female.....	-	-	2	-	1	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Diseases of urinary system (N00-N39)

Kentucky

All races, both sexes.....	31	49	65	79	146	178	231	230	153	45	9	-
Male.....	14	26	46	35	74	83	101	84	51	14	1	-
Female.....	17	23	19	44	72	95	130	146	102	31	8	-
White, both sexes.....	25	42	54	67	129	164	216	212	140	43	7	-
Male.....	10	23	37	25	66	77	94	78	48	14	1	-
Female.....	15	19	17	42	63	87	122	134	92	29	6	-
Black, both sexes.....	6	7	11	12	16	14	14	18	13	2	2	-
Male.....	4	3	9	10	8	6	7	6	3	-	-	-
Female.....	2	4	2	2	8	8	7	12	10	2	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-

Louisiana

All races, both sexes.....	51	70	76	110	139	189	262	231	153	53	13	-
Male.....	27	31	45	46	60	97	124	94	53	14	2	-
Female.....	24	39	31	64	79	92	138	137	100	39	11	-
White, both sexes.....	24	29	36	67	68	122	192	172	104	40	5	-
Male.....	15	14	23	29	31	68	96	70	41	8	1	-
Female.....	9	15	13	38	37	54	96	102	63	32	4	-
Black, both sexes.....	27	40	40	43	70	65	68	57	47	13	8	-
Male.....	12	16	22	17	29	28	27	22	11	6	1	-
Female.....	15	24	18	26	41	37	41	35	36	7	7	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	1	2	1	2	2	-	-	-
Male.....	-	1	-	-	-	1	-	2	1	-	-	-
Female.....	-	-	-	-	1	1	1	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Diseases of urinary system (N00-N39)

Maine

All races, both sexes.....	4	8	15	21	23	48	76	74	46	21	6	-
Male.....	3	8	9	9	11	19	38	30	23	9	1	-
Female.....	1	-	6	12	12	29	38	44	23	12	5	-
White, both sexes.....	4	8	15	21	23	47	75	74	44	21	6	-
Male.....	3	8	9	9	11	18	38	30	22	9	1	-
Female.....	1	-	6	12	12	29	37	44	22	12	5	-
Black, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	1	-	-	-
Male.....	-	-	-	-	-	1	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Maryland

All races, both sexes.....	28	47	67	80	96	130	223	228	117	49	9	-
Male.....	18	27	39	35	47	57	94	80	46	14	1	-
Female.....	10	20	28	45	49	73	129	148	71	35	8	-
White, both sexes.....	11	25	27	47	64	86	164	174	100	38	8	-
Male.....	8	13	17	21	29	38	69	65	37	11	-	-
Female.....	3	12	10	26	35	48	95	109	63	27	8	-
Black, both sexes.....	16	21	38	32	27	43	58	47	15	9	1	-
Male.....	9	13	21	14	15	19	24	14	7	2	1	-
Female.....	7	8	17	18	12	24	34	33	8	7	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	2	1	5	1	1	7	2	2	-	-
Male.....	1	1	1	-	3	-	1	1	2	1	-	-
Female.....	-	-	1	1	2	1	-	6	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Diseases of urinary system (N00-N39)

Massachusetts

All races, both sexes.....	24	38	59	89	136	234	350	388	286	111	20	-
Male.....	17	21	33	53	66	114	165	181	106	38	5	-
Female.....	7	17	26	36	70	120	185	207	180	73	15	-
White, both sexes.....	21	34	52	73	119	209	334	362	275	107	18	-
Male.....	14	19	29	44	59	97	159	172	102	37	5	-
Female.....	7	15	23	29	60	112	175	190	173	70	13	-
Black, both sexes.....	2	4	6	14	14	21	13	23	9	4	2	-
Male.....	2	2	3	7	6	14	6	8	4	1	-	-
Female.....	-	2	3	7	8	7	7	15	5	3	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	1	2	3	4	3	3	2	-	-	-
Male.....	1	-	1	2	1	3	-	1	-	-	-	-
Female.....	-	-	-	-	2	1	3	2	2	-	-	-

Michigan

All races, both sexes.....	56	81	92	124	163	293	438	471	273	98	20	-
Male.....	31	41	46	63	75	124	214	199	100	26	10	-
Female.....	25	40	46	61	88	169	224	272	173	72	10	-
White, both sexes.....	33	53	63	92	135	239	380	414	247	82	16	-
Male.....	18	24	31	44	61	103	185	176	91	20	8	-
Female.....	15	29	32	48	74	136	195	238	156	62	8	-
Black, both sexes.....	23	26	28	31	28	47	54	54	25	15	3	-
Male.....	13	15	14	18	14	18	26	22	8	6	1	-
Female.....	10	11	14	13	14	29	28	32	17	9	2	-
American Indian 1/, both sexes.....	-	2	1	-	-	5	1	1	-	1	-	-
Male.....	-	2	1	-	-	2	1	1	-	-	-	-
Female.....	-	-	-	-	-	3	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	2	3	2	1	-	1	-
Male.....	-	-	-	1	-	1	2	-	1	-	1	-
Female.....	-	-	-	-	-	1	1	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Diseases of urinary system (N00-N39)

Minnesota

All races, both sexes.....	13	25	33	31	69	102	196	225	167	63	13	-
Male.....	9	14	17	15	48	57	97	107	62	22	3	-
Female.....	4	11	16	16	21	45	99	118	105	41	10	-
White, both sexes.....	11	19	28	28	57	90	191	224	162	62	13	-
Male.....	8	10	15	14	40	52	95	106	61	22	3	-
Female.....	3	9	13	14	17	38	96	118	101	40	10	-
Black, both sexes.....	2	4	1	1	4	7	2	1	1	1	-	-
Male.....	1	3	-	-	3	3	1	1	-	-	-	-
Female.....	1	1	1	1	1	4	1	-	1	1	-	-
American Indian 1/, both sexes.....	-	-	2	-	4	3	1	-	1	-	-	-
Male.....	-	-	1	-	3	1	-	-	-	-	-	-
Female.....	-	-	1	-	1	2	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	2	2	4	2	2	-	3	-	-	-
Male.....	-	1	1	1	2	1	1	-	1	-	-	-
Female.....	-	1	1	1	2	1	1	-	2	-	-	-

Mississippi

All races, both sexes.....	37	39	56	78	97	132	161	140	97	34	10	-
Male.....	22	17	28	33	45	69	80	49	38	9	2	-
Female.....	15	22	28	45	52	63	81	91	59	25	8	-
White, both sexes.....	16	12	27	33	48	88	99	94	67	18	5	-
Male.....	9	5	12	15	26	53	51	36	26	3	-	-
Female.....	7	7	15	18	22	35	48	58	41	15	5	-
Black, both sexes.....	21	27	27	43	49	43	62	46	29	16	5	-
Male.....	13	12	14	17	19	16	29	13	12	6	2	-
Female.....	8	15	13	26	30	27	33	33	17	10	3	-
American Indian 1/, both sexes.....	-	-	2	1	-	-	-	-	-	-	-	-
Male.....	-	-	2	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	1	-	-	1	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	1	-	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Diseases of urinary system (N00-N39)

Nebraska

All races, both sexes.....	5	12	17	12	33	39	61	86	65	24	7	-
Male.....	2	7	12	3	21	21	27	31	23	5	1	-
Female.....	3	5	5	9	12	18	34	55	42	19	6	-
White, both sexes.....	5	9	10	9	30	36	60	86	63	24	7	-
Male.....	2	6	10	3	19	19	27	31	22	5	1	-
Female.....	3	3	-	6	11	17	33	55	41	19	6	-
Black, both sexes.....	-	3	4	3	3	2	1	-	2	-	-	-
Male.....	-	1	1	-	2	1	-	-	1	-	-	-
Female.....	-	2	3	3	1	1	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	3	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	20	23	32	43	78	80	94	84	41	13	1	-
Male.....	11	11	20	27	50	43	52	39	16	8	-	-
Female.....	9	12	12	16	28	37	42	45	25	5	1	-
White, both sexes.....	16	20	24	35	63	65	83	74	34	11	-	-
Male.....	10	9	15	23	42	36	43	34	14	7	-	-
Female.....	6	11	9	12	21	29	40	40	20	4	-	-
Black, both sexes.....	2	2	4	5	10	11	6	7	4	1	-	-
Male.....	1	1	2	2	6	4	5	4	-	-	-	-
Female.....	1	1	2	3	4	7	1	3	4	1	-	-
American Indian 1/, both sexes.....	1	-	-	2	-	1	2	-	-	-	-	-
Male.....	-	-	-	2	-	1	1	-	-	-	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	4	1	5	3	3	3	3	1	1	-
Male.....	-	1	3	-	2	2	3	1	2	1	-	-
Female.....	1	-	1	1	3	1	-	2	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Diseases of urinary system (N00-N39)

New Hampshire

All races, both sexes.....	1	2	7	8	17	27	47	53	29	11	2	-
Male.....	1	1	4	4	8	11	22	24	12	3	-	-
Female.....	-	1	3	4	9	16	25	29	17	8	2	-
White, both sexes.....	1	2	7	8	17	27	46	53	29	11	2	-
Male.....	1	1	4	4	8	11	22	24	12	3	-	-
Female.....	-	1	3	4	9	16	24	29	17	8	2	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

New Jersey

All races, both sexes.....	52	66	100	108	158	300	408	445	334	102	11	-
Male.....	26	31	58	67	83	135	208	195	126	23	4	-
Female.....	26	35	42	41	75	165	200	250	208	79	7	-
White, both sexes.....	34	40	57	76	110	237	365	402	309	89	9	-
Male.....	16	23	34	50	59	114	189	181	117	21	3	-
Female.....	18	17	23	26	51	123	176	221	192	68	6	-
Black, both sexes.....	16	23	39	31	42	53	36	37	17	13	1	-
Male.....	8	6	22	16	19	17	14	11	4	2	-	-
Female.....	8	17	17	15	23	36	22	26	13	11	1	-
American Indian 1/, both sexes.....	-	-	1	-	1	1	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	3	3	1	5	9	7	6	8	-	1	-
Male.....	2	2	2	1	5	3	5	3	5	-	1	-
Female.....	-	1	1	-	-	6	2	3	3	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of urinary system (N00-N39)												
New Mexico												
All races, both sexes.....	8	15	16	20	33	44	75	67	33	17	4	-
Male.....	4	12	10	10	15	24	33	29	22	8	3	-
Female.....	4	3	6	10	18	20	42	38	11	9	1	-
White, both sexes.....	6	13	13	17	29	41	66	63	32	15	4	-
Male.....	2	11	8	9	14	22	30	27	21	8	3	-
Female.....	4	2	5	8	15	19	36	36	11	7	1	-
Black, both sexes.....	-	1	-	1	-	1	4	-	-	-	-	-
Male.....	-	1	-	1	-	1	2	-	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes.....	2	1	2	2	4	2	5	4	1	2	-	-
Male.....	2	-	2	-	1	1	1	2	1	-	-	-
Female.....	-	1	-	2	3	1	4	2	-	2	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
New York												
All races, both sexes.....	62	110	139	200	255	495	680	813	513	197	54	-
Male.....	34	61	64	90	135	246	302	335	170	47	10	-
Female.....	28	49	75	110	120	249	378	478	343	150	44	-
White, both sexes.....	41	76	95	148	192	411	588	741	459	177	43	-
Male.....	21	41	43	74	107	207	265	309	154	46	5	-
Female.....	20	35	52	74	85	204	323	432	305	131	38	-
Black, both sexes.....	19	30	41	46	55	71	76	66	48	19	9	-
Male.....	11	17	21	11	24	35	29	23	14	1	4	-
Female.....	8	13	20	35	31	36	47	43	34	18	5	-
American Indian 1/, both sexes.....	2	2	-	2	2	-	1	1	-	-	-	-
Male.....	2	1	-	2	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	2	-	1	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	3	4	6	13	15	5	6	1	2	-
Male.....	-	2	-	3	4	4	8	3	2	-	1	-
Female.....	-	-	3	1	2	9	7	2	4	1	1	-







## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of urinary system (N00-N39)												
Oregon												
All races, both sexes.....	13	20	23	27	49	78	121	140	100	40	8	-
Male.....	11	8	9	10	24	35	54	59	27	10	6	-
Female.....	2	12	14	17	25	43	67	81	73	30	2	-
White, both sexes.....	9	15	20	25	47	77	118	134	98	40	8	-
Male.....	8	6	8	10	24	34	52	55	27	10	6	-
Female.....	1	9	12	15	23	43	66	79	71	30	2	-
Black, both sexes.....	3	2	-	-	1	-	-	4	-	-	-	-
Male.....	2	1	-	-	-	-	-	3	-	-	-	-
Female.....	1	1	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	1	1	1	1	-	-	-	1	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	1	1	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	2	1	1	1	3	1	2	-	-	-
Male.....	-	1	1	-	-	1	2	1	-	-	-	-
Female.....	-	1	1	1	1	-	1	-	2	-	-	-
Pennsylvania												
All races, both sexes.....	69	93	135	211	328	542	751	782	532	169	30	-
Male.....	36	59	71	114	150	269	323	322	175	40	7	-
Female.....	33	34	64	97	178	273	428	460	357	129	23	-
White, both sexes.....	51	71	105	165	279	484	681	728	493	153	28	-
Male.....	24	43	54	84	133	248	297	299	164	36	6	-
Female.....	27	28	51	81	146	236	384	429	329	117	22	-
Black, both sexes.....	18	22	28	43	49	55	66	52	39	16	2	-
Male.....	12	16	15	28	17	19	22	23	11	4	1	-
Female.....	6	6	13	15	32	36	44	29	28	12	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	3	-	2	4	2	-	-	-	-
Male.....	-	-	2	2	-	1	4	-	-	-	-	-
Female.....	-	-	-	1	-	1	-	2	-	-	-	-





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Diseases of urinary system (N00-N39)												
Texas												
All races, both sexes.....	155	193	255	327	446	593	727	722	507	179	46	-
Male.....	84	96	136	171	219	286	326	317	184	46	8	-
Female.....	71	97	119	156	227	307	401	405	323	133	38	-
White, both sexes.....	108	143	188	245	363	502	638	620	458	160	38	-
Male.....	64	67	96	132	180	245	289	278	169	40	5	-
Female.....	44	76	92	113	183	257	349	342	289	120	33	-
Black, both sexes.....	45	42	59	75	76	78	76	91	46	19	7	-
Male.....	18	24	33	36	33	37	30	34	13	6	3	-
Female.....	27	18	26	39	43	41	46	57	33	13	4	-
American Indian 1/, both sexes.....	-	3	1	-	1	-	-	2	-	-	-	-
Male.....	-	2	1	-	1	-	-	2	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	5	7	7	6	13	13	9	3	-	1	-
Male.....	2	3	6	3	5	4	7	3	2	-	-	-
Female.....	-	2	1	4	1	9	6	6	1	-	1	-
Utah												
All races, both sexes.....	4	15	13	17	39	46	59	54	38	21	3	-
Male.....	2	7	5	4	20	18	32	20	11	10	-	-
Female.....	2	8	8	13	19	28	27	34	27	11	3	-
White, both sexes.....	3	14	13	14	38	43	58	53	36	20	3	-
Male.....	1	7	5	4	20	17	31	20	11	10	-	-
Female.....	2	7	8	10	18	26	27	33	25	10	3	-
Black, both sexes.....	-	-	-	1	1	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	1	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	2	-	3	-	1	2	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	1	-	2	-	2	-	1	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Diseases of urinary system (N00-N39)

Vermont

All races, both sexes.....	-	4	2	4	8	9	12	21	14	4	-	-
Male.....	-	3	2	1	5	5	5	6	2	-	-	-
Female.....	-	1	-	3	3	4	7	15	12	4	-	-
White, both sexes.....	-	4	2	4	8	9	12	21	14	4	-	-
Male.....	-	3	2	1	5	5	5	6	2	-	-	-
Female.....	-	1	-	3	3	4	7	15	12	4	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	48	78	90	140	188	256	349	354	220	78	14	-
Male.....	26	46	40	67	90	133	150	145	84	14	1	-
Female.....	22	32	50	73	98	123	199	209	136	64	13	-
White, both sexes.....	24	38	48	89	119	181	278	274	177	64	7	-
Male.....	14	25	25	46	58	95	119	114	71	12	1	-
Female.....	10	13	23	43	61	86	159	160	106	52	6	-
Black, both sexes.....	23	40	40	49	61	71	66	74	41	14	7	-
Male.....	11	21	14	20	28	38	30	30	13	2	-	-
Female.....	12	19	26	29	33	33	36	44	28	12	7	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	2	2	8	4	5	5	2	-	-	-
Male.....	1	-	1	1	4	-	1	1	-	-	-	-
Female.....	-	-	1	1	4	4	4	4	2	-	-	-





































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Glomerular and renal tubulo-interstitial diseases (N00-N15)

Arkansas

All races, both sexes.....	-	1	2	1	9	4	8	8	6	1	1	-
Male.....	-	1	1	-	5	2	2	4	3	-	-	-
Female.....	-	-	1	1	4	2	6	4	3	1	1	-
White, both sexes.....	-	1	2	1	6	4	7	7	5	1	-	-
Male.....	-	1	1	-	4	2	1	4	2	-	-	-
Female.....	-	-	1	1	2	2	6	3	3	1	-	-
Black, both sexes.....	-	-	-	-	3	-	1	1	1	-	1	-
Male.....	-	-	-	-	1	-	1	-	1	-	-	-
Female.....	-	-	-	-	2	-	-	1	-	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	15	13	23	21	35	60	85	97	76	36	7	-
Male.....	9	7	14	12	21	31	41	41	34	20	2	-
Female.....	6	6	9	9	14	29	44	56	42	16	5	-
White, both sexes.....	13	9	20	15	25	48	70	76	57	31	5	-
Male.....	8	4	13	9	15	27	33	33	27	17	-	-
Female.....	5	5	7	6	10	21	37	43	30	14	5	-
Black, both sexes.....	1	3	3	6	5	8	2	13	7	2	-	-
Male.....	-	2	1	3	3	2	1	4	2	-	-	-
Female.....	1	1	2	3	2	6	1	9	5	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	1	2	-	-	-
Male.....	-	-	-	-	-	-	1	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	-	5	4	12	7	10	3	2	-
Male.....	1	1	-	-	3	2	6	4	4	3	2	-
Female.....	-	-	-	-	2	2	6	3	6	-	-	-

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Glomerular and renal tubulo-interstitial diseases (N00-N15)

Maine

All races, both sexes.....	1	2	3	3	1	5	5	2	6	1	-	-
Male.....	1	2	3	-	-	1	1	1	4	-	-	-
Female.....	-	-	-	3	1	4	4	1	2	1	-	-
White, both sexes.....	1	2	3	3	1	5	5	2	6	1	-	-
Male.....	1	2	3	-	-	1	1	1	4	-	-	-
Female.....	-	-	-	3	1	4	4	1	2	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	2	7	-	6	4	10	13	11	9	5	1	-
Male.....	1	3	-	4	2	3	5	2	4	2	-	-
Female.....	1	4	-	2	2	7	8	9	5	3	1	-
White, both sexes.....	-	4	-	4	2	9	12	7	5	4	1	-
Male.....	-	-	-	2	1	3	4	2	2	2	-	-
Female.....	-	4	-	2	1	6	8	5	3	2	1	-
Black, both sexes.....	2	3	-	2	2	1	1	3	4	1	-	-
Male.....	1	3	-	2	1	-	1	-	2	-	-	-
Female.....	1	-	-	-	1	1	-	3	2	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Glomerular and renal tubulo-interstitial diseases (N00-N15)

New Hampshire

All races, both sexes.....	-	-	1	1	1	2	1	3	1	-	-	-
Male.....	-	-	-	-	1	1	1	2	1	-	-	-
Female.....	-	-	1	1	-	1	-	1	-	-	-	-
White, both sexes.....	-	-	1	1	1	2	1	3	1	-	-	-
Male.....	-	-	-	-	1	1	1	2	1	-	-	-
Female.....	-	-	1	1	-	1	-	1	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	2	2	2	4	5	12	18	17	15	4	-	-
Male.....	2	1	-	3	2	6	10	6	6	-	-	-
Female.....	-	1	2	1	3	6	8	11	9	4	-	-
White, both sexes.....	2	1	2	3	5	8	15	15	14	2	-	-
Male.....	2	1	-	2	2	5	8	5	5	-	-	-
Female.....	-	-	2	1	3	3	7	10	9	2	-	-
Black, both sexes.....	-	1	-	1	-	3	2	2	1	2	-	-
Male.....	-	-	-	1	-	1	1	1	1	-	-	-
Female.....	-	1	-	-	-	2	1	1	-	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Glomerular and renal tubulo-interstitial diseases (N00-N15)

New Mexico

All races, both sexes.....	1	2	2	1	2	7	3	7	6	1	1	-
Male.....	1	2	1	-	1	4	1	4	5	-	1	-
Female.....	-	-	1	1	1	3	2	3	1	1	-	-
White, both sexes.....	-	2	2	1	2	7	3	6	5	1	1	-
Male.....	-	2	1	-	1	4	1	4	4	-	1	-
Female.....	-	-	1	1	1	3	2	2	1	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	1	1	-	-	-
Male.....	1	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	3	4	6	11	14	20	34	44	28	12	-	-
Male.....	-	-	5	5	7	15	13	21	9	-	-	-
Female.....	3	4	1	6	7	5	21	23	19	12	-	-
White, both sexes.....	3	1	2	10	10	18	28	37	24	12	-	-
Male.....	-	-	2	5	5	14	10	18	7	-	-	-
Female.....	3	1	-	5	5	4	18	19	17	12	-	-
Black, both sexes.....	-	3	4	1	3	2	6	6	3	-	-	-
Male.....	-	-	3	-	2	1	3	2	2	-	-	-
Female.....	-	3	1	1	1	1	3	4	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-





















































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Glomerular diseases (N00-N07)

Arkansas

All races, both sexes.....	-	-	2	1	8	-	4	7	1	-	1	-
Male.....	-	-	1	-	4	-	1	4	1	-	-	-
Female.....	-	-	1	1	4	-	3	3	-	-	1	-
White, both sexes.....	-	-	2	1	6	-	3	6	1	-	-	-
Male.....	-	-	1	-	4	-	-	4	1	-	-	-
Female.....	-	-	1	1	2	-	3	2	-	-	-	-
Black, both sexes.....	-	-	-	-	2	-	1	1	-	-	1	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	2	-	-	1	-	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	11	11	15	15	25	45	66	82	60	25	4	-
Male.....	8	7	11	8	15	20	37	35	29	16	2	-
Female.....	3	4	4	7	10	25	29	47	31	9	2	-
White, both sexes.....	10	7	12	9	17	34	53	63	44	21	2	-
Male.....	7	4	10	5	10	17	30	28	23	14	-	-
Female.....	3	3	2	4	7	17	23	35	21	7	2	-
Black, both sexes.....	-	3	3	6	4	7	2	13	6	2	-	-
Male.....	-	2	1	3	2	1	1	4	1	-	-	-
Female.....	-	1	2	3	2	6	1	9	5	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	2	-	-	-
Male.....	-	-	-	-	-	-	1	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	-	4	4	10	6	8	2	2	-
Male.....	1	1	-	-	3	2	5	3	4	2	2	-
Female.....	-	-	-	-	1	2	5	3	4	-	-	-

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Glomerular diseases (N00-N07)

Maine

All races, both sexes.....	1	2	3	3	-	5	5	1	5	1	-	-
Male.....	1	2	3	-	-	1	1	-	3	-	-	-
Female.....	-	-	-	3	-	4	4	1	2	1	-	-
White, both sexes.....	1	2	3	3	-	5	5	1	5	1	-	-
Male.....	1	2	3	-	-	1	1	-	3	-	-	-
Female.....	-	-	-	3	-	4	4	1	2	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	2	4	-	4	2	7	10	7	7	4	1	-
Male.....	1	2	-	3	1	3	4	1	4	1	-	-
Female.....	1	2	-	1	1	4	6	6	3	3	1	-
White, both sexes.....	-	2	-	2	1	6	9	5	3	3	1	-
Male.....	-	-	-	1	-	3	3	1	2	1	-	-
Female.....	-	2	-	1	1	3	6	4	1	2	1	-
Black, both sexes.....	2	2	-	2	1	1	1	1	4	1	-	-
Male.....	1	2	-	2	1	-	1	-	2	-	-	-
Female.....	1	-	-	-	-	1	-	1	2	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Glomerular diseases (N00-N07)

New Mexico

All races, both sexes.....	-	1	1	-	1	5	2	4	4	-	1	-
Male.....	-	1	1	-	1	4	1	3	3	-	1	-
Female.....	-	-	-	-	-	1	1	1	1	-	-	-
White, both sexes.....	-	1	1	-	1	5	2	3	3	-	1	-
Male.....	-	1	1	-	1	4	1	3	2	-	1	-
Female.....	-	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	2	1	4	7	8	12	26	30	17	11	-	-
Male.....	-	-	3	3	3	8	7	14	5	-	-	-
Female.....	2	1	1	4	5	4	19	16	12	11	-	-
White, both sexes.....	2	-	1	6	6	11	21	24	14	11	-	-
Male.....	-	-	1	3	2	8	5	12	4	-	-	-
Female.....	2	-	-	3	4	3	16	12	10	11	-	-
Black, both sexes.....	-	1	3	1	2	1	5	5	2	-	-	-
Male.....	-	-	2	-	1	-	2	1	1	-	-	-
Female.....	-	1	1	1	1	1	3	4	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute and rapidly progressive nephritic syndrome (N00-N01)

Arkansas

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute and rapidly progressive nephritic syndrome (N00-N01)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute and rapidly progressive nephritic syndrome (N00-N01)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute and rapidly progressive nephritic syndrome (N00-N01)

Missouri

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute and rapidly progressive nephritic syndrome (N00-N01)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute and rapidly progressive nephritic syndrome (N00-N01)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Acute and rapidly progressive nephritic syndrome (N00-N01)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Colorado

Table with 13 columns (Age groups) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Age groups) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Illinois

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

New Hampshire

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Ohio

Table with 13 columns (Age groups) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Age groups) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Rhode Island

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

South Dakota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Texas

Table with 13 columns (Age groups) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Age groups) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Acute and rapidly progressive nephritic syndrome (N00-N01)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Nephrotic syndrome (N04)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Nephrotic syndrome (N04)

Delaware

Table with 12 columns for age groups and rows for race/sex categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns for age groups and rows for race/sex categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Nephrotic syndrome (N04)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Nephrotic syndrome (N04)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows for Maine (All races, White, Black, American Indian, Asian or Pacific Islander).

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows for Maryland (All races, White, Black, American Indian, Asian or Pacific Islander).















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Nephrotic syndrome (N04)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Nephrotic syndrome (N04)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Nephrotic syndrome (N04)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Nephrotic syndrome (N04)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other and unspecified glomerular diseases (N02-N03,N05-N07)

Arkansas

All races, both sexes.....	-	-	2	1	6	-	4	6	1	-	1	-
Male.....	-	-	1	-	3	-	1	4	1	-	-	-
Female.....	-	-	1	1	3	-	3	2	-	-	1	-
White, both sexes.....	-	-	2	1	4	-	3	5	1	-	-	-
Male.....	-	-	1	-	3	-	-	4	1	-	-	-
Female.....	-	-	1	1	1	-	3	1	-	-	-	-
Black, both sexes.....	-	-	-	-	2	-	1	1	-	-	1	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	2	-	-	1	-	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	10	11	14	15	24	44	65	75	57	25	4	-
Male.....	7	7	10	8	15	19	36	32	27	16	2	-
Female.....	3	4	4	7	9	25	29	43	30	9	2	-
White, both sexes.....	9	7	11	9	16	33	52	56	41	21	2	-
Male.....	6	4	9	5	10	16	29	25	21	14	-	-
Female.....	3	3	2	4	6	17	23	31	20	7	2	-
Black, both sexes.....	-	3	3	6	4	7	2	13	6	2	-	-
Male.....	-	2	1	3	2	1	1	4	1	-	-	-
Female.....	-	1	2	3	2	6	1	9	5	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	2	-	-	-
Male.....	-	-	-	-	-	-	1	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	-	-	4	4	10	6	8	2	2	-
Male.....	1	1	-	-	3	2	5	3	4	2	2	-
Female.....	-	-	-	-	1	2	5	3	4	-	-	-

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified glomerular diseases (N02-N03,N05-N07)

Maine

All races, both sexes.....	1	2	2	3	-	5	5	1	5	1	-	-
Male.....	1	2	2	-	-	1	1	-	3	-	-	-
Female.....	-	-	-	3	-	4	4	1	2	1	-	-
White, both sexes.....	1	2	2	3	-	5	5	1	5	1	-	-
Male.....	1	2	2	-	-	1	1	-	3	-	-	-
Female.....	-	-	-	3	-	4	4	1	2	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	1	3	-	2	1	7	10	7	7	4	1	-
Male.....	1	1	-	1	1	3	4	1	4	1	-	-
Female.....	-	2	-	1	-	4	6	6	3	3	1	-
White, both sexes.....	-	2	-	1	-	6	9	5	3	3	1	-
Male.....	-	-	-	-	-	3	3	1	2	1	-	-
Female.....	-	2	-	1	-	3	6	4	1	2	1	-
Black, both sexes.....	1	1	-	1	1	1	1	1	4	1	-	-
Male.....	1	1	-	1	1	-	1	-	2	-	-	-
Female.....	-	-	-	-	-	1	-	1	2	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other and unspecified glomerular diseases (N02-N03,N05-N07)

New Mexico

All races, both sexes.....	-	1	1	-	1	5	1	4	3	-	1	-
Male.....	-	1	1	-	1	4	-	3	2	-	1	-
Female.....	-	-	-	-	-	1	1	1	1	-	-	-
White, both sexes.....	-	1	1	-	1	5	1	3	2	-	1	-
Male.....	-	1	1	-	1	4	-	3	1	-	1	-
Female.....	-	-	-	-	-	1	1	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	2	1	4	6	8	9	24	30	17	11	-	-
Male.....	-	-	3	3	3	5	6	14	5	-	-	-
Female.....	2	1	1	3	5	4	18	16	12	11	-	-
White, both sexes.....	2	-	1	6	6	8	19	24	14	11	-	-
Male.....	-	-	1	3	2	5	4	12	4	-	-	-
Female.....	2	-	-	3	4	3	15	12	10	11	-	-
Black, both sexes.....	-	1	3	-	2	1	5	5	2	-	-	-
Male.....	-	-	2	-	1	-	2	1	1	-	-	-
Female.....	-	1	1	-	1	1	3	4	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-





















































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Renal tubulo-interstitial diseases (N10-N15)

Arkansas

All races, both sexes.....	-	1	-	-	1	4	4	1	5	1	-	-
Male.....	-	1	-	-	1	2	1	-	2	-	-	-
Female.....	-	-	-	-	-	2	3	1	3	1	-	-
White, both sexes.....	-	1	-	-	-	4	4	1	4	1	-	-
Male.....	-	1	-	-	-	2	1	-	1	-	-	-
Female.....	-	-	-	-	-	2	3	1	3	1	-	-
Black, both sexes.....	-	-	-	-	1	-	-	-	1	-	-	-
Male.....	-	-	-	-	1	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	4	2	8	6	10	15	19	15	16	11	3	-
Male.....	1	-	3	4	6	11	4	6	5	4	-	-
Female.....	3	2	5	2	4	4	15	9	11	7	3	-
White, both sexes.....	3	2	8	6	8	14	17	13	13	10	3	-
Male.....	1	-	3	4	5	10	3	5	4	3	-	-
Female.....	2	2	5	2	3	4	14	8	9	7	3	-
Black, both sexes.....	1	-	-	-	1	1	-	-	1	-	-	-
Male.....	-	-	-	-	1	1	-	-	1	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	2	1	2	1	-	-
Male.....	-	-	-	-	-	-	1	1	-	1	-	-
Female.....	-	-	-	-	1	-	1	-	2	-	-	-



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Renal tubulo-interstitial diseases (N10-N15)

New Hampshire

All races, both sexes.....	-	-	-	-	1	1	-	2	-	-	-	-
Male.....	-	-	-	-	1	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
White, both sexes.....	-	-	-	-	1	1	-	2	-	-	-	-
Male.....	-	-	-	-	1	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	1	-	-	1	2	3	1	3	3	1	-	-
Male.....	1	-	-	-	-	1	-	1	2	-	-	-
Female.....	-	-	-	1	2	2	1	2	1	1	-	-
White, both sexes.....	1	-	-	1	2	2	1	2	3	1	-	-
Male.....	1	-	-	-	-	1	-	-	2	-	-	-
Female.....	-	-	-	1	2	1	1	2	1	1	-	-
Black, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Renal tubulo-interstitial diseases (N10-N15)

Wisconsin

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Wyoming

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Renal failure (N17-N19)												
Arkansas												
All races, both sexes.....	641	1	-	-	1	-	-	3	5	3	15	22
Male.....	313	-	-	-	-	-	-	2	3	3	9	10
Female.....	328	1	-	-	1	-	-	1	2	-	6	12
White, both sexes.....	511	-	-	-	1	-	-	2	5	2	9	13
Male.....	249	-	-	-	-	-	-	2	3	2	5	6
Female.....	262	-	-	-	1	-	-	-	2	-	4	7
Black, both sexes.....	126	1	-	-	-	-	-	1	-	1	6	9
Male.....	63	-	-	-	-	-	-	-	-	1	4	4
Female.....	63	1	-	-	-	-	-	1	-	-	2	5
American Indian 1/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	2,440	12	-	-	-	1	7	6	12	17	31	54
Male.....	1,242	7	-	-	-	1	4	5	5	11	18	25
Female.....	1,198	5	-	-	-	-	3	1	7	6	13	29
White, both sexes.....	1,849	10	-	-	-	-	5	4	7	10	22	34
Male.....	954	5	-	-	-	-	2	4	3	8	11	13
Female.....	895	5	-	-	-	-	3	-	4	2	11	21
Black, both sexes.....	299	2	-	-	-	-	2	1	5	7	8	14
Male.....	139	2	-	-	-	-	2	1	2	3	6	8
Female.....	160	-	-	-	-	-	-	-	3	4	2	6
American Indian 1/, both sexes.....	20	-	-	-	-	-	-	-	-	-	-	-
Male.....	7	-	-	-	-	-	-	-	-	-	-	-
Female.....	13	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	272	-	-	-	-	1	-	1	-	-	1	6
Male.....	142	-	-	-	-	1	-	-	-	-	1	4
Female.....	130	-	-	-	-	-	-	1	-	-	-	2























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Alaska												
All races, both sexes.....	1	4	4	1	5	8	3	5	5	1	-	-
Male.....	1	1	3	1	1	3	1	3	2	-	-	-
Female.....	-	3	1	-	4	5	2	2	3	1	-	-
White, both sexes.....	1	3	2	-	4	6	2	4	4	1	-	-
Male.....	1	1	2	-	1	2	1	2	2	-	-	-
Female.....	-	2	-	-	3	4	1	2	2	1	-	-
Black, both sexes.....	-	-	1	-	1	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	1	1	1	-	2	1	1	1	-	-	-
Male.....	-	-	-	1	-	1	-	1	-	-	-	-
Female.....	-	1	1	-	-	1	1	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Arizona												
All races, both sexes.....	21	13	29	49	42	71	93	76	60	19	1	-
Male.....	14	10	18	22	23	40	49	30	32	3	-	-
Female.....	7	3	11	27	19	31	44	46	28	16	1	-
White, both sexes.....	16	12	26	38	38	65	82	66	52	18	1	-
Male.....	10	9	18	19	21	35	43	27	29	3	-	-
Female.....	6	3	8	19	17	30	39	39	23	15	1	-
Black, both sexes.....	2	-	-	4	-	5	1	5	4	1	-	-
Male.....	2	-	-	1	-	5	1	3	1	-	-	-
Female.....	-	-	-	3	-	-	-	2	3	1	-	-
American Indian 1/, both sexes.....	3	1	3	6	4	1	8	5	3	-	-	-
Male.....	2	1	-	2	2	-	3	-	2	-	-	-
Female.....	1	-	3	4	2	1	5	5	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	-	2	-	1	-	-	-
Male.....	-	-	-	-	-	-	2	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	1	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Arkansas												
All races, both sexes.....	20	17	25	44	63	95	96	113	81	29	8	-
Male.....	12	13	12	27	41	49	47	49	24	9	3	-
Female.....	8	4	13	17	22	46	49	64	57	20	5	-
White, both sexes.....	11	12	14	34	49	75	86	102	68	22	6	-
Male.....	4	8	7	20	36	41	43	43	21	7	1	-
Female.....	7	4	7	14	13	34	43	59	47	15	5	-
Black, both sexes.....	9	5	11	10	12	20	9	11	12	7	2	-
Male.....	8	5	5	7	5	8	4	6	2	2	2	-
Female.....	1	-	6	3	7	12	5	5	10	5	-	-
American Indian 1/, both sexes.....	-	-	-	-	1	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
California												
All races, both sexes.....	100	153	167	188	263	354	403	373	203	72	24	-
Male.....	58	80	94	99	134	181	219	181	85	30	5	-
Female.....	42	73	73	89	129	173	184	192	118	42	19	-
White, both sexes.....	68	104	106	139	197	280	315	302	165	63	18	-
Male.....	38	54	59	71	99	151	182	149	76	27	2	-
Female.....	30	50	47	68	98	129	133	153	89	36	16	-
Black, both sexes.....	22	31	35	22	32	36	32	29	16	3	2	-
Male.....	16	15	18	9	15	13	10	13	4	1	1	-
Female.....	6	16	17	13	17	23	22	16	12	2	1	-
American Indian 1/, both sexes.....	1	2	6	4	3	3	-	1	-	-	-	-
Male.....	-	-	4	1	1	-	-	1	-	-	-	-
Female.....	1	2	2	3	2	3	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	9	16	20	23	31	35	56	41	22	6	4	-
Male.....	4	11	13	18	19	17	27	18	5	2	2	-
Female.....	5	5	7	5	12	18	29	23	17	4	2	-







## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Florida												
All races, both sexes.....	73	125	142	175	252	389	497	490	307	110	17	-
Male.....	45	70	78	97	150	224	273	263	140	47	5	-
Female.....	28	55	64	78	102	165	224	227	167	63	12	-
White, both sexes.....	54	84	100	126	193	329	436	439	281	102	16	-
Male.....	32	51	60	71	116	191	251	249	130	45	5	-
Female.....	22	33	40	55	77	138	185	190	151	57	11	-
Black, both sexes.....	17	38	40	48	55	56	52	48	22	8	1	-
Male.....	11	18	17	25	32	33	19	14	7	2	-	-
Female.....	6	20	23	23	23	23	33	34	15	6	1	-
American Indian 1/, both sexes.....	-	1	1	-	1	1	2	-	1	-	-	-
Male.....	-	-	1	-	1	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	1	2	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	2	1	1	3	3	7	3	3	-	-	-
Male.....	2	1	-	1	1	-	3	-	3	-	-	-
Female.....	-	1	1	-	2	3	4	3	-	-	-	-
Georgia												
All races, both sexes.....	68	87	117	149	171	244	239	244	121	42	10	-
Male.....	29	48	63	67	83	107	110	92	49	15	3	-
Female.....	39	39	54	82	88	137	129	152	72	27	7	-
White, both sexes.....	20	43	47	78	90	158	168	164	86	26	6	-
Male.....	9	25	26	38	48	78	87	69	35	10	2	-
Female.....	11	18	21	40	42	80	81	95	51	16	4	-
Black, both sexes.....	48	44	69	70	77	82	68	77	35	16	4	-
Male.....	20	23	36	29	31	28	22	21	14	5	1	-
Female.....	28	21	33	41	46	54	46	56	21	11	3	-
American Indian 1/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	4	4	3	3	-	-	-	-
Male.....	-	-	-	-	4	1	1	2	-	-	-	-
Female.....	-	-	-	1	-	3	2	1	-	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Renal failure (N17-N19)

Iowa

All races, both sexes.....	6	4	8	6	18	35	43	53	37	19	5	-
Male.....	2	3	5	4	9	24	22	28	15	6	1	-
Female.....	4	1	3	2	9	11	21	25	22	13	4	-
White, both sexes.....	6	4	8	6	17	34	40	53	36	19	5	-
Male.....	2	3	5	4	8	23	22	28	14	6	1	-
Female.....	4	1	3	2	9	11	18	25	22	13	4	-
Black, both sexes.....	-	-	-	-	-	1	3	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	3	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	1	-	-	-	1	-	-	-
Male.....	-	-	-	-	1	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Kansas

All races, both sexes.....	8	23	21	36	45	74	100	105	74	21	5	-
Male.....	5	13	7	21	31	37	48	47	31	12	1	-
Female.....	3	10	14	15	14	37	52	58	43	9	4	-
White, both sexes.....	6	19	16	29	40	68	93	97	71	21	4	-
Male.....	3	10	6	19	27	35	45	45	30	12	1	-
Female.....	3	9	10	10	13	33	48	52	41	9	3	-
Black, both sexes.....	2	4	3	6	4	5	5	8	3	-	1	-
Male.....	2	3	1	1	3	2	2	2	1	-	-	-
Female.....	-	1	2	5	1	3	3	6	2	-	1	-
American Indian 1/, both sexes.....	-	-	2	1	1	-	2	-	-	-	-	-
Male.....	-	-	-	1	1	-	1	-	-	-	-	-
Female.....	-	-	2	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Kentucky												
All races, both sexes.....	24	38	54	62	114	128	175	156	108	34	5	-
Male.....	12	22	40	26	57	68	77	66	40	11	-	-
Female.....	12	16	14	36	57	60	98	90	68	23	5	-
White, both sexes.....	19	31	43	54	98	116	164	140	98	33	4	-
Male.....	8	19	31	20	49	62	71	60	38	11	-	-
Female.....	11	12	12	34	49	54	93	80	60	22	4	-
Black, both sexes.....	5	7	11	8	15	12	10	16	10	1	1	-
Male.....	4	3	9	6	8	6	6	6	2	-	-	-
Female.....	1	4	2	2	7	6	4	10	8	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	1	-	-	-	-	-
Louisiana												
All races, both sexes.....	46	64	63	92	115	148	190	149	106	37	6	-
Male.....	25	28	37	41	50	82	106	67	36	11	1	-
Female.....	21	36	26	51	65	66	84	82	70	26	5	-
White, both sexes.....	22	24	29	57	54	89	140	110	68	28	2	-
Male.....	15	11	19	25	25	54	81	48	25	6	-	-
Female.....	7	13	10	32	29	35	59	62	43	22	2	-
Black, both sexes.....	24	39	34	35	60	58	50	38	36	9	4	-
Male.....	10	16	18	16	25	27	25	18	10	5	1	-
Female.....	14	23	16	19	35	31	25	20	26	4	3	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	-	1	1	-	1	2	-	-	-
Male.....	-	1	-	-	-	1	-	1	1	-	-	-
Female.....	-	-	-	-	1	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Renal failure (N17-N19)

Maine

All races, both sexes.....	2	5	11	16	18	30	54	48	29	16	3	-
Male.....	2	5	5	9	9	13	30	21	13	7	1	-
Female.....	-	-	6	7	9	17	24	27	16	9	2	-
White, both sexes.....	2	5	11	16	18	29	54	48	28	16	3	-
Male.....	2	5	5	9	9	12	30	21	13	7	1	-
Female.....	-	-	6	7	9	17	24	27	15	9	2	-
Black, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Maryland

All races, both sexes.....	22	33	61	55	56	75	137	126	57	21	6	-
Male.....	13	20	36	28	34	35	69	54	28	10	1	-
Female.....	9	13	25	27	22	40	68	72	29	11	5	-
White, both sexes.....	9	17	24	34	38	44	98	96	50	16	5	-
Male.....	6	11	16	17	22	23	49	43	24	7	-	-
Female.....	3	6	8	17	16	21	49	53	26	9	5	-
Black, both sexes.....	12	15	36	20	14	30	38	28	6	4	1	-
Male.....	6	8	20	11	10	12	19	11	3	2	1	-
Female.....	6	7	16	9	4	18	19	17	3	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	1	1	4	1	1	2	1	1	-	-
Male.....	1	1	-	-	2	-	1	-	1	1	-	-
Female.....	-	-	1	1	2	1	-	2	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Massachusetts												
All races, both sexes.....	19	33	48	73	112	175	256	262	202	77	12	-
Male.....	15	18	30	41	60	92	128	131	85	28	3	-
Female.....	4	15	18	32	52	83	128	131	117	49	9	-
White, both sexes.....	16	29	41	58	95	153	242	243	196	75	11	-
Male.....	12	16	26	33	53	76	123	125	83	27	3	-
Female.....	4	13	15	25	42	77	119	118	113	48	8	-
Black, both sexes.....	2	4	6	14	14	18	12	18	5	2	1	-
Male.....	2	2	3	7	6	13	5	5	2	1	-	-
Female.....	-	2	3	7	8	5	7	13	3	1	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	1	1	3	4	2	1	1	-	-	-
Male.....	1	-	1	1	1	3	-	1	-	-	-	-
Female.....	-	-	-	-	2	1	2	-	1	-	-	-
Michigan												
All races, both sexes.....	43	61	68	91	128	194	268	314	171	61	16	-
Male.....	25	33	36	44	61	84	151	150	67	17	9	-
Female.....	18	28	32	47	67	110	117	164	104	44	7	-
White, both sexes.....	24	37	46	68	102	149	225	269	153	49	13	-
Male.....	13	19	23	30	48	67	130	131	60	11	8	-
Female.....	11	18	23	38	54	82	95	138	93	38	5	-
Black, both sexes.....	19	23	21	22	26	39	41	43	17	12	3	-
Male.....	12	13	12	13	13	15	20	18	6	6	1	-
Female.....	7	10	9	9	13	24	21	25	11	6	2	-
American Indian 1/, both sexes.....	-	1	1	-	-	4	-	1	-	-	-	-
Male.....	-	1	1	-	-	1	-	1	-	-	-	-
Female.....	-	-	-	-	-	3	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	2	2	1	1	-	-	-
Male.....	-	-	-	1	-	1	1	-	1	-	-	-
Female.....	-	-	-	-	-	1	1	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Minnesota												
All races, both sexes.....	9	18	26	22	62	79	149	154	122	44	8	-
Male.....	5	12	12	13	43	42	73	82	46	18	3	-
Female.....	4	6	14	9	19	37	76	72	76	26	5	-
White, both sexes.....	8	13	21	20	50	68	145	153	117	44	8	-
Male.....	5	9	10	12	35	38	71	81	45	18	3	-
Female.....	3	4	11	8	15	30	74	72	72	26	5	-
Black, both sexes.....	1	3	1	-	4	6	2	1	1	-	-	-
Male.....	-	2	-	-	3	2	1	1	-	-	-	-
Female.....	1	1	1	-	1	4	1	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	2	-	4	3	-	-	1	-	-	-
Male.....	-	-	1	-	3	1	-	-	-	-	-	-
Female.....	-	-	1	-	1	2	-	-	1	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	2	2	4	2	2	-	3	-	-	-
Male.....	-	1	1	1	2	1	1	-	1	-	-	-
Female.....	-	1	1	1	2	1	1	-	2	-	-	-
Mississippi												
All races, both sexes.....	32	30	40	64	73	101	112	91	62	18	6	-
Male.....	18	13	21	27	33	51	55	38	25	7	2	-
Female.....	14	17	19	37	40	50	57	53	37	11	4	-
White, both sexes.....	13	8	17	27	34	70	71	61	41	9	3	-
Male.....	7	4	8	11	20	40	36	29	16	2	-	-
Female.....	6	4	9	16	14	30	35	32	25	7	3	-
Black, both sexes.....	19	22	21	35	39	30	41	30	21	9	3	-
Male.....	11	9	11	15	13	11	19	9	9	5	2	-
Female.....	8	13	10	20	26	19	22	21	12	4	1	-
American Indian 1/, both sexes.....	-	-	2	1	-	-	-	-	-	-	-	-
Male.....	-	-	2	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	1	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Renal failure (N17-N19)

Nebraska

All races, both sexes.....	3	9	15	10	24	30	42	51	46	15	4	-
Male.....	2	6	10	3	16	17	22	24	17	3	1	-
Female.....	1	3	5	7	8	13	20	27	29	12	3	-
White, both sexes.....	3	6	8	7	21	27	41	51	45	15	4	-
Male.....	2	5	8	3	14	15	22	24	16	3	1	-
Female.....	1	1	-	4	7	12	19	27	29	12	3	-
Black, both sexes.....	-	3	4	3	3	2	1	-	1	-	-	-
Male.....	-	1	1	-	2	1	-	-	1	-	-	-
Female.....	-	2	3	3	1	1	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	3	-	-	-	-	-	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	2	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Nevada

All races, both sexes.....	16	21	27	37	65	71	72	65	30	10	1	-
Male.....	10	11	18	23	47	36	38	32	11	7	-	-
Female.....	6	10	9	14	18	35	34	33	19	3	1	-
White, both sexes.....	13	18	19	29	52	57	64	56	24	8	-	-
Male.....	9	9	13	19	40	29	32	28	10	6	-	-
Female.....	4	9	6	10	12	28	32	28	14	2	-	-
Black, both sexes.....	1	2	4	5	9	10	4	6	4	1	-	-
Male.....	1	1	2	2	6	4	3	3	-	-	-	-
Female.....	-	1	2	3	3	6	1	3	4	1	-	-
American Indian 1/, both sexes.....	1	-	-	2	-	1	1	-	-	-	-	-
Male.....	-	-	-	2	-	1	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	1	4	1	4	3	3	3	2	1	1	-
Male.....	-	1	3	-	1	2	3	1	1	1	-	-
Female.....	1	-	1	1	3	1	-	2	1	-	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
New Hampshire												
All races, both sexes.....	1	1	5	6	13	17	36	31	21	7	2	-
Male.....	1	1	3	3	6	9	16	17	8	1	-	-
Female.....	-	-	2	3	7	8	20	14	13	6	2	-
White, both sexes.....	1	1	5	6	13	17	35	31	21	7	2	-
Male.....	1	1	3	3	6	9	16	17	8	1	-	-
Female.....	-	-	2	3	7	8	19	14	13	6	2	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
New Jersey												
All races, both sexes.....	43	53	89	91	133	236	309	316	228	61	11	-
Male.....	22	25	52	59	71	108	172	153	96	17	4	-
Female.....	21	28	37	32	62	128	137	163	132	44	7	-
White, both sexes.....	25	29	48	61	91	187	275	282	213	54	9	-
Male.....	12	17	29	44	51	92	157	141	88	15	3	-
Female.....	13	12	19	17	40	95	118	141	125	39	6	-
Black, both sexes.....	16	21	38	29	37	40	29	28	9	7	1	-
Male.....	8	6	22	14	16	12	11	9	3	2	-	-
Female.....	8	15	16	15	21	28	18	19	6	5	1	-
American Indian 1/, both sexes.....	-	-	1	-	1	1	-	-	-	-	-	-
Male.....	-	-	-	-	1	-	-	-	-	-	-	-
Female.....	-	-	1	-	1	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.	2	3	2	1	4	8	5	6	6	-	1	-
Male.....	2	2	1	1	4	3	4	3	5	-	1	-
Female.....	-	1	1	-	-	5	1	3	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Renal failure (N17-N19)

New Mexico

All races, both sexes.....	6	12	11	17	27	29	53	38	11	9	3	-
Male.....	3	9	8	9	13	17	25	18	10	6	2	-
Female.....	3	3	3	8	14	12	28	20	1	3	1	-
White, both sexes.....	5	10	9	14	23	26	46	37	11	8	3	-
Male.....	2	8	6	8	12	15	23	17	10	6	2	-
Female.....	3	2	3	6	11	11	23	20	1	2	1	-
Black, both sexes.....	-	1	-	1	-	1	3	-	-	-	-	-
Male.....	-	1	-	1	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	2	-	-	-	-	-
American Indian 1/, both sexes.....	1	1	2	2	4	2	4	1	-	1	-	-
Male.....	1	-	2	-	1	1	1	1	-	-	-	-
Female.....	-	1	-	2	3	1	3	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	44	81	91	136	182	314	435	467	280	106	29	-
Male.....	26	51	47	67	104	169	217	222	110	27	8	-
Female.....	18	30	44	69	78	145	218	245	170	79	21	-
White, both sexes.....	29	56	63	105	141	266	387	430	254	95	25	-
Male.....	17	33	33	57	86	145	193	207	104	27	4	-
Female.....	12	23	30	48	55	121	194	223	150	68	21	-
Black, both sexes.....	13	21	27	28	35	41	38	34	23	10	3	-
Male.....	7	15	14	8	14	23	17	14	5	-	3	-
Female.....	6	6	13	20	21	18	21	20	18	10	-	-
American Indian 1/, both sexes.....	2	2	-	2	1	-	-	1	-	-	-	-
Male.....	2	1	-	2	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	1	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	1	1	5	7	10	2	3	1	1	-
Male.....	-	2	-	-	4	1	7	1	1	-	1	-
Female.....	-	-	1	1	1	6	3	1	2	1	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Oregon												
All races, both sexes.....	11	16	14	20	34	43	69	87	49	13	4	-
Male.....	9	7	5	7	17	23	36	40	15	3	3	-
Female.....	2	9	9	13	17	20	33	47	34	10	1	-
White, both sexes.....	7	12	12	18	32	43	68	83	47	13	4	-
Male.....	6	5	4	7	17	23	35	37	15	3	3	-
Female.....	1	7	8	11	15	20	33	46	32	10	1	-
Black, both sexes.....	3	2	-	-	1	-	-	3	-	-	-	-
Male.....	2	1	-	-	-	-	-	2	-	-	-	-
Female.....	1	1	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	1	1	1	1	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	1	1	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	1	1	-	1	1	2	-	-	-
Male.....	-	1	1	-	-	-	1	1	-	-	-	-
Female.....	-	-	-	1	1	-	-	-	2	-	-	-
Pennsylvania												
All races, both sexes.....	47	78	109	171	268	426	579	544	359	115	20	-
Male.....	27	50	58	93	122	216	259	241	122	29	5	-
Female.....	20	28	51	78	146	210	320	303	237	86	15	-
White, both sexes.....	34	56	86	131	224	371	525	507	332	103	18	-
Male.....	17	34	45	67	107	196	239	224	112	26	4	-
Female.....	17	22	41	64	117	175	286	283	220	77	14	-
Black, both sexes.....	13	22	22	38	44	52	50	36	27	12	2	-
Male.....	10	16	12	24	15	18	16	17	10	3	1	-
Female.....	3	6	10	14	29	34	34	19	17	9	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	2	-	2	4	1	-	-	-	-
Male.....	-	-	1	2	-	1	4	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Texas												
All races, both sexes.....	124	154	198	257	352	418	469	457	308	108	25	-
Male.....	68	78	104	140	176	216	218	211	129	31	3	-
Female.....	56	76	94	117	176	202	251	246	179	77	22	-
White, both sexes.....	84	115	141	190	282	350	406	391	276	99	23	-
Male.....	51	58	71	110	141	187	194	190	117	29	3	-
Female.....	33	57	70	80	141	163	212	201	159	70	20	-
Black, both sexes.....	38	34	51	60	64	60	56	57	29	9	1	-
Male.....	15	16	28	27	29	25	21	17	10	2	-	-
Female.....	23	18	23	33	35	35	35	40	19	7	1	-
American Indian 1/, both sexes.....	-	2	1	-	1	-	-	2	-	-	-	-
Male.....	-	2	1	-	1	-	-	2	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	3	5	7	5	8	7	7	3	-	1	-
Male.....	2	2	4	3	5	4	3	2	2	-	-	-
Female.....	-	1	1	4	-	4	4	5	1	-	1	-
Utah												
All races, both sexes.....	3	11	9	8	26	29	39	37	25	10	1	-
Male.....	2	7	3	2	13	16	22	16	8	7	-	-
Female.....	1	4	6	6	13	13	17	21	17	3	1	-
White, both sexes.....	2	10	9	7	25	27	39	37	23	10	1	-
Male.....	1	7	3	2	13	15	22	16	8	7	-	-
Female.....	1	3	6	5	12	12	17	21	15	3	1	-
Black, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	-	1	-	2	-	-	2	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	1	-	1	-	1	-	-	2	-	-	-

## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
Renal failure (N17-N19)												
Vermont												
All races, both sexes.....	-	3	1	2	7	6	6	14	7	4	-	-
Male.....	-	3	1	-	4	4	4	5	1	-	-	-
Female.....	-	-	-	2	3	2	2	9	6	4	-	-
White, both sexes.....	-	3	1	2	7	6	6	14	7	4	-	-
Male.....	-	3	1	-	4	4	4	5	1	-	-	-
Female.....	-	-	-	2	3	2	2	9	6	4	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Virginia												
All races, both sexes.....	38	66	76	114	149	189	232	248	139	43	6	-
Male.....	21	39	37	57	74	105	113	105	62	9	-	-
Female.....	17	27	39	57	75	84	119	143	77	34	6	-
White, both sexes.....	17	30	40	71	93	132	180	180	111	33	4	-
Male.....	11	22	24	39	44	77	87	81	54	7	-	-
Female.....	6	8	16	32	49	55	93	99	57	26	4	-
Black, both sexes.....	21	36	35	41	48	54	47	64	26	10	2	-
Male.....	10	17	12	17	26	28	25	23	8	2	-	-
Female.....	11	19	23	24	22	26	22	41	18	8	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	2	8	3	5	4	2	-	-	-
Male.....	-	-	1	1	4	-	1	1	-	-	-	-
Female.....	-	-	-	1	4	3	4	3	2	-	-	-















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Urolithiasis (N20-N23)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Urolithiasis (N20-N23)

Minnesota

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Urolithiasis (N20-N23)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Urolithiasis (N20-N23)

Wisconsin

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Urolithiasis (N20-N23)

Arkansas

All races, both sexes.....	-	-	-	-	-	1	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	1	-	-	-
White, both sexes.....	-	-	-	-	-	1	-	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	-	3	2	3	5	5	8	6	4	-	1	-
Male.....	-	-	1	1	1	2	3	2	3	-	-	-
Female.....	-	3	1	2	4	3	5	4	1	-	1	-
White, both sexes.....	-	3	2	2	3	3	6	5	3	-	1	-
Male.....	-	-	1	1	-	2	2	1	3	-	-	-
Female.....	-	3	1	1	3	1	4	4	-	-	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	2	2	2	1	-	-	-	-
Male.....	-	-	-	-	1	-	1	1	-	-	-	-
Female.....	-	-	-	1	1	2	1	-	-	-	-	-





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
---	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Urolithiasis (N20-N23)

New Mexico

All races, both sexes.....	1	-	-	-	-	-	-	2	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-
White, both sexes.....	1	-	-	-	-	-	-	2	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	1	-	-	-	-	-	-	1	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	1	-	3	2	3	11	5	4	2	-	1	-
Male.....	-	-	-	-	-	6	3	1	2	-	-	-
Female.....	1	-	3	2	3	5	2	3	-	-	1	-
White, both sexes.....	1	-	3	2	2	9	5	4	2	-	1	-
Male.....	-	-	-	-	-	5	3	1	2	-	-	-
Female.....	1	-	3	2	2	4	2	3	-	-	1	-
Black, both sexes.....	-	-	-	-	-	2	-	-	-	-	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	-	-	-	-	-















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Urolithiasis (N20-N23)

Vermont

All races, both sexes.....	-	1	-	-	-	-	1	-	1	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	1	-	-	-
White, both sexes.....	-	1	-	-	-	-	1	-	1	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	-	-	1	-	-	-	3	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	3	1	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	3	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	3	-	-	-	-	-
Black, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified disorders of kidney (N25-N27)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified disorders of kidney (N25-N27)

Kentucky

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified disorders of kidney (N25-N27)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified disorders of kidney (N25-N27)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified disorders of kidney (N25-N27)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified disorders of kidney (N25-N27)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified disorders of kidney (N25-N27)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other and unspecified disorders of kidney (N25-N27)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified disorders of kidney (N25-N27)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified disorders of kidney (N25-N27)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified disorders of kidney (N25-N27)

Minnesota

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified disorders of kidney (N25-N27)

Nebraska

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified disorders of kidney (N25-N27)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified disorders of kidney (N25-N27)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified disorders of kidney (N25-N27)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Urinary tract infection, site not specified (N39.0)												
Arkansas												
All races, both sexes.....	161	1	-	-	-	-	-	-	-	1	-	-
Male.....	55	-	-	-	-	-	-	-	-	1	-	-
Female.....	106	1	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	137	1	-	-	-	-	-	-	-	1	-	-
Male.....	46	-	-	-	-	-	-	-	-	1	-	-
Female.....	91	1	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	24	-	-	-	-	-	-	-	-	-	-	-
Male.....	9	-	-	-	-	-	-	-	-	-	-	-
Female.....	15	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	1,527	-	1	-	-	1	1	2	-	3	5	10
Male.....	522	-	-	-	-	-	1	-	-	1	3	4
Female.....	1,005	-	1	-	-	1	-	2	-	2	2	6
White, both sexes.....	1,304	-	1	-	-	-	1	2	-	3	5	7
Male.....	452	-	-	-	-	-	1	-	-	1	3	3
Female.....	852	-	1	-	-	-	-	2	-	2	2	4
Black, both sexes.....	116	-	-	-	-	-	-	-	-	-	-	3
Male.....	33	-	-	-	-	-	-	-	-	-	-	1
Female.....	83	-	-	-	-	-	-	-	-	-	-	2
American Indian 1/, both sexes.....	7	-	-	-	-	-	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	6	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	100	-	-	-	-	1	-	-	-	-	-	-
Male.....	36	-	-	-	-	-	-	-	-	-	-	-
Female.....	64	-	-	-	-	1	-	-	-	-	-	-



























































## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

## Urinary tract infection, site not specified (N39.0)

## Arkansas

All races, both sexes.....	1	1	2	4	17	14	40	34	24	19	3	-
Male.....	-	1	1	3	8	3	14	13	5	6	-	-
Female.....	1	-	1	1	9	11	26	21	19	13	3	-
White, both sexes.....	1	1	-	3	14	14	34	28	19	19	2	-
Male.....	-	1	-	2	6	3	10	13	4	6	-	-
Female.....	1	-	-	1	8	11	24	15	15	13	2	-
Black, both sexes.....	-	-	2	1	3	-	6	6	5	-	1	-
Male.....	-	-	1	1	2	-	4	-	1	-	-	-
Female.....	-	-	1	-	1	-	2	6	4	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

## California

All races, both sexes.....	25	33	38	65	90	177	295	360	288	113	20	-
Male.....	12	15	12	24	33	58	111	120	97	29	2	-
Female.....	13	18	26	41	57	119	184	240	191	84	18	-
White, both sexes.....	18	24	31	52	74	152	260	310	249	99	16	-
Male.....	10	11	10	20	27	50	103	105	81	26	1	-
Female.....	8	13	21	32	47	102	157	205	168	73	15	-
Black, both sexes.....	5	7	3	7	11	15	18	21	17	8	1	-
Male.....	1	2	1	4	3	5	4	5	6	1	-	-
Female.....	4	5	2	3	8	10	14	16	11	7	1	-
American Indian 1/, both sexes.....	-	-	2	-	-	1	1	-	2	1	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	2	-	-	1	1	-	1	1	-	-
Asian or Pacific Islander 2/, both sexes.....	2	2	2	6	5	9	16	29	20	5	3	-
Male.....	1	2	1	-	3	3	4	10	9	2	1	-
Female.....	1	-	1	6	2	6	12	19	11	3	2	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Urinary tract infection, site not specified (N39.0)

Colorado

All races, both sexes.....	1	4	5	6	9	18	40	49	34	11	2	-
Male.....	-	2	2	2	2	7	8	18	16	1	-	-
Female.....	1	2	3	4	7	11	32	31	18	10	2	-
White, both sexes.....	1	4	5	6	9	16	39	49	33	11	2	-
Male.....	-	2	2	2	2	7	8	18	16	1	-	-
Female.....	1	2	3	4	7	9	31	31	17	10	2	-
Black, both sexes.....	-	-	-	-	-	1	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	1	-	-	-

Connecticut

All races, both sexes.....	2	2	3	3	6	16	30	48	44	21	6	-
Male.....	1	-	2	1	3	8	14	17	11	4	-	-
Female.....	1	2	1	2	3	8	16	31	33	17	6	-
White, both sexes.....	2	2	3	3	5	16	26	44	43	21	6	-
Male.....	1	-	2	1	3	8	10	17	10	4	-	-
Female.....	1	2	1	2	2	8	16	27	33	17	6	-
Black, both sexes.....	-	-	-	-	1	-	3	3	1	-	-	-
Male.....	-	-	-	-	-	-	3	-	1	-	-	-
Female.....	-	-	-	-	1	-	-	3	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Urinary tract infection, site not specified (N39.0)

Kentucky

All races, both sexes.....	3	8	7	11	18	39	38	56	31	8	3	-
Male.....	-	4	3	5	10	10	19	12	6	2	1	-
Female.....	3	4	4	6	8	29	19	44	25	6	2	-
White, both sexes.....	3	8	7	8	18	38	35	54	29	8	2	-
Male.....	-	4	3	2	10	10	18	12	5	2	1	-
Female.....	3	4	4	6	8	28	17	42	24	6	1	-
Black, both sexes.....	-	-	-	3	-	1	3	2	2	-	1	-
Male.....	-	-	-	3	-	-	1	-	1	-	-	-
Female.....	-	-	-	-	-	1	2	2	1	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Louisiana

All races, both sexes.....	3	1	7	10	15	30	51	63	36	12	6	-
Male.....	1	1	4	3	6	11	10	19	15	3	-	-
Female.....	2	-	3	7	9	19	41	44	21	9	6	-
White, both sexes.....	2	1	2	7	9	26	36	48	30	10	2	-
Male.....	-	1	-	2	4	11	7	15	14	2	-	-
Female.....	2	-	2	5	5	15	29	33	16	8	2	-
Black, both sexes.....	1	-	5	3	6	3	13	14	6	2	4	-
Male.....	1	-	4	1	2	-	2	3	1	1	-	-
Female.....	-	-	1	2	4	3	11	11	5	1	4	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	1	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	1	1	-	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Urinary tract infection, site not specified (N39.0)

Maine

All races, both sexes.....	1	-	-	1	2	10	14	24	9	4	3	-
Male.....	-	-	-	-	2	3	5	8	4	2	-	-
Female.....	1	-	-	1	-	7	9	16	5	2	3	-
White, both sexes.....	1	-	-	1	2	10	13	24	8	4	3	-
Male.....	-	-	-	-	2	3	5	8	3	2	-	-
Female.....	1	-	-	1	-	7	8	16	5	2	3	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-

Maryland

All races, both sexes.....	3	5	4	12	33	42	72	90	49	22	2	-
Male.....	3	3	1	2	11	17	20	23	13	1	-	-
Female.....	-	2	3	10	22	25	52	67	36	21	2	-
White, both sexes.....	2	3	2	7	21	30	54	71	44	17	2	-
Male.....	2	2	-	2	6	10	16	20	10	1	-	-
Female.....	-	1	2	5	15	20	38	51	34	16	2	-
Black, both sexes.....	1	2	2	5	11	12	18	16	4	4	-	-
Male.....	1	1	1	-	4	7	4	3	2	-	-	-
Female.....	-	1	1	5	7	5	14	13	2	4	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	-	-	3	1	1	-	-
Male.....	-	-	-	-	1	-	-	-	1	-	-	-
Female.....	-	-	-	-	-	-	-	3	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Urinary tract infection, site not specified (N39.0)

Massachusetts

All races, both sexes.....	2	3	6	10	17	42	68	101	72	27	8	-
Male.....	-	2	-	6	6	13	23	37	17	8	2	-
Female.....	2	1	6	4	11	29	45	64	55	19	6	-
White, both sexes.....	2	3	6	10	17	41	66	96	69	25	7	-
Male.....	-	2	-	6	6	13	22	35	15	8	2	-
Female.....	2	1	6	4	11	28	44	61	54	17	5	-
Black, both sexes.....	-	-	-	-	-	1	1	4	2	2	1	-
Male.....	-	-	-	-	-	-	1	2	2	-	-	-
Female.....	-	-	-	-	-	1	-	2	-	2	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	1	1	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	1	-	-	-

Michigan

All races, both sexes.....	6	9	13	22	21	71	119	109	77	31	3	-
Male.....	2	3	6	14	10	28	37	25	20	7	1	-
Female.....	4	6	7	8	11	43	82	84	57	24	2	-
White, both sexes.....	5	7	12	17	21	64	110	99	70	27	2	-
Male.....	2	1	5	10	10	24	32	22	19	7	-	-
Female.....	3	6	7	7	11	40	78	77	51	20	2	-
Black, both sexes.....	1	1	1	5	-	6	8	9	7	3	-	-
Male.....	-	1	1	4	-	3	4	3	1	-	-	-
Female.....	1	-	-	1	-	3	4	6	6	3	-	-
American Indian 1/, both sexes.....	-	1	-	-	-	1	1	-	-	1	-	-
Male.....	-	1	-	-	-	1	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	1	-	-	1	-
Male.....	-	-	-	-	-	-	-	-	-	-	1	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Urinary tract infection, site not specified (N39.0)

Minnesota

All races, both sexes.....	2	2	3	6	1	15	31	50	28	10	3	-
Male.....	2	-	2	2	-	10	15	15	7	2	-	-
Female.....	-	2	1	4	1	5	16	35	21	8	3	-
White, both sexes.....	2	2	3	6	1	14	31	50	28	9	3	-
Male.....	2	-	2	2	-	9	15	15	7	2	-	-
Female.....	-	2	1	4	1	5	16	35	21	7	3	-
Black, both sexes.....	-	-	-	-	-	1	-	-	-	1	-	-
Male.....	-	-	-	-	-	1	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Mississippi

All races, both sexes.....	1	3	12	11	16	24	40	43	27	14	4	-
Male.....	-	1	4	6	5	13	20	8	9	2	-	-
Female.....	1	2	8	5	11	11	20	35	18	12	4	-
White, both sexes.....	1	1	8	5	10	15	24	30	21	8	2	-
Male.....	-	-	3	4	2	10	13	6	7	1	-	-
Female.....	1	1	5	1	8	5	11	24	14	7	2	-
Black, both sexes.....	-	2	4	6	6	9	16	13	5	6	2	-
Male.....	-	1	1	2	3	3	7	2	2	1	-	-
Female.....	-	1	3	4	3	6	9	11	3	5	2	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	1	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Urinary tract infection, site not specified (N39.0)

New Hampshire

All races, both sexes.....	-	-	1	1	2	7	8	19	5	4	-	-
Male.....	-	-	1	1	-	1	4	5	2	2	-	-
Female.....	-	-	-	-	2	6	4	14	3	2	-	-
White, both sexes.....	-	-	1	1	2	7	8	19	5	4	-	-
Male.....	-	-	1	1	-	1	4	5	2	2	-	-
Female.....	-	-	-	-	2	6	4	14	3	2	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	2	6	7	9	18	45	77	103	86	34	-	-
Male.....	-	2	5	3	9	17	23	32	22	6	-	-
Female.....	2	4	2	6	9	28	54	71	64	28	-	-
White, both sexes.....	2	5	5	8	12	37	72	96	77	31	-	-
Male.....	-	2	4	2	5	15	22	31	22	6	-	-
Female.....	2	3	1	6	7	22	50	65	55	25	-	-
Black, both sexes.....	-	1	1	1	5	8	4	7	7	3	-	-
Male.....	-	-	-	1	3	2	1	1	-	-	-	-
Female.....	-	1	1	-	2	6	3	6	7	3	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	1	-	1	-	2	-	-	-
Male.....	-	-	1	-	1	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	2	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Urinary tract infection, site not specified (N39.0)

New Mexico

All races, both sexes.....	-	-	3	2	4	8	16	18	14	7	-	-
Male.....	-	-	1	1	1	3	5	4	6	2	-	-
Female.....	-	-	2	1	3	5	11	14	8	5	-	-
White, both sexes.....	-	-	2	2	4	8	15	16	14	6	-	-
Male.....	-	-	1	1	1	3	5	3	6	2	-	-
Female.....	-	-	1	1	3	5	10	13	8	4	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	2	-	1	-	-
Male.....	-	-	-	-	-	-	-	1	-	-	-	-
Female.....	-	-	-	-	-	-	1	1	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	10	22	36	46	53	142	201	284	197	76	24	-
Male.....	5	7	10	15	22	52	68	84	48	18	2	-
Female.....	5	15	26	31	31	90	133	200	149	58	22	-
White, both sexes.....	6	16	24	28	37	113	163	257	174	67	17	-
Male.....	3	5	6	11	15	41	58	76	40	17	1	-
Female.....	3	11	18	17	22	72	105	181	134	50	16	-
Black, both sexes.....	4	6	10	16	16	24	32	25	21	9	6	-
Male.....	2	2	4	2	7	8	9	7	7	1	1	-
Female.....	2	4	6	14	9	16	23	18	14	8	5	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	1	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	2	2	-	5	5	2	2	-	1	-
Male.....	-	-	-	2	-	3	1	1	1	-	-	-
Female.....	-	-	2	-	-	2	4	1	1	-	1	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Urinary tract infection, site not specified (N39.0)

Ohio

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Urinary tract infection, site not specified (N39.0)

Oregon

All races, both sexes.....	-	3	2	4	9	26	30	31	40	21	3	-
Male.....	-	-	1	1	3	8	9	9	6	2	2	-
Female.....	-	3	1	3	6	18	21	22	34	19	1	-
White, both sexes.....	-	2	1	4	9	26	30	30	40	21	3	-
Male.....	-	-	1	1	3	8	9	9	6	2	2	-
Female.....	-	2	-	3	6	18	21	21	34	19	1	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	1	1	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	1	-	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	13	9	15	21	34	81	125	178	120	43	7	-
Male.....	3	4	8	7	17	36	41	55	35	8	1	-
Female.....	10	5	7	14	17	45	84	123	85	35	6	-
White, both sexes.....	10	9	12	19	31	79	112	165	113	41	7	-
Male.....	3	4	5	6	16	36	37	50	35	8	1	-
Female.....	7	5	7	13	15	43	75	115	78	33	6	-
Black, both sexes.....	3	-	2	2	3	2	13	12	7	2	-	-
Male.....	-	-	2	1	1	-	4	5	-	-	-	-
Female.....	3	-	-	1	2	2	9	7	7	2	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	1	-	-	-	-	1	-	-	-	-
Male.....	-	-	1	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Urinary tract infection, site not specified (N39.0)

Texas

All races, both sexes.....	13	20	33	35	61	113	186	190	155	60	17	-
Male.....	7	8	16	9	23	38	68	71	40	12	4	-
Female.....	6	12	17	26	38	75	118	119	115	48	13	-
White, both sexes.....	9	15	28	26	52	98	166	167	141	52	13	-
Male.....	6	4	14	6	21	33	57	60	38	9	2	-
Female.....	3	11	14	20	31	65	109	107	103	43	11	-
Black, both sexes.....	4	4	5	9	8	10	16	21	14	8	4	-
Male.....	1	4	2	3	2	5	8	10	2	3	2	-
Female.....	3	-	3	6	6	5	8	11	12	5	2	-
American Indian 1/, both sexes.....	-	1	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	1	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	1	5	4	2	-	-	-	-
Male.....	-	-	-	-	-	-	3	1	-	-	-	-
Female.....	-	-	-	-	1	5	1	1	-	-	-	-

Utah

All races, both sexes.....	-	3	2	7	8	10	13	13	10	10	2	-
Male.....	-	-	1	2	4	-	5	2	2	3	-	-
Female.....	-	3	1	5	4	10	8	11	8	7	2	-
White, both sexes.....	-	3	2	6	8	9	12	12	10	9	2	-
Male.....	-	-	1	2	4	-	4	2	2	3	-	-
Female.....	-	3	1	4	4	9	8	10	8	6	2	-
Black, both sexes.....	-	-	-	1	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	1	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	1	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Urinary tract infection, site not specified (N39.0)

Vermont

All races, both sexes.....	-	-	-	2	-	3	4	6	5	-	-	-
Male.....	-	-	-	1	-	1	-	1	1	-	-	-
Female.....	-	-	-	1	-	2	4	5	4	-	-	-
White, both sexes.....	-	-	-	2	-	3	4	6	5	-	-	-
Male.....	-	-	-	1	-	1	-	1	1	-	-	-
Female.....	-	-	-	1	-	2	4	5	4	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	4	7	8	14	20	45	90	82	68	31	7	-
Male.....	1	5	1	5	7	13	29	27	15	5	1	-
Female.....	3	2	7	9	13	32	61	55	53	26	6	-
White, both sexes.....	2	5	6	12	14	32	76	75	57	28	3	-
Male.....	-	3	-	4	6	6	24	23	11	5	1	-
Female.....	2	2	6	8	8	26	52	52	46	23	2	-
Black, both sexes.....	1	2	1	2	6	12	14	6	11	3	4	-
Male.....	-	2	1	1	1	7	5	4	4	-	-	-
Female.....	1	-	-	1	5	5	9	2	7	3	4	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	1	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	1	-	-	1	-	-	-	-	-	-
Male.....	1	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	1	-	-	1	-	-	-	-	-	-





































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of urinary system (N28-N36,N39.1-N39.9)

Arkansas

All races, both sexes.....	1	1	1	1	2	2	2	3	2	-	-	-
Male.....	1	1	1	1	-	2	2	-	2	-	-	-
Female.....	-	-	-	-	2	-	-	3	-	-	-	-
White, both sexes.....	1	-	1	-	1	2	2	2	2	-	-	-
Male.....	1	-	1	-	-	2	2	-	2	-	-	-
Female.....	-	-	-	-	1	-	-	2	-	-	-	-
Black, both sexes.....	-	1	-	1	1	-	-	1	-	-	-	-
Male.....	-	1	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	1	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	3	6	6	11	8	7	21	18	18	3	-	-
Male.....	1	3	4	6	5	5	7	8	7	2	-	-
Female.....	2	3	2	5	3	2	14	10	11	1	-	-
White, both sexes.....	3	3	6	10	6	7	17	16	14	3	-	-
Male.....	1	2	4	6	5	5	6	7	5	2	-	-
Female.....	2	1	2	4	1	2	11	9	9	1	-	-
Black, both sexes.....	-	1	-	-	2	-	2	1	1	-	-	-
Male.....	-	-	-	-	-	-	1	1	1	-	-	-
Female.....	-	1	-	-	2	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	2	-	1	-	-	2	1	3	-	-	-
Male.....	-	1	-	-	-	-	-	-	1	-	-	-
Female.....	-	1	-	1	-	-	2	1	2	-	-	-































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Other diseases of urinary system (N28-N36,N39.1-N39.9)

New Mexico

All races, both sexes.....	-	1	-	-	-	-	3	2	2	-	-	-
Male.....	-	1	-	-	-	-	2	2	1	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	-	-	-
White, both sexes.....	-	1	-	-	-	-	2	2	2	-	-	-
Male.....	-	1	-	-	-	-	1	2	1	-	-	-
Female.....	-	-	-	-	-	-	1	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	1	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	4	3	3	4	3	8	5	14	6	3	-	-
Male.....	3	3	2	2	2	4	1	7	1	2	-	-
Female.....	1	-	1	2	1	4	4	7	5	1	-	-
White, both sexes.....	2	3	3	3	2	5	5	13	5	3	-	-
Male.....	1	3	2	1	1	2	1	7	1	2	-	-
Female.....	1	-	1	2	1	3	4	6	4	1	-	-
Black, both sexes.....	2	-	-	-	1	2	-	1	1	-	-	-
Male.....	2	-	-	-	1	2	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	1	1	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	1	-	1	-	-	-	-	-	-
Male.....	-	-	-	1	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	1	-	-	-	-	-	-





































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Diseases of male genital organs (N40-N50)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diseases of male genital organs (N40-N50)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diseases of male genital organs (N40-N50)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diseases of male genital organs (N40-N50)

Texas

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, White, Black, American Indian 1/, and Asian or Pacific Islander 2/ for both sexes and by gender.

Utah

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, White, Black, American Indian 1/, and Asian or Pacific Islander 2/ for both sexes and by gender.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Diseases of male genital organs (N40-N50)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



































































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Hyperplasia of prostate (N40)

New Hampshire

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, White, Black, American Indian 1/, Asian or Pacific Islander 2/ with Male and Female sub-rows.

New Jersey

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, White, Black, American Indian 1/, Asian or Pacific Islander 2/ with Male and Female sub-rows.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Hyperplasia of prostate (N40)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Hyperplasia of prostate (N40)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























































































































































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Disorders of breast (N60-N64)

Arkansas

All races, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	2	-	-	-	-	-	1	-	-	-	1	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	2	-	-	-	-	-	1	-	-	-	1	-
White, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	1	-	-	-	-	-	-	-	-	-	1	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	-	-	-	-	-	1	-	-	-	-	-
Male.....	-	-	-	-	-	-	-	-	-	-	-	-
Female.....	1	-	-	-	-	-	1	-	-	-	-	-













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex
All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Disorders of breast (N60-N64)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Disorders of breast (N60-N64)

Rhode Island

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Disorders of breast (N60-N64)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Disorders of breast (N60-N64)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Disorders of breast (N60-N64)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Disorders of breast (N60-N64)

Massachusetts

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Michigan

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Disorders of breast (N60-N64)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Disorders of breast (N60-N64)

Washington

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

West Virginia

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Disorders of female genital organs (N70-N98)

Arkansas

All races, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	44	1	-	-	-	-	1	1	1	2	2	2
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	44	1	-	-	-	-	1	1	1	2	2	2
White, both sexes.....	37	-	-	-	-	-	-	1	-	2	2	2
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	37	-	-	-	-	-	-	1	-	2	2	2
Black, both sexes.....	4	1	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	-	-	-	-	-	1	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	-	-	-	-	-	1	-	1	-	-	-









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Disorders of female genital organs (N70-N98)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Disorders of female genital organs (N70-N98)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Disorders of female genital organs (N70-N98)

Arkansas

All races, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	1	5	4	2	4	3	7	4	2	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	5	4	2	4	3	7	4	2	1	1	-
White, both sexes.....	1	5	3	1	3	3	7	4	1	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	5	3	1	3	3	7	4	1	1	1	-
Black, both sexes.....	-	-	1	1	1	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	1	1	1	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	1	-	-	-





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Disorders of female genital organs (N70-N98)

New Mexico

All races, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	3	1	1	5	4	5	4	5	2	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	1	1	5	4	5	4	5	2	-	-	-
White, both sexes.....	2	1	1	5	3	3	4	4	2	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	1	1	5	3	3	4	4	2	-	-	-
Black, both sexes.....	1	-	-	-	1	1	-	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	-	-	-	1	1	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	1	-	-	-	-	-	-



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
---	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Inflammatory diseases of female pelvic organs (N70-N76)

Arkansas

All races, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

California

All races, both sexes.....	17	1	-	-	-	-	1	-	1	1	1	2
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	17	1	-	-	-	-	1	-	1	1	1	2
White, both sexes.....	13	-	-	-	-	-	-	-	-	1	1	2
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	13	-	-	-	-	-	-	-	-	1	1	2
Black, both sexes.....	2	1	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	1	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	-	-	-	-	-	1	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	-	-	-	-	-	1	-	1	-	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Inflammatory diseases of female pelvic organs (N70-N76)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Inflammatory diseases of female pelvic organs (N70-N76)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Inflammatory diseases of female pelvic organs (N70-N76)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Inflammatory diseases of female pelvic organs (N70-N76)

Ohio

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Inflammatory diseases of female pelvic organs (N70-N76)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Inflammatory diseases of female pelvic organs (N70-N76)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Inflammatory diseases of female pelvic organs (N70-N76)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

Delaware

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

Kentucky

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

Maine

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

Massachusetts

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Michigan

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by sex.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

Minnesota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

North Carolina

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

Ohio

Table with 13 columns (Age groups) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Age groups) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

South Dakota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Inflammatory diseases of female pelvic organs (N70-N76)

Vermont

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Noninflammatory disorders of female genital tract (N80-N98)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Noninflammatory disorders of female genital tract (N80-N98)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Noninflammatory disorders of female genital tract (N80-N98)

Missouri

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Noninflammatory disorders of female genital tract (N80-N98)

Nebraska

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns: Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Noninflammatory disorders of female genital tract (N80-N98)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Noninflammatory disorders of female genital tract (N80-N98)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Noninflammatory disorders of female genital tract (N80-N98)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Noninflammatory disorders of female genital tract (N80-N98)

Alaska

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Noninflammatory disorders of female genital tract (N80-N98)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Noninflammatory disorders of female genital tract (N80-N98)

Massachusetts

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Noninflammatory disorders of female genital tract (N80-N98)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	----------------	-----------------------	-------------------

Noninflammatory disorders of female genital tract (N80-N98)

New Mexico

All races, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
White, both sexes.....	-	-	-	-	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	1	-	-	-
Black, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	3	-	1	2	3	4	3	4	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	-	1	2	3	4	3	4	-	-	-	-
White, both sexes.....	2	-	1	2	3	2	3	3	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	-	1	2	3	2	3	3	-	-	-	-
Black, both sexes.....	1	-	-	-	-	1	-	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	-	-	-	-	1	-	1	-	-	-	-
American Indian 1/, both sexes.....	-	-	-	-	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	-	-	-	-	1	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	-	-	-	-	1	-	-	-	-	-	-















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Noninflammatory disorders of female genital tract (N80-N98)

Vermont

Table with 13 columns (Age groups) and 13 rows (Race categories) for Vermont. All cells contain dashes (-).

Virginia

Table with 13 columns (Age groups) and 13 rows (Race categories) for Virginia. Contains numerical values such as 1 and 2.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XV. Pregnancy, childbirth and the puerperium (000-099)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XV. Pregnancy, childbirth and the puerperium (O00-O99)												
Florida												
All races, both sexes.....	52	...	...	...	-	2	12	11	10	15	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	52	...	...	...	-	2	12	11	10	15	2	-
White, both sexes.....	28	...	...	...	-	-	9	2	6	10	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	28	...	...	...	-	-	9	2	6	10	1	-
Black, both sexes.....	23	...	...	...	-	2	3	8	4	5	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	23	...	...	...	-	2	3	8	4	5	1	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	1	-	-	-	-
Georgia												
All races, both sexes.....	22	...	...	...	-	3	4	2	8	3	-	2
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	22	...	...	...	-	3	4	2	8	3	-	2
White, both sexes.....	7	...	...	...	-	1	-	-	5	-	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	7	...	...	...	-	1	-	-	5	-	-	1
Black, both sexes.....	14	...	...	...	-	2	4	2	3	2	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	14	...	...	...	-	2	4	2	3	2	-	1
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>XV. Pregnancy, childbirth and the puerperium (O00-O99)</b>												
<b>New Hampshire</b>												
All races, both sexes.....	5	...	...	...	-	-	2	1	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	5	...	...	...	-	-	2	1	1	1	-	-
White, both sexes.....	5	...	...	...	-	-	2	1	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	5	...	...	...	-	-	2	1	1	1	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
<b>New Jersey</b>												
All races, both sexes.....	36	...	...	...	-	1	7	3	11	10	3	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	36	...	...	...	-	1	7	3	11	10	3	-
White, both sexes.....	22	...	...	...	-	-	5	2	7	5	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	22	...	...	...	-	-	5	2	7	5	2	-
Black, both sexes.....	12	...	...	...	-	1	2	1	3	4	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	12	...	...	...	-	1	2	1	3	4	1	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	-	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
<b>XV. Pregnancy, childbirth and the puerperium (O00-O99)</b>												
<b>New Mexico</b>												
All races, both sexes.....	10	...	...	...	-	1	-	3	1	3	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	10	...	...	...	-	1	-	3	1	3	2	-
White, both sexes.....	9	...	...	...	-	1	-	3	1	2	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	9	...	...	...	-	1	-	3	1	2	2	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
<b>New York</b>												
All races, both sexes.....	51	...	...	...	-	1	6	11	17	7	6	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	51	...	...	...	-	1	6	11	17	7	6	1
White, both sexes.....	18	...	...	...	-	1	1	2	8	3	3	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	18	...	...	...	-	1	1	2	8	3	3	-
Black, both sexes.....	28	...	...	...	-	-	5	8	8	3	1	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	28	...	...	...	-	-	5	8	8	3	1	1
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	...	...	...	-	-	-	1	1	1	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	5	...	...	...	-	-	-	1	1	1	2	-





## Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
XV. Pregnancy, childbirth and the puerperium (O00-O99)												
Oregon												
All races, both sexes.....	10	...	...	...	-	-	2	-	3	2	1	2
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	10	...	...	...	-	-	2	-	3	2	1	2
White, both sexes.....	8	...	...	...	-	-	2	-	1	2	1	2
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	8	...	...	...	-	-	2	-	1	2	1	2
Black, both sexes.....	2	...	...	...	-	-	-	-	2	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	2	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes.....	40	...	...	...	-	2	6	8	8	11	3	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	40	...	...	...	-	2	6	8	8	11	3	1
White, both sexes.....	24	...	...	...	-	1	2	5	5	8	1	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	24	...	...	...	-	1	2	5	5	8	1	1
Black, both sexes.....	14	...	...	...	-	1	4	2	2	3	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	14	...	...	...	-	1	4	2	2	3	2	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	1	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	1	1	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XV. Pregnancy, childbirth and the puerperium (O00-O99)

Texas

All races, both sexes.....	80	...	...	...	-	8	2	16	12	16	9	10
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	80	...	...	...	-	8	2	16	12	16	9	10
White, both sexes.....	66	...	...	...	-	6	2	16	9	11	6	9
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	66	...	...	...	-	6	2	16	9	11	6	9
Black, both sexes.....	12	...	...	...	-	2	-	-	3	4	3	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	12	...	...	...	-	2	-	-	3	4	3	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	-	-	1	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	-	1	-	1

Utah

All races, both sexes.....	11	...	...	...	-	1	1	2	2	2	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	11	...	...	...	-	1	1	2	2	2	2	-
White, both sexes.....	8	...	...	...	-	1	1	1	2	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	8	...	...	...	-	1	1	1	2	1	1	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	...	-	-	-	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	1	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	-	-	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	-	1	1	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

XV. Pregnancy, childbirth and the puerperium (O00-O99)

Vermont

All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	15	...	...	...	-	-	1	7	4	2	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	15	...	...	...	-	-	1	7	4	2	-	1
White, both sexes.....	7	...	...	...	-	-	-	2	3	1	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	7	...	...	...	-	-	-	2	3	1	-	1
Black, both sexes.....	5	...	...	...	-	-	1	4	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	5	...	...	...	-	-	1	4	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	...	...	...	-	-	-	1	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	-	1	1	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XV. Pregnancy, childbirth and the puerperium (000-099)

Washington

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XV. Pregnancy, childbirth and the puerperium (000-099)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XV. Pregnancy, childbirth and the puerperium (000-099)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XV. Pregnancy, childbirth and the puerperium (000-099)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XV. Pregnancy, childbirth and the puerperium (000-099)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XV. Pregnancy, childbirth and the puerperium (000-099)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XV. Pregnancy, childbirth and the puerperium (000-099)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XV. Pregnancy, childbirth and the puerperium (000-099)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

XV. Pregnancy, childbirth and the puerperium (000-099)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XV. Pregnancy, childbirth and the puerperium (000-099)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Pregnancy with abortive outcome (000-007)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Pregnancy with abortive outcome (000-007)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Pregnancy with abortive outcome (000-007)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Pregnancy with abortive outcome (000-007)

New Hampshire

All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	2	...	...	...	-	-	-	-	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	1	1	-	-
White, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Pregnancy with abortive outcome (000-007)

New Mexico

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Pregnancy with abortive outcome (000-007)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Pregnancy with abortive outcome (000-007)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Pregnancy with abortive outcome (000-007)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Pregnancy with abortive outcome (000-007)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Ectopic pregnancy (O00)

United States

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for United States: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Alabama

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alabama: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Ectopic pregnancy (000)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Ectopic pregnancy (000)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Ectopic pregnancy (000)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows (All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female)

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows (All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female)

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Ectopic pregnancy (O00)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Ectopic pregnancy (000)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Ectopic pregnancy (O00)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Ectopic pregnancy (000)

Massachusetts

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Ectopic pregnancy (000)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Ectopic pregnancy (O00)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Ectopic pregnancy (000)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Ectopic pregnancy (O00)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Ectopic pregnancy (000)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Ectopic pregnancy (000)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Ectopic pregnancy (000)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Ectopic pregnancy (000)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Spontaneous abortion (O03)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows (All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female)

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows (All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female)

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Spontaneous abortion (O03)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Spontaneous abortion (O03)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Spontaneous abortion (O03)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Spontaneous abortion (O03)

Massachusetts

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Spontaneous abortion (O03)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Spontaneous abortion (O03)												
New Hampshire												
All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
New Jersey												
All races, both sexes.....	2	...	...	...	-	-	-	-	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	1	1	-	-
White, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Spontaneous abortion (O03)

New Mexico

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Spontaneous abortion (O03)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Spontaneous abortion (O03)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Spontaneous abortion (O03)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Spontaneous abortion (O03)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Spontaneous abortion (O03)

Colorado

Table with 13 columns (Age groups) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Age groups) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Spontaneous abortion (O03)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Spontaneous abortion (O03)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Spontaneous abortion (O03)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Spontaneous abortion (O03)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Spontaneous abortion (O03)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Spontaneous abortion (O03)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other abortion (O05)

Arkansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other abortion (O05)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other abortion (O05)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other abortion (O05)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other abortion (O05)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other abortion (O05)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Other abortion (O05)

New Mexico

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Mexico categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New York categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other abortion (O05)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other abortion (O05)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other abortion (O05)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other abortion (O05)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

Colorado

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

Hawaii

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified pregnancy with abortive outcome (001-002,006-007)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other direct obstetric deaths (O10-O92)

Arkansas

All races, both sexes.....	3	...	...	...	-	-	2	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	2	1	-	-	-	-
White, both sexes.....	3	...	...	...	-	-	2	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	2	1	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

California

All races, both sexes.....	33	...	...	...	-	-	6	6	10	6	4	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	33	...	...	...	-	-	6	6	10	6	4	1
White, both sexes.....	23	...	...	...	-	-	5	3	7	5	3	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	23	...	...	...	-	-	5	3	7	5	3	-
Black, both sexes.....	6	...	...	...	-	-	1	3	1	-	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	6	...	...	...	-	-	1	3	1	-	-	1
American Indian 1/, both sexes.....	1	...	...	...	-	-	-	-	-	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	3	...	...	...	-	-	-	-	2	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	-	-	2	1	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other direct obstetric deaths (O10-O92)												
Florida												
All races, both sexes.....	31	...	...	...	-	1	8	4	7	10	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	31	...	...	...	-	1	8	4	7	10	1	-
White, both sexes.....	20	...	...	...	-	-	7	1	5	7	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	20	...	...	...	-	-	7	1	5	7	-	-
Black, both sexes.....	11	...	...	...	-	1	1	3	2	3	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	11	...	...	...	-	1	1	3	2	3	1	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes.....	14	...	...	...	-	2	3	2	4	2	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	14	...	...	...	-	2	3	2	4	2	-	1
White, both sexes.....	6	...	...	...	-	1	-	-	4	-	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	6	...	...	...	-	1	-	-	4	-	-	1
Black, both sexes.....	7	...	...	...	-	1	3	2	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	7	...	...	...	-	1	3	2	-	1	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other direct obstetric deaths (O10-O92)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other direct obstetric deaths (O10-O92)

New Hampshire

All races, both sexes.....	2	...	...	...	-	-	1	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	1	-	-	1	-	-
White, both sexes.....	2	...	...	...	-	-	1	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	1	-	-	1	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	14	...	...	...	-	1	4	1	2	4	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	14	...	...	...	-	1	4	1	2	4	1	-
White, both sexes.....	8	...	...	...	-	-	3	1	-	2	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	8	...	...	...	-	-	3	1	-	2	1	-
Black, both sexes.....	5	...	...	...	-	1	1	-	2	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	5	...	...	...	-	1	1	-	2	1	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other direct obstetric deaths (O10-O92)

New Mexico

All races, both sexes.....	6	...	...	...	-	1	-	3	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	6	...	...	...	-	1	-	3	1	1	-	-
White, both sexes.....	6	...	...	...	-	1	-	3	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	6	...	...	...	-	1	-	3	1	1	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	31	...	...	...	-	1	3	6	12	3	4	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	31	...	...	...	-	1	3	6	12	3	4	1
White, both sexes.....	14	...	...	...	-	1	-	1	7	2	3	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	14	...	...	...	-	1	-	1	7	2	3	-
Black, both sexes.....	15	...	...	...	-	-	3	4	5	-	1	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	15	...	...	...	-	-	3	4	5	-	1	1
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	1	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	1	-	1	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other direct obstetric deaths (O10-O92)

Oregon

All races, both sexes.....	4	...	...	...	-	-	1	-	-	1	1	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	...	...	...	-	-	1	-	-	1	1	1
White, both sexes.....	4	...	...	...	-	-	1	-	-	1	1	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	...	...	...	-	-	1	-	-	1	1	1
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	16	...	...	...	-	-	4	2	3	4	2	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	16	...	...	...	-	-	4	2	3	4	2	1
White, both sexes.....	7	...	...	...	-	-	2	-	1	3	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	7	...	...	...	-	-	2	-	1	3	-	1
Black, both sexes.....	7	...	...	...	-	-	2	1	1	1	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	7	...	...	...	-	-	2	1	1	1	2	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	1	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	1	1	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Other direct obstetric deaths (O10-O92)												
Texas												
All races, both sexes.....	38	...	...	...	-	2	-	8	6	9	3	7
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	38	...	...	...	-	2	-	8	6	9	3	7
White, both sexes.....	31	...	...	...	-	2	-	8	3	6	2	7
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	31	...	...	...	-	2	-	8	3	6	2	7
Black, both sexes.....	6	...	...	...	-	-	-	-	3	2	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	6	...	...	...	-	-	-	-	3	2	1	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-
Utah												
All races, both sexes.....	5	...	...	...	-	1	-	-	2	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	5	...	...	...	-	1	-	-	2	1	1	-
White, both sexes.....	3	...	...	...	-	1	-	-	2	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	1	-	-	2	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	-	-	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	-	1	1	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other direct obstetric deaths (O10-O92)

Vermont

All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	9	...	...	...	-	-	-	5	2	1	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	9	...	...	...	-	-	-	5	2	1	-	1
White, both sexes.....	4	...	...	...	-	-	-	2	1	-	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	...	...	...	-	-	-	2	1	-	-	1
Black, both sexes.....	2	...	...	...	-	-	-	2	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	2	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	...	...	...	-	-	-	1	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	-	1	1	1	-	-

























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other direct obstetric deaths (O10-O92)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Eclampsia and pre-eclampsia (O11,O13-O16)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Eclampsia and pre-eclampsia (O11,O13-O16)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Eclampsia and pre-eclampsia (O11,O13-O16)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Eclampsia and pre-eclampsia (O11,O13-O16)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows (All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female)

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows (All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female)











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Eclampsia and pre-eclampsia (O11,O13-O16)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Eclampsia and pre-eclampsia (O11,O13-O16)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Eclampsia and pre-eclampsia (O11,O13-O16)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Eclampsia and pre-eclampsia (O11,O13-O16)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Eclampsia and pre-eclampsia (O11,O13-O16)

Iowa

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Iowa. All cells contain dashes (-).

Kansas

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Kansas. All cells contain dashes (-).









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Eclampsia and pre-eclampsia (O11,O13-O16)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Eclampsia and pre-eclampsia (O11,O13-O16)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

New Hampshire

All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

New Jersey

All races, both sexes.....	2	...	...	...	-	-	-	1	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	1	-	1	-	-
White, both sexes.....	1	...	...	...	-	-	-	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	1	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

New Mexico

All races, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-
White, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	5	...	...	...	-	-	-	2	2	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	5	...	...	...	-	-	-	2	2	-	1	-
White, both sexes.....	2	...	...	...	-	-	-	-	2	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	2	-	-	-
Black, both sexes.....	2	...	...	...	-	-	-	1	-	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	1	-	-	1	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	1	-	-	-	-























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Alaska

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Arkansas

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Colorado

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Illinois

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Massachusetts

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by Male and Female.

Michigan

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by Male and Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Minnesota

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

New Hampshire

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

New Mexico

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Ohio

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

South Dakota

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for South Dakota. All cells contain dashes.

Tennessee

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Tennessee. All cells contain dashes.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Vermont

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Hemorrhage of pregnancy and childbirth and placenta previa (O20,O44-O46,O67,O72)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Complications predominantly related to the puerperium (O85-O92)

Arkansas

All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

California

All races, both sexes.....	6	...	...	...	-	-	1	-	2	1	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	6	...	...	...	-	-	1	-	2	1	2	-
White, both sexes.....	3	...	...	...	-	-	1	-	1	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	1	-	1	-	1	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	...	-	-	-	-	-	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	-	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	1	1	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Complications predominantly related to the puerperium (O85-O92)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Complications predominantly related to the puerperium (O85-O92)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Complications predominantly related to the puerperium (O85-O92)

Florida

All races, both sexes.....	11	...	...	...	-	1	2	2	3	3	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	11	...	...	...	-	1	2	2	3	3	-	-
White, both sexes.....	7	...	...	...	-	-	2	1	2	2	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	7	...	...	...	-	-	2	1	2	2	-	-
Black, both sexes.....	4	...	...	...	-	1	-	1	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	...	...	...	-	1	-	1	1	1	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Georgia

All races, both sexes.....	3	...	...	...	-	-	1	-	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	1	-	1	1	-	-
White, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-
Black, both sexes.....	1	...	...	...	-	-	1	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	1	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Complications predominantly related to the puerperium (O85-O92)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Complications predominantly related to the puerperium (O85-O92)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Complications predominantly related to the puerperium (O85-O92)

Maine

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Complications predominantly related to the puerperium (O85-O92)

Massachusetts

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Massachusetts categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Michigan categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Complications predominantly related to the puerperium (O85-O92)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Complications predominantly related to the puerperium (O85-O92)

Oregon

All races, both sexes.....	1	...	...	...	-	-	1	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	1	-	-	-	-	-
White, both sexes.....	1	...	...	...	-	-	1	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	1	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	4	...	...	...	-	-	-	1	1	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	...	...	...	-	-	-	1	1	1	1	-
White, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-
Black, both sexes.....	1	...	...	...	-	-	-	-	-	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	1	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	1	1	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Complications predominantly related to the puerperium (O85-O92)

Texas

All races, both sexes.....	6	...	...	...	-	1	-	1	1	3	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	6	...	...	...	-	1	-	1	1	3	-	-
White, both sexes.....	4	...	...	...	-	1	-	1	-	2	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	...	...	...	-	1	-	1	-	2	-	-
Black, both sexes.....	2	...	...	...	-	-	-	-	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	1	1	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Utah

All races, both sexes.....	2	...	...	...	-	-	-	-	-	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	-	1	1	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	-	-	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	-	1	1	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49  
ages 1 year years years years years years years years years years years

Complications predominantly related to the puerperium (O85-O92)

Vermont

All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	2	...	...	...	-	-	-	-	1	-	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	1	-	-	1
White, both sexes.....	1	...	...	...	-	-	-	-	-	-	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	-	-	1
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Complications predominantly related to the puerperium (O85-O92)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Complications predominantly related to the puerperium (O85-O92)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Complications predominantly related to the puerperium (O85-O92)

Hawaii

Table with 13 columns (Age groups) and 13 rows (Race categories) for Hawaii. All cells contain dashes (-).

Idaho

Table with 13 columns (Age groups) and 13 rows (Race categories) for Idaho. All cells contain dashes (-).









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Complications predominantly related to the puerperium (O85-O92)

Maine

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Maine. All cells contain dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Maryland. All cells contain dashes (-).



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Complications predominantly related to the puerperium (O85-O92)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Complications predominantly related to the puerperium (O85-O92)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Obstetric embolism (O88)												
Arkansas												
All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	2	...	...	...	-	-	1	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	1	-	1	-	-	-
White, both sexes.....	1	...	...	...	-	-	1	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	1	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Obstetric embolism (O88)												
Florida												
All races, both sexes.....	3	...	...	...	-	-	-	-	2	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	-	-	2	1	-	-
White, both sexes.....	3	...	...	...	-	-	-	-	2	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	-	-	2	1	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Georgia												
All races, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Obstetric embolism (O88)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Obstetric embolism (O88)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Obstetric embolism (O88)

Missouri

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Obstetric embolism (O88)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Obstetric embolism (O88)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Obstetric embolism (O88)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Obstetric embolism (O88)												
Oregon												
All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Pennsylvania												
All races, both sexes.....	2	...	...	...	-	-	-	1	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	1	-	1	-	-
White, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	1	-	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Obstetric embolism (O88)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Obstetric embolism (O88)

Texas

All races, both sexes.....	3	...	...	...	-	-	-	1	1	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	-	1	1	1	-	-
White, both sexes.....	2	...	...	...	-	-	-	1	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	1	-	1	-	-
Black, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Utah

All races, both sexes.....	2	...	...	...	-	-	-	-	-	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	-	1	1	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	-	-	1	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	-	1	1	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Obstetric embolism (O88)

Vermont

All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Illinois

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Iowa

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Massachusetts

Table with 13 columns (Age groups) and rows for Massachusetts: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Michigan

Table with 13 columns (Age groups) and rows for Michigan: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

New Mexico

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

North Carolina

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Ohio

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Oregon

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Obstetric embolism (O88)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other complications predominately related to the puerperium (O85-O87,O89-O92)

United States

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for United States (All races, White, Black, American Indian, Asian or Pacific Islander) split by sex.

Alabama

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Alabama (All races, White, Black, American Indian, Asian or Pacific Islander) split by sex.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Arkansas

All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

California

All races, both sexes.....	4	...	...	...	-	-	-	-	1	1	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	...	...	...	-	-	-	-	1	1	2	-
White, both sexes.....	2	...	...	...	-	-	-	-	1	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	1	-	1	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	...	-	-	-	-	-	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Oregon

All races, both sexes.....	1	...	...	...	-	-	1	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	1	-	-	-	-	-
White, both sexes.....	1	...	...	...	-	-	1	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	1	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Pennsylvania

All races, both sexes.....	2	...	...	...	-	-	-	-	1	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	1	-	1	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	1	...	...	...	-	-	-	-	-	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	-	1	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Colorado

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Delaware

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Maine

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maine. All cells contain dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maryland. All cells contain dashes (-).

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Massachusetts

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by Male and Female.

Michigan

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by Male and Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

North Carolina

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Oregon

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Oregon. All cells contain dashes.

Pennsylvania

Table with 13 columns (Age groups) and 13 rows (All races, White, Black, American Indian, Asian or Pacific Islander) for Pennsylvania. All cells contain dashes.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other complications predominately related to the puerperium (O85-O87,O89-O92)

Wisconsin

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)

Arkansas

All races, both sexes.....	3	...	...	...	-	-	2	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	2	1	-	-	-	-
White, both sexes.....	3	...	...	...	-	-	2	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	2	1	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

California

All races, both sexes.....	20	...	...	...	-	-	4	5	5	3	2	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	20	...	...	...	-	-	4	5	5	3	2	1
White, both sexes.....	14	...	...	...	-	-	3	3	3	3	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	14	...	...	...	-	-	3	3	3	3	2	-
Black, both sexes.....	5	...	...	...	-	-	1	2	1	-	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	5	...	...	...	-	-	1	2	1	-	-	1
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	1	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	1	-	-	-

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)

New Mexico

All races, both sexes.....	3	...	...	...	-	-	-	2	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	-	2	-	1	-	-
White, both sexes.....	3	...	...	...	-	-	-	2	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	-	2	-	1	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	14	...	...	...	-	-	2	3	5	1	1	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	14	...	...	...	-	-	2	3	5	1	1	1
White, both sexes.....	4	...	...	...	-	-	-	1	2	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	...	...	...	-	-	-	1	2	-	1	-
Black, both sexes.....	9	...	...	...	-	-	2	2	3	-	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	9	...	...	...	-	-	2	2	3	-	-	1
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)

Vermont

All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

Virginia

All races, both sexes.....	4	...	...	...	-	-	-	3	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	...	...	...	-	-	-	3	-	1	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	2	...	...	...	-	-	-	2	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	2	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	2	...	...	...	-	-	-	1	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	1	-	1	-	-































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

All other direct obstetric causes (O10,O12,O21-O43,O47-O66,O68-O71,O73-O75)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Obstetric death of unspecified cause (O95)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Obstetric death of unspecified cause (O95)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Obstetric death of unspecified cause (O95)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Obstetric death of unspecified cause (O95)

Nebraska

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows (All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female)

Nevada

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and 13 rows (All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female)





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Obstetric death of unspecified cause (O95)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Obstetric death of unspecified cause (O95)

Ohio

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Obstetric death of unspecified cause (O95)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Obstetric death of unspecified cause (O95)

Washington

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Arkansas

All races, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
White, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

California

All races, both sexes.....	55	...	...	...	-	1	9	7	14	13	2	6
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	55	...	...	...	-	1	9	7	14	13	2	6
White, both sexes.....	37	...	...	...	-	1	6	3	7	10	2	5
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	37	...	...	...	-	1	6	3	7	10	2	5
Black, both sexes.....	13	...	...	...	-	-	3	2	5	2	-	1
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	13	...	...	...	-	-	3	2	5	2	-	1
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	5	...	...	...	-	-	-	2	2	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	5	...	...	...	-	-	-	2	2	1	-	-













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years years

Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)

Minnesota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Minnesota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Mississippi categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
--	----------	--------------	-----------	-----------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------

Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)

New Mexico

All races, both sexes.....	2	...	...	...	-	-	-	-	-	-	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	-	-	2	-
White, both sexes.....	2	...	...	...	-	-	-	-	-	-	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	2	...	...	...	-	-	-	-	-	-	2	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-

New York

All races, both sexes.....	12	...	...	...	-	-	1	4	3	2	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	12	...	...	...	-	-	1	4	3	2	2	-
White, both sexes.....	1	...	...	...	-	-	-	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	1	-	-	-	-
Black, both sexes.....	8	...	...	...	-	-	1	3	2	2	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	8	...	...	...	-	-	1	3	2	2	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	3	...	...	...	-	-	-	-	1	-	2	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	3	...	...	...	-	-	-	-	1	-	2	-











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)

Texas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Iowa

Table with 13 columns (Age groups) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Age groups) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)

Massachusetts

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by Male and Female.

Michigan

Table with 13 columns (Age groups) and rows for All races, White, Black, American Indian, and Asian or Pacific Islander, each split by Male and Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

North Carolina

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (O96-O97)

Ohio

Table with 13 columns (Age groups) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Age groups) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other deaths related to pregnancy, childbirth and the puerperium (096-097)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see <http://www.cdc.gov/nchs> and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	All ages	Under 1 year	1-4 years	5-9 years	10-14 years	15-19 years	20-24 years	25-29 years	30-34 years	35-39 years	40-44 years	45-49 years
Indirect obstetric deaths (O98-O99)												
Arkansas												
All races, both sexes.....	1	...	...	...	-	-	1	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	1	-	-	-	-	-
White, both sexes.....	1	...	...	...	-	-	1	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	1	-	-	-	-	-
Black, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
American Indian 1/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
Asian or Pacific Islander 2/, both sexes.....	-	...	...	...	-	-	-	-	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	-	...	...	...	-	-	-	-	-	-	-	-
California												
All races, both sexes.....	7	...	...	...	-	-	1	3	-	2	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	7	...	...	...	-	-	1	3	-	2	1	-
White, both sexes.....	4	...	...	...	-	-	1	2	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	4	...	...	...	-	-	1	2	-	1	-	-
Black, both sexes.....	1	...	...	...	-	-	-	1	-	-	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	1	-	-	-	-
American Indian 1/, both sexes.....	1	...	...	...	-	-	-	-	-	-	1	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	-	1	-
Asian or Pacific Islander 2/, both sexes.....	1	...	...	...	-	-	-	-	-	1	-	-
Male.....	...	...	...	...	...	...	...	...	...	...	...	...
Female.....	1	...	...	...	-	-	-	-	-	1	-	-



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Indirect obstetric deaths (O98-O99)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Indirect obstetric deaths (O98-O99)

Florida

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Indirect obstetric deaths (O98-O99)

Hawaii

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Indirect obstetric deaths (O98-O99)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Indirect obstetric deaths (O98-O99)

North Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Carolina categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for North Dakota categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Indirect obstetric deaths (O98-O99)

United States

Table with 13 columns for age groups and 13 rows for race/sex categories (All races, White, Black, American Indian, Asian or Pacific Islander) for both sexes and by gender.

Alabama

Table with 13 columns for age groups and 13 rows for race/sex categories (All races, White, Black, American Indian, Asian or Pacific Islander) for both sexes and by gender.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Indirect obstetric deaths (O98-O99)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Indirect obstetric deaths (O98-O99)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

XVI. Certain conditions originating in the perinatal period (P00-P96)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

XVI. Certain conditions originating in the perinatal period (P00-P96)

Colorado

Table with 13 columns (Age groups) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Age groups) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XVI. Certain conditions originating in the perinatal period (P00-P96)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

XVI. Certain conditions originating in the perinatal period (P00-P96)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XVI. Certain conditions originating in the perinatal period (P00-P96)

Maine

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maine. All cells contain dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maryland. All cells contain dashes (-).



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XVI. Certain conditions originating in the perinatal period (P00-P96)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XVI. Certain conditions originating in the perinatal period (P00-P96)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

XVI. Certain conditions originating in the perinatal period (P00-P96)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00-P04)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Newborn affected by maternal factors and by complications of pregnancy, labor and delivery (P00-P04)

Colorado

Table with 13 columns for age groups and one for 'Age not stated'. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns for age groups and one for 'Age not stated'. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Disorders related to short gestation and low birth weight, not elsewhere classified (P07)

Kentucky

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Disorders related to short gestation and low birth weight, not elsewhere classified (P07)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Disorders related to short gestation and low birth weight, not elsewhere classified (P07)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Disorders related to short gestation and low birth weight, not elsewhere classified (P07)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Disorders related to short gestation and low birth weight, not elsewhere classified (P07)

Minnesota

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Minnesota. All cells contain dashes (-).

Mississippi

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Mississippi. All cells contain dashes (-).

























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Disorders related to short gestation and low birth weight, not elsewhere classified (P07)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Disorders related to short gestation and low birth weight, not elsewhere classified (P07)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Birth trauma (P10-P15)

Delaware

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Birth trauma (P10-P15)

Hawaii

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Birth trauma (P10-P15) in Hawaii, categorized by race and sex.

Idaho

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Birth trauma (P10-P15) in Idaho, categorized by race and sex.





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Birth trauma (P10-P15)

Oregon

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Birth trauma (P10-P15)

Rhode Island

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Birth trauma (P10-P15)

South Dakota

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Birth trauma (P10-P15)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Birth trauma (P10-P15)

Washington

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Birth trauma (P10-P15) in Washington, categorized by race and sex.

West Virginia

Table with 12 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Birth trauma (P10-P15) in West Virginia, categorized by race and sex.

















































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Birth trauma (P10-P15)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intrauterine hypoxia and birth asphyxia (P20-P21)

Delaware

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.















































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Intrauterine hypoxia and birth asphyxia (P20-P21)

Wisconsin

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

Colorado

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns: Cause of death, 50-54 years, 55-59 years, 60-64 years, 65-69 years, 70-74 years, 75-79 years, 80-84 years, 85-89 years, 90-94 years, 95-99 years, 100 years and over, Age not stated. Rows include All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

North Carolina

Table with 13 columns (Age groups) and rows for North Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

North Dakota

Table with 13 columns (Age groups) and rows for North Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

Texas

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Texas. All cells contain dashes (-).

Utah

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Utah. All cells contain dashes (-).





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Intrauterine hypoxia and birth asphyxia (P20-P21)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Respiratory distress of newborn (P22)

Delaware

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Delaware categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Dist. of Columbia categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Respiratory distress of newborn (P22)

Iowa

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Iowa: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Kansas

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Respiratory distress of newborn (P22)

Kentucky

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Respiratory distress of newborn (P22)

New Hampshire

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Hampshire categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for New Jersey categories: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

















Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All Under 1-4 5-9 10-14 15-19 20-24 25-29 30-34 35-39 40-44 45-49
ages 1 year years years years years years years years years years

Respiratory distress of newborn (P22)

Vermont

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





























Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Respiratory distress of newborn (P22)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

































































































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other respiratory conditions originating in the perinatal period (P23-P28)

Arkansas

Table with 13 columns (Age groups) and rows for Arkansas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

California

Table with 13 columns (Age groups) and rows for California: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other respiratory conditions originating in the perinatal period (P23-P28)

Colorado

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Colorado. All cells contain dashes (-).

Connecticut

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Connecticut. All cells contain dashes (-).







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other respiratory conditions originating in the perinatal period (P23-P28)

Hawaii

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Hawaii. All cells contain dashes (-).

Idaho

Table with 13 columns (Age groups) and 13 rows (Race and sex categories) for Idaho. All cells contain dashes (-).





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other respiratory conditions originating in the perinatal period (P23-P28)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other respiratory conditions originating in the perinatal period (P23-P28)

Maine

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maine. All cells contain dashes (-).

Maryland

Table with 13 columns (Age groups) and 13 rows (Race categories) for Maryland. All cells contain dashes (-).



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other respiratory conditions originating in the perinatal period (P23-P28)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.









Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other respiratory conditions originating in the perinatal period (P23-P28)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other respiratory conditions originating in the perinatal period (P23-P28)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other respiratory conditions originating in the perinatal period (P23-P28)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other respiratory conditions originating in the perinatal period (P23-P28)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections specific to the perinatal period (P35-P39)

Colorado

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Colorado: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Connecticut

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Connecticut: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections specific to the perinatal period (P35-P39)

Illinois

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

All ages Under 1 year 1-4 years 5-9 years 10-14 years 15-19 years 20-24 years 25-29 years 30-34 years 35-39 years 40-44 years 45-49 years

Infections specific to the perinatal period (P35-P39)

Wisconsin

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Cause of death, All ages, Under 1 year, 1-4 years, 5-9 years, 10-14 years, 15-19 years, 20-24 years, 25-29 years, 30-34 years, 35-39 years, 40-44 years, 45-49 years) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Florida

Table with 13 columns (Age groups) and rows for Florida: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Georgia

Table with 13 columns (Age groups) and rows for Georgia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Hawaii

Table with 13 columns (Age groups) and rows for Hawaii: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Idaho

Table with 13 columns (Age groups) and rows for Idaho: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Maine

Table with 13 columns (Age groups) and rows for Maine: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Maryland

Table with 13 columns (Age groups) and rows for Maryland: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Minnesota

Table with 13 columns (Age groups) and rows for Minnesota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Mississippi

Table with 13 columns (Age groups) and rows for Mississippi: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Missouri

Table with 13 columns (Age groups) and rows for Missouri: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Montana

Table with 13 columns (Age groups) and rows for Montana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

New Hampshire

Table with 13 columns for age groups and one for 'Age not stated'. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns for age groups and one for 'Age not stated'. Rows include: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

New Mexico

Table with 13 columns (Age groups) and rows for New Mexico: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New York

Table with 13 columns (Age groups) and rows for New York: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Ohio

Table with 13 columns (Age groups) and rows for Ohio: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Oklahoma

Table with 13 columns (Age groups) and rows for Oklahoma: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Oregon

Table with 13 columns (Age groups) and rows for Oregon: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Pennsylvania

Table with 13 columns (Age groups) and rows for Pennsylvania: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Rhode Island

Table with 13 columns (Age groups) and rows for Rhode Island: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

South Carolina

Table with 13 columns (Age groups) and rows for South Carolina: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Vermont

Table with 13 columns (Age groups) and rows for Vermont: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Virginia

Table with 13 columns (Age groups) and rows for Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Washington

Table with 13 columns (Age groups) and rows for Washington: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

West Virginia

Table with 13 columns (Age groups) and rows for West Virginia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Infections specific to the perinatal period (P35-P39)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





























































Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

Alaska

Table with 13 columns (Age groups) and rows for Alaska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Arizona

Table with 13 columns (Age groups) and rows for Arizona: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

Delaware

Table with 13 columns (Age groups) and rows for Delaware: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Dist. of Columbia

Table with 13 columns (Age groups) and rows for Dist. of Columbia: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.







Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

Illinois

Table with 13 columns (Age groups) and rows for Illinois: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Indiana

Table with 13 columns (Age groups) and rows for Indiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.



Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

Kentucky

Table with 13 columns (Age groups) and rows for Kentucky: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Louisiana

Table with 13 columns (Age groups) and rows for Louisiana: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.











Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

Nebraska

Table with 13 columns (Age groups) and rows for Nebraska: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Nevada

Table with 13 columns (Age groups) and rows for Nevada: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

New Hampshire

Table with 13 columns (Age groups) and rows for New Hampshire: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

New Jersey

Table with 13 columns (Age groups) and rows for New Jersey: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.













Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 years 55-59 years 60-64 years 65-69 years 70-74 years 75-79 years 80-84 years 85-89 years 90-94 years 95-99 years 100 years and over Age not stated

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

South Dakota

Table with 13 columns (Age groups) and rows for South Dakota: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Tennessee

Table with 13 columns (Age groups) and rows for Tennessee: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex 50-54 55-59 60-64 65-69 70-74 75-79 80-84 85-89 90-94 95-99 100 years and over Age not stated

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

Texas

Table with 13 columns (Age groups) and rows for Texas: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Utah

Table with 13 columns (Age groups) and rows for Utah: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.





Worktable III. Deaths from 358 selected causes, by 5-year age groups, race, and sex: United States and each state, 2007

[Race categories are consistent with the 1977 Office of Management and Budget (OMB) standards. In 2007, multiple-race data were reported by California, Connecticut, Delaware, District of Columbia, Florida, Hawaii, Idaho, Kansas, Maine, Michigan, Minnesota, Montana, Nebraska, New Hampshire, New Jersey, New Mexico, New York, Ohio, Oklahoma, Oregon, Rhode Island, South Carolina, South Dakota, Texas, Utah, Washington, Wisconsin, and Wyoming. The multiple-race data for these areas were bridged to the single-race categories of the 1977 OMB standards for comparability with other areas; see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality." Data for specified races other than white and black should be interpreted with caution because of inconsistencies between reporting race on death certificates and on censuses and surveys. For explanation of asterisks preceding cause-of-death codes, see http://www.cdc.gov/nchs and refer to "Detailed technical notes to the United States 2007 data-mortality"]

Cause of death (Based on the International Classification of Diseases, Tenth Revision, Second Edition, 2004), area, race, and sex

	50-54 years	55-59 years	60-64 years	65-69 years	70-74 years	75-79 years	80-84 years	85-89 years	90-94 years	95-99 years	100 years and over	Age not stated
--	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	-------------	--------------------	----------------

Other and unspecified conditions originating in the perinatal period (P05,P08,P29,P50-P96)

Wisconsin

Table with 13 columns (Age groups) and rows for Wisconsin: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.

Wyoming

Table with 13 columns (Age groups) and rows for Wyoming: All races, both sexes; Male; Female; White, both sexes; Male; Female; Black, both sexes; Male; Female; American Indian 1/, both sexes; Male; Female; Asian or Pacific Islander 2/, both sexes; Male; Female.